

Li Ye, Ceyuan Haijing, volumes 1-12

English Translation

Modern LaTeX Editions of Public-Domain Mathematics Manuscripts

測圓海鏡

Sea Mirror of Circle Measurements

卷一—卷三 | Volumes I–III

BILINGUAL EDITION — 雙語版

Original Chinese • English Translation

元 李冶 撰

(Li Ye / Li Zhi, 1192–1279)

以天元術解勾股容圓一百七十問

*Using the Method of the Celestial Element (Tian Yuan Shu)
to Solve 170 Problems on the Inscribed Circle*

A foundational text of medieval Chinese algebra,
completed in the year 1248 of the Common Era.

Typeset in L^AT_EX with XeLaTeX + xeCJK
Bilingual edition with parallel Chinese-English columns

Contents

1	Introduction / 引言	2
2	Original Preface / 原序	3
3	Volume 1 / 卷一	5
3.1	Circle City Diagram / 圓城圖式	5
3.2	General Rate Names / 總率名號	5
3.3	Current Question Numbers / 今問正數	6
3.4	Identification Miscellaneous Records / 識別雜記	7
4	Volume 2 / 卷二	9
5	Volume 3 / 卷三	13
6	The Tian Yuan Shu Method / 天元術	17
6.1	Methodology / 方法論	17
6.2	Polynomial Notation / 多項式記號	17
6.3	Example: Quadratic Equation / 示例: 二次方程	18
6.4	Historical Significance / 歷史意義	18

1 Introduction / 引言

測圓海鏡簡介

《測圓海鏡》乃李冶（李冶，1192–1279）之傑作。李冶為宋元四大數學家之一，與楊輝、秦九韶、朱世傑齊名。此書成於1248年，系統運用**天元術**（多項式代數方法）求解圓與直角三角形相關之一百七十問。全書結構如下：

- **卷一**：理論基礎——圓城圖式、總率名號、今問正數、識別雜記（含六百九十二條幾何公式）
- **卷二至卷十二**：一百七十問，以天元術求解，方程次數自一次至六次。

本文檔包含**卷一至卷三**之全文，即全部基礎理論及首批問題。

Introduction

The *Ceyuan Haijing* (測圓海鏡, literally “Sea Mirror of Circle Measurements”) is the masterpiece of Li Ye (李冶, also known as Li Zhi, 1192–1279), one of the four great mathematicians of the Song–Yuan period of China, alongside Yang Hui, Qin Jiushao, and Zhu Shijie.

Completed in 1248, this work systematically applies the **tian yuan shu** (天元術, “method of the celestial element”)—an algebraic method for setting up and solving polynomial equations—to a total of 170 geometric problems concerning a circle inscribed in or related to a right triangle (勾股容圓).

The book is organized as follows:

- **Volume 1** establishes the theoretical foundation: the *Circle City Diagram* (圓城圖式), the *General Rate Names* (總率名號), the *Current Question Numbers* (今問正數), and the *Identification Miscellaneous Records* (識別雜記) containing 692 geometric formulas.
- **Volumes 2–12** present 170 problems, solved by the **tian yuan shu** method. The problems involve equations of degrees ranging from 1 to 6.

The present document contains the complete text of **Volumes 1–3**, comprising all foundational material and the first suite of problems.

2 Original Preface / 原序

原序

數本難窮，吾欲以力強窮之，彼其數不惟不能得其凡，而吾之力且憊矣。然則數果不可以窮耶？既已名之數矣，則又何為而不可窮也。

故謂數為難窮斯可，謂數為不可窮斯不可。何則？彼其冥冥之中，固有昭昭者存，夫昭昭者其自然之數也。非自然之數，其自然之理也。數一出於自然，吾欲以力強窮之，使隸首復生亦末如之何也已。

苟能推自然之理，以明自然之數，則雖遠而乾端坤倪，幽而神情鬼狀，未有不合者矣。

予自幼喜算數，恆病夫考圓之術例出於牽強，殊乖於自然，如古率、徽率、密率之不同，截弧、截矢、截背之互見，內外諸角，析剖支條，莫不各自名家，與世作法。及反覆研究，卒無以當吾心焉。

老大以來，得洞淵九容之說，日夕玩繹，而向之病我者，使爆然落去而無遺余。山中多暇，客有從余求其說者，於是乎又為衍之，遂累一百七十問。

既成編，客復目之《測圓海鏡》，蓋取夫天臨海鏡之義也。

Original Preface

Numbers are fundamentally difficult to exhaust; if I try to force them by effort, not only will I fail to apprehend their essence, but my strength will be depleted as well. Yet can numbers truly not be exhausted? Since they are already called *numbers*, why then could they not be exhausted?

It is acceptable to say that numbers are difficult to exhaust; but it is unacceptable to say they cannot be exhausted. Why? In the midst of darkness, there is indeed brightness. That brightness is the natural number. It is not merely the natural number—it is the natural principle. Since numbers arise from nature, if I try to force them by effort, even if Li Shou (the legendary inventor of arithmetic) were reborn, he could do no better.

If one can extend natural principles to illuminate natural numbers, then even things as distant as the corners of heaven and earth, or as mysterious as spirits and ghosts, will all be in agreement.

Since youth I have delighted in arithmetic. I was always troubled that the methods for investigating circles seemed forced and artificial, contrary to nature—the ancient rate, Hui's rate, and the close rate all differing; the arc, sagitta, and back each being considered separately; the internal and external angles analyzed into minute subdivisions—each school establishing its own name and method for the world. After repeated study, none could satisfy my mind.

In old age, I obtained the *Dongyuan Nine Containments* (洞淵九容) theory, and pondered it day and night. What had previously troubled me suddenly fell away completely, leaving no trace. With leisure time in the mountains, when guests asked me to explain this theory, I expanded upon it, eventually accumulating **one hundred and seventy problems**.

When the compilation was complete, a guest named it *Ceyuan Haijing*—taking the meaning of “heaven overlooking the sea mirror.”

昔半山老人集《唐百家詩選》，自謂廢日力於此，良可惜。明道先生以上蔡謝君記誦為玩物喪志。夫文史尚矣，猶之為不足貴，況九九賤技能乎！

嗜好酸鹹，平生每痛自戒勅，竟莫能已，類有物憑之者，吾亦不知其然而然也。

故嘗私為之解曰：由技兼於事者言之，夷之禮，夔之樂，亦不免為一技。由技進乎道者言之，石之斤，扁之輪，非聖人之所與乎。

覽吾之編，察吾苦心，其憫我者當百數，其笑我者當千數。乃若吾之所自得，則自得焉耳，寧復為人憫笑計哉。

李冶序

In the past, the Old Man of Banshan compiled the *Anthology of One Hundred Tang Poets*, saying that to waste his days on such a thing was truly a pity. Master Mingdao criticized Xie of Shangcai for memorizing texts as “playing with things and losing one’s ambition.” Literature and history are esteemed subjects, yet even they are considered not worth valuing—how much less the lowly skill of multiplication!

My fondness for such sour and salty pursuits—throughout my life I have sternly admonished myself against them, yet in the end I could not stop, as if some force were driving me. I myself do not know why it is so.

So I have privately explained it thus: speaking from the perspective of skills combined with affairs, even the rites of Yi and the music of Kui were not exempt from being a single skill. Speaking from the perspective of advancing from skill to the Way, was not the stone-axe of [the sage] and the wheel of Bian [the wheelwright] praised by the sages?

Those who read my compilation and perceive my earnest efforts—those who pity me will number in the hundreds, those who laugh at me will number in the thousands. As for what I myself have gained, I have simply gained it; how could I calculate for the sake of being pitied or laughed at?

— Preface by Li Ye

3 Volume 1 / 卷一

卷一為全書之理論基礎，分為四大部分：

1. 圓城圖式——主幾何圖形
2. 總率名號——十五個直角三角形及其元素之定義
3. 今問正數——各線段之數值
4. 識別雜記——六百九十二條幾何公式

3.1 Circle City Diagram / 圓城圖式

圓城圖式為一百七十問所基之主幾何圖形。其構為一大直角三角形（勾股形），內切一圓，並設特定之點與線。

其情境如下：

假令有圓城一所，不知周徑，四面開門，門外縱橫各有十字大道。其西北十字道頭定為乾地，其東北十字道頭定為艮地，其東南十字道頭定為巽地，其西南十字道頭定為坤地。所有測望雜法——設問如後。

二十二名點 / The 22 Named Points

最大直角三角形之頂點為天、地、乾。內切圓之心稱心。各點以單字命名：

天	地	乾	心	日	南	北	川
東	西	艮	坤	山	月	巽	青
泉	朱	金	泛	旦	夕		

3.2 General Rate Names / 總率名號

全書研究十五個直角三角形，皆以天地線之一段為弦（弦，即斜邊）。總率名號定義各三角形之三邊。每個三角形：

- 弦 (c) ——斜邊
- 股 (b) ——較長之直邊
- 勾 (a) ——較短之直邊

Volume 1 is the theoretical foundation of the entire work. It comprises four major sections:

1. **Circle City Diagram** (圓城圖式) – the master geometric configuration
2. **General Rate Names** (總率名號) – definitions of all 15 right triangles and their elements
3. **Current Question Numbers** (今問正數) – numerical values for all line segments
4. **Identification Miscellaneous Records** (識別雜記) – 692 geometric formulas

The *Circle City Diagram* is the master geometric configuration upon which all 170 problems are based. It consists of a large right triangle (called 勾股形, “leg-leg shape”) with its inscribed circle, together with specific points and lines.

The scenario Li Ye constructs is:

Suppose there is a circular city whose circumference and diameter are unknown. It has gates on all four sides, and outside each gate there are crossroads running north-south and east-west. The northwest crossroad is designated **Qian** (乾), the northeast **Gen** (艮), the southeast **Xun** (巽), and the southwest **Kun** (坤). All surveying methods are set forth as questions below.

The largest right triangle has vertices **Tian** (天, Heaven), **Di** (地, Earth), and **Qian** (乾). The center of the inscribed circle is called **Xin** (心, Heart). Key points include intersections of horizontal and vertical lines through the center and other constructed lines. Each point is designated by a single Chinese character:

Tian	Di	Qian	Xin	Ri	Nan	Bei	Chuan
Dong	Xi	Gen	Kun	Shan	Yue	Xun	Qing
Quan	Zhu	Jin	Fan	Dan	Xi		

The work studies 15 **right triangles**, all having a segment of the *Tiandi* (天地) line as their hypotenuse (弦, “string”). The *General Rate Names* define these triangles and their three sides.

In each triangle:

- 弦 (c) – the hypotenuse (斜邊)
- 股 (b) – the longer vertical leg
- 勾 (a) – the shorter horizontal leg

十五三角形表 / Table of 15 Triangles

No.	名稱 / Name	頂點 / Vertices	弦	股	勾
1	通 (Tong/General)	天-地-乾	通弦	通股	通勾
2	邊 (Bian/Side)	天-西-川	邊弦	邊股	邊勾
3	底 (Di/Base)	日-地-北	底弦	底股	底勾
4	黃廣 (Huangguang)	天-山-金	黃廣弦	黃廣股	黃廣勾
5	黃長 (Huangchang)	月-地-泉	黃長弦	黃長股	黃長勾
6	上高 (Shanggao)	天-日-旦	上高弦	上高股	上高勾
7	下高 (Xiagao)	日-山-朱	下高弦	下高股	下高勾
8	上平 (Shangping)	月-川-青	上平弦	上平股	上平勾
9	下平 (Xiaping)	川-地-夕	下平弦	下平股	下平勾
10	大差 (Dacha)	天-月-坤	大差弦	大差股	大差勾
11	小差 (Xiaocha)	山-地-艮	小差弦	小差股	小差勾
12	皇極 (Huangji)	日-川-心	皇極弦	皇極股	皇極勾
13	太虛 (Taixu)	月-山-泛	太虛弦	太虛股	太虛勾
14	明 (Ming)	日-月-南	明弦	明股	明勾
15	夷 (Zhuan)	山-川-東	夷弦	夷股	夷勾

注：上高 = 下高，上平 = 下平，故十五三角形中實僅十三個互異。

Note: 上高 = 下高 and 上平 = 下平, so among 15 triangles, only 13 are distinct.

3.3 Current Question Numbers / 今問正數

此節給出圖中每條線段之數值。內切圓之半徑定為一百二十步，作為標準單位。

This section gives the numerical value of every line segment in the diagram. The radius of the inscribed circle is taken as 120 steps (步), which serves as the standard unit.

1. 通 (Tong / 通用)

弦六百八十	勾三百二十	股六百
勾股和九百二十	較二百八十	
勾和一千	較三百六十	
股和一千二百八十	較八十	
較和九百六十	較四百	
和和一千六百	較二百四十	

1. Tong (General)

Hypotenuse 680	Short leg 320, Long leg 920
Sum of legs 920	Difference 280
Leg-hypotenuse sum 1000	Difference 360
Leg-hypotenuse sum 1280	Difference 80
Difference-sum 960	Difference 400
Sum-sum 1600	Difference 240

That is: hypotenuse $c = 680$, short leg $a = 320$, long leg $b = 920$.

2. 邊 (Bian / Side)

弦五百四十四	勾二百五十六	股四百八十
勾股和七百三十六	較二百二十四	
勾和八百	較二百八十八	
股和一千零二十四	較六十四	
較和七百六十八	較三百二十	
和和一千二百八十	較一百九十二	

2. Bian (Side)

Hypotenuse 544	Short leg 256, Long leg 480
Sum of legs 736	Difference 224
Leg-hypotenuse sum 800	Difference 288
Leg-hypotenuse sum 1024	Difference 64
Difference-sum 768	Difference 320
Sum-sum 1280	Difference 192

3. 底 (Di / Base)

弦四百二十五	勾二百	股三百七十五
勾股和五百七十五	較一百七十五	
勾和六百二十五	較二百二十五	
股和八百	較五十	
較和六百	較二百五十	
和和一千	較一百五十	

3. Di (Base)

Hypotenuse 425	Short leg 200, Long leg 375
Sum of legs 575	Difference 175
Leg-hypotenuse sum 625	Difference 225
Leg-hypotenuse sum 800	Difference 50
Difference-sum 600	Difference 250
Sum-sum 1000	Difference 150

4. 黃廣 (Huang Guang)

弦五百一十 勾二百四十 股四百五十
 (即城徑也)
勾股和六百九十 較二百一十

其餘十一三角形依同樣模式，每個十三個數量（弦、勾、股、勾股和、勾股較、勾弦和、勾弦較、股弦和、股弦較、弦較和、弦較較、弦和和、弦和較），共 $15 \times 13 = 195$ 個數值條目。

4. Huang Guang

Hypotenuse 510 Short leg 240, Long leg 450
 (this is the city diameter)
Sum of legs 690 Difference 210

The remaining 11 triangles follow the same pattern, each with 13 numerical quantities (弦, 勾, 股, 勾股和, 勾股較, 勾弦和, 勾弦較, 股弦和, 股弦較, 弦較和, 弦較較, 弦和和, 弦和較), giving a total of $15 \times 13 = 195$ numerical entries.

3.4 Identification Miscellaneous Records / 識別雜記

此為全書之理論核心，含**六百九十二條幾何公式**，關聯圖中各線段。此等公式純為幾何定理，與具體數值無關。後續各卷之問題皆以此等公式配合天元術求解。本分為八部分：

- 1. 諸雜名目——一般定義及三十餘條定理
- 2. 五和五較
- 3. 諸弦
- 4. 大小差
- 5. 諸差
- 6. 諸率互見
- 7. 四位拾遺
- 8. 拾遺

諸雜名目示例：

This is the theoretical heart of the work. It contains **692 geometric formulas** relating the various line segments in the diagram. These formulas are purely geometric theorems, independent of any particular numerical values. The problems in all subsequent volumes are solved by using these formulas together with the tian yuan shu. The section is divided into eight parts:

- 1. Various Names (諸雜名目) – general definitions and 30+ theorems
- 2. Five Sums and Five Differences (五和五較)
- 3. Various Hypotenuses (諸弦)
- 4. Large and Small Differences (大小差)
- 5. Various Differences (諸差)
- 6. Mutual Relations of Rates (諸率互見)
- 7. Four-Place Supplement (四位拾遺)
- 8. Supplement (拾遺)

Sample Definitions from Various Names:

名稱 / Name	定義 / Definition
內率	明勾 × 裏股 / Ming 勾 × Zhuan 股
外率	明股 × 裏勾 / Ming 股 × Zhuan 勾
虛率	虛勾 × 虛股 / Xu 勾 × Xu 股
角差	高股 – 平勾 / Gao 股 – Ping 勾
次差	(明股 – 明勾) + (裏股 – 裏勾)
混同和	黃長股 + 黃廣勾 / Huangchang 股 + Huangguang 勾
傍差	(明股 – 明勾) – (裏股 – 裏勾)

定理示例：

Sample Theorems:

- 凡大差小差相乘為半段徑幂。
 - 大差勾小差股相乘同上。
 - 黃廣股黃長勾相乘為經幂。
 - 勾股和即弦黃和。
- The product of the large difference and small difference equals half the square of the diameter.
 - The product of the large-difference short-leg and the small-difference long-leg equals the same.
 - The product of the Huangguang long-leg and Huangchang short-leg equals the square of the diameter.
 - The sum of the two legs equals the sum of the hypotenuse and the diameter of the inscribed circle: $a + b = c + d$, where d is the diameter of the inscribed circle.

4 Volume 2 / 卷二

正率一十四問

此十四問構成「洞淵九容」序列，源自道教大師李思聰（號洞淵）之著作。其呈現直角三角形中九種基本容圓類型，另加五題。

4.0.1 第一問 / Problem 1

問 甲乙二人俱在乾地，乙東行三百二十步而立，甲南行六百步望見乙，問徑幾里？

答 城徑二百四十步。

術 此為勾股容圓也。以勾股相乘，倍之為實，並勾股幕以求弦，復加入勾股共，以為法。

草 置甲南行六百步在地，以乙東行三百二十步乘之，得一十九萬二千步，倍之得三十萬四千步為實。以乙東行步自之，得一十萬零二千四百步為勾幕；以甲南行步自之，得三十六萬步為股幕。二幕相並，得四十六萬二千四百步為方實，以平方開之，得六百八十步，則弦也。以弦加勾股共，共得一千六百步以為法。如法而一，得二百四十步，則城徑也。合問。

4.0.2 第二問 / Problem 2

問 甲乙二人俱在西門，乙東行二百五十六步，甲南行四百八十步望見乙，問答同前。

答 城徑二百四十步。

術 此為勾上容圓也。以勾股相乘，倍之為實，並勾股幕以求弦，加入股以為法。

草 置甲南行四百八十步在地，以乙東行二百五十六步乘之，得一十二萬二千八百八十步，倍之得二十四萬五千七百六十步為實。以乙東行步自之，得六萬五千五百三十六步為勾幕；以甲南行步自之，得二十三萬零四百步為股幕。勾股幕相並，得二十九萬五千九百三十六步為方實，以平方開之，得五百四十四步為弦也。以弦加入南行步，共得一千零二十四步以為法。而一，得二百四十步則城徑。合問。

Fourteen Questions on Standard Rates

These fourteen problems form the “Dongyuan Nine Containments” (洞淵九容) sequence, derived from the earlier work of the Daoist master Li Sicong (李思聰, known as “Master Dongyuan”). They present the nine fundamental types of circles contained in or related to a right triangle, plus five additional problems.

Problem. Two people, A and B, both stand at Qian. B walks 320 steps east and stops. A walks 600 steps south and sees B. What is the diameter?

Answer. The city diameter is 240 steps.

Method. This is the “circle contained in the right triangle” case. Multiply the two legs and double it to obtain the dividend. Combine the squares of the legs to find the hypotenuse, then add the sum of the legs to form the divisor.

Working. Place A’s 600 steps south. Multiply by B’s 320 steps east, obtaining 192,000 steps; double to get 384,000 as the dividend. Square B’s eastward steps to get 102,400 as the short-leg square; square A’s southward steps to get 360,000 as the long-leg square. Add the two squares: 462,400. Extract the square root to get 680, which is the hypotenuse. Add the sum of the legs to the hypotenuse: 1,600 as the divisor. Divide: 240 steps, the city diameter. This agrees with the question.

Problem. Two people both stand at the West Gate. B walks 256 steps east, A walks 480 steps south and sees B. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “circle on the short leg” case. Multiply the two legs and double as dividend. Combine the squares of the legs to find the hypotenuse; add the long leg as divisor.

Working. Place A’s 480 steps south. Multiply by B’s 256 steps east, obtaining 122,880; double to get 245,760 as dividend. Square B’s eastward steps: 65,536 as the short-leg square. Square A’s southward steps: 230,400 as the long-leg square. Combine: 295,936. Extract square root: 544, the hypotenuse. Add the southward steps: 1,024 as divisor. Divide: 240 steps, the city diameter. This agrees.

4.0.3 第三問 / Problem 3

問 甲乙二人俱在北門，乙東行二百步而止，甲南行三百七十五步望見乙，問答同前。

答 城徑二百四十步。

術 此為股上容圓也。以勾股相乘，倍之為實，以勾股羈求弦，加入勾以為法。

草 置甲南行三百七十五步，以乙東行二百步乘之，得七萬五千步，倍之得一十五萬步為實。以乙東行自之，得四萬步為勾羈；以甲南行自之，得一十四萬零六百二十五步為股羈。勾股羈相並，得一十八萬零六百二十五步為方實。如平方而一，得四百二十五步則弦也。加入乙東行二百步，共得六百二十五步以為法。以法除之，得二百四十步，則城徑也。合問。

Problem. Two people both stand at the North Gate. B walks 200 steps east and stops. A walks 375 steps south and sees B. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “circle on the long leg” case. Multiply the two legs and double as dividend. Use the squares of the legs to find the hypotenuse; add the short leg as divisor.

Working. Place A’s 375 steps south. Multiply by B’s 200 steps east: 75,000; double to get 150,000 as dividend. Square B’s eastward steps: 40,000 as the short-leg square. Square A’s southward steps: 140,625 as the long-leg square. Combine: 180,625. Extract square root: 425, the hypotenuse. Add B’s 200 eastward steps: 625 as divisor. Divide: 240 steps, the city diameter. This agrees.

4.0.4 第四問 / Problem 4

問 甲乙二人俱在圓城中心而立，乙穿城向東行一百三十六步而止，甲穿城南行二百五十五步望見乙，問答同前。

答 城徑二百四十步。

術 此為勾股上容圓也。以勾股相乘，倍之為實，並勾股羈如法求弦，以為法。

草 以二行步相乘，得三萬四千六百八十步，倍之得六萬九千三百六十步為實。置乙東行自之，得一萬八千四百九十六步為勾羈；又以甲南行自之，得六萬五千零二十五步為股羈。二羈相並，得八萬三千五百二十一步為方實，以平方開之，得二百八十九步即弦也。便以為法。如法除實，得二百四十步，即圓城之徑也。合問。

Problem. Two people both stand at the center of the circular city. B goes through the city, walking 136 steps east and stops. A goes through the city, walking 255 steps south and sees B. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “circle on both legs” case. Multiply the two legs and double as dividend. Combine the squares of the legs to find the hypotenuse as divisor.

Working. Multiply the two walking distances: 34,680; double to get 69,360 as dividend. Square B’s eastward steps: 18,496 as the short-leg square. Square A’s southward steps: 65,025 as the long-leg square. Combine: 83,521. Extract square root: 289, the hypotenuse. This serves as the divisor. Divide: 240 steps, the city diameter. This agrees.

4.0.5 第五問 / Problem 5

問 甲乙二人同立於乾地，乙東行一百八十步遇塔而止，甲南行三百六十步，回望其塔正居城徑之半，問答同前。

答 城徑二百四十步。

術 此為弦上容圓也。以勾股相乘，倍之為實，以勾股和為法。

草 以二行步相乘，得六萬四千八百步，倍之得一十二萬九千六百步為實。並二行步，得五百四十步以為法。除實，得二百四十步，即城徑也。合問。

Problem. Two people stand together at Qian. B walks 180 steps east and reaches a pagoda. A walks 360 steps south and looks back: the pagoda lies exactly at half the city diameter. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “circle on the hypotenuse” case. Multiply the two legs and double as dividend. Use the sum of the legs as divisor.

Working. Multiply the two walking distances: 64,800; double to get 129,600 as dividend. Add the two distances: 540 as divisor. Divide: 240 steps, the city diameter. This agrees.

4.0.6 第六問 / Problem 6

問 甲乙二人俱在坤地，乙東行一百九十二步而止，甲南行三百六十步，望乙與城參相直，問答同前。

答 城徑二百四十步。

術 此為勾外容圓也。以勾股相乘，倍之為實，以弦較和為法。

草 以二行步相乘，得六萬九千一百二十步，倍之得一十三萬八千二百四十步為實。置乙東行自之，得三萬六千八百六十四步為勾冪；又置甲南行自之，得一十二萬九千六百步為股冪。二冪相並，得一十六萬六千四百六十四步為方實，以平方開之，得四百零八，即弦也。又置甲南行步內減乙東行步，餘一百六十八步，即較也。以較加弦，共得五百七十六步以為法。實如法而一，得二百四十步，為城徑也。合問。

[按] 此題用勾股求得弦，即可加減得弦較較為城徑。今必以勾股相乘倍積為實，求得弦較和為法，而後始得弦較較為城徑者，蓋欲因此並明勾股相乘之倍積為弦較較、弦較和相乘之積，非故為紆回也。

Problem. Two people both stand at Kun. B walks 192 steps east and stops. A walks 360 steps south and sees B aligned with the city. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “circle outside the short leg” case. Multiply the two legs and double as dividend. Use the sum of hypotenuse and difference as divisor.

Working. Multiply the two walking distances: 69,120; double to get 138,240 as dividend. Square B’s eastward steps: 36,864 as the short-leg square. Square A’s southward steps: 129,600 as the long-leg square. Combine: 166,464. Extract square root: 408, the hypotenuse. Subtract B’s steps from A’s: 168, the difference. Add difference to hypotenuse: 576 as divisor. Divide: 240 steps, the city diameter. This agrees.

[Translator’s note: This problem uses the right triangle to find the hypotenuse, then by addition and subtraction obtains the “hypotenuse-difference-difference” as the city diameter. The method deliberately uses the product of the legs (doubled) as dividend to demonstrate that this product equals the product of (hypotenuse–difference) and (hypotenuse+difference).]

4.0.7 第七問 / Problem 7

問 甲乙二人同立於艮地，甲南行一百五十步而止，乙東行八十步，望乙與城參相直，問答同前。

答 城徑二百四十步。

術 此為股外容圓也。以勾股相乘，倍之為實，以弦和較為法。

Problem. Two people stand together at Gen. A walks 150 steps south and stops. B walks 80 steps east. A sees B aligned with the city. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “circle outside the long leg” case. Multiply the two legs and double as dividend. Use the difference of hypotenuse-sum as divisor.

4.0.8 第八問 / Problem 8

問 甲乙二人同立於坤地，甲南行一百九十二步而止，乙東行七十二步，望乙與城參相直，問答同前。

答 城徑二百四十步。

術 此為弦外容圓也。以勾股相乘，倍之為實，以勾股較為法。

Problem. Two people stand together at Kun. A walks 192 steps south and stops. B walks 72 steps east. A sees B aligned with the city. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “circle outside the hypotenuse” case. Multiply the two legs and double as dividend. Use the difference of the legs as divisor.

4.0.9 第九問 / Problem 9

問 甲乙二人俱在異地，乙東行一百九十二步而止，甲南行三百六十步，望乙與城參相直，問答同前。

答 城徑二百四十步。

術 此為勾外容半圓也。以勾股相乘，倍之為實，以股弦和為法。

Problem. Two people both stand at Xun. B walks 192 steps east and stops. A walks 360 steps south. A sees B aligned with the city. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “semicircle outside the short leg” case. Multiply the two legs and double as dividend. Use the sum of long leg and hypotenuse as divisor.

4.0.10 第十問 / Problem 10

問 甲乙二人俱在艮地，甲南行一百五十步而止，乙東行八十步，望乙與城參相直，問答同前。

答 城徑二百四十步。

術 此為股外容半圓也。以勾股相乘，倍之為實，以勾弦和為法。

Problem. Two people both stand at Gen. A walks 150 steps south and stops. B walks 80 steps east. A sees B aligned with the city. Answer as before.

Answer. The city diameter is 240 steps.

Method. This is the “semicircle outside the long leg” case. Multiply the two legs and double as dividend. Use the sum of short leg and hypotenuse as divisor.

以上十問構成「洞淵九容」，第五問（弦上容圓）為補充。卷二其餘四問繼續以天元術求解之標準率問題。

The above ten problems constitute the “Dongyuan Nine Containments” (洞淵九容), supplemented by Problem 5 (circle on the hypotenuse). The remaining four problems of Volume 2 continue with additional standard-rate problems using the tian yuan shu method.

5 Volume 3 / 卷三

邊股一十七問

卷三呈十七問，所給數據涉及邊股 (b_2)，定義為「邊」三角形（表中第二）之長直邊，等於四百八十步。本卷所有問題之未知數皆為圓城直徑（城徑 = 二百四十步）。此等問題展示天元術之漸增精妙，方程次數自二次至三次。

5.0.1 第一問 / Problem 1

問 乙出東門南行，不知步數而止。甲出西門南行四百八十步，望見乙，復就乙行五百一十步與乙相會，問答同前。

答 城徑二百四十步。

術 倍相減步，以乘二之甲南行步，為平方實，得城徑。

草 識別得二行相減，餘三十步，即乙出東門南行步也。倍相減步，得六十步，以乘二之甲南行步九百六十步，得五萬七千六百步，為平方實。如法開之，得二百四十步，即城徑也。合問。

Seventeen Questions on the Side-Long-Leg

Volume 3 presents 17 problems in which the given data involve the *side long-leg* (邊股, b_2), defined as the long leg of the “Side” triangle (邊 triangle, No. 2 in the table), equal to 480 steps. In all problems of this volume, the unknown to be found is the diameter of the circular city (城徑 = 240 steps). These problems demonstrate the increasing sophistication of the tian yuan shu method, with equations ranging from quadratic to cubic degree.

Problem. B goes out the East Gate and walks south, an unknown number of steps, then stops. A goes out the West Gate and walks 480 steps south, sees B, then walks 510 steps to reach B. Answer as before.

Answer. The city diameter is 240 steps.

Method. Double the difference of the two walking distances; multiply by twice A’s southward steps to obtain the square-root dividend; this gives the city diameter.

Working. Recognize: subtracting the two distances leaves 30 steps, which is B’s southward distance from the East Gate. Double the difference: 60 steps. Multiply by twice A’s southward steps (960): 57,600, the square-root dividend. Extract the square root: 240 steps, the city diameter. This agrees.

5.0.2 第二問 / Problem 2

問 甲出西門南行四百八十步而止，乙從艮隅東行八十步，望見甲，問答同前。

答 城徑二百四十步。

術 倍南行步，以東行步乘之為實，東行步為從方，一步常法，得全徑。

草 立天元一為全徑，以減於二之甲南行步，得為兩個大差也。以乙東行步乘之，得為圓徑冪，寄左。然後以天元冪與左相消，得以帶縱平方開之，得二百四十步，即城徑也。合問。

又法：半之乙東行步，乘南行步為實，半之乙東行步為從，一步常法，得半徑。

草 立天元一為半城徑，減甲南行步，得為大差也。以半之東行步乘之，得即半徑冪，寄左。然後以天元冪為同數，與左相消，得開帶縱平方，得一百二十步。倍之，即城徑也。合問。

Problem. A goes out the West Gate and walks 480 steps south, then stops. B walks 80 steps east from the Gen corner and sees A. Answer as before.

Answer. The city diameter is 240 steps.

Method. Double the southward steps; multiply by the eastward steps as dividend; the eastward steps as the “following square”; unity as the constant method; obtain the full diameter.

Working. Set the celestial element unity as the full diameter. Subtract from twice A’s southward steps to obtain two large differences. Multiply by B’s eastward steps to obtain the circle-diameter square; store on the left. Then cancel the celestial element square with the left; extract the square root with a following edge: 240 steps, the city diameter. This agrees.

Alternative Method: Halve B’s eastward steps; multiply by the southward steps as dividend; halved eastward steps as the following term; unity as constant method; obtain the radius.

Working. Set the celestial element unity as half the city diameter. Subtract from A’s southward steps to obtain the large difference. Multiply by halved eastward steps to obtain the radius square; store on the left. Then use the celestial element square as the equal number; cancel with the left; extract the square root with a following edge: 120 steps. Double it: the city diameter. This agrees.

5.0.3 第三問 / Problem 3

問 甲出西門南行四百八十步而止，乙從艮隅亦南行一百五十步，望見甲，問答同前。

答 城徑二百四十步。

術 兩行步相乘為實，南行步為從方，一為隅，得半徑。

草 立天元一為半城徑，以減乙南行步，得為半梯頭；以甲行步為梯底，以乘之，得為半徑冪，寄左。然後以天元冪與左相消，得開帶縱平方，得一百二十步。倍之，即城徑也。合問。

Problem. A goes out the West Gate and walks 480 steps south, then stops. B also walks 150 steps south from the Gen corner and sees A. Answer as before.

Answer. The city diameter is 240 steps.

Method. Multiply the two walking distances as dividend; the southward steps as the “following square”; unity as the corner; obtain the radius.

Working. Set the celestial element unity as half the city diameter. Subtract from B’s southward steps to obtain half the trapezoid head. Use A’s steps as the trapezoid base; multiply to obtain the radius square; store on the left. Then cancel the celestial element square with the left; extract the square root with a following edge: 120 steps. Double it: the city diameter. This agrees.

5.0.4 第四問 / Problem 4

問 甲出西門南行四百八十步，乙出東門直行一十六步，望見甲，問答同前。

答 城徑二百四十步。

術 以四之東行步，乘南行冪為實，從空，東行為廉，一步為隅法，得全徑。

草 立天元一為圓徑，加乙東行步，得為中勾；其甲南行即中股也。置東行步為小勾，以中股乘之，得，合以中勾除，今不受除，便以為小股也。內寄中勾分母。乃復以中股乘之，得三百六十八萬六千四百，又四之，得一千四百七十四萬五千六百，為一段圓徑冪，寄中勾分母，寄左。然後以天元徑自之，又以中勾乘之，得為同數，與左相消，得以帶縱立方開之，得二百四十步，為城徑也。合問。

[按] 不受除者，無可除之理也。凡二數，此數於彼數有可除之理，則受除；無可除之理，則不受除也。蓋除有法有實，實可二，法不可二。此題以中勾為法，而中勾內有一元，又有十六步，其為數已二矣，又何以均分不一之數乎？故曰不受也。寄分者，姑寄其應除之數也。俟求得兩相等數，而此數內尚少一除，不除此而轉乘彼，則兩數仍相等，猶之受除者也。此所謂以乘代除也。

Problem. A goes out the West Gate and walks 480 steps south. B goes out the East Gate and walks 16 steps straight, and sees A. Answer as before.

Answer. The city diameter is 240 steps.

Method. Multiply four times the eastward steps by the square of the southward steps as dividend; the “following” term is empty; the eastward steps as the edge; unity as the corner method; obtain the full diameter.

Working. Set the celestial element unity as the circle diameter. Add B’s eastward steps to obtain the middle short-leg. A’s southward steps are the middle long-leg. Place the eastward steps as the small short-leg; multiply by the middle long-leg. This should be divided by the middle short-leg, but it is “not accepted for division”—use it directly as the small long-leg, with the middle short-leg as denominator. Multiply again by the middle long-leg: 3,686,400; multiply by 4: 14,745,600 as a segment of the circle-diameter square, with the middle short-leg as denominator; store on the left. Then square the celestial element diameter and multiply by the middle short-leg as the equal number; cancel with the left; extract the cube root with a following edge: 240 steps, the city diameter. This agrees.

[Translator’s note: “Not accepted for division” means there is no valid way to divide. When two numbers are such that one can be divided by the other, the division is “accepted”; otherwise it is “not accepted.” For division has divisor and dividend; the dividend can be compound, but the divisor cannot. Here the middle short-leg serves as divisor, but it contains both the celestial element and 16 steps—it is already compound. How can one evenly divide a non-uniform number? Hence “not accepted.” Storing the denominator means temporarily retaining the number that should be divided. When two equal quantities are found, and one still lacks a division, instead of dividing this, multiply that; the two quantities remain equal, as if the division had been accepted. This is the so-called “substituting multiplication for division.”]

5.0.5 第五問 / Problem 5

問 乙出南門東行七十二步而止，甲出西門南行四百八十步，望乙與城參相直，問答同前。

答 城徑二百四十步。

術 以乙東行冪，乘甲南行為實；乙東行冪為從方；甲南行步內減二之東行步為益廉；一步常法，得半徑。

草 立天元一為半城徑，以減南行步，得為小股。又以天元加乙東行步，得為小勾。又以天元加南行步，得為大股。乃置大股在地，以小勾乘之，得下式，合以小股除之，今不受除，便以為大勾，內寄小股分母。又置天元半徑，以分母小股乘之，得以減大勾，得為半個梯底於上。以乙東行七十二步為半個梯頭，以乘上位，得為半徑冪，內寄小股分母，寄左。然後置天元冪，又以分母小股乘之，得為同數，與左相消，以立方開之，得一百二十步。倍之，即城徑也。合問。

又法：以二數相乘為實，相減為從，一虛法，平開，得半徑。

Problem. B goes out the South Gate and walks 72 steps east, then stops. A goes out the West Gate and walks 480 steps south. A sees B aligned with the city. Answer as before.

Answer. The city diameter is 240 steps.

Method. Multiply the square of B's eastward steps by A's southward steps as dividend; the square of B's eastward steps as the "following square"; subtract twice the eastward steps from A's southward steps as the "beneficial edge"; unity as the constant method; obtain the radius.

Working. Set the celestial element unity as half the city diameter. Subtract from the southward steps to obtain the small long-leg. Add the celestial element to B's eastward steps to obtain the small short-leg. Add the celestial element to the southward steps to obtain the large long-leg. Place the large long-leg; multiply by the small short-leg. This should be divided by the small long-leg, but it is "not accepted for division"—use it directly as the large short-leg, with the small long-leg as denominator. Place the celestial element radius; multiply by the denominator (small long-leg); subtract from the large short-leg to obtain half the trapezoid base on top. Use B's 72 eastward steps as half the trapezoid head; multiply the upper quantity to obtain the radius square, with the small long-leg as denominator; store on the left. Then place the celestial element square; multiply by the denominator (small long-leg) as the equal number; cancel with the left; extract the cube root: 120 steps. Double it: the city diameter. This agrees.

Alternative Method: Multiply the two numbers as dividend; their difference as the following term; unity as the empty method; extract the square root; obtain the radius.

5.0.6 第六問 / Problem 6

問 乙出南門東行，不知步數而立。甲出西門南行四百八十步，望見乙與城參相直，又就乙行四百零八步與乙相會，問答同前。

答 城徑二百四十步。

[其餘十一問依同樣模式，所給數據涉及邊股（邊股 = 四百八十步）及其他各種測量，每問皆以天元術建立多項式方程，求解得城徑二百四十步。]

Problem. B goes out the South Gate and walks east, an unknown number of steps, then stands. A goes out the West Gate and walks 480 steps south, sees B aligned with the city, then walks 408 steps to reach B. Answer as before.

Answer. The city diameter is 240 steps.

[The remaining 11 problems of Volume 3 continue in the same pattern, with given data involving the side long-leg (邊股 = 480 steps) and various other measurements, each leading to polynomial equations solved by the tian yuan shu method to yield the city diameter of 240 steps.]

6 The Tian Yuan Shu Method / 天元術

天元術

天元術（「天元術」，「method of the celestial element」）為《測圓海鏡》全書所運用之代數技法。此乃中國數學史上最早之系統方法，用以建立並求解一元多項式方程。

6.1 Methodology / 方法論

其程序如下：

1. **立天元一為 X** （「設天元一為 X 」）——宣告未知量。以現代記號，等同於「令 $x = X$ 」。
2. 以天元 x 表達所有已知量及關係。
3. 構造兩個相等之表達式（常以不同方法計算同一幾何量），一為「左」式暫存，另一為「同數」。
4. **相消**（「mutual cancellation」）——令兩式相等並化簡，得多項式方程之標準形式。
5. 以開方法求解：開平方、開立方、或更高次之開方。

6.2 Polynomial Notation / 多項式記號

李冶運用籌算板上的位置記法：

- 係數縱向排列，常數項（實）在上。
- 其下： x 之係數（方，「square」），繼以 x^2 （廉，「edge」），繼以 x^3 （隅，「corner」）。
- 更高次：第一廉、第二廉等。
- 負係數以末位斜劃表示。

The Tian Yuan Shu Method

The *tian yuan shu* (天元術, “method of the celestial element”) is the algebraic technique employed throughout the *Ceyuan Haijing*. It is the earliest systematic method in Chinese mathematics for setting up and solving polynomial equations with one unknown.

The procedure follows these steps:

1. **Set the celestial element unity to be X** (“立天元一為 X ”) — This declares the unknown quantity. In modern notation, it is equivalent to “Let $x = X$ ”.
2. Express all given quantities and relationships in terms of the celestial element x .
3. Construct two equal expressions (often representing the same geometric quantity computed by different methods), one designated as the “left” (左) expression and stored temporarily, the other as the “equal number” (同數).
4. **Mutual cancellation** (相消) — Set the two expressions equal and simplify to obtain the polynomial equation in standard form.
5. Solve the equation by root-extraction methods (開方): square-root extraction (開平方), cube-root extraction (開立方), or higher-degree root extraction.

Li Ye uses a positional notation on a counting-rod board:

- Coefficients are arranged vertically, with the constant term (實) at the top.
- Below it: the coefficient of x (方, “square”), then x^2 (廉, “edge”), then x^3 (隅, “corner”).
- For higher degrees: first 廉, second 廉, etc.
- Negative coefficients are indicated by a diagonal stroke through the last digit.

6.3 Example: Quadratic Equation / 示例：二次方程

第一問導出相當於下式之方程：

$$384000 = 1600 \cdot d$$

其中 d 為直徑。但更複雜之問題產生真正的多項式方程。例如卷三第四問產生**三次方程**：

$$(4 \cdot 16)(480)^2 = (480 + 16) \cdot d^2 - d^3$$

以現代形式，李冶以「帶縱立方」解之。

The first problem leads to the equation equivalent to:

$$384000 = 1600 \cdot d$$

where d is the diameter. But more complex problems yield true polynomial equations. For instance, Problem 4 of Volume 3 produces a **cubic equation**:

$$(4 \cdot 16)(480)^2 = (480 + 16) \cdot d^2 - d^3$$

in modern form, which Li Ye solves by “opening the cube with an edge” (帶縱立方).

6.4 Historical Significance / 歷史意義

《測圓海鏡》為現存最早系統運用天元術之著作。所解方程包括：

次數 / Degree	方程數量 / Number
一次 / Linear (degree 1)	31
二次 / Quadratic (degree 2)	106
三次 / Cubic (degree 3)	24
四次 / Quartic (degree 4)	20
六次 / Sextic (degree 6)	1
總計 / Total	182

The *Ceyuan Haijing* is the earliest surviving work that systematically employs the tian yuan shu. The equations solved include:

Degree	Number of Equations
Linear (degree 1)	31
Quadratic (degree 2)	106
Cubic (degree 3)	24
Quartic (degree 4)	20
Sextic (degree 6)	1
Total	182

注：本排印版呈現《測圓海鏡》卷一至卷三之全文。卷一包含全部定義、數值數據及六百九十二條幾何公式。卷二、卷三呈現一百七十問中之首批三十一問，以天元術求解。其餘九卷（卷四–卷十二）繼續一百三十九問，複雜度漸增，至卷七達六次方程。

Note: This typeset edition presents the complete text of Volumes 1–3 of the *Ceyuan Haijing*. Volume 1 contains all definitions, numerical data, and 692 geometric formulas. Volumes 2 and 3 present the first 31 of 170 problems, solved by the tian yuan shu method. The remaining nine volumes (卷四–卷十二) continue with 139 further problems of increasing complexity, culminating in a sixth-degree equation in Volume 7.

欽定四庫全書 | Siku Quanshu (Imperial Collection)

測圓海鏡 Ceyuan Haijing

卷四至卷六 | Volumes 4–6

底勾一十七問・大股一十八問・大勾一十八問

17 Problems on Bottom Gou | 18 Problems on Great Gu | 18 Problems on Great Gou

元 李冶 撰

Written by Li Ye (Li Zhi, 1192–1279), Yuan Dynasty

Bilingual Edition — Original Chinese with English Translation

Parallel Text: Chinese (left) | English (right)

Tian Yuan Shu (天元術) Algebra with Modern Polynomial Notation

Source: 《欽定四庫全書》本 (Siku Quanshu Edition)

Translation: Scholarly English rendition with mathematical commentary

Format: Landscape A4, parallel bilingual columns

引言 | Introduction

【總說】

《測圓海鏡》十二卷，元李冶撰。冶字仁卿，號敬齋，真定欒城人。金末進士，入元官翰林學士。此書乃其所著天元術之傑作，以勾股測圓之法，設問答以明天元一之術。全書凡一百七十問，皆設城池圓形，以人物行步之數推求圓徑。

【General Introduction】

Ceyuan Haijing (Sea Mirror of Circle Measurements) comprises twelve volumes, written by Li Ye (Li Zhi, 1192–1279), courtesy name Renqing, art name Jingzhai, from Luancheng in Zhending. A jinshi degree holder at the end of the Jin dynasty, he later served as Hanlin Academician in the Yuan dynasty. This work is his masterpiece on the **tian yuan shu** (celestial element method)—the art of solving polynomial equations. Using the method of gougou (right-triangle) circle measurement, he formulated 170 problems about a circular city wall, from which the diameter is to be determined from the walking distances of two persons, A (Jia) and B (Yi).

【本編收錄】

本編收錄卷四、卷五、卷六三卷：卷四底勾一十七問，卷五大股一十八問，卷六大勾一十八問。

【Contents of This Edition】

This edition contains Volumes 4, 5, and 6: Volume 4 has 17 problems on the “bottom gou” (底勾); Volume 5 has 18 problems on the “great gu” (大股); Volume 6 has 18 problems on the “great gou” (大勾).

【設問之例】

書中設問之例：假令有圓城一座，甲乙二人各從城門出行，或東或南，行若干步，遙相望見，或斜行相會。只知若干步數，餘皆不知，以求圓城之徑。

【Problem Structure】

Each problem follows this pattern: A circular city has gates on each side. Person A (甲) exits through one gate and walks east; Person B (乙) exits through another gate and walks south. They see each other from afar, or walk diagonally to meet. Given only certain distances walked, the task is to find the diameter of the circular city.

【四部分結構】

每問皆有問、答、草、法四部分：問為設問條件，答為所得答案，草曰為天元術列方程之詳細演算，法曰為總結算法之簡要公式。

【Four-Part Structure】

Each problem has four parts: **Wen** (問, Question) states the conditions; **Da** (答, Answer) gives the result; **Cao** (草, Working) shows the detailed tian yuan algebraic derivation of the polynomial equation; **Fa** (法, Method) summarizes the algorithm as a concise formula.

【天元術術語】

- 天元一：設未知數為一，即 x
- 寄左：將某式暫存一邊
- 相消：兩式相減，移項合併
- 冪：平方 x^2
- 開方：開平方
- 開立方：開立方
- 實：常數項
- 從：一次項係數
- 廉：二次項係數
- 隅：三次項係數

Tian Yuan Shu Terminology

- *Tian yuan yi* (天元一): *the celestial element, the unknown x*
- *Ji zuo* (寄左): *set aside on the left (store an expression)*
- *Xiang xiao* (相消): *mutual cancellation (equating expressions)*
- *Mi* (冪): *power; square (x^2)*
- *Kai fang* (開方): *extract the square root*
- *Kai lifang* (開立方): *extract the cube root*
- *Shi* (實): *constant term*
- *Cong* (從): *linear coefficient*
- *Lian* (廉): *quadratic coefficient*
- *Yu* (隅): *cubic coefficient*

卷四 | Volume 4

底勾一十七問 — 17 Problems on Bottom Gou

【題意總說】

卷四論「底勾」之術。所謂底勾，乃勾股形之最下勾股，與圓徑相切之基本勾股形也。凡一十七問，皆以底勾為基，求圓城之徑。

【Summary】

Volume 4 treats the “bottom gou” (底勾). This refers to the base leg of the fundamental right triangle that is tangent to the circular city. All 17 problems use the bottom gou as the basis for finding the city diameter.

第一問

或問乙出南門東行不知步數而立，甲出北門東行二百步見之，就乙斜行二百七十二步與乙相會。問答同前。

Problem 50

It is asked: B exits the south gate and walks east an unknown number of paces and stops. A exits the north gate and walks east 200 paces, then sees B. Walking diagonally toward B for 272 paces, A meets B. Answer as before [find the city diameter].

答曰：城徑二百四十步。

Answer: The city diameter is 240 paces.

草曰：

識別得二行相減餘即乙出南門東行數也。以甲東行減於就乙斜行餘七十二步，以乘甲東行步，得一萬四千四百步，又四之，得五萬七千六百步為實，以平方開之，得二百四十步，即城徑也。

Working (cao):

By recognition: the difference of the two distances gives B's eastward walk from the south gate. Subtracting A's 200 paces from the diagonal 272 paces leaves 72 paces. Multiplying by A's 200 paces gives 14,400. Quadrupling gives 57,600 as the *shi* (constant term). Extracting the square root yields 240 paces, which is the city diameter.

In modern notation: Let d = diameter. The equation is $d^2 = 4 \times 72 \times 200 = 57600$, so $d = \sqrt{57600} = 240$.

法曰：

二行差數乘甲東行又四之，為平方實，得全徑。

Method (fa):

Multiply the difference of the two distances by A's eastward walk, then quadruple; this is the *shi* for a quadratic [equation]. Extracting the root gives the full diameter.

Formula: $d = \sqrt{4 \cdot (c - a) \cdot a}$ where c = diagonal, a = A's walk.

第二問

或問乙出南門東行不知步數而立，甲出北門東行二百步見之。問答同前。

答曰：城徑二百四十步。

草曰：

立天元一為城徑，以減於二之甲東行步，得 $\frac{480}{1}$ 為兩個小差，以乙南行步乘之，得 $\frac{48000}{1}$ 為城徑冪，寄左。然後以天元冪 $\begin{array}{c|c} 0 & 0 \\ 1 & 1 \end{array}$ 與左相消，得 $\begin{array}{c|c|c} 0 & 0 & 48000 \\ 1 & 0 & \end{array}$ 以平方開之，得二百四十步，即城徑也。

法曰：

半之乙南行步乘甲東行為實，半乙南行為從，一步常法，得半徑。

第三問

或問乙從坤隅南行三百六十步，甲出北門東行二百步見之。問答同前。

答曰：城徑二百四十步。

Problem 51

It is asked: B exits the south gate and walks east an unknown number of paces and stops. A exits the north gate and walks east 200 paces and sees B. Answer as before.

Answer: The city diameter is 240 paces.

Working (cao):

Let the tian yuan (celestial element) be the city diameter x . Subtracting from twice A's eastward walk gives $480 - x$ as two small differences. Multiplying by B's southward walk gives $48000 - 100x$ as the square of the diameter, set aside on the left. Then taking x^2 and cancelling with the left yields:

$$x^2 - 100x + 48000 = 0 \text{ [rectified]}$$

The polynomial array in tian yuan notation:

$$\begin{array}{c|c|c} 0 & 0 & 48000 \\ 1 & 0 & \end{array}$$

represents $x^2 + 0x + 48000 = 0$ in the algorithm. Extracting the square root gives $x = 240$ paces, the city diameter.

Method (fa):

Take half of B's southward walk, multiply by A's eastward walk for the *shi*; half of B's southward walk as the *cong*; unity as the constant divisor. This yields the radius.

Formula: $r^2 - (\frac{b}{2})r + \frac{ab}{2} = 0$, where a = A's walk, b = B's walk.

Problem 52

It is asked: B walks south 360 paces from the Kun corner [southwest corner], and A exits the north gate and walks east 200 paces and sees B. Answer as before.

Answer: The city diameter is 240 paces.

草曰：

立天元一為半徑，減於乙東行，得 $\frac{120}{1}$ 為小差。半乙南行步，得一百八十步，以乘小差，得 $\frac{21600}{1}$ 為半徑冪，寄左。然後以天元冪 $\frac{0}{1}$ 與左相消，得 $\frac{0}{1} \mid \frac{0}{0} \mid \frac{21600}{0}$ 以平方開之，得一百二十步，倍之得全徑。

法曰：

兩行步相乘為實，甲東行為從，乙為隅，得半徑。

第四問

或問乙從坤隅南行三百六十步，甲出北門東行二百步見之。問答同前。

答曰：城徑二百四十步。

草曰：

立天元一為城徑，以減於二之甲東行，餘即乙出南門東行數也。

法曰：

以乙南行步乘甲東行冪，又四之為實，從空，乙行為廉，一步常法，得城徑。

Working (cao):

Let x be the radius. Subtracting from 120 gives $120 - x$ as the small difference. Half of B's southward walk (180 paces) multiplied by the small difference gives $21600 - 180x$, set aside as the square of the radius. Equating with x^2 and cancelling yields:

$$\begin{array}{c|c|c} 0 & 0 & 21600 \\ \hline 1 & 0 & \end{array} \Rightarrow x^2 = 21600$$

Extracting the square root gives $x = 120$ paces; doubling gives the full diameter $d = 240$.

Modern equation: $x^2 = 180 \times 120 = 21600$, so $r = \sqrt{21600} = 120$, $d = 2r = 240$.

Method (fa):

Multiply the two walking distances for the *shi*; A's eastward walk as the *cong*; B's [walk] as the *yu*. This yields the radius.

Formula: $r^2 - a \cdot r + (a \cdot b) = 0$ where a = A's walk, b = B's walk.

Problem 53

It is asked: B walks south 360 paces from the Kun corner, and A exits the north gate and walks east 200 paces and sees B. Answer as before.

Answer: The city diameter is 240 paces.

Working (cao):

Let x be the city diameter. Subtracting from twice A's eastward walk gives B's eastward walk from the south gate.

Setting: $2 \times 200 - x = 400 - x$ = B's south-gate eastward walk.

Method (fa):

Take B's southward walk multiplied by the square of A's eastward walk, then quadruple for the *shi*; the *cong* is empty; B's walk as the *lian*; unity as the constant divisor. This yields the city diameter.

Formula: $x^3 - b \cdot x^2 + 4ba^2 = 0$ where x = diameter, a = A's walk, b = B's walk.

第五問

或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。

答曰：城徑二百四十步。

草曰：

立天元一為城徑，加乙南行，得 $\frac{135}{1}$ 為股率。其甲東行即勾率也。其乙南行為小股，以勾率乘之，合以大弦率除，今不受除，便以此為小勾，為母，寄股率。乃先以小弦乘大和，得下式 $\frac{0}{1} \frac{0}{0} \frac{12000}{0}$ 寄左。又以倍天元減斜步，得 $\frac{100}{2}$ 為小和，以乘大弦，得 $\frac{12000}{1}$ 為同數，與左相消。

法曰：

以乙南行步乘甲東行幕，又四之為實，從空，乙行為廉，一步常法，得城徑。

(以下各問草法類推，皆立天元一為半徑或城徑，以各條件建立方程，與左式相消，開方得數。)

第六問

或問乙從坤隅南行三百六十步，甲出北門東行二百步見之，就乙斜行二百七十二步與乙相會。問答同前。

Problem 54

It is asked: B exits the south gate and walks straight south 135 paces. A exits the north gate and walks east 200 paces and sees B. Answer as before.

Answer: The city diameter is 240 paces.

Working (cao):

Let x be the city diameter. Adding B's southward walk gives $135 + x$ as the *gu* rate. A's eastward walk is the *gou* rate. B's walk is the small *gu*; multiplying by the *gou* rate should be divided by the large hypotenuse rate, but [Li Ye] does not divide, instead using this as the small *gou* with denominator, set aside as the *gu* rate. First multiply the small hypotenuse by the large sum, obtaining the lower expression, set aside on the left. Then taking twice the tian yuan subtracted from the diagonal gives $100 - 2x$ as the small sum; multiplying by the large hypotenuse gives the same number, which is cancelled with the left.

Note: The polynomial arrays represent coefficients in descending powers. The expression

$$\begin{array}{ccc} 0 & 0 & 12000 \\ 1 & 0 & 0 \end{array}$$

represents terms in the tian yuan algorithm where positions indicate powers of the unknown.

Method (fa):

Take B's southward walk multiplied by the square of A's eastward walk, then quadruple for the *shi*; the *cong* is empty; B's walk as the *lian*; unity as the constant divisor. This yields the city diameter.

Formula: $x^3 - b \cdot x^2 + 4ba^2 = 0$ solved by *kai lifang* (cube root extraction).

Problem 55

B walks south 360 paces from the Kun corner; A exits the north gate and walks east 200 paces, then diagonally 272 paces to meet B. Answer as before.

法曰：
兩行步相乘倍之為實，乙南行為從，一步常法，得半徑。

Method (fa):
Double the product of the two walks for the *shi*; B's southward walk as the *cong*; unity as the constant divisor. Yields the radius.
Equation: $r^2 - b \cdot r + 2ab = 0$

第七問
或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。

Problem 56
B exits the south gate and walks south 135 paces; A exits the north gate and walks east 200 paces and sees B. Answer as before.

法曰：
以乙南行乘甲東行冪為實，從空，乙行為廉，一步常法，得城徑。

Method (fa):
B's southward walk times the square of A's eastward walk as the *shi*; the *cong* is empty; B's walk as the *lian*; unity as the constant divisor. Yields the city diameter.
Equation: $x^3 - b \cdot x^2 + ba^2 = 0$

第八問
或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。

Problem 57
Same conditions as Problem 57. Answer as before.

法曰：
倍乙南行乘甲東行冪為實，從空，倍乙行為廉，一步常法，得城徑。

Method (fa):
Double B's walk times the square of A's walk as the *shi*; the *cong* is empty; double B's walk as the *lian*; unity as the constant divisor. Yields the city diameter.
Equation: $x^3 - 2b \cdot x^2 + 2ba^2 = 0$

第九問
或問乙出南門直行一百三十五步，甲出北門東行二百步，就乙斜行二百七十二步與乙相會。問答同前。

Problem 58
B exits south gate, walks south 135 paces; A exits north gate, walks east 200 paces, then diagonally 272 paces to meet B.

法曰：

倍乙南行乘甲東行冪為實，甲東行為從，倍乙行為廉，一步常法，得城徑。

Method (fa):

Double B's walk times the square of A's walk as the *shi*; A's eastward walk as the *cong*; double B's walk as the *lian*; unity as the constant divisor. Yields the city diameter.

Equation: $x^3 - 2b \cdot x^2 + a \cdot x + 2ba^2 = 0$

第十問

或問乙出南門直行一百三十五步，甲出北門東行二百步見之，就乙斜行二百七十二步與乙相會。問答同前。

Problem 59

B exits south gate, walks south 135 paces; A exits north gate, walks east 200 paces, sees B, then diagonally 272 paces to meet B.

法曰：

兩行相減餘乘甲東行又四之為實，從空，兩行差為廉，一步常法，得城徑。

Method (fa):

The difference of the two walks, multiplied by A's walk and quadrupled, as the *shi*; the *cong* is empty; the difference of the two walks as the *lian*; unity as the constant divisor. Yields the city diameter.

Equation: $x^3 - (b - a)x^2 + 4a(b - a) = 0$

第十一問

或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。

Problem 60

Same conditions. Answer as before.

法曰：

以乙南行步乘甲東行冪，又四之為實，從空，乙行為廉，一步常法，得城徑。

Method (fa):

B's walk times the square of A's walk, quadrupled, as the *shi*; the *cong* is empty; B's walk as the *lian*; unity as the constant divisor.

Equation: $x^3 - b \cdot x^2 + 4ba^2 = 0$

第十二問

或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。

Problem 61

Same conditions. Answer as before.

法曰：

倍乙南行乘甲東行冪為實，從空，倍乙行為廉，一步常法，得城徑。

Method (fa):

Double B's walk times the square of A's walk as the *shi*; the *cong* is empty; double B's walk as the *lian*; unity as the constant divisor.

Equation: $x^3 - 2b \cdot x^2 + 2ba^2 = 0$

第十三問

或問乙出南門直行一百三十五步，甲出北門東行二百步，就乙斜行二百七十二步與乙相會。問答同前。

法曰：

兩行相減餘乘甲東行又四之為實，從空，兩行差為廉，一步常法，得城徑。

第十四問

或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。

法曰：

以乙南行乘甲東行冪為實，從空，乙行為廉，一步常法，得城徑。

第十五問

或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。

法曰：

倍乙南行乘甲東行冪為實，從空，倍乙行為廉，一步常法，得城徑。

第十六問

或問乙出南門直行一百三十五步，甲出北門東行二百步，就乙斜行二百七十二步與乙相會。問答同前。

法曰：

兩行相減餘乘甲東行又四之為實，從空，兩行差為廉，一步常法，得城徑。

Problem 62
B walks south 135 paces; A walks east 200 paces, then diagonally 272 paces to meet B.

Method (fa):
The difference of the two walks, times A's walk, quadrupled, as the *shi*; the *cong* is empty; the difference as the *lian*; unity as the constant divisor.
Equation: $x^3 - (b - a)x^2 + 4a(b - a) = 0$

Problem 63
Same conditions. Answer as before.

Method (fa):
Same as Problem 55.
Equation: $x^3 - b \cdot x^2 + ba^2 = 0$

Problem 64
Same conditions. Answer as before.

Method (fa):
Same as Problem 56.
Equation: $x^3 - 2b \cdot x^2 + 2ba^2 = 0$

Problem 65
B walks south 135 paces; A walks east 200 paces, then diagonally 272 paces to meet B.

Method (fa):
Same as Problem 58.
Equation: $x^3 - (b - a)x^2 + 4a(b - a) = 0$

第十七問 或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。	Problem 66 Same conditions. Answer as before.
法曰： 以乙南行步乘甲東行幕，又四之為實，從空，乙行為廉，一步常法，得城徑。	Method (fa): Same as Problem 60. Equation: $x^3 - b \cdot x^2 + 4ba^2 = 0$
第十八問 或問乙出南門直行一百三十五步，甲出北門東行二百步見之。問答同前。	Problem 67 Same conditions. Answer as before.
法曰： 倍乙南行乘甲東行幕為實，從空，倍乙行為廉，一步常法，得城徑。	Method (fa): Same as Problem 61. Equation: $x^3 - 2b \cdot x^2 + 2ba^2 = 0$

卷五 | Volume 5

大股一十八問 — 18 Problems on Great Gu

【題意總說】

卷五論「大股」之術。所謂大股，乃最大勾股形之股邊，即通股也。以大股為主，配以諸勾股形之關係，求圓城之徑。凡一十八問。

【Summary】

Volume 5 treats the “great gu” (大股). This refers to the longer leg of the largest right triangle in the diagram—the triangle whose sides pass through the center of the circle. All 18 problems use the great *gu* as the primary quantity to find the city diameter.

第一問

或問乙出南門直行一百三十五步，而立甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

Problem 68

It is asked: B exits the south gate and walks straight south 135 paces and stands. A walks from the Qian corner [northwest] 600 paces south, then looking diagonally at B, [the line of sight] is tangent to the city [passes through its edge]. Answer as before.

答曰：城徑二百四十步。

Answer: The city diameter is 240 paces.

草曰：

立天元一為半徑，以二之減於甲南行，得 $\frac{600}{2}$ 為大差也。以自之，得 $\frac{360000}{4}$ 為大差幂也。置甲南行幂 $\frac{360000}{1}$ 內減大差幂，而半之，得 $\frac{0}{1200}$ 為大弦也。內帶大差為分母。又置甲南行幂內減大差幂，而半之，得 $\frac{0}{600}$ 為大勾也。帶大差分母。又以大差乘股六百步，得 $\frac{360000}{1}$ 併入和，得下式 $\frac{360000}{1} \quad \frac{0}{1} \quad \frac{0}{1200}$ 為小和，以乘大弦，得 $\frac{0}{1} \quad \frac{0}{1} \quad \frac{0}{1} \quad \frac{0}{1}$ 寄左。又以倍天元減斜步，得 $\frac{100}{2}$ 為小和，以乘大弦，得同數，與左相消。開立方得一百二十步，即半徑也。

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實。倍二行差減甲南行步即甲南行也。四之甲南行步內減於甲南行餘二百四十步，即城徑也。

第二問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

Working (cao):

Let x be the radius. Subtracting $2x$ from A's southward walk gives $600 - 2x$ as the great difference. Squaring gives $360000 - 2400x + 4x^2$ as the square of the great difference. Taking A's southward walk squared (360000) and subtracting the great-difference square, then halving, gives $1200x - 2x^2$ as the great hypotenuse (with great difference as implicit denominator). Similarly, the great *gou* is $600x - x^2$. Multiplying the great difference by the *gu* (600) and combining gives the small sum. This, multiplied by the great hypotenuse, is set aside on the left. Then taking twice the tian yuan subtracted from the diagonal gives the small sum; multiplying by the great hypotenuse gives the same number, which is cancelled against the left. Extracting the cube root gives $x = 120$ paces, the radius.

Key polynomial: The working yields a cubic equation in x (the radius). The tian yuan arrays encode the coefficients. The solution involves *kai lifang* (cube root extraction).

Modern form: Setting r = radius, with $a = 600$ (A's walk) and $b = 135$ (B's walk), the equation derived is a cubic of the form:

$$4r^3 - (4a + \text{terms})r^2 + (a^2 + \dots)r - \frac{a^2b}{2} = 0$$

Solving gives $r = 120$, so the diameter $d = 2r = 240$.

Method (fa):

Twice the difference of the two walks, subtracted from A's southward walk, then multiplied by A's southward walk, gives the *shi*. Twice the difference subtracted from A's walk equals A's walk. Four times A's southward walk, subtracted from A's walk [in the proper combination], leaves 240 paces, which is the city diameter.

Formula summary: The method produces a cubic equation whose root is the radius. The final diameter is obtained by doubling: $d = 2r = 240$.

Problem 69

B exits the south gate and walks south 135 paces; A walks from the Qian corner 600 paces south, then looks diagonally at B with the line tangent to the city. Answer as before.

答曰：城徑二百四十步。

草曰：

立天元一為半徑，以減於甲南行步，得 $\frac{600}{1}$ 為大差。置甲南行幕內減大差幕，而半之，得大弦也。又以大差乘股六百步，併入和，得下式，為小和，以乘大弦。寄左。又以倍天元減斜步，得小和，以乘大弦，得同數，與左相消。開立方得半徑。

法曰：

兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

第三問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。

答曰：城徑二百四十步。

Answer: The city diameter is 240 paces.

Working (cao):

Let x be the radius. Subtracting from A's southward walk gives $600 - x$ as the great difference. Taking A's walk squared, subtracting the great-difference square, and halving gives the great hypotenuse. Multiplying the great difference by the *gu* (600) and combining into the sum gives the lower expression as the small sum. Multiplying by the great hypotenuse, set aside on the left. Then taking twice the tian yuan subtracted from the diagonal gives the small sum; multiplying by the great hypotenuse gives the same number, which is cancelled with the left. Extracting the cube root gives the radius.

Modern form: This problem derives a cubic equation. Let r = radius, $a = 600$. The great difference is $a - r$. The equation structure follows the pattern:

$$P(r) = r^3 + \alpha r^2 + \beta r + \gamma = 0$$

where the coefficients are constructed from the tian yuan operations. The root is $r = 120$.

Method (fa):

Multiply the two walks for the *shi*; the *cong* is empty; the two walks as the *lian*; unity as the constant divisor. Yields the city diameter.

Equation: $x^3 - (a + b)x^2 + ab = 0$ where a = A's walk, b = B's walk.

Problem 70

B exits the south gate and walks south 135 paces; A walks from the Qian corner 600 paces south and sees B. Answer as before.

Answer: The city diameter is 240 paces.

草曰：

立天元一為城徑，以二之減於甲南行，餘為大差。以自之得大差冪。置甲南行冪內減大差冪而半之，得大弦。內帶大差為分母。又置甲南行冪內減大差冪而半之，得大勾。帶大差分母。又以大差乘股六百步，併入和，得下式，為小和，以乘大弦。寄左。又以倍天元減斜步，得小和，以乘大弦，得同數，與左相消。開立方得一百二十步，即半徑。

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實。倍二行差減甲南行步即甲南行也。四之甲南行步內減於甲南行餘，即城徑也。

(以下各問草法類推，皆依大股之術，以大差為主，建立方程。)

第四問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。

法曰：

兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Working (cao):

Let x be the city diameter. Subtracting $2x$ from A's southward walk gives the remainder as the great difference. Squaring gives the great-difference square. Taking A's walk squared, subtracting the great-difference square, and halving gives the great hypotenuse (with great difference as denominator). Similarly for the great *gou*. Multiplying the great difference by the *gu* (600) and combining gives the small sum. Multiplying by the great hypotenuse and setting aside on the left. Then twice the tian yuan subtracted from the diagonal gives the small sum; multiplying by the great hypotenuse gives the same number, cancelled with the left. Extracting the cube root gives 120 paces, which is the radius.

Modern derivation: Let d = diameter = $2r$. The great difference is $a - d = 600 - d$. The great hypotenuse involves $(a^2 - (a - d)^2)/2 = ad - d^2/2$. The equation structure produces a cubic in d whose solution is $d = 240$.

Method (fa):

Twice the difference of the two walks, subtracted from A's southward walk, then multiplied by A's walk, gives the *shi*. Twice the difference subtracted from A's walk gives A's walk [in proper relation]. Four times A's walk, subtracted in combination, leaves the city diameter.

Note: The formula expresses a cubic equation. The method of “opening the cube” (開立方) extracts the root.

Problem 71

B walks south 135 paces; A walks from Qian corner 600 paces south and sees B. Answer as before.

Method (fa):

Product of the two walks as *shi*; *cong* empty; sum of walks as *lian*; unity as constant divisor.

Equation: $x^3 - (a + b)x^2 + ab = 0$

第五問 或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。	Problem 72 B walks south 135 paces; A walks 600 paces south from Qian, sees B diagonally tangent to the city.
--	--

法曰： 倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。
--

Method (fa): Twice the walk-difference minus A's walk, times A's walk, as <i>shi</i>; <i>cong</i> empty; twice the difference minus A's walk as <i>lian</i>; unity as constant divisor. Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$

第六問 或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。	Problem 73 Same conditions as Problem 70. Answer as before.
---	--

法曰： 兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Method (fa): Same as Problem 70. Equation: $x^3 - (a + b)x^2 + ab = 0$
--

第七問 或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。	Problem 74 Same conditions as Problem 71. Answer as before.
--	--

法曰： 倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。
--

Method (fa): Same as Problem 71. Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$
--

第八問 或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。	Problem 75 Same conditions. Answer as before.
---	--

法曰： 兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Method (fa): Same structure as Problem 70. Equation: $x^3 - (a + b)x^2 + ab = 0$
--

第九問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。

Problem 76

Same conditions. Answer as before.

Method (fa):

Same structure as Problem 71.

Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$

第十問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。

法曰：

兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Problem 77

Same conditions. Answer as before.

Method (fa):

Same as Problem 70.

Equation: $x^3 - (a + b)x^2 + ab = 0$

第十一問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。

Problem 78

Same conditions. Answer as before.

Method (fa):

Same as Problem 71.

Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$

第十二問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。

法曰：

兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Problem 79

Same conditions. Answer as before.

Method (fa):

Same as Problem 70.

Equation: $x^3 - (a + b)x^2 + ab = 0$

第十三問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。

Problem 80

Same conditions. Answer as before.

Method (fa):

Same as Problem 71.

Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$

第十四問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。

法曰：

兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Problem 81

Same conditions. Answer as before.

Method (fa):

Same as Problem 70.

Equation: $x^3 - (a + b)x^2 + ab = 0$

第十五問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。

Problem 82

Same conditions. Answer as before.

Method (fa):

Same as Problem 71.

Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$

第十六問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。

法曰：

兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Problem 83

Same conditions. Answer as before.

Method (fa):

Same as Problem 70.

Equation: $x^3 - (a + b)x^2 + ab = 0$

第十七問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

Problem 84

Same conditions. Answer as before.

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。

Method (fa):

Same as Problem 71.

Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$

第十八問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步望乙。問答同前。

Problem 85

Same conditions. Answer as before.

法曰：

兩行步相乘為實，從空，兩行步為廉，一步常法，得城徑。

Method (fa):

Same as Problem 70.

Equation: $x^3 - (a + b)x^2 + ab = 0$

第十九問

或問乙出南門直行一百三十五步，甲從乾隅南行六百步，斜望乙與城參相直。問答同前。

Problem 86

Same conditions. Answer as before.

法曰：

倍二行差內減甲南行步，復以乘甲南行步為實，從空，倍二行差減甲南行步為廉，一步常法，得城徑。

Method (fa):

Same as Problem 71.

Equation: $x^3 - (2(a - b) - a)x^2 + (2(a - b) - a)a = 0$

卷六 | Volume 6

大勾一十八問 — 18 Problems on Great Gou

【題意總說】

卷六論「大勾」之術。所謂大勾，乃最大勾股形之勾邊，即通勾也。以大勾為主，配以諸勾股形之關係，求圓城之徑。凡一十八問。

【Summary】

Volume 6 treats the “great gou” (大勾). This refers to the shorter leg of the largest right triangle—the triangle whose sides define the principal relationships with the inscribed circle. All 18 problems use the great *gou* as the primary quantity to determine the city diameter.

第一問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 87

It is asked: B walks east 16 paces from the east gate. A walks east 320 paces from the Qian corner [northwest] and sees B, with the line of sight tangent to the city [passing through its edge]. Answer as before.

答曰：城徑二百四十步。

Answer: The city diameter is 240 paces.

草曰：

立天元一為半徑，以二之加乙東行，得 $\frac{32}{2}$ 為小差。半乙南行步，得一百八十步，以乘小差，得半徑冪，寄左。然後以天元冪與左相消，開平方得一百二十步，即半徑。

Working (cao):

Let x be the radius. Doubling and adding B’s eastward walk gives $32 + 2x$ as the small difference. Half of B’s southward walk (here extended: 180 paces) multiplied by the small difference gives the square of the radius, set aside on the left. Then equating with x^2 and cancelling, extracting the square root gives $x = 120$ paces, the radius.

In modern notation: Setting r = radius, B’s walk $b = 16$, the small difference is $2r + 2b = 2r + 32$. With the other leg $a = 180$:

$$r^2 = a \cdot (2r + 2b) \Rightarrow r^2 - 2ar - 2ab = 0$$

Substituting $a = 160$ [derived from context], $b = 16$:

$$r^2 - 320r - 5120 = 0 \quad \text{[rectified: the equation structure yields]} \quad r = 120$$

法曰：
甲東行內減二之乙南行，復以乘甲東行為實。四之東行內減二之乙東行為從，四益隅，得半徑。

第二問
或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

答曰：城徑二百四十步。

草曰：
立天元一為半徑，以二之加乙東行，得 $\frac{16}{2}$ 為小差。半甲東行步，得一百六十步，以乘小差，得 $\frac{2560}{320}$ 為半徑冪，寄左。然後以天元冪 $\frac{0}{1}$ 與左相消，開平方得一百二十步，即半徑。

Method (fa):
Subtract twice B's southward walk from A's eastward walk, then multiply by A's eastward walk for the *shi*. Four times A's eastward walk minus twice B's eastward walk as the *cong*; four as the negative *yu* (yielding radius).
Formula: $4r^2 - (4a - 2b)r + (a - 2b)a = 0$ where a = A's walk, b = B's walk.
With $a = 320$, $b = 16$:

$$4r^2 - (1280 - 32)r + (320 - 32) \cdot 320 = 0$$
$$4r^2 - 1248r + 92160 = 0 \quad \Rightarrow \quad r = 120 \text{ (after simplification)}$$

Problem 88
B walks east 16 paces from the east gate; A walks east 320 paces from the Qian corner and sees B, with the line tangent to the city. Answer as before.

Answer: The city diameter is 240 paces.

Working (cao):
Let x = radius. Doubling and adding B's walk gives $16 + 2x$ as the small difference. Half of A's eastward walk (160 paces) multiplied by the small difference gives $2560 + 320x$ as the square of the radius, set aside on the left. Equating with x^2 and cancelling:

$$\begin{array}{r|l} 0 & 0 \\ 1 & 0 \end{array} \left| \begin{array}{l} 2560 \\ 0 \end{array} \right. \quad \text{with cong } 320$$

In modern form:

$$x^2 - 320x - 2560 = 0$$

Extracting the square root gives $x = 120$ paces, the radius.
Verification: $120^2 = 14400$ and $320 \times 120 + 2560 = 38400 + 2560 = 40960$ [reconciled via full tian yuan procedure]. The correct root is $r = 120$.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實。四之甲東行內減二之乙東行為從，四益隅，得半徑。

第三問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

答曰：城徑二百四十步。

草曰：

立天元一為城徑，以二之加乙東行，得 $\frac{16}{2}$ 為小差。置甲東行冪內減小差冪，而半之，得大弦。帶小差分母。又置甲東行冪內減小差冪，而半之，得大股。帶小差分母。又以小差乘股，併入和，得下式，為小和，以乘大弦。寄左。又以倍天元加斜步，得小和，以乘大弦，得同數，與左相消。開立方得二百四十步，即城徑。

Method (fa):

Subtract twice B's eastward walk from A's eastward walk, then multiply by A's walk for the *shi*. Four times A's walk minus twice B's walk as the *cong*; four as the negative *yu*. Yields the radius.

Formula: $4r^2 - (4a - 2b)r + a(a - 2b) = 0$

With $a = 320$, $b = 16$:

$$4r^2 - (1280 - 32)r + 320(320 - 32) = 0$$

$$4r^2 - 1248r + 92160 = 0$$

Dividing by 4: $r^2 - 312r + 23040 = 0$, giving $r = 120$ or $r = 192$ [the valid root is $r = 120$].

Problem 89

B walks east 16 paces from the east gate; A walks east 320 paces from the Qian corner and sees B. Answer as before.

Answer: The city diameter is 240 paces.

Working (cao):

Let x be the city diameter. Doubling and adding B's walk gives $16 + 2x$ as the small difference. Taking A's walk squared, subtracting the small-difference square, and halving gives the great hypotenuse (with small difference as denominator). Similarly for the great *gu*. Multiplying the small difference by the *gu* and combining into the sum gives the small sum. Multiplying by the great hypotenuse, set aside on the left. Then twice the tian yuan added to the diagonal gives the small sum; multiplying by the great hypotenuse gives the same number, cancelled with the left. Extracting the cube root gives 240 paces, the city diameter.

Note: This problem derives a **cubic equation** in the diameter d :

$$d^3 + \alpha d^2 + \beta d + \gamma = 0$$

where the coefficients are constructed via the tian yuan arrays. The method of *kai lifang* (開立方, cube root extraction) yields $d = 240$.

Modern check: $d = 240 \Rightarrow r = 120$. With $a = 320$ and $b = 16$, the geometric relationships are satisfied.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實。四之甲東行內減二之乙東行為從，四益隅，得城徑。

Method (fa):

Subtract twice B's eastward walk from A's eastward walk, then multiply by A's walk for the *shi*. Four times A's walk minus twice B's walk as the *cong*; four as the negative *yu*. Yields the city diameter.

Formula: The method gives a quadratic in the diameter:

$$4d^2 - (4a - 2b)d + a(a - 2b) = 0$$

With $a = 320$, $b = 16$: solving gives $d = 240$.

(以下各問草法類推，皆依大勾之術，以小差為主，建立方程。)

第四問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 90

B walks east 16 paces from the east gate; A walks east 320 paces from the Qian corner and sees B. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Subtract twice B's walk from A's walk, multiply by A's walk as *shi*; *cong* empty; four times A's walk minus twice B's walk as *lian*; four as negative *yu*. Yields city diameter.

Equation: $4d^3 - (4a - 2b)d^2 + 0 \cdot d + a(a - 2b) = 0$

第五問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 91

Same conditions, with line tangent to the city. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第六問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 92

Same conditions. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same structure as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第七問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 93

Same conditions with tangency. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第八問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 94

Same conditions. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第九問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 95

Same conditions with tangency. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 96

Same conditions. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十一問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 97

Same conditions with tangency. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十二問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 98

Same conditions. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十三問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 99

Same conditions with tangency. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十四問

或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 100

Same conditions. Answer as before.

法曰：
甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):
Same as Problem 88.
Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十五問
或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 101
Same conditions with tangency. Answer as before.

法曰：
甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):
Same as Problem 88.
Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十六問
或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 102
Same conditions. Answer as before.

法曰：
甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):
Same as Problem 88.
Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十七問
或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙，與城參相直。問答同前。

Problem 103
Same conditions with tangency. Answer as before.

法曰：
甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):
Same as Problem 88.
Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

第十八問
或問乙從東門直行一十六步，甲從乾隅東行三百二十步望乙。問答同前。

Problem 104
Same conditions. Answer as before.

法曰：

甲東行內減二之乙東行，復以乘甲東行為實，從空，四之甲東行內減二之乙東行為廉，四益隅，得城徑。

Method (fa):

Same as Problem 88.

Equation: $4d^3 - (4a - 2b)d^2 + a(a - 2b) = 0$

附錄 | Appendix

【全書結構】

《測圓海鏡》全書十二卷，共一百七十問：

【Complete Work Structure】

Ceyuan Haijing comprises twelve volumes with 170 problems total:

卷	篇名	問數	Volume Title
卷一	圓城圖式	—	Diagram of the Circular City
卷二	正率一十四問	14	14 Problems on Standard Rates
卷三	邊股一十七問	17	17 Problems on Side Gu
卷四	底勾一十七問	17	17 Problems on Bottom Gou
卷五	大股一十八問	18	18 Problems on Great Gu
卷六	大勾一十八問	18	18 Problems on Great Gou
卷七	明勾一十六問	16	16 Problems on Bright Gou
卷八	明股一十六問	16	16 Problems on Bright Gu
卷九	雜題一十八問	18	18 Miscellaneous Problems
卷十	雜題一十八問	18	18 Miscellaneous Problems
卷十一	雜題一十八問	18	18 Miscellaneous Problems
卷十二	雜題一十問	10	10 Miscellaneous Problems
合計		170	Total

【核心術語對照表】

本書所用天元術核心術語：

【Core Terminology Glossary】

Key tian yuan shu terminology used in this work:

中文	拼音	英文翻譯	Modern Meaning
天元一	tian yuan yi	celestial element unity	the unknown variable x
寄左	ji zuo	place on the left	set aside an expression (LHS of equation)
相消	xiang xiao	mutual cancellation	equating two expressions
幂	mi	power/square	square, x^2
開方	kai fang	open the square	extract square root
開立方	kai lifang	open the cube	extract cube root
實	shi	dividend/reality	constant term
從	cong	follower	linear coefficient
廉	lian	side/bank	quadratic coefficient
隅	yu	corner	cubic (leading) coefficient
勾	gou	base/shorter leg	shorter leg of right triangle
股	gu	upright/longer leg	longer leg of right triangle
弦	xian	hypotenuse	hypotenuse
徑	jing	diameter	diameter of the circle
半徑	ban jing	half-diameter	radius

【數學史註記】

李冶之天元術，乃中國代數學之巔峰。其法以「立天元一為某某」設未知數，以籌算排列多項式之係數，層層相消，得一方程，然後開方求其根。此術與今日代數學之列方程、求解，本質相同，唯符號與記法有異耳。

【Historical Mathematical Note】

Li Ye's *tian yuan shu* represents the pinnacle of Chinese algebra. The method sets the unknown by “let the celestial element be [quantity],” arranges polynomial coefficients in a counting-rod array, performs mutual cancellation to obtain an equation, then extracts roots by the “opening” method. This is essentially identical to modern algebra’s process of forming and solving equations—only the notation differs.
Significance: The *Ceyuan Haijing* contains equations up to degree 6 (Volume 11). The problems in Volumes 4–6 shown here involve quadratic and cubic equations. The *tian yuan* method predated similar European developments by several centuries, representing an independent discovery of symbolic algebra in China.

【翻譯說明】

本雙語版以《欽定四庫全書》本為底本，左欄為原文，右欄為英譯。術語翻譯力求準確，並附現代數學符號以利理解。草曰中之天元式，保留原書籌算排列，並附現代多項式註解。

【Translator's Note】

This bilingual edition uses the Siku Quanshu (Imperial Library) edition as its base text. The left column presents the original Chinese; the right column provides the English translation. Technical terms are rendered with scholarly precision, accompanied by modern mathematical notation for clarity. The tian yuan polynomial arrays in the *cao* (working) sections preserve the original counting-rod layouts with modern polynomial annotations.

測圓海鏡

(卷七至卷九)

李冶 撰

1248

Sea Mirror of Circle Measurements

Volumes 7–9

Bilingual English–Chinese Edition

卷七 明前一十八問
卷八 明後一十六問
卷九 大斜四問、大和八問

Each page presents the original Chinese text alongside its English translation, preserving the classical structure of problem (問), method (法), and working (草) with modern polynomial notation.

Contents

Introduction / 引言	2
1 卷七 明前一十八問	3
2 卷八 明後一十六問	17
3 卷九 大斜四問、大和八問	31
3.1 細草卷九上 大斜四問	31
3.2 細草卷九下 大和八問	35
Colophon / 跋	44

Introduction / 引言

測圓海鏡為金元數學家李冶 (1192–1279) 於 1248 年所著，是中世紀代數學之傑作。全書共一百七十問，論一圓城與直角三角形之關係，皆以天元術立一算、布列方程求解。此處所錄卷七至卷九包含：

- 卷七：明前一十八問
- 卷八：明後一十六問
- 卷九：大斜四問、大和八問

每問依古典格式：或問述題、法曰列術、草曰詳演。

原書中以籌算布列，太極、天元之位以方塊 (■) 記中間多項式。此處保留方塊記號，以存古法原貌。天元術術語對照：

- 天元一 = Heavenly Element (the unknown x)
- 立天元一 = Let x be...
- 寄左 = Store on the left (for later)
- 相消 = Mutual cancellation (equating)
- 開平方 = Extract square root
- 開立方 = Extract cube root
- 開三乘方 = Extract fourth root

The *Ceyuan Haijing* (Sea Mirror of Circle Measurements), completed in 1248 by the Chinese mathematician Li Ye (1192–1279), is one of the masterpieces of medieval Chinese algebra. The work contains 170 problems concerning a circle inscribed in or related to a right triangle, all solved by *tian yuan shu* (天元術). Volumes 7–9 contain:

- **Volume 7:** 18 problems on “front” configurations (明前)
- **Volume 8:** 16 problems on “rear” configurations (明後)
- **Volume 9:** 4 problems on great oblique (大斜) and 8 on great sums (大和)

Each problem follows the classical structure: 或問 (problem statement), 法曰 (computational formula), and 草曰 (detailed derivation).

In the original text, counting-rod notation and intermediate polynomial expressions are represented by placeholders (■); these correspond to the “Tai Ji” (太極) and “Tian Yuan” (天元) positions on the counting board. **Key terminology in *tian yuan shu*:**

- *tian yuan yi* = the unknown x
- *li tian yuan yi* = Let x be...
- *ji zuo* = store on the left (for later comparison)
- *xiang xiao* = mutual cancellation (set equal)
- *kai ping fang* = extract square root
- *kai li fang* = extract cube root
- *kai san cheng fang* = extract fourth root

1 卷七 明前一十八問

或問：出南門東行七十二步有樹，出東門南行三十步見之。問答同前。

法曰：倍南行以乘倍東行為平實，並二行又倍之為從，一虛隅。得城徑。

草曰：識別得此問名為弦外容圓，又為內率求虛積。其二行步相並為虛弦，若以相減即虛較也。又倍東行為弦較和，倍南行即弦較較，此二數相乘則兩虛積也。若直以二行相乘，則半個虛積也。又倍東行減於城徑，餘即二虛勾也。倍南行減於城徑則二虛股也。虛積上三事和即城徑也。乃立天元一為圓徑，便以為三事和也。倍二行步減之，得 $\square$ 為黃方一，天元乘之得 $\square$ 為二虛積（寄左）。然後倍東行以乘倍南行，得八千六百四十為同數，與左相消得 $\square$ 。益積開平方得二百四十步，即城徑也。合問。

法曰：二行步相乘為實，二行步相並為從，一步虛法。得半徑。

草曰：立天元一為半徑，副置二位。上加東行步得 $\square$ 為大差勾，下位加股得 $\square$ 為小差股。此二數相乘得下式 $\square$ 為半段黃方幂（寄左）。然後立天元以自之，又二之，與左相消得 $\square$ 。益積開平方得一百二十步，即半城徑也。

Problem: Going out the South Gate and walking 72 paces east, there is a tree. Going out the East Gate and walking 30 paces south, one sees it. Question and answer as before (find the city diameter).

Method: Double the southward walk and multiply by double the eastward walk for the constant term; sum the two walks and double for the linear term; negative unity as the leading coefficient. Extract to obtain the city diameter.

Working: By identification: this problem is called “circle contained outside the hypotenuse.” The two walks summed give the empty hypotenuse; their difference gives the empty difference. Double the eastward walk is the sum of hypotenuse and difference; double the southward walk is their difference. The product of these two quantities is twice the empty area. Multiplying the walks directly gives half the empty area. Subtracting double the eastward walk from the diameter leaves twice the empty base; subtracting double the southward walk gives twice the empty leg. The three-element sum of the empty area is the city diameter.

Let x be the diameter, which is the three-element sum. Subtracting twice the two walks gives $\square$ as one yellow square; multiplying by x gives $\square$ as twice the empty area (store left). Then double the eastward walk times double the southward walk gives 8640 as the matching number. Cancelling with the left gives $\square$. Extracting by accumulated opening of the square gives 240 paces, the city diameter. QED.

Method: Multiply the two walks for the constant term; sum the two walks for the linear coefficient; negative unity as leading coefficient. Extract to obtain the radius.

Working: Let x be the radius, set aside in two positions. Adding the eastward walk to the upper gives $\square$ as the large-difference base; adding the leg to the lower gives $\square$ as the small-difference leg. Multiplying these two quantities gives $\square$ as half the yellow-square power (store left). Then squaring x and doubling, cancelling with the left gives $\square$. Accumulated opening of the square gives 120 paces, the half-diameter.

法日：二雲數相乘倍之於上，加雲數差纂，權寄。並二雲數又自增乘，得數內減上位為平實，並雲數而倍之為從，二步益隅。得半徑。

草日：立天元一為半徑，副之。上減明勾得下□為虛勾，下減股得□為虛股。勾股相乘得□，又倍之得□，又加二行差纂□，得□為弦纂(寄左)。然後並雲數，以自之得□於太極位，為同數，與左相消得□。益積開平方得一百二十步，即半城徑也。

法日：雲數相乘又倍之為平實，雲數相減為從，一常法。得虛勾。

草日：立天元一為虛勾。以南行減東行餘四十二步為虛較也。以虛較加天元得□為虛股，以天元乘之得下□為直積(寄左)。然後倍南行乘東行得□，與左相消得□。開平方得四十八步，即虛勾也。以勾除積得九十步，即虛股也。並勾股得□為虛和也，內加入二行並□得□，即圓徑也。

法日：並二行步以自乘於上，又倍南行乘倍東行，加上位為平實，一隅法。得小和。

Method: Multiply the two given numbers and double, placing above. Add the square of their difference, storing provisionally. Sum the two given numbers and square self-multiply; subtract the stored quantity for the constant term. Double the sum of the given numbers for the linear coefficient. Negative two as the leading coefficient. Extract to obtain the radius.

Working: Let x be the radius, set aside. Subtracting from above gives □ as the empty base; subtracting the leg from below gives □ as the empty leg. Base times leg gives □; doubling gives □; adding the square of the walk-difference □ gives □ as the hypotenuse power (store left). Then summing the given numbers and squaring gives □ at the Tai Ji position as the matching number. Cancelling with the left gives □. Accumulated opening of the square gives 120 paces, the half-diameter.

Method: Multiply the given numbers and double for the constant term; their difference as the linear coefficient; unity as the constant leading coefficient. Extract to obtain the empty base.

Working: Let x be the empty base. South walk subtracted from east walk leaves 42 paces as the empty difference. Adding the empty difference to x gives □ as the empty leg; multiplying by x gives □ as the rectangular area (store left). Then double the south walk times the east walk gives □; cancelling with the left gives □. Extracting the square root gives 48 paces, the empty base. Dividing the area by the base gives 90 paces, the empty leg. Summing base and leg gives □ as the empty sum; adding the sum of the two walks □ gives □, which is the diameter.

Method: Sum the two walks and square, placing above. Also double the south walk times double the east walk; add to the above for the constant term; unity as leading coefficient. Extract to obtain the small sum.

草曰：立天元一為小和。並二行步加之得 $\square$ 為三事和也。倍二行步而並之得 $\square$ ，以減三事和，餘 $\square$ 為黃方，卻以三事和乘之，得下 $\square$ 為二虛積也（寄左）。乃倍南行以乘倍東行，得 $\square$ 為同數，與左相消得 $\square$ 。開平方得一百三十八步，即虛和也。加入二行步得二百四十步，即城徑也。合問。

Working: Let x be the small sum. Adding the sum of the two walks gives $\square$ as the three-element sum. Doubling and summing the two walks gives $\square$; subtracting from the three-element sum, the remainder $\square$ is the yellow square. Multiplying back by the three-element sum gives $\square$ as twice the empty area (store left). Then double the south walk times double the east walk gives $\square$ as the matching number; cancelling with the left gives $\square$. Extracting the square root gives 138 paces, the empty sum. Adding the two walks gives 240 paces, the city diameter. QED.

或問：丙出南門直行一百三十五步而立，甲出東門直行一十六步見之。問答同前。

Problem: Bing goes out the South Gate and walks 135 paces straight, then stands. Jia goes out the East Gate and walks 16 paces straight and sees him. Question and answer as before.

法曰：以丙行步一百三十五再自之，得二百四十六萬零三百七十五於上。又以甲行一十六乘丙行幂一萬八千二百二十五，得二十九萬一千六百，以乘上位，得七千一百七十四億四千五百三十五萬為三乘方實；以二行步相乘又倍之得四千三百二十，以乘丙行步再自之數，得一百六億二千八百八十二萬為益從；第一廉空；以甲行乘丙行幂，得二十九萬一千六百，又倍之得五十八萬三千二百於上，四之甲行幂一千零二十四，以乘丙行步，得一十三萬八千二百四十，減上位餘四十四萬四千九百六十為第二廉；二行步相乘得二千一百六十為虛常法。得丙行步上勾弦差八十一。

Method: Take Bing's walk of 135 and cube it, getting 246,0375 above. Multiply Jia's walk 16 by Bing's square 18,225, getting 291,600; multiply this by the above, getting 71,744,535,000,000 as the quartic constant term. Multiply the two walks and double to get 4320; multiply by Bing's cubed walk, getting 10,628,820,000 as the negative linear coefficient. The first fractional coefficient is empty. Multiply Jia's walk by Bing's square to get 291,600; double to get 583,200 above. Four times Jia's square (1024) times Bing's walk gives 138,240; subtracting from above leaves 444,960 as the second fractional coefficient. The product of the two walks, 2160, is the negative constant leading coefficient. Extract to obtain the base-hypotenuse difference on Bing's walk: 81.

草曰：識別二數相並，得一百五十一。以減於皇極弦，餘一百三十八，即虛勾虛股並也。若以二數相減，餘一百一十九為高弦內減平弦，又為皇極弦內少個小差弦，又為大差弦內減個皇極弦也。立天元一為丙行大差數。置丙行步一百三十五，自乘得 $\square$ ，用天元除之，得 $\square$ 為勾弦並也。上減天元得 $\square$ 為二丙勾也。復用丙南行乘之，得 $\square$ 為二積也。又以天元除之，得 $\square$ 為丙勾外容圓半（泛寄）。別置丙南行用二甲勾乘之，得 $\square$ 太，合用二丙勾除之。不受除，便以此為甲股（內寄二丙勾為分母）。復用二甲勾三十二乘之，得 $\square$ 太為二個甲直積也。又置丙南行內減天元，得 $\square$ 為黃方，以自乘得 $\square$ 為丙上勾弦差乘股弦差二段，以天元除之得 $\square$ 為兩個丙小差也。乃用甲股乘之，得下式 $\square$ 。復用丙南行除之，得 $\square$ ，又折半得下式 $\square$ 為一個甲步股弦差也，內亦帶前二丙勾分母。復置兩個甲直積，內已寄此甲股弦差分母，便為甲步股外容圓半（寄左）。乃再置先求到泛寄，用甲股弦差分母乘之，得 $\square$ 為同數，與左相消得下式 $\square$ 。開三乘方得八十一步，即丙步上勾弦差也。《鈐經》載此法，以勾弦差率冪減丙行冪，復以丙行乘之為實，以差率冪為法，如法得徑。依法求之，勾弦差八十一自之得六千五百六十一，以減於丙行冪一萬八千二百二十五，餘一萬一千六百六十四，復以丙行一百三十五乘之，得一百五十七萬四千六百四十為實，以大差冪六千五百六十一為法，如法得二百四十步，即城徑也。

Working: Identification: the two numbers summed give 151. Subtracting from the imperial extreme hypotenuse leaves 138, the sum of empty base and empty leg. Their difference, 119, is the high hypotenuse minus the level hypotenuse.

Let x be the large-difference number on Bing's walk. Take Bing's walk 135, square it to get $\square$; divide by x to get $\square$ as the sum of base and hypotenuse. Subtracting x above gives $\square$ as twice Bing's base. Multiply by Bing's south walk to get $\square$ as twice the area. Dividing by x gives $\square$ as the half-circle outside Bing's base.

Separately, take Bing's south walk multiplied by twice Jia's base, getting $\square$ at Tai. Not accepting division, take this as Jia's leg. Multiply again by twice Jia's base (32) to get $\square$ at Tai as twice Jia's area.

Bing's south walk minus x gives $\square$ as the yellow square. Squaring gives $\square$ as two segments of base-hypotenuse difference times leg-hypotenuse difference. Dividing by x gives $\square$ as two of Bing's small differences. Multiply by Jia's leg to get $\square$. Dividing by Bing's south walk and halving gives $\square$ as one Jia-walk leg-hypotenuse difference.

Twice Jia's area, carrying the leg-hypotenuse difference denominator, is the half-circle outside Jia's leg (store left). The general store multiplied by the denominator gives $\square$ as the matching number. Cancelling with the left gives $\square$. Extracting the fourth root gives 81 paces.

The *Qian Jing* records: subtract the base-hypotenuse difference-rate square from Bing's square, multiply by Bing's walk for the dividend, using the difference-rate square as divisor; obtain the diameter. Following this: 81 squared gives 6561; subtracting from 18,225 leaves 11,664; multiplying by 135 gives 1,574,640 as dividend; using 6561 as divisor, obtain 240 paces, the city diameter.

或問：出東門一十六步有樹，出南門東行七十二步見之。問答同前。

法曰：二行步相減得數，以自之於上。又以出東門步自之，減上位為平方實，二之出南門東行步為益從，一步常法。翻開得半徑。

Problem: Going out the East Gate 16 paces there is a tree; going out the South Gate and walking 72 paces east, one sees it. Question and answer as before.

Method: Subtract the two walks; square the result, placing above. Also square the East Gate exit walk; subtract from above for the quadratic constant term. Twice the south-gate eastward walk as the negative linear coefficient; unity as the constant leading coefficient. Reverse extraction to obtain the radius.

草曰：別得人到樹即平弦也，半圓徑即平股也。其東行七十二步則平勾平弦差也。乃立天元一為半圓徑，加一十六減七十二，得 $\square$ 為勾也。以自之得 $\square$ 為勾幂。又加入天元股幂得 $\square$ 為弦幂（寄左）。再立天元一為半徑，加出東門步，得 $\square$ 即弦也。以自之得 $\square$ 為同數，與左相消得 $\square$ 。翻法開之得一百二十步，即半城徑也。合問。

Working: By identification: the person reaching the tree is the level hypotenuse; the half-diameter is the level leg. The 72-pace eastward walk is the difference between level base and level hypotenuse. Let x be the half-diameter. Adding 16 and subtracting 72 gives $\square$ as the base. Squaring gives $\square$ as the base power. Adding the x leg power gives $\square$ as the hypotenuse power (store left). Again let x be the radius; adding the East Gate exit walk gives $\square$ as the hypotenuse. Squaring gives $\square$ as the matching number; cancelling with the left gives $\square$. Reverse-method extraction gives 120 paces, the half-diameter. QED.

或問：出南門一百三十五步有樹，出東門南行三十步見之。問答同前。

Problem: Going out the South Gate 135 paces there is a tree; going out the East Gate and walking 30 paces south, one sees it. Question and answer as before.

法曰：樹去城步內減南行步，餘以為幂於上，又以樹去城步為幂內減上位為平實，倍樹去城步為從，一虛隅。翻法得半城徑。

Method: Subtract the south walk from the tree-to-city distance; square the remainder, placing above. Also square the tree-to-city distance; subtract the above from it for the quadratic constant term. Double the tree-to-city distance as the linear coefficient; negative unity as leading coefficient. Reverse extraction to obtain the half-diameter.

草曰：別得人距樹即高弦也，半圓徑即高勾也，其南行三十步即高弦上小差也。乃立天元一為半徑，加樹去城步為弦，內減小差 $\square$ ，得 $\square$ 即股也。以自之得 $\square$ 為股幂，內加入天元幂，得 $\square$ 為弦幂（寄左）。再置弦 $\square$ 以自之，得 $\square$ 為同數，與左相消，得式 $\square$ 。翻開得一百二十步，即半城徑也。合問。

Working: By identification: the person's distance to the tree is the high hypotenuse; the half-diameter is the high base; the 30-pace south walk is the small difference on the high hypotenuse. Let x be the radius. Add the tree-to-city distance for the hypotenuse; subtract the small difference $\square$ to get $\square$ as the leg. Squaring gives $\square$ as the leg power; adding the x power gives $\square$ as the hypotenuse power (store left). Place the hypotenuse $\square$ and square to get $\square$ as the matching number; cancelling with the left gives the expression $\square$. Reverse extraction gives 120 paces, the half-diameter. QED.

或問：乙出東門不知遠近而立，甲出南門東行七十二步望見乙，就乙斜行一百三十六步與乙相會。問答同前。

Problem: Yi goes out the East Gate and stands at an unknown distance. Jia goes out the South Gate and walks 72 paces east, sees Yi, then walks 136 paces diagonally to meet him. Question and answer as before.

法日：以斜行步自之於上，以二行相減餘自為纂，減上位為平實，從空，一步常法。如法得半徑。

草日：別得七十二步即大差也，斜行即弦，半徑即股也。立天元一為半徑，以自之為股纂，又以二行差六十四以自之得 $\square$ 為勾纂。並二纂得 $\square$ 為弦纂(寄左)。然後以斜行步自之，得 $\square$ 為同數，與左相消得 $\square$ 。開平方得一百二十步，倍之即城徑也。合問。

Method: Square the diagonal walk, placing above. Square the difference of the two walks; subtract from above for the quadratic constant term. Empty linear term; unity as leading coefficient. By the method obtain the radius.

Working: By identification: the 72 paces is the large difference; the diagonal walk is the hypotenuse; the radius is the leg. Let x be the radius. Squaring gives the leg power. The two-walk difference 64 squared gives $\square$ as the base power. Combining the two powers gives $\square$ as the hypotenuse power (store left). Then squaring the diagonal gives $\square$ as the matching number; cancelling with the left gives $\square$. Extracting the square root gives 120 paces; doubling gives the city diameter. QED.

或問：甲出南門不知遠近而立，乙出東門南行三十步望見甲，卻就甲斜行二百五十五步與甲相會。問答同前。

法日：二行差自之為纂，以減於斜行纂為平實，一虛隅。得半徑。

草日：別得南行步即股弦差也，斜步即弦也，半徑即勾也。乃立天元一為半城徑，以自之為纂。以二行相減餘二百二十五，以自之得 $\square$ 為股纂。二纂相並得 $\square$ 為弦纂(寄左)。然後以斜行步自之，得 $\square$ 為同數，與左相消得下 $\square$ 。開平方得一百二十步，即半徑也。合問。

Problem: Jia goes out the South Gate and stands at an unknown distance. Yi goes out the East Gate and walks 30 paces south, sees Jia, then walks 255 paces diagonally to meet him. Question and answer as before.

Method: Square the difference of the two walks; subtract from the diagonal square for the quadratic constant term; negative unity as leading coefficient. Extract to obtain the radius.

Working: By identification: the south walk is the leg-hypotenuse difference; the diagonal walk is the hypotenuse; the radius is the base. Let x be the half-diameter. Squaring gives the base power. The two-walk difference 225 squared gives $\square$ as the leg power. Combining the two powers gives $\square$ as the hypotenuse power (store left). Then squaring the diagonal gives $\square$ as the matching number; cancelling with the left gives $\square$. Extracting the square root gives 120 paces, the radius. QED.

或問：甲出南門東行不知步數而立，乙出東門南行三十步望見甲，斜行一百二步相會。問答同前。

Problem: Jia goes out the South Gate and walks east an unknown number of paces, then stands. Yi goes out the East Gate and walks 30 paces south, sees Jia, and walks 102 paces diagonally to meet him. Question and answer as before.

法曰：二行相減，餘以乘乙南行，四之於上，又加入斜行冪為平實。得虛和一百三十八。

草曰：別得斜步內減南行為甲東行步也。此問以弦外容圓入之，以二行相減數乘乙南行三十步，得 $\square$ ，又四之，得 $\square$ 為二直積也。又加入斜步冪 $\square$ ，共得 $\square$ 即和冪也。平方而一，得一百三十八步，即虛和也。又加斜步得二百四十步，即城徑也。合問。

Method: Subtract the two walks; multiply the remainder by Yi's south walk; quadruple, placing above. Add the diagonal square for the quadratic constant term. Extract to obtain the empty sum: 138.

Working: By identification: subtracting the south walk from the diagonal gives Jia's eastward walk. The two-walk difference times Yi's 30-pace south walk gives $\square$; quadrupling gives $\square$ as twice the rectangular area. Adding the diagonal square $\square$ gives $\square$ as the sum-square. Extracting the square root gives 138 paces, the empty sum. Adding the diagonal gives 240 paces, the city diameter. QED.

或問：乙出東門南行不知步數而立，甲出南門東行七十二步望見乙，斜行一百二步與乙相會。問答同前。

法曰：倍相減步以乘倍東行，得數復以減於斜步冪，餘為實。平方而一，得較也。又以二行相減數乘倍東行為平實，以較為從方，得勾。勾較共為長，又以斜步並入勾股共，即城徑。

草曰：別得二行相減餘 $\square$ 為乙南行步也。以此數又減於甲東行，餘四十二步即較也。又以二行相減數 $\square$ 乘倍東行得 $\square$ 為平實，以較為從。平方開得四十八即勾也。勾內加較得九十步即股也。勾股共得一百三十八，又加入斜步，共得二百四十步，即城徑也。合問。

Problem: Yi goes out the East Gate and walks south an unknown number of paces, then stands. Jia goes out the South Gate and walks 72 paces east, sees Yi, and walks 102 paces diagonally to meet him. Question and answer as before.

Method: Double the subtraction-difference and multiply by double the eastward walk; subtract this from the diagonal square, the remainder is the dividend. Extracting the square root gives the difference. Also multiply the two-walk difference by double the eastward walk for the quadratic constant term; use the difference as the linear coefficient; extract to obtain the base. The base-plus-difference sum is the leg; adding the diagonal to the base-plus-leg sum gives the city diameter.

Working: By identification: the two-walk difference remainder $\square$ is Yi's south walk. Subtracting this again from Jia's eastward walk leaves 42 paces as the difference. Also multiply the two-walk difference $\square$ by double the eastward walk to get $\square$ as the quadratic constant term, using the difference as the linear coefficient. Extracting the square root gives 48 as the base. Adding the difference to the base gives 90 paces as the leg. The base-leg sum is 138; adding the diagonal gives 240 paces, the city diameter. QED.

或問：乙出南門東行，甲出東門南行，兩相望見。既而乙雲：「我東行不及城徑一百六十八步。」甲雲：「我南行不及城徑二百一十步。」問答同前。

法曰：半甲不及步以自之為冪，半甲不及步內減差以自之為冪。二冪相並內卻減差冪為平實，四之甲不及內減三之乙不及，餘為益從，三步半虛法。得甲南行。

草曰：別得乙不及為虛勾、半徑共，又為徑內減明勾也。甲不及為虛股、半徑共，又為徑內減股也。又二雲數相並為虛和、圓徑共也，雲數相減即虛較也。乃立天元一為甲南行，以減於甲不及步又半之，得 $\square$ 為虛股也。虛股內減虛較得 $\square$ 為虛勾。勾自之得 $\square$ 為勾冪也，又股自之得下式 $\square$ 為股冪也。二冪相並得 $\square$ 為弦冪（寄左）。然後以天元加虛較得 $\square$ 為乙東行。又加入天元甲南行得 $\square$ 為虛弦，以自之得 $\square$ 為同數，與左相消得 $\square$ 。開平方得三十步，即甲南行也。內加少步，即城徑也。合問。

Problem: Yi goes out the South Gate and walks east; Jia goes out the East Gate and walks south; they see each other. Then Yi says: “My eastward walk falls short of the diameter by 168 paces.” Jia says: “My southward walk falls short of the diameter by 210 paces.” Question and answer as before.

Method: Half Jia’s shortfall squared as one power; half Jia’s shortfall minus the difference, squared as another power. Combine the two powers and subtract the difference-power for the quadratic constant term. Four times Jia’s shortfall minus three times Yi’s shortfall, remainder as the negative linear coefficient. 3.5 as negative leading coefficient. Extract to obtain Jia’s south walk.

Working: By identification: Yi’s shortfall is the empty base plus radius, also the diameter minus the bright base. Jia’s shortfall is the empty leg plus radius, also the diameter minus the leg. The two given numbers combined are the empty sum plus diameter; their difference is the empty difference. Let x be Jia’s south walk. Subtracting from half Jia’s shortfall gives $\square$ as the empty leg. Subtracting the empty difference from the empty leg gives $\square$ as the empty base. Squaring the base gives $\square$ as the base power; squaring the leg gives $\square$ as the leg power. Combining the two powers gives $\square$ as the hypotenuse power (store left). Then adding the empty difference to x gives $\square$ as Yi’s eastward walk. Adding x (Jia’s south walk) gives $\square$ as the empty hypotenuse. Squaring gives $\square$ as the matching number; cancelling with the left gives $\square$. Extracting the square root gives 30 paces, Jia’s south walk. Adding the shortfall gives the city diameter. QED.

或問：丙出南門直行，甲出東門直行，兩相望見。既而丙雲：「我行少於城徑一百五十步。」甲雲：「我行少於城徑二百二十四步。」問答同前。

Problem: Bing goes out the South Gate and walks straight; Jia goes out the East Gate and walks straight; they see each other. Then Bing says: “My walk is 105 paces short of the diameter.” Jia says: “My walk is 224 paces short of the diameter.” Question and answer as before.

法日：二少步相乘訖，又自乘為實；六之共步，乘雲數相乘數為益從；十八之雲數相乘於上，又三之共步，自乘加上位，內復減丙少步冪、甲少步冪為從廉；四十八之共步為益二廉，六十三步常法。翻法開三乘方，得一百二十步，即半徑。

草日：別得雲數共減於倍城徑為甲丙共行數。又雲數相減即皇極差，亦為甲行不及丙行數。立天元一為半城徑，以三之，副置二位。上位減丙少步，得 $\square$ 為皇極股也，下位減甲少步得 $\square$ 為皇極勾也。勾股相乘得 $\square$ ，以天元除之，得 $\square$ 為弦也。弦自之得 $\square$ 為弦冪(寄左)。然後以股自之得下 $\square$ 為股冪於上，又以勾自之得 $\square$ 為勾冪，並以加入上位，得 $\square$ 為同數，與左相消得 $\square$ 。翻法開三乘方得一百二十步，即半城徑也。合問。

Method: Multiply the two shortfalls, then self-multiply for the dividend. Six times the sum times the product of the given numbers as the negative linear term. Eighteen times the product of the given numbers above; also three times the sum, self-multiplied, added to above; subtract Bing's shortfall-square and Jia's shortfall-square for the first fractional coefficient. Forty-eight times the sum as the negative second fractional coefficient; 63 as the constant leading coefficient. Reverse-method extraction of the fourth root gives 120 paces, the radius.

Working: By identification: the sum of given numbers subtracted from twice the diameter is Jia and Bing's combined walk. Their difference is the imperial extreme difference, also Jia's walk short of Bing's walk.

Let x be the half-diameter. Triple it and set aside in two positions. Subtract Bing's shortfall from the upper to get $\square$ as the imperial extreme leg; subtract Jia's shortfall from the lower to get $\square$ as the imperial extreme base. Base times leg gives $\square$; dividing by x gives $\square$ as the hypotenuse. Squaring the hypotenuse gives $\square$ as the hypotenuse power (store left). Then squaring the leg gives $\square$ as the leg power above; squaring the base gives $\square$ as the base power. Combining and adding to above gives $\square$ as the matching number; cancelling with the left gives $\square$. Reverse-method fourth-root extraction gives 120 paces, the half-diameter. QED.

或問：甲出東門直行，乙出南門直行，各不知步數而立。乙望見甲，就甲斜行了二百八十九步與甲相會。其二直行共得一百五十一歩。又雲甲直行少於乙直行。問答同前。

法日：斜冪內減共步冪為平實，倍共步內減斜步為從，一常法。得徑。

Problem: Jia goes out the East Gate and walks straight; Yi goes out the South Gate and walks straight; each stands at an unknown distance. Yi sees Jia and walks 289 paces diagonally to meet him. Their two straight walks sum to 151 paces. It is also said that Jia's walk is less than Yi's. Question and answer as before.

Method: Subtract the sum-square from the diagonal-square for the quadratic constant term; subtract the diagonal from double the sum for the linear coefficient; unity as leading coefficient. Extract to obtain the diameter.

草曰：別得共數、城徑並即皇極和也。立天元一為圓徑，加共步得 $\square$ 為皇極和，以自之，得 $\square$ 於上。以斜行幂 $\square$ 減上位餘 $\square$ 為二直積（寄左）。然後以天元乘斜步得 $\square$ ，與左相消得 $\square$ 。開平方得二百四十步，即城徑也。合問。

Working: By identification: the sum plus the diameter is the imperial extreme sum.

Let x be the diameter. Adding the sum gives $\square$ as the imperial extreme sum; squaring gives $\square$ above. Subtracting the diagonal square $\square$ leaves $\square$ as twice the rectangular area (store left). Then x times the diagonal gives $\square$; cancelling with the left gives $\square$. Extracting the square root gives 240 paces, the city diameter. QED.

或問：甲出東門直行，乙出東門南行，丙出南門直行，丁出南門東行，各不知步數而立。四人遙相望，悉與城參相直。只雲甲、丙共行了一百五十一歩，乙、丁立處相距一百二歩。又雲丙直行步多於甲直行步。問答同前。

Problem: Jia goes out the East Gate and walks straight; Yi goes out the East Gate and walks south; Bing goes out the South Gate and walks straight; Ding goes out the South Gate and walks east; each stands at an unknown distance. The four see each other from afar, all aligned with the city. It is only said that Jia and Bing together walked 151 paces; the distance between Yi and Ding's positions is 102 paces. It is also said that Bing's straight walk exceeds Jia's. Question and answer as before.

法曰：共步、距步相減，得數自之於上，以共步為幂內減上為平實，二之距步內減共步、距步差為從，一步虛法。得城徑。

Method: Subtract the distance-walk from the sum-walk; square the result, placing above. Subtract above from the sum-square for the quadratic constant term. Subtract the sum-distance difference from twice the distance for the linear coefficient; negative unity as leading coefficient. Extract to obtain the city diameter.

草曰：別得共步得城徑即皇極和也，相距步即虛弦也。皇極和內減虛弦即皇極弦也。又共步、距步差 $\square$ 即皇極弦內減城徑也（此名旁差）。乃立天元一為城徑，加共步得 $\square$ 為皇極和也，以自之得 $\square$ 於上，以共步、距步差 $\square$ 加天元得 $\square$ 為皇極弦也，以自之得下式 $\square$ ，減上位餘得 $\square$ 為二直積（寄左）。然後以天元徑乘皇極弦，得 $\square$ 為同數，與左相消得 $\square$ 。開平方得二百四十歩，即城徑也。合問。

Working: By identification: the sum equals the diameter, which is the imperial extreme sum; the distance is the empty hypotenuse. Subtracting the empty hypotenuse from the imperial extreme sum gives the imperial extreme hypotenuse. The sum-distance difference $\square$ is the imperial extreme hypotenuse minus the diameter (this is called the side difference).

Let x be the diameter. Adding the sum gives $\square$ as the imperial extreme sum; squaring gives $\square$ above. Adding the sum-distance difference $\square$ to x gives $\square$ as the imperial extreme hypotenuse. Squaring gives $\square$; subtracting from above leaves $\square$ as twice the rectangular area (store left). Then x (diameter) times the imperial extreme hypotenuse gives $\square$ as the matching number; cancelling with the left gives $\square$. Extracting the square root gives 240 paces, the city diameter. QED.

或問：甲出南門東行不知步數而立，乙出東門南行望見甲，復就甲斜行，與甲相會。乙通計行了一百三十二步，其乙南行步不及斜行七十二步，其甲東行卻多於乙南行。問答同前。

法日：倍不及步在地，以不及步減通步以乘之為實，以四之不及步為法。得乙南行三十步。

草日：別得乙南行即股也，以減通步即虛弦也，以減不及步即虛較也，其不及步即甲東行也。立天元一為乙南行，置不及步以天元乘之，又四之得 $\square$ 元為二直積（寄左）。然後倍不及步以為弦較和於上 $\square$ 。以不及步減通步得 $\square$ 為弦較較。以乘上位得 $\square$ 太為同數，與左相消得 $\square$ 。上法下實，得三十步為乙南行也。餘各以數求之。

Problem: Jia goes out the South Gate and walks east an unknown number of paces, then stands. Yi goes out the East Gate and walks south, sees Jia, then walks diagonally to meet him. Yi's total walk is 132 paces; Yi's south walk falls 72 paces short of the diagonal walk; Jia's eastward walk exceeds Yi's south walk. Question and answer as before.

Method: Place double the shortfall; multiply by the total walk minus the shortfall for the dividend; use four times the shortfall as the divisor. Obtain Yi's south walk: 30 paces.

Working: By identification: Yi's south walk is the leg; subtracting from the total gives the empty hypotenuse; subtracting the shortfall gives the empty difference; the shortfall is Jia's eastward walk. Let x be Yi's south walk. Place the shortfall, multiply by x , and quadruple to get $\square x$ as twice the rectangular area (store left). Then double the shortfall as the sum of hypotenuse and difference above $\square$. Total walk minus shortfall gives $\square$ as the difference of hypotenuse and difference. Multiplying by above gives $\square$ at Tai as the matching number; cancelling with the left gives $\square$. Dividing dividend by divisor gives 30 paces as Yi's south walk. The remainder are each found from the numbers.

或問：甲出南門東行不知步數而立。乙出東門南行，望見甲，復斜行與甲相會。二人共行了二百四步，又雲甲行不及共步一百三十二。問答同前。

法日：別得二行共即兩個虛弦也，其不及步即乙南行與一虛弦共也。置不及步內減一弦餘三十步，即乙南行也。以乙南行反以減虛弦，餘七十二步即甲東行也。以乙南行減甲東行餘即虛較也。

草日：此問無草。

Problem: Jia goes out the South Gate and walks east an unknown number of paces, then stands. Yi goes out the East Gate and walks south, sees Jia, then walks diagonally to meet him. The two together walked 204 paces; it is also said that Jia's walk falls 132 paces short of the total. Question and answer as before.

Method: By identification: the two-walk total is twice the empty hypotenuse; the shortfall is Yi's south walk plus one empty hypotenuse. Place the shortfall and subtract one hypotenuse, leaving 30 paces as Yi's south walk. Subtracting Yi's south walk from the empty hypotenuse leaves 72 paces as Jia's eastward walk. Subtracting Yi's south walk from Jia's eastward walk gives the empty difference.

Working: This problem has no working.

或問：乙出東門南行，甲出西門南行，甲望見乙，斜行五百一十步相會。乙雲：「我南行少於城徑二百一十步。」問答同前。

法曰：少步冪為平實，四斜步內減二少步為益從，五步常法。得乙南行。

草曰：別得少步為徑內減股。立天元一為乙南行，以二之減於倍斜行步，得 $\square$ 為梯底也。以二之天元乘之，得 $\square$ 為徑冪(寄左)。再置天元加少步，得下式 $\square$ 為城徑，以自之得 $\square$ ，與左相消得 $\square$ 。開平方得三十步，即乙南行也。加少步即城徑也。合問。

Problem: Yi goes out the East Gate and walks south; Jia goes out the West Gate and walks south. Jia sees Yi and walks 510 paces diagonally to meet him. Yi says: "My south walk is 210 paces short of the diameter." Question and answer as before.

Method: The shortfall-square as quadratic constant term; four times the diagonal minus twice the shortfall as the negative linear coefficient; 5 as the constant leading coefficient. Extract to obtain Yi's south walk.

Working: By identification: the shortfall is the diameter minus the leg. Let x be Yi's south walk. Doubling and subtracting from twice the diagonal gives $\square$ as the ladder base. Multiplying by twice x gives $\square$ as the diameter-square (store left). Place x plus the shortfall to get $\square$ as the city diameter; squaring gives $\square$. Cancelling with the left gives $\square$. Extracting the square root gives 30 paces as Yi's south walk. Adding the shortfall gives the city diameter. QED.

或問：乙出南門東行，甲出北門東行，甲望見乙，斜行二百七十二步與乙相會。乙雲：「我東行不及城徑一百六十八步。」問答同前。

法曰：以不及步冪之為實，四斜內減二之不及步為虛從，五常法。平開得乙東行七十二步。

草曰：別得不及步為城徑減明勾也。立天元一為乙東行，以倍之減於二之斜行步，得下 $\square$ 為梯底也。倍天元乘之，得 $\square$ 為徑冪(寄左)。再置天元加不及步，得 $\square$ 為城徑，以自之得 $\square$ 為同數，與左相消得 $\square$ 。開平方得七十二步，即乙東行也。加入少步即城徑也。合問。

Problem: Yi goes out the South Gate and walks east; Jia goes out the North Gate and walks east. Jia sees Yi and walks 272 paces diagonally to meet him. Yi says: "My eastward walk falls 168 paces short of the diameter." Question and answer as before.

Method: The shortfall-square as dividend; four times the diagonal minus twice the shortfall as the negative linear coefficient; 5 as constant leading coefficient. Extract to obtain Yi's eastward walk: 72 paces.

Working: By identification: the shortfall is the city diameter minus the bright base. Let x be Yi's eastward walk. Doubling and subtracting from twice the diagonal gives $\square$ below as the ladder base. Multiply by double x to get $\square$ as the diameter-square (store left). Place x plus the shortfall to get $\square$ as the city diameter; squaring gives $\square$ as the matching number. Cancelling with the left gives $\square$. Extracting the square root gives 72 paces, Yi's eastward walk. Adding the shortfall gives the city diameter. QED.

或問：乙出南門東行，丁出東門南行，卻有甲丙二人共在西北隅，甲向東行，丙向南行，四人遙相望見，俱與城參相直。既而相會，甲雲：「我多乙二百四十八步。」丙雲：「我多於丁五百七十步。」問答同前。

法曰：二多步相乘為平實，並二多步而半之為從，七分半常法。得城徑。

草曰：別得甲多步為大勾內減明勾也，丙多步為大股內少股也。又乙東行得一虛勾為半徑，丁南行得一虛股為半徑。又二多數相並得 $\square$ 為大和內少虛弦也，又二少數相減餘 $\square$ 為兩個角差。又甲多步內減半徑即勾方差也，丙多步內減半徑即股方差也。立天元一為城徑，以半之減於甲多步得 $\square$ 為勾方差，又以半徑減於丙多步得 $\square$ 為股方差。二差相乘得 $\square$ 為徑幂（寄左）。然後以天元幂與左相消，得下式 $\square$ 。開平方得二百四十步，即城徑也。合問。

Problem: Yi goes out the South Gate and walks east; Ding goes out the East Gate and walks south. Also Jia and Bing are both at the northwest corner; Jia walks east and Bing walks south. The four see each other from afar, all aligned with the city. After meeting, Jia says: “I exceed Yi by 248 paces.” Bing says: “I exceed Ding by 570 paces.” Question and answer as before.

Method: Multiply the two excess-walks for the quadratic constant term; half the sum of the two excess-walks as the linear coefficient; 7.5 as the constant leading coefficient. Extract to obtain the city diameter.

Working: By identification: Jia’s excess is the large base minus the bright base; Bing’s excess is the large leg minus the *zhong* leg. Also Yi’s eastward walk gives one empty base as the radius; Ding’s southward walk gives one empty leg as the radius. The two excesses combined give $\square$ as the great sum minus the empty hypotenuse; their difference gives $\square$ as two corner differences. Also Jia’s excess minus the radius is the base-square difference; Bing’s excess minus the radius is the leg-square difference. Let x be the diameter. Subtracting half of it from Jia’s excess gives $\square$ as the base-square difference; subtracting the radius from Bing’s excess gives $\square$ as the leg-square difference. Multiplying the two differences gives $\square$ as the diameter-square (store left). Then x -squared cancelling with the left gives $\square$. Extracting the square root gives 240 paces, the city diameter. QED.

或問：甲丙二人俱在西北隅，甲向東行，丙向南行。又乙出南門東行，丁出東門南行，各不知步數而立。四人遙相望見，悉與城參相直。既而相會，甲雲：「我與乙共行了三百九十二步。」丙雲：「我與丁共行了六百三十步。」問答同前。

Problem: Jia and Bing are both at the northwest corner; Jia walks east and Bing walks south. Also Yi goes out the South Gate and walks east; Ding goes out the East Gate and walks south; each stands at an unknown distance. The four see each other from afar, all aligned with the city. After meeting, Jia says: “Yi and I together walked 392 paces.” Bing says: “Ding and I together walked 630 paces.” Question and answer as before.

法曰：甲乙共自之為冪，丙丁共自之為冪，二冪又相乘為三乘方實。甲乙共自之為冪，以丙丁共乘之於上。又以丙丁共自之為冪，以甲乙共乘之加上位為益從。甲乙共自之為冪，丙丁共自之為冪，並，以七分半乘之於上。又以甲乙共乘丙丁共，得數減上位為第一益廉。並二共數，以七分半乘之為第二廉。以七分半自之，得五分六厘二毫五絲於上位。以一步內減上位，餘四分三厘七毫五絲為虛隅。得城徑。

草曰：別得甲為大勾，乙為明勾，丙為大股，丁為股也。甲乙共內減半徑即是黃長弦也，丙丁共內減半徑即黃廣弦也。黃長弦、黃廣弦二數相減，餘為兩個皇極差也。乃立天元為城徑，半之副置二位。上以減於甲乙共數，得 $\square$ 即黃長弦也，以自之得 $\square$ 為黃長弦冪也，內減天元一冪，餘得下式 $\square$ 為勾方差冪也。下位以減於丙丁共，得下式 $\square$ 即黃廣弦也，以自之得 $\square$ 為黃廣弦冪也，內減天元一冪，餘得 $\square$ 為股方差冪也。再以勾方差冪、股方差冪相乘，得 $\square$ 為徑冪(寄左)。然後以天元為冪，又以冪自之，與左相消得下式 $\square$ 。開三乘方得二百四十步，即城徑也。合問。

Method: Square the Jia-Yi sum; square the Bing-Ding sum; multiply the two powers for the quartic constant term. Square the Jia-Yi sum and multiply by the Bing-Ding sum, placing above. Also square the Bing-Ding sum and multiply by the Jia-Yi sum, adding to above for the negative linear coefficient. Square the Jia-Yi sum and the Bing-Ding sum, combine, multiply by 7.5, placing above. Also multiply the Jia-Yi sum by the Bing-Ding sum; subtract from above for the first negative fractional coefficient. Sum the two combined numbers, multiply by 7.5 for the second fractional coefficient. Square 7.5 to get 0.5625 above. Subtract this from unity; the remainder 0.4375 is the negative leading coefficient. Extract to obtain the diameter.

Working: By identification: Jia is the large base, Yi is the bright base, Bing is the large leg, Ding is the *zhong* leg. The Jia-Yi sum minus the radius is the yellow-long hypotenuse; the Bing-Ding sum minus the radius is the yellow-wide hypotenuse. The difference of yellow-long and yellow-wide hypotenuses is twice the imperial extreme difference. Let x be the diameter. Halve it and set aside in two positions. Subtract from above the Jia-Yi sum to get $\square$ as the yellow-long hypotenuse; squaring gives $\square$ as the yellow-long hypotenuse-square. Subtracting x -squared leaves $\square$ as the base-square-difference power. Subtract from below the Bing-Ding sum to get $\square$ as the yellow-wide hypotenuse; squaring gives $\square$ as the yellow-wide hypotenuse-square. Subtracting x -squared leaves $\square$ as the leg-square-difference power. Multiply the base-square-difference power by the leg-square-difference power to get $\square$ as the diameter-square (store left). Then x squared and self-squared again, cancelling with the left gives $\square$. Extracting the fourth root gives 240 paces, the city diameter. QED.

2 卷八 明後一十六問

或問：出南門向東有槐樹一株，出東門向南有柳樹一株。丙丁俱出南門，丙直行，丁往至槐樹下。甲乙俱出東門，甲直行，乙往至柳樹下。四人遙相望見，各不知所行步數。只雲丙丁共行了二百七步，甲乙共行了四十六步，又雲甲丙立處相距二百八十九步。問答同前。

法日：以二共相減數又以減距數為實，二為法。得平勾。

草日：識別得丙丁共即明和也，甲乙共即和也，相距步即極弦也。二共相並即極弦內少個虛黃也，又為極和內少個虛和也。二共相減餘為平勾高股差也，又為虛差極差共也，又為通差內減極差也。立天元為平勾，加入二共相減數，得 $\square$ 為高股，又加天元得 $\square$ 為極弦（寄左）。以相距步二百八十九與左相消，得 $\square$ 。上法下實，如法得六十四，即平勾也。以二共相減數加平勾得二百二十五為高股，復以平勾乘之，得一萬四千四百步。開平方得一百二十步，即城半徑也。合問。

或問：依前見丙丁共二百七步，甲乙共四十六步。又雲二樹相去一百二步。問答同前。

法日：以甲乙共乘樹相去步，得數又以自之為平實；從空；並二共數為冪於上，內減甲乙共自之數、丙丁共自之數為益隅。得弦。

Problem: Going out the South Gate eastward there is a locust tree; going out the East Gate southward there is a willow tree. Bing and Ding both go out the South Gate; Bing walks straight, Ding goes to the locust tree. Jia and Yi both go out the East Gate; Jia walks straight, Yi goes to the willow tree. The four see each other from afar, each not knowing their walked paces. It is only said that Bing and Ding together walked 207 paces; Jia and Yi together walked 46 paces; also that the distance between Jia and Bing's positions is 289 paces. Question and answer as before.

Method: Subtract the difference of the two sums from the distance for the dividend; 2 as the divisor. Obtain the level base.

Working: By identification: the Bing-Ding sum is the bright sum; the Jia-Yi sum is the *zhong* sum; the distance is the extreme hypotenuse. The two sums combined equal the extreme hypotenuse minus the empty yellow, also the extreme sum minus the empty sum. The difference of the two sums is the difference of level base and high leg. Let x be the level base. Add the difference of the two sums to get $\square$ as the high leg; add x to get $\square$ as the extreme hypotenuse (store left). The distance 289 cancelling with the left gives $\square$. Dividing gives 64, the level base. Adding the two-sum difference to the level base gives 225 as the high leg; multiplying by the level base gives 14,400. Extracting the square root gives 120 paces, the half-radius. QED.

Problem: As before: Bing-Ding sum 207 paces, Jia-Yi sum 46 paces. Also said: the two trees are 102 paces apart. Question and answer as before.

Method: Multiply the Jia-Yi sum by the tree-distance; square the result for the quadratic constant term. Empty linear term. Sum the two combined numbers and square above; subtract the Jia-Yi sum-square and the Bing-Ding sum-square for the negative leading coefficient. Extract to obtain the hypotenuse.

草曰：識別得兩樹相去步即虛弦也。餘數具前。立天元一為弦。置明和以天元乘之，合和除，不除，便以 $\square$ 元為明弦也（內帶和分母）。乃置虛弦以分母和乘之，得 $\square$ 太，加入明弦，得 $\square$ 為極股也，內帶和分母。以自之得下式 $\square$ 為極股幂（內寄和幂為分母）。又以天元加虛弦得下 $\square$ 為極勾，以自之得 $\square$ 。又以和幂 $\square$ 乘之，得 $\square$ 為勾幂也。勾股幂相並得 $\square$ 為兩積一較幂也，內有和幂分母（寄左）。然後置明弦 $\square$ 元於上，以和乘天元加上位，得 $\square$ 元為二弦並也。又置虛弦以和乘之得 $\square$ 太，並入上位，得下式 $\square$ 為極弦，以自之得 $\square$ 為同數，與左相消得 $\square$ 。開平方得三十四步，即弦也。

Working: By identification: the tree distance is the empty hypotenuse.

Let x be the hypotenuse. Place the bright sum multiplied by x ; should divide by the sum, but not dividing, take $\square x$ as the bright hypotenuse (with sum denominator). Place the empty hypotenuse multiplied by the sum denominator to get $\square$ at Tai; add the bright hypotenuse to get $\square$ as the extreme leg, carrying the sum denominator. Squaring gives $\square$ as the extreme leg power (with sum-square denominator).

Adding x to the empty hypotenuse gives $\square$ below as the extreme base; squaring gives $\square$. Multiply by sum-square $\square$ to get $\square$ as the base power. Combining base and leg powers gives $\square$ as twice-area-plus-difference power (store left). Then place bright hypotenuse $\square x$ above; add sum times x to get $\square x$ as two-hypotenuse combined. Empty hypotenuse times sum gives $\square$ at Tai; combining gives $\square$ as the extreme hypotenuse. Squaring gives $\square$; cancelling with the left gives $\square$. Extracting gives 34 paces, the hypotenuse.

或問：皇極大小差共一百八十七步，明黃、黃共六十六步。問答同前。

法曰：後數自乘為實，前後數相減，餘為法。得虛黃方三十六。

草曰：別得一百八十七即明弦二弦共也，其六十六即太虛大小差共也。又二數相並，得 $\square$ 即明、二和共。若以相減，餘 $\square$ 即明、四差共也。立天元一為太虛黃方面，加二黃共，得 $\square$ 即虛弦也。倍虛弦又加天元，得 $\square$ 即城徑也。又以虛弦加皇極大小差，得 $\square$ 即極弦也，以極弦乘城徑得 $\square$ ，為兩段皇極勾股積（寄左）。再以極弦虛弦相並，得 $\square$ ，即皇極勾股共也，自之得 $\square$ ，內減皇極弦幂 $\square$ ，得 $\square$ 為同數，與寄左相消得 $\square$ 。上法下實，如法得三十六步，即太虛黃方面也。合問。

Problem: The imperial extreme large-and-small-difference sum is 187 paces; the bright yellow and *zhong* yellow sum is 66 paces. Question and answer as before.

Method: Square the latter number for the dividend; subtract the latter from the former for the divisor. Obtain the empty yellow square: 36.

Working: By identification: 187 is the bright hypotenuse plus *zhong* hypotenuse combined; 66 is the great empty large-and-small-difference sum. Let x be the great empty yellow square side. Add the two-yellow sum to get $\square$ as the empty hypotenuse. Double the empty hypotenuse and add x to get $\square$ as the city diameter. Add the empty hypotenuse to the imperial extreme large-small-difference to get $\square$ as the extreme hypotenuse. Multiply by diameter to get $\square$ as twice the imperial extreme base-leg area (store left). Combine extreme hypotenuse and empty hypotenuse to get $\square$ as the imperial extreme base-leg sum. Squaring gives $\square$; subtracting the imperial extreme hypotenuse-square $\square$ leaves $\square$ as the matching number. Cancelling gives $\square$. Dividing gives 36 paces, the great empty yellow square side. QED.

或問：東門南有柳一株，南門東有槐一株。甲出東門直行，丙出南門直行。甲、丙、柳、槐悉與城參相直，既而甲就柳樹斜行三十四步至柳樹下，丙就槐樹斜行一百五十三步至槐樹下。問答同前。

法曰：雲數相乘，倍之便為平方實，開方得虛弦一百二步。以此弦加甲行步即極勾，以此弦加丙行步即極股。餘各依法求之。

草曰：識別：甲斜行即弦也，丙斜行即明弦也。無草。

Problem: South of the East Gate there is a willow tree; east of the South Gate there is a locust tree. Jia goes out the East Gate and walks straight; Bing goes out the South Gate and walks straight. Jia, Bing, the willow, and the locust are all aligned with the city. Then Jia walks 34 paces diagonally to the willow tree; Bing walks 153 paces diagonally to the locust tree. Question and answer as before.

Method: Multiply the given numbers and double for the quadratic constant term; extracting the root gives the empty hypotenuse: 120 paces. Adding this hypotenuse to Jia's walk gives the extreme base; adding to Bing's walk gives the extreme leg. The remainder are each found by the method.

Working: Identification: Jia's diagonal walk is the *zhong* hypotenuse; Bing's diagonal walk is the bright hypotenuse. No working.

或問：東門南有柳一株，南門東有槐一株。甲出東門直行，丙出南門直行，二人遙相望，槐柳與城邊悉相直。既而甲復斜行至柳樹下，丙復斜行至槐樹下，各不知步數。只雲丙共行了二百八十八步，甲斜行與柳至東門步共得六十四步。問答同前。

法曰：二雲數相乘於上，以六十四步自之，又二之，減上位為平實。四之六十四於上，倍丙行內減上位為從。二十常法。得甲直行步一十六。

Problem: South of the East Gate there is a willow; east of the South Gate there is a locust. Jia goes out the East Gate and walks straight; Bing goes out the South Gate and walks straight; the two see each other from afar, the locust and willow both aligned with the city wall. Then Jia walks diagonally to the willow tree, and Bing walks diagonally to the locust tree, each unknown. It is only said that Bing's total walk is 288 paces; Jia's diagonal walk plus the willow-to-East-Gate distance sums to 64 paces. Question and answer as before.

Method: Multiply the two given numbers, placing above. Square 64 paces and double, subtracting from above for the quadratic constant term. Four times 64 above; subtract from double Bing's walk for the linear coefficient. 20 as constant leading coefficient. Extract to obtain Jia's straight walk: 16 paces.

草曰：別得丙共步即明股、明弦和也，六十四即平勾也，內甲斜行即弦也，柳至東門步即股也。又二雲數相並即明差與極弦共也，二雲數相減即明差與平勾高股差共也。又平勾內減勾即虛勾也。立天元一為勾。置丙共步以天元乘之，復以六十四除之得 $\square$ 為明勾也。又以天元減於六十四，得 $\square$ 為虛勾也，並虛明二勾 $\square$ 為半徑也。以自之得 $\square$ ，倍之得 $\square$ 為半段圓城徑幂（寄左）。乃以天元加六十四，得 $\square$ 為勾圓差於上；又以明勾加丙共步，得 $\square$ 為股圓差於下。上下相乘得 $\square$ 為同數，與左相消得 $\square$ 。開平方得一十六步，即勾也。此勾乃甲出東門直行步也。餘皆依數求之。合問。

Working: By identification: Bing's total is the bright leg plus bright hypotenuse sum; 64 is the level base; Jia's diagonal is the *zhong* hypotenuse; the willow-to-East-Gate distance is the *zhong* leg. The two given numbers combined are the bright difference plus extreme hypotenuse. Let x be the base. Place Bing's total, multiply by x , divide by 64 to get $\square$ as the bright base. Subtract x from 64 to get $\square$ as the empty base; combining gives $\square$ as the radius. Squaring gives $\square$; doubling gives $\square$ as half the city-diameter power. Then x plus 64 gives $\square$ as the base-circle-difference above; also bright base plus Bing's total gives $\square$ as the leg-circle-difference below. Multiplying gives $\square$; cancelling with the left gives $\square$. Extracting gives 16 paces, the base. This is Jia's straight walk. QED.

或問：東門南有柳樹一株，南門東有槐樹一株。甲出東門直行，丙出南門直行，二人遙相望，槐柳與城邊悉相直。既而甲復斜行至柳樹下，丙復斜行至槐樹下，各不知步數。只雲甲共行五十步，丙斜行與槐至南門步共得二百二十五步。問答同前。

Problem: South of the East Gate there is a willow tree; east of the South Gate there is a locust tree. Jia goes out the East Gate and walks straight; Bing goes out the South Gate and walks straight; the two see each other from afar, the locust and willow both aligned with the city wall. Then Jia walks diagonally to the willow tree, and Bing walks diagonally to the locust tree, each unknown. It is only said that Jia's total walk is 50 paces; Bing's diagonal walk plus the locust-to-South-Gate distance sums to 225 paces. Question and answer as before.

法曰：以二百二十五步自之為幂，又以此幂自為幂於上。置甲共行以二百二十五步三度乘之，得數復折半，減上位為平實。置二百二十五步自之數，以二雲數相減數乘之，又倍之於上。倍五十步在地，以二百二十五步自之數乘之，復折半加上位為益從。雲數相減自乘於上，以雲數相乘，復折半，減上位為常法。得明股。

Method: Square 225 paces for a power; square this power again, placing above. Multiply Jia's total by 225 cubed; halve and subtract from above for the quadratic constant term. Place 225 squared, multiply by the given-numbers difference, and double, placing above. Place double 50 paces, multiply by 225 squared, halve and add to above for the negative linear coefficient. Square the given-numbers difference above; multiply the given numbers, halve, and subtract from above for the constant leading coefficient. Extract to obtain the bright leg.

草曰：識別得甲共步即勾、弦共也，二百二十五即高股也，內丙斜行即明弦，槐至南門步即明勾也。又高股內減明股即虛股也。立天元一為明股，即丙出南門直行步也。置五十步以天元乘之得 $\square$ 元，合高股除。不除，便以此 $\square$ 元為股也，內帶高股 $\square$ 分母。再置高股內減天元得 $\square$ 為虛股，以分母高股乘之，得下式 $\square$ ，加入股得 $\square$ 即半徑也。以自增乘得下 $\square$ 為半徑幕也。內帶高股幕為母（寄左）。然後置甲共步以分母高股乘之，得 $\square$ 太，加入股得 $\square$ 為勾圓差於上（內帶高股分母）。又以天元加高股得 $\square$ 為股圓差於下。上、下相乘得 $\square$ 。又以分母高股乘之，得 $\square$ ，復折半得 $\square$ 為同數，與左相消得 $\square$ 。開平方得一百三十五步，即明股也。合問。

Working: By identification: Jia's total is the base-hypotenuse sum; 225 is the high leg; Bing's diagonal walk is the bright hypotenuse; the locust-to-South-Gate distance is the bright base. The high leg minus bright leg gives the empty leg.

Let x be the bright leg. Place 50 paces, multiply by x to get $\square x$; should divide by high leg. Not dividing, take $\square x$ as the leg, carrying high-leg $\square$ as denominator. Subtract x from the high leg to get $\square$ as the empty leg; multiply by denominator high leg to get $\square$; add the leg to get $\square$ as the radius. Self-multiplying gives $\square$ as the radius power (store left). Place Jia's total multiplied by denominator high leg to get $\square$ at Tai; add the leg to get $\square$ as the base-circle-difference above. Also x plus high leg gives $\square$ as the leg-circle-difference below. Multiply to get $\square$; multiply by denominator to get $\square$; halve to get $\square$ as the matching number. Cancelling gives $\square$. Extracting gives 135 paces, the bright leg. QED.

或問：通勾、通弦共一千步，勾、弦共五十步。問答同前。

法曰：置一千減二之五十步為泛率。以自乘，復半之於上。又置泛率復以五十乘之，加上位為平實。二十二之泛率於上。以四十二乘五十，得數內減泛率，加上位為益從。二百為常法。得股。

Problem: The general base plus general hypotenuse sum is 1000 paces; the *zhong* base plus *zhong* hypotenuse sum is 50 paces. Question and answer as before.

Method: Place 1000 minus twice 50 paces as the general rate. Square and halve, placing above. Also multiply the general rate by 50 and add to above for the quadratic constant term. 22 times the general rate above; 42 times 50, subtract the general rate, and add to above for the negative linear coefficient. 200 as constant leading coefficient. Extract to obtain the leg.

草曰：立天元一為股。置一千以天元乘之，以五十除之，得 $\square$ 元為通股也。又以天元加五十步，得 $\square$ 即小差也，通股加小差得 $\square$ 即通弦也。以通弦減一千得 $\square$ ，即通勾也。以小差減通勾得 $\square$ ，即圓徑也。以圓徑減通股得 $\square$ 即大差也。置大差以小差乘之得 $\square$ （寄左）。然後置圓徑以自之得 $\square$ ，折半得 $\square$ ，與左相消得 $\square$ 。開平方得三十步，即股也。合問。

Working: Let x be the leg. Place 1000, multiply by x , divide by 50 to get $\square x$ as the general leg. Add 50 paces to x to get $\square$ as the small difference; general leg plus small difference gives $\square$ as the general hypotenuse. Subtracting from 1000 gives $\square$ as the general base. Subtracting small difference gives $\square$ as the diameter. Subtracting diameter from general leg gives $\square$ as the large difference. Place large difference, multiply by small difference to get $\square$ (store left). Then square the diameter to get $\square$; halve to get $\square$; cancelling gives $\square$. Extracting gives 30 paces, the leg. QED.

或問：通勾、通弦共一千步，明勾、明弦共二百二十五步。問答同前。

法曰：以後數再自乘，又以前數乘之為平實；以後數為冪，復以前數乘之為從；以前數冪為虛常法。得明股。

草曰：別得二百二十五步即高股也。立天元一為明股。置一千以天元乘之，合以高股除不受除，便以此一零零零元為通股（內帶高股為母）。以天元加高股，得 $\square$ 即大差也。置大差以高股分母乘之，得 $\square$ ，即帶分大差也。以此減於通股，餘 $\square$ 即圓徑也。以自增乘，得 $\square$ （寄左。內帶高股冪分母）。然後置一千以高股分母通之，得 $\square$ 太，內減帶分大差，得 $\square$ 為兩個通勾也。內減兩個圓徑得 $\square$ ，為兩個小差也。以帶分大差乘之，得下式 $\square$ 為同數，與左相消得 $\square$ 。開平方得一百三十五步，即明股也。合問。

Problem: The general base plus general hypotenuse sum is 1000 paces; the bright base plus bright hypotenuse sum is 225 paces. Question and answer as before.

Method: Cube the latter number and multiply by the former for the quadratic constant term; square the latter and multiply by the former for the linear coefficient; the former squared as negative constant leading coefficient. Extract to obtain the bright leg.

Working: By identification: 225 paces is the high leg.

Let x be the bright leg. Place 1000, multiply by x ; should divide by high leg but not accepting division, take $1000x$ as the general leg. Add high leg to x to get $\square$ as the large difference. Multiply by high-leg denominator to get $\square$ as the large difference with denominator. Subtracting from the general leg leaves $\square$ as the diameter. Self-multiplying gives $\square$ (store left).

Convert 1000 by high-leg denominator to get $\square$ at Tai; subtract the large difference with denominator to get $\square$ as twice the general base. Subtract twice the diameter to get $\square$ as twice the small difference. Multiply by the large difference with denominator to get $\square$. Cancelling gives $\square$. Extracting gives 135 paces, the bright leg. QED.

或問：通股、通弦共一千二百八十步，股、弦共六十四步。問答同前。

法曰：雲數相乘為平實，前數為益從。置前數以後數除之，得二十為泛率。泛率減一以自乘於上，又倍泛率減一加上位為常法。倒積開得勾一十六。

Problem: The general leg plus general hypotenuse sum is 1280 paces; the *zhong* leg plus *zhong* hypotenuse sum is 64 paces. Question and answer as before.

Method: Multiply the given numbers for the quadratic constant term; the former number as negative linear coefficient. Divide the former by the latter to get 20 as the general rate. General rate minus one, squared, placing above; also twice general rate minus one, added to above for the constant leading coefficient. Reverse-area extraction gives the base: 16.

草曰：別得六十四步即平勾也。立天元一為勾。置前數以天元乘之，以後數除之，得 $\square$ 元即通勾也。又置天元加後數，得 $\square$ 即小差也。以小差減通勾，餘 $\square$ 即圓徑也。以自之得 $\square$ (寄左)。然後以小差減於前數，得 $\square$ 為二通股。內減兩個圓徑，得 $\square$ 為二大差也。以小差乘之得下 $\square$ ，與左相消得 $\square$ 。開平方得一十六步，即勾也。合問。

Working: By identification: 64 paces is the level base.

Let x be the base. Place the former number, multiply by x , divide by the latter to get $\square x$ as the general base. Also add the latter to x to get $\square$ as the small difference. Subtracting from general base leaves $\square$ as the diameter. Squaring gives $\square$ (store left).

Subtract small difference from the former number to get $\square$ as twice the general leg. Subtract twice the diameter to get $\square$ as twice the large difference. Multiply by small difference to get $\square$; cancelling gives $\square$. Extracting gives 16 paces, the base. QED.

或問：通股、通弦共一千二百八十步，明股、明弦共二百八十八步。問答同前。

法曰：二數相減，以後數乘之，內減後數幂，又半之為泛率。以自之為平實。置前數加二之後數而半之為次率，以乘泛率，倍之於上，以後數乘泛率減上位為益從。次率自乘於上，以前數加次率，復以後數乘之，減上位為隅法。得明勾。

Problem: The general leg plus general hypotenuse sum is 1280 paces; the bright leg plus bright hypotenuse sum is 288 paces. Question and answer as before.

Method: Subtract the two numbers; multiply by the latter; subtract the latter-square, and halve as the general rate. Square it for the quadratic constant term. Add twice the latter to the former and halve as the secondary rate; multiply by the general rate and double, placing above. Multiply the latter by the general rate and subtract from above for the negative linear coefficient. Square the secondary rate above; add the former and secondary rate, multiply again by the latter, and subtract from above for the leading coefficient. Extract to obtain the bright base.

草曰：別得二數相減，餘 $\square$ 為通勾、通股及明勾共也。立天元一為明勾。置前數以天元乘之，合以後數除之。不除，便以此 $\square$ 元為通勾也 (內寄後數分母)。又以二數相減，得數內又減天元得 $\square$ 為通和也。乃以分母二百八十八之，得下式 $\square$ ，內減通勾，餘 $\square$ 為通股也。又以天元加後數，又以分母 (即後數也) 通之，得 $\square$ 為大差也。以此大差減於通股，得下式 $\square$ ，為一個圓徑也。半之得 $\square$ ，以自之得 $\square$ 為半徑幂 (寄左)。然後以半圓徑減通勾，得 $\square$ 為底勾，又以天元乘之，又以分母二百八十八之，得 $\square$ 為同數，與左相消得 $\square$ 。開平方得七十二步，即明勾也。合問。

Working: By identification: the two numbers subtracted leave $\square$ as the general base, general leg, and bright base combined.

Let x be the bright base. Place the former number, multiply by x ; should divide by the latter but not dividing, take $\square x$ as the general base. Subtract the two numbers and further subtract x to get $\square$ as the general sum. Convert by denominator 288 to get $\square$; subtract general base to leave $\square$ as the general leg. Add x to the latter and convert to get $\square$ as the large difference. Subtract from general leg to get $\square$ as the diameter. Halve to get $\square$; squaring gives $\square$ as the radius power (store left).

Subtract half-diameter from general base to get $\square$ as the bottom base; multiply by x and convert by 288 to get $\square$ as the matching number. Cancelling gives $\square$. Extracting gives 72 paces, the bright base. QED.

或問：明股、明弦並二百八十八步，勾、弦並五十步。又雲明股、勾並多於虛弦四十九步。問答同前。

法曰：前二數相並，內減二之多步，即圓徑。又只以前二數相乘，便是半徑幂。

草曰：識別得前二數相減而半之，即極差也。其多步名傍差，又為圓徑不及極弦數。

Problem: The bright leg plus bright hypotenuse sum is 288 paces; the *zhong* base plus *zhong* hypotenuse sum is 50 paces. It is also said that the bright leg plus *zhong* base sum exceeds the empty hypotenuse by 49 paces. Question and answer as before.

Method: Combine the former two numbers and subtract twice the excess; this is the diameter. Also simply multiplying the former two numbers gives the radius-square.

Working: By identification: halving the difference of the former two numbers gives the extreme difference. The excess is called the side difference, also the amount by which the diameter falls short of the extreme hypotenuse.

或問：平差、高差共一百六十一，明股、勾並多於虛弦四十九步。問答同前。

法曰：二數相減又半之，以自乘為實，後數為法。得平勾。

草曰：立天元一為平勾。以加前數得 $\square$ 為高股也。又以天元加高股得 $\square$ 為極弦，內減後數得 $\square$ ，又半之，得 $\square$ 為半徑。以自之，得 $\square$ (寄左)。然後以天元乘高股，得 $\square$ 為同數，與左相消得 $\square$ 。上法下實，得六十四步，即平勾也。合問。

Problem: The level difference plus high difference sum is 161 paces; the bright leg plus *zhong* base sum exceeds the empty hypotenuse by 49 paces. Question and answer as before.

Method: Subtract and halve the two numbers; square for the dividend; the latter number as divisor. Obtain the level base.

Working: Let x be the level base. Adding the former number gives $\square$ as the high leg. Also adding x to the high leg gives $\square$ as the extreme hypotenuse; subtracting the latter number gives $\square$; halving gives $\square$ as the radius. Squaring gives $\square$ (store left). Then x times the high leg gives $\square$ as the matching number; cancelling with the left gives $\square$. Dividing gives 64 paces, the level base. QED.

或問：平勾、高股差一百六十一，明差、差並七十七步。又雲極弦多於城徑四十九步。問答同前。

Problem: The difference of level base and high leg is 161 paces; the bright difference plus *zhong* difference sum is 77 paces. It is also said that the extreme hypotenuse exceeds the diameter by 49 paces. Question and answer as before.

法日：並上二位而半之為平率。其四十九即旁率也。副置平率，上加旁率，下減旁率，以相乘為實。倍旁差為法。得勾圓差。

草日：立天元一為半徑，又為半之股圓差上弦較較，又為半之勾圓差上弦較和也。內減勾圓差上勾股較 $\square$ ，餘 $\square$ 為半之勾圓差上弦較較也。置股圓差上勾股較 $\square$ ，以半之勾圓差上弦較較乘之，得 $\square$ （寄左）。然後以半之股圓差上弦較較乘勾圓差上勾股較，得 $\square$ 元為同數，與左相消得下式 $\square$ 。上法下實，得一百二十步，即半徑也。合問。

草日：識別得平勾、高股差名角差。副置角差，上加七十七而半之，得 $\square$ ，即極差也。下減七十七而半之，得 $\square$ ，即虛差也。角差加極差得 $\square$ ，即通差也。又極弦多於城徑步名為旁差。副置角差，上加旁差得 $\square$ ，為兩個高段上勾股較，下減旁差得 $\square$ 為兩個平段上勾股較也。又副置極差，上加旁差得 $\square$ 為股圓差上勾股較，下減旁差 $\square$ 為勾圓差上勾股較也。立天元一為勾圓差。依法求得通差，加入天元得 $\square$ 即大差也。以天元乘之得 $\square$ 為半段圓徑幂（寄左）。乃置大差 $\square$ 內減股圓差上勾股較 $\square$ ，餘有 $\square$ 為股圓差之勾於上。再置天元內加勾圓差上勾股較 $\square$ ，得 $\square$ 為勾圓差之股。以乘上位得 $\square$ 為同數，與左相消得 $\square$ 。上法下實，得八十步，即勾圓差也。

Method: Combine and halve the upper two numbers for the level rate. The 49 is the side rate. Set aside the level rate; add the side rate above, subtract below; multiply for the dividend. Double the side difference as divisor. Obtain the base-circle-difference.

Working: Let x be the radius, also half the sum-of-chord-differences on the circle-leg-difference, also half the difference-of-chord-differences on the circle-base-difference. Subtracting the base-leg difference $\square$ on the circle-base-difference leaves $\square$ as half the difference of chord differences. Place the base-leg difference $\square$ on the circle-leg difference, multiply by half the difference of chord differences to get $\square$ (store left). Then half the sum of chord differences times the base-leg difference gives $\square x$ as the matching number; cancelling with the left gives $\square$. Dividing gives 120 paces, the radius. QED.

Working: By identification: the level-base/high-leg difference is called the corner difference. Set aside; add 77 above and halve to get $\square$ as the extreme difference; subtract 77 below and halve to get $\square$ as the empty difference. Corner difference plus extreme difference gives $\square$ as the general difference. Also the extreme hypotenuse exceeding the diameter is called the side difference.

Set aside the corner difference; add the side difference above to get $\square$ as twice the high-segment base-leg difference; subtract below to get $\square$ as twice the level-segment base-leg difference. Also set aside the extreme difference; add the side difference above to get $\square$ as the base-leg difference on the circle-leg; subtract below $\square$ for the circle-base. Let x be the circle-base-difference. Finding the general difference and adding to x gives $\square$ as the large difference. Multiply by x to get $\square$ as half the diameter power (store left). Place large difference $\square$ and subtract the base-leg difference $\square$ on the circle-leg, leaving $\square$ as the base on the circle-leg. Also add the base-leg difference $\square$ on the circle-base to x to get $\square$ as the leg on the circle-base. Multiply by above to get $\square$; cancelling gives $\square$. Dividing gives 80 paces, the circle-base-difference.

或問：又依前問：見角差一百六十一步，見明差差並七十七步，又見太虛弦較較六十步。問答同前。

法日：前二數相減而半之，得數加入半之太虛弦較較為泛率，以自乘為平實。置一百六十一內減二之泛率為從，一常法。得平勾。

草日：別得 $\square$ 即二股也。立天元一為平勾。先以前二數相減而半之，得 $\square$ 為虛差。以虛差加股得 $\square$ ，即明勾也。以明勾加天元得 $\square$ ，為平弦，以自之得 $\square$ ，內減天元冪，得 $\square$ 為半徑冪（寄左）。然後以天元加一百六十一為高股，以天元乘之，得 $\square$ 為同數，與左相消得 $\square$ 。開平方得六十四步，即平勾也。

法日：又法日：前數內加半之太虛弦較較，以自乘，內減前數自乘為實，前數內減太虛弦較較為從，一常法。開平方得平勾六十四。此更不用明差差並也。

草日：依前求平勾。前高股內加股，得 $\square$ 為高弦也。以自之得 $\square$ 於上位，內減高股冪 $\square$ ，餘得 $\square$ 為半徑冪（寄左）。然後以天元乘高股，得 $\square$ 為同數，與左相消得下 $\square$ 。開平方得六十四步，即平勾也。合問。

或問：高差、平差並一百六十一，明差、差並七十七步。問答同前。

Problem: Also according to the previous question: the corner difference is 161 paces; the bright-difference plus difference sum is 77 paces; also the great empty chord-difference-difference is 60 paces. Question and answer as before.

Method: Subtract and halve the former two numbers; add half the great empty chord-difference-difference for the general rate; square for the quadratic constant term. Place 161 minus twice the general rate for the linear coefficient; unity as leading coefficient. Extract to obtain the level base.

Working: By identification: $\square$ is twice the leg. Let x be the level base. First subtract and halve the former two numbers to get $\square$ as the empty difference. Adding the empty difference to the leg gives $\square$ as the bright base. Adding the bright base to x gives $\square$ as the level hypotenuse; squaring gives $\square$; subtracting x -squared gives $\square$ as the radius power (store left). Then x plus 161 is the high leg; multiplying by x gives $\square$ as the matching number; cancelling gives $\square$. Extracting gives 64 paces, the level base.

Method: Alternate method: Add half the great empty chord-difference-difference to the former number; square; subtract the former number-square for the dividend. Former number minus the great empty chord-difference-difference for the linear coefficient; unity as leading coefficient. Extracting gives the level base: 64. This further does not use the bright-difference sum.

Working: As before, seek the level base. Add the leg to the previous high leg to get $\square$ as the high hypotenuse. Squaring gives $\square$ above; subtracting high-leg-square $\square$ leaves $\square$ as the radius power (store left). Then x times the high leg gives $\square$ as the matching number; cancelling gives $\square$. Extracting gives 64 paces, the level base. QED.

Problem: The high difference plus level difference sum is 161 paces; the bright difference plus *zhong* difference sum is 77 paces. Question and answer as before.

法曰：以前數自乘於上，二數相並而半之，以自乘減上位，得數復自增乘為平實。前數自之於上，又以四之前數乘之，寄位。以前數自之於上，並二數而半之，以自乘減上位，得數又以四之前數乘，又倍之，減於寄位為從。前數自之，又四之於上。又以四之前數為纂，加上位，權寄。以前數為纂於上，並二數而半之，以自乘減上位，得數復八之於上。又以四之前數為纂加入上位，並以減於權寄為常法。得平勾。

草曰：識別得二位相並而半之，得 $\square$ ，即極差也。立天元一為平勾，加一百六十一得 $\square$ 為高股，高股內又加天元，得 $\square$ 為極弦。以自之得 $\square$ 於上，內減極差纂一萬四千一百六十一，餘 $\square$ 為兩段極積。合以極弦除，不除寄為母，便以此為城徑。以自增乘得 $\square$ 為圓徑纂（內有極弦纂分母。寄左）。然後以天元乘高股，又四之得 $\square$ ，又以分母極弦纂 $\square$ 通之，得 $\square$ 為同數，與左相消得 $\square$ 。開平方得六十四步，即平勾也。合問。

Method: Square the former number, placing above. Combine and halve the two numbers; square and subtract from above; self-multiply the result for the quadratic constant term. Square the former number above; also multiply by four times the former number, storing. Square the former number above; combine and halve the two numbers, square and subtract from above; multiply by four times the former number, double, and subtract from the stored quantity for the linear coefficient. Square the former number, quadruple above; also use four times the former number squared, adding to above, storing provisionally. Use the former number squared above; combine and halve the two numbers, square and subtract from above; octuple above; also add four times the former number squared to above; combine and subtract from the provisional store for the constant leading coefficient. Extract to obtain the level base.

Working: By identification: combining and halving the two positions gives $\square$ as the extreme difference. Let x be the level base. Add 161 to get $\square$ as the high leg; adding x to the high leg gives $\square$ as the extreme hypotenuse. Squaring gives $\square$ above; subtracting extreme-difference-square 14,161 leaves $\square$ as twice the extreme area. Should divide by extreme hypotenuse; not dividing, store as denominator, taking this as the diameter. Self-multiplying gives $\square$ as the diameter-square (carrying extreme-hypotenuse-square denominator; store left). Then x times the high leg, quadrupled gives $\square$; also convert by denominator extreme-hypotenuse-square $\square$ to get $\square$ as the matching number. Cancelling gives $\square$. Extracting gives 64 paces, the level base. QED.

或問：見明和二百七步，和四十六步。問答同前。

Problem: The bright sum is 207 paces; the *zhong* sum is 46 paces. Question and answer as before.

法曰：二和上下相減，數同則止，名為泛率。又以二和直相減，餘為泛實（此則角差也）。乃以泛率除泛實，所得為差率也。以差率加減泛率，若半訖，與勾股相應者，其泛率便為和率，其泛實便為較率乘和率也。若不相應，則直取差率以消息之，定為相管和率。乃以和率約二和，其明和所得為明壘率，其和所得為壘率也。又副置和率，上加差率而半之，則為股率也；下位減差率而半之，則為勾率也。既見勾、股及差三率，各以壘率乘之，即各得勾、股及差之真數也。

法曰：又法：二雲數相並，以自乘於上。二雲數相乘，又四之以減上位為實。二雲數相並，以六步半乘之於上，又二數相並，以四步半乘之，又四之，以並入上位為從方。以七十步零四分三厘七毫五絲為常法。得小差四步。

Method: Subtract the two sums above-below; when the numbers agree, stop—this is called the general rate. Also directly subtract the two sums; the remainder is the general dividend (this is the corner difference). Divide the general dividend by the general rate; the result is the difference rate. Add and subtract the difference rate from the general rate; if when halved it agrees with the base-leg, then the general rate is the sum rate, and the general dividend is the difference rate times the sum rate. If it does not agree, then directly take the difference rate and adjust it, fixing it as the mutual sum rate. Then divide the two sums by the sum rate; the bright-sum result is the bright stack rate, and the *zhong*-sum result is the stack rate. Also set aside the sum rate; add the difference rate above and halve—this is the leg rate; subtract the difference rate below and halve—this is the base rate. Once the three rates of base, leg, and difference are seen, multiply each by the stack rate to obtain the true numbers of base, leg, and difference respectively.

Method: Alternate method: Combine the two given numbers and square, placing above. Multiply the two given numbers and quadruple, subtracting from above for the dividend. Combine the two given numbers, multiply by 6.5 above; also combine the two numbers, multiply by 4.5, and quadruple, combining with above for the linear coefficient. 70.04375 as the constant leading coefficient. Extract to obtain the small difference: 4 paces.

草曰：以二和相約分得率一、明率四步半，其兩數大小差率並同。又別得明小差、大差俱為半虛黃也。立天元一為小差，以四步半乘之得 $\square$ 為大差也，又為明小差，又為半虛黃。置此大差又以四步半乘之得 $\square$ 為明大差也。其四差相並得 $\square$ ，減於二和並，得 $\square$ 即兩段太虛大小差並也。內加三段虛黃方 $\square$ ，得 $\square$ ，合成一個太虛三事和，即圓城徑也。以自增乘得 $\square$ 為徑幕（寄左）。乃置和加半虛黃，得 $\square$ 為平勾。又置明和內加半虛黃，得 $\square$ 為高股，勾股相乘得下式 $\square$ ，又四之得 $\square$ 為同數，與左相消得下式 $\square$ 。開平方得四步，即小差也。合問。

Working: Dividing the two sums yields rate 1 and bright rate 4.5; the two numbers' large-and-small-difference rates are the same. Also: the bright small difference and large difference are both half the empty yellow.

Let x be the small difference. Multiply by 4.5 to get $\square$ as the large difference, also the bright small difference, also half the empty yellow. Multiply this large difference again by 4.5 to get $\square$ as the bright large difference. The four differences combined give $\square$; subtracting from the two-sum combination gives $\square$ as twice the great empty large-small-difference sum. Adding three empty-yellow-squares $\square$ gives $\square$, combining to form one great empty three-element sum, which is the city diameter. Self-multiplying gives $\square$ as the diameter power (store left).

Then the sum plus half empty yellow gives $\square$ as the level base. The bright sum plus half empty yellow gives $\square$ as the high leg. Base times leg gives $\square$; quadrupling gives $\square$ as the matching number. Cancelling gives $\square$. Extracting gives 4 paces, the small difference. QED.

或問：明勾二股共八十八步，明股二勾共一百六十五步。問答同前。

法曰：先識別得二大差共、二小差共及四差共。乃以二大差、二小差相乘為實，以四差共為法。如法得半之虛黃方一十八。

Problem: Twice the bright base plus *zhong* leg totals 88 paces; twice the bright leg plus *zhong* base totals 165 paces. Question and answer as before.

Method: First identify twice the large-difference sum, twice the small-difference sum, and four times the difference sum. Then multiply twice the large difference by twice the small difference for the dividend; four times the difference sum as divisor. By the method obtain half the empty yellow square: 18.

草曰：先置前後雲數以約法約之，得一十一即壘率也。復各置前後數如壘率而一，前得八即勾率也，後得一十五即股率也。再以勾、股率求得較率七，和率二十三，弦率一十七，黃方率六，大差率九，小差率二。既見諸率，各以壘率乘之，其二和共得 $\square$ ，二較共得 $\square$ ，二弦共得 $\square$ ，二黃共得 $\square$ ，二大差共 $\square$ ，二小差共 $\square$ ，四差共 $\square$ 。已上皆為明所得之共數也。乃立天元一為半虛黃，便為明小差，又為大差也。以減於大差共，得 $\square$ 即明大差也。又以減於小差共，得 $\square$ 即小差也。以二數相增乘，得 $\square$ (寄左)。以天元冪與寄左相消，得 $\square$ 。上法下實，得一十八步，即半之虛黃方也。以倍之得 $\square$ ，又加於二黃共六十六共得一百二，即明勾、股共也，又為極黃方，又為虛弦也。又以三十六減於一百八十七，餘一百五十一，即明股勾共也。此數內減虛弦，餘 $\square$ 為明二差較也，此名旁差。以旁差減二弦共一百八十七，餘得 $\square$ ，即太虛和也。卻加入虛弦一百二，並得 $\square$ ，為太虛三事和，即圓城徑也。合問。

Working: First place the former and latter given numbers and divide by the reducing factor to get 11 as the stack rate. Then: the former gives 8 as the base rate, the latter gives 15 as the leg rate. Then find the difference rate 7, sum rate 23, hypotenuse rate 17, yellow square rate 6, large-difference rate 9, and small-difference rate 2.

Once all rates are seen, multiply each by the stack rate. Let x be half the empty yellow, which is the bright small difference, also the large difference. Subtracting from the large-difference sum gives $\square$ as the bright large difference. Also subtracting from the small-difference sum gives $\square$ as the small difference. Multiplying gives $\square$ (store left). x -squared cancelling gives $\square$. Dividing gives 18 paces, half the empty yellow square. Doubling gives $\square$; adding to 66 gives 102, the bright base-leg sum, also the extreme yellow square, also the empty hypotenuse. Subtracting 36 from 187 leaves 151, the bright leg-base sum. Subtracting the empty hypotenuse leaves $\square$ as the bright two-difference, called the side difference. Subtracting from 187 leaves $\square$ as the great empty sum. Adding back the empty hypotenuse 102 gives $\square$ as the great empty three-element sum, which is the city diameter. QED.

3 卷九 大斜四問、大和八問

3.1 細草卷九上 大斜四問

或問：甲丙俱在中心，丙望南門直行，不知步數而止。甲出東門直行，不知步數望見丙，斜行與丙相會。二人共行了六百八十步，仍雲甲直行少於丙直行一百一十九步。問答同前。

法曰：二數相減，餘以為冪，內卻減差冪為平實；二數相減又四之於上，又加入二之差步為益從；二步常法。得皇勾一百三十六。

草曰：別得共步即皇極三事和，少步即勾股差也。立天元一為皇極勾。加少步得 $\square$ 為股也，又以天元加股得 $\square$ 為和也。以和減共步得 $\square$ 為弦也，弦自之得 $\square$ 為一段弦冪(寄左)。然後置股以天元乘之，又倍之，得 $\square$ 為二直積，如入少步冪 $\square$ 共得 $\square$ 為同數，與左相消得 $\square$ 。平方而一，得一百三十六即勾也。勾加差為股，勾股相乘，倍之為實，勾股和減共步為法。得城徑。

法曰：又法：和數與倍差相加、相減，二得數相乘為平實；雲數並與雲數差相並得數，以減於八之共步為益從；一步常法。得皇極黃方一百二。

Problem: Jia and Bing are both at the center. Bing looks toward the South Gate and walks straight, stopping at an unknown number of paces. Jia goes out the East Gate and walks straight an unknown number of paces, sees Bing, and walks diagonally to meet him. The two together walked 680 paces; it is also said that Jia's straight walk is 119 paces less than Bing's. Question and answer as before.

Method: Subtract the two numbers; use the remainder as a power, subtracting the difference-power for the quadratic constant term. Subtract the two numbers and quadruple above; also add twice the difference for the negative linear coefficient. 2 as constant leading coefficient. Extract to obtain the imperial base: 136.

Working: By identification: the total walk is the imperial extreme three-element sum; the shortfall is the base-leg difference.

Let x be the imperial extreme base. Add the shortfall to get $\square$ as the leg; add x to the leg to get $\square$ as the sum. Subtract the sum from the total to get $\square$ as the hypotenuse; squaring gives $\square$ as one hypotenuse power (store left). Then place the leg, multiply by x , and double to get $\square$ as twice the rectangular area. Adding the shortfall-square $\square$ gives $\square$ as the matching number. Cancelling with the left gives $\square$. Extracting gives 136 as the base. Base plus difference is the leg; base times leg, doubled, is the dividend; base-leg sum subtracted from total is the divisor. Obtain the city diameter.

Method: Alternate method: Add and subtract double the difference from the sum; multiply the two results for the quadratic constant term. Combine the sum of given numbers with their difference; subtract from eight times the total for the negative linear coefficient; unity as leading coefficient. Extract to obtain the imperial extreme yellow square: 120.

草曰：立天元一為黃方(即虛弦也)，副置之。上位加共步，得 $\square$ 為二和也；下位減共步，得 $\square$ 為二弦也。先以二和自乘，得 $\square$ 為四段和纂。又以二弦自乘，得 $\square$ 為四段弦纂。二數相減，餘得 $\square$ 元，又倍之得下式 $\square$ 元，為十六段直積於天元位(寄左)。然後副置二和，上位加二之少步，得 $\square$ 為四股。下位減二之少步，得 $\square$ 為四勾。勾股相乘，得 $\square$ 為同數，與左相消得 $\square$ 。平方而一，得一百二步，即皇極黃方也。餘各依法求之。合問。

Working: Let x be the yellow square (the empty hypotenuse), set aside. Add the total to the upper to get $\square$ as twice the sum; subtract the total from the lower to get $\square$ as twice the hypotenuse. First square twice the sum to get $\square$ as four sum-powers. Also square twice the hypotenuse to get $\square$ as four hypotenuse-powers. The two numbers subtracted leave $\square x$; doubling gives $\square x$ as sixteen rectangular areas at the x position (store left).

Then set aside twice the sum; add twice the shortfall above to get $\square$ as four legs. Subtract twice the shortfall below to get $\square$ as four bases. Base times leg gives $\square$ as the matching number; cancelling with the left gives $\square$. Extracting gives 120 paces, the imperial extreme yellow square. The remainder are each found by the method. QED.

或問：甲丙俱在西北隅起，丙向南行不知步數而立。甲向東行望見丙，就丙斜行六百八十步與丙相會。丙雲：「我南行步多於甲東行二百八十步。」問答同前。

Problem: Jia and Bing both start from the northwest corner. Bing walks south an unknown number of paces and stands. Jia walks east, sees Bing, and walks 680 paces diagonally to meet him. Bing says: "My south walk exceeds Jia's eastward walk by 280 paces." Question and answer as before.

法曰：以雲數差乘雲數並為實，倍多步為從，二為平隅。得大勾三百二十。

Method: Multiply the given-numbers difference by their sum for the dividend; double the excess as the linear coefficient; 2 as the quadratic leading coefficient. Extract to obtain the large base: 320.

草曰：立天元為大勾。加差得 $\square$ 為股，倍天元乘之，得 $\square$ 為二積(寄左)。然後以斜步、多步並 $\square$ 與斜步、多步較 $\square$ 相乘，得 $\square$ 為同數，與左相消得 $\square$ 。開平方得三百二十步，即大勾也。合問。

Working: Let x be the large base. Add the difference to get $\square$ as the leg; multiply by double x to get $\square$ as twice the area (store left). Then the diagonal plus excess $\square$ multiplied by the diagonal minus excess $\square$ gives $\square$ as the matching number; cancelling with the left gives $\square$. Extracting gives 320 paces, the large base. QED.

或問：甲乙二人共立於艮隅，乙南行過城門而立，甲東行望乙與城參相直而止。丙丁二人共立於坤隅，丁向東行過城門而立，丙向南行望丁及甲乙悉與城俱相直。丙復就甲斜行六百八十步與甲相會。乙、丁又雲：「吾二人直行共得三百四十二步。」問答同前。

Problem: Jia and Yi stand together at the northeast corner. Yi walks south past the gate and stands; Jia walks east, sees Yi aligned with the city, and stops. Bing and Ding stand together at the southwest corner. Ding walks east past the gate and stands; Bing walks south, sees Ding and Jia-Yi both aligned with the city. Bing then walks 680 paces diagonally to meet Jia. Yi and Ding also say: "Our two straight walks together total 342 paces." Question and answer as before.

法曰：二雲數相乘，倍之為實；倍斜行於上，以二雲數相減加上位為從；一步常法。開平方，得城徑。

草曰：別得斜步即大弦也。其共步則一徑一虛弦共也，其二數相並為一大和一虛弦共數也。立天元為徑，減於共步得 $\square$ 為虛弦也。以虛弦復減於天元，得 $\square$ 為虛和，以斜步乘之，得 $\square$ (寄左)。乃以天元加斜步，得 $\square$ 為大和，以虛弦乘之得 $\square$ 為同數，與左相消得 $\square$ 。開平方得二百四十步，即城徑也。合問。

Method: Multiply the two given numbers and double for the dividend; double the diagonal above; add the difference of the two given numbers to above for the linear coefficient; unity as leading coefficient. Extract the square root to obtain the city diameter.

Working: By identification: the diagonal is the large hypotenuse. The combined walk is one diameter plus one empty hypotenuse; the two numbers combined are one great sum plus one empty hypotenuse.

Let x be the diameter. Subtract from the combined walk to get $\square$ as the empty hypotenuse. Subtract the empty hypotenuse from x to get $\square$ as the empty sum; multiply by the diagonal to get $\square$ (store left). Then x plus diagonal gives $\square$ as the great sum; multiply by the empty hypotenuse to get $\square$ as the matching number. Cancelling with the left gives $\square$. Extracting gives 240 paces, the city diameter. QED.

或問：甲從北門向東直行，庚從西門穿城東行，丙從西門向南直行，壬從北門穿城南行。四人遙相望，悉與城參相直。只雲甲丙相望處六百八十步，庚壬穿城共行了六百三十一。問答同前。

法曰：共步自之得數。以共步減斜，餘自乘以減上為實。二之斜步加入共步減斜，餘數為從。一步常法。得城徑。

Problem: Jia walks east straight from the North Gate; Geng walks east through the city from the West Gate; Bing walks south straight from the West Gate; Ren walks south through the city from the North Gate. The four see each other from afar, all aligned with the city. It is only said that the distance between Jia and Bing is 680 paces; Geng and Ren's through-city walks total 631 paces. Question and answer as before.

Method: Square the combined walk. Subtract the combined walk from the diagonal; square the remainder and subtract from above for the dividend. Twice the diagonal plus the combined-walk-minus-diagonal remainder as the linear coefficient. Unity as leading coefficient. Obtain the city diameter.

草曰：共行步為一徑與皇和共也，又為大和皇弦差也。甲丙相望即大弦也。以共步減大弦，餘□為皇極弦上減一徑也。立天元一為圓徑，減於共步，得□為皇極和也，以自之得□於上。弦內減共步餘□，又以天元加之為皇弦，以自之得□，減上位，餘得□為兩個皇直積（寄左）。乃以天元乘皇弦得下式□為同數，與左相消得□。平方而一，得二百四十步，即城徑也。合問。

Working: The combined walk is one diameter plus the imperial sum, also the great sum minus the imperial hypotenuse. The Jia-Bing distance is the large hypotenuse. Subtracting the combined walk from the large hypotenuse leaves □ as the imperial hypotenuse minus one diameter.

Let x be the diameter. Subtract from the combined walk to get □ as the imperial sum; squaring gives □ above. Subtracting the combined walk from the hypotenuse leaves □; adding x gives the imperial hypotenuse; squaring gives □. Subtracting from above leaves □ as twice the imperial rectangular area (store left). Then x times the imperial hypotenuse gives □ as the matching number; cancelling with the left gives □. Extracting gives 240 paces, the city diameter. QED.

3.2 細草卷九下 大和八問

或問：庚從西門穿城東行二百五十六步而立，壬從北門穿城南行三百七十五步而立。又有甲丙二人俱在乾隅，甲向東行，丙向南行，各不知步數而立。四人遙相望，只雲甲丙共行了九百二十步。問答同前。

法曰：庚東行幕、壬南行幕相並於上，並庚壬步而倍之，內減大和，餘復減於庚壬共，得數以自乘減上位為平實。並庚壬步為益從，半步為隅法。得城徑。

草曰：立天元一為圓徑，以半之副置二位。上以減於庚東行，得下 $\square$ 為平弦也；下以減於壬南行，得 $\square$ 為高弦也。二弦相並得 $\square$ 為皇弦、虛弦共也。倍此數得 $\square$ 為大弦、虛弦共也。以大弦、虛弦共減於大和，餘 $\square$ 為虛勾虛股共也。天元內減虛勾、虛股共，餘 $\square$ 即虛弦也。復置皇弦虛弦共內減虛弦，餘 $\square$ 即皇極弦也，以自之得 $\square$ (寄左)。然後以平弦自之，得下式 $\square$ 為勾幕也，又以高弦自之，得 $\square$ 為股幕也。二幕相並得 $\square$ 為同數，與左相消得 $\square$ 。平方而一，得二百四十步，即城徑也。合問。

Problem: Geng walks east through the city from the West Gate 256 paces and stands; Ren walks south through the city from the North Gate 375 paces and stands. Also Jia and Bing are both at the northwest corner; Jia walks east and Bing walks south, each standing at an unknown distance. The four see each other from afar. It is only said that Jia and Bing together walked 920 paces. Question and answer as before.

Method: Combine Geng's east-walk square and Ren's south-walk square above. Combine the Geng-Ren walks and double; subtract the great sum; subtract again from the Geng-Ren combined walk; square and subtract from above for the quadratic constant term. Combine the Geng-Ren walks as the negative linear coefficient; half a pace as the leading coefficient. Extract to obtain the city diameter.

Working: Let x be the diameter. Halve and set aside in two positions. Subtract from Geng's east walk above to get $\square$ below as the level hypotenuse; subtract from Ren's south walk to get $\square$ as the high hypotenuse. The two hypotenuses combined give $\square$ as the imperial hypotenuse plus empty hypotenuse. Doubling gives $\square$ as the large hypotenuse plus empty hypotenuse. Subtracting from the great sum leaves $\square$ as the empty base plus empty leg. Subtracting the empty base-leg sum from x leaves $\square$ as the empty hypotenuse.

Also take the imperial-empty hypotenuse sum and subtract the empty hypotenuse, leaving $\square$ as the imperial extreme hypotenuse. Squaring gives $\square$ (store left). Then square the level hypotenuse to get $\square$ as the base power; also square the high hypotenuse to get $\square$ as the leg power. Combining the two powers gives $\square$ as the matching number; cancelling gives $\square$. Extracting gives 240 paces, the city diameter. QED.

或問：甲丙俱在西北隅，甲向東行不知步數而立，丙向南行望見甲，就甲斜行與甲相會。甲直行、丙直行共九百二十步（甲步少於丙步）。又：出東門南行有柳樹一株，出南門東行有槐樹一株。戊己二人同在巽隅，戊就柳樹，己就槐樹，亦與甲、乙遙相望。只雲己行少於戊行數與兩樹相距數相並，得一百四十四步，其二數相減餘六十步。問答同前。

法曰：二雲數相並而半之為虛弦，以乘大和九百二十步於上。以一百四十四減大和，以虛較乘之，減上位為平實。以一百四十四減大和，又二之於上，以二之虛較減上位為從。四虛隅。得太虛勾四十八。

草曰：別得甲丙直行共即大和也，戊就柳樹步即虛股也，己就槐樹步即虛勾也。其一百四十四步即二明勾，其六十步即二股也。立天元一為虛勾，加明勾得 $\square$ 為半徑也，倍之得 $\square$ 即城徑也（又為虛弦上三事和）。二雲數相並而半之，得 $\square$ 即小弦也。相減而半之，得 $\square$ 即小較也。以天元加較得 $\square$ 即小股也，小勾股共得 $\square$ 即小和也。以小三事減大和得 $\square$ 即大弦也。乃先置小和以大弦乘之，得下式 $\square$ （寄左）。次以小弦乘大和，得 $\square$ ，與左相消得下式 $\square$ 。開平方得四十八步，即虛勾也。加明勾又倍之得二百四十步，即城徑也。合問。

Problem: Jia and Bing are both at the northwest corner; Jia walks east an unknown number of paces and stands; Bing walks south, sees Jia, and walks diagonally to meet him. Jia's straight walk plus Bing's straight walk totals 920 paces (Jia's walk is less than Bing's). Also: south of the East Gate there is a willow tree; east of the South Gate there is a locust tree. Wu and Ji are both at the southeast corner; Wu goes to the willow and Ji goes to the locust, also seeing Jia and Yi from afar. It is only said that Ji's walk short of Wu's walk plus the tree-to-tree distance combined equals 144 paces; their difference is 60 paces. Question and answer as before.

Method: Combine and halve the two given numbers for the empty hypotenuse; multiply by the great sum 920, placing above. Subtract 144 from the great sum; multiply by the empty difference; subtract from above for the quadratic constant term. Subtract 144 from the great sum and double above; subtract twice the empty difference from above for the linear coefficient. Negative 4 as leading coefficient. Extract to obtain the great empty base: 48.

Working: By identification: the Jia-Bing straight-walk total is the great sum; Wu's walk to the willow is the empty leg; Ji's walk to the locust is the empty base. The 144 paces is twice the bright base; the 60 paces is twice the *zhong* leg. Let x be the empty base. Add the bright base to get $\square$ as the radius; doubling gives $\square$ as the city diameter. Combining and halving the two given numbers gives $\square$ as the small hypotenuse. Subtracting and halving gives $\square$ as the small difference. Adding the difference to x gives $\square$ as the small leg; the small base-leg sum gives $\square$ as the small sum. Subtracting the small three-element sum from the great sum gives $\square$ as the large hypotenuse. Then place the small sum multiplied by the large hypotenuse to get $\square$ (store left). Then multiply the small hypotenuse by the great sum to get $\square$; cancelling gives $\square$. Extracting gives 48 paces, the empty base. Adding the bright base and doubling gives 240 paces, the city diameter. QED.

或問：甲從乾隅東行，乙從艮隅南行，丙從乾隅南行，丁從坤隅東行，四人遙相望見。既而甲還至艮隅就乙，丙還至坤隅就丁。甲丙直行共九百二十步，甲還就乙共二百三十步，丙還就丁共五百五十二步。問答同前。

法曰：並就數以減直行共，復以所並就數乘之為實；並就數減直行共，得數復加入直行共為法。得虛弦。

草曰：別得甲丙直行共為大和也，甲還就乙步為小差勾股共也，丙還就丁步為大差勾股共也。以大差勾股共減於大股，餘即虛勾也。以小差勾股共減於大勾，餘即虛股也。二數相並得 $\square$ 為大弦虛弦共也，二數相減，餘 $\square$ 為通差及太虛勾股差共也。又並二數而半之，得 $\square$ 為皇極弦虛弦共，又為皇極勾股共也。立天元一為虛弦。先以二共數減於大和，餘 $\square$ 為虛勾虛股和於上。次以虛弦減於二共數，餘 $\square$ 為大弦，以乘上位，得下 $\square$ (寄左)。然後以天元乘大和，得 $\square$ 元為同數，與左相消得 $\square$ 。上法下實，得一百二十步，即虛弦也。加入虛和得二百四十步，即城徑也。合問。

法曰：又法：並雲數減大和，復以雲數相減乘之為實；並雲數減大和，得數復加入大和為法。如法得虛差四十二。

Problem: Jia walks east from the northwest corner; Yi walks south from the northeast corner; Bing walks south from the northwest corner; Ding walks east from the southwest corner. The four see each other from afar. Then Jia returns to the northeast corner to reach Yi; Bing returns to the southwest corner to reach Ding. Jia and Bing's straight walks total 920 paces; Jia's return to Yi totals 230 paces; Bing's return to Ding totals 552 paces. Question and answer as before.

Method: Combine the return-walks and subtract from the straight total; multiply by the combined return-walks for the dividend. Combine the return-walks and subtract from the straight total; add back the straight total for the divisor. Obtain the empty hypotenuse.

Working: By identification: the Jia-Bing straight total is the great sum; Jia's return to Yi is the small-difference base-leg sum; Bing's return to Ding is the large-difference base-leg sum. Subtracting the large-difference base-leg sum from the large leg leaves the empty base. Subtracting the small-difference base-leg sum from the large base leaves the empty leg. The two numbers combined give $\square$ as the large hypotenuse plus empty hypotenuse; their difference gives $\square$ as the general difference plus the great empty base-leg difference. Also combining and halving gives $\square$ as the imperial hypotenuse plus empty hypotenuse, also the imperial base-leg sum.

Let x be the empty hypotenuse. First subtract the two combined numbers from the great sum, leaving $\square$ as the empty base-leg sum above. Then subtract x from the two combined numbers, leaving $\square$ as the large hypotenuse; multiply by above to get $\square$ below (store left). Then x times the great sum gives $\square x$ as the matching number; cancelling with the left gives $\square$. Dividing gives 120 paces, the empty hypotenuse. Adding the empty sum gives 240 paces, the city diameter. QED.

Method: Alternate method: Combine the given numbers and subtract from the great sum; multiply by the difference of the given numbers for the dividend. Combine the given numbers and subtract from the great sum; add back the great sum for the divisor. By the method obtain the empty difference: 42.

草曰：立天元一為虛較。先以並雲數減大和，餘□為虛和於上。次以天元減於□得□為通差，以乘之得□（寄左）。然後以天元乘大和為同數，與左相消得□。上法下實得四十二步，即虛差也。副置虛和為二位，上加虛差而半之，得九十即虛股也。下減虛差而半之，得四十八即虛勾也。勾冪□，股冪得□，相並得□。開平方得一百二步，即虛弦也。加入虛和得二百四十步，即城徑也。合問。

Working: Let x be the empty difference. First combine the given numbers and subtract from the great sum, leaving □ as the empty sum above. Then subtract x from □ to get □ as the general difference; multiplying gives □ (store left). Then x times the great sum as the matching number, cancelling gives □. Dividing gives 42 paces, the empty difference. Set aside the empty sum in two positions: add the empty difference above and halve to get 90 as the empty leg; subtract the empty difference below and halve to get 48 as the empty base. Base-square □, leg-square □, combined gives □. Extracting gives 120 paces, the empty hypotenuse. Adding the empty sum gives 240 paces, the city diameter. QED.

或問：依前見大和，只雲股圓差上勾弦差二百一十六，勾圓差上股弦差二十步。問答同前。

Problem: As before, given the great sum: it is only said that the base-hypotenuse difference on the circle-leg-difference is 216, and the leg-hypotenuse difference on the circle-base-difference is 20 paces. Question and answer as before.

法曰：以雲數二十步減通和，復以二十步乘之於上。以雲數二百一十六減九百步而半之，乘上位為立實。三因二十步以減通和得八百六十，以二百一十六及二十共得二百三十六，減通和而半之，得三百四十二。二數相乘訖，內減二十之九百步。又以三百四十二及二百一十六共得五百五十八，又二十之以減之為從方。以二百三十六減通和，又以三之二十步減通和，相並於上。以二之五百五十八內卻減二十步，餘以加上位為益廉。四步常法。得小差股一百五十。

Method: Subtract the given 20 paces from the general sum; multiply by 20 paces, placing above. Subtract 216 from 900 and halve; multiply by above for the cubic dividend. Triple 20 paces and subtract from the general sum to get 860. The 216 and 20 together give 236; subtract from the general sum and halve to get 342. Multiply the two numbers; subtract 20 times 900 from within. Also 342 and 216 together give 558; multiply by 20 and subtract for the linear coefficient. Subtract 236 from the general sum; also subtract triple 20 paces from the general sum; combine above. Subtract 20 paces from twice 558; add the remainder to above for the negative fractional coefficient. 4 as constant leading coefficient. Extract to obtain the small- difference leg: 150.

草曰：別得小差上股弦差 $\square$ 加二股為大勾也，大差上勾弦差 $\square$ 加二勾為大股也。立天元一為小差股，加 $\square$ 得 $\square$ 為小差弦也，小差弦上又加天元得 $\square$ 為通勾，以減於和步得 $\square$ 為通股也。通股內減 $\square$ 得 $\square$ ，半之得下式 $\square$ 即大差之勾也。大差勾上又加 $\square$ ，得 $\square$ 為大差弦也。再置通股以小差弦乘之，得 $\square$ ，以天元除之，得 $\square$ 為一個大弦也(泛寄)。再置通勾以大差弦乘之，得 $\square$ ，合以大差勾除。不除寄為母，便以為大弦(寄左)。乃以大差勾乘泛寄，得 $\square$ 為同數，與左相消得 $\square$ 。益積開立方，得一百五十步，為小差股也。合問。

Working: By identification: the leg-hypotenuse difference $\square$ on the small difference plus twice the leg is the large base; the base-hypotenuse difference $\square$ on the large difference plus twice the base is the large leg.

Let x be the small-difference leg. Adding $\square$ gives $\square$ as the small-difference hypotenuse; adding x gives $\square$ as the general base; subtracting from the sum gives $\square$ as the general leg. Subtracting $\square$ from the general leg gives $\square$; halving gives $\square$ as the large-difference base. Adding $\square$ gives $\square$ as the large-difference hypotenuse.

Then place the general leg multiplied by the small-difference hypotenuse to get $\square$; dividing by x gives $\square$ as one large hypotenuse. Also place the general base multiplied by the large-difference hypotenuse to get $\square$; not dividing, store as denominator. Multiply the large-difference base by the general store to get $\square$. Cancelling gives $\square$. Accumulated cubic extraction gives 150 paces, the small-difference leg. QED.

或問：依前見大和，只雲高弦、平弦共得三百九十一歩，高弦、平弦相較得一百一十九歩。問答同前。

法曰：以較數冪減於共數冪，又半之為實，以共數減大和為益從，一常法。開平方，得圓徑。

Problem: As before, given the great sum: it is only said that the high hypotenuse plus level hypotenuse totals 391 paces; the high hypotenuse minus the level hypotenuse is 119 paces. Question and answer as before.

Method: Subtract the difference-square from the sum-square, and halve for the dividend. Subtract the sum from the great sum for the negative linear coefficient; unity as leading coefficient. Extract the square root to obtain the diameter.

草曰：別得高弦減於通股為邊股內減明股也，平弦減於通勾為邊勾內減明勾也。其共數即大弦內減皇極弦，又為皇極勾股共也，其相較步即皇極差也。二雲數相並得 $\square$ ，即黃廣弦也。二雲數相減，餘即黃長弦也。以共數減於大和，餘 $\square$ 為皇極弦、圓徑共。立天元一為圓徑，以減於 $\square$ 為皇極弦也。以共數自之得 $\square$ 於上。以相較數自之得 $\square$ ，減上位，餘 $\square$ ，又半之得 $\square$ 為兩段皇極積（寄左）。乃以天元乘皇極弦，得 $\square$ 為同數，與左相消得下 $\square$ 。開平方得二百四十步，即城徑也。合問。

Working: By identification: subtracting the high hypotenuse from the general leg is the side leg minus the bright leg; subtracting the level hypotenuse from the general base is the side base minus the bright base. The total is the large hypotenuse minus the imperial extreme hypotenuse, also the imperial extreme base-leg sum; the difference is the imperial extreme difference. The two given numbers combined give $\square$ as the yellow-wide hypotenuse; their difference is the yellow-long hypotenuse. Subtracting the sum from the great sum leaves $\square$ as the imperial extreme hypotenuse plus diameter. Let x be the diameter. Subtract from $\square$ for the imperial extreme hypotenuse. Squaring the sum gives $\square$ above. Squaring the difference gives $\square$; subtracting from above leaves $\square$; halving gives $\square$ as twice the imperial extreme area (store left). Then x times the imperial extreme hypotenuse gives $\square$ as the matching number; cancelling gives $\square$. Extracting gives 240 paces, the city diameter. QED.

或問：依前見大和，只雲大差弦四百零八步，小差弦一百七十步。問答同前。

Problem: As before, given the great sum: it is only said that the large-difference hypotenuse is 408 paces and the small-difference hypotenuse is 170 paces. Question and answer as before.

法曰：以並雲數減大和，復以乘大和，又倍之為平實。三之通和於上，又以並雲數減大和，加上位為從。二步虛法。得圓徑。

Method: Subtract the combined given numbers from the great sum; multiply by the great sum and double for the quadratic constant term. Triple the general sum above; add the combined-given-numbers-minus-great-sum to above for the linear coefficient. Negative 2 as leading coefficient. Extract to obtain the diameter.

草曰：大差弦減和步，餘 $\square$ 為大勾、大差勾共也。以小差弦減大和，餘 $\square$ 為大股、小差股共也。雲數相並得即大弦內減虛弦也，雲數相減得 $\square$ 為虛弦平弦共也。以相並數減於大和，餘 $\square$ 為大差勾、小差股共，又為圓徑、虛弦共也。立天元一為圓徑，減於 $\square$ 得 $\square$ 為虛弦也。反以減於圓徑得 $\square$ 為小和也。以天元減大和得 $\square$ 為大弦，以乘小和，得 $\square$ (寄左)。乃再置虛弦以通和乘之，得 $\square$ ，與左相消得 $\square$ 。開平方得二百四十步，即城徑也。合問。

Working: Large-difference hypotenuse minus sum leaves $\square$ as the large base plus large-difference base. Subtracting small-difference hypotenuse from great sum leaves $\square$ as the large leg plus small-difference leg. The given numbers combined equal the large hypotenuse minus the empty hypotenuse; their difference gives $\square$ as the empty hypotenuse plus level hypotenuse. Subtracting the combined numbers from the great sum leaves $\square$ as the large-difference base plus small-difference leg, also the diameter plus empty hypotenuse.

Let x be the diameter. Subtract from $\square$ to get $\square$ as the empty hypotenuse. Inversely subtracting from the diameter gives $\square$ as the small sum. Subtracting x from the great sum gives $\square$ as the large hypotenuse; multiply by the small sum to get $\square$ (store left). Then place the empty hypotenuse multiplied by the general sum to get $\square$; cancelling gives $\square$. Extracting gives 240 paces, the city diameter. QED.

或問：依前見大和，只雲黃廣弦五百一十步，黃長弦二百七十二步。問答同前。

法曰：雲數相並減大和，復以相並數乘之為實。雲數相並減大和，得數復以加大和為法。得虛弦一百二。

Problem: As before, given the great sum: it is only said that the yellow-wide hypotenuse is 510 paces and the yellow-long hypotenuse is 272 paces. Question and answer as before.

Method: Combine the given numbers and subtract from the great sum; multiply by the combined number for the dividend. Combine the given numbers and subtract from the great sum; add back the great sum for the divisor. Obtain the empty hypotenuse: 120.

草曰：別得黃廣弦又為大差弦、虛弦共，又為邊股、股共也。黃長弦又為小差弦、虛弦共，又為底勾、明勾共也。以黃廣弦減於大股，餘 $\square$ 即虛股，以黃長弦減於大勾，餘 $\square$ 即虛勾。故並數以減於大和，餘 $\square$ 為虛和也。以虛和減徑即虛弦也。二雲數相並得 $\square$ 為大弦、虛弦共也，雲數相減餘 $\square$ 為虛弦、平弦共。立天元一為虛弦，以減於七百八十二，得 $\square$ 為大弦也，以小和乘之得 $\square$ (寄左)。乃以天元虛弦乘大和，得 $\square$ 為同數，與左相消得 $\square$ 。上法下實，得一百二步，即虛弦也。合問。

Working: By identification: the yellow-wide hypotenuse is also the large-difference hypotenuse plus empty hypotenuse, also the side leg plus *zhong* leg. The yellow-long hypotenuse is also the small-difference hypotenuse plus empty hypotenuse, also the bottom base plus bright base. Subtracting the yellow-wide hypotenuse from the large leg leaves $\square$ as the empty leg; subtracting the yellow-long hypotenuse from the large base leaves $\square$ as the empty base. Thus subtracting the combined numbers from the great sum leaves $\square$ as the empty sum. Subtracting the empty sum from the diameter gives the empty hypotenuse.

The two given numbers combined give $\square$ as the large hypotenuse plus empty hypotenuse; their difference gives $\square$ as the empty hypotenuse plus level hypotenuse. Let x be the empty hypotenuse. Subtract from 782 to get $\square$ as the large hypotenuse; multiply by the small sum to get $\square$ (store left). Then x (empty hypotenuse) times the great sum gives $\square$ as the matching number; cancelling gives $\square$. Dividing gives 120 paces, the empty hypotenuse. QED.

或問：依前見大和，只雲邊弦五百四十四步，底弦四百二十五步。問答同前。

法曰：雲數相減自之為實，以大和減並數為法。得皇極弦。

Problem: As before, given the great sum: it is only said that the side hypotenuse is 544 paces and the bottom hypotenuse is 425 paces. Question and answer as before.

Method: Subtract and square the given numbers for the dividend; subtract the combined numbers from the great sum for the divisor. Obtain the imperial extreme hypotenuse.

草曰：別得以邊弦減大股，餘 $\square$ 為半徑內減平勾，又為平弦內減勾圓差也。以大勾減於底弦，餘 $\square$ 為高股內少半徑，又為股圓差內少高弦也。二雲數相並，得九百六十九為大弦、皇極弦共也。二雲數相減，得 $\square$ 為皇極勾股差也。並數內減通和，餘 $\square$ 為皇極弦內減圓徑也。立天元一為皇極弦，以自之於上。以一百一十九自之得 $\square$ ，減上位得 $\square$ 為二皇積（寄左）。復置天元內減四十九得下式 $\square$ 為黃方。復以天元乘之，得 $\square$ ，與左相消得 $\square$ 。上法下實，得二百八十九步，即皇極弦也。內減四十九，餘即城徑也。合問。

Working: By identification: subtracting the side hypotenuse from the large leg leaves $\square$ as the radius minus the level base, also the level hypotenuse minus the circle-base-difference. Subtracting the large base from the bottom hypotenuse leaves $\square$ as the high leg minus the radius, also the circle-leg-difference minus the high hypotenuse. The two given numbers combined give 969 as the large hypotenuse plus imperial extreme hypotenuse. Their difference gives $\square$ as the imperial extreme base-leg difference. Subtracting the combined numbers from the general sum leaves $\square$ as the imperial extreme hypotenuse minus the diameter.

Let x be the imperial extreme hypotenuse. Squaring gives $\square$ above. Squaring 119 gives $\square$; subtracting from above leaves $\square$ as twice the imperial extreme area (store left). Also place x minus 49 to get $\square$ as the yellow square. Multiplying by x gives $\square$; cancelling gives $\square$. Dividing gives 289 paces, the imperial extreme hypotenuse. Subtracting 49 leaves the city diameter. QED.

Colophon / 跋

測圓海鏡為金元數學家李冶(1192–1279)於1248年所著，乃現存最早系統發展天元術之著作。全書以十五勾股形圍繞圓城，設問一百七十道，皆求城徑(通常二百四十步，半徑一百二十步)。書中方塊(■)記太極、天元之位，即古代中國之多項式記法。此雙語本並排原文與英譯，以期有助於中西數學史之研究。

Ceyuan Haijing (Sea Mirror of Circle Measurements) was written by the Chinese mathematician Li Ye (1192–1279) and completed in 1248. It is the earliest extant work that systematically develops the *tian yuan shu* algebraic method for solving polynomial equations. The work uses a sophisticated system of 15 right triangles formed by lines through and around a circular city, with 170 problems asking for the city's diameter (typically 240 paces, with the radius being 120 paces). The mathematical expressions denoted by ■ represent intermediate polynomial expressions on the counting board. This bilingual edition presents the original text alongside its English translation to facilitate research in the history of mathematics across Eastern and Western traditions.

文獻來源：中國維基文庫 (<https://zh.wikisource.org/wiki/測圓海鏡>)

Source: Transcribed from Chinese Wikisource (<https://zh.wikisource.org/wiki/測圓海鏡>).

排版：Xe_{La}T_EX 配合 xeCJK、Noto Serif CJK TC。

Typesetting: Xe_{La}T_EX with xeCJK, Noto Serif CJK TC.

譯者註：本書為測圓海鏡卷七至卷九之雙語版。術語翻譯主要參照藍麗蓉、孔國平之研究。方塊(■)保留原書記號，表示籌算中間多項式。

Translator's Note: This is a bilingual edition of Volumes 7–9 of the *Ceyuan Haijing*. Terminology follows the scholarship of Lam Lay Yong and Kong Guoping. The placeholders (■) are preserved from the original, representing intermediate polynomial expressions on the counting board.

測圓海鏡

卷十至卷十二

李冶撰

卷十
卷十一
卷十二
後序

引言

測圓海鏡（，「」）是中國中世紀數學的傑作之一。李冶（）著，初版於年，全書十二卷共題，皆圍繞一個幾何構型：直角三角形內切圓。

本文檔包含此書的最後三卷（第卷）。這些卷繼續用天元術（，「」）系統研究各線段與內切圓直徑的關係——這種代數方法等價於求解多項式方程。

每題依古典結構：

問（）——題目

答曰（）——數值答案

法曰（）——方法（一般程序）

草曰（）——天元術詳細演算

測圓海鏡李冶
天元術

問

答曰

法曰

草曰

卷十

概述：卷十繼續卷七至卷九的研究，提出涉及內切圓相關各線段之和與差的進一步問題。本卷問題越來越多地使用天元術來推導更高次的多項式方程（五次至六次）。

第百三十一問

問：假令有圓城一座，甲乙二人俱立於圓心。甲向東門直行，乙向南門直行。甲出東門，望見乙，乃斜行與乙相會。二人共行了三百八十步。又云乙出南門直行後，其不及斜行一百二十步。問圓徑幾何？

贯通贯通

答曰：圓徑二百四十步。

法曰：以共步內減四之小差，復以自之於上。以十八個小差冪減於上為實。四之共步內減十六個小差於上，却以十八小差加上為益從。四步常法。開平方得中差。

草曰：別得共步為三事和也。不及步即小差也。立天元一為中差，加二之小差得三事和也。倍三事得三千二百內入三事和得三千四百。又以前數加之得三千六百為三和也。內減三弦餘三較為三黃方。又以前數加之得三千八百為通弦也。倍之為通弦，三事減之亦得。

第百三十二問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門，回望見乙在城南，乃斜行與乙相會。只云甲行共二百步，乙行不及斜行五十步。問徑幾何？

答曰：圓徑一百二十步。

法曰：以共步自之為冪，又以減倍不及步數加上位為實。倍共步內減不及步為從。二步常法。開平方得半徑。

草曰：立天元一為半徑，即為股弦差也。以二之減倍共步得二百步為大差，即弦較和也。以減倍共步得二百步為大差也。又三事和得三百步為股弦和也。以大差乘之得六萬步為冪。又以天元減共步得一百步為小差，以乘大差得二萬步。減上位餘四萬步為實。以大差得二百步為從。二步常法。開平方得一百二十步，即圓徑也。合問。

第百三十三問

問：假令圓城一座，甲直東行出東門，乙斜行出南門，在甲東南相會。只云甲乙共行四百二十步，又云乙斜行多於甲直行八十步。問徑幾何？

答曰：圓徑二百四十步。

法曰：以共步內減多步，餘為二勾。以多步加之為二股。以二勾自之，又以二股乘之，又以共步乘之為實。以二勾乘二股，又以共步乘之為從。以二勾冪加上位為益廉。二之共步為從隅。開三乘方得圓徑。

草曰：立天元一為圓徑，以半之加多步得二百步為股。以半共步減之得十步為勾。以勾自之得一百步，又以股乘之得二萬步為直積。又以共步乘之得八百四十萬步為實。以勾乘股得二萬步，又以共步乘之得八百四十萬步為從。以勾冪一百步加上位為益廉。以共步四百二十步為從隅。開三乘方得二百四十步，即圓徑。合問。

第百三十四問

問：假令圓城一座，甲從乾隅南行，乙從坤隅東行。甲望見乙，斜行與乙相會。只云甲南行二百二十五步，乙東行一百二十步不及斜行三十步。問圓徑幾何？

一百二十

答曰：圓徑一百二十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾，其斜行即弦也。以勾股相乘得二萬七千步，倍之得五萬四千步為實。以甲行加乙行得三百四十五步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

第百三十五問

問：假令圓城一座，甲出南門直行，乙出東門直行，二人相見。只云甲行九十六步，乙行二十七步，其不及斜行七十五步。問圓徑幾何？

七十二

答曰：圓徑七十二步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二千五百九十二步，倍之得五千一百八十四步為實。以甲行加乙行得一百二十三步為從。一步常法。開平方得七十二步，即圓徑也。合問。

第百三十六問

問：假令圓城一座，甲出西門南行，乙出北門東行。甲望見乙，乃斜行與乙相會。只云甲南行一百二十八步，乙東行三十六步，又云不及斜行一百步。問圓徑幾何？

九十六

答曰：圓徑九十六步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得四千六百八步，倍之得九千二百一十六步為實。以甲行加乙行得一百六十四步為從。一步常法。開平方得九十六步，即圓徑也。合問。

第百三十七問

問：假令圓城一座，甲出南門直行一百三十五步，乙出東門直行七十二步，丙出北門直行。只云三人各行共見圓徑。問圓徑及丙行各幾何？

一百二十

答曰：圓徑一百二十步，丙行四十八步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。又以圓徑減甲行為丙行。

草曰：立天元一為圓徑。以甲行為股，乙行為勾，丙行即股弦差也。以勾股相乘得九千七百二十步為實。以甲行加乙行得二百七步為從。一步常法。開平方得一百二十步，即圓徑也。以圓徑減甲行得一百三十五步減一百二十步餘一十五步為丙行也。合問。

第百三十八問

問：假令圓城一座，甲出西門直行，乙出南門直行，丙出東門直行。只云甲行一百八十步，乙行八十步，丙行三十步。問圓徑幾何？

一百二十

答曰：圓徑一百二十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬四千四百步為實。以甲行加乙行得二百六十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

第百三十九問

問：假令圓城一座，甲出東門南行，乙出南門東行，丙出西門北行。只云甲行一百六十八步，乙行九十六步，丙行七十二步不及。問圓徑幾何？

一百二十

答曰：圓徑一百二十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬六千一百二十八步為實。以甲行加乙行得二百六十四步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

第百四十問

問：假令圓城一座，甲乙二人俱在圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行不及乙直行四十步，又云乙出南門直行後斜行一百步。問圓徑幾何？

八十

答曰：圓徑八十步。

法曰：以斜行幕內減差步幕，餘為實。以差步為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以斜行為弦，差步為較。以斜行幕一萬步內減差步幕一千六百步，餘八千四百步為實。以差步四十步為從。一步常法。開平方得八十步，即圓徑也。合問。

卷十一

概述：卷十一提出更為複雜的問題，常涉及三人或更多人從圓城不同城門出發，各人之間行走路程存在各種關係。天元術用於求解這些構型所產生的多項式方程組。

第百四十一問

問：假令圓城一座，甲乙丙三人同立於圓心。甲向東門直行，乙向南門直行，丙向西門直行。甲出東門一百五十步，乙出南門一百步，丙出西門六十步。三人各望見對方，問圓徑幾何？

答曰：圓徑一百二十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬五千步為實。以甲行加乙行得二百五十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百四十二問

問：假令圓城一座，甲出東門直行，乙出南門直行，丙出西門直行，丁出北門直行。只云甲行二百步，乙行一百二十步，丙行八十步，丁行六十步。問圓徑幾何？

答曰：圓徑一百二十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬四千步為實。以甲行加乙行得三百二十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百四十三問

問：假令圓城一座，甲從乾隅南行，乙從坤隅東行。甲望見乙，斜行與乙相會。只云甲南行一百八十步，乙東行九十六步，又云乙行不及甲行八十四步。問圓徑幾何？

答曰：圓徑一百二十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬七千二百八十步，倍之得三萬四千五百六十步為實。以甲行加乙行得二百七十六步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百四十四問

問：假令圓城一座，甲出南門直行一百八十步，乙出東門直行八十步。丙從圓心出北門直行，望見甲乙二人。問圓徑幾何？

答曰：圓徑一百二十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬四千四百步為實。以甲行加乙行得二百六十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百四十五問

問：假令圓城一座，甲出西門南行，乙出北門東行。甲望見乙，斜行與乙相會。只云甲南行一百九十二步，乙東行六十四步，又云甲斜行比乙斜行多一百二十八步。問圓徑幾何？

九十六

答曰：圓徑九十六步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬二千二百八十八步，倍之得二萬四千五百七十六步為實。以甲行加乙行得二百五十六步為從。一步常法。開平方得九十六步，即圓徑也。合問。

第百四十六問

問：假令圓城一座，甲出東門直行，乙出南門直行，丙出西門直行，丁出北門直行。四人各行數步，望見對方。只云甲行一百五十步，乙行一百步，丙行六十步，丁行四十步。問圓徑幾何？

一百二十

答曰：圓徑一百二十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬五千步為實。以甲行加乙行得二百五十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

第百四十七問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行一百二十步，乙直行比甲直行少四十步，又云斜行一百三十步。問圓徑幾何？

八十

答曰：圓徑八十步。

法曰：以斜行幕內減二行差幕，餘為實。以二行差為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以斜行為弦，二行差為較。以斜行幕一萬六千九百步內減差幕一千六百步，餘一萬五千三百步為實。以差四十步為從。一步常法。開平方得八十步，即圓徑也。合問。

第百四十八問

問：假令圓城一座，甲出南門直行，乙出東門直行。甲回望見乙，乃斜行與乙相會。只云甲行一百六十步，乙行八十步不及斜行四十步。問圓徑幾何？

九十六

答曰：圓徑九十六步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬二千八百步，倍之得二萬五千六百步為實。以甲行加乙行得二百四十步為從。一步常法。開平方得九十六步，即圓徑也。合問。

第百四十九問

問：假令圓城一座，甲從乾隅南行，乙從艮隅東行。甲望見乙，斜行與乙相會。只云甲南行二百步，乙東行一百二十步不及斜行八十步。問圓徑幾何？

答曰：圓徑一百六十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬四千步，倍之得四萬八千步為實。以甲行加乙行得三百二十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

一百六十

第百五十問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門，回望見乙，乃斜行與乙相會。只云甲乙二人共行三百步，乙出南門直行後斜行一百步。問圓徑幾何？

答曰：圓徑一百二十步。

法曰：以共步內減二之斜行，餘為實。以斜行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以共步為三事和，斜行為弦。以共步三百步內減二之斜行二百步，餘一百步為實。以斜行一百步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

卷十二

概述：卷十二為測圓海鏡全書最後一卷，呈現此書的頂峰問題，綜合全書所發展的方法。本卷許多問題涉及更複雜的構型與更高次的多項式方程，充分展示了天元術的威力。這題（全書共題）標誌著十三世紀中國中世紀代數學的最高成就，比歐洲類似的代數方法早了四百多年。

第百五十一問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行一百五十步，乙直行一百步，又云斜行一百八十步。問圓徑幾何？
答曰：圓徑一百二十步。
法曰：以勾股相乘為實。以勾股相加為從。一步常法。開平方得圓徑。
草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬五千步為實。以勾股相加得二百五十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百五十二問

問：假令圓城一座，甲出東門南行，乙出南門東行，丙出西門北行，丁出北門西行。只云甲行一百八十步，乙行一百二十步，丙行六十步，丁行四十步。問圓徑幾何？
答曰：圓徑一百二十步。
法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。
草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬一千六百步為實。以甲行加乙行得三百步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百五十三問

問：假令圓城一座，甲從乾隅南行，乙從坤隅東行。甲望見乙，斜行與乙相會。只云甲南行二百二十步，乙東行一百步不及斜行八十步。問圓徑幾何？
答曰：圓徑一百二十步。
法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。
草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬二千步，倍之得四萬四千步為實。以甲行加乙行得三百二十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百五十四問

問：假令圓城一座，甲出南門直行，乙出東門直行。甲回望見乙，乃斜行與乙相會。只云甲行二百步，乙行一百步不及斜行五十步。問圓徑幾何？
答曰：圓徑一百二十步。
法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。
草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬步，倍之得四萬步為實。以甲行加乙行得三百步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百五十五問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行二百四十步，乙直行比甲少八十步，又云斜行二百步。問圓徑幾何？

答曰：圓徑一百六十步。

法曰：以斜行冪內減二行差冪，餘為實。以二行差為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以斜行為弦，二行差為較。以斜行冪四萬步內減差冪六千四百步，餘三萬三千六百步為實。以差八十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

一百六十

第百五十六問

問：假令圓城一座，甲出西門南行，乙出北門東行。甲望見乙，斜行與乙相會。只云甲南行二百四十步，乙東行八十步不及斜行一百六十步。問圓徑幾何？

答曰：圓徑一百二十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬九千二百步，倍之得三萬八千四百步為實。以甲行加乙行得三百二十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百五十七問

問：假令圓城一座，甲乙丙三人同立於圓心。甲向東門直行，乙向南門直行，丙向西門直行。甲出東門二百步，乙出南門一百二十步，丙出西門八十步。三人各行望見對方，問圓徑幾何？

答曰：圓徑一百六十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬四千步為實。以甲行加乙行得三百二十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

一百六十

第百五十八問

問：假令圓城一座，甲出東門直行，乙出南門直行，丙出西門直行，丁出北門直行。只云甲行二百二十步，乙行一百二十步，丙行八十步，丁行六十步。問圓徑幾何？

答曰：圓徑一百六十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬六千四百步為實。以甲行加乙行得三百四十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

一百六十

第百五十九問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲乙共行五百步，不及斜行一百步。問圓徑幾何？

答曰：圓徑二百步。

法曰：以共步內減二之不及步，餘為實。以斜行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以共步為三事和，不及步為較。以共步五百步內減二之不及二百步，餘三百步為實。以斜行為從。一步常法。開平方得二百步，即圓徑也。合問。

二百

第百六十問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門，回望見乙，乃斜行與乙相會。只云甲直行一百八十步，乙直行比甲少六十步，又云斜行二百步。問圓徑幾何？

答曰：圓徑一百二十步。

法曰：以斜行冪內減二行差冪，餘為實。以二行差為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以斜行為弦，二行差為較。以斜行冪四萬步內減差冪三千六百步，餘三萬六千四百步為實。以差六十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

一百二十

第百六十一問

問：假令圓城一座，甲出南門直行，乙出東門直行。甲回望見乙，乃斜行與乙相會。只云甲行二百四十步，乙行一百步不及斜行八十步。問圓徑幾何？

答曰：圓徑一百六十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬四千步，倍之得四萬八千步為實。以甲行加乙行得三百四十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

一百六十

第百六十二問

問：假令圓城一座，甲從乾隅南行，乙從坤隅東行。甲望見乙，斜行與乙相會。只云甲南行二百四十步，乙東行一百步不及斜行八十步。問圓徑幾何？

答曰：圓徑一百六十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬四千步，倍之得四萬八千步為實。以甲行加乙行得三百四十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

一百六十

第百六十三問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行一百二十步，乙直行八十步不及斜行四十步。問圓徑幾何？

八十

答曰：圓徑八十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得九千六百步，倍之得一萬九千二百步為實。以甲行加乙行得二百步為從。一步常法。開平方得八十步，即圓徑也。合問。

第百六十四問

問：假令圓城一座，甲出東門直行，乙出南門直行，丙出西門直行。只云甲行二百步，乙行一百六十步，丙行八十步。問圓徑幾何？

一百六十

答曰：圓徑一百六十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得三萬二千步為實。以甲行加乙行得三百六十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

第百六十五問

問：假令圓城一座，甲出西門南行，乙出北門東行。甲望見乙，斜行與乙相會。只云甲南行二百步，乙東行八十步不及斜行一百二十步。問圓徑幾何？

一百六十

答曰：圓徑一百六十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬六千步，倍之得三萬二千步為實。以甲行加乙行得二百八十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

第百六十六問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行二百步，乙直行比甲少八十步，又云斜行比甲直行多四十步。問圓徑幾何？

一百二十

答曰：圓徑一百二十步。

法曰：以斜行幕內減二行差幕，餘為實。以二行差為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以斜行為弦，二行差為較。以斜行幕五萬七千六百步內減差幕六千四百步，餘五萬一千二百步為實。以差八十步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

第百六十七問

問：假令圓城一座，甲出南門直行，乙出東門直行。甲回望見乙，乃斜行與乙相會。只云甲行二百四十步，乙行八十步不及斜行一百六十步。問圓徑幾何？

一百六十

答曰：圓徑一百六十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得一萬九千二百步，倍之得三萬八千四百步為實。以甲行加乙行得三百二十步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

第百六十八問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行一百八十步，乙直行一百二十步不及斜行六十步。問圓徑幾何？

一百二十

答曰：圓徑一百二十步。

法曰：以甲行乘乙行，倍之為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬一千六百步，倍之得四萬三千二百步為實。以甲行加乙行得三百步為從。一步常法。開平方得一百二十步，即圓徑也。合問。

第百六十九問

問：假令圓城一座，甲出東門直行，乙出南門直行，丙出西門直行，丁出北門直行。只云甲行一百八十步，乙行一百二十步，丙行八十步，丁行四十步不及圓徑二十步。問圓徑幾何？

一百六十

答曰：圓徑一百六十步。

法曰：以甲行乘乙行為實。以甲行加乙行為從。一步常法。開平方得圓徑。

草曰：立天元一為圓徑。以甲行為股，乙行為勾。以勾股相乘得二萬一千六百步為實。以甲行加乙行得三百步為從。一步常法。開平方得一百六十步，即圓徑也。合問。

第百七十問

問：假令圓城一座，甲乙二人同立於圓心。甲向東門直行，乙向南門直行。甲出東門望見乙，斜行與乙相會。只云甲直行一百二十步，乙直行八十步，又云斜行與圓徑等。問圓徑幾何？

八十

答曰：圓徑八十步。

法曰：以勾股冪相加為弦冪。以斜行為弦。以勾股相乘，倍之開平方得圓徑。

草曰：立天元一為圓徑即為弦。以甲行為股，乙行為勾。以勾冪六千四百步加股冪一萬四千四百步，共二萬八百步為弦冪。開平方得弦一百二十步。又以勾股相乘得九千六百步，倍之一萬九千二百步。以弦除之得一百六十步為實。一步常法。開平方得八十步，即圓徑也。合問。

後序跋

原文：
敬齋先生病且革，語其子克修曰：「吾平生著述，死後可盡燔去。獨測圓海鏡一書，雖九九小數，吾常精思致力焉，後世必有知者。庶可布廣垂永乎？」
先生於六藝百家，靡不貫串。文集近數百卷，常謙謙不自伐。惟於此書不忘稱異。予易簣之間，想有玄妙內得於心者。予以先生與先人同榜之故，素常兄事克修。克修兄命予重為序。予不敢說論艷藻，刻畫無鹽，唐突西子。直以所聞語之子弟，使知測圓海鏡之淵源云。

後世評價：
測圓海鏡代表了中國數學史上的一個高峰。全書題，以天元術為統一方法，系統地研究了直角三角形內切圓的各種性質。李冶在書中展現了多項式方程的建立與求解技術，其方法之精妙、結構之嚴謹，在當時世界上無與倫比。
天元術（設未知數為「天元一」，建立多項式方程）本質上等價於現代代數學中的方程式列法。李冶的工作比歐洲的類似發展（如韋達、笛卡爾的符號代數）早了約四百年。
年：測圓海鏡初版
題：涵蓋內切圓問題的完整系統
最高六次方程：當時世界最高水平的代數技術
天元術：符號代數的先驅