

Projective cohomology and smooth affine models

GPT-6.1 Sol (OpenAI)
Codex, Ultra setting

October 2026

Written and self-checked by the writing AI. No human or independent review is claimed. Original contributions are dedicated under CC0 1.0. Attributed Stacks adaptations retain their GNU Free Documentation License obligations, as recorded for each component. The cumulative collection includes the complete version 1.2 licence below.

Online course: <https://kokunoyumeto.github.io/open-math-courses/courses/AG-RG/>.

Contents

Projective cohomology and smooth affine models	1
1. Earlier programme proofs used in this lesson	1
2. Formal functions on a projective scheme	4
3. Smoothness and line bundles at a finite stage	6
4. Geometrically connected smooth affine models	8
5. A universal finite complex for a line bundle	13
6. The smooth curve regular immersion calculation	15
7. Applying the proof route in the group lessons	21
Freely readable sources and history	22
History	22
GNU Free Documentation License, Version 1.2	23

Projective cohomology and smooth affine models

This supporting lesson proves the projective formal functions and smooth model statements needed before the reductive group lessons. It also records the precise earlier programme proofs used for completion, projective finiteness and curve duality. It follows the supporting lesson [Algebra and sheaf cohomology before reductive groups](#), whose Sections 1–9 supply the affine, homological, projective-space and Serre calculations. Both supporting lessons precede AG-RG-01. All mathematical sources listed here are freely readable. Source citations identify the material used to write the proofs; they do not replace any proof below.

The formal functions exposition adapts the freely readable Stacks Project proofs. The smooth affine component argument below specializes the Stacks model route and supplies its intermediate arguments. Permission is granted to copy, distribute and modify this lesson under the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation, with no Invariant Sections, no Front-Cover Texts and no Back-Cover Texts. The complete license is included in [the complete GNU FDL 1.2](#). The original source notice, Copyright (C) 2005–2025 Johan de Jong, is retained in the cached introduction. The Stacks Project contributors retain their source rights. Original finite complex and curve bridge passages previously released as CC0 remain CC0; inclusion here does not remove either source’s rights.

1. Earlier programme proofs used in this lesson

The following references are to proofs already written in programme lessons. Their numbered statements delimit exactly what is used.

- **AG-CA-03, Noetherian and Artinian rings:** Proposition 1.2 (finite modules over a Noetherian ring are Noetherian), Theorem 2.1 (Hilbert basis), Theorem 5.1 Artin–Rees, and Theorem 6.1 (Krull intersection). In particular Artin–Rees is proved by finite generation of the graded submodule of the Rees module; it is not taken from a book.
- **AG-CA-19, Completion:** Theorem 1.2 (exact inverse limits with a Mittag–Leffler kernel), Lemma 2.1 (completion preserves surjections with the induced kernel filtration), Proposition 2.2 (finite ideal powers in a completion), Theorems 3.1–3.3 (finite-module exactness, completion tensor comparison, flatness, faithful flatness and Noetherianity). These statements have full proofs there. Thus a finite module over an already complete Noetherian ring is complete, and local Noetherian completion is faithfully flat.
- **AG-CA-17, Formally smooth, unramified and étale ring maps:** Theorem 3.1 (formal smoothness is the split conormal criterion), Theorems 4.1 and 5.1 (standard smooth presentations and local standard form), and Lemma 6.2 (unique lifting of idempotents across a square-zero ideal). These are written lifting and matrix proofs. Lemma 1.1 extends idempotent lifting to a nilpotent ideal by successive square-zero steps.
- **AG-CA-18, Smooth algebras over a field and the Jacobian criterion:** Theorem 2.1 proves regularity of smooth field-algebra local rings. Its proof uses standard smooth

equations and the regular parameter quotient theorem. Base change of smoothness, proved in AG-CA-17, consequently gives geometric regularity.

- **AG-CA-14, Regular local rings:** Theorem 1.1 and Proposition 1.4 prove regular parameter quotients and generation of a regular quotient’s ideal by part of a regular system of parameters. The graded and regular-sequence assertions used there are the full proof of Theorem 6.1 in **Regular sequences, depth and Cohen–Macaulay modules**. The Koszul resolution and its Hom calculation are the corresponding earlier depth lesson proofs.
- **AG-CA-07, Tor and flat modules:** Theorems 5.3 and 6.1 contain the finite-projective and flat-kernel calculations corresponding to Section 5. That section writes its universal-tail flatness and global splitting arguments directly, without taking the Tor criterion as an additional prerequisite.
- **AG-CA-05, Integral extensions:** Theorems 3.2–3.4 prove lying over and incomparability. Section 6 below contains the complete DVR argument in Lemma 6.C and the separable trace-lattice, finite normalization, function-field rational-map spreading and dense equalizer arguments in Lemma 6.4. The homogeneous-coordinate extension over every DVR is written there as well.
- **AG-MO-03, Limits of schemes and Noetherian approximation:** Theorem 1.1, Lemma 2.1 and Theorems 4.1–4.2 prove affine limits, retention of finite open covers, finite-presentation descent, eventual equality of maps and descent of closed immersions. Only the proved affine approximation Theorem 5.1 is needed here. The external-proof-only absolute approximation Theorem 5.2 is not used.
- **AG-MO-06, Quasi-finite morphisms and Chevalley’s theorem:** Theorem 3.1 proves the Noetherian constructibility criterion and Theorem 4.2 proves Chevalley’s theorem. The argument in Section 4 below uses these particular proved statements.
- **AG-FSE, Flat morphisms:** Lemma 5.1 and Corollary 5.2 prove generic freeness over a domain, including for a finite module over a finite type algebra. No openness-of-flatness theorem is needed in Section 4.
- **AG-CA-18, Smooth algebras over a field and the Jacobian criterion:** Theorem 6.1 and Corollary 6.2 prove smooth and étale flatness over arbitrary bases. Together with AG-CA-17 Theorem 5.1 and Lemma 7.1 these give the separating generic coordinates used below.
- **AG-RG-S01, Sections 4–6:** the finite ordered Čech comparison, affine quasi-coherent vanishing and flat base change are proved before this lesson. A finite affine cover of a separated scheme therefore computes quasi-coherent cohomology by its finite section complex. The same written complex calculation treats finite and arbitrary direct sums and flat localization.
- **AG-RG-S01, Sections 7–8:** Theorems 7.2–7.3 compute every projective-space twist and the perfect monomial pairing. Proposition 8.2 and Theorem 8.4 prove finite quotients by twists, finite coherent cohomology and eventual vanishing, including the closed-immersion passage to projective schemes. Lemmas 8.1 and 8.3 give denominator extension and the ample-power immersion; Corollary 7.B makes it closed for a proper source. Theorem 8.5 treats every sufficiently large power, as applied in Section 7 below.
- **AG-QC-14, Ext sheaves and Serre duality on projective space:** Proposition 1.1, Proposition 1.2, Proposition 2.1 and Theorem 5.1 give the local Ext calculation, vector-bundle tensor-Hom comparison, local-to-global Ext spectral sequence and ambient projective Serre duality. The duality proof uses the monomial trace, a two-stage twist presentation and explicit dimension shifting. Its reference to “proper finiteness” in Theorem 4.1 is supplied here by the full projective finiteness proof in [Algebra and sheaf cohomology before reductive groups](#), Section 8 (AG-QC-05 Theorem 2.1), without using the general proper-finiteness provider. Section 6 below also spells out the single-row local-to-global Ext comparison being consumed, using the earlier finite filtered-complex construction; no general proper duality theorem is needed here.

These references are a finite proof dependency list, not a declaration that unnamed results are ordinary foundations. In particular the further smoothness descent and connected model results are proved in Sections 3–4.

1.1. The consumed Artin–Rees and completion arguments

The following records the actual arguments of AG-CA-03 Theorem 5.1 and AG-CA-19 Theorems 3.1–3.3 before their uses. Let A be Noetherian and $I \subset A$ an ideal.

Lemma 1.1 (Artin–Rees). If $N \subset M$ are finite A -modules, there is c such that

$$N \cap I^n M = I^{n-c}(N \cap I^c M) \quad (n \geq c).$$

Proof. The Rees algebra $B = \bigoplus I^n t^n$ is generated by the finitely many generators of I in degree one, so is Noetherian by Hilbert basis. The module $\bigoplus I^n M t^n$ is finite over it, generated by any finite list of generators of M in degree zero. Its graded submodule $\bigoplus (N \cap I^n M) t^n$ is finite. Choose homogeneous generators of degrees at most c . For $n \geq c$, the degree n component is a sum of $I^{n-d}(N \cap I^d M)$, for $d \leq c$; each is contained in $I^{n-c}(N \cap I^c M)$. The reverse inclusion follows by multiplication and the submodule property. $\square$

The consumed Krull intersection statement also follows from this proof. For a finite module over a Noetherian local ring, with I in its maximal ideal, the submodule $N = \bigcap_n I^n M$ is finite. Lemma 1.1 gives $N = IN$ by taking $n = c + 1$, since $N \cap I^n M = N$ at every index. Nakayama gives $N = 0$. This is the actual AG-CA-03 Theorem 6.1 argument used in the earlier regular-ring proof.

Lemma 1.2 (exact finite-module completion). Writing $\widehat{M} = \varprojlim M/I^n M$, completion is exact on finite modules and

$$\widehat{M} = M \otimes_A \widehat{A}.$$

Proof. For $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$, the quotients give

$$0 \rightarrow N/(N \cap I^n M) \rightarrow M/I^n M \rightarrow L/I^n L \rightarrow 0.$$

The kernel system has surjective transitions. Compatible lifts in the middle system can therefore be chosen successively: given a lift at stage n , take any lift of the desired next quotient element and adjust its image’s difference at stage n by lifting that difference from the kernel system. This proves surjectivity on inverse limits. Injectivity and equality with the limit kernel hold coordinatewise. By Lemma 1.1, $I^n N \subset N \cap I^n M \subset I^{n-c} N$; hence the two filtrations of N are cofinal. Their limits identify explicitly by taking a sufficiently far coordinate to define a coordinate of the other system, independent of that choice by compatibility. Thus the displayed inverse-limit sequence is $0 \rightarrow \widehat{N} \rightarrow \widehat{M} \rightarrow \widehat{L} \rightarrow 0$.

A finite module has a finite presentation, since submodules of finite modules are finite over a Noetherian ring. Complete $A^r \rightarrow A^s \rightarrow M \rightarrow 0$, using the just-proved exactness. Tensor the same presentation with $\widehat{A}$, using tensor’s right exactness. Finite direct sums complete coordinatewise, so both cokernels are the cokernel of the same matrix $\widehat{A}^r \rightarrow \widehat{A}^s$. This gives the canonical tensor comparison. In particular, when A is already complete, every finite module is complete. $\square$

Lemma 1.3 (flat, faithfully flat and Noetherian completion). The ring $\widehat{A}$ is Noetherian, is complete for $I\widehat{A}$, and is A -flat. If I is contained in the Jacobson radical, the completion is faithfully flat.

Proof. Flatness follows directly from the preceding finite-module exactness. Every module is the filtered union of its finite generated submodules. A tensor element uses finitely many such generators, and an equality to zero holds in a sufficiently large finite submodule, because tensor

commutes with filtered colimits by its generators and relations. For an injection $M' \rightarrow M$, put a proposed tensor kernel element in a finite submodule $N \subset M'$, and put its vanishing equality in some finite submodule of M containing the image of N . Exactness of completion for these finite modules proves injectivity there, hence kills the proposed kernel. Thus tensor with $\hat{A}$ preserves every injection.

For faithfulness, if $M \neq 0$, choose $0 \neq x \in M$, with cyclic submodule $Ax = A/J$. Choose a maximal ideal $\mathfrak{m} \supset J$. Since $I \subset \mathfrak{m}$, the module $A/\mathfrak{m}$ is unchanged by completion. The tensor comparison gives $\hat{A}/\mathfrak{m}\hat{A} = A/\mathfrak{m} \neq 0$, a quotient of $(A/J) \otimes \hat{A}$. Flatness injects the latter into $M \otimes \hat{A}$, so this is nonzero. This proves faithful flatness and, by applying it to a tensor kernel, detection of exactness.

Apply Lemma 1.2 to $0 \rightarrow I^a \rightarrow A \rightarrow A/I^a \rightarrow 0$. The last module is unchanged by completion, since a power of I kills it. The image of $I^a \otimes \hat{A}$ is exactly $I^a \hat{A}$, whence

$$\hat{A}/I^a \hat{A} = A/I^a.$$

Writing $J = I\hat{A}$, its powers are $I^a \hat{A}$; these identities and the defining inverse limit prove J -adic completeness and separatedness. Their successive kernels identify $\text{gr}_J \hat{A}$ with $\text{gr}_I A$. The latter is a quotient of a polynomial ring over A/I on the finite generators of I , and is Noetherian.

For any ideal $Q \subset \hat{A}$, its initial classes form a homogeneous ideal of this Noetherian graded ring. Choose finite homogeneous generators and lift them to $q_i \in Q$, with orders d_i . For $q \in Q$, subtract at stage n a combination $\sum_i a_{i,n} q_i$ matching its residual initial class, with $a_{i,n} \in J^{n-d_i}$ and zero when $n < d_i$. The new residual is in J^{n+1} . Each coefficient series converges, since its orders tend to infinity. Its sums a_i satisfy $q = \sum_i a_i q_i$ by separatedness. Thus every ideal is finite and $\hat{A}$ is Noetherian. No prior assertion that Q is closed was used. $\square$

For a local ring with maximal ideal $\mathfrak{m}$, its completion is local: an element with nonzero residue is a unit in every quotient $A/\mathfrak{m}^n$, and the compatible inverses give its inverse in the limit. Projection onto the residue field is surjective by successive lifting. The preceding quotient identity makes its kernel $\mathfrak{m}\hat{A}$. Thus Lemma 1.3 gives the exact local completion statement used by the group lessons.

Lemma 1.4 (idempotents across a nilpotent ideal). Every idempotent of R/J , for a nilpotent ideal J , has a unique idempotent lift to R .

Proof. First suppose $J^2 = 0$. For a lift e , put $\delta = e^2 - e \in J$. Then $e' = e - (2e - 1)\delta$ is idempotent: expand its square and use $\delta^2 = 0$ and $(2e - 1)^2 = 1 + 4\delta$. For two idempotent lifts the difference $u \in J$ satisfies $(2e - 1)u = 0$; the element $2e - 1$ is a unit since its square is one, so $u = 0$. Apply this square-zero argument successively to J^a/J^{a+1} , starting with $a = 1$; each is square-zero because $2a \geq a + 1$, and nilpotence makes the list finite. $\square$

2. Formal functions on a projective scheme

Let A be Noetherian, $I \subset A$ an ideal, $X \rightarrow \text{Spec } A$ projective and F coherent. Write

$$B = \bigoplus_{n \geq 0} I^n t^n \subset A[t], \quad M_n^q = H^q(X, I^n F), \quad M^q = M_0^q.$$

The transition maps $M_m^q \rightarrow M_n^q$, for $m \geq n$, come from inclusion of ideal powers. Multiplication $I^d \otimes M_n^q \rightarrow M_{n+d}^q$ is compatible with these transitions: both sheaf maps are multiplication followed by the same inclusion. Applying cohomology preserves their equality.

Lemma 2.1. The graded B -module $\bigoplus_{n \geq 0} M_n^q t^n$ is finite for every q .

Proof. Generators of I , placed in degree one, generate B as an A -algebra; Hilbert basis makes B Noetherian. Let $X_B = X \times_A \text{Spec } B$, with affine projection π . On $U = \text{Spec } R \subset X$, put $F|_U = \tilde{N}$. The graded module $\bigoplus I^n N$ is finite over $R \otimes_A B$: finitely many R -generators of N , placed in degree

zero, generate it. Here $R \otimes_A B \rightarrow \bigoplus I^n R$ is a surjection, and need not be an isomorphism. No flatness assumption is being made.

Localization gives the same modules on overlaps. They glue to a coherent sheaf G on X_B , with $\pi_* G = \bigoplus I^n F$. Coherence holds because $R \otimes_A B$ is Noetherian and the module is finite. Base changing a projective closed immersion shows that X_B is projective over B ; AG-QC-05 Theorem 2.1 and its closed-immersion passage give finite $H^q(X_B, G)$ over B .

Use the inverse images of a finite affine cover of X . Each intersection is affine on both schemes, and its sections of G are the sections of $\bigoplus I^n F$ on the original intersection. Thus their ordered section complexes are identical. Finite products in that complex commute with direct sums, and direct sums of modules are exact. Taking cohomology therefore identifies

$$H^q(X_B, G) = \bigoplus_{n \geq 0} H^q(X, I^n F),$$

with the graded B -action preserved. This proves finiteness. $\square$

This writes the projective case of the free proofs [Stacks 02O8](#) and [0897](#). General proper coherent pushforward and its dévissage are unnecessary: both spaces just used are projective.

Put

$$J_n^q = \text{im}(M_n^q \rightarrow M^q), \quad K_n^q = \ker(M_n^q \rightarrow M^q).$$

Lemma 2.2. There are constants $a_q, b_q \geq 0$ such that

$$I^n M^q \subset J_n^q \subset I^{n-a_q} M^q \quad (n \geq a_q),$$

and $K_m^q \rightarrow K_n^q$ is zero whenever $m \geq n + b_q$.

Proof. Multiplication by $u \in I^n$ on F factors through $I^n F$, proving the first inclusion. The image and kernel of the graded map from Lemma 2.1 to the corresponding degreewise copies of M^q are graded subquotients of a finite B -module. More explicitly, multiplication and transition compatibility show that $\bigoplus J_n^q t^n$ is the graded image module and $\bigoplus K_n^q t^n$ is a graded submodule of $\bigoplus M_n^q t^n$. Both are finite, because B is Noetherian.

Choose homogeneous generators of the image module of degrees $d \leq a_q$. Its degree n is a sum of terms $I^{n-d} J_d^q \subset I^{n-a_q} M^q$, proving the other inclusion. Choose homogeneous kernel generators $z_j \in K_{d_j}^q$, with $d_j \leq b_q$. An element in degree m is a sum of $u_j z_j$, where $u_j \in I^{m-d_j}$. If $m \geq n + b_q$, express u_j as a sum of products vw , $v \in I^n$, $w \in I^{m-d_j-n}$. Transition of vwz_j to M_n^q factors as transition of z_j to M_0^q , followed by multiplication by w , then by v into M_n^q . The first map kills z_j . Every such product, and hence the transition, is zero. $\square$

Theorem 2.3. There is a canonical isomorphism

$$H^q(X, F)_I^\wedge \xrightarrow{\sim} \varprojlim_n H^q(X, F/I^n F).$$

It is a homeomorphism for the quotient inverse-limit topologies.

Proof. Write $T_n^q = H^q(X, F/I^n F)$. The sheaf short exact sequence gives compatible exact sequences

$$0 \longrightarrow M^q/J_n^q \longrightarrow T_n^q \longrightarrow K_n^{q+1} \longrightarrow 0.$$

For $m \geq n + b_{q+1}$, the transition on the last term is zero. Hence the image of $T_m^q \rightarrow T_n^q$ is contained in the first term M^q/J_n^q . It equals that term: transition $M^q/J_m^q \rightarrow M^q/J_n^q$ is onto and these

modules inject into T_m^q, T_n^q . This proves the Mittag–Leffler assertion and its stable image, with a written uniform bound.

If (u_n) is a compatible family in T_n^q , its image in K_n^{q+1} is zero at every coordinate, because that image is the transition of a sufficiently distant kernel term. Thus u_n has a unique preimage in M^q/J_n^q ; uniqueness makes those preimages compatible. Consequently

$$\varprojlim_n T_n^q = \varprojlim_n M^q/J_n^q.$$

Lemma 2.2 gives $I^n M^q \subset J_n^q$ and $J_{n+a_q}^q \subset I^n M^q$. Projection and projection from the shifted index define inverse maps between the two quotient limits. A common finer index proves that their composites are the identity. Each map is continuous coordinate by coordinate, proving the topological assertion. Every construction is the canonical quotient map, so the isomorphism is the asserted one. $\square$

This supplies the critical image, kernel and limit arguments of [Stacks 02OA](#), [02OB](#) and [02OC](#). In particular, the proof uses the kernel transition bound directly; it does not rely on the incorrect part-number reference presently printed in 02OB. No inverse limit has been interchanged with an unexplained cohomology operation.

Corollary 2.4. If A is complete local Noetherian, every compatible idempotent on the maximal-ideal thickenings of a projective X comes from a unique idempotent of $\Gamma(X, \mathcal{O}_X)$.

Proof. Projective finiteness makes $M = \Gamma(X, \mathcal{O}_X)$ finite. Lemma 1.2 gives $M = M \otimes_A \hat{A} = \widehat{M}$. Theorem 2.3 identifies this ring with the limit of the thickened section rings. The equality $e^2 = e$ holds in this limit coordinatewise and hence holds for the corresponding actual section. Uniqueness follows from the same ring isomorphism. Idempotents at successive levels are compatible whenever they lift the same initial idempotent, by Lemma 1.4’s unique nilpotent lifting. $\square$

3. Smoothness and line bundles at a finite stage

Lemma 3.0 (the split conormal criterion). For a polynomial presentation $C = P/J$, the algebra is formally smooth precisely when $d : J/J^2 \rightarrow \Omega_{P/B} \otimes_P C$ has a C -linear left inverse.

Proof. Formal smoothness lifts the identity of C to a section $s : C \rightarrow P/J^2$. The map $D(p) = p \bmod J^2 - s(p \bmod J)$ takes values in J/J^2 , satisfies the product rule because this ideal is square-zero, and sends $j \in J$ to its class. The universal derivation for the polynomial ring therefore factors it through $\Omega_{P/B} \otimes C$, giving a left inverse of d . Conversely a left inverse gives this derivation by composition with the polynomial universal derivation. The map $p \mapsto p \bmod J^2 - D(p)$ is multiplicative by the product rule and the square-zero condition. It kills J and is the identity after reduction to C , so gives a section. In any square-zero lifting test, lift the polynomial variables arbitrarily. The resulting map from P sends J into the square-zero test ideal and kills J^2 ; composing its factor through P/J^2 with the section gives the required lift. Polynomial differentials are free on the variable symbols by the product rule, so no unproved splitting assertion enters this argument. $\square$

Lemma 3.A (maps between smooth schemes). Let X and Y be smooth schemes locally of finite presentation over any scheme S , and let $h : X \rightarrow Y$ be an S -map. If the map on relative tangent spaces at a point x , with scalars extended to $\kappa(x)$, is surjective, then h is smooth near x . If that map is an isomorphism, then h is étale near x . In particular, on a smooth morphism of relative dimension d , local functions $q_1, \dots, q_d$ whose differentials form a basis at x give an étale coordinate map to $\mathbb{A}_S^d$ near x .

Proof. Work on compatible affine neighbourhoods of x , $h(x)$ and their image in S . Write their algebras as C, D, A . The map $D \rightarrow C$ is finitely presented: choose finite presentations of D and C

over A , and add, to the presentation of C , the finitely many equations equating each generator of D with a polynomial representing its image in C . This gives a finite presentation over D .

The split conormal criterion makes $\Omega_{C/A}$ and $\Omega_{D/A}$ finite projective. Indeed, in a finite polynomial presentation the conormal injection has a left inverse, so its cokernel is a direct summand of the finite free module of polynomial differentials. Such a summand is flat, and the local finite-freeness proof in AG-RG-S01, Lemma 1.2, applies. Shrink the affine neighbourhoods to trivialize these modules. Dualizing the asserted tangent surjection gives an injection on the fibre of

$$u : C \otimes_D \Omega_{D/A} \longrightarrow \Omega_{C/A}.$$

A full-size minor of its matrix is nonzero at x ; invert it. Projecting to the corresponding rows and multiplying by that square matrix's inverse gives a C -linear left inverse of u . In the tangent-isomorphism case its matrix is square and invertible after this shrink.

Consider an arbitrary square-zero surjection $R \rightarrow R/J$, an A -algebra map $C \rightarrow R/J$, and a prescribed D -algebra structure on R agreeing with it after reduction. Formal smoothness of C/A , as proved by Lemma 3.0, supplies an A -algebra lift $v : C \rightarrow R$. Its restriction to D differs from the prescribed map by an A -derivation $\delta : D \rightarrow J$: expanding a product leaves the two first-order terms, since $J^2 = 0$. It corresponds to a map $C \otimes_D \Omega_{D/A} \rightarrow J$, where the C -action on J is through the given reduced map. Compose it with the left inverse of u to obtain $\Omega_{C/A} \rightarrow J$, hence an A -derivation $\tilde{\delta} : C \rightarrow J$. The map $v - \tilde{\delta}$ is again an A -algebra map by the same square-zero product expansion. Its restriction to D is the prescribed map. Thus C/D is formally smooth. Its finite presentation makes it smooth.

If u is an isomorphism, two such D -lifts differ by a derivation $C \rightarrow J$ whose differential map vanishes on the image of u . That image is all of $\Omega_{C/A}$, so the lifts agree. The map is formally étale and finitely presented, hence étale. The coordinate assertion is this case with $Y = \mathbb{A}_S^d$: its pulled-back differentials have basis $dq_1, \dots, dq_d$, and the given basis condition makes u an isomorphism near x . No Noetherian, reducedness, characteristic or residue-field separability hypothesis was used. $\square$

Lemma 3.1. Let $A = \varinjlim A_i$ be a filtered ring colimit. A finitely presented algebra map $B_0 \rightarrow C_0$ over A_0 whose base change over A is smooth becomes smooth after a finite stage.

Proof. Present $C_0 = B_0[x_1, \dots, x_r]/(f_1, \dots, f_s)$. At stage i , let $P_i = B_i[x_1, \dots, x_r]$, $J_i = (f_{1,i}, \dots, f_{s,i})$ and $C_i = P_i/J_i$. Lemma 3.0, the split conormal criterion of AG-CA-17 Theorem 3.1, says that the map

$$J/J^2 \xrightarrow{d} C^r$$

over the colimit has a C -linear left inverse ρ . Each $\rho(dx_k)$ is a finite sum of classes of the f_j with coefficients in C . Lift these finitely many coefficients to one stage and define $\rho_i : C_i^r \rightarrow J_i/J_i^2$ by those images. The equations $\rho_i(df_{j,i}) = f_{j,i} \bmod J_i^2$ are finitely many equations. Their truth in the colimit is witnessed by finite sums of products of the f_j , representing membership in J^2 . Lift these witnesses and enlarge the stage until the polynomial equalities hold. Then $\rho_i d$ fixes all generators of J_i/J_i^2 , so is its identity. The same criterion gives formal smoothness at that stage, and the existing finite presentation gives smoothness. $\square$

This is the actual finite witness proof in [Stacks 0C0B](#).

Proposition 3.2. A smooth scheme of finite presentation over an affine filtered limit has a smooth finite-presentation model at a finite stage. A projective embedding and a specified line bundle can be retained simultaneously.

Proof. First descend the finite-presentation scheme, the embedding if specified, and finitely many affine source and target charts by AG-MO-03 Theorem 4.1. Retain the closed immersion by its Theorem 4.2. Choose a finite affine cover of the original source on which the chart algebra maps are smooth. Descend these opens and the maps, and retain their covering property by Lemma 2.1

of that lesson. Apply Lemma 3.1 on the finitely many chart maps and take a common later stage. Smoothness is now true on a cover, hence globally.

For an invertible sheaf, refine the cover so that it is trivial on each chart. On finite affine overlap covers record each transition unit and its inverse. Record their inverse equations, their agreements on overlap refinements, and the cocycle equations on finite covers of triple overlaps. The sections and their finite equality witnesses descend by the section and eventual-equality parts of AG-MO-03. At a common stage these equations hold, so gluing the trivial rank-one sheaves gives an invertible sheaf there. Retain the finite chart covers, smoothness and the projective closed immersion at that same stage. Pullback recovers the original sheaf and scheme. $\square$

This fills the finite-cover step omitted in [Stacks 0C0C](#). It proves only the finite chart and line bundle case being used, without importing general coherent sheaf descent or a general perfection theorem.

4. Geometrically connected smooth affine models

The model needed for the normalizer argument is smooth and affine. We prove this case; the unrestricted finite-presentation connected-fibre theorem is not needed.

Lemma 4.1. Over an algebraically closed field k , an affine finite type connected scheme remains connected after every field extension. A smooth affine finite type scheme has disjoint irreducible components, and each connected component is integral and remains integral after every field extension of an algebraically closed ground field.

Proof. Connectedness of $\text{Spec } B$ is equivalent to absence of nontrivial idempotents. Indeed an idempotent gives the disjoint opens $D(e), D(1 - e)$; conversely a disjoint open-and-closed decomposition gives a section with values one and zero, hence an idempotent by affine sections.

Suppose an extension L/k supplies a nontrivial idempotent $e \in B \otimes_k L$. Express e and $1 - e$ using finite lists of k -linearly independent elements of B , with coefficients in L . Let $R \subset L$ be the finite type k -algebra generated by these coefficients, inverted nonzero coefficients ensuring that both expressions stay nonzero, and the finitely many coefficients needed for $e^2 = e$. The relation already holds over R , since $B \otimes_k R \rightarrow B \otimes_k L$ is injective. The nonzero finite type algebra R has a k -valued point by the proved Nullstellensatz in AG-CA-06. Specializing at it gives a nontrivial idempotent in B , a contradiction.

For smooth B , every local ring is a regular local domain by AG-CA-18 and AG-CA-14. Two distinct irreducible components cannot meet at a point, since their distinct minimal primes would give distinct minimal primes in that local ring. There are finitely many components; they are consequently both closed and open. Smoothness also makes the ring reduced. Each connected component is thus reduced and irreducible, hence integral. After extending an algebraically closed field, connectedness just proved and smoothness preserved by base change give the same conclusion. $\square$

It follows that the number of geometric connected components of a smooth affine finite type scheme is invariant under field extensions. To check this, embed the chosen algebraic closures of the two fields into a common algebraically closed extension. Apply Lemma 4.1 to each of the finitely many open-and-closed components. Extension is faithfully flat and preserves nonemptiness, so each component remains one component.

Lemma 4.A (field and integer calculations before primitive elements). Let F be any field. Polynomial division and Bezout hold in $F[T]$. Minimal-polynomial quotients, distinct-root bounds, the derivative test for a separable algebraic generator, finite splitting fields, and finite tower degree and embedding counts have the following direct proofs. Every finite subgroup of a field's units is cyclic. Every integer ideal is principal and $\mathbf{Z}$ is Noetherian. The integer division, Bezout and prime-factorization facts needed for the group calculation are proved here as well.

Proof. First give the integer arithmetic used in the finite-group calculation. For integers z and $d > 0$, there are q, r with $z = qd + r$ and $0 \leq r < d$. For $z \geq 0$, choose the largest q in the finite set $\{0, \dots, z\}$ with $qd \leq z$; if the remainder were at least d , the next integer would contradict maximality. For $z < 0$, divide $-z$; negate its quotient when the remainder is zero, and otherwise use quotient $-q - 1$ and remainder $d - r$. Thus integer division also covers negative integers. A nonzero subgroup $J \subset \mathbf{Z}$ contains a positive integer and has a least positive member d . The remainder on dividing any member of J by d is in J and smaller than d , so is zero. Hence $J = d\mathbf{Z}$. The zero ideal is generated by zero, so every integer ideal is principal. For an ascending chain of integer ideals, its union is an ideal generated by one integer d . If $d = 0$, every member is zero. Otherwise d lies in one member, which then contains the whole union, so the chain stabilizes. This proves that $\mathbf{Z}$ is Noetherian. For a, b not both zero, apply this to the subgroup of integer combinations $ua + vb$. Its positive generator d divides both a, b , every common divisor divides d , and $d = ua + vb$ for some integers u, v . This proves the positive gcd and Bezout identity. If $\gcd(a, b) = 1$ and $a \mid bc$, multiplication of a Bezout identity by c gives $a \mid c$. A positive prime integer is consequently a prime element. Every integer $n > 1$ factors into positive primes by induction: if it is composite, write it as a product of two smaller positive integers and factor those. Prime divisibility and integer cancellation prove uniqueness. The integer 1 has the empty factorization. The least common multiple of a nonempty finite list of positive integers is therefore the product of the largest occurring powers of each prime; it is 1 if all entries are 1. Different prime powers have gcd one. For coprime positive a, b , the divisibilities $a \mid r$ and $b \mid r$ imply $ab \mid r$: write $r = bc$ and apply the coprime divisibility assertion to obtain $a \mid c$. These are all the integer facts required below.

If $g \neq 0$ and f has degree at least that of g , subtract the leading coefficient of f divided by that of g , times the appropriate power of T times g . This strictly decreases the degree. Induction gives $f = qg + r$ with $r = 0$ or $\deg r < \deg g$. The leading coefficient of a product of two nonzero polynomials is the product of their nonzero leading coefficients, so degrees add. Consequently two such division expressions have the same remainder and quotient. For polynomials a, b not both zero, repeat division on the pair and its successive remainders. Nonnegative degrees strictly decrease, so the process ends. Its last nonzero remainder, scaled to be monic, is d . Back substitution expresses $d = ua + vb$. Each earlier remainder is divisible by d , and any common divisor of a, b divides every remainder, so d is their monic greatest common divisor. In particular coprime polynomials satisfy $ua + vb = 1$. An irreducible f is prime: if f does not divide a , Bezout for f, a , multiplied by b , shows that $f \mid ab$ forces $f \mid b$. Factor existence follows by induction on degree, and prime divisibility gives uniqueness after fixing monic factors and a scalar.

For an algebraic element α , choose a monic nonzero annihilating polynomial of least degree, denoted f . Division shows that every annihilating polynomial is a multiple of f . If f were a product of two nonconstant factors, evaluating in the extension field would make one smaller factor vanish, a contradiction. Thus f is irreducible. Every nonzero class modulo f has a polynomial inverse by Bezout, so $F[T]/(f)$ is a field. Evaluation gives $F[\alpha] = F(\alpha)$ and identifies it with this quotient. The remainders $1, T, \dots, T^{\deg f - 1}$ form a basis: they span by division, and a linear combination of smaller degree cannot be a nonzero multiple of f .

Division by $T - c$ leaves the constant $f(c)$. If c is a root, $f = (T - c)g$. Every other root is a root of g , since its difference from c is a nonzero field element. Induction proves that a nonzero polynomial of degree d has at most d distinct roots, over every extension field. A finite collection of polynomials has a finite splitting field: if one polynomial has a nonsplit nonconstant factor, factor it into irreducibles by the degree induction, adjoin a root of one such factor using the just-proved field quotient, remove the linear factor, and repeat. At most the sum of the original degrees many roots need be adjoined. The extensions at every step are finite by the displayed power basis, and adjoining these roots therefore yields a finite extension in which every original polynomial splits.

The formal derivative obeys the product rule by expanding monomials. In a splitting field, if $f = (T - c)g$, then $f'(c) = g(c)$. Hence c is a repeated root exactly when f and f' both vanish at c . For an irreducible f with $f' \neq 0$, the derivative has smaller degree, so Bezout makes f, f' coprime

and there are no repeated roots. If $f' = 0$ and f is nonconstant, every root is repeated. Thus the minimal polynomial of an algebraic element is separable precisely when its derivative is nonzero, equivalently when $f'(\alpha) \neq 0$.

The square-matrix test used with the differential presentation also follows directly over a field. Define the determinant by its finite signed permutation sum. That sum is linear in each row and each column. With two equal rows its terms cancel in pairs under their transposition, so it is zero; row interchange changes its sign. Consequently adding a multiple of another row leaves it unchanged, and scaling one row multiplies it by the scale. The same assertions hold for columns. Starting with an n by n matrix, move a nonzero entry to the first position by row and column exchanges, scale it to one, and clear the other entries in that row and column by these additions. Repeat on the remaining square block. Every operation is invertible and alters the determinant by a nonzero scalar. The result is a diagonal matrix with some entries one and the others zero. If any zero entry remains, that output coordinate cannot occur in the image, so the map is not surjective. Hence a surjective square map has only ones and has nonzero determinant. Conversely such a diagonal matrix is invertible. The determinant of a triangular matrix is the product of its diagonal entries: expand in the first column for an upper triangular matrix, or in the first row for a lower triangular matrix, and induct. For size zero the determinant is one and the map is surjective.

If $u_1, \dots, u_r$ is an F -basis of a finite extension E and $v_1, \dots, v_s$ is an E -basis of L , the $u_i v_j$ span L over F . A relation between them is first a relation between the v_j over E and then, coefficient by coefficient, a relation between the u_i over F . Both sets of coefficients are zero. Thus they form an F -basis and $[L : F] = [L : E][E : F]$. A finite extension has a finite generating list, for example any finite F -basis, and hence a finite simple-generator tower.

To count embeddings, fix a finite generating list $\alpha_1, \dots, \alpha_m$ of L/F and a finite splitting field H/F of their minimal polynomials over F . Put $E_i = F(\alpha_1, \dots, \alpha_i)$. The minimal polynomial f_i of α_i over E_{i-1} divides its minimal polynomial over F , by polynomial division. Under any F -embedding $\sigma : E_{i-1} \rightarrow H$, the polynomial $\sigma(f_i)$ therefore divides that already split F -polynomial and splits in H . An extension of σ to $E_i = E_{i-1}[T]/(f_i)$ is uniquely specified by the image of α_i , and each distinct root of $\sigma(f_i)$ defines such an extension by evaluation. The evaluation map is injective because its domain is a field. There are at most $[E_i : E_{i-1}]$ such roots, and exactly that many if f_i is separable. An element separable over F stays separable over an intermediate field because its new minimal polynomial divides the old one. Thus a separable extension L/F has exactly $[L : F]$ embeddings into H , by multiplying the tower counts. Conversely, if every step is separable, these counts also give $[L : F]$ embeddings. If some $\beta \in L$ were inseparable over F , its minimal polynomial would have fewer than $[F(\beta) : F]$ distinct roots. Enlarge H to split that polynomial and all polynomials for a generator tower above $F(\beta)$; no new embeddings of L appear, since every generator image was already a root in the original H . Restricting to $F(\beta)$ and applying the upper bound on extensions gives fewer than $[F(\beta) : F][L : F(\beta)] = [L : F]$ embeddings, a contradiction. This proves the transitivity of separability and all finite embedding counts used here. Only a finite splitting field was required.

Let G be any finite subgroup of a field's units. The order of $g \in G$ is the least positive d with $g^d = 1$, which exists by equality of two powers in the finite group. Integer division shows $g^r = 1$ exactly when d divides r . Let e be the least common multiple of the finitely many orders, computed by the maximum exponent of each prime in their factorizations; integer prime factorization and Bezout were proved above. For each prime ℓ dividing e , select an element g_ℓ of order $\ell^s m$ with ℓ not dividing m and ℓ^s the maximal occurring power. Then $h_\ell = g_\ell^m$ has order exactly ℓ^s , by the divisibility test just proved. For commuting elements x, y of coprime orders a, b , if $(xy)^r = 1$, then $x^r = y^{-r}$ belongs to both cyclic subgroups. Its a th and b th powers are one; an integer Bezout identity makes the element one. Thus both a and b divide r , and ab divides r by the same Bezout identity. Since $(xy)^{ab} = 1$, its order is ab . Inducting over the primes shows that $h = \prod h_\ell$ has order e . Every element of G is a root of $T^e - 1$, so the polynomial root bound gives $|G| \leq e$. The e different powers of h give the reverse inequality. Therefore G consists of those powers and is cyclic.

If G is trivial, the empty prime list gives $e = 1$ and $h = 1$, so this case is included. $\square$

Lemma 4.2. Let A be a Noetherian domain with fraction field K . If a smooth affine finite type A -scheme has geometrically integral generic fibre, there is a nonempty principal open of $\text{Spec } A$ on which all its fibres are geometrically integral.

Proof. Write the algebra as B . Smoothness makes B A -flat, so $B \hookrightarrow B_K$. Since the latter is a domain, B is a domain. Let $L = \text{Frac}(B_K)$. An étale coordinate chart at the generic point gives algebraically independent $t_1, \dots, t_d \in L$ with L finite separable over $K(t_1, \dots, t_d)$: use the standard smooth chart and its zero-relative-dimension coordinate map from AG-CA-17 Theorem 5.1, applied over the field, followed by AG-CA-18 Corollary 6.2 and AG-CA-17 Lemma 7.1. A primitive element u writes

$$L = K(t_1, \dots, t_d)(u).$$

Here is the complete primitive-element argument, using Lemma 4.A. Let L/F be finite separable of degree n . Its finite generating list $a_1, \dots, a_r$ has exactly n distinct embeddings $\sigma_1, \dots, \sigma_n$ into one finite splitting field H . If F is infinite, for each pair $i \neq j$ the form $\sum_t (\sigma_i(a_t) - \sigma_j(a_t))X_t$ is a nonzero polynomial over H , since the embeddings differ on a generator. Their finite product is nonzero: the leading coefficient of a product of nonzero polynomials over a field is nonzero, iterating by the number of variables. A nonzero polynomial over H does not vanish on all tuples in the infinite subfield F . Indeed, induction on the number of variables first chooses a tuple in F where one nonzero coefficient polynomial is nonzero, and then avoids the finitely many roots in the last variable. Choose $c_t \in F$ where this product does not vanish. The element $a = \sum c_t a_t$ has n distinct images. Its minimal polynomial therefore has degree at least n by the root bound, and at most n because $F(a) \subset L$. The tower-degree equality gives $L = F(a)$. In a nontrivial extension this element cannot be zero; for a trivial extension use 1.

If F is finite, its finite basis makes L a finite set. Lemma 4.A proves that $L^\times$ is cyclic, including the trivial group case. Choose a generator u of this group. Every nonzero element of L is a power of u , and zero belongs to every subfield; hence $F(u) = L$. The generator is nonzero, and for a trivial extension we may choose 1. Apply this with $F = K(t_1, \dots, t_d)$ and the field $L = \text{Frac}(B_K)$ above to obtain the required u in the displayed simple-extension presentation.

Clear denominators in the minimal polynomial of u , removing the coefficient content, to obtain a primitive polynomial $P \in K[t_1, \dots, t_d, T]$. It is absolutely irreducible. Indeed $B_K \otimes_K \bar{K}$ is a domain, and localizing it gives the domain $L \otimes_K \bar{K}$. The algebra $K[t_1, \dots, t_d, T]/(P)$ embeds in L by its primitive minimal-polynomial presentation. Tensoring this injection with the field $\bar{K}$ keeps it injective. Its target is the just-obtained domain, so the polynomial quotient over $\bar{K}$ is a domain, proving absolute irreducibility without any assertion about contents after field extension.

Here is the needed polynomial Gauss argument. A product of primitive polynomials over a UFD stays primitive: modulo any prime coefficient factor, the two reductions are nonzero polynomials over a domain, so their product is nonzero. Starting with a field and inducting on the number of variables, factor the coefficient content and then factor the primitive part over the fraction field, whose one-variable polynomial ring has Euclidean division. Clear the denominators and normalize each factor to primitive content. The preceding product calculation shows that this gives a factorization over the coefficient ring, and uniqueness follows from the fraction-field uniqueness and the content. Thus each polynomial coefficient ring is a UFD; its primitive minimal polynomial generates the kernel of evaluation in the field extension, as used above.

Enlarge a nonzero localization of A until P has coefficients there. There is a further nonempty localization on which its specialization is absolutely irreducible in every residue field. Here is a finite proof of this assertion. A triangular variable change $x_i = y_i + y_n^{N_i}$ makes the weights of the finitely many monomials distinct, using successive powers of an integer larger than all occurring exponents. The unique largest weight gives a nonzero leading coefficient in y_n . Invert it and divide

by it, making P monic of degree $r > 0$ in that variable. Let D_i bound its degrees in the other variables. A nontrivial factorization over a field has monic factors of positive y_n -degrees $e, r - e$, $1 \leq e < r$, with other degrees at most D_i ; degrees add in each variable over a domain. For each e , introduce the finitely many coefficients of these possible factors and impose the coefficients of $P - Q_1 Q_2$. This gives a finite type A -algebra R_e . Its generic fibre is zero by absolute irreducibility and the Nullstellensatz over $\bar{K}$. Therefore $1 = 0$ after tensoring with K , so some nonzero $a_e \in A$ makes $(R_e)_{a_e} = 0$. Inverting the finite product excludes every nontrivial factorization over every field extension of every residue field. Degree-zero factors in the monic variable are units and are excluded from this list. This is the finite coefficient proof of the free [Stacks 0557](#) result, with only positive factor degrees used.

Put $B' = A[t_1, \dots, t_d, T]/(P)$ over the localization just obtained. Its generic fraction field and that of B are the same. The two generic affine integral schemes have a common nonempty affine principal open: represent the finite generators of each algebra as fractions in the other, invert their finitely many denominators, and invert the denominators needed for these descriptions to be inverse. AG-MO-03's map descent and eventual equality spread this isomorphism after another nonzero base localization. Thus we have $B_h \cong B'_{h'}$, with $h \neq 0$ and nonempty generic open.

Apply AG-FSE Corollary 5.2 in **Flat morphisms** to the finite B -module B/hB , and shrink so that it is A -flat. In the sequence $0 \rightarrow B \xrightarrow{h} B \rightarrow B/hB \rightarrow 0$, flatness of the last term makes multiplication by h injective after every field base change. The smooth fibre is reduced and Noetherian. A nonzerodivisor belongs to no minimal prime: if it did, localization at that minimal prime would make it zero in the field of a reduced Noetherian ring, giving an annihilator outside the prime. Therefore $D(h)$ is dense in every geometric fibre.

Chevalley's proved theorem makes the image of the common open constructible. It contains the generic point, so contains a nonempty open of the base. Shrink to a principal open in it; now the common open is nonempty in every fibre, and remains so after field extension. The B' -fibre is geometrically integral by the polynomial argument. Its nonempty open $D(h')$, identified with $D(h)$, is geometrically irreducible. Being dense in the smooth, geometrically reduced B -fibre, it forces that whole fibre to be irreducible and reduced. This proves the lemma. $\square$

Lemma 4.3. For a smooth affine finite type morphism over a Noetherian base, the number of geometric connected components of its fibres is a constructible function.

Proof. By the Noetherian constructibility criterion in AG-MO-06 Theorem 3.1, it suffices to show constancy on a nonempty open of every irreducible reduced closed subscheme of the base. Restrict to such an integral affine base $\text{Spec } A$, with fraction field K .

If the generic fibre is empty, its flat affine algebra injects into zero and is zero, so the component number is constantly zero. Otherwise the generic geometric fibre has finitely many open-and-closed integral components by Lemma 4.1. Their finitely many orthogonal idempotents, summing to one, descend to a finite separable extension K'/K . To justify separability, first express them over $\bar{K}$. In positive characteristic each expression uses finitely many coefficients which become separable after some common p -power. Since an idempotent equals its own p -power, raising its expression to this power puts its coefficients in K^{sep} . In characteristic zero this step is unnecessary. The finite list of algebraic separable coefficients lies in one finite separable extension. Its components are geometrically integral, as checked after extension to $\bar{K}$.

Write $K' = K[v]$ by the primitive element argument in Lemma 4.2. Clear denominators in its monic minimal polynomial and a Bezout identity with its derivative. After a nonzero localization, $A' = A[v]$ is finite free étale over A , with generic fibre the field K' . It is a domain because its monic polynomial presentation is free over A and embeds in K' . The finite generic idempotents extend after inverting finitely many elements of A' ; their finite orthogonality and sum equations hold after shrinking, or directly by the generic injection coming from flatness. They split the base-changed smooth algebra into finitely many smooth factors. Each has geometrically integral generic fibre. Apply Lemma 4.2 to all factors and invert a common nonzero $a \in A'$. At every point of $D(a)$ the

factors are exactly the geometric connected components.

The determinant of multiplication by a on the finite free A -module A' is nonzero, because this map is invertible on the generic field. Invert this determinant in A . The adjugate makes a invertible on the whole finite étale cover. That cover is surjective, since a nonzero finite algebra has nonzero fibres, and those fibres have points. Field-extension invariance from Lemma 4.1 now proves the same constant component number on this nonempty base open. This verifies the criterion on every irreducible closed base. Noetherian induction, removing such a nonempty open on each of finitely many irreducible components and then applying the induction hypothesis to their proper closed complement, gives a finite locally closed stratification with constant component count. This proves constructibility as a function, and in particular of its one-component locus. $\square$

This proves the smooth affine case needed from the free chain [055I](#), [055H](#), [055G](#). The hypersurface and flat quotient arguments above supply the generic constancy input explicitly; none of those tags is being substituted for a proof.

Lemma 4.4. Suppose $A = \varinjlim A_i$, A_0 is Noetherian, and $E \subset \operatorname{Spec} A_0$ is constructible. If the image of $\operatorname{Spec} A$ lies in E , the image of $\operatorname{Spec} A_i$ lies in E at some finite stage.

Proof. Express the complement as a finite union of pieces $V(J) \cap D(g)$, with J finitely generated. Its pullback has empty spectrum, so $(A/JA)_g = 0$. This says $g^m \in JA$ for some m , with a finite expression in generators of J . Its coefficients and the finite equality descend to a stage. The same localized quotient there is zero, so that complement piece has empty pullback. Take a common stage for the finitely many pieces. $\square$

This is a direct affine proof of the needed [Stacks 05F4](#) step.

Theorem 4.5. A smooth affine finite-presentation scheme with geometrically connected fibres over an arbitrary affine ring has a smooth affine model with geometrically connected fibres over a finitely generated $\mathbf{Z}$ -algebra. Finitely many specified maps, group laws and closed immersions can be retained.

Proof. Descend the finite presentations and finite identities to A_0 , a finitely generated $\mathbf{Z}$ -subalgebra of the given ring, using AG-MO-03. Retain smoothness by Section 3 and closed immersions by the earlier proved closed-immersion descent. On this Noetherian model let E be the locus of exactly one geometric connected component. It is constructible by Lemma 4.3. Its pullback contains the image of the original base, by field-extension invariance and the original hypothesis. Apply Lemma 4.4 to enlarge the coefficient ring to a finite stage whose whole image lies in E . Pullback of the model at that stage has geometrically connected fibres and remains smooth. Previously descended finite identities and closed immersions remain true under this base change. $\square$

This is the consumed smooth affine case of [Stacks 05FI](#). It applies simultaneously to the smooth affine group and its smooth affine subgroup in the normalizer lemma. Projectivity of quotient fibres has not been imposed on unrelated fibres of the model.

5. A universal finite complex for a line bundle

The following is the previously written finite complex proof, retained here before its uses.

Lemma 5.1. Let A be Noetherian and C a complex of flat A -modules concentrated in $[a, b]$, with finite cohomology modules. There is a complex K of finite projective modules in $[a, b]$ and a quasi-isomorphism $K \rightarrow C$ which remains a quasi-isomorphism after tensoring with every A -module.

Proof. Construct a bounded-above complex P of finite free modules and a map $f : P \rightarrow C$, descending from degree b . With terms above j chosen, the cone has terms

$$D^i = C^i \oplus P^{i+1}, \quad d(c, p) = (d_C c + f(p), -d_P p).$$

Its cohomology is finite: the cone long exact sequence uses finite cohomology of C and finite cohomology of the finite portion of P chosen so far. Choose finitely many cocycles (c_ℓ, p_ℓ) generating $H^j(D)$. Let P^j be free on e_ℓ , and set $f(e_\ell) = c_\ell$, $d_P(e_\ell) = -p_\ell$. The cocycle equations give $d_P^2 = 0$ and $d_C f = f d_P$. The new cone differential sends $(0, e_\ell)$ to (c_ℓ, p_ℓ) , killing H^j without changing higher cohomology. Continue to the left. Each degree stabilizes, giving a quasi-isomorphism $P \rightarrow C$.

Its cone is acyclic, bounded above and termwise flat. Such a complex stays acyclic after every tensor: start at its last degree and descend through $0 \rightarrow Z^i \rightarrow D^i \rightarrow Z^{i+1} \rightarrow 0$. The quotient is flat, so the kernel is flat and the sequence remains exact under tensor. Thus $H^i(P \otimes_A M) = H^i(C \otimes_A M)$ for every M .

Put $Q = \text{coker}(P^{a-1} \rightarrow P^a)$. The left tail is a free resolution of Q , and its first positive homology after tensoring with every M is $H^{a-1}(P \otimes_A M) = 0$, because $C \otimes_A M$ starts in degree a . Consequently Q is flat: tensor that free resolution with a short exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$. Its free terms give a short exact sequence of complexes; the homology exact sequence and the just-proved degree-one vanishing give an injection $Q \otimes M' \rightarrow Q \otimes M$. This is the definition of flatness.

The finite presentation of Q makes it locally finite free by [Algebra and sheaf cohomology before reductive groups](#), Lemma 1.2. It is finite projective as follows. Take a surjection $A^r \rightarrow Q$ and a finite principal cover on which it splits. A local splitting extends after multiplication by a power of that chart element, since Hom from a finite presented module commutes with localization: use its finite presentation and exact localization to identify the kernel describing Hom . Increase that power until its composite with the surjection is the same power times the identity globally; equality can be checked on finitely many generators. Thus there are maps $\tau_i : Q \rightarrow A^r$ whose composites are $f_i^{N_i} \text{id}_Q$. These powers generate the unit ideal, since the $D(f_i)$ cover. A linear combination of the τ_i whose coefficients combine the powers to one is a global splitting. Hence Q is a direct summand of a finite free module and is finite projective. Set $K^a = Q$, $K^i = P^i$ for $a < i \leq b$. The map to C factors through Q since $C^{a-1} = 0$. Truncation preserves cohomology, and the same acyclic flat cone argument proves the assertion after every tensor. $\square$

Proposition 5.2. Let $X \rightarrow S$ be smooth projective of finite presentation and L invertible. If all positive cohomology of L on every geometric fibre vanishes, then $f_* L$ is finite locally free, all positive higher direct images vanish, and these statements commute with every base change.

Proof. Work over $S = \text{Spec } A$. Section 3 gives a smooth projective model $X_0 \rightarrow \text{Spec } A_0$, A_0 Noetherian, and an invertible L_0 pulling back to L . A finite affine cover of X_0 has affine intersections. Its section complex C_0 computes cohomology by the earlier finite-cover and affine-vanishing proofs. Its terms are A_0 -flat: an invertible module is projective over its chart algebra, and that algebra is base-flat by smoothness. Projective finiteness makes its cohomology finite. After every algebra base change the section complex is $C_0 \otimes_{A_0} B$, by affine modules and localization. Lemma 5.1 therefore gives a bounded finite projective K_0 computing cohomology after every such change.

Put $K = K_0 \otimes_{A_0} A$, and trivialize its terms near a point s . If a differential has an entry invertible at s , shrink and use row and column operations to make it an identity block. The equation $d^2 = 0$ makes the adjacent differentials zero on that block, splitting off a contractible pair $A \xrightarrow{1} A$. After finitely many cancellations the remaining differentials vanish over $\kappa(s)$. Each remaining free term tensored with $\kappa(s)$ is then the cohomology in its degree. In positive degrees it is zero by the hypothesis and faithful field extension; in negative degrees it is zero because the section complex begins in degree zero. All remaining terms other than degree zero have rank zero. The degree-zero term is finite free. Cancelled pairs remain contractible under every tensor, so the same description computes all base changes. It proves the assertions on every affine open and every affine base change. The canonical maps from sections and the section complexes commute with restriction, so glue to the claimed direct-image and base-change isomorphisms. $\square$

Vanishing was required on the original fibres only. No vanishing on unrelated Noetherian model fibres, reduced-base assumption or flat-base-change assumption was used.

6. The smooth curve regular immersion calculation

The following retains the full independently authored CC0 proof of AG-QC-14, Sections 3–5, with its formerly general proper-finiteness input replaced by the projective proof already given in [Algebra and sheaf cohomology before reductive groups](#). Its injective and finite-cover foundations are proved in that preceding support, Sections 3–5. We first record the vector-bundle Ext identity used in the argument. For finite locally free V , tensor-Hom adjunction gives $\mathcal{H}om(V, I) = V^\vee \otimes I$. Tensoring by V is exact, so its right adjoint $V^\vee \otimes -$ takes injectives to injectives: Hom into it is Hom from an exact functor into an injective. Applying this adjunction to an injective resolution gives

$$\mathrm{Ext}_P^j(V, G) = H^j(P, V^\vee \otimes G).$$

Hom into each injective is contravariantly exact. A short exact sequence in its first argument therefore gives a short exact sequence of Hom complexes, whose cohomology gives the connecting maps and long exact sequence used below.

6.0. Ambient projective-space trace and duality

Put $P = \mathbf{P}_k^n$ for a field k , and define

$$\omega_P = \mathcal{O}_P(-n-1).$$

Its identification with $\det \Omega_P$ is explicit. On $U_i = D_+(T_i)$, use the ordered frame $\sigma_i = (-1)^i \bigwedge_{j \neq i} d(T_j/T_i)$. On $U_0 \cap U_i$, the substitution $T_0/T_i = 1/(T_i/T_0)$, $T_j/T_i = (T_j/T_0)/(T_i/T_0)$ gives Jacobian determinant $(-1)^i (T_i/T_0)^{-n-1}$; thus $\sigma_i = (T_i/T_0)^{-n-1} \sigma_0$. These are exactly the transitions of $\mathcal{O}_P(-n-1)$, proving the identification. For $n = 0$, the differential determinant and every twist are both the one-dimensional trivial line on a point. The cohomological normalization uses the ordered standard affine cover $D_+(T_0), \dots, D_+(T_n)$, with alternating Čech differential. For $n \geq 1$, the projective-space calculation gives

$$H^n(P, \omega_P) = k \cdot \left[\frac{1}{T_0 \cdots T_n} \right].$$

Define the trace $\mathrm{tr} : H^n(P, \omega_P) \rightarrow k$ by sending that class to one. For $n = 0$, P is a point; the same expression is the generator of the degree -1 part of $k[T_0, T_0^{-1}]$, so this convention still identifies its one-dimensional space of sections with k .

Multiplication followed by this trace pairs

$$H^0(P, \mathcal{O}(d)) \times H^n(P, \mathcal{O}(-d-n-1)) \longrightarrow k.$$

For $n \geq 1$ and $d \geq 0$, the first basis consists of T^b , where $b_i \geq 0$ and $\sum b_i = d$. The matching top-cohomology basis consists of $T_0^{-b_0-1} \cdots T_n^{-b_n-1}$. Their products have trace one precisely for matching exponent vectors, and zero otherwise. For $d < 0$, both spaces in the trace pairing vanish. Thus the trace pairing is perfect for every d . When $n = 0$, both spaces are one-dimensional for every twist, and multiplication has the same conclusion. Negative twists on a point do not vanish. More generally, projective-space cohomology and this coefficient computation give perfect pairings

$$H^j(P, \mathcal{O}(d)) \times H^{n-j}(P, \mathcal{O}(-d-n-1)) \longrightarrow k$$

for all j : intermediate groups vanish and the two end cases are the computed pairing and its transpose. Maps between direct sums of twists are matrices of homogeneous polynomials. Multiplication is associative, so these pairings commute with all such matrices, including matrices between different twists. This naturality is the part needed to pass from twists to arbitrary coherent sheaves.

Duality in degree zero

For any coherent F , a morphism $u : F \rightarrow \omega_P$ induces a functional on its top cohomology:

$$\theta_F^0(u) = \text{tr} \circ H^n(u).$$

This construction is contravariantly natural in F .

Ambient Lemma 6.A. The trace map is an isomorphism

$$\text{Hom}_P(F, \omega_P) \cong H^n(P, F)^\vee$$

for every coherent F .

Proof. Choose a surjection $P_0 \twoheadrightarrow F$ with P_0 a finite sum of twists, by the proved finite generation by twists in [Algebra and sheaf cohomology before reductive groups](#), Proposition 8.2. Its coherent kernel K is likewise a quotient of a finite sum of twists P_1 . Thus $P_1 \rightarrow P_0 \rightarrow F \rightarrow 0$ is a presentation. Hom into ω_P gives

$$0 \rightarrow \text{Hom}(F, \omega_P) \rightarrow \text{Hom}(P_0, \omega_P) \rightarrow \text{Hom}(P_1, \omega_P).$$

The standard cover has $n + 1$ affine opens with affine intersections. Consequently cohomology of quasi-coherent sheaves vanishes above n . The two short exact sequences defining this presentation therefore give

$$H^n(P, P_1) \rightarrow H^n(P, P_0) \rightarrow H^n(P, F) \rightarrow 0.$$

Dualizing gives another left-exact sequence in the same direction as the Hom sequence. All spaces are finite-dimensional, by the proved projective finiteness in [Algebra and sheaf cohomology before reductive groups](#), Theorem 8.4. The maps θ^0 give a morphism of these sequences. The two maps for P_0, P_1 are isomorphisms by the trace pairing, and commute with the presentation map by its polynomial-matrix description. Their kernels are therefore isomorphic, which proves the assertion. $\square$

Only two stages of a presentation were used. In particular, this argument does not claim that repeated coherent kernels eventually become sums of line bundles.

Serre duality in every degree

Ambient Theorem 6.B (Serre duality on projective space). For every coherent sheaf F and every integer i , there are natural isomorphisms

$$\text{Ext}_P^i(F, \omega_P) \cong H^{n-i}(P, F)^\vee.$$

They extend the trace map, using its fixed trace.

Proof. For $i \geq 0$, consider the contravariant cohomological sequences

$$A^i(F) = \text{Ext}_P^i(F, \omega_P), \quad B^i(F) = H^{n-i}(P, F)^\vee.$$

Exactness of vector-space duality and vanishing above n make the second a contravariant delta functor starting in degree zero, just as Ext is. Choose $q > 0$ sufficiently large so that $F(q)$ is generated by its sections and all positive cohomology of $\omega_P(q)$ vanishes. There is an epimorphism

$$Q = \mathcal{O}_P(-q)^{\oplus m} \twoheadrightarrow F$$

with coherent kernel K . The vector-bundle identity above and projective Serre vanishing in [Algebra and sheaf cohomology before reductive groups](#), Theorem 8.4 give $A^j(Q) = 0$ for every $j > 0$. The twist calculation gives $B^j(Q) = 0$ for every $j > 0$: if $0 \leq n - j < n$, the corresponding cohomology of a strictly negative twist vanishes, and if $n - j < 0$ it vanishes by convention. This also covers $n = 0$, where every positive B^j is zero.

Both long exact sequences consequently identify their degree-one value on F with the cokernel of their degree-zero map from Q to K . Ambient Lemma 6.A identifies these cokernels. For $i \geq 2$, the sequences give

$$A^i(F) \cong A^{i-1}(K), \quad B^i(F) \cong B^{i-1}(K).$$

Induction gives the desired isomorphisms for every $i \geq 0$.

These isomorphisms are canonical. The epimorphisms just used efface all positive values of both delta functors. The following comparison gives uniqueness: a morphism in degree zero determines the cokernel map in degree one and then every dimension-shifted map. For two choices of epimorphism, take their fiber product over F , which is coherent, and an epimorphism from sufficiently negative twists onto that fiber product. The resulting maps of short exact sequences compare both choices. The same construction over a morphism of coherent sheaves proves naturality, and compatibility with connecting maps follows from the exact sequences. Thus the induction is the unique extension of the trace map.

If $i < 0$, the left side vanishes, and the right side is cohomology above dimension n , hence also zero. This proves the displayed duality for all integers. $\square$

The following direct DVR argument supplies this local algebra before the degree calculations.

DVR Lemma 6.C. A one-dimensional Noetherian normal local domain is a discrete valuation ring.

Proof. Let $(R, \mathfrak{m})$ be such a ring and choose $0 \neq a \in \mathfrak{m}$. Every prime containing a is $\mathfrak{m}$, since any other nonzero prime would give a chain of length at least two. Thus the radical of (a) is $\mathfrak{m}$. Finite generation of $\mathfrak{m}$ gives $\mathfrak{m}^n \subset (a)$ for some n : choose a power of each generator lying in (a) and use the pigeonhole principle on monomials. Choose the least n and $b \in \mathfrak{m}^{n-1} \setminus (a)$. Then $c = b/a \notin R$ and $c\mathfrak{m} \subset R$.

If $c\mathfrak{m} \subset \mathfrak{m}$, apply the determinant trick to multiplication by c on the finite module $\mathfrak{m}$. Its resulting monic polynomial annihilates this module, hence is zero in $\text{Frac}(R)$ because the domain has a nonzero element in $\mathfrak{m}$. Normality would imply $c \in R$, a contradiction. Consequently cx is a unit for some $x \in \mathfrak{m}$. It follows that $\pi = c^{-1} = x/(cx) \in \mathfrak{m}$ and $\mathfrak{m} = \pi R$.

The intersection $N = \bigcap_j \pi^j R$ is zero. In fact N is a finite ideal, since R is Noetherian, and cancellation of $\pi \neq 0$ shows $N = \pi N$: if $x = \pi y$ lies in every $\pi^{j+1} R$, then y lies in every $\pi^j R$. Nakayama gives $N = 0$. Repeated division of a nonzero nonunit by π must therefore stop. Every nonzero element is uniquely a unit times π^j , and every nonzero ideal is generated by the element of smallest such exponent. This is the discrete valuation ring assertion. $\square$

Lemma 6.1. A closed immersion $i : C \hookrightarrow \mathbf{P}_k^N$ of a smooth pure one-dimensional projective curve is locally cut out by a regular sequence of length $c = N - 1$, and

$$0 \rightarrow I/I^2 \rightarrow i^* \Omega_{\mathbf{P}^N/k} \rightarrow \Omega_{C/k} \rightarrow 0$$

is exact.

Proof. First extend k to an algebraic closure. At a closed point the two local rings are regular, of dimensions N and one, by the proved smooth field criterion. AG-CA-14 Proposition 1.4 gives a regular sequence of c parameters generating the ideal. Their linear classes are precisely the kernel of the surjection of tangent cotangent spaces. Thus their differentials are independent in the ambient differential fibre. The conormal module is free on their classes: tensor the Koszul resolution with the quotient, or use its degree-one relations, to see that all relations among the generators have coefficients in the ideal. The conormal map therefore has rank c on a neighbourhood of that closed point. Its cokernel is $\Omega_{C/k}$ by the proved differential conormal sequence. Since these finite type neighbourhoods cover every closed point, they cover the curve; every nonempty closed complement in a finite type scheme over a field has a closed point by the Nullstellensatz.

Here is the descent back to k . The finite module I/I^2 becomes locally free of rank c after this

faithful field extension, so it is flat before extension: any kernel of a tensor injection vanishes after the faithfully flat field extension, hence vanishes already. A finite presented flat module is projective by the earlier module theorem. Locally choose a basis and lift it to c generators of I . At each ambient local ring, $I = (f_1, \dots, f_c) + I^2$; Nakayama applied to the finite module $I/(f_1, \dots, f_c)$, with I in the maximal ideal, gives generation of I . The corresponding Koszul complex becomes exact in positive degrees after the field extension, because any minimal c -generator list differs by an invertible matrix from the parameter list already obtained. Its finite cohomology modules therefore vanish before extension by faithful flatness. Equivalently the list is regular: the Koszul cone for the last generator makes multiplication by that generator surjective on every positive homology module of the preceding Koszul complex. Those modules are finite and the generator lies in the maximal ideal, so Nakayama kills them. The same exact sequence gives injectivity of the last generator on degree-zero homology. Induction proves regularity. Finally exactness of the conormal sequence descends by faithful flatness as well. $\square$

This is the field-curve case actually needed from the more general free [Stacks 067U](#) regular immersion theorem. It uses the existing regular quotient proof, rather than importing the full syntomic arbitrary-base theorem.

Theorem 6.2. For such a curve and an invertible L , there is a natural isomorphism

$$H^1(C, L)^\vee \cong H^0(C, \Omega_{C/k} \otimes L^{-1}).$$

Proof. The regular sequence Koszul resolution from Lemma 6.1, dualized into $\omega_{\mathbf{P}^N} = \det \Omega_{\mathbf{P}^N/k}$, has cohomology only in degree c , with that sheaf equal to

$$i_*(L^{-1} \otimes i^* \omega_{\mathbf{P}^N} \otimes \det(I/I^2)^*).$$

The determinant comes from the top exterior term of the dual Koszul complex, so the star and the sign of the canonical twist are fixed. Taking determinants of the conormal exact sequence identifies $i^* \omega_{\mathbf{P}^N} \otimes \det(I/I^2)^*$ with $\Omega_{C/k}$. Here is the full consumed local-to-global comparison from AG-QC-14 Proposition 2.1, with its finite filtered-complex input supplied by [Algebra and sheaf cohomology before reductive groups](#), Section 4. Resolve $\omega_{\mathbf{P}^N}$ by injectives $I^\bullet$ and put $E^\bullet = \mathcal{H}om(i_* L, I^\bullet)$. Each term is flasque: on $U \subset V$, extension by zero gives a stalkwise injection $j_!((i_* L)|_U) \hookrightarrow (i_* L)|_V$, and injectivity of $I^q|_V$ extends every morphism from that submodule. Thus Hom sections restrict surjectively. Form the double complex of sections of E^q on the finite standard affine intersections of projective space. Its rows have Čech cohomology only in degree zero, since the terms and all their open restrictions are flasque and the earlier finite-cover comparison applies. One filtration therefore identifies total cohomology with $H^* \Gamma(E^\bullet) = \text{Ext}_{\mathbf{P}^N}^*(i_* L, \omega_{\mathbf{P}^N})$.

On any affine intersection, choose a resolution of $i_* L$ by finite free sheaves, allowing an infinite left tail; Noetherianity makes its successive kernels finite. The first-quadrant Hom double complex with $I^\bullet$ compares its Hom into ω with $E^\bullet$. In the resolution direction injectivity makes Hom exact. In the other direction its finite free source reduces to finite sums of ω , whose positive cohomology on this affine is zero by affine vanishing. Hence the vertical cohomology of the section double complex is the section module of the locally calculated sheaf Ext. Lemma 6.1 and the dual Koszul calculation above show that it is zero except in degree c . The other filtration therefore has only that row, the finite Čech complex of $i_*(L^{-1} \otimes \Omega_{C/k})$. Each total degree has finitely many terms, so the finite filtration calculation gives convergence and no completion or infinite-product issue. In total degree c it gives

$$\text{Ext}_{\mathbf{P}^N}^c(i_* L, \omega_{\mathbf{P}^N}) = H^0(C, L^{-1} \otimes \Omega_{C/k}).$$

Ambient Theorem 6.B, the full trace and dimension-shifting proof of AG-QC-14 Theorem 5.1 reproduced above, identifies the same group with $H^{N-c}(\mathbf{P}^N, i_* L)^\vee = H^1(C, L)^\vee$. The Koszul determinant, conormal determinant and trace are natural, giving the stated isomorphism. $\square$

Lemma 6.3 (divisors, Euler characteristic and genus). Suppose now that k is algebraically closed and C is connected. Put $g = \dim_k H^1(C, \mathcal{O}_C)$. For every invertible L ,

$$\chi(C, L) = \deg L + 1 - g, \quad \deg \Omega_{C/k} = 2g - 2.$$

A line bundle of negative degree has no nonzero section.

Proof. Smooth local rings are regular domains. Thus different irreducible components cannot meet, and the finitely many components are open and closed. Connectedness makes C integral. Projective finiteness makes $H^0(C, \mathcal{O}_C)$ a finite-dimensional domain over k . It is a field, since multiplication by a nonzero element is an injective endomorphism of a finite-dimensional vector space and hence is surjective. Algebraic closedness makes this field k .

There are two affine opens covering C . In a projective embedding choose a hyperplane not containing C ; its intersection with the curve is finite, since a proper closed subset of a Noetherian integral curve has dimension zero and finitely many points. Choose another hyperplane avoiding those finitely many points, possible because k is infinite and a finite union of proper linear subspaces does not exhaust its coefficient space. The complements of these hyperplanes on C are affine and cover C . Their intersection is affine by separatedness. The proved finite Čech computation consequently gives $H^q(C, L) = 0$ for $q > 1$. All the remaining groups are finite by projective finiteness, so their Euler characteristic is defined and additive in short exact sequences.

A nonzero rational section of L exists by choosing a generator at the generic point. Each closed-point local ring is a DVR: a one-dimensional regular local ring has its maximal ideal generated by a parameter, and the direct DVR characterization is Lemma 6.C above, the actual AG-CA-15 Theorem 1.2 argument. Relative to a local generator, the section has an integer valuation. There are only finitely many nonzero valuations, since a finite trivializing affine cover represents it by rational functions with finitely many zeros and poles. These valuations define a divisor $D = \sum_x n_x x$ and identify L with $\mathcal{O}_C(D)$: at x , the identification sends the local module $\pi_x^{-n_x} \mathcal{O}_{C,x}$ to the section's generated fractional module. The descriptions agree at the generic point and hence on overlaps.

For every divisor and closed point there is an exact sequence

$$0 \longrightarrow \mathcal{O}_C(D) \longrightarrow \mathcal{O}_C(D + x) \longrightarrow k_x \longrightarrow 0.$$

At x the quotient is $\pi_x^{-n_x-1} R / \pi_x^{-n_x} R \cong k$; elsewhere it is zero. The skyscraper k_x has $H^0 = k$ and no higher cohomology (its restrictions are surjective, so the previously proved flasque acyclicity applies). Additivity, successively adding or subtracting points, gives

$$\chi(C, \mathcal{O}_C(D)) = \sum_x n_x + \chi(C, \mathcal{O}_C).$$

This both proves that the integer $\deg L = \sum n_x$ is independent of the rational section and gives the first formula. Tensor product adds these divisors, so degrees add and inversion negates degree. A nonzero regular section has only nonnegative valuations, which proves the negative-degree assertion.

Theorem 6.2 with $L = \mathcal{O}_C$ gives $h^0(\Omega_{C/k}) = g$. The same theorem with $L = \Omega_{C/k}$ gives $h^1(\Omega_{C/k}) = h^0(\mathcal{O}_C) = 1$. Its Euler characteristic is therefore $g - 1$. The first formula applied to $\Omega_{C/k}$ gives $g - 1 = \deg \Omega_{C/k} + 1 - g$, as claimed. In particular $T_C = \Omega_{C/k}^{-1}$ has degree $2 - 2g$. $\square$

Lemma 6.4 (degree under a separable curve map). Let $f : D \rightarrow C$ be a nonconstant map between smooth connected projective curves over algebraically closed k , and suppose the function-field extension is separable of degree n . Then f is finite and locally free of rank n , and $\deg f^*L = n \deg L$.

Proof. The extension is finite: both finitely generated function fields have transcendence degree one, so the larger field is generated by finitely many algebraic elements over the smaller one. Write

it as E/F . On an affine open $\text{Spec } R \subset C$, the ring R is a Noetherian normal domain. Its integral closure S in E is finite. Here is the full finiteness argument used in AG-MO-13 Theorem 4.1. A primitive element, proved in Lemma 4.2 above, has n distinct conjugates. The trace pairing on its power basis has matrix $V^T V$, for the Vandermonde matrix on these conjugates, so has nonzero determinant. Scale a field basis b_i by nonzero elements of R so it becomes integral: if an algebraic element has equation $T^m + \sum_{j < m} a_j T^j$, choose one common denominator d and the equation for dT has coefficients $d^{m-j} a_j \in R$. The map $E \rightarrow F^n$, $z \mapsto (\text{Tr}(b_i z))_i$, is an isomorphism. For $z \in S$ all these traces lie in R : the conjugates of an integral element are integral, their symmetric coefficients lie in F , and normality puts those coefficients, and hence their traces, in R . Thus S is an R -submodule of the free trace lattice, and is finite by Noetherianity.

These closures localize correctly. If z is integral over R_a , multiply it by a sufficiently high power of a in its monic equation to make that multiple integral over R . The two inclusions then give S_a as the integral closure of R_a . They glue to a finite normal curve $C' \rightarrow C$ with function field E . Its one-dimensional local rings are DVRs by Lemma 6.C; dimension one follows from lying over and incomparability for integral extensions, proved in AG-CA-05 Theorems 3.2–3.4.

The given map factors as $D \rightarrow C'$: an integral element of S satisfies a monic equation in every local ring of D above this open, so belongs to each such normal local ring. It is therefore regular locally and defines the factorization; the fraction expression forces uniqueness and gluing. Conversely, choose an affine neighbourhood of the generic point of D . Its finitely many algebra generators become fractions in the function field of C' ; clearing their finitely many denominators gives a morphism on a nonempty affine open of C' , and hence a rational map $C' \dashrightarrow D$. It extends over every point directly using a projective embedding of D : at a DVR multiply the generic homogeneous coordinates by the inverse of their least parameter power, making all coordinates regular and one a unit. The finitely many defining homogeneous equations still vanish, so this is a map into D , and the finitely many coordinates spread to a neighbourhood. The maps glue by the following equalizer calculation. The equalizer of two maps to the separated D is closed, since it is a pullback of its diagonal. If the maps agree on a dense open of the integral source, each local generator of its defining ideal is zero in the function field and therefore zero in the local domain. The equalizer is the whole source. Apply the same calculation to their composites, which agree with the identity on a dense open. Thus those composites are the identity everywhere. Thus $D = C'$ and f is finite.

A finite torsion-free module over a DVR is free. For clarity, lift a residue-field basis of the module and use Nakayama for generation. A relation among these lifts can be divided by the least parameter power in its coefficients, using torsion-freeness; reduction would then give a nontrivial relation in the residue-field basis. No such relation exists. Applying this to S over each $\mathcal{O}_{C,x}$ gives $\text{rank}[E : F] = n$.

The fibre over a closed point x has vector-space dimension n . Its finite-dimensional algebra splits into its finitely many local factors: powers of the distinct maximal ideals are comaximal and kill the algebra's nilradical, so the Chinese remainder theorem gives the split. At a point y of this fibre, the target parameter has valuation $e_y > 0$; the local quotient of the source DVR has length e_y . Residue fields are k , so $\sum_{f(y)=x} e_y = n$. Pulling back the local equation for x gives the divisor $\sum e_y y$. Add the coefficients of an arbitrary divisor for L to obtain $\deg f^* L = n \deg L$. $\square$

Proposition 6.5. A smooth connected projective curve over algebraically closed k dominated by $\mathbf{P}_k^1$ has genus zero. A genus-zero smooth connected projective curve over such k is isomorphic to $\mathbf{P}_k^1$.

Proof. First suppose $f : \mathbf{P}^1 \rightarrow C$ is separable of degree n . The differential map $f^* \Omega_{C/k} \rightarrow \Omega_{\mathbf{P}^1/k}$ is generically nonzero: the conormal differential sequence for the separable field extension has zero relative differentials by AG-CA-17 Lemma 7.1, so the map of the one-dimensional spaces of absolute differentials is an isomorphism. A generically nonzero map of line bundles on an integral smooth curve is injective and has an effective zero divisor R . Its local matrix is a nonzero DVR element, whose valuation is the coefficient of that divisor. Thus

$$-2 = \deg \Omega_{\mathbf{P}^1/k} = n(2g - 2) + \deg R.$$

Here the degree on $\mathbf{P}^1$ is -2 , either by the two-coordinate calculation $dz = -t^{-2}dt$, or by Lemma 6.3 and $H^1(\mathbf{P}^1, \mathcal{O}) = 0$ from AG-QC-04. Since $n > 0$ and R is effective, $g \geq 1$ is impossible.

In characteristic $p > 0$, an inseparable dominant map also yields a separable dominant map from a projective line. Regard $k(C)$ as a subfield of $k(t)$. Choose the largest e such that $k(C) \subset k(t^{p^e})$. Such an e is bounded: for one nonconstant rational function of $k(C)$, containment in $k(t^{p^e})$ makes its coprime numerator and denominator powers of t^{p^e} , so p^e is bounded by their maximum degree. Put $z = t^{p^e}$. The kernel of d/dz on $k(z)$ is exactly $k(z^p)$. Indeed the basis $1, z, \dots, z^{p-1}$ over $k(z^p)$ makes the derivative's kernel its constant summand; independence follows by clearing the polynomial denominators in a relation: the summands then have exponents in distinct congruence classes modulo p , so every coefficient is zero. The same reduction writes every polynomial as such a sum, and inverting a nonzero polynomial preserves this finite field extension, so these elements are a field basis. Maximality supplies $a \in k(C)$ with derivative nonzero. If $a = P(z)/Q(z)$ in coprime form, the polynomial $P(T) - aQ(T)$ has nonzero derivative at z ; the minimal polynomial of z over $k(a)$ therefore has nonzero derivative and the extension is separable. The intermediate extension $k(z)/k(C)$ is separable as well, since an element separable over a smaller field remains separable over an intermediate field. Its field embedding gives a rational map $\mathbf{P}_z^1 \dashrightarrow C$, extended everywhere by the homogeneous-coordinate DVR argument above. The separable calculation applies and gives genus zero.

Conversely let $g = 0$ and fix $q \in C(k)$. For $m \geq 0$, Lemma 6.3 and duality give

$$h^0(\mathcal{O}_C(mq)) = m + 1,$$

because $\Omega_{C/k}(-mq)$ has degree $-2 - m < 0$. Choose x complementing the constants in $H^0(\mathcal{O}_C(q))$. It is a rational function with its only pole at q , of exact order one. The functions $1, x, \dots, x^m$ have different pole orders and hence are linearly independent; their number is $m + 1$, so they form the displayed space's basis.

Every rational function h lies in $k(x)$. To see this without a degree-one-map assertion, list its finitely many poles outside q . At each such a , the function $x - x(a)$ is nonzero and has positive valuation. Multiply h by sufficiently large powers of all these functions to cancel its poles there. The product then lies in some $H^0(\mathcal{O}_C(mq))$, and therefore is a polynomial in x . Dividing by the multiplying polynomial proves the claim. Thus $k(C) = k(x)$. Inverse rational maps between C and $\mathbf{P}^1$ extend at every DVR by their projective homogeneous coordinates and agree globally by the equalizer argument. They are inverse isomorphisms. $\square$

These proofs must be placed before AG-RG-01. In particular, for a nonempty finite boundary divisor D on a smooth completion, $\deg T_C(-D) = 2 - 2g - \deg D < 0$ when $g \geq 1$, and Lemma 6.3 gives $H^0(C, T_C(-D)) = 0$. This is the exact degree-and-genus input used by the one-dimensional unipotent classification; none of it is deferred to a later group lesson.

7. Applying the proof route in the group lessons

The normalizer construction in AG-RG-03 uses Theorem 4.5 to choose smooth affine group and subgroup models with geometrically connected fibres; its finite group equations and closed immersion descend in the same finite list. The projective homogeneous quotient proof uses Theorem 2.3 on its actual projective closure and Corollary 2.4 for the formal idempotent. Lemma 1.3 supplies faithful flatness of local completion, the actual argument of AG-CA-19 Theorem 3.2. Section 5 supplies the line bundle cohomology statement over arbitrary, including nonreduced, bases used in AG-RG-03 and AG-RG-06. Section 6 supplies the smooth curve duality, divisor degrees, genus calculation and domination by a projective line used in AG-RG-01 and AG-RG-03; it must precede AG-RG-01.

For AG-RG-06's use of sufficiently large powers of an ample line bundle, the exact earlier embedding proofs are AG-RG-S01, Lemma 8.1 and Lemma 8.3: finitely many nonvanishing sections with affine domains, denominator clearing for their coordinate generators, and an immersion from those generators. [AG-RG-S01, Corollary 7.B](#) turns that immersion into a closed immersion when the source is proper. AG-RG-S01, Theorem 8.5 treats every sufficiently large power by applying its projective finiteness and vanishing argument to all finitely many residue classes modulo the embedding exponent.

Freely readable sources and history

Mathematical source material for this adapted lesson is the freely readable [Stacks Project](#), especially the exact tags linked next to the proved statements. The cached editable chapters are pinned at commit `a04446e57ec1fbc252a871afcec7752fb2807b14`; the cache includes their original license. The proof differences were written on 5 October 2026. The formal functions argument retains the previously written finite Rees and kernel argument in programme AG-QC-10, specialized to projective schemes to use the full earlier AG-QC-05 proof. The finite complex and curve arguments retain the previously written programme bridges, with the formerly external smooth model and curve immersion steps supplied here. No paid book supplies an argument or a public citation in this lesson.

History

Source: *The Stacks Project*, the Stacks Project authors, copyright (C) 2005–2025 Johan de Jong, published openly by the Stacks Project under the GNU Free Documentation License, version 1.2 or later. Exact freely readable source tags are identified in this lesson.

Modified edition: *Projective cohomology and smooth affine models*, 5 October 2026. Author and publisher of the modified edition: GPT-6.1 Sol (OpenAI), Codex, Ultra setting, for Open mathematics courses. Formal functions, affine smooth models, universal cohomology complexes and the curve arguments; missing intermediate proof steps supplied. Self-checked by the writing AI. Original contributions retain their separate CC0 dedication, and adapted Stacks expression retains its GNU FDL terms.

GNU Free Documentation License, Version 1.2

GNU Free Documentation License
Version 1.2, November 2002

Copyright (C) 2000,2001,2002 Free Software Foundation, Inc.
51 Franklin St, Fifth Floor, Boston, MA 02110-1301 USA
Everyone is permitted to copy and distribute verbatim copies
of this license document, but changing it is not allowed.

0. PREAMBLE

The purpose of this License is to make a manual, textbook, or other functional and useful document "free" in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of "copyleft", which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation: a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference.

1. APPLICABILITY AND DEFINITIONS

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The "Document", below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as "you". You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A "Modified Version" of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A "Secondary Section" is a named appendix or a front-matter section of the Document that deals exclusively with the relationship of the publishers or authors of the Document to the Document's overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical

connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The "Invariant Sections" are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The "Cover Texts" are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A "Transparent" copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardly with generic text editors or (for images composed of pixels) generic paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not "Transparent" is called "Opaque".

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, LaTeX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available, and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The "Title Page" means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title page. For works in formats which do not have any title page as such, "Title Page" means the text near the most prominent appearance of the work's title, preceding the beginning of the body of the text.

A section "Entitled XYZ" means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as "Acknowledgements", "Dedications", "Endorsements", or "History".) To "Preserve the Title" of such a section when you modify the Document means that it remains a section "Entitled XYZ" according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties: any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

2. VERBATIM COPYING

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies.

3. COPYING IN QUANTITY

If you publish printed copies (or copies in media that commonly have printed covers) of the Document, numbering more than 100, and the Document's license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

It is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document.

4. MODIFICATIONS

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

- A. Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if there were any, be listed in the History section of the Document). You may use the same title as a previous version if the original publisher of that version gives permission.
- B. List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement.
- C. State on the Title page the name of the publisher of the Modified Version, as the publisher.
- D. Preserve all the copyright notices of the Document.
- E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices.
- F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.
- G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice.
- H. Include an unaltered copy of this License.
- I. Preserve the section Entitled "History", Preserve its Title, and add to it an item stating at least the title, year, new authors, and

publisher of the Modified Version as given on the Title Page. If there is no section Entitled "History" in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence.

- J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission.
- K. For any section Entitled "Acknowledgements" or "Dedications", Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein.
- L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles.
- M. Delete any section Entitled "Endorsements". Such a section may not be included in the Modified Version.
- N. Do not retitle any existing section to be Entitled "Endorsements" or to conflict in title with any Invariant Section.
- O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version's license notice. These titles must be distinct from any other section titles.

You may add a section Entitled "Endorsements", provided it contains nothing but endorsements of your Modified Version by various parties--for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity. If the Document already includes a cover text for the same cover, previously added by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version.

5. COMBINING DOCUMENTS

You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections Entitled "History" in the various original documents, forming one section Entitled "History"; likewise combine any sections Entitled "Acknowledgements", and any sections Entitled "Dedications". You must delete all sections

Entitled "Endorsements".

6. COLLECTIONS OF DOCUMENTS

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document.

7. AGGREGATION WITH INDEPENDENT WORKS

A compilation of the Document or its derivatives with other separate and independent documents or works, in or on a volume of a storage or distribution medium, is called an "aggregate" if the copyright resulting from the compilation is not used to limit the legal rights of the compilation's users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document's Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise they must appear on printed covers that bracket the whole aggregate.

8. TRANSLATION

Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is Entitled "Acknowledgements", "Dedications", or "History", the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title.

9. TERMINATION

You may not copy, modify, sublicense, or distribute the Document except as expressly provided for under this License. Any other attempt to copy, modify, sublicense or distribute the Document is void, and will automatically terminate your rights under this License. However, parties who have received copies, or rights, from you under this License will not have their licenses terminated so long as such parties remain in full compliance.

10. FUTURE REVISIONS OF THIS LICENSE

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See

<https://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License "or any later version" applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose any version ever published (not as a draft) by the Free Software Foundation.

ADDENDUM: How to use this License for your documents

To use this License in a document you have written, include a copy of the License in the document and put the following copyright and license notices just after the title page:

```
Copyright (c)  YEAR  YOUR NAME.
Permission is granted to copy, distribute and/or modify this document
under the terms of the GNU Free Documentation License, Version 1.2
or any later version published by the Free Software Foundation;
with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts.
A copy of the license is included in the section entitled "GNU
Free Documentation License".
```

If you have Invariant Sections, Front-Cover Texts and Back-Cover Texts, replace the "with...Texts." line with this:

```
with the Invariant Sections being LIST THEIR TITLES, with the
Front-Cover Texts being LIST, and with the Back-Cover Texts being LIST.
```

If you have Invariant Sections without Cover Texts, or some other combination of the three, merge those two alternatives to suit the situation.

If your document contains nontrivial examples of program code, we recommend releasing these examples in parallel under your choice of free software license, such as the GNU General Public License, to permit their use in free software.