

Roots and reductive groups of rank one

GPT-6.1 Sol (OpenAI)
Codex, Ultra setting

October 2026

Written and self-checked by the writing AI. No human or independent review is claimed. Original contributions are dedicated under CC0 1.0. Attributed Stacks adaptations retain their GNU Free Documentation License obligations, as recorded for each component. The cumulative collection includes the complete version 1.2 licence below.

Online course: <https://kokunoyumeto.github.io/open-math-courses/courses/AG-RG/>.

Contents

3	Roots and reductive groups of rank one	1
1.	The centre and the two ranks	1
2.	Why rank one has only two moving directions	3
3.	Subgroups obtained by taking a limit	6
4.	Root groups over the base	6
5.	A projective quotient and its rank-one form	7
6.	Splitting a central extension of the rank-one cover	14
7.	Coroots from multiplication	16
8.	The three rank-one groups	17
9.	Exercises and solutions	19
	References and prerequisite proofs	20
	History	21
	GNU Free Documentation License, Version 1.2	22

Lesson 3

Roots and reductive groups of rank one

Written by GPT-6.1 Sol (OpenAI) in Codex, Ultra setting, October 2026. Self-checked by the writing AI, GPT-6.1 Sol (OpenAI). Original exposition is public domain (CC0). The collected lesson, including the explicitly attributed Stacks passage, is also distributed under the GNU Free Documentation License 1.2.

The diagonal entries of a matrix act on its off-diagonal entries through ratios. Those ratios are the roots of a general linear group. For an arbitrary reductive group, roots are still characters of a maximal torus, but their one-dimensional spaces need not be globally trivial over the base. This lesson constructs the corresponding groups without using a power-series exponential, and shows why every root carries a copy of the rank-one geometry of SL_2 .

We use torus conjugacy, torus deformation, and the centralizer theorem from the preceding two lessons. A reductive S -group is smooth affine of finite presentation with connected reductive geometric fibres. A semisimple S -group has semisimple geometric fibres. Statements about line bundles and subgroup schemes are made over arbitrary schemes, including nonreduced ones.

1. The centre and the two ranks

Let T be a split maximal torus of G , and, on an open set where the ranks are constant, write

$$\mathfrak{g} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_{\alpha}.$$

For now Φ means the nonzero weights; we will prove that they form a reduced root system. Put

$$Z = \bigcap_{\alpha \in \Phi} \ker(\alpha : T \rightarrow \mathbf{G}_m).$$

The group Z is of multiplicative type. It is precisely the scheme-theoretic centre of G . Indeed, its conjugation action on $\mathfrak{g}$ is trivial. Exactness of invariants for a multiplicative-type group makes $C_G(Z)$ smooth with Lie algebra $\mathfrak{g}$. Its geometric fibres are therefore all of G ; the smooth fibrewise isomorphism argument of the preceding lesson gives $C_G(Z) = G$. Conversely any central point belongs to $C_G(T) = T$ and acts trivially on every weight summand, hence lies in Z . These constructions agree on overlaps and descend when T is only étale locally split.

The centre need not be smooth: $Z(\mathrm{SL}_n) = \mu_n$, including when the characteristic divides n . Its largest subtorus is called the **radical**, denoted $\mathrm{Rad}(G)$. For a group of multiplicative type with character sheaf M , this subtorus has character sheaf M/M_{tors} . Thus its rank is locally constant

and its formation commutes with base change. The **rank** of G is the dimension of a maximal torus; its **semisimple rank** is

$$\text{rank}(G) - \dim \text{Rad}(G).$$

Over an algebraically closed field, the radical also equals the largest smooth connected solvable normal subgroup. Its unipotent radical is normal in G and is trivial by reductivity, so it is a torus; rigidity of torus automorphisms makes it central. Conversely a central torus is connected solvable normal. The quotient by this torus is semisimple: the inverse image of a connected solvable normal subgroup of the quotient would enlarge the radical.

The derived group will be constructed for arbitrary bases after the pinned existence theorem in [Pinnings and the classification of split reductive groups](#). It is semisimple and the multiplication map

$$\text{Rad}(G) \times G_{\text{der}} \longrightarrow G$$

is a central isogeny. Its construction uses the root groups established here. We do not assume a relative derived group in constructing these groups. Central quotients are constructed by graded invariant algebras in Theorem 1.1 below, without using coroots. In rank one we will construct the simply connected cover explicitly below.

Central quotients before the rank-one argument

The rank-one argument below needs the affine quotient even when the centre is not smooth. The invariant-algebra construction is independent of coroots and root groups.

Theorem 1.1 (affine central quotient). Let G be a reductive group over a scheme S , and let $H \subset Z(G)$ be a closed subgroup of multiplicative type of finite presentation. The fppf quotient G/H is an affine reductive S -group. Formation of the quotient commutes with arbitrary base change.

Proof. Work étale locally on $S = \text{Spec } R$, with $H = D_R(M)$. Its character group M is finitely generated. Right translation gives a grading $A = \Gamma(G, \mathcal{O}_G) = \bigoplus_{m \in M} A_m$. Put $B = A_0$ and $Q = \text{Spec } B$. Invariants commute with any base change because they are a direct grading summand. The same summand proves that B is flat over R .

We check finite presentation. The data descend to a finitely generated $\mathbf{Z}$ -algebra, including the smooth group, the closed embedding of $D(M)$, and centrality; all are finite-presentation equations. We may therefore assume R Noetherian. Choose finitely many homogeneous generators of A as an R -algebra and a graded surjection $P = R[z_1, \dots, z_n] \rightarrow A$. Its invariant part surjects onto B . The monomials of weight zero in P are finitely generated as a semigroup: their indecomposable exponent vectors form an antichain in $\mathbf{N}^n$, since if $u \leq v$ are two such vectors, $v - u$ also has weight zero. Every antichain is finite. The latter assertion follows inductively on n by extracting a nondecreasing subsequence of the first coordinates from any infinite sequence, then applying the induction to the remaining coordinates. Every zero-weight vector is a sum of indecomposables by induction on its coordinate sum. Thus P_0 , and then B , are finitely generated. They are finitely presented over Noetherian R . Descent and base change now give finite presentation in general.

Next we prove that $q : G \rightarrow Q$ is an H -torsor. First suppose $R = k$ is algebraically closed. The closed embedding $H \hookrightarrow G$ gives a graded surjection $A \rightarrow k[M]$. For each of finitely many generators m_i of M , choose elements $a_i \in A_{m_i}$ and $b_i \in A_{-m_i}$ restricting to the corresponding characters of H . Then $f = \prod_i a_i b_i \in B$ has value one at the identity. In A_f , every a_i and b_i is a unit. Each homogeneous component is therefore a free rank-one B_f -module generated by a homogeneous unit; multiplication of components is an isomorphism. Explicitly the map

$$A_f \otimes_{B_f} A_f \longrightarrow A_f \otimes_k k[M], \quad a \otimes c_m \longmapsto ac_m \otimes e_m$$

is an isomorphism: for a homogeneous unit u_m its inverse sends $a \otimes e_m$ to $au_m^{-1} \otimes u_m$. Also $A_f = \bigoplus_m B_f u_m$ is faithfully flat over B_f . This proves the torsor assertion on $D(f)$.

Left translation by $g \in G(k)$ preserves the grading and gives the same proof on $D(f_g)$, where $f_g(x) = f(g^{-1}x)$. These opens cover Q . Indeed $B \hookrightarrow A$ has a B -linear retraction, so $A \otimes_B \kappa(y)$ is nonzero for every prime y of B and q is surjective. A closed point of Q has a k -point g above it, and $f_g(g) = 1$. Since Q is of finite type over k , an uncovered nonempty closed subset would contain a closed point. Hence there is no uncovered point.

Return to general R . Both G and Q are flat of finite presentation over R , and the geometric-fibre morphisms are faithfully flat torsors by the preceding calculation. The fibre criterion for flatness gives that q is flat and of finite presentation; it is surjective by the invariant retraction. The natural morphism $G \times H \rightarrow G \times_Q G$ is an isomorphism on geometric fibres. Both sides are flat of finite presentation over R . The fibre criterion applied to this morphism makes it étale, and its fibrewise bijectivity makes it an isomorphism. Thus q is an fppf torsor. Centrality permits multiplication and inversion to descend, making Q a group; this also identifies Q with the fppf quotient.

Its geometric fibres are smooth connected reductive groups. Here is a proof of smoothness and reductivity that does not assume a quotient theorem. Over an algebraically closed field, A is a normal domain, and its invariant subring B is normal: an element of $\text{Frac } B$ integral over B is integral over A , hence in A , and is invariant, hence in B . A finite-type normal group over a perfect field has a nonempty smooth open subset; translation carries a smooth point to every closed point, so it is smooth. It is connected because $G \rightarrow Q$ is surjective. If its smooth connected unipotent radical were nontrivial, its inverse image would be a solvable normal subgroup of G . The reduced identity component of that inverse image is smooth, connected, solvable and normal, so is a central torus by reductivity. It surjects onto the radical: the finitely many components of the inverse image cannot map onto a connected group with the identity component having smaller dimension. A torus has trivial image in a unipotent group, a contradiction. Flatness, finite presentation and these smooth geometric fibres prove smoothness of Q/S .

□

2. Why rank one has only two moving directions

We first prove the geometric rank-one statement, rather than assuming a root classification. Work over an algebraically closed field k .

Lemma 2.1. If a torus acts linearly on an irreducible projective variety X , it has at least $\dim X + 1$ fixed points.

Proof. Choose a cocharacter separating the finitely many weights of an equivariant projective embedding. Induct on $\dim X$. Choose a lowest-weight basis vector and the complementary coordinate hyperplane. If X lies in that hyperplane, remove the basis vector and repeat. Otherwise the hyperplane section has dimension $\dim X - 1$ and has at least $\dim X$ fixed points by induction on an irreducible component. The affine complementary chart has nonnegative cocharacter weights. Sending its cocharacter parameter to zero produces another fixed point, outside the hyperplane. Separation of the torus weights makes it torus-fixed, not merely cocharacter-fixed. The base case is a point. □

The curve duality used in rank one

Lemma (curve duality). If C is a smooth projective pure one-dimensional curve over a field k and L is invertible, then naturally

$$H^1(C, L)^* \simeq H^0(C, \Omega_{C/k} \otimes L^{-1}).$$

Proof. Embed $i : C \hookrightarrow P = \mathbf{P}_k^N$, and put $c = N - 1$. The regular-immersion calculation in [Projective cohomology and smooth affine models](#), Lemma 6.1, proves over the original field, including imperfect fields, that I/I^2 is locally free of rank c , that the ideal is locally generated by a regular sequence, and that

$$0 \longrightarrow I/I^2 \longrightarrow i^* \Omega_{P/k} \longrightarrow \Omega_{C/k} \longrightarrow 0$$

is exact. Its proof first calculates at geometric closed points, then descends the conormal module and the exact Koszul complex by faithful field extension.

Dualizing that Koszul resolution, with a local frame of L , gives

$$\mathcal{E}xt_P^q(i_* L, \omega_P) = 0 \quad (q \neq c), \quad \mathcal{E}xt_P^c(i_* L, \omega_P) = i_*(L^{-1} \otimes i^* \omega_P \otimes \det(I/I^2)^*).$$

Here $\omega_P = \det \Omega_{P/k} = \mathcal{O}_P(-N - 1)$. Taking determinants of the conormal sequence identifies the sheaf in degree c with $i_*(L^{-1} \otimes \Omega_{C/k})$. The finite-cover Hom double complex written in S02 Theorem 6.2 has only this nonzero sheaf-Ext row, and therefore gives

$$\mathrm{Ext}_P^c(i_* L, \omega_P) = H^0(C, L^{-1} \otimes \Omega_{C/k}).$$

S02 Ambient Theorem 6.B proves projective-space duality by the monomial trace, a two-stage twist presentation and dimension shifting; it identifies the same Ext group with $H^{N-c}(P, i_* L)^* = H^1(C, L)^*$. Its trace sends the class of $T_0^{-1} \cdots T_N^{-1}$ to one. All the Koszul, determinant, finite-cover and trace comparisons are natural. This proves the asserted isomorphism. $\square$

Lemma 2.2 (the curve argument). Over an algebraically closed field, a smooth projective curve dominated by $\mathbf{P}^1$ is isomorphic to $\mathbf{P}^1$.

Proof. The preceding curve-duality lemma identifies $H^1(C, L)^*$ with $H^0(C, \Omega_{C/k} \otimes L^{-1})$.

Here are the remaining curve calculations. For any divisor D , adding one point in the divisor exact sequence changes the Euler characteristic of $\mathcal{O}_C(D)$ by one. Thus, with $g = h^1(\mathcal{O}_C)$,

$$\chi(L) = \deg L + 1 - g.$$

Every line bundle on a smooth curve has a rational section and is of this form. Duality gives $h^0(\omega_C) = g$ and $h^1(\omega_C) = 1$, so $\deg \omega_C = 2g - 2$. A nonzero section of a line bundle has an effective zero divisor; hence a negative-degree line bundle has no nonzero section.

For a nonconstant separable morphism $f : \mathbf{P}^1 \rightarrow C$ of degree n , the map $f^* \Omega_C \rightarrow \Omega_{\mathbf{P}^1}$ is a nonzero map of line bundles. Its zero divisor R is effective, and taking degrees gives

$$-2 = n(2g - 2) + \deg R.$$

Pullback multiplies degree by n : for a point divisor this is the sum of the lengths of the finite flat fibres over the discrete valuation ring, and the general assertion follows by addition. The displayed equality forces $g = 0$.

The inseparable case reduces to this one without any genus assertion about inseparable maps. In characteristic p , view $k(C)$ inside $k(t)$, and choose the largest e such that $k(C) \subset k(t^{p^e})$. It exists, since any fixed nonconstant element has finite rational-function degree, divisible by p^e if it lies in that subfield. Put $z = t^{p^e}$. Some element $f(z)$ of $k(C)$ has nonzero derivative: the kernel of the derivative on $k(z)$ is $k(z^p)$, as follows by writing $k(z)$ in the basis $1, z, \dots, z^{p-1}$ over $k(z^p)$. The extension $k(z)/k(f)$ is separable, by differentiating its rational-function equation, so $k(z)/k(C)$ is

separable. It gives a nonconstant separable morphism from the smooth projective curve with field $k(z)$, namely $\mathbf{P}^1$, to C . We have again proved $g = 0$.

Choose $q \in C(k)$. The line bundle $L = \mathcal{O}_C(q)$ has degree one. Duality and $\deg \omega_C = -2$ give $h^1(L) = 0$ and $h^0(L) = 2$. For any point x , the same calculation for $L(-x)$ gives $h^0(L(-x)) = 1$. Thus L is generated by its two sections. They define a nonconstant map $C \rightarrow \mathbf{P}^1$ whose pullback of $\mathcal{O}(1)$ has degree one. Its degree is one, so it is a finite birational morphism between smooth curves and hence an isomorphism, by their integrally closed local rings. $\square$

Proposition 2.3. A semisimple group of rank one has exactly two Borel subgroups containing a maximal torus, its Borel quotient is $\mathbf{P}^1$, and the action on that quotient has image PGL_2 and kernel its centre.

Proof. The normalizer quotient acts faithfully on the rank-one torus, whose automorphism group is $\{1, -1\}$. It acts simply transitively on the torus-fixed points of the homogeneous projective quotient G/B , as proved in the preceding lesson without assuming the full Borel-normalizer theorem. There are at most two such points. There cannot be one: Lemma 2.1 would then make this quotient zero-dimensional, so the connected group would be solvable, contradicting nontrivial semisimplicity. There are therefore two, and Lemma 2.1 makes the quotient a smooth projective curve.

The kernel of the action is the intersection of the conjugates of a Borel. Its reduced identity component is solvable and normal, so is trivial by semisimplicity. Thus the action kernel is finite. In particular a positive-dimensional Borel acts nontrivially on the curve. A connected solvable group over k is a rational variety: its torus times its split unipotent radical has an open affine parametrization, as in the solvable-group discussion in the first lesson. A nonconstant orbit map therefore restricts to a nonconstant rational map from an affine line, by fixing all but one of these coordinates. Since k is infinite, the fixed coordinates can be chosen outside the finitely many vanishing conditions in its rational functions. Properness extends this map to $\mathbf{P}^1$. Lemma 2.2 identifies the curve with $\mathbf{P}^1$.

The natural map $\mathrm{PGL}_2 \rightarrow \mathrm{Aut}(\mathbf{P}^1)$ is an isomorphism. To verify it over a ring, an automorphism carries the three pairwise fibrewise disjoint sections $0, 1, \infty$ to another such triple. Locally, a 2×2 change of basis carries $0, \infty$ to the first and third members; diagonal scaling then carries 1 to the second. The resulting projective matrix glues uniquely. An automorphism fixing 0 and ∞ preserves their Cartier divisors; the ratio of its pullback of the coordinate to that coordinate is a global unit on $\mathbf{P}^1$, hence a unit of the base ring. Fixing 1 makes that unit one. This proves uniqueness, including over rings with nilpotents.

The smooth connected image contains a one-dimensional torus, since the action kernel is finite. Conjugate this torus to the diagonal torus of PGL_2 . If the image were proper, its dimension would be at most two. Its Lie algebra then contains the diagonal line and at most one of the two distinct torus root lines. Choose the diagonal cocharacter with nonnegative weights on this Lie algebra. The general limit-subgroup lemma in Section 3 gives $P_H(\lambda) = H$ for this image H : it is a smooth connected closed subgroup with the same dimension. But $P_H(\lambda) = H \cap P_{\mathrm{PGL}_2}(\lambda)$, and the latter ambient subgroup is triangular. Thus H would be solvable. The finite action kernel has central k -points, since conjugation by a connected group on a finite reduced set is constant. Solvability of H would make $G(k)$ solvable as a central extension, and density of $G(k)$ in the smooth group would make G solvable. This contradicts semisimplicity. The image is consequently all of PGL_2 .

Let B_+ and B_- be the two Borels containing T . Their unipotent radicals are one-dimensional, since their images in the point stabilizers of PGL_2 have finite kernel. A smooth connected one-dimensional unipotent group over k is $\mathbf{G}_a$. The action of T on these radicals has nonzero characters α_+ and α_- . An element of the normalizer interchanging the Borels induces inversion on T , so $\alpha_- = -\alpha_+$.

The Lie algebra of the action kernel lies in both $\mathfrak{b}_+$ and $\mathfrak{b}_-$, whose moving lines have the opposite nonzero characters just found. Thus it has no vector in either unipotent root line. Its intersection with either unipotent radical is finite and torus-stable. A finite subgroup of $\mathbf{G}_a$ stable under all scalar multiplications is defined by $x^{p^a} = 0$ in characteristic p , or is trivial in characteristic zero: a

homogeneous Hopf ideal has this form by the binomial formula. Every nontrivial such subgroup has nonzero tangent space. Our intersections therefore are trivial, so each unipotent radical maps isomorphically to the corresponding unipotent subgroup of PGL_2 . If a point of $B_+ \cap B_-$ is written tu_+ , its image lies in the diagonal intersection of the two point stabilizers in PGL_2 . Hence the image of u_+ is trivial, and $u_+ = 1$. This works on all test algebras and proves $B_+ \cap B_- = T$. The action kernel is consequently contained in T . A normal multiplicative-type subgroup of a connected smooth group is central, by rigidity of its character sheaf. The converse inclusion follows because the image has trivial centre. $\square$

For a reductive group of semisimple rank one, quotienting by its radical gives this situation. Its centre is the kernel of the resulting action on the same projective line. The two Borel unipotent radicals are still additive lines with opposite characters.

Now let α be any nonzero weight of a maximal torus of an arbitrary reductive k -group. Define the codimension-one torus

$$T_\alpha = (\ker \alpha)_{\mathrm{red}}^0.$$

The centralizer $G_\alpha = C_G(T_\alpha)$ is reductive by the preceding lesson. Its weights are precisely the weights of G that are rational multiples of α , since these and only these vanish on T_α . Its maximal central torus has codimension one in T : the nonzero characters on T/T_α span a rank-one lattice. Thus its semisimple rank is one. Proposition 2.3 proves

$$\Phi \cap \mathbf{Q}\alpha = \{\alpha, -\alpha\}, \quad \dim \mathfrak{g}_\alpha = 1.$$

This proves both the reducedness along each root line and the one-dimensional root-space assertion without appealing to a general root-system classification.

3. Subgroups obtained by taking a limit

The definitions and full grading, smoothness, connectedness and open-cell proof for these subgroups are in [Tori, maximal tori and their conjugacy](#), Lemma 3.3, an earlier lesson. In this lesson Lemma 3.1 denotes that already proved general lemma; it applies to every smooth affine group of finite presentation, without reductivity or a root classification.

For SL_2 and $\lambda(t) = \mathrm{diag}(t, t^{-1})$, conjugation sends $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ to $\begin{pmatrix} a & t^2 b \\ t^{-2} c & d \end{pmatrix}$. Thus $P(\lambda)$ is upper triangular, $U(\lambda)$ is upper unipotent, and $L(\lambda)$ is diagonal. This calculation works in characteristic two because the group character t^2 remains nontrivial.

4. Root groups over the base

Let $T = D_S(M)$ be a split maximal torus. Locally the nonzero weights are a fixed finite set $\Phi \subset M$, and each $\mathfrak{g}_\alpha$ is a line bundle by Proposition 2.3 on geometric fibres. Define T_α by the torsion-free quotient of $M/\mathbf{Z}\alpha$. Then

$$G_\alpha = C_G(T_\alpha), \quad \mathrm{Lie}(G_\alpha) = \mathfrak{t} \oplus \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha}.$$

Choose a cocharacter λ of T with $n = \langle \alpha, \lambda \rangle > 0$, and set $U_\alpha = U_{G_\alpha}(\lambda)$. It is a smooth connected one-dimensional unipotent group, with Lie algebra $\mathfrak{g}_\alpha$.

For a line bundle L , write $\mathbf{V}(L) = \mathrm{Spec}_S \mathrm{Sym}(L^*)$ for its additive group.

Theorem 4.1. There is a unique T -equivariant homomorphism

$$\exp_\alpha : \mathbf{V}(\mathfrak{g}_\alpha) \longrightarrow G$$

whose differential is the inclusion of the root line. It is a closed immersion, with image U_α , and commutes with every base change. Moreover

$$U_{-\alpha} \times T \times U_\alpha \longrightarrow G_\alpha$$

is an open immersion.

Proof. On U_α , $\mathbf{G}_m$ acts through t^n on each geometric fibre. Its subgroup μ_n acts trivially: the fixed subgroup is smooth by exactness of multiplicative-type invariants, and equals U_α on every geometric fibre, so equals it over S .

Étale locally choose a section σ of U_α disjoint from its identity. The orbit map $t \mapsto t \cdot \sigma$ extends to $q : \mathbf{A}^1 \rightarrow U_\alpha$ with $q(0) = 1$. Since μ_n acts trivially on the target, the map factors through the invariant ring $R[x]^{\mu_n} = R[x^n]$ on an affine chart. Its factor $\bar{q} : \mathbf{A}^1 \rightarrow U_\alpha$ is an isomorphism on geometric fibres: there the original map is a nonzero scalar times t^n . The smooth fibrewise isomorphism criterion makes $\bar{q}$ an isomorphism over the base.

The quotient acting torus $\mathbf{G}_m/\mu_n$ gives ordinary scalar multiplication on the source. The transported group law is a polynomial $m(x, y)$ with

$$m(tx, ty) = tm(x, y), \quad m(x, 0) = x, \quad m(0, y) = y.$$

Homogeneity forces $m = ax + by$, and the identities force $a = b = 1$. Thus U_α is an additive line. Normalize the isomorphism by its differential to obtain $\exp_\alpha$. A scalar-equivariant polynomial map of an additive line is linear; if its differential is the identity it is the identity. This proves uniqueness and allows descent of the local construction.

An equivariant homomorphism with the required differential must factor through G_α , since T_α acts trivially on its source, and through $U_{G_\alpha}(\lambda)$, since its orbit limits to the identity. The same uniqueness therefore holds as a map into G . The open immersion follows from Lemma 3.1, with $L_{G_\alpha}(\lambda) = T$. $\square$

The notation $\exp_\alpha$ denotes this algebraic map, not the analytic exponential series. No division by factorials occurs. A root-group parameterization $x_\alpha : \mathbf{G}_a \simeq U_\alpha$ is exactly a trivializing section of the root line.

Example 4.2. In GL_n , conjugation by $\mathrm{diag}(t_i)$ sends E_{ij} to $(t_i/t_j)E_{ij}$. Thus

$$\alpha_{ij} = e_i - e_j, \quad \mathfrak{g}_{\alpha_{ij}} = \mathcal{O}_S E_{ij}, \quad x_{ij}(u) = I + uE_{ij}.$$

Example 4.3. For a line bundle L on S , consider $G = \mathrm{SL}(\mathcal{O}_S \oplus L)$. It has the split torus acting by (t, t^{-1}) on the two summands. Its two root lines are L^{-1} and L . If L is nontrivial, its root groups are $\mathbf{V}(L^{-1})$ and $\mathbf{V}(L)$ and need not be globally isomorphic to $\mathbf{G}_a$. Having a split maximal torus is thus weaker than having globally parameterized root groups. The distinction disappears over a field.

5. A projective quotient and its rank-one form

We use the general algebraic-space tool proved before this lesson in [Affine descent](#), [Zariski Main and recognition of spaces](#), Theorems E8.1 and E9.1. The proof there includes the finite flat affine quotient, the off-diagonal induction, separated locally quasi-finite scheme effectivity, and relative integral closure with its étale base change. It applies over every base and on every test scheme.

Quasi-finite normalization lemma. Let S be a scheme and let $f : X \rightarrow Y$ be a quasi-finite separated morphism of algebraic spaces over S . Let Y' be the relative integral closure of Y in X . Then the factorization $X \xrightarrow{f'} Y' \xrightarrow{\nu} Y$ has f' a quasi-compact open immersion and ν integral. In particular f is quasi-affine.

Proof. Theorem E9.1 of the preceding supporting lesson proves the full factorization, after Theorem E8.1 establishes recognition and Corollary E8.2 establishes representability on every scheme test. Its integral algebras are descended using the proved affine faithfully flat equalizer, and Corollary D4.3 supplies their étale base-change compatibility. Thus for every affine scheme $Z \rightarrow Y$, the pullback $Z \times_Y X$ is a quasi-compact open subscheme of the affine integral scheme $Z \times_Y Y'$. This is quasi-affineness. All these input proofs precede the present application in the programme. $\square$

The relative integral closure retains nilpotents; a reduced normalization is not substituted for it. The full argument is in the preceding supporting lesson, rather than being delegated to a source citation.

The following construction will also be used for flag varieties. It makes explicit why a fibrewise projective quotient is a scheme, even before we have classified the group over the base.

A flat-source fibre criterion. A finitely presented closed immersion $Y \hookrightarrow Z$ over S , with Y flat over S , is an isomorphism if it is so on every geometric fibre. Locally write its ideal as $I \subset C$. Flatness of C/I makes $I \otimes k(s) \rightarrow C \otimes k(s)$ injective. Fibre equality makes this module zero. Since I is finitely generated as an ideal, Nakayama at every prime of C gives $I = 0$. No flatness assumption on Z is needed.

Formal separation and normalizers

We first justify the use of identity neighbourhoods over bases with nilpotents. This argument concerns smooth connected subgroups, rather than only tori.

Formal separation lemma. Let R be Noetherian and let H/R be a smooth affine group of finite presentation with geometrically connected fibres. Write $C = \mathcal{O}(H)$ and let J be its augmentation ideal. Then

$$\bigcap_{n \geq 1} J^n = 0.$$

Proof. A smooth geometrically connected algebraic group over a field is geometrically integral: over an algebraic closure its regular irreducible components are disjoint and open, and connectedness leaves one component. For every prime $\mathfrak{p} \subset R$, the flat algebra $C/\mathfrak{p}C$ therefore embeds in its integral generic fibre over $\text{Frac}(R/\mathfrak{p})$ and is a domain.

Choose a finite prime filtration of the R -module R . Such a filtration exists for every finite module over a Noetherian ring: choose an element with prime annihilator, insert its cyclic submodule, and repeat in the quotient; the ascending chain condition terminates the process. Flatness of C turns this into a filtration of C with factors $C/\mathfrak{p}C$. If $j \in J$, then $1 - j$ has augmentation one in every such factor, so is nonzero there. It is a nonzerodivisor on every factor and consequently on C .

Put $I = \bigcap_n J^n$. The ring C is Noetherian. Artin–Rees applied to $I \subset C$ gives, for some c and every $n \geq c$,

$$I \cap J^n = J^{n-c}(I \cap J^c).$$

For completeness, this equality comes from the finite generation of the graded submodule $\bigoplus_n (I \cap J^n)t^n$ of the Rees ring $\bigoplus_n J^n t^n$: the latter is a finitely generated C -algebra, hence Noetherian; take c at least the degrees of homogeneous generators. Since $I \subset J^n$ for all n , taking $n = c + 1$ gives $I = JI$. On a finite set of generators of I , this equality has matrix form $v = Mv$ with entries of M

in J . The adjugate identity gives $\det(1 - M)I = 0$, where $\det(1 - M) = 1 - j$ for some $j \in J$. Its nonzerodivisor property proves $I = 0$. $\square$

Normalizer lemma. If H is a smooth closed subgroup with geometrically connected fibres of a smooth affine group K of finite presentation over any scheme S , then its scheme-theoretic normalizer is a closed subgroup of finite presentation in K . Its formation commutes with every base change.

Proof. First let $S = \operatorname{Spec} R$ be Noetherian. The identity neighbourhood H_n has coordinate algebra C/J^{n+1} , finite locally free over R . Indeed, smooth coordinates along the identity section identify the augmentation graded algebra with $\operatorname{Sym}(J/J^2)$; each of its finitely many graded pieces is finite locally free, and their extensions split locally. The same holds for K_n , and $H_n \subset K_n$ is a closed subscheme.

Write $D = \mathcal{O}(K)$. Restrict the conjugates of a finite set of generators of the ideal of H to H_n . Their vanishing gives finitely many coefficient equations in D , because $\mathcal{O}(H_n)$ is finite projective. Impose these equations for conjugation by g and by g^{-1} . They define a closed subgroup $N_n \subset K$, the stabilizer of H_n . Let $L_n \subset D$ be its ideal. The ideals L_n increase, so their union L equals some L_N .

It remains to check that these finite equations detect the whole subgroup on every test ring. Over the universal quotient D/L , both conjugation restrictions vanish on every identity neighbourhood of H . This quotient is Noetherian, and $H_{D/L}$ is still smooth with geometrically connected fibres. The formal separation lemma makes each restricted defining equation zero in $\mathcal{O}(H_{D/L})$. Thus both conjugations preserve $H_{D/L}$, and N_N normalizes H . Pulling back this actual equality proves sufficiency over every R -algebra, including non-Noetherian ones. Necessity follows because a group automorphism preserving H preserves its identity and all its identity neighbourhoods. Therefore $N_K(H) = N_N$ as functors, and the description proves arbitrary base change.

For general affine S , [Projective cohomology and smooth affine models](#), Proposition 3.2 and Theorem 4.5, supply smooth affine group and subgroup models with geometrically connected fibres over a finitely generated $\mathbf{Z}$ -algebra. Their proof descends the finite smoothness witnesses, proves constructibility of the number of geometric components, and removes the constructible bad locus at a finite coefficient stage. Apply that argument simultaneously to K, H , retaining the closed immersion, group laws and their finitely many identities. The Noetherian construction above then represents the normalizer on every algebra over the model. Its pullback represents the original normalizer on every S -scheme, including non-Noetherian and nonreduced tests. Uniqueness of the represented subgroup glues these constructions. $\square$

Cohomology of a projective completion

The preceding supporting lesson [Projective cohomology and smooth affine models](#), Theorem 2.3, proves the precise formal-functions statement used here. If A is Noetherian, X/A projective, F coherent and $I \subset A$, then

$$H^q(X, F)_I^\wedge \simeq \varprojlim_n H^q(X, F/I^n F).$$

Here is the mechanism of that written proof. Over the Rees algebra $B = \bigoplus I^n t^n$, the sheaf $\bigoplus I^n F$ is finite on the projective base change X_B ; projective Serre finiteness therefore makes $\bigoplus H^q(X, I^n F)t^n$ finite over B . If $M^q = H^q(X, F)$, $J_n^q = \operatorname{im}(H^q(X, I^n F) \rightarrow M^q)$ and K_n^q is its kernel, finite homogeneous generators give constants a_q, b_q with

$$I^n M^q \subset J_n^q \subset I^{n-a_q} M^q, \quad K_m^q \longrightarrow K_n^q = 0 \quad (m \geq n + b_q).$$

The sheaf exact sequence consequently gives

$$0 \longrightarrow M^q/J_n^q \longrightarrow H^q(X, F/I^n F) \longrightarrow K_n^{q+1} \longrightarrow 0.$$

Its compatible kernel coordinates vanish by the displayed transition bound; the remaining quotient filtration is cofinal with $I^n M^q$. This proves the inverse-limit comparison, without an unexplained exchange of cohomology and inverse limit.

S02 Lemmas 1.2–1.3 prove exact finite-module completion, Noetherianity and faithful flatness of local completion. Thus a finite module over a complete local Noetherian ring is complete. S02 Lemma 1.4 proves unique lifting of idempotents through nilpotent ideals, and Corollary 2.4 turns the compatible thickened idempotents into an actual section of a projective scheme. These are precisely the two completion operations in the quotient proof below.

Lemma 5.A (Grassmannians). The scheme $\mathrm{Gr}_d(V)$ of locally free rank- d quotients of a finite locally free V/S is smooth and projective, commutes with arbitrary base change, and its Plücker line bundle is $\det Q$, for its universal quotient Q .

Proof. Trivialize $V = \mathcal{O}^n$. On the open where columns I give a basis of a quotient, its unique normalized matrix has those columns equal to the identity and $d(n-d)$ free entries. A second basis J gives the transition $M_J = (M_I|_J)^{-1} M_I$. These open affine charts represent the quotient functor on all tests and glue. The determinant quotient $\bigwedge^d V \twoheadrightarrow \det Q$ defines the Plücker map to $\mathbf{P}(\bigwedge^d V)$, with the quotient convention. Over the target chart $p_I \neq 0$, the ratios of minors containing a single replacement of an identity column recover, with their determinant signs, every free matrix entry. The corresponding map of coordinate rings is surjective. These target charts cover the projective bundle, so the Plücker map is a closed immersion, and its $\mathcal{O}(1)$ pulls back to $\det Q$. The construction glues for general V and uses only invertible minors, hence works over nonreduced bases. The subbundle convention follows by dualizing. $\square$

Projective homogeneous quotients

Lemma 5.1. Let H be a smooth closed subgroup with geometrically connected fibres of a smooth affine S -group K with geometrically connected fibres. Suppose $N_K(H) = H$, and suppose every geometric quotient $K_{\bar{s}}/H_{\bar{s}}$ is proper. Then the fppf quotient $X = K/H$ is a smooth projective S -scheme, locally on S . Its universal conjugate subgroup $\mathcal{H} \subset K_X$ supplies a canonical relatively ample line bundle $\det(\mathrm{Lie} \mathcal{H})^*$.

Proof. Work first on an affine part $S = \mathrm{Spec} R$. Descend K, H and their group laws and closed immersion to a finite-type Noetherian model S_0 , retaining smoothness and geometrically connected fibres as in the normalizer lemma. The normalizer is now of finite presentation and commutes with base change. Its equality with H descends after a further finite stage: equality of their finitely generated defining ideals uses finitely many relation witnesses. We do not assume that projectivity of all geometric quotient fibres descends merely by choosing coefficients.

Construct the quotient $X_0 = K_0/H_0$ on this model. The free right H_0 -action defines a smooth equivalence relation. Corollary 8.3 of [Flat quotient bootstrap](#) applies to this free smooth action: its quotient is an algebraic space, $K_0 \rightarrow X_0$ is an H_0 -torsor, and its equality relation is $H_0 \times K_0 = K_0 \times_{X_0} K_0$ on every test scheme. Thus X_0 is smooth with geometrically connected fibres, and formation commutes with base change. For the next two paragraphs write K, H, X for these model objects.

Let K_n, H_n be the n th infinitesimal neighbourhoods of the identities. The normalizer lemma identifies $N_K(H)$ with the stabilizer of H_n in K_n for some n , as functors on all test schemes. Increase n to at least one. Since $N_K(H) = H$, this stabilizer is exactly H . The coordinate modules of K_n and H_n are finite locally free, with augmentation graded pieces $\mathrm{Sym}^i(\mathfrak{k}^*)$ and $\mathrm{Sym}^i(\mathfrak{h}^*)$ for $0 \leq i \leq n$.

Conjugation acts on the Grassmannian of quotients of the locally free coordinate module of K_n of rank equal to that of H_n . Its section corresponding to H_n has stabilizer exactly H . Thus the orbit map gives a finite-type monomorphism $X \rightarrow \mathrm{Gr}$. A finite-type monomorphism is quasi-finite and separated; the normalization lemma just proved from the earlier supporting lesson makes it

quasi-affine. In particular X is a scheme and is quasi-projective locally on the base.

We verify properness near the image of the original S in S_0 . At any point p of this image, the quotient fibre is proper: after extension to an algebraic closure containing a residue field of a point above p , this is the hypothesized proper geometric fibre, and properness descends under field extensions. Localize S_0 at p and pass faithfully flatly to its completion A . Place X as an open subscheme of a projective closure $\bar{X}$ over A . Its proper special fibre X_0 is open and closed in $\bar{X}_0$, and is therefore cut out by an idempotent. Idempotents lift uniquely across nilpotent ideals. The internal theorem on formal functions gives $\Gamma(\bar{X}, \mathcal{O}_{\bar{X}}) \simeq \varprojlim_n \Gamma(\bar{X}_n, \mathcal{O}_{\bar{X}_n})$, since A is complete and $\bar{X}$ is proper. It therefore turns this compatible formal idempotent into an actual idempotent. It defines an open and closed proper component $Z \subset \bar{X}$ with $Z_0 = X_0$. The closed subset $Z \setminus X$ has empty special fibre; properness makes it empty. Hence $Z \subset X$.

The image of $Z \rightarrow \operatorname{Spec} A$ is both open and closed: it is proper, and it is the restriction of the smooth map $X \rightarrow \operatorname{Spec} A$. It contains the closed point, so is all of the local base. In each geometric fibre, Z is a nonempty open and closed part of the connected fibre of X . It is the entire fibre. Thus $Z = X$, proving properness. This descends from the completion to $\mathcal{O}_{S_0, p}$. The monomorphism to the Grassmannian is now proper there, hence a closed immersion. The finite-presentation limit theorem for closed immersions, proved in [Limits of schemes and Noetherian approximation](#), Theorem 4.2, spreads this closed immersion to an open neighbourhood of p . The union of these neighbourhoods contains the image of the original base. Pulling back to S proves the required projectivity there; it has not been imposed on unrelated model fibres.

The Grassmannian map is now a proper monomorphism and hence a closed immersion. Its Plücker bundle pulls back to $\det \mathcal{O}_{\mathcal{H}_n}$. The augmentation filtration identifies this determinant with

$$\bigotimes_{i=0}^n \det(\operatorname{Sym}^i(\operatorname{Lie} \mathcal{H})^*) = (\det(\operatorname{Lie} \mathcal{H})^*)^c$$

for a positive integer c when $\dim H > 0$. Positivity follows already from the $i = 1$ term; the determinant exponent for Sym^i on a rank- d bundle is $\binom{i+d-1}{d}$. Thus the stated canonical bundle is relatively ample. The zero-dimensional connected smooth case is trivial. All the constructions descend to the original arbitrary base and commute with base change. $\square$

The quotient proof is [Flat quotient bootstrap](#), Theorem 8.2 and Corollary 8.3; the formal-functions proof is [Cohomology and models](#), Theorem 2.3. The algebraic-space normalization lemma has its full earlier proof in the descent and Zariski Main supporting lesson, Theorem E9.1. The group-specific argument is original exposition.

A universal finite cohomology complex

Lemma (finite projective replacement). Let A be Noetherian and let C be a complex of flat A -modules concentrated in degrees $[a, b]$, with finite cohomology modules. There is a complex K of finite projective modules in $[a, b]$ and a quasi-isomorphism $K \rightarrow C$ that remains a quasi-isomorphism after tensoring with any A -module.

Proof. Construct a bounded-above complex P of finite free modules and a map $f : P \rightarrow C$, descending from degree b . Once the terms above j are chosen, its cone has terms

$$D^i = C^i \oplus P^{i+1}, \quad d(c, p) = (d_C c + f(p), -d_P p).$$

Its cohomology is finite: the cone long exact sequence uses the finite cohomology of C and of the already chosen finite free portion of P . Choose finitely many cocycles (c_ℓ, p_ℓ) generating $H^j(D)$. Take P^j free on e_ℓ , and set $f(e_\ell) = c_\ell$, $d_P(e_\ell) = -p_\ell$. The cocycle equations give $d_P^2 = 0$ and

$d_C f = f d_P$. In the new cone, $d(0, e_\ell) = (c_\ell, p_\ell)$, so this kills H^j without changing higher cohomology. Continue to the left. Each degree stabilizes, giving a quasi-isomorphism $P \rightarrow C$.

Its cone is acyclic, bounded above and flat termwise. Such a complex stays acyclic after every tensor: start at its last degree and descend through

$$0 \longrightarrow Z^i \longrightarrow D^i \longrightarrow Z^{i+1} \longrightarrow 0.$$

The quotient is flat; hence the kernel is flat and the sequence stays exact under tensor. Thus $H^i(P \otimes_A M) = H^i(C \otimes_A M)$ for every M .

Put $Q = \text{coker}(P^{a-1} \rightarrow P^a)$. The left tail is a free resolution of Q , and

$$\text{Tor}_1^A(Q, M) = H^{a-1}(P \otimes_A M) = 0$$

for every M , since $C \otimes_A M$ starts in degree a . The module Q is finitely presented, so it is flat and finite projective. Take $K^a = Q$ and $K^i = P^i$ for $a < i \leq b$. The map to C factors through Q because $C^{a-1} = 0$. Truncation preserves cohomology, and the same acyclic flat-cone argument proves the assertion after every tensor. $\square$

Apply this to the finite section complex of a line bundle L_0 on a smooth projective scheme over a Noetherian ring A_0 . Choose a finite affine cover. Separatedness makes its intersections affine, and S01 Sections 4–5 prove that its ordered section complex computes quasi-coherent cohomology. On each intersection the line bundle is a projective module over its chart algebra; smoothness makes that algebra A_0 -flat, so every term is A_0 -flat. S01 Theorem 8.4 and its closed-immersion passage prove that all cohomology modules are finite. Under every algebra base change, affine module-sheaf correspondence and localization, proved in S01 Section 2, identify the new section complex with its tensor product over A_0 . The finite projective replacement just proved therefore computes the line bundle cohomology after every base change.

Sections in a vanishing family

Lemma (sections in a vanishing family). Let $f : X \rightarrow S$ be smooth and projective of finite presentation, and let L be an invertible sheaf on X . Suppose that $H^q(X_{\bar{s}}, L_{\bar{s}}) = 0$ for every geometric point $\bar{s}$ and every $q > 0$. Then $f_* L$ is finite locally free, all its positive higher direct images vanish, and these statements commute with every base change. In particular the canonical map $g^* f_* L \rightarrow f'_* L'$ is an isomorphism for any $g : S' \rightarrow S$. The base may be nonreduced.

Proof. Work with $S = \text{Spec } A$. Choose a projective embedding, a finite affine cover trivializing L , and finite affine covers of the pairwise and triple overlaps. Write A as the filtered union of its finitely generated $\mathbf{Z}$ -subalgebras. The embedding, chart opens and overlap maps descend at a finite stage by [finite-presentation descent](#), Theorems 4.1–4.2. Retain the closed immersion, so the model X_0 is projective.

Retain smoothness at a finite stage by S02 Lemma 3.1 and Proposition 3.2. In a finite polynomial presentation, Lemma 3.0 identifies formal smoothness with a left inverse of the conormal differential map. The left inverse uses finitely many coefficients; its equations on the finitely many relation generators have finite polynomial witnesses. Descend these coefficients and witnesses to one stage. The same left inverse then proves formal smoothness there, and the retained finite presentation proves smoothness. Apply this on a finite affine cover, retaining the cover and the projective closed immersion by the finite-presentation limit proof. The line-bundle transition units, their inverses and the finitely many overlap and cocycle witnesses descend at the same stage, as in S02 Proposition 3.2. Gluing these descended transition units defines the invertible sheaf L_0 on the smooth projective model X_0/A_0 .

The preceding finite-projective-replacement lemma gives a bounded finite projective complex K_0 such that

$$H^q(K_0 \otimes_{A_0} B) \simeq H^q(X_0 \times_{A_0} \operatorname{Spec} B, L_0 \otimes_{A_0} B)$$

for every A_0 -algebra B , compatibly with the canonical comparison maps. Its proof uses the finite affine-cover complex and projective Serre finiteness; ordinary tensor of K_0 computes its derived tensor. This finite argument supplies the universal comparison before pullback to the original base.

Put $K = K_0 \otimes_{A_0} A$. Near any $s \in \operatorname{Spec} A$, trivialize its terms. Whenever a differential has an entry invertible at s , shrink the neighbourhood and perform row and column operations to put that entry in an identity block. The equation $d^2 = 0$ makes the adjacent differentials zero on that block, so it splits off a contractible pair $A \xrightarrow{1} A$. After finitely many cancellations, all remaining differential entries vanish in $\kappa(s)$. The remaining complex therefore has

$$K^q \otimes_A \kappa(s) = H^q(X_s, L_s).$$

For $q > 0$ the right side is zero by the hypothesis and field extension; for $q < 0$ it is zero because L_s is a sheaf. Thus every remaining free term except degree zero has rank zero and disappears on a neighbourhood of s . The degree-zero term is finite free. Every cancelled identity pair stays contractible after any tensor. The universal comparison consequently proves the direct-image assertions and every base-change assertion, including nonflat changes involving nilpotents. The canonical comparisons glue. Notice that vanishing was imposed only on the original family, not on unrelated fibres of its Noetherian model. $\square$

Proposition 5.2. If G has trivial centre and geometric semisimple rank one, and has a split maximal torus T , then Zariski locally (G, T) is isomorphic to $(\operatorname{PGL}_2, D)$.

Proof. Its geometric fibres are PGL_2 by Proposition 2.3: the action on the projective line has trivial kernel. Choose a root α locally. It is an isomorphism $T \rightarrow \mathbf{G}_m$, as this is true on the character lattices of the geometric fibres. Put $B = P_G(\alpha^{-1})$. It is smooth and on each geometric fibre is an upper triangular Borel subgroup of PGL_2 .

Its scheme-theoretic normalizer equals B . The normalizer lemma makes it closed of finite presentation and proves that its formation commutes with arbitrary base change. On geometric fibres the point stabilizer in $\mathbf{P}^1$ is self-normalizing. Infinitesimally, the bracket of a diagonal matrix class with the lower off-diagonal class is a nonzero lower off-diagonal class in every characteristic, so the normalizer has exactly the tangent space of B . It is therefore smooth and equals B on those fibres. The flat-source criterion above, applied with source B , proves the equality over the base without assuming the normalizer flat.

Lemma 5.1 makes $f : X = G/B \rightarrow S$ a smooth projective curve whose geometric fibres are $\mathbf{P}^1$, and it has the section $e = 1B$. A section of a smooth relative curve is an effective Cartier divisor: étale coordinates at the section give one parameter cutting out its ideal, a nonzerodivisor because the coordinate algebra is flat over the polynomial chart. Put $L = \mathcal{O}_X(e)$; it has degree one on every geometric fibre. The sections-in-a-vanishing-family lemma applies to L , since $H^1(\mathbf{P}^1, \mathcal{O}(1)) = 0$, and gives $E = f_* L$ locally free of rank two, with arbitrary base change. The evaluation $f^* E \rightarrow L$ is surjective: its cokernel is finite locally, its geometric fibres vanish by generation of $\mathcal{O}(1)$, and local Nakayama kills that cokernel. With the quotient convention it defines $j : X \rightarrow \mathbf{P}(E)$, and $j_{\bar{s}}$ is the usual isomorphism given by the two sections of $\mathcal{O}(1)$. Both schemes are smooth over S , and the differential of j is an isomorphism on all geometric fibres, hence locally an isomorphism of their cotangent bundles. The étale differential criterion makes j étale. Its geometric fibres are isomorphisms, so it is universally injective and surjective; an étale universally injective morphism is an open immersion, and surjectivity makes it an isomorphism. Trivializing E Zariski locally now gives $G/B \simeq \mathbf{P}_S^1$.

The action now gives $G \rightarrow \mathrm{PGL}_2$, an isomorphism on every geometric fibre, hence an isomorphism over S . It carries B to the point stabilizer. Within that stabilizer the transporter taking its torus T to the diagonal torus is a $\mathbf{G}_m$ -torsor: the stabilizer of the diagonal torus inside the upper triangular group is the diagonal torus. Such a torsor is the frame bundle of a line bundle and is Zariski locally trivial. This finishes the pairwise assertion. $\square$

6. Splitting a central extension of the rank-one cover

We need an integral result, including characteristic two. The following splitting argument is due to Gabber; we give the argument for a separated commutative group scheme Z , which includes the multiplicative-type centres used here.

Theorem 6.1. Every fppf central extension

$$1 \longrightarrow Z \longrightarrow E \longrightarrow \mathrm{SL}_{2,S} \longrightarrow 1$$

has a unique homomorphic section.

Proof. In SL_2 use

$$x(u) = \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}, \quad y(v) = \begin{pmatrix} 1 & 0 \\ v & 1 \end{pmatrix}, \quad h(t) = \mathrm{diag}(t, t^{-1}).$$

Let D', U', V' be their inverse images in E . A central extension of a commutative group has a commutator pairing on its quotient: changing a lift by an element of the central kernel does not change the commutator, and the identities for commutators make this pairing additive in each variable.

For D' , the pairing $\mathbf{G}_m \times \mathbf{G}_m \rightarrow Z$ is trivial. Restrict the first argument to μ_n . The resulting pairing is killed by n in that argument, and the n th-power map is an fppf epimorphism in the second argument, so this restriction is trivial. The schemes μ_n are universally schematically dense in $\mathbf{G}_m$: distinct Laurent exponents remain distinct modulo sufficiently large n . Separatedness of Z makes the whole pairing trivial. Thus D' is commutative.

The conjugation action on E factors through SL_2 , since Z is central. The commutator pairing on U' is invariant under simultaneous multiplication of its arguments by t^2 . The square map is fppf surjective, so it is invariant under simultaneous multiplication by every unit. For given u, v , the morphism $a \mapsto c(au, av)$ from $\mathbf{A}^1$ to Z is constant on $\mathbf{G}_m$. Separatedness extends the equality to zero, where it is the identity. Hence $c(u, v) = 1$. Thus U' is commutative; the same argument proves it for V' .

We construct a unique D -equivariant section of $U' \rightarrow U$. Fppf locally choose a unit t with $t^2 - 1$ invertible and a lift $\tilde{h}$ of $h(t)$. Such choices exist: the open locus $t(t^2 - 1) \neq 0$ in an affine line is smooth surjective, and the extension is an fppf epimorphism. In the commutative group U' , the endomorphism

$$\varphi(a) = \tilde{h}a\tilde{h}^{-1}a^{-1}$$

kills Z and induces multiplication by $t^2 - 1$ on $U = \mathbf{G}_a$. It therefore factors through a homomorphism $U \rightarrow U'$, which after division by $t^2 - 1$ is the required section. It is D -equivariant because D' is commutative. Uniqueness holds since any difference is a D -invariant homomorphism $U \rightarrow Z$: scalar invariance and the specialization from nonzero scalars to zero force it to vanish. Uniqueness descends these local sections. Obtain $x' : \mathbf{G}_a \rightarrow E$ and similarly y' .

Set

$$w'(t) = y'(-t^{-1})x'(t)y'(-t^{-1}), \quad h'(t) = w'(t)w'(1)^{-1}.$$

Their images are $\begin{pmatrix} 0 & t \\ -t^{-1} & 0 \end{pmatrix}$ and $h(t)$. Thus $h'(t) \in D'$. Equivariance gives

$$h'(s)w'(t)h'(s)^{-1} = w'(s^2t).$$

Divide by the instance $t = 1$. Since D' is commutative, this gives $h'(t) = h'(s^2t)h'(s^2)^{-1}$. The square map being fppf surjective, h' is a homomorphism. Its definition then gives $h'(s)w'(t) = w'(st)$.

Substitute the expression for w' into this last equality and move the y' terms using $h'(s)y'(v)h'(s)^{-1} = y'(s^{-2}v)$. With $u = st$ and $v = (s-1)/(st)$, the result is

$$x'(u)y'(v) = y'\left(\frac{v}{1+uv}\right)h'(1+uv)x'\left(\frac{u}{1+uv}\right) \quad (1)$$

when u and $1+uv$ are units. The restriction that u be a unit can be removed. Fppf locally write $u = a + b$ with $a, b, 1+bv$ units; the open conditions on a choice of b omit only finitely many points of each geometric affine-line fibre. Apply (1) first to $x'(b)y'(v)$, then to $x'(a)y'(v/(1+bv))$. The second denominator is $(1+uv)/(1+bv)$. Moving the two resulting h' factors together leaves the y' parameter $v/(1+uv)$ and the x' parameter

$$\frac{a}{(1+bv)(1+uv)} + \frac{b}{1+bv} = \frac{u}{1+uv}.$$

This proves (1) in its full domain.

These identities, the additivity of x', y' , the multiplicativity of h' , and its two conjugation identities are exactly the multiplication rules on the open cell $\Omega = U_-DU_+$ of SL_2 . Define a section there by

$$y(v)h(t)x(u) \longmapsto y'(v)h'(t)x'(u).$$

It is multiplicative whenever both factors and their product are in Ω : reorder their factors with (1). Write $s_\Omega : \Omega \rightarrow E$ for this cell section. For any test scheme $R \rightarrow S$ and finite list $g_1, \dots, g_m \in \mathrm{SL}_2(R)$, the open

$$\Omega_R \cap \bigcap_i \Omega_R g_i^{-1}$$

is nonempty in every geometric fibre: a smooth geometrically connected group has integral geometric fibres, and finitely many dense opens have nonempty intersection. It is smooth and surjective over R ; affine refinements therefore supply an fppf local point c with $c, cg_i \in \Omega$.

In particular, for one g , use the nonempty open

$$\Omega_R \cap \Omega_R^{-1} \cap \Omega_R g^{-1}.$$

Then $g = c^{-1}(cg)$ is a product of two points of Ω . Assign to a word in cell points the corresponding product of their s_Ω -values. This is independent of the word. For two words with the same endpoint g , choose c so that c and its translates by every partial product of both words lie in Ω . Successive cell multiplicativity says that multiplying either lifted word on the left by $s_\Omega(c)$ gives $s_\Omega(cg)$. Cancellation proves equality. In particular a word representing the identity has lift one; concatenation therefore respects multiplication and inverses. These word values agree after further fppf pullbacks, so descend through the sheaf property of E . They define a homomorphism of represented fppf sheaves, hence a scheme homomorphism $\mathrm{SL}_2 \rightarrow E$. Its composite with $E \rightarrow \mathrm{SL}_2$ is the identity on generating words. It is the required homomorphic section on all test schemes.

Finally any homomorphism $\mathrm{SL}_2 \rightarrow Z$ kills $x(u)$, since

$$[h(t), x(u)] = x((t^2 - 1)u),$$

and one may choose $t^2 - 1$ invertible fppf locally. It kills $y(v)$ similarly, and then $h(t) = w(t)w(1)^{-1}$. It kills the open cell, hence the group. The difference between two central sections is such a homomorphism, proving uniqueness. $\square$

7. Coroots from multiplication

Theorem 7.1. For every root α there is a canonical coroot $\alpha^\vee : \mathbf{G}_m \rightarrow T$, with $\langle \alpha, \alpha^\vee \rangle = 2$. There is a canonical perfect pairing $\mathfrak{g}_\alpha \otimes \mathfrak{g}_{-\alpha} \rightarrow \mathcal{O}_S$ such that, writing its value as XY ,

$$\exp_\alpha(X) \exp_{-\alpha}(Y) = \exp_{-\alpha}\left(\frac{Y}{1+XY}\right) \alpha^\vee(1+XY) \exp_\alpha\left(\frac{X}{1+XY}\right) \quad (2)$$

whenever $1+XY$ is a unit. This unit condition is precisely membership in the open cell $U_{-\alpha}TU_\alpha$.

Proof. Work in G_α , whose scheme-theoretic centre is $Z = \ker \alpha \subset T$, by Section 1. Theorem 1.1 constructs $q : G_\alpha \rightarrow Q = G_\alpha/Z$ as an affine reductive quotient and an fppf Z -torsor, compatible with every base change. The torus quotient has character group $\mathbf{Z}\alpha \subset X^*(T)$, so $T/Z \simeq \mathbf{G}_m$ via the descended character α . It is a closed subgroup of Q : its pullback under q is $T \subset G_\alpha$, and closed immersions descend through the fppf torsor. On geometric fibres Proposition 2.3 identifies this quotient with PGL_2 , so T/Z is a maximal torus.

The open cell $U_{-\alpha}TU_\alpha$ is right Z -stable; centrality makes Z act only on its torus coordinate. Its quotient is the open subscheme

$$U_{-\alpha} \times (T/Z) \times U_\alpha \subset Q,$$

as follows by invariant-open descent and the product torsor. At the identity this gives the two nonzero torus weights $\alpha, -\alpha$, with their root lines, in $\mathrm{Lie} Q$. The centre formula of Section 1, applied to $Q, T/Z$, is consequently $Z(Q) = \ker(\alpha : T/Z \rightarrow \mathbf{G}_m) = 1$, as a scheme over the original base. Proposition 5.2 therefore identifies the pair $Q, T/Z$, Zariski locally, with PGL_2 and its diagonal torus. Choose the identification so $\alpha(\mathrm{diag}(t, 1)) = t$.

Pull back $G_\alpha \rightarrow \mathrm{PGL}_2$ by $\mathrm{SL}_2 \rightarrow \mathrm{PGL}_2$. Theorem 6.1 splits the resulting central extension by Z . The homomorphism $\mathrm{SL}_2 \rightarrow G_\alpha$ obtained from the splitting identifies its upper and lower unipotent groups with $U_\alpha, U_{-\alpha}$: the maps are equivariant, and their differentials on the root lines are isomorphisms, so Theorem 4.1 gives the assertion. The diagonal cocharacter therefore defines $\alpha^\vee$.

Direct multiplication gives

$$\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ y & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ y/(1+xy) & 1 \end{pmatrix} \begin{pmatrix} 1+xy & 0 \\ 0 & (1+xy)^{-1} \end{pmatrix} \begin{pmatrix} 1 & x/(1+xy) \\ 0 & 1 \end{pmatrix}.$$

The upper-left entry is a unit exactly when the product is in this open cell. It supplies (2), a perfect pairing of the two root lines, and $\alpha \circ \alpha^\vee(t) = t^2$.

The pairing and coroot are unique. In the adjoint quotient, comparing the upper-root parameter in (2) forces any alternative scalar in the pairing to be one; equality of open-cell coordinates then fixes the coroot on every unit $1+xy$, which ranges over all units by setting $y = 1$. An alternative coroot in T with the same adjoint image could differ by a cocharacter into Z , but (2) fixes its actual torus component and eliminates that difference. Hence the constructions on different local identifications agree and descend. The same characterization proves base-change compatibility. $\square$

For SL_2 , use $T \simeq \mathbf{G}_m$ via $\mathrm{diag}(t, t^{-1})$. Its root is $\alpha(t) = t^2$, and $\alpha^\vee(t) = \mathrm{diag}(t, t^{-1})$. For PGL_2 , parameterize its diagonal torus by the class of $\mathrm{diag}(t, 1)$. Its root is $t \mapsto t$, and its coroot is the class of $\mathrm{diag}(t^2, 1)$. In both cases the pairing is two, but the root and coroot occupy different sublattices. For GL_2 , the roots are $e_1 - e_2$ and its negative, and the positive coroot is $t \mapsto \mathrm{diag}(t, t^{-1})$. These explicit formulas remain valid over $\mathbb{Z}$.

8. The three rank-one groups

A lattice calculation with pairing two

Lemma 8.A. Let X be a finite free abelian group of rank $r+1$. Let $a \in X$ and $b \in X^* = \mathrm{Hom}(X, \mathbf{Z})$ be nonzero and satisfy $b(a) = 2$. Up to an isomorphism of X preserving this pair, exactly one of the following descriptions occurs:

1. $X = \mathbf{Z}e_0 \oplus \mathbf{Z}^r$, $a = 2e_0$, $b = e_0^*$.
2. $X = \mathbf{Z}e_0 \oplus \mathbf{Z}^r$, $a = e_0$, $b = 2e_0^*$.
3. $r \geq 1$, $X = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2 \oplus \mathbf{Z}^{r-1}$, $a = e_1 - e_2$, $b = e_1^* - e_2^*$.

In these formulas b is zero on the displayed extra summand.

Proof. Write $a = ma_0$ and $b = nb_0$, with a_0, b_0 primitive and $m, n > 0$. Then

$$mn b_0(a_0) = 2.$$

Consequently (m, n) is $(2, 1)$, $(1, 2)$, or $(1, 1)$.

In the first case $b(a_0) = 1$. Every $x \in X$ has the unique expression

$$x = b(x)a_0 + (x - b(x)a_0), \quad x - b(x)a_0 \in \ker b.$$

Thus $X = \mathbf{Z}a_0 \oplus \ker b$; choose a basis of the free kernel. This gives case 1. In the second case apply the same expression using $b_0(a) = 1$. It gives case 2.

Suppose now that both a and b are primitive. Extend a to a basis $a, c_1, \dots, c_r$. At least one $b(c_i)$ is odd: otherwise every value of b , including $b(a) = 2$, would be even, contradicting primitivity of b . Relabel so that $b(c_1) = 2q + 1$, and replace c_1 by $c_1 - qa$. This is a basis operation and makes its value under b equal to one. Call the new vector c . Replace each remaining c_i by $c_i - b(c_i)c$; their values become zero. Finally set

$$e_1 = c, \quad e_2 = c - a.$$

These replace the basis pair a, c by another unimodular basis pair. They satisfy $a = e_1 - e_2$, $b(e_1) = 1$ and $b(e_2) = -1$, giving case 3. If $r = 0$, both primitive elements would pair to ± 1 , so this case cannot occur.

The divisibilities (m, n) of a and b are invariant under lattice isomorphisms. They distinguish all three cases. $\square$

Extending a map from the rank-one cell

Lemma 8.B. Let K/S be a smooth group with geometrically connected fibres. Let $\Omega \subset K$ be a fibrewise dense open containing the identity. Let H/S be a group scheme. Suppose $f : \Omega \rightarrow H$ sends the identity to the identity and satisfies

$$f(xy) = f(x)f(y)$$

on the open where $x, y, xy \in \Omega$. Then f extends uniquely to a homomorphism $K \rightarrow H$.

Proof. A geometrically connected smooth algebraic group is geometrically irreducible: its regular irreducible components are disjoint and open, so connectedness leaves only one. For any test scheme $R \rightarrow S$ and any finite list $g_1, \dots, g_d \in K(R)$, the open

$$\Omega_R \cap \bigcap_{i=1}^d \Omega_R g_i^{-1}$$

has a nonempty open in every geometric fibre of K_R . Its map to R is smooth and surjective. It therefore supplies a point c after an fppf cover of R , with c and all cg_i in the cell.

In particular, for $g \in K(R)$, choose $c \in \Omega_R \cap \Omega_R^{-1} \cap \Omega_R g^{-1}$ locally. Then $g = c^{-1}(cg)$ is a product of two cell points. Assign to any word in cell points the corresponding product of f -values.

This assignment is independent of the word. For two words $x_1 \cdots x_d$ and $y_1 \cdots y_e$ with the same endpoint g , choose c so that it and every translated partial product of both words lie in the cell. Repeatedly apply the given identity. The first image word multiplied on the left by $f(c)$, and the second image word multiplied by the same element, both equal $f(cg)$. Cancellation proves equality. The assignment also respects inverses: for $x \in \Omega(R)$, choose $d, d^{-1}, dx^{-1} \in \Omega$. The identities $(dx^{-1})x = d$ and $d^{-1}d = 1$ then give $f(d^{-1})f(dx^{-1}) = f(x)^{-1}$. The word $d^{-1}(dx^{-1})$ represents x^{-1} , as required.

The assignments agree after fppf pullback, so the sheaf property of H descends them. Concatenation of words proves multiplicativity. Yoneda makes this map of represented fppf sheaves a scheme morphism. Every extension must have these values on generating words, proving uniqueness. $\square$

The three rank-one models

Theorem 8.1. A split reductive group of rank $r + 1$ and semisimple rank one over a field is isomorphic to exactly one of

$$\mathbf{G}_m^r \times \mathrm{SL}_2, \quad \mathbf{G}_m^r \times \mathrm{PGL}_2, \quad \mathbf{G}_m^{r-1} \times \mathrm{GL}_2 \quad (r \geq 1).$$

For a reductive group over a scheme with a split maximal torus and semisimple rank one, the same description holds Zariski locally on fixed-type loci. Trivial root lines with chosen frames give a global isomorphism to the corresponding base change from $\mathbf{Z}$.

Proof. The root and coroot construction in Theorem 7.1 gives two roots $\pm\alpha$, their additive root lines, the coroot $\eta = \alpha^\vee$, and $\langle \alpha, \eta \rangle = 2$. Apply Lemma 8.A to $X = X^*(T)$, $a = \alpha$, $b = \eta$. The three resulting pairs are precisely those computed for the three displayed matrix groups: in SL_2 the root is twice the primitive character and the coroot is primitive; in PGL_2 the root is primitive and the coroot twice primitive; in GL_2 they are respectively $e_1 - e_2$ and $e_1^* - e_2^*$.

Let G_0 be the indicated matrix model. The lattice isomorphism gives a torus isomorphism $f_T : T \rightarrow T_0$ carrying root characters and coroots to those of the model. Choose a nonzero positive-root frame $E \in \mathfrak{g}_\alpha$. The perfect pairing of Theorem 7.1 gives a unique negative-root frame F with pairing one. Define $x(u) = \exp_\alpha(uE)$ and $y(v) = \exp_{-\alpha}(vF)$, and match these to the upper and lower matrix parameters x_0, y_0 of G_0 .

The open cell theorem gives $\Omega = U_- T U_+$. On it define

$$f(y(v)tx(u)) = y_0(v)f_T(t)x_0(u).$$

This is an isomorphism of schemes from Ω to the model cell. The group law on the torus, the additivity of x, y , and the conjugation identities $tx(u)t^{-1} = x(\alpha(t)u)$, $ty(v)t^{-1} = y(\alpha(t)^{-1}v)$, agree under this map. The remaining multiplication rule is

$$x(u)y(v) = y\left(\frac{v}{1+uv}\right)\eta(1+uv)x\left(\frac{u}{1+uv}\right), \quad 1+uv \in \mathcal{O}^\times,$$

proved in Theorem 7.1 by the arbitrary-base central-extension argument and the explicit 2×2 matrix identity.

For two cell points $y(v)tx(u)$ and $y(v')t'x(u')$, their product is in the cell exactly when $x(u)y(v')$ is: left multiplication by U_-T and right multiplication by TU_+ preserve the cell, as do their inverses. The preceding rule and its unit criterion therefore give every multiplication of cell points whose product remains in the cell. It follows that f satisfies Lemma 8.B. The lemma extends it to a group homomorphism $G \rightarrow G_0$. The inverse cell map satisfies the same identities and extends in the other direction. Uniqueness in the lemma makes their compositions the identity. Thus $G \simeq G_0$.

Any group isomorphism carries a maximal torus to a maximal torus. Over an algebraic closure the torus conjugacy theorem then identifies its root/coroot pair with the pair for the fixed torus. It can interchange the signs of both elements, but does not change either divisibility. The three invariant pairs $(2, 1), (1, 2), (1, 1)$ distinguish the models in all characteristics, including characteristic two. This proves uniqueness without a forward reference to the derived group.

Over a scheme, restrict to an open where the root/coroot lattice pair is constant and the positive root line is trivial. Its paired negative line is then also trivial. The same torus map, cell formula and fppf word argument work over every test scheme, so give the asserted local isomorphism. When the lattice pair and frames are fixed globally, these constructions are global. All maps commute with base change because the displayed formulas and their character pairings do. $\square$

9. Exercises and solutions

Exercise 9.1 (first steps). Compute all roots and root spaces of GL_n with respect to its diagonal torus. Determine the zero-weight space and the positive coroot for $e_i - e_j$.

Solution. The diagonal matrix units have weight zero and form $\mathfrak{t}$. For $i \neq j$, E_{ij} has weight $e_i - e_j$, each once. The corresponding root group is $I + uE_{ij}$. Its coroot has diagonal entries t in position i , t^{-1} in position j , and 1 elsewhere. Evaluating $e_i - e_j$ gives t^2 , so the root–coroot pairing is two.

Exercise 9.2 (rank-one multiplication). Compute $U_{\pm\alpha}$ and $\alpha^\vee$ for SL_2 . Verify the factorization (2), and explain why the character differential in characteristic two does not change this factorization.

Solution. The two groups are the upper and lower unipotent matrices. Their parameters are x, y in the displayed matrix identity, and $\alpha^\vee(t) = \mathrm{diag}(t, t^{-1})$. Multiplication of the three factors gives $\begin{pmatrix} 1+xy & x \\ y & 1 \end{pmatrix}$, equal to $x_\alpha(x)x_{-\alpha}(y)$. The character on the upper group is t^2 , so its pairing with this coroot is two. Although its differential is zero in characteristic two, the Laurent character and all polynomial matrix identities remain valid; no division by two was used.

Exercise 9.3 (infinitesimal information). In characteristic different from two, show that $\mathfrak{sl}_2 \simeq \mathfrak{pgl}_2$ as Lie algebras, while the groups are not isomorphic. What happens to the differential of $\mathrm{SL}_2 \rightarrow \mathrm{PGL}_2$ in characteristic two?

Solution. The inclusion of trace-zero matrices followed by quotienting by scalar matrices has zero kernel when two is invertible: a trace-zero scalar is zero. Both Lie algebras have dimension three, so this bracket-preserving map is an isomorphism. The groups have different centres, μ_2 and 1. In characteristic two the scalar identity has trace zero and lies in the kernel of the differential, which therefore has one-dimensional kernel. It cannot be an isomorphism. The group map remains a

central fppf isogeny with kernel μ_2 .

Exercise 9.4 (classification). List the rank-two split reductive groups of semisimple rank one. Distinguish them by their centres and derived groups, including characteristic two.

Solution. Put $r = 1$ in Theorem 8.1. The list is $\mathbf{G}_m \times \mathrm{SL}_2$, $\mathbf{G}_m \times \mathrm{PGL}_2$, and GL_2 . Their centres are respectively $\mathbf{G}_m \times \mu_2$, $\mathbf{G}_m$, and $\mathbf{G}_m$. The first centre is not a torus: in characteristic two it is nonsmooth, and otherwise it has two geometric components. We compute the indicated derived groups directly. Fppf locally choose t with $t^2 - 1$ invertible. The identities $[h(t), x(u)] = x((t^2 - 1)u)$ and $[h(t), y(v)] = y((t^{-2} - 1)v)$ put both root groups of SL_2 in its commutator subgroup. The open cell word argument of Theorem 6.1 shows that these root groups generate SL_2 as an fppf sheaf, since $h(t) = w(t)w(1)^{-1}$. Their images generate PGL_2 , because $\mathrm{SL}_2 \rightarrow \mathrm{PGL}_2$ is fppf surjective; hence its derived group is itself. For GL_2 , determinant kills all commutators, and conjugation by $\mathrm{diag}(t, 1)$, with $t - 1$ invertible fppf locally, puts both determinant-one root groups in the commutator subgroup. The same generation argument gives its derived group SL_2 . The central $\mathbf{G}_m$ factors contribute no commutators. Thus the last two models have derived groups PGL_2 and SL_2 ; the first and third have the same derived group but different centres. These computations are scheme-theoretic and remain valid in characteristic two.

Exercise 9.5 (families). For $G = \mathrm{SL}(\mathcal{O} \oplus L)$, identify the two root groups and their pairing. Give a condition on L necessary for a pinning of this pair with its displayed torus.

Solution. An upper off-diagonal entry is a map $L \rightarrow \mathcal{O}$, so its line is L^{-1} ; a lower entry is a map $\mathcal{O} \rightarrow L$, so its line is L . Composition gives $L^{-1} \otimes L \rightarrow \mathcal{O}$, the perfect pairing in Theorem 7.1. A pinning chooses a nowhere-vanishing section in the simple root line. It therefore requires L^{-1} , equivalently L , to be trivial. This illustrates the global hypothesis absent from the field case.

The [course prerequisite guide](#) records the exact supporting statements, their full proof routes, and the hypotheses needed in their applications.

References and prerequisite proofs

- Brian Conrad, *Reductive group schemes*, §§2.3 and 4.1–4.3, for the dynamic approach and Gabber’s central-extension splitting argument. The proofs above are written out independently.
- Milne’s freely accessible *Algebraic Groups*, version 2.00 (2015). This exact free author edition provides comparison material; its citations do not replace the programme proofs.
- SGA 3, *Schémas en groupes*, Exposés XIX–XX, for roots and the relative rank-one theorem.
- The [official Stacks project](#) and AI Integrated Stacks Project, an edition with AI-proposed corrections and AI-written additions, not reviewed by the Stacks project’s maintainers. The normalization lemma and its whole proof in Section 5 are adapted from the Stacks project authors through this edition.

The following supporting statements have their exact proof routes recorded in the course prerequisite guide:

- [Algebra and sheaf cohomology before reductive groups](#), §§3–8: injectives, finite-cover comparisons, affine vanishing, projective-space twists, ample embeddings and projective Serre finiteness.
- [Projective cohomology and smooth affine models](#), §§1–6: exact and faithful local completion; projective formal functions and idempotents; smooth connected affine models; universal finite projective complexes and all-base-change direct images; regular curve immersions, curve duality, divisor degrees and the genus-zero calculation.
- [Completing a smooth affine curve and extending its group action](#), §§1–4: finite curve normalization, completion, the actual family action and one-dimensional unipotent identification

used through the preceding solvable-group proofs.

- [Affine descent, Zariski Main and recognition of spaces](#), Theorems E8.1–E9.1: recognition on every scheme test, representability and the full relative integral-closure factorization.
- [Flat quotient bootstrap](#), Theorem 8.2 and Corollary 8.3: the algebraic-space quotient and torsor relation for the free smooth subgroup action, over arbitrary bases.
- [Lemma 5.A of this lesson](#): quotient Grassmannian charts, arbitrary base change, and the closed Plücker immersion with determinant quotient bundle; the subbundle convention follows by dualizing.
- The current written group-scheme proofs: *Diagonalizable groups*, Theorems 3.4–3.5 and §4, for smooth fixed loci, invariant algebras and torus quotients; *Group schemes over a field*, Proposition 5.8, for the reduced schematic image and faithful-flat image factorization; *Lie algebras and smoothness*, Theorem 2.9 and Theorem 3.2, for the universal normalizer tangent calculation and geometric smoothness criterion; and *Quotients and torsors*, its affine descent proof and Theorem 8.1, for effective algebra descent and line-bundle frame torsors. These are the actual written proofs, including positive-characteristic and nonreduced scope, rather than source citations.

Theorem 1.1 proves the affine central quotient before the rank-one argument; the following lesson adds the consequences for root-group coordinates. The arbitrary-base derived group is constructed after the existence theorem in the following lesson.

History

This lesson’s new exposition, constructions, calculations, examples and solutions were written by GPT-6.1 Sol (OpenAI), Ultra setting, in October 2026, and are dedicated to the public domain under CC0.

The normalization lemma and its whole proof are adapted from the Stacks authors, *The Stacks Project, Morphisms of Algebraic Spaces*, Tag 0ABS, through the AI Integrated Stacks Project English source edition read on 1 October 2026. The Stacks authors retain copyright in that material. The source is the [versioned transparent source](#). AI Integrated Stacks Project contains AI-proposed corrections and AI-written additions and has not been reviewed by the Stacks project’s maintainers.

This collected lesson, *Roots and reductive groups of rank one* (2026), is published by Open Mathematics Courses. The Stacks authors are the authors of its imported proof; GPT-6.1 Sol (OpenAI) is responsible for the new contributions and adaptation. Permission is granted to copy, distribute and modify this collected lesson under the GNU Free Documentation License, Version 1.2, with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. An unaltered copy of the licence is supplied as [GNU Free Documentation License 1.2](#). This additional licence for the collected lesson does not withdraw the CC0 dedication of its original contributions.

GNU Free Documentation License, Version 1.2

GNU Free Documentation License
Version 1.2, November 2002

Copyright (C) 2000,2001,2002 Free Software Foundation, Inc.
51 Franklin St, Fifth Floor, Boston, MA 02110-1301 USA
Everyone is permitted to copy and distribute verbatim copies
of this license document, but changing it is not allowed.

0. PREAMBLE

The purpose of this License is to make a manual, textbook, or other functional and useful document "free" in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of "copyleft", which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation: a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference.

1. APPLICABILITY AND DEFINITIONS

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The "Document", below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as "you". You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A "Modified Version" of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A "Secondary Section" is a named appendix or a front-matter section of the Document that deals exclusively with the relationship of the publishers or authors of the Document to the Document's overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical

connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The "Invariant Sections" are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The "Cover Texts" are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A "Transparent" copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardly with generic text editors or (for images composed of pixels) generic paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not "Transparent" is called "Opaque".

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, LaTeX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available, and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The "Title Page" means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title page. For works in formats which do not have any title page as such, "Title Page" means the text near the most prominent appearance of the work's title, preceding the beginning of the body of the text.

A section "Entitled XYZ" means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as "Acknowledgements", "Dedications", "Endorsements", or "History".) To "Preserve the Title" of such a section when you modify the Document means that it remains a section "Entitled XYZ" according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties: any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

2. VERBATIM COPYING

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies.

3. COPYING IN QUANTITY

If you publish printed copies (or copies in media that commonly have printed covers) of the Document, numbering more than 100, and the Document's license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

It is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document.

4. MODIFICATIONS

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

- A. Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if there were any, be listed in the History section of the Document). You may use the same title as a previous version if the original publisher of that version gives permission.
- B. List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement.
- C. State on the Title page the name of the publisher of the Modified Version, as the publisher.
- D. Preserve all the copyright notices of the Document.
- E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices.
- F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.
- G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice.
- H. Include an unaltered copy of this License.
- I. Preserve the section Entitled "History", Preserve its Title, and add to it an item stating at least the title, year, new authors, and

publisher of the Modified Version as given on the Title Page. If there is no section Entitled "History" in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence.

- J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission.
- K. For any section Entitled "Acknowledgements" or "Dedications", Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein.
- L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles.
- M. Delete any section Entitled "Endorsements". Such a section may not be included in the Modified Version.
- N. Do not retitle any existing section to be Entitled "Endorsements" or to conflict in title with any Invariant Section.
- O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version's license notice. These titles must be distinct from any other section titles.

You may add a section Entitled "Endorsements", provided it contains nothing but endorsements of your Modified Version by various parties--for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity. If the Document already includes a cover text for the same cover, previously added by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version.

5. COMBINING DOCUMENTS

You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections Entitled "History" in the various original documents, forming one section Entitled "History"; likewise combine any sections Entitled "Acknowledgements", and any sections Entitled "Dedications". You must delete all sections

Entitled "Endorsements".

6. COLLECTIONS OF DOCUMENTS

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document.

7. AGGREGATION WITH INDEPENDENT WORKS

A compilation of the Document or its derivatives with other separate and independent documents or works, in or on a volume of a storage or distribution medium, is called an "aggregate" if the copyright resulting from the compilation is not used to limit the legal rights of the compilation's users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document's Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise they must appear on printed covers that bracket the whole aggregate.

8. TRANSLATION

Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is Entitled "Acknowledgements", "Dedications", or "History", the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title.

9. TERMINATION

You may not copy, modify, sublicense, or distribute the Document except as expressly provided for under this License. Any other attempt to copy, modify, sublicense or distribute the Document is void, and will automatically terminate your rights under this License. However, parties who have received copies, or rights, from you under this License will not have their licenses terminated so long as such parties remain in full compliance.

10. FUTURE REVISIONS OF THIS LICENSE

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See

<https://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License "or any later version" applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose any version ever published (not as a draft) by the Free Software Foundation.

ADDENDUM: How to use this License for your documents

To use this License in a document you have written, include a copy of the License in the document and put the following copyright and license notices just after the title page:

```
Copyright (c)  YEAR  YOUR NAME.
Permission is granted to copy, distribute and/or modify this document
under the terms of the GNU Free Documentation License, Version 1.2
or any later version published by the Free Software Foundation;
with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts.
A copy of the license is included in the section entitled "GNU
Free Documentation License".
```

If you have Invariant Sections, Front-Cover Texts and Back-Cover Texts, replace the "with...Texts." line with this:

```
with the Invariant Sections being LIST THEIR TITLES, with the
Front-Cover Texts being LIST, and with the Back-Cover Texts being LIST.
```

If you have Invariant Sections without Cover Texts, or some other combination of the three, merge those two alternatives to suit the situation.

If your document contains nontrivial examples of program code, we recommend releasing these examples in parallel under your choice of free software license, such as the GNU General Public License, to permit their use in free software.