The Claimed Complex Structure on S^6

What the manuscript and the proof reconstruction establish

A standalone mathematical orientation to the canonical workbench

Document map. This file is the equation-first orientation to the cumulative `workbench.tex`. Readers following one theorem lane can instead use the standalone drafts below; the cumulative workbench remains the source of record for their shared hypotheses and comparison maps.

`koide_arithmetic_preprint.tex`: exact Koide arithmetic.
`s6_spectral_degeneration_preprint.tex`: cusp spectral limits.
`s6_compact_link_gaussian_preprint.tex`: compact-link Gaussian model.
`s6_cone_sphere_routes_preprint.tex`: sphere-carrier topology.
`s6_erdos_straus_koide_bridge_preprint.tex`: Erdős–Straus–Koide incidence.
`s6_higher_rung_su3_gram_cone_preprint.tex`: rank-three Gram and toroidal-successor calculations.
`s6_worldline_presymplectic_reduction_preprint.tex`: exact massive, massless, and hard-threshold worldline reduction.

1. The candidate, its fillings, and the original audit

The construction starts from

$$B^\circ = \mathbb{P}^1 \setminus \{p_0, p_1, p_2\}, \qquad \Delta \simeq (\mathbb{Z}/3) * (\mathbb{Z}/4), \qquad g_0 = (g_1 g_2)^{-1},$$

and the marked two-tori

$$\Pi = [Z \mid I_2], \quad Z = \begin{pmatrix} 6\mu & \tau \\ \beta & \mu \end{pmatrix}, \quad D = \operatorname{Im}\beta - \frac{6(\operatorname{Im}\mu)^2}{\operatorname{Im}\tau} < 0, \quad \det_{\mathbb{R}} \Pi = \operatorname{Im}\tau\, D < 0.$$

The elliptic orientations are simultaneous. With s_j the Cayley coordinates and $S_1(w) = (w-1)/w$, $S(w) = -1/w$, $P(w) = w - 1$,

$$s_1 \circ g_1 = e^{-2\pi i/3} s_1, \qquad s_2 \circ g_2 = -i s_2, \qquad g_0 \text{ parabolic},$$
$$\tau \circ g = \varphi(g) \circ \tau, \qquad (\varphi(g_1), \varphi(g_2), \varphi(g_0)) = (S_1, S, P), \qquad S_1 S P = 1.$$

This normalization is forced at the cusp. On a g_0-stable component U_0, choose the j-cusp component K containing $\tau(U_0)$ and put

$$P(K) = K, \qquad \tau' = \gamma_0^{-1} \tau|_{U_0}, \qquad P' = \gamma_0^{-1} P \gamma_0, \qquad \tau' \circ g_0 = P' \circ \tau'.$$

If $P' = 1$, then $\exp(2\pi i \tilde{\tau}')$ has a single-valued logarithm and winding zero, whereas the pole of $j(\tau)$ forces a zero and positive winding. Hence P' is nontrivial parabolic. Returning to $\tau(z_1) = \rho$, $\varphi(g_1) = S_1$, the order-two condition gives

$$0 = \operatorname{tr}(\widetilde{S}_1^{-1} \widetilde{P}^{-1}) = 1 + k(p^2 - pr + r^2), \qquad k = -1, \qquad p^2 - pr + r^2 = 1,$$

so conjugation by a power of S_1 gives the displayed triple. Only then is $q := e^{2\pi i \tau} = t_c u_1(t_c)$, $u_1(0) \neq 0$, redefined. In the resulting cusp coordinate $e^{2\pi i s} = t_c$.

For $M_g = \rho_V(g)^{\mathsf{t}}$ and $A_g = M_g^{-1}$, the exact deck identity and determinant signs are

$$\Pi(gz) = R_g(z) \Pi(z) M_g, \quad R_{g_1} = \begin{pmatrix} -1/\tau & 0 \\ (1-\mu)/\tau & 1 \end{pmatrix}, \quad R_{g_2} = \begin{pmatrix} 1/\tau & 0 \\ -\mu/\tau & 1 \end{pmatrix},$$

$$\det R_{g_1} = -\tau^{-1}, \qquad \det R_{g_2} = \tau^{-1}, \qquad \det R_{g_0} = 1.$$

At the cusp,

$$Z = sB_0 + C(t_c), \quad B_0 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \Psi_{\bar{\lambda}}(x, t_c) = \left(e^{2\pi i C(t_c)\bar{\lambda}} t_c^{B_0 \bar{\lambda}} x, t_c\right), \quad N_0 = Y_\epsilon / \bar{\Lambda}.$$

The reduced irreducible fibre W has local equations z_0, $z_0 z_1$, $z_0 z_1 z_2$ and normalization dP_6 with opposite hexagon sides identified. The invariant toric form gives

$$\Omega = \frac{dx_1 \wedge dx_2 \wedge dt_c}{x_1 x_2} = \pm dz_0 \wedge dz_1 \wedge dz_2, \qquad \Psi_{\bar{\lambda}}^* \Omega = \Omega, \qquad \omega_X|_{N_0} \simeq \omega_{N_0} \simeq \mathcal{O}_{N_0}.$$

For $(m_1, m_2) = (3, 4)$ and $(v_1, v_2) = (\varepsilon, -\varepsilon')$, the finite lifts are

$$\widetilde{g}_j^b(z, \zeta) = (g_j z, R_{g_j}(z)\zeta + \Pi(g_j z) b), \qquad Q_{j,\log} = \widetilde{g}_j^{v_j/m_j}, \qquad Q_{j,\mathrm{lin}} = \widetilde{g}_j^0,$$

$$Q_{j,\log} \tau_\lambda Q_{j,\log}^{-1} = \tau_{A_j \lambda}, \qquad (Q_{j,\log})^{m_j} = \tau_{v_j}, \qquad (Q_{j,\mathrm{lin}})^{m_j} = 1.$$

The invariant covector satisfies $\gamma A_j = \gamma$, $\gamma(kv_1/3) = k/3$ and $\gamma(kv_2/4) = -k/4$ for $0 < k < m_j$; hence every nonidentity power of the descended $q_{j,\log}$ is fixed-point free. Thus

$$N_j = J_j'/\langle q_{j,\log}\rangle, \qquad f_j^*(p_j) = m_j S_j,$$

and the punctured conjugacy is $h_j q_{j,\log} h_j^{-1} = q_{j,\lin}$ on $J_j'^*$. These maps glue a compact smooth threefold $f : X = N_0 \cup J \cup N_1 \cup N_2 \to \mathbb{P}^1$. The integer

$$p = 12\ell_0 - 4\ell_1 - 3\ell_2, \qquad (\ell_0, \ell_1, \ell_2) = (0, 1, -1) \Longrightarrow p = -1$$

controls the topology. The manuscript claimed $X \cong_{\mathrm{diff}} S^6$, $a(X) = 1$, and $\langle c_3(T_X), [X]\rangle = 2$; the first audits verified the integer algebra but had not yet identified its collapse and matrices with the actual specialization and oriented collar maps.

2. Two integral topology certificates and recognition of S^6

Let F be a smooth fibre, $V = H^1(F;\mathbb{Z}) = \mathbb{Z}\langle\gamma, u, w, \delta\rangle$, and $q = uw + 6\gamma\delta$. Integral nearby cycles prove

$$\mathrm{sp}^r : H^r(W;\mathbb{Z}) \longrightarrow H^r(F;\mathbb{Z}), \qquad \operatorname{im}\mathrm{sp}^r = H^r(F;\mathbb{Z})^{T_0} \quad (0 \le r \le 4),$$

with saturated image. In $0 \to R^r f_*\mathbb{Z} \to j_*L^r \to Q^r \to 0$,

$$Q^1 = (\mathbb{Z}/3)_{p_1} \oplus (\mathbb{Z}/4)_{p_2}, \quad Q^2 = Q^3 = (\mathbb{Z}/2)_{p_2}, \quad Q^4 = (\mathbb{Z}/3)_{p_1} \oplus (\mathbb{Z}/4)_{p_2},$$

and therefore

$$H^0(R^1 f_*\mathbb{Z}) = \mathbb{Z}(12\gamma), \quad H^0(R^2 f_*\mathbb{Z}) = \mathbb{Z}(2q), \quad H^0(R^3 f_*\mathbb{Z}) = \mathbb{Z}(2\gamma uw).$$

Canonical truncation identifies the disc–pair-of-pants discrepancy with Leray d_2:

$$d_2(12\gamma) = -p\omega, \qquad d_2(2q) = -p\omega[\delta], \qquad d_2(2\gamma uw) = -p\omega[\gamma\delta].$$

For $p = -1$ these maps are primitive, so $H^1(X;\mathbb{Z}) = H^2(X;\mathbb{Z}) = H^3(X;\mathbb{Z}) = 0$ and $H_2(X) = 0$. The actual based overlap maps give

$$\pi_1(X) = \langle c, x, y \mid c \text{ central},\ xy = 1,\ x^3 = c,\ y^4 = c^{-1}\rangle = 1.$$

Consequently

$$H_k(X;\mathbb{Z}) = \begin{cases} \mathbb{Z}, & k = 0, 6, \\ 0, & 1 \le k \le 5, \end{cases} \qquad X \simeq S^6, \qquad X \cong_{\mathrm{diff}} S^6, \qquad \langle c_3(T_X), [X]\rangle = \chi(X) = 2,$$

where the last diffeomorphism uses $\Theta_6 = 0$.

The independent cellular certificate starts from the equivariant honeycomb collapse:

$$0 \longrightarrow \operatorname{im}(\textstyle\bigwedge^r M_0 - I) \longrightarrow \bigwedge^r \Lambda \xrightarrow{c_*} H_r(W;\mathbb{Z}) \longrightarrow 0, \qquad \mathrm{sp}^r = i_F^* r^* = c^*.$$

The finite-band attachment has the oriented map

$$\alpha_r = (\pi_{1*}, -\pi_{2*}) : H_r(F) \longrightarrow H_r(S_1) \oplus H_r(S_2),$$

and, after splitting primitive cusp kernels, the only global blocks are

$$D_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -12 & 0 & -p \end{pmatrix}, \qquad D_2^{\mathrm{red}} = \begin{pmatrix} -p & 0 \\ 0 & 1 \end{pmatrix}, \qquad \det D_1 = \det D_2^{\mathrm{red}} = -p.$$

They are unimodular for $p = -1$ and reproduce the same integral homology. Dualizing these oriented collar maps gives the three displayed transgressions, so the Leray and cellular proofs agree map-by-map without depending on one another.

The explicit candidate and terminal algebra remain attributed to the Claude/Fabel source. The audits isolated the missing geometric identifications, and the present AI-produced verification aid reconstructs the gluing, both topology certificates, meromorphic descent, Picard theory, conductor mechanism, Hodge and Bott–Chern calculations, and the CDP repair as checkable proofs. Its purpose is to reduce the reconstruction burden for human experts, not to claim priority for the candidate or to substitute for independent specialist adjudication. Root `main.tex` is the annotated edition and `workbench.tex` contains the complete proofs and sources.

3. Algebraic reduction, Picard, Hodge, and Bott–Chern invariants

For a very general fibre,

$$\mathrm{NS}(F_b) = \mathbb{Z}\eta, \qquad \eta = u \wedge w + 6\gamma \wedge \delta, \qquad \eta^2 = 12\gamma \wedge u \wedge w \wedge \delta.$$

Because $\det_{\mathbb{R}} \Pi < 0$, complex and marked orientations are opposite:

$$\int_{(F_b, \mathrm{or}_{\mathrm{mark}})} \eta^2 = 12, \qquad \int_{(F_b, \mathrm{or}_{\mathbb{C}})} \eta^2 = -12.$$

The Appell–Humbert form has signature $(1,1)$, hence $a(F_b) = 0$; the meromorphic descent theorem gives

$$\mathcal{M}(X) = f^*\mathbb{C}(t), \qquad a(X) = 1.$$

Writing $\mathfrak{K} = f_*\omega_{X/B}$, one has

$$\mathfrak{K} \simeq \mathcal{O}_B(1), \quad f_*\mathcal{O}_X = \mathcal{O}, \quad R^1 f_*\mathcal{O}_X = \mathcal{O} \oplus \mathcal{O}(-1), \quad R^2 f_*\mathcal{O}_X = \mathfrak{K}^\vee = \mathcal{O}(-1),$$

and $\bigwedge^2 R^1 f_*\mathcal{O}_X \simeq R^2 f_*\mathcal{O}_X$. The full canonical-bundle chain is

$$L := K_X \otimes \mathcal{O}_X(-2S_2), \qquad M := f_*L, \qquad f^*M \simeq L, \qquad \mathfrak{K} = M(2), \qquad M \simeq \mathcal{O}_B(-1),$$
$$K_X \simeq f^*\mathcal{O}_B(-1) \otimes \mathcal{O}_X(2S_2), \qquad K_X^{\otimes 4} \simeq f^*\mathcal{O}_B(-2).$$

Leray and the exponential sequence give

$$H^1(X, \mathcal{O}_X) = \mathbb{C}, \quad H^2(X, \mathcal{O}_X) = 0, \quad \mathrm{Pic}(X) \simeq H^1(X, \mathcal{O}_X) \simeq \mathbb{C}.$$

For $D = W + S_1 + S_2$, $\Sigma = \{p_0, p_1, p_2\}$ and $\mathcal{E} = \Omega^1_X(\log D)/f^*\Omega^1_B(\log \Sigma)$, the cyclic invariant lattices give

$$f_*\mathcal{E} = \mathcal{O} \oplus \mathcal{O}(-1), \quad R^1 f_*\mathcal{E} = \mathcal{O}^{\oplus 2} \oplus \mathcal{O}(-1)^{\oplus 2}, \quad R^2 f_*\mathcal{E} = \mathcal{O} \oplus \mathcal{O}(-1).$$

The typed Kodaira–Spencer maps

$$\kappa_0 : f_*\mathcal{E} \to \Omega^1_B(\log \Sigma) \otimes R^1 f_*\mathcal{O}_X, \qquad \kappa_1 : R^1 f_*\mathcal{E} \to \mathcal{O}_B$$

are certified by the load-bearing local equations

$$\mathrm{div}(\det \kappa_0) = p_1 + p_2, \qquad \mathrm{coker}\, \kappa_0 = \mathbb{C}_{p_1} \oplus \mathbb{C}_{p_2}, \qquad \phi \longmapsto (\phi(p_0), 3\phi(p_1), 4\phi(p_2)),$$
$$\rho_* = (0, 1, -1, 0) : \mathbb{C}^4 \to \mathbb{C}\theta, \qquad \kappa_1(h \otimes \gamma)(p_2) = -\tfrac{1}{4}\dot{\mu}(z_2) \neq 0.$$

The first row is the normalized residue cancellation, the second the cusp Čech sign and invariant order-four scalar. Thus

$$f_*\Omega^1_X = \mathcal{O}(-2), \qquad R^1 f_*\Omega^1_X = \mathcal{O} \oplus \mathcal{O}(-1)^{\oplus 2}, \qquad R^2 f_*\Omega^1_X = \mathcal{O} \oplus \mathcal{O}(-1),$$

and coherent Leray yields

$q\backslash p$	0	1	2	3
3	0	0	0	1
2	0	1	2	1
1	1	2	1	0
0	1	0	0	0.

The two nontrivial Frölicher rows are $0 \to \mathbb{C} \to \mathbb{C}^2 \to \mathbb{C} \to 0$; $\dim E_1 = 10$, $\dim E_2 = 2$, and $E_2 = E_\infty$. McHugh's formulas give

$$(h^k_{BC}) = (1, 0, 2, 2, 2, 2, 1), \quad (h^k_A) = (1, 2, 2, 2, 2, 0, 1), \quad (\Delta^k) = (0, 2, 4, 4, 4, 2, 0).$$

Let $\mathfrak{g} = [G]_{\bar{\partial}}$ be induced by γ, and choose a coherent-Leray lift $\mathfrak{q} = [Q]_{\bar{\partial}}$ of η. In complex orientation the decisive fibre pairings are

$$\int_{F_b} Q_F^2 = -12, \qquad \int_{F_b} Q_F \wedge \Gamma \wedge d\zeta_2 = -1.$$

For $\exp : H^1(X, \mathcal{O}_X) \to \mathrm{Pic}(X)$ induced by $z \mapsto e^{2\pi i z}$,

$$H^0(B, R^1 f_*\mathbb{Z}) = \mathbb{Z}(12\gamma), \qquad \exp(12\mathfrak{g}) = f^*\mathcal{O}_B(1).$$

Put $\mathsf{A}=[\partial G]_{BC}$, $\mathsf{B}=[\bar\partial\bar G]_{BC}$. In a unit frame,

$$\bar\partial_z s=-2\pi i z G\otimes s,\qquad \nabla_z=d-2\pi i(zG+\bar z\,\bar G),\qquad c_{1,BC}(\exp(z\mathfrak{g}))=z\mathsf{A}+\bar z\,\mathsf{B},$$

so $c_{1,BC}(f^*\mathcal{O}_B(1))=12(\mathsf{A}+\mathsf{B})$. If N is spanned by

$$\mathsf{A},\mathsf{B},\ [\partial Q]_{BC},[\bar\partial\bar Q]_{BC},\ [\partial(G\wedge Q)]_{BC},[\bar\partial(\bar G\wedge\bar Q)]_{BC},\ [\partial(Q^2)]_{BC},[\bar\partial(\bar Q^2)]_{BC},\ [\nu_{\mathrm{top}}]_{BC},$$

with $\int_X\nu_{\mathrm{top}}=1$, then

$$\mathbf{1}=[1]_{BC},\qquad H^{*,*}_{BC}(X)=\mathbb{C}\mathbf{1}\oplus N,\qquad N^2=0.$$

After fixing $\mathfrak{g}$ and one coherent-Leray lift $\mathfrak{q}$, smooth gauge changes add Dolbeault coboundaries and preserve all displayed classes, pairings, nonvanishing arguments and products. The invariant-basis, residue, Čech and scalar computations are in `supporting_materials/workbench/research/hodge_frolicher.tex`; the product proofs are in `supporting_materials/workbench/research/bott_chern_algebra.tex`.

4. CDP20, the analytic Picard certificate, and the corrected theorem

For a reduced nonnormal fibre S, put $\mathcal{K}_S=\ker(\Omega^1_X|_S\to\widetilde{\Omega}^1_S)$. The exact sequence and the group surviving the two vanishings used in CDP20 are

$$0\to N^*_{S/X}\to\mathcal{K}_S\to\operatorname{tors}\Omega^1_S\to 0,$$

$$H^0(S,\Omega^1_X|_S\otimes A)=\ker\Big\{H^0(S,\operatorname{tors}\Omega^1_S\otimes A)\xrightarrow{\delta_A}H^1(S,N^*_{S/X}\otimes A)\Big\}.$$

CDP20 does not prove δ_A injective. Its second imported step must also use coinvariants:

$$(K\rtimes Q)^{\mathrm{ab}}\simeq Q^{\mathrm{ab}}\oplus(K^{\mathrm{ab}})_Q.$$

Thus the completed candidate, with $a(X)=1$ and $c_3(X)=2$, contradicts Theorem 2.2 and its S^6 corollary as stated.

It remains to prove the Picard restriction needed by the cusp conductor without recycling global topology. For

$$E=c_1(L|_F)=kq\in\mathrm{Alt}^2(\Lambda,\mathbb{Z}),\qquad V_{\mathbb{R}}=\mathrm{Hom}(\Lambda,\mathbb{R}),$$

the global factors give

$$d:\pi_1(P)\to V_{\mathbb{R}},\qquad d(gh)=d(g)+T_g d(h),\qquad \bar d_1+\bar d_2+\bar d_0=0\quad\text{in } V_G\otimes\mathbb{R}.$$

Distinguish every finite covering level:

$$\widetilde{J}_j=\Delta_j\times\mathbb{C}^2\xrightarrow{r_j}J'_j,\qquad q_{j,\bullet}\circ r_j=r_j\circ Q_{j,\bullet},\quad \bullet\in\{\log,\mathrm{lin}\},$$

$$Q_{j,\log}\tau_\lambda Q_{j,\log}^{-1}=\tau_{A_j\lambda},\quad (Q_{j,\log})^{m_j}=\tau_{v_j},$$
$$Q_{j,\mathrm{lin}}\tau_\lambda Q_{j,\mathrm{lin}}^{-1}=\tau_{A_j\lambda},\quad (Q_{j,\mathrm{lin}})^{m_j}=1.$$

On $J'^*_j=J'_j|_{\Delta^*_j}$,

$$h_j q_{j,\log}h_j^{-1}=q_{j,\mathrm{lin}},$$
$$p_{j,\log}:J'^*_j\to N^*_j:=J'^*_j/\langle q_{j,\log}\rangle,\qquad p_{j,\mathrm{lin}}:J'^*_j\to J^*_j:=J'^*_j/\langle q_{j,\mathrm{lin}}\rangle,$$
$$\bar h_j:N^*_j\xrightarrow{\sim}J^*_j,\qquad p_{j,\mathrm{lin}}h_j=\bar h_j p_{j,\log},\qquad p^*_{j,\log}(L|_{N^*_j})\simeq h_j^*p^*_{j,\mathrm{lin}}(L|_{J^*_j}).$$

Put

$$\widetilde{J}^*_j=\Delta^*_j\times\mathbb{C}^2,\quad \Phi_{j,\log}=p_{j,\log}r_j,\quad \Phi_{j,\mathrm{glob}}=p_{j,\mathrm{lin}}h_jr_j=\bar h_j\Phi_{j,\log}.$$

The pulled gluing isomorphism

$$\Theta_j:\Phi^*_{j,\log}(L|_{N^*_j})\xrightarrow{\sim}\Phi^*_{j,\mathrm{glob}}(L|_{J^*_j})$$

intertwines descent and induces a centre-preserving isomorphism of the two $\Gamma^{\log}_j$-factor extensions, where

$$\Gamma^{\log}_j=\langle\tau_\lambda,Q_{j,\log}:\lambda\in\Lambda\rangle.$$

For normalized vertical lifts

$$U^{(j)}_\lambda U^{(j)}_\mu=\mathbf{c}\big(\tfrac12E(\lambda,\mu)\big)\,U^{(j)}_{\lambda+\mu},$$

define

$$\widetilde{G}_\alpha U^{(j)}_{A_\alpha^{-1}\lambda}\widetilde{G}_\alpha^{-1}=\mathbf{c}(\langle\widetilde{d}_j(\alpha),\lambda\rangle)U^{(j)}_\lambda.\tag{5.17-A14c}$$

Then

$$\widetilde{d}_j(\alpha\beta) = \widetilde{d}_j(\alpha) + T_\alpha \widetilde{d}_j(\beta).$$

Compatible basepoints and $\ell'(w) = \ell(g_j^{-1}w) - 2\pi i/m_j$ identify the clockwise zero-section endpoint with exactly $Q_{j,\log}$, not an inverse or lattice translate. Thus, for a comparison character a_j,

$$\widetilde{d}_j(Q_{j,\log}) = d_j + (T_j - I)a_j, \qquad \widetilde{d}_j(\tau_v) = \iota_v E.$$

Since $N_{T_j}(T_j - I) = 0$,

$$N_{T_j} d_j = N_{T_j}\widetilde{d}_j(Q_{j,\log}) = \widetilde{d}_j((Q_{j,\log})^{m_j}) = \widetilde{d}_j(\tau_{v_j}) = \iota_{v_j} E. \tag{5.17-A15}$$

The comparison is independent of frames, lifts, normalized vertical systems, and logarithm branch; branch change $H_j \mapsto \tau_{nv_j} H_j$ changes a_j by $n\iota_{v_j}E$, which is T_j-invariant.

At the cusp the argument principle gives $\bar{d}_0 = 0$. With

$$\eta_1 = 2u + w + 3\delta, \qquad \eta_2 = u + w + 2\delta,$$

$$\iota_{v_1} q = 2\eta_1, \quad \iota_{v_2} q = -3\eta_2, \quad [\eta_1] = 3[\delta], \quad [\eta_2] = 2[\delta],$$

equation (5.17-A15) yields

$$\bar{d}_1 = 2k[\delta], \qquad \bar{d}_2 = -\tfrac{3}{2}k[\delta], \qquad 0 = \bar{d}_1 + \bar{d}_2 + \bar{d}_0 = \tfrac{k}{2}[\delta].$$

Hence $k = 0$ and $E = 0$.

For central triviality, put

$$M = L|_{N_0}, \qquad \mathcal{M} = \pi^* M, \qquad \mathcal{R} = \mathcal{M} \otimes \mathcal{O}(-D_a).$$

The negative-divisor linearization is the inverse-dual

$$\mu_\lambda^- = ((\mu_\lambda^+)^{-1})^\vee, \qquad \theta_\lambda^{\mathcal{R}} = \theta_\lambda^{\mathcal{M}} \otimes \mu_\lambda^-.$$

In the stated lattice bases,

$$B_0 : \bar{\Lambda} \xrightarrow{\sim} \Lambda_{\mathrm{tor}}, \qquad [B_0] = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

If $A = \nu^*(M|_W)$ has degrees $(d_1, d_2, d_3, d_1, d_2, d_3)$ and support matrix

$$[B] = \begin{pmatrix} d_2 + d_3 & d_3 \\ d_3 & d_1 + d_3 \end{pmatrix},$$

then

$$E_M((\lambda, 0), (0, m)) = -B(B_0\lambda, m), \qquad E_M = c_1(M|_{F_t}) = c_1(L|_F) = E = 0.$$

Unimodularity of B_0 gives $B = 0$, hence $d_1 = d_2 = d_3 = 0$ and $A \simeq \mathcal{O}_{\mathrm{dP}_6}$. Therefore $\nu^* i^* : \mathrm{Pic}(X) \to \mathrm{Pic}(\mathrm{dP}_6)$ is zero.

For $F_L = T_X \otimes L$ and $A_L = (L^\vee \otimes \omega_X)|_W$, duality and base change give

$$H^2(W, F_L|_W)^\vee \simeq H^0(W, \Omega^1_X|_W \otimes A_L), \qquad (R^2 f_* F_L)_{p_0} \otimes \mathbb{C} \simeq H^2(W, F_L|_W).$$

The conductor section and the just-proved Picard restriction show $R^2 f_*(T_X \otimes L) \neq 0$ for every $L \in \mathrm{Pic}(X)$.

A corrected CDP statement is: if one L_0 satisfies $H^2(X, T_X \otimes L_0) = 0$, then for general $M \in \mathrm{Pic}(X)$,

$$H^i(X, T_X \otimes M) = 0 \ (i \neq 1), \qquad \tfrac{1}{2}\langle c_3(T_X), [X]\rangle = -h^1(X, T_X \otimes M) \leq 0.$$

Relatively, one may instead assume one $R^2 f_*(T_X \otimes L) = 0$; for reduced full fibres, the exact replacement for the invalid local step is injectivity of the conductor connecting map δ_c on every whole fibre. The candidate fails this relative sufficient condition by the preceding conductor nonvanishing. Separately, because $c_3(X) = 2$, the contrapositive of the corrected global statement gives $H^2(X, T_X \otimes L_0) \neq 0$ for every L_0. Full local calculations are in `supporting_materials/workbench/research/analytic_picard_completion.tex` and `supporting_materials/workbench/research/cdp_program.tex`.

5. New cusp mathematics: Kummer extension, sphere carriers, and finite gauge systems

The subsequent work adds several precise results which are logically separate from the construction and recognition of X. They concern the local toric cusp, its metric limits, topological carriers for its expanding quotients, and a finite classical/Gaussian gauge model. None changes the provenance statement in Section 2: these are reconstruction and extension results in the AI-produced workbench, not claims of priority for the original candidate.

The local toric Kummer map and its two degrees. Write the completed cusp as $f_0 : N_0 = Y/\bar{\Lambda} \to \Delta_t$, with dense-torus action

$$\Psi_{\bar{\lambda}}(x,t) = \big(e^{2\pi i C(t)\bar{\lambda}} t^{B_0\bar{\lambda}} x, t\big), \qquad B_0 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

For an integer $N \geq 1$, let $\rho_N : \Delta_v \to \Delta_t$ be $\rho_N(v) = v^N$, take a copy $Y^{(N)}$ of the same toric variety, and replace the source multiplier by the holomorphic root

$$C_N(v) = \frac{1}{N} C(v^N), \qquad \Psi^{(N)}_{\bar{\lambda}}(\xi, v) = \big(e^{2\pi i C_N(v)\bar{\lambda}} v^{B_0\bar{\lambda}} \xi, v\big).$$

Its quotient and reduced central fibre are

$$f_N : N_0^{\langle N\rangle} := Y^{(N)}/\bar{\Lambda} \longrightarrow \Delta_v, \qquad W_N := \big(f_N^{-1}(0)\big)_{\mathrm{red}}.$$

Then the toric power map

$$P_N(\xi_1, \xi_2, v) = (\xi_1^N, \xi_2^N, v^N)$$

is equivariant and descends to a finite flat map

$$\kappa_N : N_0^{\langle N\rangle} \longrightarrow N_0, \qquad f_0 \circ \kappa_N = \rho_N \circ f_N, \qquad \deg \kappa_N = N^3, \qquad \kappa_N^* W = N W_N, \qquad \mathrm{Ram}(\kappa_N) = (N-1) W_N.$$

The exponent 3 is the degree over the *unchanged* cusp disc: two fibre coordinates and the base coordinate are all raised to the Nth power. After replacing the target by the base change $N_0[N] = N_0 \times_{\Delta_t} \Delta_v$, the factorization

$$N_0^{\langle N\rangle} \xrightarrow{Q_N} N_0[N] \longrightarrow N_0$$

has generic degree N^2. For $N > 1$, Q_N is quasi-étale but not flat; its non-étale and nonflat loci lie exactly over the double and triple strata. At a triple point its closed-fibre length is $3N^2 - 3N + 1 > N^2$, and at a double stratum the transverse closed-fibre length is $2N - 1 > N$. On the reduced central fibre, the generic degree is N^2 and after normalizing the reduced source and target central fibres, the induced map is toric multiplication $[N] : \mathrm{dP}_6 \to \mathrm{dP}_6$, ramified with index N along all six boundary curves. For $v \neq 0$, the relative fibre map $Q_{N,v}$ has deck group $(\mathbb{Z}/N\mathbb{Z})^2$, and

$$S_N^{-1} M_0^N S_N = M_0, \qquad S_N = \mathrm{diag}(1, 1, N, N).$$

This is a completed *local* cusp construction. It is not a degree-N^2 self-map over the original disc, not a normalization of the base change, and not yet a cover over the full three-point base.

Native and auxiliary pointed limits. Put $t_c = e^{-2\pi T}$ on the positive radial ray and write the marked period matrix and base point as

$$P_T = (v_{\gamma,T}\ v_{u,T}\ e_1\ e_3), \qquad F_T = \mathbb{R}^4/P_T\mathbb{Z}^4, \qquad o_T = [0].$$

Two primitive periods remain of length one, while the other two escape without hidden short integer cancellations: for constants $c_-, c_+ > 0$ independent of T,

$$\rho_T := \min_{(a,b)\neq(0,0)} \mathrm{dist}\big(a v_{\gamma,T} + b v_{u,T}, \mathbb{Z}\langle e_1, e_3\rangle\big) \geq T - c_-.$$

Consequently the exact native limit of the unmodified four-torus fibres F_T, in the unscaled Euclidean quotient metric, is

$$(F_T, o_T) \longrightarrow (\mathbb{T}^2 \times \mathbb{R}^2, (0,0)),$$

and every fixed-radius ball is eventually isometric to the corresponding ball in that product. An $\mathbb{R}^4$ pointed limit of those four-torus fibres themselves appears only after the added degree-M^2 local cover obtained by replacing the two short periods by M times those periods:

$$\Lambda_{T,M} = \mathbb{Z}\langle v_{\gamma,T}, v_{u,T}, Me_1, Me_3\rangle, \qquad \widetilde{F}_{T,M} = \mathbb{R}^4/\Lambda_{T,M},$$
$$\min\{T - c_-, M\} \leq \mathrm{sys}(\Lambda_{T,M}) \leq \min\{T + c_+, M\}.$$

For every integer-valued $M(T) \in \mathbb{Z}_{>0}$ with $M(T) \to \infty$, the pointed cover tori converge locally isometrically and in pointed Gromov–Hausdorff topology to $\mathbb{R}^4$; on each eventual Euclidean base-point neighbourhood this convergence is smooth. This

auxiliary cover is the regular-fibre restriction of the relative degree-M^2 Kummer factor above; the completed map to the original cusp family still has degree M^3.

Exact scalar counting and a fibrewise magnetic gap. Let

$$D_T := -\det_{\mathbb{R}} P_T = \mathrm{Vol}(F_T) = T^2 + O(T), \qquad N_T(E) := \#\{k \in \mathbb{Z}^4 : 4\pi^2 \|P_T^{-\mathsf{T}} k\|^2 \le E\}.$$

Counting Fourier modes with multiplicity gives, for every fixed $E \ge 0$,

$$N_T(E) = \pi D_T \sum_{n \in \mathbb{Z}^2} \left(\frac{E}{4\pi^2} - \|n\|^2 \right)_+ + O_E(T), \qquad (x)_+ := \max\{x, 0\}.$$

In particular, below the first transverse threshold $0 < E < 4\pi^2$,

$$N_T(E) = \frac{E D_T}{4\pi} + O_E(T) = \frac{E T^2}{4\pi} + O_E(T).$$

Thus the fixed-energy population has a two-dimensional leading density, even though every individual compact fibre has discrete spectrum.

There is also an exact full-four-torus statement which does not pass through an elliptic quotient. Fix a nondegenerate integral alternating form $\mathsf{E} = (E_{jk})$ on the marked lattice, let

$$d := |\operatorname{Pf}(\mathsf{E})| = |E_{12}E_{34} - E_{13}E_{24} + E_{14}E_{23}| > 0,$$

and let $\Delta_{\mathsf{E},T}$ be the Friedrichs magnetic Laplacian for the constant-curvature connection with curvature $2\pi i \mathsf{E}$. If

$$\operatorname{Spec}\Big((P_T^{\mathsf{T}} P_T)^{-1/2} (2\pi \mathsf{E}) (P_T^{\mathsf{T}} P_T)^{-1/2} \Big) = \{\pm i b_1, \pm i b_2\}, \qquad b_1, b_2 > 0,$$

then

$$\operatorname{Spec}(\Delta_{\mathsf{E},T}) = \{(2n_1+1)b_1 + (2n_2+1)b_2 : n_1, n_2 \in \mathbb{Z}_{\ge 0}\},$$

$$b_1 b_2 = \frac{4\pi^2 d}{D_T}, \qquad \operatorname{gap}(\Delta_{\mathsf{E},T}) = 2\min\{b_1, b_2\} \le 4\pi \sqrt{\frac{d}{D_T}} = O(T^{-1}).$$

This magnetic theorem is fibrewise for a fixed marked integral flux; it asserts neither full-monodromy invariance of that flux nor a Yang–Mills mass-gap result.

Exact S^{12} and S^{24} source maps and their metrics. Let

$$K_\Lambda = \mathbb{Z}\langle \widehat{w}, \widehat{\delta} \rangle, \qquad \overline{\mathbb{T}}^{\,2} = \frac{(\Lambda/K_\Lambda) \otimes \mathbb{R}}{\Lambda/K_\Lambda}.$$

Put

$$Y_T = \begin{pmatrix} 6 \operatorname{Im} \mu_T & \operatorname{Im} \tau_T \\ \operatorname{Im} \beta_T & \operatorname{Im} \mu_T \end{pmatrix}, \qquad Q_T = \mathbb{R}^2 / Y_T \mathbb{Z}^2, \qquad \iota_T : \overline{\mathbb{T}}^{\,2} \xrightarrow{\sim} Q_T, \quad [y] \longmapsto [Y_T y].$$

The quotient $\pi_T : F_T \twoheadrightarrow Q_T$ is the radial restriction of a quotient over the whole punctured cusp. Collapsing the central fibre and the exterior boundary gives the continuous surjective based quotient denoted q_2 in the proof; in this overview write it as

$$\rho_{2,\mathrm{cone}} : (X, x_0) \longrightarrow (\mathcal{A}_{\mathrm{cone}}, v_0), \qquad \mathcal{A}_{\mathrm{cone}} = \mathrm{Susp}(S^1 \times \overline{\mathbb{T}}^{\,2}).$$

The full-monodromy-invariant character lattice is $\mathbb{Z}\varepsilon_\gamma$; fix the displayed primitive generator. It gives a circle quotient on the regular base and explicit Seifert targets at the two finite fillings. Gluing them gives a compact orientable three-manifold $\mathcal{M}_{\mathrm{Seif}}$ with boundary $\mathbb{T}^2$; coning that entire boundary gives the map denoted q_1 in the proof, written here as

$$\mathcal{C}_{\mathrm{cone}} = \mathcal{M}_{\mathrm{Seif}} \cup_{\mathbb{T}^2} \mathrm{Cone}(\mathbb{T}^2), \qquad \rho_{1,\mathrm{cone}} : (X, x_0) \longrightarrow (\mathcal{C}_{\mathrm{cone}}, c_0).$$

There is also the cusp-supported target $\mathcal{B}_{\mathrm{cone}} = \mathrm{Susp}(S^1 \times S^1)$ and a based quotient $\rho_{1,\mathrm{cusp}} : X \to \mathcal{B}_{\mathrm{cone}}$. After a based choice $X \cong_{\mathrm{diff}} S^6$, basedness induces

$$\rho_{2,\mathrm{cone}}^{\wedge 2} : S^{12} \cong X^{\wedge 2} \longrightarrow \mathcal{A}_{\mathrm{cone}}^{\wedge 2}, \qquad \rho_{1,\mathrm{cone}}^{\wedge 4} : S^{24} \cong X^{\wedge 4} \longrightarrow \mathcal{C}_{\mathrm{cone}}^{\wedge 4}.$$

The corresponding Cartesian restrictions, before passage to the smash quotient and after identifying their radial images, are exactly

$$\rho_{2,\mathrm{cone}}^{\times 2}\Big|_{F_T^{\times 2}} = \pi_T^{\times 2} : F_T^{\times 2} \to Q_T^{\times 2}, \qquad \rho_{1,\mathrm{cone}}^{\times 4}\Big|_{F_T^{\times 4}} = \chi_{\gamma,T}^{\times 4} : F_T^{\times 4} \to C_T^{\times 4},$$

where

$$C_T = \mathbb{R}/\ell_T\mathbb{Z}, \qquad \ell_T = T + H_{\mathrm{cusp}} - B_{\mathrm{cusp}} + O(T^{-1}), \qquad H_{\mathrm{cusp}} = \operatorname{Im} h(0), \quad B_{\mathrm{cusp}} = \operatorname{Im} b(0).$$

Here are the target-slice embeddings, with no implicit identification. Put $r_T = e^{-2\pi T}$ and

$$\vartheta_T : \mathbb{R}/\mathbb{Z} \xrightarrow{\sim} C_T, \qquad [a] \longmapsto [\ell_T a].$$

In the suspension/cone coordinates used above,

$$\begin{aligned} j_T(y_1, y_2) &= \big[[r_T, 1, \iota_T^{-1}(y_1)], [r_T, 1, \iota_T^{-1}(y_2)]\big]_\wedge \in \mathcal{A}^{\wedge 2}_{\mathrm{cone}}, \\ k_T^{\mathcal{B}}(c_1, \dots, c_4) &= \big[[r_T, 1, \vartheta_T^{-1}(c_1)], \dots, [r_T, 1, \vartheta_T^{-1}(c_4)]\big]_\wedge \in \mathcal{B}^{\wedge 4}_{\mathrm{cone}}, \\ k_T^{\mathcal{C}}(c_1, \dots, c_4) &= \bigwedge_{j=1}^{4} [r_T, (1, \vartheta_T^{-1}(c_j))] \in \mathcal{C}^{\wedge 4}_{\mathrm{cone}}. \end{aligned}$$

Each is a topological embedding. With the intrinsic product quotient metrics,

$$(Q_T^{\times 2}, 0) \longrightarrow (\mathbb{R}^4, 0), \qquad (C_T^{\times 4}, 0) \longrightarrow (\mathbb{R}^4, 0)$$

pointed locally isometrically and in pointed Gromov–Hausdorff topology, with smooth convergence on the eventual Euclidean base-point neighbourhoods.

These copy numbers are sharp for the unrescaled quotient norm induced by the marked Euclidean fibre metric. For a connected torus quotient of r copies over the punctured cusp, divergence of the quotient systole forces the bounded lattice $K_\Lambda^{\oplus r}$ into the kernel, so the quotient rank is at most $2r$; rank four requires $r \geq 2$ and is attained by $Q_T^{\times 2}$. For a quotient over the full regular base, even after a finite base cover, monodromy also forces $\widehat{u}^{\oplus r}$ into the primitive kernel, so the quotient rank is at most r; rank four requires $r \geq 4$ and is attained by $C_T^{\times 4}$. Hence S^{12} and S^{24} are the smallest smash-sphere sources supplied by the native cusp-local and full-base quotient routes, respectively.

There are also genuine sphere targets. Collapsing the one-skeleton

$$K^{(1)} = ((\mathbb{R}/\mathbb{Z}) \times \{0\}) \cup (\{0\} \times (\mathbb{R}/\mathbb{Z})) \subset \overline{\mathbb{T}}^2$$

and forgetting the base angle give continuous surjective based quotient maps $\widehat{\rho}_3 : X \to S^3$ and $\widehat{\rho}_\gamma : X \to S^2$. Their smash powers are

$$\widehat{\rho}_3^{\wedge 2} : S^{12} \twoheadrightarrow S^6, \qquad \widehat{\rho}_\gamma^{\wedge 4} : S^{24} \twoheadrightarrow S^8.$$

Here $\widehat{\rho}_3 = \widehat{q}_3$ and $\widehat{\rho}_\gamma = \widehat{q}_\gamma$ in the detailed construction. Both full sphere carriers are based null-homotopic, by two distinct exact calculations. For every based map $f : S^6 \to S^2$,

$$S^{12} = S^6 \wedge S^6 \xrightarrow{f \wedge \mathrm{id}_{S^6} = \Sigma^6 f} S^2 \wedge S^6 = S^8 \xrightarrow{\mathrm{id}_{S^2} \wedge f} S^2 \wedge S^2 = S^4.$$

Since $\pi_{12}(S^8) \cong \pi_4^{\mathrm{S}} = 0$, the first arrow is based null-homotopic, and hence so is $f \wedge f$. Therefore

$$[f^{\wedge r}] = 0 \in \pi_{6r}(S^{2r}) \qquad \text{for every } r \geq 2,$$

because $f^{\wedge r} = (f \wedge f) \wedge f^{\wedge(r-2)}$. Taking $f = \widehat{\rho}_\gamma$ proves that the entire surjective quotient $\widehat{\rho}_\gamma^{\wedge 4} : S^{24} \twoheadrightarrow S^8$ is based null-homotopic.

For every based map $g : S^6 \to S^3$, the exact smash factorization is

$$g \wedge g = (\Sigma^3 g) \circ (\Sigma^6 g) : S^{12} \longrightarrow S^9 \longrightarrow S^6.$$

Let st denote stabilization. The exact low-stem data are

$$\pi_6(S^3) \cong \mathbb{Z}/12, \quad \pi_3^{\mathrm{S}} = \mathbb{Z}/24\langle\nu\rangle, \quad \pi_6^{\mathrm{S}} = \mathbb{Z}/2\langle\nu^2\rangle,$$

$$\pi_{12}(S^6) = \mathbb{Z}/2\langle\nu_6 \circ \nu_9\rangle, \qquad \mathrm{st} : \pi_{12}(S^6) \xrightarrow{\cong} \pi_6^{\mathrm{S}}.$$

Here the full group table is Hatcher, p. 339, and the signed smash formula and unstable stabilization calculation are Toda, Propositions 3.1 and 5.11. Since $12[g] = 0$, one has $\mathrm{st}([g]) = 2a\nu$ for some $a \in \mathbb{Z}/12$. Multiplicativity gives

$$\mathrm{st}([g \wedge g]) = \mathrm{st}([g])^2 = 4a^2\nu^2 = 0 \in \pi_6^{\mathrm{S}}.$$

The stabilization isomorphism therefore gives $[g \wedge g] = 0 \in \pi_{12}(S^6)$. Thus $g^{\wedge r} \simeq *$ for every $r \geq 2$, and taking $g = \widehat{\rho}_3$ proves

$$[\widehat{\rho}_3^{\wedge 2}] = 0 \in \pi_{12}(S^6), \qquad [\widehat{\rho}_\gamma^{\wedge 4}] = 0 \in \pi_{24}(S^8).$$

The one-copy classes of $\widehat{\rho}_3$ and $\widehat{\rho}_\gamma$ remain uncomputed. The unstable classes of the conical-target maps $\rho^{\wedge 2}_{2,\mathrm{cone}}$, $\rho^{\wedge 4}_{1,\mathrm{cusp}}$, and $\rho^{\wedge 4}_{1,\mathrm{cone}}$ also remain open: nullity after postcomposition with the angle-forgetting quotient does not imply nullity of the maps before that postcomposition. For each sufficiently large T, their targets contain moving topologically embedded four-spheres, equipped separately with their intrinsic quotient length metrics,

$$\mathbb{S}^4_{12,T} = S^2_T \wedge S^2_T \hookrightarrow S^6, \qquad \mathbb{S}^4_{24,T} = C^{\wedge 4}_T \hookrightarrow S^8,$$

where $S^2_T = \overline{\mathbb{T}}^2/K^{(1)}$ carries the quotient length metric pulled back from Q_T. Pointing these spaces at the midpoints $\zeta_{12,T}, \zeta_{24,T}$, rather than at their smash basepoints, every radius-R ball is exactly a Euclidean four-ball once $T - c_- > 4R$, respectively once $\ell_T > 4R$. Thus both moving four-spheres converge to $\mathbb{R}^4$ pointed locally isometrically and in pointed Gromov–Hausdorff topology, again smoothly on the eventual Euclidean neighbourhoods. The based inclusion maps

$$\mathbb{S}^4_{12,T} \hookrightarrow S^6, \qquad \mathbb{S}^4_{24,T} \hookrightarrow S^8$$

are null-homotopic: sliding the fixed suspension coordinate to an endpoint contracts every factor through based maps. Their four-dimensional fundamental classes therefore die in the ambient targets. These two slice-inclusion null-homotopies are separate from the global equalities $[\widehat{\rho}_3^{\wedge 2}] = 0 \in \pi_{12}(S^6)$ and $[\widehat{\rho}_\gamma^{\wedge 4}] = 0 \in \pi_{24}(S^8)$.

Deleting the common smash basepoint gives literal nonclosed topological four-cells in the sphere targets:

$$(Q_T \setminus K^{(1)}_T)^2 \xrightarrow{\cong} \mathbb{S}^4_{12,T} \setminus \{*\} \cong \mathbb{R}^4 \hookrightarrow S^6,$$

$$(C_T \setminus \{0\})^4 \xrightarrow{\cong} \mathbb{S}^4_{24,T} \setminus \{*\} \cong \mathbb{R}^4 \hookrightarrow S^8.$$

They are open only inside their embedded four-spheres, not in the ambient sphere, and no fixed compatible compact-target metric makes them globally isometric to Euclidean $\mathbb{R}^4$.

The metric scope is exact. For every unrescaled fixed metric compatible with any compact cone-target topology, the slices $j_T(Q_T^{\times 2})$ and $k^{\mathcal{B}}_T(C_T^{\times 4})$, and $k^{\mathcal{C}}_T(C_T^{\times 4})$ converge to the cone base point. Fixed cusp-adapted length metrics do admit moving-base-point blow-ups: after factors T/r_T and ℓ_T/r_T, the displayed slices converge to $\mathbb{R}^4$, while the full conical ambient bubbles are respectively $\mathbb{R}^8$ and $\mathbb{R}^{12}$. Their conic vertex links have dimensions 7 and 11, so the vertex cone dimensions are 8 and 12, not 4. By contrast, an independently chosen smooth Riemannian metric on the source sphere has ordinary tangent space $\mathbb{R}^{12}$ or $\mathbb{R}^{24}$. The quotient construction supplies no Riemannian metric on either source sphere.

Thus $\mathbb{R}^4$ is neither the whole compact source nor the whole compact target, and it cannot be a continuous quotient target of a compact sphere. What is proved is a literal nonclosed topological open cell, an intrinsic pointed limit of moving flat slices (not Sormani–Wenger intrinsic-flat convergence) and, for the fixed cusp-adapted target metrics, a four-plane inside a higher-dimensional moving-point bubble. No auxiliary finite-cover unspooling is used in these sphere routes, and no compact complex structure on S^{12} or S^{24} follows. Cartesian products also remain distinct from smash spheres: $X^{\times 2} \simeq S^6 \times S^6 \neq S^{12}$ and $X^{\times 4} \simeq (S^6)^{\times 4} \neq S^{24}$.

The compact carrier subspaces swept out by the synchronized slices are

$$\mathcal{D}_{12} \cong \mathcal{D}_{24} \cong \mathrm{Cone}(\mathbb{T}^4).$$

At their vertices the local integral homology has ranks $4, 6, 4, 1$ in degrees $2, 3, 4, 5$, respectively. Thus a vertex is not a homology-manifold point, and hence not locally a topological manifold (with or without boundary) of any dimension. The full smash targets retain still more topology:

$$\mathcal{A}^{\wedge 2}_{\mathrm{cone}} \simeq \bigvee^{9} S^4 \vee \bigvee^{18} S^5 \vee \bigvee^{15} S^6 \vee \bigvee^{6} S^7 \vee S^8,$$

$$\mathcal{C}^{\wedge 4}_{\mathrm{cone}} \simeq S^8 \vee \bigvee^{4} S^9 \vee \bigvee^{6} S^{10} \vee \bigvee^{4} S^{11} \vee S^{12}.$$

These are homotopy equivalences, not homeomorphisms.

There is nevertheless a space in which the limiting Euclidean fibre is literal rather than merely asymptotic. For $\nu \in \{12, 24\}$ put

$$A^{(12)}_T = \mathrm{diag}(Y_T, Y_T), \qquad A^{(24)}_T = \ell_T I_4,$$

and quotient $[0, s_0] \times \mathbb{R}^4$ fibrewise by $A^{(\nu)}_{1/s}\mathbb{Z}^4$ for $s > 0$, while imposing no identifications at $s = 0$. The resulting $\mathfrak{E}_\nu$ is a Hausdorff, second-countable topological five-manifold with boundary, with

$$p_{12}^{-1}(s) \cong Q^{\times 2}_{1/s}, \quad p_{24}^{-1}(s) \cong C^{\times 4}_{1/s} \quad (s > 0), \qquad p_\nu^{-1}(0) \cong \mathbb{R}^4.$$

There are continuous surjective comparison maps $\beta_\nu : \mathfrak{E}_\nu \twoheadrightarrow \mathcal{D}_\nu$ which identify every positive fibre with its radial carrier slice and collapse the entire $\mathbb{R}^4$ boundary fibre to the cone vertex. They are homeomorphisms off that boundary, but are

nonproper and nonquotient; in particular they are not topological blow-downs or resolutions, and $\mathcal{D}_\nu \not\cong \mathfrak{E}_\nu/\mathbb{R}^4$ via β_ν. Thus the requested Euclidean object is an intrinsic moving-slice limit and an actual boundary fibre of a noncompact comparison manifold, not the whole compact S^{12}, S^{24}, or compact quotient target.

Repaired cubical $SU(2)$ flux and the Gaussian theorem. The finite model keeps independent integers $M_{\mathrm{cov}} \geq 1$, the covering index, and $N_{\mathrm{mesh}} \geq 2$, the number of cubical subdivisions in each direction. Put

$$S_{M_{\mathrm{cov}}} = \mathrm{diag}(1, 1, M_{\mathrm{cov}}, M_{\mathrm{cov}}), \qquad P_{T,M_{\mathrm{cov}}} = P_T S_{M_{\mathrm{cov}}},$$

$$V_{N_{\mathrm{mesh}}} = (\mathbb{Z}/N_{\mathrm{mesh}}\mathbb{Z})^4, \qquad E_{N_{\mathrm{mesh}}} = V_{N_{\mathrm{mesh}}} \times \{1, 2, 3, 4\}.$$

Recall that $D_T = -\det_{\mathbb{R}} P_T = T^2 + O(T)$, so the cover covolume is $M_{\mathrm{cov}}^2 D_T$. With the normalization

$$\langle X, Y\rangle = -\tfrac{1}{2}\,\mathrm{tr}(XY), \qquad H_{\mathrm{link}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \qquad \|H_{\mathrm{link}}\|^2 = 1, \qquad \exp(2\pi H_{\mathrm{link}}) = I_2,$$

the periodic inverse-transport links

$$\begin{aligned} U_1(n) &= \exp\left(-\frac{2\pi n_2}{N_{\mathrm{mesh}}^2} H_{\mathrm{link}}\right), \\ U_2(n) &= \begin{cases} I_2, & 0 \leq n_2 \leq N_{\mathrm{mesh}} - 2, \\ \exp(2\pi n_1 H_{\mathrm{link}}/N_{\mathrm{mesh}}), & n_2 = N_{\mathrm{mesh}} - 1, \end{cases} \\ U_3(n) &= U_4(n) = I_2, \qquad n \in V_{N_{\mathrm{mesh}}}. \end{aligned}$$

give every $(1,2)$ plaquette the same value $\exp(2\pi H_{\mathrm{link}}/N_{\mathrm{mesh}}^2)$ and all other positively oriented coordinate plaquettes the identity. The principal-log curvature is exactly $2\pi H_{\mathrm{link}}\, dy^1 \wedge dy^2$, and the metric-weighted logarithmic action is

$$S^{\log}_{T,M_{\mathrm{cov}},N_{\mathrm{mesh}}} = \frac{2\pi^2 M_{\mathrm{cov}}^2}{g_{\mathrm{YM}}^2 D_T},$$

independent of the mesh. The explicitly defined continuum-matched Wilson sector has value

$$\frac{M_{\mathrm{cov}}^2 N_{\mathrm{mesh}}^4}{2 g_{\mathrm{YM}}^2 D_T}\left(2 - 2\cos\frac{2\pi}{N_{\mathrm{mesh}}^2}\right) \longrightarrow \frac{2\pi^2 M_{\mathrm{cov}}^2}{g_{\mathrm{YM}}^2 D_T}.$$

For $k = (k_1, \ldots, k_4) \in V_{N_{\mathrm{mesh}}}$, define the componentwise Fourier symbol

$$\mathfrak{d}_{N_{\mathrm{mesh}}}(k)_j = N_{\mathrm{mesh}}\left(e^{2\pi i k_j/N_{\mathrm{mesh}}} - 1\right), \qquad 1 \leq j \leq 4.$$

The reduced finite Coulomb space $\ker d^*_{N_{\mathrm{mesh}}} \cap (\ker \Delta^{(1)}_{N_{\mathrm{mesh}}})^\perp$ removes the complete harmonic $k = 0$ fibre. On that space, Fourier diagonalization of $\mathcal{L} = g_{\mathrm{YM}}^{-2} d^*_{N_{\mathrm{mesh}}} d_{N_{\mathrm{mesh}}}$ gives

$$\lambda_{T,M_{\mathrm{cov}},N_{\mathrm{mesh}}}(k) = \frac{1}{g_{\mathrm{YM}}^2}\left\| P_T^{-\mathsf{T}} S^{-1}_{M_{\mathrm{cov}}} \mathfrak{d}_{N_{\mathrm{mesh}}}(k)\right\|^2, \qquad k \neq 0,$$

with transverse complex multiplicity 9. Its Friedrichs Gaussian oscillator

$$\widehat{H}^{\mathrm{Gauss}} = -\tfrac{1}{2}\Delta + \tfrac{1}{2}\langle a, \mathcal{L} a\rangle$$

has a unique Gaussian ground state and exact gap $\sqrt{\lambda_{\min}(\mathcal{L})}$. The explicit sequence

$$T_j^{\mathrm{rad}} = j^2, \qquad M_{\mathrm{cov},j} = j, \qquad N_{\mathrm{mesh},j} = j^3$$

simultaneously gives a pointed local-isometric $\mathbb{R}^4$ cover limit, vanishing mesh size, vanishing Gaussian gap, and vanishing logarithmic classical action.

This last statement is a finite quadratic-oscillator theorem, not a four-dimensional interacting Yang–Mills mass-gap result. Its Hilbert space is L^2 of a noncompact linear Coulomb slice with Lebesgue measure, whereas compact-link lattice theory uses Haar measure on $SU(2)^{E_{N_{\mathrm{mesh}}}}$ followed by the nonlinear gauge constraint. The exponential is only a local chart, no unitary identification of these Hilbert spaces is constructed, a link configuration defines a Dirac distribution rather than an L^2 state, and there is no nonlinear Hamiltonian, transfer matrix, renormalized continuum limit, or Osterwalder–Schrader reconstruction in the result.

6. Split $SU(r)$, the $SU(3)$ Gram and Lorentz carriers, and the exact Erdős–Straus–Koide incidence bridge

The 3–4 September 2026 continuation separates the structures that first appeared together as a heuristic and then computes the strongest exact maps among them: a gauge field on each native four-torus; the two- and three-column $SU(3)$ orbit cones; a split Lorentz observation attached to the toroidal angular epimorphism; the full retained four-vector–spin-representation–scalar fibre; and the arithmetic incidence correspondence. Each now has a complete theorem, but the maps which would identify them with one another either fail by an explicit invariant or have not been constructed.

The full split-Cartan $SU(r)$ theorem on the native fibres. Fix $r \geq 2$ and a nonzero integral weight

$$\mathbf{m} = (m_1, \dots, m_r), \qquad \sum_i m_i = 0, \qquad s_2(\mathbf{m}) = \sum_i m_i^2.$$

For the full-flux line bundle $\mathscr{L}_{\eta,T} \to F_T$, put

$$\mathscr{E}_{\mathbf{m},T} = \bigoplus_{i=1}^{r} \mathscr{L}_{\eta,T}^{m_i}, \qquad H_{\mathbf{m}} = i\,\mathrm{diag}(m_1, \dots, m_r).$$

The ordered splitting gives a parallel determinant trivialization, and

$$c_1(\mathscr{E}_{\mathbf{m},T}) = 0, \qquad c_2(\mathscr{E}_{\mathbf{m},T}) = -\frac{s_2(\mathbf{m})}{2}[\eta_T]^2.$$

Since the marked and physical complex orientations are opposite,

$$\langle c_2, [F_T]_{\mathrm{mark}} \rangle = -6s_2(\mathbf{m}), \qquad k_{\mathbf{m}} := \big\langle c_2, [F_T]_{\mathrm{phys}} \big\rangle = 6s_2(\mathbf{m}).$$

The congruence $s_2(\mathbf{m}) \equiv \sum_i m_i = 0 \pmod 2$ shows that this split family realizes precisely a subset of the sectors $k_{\mathbf{m}} \in 12\mathbb{Z}_{>0}$.

The diagonal connection and curvature are

$$\nabla^{\mathbf{m},T} = d - 2\pi(6y^1\,dy^4 + y^2\,dy^3)H_{\mathbf{m}}, \qquad F_{\mathbf{m},T} = -2\pi H_{\mathbf{m}}\eta_T.$$

They solve the Yang–Mills equation on the flat torus. With $\langle X, Y \rangle = -\frac{1}{2}\,\mathrm{Tr}(XY)$, $D_T = \mathrm{Vol}(F_T)$, and $S_T = 6L_T - q_T$, the exact action is

$$\mathrm{YM}_T(\nabla^{\mathbf{m},T}) = \frac{2\pi^2 k_{\mathbf{m}}}{g_{\mathrm{YM}}^2} + \frac{\pi^2 s_2(\mathbf{m}) S_T^2}{g_{\mathrm{YM}}^2 D_T}.$$

Let $\mathcal{B}_{T,\min} \leq \mathcal{B}_{T,\max}$ be the two positive charge-one normal-form frequencies and

$$\rho_T^* = \frac{\mathcal{B}_{T,\max}}{\mathcal{B}_{T,\min}}, \qquad N_-(T) = \left\lceil \frac{\rho_T^* - 1}{2} \right\rceil.$$

For every positive root $i < j$, write $q_{ij} = |m_i - m_j|$. The complete negative spectrum of the normalized Coulomb Jacobi operator is

$$q_{ij}\big((2n+1)\mathcal{B}_{T,\min} - \mathcal{B}_{T,\max}\big), \qquad 0 \leq n < N_-(T),$$

and the real multiplicity contributed by this root pair is $12q_{ij}^2$. Coincident numerical bands are summed. Consequently

$$\mathrm{ind}_{\mathbb{R}}\,\mathcal{J}_{\mathbf{m},T} = 12N_-(T)\sum_{i<j}(m_i - m_j)^2 = 12rN_-(T)s_2(\mathbf{m}).$$

If the distinct weights have multiplicities $r_1, \dots, r_\ell$, the neutral Coulomb nullity away from a wall is

$$4\left(\sum_{a=1}^{\ell} r_a^2 - 1\right).$$

At an odd wall $\rho_T^* = 2n + 1 > 1$, charged zero modes add $12rs_2(\mathbf{m})$; at $S_T = 0$, where both favourable polarizations vanish, they add $24rs_2(\mathbf{m})$. On the deep cusp tail, $N_-(T) = 3$, so the index is $36rs_2(\mathbf{m}) = 6rk_{\mathbf{m}}$. For example, $\mathbf{m} = (1, 0, -1)$ in $SU(3)$ has

$$(q_{12}, q_{13}, q_{23}) = (1, 2, 1), \qquad k_{\mathbf{m}} = 12, \qquad \mathrm{ind}_{\mathbb{R}} = 216, \qquad \mathrm{null}_{\mathrm{neutral}} = 8$$

on that tail.

These negative eigenspaces are descent directions, not bifurcation kernels; only the wall zero modes are candidates for a Lyapunov–Schmidt analysis. The bundle itself cannot extend over X, S^{12}, or S^{24}: each proposed carrier has $H^4(-;\mathbb{Z}) = 0$, whereas $c_2(\mathscr{E}_{\mathbf{m},T}) \neq 0$.

The four-dimensional $SU(3)$ Gram quotient and its exact boundary. Let

$$\mathscr{V} = \mathrm{Mat}_{3\times 2}(\mathbb{C}), \qquad \Gamma(A) = A^\dagger A,$$

with $SU(3)$ acting by left multiplication. Equality of Gram matrices is exactly equality of $SU(3)$-orbits: after matching the two column spans by a unitary map, the unused complex line corrects its determinant uniquely. Thus

$$\mathscr{V}/SU(3) \cong \mathrm{Herm}_2^{\geq 0}.$$

The rank-two stratum is free and has the explicit trivialization

$$\mathscr{V}_{(2)} \cong SU(3) \times \mathrm{Herm}_2^{>0};$$

the nonzero rank-one stratum has stabilizer $SU(2)$, and the zero matrix has stabilizer $SU(3)$. If $A = (a\ b)$, then

$$\det \Gamma(A) = \|a\|^2\|b\|^2 - |\langle a, b\rangle|^2 = \|a \wedge b\|^2 = \sum_{i<j} |a_i b_j - a_j b_i|^2.$$

In Pauli coordinates this is

$$t^2 - x^2 - y^2 - z^2,$$

but the quotient is only the closed future cone $t \geq \sqrt{x^2+y^2+z^2}$, not all of Minkowski space. Its fixed-trace link is the Bloch ball B^3, with rank-one boundary $\mathbb{C}P^1 \cong S^2$. If $S^{12}_{\mathrm{col}} = \mathscr{V}^+$ is the one-point compactification of the representation, then

$$S^{12}_{\mathrm{col}}/SU(3) \cong \left(\mathrm{Herm}_2^{\geq 0}\right)^+ \cong B^4,$$

and the rank-one, zero, and infinity orbits together form its boundary S^3. This S^{12}_{col} is a representation sphere; it is not the earlier cusp-induced smash construction.

The four-variable identity

$$Q_4(r) = \frac{1}{2}\left(\sum_{i=1}^4 r_i\right)^2 - \sum_{i=1}^4 r_i^2 = h^2 - x^2 - y^2 - z^2$$

under the orthogonal Hadamard transform is exact. More generally,

$$Q_{4,\lambda} = (4\lambda - 1)h^2 - x^2 - y^2 - z^2;$$

therefore every $\lambda > 1/4$ has Lorentz signature after rescaling time. The coefficient $\lambda = 1/2$ is the unique choice retaining unit coefficients in this particular orthogonal coordinate system and the $\pi/4$ cone angle, but it is not forced by the $SU(3)$ orbit theorem.

The quotient does not supply a nonzero $SU(3)$ instanton sector: its principal bundle over $\mathrm{Herm}_2^{>0} \cong \mathbb{R}^4$ is trivial, and compactifying that open cone to S^4 requires an additional clutching class in $\pi_3(SU(3)) \cong \mathbb{Z}$. Conversely, the native torus bundle above is topologically the pullback of an S^4-bundle along a chosen degree-one collapse $c_T : F_T \to S^4$, because c_2 classifies $SU(3)$-bundles on four-complexes. Its native connection cannot be such a pullback. Indeed

$$\mathrm{Tr}(F_{\mathbf{m},T} \wedge F_{\mathbf{m},T}) = -4\pi^2 s_2(\mathbf{m})\eta_T^2$$

is nowhere zero. If it were $c^* \mathrm{Tr}(F_B \wedge F_B)$, then dc would have rank four everywhere, so $c : T^4 \to S^4$ would be a covering map, contradicting $\pi_1(T^4) = \mathbb{Z}^4$ and $\pi_1(S^4) = 0$.

Three columns: the cubic carrier, its rank-loss sphere, and the exact limit of the analogy. Now let

$$\mathscr{V}_3 = \mathrm{Mat}_{3\times 3}(\mathbb{C}).$$

For the left $U(3)$-action, $A \mapsto A^\dagger A$ is a complete orbit invariant and gives

$$\mathscr{V}_3/U(3) \cong \mathrm{Herm}_3(\mathbb{C})_{\geq 0}.$$

This is a nine-real-dimensional rank-three Euclidean Jordan cone. Its determinant is cubic, and its trace-one section is an eight-ball. After one-point compactification,

$$\mathscr{V}_3^+/U(3) \cong B^9,$$

whose boundary is exactly the compactified rank-deficient locus

$$\{\operatorname{rank} A \le 2\} \cup \{\infty\} \cong S^8.$$

Thus the passage from two to three columns replaces the quadratic $3+1$ Lorentz cone and its S^3 boundary by a cubic rank-three cone and an S^8 boundary at the level of the full orbit space. The full nine-dimensional cone is not itself a Lorentz cone; the marked angular epimorphism nevertheless gives a split positive observation onto the ordinary $3+1$ cone and identifies all five retained fibre directions.

Strict $SU(3)$ remembers one more invariant. The complete quotient map is

$$A \longmapsto (H, z) = (A^\dagger A, \det A), \qquad |z|^2 = \det H,$$

and its image is homeomorphic to $\mathbb{R}^{10}$. Hence

$$S^{17}/SU(3) \cong S^9, \qquad \mathscr{V}_3^+/SU(3) \cong S^{10}.$$

In this S^{10}, the same compactified rank-loss set is the standard topologically locally flat codimension-two S^8, and

$$S^{10} \setminus S^8 \cong \mathbb{R}^8 \times \mathbb{C}^* \simeq S^1.$$

The positive generator is the determinant phase $\det A/|\det A|$. The rank-one trace-one stratum is $\mathbb{C}P^2$. Quotienting the residual determinant-phase circle recovers B^9, with the S^8 as its fixed-set boundary. This supplies a precise three-input carrier and alternating invariant, but not by itself a principal bundle, connection, particle assignment, or Yang–Mills operator.

The exact Lorentz observation and all retained directions. The angular map already present in the toroidal quotient,

$$P = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix}, \qquad u = (1,1,1)^{\mathsf{T}}, \qquad Pu = 0,$$

has the standard-Hermitian right inverse

$$R = P^*(PP^*)^{-1} = \frac{1}{3}\begin{pmatrix} 2 & 1 \\ -1 & 1 \\ -1 & -2 \end{pmatrix}.$$

With $\Pi = uu^*/3$ and $Q_u = I_3 - \Pi$, define

$$\mathcal{O}_P(H) = PHP^*, \qquad \jmath_P(G) = RGR^*.$$

Both maps preserve positive semidefiniteness, and

$$\boxed{\mathcal{O}_P \jmath_P = \mathrm{id}, \qquad \jmath_P \mathcal{O}_P(H) = Q_u H Q_u.}$$

The complete linear kernel is

$$\ker \mathcal{O}_P = \{a\, uu^* + uv^* + vu^* : a \in \mathbb{R},\ v \in u^\perp\},$$

and its positive zero fibre is precisely $\{\lambda uu^* : \lambda \ge 0\}$.

Set $X = (R\ u)$. Since

$$\det X = 1, \qquad X^{-1} = \begin{pmatrix} P \\ u^*/3 \end{pmatrix},$$

every $H \in \mathrm{Herm}_3(\mathbb{C})$ has the unique form

$$H = X \begin{pmatrix} G & \psi \\ \psi^* & \lambda \end{pmatrix} X^*, \qquad G = PHP^*, \quad \psi = \frac{1}{3}PHu, \quad \lambda = \frac{1}{9}u^*Hu.$$

This is the complete real decomposition

$$\boxed{\mathrm{Herm}_3(\mathbb{C}) \cong \mathrm{Herm}_2(\mathbb{C}) \oplus \mathbb{C}^2 \oplus \mathbb{R}, \qquad 9 = 4+4+1,}$$

with cubic and positive fibre

$$\boxed{\det H = \lambda \det G - \psi^* \operatorname{adj}(G)\psi,}$$

$$H \succeq 0 \iff G \succeq 0, \quad \psi \in \operatorname{im} G, \quad \lambda \ge \psi^* G^+ \psi.$$

The embedded spin action is

$$T_P(g) = RgP + \Pi = X(g \oplus 1)X^{-1}, \qquad (G, \psi, \lambda) \longmapsto (gGg^*, g\psi, \lambda).$$

For $g \in SL_2(\mathbb{C})$, the summands are a Lorentz four-vector, the defining complex two-dimensional spin representation, and a real scalar. The central element $-I_2$ fixes G, λ and negates ψ; hence the four-vector action factors through $SO^+(1,3)$, while the full nine-dimensional carrier remembers the spin double cover.

The relation survives the relevant compactification in one direction:

$$\boxed{\jmath_P^+ : B^4 \cong \mathcal{P}_2^+ \hookrightarrow (\partial\mathcal{P}_3)^+ \cong S^8,} \qquad e_2(\jmath_P(G)) = \frac{1}{3}\det G,$$

and its normal opening at uu^* is

$$\boxed{\det(uu^* + \varepsilon \jmath_P(G)) = \varepsilon^2 \det G.}$$

The reverse boundary observation is nonproper. In the strict $SU(3)$ quotient the determinant phase satisfies

$$|\omega|^2 = \lambda \det G - \psi^* \operatorname{adj}(G)\psi,$$

and $\omega = 0$ is exactly the rank-loss locus compactifying to the distinguished $S^8 \subset S^{10}$.

The four-branch chart reaches this Lorentz sector through the holomorphic seed

$$K(z) = \operatorname{diag}(z_0 z_3, z_1 z_2), \qquad \det K = z_0 z_1 z_2 z_3.$$

Its positive $SL_2(\mathbb{C})$-orbit supplies every future causal direction. For $Z \in \mathrm{Mat}_{3\times 2}(\mathbb{C})$ with the commuting colour and spin actions, retain $\alpha, c > 0$ and put

$$p_\alpha(Z) = \alpha Z^* Z, \qquad m^2_{\alpha,c}(Z) = \frac{\alpha^2}{c^2} \sum_{a<b} |Z_{a1}Z_{b2} - Z_{a2}Z_{b1}|^2.$$

The map p_α separates $SU(3)$-orbits and is Lorentz covariant; the mass is positive exactly at complex rank two. Since

$$m^2_{\alpha,c}(rZ) = r^4 m^2_{\alpha,c}(Z),$$

the unrestricted pointwise carrier is gapless. Exact presymplectic reduction now gives four distinct results. For $m_0 > 0$, the rank-two colour quotient and its characteristic reduction are

$$(\mathcal{J}_2 \times \mathcal{H}^+_{m_0,c})/((x,p) \sim (x + sp, p)) \cong_{\mathrm{symp}} T^*\mathcal{H}^+_{m_0,c},$$

exactly the future-timelike spinless particle. On nonzero rank one, the intrinsic characteristic quotient is $T^*\mathcal{N}^\times_+$, but the written massless multiplier action is irregular because

$$d(\det \circ p_\alpha)_Z = 0.$$

It yields $\dot{x} = \lambda p$ without $\lambda = e$, has four transverse rank-opening directions which are first-order null but not colour gauge, and at $Z = 0$ admits arbitrary $x(\tau)$. For the separate hard constraint with $\omega_0 \neq 0$, the momentum map has rank four above its sharp threshold and rank three at the endpoint; its action therefore forces $\dot{x} = 0$ above threshold and has the ordinary $\dot{x} \in \mathbb{R}p$ characteristic only at threshold. The accompanying multiplication/Fock spectrum is exact as a free operator calculation but is not the Hamiltonian generating this action, so its positive endpoint is a constraint-selected threshold rather than a dynamically generated interacting gap. Complete definitions, maps, stabilizers, reduced forms, equations, and singular counterexamples are proved in the canonical module `supporting_materials/workbench/research/worldline_presymplectic_reduction.tex`.

The affine normal slice is now developed as an exact field-theoretic host. Indeed,

$$\mathcal{N}_P = \{Y \in \mathrm{Herm}_3(\mathbb{C}) : Yu = 0\} = \jmath_P\big(\mathrm{Herm}_2(\mathbb{C})\big), \qquad \jmath_P^{-1}(Y) = PYP^*,$$

and polarization of

$$\det(uu^* + \varepsilon \jmath_P(G)) = \varepsilon^2 \det G$$

gives the constant Minkowski form of signature $(1,3)$. Positive semidefinite G are the future cone, rank-one positive G its null boundary, and the lifted congruence action realizes the spin cover $SL_2(\mathbb{C}) \to SO^+(1,3)$. The standard metrics used in the Moore–Penrose section further select

$$W_P = R^*R = (PP^*)^{-1}, \qquad K_P = \{a \in SL_2(\mathbb{C}) : a^* W_P a = W_P\} \cong SU(2).$$

Its real normal representation is exactly $\mathbf{1} \oplus \mathbf{3}$, with trivial central action. Promoting this pointwise action to a local gauge symmetry gives the explicit nonnegative Euclidean model

$$\int_M \left(\frac{|F_{\mathcal{A}}|^2}{2g_{\mathrm{YM}}^2} + \frac{Z_G}{2} |D_{\mathcal{A}} G|^2_{E_P} + \frac{m_G^2}{2} |G|^2_{E_P} + \frac{\lambda_4}{4} (\det G)^2 \right) d\,\mathrm{vol}_g,$$

where $E_P(G,K) = \mathrm{tr}(GW_PKW_P)$ and $D_{R\mathcal{A}P\jmath P}(G) = \jmath_P(D_{\mathcal{A}}G)$. The normal field cannot screen a fundamental centre charge; the retained spinor ψ is centre-nontrivial, but a kinetic and interaction theory for that additional direction has not yet been constructed.

The distinction between the candidate cusp matrix and the Gram momentum is also now an equation:

$$\det G_{\eta,T} = -\frac{2}{3} D_T < 0, \qquad \mathrm{im}\, p_\alpha = \mathrm{Herm}_2(\mathbb{C})_{\geq 0}.$$

Positive scaling and complex congruence preserve their different inertia types, so any common physical model requires a new morphism rather than an identification. After an independent length/time dictionary is chosen, the same Lorentzian host carries a standard Klein–Gordon/Fock model with exact gap m_{KG} for $m_{\mathrm{KG}} > 0$ and gap zero for $m_{\mathrm{KG}} = 0$. The massless transverse gauge Gaussian has lowest nonzero frequency $2\pi/L$, which closes as $L \to \infty$, and explicit Hamiltonian countermodels show that entanglement alone determines neither the presence nor the size of a gap. An auxiliary positive gap transfers to a physical Hamiltonian only when a comparison map controls the entire excitation form domain:

$$\|Ru\| \geq c_R \|u\|, \quad \mathfrak{b}[Ru] \leq C_R(\mathfrak{a}_{\mathrm{phys}}[u] - E_{\mathrm{vac}}\|u\|^2) \Longrightarrow \mathrm{gap}(A_{\mathrm{phys}}) \geq \frac{\gamma c_R^2}{C_R}.$$

An intertwiner on a single selected subspace is insufficient. Thus a field-theoretic kinetic operator has now been constructed on the exact Lorentz slice, but dynamics for the complete $(G, \psi, \lambda, \omega)$ carrier, a gauge-invariant interacting continuum spectrum, and the comparison map required for a nonlinear mass-gap theorem remain exact construction problems; the present theorem neither proves nor disproves the four-dimensional Yang–Mills mass gap.

The actual toroidal successor and its elliptic quotient. Independently of the Gram quotient, the rank-three Kuhn fan produces a proper analytic degeneration

$$\pi_3 : N_3 \longrightarrow \Delta, \qquad N_{3,t} \cong E_\tau^3 \quad (t \neq 0),$$

with cusp monodromy

$$M_3 = \begin{pmatrix} I_3 & I_3 \\ 0 & I_3 \end{pmatrix}$$

and deepest local equation $t = z_0 z_1 z_2 z_3$. The irreducible central fibre has one dense three-torus, seven double surfaces, twelve triple curves, and six quadruple points:

$$(f_0, f_1, f_2, f_3) = (1, 7, 12, 6), \qquad \chi(W_3) = 6.$$

Its normalization is the permutohedral toric threefold, with orbit vector

$$(1, 14, 36, 24) \longrightarrow (1, 7, 12, 6);$$

the positive-codimension strata map with degrees $2, 3, 4$, respectively. Consequently, any real S^8-total-space fibration retaining W_3 must have its other singular fibres contribute Euler sum -4. For a full-rank integral shear $\mathcal{B} : \bar{L}_3 \hookrightarrow L_3$, the calculation retains the entire lattice index:

$$\chi(W_{3,\mathcal{B}}) = 3!\,[L_3 : \mathcal{B}\bar{L}_3] = 3!\,|\det \mathcal{B}|.$$

This is the precise content of the observed $6 = 3! \det I_3$. It is a toroidal volume/index theorem, not a proof that the Hermitian Jordan cone is a globalization of the degeneration.

The selectively corrected Jordan determinant/Euler calculation adds an exact algebraic bridge at the level of determinant forms. Fixing the orientation and writing $[a]_\times v = a \times v$, every 3×3 Hermitian matrix is uniquely $S + i[a]_\times$, with

$$\boxed{\det\big(S + i[a]_\times\big) = \det S - a^{\mathsf{T}} S a.}$$

For a marked positive-definite symmetric monodromy form B, the section

$$\ell_B(\lambda, a) = \lambda B + i[a]_\times$$

meets the positive-semidefinite cone in exactly

$$\lambda \geq 0, \qquad a^{\mathsf{T}} B a \leq \lambda^2 \det B,$$

the future cone of a $(3+1)$-Lorentz form. There is also an explicit real-linear order-cone bijection

$$\Theta : \mathrm{Herm}_2(\mathbb{C})_{\geq 0} \longrightarrow (\mathbb{R} I_3 \oplus i\mathfrak{so}(3)) \cap \mathrm{Herm}_3(\mathbb{C})_{\geq 0}$$

such that

$$\det_3 \Theta(H) = \frac{\operatorname{tr} H}{2} \det_2 H.$$

It sends positive-semidefinite ranks $2, 1, 0$ to $3, 2, 0$. Separately, if P is a rank-one projection in $\mathrm{Herm}_2(\mathbb{C})$, then $\Theta(P)$ has eigenvalues $0, 1/2, 1$ and is not idempotent; hence Θ is not a Jordan-algebra homomorphism. These identities exhibit literal Lorentz sections inside the cubic cone, but they contain no torus lattice, monodromy, period family, or filling map. The toric parameter tends to zero with $B = I_3$ fixed; this is not the same as moving to $\det B = 0$.

There is nevertheless a literal map to the reconstructed rank-two cusp. In the marked angular and deck lattices, set

$$P = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix}, \qquad Q = \begin{pmatrix} 0 & -1 & 1 \\ 1 & -1 & 0 \end{pmatrix}, \qquad \Phi = P \oplus Q.$$

Then $\Phi : H_1(T^6) \twoheadrightarrow H_1(T^4)$ satisfies

$$\boxed{\Phi M_3 = M_F \Phi, \qquad c_F \Phi = Q c_3.}$$

If the candidate cusp period is $Z_2(t) = sB_0 + C(t)$, the source period

$$Z_3(t) = sI_3 + S\, C(t)\, Q, \qquad S = \begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 0 & 0 \end{pmatrix},$$

obeys

$$\boxed{P Z_3(t) = Z_2(t) Q.}$$

Consequently $P : \mathbb{C}^3 \to \mathbb{C}^2$ descends to a surjective holomorphic map of every smooth cusp fibre, has connected elliptic kernel, and extends over the toric boundary. An explicit unimodular basis change splits the source cusp integrally as

$$M_3 \cong M_F \oplus \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

The candidate's free affine order-three and order-four peripheral actions also lift on this added elliptic factor. The remaining problem is analytic, not merely numerical: one still has to build a single period and overlap family across both finite-order discs.

This successful lift is not the diagonal local-system candidate

$$M_1 = I_3 \otimes S, \qquad M_2 = I_3 \otimes Q, \qquad M_0 = I_3 \otimes T, \qquad M_1 M_2 M_0 = I.$$

Although its finite factors have orders 4 and 6, each $I - M_j : T^6 \to T^6$ is a surjective isogeny for $j = 1, 2$. Thus every affine lift of either finite action has a fixed point.

The six Kuhn tetrahedra also meet the Gram construction exactly at the level of marked fans. With $\nu_i = E_{ii} - I_3/3$, the π-th tetrahedron maps to the chamber

$$\mathbb{R}_{\geq 0} \nu_{\pi_1} + \mathbb{R}_{\geq 0}(-\nu_{\pi_3}) = \{\lambda_{\pi_1} \geq \lambda_{\pi_2} \geq \lambda_{\pi_3}\}.$$

Thus the six deepest toroidal strata and the six A_2 Weyl chambers form the same S_3-torsor after marking. This Coxeter correspondence does not identify the analytic germs: the toroidal germ has four branches, while the rank-one Gram normal determinant is an irreducible quadratic.

The tempting cyclic compression of the six Kuhn labels to two also needs this distinction. The three even labels and three odd labels do map, through the saturated rank-lowering homomorphism $P : \mathbb{Z}^3 \twoheadrightarrow \mathbb{Z}^2$, to the two unimodular triangle orbits of the target fan. They do not arise from the asserted index-three translation superlattice: any unimodular triangulation of a rank-three lattice fundamental domain again has six maximal tetrahedra.

The free $\mathbb{Z}^6$ belonging to this nine-dimensional-cone program is

$$\Lambda_{\mathrm{tor}} = H_1(T^6; \mathbb{Z}) = \mathbb{Z}^3_{\mathrm{angular}} \oplus \mathbb{Z}^3_{\mathrm{deck}}.$$

It must not be confused either with $\mathbb{Z}/6$ or with the different rank-six bundle-defect group on the original regular locus. Under diagonal coordinate permutation, $P \oplus Q$ gives

$$0 \longrightarrow \mathbb{Z}(k, 0) \oplus \mathbb{Z}(0, k) \longrightarrow \Lambda_{\mathrm{tor}} \longrightarrow A_F \oplus B_F \longrightarrow 0, \qquad k = (1, 1, 1).$$

This sequence splits as an integral lattice with monodromy, but not as an integral S_3-module. The quotient is two copies of the A_2 weight lattice, and the associated root lattices have index three. The six homology basis elements are two three-element permutation orbits; the six Kuhn/Weyl chambers are one regular S_3-orbit. Thus $\operatorname{rk}\Lambda_{\mathrm{tor}} = 6$, $|S_3| = 3! = 6$, and $\chi(W_{3,\mathcal{B}}) = 3!|\det\mathcal{B}|$ are not three names for one object. The last two share a toric cell count; the first comes from angular plus deck homology.

The regular six-set does, however, have an exact home in the associated nine-dimensional carrier. Conjugation by permutation matrices gives

$$\operatorname{Hom}(\overline{L}_3, L_3) = \bigoplus_i \mathbb{Z}E_{ii} \oplus \bigoplus_{i\neq j} \mathbb{Z}E_{ij},$$

and the six off-diagonal units form one regular S_3-orbit. The marked chamber $\pi = (\pi_1\pi_2\pi_3)$ corresponds to E_{π_1,π_3}, or equivalently to the mixed angular–deck pair whose alternating pairing is zero. The three excluded diagonal pairs have pairing one. A real-linear equivariant isomorphism identifies the full matrix representation with $\mathrm{Herm}_3(\mathbb{C})$, but the symmetric/skew integral splitting has index 2^3, and the six off-diagonal images are indefinite. This is therefore a precise representation/incidence bridge, not a positive, determinant, Jordan, or toric morphism.

The only map presently reaching the positive-semidefinite Hermitian Gram boundary is the piecewise-linear chamber map $\Phi_{\mathrm{G}} \circ \jmath \circ P$, which uses the angular half, is not additive, and does not descend to a torus fibre. Full proofs are in the workbench sections *The rank-six cusp lattice inside the nine-dimensional cone program* and *The chamber torsor inside the nine-dimensional matrix carrier.*

Why the higher-rung object cannot be a literal complex fourfold S^8. If a homotopy S^8 admitted an almost-complex tangent bundle E, then

$$c_1(E) = c_2(E) = c_3(E) = 0, \qquad c_4(E) = 2u,$$

and therefore

$$p_2(TS^8) = c_2^2 - 2c_1c_3 + 2c_4 = 4u.$$

But $p_1 = 0$ and $\operatorname{sign}(S^8) = 0$, so Hirzebruch's formula

$$\operatorname{sign}(S^8) = \left\langle \frac{7p_2 - p_1^2}{45}, [S^8] \right\rangle$$

forces $p_2 = 0$, a contradiction. Thus the rank-loss S^8 above is a real stratified boundary, not a smooth complex-fourfold analogue of the candidate. A nonsingular fibration $T^6 \to S^8 \to S^2$ is also impossible: its homotopy sequence would require a surjection $\mathbb{Z} \to \mathbb{Z}^6$, while Euler characteristic would require $2 = 0$. Any real singular replacement must instead derive the actual local filling maps; its special fibres obey

$$\sum_i \chi(F_i) = 2.$$

For regular-piece monodromy ρ, the corrected homological Serre filtration is

$$0 \longrightarrow (\textstyle\bigwedge^q \Lambda)_\rho \longrightarrow H_q(E;\mathbb{Z}) \longrightarrow H_1(B^\circ; \bigwedge^{q-1}\Lambda) \longrightarrow 0,$$

whereas the cohomological edge map has invariant quotient

$$H^q(E;\mathbb{Z}) \twoheadrightarrow (\textstyle\bigwedge^q \Lambda^\vee)^\rho.$$

Consequently its van Kampen, Wang, and Mayer–Vietoris maps must kill the regular fibre group and all intermediate integral homology while respecting the coinvariant subobject and the base-one-cycle quotient; using the reverse Serre arrows does not meet this requirement. Every effective torus action on S^8 has rank at most four, and this bound is attained by the coordinate action

$$S^8/T^4 \cong \operatorname{Susp}(\Delta^3) \cong B^4.$$

This is a real two-pole singular compactification with generic T^4 orbit, not a three-special-fibre T^6-fibration over S^2.

The exact interface with the Fabel–Alpöge degeneration. The present interface does not identify the rank-loss S^8 with the candidate threefold. It occurs on the regular locus of the reconstructed S^6. Put

$$D = W \sqcup S_1 \sqcup S_2, \qquad U = X \setminus D,$$

where W is the non-normal central fibre and S_1, S_2 are the two multiple bielliptic fibres. Their computed first homology and Alexander duality give the exact map

$$\boxed{H_1(W;\mathbb{Z}) \oplus H_1(S_1;\mathbb{Z}) \oplus H_1(S_2;\mathbb{Z}) \xrightarrow{\sim} H^4(U;\mathbb{Z}) \cong \mathbb{Z}^2 \oplus \mathbb{Z}^2 \oplus \mathbb{Z}^2.}$$

Because U has CW dimension at most five, this complete rank-six group classifies principal $SU(3)$-bundles on U by c_2. The pair sequence gives another isomorphism

$$\delta : H^4(U;\mathbb{Z}) \xrightarrow{\sim} H^5(X, U;\mathbb{Z}),$$

so an unframed bundle extends across all three special fibres exactly when its six-component class vanishes. Global $SU(3)$-bundles on S^6 carry instead the separate clutching integer

$$k \in \pi_5(SU(3)) \cong \mathbb{Z}, \qquad c_3 = \pm 2ku,$$

and every such global bundle restricts trivially to U.

The displayed $\mathbb{Z}^6$ is a free lattice consisting of three geometrically localized $\mathbb{Z}^2$ summands. It is not itself a cyclic $\mathbb{Z}/6\mathbb{Z}$. Write its marked basis as

$$(a_0, b_0, a_1, b_1, a_2, b_2) = (\bar{\gamma}, \bar{u}, g_1, h_1, g_2, h_2).$$

The calculations below are proved in the integrated workbench section *Fibre degree and the cyclic-six centre*, including the one-shot collar matrix and the complete kernel, coefficient, tensor-centre, and common-centre arguments. They have the workbench's internal *Verified* status as lattice and matrix-group statements; that status does not extend them to a gauge-bundle or spectral conclusion. With the complex orientation of a smooth fibre, outward-first boundary orientation, and the manuscript's clockwise meridians, the transpose of the actual collar matrix gives

$$\operatorname{res}_F\left(\sum_{i=0}^{2}(n_i a_i + r_i b_i)\right) = (12n_0 + 4n_1 - 3n_2)\omega_F, \qquad [\deg_F] = (12, 0, 4, 0, -3, 0),$$

where $\langle \omega_F, [F] \rangle = 1$. This primitive map has saturated kernel

$$\ker \deg_F = \mathbb{Z}\langle b_0, b_1, b_2,\ a_0 - 3a_1,\ 3a_1 + 4a_2 \rangle.$$

It is the pullback of the topology-killing functional, not a new numerical coincidence. Define

$$q_{\mathrm{col}} : L \longrightarrow H_1(X^\circ; \mathbb{Z}), \qquad a_0 \mapsto c, \quad a_1 \mapsto \bar{g}_1, \quad a_2 \mapsto \bar{g}_2, \quad b_0, b_1, b_2 \mapsto 0,$$

and let $\Phi_1 : H_1(X^\circ; \mathbb{Z}) \to \mathbb{Z}$ have row $(12, 4\ell_1, 3\ell_2)$ in the basis $(c, \bar{g}_1, \bar{g}_2)$. Then

$$\deg_F = \Phi_1 \circ q_{\mathrm{col}}, \qquad \Phi_1(j_*[\widetilde{a}_0]) = 12\ell_0 - 4\ell_1 - 3\ell_2 = p.$$

The order-three and order-four normal-line holonomies give more than the formal reduction of an arbitrary rank-six lattice. If $\vartheta_1(a_1) = e^{-2\pi i/3}$, $\vartheta_2(a_2) = e^{-2\pi i/4}$, and both characters are trivial on the other marked generators, and $\zeta_{12} = e^{-2\pi i/12}$, then the candidate twists $(\ell_1, \ell_2) = (1, -1)$ select

$$\chi_{12}(x) = \vartheta_1(n_1 a_1 + r_1 b_1)\vartheta_2(n_2 a_2 + r_2 b_2)^{-1} = \zeta_{12}^{4n_1 - 3n_2}, \qquad \chi_6 = \chi_{12}^2,$$

and, for $\rho_m(k) = e^{-2\pi i k/m}$,

$$\boxed{\chi_{12} = \rho_{12} \circ \deg_F, \qquad \chi_6 = \rho_6 \circ \deg_F\,.}$$

Consequently

$$H^4(U; \mathbb{Z})/\ker \chi_{12} \cong \mathbb{Z}/12\mathbb{Z}, \qquad H^4(U; \mathbb{Z})/\ker \chi_6 \cong \mathbb{Z}/6\mathbb{Z}.$$

This distinguished cyclic quotient is not ordinary coefficient reduction:

$$H^4(U; \mathbb{Z}/6) \cong (\mathbb{Z}/6)^6.$$

For a principal $SU(3)$-bundle $P \to U$, this last identity is the exact topological congruence

$$\chi_6(c_2(P)) = \exp\Big(-\frac{2\pi i}{6}\langle c_2(P)|_F, [F]\rangle\Big)\,.$$

The full order-four normal holonomy is essential before squaring: the natural combined character has order twelve, not six.

There is a separate exact group-theoretic occurrence of the same finite centre. Tensor product defines the proper closed embedding

$$\iota : SU(2) \times SU(3) \hookrightarrow SU(6), \qquad \iota(A, B) = A \otimes B,$$

and its restriction $\mu_2 \times \mu_3 \xrightarrow{\sim} \mu_6 = Z(SU(6))$ sends (λ, μ) to $\lambda\mu$. Since U has homotopy dimension at most five, c_2 classifies $SU(n)$-bundles for $n \geq 3$. If $e = c_2(E)$ and $f = c_2(F_3)$ for determinant-trivial bundles of ranks two and three, then the splitting principle gives

$$c_2(E \otimes F_3) = 3e + 2f.$$

Hence every $SU(6)$-bundle class $h \in H^4(U; \mathbb{Z})$ admits at least one tensor-subgroup reduction, for example $(e, f) = (h, -h)$; all solutions at the c_2-level are

$$(e, f) = (h + 2t, -h - 3t), \qquad t \in H^4(U; \mathbb{Z}),$$

so neither a nonzero h nor a reduction is canonically selected. Moreover

$$H^5(U;\mathbb{Z}) \cong \ker\{\mathbb{Z}^3 \xrightarrow{(1,1,1)} \mathbb{Z}\} \cong Q(A_2),$$

after equipping the distinguished three-component support basis with the combinatorial orthonormal form. The resulting simple-root Gram matrix is $\left(\begin{smallmatrix} 2 & -1 \\ -1 & 2 \end{smallmatrix}\right)$; it is not a cup-product or intersection form. This support-redistribution lattice is not the defect lattice and does not make its rank-five fibre-degree kernel an A_5 root lattice.

The isomorphism $H^4(U)/\ker\chi_6 \cong Z(SU(6))$ is an isomorphism of finite groups, not a system of $SU(6)$ transition functions or a connection. In particular, the injective tensor map does not factor through $(SU(2)\times SU(3))/\mu_6$: its central μ_6 maps to nontrivial scalar matrices. The distinct common-centre construction is

$$\Gamma_6 = \{((\lambda\mu)^{-1}, \lambda I_2, \mu I_3) : \lambda \in \mu_2,\ \mu \in \mu_3\}, \qquad \big(U(1)\times SU(2)\times SU(3)\big)/\Gamma_6 \hookrightarrow U(6), \qquad [z,A,B] \longmapsto z(A\otimes B),$$

where $\Gamma_6 \cong \mu_6$. Its determinant is z^6, and its image meets $SU(6)$ in exactly $\iota(SU(2)\times SU(3))$; the image in $U(6)$ is not generally contained in $SU(6)$. What remains unproved is a degree-changing morphism identifying this degree-four number with the degree-two lifting obstruction of a common-centre quotient gauge bundle and with its electric–magnetic line pairing. No mass-gap statement follows before that bundle, a connection, operator, domain, and spectral comparison have been constructed.

The audited Erdős–Straus selector and its Koide boundary model. Let $P \equiv 1 \pmod 4$ be prime and let $\mathscr{S}(P)$ be the set of sorted positive solutions of

$$\frac{4}{P} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

Every solution has a unique least denominator

$$x = a_j = \frac{P + R_j}{4}, \qquad R_j = 4j+3, \qquad 0 \le j < \frac{P-1}{4}.$$

For $a = a_j$, every divisor $u \mid a^2$ belongs to one of two tagged selector channels,

$$E_j:\ R_j \mid 4u+1, \qquad M_j:\ R_j \mid 4u+P.$$

For $\varepsilon = 0$ on E_j and $\varepsilon = 1$ on M_j, put

$$T = \frac{4u + P^{\varepsilon}}{R_j}, \qquad K = \frac{P^{1-\varepsilon}T + 1}{4}.$$

Then

$$\frac{4}{P} = \frac{1}{a} + \frac{1}{P^{\varepsilon}aK/u} + \frac{1}{PK}.$$

Middle-channel complementation $\sigma(u) = a^2/u$ exchanges the last two denominators and has no fixed point. The corrected selector theorem is the bijection

$$\mathcal{E}(P) \ \sqcup\ \big(\mathcal{M}(P)/\langle\sigma\rangle\big) \xrightarrow{\sim} \mathscr{S}(P).$$

The missing point in the supplied proof was that a_j remains the unique least denominator: if another denominator $b < a_j$, then $1/a_j + 1/b > 2/a_j \ge 4/P$, contradicting positivity of the third summand.

Writing $v = a^2/u$ gives the positive rank-one matrix

$$\Gamma(a,u) = \begin{pmatrix} u & a \\ a & v \end{pmatrix}.$$

It has the unique primitive factorization

$$u = hr^2, \qquad a = hrs, \qquad v = hs^2, \qquad \gcd(r,s) = 1,$$

and the two selector congruences become

$$E_j:\ R_j \mid Pr + s, \qquad M_j:\ R_j \mid r + s.$$

Consequently the unresolved prime case of the Erdős–Straus conjecture is equivalently the assertion that for every prime $P \equiv 1 \pmod{12}$ there exist $h, r, s > 0$ with

$$\gcd(r,s) = 1, \qquad \frac{P}{4} < hrs < \frac{P}{2}, \qquad R = 4hrs - P, \qquad R \mid Pr + s \ \text{ or } \ R \mid r + s.$$

This is an exact reformulation; it does not prove the required nonemptiness.

The accompanying terminal matrix is also exact. With

$$\ell_0 = \frac{T+1}{4}, \qquad \ell_1 = u - jT, \qquad \ell_2 = 1,$$

$$\lambda = 12\ell_0 - 4\ell_1 - 3\ell_2 = P^{\varepsilon}, \qquad \mathsf{R} = \begin{pmatrix} 3 & -\ell_1 \\ 4 & 1 - 4\ell_0 \end{pmatrix}, \qquad \det \mathsf{R} = -P^{\varepsilon},$$

and $\mathrm{SNF}(\mathsf{R}) = \mathrm{diag}(1, P^{\varepsilon})$. The pair (Γ, R) is a faithful arithmetic certificate, whereas R alone has explicit collisions. For exterior states $\lambda = +1$ and $\det \mathsf{R} = -1$; the actual reconstructed S^6 candidate uses

$$(\ell_0, \ell_1, \ell_2) = (0, 1, -1), \qquad \lambda = -1, \qquad \det \mathsf{R} = +1.$$

No $SL_2(\mathbb{Z})$ row operation identifies these two oriented presentations, since it preserves the signed determinant.

Finally,

$$G(t, m, b) = \begin{pmatrix} 2t & 2m \\ 2m & b/3 \end{pmatrix}, \qquad (t, m, b) = \left(\frac{u}{2}, \frac{a}{2}, 3v\right)$$

gives $G = \Gamma$ and

$$\det G = \left(\frac{u+v}{2}\right)^2 - a^2 - \left(\frac{u-v}{2}\right)^2 = 0.$$

The inverse orthogonal Koide transform therefore produces a real Koide-null vector. It lies in the positive three-coordinate chamber only when, for $\theta_K = u/a = r/s$,

$$0 < \theta_K < \frac{\sqrt{3} - \sqrt{2}}{1 + \sqrt{2}} \quad \text{or} \quad \frac{\sqrt{3} + \sqrt{2}}{1 + \sqrt{2}} < \theta_K < 1 + \sqrt{2}.$$

Most selector states are not automatically positive mass points.

This coefficient correspondence lies on the discriminant of the ambient period-matrix space:

$$\det P_{\text{selector}} = tb - 6m^2 = 0,$$

whereas a regular torus fibre requires $\det P_T = -D_T \neq 0$, with $\det P_T \sim -T^2$ on the radial cusp. Thus the displayed arithmetic map cannot be a map to a regular fibre. Together with the signed terminal mismatch, this proves the exact current boundary: the arithmetic incidence and Koide/Lorentz identities are real, but they do not define a geometric morphism to the candidate threefold. The complete proofs and correction ledger are in `supporting_materials/workbench/research/` and the Erdős–Straus bridge audit supplied with the workbench.

Exact Koide-shell parametrization, hyperbolic realization, and the remaining analytic problem. The arithmetic count isolated by the reconstruction is

$$R(X) = \#\left\{(a, q, n) \in \mathbb{Z}^3 : \begin{array}{c} a^2 - 3q^2 - 18n^2 = -2, \quad a > 0,\ q, n \geq 0, \\ a + 4n > q, \quad 4a + 17n \leq X \end{array}\right\}.$$

It now has a multiplicity-one integral parametrization. Put

$$\mathcal{O}_K = \mathbb{Z}[\sqrt{3}], \qquad \epsilon = 2 + \sqrt{3}, \qquad m_n = 18n^2 - 2 \quad (n \geq 1).$$

The ideals of norm m_n are in bijection with

$$\mathcal{P}(m_n) = \{(C, D, E) : C, D > 0,\ C^2 D = m_n,\ 0 \leq E < D,\ E^2 \equiv 3 \pmod{D}\}$$

by

$$(C, D, E) \longmapsto C\big(D\mathbb{Z} + (E + \sqrt{3})\mathbb{Z}\big).$$

For every such triple there is one reduced generator

$$C(A + B\sqrt{3}), \qquad A^2 - 3B^2 = D, \quad A \equiv EB \pmod{D}, \quad A > 2B \geq 0,$$

and all nonnegative-q points in its orbit are obtained, exactly once, from

$$a_{n,C,D,E,k} + q_{n,C,D,E,k}\sqrt{3} = C(A + B\sqrt{3})\epsilon^k, \qquad k \geq 0.$$

The inverse is equally integral: form the HNF data of $(a + q\sqrt{3})$, then apply

$$(a, q) \longmapsto (2a - 3q,\ 2q - a)$$

while $a \le 2q$. The number of steps is k, and the terminal pair is the unique reduced generator. The omitted face is not hidden in this description: for $n = 0$ the solutions are precisely

$$a_k + q_k\sqrt{3} = (1+\sqrt{3})\epsilon^k, \qquad k \ge 1.$$

Writing

$$\rho(m) = \sum_{d \mid m} \left(\frac{12}{d}\right),$$

the number of unit orbits at fixed $n \ge 1$ is

$$\#\mathcal{P}(m_n) = \rho(m_n) = \rho(3n-1)\rho(3n+1).$$

The original sharp height is retained rather than replaced by a radial norm. For the reduced generator $\alpha_{n,\mathbf{p}}$, set

$$\tau_{n,\mathbf{p}} = \frac{\log(\alpha_{n,\mathbf{p}}/\sqrt{m_n})}{\log \epsilon} \in [0,1), \qquad Y_n(X) = \frac{X - 17n}{4},$$

$$L_n(X) = \frac{1}{\log \epsilon} \log\left(\frac{Y_n(X) + \sqrt{Y_n(X)^2 - m_n}}{\sqrt{m_n}}\right), \quad \mathcal{N}(X) = \{n \ge 1 : Y_n(X) \ge \sqrt{m_n}\}.$$

Then the endpoint-exact formula is

$$R(X) = R_0(X) + \sum_{n \in \mathcal{N}(X)} \sum_{\mathbf{p} \in \mathcal{P}(m_n)} \big(\lfloor L_n(X) - \tau_{n,\mathbf{p}} \rfloor + 1\big),$$

where $R_0(X)$ is the preceding $n = 0$ Pell count. Thus the strict chamber wall and every equality $4a + 17n = X$ remain visible. Fourier expansion of the phases gives unitary Hecke coefficients

$$S_\ell(m) = \sum_{N\mathfrak{a} = m} \xi_\ell(\mathfrak{a}), \qquad S_\ell(18n^2 - 2) = (-1)^\ell S_\ell(3n-1)S_\ell(3n+1),$$

so the unproved remainder is an explicit sharply truncated shifted correlation, rather than an unspecified equidistribution assumption.

There is also an exact quaternionic realization. In

$$B = (18,-6)_{\mathbb{Q}} = \mathbb{Q}\langle I, J\rangle/(I^2 - 18,\ J^2 + 6,\ IJ + JI),$$

the lattice

$$\mathcal{O} = \mathbb{Z}1 \oplus \mathbb{Z}\frac{I+J}{2} \oplus \mathbb{Z}\frac{I-J}{2} \oplus \mathbb{Z}\frac{IJ}{2}$$

is an order of reduced discriminant 108, of index 18 in a maximal overorder. The algebra has discriminant 6, ramifies exactly at 2 and 3, and splits over $\mathbb{R}$. The map

$$(a,q,n) \longmapsto \gamma = 2 + \frac{qI + aJ + nIJ}{2}$$

is a bijection from the parity-restricted ternary shell to the locus

$$\{\gamma \in \mathcal{O} : \mathrm{trd}(\gamma) = 4,\ \mathrm{nrd}(\gamma) = 1\}.$$

Every image is primitive hyperbolic, has eigenvalues $2 \pm \sqrt{3}$, and has translation length $2\log(2+\sqrt{3})$. Its centralizer order is

$$\mathcal{O} \cap \mathbb{Q}(\gamma) = \mathbb{Z}[\gamma - 2] \cong \mathbb{Z}[\sqrt{3}].$$

The full shell is a finite union of conjugacy classes under $\Gamma = \mathcal{O}^1$, and $R(X)$ is exactly the sum over their left cosets by the centralizers subject to the same four chamber inequalities and the same height $4a + 17n \le X$. No factor two is introduced on passing to $PSL_2(\mathbb{R})$.

This realization explains both what is now proved and what is still open. The standard radial Huber parameter is

$$X_i(\gamma)^2 = 1 + \frac{(7a + 15\sqrt{2}\,n)^2}{48},$$

not the Koide height $4a + 17n$. The two heights are comparable but their sharp regions have a main-order symmetric difference, so a radial remainder does not transfer to the original count. Benoist–Oh's effective theorem does apply directly to the literal five-wall sector and proves, for some $\eta > 0$,

$$R(X) = C_0 X + O(X^{1-\eta}), \qquad C_0 = \frac{9}{2\pi^2}\log(2+\sqrt{3})\log\left(\frac{3}{2\sqrt{2}}\right).$$

The strongest audited nearby pointwise sharp radial theorem has exponent $2/3$; the smooth radial spectral estimate has exponent $39/64$, but neither theorem has the target sector and height. Therefore

$$R(X) - C_0 X = O_\varepsilon(X^{1/2+\varepsilon})$$

remains **OPEN**, not a polite substitute for a theorem: its exact missing object is the displayed sharp shifted Hecke correlation, including all endpoint and exceptional-spectrum terms.

The exact census independently verifies every point through $X = 10^8$:

$$R(10^8) = 3{,}536{,}002.$$

For integer $10^4 \le X \le 10^8$, the largest observed value of $|R(X) - C_0 X|/\sqrt{X}$ is $0.1714001928088145\ldots$, attained at $X = 867{,}240$; this is finite evidence for the open estimate, not its proof. These arithmetic and hyperbolic results do not by themselves identify the counted triples with physical lepton masses or a quantum-field-theoretic spectrum. They instead give the exact morphisms and analytic target that any such identification must respect.