Editorial preface

This edition reconstructs the English text against the reviewed French source, including all printed folios 1–700 and the original front and rear matter. The Introduction, the authors' and editors' notes, and the original additions are retained in translation. Original source folios remain marked throughout; the comments for the certified blank leaves are preserved without printing an empty-leaf label beside subsequent text.

The three reader forms come from this single annotated master. The diplomatic form follows the printed readings wherever they affect the English. The corrected form gives the adjudicated readings and their source-linked apparatus. French spelling or agreement corrections that leave the English unchanged are recorded in the transfer ledger and create no English variant.

Some source assertions remain as printed because the examination did not establish a unique textual emendation. In particular, the accompanying dossier gives explicit counterexamples for the passages on source folios 431, 443 and 590–592 while retaining the authors' wording in translation. The printed blank locators on folio 464 and the integral membership assertions on folio 465 are likewise retained, with the exact computations and their limits recorded. These unchanged passages have no variant entries. The decision records also identify the other retained mathematical issues and give their source evidence.

About this edition

Théorie des intersections et théorème de Riemann–Roch, Séminaire de Géométrie Algébrique du Bois-Marie 1966–67 (SGA 6), directed by P. Berthelot, A. Grothendieck and L. Illusie, with the collaboration of D. Ferrand, J.-P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud and J.-P. Serre. Lecture Notes in Mathematics 225, Springer, 1971.

The French layers are AI-transcribed. The English is AI-translated. Codex's layers and corrections of August 2026, Claude Opus 5.5's check at effort max of every numbered page on 10–13 November 2026, and the corrections and assembly in this build are AI-produced corrections. This build is credited to GPT-6 Astra (OpenAI, ChatGPT Pro mode). The accompanying decision records state the actual verification status of this build.

The obscured part of the original publisher notice has been supplied from the matching inside-cover notice in Springer's *Lecture Notes in Mathematics* 228 (1971). This is an attributed restitution from that contemporary template. The hidden physical strokes in the SGA 6 scan were not recovered.

Sources and authorship retained from the supplied records

The original SGA volumes retain their authors, seminar contributors and editors, named in the corresponding texts. The collection includes work of Alexander Grothendieck, Michèle Raynaud, Michel Raynaud, Pierre Deligne, Nicholas Katz, Michel Demazure, Michael Artin, Jean-Louis Verdier, Luc Illusie and their collaborators; this list does not replace the detailed credits in each volume.

SGA 1 retains the French retypesetting associated with Bas Edixhoven's volunteer project, the Société mathématique de France and Michel Raynaud's notes, from arXiv math/0206203v2.

SGA 2 is AI-translated from the corrected French TeX of the re-edition edited by Yves Laszlo, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, Documents Mathématiques 4, Société mathématique de France, 2005 (arXiv math/0511279v1),

and follows its numbering and labels. Most of the initial French typesetting was done by volunteers of Bas Edixhoven's project (L. Bayle, N. Borne, O. Brinon, J. Buresi, M. Chardin, F. Ducrot, P. Graftiaux, F. Han, P. Karwasz, L. Koelblen, D. Madore, S. Morel, D. Naie, B. Osserman, J. Riou and V. Sécherre); C. Sabbah adapted it to the SMF format.

The SGA 2 and SGA 3 English sources acknowledge Jacob C. Reinhold's English translation, snapshot `e7a259f3f8608ad3edf9bf6eead3fd504dd2d23e`, used for comparison and drafting. His stated CC BY 4.0 notice applies to that contribution and is preserved; it does not license the underlying French works or every contribution here. Inherited English wording remains attributed to its human source. In SGA 3, lemmas, proofs and corrections added in the Philippe Gille–Patrick Polo re-edition are given in Translator's notes and Additions that cite it.

SGA 4 is AI-translated from the French LaTeX edition of SGA 4, Tomes 1–3, distributed by Fabrice Orgogozo (source snapshot at git commit `71766d9`), and uses its TeX configuration files. Its footnotes headed "N.D.E." are that edition's editorial notes, translated.

SGA 4½ preserves Daniel Miller's LaTeX typesetting and its original typesetter's note. It is not attributed to AI. His repository (`https://github.com/dkmiller/sga4.5`) is no longer online; the French text is in sga-fr, and a copy survives at NomiL/sga4.5.

Additional English translation is AI-translated; subsequent corrections, editing, reconstruction and typesetting are AI-produced. The publisher did not translate the text. The source-specific human credits and genuine third-party notices remain applicable; attribution does not establish permission or endorsement.

Rights

The rights in the original work belong to its author or the author's representatives or heirs. Where we hold rights, in the AI-produced transcription, translation, corrections and typesetting, they are dedicated to the public domain under CC0 1.0. This edition is published for educational, non-profit purposes, and we claim no rights we do not hold. If you hold rights in this work and object to its publication here, please open a GitHub issue and we will take it down.

[print cover]

Lecture Notes in Mathematics

A collection of informal reports and seminars

Edited by A. Dold, Heidelberg and B. Eckmann, Zürich

225

Algebraic Geometry Seminar of Bois Marie 1966/67

SGA 6

Directed by

P. Berthelot, A. Grothendieck, L. Illusie

With the collaboration of

D. Ferrand, J.P. Jouanolou, O. Jussila,

S. Kleiman, M. Raynaud, J.P. Serre

Intersection Theory and the Riemann–Roch Theorem

Springer-Verlag

Berlin · Heidelberg · New York

[print cover-verso]

This series aims to report new developments in mathematical research and teaching – quickly, informally and at a high level. The type of material considered for publication includes:

1. Preliminary drafts of original papers and monographs
2. Lectures on a new field, or presenting a new angle on a classical field
3. Seminar work-outs
4. Reports of meetings

Texts which are out of print but still in demand may also be considered if they fall within these categories.

The timeliness of a manuscript is more important than its form, which may be unfinished or tentative. Thus, in some instances, proofs may be merely outlined and results presented which have been or will later be published elsewhere.

Manuscripts should comprise not less than 100 pages.

Publication of *Lecture Notes* is intended as a service to the international mathematical community, in that a commercial publisher, Springer-Verlag, can offer a wider distribution to documents which would otherwise have a restricted readership. Once published and copyrighted, they can be documented in the scientific literature. –

Manuscripts

Manuscripts are reproduced by a photographic process; they must therefore be typed with extreme care.

Symbols not on the typewriter should be inserted by hand in indelible black ink. Corrections to the typescript should be made by sticking the amended text over the old one, or by obliterating errors with white correcting fluid. Authors receive 75 free copies.

The typescript is reduced slightly in size during reproduction; best results will not be obtained unless the text on any one page is kept within the overall limit of 18×26.5 cm ($7 \times 10\frac{1}{2}$ inches). The publishers will be pleased to supply on request special stationery with the typing area outlined.

Manuscripts in English, German or French should be sent to Prof. Dr. A. Dold, Mathematisches Institut der Universität Heidelberg, 69 Heidelberg/Germany, Tiergartenstraße or Prof. Dr. B. Eckmann, Eidgenössische Technische Hochschule, CH-8006 Zürich/Switzerland.

Lecture Notes in Physics

Bisher erschienen/Already published

Vol. 1: J. C. Erdmann, Wärmeleitung in Kristallen, theoretische Grundlagen und fortgeschrittene experimentelle Methoden. II, 283 Seiten. 1969. DM 20,–

Vol. 2: K. Hepp, Théorie de la renormalisation. III, 215 pages. 1969. DM 18,–

Vol. 3: A. Martin, Scattering Theory: Unitarity, Analytic and Crossing. IV, 125 pages. 1969. DM 14,–

Vol. 4: G. Ludwig, Deutung des Begriffs physikalische Theorie und axiomatische Grundlegung der Hilbertraumstruktur der Quantenmechanik durch Hauptsätze des Messens. XI, 469 Seiten. 1970. DM 28,–

Vol. 5: M. Schaaf, The Reduction of the Product of Two Irreducible Unitary Representations of the Proper Orthochronous Quantummechanical Poincaré Group. IV, 120 pages. 1970. DM 14,–

Vol. 6: Group Representations in Mathematics and Physics. Edited by V. Bargmann. V, 340 pages. 1970. DM 24,–

Vol. 7: R. Balescu, J. L. Lebowitz, I. Prigogine, P. Résibois, Z. W. Salsburg, Lectures in Statistical Physics. V, 181 pages. 1971. DM 18,–

Vol. 8: Proceedings of the Second International Conference on Numerical Methods in Fluid Dynamics. Edited by M. Holt. IX, 462 pages. 1971. DM 28,–

[print I]

Lecture Notes in Mathematics

A collection of informal reports and seminars

Edited by A. Dold, Heidelberg and B. Eckmann, Zürich

225

Intersection Theory and the Riemann–Roch Theorem

Springer-Verlag

Berlin · Heidelberg · New York 1971

[print II]AMS Subject Classifications (1970): 14-02, 14C15, 14F15, 14K30, 18E30

ISBN 3-540-05647-5 Springer-Verlag Berlin · Heidelberg · New York

ISBN 0-387-05647-5 Springer-Verlag New York · Heidelberg · Berlin

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically those of translation, reprinting, re-use of illustrations, broadcasting, reproduction by photocopying machine or similar means, and storage in data banks.

Under § 54 of the German Copyright Law where copies are made for other than private use, a fee is payable to the publisher, the amount of the fee to be determined by agreement with the publisher.

© by Springer-Verlag Berlin · Heidelberg 1971. Library of Congress Catalog Card Number 73-178287. Printed in Germany.

Offsetdruck: Julius Beltz, Hemsbach/Bergstr.

[print III]

Algebraic Geometry Seminar of Bois Marie

1966–67

Intersection Theory and the Riemann–Roch Theorem

(SGA 6)

a Seminar directed by

P. Berthelot, A. Grothendieck, L. Illusie

with the collaboration of

D. Ferrand, J.-P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud, J.-P. Serre

[print V]

Preface

The present edition of the Algebraic Geometry Seminar 6 reproduces, with very few changes, the first edition, distributed by the I.H.E.S. as mimeographed exposés. The corrections made concern essentially printing errors, except in Exposés I and III, where L. Illusie made slight modifications to some statements that were incorrect in the original version. In addition, footnotes have been added, in particular in Exposé XIV by A. Grothendieck, to indicate recent progress on questions that were still open at the time of the Seminar.

Finally, we point out that Exposé XI, by D. Ferrand, concerning the theory of the determinant for perfect complexes, has not been written up and will therefore not be published; to avoid errors in references, we have kept the numbering of the subsequent exposés unchanged.

June 1971

[print VII] # Introduction

The aim of the present Seminar is to develop a global intersection theory, and a Riemann–Roch formula, for arbitrary schemes. The reader will find a description of the Seminar's program in Exposé 0. The possibility, in principle, of proving a Riemann–Roch formula for general regular schemes, along the lines of the well-known report of Borel–Serre in 1958, was clear from that time, at least for a projective morphism. The extension to the general case is less evident; the program of which it forms a part (and which is partially carried out in our seminar) dates from 1960. As was also the case for the duality theory of coherent sheaves (cf. R. Hartshorne, Residues and Duality, Lecture Notes in Mathematics n° 20, Springer), an essential tool for formulating a satisfactory theory is Verdier's theory of derived categories, knowledge of which is indispensable for understanding the Seminar.

Apart from this theory, studying the Seminar requires scarcely more than a general knowledge of the foundations of Algebraic Geometry, as set out in EGA, Chapters I, II, III; in addition, we shall have only occasional references to make to EGA IV, for certain facts concerning flat or smooth morphisms, which for the most part also appear in the first exposés of SGA 1. Finally, to develop certain results with all the generality desirable for applications, we sometimes use the notions of a ringed site and a ringed topos, for which we refer to SGA 4, Exposés I to IV. A reader unfamiliar with the language of sites and topoi may everywhere replace these objects by ordinary topological spaces, the objects of the topos then being replaced by open subsets of these spaces; nevertheless, we advise the reader, preferably, to assimilate the language of topoi, which provides an extremely convenient principle of unification.

The fundamental notion for the theory presented here is that of a perfect complex of Modules, and its various variants, developed in Exposés I, II, III. It seems clear that these notions, which are also important in other contexts in Algebraic Geometry (notably for formulas of Lefschetz–Verdier type in cohomology with discrete (SGA 5) or coherent coefficients), will also have their part to play outside Algebraic Geometry, notably for the formulation of a complex[print VIII]analytic variant of the Riemann–Roch theorem presented here, or suitable variants of the Atiyah–Singer index theorem. Some indications in this direction will be given in Exp. II. Unfortunately, at the present time there is still no statement, even a heuristic one, that would encompass these two theorems (the first of which, for the moment, remains conjectural). A new idea is apparently lacking, as one is also lacking in Algebraic Geometry to obtain a proof of the Riemann–Roch theorem beyond projectivity hypotheses (cf. Exp. XIV, n° 2).

We shall make no use in this seminar of the local theory of intersections, set out in J.P. Serre, Algèbre Locale. Multiplicités (Lecture Notes in Mathematics[1] n° 11, Springer), and shall simply content ourselves with pointing out here that this course of Serre had an evident influence on the genesis of the theory in 1957. Nor shall we use the theory of the Chow ring developed in the Chevalley Seminar 1958, Exposés 2 and 3 (École Normale Supérieure). This theory is essentially linked to nonsingularity hypotheses, whereas the aim of our seminar, on the contrary, is to develop an intersection theory on general schemes (and even on general locally ringed topoi) see on this subject the comments in Exp. XIV n° 4 and n° 8, giving the relations between our theory and that of the Chow ring, and raising the question of a generalization of the latter. We may say that the Seminar presents a coherent and self-contained intersection theory, but by no means exhausts the different points of view that are useful (or indeed indispensable) in intersection theory, and that it should consequently be supplemented by the cited exposés of Serre and Chevalley.

We have attached to the seminar (as an Appendix to Exp. 0) Grothendieck's 1957 report "classes de Faisceaux et théorème de Riemann–Roch", which served as the basis for the seminar of Borel and Serre at Princeton in the same year, as well as for their already cited article. This report

outlines two proofs of the Riemann–Roch theorem for nonsingular quasi-projective varieties; the first, valid at present only in characteristic zero, but giving in return a slightly more precise result in the case of an immersion, does not yet appear in the literature (including the work of Borel–Serre and the present seminar). Reading this report of course presupposes none of the other exposés of the seminar, and may even serve as an introduction to its study on the same footing as Exposé 0, especially for a reader not yet familiar with Borel–Serre. Moreover, the proof to which we have just alluded applies essentially as it stands to the case [print IX]of a projective morphism of complex analytic spaces, and will undoubtedly also apply in arbitrary characteristic once the problem of the "exterior power" operations in the derived category has been solved (cf. Exp. XIV, n°s 1, 2, 3). The absence of a study of this question undoubtedly constitutes the most serious gap in this Seminar, even from the point of view we have adopted.

One will notice the absence, from the table of contents of the present Seminar, of two of the participants mentioned on the cover page. One of them, J.P. Jouanolou, had taken an active part in the technical development of the first part of the seminar, but had been prevented from taking part in the oral exposés; to him we owe, in particular, the linear independence assertion contained in the important statement VI 1.1. giving the structure of the ring $K^\bullet$ of classes of locally free Modules on a projective bundle (which had previously been proved only under the assumption that the base scheme admits an ample invertible Module). The other, J.P. Serre, had given two oral exposés, logically independent of the rest of the Seminar, which he therefore preferred to publish as a separate article (cf. J.P. Serre, Groupes de Grothendieck des schémas en groupes réductifs déployés, to appear in Publications Mathématiques, n° 34).

As in the other parts of the Algebraic Geometry Seminar of Bois Marie, the sigla SGA 1 to SGA 6 refer to the different parts of that Seminar, the Roman numeral following the siglum indicating the number of the exposé, and the Arabic numerals that follow it corresponding to the internal numbering of the exposé in question; for internal references within the present Seminar, the siglum SGA 6 is omitted, using references of the form I 4.9 (meaning: Exposé I, statement 4.9). The siglum EGA refers to the Éléments de Géométrie Algébrique of J. Dieudonné and A. Grothendieck.

[print XI]

Table of contents

EXPOSÉ 0
Outline of a Program for an Intersection Theory on General Schemes
by A. Grothendieck 1

Classes of Sheaves and the Riemann–Roch Theorem
by A. Grothendieck 20

EXPOSÉ I
Generalities on Finiteness Conditions in Derived Categories
by L. Illusie 78

EXPOSÉ II
Existence of Global Resolutions
by L. Illusie 160

EXPOSÉ III
Relative Finiteness Conditions
by L. Illusie 222

EXPOSÉ IV
Grothendieck Groups of Ringed Topoi
by L. Illusie 274

EXPOSÉ V
Generalities on λ-Rings
by P. Berthelot 297

EXPOSÉ VI
The $K^\bullet$ of a Projective Bundle: Calculations and Consequences
by P. Berthelot 365

[print XII]EXPOSÉ VII
Regular Immersions and Calculation of the $K^\bullet$ of a Blown-up Scheme
by P. Berthelot 416

EXPOSÉ VIII
The Riemann–Roch Theorem
by P. Berthelot 466

EXPOSÉ IX
Some Calculations of K Groups.
by P. Berthelot 498

EXPOSÉ X
Intersection Formalism on Proper Algebraic Schemes
by O. Jussila
With an Appendix by A. Grothendieck
Specialization in Intersection Theory 519

EXPOSÉ XI – Not written up

EXPOSÉ XII
A Relative Representability Theorem for the Picard Functor
by M. Raynaud (written up by S. Kleiman) 595

EXPOSÉ XIII
Finiteness Theorems for the Picard Functor
by S. Kleiman 616

EXPOSÉ XIV
Open Problems in Intersection Theory
by A. Grothendieck 667

Terminological Index 691

Index of Notations 696

[print 1]

Exposé 0

OUTLINE OF A PROGRAMME FOR A THEORY OF INTERSECTIONS ON GENERAL SCHEMES

by A. Grothendieck

The present exposé is introductory in nature, and reading it is not logically indispensable for the study of the Seminar. It is addressed more particularly to readers familiar with the provisional version of the Riemann–Roch theorem, as it is presented in the report of Borel–Serre [2] or in the report of Grothendieck already mentioned in the Introduction (subsequently cited as [RRR]), which is reproduced as an appendix at the end of the present exposé.

1.

Recall the Riemann–Roch formula for a proper morphism

$$f : X \longrightarrow Y^{[2]}$$

of quasi-projective smooth schemes over a field k, and a coherent sheaf F on X, whose class in the group $K(X)$ of classes of coherent sheaves on X is denoted by $\mathrm{cl}(F)$:

$$\mathrm{Todd}(T_Y)\ \mathrm{ch}_Y\big(f_*(\mathrm{cl}(F))\big) = f_*\big(\mathrm{Todd}(T_X)\ \mathrm{ch}_X(F)\big). \tag{1.1}$$

This formula holds in $A(Y) \otimes_{\mathbb{Z}} \mathbb{Q}$, where $A(Y)$ is the Chow ring of Y, the f_* on the right-hand side being obtained by tensoring with $\mathbb{Q}$ the homomorphism "direct image of cycles"

$$f_* : A(X) \longrightarrow A(Y),$$

and that on the left-hand side being the Euler–Poincaré characteristic of F relative to f:

$$f_*(\mathrm{cl}(F)) = \sum_i (-1)^i\, \mathrm{cl}\big(R^i f_*(F)\big)\ ;$$

ch_X, ch_Y denote the Chern characters on X, respectively on Y, and T_X, T_Y the tangent bundles of X, respectively of Y. As is known, $\mathrm{Todd}(-)$ and $\mathrm{ch}(-)$ are universal polynomials with coefficients in $\mathbb{Q}$ in the Chern classes of the argument. Since the constant term of $\mathrm{Todd}(-)$ is 1, it is an invertible element for every value of the argument, so that, after multiplication by $\mathrm{Todd}(T_Y)^{-1}$, formula (1.1) takes the equivalent form, more convenient for our purpose,

$$\mathrm{ch}_Y\big(f_*(\mathrm{cl}(F))\big) = f_*\big(\mathrm{Todd}(T_f) \cdot \mathrm{ch}_X(F)\big),^{[3]} \tag{1.2}$$

where we have put

$$T_f = T_X - f^*(T_Y) \in K(X), \tag{1.3}$$

[print 2] so that T_f plays the role of a virtual relative tangent bundle of X over Y. When the morphism f is smooth (i.e. its tangent map is everywhere surjective), we simply have

$$T_f = T_{X/Y} \qquad \text{(tangent bundle along the fibres)},$$

whereas when $f : X \to Y$ is an immersion, we find

$$T_f = -\check{N}_{X/Y},$$

where $\check{N}_{X/Y}$ denotes the normal sheaf of X in Y.

One of the main aims of the Seminar is to generalize (1.2) simultaneously in two directions:

a) To remove the hypothesis of the existence of a base field k.

b) To replace the regularity hypotheses on Y, X by a hypothesis of "local regularity" of f.

Finally, along the way, we shall also consider the problem:

c) To eliminate the quasi-projectivity hypotheses which, in the absence of a base field, are expressed by the existence of ample invertible Modules on X and on Y.

2.

Let us first examine generalization a), while retaining the hypotheses of regularity and of the existence of ample invertible Modules on X and on Y.

The definition of $K(X)$, $K(Y)$ and of the homomorphism $f_* : K(X) \to K(Y)$ then presents no new problem, because X and Y are regular. The most natural way of giving a meaning to (1.2) therefore seems to consist in defining Chow rings $A(X), A(Y)$ and a group homomorphism

$$f_* : A(X) \longrightarrow A(Y),$$

in establishing a theory of Chern classes, providing maps

$$c_i : K(X) \longrightarrow A(X),$$

and similarly for Y, and finally in describing a virtual relative tangent bundle element

$$T_f \in K(X).$$

[print 3]

2.1

As for the latter, a definition presents itself fairly evidently. We use the fact that, by the hypothesis of the existence of an ample invertible Module on X and on Y, the morphism f can be factored as

$$X \xrightarrow{i} X' \xrightarrow{f'} Y, \tag{2.1}$$

where i is a closed immersion and f' is a smooth morphism; for example, we may take $X' = \mathbb{P}^r_Y$ for suitable r. We then put

$$\check{T}_f = \mathrm{cl}(\underline{\Omega}^1_{X'/Y}) - \mathrm{cl}(\underline{N}_{X/X'}), \tag{2.2}$$

where $\underline{\Omega}^1_{X'/Y}$ is the locally free Module of relative 1-differentials of X' over Y (or relative cotangent Module of X' over Y), and $\underline{N}_{X/X'}$ is the conormal Module of X in X', equal by definition to $\underline{J}/\underline{J}^2$, where $\underline{J}$ is the Ideal on X' defining the closed subscheme X; since X and X' are regular, we also conclude that $\underline{N}_{X/X'}$ is locally free. Finally, $\check{T}_f$ denotes the dual of T_f. We readily verify (Exp. VIII) that the element T_f defined by (2.1) does not depend on the chosen factorization of f.

2.2.

As for a definition, under the conditions considered, of a Chow ring and a corresponding theory of Chern classes, although it is plausible that such a theory should be developable on the model of

the theory already known for smooth quasi-projective algebraic schemes, it seems that this work has not been done at present, and a fortiori it has not been undertaken in the Seminar. Instead, we introduce another ring which will play a role analogous to that of the Chow ring, and whose tensor product with Q is indeed isomorphic to $A(X)\otimes_{\mathbb{Z}}\mathbb{Q}$ when X is quasi-projective and smooth over a field k. This ring is the associated graded ring of $K(X)$ for a suitable filtration of that ring. A first natural filtration of $K(X)$, the one considered in [RRR], is the "topological filtration" by the codimension of the support of coherent sheaves, $\mathrm{Filt}^i_{\mathrm{top}}\,K(X)$ being generated by the $\mathrm{cl}(F)$ with F coherent and $\mathrm{codim}(\mathrm{supp}\,F, X)\geq i$. A first question is the compatibility of this filtration with the ring structure, a question resolved in [RRR], when X is smooth over a field k, by means of Chow's "moving lemma". Granting this point, we therefore obtain an associated graded ring, which we shall denote here by $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$. In [RRR] a surjective ring homomorphism is defined:

$$\phi : A(X) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(X),$$

[print 4] (sending the class of a prime cycle Z to $\mathrm{cl}(\underline{O}_Z)$,) whose kernel subsequently turns out (Exp. XIV 4.2) to be a torsion subgroup. Moreover, [RRR] shows how the images in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ of the Chern classes $c_i(F)$ can be calculated directly in terms of $\mathrm{cl}(F)\in K(X)$ and the exterior-power operations λ^i in $K(X)$.

This shows that, for X a noetherian regular scheme equipped with an ample invertible Module, the ring $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ is a convenient substitute for the hypothetical Chow ring $A(X)$; it also has the advantage over the latter that certain calculations in [RRR] take place directly in this ring and lead to important formulas in it, formulas which, even for X projective and smooth over a field, have not at present been established in $A(X)$ itself (Exp. XIV, 4.3 and 4.4). To justify this point of view completely, however, one would have to resolve the problem mentioned above of the compatibility of the topological filtration of $K(X)$ with its ring structure, a problem which seems essentially connected with Chow's "moving lemma", which is itself the essential technical ingredient of the theory of the Chow ring that we have claimed to be able to avoid.

2.3.

To avoid this problem, we give $K(X)$ a second filtration, finer than the topological filtration, which can be described in purely algebraic terms from the augmented λ-ring structure of $K(X)$. The definition of this filtration is suggested by the expression given in [RRR] for the Chern classes of an $x\in K(X)$ in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$,

$$c_i(x) = \gamma^i(x-\epsilon(x)) \mod \mathrm{Filt}^{i+1}_{\mathrm{top}}\,K(X), \tag{2.3}$$

where γ^i denotes a certain linear combination of the λ^j $(j\leq i)$[4] made explicit in loc. cit., ϵ being the augmentation. We show that $\gamma^i(x-\epsilon(x))$ does have topological filtration $\geq i$, which gives a meaning to formula (2.3). We then define the λ-filtration of $K(X)$ by the ideals $\mathrm{Filt}^i\,K(X)$, where $\mathrm{Filt}^i\,K(X)$ consists of sums of expressions of the form

$$\gamma^{i_1}(x_1-\epsilon(x_1))\cdots\gamma^{i_k}(x_k-\epsilon(x_k)), \qquad i_1+\cdots+i_k\geq i.$$

The associated graded ring of this filtration will be denoted by $\mathrm{Gr}^\bullet(X)$, and it is this ring that will replace the Chow ring of X in the Seminar. We can moreover show that, in the case considered here, the canonical homomorphism

$$\mathrm{Gr}^\bullet(X) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(X) \tag{2.4}$$

[print 5] is an isomorphism modulo torsion, which justifies the point of view that this ring tensored with $\mathbb{Q}$ can play in every respect the same role as the Chow ring tensored with $\mathbb{Q}$. On the other

hand, we have done exactly what is needed for formula (2.3) (with $\mathrm{Filt}^i_{\mathrm{top}}$ replaced by Filt^i) to define a theory of Chern classes with values in $\mathrm{Gr}^\bullet(X)$, from which the Chern classes in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ are deduced using the homomorphism (2.4). We show (Exp. V), using the fact (Exp. VI) that $K(X)$ is a special λ-ring in the sense of [RRR] (i.e. a λ-ring in the sense of Exp. V), that this notion of Chern class satisfies all the usual formal properties reviewed in [RRR].

2.4.

We note a very appreciable advantage of the definition of $\mathrm{Gr}^\bullet(X)$ over that of $A(X)$ and $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$: it remains meaningful for an absolutely arbitrary scheme X, and more generally for an arbitrary locally ringed topos X, taking for $K(X)$ the group of classes constructed from locally free Modules on X, which is still an augmented λ-ring, hence equipped with a corresponding filtration allowing an associated graded $\mathrm{Gr}^\bullet(X)$ to be defined. As the price of this generality, however, we must observe that, when we are unwilling to disregard torsion phenomena, i.e. to tensor with $\mathbb{Q}$, the properties of $\mathrm{Gr}^\bullet(X)$ are less satisfactory in various respects than those of its two competitors (Exp. XIV no. 4). Thus, supposing again that $f : X \to Y$ is a proper morphism of noetherian regular schemes equipped with ample invertible Modules, it is not true in general that the homomorphism $f_* : K(X) \longrightarrow K(Y)$ maps $\mathrm{Filt}^i K(X)$ into $\mathrm{Filt}^{i-d} K(Y)$, where d is the relative dimension of X over Y, and hence defines, by passage to the associated graded rings, a homomorphism of degree $-d$ from $\mathrm{Gr}^\bullet(X)$ to $\mathrm{Gr}^\bullet(Y)$. However, we can prove (Exp. VII) that this holds after tensoring with $\mathbb{Q}$, so that we obtain a homomorphism of degree $-d$

$$f_* : \mathrm{Gr}^\bullet(X)_{\mathbb{Q}} \longrightarrow \mathrm{Gr}^\bullet(Y)_{\mathbb{Q}}, \tag{2.5}$$

playing the role of the homomorphism "direct image of cycles" $A(X) \to A(Y)$.

With this understood, we see that we have all the structural elements needed to give a meaning to formula (1.2) without postulating the existence of a base field, $f : X \to Y$ being a proper morphism of noetherian regular schemes admitting ample invertible Modules.

[print 6]

3.

Let us now show how we can still give a meaning to formula (1.2) when we abandon the regularity hypothesis on X, Y, replacing it by a hypothesis of a local nature on the morphism f, which we shall specify.

3.1

A first difficulty arises because the observation that served as the starting point of the theory [RRR], namely that the group $K(X)$ of classes of sheaves on X is the same whether it is defined in terms of coherent sheaves or in terms of locally free sheaves, is essentially tied to the regularity hypothesis on X. When X is a general scheme (say noetherian, for simplicity), we must distinguish these two groups, which we shall denote respectively by $K_\bullet(X)$ and $K^\bullet(X)$, the first being covariant in X for proper morphisms, and the second contravariant in X for arbitrary morphisms. We note that $K^\bullet(X)$ has a natural ring structure (and even an augmented λ-ring structure), whereas $K_\bullet(X)$ in general no longer has a ring structure, but only a module structure over $K^\bullet(X)$. The homomorphism $x \to x\ \mathrm{cl}_\bullet(\underline{O}_X)$ (where $\mathrm{cl}_\bullet(\underline{F})$ denotes the class of $\underline{F}$ in $K_\bullet(X)$) from $K^\bullet(X)$ to $K_\bullet(X)$ is in general no longer an isomorphism. For our formulation of the generalized Riemann–Roch theorem, we shall work exclusively with $K^\bullet(X)$, $K^\bullet(Y)$, to the

exclusion of $K_\bullet(X)$, $K_\bullet(Y)$. As we noted in 2.3., these augmented λ-rings, equipped with their natural λ-filtration, give rise to graded rings $\mathrm{Gr}^\bullet(X)$, $\mathrm{Gr}^\bullet(Y)$, and "Chern class" homomorphisms

$$c_i : K^\bullet(X) \longrightarrow \mathrm{Gr}^\bullet(X),$$

and similarly for Y. The difficulty, however, arises in defining a direct-image homomorphism

$$f_* : K^\bullet(X) \longrightarrow K^\bullet(Y), \tag{3.1}$$


which can no longer be defined by the naive formula

$$f_*(\mathrm{cl}^\bullet(\underline{F})) = \sum_i (-1)^i \,\mathrm{cl}^\bullet(R^i f_*(\underline{F})),$$ [5]

because, even when $\underline{F}$ is locally free, the Modules $R^i f_*(\underline{F})$ are in general no longer locally free, nor even of finite tor-dimension, and the preceding formula loses its meaning. A solution to this difficulty is provided by the point of view of derived categories. Indeed, although the $R^i f_*(\underline{F})$ are in general "bad" sheaves, say of infinite tor-dimension, [print 7] to which we cannot associate a class in $K^\bullet(Y)$, the complex $Rf_*(\underline{F})$ on Y, the direct image of $\underline{F}$ in the sense of derived categories, whose cohomology sheaves are the $R^i f_*(\underline{F})$, nevertheless tends to be excellent. More precisely, suppose that $\underline{F}$ has finite tor-dimension over $\underline{O}_Y$, which is the case when $\underline{O}_X$ has finite tor-dimension over $\underline{O}_Y$ and, in addition, F is locally free. We then show (Exp. III) that the complex $Rf_*(F)$ is "perfect", by which we mean here: it has coherent cohomology sheaves (which we already knew in any case) and finite global tor-dimension; or, equivalently: on every affine open U of Y, this complex is isomorphic, in the derived category $D(U)$, to a complex bounded in degree and locally free of finite type in each degree. From the point of view of Homological Algebra, perfect complexes are "as good" as locally free Modules, of which they are the natural generalization. They moreover form a "triangulated subcategory" of the derived category $D(Y)$, i.e. a subcategory stable under cones (or "mapping cylinders") of morphisms. On the other hand, to every triangulated category C we associate a group $K(C)$ of classes, a natural variant of the construction of the group of classes of an abelian category, by taking the free group generated by the objects of C, modulo the relations

$$\mathrm{cl}(K) - \mathrm{cl}(K') - \mathrm{cl}(K'')$$

for every distinguished triangle



What replaces the fundamental observation of [RRR] in the present case is the fact that, when the scheme Y admits an ample invertible Module, every perfect complex $L_\bullet$ on Y is globally (and not only locally) isomorphic in $D(Y)$ to a complex $L'_\bullet$ bounded in degree and locally free of finite type in each degree (Exp. II); it follows readily (Exp. IV) that the canonical homomorphism

$$K^\bullet(Y) \longrightarrow K(\mathrm{Parf}(Y)),$$ [6]

where $\mathrm{Parf}(Y)$ is the triangulated category of perfect complexes on Y, is an isomorphism. Thus to every perfect complex $L_\bullet$ on Y we can associate a class

$$\mathrm{cl}^\bullet(L_\bullet) \in K^\bullet(Y),$$

[print 8] which is moreover simply

$$\mathrm{cl}^\bullet(L_\bullet)^{[7]} = \sum_i (-1)^i \,\mathrm{cl}^\bullet(L'_i).$$

where $L'_\bullet$ is as above, so that the L'_i are indeed locally free Modules on Y whose classes we can take in $K^\bullet(Y)$.

With this understood, we define a homomorphism (3.1) whenever $f : X \to Y$ is a proper morphism of finite tor-dimension of noetherian schemes such that Y admits an ample invertible Module, simply by putting

$$f_*(\mathrm{cl}^\bullet(\underline{F})) = \mathrm{cl}^\bullet(Rf_*(\underline{F})) \tag{3.2}$$

for every locally free Module $\underline{F}$ on X (so that $Rf_*(\underline{F})$ is indeed a perfect complex on Y, and the right-hand side is defined). When X also admits an ample invertible Module, $K^\bullet(X)$ can likewise be interpreted as $K(\mathrm{Parf}(X))$, and the homomorphism (3.1) is precisely the one associated, by functoriality, with the exact functor of triangulated categories induced by

$$Rf_* : \mathrm{Parf}(X) \longrightarrow \mathrm{Parf}(Y).$$

3.2.

To give a meaning to (1.2), we must still impose on f a hypothesis allowing a meaning to be given to the homomorphism (2.5)

$$f_* : \mathrm{Gr}^\bullet(X)_\mathbb{Q} \longrightarrow \mathrm{Gr}^\bullet(Y)_\mathbb{Q}, \tag{3.3}$$

i.e. ensuring that the homomorphism (3.1) just defined is (up to a shift and modulo torsion) compatible with the filtrations of these rings; and finally we must still define a "virtual relative tangent bundle" element

$$T_f \in K^\bullet(X).$$

For the latter, returning to the definition proposed in 2.1, we see that we should make a hypothesis ensuring that (with the notation of loc. cit.) the conormal Module $\underline{N}_{X/X'}$ is locally free. The most natural hypothesis for this purpose is that i is a "regular immersion", i.e. identifies X with a subscheme of X' defined by an Ideal which locally can be defined by an $\underline{O}_{X'}$-regular sequence of sections of $\underline{O}_{X'}$. This hypothesis, of a purely local nature, does not in fact depend on the choice of the factorization of f into an immersion followed by a smooth morphism (Exp. VIII), and, as above, the same holds for the class T_f defined by formula (2.2)[8]. When the hypothesis just considered holds, we say that the morphism f is a "locally complete intersection" morphism(*). Such a morphism is moreover locally of finite tor-dimension.

[print 9] For a locally complete intersection morphism f, we introduce a natural notion of "virtual relative dimension" $d(f)$ by the formula

$$d(f)(x) = \dim_x\big(f'^{-1}f(x)\big) - \mathrm{codim}_x(X, X').$$

This is an integer-valued function, locally constant in the point $x \in X$, hence constant when X is connected; it likewise does not depend on the choice of the factorization (2.1). When this virtual relative dimension d is constant, f is proper, and X and Y admit ample invertible Modules, we can show (Exp. VIII), although this is far from a trivial result, that

$$f_*(\mathrm{Filt}^i K^\bullet(X)_\mathbb{Q}) \subset \mathrm{Filt}^{i-d} K^\bullet(Y)_\mathbb{Q}, \tag{3.4}$$

hence a homomorphism (3.3) of degree $-d$.

(*) We shall also allow ourselves to omit the word "locally" in this terminology, which entails no risk of confusion.

3.3.

We have thus assembled all the structural elements needed to give a meaning to formula (1.2) for a proper locally complete intersection morphism of noetherian schemes X, Y both admitting ample invertible Modules. With a little additional care, this formulation moreover remains meaningful independently of any noetherian hypothesis.

In this general form, this Riemann–Roch formula will be proved in Exp. VIII. The reader will see that the proof given in this Seminar combines elements of the two proofs in [RRR] which were discussed in the Introduction. The need to use the methods of the first of these proofs, involving more λ-formalism than the second (followed in Borel–Serre), is due to the need for a proof of inclusion (3.4), a question that did not arise in the more restricted setting of [RRR] and Borel–Serre [2], where we could work with the Chow ring or the ring $\mathrm{Gr}^\bullet_{\mathrm{top}}$. From the second proof, we retain the introduction of the blow-up of Y along X when f is an immersion. This technical device will doubtless disappear, and the proof for an immersion will extend to the case where we no longer postulate the existence of an ample invertible Module on Y, once the question raised in Exp. XIV no. 1 concerning the introduction of exterior-power operations in the derived category $D^-(Y)$ has been resolved.

4.

Finally, we must give some indications of how a meaning can still be given to (1.2) without a hypothesis of the existence [print 10] of ample invertible Modules on X and Y.

4.1.

First, we note that for an arbitrary scheme X (or, more generally, locally ringed topos), it seems beyond doubt that the "right" invariant to replace the ring $K(X)$ of [RRR] is not the ring $K^\bullet(X)$ considered above, defined in terms of locally free Modules on X, but the group $K(\mathrm{Parf}(X))$, considered in 3.1., defined in terms of perfect complexes on X, a group that we shall denote here by $K^\bullet(X)$, and which we shall distinguish from the "naive" group of 3.1, denoted by $K^\bullet_{\mathrm{naive}}(X)$. This point of view is forced by the need to define a direct-image homomorphism (3.1) for a proper morphism $f : X \to Y$ of finite tor-dimension between, say, noetherian schemes (the non-noetherian case requires somewhat stronger local finiteness properties of f, cf. Exp. III). With the definition of $K^\bullet(X)$ now adopted, this homomorphism f_* is indeed defined trivially by formula (3.2), where $\underline{F}$ is now an arbitrary perfect complex on X, which implies that $Rf_*(\underline{F})$ is also perfect.

4.2

This new definition of $K^\bullet(X)$ nevertheless raises a new problem of homological algebra: that of defining a λ-structure on $K^\bullet(X)$ compatible with its natural ring structure arising from the tensor product in the derived category. (NB When $L_\bullet$ and $L'_\bullet$ are two perfect complexes, their total tensor product $L_\bullet \otimes^{\mathbf{L}} L'_\bullet$ is also perfect). More precisely, we should define "exterior-power" operations in the derived category $D^-(X)$, taking perfect complexes to perfect complexes. This question has not yet been clarified at present, and, as was stated in the Introduction, it constitutes the most serious gap in the Seminar. Once this question has been resolved, and proceeding as in 2.3, we obtain a natural filtration on $K^\bullet(X)$, allowing us to form the associated graded ring $\mathrm{Gr}^\bullet(X)$, which will play the role of the Chow ring, and to define a theory of Chern

classes

$$c_i : K^\bullet(X) \longrightarrow \mathrm{Gr}^\bullet(X).$$

4.3

To give a meaning to formula (1.2) for a proper locally complete intersection morphism $f : X \to Y$ of noetherian schemes, we must still include in the statement of this formula the nontrivial inclusions (3.4) (where d is the virtual relative dimension) to allow (3.3) to be defined, and finally define the virtual relative tangent class [print 11]

$$T_f \in K^\bullet(X).$$

When f admits a factorization (2.1) through an immersion into a smooth relative scheme, the definition of T_f presents no new difficulty. We can moreover reformulate it in this case, in a manner more in keeping with the "derived categories" point of view, by considering the familiar canonical homomorphism

$$\underline{N}_{X/X'} \xrightarrow{d} \Omega^1_{X'/Y} \otimes_{\underline{O}_{X'}} \underline{O}_X \tag{4.1}$$

induced by the exterior differential

$$d_{X'/Y} : \underline{J} \subset \underline{O}_{X'} \longrightarrow \Omega^1_{X'/Y}, \text{ [9]}$$

and the chain complex

$$L_\bullet^{X/Y} \in \mathrm{Ob}\, D(X) \tag{4.2}$$

defined from (4.1) "by extending by zeros", a complex equal in degree 1, respectively 0, to the source, respectively the target, of the homomorphism (4.1). Formula (2.2) then takes the more pleasant form

$$\check{T}_f = \mathrm{cl}^\bullet(L_\bullet^{X/Y}), \tag{4.3}$$

a formula which makes sense in $K^\bullet(X)$ when f is a locally complete intersection morphism, since in this case the complex $L_\bullet^{X/Y}$ is manifestly perfect. The assertion of invariance of T_f relative to the chosen factorization of f can then readily be made more precise as an assertion of invariance of the complex $L_\bullet^{X/Y}$: up to canonical isomorphism in the derived category $D(X)$, this complex is likewise independent of the chosen factorization (Exp. VIII).

4.4.

When $f : X \to Y$ is an arbitrary morphism of schemes, we can still construct a canonical object $L_\bullet^{X/Y}$ of the derived category $D(X)$, defined up to unique isomorphism, which on each affine open of X over an affine open of Y is isomorphic to the complex constructed according to the principle explained in the preceding paragraph (where, however, in the general case "smooth" should be replaced by "formally smooth", so that every algebra B over a ring A occurs as a quotient of a formally smooth algebra, for example a polynomial algebra). For such a definition, we cannot proceed by simple "gluing" from the affine constructions, because the objects of the derived category $D(X)$ are by their nature [print 12] essentially not glueable. Here is a general construction of $L_\bullet^{X/Y}$, valid more generally for every morphism $f : X \to Y$ of (commutatively) ringed topoi: let $B = \underline{O}_X$, $A = f^{-1}(\underline{O}_Y)$, so that A is a Ring on the space X, and B is an A-Algebra. Let $T \to B$ be a homomorphism of sheaves of sets generating B as an A-Algebra, i.e. such that the corresponding homomorphism of A-Algebras

$$\phi : C = A[T] \longrightarrow B$$

is an epimorphism of sheaves of sets, where $A[T]$ denotes the free commutative A-Algebra generated by T (i.e. the sheaf associated with the presheaf $U \longmapsto A(U)[T(U)]$ = free commutative $A(U)$-algebra generated by $T(U)$). Let J be the kernel of ϕ, and consider, as in 4.3, the chain complex of length 1 defined by the natural homomorphism

$$J/J^2 \xrightarrow{d} \Omega^1_{C/A} \otimes_C B.$$

It is not difficult to see that, up to canonical isomorphism in the category $D(X)$, the complex so constructed is independent of the choice of T and of $T \to B$, so that it deserves the name relative cotangent complex $L_\bullet^{X/Y}$ of X over Y. We also verify that, on every affine open of X over an affine open of Y, it is given (up to canonical isomorphism) by the procedure of 4.3.

When f is locally complete intersection, we moreover see that we even obtain a perfect complex, which then allows us again to define an element T_f in $K^\bullet(X)$ by formula (4.3). We point out that here again, in general, it would not have been possible to define T_f as an element of $K^\bullet_{\text{naive}}(X)$, and nothing permits us to suppose that T_f is globally isomorphic in $D(X)$ to a bounded complex with locally free components of finite type. This therefore provides further justification for the point of view advanced in 4.1 in favour of the "sophisticated" $K^\bullet(X)$, in preference to $K^\bullet_{\text{naive}}(X)$.

4.5.

Subject to the question raised in 4.2, we have therefore again assembled all the structural elements needed to give a meaning to (1.2) for a proper locally complete intersection morphism of arbitrary schemes (the noetherian hypothesis implicitly made in the preceding sections can in fact be removed without essential difficulty). Unfortunately, this does not mean that, even subject to this reservation, the formula in question has been established, even when Y is the spectrum of a field [print 13] that is algebraically closed of characteristic $p > 0$, and X is, say, a proper smooth algebraic scheme of dimension 3. See the comments in Exp. XIV no. 2.

5.

In the form presented in no. 4, the Riemann–Roch formula (1.2) also has a meaning for a proper locally complete intersection morphism of complex analytic spaces (or of rigid analytic spaces in the sense of Tate). But of course the proof given in [RRR] applies to this case only when we also assume that the morphism f is projective, and, as in the algebraic case, it seems that an essentially new idea is needed to treat the general case. The formula we envisage here has values in an "analytic Chow ring" $\mathrm{Gr}^\bullet(X)$, or rather $\mathrm{Gr}^\bullet(X)_{\mathbb{Q}}$, constructed according to the principle explained in 4.2 (the difficulty encountered there moreover disappears because we are in characteristic zero, as noted in Exp. XIV no. 1).

When X and Y are nonsingular, another version of the Riemann–Roch formula can be given as follows. First let X be an arbitrary complex analytic space, and X^{top} the same space equipped with the sheaf of continuous complex-valued functions. We thus have a canonical morphism of ringed spaces

$$X^{\text{top}} \longrightarrow X,$$

which, by the evident contravariance of the functor $K^\bullet(X)$ in the variable ringed space X, defines a ring homomorphism

$$K^\bullet(X) \longrightarrow K^\bullet(X^{\text{top}}). \tag{5.1}$$

For simplicity, suppose that X^{top} is compact; then, as in the case (already mentioned in 3.1) of a scheme admitting an ample invertible module, we see (Exp. II) that every perfect complex on X^{top} is globally isomorphic to a bounded complex locally free in each degree, i.e. to a complex

of complex vector bundles on X^{top}, and consequently $K^\bullet(X^{\mathrm{top}})$ is canonically isomorphic to the topologists' familiar group $K(X^{\mathrm{top}})$ [1], defined in terms of complex vector bundles on the compact space X^{top}. Thus we can interpret the homomorphism (5.1) as a homomorphism from $K^\bullet(X)$, defined in terms of the analytic structure, to the topologists' familiar ring $K(X^{\mathrm{top}})$. Now, for a morphism $f : X \longrightarrow Y$ of compact nonsingular analytic spaces, we know how to associate, by Atiyah–Hirzebruch [print 14] [1], a group homomorphism

$$f_*^{\mathrm{top}} : K^\bullet(X^{\mathrm{top}}) \longrightarrow K^\bullet(Y^{\mathrm{top}}). \tag{5.2}$$

Using likewise the homomorphism f_* of (3.1) defined by formula (3.2) (which here implicitly uses Grauert's finiteness theorem [3][10] for a proper morphism of analytic spaces), we obtain a square of natural homomorphisms involving only groups of type $K^\bullet$ (to the exclusion of rings of Chow-ring or cohomology type):

$$\begin{array}{ccc} K^\bullet(X) & \longrightarrow & K^\bullet(X^{\mathrm{top}}) \\ f_* \downarrow & & \downarrow f_*^{\mathrm{top}} \\ K^\bullet(Y) & \longrightarrow & K^\bullet(Y^{\mathrm{top}}). \end{array} \tag{5.3}$$

The Riemann–Roch type formula to which we alluded consists in asserting that this square commutes. We note the difference in nature between this assertion (which does not disregard torsion phenomena, and lies in a group $K^\bullet(Y^{\mathrm{top}})$ of an essentially discrete nature) and the Riemann–Roch formula of the type considered in no. 4, which lies in a group of "Chow" type that is no longer discrete in nature but must instead be tensored with $\mathbb{Q}$. We note that formula (5.3) is proved, in any case, when X and Y are nonsingular projective varieties [7], as is the formula of the type considered in no. 4, which is proved in this case in [RRR]. This is a plausible variant of the Atiyah–Singer index theorem (and of its generalizations due to Shih [6][11] and Atiyah), which plainly calls for a common generalization. Once the commutativity of (5.3) has been proved, the "differentiable Riemann–Roch theorem" of Atiyah–Hirzebruch [1] would immediately give a cohomological variant of the complex analytic Riemann–Roch formula, namely the commutativity of the square

$$\begin{array}{ccc} K^\bullet(X) & \xrightarrow{x \longrightarrow \mathrm{ch}(x)\,\mathrm{Todd}(T_X)} & H^{2*}(X, \mathbb{Q}) \\ \downarrow & & \downarrow \\ K^\bullet(Y) & \xrightarrow{y \longrightarrow \mathrm{ch}(y)\,\mathrm{Todd}(T_Y)} & H^{2*}(Y, \mathbb{Q}), \end{array} \tag{5.4}$$

where the horizontal arrows are obtained from the cohomological Chern classes, obtained from the homomorphism (5.1) and the ordinary Chern class homomorphism

$$K^\bullet(X^{\mathrm{top}}) \longrightarrow H^{2*}(X, \mathbb{Z}),$$

[print 15] the first vertical arrow is the same as in (5.3), and the second is the Gysin homomorphism in rational cohomology.

6.

Throughout this discussion, we have considered almost exclusively the contravariant invariant $K^\bullet(X)$, to the exclusion of the covariant invariant $K_\bullet(X)$ whose existence was mentioned in 3.1. Regarding $K^\bullet(X)$ as an analogue of the integral cohomology ring of a topological space, we must regard $K_\bullet(X)$ as an analogue of integral homology, the Module structure of $K_\bullet(X)$ over $K^\bullet(X)$ then corresponding to the cap-product operation(*). The fact that, when X is

(*) This analogy is moreover not purely verbal: after passing to the associated graded objects $\mathrm{Gr}^\bullet(X)$, $\mathrm{Gr}_\bullet(X)$, it can be made precise by an explicit compatibility relation between the module structure on $\mathrm{Gr}_\bullet(X)$ and the "cap" operation.

regular, the natural homomorphism $K^\bullet(X) \to K_\bullet(X)$ is an isomorphism must be regarded as the analogue of the Poincaré duality theorem on oriented topological manifolds, asserting that cap product with the fundamental homology class defines an isomorphism from cohomology to homology. These analogies also suggest the desirability of defining, say for a finite polyhedron X, an invariant $K_\bullet(X)$ of a covariant nature, whose relations to integral homology (spectral sequence, Chern homomorphism...) would be analogous to those between $K^\bullet(X)$ and integral cohomology; nor does it seem excluded that there might exist, in terms of these groups $K_\bullet(X)$, variants of the differentiable Riemann–Roch theorem valid for spaces more general than compact differentiable manifolds.

Returning to an arbitrary noetherian scheme X, we should equip $K_\bullet(X)$ with an increasing filtration, $\mathrm{Filt}_i\, K_\bullet(X)$ being generated by classes $\mathrm{cl}_\bullet(F)$ with F a coherent sheaf on X such that $\dim \mathrm{supp}\, .F \leq i$. We shall prove (Exp. X) the compatibility of this filtration with the module structure and the filtration of $K^\bullet(X)$, i.e. the formula

$$\mathrm{Filt}^i\, K^\bullet(X) \cdot \mathrm{Filt}_j\, K_\bullet(X) \subset \mathrm{Filt}_{j-i}\, K_\bullet(X), \text{ [12]} \tag{6.1}$$

which equips the associated graded object of the filtered module $K_\bullet(X)$, denoted by

$$\mathrm{Gr}_\bullet(X),$$

with a module structure over $\mathrm{Gr}^\bullet(X)$. Its elements can moreover be interpreted as classes of algebraic cycles on X, (for an equivalence relation less fine than rational equivalence, when X is of finite type over a base field K). If [print 16]

$$f : X \longrightarrow Y$$

is a proper morphism of noetherian schemes, then

$$f_* : K_\bullet(X) \longrightarrow K_\bullet(Y) \tag{6.2}$$

is compatible with the filtrations and defines a homomorphism of graded groups

$$f_* : \mathrm{Gr}_\bullet(X) \longrightarrow \mathrm{Gr}_\bullet(Y), \tag{6.3}$$

giving a "projection formula", i.e. it is $\mathrm{Gr}^\bullet(Y)$-linear, a formula which is deduced by passage to the associated graded objects from[13] the analogous projection formula for the homomorphism (6.2) (which is $K^\bullet(Y)$-linear).

From the point of view proposed in the Seminar, a workable theory of intersections, on noetherian schemes X that are not necessarily regular, must consist in the combined study of the contravariant invariants $K^\bullet(X), \mathrm{Gr}^\bullet(X)$ and the covariant invariants $K_\bullet(X), \mathrm{Gr}_\bullet(X)$, which play roles of an essentially different nature, and whose properties complement one another. Thus the covariant groups $K_\bullet(X), \mathrm{Gr}_\bullet(X)$ can be calculated more readily for certain nonproper fibrations, by means of the "homotopy exact sequence" that can be established for them ([RRR] and Exp. IX); for example, we prove (Exp. IX) canonical isomorphisms such as

$$K_\bullet(X[t]) \simeq K_\bullet(X), \qquad \mathrm{Gr}_\bullet(X[t]) \simeq \mathrm{Gr}_\bullet(X),$$

which fail in general for $K^\bullet$ when X is a nonregular scheme. On the other hand, $K^\bullet(X)$ and $\mathrm{Gr}^\bullet(X)$ lend themselves well to calculation because of their ring structures.

For schemes X proper over a field, the projection

$$f^X : X \longrightarrow (\text{point})$$

defines an "Euler–Poincaré characteristic" homomorphism

$$\lambda_X = f_*^X : K_\bullet(X) \longrightarrow K_\bullet(\text{point}) = \mathbb{Z}, \tag{6.4}$$

inducing on the subgroup $\mathrm{Gr}_0(X) = \mathrm{Filt}_0 K_\bullet(X)$ the "degree of 0-cycles" homomorphism

$$\deg_X : \mathrm{Gr}_0(X) \longrightarrow \mathbb{Z}. \tag{6.5}$$

The multiplication homomorphism

$$\mathrm{Gr}^i(X) \times \mathrm{Gr}_i(X) \longrightarrow \mathrm{Gr}_0(X),$$

composed with the degree homomorphism (6.5), then gives a pairing

$$\mathrm{Gr}^i(X) \times \mathrm{Gr}_i(X) \longrightarrow \mathbb{Z}, \tag{6.6}$$

which explains why, in certain respects, $\mathrm{Gr}^\bullet(X)$ and $\mathrm{Gr}_\bullet(X)$ play dual roles, [print 17] just as cohomology and homology do. Thus, for a morphism $f : X \to Y$ of proper algebraic schemes, the two homomorphisms

$$f^* : \mathrm{Gr}^\bullet(Y) \longrightarrow \mathrm{Gr}^\bullet(X) \qquad \text{and} \qquad f_* : \mathrm{Gr}_\bullet(X) \longrightarrow \mathrm{Gr}_\bullet(Y)$$

are formally transposes of one another, with respect to the pairings (6.6).

The homomorphisms (6.4) and (6.5), and the pairing (6.6) obtained from them, can be regarded as the source of numerical relations in intersection theory on proper algebraic schemes over a given field k. This point of view will be developed in some detail in Exp. X, which is largely independent of Exp. VI, VII, VIII (centred on the proof of the Riemann–Roch theorem) and of Exp. IX. Some notions concerning the behaviour of $K^\bullet(X), \mathrm{Gr}^\bullet(X)$ under specialization will also be given there.

7.

As in every theory of intersections, it is important to specify the place of the Picard group in the theory (or the group of divisor classes in the classical terminology). Subject to a slight restriction, automatically satisfied for example when X is quasi-compact, Exp. X establishes a canonical isomorphism

$$c_1 : \mathrm{Pic}(X) \xrightarrow{\sim} \mathrm{Gr}^1(X), \tag{7.1}$$

which shows the exact role of the Picard group in the theory we propose. This role justifies a more detailed study of this Picard group in Exposés XII, XIII, which are logically independent of the text of the Seminar. A study of numerical equivalence and certain finiteness theorems for the Picard group (announced without proof in [5][14]) is made by Kleiman in Exp. XIII, using techniques of a purely numerical nature (which do not use the theory of the Picard functor and its representability), due to himself and to Mumford, markedly simpler than Grothendieck's initial proofs. The only result refractory to Kleiman's methods (and which he is obliged to use) is [5, C-07 (i)][15], whose proof and certain refinements (including certain representability theorems for the Picard functor) occupy Exposé XII. The latter moreover has a spirit and a technique entirely foreign to the rest of the Seminar, being connected with [5]; this ex[print 18]posé is logically independent of all the others.

Finally, in Exp. XI, of a distinctly more general nature than XII and XIII and closer to the general spirit of the Seminar, we study, on an arbitrary locally ringed topos, a remarkable functor called the "determinant functor":

$$\overset{*}{\det} : \underline{\mathrm{Parfis}}(X) \longrightarrow \underline{\mathrm{Inv}}(X) \tag{7.2}$$

from the category of perfect complexes on X, whose morphisms consist only of the isomorphisms in the sense of the derived category $D(X)$, to the category of invertible Modules on X. This functor is described almost completely by the requirement that it be of a "local nature" and that, for a bounded complex $L_\bullet$ with locally free components, it be given by the formula

$$\overset{*}{\det}(L_\bullet) \simeq \bigotimes_i \det(L_i)^{(-1)^i}, \tag{7.3}$$

where $\det(L_i)$ denotes the exterior power of maximal degree of the locally free Module L_i. Although this exposé too is logically independent of the rest of the Seminar except for Exposé I, it nevertheless forms an important part of the general "yoga" proposed in the Seminar and developed mainly in Exposés I to XI.

[print 19]

BIBLIOGRAPHY

[1] M. Atiyah and F. Hirzebruch, Vector bundles and homogeneous spaces, Symposia in Pure Mathematics, vol. 3, Differential Geometry[16] p. 7–38, 1961.

[2] A. Borel and J.P. Serre, Le théorème de Riemann–Roch, Bull. Soc. math. France[17], t. 86, p. 97–136 (1958).

[3] H. Grauert, Ein Theorem der analytischen Garbentheorie, Pub. Math. no. 5, p. 5–64 (1960).

[4] A. Grothendieck, Classes de faisceaux et théorème de Riemann–Roch, mimeographed notes, Princeton 1957, (cited as [RRR] and reproduced as an Appendix to Exp. 0 of the Seminar).

[5] A. Grothendieck, Technique de descente et théorèmes d'existence en Géométrie Algébrique, in Fondements de la Géométrie Algébrique, Extracts from the Bourbaki Seminar 1957–1962, Secrétariat Mathématique, Paris.

[6] W. Shih, Séminaire de Topologie à l'IHES, 1966/67.

[7] M. Atiyah and F. Hirzebruch, The Riemann–Roch theorem for analytic embeddings, Topology, vol. 1, p. 151–166 (1962).

[print 20]

CLASSES OF SHEAVES AND THE RIEMANN–ROCH THEOREM

by A. GROTHENDIECK(*)

CHAP. I – λ-Rings (formal preliminaries)

1. Definitions 1
2. Examples 4
3. The λ-ring defined by a graded ring 8
4. The operations $\lambda^p(N, x)$ 13

CHAP. II – Classes of coherent algebraic sheaves and Chern classes 18

1. Chow's theory 18
2. Definition of the Chern classes of coherent algebraic sheaves 22
3. Functorial generalities on $K(X)$ 27
4. Some technical results 34
5. Sheaf-theoretic definition of Chern classes. Application to the study of injection morphisms 43
6. The Riemann–Roch theorem 49

Proof of the Riemann–Roch theorem (1 November 1957).

(*) This is the verbatim reproduction of the report cited in the Introduction to the present Seminar. The footnotes indicated by the signs (*), (**) were added in October 1967.

[print 21]

CHAPTER I

λ-Rings (Formal preliminaries)

§1. Definitions

A λ-ring(*) is a commutative ring K with unit, equipped with a family of maps

$$\lambda^i : K \longrightarrow K \tag{1.1}$$

(i an integer ≥ 0) satisfying the following conditions:

$$\lambda^0 x = 1, \qquad \lambda^1 x = x, \qquad \lambda^n(x+y) = \sum_{i=0}^{n} \lambda^i x \cdot \lambda^{n-i} y. \tag{1.2}$$

For every $x \in K$, put

$$\lambda_t(x) = \sum_{n \geq 0} \lambda^n(x) t^n \in K[[t]]; \tag{1.3}$$

the preceding relations then express that $x \longmapsto \lambda_t(x)$ is a homomorphism from the additive group K to the multiplicative group $1 + K[[t]]^+$ of formal series over K with augmentation 1, lifting the canonical homomorphism $1 + \sum_{i>0} x_i t^i \longrightarrow x_1$.

Thus, on the ring $\mathbb{Z}$, there exists a unique λ-structure compatible with its ring structure and such that [print 22]

$$\lambda_t(1) = 1 + t. \tag{1.4}$$

We then have

$$\lambda_t(n) = (1+t)^n, \tag{1.5}$$

hence $\lambda^i(n) = \binom{n}{i}$.

The notion of a λ-homomorphism of λ-rings is clear; an augmented λ-ring is a λ-ring equipped with a λ-homomorphism to $\mathbb{Z}$.

To define the notion of a special λ-ring, the following considerations are useful. Let k be a commutative ring with unit, and let $K = 1 + k[[t]]^+$ be the group of formal series with coefficients in k and augmentation 1; we shall introduce on it a λ-ring law whose additive structure is multiplication of formal series. The multiplication of this ring will be denoted by $f \circ g$. It is completely determined by the condition that the coefficients $c_n = c_n(f \circ g)$ of $f \circ g$ are given by universal polynomials with integer coefficients in the coefficients $a_i = c_i(f)$ and $b_i = c_i(g)$ of f and g, respectively:

$$c_n = P_n(a_1, \ldots, a_n; b_1, \ldots, b_n), \tag{1.6}$$

that it is distributive with respect to the "addition" fg, and is given for linear factors by:

$$(1+at) \circ (1+bt) = 1 + abt. \tag{1.7}$$

The P_n are calculated by calculations of symmetric functions; P_n is isobaric of weight n in the a_i and isobaric of weight n in the b_i (a_i and b_i having weight i), and is moreover symmetric in $a = (a_i)$ and $b = (b_i)$. Likewise, the $\lambda^i(f)$ are determined unambiguously by the condition that

(*) In this seminar we instead say pre-λ-ring (V 2.1), reserving the name λ-ring for those satisfying the additional condition on page 3 (cf V 2.4).

[print 23] the coefficients $c_n(\lambda^i(f)) = c_n$ of $\lambda^i(f)$ are calculated using universal polynomials with integer coefficients in the coefficients $c_n(f) = a_n$ of f:

$$c_n = P_{i,n}(a_1, \ldots, a_{in}), \tag{1.8}$$

that relations (1.2) hold, and that, for $i > 1$, we have:

$$\lambda^i(1 + at) = 1 \tag{1.9}$$

(1 is the zero element of the ring $K = 1 + k[[t]]^+$). The $P_{i,n}$ are again calculated by calculations of symmetric functions, and we find that $P_{i,n}$ is an isobaric polynomial of weight in in the a_i. Equipped with the preceding operations, K becomes a λ-ring; its zero element is 1, and its unit element is $1 + t$. We also note the relation:

$$(1 + at) \circ f = f(at) \tag{1.7 bis}$$

Now let K be an arbitrary λ-ring. We say that K is special when the additive homomorphism λ_t from K to $1 + K[[t]]^+$ is a λ-homomorphism. This therefore means that we have the formulas:

$$\lambda_t(1) = 1 + t, \quad \lambda_t(xy) = \lambda_t(x) \circ \lambda_t(y), \quad \lambda_t(\lambda^i(x)) = \lambda^i(\lambda_t(x)), \text{ [18]} \tag{1.10}$$

or, explicitly:

$$\begin{cases} \lambda^i(1) = 0 & \text{for } i > 1 \text{ [19]} \\ \lambda^n(xy) = P_n(\lambda^1 x, \ldots, \lambda^n x; \lambda^1 y, \ldots, \lambda^n y) \\ \lambda^n(\lambda^i(x)) = P_{i,n}(\lambda^1 x, \ldots, \lambda^{in} x), \text{ [20]} \end{cases} \tag{1.11}$$

where the P_n and $P_{i,n}$ are the universal polynomials considered above. In particular, formulas (1.5) follow. (We can show that the λ-ring $1 + k[[t]]^+$ constructed from a commutative ring k is spe[print 24]cial; but we shall not need this fact).

§2. Examples

1) Let G be a group and k a commutative ring; we consider the group $\underline{K}(G)$, the quotient of the free group generated by the classes of representations of G in k-modules of finite type by the subgroup generated by elements of the form $(\rho \oplus \rho') - (\rho) - (\rho')$ (we write $\oplus$ for direct sum). Tensor product and exterior powers define a λ-ring structure on $\underline{K}(G)$. We can also pass to the quotient by elements of the form $(\rho) - (\rho') - (\rho'')$, where ρ is a representation that is an extension of ρ' by ρ'', trivial as an extension of k-modules. We obtain the group $\underline{K}_r(G)$ of reduced classes of representations of G, and we find that the λ-ring law passes to the quotient.

When k is an algebraically closed field of characteristic 0(*), we can show that $\underline{K}(G)$, and a fortiori $\underline{K}_r(G)$, is a special λ-ring; this no longer holds when the characteristic is $\neq 0$, but nevertheless, when k is algebraically closed(*), $\underline{K}_r(G)$ is still a special λ-ring. To prove this, we reduce to the case where G is a product $\mathrm{Gl}(m, k) \times \mathrm{Gl}(n, k)$ and only rational representations of G are considered; our assertion follows from the explicit determination of $\underline{K}_r(G)$ to be made below (example 3); finally, when k has characteristic 0, complete reducibility of the rational representations of G shows that $\underline{K}(G) = \underline{K}_r(G)$.

[print 25] Variants of the preceding example: G is an algebraic group defined over k, and only the rational representations of G defined over k are considered; we can consider a topological group,

(*) It is unnecessary for k to be algebraically closed, cf. (VI 3.3) for a still more general result (including all those that will be used in the present report).

or a Lie group, and continuous representations, etc... ([1])

2) Let X be an algebraic variety, irreducible or not, defined over the field k. We can consider the quotient of the free group generated by the classes of vector bundles on X (defined over k) by the subgroup generated by elements of the form $(E \oplus E') - (E) - (E')$, on which the λ-ring operations will be defined by tensor product and exterior powers. When X is complete, this is a free group generated by the indecomposable vector bundles on X (by Krull–Remak–Schmidt); knowing this ring and its basis is equivalent to the complete classification of vector bundles on X and knowing the elementary tensor operations on those bundles. When k is algebraically closed of characteristic 0, all the other tensor operations on classes of bundles are then known, using what follows in 3). We can make a more drastic passage to the quotient by the group generated by the elements $(E) - (E') - (E'')$, where E is a vector bundle that is an extension of E' by E''; we then obtain a λ-ring, denoted by $\underline{K}(\xi(X))$, where $\xi(X)$ denotes the additive category of vector bundles on X defined over k. Using the results of example 1), we show that, when k is algebraically closed(*), the λ-ring $\underline{K}(\xi(X))$ is special, and the same holds for the [print 26]first λ-ring defined here, if the characteristic of k is 0, but not in the general case.

3) Determination of $\underline{K}(G)$ and of $\underline{K}_r(G)$ in certain important cases: We suppose that the field k is algebraically closed, that the group G is algebraic, affine connected and "globally reductive" (the radical of G is isomorphic to k^{*s}), and we consider only rational linear representations. In characteristic 0, the complete reducibility of the rational linear representations of G shows that $\underline{K}(G) = \underline{K}_r(G)$ is the free group generated by the classes of irreducible rational representations. Now, if T is a maximal torus of G, we have $\underline{K}(T) = \underline{K}_r(T) = \mathbb{Z}(\widehat{T})$, the algebra of the group $\widehat{T}$ of rational characters of T; we have $\widehat{T} \simeq \mathbb{Z}^r$ if r is the rank of G (this is independent of the characteristic). The Weyl group W operates on T, hence on $\underline{K}(T)$, in a way that depends only on the operations of W on the free group $\widehat{T}$. We have an evident homomorphism of λ-rings, deduced from the injection of T into G:

$$\underline{K}_r(G) \longrightarrow \underline{K}_r(T)^W = \underline{K}(T)^W, \tag{1.13}$$

where $\underline{K}(T)^W$ denotes the set of fixed points of W in $\underline{K}(T)$. The theory of linear representations of G(*) then proves the following theorem:

Theorem 1.1. The homomorphism (1.13) is an isomorphism.

We thus see that $\underline{K}_r(G)$, once the "type" of G is known (characterized by W operating on a group $\mathbb{Z}^r$), does not depend on the characteristic, and that it is a special λ-ring. We have a canonical basis of $\underline{K}_r(G)$ corresponding to the orbits of W in $\widehat{T}$; another basis is obtained as follows: [print 27]let $(\rho_i)_{1 \leq i \leq r'}$ be the representation classes of G corresponding to the fundamental representations of $G/\operatorname{rad} G$, and let $(\sigma_i)_{1 \leq i \leq r''}$ be a basis of the group of rational homomorphisms from G to k^*; we thus have $r' + r'' = r$ (the rank of G).

Corollary: The ring $\underline{K}_r(G)$ is identified with the subring

$$\mathbb{Z}[\rho_1, \ldots, \rho_{r'}; \sigma_1, \ldots, \sigma_{r''}; \sigma_1^{-1}, \ldots, \sigma_{r''}^{-1}]$$

of the field of rational functions in the ρ_i and the σ_j with coefficients in $\mathbb{Q}$(*).

In particular, $\underline{K}_r(G)$ has no divisors of 0; if $G = \prod_i \mathrm{Gl}(n_i, k)$, the ring $\underline{K}_r(G)$ is made explicit as follows: let ρ_i be the representation of G deduced from the identity representation of $\mathrm{Gl}(n_i, k)$;

([1]) Other examples of λ-rings can be constructed, for example the set of equivalence classes of quadratic forms, the set of classes of representations of a Lie algebra, etc... We promise that we shall not need all these examples to treat the Riemann–Roch theorem!

(*) Unnecessary condition, cf. the preceding footnote.

(*) Cf. Séminaire Chevalley 1956/58 Groupes de Lie algébriques[21] (École Normale Supérieure).

(*) The derived group G' of G must be assumed simply connected for this statement to be correct.

if we then consider the field of rational functions over $\mathbb{Q}$ in indeterminates $\lambda^j(\rho_i)$ $(1 \leq j \leq n_i)$, then $\underline{K}_r(G)$ is identified with the subring generated by these indeterminates and the inverses of the $\lambda^{n_i}(\rho_i)$. This justifies our assertion that, at least in characteristic 0, the operations λ^i allow all tensor operations on vector bundles to be reconstructed; this is still true over an algebraically closed field of arbitrary characteristic, provided reduced classes are taken.

Remarks: a) In example 1), the ring $\underline{K}(G)$ is augmented, the augmentation being defined by the degree of the representations, and, by passage to the quotient, this augmentation defines an augmentation on $\underline{K}_r(G)$. Likewise, the λ-rings considered in example 2) are augmented when X is k-irreducible; in $\underline{K}(\xi(X))$, the augmentation is defined by the rank of a vector bundle, that is, the dimension of its fibres.

[print 28]b) When the field k is not algebraically closed, it is plausible that $\underline{K}(G)$ is a special λ-ring in characteristic 0, and likewise for $\underline{K}_r(G)$ in all characteristics.

c) In nonzero characteristic, theorem 1.1 was kindly supplied to me by Chevalley.

§3. The λ-ring defined by a graded ring

Let A be a commutative graded ring with unit; for simplicity we shall assume that the degrees of A are positive and that $A^0 = \mathbb{Z}$. We denote by $\widehat{A}$ the product of the A^i (it is a commutative ring with unit), and by $\widehat{A}^+$ the kernel of the evident augmentation $\widehat{A} \to A^0 = \mathbb{Z}$. Then $1 + \widehat{A}^+$ is a subgroup of the multiplicative group of invertible elements of $\widehat{A}$. Consider the abelian group

$$\widetilde{A} = \mathbb{Z} \times (1 + \widehat{A}^+), \tag{1.14}$$

whose elements will be denoted $[n, x]$ with $n \in \mathbb{Z}$ and $x = 1 + \sum_{i \geq 1} x^i \in 1 + \widehat{A}^{+[22]}$ $(x^i \in A^i)$. We shall write this group additively, thus:

$$[n, x] + [n', x'] = [n + n', xx']. \tag{1.15}$$

The zero element of this group is $[0, 1]$. We are going to introduce on it an augmented λ-ring structure, compatible with its additive structure, the augmentation being $\varepsilon : [n, x] \longrightarrow n$. The product $[m, x][n, y]$ is completely characterized by being written $[mn, x^n y^m (x * y)]$, where $x * y$ is an element of $1 + \widehat{A}^+$ whose components are expressed as universal polynomials with integer coefficients in the x^i and the y^i:

$$(x * y)^i = Q_i(x^1, \ldots, x^i; y^1, \ldots, y^i) \qquad (i \geq 1) \tag{1.16}$$

[print 29]the polynomial Q_i being isobaric of weight i. Moreover, $x * y$ must be distributive with respect to the "addition" xy and, for "linear" factors, must reduce to the following:

$$[1, 1 + x^1][1, 1 + y^1] = [1, 1 + x^1 + y^1]. \tag{1.17}$$

The polynomials Q_i are evaluated by calculations with elementary symmetric functions (this is the calculation of the Chern classes of a tensor product of vector bundles by means of the Chern classes x^i and y^i of the factors and their respective dimensions m and n). From (1.17) we deduce:

$$(1 + x^1) * (1 + y^1) = (1 + x^1 + y^1)/(1 + x^1)(1 + y^1), \tag{1.17 bis}$$

and an arbitrary product $x * y$ is calculated term by term by decomposing "formally" x into a product $\prod_{i=1}^n (1 + \alpha_i)$ and y into a product $\prod_{i=1}^n (1 + \beta_i)$, where the α_i and the β_i have degree 1.

In this way, $\widetilde{A}$ becomes an augmented commutative ring, with a unit $[1, 1]$, and it can be interpreted as the ring obtained by adjoining a unit to the ring $1 + \widehat{A}^+$. Moreover, on $\widetilde{A}$, we

define a filtration by putting $\widetilde{A}^0 = \widetilde{A}$ and denoting by $\widetilde{A}^n$ the set of the $[0,x]$ with $x^i = 0$ for $1 \leq i < n$; this filtration is compatible with the product, for we have

$$\left(1 + \sum_{i \geq m} x^i\right) * \left(1 + \sum_{j \geq n} y^j\right) = 1 - \frac{(m+n-1)!}{(m-1)!(n-1)!} x^m y^n + \cdots , \tag{1.18}$$

the unwritten terms having degree $> m+n$. This formula shows that the graded ring $G(\widetilde{A})$ associated with $\widetilde{A}$ is identified with A from the additive point of view, but not from the multiplicative point of view; we can only say that the map η' from A to $G(\widetilde{A})$ defined by [print 30]

$$\eta'(x^i) = (-1)^{i-1}(i-1)!\, x^i \qquad (x^i \in A^i) \tag{1.19}$$

is a ring homomorphism.

We introduce on $\widetilde{A}$ operations λ^i satisfying the axioms of λ-rings (§1, formulas (1.2)), well determined by the condition that the components of $\lambda^i[0,x]$ be deduced from the components x^i of x by universal polynomials with integer coefficients

$$\lambda^i[0,x] = [0, \lambda^i x] \qquad (\lambda^i x)^{(n)} = Q_{i,n}(x^1, \ldots, x^n) \tag{1.20}$$

($Q_{i,n}$ is isobaric of weight n), and that we have

$$\lambda^i[1, 1+x^1] = 0 \qquad \text{for } i > 1. \tag{1.21}$$

The polynomials $Q_{i,n}$ are evaluated by a calculation with elementary symmetric functions, which is the calculation of the Chern classes of the exterior powers of a vector bundle, by means of the Chern classes and the rank of the latter. We then find that $\widetilde{A}$ is a special augmented λ-ring and that the $\widetilde{A}^i$ are stable under the operations λ^j.

The augmented λ-ring $\widetilde{A}$ may be considered as a functor in A; consequently, the principal automorphism

$$x = \sum_{i \geq 0} x^i \longrightarrow \check{x} = \sum_{i \geq 0} (-1)^i x^i$$

of A defines an automorphism of $\widetilde{A}$, denoted by the same symbol $\check{}$.

Proposition 1.2. For $x \in \widetilde{A}$ of the form $[n, 1 + x^1 + \cdots + x^n]$ (we therefore have $x^i = 0$ for $i > n$), we have $\lambda^i(x) = 0$ for $i > n$, we have $\lambda^n(x) = [1, 1+x^1]$, and finally

$$\sum_{i=0}^{n} (-1)^i \lambda^i(x) = [0, 1 - (n-1)! x^n + \cdots] \tag{1.22}$$

From this proposition and from the fact that the $\widetilde{A}^i$ are stable under the operations λ^j, we deduce that formula (1.22) is valid whenever $\varepsilon(x) = n$, whether or not we have $x^i = 0$ for $i > n$.

[print 31]Formula (1.22) admits the following generalizations: let K be any λ-ring, otherwise arbitrary; for $x \in K$, put

$$\gamma^n(x) = (-1)^n \sum_{i=0}^{n} (-1)^i \lambda^i(x+n) = \lambda^n(x+n-1) \tag{1.23}$$

(the identity of the last two members follows from an immediate induction). In particular, we have $\gamma^0(x) = 1$ and $\gamma^1(x) = x$; next put:

$$\gamma_t(x) = \sum_{n \geq 0} \gamma^n(x) t^n \in K[[t]] \tag{1.24}$$

whence we deduce the formula:

$$\gamma_t(x) = \lambda_{t/(1-t)}(x), \tag{1.25}$$

which is equivalent to

$$\lambda_s(x) = \gamma_{s/(1+s)}(x). \tag{1.25 bis}$$

This proves that the λ^i are expressed by means of the γ^i and that $\gamma_t(x+y) = \gamma_t(x)\gamma_t(y)$, and consequently that the maps γ^i define a new λ-ring structure on K. Formula (1.22) then shows that, for $K = \widetilde{A}$ and every $x \in \widetilde{A}$, we have

$$\gamma^n(x - \varepsilon(x)) = [0, 1 + (-1)^{n-1}(n-1)!x^n + \cdots] \tag{1.22 bis}$$

and consequently, the component x^n is obtained, up to the factor $(-1)^{n-1}(n-1)!$, by reducing modulo $\widetilde{A}^{n+1}$ the element $\gamma^n(x - \varepsilon(x))$ of $\widetilde{A}^n$.

The Chern homomorphism: For every graded ring A satisfying the preceding conditions, we are going to define a homomorphism of augmented rings:

$$\mathrm{ch} : \widetilde{A} \longrightarrow {}^{[23]}\widehat{A \otimes \mathbb{Q}}. \tag{1.26}$$

This homomorphism is defined unambiguously by the conditions that it be an additive homomorphism, take 1 to 1, be such that the com[print 32]ponents of $\mathrm{ch}(x)$ in degree > 0 be given by universal isobaric polynomials with rational coefficients in the components of x, and that

$$\mathrm{ch}([1, 1 + x^1]) = \exp x^1 = \sum_{n \geq 0} (x^1)^n / n!. \tag{1.27}$$

We then immediately obtain the explicit expression for ch: let η be the additive endomorphism of $\widehat{A \otimes \mathbb{Q}}$ defined, for x^i of degree i, by

$$\eta(x^i) = 1/(-1)^{i-1}(i-1)!\, x^i \tag{1.28}$$

we then have

$$\mathrm{ch}\left([n, 1 + \sum_{i \geq 1} x^i]\right) = n + \eta\left(\log\left(1 + \sum_{i \geq 1} x^i\right)\right). \tag{1.29}$$

As we can "invert" this formula, at least when A has only finitely many homogeneous components, we see that, in this case, we have an isomorphism ch from the ring $\widetilde{A} \otimes \mathbb{Q}$ onto the ring $\widehat{A \otimes \mathbb{Q}}$ (which completely elucidates the structure of $\widetilde{A} \otimes \mathbb{Q}$).

Recall that, according to Hirzebruch [1], a formal series $f \in \mathbb{Q}[[t]]$ defines, by universal polynomial formulas, an additive homomorphism $1 + \widehat{A}^+ \longrightarrow 1 + (\widehat{A \otimes \mathbb{Q}})^+$, denoted $\mathcal{T}_f$; when $f(t) = t/(1 - \exp(-t))$, we simply put $\mathcal{T}_f = \mathcal{T}$ ($\mathcal{T}$ is the initial of Todd). We extend the definition of $\mathcal{T}(x)$ to $x \in \widetilde{A}$. This said, we have the following result:

Proposition 1.3. Let $N \in \widetilde{A}$ be of the form $[q, 1 + N^1 + \cdots + N^q]$ (we have $N^i = 0$ for $i > q$). If we put:

$$\lambda_{-1}(N) = \sum_{i \geq 0} (-1)^i \lambda^i(N) = \sum_{i=0}^{q} (-1)^i \lambda^i(N)^{[24]},$$

we shall have:

$$\mathrm{ch}(\lambda_{-1}(N)) = (-1)^q N^q \mathcal{T}(\check{N})^{-1}, \tag{1.30}$$

[print 33]which is equivalent to:

$$\mathrm{ch}(\lambda_{-1}(\check{N})) = N^q \mathcal{T}(N)^{-1}. \tag{1.30 bis}$$

§4. The operations $\lambda^p(N,x)$

Let A be the ring of formal series, with coefficients in $\mathbb{Z}$, in two infinite sequences of indeterminates, denoted $\lambda^i N$ and $\lambda^i x$ ($i \geq 1$). Put $\lambda^1 N = N$ and $\lambda^1 x = x$, and introduce on A the unique special λ-ring structure for which we have $\lambda^i(N) = \lambda^i N$ and $\lambda^i(x) = \lambda^i x$ and for which the operations λ^i are "continuous" (if we restricted ourselves to polynomials in indeterminates $\lambda^i N$ and $\lambda^i x$, we would have the "free λ-ring generated by N and x"; our ring A is a completion of it). We can then form

$$\lambda_{-1}(N) = \sum_{i \geq 0} (-1)^i \lambda^i N, \tag{1.31}$$

which is a formal series with constant term 1, hence invertible; we can therefore define an element $\lambda^p(N,x)$ of A unambiguously by putting

$$\lambda^p(N,x)\lambda_{-1}(N) = \lambda^p(x\lambda_{-1}(N)) \tag{1.32}$$

Theorem 1.4. Let J_q be the closed ideal of A generated by the $\lambda^i N$ with $i > q$; identify A/J_q with the ring of formal series in the $\lambda^i x$ and the $\lambda^j N$ ($1 \leq j \leq q$). Then $\lambda^p(N,x)$ reduced modulo J_q is a polynomial, isobaric of weight p with respect to the variables $\lambda^i x$.

Let N and F be two vector spaces of finite dimensions q, q' over an algebraically closed field k of characteristic 0; let $N^{(p)}$ be the kernel of the map $(a_1, \ldots, a_p) \longrightarrow a_1 + \cdots + a_p$ from N^p to N. We [print 34]let the symmetric group $\mathfrak{S}_p$ operate on N^p, hence on $N^{(p)}$ hence on $\Lambda(N^{(p)})$, then on $\bigotimes^p F$ and on $\Lambda(N^{(p)}) \otimes \bigotimes^p F$. Consider this vector space as a graded module over $\mathfrak{S}_p \times \mathrm{Gl}(N) \times \mathrm{Gl}(F)$ and take its "alternating part", which is a graded module over $\mathrm{Gl}(N) \times \mathrm{Gl}(F) =: G$. We then have

$$\lambda^p(N,F) = \underline{E}_G\left(\left(\Lambda(N^{(p)}) \otimes \bigotimes^p F\right)^{\mathrm{alt}}\right). \tag{1.33}$$

N.B. If M is a complex of G-modules, we denote by $\underline{E}_G(M)$ the alternating sum in $\underline{K}(G)$ of the classes of the components of M; this is the Euler–Poincaré characteristic of M; in formula (1.33), the first member $\lambda^p(N,F)$ denotes the element of the ring $\underline{K}(G)$ obtained by substituting for the variables $\lambda^i N$ and $\lambda^i F$ the classes in $\underline{K}(G)$ of the G-modules $\Lambda^i(N)$ and $\Lambda^i(F)$ respectively.

Proof: By expanding the second member of (1.32) by the second formula (1.11), we see that the first assertion is proved if we prove the analogous assertion obtained by putting $x = 1$, i.e. if we prove that, in the λ-ring $\mathbb{Z}[\lambda^1 N, \ldots, \lambda^q N]$ where $\lambda^i N = 0$ for $i > q$, the element $\lambda^p(\lambda_{-1}(N))$ is divisible by $\lambda_{-1}(N)$. A priori, the quotient $\lambda^p(N,1) = \lambda^p(\lambda_{-1}(N))/\lambda_{-1}(N)$ is a formal series contained in the field of fractions K of the ring of formal series in the $\lambda^i N$ for $1 \leq i \leq q$. On the other hand, also denote by N a vector space of dimension q over an algebraically closed field k of characteristic 0, and consider the element $\ell^p(N) = \underline{E}_G((\Lambda(N^{(p)}))^{\mathrm{alt}})$[25] $\in \underline{K}(G)$, where we now have $G = \mathrm{Gl}(N)$. We know (§2, theorem 1.1)[26] that $\underline{K}(G)$ is identified [print 35]with the subring of K generated by the $\lambda^i N$ for $1 \leq i \leq q$ and by the inverse of $\lambda^q N$. On the other hand, we readily verify in this ring that we have $\ell^p(N)\lambda_{-1}(N) = \lambda^p(\lambda_{-1}(N))$. As we are in a large field K and $\lambda_{-1}(N) \neq 0$, we conclude that $\ell^p(N) = \lambda^p(N,1)$; both members are in the intersection of $\underline{K}(G)$ and the ring of formal series in the $\lambda^i N$, and are therefore polynomials in the $\lambda^i N$.

To prove (1.33) in the general case, we verify that the second member satisfies in $\underline{K}(G)$ the formula analogous to (1.32), that is, its product by $\lambda_{-1}(N)$ is $\lambda^p(F\lambda_{-1}(N))$. Likewise, the product of $\lambda^p(N,F) \in \underline{K}(G)$ by $\lambda_{-1}(N)$ is $\lambda^p(F\lambda_{-1}(N))$, as follows from the fact that $\underline{K}(G)$ is special. As $\underline{K}(G)$ is an integral domain (cf. §2) and $\lambda_{-1}(N) \neq 0$ (same reference), we have indeed proved formula (1.33). q.e.d.

Remarks: a) When the dimension q' of F is $\geq p$, then (1.33) characterizes the polynomial $\lambda^p(N,x)$ in the $\lambda^i N$ $(1 \leq i \leq q)$ and the $\lambda^j x$ $(1 \leq j \leq p)$.

b) It is plausible that (1.33) still holds when the field k is not algebraically closed(*). This question would have to be resolved to prove the Riemann–Roch theorem over a field of characteristic 0 that is not algebraically closed by the method we are going to explain. Moreover, the passage from an $\mathfrak{S}_p$-module to its alternating part has a reasonable meaning only when the base field has characteristic 0.

Now let K be any special λ-ring and let N be an ele[print 36]ment of K such that $\lambda^i N = 0$ for sufficiently large i. We can then, for every $x \in K$ and every integer $p \geq 1$, consider the elements $\lambda^p(N,x)$, which are expressed by universal polynomials with integer coefficients in the $\lambda^i N$ and the $\lambda^j x$. We have $\lambda^1(N,x) = x$ and we have formula (1.32) in K. To express the properties of the operations $x \mapsto \lambda^p(N,x)$, it is convenient to introduce a new λ-ring denoted K_N. As a commutative ring, K_N is obtained by adjoining a unit to the ring whose underlying additive group is K and whose multiplication is given by

$$x_N y = xy\mu \qquad (\mu = \lambda_{-1}(N)). \tag{1.34}$$

The map λ_t from K_N to $K_N[[t]]$ is defined by:

$$\lambda_t(n,x) = (1+t)^n \sum_{p \geq 0} \lambda^p(N,x) t^p, \tag{1.35}$$

where we have put $\lambda^0(N,x) = (1,0)$ (the unit element of K_N). I say that, equipped with these operations, K_N is a special λ-ring. We therefore have to verify the formulas $\lambda_t(X+Y) = \lambda_t(X)\lambda_t(Y)$, $\lambda_t(XY) = \lambda_t(X).\lambda_t(Y)$, and $\lambda_t(\lambda^i(X)) = \lambda^i(\lambda_t(X))$ for $X, Y \in K_N$. We can restrict ourselves to verifying these formulas when X and Y are in K, and we find that it suffices to verify them when K is the free special λ-ring generated by X, Y and N subject to the relations $\lambda^i N = 0$ for $i > q$. But the group homomorphism

$$K_N \longrightarrow K, \tag{1.36}$$

which takes unit to unit and on K reduces to the homomorphism $x \longrightarrow x\mu$, is a homomorphism compatible with the ring structures and the maps λ^i, as we verify immediately. In the present case, [print 37]its restriction to K is injective; as K is a special λ-ring, it follows readily that the above relations do hold for $X, Y \in K$, and this completes the proof of our assertion.

According to formula (1.23), it is natural, for $x \in K$ and $n \geq 1$, to put

$$\gamma^n(N,x) = \sum_{i=0}^{n-1} \binom{n-1}{i} \lambda^{n-i}(N,x)^{[27]}, \tag{1.37}$$

in other words, we put $\gamma^n(N,x) = \gamma^n(x_N)$, where x_N denotes x considered as an element of the λ-ring K_N.

Proposition 1.5. Let A be a commutative graded ring with unit, N an element of $\widetilde{A}$ of the form $N = [q, 1 + N^1 + \cdots + N^q]$, and x an arbitrary element of $\widetilde{A}$: $x = [\nu, 1 + \sum_{i \geq 1} x^i]$. Then, for every integer $j \geq 1$[28], we have $\gamma^{q+j}(N,x) \in \widetilde{A}^j$, and the component of degree j of $\gamma^{q+j}(N,x)$ is of the form

$$\Big(\gamma^{q+j}(N,x)\Big)^{(j)} = (-1)^{j-1}(j-1)! G_{q,j}(\nu, x^1, \ldots, x^j; N^1, \ldots, N^{j\,[29]}), \tag{1.38}$$

where $G_{q,j}$ is a universal polynomial with integer coefficients characterized by the condition

$$G_{q,j}(\nu, x^1, \ldots, N^{j\,[30]}) N^q = (-1)^q (x * \lambda_{-1}(N))^{(q+j)}. \tag{1.39}$$

(*) This is indeed proved by the same method.

Proof. We can assume that A is the ring of polynomials with integer coefficients in indeterminates x^i and N^j (with $1 \leq j \leq q$). As we evidently have, combining (1.32) and (1.37),

$$\gamma^n(N, x)\lambda_{-1}(N) = \gamma^n(x\lambda_{-1}(N)), \tag{1.40}$$

and, for $n = q + j$, the second member has filtration $\geq q + j$ (use formula (1.22 bis), taking into account that $x\lambda_{-1}(N)$ has [print 38]zero augmentation), whereas $\lambda_{-1}(N)$ has filtration q (formula (1.22)), it follows that $\gamma^n(N, x)$ has filtration j, since, by (1.18), the associated graded ring of A is here an integral domain. Using the preceding formulas, we immediately obtain the indicated form for $(\gamma^{q+j}(N, x))^{(j)}$.

q.e.d.

CHAPTER II

Classes of coherent algebraic sheaves

and Chern classes

§1. Chow's theory

Throughout this chapter, we assume a base field k fixed, which we shall assume algebraically closed, for simplicity. A large part of what follows, and perhaps all of it, is nevertheless valid without this restriction. The algebraic spaces and regular maps considered will be defined over k.

An algebraic space is called quasi-projective if it is isomorphic to a locally closed subset of a projective space.

Let X be a connected nonsingular quasi-projective variety of dimension n. We denote by $A(X)$, and call the Chow ring of X, the graded ring of classes of cycles on X modulo rational equivalence. ($A^i(X)$ consists of the classes of cycles of dimension $n - i$). The fact that $A(X)$ [print 39]is indeed a graded ring follows from the following theorem of Chow:

Theorem 2.1.(*) Every cycle on X is rationally equivalent to a nonsingular cycle. Given elements of $A^i(X)$ and $A^j(X)$, we can find nonsingular representatives of these elements, say Y^{n-i} and Z^{n-j}, which meet transversely everywhere.

(A cycle is called nonsingular if its components are nonsingular subvarieties. We say that two cycles meet transversely everywhere if every component of one meets every component of the other transversely everywhere, i.e. if the two intersecting varieties are nonsingular at their points of intersection and their tangent varieties there are in general position ...)

Let f be a morphism (i.e. a regular map) from X to Y (X and Y being quasi-projective and nonsingular); then f defines (using the preceding theorem) a homomorphism of graded rings:

$$f^* : A(Y) \longrightarrow A(X) \tag{2.1}$$

(inverse image of cycle classes). Thus $A(X)$ becomes a contravariant functor in X.

(*) As a result of a misunderstanding on the author's part, the theorem states more than is currently known. In fact, only Chow's "moving lemma" is used, saying that two cycles can be moved within their classes so that their intersection has no excess; cf. Séminaire Chevalley 1958, Anneau de Chow et applications.

Let f be a proper morphism from X^n to Y^m (i.e. in Weil's terminology, the graph of f is complete over Y, cf. [3]). Then f also defines an additive homomorphism

$$f_* : A(X^n) \longrightarrow A(Y^m) \tag{2.2}$$

preserving the dimension of cycles, and therefore increasing the degree by $m - n$.

[print 40]Thus, with respect to proper morphisms, and disregarding the multiplicative structures and gradings, $A(X)$ is a covariant functor in X. Let us note the classical formula

$$f_*(x.f^*(y)) = f_*(x).y. \tag{2.3}$$

We shall also use Chow's theory of Chern classes. It consists in attaching to every vector bundle E on X (still assumed quasi-projective and nonsingular) Chern classes $C^i(E) \in A^i(X)$, $(i \geq 1)$, in such a way as to satisfy the following conditions; put $C^0(E) = 1$ and

$$\widetilde{C}(E) = \left[\operatorname{rang} E, \sum_{i\geq 0} C^i(E)\right] \in \widetilde{A}(X) \tag{2.4}$$

(see chapter I, §3 for the definition of $\widetilde{A}(X)$). The conditions to be satisfied by a theory of Chern classes are then expressed as follows:

$$\widetilde{C}(E) = \widetilde{C}(E') + \widetilde{C}(E'') \tag{2.5}$$

if the vector bundle E is an extension of E' by E'' (putting $C(E) = \sum_{i\geq 0} C^i(E)$, this condition is expressed by $C(E) = C(E').C(E'')$). Formula (2.5) allows $\widetilde{C}$ to be extended to an additive homomorphism from $\underline{K}(\xi(X))$ (cf. Chap. I, §2, example 2) to $\widetilde{A}(X)$:

$$\widetilde{C} : \underline{K}(\xi(X)) \longrightarrow \widetilde{A}(X), \tag{2.6}$$

and we want $\widetilde{C}$ to be a λ-homomorphism of λ-rings; this last condition is also written

$$\widetilde{C}(E \otimes F) = \widetilde{C}(E)\widetilde{C}(F) \qquad \widetilde{C}(\Lambda^i E) = \lambda^i \widetilde{C}(E), \tag{2.7}$$

for two vector bundles E and F (of course, $\widetilde{C}$ is automatically compatible with the augmentations). Let D be a divisor on X and $L(D)$ the vector bundle it defines; we want [print 41]

$$C^1(L(D)) = \ell(D) \qquad C^i(L(D)) = 0 \quad (i > 1)^{[31]} \tag{2.8}$$

(where, for an arbitrary cycle Z, we denote by $\ell(Z)$ its class in $A(X)$). Finally, we require the C^i to be functorial, i.e. that, when f is a morphism from X to Y and E is a vector bundle on Y, we have

$$C^i(f^!(E)) = f^*(C^i(E)), \tag{2.9}$$

where $f^!(E)$ is the inverse image of E on X. We can also write (2.9) in the following form: the notion of inverse image of a bundle defines a λ-homomorphism of augmented λ-rings

$$f^! : \underline{K}(Y) \longrightarrow \underline{K}(X)(*) \tag{2.10}$$

(here we write $\underline{K}(X)$ instead of $\underline{K}(\xi(X))^{[32]}$), and (2.9) means that the diagram commutes:

$$\begin{array}{ccc} K(Y) & \xrightarrow{f^!} & K(X) \\ \downarrow{\scriptstyle \widetilde{C}_Y} & & \downarrow{\scriptstyle \widetilde{C}_X} \\ \widetilde{A}(Y)^{[33]} & \xrightarrow{f^*} & \widetilde{A}(X)^{[34]}. \end{array}$$

(*) The notation $f^!$ used here conflicts with that used in duality theory (cf. for example R. Hartshorne, Residues and Duality, Lecture Notes in Math. no. 20, Springer, 1966). This is why in this Seminar we replace it by the notation f^*.

(The space on which the symbol $\widetilde{C}$ is considered has been put as a subscript.)

Theorem 2.2. There exists a theory of Chern classes satisfying conditions (2.5), (2.7), (2.8) and (2.9).

We shall also admit the following proposition:

Proposition 2.3. Let Y be a closed subset of X and U its complement; if x is an element of $A(X)$ inducing 0 on U, there exists a representative of x that is a cycle supported on Y.

[print 42]**Corollary.** Let $p = \dim X - \dim Y$. Then, if $i < p$, the homomorphism $A^i(X) \longrightarrow A^i(U)$ is bijective. Every element of the kernel of $A^p(X) \longrightarrow A^p(U)$ is a linear combination of components of dimension $n - p$ of Y (and is therefore proportional to $\ell(Y)$ if Y is irreducible)

§2. Definition of the Chern classes of coherent algebraic sheaves

Let $\mathbb{C}$ be an abelian category, and let $\mathbb{C}_0$ be a subclass of $\mathbb{C}$. We denote by $K(\mathbb{C}_0)$ the quotient group of the free group generated by the classes (up to isomorphism) of elements of $\mathbb{C}_0$, by the subgroup generated by elements of the form $(E) - (E') - (E'')$, where E is an extension of E' by E'' ($E, E', E'' \in \mathbb{C}_0$). We must assume that the classes of elements of $\mathbb{C}_0$ for the isomorphism relation form a set. We denote by $\gamma_{\mathbb{C}_0}(E)$ the class of an $E \in \mathbb{C}_0$ in $K(\mathbb{C}_0)$, and omit the subscript $\mathbb{C}_0$ when there is no risk of confusion. The group $\underline{K}(\mathbb{C}_0)$ is the solution of a universal problem concerning functions $E \longrightarrow f(E)$ defined on $\mathbb{C}_0$, with values in an arbitrary abelian group A, which are additive in the sense that $f(E) = f(E') + f(E'')$ whenever E is an extension of E' by E''.

Let F be an additive functor from $\mathbb{C}$ to $\mathbb{C}'$, and $\mathbb{C}'_0$ a subclass of $\mathbb{C}'$ satisfying the same condition as $\mathbb{C}_0$, such that, for every $A \in \mathbb{C}_0$, the $R^qF(A)$ are in $\mathbb{C}'_0$ and are zero for sufficiently large q (we assume that injective resolutions exist in $\mathbb{C}$, so that the derived functors R^qF of F exist); then F defines a homomorphism from $\underline{K}(\mathbb{C}_0)$ to $\underline{K}(\mathbb{C}'_0)$, still denoted F, by the formula

$$F(\gamma_{\mathbb{C}_0}(E)) = \sum_{i \geq 0} (-1)^i \gamma_{\mathbb{C}'_0}(R^iF(E)). \tag{2.11}$$

More generally, when we have a cohomological functor (F^i) from $\mathbb{C}$ to $\mathbb{C}'$ [print 43]such that, for every $E \in \mathbb{C}_0$, the $F^i(E)$ are in $\mathbb{C}'_0$ and are zero except for finitely many, we deduce a homomorphism from $\underline{K}(\mathbb{C}_0)$ to $\underline{K}(\mathbb{C}'_0)$. These remarks extend to multifunctors; moreover, the "classical" spectral sequences give transitivity properties that we shall not make explicit.

In particular, when F is an exact functor, formula (2.11) simplifies:

$$F(\gamma_{\mathbb{C}_0}(E)) = \gamma_{\mathbb{C}'_0}(F(E)). \tag{2.11 bis}$$

This applies in particular when $\mathbb{C} = \mathbb{C}'$, when F is the identity functor, and when $\mathbb{C}_0 \subset \mathbb{C}'_0$: we have a canonical homomorphism from $\underline{K}(\mathbb{C}_0)$ to $\underline{K}(\mathbb{C}'_0)$.

Theorem 2.3. Let $\mathbb{C}$ be an abelian category and $\mathbb{C}_0 \subset \mathbb{C}_1$ two subclasses such that the classes (up to isomorphism) of objects of $\mathbb{C}_1$ form a set. We assume that the following conditions hold

(i) For every exact sequence $0 \to A' \to A \to A'' \to 0$ in $\mathbb{C}$ such that A and A'' are in $\mathbb{C}_0$ (resp. $\mathbb{C}_1$), we have $A' \in \mathbb{C}_0$ (resp. $A' \in \mathbb{C}_1$).

(ii) Let $u : A \to B$ and $v : C \to B$ with $A, B, C \in \mathbb{C}_1$ and u surjective; then there exist $L \in \mathbb{C}_0$ and homomorphisms $u' : L \to C$ and $v' : L \to A$ such that u' is surjective and $vu' = uv'$.

(iii) For every $A \in \mathbb{C}_1$, there exists an integer $d = d(A)$ such that, for every left resolution $\underline{L}$[35] of A by objects of $\mathbb{C}_0$, we have $Z_d(\underline{L}) \in \mathbb{C}_0$ (Z_d denotes the cycles of dimension d).

Under these conditions, the canonical homomorphism from $K(\mathbb{C}_0)$ to $K(\mathbb{C}_1)$ is an isomorphism.

Principle of the proof: let $A \in \mathbb{C}_1$; by (ii) and (iii) there exists a finite resolution $\underline{L}$ of A by objects of $\mathbb{C}_0$; put [print 44]then $f(A) = \sum_{i\geq 0}(-1)^i \gamma_{\mathbb{C}_0}(L^i)$. We show that the element of $\underline{K}(\mathbb{C}_0)$ thus constructed does not depend on the chosen resolution $\underline{L}$: given a second resolution $\underline{L}'$, we "cover" $\underline{L}$ and $\underline{L}'$ by a third resolution $\underline{L}''$, with surjective homomorphisms from $\underline{L}''$ to $\underline{L}$ and $\underline{L}'$ (which is possible by (i) and (ii));[36] we are then reduced to proving $\underline{E}_{\mathbb{C}_0}(\underline{L}) = \underline{E}_{\mathbb{C}_0}(\underline{L}'')$. Now the difference of the two is $\underline{E}_{\mathbb{C}_0}(\underline{N})$, where $\underline{N}$ is the kernel of the surjective homomorphism from $\underline{L}''$ onto $\underline{L}$, a kernel that is in $\mathbb{C}_0$ by (i). Now $\underline{N}$ is an acyclic complex in all dimensions, whence $\underline{E}_{\mathbb{C}_0}(\underline{N}) = 0$[37], whence the conclusion (N.B. $\underline{E}_{\mathbb{C}_0}(\underline{L})$, for a finite complex with values in $\mathbb{C}_0$, denotes the Euler–Poincaré characteristic $\sum_{i\geq 0}(-1)^i \gamma_{\mathbb{C}_0}(L^i)$ of $\underline{L}$).

We prove analogously that the function $f : \mathbb{C}_1 \longrightarrow \underline{K}(\mathbb{C}_0)$ thus constructed is additive, and therefore defines a homomorphism from $\underline{K}(\mathbb{C}_1)$ to $\underline{K}(\mathbb{C}_0)$, and it is immediate that this homomorphism is inverse to the canonical homomorphism from $\underline{K}(\mathbb{C}_0)$ to $\underline{K}(\mathbb{C}_1)$. q.e.d.

Remark: To verify (ii), it suffices to verify that $\mathbb{C}_1$ is contained in a subclass $\mathbb{C}_2$ of $\mathbb{C}$ satisfying the two conditions

(ii a) Every $A \in \mathbb{C}_2$ is isomorphic to a quotient of an $L \in \mathbb{C}_0$.

(ii b) If $u : A \to B$, with $A, B \in \mathbb{C}_2$, then $\operatorname{Ker} u \in \mathbb{C}_2$.

Corollary: Let X be a quasi-projective algebraic space. Let $\mathscr{F}(X)$ be the category of coherent algebraic sheaves on X, $\xi(X)$ the category of locally free coherent algebraic sheaves (equivalent to the category of vector bundles on X), and $\mathscr{F}_0(X)$ the category of coherent algebraic sheaves F such that, for every $x \in X$, F_x is an $\underline{O}_x$-module [print 45]of finite cohomological dimension (we therefore have $\xi(X) \subset \mathscr{F}_0(X) \subset \mathscr{F}(X)$). Then the canonical homomorphism

$$K(\xi(X)) \longrightarrow K(\mathscr{F}_0(X)) \tag{2.12}$$

is an isomorphism.

We are going to verify the conditions of theorem 2.3. Conditions (i) and (iii) follow from the definition of cohomological dimension and from the fact that a projective module of finite type over the noetherian local ring $\underline{O}_x$ is free. To verify (ii), we shall show that $\mathbb{C} = \mathscr{F}(X)$ satisfies conditions (ii a) and (ii b) of the remark. The second is trivial; it therefore remains to prove the following result:

Proposition 2.4. (Serre). Let X be a quasi-projective algebraic space. Then every coherent algebraic sheaf on X is isomorphic to the quotient of a locally free algebraic sheaf.

This proposition holds when X is a closed subset of projective space, by [2], since, for sufficiently large n, $F(n)$ is generated by its sections, which means that F is a quotient of $\underline{O}(-n)^k$. In the general case, X is an open subset of a closed subset of a projective space, and it suffices to use the following result (due to Cartier and Serre in particular cases):

Proposition 2.5. Let F be a coherent algebraic sheaf on an open subset U of an algebraic space X. Then F is isomorphic to the restriction of a coherent algebraic sheaf on X.

By applying Zorn's lemma to the open subsets containing U to which F can be extended, we are reduced to the case where X is affine. Let $\overline{F}$ be the alge[print 46]braic sheaf on X whose sections on an open subset V are the sections of F on $U \cap V$. We readily find that $\overline{F}$ is quasi-coherent (i.e. defined on every affine open subset V – hence here on all of X – by a module, of finite type or otherwise, over the coordinate ring of the open subset in question). This is immediate when U itself is affine, for $\overline{F}$ is then defined on the open subset V by $\Gamma(U \cap V, F)$ considered as a

module over $\Gamma(V, \underline{O}_X)$. In the general case, U is a finite union of affine open subsets U_i, so $\overline{F}$ is the kernel of an evident homomorphism of quasi-coherent sheaves:

$$\sum_i \overline{F|U_i} \longrightarrow \sum_{i,j} \overline{F|U_i \cap U_j}$$

and is therefore itself quasi-coherent. As X is assumed affine, $\overline{F}$ is therefore generated by its sections. As $\overline{F}|U = F$ is coherent, there exist finitely many sections of $\overline{F}$ generating $\overline{F}|U$. These sections generate a coherent subsheaf of $\overline{F}$, whose restriction to U is isomorphic to F.

q.e.d.

Now assume X quasi-projective and nonsingular. Then, by the syzygy theorem, we have, with the notation of the corollary to theorem 2.3, $\mathscr{F}_0(X) = \mathscr{F}(X)$. In this case we therefore have:

$$\underline{K}(\xi(X)) = \underline{K}(\mathscr{F}(X)). \tag{2.13}$$

We can express this isomorphism as follows:

Corollary 2 to theorem 2.3. Let X be a nonsingular quasi-projective variety. For every abelian group A, there is a one-to-one correspondence between additive functions f of vector bundles on X, with values in A, and additive functions g of coherent algebraic sheaves on X, with values in A. To every f corresponds the unique g such that we have: [print 47]

$$f(E) = g(\underline{O}_X(E)) \tag{2.14}$$

for every vector bundle E on X, denoting by $\underline{O}_X(E)$ the sheaf of germs of regular sections of E on X.

To calculate $g(F)$ for an arbitrary coherent algebraic sheaf, we consider a finite resolution of F by sheaves of the form $\underline{O}_X(E_i)$, and we then have:

$$g(F) = \sum_{i \geq 0} (-1)^i f(E_i). \tag{2.14 bis}$$

In particular, when X is connected, we considered in §1 an additive function $\widetilde{C}(E)$ of the vector bundle E, with values in the group $\widetilde{A} = \widetilde{A}(X)$. We deduce from it an additive function of coherent sheaves, say $\widetilde{C}(F)$, and, by passing to components, functions $F \longrightarrow C^i(F) \in A^i(X)$. The $C^i(F)$ are called the Chern classes of the coherent sheaf F; they are theoretically calculated from the Chern classes of vector bundles by (2.14 bis) (where, however, we must pass to multiplicative notation in the second member). As for the component of $\widetilde{C}(F)$ along $\mathbb{Z}$, it is the rank of the sheaf F, which corresponds to the rank of a module over an integral domain: when X is affine (a case to which we can always reduce), F is defined by a module of finite type over the coordinate ring of X, and we take the rank of this module....

§3. Functorial generalities on $\underline{K}(X)$

When X is an algebraic space, we put, for brevity, $\underline{K}(X) = \underline{K}(\mathscr{F}(X))$, and, when X is nonsingular and quasi-projective, we identify this group with $\underline{K}(\xi(X))$ by corollary 2 to theorem 2.3. As $\underline{K}(\xi(X))$ is [print 48]not only an abelian group, but a λ-ring, it follows by transport of structure that $\underline{K}(X)$ is also a λ-ring, and is therefore equipped with a multiplicative structure and maps λ^i. It is important to define these structures directly, without passing to vector bundles; this will be done below.

If E is a vector bundle on X, we denote by $\gamma(E)$ its class in $\underline{K}(X)$; if F is a coherent algebraic sheaf on X, $\gamma(F)$ likewise denotes[38] its class in $\underline{K}(X)$. If Y is an irreducible subvariety of X, we put $\gamma(Y) = \gamma(\underline{O}_Y)$, where $\underline{O}_Y$ denotes the sheaf of local rings of Y, considered as a coherent algebraic sheaf on X. We define the symbol $\gamma(Y)$ by linearity when Y is an arbitrary cycle on X. When there is a risk of confusion, we shall write γ_X for γ.

Let X be an arbitrary algebraic space. We introduce a filtration on $\underline{K}(X)$, denoting by $\underline{K}(X)^i$ the subgroup of $\underline{K}(X)$ generated by the $\gamma(F)$ with dim.supp. $F \leq n-i$ (where n is the dimension of X). From Serre's "dévissage lemma", in the form given in [3], follows the result:

Proposition 2.6. The group $\underline{K}(X)^i$ is generated by the $\gamma(Y)$, where Y runs through the set of subvarieties of X of dimension $\leq n-i$. If X is irreducible, we have $G^0(\underline{K}(X)) = \mathbb{Z}$.

The latter isomorphism is given by the homomorphism from $\underline{K}(X)$ to $\mathbb{Z}$ defined by means of the notion of rank of a coherent algebraic sheaf.

Now assume X nonsingular. Following the developments [print 49]at the beginning of §2, we define a multiplication on $\underline{K}(X)$ by putting

$$\gamma(F)\gamma(G) = \sum_{i\geq 0}(-1)^i\gamma(\underline{\mathrm{Tor}}_i(F,G)) \tag{2.15}$$

(the $\underline{\mathrm{Tor}}_i$ being taken, of course, with respect to the local rings of X). The exact sequence of the $\underline{\mathrm{Tor}}_i$ shows that formula (2.15) does define a biadditive map from $\underline{K}(X) \times \underline{K}(X)$ to $\underline{K}(X)$; associativity follows from the spectral sequence of multiple Tor. We also verify in this way that the product of several factors can be calculated by the formula:

$$\gamma(F_1)....\gamma(F_p) = \sum_{i\geq 0}(-1)^i\gamma(\underline{\mathrm{Tor}}_i(F_1,\ldots,F_p)) \tag{2.16}$$

where, in the second member, the $\underline{\mathrm{Tor}}_i$ are "simultaneous Tor", calculated by simultaneous projective resolution of the arguments F_i.

If, in (2.15), we assume G locally free, we find:

$$\gamma(F)\gamma(G) = \gamma(F \otimes G), \tag{2.15 bis}$$

which shows in particular that $\gamma(\underline{O}_X)$ is the unit element of $\underline{K}(X)$, and that the canonical homomorphism from $\underline{K}(\xi(X))$[39] to $\underline{K}(X)$ is a ring homomorphism.

Formula (2.15) is evidently suggested by Serre's "Tor formula" in intersection theory. This formula proves the second part of the following proposition, the first following immediately from formula (2.15) and a calculation of local $\underline{\mathrm{Tor}}_i$:

Proposition 2.7. Let X be a nonsingular variety, and Y and Z two cycles in X. If Y and Z meet transversely everywhere, we have:

$$\gamma(Y)\gamma(Z) = \gamma(Y.Z). \tag{2.17}$$

If Y and Z are homogeneous and have respective dimensions $n-i$ and $n-j$, and if they meet without an excess component, we then have: [print 50]

$$\gamma(Y)\gamma(Z) = \gamma(Y.Z) \qquad \text{mod. } \underline{K}(X)^{i+j+1}. \tag{2.18}$$

We must take care that equality does not hold in general in (2.18). We shall see in §4 that, when X is nonsingular and quasi-projective, the ring structure of $\underline{K}(X)$ is compatible with its filtration and that the associated graded ring $G(\underline{K}(X))$ is identified with a quotient of $A(X)$ by a torsion subgroup (perhaps always reduced to 0(*)).

(*) No; for a counterexample cf. this Seminar XIV 4.7.

I know how to define the $\lambda^i(F)$ directly in terms of sheaves only when the field k has characteristic 0 (this is the reason that has prevented us from proving the Riemann–Roch theorem when k is not of characteristic 0[40]). Let F be a coherent algebraic sheaf on X on which the group $\mathfrak{S}_p$ of permutations of p letters operates; we then denote by $(F)^{\text{alt}}$ the set of the $x \in F$ such that $s.x = \varepsilon(s)x$ for every $s \in \mathfrak{S}_p$ ($\varepsilon(s)$ is the signature of s). This is evidently a coherent algebraic subsheaf of F. This said, we have:

Theorem 2.8. Let X be a nonsingular quasi-projective variety over the field k of characteristic 0. Let F be a coherent algebraic sheaf on X; we then have:

$$\lambda^p(\gamma(F)) = \sum_{i\geq 0}(-1)^i\gamma\left((\underline{\mathrm{Tor}}_i(F,F,\ldots,F))^{\text{alt}}\right) \tag{2.19}$$

N.B. It is clear that the group $\mathfrak{S}_p$ acts on the sheaf $\underline{\mathrm{Tor}}_i(F,\ldots,F)$[41].

Principle of the proof: we consider a finite resolution $\underline{L}$ of F by locally free sheaves L_i, and use the following general lemma:

[print 51]**Lemma 2.9.** Let $\underline{L} = (L_i)$ be a graded locally free coherent algebraic sheaf, with finitely many nonzero components. In $\underline{K}(X)$ we have the formula:

$$\underline{E}\left((\overset{p}{\otimes}\underline{L})^{\text{alt}}\right) = \wedge^p(\underline{E}(\underline{L})). \tag{2.20}$$

In this formula, we consider $\overset{p}{\otimes}\underline{L}$ as a sheaf on which the group $\mathfrak{S}_p$ operates, taking the gradings into account (which therefore introduce signs in the well-known way, cf. Cartan–Eilenberg); moreover, if $\underline{M}$ is a graded coherent algebraic sheaf on X, we put $\underline{E}(\underline{M}) = \sum_{i\geq 0}(-1)^i\gamma(M_i)$.

To prove formula (2.20), we are reduced to proving the analogous formula in $\underline{K}(G)$, where G is an algebraic group (a product of groups $\mathrm{Gl}(n,k)$, if desired) and $\underline{L}$ a G-module of finite dimension over k (cf. Chap. I, §2, example 1, for the definition[42] of $\underline{K}(G)$). In this case, this formula has already been used in theorem 1.4 to show that the second member of (1.33) satisfies the same functional equation as the first member; its verification is elementary.

q.e.d.

Let f be a morphism from X to Y, where Y is an algebraic space. The notion of inverse image of a vector bundle defines a homomorphism $f^! : \underline{K}(\xi(Y)) \longrightarrow \underline{K}(\xi(X))$ of λ-rings. Identifying vector bundles with locally free algebraic sheaves, the notion of inverse image of a vector bundle corresponds to the notion of inverse image of an algebraic sheaf (already used in [3, no. 8]). When Y is nonsingular, we define more generally a group homomorphism:

$$f^! : \underline{K}(Y) \longrightarrow \underline{K}(X) \tag{2.21}$$

[print 52]by a formula involving an alternating sum of Tor, inspired by the developments at the beginning of §2, which the reader will make explicit. The following diagram then commutes:

$$\begin{array}{ccc} \underline{K}(\xi(Y)) & \xrightarrow{f^!} & \underline{K}(\xi(X)) \\ \downarrow & & \downarrow \\ \underline{K}(Y) & \xrightarrow{f^!} & \underline{K}(X), \end{array}$$

which shows in particular, when X and Y are both nonsingular and quasi-projective, that the two definitions of $f^!$ coincide under the identification of $\underline{K}(X)$ with $\underline{K}(\xi(X))$ and of $\underline{K}(Y)$ with $\underline{K}(\xi(Y))$.

Another case in which a homomorphism (2.21) can be defined, without Y necessarily being nonsingular, is that in which the functor $F \to f^!(F)$ from $\mathscr{F}(Y)$ to $\mathscr{F}(X)$ is exact, i.e. when the local rings of X are flat modules over the local rings of Y. It then suffices to put $f^!(\gamma_Y(F)) = \gamma_X(f^!(F))$. This case occurs in particular when X is a locally trivial fibre space over Y (to see this, we are reduced to the case of a product $X = T \times Y$). Note that we also verify that, in this case, we have $f^!(\underline{O}_Z) = \underline{O}_{f^{-1}(Z)}$ if Z is an irreducible subvariety of Y, whence:

$$f^!(\gamma_Y(Z)) = \gamma_X(f^{-1}(Z)) \tag{2.22}$$

if X is "flat" over Y. Without this last condition, formula (2.22) is in general incorrect, but we have the following analogue of proposition 2.7:

Proposition 2.8. Let $f : X \longrightarrow Y$ be a morphism of nonsingular varieties, and Z a cycle on Y. If Z is everywhere transverse to f, formula [print 53](2.22) is exact; if Z is homogeneous of codimension i and $f^{-1}(Z)$ has no excess component, we have:

$$f^!(\gamma_Y(Z)) \equiv \gamma_X(f^{-1}(Z)). \qquad \text{mod. } \underline{K}(X)^{i+1} \tag{2.23}$$

Proof analogous to that of proposition 2.7.

Finally, let us mention that we shall see in §4 that, when X and Y are quasi-projective and nonsingular, $f^!$ is compatible with the filtrations of $\underline{K}(X)$ and of $\underline{K}(Y)$.

Again let X and Y be two arbitrary algebraic spaces, and f a proper morphism from X to Y. For every coherent algebraic sheaf F on X, the direct image sheaf $f_!(F)$ is coherent algebraic, and the same holds more generally for the sheaves $R^q f_!(F)$ (cf. [3]). Recall that, by definition, $R^q f_!(F)$ is the sheaf on Y associated with the presheaf assigning to an open subset U the group $H^q(f^{-1}(U), F)$; in particular, we have:

$$R^0 f_!(F) = f_!(F), \qquad R^q f_!(F) = 0 \quad (q > \dim X), \tag{2.24}$$

moreover:

$$\Gamma(U, R^q f_!(F)) = H^q(f^{-1}(U), F) \tag{2.25}$$

if the open subset U is affine. By §2, we define a homomorphism from $\underline{K}(X)$ to $\underline{K}(Y)$, still denoted $f_!$, by the formula

$$f_!(\gamma_X(F)) = \sum_{q \geq 0} (-1)^q \gamma_Y(R^q f_!(F)). \tag{2.26}$$

Thus $\underline{K}(X)$ becomes a covariant functor in X, with respect to proper morphisms. The relation $(fg)_! = f_! g_!$[43] (just like the relation $(fg)^! = g^! f^!$ that we forgot to mention in its place) is a parti[print 54]cular case of the transitivity relations to which we alluded in §2, which follow from the spectral sequences of composite functors.

Let us note the following formula, valid for Y nonsingular (and particularly easy to verify when Y is also quasi-projective, by taking for y the class of a vector bundle on Y):

$$f_!(x.f^!(y)) = f_!(x).y \qquad (x \in \underline{K}(X)^{[44]},\ y \in \underline{K}(Y)). \tag{2.27}$$

In the particular case where the inverse images under f of the points of Y are finite and contained in an affine open subset(*), we find that the functor $F \longrightarrow f_!(F)$ is exact and that $R^q f_!(F) = 0$ for $q > 0$; in this case, we therefore have $f_!(\gamma_X(F)) = \gamma_Y(f_!(F))$. In particular, if X is an algebraic subspace of Y and i is the canonical injection from X to Y[45], we have:

$$i_!(\gamma_X(F)) = \gamma_Y(F), \tag{2.28}$$

where, in the second member, F is considered as a sheaf on Y that is zero outside X.

(*) This last condition is evidently a consequence of the first (something the author still did not know in 1957!).

§4. Some technical results

Let X be an algebraic space; we saw in §3 that the projection p from $X \times k$ onto X defines a homomorphism

$$p^! : \underline{K}(X) \longrightarrow \underline{K}(X \times k). \tag{2.29}$$

Proposition 2.9. The homomorphism (2.29) is surjective.

We argue by induction on $n = \dim X$; the assertion is trivial if $n < 0$; assume $n \geq 0$; by proposition 2.6, it suffices to [print 55]verify that, for every irreducible closed subset Y of $X \times k$, $\gamma(Y)$ is in the image of $p^!$. It suffices to continue the reasoning assuming X irreducible; if $p(Y)$ is not dense in X, our assertion follows from the induction hypothesis; otherwise, either $Y = X \times k$ and the assertion is trivial, or Y is a hypersurface in $X \times k$. Now let U be an affine open subset of X consisting of nonsingular points; if D is the divisor on $U \times k$ induced by Y, we know (commutative algebra) that it is linearly equivalent to a divisor of the form $D' \times k$, where D' is a divisor on U. Then $\gamma_{U\times k}(D) - \gamma_{U\times k}(D' \times k)$ belongs to $\underline{K}(U \times k)^2$ (corollary of proposition 2.10), and, by what we have already proved, this element is of the form $p^!(x')$ with $x' \in \underline{K}(U)$. Thus $\gamma_{U\times k}(D) = p^!(y')$ with $y' = x' + \gamma_U(D')$; but y' is the restriction of an element y of $\underline{K}(X)$ (prop. 2.10), and consequently the restriction of $\gamma_{X\times k}(Y) - p^!(y)$[46] to $U \times k$ is zero; by proposition 2.10, $\gamma_{X\times k}(Y) - p^!(y)$[47] comes from an element of $\underline{K}(X \times k - U \times k) = \underline{K}(X' \times k)$ with $X' = X - U$. The proof is completed by applying the induction hypothesis again.

q.e.d.

Corollary: Let X be an algebraic space and p the projection from $X \times k^n$ onto X; then the homomorphism $p^! : \underline{K}(X) \longrightarrow \underline{K}(X \times k^n)$ is surjective; it is even bijective when X is nonsingular.

Induction on n for the first assertion. For the second, we put $f(x) = (x, 0)$, whence $pf = \underline{1}$ and $f^! p^! = 1$, which proves that $p^!$ is injective.

Proposition 2.10. Let X be an algebraic space, U an open subset of X, and Y its complement in X. Then the sequence of natural homomorphisms: [print 56]

$$\underline{K}(Y) \longrightarrow \underline{K}(X) \longrightarrow \underline{K}(U) \longrightarrow 0 \tag{2.30}$$

is exact.

N.B. Compare with proposition 2.3.

The fact that $\underline{K}(X) \longrightarrow \underline{K}(U)$ is surjective follows immediately from proposition 2.5 (or also from proposition 2.6). It remains to prove that, if $x \in \underline{K}(X)$ has zero restriction to $\underline{K}(U)$, it is in the image of $\underline{K}(Y)$. For this, we return to the explicit definition of $\underline{K}(X)$ and $\underline{K}(U)$: we have two coherent algebraic sheaves F and G on X with $x = \gamma_X(F) - \gamma_X(G)$ and $\gamma_U(F) - \gamma_U(G) = 0$; this last relation means that we can find coherent sheaves P' and Q' on U, equipped with filtrations whose factors are pairwise isomorphic on U, and such that we have an isomorphism on U:

$$P = F + Q' \longrightarrow Q = G + P'.$$

By proposition 2.5, the sheaves P' and Q' are restrictions to U of coherent sheaves on X (denoted by the same letters); likewise, their filtrations come from filtrations on the corresponding sheaves on X, by lemma 2.11 below. We then have:

$$\gamma_X(F) - \gamma_X(G) = (\gamma_X(P) - \gamma_X(Q)) + (\gamma_X(P') - \gamma_X(Q')).$$

Introducing the factors P'_i and Q'_j of the filtrations of P' and Q' on X, the last term is a sum of terms $\gamma_X(P'_i) - \gamma_X(Q'_i)$, where $P'_i|U \simeq Q'_i|U$. We are thus reduced to proving the

following particular case: if R and S are coherent sheaves on X such that $R|U \simeq S|U$, then $\gamma_X(R) - \gamma_X(S)$ is in the image of $\underline{K}(Y)$. Now let T be the graph of the given isomorphism from $R|U$ onto $S|U$; it is a coherent subsheaf of $(R \times S)|U$, and therefore, by lemma 2.11, it is the restriction to U of a coherent subsheaf, [print 57]again denoted T, of $R \times S$. We then have $\gamma_X(R) - \gamma_X(S) = (\gamma_X(R) - \gamma_X(T)) - (\gamma_X(S) - \gamma_X(T))$, and it suffices to prove, for example, that the first term of the second member is in the image of $\underline{K}(Y)$. Now the projection $u : T \longrightarrow R$ is an isomorphism over U, so $\gamma_X(R) - \gamma_X(T) = \gamma_X(\mathrm{Coker}\, u) - \gamma_X(\mathrm{Ker}\, u)$, and $\mathrm{Ker}\, u$ and $\mathrm{Coker}\, u$ have their support in Y.

It therefore remains to prove the following lemma:

Lemma 2.11. Let X be an algebraic space, F a coherent algebraic sheaf on X, U an open subset of X, and G a coherent subsheaf of $F|U$. Then G is the restriction to U of a coherent subsheaf of F.

Let $\overline{G}$ be the subsheaf of F whose sections on the open subset V of X are the sections of F on V whose restriction to $U \cap V$ is a section of G[48]. Everything reduces to proving that this sheaf $\overline{G}$ is coherent. For this, we can assume X and U affine (for U is a union of affine open subsets U_i, and $\overline{G}$ is then the intersection of the analogous sheaves $\overline{G|U_i}$). Then the coordinate ring of U is a ring of fractions A_S of the coordinate ring A of X, and we verify immediately that, if F is defined by the A-module M, then $F|U$ is defined by the A_S-module $M_S = A_S \otimes_A M$. The subsheaf G of $F|U$ is then defined by a submodule N' of M_S, and we see that $\overline{G}$ is precisely defined by the submodule N of M that is the inverse image of N' under the canonical homomorphism from M to M_S.

q.e.d.

Corollary 1 to proposition 2.10: Let $x \in \underline{K}(X)$. For x to have filtration $\geq i$[49], it is necessary and sufficient that there exist a closed subset Y of X, of dimension $\leq n - i$, such that the restriction of x to $X - Y$ is zero.

[print 58]**Corollary 2.** Assume X connected of dimension n, without singularities in dimension $n - 1$. If D and D' are two linearly equivalent divisors on X, we have $\gamma(D) \equiv \gamma(D') \mod \underline{K}(X)^2$.

By corollary 1, we can assume X nonsingular; the corollary then follows from the general formula:

$$\gamma(D) = 1 - \gamma(L(-D)) \qquad \mod \underline{K}(X)^2, \tag{2.31}$$

where $L(-D)$ denotes the vector bundle of rank 1 defined by the divisor $-D$. To verify (2.31), we write it in the form:

$$\gamma(L(-D)) = 1 - \gamma(D) \qquad \mod \underline{K}(X)^2, \tag{2.31 bis}$$

and note that both members are homomorphisms from the group of divisors to the multiplicative group of invertible elements of the ring $\underline{K}(X)/\underline{K}(X)^2$ (note that $\underline{K}(X)^1/\underline{K}(X)^2$ has square zero, which is very easily verified from the explicit formula (2.15), using corollary 1). We are thus reduced to the case where D is an irreducible hypersurface. More generally, if D is a hypersurface whose components are disjoint, we have the equality:

$$\gamma(D) = 1 - \gamma(L(-D)) \tag{2.32}$$

as follows from the exact sequence of sheaves

$$0 \longrightarrow \underline{O}_X(L(-D)) \longrightarrow \underline{O}_X \longrightarrow \underline{O}_D \longrightarrow 0.$$

Theorem 2.12. Let X be a nonsingular quasi-projective variety, and Z and Z' two homogeneous cycles of codimension p on X that are linearly equivalent. We then have:

$$\gamma_X(Z) \equiv \gamma_X(Z') \qquad \text{mod } \underline{K}(X)^{p+1}. \tag{2.33}$$

By hypothesis, we can find a cycle Y of dimension $n-p+1$ on [print 59]$X \times k$ (n is the dimension of X), and two points $a, a' \in k$ such that we have:

$$Z = f_a^{-1}(Y), \qquad Z' = f_{a'}^{-1}(Y),$$

where we put $f_u(x) = (x, u)$. By formula (2.23), we then have:

$$\gamma_X(Z) \equiv f_a^!(\gamma_{X\times k}(Y)) \qquad \text{mod } \underline{K}(X)^{p+1}$$

and an analogous formula for $\gamma_X(Z')$. It then suffices to prove that $f_a^! = f_{a'}^{![50]}$, which follows immediately from proposition 2.9.

q.e.d.

Corollary 1: The filtration of $\underline{K}(X)$ is compatible with its ring structure, i.e. we have $\underline{K}(X)^i\underline{K}(X)^j \subset \underline{K}(X)^{i+j}$ for all integers $i, j \geq 0$. Moreover, we have a natural surjective homomorphism of graded rings:

$$\varphi : A(X) \longrightarrow G(\underline{K}(X))^{[51]} \tag{2.34}$$

(where the second member denotes the graded ring associated with $\underline{K}(X)$ equipped with the filtration $(\underline{K}(X)^i)_{i\geq 0}$) defined by the condition:

$$\varphi(Z^{n-p}) = \gamma(Z^{n-p}) \qquad \text{mod } \underline{K}(X)^{p+1}.$$

This corollary follows immediately from theorem 2.12, propositions 2.6 and 2.7, and Chow's theorem 2.1, by induction on the integer $i + j$.

Corollary 2: Let X and Y be two nonsingular quasi-projective varieties and f a morphism from X to Y. Then the homomorphism $f^!$ from $\underline{K}(Y)$ to $\underline{K}(X)^{[52]}$ is compatible with the filtrations, and we have a commutative diagram:

$$\begin{array}{ccc} G(\underline{K}(Y)) & \xrightarrow{G(f^!)} & G(\underline{K}(X)) \\ \uparrow{\scriptstyle \varphi_Y} & & \uparrow{\scriptstyle \varphi_X} \\ A(Y) & \xrightarrow{f^*} & A(X). \end{array}$$

[print 60]Using the graph of f, we decompose f into the product of an immersion of X into $X \times Y$ and a projection from $X \times Y$ onto Y. For the latter the corollary is trivial, and for an immersion we proceed as for corollary 1, using proposition 2.8 in place of proposition 2.7.

Proposition 2.13 ("cellular decomposition"): Let P be an algebraic space (irreducible or not), equipped with an increasing family $P^0 \subset P^1 \subset \cdots \subset P^n = P$ of closed subsets such that, for every i with $0 \leq i \leq n$, the connected components of $P^i - P^{i-1}$ are algebraic spaces $V_{i,\alpha}$ isomorphic to k^i (put $P^{-1} = \varnothing$). Then:

a) For every algebraic space X, the natural map from $\underline{K}(X) \otimes \underline{K}(P)$ to $\underline{K}(X \times P)$ deduced from the tensor product of sheaves on X and P is surjective.

b) The $\gamma_P(V_{i,\alpha})$ form a system of generators of the group $\underline{K}(P)$.

The proof proceeds by induction on n, using the corollary of proposition 2.9 and proposition 2.10. (N.B. Unfortunately, I do not know whether the $\gamma_P(V_{i,\alpha})$ form a basis[53] of $\underline{K}(P)$ and whether the homomorphism considered in a) is bijective. We shall not need to know this in what follows).

Theorem 2.14. Let x_i $(1 \leq i \leq q)$ and E_j $(1 \leq j \leq r)$ be letters, and n_j $(1 \leq j \leq r)$ integers > 0; consider the symbols $\lambda^k(x_i)$ with $k \geq 1$ and $\lambda^k(E_j)$ with $1 \leq k \leq n_j$ as indeterminates, and let R be a polynomial with integer coefficients in these indeterminates. Assume that, for every graded ring A, when in R we substitute for the x_i elements ξ_i and for the E_j elements η_j of [print 61]$\widetilde{A}$ subject to the restrictions $\varepsilon(\xi_i) = 0$ and $\eta_j = [n_j, 1 + \eta_j^{(1)} + \cdots + \eta_j^{(n_j)}$[54]] (thus $\eta_j^{(k)} = 0$ for $k > n_j$), we have:

$$R((\lambda^k(\xi_i)), (\lambda^k(\eta_j))) \in \widetilde{A}^p$$

Now let X be an irreducible nonsingular quasi-projective variety. For arbitrary elements x'_i of $\underline{K}(X)$[1] and vector bundles E'_j of dimension n_j on X $(1 \leq i \leq q,\ 1 \leq j \leq r)$, we have

$$R((\lambda^k(x'_i)), (\lambda^k(\gamma(E'_j))))^{[55]} \in \underline{K}(X)^p \qquad (*)\ .$$

Proof: By §2, the x'_i themselves are differences of elements of the form $\gamma(E)$, where E is a vector bundle. On the other hand, assuming X locally closed in a projective space P, it follows from §2 and proposition 2.5 that, for every $F \in \mathscr{F}(X)$, the sheaf $F(n)$ is generated by its sections for sufficiently large n. If F is defined by a vector bundle E, this proves that $E(n)$ is the inverse image of the canonical vector bundle on a Grassmannian variety G under a morphism from X to G. Considering the corresponding morphism from X to $P \times G$, we see that E is the inverse image of a vector bundle on $P \times G$. Thus, applying this to all the vector bundles involved, and using corollary 2 of theorem 2.12, we see that we are reduced to the case where X is a product of Grassmannians. In this case, the conclusion will follow from the following lemma:

Lemma 2.15. Assume that X is a product of Grassmannians. Then we can find a λ-homomorphism from $\underline{K}(X)$ to a λ-ring without torsion of the form [print 62]$\widetilde{A}$ (where A is a graded ring), compatible with the filtrations, and such that the corresponding homomorphism $G(\underline{K}(X)) \longrightarrow G(A)$[57] is injective.

To prove the lemma, we shall use Chow's theory of Chern classes; a direct proof of the lemma, or of theorem 2.14, would yield a purely "sheaf-theoretic" theory of Chern classes (cf. §5).

We shall consider the homomorphism $\widetilde{C}$ from $\underline{K}(X)$ to $\widetilde{A}(X)$ and put $A = A(X)$; we must prove that the homomorphism $\psi = G(\widetilde{C})$ from $G(\underline{K}(X))$ to $G(A(X)) \simeq A(X)$ is injective. The fact that $\widetilde{C}$ is compatible with the filtrations holds for every nonsingular quasi-projective variety and follows readily from the corollary to proposition 2.3. On the other hand, as X is a product of Grassmannians C_j, we have on X vector bundles E_j, inverse images of the canonical bundles on the factors. We shall use the following (known) facts:

a) On X, linear equivalence of cycles is equivalent to numerical equivalence, and $A(X)$ is a group of finite rank;

b) The $C^i(E_j)$ generate the ring $A(X)$.

(*) We readily prove that the hypothesis made implies that, if R is expressed in terms of the $\gamma^k(\xi_i)$, $\gamma^k(E_j - \varepsilon(E_j))$[56], then all terms of R have weight $\geq p$ (where γ^k has weight k). Then 2.14 is a particular case of X 1.3.2.

From a) it follows that $\varphi : A(X) \longrightarrow G(\underline{K}(X))$ is injective (hence bijective), by the following lemma:

Lemma 2.15. Let X^n be a nonsingular projective variety, and Z^{n-p} a cycle on X such that $\varphi(Z) \in G^p(\underline{K}(X))$ is zero; then Z^{n-p} is numerically equivalent to zero.

To prove this lemma, it suffices to use the additive function $\chi(F) = \sum_{i\geq 0}(-1)^i \dim H^i(X, F)$ and the fact that this function equals [print 63]1 for $F = \underline{O}_{(x)}$ for every $x \in X$.

Thus, in the case considered, $A(X)$ and $G(\underline{K}(X))$ have the same finite rank in every degree, and $G(\underline{K}(X))$ has no torsion, since this holds for $\widetilde{A(X)}$. From assertion b) above it follows, in a "formal" way, that the map $\widetilde{C}$ from $\underline{K}(X)$ to $\widetilde{A(X)}$ is surjective, modulo finite groups. It readily follows, since $G(\underline{K}(X))$ and $A(X)$ have the same finite rank in every degree, that the homomorphism $G(\widetilde{C})$ from $G(\underline{K}(X))$ to $G(\widetilde{A(X)})$ is bijective modulo finite groups in every degree (use induction on the degree). As $G(\underline{K}(X))$ has no torsion, this homomorphism $G(\widetilde{C})$ is therefore injective.

q.e.d.

N.B. The homomorphism $\widetilde{C}$ is not surjective in general, even when X is a projective space of dimension 2.

§5. Sheaf-theoretic definition of Chern classes. Application to the study of injection morphisms.

Theorem 2.16. Let X be a nonsingular quasi-projective variety. Then, for every $x \in \underline{K}(X)$[58] and every integer $i \geq 0$, the element $\gamma^i(x - \varepsilon(x))$ has filtration $\geq i$. Let $c^i(x) \in G^i(\underline{K}(X))$ be the class of this element modulo $\underline{K}(X)^{i+1}$; if we put:

$$\widetilde{c}(x) = \left[\varepsilon(x), 1 + \sum_{i\geq 1} c^i(x)^{[59]}\right] \in G(\widetilde{\underline{K}(X)}),$$

then $\widetilde{c}$ satisfies the conditions analogous to those of theorem 2.2.

The first assertion follows from theorem 2.14, as does the fact that $x \longrightarrow \widetilde{c}(x)$ is a homomorphism of λ-rings (formula (1.25) implies $\gamma_t(x+y) = \gamma_t(x)\gamma_t(y)$, which already shows that $\widetilde{c}$ is an additive homomorphism). The "functoriality" of $\widetilde{c}$ is verified immediately, and [print 64](2.8) follows from (2.31).

q.e.d.

Corollary 1: We have $c^i(x) = \varphi(C^i(x))$.

This follows from a uniqueness theorem for Chern classes, proved as in Hirzebruch by passage to flag varieties. We need only show that, if Y is a flag variety associated with a vector bundle on X, then $G(\underline{K}(X)) \longrightarrow G(\underline{K}(Y))$ is injective, which is verified by explicitly determining $\underline{K}(Y)$ with all its structures from $\underline{K}(X)$ (I shall not give the details here).

Corollary 2: Assume that, for every element $Z \in A(X)$, there exists a morphism f from X to a product X' of Grassmannians such that Z is in the image of f^*. Then the canonical homomorphisms

$$\varphi : A(X) \longrightarrow G(\underline{K}(X)), \qquad \psi : G(\underline{K}(X)) \longrightarrow A(X),$$

satisfy in dimension i the following relations:

$$\psi\varphi = (-1)^{i-1}(i-1)!\,\underline{1}, \qquad \varphi\psi = (-1)^{i-1}(i-1)!\,\underline{1}. \tag{2.35}$$

As φ is surjective, it evidently suffices to prove the first formula, and for this we can assume X a product of Grassmannians. As φ is then injective, it suffices to prove the second formula (2.35), which, by corollary 1, is written

$$\gamma^i(x - \varepsilon(x))^{[60]} \equiv (-1)^{i-1}(i-1)!\,x \qquad \mod \underline{K}(X)^{i+1}$$

for every $x \in \underline{K}(X)^i$. As ψ is injective, it suffices to prove the analogous relation for $\widetilde{C}(x) \in \widetilde{A}(X)^{i\,[61]}$, and this is then a consequence of (1.22 bis).

Remark: Corollary 2 makes the validity of (2.35) in all cases plausible; there should be a direct proof of it. In [print 65]any case, we readily prove the second formula (2.35) modulo the torsion subgroup of $G(\underline{K}(X))$, and, in characteristic 0, this same formula is a consequence of the Riemann–Roch theorem. The interest of (2.35) (when it holds) is that it shows that φ is injective modulo torsion groups (perhaps always zero!(*)), so that we "do not lose much" by defining the Chern classes as elements of $G(\underline{K}(X))$; after taking the tensor product with $\mathbb{Q}$, as we shall do later for the Riemann–Roch theorem, the two definitions of Chern classes would then be identical.

Let X still be nonsingular and quasi-projective; let also Y be a nonsingular subvariety of X, and i the injection of Y into X. For $i^!$ and $i_!$, we then have the following formulas:

(i) $i^!$ is a homomorphism of λ-rings

(ii) $i_!(x.i^!(y)) = i_!(x).y$, whence in particular $i_!i^!(y) = \xi y$ with $\xi = i_!i^!(1) = \gamma_X(Y)$,

(iii) $i^!(i_!(x)) = x.\lambda_{-1}(N)$, whence in particular $i^!(\xi) = i^!i_!(1) = \lambda_{-1}(N)$,

(iv) $i_!(x)i_!(x') = i_!(xx'\lambda_{-1}(N))^{[62]}$,

where, by definition, N is the class in $\underline{K}(Y)$ of the conormal bundle of Y in X. Formulas (i) and (ii) hold for arbitrary morphisms, not necessarily injections, and (iii) and (iv) are verified from definition (2.15) and the analogous formula for inverse images (taking for x and x' classes of vector bundles, and using the fact that

$$\mathrm{Tor}_i^{O_X}(\underline{O}_Y, \underline{O}_Y) = \underline{O}_Y(\wedge^i N). \tag{2.36}$$

) [print 66]Formula (iv) expresses that $i_!$ is a ring homomorphism when we introduce on $\underline{K}(Y)$ a new multiplicative structure, by the definition $x_N x' = xx'\lambda_{-1}(N)$ (cf. Chapter I, §4). It is completed by the formula:

(v) $i_!(\lambda^p(N, x)) = \lambda^p(i_!(x))$ for $p \geq 1$,

which means that $i_!$ defines a λ-homomorphism from $\underline{K}(Y)_N$ to $\underline{K}(X)$. This formula (v), certainly true in all cases, has for the moment been proved only in characteristic 0, since it follows from the conjunction of theorems 1.4 and 2.8.

We have analogous formulas for i_* and i^*:

(i bis) i^* is a homomorphism of graded rings.

(*) No, cf. the note on page 30.

(ii bis) $i_*(x.i^*(y)) = i_*(x).y$, whence in particular $i_*(i^*(y)) = \xi' y$, with $\xi' = i_*(1)$,

(iii bis) $i^*(i_*(x)) = x.(-1)^q N^q$, whence in particular $i^*(\xi') = (-1)^q N^q$,

(iv bis) $i_*(x)i_*(x') = i_*(xx'(-1)^q N^q)$,

(v bis) i_* increases degrees by q units.

In these formulas, q is the codimension of Y in X, and we put $N^i = C^i(N)$. Only formulas (iii bis) and (iv bis) require a proof; I do not know whether they are well known (or even whether they are true!!!), and merely note that they follow by reduction of formulas (iii) and (iv)[63], provided that we work in $G(\underline{K}(Y))$ and $G(\underline{K}(X))$ and not in $A(Y)$ and $A(X)$. We then use the fact that i_* and i^* arise by passage to the quotient from $i_!$ and $i^!$, noting that $i_!$ shifts the filtration by q units.

[print 67]N.B. All the "universal" formulas relating $i_!$ and $i^!$ (resp. i_* and i^*) are formal consequences of the preceding ones, as one can convince oneself without much difficulty.

From formula (v) follows the formula

$$i_!(\gamma^p(N,x)) = \gamma^p(i_!(x)).$$

The conjunction of theorem 2.14 and proposition 1.5 shows that $\gamma^p(N,x)$ has, for $p = q+j$, filtration $\geq j$; denoting by $G_p(N,x)$ its class in $G^j(\underline{K}(Y))$, theorem 2.16 shows that we have

$$i_*(G_p(N,x)) = c^p(i_!(x)).$$

Finally, proposition 1.5 together with theorem 2.14 allows us to express $G_p(N,x)$ as a polynomial with integer coefficients in the $c^i(N)$, the $c^i(x)$ and $\varepsilon(x)$; we deduce the theorem:

Theorem 2.17. Assume that the field k has characteristic 0 and let $i : Y \longrightarrow X$ be an injection morphism of nonsingular quasi-projective varieties. We then have formulas (i) to (v), and, moreover, for every integer $j \geq 0$, the formula:

$$c^{q+j}(i_!(x)) = i_*\left(\frac{\widetilde{c}(x) * \lambda_{-1}(\widetilde{c}(N))}{(-1)^q N^q}\right)^{(j)}. \tag{2.37}$$

The right-hand side of formula (2.37) is an abbreviated way of denoting the polynomial in the components of $\widetilde{c}(x)$ and $\widetilde{c}(N)$ described in proposition 1.5.

Corollary: Under the preceding conditions, we have:

$$\mathrm{ch}(\widetilde{c}(i_!(x))) = i_*\left(\mathrm{ch}(\widetilde{c}(x))\,\mathcal{T}(\check{N})^{-1}\right) \tag{2.38}$$

where ch is the Chern homomorphism defined by formula (1.26).

[print 68]It suffices to use (2.37) written in the form:

$$\widetilde{c}(i_!(x)) = 1 + i_*\left(\frac{\widetilde{c}(x) * \lambda_{-1}(\widetilde{c}(N))^{[64]} - 1}{(-1)^q N^q}\right), \tag{2.37 bis}$$

the explicit formula (1.29), formula (iv bis), and finally (1.30).

We can write (2.38) in the form

$$\mathrm{ch}_X(F) = i_*\left(\mathrm{ch}_Y(F)\,\mathcal{T}(T_{X/Y})^{-1}\right), \tag{2.38 bis}$$

where F is a coherent algebraic sheaf on Y (considered on the left-hand side as a sheaf on X zero outside Y), ch_X and ch_Y are the Chern homomorphisms taken respectively on X and Y, and $T_{X/Y}$ is the normal bundle of Y in X.

Remarks: Unlike (2.38), formula (2.37) is a "formula without denominators". It has been proved (even in characteristic 0) only by working in $G(\underline{K}(X))$ and not in $A(X)$, which may not be the same thing. It is certainly true in every characteristic and in $A(X)$. Let us note that we immediately verify that the images of the two sides under i^* are the same, whence it follows, on applying i_*, that the product of their difference by ξ'[65] is zero. This suggests a completely different method of proof, by trying to obtain Y as the transverse inverse image of a subvariety Y' in a variety X' such that, in dimension $\leq n = \dim X$, the class of Y' in $A(X')$ is not a divisor of 0.... Unfortunately, we cannot hope that X' is a Grassmannian or something close to one.

We shall note that, if we disregard torsion elements, formula (2.38) is equivalent to (2.37).

[print 69]

§6. The Riemann–Roch theorem

Its statement is as follows: let f be a proper morphism from a nonsingular quasi-projective variety Y to a variety X of the same kind. Then we have:

$$f_*(\mathrm{ch}_Y(x)\,\mathcal{T}(Y)) = \mathrm{ch}_X(f_!(x))^{[66]}\,\mathcal{T}(X). \tag{2.39}$$

(Both sides are in $A(X)\otimes\mathbb{Q}$ and $\mathcal{T}(X) = \mathcal{T}(C(T_X))$.)

It allows us to calculate, up to torsion elements, the Chern classes of $f_!(x)$ from the Chern classes of x, those of T_X and T_Y, and the homomorphism f_*. Or, again, it allows us to determine the homomorphism $f_!$ from $\underline{K}(Y)\otimes\mathbb{Q}$ to $\underline{K}(X)\otimes\mathbb{Q}$, when we identify these groups by means of the homomorphisms ch_Y and ch_X respectively with $A(Y)\otimes\mathbb{Q}$ and $A(X)\otimes\mathbb{Q}$ (if formula (2.35) is true!), using f_* and the Chern classes of T_X and T_Y.

I know how to prove formula (2.39) only in characteristic 0(*), and as an equality in $G(\underline{K}(X))\otimes\mathbb{Q}$. For this, since Y is locally closed in a projective space P, we decompose f into the product of an injection morphism from Y to $P\times X$, and a projection morphism from $P\times X$ onto X, and prove (2.39) for each of these two morphisms.[67]

For an injection morphism, we see immediately that formulas (2.38) and (2.39) are equivalent (the characteristic 0 hypothesis entered only through theorem 2.8, which allows us to express the $\lambda^i(\gamma(F))$, for a coherent algebraic sheaf F, by local cohomological invariants of F).

For a projection morphism $f : P\times X \longrightarrow X$, we can, by proposition 2.13, assume that x is of the form $y\otimes z$, with $y\in\underline{K}(P)$ and [print 70]$z\in\underline{K}(X)$. Then the Künneth[68] formula gives $f_!(x) = \chi(y)z$, where, for a coherent algebraic sheaf F on P, we put:

$$\chi(F) = \sum_{i\geq 0}(-1)^i \dim H^i(P,F).$$

We are then immediately reduced to proving the formula

$$\chi(F) = \varkappa_r(\mathrm{ch}_P(F)\,\mathcal{T}(P)) \tag{2.40}$$

where $\varkappa_r$ denotes the component of degree $r = \dim P$. By proposition 2.13, we can assume that $F = \underline{O}_Q$, where Q is a linear subvariety of P. Then $\chi(F) = 1$, since it is the arithmetic genus of Q; it remains to see that the right-hand side of formula (2.40) equals 1.

(*) See below for a proof valid in every characteristic, based on a different principle (p. 51).

Indeed, if V is a hyperplane of P and we put $\xi = \gamma_P(\underline{O}_V)$ and $L = \ell(V)$, we shall have:

$$1 - \xi = \gamma_P(E(-L))$$

(where $E(-L)$ is the vector bundle defined by $-L$), whence:

$$C_P(1 - \xi) = [1, 1 - L],$$

and consequently

$$\mathrm{ch}_P(1 - \xi) = e^{-L}, \qquad \text{whence} \qquad \mathrm{ch}_P(\xi) = 1 - e^{-L},$$

and therefore

$$\mathrm{ch}_P(\xi^p) = (1 - e^{-L})^p.$$

Now, if p is the codimension of Q in P[69], we have $\xi^p = \gamma_P(Q)$; we deduce

$$\mathrm{ch}_P(F)\,\mathcal{T}(P) = \big(L/(1 - e^{-L})\big)^{r+1}(1 - e^{-L})^p$$

and we must verify that the coefficient of L^r in this formal series in L equals 1, which follows from the characteristic property of the formal series $L/(1 - e^{-L})$, namely that the coefficient of L^q in $\big(L/(1 - e^{-L})\big)^{q+1}$ equals 1 (put $q = r - p$[70]).

[print 71]

Bibliographical references:

[1] Hirzebruch: Neue topologische Methoden[71] in der algebraischen Geometrie.

[2] Serre: Faisceaux algébriques cohérents, Ann. of Maths.

[3] Grothendieck: Sur les faisceaux algébriques cohérents, in Séminaire Cartan 1956/57.

Proof of the Riemann–Roch theorem

(Grothendieck, 1 November 1957)

* Reminder of notation

[These are those of the "report on Riemann–Roch", cited as R.R.R. in what follows.]

All varieties considered are nonsingular and quasi-projective (i.e. isomorphic to a locally closed subvariety of a projective space). The base field is algebraically closed, of arbitrary characteristic.

For such a variety X, we denote by $K(X)$ the abelian group generated by the classes of coherent sheaves on X (modulo the identification of an extension with a sum); the alternating sum of Tor makes $K(X)$ a ring. If $\underline{F}$ is a sheaf (resp. a vector bundle E, resp. a sub[print 72]variety Y of X), we denote by $\gamma_X(\underline{F})$ (resp. $\gamma_X(E)$, resp. $\gamma_X(Y)$) the element of $K(X)$ corresponding to $\underline{F}$ (resp. to $\underline{E}$, resp. to $\underline{O}_Y$). (In fact, we often write $\underline{F}$, E instead of $\gamma_X(\underline{F})\ldots$).

We denote by $A(X)$ the graded ring of classes of cycles on X (for rational equivalence—cf. Chow); $A^i(X)$ consists of cycles of codimension i.

The passages between asterisks $^*\ldots*$ are interpolations by the copyist.

For a vector bundle E, we denote by $C(E) = \sum C^i(E)$, $C^i(E) \in A^i(X)$, its Chern class (defined as the inverse image of Schubert cycles); by linearity we deduce the $C^i(y)$, for every $y \in K(X)$. If $C(y) = \prod(1 + a_i)$, we put, following Hirzebruch:

$$\mathrm{ch}(y) = \sum e^{a_i}, \qquad T(y) = \prod \frac{a_i}{1 - e^{-a_i}}.$$

These are elements of $A(X) \otimes \mathbb{Q}$.

When y is the tangent bundle of X, we write $T(X)$ instead of $T(y)$.

For a vector bundle G, we denote by $\lambda^p G$ its p-th exterior power, and put $\lambda_{-1}(G) = \sum(-1)^p \lambda^p(G)$; this is an element of $K(X)$.

For a proper regular map f from a variety Y to a variety X, we define (additive) homomorphisms:

$$f^* : A(X) \longrightarrow A(Y) \qquad \text{and} \qquad f_* : A(Y) \longrightarrow A(X).$$

The first is homogeneous of degree 0 and multiplicative. The second is homogeneous of degree $\dim .X - \dim .Y$.

We also define homomorphisms:

$$f^! : K(X) \longrightarrow K(Y) \qquad \text{and} \qquad f_! : K(Y) \longrightarrow K(X).$$

For a coherent sheaf $\underline{F}$ on X, $f^!(\underline{F})$ equals, by definition, the alternating sum of the $\mathrm{Tor}_p^{O_X}(\underline{O}_Y, \underline{F})$. For vector bundles, this is simply the inverse-image operation.

[print 73]For a coherent sheaf G on Y, $f_*(\underline{G})$ is coherent on X. Denoting by $R^p f_*$ the p-th derived functor of the functor f_* thus defined[72], we have, by definition, $f_!(\underline{G}) = \sum(-1)^p R^p f_*(\underline{G})$. $*$

RIEMANN–ROCH THEOREM. Let $f : Y \longrightarrow X$ be a proper morphism (* regular map $*$) from a nonsingular quasi-projective variety Y to another X, and let $y \in K(Y)$. We have:

$$f_*(\mathrm{ch}(y)T(Y)) = \mathrm{ch}(f_!(y))T(X) \quad \text{in } A(X) \otimes \mathbb{Q}. \tag{1}$$

Proof (* abridged $*$).

LEMMA 1. Consider proper morphisms of varieties $Z \xrightarrow{g} Y \xrightarrow{f} X$. If the theorem R.R. is true for f and g, it is true for $f \circ g$. If it is true for fg and g, it is true for f. If f_* is injective, and the theorem R.R. is true for fg and f, it is true for g.

Trivial. (* In fact, when we assume R.R. true for fg and g, we must also assume that $g_!$ is surjective to be able to assert that R.R. is true for f. $*$)

As a proper morphism is the composite of an injection and a projection $X \times P \longrightarrow X$ (where P is a projective space), we are reduced to treating these two cases separately. The second, as we have seen, reduces to a standard calculation, by proposition 2.13 ("cellular decomposition") of R.R.R. We shall therefore assume $Y \subset X$, with the injection morphism i. Formula (1) then becomes:

$$\mathrm{ch}(i_!(y)) = i_*(\mathrm{ch}(y)T(E)^{-1}), \tag{2}$$

where $E = T_{X/Y}$ is the normal bundle of Y in X.

[print 74]**LEMMA 2.** Assume that there exists on X a vector bundle G of rank $p = \mathrm{codim}_X Y$, such that:

$$\gamma_X(Y) = i_!(1) = \lambda_{-1}(G)\,; \qquad i^!(G) = \check{E}\,; \qquad (-1)^p C^p(G) = Y. \tag{3}$$

If we then have:

$$y = i^!(x), \qquad \text{with } x \in K(X), \tag{4}$$

formula (2) is valid.

The proof is immediate from the fact that ch is a ring homomorphism, and that we have:

$$\mathrm{ch}(\lambda_{-1}(G)) = (-1)^p C^p(G) T(\check{G})^{-1}. \tag{5}$$

COROLLARY. R.R. is true if Y is a hypersurface and y is of the form $i^!(x)$, with $x \in K(X)$.

LEMMA 3. Assume that $X = Y \times P$, with P a projective space, and that i is the injection $u \longrightarrow (u, v_0)$ $v_0 \in P$. Then i_* is injective and R.R. is true.

This is pure standard calculation.

Assume again that X, Y, y are arbitrary (still with $Y \subset X$). Let X' be the variety obtained by blowing up X along Y, and Y' the inverse-image hypersurface of Y. We have a diagram of canonical maps

$$\begin{array}{ccc} Y & \xrightarrow{\ i\ } & X \\ {\scriptstyle g}\uparrow & & \uparrow{\scriptstyle f} \\ Y' & \xrightarrow{\ j\ } & X' \end{array}$$

(j being the canonical map). The inverse image of E on Y' has a subbundle of rank 1, whose dual is denoted by L. Moreover, L^{-1} is also the restriction to Y' of the bundle $L(Y')$ defined on X' by the divisor Y'. We put $F = E/L^{-1}$. Elementary calculations give:

[print 75]**LEMMA 4.** We have the formulas (where $p = \mathrm{codim}_X Y$):

$$f^! i_!(y) = j_!(\lambda_{-1}(\check{F}) y). \tag{6}$$

$$\lambda_{-1}(\check{F}) \equiv \gamma^{p-1}(E) \lambda^p(\check{E}) \mod.(1-L)(*). \tag{7}$$

$$f_*(1) = 1, \quad \text{whence } f_* f^* = 1. \tag{8}$$

$$f^* i_*(\alpha) \equiv j_*(\alpha C^{p-1}(F)) \mod. \mathrm{Ker}\, f_* \quad \text{for every } \alpha \in A(Y). \tag{9}$$

[To simplify notation, we have made $K(Y)$ operate on $K(Y')$ and $A(Y)$ on $A(Y')$ by means of the homomorphisms $g^!$ and g^*. *In formula (7), it seems that $\gamma^{p-1}(E)$ denotes $\lambda^{p-1}(E-2)$, that is, $\sum_{i+j=p-1} \binom{-p}{j} \lambda^i(E)$; the congruence holds in $K(Y')$.*]

Formula (6) follows from the equality:

$$\mathrm{Tor}_i^{\underline{O}_X(v)}(\underline{O}_Y(v), \underline{O}_{Y'}(v')) = \wedge^i \check{F}(v') \quad (v' \in Y',\ v = g(v')). \tag{6 bis}$$

Formula (7) follows from the formulas

$$g_!(L^{-i}) = 0 \quad \text{for } 0 < i \leq p-1, \qquad g_!(1) = 1, \tag{10}$$

which themselves follow from Künneth[73] and the cohomology of projective spaces.

Formula (8) is trivial; finally, (9) follows from $g_*(C^{p-1}(F)) = 1$, which is deduced, for example, from the immediate formulas:

$$g_*(\xi^i) = 0 \quad \text{if } 0 \leq i < p-1, \qquad g_*(\xi^{p-1}) = 1, \tag{10 bis}$$

with $\xi = C^1(L) \in A^1(Y')$.

[In fact, (9) is very probably an identity, and not merely a congruence.]

(*) This unnecessarily complicated formula can be replaced, for example, by VII 4.1.

[print 76]Formula (7) immediately gives $y\lambda_{-1}(\check{F}) \in j^!(K(X'))$, provided that we have:

$$y\,\gamma^{p-1}(E)\lambda^p(\check{E}) \in i^!(K(X)). \tag{11}$$

Assume that this condition (11) is satisfied. By the corollary to lemma 2, R.R. is true for $(Y', X', y\lambda_{-1}(\check{F}))$. On the other hand, to prove (2), it suffices, by formula (8), to prove that

$$f^* \operatorname{ch}(i_!(y)) \equiv f^*(i_*(\operatorname{ch}(y)T(E)^{-1})) \quad \text{mod. Ker } f_*. \tag{12}$$

Now, this last formula is verified immediately, using (6), (9), R.R. for $(Y', X', y\lambda_{-1}(\check{F}))$[74] and the formula:

$$\operatorname{ch}(\lambda_{-1}(\check{F})) = C^{p-1}(F)T(F)^{-1} \quad \text{(cf. formula (5))}. \tag{13}$$

Thus, R.R. is proved whenever $y\,\gamma^{p-1}(E) = 0$, in particular (since $\gamma^{p-1}(E)$ has filtration $\geq p-1$ *in $K(Y)$ filtered by the codimension of supports*) whenever $p - 1 > \dim .Y$, i.e. whenever $\dim .Y < \frac{\dim .X - 1}{2}$. Now, we reduce to this case by combining lemma 1 and lemma 3 (taking for P a projective space of sufficiently large dimension r, for example $r > \dim .X + 1$). Q.E.D.

N.B. We have used the fact that $\gamma^{p-1}(E)$ has filtration $\geq p - 1$. We can prove this fact directly, without passing through Grassmannians as in R.R.R. 2.16, by studying the bundle Y' of projective spaces associated with E. This method gives, more generally, th. 2.14 of R.R.R.

Extract from a letter of Grothendieck

... "morally", the new method rests on the determination of $K(X)$ and $A(X)$ when X is either a projective bundle associated with a vector bundle, or a variety obtained by blowing up a non[print 77]singular subvariety. In this second case, I have not made the complete determination; a little something is missing. With the notation of the short paper, we must prove formula (9) (but as an <u>equality</u>):

$$f^*(i_*(\alpha)) = j_*(\alpha C^{p-1}(F)), \qquad \alpha \in A(Y). \tag{9}$$

We observe that the two sides have the same image under f_*, and the same image under j^*, so that the question is equivalent to:

$$\operatorname{Ker} f_* \cap \operatorname{Ker} j^* = 0. \tag{9 bis}$$

In any case, (9) is true upon reduction to $G(K(X'))$ (which destroys a little torsion, at most), since it then follows from (6); there is therefore hardly any doubt that it is true.

...I also point out to you that the <u>equality</u> (9) follows very easily from (6) without passing through the troublesome (6 bis)...

p.c.c.J-P.S.

[print 78]

GENERALITIES ON FINITENESS CONDITIONS IN DERIVED CATEGORIES

by L. ILLUSIE

0. Introduction

The purpose of Exposés I to IV is to develop, with the generality appropriate to this Seminar, the formalism of finiteness conditions in the derived categories of ringed topoi.

As was said in Exp. 0, it was the need to define, on arbitrary schemes, "Grothendieck groups" with good variance properties that required us to generalize and relax the notions of finiteness used until now.

A classical notion such as that of a coherent sheaf on a ringed space $(X, \underline{O}_X)$ becomes uninteresting as soon as $\underline{O}_X$ ceases to be coherent. One might think of replacing it by the notion of finite presentation, but this has the drawback that the kernel of an epimorphism of Modules of finite presentation is in general no longer of finite presentation. We arrive at a satisfactory notion by observing that, when $\underline{O}_X$ is coherent, a coherent sheaf F is not only of finite presentation but of n-finite presentation for every $n \in \mathbb{N}$, "n-finite presentation" meaning that there locally exists an exact sequence [print 79]

$$L_n \longrightarrow L_{n-1} \longrightarrow \ldots \longrightarrow L_0 \longrightarrow F \longrightarrow 0,$$


where the L_i are free of finite type. With the sheaf $\underline{O}_X$ no longer assumed coherent, let us say that an $\underline{O}_X$-Module F is pseudo-coherent when it satisfies the preceding condition, i.e. is of n-finite presentation for every n. Then the notion of pseudo-coherence has, with respect to short exact sequences, the same stability property as coherence: when two of the terms of a short exact sequence of $\underline{O}_X$-Modules are pseudo-coherent, the third is too. We readily generalize the preceding notion to complexes of sheaves. A complex F of $\underline{O}_X$-Modules is called pseudo-coherent when it is n-pseudo-coherent for every $n \in \mathbb{Z}$, "n-pseudo-coherent" meaning that there locally exists a quasi-isomorphism $L \to F$, where L is a complex with degrees bounded above and components free of finite type in degree $\geq n$. For a complex F concentrated in degree 0, it amounts to the same thing to say that F is pseudo-coherent as a complex or as a Module. The notion of a pseudo-coherent complex has excellent stability properties, described in Exp. I §§ 1 and 2. First, it is stable under isomorphism in the derived category $\mathrm{D}(X)$ of the category of $\underline{O}_X$-Modules; moreover, when two of the objects of a distinguished triangle of $\mathrm{D}(X)$ are pseudo-coherent, the third is too: in other words, the full subcategory $\mathrm{D}(X)_{\mathrm{coh}}$ of $\mathrm{D}(X)$ formed by pseudo-coherent complexes is a triangulated subcategory. Furthermore, pseudo-coherence is preserved by derived inverse image and tensor product. When $\underline{O}_X$ is coherent, saying that a complex F is pseudo-coherent simply means that F has locally bounded-above cohomology and that all the $\underline{H}^i(F)$ are coherent.

When $\underline{O}_X$ is a sheaf of regular local rings, every $\underline{O}_X$-Module [print 80] F that is coherent locally admits a finite resolution to the left by free Modules of finite type, i.e. there locally exists an exact sequence

$$0 \longrightarrow L_n \longrightarrow \ldots \longrightarrow L_0 \longrightarrow F \longrightarrow 0,$$

where the L_i are free of finite type. It is therefore natural, when $\underline{O}_X$ is an arbitrary sheaf of local rings, to consider Modules with the preceding property. Such Modules are called

perfect. More generally, a complex F of $\underline{O}_X$-Modules is called perfect when there locally exists a quasi-isomorphism $L \to F$, where L is a complex with bounded degrees and components free of finite type. It is shown in Exp. I § 3 that the full subcategory $\mathrm{D}(X)_{\mathrm{parf}}$ of $\mathrm{D}(X)$ formed by perfect complexes is, like $\mathrm{D}(X)_{\mathrm{coh}}$, a triangulated subcategory, "stable" under derived inverse image and tensor product. We evidently have $\mathrm{D}(X)_{\mathrm{parf}} \subset \mathrm{D}(X)_{\mathrm{coh}}$, but the inclusion is strict in general. Pseudo-coherence can be redefined "by passage to the limit" from perfection: a complex F is pseudo-coherent precisely when, for every $n \in \mathbb{Z}$, F can be "locally approximated to order n by a perfect complex", by which we mean that there locally exists a distinguished triangle



where L is perfect and R acyclic in degree $\geq n$. Conversely, perfection is recovered from pseudo-coherence through an additional regularity condition: a complex is perfect precisely when it is pseudo-coherent and locally of finite tor-dimension (Exp. I § 5). We thus recover the fact that, when the local rings are regular, every coherent sheaf is perfect, and, more generally, every pseudo-coherent complex with locally bounded cohomology is perfect.

[print 81]Pseudo-coherence and perfection are the two fundamental notions of finiteness with which we shall work in this Seminar. Sections 1 to 5 of Exposé I are devoted to their definition and to establishing their elementary stability properties. For this, we use only two or three local properties of the category of free Modules of finite type with respect to the category of all Modules. It therefore proves convenient—and useful, particularly in Exp. II—to axiomatize the situation by introducing a notion of pseudo-coherence (resp. perfection) in a category fibered over a site relative to a fibered subcategory satisfying suitable conditions.

Sections 6 to 8 of Exp. I generalize to perfect complexes a number of familiar notions for locally free sheaves of finite type: rank, duality, trace of an endomorphism.

In Exp. II, we examine the problem of "the existence of global resolutions": under what conditions, on a ringed topos $(X, \underline{O}_X)$, is a pseudo-coherent (resp. perfect) complex globally isomorphic in $\mathrm{D}(X)$ to a complex of locally free Modules of finite type? The importance of this question lies essentially in the fact that, at present, we do not know how to generalize to perfect complexes certain usual constructions on locally free Modules: exterior and symmetric powers, Chern classes (for more details on this subject, see Exp. XIV). This is why it is convenient to have workable sufficient criteria for the existence of global resolutions, which allow us to reduce certain questions about pseudo-coherent (resp. perfect) complexes to analogous questions about locally free sheaves of finite type. Such criteria [print 82]are given in Exp. II. In particular, it is shown that global resolutions exist in the following cases: X is the Zariski topos of an affine scheme, or more generally of a quasi-compact scheme with an ample invertible sheaf, or even an ample family of invertible sheaves in Kleiman's sense (for example a regular scheme); or again X is the topos of sheaves of sets on a compact topological space, $\underline{O}_X$ the sheaf of continuous complex-valued functions. As an illustration—and to bring out the flexibility of the notions introduced—the appendix gives a "purely sheaf-theoretic" definition of the index of a family of elliptic operators.

Exposé III studies the stability of finiteness conditions under derived direct image. To obtain reasonable statements, we must "relativize" pseudo-coherence and perfection. We work here in the setting of ordinary schemes, which suffices for the Seminar, but there is no doubt that sooner or later an analogous theory will have to be developed for relative analytic schemes or spaces. Let $p : X \to S$ be an S-scheme locally of finite type and F a complex of $\underline{O}_X$-Modules; we say that F is pseudo-coherent (resp. perfect) relative to p when we can locally embed (by a closed immersion) X in a smooth S-scheme X' so that F, extended by zero to X', is pseudo-coherent

(resp. perfect). Pseudo-coherence relative to p is especially interesting when S is not locally noetherian, since in the opposite case it coincides with ordinary pseudo-coherence. Moreover, as one might expect, saying that F is perfect rel. to p [print 83]amounts to saying that F is pseudo-coherent rel. to p and locally of finite tor-dimension rel. to p (i.e. rel. to the sheaf of rings $p^{-1}(\underline{O}_S)$). The central theorem of Exp. III is the finiteness theorem, which asserts (in a slightly more precise form) that, for a proper morphism $f : X \to Y$ of S-schemes locally of finite type, the functor $\mathbb{R}f_*$ sends complexes pseudo-coherent rel. to S to complexes pseudo-coherent rel. to S; this theorem is in fact a conjecture, but is nevertheless proved in the following two special cases: a) S is locally noetherian b) f is projective; unfortunately, it seems that extending it to the general case is of the same order of difficulty as the analogous theorem in Analytic Geometry (Grauert's "theorem"). Combining the finiteness theorem with an essentially trivial formula called the projection formula, we obtain workable criteria for the stability of relative perfection under direct image. As a corollary, we recover "Grauert's continuity and semicontinuity theorems" (EGA III 7.6).

Exposé IV translates the results of Exposés I to III into the language of "Grothendieck groups". On a ringed topos $(X, \underline{O}_X)$, we denote by $K^\bullet(X)$ (resp. $K_\bullet(X)$) the "Grothendieck group" of the category of complexes that are perfect and of finite tor-dimension (resp. pseudo-coherent and with bounded cohomology). The group $K^\bullet(X)$ is a ring, and $K_\bullet(X)$ a module over $K^\bullet(X)$. $K^\bullet$ is a contravariant functor (as the raised dot suggests), while $K_\bullet$ is a covariant functor for proper pseudo-coherent morphisms (of schemes or analytic spaces, and subject to having the finiteness theorem...). There are also unusual variances, [print 84]covariance of $K^\bullet$, contravariance of $K_\bullet$, for morphisms satisfying suitable regularity hypotheses. Finally, the globalization criteria of Exp. II allow us to relate these to the "Grothendieck groups" defined naively from coherent sheaves or locally free sheaves of finite type.

1. Preliminary definitions

1.1. Fibered categories with additive (resp. abelian, resp. triangulated) fibres

1.1.1. Let $\underline{S}$ be a category, and $\underline{C}$ a fibered $\underline{S}$-category (SGA 1 VI 6.1) whose fibres are additive (resp. triangulated) categories. We shall say that $\underline{C}$ is an additive $\underline{S}$-category (resp. triangulated) (or that $\underline{C}$ is additive (resp. triangulated) over $\underline{S}$) when, for every arrow $f : X \to Y$ of $\underline{S}$, the inverse-image functor (determined up to unique isomorphism) $f^* : \underline{C}_Y \longrightarrow \underline{C}_X$ is additive (resp. exact). We shall say that $\underline{C}$ is an abelian $\underline{S}$-category when $\underline{C}$ is an additive $\underline{S}$-category with abelian fibres, and that $\underline{C}$ is a flat abelian $\underline{S}$-category when, moreover, the inverse-image functors are exact.

1.1.2. Let $\underline{C}$ be a triangulated $\underline{S}$-category. We define, for example using a cleavage of $\underline{C}$ (SGA 1 VI 7.1)[75], a fibered $\underline{S}$-category $\mathrm{Tr}(\underline{C})$ whose fibre at each $X \in \mathrm{ob}(\underline{S})$ is the category $\mathrm{Tr}(\underline{C}_X)$ of distinguished triangles of $\underline{C}_X$.

1.1.3. Let $\underline{C}$ and $\underline{C}'$ be additive (resp. triangulated) $\underline{S}$-categories and $F : \underline{C} \to \underline{C}'$ a functor. We call F an additive (resp. exact) $\underline{S}$-functor when F is a cartesian $\underline{S}$-functor that is additive (resp. exact) on each fibre. [print 85]When $\underline{C}$ and $\underline{C}'$ are triangulated $\underline{S}$-categories and F is an exact $\underline{S}$-functor, F defines a cartesian $\underline{S}$-functor $\mathrm{Tr}(\underline{C}) \longrightarrow \mathrm{Tr}(\underline{C}')$.

1.1.4. Let $\underline{C}$ be a triangulated $\underline{S}$-category, and $\underline{C}'$ a full fibered sub-$\underline{S}$-category of $\underline{C}$. We call $\underline{C}'$ a triangulated sub-$\underline{S}$-category of $\underline{C}$ when, for every $X \in \mathrm{ob}\,\underline{S}$, $\underline{C}'_X$ is a triangulated subcategory of $\underline{C}_X$ ([V] chap. I § 1 2.3).

1.1.5. Let $\underline{C}$ be an additive $\underline{S}$-category. We define (SGA 4 XVII 2.2, 2.4) an additive (resp. triangulated) $\underline{S}$-category $\mathrm{C}(\underline{C})$, (resp. $\mathrm{K}(\underline{C})$), whose fibre at $X \in \mathrm{ob}\,\underline{S}$ is the category $\mathrm{C}(\underline{C}_X)$

(resp. $\mathrm{K}(\underline{C}_X)$) of complexes([1]) (resp. complexes "up to homotopy") of $\underline{C}_X$. When $\underline{C}$ is a flat abelian $\underline{S}$-category, we define (loc. cit.) a triangulated $\underline{S}$-category $\mathrm{D}(\underline{C})$, whose fibre at $X \in \mathrm{ob}\,\underline{S}$ is the derived category $\mathrm{D}(\underline{C}_X)$; for an arrow $f : X \to Y$ of $\underline{S}$, we shall denote by $f^* : \mathrm{D}(\underline{C}_Y) \longrightarrow \mathrm{D}(\underline{C}_X)$ the inverse-image functor, obtained from $f^* : \mathrm{K}(\underline{C}_Y) \longrightarrow \mathrm{K}(\underline{C}_X)$ by simple passage to the quotient([2]). We define full triangulated sub-$\underline{S}$-categories $\mathrm{K}^*(\underline{C})$ (resp. $\mathrm{D}^*(\underline{C})$) $(* = +, -, b)$ by the usual degree conditions.

1.2. Terminology. Let $\underline{S}$ be a site, $\underline{C}$ a flat abelian $\underline{S}$-category (1.1), and $\underline{C}_0$ a strictly full fibered sub-$\underline{S}$-category.

Let $F \in \mathrm{ob}\,\underline{C}_X$, for $X \in \mathrm{ob}\,\underline{S}$. The set of $f : U \to X$ in $\underline{S}_{/X}$ such that there exists an epimorphism $E \longrightarrow f^*F$ with $E \in \mathrm{ob}\,\underline{C}_{0U}$ is a sieve on X. When it is covering, we shall say that F is of $\underline{C}_0$-finite type (or simply of finite type when there is no risk of confusion).

[print 86]We shall say that $\underline{C}_0$ is locally stable under kernels of epimorphisms when the following condition is satisfied: for every $X \in \mathrm{ob}\,\underline{S}$ and every epimorphism $u : E \longrightarrow F$ of $\underline{C}_X$ with E and $F \in \mathrm{ob}\,\underline{C}_{0X}$, the kernel of u is locally in $\underline{C}_0$ (i.e. the set of $f : U \to X$ in $\underline{S}_{/X}$ such that $f^*\,\mathrm{Ker}(u) \in \mathrm{ob}\,\underline{C}_{0U}$ is a refinement of X).

By the standard procedure of decomposing a long exact sequence into short exact sequences, we see that the preceding condition is equivalent to the following (apparently stronger) condition: for every $X \in \mathrm{ob}\,\underline{S}$ and every exact sequence of $\underline{C}_X$

$$0 \longrightarrow E^0 \longrightarrow E^1 \longrightarrow \ldots \longrightarrow E^n \longrightarrow 0 \qquad (*)$$

with $E^i \in \mathrm{ob}\,\underline{C}_{0X}$ for $1 \leq i \leq n$, E^0 is locally in $\underline{C}_0$.

We shall say that $\underline{C}_0$ is locally quasi-stable under kernels of epimorphisms when, for every exact sequence such as $(*)$, E^0 is of $\underline{C}_0$-finite type.

On the other hand, we shall say that $\underline{C}_0$ is locally liftable in $\underline{C}$ when, for every $X \in \mathrm{ob}\,\underline{S}$, every epimorphism $u : E \longrightarrow F$ of $\underline{C}_X$ with $F \in \mathrm{ob}\,\underline{C}_{0X}$ locally admits a section. For $\underline{C}_0$ to be locally liftable in $\underline{C}$, it is necessary and sufficient that, for every diagram of $\underline{C}_X$ $(X \in \mathrm{ob}\,\underline{S})$

$$\begin{array}{ccc} & & H \\ & & \downarrow \\ E & \xrightarrow{u} & F \end{array}$$

where u is an epimorphism and $H \in \mathrm{ob}\,\underline{C}_0$, the set of U over X such that there exists a commutative diagram [76]

$$\begin{array}{ccc} & & H_U \\ & \swarrow & \downarrow \\ E_U & \xrightarrow{u_U} & F_U \end{array}$$

is a refinement of X.

[print 87]We shall say that $\underline{C}_0$ is locally quasi-liftable in $\underline{C}$ when the following weaker condition is satisfied: for every $X \in \mathrm{ob}\,\underline{S}$, and every epimorphism $u : E \longrightarrow F$ of $\underline{C}_X$ with $F \in \mathrm{ob}\,\underline{C}_{0X}$, the set of $f : U \to X$ in $\underline{S}_{/X}$ such that there exists $v : F' \longrightarrow f^*E$ with $F' \in \mathrm{ob}\,\underline{C}_{0U}$ such that $f^*(u)v$ is an epimorphism is a refinement of X.

([1]) with differential of degree $+1$.
([2]) for $a \in \mathrm{ob}\,\mathrm{D}(\underline{C}_Y)$ (resp. $\mathrm{Fl}\,\mathrm{D}(\underline{C}_Y)$), we shall often write, in abbreviated form, a_X instead of $f^*(a)$.

This condition is equivalent to the following: for every diagram of $\underline{C}_X$

$$\begin{array}{ccc} & & H \\ & & \downarrow \\ E & \xrightarrow{u} & F, \end{array}$$

where u is an epimorphism, and $H \in \mathrm{ob}\,\underline{C}_0$, the set of U over X such that there exists a commutative diagram

$$\begin{array}{ccc} G & \xrightarrow{v} & H_U \\ \downarrow & & \downarrow \\ E_U & \xrightarrow{u_U} & F_U, \end{array}$$

with $G \in \mathrm{ob}\,\underline{C}_0$ and v an epimorphism is a refinement of X.

When $\underline{S}$ is a chaotic site, for example the punctual site (one object, one morphism), we shall omit the adverb "locally" in the preceding definitions.

1.3. Examples

a) Let $\underline{S}$ be a punctual site, A a ring, and $\underline{C}$ the category of left A-modules. The full subcategory $\underline{C}_0$ of $\underline{C}$ formed by the projective (left) A-modules of finite type is stable under kernels of epimorphisms and liftable in $\underline{C}$; an A-module is of $\underline{C}_0$-finite type if and only if it is of finite type in the [print 88]ordinary sense. The full subcategory $\underline{C}_1$ of $\underline{C}$ formed by the A-modules of finite type is quasi-liftable in $\underline{C}$ (but not liftable in general); when A is noetherian, $\underline{C}_1$ is stable under kernels of epimorphisms. The full subcategory $\underline{C}'_0$ of $\underline{C}$ formed by the free A-modules of finite type is liftable in $\underline{C}$ and quasi-stable (not stable in general) under kernels of epimorphisms.

b) Let $(S, \underline{O}_S)$ be a space ringed by local rings, $\underline{S}$ the site of open subsets of S, and $\underline{C}$ the fibered $\underline{S}$-category (even split (SGA 1 VI 9)) defined by the functor $U \longmapsto$ (category of left $\underline{O}_U$-Modules); $\underline{C}$ is a flat abelian $\underline{S}$-category (1.1). Let $\underline{C}_0$ be the strictly full additive sub-$\underline{S}$-category of $\underline{C}$ defined by $U \longmapsto$ (category of free $\underline{O}_U$-Modules of finite type). An $\underline{O}_S$-Module M is locally of $\underline{C}_0$-finite type if and only if M is of finite type in the ordinary sense (EGA 0_{I} 5.2). It follows from (EGA 0_{I} 5.2.7 and 5.4.9) that $\underline{C}_0$ is locally stable under kernels of epimorphisms and locally liftable in $\underline{C}$.

c) Let G be a group, $\underline{S}$ the topos of G-sets (SGA 4 V 3.11), k[77] a field (regarded as a constant Ring of $\underline{S}$), and $\underline{C}$ the split $\underline{S}$-category defined by the functor $U \longmapsto$ (category of left k_U-Modules); $\underline{C}$ is a flat abelian $\underline{S}$-category. Let $\underline{C}_0$ be the strictly full additive sub-$\underline{S}$-category of $\underline{C}$ defined by $U \longmapsto$ (category of free k_U-Modules of finite type). Since localization in $\underline{S}$ is equivalent to forgetting the operations of G, it is clear that a G-k-module M is locally of $\underline{C}_0$-finite type if and only if M is finite-dimensional as a vector space over k, and that $\underline{C}_0$ is locally stable under kernels of epimorphisms and locally liftable in $\underline{C}$.

The reader is left to examine the "Galois" variant of c) in which G is a profinite group and $\underline{S}$ the topos of sets on which G operates continuously.

[print 89]In fact, examples b) and c) are special cases of the following example:

d) Let $(\underline{S}, \underline{O}_{\underline{S}})$ be a ringed topos (cf. SGA 4 IV 2.10)(*), and $\underline{C}$ the split $\underline{S}$-category defined by $U \longmapsto$ (category of left $\underline{O}_U$-Modules); $\underline{C}$ is a flat abelian $\underline{S}$-category. Let $\underline{C}_0$ be the strictly full additive sub-$\underline{S}$-category of $\underline{C}$ defined by $U \longmapsto$ (category of free $\underline{O}_U$-Modules of finite type). An $\underline{O}_{\underline{S}}$-Module locally of $\underline{C}_0$-finite type will simply be said to be of finite type. The category $\underline{C}_0$ is locally quasi-stable under kernels of epimorphisms and locally liftable in $\underline{C}$, as follows from

(*) Here $\underline{O}_{\underline{S}}$ is a ring of $\underline{S}$, with identity element, but not necessarily commutative.

Lemma 1.3.1.[78] Let P be an $\underline{O}_S$-Module. Equivalent conditions[79]:

(i) P is of finite type and the functor $\underline{\mathrm{Hom}}(P, .)$ is exact;

(ii) P is of finite type and every epimorphism $M \to P$ locally admits a section;

(iii) P is locally a direct factor of a free Module of finite type.

The proof is immediate and is left as an exercise to the reader.

1.4. An "extension" lemma (cf. EGA 0_{III} 11.9.1)

Proposition 1.4.1. Let S be a site,[80] C a flat abelian[81] S-category (1.1),[82] C_0 a strictly full additive sub-S[83]-category (1.1) of C; assume that[84] C_0 is locally quasi-liftable in C (1.2). Let[85] $S \in \mathrm{ob}\, S$, $u : E \longrightarrow F$ be a morphism (of degree 0) of complexes of C_S (differentials of degree $+1$), G the cone of u (cf. [V] chap. I § 1 2.1), $n \in Z$; assume that $\mathrm{H}^n(G)$ is [print 90]of C_0-finite type (1.2). Then the set of $X \in \mathrm{ob}\, S_{/S}$ satisfying property (P) below is a refinement of S.

(P): there exist an object M of C_{0X} and arrows of C_X

$$r : M \longrightarrow Z^{n+1}(E_X) = \mathrm{Ker}\left(E_X^{n+1} \xrightarrow{d^{n+1}} E_X^{n+2}\right)$$

$$\text{and} \quad s : M \longrightarrow F_X^n$$

such that the following diagram of C_X

$$\begin{array}{ccccccccc}
\ldots & \longrightarrow & E_X^{n-1} & \xrightarrow{\binom{d^{n-1}_{E_X}}{0}} & E_X^n \oplus M & \xrightarrow{(d^n_{E_X}, r)} & E_X^{n+1} & \longrightarrow & \ldots \\
 & & \downarrow{\scriptstyle u_X^{n-1}} & & \downarrow{\scriptstyle (u_X^n, s)} & & \downarrow{\scriptstyle u_X^{n+1}} & & \\
\ldots & \longrightarrow & F_X^{n-1} & \longrightarrow & F_X^n & \longrightarrow & F_X^{n+1} & \longrightarrow & \ldots
\end{array}$$

is a morphism of complexes $u' : E' \to F_X$ (i.e. $d^n_{F_X} s = u_X^{n+1} r$), and that

(i) $\mathrm{H}^{n+1}(u')$ is a monomorphism, with $\mathrm{Im}\, \mathrm{H}^{n+1}(u') = \mathrm{Im}\, \mathrm{H}^{n+1}(u_X)$,

(ii) $\mathrm{H}^n(u')$ is an epimorphism.

In pictorial terms, we may say that, by modifying E and u locally, in degree n only, by adding to E^n an object of C_0, we can locally make $\mathrm{H}^{n+1}(u)$ injective, without changing its image, and $\mathrm{H}^n(u)$ surjective.

Proof. The procedure is as follows: first we show that we can choose r and s locally so as to make $\mathrm{H}^{n+1}(u)$ injective; we verify [print 91]that the finiteness hypothesis on $\mathrm{H}^n(G)$ is preserved; a new local correction in degree n then allows us to make $\mathrm{H}^n(u)$ surjective.

The exact cohomology sequence

$$\longrightarrow \mathrm{H}^n(G) \longrightarrow \mathrm{H}^{n+1}(E) \longrightarrow \mathrm{H}^{n+1}(F) \longrightarrow$$

shows that the kernel of $\mathrm{H}^{n+1}(u)$ is of $\underline{C}_0$-finite type. In other words, the set of X over S such that there exists an exact sequence

$$M \longrightarrow \mathrm{H}^{n+1}(E_X) \xrightarrow{\mathrm{H}^{n+1}(u)} \mathrm{H}^{n+1}(F_X),$$

with $M \in \mathrm{ob}\,\underline{C}_{0X}$, is a refinement of S. Since $\underline{C}_0$ is locally quasi-liftable in $\underline{C}$, we can moreover require the arrow $M \longrightarrow \mathrm{H}^{n+1}(E_X)$ to factor through $Z^{n+1}(E_X)$ as

$$M \xrightarrow{r} Z^{n+1}(E_X) \longrightarrow \mathrm{H}^{n+1}(E_X).$$

The composite arrow

$$M \xrightarrow{u_X^{n+1} r} Z^{n+1}(F_X) \longrightarrow \mathrm{H}^{n+1}(F_X)$$

is zero by construction, so $u_X^{n+1} r$ factors through $B^{n+1}(F_X) = \mathrm{Im}\Big(F_X^n \xrightarrow{d^n} F_X^{n+1}\Big)$. Applying quasi-liftability again, we can require $u_X^{n+1} r$ to factor through F_X^n as


$$M \xrightarrow{s} F_X^n \longrightarrow B^{n+1}(F_X) \longrightarrow Z^{n+1}(F_X).$$

We then check trivially that u', defined as in (1.4.1), is a morphism of complexes satisfying (i).

Let G' be the cone of $u' : E' \to F_X$. Let us show that $\mathrm{H}^n(G')$ is of $\underline{C}_0$-finite type. By construction, there is an exact sequence

$$0 \longrightarrow E_X \xrightarrow{v} E' \longrightarrow M[-n] \longrightarrow 0,$$

[print 92]with $u'v = u_X$. To compare G_X and G', we can, for example, form the "octahedron" of $\mathrm{K}(\underline{C}_X)$ [86]



.

The exact cohomology sequence of the distinguished triangle $(*)$ shows that $\mathrm{H}^n(G_X) \longrightarrow \mathrm{H}^n(G')$ is an epimorphism, hence that $\mathrm{H}^n(G')$ is indeed of $\underline{C}_0$-finite type.

The preceding correction therefore reduces us to the case where $\mathrm{H}^{n+1}(u)$ is a monomorphism, which we shall now assume. The exact cohomology sequence of $E \to F \to G$ then gives an exact sequence

$$\mathrm{H}^n(E) \xrightarrow{\mathrm{H}^n(u)} \mathrm{H}^n(F) \longrightarrow \mathrm{H}^n(G) \longrightarrow 0. \qquad (.)$$

Since $\mathrm{H}^n(G)$ is of $\underline{C}_0$-finite type, the set of X over S such that there exists an epimorphism $M \longrightarrow \mathrm{H}^n(G_X)$ with $M \in \mathrm{ob}\,\underline{C}_{0X}$ is, by definition, a refinement of S. Using the quasi-liftability of $\underline{C}_0$ in $\underline{C}$, we can require this epimorphism to factor through $Z^n(F_X)$ as

$$M \xrightarrow{s} Z^n(F_X) \longrightarrow \mathrm{H}^n(F_X) \longrightarrow \mathrm{H}^n(G_X).$$

It is clear that u', defined as in (1.4.1) with $(r = 0)$, is a morphism of complexes and that $\mathrm{H}^{n+1}(u') = \mathrm{H}^{n+1}(u_X)$. On the other hand, it follows immediately from the exact sequence (.) that [print 93]

$$\mathrm{H}^n(u') : \mathrm{H}^n(E_X) \oplus M \longrightarrow \mathrm{H}^n(F_X)$$

is an epimorphism. This completes the proof of (1.4.1).

Lemma 1.4.2. Let $\underline{C}$[87] be an abelian category, $u : E \to F$ a morphism of complexes of $\underline{C}$[88] (resp. an arrow of $\mathrm{D}(\underline{C})$), G the cone of u. Equivalent conditions for a given $n \in \mathbb{Z}$:

(i) $\mathrm{H}^i(G) = 0$ for $i \geq n$;

(ii) $\mathrm{H}^i(u)$ is an isomorphism for $i > n$ and an epimorphism for $i = n$.

The proof follows immediately from the exact cohomology sequence.

Definition 1.4.3. We shall say that u is an n-quasi-isomorphism (resp. an n-isomorphism) when u satisfies the equivalent conditions of (1.4.2).

That being so, we deduce at once from (1.4.1) the

Corollary 1.4.4. Let $S, S, \underline{C}, \underline{C}_0$ be as in (1.4.1). Let $u : E \to F$ be an $(n+1)$-quasi-isomorphism of complexes of $\underline{C}_S$ ($n \in \mathbb{Z}$ given) such that $\mathrm{H}^n(G)$ is of $\underline{C}_0$-finite type. Then the set of X over S such that there exist (M, r, s) as in (1.4.1) such that u' is an n-quasi-isomorphism is a refinement of S.

In pictorial terms, using the finiteness condition on $\mathrm{H}^n(G)$, we can locally "gain one notch", i.e. modify u locally in degree n by adding to E^n an object of $\underline{C}_0$ so as to make u an n-quasi-isomorphism.

Corollary 1.4.5. Let $\underline{C}$ be an abelian category, $n \in \mathbb{Z}$, and $u : E \to F$ an n-quasi-isomorphism of complexes of $\underline{C}$. There exists a commutative diagram of $\mathrm{C}(\underline{C})$

$$\begin{array}{ccc} E & \xrightarrow{v} & E' \\ {\scriptstyle u}\downarrow & \swarrow_{u'} & \\ F & & \end{array},$$

[print 94]where u' is a quasi-isomorphism and v a monomorphism inducing the identity in degree $\geq n$ (i.e. such that $E'^i = E^i$ and $v^i = \mathrm{Id}(E^i)$ for $i \geq n$).

Proof. Applying (1.4.4) in the case where S is the punctual site and $\underline{C}_0 = \underline{C}$, we obtain a commutative triangle of $\mathrm{C}(\underline{C})$

$$\begin{array}{ccc} E & \xrightarrow{v_{n-1}} & E'_{n-1} \\ {\scriptstyle u}\downarrow & \swarrow_{u'_{n-1}} & \\ F & & \end{array},$$

where u'_{n-1} is an $(n-1)$-quasi-isomorphism and v_{n-1} a monomorphism inducing the identity in degree $\geq n$. Iterating the procedure, we find, for $i \leq n-1$, a sequence of commutative triangles of $\mathrm{C}(\underline{C})$

$$\begin{array}{ccc} E'_i & \xrightarrow{v_{i-1}} & E'_{i-1} \\ {\scriptstyle u'_i}\downarrow & \swarrow_{u'_{i-1}} & \\ F & & \end{array},$$

where u'_i is an i-quasi-isomorphism and v_{i-1} a monomorphism inducing the identity in degree $\geq i$. The corollary follows by passage to the limit. More precisely, denoting by d'_i the differential of E'_i, we define E' by

$$E'^i = E^i, \quad d'^i = d^i \quad \text{for } i \geq n,$$

and

$$E'^i = (E'_i)^i, \quad d'^i = (d'_i)^i \quad \text{for } i < n,$$

then v and u' by

$$v^j = (v_i v_{i+1} \dots v_{n-1})^{j\,[89]}, \qquad u'^j = (u'_i)^j \qquad \text{for } i \leq j,$$

and we check at once that E', v, u' so defined answer the question.

[print 95]In other words, briefly, we can extend an n-quasi-isomorphism to a quasi-isomorphism "without changing anything given in degree $\geq n$".

2. Pseudo-coherent complexes

2.0. In this section, $\underline{S}$ denotes a site, S an object of $\underline{S}$, $\underline{C}$ a flat abelian $\underline{S}$-category (1.1), and $\underline{C}_0$ a strictly full additive sub-$\underline{S}$-category of $\underline{C}$. We assume that $\underline{C}_0$ is locally quasi-stable under kernels of epimorphisms (1.2) and locally quasi-liftable in $\underline{C}$ (1.2).

Definition 2.1. Let $X \in \mathrm{ob}\,\underline{S}$, E be a complex of $\underline{C}_X$, and $n \in \mathbb{Z}$. We shall say that E is strictly $\underline{C}_0$-n-pseudo-coherent (resp. $\underline{C}_0$-pseudo-coherent, resp. $\underline{C}_0$-perfect) when $E^i = 0$ for i sufficiently large and $E^i \in \mathrm{ob}\,\underline{C}_{0X}$[90] for $i \geq n$ (resp. for every i, resp. for every i and $E^i = 0$ for almost every i).

When there is no risk of confusion, we shall sometimes say "strictly n-pseudo-coherent (resp...)" instead of "strictly $\underline{C}_0$-n-pseudo-coherent (resp...)".

Lemma 2.1.1. Let $E \in \mathrm{ob}\,\mathrm{C}(\underline{C}_S)$ and $n \in \mathbb{Z}$. When E is strictly n-pseudo-coherent and $\mathrm{H}^i(E) = 0$ for $i > n$, then $\mathrm{H}^n(E)$ is of finite type.

Proof. The sequence

$$0 \longrightarrow Z^n(E) \longrightarrow E^n \longrightarrow E^{n+1} \longrightarrow \ldots \longrightarrow 0$$

is exact and $E^i \in \mathrm{ob}\,\underline{C}_0$ for $i \geq n$. Since $\underline{C}_0$ is locally quasi-stable under kernels of epimorphisms, it follows that $Z^n(E)$ is of finite type, hence so is $\mathrm{H}^n(E)$, which is a quotient of it.

[print 96]**Proposition 2.2.** Let $F \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$ and $n \in \mathbb{Z}$. Then:

a) When F is strictly n-pseudo-coherent (relative to $\underline{C}_0$) and $u : E \to F$ is a quasi-isomorphism, the set of X over S such that there exists a quasi-isomorphism $v : F' \longrightarrow E_X$ with F' strictly n-pseudo-coherent is a refinement of S.

b) The following conditions are equivalent:

(i) the set of X over S such that there exists a quasi-isomorphism $u : E \longrightarrow F_X$ of $\mathrm{C}(\underline{C}_X)$, with E strictly n-pseudo-coherent, is a refinement of S;

(i′) the same condition as (i), but with u an isomorphism of $\mathrm{D}(\underline{C}_X)$;

(ii) the set of X over S such that there exists an n-quasi-isomorphism (1.4.3) $u : E \longrightarrow F_X$ of $\mathrm{C}(\underline{C}_X)$, with E strictly perfect, is a refinement of S;

(ii′) the same condition as (ii), but with u an n-isomorphism of $\mathrm{D}(\underline{C}_X)$.

Proof. a) For an integer i sufficiently large, there exists an i-quasi-isomorphism $v_i : F'_i \longrightarrow E$, with F'_i strictly n-pseudo-coherent (take $F'_i = 0$). Let us show that, for $i > n$, there exists, after refining S, an $(i-1)$-quasi-isomorphism $v_{i-1} : F'_{i-1} \longrightarrow E$, with F'_{i-1} strictly n-pseudo-coherent. By (1.4.4), it suffices to show that $\mathrm{H}^{i-1}(M')$, where M' is the cone of v_i, is of $\underline{C}_0$-finite type. Now, with M'' denoting the cone of uv_i, the hypothesis that u is a quasi-isomorphism and (1.4.2) imply that

$$\mathrm{H}^j(M') \xrightarrow{\sim} \mathrm{H}^j(M'') \qquad \text{for every } j$$

and $\mathrm{H}^j(M') = 0$ for $j \geq i$. [print 97]The cone formula $M''^j = F_i'^{\,j+1} \oplus F^j$ shows, on the other hand, that M'' is strictly $(i-1)$-pseudo-coherent. Thus, by (2.1.1), $\mathrm{H}^{i-1}(M'') = \mathrm{H}^{i-1}(M')$ is of finite

type, whence our assertion. By iteration, and after refining S, we deduce an n-quasi-isomorphism $v_n : F'_n \longrightarrow E$, where F'_n is strictly n-pseudo-coherent. But, by (1.4.5), we can extend v_n to a quasi-isomorphism $v : F' \longrightarrow E$ without changing anything in degree $\geq n$, hence in particular with F' strictly n-pseudo-coherent, which proves a).

b) The equivalence of conditions (i) and (i′) follows at once from a) and the definition of the isomorphisms of $\mathrm{D}^-(\underline{C}_X)$. For $E \in \mathrm{ob}\,\mathrm{D}^-(\underline{C}_S)$ strictly n-pseudo-coherent, there exists an exact sequence

$$0 \longrightarrow E' \longrightarrow E \longrightarrow E'' \longrightarrow 0,$$

where E' is strictly perfect, and $E''^i = 0$ for $i \geq n$; consequently $E' \to E$ is an n-quasi-isomorphism, giving the implication (i) $\Rightarrow$ (ii). The implication (ii) $\Rightarrow$ (ii′) is trivial, so it remains to prove (ii′) $\Rightarrow$ (i). Let, then, $u : E \longrightarrow F$ be an n-isomorphism of $\mathrm{D}^-(\underline{C}_S)$, with E strictly perfect; u is therefore defined by a diagram of $\mathrm{C}^-(\underline{C}_S)$

$$\begin{array}{ccccc} & & E' & & \\ & \swarrow^{s} & & \searrow^{u'} & \\ E & & & & F, \end{array}$$

where s is a quasi-isomorphism and u' an n-quasi-isomorphism. By a), there exists, after refining S, a quasi-isomorphism $t : E'' \longrightarrow E'$, with E'' strictly n-pseudo-coherent. We then apply (1.4.5) to extend $u't : E'' \to F$ to a quasi-isomorphism $E''' \longrightarrow F$ with E''' strictly n-pseudo-coherent, which proves (ii′) $\Rightarrow$ (i) and completes the proof of (2.2).

[print 98]**Definition 2.3.** Under the hypotheses of (2.2), we shall say that $F \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$ is n-$\underline{C}_0$-pseudo-coherent when F satisfies the equivalent conditions b) of (2.2). We shall say that F is $\underline{C}_0$-pseudo-coherent when F is n-$\underline{C}_0$-pseudo-coherent for every n.

When this will cause no confusion, we shall sometimes allow ourselves to say n-pseudo-coherent, pseudo-coherent, instead of n-$\underline{C}_0$-pseudo-coherent, $\underline{C}_0$-pseudo-coherent.

Remark 2.3.1. Let $F \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$ be a pseudo-coherent complex. The set of X over S such that there exists a quasi-isomorphism $E \longrightarrow F_X$ with E strictly pseudo-coherent (2.1) is in general not a refinement of S. However, we shall see later (2.7. a) and Exp.II) important cases where it is.

The proof of (2.2 a)) gives the following result:

Proposition 2.4. Let $n, i \in \mathbb{Z}$, and let $u : E \to F$ be an i-quasi-isomorphism of $\mathrm{C}(\underline{C}_S)$, with F n-pseudo-coherent and E strictly n-pseudo-coherent. Then, after refining S, there exists a commutative triangle of $\mathrm{C}(\underline{C}_S)$

$$\begin{array}{ccc} E & \xrightarrow{v} & E' \\ {\scriptstyle u}\downarrow & \swarrow_{u'} & \\ F & & \end{array},$$

where E' is strictly n-pseudo-coherent, u' a quasi-isomorphism and v a monomorphism inducing the identity in degree $\geq i$.

Proposition 2.5. a) When $F \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$ is n-pseudo-coherent (resp. pseudo-coherent), $F[m]$ is $(n-m)$-pseudo-coherent (resp. pseudo-coherent).


[print 99]b) Let

$$\begin{array}{ccccc} & & G & & \\ & \swarrow^{+1} & & \nwarrow & \\ E & & \longrightarrow & & F \end{array} \qquad (*)$$

be a distinguished triangle of $\mathrm{D}(\underline{C}_S)$. When E and G are n-pseudo-coherent (resp. pseudo-coherent), then F is n-pseudo-coherent (resp. pseudo-coherent).


Proof. a) Trivial from the definitions.

b) It of course suffices to prove the assertion without the "resp." alternatives. The triangle $(*)$ can also be written



.

By a), $G[-1]$ is $(n+1)$-pseudo-coherent. We are therefore reduced to proving that, when E is $(n+1)$-pseudo-coherent and F n-pseudo-coherent, then G is n-pseudo-coherent. After localizing, we can assume that E (resp. F) is strictly $(n+1)$-pseudo-coherent (resp. n-pseudo-coherent). The arrow $E \xrightarrow{u} F$ of $\mathrm{D}^-(\underline{C}_S)$ is defined by a pair of arrows of $\mathrm{C}^-(\underline{C}_S)$



where s is a quasi-isomorphism. By (2.2 a)), there exists, after localizing, a quasi-isomorphism $t : E'' \to E'$, where E'' is strictly $(n+1)$-pseudo-coherent. We are therefore reduced to the case where u is defined by an [print 100]"actual" arrow of $\mathrm{C}^-(\underline{C}_S)$. The assertion then follows from the definitions and the formula $C(u)^i = E^{i+1} \oplus F^i$, where $C(u)$ denotes the cone of u.

Scholium 2.6. By "rotation" of the triangles, we see that pseudo-coherence properties of two of the objects of the triangle $(*)$ imply another such property of the third. The different cases can be summarized in the table below (the lowercase letters denote degrees of pseudo-coherence and each row corresponds to an implication whose conclusion is underlined):

E	F	G
$n+1$	n	$\underline{n}$
n	$\underline{n}$	n
$\underline{n}$	n	$n-1$.

We shall denote by $\mathrm{D}(\underline{C})_{\underline{C}_0\text{-}n\text{-coh}}$ (resp. $\mathrm{D}(\underline{C})_{\underline{C}_0\text{-coh}}$) the strictly full fibered subcategory of $\mathrm{D}(\underline{C})$ whose fibre at each $X \in \mathrm{ob}\,\underline{S}$ consists of the objects of $\mathrm{D}(\underline{C}_X)$ that are n-pseudo-coherent (resp. pseudo-coherent) relative to $\underline{C}_0$. By (2.5), the category $\mathrm{D}(\underline{C})_{\underline{C}_0\text{-coh}}$ is a triangulated sub-$\underline{S}$-category of $\mathrm{D}(\underline{C})$ (1.1).

Proposition 2.7.[91] Assume that $\underline{S}$ is a punctual site, so that $\underline{C} = \underline{C}_S$, $\underline{C}_0 = \underline{C}_{0S}$. Then:

a) When $F \in \mathrm{ob}\,\mathrm{D}(\underline{C})$ is pseudo-coherent, there exists a quasi-isomorphism $E \longrightarrow F$, with E strictly pseudo-coherent (2.1).

b) Denote by $\mathrm{K}^{-,\varnothing}(\underline{C}_0)$ the full triangulated subcategory of $\mathrm{K}^-(\underline{C}_0)$ consisting of the acyclic complexes. The canonical functor

$$\mathrm{K}^-(\underline{C}_0)/\mathrm{K}^{-,\varnothing}(\underline{C}_0) \longrightarrow \mathrm{D}(\underline{C})$$

is fully faithful and has essential image $\mathrm{D}(\underline{C})_{\underline{C}_0\text{-coh}}$.

Proof. a) Let $F \in \mathrm{ob}\,\mathrm{D}(\underline{C})_{\underline{C}_0\text{-coh}}$, and suppose that an i-quasi-isomorphism (of $\mathrm{C}(\underline{C})$) $u_i : E_i \longrightarrow F$ has been constructed with E_i strictly pseudo-coherent (this is possible for i sufficiently large

with $E_i = 0$). Let G_i be the cone of u_i. It follows from (1.4.2) and (2.1.1) that $\mathrm{H}^{i-1}(G_i)$ is of $\underline{C}_0$-finite type. Thus, by (1.4.4), there exists a commutative triangle of $\mathrm{C}(\underline{C})$

$$\begin{array}{ccc} E_i & \xrightarrow{v_{i-1}} & E_{i-1} \\ {\scriptstyle u_i}\downarrow & \swarrow_{u_{i-1}} & \\ F & & \end{array},$$

where E_{i-1} is strictly pseudo-coherent, u_{i-1} an $(i-1)$-quasi-isomorphism, and v_{i-1} a monomorphism inducing the identity in degree $\geq i$. By passage to the limit (cf. proof of (1.4.5)), we deduce a quasi-isomorphism $u : E \to F$ with E strictly pseudo-coherent.

b) Apply the criterion ([V] chap I § 2 4.2 b)(i)) and a).

Observe that the proof of a) gives the following more precise result, which generalizes (1.4.5):

Proposition 2.7.1. Under the hypotheses of (2.7), let $u : E \to F$ be an n-quasi-isomorphism of $\mathrm{C}(\underline{C})$, with F pseudo-coherent and E strictly pseudo-coherent. There then exists a commutative triangle of $\mathrm{C}(\underline{C})$

$$\begin{array}{ccc} E & \xrightarrow{v} & E' \\ {\scriptstyle u}\downarrow & \swarrow_{u'} & \\ F & & \end{array},$$

where E' is strictly pseudo-coherent, u' a quasi-isomorphism, and v a [print 102]monomorphism inducing the identity in degree $\geq n$.

Definition 2.8. Let $n \in \mathbb{N}$ and $F \in \mathrm{ob}\,\underline{C}_S$. We shall say that F is of n-$\underline{C}_0$-finite presentation when the set of X over S such that there exists an exact sequence

$$E^{-n} \longrightarrow \ldots \longrightarrow E^0 \longrightarrow F_X \longrightarrow 0,$$

with $E^i \in \mathrm{ob}\,\underline{C}_{0X}$ for $-n \leq i \leq 0$, is a refinement of S. We shall say that F is of ∞-$\underline{C}_0$-finite presentation when F is of n-$\underline{C}_0$-finite presentation for every $n \in \mathbb{N}$.

Of course, we shall sometimes simply say n-finite presentation when no ambiguity concerning $\underline{C}_0$ is possible.

We shall say "of finite presentation" for "of 1-finite presentation", and "of finite type" for "of 0-finite presentation", the latter notion coinciding with the one already introduced in (1.2).

Proposition 2.9. Let $n \in \mathbb{N}$ and $F \in \mathrm{ob}\,\underline{C}_S$. Equivalent conditions:

(i) F is of n-finite presentation (resp. of ∞-finite presentation);

(ii) F is $(-n)$-pseudo-coherent (resp. pseudo-coherent).

In the case where S is a punctual site, F is pseudo-coherent if and only if there exists an exact sequence

$$\ldots \longrightarrow E^i \longrightarrow \ldots \longrightarrow E^0 \longrightarrow F \longrightarrow 0,$$

with $E^i \in \mathrm{ob}\,\underline{C}_0$ for every i.

Proof. For the equivalence (i) $\Leftrightarrow$ (ii), it clearly suffices to treat the case without the "resp." alternatives. The implication (i) $\Rightarrow$ (ii) follows immediately from the definitions, and [print 103]the implication (ii) $\Rightarrow$ (i) follows at once from (2.4): the arrow $0 \to F$ is a 1-quasi-isomorphism; we deduce (after localizing) a quasi-isomorphism $E \longrightarrow F$ with E strictly $(-n)$[92]-pseudo-coherent and $E^i = 0$ for $i \geq 1$, i.e. an n-finite presentation of F. The second assertion is proved in a similar way (use (2.7.1)).

Proposition 2.10. a) Let $m, n \in \mathbb{Z}$, with $n \leq m$. When $F \in \mathrm{ob}\,\mathrm{D}^-(\underline{C}_S)$ is n-pseudo-coherent and acyclic in degree $\geq m$, there locally exists a quasi-isomorphism $E \longrightarrow F$, with E strictly n-pseudo-coherent and $E^i = 0$ for $i \geq m$. When S is punctual, and F pseudo-coherent and acyclic in degree $\geq m$, there exists a quasi-isomorphism $E \longrightarrow F$ with E strictly pseudo-coherent and $E^i = 0$ for $i \geq m$.

b) Let $F \in \mathrm{ob}\,\mathrm{D}^-(\underline{C}_S)$ be acyclic in degree $> n$. For F to be n-pseudo-coherent, it is necessary and sufficient that $\mathrm{H}^n(F)$ be of finite type.

Proof. a) Since F is acyclic in degree $\geq m$, the arrow $0 \to F$ is an m-quasi-isomorphism. The assertion follows from (2.4) and (2.7.1).

b) There exists a distinguished triangle of $\mathrm{D}^-(\underline{C}_S)$:

$$\begin{array}{ccc} & F' & \\ {\scriptstyle +1}\swarrow & & \nwarrow \\ F & \longrightarrow & \mathrm{H}^n(F)[-n] \end{array} \quad ,$$

(to see this, begin by truncating F, replacing F^n by $Z^n(F)$ and F^i by 0 for $i > n$). The exact cohomology sequence implies $\mathrm{H}^i(F') = 0$ for $i \geq n-1$, so F' is $(n-1)$-pseudo-coherent. We conclude using (2.5) and (2.6).

[print 104]**Proposition 2.11.** Let $n \in \mathbb{Z}$.

a) Let $F \in \mathrm{ob}\,\mathrm{C}^-(\underline{C}_S)$. When, for every $i \geq n$ (resp. for every i), F^i is of $(i-n)$-finite presentation (resp. of ∞-finite presentation), then F is n-pseudo-coherent (resp. pseudo-coherent).

b) Let $F \in \mathrm{ob}\,\mathrm{D}^-(\underline{C}_S)$. When, for every $i \geq n$ (resp. for every i), $\mathrm{H}^i(F)$ is of $(i-n)$-finite presentation (resp. of ∞-finite presentation), then F is n-pseudo-coherent (resp. pseudo-coherent).[94]

Proof. It suffices to treat the assertions without the "resp." alternatives.

a) We argue by induction on N such that $F^i = 0$[93] for $i > n+N$. For $N = 0$, the assertion follows at once from (2.10 b)). Suppose $N \geq 1$. There is an exact sequence

$$0 \longrightarrow F^{n+N}[-n-N] \longrightarrow F \longrightarrow F' \longrightarrow 0,$$

By the induction hypothesis, F' is n-pseudo-coherent; on the other hand, $F^{n+N}[-n-N]$ is n-pseudo-coherent by hypothesis (2.9 and 2.5.a)); thus, by (2.5.b)), F is n-pseudo-coherent.

b) We argue by induction on N such that $\mathrm{H}^i(F) = 0$ for $i > n+N$. For $N = 0$, the assertion follows from (2.10 b)). Suppose $N \geq 1$. There is a distinguished triangle of $\mathrm{D}^-(\underline{C}_S)$ (cf. proof of (2.10.b))):

$$\begin{array}{ccc} & F' & \\ {\scriptstyle +1}\swarrow & & \nwarrow \\ F & \longrightarrow & \mathrm{H}^{n+N}(F)[-n-N] \end{array} \quad .$$

The exact cohomology sequence implies

$$\mathrm{H}^i(F') = 0 \quad \text{for } i \geq n+N-1, \quad \text{i.e. } i > (n-1)+(N-1),$$

and

$$\mathrm{H}^i(F') \xrightarrow{\sim} \mathrm{H}^{i+1}(F) \quad \text{for } i < n+N-1.$$

[print 105]By the induction hypothesis (applied with $n-1$ in place of n), F' is therefore $(n-1)$-pseudo-coherent. On the other hand, $\mathrm{H}^{n+N}(F)[-n-N]$ is n-pseudo-coherent (2.9 and 2.5.a)). Thus, by (2.6), F is n-pseudo-coherent.

Remark 2.11.1. When F is pseudo-coherent, the F^i (resp. $\mathrm{H}^i(F)$) are in general not pseudo-coherent, nor even of finite type. However, in the next paragraph, in the sorites on notions of coherence, we shall see important examples where the pseudo-coherence of F implies that of the $\mathrm{H}^i(F)$.

Proposition 2.12. Let $F', F'' \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$, and $F = F' + F''$. For F to be n-pseudo-coherent (resp. pseudo-coherent), it is necessary and sufficient that F' and F'' be n-pseudo-coherent (resp. pseudo-coherent).

Proof. It suffices to prove the assertion without the "resp." alternatives. Sufficiency is clear; let us prove necessity. By induction, we may suppose we have already proved that F' and F'' are $(n+1)$-pseudo-coherent. After localizing, we can even suppose that F' and F'' are strictly $(n+1)$-pseudo-coherent and that $F = F' + F''$ in $\mathrm{C}^-(\underline{C}_S)$. Denote by $F'_{>n}$ (resp. $F'_{\leq n}$) the complex obtained from F' by replacing F'^i by 0 for $i \leq n$ (resp. $i > n$), and similarly $F''_{>n}$ (resp. $F''_{\leq n}$)... There is an exact sequence

$$0 \longrightarrow F'_{>n} + F''_{>n} \longrightarrow F \longrightarrow F'_{\leq n} + F''_{\leq n} \longrightarrow 0.$$

$F'_{>n} + F''_{>n}$ is strictly perfect and F is n-pseudo-coherent, so (2.6) $F'_{\leq n} + F''_{\leq n}$ is n-pseudo-coherent. By (2.10 b)), this means that $\mathrm{H}^n(F'_{\leq n} + F''_{\leq n})$ is of finite type. But we have

$$\mathrm{H}^n(F'_{\leq n} + F''_{\leq n}) = \mathrm{H}^n(F'_{\leq n}) + \mathrm{H}^n(F''_{\leq n}),$$

[print 106]so $\mathrm{H}^n(F'_{\leq n})$ (resp. $\mathrm{H}^n(F''_{\leq n})$) is of finite type, so again, by (2.10.b)), $F'_{\leq n}$ (resp.[95] $F''_{\leq n}$) is n-pseudo-coherent. Applying (2.5) to the exact sequence

$$0 \longrightarrow F'_{>n} \longrightarrow F' \longrightarrow F'_{\leq n} \longrightarrow 0,$$

(resp.. . .), we deduce that F' (resp. F'') is n-pseudo-coherent, QED.

Proposition 2.13. Let $(\underline{S}', \underline{C}', \underline{C}'_0)$ be a triple satisfying the same hypotheses (2.0) as $(\underline{S}, \underline{C}, \underline{C}_0)$. Let $f : \underline{S}' \to \underline{S}$ be a morphism of sites ([G] I 1.4), corresponding to an underlying functor $f^{-1} : \underline{S} \to \underline{S}'$, and let $u : \mathrm{D}^*(\underline{C}) \longrightarrow \mathrm{D}^*(\underline{C}')$ $(* = +, -, \varnothing)$ be a cartesian functor over f, i.e. a cartesian $\underline{S}$-functor $\mathrm{D}^*(\underline{C}') \times_{\underline{S}'} \underline{S} \longleftarrow \mathrm{D}^*(\underline{C})$. Assume that u is exact on each fibre and that there exist $a, b \in \mathbb{Z}$ such that the following conditions hold for every $S \in \mathrm{ob}\,\underline{S}$:

(i) when $E \in \mathrm{ob}\,\underline{C}_{0S}$, $u(E)$ is a-$\underline{C}'_0$-pseudo-coherent (resp. $\underline{C}'_0$-pseudo-coherent)

(ii) when $F \in \mathrm{ob}\,\mathrm{D}^*(\underline{C}_S)$ is acyclic in degree > 0, $u(F)$ is acyclic in degree $> b$.

Then, when $F \in \mathrm{ob}\,\mathrm{D}^*(\underline{C}_S)$ is m-pseudo-coherent rel. to $\underline{C}_0$ and acyclic in degree $> n$, $u(F) \in \mathrm{ob}\,\mathrm{D}^*(\underline{C}'_{S'})$ (where $S' = f^{-1}(S)$) is $\sup(m+b, n+a)$-pseudo-coherent (rel. to $\underline{C}'_0$) (resp. $(m+b)$-pseudo-coherent) and acyclic in degree $> n+b$.

In particular, in the case of the "resp." alternative, when $F \in \mathrm{ob}\,\mathrm{D}^*(\underline{C}_S)$ is pseudo-coherent (rel. to $\underline{C}_0$), $u(F)$ is pseudo-coherent (rel. to $\underline{C}'_0$).

Proof. We can restrict ourselves to the case without the "resp." alternative. Let us first show that, for $F \in \mathrm{ob}\,\mathrm{D}^*(\underline{C}_S)$ strictly perfect (2.1) with $F^i = 0$ for $i > n$, $u(F)$ is $(n+a)$-$\underline{C}'_0$-pseudo-coherent. We argue by induction on the number N of indices i such that $F^i \neq 0$. For $N = 1$, [print 107]we have $F = F^p[-p]$ for some $p \leq n$, so $u(F)$ is $(p+a)$-$\underline{C}'_0$-pseudo-coherent, hence a fortiori $(n+a)$-pseudo-coherent. For $N > 1$, there is an exact sequence

$$0 \longrightarrow F' \longrightarrow F \longrightarrow F'' \longrightarrow 0,$$

where F' and F'' are strictly perfect of length $\leq N-1$ and satisfy $F'^i = F''^i = 0$ for $i > n$. Applying (2.5.b)) to the distinguished triangle

$$u(F'') \xrightarrow{+1} u(F') \longrightarrow u(F) \quad ,$$

we deduce from the induction hypothesis that $u(F)$ is $(n+a)$-pseudo-coherent.

Let us now pass to the general case: F m-$\underline{C}_0$-pseudo-coherent and acyclic in degree $> n$. After localizing, we can, by (2.10.a)), suppose that F is strictly m-pseudo-coherent and satisfies $F^i = 0$ for $i > n$. There is then an exact sequence

$$0 \longrightarrow F' \longrightarrow F \longrightarrow F'' \longrightarrow 0,$$

where F'' is zero in degree $\geq m$, and F' strictly perfect and zero in degree $> n$. In the distinguished triangle

$$u(F'') \xrightarrow{+1} u(F') \longrightarrow u(F) \quad ,$$

$u(F'')$ is acyclic in degree $\geq m+b$, hence $(m+b)$-pseudo-coherent, and $u(F')$ is $(n+a)$-pseudo-coherent by what we have seen. We conclude by (2.5.b)).

Corollary 2.14. Let $\underline{C}_1$ be a strictly full additive sub-$\underline{S}$-category of $\underline{C}$ such that $\underline{C}_0 \subset \underline{C}_1 \subset \underline{C}$. Assume that every object of $\underline{C}_1$ is $\underline{C}_0$-pseudo-coherent. [print 108]Then:

(i) $\underline{C}_1$ is locally quasi-liftable in $\underline{C}$ (1.2) and locally quasi-stable under kernels of epimorphisms (1.2).

(ii) Let $S \in \operatorname{ob} \underline{S}$ and $F \in \operatorname{ob} \mathrm{D}(\underline{C}_S)$. For F to be n-pseudo-coherent (resp. pseudo-coherent) relative to $\underline{C}_1$, it is necessary and sufficient that it be so relative to $\underline{C}_0$.

Proof. (i) Let $E \to F$ be an epimorphism of $\underline{C}_S$, with $F \in \operatorname{ob} \underline{C}_{1S}$. Since F is $\underline{C}_0$-pseudo-coherent (hence a fortiori of $\underline{C}_0$-finite type), there exists (after localizing) an epimorphism $E' \to F$ with $E' \in \operatorname{ob} \underline{C}_{0S}$. Using the quasi-liftability of $\underline{C}_0$ in $\underline{C}$, we can suppose (after localizing again) that this factors through E, whence the quasi-liftability of $\underline{C}_1$. On the other hand, when $E \in \operatorname{ob} \underline{C}_S$ admits a finite right resolution by objects of $\underline{C}_1$, E is $\underline{C}_0$-pseudo-coherent by (2.11.a)), hence a fortiori of $\underline{C}_1$-finite type.

(ii) It suffices to prove necessity. For this we can apply, as we choose, (2.11.a)) or (2.13) with $\underline{S}' = \underline{S}$, $\underline{C}' = \underline{C}$, $f =$ Identity, $u =$ Identity, replacing $\underline{C}'_0$ and $\underline{C}_0$ respectively by $\underline{C}_0$ and $\underline{C}_1$.[96]

Scholium 2.14.1. There thus exists a largest strictly full additive sub-$\underline{S}$-category of $\underline{C}$ giving the same notion of n-pseudo-coherence (resp. pseudo-coherence) as $\underline{C}_0$, namely the category formed by the $\underline{C}_0$-pseudo-coherent objects. The latter is locally stable under kernels of epimorphisms.

Example 2.15. Let $(\underline{S}, A)$ be a ringed topos and $\underline{C}, \underline{C}_0$ the $\underline{S}$-categories considered in (1.3.d)). The fibered category $\mathrm{D}(\underline{C})$ will often be denoted by $\mathrm{D}(\underline{S}, A)$, or $\mathrm{D}({}_A\underline{S})$ (to [print 109]suggest that these are left A-Modules), or even $\mathrm{D}(\underline{S})$ when there is no ambiguity concerning A. We shall denote by S the final object of $\underline{S}$ and put $\mathrm{D}(\underline{C}_S) = \mathrm{D}({}_AS)$ (or $\mathrm{D}(S)$). An object of $\mathrm{D}(\underline{S})$ will be called $\underline{n}$-pseudo-coherent (resp. pseudo-coherent) when it is n-$\underline{C}_0$-pseudo-coherent (resp. $\underline{C}_0$-pseudo-coherent) (likewise for the strict notions). The fibered category $\mathrm{D}(\underline{C})_{n\text{-}\underline{C}_0\text{-}\mathrm{coh}}$ (resp. $\mathrm{D}(\underline{C})_{\underline{C}_0\text{-}\mathrm{coh}}$) (2.6) will be denoted by $\mathrm{D}({}_A\underline{S})_{n\text{-}\mathrm{coh}}$ (resp. $\mathrm{D}({}_A\underline{S})_{\mathrm{coh}}$). Observe that, by (2.14), we obtain the same notion of pseudo-coherence on replacing $\underline{C}_0$ by the $\underline{S}$-category $\underline{C}_1$ defined by

$U \longmapsto$ (category of left A_U-Modules locally direct factors of free Modules of finite type). When an A-Module M is a direct factor of a free A-Module of finite type, it does not follow, in general, that M is locally free of finite type. However, we have:

Lemma 2.15.1. Assume that $\underline{S}$ satisfies one of the following two hypotheses:

(i) $\underline{S}$ "has enough points" (SGA 4 IV 6.4) and, for every point s of $\underline{S}$, the projective A_s-modules (on the left) of finite type are free, this last condition being satisfied for example (BOURBAKI Alg. Comm. chap. II § 3 n° 2 cor. 2 of prop. 5) when $A_s/\mathrm{radical}(A_s)$ is a field.[97]

(ii) $(\underline{S}, A)$ is a "locally ringed" topos in the sense of [1], which means that A is commutative and that, for every $X \in \mathrm{ob}\,\underline{S}$ and every $f \in \Gamma(X, A)$, we have $X = X_f \cup X_{1-f}$, where X_f (resp. X_{1-f}) denotes the largest subobject of X on which f (resp. $1-f$) is invertible.

Then every direct factor of a free A-module (on the left) of finite type is locally free of finite type.

Proof. Suppose (i). We first observe that, for two A-Modules E and F, with E of finite presentation, the canonical arrow [print 110]

$$\underline{\mathrm{Hom}}(E, F)_s \longrightarrow \mathrm{Hom}(E_s, F_s)$$

is an isomorphism for every point s of $\underline{S}$. It follows that, when E and F are both of finite presentation, and $E_s \simeq F_s$ at a point s of $\underline{S}$, there exists an s-pointed object U of $\underline{S}$(*) such that $E|U \simeq F|U$. The conclusion follows at once.

Suppose (ii). Let[98] $n \in \mathbb{N}$, and denote by $\mathbb{D}^n$ the functor on the category of locally ringed topoi (belonging to a fixed universe...) defined by $T \longmapsto \Gamma(T, \underline{O}_T)^n$. We know [1] that the functor $\underline{\mathrm{End}}_{\mathbb{D}}(\mathbb{D}^n)$ is represented by $M_n = \mathrm{Spec}\,\mathbb{Z}[T_{ij}]$ $(1 \leq i \leq n,\ 1 \leq j \leq n)$, equipped with the endomorphism u_n[99] of $(\underline{O}_{M_n})^n$ defined by the matrix (T_{ij}). Now let $p : A^n \longrightarrow A^n$ be a projector; let us show that $p(A^n)$ is locally free of finite type. By what we have just said, there exists a morphism $s : \underline{S} \longrightarrow M_n$ of locally ringed topoi such that $s^*(u_n) = p$. We can interpret s as a point of M_n with values in $\Gamma(S, A)$, i.e. a matrix of order n with coefficients in $\Gamma(S, A)$; since this matrix is a projector, s factors through the closed subscheme $i : P_n \to M_n$ corresponding to projectors as $\underline{S} \xrightarrow{s'} P_n \xrightarrow{i} M_n$. Thus $p = s'^* i^*(u_n)$, and we are done, because by the assertion under hypothesis (i), the image of $i^*(u_n)$ is a locally free sheaf of finite type.

Corollary 2.16. Let $f : (\underline{S}', A') \to (\underline{S}, A)$ be a morphism of ringed topoi (defined by a functor $f^{-1} : \underline{S} \to \underline{S}'$ and a morphism of sheaves of rings $f^{-1}(A) \to A'$). Denote by $\mathrm{D}({}_{A'}\underline{S}'_A)$ the derived category of the category of A'-A-biModules (on the left over A', on the right over A), and by

$$\otimes^{\mathrm{L}}_A : \mathrm{D}^-({}_{A'}\underline{S}'_A) \times_{\underline{S}'} \mathrm{D}^-({}_A\underline{S}) \longrightarrow \mathrm{D}^-({}_{A'}\underline{S}')$$

[print 111]the derived functor of the tensor product $(E', E) \longmapsto E' \otimes_{f^{-1}(A)} f^{-1}(E)$. Let $X \in \mathrm{ob}\,\underline{S}$, $X' = f^{-1}(X)$[100], $E' \in \mathrm{ob}\,\mathrm{D}^-({}_{A'}X'_A)$, $E \in \mathrm{ob}\,\mathrm{D}^-({}_AX)$. Then, when E' is m'-pseudo-coherent (as an object of $\mathrm{D}^-({}_{A'}\underline{S}')$) and acyclic in degree $> n'$ and E is m-pseudo-coherent and acyclic in degree $> n$, $E' \otimes^{\mathrm{L}}_A E$ is $\sup(m + n', m' + n)$-pseudo-coherent and acyclic in degree $> n + n'$.

Proof. Replacing $\underline{S}$ by $\underline{S}_{/X}$, we can suppose that $X = S$, and we apply (2.13) with $u = E' \otimes^{\mathrm{L}}_A : \mathrm{D}^-({}_A\underline{S}) \to \mathrm{D}^-({}_{A'}\underline{S}')$. Let $U \in \mathrm{ob}\,\underline{S}$, $U' = f^{-1}(U)$, and E be a complex of left A_U-Modules. When E is acyclic in degree $> n$, $E' \otimes^{\mathrm{L}}_A E$ is acyclic in degree $> n + n'$ (definition of $\otimes^{\mathrm{L}}_A$); on the other hand, for $E = (A_U)^p$, we have $E' \otimes^{\mathrm{L}}_A E = (E'|U')^p$, which is m'-pseudo-coherent by hypothesis (and (2.12)). The conditions of (2.13) are therefore satisfied, with $a = m'$, $b = n'$, whence the conclusion.

In particular:

(*) i.e. such that s factors through $\underline{S}_{/U}$.

Corollary 2.16.1. Under the hypotheses of (2.16), we have:

a) The functor $\mathbb{L}f^* : \mathrm{D}^-({}_A\underline{S}) \to \mathrm{D}^-({}_{A'}\underline{S}')$ (which is simply the functor $A' \otimes^{\mathrm{L}}_A$) induces a functor

$$\mathbb{L}f^* : \mathrm{D}^-({}_A\underline{S})_{\mathrm{coh}} \longrightarrow \mathrm{D}^-({}_{A'}\underline{S}')_{\mathrm{coh}}$$

(notation of (2.6)).

b) When A is commutative, the functor $\otimes^{\mathrm{L}}_A : \mathrm{D}^-(\underline{S}) \times_{\underline{S}} \mathrm{D}^-(\underline{S}) \longrightarrow \mathrm{D}^-(\underline{S})$ induces a functor

$$\otimes^{\mathrm{L}}_A : \mathrm{D}^-(\underline{S})_{\mathrm{coh}} \times_{\underline{S}} \mathrm{D}^-(\underline{S})_{\mathrm{coh}} \longrightarrow \mathrm{D}^-(\underline{S})_{\mathrm{coh}}.$$

[print 112]**Exercise 2.16.2.** Let $\underline{S}$ be a topos, and $A \to A'$ a morphism of Rings of $\underline{S}$. We make one of the following hypotheses:

(i) $A \to A'$ is surjective with kernel a nilpotent Ideal

(ii) the functor $A' \otimes_A -$ on the category of left A-modules is exact and faithful.

Show that $E \in \mathrm{ob}\,\mathrm{D}^-({}_A\underline{S})$ is n-pseudo-coherent (resp. pseudo-coherent) if and only if $A' \otimes^{\mathrm{L}}_A E \in \mathrm{ob}\,\mathrm{D}^-({}_{A'}\underline{S})$ is n-pseudo-coherent (resp. pseudo-coherent).

Hints: First show that a left A-Module L is of finite type if and only if $A' \otimes_A L$ is (cf. Bourbaki, Alg. commutative, chap. I, § 3, prop. 11 and chap. II, § 3, prop. 4), then reduce to this case by truncation, using (2.5) and (2.10 b)).

3. Link with the classical notion of coherence

3.0. The hypotheses and notation are those of (2.0), except that $\underline{C}_0$ is no longer assumed locally quasi-stable under kernels of epimorphisms.

Definition 3.1. Let $F \in \mathrm{ob}\,\underline{C}_S$. We shall say that F is precoherent (relative to $\underline{C}_0$)(*) when, for every U over S and every morphism $u : E \to F|U$ with $E \in \mathrm{ob}\,\underline{C}_{0U}$, $\mathrm{Ker}(u)$ is of $\underline{C}_0$-finite type (1.2). We shall say that F is coherent (relative to $\underline{C}_0$) when F is precoherent and of finite type.

Remark 3.2. Let $F \in \mathrm{ob}\,\underline{C}_S$. It is clear from the definitions that, when F is precoherent (resp. of finite type), every subobject (resp. quotient) of F is precoherent (resp. of finite type).

Lemma 3.3.[101] a) Let

$$0 \longrightarrow F \longrightarrow G \longrightarrow H$$

be an exact sequence of $\underline{C}_S$. When G is of finite type and H precoherent, F is of finite type.

b) Let

$$F \longrightarrow G \longrightarrow H \longrightarrow 0$$

be an exact sequence of $\underline{C}_S$. When F is of finite type and G precoherent, H is precoherent.

c) Let

$$0 \longrightarrow F \longrightarrow G \longrightarrow H \longrightarrow 0$$

be an exact sequence of $\underline{C}_S$. When F and H are precoherent (resp. of finite type), G is too.

(*) We introduce this notion here as a technical intermediate step; the only notion to be retained is that of coherence.

Proof. a) After localizing, we have an epimorphism $G' \to G$ with $G' \in \operatorname{ob} \underline{C}_0$. Hence a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & F' & \longrightarrow & G' & \longrightarrow & H \\ & & \downarrow & & \downarrow & & \downarrow{\scriptstyle \mathrm{Id}} \\ 0 & \longrightarrow & F & \longrightarrow & G & \longrightarrow & H. \end{array}$$

F' is of finite type because H is precoherent, but $F' \to F$ is an epimorphism, so F is of finite type.

b) Let U be over S, and $H' \xrightarrow{f} H|U$ a morphism of $\underline{C}_U$ with $H' \in \operatorname{ob} \underline{C}_{0U}$. Changing notation, we can suppose that $U = S$. By the quasi-liftability of $\underline{C}_0$ in $\underline{C}$, there exists (after localizing) a commutative square

$$\begin{array}{ccc} G' & \xrightarrow{p} & H' \\ \downarrow & & \downarrow{\scriptstyle f} \\ G & \longrightarrow & H, \end{array}$$

where the horizontal arrows are epimorphisms and $G' \in \operatorname{ob} \underline{C}_0$. Since $\operatorname{Ker}(f)$ is a quotient of $\operatorname{Ker}(fp)$, it suffices to prove that $\operatorname{Ker}(fp)$ is of finite type, so we can suppose that $H' = G'$. The arrows $F \to G$ and $H' \to G$ define $g : F \oplus H' \to G$. Putting $K = \operatorname{Ker}(G \to H)$, we therefore have a commutative diagram in which the rows are exact and q is an epimorphism:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & F & \longrightarrow & F \oplus H' & \longrightarrow & H' & \longrightarrow & 0 \\ & & {\scriptstyle q}\downarrow & & \downarrow{\scriptstyle g} & & \downarrow{\scriptstyle f} & & \\ 0 & \longrightarrow & K & \longrightarrow & G & \longrightarrow & H & \longrightarrow & 0. \end{array}$$

The snake diagram shows that $\operatorname{Ker}(g) \to \operatorname{Ker}(f)$ is an epimorphism. But, since $F \oplus H'$ is of finite type and G precoherent, $\operatorname{Ker}(g)$ is of finite type by a), and we are done.

c) Let us first prove the assertion without the "resp." alternative. Up to a change of notation, we must show that, for a morphism $g : G' \to G$ of $\underline{C}_S$ with $G' \in \operatorname{ob} \underline{C}_0$, $\operatorname{Ker}(g)$ is of finite type. Denoting by H' the image of the composite arrow $G' \to H$, we have a commutative diagram whose rows are exact

$$\begin{array}{ccccccccc} 0 & \longrightarrow & F' & \longrightarrow & G' & \longrightarrow & H' & \longrightarrow & 0 \\ & & {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle g} & & \downarrow{\scriptstyle h} & & \\ 0 & \longrightarrow & F & \longrightarrow & G & \longrightarrow & H & \longrightarrow & 0. \end{array}$$

Since h is a monomorphism, the snake diagram gives an isomorphism $\operatorname{Ker}(f) \xrightarrow{\sim} \operatorname{Ker}(g)$. But F' is of finite type by a), so (again by a)) $\operatorname{Ker}(f)$ is of finite type, which proves the assertion without the "resp." alternative. Let us pass to the "resp." assertion (strictly speaking, we cannot invoke (2.5 b)) because we do not assume that $\underline{C}_0$ is locally quasi-stable under kernels of epimorphisms). After localizing, there exist epimorphisms $F' \xrightarrow{u} F$, $H' \xrightarrow{v} H$, with $F' \in \operatorname{ob} \underline{C}_0$, $H' \in \operatorname{ob} \underline{C}_0$. After localizing further, we can suppose that v factors through G, whence a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & F' & \longrightarrow & F' \oplus H' & \longrightarrow & H' & \longrightarrow & 0 \\ & & {\scriptstyle u}\downarrow & & \downarrow{\scriptstyle w} & & \vdots{\scriptstyle v} & & \\ 0 & \longrightarrow & F & \longrightarrow & G & \longrightarrow & H & \longrightarrow & 0. \end{array}$$

Since u and v are epimorphisms, so is w, and we are done. This completes the proof of (3.3).

We deduce immediately the

Corollary 3.4. The (strictly) full subcategory of $\underline{C}_S$ formed by the coherent objects is stable under finite projective (resp. inductive) limits. In particular, the kernel, cokernel and image of an arrow between two coherent objects are coherent. Let

$$E^0 \longrightarrow E^1 \longrightarrow E^2 \longrightarrow E^3 \longrightarrow E^4$$

be an exact sequence of $\underline{C}_S$. When E^0, E^1, E^3, E^4 are coherent, E^2 is too.

Corollary 3.5. Assume that the objects of $\underline{C}_0$ are $\underline{C}_0$-coherent (which implies in particular that $\underline{C}_0$ is locally quasi-stable under kernels of epimorphisms, i.e. that the conditions of (2.0) hold).

a) Equivalent conditions on $F \in \mathrm{ob}\,\underline{C}_S$:

(i) F is of finite presentation;

(ii) F is coherent;

(iii) F is of ∞-finite presentation (2.8).

b) Let $F \in \mathrm{ob}\,\mathrm{D}^-(\underline{C}_S)$, $n \in \mathbb{Z}$. We have:

$$F \text{ } n\text{-pseudo-coherent} \iff \mathrm{H}^i(F) \text{ coherent for } i \geq n+1, \text{ and } \mathrm{H}^n(F) \text{ of finite type.}$$

[print 116]In particular, for F to be pseudo-coherent, it is necessary and sufficient that $\mathrm{H}^n(F)$ be coherent for every n.

Proof. Assertion a) and the implication $\Rightarrow$ of b) follow trivially from (3.4). The implication $\Leftarrow$ of b) follows at once from a) and (2.11.b)).

Examples 3.6. Let $(\underline{S}, A)$ be a ringed topos, and $\underline{C}, \underline{C}_0$ the $\underline{S}$-categories of (1.3.d)) (cf. also (2.15)). For the objects of $\underline{C}_0$ to be coherent, it is necessary and sufficient that A be coherent (i.e. precoherent). Here are examples where this holds:

- A is constant with value a left noetherian ring (SGA 4 IX 2.1).
- $\underline{S}$ is the topos of sheaves of sets (for the Zariski topology) on a locally noetherian scheme S(*), and $A = \underline{O}_S$.
- $\underline{S}$ is the topos of sheaves of sets on a complex analytic space S, $A = \underline{O}_S$.

On the other hand, the sheaf of continuous complex-valued functions on a topological space is in general not coherent.

Of course, when $(\underline{S}, A)$ is defined by a ringed space, the notion of coherence considered here coincides with Serre's usual notion (EGA 0_I 5.3.1).

Proposition 3.7. a) Let $f : (\underline{S}', A') \to (\underline{S}, A)$ be a morphism of ringed topoi. Assume that f is flat on the right, i.e. that the functor $f^* = A' \otimes_A .$ is exact. For $F \in \mathrm{ob}\,\mathrm{D}({}_A S)$, $G \in \mathrm{ob}\,\mathrm{D}^+({}_A S)$, there is a functorial canonical arrow:

$$f^* \mathbb{R}\underline{\mathrm{Hom}}(F, G) \longrightarrow \mathbb{R}\underline{\mathrm{Hom}}(f^*(F), f^*(G)).$$

[print 117]This is an isomorphism when F is pseudo-coherent.

(*) In accordance with the new conventions, we shall say "scheme" for "prescheme", and "separated scheme" for "scheme".

b) Suppose that A is commutative and coherent, and let $F \in \mathrm{ob}\,\mathrm{D}^-(S)$, $G \in \mathrm{ob}\,\mathrm{D}^+(S)$. If $\underline{\mathrm{H}}^i(F)$ and $\underline{\mathrm{H}}^i(G)$ are coherent for every i, the same is true of $\underline{\mathrm{Ext}}^i(F,G)$.

Proof. a) To define the arrow (cf. SGA 4 XVII), we can suppose that G has injective components, and take the composite arrow

$$f^*\underline{\mathrm{Hom}}(F,G) \longrightarrow \underline{\mathrm{Hom}}(f^*F, f^*G) \longrightarrow \mathbb{R}\underline{\mathrm{Hom}}(f^*F, f^*G).$$

Let us show that it is an isomorphism for F pseudo-coherent. For brevity, write T (resp. T') for the functor $f^*\mathbb{R}\underline{\mathrm{Hom}}(.,G)$ (resp. $\mathbb{R}\underline{\mathrm{Hom}}(f^*.,f^*G)$), and let $a \in \mathbb{Z}$ be such that $\mathrm{H}^i(G) = 0$ for $i < a$. It is clear that $TA \xrightarrow{\sim} T'A$, hence, by dévissage, $TF \xrightarrow{\sim} T'F$ for F strictly perfect (2.1). On the other hand, when F is acyclic in degree $> n$, TF and $T'F$ are acyclic in degree $< a - n$. To verify that $TF \xrightarrow{\sim} T'F$ for F pseudo-coherent, we can, since the question is local, suppose that F is strictly n-pseudo-coherent, i.e. that there is a distinguished triangle

$$\begin{array}{ccc} & F'' & \\ {\scriptstyle +1}\swarrow & & \nwarrow \\ F' & \longrightarrow & F \end{array} ,$$

with F' strictly perfect and F'' acyclic in degree $> n-1$. We deduce a morphism of distinguished triangles

$$\begin{array}{ccc} & TF'' & \\ & {\scriptstyle -1}\nwarrow\ \searrow & \\ TF' & \longleftarrow & TF \end{array} \longrightarrow \begin{array}{ccc} & T'F'' & \\ & {\scriptstyle -1}\nwarrow\ \searrow & \\ T'F' & \longleftarrow & T'F \end{array} .$$

Now $TF' \xrightarrow{\sim} T'F'$ since F' is strictly perfect, and TF'' and $T'F''$ are [print 118]acyclic in degree $\leq a - n$; writing the induced morphism on the exact cohomology sequences of the preceding triangles, we deduce that $TF \to T'F$ induces an isomorphism $\mathrm{H}^i(TF) \xrightarrow{\sim} \mathrm{H}^i(T'F)$ for $i \leq a - n$. Since we can choose n arbitrarily negative, we are done.

b) The proof is left as an exercise to the reader: see, for example, ([H] II 3.3).

Remark 3.7.1. We shall see below (7.1.2) another case of "compatibility" of $\mathbb{R}\underline{\mathrm{Hom}}$ with $\mathbb{L}f^*$.

Corollary 3.7.2. (cf. EGA 0_I 5.2.7). Let $(\underline{S}, A)$ be a ringed topos, s a point of $\underline{S}$ (SGA 4 VIII 7.8)[102], $E, F \in \mathrm{ob}\,\mathrm{D}^b({}_A S)_{\mathrm{coh}}$. If $E_s \simeq F_s$, there exists an s-pointed object U of $\underline{S}$ such that $E|U \simeq F|U$.

Proof. Let $\alpha : E_s \to F_s$ and $\beta : F_s \to E_s$ be reciprocal isomorphisms. By (3.7 a)),[103] we have

$$\underline{\mathrm{Hom}}(E,F)_s \xrightarrow{\sim} \mathrm{Hom}(E_s, F_s) \qquad (\text{resp. } \underline{\mathrm{Hom}}(F,E)_s \xrightarrow{\sim} \mathrm{Hom}(F_s, E_s)).$$

There are therefore a "neighbourhood" U of s and a section u (resp. v) of $\underline{\mathrm{Hom}}(E,F)$ (resp. $\underline{\mathrm{Hom}}(F,E)$) over U such that $u_s = \alpha$ (resp. $v_s = \beta$). The homomorphisms $vu : E|U \to E|U$ and $uv : F|U \to F|U$ induce the identity at s, hence over a suitable V over U, as required.

[print 119]

4. Perfect complexes

4.0. In this section, $\underline{S}$ denotes a site, S an object of $\underline{S}$, $\underline{C}$ a flat abelian $\underline{S}$-category, and $\underline{C}_0$ a strictly full additive sub-$\underline{S}$-category of $\underline{C}$ (1.1). Unless otherwise mentioned, we assume that $\underline{C}_0$ is locally liftable in $\underline{C}$ and locally stable under kernels of epimorphisms (1.2).

Lemma 4.1(*). Let $u : E \to F$ be an arrow of $\mathrm{C}(\underline{C}_S)$. If E is strictly $\underline{C}_0$-perfect (2.1) and F is acyclic, then u is locally homotopic to zero.

Proof. We must construct locally a homotopy k such that $u = dk + kd$. Since E is strictly perfect, there exist $a, b \in \mathbb{Z}$, with $a \leq b$, such that $E^i = 0$ for $i \notin [a, b]$. We can therefore take $k^i = 0$ for $i \notin [a, b]$. We then construct k^i for $i \geq a$ by descending induction on i. Suppose that $k^i : E^i \to F^{i-1}$ has been constructed so that $u^i = k^{i+1}d_E^i + d_F^{i-1}k^i$ for $i \geq n > a$, and let us show that, after localizing, there exists $k^{n-1} : E^{n-1} \to F^{n-2}$ such that $u^{n-1} = k^n d_E^{n-1} + d_F^{n-2}k^{n-1}$. By the induction hypothesis, the arrow

$$v^{n-1} = u^{n-1} - k^n d_E^{n-1} : E^{n-1} \to F^{n-1}$$

factors through $Z^{n-1}(F)$. But $Z^{n-1}(F) = B^{n-1}(F)$ since F is acyclic. Since $\underline{C}_0$ is locally liftable in $\underline{C}$, it follows that, after localizing, there exists $k^{n-1} : E^{n-1} \to F^{n-2}$ such that $v^{n-1} = d_F^{n-2}k^{n-1}$, as required.

Corollary 4.2. Let

$$\begin{array}{ccc} & & P \\ & & \downarrow{\scriptstyle v} \\ E & \xrightarrow{f} & F \end{array}$$

[print 120]be a diagram of $\mathrm{C}(\underline{C}_S)$, where P is strictly $\underline{C}_0$-perfect and f a quasi-isomorphism. The set of X over S for which there exists an arrow $u : P_X \to E_X$ of $\mathrm{C}(\underline{C}_X)$ such that the diagram

$$\begin{array}{ccc} & & P_X \\ & {\scriptstyle u}\swarrow & \downarrow{\scriptstyle v_X} \\ E_X & \xrightarrow{f_X} & F_X \end{array}$$

is commutative in $\mathrm{K}(\underline{C}_X)$ is a refinement of S.

Proof. Let G be the cone of f. We have the exact sequence

$$\longrightarrow \mathrm{Hom}_{\mathrm{K}(\underline{C}_S)}(P, E) \longrightarrow \mathrm{Hom}_{\mathrm{K}(\underline{C}_S)}(P, F) \longrightarrow \mathrm{Hom}_{\mathrm{K}(\underline{C}_S)}(P, G) \longrightarrow .$$

Since G is acyclic, the image of v in $\mathrm{Hom}_{\mathrm{K}(\underline{C}_S)}(P, G)$ is locally zero by (4.1). Thus, after localizing, there exists $u : P \to E$ such that $v = fu$ in $\mathrm{K}(\underline{C}_S)$, as required.

Corollary 4.3. Let $f : E \to F$ be a quasi-isomorphism of $\mathrm{C}(\underline{C}_S)$, with F strictly perfect (relative to $\underline{C}_0$). The set of X over S for which there exists a quasi-isomorphism $g : F_X \to E_X$ such that $f_X g$ is homotopic to Id_{F_X} is a refinement of S.

Proof. This is a special case of (4.2).

Scholium 4.4. Let $F \in \mathrm{ob}\,\mathrm{K}(\underline{C}_S)$. We can define (for example using a cleavage of $\underline{C}$ (SGA 1 VI 7.1)) a category $(\mathrm{Qis}\,/F)$ fibered over $\underline{S}_{/S}$, whose fibre at X over S is the category $(\mathrm{Qis}\,/F_X)^\circ$ (opposite to the category of quasi-isomorphisms with target F_X). We can then express (4.3) by saying that, [print 121]when F is strictly perfect, the arrow Id_F is locally cofinal in $(\mathrm{Qis}\,/F)$.

Notation 4.5. Let $\underline{M}$ be a fibered $\underline{S}$-category. Let $X \in \mathrm{ob}\,\underline{S}$, and $E, F \in \mathrm{ob}\,\underline{M}_X$. We shall denote by $\underline{\mathrm{Hom}}_{\underline{M}}(E, F)$ the sheaf on $\underline{S}_{/X}$ associated to the presheaf of $\underline{S}$-morphisms from E to F in $\underline{M}$ (for the definition of this presheaf, see [G] I 2.6; take care that, even when $\underline{C}$ is a stack ([G]), the corresponding presheaf for $\underline{M} = \mathrm{K}(\underline{C})$ (or $\mathrm{D}(\underline{C})$) is in general not a sheaf).

Corollary 4.6. Let $E, F \in \mathrm{ob}\,\mathrm{C}(\underline{C}_S)$, with E strictly perfect. The canonical arrow

$$\underline{\mathrm{Hom}}_{\mathrm{K}(\underline{C})}(E, F) \longrightarrow \underline{\mathrm{Hom}}_{\mathrm{D}(\underline{C})}(E, F)$$

(*) This lemma was communicated to the editor by D. Ferrand.

is an isomorphism.

Proof. This is an immediate consequence of (4.4) and of the formula

$$\mathrm{Hom}_{\mathrm{D}(\underline{C}_X)}(E_X, F_X) = \varinjlim_{(\mathrm{Qis}\,/E_X)^\circ} \mathrm{Hom}_{\mathrm{K}(\underline{C}_X)}(\,.\,, F_X)$$

for X over S.

Definition 4.7. Let $F \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$ and let $a, b \in \mathbb{Z}$ be such that $a \leq b$. We shall say that F has $\underline{C}_0$-perfect amplitude contained in the interval $[a, b]$, and we shall write

$$\underline{C}_0\text{-parf-}\,\mathrm{amp}(F) \subset [a, b],$$

when the set of X over S for which there exists a quasi-isomorphism [print 122] $F' \to F$ of $\mathrm{C}(\underline{C}_X)$ with F' strictly $\underline{C}_0$-perfect (2.1), and such that $F'^i = 0$ for $i \notin [a, b]$, is a refinement of S. (By (4.3) this condition is equivalent to the one obtained by replacing the expression "quasi-isomorphism of $\mathrm{C}(\underline{C}_X)$" by "isomorphism of $\mathrm{D}(\underline{C}_X)$".) We shall say that F has finite $\underline{C}_0$-perfect amplitude when there exist integers a and b such that $\underline{C}_0$-parf-amp$(F) \subset [a, b]$. We shall say that F is $\underline{C}_0$-perfect when F locally has finite $\underline{C}_0$-perfect amplitude(*).

Example 4.8. Let $(\underline{S}, A)$ be a ringed topos, $\underline{C}$ the $\underline{S}$-category defined by $U \longmapsto$ (category of left A_U-Modules), and $\underline{C}_0$ the sub-$\underline{S}$-category defined by $U \longmapsto$ (category of left A_U-Modules that are direct factors of free Modules of finite type). By (1.3.1), the pair $(\underline{C}, \underline{C}_0)$ satisfies the hypotheses (4.0). An object F of $\mathrm{D}({}_A\underline{S})$ (cf. (2.15)) will be said to have perfect amplitude contained in $[a, b]$ (resp. finite perfect amplitude, resp. to be perfect) when F has $\underline{C}_0$-perfect amplitude contained in $[a, b]$ (resp.. . .). When the direct factors of free Modules of finite type are locally free (cf. (2.15.1)), saying that F is perfect therefore means that F is locally isomorphic (in $\mathrm{D}(\underline{S})$) to a complex with bounded degrees and components free of finite type.

Notation 4.9. We shall denote by $\mathrm{D}(\underline{C})_{\underline{C}_0\text{-parf}}$ or $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ the strictly full sub-$\underline{S}$-category of $\mathrm{D}(\underline{C})$ formed by the $\underline{C}_0$-perfect objects. In the case of Example (4.8), we shall write $\mathrm{D}({}_A\underline{S})_{\mathrm{parf}}$, or $\mathrm{Parf}({}_A\underline{S})$, or simply $\mathrm{Parf}(\underline{S})$ when there is no possible ambiguity about A.

[print 123]**Proposition 4.10.**[104] The category $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ is a triangulated sub-$\underline{S}$-category of $\mathrm{D}(\underline{C})$ (1.1.4). The canonical functor

$$\mathrm{K}^b(\underline{C}_0) \longrightarrow \mathrm{Parf}(\underline{C}, \underline{C}_0)$$

(resp. $\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0)) \to \mathrm{Tr}(\mathrm{Parf}(\underline{C}, \underline{C}_0))$ (1.1.3)) induces an $\underline{S}$-equivalence on the associated stacks ([G] II 2.2.4), which means:

(i) for every $X \in \mathrm{ob}\,\underline{S}$[105] and all $E, F \in \mathrm{ob}\,\mathrm{K}^b(\underline{C}_{0X})$ (resp. $\mathrm{Tr}(\mathrm{K}^b(\underline{C}_{0X}))$), the canonical arrow

$$\underline{\mathrm{Hom}}_{\mathrm{K}^b(\underline{C}_0)}(E, F) \longrightarrow \underline{\mathrm{Hom}}_{\mathrm{D}(\underline{C})}(E, F)$$

(resp. $\underline{\mathrm{Hom}}_{\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0))}(E, F) \to \underline{\mathrm{Hom}}_{\mathrm{Tr}(\mathrm{D}(\underline{C}))}(E, F)$) is an isomorphism, and

(ii) every object of $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ (resp. $\mathrm{Tr}(\mathrm{Parf}(\underline{C}, \underline{C}_0))$) is locally isomorphic in $\mathrm{D}(\underline{C})$ (resp. $\mathrm{Tr}(\mathrm{D}(\underline{C}))$) to an object of[106] $\mathrm{K}^b(\underline{C}_0)$ (resp. $\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0))$).[107]

Proof. Assertion (i) without the "resp." alternative is a special case of (4.6), and assertion (ii) without the "resp." alternative is true by definition ((4.7) and (4.9)). It is also clear that

(*) Of course, when no confusion can arise, we shall omit $\underline{C}_0$ from the preceding expressions.

$\mathrm{Parf}(\underline{C}, \underline{C}_0)$ is stable under translation of degrees. To prove that $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ is a triangulated sub-$\underline{S}$-category of $\mathrm{D}(\underline{C})$, it therefore suffices to show that, when

$$\begin{array}{ccccc} & & G & & \\ & {\scriptstyle +1}\swarrow & & \nwarrow & \\ E & & \xrightarrow{\quad u \quad} & & F \end{array} \qquad (*)$$

is a distinguished triangle of $\mathrm{D}(\underline{C})$ such that E and F are perfect, G is perfect. After localizing, we can suppose that E and F are objects of $\mathrm{K}^b(\underline{C}_0)$. After localizing further, we can suppose, by assertion (i) without the "resp." alternative, that u is an arrow of $\mathrm{K}^b(\underline{C}_0)$; then G is perfect because it is isomorphic to the cone of u, which is strictly perfect. Thus $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ is indeed a triangulated sub-$\underline{S}$-category of $\mathrm{D}(\underline{C})$. Moreover, the preceding argument also proves the "resp." assertion (ii). It remains to prove the "resp." assertion (i). Let $X \in \mathrm{ob}\,\underline{S}$, and let $E = (E_1 \xrightarrow{a_1} E_2 \xrightarrow{a_2} E_3 \xrightarrow[+1]{a_3} E_1)$, $F = (F_1 \xrightarrow{b_1} F_2 \xrightarrow{b_2} F_3 \xrightarrow[+1]{b_3} F_1)$ be distinguished triangles of $\mathrm{K}^b(\underline{C}_{0X})$. Let us show that the arrow

$$\underline{\mathrm{Hom}}_{\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0))}(E, F) \longrightarrow \underline{\mathrm{Hom}}_{\mathrm{Tr}(\mathrm{D}(\underline{C}))}(E, F) \qquad (**)$$

is an isomorphism. By assertion (i) without the "resp." alternative, we already know that $(**)$ is injective. Let $(u_i) : E \to F$ be a morphism of triangles in $\mathrm{D}(\underline{C})$. After localizing, we can suppose, by assertion (i) without the "resp." alternative, that u_i $(1 \leq i \leq 3)$ is an arrow of $\mathrm{K}^b(\underline{C}_0)$. The arrow $u_{i+1}a_i - b_i u_i$, for $1 \leq i \leq 3$ (with the convention $u_4 = u_1$), is zero in $\mathrm{D}(\underline{C})$, hence locally zero in $\mathrm{K}^b(\underline{C}_0)$ by assertion (i) without the "resp." alternative. In other words, (u_i) locally defines a morphism of triangles in $\mathrm{K}^b(\underline{C}_0)$. This proves that $(**)$ is surjective and completes the proof of (4.10).

Remark 4.10.1. In (4.10), we can replace $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ by $\mathrm{D}^*(\underline{C})_{\underline{C}_0\text{-parf}} = \mathrm{D}^*(\underline{C}) \cap \mathrm{Parf}(\underline{C}, \underline{C}_0)$ $(* = -$ or $b)$; nothing in the proof needs to be changed.

Complement 4.11. We can describe more precisely the triangulated structure of $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ in terms of perfect amplitude (4.7). When $E \in \mathrm{ob}\,\mathrm{D}(\underline{C})$ has perfect amplitude contained in $[a, b]$, $E[n]$ has perfect amplitude contained in $[a - n, b - n]$.

[print 125]Let

$$\begin{array}{ccccc} & & G & & \\ & {\scriptstyle +1}\swarrow & & \nwarrow & \\ E & & \xrightarrow{\qquad} & & F \end{array}$$

be a distinguished triangle of $\mathrm{D}(\underline{C})$. We can draw up a table analogous to (2.6), in which the entries indicate perfect amplitudes and each row corresponds to an implication whose conclusion is underlined.

E	F	G
$[a+1, b+1]$	$[a, b]$	$\underline{[a, b]}$
$[a, b]$	$\underline{[a, b]}$	$\underline{[a, b]}$
$\underline{[a, b]}$	$[a, b]$	$[a-1, b-1]$.

From (4.10) we immediately deduce the

Corollary 4.12. Suppose that $\underline{S}$ is a punctual site. For $F \in \mathrm{ob}\,\mathrm{D}(\underline{C})$ to have perfect amplitude (rel. to $\underline{C}_0$) contained in $[a, b]$, it is necessary and sufficient that there exist a quasi-isomorphism $F' \to F$, with F' strictly perfect and such that $F'^i = 0$ for $i \notin [a, b]$. The canonical functors

$$\mathrm{K}^b(\underline{C}_0) \longrightarrow \mathrm{Parf}(\underline{C}, \underline{C}_0)$$

and

$$\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0)) \longrightarrow \mathrm{Tr}(\mathrm{Parf}(\underline{C}, \underline{C}_0))$$

are equivalences of categories.

Remark 4.12.1. When $\underline{S}$ is a punctual site, the objects of $\underline{C}_0$ are projective. When $\underline{C}_0$ contains all projective objects of $\underline{C}$, one says "projective amplitude" instead of "$\underline{C}_0$-perfect amplitude".

[print 126]**Remark 4.12.2.** Let $\underline{C}$ be an abelian category (which can be regarded as fibered over a punctual site), and let $\underline{C}_0$ be a strictly full additive subcategory. Suppose that $\underline{C}_0$ is stable under kernels of epimorphisms and quasi-liftable in $\underline{C}$. Then an acyclic object of $\mathrm{K}^b(\underline{C}_0)$ is in general not zero, so there is no hope that the functor $\mathrm{K}^b(\underline{C}_0) \longrightarrow \mathrm{D}(\underline{C})$[108] is faithful. However, when $\mathrm{K}^{b,\varnothing}(\underline{C}_0)$ denotes the thick subcategory of $\mathrm{K}^b(\underline{C}_0)$ formed by the acyclic objects (in $\underline{C}$), the canonical functor

$$\mathrm{K}^b(\underline{C}_0)/\mathrm{K}^{b,\varnothing}(\underline{C}_0) \longrightarrow \mathrm{D}(\underline{C})$$

is fully faithful.

Hints: We already know (2.7) that the functor

$$\mathrm{K}^-(\underline{C}_0)/\mathrm{K}^{-,\varnothing}(\underline{C}_0) \longrightarrow \mathrm{D}^-(\underline{C})$$

is fully faithful. Show, for example using the criterion ([V] 4.2.b)(ii)'), that the functor

$$\mathrm{K}^b(\underline{C}_0)/\mathrm{K}^{b,\varnothing}(\underline{C}_0) \longrightarrow \mathrm{K}^-(\underline{C}_0)/\mathrm{K}^{-,\varnothing}(\underline{C}_0)^{[109]}$$

is fully faithful.

Lemma 4.13. Let $F \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$, $a, b, c \in \mathbb{Z}$ be such that $a \leq b$ and $a \leq c$. If F has perfect amplitude contained in $[a, c]$ (4.7) and $\mathrm{H}^i(F) = 0$ for $i > b$, then F has perfect amplitude contained in $[a, d]$, where $d = \inf(b, c)$.

Proof. We can suppose that $b \leq c$, and that F is strictly perfect with $F^i = 0$ for $i \notin [a, c]$. Then F is isomorphic to its truncation F' defined by [print 127]$F'^i = F^i$ for $i < b$, $F'^b = Z^b(F)$, $F'^i = 0$ for $i > b$. But $Z^b(F)$ admits a finite right resolution by objects of $\underline{C}_0$, so it is locally an object of $\underline{C}_0$, whence the assertion.

Corollary 4.14. Let $F \in \mathrm{ob}\,\underline{C}_S$ and $n \in \mathbb{N}$. Equivalent conditions:

(i) $\mathrm{parf.\,amp}(F) \subset [-n, 0]$;

(ii) F locally admits a left resolution of length n by objects of $\underline{C}_0$, i.e. the set of X over S for which there exists an exact sequence

$$0 \longrightarrow L^{-n} \longrightarrow L^{-n+1} \longrightarrow \ldots \longrightarrow L^0 \longrightarrow F_X \longrightarrow 0,$$

with $L^i \in \mathrm{ob}\,\underline{C}_{0X}$ for every i, is a refinement of S.

Proof. This is an immediate consequence of (4.7) and (4.13).

Definition 4.14.1. We shall say that F has perfect dimension $\leq n$, and we shall write $\mathrm{parf.\,dim}(F) \leq n$, when F satisfies the equivalent conditions of (4.14). We shall call the smallest integer n such that $\mathrm{parf.\,dim}(F) \leq n$ the perfect dimension of F (when there is no such integer, we shall say that F has infinite perfect dimension).

When $\underline{S}$ is the punctual site and $\underline{C}_0$ is the full subcategory of $\underline{C}$ formed by the projective objects, perfect dimension is the usual projective dimension.

Proposition 4.15. Let $a, b, n \in \mathbb{Z}$ ($a \leq b$, $n \geq 0$), and let $F \in \mathrm{ob}\,\mathrm{D}(\underline{C})$ be such that $F^i = 0$ (resp. $\mathrm{H}^i(F) = 0$) for $i \notin [a, b]$. If, for every $i \in [a, b]$, F^i (resp. $\mathrm{H}^i(F)$) has perfect dimension

$\leq n + (i - a)$, then F has perfect amplitude contained in $[a - n, b]$. Consequently, if for every $i \in [a, b]$, F^i (resp. $\mathrm{H}^i(F)$) is perfect, F is too.

[print 128]Proof. Proceed by induction on b (with a and n fixed), truncating F and applying (4.11) (cf. the proof of (2.11)).

Lemma 4.16. Let $a, b \in \mathbb{Z}$ ($a \leq b$), and $F \in \mathrm{ob}\,\mathrm{D}(\underline{C})$ be such that $F^i = 0$ for $i \notin [a, b]$. Assume that $\mathrm{parf.amp}(F) \subset [a, b]$, and that $F^i \in \mathrm{ob}\,\underline{C}_0$ for $i > a$. Then F^a is locally an object of $\underline{C}_0$.

Proof. We have an exact sequence (obtained by truncating F)

$$0 \longrightarrow F' \longrightarrow F \longrightarrow F^a[-a] \longrightarrow 0,$$

with $\mathrm{parf.amp}(F') \subset [a + 1, b + 1]$. From (4.11) and (4.13) we deduce that $\mathrm{parf.dim}(F^a) = 0$, i.e. that F^a is locally an object of $\underline{C}_0$.

Proposition 4.17. Let $F', F'' \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$, and $a, b \in \mathbb{Z}$, with $a \leq b$. Consider the conditions:

(i) $\mathrm{parf.amp}(F') \subset [a, b]$ and $\mathrm{parf.amp}(F'') \subset [a, b]$;

(ii) $\mathrm{parf.amp}(F' + F'') \subset [a, b]$.

We have (i) $\Rightarrow$ (ii), and, when $\underline{C}_0$ is locally stable under direct factors (i.e. the relation $E' + E'' \in \mathrm{ob}\,\underline{C}_0$, for $E', E'' \in \mathrm{ob}\,\underline{C}_X$ and $X \in \mathrm{ob}\,\underline{S}$, implies that locally E' and E'' are objects of $\underline{C}_0$), we have (i) $\Leftrightarrow$ (ii).

Proof. The implication (i) $\Rightarrow$ (ii) is a special case of (4.11). Let us prove the converse. Put $F = F' + F''$. Since F is pseudo-coherent, F' and F'' are too (2.12). On the other hand, the relation $\mathrm{H}^i(F) = 0$ for $i \notin [a, b]$ implies $\mathrm{H}^i(F') = \mathrm{H}^i(F'') = 0$ for $i \notin [a, b]$. By (2.10 a)), we can therefore suppose (after localizing) that F'^i (resp. F''^i) $\in \mathrm{ob}\,\underline{C}_0$ for $i \geq a+1$ and $F'^i = F''^i = 0$ for $i > b$, then, by truncating, that [print 129]$F'^i = F''^i = 0$ for $i < a$. Since $\mathrm{parf.amp}(F) \subset [a, b]$[110], we deduce from (4.16) that F^a is locally an object of $\underline{C}_0$. But $F^a = F'^a + F''^a$, so F'^a and F''^a are locally objects of $\underline{C}_0$, since $\underline{C}_0$ is locally stable under direct factors.

Proposition 4.18. Let $(\underline{S}', \underline{C}', \underline{C}'_0)$ be a triple satisfying the same hypotheses (4.0) as $(\underline{S}, \underline{C}, \underline{C}_0)$. Let $f : \underline{S}' \to \underline{S}$ be a morphism of sites (defined by $f^{-1} : \underline{S} \to \underline{S}'$), and let $u : \mathrm{D}^*(\underline{C}) \longrightarrow \mathrm{D}^*(\underline{C}')$ $(* = -, +, \varnothing)$ be a functor over f (cf. (2.13)). If u takes objects of $\underline{C}_0$ to $\underline{C}'_0$-perfect complexes, then u defines a functor $\mathrm{D}^*(\underline{C})_{\underline{C}_0\text{-parf}} \longrightarrow \mathrm{D}^*(\underline{C}')_{\underline{C}'_0\text{-parf}}$.

Proof. Let $F \in \mathrm{ob}\,\mathrm{D}^*(\underline{C})_{\underline{C}_0\text{-parf}}$; let us show that $u(F) \in \mathrm{ob}\,\mathrm{D}^*(\underline{C}')_{\underline{C}'_0\text{-parf}}$. Since the question is local, we can suppose that F is strictly perfect. We need only use dévissage, taking into account that $\mathrm{D}^*(\underline{C}')_{\underline{C}'_0\text{-parf}}$ is a triangulated subcategory of $\mathrm{D}^*(\underline{C}')$ (4.10).[111]

Remark 4.18.1. Suppose that there exist $m, n \in \mathbb{Z}$ ($m \leq n$) such that $F \in \mathrm{ob}\,\underline{C}_0$ implies $\underline{C}'_0\text{-}\mathrm{parf.amp}(u(F)) \subset [m, n]$. Then, for $F \in \mathrm{ob}\,\mathrm{D}^*(\underline{C})$, we have

$$\underline{C}_0\text{-}\mathrm{parf.amp}(F) \subset [a, b] \quad \Rightarrow \quad \underline{C}'_0\text{-}\mathrm{parf.amp}(u(F)) \subset [a + m, b + n].$$

The proof is left as an exercise to the reader (use (4.11)).

Corollary 4.19. Under the hypotheses of (2.16), if $E' \in \mathrm{ob}\,\mathrm{D}^-({}_{A'}\underline{S}'_A)$ is perfect as an object of $\mathrm{D}^-({}_{A'}\underline{S}')$ and $E \in \mathrm{ob}\,\mathrm{D}^-({}_A\underline{S})$ is perfect, then $E' \overset{\mathrm{L}}{\otimes}_A E$ is perfect.

Proof. We can suppose that $E' \in \mathrm{ob}\,\mathrm{D}^-({}_{A'}S'_A)$ (S'= final object of $\underline{S}'$), and apply (4.18) to $u = E' \overset{\mathrm{L}}{\otimes}_A : \mathrm{D}^-({}_A\underline{S}) \longrightarrow \mathrm{D}^-({}_{A'}\underline{S}')$. In particular:

[print 130]**Corollary 4.19.1.** a) The functor $\mathbb{L}f^* : \mathrm{D}^-({}_A\underline{S}) \longrightarrow \mathrm{D}^-({}_{A'}\underline{S}')$ induces a functor

$$\mathbb{L}f^* : \mathrm{D}^-({}_A\underline{S})_{\mathrm{parf}} \longrightarrow \mathrm{D}^-({}_{A'}\underline{S}')_{\mathrm{parf}}.$$

b) If A is commutative, the functor $\overset{\mathrm{L}}{\otimes}_A : \mathrm{D}^-(\underline{S}) \times_{\underline{S}} \mathrm{D}^-(\underline{S}) \longrightarrow \mathrm{D}^-(\underline{S})$ induces a functor

$$\overset{\mathrm{L}}{\otimes}_A : \mathrm{D}^-(\underline{S})_{\mathrm{parf}} \times_{\underline{S}} \mathrm{D}^-(\underline{S})_{\mathrm{parf}} \longrightarrow \mathrm{D}^-(\underline{S})_{\mathrm{parf}}.$$

Remark 4.19.2. We can make (4.19) more precise in terms of perfect amplitude:

$$\mathrm{parf.amp}(E) \subset [a,b] \quad \text{and} \quad \mathrm{parf.amp}(E') \subset [a',b'] \quad \Rightarrow \quad \mathrm{parf.amp}(E' \overset{\mathrm{L}}{\otimes}_A E) \subset [a+a', b+b'].$$

In particular:

$$\mathrm{parf.amp}(E) \subset [a,b] \quad \Rightarrow \quad \mathrm{parf.amp}(\mathbb{L}f^*(E)) \subset [a,b].$$

5. Finite Tor-dimension and perfection

5.0. The reader will surely have asked: what does a pseudo-coherent complex E lack in order to be perfect? Is it enough, for example, that E have bounded cohomology? The answer is no. Indeed, take for $\underline{C}$ the category of modules over a commutative noetherian ring A, and for $\underline{C}_0$ the full subcategory of projective A-modules of finite type. Any A-module of finite type E is pseudo-coherent and has bounded cohomology, but E is perfect only when E also has finite projective dimension (or, equivalently, finite tor-dimension). This example suggests that the difference between a pseudo-coherent complex and a perfect complex lies in a question of cohomological dimension. This is what we shall now examine [print 131]in the case of the category of Modules of a ringed topos (cf. (3.3)). We shall need some preliminary sorites on the notion of finite tor-dimension. We shall be brief, since this question already appears in the literature: cf. ([H] II 4) and (SGA 4 XVII 4.1.9.).

Proposition 5.1. Let $(\underline{S}, A)$ be a ringed topos, $X \in \mathrm{ob}\,\underline{S}$, $F \in \mathrm{ob}\,\mathrm{D}^-({}_AX)$, $m, n \in \mathbb{Z}$ be such that $m \leq n$. Equivalent conditions:


(i) F is isomorphic, in $\mathrm{D}^-({}_AX)$, to a complex F' such that F'^i is flat for every i and zero for $i \notin [m,n]$;

(ii) for every right A_X-Module E, we have $\underline{\mathrm{H}}^i(E \overset{\mathrm{L}}{\otimes}_A F) = 0$ for $i \notin [m,n]$.

Proof. The implication (i) $\Rightarrow$ (ii) is trivial; the implication (ii) $\Rightarrow$ (i) follows, by truncation, from the

Lemma 5.1.1. (cf. (4.16)). Let $F \in \mathrm{ob}\,\mathrm{D}^-({}_AX)$. Assume that F satisfies hypothesis (ii) of (5.1), that $F^i = 0$ for $i \notin [m,n]$, and that F^i is flat for $i \neq m$. Then F^m is flat.

Proof. We have a distinguished triangle



,

where F' has flat components that vanish outside the interval $[m+1, n]$ (suppose $m < n$). Apply the functor $E \overset{\mathrm{L}}{\otimes}_A .$ for E an arbitrary right A_X-Module, and write the exact cohomology sequence.

[print 132]**Definition 5.2.** An object F of $\mathrm{D}^-({}_A\underline{S})$ satisfying the equivalent conditions of (5.1) will be said to have flat amplitude contained in the interval $[m,n]$, and we shall write

$$\mathrm{tor.amp}(F) \subset [m,n].$$

We shall say that F has finite flat amplitude (or finite tor-dimension) when there exist integers m and n such that the preceding condition holds.

5.3. Let $X \in \mathrm{ob}\,\underline{S}$. We shall denote by $\mathrm{D}^-({}_AX)_{\mathrm{torf}}$ the full subcategory of $\mathrm{D}^-({}_AX)$ formed by the objects of finite tor-dimension. By criterion (ii) of (5.1), it is a triangulated subcategory of $\mathrm{D}^-({}_AX)$. As X varies, the categories $\mathrm{D}^-({}_AX)_{\mathrm{torf}}$ form a triangulated sub-$\underline{S}$-category (1.1.4) of $\mathrm{D}^-({}_A\underline{S})$, which we shall denote by $\mathrm{D}^-({}_A\underline{S})_{\mathrm{torf}}$. The functor $\overset{\mathrm{L}}{\otimes}_A$ induces an exact $\underline{S}$-functor

$$\overset{\mathrm{L}}{\otimes}_A : \mathrm{D}^b(\underline{S}_A) \times_{\underline{S}} \mathrm{D}({}_A\underline{S})_{\mathrm{torf}} \longrightarrow \mathrm{D}^b(\underline{S}_{\mathbb{Z}}).$$

5.4. For a left A-Module F to have flat amplitude contained in $[-n, 0]$ ($n \in \mathbb{N}$), it is necessary and sufficient that F have tor-dimension $\leq n$ in the usual sense, i.e. that F admit a left resolution of length n by flat A-Modules.

[print 133]5.5. Let $X \in \mathrm{ob}\,\underline{S}$, and $m, n \in \mathbb{Z}$ ($m \leq n$). For $F \in \mathrm{ob}\,\mathrm{D}^-({}_AX)$ to have flat amplitude contained in $[m, n]$, it is necessary and sufficient that there exist a covering family $(X_i \to X)$ such that $F|X_i$ has flat amplitude contained in $[m, n]$. In particular, when $\underline{S}_{/X}$ is of finite type (SGA 4 VI 1.7), and F is locally of finite tor-dimension, F has finite tor-dimension.

Proposition 5.6. Under the hypotheses of (2.16), if E' has flat amplitude contained in $[m', n']$ as an object of $\mathrm{D}^-({}_{A'}\underline{S}')$ and E has flat amplitude contained in $[m, n]$, then $E' \overset{\mathrm{L}}{\otimes}_A E$ has flat amplitude contained in $[m + m', n + n']$.

Proof. This is immediate: use form (5.1 (i)) of the notion of finite tor-dimension and the fact that, when E is a flat left A-Module, $f^{-1}(E)$ is a flat left $f^{-1}(A)$-Module (SGA 4 IV App. II).

In particular:

Corollary 5.6.1. a) The functor $\mathbb{L}f^* : \mathrm{D}^-({}_A\underline{S}) \longrightarrow \mathrm{D}^-({}_{A'}\underline{S}')$ induces a functor

$$\mathbb{L}f^* : \mathrm{D}({}_A\underline{S})_{\mathrm{torf}} \longrightarrow \mathrm{D}({}_{A'}\underline{S}')_{\mathrm{torf}}.$$

b) If A is commutative, the functor $\overset{\mathrm{L}}{\otimes}_A : \mathrm{D}^-(\underline{S}) \times_{\underline{S}} \mathrm{D}^-(\underline{S}) \longrightarrow \mathrm{D}^-(\underline{S})$ induces a functor

$$\overset{\mathrm{L}}{\otimes}_A : \mathrm{D}(\underline{S})_{\mathrm{torf}} \times_{\underline{S}} \mathrm{D}(\underline{S})_{\mathrm{torf}} \longrightarrow \mathrm{D}(\underline{S})_{\mathrm{torf}}.$$

Remark 5.6.2. Let $E \in \mathrm{ob}\,\mathrm{D}^-({}_AS)$ (S= final object of $\underline{S}$), and $m, n \in \mathbb{Z}$ ($m \leq n$). When E has flat amplitude contained in $[m, n]$, so does E_s, by (5.6), for every point s of $\underline{S}$. The converse is true when $\underline{S}$ has enough points (use form (5.1.(ii)) of the notion of flat [print 134]amplitude, and the fact that $\overset{\mathrm{L}}{\otimes}_A$ commutes with passage to the fibres).

Exercises 5.7. a) Suppose that A is commutative and $(\underline{S}', A')$ faithfully flat over $(\underline{S}, A)$ (cf. (2.16.2)). For $E \in \mathrm{ob}\,\mathrm{D}^-({}_AS)$([1]) to have flat amplitude contained in $[m, n]$, it is necessary and sufficient that $f^*(E)$ have that property.

b) Let $F \in \mathrm{ob}\,\mathrm{D}^+({}_AS)$([1]). Show the equivalence of the conditions:

(i) F is isomorphic, in $\mathrm{D}^+({}_AS)$, to a complex F' such that F'^i is injective for every i and zero for $i \notin [m, n]$;

(ii) for every left A_S-Module E, we have $\mathrm{Ext}^i(E, F) = 0$ for $i \notin [m, n]$.

([1]) S = final object of $\underline{S}$

When F satisfies the preceding conditions, we say that F has injective amplitude contained in $[m,n]$. The full subcategory of $\mathrm{D}^+({}_AS)$ formed by the objects of finite injective amplitude is a triangulated subcategory, denoted by $\mathrm{D}^+({}_AS)_{\mathrm{injf}}$. Show that, when A is commutative, the functor $\mathbb{R}\underline{\mathrm{Hom}} : \mathrm{D}(S)^\circ \times \mathrm{D}^+(S) \longrightarrow \mathrm{D}(S)$ induces a functor

$$\mathbb{R}\underline{\mathrm{Hom}} : \mathrm{D}(S)^\circ_{\mathrm{torf}} \times \mathrm{D}(S)_{\mathrm{injf}} \longrightarrow \mathrm{D}(S)_{\mathrm{injf}}.$$

We can now answer the question of (5.0) and state the main result of this section.

Theorem 5.8. Let $(\underline{S}, A)$ be a ringed topos, $E \in \mathrm{ob}\,\mathrm{D}({}_A\underline{S})$, $m, n \in \mathbb{Z}$, with $m \leq n$. Equivalent conditions:

(i) $\mathrm{parf.amp}(E) \subset [m,n]$ (4.8);

(ii) E is $(m-1)$-pseudo-coherent (2.15) and we have $\mathrm{tor.amp}(E) \subset [m,n]$ (5.2).

Since "perfect" obviously implies "pseudo-coherent", we deduce:

[print 135]**Corollary 5.8.1.** For E to be perfect (4.8), it is necessary and sufficient that E be pseudo-coherent (2.15)[112] and locally of finite tor-dimension (5.2).

For the proof, we shall need some lemmas.

Lemma 5.8.2. Let B be a ring of $\underline{S}$. If E is a left A-Module, G a right B-Module, and F an A-B-Bimodule (on the left over A, on the right over B), there is a functorial canonical homomorphism

$$\underline{\mathrm{Hom}}_B(F,G) \otimes_A E \longrightarrow \underline{\mathrm{Hom}}_B(\underline{\mathrm{Hom}}_A(E,F),G). \tag{5.8.2.1}$$

This is an isomorphism for G injective and E of finite presentation.

Proof. Let us first define the arrow. By the adjunction formula dear to Cartan, it amounts to defining a homomorphism of B-Modules:

$$\underline{\mathrm{Hom}}_B(F,G) \otimes_A E \otimes_{\mathbb{Z}_S} \underline{\mathrm{Hom}}_A(E,F) \longrightarrow G(^1). \tag{*}$$

Now we have a canonical homomorphism of A-B-biModules

$$E \otimes_{\mathbb{Z}_S} \underline{\mathrm{Hom}}_A(E,F) \longrightarrow F \tag{**}$$

(coming by adjunction from the identity arrow of $\underline{\mathrm{Hom}}_A(E,F)$). Likewise, we have a canonical homomorphism of B-Modules

$$\underline{\mathrm{Hom}}_B(F,G) \otimes_A F \longrightarrow G. \tag{***}$$

We then define (*) as the composite of (**) and (***). To see that the arrow (5.8.2.1) is an isomorphism when G is injective and E of finite presentation, it suffices to observe that, when G is injective, the source and target of the arrow are right-exact functors in E, and that the arrow induces the identity for $E = A$.

[print 136]Here is now a special case of (5.8).

Lemma 5.8.3. Let E be a left A-Module. Equivalent conditions:

(i) E is locally a direct factor of a free Module of finite type;

(ii) E is flat and of finite presentation.

(1) we write S for the final object of $\underline{S}$ and $\mathbb{Z}_S$ for the constant sheaf with value $\mathbb{Z}$.

Proof (after BOURBAKI Alg. Comm. chap. I, § 2 Exer 15). It is clear that (i) implies (ii). Let us prove (ii) ⇒ (i). By (1.3.1), we must show that the functor $\underline{\mathrm{Hom}}_A(E,.)$ is exact. Let $F \longrightarrow F'' \longrightarrow 0$ be an exact sequence of A-Modules (on the left). We must show that the sequence

$$\underline{\mathrm{Hom}}_A(E,F) \longrightarrow \underline{\mathrm{Hom}}_A(E,F'') \longrightarrow 0$$

is exact. For this it suffices to prove that, for every injective $\mathbb{Z}_S$-Module I, the sequence obtained by applying the functor $\underline{\mathrm{Hom}}_{\mathbb{Z}}(.,I)$ is exact. Now we have a commutative diagram, in which the vertical arrows are the canonical arrows of (5.8.2.1)[113]:


$$\begin{array}{ccccc} 0 & \to & \underline{\mathrm{Hom}}_{\mathbb{Z}_S}(F'',I)\otimes_A E & \to & \underline{\mathrm{Hom}}_{\mathbb{Z}_S}(F,I)\otimes_A E \\ & & \downarrow & & \downarrow \\ 0 & \to & \underline{\mathrm{Hom}}_{\mathbb{Z}_S}(\underline{\mathrm{Hom}}_A(E,F''),I) & \to & \underline{\mathrm{Hom}}_{\mathbb{Z}_S}(\underline{\mathrm{Hom}}_A(E,F),I). \end{array}$$

The vertical arrows are isomorphisms by (5.8.2). The top row is exact because E is flat. Thus the bottom row is exact, as required.

Proof of (5.8). The implication (i) ⇒ (ii) follows trivially from the definitions. Let us prove (ii) ⇒ (i). Since the question is local, we can suppose that E is strictly $(m-1)$-pseudo-coherent. Since, on the other hand, E is acyclic in degree $< m$, the [print 137]truncation arrow

$$\begin{array}{ccccccc} \to & E^{m-1} & \to & E^m & \to & E^{m+1} & \to \dots \\ & \downarrow & & \downarrow & & \downarrow & \\ \to & 0 & \to & E^m/B^m & \to & E^{m+1} & \to \dots \end{array}$$

is a quasi-isomorphism. Denote the truncated complex by E'. By construction, E' has flat amplitude $\subset [m,n]$, and E'^i is free of finite type for $i > m$. Thus, by (5.1.1), E'^m is flat. On the other hand, we have a distinguished triangle



,

where E'' is strictly perfect. By (2.6), $E'^m[-m]$ is $(m-1)$-pseudo-coherent, i.e. (2.9) E'^m is of finite presentation. By (5.8.3), we conclude that E'^m is locally a direct factor of a free Module of finite type, hence that $\mathrm{parf.amp}(E) \subset [m,n]$.

Remarks 5.8.4. a) Using (2.5), we recover the fact that perfect complexes form a triangulated subcategory of $\mathrm{D}({}_A\underline{S})$.

b) A perfect complex, even with bounded cohomology, does not necessarily have finite tor-dimension. It does, however, when $\underline{S}$ is of finite type (5.5). Denote by $\mathrm{D}({}_A\underline{S})_{\underline{\mathrm{parf}}}$ the full sub-$\underline{S}$-category of $\mathrm{D}^b({}_A\underline{S})$ formed by the perfect complexes of finite tor-dimension. It follows at once from (2.16.1 b)) and (5.3) that, when A is commutative, the functor $\overset{\mathrm{L}}{\otimes}_A$ induces a functor

$$\overset{\mathrm{L}}{\otimes}_A : \mathrm{D}(\underline{S})_{\underline{\mathrm{parf}}} \times_{\underline{S}} \mathrm{D}^b(\underline{S})_{\mathrm{coh}} \longrightarrow \mathrm{D}^b(\underline{S})_{\mathrm{coh}}.$$

[print 138]This formula is the starting point of the theory of intersections on schemes. Observe, on the other hand, that the (derived) tensor product of two pseudo-coherent complexes with bounded cohomology does not in general have bounded cohomology.

Corollary 5.9. Suppose that $\underline{S}$ = (Ens) (so A "is" an ordinary ring) and that every left A-module has finite tor-dimension (this holds, for example, when A has finite global cohomological dimension, see (MULT VI 2.6) and (EGA 0_{IV} 17.2.8)). Then we have

(i) $\mathrm{D}^b({}_A\underline{S}) = \mathrm{D}({}_A\underline{S})_{\mathrm{torf}}$, and

(ii) $\mathrm{D}^b({}_A\underline{S})_{\mathrm{coh}} = \mathrm{D}({}_A\underline{S})_{\underline{\mathrm{parf}}}$.

In other words, for $E \in \mathrm{ob}\,\mathrm{D}({}_A\underline{S})$ to have finite tor-dimension (resp. finite tor-dimension and be perfect), it is necessary and sufficient that E have bounded cohomology (resp. bounded cohomology and be pseudo-coherent).

Proof. By (5.8), it suffices to prove (i). Let $E \in \mathrm{ob}\,\mathrm{D}^b({}_A\underline{S})$. We can suppose (extending E by 0) that $E \in \mathrm{ob}\,\mathrm{D}^b({}_AS)$, where S is the final object of $\underline{S}$. Proceeding by induction on the number of nonzero cohomology objects of E, we reduce by dévissage to the case where E is an ordinary A-Module, and we are done.

More generally, we can state:

Corollary 5.10 Let $X \in \mathrm{ob}\,\underline{S}$. Suppose that $\underline{S}_{/X}$ has enough points (SGA 4 IV 6.4.1.) and that, for every point s of $\underline{S}_{/X}$, every left A_s-module has finite tor-dimension. Then we have

$$\mathrm{D}^b({}_AX)_{\mathrm{coh}} = \mathrm{D}^b({}_AX)_{\mathrm{parf}}.$$

[print 139]In other words, for $E \in \mathrm{ob}\,\mathrm{D}^b({}_AX)$ to be pseudo-coherent, it is necessary and sufficient that E be perfect.

Proof. Let $E \in \mathrm{ob}\,\mathrm{D}^b({}_AX)_{\mathrm{coh}}$ and s be a point of $\underline{S}_{/X}$. We must show that E is perfect "in a neighbourhood of s", i.e. that there exists an s-pointed object U of $\underline{S}$ over X such that $E|U$ is perfect. Since the functor s^* is exact, by (2.16.1 a)) we have $E_s \in \mathrm{ob}\,\mathrm{D}^b(A_s)_{\mathrm{coh}}$. Thus, by (5.9), E_s is perfect. The assertion will therefore follow from (3.7.2) and the following lemma:

Lemma 5.10.1. Let s be a point of $\underline{S}$ and let $F \in \mathrm{ob}\,\mathrm{D}(A_s)_{\mathrm{parf}}$. There exist an s-pointed object U of $\underline{S}$ and $F' \in \mathrm{ob}\,\mathrm{D}({}_AU)_{\mathrm{parf}}$ such that $F'_s = F$.

Proof. This is evident when F is a free A_s-module of finite type. Now suppose that F is projective of finite type, hence the image of a projector $p : L \longrightarrow L$, where L is free of finite type. There then exist a "neighbourhood" U of s, a free A_U-Module L', and a homomorphism $p' : L' \to L'$ such that $p'_s = p$. Since ${p'_s}^2 = p'_s$, we can suppose, after shrinking U, that $p'^2 = p'$. Thus $F' = \mathrm{Im}(p')$ is a direct factor of L' and satisfies $F'_s = F$. This generalizes immediately to the case where F is strictly perfect, i.e. a complex with bounded degrees made up of projective modules of finite type.

Examples 5.11. Corollary (5.10) applies in particular when $(\underline{S}, A)$ is defined by a ringed space with regular local rings, for example a regular scheme, or a complex analytic space smooth over $\mathbb{C}$.

[print 140]

6. Rank of a perfect complex

6.0. Let $(X, \underline{O}_X)$ be a ringed space with local rings. It is well known that to every locally free sheaf of finite type L on an open set U of X we can attach a function $\mathrm{rg}_U(L)$, locally constant on U and with integer values, called the rank of L, the family of functions $\mathrm{rg}_U(L)$ being characterized by the following properties:

(i) for an inclusion $V \subset U$ of open sets, we have $\mathrm{rg}_V(L|V) = \mathrm{rg}_U(L)|V$;

(ii) for every exact sequence of locally free $\underline{O}_U$-Modules of finite type


$$0 \longrightarrow L' \longrightarrow L \longrightarrow L'' \longrightarrow 0,$$

we have $\mathrm{rg}_U(L) = \mathrm{rg}_U(L') + \mathrm{rg}_U(L'')$;

(iii) $\mathrm{rg}_X(\underline{O}_X) = 1$.

In this section, we propose to extend the notion of rank to perfect complexes.

Definition 6.1 (cf. SGA 5 VIII 2). Let C be an additive (resp. triangulated) category and F an abelian group. A function $f : \mathrm{ob}(C) \longrightarrow F$ is called an additive function from C to F when for all $L, L', L'' \in \mathrm{ob}(C)$ such that $L \simeq L' + L''$ (resp. there is a distinguished triangle



) we have $f(L) = f(L') + f(L'')$.

The additive functions from C to F form an abelian group, denoted by $\mathrm{Add}(C, F)$.

[print 141]6.2. Let C be a triangulated category and f an additive function from C to an abelian group F. Then f induces an additive function on the additive category C^{ad} underlying C, hence an inclusion $\mathrm{Add}(C, F) \subset \mathrm{Add}(C^{\mathrm{ad}}, F)$. In particular, we have $f(0) = 0$ and $f(L) = f(M)$ when $L \simeq M$. On the other hand, we have $f(L[n]) = (-1)^n f(L)$ for every $L \in \mathrm{ob}(C)$ and $n \in \mathbb{Z}$.

6.3. Let C be an additive (resp. triangulated) category. The group $\mathrm{Add}(C, F)$ depends functorially on F. The functor $\mathrm{Add}(C, .)$ is representable (cf. SGA 5 VIII 2) by an abelian group denoted by $k(C)$(*) endowed with an additive map $\mathrm{cl}_C : \mathrm{ob}(C) \longrightarrow k(C)$ (when C is additive, we take for $k(C)$ the symmetrization of the monoid of isomorphism classes of objects of C; when C is triangulated, we take for $k(C)$ the quotient of the free abelian group generated by $\mathrm{ob}(C)$ by the relations identifying L with $L' + L''$ whenever there is a distinguished triangle $L' \longrightarrow L \longrightarrow L'' \longrightarrow L'[1]$).

The group $k(C)$ depends functorially on C (with respect to additive (resp. exact) functors). When $u, u' : C \to D$ are isomorphic additive (resp. exact) functors, we have $k(u) = k(u')$.

When C is a triangulated category, we have a canonical epimorphism $k(C^{\mathrm{ad}}) \longrightarrow k(C)$ (cf. (6.2)).

Lemma 6.4. Let C be an additive category. The canonical functor $C \longrightarrow \mathrm{K}^b(C)$ induces an isomorphism (functorial in C) $k(C) \longrightarrow k(\mathrm{K}^b(C))$.

Proof. This is an immediate consequence of the following formula (which follows trivially from (6.2)):

$$f(L) = \sum (-1)^i f(L^i) \tag{6.4.1}$$

[print 142]for $L \in \mathrm{ob}\,\mathrm{K}^b(C)$ and f an additive function on $\mathrm{K}^b(C)$.

Definition 6.5. Let $\underline{S}$ be a category, $\underline{C}$ an additive (resp. triangulated) $\underline{S}$-category (1.1.1), and F an abelian presheaf on $\underline{S}$. A family of functions $f_X : \mathrm{ob}(\underline{C}_X) \longrightarrow F(X)$ indexed by $X \in \mathrm{ob}\,\underline{S}$ is called an additive $\underline{S}$-function from $\underline{C}$ to F when the following conditions are satisfied:

(i) for every $X \in \mathrm{ob}\,\underline{S}$[114], f_X is an additive function (6.1);

(ii) for every arrow $u : X \to Y$ of $\underline{S}$, the following diagram is commutative [115]

$$\begin{array}{ccc} \mathrm{ob}\,\underline{C}_Y & \xrightarrow{f_Y} & F(Y) \\ {\scriptstyle u^*}\downarrow & & \downarrow{\scriptstyle F(u)} \\ \mathrm{ob}\,\underline{C}_X & \xrightarrow{f_X} & F(X). \end{array}$$

The additive $\underline{S}$-functions from $\underline{C}$ to F form an abelian group, denoted by $\mathrm{Add}_{\underline{S}}(\underline{C}, F)$.

(*) we shall not use here the usual notation $\mathrm{K}(C)$, since $\mathrm{K}(C)$ already denotes, when C is additive, the category of complexes of C "up to homotopy".

Remarks 6.6. a) Denote by $\underline{C}^{\mathrm{is}}$ the fibered $\underline{S}$-category whose fiber over $X \in \mathrm{ob}\,\underline{S}$ is the category with the same objects as $\underline{C}_X$, but whose arrows are the isomorphisms of $\underline{C}_X$. On the other hand, denote by $\underline{F}$ the fibered $\underline{S}$-category whose fiber over $X \in \mathrm{ob}\,\underline{S}$ is the discrete category (i.e. the only arrows are the identity arrows) $F(X)$, the inverse-image functor by $u \in \mathrm{Fl}(\underline{S})$ being $F(u)$. Then an additive $\underline{S}$-function from $\underline{C}$ to F is precisely a cartesian $\underline{S}$-functor from $\underline{C}^{\mathrm{is}}$ to $\underline{F}$, inducing an additive function in the sense of (6.1) on each fiber. It will be convenient for us to interpret additivity as follows. To fix ideas, suppose that $\underline{C}$ is triangulated. The category $\underline{\mathrm{Cart}}_{\underline{S}}(\underline{C}^{\mathrm{is}}, \underline{F})$ (resp. $\underline{\mathrm{Cart}}_{\underline{S}}(\mathrm{Tr}(\underline{C})^{\mathrm{is}}, \underline{F})$ (1.1.2)) of cartesian $\underline{S}$-functors from $\underline{C}^{\mathrm{is}}$ (resp. $\mathrm{Tr}(\underline{C})^{\mathrm{is}}$) to $\underline{F}$ is the discrete category associated to an abelian [print 143]group, and we have a homomorphism

$$\phi : \underline{\mathrm{Cart}}_{\underline{S}}(\underline{C}^{\mathrm{is}}, \underline{F}) \longrightarrow \underline{\mathrm{Cart}}_{\underline{S}}(\mathrm{Tr}(\underline{C})^{\mathrm{is}}, \underline{F}),$$

defined by $\phi(f)(T) = f(L) - f(L') - f(L'')$ for $f \in \underline{\mathrm{Cart}}_{\underline{S}}(\underline{C}^{\mathrm{is}}, \underline{F})$ and[116] $T = (L' \to L \to L'' \to L'[1]) \in \mathrm{ob}\,\mathrm{Tr}(\underline{C})$. To say that $f \in \underline{\mathrm{Cart}}_{\underline{S}}(\underline{C}^{\mathrm{is}}, \underline{F})$ is additive means that $\phi(f) = 0$; in other words, we have a (canonical functorial) exact sequence

$$0 \longrightarrow \mathrm{Add}_{\underline{S}}(\underline{C}, F) \longrightarrow \underline{\mathrm{Cart}}_{\underline{S}}(\underline{C}^{\mathrm{is}}, \underline{F}) \xrightarrow{\phi} \underline{\mathrm{Cart}}_{\underline{S}}(\mathrm{Tr}(\underline{C})^{\mathrm{is}}, \underline{F}). \tag{6.6.1}$$

b) Let $u : X \to Y$ be an arrow of $\underline{S}$. The "inverse-image by u" functors, being all isomorphic to one another, define (6.3) a unique homomorphism $k(Y) \longrightarrow k(X)$. Consequently, $X \mapsto k(X)$ defines an abelian presheaf on $\underline{S}$, denoted by $k(\underline{C})$. An additive $\underline{S}$-function from $\underline{C}$ to F can then be interpreted simply as a homomorphism of presheaves $k(\underline{C}) \longrightarrow F$; in other words, we have

$$\mathrm{Add}_{\underline{S}}(\underline{C}, F) = \mathrm{Hom}(k(\underline{C}), F).$$

The presheaf $k(\underline{C})$ of course depends functorially on $\underline{C}$ with respect to additive (resp. exact) $\underline{S}$-functors.

Proposition 6.7. Let $\underline{S}$ be a site, $\underline{C}$ a flat abelian $\underline{S}$-category, and $\underline{C}_0$ a strictly full additive sub-$\underline{S}$-category of $\underline{C}$ satisfying the hypotheses (4.0). Then the morphism of presheaves

$$k(\underline{C}_0) \longrightarrow k(\mathrm{Parf}(\underline{C}, \underline{C}_0)) \qquad (\text{cf. } (4.9)),$$

induced by the inclusion of $\underline{C}_0$ in $\mathrm{Parf}(\underline{C}, \underline{C}_0)$, defines an isomorphism on the associated sheaves. In other words, for every abelian sheaf F on $\underline{S}$, [print 144]the restriction homomorphism

$$\mathrm{Add}_{\underline{S}}(\mathrm{Parf}(\underline{C}, \underline{C}_0), F) \longrightarrow \mathrm{Add}_{\underline{S}}(\underline{C}_0, F)$$

is an isomorphism.

Proof. Let F be an abelian sheaf on $\underline{S}$; let us show that the arrow

$$\mathrm{Add}_{\underline{S}}(\mathrm{Parf}(\underline{C}, \underline{C}_0), F) \longrightarrow \mathrm{Add}_{\underline{S}}(\underline{C}_0, F)$$

is an isomorphism. It factors as

$$\mathrm{Add}_{\underline{S}}(\mathrm{Parf}(\underline{C}, \underline{C}_0), F) \xrightarrow{a} \mathrm{Add}_{\underline{S}}(\mathrm{K}^b(\underline{C}_0), F) \xrightarrow{b} \mathrm{Add}_{\underline{S}}(\underline{C}_0, F).$$

Now, by (6.4), the arrow b is an isomorphism. We are therefore reduced to proving that a is an isomorphism.

Consider the following commutative diagram, in which the rows are exact (6.6.1), and the vertical arrows are the "restriction" arrows:

$$\begin{array}{ccccccc}
0 & \to & \mathrm{Add}_{\underline{S}}(\mathrm{Parf}(\underline{C}), F) & \to & \underline{\mathrm{Cart}}_{\underline{S}}(\mathrm{Parf}(\underline{C})^{\mathrm{is}}, \underline{F}) & \to & \underline{\mathrm{Cart}}_{\underline{S}}(\mathrm{Tr}(\mathrm{Parf}(\underline{C}))^{\mathrm{is}}, \underline{F}) \\
 & & \downarrow a & & \downarrow & & \downarrow \\
0 & \to & \mathrm{Add}_{\underline{S}}(\mathrm{K}^b(\underline{C}_0), F) & \to & \underline{\mathrm{Cart}}_{\underline{S}}(\mathrm{K}^b(\underline{C}_0)^{\mathrm{is}}, \underline{F}) & \to & \underline{\mathrm{Cart}}_{\underline{S}}(\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0))^{\mathrm{is}}, \underline{F}).
\end{array}$$

Since F is a sheaf, $\underline{F}$ is a stack (G II 2.2.4); thus, by (4.10), the two vertical arrows on the right are isomorphisms.

The same is therefore true of a, q.e.d.

Remark 6.7.1. (cf. 4.10.1). In (6.7), we can replace $\mathrm{Parf}(\underline{C}, \underline{C}_0)$ by $\mathrm{D}^*(\underline{C})_{\underline{C}_0\text{-parf}}$ ($* = -$ or b).

[print 145]6.8. Let $(\underline{S}, A)$ be a ringed topos and $\underline{C}$ and $\underline{C}_0$ the fibered $\underline{S}$-categories considered in (4.8). We have homomorphisms of abelian presheaves

$$k(\underline{C}_0) \xrightarrow{\alpha} k(\mathrm{D}^-({}_A\underline{S})_{\mathrm{parf}}) \xrightarrow{\beta} k(\mathrm{Parf}({}_A\underline{S})),$$

which, by (6.7), induce isomorphisms on the associated sheaves. The homomorphism α is compatible with inverse images, i.e. when $u : (\underline{S}', A') \longrightarrow (\underline{S}, A)$ is a morphism of ringed topoi, we have a commutative diagram, in which the vertical arrows are induced by the functor $\mathbb{L}u^*$ (cf. (4.19.1)):

$$\begin{array}{ccc} k(\underline{C}_0) & \longrightarrow & k(\mathrm{D}^-({}_A\underline{S})_{\mathrm{parf}}) \\ \downarrow & & \downarrow \\ k(\underline{C}'_0) & \longrightarrow & k(\mathrm{D}^-({}_{A'}\underline{S}')_{\mathrm{parf}}) \end{array}$$

($\underline{C}'_0$ being defined analogously to $\underline{C}_0$). When A is commutative, the tensor product $\overset{\mathrm{L}}{\otimes}$ equips $k(\underline{C}_0)$ and $k(\mathrm{D}^-({}_A\underline{S})_{\mathrm{parf}})$ with structures of presheaves of rings (cf. (4.19.1)), and α is a homomorphism of presheaves of rings.

6.9. In the situation of (6.8), suppose that $(\underline{S}, A)$ is locally ringed (cf. (2.15.1)). Then (2.15.1), for every $X \in \mathrm{ob}\,\underline{S}$, $\underline{C}_{0X}$ is the category of locally free A_X-Modules of finite type. Denote by $\underline{C}_1$ the fibered $\underline{S}$-category defined by $X \longmapsto$ (the category of free A_X-Modules of finite type). It is clear that $\underline{C}_1$ satisfies the hypotheses (4.0). An object of $\mathrm{D}(\underline{S})$ is perfect if and only if it is perfect relative to $\underline{C}_1$. We have homomorphisms of abelian presheaves

$$k(\underline{C}_1) \longrightarrow k(\underline{C}_0) \longrightarrow k(\mathrm{D}^-(\underline{S})_{\mathrm{parf}}) \longrightarrow k(\mathrm{Parf}(\underline{S}))$$

[print 146](the two on the left being homomorphisms of presheaves of rings), which, by (6.7), induce isomorphisms on the associated sheaves. We shall show that the associated sheaves are in fact canonically isomorphic to the constant sheaf $\mathbb{Z}_{\underline{S}}$.

Lemma 6.10. Let $(\underline{S}, A)$ be a locally ringed topos, U an object of $\underline{S}$ distinct from the empty object, and $p, q \in \mathbb{N}$. If we have $A_U^p \simeq A_U^q$, then $p = q$.

Proof. By an argument similar to the one given in the proof of (2.15.1), we reduce to the case where $(\underline{S}, A)$ is the locally ringed topos defined by a prescheme, and in this case the assertion is trivial (take a point in U).

Definition 6.11. We call rank, and denote by rg, the additive $\underline{S}$-function from $\underline{C}_1$ to $\mathbb{Z}_{\underline{S}}$ defined by

$$\mathrm{rg}(L) = p \quad \text{if } L \simeq A_X^p, \qquad X \neq \varnothing,$$
$$\mathrm{rg}(L) = 0 \quad \text{if } L \in \mathrm{ob}\,\underline{C}_{1X}, \qquad X = \varnothing.$$

By (6.7), the rank function extends uniquely to an additive $\underline{S}$-function from $\mathrm{Parf}(\underline{S})$ to $\mathbb{Z}_{\underline{S}}$ (which we shall again call rank and denote by rg).

Proposition 6.12. a) For $L, M \in \mathrm{ob}\,\mathrm{D}^-(X)_{\mathrm{parf}}$ ($X \in \mathrm{ob}\,\underline{S}$), we have $\mathrm{rg}(L \overset{\mathrm{L}}{\otimes} M) = \mathrm{rg}(L)\ \mathrm{rg}(M)$.

b) Let $u : (\underline{S}', A') \longrightarrow (\underline{S}, A)$ be a morphism of locally ringed topoi. For $M \in \mathrm{ob}\,\mathrm{D}^-(\underline{S})_{\mathrm{parf}}$, we have $\mathrm{rg}(\mathbb{L}u^*(M)) = u^*(\mathrm{rg}(M))$. In other words, the rank homomorphism: $k(\mathrm{D}^-(\underline{S})_{\mathrm{parf}}) \longrightarrow \mathbb{Z}_{\underline{S}}$ is a homomorphism of presheaves of rings, compatible with inverse images.

[print 147]Proof. It suffices to show that the homomorphism induced by rank on the sheaf associated to $k(\mathrm{D}^-(\underline{S})_{\mathrm{parf}})$ is a homomorphism of sheaves of rings compatible with inverse images. By (6.9), we can replace $k(\mathrm{D}^-(\underline{S})_{\mathrm{parf}})$ by $k(\underline{C}_1)$. We are therefore reduced to checking a) and b) for L and M free of finite type, which is trivial.

6.13. With $(\underline{S}, A)$ still denoting a locally ringed topos, let $X \in \mathrm{ob}\,\underline{S}$ and $\underline{n} \in H^0(X, \mathbb{Z}_{\underline{S}})$. We define an object of $\mathrm{Parf}(X)$, denoted by $A_X^{\underline{n}}$, as follows. When $\underline{n}$ is constant with value an integer $n \geq 0$ (resp. $n < 0$), we put $A_X^{\underline{n}} = A_X^n$ (resp. $A_X^{\underline{n}} = A_X^{-n}[1]$); in the general case, we decompose X into disjoint pieces X_i such that $\underline{n}|X_i$ is constant, and $A_X^{\underline{n}}$ is defined as the unique object of $\mathrm{Parf}(X)$ inducing $A_{X_i}^{\underline{n}|X_i}$ on each X_i. (More generally, we can define $L^{\underline{n}}$ for $L \in \mathrm{ob}\,\mathrm{Parf}(X)$, by putting $L^{\underline{n}} = A_X^{\underline{n}} \otimes L$.). The map $i_X : H^0(X, \mathbb{Z}_{\underline{S}}) \longrightarrow k(\mathrm{Parf}(X))$, defined by $i_X(\underline{n}) = \mathrm{cl}(A_X^{\underline{n}})$, is a ring homomorphism (such that $i_X(1) = 1$), and defines, as X varies, a homomorphism of presheaves of rings

$$i : \mathbb{Z}_{\underline{S}} \longrightarrow k(\mathrm{Parf}(\underline{S})).$$

Proposition 6.14. Under the hypotheses of (6.13), we have

$$\mathrm{rg}\ i = \mathrm{Id}(\mathbb{Z}_{\underline{S}})$$

(in other words, $k(\mathrm{Parf}(\underline{S}))$, considered as a presheaf of $\mathbb{Z}_{\underline{S}}$-algebras by means of i, is augmented toward $\mathbb{Z}_{\underline{S}}$ by rank). Moreover, rank

$$\mathrm{rg} : k(\mathrm{Parf}(\underline{S})) \longrightarrow \mathbb{Z}_{\underline{S}}$$

defines an isomorphism on the associated sheaves.

[print 148]Proof. The first assertion follows trivially from the relations $\mathrm{cl}(L[1]) = -\mathrm{cl}(L)$, $\mathrm{cl}(L^n) = n\,\mathrm{cl}(L)$ for $L \in \mathrm{ob}\,\mathrm{Parf}(\underline{S})$. Let us prove the second. First observe that i factors in the evident way through $k(\underline{C}_0)$. Since rg i is the identity, we are reduced (by (6.9)) to proving that the homomorphism $i\ \mathrm{rg} : k(\underline{C}_0) \longrightarrow k(\underline{C}_0)$ induces an isomorphism on the associated sheaves, i.e. that, for $L \in \mathrm{ob}\,\underline{C}_0$, $i\ \mathrm{rg}(L)$ is locally isomorphic to L, which is immediate.

We deduce, by (6.6.b)),

Corollary 6.15. Under the hypotheses of (6.13), for every abelian sheaf F on $\underline{S}$, the group of additive $\underline{S}$-functions from $\mathrm{Parf}(\underline{S})$ to F is canonically identified with $H^0(S, F)$ (S = final object of $\underline{S}$), the rank function corresponding to $F = \mathbb{Z}_{\underline{S}}$ and to the unit section of $\mathbb{Z}_{\underline{S}}$.

7. Duality of perfect complexes

7.0. In this section, $(\underline{S}, A)$ denotes a topos ringed in commutative rings; we denote by S the final object of $\underline{S}$. We propose to generalize to perfect complexes the notions of duality well known for locally free sheaves of finite type.

Proposition 7.1. Let $E \in \mathrm{ob}\,\mathrm{D}(S)_{\mathrm{parf}}$, $F \in \mathrm{ob}\,\mathrm{D}^+(S)$.

a) When F has locally bounded cohomology, the same holds for $\mathbb{R}\underline{\mathrm{Hom}}(E, F)$. When $F \in \mathrm{ob}\,\mathrm{D}(S)$ is pseudo-coherent (resp. locally of finite tor-dimension, resp. perfect), the same holds for $\mathbb{R}\underline{\mathrm{Hom}}(E, F)$.

[print 149]b) Suppose that E is moreover of finite tor-dimension (a condition satisfied when $\underline{S}$ is of finite type). Then, when F has bounded cohomology (resp. finite tor-dimension), the same holds for $\mathbb{R}\underline{\mathrm{Hom}}(E, F)$. More precisely, let a, b, m, n be integers such that $\mathrm{tor.amp}(E) \subset [a, b]$ (or, equivalently, by (5.8), $\mathrm{parf.amp}(E) \subset [a, b]$) and $\mathrm{H}^i(F) = 0$ for $i \notin [m, n]$ (resp. $\mathrm{tor.amp}(F) \subset [m, n]$). Then we have $\mathrm{H}^i(\mathbb{R}\underline{\mathrm{Hom}}(E, F)) = 0$ for $i \notin [m - b, n - a]$ (resp. $\mathrm{tor.amp}(\mathbb{R}\underline{\mathrm{Hom}}(E, F)) \subset [m - b, n - a]$).

Proof. Assertion a) is local, so we can suppose that E is strictly perfect, and we conclude by dévissage on E (reducing to $E = A$). For assertion b), we can suppose that E is strictly perfect with $E^i = 0$ for $i \notin [a, b]$, and $F^i = 0$ for $i \notin [m, n]$ (resp. F^i flat for every i and $F^i = 0$ for $i \notin [m, n]$). It then suffices to apply the following lemma:

Lemma 7.1.1. Let $E \in \mathrm{ob}\,\mathrm{D}^-(S)$, $F \in \mathrm{ob}\,\mathrm{D}^+(S)$. When E is strictly pseudo-coherent (i.e. (2.1) with components direct factors of free Modules of finite type), the canonical arrow of $\mathrm{D}^+(S)$

$$\underline{\mathrm{Hom}}^\bullet(E, F) \longrightarrow \mathbb{R}\underline{\mathrm{Hom}}(E, F)$$

is an isomorphism.

Proof. The spectral sequence

$$E_2^{pq} = \underline{\mathrm{H}}^p(\underline{\mathrm{Ext}}^q(E, F)^\bullet) \Longrightarrow \underline{\mathrm{Ext}}^*(E, F),$$

where

$$\underline{\mathrm{Ext}}^q(E, F)^i = \coprod_{i_2 - i_1 = i} \underline{\mathrm{Ext}}^q(E^{i_1}, F^{i_2}),$$

degenerates since $\underline{\mathrm{Ext}}^q(E, F)^\bullet = 0$ for every $q > 0$ (1.3.1). We deduce isomorphisms [print 150]

$$E_2^{p0} = \underline{\mathrm{H}}^p(\underline{\mathrm{Hom}}^\bullet(E, F)) \xrightarrow{\sim} \underline{\mathrm{Ext}}^p(E, F) = \underline{\mathrm{H}}^p(\mathbb{R}\underline{\mathrm{Hom}}(E, F)),$$

which proves the assertion.

Proposition 7.1.2. Let $f : (S', A') \longrightarrow (S, A)$ be a morphism of ringed topoi. Let $E \in \mathrm{ob}\,\mathrm{D}^-(S)$ and $F \in \mathrm{ob}\,\mathrm{D}^b(S)$ be such that $\mathbb{R}\underline{\mathrm{Hom}}(E, F) \in \mathrm{ob}\,\mathrm{D}^b(S)$ and $\mathbb{L}f^*(F) \in \mathrm{ob}\,\mathrm{D}^b(S')$. There exists a canonical functorial arrow

$$\mathbb{L}f^*\mathbb{R}\underline{\mathrm{Hom}}(E, F) \longrightarrow \mathbb{R}\underline{\mathrm{Hom}}(\mathbb{L}f^*E, \mathbb{L}f^*F).$$

It is an isomorphism when E is perfect.

Proof. By the adjunction of the functors $\mathbb{L}f^*$ and $\mathbb{R}f_*$, it suffices to define an arrow

$$\mathbb{R}\underline{\mathrm{Hom}}(E, F) \longrightarrow \mathbb{R}f_*\mathbb{R}\underline{\mathrm{Hom}}(\mathbb{L}f^*E, \mathbb{L}f^*F).$$

For this we use the adjunction arrow $F \to \mathbb{R}f_*\mathbb{L}f^*F$ and the duality isomorphism

$$\mathbb{R}f_*\mathbb{R}\underline{\mathrm{Hom}}(\mathbb{L}f^*E, \mathbb{L}f^*F) \simeq \mathbb{R}\underline{\mathrm{Hom}}(E, \mathbb{R}f_*\mathbb{L}f^*F).$$

To see that the arrow of (7.1.2) is an isomorphism for E perfect, we reduce trivially, by localization and dévissage, to the case $E = A$, in which it is a tautology.

Proposition 7.2. There exists (cf. SGA 5 I 1.6), for $E \in \mathrm{ob}\,\mathrm{D}(S)$ and $F \in \mathrm{ob}\,\mathrm{D}^+(S)$, a canonical functorial arrow

$$E \longrightarrow \mathbb{R}\underline{\mathrm{Hom}}(\mathbb{R}\underline{\mathrm{Hom}}(E, F), F).$$

It is an isomorphism when $E \in \mathrm{ob}\,\mathrm{D}(S)_{\mathrm{parf}}$ and $F = A$.

[print 151]Proof. We define the arrow by resolving F injectively and using the canonical arrow of $\mathrm{K}(S)$

$$E \longrightarrow \underline{\mathrm{Hom}}^\bullet(\underline{\mathrm{Hom}}^\bullet(E, F), F).$$

To see that the arrow in question is an isomorphism when E is perfect and $F = A$, we reduce, by localization and dévissage, to $E = A$, in which case it is evident.

Definition 7.3. Let $E \in \mathrm{ob}\,\mathrm{D}(S)$. We shall put $\mathbb{R}\underline{\mathrm{Hom}}(E, A) = E^\vee$, and say that $E^\vee$ is the dual of E.

By (7.1) and (7.2), when $E \in \mathrm{ob}\,\mathrm{D}(S)$ is perfect (resp. perfect and of finite tor-dimension), $E^\vee$ is also, and the canonical arrow $E \to E^{\vee\vee}$ is an isomorphism. In other words, the functor $\mathbb{R}\underline{\mathrm{Hom}}(\ \ , A)$ defines an anti-auto-equivalence of the category $\mathrm{D}(S)_{\mathrm{parf}}$ (resp. $\mathrm{D}(S)_{\underline{\mathrm{parf}}}$ (5.8.4.b))) with itself.

Lemma 7.4 (CARTAN isomorphism) (cf. ([H] II 5.15)). For $E \in \mathrm{ob}\,\mathrm{D}(S)$, $F \in \mathrm{ob}\,\mathrm{D}^-(S)$, $G \in \mathrm{ob}\,\mathrm{D}^+(S)$, there exist canonical functorial isomorphisms:

$$\mathrm{R}\underline{\mathrm{Hom}}(E \overset{\mathrm{L}}{\otimes} F, G) \xrightarrow{\sim} \mathrm{R}\underline{\mathrm{Hom}}(E, \mathrm{R}\underline{\mathrm{Hom}}(F, G)),$$
$$\mathrm{RHom}(E \overset{\mathrm{L}}{\otimes} F, G) \xrightarrow{\sim} \mathrm{RHom}(E, \mathrm{R}\underline{\mathrm{Hom}}(F, G)),$$
$$\mathrm{Hom}(E \overset{\mathrm{L}}{\otimes} F, G) \xrightarrow{\sim} \mathrm{Hom}(E, \mathrm{R}\underline{\mathrm{Hom}}(F, G)).$$

[print 152]Proof: This is standard: resolve G injectively and F flatly.

Lemma 7.5. For $E \in \mathrm{ob}\,\mathrm{D}^-(S)$, $F \in \mathrm{ob}\,\mathrm{D}^+(S)$, there exists a canonical functorial homomorphism

$$E \overset{\mathrm{L}}{\otimes} \mathrm{R}\underline{\mathrm{Hom}}(E, F) \longrightarrow F$$

Proposition 7.6 (cf. ([H] II 5.13)). Let $E \in \mathrm{ob}\,\mathrm{D}(S)$, $F \in \mathrm{ob}\,\mathrm{D}^+(S)$, $G \in \mathrm{ob}\,\mathrm{D}^+(S)$. Under one of the following hypotheses

i) G of finite tor-dimension,

ii) $E \in \mathrm{ob}\,\mathrm{D}^-(S)$, F of finite tor-dimension and $G \in \mathrm{ob}\,\mathrm{D}^b(S)$,

there exists a canonical functorial homomorphism

$$\mathrm{R}\underline{\mathrm{Hom}}(E, F) \overset{\mathrm{L}}{\otimes} G \longrightarrow \mathrm{R}\underline{\mathrm{Hom}}(E, F \overset{\mathrm{L}}{\otimes} G).$$

It is an isomorphism when E is perfect.

Proof. Let us define the arrow. Under hypothesis (i), it suffices to resolve F injectively and G flatly. Under hypothesis (ii), it is equivalent, by (7.4), to define an arrow

$$E \overset{\mathrm{L}}{\otimes} \mathbb{R}\underline{\mathrm{Hom}}(E, F) \overset{\mathrm{L}}{\otimes} G \longrightarrow F \overset{\mathrm{L}}{\otimes} G.$$

We apply the functor $\otimes^{\mathrm{L}} G$ to the arrow of (7.5).

To see that the arrow is an isomorphism when E is perfect, we use the standard method (cf. the proof of (7.2)): we reduce, by localization and dévissage, to the case $E = A$.

[print 153]**Corollary 7.7.** There exists, for $E \in \mathrm{ob}\,\mathrm{D}(S)_{\underline{\mathrm{parf}}}$ and $F \in \mathrm{ob}\,\mathrm{D}^+(S)$, a functorial isomorphism

$$E^\vee \overset{\mathrm{L}}{\otimes} F \xrightarrow{\sim} \mathbb{R}\underline{\mathrm{Hom}}(E, F),$$

with the notation of (7.3).

Proof. Apply (7.6), replacing F and G respectively by A and F, and use the fact (7.1) that $E^\vee \in \mathrm{ob}\,\mathrm{D}(S)_{\underline{\mathrm{parf}}}$.

Corollary 7.8. Let $E, F \in \mathrm{ob}\,\mathrm{D}(S)_{\underline{\mathrm{parf}}}$. There exists a natural arrow

$$\mathbb{R}\underline{\mathrm{Hom}}(E, F) \overset{\mathrm{L}}{\otimes} \mathbb{R}\underline{\mathrm{Hom}}(F, E) \longrightarrow A,$$

which is a "perfect duality", i.e. identifies, modulo (7.4), each of the complexes $\mathbb{R}\underline{\mathrm{Hom}}(E,F)$, $\mathbb{R}\underline{\mathrm{Hom}}(F,E)$ with the dual of the other (in the sense of (7.3)).

Proof. By (7.5), we have natural pairings $E^\vee \otimes^{\mathrm{L}} E \to A$, $F^\vee \otimes^{\mathrm{L}} F \to A$, hence, by tensor product, a natural pairing

$$E^\vee \overset{\mathrm{L}}{\otimes} F \overset{\mathrm{L}}{\otimes} F^\vee \overset{\mathrm{L}}{\otimes} E \longrightarrow A,$$

which, in view of (7.7), gives the stated pairing on the $\mathbb{R}\underline{\mathrm{Hom}}$. Let us show that it is a perfect duality. We have the canonical isomorphisms

$$\begin{aligned} \mathbb{R}\underline{\mathrm{Hom}}(F^\vee \overset{\mathrm{L}}{\otimes} E, A) &\simeq \mathbb{R}\underline{\mathrm{Hom}}(F^\vee, E^\vee) && (7.4)\\ &\simeq F^{\vee\vee} \overset{\mathrm{L}}{\otimes} E^\vee && (7.6)\\ &\simeq F \overset{\mathrm{L}}{\otimes} E^\vee && (7.2). \end{aligned}$$

Thus, after checking compatibility, we are done. We could also localize and use dévissage to reduce to $E = F = A$.

[print 154]**Exercise 7.9.** Let $E, F \in \mathrm{ob}\,\mathrm{D}(S)_{\underline{\mathrm{parf}}}$. Show that there exists a canonical isomorphism

$$E^\vee \overset{\mathrm{L}}{\otimes} F^\vee \xrightarrow{\sim} (E \overset{\mathrm{L}}{\otimes} F)^\vee.$$

8. Traces and cup-products

8.0. In this section, we generalize to perfect complexes the notion of the trace of an endomorphism of a projective module of finite type (see SGA 5 III for applications). The hypotheses and notation are those of (7.0).

8.1. Let $E \in \mathrm{ob}\,\mathrm{D}(S)_{\underline{\mathrm{parf}}}$ (5.8.4.b)). There exists a natural arrow

$$\tau : \mathbb{R}\underline{\mathrm{Hom}}(E,E) \longrightarrow A,$$

composed of the inverse isomorphism of (7.7), $\mathbb{R}\underline{\mathrm{Hom}}(E,E) \xrightarrow{\sim} E^\vee \otimes^{\mathrm{L}} E$, and the canonical arrow (7.5) $E^\vee \otimes^{\mathrm{L}} E \to A$.

Let $U \in \mathrm{ob}\,\underline{S}$. Applying the functor $H^0(U,.)$ to τ, and using the fact that $H^0(U, \mathbb{R}\underline{\mathrm{Hom}}(E,E))$ is identified with $\mathrm{Hom}_{\mathrm{D}(U)}(E|U, E|U)$, we deduce a homomorphism of $H^0(U,A)$-modules,

$$\mathrm{Tr}_{E|U} : \mathrm{Hom}_{\mathrm{D}(U)}(E|U, E|U) \longrightarrow H^0(U,A),$$

which is called the trace homomorphism(*).

Before going further, let us give some properties of the trace that follow immediately from the definition.

[print 155]8.1.1. When $E = A$, $\mathrm{Tr}_{A|U} : \mathrm{Hom}(A|U, A|U) \longrightarrow H^0(U,A)$ is precisely the well-known isomorphism. When E is a direct factor of a free Module of finite type, a homomorphism from E to E over U is locally defined by

$$e \longmapsto \sum \langle e, x_i^\vee \rangle x_i,$$

where x_i (resp. $x_i^\vee$) is a local section of E (resp. $E^\vee$); the corresponding trace is $\sum \langle x_i, x_i^\vee \rangle$. We easily deduce that, when E and F are direct factors of free Modules of finite type, and

(*) The finite tor-dimension hypothesis on E is in fact superfluous for the definition of $\mathrm{Tr}_{E|U}$, see (8.5).

$u : (E+F)|U \longrightarrow (E+F)|U$ is defined by a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$, we have

$$\mathrm{Tr}_{(E+F)|U}(u) = \mathrm{Tr}_{E|U}(a) + \mathrm{Tr}_{F|U}(d).$$

8.1.2. Let E be a strictly perfect complex, $U \in \mathrm{ob}\,\underline{S}$, and $u : E|U \longrightarrow E|U$ a homomorphism of complexes, with components $u^i : E^i|U \to E^i|U$. We then have

$$\mathrm{Tr}_{E|U}(u) = \sum (-1)^i \,\mathrm{Tr}_{E^i|U}(u^i).$$

Indeed, since the question is local, we can suppose that u is defined by a section of $\underline{H}^0(E^\vee \otimes E)$, and even that this section is of the form $\coprod_i \sum_j x_{ij}^\vee \otimes x_{ij}$, where x_{ij} (resp. $x_{ij}^\vee$) is a section of E^i (resp. $E^{i\vee}$). Making the arrow (7.5) $E^\vee \otimes E \to A$ explicit, we find

$$\mathrm{Tr}_{E|U}(u) = \sum (-1)^i \langle x_{ij}, x_{ij}^\vee \rangle = \sum (-1)^i \,\mathrm{Tr}_{E^i|U}(u^i),$$

q.e.d.

8.2. Let $E, F \in \mathrm{ob}\,\mathrm{D}(S)_{\underline{\mathrm{parf}}}$. We have seen (7.8) that there exists a canonical homomorphism

$$\gamma : \mathbb{R}\underline{\mathrm{Hom}}(E,F) \overset{\mathrm{L}}{\otimes} \mathbb{R}\underline{\mathrm{Hom}}(F,E) \longrightarrow A.$$

Let $U \in \mathrm{ob}\,\underline{S}$, and take $u \in H^0(U, \mathbb{R}\underline{\mathrm{Hom}}(E,F))$ and $v \in H^0(U, \mathbb{R}\underline{\mathrm{Hom}}(F,E))$. [print 156]We can interpret u (resp. v) as a homomorphism, in $\mathrm{D}(U)$, from $A|U$ to $\mathbb{R}\underline{\mathrm{Hom}}(E,F)|U$ (resp. $\mathbb{R}\underline{\mathrm{Hom}}(F,E)|U$), and form

$$u \overset{\mathrm{L}}{\otimes} v : A|U \longrightarrow (\mathbb{R}\underline{\mathrm{Hom}}(E,F) \overset{\mathrm{L}}{\otimes} \mathbb{R}\underline{\mathrm{Hom}}(F,E))|U.$$

The map $(u,v) \mapsto \langle u, v\rangle = \gamma \circ (u \otimes^{\mathrm{L}} v)$ defines an $H^0(U,A)$-linear map,

$$\mathrm{Hom}_{\mathrm{D}(U)}(E|U, F|U) \otimes_{H^0(U,A)} \mathrm{Hom}_{\mathrm{D}(U)}(F|U, E|U) \longrightarrow H^0(U,A),$$

which we shall call the cup-product(*).

The relation between trace and cup-product is expressed in

Proposition 8.3. In the situation of (8.2), for $u \in \mathrm{Hom}(E|U, F|U)$ and $v \in \mathrm{Hom}(F|U, E|U)$, we have

$$\mathrm{Tr}_{E|U}(vu) = \mathrm{Tr}_{F|U}(uv) = \langle u, v\rangle.$$


In particular, when $E = F$, we have

$$\mathrm{Tr}_{E|U}(u) = \langle u, \mathrm{Id}\rangle = \langle \mathrm{Id}, u\rangle.$$

Proof. For simplicity, we shall suppose that $U = S$. We therefore have two arrows $u : A \longrightarrow E^\vee \otimes^{\mathrm{L}} F$ and $v : A \longrightarrow F^\vee \otimes^{\mathrm{L}} E$, hence

$$u \overset{\mathrm{L}}{\otimes} v : A \longrightarrow E^\vee \overset{\mathrm{L}}{\otimes} F \overset{\mathrm{L}}{\otimes} F^\vee \overset{\mathrm{L}}{\otimes} E.$$

On the other hand, we have $vu : A \to E^\vee \otimes^{\mathrm{L}} E$ and $uv : A \to F^\vee \otimes^{\mathrm{L}} F$. The proposition follows from the commutativity of the following diagrams (whose verification is "purely mechanical"): [print 157]



(*) the same remark as for the definition of $\mathrm{Tr}_{E|U}$

[117]

$$\begin{array}{ccc} E^\vee \otimes^{\mathrm{L}} F \otimes^{\mathrm{L}} F^\vee \otimes^{\mathrm{L}} E & \xrightarrow{\mathrm{Id} \otimes^{\mathrm{L}} \tau_E} & F^\vee \otimes^{\mathrm{L}} F \\ \downarrow{\scriptstyle \mathrm{Id} \otimes^{\mathrm{L}} \tau_F} & & \downarrow{\scriptstyle \tau_F} \\ E^\vee \otimes^{\mathrm{L}} E & \xrightarrow[\tau_E]{} & A. \end{array}$$

Corollary 8.4. E, F, U being given as in (8.2), let $u \in \mathrm{Hom}(E|U, E|U)$, $s : E|U \to F|U$ be an isomorphism, so that $sus^{-1} \in \mathrm{Hom}(F|U, F|U)$. We then have

$$\mathrm{Tr}_{F|U}(sus^{-1}) = \mathrm{Tr}_{E|U}(u).$$

Proof. This is immediate.

Remark 8.5. When $\underline{C}$ is a fibered $\underline{S}$-category, denote by $E(\underline{C})$ the fibered $\underline{S}$-category whose fiber over $U \in \mathrm{ob}\,\underline{S}$ is the category with objects the pairs (L, u), where $L \in \mathrm{ob}\,\underline{C}_U$ and $u \in \mathrm{End}_{\underline{S}}(L)$, and with arrows from (L, u) to (M, v) the arrows $f : L \to M$ such that $fu = vf$. By (8.4), we can regard the trace homomorphism as an $\underline{S}$-functor from $E(\mathrm{D}(\underline{S})_{\mathrm{parf}})^{\mathrm{is}}$ to $\underline{A}$ (where the exponent is means that the only arrows considered are the isomorphisms, and $\underline{A}$ is the $\underline{S}$-category with discrete fibers defined by A (cf. (6.6.a))). Since every perfect complex is locally of finite tor-dimension, the forgetful $\underline{S}$-functor

$$E(\mathrm{D}(\underline{S})_{\mathrm{parf}})^{\mathrm{is}} \longrightarrow E(\mathrm{D}(\underline{S})_{\mathrm{parf}})^{\mathrm{is}}$$

induces an isomorphism on the associated stacks. Consequently, the trace [print 158]homomorphism extends uniquely to an $\underline{S}$-functor from $E(\mathrm{D}(\underline{S})_{\mathrm{parf}})^{\mathrm{is}}$ to $\underline{A}$.

For $E \in \mathrm{ob}\,\mathrm{Parf}(U)$ ($U \in \mathrm{ob}\,\underline{S}$), we again denote by

$$\mathrm{Tr}_E : \mathrm{Hom}(E, E) \longrightarrow H^0(U, A)$$

the corresponding homomorphism, which is a homomorphism of $H^0(U, A)$-modules. We see similarly that the definition of the cup-product (8.2), as well as assertions (8.3) and (8.4), generalize to the case of perfect complexes (not necessarily of finite tor-dimension).

Proposition 8.6. Let $U \in \mathrm{ob}\,\underline{S}$, $E, F \in \mathrm{Parf}(U)$, $u \in \mathrm{Hom}(E, E)$, $v \in \mathrm{Hom}(F, F)$, so that $u \otimes^{\mathrm{L}} v \in \mathrm{Hom}(E \otimes^{\mathrm{L}} F, E \otimes^{\mathrm{L}} F)$. We have

$$\mathrm{Tr}_{E \otimes^{\mathrm{L}} F}(u \overset{\mathrm{L}}{\otimes} v) = \mathrm{Tr}_E(u)\ \mathrm{Tr}_F(v).$$

Proof. Since the question is local, by (8.4) and (4.10) we can suppose that u and v are arrows of $K^b(\underline{C}_0)$. Using (8.1.2), we reduce to the case where u and v are arrows of $\underline{C}_0$. We conclude by (8.1.1).

[print 159]

BIBLIOGRAPHY

[H] Hartshorne, R. Residues and Duality, Lecture Notes no. 20, Springer, 1966.

[V] Verdier, J-L. Catégories dérivées, I.H.E.S., 1963.

[G] Giraud, J. Cohomologie non abélienne, to appear in North Holl. Pub. Comp.

SGA 4 Cohomologie étale des schémas, by M. Artin, A. Grothendieck, and J-L. Verdier, to appear in North Holl. Pub. Comp.

SGA 5 Cohomologie ℓ-adique et fonctions L, by A. Grothendieck, to appear in North Holl. Pub. Comp.

[**1**] Hakim, M. Topos annelés et schémas relatifs, thesis (in preparation).

[**2**] Verdier, J-L. Les catégories dérivées des catégories abéliennes, to appear in North Holl. Pub. Comp.

[**MULT**] Cartan, H. and Eilenberg, S. Homological Algebra, Princeton, 1956.

[print 160]

Exposé II

EXISTENCE OF GLOBAL RESOLUTIONS

by L. ILLUSIE

1. General globalization criteria

1.0. In this section, we are given a site $\underline{S}$, a flat abelian $\underline{S}$-category $\underline{C}$ (I 1.1), and a strictly full additive sub-$\underline{S}$-category $\underline{C}_0$ (I 1.1). We assume that $\underline{C}_0$ is locally quasi-liftable in $\underline{C}$ (I 1.2) and locally quasi-stable under kernels of epimorphisms (I 1.2).

Proposition 1.1. Let $S \in \mathrm{ob}\,\underline{S}$, and suppose that:

(i) $\underline{C}_{0S}$ is quasi-liftable in $\underline{C}_S$ (I 1.2) ($\underline{C}_S$ being regarded as a category fibered over a punctual site);

(ii) every object of $\underline{C}_S$ which is of $\underline{C}_0$-finite type (I 1.2) (resp. of $\underline{C}_0$-finite presentation (I 2.8)) is of $\underline{C}_{0S}$-finite type.

Then:

a) $\underline{C}_{0S}$ is quasi-stable under kernels of epimorphisms (I 1.2) ($\underline{C}_{0S}$ being regarded as a category fibered over a punctual site).

b) Let $E \in \mathrm{ob}\,\mathrm{D}^-(\underline{C}_S)$ and $n \in \mathbb{Z}$. If E is n-$\underline{C}_0$-pseudo-coherent (I 2.3), E is n-$\underline{C}_{0S}$-pseudo-coherent (resp. $(n+1)$-$\underline{C}_{0S}$-pseudo-coherent). In other words, we have

$$\mathrm{D}^-(\underline{C}_S)_{n\text{-}\underline{C}_0\text{-coh}} = \mathrm{D}^-(\underline{C}_S)_{n\text{-}\underline{C}_{0S}\text{-coh}}$$

(resp.

$$\mathrm{D}^-(\underline{C}_S)_{n\text{-}\underline{C}_0\text{-coh}} \subset \mathrm{D}^-(\underline{C}_S)_{(n+1)\text{-}\underline{C}_{0S}\text{-coh}} \subset \mathrm{D}^-(\underline{C}_S)_{(n+1)\text{-}\underline{C}_0\text{-coh}},$$

inclusions of strictly full subcategories of $\mathrm{D}(\underline{C}_S)$, the inclusion on the right being trivial, moreover).

c) For $E \in \mathrm{ob}\,\mathrm{D}^-(\underline{C}_S)$ to be pseudo-coherent relative to $\underline{C}_0$ (I 2.3), it is necessary and sufficient that it be so relative to $\underline{C}_{0S}$; in other words, we have

$$\mathrm{D}^-(\underline{C}_S)_{\underline{C}_0\text{-coh}} = \mathrm{D}^-(\underline{C}_S)_{\underline{C}_{0S}\text{-coh}}.$$

Moreover, the canonical functor

$$\mathrm{K}^-(\underline{C}_{0S})/\mathrm{K}^{-,\varnothing}(\underline{C}_{0S}) \longrightarrow \mathrm{D}(\underline{C}_S),$$

where $\mathrm{K}^{-,\varnothing}(\underline{C}_{0S})$ denotes the thick subcategory of $\mathrm{K}^-(\underline{C}_{0S})$ formed by [print 161]acyclic complexes, is fully faithful and has as essential image $\mathrm{D}^-(\underline{C}_S)_{\underline{C}_0\text{-coh}}$. If $\underline{C}_{0S}$ is liftable in $\underline{C}_S$ (I 1.2) (i.e. if the objects of $\underline{C}_{0S}$ are projective), the category $\mathrm{K}^{-,\varnothing}(\underline{C}_{0S})$ is reduced to zero objects, and consequently the functor

$$\mathrm{K}^-(\underline{C}_{0S}) \longrightarrow \mathrm{D}(\underline{C}_S)$$

is fully faithful (and has as essential image $\mathrm{D}^-(\underline{C}_S)_{\underline{C}_0\text{-coh}}$).

Proof. a) Let

$$0 \longrightarrow L \longrightarrow L^0 \longrightarrow \ldots \longrightarrow L^m \longrightarrow 0$$

be an exact sequence in $\underline{C}_S$, with $L^i \in \mathrm{ob}\,\underline{C}_{0S}$ for every i; we have to show that L is of $\underline{C}_{0S}$-finite type. Now, by (I 2.2), L is $\underline{C}_0$-pseudo-coherent, hence (I 2.9) of $\underline{C}_0$-finite presentation, and the assertion follows from (ii).

b) Let us show, by descending induction on i, that E is i-$\underline{C}_{0S}$-pseudo-coherent for $i \geq n$ (resp. $i \geq n+1$). We may already assume that E is strictly $(n+1)$-$\underline{C}_{0S}$-pseudo-coherent (resp. $(n+2)$-$\underline{C}_{0S}$-pseudo-coherent) (I 2.1), and we have to prove that E is n-$\underline{C}_{0S}$-pseudo-coherent (resp. $(n+1)$-$\underline{C}_{0S}$-pseudo-coherent). We therefore have an exact sequence

$$0 \longrightarrow E' \longrightarrow E \longrightarrow E'' \longrightarrow 0,$$

where E' is strictly $\underline{C}_{0S}$-perfect (I 2.1) and $E''^i = 0$ for $i > n$ (resp. $i > n+1$). Replacing E by E'', we may therefore assume (I 2.6) that $E^i = 0$ for $i > n$ (resp. $i > n+1$). Then, by (I 2.10 a)), $H^n(E)$ (resp. $H^{n+1}(E)$) is of $\underline{C}_0$-finite type (resp. of $\underline{C}_0$-finite presentation). Thus, by (ii), $H^n(E)$ (resp. $H^{n+1}(E)$) is of $\underline{C}_{0S}$-finite type. Applying (I 2.10 b)) again, we deduce that E is n-$\underline{C}_{0S}$-pseudo-coherent (resp. $(n+1)$-$\underline{C}_{0S}$-pseudo-coherent), q.e.d. (N.B. hypothesis (i) and assertion a) were used implicitly in the references to (I 2.6 and 2.10)).

c) The first assertion follows from b), the second from (I 2.7); the last is evident.

Proposition 1.2. Suppose that $\underline{C}_0$ is locally liftable in $\underline{C}$ (I 1.2) and locally stable under kernels of epimorphisms (I 1.2), and make the [print 162]hypotheses (1.1 (i) and (ii) in the resp. case). Denote by $\underline{C}_{1S}$ the strictly full subcategory of $\underline{C}_S$ formed by objects which are locally in $\underline{C}_0$. Then:

a) $\underline{C}_{1S}$ is stable under kernels of epimorphisms (I 1.2) and quasi-liftable in $\underline{C}_S$ (I 1.2) (as a category fibered over a punctual site).

b) Let $E \in \mathrm{ob}\,\mathrm{D}(\underline{C}_S)$ and $[m,n]$ be an interval in $\mathbb{Z}$. For E to have $\underline{C}_0$-perfect amplitude contained in $[m,n]$, it is necessary and sufficient that E be isomorphic, in $\mathrm{D}(\underline{C}_S)$, to a complex F such that $F^i = 0$ for $i \notin [m,n]$, $F^i \in \mathrm{ob}\,\underline{C}_{0S}$ for $i > m$, and $F^m \in \mathrm{ob}\,\underline{C}_{1S}$.

c) The canonical functor

$$\mathrm{K}^b(\underline{C}_{1S})/\mathrm{K}^{b,\varnothing}(\underline{C}_{1S}) \longrightarrow \mathrm{D}(\underline{C}_S)$$

is fully faithful and has as essential image the full subcategory of $\mathrm{D}(\underline{C}_S)$ formed by complexes of finite $\underline{C}_0$-perfect amplitude.

d) If $\underline{C}_{1S}$ is not only quasi-liftable but liftable in $\underline{C}_S$, one may replace in b) the expression "E is isomorphic to F in $\mathrm{D}(\underline{C}_S)$" by "there exists a quasi-isomorphism $F \to E$", and in c) the category $\mathrm{K}^{b,\varnothing}(\underline{C}_{1S})$ is reduced to zero objects.

Proof. a) It is clear that $\underline{C}_{1S}$ is stable under kernels of epimorphisms. On the other hand, the objects of $\underline{C}_{1S}$ are $\underline{C}_0$-pseudo-coherent, hence $\underline{C}_{0S}$-pseudo-coherent by (1.1 c)); the second assertion therefore follows from (I 2.14) (where we take the punctual site for $\underline{S}$).

b) It suffices to prove necessity. We already know, by (1.1 c)), that E is $\underline{C}_{0S}$-pseudo-coherent. Since E is acyclic in degree $> n$, we may therefore assume (I 2.10), after replacing E by an isomorphic complex in $\mathrm{D}^-(\underline{C}_S)$, that E is strictly $(m+1)$-$\underline{C}_{0S}$-pseudo-coherent and that $E^i = 0$ for $i > n$. Since E is acyclic in degree $< m$, we may further assume, after replacing E by its truncation

$$0 \longrightarrow E^m/B^m \longrightarrow E^{m+1} \longrightarrow \ldots \longrightarrow E^n \longrightarrow 0,$$

that $E^i = 0$ for $i < m$. Then, by (I 4.16), E^m is locally an object of $\underline{C}_0$, hence an object of $\underline{C}_{1S}$, which proves the assertion.

[print 163]c) In view of (1.2 a)), the assertion follows from (1.2 b)) and (I 4.12.2).

d) The fact that $\mathrm{K}^{b,\varnothing}(\underline{C}_{1S})$ is reduced to zero objects is evident. The other assertion follows from (I 4.6) (applied over the punctual site).

2. Application to certain categories of Modules

2.0. Notation

Let $(\underline{S}, A)$ be a ringed topos. For $U \in \mathrm{ob}\,\underline{S}$, we write $\mathrm{Mod}(U)$ for the category of left A_U-Modules, and $\mathrm{Lib}(U)$ (resp. $\mathrm{Loclib}(U)$) for the full subcategory of $\mathrm{Mod}(U)$ formed by free A_U-Modules (resp. Modules locally direct factors of free Modules) of finite type.

2.1. Lifting lemmas

Lemma 2.1.1 (cf. EGA 0_{I} 5.2.3). Let $(\underline{S}, A)$ be a ringed topos, and $S \in \mathrm{ob}\,\underline{S}$. We assume that S is quasi-compact[118], i.e. that every covering family $(S_i \to S)$ is dominated by a finite covering family.

a) Let F be a left A_S-Module which is the inductive limit, over a filtered category Λ, of A_S-Modules F_λ, and let

$$u = \varinjlim u_\lambda : F \longrightarrow G$$

be an epimorphism, where G is an A_S-Module of finite type. Then there exists $\lambda \in \Lambda$ such that u_λ is an epimorphism.

b) Let F be a left A_S-Module generated by its sections (i.e. such that there exists an epimorphism $A_S^{(I)} \to F$ for a suitable index set I). Then, if F is of finite type, there exists an epimorphism $A_S^p \to F$ for $p \in \mathbb{N}$.

The proof of a) is analogous to that of (loc. cit.) and is left as an exercise to the reader, and b) follows immediately from a).

Lemma 2.1.2. Let $(\underline{S}, A)$ be a ringed topos, $\underline{C}$ the fibered $\underline{S}$-category of left A-Modules, $\underline{C}_0$ the fibered subcategory of free A-Modules of finite type (I 1.3 d)), $S \in \mathrm{ob}\,\underline{S}$, $\underline{C}'_S$[119] a strictly full additive subcategory of $\underline{C}_S$ containing $\underline{C}_{0S}$ and stable under kernels, cokernels and extensions. We assume that [print 164]

$$H^1(S, F) = 0 \quad \text{for every } F \in \mathrm{ob}\,\underline{C}'_S.$$

Then:

a) Every exact sequence in $\underline{C}'_S$:

$$0 \longrightarrow E \longrightarrow F \longrightarrow G \longrightarrow 0,$$

such that $\underline{\mathrm{Hom}}(G, E) \in \mathrm{ob}\,\underline{C}'_S$ and such that G is locally a direct factor of a free Module of finite type, is split. The category $\underline{C}_{0S}$ is liftable in $\underline{C}'_S$ (I 1.2).

b) Denote by $\mathrm{Mod}(A(S))$ the category of $A(S)$-modules. The functor

$$\Gamma_S : \underline{C}'_S \longrightarrow \mathrm{Mod}(A(S))$$

is exact. Moreover:

(i) If every object F of $\underline{C}'_S$ admits a finite global presentation

$$A_S^p \longrightarrow A_S^q \longrightarrow F \longrightarrow 0,$$

the functor $\Gamma_S : \underline{C}'_S \longrightarrow \mathrm{Mod}(A(S))$ is fully faithful and has as essential image the full subcategory of $\mathrm{Mod}(A(S))$ formed by $A(S)$-modules of finite presentation;

(ii) If $\underline{C}'_S$ contains the free A_S-Modules, if every object F of $\underline{C}'_S$ admits a global presentation

$$A_S^{(I)} \longrightarrow A_S^{(J)} \longrightarrow F \longrightarrow 0,$$

and if the functor Γ_S commutes with filtered inductive limits([1]), the functor $\Gamma_S : \underline{C}'_S \longrightarrow \mathrm{Mod}(A(S))$ is an equivalence of categories.

Proof. a) Since G is locally a direct factor of a free Module of finite type, the spectral sequence

$$E_2^{pq} = H^p(S, \underline{\mathrm{Ext}}^q(G, E)) \Longrightarrow \mathrm{Ext}^*(G, E)$$

degenerates (I 1.3.1) and gives an isomorphism

$$H^1(S, \underline{\mathrm{Hom}}(G, E)) \simeq \mathrm{Ext}^1(G, E),$$

[print 165]whence the first assertion. The second is a particular case, since, for $G \in \mathrm{ob}\,\underline{C}_{0S}$, i.e. G free and of finite type, and $E \in \mathrm{ob}\,\underline{C}'_S$, we have $\underline{\mathrm{Hom}}(G, E) \in \mathrm{ob}\,\underline{C}'_S$.

b) The first assertion is trivial. Let us now prove, in case (i) or (ii), the full faithfulness of $\Gamma_S : \underline{C}'_S \longrightarrow \mathrm{Mod}(A(S))$. Let $F, G \in \mathrm{ob}\,\underline{C}'_S$, and let

$$A_S^{(I)} \longrightarrow A_S^{(J)} \longrightarrow F \longrightarrow 0 \qquad (*)$$

be a global presentation of F (with I and J finite in case (i)). Applying to $(*)$ the functor Γ_S, which is exact on $\underline{C}'_S$ and commutes, in case (ii), with filtered inductive limits, we obtain an exact sequence

$$A(S)^{(I)} \longrightarrow A(S)^{(J)} \longrightarrow \Gamma_S(F) \longrightarrow 0. \qquad (**)$$

On the other hand, applying to $(*)$ (resp. $(**)$) the left-exact functor $\mathrm{Hom}_{A_S}(., G)$ (resp. $\mathrm{Hom}_{A(S)}(., \Gamma_S(G))$[120]), we obtain a commutative diagram with exact rows: [121]

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathrm{Hom}_{A_S}(F, G) & \longrightarrow & \mathrm{Hom}_{A_S}(A_S^{(J)}, G) & \longrightarrow & \mathrm{Hom}_{A_S}(A_S^{(I)}, G) \\
 & & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathrm{Hom}_{A(S)}(\Gamma_S(F), \Gamma_S(G)) & \longrightarrow & \mathrm{Hom}_{A(S)}(A(S)^{(J)}, \Gamma_S(G)) & \longrightarrow & \mathrm{Hom}_{A(S)}(A(S)^{(I)}, \Gamma_S(G)).
\end{array}$$

The two vertical arrows on the right are visibly isomorphisms; the same[122] is therefore true of the vertical arrow on the left, which proves the full faithfulness of $\Gamma_S : \underline{C}'_S \longrightarrow \mathrm{Mod}(A(S))$. The characterization of the essential image, in case (i) or (ii), is immediate and is left to the reader.

Lemma 2.1.3. Let A and B be abelian categories, and $\phi : A \to B$ an exact functor.

([1]) a hypothesis satisfied when $\underline{S}$ is locally of finite type (SGA 4 VI 1.2) or when $\underline{S}$ is the topos of sheaves of sets on a compact space and S the final object (T.F II 3.10.1).

a) For $M \in \mathrm{ob}\, A$, the following conditions are equivalent:

(i) for every $L \in \mathrm{ob}\, A$, the canonical arrow (induced by ϕ)

$$\mathrm{Ext}^n_A(L, M) \longrightarrow \mathrm{Ext}^n_B(\phi(L), \phi(M))$$

is an isomorphism for every n;

(ii) for every $L \in \mathrm{ob}\, A$, the canonical arrow [print 166]

$$\mathrm{Hom}_A(L, M) \longrightarrow \mathrm{Hom}_B(\phi(L), \phi(M))$$

is an isomorphism, and for every $n \geq 1$ and every $x \in \mathrm{Ext}^n_B(\phi(L), \phi(M))$ there exists an epimorphism $L' \to L$ in A which effaces x.

b) The following conditions are equivalent:

(i) for all $L, M \in \mathrm{ob}\, A$ and every $n \in \mathbb{Z}$, the canonical arrow

$$\mathrm{Ext}^n_A(L, M) \longrightarrow \mathrm{Ext}^n_B(\phi(L), \phi(M))$$

is an isomorphism;

(ii) the canonical functor (induced by ϕ)

$$\mathrm{D}^b(A) \longrightarrow \mathrm{D}^b(B)$$

is fully faithful and has as essential image the full subcategory of $\mathrm{D}^b(B)$ formed by complexes F such that $H^i(F)$ is in the essential image of ϕ for every i.

Proof. a) For the implication (i)⇒(ii), it suffices to show the following: given $n \geq 1$, $L \in \mathrm{ob}\, A$, and $y \in \mathrm{Ext}^n_A(L, M)$, there exists an epimorphism $L' \to L$ which effaces y. For this, one can use Yoneda's interpretation ([18] III 3.2): y is the class of an exact sequence

$$0 \longrightarrow M \longrightarrow E_{n-1} \longrightarrow \ldots \longrightarrow E_0 \longrightarrow L \longrightarrow 0;$$

the epimorphism $E_0 \to L$ effaces y ([18] III 3.2.5 and 3.2.8). Let us prove (ii)⇒(i). For brevity, put

$$\mathrm{Ext}^n_A(., M) = F^n, \qquad \mathrm{Ext}^n_B(\phi(.), \phi(M)) = G^n.$$

The effaceability hypothesis means that, for $n \geq 1$ and $L \in \mathrm{ob}\, A$, we have

$$G^n(L) = \varinjlim \mathrm{Coker}(G^{n-1}(L_0) \to G^{n-1}(L_1)),$$

the inductive limit being taken over the set of exact sequences [print 167] $0 \to L_1 \to L_0 \to L \to 0$. From what we saw above, we likewise have

$$F^n(L) = \varinjlim \mathrm{Coker}(F^{n-1}(L_0) \to F^{n-1}(L_1)).$$

since $\phi = \phi^0 : F^0 \to G^0$ is an isomorphism, it follows, by induction on n, that the morphism $\phi^n : F^n \to G^n$ induced by ϕ is an isomorphism for every n.

b) The equivalence (i)⇔(ii) follows from an immediate dévissage, left to the reader.

Remarks 2.1.3.1. There is of course a dual equivalence to a), which we leave the reader to state. On the other hand, when A and B have enough injectives, conditions (i) and (ii) of b) are equivalent to

(iii) for $L \in \mathrm{ob}\, \mathrm{D}^-(A)$ and $M \in \mathrm{ob}\, \mathrm{D}^+(A)$, the canonical arrow

$$\underline{\underline{R}}\mathrm{Hom}(L, M) \longrightarrow \underline{\underline{R}}\mathrm{Hom}(\phi(L), \phi(M))$$

is an isomorphism,

as one sees immediately by a spectral-sequence argument.

2.2. Noetherian schemes and divisorial schemes

Notation 2.2.1. Let S be a scheme. We write $\mathrm{Qcoh}(S)$ (resp. $\mathrm{Coh}(S)$) for the (abelian) category of quasi-coherent (resp. coherent) $\underline{O}_S$-Modules. Let $[m,n]$ be an interval in $\mathbb{Z}$. We shall say that an object E of $\mathrm{D}(\mathrm{Qcoh}(S))$ is n-pseudo-coherent (resp. pseudo-coherent, resp. of perfect amplitude contained in $[m,n]$, resp. perfect) when E is n-pseudo-coherent (resp. pseudo-coherent, resp. of perfect amplitude contained in $[m,n]$, resp. perfect) relative to the category fibered over the Zariski site of S defined by $U \mapsto \mathrm{Loclib}(U)$ ((I 2.3), (I 4.7) and (2.0)). We denote by $\mathrm{D}^-(\mathrm{Qcoh}(S))_{\mathrm{coh}}$ (resp. $\mathrm{D}(\mathrm{Qcoh}(S))_{\mathrm{parf}}$) the full triangulated subcategory of $\mathrm{D}(\mathrm{Qcoh}(S))$ formed by pseudo-coherent (resp. perfect) complexes.

Proposition 2.2.2(*). Let S be a noetherian scheme. The canonical functor [print 168]

$$\mathrm{D}^-(\mathrm{Coh}(S)) \longrightarrow \mathrm{D}(\mathrm{Qcoh}(S))$$

is fully faithful and has as essential image $\mathrm{D}^-(\mathrm{Qcoh}(S))_{\mathrm{coh}}$[123].

Proof. We apply (1.1), taking for $\underline{S}$ the Zariski site of S, and for $\underline{C}$ (resp. $\underline{C}_0$) the fibered category $U \mapsto \mathrm{Qcoh}(U)$ (resp. $\mathrm{Coh}(U)$). Since $\underline{O}_S$ is coherent, the objects of $\underline{C}_0$ are pseudo-coherent (I 3.5), so an object of $\mathrm{D}(\mathrm{Qcoh}(U))$ (U = open subset of S) is pseudo-coherent (in the sense of (2.2.1)) precisely when it is so relative to $\underline{C}_0$ (I 2.14.1). It therefore suffices, by (1.1 c)), to show that conditions (i) and (ii) in the non-resp. case of (1.1) are satisfied. For (ii), this is trivial, since a quasi-coherent Module which is a quotient of a coherent Module is coherent, and the notion of coherence is local. For (i), we use (2.1.1 a)) and the fact that every quasi-coherent $\underline{O}_S$-Module is a filtered inductive limit of coherent $\underline{O}_S$-Modules (EGA I 9.4.9).

Admitting (3.5) and (3.7), which are independent of the present section, one easily deduces:

Corollary 2.2.2.1. Let S be a noetherian scheme (resp. noetherian and of finite dimension). The canonical functor

$$\mathrm{D}^b(\mathrm{Coh}(S)) \longrightarrow \mathrm{D}(S)$$
$$(\text{resp. } \mathrm{D}^-(\mathrm{Coh}(S)) \longrightarrow \mathrm{D}(S))$$

is fully faithful and has as essential image $\mathrm{D}^b(S)_{\mathrm{coh}}$ (resp. $\mathrm{D}^-(S)_{\mathrm{coh}}$) (I 2.15).

Proposition 2.2.3 (cf. EGA II 4.5.2 and 4.5.5). Let S be a quasi-compact and quasi-separated scheme, and let $(L_i)_{i\in I}$ be a family of invertible sheaves on S. For every $\underline{O}_S$-Module F, and every pair $(i,n) \in I \times \mathbb{Z}$, we denote by $F_{(i,n)}$ the sub-$\underline{O}_S$-Module of $F \otimes L_i^{\otimes n}$ generated by the sections of $F \otimes L_i^{\otimes n}$ over S. The following conditions are equivalent:

(i) [print 169]The open sets S_f, for $f \in \Gamma(S, L_i^{\otimes n})$, $i \in I$ and $n \in \mathbb{N}$, form a basis for the topology of S.

(i′) Those of the S_f which are affine cover S.

(ii) For every quasi-coherent $\underline{O}_S$-Module F of finite type, there exists a family $(r_i)_{i\in I}$ (resp. $(n_i)_{i\in I}$) of integers ≥ 0 (resp. > 0), and an epimorphism

$$\coprod_{i\in I} \underline{O}_S^{r_i} \otimes L_i^{\otimes -n_i} \longrightarrow F.$$

(iii) For every $F \in \mathrm{ob}\,\mathrm{Qcoh}(S)$, we have

$$F = \sum_{(i,n)\in I\times\mathbb{N}^+} F_{(i,n)} \otimes L_i^{\otimes -n}$$

(sum (not direct) of subsheaves of F).

(*) This ancillary result will not be used in the remainder of the Seminar: the reader is advised to skip it on a first reading.

(iii′) The same condition as (iii), but with F a quasi-coherent Ideal of $\underline{O}_S$.[124]

Proof. It consists of a simple paraphrase of (loc. cit.); we give it briefly for the reader's convenience.

(i)⇒(i′): For $x \in S$, choose an affine open subset U containing x. An open subset S_f such that $x \in S_f \subset U$, with $f \in \Gamma(S, L_i^{\otimes n})$ $(n \geq 0)$, is necessarily affine, as follows from

Lemma 2.2.3.1. (EGA II 5.5.8). Let X be a scheme, L an invertible $\underline{O}_X$-Module, $f \in \Gamma(X, L)$[125]. The canonical immersion $i : X_f \to X$ is affine.

Proof. This is trivial: the question is local on X, so we may assume that X is affine and $L \simeq \underline{O}_X$, and we are done.

(i′)⇒(ii): Take a finite covering of S by affine open subsets $S_{f_{ij}}$, where $f_{ij} \in \Gamma(S, L_i^{\otimes m_{ij}})$; put $S_i = \bigcup_j S_{f_{ij}}$. Replacing the f_{ij} by suitable powers, we may assume that the m_{ij} corresponding to the same index i are equal to an integer m_i. The sheaf [print 170]$F \mid S_{f_{ij}}$ is generated by finitely many sections h_{ijk} over $S_{f_{ij}}$. There exists (EGA I 9.3.1 and IV 1.7.5) an integer $q_i \geq 0$ such that $h_{ijk} \otimes f_{ij}^{\otimes q_i}$ extends to a section of $F \otimes L_i^{\otimes q_i m_i}$ over S. In other words, $F \otimes L_i^{\otimes q_i m_i}$ is generated over S_i by finitely many of its sections over S. Proceeding as in the proof of (EGA II 4.5.5), we deduce that there exists an integer n_i, which we may take to be > 0, such that, for every $n \geq n_i$, $F \otimes L_i^{\otimes n}$ is generated over S_i by finitely many of its sections over S. In particular, there exist an integer $r_i \geq 0$ and a morphism

$$\underline{O}_S^{r_i} \otimes L_i^{\otimes -n_i} \longrightarrow F$$

which induces an epimorphism over S_i. The implication (i′)⇒(ii) follows immediately.

(ii)⇒(iii): Since every quasi-coherent sheaf on S is an inductive limit of its quasi-coherent subsheaves of finite type (EGA I 9.4.9 and IV 1.7.7), we may restrict ourselves to proving (iii) for F of finite type. There then exist, for $i \in I$, morphisms

$$u_i : \underline{O}_S^{r_i} \otimes L_i^{\otimes -n_i} \longrightarrow F$$

such that $\sum u_i$ is an epimorphism. Now we have

$$\begin{aligned} \operatorname{Im}(u_i) &= \operatorname{Im}\big(u_i \otimes \operatorname{Id}(L_i^{\otimes n_i}) \otimes \operatorname{Id}(L_i^{\otimes -n_i})\big) \\ &= \operatorname{Im}\big(u_i \otimes \operatorname{Id}(L_i^{\otimes n_i})\big) \otimes L_i^{\otimes -n_i} \end{aligned}$$

But $\operatorname{Im}(u_i \otimes \operatorname{Id}(L_i^{\otimes n_i}))$ is a subsheaf of $F_{(i,n_i)}$, whence the assertion.

(iii)⇒(iii′): This is trivial.

(iii′)⇒(i): Let $x \in S$, U be an open subset containing x, and F a [print 171]quasi-coherent Ideal defining $S - U$. By (iii′), there exist $(i, n) \in I \times \mathbb{N}^+$ and a section f of $F \otimes L_i^{\otimes n}$ over S such that $f(x) \neq 0$. We have $x \in S_f \subset U$, which proves (i).

Definition 2.2.4. Let S be a quasi-compact and quasi-separated scheme, and let $(L_i)_{i \in I}$ be a family of invertible sheaves on S. We say that the L_i form an ample family(*) when the equivalent conditions of (2.2.3) hold. (In particular, a family consisting of a single sheaf L is ample precisely when L is ample in the usual sense (EGA II 4.5.3)).

Definition 2.2.5. A scheme S is said to be divisorial when S is quasi-compact and quasi-separated and there exists an ample family (2.2.4) of invertible $\underline{O}_S$-Modules.

(*) This notion, in form (i′), was introduced by M. BORELLI and S. L. KLEIMAN; the terminology "ample family" is proposed by the editor.

Lemma 2.2.6. Let X be a locally noetherian scheme. Let U be an open subset of X such that the inclusion $U \to X$ is affine (for example U affine and X separated). Then $X - U$ is purely of codimension 0 or 1 (i.e. at every maximal point x of $X - U$ we have $\dim(\underline{O}_{X,x}) = 0$ or 1), hence purely of codimension 1 when U contains the maximal points of X.

Proof. Put $Y = X - U$. Let x be a maximal point of Y, and make the base change $X' = \mathrm{Spec}(\underline{O}_{X,x}) \to X$: we obtain a cartesian diagram:

$$\begin{array}{ccccc} U' & \longrightarrow & X' & \longleftarrow & Y' \\ \downarrow & & \downarrow & & \downarrow \\ U & \longrightarrow & X & \longleftarrow & Y \end{array},$$

where $Y' = X' - U'$[126]. Since x is a maximal point of Y, Y' is precisely the closed point of X'. On the other hand, since X' is affine, U' is an affine open subset of X'. Let us show that the dimension n of X' is equal to 0 or 1. Indeed, we know (SGA 2 V 5.7) that $H^n_{Y'}(X', \underline{O}_{X'}) \neq 0$. But, for $n \geq 2$, we have

$$H^n_{Y'}(X', \underline{O}_{X'}) \simeq H^{n-1}(U', \underline{O}_{U'}),$$

whereas

$$H^{n-1}(U', \underline{O}_{U'}) = 0,$$

[print 172]since U' is affine and $n - 1 \geq 1$, whence a contradiction.

Remark 2.2.6.1. Without the separation hypothesis on X, (2.2.6) for an affine open subset U of X would be false, as shown by the example of the affine plane with the origin doubled (EGA I 5.5.11).

Proposition 2.2.7. Let S be a noetherian, separated, locally factorial scheme (i.e. one whose local rings are factorial). Then S is divisorial (2.2.5).

In particular (EGA IV 21.11.1):

Corollary 2.2.7.1. Every noetherian regular separated scheme is divisorial.

Proof of 2.2.7. We may assume that S is integral. Let $(S_i)_{i \in I}$ be a finite covering of S by nonempty affine open subsets. By (2.2.6), $S - S_i$ is purely of codimension 1, hence, by (EGA IV 21.7.4), is the support of a positive divisor. In other words, for each $i \in I$, there exist an invertible sheaf L_i and a section $f_i \in \Gamma(S, L_i)$ such that $S_i = S_{f_i}$. The family (L_i) is then ample (2.2.4) by (2.2.3 (i′)). This completes the proof.

Remark 2.2.7.2. As in (2.2.6), the separation hypothesis on S is essential. Indeed, again take for S the affine plane with the origin doubled. Recall that S is obtained by gluing two copies of the affine plane $\underline{\underline{E}}^2_k$ (k, a field) by the identity on the complement U of the origin O; we therefore have a canonical projection

$$p : S \longrightarrow \underline{\underline{E}}^2_k,$$

which induces an isomorphism over U, while $p^{-1}(O)$ consists of two points O_1 and O_2. An invertible sheaf on S is obtained by gluing two invertible sheaves on $\underline{\underline{E}}^2_k$ by an isomorphism over U; since every invertible sheaf on $\underline{\underline{E}}^2_k$ is trivial, an invertible sheaf on S is therefore defined by an element $g \in \Gamma(U, \underline{O}^*_U)$; but g is necessarily a constant, since the restriction $\Gamma(\underline{\underline{E}}^2_k, \underline{O}_{\underline{\underline{E}}^2_k}) \to \Gamma(U, \underline{O}_U)$ is an isomorphism, O being of depth 2 in the plane; thus every [print 173]invertible sheaf on S is trivial. On the other hand, it is immediate that p induces an isomorphism

$$p^* : \Gamma(\mathbb{E}^2_k, \underline{O}_{\mathbb{E}^2_k}) \longrightarrow \Gamma(S, \underline{O}_S).$$

Consequently, every global section of $\underline{O}_S$ which vanishes at O_1 also vanishes at O_2, so the S_f, for $f \in \Gamma(S, \underline{O}_S)$, do not form a basis for the topology of S.

Lemma 2.2.8. Let S be a divisorial scheme (2.2.5), $(L_i)_{i\in I}$ an ample family of invertible $\underline{O}_S$-Modules. Denote by $\underline{L}(S)$ the strictly full additive subcategory of $\mathrm{Qcoh}(S)$ (2.2.1) generated by the $L_i^{\otimes -n}$ for $(i,n)\in I\times\mathbb{N}^+$. Let $E\in \mathrm{ob}\,\mathrm{D}(\mathrm{Qcoh}(S))$.

a) The categories $\underline{L}(S)$ and $\mathrm{Loclib}(S)$ (2.0) are quasi-liftable in $\mathrm{Qcoh}(S)$ (I 1.2) and quasi-stable under kernels of epimorphisms (I 1.2).

b) Let $n\in\mathbb{Z}$. For E to be n-pseudo-coherent (resp. pseudo-coherent) (2.2.1), it is necessary and sufficient that E be n-pseudo-coherent (resp. pseudo-coherent) relative to $\underline{L}(S)$ or $\mathrm{Loclib}(S)$ (I 2.3). The canonical functors

$$\mathrm{K}^-(\underline{L}(S))/\mathrm{K}^{-,\varnothing}(\underline{L}(S))\longrightarrow \mathrm{D}(\mathrm{Qcoh}(S))$$

and

$$\mathrm{K}^-(\mathrm{Loclib}(S))/\mathrm{K}^{-,\varnothing}(\mathrm{Loclib}(S))\longrightarrow \mathrm{D}(\mathrm{Qcoh}(S))$$

are fully faithful and have as essential image $\mathrm{D}^-(\mathrm{Qcoh}(S))_{\mathrm{coh}}$ (2.2.1).

c) Let $[m,n]$ be an interval in $\mathbb{Z}$. For E to have perfect amplitude contained in $[m,n]$ (2.2.1), it is necessary and sufficient that E be isomorphic, in $\mathrm{D}(\mathrm{Qcoh}(S))$, to a complex F such that $F^i=0$ for $i\notin[m,n]$, $F^i\in\mathrm{ob}\,\underline{L}(S)$ for $i>m$, with F^m locally free of finite type. The canonical functor

$$\mathrm{K}^b(\mathrm{Loclib}(S))/\mathrm{K}^{b,\varnothing}(\mathrm{Loclib}(S))\longrightarrow \mathrm{D}(\mathrm{Qcoh}(S))$$

[print 174]is fully faithful and has as essential image $\mathrm{D}(\mathrm{Qcoh}(S))_{\mathrm{parf}}$ (2.2.1).

Proof. Let $F\to G$ be an epimorphism in $\mathrm{Qcoh}(S)$, where G is of finite type. Since F is an inductive limit of its quasi-coherent subsheaves of finite type (EGA I 9.4.9), there exists, by (2.1.1 a)), a quasi-coherent subsheaf F' of F of finite type such that the composite $F'\to F\to G$ is an epimorphism. By (2.2.3 (ii)), there exists an epimorphism $E\to F'$, with $E\in\mathrm{ob}\,\underline{L}(S)$, so the composite $E\to F'\to F$, composed with $F\to G$, is an epimorphism, which proves that the category of quasi-coherent $\underline{O}_S$-Modules of finite type is quasi-liftable in $\mathrm{Qcoh}(S)$; a fortiori, the same holds for the category $\mathrm{Loclib}(S)$ or $\underline{L}(S)$. This proves the first assertion of a). For the second, we apply (1.1), taking for $\underline{S}$ the Zariski site of S, and for $\underline{C}$ (resp. $\underline{C}_0$) the fibered category $U\mapsto\mathrm{Qcoh}(U)$ (resp. $\underline{L}(U)$) (U denotes a variable open subset of S, and $\underline{L}(U)$ the strictly full additive subcategory of $\mathrm{Qcoh}(U)$ generated by the $(L_i^{\otimes -n}\,|\,U)$). Since every object of $\underline{L}$ is pseudo-coherent, the notions of n-pseudo-coherence (resp. pseudo-coherence) (2.2.1) in the fibered category $\mathrm{D}(\mathrm{Qcoh})$ coincide with the same notions relative to $\underline{L}$ (I 2.14). Condition (ii) (in the non-resp. case) of (1.1) being satisfied by (2.2.3 (ii)), we deduce, by (1.1 a)), the second part of a) concerning $\underline{L}(S)$; the assertion concerning $\mathrm{Loclib}(S)$ is proved similarly. Assertions b) follow immediately from (1.1 b) and c)), taking into account the fact that, when $E\in\mathrm{ob}\,\mathrm{D}(\mathrm{Qcoh}(S))$ is n-pseudo-coherent, E is an object of $\mathrm{D}^-(\mathrm{Qcoh}(S))$, since S is quasi-compact. An object of the fibered category Qcoh is locally in $\underline{L}$ precisely when it is locally free of finite type: the category $\underline{L}$ is therefore locally liftable in Qcoh and locally stable under kernels of epimorphisms. Consequently, assertions c) follow[127] from (1.2) applied with $\underline{C}=\mathrm{Qcoh}$, $\underline{C}_0=\underline{L}$, $\underline{C}_{1S}=\mathrm{Loclib}(S)$. This completes the proof.[128]

Proposition 2.2.9. The hypotheses and notation are those of (2.2.8), but we further assume that S is separated or noetherian.

a) The canonical functors

$$\mathrm{K}^{-,b}(\underline{L}(S))/\mathrm{K}^{-,\varnothing}(\underline{L}(S))\longrightarrow \mathrm{D}(S)$$

and

$$\mathrm{K}^{-,b}(\mathrm{Loclib}(S))/\mathrm{K}^{-,\varnothing}(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}(S)$$

[print 175]are fully faithful and have as essential image $\mathrm{D}^b(S)_{\mathrm{coh}}$ (I 2.15). Moreover, when S has "finite cohomological dimension" in the sense of (3.7)([1]) (for example, when the underlying space of S is noetherian and of finite Krull dimension), the functors

$$\mathrm{K}^{-}(\underline{L}(S))/\mathrm{K}^{-,\varnothing}(\underline{L}(S)) \longrightarrow \mathrm{K}^{-}(\mathrm{Loclib}(S))/\mathrm{K}^{-,\varnothing}(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}^{-}(S)_{\mathrm{coh}}$$

are equivalences of categories.

b) The canonical functor

$$\mathrm{K}^{b}(\mathrm{Loclib}(S))/\mathrm{K}^{b,\varnothing}(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}(S)$$

is fully faithful and has as essential image $\mathrm{D}(S)_{\mathrm{parf}}$ (I 4.9). For $E \in \mathrm{ob}\,\mathrm{D}(S)$ to have perfect amplitude contained in $[m,n]$ (I 4.8), it is necessary and sufficient that E be isomorphic, in $\mathrm{D}(S)$, to a complex F such that $F^i = 0$ for $i \notin [m,n]$, $F^i \in \mathrm{ob}\,\underline{L}(S)$[129] for $i > m$, with F^m locally free of finite type.

Proof. In view of (3.5) and (3.7)([1]), this is an immediate corollary of (2.2.8): the only point to verify, which follows trivially from (3.5) (resp. (3.7)), is that, for an object E of $\mathrm{D}^b(\mathrm{Qcoh}(S))$ (resp. $\mathrm{D}(\mathrm{Qcoh}(S))$ when S has finite cohomological dimension in the sense of (3.7)), the notions of pseudo-coherence and perfection of (2.2.1) coincide with the usual notions (I 2.15 and 4.9) when E is regarded as an object of $\mathrm{D}(S)$.

Remark 2.2.10. a) When S is an affine scheme with ring A, the family consisting of the sheaf $\underline{O}_S$ is ample. For this family, we have $\underline{L}(S) = \mathrm{Lib}(S)$. It is also clear that the categories $\mathrm{K}^{-,\varnothing}(\mathrm{Lib}(S))$ and $\mathrm{K}^{-,\varnothing}(\mathrm{Loclib}(S))$ are reduced to zero objects. We therefore conclude from (2.2.9) that, writing $\mathrm{Mod}(A)$ for the category of A-modules and $\mathrm{D}(A) = \mathrm{D}(\mathrm{Mod}(A))$, the functor $\underline{\underline{R}}\Gamma_S : \mathrm{D}^+(S) \longrightarrow \mathrm{D}^+(A)$ induces a fully faithful functor $\mathrm{D}^b(S)_{\mathrm{coh}} \longrightarrow \mathrm{D}(A)$ (resp. $\mathrm{D}(S)_{\mathrm{parf}} \longrightarrow \mathrm{D}(A)$) whose essential image is [print 176]the triangulated subcategory $\mathrm{D}^b(A)_{\mathrm{coh}}$ (resp. $\mathrm{D}(A)_{\mathrm{parf}}$) formed by complexes with bounded cohomology and pseudo-coherent (resp. perfect) relative to the subcategory $\mathrm{Lib}(A)$ (resp. $\mathrm{Loclib}(A)$) of free (resp. projective) modules of finite type. We have commutative diagrams of equivalences of categories

$$\begin{array}{ccccc}
\mathrm{K}^{-,b}(\mathrm{Lib}(S)) & \longrightarrow & \mathrm{K}^{-,b}(\mathrm{Loclib}(S)) & \longrightarrow & \mathrm{D}^b(S)_{\mathrm{coh}} \\
\downarrow{\scriptstyle \Gamma_S} & & \downarrow{\scriptstyle \Gamma_S} & & \downarrow{\scriptstyle \mathbb{R}\Gamma_S} \\
\mathrm{K}^{-,b}(\mathrm{Lib}(A)) & \longrightarrow & \mathrm{K}^{-,b}(\mathrm{Loclib}(A)) & \longrightarrow & \mathrm{D}^b(A)_{\mathrm{coh}},
\end{array}$$

$$\begin{array}{ccc}
\mathrm{K}^{b}(\mathrm{Loclib}(S)) & \longrightarrow & \mathrm{D}(S)_{\mathrm{parf}} \\
\downarrow{\scriptstyle \Gamma_S} & & \downarrow{\scriptstyle \mathbb{R}\Gamma_S} \\
\mathrm{K}^{b}(\mathrm{Loclib}(A)) & \longrightarrow & \mathrm{D}(A)_{\mathrm{parf}},
\end{array}$$

(where, in the upper diagram, one may replace b by $-$ when S has finite cohomological dimension in the sense of (3.7)).

b) Let S be a scheme. It follows from a) that a complex of $\underline{O}_S$-Modules which is pseudo-coherent and has bounded cohomology is locally isomorphic (in $\mathrm{D}(S)$) to a strictly pseudo-coherent complex, which was by no means evident a priori (cf. (I 2.3.1)).

([1]) Section 3 is independent of the rest of the exposé.

2.3. Soft sheaves

2.3.1. Recall some definitions and results from [T.F]. Let X be a paracompact space. A sheaf of sets F on X is said to be soft when, for every closed subset Y of X, every section of F over Y extends to a section of F over X. Soft abelian sheaves are acyclic, i.e. when F is soft, we have $H^i(X, F) = 0$ for $i > 0$ (T.F. II 4.4.3).

Let A be a sheaf of rings on X. When F is a soft A-Module, F is generated by its sections (cf. 2.1.1): indeed, the homomorphism $A_X^{(I)} \to F$ defined by $I = H^0(X, F)$ is clearly an epimorphism. [print 177]On the other hand, when A is soft, every A-Module is soft (T.F II 3.7.1). As examples of soft sheaves of rings, we mention the sheaf of continuous functions (real or complex) on a paracompact space, and the sheaf of C^r functions on a manifold of class C^r.

Proposition 2.3.2. Let S be a compact space, equipped with a soft sheaf of rings A. We write $\mathrm{Mod}(S)$ for the category of left A-Modules, $\mathrm{D}(S) = \mathrm{D}(\mathrm{Mod}(S))$.

a) The category $\mathrm{Lib}(S)$ (resp. $\mathrm{Loclib}(S)$) (2.0) is liftable in $\mathrm{Mod}(S)$ and quasi-stable (resp. stable) under kernels of epimorphisms (I 1.2)([1]).

b) Let $E \in \mathrm{ob}\,\mathrm{D}(S)$ and $n \in \mathbb{Z}$. For E to be n-pseudo-coherent (resp. pseudo-coherent), it is necessary and sufficient that E be so relative to $\mathrm{Lib}(S)$ or to $\mathrm{Loclib}(S)$. The canonical functors

$$\mathrm{K}^-(\mathrm{Lib}(S)) \longrightarrow \mathrm{D}(S)$$

and

$$\mathrm{K}^-(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}(S)$$

are fully faithful and have as essential image $\mathrm{D}^-(S)_{\mathrm{coh}}$ (I 2.15).

c) Let $E \in \mathrm{ob}\,\mathrm{D}(S)$, and $[m, n]$ be an interval in $\mathbb{Z}$. For E to have perfect amplitude contained in $[m, n]$ (I 4.8)[130], it is necessary and sufficient that there exist a quasi-isomorphism $F \to E$, where F is a complex such that $F^i = 0$ for $i \notin [m, n]$, $F^i \in \mathrm{ob}\,\mathrm{Lib}(S)$ for $i > m$ and $F^m \in \mathrm{ob}\,\mathrm{Loclib}(S)$. The canonical functor

$$\mathrm{K}^b(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}(S)$$

is fully faithful and has as essential image $\mathrm{D}(S)_{\mathrm{parf}}$ (I 4.9)[131].

d) Denote by $\mathrm{Mod}(A(S))$ the category of left $A(S)$-modules, and by $\mathrm{D}(A(S))$ the corresponding derived category. The functor

$$H^0(S, .) : \mathrm{Mod}(S) \longrightarrow \mathrm{Mod}(A(S))$$

is exact and an equivalence of categories. Moreover, it defines [print 178]equivalences of categories:

$$\mathrm{D}(S) \longrightarrow \mathrm{D}(A(S))$$
$$\mathrm{D}^-(S)_{\mathrm{coh}} \longrightarrow \mathrm{D}^-(A(S))_{\mathrm{coh}}$$
$$\mathrm{D}(S)_{\mathrm{parf}} \longrightarrow \mathrm{D}(A(S))_{\mathrm{parf}},$$

the notions of pseudo-coherence (resp. perfection) in $\mathrm{D}(A(S))$ being understood relative to the category of free (resp. projective) modules of finite type. It also defines an equivalence from the full subcategory of $\mathrm{D}(S)$ formed by n-pseudo-coherent objects (resp. objects of perfect amplitude contained in $[m, n]$) onto the analogous full subcategory of $\mathrm{D}(A(S))$.

([1]) This assertion holds more generally for S paracompact.

Proof. The first assertion of a) follows from (2.1.2 a)) applied by taking for $\underline{S}$ the topos of sheaves of sets on S, and $\underline{C}'_S = \mathrm{Mod}(S)$ (in view of (2.3.1)). The second follows from (1.1 a)) (resp. (1.2 a))) with $\underline{C} = \mathrm{Mod}$ (the fibered category of A-Modules on variable open subsets), $\underline{C}_0 = \mathrm{Lib}$ (resp. Loclib), condition (ii) of (1.1) being verified by (2.3.1) and (2.1.1.b)). For assertion b), we apply (1.1 b),c)) (with $\underline{C}$ and $\underline{C}_0$ as before). Assertion c) follows from (1.2 b),c),d))[132], where $\underline{C} = \mathrm{Mod}$ and $\underline{C}_0 = \mathrm{Lib}$. The first assertion of d) follows, by (2.3.1), from (2.1.2 b) (ii)), where we take $\underline{C}'_S = \underline{C}_S = \mathrm{Mod}(S)$. The other assertions of d) follow from the preceding one, in view of b) and c), and the fact that the functor Γ_S sends free Modules (resp. direct factors of free Modules) of finite type[133] to free modules (resp. projective modules of finite type).

2.4. Stein spaces

Definition 2.4.1. Let $\underline{S}$ be a locally ringed topos (I 2.15.1 (ii)), with final object S. We say that $\underline{S}$ is Stein when the structure sheaf $\underline{O}_S$ is coherent (I 3.1) and the following properties hold:

"theorem A" : every coherent $\underline{O}_S$-Module (I 3.1) is generated by its sections,

"theorem B" : for every coherent $\underline{O}_S$-Module F, we have $H^i(S, F) = 0$ for $i > 0$.

[print 179]**Remark 2.4.1.1.** When $\underline{S}$ has "enough closed points"([1]) s such that the Ideal $\underline{m}_s$ defined by s([2]) is coherent, the relation "$H^1(S, F) = 0$ for every coherent sheaf F" implies theorem A. Indeed, let F be a coherent $\underline{O}_S$-Module and s a closed point of S. We have the exact sequence

$$0 \longrightarrow \underline{m}_s F \longrightarrow F \longrightarrow F/\underline{m}_s F \longrightarrow 0,$$

where $F/\underline{m}_s F$ is a coherent sheaf concentrated at s. Applying the functor $H^0(S, .)$ and writing the cohomology exact sequence, we find that $H^0(S, F) \longrightarrow H^0(S, F/\underline{m}_s F)$ is an epimorphism. Since $H^0(S, F/\underline{m}_s F) \simeq F_s/m_s F_s$, where m_s is the maximal ideal of $\underline{O}_{S,s}$, we conclude by Nakayama.

The editor does not know whether the preceding implication is true in general (when one assumes only that $\underline{O}_S$ is a coherent sheaf of local rings).

The editor also does not know whether the relation "$H^1(S, F) = 0$ for every coherent sheaf F" implies theorem B.

Examples 2.4.2. A complex analytic Stein space [3], a compact convex subset of $\mathbb{C}^n$ equipped with the sheaf induced by the sheaf of holomorphic functions [3], the spectrum of a noetherian ring, and an affine rigid analytic space [13] define Stein topoi.

Lemma 2.4.3. Let $\underline{S}$ be a Stein topos (2.4.1). We assume that the final object S is quasi-compact (2.1.1), and we denote by $\mathrm{Coh}(S)$ the abelian category of coherent $\underline{O}_S$-Modules, i.e. of finite presentation. Then:

a) The category $\mathrm{Lib}(S)$ (resp. $\mathrm{Loclib}(S)$) (2.0) is liftable in $\mathrm{Coh}(S)$ and quasi-stable (resp. stable) under kernels of epimorphisms (I 1.2).

b) The canonical functors

$$\mathrm{K}^-(\mathrm{Lib}(S)) \longrightarrow \mathrm{K}^-(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}^-(\mathrm{Coh}(S))$$

are equivalences of categories.

([1]) i.e. enough fiber functors (SGA 4 IV 6.4.1) of the type $i_s : s \to \underline{S}$, where i_s is the inclusion of a closed subtopos of $\underline{S}$.

([2]) i.e. formed by sections of $\underline{O}_S$ whose fiber at s belongs to the maximal ideal m_s of $\underline{O}_{S,s}$.

c) [print 180]Put $A = \Gamma(S, \underline{O}_S)$, and denote by $\mathrm{Mod}(A)$ the category of A-modules, and by $\mathrm{D}(A)$ the corresponding derived category. The functor

$$\Gamma_S : \mathrm{Coh}(S) \longrightarrow \mathrm{Mod}(A)$$

is exact, fully faithful and has as essential image the strictly full additive subcategory of $\mathrm{Mod}(A)$ formed by A-modules of finite presentation. It induces a fully faithful functor

$$\mathrm{D}^-(\mathrm{Coh}(S)) \longrightarrow \mathrm{D}(A),$$

whose essential image is the full subcategory $\mathrm{D}^-(A)_{\mathrm{coh}}$ formed by pseudo-coherent complexes (relative to the subcategory of free modules of finite type).

Proof. For the first assertion of a), we apply (2.1.2 a)) with $\underline{C}'_S = \mathrm{Coh}(S)$, the hypothesis of (2.1.2) being satisfied by theorem B: with the notation of (2.1.2 a)), when E is coherent and G locally free of finite type, $\underline{\mathrm{Hom}}(G, E)$ is coherent, so the category $\mathrm{Loclib}(S)$ (and a fortiori $\mathrm{Lib}(S)$) is liftable in $\mathrm{Coh}(S)$. The second assertion of a) follows from (1.1 a)) (resp.(1.2 a))) applied with $\underline{C}_0 = \mathrm{Lib}$ (resp. Loclib)([1]), the resp. condition (ii) of (1.1) being satisfied by theorem A and (2.1.1 b)). Assertion b) follows from (1.1 c)) applied successively with $\underline{C}_0 = \mathrm{Lib}$ and $\underline{C}_0 = \mathrm{Loclib}$. The first assertion of c) follows from (2.1.2 b)(ii)) applied with $\underline{C}'_S = \mathrm{Coh}(S)$. To prove that the functor $\Gamma_S : \mathrm{D}^-(\mathrm{Coh}(S)) \to \mathrm{D}(A)$ is fully faithful, it suffices, in view of b), to prove that the composite

$$\mathrm{K}^-(\mathrm{Lib}(S)) \longrightarrow \mathrm{D}^-(\mathrm{Coh}(S)) \xrightarrow{\Gamma_S} \mathrm{D}(A)$$

is so. But this factors as

$$\mathrm{K}^-(\mathrm{Lib}(S)) \xrightarrow{(*)} \mathrm{K}^-(\mathrm{Lib}(A)) \xrightarrow{(**)} \mathrm{D}(A),$$

[print 181]where $\mathrm{Lib}(A)$ denotes the category of free A-modules of finite type and $(*)$ is induced by Γ_S. It is trivial that $(*)$ is an equivalence of categories. On the other hand, (I 2.7 b)) applied with $\underline{C} = \mathrm{Mod}(A)$, $\underline{C}_0 = \mathrm{Lib}(A)$ (and taking account of the fact that $\mathrm{K}^{-,\varnothing}(\mathrm{Lib}(A)) = 0$) shows that $(**)$ is fully faithful and has as essential image $\mathrm{D}^-(A)_{\mathrm{coh}}$. This proves the last assertion of c) and completes the proof of (2.4.3).

Lemma 2.4.4. Under the hypotheses of (2.4.3), the canonical functor

$$\mathrm{D}^b(\mathrm{Coh}(S)) \longrightarrow \mathrm{D}(S)$$

is fully faithful and has as essential image $\mathrm{D}^b(S)_{\mathrm{coh}}$ (I 2.15).

Proof We apply (2.1.3) with $A = \mathrm{Coh}(S)$, $B = \mathrm{Mod}(S)$, $\phi : A \to B$ being the inclusion functor. Since this functor is fully faithful, it suffices to check, by (2.1.3 a)(ii)), that for fixed $M \in \mathrm{ob}\,\mathrm{Coh}(S)$ and $n > 0$ the functor $L \longmapsto \mathrm{Ext}^n_{\mathrm{Mod}(S)}(L, M)$ is coeffaceable in $L \in \mathrm{ob}\,\mathrm{Coh}(S)$. Now, by theorem A and (2.1.1 b)), every coherent $\underline{O}_S$-Module is a quotient of a free Module of finite type, and, when L is free of finite type, we have $\mathrm{Ext}^n_{\mathrm{Mod}(S)}(L, M) = 0$ by theorem B, whence our assertion. It follows that the functor $\mathrm{D}^b(\mathrm{Coh}(S)) \to \mathrm{D}(S)$ is fully faithful and has as essential image the full subcategory of $\mathrm{D}(S)$ formed by complexes with bounded and coherent cohomology, which is precisely $\mathrm{D}^b(S)_{\mathrm{coh}}$ since $\underline{O}_S$ is coherent (I 3.5. b)).

Proposition 2.4.5. With the hypotheses and notation of (2.4.3), we have:

([1]) Lib and Loclib are locally stable under kernels of epimorphisms by (I 2.15.1).

a) The canonical functors

$$\mathrm{K}^{-,b}(\mathrm{Lib}(S)) \longrightarrow \mathrm{K}^{-,b}(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}^b(S)_{\mathrm{coh}}$$

are equivalences of categories. Moreover, the functor $\underline{\underline{R}}\Gamma_S : \mathrm{D}^+(S) \to \mathrm{D}^+(A)$ induces a fully faithful functor

$$\underline{\underline{R}}\Gamma_S : \mathrm{D}^b(S)_{\mathrm{coh}} \longrightarrow \mathrm{D}^+(A)^{[134]},$$

[print 182]whose essential image is $\mathrm{D}^b(A)_{\mathrm{coh}}$ (the triangulated subcategory of $\mathrm{D}(A)$ formed by complexes with bounded cohomology and pseudo-coherent relative to the category of free modules of finite type).

b) Let $E \in \mathrm{ob}\,\mathrm{D}(S)$, and let $[m,n]$ be an interval of $\mathbb{Z}$. When E has perfect amplitude contained in $[m,n]$ (I 4.8), E is isomorphic to a complex F such that $F^i = 0$ for $i \notin [m,n]$[135], $F^i \in \mathrm{ob}\,\mathrm{Lib}(S)$ for $i > m$ and $F^m \in \mathrm{ob}\,\mathrm{Loclib}(S)$.

c) The canonical functor

$$\mathrm{K}^b(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}(S)_{\mathrm{parf}} \qquad \text{(I 4.9)}$$

is an equivalence of categories. Moreover, the functor $\underline{\underline{R}}\Gamma_S : \mathrm{D}^+(S) \to \mathrm{D}^+(A)$ induces a fully faithful functor

$$\underline{\underline{R}}\Gamma_S : \mathrm{D}(S)_{\mathrm{parf}} \longrightarrow \mathrm{D}^+(A),$$

whose essential image is $\mathrm{D}(A)_{\mathrm{parf}}$ (the triangulated subcategory of $\mathrm{D}(A)$ formed by complexes perfect relative to the category of projective modules of finite type).

Proof. a) The functors

$$\mathrm{K}^{-,b}(\mathrm{Lib}(S)) \longrightarrow \mathrm{K}^{-,b}(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}^b(\mathrm{Coh}(S)) \xrightarrow{\Gamma_S} \mathrm{D}^b(A)_{\mathrm{coh}}$$

are equivalences of categories by (2.4.3 b) and c)). Since the (fully faithful) functor $\Gamma_S : \mathrm{D}^b(\mathrm{Coh}(S)) \to \mathrm{D}(A)$ factors as

$$\mathrm{D}^b(\mathrm{Coh}(S)) \longrightarrow \mathrm{D}^b(S)_{\mathrm{coh}} \xrightarrow{\underline{\underline{R}}\Gamma_S} \mathrm{D}(A),$$

we conclude by (2.4.4).

b) Suppose that $\mathrm{parf.amp}(E) \subset [m,n]$. In particular, E is pseudo-coherent; by a), after replacing E by an isomorphic object, we may therefore assume that $E^i \in \mathrm{ob}\,\mathrm{Lib}(S)$ for $i > m$ and $E^i = 0$ for $i > n$. [print 183]After truncation, we may even assume that $E^i = 0$ for $i < m$. By (I 4.16) applied with $\underline{C} = \mathrm{Mod}$, $\underline{C}_0 = \mathrm{Lib}$, we then have $E^m \in \mathrm{ob}\,\mathrm{Loclib}(S)$[136], whence the assertion.

c) Hypothesis (1.1 (i))(resp. (1.1 (ii) resp.)) being satisfied for $\underline{C}_S = \mathrm{Coh}(S)$, $\underline{C}_0 = \mathrm{Lib}$ by (2.4.3 a)) (resp. (2.1.1 b)) and theorem A), we know from (1.2 c) and d)) that the functor

$$\mathrm{K}^b(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}^b(\mathrm{Coh}(S))$$

is fully faithful. Thus, by (2.4.4), the same holds for the functor

$$\mathrm{K}^b(\mathrm{Loclib}(S)) \longrightarrow \mathrm{D}(S),$$

whose essential image is $\mathrm{D}(S)_{\mathrm{parf}}$ by b). By an argument analogous to that in the proof of (2.4.3 c)), we see that the functor $\mathrm{K}^b(\mathrm{Loclib}(S)) \to \mathrm{D}(A)^{[137]}$ induced by Γ_S is fully faithful and has as essential image $\mathrm{D}(A)_{\mathrm{parf}}$[138], and we complete the proof as in a).

Remark 2.4.6. The editor does not know whether, in (2.4.4), one may replace D^b by D^-, and consequently whether (2.4.5 a)) remains true when one replaces $\mathrm{K}^{-,b}$ by K^- and D^b by D^-.

To finish, here is an interesting complement to (2.4.5). Let us first give a definition:

Definition 2.4.7. A complex pseudo-analytic space is a ringed space locally isomorphic to a ringed space of the form $(Y, \underline{O}_X|Y)$, where X is a complex analytic space and Y a closed subset of X. The structure sheaf of a complex pseudo-analytic space is therefore coherent. A complex pseudo-analytic space is said to be Stein when the corresponding topos is Stein (2.4.1).

[print 184]**Proposition 2.4.8.**([1]). Let S be a compact complex pseudo-analytic Stein space (2.4.7). We assume that, for every open subset U of S and every reduced closed subspace X of U, the normalization of X is locally connected([2]) (a condition satisfied, for example, when S is locally defined by real analytic equations or inequalities in a space $\underline{\underline{R}}^n$ [15]). Then the ring $A = \Gamma(S, \underline{O}_S)$ is a noetherian ring.

Proof. We have to show that, for an A-module M of finite presentation, the set of submodules (or, equivalently, the set of submodules of finite type) of M satisfies the maximal condition. Now, when N is a submodule of finite type of M, M/N is of finite presentation by (I 2.6 and 2.9), so N is of finite presentation by the equivalence (2.4.3 c)). Using again the equivalence (2.4.3 c)) between the category of coherent $\underline{O}_S$-Modules and the category of A-modules of finite presentation, and the compactness of S, we are reduced to proving the following lemma:

Lemma 2.4.8.1. Let S be a complex pseudo-analytic space satisfying the local connectedness condition of (2.4.8). Let F be a coherent $\underline{O}_S$-Module, (E_n) an increasing sequence of coherent sub-$\underline{O}_S$-Modules of F, s a point of S. Then there exists an open neighbourhood U of s such that the sequence $(E_n|U)$ is stationary.

Proof. Since the ring $\underline{O}_{S,s}$ is noetherian, there exists an integer n_0 such that $(E_n)_s = (E_{n_0})_s$ for $n \geq n_0$. Replacing the sequence (E_n) by the sequence (E_n/E_{n_0}) for $n \geq n_0$, we may therefore assume that $(E_n)_s = 0$ for every n. On the other hand, $F_s = M$ is an $\underline{O}_s$-module of finite type, so there exists (Bourbaki Alg. com. chap. IV §1 th.1) a finite filtration

$$M = M_0 \supset M_1 \supset \ldots \supset M_q = 0$$

such that $M_i/M_{i+1} \simeq \underline{O}_s/p_i$, where p_i is a prime ideal of $\underline{O}_s$. Since the category of $\underline{O}_s$-modules of finite type is equivalent to the category of germs at s of coherent $\underline{O}_S$-Modules, we may suppose, after [print 185]localizing in a neighbourhood of s, that the M_i (resp. p_i) are the fibers at s of coherent sub-$\underline{O}_S$-Modules F_i (resp. coherent Ideals $\underline{p}_i$) of F (resp. $\underline{O}_S$) such that

$$F = F_0 \supset F_1 \supset \ldots \supset F_q = 0$$

and $F_i/F_{i+1} \simeq \underline{O}_S/\underline{p}_i$. Proving that the sequence (E_n) is stationary in a neighbourhood of s is evidently equivalent to proving that the sequences induced on each graded piece of the filtration (F_i) of F are stationary. Thus we may assume that $F = \underline{O}_S$, and that $\underline{O}_{S,s}$ is an integral domain, the E_n forming an increasing sequence of Ideals of $\underline{O}_S$ such that $(E_n)_s = 0$. Since $\underline{O}_s$ is an integral domain, hence in particular reduced, Oka's theorem ([10] IV §3) further allows us to assume, after localizing at s, that S is reduced. Let us show that there exists an open neighbourhood U of s such that $E_n|U = 0$ for sufficiently large n. For this, consider the normalization $\widetilde{S}$ of S ([10] IV §4). We know (loc. cit.) that $\widetilde{S}$ is the analytic spectrum[140] ([10] II §3) of a finite $\underline{O}_S$-Algebra $\widetilde{\underline{O}}_S$ whose fiber at each $t \in S$ is the integral closure of $\underline{O}_{S,t}$ (cf. note ([2]) to (2.4.8)). We have $E_n \subset E_n \cdot \widetilde{\underline{O}}_S$, so it suffices to prove that $E_n \cdot \widetilde{\underline{O}}_S = 0$ for sufficiently large n. Now we have ([10] II §3 Prop.8)[141]

$$E_n \cdot \widetilde{\underline{O}}_S = p_*(\widetilde{E_n \cdot \widetilde{\underline{O}}_S}),$$

([1]) This result, together with its proof, was kindly communicated to the editor by A. Grothendieck; it was obtained independently by J. Frisch [5][139].

([2]) We shall admit that the notions of local Analytic Geometry in [10] extend to complex pseudo-analytic spaces; the reduction to the usual case is indeed immediate each time.

where $p : \widetilde{S} \longrightarrow S$ is the canonical projection, and $\widetilde{E_n \cdot \underline{O}_S}$ the Ideal of $\underline{O}_{\widetilde{S}}$ canonically defined by $E_n \cdot \underline{\widetilde{O}}_S$. It therefore suffices to show that there exists an open neighbourhood V of s in $\widetilde{S}$ such that $\widetilde{E_n \cdot \underline{O}_S}|V = 0$[142] for sufficiently large n, which reduces us to the case where S is normal, as we shall now assume. Since $(E_n)_s = 0$, the closed immersion $i_n : S_n \longrightarrow S$ defined by E_n is étale at s. But, since S is normal, the set of points where i_n is étale is both open and closed (see the lemma below), so its image in S is a subset of S that is both open and closed. Thus the restriction of E_n to the connected component of s in S is zero; this connected component is an open set, since S is locally connected, and we are done. It therefore remains to establish the following

[print 186]**Lemma 2.4.8.2.** (cf. EGA IV 18.10.3) Let Y be a normal pseudo-analytic space([1]), and $f : X \longrightarrow Y$ an unramified morphism. The set of points of X where f is étale is both open and closed.

Proof. It is immediate that the set of points where f is étale is open (since saying that f is étale at a point means that f is an isomorphism in a neighbourhood of that point ([8] VI)). It remains to show that the set of points where f is not étale is open. Since an unramified morphism is locally an immersion ([8] VI), we may assume that f is a closed immersion, defined by a coherent Ideal I. Let x be a point of X such that $I_x \neq 0$. Let $\underline{p}$ be a prime ideal of $\underline{O}_{Y,x}$ containing I_x. We have $\dim(\underline{O}_{Y,x})_{\underline{p}} > 0$, since otherwise we would have $\underline{p} = 0$, as $\underline{O}_{Y,x}$ is an integral domain. Thus X has codimension[143] > 0 at x, hence codimension > 0 in a neighbourhood of x ([10] III A Cor 1 to Lemma 3), and we are done.

3. Complements on quasi-coherent sheaves on schemes([2])

3.0. In this section, we fix a quasi-compact and quasi-separated scheme S. The canonical (fully faithful) inclusion functor

$$\phi : \mathrm{Qcoh}(S) \longrightarrow \mathrm{Mod}(S) \qquad ([3])$$

extends to a functor, still denoted by ϕ, from $\mathrm{D}(\mathrm{Qcoh}(S))$ to $\mathrm{D}(S) = \mathrm{D}(\mathrm{Mod}(S))$, which we propose to study([4]).

Lemma 3.1. The category $\mathrm{Qcoh}(S)$ is an abelian category satisfying (AB 5) and for which the quasi-coherent sheaves of finite type form a "small" family of generators. In particular (Tohoku 1.10.1), $\mathrm{Qcoh}(S)$ has enough injectives.

[print 187]Proof. The assertion concerning generators follows immediately from the fact that every quasi-coherent sheaf on S is the filtered inductive limit of its quasi-coherent subsheaves of finite type (EGA I 9.4.9 and IV 1.7.7). The other assertions are evident.

Remark 3.1.1. The editor does not know whether $\mathrm{Qcoh}(S)$ still has enough injectives when one assumes only that S is quasi-separated.

Lemma 3.2. The functor $\phi : \mathrm{Qcoh}(S) \longrightarrow \mathrm{Mod}(S)$ has a right adjoint

$$Q : \mathrm{Mod}(S) \longrightarrow \mathrm{Qcoh}(S),$$

(called the coherer).

Proof. When S is affine, the existence of Q is immediate: take $Q = \widetilde{\Gamma}_S$ (EGA $\mathrm{Err_I}$ 1.8.7). Now let $f : U \longrightarrow S$ be an affine open subset of S; it is clear that, for $E \in \mathrm{ob}\,\mathrm{Qcoh}(S)$ and

([1]) It suffices that the local rings of Y be integral domains.
([2]) This section is essentially due to J.L. Verdier.
([3]) $\mathrm{Mod}(S)$ (resp. $\mathrm{Qcoh}(S)$) denotes the category of $\underline{O}_S$-Modules (resp. quasi-coherent $\underline{O}_S$-Modules).
([4]) We shall omit ϕ in formulas where there is no risk of confusion.

$F \in \mathrm{ob}\,\mathrm{Mod}(U)$, we have a canonical isomorphism

$$\mathrm{Hom}(E, f_*(F)) \simeq \mathrm{Hom}(E, f_*Q(F)). \tag{1}$$

Then let $(f_i : U_i \longrightarrow S)$ be a finite covering of S by affine open subsets, and for each (i,j) let $(U_{ijk} \longrightarrow U_{ij})$ be a finite covering of $U_{ij} = U_i \cap U_j$ by affine open subsets (U_{ij} is quasi-compact since S is quasi-separated). Let $F \in \mathrm{ob}\,\mathrm{Mod}(S)$. We have an exact sequence

$$0 \longrightarrow F \longrightarrow \coprod_i f_{i*}(F|U_i) \longrightarrow \coprod_{ijk} f_{ijk*}(F|U_{ijk}), \tag{2}$$

(where f_{ijk} denotes the inclusion of U_{ijk} into S). Note that, for an inclusion $g : V \longrightarrow U$ of affine open subsets of S, we have an evident natural arrow $Q(F|U) \longrightarrow g_*Q(F|V)$. Consequently, (2) gives an arrow

$$\coprod_i f_{i*}Q(F|U_i) \longrightarrow \coprod_{ijk} f_{ijk*}Q(F|U_{ijk}), \tag{3}$$

whose kernel we shall denote by QF. Let $E \in \mathrm{ob}\,\mathrm{Qcoh}(S)$. Applying the functor $\mathrm{Hom}(E, .)$ to the sequence (2), we obtain, in view of (1), an [print 188]isomorphism functorial in E:

$$\mathrm{Hom}(E, F) \simeq \mathrm{Hom}(E, QF).$$

This proves the existence of Q; uniqueness is evident, since it is an adjoint.

Lemma 3.3.

a) The coherer is an additive functor taking injectives to injectives.

b) Write

$$\underline{\underline{R}}Q : \mathrm{D}^+(S) \longrightarrow \mathrm{D}^+(\mathrm{Qcoh}(S))$$

for the right derived functor of the coherer. For $E \in \mathrm{ob}\,\mathrm{D}(\mathrm{Qcoh}(S))$ and $F \in \mathrm{ob}\,\mathrm{D}^+(S)$, there are canonical functorial isomorphisms

$$\underline{\underline{R}}\,\mathrm{Hom}_{\mathrm{Mod}}(E, F) \simeq \underline{\underline{R}}\,\mathrm{Hom}_{\mathrm{Qcoh}}(E, \underline{\underline{R}}Q(F)) \qquad ([1]) \tag{i}$$

$$\mathrm{Hom}_{\mathrm{Mod}}(E, F) \simeq \mathrm{Hom}_{\mathrm{Qcoh}}(E, \underline{\underline{R}}Q(F)). \tag{ii}$$

Proof. Assertion a) is an immediate consequence of the definition. For b), it suffices to establish the isomorphism (i), since (ii) will follow by passing to cohomology. For this, we may assume that F has injective components; QF then has injective components by a), and the isomorphism (i) comes simply from the definition of Q.

Lemma 3.4. Let $f : X \longrightarrow Y$ be a morphism of quasi-compact and quasi-separated schemes. For $E \in \mathrm{ob}\,\mathrm{Mod}(X)$, there is a canonical functorial isomorphism

$$Qf_*(E) \simeq f_*Q(E).$$

[print 189]Proof. Since f is quasi-compact and quasi-separated (EGA IV 1.2.3 and 1.2.4), f_* takes quasi-coherent sheaves to quasi-coherent sheaves (III 1.4.10 and IV 1.7.21). The functors Qf_* and f_*Q from $\mathrm{Mod}(X)$ to $\mathrm{Qcoh}(Y)$ are both right adjoints to the functor $f^* : \mathrm{Qcoh}(Y) \longrightarrow \mathrm{Mod}(X)$, and hence canonically isomorphic.

([1]) The subscript Mod (resp. Qcoh) is an abbreviation for $\mathrm{D}(S)$ (resp. $\mathrm{D}(\mathrm{Qcoh}(S))$).

Proposition 3.5. When S is noetherian, or quasi-compact and separated, the functor

$$\phi : \mathrm{D}^+(\mathrm{Qcoh}(S)) \longrightarrow \mathrm{D}^+(S) \tag{3.0}$$

is fully faithful and has as essential image the full subcategory $\mathrm{D}^+(S)_{\mathrm{qcoh}}$ of $\mathrm{D}^+(S)$ formed by complexes with quasi-coherent cohomology.

Proof. We may restrict ourselves to proving the assertion in the case where S is separated, the case where S is noetherian being well known ([H] II 7.19). It is evidently equivalent to prove the following assertions:

(3.5.1) : for $E \in \mathrm{ob}\,\mathrm{D}^+(\mathrm{Qcoh}(S))$, the adjunction arrow

$$E \longrightarrow \underline{\underline{R}}Q(E)$$

(deduced from (3.3 b) (ii)) is an isomorphism[144];

(3.5.2) : for $E \in \mathrm{ob}\,\mathrm{D}^+(S)_{\mathrm{qcoh}}$, the adjunction arrow

$$\underline{\underline{R}}Q(E) \longrightarrow E$$

(deduced from (3.3 b)(ii)) is an isomorphism[145].

Proof of (3.5.1). By an immediate dévissage([1]), we are reduced to proving that quasi-coherent sheaves are acyclic for the coherer, (i.e. that, for $E \in \mathrm{ob}\,\mathrm{Qcoh}(S)$, we have $R^iQ(E) = 0$ for $i > 0$), and that, on the other hand, for $E \in \mathrm{ob}\,\mathrm{Qcoh}(S)$, the adjunction arrow $E \longrightarrow QE$ is an isomorphism. When S is affine, this follows from the formula $Q = \widetilde{\Gamma}_S$ (cf. proof of (3.2)). In the general case, cover S by finitely many affine open subsets U_i. Since S is separated, the inclusions $f_i : U_i \longrightarrow S$ are affine morphisms, [print 190]as are the inclusions $f_{i_0\ldots i_p} : U_{i_0\ldots i_p} = U_{i_0} \cap \ldots \cap U_{i_p} \longrightarrow S$. Let $E \in \mathrm{ob}\,\mathrm{Qcoh}(S)$, and consider the resolution of E defined by the covering (U_i) (T.F. II 5.2):

$$0 \longrightarrow E \longrightarrow \coprod f_{i*}f_i^*(E) \longrightarrow \ldots \longrightarrow \coprod f_{i_0\ldots i_p*}f_{i_0\ldots i_p}^*(E) \longrightarrow . \tag{*}$$

It gives a spectral sequence

$$E_1^{pq} = \coprod_{i_0\ldots i_p} R^qQ\big(f_{i_0\ldots i_p*}f_{i_0\ldots i_p}^*(E)\big) \Longrightarrow R^*Q(E). \tag{**}$$

Since $f_{i_0\ldots i_p}$ is affine, (3.4) gives:

$$R^qQ\big(f_{i_0\ldots i_p*}f_{i_0\ldots i_p}^*(E)\big) \simeq f_{i_0\ldots i_p*}R^qQ\big(f_{i_0\ldots i_p}^*(E)\big) \tag{***}$$

for every q, whence in particular

$$E_1^{pq} = 0 \quad \text{for } q > 0,$$

by the affine case, already treated. From (**) we deduce isomorphisms

$$R^nQ(E) \simeq H^n(QE^\bullet) \quad \text{for } n \geq 0,$$

where $E^\bullet$ is the resolution of E defined by the exact sequence (*). But (***) gives $QE^n \simeq E^n$ for $n \geq 0$, whence finally

$$R^nQ(E) = 0 \quad \text{for } n > 0,$$

and $Q(E) \simeq E$, q.e.d.

([1]) Using, for example, the spectral sequence $R^pQ(H^q(E)) \Longrightarrow R^*Q(E)$.

Proof of (3.5.2). The spectral sequence

$$R^pQ(H^q(E)) \Longrightarrow R^*Q(E),$$

[print 191]immediately reduces us to the case where $E \in \mathrm{ob}\,\mathrm{Qcoh}(S)$. Since the composite

$$E \longrightarrow \underline{\underline{R}}QE \longrightarrow E$$

of the adjunction arrows is the identity, the assertion therefore follows from (3.5.1). This completes the proof of (3.5).

Remark 3.6. The conclusion of (3.5) would be false with the hypothesis "separated or noetherian" omitted, as a neat counterexample of J.L. Verdier shows (see Appendix I to this exposé).

Proposition 3.7. Under the hypothesis of (3.5), suppose further that there exists an integer N such that, for every $F \in \mathrm{ob}\,\mathrm{Mod}(S)$ and every quasi-compact open subset U of S,

$$H^i(U, F) = 0 \quad \text{for } i > N \qquad ([1]) .$$

Then:

a) The coherer has finite cohomological dimension.

b) The functor

$$\phi : \mathrm{D}(\mathrm{Qcoh}(S)) \longrightarrow \mathrm{D}(S)$$

is fully faithful and has as essential image the full subcategory $\mathrm{D}(S)_{\mathrm{qcoh}}$ of $\mathrm{D}(S)$ formed by complexes with quasi-coherent cohomology.

Proof of a). When S is affine, the formula $Q = \widetilde{\Gamma}_S$ (cf. proof of (3.2)) implies that Q has cohomological dimension $\leq N$. We pass to the general case. Let us show that there exists an integer a such that $R^qQ(E) = 0$ for every $E \in \mathrm{ob}\,\mathrm{Mod}(S)$ and every $q > a$. The hypothesis implies that every $E \in \mathrm{ob}\,\mathrm{Mod}(S)$ admits a finite right resolution $E^\bullet$ of length N such that the E^i are acyclic, i.e. satisfy $H^q(U, E^i) = 0$ for every open subset U of S and every $q > 0$. It therefore suffices to prove that there exists an integer b such that $R^qQ(E) = 0$ for every acyclic $\underline{O}_S$-Module E and every $q > b$. (We may then take $a = b + N$). Let (U_i) be a finite covering of S by N' affine [print 192]open subsets. Let E be an acyclic $\underline{O}_S$-Module, and consider the resolution $C^\bullet(E)$ of E defined by the covering (U_i) (T.F. II 5.2). There are two cases to examine:

1°) S is separated. The components of $C^\bullet(E)$ are then sums of terms of the form $f_*f^*(E)$, where $f : U \longrightarrow S$ is the inclusion of an affine open subset (hence an affine morphism, since S is separated). We have

$$\underline{\underline{R}}(Qf_*)f^*E \simeq \underline{\underline{R}}Q\,\underline{\underline{R}}f_*f^*E \simeq \underline{\underline{R}}Q\,f_*f^*E,$$

since f^*E is acyclic. On the other hand, by (3.4), we have

$$\underline{\underline{R}}(Qf_*)f^*E \simeq \underline{\underline{R}}(f_*Q)f^*E \simeq f_*\underline{\underline{R}}Qf^*E,$$

since f is affine. We deduce that

$$R^nQ(f_*f^*E) \simeq f_*R^nQ(f^*E) \quad \text{for every } n \geq 0,$$

whence, by the affine case,

$$R^nQ(f_*f^*E) = 0 \quad \text{for } n > N.$$

([1]) By (Tohoku 3.6.5), this condition holds when the underlying space of S is noetherian and has Krull dimension $\leq N$.

Since $C^\bullet(E)$ has length N', we finally have

$$R^nQ(E) = 0 \quad \text{for } n > N + N',$$

and we are done.

2°) S is noetherian. The components of $C^\bullet(E)$ are sums of terms of the form $f_*f^*(E)$, where $f : U \longrightarrow S$ is the inclusion of a quasi-compact and separated open subset. As in case 1°), we have [print 193]

$$\underline{\underline{R}}(Qf_*)f^*E \simeq \underline{\underline{R}}Q\, f_*f^*E \simeq \underline{\underline{R}}(f_*Q)f^*E. \tag{*}$$

Since U is noetherian, $\mathrm{Qcoh}(U)$ has enough objects that are injective in $\mathrm{Mod}(U)$ ([H] II 7.18), so the right derived functor of $f_* : \mathrm{Qcoh}(U) \longrightarrow \mathrm{Qcoh}(S)$ is "induced" by $\underline{\underline{R}}f_* : \mathrm{D}^+(U) \longrightarrow \mathrm{D}^+(S)$ when we identify $\mathrm{D}^+(\mathrm{Qcoh}(U))$ (resp. $\mathrm{D}^+(\mathrm{Qcoh}(S))$) with a full subcategory of $\mathrm{D}^+(U)$ (resp. $\mathrm{D}^+(S)$) by (3.5). Consequently, we have

$$\underline{\underline{R}}(f_*Q)f^*E \simeq \underline{\underline{R}}f_*\underline{\underline{R}}Qf^*E.$$

We deduce a spectral sequence

$$E_2^{pq} = R^pf_*R^qQf^*E \Longrightarrow R^*(f_*Q)f^*E.$$

By case 1°), there exists an integer $n(U)$ depending only on U such that Q has cohomological dimension $\leq n(U)$ on $\mathrm{Mod}(U)$. Thus we have

$$E_2^{pq} = 0 \quad \text{for } q > n(U).$$

But we have

$$E_2^{pq} = 0 \quad \text{for } p > N,$$

because of the "cohomological dimension" hypothesis on S. We therefore have

$$R^n(f_*Q)f^*E = 0 \quad \text{for } n > N + N'',$$

where N'' is an integer greater than all the $n(U)$, with U ranging over the set of intersections $U_{i_0} \cap \ldots \cap U_{i_p}$. Taking account of (*), we obtain

$$R^nQ(f_*f^*E) = 0 \quad \text{for } n > N + N'',$$

whence finally

$$R^nQ(E) = 0 \quad \text{for } n > N + N' + N'',$$

[print 194]which completes the proof of a).

Proof of b). The coherer, having finite cohomological dimension by a), admits ([H] I 5.3) a right derived functor

$$\underline{\underline{R}}Q : \mathrm{D}(S) \longrightarrow \mathrm{D}(\mathrm{Qcoh}(S))$$

(whose restriction to $\mathrm{D}^+(S)$ coincides with the functor $\underline{\underline{R}}Q$ defined above). For $E \in \mathrm{ob}\,\mathrm{D}(\mathrm{Qcoh}(S))$, there is a canonical functorial isomorphism

$$E \xrightarrow{\sim} \underline{\underline{R}}Q(E), \tag{1}$$

since, by (3.5.1), the E^i are acyclic for Q and satisfy $E^i \xrightarrow{\sim} Q(E^i)$. On the other hand, for $E \in \mathrm{ob}\,\mathrm{D}(S)$, we have a canonical functorial homomorphism

$$\underline{\underline{R}}Q(E) \longrightarrow E \tag{2}$$

(coming, in the case where the E^i are Q-acyclic, from the ordinary adjunction homomorphism between ϕ and Q). We check trivially that the homomorphisms (1) and (2) make $\underline{\underline{R}}Q : \mathrm{D}(S) \longrightarrow \mathrm{D}(\mathrm{Qcoh}(S))$ a right adjoint to $\phi : \mathrm{D}(\mathrm{Qcoh}(S)) \longrightarrow \mathrm{D}(S)$. The fact that (1) is an isomorphism implies that ϕ is fully faithful. Finally, using the spectral sequence

$$R^pQ(H^q(E)) \Longrightarrow R^*Q(E),$$

(which converges since Q has finite cohomological dimension), we see that (2) is an isomorphism for $E \in \mathrm{ob}\,\mathrm{D}(S)_{\mathrm{qcoh}}$, so ϕ has as essential image $\mathrm{D}(S)_{\mathrm{qcoh}}$. This proves b) and completes the proof of (3.7).

[print 195]

APPENDIX I
A COUNTEREXAMPLE OF VERDIER(1)

0. Introduction

For a scheme S, we denote by $\mathrm{Mod}(S)$ (resp. $\mathrm{Qcoh}(S)$) the category of $\underline{O}_S$-Modules (resp. quasi-coherent $\underline{O}_S$-Modules). The purpose of this appendix is to draw the reader's attention to certain pathological properties of the category $\mathrm{D}(\mathrm{Qcoh}(S))$, and in particular to prove the following facts:

0.1. Let X be an affine scheme with ring A, and let I be an injective A-module. We know ([H] II 7.14) that, when A is noetherian, the sheaf $\widetilde{I}$ is an injective $\underline{O}_X$-Module. When A is not noetherian, $\widetilde{I}$ need not be injective, or even flasque, even when the underlying space of X is noetherian and of finite Krull dimension. Moreover, the restriction of $\widetilde{I}$ to an open subset U of X is not in general an injective object of $\mathrm{Qcoh}(U)$.

0.2. Let $f : X \longrightarrow Y$ be a morphism of quasi-compact and quasi-separated schemes. The functor $\underline{\underline{R}}f_* : \mathrm{D}^+(X) \longrightarrow \mathrm{D}^+(Y)$ does not in general induce on $\mathrm{D}^+(\mathrm{Qcoh}(X))$ (a full subcategory of $\mathrm{D}^+(X)$ by (II 3.5)) the right derived functor of $f_* : \mathrm{Qcoh}(X) \longrightarrow \mathrm{Mod}(Y)$. More precisely, an injective object of $\mathrm{Qcoh}(X)$ need not be acyclic for $f_* : \mathrm{Mod}(X) \longrightarrow \mathrm{Mod}(Y)$.

0.3. Let S be a quasi-compact and quasi-separated scheme. An object of $\mathrm{Qcoh}(S)$, even an injective one, need not be acyclic for the coherer (II 3.2). The functor $\phi : \mathrm{D}^b(\mathrm{Qcoh}(S)) \longrightarrow \mathrm{D}^b(S)$ (II 3.0) is not in general fully faithful.

1. The counterexamples

We first recall the following lemma (SGA 2 II 9):

Lemma 1.1. Let A be a ring, $X = \mathrm{Spec}(A)$, $\underline{f} = (f_\alpha)$ a finite system of elements of A, Y the closed subset defined by $\underline{f}(x) = 0$. For $i > 0$, the following conditions are equivalent:

(i) for every injective A-module I, we have $H^i_Y(X, \widetilde{I}) = 0$;

(ii) the pro-object $n \mapsto H_i(\underline{f}^n, A)$ is zero, i.e. for every n, there exists $n' \geq n$ [print 196] such that

$$H_i(\underline{f}^{n'}, A) \longrightarrow H_i(\underline{f}^n, A)$$

is zero.

(1) Apart from a few details of presentation, this appendix reproduces two letters from J.-L. Verdier (30/9/67 and 11/10/67) to the editor, in reply to a question from the latter.

1.2 We shall use (1.1) to prove (0.1). Let k be a field, $B = k[[x, y]]$, $\underline{f}$ the system (x, y), $M = \coprod_{n>0} B/(\underline{f}^n)$[146], A the B-algebra $D_B(M)$ (EGA 0_{IV} 18.2.3). We have trivially

$$H_2^A(\underline{f}^n, A) \simeq H_2^B(\underline{f}^n, A). \qquad (^1)$$

On the other hand, we have

$$H_2^B(\underline{f}^n, A) \simeq H_2^B(\underline{f}^n, B) \oplus H_2^B(\underline{f}^n, M)$$

but $H_2^B(\underline{f}^n, B) = 0$ since $\underline{f}$ is a regular sequence (EGA III 1.1.4 and 0_{IV} 15.1.20), so we have

$$H_2^B(\underline{f}^n, A) \simeq H_2^B(\underline{f}^n, M).$$[147]

Finally, we have (EGA III 1.1.3.3)[148]

$$H_2^B(\underline{f}^n, M) \simeq H_B^0(\underline{f}^n, M),$$

where $H_B^0(\underline{f}^n, M)$ is canonically identified with the submodule ${}_{(\underline{f}^n)}M$ of M annihilated by $(\underline{f}^n)$. Hence finally:

$$H_2^A(\underline{f}^n, A) \simeq {}_{(\underline{f}^n)}M.$$[149] $\qquad (*)$

Since the transition morphism $H_2^A(\underline{f}^{n'}, A) \longrightarrow H_2^A(\underline{f}^n, A)$ is induced by multiplication by $\underline{f}^{n'-n}$ (EGA III 1.1.6), we check immediately, using (*), that the pro-object $n \mapsto H_2^A(\underline{f}^n, A)$ is not zero. From (1.1) we then deduce that there exists an injective A-module I such that $H_Y^2(X, \widetilde{I}) \neq 0$, and thus such that $\widetilde{I}$ is not flasque (SGA 2 I 2.12). We note in passing that the projection $\mathrm{Spec}(A) \longrightarrow \mathrm{Spec}(B)$ is a homeomorphism, [print 197]so the underlying space of X is noetherian of dimension 2. Let I be an injective A-module such that $H_Y^2(X, \widetilde{I}) \neq 0$, and denote by U the open complement of Y in X. We have (SGA 2 II 4)

$$H_Y^2(X, I) \simeq H^1(U, I) \neq 0.$$

Now, since U is separated, the functor $\mathrm{D}^+(\mathrm{Qcoh}(U)) \to \mathrm{D}^+(U)$ is fully faithful (II 3.5), so $\widetilde{I}|U$ is not injective in $\mathrm{Qcoh}(U)$. We have therefore proved assertion (0.1).

Denote by Z the quasi-compact and quasi-separated scheme obtained by gluing two copies of X along U, and by j_1, j_2 the two canonical open immersions of X into Z, so that we have a cartesian square

$$\begin{array}{ccc} U & \xrightarrow{i} & X \\ {\scriptstyle i}\downarrow & & \downarrow{\scriptstyle j_1} \\ X & \xrightarrow{j_2} & Z. \end{array}$$

We have

$$j_2^* R^1 j_{1*}(\widetilde{I}) \simeq R^1 i_*(i^* \widetilde{I}),$$

whence

$$H^0(X, R^1 j_{1*}(\widetilde{I})) \simeq H^1(U, \widetilde{I}) \neq 0,$$

so $R^1 j_{1*}(\widetilde{I}) \neq 0$, which proves (0.2).

On the other hand, by (II 3.4 and 3.5), we have

$$\underline{\underline{R}}Q\, \underline{\underline{R}}j_{1*}(\widetilde{I}) \simeq j_{1*}(\widetilde{I}).$$

We deduce a spectral sequence whose exact sequence of low-degree terms gives the isomorphism

$$R^1 j_{1*}(\widetilde{I}) \simeq R^2 Q(j_{1*}\widetilde{I}).$$

(^{1}) The superscript A or B indicates the ring over which the Koszul complex is computed.

Since the sheaf $j_{1*}\widetilde{I}$ is injective in $\mathrm{Qcoh}(Z)$, we have proved the first [print 198]part of (0.3).

Finally, let $E \in \mathrm{ob}\,\mathrm{Qcoh}(Z)$. We have the duality isomorphism

$$\underline{\underline{R}}\,\mathrm{Hom}_{\mathrm{Mod}}(E, \underline{\underline{R}}j_{1*}(\widetilde{I})) \simeq \underline{\underline{R}}\,\mathrm{Hom}_{\mathrm{Mod}}(j_1^*E, \widetilde{I}),$$

but, since X is affine, we have (II 3.5)

$$\underline{\underline{R}}\,\mathrm{Hom}_{\mathrm{Mod}}(j_1^*E, \widetilde{I}) \simeq \underline{\underline{R}}\,\mathrm{Hom}_{\mathrm{Qcoh}}(j_1^*E, I) \simeq \mathrm{Hom}(j_1^*E, \widetilde{I}).$$

We deduce

$$\underline{\underline{R}}\,\mathrm{Hom}_{\mathrm{Mod}}(E, \underline{\underline{R}}j_{1*}(\widetilde{I})) \simeq \mathrm{Hom}(j_1^*E, \widetilde{I}),$$

whence a spectral sequence whose exact sequence of low-degree terms gives the isomorphism

$$\mathrm{Ext}^2_{\mathrm{Mod}}(E, j_{1*}\widetilde{I}) \simeq \mathrm{Hom}(E, R^1j_{1*}\widetilde{I}).$$

Consequently, the identity morphism of $E = R^1j_{1*}(\widetilde{I})$ gives a nonzero element of $\mathrm{Ext}^2_{\mathrm{Mod}}(E, j_{1*}\widetilde{I})$, so the functor $\mathrm{D}^b(\mathrm{Qcoh}(Z)) \longrightarrow \mathrm{D}^b(Z)$ is not fully faithful, which completes the proof of (0.3).

[print 199]

Appendix II
Definition of the analytic index of a relative elliptic complex

In this appendix, we propose to show that the notion of a perfect complex leads to a simple and natural definition of the "analytic index" of a continuous family of elliptic differential operators, or more generally of a "relative elliptic complex". For the initiated reader, let us say that the definition given here has the advantage over the classical definitions (see for example [2][17]) of requiring no deformation of the elliptic complex under consideration.

Sections 1 and 2 consist of reminders about the classical notion of a "mixed structure"([1]). We give, of course, only the minimum of definitions and results required to state and prove, in §3, the central theorem we have in view, which says, a little more precisely, that the direct image, under the projection (assumed proper) onto the base (assumed locally compact), of a relative elliptic complex is a perfect complex. When the base S is compact, the definition of the analytic index as an element of $K(S)$ ("Grothendieck group" of complex vector bundles of finite rank on S) then follows from the globalization sorite of (II 2.3.2).

Throughout what follows, for a topological space S, we shall denote by $\underline{O}_S$ the sheaf of continuous real-valued functions on S.

1. The sorite of mixed structures

In this section, we fix a topological space S.

1.1. Definition of mixed varieties

Let V be an open subset of $\underline{\underline{R}}^n$. For $r \in \underline{\underline{N}}$, we denote by $C^r(V)$ the Fréchet space of real-valued functions of class C^r on V. We put

$$C^\infty(V) = \varprojlim_r C^r(V).$$

Let U be an open subset of S. For $r \in \underline{\underline{N}} \cup \{\infty\}$, we denote by ${}^r\underline{O}_{U\times V}$ the sheaf of rings on $U \times V$ defined by [print 200]

$$\Gamma(U' \times V', {}^r\underline{O}_{U'\times V'}) = C(U', C^r(V')),$$

where U' (resp. V') is an open subset of U (resp. V), and $C(U', C^r(V'))$ denotes the space of continuous maps from U' to $C^r(V')$. There is a canonical injection

$$\mathrm{pr}_1^{-1}(\underline{O}_U) \longrightarrow {}^r\underline{O}_{U\times V},$$

which makes $(U \times V, {}^r\underline{O}_{U\times V})$ a ringed space over U.

A ringed space X over S is called a mixed variety of class C^r ($r \in \underline{\underline{N}} \cup \{\infty\}$) when X is locally isomorphic (as a ringed space over S) to a space of the form $(U \times V, {}^r\underline{O}_{U\times V})$. (When S consists of one point, a mixed variety of class C^r over S is a variety of class C^r in the usual sense).

Let $f : X \to S$ be a mixed variety of class C^r. For $s \in S$, the fiber $X_s = f^{-1}(s)$, equipped with the sheaf $i_s^{-1}(\underline{O}_X)/I_s$, where $i_s : X_s \to X$ is the injection and I_s is the Ideal of sections that "vanish" on X_s, is a variety of class C^r.

([1]) The informed reader is advised to skip them.

1.2. "Relative" differential operators

It is easy to make an infinitesimal study of mixed varieties on the model of Algebraic Geometry (EGA IV 16, SGA 3 VII). We shall limit ourselves to a brief review of the notions we shall need.

1.2.1. Let $f : X \to S$ be a mixed variety of class C^∞ (1.1). For every $n \in \underline{\underline{N}}$, we define the sheaf $P^n_{X/S}$ of principal parts of order n on X. In two different ways, corresponding to the two projections pr_1 and pr_2 from $X \times_S X$ onto X, it is an $\underline{O}_X$-Algebra augmented to $\underline{O}_X$, locally free of finite type as an $\underline{O}_X$-Module. When we do not specify the $\underline{O}_X$-Module structure considered on $P^n_{X/S}$, it will be understood to be the one coming from pr_1. There are canonical isomorphisms [print 201]

$$P^0_{X/S} \simeq \underline{O}_X,$$
$$P^1_{X/S} \simeq \underline{O}_X \oplus \Omega^1_{X/S},$$

where $\Omega^1_{X/S}$ is the sheaf of relative differentials of X over S, or relative cotangent sheaf. We denote by $T_{X/S}$ the dual of $\Omega^1_{X/S}$, called the relative tangent sheaf. The sheaf $T_{X/S}$ (resp. $\Omega^1_{X/S}$) is the sheaf of sections of a vector bundle of finite rank on X, called the relative tangent bundle (resp. cotangent bundle), which is a mixed S-variety over X.

The $f^{-1}(\underline{O}_S)$-linear homomorphism (corresponding to the $\underline{O}_X$-Algebra structure defined by pr_2)

$$d^n_{X/S} : \underline{O}_X \longrightarrow P^n_{X/S},$$

which associates to a section of $\underline{O}_X$ its principal part of order n, is called the universal differential operator of order n on X relative to S.

1.2.2. More generally, for a locally free $\underline{O}_X$-Module E of finite type, we can consider the sheaf $P^n_{X/S}(E) = P^n_{X/S} \otimes_{\underline{O}_X} E$ [150] of principal parts of order n of E (the tensor product being defined by means of the second $\underline{O}_X$-Module structure on $P^n_{X/S}$), and we have a universal differential operator

$$d^n_{X/S}(E) : E \longrightarrow P^n_{X/S}(E).$$

Let E, F be locally free $\underline{O}_X$-Modules of finite type. By definition, a differential operator from E to F of order $\leq n$ is an $f^{-1}(\underline{O}_S)$-linear homomorphism [151] from E to F that factors through $d^n_{X/S}(E)$ as an $\underline{O}_X$-linear homomorphism from $P^n_{X/S}(E)$ to F (the factorization is then unique). In other words, the operator $d^n_{X/S}(E)$ gives a bijection

$$\mathrm{Hom}_{\underline{O}_X}\left(P^n_{X/S}(E), F\right) \xrightarrow{\sim} \mathrm{Diff}^n(E, F), \tag{1.2.2.1}$$

where $\mathrm{Diff}^n(E, F)$ denotes the set of differential operators from E [print 202] to F of order $\leq n$.

1.2.3. We shall also have to consider differential operators with complex coefficients. Denote by $\underline{O}_S \otimes \underline{\underline{C}}$ (resp. $\underline{O}_X \otimes \underline{\underline{C}}$) the complexification of $\underline{O}_S$ (resp. $\underline{O}_X$). Let E, F be locally free $(\underline{O}_X \otimes \underline{\underline{C}})$-Modules of finite type. By a differential operator from E to F of order $\leq n$, we shall mean an $f^{-1}(\underline{O}_S \otimes \underline{\underline{C}})$-linear homomorphism from E to F that factors through $d^n_{X/S}(E)$ as an $(\underline{O}_X \otimes \underline{\underline{C}})$-linear homomorphism from $P^n_{X/S}(E)$ to F. Denoting the set of these operators by $\mathrm{Diff}^n_{\underline{\underline{C}}}(E, F)$, the operator $d^n_{X/S}(E)$ therefore defines a bijection

$$\mathrm{Hom}_{\underline{O}_X \otimes \underline{\underline{C}}}\left(P^n_{X/S}(E), F\right) \xrightarrow{\sim} \mathrm{Diff}^n_{\underline{\underline{C}}}(E, F). \tag{1.2.3.1}$$

1.2.4. For $n \geq 1$, there is a canonical exact sequence

$$0 \longrightarrow S^n(\Omega^1_{X/S}) \longrightarrow P^n_{X/S} \longrightarrow P^{n-1}_{X/S} \longrightarrow 0.(^1)$$

Hence, tensoring over $\underline{O}_X$(2) with a locally free $\underline{O}_X$-Module (resp. $(\underline{O}_X \otimes \underline{\underline{C}})$-Module) E of

(1) $S^n(\ )$ denotes the nth symmetric power.

(2) For the second $\underline{O}_X$-Module structure of $P^n_{X/S}$ and $P^{n-1}_{X/S}$.

finite type, we obtain an exact sequence of $\underline{O}_X$ (resp. $(\underline{O}_X \otimes \underline{\underline{C}})$)-Modules:

$$0 \longrightarrow S^n(\Omega^1_{X/S}) \otimes_{\underline{O}_X} E \longrightarrow P^n_{X/S}(E) \longrightarrow P^{n-1}_{X/S}(E) \longrightarrow 0. \tag{1.2.4.1}$$

Let F be a locally free $\underline{O}_X$ (resp. $(\underline{O}_X \otimes \underline{\underline{C}})$)-Module of finite type. Applying the functor $\mathrm{Hom}_{\underline{O}_X}(.,F)$ (resp. $\mathrm{Hom}_{\underline{O}_X \otimes \underline{\underline{C}}}(.,F)$) to (1.2.4.1), we deduce, taking account of the identifications (1.2.2.1) and (1.2.3.1)[152], an exact sequence

$$\begin{aligned} 0 \longrightarrow \mathrm{Diff}^{n-1}(E,F) \longrightarrow \mathrm{Diff}^n(E,F) \xrightarrow{\sigma_n} \mathrm{Hom}_{\underline{O}_X}(S^n(\Omega^1_{X/S}) \otimes_{\underline{O}_X} E, F) \\ (\text{resp. } 0 \longrightarrow \mathrm{Diff}^{n-1}_{\underline{\underline{C}}}(E,F) \longrightarrow \mathrm{Diff}^n_{\underline{\underline{C}}}(E,F) \longrightarrow \mathrm{Hom}_{\underline{O}_X \otimes \underline{\underline{C}}}(S^n(\Omega^1_{X/S}) \otimes_{\underline{O}_X} E, F)). \end{aligned} \tag{1.2.4.2}$$

[print 203](When X is paracompact and $\underline{O}_X$ is soft (cf. II 2.3.1), we can complete the exact sequence (1.2.4.2) by adding a zero on the right). The homomorphism σ_n is called the <u>symbol</u>; it associates to a differential operator from E to F of order $\leq n$ a "homogeneous polynomial map of degree n from $\Omega^1_{X/S}$ to $\underline{\mathrm{Hom}}_{\underline{O}_X}(E,F)$ (resp. $\underline{\mathrm{Hom}}_{\underline{O}_X \otimes \underline{\underline{C}}}(E,F)$)", which we may also regard, if we wish, as a homomorphism $p^*(E) \longrightarrow p^*(F)$ (where p is the projection onto X from the relative cotangent bundle (1.2.1)), satisfying a certain homogeneity condition.

Let E, F, G be locally free $\underline{O}_X$-Modules of finite type, and $u \in \mathrm{Diff}^m(E,F)$, $v \in \mathrm{Diff}^n(F,G)$, whence $vu \in \mathrm{Diff}^{m+n}(E,G)$. We easily check the relation

$$\sigma_{m+n}(vu) = \sigma_n(v)\sigma_m(u). \tag{1.2.4.3}$$

(The same relation holds for differential operators with complex coefficients). In particular, when the composite of two differential operators is zero, the composite of their symbols is also zero.

1.2.5. Let E be a complex, with bounded degrees, of differential operators with complex coefficients on X relative to S (1.2.3), with $d^i : E^i \longrightarrow E^{i+1}$ of order $\leq n_i$. By (1.2.4.3), we have

$$\sigma_{n_{i+1}}(d^{i+1})\, \sigma_{n_i}(d^i) = 0.$$

We call E an <u>elliptic complex</u> of order $\underline{n} = (n_i)$ when the complex $(p^*(E^i), \sigma_{n_i}(d^i))$ (where p is the projection onto X from the cotangent bundle) is acyclic outside the zero section.

When S consists of one point, the preceding notion of an elliptic complex on X coincides with the usual notion, studied in detail in [4] or [16].

1.2.6. Let $s \in S$, and let i_s be the inclusion of the fiber X_s into X (considered as a monomorphism of ringed spaces) cf. (1.1).[153] For $n \in \underline{\underline{N}}$, there is a canonical isomorphism [print 204]

$$i_s^*(P^n_{X/S}) \simeq P^n_{X_s},$$

whence a restriction homomorphism

$$i_s^* : \mathrm{Diff}^n(E,F) \longrightarrow \mathrm{Diff}^n(i_s^*(E), i_s^*(F)),$$

for locally free $\underline{O}_X$-Modules E, F of finite type (and an analogous homomorphism for operators with complex coefficients). Let E be a complex, with bounded degrees, of differential operators with complex coefficients on X; for E to be elliptic (1.2.5), it is necessary and sufficient that, for every $s \in S$, $i_s^*(E)$ be an elliptic complex (verification left to the reader).

2. Rigidity lemmas

2.0. It is "well known" that compact differentiable varieties are "rigid", i.e. admit no continuous deformation other than a trivial one. However, since this result will be essential to us in §3, and references seem rather scarce, it may be useful to give precise statements and a few proofs.

We fix a topological space S. The mixed varieties considered will be assumed to be of class C^r, with $r \geq 1$ (possibly $r = \infty$).

2.1. Notation. Let X, Y be mixed S-varieties. We denote by $\mathrm{Hom}(X, Y)$ the set of S-morphisms from X to Y, and by $\underline{\mathrm{Hom}}(X, Y)$ the sheaf on X of germs of S-morphisms from X to Y (defined by

$$\underline{\mathrm{Hom}}(X, Y)(U) = \mathrm{Hom}(U, Y)$$

for U open in X).

Let $s \in S$. We denote by i_s the inclusion of X_s into X. There is a canonical induction homomorphism

$$i_s^{-1}\underline{\mathrm{Hom}}(X, Y) \longrightarrow \underline{\mathrm{Hom}}(X_s, Y_s), \tag{2.1.1}$$

[print 205]which is an epimorphism, as we see immediately. When $Y = S \times \underline{R}$, the sheaf $\underline{\mathrm{Hom}}(X, Y)$ is canonically identified with $\underline{O}_X$, and the epimorphism (2.1.1) with the epimorphism

$$i_s^{-1}(\underline{O}_X) \longrightarrow i_s^*(\underline{O}_X) = \underline{O}_{X_s}$$

considered in (1.1).

Proposition([1]) **2.2.** Let X, Y be mixed S-varieties, and $s \in S$. Suppose that the fiber X_s is paracompact and that s has a fundamental system of paracompact neighborhoods. Let A be a closed subset of X_s, and $f \in H^0(A, \underline{\mathrm{Hom}}(X_s, Y_s))$. The subsheaf F of $i_s^{-1}\underline{\mathrm{Hom}}(X, Y)$ formed by the germs "that extend f" (i.e. whose image under (2.1.1) coincides with f over A) is soft.

Proof. Let X' be an open neighborhood of x in X, isomorphic (as a mixed variety) to $S' \times U$, where S' is a paracompact neighborhood of s and U a relatively compact open subset of $\underline{\underline{R}}^m$, and such that the image under f of $X'_s \cap A$ is contained in an open subset of Y isomorphic to $S' \times V$, where V is a convex open subset of $\underline{\underline{R}}^n$ containing the origin. By (T.F. II 3.4.1), it suffices to show that every section of F over a closed subset A_0 of X_s contained in X'_s extends to X'_s. Now X'_s ($\simeq \{s\} \times U$) is relatively compact, so a closed subset of X_s contained in X'_s is compact. Identify X' (resp. Y') with $S' \times U$ (resp. $S' \times V$), put $A' = A \cap X'_s$, and let A_0 be a compact subset of $X'_s = \{s\} \times U$. Since A' is closed in $\{s\} \times U$, which is paracompact, we have ([T.F.] II 3.3.1)([1])

$$H^0(A', \underline{\mathrm{Hom}}(X_s, Y_s)) = \varinjlim_{U' \supset A'} \mathrm{Hom}(U', Y'_s),$$

over the open subsets U' of X'_s containing A'. Similarly, since A_0 is closed in $S' \times \overline{U}$, which is paracompact (a product of a paracompact space and a compact space), we have

$$H^0(A_0, i_s^{-1}\underline{\mathrm{Hom}}(X, Y)) = \varinjlim_{U_0 \supset A_0} \mathrm{Hom}(U_0, Y'),$$

over the open subsets U_0 of X' containing A_0. The proposition will therefore be a [print 206]consequence of the following lemma:

Lemma 2.2.1. Let U be an open subset of $\underline{\underline{R}}^m$, V a convex open subset of $\underline{\underline{R}}^n$ containing 0. Let A_0, A_1 be closed subsets of $\{s\} \times U$, let U_0 (resp. U_1) be an open subset of $S \times U$ (resp. $\{s\} \times U$) containing A_0 (resp. A_1). Let $f_1 : U_1 \to V$ be a morphism and let $f_0 : U_0 \to S \times V$ be an S-morphism inducing on $U_0 \cap (\{s\} \times U)$ a morphism defining the same germ as f_1 in a neighborhood of A_1. There then exists an open subset U' of $S \times U$ containing $s \times U$, and an S-morphism $\bar{f} : U' \longrightarrow S \times V$ such that

(i) the germ of $\bar{f}$ along $\{s\} \times U$ extends the germ defined by f_0;

([1]) Communicated to the editor by A. Douady.

([1]) A closed subset of a paracompact space has a fundamental system of paracompact neighborhoods.

(ii) the morphism induced by $\bar{f}$ on $\{s\} \times U$ defines the same germ as f_1 in a neighborhood of A_1.

Proof. Replacing f_1 by af_1, where a is a C^r function on U, equal to 1 in a neighborhood of A_1 and with support in U_1, we may assume that $U_1 = U$. Let b be a C^r function on U, equal to 1 on A_0 and with support in $U_0 \cap (\{s\} \times U)$. Then

$$U' = U_0 \cup (S \times (U - \operatorname{Supp} b))$$

is an open subset of $S \times U$ containing $\{s\} \times U$. The morphism $\bar{f} : U' \to S \times V$ defined by

$$\bar{f}(t, x) = (t,\ b(x) \operatorname{pr}_2 f_0(t, x) + (1 - b(x)) f_1(x))$$

visibly satisfies conditions (i) and (ii).

Corollary 2.3. Under the hypotheses of (2.2), the sheaves $i_s^{-1}\underline{\mathrm{Hom}}(X, Y)$, $\underline{\mathrm{Hom}}(X_s, Y_s)$ are soft. In particular, the sheaves $i_s^{-1}(\underline{O}_X)$, $\underline{O}_{X_s}$ are soft(1). For every closed subset A of X_s, the map

$$H^0(X_s, i_s^{-1}\underline{\mathrm{Hom}}(X, Y)) \longrightarrow H^0(A, \underline{\mathrm{Hom}}(X_s, Y_s))$$

deduced from (2.1.1) is surjective.

[print 207]Proof. The softness of $i_s^{-1}\underline{\mathrm{Hom}}(X, Y)$ follows from (2.2) applied with $A = \varnothing$; taking $S = s$, we deduce that $\underline{\mathrm{Hom}}(X_s, Y_s)$ is soft. The last assertion follows from (2.2).

Corollary 2.4. Under the hypotheses of (2.2), let E be an $\underline{O}_X$-Module. For every closed subset A of X_s, the canonical homomorphism

$$H^0(X_s, i_s^{-1}(E)) \longrightarrow H^0(A, i_s^*(E)),$$

deduced from the epimorphism

$$i_s^{-1}(E) \longrightarrow i_s^*(E) = i_s^{-1}(E) \otimes_{i_s^{-1}(\underline{O}_X)} \underline{O}_{X_s},$$

is an epimorphism.

Proof. Since $i_s^{-1}(\underline{O}_X)$ is soft (2.3), the functor $H^0(A, .)$ is exact on the category of $i_s^{-1}(\underline{O}_X)$-Modules (T.F. II 3.7.1, 3.4.2, 4.4.3), so the homomorphism

$$H^0(A, i_s^{-1}(E)) \longrightarrow H^0(A, i_s^*(E))$$

is an epimorphism. Since $i_s^{-1}(E)$ is soft, the homomorphism

$$H^0(X_s, i_s^{-1}(E)) \longrightarrow H^0(A, i_s^{-1}(E))$$

is an epimorphism, and we are done.

Corollary 2.5. Let $p : X \to S$ be a mixed variety, and $s \in S$. Suppose that s has a fundamental system of paracompact neighborhoods S' such that the $p^{-1}(S')$ form a fundamental system of paracompact neighborhoods of X_s (a condition satisfied, for example, when S is locally compact and p is a proper map of topological spaces). Then, for every $\underline{O}_X$-Module E, we have

$$R^i p_*(E)_s = 0 \quad \text{for } i > 0.$$

(1) For $\underline{O}_{X_s}$, this is "very well known". The editor does not know whether $\underline{O}_X$ is soft when the hypotheses of (2.2) hold for every $s \in S$ and X is paracompact. Under those hypotheses, this would trivially imply the softness of $i_s^{-1}(\underline{O}_X)$ as well as (2.5) below.

[print 208]Proof. By (T.F. II 4.17.1), we have

$$R^i p_*(E)_s = H^i(X_s, i_s^{-1}(E)).$$

But, since $i_s^{-1}(\underline{O}_X)$ is soft (2.3), we have

$$H^i(X_s, i_s^{-1}(E)) = 0 \quad \text{for } i > 0,$$

whence the assertion.

Corollary 2.6. Let X, Y be mixed S-varieties, and $s \in S$. Suppose that X satisfies the hypotheses of (2.5). For every morphism $f : X_s \to Y_s$, there exists an open neighborhood S' of s and a morphism $\bar{f} : X|S' \longrightarrow Y|S'$ such that $\bar{f}_s = f$.

Proof. This follows immediately from (2.3) (with $A = X_s$) and the fact that, by the hypotheses on X, we have

$$H^0(X_s, i_s^{-1}\underline{\mathrm{Hom}}(X, Y)) = \varinjlim_{S' \ni s} \mathrm{Hom}(X|S', Y) \qquad (S' \text{ open})$$

(T.F. II 3.3.1).

Proposition 2.7. Let X, Y be mixed S-varieties, $f : X \to Y$ a morphism, $s \in S$, $x \in X_s$. If the tangent linear map of $f_s : X_s \to Y_s$ is invertible at x, then f is a local isomorphism at x, i.e. there exists an open neighborhood U of x in X and an open subset V of Y such that f induces an isomorphism from U onto V.

Proof. This is an easy consequence of the implicit function theorem. The details are left to the reader.

Corollary 2.8. Let X, Y be mixed S-varieties, and $s \in S$. Suppose that s has a fundamental system of neighborhoods S' such that the $X|S'$ (resp. $Y|S'$) form a fundamental system of neighborhoods of X_s (resp. Y_s) (a condition satisfied, for example, when S is locally compact and X (resp. Y) is proper over S). Let $f : X \to Y$ be an S-morphism such that $f_s : X_s \to Y_s$ is an isomorphism. There then exists a neighborhood S' [print 209]of s such that $f|X|S' : X|S' \longrightarrow Y|S'$ is an isomorphism.

Proof. By (2.7), there exists an open neighborhood U of X_s in X such that $f|U$ is étale (i.e. a local isomorphism). We may assume that $U = X|S'$, where S' is a neighborhood of s in S; in other words, after restricting S, we may assume that f is étale. We then easily check that the fiber product $Z = X \times_Y X$ is naturally equipped with the structure of a mixed S-variety (étale over X for each projection), and that the diagonal morphism $\Delta : X \to Z$ is an open immersion([1]). Moreover, it is immediate that Z satisfies the same hypothesis as X and Y, namely that the open subsets of the form $Z|S'$, for S' an open neighborhood of s in S, form a fundamental system of neighborhoods of Z_s. Since $\Delta_s : X_s \to Z_s$ is an isomorphism, it follows, after restricting S, that Δ is an isomorphism. Thus f is an open immersion, hence an isomorphism after shrinking S again, and we are done.

Corollary 2.9. Let X be a mixed S-variety, and $s \in S$. Suppose that the hypothesis of (2.5) holds. There then exists a neighborhood S' of s in S such that $X|S'$ is S'-isomorphic to $S' \times X_s$.

Proof. Apply (2.6) and (2.8).

Corollary 2.10. Let $s \in S$ and let V be a paracompact variety. Put $X = S \times V$, and denote by i_s the injection of V into X defined by s. Suppose that the hypothesis of (2.5) holds. Let E be a locally free $\underline{O}_X$ (resp. $\underline{O}_X \otimes \underline{C}$)-Module of finite type. There then exists a neighborhood S' of s in S such that $E|X|S'$ is isomorphic to $\mathrm{pr}_2^* i_s^*(E)$ ($\mathrm{pr}_2 : S' \times V \to V$ being the second projection).

([1]) Note the analogy with Algebraic Geometry (SGA 1 I 5.1).

Proof. Put $F = \mathrm{pr}_2^*\, i_s^*(E)$. By (2.4), applied to the Module $\underline{\mathrm{Hom}}(E,F)$, the induction homomorphism

$$\mathrm{Hom}\big(i_s^{-1}(E), i_s^{-1}(F)\big) \longrightarrow \mathrm{Hom}\big(i_s^*(E), i_s^*(F)\big)$$

[print 210]is an epimorphism. On the other hand, we have (T.F. II 3.3.1):

$$\begin{aligned}\mathrm{Hom}\big(i_s^{-1}(E), i_s^{-1}(F)\big) &\simeq H^0\big(X_s, i_s^{-1}\underline{\mathrm{Hom}}(E,F)\big)\\ &= \varinjlim_{S'\ni s} H^0\big(X|S', \underline{\mathrm{Hom}}(E,F)\big) \qquad (S' \text{ open})\\ &\simeq \varinjlim_{S'\ni s} \mathrm{Hom}(E|X|S', F|X|S').\end{aligned}$$

Thus the canonical isomorphism $i_s^*(E) \simeq i_s^*(F)$ is induced by a homomorphism $f : E \to F$ (after shrinking S). Since f_s is an isomorphism, f is an isomorphism in a neighborhood of X_s by Nakayama, hence over $X|S'$, where S' is a suitable neighborhood of s, q.e.d.

3. The finiteness theorem

3.0. Conventions and notation

3.0.1. Let X and Y be topological spaces. We denote by $C(X,Y)$ the set of continuous maps from X to Y, and by $\underline{C}(X,Y)$ the sheaf on X defined by $U \mapsto C(U,Y)$.

3.0.2. Let E, F be Banach spaces (real or complex). We denote by $\mathrm{L}(E,F)$ the Banach space of continuous linear maps from E to F.

3.0.3. The mixed varieties considered in this section will be assumed to be of class C^∞. Let X be a mixed S-variety; we shall often identify a locally free $\underline{O}_X$ (resp. $\underline{O}_X \otimes \underline{\underline{C}}$)-Module of finite type with the corresponding real (resp. complex) vector bundle.

3.0.4. Let X be a compact (ordinary) variety, and let E be a vector bundle (real or complex) of finite rank on X. We denote by $C^\infty(X,E)$ the Fréchet space of sections (of class C^∞) of E on X. For $r \in \underline{\underline{Z}}$, we denote by $C_2^r(X,E)$ the Hilbert space of sections of E that are r times differentiable in L^2 ([16] IX). For $s \geq r$, $C_2^s(X,E)$ maps into $C_2^r(X,E)$ by a continuous injective linear map with dense image, and we have [print 211](loc.cit.)

$$C^\infty(X,E) = \varprojlim_r C_2^r(X,E).$$

3.0.5. Let S be a topological space. We shall need the notion of a "Hilbert bundle" over S, for which we refer the reader to [12] or [14]. There is an additive functor from the category of Hilbert bundles (real resp. complex) over S to that of $\underline{O}_S$ resp.[154] $\underline{O}_S \otimes \underline{\underline{C}}$-Modules, associating to each Hilbert bundle E the sheaf $\underline{E}$ of continuous sections of E.

3.1. Auxiliary technical results

Lemma 3.1.1. Let X, Y be topological spaces, with X locally compact. Denote by $\underline{C}'(X,Y)$ the presheaf defined on the basis of relatively compact open subsets of X by $\underline{C}'(X,Y)(U) = C(\bar{U}, Y)$ (3.0.1). The canonical restriction morphism

$$\underline{C}'(X,Y) \longrightarrow \underline{C}(X,Y)$$

induces an isomorphism on the associated sheaves.

Proof. This is immediate.

Lemma 3.1.2. Let E, F be locally convex topological vector spaces (over $\underline{\underline{C}}$, to fix ideas), and $u : E \to F$ a continuous linear map (resp. continuous with dense image). Let X be a compact space. Equip $C(X, E)$, $C(X, F)$ with the topology of uniform convergence. Then the linear map

$$u_X : C(X, E) \longrightarrow C(X, F)$$

induced by u is continuous (resp. continuous with dense image).

[print 212]Proof. It is clear that u_X is continuous. On the other hand (Bourbaki, Intégration, chap. III § 2, lemmas 1 and 2), the subspace $C(X, \underline{\underline{C}}) \otimes_{\underline{\underline{C}}} E$ of $C(X, E)$ (formed by the continuous maps from X to E whose values lie in a finite-dimensional vector subspace) is dense; similarly, $C(X, \underline{\underline{C}}) \otimes_{\underline{\underline{C}}} F$ is dense in $C(X, F)$. Suppose u is continuous with dense image. Then

$$u_X(C(X, \underline{\underline{C}}) \otimes_{\underline{\underline{C}}} E) = C(X, \underline{\underline{C}}) \otimes_{\underline{\underline{C}}} u(E)$$

is dense in $C(X, \underline{\underline{C}}) \otimes_{\underline{\underline{C}}} F$, whence, taking closures, we deduce that $u_X(C(X, E))$ is dense in $C(X, F)$, q.e.d.

3.1.3. Let V be a compact variety, E and F complex vector bundles([1]) of finite rank on V, and $n \in \underline{\underline{N}}$. The space $\mathrm{Diff}^n_{\underline{\underline{C}}}(E, F)$, identified by (1.2.3.1) with

$$\mathrm{Hom}(P^n(E), F) \simeq C^\infty(V, \underline{\mathrm{Hom}}(P^n(E), F)),$$

is naturally equipped with the structure of a Fréchet space. Let $u \in \mathrm{Diff}^n_{\underline{\underline{C}}}(E, F)$; we easily check that the linear map from $C^\infty(V, E)$ to $C^\infty(V, F)$ defined by u is continuous. Denoting by $\mathrm{L}(C^\infty(V, E), C^\infty(V, F))$ the space of continuous linear maps from $C^\infty(V, E)$ to $C^\infty(V, F)$, we therefore have a linear map

$$\phi : \mathrm{Diff}^n_{\underline{\underline{C}}}(E, F) \longrightarrow \mathrm{L}(C^\infty(V, E), C^\infty(V, F)).$$

On the other hand, for $r \in \underline{\underline{Z}}$, u acts by extension of scalars ([16] IX Th. 12) on the C^r_2 sections of E and defines a linear map

$$\phi_r : \mathrm{Diff}^n_{\underline{\underline{C}}}(E, F) \longrightarrow \mathrm{L}(C^r_2(V, E), C^{r-n}_2(V, F)).$$

We readily check that:

(i) the ϕ_r are continuous (the target is a Banach space [print 213](3.0.2));

(ii) $\phi = \varprojlim \phi_r$ (i.e. for $u \in \mathrm{Diff}^n_{\underline{\underline{C}}}(E, F)$, $\phi(u)$ is the limit of the projective system of the $\phi_r(u)$).

(The preceding assertions follow easily from the results of (loc.cit.):[155] we reduce immediately to $n = 0$; (ii) then follows directly from the definitions, while (i) follows (after localizing the problem by a partition of unity) from the formula giving the successive derivatives of a product).

Lemma 3.1.4. Let S be a topological space, V a compact variety, E and F complex vector bundles of finite rank on V, and $n \in \underline{\underline{N}}$. Denote by X the mixed S-variety $S \times V$. For $u \in \mathrm{Diff}^n_{\underline{\underline{C}}}(\mathrm{pr}_2^*(E), \mathrm{pr}_2^*(F))$, the map $s \mapsto i_s^*(u)$ from S to $\mathrm{Diff}^n_{\underline{\underline{C}}}(E, F)$ is continuous. The linear map

$$\mathrm{Diff}^n_{\underline{\underline{C}}}(\mathrm{pr}_2^*(E), \mathrm{pr}_2^*(F)) \longrightarrow C(S, \mathrm{Diff}^n_{\underline{\underline{C}}}(E, F))$$

thus defined is an isomorphism.

([1]) To fix ideas (analogous results hold in the real case).

Proof. Using the identification (1.2.3.1), we may restrict to $n = 0$. Putting $\underline{\mathrm{Hom}}(E, F) = G$, we are then reduced to showing that, for $u \in \Gamma(X, \mathrm{pr}_2^*(G))$, the map $s \mapsto u_s = i_s^*(u)$ from S to $C^\infty(V, G)$ is continuous, and that the linear map

$$\Gamma(X, \mathrm{pr}_2^*(G)) \longrightarrow C(S, C^\infty(V, G))$$

so obtained is an isomorphism. But this follows immediately from the definition of mixed varieties (1.1).

3.1.5. Under the hypotheses of (3.1.4), let

$$u \in \mathrm{Diff}^n(\mathrm{pr}_2^*(E), \mathrm{pr}_2^*(F)).(^1)$$

[print 214]Identifying u, by (3.1.4), with a continuous map from S to $\mathrm{Diff}^n(E, F)$, we see that, for U open in S, the morphism

$$\mathrm{pr}_{1*}(u)(U) : \Gamma(X|U, E) \longrightarrow \Gamma(X|U, F)$$

is canonically identified with the linear map

$$C(U, C^\infty(V, E)) \longrightarrow C(U, C^\infty(V, F))$$

deduced from

$$\phi u : U \longrightarrow \mathrm{L}(C^\infty(V, E), C^\infty(V, F)). \tag{3.1.3}$$

Suppose S is locally compact. Then (3.1.1) the morphism

$$\mathrm{pr}_{1*}(u) : \mathrm{pr}_{1*}\,\mathrm{pr}_2^*(E) \longrightarrow \mathrm{pr}_{1*}\,\mathrm{pr}_2^*(F)$$

is the morphism of sheaves associated to the morphism of presheaves

$$u_\infty : \underline{C}'(S, C^\infty(V, E)) \longrightarrow \underline{C}'(S, C^\infty(V, F))$$

defined by ϕu. On the other hand, for $r \in \underline{\underline{Z}}$,

$$\phi_r u : S \longrightarrow \mathrm{L}(C_2^r(V, E), C_2^{r-n}(V, F))$$

defines a morphism of presheaves

$$u_r : \underline{C}'(S, C_2^r(V, E)) \longrightarrow \underline{C}'(S, C_2^{r-n}(V, F)).^{[156]}$$

By (3.1.3 (ii)), the u_r form a projective system whose limit is u_∞. It will be convenient to interpret $\phi_r u$ as a morphism of trivial Hilbert bundles (3.0.5)

$$S \times C_2^r(V, E) \longrightarrow S \times C_2^{r-n}(V, F),$$

[print 215]inducing on the sheaves of sections the morphism of sheaves associated to u_r.

3.2. Definition of the index

Theorem 3.2.1. Let S be a locally compact space, $p : X \to S$ a proper mixed variety, $E^\bullet$ an elliptic complex of order n of complex vector bundles on X (1.2.5). Then the complex $\underline{\underline{R}}p_*(E^\bullet)$ (which, by (2.5), is isomorphic, to $p_*(E^\bullet)$ in the derived category of that of $\underline{O}_S \otimes \underline{\underline{C}}$-Modules) is perfect.

(1) We omit the subscript $\underline{\underline{C}}$ for brevity.

Proof. The question being local on S, we may assume (2.9) that $X = S \times V$, where V is a compact variety, and $p = \mathrm{pr}_1$. Moreover, we may assume (2.10) that, for every i, $E^i = \mathrm{pr}_2^*(F^i)$, where F^i is a complex vector bundle of finite rank on V. By (3.1.5), the complex $p_*(E)$ is associated to the complex of presheaves $(\underline{C}'(S, C^\infty(V, F^i)), d^i_\infty)$ (where the $d^i : E^i \longrightarrow E^{i+1}$ are the given differential operators), which we shall denote simply by $\underline{C}'(S, C^\infty(V, F^\bullet))$. Put, for each $i \in \underline{\underline{Z}}$,

$$m_i = \sum_{j<i} n_j \qquad \text{(we agree that } n_i = 0 \text{ when } E^i = 0\text{)}.$$

For each $r \in \underline{\underline{Z}}$, we have (3.1.5) a complex of presheaves $(\underline{C}'(S, C_2^{r-m_i}(V, F^i)), d^i_{r-m_i})$, which we shall denote by $\underline{C}'(S, C_2^r(V, F^\bullet))$. The $\underline{C}'(S, C_2^r(V, F^\bullet))$ form a projective system, and we have (3.1.5):

$$\underline{C}'(S, C^\infty(V, F^\bullet)) = \varprojlim_r \underline{C}'(S, C_2^r(V, F^\bullet)). \tag{3.2.1.1}$$

Finally, we shall denote by $F_r^\bullet$ the complex of trivial Hilbert bundles over S defined by $\phi_{r-m_i} d^i$, and by $\underline{F}_r^\bullet$ the corresponding complex of sheaves of sections, which is precisely (3.1.5) the complex of sheaves associated to $\underline{C}'(S, C_2^r(V, F^\bullet))$. With this notation, to prove the theorem it suffices to prove the following assertions:

(a) For $r \in \underline{\underline{Z}}$, the canonical morphism [print 216]

$$\underline{C}'(S, C^\infty(V, F^\bullet)) \longrightarrow \underline{C}'(S, C_2^r(V, F^\bullet))$$

is a quasi-isomorphism of complexes of presheaves, i.e. induces isomorphisms on the cohomology presheaves.

(b) For $r \in \underline{\underline{Z}}$, the complex $\underline{F}_r^\bullet$ is perfect.

Proof of (b). The ellipticity of $E^\bullet$ implies ([16] XI Th.6([1])) that, for every $s \in S$, the differential of $(F_r^\bullet)_s$ has closed image and the $H^i((F_r^\bullet)_s)$ are finite-dimensional vector spaces. In other words, to use the terminology of [11], $F_r^\bullet$ is a direct quasi-acyclic complex of Hilbert bundles. Consequently, by ([12] 2.3), $F_r^\bullet$ is locally homotopically equivalent to a bounded complex of vector bundles of finite rank. Since the "sheaf of sections" functor is additive (3.0.5), $\underline{F}_r^\bullet$ is locally homotopically equivalent to a bounded complex of locally free sheaves of finite type, and hence is perfect.

For the proof of (a), we shall need the following intermediate assertion:

(a') For $p \geq q$, the canonical morphism

$$\underline{F}_p^\bullet \longrightarrow \underline{F}_q^\bullet$$

is a quasi-isomorphism.

Proof of (a'). Denote by $M_{pq}^\bullet$ the complex of Hilbert bundles that is the cone of the morphism $F_p^\bullet \longrightarrow F_q^\bullet$. The ellipticity of E implies that, for every $s \in S$, the morphism $(F_p^\bullet)_s \longrightarrow (F_q^\bullet)_s$ is a quasi-isomorphism ([16] XI Cor. 2 of Th. 6). In other words, $M_{pq}^\bullet$ is a direct acyclic complex. It follows ([11] 2.3) that $M_{pq}^\bullet$ is locally homotopically trivial, and hence that $\underline{M}_{pq}^\bullet$, which is precisely the cone of $\underline{F}_p^\bullet \longrightarrow \underline{F}_q^\bullet$, is acyclic, q.e.d.

Proof of (a). We must show that, for every relatively compact open subset U of S, the morphism [print 217]

$$C(\overline{U}, C^\infty(V, F^\bullet)) \longrightarrow C(\overline{U}, C_2^r(V, F^\bullet))$$

is a quasi-isomorphism. Replacing S by $\overline{U}$ (and $E^\bullet$ by the complex induced over $\overline{U}$), we are reduced to showing that

$$C(S, C^\infty(V, F^\bullet)) \longrightarrow C(S, C_2^r(V, F^\bullet))$$

([1]) which readily generalizes to elliptic complexes.

is a quasi-isomorphism, with S assumed compact. By (3.2.1.1), we have

$$C(S, C^\infty(V, F^\bullet)) = \varprojlim_r C(S, C_2^r(V, F^\bullet))$$

For each i, $C(S, C_2^r(V, F^i))$ is a Banach space (for the topology of uniform convergence), and the transition morphisms

$$C(S, C_2^p(V, F^i)) \longrightarrow C(S, C_2^q(V, F^i))$$

are, by (3.0.4) and (3.1.2), continuous linear maps with dense image. On the other hand, applying to the quasi-isomorphisms (a') $\underline{F}_p^\bullet \longrightarrow \underline{F}_q^\bullet$ the functor Γ_S, which is exact since $\underline{O}_S$ is soft, we see that the transition morphisms

$$C(S, C_2^p(V, F^\bullet)) \longrightarrow C(S, C_2^q(V, F^\bullet))$$

are quasi-isomorphisms. Assertion (a) then follows from (EGA 0_{III} 13.2.3, "topological" variant). This completes the proof of (3.2.1).

3.2.2. For a topological space S, we shall denote by $\mathrm{Parf}(S)$ the triangulated category of perfect complexes of $(\underline{O}_S \otimes \underline{\underline{C}})$-Modules, and by $K^\bullet(S)$ the corresponding "Grothendieck group"([1]).

Let S, X, $E^\bullet$ be as in (3.2.1); the analytic index of $E^\bullet$ is the element

$$\mathrm{ind.\,an}(E^\bullet) = C\ell(\underline{\underline{R}}p_*(E^\bullet)) \in K^\bullet(S).$$

[print 218]For compact S, $\mathrm{ind.\,an}(E^\bullet)$ canonically defines an element in the usual group $K(S)$ of topologists. Indeed, in general, for every topological space S, denote by $\mathrm{Loclib}(S)$ the additive category of locally free $(\underline{O}_S \otimes \underline{\underline{C}})$-Modules of finite type, and by $K(S)$ the "Grothendieck group" of the triangulated category $\overline{K^b}(\mathrm{Loclib}(S))/K^{b,\varnothing}(\mathrm{Loclib}(S))$ (the latter is, moreover, identified with $K^b(\mathrm{Loclib}(S))$ when S is paracompact (II 2.3.2 (a)), so that in this case (I 6.4) $K(S)$ is canonically isomorphic to the "Grothendieck group" of the additive category $\mathrm{Loclib}(S)$). There is a natural homomorphism

$$K(S) \longrightarrow K^\bullet(S).$$

But, by (II 2.3.2), this is an isomorphism for compact S, proving our assertion. More generally, $\mathrm{ind.\,an}(E^\bullet)$ defines an element in the completed group

$$\widehat{K}(S) = \varprojlim K(A),$$

the projective limit being taken over the compact subsets A of S.

3.2.3. The writer does not know whether the finiteness theorem (3.2.1) remains valid when, instead of assuming S locally compact, one assumes only that every point of S has a fundamental system of paracompact neighborhoods. Moreover, the local compactness hypothesis on S does not seem very natural. The space of "universal local parameters" for elliptic complexes of fixed order is indeed an (infinite-dimensional) subspace of a Fréchet space. Let us make this precise. Let V be a compact variety, $F = (F^i)_{i \in \underline{\underline{Z}}}$ a sequence of complex vector bundles of finite rank on V, all but finitely many zero, and $\underline{n} = (n_i)$ a sequence of integers ≥ 0. Denote by $\mathrm{Comp.\,diff}^{\underline{n}}(F)$ the space of complexes of differential operators "with base F" and of order $\underline{n}$, i.e. the closed subspace of $\prod \mathrm{Diff}^{n_i}(F^i, F^{i+1})$ defined by the equations $d^{i+1}d^i = 0$. (the space $\prod \mathrm{Diff}^{n_i}(F^i, F^{i+1})$ is a Fréchet space). The subset [print 219]$\mathrm{Ell}^{\underline{n}}(F)$ of $\mathrm{Comp.\,diff}^{\underline{n}}(F)$ formed by the elliptic complexes of order $\underline{n}$ is open; it is this space (which in general is not locally compact) that plays the role of a space of universal local parameters.

([1]) see (I 6.3). For a general review of the functors K, see Exp. IV.

3.2.4. Nevertheless, even should the generalization of (3.2.1) mentioned above prove impossible, we can always associate to $E^\bullet$, under the sole hypothesis that the fibration p is proper and locally trivial, an element of $K^\bullet(S)$ deserving the name analytic index of $E^\bullet$. Indeed, suppose (for simplicity([1])) that we are in the "local" situation where $X = S \times V$ (V, a compact variety), with $E^\bullet$ given by a continuous map from S to $\mathrm{Ell}^{\underline{n}}(F)$ (3.2.3). Assertions (a') and (b) in the proof of (3.2.1) are in fact valid without any hypothesis on S: it is therefore reasonable to put, with the notation of (loc.cit.):

$$\text{ind. an}(E^\bullet) = C\ell(\underline{F}_r^\bullet) \in K^\bullet(S),$$

this element being independent of r.

3.2.5. Suppose S is paracompact, and denote by H a complex Hilbert space of infinite Hilbert dimension, and by $I(H)$ the subspace of $L(H,H)$ formed by the operators "with index". Then the complexes $F_r^\bullet$ define [12] a canonical element in the group $[S, I(H)]$ of homotopy classes of continuous maps from S to $I(H)$, but the writer does not know, for noncompact S (even when S is locally compact), the relations between the groups $[S, I(H)]$ and[157] $K^\bullet(S)$ (is there a natural arrow from one to the other? are they isomorphic?), and therefore a fortiori between the two indices thus defined.

For a generalization, under the hypotheses of (3.2.1), of the Atiyah–Singer index theorem, we refer to the recent work of Shih [17], or to work in preparation by M. F. Atiyah and I. M. Singer.

[print 220]

BIBLIOGRAPHY

[1] Artin, M., Grothendieck, A., and Verdier, J-L., Cohomologie étale des schémas, to appear in North Holl. Pub. Comp., (cited as SGA 4).

[2] Atiyah, M. and Segal, G., secret papers.

[3] Cartan, H. Funzioni e varietà complesse, Faisceaux analytiques cohérents, Course at the Centro Internazionale Matematico Estivo, 1964, Roma.

[4] Cartan, H., and Schwartz, L., Théorème d'Atiyah–Singer, Seminar 1963-64, Institut Henri Poincaré, Paris.

[5] Frisch, J., Fonctions analytiques sur un ensemble semi-analytique[158], note in CRAS Paris, t. 260, p. 2974-2976 (1965).

[6] Godement, R., Théorie des Faisceaux, Hermann (1958), (cited as T.F).

[7] Grothendieck, A., Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux, Séminaire de Géométrie Algébrique de l'IHES 1962, to appear in North Holl. Pub. Comp., (cited as SGA 2).

[8] Grothendieck, A., Fondements de la Géométrie Analytique, Séminaire Cartan 1959-60, Institut Henri Poincaré, Paris.

[9] Hartshorne, R., Residues and Duality, Lecture Notes, Springer n° 20, cited as [H].

[10] Houzel, C., Géométrie Analytique Locale, Séminaire Cartan 1959-60, Institut Henri Poincaré, Paris.

([1]) for the general case, use the notion of a "weak Hilbert bundle" or continuous field of Hilbert spaces from [12].

[11] Illusie, L., Complexes de fibrés et définitions du foncteur K, Séminaire Shih Weishu, IHES 1964-65.

[12] Illusie, L., Complexes quasi-acycliques directs de fibrés banachiques, note in CRAS Paris, t. 260, p. 6499-6502.

[13] Kiehl, R., Theorem A und Theorem B in der nichtarchimedischen Funktionentheorie, Invent. math. 2, p. 256-273 (1967). [print 221]

[14] Lang, S., Introduction to differentiable manifolds,

[15] Łojasiewicz, S., Ann. della Scuola Norm. Sup. Pisa, serie III, 18, fasc. IV, 1964.

[16] Palais, R., Lectures at the Princeton Seminar 1963-64 on the index theorem.

[17] Shih, W., Fiber cobordism and the index of a family of elliptic differential operators,

[18] Verdier, J.-L., Les catégories dérivées des catégories abéliennes, to appear in North Holl. Pub. Comp.

[print 222]

EXPOSÉ III
RELATIVE FINITENESS CONDITIONS

by L. ILLUSIE

1. Relative pseudo-coherence

Theorem 1.1. Let $n \in \mathbb{Z}$. For each locally finite type morphism of schemes $f : X \longrightarrow Y$, there exists one and only one full subcategory $D(f)_{n\text{-coh}}$ of $D(X)$([1]) such that the following conditions are satisfied:

(i) If $E \in \mathrm{Ob}\, D(f)_{n\text{-coh}}$, then $i^*(E) \in \mathrm{Ob}\, D(fi)_{n\text{-coh}}$ for every open immersion $i : U \to X$.

(ii) If $E \in \mathrm{ob}\, D(X)$ and there exists an open covering $(i_\alpha : U_\alpha \to X)_{\alpha\in A}$ such that $i_\alpha^*(E) \in \mathrm{ob}\, D(fi_\alpha)_{n\text{-coh}}$ for every $\alpha \in A$, then $E \in \mathrm{ob}\, D(f)_{n\text{-coh}}$.

(iii) If

$$\begin{array}{ccc} X & \xrightarrow{\quad f \quad} & Y \\ & {\scriptstyle h}\searrow \quad \swarrow{\scriptstyle g} & \\ & Z & \end{array}$$

is a commutative triangle of locally finite type morphisms, where f is a closed immersion and g a smooth morphism, then we have

$$E \in \mathrm{Ob}\, D(h)_{n\text{-coh}} \iff f_*(E) \in D(Y)_{n\text{-coh}} \qquad ([2]) \ .$$

Proof. Uniqueness is evident. Indeed, let $f : X \to Y$ be a locally finite type morphism. Choose an open covering $(i_\alpha : X_\alpha \to X)_{\alpha\in A}$ such that, for each $\alpha \in A$, fi_α factors as

$$X_\alpha \xrightarrow{i'_\alpha} Y_\alpha \xrightarrow{f'_\alpha} Y,$$

where i'_α is a closed immersion and f'_α a smooth morphism. It follows at once from (i), (ii), (iii) that the objects of $D(f)_{n\text{-coh}}$ are the objects E of $D(X)$ such that

$$i'_{\alpha*}i_\alpha^*(E) \in \mathrm{ob}\, D(Y_\alpha)_{n\text{-coh}};$$


thus $D(f)_{n\text{-coh}}$ [print 223] is indeed determined (since it is a full subcategory of $D(X)$). To prove existence, we shall need a few lemmas.

Lemma 1.1.1. Let $i : X \to Y$ be a closed immersion of schemes such that $i_*(\underline{O}_X)$ is pseudo-coherent, and let $E \in \mathrm{ob}\, D(X)$. For E to be n-pseudo-coherent, it is necessary and sufficient that $i_*(E)$ be so.


Proof. The "necessity" follows from the general criterion for stability of pseudo-coherence under application of a functor (I 2.13). Conversely, suppose that $i_*(E)$ is n-pseudo-coherent, and prove

([1]) $D(X)$ = derived category of the category of $\underline{O}_X$-Modules.
([2]) $D(Y)_{n\text{-coh}}$ = full subcategory of $D(Y)$ formed by the n-pseudo-coherent objects (I 2.15).

that the same is true of E. Since $i_*(E)$ has locally bounded-above cohomology, the same is true of E; thus, the question being local, we may suppose that E is m-pseudo-coherent for sufficiently large m. Let us show, by descending induction on m, that E is m-pseudo-coherent for every $m \geq n$. We may suppose it already proved that E is $(n+1)$-pseudo-coherent. After localizing, we therefore have a distinguished triangle



where E' is perfect and $E''^i = 0$ for $i > n$. By (I 2.6), we are reduced to showing that E'' is n-pseudo-coherent. Now, applying i_* to the preceding triangle, we obtain a distinguished triangle



By the first part of the proof, $i_*(E')$ is pseudo-coherent, so by (I 2.6), $i_*(E'')$ is n-pseudo-coherent. Since $i_*(E'')$ is zero in degree $> n$, this means, by (I 2.10 b)), that $H^n(i_*(E''))$[159] is of finite type. Since $H^n(i_*(E'')) \simeq i_*H^n(E'')$, we are reduced to showing, by (I 2.10 b)), that if M is an $\underline{O}_X$-Module such that $i_*(M)$ is of finite type, then M is of finite type, which is immediate (and independent of the hypothesis on $i_*(\underline{O}_X)$[160]).

[print 224] **Remark 1.1.2.** The preceding lemma applies in particular when i is a regular immersion (VII 1.4). Indeed, by means of the Koszul complex[161], we have locally a finite left resolution of $i_*(\underline{O}_X)$ by free $\underline{O}_Y$-Modules of finite type, so $i_*(\underline{O}_X)$ is perfect (I 4.8 and 4.14) (and therefore a fortiori pseudo-coherent).

Lemma 1.1.3. Let there be a Cartesian square of schemes

$$\begin{array}{ccc} X & \xleftarrow{f} & X' \\ {\scriptstyle i}\downarrow & & \downarrow{\scriptstyle i'} \\ Y & \xleftarrow{g} & Y', \end{array}$$

where i is a closed immersion, and let E be a complex of $\underline{O}_X$-Modules. Then the canonical base-change arrow (EGA III 1.4.15.4)

$$g^*i_*(E) \longrightarrow i'_*f^*(E)$$

is an isomorphism.

(ii) Put $E' = f^*(E)$. If g is flat and $i_*(E)$ n-pseudo-coherent, then $i'_*(E')$ is n-pseudo-coherent. Conversely, if g is faithfully flat and $i'_*(E')$ n-pseudo-coherent, then $i_*(E)$ is n-pseudo-coherent.

Proof. The assertion is checked trivially on the stalks.

[print 225] **Lemma 1.1.4.** Let there be a commutative diagram of schemes

$$\begin{array}{ccccc} X' & \xleftarrow{i'} & X & \xrightarrow{i''} & X'' \\ & {\scriptstyle f'}\searrow & & \swarrow{\scriptstyle f''} & \\ & & Y, & & \end{array}$$

where i' (resp. i'') is a closed immersion, and f' (resp. f'') a smooth morphism, and let $E \in \operatorname{ob} D(X)$. We then have:

$$i'_*(E) \text{ } n\text{-pseudo-coherent} \iff i''_*(E) \text{ } n\text{-pseudo-coherent}^{[162]}.$$

Proof. Following a device due to Lichtenbaum, we send X diagonally into $X' \times_Y X''$, and are at once reduced to proving the assertion when one of the morphisms f', f'' is the identity. Changing notation, we thus have a commutative triangle

$$\begin{array}{ccc} & & Y \\ & \overset{i}{\nearrow} & \downarrow f \\ X & \xrightarrow[j]{} & Z, \end{array} \qquad (*)$$

where i and j are closed immersions and f a smooth morphism, and we must show that "$i_*(E)$ n-pseudo-coherent" is equivalent to "$j_*(E)$ n-pseudo-coherent". Since the question is local in Y and in Z near a point of X, we may suppose that X, Y, Z are affine and that we have a factorization

$$\begin{array}{ccc} Y & \xrightarrow{i'} & Z[T_1, \ldots, T_r] \\ {\scriptstyle f}\downarrow & \swarrow_{f'} & \\ Z, & & \end{array}$$

where i' is a closed immersion, and the oblique arrow the canonical projection. Then (VII 1.10) i' is a regular immersion, so (1.1.1) an object F of $D(Y)$ is n-pseudo-coherent if and only if $i'_*(F)$ is so.

[print 226] We may therefore suppose that $Y = Z[T_1, \ldots, T_r]$, with f the canonical projection. The diagram $(*)$ then corresponds to a diagram of rings

$$\begin{array}{ccc} & & B = C[T_1, \ldots, T_r] \\ & \overset{u}{\swarrow} & \uparrow \varepsilon \\ A & \xleftarrow[v]{} & C \end{array} \quad ,$$

where u and v are epimorphisms, and ε the canonical inclusion. For $1 \leq i \leq r$, put $u(T_i) = x_i$, and let $z_i \in C$ satisfy $v(z_i) = x_i$. The retraction $\sigma : B \longrightarrow C$ of ε defined by $\sigma(T_i) = z_i$ satisfies $v\sigma = u$. In geometric language, it corresponds to a section s of f such that $i = sj$. By (VII 1.10), s is a regular immersion, so the assertion follows from (1.1.1).

Proof of (1.1)(end). Let $f : X \to Y$ be a locally finite type morphism. There exists an open covering (X_α) of X such that each X_α embeds by a closed immersion into a scheme Y_α smooth over Y. Putting $\coprod X_\alpha = X'$, $\coprod Y_\alpha = Y'$, we thus have a commutative diagram [163]

$$\begin{array}{ccc} X' & \xrightarrow{i'} & Y' \\ {\scriptstyle p'}\downarrow & & \\ X & \overset{f'}{\searrow} & \\ {\scriptstyle f}\downarrow & & \\ Y & & \end{array} \qquad (a')$$

where p' is a morphism that is the sum of a surjective family of open immersions (we shall say special for short), i' a closed immersion, and f' a smooth morphism. We shall define $D(f)_{n\text{-coh}}$ as the full subcategory of $D(X)$ formed by the objects E such that $i'_* p'^*(E)$ is n-pseudo-coherent.

Let us first show that this definition makes sense, i.e. that $D(f)_{n\text{-coh}}$ is independent of the choice of i', p', f'. Thus, let [print 227]

$$\begin{array}{ccc} X'' & \xrightarrow{i''} & Y'' \\ {\scriptstyle p''}\downarrow & \swarrow{\scriptstyle f''} & \\ X & & \\ {\scriptstyle f}\downarrow & & \\ Y & & \end{array} \tag{a''}$$

be a commutative diagram analogous to (a') (p'' a special morphism, i'' a closed immersion, f'' a smooth morphism). From (a') and (a''), we deduce a commutative diagram

$$\begin{array}{ccccc} X' \times_X X'' & \hookrightarrow & Y' \times_Y Y'' & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ & X' & \xhookrightarrow{i'} & & Y' \\ & \downarrow & & & \downarrow \\ X'' & \xhookrightarrow{i''} & Y'' & & {\scriptstyle f'}\downarrow \\ & \searrow & & \searrow & \\ & & X & \xrightarrow{f} & Y, \end{array}$$

where the horizontal arrows, except for f, are closed immersions, and the arrows of the left (resp. right) Cartesian square are special morphisms (resp. smooth morphisms). We are therefore reduced to proving the following assertion:

(A): Let

$$\begin{array}{ccc} X_1 & \xhookrightarrow{i_1} & Y_1 \\ {\scriptstyle p}\downarrow & & \downarrow{\scriptstyle q} \\ X_2 & \xhookrightarrow{i_2} & Y_2 \end{array}$$

be a commutative square of schemes, where p is a special morphism, q a smooth morphism, i_1 and i_2 closed immersions, and let $E \in \operatorname{ob} D(X_2)$. Then, for $i_{2*}(E)$ to be n-pseudo-coherent, it is necessary and sufficient that $i_{1*}p^*(E)$ be so.

[print 228] Proof of (A). By definition, p is the sum of a family of open subsets U_α of X_2 that cover X_2. Each U_α is induced by an open subset V_α of Y_2, and the V_α cover an open neighborhood X_2' of X_2 in Y_2. We therefore have a commutative diagram with a Cartesian square

$$\begin{array}{ccccc} X_1 & \xhookrightarrow{i_1'} & X_1' & & \\ {\scriptstyle p}\downarrow & & \downarrow{\scriptstyle p'} & & \\ X_2 & \xhookrightarrow{i_2'} & X_2' & \xhookrightarrow{j} & Y_2, \\ & \searrow^{i_2} & & \nearrow & \end{array}$$

where i_1' and i_2' are closed immersions, p and p' special morphisms, and j an open immersion. The condition "$i_{2*}(E)$ n-pseudo-coherent" is clearly equivalent to "$i'_{2*}(E)$ n-pseudo-coherent". On the other hand, it follows at once from (1.1.3) that the condition "$i'_{2*}(E)$ n-pseudo-coherent" is equivalent to "$i'_{1*}p^*(E)$ n-pseudo-coherent". Applying (1.1.4) to the factorizations $(jp')i_1'$ and qi_1 of i_2p, and to $p^*(E) \in \operatorname{ob} D(X_1)$, we finally conclude that "$i_{2*}(E)$ n-pseudo-coherent" is equivalent to "$i_{1*}p^*(E)$ n-pseudo-coherent", q.e.d.

We have thus shown that the category $D(f)_{n\text{-coh}}$ defined above by means of the diagram (a') in fact depends only on f. Moreover, it is trivial from the definition that it does satisfy conditions (i) to (iii) of (1.1). This therefore completes the proof of (1.1).

Definition 1.2. Let $f : X \to Y$ be a locally finite type morphism, $E \in \operatorname{ob} D(X)$, $n \in \mathbb{Z}$. We shall say that E is n-pseudo-coherent relative to f if $E \in \operatorname{ob} D(f)_{n\text{-coh}}$ (cf. (1.1)). We shall say that E is pseudo-coherent (or equivalently $(-\infty)$-pseudo-coherent) relative to f if E is n-pseudo-coherent relative to f for every $n \in \mathbb{Z}$. We shall write $D(f)_{\text{coh}}$ for the full subcategory [print 229] of $D(X)$ formed by the objects pseudo-coherent relative to f. We shall say that f is n-pseudo-coherent (resp. pseudo-coherent) if $\underline{O}_X$ is n-pseudo-coherent (resp. pseudo-coherent) relative to f. We shall sometimes say, when this cannot lead to confusion, "n-pseudo-coherent (resp. pseudo-coherent) relative to Y" instead of "relative to f", and shall also write $D(X)_{Y\text{-}n\text{-coh}}$ (resp. $D(X)_{Y\text{-coh}}$) instead of $D(f)_{n\text{-coh}}$ (resp. $D(f)_{\text{coh}}$). Finally, for $* = +, -, b$, we shall put

$$D^*(f)_{n\text{-coh}} = D^*(X) \cap D(f)_{n\text{-coh}},$$
$$D^*(f)_{\text{coh}} = D^*(X) \cap D(f)_{\text{coh}}.$$

Remarks 1.2.1.


a) The categories $D(f)_{n\text{-coh}}$, $D(f)_{\text{coh}}$ are not in general contained in $D^-(X)$. This is so, however, when X is quasi-compact, since an object of $D(X)$ with locally bounded-above cohomology then has bounded-above cohomology.

b) The objects of $D(f)_{\text{coh}}$ have quasi-coherent cohomology. Indeed, let $E \in \operatorname{ob} D(f)_{\text{coh}}$; the question being local, we may suppose that f is a closed immersion. Then $f_*(E)$ is pseudo-coherent, hence has quasi-coherent cohomology, and the result follows from the fact that an $\underline{O}_X$-Module M is quasi-coherent if and only if $f_*(M)$ is so.

c) When speaking of an $\underline{O}_X$-Module, we shall use the terminology "of finite presentation rel. to f" in preference to "(-1)-pseudo-coherent rel. to f". For an $\underline{O}_X$-Module E to be of finite presentation rel. to f, it is necessary and sufficient, in view of (I 2.9), that there exist locally a closed immersion $i : X \to X'$, with X' smooth over Y, such that $i_*(E)$ is of finite presentation (in the ordinary sense). In particular, saying that f is [print 230] (-1)-pseudo-coherent is equivalent to saying that f is locally of finite presentation in the sense of (EGA IV 1.4).

Examples 1.3. A smooth morphism and a regular immersion (1.1.2) are pseudo-coherent morphisms. The same holds for any locally finite type morphism $f : X \to Y$ when Y is locally noetherian.

Proposition 1.4. Let $f : X \to Y$ be a locally finite type morphism, and let



be a distinguished triangle in $D(X)$. We then have the following table, where $n \in \mathbb{Z} \cup \{-\infty\}$ indicates a degree of pseudo-coherence relative to f, and each row represents an implication whose conclusion is underlined:

E	F	G
$n+1$	n	$\underline{n}$
n	$\underline{n}$	n
$\underline{n}$	n	$n-1$.

In particular, $D(f)_{\mathrm{coh}}$ is a triangulated subcategory of $D(X)$.

Proof. By properties (i) to (iii) of (1.1), we may suppose that f is a closed immersion. We then have a distinguished triangle

$$\begin{array}{ccc} & f_*(G) & \\ \swarrow & & \nwarrow \\ f_*(E) & \longrightarrow & f_*(F) \end{array},$$

and the assertion follows from (I 2.6).

Definition 1.5. Let X and Y be two S-schemes. We shall say that X and Y are tor-independent over S if we have

$$\mathrm{Tor}_i^S(\underline{O}_X, \underline{O}_Y)^{[164]} = 0 \quad \text{for } i \neq 0,$$

where the Tor_i^S are the local hyper-tor defined in (EGA III 6.5). [print 231] This is a condition of a local nature; when X, Y, S are affine with respective rings B, C, A, it means that the canonical arrow

$$B \otimes_A^L C \longrightarrow B \otimes_A C$$

is an isomorphism. It is therefore satisfied in particular when X or Y is flat over S.

Lemma 1.5.1. Let there be Cartesian squares of schemes

$$\begin{array}{ccccc} X & \longleftarrow & X' & \longleftarrow & X'' \\ \downarrow & & \downarrow & & \downarrow \\ Y & \longleftarrow & Y' & \longleftarrow & Y''. \end{array}$$

If X and Y' are tor-independent over Y, and if X' and Y'' are tor-independent over Y', then X and Y'' are tor-independent over Y.

Proof. An easy exercise on associativity of the tensor product.

Notation 1.6. Let X and Y be two S-schemes, $E \in \mathrm{ob}\, D^-(X)$, $F \in \mathrm{ob}\, D^-(Y)$. We shall put

$$E \otimes_S^L F = L\,\mathrm{pr}_1^*(E) \otimes^L L\,\mathrm{pr}_2^*(F),$$

where $\mathrm{pr}_1, \mathrm{pr}_2$ are the two projections from $X \times_S Y$.

Lemma 1.7. Let S be a scheme, and, for $i = 1, 2$, $f_i : X_i \longrightarrow Y_i$ be a closed S-immersion. Suppose that X_1 and X_2, as well as Y_1 and Y_2, are tor-independent over S. Then, for $E_i \in \mathrm{ob}\, D^-(X_i)_{\mathrm{qcoh}}$([1]), there exists a canonical isomorphism

$$f_{1*}(E_1) \otimes_S^L f_{2*}(E_2) \xrightarrow{\sim} (f_1 \times_S f_2)_*(E_1 \otimes_S^L E_2).$$

Proof. Put $f_1 \times_S f_2 = f$. By adjunction, one easily defines a canonical arrow [print 232]

$$f_{1*}(E_1) \otimes_S^L f_{2*}(E_2) \longrightarrow f_*(E_1 \otimes_S^L E_2).$$

Let us show that it is an isomorphism. The question being local, we may suppose that S, X_i, Y_i are affine with respective rings C, A_i, B_i. By dévissage, we reduce to $E_i = \underline{O}_{X_i}$. We must then check that

$$A_1 \otimes_{B_1}^L B \otimes_{B_2}^L A_2 \longrightarrow A$$

([1]) $D(X_i)_{\mathrm{qcoh}}$ denotes the full subcategory of $D(X_i)$ formed by complexes with quasi-coherent cohomology.

is an isomorphism (we have put $A = A_1 \otimes_C A_2$, $B = B_1 \otimes_C B_2$). But, taking account of the tor-independence hypotheses, this follows at once from associativity of the tensor product.

Proposition 1.8. Let S be a scheme, and, for $i = 1, 2$, $f_i : X_i \longrightarrow Y_i$ be a locally finite type S-morphism. Suppose X_1 and X_2, as well as Y_1 and Y_2, tor-independent over S. Let $[a_i, b_i]$ be an interval of $\mathbb{Z}$, and $E_i \in \operatorname{ob} D^-(X_i)_{\mathrm{qcoh}}$ a complex that is a_i-pseudo-coherent rel. to f_i and acyclic in degree $> b_i$. Then $E_1 \otimes_S^L E_2$ is a-pseudo-coherent rel. to f and acyclic in degree $> b$, with $f = f_1 \times_S f_2$, $a = \operatorname{Sup}(a_1 + b_2, a_2 + b_1)$, $b = b_1 + b_2$.

Proof. The question being local on $X_1 \times_S X_2$, we may suppose that f_i factors as

$$X_i \xrightarrow{f_i'} Y_i' \xrightarrow{p_i} Y_i,$$

where f_i' is a closed immersion and p_i a smooth morphism. Putting $f' = f_1' \times_S f_2'$, we must show that $f_{1*}'(E_1 \otimes_S^L E_2)$ is a-pseudo-coherent and acyclic in degree $> b$. By (1.5.1), Y_1' and Y_2' are tor-independent over S: after changing notation, we may therefore suppose that $f_i = f_i'$, in other words that f_i is a closed immersion. By hypothesis, $f_{i*}(E_i)$ is a_i-pseudo-coherent and acyclic in degree $> b_i$, so (I 2.16) $f_{1*}(E_1) \otimes_S^L f_{2*}(E_2)$ is a-pseudo-coherent and acyclic in degree $> b$. We conclude by (1.7).

[print 233] **Corollary 1.9.** Under the hypotheses of (1.8), the functor

$$\otimes_S^L : D^-(X_1) \times D^-(X_2) \longrightarrow D^-(X)$$

induces a functor

$$\otimes_S^L : D^-(f_1)_{\mathrm{coh}} \times D^-(f_2)_{\mathrm{coh}} \longrightarrow D^-(f)_{\mathrm{coh}} \qquad \text{(cf. (1.2))}.$$

In particular, if f_1 and f_2 are pseudo-coherent (1.2), the same holds for f.

Corollary 1.10. Let

$$\begin{array}{ccc} X & \xleftarrow{g} & X' \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle f'} \\ Y & \xleftarrow{h} & Y' \end{array}$$

be a Cartesian square, with f locally of finite type. Then:

a) Suppose X and Y' tor-independent over Y (1.5). For $n \in \mathbb{Z} \cup \{-\infty\}$, the functor $Lg^* : D^-(X) \to D^-(X')$ induces a functor

$$Lg^* : D^-(f)_{n\text{-}\mathrm{coh}} \longrightarrow D^-(f')_{n\text{-}\mathrm{coh}}.$$

In particular, if f is n-pseudo-coherent (1.2), the same holds for f'.

b) Suppose h faithfully flat, let $n \in \mathbb{Z} \cup \{-\infty\}$ and $E \in \operatorname{ob} D^-(X)$ be a complex with quasi-coherent cohomology. For E to be n-pseudo-coherent rel. to f, it is necessary and sufficient that $g^*(E)$ be n-pseudo-coherent relative to f'. In particular, for f to be n-pseudo-coherent, it is necessary and sufficient that f' be so.

Proof. Taking account of (1.2.1 b)), all the assertions of (1.9) and (1.10) follow trivially from (1.8), except the "sufficiency" part of (1.10 b)). For this, the question being local on X, we may suppose that f factors as a closed immersion followed by a smooth morphism. By (1.1.3), we are then reduced to the case where f (resp. f') is the identity of X (resp. X'). The assertion then follows, by a standard dévissage (cf. I 2.16.2), from the fact that a quasi-coherent sheaf L on X is of finite type if and only if g^*L is of finite type.

[print 234] **Proposition 1.11.** Let [165]

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow^{h} & \downarrow{\scriptstyle g} \\ & & Z \end{array}$$

be a commutative triangle of locally finite type morphisms, let $E \in \operatorname{ob} D(X)$, $F \in \operatorname{ob} D(Y)$, and $[a,b]$, $[c,d]$ be intervals of $\mathbb{Z}$.

a) If E is a-pseudo-coherent rel. to f and acyclic in degree $> b$ and if F is c-pseudo-coherent rel. to g and acyclic in degree $> d$, then $E \otimes^L_Y F$ (cf. (1.6)) is m-pseudo-coherent rel. to h and acyclic in degree $> n$, where $m = \operatorname{Sup}(a+d, b+c)$, and $n = b+d$.

b) If E is a-pseudo-coherent rel. to h and acyclic in degree $> b$, and if g is $(a+1-b)$-pseudo-coherent, then E is a-pseudo-coherent rel. to f.

Proof. The question being local on X, we may suppose that we have a commutative diagram with a Cartesian square

$$\begin{array}{ccccc} & & & & E_Z^{r+s} \\ & & & \overset{j}{\nearrow} & \\ & & E_Y^r & & \searrow \\ & \overset{i}{\nearrow} & \downarrow{\scriptstyle p} & & \downarrow{\scriptstyle q} \\ X & \xrightarrow{f} & Y & \xrightarrow{g} & Z, \end{array} \qquad \text{with } E_Y^r \xrightarrow{k} E_Z^s$$

where i, j, k are closed immersions, and p, q the canonical projections of the standard affine spaces. If F is c-pseudo-coherent rel. to Z, then $k_*(F)$ is c-pseudo-coherent by (1.1 (iii)), and $j_*(p^*(F))$ is c-pseudo-coherent by (1.1.3). Similarly, if g is x-pseudo-coherent, then $j_*(\underline{O}_{E_Y^r})$ is x-pseudo-coherent. By the definition of relative pseudo-coherence (1.2), we are therefore reduced to proving a) and b) when f and g are closed immersions.

[print 235] Proof of a). By (1.1 (iii)), we must show the following: suppose $f_*(E)$ is a-pseudo-coherent and acyclic in degree $> b$, and $g_*(F)$ is c-pseudo-coherent and acyclic in degree $> d$; then $h_*(E \otimes^L_Y F)$ is m-pseudo-coherent and acyclic in degree $> n$. Now we have a canonical isomorphism

$$f_*(E) \otimes^L F \xrightarrow{\sim} f_*(E \otimes^L_Y F)$$

(a special case of the isomorphism in (1.7)[166]). Since the functor $g_*(. \otimes^L F)$ takes locally free sheaves of finite type to c-pseudo-coherent complexes, and complexes acyclic in degree > 0 to complexes acyclic in degree $> d$, the assertion therefore follows from (I 2.13).

Proof of b). We must prove the following assertion: suppose $h_*(E)$[167] is a-pseudo-coherent and acyclic in degree $> b$, and $g_*(\underline{O}_Y)$ is $(a+1-b)$-pseudo-coherent; then $f_*(E)$ is a-pseudo-coherent. Since the assertion concerns $f_*(E)$, we may suppose $X = Y$ and $f = \mathrm{Id}_X$. We then proceed as in the proof of (1.1.1), proving by descending induction that E is pseudo-coherent in degree $\geq a$. We suppose it already proved that E is $(a+1)$-pseudo-coherent, whence (I 2.10), after localizing, a distinguished triangle

$$\begin{array}{ccccc} & & E'' & & \\ & \swarrow & & \nwarrow & \\ E' & & \longrightarrow & & E, \end{array}$$

where E'^i is locally free for $i \in [a+1, b]$, $E'^i = 0$ for $i \notin [a+1, b]$, and $E''^i = 0$ for $i > a$. The hypothesis that $g_*(\underline{O}_Y)$ is $(a+1-b)$-pseudo-coherent then implies (I 2.11 a)) that $g_*(E')$ is $(a+1)$-pseudo-coherent. Since $g_*(E)$ is a-pseudo-coherent, it follows from (I 2.6) that $g_*(E'')$ is a-pseudo-coherent. We conclude by (I 2.10 b)) and the fact that an $\underline{O}_Y$-Module M is of finite type if and only if $g_*(M)$ is so.

The following corollaries are deduced trivially from (1.11):

[print 236] **Corollary 1.12.** In the situation of (1.11), if g is pseudo-coherent, we have, for $r \in \mathbb{Z} \cup \{-\infty\}$,

$$D^-(f)_{r\text{-coh}} = D^-(h)_{r\text{-coh}};$$

in particular (taking $X = Y$), the notions of r-pseudo-coherence over Y, "in the absolute sense" or rel. to g, coincide.

Corollary 1.13. In the situation of (1.11), if f is pseudo-coherent, the functor $\underline{O}_X \otimes^{\mathrm{L}}_Y \cdot = Lf^* : D^-(Y) \longrightarrow D^-(X)$ induces, for $r \in \mathbb{Z} \cup \{-\infty\}$, a functor

$$Lf^* : D^-(g)_{r\text{-coh}} \longrightarrow D^-(h)_{r\text{-coh}}.$$

Corollary 1.14. In the situation of (1.11), if f is r-pseudo-coherent and g is s-pseudo-coherent $(r, s \in \mathbb{Z} \cup \{-\infty\})$, then h is $\mathrm{Sup}(r, s)$-pseudo-coherent. If h is r-pseudo-coherent and g is $(r+1)$-pseudo-coherent, f is r-pseudo-coherent.

Remark 1.14.1. In particular, for $r = s = -1$, we recover (cf. (1.2.1 c))) assertions (ii) and (v) of (EGA IV 1.4.3).

2. The finiteness theorem.

Conjecture 2.1. Let S be a scheme, $f : X \to Y$ be a proper S-morphism of schemes locally of finite type over S, $E \in \mathrm{ob}\, D^+(X)$. Then, if E is pseudo-coherent rel. to S (1.2), the same holds for $Rf_*(E)$.

Theorem 2.2. The preceding conjecture is true in the following cases:

(i) S is locally noetherian;

(ii) f is projective.

[print 237] Proof in case (i). Since S is locally noetherian, the projection of X (resp. Y) onto S is a pseudo-coherent morphism (1.3), and the notions of pseudo-coherence over X (resp. Y) and relative to S coincide (1.12). On the other hand, since X (resp. Y) is locally noetherian, hence $\underline{O}_X$ (resp. $\underline{O}_Y$) coherent, an object M of $D(X)$ (resp. $D(Y)$) is pseudo-coherent if and only if (I 3.5) the $H^i(M)$ are coherent and M is locally in D^-. The theorem therefore follows from the following one:

Theorem 2.2.1. Let $f : X \to Y$ be a proper morphism of locally noetherian schemes, and let $E \in \mathrm{ob}\, D^+(X)$. If the $H^i(E)$ are coherent, the same holds for the $R^i f_*(E)$. If, moreover, E is locally in D^-, the same holds for $Rf_*(E)$.


Proof. The first assertion follows from the spectral sequence

$$R^p f_*(H^q(E)) \Longrightarrow R^* f_*(E) \qquad (*)$$

and the finiteness theorem (EGA III 3.2.1). For the second, we may suppose Y affine; X is then quasi-compact and separated. Suppose it covered by r affine open subsets. We know (EGA III

1.4.12) that, under these conditions, $R^i f_*(M) = 0$ for every quasi-coherent $\underline{O}_X$-Module M and every $i > r$. If E is locally in D^-, E is in D^-, since X is quasi-compact; say $H^i(E) = 0$ for $i > N$. From the spectral sequence (*) we then deduce that $R^i f_*(E) = 0$ for $i > N + r$, whence the assertion.

Proof in case (ii). The question being local on Y, we may suppose S and Y affine. We then have a commutative diagram with a Cartesian square



[print 238] where i, j, j' are closed immersions, g' a smooth affine morphism, and p, p' the projections of the standard projective spaces. Suppose E pseudo-coherent rel. to S. By (1.1 (iii)), this means that $(ji)_*(E)$ is pseudo-coherent. We must show that $Rf_*(E)$ is pseudo-coherent rel. to S, i.e. (1.1 (iii)) that $j'_* Rf_*(E)$ is pseudo-coherent. But we have

$$j'_* Rf_*(E) \simeq Rp'_*(ji)_*(E),$$

so the assertion will follow from the following more precise result:

Theorem 2.2.2. Let Y be an affine scheme, $f : X = P^r_Y \longrightarrow Y$ be the projection of a standard projective space, $E \in \operatorname{ob} D^b(X)_{\mathrm{coh}}$, $a \in \mathbb{Z}$. Then $Rf_*(E)$ is pseudo-coherent. Moreover, if E is acyclic in degree $> a$, $Rf_*(E)$ is acyclic in degree $> a + r$, and there exists an integer N such that, for every $n \geq N$, $Rf_*(E(n))$ (where $E(n) = E \otimes \underline{O}_X(n)$) is acyclic in degree $> a$.

Proof. Since Y is affine, X is quasi-compact and separated, so (II 2.2.9 a)) E is isomorphic, in $D(X)$, to a complex F bounded above in degree such that, for every i, $F^i \in \operatorname{ob} \underline{L}(X)$, i.e. (loc. cit.) is a finite direct sum of sheaves of the type $\underline{O}_X(-n)$ $(n > 0)$. Moreover, if E is acyclic in degree $> a$, we may choose F so that $F^i = 0$ for $i > a$: this follows formally from the preceding assertion, taking account of (I 2.10 a) and II 2.2.8 a)). Replacing E by F, we may therefore suppose that $E^i \in \operatorname{ob} \underline{L}(X)$ for every i, and $E^i = 0$ for $i > a$. Since X is covered by $(r + 1)$ affine open subsets, we have, by (EGA III 1.4.10) and the hypercohomology spectral sequence, $R^q f_*(E) = 0$ for $q > a + r$[168]. Let t be an integer $\leq a$, and put $E' = (0 \to E^t \to \ldots \to E^a \to 0)$. We have an exact sequence

$$0 \longrightarrow E' \longrightarrow E \longrightarrow E'' \longrightarrow 0,$$

[print 239] where E'' satisfies the same conditions as E, but with a replaced by $t - 1$. Applying Rf_*, we obtain a distinguished triangle

$$\begin{array}{ccc} & Rf_*(E'') & \\ \swarrow & & \nwarrow \\ Rf_*(E') & \longrightarrow & Rf_*(E) \end{array} \qquad (*)$$

Since t can be chosen arbitrarily far in the negative direction, it suffices, for the first assertion, to show that $Rf_*(E)$ is $(t + r)$-pseudo-coherent. Now $Rf_*(E'')$ is acyclic in degree $\geq t + r$, hence $(t + r)$-pseudo-coherent. Thus, by (I 2.6), the result will follow once we prove that $Rf_*(E')$ is perfect. For this, we reduce by dévissage (I 4.10) to the case where E' is concentrated in degree

0 and equal to $\underline{O}_X(-n)$, $n > 0$. Then, by (EGA III 2.1.16), $R^r f_*(\underline{O}_X(-n))$ is locally free of finite type, and we have

$$Rf_*(\underline{O}_X(-n)) \simeq R^r f_*(\underline{O}_X(-n))[-r].$$

Thus $Rf_*(\underline{O}_X(-n))$ is perfect, and we deduce the first assertion of (2.2.2). We have already seen that $Rf_*(E)$ is acyclic in degree $> a + r$. It remains to prove the last assertion. For $t < -r + a$, $Rf_*(E'')$ is acyclic in degree $> a$, so, taking account of the triangle $(*)$, we are reduced to proving the result for E' in place of E. Since E' is bounded in degree and its components are objects of $\underline{L}(X)$, there exists an integer N such that, for $n \geq N$, the components of $E'(n)$ are finite direct sums of sheaves of the type $\underline{O}_X(p)$, with $p \geq 0$. Then, by (EGA III 2.1.16) and the hypercohomology spectral sequence, $Rf_*(E'(n))$ is acyclic in degree $> a$ for every $n \geq N$, which completes the proof of (2.2.2), and hence of (2.2).

We have a "classical" complement to (2.2 (ii)), which we shall give for later reference:

[print 240] **Corollary 2.3** (EGA III 2.2.1). Let S be a quasi-compact scheme, $f : X \to Y$ be a projective S-morphism of schemes of finite type over S, $L = i^*\underline{O}_P(1)$ be the invertible sheaf defined by an embedding $i : X \to P = P(F)$, where F is a quasi-coherent $\underline{O}_Y$-Module of finite type. For $M \in \operatorname{ob} D(X)$ and $n \in \mathbb{Z}$, put $M(n) = M \otimes^{\mathrm{L}} L^{\otimes n}$. Then, if $E \in \operatorname{ob} D^+(X)$ is pseudo-coherent rel. to S and acyclic in degree $> a$ ($a \in \mathbb{Z}$), there exists an integer N such that, for every $n \geq N$, $Rf_*E(n)$ satisfies the same conditions.

Proof. We reduce trivially, as in the proof of (2.2 (ii)), to the situation of (2.2.2): $Y = S$, $f : X = P^r_Y \to Y$ the canonical projection, $L = \underline{O}_X(1)$ the canonical sheaf, $E \in \operatorname{ob} D^b(X)$ a pseudo-coherent complex acyclic in degree $> a$, and the result follows by (2.2.2).

Corollary 2.4. Let S be a quasi-compact scheme, and $f : X \to Y$ be a proper S-morphism of schemes of finite type over S. If S is noetherian or f is projective, the functor $Rf_* : D^+(X) \longrightarrow D^+(Y)$ induces a functor

$$Rf_* : D^b(X)_{S\text{-coh}} \longrightarrow D^b(Y)_{S\text{-coh}}$$

Proof. A consequence of (2.2) and the fact that there is then (EGA III 1.4.12 and IV 1.7.21) an integer N such that $R^i f_*(E) = 0$ for every $i > N$ and every quasi-coherent $\underline{O}_X$-Module E.

Corollary 2.5. Let $f : X \to Y$ be a proper and pseudo-coherent morphism (1.2), and $E \in \operatorname{ob} D^+(X)$. Suppose Y locally noetherian or f projective. Then, if E is pseudo-coherent (in the absolute sense), the same holds for $Rf_*(E)$. If Y is quasi-compact, the functor Rf_* induces a functor

$$Rf_* : D^b(X)_{\mathrm{coh}} \longrightarrow D^b(Y)_{\mathrm{coh}}.$$

[print 241]Proof. A consequence of (2.2), (2.4) and (1.12)[169].

Scholium 2.6. To prove conjecture (2.1) in the general case, it will no doubt be necessary to introduce new techniques. One may hope that these will one day help to prove the analogous conjecture in Analytic Geometry, for a proper morphism of analytic spaces over a locally ringed topos in suitably topologized $\mathbb{C}$-algebras, a conjecture that is at present "proved" only over a complex analytic space (Grauert's theorem [1]).([1])

3. Relative finite Tor-dimension

Definition 3.1. Let $f : (X, \underline{O}_X) \longrightarrow (Y, \underline{O}_Y)$ be a morphism of ringed topoi, $E \in \operatorname{ob} D^-(X)$, $[m, n]$ an interval of $\mathbb{Z}$. We say that E is of flat amplitude relative to f (or Y, when no confusion

([1]) (added in April 1971) see an article in preparation by R. Kiehl and J.L. Verdier, in which finiteness theorem 1 is proved by different, simpler techniques and under more general hypotheses.

is possible) contained in $[m,n]$, and write

$$\operatorname{tor.amp}_f(E) \subset [m,n] \qquad (\text{or } \operatorname{tor.amp}_Y(E) \subset [m,n])$$

if E, considered as a complex of $f^{-1}(\underline{O}_Y)$-Modules, is of flat amplitude contained in $[m,n]$ (I 5.2). We say that f is of flat amplitude contained in $[-a,0]$ $(a \geq 0)$, or of tor-dimension $\leq a$, if $\underline{O}_X$ is of flat amplitude rel. to f, contained in $[-a,0]$. We say that E is of finite flat amplitude rel. to f if there is an interval $[m,n]$ such that $\operatorname{tor.amp}_f(E) \subset [m,n]$.

Notation 3.2. We denote by $D(f)_{\text{torf}}$, or $D(X)_{f\text{-torf}}$, or $D(X)_{Y\text{-torf}}$ when no confusion is possible, the full subcategory of $D^-(X)$ formed by the complexes of finite flat amplitude relative to f. By (I 5.3), this is a triangulated subcategory of $D^b(X)$.

[print 242]**Proposition 3.3.** With the data of (3.1), suppose that X has enough points (SGA 4 IV 6.4)[170]. The following conditions are then equivalent:

(i) $\operatorname{tor.amp}_f(E) \subset [m,n]$;

(ii) for every point x of X, we have

$$\operatorname{tor.amp}_y(E_x) \subset [m,n],$$

where y is the image point of x under f, ringed by $\underline{O}_{Y,y}$;

(iii) for every object V of Y and every $\underline{O}_V$-Module M, we have

$$H^i\left((E|f^{-1}(V)) \otimes^{\mathrm{L}}_V M\right) = 0 \quad \text{for } i \notin [m,n],$$

where $(E|f^{-1}(V)) \otimes^{\mathrm{L}}_V M = (E|f^{-1}(V)) \otimes^{\mathrm{L}} \mathrm{L}f^*(M)$;

(iv) the same condition as (iii), but with M the cokernel of a homomorphism $\underline{O}_V^q \longrightarrow \underline{O}_V^p$, where $p, q \in \mathbb{N}$.

Proof. The equivalence (i)$\Longleftrightarrow$(ii) follows from (I 5.6.2), and the implications (ii)$\Longrightarrow$(iii)$\Longrightarrow$(iv) are trivial. Let us prove (iv)$\Longrightarrow$(ii). Let x be a point of X, $y = f(x)$; we must show that

$$H^i(E_x \otimes^{\mathrm{L}}_{\underline{O}_{Y,y}} M) = 0 \qquad (*)$$


for every $\underline{O}_{Y,y}$-module M and every $i \notin [m,n]$. Since every $\underline{O}_{Y,y}$-module is a filtered inductive limit of modules of finite presentation, we may restrict ourselves to checking the preceding relation for M of finite presentation. Now, when M is an $\underline{O}_{Y,y}$-module of finite presentation, the same argument as for ordinary ringed spaces (EGA 0_{I} 5.3.9) shows that there is a y-pointed object V of Y and an $\underline{O}_V$-Module P, the cokernel of a homomorphism $\underline{O}_V^q \longrightarrow \underline{O}_V^p$ $(p, q \in \mathbb{N})$, such that $P_y \simeq M$. We then have

$$\left((E|f^{-1}(V)) \otimes^{\mathrm{L}}_V P\right)_x \simeq E_x \otimes^{\mathrm{L}}_{\underline{O}_{Y,y}} M,$$

[print 243]by the commutation of tensor product with passage to stalks, and hypothesis (iv) implies (*), q.e.d.

Remark 3.3.1. The writer does not know whether the equivalences (i)$\Longleftrightarrow$(iii)$\Longleftrightarrow$(iv) hold without the hypothesis that X has enough points.

Corollary 3.4. Let



be a commutative triangle of ringed topoi, where we suppose that X has enough points, and let $E \in \mathrm{ob}\, D(X)$, $F \in \mathrm{ob}\, D(Y)$, $[a,b]$, $[c,d]$ be intervals of $\mathbb{Z}$. Then we have:

$$\mathrm{tor.amp}_f(E) \subset [a,b] \quad \text{and} \quad \mathrm{tor.amp}_g(F) \subset [c,d]$$

$$\Longrightarrow \mathrm{tor.amp}_h(E \otimes^{\mathrm{L}}_Y F) \subset [a+c, b+d].$$

Proof. By (3.3), we must show that the relations on the left imply

$$H^i\left(((E \otimes^{\mathrm{L}}_Y F)|h^{-1}(Z')) \otimes^{\mathrm{L}}_{Z'} M\right) = 0$$

for every $i \notin [a+c, b+d]$ and every $\underline{O}_{Z'}$-Module M, where Z' is an object of Z. Now, putting $Y' = g^{-1}(Z')$, $X' = f^{-1}(Y') = h^{-1}(Z')$, we have

$$((E \otimes^{\mathrm{L}}_Y F)|X') \otimes^{\mathrm{L}}_{Z'} M \simeq (E|X') \otimes^{\mathrm{L}}_{Y'} ((F|Y') \otimes^{\mathrm{L}}_{Z'} M),$$

and the assertion follows at once from the hypotheses.

In particular:

Corollary 3.4.1. Under the hypotheses of (3.4),

a) if g is of tor-dimension $\leq n$, we have

$$\mathrm{tor.amp}_f(E) \subset [a,b] \Longrightarrow \mathrm{tor.amp}_h(E) \subset [a-n, b];$$

[print 244]

b) if f is of tor-dimension $\leq m$, we have

$$\mathrm{tor.amp}_g(F) \subset [c,d] \Longrightarrow \mathrm{tor.amp}_h(\mathrm{L}f^*(F)) \subset [c-m, d].$$

Remark 3.4.2. Let $f : X \to Y$ be a morphism of finite tor-dimension and $E \in \mathrm{ob}\, D(X)$. Then, by (3.4.1 a)), when E is of finite flat amplitude (in the absolute sense), E is of finite flat amplitude rel. to f. But the converse is false, even when f is flat, as the example of a singularity of a scheme over a field shows.

Proposition 3.5. Let S be a scheme, and, for $i = 1, 2$, $f_i : X_i \longrightarrow Y_i$ an S-morphism, $E_i \in \mathrm{ob}\, D^-(X_i)$, $[a_i, b_i]$ an interval of $\mathbb{Z}$. Suppose X_1 and X_2, as well as Y_1 and Y_2, are tor-independent over S (1.5). Put $a = a_1 + a_2$, $b = b_1 + b_2$, $f = f_1 \times_S f_2 : X \to Y$. Then:

a) If we have $\mathrm{tor.amp}_{f_1}(E_1) \subset [a_1, b_1]$ and if E_2 is acyclic outside $[a_2, b_2]$, $E_1 \otimes^{\mathrm{L}}_S E_2$ (cf. (1.6)) is acyclic outside $[a,b]$.

b) If we have $\mathrm{tor.amp}_{f_i}(E_i) \subset [a_i, b_i]$ for $i = 1, 2$, we have

$$\mathrm{tor.amp}_f(E_1 \otimes^{\mathrm{L}}_S E_2) \subset [a,b].$$

Proof. By (3.3), it suffices to make the verification "on the stalks". Let s be a point of S, x a point of X over s, $y = f(x)$, x_i (resp. y_i) the canonical projection of x (resp. y) onto X_i (resp. Y_i). Put

$$(E_i)_{x_i} = M_i, \quad \underline{O}_{X_i,x_i} = A_i, \quad \underline{O}_{Y_i,y_i} = B_i, \quad \underline{O}_{S,s} = C,$$
$$\underline{O}_{X,x} = A = A_1 \otimes_C A_2, \qquad \underline{O}_{Y,y} = B = B_1 \otimes_C B_2.$$

We have

$$(E_1 \otimes^{\mathrm{L}}_S E_2)_x \simeq M_1 \otimes^{\mathrm{L}}_{A_1} A \otimes^{\mathrm{L}}_{A_2} M_2. \qquad (*)$$

The tor-independence hypotheses imply, on the other hand:

$$A \simeq A_1 \otimes^{\mathrm{L}}_{B_1} B \otimes^{\mathrm{L}}_{B_2} A_2.$$

Consequently, the right-hand side of (*) is canonically isomorphic, in the [print 245]derived category $D(B)$ of B-modules, to $M_1 \otimes^{\mathrm{L}}_{B_1} B \otimes^{\mathrm{L}}_{B_2} M_2$. The relations $\mathrm{tor.amp}_{f_i}(E_i) \subset [a_i, b_i]$ $(i = 1, 2)$ imply $\mathrm{tor.amp}_{B_i}(M_i) \subset [a_i, b_i]$, hence (I 5.6) $\mathrm{tor.amp}_B(M_1 \otimes^{\mathrm{L}}_{B_1} B \otimes^{\mathrm{L}}_{B_2} M_2) \subset [a, b]$, giving assertion b). On the other hand, when we have $\mathrm{tor.amp}_{f_1}(E_1) \subset [a_1, b_1]$, M_1 is isomorphic to a complex of flat B_1-modules, zero outside $[a_1, b_1]$, and consequently when E_2 (hence M_2) is acyclic outside $[a_2, b_2]$, $M_1 \otimes^{\mathrm{L}}_{B_1} B \otimes^{\mathrm{L}}_{B_2} M_2$ is acyclic outside $[a, b]$, giving assertion a), which completes the proof.

Corollary 3.5.1. Under the hypotheses of (3.5), if f_1 and f_2 are of finite tor-dimension (3.1), the same holds for f.

Corollary 3.5.2. Let

$$\begin{array}{ccc} X & \xleftarrow{g} & X' \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle f'} \\ Y & \xleftarrow{h} & Y' \end{array}$$

be a Cartesian square of schemes, where X and Y' are supposed tor-independent over Y, and let $E \in \mathrm{ob}\, D(X)$. Then we have

$$\mathrm{tor.amp}_f(E) \subset [a, b] \Longrightarrow \mathrm{tor.amp}_{f'}(\mathrm{L}g^*(E)) \subset [a, b],$$

and consequently the functor $\mathrm{L}g^* : D^-(X) \to D^-(X')$ induces a functor

$$\mathrm{L}g^* : D(f)_{\mathrm{torf}} \longrightarrow D(f')_{\mathrm{torf}}.$$

[print 246]**Proposition 3.6.** Let

$$\begin{array}{ccc} X & \xhookrightarrow{i} & X' \\ {\scriptstyle f}\downarrow & \swarrow{\scriptstyle f'} & \\ Y & & \end{array}$$

be a commutative triangle of schemes, where i is a closed immersion and f' a smooth morphism, and let $E \in \mathrm{ob}\, D(X)$. For E to be locally of finite flat amplitude rel. to f, it is necessary and sufficient that $i_*(E)$ be locally of finite flat amplitude (in the absolute sense). More precisely, if f' is of relative dimension $\leq d$, we have

(i) $\mathrm{tor.amp}(i_*(E)) \subset [a, b] \Longrightarrow \mathrm{tor.amp}_f(E) \subset [a, b]$,

(ii) $\mathrm{tor.amp}_f(E) \subset [a, b] \Longrightarrow \mathrm{tor.amp}(i_*(E)) \subset [a - d, b]$.

[print 247]Proof. For an interval $[m, n]$ of $\mathbb{Z}$, we evidently have

$$\mathrm{tor.amp}_f(E) \subset [m, n] \Longleftrightarrow \mathrm{tor.amp}_{f'}(i_*(E)) \subset [m, n],$$

so that we may suppose $X = X'$, $i = \mathrm{Id}_X$, $f = f'$. Relation (i) follows trivially from (3.4.1 a)). Let us prove (ii). We have a Cartesian square

$$\begin{array}{ccc} X & \xleftarrow{\mathrm{pr}_2} & X \times_Y X \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle \mathrm{pr}_1} \\ Y & \xleftarrow{f} & X \end{array},$$

and the diagonal morphism $s : X \longrightarrow X \times_Y X$ is a section of pr_1 and pr_2. Suppose f is smooth of relative dimension d. Then (VII 1.10) s is a regular immersion of codimension $\leq d$, hence is of tor-dimension $\leq d$ (as is seen using the Koszul complex). By (3.5.2), we have

$$\mathrm{tor.amp}_f(E) \subset [a,b] \Longrightarrow \mathrm{tor.amp}_{\mathrm{pr}_1}(\mathrm{pr}_2^*(E)) \subset [a,b],$$

and by (3.4.1 b)) we have

$$\mathrm{tor.amp}_{\mathrm{pr}_1}(\mathrm{pr}_2^*(E)) \subset [a,b] \Longrightarrow \mathrm{tor.amp}(\mathrm{L}s^* \mathrm{pr}_2^*(E)) \subset [a-d,b],$$

giving implication (ii), since $\mathrm{L}s^* \mathrm{pr}_2^* \simeq \mathrm{Id}$.

Remark 3.6.1. Statement (3.6) would be false with only the assumption that f' is flat; see (3.4.2).

Proposition 3.7. ("Projection formula")([1]). Let S be a quasi-compact scheme, $f : X \to Y$ a quasi-compact and quasi-separated S-morphism of quasi-compact schemes over S, $E \in \mathrm{ob}\, D^b(X)_{\mathrm{qcoh}}$, $M \in \mathrm{ob}\, D^b(S)_{\mathrm{qcoh}}$([2]). If M is of finite flat amplitude, or if E is of finite flat amplitude rel. to S, there [print 248]is a canonical functorial isomorphism in $D(Y)_{\mathrm{qcoh}}$

$$Rf_*(E) \otimes^{\mathrm{L}}_S M \xrightarrow{\sim} Rf_*(E \otimes^{\mathrm{L}}_S M) \qquad \text{(cf. (1.6))}.$$

Proof. First note that there is (EGA III 1.4.12 and IV 1.7.21) an integer N such that $R^i f_*(F) = 0$ for every $i > N$ and every quasi-coherent $\underline{O}_X$-Module F. Consequently the functor Rf_* sends $D^b(X)_{\mathrm{qcoh}}$ into $D^b(Y)$ (and even $D^b(Y)_{\mathrm{qcoh}}$ (EGA III 1.4.10 and IV 1.7.21)). Suppose E (resp. M) is of finite flat amplitude rel. to S (resp. finite flat amplitude).[171] The adjunction arrow $Lf^*Rf_*(E) \longrightarrow E$ defines, by tensoring with M, a canonical arrow of $D^-(X)$

$$Lf^*(Rf_*(E) \otimes^{\mathrm{L}}_S M) \longrightarrow E \otimes^{\mathrm{L}}_S M,$$

where $E \otimes^{\mathrm{L}}_S M \in \mathrm{ob}\, D^b(X)$. By adjunction we deduce a canonical arrow of $D^-(Y)$

$$Rf_*(E) \otimes^{\mathrm{L}}_S M \longrightarrow Rf_*(E \otimes^{\mathrm{L}}_S M). \qquad (*)$$

Its target and source moreover have quasi-coherent cohomology, since Rf_* sends $D^b(X)_{\mathrm{qcoh}}$ into $D^b(Y)_{\mathrm{qcoh}}$ (EGA III 1.4.10 and IV 1.7.21), and $\otimes^{\mathrm{L}}_S$ sends $D^-(X)_{\mathrm{qcoh}} \times D^-(S)_{\mathrm{qcoh}}$ into $D^-(X)_{\mathrm{qcoh}}$ (likewise with Y in place of X). Let us prove that $(*)$ is an isomorphism. The question being local on Y, we may suppose S and Y affine. By dévissage, in the first case we immediately reduce to the case where M is an ordinary quasi-coherent $\underline{O}_S$-Module. Noting that the functors $Rf_*(E) \otimes^{\mathrm{L}}_S .$ and $Rf_*(E \otimes^{\mathrm{L}}_S .)$ are "way-out to the left" on the category $D^b(S)_{\mathrm{qcoh}}$ ([H] I 7), we then reduce, by taking a left resolution of M by free Modules and truncating, to the case where M is free. Since the $R^i f_*$ commute with inductive limits (f being quasi-compact and quasi-separated), we may even suppose M free of finite type, and then $M = \underline{O}_S$. The arrow $(*)$ is then tautologically an isomorphism. In the second case, by dévissage we reduce to $M = P$, where P is a flat A-module [print 249]$(S = \mathrm{Spec}(A))$. Putting $A' = D_A(P)$ (EGA 0_{IV} 18.2.3), $S' = \mathrm{Spec}(A')$, everything reduces to proving that $Rf_*(E)$ commutes with the flat base change $S' \to S$, i.e. that the canonical arrow

$$q^* Rf_*(E) \longrightarrow Rf'_* p^*(E)$$

is an isomorphism (we have put $f_{(S')} = f'$ and denoted by p (resp. q) the canonical projection $X_{(S')} \longrightarrow X$ (resp. $Y_{(S')} \longrightarrow Y$)). But this follows from (EGA III 1.4.15 and IV 1.7.21). This completes the proof of (3.7).

([1]) cf. (H II 5.6).

([2]) The subscript "qcoh" indicates the full subcategory formed by complexes with quasi-coherent cohomology.

Corollary 3.7.1. Let S be a scheme, $f : X \to Y$ a quasi-compact and quasi-separated S-morphism of quasi-compact schemes over S, $E \in \mathrm{ob}\, D^b(X)_{\mathrm{qcoh}}$. If E is locally of finite flat amplitude rel. to S, the same holds for $Rf_*(E)$. More precisely, if S is quasi-compact, and if N is an integer such that $R^i f_*(F) = 0$ for every $i > N$ and every quasi-coherent $\underline{O}_X$-Module F (EGA III 1.4.12 and IV 1.7.21), then we have

$$\mathrm{tor.amp}_S(E) \subset [a, b] \Longrightarrow \mathrm{tor.amp}_S(Rf_*(E)) \subset [a, b + N].$$

Proof. We may evidently restrict ourselves to proving the last assertion. By (3.3 (i)$\Longleftrightarrow$(iv)), and taking into account that Rf_* "commutes with restriction to an open subset of S", we must prove that the hypothesis $\mathrm{tor.amp}_S(E) \subset [a, b]$ implies

$$H^i(Rf_*(E) \otimes^{\mathrm{L}}_S M) = 0$$

for every $i \notin [a, b + N]$ and every quasi-coherent $\underline{O}_S$-Module M. Now the preceding hypothesis means that $H^i(E \otimes^{\mathrm{L}}_S M) = 0$ for every $i \notin [a, b]$ and every quasi-coherent $\underline{O}_S$-Module M, and (3.7) gives the result.

[print 250]**Corollary 3.7.2.** Let $f : X \to Y$ be a quasi-compact, quasi-separated morphism, locally of finite tor-dimension (3.1), and let $E \in \mathrm{ob}\, D^b(X)_{\mathrm{qcoh}}$. If E is locally of finite flat amplitude (in the absolute sense), the same holds for $Rf_*(E)$.

Proof. Combine (3.4.1 a)) and (3.7.1).

4. Relative perfection

Definition 4.1. Let $f : X \to Y$ be a locally finite type morphism of schemes, $E \in \mathrm{ob}\, D(X)$, $[a, b]$ an interval of $\mathbb{Z}$. We say that E is of perfect amplitude relative to f (or Y when there is no ambiguity), contained in $[a, b]$, and write

$$\mathrm{parf.amp}_f(E) \subset [a, b] \qquad (\text{or } \mathrm{parf.amp}_Y(E) \subset [a, b])$$

if, relative to f, E is pseudo-coherent (1.2) and of flat amplitude contained in $[a, b]$ (3.1). We say that f is of perfect amplitude contained in $[-n, 0]$, or of perfect dimension $\leq n$ ($n \geq 0$), if $\underline{O}_X$ is of perfect amplitude contained in $[-n, 0]$. We say that E is of finite perfect amplitude relative to f if there is an interval $[a, b]$ such that $\mathrm{parf.amp}_f(E) \subset [a, b]$. We say that E is perfect rel. to f if E is locally of finite perfect amplitude rel. to f. We say that f is perfect if f is locally of finite perfect amplitude.

Example 4.1.1. A regular immersion of codimension d (VII 1.4) is of perfect dimension $\leq d$ (as is seen from the Koszul complex). A locally complete-intersection morphism (i.e. locally a composite [print 251]of a regular immersion and a smooth morphism) is a perfect morphism.[172]

Notation 4.2. Let $f : X \to Y$ be a locally finite type morphism of schemes. We denote by $D(f)_{\mathrm{parf}}$, or $D(X)_{f\text{-}\mathrm{parf}}$, or $D(X)_{Y\text{-}\mathrm{parf}}$ when no confusion is to be feared (resp. $D(f)_{\underline{\mathrm{parf}}}$, $D(X)_{f\text{-}\underline{\mathrm{parf}}}$, $D(X)_{Y\text{-}\underline{\mathrm{parf}}}$) the full subcategory of $D(X)$ formed by the complexes perfect (resp. of finite perfect amplitude) rel. to f. By (I 5.3) and (1.4), $D(f)_{\mathrm{parf}}$ and $D(f)_{\underline{\mathrm{parf}}}$ are triangulated subcategories of $D(X)$. When X is quasi-compact, we have $D(f)_{\mathrm{parf}} = D(f)_{\underline{\mathrm{parf}}}$.

Proposition 4.3 (cf. I 5.8). Let $f : X \to Y$ be a locally finite type morphism of schemes, $E \in \mathrm{ob}\, D(X)$, $[a, b]$ an interval of $\mathbb{Z}$, x a point of X, $y = f(x)$. The following conditions are equivalent:

(i) $\mathrm{parf.amp}_f(E) \subset [a, b]$ in a neighborhood of x:

(ii) E is, in a neighborhood of x, $(a-1)$-pseudo-coherent rel. to f and acyclic outside $[a,b]$, and we have $\mathrm{tor.amp}_{\underline{O}_{Y,y}}(E_x) \subset [a,b]$.

Proof. The implication (i) $\Longrightarrow$ (ii) is trivial; let us prove (ii) $\Longrightarrow$ (i). The question being local in a neighborhood of x, we immediately reduce, by the definitions, to the case where f is smooth. Then E is $(a-1)$-pseudo-coherent in a neighborhood of x; hence, after localizing, we may suppose E strictly $(a-1)$-pseudo-coherent (I 2.1) and zero in degree $> b$, and after truncating (i.e. replacing E by the complex $(0 \to E^a/B^a \to E^{a+1} \to \dots E^b \to 0)$), we may suppose that E^i is free of finite type for $i > a$, E^a of finite presentation, and $E^i = 0$ for $i \notin [a,b]$. By (I 5.1.1), E^a_x is flat rel. to $\underline{O}_{Y,y}$, hence (EGA IV 11.3.1) E^a is f-flat in a neighborhood of x. Write $E^a = F$ for brevity; we must show that F is pseudo-coherent (in a neighborhood of x). After embedding X in a standard affine space over Y, we may [print 252]suppose that $X = Y[t_1, \dots, t_r]$, with Y affine and f the canonical projection. Then Y is a projective limit of noetherian affine schemes Y_α. Denote by $f_\alpha : X_\alpha = Y_\alpha[t_1, \dots, t_r] \to Y_\alpha$ the canonical projection. There is (EGA IV 8.5.2 and 11.2.6) an index α and an $\underline{O}_{X_\alpha}$-Module F_α of finite presentation and f_α-flat such that $g_\alpha^*(F_\alpha) \simeq F$, where $g_\alpha : X \to X_\alpha$ is the canonical morphism. Since X_α is noetherian, F_α is coherent, hence pseudo-coherent rel. to f_α. Since, on the other hand, F_α is f_α-flat and f_α is flat, we have $g_\alpha^*(F_\alpha) \simeq F_\alpha{}^{[173]} \otimes^{\mathrm{L}}_{Y_\alpha} \underline{O}_Y$, and it follows from (1.10 a)) that F is pseudo-coherent rel. to Y, hence pseudo-coherent, q.e.d.

Corollary 4.3.1. Let $f : X \to Y$ be a locally finite type morphism, and E an $\underline{O}_X$-Module. For $\mathrm{parf.amp}_f(E) \subset [0,0]$ to hold, it is necessary and sufficient that E be, rel. to f, flat and of finite presentation (1.2.1 c)). In particular, for f to have perfect dimension zero (4.1), it is necessary and sufficient that f be flat and locally of finite presentation.

Proof. Apply (4.3 (ii) $\Longrightarrow$ (i)) with $a = b = 0$.

Corollary 4.3.2. Let $f : X \to Y$ be a locally finite type morphism, and $E \in \mathrm{ob}\, D(X)$ a complex of $\underline{O}_X$-Modules flat and of finite presentation rel. to f (1.2.1 c)), and $[a,b]$ an interval of $\mathbb{Z}$. If $E^i = 0$ for $i \notin [a,b]$, we have $\mathrm{parf.amp}_f(E) \subset [a,b]$. If E is bounded above in degree, E is pseudo-coherent rel. to f.

Proof. A consequence of (4.3.1) and (I 4.15 and 2.11).

Proposition 4.4. Let there be a commutative triangle of schemes

$$\begin{array}{ccc} X & \xrightarrow{\ i\ } & Y \\ {\scriptstyle f}\downarrow & \swarrow_{g} & \\ S & & \end{array} ,$$

[print 253]where i is a closed immersion and g a smooth morphism, and let $E \in \mathrm{ob}\, D(X)$. For E to be perfect rel. to f, it is necessary and sufficient that $i_*(E)$ be perfect (in the absolute sense (I 4.8)). More precisely, suppose g is of relative dimension $\leq d$, and let $[a,b]$ be an interval of $\mathbb{Z}$. Then we have:

(i) $\mathrm{parf.amp}(i_*(E)) \subset [a,b] \Longrightarrow \mathrm{parf.amp}_f(E) \subset [a,b]$[174];

(ii) $\mathrm{parf.amp}_f(E) \subset [a,b] \Longrightarrow \mathrm{parf.amp}(i_*(E)) \subset [a-d,b]$.

Proof. An immediate consequence of (3.6).

Proposition 4.5. Let

$$\begin{array}{ccc} X & \xrightarrow{\quad f\quad} & Y \\ & {\scriptstyle h}\searrow \quad \swarrow{\scriptstyle g} & \\ & S & \end{array}$$

be a commutative triangle of locally finite type morphisms of schemes, and let $E \in \operatorname{ob} D^-(X)$, $F \in \operatorname{ob} D^-(Y)$, $[a,b]$, $[c,d]$ be intervals of $\mathbb{Z}$. Then we have:

$$\operatorname{parf.amp}_f(E) \subset [a,b] \text{ and } \operatorname{parf.amp}_g(F) \subset [c,d] \Longrightarrow \operatorname{parf.amp}_h(E \otimes^{\mathrm{L}}_Y F) \subset [a+c, b+d].$$

Consequently, if E is perfect rel. to f and F perfect rel. to g, $E \otimes^{\mathrm{L}}_Y F$ is perfect rel. to h.

Proof. Combine (1.11) and (3.4).

Corollary 4.5.1. Under the hypotheses of (4.5):

a) if g has perfect dimension $\leq n$, we have

$$\operatorname{parf.amp}_f(E) \subset [a,b] \Longrightarrow \operatorname{parf.amp}_h(E) \subset [a-n, b];$$

consequently, if g is perfect, E perfect rel. to f implies E perfect rel. to h;

b) if f has perfect dimension $\leq m$, we have

$$\operatorname{parf.amp}_g(F) \subset [c,d] \Longrightarrow \operatorname{parf.amp}_h(\mathrm{L}f^*(F)) \subset [c-m, d];$$

[print 254]consequently, if f is perfect, F perfect rel. to g implies $\mathrm{L}f^*(F)$ perfect rel. to h.

Remark 4.5.2. With the preceding notation, when g is perfect, E perfect rel. to h does not necessarily imply E perfect rel. to f, as the ever-recurring example of a singular point of a scheme over a field shows. We have the following criterion:

Proposition 4.6 (criterion of perfection by fibers)(1). Let

$$\begin{array}{ccc} X & \xrightarrow{\quad f \quad} & Y \\ & {\scriptstyle h}\searrow \quad \swarrow{\scriptstyle g} & \\ & S & \end{array}$$

be a commutative triangle of morphisms of schemes. We suppose f, h locally of finite type, and g locally of finite presentation and flat. Let $x \in X$, $y = f(x)$, $s = h(x) = g(y)$. Finally, let $E \in \operatorname{ob} D^-(X)$, and let $[a,b]$ be an interval of $\mathbb{Z}$. We denote by $f_s : X_s \to Y_s$ the fiber of f at s, i.e. the morphism obtained from f by the base change $\operatorname{Spec}(k(s)) \to S$. Then the following conditions are equivalent:

(i) $\operatorname{parf.amp}_h(E) \subset [a,b]$ in a neighborhood of x, and

$$\operatorname{tor.amp}_{\underline{O}_{Y_s,y}}(E_x \otimes^{\mathrm{L}}_{\underline{O}_{Y,y}} \underline{O}_{Y_s,y}) \subset [a,b];$$

(ii) $\operatorname{parf.amp}_f(E) \subset [a,b]$ in a neighborhood of x.

Proof. (ii) $\Longrightarrow$ (i). We already know (4.3.1) that g has perfect dimension zero. Thus (4.5.1 a)), condition (ii) implies $\operatorname{parf.amp}_h(E) \subset [a,b]$ in a neighborhood of x. It also implies (by 4.3),[175] $\operatorname{tor.amp}_{\underline{O}_{Y,y}}(E_x) \subset [a,b]$, hence (I 5.1 (i)), $\operatorname{tor.amp}_{\underline{O}_{Y_s,y}}(E_x \otimes^{\mathrm{L}}_{\underline{O}_{Y,y}} \underline{O}_{Y_s,y}) \subset [a,b]$, which [print 255]completes the proof of (ii) $\Longrightarrow$ (i).

(i) $\Longrightarrow$ (ii). The question being local in a neighborhood of x, we may suppose that f factors as $X \overset{i}{\hookrightarrow} X' \xrightarrow{f'} Y$, where i is a closed immersion and f' a smooth morphism. It is immediate that conditions (i) and (ii) are unchanged when we replace E by $i_*(E)$, f by f' (and h by gf'),

(1) Analogous to the "criterion of flatness by fibers" (EGA IV 11.3.10).

so that we may suppose f smooth. Since g is perfect (4.3.1), h is perfect, hence (1.12) E is pseudo-coherent. After localizing, by (II 2.2.10 b)) and (I 2.10 a)), we may suppose that E is strictly pseudo-coherent (I 2.1) and zero in degree $> b$, and that the truncation arrow

$$E \longrightarrow (0 \longrightarrow E^a/B^a \longrightarrow E^{a+1} \longrightarrow \ldots \longrightarrow E^b \longrightarrow 0) \qquad t)$$

is a quasi-isomorphism (hypothesis (i) indeed implies that E is, in a neighborhood of x, acyclic outside[176] $[a,b]$). We must prove that E^a/B^a is f-flat in a neighborhood of x. Denote by $i : X_s \longrightarrow X$ the canonical arrow. We have:

$$E_x \otimes^{\mathrm{L}}_{\underline{O}_{Y,y}} \underline{O}_{Y_s,y} \simeq i^*(E)_x.$$

Since h is flat, hypothesis (i) implies, by (3.5.2), $\mathrm{tor.amp}_{h_s}(i^*(E)) \subset [a,b]$ in a neighborhood of x. Thus, after localizing, we may suppose[177] that the truncation arrow

$$i^*(E) \longrightarrow (0 \longrightarrow i^*(E^a)/B^a(i^*(E)) \longrightarrow \ldots \longrightarrow i^*(E^b) \longrightarrow 0) \qquad i^*(t)$$

is a quasi-isomorphism. The hypothesis $\mathrm{tor.amp}_{\underline{O}_{Y_s,y}} i^*(E)_x \subset [a,b]$ then implies, by (I 5.1.1), that $i^*(E^a/B^a)_x$ is a flat $\underline{O}_{Y_s,y}$-module. On the other hand, since the arrow (t)[178] is a quasi-isomorphism, the hypothesis $\mathrm{tor.amp}_h(E) \subset [a,b]$ implies, again by (I 5.1.1), [print 256]that E^a/B^a is h-flat. By the criterion of flatness by fibers (EGA IV 11.3.10), we conclude that E^a/B^a is f-flat in a neighborhood of x, q.e.d.

Corollary 4.6.1. Let $f : X \to S$ be a flat morphism locally of finite presentation, $x \in X$, $s = f(x)$, $E \in \mathrm{ob}\, D^-(X)$, $[a,b]$ an interval of $\mathbb{Z}$. Denote by $i_s : X_s = X \times_S \mathrm{Spec}(k(s)) \to X$ the canonical projection, and put $E_{X_s} = \mathrm{L}i_s^*(E)$. Then we have

$$\mathrm{parf.amp}_f(E) \subset [a,b] \text{ in a neighborhood of } x \text{ and } \mathrm{tor.amp}(E_{X_s})_x \subset [a,b]$$
$$\Longleftrightarrow \mathrm{parf.amp}(E) \subset [a,b] \text{ in a neighborhood of } x.$$

Consequently, E is perfect if and only if E is perfect rel. to f and E_{X_s} is perfect for every $s \in S$.

Proof. Apply (4.6) with $X = Y$, $f = \mathrm{Id}_X$.

Remark 4.6.2. From (4.6.1) and (I 5.10) it follows that, when f is locally of finite presentation, flat and has regular schemes as fibers, E perfect rel. to f is equivalent to E perfect (this equivalence had already been obtained (4.4) for f smooth).

Proposition 4.7. Let S be a scheme, and for $i = 1, 2$, $f_i : X_i \to Y_i$ an S-morphism locally of finite type, $E_i \in \mathrm{ob}\, D(X_i)$, $[a_i, b_i]$ an interval of $\mathbb{Z}$. Suppose X_1 and X_2, as well as Y_1 and Y_2, are tor-independent over S (1.5). Put $a = a_1 + a_2$, $b = b_1 + b_2$, $f = f_1 \times_S f_2$. Then we have:

a) If $\mathrm{parf.amp}_{f_1}(E_1) \subset [a_1, b_1]$, and if E_2 is pseudo-coherent rel. to f_2 and acyclic in degree outside $[a_2, b_2]$, $E_1 \otimes^{\mathrm{L}}_S E_2$ (cf. (1.6)) is pseudo-coherent rel. to f and acyclic in degree outside $[a,b]$. [print 257]

b) $\mathrm{parf.amp}_{f_i}(E_i) \subset [a_i, b_i]$ for $i = 1, 2$

$$\Longrightarrow \mathrm{parf.amp}_f(E_1 \otimes^{\mathrm{L}}_S E_2) \subset [a,b].$$

Proof. Combine (1.9) and (3.5).

Corollary 4.7.1. Under the hypotheses of (4.7)[179], if, for $i = 1, 2$, f_i has perfect dimension $\leq n_i$, f has perfect dimension $\leq n_1 + n_2$. Consequently, if f_1 and f_2 are perfect, the same holds for f.

Corollary 4.7.2. Let

$$\begin{array}{ccc} X & \xleftarrow{g} & X' \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle f'} \\ Y & \xleftarrow{h} & Y' \end{array}$$

be a Cartesian square of schemes, where f is locally of finite type and X and Y' are supposed tor-independent over Y, and let $E \in \operatorname{ob} D(X)$, and $[a,b]$ be an interval of $\mathbb{Z}$. We have:

$$\operatorname{parf.amp}_f(E) \subset [a,b] \Longrightarrow \operatorname{parf.amp}_{f'}(\mathrm{L}g^*(E)) \subset [a,b],$$

and consequently the functor $\mathrm{L}g^* : D^-(X) \to D^-(X')$ sends $D(f)_{\mathrm{parf}}$ (resp. $D(f)_{\mathrm{parf}}$) into $D(f')_{\mathrm{parf}}$ (resp. $D(f')_{\mathrm{parf}}$) (4.2).

Proposition 4.8. Let S be a scheme, $f : X \to Y$ a proper S-morphism of S-schemes locally of finite type, $E \in \operatorname{ob} D^b(X)$. Then, subject to the validity of (2.1) for f (hence in any case when S is locally noetherian or when f is projective):

a) If E is perfect rel. to S, the same holds for $Rf_*(E)$.

b) Suppose S quasi-compact, X and Y of finite type over S; let $[a,b]$ be an interval of $\mathbb{Z}$, and N an integer such that $R^i f_*(M) = 0$ for every $i > N$ and [print 258]every O_X-Module M that is quasi-coherent (such an integer exists by (EGA III 1.4.12 and IV 1.7.21)) ; we then have

$$\operatorname{parf.amp}_S(E) \subset [a,b] \Longrightarrow \operatorname{parf.amp}_S(Rf_*(E)) \subset [a, b+N].$$

Consequently, the functor Rf_* sends $D(X)_{S\text{-parf}}$ into $D(Y)_{S\text{-parf}}$(1).

Proof. Assertion a) follows from b), and b) is obtained by combining (2.1) and (3.7.1).

Corollary 4.8.1. Let $f : X \to Y$ be a proper and perfect morphism, and $E \in \operatorname{ob} D^+(X)$. Then, subject to the validity of (2.1) for f, if E is perfect, the same holds for $Rf_*(E)$; moreover, if Y is quasi-compact, the functor Rf_* sends $D(X)_{\mathrm{parf}}$ into $D(Y)_{\mathrm{parf}}$(2).

Proof. Indeed, when E is perfect, E is perfect rel. to f by (4.5.1 a)).

Proposition 4.9(3). Let S be a noetherian scheme, X and Y be S-schemes of finite type, $f : X \to Y$ a compactifiable S-morphism (i.e. a composite of an open immersion and a proper morphism), $E \in \operatorname{ob} D(X)$, $F \in \operatorname{ob} D^+(Y)$. If E is perfect rel. to f, and F perfect rel. to S, $\mathrm{R}\underline{\mathrm{Hom}}(E, f^!(F))$ is perfect rel. to S(4).

Proof. The question being local on X, we may suppose that f factors as $X \xrightarrow{i} X' \xrightarrow{f'} Y$, where i is a closed immersion and f' a smooth morphism. When E is perfect rel. to f, E is perfect rel. to i (4.4), and when F is perfect rel. to S, $f'^!(F)$ is perfect rel. to S by (4.5.1 b)), since $f'^!(F)$ is locally isomorphic to $f'^*(F)$ up to a shift. Since $f^! \simeq i^! f'^!$, we therefore reduce to proving the assertion when f is a closed immersion. Thus suppose E perfect rel. to f, i.e. $f_*(E)$ perfect, and F perfect rel. to S. We must show that $f_*\mathrm{R}\underline{\mathrm{Hom}}(E, f^!F)$ is [print 259]perfect rel. to S. But by the duality theorem ([H] III 6.7), we have

$$f_*\mathrm{R}\underline{\mathrm{Hom}}(E, f^!F) \simeq \mathrm{R}\underline{\mathrm{Hom}}(f_*(E), F).$$

(1) moreover, $D(X)_{S\text{-parf}} = D(X)_{S\text{-parf}}$ (and likewise with Y), since X and Y are quasi-compact (cf. (4.2)).

(2) We have $D(X)_{\mathrm{parf}} = D(X)_{\mathrm{parf}}$ (and likewise with Y).

(3) This result and its corollaries will not be used in the remainder of this Seminar, but play an important role in the Lefschetz–Verdier formula for coherent sheaves; see SGA 5 Exp. III.

(4) For the definition of the functor $f^!$, see [H] App. n° 4.

Since $f_*(E)$ is perfect, we have, by (I 7.7),

$$\mathrm{R}\underline{\mathrm{Hom}}(f_*(E), F) \simeq (f_*(E))^\vee \otimes^{\mathrm{L}} F,$$

and (4.5) (where $X = Y$ and $f = \mathrm{Id}_X$) gives the result.

Corollary 4.9.1. Under the hypotheses of (4.9)[180], if f is perfect, $f^!$ sends $D(Y)_{S\text{-parf}}$ into $D(X)_{S\text{-parf}}$.

Proof. Take $E = \underline{O}_X$.

Corollary 4.9.2. Let S be a noetherian scheme, $f : X \to S$ a compactifiable morphism. Put $f^!(\underline{O}_S) = K_X$, $\mathrm{R}\underline{\mathrm{Hom}}(., K_X) = \underline{D}_X$. For $E \in \mathrm{ob}\, D(f)_{\mathrm{parf}}$, we have $\underline{D}_X(E) \in \mathrm{ob}\, D(f)_{\mathrm{parf}}$, and the canonical arrow (I 7.2)

$$E \longrightarrow \underline{D}_X \underline{D}_X(E)$$

is an isomorphism.

Proof. For the first assertion, apply (4.9)[181] with $Y = S$, $F = \underline{O}_S$. For the second, we easily reduce to the case where f is a closed immersion. It then suffices to prove that the arrow obtained from the canonical arrow by applying f_* is an isomorphism. By global duality ([H] III 6.7), we have

$$f_* \underline{D}_X \simeq \underline{D}_S f_*.$$

We therefore reduce to showing that the canonical arrow

$$f_*(E) \longrightarrow \underline{D}_S \underline{D}_S f_*(E)$$

is an isomorphism for E perfect rel. to f, i.e. $f_*(E)$ perfect, which follows from ordinary biduality for perfect complexes (I 7.2).

5. Applications: exchange and semi-continuity theorems(1)

Lemma 5.1. Let $(X, \underline{O}_X)$ be a ringed topos (commutative), denote by $\mathrm{Mod}(X)$ the category of $\underline{O}_X$-Modules, $D(X) = D(\mathrm{Mod}(X))$. Let $E \in \mathrm{ob}\, D^-(X)$, and $p \in \mathbb{Z}$. Consider the following conditions: [print 260]

a) The functor $M \mapsto H^p(E \otimes^{\mathrm{L}} M)$ from $\mathrm{Mod}(X)$ to $\mathrm{Mod}(X)$ is right exact.

a') The canonical arrow

$$H^p(E) \otimes M \longrightarrow H^p(E \otimes^{\mathrm{L}} M)$$

is an isomorphism for every $M \in \mathrm{ob}\, \mathrm{Mod}(X)$.

b) The functor $M \mapsto H^{p+1}(E \otimes^{\mathrm{L}} M)$ from $\mathrm{Mod}(X)$ to $\mathrm{Mod}(X)$ is left exact.

b') The canonical arrow defined below

$$H^{p+1}(E \otimes^{\mathrm{L}} M) \longrightarrow \underline{\mathrm{Hom}}(H^{-p-1}(E^\vee), M) \qquad (*)$$

is an isomorphism for every $M \in \mathrm{ob}\, \mathrm{Mod}(X)$.

Then we have:

$$\mathrm{a}) \Longleftrightarrow \mathrm{a}') \Longleftrightarrow \mathrm{b}) \Longleftarrow \mathrm{b}').$$

Moreover, if E is pseudo-coherent, the conditions a), a'), b), b') are equivalent.

(1) The results of this section will not be used in the remainder of the Seminar.

Definition of $(*)$: The canonical arrow (I 7.5) $E^\vee \otimes^{\mathrm{L}} E \longrightarrow \underline{O}_X$, where $E^\vee = \mathrm{R}\underline{\mathrm{Hom}}(E, \underline{O}_X)$, gives, by tensoring with $M \in \mathrm{ob}\,\mathrm{Mod}(X)$, an arrow $E^\vee \otimes^{\mathrm{L}} E \otimes^{\mathrm{L}} M \longrightarrow M$; hence, by the adjunction formula dear to Cartan (I 7.4), a canonical arrow

$$E \otimes^{\mathrm{L}} M \longrightarrow \mathrm{R}\underline{\mathrm{Hom}}(E^\vee, M). \qquad (**)$$

Composing the arrow induced by $(**)$ on H^{p+1} with the canonical arrow

$$\underline{\mathrm{Ext}}^{p+1}(E^\vee, M) \longrightarrow \underline{\mathrm{Hom}}(H^{-p-1}(E^\vee), M), \qquad (***)$$

we obtain the arrow $(*)$.

Proof. The implications

$$\mathrm{a}') \Longrightarrow \mathrm{a}) \Longleftrightarrow \mathrm{b}) \Longleftarrow \mathrm{b}')$$

are trivial. Let us prove a) $\Longrightarrow$ a'). Let M be an $\underline{O}_X$-Module. There is an exact sequence $M^1 \longrightarrow M^0 \longrightarrow M \longrightarrow 0$, where M^0 and M^1 are direct sums of sheaves [print 261]of the type $\underline{O}_{U,X}$, where U is an object of X. Using the right exactness and commutation with direct sums of the functors $H^p(E) \otimes \,.\,$ and $H^p(E \otimes^{\mathrm{L}} \,.\,)$, we reduce to proving that the arrow of a') is an isomorphism for $M = \underline{O}_{U,X}$, $U \in \mathrm{ob}\,X$. Now $H^p(E) \otimes \underline{O}_{U,X}$ (resp. $H^p(E \otimes^{\mathrm{L}} \underline{O}_{U,X})$) identifies canonically (SGA 4 IV 2.13) with $i_!(H^p(E)|U)$ (resp. $H^p(i_!(E|U))$), where $i_!$ is extension by 0 outside U, and the result follows because $i_!$ is exact.

Suppose E pseudo-coherent, and let us prove b) $\Longrightarrow$ b'). It clearly suffices to show that $(*)$ is an isomorphism for M injective. Since the arrow $(***)$ is an isomorphism for M injective, the assertion will therefore follow from the following one:

(5.1.1): the arrow $(**)$ is an isomorphism if E is pseudo-coherent and M injective.

Proof of (5.1.1). Denote by T (resp. T') the functor from $D^-(X)$ to $D(X)$ defined by $F \mapsto F \otimes^{\mathrm{L}} M$ (resp. $\mathrm{R}\underline{\mathrm{Hom}}(F^\vee, M)$). Let $n \in \mathbb{Z}$; if $H^i(F) = 0$ for $i > n$, then $H^i T(F) = 0$ for $i > n$, $H^i(F^\vee) = 0$ for $i < -n$, and $H^i T'(F) = 0$ for $i > n$ if M is injective. Since, on the other hand, the arrow $(**)$ commutes with restriction to an open, and an injective restricted to an open is still injective, we may, after localizing and truncating, suppose E strictly perfect (I 2.1)[182]. By dévissage we then reduce to $E = \underline{O}_X$, and the result follows.

Remarks 5.2. (i) Condition a) (resp. a'), b), b')) is local in nature: if (X_α) is an open covering of the final object of X, and if, for each α, condition a) (resp. ...) holds for X_α and $E \,|\, X_\alpha$, then a) (resp. ...) holds.

(ii) If x is a point of X, denote by a_x) (resp. a'_x), etc.) the condi[print 262]tion a) (resp. a'), etc.) relative to E_x and $\mathrm{Mod}(\underline{O}_{X,x})$. Then, if X has enough points, condition a) (resp. a'), b)) is equivalent to the condition "for every point x of X, a_x) (resp. a'_x), b_x))", and is equivalent to the condition "for every point x of X, b'_x)" if E is pseudo-coherent; moreover, condition a') is equivalent to the analogous condition in which "for every $M \in \mathrm{ob}\,\mathrm{Mod}(X)$" is replaced by "for every $\underline{O}_X$-Module that is the cokernel of a homomorphism $\underline{O}^p_{U,X} \longrightarrow \underline{O}^q_{U,X}$, where U is an object of X".

(iii) The criterion (EGA III 7.4.2) extends word for word to our situation: if E has flat components, condition a) simply means that $\mathrm{Coker}(E^p \longrightarrow E^{p+1})$ is flat, hence locally a direct factor of a free Module of finite type when E^p and E^{p+1} are also supposed of finite presentation (I 5.8.3). In particular, we deduce that, if E is perfect, then E satisfies condition a) relative to the integer p if and only if $E^\vee$ satisfies condition a) relative to the integer $-p-1$.

(iv) Let X be a quasi-separated scheme, $\underline{O}_X$ the structure sheaf, $E \in \mathrm{ob}\,D^-(X)$ a complex with locally bounded and quasi-coherent cohomology. Denote by $\mathrm{a}_{\mathrm{qcoh}}$) (resp. $\mathrm{a}'_{\mathrm{qcoh}}$), etc.) the condition analogous to a) (resp. a'), etc.) in which[183] $\mathrm{Mod}(X)$ is replaced by the category

$\mathrm{Qcoh}(X)$ of quasi-coherent $\underline{O}_X$-Modules. Then we have

$$\mathrm{a}) \Longleftrightarrow \mathrm{a}_{\mathrm{qcoh}}) \Longleftrightarrow \mathrm{a}'_{\mathrm{qcoh}}) \Longleftrightarrow \mathrm{b}_{\mathrm{qcoh}}) \Longleftarrow \mathrm{b}'_{\mathrm{qcoh}}),$$

and the preceding conditions are equivalent if E is pseudo-coherent.

Indeed, since X is quasi-separated, every quasi-coherent sheaf on an affine open of X is induced by a quasi-coherent sheaf on X (EGA I 9.4.2 and IV 1.2.7). It follows that condition $\mathrm{a}_{\mathrm{qcoh}}$) (resp. $\mathrm{a}'_{\mathrm{qcoh}}$), $\mathrm{b}_{\mathrm{qcoh}}$), $\mathrm{b}'_{\mathrm{qcoh}}$)) holds if and only if it holds over every affine open. To prove the implications above, we may therefore suppose X affine. Then (II 3.5) E is isomorphic in $D(X)$ to a complex of quasi-coherent Modules, and the implications

$$\mathrm{a}_{\mathrm{qcoh}}) \Longleftrightarrow \mathrm{a}'_{\mathrm{qcoh}}) \Longleftrightarrow \mathrm{b}_{\mathrm{qcoh}}) \Longleftarrow \mathrm{b}'_{\mathrm{qcoh}})$$

[print 263]follow from (5.1) applied to the "one-point" topos ringed by $\Gamma(X, \underline{O}_X)$; the same holds, by (II 2.2.10 a)), for the implication $\mathrm{b}_{\mathrm{qcoh}}) \Longrightarrow \mathrm{b}'_{\mathrm{qcoh}})$ when E is pseudo-coherent. Finally, the equivalence a) $\Longleftrightarrow \mathrm{a}_{\mathrm{qcoh}})$, or, equivalently, a') $\Longleftrightarrow \mathrm{a}'_{\mathrm{qcoh}})$ follows from the last remark of (ii) (and the fact that quasi-coherent sheaves "extend", since X is quasi-separated).

Proposition 5.3. (exchange property) (EGA III 7.7.5 II). Let $f : X \to Y$ be a morphism of quasi-compact and quasi-separated schemes, $E \in \mathrm{ob}\, D^b(X)$ a complex with quasi-coherent cohomology and finite flat amplitude rel. to f, and $p \in \mathbb{Z}$. Consider the following conditions:

a) The functor $M \mapsto R^p f_*(E \otimes^{\mathrm{L}}_Y M)$ from $\mathrm{Qcoh}(Y)(^1)$ to $\mathrm{Qcoh}(Y)(^2)$ is right exact.

a') The canonical arrow

$$R^p f_*(E) \otimes M \longrightarrow R^p f_*(E \otimes^{\mathrm{L}}_Y M)$$

(the composite of the canonical arrow $R^p f_*(E) \otimes M \longrightarrow H^p(Rf_*(E) \otimes^{\mathrm{L}} M)$ and the isomorphism (3.7) $H^p(Rf_*(E) \otimes^{\mathrm{L}} M) \xrightarrow{\sim} R^p f_*(E \otimes^{\mathrm{L}}_Y M)$) is an isomorphism for every $M \in \mathrm{ob}\, \mathrm{Qcoh}(Y)$.

b) The functor $M \mapsto R^{p+1} f_*(E \otimes^{\mathrm{L}}_Y M)$ from $\mathrm{Qcoh}(Y)$ to $\mathrm{Qcoh}(Y)$ is left exact.

b') The canonical arrow

$$R^{p+1} f_*(E \otimes^{\mathrm{L}}_Y M) \longrightarrow \underline{\mathrm{Hom}}(H^{-p-1}(Rf_*(E)^\vee), M)$$

(the composite of the isomorphism (3.7) $R^{p+1} f_*(E \otimes^{\mathrm{L}}_Y M) \simeq H^{p+1}(Rf_*(E) \otimes^{\mathrm{L}} M)$ and the canonical arrow (5.1 $(*)$) $H^{p+1}(Rf_*(E) \otimes^{\mathrm{L}} M) \longrightarrow \underline{\mathrm{Hom}}(H^{-p-1}(Rf_*(E)^\vee), M)$) is an isomorphism for every $M \in \mathrm{ob}\, \mathrm{Qcoh}(Y)$. [print 264]

c) For every base change $Y' \to Y$, the canonical arrow

$$R^p f_*(E) \otimes_Y \underline{O}_{Y'} \longrightarrow H^p(Rf_*(E) \otimes^{\mathrm{L}}_Y \underline{O}_{Y'})$$

is an isomorphism.

Then we have:

$$\mathrm{a}) \Longleftrightarrow \mathrm{a}') \Longleftrightarrow \mathrm{b}) \Longleftrightarrow \mathrm{c}) \Longleftarrow \mathrm{b}').$$

Moreover, if f is proper, E pseudo-coherent rel. to f (hence perfect rel. to f), and subject to the validity of (2.1) for f (hence in particular if f is projective or if Y is noetherian), the preceding conditions are equivalent.

(^{1}) $\mathrm{Qcoh}(Y)$ = category of quasi-coherent $\underline{O}_Y$-Modules.
(^{2}) by (3.7).

Proof. The implications a) $\iff$ a′) $\iff$ b) $\impliedby$ b′) follow at once from (3.7), (5.1) and (5.2 (iv)), as does the implication b′) $\impliedby$ b) (for f proper and E pseudo-coherent rel. to f) by (2.1). It remains to prove that a) $\iff$ c). This will follow from the first assertion of the following lemma:

Lemma 5.3.1. Let X be a quasi-separated scheme, and let $E \in \operatorname{ob} D^-(X)$ be a complex with locally bounded and quasi-coherent cohomology. Then condition (5.1 a)) is equivalent to the condition

c) For every scheme X' over X, the canonical arrow

$$H^p(E) \otimes_X \underline{O}_{X'} \longrightarrow H^p(E \otimes_X^{\mathrm{L}} \underline{O}_{X'})$$

is an isomorphism.

Moreover, if E is pseudo-coherent, condition (5.1 a)) implies that, for every morphism $X' \to X$, the canonical arrow

$$H^{-p-1}(E^\vee) \otimes_X \underline{O}_{X'} \longrightarrow H^{-p-1}((E \otimes_X^{\mathrm{L}} \underline{O}_{X'})^\vee), \tag{5.3.1.1}$$

[print 265]the composite of $H^{-p-1}(E^\vee) \otimes_X \underline{O}_{X'} \longrightarrow H^{-p-1}(E^\vee \otimes_X^{\mathrm{L}} \underline{O}_{X'})$ and the canonical arrow (I 7.1.2) $H^{-p-1}(E^\vee \otimes_X^{\mathrm{L}} \underline{O}_{X'}) \longrightarrow H^{-p-1}((E \otimes_X^{\mathrm{L}} \underline{O}_{X'})^\vee)$, is an isomorphism.

Proof. a) $\Longrightarrow$ c). The question being local on X', we may suppose X (resp. X') affine with ring A (resp. A'). By (II 3.5), we may also suppose that $E = \widetilde{F}$, where F is a complex of A-modules. By hypothesis (condition a')), the arrow

$$H^p(F) \otimes_A A' \longrightarrow H^p(F \otimes_A^{\mathrm{L}} A')$$

is an isomorphism, and the result follows since $H^p(E) \otimes_X \underline{O}_{X'}$ (resp. $H^p(E \otimes_X^{\mathrm{L}} \underline{O}_{X'})$) identifies canonically with the sheaf on X' defined by the A'-module $H^p(F) \otimes_A A'$ (resp. $H^p(F \otimes_A^{\mathrm{L}} A')$).

c) $\Longrightarrow$ a). In view of (5.2 (iv)), it suffices to prove c) $\Longrightarrow$ $\mathrm{a}'_{\mathrm{qcoh}}$). The question being local (since X is quasi-separated), we may suppose X affine with ring A, and $E = \widetilde{F}$, where F is a complex of A-modules. Let $M = \widetilde{N}$ be a quasi-coherent $\underline{O}_X$-Module. We must show that the arrow

$$H^p(F) \otimes_A N \longrightarrow H^p(F \otimes_A^{\mathrm{L}} N)$$

is an isomorphism. But, putting $A' = D_A(N)$ (EGA O_{IV} 18.2.3), $X' = \operatorname{Spec}(A')$, this follows from c) applied to the canonical morphism $X' \to X$.

Let us prove the last assertion. The question being local on X', we may, by (II 2.2.10 b)), suppose E strictly pseudo-coherent. Moreover, since the assertion involves only a part of E of bounded length, we may, after truncating, suppose E strictly perfect. Then $(E \otimes_X^{\mathrm{L}} \underline{O}_{X'})^\vee$ identifies canonically (I 7.1.2) with $E^\vee \otimes_X^{\mathrm{L}} \underline{O}_{X'}$, and the arrow (5.3.1.1) may be rewritten as the canonical arrow

$$H^{-p-1}(E^\vee) \otimes_X \underline{O}_{X'} \longrightarrow H^{-p-1}(E^\vee \otimes_X^{\mathrm{L}} \underline{O}_{X'}). \tag{$*$}$$

[print 266]If E satisfies condition (5.1 a)) for the integer p, $E^\vee$ satisfies the same condition for the integer $-p-1$ by (5.2 (iii)), and the arrow $(*)$ is an isomorphism by the first assertion of (5.3.1). This completes the proof of (5.3.1), and hence of (5.3).

Remark 5.3.2. Proposition (5.3) applies in particular when E is a complex of bounded degrees of quasi-coherent $\underline{O}_X$-Modules that are flat and of finite presentation rel. to f: indeed, E then has finite perfect amplitude rel. to f by (4.3.2).

[print 267]**Definition 5.4.** Let X be a topos, and denote by $\mathrm{Pt}(X)$ the set of its points (SGA 4 IV 6.1). Let M be an ordered set, $f : \mathrm{Pt}(X) \to M$ a map, and $x \in \mathrm{Pt}(X)$. We say that f is upper

semi-continuous (resp. lower semi-continuous, resp. locally constant) at x if there exists an x-pointed object U of X such that for every point y of U we have $f(y) \leq f(x)$ (resp. $f(y) \geq f(x)$, resp. $f(y) = f(x)$). We say that f is upper semi-continuous (resp. lower semi-continuous, resp. locally constant) if it is so at every point([1]).

Examples 5.4.1. a) Denote by M_X the constant sheaf with value M on X, and let $s \in H^0(X, M)$. The map $x \mapsto s(x) = x^*(s)$ from $\mathrm{Pt}(X)$ to M is locally constant. In particular, let E be a perfect complex on X, supposed locally ringed: the rank of E (I 6.11) defines a locally constant function on $\mathrm{Pt}(X)$ with values in $\mathbb{Z}$.

b) Suppose X locally ringed, and let P be an $\underline{O}_X$-Module of finite type. For $x \in \mathrm{Pt}(X)$, denote by $\underline{O}_{X,x}$ the local stalk ring of $\underline{O}_X$ at x, and by $k(x)$ its residue field. Then, for $x \in \mathrm{Pt}(X)$, $P_x \otimes_{\underline{O}_{X,x}} k(x)$ has finite rank over $k(x)$, and the function $x \mapsto \mathrm{rg}_{k(x)}(P \otimes_{\underline{O}_{X,x}} k(x))$ is upper semi-continuous. Indeed, if, for $x \in \mathrm{Pt}(X)$, $P_x \otimes_{\underline{O}_{X,x}} k(x)$ has rank n, there is an epimorphism $\underline{O}_{X,x}^n \to P_x$ by Nakayama, and, since P is of finite type, there is an x-pointed object U of X and an epimorphism $\underline{O}_U^n \to P\,|\,U$, giving the assertion.

Lemma 5.5. Let $(X, \underline{O}_X)$ be a locally ringed topos, $E \in \mathrm{ob}\, D^-(X)$, and $p \in \mathbb{Z}$. Then we have, with the notation of (5.4.1 b)):

(i) [print 268]If E is perfect, the $H^i(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x))$, for $x \in \mathrm{Pt}(X)$, are vector spaces of finite rank over $k(x)$, all but finitely many zero, and the function

$$x \longmapsto \sum (-1)^i \,\mathrm{rg}_{k(x)}\, H^i(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x))$$

is locally constant on $\mathrm{Pt}(X)$.

(ii) If E is $(p-1)$-pseudo-coherent, $H^p(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x))$ is, for $x \in \mathrm{Pt}(X)$, a vector space of finite rank over $k(x)$, and the function

$$x \longmapsto \mathrm{rg}_{k(x)}\, H^p(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x))$$

is upper semi-continuous on $\mathrm{Pt}(X)$. If moreover the functor $M \mapsto H^p(E \otimes^{\mathrm{L}} M)$ from $\mathrm{Mod}(X)$ to $\mathrm{Mod}(X)$ is exact, $H^p(E)$ is locally free of finite type and the preceding function is locally constant.

(iii) Suppose E p-pseudo-coherent (resp. $(p-1)$-pseudo-coherent), and let $x \in \mathrm{Pt}(X)$. If the functor $M \mapsto H^p(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} M)$ is right exact (resp. left exact) on the category $\mathrm{Mod}(x)$ of $\underline{O}_{X,x}$-Modules, there is an x-pointed open U of X such that the functor $M \mapsto H^p(E\,|\,U \otimes^{\mathrm{L}} M)$ is right exact (resp. left exact) on $\mathrm{Mod}(U)$.

Proof. (i) By (I 4.19.1), $E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x)$ is a perfect complex of $k(x)$-modules, so (I 5.9 and 3.5) the $H^i(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x))$ have finite rank over $k(x)$ and all but finitely many are zero. We have

$$\mathrm{rg}(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x)) = \sum (-1)^i \,\mathrm{rg}_{k(x)}\, H^i(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x)), \qquad ([1]) \tag{5.5.1}$$

as is seen by an immediate dévissage using the additivity of the rank function. In view of (5.4.1 a)), assertion (i) therefore follows from [print 269]the compatibility of rank with inverse images (I 6.12).

([1]) When X comes from a topological space, we recover the usual notions.
([1]) rg( ) denotes the rank of a perfect complex (I 6.11).

(ii) After truncating and shifting, we may, for the first assertion, suppose E strictly perfect, $p = 0$, and $E^i = 0$ for $i \notin [-1, +1]$. For $x \in \mathrm{Pt}(X)$, put $E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} k(x) = E_{k(x)}$. By (5.5.1) we have

$$\begin{aligned}\operatorname{rg}_{k(x)} H^0(E_{k(x)}) = \operatorname{rg}_{k(x)} H^{-1}(E_{k(x)}) + \operatorname{rg}_{k(x)} H^1(E_{k(x)}) \\ + \operatorname{rg}(E_x),\end{aligned}$$

so it suffices to prove that the function $x \mapsto \operatorname{rg}_{k(x)} H^{-1}(E_{k(x)})$ (resp. $\operatorname{rg}_{k(x)} H^1(E_{k(x)})$) is upper semi-continuous[184] (since the function $x \mapsto \operatorname{rg}(E_x) = \operatorname{rg}(E_{k(x)})$[185] is locally constant). For the function $x \mapsto \operatorname{rg}_{k(x)} H^1(E_{k(x)})$, this is an immediate consequence of (5.4.1 b))(*). For the function $x \mapsto \operatorname{rg}_{k(x)} H^{-1}(E_{k(x)})$, we may suppose $E^1 = 0$, i.e. replace E by $E' = (0 \to E^{-1} \to E^0 \to 0)$. We already know that the function $x \mapsto \operatorname{rg}_{k(x)} H^0(E'_{k(x)})$ is upper semi-continuous, so the same holds for the function $x \mapsto \operatorname{rg}_{k(x)} H^{-1}(E'_{k(x)}) = \operatorname{rg}_{k(x)} H^{-1}(E_{k(x)})$, since the function $x \mapsto \operatorname{rg}_{k(x)} H^0(E'_{k(x)}) - \operatorname{rg}_{k(x)} H^{-1}(E'_{k(x)}) = \operatorname{rg}(E'_x)$ is locally constant. Let us now prove the second assertion. After truncating, we may suppose E strictly perfect and $E^i = 0$ for $i < p - 1$. The $\underline{O}_X$-Module $Z'^i(E) = \operatorname{Coker}(E^{i-1} \to E^i)$ is therefore of finite presentation. We have an exact sequence:

$$0 \to H^p(E) \to Z'^p(E) \to E^{p+1} \to Z'^{p+1}(E) \to 0. \qquad (*)$$

Since the functor $M \mapsto H^p(E \otimes^{\mathrm{L}} M)$ is right and left exact, $Z'^{p+1}(E)$ and $Z'^p(E)$ are flat[187] by (5.2 (iii)), hence (I 5.8.3) locally free of finite type. The exactness of (*) then implies that $H^p(E)$ is locally free of finite type, and consequently that the function $x \mapsto \operatorname{rg}_{k(x)} H^p(E_{k(x)}) = \operatorname{rg}(H^p(E)_x)$ [print 270]is locally constant, which completes the proof of (ii).

(iii) We may suppose E strictly perfect and $E^i = 0$ for $i < p$ (resp. $p-1$). Saying that the functor $M \mapsto H^p(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} M)$ is right exact (resp. left exact) on $\mathrm{Mod}(x)$ is equivalent, by (5.2 (iii)), to saying that $Z'^{p+1}(E_x)$ (resp. $Z'^p(E_x)$) (with the notation of (ii)) is flat, hence free of finite type (since it is of finite presentation). Since $Z'^{p+1}(E_x) \simeq Z'^{p+1}(E)_x$ (resp. $Z'^p(E_x) \simeq Z'^p(E)_x$), the usual argument (cf. EGA O_I 5.2.7) shows that there is an x-pointed object U of X such that $Z'^{p+1}(E\,|\,U)$ (resp. $Z'^p(E\,|\,U)$) is free of finite type, and the result follows.

Corollary 5.6 (EGA III 7.6.9). Let X be a scheme, $p \in \mathbb{Z}$, and $E \in \operatorname{ob} D^-(X)$ a $(p-1)$-pseudo-coherent complex. Then:

(i) For $x \in X$, $H^p(E \otimes^{\mathrm{L}}_X k(x))$ has finite rank over $k(x)$ and the function

$$x \longmapsto \operatorname{rg}_{k(x)} H^p(E \otimes^{\mathrm{L}}_X k(x))$$

is constructible and upper semi-continuous on X.

(ii) Suppose X quasi-separated and E with quasi-coherent and locally bounded cohomology. If the functor $M \mapsto H^p(E \otimes^{\mathrm{L}} M)$ is exact on the category of quasi-coherent $\underline{O}_X$-Modules, the function $x \mapsto \operatorname{rg}_{k(x)} H^p(E \otimes^{\mathrm{L}}_X k(x))$ is locally constant and $H^p(E)$ is locally free of finite type. Conversely, if X is locally noetherian and reduced and the preceding function is locally constant, the functor $M \mapsto H^p(E \otimes^{\mathrm{L}} M)$ is exact on the category of quasi-coherent $\underline{O}_X$-Modules.

Proof. If X is integral with generic point x, the functor $M \mapsto H^p(E_x \otimes^{\mathrm{L}}_{\underline{O}_{X,x}} M)$ is exact on the category of $\underline{O}_{X,x}$-modules (since $\underline{O}_{X,x} = k(x)$), so (5.5 (iii)) there is an open neighborhood U of x such that the functor $M \mapsto H^p(E\,|\,U \otimes^{\mathrm{L}} M)$ is exact on $\mathrm{Mod}(U)$, and, on U, the function [print 271]$y \mapsto \operatorname{rg}_{k(y)} H^p(E \otimes^{\mathrm{L}}_X k(y))$[188] is locally constant by (5.5 (ii)). The constructibility assertion of (i) follows at once. The other assertions follow from (5.5 (ii) and (iii)) and (5.2

(*) taking into account that $H^1(E_{k(x)}) \simeq H^1(E)_{k(x)}$[186].

(iv)), except for the converse of (ii), for which it suffices, after reducing to E strictly perfect (by localization and truncation), to apply (EGA III 7.6.2 and 7.6.5) to the functor $T_{-p} = H^p(E \otimes .)$.

Proposition 5.7. (EGA III 7.7.5 and 7.9.4). Let $f : X \to Y$ be a proper morphism of quasi-compact schemes, $E \in \mathrm{ob}\, D^b(f)_{\mathrm{parf}}$ (4.2), and $p \in \mathbb{Z}$. Then, subject to the validity of (2.1), hence for example if f is projective or if Y is noetherian:

(i) For $y \in Y$, the $H^i(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y))$ are vector spaces of finite rank over[189] $k(y)$, all but finitely many zero, and the function

$$y \longmapsto \sum (-1)^i \operatorname{rg}_{k(y)} H^i(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y))^{[190]}$$

is locally constant on Y.

(ii) The function

$$y \longmapsto \operatorname{rg}_{k(y)} H^p(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y))^{[191]}$$

is constructible and upper semi-continuous on Y.

(iii) Suppose Y quasi-separated. Then, if the functor $M \mapsto R^p f_*(E \otimes^{\mathrm{L}}_Y M)$ is exact on the category of quasi-coherent $\underline{O}_Y$-Modules, $R^p f_*(E)$ is a locally free $\underline{O}_Y$-Module of finite type, and the function of (ii) is locally constant. Conversely, if Y is noetherian and reduced, and if the function of (ii) is locally constant, the functor $M \mapsto R^p f_*(E \otimes^{\mathrm{L}}_Y M)$ is exact on the category of quasi-coherent $\underline{O}_Y$-Modules.

[print 272]Proof. By (4.8), $Rf_*(E)$ is a perfect complex.

Assertion (i) (resp. (ii)) therefore follows from (5.5 (i)) (resp. (5.6 (i))). Finally, if M is a quasi-coherent $\underline{O}_Y$-Module, the canonical arrow

$$Rf_*(E) \otimes^{\mathrm{L}}_Y M \longrightarrow Rf_*(E \otimes^{\mathrm{L}}_Y M)$$

is an isomorphism by (3.7), so assertion (iii) follows from (5.6 (ii)).

Remarks 5.8.

a) In the situation of (5.7), denote by $i_y : X_y \to X$ the inclusion of the fiber of X over a point y of Y. One can show (cf. (EGA III 6) and (IV 3.1.0))[192] that, for f flat (resp. E bounded with quasi-coherent and Y-flat components), we have the Künneth formula:

$$H^i(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y)) \xrightarrow{\sim} H^i(X_y, E_y),$$

where $E_y = Li_y^*(E)$ (resp. $i_y^*(E)$). [print 273]

b) One can develop for complex analytic spaces a body of results analogous to that of this section (exchange, semi-continuity, continuity properties)([1]). The essential ingredients are: 1) the finiteness theorem for proper morphisms ("Grauert's theorem"); 2) an analogue of the projection formula (3.7), which may be stated as follows: let $f : X \to Y$ be a morphism of analytic spaces, where X is supposed of bounded dimension, and let $E \in \mathrm{ob}\, D^-(X)$, $M \in \mathrm{ob}\, D^-(Y)_{\mathrm{coh}}$ ("coh" means "with coherent cohomology"). The canonical arrow

$$Rf_*(E) \otimes^{\mathrm{L}} M \longrightarrow Rf_*(E \otimes^{\mathrm{L}}_Y M)^{[193]}$$

is an isomorphism (the hypothesis on X ensures (T.F II 5.12) that f_* has finite cohomological dimension, hence that Rf_* is defined on $D(X)$).

Unlike statement 1), statement 2) is essentially trivial, and follows from a dévissage analogous to that of the proof of (3.7); we moreover have greater freedom since Rf_* is defined on $D(X)$, and in particular no hypothesis on E is needed (apart from $E \in \mathrm{ob}\, D^-(X)$).

([1]) It was moreover in this setting that these properties were first formulated (see for example Grauert [1]), before being transposed to Algebraic Geometry in (EGA III 7).

BIBLIOGRAPHY

[1] Grauert, H., Ein Theorem der Analytischen Garbentheorie ..., Pub. IHES, n°5.

[H] Hartshorne, R., Residues and Duality, Lecture Notes n°20, Springer, 1966.

[T.F] Godement, R., Théorie des faisceaux, Hermann, 1958.

[print 274]

EXPOSÉ IV
GROTHENDIECK GROUPS OF RINGED TOPOI

by L. ILLUSIE

1. Reminders and generalities on Grothendieck groups

1.1. Let C be a triangulated category. Recall (I 6.3 and SGA 5 VIII 2) that a map f from $\operatorname{ob} C$ to an abelian group G is called additive if $f(E) = f(E') + f(E'')$ for every distinguished triangle $E' \longrightarrow E \longrightarrow E'' \longrightarrow E'[1]$. The additive maps from $\operatorname{ob} C$ to G form an abelian group, depending functorially on G. The functor thus obtained is represented by an abelian group $k(C)$ and a universal additive map $\mathrm{cl}_C : \operatorname{ob} C \longrightarrow k(C)$ (denoted by cl when there is no risk of confusion).

The group $k(C)$ depends functorially on C with respect to exact functors. If C, C' are triangulated categories, two isomorphic exact functors from C to C' induce the same homomorphism $k(C) \to k(C')$; in particular, if $u : C \to C'$ is an exact functor that is an equivalence of categories, $k(u) : k(C) \to k(C')$ is an isomorphism.

Lemma 1.2. Let C be a triangulated category, and let $L \in \operatorname{ob} C$.

a) For every $n \in \mathbb{Z}$, we have
$$\mathrm{cl}(L[n]) = (-1)^n \,\mathrm{cl}(L).$$
If L', L'' are objects of C such that $L \simeq L' \oplus L''$, then $\mathrm{cl}(L) = \mathrm{cl}(L') + \mathrm{cl}(L'')$.

b) Suppose there exists $n \in \mathbb{Z}$ such that $\coprod_{i\geq 0} L[2ni]$ is representable. Then $\mathrm{cl}(L) = 0$.

Proof. Assertion a) follows directly from the definitions (cf. SGA 5 VIII 2). For b), it suffices to observe that we have [print 275]

$$\coprod_{i\geq 0} L[2ni] \simeq L \oplus \left(\coprod_{i\geq 0} L[2ni]\right)[2n],$$

and to apply a).

Lemma 1.3. Let C be a triangulated category, and A a thick subcategory ([V] I 2.1). The inclusion and quotient functors define an exact sequence

$$k(A) \longrightarrow k(C) \longrightarrow k(C/A) \longrightarrow 0.$$

Proof. See (SGA 5 VIII 3.1).

Lemma 1.4. Let A be an additive (resp. abelian) category, and f an additive function (1.1) on $\operatorname{ob} K^b(A)$ (resp. $\operatorname{ob} D^b(A)$). For $E \in \operatorname{ob} K^b(A)$ (resp. $D^b(A)$) we have

$$f(E) = \sum (-1)^i f(E^i)$$

(resp.
$$f(E) = \sum (-1)^i f(H^i(E))$$

).

Proof. Left as an exercise to the reader.

1.5. Let C be an abelian category and A a full subcategory of C, stable under finite sums. A map f from $\operatorname{ob} A$ to an abelian group G is called additive if $\sum(-1)^i f(E^i) = 0$ for every acyclic complex E of C, with bounded degrees and components belonging to A. When A is stable under kernels of epimorphisms (I 1.2), this amounts to saying that $f(L) = f(L') + f(L'')$ for every exact sequence $0 \longrightarrow L' \longrightarrow L \longrightarrow L'' \longrightarrow 0$ of C whose terms belong to A. The additive maps from $\operatorname{ob} A$ to G form an abelian group depending functorially on G. Denote by $K^{b,\emptyset}(A)$ the thick subcategory of $K^b(A)$ formed by the acyclic complexes. It follows easily from (1.3) and (1.4) that the map

$$\mathrm{cl}_A : \operatorname{ob} A \longrightarrow k\big(K^b(A)/K^{b,\emptyset}(A)\big) \tag{1.5.1}$$

[print 276]induced by $\mathrm{cl}_{K^b(A)/K^{b,\emptyset}(A)}$ (1.1) is a universal additive map, i.e. the pair $\Big(k\big(K^b(A)/K^{b,\emptyset}(A)\big), \mathrm{cl}_A\Big)$ represents the functor assigning to each abelian group G the abelian group of additive maps from $\operatorname{ob} A$ to G. We shall put

$$k\big(K^b(A)/K^{b,\emptyset}(A)\big) = k(A) \tag{1.5.2}$$

(we shall write $k(A, C)$ instead of $k(A)$ when we wish to avoid any ambiguity about the ambient abelian category)([1]). In particular, for $A = C$, we have

$$k(D^b(C)) = k(C). \tag{1.5.3}$$

Lemma 1.6. Let C be an abelian category, and A a thick subcategory (i.e. a full subcategory stable under kernels, cokernels, and extensions). Denote by $D_A(C)$ the full subcategory of $D(C)$ formed by the objects E such that $H^i(E) \in \operatorname{ob} A$ for every i. Then $D_A(C)$ is a triangulated subcategory ([V] I 1.2.3) of $D(C)$, and the canonical functor $D^b(A) \longrightarrow D^b_A(C) = D^b(C) \cap D_A(C)$ induces an isomorphism

$$k(A) \xrightarrow{\sim} k(D^b_A(C)),$$

where $k(A)$ is defined by (1.5.2).

Proof. The first assertion is an immediate consequence of the long exact cohomology sequence. For the second, consider the map $f : \operatorname{ob} D^b_A(C) \longrightarrow k(A)$ defined by

$$f(E) = \sum (-1)^i \,\mathrm{cl}_A(H^i(E)),$$

where cl_A is the map (1.5.1). By the long exact cohomology sequence, f is additive (1.1), hence defines a homomorphism

$$\varphi : k(D^b_A(C)) \longrightarrow k(A).$$

[print 277]It follows at once from (1.4) that φ is inverse to the canonical homomorphism $k(A) \longrightarrow k(D^b_A(C))$.

Remark 1.6.1. In the situation of (1.6), the canonical functor $D^b(A) \longrightarrow D^b_A(C)$ is not in general an equivalence of categories, as shown by the example where S is a scheme, C the category of $\underline{O}_S$-Modules, and A the category of quasi-coherent $\underline{O}_S$-Modules (II Ap. I 0.3). See, however, the criterion (II 2.1.3).

Definition 1.7. Let C be a triangulated (resp. abelian) category, and C' a subset of $\operatorname{ob} C$. We shall say that C' is an exact subset (cf. EGA III 3.1.1) if, for every distinguished triangle

([1]) The notation $k(A)$ was introduced in (I 6.3) to denote the group $k(K^b(A))$, which is not in general isomorphic to the group defined by (1.5.2). We therefore change convention here: in the present exposé, the notation $k(A)$ will denote exclusively the group defined by (1.5.2).

$E' \to E \to E'' \to E'[1]$ (resp. every exact sequence $0 \to E' \to E \to E'' \to 0$) such that two of the terms E', E, E'' belong to C', the third also belongs to it. We shall say that C' is dense in $\mathrm{ob}\, C$ if $\mathrm{ob}\, C$ is the smallest exact subset of $\mathrm{ob}\, C$ containing C'.

Lemma 1.8. Let C be a triangulated (resp. abelian) category, and C' a dense subset of $\mathrm{ob}\, C$. Then the elements $\mathrm{cl}_C(E)$, for $E \in C'$, generate $k(C)$.

Proof. Put $C' = C_0$, and, for $i \geq 1$, define C_i inductively as the set of objects of C appearing in a distinguished triangle (resp. a short exact sequence) two of whose terms belong to C_{i-1}. The C_i form an increasing filtration of $\mathrm{ob}\, C$, which is exhaustive by hypothesis. For $i \geq 0$, denote by $k(C)_i$ the subgroup of $k(C)$ generated by the elements $\mathrm{cl}_C(E)$ for $E \in C_i$. It is immediate from the definitions that, for every $i \geq 0$, $k(C)_i = k(C)_{i+1}$, hence $k(C)_i = k(C)_0$. Now let $x \in k(C)$; [print 278]as is easily seen, we can write $x = \mathrm{cl}(E') - \mathrm{cl}(E'')$ for $E', E'' \in \mathrm{ob}\, C$ (in the first case, we can even arrange that $E'' = 0$). Since the filtration (C_i) is exhaustive, there is an integer $i \geq 0$ such that E' and E'' belong to C_i; then $x \in k(C)_i = k(C)_0$, as required.

Exercise 1.9. Let C be an abelian category, and A a full subcategory stable under finite sums, subobjects, and quotients, such that $\mathrm{ob}\, A$ is dense in $\mathrm{ob}\, C$. Then the homomorphism $k(A, C) \longrightarrow k(C)$ defined by the additive function $E \mapsto \mathrm{cl}_C(E)$ on $\mathrm{ob}\, A$ (cf. (1.5)) is an isomorphism.

Hint: Put $A = A_0$, and define A_i, for $i \geq 1$, as the full subcategory of C formed by the objects of C that are extensions of two objects of C_{i-1}. Prove that A_i satisfies the same hypotheses as A, then show that the canonical homomorphism $\varinjlim k(A_i, C) \longrightarrow k(C)$ is an isomorphism (the desired conclusion follows at once).

2. The functors $K_\bullet$ and $K^\bullet$ of a ringed topos

2.1. Let (X, A) be a ringed topos. Recall the following notation:

$D^b({}_AX)_{\mathrm{coh}}$ = category of pseudo-coherent complexes with bounded cohomology of left A-Modules (I 2.15).

$D({}_AX)_{\mathrm{parf}}$ = category of complexes of left A-Modules of finite perfect amplitude (I 5.8.4).

$\mathrm{Loclib}({}_AX)$ = category of left A-Modules locally direct summands of free Modules of finite type (II 2.0).

$\mathrm{Coh}({}_AX)$ = category of coherent left A-Modules (relative to the category of free A-Modules of finite type) (I 3.1).

[print 279]The last two categories are full subcategories of the category $\mathrm{Mod}({}_AX)$ of left A-Modules, and the first two are full triangulated subcategories of $D({}_AX)$ $(= D(\mathrm{Mod}({}_AX)))$.

Definition 2.2. Let (X, A) be a ringed topos. Unless otherwise stated, we shall put

$$K_\bullet({}_AX) = k(D^b({}_AX)_{\mathrm{coh}}),$$
$$K^\bullet({}_AX) = k(D({}_AX)_{\mathrm{parf}}),$$

where $k(\ )$ is defined in (1.1), and

$$K_\bullet({}_AX)_{\mathrm{naïf}} = k(\mathrm{Coh}({}_AX)),$$
$$K^\bullet({}_AX)_{\mathrm{naïf}} = k(\mathrm{Loclib}({}_AX)),$$

where $k(\ )$ is defined by (1.5.2), the ambient abelian category being $\mathrm{Mod}({}_AX)$([1]). We shall

([1]) We shall allow ourselves to omit the index A when no confusion can result. Sometimes we shall also drop the index "naïf", but shall indicate this explicitly.

say that $K_\bullet({}_AX)$ (resp. $K^\bullet({}_AX)$) is the Grothendieck group of pseudo-coherent (resp. perfect) complexes of left A-Modules, and that $K_\bullet({}_AX)_{\text{naïf}}$ (resp. $K^\bullet({}_AX)_{\text{naïf}}$) is the Grothendieck group of coherent (resp. locally free) left A-Modules.

Remarks 2.3.

a) One can define variants of $K_\bullet(\ )$ and $K^\bullet(\ )$ by putting

$$K^{\text{loc}}_\bullet({}_AX) = k(D^{\text{loc}b}({}_AX)_{\text{coh}})$$

(where "loc$b$" means "with locally bounded cohomology"), and

$$K^\bullet_{\text{loc}}({}_AX) = k(D({}_AX)_{\text{parf}}).$$

These groups coincide with those defined in (2.2) when the final object of X is quasi-compact, but in general are less suited to intersection calculations.

b) [print 280]The group $K_\bullet({}_AX)_{\text{naïf}}$ is practically of interest only when A is coherent (otherwise there may well be very few coherent sheaves ...).

c) Considering the k of triangulated categories that are "too large" is of no interest. Thus, from (1.2 b)) one easily deduces the relations

$$k(D({}_AX)) = k(D^-({}_AX)) = k(D^+({}_AX)) = k(D^b({}_AX)) = k(D^-({}_AX)_{\text{coh}}) = 0$$

(for the last equality, observe that $L \in \operatorname{ob} D^-({}_AX)_{\text{coh}}$ implies $\coprod_{i\geq 0} L[2i] \in \operatorname{ob} D^-({}_AX)_{\text{coh}}$).

2.4. Case of a coherent sheaf of rings. Let (X, A) be a ringed topos, with A supposed coherent as a left A-Module (I 3.1 and 3.6). Every coherent left A-Module is a pseudo-coherent complex (I 3.5), so we have a fully faithful functor

$$\operatorname{Coh}({}_AX) \hookrightarrow D^b({}_AX)_{\text{coh}}. \tag{2.4.1}$$

Now (I 3.4) $\operatorname{Coh}({}_AX)$ is a thick subcategory of $\operatorname{Mod}({}_AX)$, and, by (I 3.5), $D^b({}_AX)_{\text{coh}}$ is the full subcategory of $D^b({}_AX)$ formed by the objects L such that $H^i(L) \in \operatorname{ob}\operatorname{Coh}({}_AX)$ for every i. Consequently, it follows from (1.6) that the homomorphism

$$K_\bullet({}_AX)_{\text{naïf}} \longrightarrow K_\bullet({}_AX) \tag{2.4.2}$$

defined by (2.4.1) is an isomorphism.

2.5. The homomorphism $K^\bullet \to K_\bullet$. Let (X, A) be a ringed topos. The inclusion functor $D({}_AX)_{\text{parf}} \hookrightarrow D^b({}_AX)_{\text{coh}}$ defines a homomorphism

$$K^\bullet({}_AX) \longrightarrow K_\bullet({}_AX). \tag{2.5.1}$$

This is in general neither injective nor surjective. However, make the following hypotheses:

(i) [print 281]the final object of X is quasi-compact;

(ii) X has enough points, and, for every point x of X, every left A_x-module has finite tor-dimension.

Then, by (I 5.10), $D({}_AX)_{\text{parf}} = D^b({}_AX)_{\text{coh}}$, and the homomorphism (2.5.1) is an isomorphism.

Hypotheses (i) and (ii) are satisfied, for example, by a compact space ringed by regular local rings.

2.6. Pairings between $K^\bullet$ and $K^\bullet$ (noncommutative case). Let

$$f : (X', A') \longrightarrow (X, A)$$

be a morphism of ringed topoi. Denote by $\mathrm{Mod}({}_{A'}X'_A)$ the category of bi-Modules, left over A' and right over $f^{-1}(A)$, and by $\mathrm{Loclib}_{A'}({}_{A'}X'_A)$ the full subcategory formed by the objects belonging to $\mathrm{Loclib}({}_{A'}X')$. By analogy with (2.2), we shall put

$$K^\bullet_{\mathrm{naïf}}({}_{A'}X'_A)_{A'} = k(\mathrm{Loclib}_{A'}({}_{A'}X'_A)).$$

The tensor product

$$\otimes_A : \mathrm{Mod}({}_{A'}X'_A) \times \mathrm{Mod}({}_AX) \longrightarrow \mathrm{Mod}({}_{A'}X')$$
$$(E', E) \longmapsto E' \otimes_{f^{-1}(A)} f^{-1}(E)$$

induces a functor

$$\mathrm{Loclib}_{A'}({}_{A'}X'_A) \times \mathrm{Loclib}({}_AX) \longrightarrow \mathrm{Loclib}({}_{A'}X'). \tag{2.6.1}$$

The map $(E', E) \mapsto \mathrm{cl}(E' \otimes_A E)$ deduced from (2.6.1) is biadditive in the sense of (1.5), and therefore defines a homomorphism

$$K^\bullet_{\mathrm{naïf}}({}_{A'}X'_A)_{A'} \otimes K^\bullet_{\mathrm{naïf}}({}_AX) \longrightarrow K^\bullet_{\mathrm{naïf}}({}_{A'}X'). \tag{2.6.2}$$

Denote by $D({}_{A'}X'_A)$ the derived category of $\mathrm{Mod}({}_{A'}X'_A)$, and by $D({}_{A'}X'_A)_{A'\text{-parf}}$ the full subcategory formed by the complexes of finite perfect amplitude rel. to A' (i.e. as objects of $D({}_{A'}X')$), and put, as in (2.2),

$$K^\bullet({}_{A'}X'_A)_{A'} = k(D({}_{A'}X'_A)_{A'\text{-parf}}).^{[194]}$$

[print 282]The inclusion functors plainly define homomorphisms

$$K^\bullet_{\mathrm{naïf}}({}_{A'}X'_A)_{A'} \longrightarrow K^\bullet({}_{A'}X'_A)_{A'}, \tag{2.6.3.1}$$
$$K^\bullet_{\mathrm{naïf}}({}_AX) \longrightarrow K^\bullet({}_AX), \tag{2.6.3.2}$$
$$K^\bullet_{\mathrm{naïf}}({}_{A'}X') \longrightarrow K^\bullet({}_{A'}X'). \tag{2.6.3.3}$$

On the other hand, we know (I 4.18, 4.19) that the derived functor of $\otimes_A$,

$$\otimes^{\mathrm{L}}_A : D^-({}_{A'}X'_A) \times D^-({}_AX) \longrightarrow D^-({}_{A'}X'),$$

induces an (exact) functor

$$D({}_{A'}X'_A)_{A'\text{-parf}} \times D({}_AX)_{\mathrm{parf}} \longrightarrow D({}_{A'}X')_{\mathrm{parf}}. \tag{2.6.4}$$

The latter defines a homomorphism

$$K^\bullet({}_{A'}X'_A)_{A'} \otimes K^\bullet({}_AX) \longrightarrow K^\bullet({}_{A'}X'). \tag{2.6.5}$$

Since the functor (2.6.1) is induced by (2.6.4), the homomorphisms (2.6.2) and (2.6.5) are compatible with the homomorphisms (2.6.3.1), (2.6.3.2), and (2.6.3.3); in other words, we have a commutative diagram

$$\begin{array}{ccc} K^\bullet_{\mathrm{naïf}}({}_{A'}X'_A)_{A'} \otimes K^\bullet_{\mathrm{naïf}}({}_AX) & \to & K^\bullet_{\mathrm{naïf}}({}_{A'}X') \\ \downarrow & & \downarrow \\ K^\bullet({}_{A'}X'_A)_{A'} \otimes K^\bullet({}_AX) & \longrightarrow & K^\bullet({}_{A'}X'), \end{array} \tag{2.6.6}$$

where the upper and lower horizontal arrows are respectively (2.6.2) and (2.6.5), and the vertical arrows are defined by (2.6.3.1), (2.6.3.2), and (2.6.3.3).

[print 283]2.7. Contravariant law and ring structure on $K^\bullet$ We shall spell out some important special cases of the pairings (2.6) (their direct study is moreover immediate).

a) Let $f : (X', A') \longrightarrow (X, A)$ be a morphism of ringed topoi. The functor $Lf^* : D^-({}_AX) \longrightarrow D^-({}_{A'}X')$ sends (I 4.19.1) $D({}_AX)_{\underline{\text{parf}}}$ to $D({}_{A'}X')_{\underline{\text{parf}}}$ (resp. $\text{Loclib}({}_AX)$ to $\text{Loclib}({}_{A'}X')$), and defines a homomorphism

$$f^* : K^\bullet({}_AX) \longrightarrow K^\bullet({}_{A'}X') \tag{2.7.1}$$

(resp.

$$f^* : K^\bullet({}_AX)_{\text{naïf}} \longrightarrow K^\bullet({}_{A'}X')_{\text{naïf}} \tag{2.7.2}$$

), which can also be interpreted as multiplication by $\text{cl}(A')$ in (2.6.5) (resp. (2.6.2))[195]. We have a commutative diagram (a special case of (2.6.6))

$$\begin{array}{ccc} K^\bullet({}_AX)_{\text{naïf}} & \xrightarrow{f^*} & K^\bullet({}_{A'}X')_{\text{naïf}} \\ \downarrow & & \downarrow \\ K^\bullet({}_AX) & \xrightarrow{f^*} & K^\bullet({}_{A'}X'), \end{array}$$

where the left and right vertical arrows are respectively (2.6.3.2) and (2.6.3.3).

Let $g : (X, A) \to (X'', A'')$ be a morphism of ringed topoi. The transitivity of inverse-image functors implies that the homomorphisms of type (2.7.1) (resp. 2.7.2) satisfy the relation

$$(gf)^* = f^*g^*. \tag{2.7.3}$$

Thus $K^\bullet({}_AX)$ (resp. $K^\bullet({}_AX)_{\text{naïf}}$) is a contravariant functor of (X, A).

b) Let $(X, \underline{O}_X)$ be a commutatively ringed topos. By the associativity and commutativity of the tensor product, the homomorphism (2.6.5) (resp. (2.6.2)) (where we put $A = \underline{O}_X$, $(X', A') = (X, A)$, $f = \text{Id}$) endows $K^\bullet(X)$ (resp. $K^\bullet(X)_{\text{naïf}}$) with the structure of a commutative ring with unit [print 284]$(1 = \text{cl}(\underline{O}_X))$, and the canonical homomorphism (2.6.3.2)

$$K^\bullet(X)_{\text{naïf}} \longrightarrow K^\bullet(X)$$

is a homomorphism of unital rings.

Let $f : (X, \underline{O}_X) \to (Y, \underline{O}_Y)$ be a morphism of topoi ringed by commutative rings. Then compatibility of the tensor product with inverse images shows that the homomorphism

$$f^* : K^\bullet(Y) \longrightarrow K^\bullet(X)$$

(resp.

$$f^* : K^\bullet(Y)_{\text{naïf}} \longrightarrow K^\bullet(X)_{\text{naïf}}$$

) is a homomorphism of unital rings.

2.8. Augmentation of $K^\bullet$ to $H^0(\ , \mathbb{Z})$. Let $(X, \underline{O}_X)$ be a locally ringed topos (I 2.15.1). Recall that we have defined (I 6.11 to 6.14) a homomorphism of (unital) rings

$$i : H^0(X, \mathbb{Z}) \longrightarrow K^\bullet(X)_{\text{naïf}},$$

thus giving $K^\bullet(X)_{\text{naïf}}$ (and consequently also $K^\bullet(X)$ via the canonical homomorphism $K^\bullet(X)_{\text{naïf}} \to K^\bullet(X)$) the structure[196] of an algebra (with unit) over $H^0(X, \mathbb{Z})$. Moreover (loc.cit.), the algebra $K^\bullet(X)_{\text{naïf}}$ (resp. $K^\bullet(X)$) is augmented to $H^0(X, \mathbb{Z})$ by the rank, and the canonical homomorphism $K^\bullet(X)_{\text{naïf}} \to K^\bullet(X)$ is a homomorphism of augmented algebras.

Let $f : (X, \underline{O}_X) \to (Y, \underline{O}_Y)$ be a morphism of locally ringed topoi. The homomorphisms $f^* : K^\bullet(Y) \longrightarrow K^\bullet(X)$ and $f^* : K^\bullet(Y)_{\text{naïf}} \longrightarrow K^\bullet(X)_{\text{naïf}}$ are compatible (I 6.12, 6.13) with the

preceding augmented algebra structures; this is expressed by the commutativity of the diagram (where rg denotes rank): [print 285]

$$\begin{array}{ccccccc} H^0(Y,\mathbb{Z}) & \xrightarrow{i} & K^\bullet(Y)_{\text{naïf}} & \to & K^\bullet(Y) & \xrightarrow{\text{rg}} & H^0(Y,\mathbb{Z}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H^0(X,\mathbb{Z}) & \xrightarrow{i} & K^\bullet(X)_{\text{naïf}} & \to & K^\bullet(X) & \xrightarrow{\text{rg}} & H^0(X,\mathbb{Z}). \end{array}$$

We shall see later in the Seminar (V 3.9.1 and VI)[197] that $K^\bullet(X)_{\text{naïf}}$ can be given a richer structure than that of an algebra augmented to $H^0(X,\mathbb{Z})$, namely that of a λ-algebra augmented to $H^0(X,\mathbb{Z})$ regarded as a binomial ring (V 2.8), this structure being compatible with inverse-image homomorphisms. Deligne [1][198] recently proved that $K^\bullet(X)$ can be given an analogous structure (at least when X has characteristic $p \geq 0$) and that the canonical homomorphism $K^\bullet(X)_{\text{naïf}} \longrightarrow K^\bullet(X)$ is a homomorphism of augmented λ-algebras, thus answering a conjecture of Grothendieck (XIV 1) in the affirmative.

2.9. Comparison of $K^\bullet$ and $K^\bullet_{\text{naïf}}$. Let (X,A) be a ringed topos. In general there is no reason for the canonical homomorphism

$$K^\bullet(X)_{\text{naïf}} \longrightarrow K^\bullet(X)$$

to be injective or surjective. However, the "criteria for the existence of global resolutions" of Exposé II show that it is an isomorphism in the following cases:

(i) X is the Zariski topos of a divisorial scheme, separated or noetherian (and $A = \underline{O}_X$) (II 2.2.9);

(ii) X is the topos defined by a compact space and A is a soft sheaf (II 2.3.2);

(iii) X is a Stein topos whose final object is quasi-compact and $A = \underline{O}_X$ (II 2.4.5).

[print 286]2.10. Pairing between $K^\bullet$ and $K_\bullet$. Let $(X,\underline{O}_X)$ be a commutatively ringed topos. We know (I 5.8.4 b)) that there is a pairing

$$\otimes^{\mathrm{L}}_A : D(X)_{\underline{\text{parf}}} \times D^b(X)_{\text{coh}} \longrightarrow D^b(X)_{\text{coh}}.$$

This defines a homomorphism

$$K^\bullet(X) \otimes K_\bullet(X) \longrightarrow K_\bullet(X). \tag{2.10.1}$$

Thanks to the associativity[199] of the tensor product, we see that (2.10.1) endows $K_\bullet(X)$ with the structure of a $K^\bullet(X)$-module. More generally, if $f : (X,\underline{O}_X) \longrightarrow (Y,\underline{O}_Y)$ is a morphism of commutatively ringed topoi, we can regard $K_\bullet(X)$ as a $K^\bullet(Y)$-module via the homomorphism $f^* : K^\bullet(Y) \longrightarrow K^\bullet(X)$.

2.11. Covariant law on $K_\bullet$ and projection formula. Let (X,A) be a ringed topos. Unlike $K^\bullet({}_AX)$, the group $K_\bullet({}_AX)$ has no natural variance with respect to (X,A). However, in Algebraic Geometry and Analytic Geometry, $K_\bullet(X)$ tends to be covariant in X for proper morphisms. Let us make this precise:

2.11.1. Let $f : X \to Y$ be a proper and pseudo-coherent morphism (III 1.2) of quasi-compact schemes. Then, subject to the validity of the finiteness conjecture for f (III 2.1), hence in particular (III 2.2) when Y is noetherian (in which case the pseudo-coherence hypothesis on f is automatically satisfied (III 1.3)) or when f is projective, the functor Rf_* induces, by (III 2.6), a functor

$$Rf_* : D^b(X)_{\text{coh}} \longrightarrow D^b(Y)_{\text{coh}}.$$

This defines a homomorphism

$$f_* : K_\bullet(X) \longrightarrow K_\bullet(Y) \tag{2.11.1.1}$$

[print 287]This homomorphism is $K^\bullet(Y)$-linear (cf. (2.10)); in other words, for $y \in K^\bullet(Y)$ and $x \in K_\bullet(X)$, we have the so-called projection formula

$$f_*(yx) = y f_*(x). \tag{2.11.1.2}$$

Indeed, if $x = \mathrm{cl}(E)$ for $E \in \mathrm{ob}\, D^b(X)_{\mathrm{coh}}$ and $y = \mathrm{cl}(M)$ for $M \in \mathrm{ob}\, D(Y)_{\underline{\mathrm{parf}}}$, we have (III 3.7)

$$Rf_*(E) \otimes^{\mathrm{L}} M \simeq Rf_*(E \otimes^{\mathrm{L}}_Y M).$$

Finally, if $g : Y \to Z$ is a morphism satisfying the same hypotheses as f and for which the finiteness conjecture (III 2.1) holds, the transitivity of direct-image functors gives, for the homomorphisms (2.11.1.1), the formula

$$g_* f_* = (gf)_*. \tag{2.11.1.3}$$

2.11.2. Let $f : X \to Y$ be a proper morphism of finite-dimensional analytic spaces. Then, by Grauert's finiteness theorem [2], Rf_* sends $D^b(X)_{\mathrm{coh}}$ to $D^b(Y)_{\mathrm{coh}}$, and therefore defines a homomorphism

$$f_* : K_\bullet(X) \longrightarrow K_\bullet(Y). \tag{2.11.2.1}$$

Like its algebraic analogue (2.11.1.1), this is $K^\bullet(Y)$-linear, i.e. formula (2.11.1.2) is valid, as follows from the projection formula at the level of derived categories (III 5.8 b))[200]. We likewise have the transitivity formula (2.11.1.3).

2.11.3. Once we leave the algebraic or analytic setting, this covariance of $K_\bullet$ for proper morphisms disappears. Thus, let $f : X \to Y$ be a proper morphism of topological spaces ringed by the sheaf of complex continuous functions. Then, for $E \in \mathrm{ob}\, D^b(X)_{\mathrm{coh}}$, in general we do not have $Rf_*(E) \in \mathrm{ob}\, D^b(Y)_{\mathrm{coh}}$, as is already evident for $E = \underline{O}_X$ and Y reduced to a point. In fact, definition (2.2) of $K_\bullet$ [print 288]appears especially suited to the needs of Algebraic Geometry and Analytic Geometry, and a good definition of the $K_\bullet$ of a topological space remains to be found. It seems moreover that we shall have to leave the setting of $\underline{O}_X$-linear complexes and define, for a morphism $f : X \to Y$ (of topological spaces, resp. of mixed varieties (II App. II), etc.), a suitable class of complexes of topological vector sheaves on X, which should include, for example, when f is the structural morphism of a mixed variety, the elliptic complexes on X relative to Y. This is suggested by the finiteness theorem (II App. II), as well as by a similar theorem (where Y is supposed reduced to a point) proved by Grothendieck in [3].

2.12. Unusual variances of $K^\bullet$ and $K_\bullet$. Let $f : (X, A) \longrightarrow (Y, B)$ be a morphism of finite tor-dimension of ringed topoi (III 3.1). The functor $Lf^* : D^-({}_BY) \longrightarrow D^-({}_AX)$ sends $D^b({}_BY)$ to $D^b({}_AX)$, hence (I 2.16.1) $D^b({}_BY)_{\mathrm{coh}}$ to $D^b({}_AX)_{\mathrm{coh}}$, and therefore defines a homomorphism

$$f^* : K_\bullet({}_BY) \longrightarrow K_\bullet({}_AX). \tag{2.12.1}$$

If A and B are commutative, commutation of inverse image with the tensor product shows that the homomorphism (2.12.1) is compatible with the module structures of $K_\bullet(Y)$ and $K_\bullet(X)$ over $K^\bullet(Y)$ and $K^\bullet(X)$ respectively (2.10); in other words, for $a \in K^\bullet(Y)$ and $y \in K_\bullet(Y)$, we have the formula

$$f^*(ay) = a f^*(y). \tag{2.12.2}$$

If $g : (Y, B) \to (Z, C)$ is a morphism of finite tor-dimension of ringed topoi, transitivity of inverse-image functors implies that the homomorphisms (2.12.1) satisfy the formula:

$$f^* g^* = (gf)^*. \tag{2.12.1.1}$$

[print 289]Suppose now that we are in situation (2.11.1) (resp. (2.11.2)), with f of finite tor-dimension. Then (III 4.8.1 and 5.8 b))[201] the functor Rf_* sends $D(X)_{\underline{\text{parf}}}$ to $D(Y)_{\underline{\text{parf}}}$, and therefore defines a homomorphism

$$f_* : K^\bullet(X) \longrightarrow K^\bullet(Y). \tag{2.12.3}$$

This is not in general a ring homomorphism. However, for $y \in K_\bullet(Y)$ (resp. $K^\bullet(Y)$) and[202] $x \in K^\bullet(X)$, we have the formula

$$f_*(f^*(y)x) = yf_*(x), \tag{2.12.4}$$

where f^* is the homomorphism (2.12.1) (resp. (2.7.1)). The proof is analogous to that of (2.11.1.2). If $g : Y \to Z$ is a morphism satisfying the same hypotheses as f, the homomorphisms (2.12.3) satisfy the transitivity formula:

$$g_*f_* = (gf)_*. \tag{2.12.3.1}$$

The homomorphism (2.12.3) is one of the essential ingredients of the Riemann–Roch theorem, in both Algebraic Geometry and Analytic Geometry (for indications in the analytic case, see (Exposé 0, 5)).

Suppose that X and Y are separated and divisorial schemes (II 2.2.5) (as is the case, for example, when X and Y possess ample invertible sheaves). Then (2.9 (i)) $K^\bullet(X)$ (resp. $K^\bullet(Y)$) is identified with $K^\bullet(X)_{\text{naïf}}$ (resp. $K^\bullet(Y)_{\text{naïf}}$), and (2.12.3) therefore defines a homomorphism

$$f_* : K^\bullet(X)_{\text{naïf}} \longrightarrow K^\bullet(Y)_{\text{naïf}}. \tag{2.12.3'}$$

As already explained in (Exposé 0, 3.1), the definition of (2.12.3') was by no means evident a priori. Indeed, from the standpoint of our Seminar, it is one of the main motivations for the sorites of Exposé I through Exposé IV.

[print 290]

3. Supplements on the Grothendieck groups of schemes(1)

Proposition 3.1.0. Let

$$\begin{array}{ccc} X & \xleftarrow{g} & X' \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle f'} \\ Y & \xleftarrow{h} & Y' \end{array}$$

be a Cartesian square of schemes, where Y is quasi-compact and f quasi-compact and quasi-separated, and let $E \in \operatorname{ob} D^b(X)$ be a complex with quasi-coherent cohomology. Suppose X and Y' tor-independent over Y (III 1.5). Suppose moreover either h of finite tor-dimension or E of finite flat amplitude rel. to f. Under these conditions, the base-change arrow (cf. SGA 4 XVII 4)

$$Lh^*\, Rf_*E \longrightarrow Rf'_*\, Lg^*E$$

is defined and is an isomorphism.

Proof: To define the arrow, we use the trivial duality formula (SGA 4 XVII 2.3) for h, which says that there is a canonical functorial isomorphism

$$\operatorname{Hom}(Lh^*L, M) \xrightarrow{\sim} \operatorname{Hom}(L, Rh_*M) \tag{$*$}$$

(1) The results of this section will not be used in Exposés V through VIII.

for $L \in \operatorname{ob} D^-(Y)$, $M \in \operatorname{ob} D^+(Y')$. Using (III 3.5.2) and (III 3.7.1), we easily verify that $Lg^*E \in \operatorname{ob} D^b(X')$ and $Rf_*E \in \operatorname{ob} D^-(Y)$, so, applying (*) with $L = Rf_*E$ and $M = Rf'_*Lg^*E$, we are reduced to defining an arrow

$$Rf_*E \longrightarrow Rf_*Rg_*Lg^*E.$$

For this we take the image under Rf_* of the adjunction arrow

$$E \longrightarrow Rg_*Lg^*E,$$

which comes from the trivial duality formula for g. The base-change arrow [print 291]is therefore defined; it remains to show that it is an isomorphism. For this, we may first suppose Y and Y' affine. Then, by the Leray spectral sequence for an open covering of X, we reduce to the case where X and X' are affine. We therefore have a cocartesian square of rings

$$\begin{array}{ccc} B & \longrightarrow & B' \\ \uparrow & & \uparrow \\ A & \longrightarrow & A', \end{array}$$

and we may suppose that E is defined by a bounded-above complex of B-modules. The tor-independence of A' and B over A, which is written

$$B \otimes^{\mathrm{L}}_A A' \xrightarrow{\sim} B',$$

then implies that in $D(A')$ we have

$$E \otimes^{\mathrm{L}}_A A' \xrightarrow{\sim} E \otimes^{\mathrm{L}}_B B',$$

whence the proposition.

Translating (3.1.0) into the language of Grothendieck groups, we obtain:

Proposition 3.1.1. Let

$$\begin{array}{ccc} X & \xleftarrow{g} & X' \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle f'} \\ Y & \xleftarrow{h} & Y' \end{array}$$

[print 292]be a Cartesian square of schemes. Suppose Y and Y' quasi-compact, X and Y' tor-independent over Y (III 1.5), f proper, and pseudo-coherent, and the finiteness conjecture (III 2.1) satisfied for f and f' (for example Y and Y' noetherian, or f projective). Then, if f is perfect (resp. h of finite tor-dimension), we have, for $x \in K^\bullet(X)$ (resp. $K_\bullet(X)$),

$$h^*f_*(x) = f'_*g^*(x),$$

where h^*, g^* are defined by (2.7.1) (resp. (2.12.1)), and f_*, f'_* by (2.12.3) (resp. (2.11.1.1)) (note that by (III 4.7.1) f' is perfect if f is, and that by (III 3.5.1) g has finite tor-dimension if h does).

3.2. Passage to the limit for $K^\bullet_{\text{naïf}}$ and $K_\bullet$[204] Let $(X_i, f_{ij})_{i\in I}$ be a filtered projective system of quasi-compact and quasi-separated schemes, with affine transition morphisms (cf. EGA IV 8.2.2). Put $X = \varprojlim X_i$, and denote by $f_i : X \to X_i$ the canonical homomorphism.

3.2.1. The homomorphisms (2.7.2) $f_i^* : K^\bullet(X_i)_{\text{naïf}} \longrightarrow K^\bullet(X)_{\text{naïf}}$ define, by the transitivity formula (2.7.3), a homomorphism

$$\varinjlim K^\bullet(X_i)_{\text{naïf}} \longrightarrow K^\bullet(X)_{\text{naïf}}. \tag{3.2.1.1}$$

I claim that the homomorphism (3.2.1.1) is an isomorphism. Indeed, let E be a locally free $\underline{O}_X$-Module of finite type. By (EGA IV 8.5.2 and 8.5.5), there is an index i and a locally free $\underline{O}_{X_i}$-Module of finite type E_i such that $E \simeq f_i^*(E_i)$. By (EGA IV 8.5.2.5), the image of $\mathrm{cl}(E_i)$ in $\varinjlim K^\bullet(X_i)_{\mathrm{naïf}}$ is independent of the choice of E_i; denote it by $u(E)$. By (EGA IV 8.5.2 and 8.5.6), the map $u : \mathrm{ob}\,\mathrm{Loclib}(X) \longrightarrow \varinjlim K^\bullet(X_i)_{\mathrm{naïf}}$ [203] thus defined is additive (1.5); [print 293]it therefore defines a homomorphism $K^\bullet(X)_{\mathrm{naïf}} \longrightarrow \varinjlim K^\bullet(X_i)_{\mathrm{naïf}}$, which is immediately seen, by (EGA IV 8.5.2), to be inverse to (3.2.1.1), proving our assertion.

3.2.2. In the same way, one defines a canonical homomorphism

$$\varinjlim K^\bullet(X_i) \longrightarrow K^\bullet(X). \tag{3.2.2.1}$$

Using Deligne's localization technique (SGA 4 XVII), which implies that the category of perfect complexes on X is equivalent to the category $\varinjlim$ of the analogous categories $\mathrm{Parf}(X_i)$, one again finds that (3.2.2.1) is an isomorphism.

3.2.3. Suppose the X_i and X noetherian, and the morphisms f_{ij} flat (the f_i then are as well). Then the homomorphisms (2.12.1) $f_i^* : K_\bullet(X_i) \longrightarrow K_\bullet(X)$ define an isomorphism [205]

$$\varinjlim K_\bullet(X_i) \xrightarrow{\sim} K_\bullet(X). \tag{3.2.3.1}$$

The proof is analogous to that of (3.2.1), using the fact (2.4) that $K_\bullet(X) \simeq K_\bullet(X)_{\mathrm{naïf}}$ is computed in terms of coherent sheaves on X.

Since the module structures of the $K_\bullet(X_i)$ over the $K^\bullet(X_i)_{\mathrm{naïf}}$ are compatible with the inverse images f_{ij}^* by (2.7 a)) and (2.12.2), we find, by passing to the limit, that $\varinjlim K_\bullet(X_i)$ has a module structure over $\varinjlim K^\bullet(X_i)_{\mathrm{naïf}}$; this is compatible, via the isomorphisms (3.2.1.1) and (3.2.3.1) with[206] the module structure of $K_\bullet(X)$ over $K^\bullet(X)_{\mathrm{naïf}}$. In other words, we have a commutative diagram, where the vertical arrows are defined by (3.2.1.1) and (3.2.3.1)[207]:

$$\begin{array}{ccc} \varinjlim K^\bullet(X_i)_{\mathrm{naïf}} \otimes \varinjlim K_\bullet(X_i) & \to & \varinjlim K_\bullet(X_i) \\ \downarrow\simeq & & \downarrow\simeq \\ K^\bullet(X)_{\mathrm{naïf}} \otimes K_\bullet(X) & \longrightarrow & K_\bullet(X). \end{array}$$

[print 294]3.3. Relative Grothendieck groups


3.3.1. Let $f : X \longrightarrow Y$ be a locally finite type morphism of schemes. We shall put:

$$K^\bullet(f) = k(D(f)_{\mathrm{parf}}),$$
$$K_\bullet(f) = k(D^b(f)_{\mathrm{coh}}),$$

where $D^b(f)_{\mathrm{coh}}$ denotes the category of complexes pseudo-coherent rel. to f and with bounded cohomology (III 1.2), and $D(f)_{\mathrm{parf}}$ the category of complexes of finite perfect amplitude rel. to f (III 4.2). When no ambiguity is possible, we shall sometimes write $K^\bullet(X/Y)$ (resp. $K_\bullet(X/Y)$) instead of $K^\bullet(f)$ (resp. $K_\bullet(f)$).

The groups $K^\bullet(f)$ and $K_\bullet(f)$ do not possess the rich structure of their "absolute" analogues $K^\bullet(X)$ and $K_\bullet(X)$, since the tensor product does not in general define a homomorphism $K^\bullet(f) \otimes K^\bullet(f) \longrightarrow K^\bullet(f)$ (resp. $K^\bullet(f) \otimes K_\bullet(f) \longrightarrow K_\bullet(f)$). On the other hand, they satisfy interesting variance properties, which we shall review.

3.3.2. Let



be a commutative triangle of locally finite type morphisms of schemes.

3.3.2.1. If g is pseudo-coherent, we have $K_\bullet(f) = K_\bullet(h)$ (III 1.12). If g is smooth and X quasi-compact, we have $K^\bullet(f) = K^\bullet(h)$ (III 4.4.1).

3.3.2.2. By (III 1.11 and 4.5), the functor $\otimes^{\mathrm{L}}_Y$ defines pairings

$$D(f)_{\underline{\mathrm{parf}}} \times D^b(g)_{\mathrm{coh}} \longrightarrow D^b(h)_{\mathrm{coh}},$$
$$D(f)_{\underline{\mathrm{parf}}} \times D(g)_{\underline{\mathrm{parf}}} \longrightarrow D(h)_{\underline{\mathrm{parf}}},$$

which induce homomorphisms

$$K^\bullet(f) \otimes K_\bullet(g) \longrightarrow K_\bullet(h),$$
$$K^\bullet(f) \otimes K^\bullet(g) \longrightarrow K^\bullet(h).$$

[print 295]We deduce in particular (by putting $X = Y$, $f = \mathrm{Id}$) that $K_\bullet(g)$ is a $K^\bullet(Y)$-module. On the other hand, if f is perfect, multiplication by $\mathrm{cl}(\underline{O}_X) \in K^\bullet(f)$ defines $K^\bullet(Y)$-linear homomorphisms

$$f^* : K_\bullet(Y/S) \longrightarrow K_\bullet(X/S),$$

and

$$f^* : K^\bullet(Y/S) \longrightarrow K^\bullet(X/S).$$

3.3.2.3. Suppose S quasi-compact, X and Y of finite type over S, and f proper. Then, subject to the validity of the finiteness conjecture (III 2.1), hence in particular when Y is noetherian or f is projective, the functor Rf_* sends $D^b(h)_{\mathrm{coh}}$ to $D^b(g)_{\mathrm{coh}}$, and therefore defines a homomorphism

$$f_* : K_\bullet(X/S) \longrightarrow K_\bullet(Y/S).$$

If moreover f is perfect, Rf_* sends $D(h)_{\underline{\mathrm{parf}}}$ to $D(g)_{\underline{\mathrm{parf}}}$ (III 4.8), and defines a homomorphism

$$f_* : K^\bullet(X/S) \longrightarrow K^\bullet(Y/S).$$

From the projection formula (III 3.7) we deduce the formula

$$f_*(ax) = af_*(x)$$

for f proper, $a \in K^\bullet(S)$, $x \in K_\bullet(X/S)$[208] (resp. f proper and perfect, $a \in K_\bullet(S)$, $x \in K^\bullet(X/S)$).

3.3.2.4. Suppose S noetherian, X and Y of finite type over S, and f compactifiable and perfect. Then (III 4.9.1) the functor $f^!$ sends $D(g)_{\underline{\mathrm{parf}}}$ to $D(h)_{\underline{\mathrm{parf}}}$, and defines a homomorphism

$$f^! : K^\bullet(Y/S) \longrightarrow K^\bullet(X/S).$$

3.3.3. Under the hypotheses of (III 4.7), the functor $\otimes^{\mathrm{L}}_S$ defines pairings

$$D(f_1)_{\underline{\mathrm{parf}}} \times D^b(f_2)_{\mathrm{coh}} \longrightarrow D^b(f)_{\mathrm{coh}},$$
$$D(f_1)_{\underline{\mathrm{parf}}} \times D(f_2)_{\underline{\mathrm{parf}}} \longrightarrow D(f)_{\underline{\mathrm{parf}}},$$

which induce homomorphisms

$$K^\bullet(f_1) \otimes K_\bullet(f_2) \longrightarrow K_\bullet(f),$$
$$K^\bullet(f_1) \otimes K^\bullet(f_2) \longrightarrow K^\bullet(f).$$

[print 296](Of course, for $X_i = Y_i = S$, we recover the $K^\bullet(S)$-module structure of $K_\bullet(S)$.)

If X is flat over S, we deduce, for $n \geq 1$, homomorphisms

$$K^\bullet(X/S)^{\otimes n} \longrightarrow K^\bullet((X/S)^n),$$

which should in a sense replace the absent ring structure on $K^\bullet(X/S)$.

BIBLIOGRAPHY

[1] Deligne, P., papiers secrets.

[2] Grauert, H., Ein Theorem der Analytischen Garbentheorie ..., Pub. Math. IHES, n°5 (1960).

[3] Grothendieck, A., Théorèmes de finitude pour la cohomologie des faisceaux, Bull. Soc. Math. France, 84, 1956, p.1-7.

[V] Verdier, J.-L., Catégories dérivées, IHES (1964), miméographié.

Exposé V
GENERALITIES ON λ-RINGS

by P. BERTHELOT

The theory of λ-rings is useful whenever one uses rings such as the ring of classes of vector bundles on a ringed site, the representation ring of a group, etc..., for which the exterior-power operations define maps λ^i. We shall use it extensively in the remainder of the Seminar, so it is necessary first to develop its general formalism. This exposé is thus entirely independent of the rest of the seminar and can serve users of λ-rings outside algebraic geometry.

The first section essentially gives Hirzebruch's theory of "multiplicative sequences of polynomials" (cf. [2]); the functorial viewpoint adopted here has the advantage, besides its generality, of considerably reducing the calculations. The following sections will show how, to every augmented λ-ring, one can associate a "Chern ring" that permits the development of a theory of Chern classes. This is the theory that will enable us to state the Riemann–Roch theorem in the general form of Exp. VIII.

1. Universal polynomials

1.1. Let K be a commutative ring with unit, fixed throughout this section. We denote by $\mathcal{C}$ the category of filtered commutative K-algebras with unit, whose filtration is decreasing and such that the group of elements of filtration zero is the whole algebra. We define a functor on $\mathcal{C}$, with values in the category of groups [print 298]that are abelian, as follows: for $A \in \mathrm{Ob}(\mathcal{C})$, we associate to A the set of formal series $\psi \in A[[t]]$ satisfying the following conditions: if $\psi = a_0 + a_1 t + \ldots + a_n t^n + \ldots$, then

$$a_0 = 1; \qquad \forall i \geq 1, \quad a_i \in \mathrm{Fil}^i A. \tag{1.1.1}$$

This set will be denoted by $\widehat{G}_K(A)$, or by $\widehat{G}(A)$ when no confusion is possible; it is clear that it is an abelian group under multiplication of formal series, and that this does define a functor

$$\widehat{G}_K : \underline{K\text{-Alg.fil.}} \longrightarrow \underline{\mathrm{Ab}}.$$

For A endowed with the coarse filtration, $\widehat{G}(A)$ is the group of formal series of augmentation 1, which will also be denoted by $1 + A[[t]]^+$.

Lemma 1.2. The functor $\widehat{G}_K$ is representable.

Let $(X_i)_{i\in N,\ i\geq 1}$ be a family of indeterminates; we assign the weight i to X_i, and filter the ring $K[X_i]_{i\geq 1}$ by weight with respect to the X_i. We then define a bijection:

$$\widehat{G}_K(A) \xrightarrow{\sim} \mathrm{Hom}_{K\text{-Alg.fil.}}(K[X_i]_{i\geq 1}, A)$$

by associating to every series $\psi = 1 + a_1 t + \ldots + a_n t^n + \ldots \in \widehat{G}_K(A)$ the homomorphism f_ψ determined by $f_\psi(X_i) = a_i$. This homomorphism is compatible with the filtrations, and we thus obtain an isomorphism of functors.

Put $\phi = 1 + X_1 t + \ldots + X_n t^n + \ldots$; for every filtered K-algebra A and every series $\psi \in \widehat{G}(A)$, there exists exactly one homomorphism [print 299]f_ψ from $K[X_i]_{i\geq 1}$ to A such that $\widehat{G}(f_\psi)(\phi) = \psi$.

1.3. For every $A \in \mathrm{Ob}(\mathcal{C})$ we define a filtration on $\widehat{G}(A)$ as follows: a series $\psi = 1 + a_1 t + \ldots$ will be said to be of filtration $\geq k$ if $a_i = 0$ for $1 \leq i < k$. We denote the subgroup so obtained by $F^k\widehat{G}(A)$; this is a functor on $\mathcal{C}$, and from now on we shall regard $\widehat{G}$ as a functor with values in the category Ab.fil. of filtered abelian groups by means of this filtration (with $F^0\widehat{G} = \widehat{G}$).

Let H be a functor on $\mathcal{C}$ with values in Ab fil; we shall assume that H satisfies the following two conditions:

i) $\forall A \in \mathrm{Ob}(\mathcal{C})$, $H(A)$ is separated and complete.

ii) If $A \in \mathrm{Ob}(\mathcal{C})$ is a graded K-algebra, endowed with the filtration associated with its grading, and if G is a finite group acting on A by graded morphisms, then $H(A^G) \xrightarrow{\sim} H(A)^G$ (the exponent G denoting invariants under G)

In fact, we shall need these conditions only when A is a polynomial ring and G is a finite product of symmetric groups acting by permutation of the indeterminates.

Examples: 1) $H(A) = \widehat{G}(A)$. 2) $H(A) = 1 + A[[t]]^+$; this functor will be denoted by $\widehat{G}_g$ (g, the initial of "grossière"). 3) $H(A) = \widehat{G}_a(A)$, where $\widehat{G}_a(A)$ denotes the group, under addition, of formal series such that $a_0 = 0$, $a_i \in \mathrm{Fil}^i A$ for $i \geq 1$, endowed with the filtration analogous to that of $\widehat{G}$.

1.4. For every n, we shall denote by f_n the endomorphism of $K[X_1, \ldots, X_n]$ [print 300]defined by $f_n(X_i) = X_i$ for $i < n$, $f_n(X_n) = 0$. It factors through the inclusion of $K[X_1, \ldots, X_{n-1}]$ in $K[X_1, \ldots, X_n]$, which shows that the homomorphism from $H(K[X_1, \ldots, X_{n-1}])$ to $H(K[X_1, \ldots, X_n])$ is injective, and will allow us to identify the former with a subgroup of the latter; similarly, we shall regard $H(K[X_1, \ldots, X_n])$ as contained in $H(K[X_i]_{i \geq 1})$.

Notice that f_1 is precisely the augmentation of $K[X]$.

1.5. We embed $K[X_1, \ldots, X_n]$ in $K[T_1, \ldots, T_n]$ by sending X_i to the i-th symmetric function in the T_j; letting the symmetric group $\mathfrak{S}_n$ act by permutation of the T_j, and using the theorem on symmetric polynomials and (1.3.ii)), we see that the resulting homomorphism from $H(K[X_1, \ldots, X_n])$ to $H(K[T_1, \ldots, T_n])$ is injective, which will again allow us to identify the former with a subgroup of the latter:

$$H(K[X_1, \ldots, X_n]) \simeq H(K[T_1, \ldots, T_n]^{\mathfrak{S}_n}) \xrightarrow{\sim} H(K[T_1, \ldots, T_n])^{\mathfrak{S}_n} \subset H(K[T_1, \ldots, T_n])$$

We denote by

$$u_i^n : K[X] \longrightarrow K[T_1, \ldots, T_n]$$

the homomorphism defined by $u_i^n(X) = T_i$, and put

$$u_H^n = \sum_{i=1}^{n} H(u_i^n) : H(K[X]) \longrightarrow H(K[T_1, \ldots, T_n]).$$

The symmetric group permutes the morphisms $H(u_i^n)$, and therefore leaves the morphism u_H^n invariant, which consequently takes its values in [print 301]$H(K[X_1, \ldots, X_n])$ by (1.3.ii)).

Finally, observe that for $H = \widehat{G}$, $u_H^n(1 + Xt) = \prod_{i=1}^{n} (1 + T_i t) = 1 + X_1 t + \ldots + X_n t^n$.

Theorem 1.6. The map from $\mathrm{Hom}_{\mathrm{Ab\,fil}}(\widehat{G}_K, H)$ to $H(K[X])$ which, to a morphism of functors, assigns its value on the element $1 + Xt$ of $\widehat{G}_K(K[X])$, is injective; its image is the subgroup of $H(K[X])$ formed by the elements θ such that (with the notation of 1.4. for the f_n):

i) $H(f_1)(\theta) = 0$.

ii) $\forall n \geq 1, \quad H(f_n)(u_H^n(\theta)) \equiv u_H^n(\theta) \bmod F^n H(K[X_1, \ldots, X_n])$.

Let us first prove injectivity. By Lemma 1.2, it is enough to see that a morphism of functors Φ vanishing on $1+Xt$ vanishes on $\varphi = 1 + X_1 t + \ldots + X_n t^n + \ldots$. The commutative diagram:

$$\begin{array}{ccc} \widehat{G}(K[X]) & \xrightarrow{u_{\widehat{G}}^n} & \widehat{G}(K[X_1,\ldots,X_n]) \\ {\scriptstyle\Phi}\downarrow & & \downarrow{\scriptstyle\Phi} \\ H(K[X]) & \xrightarrow{u_H^n} & H(K[X_1,\ldots,X_n]) \end{array} \tag{1.6.1}$$

shows that for every n, $\Phi(1 + X_1 t + \ldots + X_n t^n) = 0$.

Now $\varphi \equiv 1 + X_1 t + \ldots + X_n t^n$ mod. $F^{n+1}\widehat{G}$; hence $\Phi(\varphi) \equiv \Phi(1 + \ldots + X_n t^n)$ mod. $F^{n+1}H$, and consequently $\Phi(\varphi) \equiv 0$ mod. $F^{n+1}H$, for every n; since $H(K[X_i]_{i\geq 1})$ is separated and complete, $\Phi(\varphi) = 0$.

Now let Φ be any element of $\mathrm{Hom}(\widehat{G}, H)$, and put [print 302]$\theta = \Phi(1+Xt)$; let us show that θ satisfies conditions i) and ii).

i): $H(f_1)(\theta) = H(f_1) \circ \Phi(1+Xt) = \Phi \circ \widehat{G}(f_1)(1+Xt) = \Phi(1) = 0$.

ii): $u_H^n(\theta) = \Phi(1 + \ldots + X_n t^n)$ by (1.6.1); now $1 + \ldots + X_n t^n \equiv 1 + \ldots + X_{n-1} t^{n-1}$ mod. $F^n\widehat{G}$, and $1 + \ldots + X_{n-1} t^{n-1} = \widehat{G}(f_n)(1 + \ldots + X_n t^n)$. Since $\Phi \circ \widehat{G}(f_n) = H(f_n) \circ \Phi$, and since Φ is a morphism of filtered functors, we obtain ii).

Finally, we must show that if $\theta \in H(K[X])$ satisfies i) and ii), there exists $\Phi \in \mathrm{Hom}(\widehat{G}, H)$ such that $\theta = \Phi(1+Xt)$.

We first determine the value of Φ on the element $1 + \ldots + X_n t^n$ of $\widehat{G}(K[X_1,\ldots,X_n])$ by means of (1.6.1); thus, by definition, $\Phi(1 + \ldots + X_n t^n) = \sum_{i=1}^{n} \Phi(1 + T_i t)$, the $\Phi(1+T_i t)$ being known by functoriality through the morphisms u_i^n.

We then glue the $\Phi(1 + \ldots + X_n t^n)$ for varying n by means of the commutative diagram:

$$\begin{array}{ccc} K[X_1,\ldots,X_n] & \xrightarrow{f_n} & K[X_1,\ldots,X_n] \\ {\scriptstyle\sigma_n}\downarrow & & \downarrow{\scriptstyle\sigma'_n} \\ K[T_1,\ldots,T_n] & \xrightarrow{f'_n} & K[T_1,\ldots,T_n], \end{array} \tag{1.6.2}$$

where f'_n is determined by $f'_n(T_i) = T_i$ if $i < n$, $f'_n(T_n) = 0$, where σ_n sends X_i to the i-th symmetric function of the T_j, and σ'_n sends X_i to the i-th symmetric function of $T_1, \ldots, T_{n-1}$ for $i < n$, and X_n to 0.

Indeed, this gives:

$$\begin{aligned} H(f_n)\big(\Phi(1 + \ldots + X_n t^n)\big) &= H(f'_n)\left(\sum_{i=1}^{n} \Phi(1+T_i t)\right) = \sum_{i=1}^{n-1} \Phi(1+T_i t) \text{ (by 1.6.i))} \\ &= \Phi(1 + \ldots + X_{n-1} t^{n-1}) \end{aligned}$$

[print 303](working in $H(K[T_1,\ldots,T_{n-1}])$ by means of the identifications 1.4).

In view of 1.6.ii), $\Phi(1 + \ldots + X_n t^n) \equiv \Phi(1 + \ldots + X_{n-1} t^{n-1})$ mod. $F^n H(K[X_i]_{i\geq 1})$, and the $\Phi(1 + \ldots + X_n t^n)$ define an element ψ of $H(K[X_i]_{i\geq 1})$, since this is a separated complete group; we therefore put $\Phi(\varphi) = \psi$.

Finally, let $A \in \mathrm{Ob}(\mathcal{C})$, and $\chi \in \widehat{G}(A)$; we put $\Phi(\chi) = H(f_\chi)(\psi)$, where f_χ is the homomorphism from $K[X_i]_{i\geq 1}$ to A defined in 1.2. Since we have already defined Φ on the elements $1 + \ldots + X_n t^n = \varphi_n$, we must verify that the two definitions give the same result, that is:

$$H(f_{\varphi_n})(\psi) = \Phi(1 + \ldots + X_n t^n).$$

Now, for every p, we have: $\psi \equiv \Phi(1+\ldots+X_p t^p)$ mod. $F^{p+1}H(K[X_i]_{i\geq 1})$. Hence, for every $p \geq n$; $H(f_{\varphi_n})(\psi) \equiv H(f_{\varphi_n})(\Phi(1+\ldots+X_p t^p))$ mod. $F^{p+1}H(K[X_i]_{i\geq 1})$. Using the identifications of 1.4 and the gluing of the $\Phi(\varphi_m)$ for varying m, we obtain: $H(f_{\varphi_n})(\Phi(\varphi_p)) = \Phi(\varphi_n)$. Thus, finally:

$$\forall p \geq n, \quad H(f_{\varphi_n})(\psi) \equiv \Phi(1+\ldots+X_n t^n) \text{ mod. } F^p H(K[X_i]_{i\geq 1}).$$

Since H is separated, we obtain the required equality.

We have thus defined a system of maps $\Phi_A : \widehat{G}(A) \to H(A)$; to verify that they constitute a morphism of functors (with values in $\underline{\text{Ens}}$), we must show that for every morphism $h : A \to A'$, the diagram:

$$\begin{array}{ccc} \widehat{G}(A) & \xrightarrow{\widehat{G}(h)} & \widehat{G}(A') \\ {\scriptstyle \Phi_A}\downarrow & & \downarrow{\scriptstyle \Phi_{A'}} \\ H(A) & \xrightarrow{H(h)} & H(A') \end{array}$$

[print 304]is commutative. Let $\chi \in \widehat{G}(A)$; then: $h \circ f_\chi = f_{\widehat{G}(h)(\chi)}$; hence: [209]

$$H(h)(\Phi_A(\chi)) = H(h) \circ H(f_\chi)(\psi) = H(f_{\widehat{G}(h)(\chi)})(\psi) = \Phi_{A'}(\widehat{G}(h)(\chi)).$$

It therefore remains to show that the morphism of functors thus constructed is compatible with the structures of filtered groups.

Suppose that $\chi \in F^n\widehat{G}(A)$. To say that χ has filtration $\geq n$ is equivalent to saying that $f_\chi : K[X_i]_{i\geq 1} \to A$ factors through the endomorphism g_n of $K[X_i]_{i\geq 1}$ defined by: $g_n(X_i) = 0$ if $1 \leq i \leq n-1$, and $g_n(X_i) = X_i$ if $i \geq n$. To show that Φ is compatible with the filtrations, it therefore suffices to show that $H(g_n)(\Phi(\varphi)) \in F^n H$. Now, by construction:

$$\Phi(\varphi) \equiv \Phi(1+\ldots+X_{n-1}t^{n-1}) \text{ mod. } F^n H,$$

hence:

$$H(g_n)(\Phi(\varphi)) \equiv H(g_n)(\Phi(1+\ldots+X_{n-1}t^{n-1})) = 0 \text{ mod. } F^n H.$$

To verify the additivity of Φ, it suffices to do this for the elements of finite degree of $\widehat{G}(A)$, and then pass to the limit. Thus let χ_1, χ_2 be two elements of degree n of $\widehat{G}(A)$, and χ their product. Consider the commutative diagram:

$$\begin{array}{ccc} K[X_1,\ldots,X_{2n}] & \xrightarrow{f_\chi} & A \\ \downarrow & \nearrow_{f'_\chi} & \\ (K[T_1,\ldots,T_{2n}])^{\mathfrak{S}_n\times\mathfrak{S}_n} & & \end{array},$$

where the arrows are defined as follows: $\mathfrak{S}_n \times \mathfrak{S}_n$ acts on $K[T_1,\ldots,T_{2n}]$ [print 305]in such a way that the first factor permutes $T_1,\ldots,T_n$ and the second $T_{n+1},\ldots,T_{2n}$; $K[X_1,\ldots,X_{2n}]$ is embedded in the ring of invariants by sending X_j to the j-th symmetric function of the T_i; f' sends the i-th symmetric function of $T_1,\ldots,T_n$ (resp. $T_{n+1},\ldots,T_{2n}$) to the i-th coefficient of χ_1 (resp. χ_2). Then:

$$\Phi(\chi) = H(f_\chi)\left(\Phi\left(\prod_{i=1}^{2n}(1+T_i t)\right)\right) = H(f'_\chi)\left(\Phi\left(\prod_{i=1}^{2n}(1+T_i t)\right)\right);$$

$$\Phi(\chi_1) = H(f'_\chi)\left(\Phi\left(\prod_{i=1}^{n}(1+T_i t)\right)\right) \quad (\text{resp. } \chi_2,\ i = n+1,\ldots,2n).$$

Since $H(f'_\chi)$ is a homomorphism of groups, we are reduced to the case where $\chi = \prod_{i=1}^{2n}(1+T_it)$, and in this case the additivity of Φ follows from the definition of the $\Phi(1+X_1t+\ldots+X_kt^k)$.

1.7. We can generalize the preceding considerations to a product of functors $\widehat{G}$; the functor $(\widehat{G})^k$ is representable by the ring $K[(X_i^{(j)})]$, $j=1,\ldots,k$, and $i\geq 1$, where $X_i^{(j)}$ has weight i. We denote by $\varphi^{(j)}$ the series $1+X_1^{(j)}t+\ldots+X_n^{(j)}t^n+\ldots$; the element $(\varphi^{(j)})_j$ of $\left(\widehat{G}(K[(X_i^{(j)})])\right)^k$ has the universal property analogous to that of φ (cf. 1.2).

We can introduce the morphisms $f_n^{(j)}$, $u_H^{n,(j)}$ as before, by repeating the construction for each family $(X_i^{(j)})_{i\geq 1}$ separately. We obtain a result analogous to Theorem 1.6 by seeking the homomorphisms of functors from $(\widehat{G})^k$ to H that are $\mathbb{Z}$-multilinear and compatible with the filtrations in each of the k arguments separately: they are determined by their value on the element [print 306]$(1+X^{(j)}t)_{j=1,\ldots,k}$[210] of $\left(\widehat{G}(K[(X^{(j)})_j])\right)^k$; an element θ of $H(K[(X^{(j)})_j])$ determines a morphism of functors if and only if it satisfies:

$$\text{i)}\quad \forall j,\quad H(f_1^{(j)})(\theta)=0$$
$$\text{ii)}\quad \forall n,j,\quad H(f_n^{(j)})(u_H^{n,(j)}(\theta))\equiv u_H^{n,(j)}(\theta)\ \text{mod. } F^nH.$$

The reader will readily verify that, apart from complications of notation, it suffices to copy the proof of 1.6.

1.8. We denote by $\mathcal{C}_0$ the category of K-algebras, and by

$$i:\mathcal{C}_0\longrightarrow\mathcal{C},$$

the functor that assigns to a K-algebra A the K-algebra A endowed with the coarse filtration; we denote by

$$g:\mathcal{C}\longrightarrow\mathcal{C}_0,$$

the functor "forget the filtration." These functors satisfy the following relations:

$$g\circ i=\mathrm{Id}_{\mathcal{C}_0};\qquad \widehat{G}\circ i\circ g=\widehat{G}_g;\qquad \widehat{G}_g\circ i\circ g=\widehat{G}_g$$

(cf. 1.3, ex. 2)

Lemma 1.9. Let $H:\mathcal{C}_0\to \underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}$ be a functor on $\mathcal{C}_0$; then the canonical homomorphism:

$$\mathrm{Hom}_{\underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}}(\widehat{G},H\circ g)\longrightarrow\mathrm{Hom}_{\underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}}(\widehat{G}\circ i,H)$$

[print 307]is an isomorphism.

To see that it is injective, we use the sequence of canonical homomorphisms:

$$\mathrm{Hom}(\widehat{G},H\circ g)\longrightarrow\mathrm{Hom}(\widehat{G}\circ i,H)\longrightarrow\mathrm{Hom}(\widehat{G}_g,H\circ g)\longrightarrow\mathrm{Hom}(\widehat{G},H\circ g),$$

the last arrow coming from the fact that $\widehat{G}$ is a subfunctor of $\widehat{G}_g$. To verify that the composite homomorphism is the identity, we observe that a morphism of functors: $\widehat{G}\longrightarrow H\circ g$ is determined by its value on the objects of $\mathcal{C}$ whose filtration is coarse; indeed, let $A\in\mathrm{Ob}(\mathcal{C})$, and let A_g be the algebra A endowed with the coarse filtration; the canonical morphism $A\to A_g$ defines a commutative diagram:

$$\begin{array}{ccc}\widehat{G}(A) & \xrightarrow{\Phi_A} & H\circ g(A)\\ \downarrow & & \downarrow\\ \widehat{G}(A_g) & \xrightarrow{\Phi_{A_g}} & H\circ g(A_g),\end{array}$$

showing that Φ_A is the restriction of Φ_{A_g} to $\widehat{G}(A)$.

We next consider the composite homomorphism:

$$\mathrm{Hom}(\widehat{G}\circ i, H) \longrightarrow \mathrm{Hom}(\widehat{G}_g, H\circ g) \longrightarrow \mathrm{Hom}(\widehat{G}, H\circ g) \longrightarrow \mathrm{Hom}(\widehat{G}\circ i, H),$$

and it is clear that this is the identity, giving the asserted result.

Theorem 1.10. Let H be a functor on the category $\mathcal{C}_0$ of K-algebras, with values in the category of separated complete filtered abelian groups, and commuting with invariants under a finite group. Then the map from $\mathrm{Hom}_{\underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}}(\widehat{G}\circ i, H)$ to $H(K[X])$ which, to a morphism of functors, assigns [print 308]its value on the element $1+Xt$ of $\widehat{G}\circ i(K[X])$, is injective; its image is the subgroup of $H(K[X])$ formed by the elements θ such that (with notation analogous to that of 1.4 and 1.5):

i) $H(f_1)(\theta) = 0$;

ii) $\forall n \geq 1, \quad H(f_n)(u_H^n(\theta)) \equiv u_H^n(\theta) \bmod F^n H(K[X_1,\ldots,X_n])$.

The theorem follows immediately from Theorem 1.6 and Lemma 1.9, since $H\circ g(K[X])$ does not depend on the filtration chosen on $K[X]$.

The analogous theorem obtained by replacing $\widehat{G}$ by $(\widehat{G})^k$ is also true (cf. 1.7)

When there is no risk of confusion, we shall denote the functor $\widehat{G}\circ i$ by $\widehat{G}$; this is legitimate, since i identifies $\mathcal{C}_0$ with a subcategory of $\mathcal{C}$.

1.11. We denote by $\mathcal{G}$ the category of graded K-algebras, and by j the functor from $\mathcal{G}$ to $\mathcal{C}$ that assigns to a graded K-algebra B the K-algebra B endowed with the filtration canonically defined by its grading; the "associated graded" functor: $\mathcal{C}\to\mathcal{G}$ will be denoted by Gr. There is an isomorphism of functors $\varepsilon : \mathrm{Id}_{\mathcal{G}} \xrightarrow{\sim} \mathrm{Gr}\circ j$.

Let $B \in \mathrm{Ob}(\mathcal{G})$; we denote by $1+\widehat{B}^+$ the multiplicative group of elements of $\prod_{i=0}^{\infty} B^i$ with augmentation 1. Endowing it with the filtration analogous to that of $\widehat{G}$, we thus define a functor: $\mathcal{G}\to \underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}$.

Furthermore, for every $A\in\mathrm{Ob}(\mathcal{C})$, we define a homomorphism of filtered groups: $\widehat{G}(A) \longrightarrow 1+\widehat{\mathrm{Gr}\,A}^+$, by sending the series $1+\ldots+a_nt^n+\ldots$, [print 309]to the series $1+\ldots+\bar{a}_n+\ldots$, where $\bar{a}_n$ is the class of a_n in $\mathrm{Gr}^n A$. Its kernel is the group $\widehat{G}'(A)$ of formal series such that:

$$a_0 = 1; \qquad \forall i\geq 1, \quad a_i \in \mathrm{Fil}^{i+1} A;$$

we therefore have an isomorphism, functorial in A:

$$\widehat{G}(A)/\widehat{G}'(A) \xrightarrow{\sim} 1+\widehat{\mathrm{Gr}\,A}^+.$$

Lemma 1.12. Let $H : \mathcal{G}\to \underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}$ be a functor on $\mathcal{G}$; then the canonical homomorphism:

$$\rho : \mathrm{Hom}_{\underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}}(1+\widehat{\cdot}^+, H) \longrightarrow \mathrm{Hom}_{\underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}}(\widehat{G}, H\circ \mathrm{Gr}),$$

is an isomorphism.

The homomorphism ρ is defined as the composite:

$$\mathrm{Hom}(1+\widehat{\cdot}^+, H) \longrightarrow \mathrm{Hom}(1+\widehat{\mathrm{Gr}\,\cdot}^+, H\circ\mathrm{Gr}) \longrightarrow \mathrm{Hom}(\widehat{G}, H\circ\mathrm{Gr}),$$

the last arrow being defined by the homomorphism $\widehat{G} \longrightarrow 1+\widehat{\mathrm{Gr}\,\cdot}^+$

We shall construct a morphism in the opposite direction:

$$\rho' : \mathrm{Hom}(\widehat{G}, H\circ\mathrm{Gr}) \longrightarrow \mathrm{Hom}(1+\widehat{\cdot}^+, H).$$

For this, we shall show that every morphism of functors from $\widehat{G}$ to $H \circ \mathrm{Gr}$ factors through the canonical morphism $\widehat{G} \longrightarrow 1 + \widehat{\mathrm{Gr}_{\cdot}}^{+}$, which, by 1.11, is equivalent to showing that it vanishes on $\widehat{G}'$.

Let $A \in \mathrm{Ob}(\mathcal{C})$, $\chi \in \widehat{G}'(A)$; we define a homomorphism of [print 310]filtered K-algebras $K[X_i]_{i\geq 1} \longrightarrow A$ by sending X_1 to 0 and X_i to the $(i-1)$-th coefficient of χ for $i > 1$; using this homomorphism, we see that it suffices to verify that the morphism of functors under consideration vanishes on the element $1 + X_2 t + \ldots + X_{n+1}t^n + \ldots$. We then use the endomorphism f of $K[X_i]_{i\geq 1}$ defined by $f(X_i) = X_{i+1}$ for every i, and the commutative diagram:

$$\begin{array}{ccc} \widehat{G}(K[(X_i)]) & \longrightarrow & H \circ \mathrm{Gr}(K[(X_i)]) \\ {\scriptstyle \widehat{G}(f)}\downarrow & & \downarrow{\scriptstyle H(\mathrm{Gr}(f))} \\ \widehat{G}(K[(X_i)]) & \longrightarrow & H \circ \mathrm{Gr}(K[(X_i)]); \end{array}$$

now $\mathrm{Gr}(f) = \mathrm{Gr}(f_1)$, where f_1 is the augmentation homomorphism of $K[X_i]_{i\geq 1}$; since the latter annihilates φ, the analogous diagram written for f_1 shows that $H(\mathrm{Gr}(f))$ annihilates the image of φ in $H \circ \mathrm{Gr}(K[X_i]_{i\geq 1})$, giving the result.

Passing to the quotient, we therefore associate to every morphism of functors: $\widehat{G} \longrightarrow H \circ \mathrm{Gr}$ a morphism of functors: $1 + \widehat{\mathrm{Gr}_{\cdot}}^{+} \longrightarrow H \circ \mathrm{Gr}$; by composing with j, and using the isomorphism ε, we obtain a morphism: $1 + \hat{\cdot}^{+} \longrightarrow H$, which defines the required homomorphism ρ'. It is immediate to verify that ρ and ρ' are inverses of one another.

Theorem 1.13. Let H be a functor on $\mathcal{G}$, with values in the category of separated complete filtered abelian groups, and commuting with invariants under a finite group. Then the map from $\mathrm{Hom}_{\underline{\mathrm{Ab\,fil}}}(1 + \hat{\cdot}^{+}, H)$ to $H(K[X])$ which, to a morphism of functors, assigns its value on the element [print 311]$1 + X$ of $1 + \widehat{K[X]}^{+}$ is injective; its image is the subgroup of $H(K[X])$ formed by the elements θ such that (notation of 1.4,1.5): [211]

i) $H(f_1)(\theta) = 0$;

ii) $\forall n \geq 1, \quad H(f_n)(u_H^n(\theta)) \equiv u_H^n(\theta) \bmod F^n H(K[X_1, \ldots, X_n])$.

Immediate from 1.6 and 1.12.

There is an analogous statement on replacing $\widehat{G}$ by $(\widehat{G})^k$.

Remark 1.14. If H is a functor on $\mathcal{G}$, with values in the category of abelian groups, commuting with invariants under a finite group, and satisfying the condition:

$$\forall B \in \mathrm{Ob}(\mathcal{G}), \qquad H(B) \xrightarrow{\sim} \varprojlim_n H(B/B^{\geq n}) \quad (\text{where } B^{\geq n} = \prod_{i\geq n} B^i),$$

then condition ii) of 1.13 is always satisfied when we take as the filtration on H the filtration by the kernels of the $H(B) \longrightarrow H(B/B^{\geq n})$ (which is separated and complete). Indeed, let p_n be the canonical homomorphism

$$K[X_1, \ldots, X_n] \longrightarrow K[X_1, \ldots, X_n]/K[X_1, \ldots, X_n]^{\geq n};$$

we have: $p_n \circ f_n = p_n$, hence: $H(p_n)[H(f_n) \circ u_H^n(\theta) - u_H^n(\theta)] = 0$, that is:

$$H(f_n) \circ u_H^n(\theta) - u_H^n(\theta) \in F^n H(K[X_1, \ldots, X_n]).$$

Thus, to specify a morphism of functors from $1 + \hat{\cdot}^{+}$ to H, it suffices to specify an element of $H(K[X])$ satisfying only condition i) of 1.13.

1.15. Let $H : \mathcal{C} \to \underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}$ (resp. $\mathcal{C}_0 \to \underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}$, $\mathcal{G} \to \underline{\mathrm{Ab}}\,\underline{\mathrm{fil}}$) be a functor satisfying the conditions of 1.3 (resp. 1.10, 1.13), and let $\Phi : (\widehat{G})^k \to H$ (resp. $(\widehat{G} \circ i)^k \to H$, $(1+\hat{\cdot}^+)^k \to H$) be a morphism of functors. We have seen [print 312](1.7) that Φ is determined by its value on the element $(\varphi^{(j)})_{j=1..k}$ of $(\widehat{G}(K[X_i^{(j)}]_{\substack{i\geq 1 \\ j=1..k}}))^k$ (resp. ...).. Suppose that H is a functor such as $\widehat{G}, \widehat{G}_g, \widehat{G}_a, 1+\widehat{\mathrm{Gr}\,\cdot}^+$, etc..; the element $\Phi((\varphi^{(j)})_j)$ is a formal series with coefficients in $K[X_i^{(j)}]_{i\geq 1,\ j=1..k}$; the n-th coefficient of this series will be called the n-th universal polynomial of Φ; knowing these polynomials is therefore equivalent to knowing Φ. By the proof of 1.6, these polynomials are themselves determined by the element $\Phi(1 + X^{(1)}t, \ldots, 1 + X^{(k)}t)$, the calculation following the method of the proof: we calculate the elements $\Phi(1 + X_1^{(1)}t + \ldots + X_n^{(1)}t^n, \ldots, 1 + X_1^{(k)}t + \ldots + X_n^{(k)}t^n)$ by decomposing $1 + X_1^{(i)}t + \ldots + X_n^{(i)}t^n$ into $\prod_{j=1}^n (1 + T_j^{(i)}t)$ and using the additivity of Φ; the polynomials thus obtained are symmetric in the $T_j^{(i)}$ (i fixed and j varying) and can be expressed in terms of the $X_j^{(i)}$; by gluing with the aid of the filtration of H, we thus determine the universal polynomials of Φ. If $(\psi^{(1)}, \ldots, \psi^{(k)}) \in (\widehat{G}(A))^k$, the calculation of $\Phi(\psi^{(1)}, \ldots, \psi^{(k)})$ is performed by substituting the coefficients $a_j^{(i)}$ of the $\psi^{(i)}$ for the $X_j^{(i)}$ in the universal polynomials.

Suppose there is an integer m such that the coefficient of t^n in $\theta = \Phi(1 + X^{(1)}t, \ldots, 1 + X^{(k)}t)$ is a polynomial whose isobaric components have weights ranging between n and mn; then the k-th universal polynomial of Φ has isobaric components whose weights range between k and km[212]: this follows immediately from the calculation described above and from the theorem on symmetric functions.

[print 313]Suppose in particular that $H = \widehat{G}$; then the n-th universal polynomial of Φ has no components of weight $< n$. Now consider the canonical morphism: $H \longrightarrow 1 + \widehat{\mathrm{Gr}\,\cdot}^+$; composing with Φ gives a morphism: $(\widehat{G})^k \longrightarrow 1 + \widehat{\mathrm{Gr}\,\cdot}^+$, which, by the proof of 1.12, factors through $(1 + \widehat{\mathrm{Gr}\,\cdot}^+)^k$ and defines a morphism $\Phi : (1 + \hat{\cdot}^+)^k \longrightarrow 1 + \hat{\cdot}^+$; we immediately see that the n-th universal polynomial of Φ is the same as that of $(\widehat{G})^k \longrightarrow 1 + \widehat{\mathrm{Gr}\,\cdot}^+$, that is, the isobaric component of weight n of the n-th universal polynomial of Φ.

The same remarks apply to the case $H = \widehat{G}_a$: we can define a morphism of functors $H \longrightarrow \widehat{\mathrm{Gr}\,\cdot}^+$ (with $\widehat{\mathrm{Gr}\,\cdot}^+ = \hat{\cdot}^+ \circ \mathrm{Gr}$, where $\hat{\cdot}^+$ is the functor defined on $\mathcal{G}$ by $\widehat{B}^+ = \prod_{i\geq 1} B^i$) analogously to 1.11; we then repeat the reasoning and conclusions of the preceding paragraph.

1.16. Let us finish with an example. Suppose that $K = Q$. We define an endomorphism of the functor $1 + \hat{\cdot}^+$, denoted by Todd and called the Todd character, by putting:

$$\mathrm{Todd}(1 + X) = \frac{X}{1 - e^{-X}}.$$

Conditions i) and ii) of Theorem 1.13 are verified immediately, showing that we have indeed defined an endomorphism of $1 + \hat{\cdot}^+$.

2. Definition of λ-rings; examples

In this section, the base ring of Section 1 will be $\mathbb{Z}$ (except in 2.9); we work in the category $\mathcal{C}_0$ (1.8) and denote by $\widehat{G}$ the restriction of $\widehat{G}$ to $\mathcal{C}_0$ (1.10).

[print 314]**Definition 2.1.**: By a pre-λ-ring we mean a ring K, commutative with unit, endowed with a homomorphism of abelian groups:

$$\lambda_t : K \longrightarrow 1 + K[[t]]^+,$$

such that for every $x \in K$, $\lambda_t(x) = 1 + xt + \ldots$

We put $\lambda_t(x) = \sum_{i\geq 0}^{\infty} \lambda^i(x)t^i$; giving a pre-$\lambda$-ring structure on a ring K is therefore equivalent to giving a family of maps λ^i, $i \in N$, from K to K, such that:

$$\begin{aligned}
&\text{i)} \quad \forall x \in K, \quad \lambda^0(x) = 1,\\
&\text{ii)} \quad \forall x \in K, \quad \lambda^1(x) = x,\\
&\text{iii)} \quad \forall x, y \in K,\ \forall n \in N, \quad \lambda^n(x+y) = \sum_{p+q=n} \lambda^p(x)\,.\,\lambda^q(y).
\end{aligned}$$

Let K, K' be two pre-λ-rings, and $f : K \to K'$ a ring homomorphism; we say that f is a homomorphism of pre-λ-rings, or a λ-homomorphism, when the diagram:

$$\begin{array}{ccc}
K & \xrightarrow{\lambda_t} & 1 + K[[t]]^+ \\
{\scriptstyle f}\downarrow & & \downarrow{\scriptstyle \widehat{G}(f)} \\
K' & \xrightarrow{\lambda_t} & 1 + K'[[t]]^+,
\end{array}$$

is commutative.

This is also equivalent to the following relations:

$$\forall x \in K,\ \forall i \in N, \qquad \lambda^i(f(x)) = f(\lambda^i(x)).$$

[print 315]**Examples 2.2.**: a) We can give all the pre-λ-ring structures on $\mathbb{Z}$. For this it suffices to specify $\lambda_t(1)$, which will therefore be a formal series with integer coefficients, subject only to the condition that it begin with $1 + t$.

b) Let $(T, \underline{A})$ be a commutatively ringed topos, and let K be the "Grothendieck group" of $\underline{A}$-Modules locally isomorphic to a direct summand of a free $\underline{A}$-Module of finite type; we know that K can be endowed with a ring structure by means of the tensor product (Exp. IV), and we shall endow it with a pre-λ-ring structure as follows: for every A-Module $\underline{E}$ that is locally a direct summand of a free Module of finite type, we denote by $\lambda^i(E)$ the class in K of the i-th exterior power of $\underline{E}$ (which is also locally a direct summand of a free Module of finite type); this defines a function on the category of $\underline{A}$-Modules that are locally direct summands of a free Module of finite type, with values in the abelian group $1 + K[[t]]^+$; we must then show that this function is additive, which will imply that it factors through K, giving the required structure.

Lemma 2.2.1. Let $0 \longrightarrow \underline{E}' \xrightarrow{u} \underline{E} \xrightarrow{v} \underline{E}'' \longrightarrow 0$ be an exact sequence of $\underline{A}$-Modules that is locally split (for example such that E'' is locally a direct summand of a free Module of finite type). There exists on $\wedge^n(\underline{E})$ a finite filtration, and isomorphisms:

$$\wedge^{n-p}(\underline{E}') \otimes_A \wedge^p(\underline{E}'') \xrightarrow{\sim} \mathrm{Gr}^p \wedge^n(\underline{E}).$$

We take as $F^p \wedge^n (\underline{E})$ the image of the canonical morphism:

$$\wedge^{n-p}(\underline{E}') \otimes_A \wedge^p(\underline{E}) \longrightarrow \wedge^n(\underline{E}) \quad \text{for } 0 \leq p \leq n\ ; \quad \text{if } p \geq n,$$

[print 316]$F^p \wedge^n (\underline{E}) = \wedge^n(\underline{E})$, and if $p < 0$, $F^p \wedge^n (\underline{E}) = 0$. This does define a finite filtration on $\wedge^n(\underline{E})$.

We define a homomorphism:

$$f_p : \wedge^{n-p}(\underline{E}') \otimes_A \wedge^p(\underline{E}'') \longrightarrow F^p \wedge^n (\underline{E})/F^{p-1} \wedge^n (\underline{E})$$

by putting $f_p(x'_1 \wedge \ldots \wedge x'_{n-p} \otimes x''_1 \wedge \ldots \wedge x''_p) = \mathrm{cl}_{\mathrm{Gr}^p}\big(u(x'_1) \wedge \ldots \wedge u(x'_{n-p}) \wedge x_1 \wedge \ldots \wedge x_p\big)$, which visibly does not depend on the choice of the x_i lifting the given sections x''_i of $\underline{E}''$. When the exact sequence is locally split, f_p is an isomorphism, since locally $\wedge^n(\underline{E}) \simeq \bigoplus_{p+q=n} \wedge^p(\underline{E}') \otimes_A \wedge^q(\underline{E}'')$.

When $\underline{E}', \underline{E}, \underline{E}''$ are locally direct summands of free Modules of finite type, the $\mathrm{Gr}^p \wedge^n(\underline{E})$ are locally direct summands of free Modules of finite type, and therefore so are the $F^p \wedge^n (\underline{E})$; we may thus write in K: $\lambda^n(E) = \sum_{p=0}^n \mathrm{cl}(\mathrm{Gr}^p \wedge^n(\underline{E})) = \sum_{p=0}^n \lambda^{n-p}(E') \,.\, \lambda^p(E'')$, giving additivity.

Let $(T, \underline{A})$, $(T', \underline{A}')$ be two commutatively ringed topoi, and $f : (T, \underline{A}) \longrightarrow (T', \underline{A}')$[213] a morphism of ringed topoi. We know (Exp. IV, 2.7) that f defines a ring morphism on the corresponding K, namely $f^* : K^\bullet_{\text{naïf}}(T', \underline{A}') \longrightarrow K^\bullet_{\text{naïf}}(T, \underline{A})$. Since inverse image commutes with exterior powers, the corresponding morphism f^* on the K is a λ-homomorphism for the structure just defined.

Let us mention two special cases:

- Let X be a scheme; $K^\bullet_{\text{naïf}}(X)$ is thus endowed with a pre-λ-ring structure; this is the case that will be used in the rest of the seminar, and it may be regarded as a "model" for the general theory of [print 317]λ-rings.
- Let A be a ring, $X = \operatorname{Spec} A$; also let G be a group, acting trivially on A; taking for (T, A) the topos of sheaves of sets on X on which G acts, the ring K thus obtained is the representation ring of G in the projective A-modules of finite type.

2.3. Let K be a commutative ring with unit; we propose to endow the abelian group $\widehat{G}(K) = 1 + K[[t]]^+$ with a pre-λ-ring structure, functorial in K. For this, we shall apply the results of (1.10) to the functor $\widehat{G}$.

We first define multiplication by specifying a $\mathbb{Z}$-bilinear morphism of functors: $\widehat{G} \times \widehat{G} \longrightarrow \widehat{G}$, by means of the element $1 + XYt$ of $\widehat{G}(\mathbb{Z}[X, Y])$; this element indeed satisfies 1.7.i), and to see that it satisfies 1.7.ii), we must show that in $\prod_{i=1}^n (1 + T_i Y t)$, expressed in terms of the elementary symmetric functions X_j of the T_i, X_n does not occur in the coefficients of t^k for $k < n$, which is evident for reasons of degree.

The law thus defined will be denoted by the sign $\circ$; we therefore have, by definition:

$$(1 + Xt) \circ (1 + Yt) = 1 + XYt. \tag{2.3.1}$$

It is distributive by construction, and associative and commutative by the diagrams: [214]

$$\begin{array}{ccc} \widehat{G} \times \widehat{G} \times \widehat{G} & \to & \widehat{G} \times \widehat{G} \\ \downarrow & & \downarrow \\ \widehat{G} \times \widehat{G} & \longrightarrow & \widehat{G} \end{array} \qquad \begin{array}{ccc} \widehat{G} \times \widehat{G} & \to & \widehat{G} \times \widehat{G} \\ \downarrow & & \downarrow \\ \widehat{G} & \xrightarrow{\mathrm{Id}} & \widehat{G}, \end{array}$$

[print 318]where the meaning of the arrows is evident, and whose commutativity, by 1.10, need only be verified on the elements $1 + Xt, 1 + Yt, 1 + Zt$ of $\widehat{G}(\mathbb{Z}[X, Y, Z])$ (resp. $1 + Xt, 1 + Yt$[215] of $\widehat{G}(\mathbb{Z}[X, Y])$), a verification that is immediate.

By 1.8, the product of two series under this law is calculated by substituting their coefficients in the universal polynomials of the law. We denote the n-th universal polynomial by P_n; it is a polynomial in $\mathbb{Z}[X_i, Y_j]_{1 \le i,j \le n}$, isobaric of weight n in each of the two groups of variables (see 1.15), and symmetric with respect to these two groups. As an example, let us give the value of P_2[216]:

$$P_2(X_1, X_2; Y_1, Y_2) = X_2 Y_1^2 + Y_2 X_1^2 - 2X_2 Y_2.$$

We also have the following relation: let $K \in \mathrm{Ob}(\mathcal{C}_0)$, $\psi \in \widehat{G}(K)$, $a \in K$, then

$$\psi(t) \circ (1 + at) = \psi(at). \tag{2.3.2}$$

As usual, the standard functorial argument reduces it to the case $K = \mathbb{Z}[X, Y]$, $\psi = 1 + Xt$, $a = Y$, where it is precisely 2.3.1.

Applying 2.3.2 to $\widehat{G}(K) = 1 + K[[t]]^+$, and to $a = 1$, we see that the element $1 + t$ is the unit for this multiplication. To avoid confusion, we sometimes denote it by $\underline{1}$, or by $\underline{1}_\circ$.

The functor $\widehat{G}$ can now be regarded as a functor from $\mathcal{C}_0$ to the category of commutative rings with unit, and can therefore be composed with itself; to avoid confusion, we [print 319]shall choose different indeterminates for each factor in the product of $\widehat{G}$ with itself, and indicate the indeterminate as a superscript when necessary: $\widehat{G}^t$, $\widehat{G}^u$, etc. For $A \in \mathrm{Ob}(\mathcal{C}_0)$, we therefore have $\widehat{G}^u \circ \widehat{G}^t(A) = 1 + \{1 + A[[t]]^+\}[[u]]^+$. To define the operations λ^i on $1 + K[[t]]^+$, we shall therefore specify a morphism of functors: $\widehat{G} \longrightarrow \widehat{G}^u \circ \widehat{G}^t$. For this, we must first define a filtration of the functor $\widehat{G}^u \circ \widehat{G}^t$ that permits the gluing of 1.6.

We first observe that $F^k\widehat{G}$ is an ideal for the law $\circ$; for this, functoriality reduces us to the case $A = \mathbb{Z}[X_i, Y]_{i \geq 1}$ and the product $\left(1 + \sum_{i \geq k} X_i t^i\right) \circ (1 + Yt)$, and the assertion follows from 2.3.2. We then take as $F^n(\widehat{G}^u \circ \widehat{G}^t)$ the set of series of the form $1 + \psi_1 u + \ldots + \psi_i u^i + \ldots$, $\psi_i \in F^{[n/i]}\widehat{G}$, where $[n/i]$ denotes the integer part of n/i.

The element defining the morphism λ_u is $\underline{1} + \{1 + Xt\}u$; the reader will verify as an exercise that the conditions of 1.6 are indeed satisfied (in fact, the filtration is chosen ad hoc).

Here again, the operations λ^i are calculated by substitution from $\lambda_u(1 + \ldots + X_n t^n + \ldots)$, and are therefore determined by universal polynomials. We denote by $Q_{i,j}(X_1, \ldots, X_{ij})$ the coefficient of t^j in $\lambda^i(1 + \ldots + X_n t^n + \ldots)$; it is an isobaric polynomial of weight ij in the variables $X_1, \ldots, X_{ij}$ (a verification analogous to that of 1.15). It is calculated by symmetric functions from the defining element:

$$\lambda_u(1 + Xt) = \underline{1} + \{1 + Xt\}u. \tag{2.3.3}$$

As an example, let us give the value of $Q_{2,2}$: [print 320]

$$Q_{2,2}(X_1, X_2, X_3, X_4) = X_1X_3 - X_4.$$

Finally, let us indicate the origin of formulas (2.3.1) and (2.3.3). We consider Example 2.2.b), where the operations λ^i are defined by the exterior powers of A-Modules. We want $\lambda_t : K \longrightarrow \widehat{G}(K)$ to be a λ-homomorphism; we must therefore define laws on $\widehat{G}$ such that for all A-Modules $\underline{E}, \underline{F}$, we have:

$$\lambda_t(EF) = \lambda_t(E) \circ \lambda_t(F)\ ; \qquad \lambda_u(\lambda_t(E)) = \widehat{G}^u(\lambda_t)(\lambda_u(E))\ ;$$

now relations (2.3.1) and (2.3.3) are precisely the preceding relations applied to the case where $\underline{E}$ and $\underline{F}$ are invertible modules.

We shall see in Exp. VI to what extent the desired property is satisfied.

Definition 2.4. A pre-λ-ring K will be called a λ-ring if the homomorphism of abelian groups $\lambda_t : K \longrightarrow 1 + K[[t]]^+$ is a λ-homomorphism for the pre-λ-ring structure on the second member defined in 2.3.

This therefore means that we have the relations: $\forall x, y \in K$,

$$\lambda_t(1) = 1 + t\ ; \quad \lambda_t(xy) = \lambda_t(x) \circ \lambda_t(y)\ ; \quad \lambda_u(\lambda_t(x)) = \widehat{G}(\lambda_t)(\lambda_u(x)). \tag{2.4.1}$$

In terms of the maps λ^i, the last two relations are equivalent to: $x, y \in K$, $i, j \in \mathbb{N}$[217],

$$\begin{aligned} \lambda^i(xy) &= P_i(\lambda^1(x), \ldots, \lambda^i(x); \lambda^1(y), \ldots, \lambda^i(y)) \\ \lambda^j(\lambda^i(x)) &= Q_{i,j}(\lambda^1(x), \ldots, \lambda^{ij}(x)) \end{aligned} \tag{2.4.2}$$

[print 321]We note that here we depart from the usual terminology (cf. [1]), where the name λ-ring was used for what we call a pre-λ-ring, and "special" λ-rings for our λ-rings; the reason is that only "special" λ-rings are usable in practice.

Examples 2.5.

a) We have determined all the pre-λ-ring structures on $\mathbb{Z}$; it follows from 2.4.1 that only the one defined by $\lambda_t(1) = 1 + t$ can be a λ-ring structure. We then have $\lambda_t(n) = (1+t)^n$, hence:

$$\forall n, n' \in \mathbb{Z}, \quad \lambda_t(n) \circ \lambda_t(n') = (1+t)^n \circ (1+t)^{n'} = (1+t)^{nn'} = \lambda_t(nn') \ ;$$

$$\lambda_u((1+t)^n) = (\lambda_u(1+t))^n = (\underline{1} + \{1+t\}u)^n = \widehat{G}(\lambda_t)((1+u)^n) = \widehat{G}(\lambda_t)(\lambda_u(n)).$$

Consequently, this is indeed a λ-ring structure. Denoting by $\binom{X}{k}$ the polynomial with rational coefficients $\frac{1}{k!}X(X-1)\dots(X-k+1)$, we therefore have:

$$\lambda^i(n) = \binom{n}{i}. \tag{2.5.1}$$

b) For every ring K, the pre-λ-ring $1 + K[[t]]^+$ is a λ-ring. First, $\lambda_u(\underline{1}) = \lambda_u(1+t) = \underline{1} + \{1+t\}u = \underline{1} + \underline{1}.u$. We verify the other relations by the commutativity of the diagrams: [218]

$$\begin{array}{ccc} \widehat{G}^t \times \widehat{G}^t & \xrightarrow{\lambda_u \times \lambda_u} & (\widehat{G}^u \circ \widehat{G}^t) \times (\widehat{G}^u \circ \widehat{G}^t) \\ {\scriptstyle\circ}\downarrow & & \downarrow{\scriptstyle\circ} \\ \widehat{G}^t & \xrightarrow{\lambda_u} & \widehat{G}^u \circ \widehat{G}^t, \end{array} \qquad \begin{array}{ccc} \widehat{G}^t & \xrightarrow{\lambda_v} & \widehat{G}^v \circ \widehat{G}^t \\ {\scriptstyle\lambda_u}\downarrow & & \downarrow{\scriptstyle\widehat{G}^v(\lambda_u)} \\ \widehat{G}^u \circ \widehat{G}^t & \xrightarrow{\lambda_v} & \widehat{G}^v \circ \widehat{G}^u \circ \widehat{G}^t, \end{array}$$

[print 322]whose commutativity is verified, by 1.10, on the elements $1 + Xt, 1 + Yt$. For the first, [219]

$$\begin{aligned} \lambda_u(1+Xt) \circ \lambda_u(1+Yt) &= (\underline{1} + \{1+Xt\}u) \circ (\underline{1} + \{1+Yt\}u) \\ &= \underline{1} + \{(1+Xt) \circ (1+Yt)\}u \\ &= \underline{1} + \{1+XYt\}u \\ &= \lambda_u((1+Xt) \circ (1+Yt)). \end{aligned}$$

For the second:

$$\lambda_v(\lambda_u(1+Xt)) = \underline{1} + \{\underline{1} + \{1+Xt\}u\}v = \widehat{G}^v(\lambda_u)(\lambda_v(1+Xt)).$$

c) The geometric examples of 2.2 will be treated later.

2.6. We can generalize Example 2.5.a) as follows: let K be a commutative ring with unit, such that: [220]

i) K is torsion-free,
ii) $\forall x \in K, \ \forall n \in \mathbb{N}, \quad$ we have $\binom{x}{n} \in K$

We then put, for every $x \in K$, and every $n \in \mathbb{N}$, $\lambda^n(x) = \binom{x}{n}$, that is:

$$\lambda_t(x) = \sum_{n=0}^{\infty} \binom{x}{n} t^n = (1+t)^x = \exp(x \operatorname{Log}(1+t)), \tag{2.6.1}$$

where the last identity is considered in $(K \otimes_{\mathbb{Z}} \mathbb{Q})[[t]]$. By the multiplicativity of the exponential, this defines a pre-λ-ring structure on K.

To see that it is a λ-ring structure, we must prove relations 2.4.2, and it suffices to prove them in $K \otimes_{\mathbb{Z}} \mathbb{Q}$, by 2.6.i). Functoriality then reduces us to the case $K = \mathbb{Q}[X, Y]$ and the elements X, Y. In view of the definition of the λ^i, the two members of the [print 323]equalities 2.4.2 are polynomials in X, Y, with rational coefficients, and to prove their equality it therefore suffices to show that they take the same value at all integer values of X, Y. But the relations obtained by substituting integer values for X, Y are those expressing that the pre-λ-ring Z[221] (2.5.a)) is a λ-ring;[222] they are therefore satisfied.

Definition 2.7. By a binomial ring we mean a ring satisfying conditions 2.6.i) and ii), endowed with the λ-ring structure defined by 2.6.1.

Example 2.8. Let X be a site, K a binomial ring, and $\underline{K}$ the sheaf associated to the constant presheaf with value K on X; then $H^0(X, \underline{K})$ is a binomial ring.

Indeed, conditions 2.6 are preserved under passage to an inductive limit.

In particular, $H^0(X, \underline{\mathbb{Z}})$ is a binomial ring.

2.9. Let K be a binomial ring, and A a K-algebra. For every series $\varphi \in \widehat{G}(A)$, and every $a \in K$, we can define a series φ^a in the usual way: we write $\varphi = 1 + \psi$, and put:

$$\varphi^a = (1+\psi)^a = \sum_{n=0}^{\infty} \binom{a}{n} \psi^n \, (= \exp(a \operatorname{Log} \varphi)) \,, \tag{2.9.1}$$

the last equality being writable only modulo torsion. These formulas are obtained by substitution from the analogous formulas for $\varphi = 1 + t$; the following relations, the classical properties of the exponential, therefore remain valid for $a, b \in K$, $\varphi, \psi \in \widehat{G}(A)$:

$$\varphi^a \varphi^b = \varphi^{a+b}; \qquad \varphi^{ab} = (\varphi^a)^b; \qquad \varphi^a \psi^a = (\varphi\psi)^a. \tag{2.9.2}$$

[print 324]We still suppose that $a \in K$. Multiplication by $\lambda_t(a)$ under the law $\circ$ defines a morphism of functors: $\widehat{G}_K \to \widehat{G}_K$, characterized by its value on the element $1 + Xt$ of $\widehat{G}(K[X])$. We therefore have:

$$\lambda_t(a) \circ (1 + Xt) = \lambda_{Xt}(a) \qquad \text{by (2.3.2), hence:}$$

$$\lambda_t(a) \circ (1 + Xt) = (1 + Xt)^a = \exp(a \operatorname{Log}(1 + Xt)) = \sum_{n=0}^{\infty} \binom{a}{n} X^n t^n. \tag{2.9.3}$$

We deduce:

$$\lambda_t(a) \circ \varphi = \varphi^a \quad (= \exp(a \operatorname{Log} \varphi)); \tag{2.9.4}$$

Indeed, the two members are additive functors equal on $1 + Xt$. From (2.9.3) and 1.15, it follows that the n-th universal polynomial defining multiplication by $\lambda_t(a)$ is isobaric of weight n; it will be denoted by $R_{n,a}$.

Finally, note that if A is a K-λ-algebra, i.e. a λ-ring endowed with a λ-homomorphism: $K \to A$, and if $b \in A$, we have:

$$\lambda_t(ab) = (\lambda_t(b))^a. \tag{2.9.5}$$

This follows from $\lambda_t(ab) = \lambda_t(a) \circ \lambda_t(b)$, and from (2.9.4).

3. The operations γ

We still work in the category $\mathcal{C}_0$.

3.1. Let K be a λ-ring. From the operations λ^i of K, we shall define new operations, denoted by γ^i. They are intended to introduce a notion of Chern classes under conditions that will be specified later. For the moment, let us give a "justification" of the following definitions: if $\underline{E}$ is a vector bundle of rank d, the d-th Chern class of $\underline{E}$ will have to be (whatever theory of Chern classes is considered) the "alternating sum of the exterior powers [print 325]of $\underline{E}$"; to define the n-th Chern class, we shall proceed analogously, bringing $\underline{E}$ to rank n by considering $(\mathrm{cl}(\underline{E}) - d) + n$, and we shall take the alternating sum of the exterior powers of the element thus obtained; the following proposition is to be "interpreted" by setting $x = \mathrm{cl}(\underline{E}) - d$:

Proposition 3.2. Let K be a λ-ring.

i) $\forall x \in K,\ \forall n \in \mathbb{N}, \quad \lambda^n(x+n-1) = (-1)^n \sum_{i=0}^{n} (-1)^i \lambda^i(x+n)$.

ii) Putting $\gamma^n(x) = \lambda^n(x+n-1)$, and $\gamma_t(x) = \sum_{n\geq 0} \gamma^n(x) t^n$, we have: [223]

$$\gamma_t(x) = \lambda_{t/(1-t)}(x), \qquad \text{hence}$$

[224]

$$\lambda_t(x) = \gamma_{t/(1+t)}(x).$$

We therefore see that knowing the operations γ is equivalent to knowing the operations λ.

Proof.

i) We have

$$\lambda^n(x+n-1) = \sum_{i=0}^{n} \lambda^i(x+n)\lambda^{n-i}(-1).$$

From $\lambda_t(1) = 1+t$, we obtain: $\lambda_t(-1) = 1/(1+t)$[225], hence: $\lambda^k(-1) = (-1)^k$, which gives the required relation.

ii) $\lambda_{t/(1-t)}(x) = \sum_{n\geq 0} \lambda^n(x).t^n(1-t)^{-n}$.[226] We expand $t^n(1-t)^{-n} = t^n \sum_{k\geq 0} \binom{-n}{k}(-1)^k t^k = \sum_{p\geq n} \binom{p-1}{p-n} t^p$. $\lambda_{t/(1-t)}(x) = \sum_{n\geq 0} [\lambda^n(x). \sum_{p\geq n} \binom{p-1}{p-n} t^p]$[227]; the coefficient of t^p is therefore:

$$\sum_{n=0} \lambda^n(x). \binom{p-1}{p-n} = \sum_{n=0} \lambda^n(x)\lambda^{p-n}(p-1) = \lambda^p(x+p-1).$$

[print 326]3.3. Let K, K' be two λ-rings, and $f : K \to K'$ a ring homomorphism. By Definition 3.2.ii), f is a λ-homomorphism if and only if, for every $i \in \mathbb{N}$, and every $x \in K$, $f(\gamma^i(x)) = \gamma^i(f(x))$, or equivalently $\gamma_t(f(x)) = \widehat{G}_g(f) \circ \gamma_t(x)$.

3.4. From relation 3.2.ii), we deduce the relations:

$$\gamma^0(x) = 1, \qquad \gamma^1(x) = x, \qquad \gamma^n(x+y) = \sum_{p+q=n} \gamma^p(x)\gamma^q(y). \tag{3.4.1}$$

They show that the homomorphism γ_t defines on K a new pre-λ-ring structure; we shall see that in general it is not a λ-ring structure. Denoting by ρ_t the substitution in the ring of formal series in t with coefficients in K defined by $\rho_t(t) = t/(1-t)$[228], we have:

$$\gamma_t = \rho_t \circ \lambda_t.$$

The automorphism ρ_t is functorial in K, and defines an automorphism of the functor $\widehat{G}$ considered as a functor to abelian groups; with this automorphism, by transport of structure, we can define on $\widehat{G}$ a new structure as a functor to λ-rings, such that for every λ-ring K, the homomorphism

$\gamma_t : K \to \widehat{G}(K)$ is a λ-homomorphism; we shall make explicit the multiplication and the operations λ^i of this second structure.

3.5. The multiplication is defined by the homomorphism of functors: [229]

$$\widehat{G} \times \widehat{G} \xrightarrow{(\rho^{-1},\rho^{-1})} \widehat{G} \times \widehat{G} \xrightarrow{\circ} \widehat{G} \xrightarrow{\rho} \widehat{G},$$

where the sign $\circ$ denotes the multiplication defined in 2.3. This new multiplication will be denoted by v. As usual, it is defined by its value on $(1 + Xt, 1 + Yt)$; we therefore have: [print 327][230]

$$\begin{aligned}(1 + Xt)\nabla(1 + Yt) &= \rho\Big(\rho^{-1}(1 + Xt) \circ \rho^{-1}(1 + Yt)\Big) \\ &= \rho\Big(\frac{1 + (X + 1)t}{1 + t} \circ \frac{1 + (Y + 1)t}{1 + t}\Big) \\ &= \rho\Big(\frac{(1 + (X + 1)(Y + 1)t)(1 + t)}{(1 + (X + 1)t)(1 + (Y + 1)t)}\Big) \\ &= \frac{1 + (X + Y + XY)t}{(1 + Xt)(1 + Yt)}. \end{aligned} \tag{3.5.1}$$

If K is a λ-ring, then for all $x, y \in K$ we have:

$$\gamma_t(xy) = \gamma_t(x)\nabla\gamma_t(y). \tag{3.5.2}$$

We denote by P'_n the n-th universal polynomial of the law ∇; it is a polynomial in $\mathbb{Z}[X_i, Y_j]_{1\leq i,j\leq n}$ whose isobaric components have weights ranging from n to $2n$ with respect to all the variables X_i, Y_j, (1.15) and which is symmetric with respect to these two groups of variables. For $x, y \in K$, we therefore have:

$$n \in \mathbb{N}, \quad \gamma^n(xy) = P'_n(\gamma^1(x), \ldots, \gamma^n(x), \gamma^1(y), \ldots, \gamma^n(y)). \tag{3.5.3}$$

Finally, note that the law ∇ has a unit element, namely $\rho_t(1 + t)$, that is, $1/(1 - t)$[231]; it will sometimes be denoted by $\underline{1}_\nabla$.

3.6. The "operations λ" are defined by the commutative diagram:

$$\begin{array}{ccc} 1 + K[[t]]^+ & \xrightarrow{\lambda_u} & 1 + \{1 + K[[t]]^+\}[[u]]^+ \\ {\scriptstyle\rho_t}\downarrow & & \downarrow{\scriptstyle\widehat{G}(\rho_t)} \\ 1 + K[[t]]^+ & \xrightarrow{\lambda'_u} & 1 + \{1 + K[[t]]^+\}[[u]]^+, \end{array} \tag{3.6.1}$$

i.e. by the composite morphism of functors:

$$\lambda' : \widehat{G} \xrightarrow{\rho_t^{-1}} \widehat{G} \xrightarrow{\lambda_u} \widehat{G} \circ \widehat{G} \xrightarrow{\widehat{G}(\rho_t)} \widehat{G} \circ \widehat{G},$$

and will be denoted by λ'. They are defined by their value on $1 + Xt$; we have: [print 328][232]

$$\begin{aligned}\lambda'_u(1 + Xt) &= \widehat{G}(\rho_t)(\lambda_u(\rho_t^{-1}(1 + Xt))) \\ &= \widehat{G}(\rho_t)\left(\lambda_u\left(\frac{1 + (X + 1)t}{1 + t}\right)\right) \\ &= \widehat{G}(\rho_t)\left(\frac{\underline{1}_o + \{1 + (X + 1)t\}u}{\underline{1}_o + \{1 + t\}u}\right) \\ &= \underline{1}_\nabla + \frac{\{1 + Xt\}u}{\underline{1}_\nabla + \underline{1}_\nabla.u} \\ &= \rho_u^{-1}(\underline{1}_\nabla + \{1 + Xt\}u). \end{aligned}$$

Denoting by γ' the operations γ corresponding to the operations λ', we therefore have:

$$\gamma'_u(1+Xt) = \underline{1}_\nabla + \{1+Xt\}u. \tag{3.6.2}$$

We denote by $Q'_{i,j}$ the coefficient of t^j in $\gamma'^i(1+\ldots+X_nt^n+\ldots)$; the reader can verify that it is a polynomial in $Z[X_1,\ldots,X_{ij}]$ whose isobaric components have weights ranging between j and ij.

Let K be a λ-ring, $x \in K$. From diagram (3.6.1), we deduce the relations:

$$\forall i \in \mathbb{N}, \quad \gamma_t(\lambda^i(x)) = \lambda'^i(\gamma_t(x)) \ ; \qquad \gamma_t(\gamma^i(x)) = \gamma'^i(\gamma_t(x)). \tag{3.6.3}$$

$$\gamma^j(\gamma^i(x)) = Q'_{i,j}(\gamma^1(x),\ldots,\gamma^{ij}(x)). \tag{3.6.4}$$

Finally, we observe that the pre-λ-ring structure just defined on $\widehat{G}$ is a λ-ring structure, since it is obtained from the one defined in 2.3 by the isomorphism ρ.

3.7. To avoid confusion, the functor $\widehat{G}$, endowed with the structure of a functor to λ-rings defined in §2 (resp. in §3), will be denoted by $\widehat{G}_o$ (resp. $\widehat{G}_\nabla$). It follows from the preceding considerations that to specify a [print 329]λ-ring structure on a ring A, it is equivalent to specify a ring homomorphism $A \to \widehat{G}(A)_o$ such that diagram 3.7.1 is commutative, or a ring homomorphism $A \to \widehat{G}(A)_\nabla$ such that diagram 3.7.2 is commutative; the first defines the operations λ, the second the operations γ.

$$\begin{array}{ccc} A & \xrightarrow{\lambda_u} & \widehat{G}^u(A)_o \\ {\scriptstyle\lambda_t}\downarrow & & \downarrow{\scriptstyle\widehat{G}^u(\lambda_t)} \\ \widehat{G}^t(A)_o & \overset{\lambda_u}{\to} & \widehat{G}^u(\widehat{G}^t(A)_o)_o \end{array} \tag{3.7.1}$$

[233]

$$\begin{array}{ccc} A & \xrightarrow{\gamma_u} & \widehat{G}^u(A)_\nabla \\ {\scriptstyle\gamma_t}\downarrow & & \downarrow{\scriptstyle\widehat{G}^u(\gamma_t)} \\ \widehat{G}^t(A)_\nabla & \overset{\gamma'_u}{\to} & \widehat{G}^u(\widehat{G}^t(A)_\nabla)_\nabla. \end{array} \tag{3.7.2}$$

3.8. Let K be a binomial ring (2.7); from (2.6.1) we deduce, $\forall x \in K$, the relations: [234]

$$\gamma_t(x) = (1/(1-t))^x = \exp(-x\,\mathrm{Log}(1-t)) \tag{3.8.1}$$

Let a be an element of K; multiplication by $\gamma_t(a)$ under the law ∇ is characterized by its value on $1+Xt$, namely:

$$\gamma_t(a)\nabla(1+Xt) = \rho_t\left(\lambda_t(a) \circ \frac{1+(X+1)t}{1+t}\right) = \rho_t\left(\left(\frac{1+(X+1)t}{1+t}\right)^a\right)$$

$$\gamma_t(a)\nabla(1+Xt) = (1+Xt)^a = \exp(a\,\mathrm{Log}(1+Xt)) = \sum_{n=0}^{\infty}\binom{a}{n}X^nt^n. \tag{3.8.2}$$

In particular, we see that the n-th universal polynomial defining multiplication by $\gamma_t(a)$ under the law ∇ is the polynomial $R_{n,a}$ (2.9).

[print 330]**Definition 3.9.** Let K be a λ-ring; by an augmented K-λ-algebra we mean the data of a K-λ-algebra A and a λ-homomorphism of K-algebras $A \to K$; a homomorphism of augmented K-λ-algebras is a λ-homomorphism compatible with the augmentations.

Example 3.9.1.: Let (X, A) be a commutatively locally ringed topos, and $K^\bullet(X)$ the Grothendieck group of the category of sheaves of locally free A-modules of finite type on X. We have seen that $K^\bullet(X)$ can be endowed with a pre-λ-ring structure (2.2.b, since every direct summand of a free module of finite type is then locally free of finite type) and that $H^0(X, \underline{\mathbb{Z}})$ is canonically endowed with a binomial ring structure. We define a λ-homomorphism from $H^0(X, \underline{\mathbb{Z}})$ to $K^\bullet(X)$ as follows: let a be a section of the constant sheaf $\underline{\mathbb{Z}}$ on X; it defines a partition of the final object of X into a sum of disjoint opens U_n, the section a taking the value n on the open U_n; let $\underline{F}$ be the sheaf equal to A^n on U_n for $n \geq 0$, and to 0 on U_n for $n \leq 0$, and let $\underline{F}'$ be the sheaf equal to A^{-n} on U_n for $n \leq 0$, and to 0 on U_n for $n \geq 0$; we associate to a the element $F - F'$ of $K^\bullet(X)$.

It is immediate to verify that this does give a λ-homomorphism from $H^0(X, \underline{\mathbb{Z}})$ to $K^\bullet(X)$; we can then define the augmentation from $K^\bullet(X)$ to $H^0(X, \underline{\mathbb{Z}})$ by assigning to every locally free sheaf of finite type its rank, which is a section of $\underline{\mathbb{Z}}$; additivity of rank implies that this defines a function on $K^\bullet(X)$ with values in $H^0(X, \underline{\mathbb{Z}})$. We have thus endowed $K^\bullet(X)$ with an augmented pre-λ-algebra structure. This is always the structure referred to below.

[print 331]3.10 Let A be an augmented K-λ-algebra. We define a ring filtration on A as follows: for every $n \geq 0$, we denote by $\mathrm{Fil}^n A$ the ideal generated by the products:

$$\gamma^{i_1}(x_1) \ldots \gamma^{i_k}(x_k), \qquad \text{with } i_1 + \ldots + i_k \geq n, \text{ and } \forall i,\ \epsilon(x_i) = 0 \tag{3.10.1}$$

($\epsilon : A \to K$ being the augmentation).

Suppose that A is generated as an abelian group by a family of elements $(x_\alpha)_{\alpha \in I}$; then $\mathrm{Fil}^n A$ is generated as a K-module by the products:

$$\gamma^{i_1}(x_{\alpha_1} - \epsilon(x_{\alpha_1})) \ldots \gamma^{i_k}(x_{\alpha_k} - \epsilon(x_{\alpha_k})), \qquad \text{with } i_1 + \ldots + i_k \geq n, \text{ and } \alpha_i \in I \tag{3.10.2}$$

Indeed, if x is an element of A with zero augmentation and we write $x = \sum_i x_{\alpha_i}$, we have $\sum_i \epsilon(x_{\alpha_i}) = 0$, hence $x = \sum_i x_{\alpha_i} - \epsilon(x_{\alpha_i})$; furthermore, if $a \in A$, we can write $a = (a - \epsilon(a)) + \epsilon(a) = \gamma^1(a - \epsilon(a)) + \epsilon(a)$; using relations (3.4.1), we therefore see that the elements of $\mathrm{Fil}^n A$ are linear combinations with coefficients in K of the elements (3.10.2); since these belong to $\mathrm{Fil}^n A$, we obtain the required result.

The filtration thus defined is a ring filtration; it is decreasing, and we have:

$$\mathrm{Fil}^0 A = A\ ; \qquad \mathrm{Fil}^1 A = \mathrm{Ker}\, \epsilon.$$

Moreover, it is functorial in A: every λ-homomorphism of augmented K-λ-algebras is compatible with the filtrations. When we regard an augmented λ-ring as endowed with a filtration, this will mean the filtration above, unless expressly stated otherwise.

[print 332]**Proposition 3.11.** Let A be a λ-ring, and y_i, $i = 1, \ldots, n$[235] elements of A such that $\lambda^j(y_i) = 0$ for $j \geq 2$. Put $x = \sum_i y_i$; then:

i) for $k > n$, we have $\gamma^k(x - n) = 0$

ii) for $1 \leq k \leq n$, $\gamma^k(x - n)$ is the k-th symmetric function of the $y_i - 1$.

Let $j \geq 2$. We then have $\gamma^j(y_i - 1) = \lambda^j(y_i + j - 2) = \sum_{p+q=j} \lambda^p(y_i)\lambda^q(j-2)$. Since $\lambda^p(y_i) = 0$ for $p \geq 2$, and $\lambda^q(j-2) = 0$ for $q > j - 2$, $\gamma^j(y_i - 1) = 0$. The proposition then follows from $\gamma_t(x - n) = \prod_{i=1}^n \gamma_t(y_i - 1)$.

4. λ-rings generated by generators and relations

We still work over $\mathcal{C}_0$.

4.1. Let K be a λ-ring. Consider a family of indeterminates $(X_i)_{i\in I}$, and let $A = K[X_i, \lambda^n X_i]_{i\in I,\ n\geq 2}$; we wish to define on A a λ-ring structure extending that of K.

To specify a homomorphism $\lambda_t : A \to \widehat{G}(A)$, it suffices to specify the images of the X_i, $\lambda^n X_i$, which defines a ring homomorphism: $A \to \widehat{G}(A)_o$; for it to define a λ-ring structure on A, we must verify the commutativity of diagram (3.7.1). We therefore put:

$$\lambda_t(X_i) = 1 + X_i t + \sum_{n\geq 2} \lambda^n X_i\, t^n\ ; \qquad \lambda_t(\lambda^n X_i) = \lambda^n(\lambda_t(X_i)). \tag{4.1.1}$$

Since diagram (3.7.1) is a diagram of ring homomorphisms, it suffices to verify its commutativity on the X_i, $\lambda^n X_i$ (it is known by hypothesis on the elements of K); this follows readily from definitions (4.1.1) and the fact that λ_u and $\widehat{G}^u(\lambda_t)$ are λ-homomorphisms.

[print 333]The λ-ring A is characterized by the following universal property:

Proposition 4.2. For every K-λ-algebra B, and every family $(b_i)_{i\in I}$ of elements of B, there exists exactly one K-λ-homomorphism f from A to B such that for every $i \in I$, $f(X_i) = b_i$.

For f to be a λ-homomorphism, we must have $f(\lambda^j X_i) = \lambda^j(b_i)$ for all i, j; this defines f uniquely. The homomorphism thus obtained is indeed a λ-homomorphism, since to verify that $\lambda_t \circ f = \widehat{G}(f) \circ \lambda_t$ it suffices to verify it on a system of generators of A over K, which follows from the definition of f and relations (2.4.2).

4.3. We can make the same construction using the γ in place of the λ. Let $A' = K[X_i, \gamma^n X_i]_{i\in I,\ n\geq 2}$; to define the λ-ring structure on A', we specify a ring homomorphism: $A' \to \widehat{G}(A')_\nabla$, by:

$$\gamma_t(X_i) = 1 + X_i t + \sum_{n\geq 2} \gamma^n X_i\,.\,t^n\ ; \qquad \gamma_t(\gamma^n X_i) = \gamma'^n(\gamma_t(X_i)). \tag{4.3.1}$$

We then verify the commutativity of (3.7.2), which shows that we do obtain a λ-ring structure.

The λ-ring A' also satisfies the universal property 4.2; it is therefore canonically isomorphic (as a λ-ring) to the λ-ring A.

Definition 4.4. The λ-ring A will be called the free λ-ring generated over K by the family of generators $(X_i)_{i\in I}$.

Definition 4.5. Let A be a λ-ring, J an ideal of A. We say that J is a λ-ideal of A if for every $x \in J$, and every $n \geq 1$, $\lambda^n(x) \in J$.

If $f : A \to B$ is a λ-homomorphism, the kernel of f is a λ-ideal.

Let J be a λ-ideal; the composite homomorphism: $A \xrightarrow{\lambda_t} \widehat{G}(A) \longrightarrow \widehat{G}(A/J)$ vanishes on J and therefore factors through A/J, which is thus endowed [print 334]with a λ-ring structure, relations (2.4.2) passing to the quotient.

Every intersection of λ-ideals is a λ-ideal. In particular, there exists a smallest λ-ideal containing a given family of elements $(x_i)_{i\in I}$; by relations (2.4.2), it is the (ordinary) ideal generated by the $\lambda^n(x_i)$, $(n \geq 1)$.

Again let K be a λ-ring, $(X_i)_{i\in I}$ a family of indeterminates, $A = K[(X_i), (\lambda^n X_i)]$, and let $(P_{i'})_{i'\in I'}$ be a family[236] of elements of A: let J be the λ-ideal generated by the $P_{i'}$.

Definition 4.6. The λ-ring A/J will be called the λ-ring generated over K by the family of generators $(X_i)_{i\in I}$ subject to the relations $(P_{i'})_{i'\in I'}$.

We can also specify relations between the X_i, $\gamma^n X_i$ and make the same construction, observing that an ideal is a λ-ideal if and only if it is stable under the operations γ^n, $n \geq 1$, as follows from relations 3.2.[237]

Let B be a K-λ-algebra, $(b_i)_{i\in I}$ a family of elements of B. Since the $P_{i'}$ are polynomials in the indeterminates X_i, $\lambda^n X_i$, we can substitute the b_i, $\lambda^n(b_i)$ for these indeterminates; if the resulting element is zero, we say that the b_i satisfy the relations $P_{i'}$. The K-λ-algebra A/J is then characterized by the following universal property:

Proposition 4.7. For every K-λ-algebra B and every family $(b_i)_{i\in I}$ of elements of B, satisfying the relations $(P_{i'})_{i'\in I'}$, there exists exactly one K-λ-homomorphism $f' : A/J \to B$[238] such that for every i, we have $f'(X_i) = b_i$.[239]

Indeed, the kernel of the homomorphism f defined in 4.2 contains the $P_{i'}$, hence J since it is a λ-ideal; f therefore passes to the quotient and defines [print 335] f', which is plainly unique.

Proposition 4.8. Let n be an integer ≥ 0; in the free λ-ring generated over K by an indeterminate X, the ideal generated by the $\lambda^k X$, $k > n$, is a λ-ideal.

Put $A = K[X, \lambda^k X]_{k\geq 2}$, and let J be the ideal generated by the $\lambda^k X$ for $k > n$; put $B = A/J$. We must show that the composite homomorphism: $A \xrightarrow{\lambda_t} \widehat{G}(A) \longrightarrow \widehat{G}(B)$ vanishes on J, i.e. that $\lambda_t(\lambda^k X)$, $k > n$, has image 1 in $\widehat{G}(B)$. Now $\lambda_t(\lambda^k X) = \lambda^k(\lambda_t(X))$, and $\widehat{G}(A) \to \widehat{G}(B)$ is a λ-homomorphism; it therefore suffices to show that for every ring C and every series $\varphi \in \widehat{G}(C)$ whose coefficients vanish in degrees $> n$, $\lambda^k(\varphi) = 1$ when $k > n$. But to prove this last property it suffices to prove it for $C = K[X_1, \ldots, X_n]$, and $\varphi = 1 + X_1 t + \ldots + X_n t^n$. Embedding $K[X_1, \ldots, X_n]$ in $K[T_1, \ldots, T_n]$ by means of symmetric functions, we can write: $1 + X_1 t + \ldots + X_n t^n = \prod_{i=1}^n (1 + T_i t)$[240], hence

$$\lambda_u(1 + X_1 t + \ldots + X_n t^n) = \prod_{i=1}^n \lambda_u(1 + T_i t) = \prod_{i=1}^n (\underline{1} + \{1 + T_i t\}u),$$

which has degree n in u.

Corollary 4.9. The λ-ring generated over K by the generator X subject to the relations $\lambda^k(X) = 0$ for $k > n$, is isomorphic to $K[X, \ldots, \lambda^n X]$.

4.9.1. We would have the same statements on replacing the λ by the γ. We can also replace X by a family of indeterminates $(X_i)_{i\in I}$, subject to the relations $\lambda^k(X_i) = 0$ for $k > n_i$, where the n_i are integers ≥ 2; the resulting λ-ring is then $K[X_i, \lambda^k X_i]_{i\in I,\ 2\leq k\leq n_i}$.

4.10. Let K be a binomial ring, and A an augmented K-λ-algebra. We denote by $B = A[X, \gamma^k X]_{k\geq 2}$ the free λ-ring generated by X over A; [print 336] we define a λ-homomorphism: $B \to K$ extending the augmentation of A by sending X to 0, which endows B with an augmented K-λ-algebra structure. We propose to determine its filtration (3.10) in terms of that of A.

We grade B by weight with respect to the $\gamma^k X$, assigning $\gamma^k X$ the weight k; we denote by M_k the set of monomials of weight k in the indeterminates $\gamma^i X$.

Lemma 4.11. Let $a \in A$; the coefficient of a monomial of weight n with respect to the $\gamma^j X$ in $\gamma^k(aX) = P'_k(\gamma^i(a), \gamma^j X)$ has filtration at least $k - n$ (in A)

The polynomial P'_k has no isobaric components of weight $< k$ in the $\gamma^i(a)$, $\gamma^j X$ (3.5), so the lemma is true when a has zero augmentation.

If ϵ denotes the augmentation homomorphism: $A \to K$, we can write in the general case:

$$\gamma_t(aX) = \gamma_t((a - \epsilon(a))X) \,.\, \gamma_t(\epsilon(a)X).$$

Now $\gamma_t(\epsilon(a)X) = (\gamma_t(X))^{\epsilon(a)} = \sum_{n=0}^\infty R_{n,\epsilon(a)}((\gamma^j X)) \,.\, t^n$[241], the $R_{n,\epsilon(a)}$[242] being isobaric of weight n in the $\gamma^j X$ (3.8); we deduce the lemma by taking the product.

Proposition 4.12. $\mathrm{Fil}^k B = \mathrm{Fil}^k A + \mathrm{Fil}^{k-1} A \,.\, M_1 + \ldots + \mathrm{Fil}^0 A \,.\, M_k + B \,.\, M_{k+1}$.

It is clear that the second ideal is contained in the first; it therefore remains only to prove the reverse inclusion.

Since the filtration defined by the family of ideals on the right-hand side for varying k is a ring filtration, it suffices to prove the reverse inclusion for an element of the form $\gamma^k(b)$, with $\epsilon(b) = 0$, these elements [print 337]generating the canonical filtration.

Writing $b = \sum a_i m_i$, with $a_i \in A$ and the m_i monomials in the $\gamma^j X$, and using the additivity of γ_t, we see that it suffices to give the proof for $\gamma^k(am_i)$, with $a \in A$ and $m_i \in M_i$.

Using Lemma 4.11. and the A-λ-homomorphism from B to B which sends X to m_i, we see that it suffices to give the proof for an element of the form $\gamma^k(m_i)$.

Finally, since the polynomial P'_k has no isobaric components of weight $< k$, we are reduced to the case $\gamma^k(\gamma^j X)$, which itself follows from the fact that the polynomials $Q'_{j,k}$ have no isobaric components of weight $< k$ (3.6).

Remarks 4.13. i) The decomposition of $\mathrm{Fil}^k B$ given in 4.12 is a direct-sum decomposition of A-modules. In other words, if $b \in \mathrm{Fil}^k B$, and if we write $b = \sum_{j \leq k} a_{i,j} m_{i,j} + b'$, with $a_{i,j} \in A$, $m_{i,j} \in M_j$, and $b' \in B \,.\, M_{k+1}$, then $a_{i,j} \in \mathrm{Fil}^{k-j} A$.

ii) Statement 4.12. remains true for every quotient of B by a λ-ideal contained in the augmentation ideal.

iii) The statement analogous to 4.12., obtained by replacing the indeterminate X by a family of indeterminates (X_i), is also true.

Corollary 4.14. $\mathrm{Gr}^k B \approx \mathrm{Gr}^k A \oplus \mathrm{Gr}^{k-1} A \,.\, M_1 \oplus \ldots \oplus \mathrm{Gr}^0 A \,.\, M_k$.

We define an arrow from right to left by associating to each element $\bar{a} \,.\, m$, where $\bar{a}$ is the class in $\mathrm{Gr}^{k-i} A$ of an element a of $\mathrm{Fil}^{k-i} A$ and $m \in M_i$, the class in $\mathrm{Gr}^k B$ of am. We define an arrow in the opposite direction by decomposition according to Proposition 4.12 and Remark 4.13.i). It is clear that these two arrows are inverse to one another.

[print 338]**Corollary 4.15.** Let B' be the λ-ring generated over K by the indeterminate X subject to the relations $\gamma^k(X) = 0$ for $k > n$, i.e. $B' = K[X, \ldots, \gamma^n X]$ (4.9). Then the λ-ring filtration of B' is identical to the weight filtration with respect to the $\gamma^j X$.

Taking account of Proposition 4.12, Remark 4.13.ii), and the fact that $A = K$, we have $\mathrm{Fil}^k B' = K \,.\, M_k + B' \,.\, M_{k+1}$, so $\mathrm{Fil}^k B'$ is the ideal generated by the monomials of weight $\geq k$ with respect to the $\gamma^j X$.

5. The $\lambda^p(N, x)$.

This section will be used only in Exp.VII 4, for the study of regular immersions; it may therefore be omitted on first reading.

5.1. Let d be an integer ≥ 1. We denote by A the $\mathbb{Z}$-λ-algebra generated by the generators N, x subject to the relations $\lambda^i(N) = 0$ for $i > d$; we therefore have:

$$A = \mathbb{Z}[N, \ldots, \lambda^d N; x, \ldots, \lambda^j x, \ldots].$$

We put $\lambda_{-1}(N) = \sum_{i=0}^{d} (-1)^i \lambda^i(N)$.

Lemma 5.2. For every $p \geq 1$, $\lambda^p(x \,.\, \lambda_{-1}(N))$ is divisible by $\lambda_{-1}(N)$.

We know (2.3) that $\lambda^p(x \,.\, \lambda_{-1}(N)) = P_p(\lambda^i x, \lambda^j(\lambda_{-1}(N)))$ is an isobaric polynomial of weight p with respect to the $\lambda^j(\lambda_{-1}(N))$; it therefore suffices to prove the lemma for the $\lambda^p(\lambda_{-1}(N))$.

Let B be the λ-ring generated over $\mathbb{Z}$ by the indeterminates $N_1, \ldots, N_d$, subject to the relations $\lambda^i(N_j) = 0$ if $i \geq 2$. It is clear that $\lambda^k(\sum_{i=1}^d N_i) = 0$ for $k > d$. We can therefore define a λ-homomorphism f from A to B by sending N to $\sum_{i=1}^d N_i$; we then have $\widehat{G}(f)(\lambda_t(N)) = \prod_{i=1}^d \lambda_t(N_i) = \prod_{i=1}^d (1 + N_i t)$, [print 339]which shows that, for $1 \leq k \leq d$, $\lambda^k N$ maps to the k-th symmetric function of the N_i, and consequently that f is injective and identifies A with the ring of symmetric polynomials in the N_i.

We then have $\lambda_{-1}(N) = \prod_{i=1}^d (1 - N_i)$, and $\lambda_t(\lambda_{-1}(N)) = \lambda_t(\prod_{i=1}^d (1 - N_i))$. The $\lambda^p(\lambda_{-1}(N))$ are polynomials in the N_i; if, for some index i, we set $N_i = 1$, we see that $\lambda_t(\lambda_{-1}(N))$ becomes $\lambda_t(0) = 1$, hence that the $\lambda^p(\lambda_{-1}(N))$ vanish for $N_i = 1$, and therefore that they contain $1 - N_i$ as a factor. Thus they contain the product of the $1 - N_i$ as a factor, i.e. they are divisible by $\lambda_{-1}(N)$ in B, and hence also in A, since the quotient of two symmetric polynomials is evidently symmetric.

5.3. Since A is an integral domain, we define an element $\lambda^p(N, x)$ unambiguously by putting ($p \geq 1$):

$$\lambda^p(N, x) \,.\, \lambda_{-1}(N) = \lambda^p(x \,.\, \lambda_{-1}(N)). \tag{5.3.1}$$

The element $\lambda^p(N, x)$ is therefore a polynomial in the $\lambda^i(x)$, $\lambda^j(N)$, isobaric of weight p with respect to the $\lambda^i(x)$, since the same is true of $\lambda^p(x \,.\, \lambda_{-1}(N))$ by 2.3.

Let d' be an integer $\geq d$, and let A' be the $\mathbb{Z}$-λ-algebra generated by the generators N' and x' subject to the relations $\lambda^i(N') = 0$ for $i > d'$; by Proposition 4.7, there exists exactly one λ-homomorphism $f : A' \longrightarrow A$, sending N' to N and x' to x. Definition (5.3.1) implies that $f(\lambda^p(N', x')) = \lambda^p(N, x)$ for every $p \geq 1$.

If B is a λ-ring and N'' is an element of B such that $\lambda^k(N'') = 0$ for $k > d$, we can, for every $x'' \in B$, define an element $\lambda^p(N'', x'')$ by substitution, or by using the unique λ-homomorphism from A to [print 340]B which sends N to N'', and x to x''; by the preceding remark, the element obtained does not depend on the integer d such that $\lambda^k(N'') = 0$ for $k > d$. Using the same homomorphism, we see that relation (5.3.1) is satisfied by N'' and x''.

5.4. Consider the λ-ring generated over $\mathbb{Z}$ by the indeterminates N, x, y, subject to the relations $\lambda^k(N) = 0$ for $k > d$. From (5.3.1) we obtain:

$$\begin{aligned}
\lambda^p(N, x + y) \,.\, \lambda_{-1}(N) &= \lambda^p\big((x + y) \,.\, \lambda_{-1}(N)\big) \\
&= \sum_{i+j=p} \lambda^i(x \,.\, \lambda_{-1}(N)) \,.\, \lambda^j(y \,.\, \lambda_{-1}(N)) \\
&= \sum_{\substack{i+j=p \\ i,j \neq 0}} \lambda^i(N, x) \,.\, \lambda^j(N, y) \,.\, (\lambda_{-1}(N))^2 + (\lambda^p(N, x) + \lambda^p(N, y))\lambda_{-1} N
\end{aligned}$$

Since the ring under consideration is an integral domain, we deduce the following relation, valid by substitution in every λ-ring, and for every element N such that $\lambda^k(N) = 0$ for $k > d$:

$$\lambda^p(N, x + y) = \sum_{\substack{i+j=p \\ i,j \neq 0}} \lambda^i(N, x) \,.\, \lambda^j(N, y) \,.\, \lambda_{-1}(N) + \lambda^p(N, x) + \lambda^p(N, y). \tag{5.4.1}$$

Now consider the λ-ring generated over $\mathbb{Z}$ by indeterminates N, N', x subject to the relations $\lambda^k(N) = 0$ for $k > d$, and $\lambda^k(N') = 0$ for $k > d'$. We then have $\lambda^k(N + N') = 0$ for $k > d + d'$; we can therefore define the $\lambda^p(N + N', x)$. We obtain:

$$\lambda^p(N + N', x) \,.\, \lambda_{-1}(N + N') = \lambda^p(x \,.\, \lambda_{-1}(N + N')).$$

By the additivity of λ_t, we have: $\lambda_{-1}(N+N') = \lambda_{-1}(N)\,.\,\lambda_{-1}(N')$. We deduce:

$$\begin{aligned}\lambda^p(N+N',x)\,.\,\lambda_{-1}(N)\,.\,\lambda_{-1}(N') &= \lambda^p(x\,.\,\lambda_{-1}(N)\,.\,\lambda_{-1}(N')) \\ &= \lambda^p(N', x\,.\,\lambda_{-1}(N))\,.\,\lambda_{-1}(N').\end{aligned}$$

[print 341]Since the ring under consideration is an integral domain, we deduce the following relation, valid by substitution in every λ-ring, and for all elements N, N' such that $\lambda^k(N) = 0$ for $k > d$, $\lambda^k(N') = 0$ for $k > d'$;

$$\lambda^p(N+N',x)\,.\,\lambda_{-1}(N) = \lambda^p(N', x\,.\,\lambda_{-1}(N)). \tag{5.4.2}$$

5.5. Now let A be any λ-ring, and let N be an element of A such that $\lambda^k(N) = 0$ for $k > d$. We shall associate to A a new λ-ring A_N.

We first define a new multiplication on A by putting, for $x, y \in A$:

$$x \,._N\, y = x\,.\,y\,.\,\lambda_{-1}(N). \tag{5.5.1}$$

The ring A with this multiplication is no longer unital; we therefore consider the ring $\mathbb{Z} \times A$, obtained by adjoining a unit to A. Its operations are thus defined by: [243]

$$\begin{aligned}(n,x) + (n',x') &= (n+n', x+x') \\ (n,x)\,.\,(n',x') &= (nn', nx' + n'x + x \,._N\, x')\end{aligned} \tag{5.5.2}$$

We regard A as embedded in this ring by $x \rightsquigarrow (0,x)$. We then define a pre-λ-ring structure by:

$$\lambda_t(n,x) = (1+t)^n\,.\,\sum_{p=0}^{\infty} \lambda^p(N,x)\,.\,t^p. \tag{5.5.3}$$

This is a pre-λ-ring structure by (5.4.1). We shall denote by A_N the pre-λ-ring thus obtained. To see that it is a λ-ring, it suffices by substitution to give the proof in the case where A is the λ-ring generated over $\mathbb{Z}$ by the generators N, X, Y, subject to the relations $\lambda^k(N) = 0$ for $k > d$. We then define a homomorphism $A_N \to A$ by [print 342]sending $(1,0)$ to 1, and $(0,x)$ to $x\,.\,\lambda_{-1}(N)$. It follows immediately from (5.5.1) and (5.3.1) that this is a λ-homomorphism, and, since A is an integral domain in the case considered, one immediately verifies that it is injective; since A is a λ-ring, we deduce the result.

In the λ-ring A_N, we can define operations γ (3.2). For $x \in A$[244], the value of γ^p calculated on the element $(0,x)$ of A_N is given by: [245]

$$\gamma^p(0,x) = \sum_{i=0}^{p-1} \binom{p-1}{i} \lambda^{p-i}(N,x),$$

it is therefore an element of A; it is defined by the relation:

$$\gamma^p(N,x)\,.\,\lambda_{-1}(N) = \gamma^p(x\,.\,\lambda_{-1}(N)), \tag{5.5.4}$$

and the universal polynomials resulting from it.

5.6. Let A be the λ-ring generated over $\mathbb{Z}$ by the generators N, L, x, subject to the relations $\lambda^k(N) = 0$ for $k > d$, $\lambda^k(L) = 0$ for $k > 1$. We propose to calculate the $\lambda^p(N+L,x)$ in terms of the $\lambda^q(N,x)$.

By definition:

$$\lambda^p(N+L,x)\,.\,\lambda_{-1}(N)\,.\,\lambda_{-1}(L) = \lambda^p(x\,.\,\lambda_{-1}(N)\,.\,\lambda_{-1}(L)),$$

that is: [246]

$$
\begin{aligned}
&1+\lambda_{-1}(N)\,.\,\lambda_{-1}(L)\,.\sum_{k\geq 1}\lambda^k(N+L,x)\,.\,t^k\\
&=\lambda_t(x\,.\,\lambda_{-1}(N))\circ\lambda_t(1-L)\\
&=\frac{\lambda_t(x\,.\,\lambda_{-1}(N))}{\lambda_{Lt}(x\,.\,\lambda_{-1}(N))}\qquad\Big(\text{since }\lambda_t(1-L)=\frac{\lambda_t(1)}{\lambda_t(L)}=\frac{1+t}{1+Lt},\text{ and we apply (2.3.2)}\Big)\\
&=\frac{1+\ldots+\lambda^k(x\,.\,\lambda_{-1}(N))t^k+\ldots}{1+\ldots+\lambda^k(x\,.\,\lambda_{-1}(N))L^kt^k+\ldots}\\
&=1+\frac{\sum_{k\geq 1}\lambda^k(N,x)\,.\,\lambda_{-1}(N)\,.\,(1-L^k)t^k}{1+\sum_{k\geq 1}\lambda^k(N,x)\,.\,\lambda_{-1}(N)\,.\,L^kt^k}
\end{aligned}
$$

[print 343]Thus $\lambda^p(N+L,x)$ is the coefficient of t^p in the series:

$$\frac{\displaystyle\sum_{k\geq 1}\lambda^k(N,x)(1+\ldots+L^{k-1})t^k}{\displaystyle 1+\sum_{k\geq 1}\lambda^k(N,x)\lambda_{-1}(N)L^kt^k}.$$

Proposition 5.7. Let K be a binomial ring, A an augmented K-λ-algebra, and N an element of A such that $\lambda^k(N)=0$ for $k>d$. Then for every $x\in A$ and every $p\geq 1$, we have:

$$\gamma^p(N,x)\in \mathrm{Fil}^{p-d}A. \tag{5.7.1}$$

Let A_0 be the K-λ-algebra generated by generators N_0', X_0', subject to the relations $\gamma^k(N_0')=0$ for $k>d$. We define the augmentation homomorphism $A_0\xrightarrow{\epsilon}K$ by $\epsilon(N_0')=\epsilon(X_0')=0$. We know that A_0 is the ring of polynomials with coefficients in K in the indeterminates $\gamma^i(N_0')$, with $1\leq i\leq d$, and $\gamma^j(X_0')$, with $j\geq 1$, and that the λ-filtration on A_0 is identical to the weight filtration, assigning the weight i to $\gamma^i(N_0')$, $\gamma^i(X_0')$ (4.12 and 4.15).

Put $N_0=N_0'+d$, $X_0=X_0'+\epsilon(x)$. We shall first prove 5.7 for N_0 and X_0. By the preceding discussion, (5.7.1) is equivalent to:

$$\gamma^p(N_0,X_0)\,.\,\gamma^d(N_0')\in\mathrm{Fil}^pA_0.$$

Now $\gamma^d(N_0')=(-1)^d\lambda_{-1}(N_0)$; this relation can therefore be written:

$$(-1)^d\gamma^p(X_0\gamma^d(N_0))\in\mathrm{Fil}^pA_0$$

by (5.5.4). This relation does hold, since $\gamma^d(N_0)$ has zero augmentation.

By 4.7, there exists a K-λ-homomorphism $A_0\longrightarrow A$ compatible with the augmentations, sending N_0' to $N-d$, and X_0' to $x-\epsilon(x)$, hence N_0 to N and X_0 [print 344]to x, whence the result, since the homomorphism is compatible with the filtrations.

6. Chern ring

In this section, the base ring will be a binomial ring (2.7) K, fixed once and for all. We shall use the category $\mathcal{C}$ (1.1), and therefore return to the notation of Section 1, reserving the notation $\widehat{G}$ for the functor defined on $\mathcal{C}$ (1.1), and denoting by $\widehat{G}\circ i$ its restriction to $\mathcal{C}_0$ (1.8).

6.1. In 3.5 and 3.6 we defined morphisms of functors:

$$(\widehat{G}\circ i)\times(\widehat{G}\circ i)\xrightarrow{\nabla}\widehat{G}\circ i,\qquad \widehat{G}^t\circ i\xrightarrow{\gamma'}(\widehat{G}^u\circ i)\circ(\widehat{G}^t\circ i)_\nabla,\qquad \widehat{G}\circ i\xrightarrow{a}\widehat{G}\circ i,$$

the last being multiplication under the law ∇ by $\gamma_t(a)$, with $a \in K$.

By 1.9, we deduce morphisms:

$$\widehat{G} \times \widehat{G} \xrightarrow{\nabla} \widehat{G}_g, \qquad \widehat{G}^t \xrightarrow{\gamma'} \widehat{G}^u \circ i \circ \widehat{G}^t_{g\nabla}, \qquad \widehat{G} \xrightarrow{a} \widehat{G}_g;$$

and it follows from the isobaricity properties of the polynomials P'_n, $Q'_{i,j}$, $R_{n,a}$ (3.5, 3.6, 2.9) that the image of the first and third functors is in $\widehat{G}$, and that of the second in the subgroup of formal series in u whose coefficients in nonzero degree are in $\widehat{G}^t$.

Composing the first and third with the canonical morphism: $\widehat{G} \longrightarrow 1 + \widehat{\mathrm{Gr}}.^+$ (1.11), we obtain, by 1.12, two morphisms of functors on $\mathcal{G}$: [print 345][247]

$$(1 + \hat{.}^+) \times (1 + \hat{.}^+) \xrightarrow{*} 1 + \hat{.}^+,$$

$$1 + \hat{.}^+ \xrightarrow{a} 1 + \hat{.}^+,$$

defining on the functor $1 + \hat{.}^+$ a multiplication and operations of K. By 1.13, the multiplication is defined by its value on the element $(1 + X, 1 + Y)$ of $(1 + \widehat{K[X,Y]}^+)^2$; it will be denoted by $*$; we therefore have:

$$(1 + X) * (1 + Y) = \frac{1 + X + Y}{(1 + X)(1 + Y)}. \tag{6.1.1}$$

The n-th universal polynomial of this law is the isobaric component of weight n of P'_n (1.15), and will be denoted by P''_n.

The operations of K will again be denoted by exponentiation (3.8). The universal polynomials defining these operations are the isobaric components of lowest degree of the $R_{n,a}$ (1.15); since these are isobaric, they are the $R_{n,a}$ themselves.

Let $A \in \mathrm{Ob}(\mathcal{G})$; the ring $1 + \widehat{A}^+$ defined by the preceding laws is in general without a unit; we therefore extend this ring by adjoining the scalars of K, introducing the new ring $K \times (1 + \widehat{A}^+)$, endowed with the composition laws defined by: $\forall\, a, a' \in K$[248], $\varphi, \varphi' \in 1 + \widehat{A}^+$,

$$(a, \varphi) + (a', \varphi') = (a + a', \varphi\varphi') \ ;$$

$$(a, \varphi).(a', \varphi') = (aa', \varphi^{a'} \,.\, \varphi'^{\,a} \,.\, (\varphi * \varphi')). \tag{6.1.2}$$

The ring thus constructed admits (1,1) as its unit[249], and $1 + \widehat{A}^+$ is identified with the ideal $(0) \times (1 + \widehat{A}^+)$. It is naturally endowed with a K-algebra structure such that:

$$a.(a', \varphi) = (a, 1).(a', \varphi) = (aa', \varphi^a). \tag{6.1.3}$$

[print 346]We endow it with a K-λ-algebra structure as follows: we observed above that the image of the functor $\gamma' \colon \widehat{G}^t \longrightarrow \widehat{G}^u \circ i \circ \widehat{G}^t_g$ is in the subgroup of formal series in u whose coefficients in nonzero degree are in $\widehat{G}^t$. Using the canonical morphism: $\widehat{G} \longrightarrow 1 + \widehat{\mathrm{Gr}}.^+$, which gives: $\widehat{G} \longrightarrow K \times (1 + \widehat{\mathrm{Gr}}.^+)$[250], we finally deduce a morphism:

$$\widehat{G}^t \xrightarrow{\gamma'} \widehat{G}^u \circ i \circ (K \times (1 + \widehat{\mathrm{Gr}}.^+)),$$

which, by 1.12, defines a morphism of functors on $\mathcal{G}$:[251] [252]

$$1 + \hat{.}^+ \xrightarrow{\gamma'} \widehat{G}^u \circ i \circ (K \times (1 + \hat{.}^+)).$$

The universal polynomials defining this morphism are the coefficients of the $\gamma'^i(1 + X_1 + \ldots + X_n + \ldots)$; the j-th coefficient is the isobaric component of weight j of $Q'_{i,j}$ (a proof analogous to that of 1.15), and will be denoted by $Q''_{i,j}$.

Finally, we extend the morphism thus constructed to $K \times (1 + \widehat{\cdot}^+)$ naturally, using the λ-structure of K. We therefore obtain a pre-λ-ring structure on $K \times (1 + \widehat{\cdot}^+)$, defined by the corresponding operations γ, which satisfy:

$$\gamma_t(a, \varphi) = \gamma_t(a).\Big(1 + \sum_{p \geq 1} \gamma'^p(\varphi) t^p\Big), \tag{6.1.4}$$

and which will be denoted by γ instead of γ', since no confusion is possible.

To verify that this is a λ-ring structure, we verify the commutativity of (3.7.2); it is known for the elements of K; for the elements of $1 + \widehat{\cdot}^+$, it follows from that of (3.7.2) for $\widehat{G}$ by passage to the quotient. The result follows at once.

[print 347]**Definition 6.2.** The functor defined in 6.1 on the category of graded K-algebras, with values in the category of K-λ-algebras, is called the Chern ring; its value on $A \in \mathrm{Ob}(\mathcal{G})$ is denoted by $\mathrm{Ch}(A)$ (or, if there is a risk of confusion, $\mathrm{Ch}_K(A)$).

From formulas (6.1.1), (6.1.2), (6.1.4), and (3.6.2), we deduce the relations:

$$\begin{gathered}(1, 1 + X).(1, 1 + Y) = (1, 1 + X + Y) \\ \lambda^i(1, 1 + X) = 0 \quad \text{for } i \geq 2,\end{gathered} \tag{6.2.1}$$

relations which served to define the Chern ring in [1]. Conversely, from these formulas one can recover relations (6.1.1) and (6.1.4).

Let us indicate why the Chern ring is introduced. We place ourselves in the situation of Example 3.9, and denote by A the associated graded ring (for the filtration defined in 3.10) of the $K^\bullet$ of a commutatively locally ringed topos (which we suppose to be a λ-ring for simplicity); K is the augmentation ring. We shall see below that, for every element of an augmented λ-ring, one can define Chern classes with values in the associated graded ring (6.7). To every locally free sheaf of finite type, we can therefore associate an element of $\mathrm{Ch}(A)$ whose component on the factor K is the rank of the sheaf, and whose component on the factor $1 + \widehat{A}^+$ is the sum of the Chern classes. Formulas (6.2.1) are then the usual formulas giving the Chern classes and the rank of the tensor product of two invertible sheaves and of the exterior powers of an invertible sheaf (that is, the map just defined from $K^\bullet$ to $\mathrm{Ch}(A)$ is a λ-homomorphism).

[print 348]6.3. For every graded K-algebra A, we denote by $A_\mathbb{Q}$ the K-algebra $A \otimes_\mathbb{Z} \mathbb{Q}$, graded in the evident way. By 1.13, we define a functorial homomorphism: $1 + \widehat{A}^+ \xrightarrow{\mathrm{ch}} \widehat{A}^+_\mathbb{Q}$ by specifying its value on $1 + X$, namely

$$\mathrm{ch}(1 + X) = \exp(X) - 1. \tag{6.3.1}$$

This homomorphism is compatible with the multiplication defined in 6.1 on $1 + \widehat{A}^+$: by 1.13, it suffices to show this for the elements $1 + X, 1 + Y$ of $1 + \widehat{K[X, Y]}^+$; now we have:

$$\begin{aligned}\mathrm{ch}\big((1 + X) * (1 + Y)\big) &= \mathrm{ch}\Big(\frac{1 + X + Y}{(1 + X)(1 + Y)}\Big) \\ &= \mathrm{ch}(1 + X + Y) - \mathrm{ch}(1 + X) - \mathrm{ch}(1 + Y) \\ &= \exp(X + Y) - \exp(X) - \exp(Y) + 1 \\ &= \mathrm{ch}(1 + X).\,\mathrm{ch}(1 + Y).\end{aligned}$$

Let η denote the endomorphism (functorial in A) of $\widehat{A}^+_\mathbb{Q}$ defined by:

$$\eta\left(\sum_{k \geq 1} a_k t^k\right) = \sum_{k \geq 1} \frac{(-1)^{k-1}}{(k-1)!} . a_k t^k.$$

Then, $\forall \varphi \in 1 + \widehat{A}^+$,

$$\mathrm{ch}(\varphi) = \eta(\mathrm{Log}\,\varphi). \tag{6.3.2}$$

Indeed, ch and $\eta \circ \mathrm{Log}$ are two functorial homomorphisms of abelian groups, and for them to be equal it suffices that they agree on $1 + X$, which is evident.

We deduce that ch commutes with the operations of K: indeed, here again it suffices to show this for $1 + X$, and we have: [253]

$$\begin{aligned} \forall a \in K, \quad \mathrm{ch}((1+X)^a) &= \mathrm{ch}(\exp(a\,\mathrm{Log}(1+X))) \\ &= \eta(a\,\mathrm{Log}(1+X)) \\ &= a.\eta(\mathrm{Log}(1+X)) \\ &= a.\,\mathrm{ch}(1+X). \end{aligned}$$

[print 349]We can therefore extend ch in the evident way to a homomorphism of K-algebras: $K \times (1 + \widehat{A}^+) \longrightarrow K \oplus \widehat{A}^+_{\mathbb{Q}}$. We therefore have:

$$\mathrm{ch}(a, \varphi) = a + \eta(\mathrm{Log}\,\varphi). \tag{6.3.3}$$

Definition 6.4. The homomorphism $\mathrm{ch} : \mathrm{Ch}\,A \longrightarrow K \oplus \widehat{A}^+_{\mathbb{Q}}$ just defined is called the Chern character.

If A is of characteristic 0 (i.e. is a $\mathbb{Q}$-algebra), the Chern character is an isomorphism: indeed, let $\psi \in K \oplus \widehat{A}^+$; we can write $\psi = a + \psi_1$, where ψ_1 has no term of degree zero; we then define an inverse morphism of ch by $\psi \rightsquigarrow (a, \exp(\eta^{-1}(\psi_1)))$.

Proposition 6.5. Let A be a graded K-algebra; on $\mathrm{Ch}(A)$ consider the filtration defined as follows (degree filtration):

$$(\mathrm{Ch}(A))_0 = \mathrm{Ch}(A)\ ; \qquad \forall n \geq 1,\ (\mathrm{Ch}(A))_n = 0 \times 1 + \widehat{A}^{\geq n}, \tag{6.5.1}$$

where $\widehat{A}^{\geq n} = \prod_{i \geq n} A^i$. We have the following congruences[254]:

i) Let $n \geq 1$, and $\varphi \in (\mathrm{Ch}\,A)_n$; put $\varphi = 1 + a_n + \ldots$; then: [255]

$$\mathrm{ch}(\varphi) = \frac{(-1)^{n-1}}{(n-1)!}.a_n + \ldots\ ; \tag{6.5.2}$$

ii) Let $\varphi \in (\mathrm{Ch}\,A)_m$, $\psi \in (\mathrm{Ch}\,A)_n$; put $\varphi = 1 + a_m + \ldots$, $\psi = 1 + b_n + \ldots$; then: [256]

$$\varphi * \psi = 1 - \frac{(m+n-1)!}{(m-1)!(n-1)!}.a_m b_n + \ldots, \tag{6.5.3}$$

and consequently, the degree filtration is a ring filtration.

i) The Chern character is by construction a morphism of filtered functors; from the congruence $\varphi \equiv 1 + a_n$ mod. $(\mathrm{Ch}\,A)_{n+1}$, we deduce [print 350]therefore $\mathrm{ch}(\varphi) \equiv \mathrm{ch}(1 + a_n)$ mod. $\widehat{A}^{\geq n+1}_{\mathbb{Q}}$[257]; now $\mathrm{ch}(1 + a_n) = \eta(\mathrm{Log}(1 + a_n))$

$$= \frac{(-1)^{n-1}}{(n-1)!}\,a_n + \ldots$$

ii) By substitution, it suffices to prove formula (6.5.3) for $A = K[X_i; Y_j]_{i,j \in \mathbb{N}}$ and for $\varphi = 1 + \sum_{k \geq m} X_k$, $\psi = 1 + \sum_{k \geq n} Y_k$. Since K is torsion-free, we can embed this ring in its tensor

product with $\mathbb{Q}$, and thus see that ch is injective. Put $\varphi * \psi = 1 + Z_p + \ldots$, where Z_p is the first nonzero term, $\neq 1$, of $\varphi * \psi$. The multiplicativity of ch gives, by (6.5.2): [258]

$$\frac{(-1)^{p-1}}{(p-1)!}.Z_p + \ldots = \left(\frac{(-1)^{m-1}}{(m-1)!}.X_m + \ldots\right)\left(\frac{(-1)^{n-1}}{(n-1)!}.Y_n + \ldots\right),$$

whence the value of Z_p by identifying the terms of lowest degree.

6.6. The λ-ring $\mathrm{Ch}\, A$ is augmented over K: indeed, we define a λ-homomorphism from $\mathrm{Ch}\, A$ to K by $(a, \varphi) \rightsquigarrow a$. It is therefore endowed with a canonical filtration of an augmented λ-ring. We shall compare it with the one just defined.

Proposition 6.6.1. Let $x \in \mathrm{Ch}\, A$ be an element of the form $x = (n, 1 + \ldots + a_n)$. Then: [259]

i) $\lambda^i(x) = 0$ for $i > n$;

ii) $\lambda^n(x) = (1, 1 + a_1)$

iii) $\sum_{i=0}^{n} (-1)^i \lambda^i(x) = (0, 1 - (n-1)!a_n + \ldots)$,

the unwritten terms being of degree $> n$.

By functoriality, it suffices to give the proof in the case [print 351] where A is the ring $K[X_1, \ldots, X_n]$, filtered by weight with respect to the X_i, X_i having weight i, and for the element $(n, 1 + X_1 + \ldots + X_n)$.

We embed A in $A' = K[T_1, \ldots, T_n]$ by the symmetric functions, and thus obtain an injective homomorphism: $\mathrm{Ch}\, A \longrightarrow \mathrm{Ch}\, A'$; we then have:

$$(n, 1 + X_1 + \ldots + X_n) = \left(n, \prod_{i=1}^{n}(1 + T_i)\right) = \sum_{i=1}^{n}(1, 1 + T_i)$$

We deduce, by (6.2.1):

$$\lambda_t(x) = \prod_{i=1}^{n}(1 + (1, 1 + T_i)t),$$

and obtain:

$$\lambda^k(x) = 0 \text{ for } k > n, \quad \text{and} \quad \lambda^n(x) = \prod_{i=1}^{n}(1, 1 + T_i) = \left(1, 1 + \sum_{i=1}^{n} T_i\right) = (1, 1 + X_1)$$

Finally, [260]

$$\sum_{i=0}^{n}(-1)^i \lambda^i(x) = \lambda_{-1}(x) = \prod_{i=1}^{n}(1 - (1, 1 + T_i)) = (-1)^n . \prod_{i=1}^{n}(0, 1 + T_i).$$

Now, by (6.5.3), $\prod_{i=1}^{n}(0, 1 + T_i) = \left(0, 1 + (-1)^{n-1}(n-1)! \prod_{i=1}^{n} T_i + \ldots\right)$ that is: $\lambda_{-1}(x) = (0, 1 - (n-1)!X_n + \ldots)$.

Corollary 6.6.2. Relation 6.6.1.iii) holds for every element X of augmentation n.

To prove this, we shall use the lemma:

Lemma 6.6.3. For every n, $(\mathrm{Ch}\, A)_n$ is a λ-ideal of $\mathrm{Ch}\, A$.

We may suppose that $n \geq 1$. Let $x = (0, \varphi) \in (\mathrm{Ch}\, A)_n$; we can write $\gamma^i(0, \varphi) = (0, 1 + Q''_{i,1} + \ldots + Q''_{i,j} + \ldots)$ [261] where the $Q''_{i,j}$ are the isobaric components of weight j of the $Q'_{i,j}$ (3.6). For every j, the polynomials $Q''_{i,j}$ are therefore polynomials in the first j coefficients of φ. If $j < n$,

they are therefore zero, and consequently the $\gamma^i(x)$ belong to $(\mathrm{Ch}\,A)_n$, and [print 352]$(\mathrm{Ch}\,A)_n$ is a λ-ideal.

Now let $x = (n, 1 + a_1 + \ldots + a_n + \ldots)$ be an element of augmentation n of $\mathrm{Ch}\,A$; x is congruent mod. $(\mathrm{Ch}\,A)_{n+1}$ to $(n, 1+\ldots+a_n)$, and consequently $\lambda_{-1}(x)$ is congruent to $\lambda_{-1}(n, 1+\ldots+a_n) = (0, 1-(n-1)!a_n + \ldots)$, whence:

$$\lambda_{-1}(x) = (0, 1-(n-1)!a_n + \ldots).$$

Corollary 6.6.4. Let ϵ be the augmentation homomorphism of $\mathrm{Ch}\,A$; then, for every $x \in \mathrm{Ch}\,A$, we have:

$$\gamma^n(x - \epsilon(x)) = (0, 1 + (-1)^{n-1}(n-1)!a_n + \ldots)$$

By definition, $\gamma^n(x-\epsilon(x)) = (-1)^n \sum_{i=0}^n (-1)^i \lambda^i(x + n - \epsilon(x))$[262]. Since $x + n - \epsilon(x)$ has augmentation n, we can apply Corollary 6.6.2 to it, whence the result.

Corollary 6.6.5. With the notation of 3.10, $\mathrm{Fil}_n\,\mathrm{Ch}\,A \subset (\mathrm{Ch}\,A)_n$.

By the preceding corollary, $\gamma^n(x-\epsilon(x)) \in (\mathrm{Ch}\,A)_n$. The corollary then follows from the fact that the two filtrations are ring filtrations, and that the $\gamma^n(x-\epsilon(x))$, for varying n and x, generate the λ-ring filtration.

Corollary 6.6.7. Let A be a graded K-algebra of characteristic zero, with bounded degrees; then the two filtrations defined on $\mathrm{Ch}\,A$ are identical.

Suppose that $\mathrm{Fil}^p A = 0$, and let $x = (0, 1 + a_n + \ldots + a_p)$ be an element of $(\mathrm{Ch}\,A)_n$. Put $x' = \left(0, 1 + \frac{(-1)^{n-1}}{(n-1)!}a_n + a_{n+1} + \ldots + a_p\right)$; we then have: $\gamma^n(x') = (0, 1 + a_n + \ldots + a'_p)$, since x' has zero augmentation. Consequently, [print 353]$x - \gamma^n(x') \in (\mathrm{Ch}\,A)_{n+1}$. Iterating the operation $p-n$ times, we reach $(\mathrm{Ch}\,A)_{p+1}$, which is zero, and have therefore written x as an element of $\mathrm{Fil}^n\,\mathrm{Ch}\,A$. In view of (6.6.5), the two filtrations are therefore identical.

6.7. We now suppose that A is a λ-ring augmented over K, the augmentation homomorphism being denoted by ϵ. We therefore have on A the canonical filtration of an augmented λ-ring, and can thus associate to $\mathrm{Gr}\,A$ its Chern ring.

If $x \in A$, then $\gamma_t(x - \epsilon(x)) \in \widehat{G}(A)$ by definition of the filtration of an augmented λ-ring. We denote by $c(x)$, and call the total Chern class of x, the image of $\gamma_t(x-\epsilon(x))$ in $\widehat{G}(A)/\widehat{G}'(A) = 1 + \widehat{\mathrm{Gr}\,A}^+$; we denote by $c^i(x)$, and call the i-th Chern class of x, the i-th coefficient of $c(x)$; we therefore have:

$$c^i(x) = \mathrm{cl}_{\mathrm{Gr}^i A}\Big(\gamma^i(x-\epsilon(x))\Big)\ ; \qquad c(x) = 1 + c^1(x) + \ldots \tag{6.7.1}$$

We then put:

$$\widetilde{c}(x) = (\epsilon(x), c(x)), \tag{6.7.2}$$

and call $\widetilde{c}(x)$ the completed Chern class of x.

Proposition 6.8. The completed Chern class $\widetilde{c} : A \longrightarrow \mathrm{Ch}(\mathrm{Gr}\,A)$[263], is a K-λ-homomorphism of augmented K-λ-algebras.

It is clear that the homomorphism thus defined is additive; to verify that it is multiplicative, we write:

$$\begin{aligned}\gamma_t(xy - \epsilon(xy)) &= \gamma_t(\epsilon(y)(x-\epsilon(x))) \,.\, \gamma_t(\epsilon(x)(y-\epsilon(y))) \,.\, \gamma_t((x-\epsilon(x))(y-\epsilon(y)))\\ &= \big(\gamma_t(x-\epsilon(x))\big)^{\epsilon(y)} \,.\, \big(\gamma_t(y-\epsilon(y))\big)^{\epsilon(x)} \,.\, \big(\gamma_t(x-\epsilon(x))\nabla\gamma_t(y-\epsilon(y))\big)\end{aligned}$$

[print 354]whence the result by (6.1.2), after passage to the quotient.

If $a \in K$, $\widetilde{c}(a) = (a, 1)$, so $\widetilde{c}$ is a homomorphism of K-algebras; moreover, it is compatible with the augmentations.

Finally, it remains to see that it is a λ-homomorphism. We must therefore prove the commutativity of the diagram: [264]

$$\begin{array}{ccc} A & \xrightarrow{\gamma_t} & \widehat{G}_g(A) \\ \downarrow{\scriptstyle \widetilde{c}} & & \downarrow{\scriptstyle \widehat{G}_g(\widetilde{c})} \\ \mathrm{Ch}(\mathrm{Gr}\, A) & \xrightarrow{\gamma_t} & \widehat{G}_g(\mathrm{Ch}(\mathrm{Gr}\, A)) \end{array}$$

By additivity, it suffices to prove this for the elements of K, for which it is evident, and for the elements of zero augmentation of A, for which it follows, by the definition of c and (6.1.4), from that of diagram (3.7.2).

Proposition 6.9. Let A be an augmented K-λ-algebra, $z \in \mathrm{Fil}^m A$; then:

$$\gamma^m(z) \equiv (-1)^{m-1}(m-1)!z \text{ mod. } \mathrm{Fil}^{m+1} A,$$

i.e.

$$c^m(z) = \mathrm{cl}_{\mathrm{Gr}^m A}\Big((-1)^{m-1}(m-1)!z\Big).$$

We must see that these two elements give the same image in $\mathrm{Gr}^m A$, that is, that the m-th Chern class of z is the image of $(-1)^{m-1}(m-1)!z$.

We can write z as a sum $\sum a \,.\, \gamma^{i_1}(x_1) \ldots \gamma^{i_k}(x_k)$, with $a \in A$, $\epsilon(x_i) = 0$ [265] for every i, and $i_1 + \ldots + i_k \geq m$. We then have:

$$\widetilde{c}(z) = \sum \widetilde{c}(a) \,.\, \widetilde{c}(\gamma^{i_1}(x_1)) \ldots \widetilde{c}(\gamma^{i_k}(x_k))$$

Since $\widetilde{c}$ is a λ-homomorphism, we can write:

$$\widetilde{c}(\gamma^{i_j}(x_j)) = \gamma^{i_j}(\widetilde{c}(x_j)).$$

[print 355]Taking account of the definition of c, and of (6.6.4), we obtain:

$$\widetilde{c}(\gamma^{i_j}(x_j)) = \Big(0, 1 + (-1)^{i_j - 1}(i_j - 1)!\,\gamma^{i_j}(x_j) + \ldots\Big)$$

By (6.5.3), and the fact that $\widetilde{c}(a) = (a, 1)$, we finally obtain:

$$\begin{aligned} \widetilde{c}(z) &= \sum \Big(0, \Big(1 + (-1)^{i_1 + \ldots + i_k - 1}(i_1 + \ldots + i_k - 1)!\,\gamma^{i_1}(x_1) \ldots \gamma^{i_k}(x_k) + \ldots\Big)^a\Big) \\ &= \left(0, 1 + \sum_{\substack{(x_1, \ldots, x_k) \\ i_1 + \ldots + i_k = m}} (-1)^{m-1}(m-1)!a\,\gamma^{i_1}(x_1) \ldots \gamma^{i_k}(x_k) + \ldots\right) \\ &= (0, 1 + (-1)^{m-1}(m-1)!\bar{z} + \ldots), \end{aligned}$$

whence the result.

Proposition 6.10. Let K be a binomial ring, R an augmented K-λ-algebra, N an element of R such that $\lambda^k(N) = 0$ for $k > d$, and $x \in \mathrm{Fil}^i R$. Then:

$$\gamma^{d+i}(N, x) - (-1)^{d+i-1}(d+i-1)!x \in \mathrm{Fil}^{i+1} R.$$

We can find elements $x_{\alpha,j} \in R$[266], and $a_\alpha \in K$ such that:

$$x = \sum_\alpha a_\alpha \, \gamma^{i_1}(x_{\alpha,1}) \dots \gamma^{i_k}(x_{\alpha,k}),$$

with $i_1 + \dots + i_k \geq i$, and $\epsilon(x_{\alpha,j}) = 0$ for every α and every j. Let R_0 be the K-λ-algebra generated by generators N_0', $X_{\alpha,j}$ subject to the relations:

$$\gamma^k(N_0') = 0 \quad \text{for } k > d.$$

We define an augmentation homomorphism $R_0 \to K$ by $\epsilon(N_0') = \epsilon(X_{\alpha,j}) = 0$. By (4.9.1), R_0 is the ring of polynomials with coefficients in K in the indeterminates $\gamma^n(N_0')$, with $1 \leq n \leq d$, and $\gamma^p(X_{\alpha,j})$ with $p \geq 1$, and the λ-filtration of R_0 is identical to its weight filtration, assigning the weight k to $\gamma^k(N_0')$ and $\gamma^k(X_{\alpha,j})$ (4.12 and 4.15).

We put:

$$X_0 = \sum_\alpha a_\alpha \, \gamma^{i_1}(X_{\alpha,1}) \dots \gamma^{i_k}(X_{\alpha,k}),$$

[print 356]and shall first prove 6.10 for X_0 and $N_0 = N_0' + d$[267] (which do satisfy the hypotheses of 6.10). In view of the preceding remark on the filtration of R_0, it suffices for this to show that:

$$\gamma^{d+i}(N_0, X_0)\gamma^d(N_0') - (-1)^{d+i-1}(d+i-1)!X_0\gamma^d(N_0') \in \mathrm{Fil}^{d+i+1} R_0.$$

Now we have $\lambda_{-1}(N_0) = (-1)^d\gamma^d(N_0')$; this relation can therefore be written:

$$(-1)^d\gamma^{d+i}(X_0 \ (-1)^d\gamma^d(N_0')) - (-1)^{d+i-1}(d+i-1)!X_0\gamma^d(N_0') \in \mathrm{Fil}^{d+i+1} R_0.$$

Put $z = (-1)^d X_0\gamma^d(N_0')$; then $z \in \mathrm{Fil}^{d+i} R_0$, and we are reduced to showing:

$$\gamma^{d+i}(z) - (-1)^{d+i-1}(d+i-1)!z \in \mathrm{Fil}^{d+i+1} R_0.$$

Now this follows from 6.9.

To deduce 6.10 from this particular case, we consider the K-λ-homomorphism $R_0 \to R$ which sends N_0' to $N - d$ and $X_{\alpha,j}$ to $x_{\alpha,j}$ for every α and every j (4.7). This homomorphism is compatible with the λ-filtrations, and sends N_0 to N, and X_0 to x[268], whence the result.

Proposition 6.11. Suppose that K is of characteristic zero. Let $n \geq 0$ be an integer. Then the functor Ch is an equivalence of categories from the category of graded K-algebras A such that $A_0 = K$, $A_k = 0$ for $k > n$, to the category of augmented K-λ-algebras $\wedge$ such that $\mathrm{Fil}^{n+1} \wedge = 0$; a quasi-inverse functor is given by $\wedge \rightsquigarrow \mathrm{Gr} \wedge$.

Let A be a graded K-algebra such that $A_k = 0$ for $k > n$, endowed with the filtration associated to its grading; then $(\mathrm{Ch}\, A)_{n+1} = 0$, hence $\mathrm{Fil}^{n+1} \mathrm{Ch}\, A = 0$ (6.6.5). Conversely, if $\wedge$ is an augmented K-λ-algebra such that $\mathrm{Fil}^{n+1} \wedge = 0$, then $\mathrm{Gr}^0 \wedge = K$, $\mathrm{Gr}^k \wedge = 0$ for $k > n$.

We shall define isomorphisms of functors:

$$A \longrightarrow \mathrm{Gr}(\mathrm{Ch}\, A), \qquad \wedge \longrightarrow \mathrm{Ch}(\mathrm{Gr} \wedge).$$

[print 357]Since A is of finite grading, the graded rings associated with the λ-ring filtration and the degree filtration on $\mathrm{Ch}\, A$ are identical (6.6.7). We then define $u : A \longrightarrow \mathrm{Gr}_{\deg}(\mathrm{Ch}\, A)$ by the identity on A_0 and by: [269]

$$\forall\, i \geq 1, \quad \forall\, a_i \in A_i, \qquad u(a_i) = \mathrm{cl}_{\mathrm{Gr}^i}\left(0, 1 + (-1)^{i-1}(i-1)!a_i\right),$$

and extend by additivity. By relation (6.5.3), this is a homomorphism of graded rings, and it is easy to see that it is compatible with the operations of K. The inverse morphism u^{-1} can be

defined as follows: $\forall\, i \geq 1$, $\forall\, \bar{\varphi} \in \mathrm{Gr}^i(\mathrm{Ch}\, A)$, the i-th coefficient a_i of φ does not depend on the way in which $\bar{\varphi}$ has been lifted to $(\mathrm{Ch}\, A)_i$, and we put:

$$u^{-1}(\bar{\varphi}) = \frac{(-1)^{i-1}}{(i-1)!} a_i.$$

The homomorphism u is therefore an isomorphism.

The isomorphism $\wedge \to \mathrm{Ch}(\mathrm{Gr}\wedge)$[270] is defined by the completed Chern class $\widetilde{c}$. To verify that it does define an isomorphism, it suffices to verify that the homomorphism obtained by passing to the associated graded rings is one, since the filtrations are discrete. We therefore obtain:

$$\mathrm{Gr}\wedge \xrightarrow{\mathrm{Gr}(\widetilde{c})} \mathrm{Gr}_\lambda(\mathrm{Ch}(\mathrm{Gr}\wedge)) \simeq \mathrm{Gr}_{\deg}(\mathrm{Ch}(\mathrm{Gr}\wedge)) \xrightarrow{u^{-1}} \mathrm{Gr}\wedge.$$

It suffices to show that the composite is the identity. Since the homomorphisms are homomorphisms of graded K-algebras, it suffices, by the definition of the λ-ring filtration on $\wedge$, to show that it is the identity on the $\overline{\gamma^i(x)}$, with $\epsilon(x) = 0$. We then obtain:

$$\mathrm{Gr}(\widetilde{c})(\overline{\gamma^i(x)}) = \mathrm{cl}_{\mathrm{Gr}^i}(\gamma^i(\widetilde{c}(x))).$$

Now $\gamma^i(\widetilde{c}(x)) = (0, 1 + (-1)^{i-1}(i-1)!c^i(x) + \ldots)$, by (6.6.4); hence:

$$\mathrm{cl}_{\mathrm{Gr}^i}(\gamma^i(\widetilde{c}(x))) = u(c^i(x)),$$

[print 358]which shows that $u^{-1} \circ \mathrm{Gr}(\widetilde{c}) = \mathrm{Id}$.

Corollary 6.12. If K is of characteristic zero, and $\wedge$ is a K-λ-algebra with discrete filtration, then the completed Chern class:

$$\widetilde{c} : \wedge \longrightarrow \mathrm{Ch}(\mathrm{Gr}\wedge)$$

is an isomorphism.

The proof is contained in that of 6.11.

Definition 6.13. Let A be a graded K-algebra, and $\wedge$ an augmented K-λ-algebra; we call a Chern theory for $\wedge$ with values in A any λ-homomorphism: $\wedge \longrightarrow \mathrm{Ch}(A)$, compatible with the augmentations.

We define a functor on the category of graded K-algebras, with values in the category of sets, by associating with every graded K-algebra A the set of Chern theories for $\wedge$ with values in A, namely $\mathrm{Hom}_\lambda(\wedge, \mathrm{Ch}(A))$.

Proposition 6.14. The functor defined in 6.13 is representable. If K is of characteristic zero, and $\wedge$ has discrete filtration, it is represented by the graded ring associated with $\wedge$.

Let us first prove the first assertion. We consider the graded K-algebra generated by generators $c^i(x)$, with $i \geq 1$, $x \in \mathrm{Fil}^1 \wedge$, where $c^i(x)$ has degree i, and take its quotient by the following relations: [271]

i) $\quad c^k(x+y) = \sum_{i+j=k} c^i(x)c^j(y), \quad$ for $k \geq 1,\ x, y \in \mathrm{Fil}^1 \wedge$;

ii) $\quad c^k(xy) = P''_k(c^i(x), c^j(y)) \quad$ (cf. 6.1), $\quad$ for $k \geq 1,\ x, y \in \mathrm{Fil}^1 \wedge$;

iii) $\quad c^k(ax) = R_{k,a}(c^i(x)) \quad$ (cf. 6.1), $\quad$ for $k \geq 1,\ a \in K,\ x \in \mathrm{Fil}^1 \wedge$;

iv) $\quad c^k(\gamma^j(x)) = Q''_{j,k}(c^i(x)) \quad$ (cf. 6.1), $\quad$ for $j, k \geq 1,\ x \in \mathrm{Fil}^1 \wedge$,

[print 359]These relations being isobaric, the quotient is a graded K-algebra Ω. We define a morphism $\wedge \xrightarrow{\widetilde{c}_0} \mathrm{Ch}(\Omega)$ by:

$$x \rightsquigarrow (\epsilon(x), 1 + c^1(x - \epsilon(x)) + \ldots).$$

The imposed relations show that this is a Chern theory for $\wedge$ with values in Ω. If in addition we have a Chern theory f, with values in a graded K-algebra A, there is a unique homomorphism: $\Omega \to A$, such that f factors through $\mathrm{Ch}(\Omega)$; it is defined by sending, for every $i \geq 1$ and every $x \in \mathrm{Fil}^1 \wedge$, the element $c^i(x)$ to the i-th coefficient of $f(x)$ (which is well defined since f is a λ-homomorphism).

Now suppose that K is of characteristic zero and $\wedge$ has discrete filtration; let n be such that $\mathrm{Fil}^n \wedge = 0$. Then the graded algebra Ω is zero in degrees $\geq n$. To prove this, we must show that for $x_1, \ldots, x_p \in \mathrm{Fil}^1 \wedge$ and $k_1 + \ldots + k_p \geq n$, we have:

$$c^{k_1}(x_1) \ldots . c^{k_p}(x_p) = 0.$$

Now, by hypothesis, we have:

$$\gamma^{k_1}(x_1) \ldots . \gamma^{k_p}(x_p) = 0 \ ;$$

Applying the λ-homomorphism $\widetilde{c}_0$, we obtain:

$$\gamma^{k_1}(\widetilde{c}_0(x_1)) \ldots . \gamma^{k_p}(\widetilde{c}_0(x_p)) = (0,1).$$

Now, by (6.6.4) and (6.5.3), this implies:

$$(k_1 + \ldots + k_p)!\, c^{k_1}(x_1) \ldots c^{k_p}(x_p) = 0.$$

Since K is of characteristic zero, we have the desired result.

The universal property of Ω, applied to the canonical homomorphism $\widetilde{c}$, gives a commutative diagram: [print 360]

$$\begin{array}{ccc} \wedge & \xrightarrow{\widetilde{c}} & \mathrm{Ch}(\mathrm{Gr}\wedge) \\ {\scriptstyle \widetilde{c}_0}\downarrow & \nearrow & \\ \mathrm{Ch}(\Omega) & & \end{array}$$

By Proposition 6.11, the homomorphism: $\Omega \to \mathrm{Gr}\wedge$ is an isomorphism if and only if the homomorphism: $\mathrm{Ch}(\Omega) \to \mathrm{Ch}(\mathrm{Gr}\wedge)$ is one. Since it is visibly surjective, the latter is also surjective, and it therefore suffices to show that $\mathrm{Ch}(\Omega) \to \mathrm{Ch}(\mathrm{Gr}\wedge)$ is injective. By functoriality of Ch, we obtain the diagram: [272]

$$\begin{array}{ccc} \wedge & \xrightarrow{\widetilde{c}_0} & \mathrm{Ch}(\Omega) \\ {\scriptstyle \widetilde{c}}\downarrow & & \downarrow{\scriptstyle \widetilde{c}} \\ \mathrm{Ch}(\mathrm{Gr}\wedge) & \xrightarrow{\mathrm{Ch}(\widetilde{c}_0)} & \mathrm{Ch}\big(\mathrm{Gr}_\lambda(\mathrm{Ch}(\Omega))\big); \end{array}$$

since Ω has bounded degrees, the right vertical arrow is an isomorphism (6.12), and using 6.11, we therefore obtain a morphism: $\mathrm{Gr}\wedge \longrightarrow \Omega$, the composite of the two being the identity by the universal property of Ω.

APPENDIX: The Adams operations ψ^k

We follow here a presentation due to J.P. Serre.

7.1. Let K be a pre-λ-ring. For every $x \in K$, we denote by $\frac{d}{dt}(\lambda_t(x))$ the derivative with respect to t of $\lambda_t(x)$. We then define elements of K, denoted by $\psi^k(x)$, by putting:

$$\frac{d}{dt}(\lambda_t(x))/\lambda_t(x) = \sum_{k=1}^{\infty} (-1)^{k-1} \psi^k(x) t^{k-1}. \tag{7.1.1}$$

The maps $\psi^k : K \to K$ are additive; indeed, we have: [273]

$$\frac{d}{dt}(\lambda_t(x+y))/\lambda_t(x+y) = \frac{d}{dt}(\lambda_t(x)\lambda_t(y))/(\lambda_t(x)\lambda_t(y)) \\ = \frac{d}{dt}(\lambda_t(x))/\lambda_t(x) + \frac{d}{dt}(\lambda_t(y))/\lambda_t(y)\ ;$$

[print 361]and therefore, for every k:

$$\psi^k(x+y) = \psi^k(x) + \psi^k(y).$$

By (7.1.1), the ψ^k are isobaric polynomials of weight k in the λ^i, with $i = 1, \ldots, k$. For example, let us give the values of ψ^1 and ψ^2:

$$\psi^1(x) = \lambda^1(x) = x, \qquad \psi^2(x) = x^2 - 2\lambda^2(x).$$

7.2. Let K be a ring of characteristic 0; suppose given additive maps ψ^k from K to K, such that $\psi^1 = \mathrm{Id}_K$. We then define a pre-λ-structure on K by putting:

$$\lambda_t(x) = \exp\left(\sum_{k=1}^{\infty}(-1)^{k-1}\psi^k(x)t^k/k\right). \tag{7.2.1}$$

Multiplicativity follows from the additivity of the ψ^k. There is therefore an equivalence between giving a pre-λ-structure on K and giving the corresponding operations ψ^k.

7.3. It is clear from formula (7.1.1) that the ψ^k commute with λ-homomorphisms. If K, K' are pre-λ-rings of characteristic zero, it is necessary and sufficient that a homomorphism $f : K \to K'$ commute with the ψ^k for it to be a λ-homomorphism: this follows from 7.2, the λ^i being expressed as polynomials in the ψ^k.

Examples 7.4. i) For $K = \mathbb{Z}$, endowed with its λ-ring structure, we have:

$$\frac{d}{dt}(\lambda_t(n))/\lambda_t(n) = n(1+t)^{n-1}/(1+t)^n = n.\sum_{k=1}(-1)^{k-1}t^{k-1}.$$

Consequently, the ψ^k are all equal to the identity map of $\mathbb{Z}$.

ii) Let us work in the category $\mathcal{C}_0$ of $\mathbb{Z}$-algebras (1.8). The [print 362]operations ψ^k define, by 7.3, additive endomorphisms of the functor $\widehat{G}$ with values in λ-rings; by 1.10, they are characterized by their value on $1 + Xt$. We have: [274]

$$\frac{d}{du}(\lambda_u(1+Xt))/\lambda_u(1+Xt) = \frac{1+Xt}{\underline{1} + \{1+Xt\}u} = \sum_{k=1}^{\infty}(-1)^{k-1}\{1+Xt\}^{\circ k}u^{k-1}.$$

Consequently:

$$\psi^k(1+Xt) = 1 + X^k t. \tag{7.4.1}$$

Proposition 7.5. i) <u>If</u> K <u>is a</u> λ-<u>ring</u>, <u>the operations</u> ψ^k <u>satisfy</u> <u>the identities</u>:

$$\psi^k(1) = 1\ ; \qquad \psi^k(xy) = \psi^k(x)\psi^k(y)\ ; \qquad \psi^k \circ \psi^{k'} = \psi^{kk'}. \tag{7.5.1}$$

ii) <u>Conversely</u>, <u>if the</u> ψ^k <u>satisfy these identities</u>[275], <u>and if</u> K <u>is of characteristic zero</u>, K <u>is a</u> λ-<u>ring</u>.

i) If K is a λ-ring, we have $\lambda_t(1) = 1 + t$, whence: $\frac{d}{dt}(\lambda_t(1))/\lambda_t(1) = 1/(1+t) = \sum_{k=1}^{\infty}(-1)^{k-1}t^{k-1}$[276], and consequently:

$$\psi^k(1) = 1.$$

To prove the second of relations (7.5.1), we consider the diagram:

$$\begin{array}{ccccc} K \times K & \xrightarrow{\lambda_t \times \lambda_t} & \widehat{G}(K) \times \widehat{G}(K) & \xrightarrow{\Phi^k \times \Phi^k} & K \times K \\ \downarrow & & \downarrow \circ & & \downarrow \\ K & \xrightarrow{\lambda_t} & \widehat{G}(K) & \xrightarrow{\Phi^k} & K, \end{array} \tag{7.5.2}$$

where the vertical arrows are defined by multiplication, and the maps Φ^k as follows: the "logarithmic derivative" map is [print 363]a group homomorphism from $\widehat{G}(K)$ to $K[[t]]$; the map which assigns to a series its $(k-1)$-st coefficient, multiplied by $(-1)^{k-1}$, is an additive homomorphism from $K[[t]]$ to K; the morphism Φ^k will be the composite of the two. It is clear that $\Phi^k \circ \lambda_t = \psi^k$ by definition (7.1.1).

The second of formulas (7.5.1) therefore expresses the commutativity of the diagram written above. If we suppose that K is a λ-ring, the left square is commutative, and it suffices to verify the commutativity of the right one. Now this is a diagram of homomorphisms of functors on $\mathcal{C}_0$, with source $\widehat{G} \times \widehat{G}$; it therefore suffices, by 1.10, to verify commutativity for $(1+Xt, 1+Yt)$; we then have: [277]

$$\Phi^k(1+Xt) = X^k \qquad \big((k-1)\text{-st coefficient of } (-1)^{k-1} X/(1+Xt)\big)\ ; \qquad \Phi^k(1+Yt) = Y^k\ ;$$

$\Phi^k\big((1+Xt) \circ (1+Yt)\big) = \Phi^k(1+XYt) = (XY)^k = \Phi^k(1+Xt)\Phi^k(1+Yt)$, which gives the result.

To prove the last one, we first observe that the diagram: [278]

$$\begin{array}{ccc} \widehat{G}^t(K) & \xrightarrow{\lambda_u} & \widehat{G}^u \circ \widehat{G}^t(K) \\ {\scriptstyle \Phi^{kk'}}\downarrow & & \downarrow{\scriptstyle \Phi_u^{k'}} \\ K & \xleftarrow{\Phi^k} & \widehat{G}^t(K), \end{array} \tag{7.5.3}$$

is commutative. As above, it indeed suffices to see this for $1+Xt$, and we have:

$$\Phi_u^{k'}(\lambda_u(1+Xt)) = \Phi_u^{k'}(1+\{1+Xt\}u) = (1+Xt)^{\circ k'} = 1+X^{k'}t\ ;$$

$$\Phi^k(1+X^{k'}t) = X^{kk'} = \Phi^{kk'}(1+Xt).$$

Consequently, $\psi^{kk'} = \Phi^k \circ \Phi_u^{k'} \circ \lambda_u \circ \lambda_t = \Phi^k \circ \psi^{k'} \circ \lambda_t$.

Moreover, $\psi^k \circ \psi^{k'} = \Phi^k \circ \lambda_t \circ \Phi^{k'} \circ \lambda_t = \Phi^k \circ \lambda_t \circ \psi^{k'}$.

[print 364]Now, if K is a λ-ring, the diagram:

$$\begin{array}{ccc} K & \xrightarrow{\lambda_t} & \widehat{G}(K) \\ {\scriptstyle \psi^{k'}}\downarrow & & \downarrow{\scriptstyle \psi^{k'}} \\ K & \xrightarrow{\lambda_t} & \widehat{G}(K) \end{array} \tag{7.5.4}$$

is commutative by 7.3, since λ_t is a λ-homomorphism. The desired relations follow at once.

ii) If K is of characteristic zero, the "logarithmic derivative" map is an isomorphism from $\widehat{G}(K)$ onto $K[[t]]$, since one can integrate. Consequently, two elements of $\widehat{G}(K)$ are equal if and only if, for every k, they have the same image under Φ^k.

To show that λ_t is a ring homomorphism, we must show the commutativity of the left square of diagram (7.5.2), and by the preceding remark, it suffices to show that for every k, diagram (7.5.2) is commutative, since the right square always is. Now these are precisely the relations $\psi^k(xy) = \psi^k(x)\psi^k(y)$.

To show that λ_t is a λ-homomorphism, it suffices to show that diagram (7.5.4) is commutative for every k', by 7.3. Now, by the preceding remark, this is equivalent to showing for all k, k' that $\Phi^k \circ \psi^{k'} \circ \lambda_t = \Phi^k \circ \lambda_t \circ \psi^{k'}$, that is, the relations $\psi^{kk'} = \psi^k \circ \psi^{k'}$.

Finally, the relation $\lambda_t(1) = 1 + t$ is obtained by integration from the relations $\psi^k(1) = 1$.

BIBLIOGRAPHY

[1] A. Grothendieck, La théorie des classes de Chern, Bull. Soc. math. France, 86, 1958, p. 137 à 154.

[2] F. Hirzebruch, Neue topologische Methoden in der algebraischen Geometrie.

[print 365]

Exposé VI

THE $K^\bullet$ OF A PROJECTIVE BUNDLE: COMPUTATIONS AND CONSEQUENCES

by Pierre BERTHELOT

1. Computation of the $K^\bullet$ of a projective bundle: the case of locally free sheaves of finite type

In this section, the notation $K^\bullet(X)$ will denote the Grothendieck group of locally free sheaves of finite type on a scheme X.

The aim of this section is to prove the following theorem:

Theorem 1.1. Let S be a quasi-compact scheme, $\underline{E}$ a $\underline{O}_S$-Module locally free of rank $r+1$, $X = \mathbf{P}(\underline{E})$[279] the associated projective bundle; the structural morphism: $X \xrightarrow{f} S$ defines a ring homomorphism: $K^\bullet(S) \xrightarrow{f^*} K^\bullet(X)$ endowing $K^\bullet(X)$ with the structure of a $K^\bullet(S)$-algebra. Let $\xi \in K^\bullet(X)$ be the class of the sheaf $\underline{O}_X(1)$. Then $K^\bullet(X)$ is a free $K^\bullet(S)$-module with basis $1, \dots, \xi^r$.

Plan of the proof:

1°) The powers of ξ generate $K^\bullet(X)$ as a $K^\bullet(S)$-module.

To prove this, we shall establish the following points:

a) If $\underline{F}$ is a $\underline{O}_X$-Module locally free of finite type, there exists a graded $\mathbf{S}(\underline{E})$-Module $\underline{M}$ of finite presentation[280], flat over $\underline{O}_S$, such that $\underline{F} = \underline{\mathrm{Proj}}_0(\underline{M})$ ($\mathbf{S}(\underline{E})$ being the symmetric algebra of $\underline{E}$; for the notation $\underline{\mathrm{Proj}}_0$, see EGA II 8.12.1).[281]

b) There exists a finite graded resolution of $\underline{M}$ by graded $\mathbf{S}(\underline{E})$-Modules $\underline{L}_i$ locally free of finite type: $0 \longrightarrow \underline{L}_{r+1} \longrightarrow \dots \longrightarrow \underline{L}_0 \longrightarrow \underline{M} \longrightarrow 0$.

c) The $\underline{L}_i$ admit a finite filtration whose quotients are of the form $\underline{N}_j \otimes_{\underline{O}_S} \mathbf{S}(\underline{E})(j)$, where the $\underline{N}_j$ are locally free of finite type on S.

[print 366]2°) $1, \dots, \xi^{r+1}$ are linearly dependent over $K^\bullet(S)$.

3°) $1, \dots, \xi^r$ are linearly independent over $K^\bullet(S)$.

1.2. For convenience, we shall recall here a few classical definitions and properties of graded Modules.

Let S be a scheme, $\underline{B}$ a graded $\underline{O}_S$-Algebra, $\underline{M}$ a graded $\underline{B}$-Module, For every graded $\underline{B}$-Module $\underline{N}$, we denote as usual by $\underline{N}(n)$ the graded $\underline{B}$-Module defined by:

$$(\underline{N}(n))_k = \underline{N}_{n+k} \quad \text{for every } k.$$

We say that $\underline{M}$ is a locally free graded $\underline{B}$-Module (resp. locally free of finite type) if every point of S has an open neighborhood U such that $\underline{M}|U$ is isomorphic, as a graded $\underline{B}$-Module, to a direct sum of Modules of the form $\underline{B}(n)|U$ (resp. a finite direct sum).

We say that $\underline{M}$ is a graded $\underline{B}$-Module of finite presentation if every point of S has an open neighborhood U such that there is an exact sequence $\underline{P} \longrightarrow \underline{Q} \longrightarrow \underline{M}|U \longrightarrow 0$, where $\underline{P}$ and $\underline{Q}$

are free graded $(\underline{B}|U)$-Modules of finite type, and the homomorphisms are compatible with the gradings.

1.2.1. For every integer n, we can define a canonical homomorphism of graded $\underline{B}$-Modules: [282]

$$\underline{M}_n \otimes_{\underline{O}_S} \underline{B}(-n) \longrightarrow \underline{M}, \tag{1.2.1}$$

by sending $x \otimes a$, where x is a section of $\underline{M}_n$, and a a section of $(\underline{B}(-n))_k = \underline{B}_{k-n}$, to the element $a.x$ of $\underline{M}_k$.

1.2.2. We say that a graded $\underline{B}$-Module $\underline{M}$ is free of finite type and homogeneous of degree n if it is isomorphic as a graded $\underline{B}$-Module to $(\underline{B}(-n))^k$ for some integer k.

[print 367]Let $\underline{M}$ be a free graded $\underline{B}$-Module of finite type. By 1.2, we can find a graded isomorphism:

$$\underline{M} \simeq \bigoplus_{n=n_0}^{n_1} \big(\underline{B}(-n)\big)^{k_n}. \tag{1.2.2}$$

We thus obtain a decomposition of $\underline{M}$ as a direct sum of free graded $\underline{B}$-Modules of finite type that are homogeneous, namely:

$$\underline{M} \simeq \bigoplus_{n=n_0}^{n_1} \underline{M}^{(n)},$$

where $\underline{M}^{(n)}$ is free homogeneous of degree n and of rank k_n.

Proposition 1.2.3. The integers k_n occurring in decomposition (1.2.2) do not depend on the chosen decomposition.

Suppose given two decompositions, hence a graded isomorphism:

$$\bigoplus_{n=n_0}^{n_1} \big(\underline{B}(-n)\big)^{k_n} \xrightarrow{\sim} \bigoplus_{n=n'_0}^{n'_1} \big(\underline{B}(-n)\big)^{k'_n}.$$

The proof proceeds by induction on the number of nonzero free homogeneous direct summands occurring in the decomposition on the left. If this is 0, then $\underline{M} = 0$, and the proposition is evident. Suppose it proved for $n_1 - n_0 = m$, and let us prove it for $n_1 - n_0 = m + 1$. The given isomorphism, being compatible with the gradings, gives an isomorphism on the homogeneous components of the two sides, and in particular an isomorphism of the homogeneous components of degree $< n_0$, which shows that for $n < n_0$, we have $k'_n = 0$, and on the homogeneous components of degree n_0, which shows that they have the same rank, and consequently that $k_{n_0} = k'_{n_0}$. The given isomorphism therefore induces an isomorphism of the free homogeneous direct summands of lowest degree in the two decompositions, and hence passes to the quotient by these direct summands; we then apply the induction hypothesis to the quotient Module, whence the result.

[print 368]It follows in particular from 1.2.3 that for every free graded Module of finite type $\underline{M}$, the number of elements of a homogeneous basis of $\underline{M}$ having a given degree does not depend on the chosen basis. In particular, the integers n_0 and n_1 occurring in decomposition (1.2.2), defined by the conditions $k_{n_0} \neq 0$, $k_{n_1} \neq 0$, do not depend on the decomposition.[283]

Proposition 1.3. Let S be a quasi-compact scheme, $\underline{E}$ a $\underline{O}_S$-Module locally free of finite type, X the associated projective bundle. Let $\underline{F}$ be a $\underline{O}_X$-Module of finite presentation, flat over S. There exists an integer n_0, such that for $n \geq n_0$, we have:

i) $R^i f_*(\underline{F}(n)) = 0$ for $i \geq 1$;

ii) $f_*(\underline{F}(n))$ is locally free of finite type on S;

iii) $\underline{M} = \bigoplus_{n \geq n_0} f_*(\underline{F}(n))$ is a graded $\mathbf{S}(\underline{E})$-Module of finite presentation (1.2).

The desired properties are local on S; since S is quasi-compact, we can, by taking the largest of the integers found, reduce to the case where S is affine.

First suppose that S is noetherian.

Property i) then follows from Serre's theorem.

Property ii) follows from EGA III 7.9.10, with n_0 chosen so that property i) holds.

By EGA II 3.4.5, there exists a quasi-coherent graded $\mathbf{S}(\underline{E})$-Module $\underline{M}'$, of finite type, such that $\underline{F} = \underline{\mathrm{Proj}}_0(\underline{M}')$. Moreover, $\Gamma_*(\underline{F}) = \Gamma_*(\underline{\mathrm{Proj}}_0\underline{M}') = \bigoplus_{n\in\mathbb{Z}} f_*(\underline{F}(n))$ is TN-isomorphic to $\underline{M}'$ by EGA III 2.3.1; thus there exists an integer n_1 such that $\bigoplus_{n\geq n_1} f_*(\underline{F}(n)) \simeq \bigoplus_{n\geq n_1} \underline{M}'_n$, [print 369]and hence such that $\bigoplus_{n\geq n_1} f_*(\underline{F}(n))$ is a $\mathbf{S}(\underline{E})$-Module of finite type, and therefore of finite presentation by the noetherian hypotheses; the result moreover holds for every integer n_1, since the $f_*(\underline{F}(n))$ are coherent by Serre's theorem.

The proposition is therefore proved in the case where S is noetherian.

Suppose now that we have a Cartesian square:

$$\begin{array}{ccc} X & \xrightarrow{g} & X_0 \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle f_0} \\ S & \xrightarrow{g_0} & S_0, \end{array}$$

with S_0 noetherian, $X_0 = \mathbf{P}(\underline{E}_0)$[284]; let $\underline{F}_0$ be a $\underline{O}_{X_0}$[285]-Module of finite presentation, flat over S_0, and let $\underline{F} = g^*(\underline{F}_0)$. By the preceding argument the proposition holds for $\underline{F}_0$. By EGA III 7.7.5 and 7.8.5, the base-change theorem holds under the preceding hypotheses; we immediately deduce that the theorem holds for $\underline{F}$.

To complete the proof, it therefore suffices to show that the general case (with S affine) can be obtained from the noetherian case by base change.

Since S is affine, S is a filtered projective limit of noetherian affine schemes (if $S = \mathrm{Spec}(A)$, we write A as the inductive limit of its subalgebras of finite type over $\mathbb{Z}$). Thus let $S = \varprojlim S_\lambda$. By EGA IV 8.5.2 and 8.5.5, there exists an index λ and a $\underline{O}_{S_\lambda}$-Module locally free of finite type $\underline{E}_\lambda$ such that $\underline{E} = u_\lambda^*(\underline{E}_\lambda)$ (denoting by u_λ the morphism $S \to S_\lambda$, and by $u_{\lambda\mu}$ the transition morphisms $S_\mu \to S_\lambda$).

If we put, for $\mu \geq \lambda$, $\underline{E}_\mu = u_{\lambda\mu}^*(\underline{E}_\lambda)$, and $X_\mu = \mathbf{P}(\underline{E}_\mu)$[286], then for every μ, we have $X = X_\mu \times_{S_\mu} S$, and $X = \varprojlim X_\mu$. Consequently, there exists an index $\lambda' \geq \lambda$ [print 370]and a sheaf $\underline{F}_{\lambda'}$ of finite presentation on $X_{\lambda'}$, such that $\underline{F} = u_{\lambda'}'^*(\underline{F}_{\lambda'})$ ($u'_{\lambda'}$ being the morphism $X \to X_{\lambda'}$). Since the X_μ and the $\underline{F}_\mu$ are of finite presentation, we can, by EGA IV 11.2.6, find an index λ'' such that $\underline{F}_{\lambda''}$ is flat over $S_{\lambda''}$. All the hypotheses of 1.3 are therefore satisfied, which completes the proof.

Corollary 1.4. Under the hypotheses of 1.3, there exists a graded $\mathbf{S}(\underline{E})$-Module $\underline{M}$, of finite presentation, and flat over $\underline{O}_S$, such that $\underline{F} = \underline{\mathrm{Proj}}_0(\underline{M})$.

Let n_0 be an integer such that the properties of 1.3 hold. If $\underline{M} = \bigoplus_{n\geq n_0} f_*(\underline{F}(n))$, we have seen that $\underline{M}$ is of finite presentation, and it is flat over S, since the $f_*(\underline{F}(n))$, for $n \geq n_0$, are locally free on S.

Moreover, $\underline{F} = \underline{\mathrm{Proj}}_0(\underline{M})$ by EGA II 3.4.

Let us turn to part b) of the proof.

Lemma 1.5.1. Let A be a ring, B a positively graded A-algebra such that $B_0 = A$, M a graded B-module. We suppose that M is projective of finite type, as an ungraded B-module. Let $\widetilde{B}$ be the sheaf of algebras on $\mathrm{Spec}(A)$ defined by B, $\widetilde{M}$ the $\widetilde{B}$-Module associated with M; then $\widetilde{M}$ is, locally on $\mathrm{Spec}(A)$, free of finite type over $\widetilde{B}$, as a graded $\widetilde{B}$-Module (1.2).

Since M is of finite type, its degrees are bounded below. Moreover, $M \otimes_B A$ is projective of finite type over A, and consequently has only finitely many nonzero homogeneous components, themselves projective of finite type over A; by localizing on $\mathrm{Spec}(A)$, we can therefore suppose that $M \otimes_B A$ is free of finite type over A. Finally, since M is projective, $\mathrm{Tor}_1^B(M, A) = 0$. These properties imply that M is free of finite type over B as a graded module (see Bourbaki, Alg. ch. II §11 prop. 7), so $\widetilde{M}$ is free of finite type over $\widetilde{B}$, q.e.d.

[print 371]**Lemma 1.5.2.** Let A be a ring, B an A-algebra of finite presentation, M a B-module of finite presentation. We suppose B and M flat over A, and that for every residue field k of A, $M \otimes_A k$ has projective dimension $\leq n$ over $B \otimes_A k$ (which is the case if $B = A[t_0, \ldots, t_r]$). Under these conditions, M admits a resolution of length n by projective B-modules of finite type:

$$0 \longrightarrow L_n \longrightarrow \ldots \longrightarrow L_0 \longrightarrow M \longrightarrow 0.$$

This lemma follows immediately from EGA IV 12.3.2.

Proposition 1.6. Let S be a quasi-compact scheme, $\underline{E}$ a $\underline{O}_S$-Module locally free of rank $r+1$, $\underline{B}$ its symmetric algebra, with its usual grading, $\underline{M}$ a $\underline{B}$-Module that is graded and of finite presentation, flat over S. Then there exists a graded resolution:

$$0 \longrightarrow \underline{L}_{r+1} \longrightarrow \ldots \longrightarrow \underline{L}_0 \longrightarrow \underline{M} \longrightarrow 0,$$

where the $\underline{L}_i$ are $\underline{B}$-Modules that are graded and locally free of finite type (as graded $\underline{B}$-Modules).

Let us first show that for every graded $\underline{B}$-Module $\underline{M}$ satisfying the conditions of the proposition, there exists a surjective graded homomorphism: $\underline{L} \to \underline{M}$, where $\underline{L}$ is a $\underline{B}$-Module locally free of finite type.

Since $\underline{M}$ is of finite type, there exists an integer N such that $\underline{M}$ is generated over $\underline{B}$ by its components of degree $\leq N$. We then put:

$$\underline{L} = \bigoplus_{i \leq N} \underline{M}_i \otimes_{\underline{O}_S} \underline{B}(-i);$$

the homomorphism from $\underline{L}$ to $\underline{M}$ is defined using (1.2.1), and it is clear that it is surjective.[287]

By 1.5.2, $\underline{M}$ admits an (ungraded) resolution by $\underline{B}$-Modules locally free of finite type on every affine open set. Consequently, if [print 372]$\underline{M}$ has a global partial resolution:

$$\underline{L}_i \longrightarrow \underline{L}_{i-1} \longrightarrow \ldots \longrightarrow \underline{L}_0 \longrightarrow \underline{M} \longrightarrow 0,$$

by $\underline{B}$-Modules locally free of finite type, the kernel $\underline{Z}_i$ of $\underline{L}_i \to \underline{L}_{i-1}$ is again a $\underline{B}$-Module of finite type (I. 2.11). It is even of finite presentation (by making the same argument with $i+1$ instead of i). Finally, it is flat over S. It therefore satisfies the same hypotheses as $\underline{M}$, which allows us to construct step by step a global graded resolution of $\underline{M}$ by graded $\underline{B}$-Modules locally free of finite type, by the initial remark. For $i \geq r+1$, $\underline{Z}_i$ is a projective $\underline{B}$-Module on every affine open set (1.5.2), and consequently is locally free as a graded $\underline{B}$-Module, by 1.5.1. We can then, in the graded resolution constructed, replace $\underline{L}_{r+1}$ by $\underline{Z}_r$[288], and the following terms by 0, whence the asserted resolution.

Let us finally prove point c) of the proof.

Proposition 1.7. Let S be a quasi-compact scheme, $\underline{E}$ a $\underline{O}_S$-Module locally free of finite type, $\underline{B} = \mathbf{S}(\underline{E})$, and $\underline{L}$ a graded $\underline{B}$-Module, locally free of finite type (1.2). Then there exists a decreasing filtration of $\underline{L}$ by graded sub-$\underline{B}$-Modules $\underline{L}^{(m)}$, such that: [289]

i) $\underline{L}^{(m)} = \underline{L}$ for $m \leq m_0$;

ii) $\underline{L}^{(m)} = 0$ for $m \geq m_1$;

iii) for every m, $\mathrm{gr}^m(\underline{L}) = \underline{L}^{(m)}/\underline{L}^{(m+1)}$ is isomorphic as a graded $\underline{B}$-Module to a graded $\underline{B}$-Module of the form $\underline{N}_m \otimes_{\underline{O}_S} \underline{B}(m)$, where $\underline{N}_m$ is an $\underline{O}_S$-Module locally free of finite type.

Since S is quasi-compact, we can find integers n_0 and n_1 such that in a neighborhood of every point of S, the elements of a homogeneous basis [print 373]of $\underline{L}$ have degrees between n_0 and n_1. We then proceed by induction on $N = n_1 - n_0$.

When $N = 0$, i.e. $n_1 = n_0$, let n be the common value. The canonical homomorphism (1.2.1):

$$\underline{M}_n \otimes_{\underline{O}_S} \underline{B}(-n) \longrightarrow \underline{M}$$

is then an isomorphism. Indeed, the verification is local, and by (1.2.3), we can then suppose that $\underline{M} \simeq (\underline{B}(-n))^k$, and the result is trivial.

In the case $N > 0$, we again consider the homomorphism (1.2.1): [290]

$$\underline{M}_{n_0} \otimes_{\underline{O}_S} \underline{B}(-n_0) \longrightarrow \underline{M}.$$

We verify locally that this homomorphism is injective, and that its cokernel $\underline{M}'$ is locally free of finite type, with local generators of degree between $n_0 + 1$ and n_1: indeed, in a neighborhood of every point of S, $\underline{M}_{n_0}$ is either zero or the nonzero homogeneous component of $\underline{M}$ of lowest degree; using a decomposition of $\underline{M}$ into direct summands of the form (1.2.2), we see in the latter case, using (1.2.3), that $\underline{M}_{n_0} \otimes_{\underline{O}_S} \underline{B}(-n_0)$ is isomorphic to the direct summand $(\underline{B}(-n_0))^{k_{n_0}}$ of $\underline{M}$, whence the asserted properties. We then apply the induction hypothesis to $\underline{M}'$, whence the desired conclusion follows at once.

1.8. Returning to the proof of Theorem 1.1, let us show that $K^\bullet(X)$ is generated as a $K^\bullet(S)$-module by the powers of ξ.

If $\underline{F}$ is a $\underline{O}_X$-Module locally free of finite type, there exists, by 1.4, a graded $\underline{B}$-Module of finite presentation $\underline{M}$, flat over S, such that $\underline{F} = \underline{\mathrm{Proj}}_0(\underline{M})$. By 1.6, $\underline{M}$ admits a graded resolution by graded $\underline{B}$-Modules locally free of finite type. We deduce an exact sequence: [print 374]

$$0 \longrightarrow \underline{\mathrm{Proj}}_0(\underline{L}_{r+1}) \longrightarrow \ldots \longrightarrow \underline{\mathrm{Proj}}_0(\underline{L}_0) \longrightarrow \underline{\mathrm{Proj}}_0(\underline{M}) \longrightarrow 0,$$

and the following equality in $K^\bullet(X)$:

$$\mathrm{cl}(\underline{F}) = \sum_i (-1)^i \,\mathrm{cl}\big(\underline{\mathrm{Proj}}_0(\underline{L}_i)\big).$$

Using the filtration on the $\underline{L}_i$ defined in 1.7, we find:

$$\mathrm{cl}\big(\underline{\mathrm{Proj}}_0(\underline{L}_i)\big) = \sum_j \mathrm{cl}\big(\underline{\mathrm{Proj}}_0(\underline{N}_{ij} \otimes_{\underline{O}_S} \underline{B}(j))\big)$$

Now $\underline{\mathrm{Proj}}_0(\underline{N}_{ij} \otimes_{\underline{O}_S} \underline{B}(j)) = f^*(\underline{N}_{ij})(j)$; we therefore find:

$$\begin{aligned}\mathrm{cl}\big(\underline{\mathrm{Proj}}_0(\underline{L}_i)\big) &= \sum_j \mathrm{cl}\big(f^*(\underline{N}_{ij})\big).\xi^j \\ &= \sum_j f^*\big(\mathrm{cl}(\underline{N}_{ij})\big).\xi^j.\end{aligned}$$

We have thus finally expressed $\mathrm{cl}(\underline{F})$ as a linear combination, with coefficients in $K^\bullet(S)$, of the powers of ξ.

1.9. To see that $1, \xi, \ldots, \xi^{r+1}$ are linearly dependent over $K^\bullet(S)$ we consider the canonical surjective homomorphism: $f^*(\underline{E}) \to \underline{O}_X(1)$, whose kernel $\underline{F}$ is locally free of rank r. We can therefore write:

$$\lambda^{r+1}\big(f^*(E) - \xi\big) = 0, \qquad \text{with } E = \mathrm{cl}\,\underline{E},$$

the operations λ^i on $K^\bullet(X)$ being defined in V 2.2. We therefore obtain:

$$\sum_{i=0}^{r+1} \lambda^i(f^*(E))\lambda^{r+1-i}(-\xi) = 0.$$

Since $\underline{O}_X(1)$ is an invertible sheaf, $\lambda_t(\xi) = 1 + \xi t$, and consequently, $\lambda_t(-\xi) = 1/(1+\xi t)$[291], whence finally the relation:

$$\sum_{i=0}^{r+1} (-1)^{r+1-i} \xi^{r+1-i} \lambda^i(E) = 0. \tag{1.9.1}$$

Let us finally show that $1, \xi, \ldots, \xi^r$ are independent over $K^\bullet(S)$(*).

[print 375]**Lemma 1.10.** With the notation and hypotheses of 1.1, let $\underline{F}$ and $\underline{G}$ be two locally free sheaves of finite type on X, such that:

$$\mathrm{cl}_{K^\bullet(X)}(\underline{F}) = \mathrm{cl}_{K^\bullet(X)}(\underline{G});$$

then there exists an integer n_0, such that for $n \geq n_0$, we have:

i) $f_*(\underline{F}(n))$, $f_*(\underline{G}(n))$ are locally free of finite type on S;
ii) $\mathrm{cl}_{K^\bullet(S)}\big(f_*(\underline{F}(n))\big) = \mathrm{cl}_{K^\bullet(S)}\big(f_*(\underline{G}(n))\big)$.

Let A be the free abelian group generated by the classes (up to isomorphism) of locally free sheaves of finite type on X. To say that, in $K^\bullet(X)$, $\underline{F}$ and $\underline{G}$ have the same class is to say, by the construction of $K^\bullet$, that there exist exact sequences of $\underline{O}_X$-Modules locally free of finite type:

$$0 \longrightarrow \underline{L}_i \longrightarrow \underline{M}_i \longrightarrow \underline{N}_i \longrightarrow 0, \qquad i = 1, \ldots, k,$$

and integers n_i, such that in A we have the equality:

$$(\underline{F}) - (\underline{G}) = \sum_i n_i\big((\underline{L}_i) - (\underline{M}_i) + (\underline{N}_i)\big).$$

For every n, tensoring with $\underline{O}_X(n)$ defines an automorphism of A, whence: [292]

$$(\underline{F}(n)) - (\underline{G}(n)) = \sum_i n_i\big((\underline{L}_i(n)) - (\underline{M}_i(n)) + (\underline{N}_i(n))\big),$$

the sequences $0 \to \underline{L}_i(n) \to \underline{M}_i(n) \to \underline{N}_i(n) \to 0$ remaining exact.

Let B be the free abelian group generated by the classes (up to isomorphism) of quasi-coherent sheaves on S, and $A' \subset B$ the subgroup generated by the classes of locally free sheaves of finite type on S. Direct image defines a homomorphism $A \to B$, and we therefore still have: [print 376][293]

$$(f_*(\underline{F}(n))) - (f_*(\underline{G}(n))) = \sum_i n_i\big((f_*(\underline{L}_i(n))) - (f_*(\underline{M}_i(n))) + (f_*(\underline{N}_i(n)))\big).$$

(*) This proof is due to J.P. Jouanolou.

By 1.3, for sufficiently large[294] n, the following conditions will hold:

$$\forall k \geq 1,\ \forall i, \quad R^k f_*(\underline{L}_i(n)) = 0;$$

$$\forall i, \quad f_*(\underline{L}_i(n)),\ f_*(\underline{M}_i(n)),\ f_*(\underline{N}_i(n)),\ f_*(\underline{F}(n)),\ f_*(\underline{G}(n)) \text{ are locally free of finite type.}$$

It follows that for sufficiently large n, these elements all belong to A' and satisfy the relation written above, and that the sequences:

$$0 \longrightarrow f_*(\underline{L}_i(n)) \longrightarrow f_*(\underline{M}_i(n)) \longrightarrow f_*(\underline{N}_i(n)) \longrightarrow 0$$

are exact. This implies that in $K^\bullet(S)$, $f_*(\underline{F}(n))$ and $f_*(\underline{G}(n))$ have the same class.

1.11. Suppose given an equation of linear dependence over $K^\bullet(S)$ among $1, \xi, \ldots, \xi^r$:

$$f^*(x_0) + f^*(x_1)\xi + \ldots + f^*(x_r)\xi^r = 0. \tag{1.11.1}$$

We can write each of the x_j as the difference of the classes of two sheaves, namely: $x_j = \mathrm{cl}(\underline{E}_j) - \mathrm{cl}(\underline{E}'_j)$, where $\underline{E}_j$ and $\underline{E}'_j$ are locally free of finite type on S. Put

$$\underline{F} = \bigoplus_{j=0}^{r} f^*(\underline{E}_j)(j), \qquad \underline{F}' = \bigoplus_{j=0}^{r} f^*(\underline{E}'_j)(j).$$

Equality (1.11.1) then reads:

$$\mathrm{cl}_{K^\bullet(X)}(\underline{F}) = \mathrm{cl}_{K^\bullet(X)}(\underline{F}').$$

By Lemma 1.10, we obtain for sufficiently large n:

$$\mathrm{cl}_{K^\bullet(S)}\big(f_*(\underline{F}(n))\big) = \mathrm{cl}_{K^\bullet(S)}\big(f_*(\underline{F}'(n))\big).$$

[print 377]Now

$$\begin{aligned} f_*(\underline{F}(n)) &= \bigoplus f_*(f^*(\underline{E}_j) \otimes_{\underline{O}_X} \underline{O}_X(n+j)) \\ &= \bigoplus \underline{E}_j \otimes_{\underline{O}_S} f_*(\underline{O}_X(n+j)). \end{aligned}$$

Moreover, $f_*(\underline{O}_X(n+j)) = \mathbf{S}(\underline{E})^{n+j}$; we shall denote by σ_p the class in $K^\bullet(S)$ of $\mathbf{S}(\underline{E})^p$. Relation (1.11.1) therefore finally gives:

$$\forall n \geq n_0, \qquad \sum_{j=0}^{r} x_j \sigma_{n+j} = 0. \tag{1.11.2}$$

We then consider the complex:

$$0 \longrightarrow \bigwedge^{r+1}(\underline{E}) \otimes_{\underline{O}_S} \mathbf{S}(\underline{E}) \longrightarrow \bigwedge^{r}(\underline{E}) \otimes_{\underline{O}_S} \mathbf{S}(\underline{E}) \longrightarrow \ldots \longrightarrow \underline{E} \otimes_{\underline{O}_S} \mathbf{S}(\underline{E}) \longrightarrow \mathbf{S}(\underline{E}) \longrightarrow 0,$$

whose arrows are defined by:

$$\begin{aligned} &(x_1 \wedge \ldots \wedge x_n) \otimes (y_1 \otimes \ldots \otimes y_p) \rightsquigarrow \\ &\qquad \sum (-1)^{i-1} (x_1 \wedge \ldots \wedge \widehat{x_i} \wedge \ldots \wedge x_n) \otimes (x_i \otimes y_1 \otimes \ldots \otimes y_p) \end{aligned}$$

If we regard $\bigwedge^n(\underline{E})$ as a graded $\underline{O}_S$-Module concentrated in degree n, this is a complex of graded $\mathbf{S}(\underline{E})$-Modules, the morphisms themselves being graded.

This complex is a resolution of $\underline{O}_S$: on an open set where $\underline{E}$ is free, $\mathbf{S}(\underline{E})$ identifies with a polynomial Algebra $\underline{O}_S[T_0, \ldots, T_r]$, and the complex with the Koszul complex defined by the $\mathbf{S}(\underline{E})$-regular sequence $T_0, \ldots, T_r$; we know that the latter is a resolution of $\mathbf{S}(\underline{E})/(T_0, \ldots, T_r) = \underline{O}_S$.

Writing that each homogeneous component of the complex of degree ≥ 1 is acyclic, we therefore obtain relations in $K^\bullet(S)$: [295]

$$\sum_{i=0}^{r+1} (-1)^i \lambda^i(E) \sigma_{p-i} = 0 \qquad \text{for every } p \geq 1. \tag{1.11.3}$$

(putting $\sigma_k = 0$ for $k < 0$).

Suppose relation (1.11.2) holds for $n \geq n_0$; we shall show that it still holds for $n_0 - 1$, which will allow us to descend by induction to $n_0 = -r$ (we shall see later that this suffices to imply the vanishing of the x_j).

[print 378]By (1.11.3), we can write [296]

$$\sum_{j=0}^{r} x_j \left(\sum_{i=0}^{r+1} (-1)^i \lambda^i(E) \sigma_{p(j)-i} \right) = 0, \tag{1.11.4}$$

with $p(j) \geq 1$. We take $p(j) = n_0 + r + j$, which is indeed ≥ 1 if $n_0 > -r$. We then obtain:

$$\sum_{j=0}^{r} x_j \left(\sum_{i=0}^{r+1} (-1)^i \lambda^i(E) \sigma_{n_0+r+j-i} \right) = \sum_{i=0}^{r+1} (-1)^i \lambda^i(E) \left(\sum_{j=0}^{r} x_j \sigma_{n_0+r+j-i} \right).$$

For $i \neq r+1$, $n_0 + r - i \geq n_0$, so by (1.11.2):

$$\sum_{j=0}^{r} x_j \sigma_{n_0+r+j-i} = 0.$$

Consequently, relation (1.11.4) reduces to:

$$\lambda^{r+1}(E). \left(\sum_{j=0}^{r} x_j \sigma_{n_0-1+j} \right) = 0.$$

Now $\underline{E}$ has rank $r+1$, so $\lambda^{r+1}(E)$ is the class of an invertible sheaf, and is therefore invertible in $K^\bullet(S)$. Consequently, relation (1.11.4) reduces to relation (1.11.2) for $n_0 - 1$.

Thus relation (1.11.2) holds for $n \geq -r$; for $n = -r$, it gives:

$$x_r \sigma_0 = 0.$$

Since $\sigma_0 = 1$, we obtain $x_r = 0$. We thus show by descending induction on j that the x_j are all zero, using relations (1.11.2) for n ranging from $-r$ to 0. This shows that $1, \ldots, \xi^r$ are independent over $K^\bullet(S)$, and completes the proof of Theorem 1.1.

Remark 1.12. For every $\underline{O}_S$-Module locally free of finite type $\underline{E}$, let $\sigma_i(E)$ be the class in $K^\bullet(S)$ of the i-th symmetric power of $\underline{E}$; put $\sigma_t(E) = 1 + \sum_{i \geq 1} \sigma_i(E) t^i$; relations (1.11.3) can also be written: [print 379]

$$\lambda_t(E).\sigma_{-t}(E) = 1. \tag{1.12.1}$$

This relation itself shows the additivity of the map σ_t, which can therefore be extended to a homomorphism from $K^\bullet(S)$ to $1 + K^\bullet(S)[[t]]^+$:

Remark 1.13. We can take as a basis of $K^\bullet(X)$ over $K^\bullet(S)$ any sequence of powers of ξ; $\xi^k, \ldots, \xi^{k+r}$[297], since ξ is invertible, and such a sequence therefore consists of elements linearly independent over $K^\bullet(S)$; moreover, the other powers of ξ can be expressed explicitly as linear combinations of these with coefficients in $K^\bullet(S)$, by equality (1.9.1) and the invertibility of ξ.

Put $x = 1 - \xi^{-1}$ (x is interpreted as the "class of a hyperplane section"). We can then take as a basis of $K^\bullet(X)$ over $K^\bullet(S)$ the elements $1, x, \ldots, x^r$. To obtain the equation of dependence of $1, \ldots, x^{r+1}$ we consider the exact sequence:

$$0 \longrightarrow \underline{F} \longrightarrow f^*(\underline{E}) \longrightarrow \underline{O}_X(1) \longrightarrow 0,$$

and the exact sequence obtained by passing to the duals:

$$0 \longrightarrow \underline{O}_X(-1) \longrightarrow f^*(\check{\underline{E}}) \longrightarrow \check{\underline{F}} \longrightarrow 0.$$

We then observe that $\gamma^{r+1}(\check{F} - r) = 0$, from which we deduce:

$$\gamma^{r+1}(\check{F} - r) = \gamma^{r+1}(\check{E} - r - 1 + x) = 0,$$

that is:

$$\sum_{k=0}^{r+1} \gamma^{r+1-k}(\check{E} - r - 1).x^k = 0. \tag{1.13.1}$$

We obtain an analogous result by introducing $x' = \xi - 1$.

Finally, note that if $\underline{E} = (\underline{O}_S)^{r+1}$, and hence $X = \mathbf{P}_S^r$[298], we have $E = r + 1$, and equality (1.13.1) reduces to: [print 380]

$$x^{r+1} = 0. \tag{1.13.2}$$

Remark 1.14. Using the notion of a "relative scheme" (cf [3]), we can still give a meaning to 1.1 by assuming that S is a locally ringed topos. If we assume that the final object of the topos is quasi-compact, the proof given in the case of schemes carries over without difficulty, and the theorem therefore remains valid.

2. Computation of the $K^\bullet$ of a projective bundle: the case of perfect complexes

The present section will not be used in the rest of the seminar.

In this section, the notation $K^\bullet(X)$ will denote the Grothendieck group of perfect complexes on X (IV 2.2). We propose to prove the following theorem, analogous to 1.1:

Theorem 2.1. Let S be a quasi-compact scheme, $\underline{E}$ a locally free $\underline{O}_S$-Module of rank $r + 1$, $X = \mathbf{P}(\underline{E})$[299] the associated projective bundle; let ξ be the class in $K^\bullet(X)$ of the complex equal to $\underline{O}_X(1)$ in degree zero, and zero in the other degrees. Then $K^\bullet(X)$, endowed with its natural $K^\bullet(S)$-algebra structure, is a free $K^\bullet(S)$-module with basis $1, \xi, \ldots, \xi^r$.

2.2. Let us fix the notation: $\underline{B}$ will denote the symmetric algebra of $\underline{E}$; we shall write $\mathcal{A}$ for the category of $\underline{O}_X$-Modules, $\mathcal{C}$ for the category of graded $\underline{B}$-Modules with graded morphisms as morphisms, $\mathcal{C}_0$ for the full subcategory of $\mathcal{C}$ whose objects are the $\underline{B}$-Modules locally free of finite type as graded $\underline{B}$-Modules, and $\mathcal{C}'$ for the category of $\underline{O}_S$-Modules. The categories $\mathcal{A}$, $\mathcal{C}$, and $\mathcal{C}'$ are abelian.

For every n, and every complex $\underline{L}^\bullet$ in $D^+(\mathcal{A})$, we shall write $\underline{L}^\bullet(n)$ for the complex $\underline{L}^\bullet \otimes_{\underline{O}_X} \underline{O}_X(n)$.

[print 381]2.3. We shall need to apply the functor $\underline{\mathrm{Proj}}_0$ to arbitrary graded $\underline{B}$-Modules, and not only to quasi-coherent graded $\underline{B}$-Modules as in the construction of EGA II 3.2. We shall therefore indicate briefly how this construction can be carried out in the general case.

2.3.1. First let S be a scheme, $\underline{A}$ a quasi-coherent $\underline{O}_S$-algebra, $Y = \mathrm{Spec}(\underline{A})$, and $f : Y \to S$ the structural morphism. From the isomorphism:

$$\underline{A} \xrightarrow{\sim} f_*(\underline{O}_Y)$$

we deduce a canonical homomorphism:

$$f^{-1}(\underline{A}) \longrightarrow \underline{O}_Y \tag{2.3.1.1}$$

where f^{-1} denotes inverse image in the sense of sheaves of sets; in other words, we have a morphism of ringed spaces:

$$\bar{f} : (Y, \underline{O}_Y) \longrightarrow (S, \underline{A}) \tag{2.3.1.2}$$

From it we deduce an "inverse-image" functor $\bar{f}^*$ from the category of $\underline{A}$-Modules to the category of $\underline{O}_Y$-Modules, left adjoint to the "direct-image" functor $\bar{f}_*$ from the category of $\underline{O}_Y$-Modules to the category of $f_*(\underline{O}_Y)$-Modules. We therefore have:

$$\bar{f}^*(\underline{M}) = f^{-1}(\underline{M}) \otimes_{f^{-1}(\underline{A})} \underline{O}_Y \tag{2.3.1.3}$$

for every $\underline{A}$-Module $\underline{M}$.

Since the homomorphism (2.3.1.1) is flat (since on each stalk it corresponds to a localization), the functor $\bar{f}^*$ thus defined is an exact functor (cf [4], p. 164).

If $\underline{M}$ is a quasi-coherent $\underline{A}$-Module, this construction gives the Module $\widetilde{\underline{M}}$ defined in EGA II 1.4.3. Indeed, the isomorphism: [print 382]

$$\underline{M} \xrightarrow{\sim} f_*(\widetilde{\underline{M}}) = \bar{f}_*(\widetilde{\underline{M}})$$

gives, by adjunction, a homomorphism: [300]

$$\bar{f}^*(\underline{M}) \longrightarrow \widetilde{\underline{M}}$$

and we check immediately on each stalk that it is an isomorphism.

2.3.2. Now let $\underline{B}$ be a quasi-coherent graded $\underline{O}_S$-Algebra, $\underline{M}$ a graded $\underline{B}$-Module, and $X = \mathrm{Proj}(\underline{B})$. For every affine open U of S, and every section s of $\underline{B}$ over U, homogeneous of degree $d > 0$, we write $\underline{M}_{(s),U}$ for the homogeneous component of degree 0 of the Module of fractions $\underline{M}_s$; we therefore have:

$$\underline{M}_{(s),U} = \big((\underline{M}|U) \otimes_{(\underline{B}|U)} (\underline{B}|U)_s\big)_0.$$

If V is an affine open of U, then $\underline{M}_{(s),V}$ is the restriction to V of $\underline{M}_{(s),U}$.

We consider the basis of open sets of X formed by the affine opens:

$$D_+(U, s) \simeq \mathrm{Spec}\big((\underline{B}|U)_{(s)}\big).$$

We write f for the structural morphism $X \to S$, and $f_{U,s}$ for the affine morphism:


$$f_{U,s} : D_+(U, s) \longrightarrow U.$$

We then put:

$$\underline{\mathrm{Proj}}_0(\underline{M})_{U,s} = \bar{f}^*_{U,s}(\underline{M}_{(s),U}),$$

with the notation of 2.3.1. We shall show that the $\underline{\mathrm{Proj}}_0(\underline{M})_{U,s}$ glue as U and s vary, thereby defining an $\underline{O}_X$-Module denoted $\underline{\mathrm{Proj}}_0(\underline{M})$. The gluing for variable U and fixed s being evident,

we are reduced (cf. EGA II 2.5.2) to showing that, if t is a section of $\underline{B}$ over U, homogeneous of degree $e > 0$[301], then the restriction of $\underline{\mathrm{Proj}}_0(\underline{M})_{U,s}$ to $D_+(U, st)$ is canonically isomorphic to $\underline{\mathrm{Proj}}_0(\underline{M})_{U,st}$. We consider: [print 383]



we have to show that the Modules $i^* \circ \bar{f}^*_{U,s}(\underline{M}_{(s),U})$ and $\bar{f}^*_{U,st}(\underline{M}_{(st),U})$ are canonically isomorphic. On the one hand, we have the canonical isomorphisms: [302]

$$\begin{aligned} i^* \circ \bar{f}^*_{U,s}(\underline{M}_{(s),U}) &\simeq i^*\Big(f_{U,s}^{-1}(\underline{M}_{(s),U}) \otimes_{f_{U,s}^{-1}((\underline{B}|U)_{(s)})} \underline{O}_{D_+(s)}\Big) \\ &\simeq i^{-1}\Big(f_{U,s}^{-1}(\underline{M}_{(s),U}) \otimes_{f_{U,s}^{-1}((\underline{B}|U)_{(s)})} \underline{O}_{D_+(s)}\Big) \otimes_{i^{-1}(\underline{O}_{D_+(s)})} \underline{O}_{D_+(st)} \\ &\simeq f_{U,st}^{-1}(\underline{M}_{(s),U}) \otimes_{f_{U,st}^{-1}((\underline{B}|U)_{(s)})} \underline{O}_{D_+(st)}. \end{aligned}$$

On the other hand:

$$\bar{f}^*_{U,st}(\underline{M}_{(st),U}) \simeq f_{U,st}^{-1}(\underline{M}_{(st),U}) \otimes_{f_{U,st}^{-1}((\underline{B}|U)_{(st)})} \underline{O}_{D_+(st)}.$$

Finally, it follows from EGA II 2.2.2, translated into sheaf language, that we have the canonical isomorphisms:

$$\underline{M}_{(st),U} \simeq (\underline{M}_{(s),U})_{t^d/s^e} \simeq \underline{M}_{(s),U} \otimes_{(\underline{B}|U)_{(s)}} (\underline{B}|U)_{(st)},$$

whence, finally, the desired isomorphism.

It is clear that, if $\underline{M}$ is quasi-coherent, the Module $\underline{\mathrm{Proj}}_0(\underline{M})$ thus constructed is indeed the one defined in EGA II.

2.3.3. It follows from 2.3.1 and 2.3.2 that the functor $\underline{\mathrm{Proj}}_0$ is exact. Returning to the notation of 2.2, by passage to the derived categories we therefore deduce a functor still denoted $\underline{\mathrm{Proj}}_0$:

$$\underline{\mathrm{Proj}}_0 : D^+(\mathcal{C}) \longrightarrow D^+(\mathcal{A}).$$

We also see from the preceding constructions that the functor [print 384]$\underline{\mathrm{Proj}}_0$ on the category of graded $\underline{B}$-Modules is compatible with tensor product; the same remains true of the functor obtained on the derived categories.

2.4. For every integer k, and every $\underline{O}_X$-Module $\underline{L}$, put: [303]

$$F_k(\underline{L}) = \bigoplus_{n \geq k} f_*(\underline{L}(n)). \tag{2.4.1}$$

The functor $F_k : \mathcal{A} \to \mathcal{C}$ thus defined is left exact. Since the category of $\underline{O}_X$-Modules has enough injectives, we can define a derived functor: [304]

$$\mathbb{R}F_k : D^+(\mathcal{A}) \longrightarrow D^+(\mathcal{C}). \tag{2.4.2}$$

If $\underline{L}$ is a locally free $\underline{O}_X$-Module of finite type, there exists an integer k_0 such that, for $k \geq k_0$, $\underline{L}$ is F_k-acyclic. To see this, it suffices to show that, for n sufficiently large, $\underline{L}$ is acyclic for the functors $\underline{L} \rightsquigarrow f_*(\underline{L}(n))$[305]. Since tensoring with $\underline{O}_X(n)$ is an exact functor, this follows immediately from the spectral sequence of composite functors and from 1.3.

2.5. We propose to construct, for every $\underline{O}_X$-Module $\underline{F}$, a homomorphism functorial in $\underline{F}$:

$$\beta : \underline{\mathrm{Proj}}_0(F_k(\underline{F})) \longrightarrow \underline{F}. \tag{2.5.1}$$

For this, we define the restriction of β to the $D_+(U,s)$ (2.3.2). By construction we have, on putting $\underline{M} = F_k(\underline{F})$:

$$\underline{\mathrm{Proj}}_0(\underline{M})|D_+(U,s) = \bar{f}^*_{U,s}(\underline{M}_{(s),U}).$$

To define a homomorphism:

$$\bar{f}^*_{U,s}(\underline{M}_{(s),U}) \longrightarrow \underline{F}|D_+(U,s),$$

[print 385]it suffices, by the definition of $\bar{f}^*_{U,s}$ as left adjoint to the direct-image functor, to define an $f_{U,s*}(\underline{O}_X)$-linear homomorphism

$$\underline{M}_{(s),U} \longrightarrow f_{U,s*}(\underline{F}|D_+(U,s)).$$

For this, we must give, for every open V of U, a homomorphism:

$$\Gamma(V, \underline{M})_{(s)} \longrightarrow \Gamma(D_+(V,s), \underline{F}) \tag{2.5.2}$$

compatible with restrictions to the opens of U, and take the corresponding homomorphism on the associated sheaves. Homomorphism (2.5.2) is defined as follows: let $z/s^n \in \Gamma(V, \underline{M})_{(s)}$, where, s being of degree d, $z \in \Gamma(V, \underline{M}_{nd}) = \Gamma(D_+(V,s), \underline{F}(nd))$; we then associate to z/s^n the section $z.\alpha_{nd}(s^n)^{-1}$[306] of $\underline{F}$ over $D_+(V,s)$, writing α_d for the canonical homomorphism $\Gamma(V, \underline{B}_d) \longrightarrow \Gamma(f^{-1}(V), \underline{O}_X(d))$ (cf. EGA II 2.6.3 and 2.6.4). We easily check the necessary compatibilities, whence, by gluing, homomorphism (2.5.1).

It is clear that, in the case where $\underline{F}$ is quasi-coherent, the homomorphism thus constructed is the one in EGA II (3.3.4.1).

The homomorphism of functors $\underline{\mathrm{Proj}}_0 \circ F_k \to \mathrm{Id}$ on the category of $\underline{O}_X$-Modules gives, by passage to the derived categories, a homomorphism of functors on $D^+(\mathcal{A})$: [307]

$$\underline{\mathrm{Proj}}_0 \circ \mathbb{R}F_k \longrightarrow \mathrm{Id}_{D^+(\mathcal{A})}\,. \tag{2.5.3}$$

Proposition 2.6. Let S be a quasi-compact scheme, $\underline{E}$ a locally free $\underline{O}_S$-Module of finite type, $X = \mathbf{P}(\underline{E})$[308], and $\underline{L}^\bullet$[309] a perfect complex on X. Then there exists an integer k_0 such that, for $k \geq k_0$, the following properties hold: [print 386][310]

i) $\mathbb{R}F_k(\underline{L}^\bullet)$ is $\mathcal{C}_0$-perfect;

ii) The canonical morphism (2.5.3) : $\underline{\mathrm{Proj}}_0 \circ \mathbb{R}F_k(\underline{L}^\bullet) \longrightarrow \underline{L}^\bullet$ is an isomorphism in $D^+(\mathcal{A})$.

The notion of a $\mathcal{C}_0$-perfect complex considered here is local, that is, it is perfection relative to the category fibered over S defined by $\mathcal{C}_0$. This notion of perfection does have a meaning, since $\mathcal{C}_0$ satisfies the axioms of (I 4.0): only stability under kernels of epimorphisms is not immediate; it follows from 1.5.1.

Since S is quasi-compact, it suffices to check the two assertions of 2.6 locally on S, so that we can assume S affine. The invertible sheaf $\underline{O}_X(1)$ is then ample, and global resolutions on X are available (II 2.2.9). There therefore exists a strictly perfect complex $\underline{L}'^\bullet$ in $D_{\mathrm{parf}}(\mathcal{A})$, and an isomorphism in $D_{\mathrm{parf}}(\mathcal{A})$: $\underline{L}^\bullet \xleftarrow{\sim} \underline{L}'^\bullet$, which allows us to assume $\underline{L}^\bullet$ strictly perfect. We saw in 2.4 that, for k sufficiently large, the $\underline{L}^i$, all but finitely many of which are zero, are F_k-acyclic; we consequently have: [311]

$$\mathbb{R}F_k(\underline{L}^\bullet) = F_k(\underline{L}^\bullet)$$

for k sufficiently large. By 1.3 and 1.6, we can find k_0 such that, for $k \geq k_0$, the preceding condition is satisfied, and the $F_k(\underline{L}^i)$ are $\mathcal{C}_0$-perfect $\underline{B}$-Modules for every i. The complex $F_k(\underline{L}^\bullet)$ is then $\mathcal{C}_0$-perfect, whence i).

For k chosen as above, homomorphism (2.5.3) is then the homomorphism: [print 387][312]

$$\underline{\mathrm{Proj}}_0(F_k(\underline{L}^\bullet)) \longrightarrow \underline{L}^\bullet,$$

and it is an isomorphism on each component, since $\underline{L}^\bullet$ was assumed strictly perfect, whence ii).

2.7. On $\mathcal{C}$, the functor which associates to a graded $\underline{B}$-Module its n-th component is an exact functor with values in $\mathcal{C}'$; it therefore defines a functor:

$$\mathrm{Comp}_n : D^+(\mathcal{C}) \longrightarrow D^+(\mathcal{C}'). \tag{2.7.1}$$

For every n, we can consider the functor $\mathcal{C}' \to \mathcal{C}$ defined by tensoring with $\underline{B}(-n)$. Homomorphism (1.2.1) defines a morphism of functors on $\mathcal{C}$, whence a morphism for the corresponding functors between derived categories:

$$\mathrm{Comp}_n \otimes_{\underline{O}_S} \underline{B}(-n) \longrightarrow \mathrm{Id}_{D^+(\mathcal{C})}\,. \tag{2.7.2}$$

Let $\underline{M}^\bullet$ be a $\mathcal{C}_0$-perfect complex. We shall show that there exists a sequence of distinguished triangles:

$$\begin{array}{ccccc} & & \underline{P}_1^\bullet & & \\ & \swarrow & & \nwarrow & \\ \underline{N}_1^\bullet & & \longrightarrow & & \underline{M}^\bullet, \end{array} \qquad \begin{array}{ccccccccc} & & \underline{P}_2^\bullet & & & & \underline{P}_k^\bullet & & \\ & \swarrow & & \nwarrow & & \swarrow & & \nwarrow & \\ \underline{N}_2^\bullet & & \longrightarrow & & \underline{P}_1^\bullet, & \ldots, & \underline{N}_k^\bullet & \longrightarrow & \underline{P}_{k-1}^\bullet, \end{array}$$

where $\underline{P}_k^\bullet$ is acyclic, and the $\underline{N}_i^\bullet$ are of the form $\underline{K}_i^\bullet \otimes_{\underline{O}_S} \underline{B}(d)$, the $\underline{K}_i^\bullet$ being perfect complexes in the ordinary sense on S.

Since $\underline{M}^\bullet$ is $\mathcal{C}_0$-perfect, and S is quasi-compact, there exists a finite covering of S, and an integer d_0 such that, on every open of this covering, $\underline{M}^\bullet$ is isomorphic to a bounded complex whose terms are locally free of finite type over $\underline{B}$, say ${\underline{M}'_U}^\bullet$, such that $\mathrm{Comp}_d({\underline{M}'_U}^\bullet) = 0$ for $d < d_0$.

[print 388]We then consider the canonical morphism (2.7.2):

$$\underline{N}_1^\bullet = \mathrm{Comp}_{d_0}(\underline{M}^\bullet) \otimes_{\underline{O}_S} \underline{B}(-d_0) \longrightarrow \underline{M}^\bullet;$$

let $\underline{P}_1^\bullet$ be the cone of this morphism. It is again a $\mathcal{C}_0$-perfect complex: indeed, for every n, $\mathrm{Comp}_n(\underline{M}^\bullet)$ is a perfect complex on S, since this holds when $\underline{M}^\bullet$ is bounded with terms locally free of finite type over $\underline{B}$; consequently, the complex $\mathrm{Comp}_n(\underline{M}^\bullet) \otimes_{\underline{O}_S} \underline{B}(-d_0)$ is $\mathcal{C}_0$-perfect, and the same therefore holds for the cone (I 4.10).

Moreover, on every open U of the covering we can write the commutative diagram:

$$\begin{array}{ccc} \mathrm{Comp}_{d_0}(\underline{M}^\bullet) \otimes_{\underline{O}_S} \underline{B}(-d_0) & \longrightarrow & \underline{M}^\bullet \\ \big\downarrow \wr & & \big\downarrow \wr \\ \mathrm{Comp}_{d_0}({\underline{M}'_U}^\bullet) \otimes_{\underline{O}_S} \underline{B}(-d_0) & \longrightarrow & {\underline{M}'_U}^\bullet, \end{array}$$

which induces an isomorphism between the cones; the lower homomorphism is an isomorphism in degree d_0; therefore $\underline{P}_1^\bullet$ is isomorphic over U to a bounded complex with terms locally free of finite type over $\underline{B}$, whose components Comp_d are zero for $d < d_0 + 1$. Consequently, $\underline{P}_1^\bullet$ has the same properties as $\underline{M}^\bullet$, with d_0 replaced by $d_0 + 1$. We can therefore iterate the process, and it suffices to see that, after finitely many operations, the resulting complex $\underline{P}_k^\bullet$ is acyclic. Here again the question is local, so it suffices to check this for ${\underline{M}'_U}^\bullet$. For each term ${\underline{M}'_U}^i$, the

morphisms considered above are those defining the filtration of Proposition 1.7; this filtration is finite, so, starting from $\underline{M'_U}^\bullet$, we reach the zero complex after finitely many operations, that is, the corresponding complex $\underline{P}_k^\bullet$ is acyclic.

[print 389]2.8. Let $\underline{L}^\bullet$ be a perfect complex on X. There exists an integer k such that $\underline{M}^\bullet = \mathbb{R}F_k(\underline{L}^\bullet)$[313] is a $\mathcal{C}_0$-perfect complex, and $\underline{L}^\bullet$ is isomorphic to $\underline{\mathrm{Proj}}_0(\mathbb{R}F_k(\underline{L}^\bullet))$[314]. Applying 2.7 to the complex $\underline{M}^\bullet$, the sequence of distinguished triangles constructed gives, on applying the functor $\underline{\mathrm{Proj}}_0$, a sequence of distinguished triangles in $D_{\mathrm{parf}}(\mathcal{A})$ (it is clear that $\underline{\mathrm{Proj}}_0$ transforms a $\mathcal{C}_0$-perfect complex into a perfect complex). Moreover, we have:

$$\underline{\mathrm{Proj}}_0(\underline{K}^\bullet \otimes_{\underline{O}_S} \underline{B}(d)) = f^*(\underline{K}^\bullet)(d)$$

(note that $f^* = \mathbb{L}f^*$[315] by the flatness of f) for $\underline{K}^\bullet$ in $D^+(\mathcal{C}')$.

We therefore obtain in $K^\bullet(X)$ the equality:

$$\mathrm{cl}(\underline{L}^\bullet) = \sum_i \mathrm{cl}(f^*(\underline{K}_i^\bullet)).\xi^{d_i} = \sum_i f^*(\mathrm{cl}(\underline{K}_i^\bullet)).\xi^{d_i},$$

where the $\underline{K}_i^\bullet$ are the perfect complexes[316] on S defined in 2.7. It follows that $K^\bullet(X)$ is generated as a $K^\bullet(S)$-module by the powers of ξ.

To see that $1, \xi, \ldots, \xi^{r+1}$ are linearly dependent, we consider the homomorphism: $K^\bullet_{\mathrm{naïf}}(X) \to K^\bullet_{\mathrm{parf}}(X)$ which associates to the class of a locally free sheaf of finite type $\underline{L}$ the class of the complex equal to $\underline{L}$ in degree 0, and zero in the other degrees. Equation (1.9.1) then gives a relation of dependence in $K^\bullet_{\mathrm{parf}}(X)$.

To see that $1, \xi, \ldots, \xi^r$ are linearly independent, we can use the $K^\bullet(S)$-linear morphism $f_* : K^\bullet(X) \to K^\bullet(S)$ (IV 2.11.1). We have:

$$f_*(1) = 1, \quad \text{since } f_*(\underline{O}_X) = \underline{O}_S, \text{ and } R^i(f_*(\underline{O}_X)) = 0 \text{ for } i \geq 1;$$

$$f_*(\xi^n) = 0 \quad \text{for } -r \leq n < 0, \quad \text{since } R^i\big(f_*(\underline{O}_X(n))\big) = 0 \text{ for } i \geq 0.$$

Given a relation of the form:

$$f^*(x_0) + f^*(x_1)\xi + \ldots + f^*(x_r)\xi^r = 0,$$

we multiply it by ξ^{-r}, and apply f_*; we then find $x_r = 0$; we [print 390] complete the induction, noting that, if $0 \leq s \leq r$, and if $x_j = 0$ for $s < j \leq r$, we have:

$$\sum_{i=0}^{s} f^*(x_i)\xi^i = 0,$$

whence:

$$\sum_{i=0}^{s} f^*(x_i)\xi^{i+r-s} = 0,$$

whence $x_s = 0$ by what precedes.

3. Consequences of the structure theorem for the $K^\bullet$ of a projective bundle

Proposition 3.1. Let S be a quasi-compact scheme, $\underline{E}_1, \ldots, \underline{E}_n$ locally free $\underline{O}_S$-Modules of finite type, $\mathbf{D}(\underline{E}_i)$[317] the flag scheme of $\underline{E}_i$,[318] $X = \mathbf{D}(\underline{E}_1) \times_S \ldots \times_S \mathbf{D}(\underline{E}_n)$[319]. Then the canonical homomorphism:

$$K^\bullet(S) \longrightarrow K^\bullet(X)$$

is injective ($K^\bullet$ being defined, either from locally free $\underline{O}_S$-Modules of finite type, or from perfect complexes).

We check immediately, using the universal property of flag schemes, that X can be constructed as follows: we consider the sequence of S-schemes $X_1, \ldots, X_n$, defined inductively by taking X_k to be the flag scheme of the inverse image of $\underline{E}_k$ over X_{k-1}; then $X = X_n$. We then see at once that, to prove the proposition, it suffices to prove it in the case $n = 1$, the general case following by induction. We can therefore assume that X is the flag scheme of a locally free $\underline{O}_S$-Module of finite type $\underline{E}$.

We know that X can then be constructed as follows: we put $\underline{E} = \underline{F}_0$, $S = X_0$, and define inductively schemes $X_1, \ldots, X_n$ and [print 391]for every i a locally free $\underline{O}_{X_i}$-Module of finite type $\underline{F}_i$ by putting: [320]

$$X_k = \mathbf{P}(\underline{F}_{k-1}); \qquad \underline{F}_k = \mathrm{Ker}(f_k^*(\underline{F}_{k-1}) \longrightarrow \underline{O}_{X_k}(1)),$$

where f_k is the structural morphism $X_k \to X_{k-1}$.

We then see that it suffices to prove the proposition in the case where X is a projective bundle over S, the case of a flag bundle following by induction. This case follows immediately from 1.1 (resp. 2.1).

Theorem 3.2. Let S be a quasi-compact scheme. The pre-λ-ring $K^\bullet(S)$, defined from locally free $\underline{O}_S$-Modules of finite type, is a λ-ring (V 2.4).

The homomorphism $\lambda_t : K^\bullet(S) \longrightarrow 1 + \widehat{K^\bullet(S)[[t]]}^+$ being additive, it suffices, to check that it is a λ-homomorphism, to make the verification for elements of the form $\mathrm{cl}(\underline{E})$, where $\underline{E}$ is a locally free $\underline{O}_S$-Module of finite type. We are therefore reduced to showing that, for every pair of elements E, F of $K^\bullet(S)$, classes of locally free $\underline{O}_S$-Modules of finite type $\underline{E}, \underline{F}$, relations V 2.4.2 hold:

$$\lambda^i(EF) = P_i(\lambda^1(E), \ldots, \lambda^i(E); \lambda^1(F), \ldots, \lambda^i(F));$$
$$\lambda^j(\lambda^i(E)) = Q_{i,j}(\lambda^1(E), \ldots, \lambda^{ij}(E)).$$

Since S is a quasi-compact scheme, we can find a finite covering of S by disjoint opens such that, on each, $\underline{E}$ and $\underline{F}$ have constant rank. The $K^\bullet$ of a finite direct sum of opens is the product of the $K^\bullet$ of these opens. It therefore suffices to check relations V 2.4.2 in each of the $K^\bullet$ of the opens of the covering. We can therefore assume $\underline{E}$ and $\underline{F}$ to have constant rank on S.

[print 392]Now let $X = \mathbf{D}(\underline{E}) \times_S \mathbf{D}(\underline{F})$[321]. The λ-homomorphism $K^\bullet(S) \to K^\bullet(X)$ is injective by 3.1. Consequently, to prove relations V 2.4.2, we can work in $K^\bullet(X)$. Since in $K^\bullet(X)$ the elements E and F decompose into sums of classes of invertible sheaves, it suffices, by additivity, to prove these same relations for elements ξ, η, classes of invertible sheaves; they can also be written:

$$1 + \xi\eta t = (1 + \xi t) \circ (1 + \eta t); \qquad \lambda_u(1 + \xi t) = 1 + \{1 + \xi t\}u,$$

by putting them in the form V 2.4.1. They are then precisely relations V 2.3.1 and 2.3.3 defining the λ-ring structure of $1 + \widehat{K^\bullet(S)[[t]]}^+$.

Let us state the following more general result:

Theorem 3.3.[322] Let $(T, \underline{A})$ be a commutatively ringed topos; write K for the Grothendieck group of $\underline{A}$-Modules locally free[323] of bounded rank on T. Then, K, endowed with its canonical pre-λ-structure (V 2.2 b)), is a λ-ring.

To show that K is a λ-ring, we must prove relations V 2.4.1, and, by additivity, it suffices to prove them for elements E, F which are classes of locally free $\underline{A}$-Modules of bounded rank. Since

the ranks of $\underline{E}$ and $\underline{F}$ are bounded, we can decompose T into a finite "disjoint sum" of opens over which $\underline{E}$ and $\underline{F}$ have constant rank. Using the commutation of $K^\bullet$ with finite direct sums, and putting relations V 2.4.1 in the form V 2.4.2, we see that it suffices to prove relations V 2.4.2 in each of the $K^\bullet$ of the opens considered, that is, it suffices to prove relations V 2.4.2 for elements E, F, classes of $\underline{A}$-Modules $\underline{E}, \underline{F}$ locally free of constant rank.

[print 393]Let $\underline{E}$ be a locally free $\underline{A}$-Module of finite rank n. There exists a torsor $\underline{P}$ under $\underline{\mathrm{Gl}}(n, \underline{A})$ such that $\underline{E} = \underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A})} (\underline{A})^n$, where $(\underline{A})^n$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A})$-Module by the standard representation. We can then define a λ-homomorphism from $R_{\mathbb{Z}}(\mathbf{Gl}(n))$ to K by associating to every representation ρ of $\mathbf{Gl}(n)$ in a free $\mathbb{Z}$-module of rank p the class of the locally free $\underline{A}$-Module $\underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A})} (\underline{A})^p$, where $(\underline{A})^p$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A})$-module by the representation ρ. Under this homomorphism, the class of $\underline{E}$ is therefore the image of the standard representation. To check a polynomial relation among the $\lambda^i(E)$, it therefore suffices that this relation hold for the standard representation in $R_{\mathbb{Z}}(\mathbf{Gl}(n))$.[324]

We can argue similarly with two locally free $\underline{A}$-Modules $\underline{E}, \underline{F}$ of ranks n, p; we can then find a torsor $\underline{P}$ under $\underline{\mathrm{Gl}}(n, \underline{A}) \times \underline{\mathrm{Gl}}(p, \underline{A})$ such that $\underline{E} = \underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A}) \times \underline{\mathrm{Gl}}(p,\underline{A})} (\underline{A})^n$, where $(\underline{A})^n$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A}) \times \underline{\mathrm{Gl}}(p, \underline{A})$-Module by letting $\underline{\mathrm{Gl}}(n, \underline{A})$ act by the standard representation and $\underline{\mathrm{Gl}}(p, \underline{A})$ by the trivial representation, and conversely for $\underline{F}$. We then define a λ-homomorphism from $R_{\mathbb{Z}}(\mathbf{Gl}(n) \times \mathbf{Gl}(p))$ to K in the same way as above, and are thus reduced to checking the relations among the $\lambda^i(E)$, $\lambda^j(F)$ for the corresponding elements of $R_{\mathbb{Z}}(\mathbf{Gl}(n) \times \mathbf{Gl}(p))$.[325]

We are therefore reduced to showing that, for every group scheme G over $\operatorname{Spec} \mathbb{Z}$ which is a finite product of linear group schemes, the pre-λ-ring $R_{\mathbb{Z}}(G)$ is a λ-ring. For this, we can refer to [5], where $R_{\mathbb{Z}}(G)$ is computed explicitly. To avoid using this result, we can reason in a manner analogous to 3.2, using 1.14.

[print 394]

4. Computation of the $K^\bullet$ of a flag bundle

4.1. Let S be a scheme, and let $\underline{E}$ be an $\underline{O}_S$-Module locally free of rank n. Let $\pi = (p_1, \ldots, p_k)$ be a sequence of integers ≥ 1 such that:

$$\sum_{i=1}^{k} p_i = n.$$

Recall that the flag functor of type π of $\underline{E}$ is the functor on the category of S-schemes, with values in the category of sets, which associates to an S-scheme $X \xrightarrow{f} S$ the set of sequences of quotients of $f^*(\underline{E})$:

$$(\underline{L}_{k-1}, \ldots, \underline{L}_1),$$

where, for every i, $\underline{L}_i$ is a quotient of $\underline{L}_{i+1}$, locally free of rank $p_1 + \ldots + p_i$.

Such a functor is representable by an S-scheme $\mathbf{D}_\pi(\underline{E})$[326], endowed with a canonical sequence of quotients of the inverse image of $\underline{E}$, locally free of ranks $p_1, p_1 + p_2, \ldots$

If $\pi = (1, n-1)$, then $\mathbf{D}_\pi(\underline{E}) = \mathbf{P}(\underline{E})$[327].

If $\pi = (r, n-r)$, then $\mathbf{D}_\pi(\underline{E}) = \mathbf{G}_{r,n}(\underline{E})$[328], the Grassmannian of type $(r, n-r)$ of $\underline{E}$.

If all the p_i are equal to 1, one obtains the usual flag bundle of $\underline{E}$, denoted by $\mathbf{D}(\underline{E})$[329].

Lemma 4.2. Let S be a scheme, $\underline{E}$ an $\underline{O}_S$-Module locally free of rank n, $\pi = (p_1, \ldots, p_i, \ldots, p_k)$, with $p_1 + \ldots + p_k = n$; let p', p'' be two integers ≥ 1 such that $p' + p'' = p_i$; put $\pi' = (p_1, \ldots, p_{i-1}, p', p'', p_{i+1}, \ldots, p_k)$, $\pi'' = (p', p'')$. Let $\underline{L}_1, \ldots, \underline{L}_{k-1}$ be the canonical quotients of

the inverse image of $\underline{E}$ on $\mathbf{D}_\pi(\underline{E})$[330], and let $\underline{F}$ be the kernel of $\underline{L}_i \to \underline{L}_{i-1}$. Then there is a canonical S-isomorphism: [print 395][331]

$$\mathbf{D}_{\pi'}(\underline{E}) \simeq \mathbf{D}_{\pi''}(\underline{F}).$$

The proof, immediate from the functorial definition of flag schemes, is left to the reader.

Lemma 4.3. Let A be a commutative ring with unit, n an integer > 0, $(c_i)_{1\leq i\leq n}$ a family of elements of A; let C be the A-algebra generated by generators ξ_i, $i = 1, \ldots, n$, subject to the relations:

$$\sigma_i(\xi_1, \ldots, \xi_n) = c_i, \qquad i = 1, \ldots, n,$$

where the σ_i denote the elementary symmetric functions. Let $p_1, \ldots, p_k$ be integers ≥ 1 with sum n, and for $1 \leq j \leq k$,

$$\xi^{(j)} = (\xi_1^{(j)}, \ldots, \xi_{p_j}^{(j)}),$$

with

$$\xi_i^{(j)} = \xi_{\sum_{\alpha<j} p_\alpha + i}.$$

Put:

$$c_i^{(j)} = \sigma_i(\xi_1^{(j)}, \ldots, \xi_{p_j}^{(j)}), \qquad 1 \leq j \leq k, \quad 1 \leq i \leq p_j,$$

and let B be the subalgebra of C generated by the $c_i^{(j)}$. Then:

i) the elements of the form:

$$(\xi^{(1)})^{\alpha(1)} \ldots (\xi^{(k)})^{\alpha(k)},$$

where $\alpha(j) = (\alpha_1^{(j)}, \ldots, \alpha_{p_j}^{(j)})$, and $0 \leq \alpha_i^{(j)} \leq p_j - i$, form a basis of C considered as a B-module.

ii) the generators $c_i^{(j)}$ of the A-algebra B satisfy the n isobaric relations deduced from the relation:[332] [333]

$$1 + \sum_{i=1}^{n} c_i = \prod_{j=1}^{k} \left(1 + \sum_{i=1}^{p_j} c_i^{(j)}\right)$$

[print 396](c_i and $c_i^{(j)}$ being assigned weight[334] i), and these n relations generate the ideal of relations among the $c_i^{(j)}$.

Corollary 4.4. The elements of C of the form:

$$\xi_1^{\alpha_1} \ldots \xi_{n-1}^{\alpha_{n-1}}, \qquad 0 \leq \alpha_i \leq n - i,$$

form a basis of C considered as a module over A.

For the proof of the lemma and the corollary, we refer to [2], Exposé 4, pages 4.19, from which they are borrowed.

4.5. Let S be a quasi-compact scheme, $\underline{E}$ an $\underline{O}_S$-Module locally free of rank n, and $\pi = (p_1, \ldots, p_k)$ a sequence of integers ≥ 1 with sum n. Put $X = \mathbf{D}_\pi(\underline{E})$[335]; X is endowed with a canonical sequence of quotients $\underline{L}_i$; put $\underline{L}_0 = 0$, $\underline{L}_k = f^*(\underline{E})$, where f is the structural morphism $X \to S$. For $1 \leq i \leq k$, denote by $\underline{F}_i$ the kernel of $\underline{L}_i \to \underline{L}_{i-1}$. Let $K^\bullet(X)$ be the Grothendieck group of locally free sheaves of finite type on X (resp. of perfect complexes). Denote by F_j the class of $\underline{F}_j$, and by $\lambda_i^{(j)}$ the class of $\wedge^i(\underline{F}_j)$. In the case where $K^\bullet(X)$ is defined in terms of perfect

complexes, the series $\lambda_t(E)$, $\lambda_t(F_j)$ still have an evident meaning, where E is the class of $\underline{E}$. Finally, note that:

$$\sum_{j=1}^{k} F_j = E.$$

Theorem 4.6. With the hypotheses and notation of 4.5, the $K^\bullet(S)$-algebra $K^\bullet(X)$ is generated by the $\lambda_i^{(j)}$; the ideal of relations among the $\lambda_i^{(j)}$ is generated by the relations deduced from:

$$\lambda_t(E) = \prod_{j=1}^{k} \lambda_t(F_j).$$

[print 397]In other words, $K^\bullet(X)$ is the $K^\bullet(S)$-λ-algebra generated by the generators $F_1, \ldots, F_k$ subject to the relations (cf. V 4.6):

i) $\lambda^p(F_i) = 0$ for $p > p_i$;

ii) $\sum_i F_i = E$.

We begin with the case where all the p_i are equal to 1; the theorem then reads:

Corollary 4.7. Let S be a quasi-compact scheme, $\underline{E}$ an $\underline{O}_S$-Module locally free of rank[336] n, X the flag bundle of $\underline{E}$, and $\xi_1, \ldots, \xi_n$ the classes of the invertible sheaves $\underline{F}_1, \ldots, \underline{F}_n$[337] (notation of 4.5)[338]. Then $K^\bullet(X)$ is the $K^\bullet(S)$-algebra generated by the ξ_i, subject to the relations:

$$\sigma_i(\xi_1, \ldots, \xi_n) = \lambda^i(E).$$

We proceed by induction on n. For $n = 1$, the theorem is trivial. Suppose it has been proved for $n-1$; we then consider $\mathbf{P}(\underline{E})$[339], and on $\mathbf{P}(\underline{E})$[340] the exact sequence: [341]

$$0 \longrightarrow \underline{F} \longrightarrow g^*(\underline{E}) \longrightarrow \underline{O}_{\mathbf{P}(\underline{E})}(1) \longrightarrow 0,$$

where g is the structural morphism of $\mathbf{P}(\underline{E})$[342]. By Lemma 4.2, X is identified with the usual flag bundle $\mathbf{D}(\underline{F})$[343] over $\mathbf{P}(\underline{E})$[344], the canonical sheaves corresponding under this identification. The diagram: [345]

$$\begin{array}{ccc} \mathbf{D}(\underline{E}) = \mathbf{D}(\underline{F}) & \xrightarrow{\ h\ } & \mathbf{P}(\underline{E}) \\ & \searrow^{f} & \downarrow{\scriptstyle g} \\ & & S, \end{array}$$

shows that $K^\bullet(S) \to K^\bullet(X)$ factors through $K^\bullet(\mathbf{P}(\underline{E}))$[346]. By the induction hypothesis applied to $\mathbf{D}(\underline{F})$[347], and Corollary 4.4[348], $K^\bullet(X)$ has as a basis over $K^\bullet(\mathbf{P}(\underline{E}))$[349] the elements of the form:

$$\xi_2^{\alpha_2} \ldots \xi_{n-1}^{\alpha_{n-1}}, \qquad \text{with } 0 \le \alpha_i \le n - i;$$

moreover, $K^\bullet(\mathbf{P}(\underline{E}))$[350] has as a basis over $K^\bullet(S)$ the elements $1, \xi_1, \ldots, \xi_1^{n-1}$; thus $K^\bullet(X)$ has as a basis over $K^\bullet(S)$ the elements: [print 398]

$$\xi_1^{\alpha_1} \ldots \xi_{n-1}^{\alpha_{n-1}}, \qquad \text{with } 0 \le \alpha_i \le n - i.$$

Moreover, the additivity of λ_t implies that $\prod_i (1 + \xi_i t) = \lambda_t(E)$, and the relations of the corollary are satisfied.

If C is the $K^\bullet(S)$-algebra generated by generators ξ_i subject to the relations:

$$\sigma_i(\xi_1, \ldots, \xi_n) = \lambda^i(E),$$

there is therefore a surjective homomorphism from C onto $K^\bullet(X)$. By 4.4, this homomorphism sends a basis of C to a basis of $K^\bullet(X)$, so it is an isomorphism, which completes the proof of the corollary.

Let us treat the general case of a flag bundle of type π. The usual flag bundle $\mathbf{D}(\underline{E})$[351] can be obtained from $\mathbf{D}_\pi(\underline{E})$[352] by successively constructing the flag bundles of the $\underline{F}_j$ (4.2). It follows that $K^\bullet(\mathbf{D}(\underline{E}))$[353], considered as a module over $K^\bullet(X)$, has a basis formed by the elements considered in 4.3.i), the $\xi_i^{(j)}$ being the classes of the sheaves analogous to the $\underline{F}_j$ occurring in the splitting of the inverse images of the $\underline{F}_j$ on $\mathbf{D}(\underline{E})$[354] (notation of 4.5 and 4.7)[355]. Thus $K^\bullet(X)$ is a subring of $K^\bullet(\mathbf{D}(\underline{E}))$[356]; moreover, the additivity of λ_t implies that:

$$\lambda_i^{(j)} = \sigma_i(\xi_1^{(j)}, \ldots, \xi_{p_j}^{(j)}),$$

and $K^\bullet(X)$ contains the sub-$K^\bullet(S)$-algebra of $K^\bullet(\mathbf{D}(\underline{E}))$[357] generated by the $\lambda_i^{(j)}$. Let B be this subalgebra; by 4.3.i), and what has just been said, there is in $K^\bullet(\mathbf{D}(\underline{E}))$[358] a family of elements that forms a basis both for the B-module structure and for the $K^\bullet(X)$-module structure; consequently, $K^\bullet(X) = B$, which shows that $K^\bullet(X)$ is generated by the $\lambda_i^{(j)}$. The ideal of relations among the $\lambda_i^{(j)}$ is then given by 4.3.ii), taking [print 399]into account the structure of $K^\bullet(\mathbf{D}(\underline{E}))$[359] over $K^\bullet(S)$, determined by 4.7, and the fact that the $\xi_i^{(j)}$ are the classes of the sheaves occurring in the splitting of the inverse image of $\underline{E}$ on $\mathbf{D}(\underline{E})$[360]. This completes the proof of the first form of 4.6.

To deduce the second form, note that the elements F_i of the $K^\bullet(S)$-λ-algebra $K^\bullet(X)$ satisfy relations i) and ii). If A is the $K^\bullet(S)$-λ-algebra generated by elements F_i' satisfying i) and ii), there is therefore a $K^\bullet(S)$-λ-homomorphism from A to $K^\bullet(X)$ sending F_i' to F_i (V 4.7). By 4.6, this homomorphism is surjective. Since the λ-ideal generated by relations i) and ii) in $K^\bullet(S)[F_1, \ldots, F_k]$ contains the usual ideal generated by the relations of 4.6, this homomorphism is also injective, hence is an isomorphism.

Corollary 4.8. With the hypotheses and notation of 4.5, put:

$$c_i^{(j)} = \gamma^i(F_j - p_j), \qquad 1 \leq i \leq p_j;$$

then the $K^\bullet(S)$-algebra $K^\bullet(X)$ is generated by the $c_i^{(j)}$; the ideal of relations among the $c_i^{(j)}$ is generated by the relations deduced from:

$$\gamma_t(E - n) = \prod_{j=1}^{k} \gamma_t(F_j - p_j).$$

This is just another statement of 4.6. Indeed, the $c_i^{(j)}$ can be expressed in terms of the $\lambda_i^{(j)}$, and conversely. Moreover, the relation: $\lambda_t(E) = \prod \lambda_t(F_j)$ is equivalent to the relation: $\gamma_t(E - n) = \prod \gamma_t(F_j - p_j)$.

4.9. Let S be a scheme; denote by $\mathbf{G}_S^{n,r}$[361] the Grassmannian of type $(n, r-n)$ of $(\underline{O}_S)^r$, with $r \geq n$. As r varies, the $\mathbf{G}_S^{n,r}$ form an inductive system[362], the morphism $\mathbf{G}_S^{n,r} \longrightarrow \mathbf{G}_S^{n,r+1}$[363] being defined by considering $(\underline{O}_{\mathbf{G}_S^{n,r}})^r$[364] as the [print 400]quotient of $(\underline{O}_{\mathbf{G}_S^{n,r}})^{r+1}$[365] by the $(r+1)$-st factor. In this system of morphisms, the canonical quotients therefore correspond by inverse image. The inductive system thus defined will be denoted by $\mathbf{G}_S^{n,\infty}$[366].

Its $K^\bullet$ is defined by the formula: [367]

$$K^\bullet(\mathbf{G}_S^{n,\infty}) = \varprojlim_r (K^\bullet(\mathbf{G}_S^{n,r})). \tag{4.9.1}$$

Let $\underline{F}_1, \underline{F}_2$ be the canonical sheaves on $\mathbf{G}_S^{n,r}$[368] (4.5), and F_1, F_2 their class in $K^\bullet(\mathbf{G}_S^{n,r})$[369]. Put $c_i = \gamma^i(F_1 - n)$. As r varies, the c_i correspond under the transition morphisms, and therefore define elements, still denoted by c_i, of $K^\bullet(\mathbf{G}_S^{n,\infty})$[370].

Proposition 4.10. Let S be a quasi-compact scheme. Then, with the notation of 4.9, there is an isomorphism:[371] [372]

$$K^\bullet(\mathbf{G}_S^{n,\infty}) \xrightarrow{\sim} K^\bullet(S)[[c_1, \ldots, c_n]].$$

By 4.8, $K^\bullet(\mathbf{G}_S^{n,r})$[373] is the $K^\bullet(S)$-algebra generated by the $c_i = \gamma^i(F_1 - n)$, $1 \leq i \leq n$, and the $\gamma^j(F_2 - r + n)$, $1 \leq j \leq r - n$, subject to the relations deduced from:

$$\gamma_t(F_1 - n).\gamma_t(F_2 - r + n) = \gamma_t(r - r) = 1,$$

or again:

$$\gamma_t(F_2 - r + n) = 1/\gamma_t(F_1 - n).$$

This relation expresses the $\gamma^j(F_2 - r + n)$ in terms of the c_i, and gives a series of isobaric relations among the c_i (c_i being assumed to have weight i) by writing that the coefficients of $1/\gamma_t(F_1 - n)$ are zero in degree $> r - n$; moreover, these relations generate the ideal of relations among the c_i. We may therefore write: [print 401][374]

$$K^\bullet(\mathbf{G}_S^{n,r}) = K^\bullet(S)[c_1, \ldots, c_n]/I_r,$$

where I_r is a homogeneous ideal of $K^\bullet(S)[c_1, \ldots, c_n]$, of degree $> r - n$. It follows from definition (4.9.1) that $K^\bullet(\mathbf{G}_S^{n,\infty})$[375] is the separated completion of $K^\bullet(S)[c_1, \ldots, c_n]$ for the topology defined by the I_r. This topology is finer than the topology defined by weight; we shall show that it is equivalent to it, which will imply the proposition.

Put $A = K^\bullet(S)[c_1, \ldots, c_n]/I_r$, and let C be the A-algebra generated by generators $x_1, \ldots, x_n$ subject to the relations:

$$\sigma_j(x_1, \ldots, x_n) = c_j,$$

where the σ_j denote the elementary symmetric functions. By 4.4, the canonical homomorphism $A \to C$ is injective. We obtain:

$$\gamma_t(F_1 - n) = 1 + c_1 t + \ldots + c_n t^n = \prod_{i=1}^{n} (1 + x_i t).$$

Since $1/\gamma_t(F_1 - n)$ is a polynomial, so is $1/(1 + x_i t)$ for every i. Consequently, the x_i are nilpotent, and there is an integer N such that every polynomial of degree $> N$ in the x_i, with coefficients in $K^\bullet(S)$, is zero. Thus every polynomial of A, of weight $> N$, is zero, i.e. every polynomial of $K^\bullet(S)[c_1, \ldots, c_n]$ of weight $> N$ belongs to I_r. It follows that the two topologies under consideration are equivalent, which completes the proof.

5. Applications to projective bundles; study of f_*

5.1. In this section, S is a quasi-compact scheme, $\underline{E}$ an $\underline{O}_S$-Module locally free of rank $r + 1$, and X the projective bundle associated to $\underline{E}$. On X, there is the canonical exact sequence:

$$0 \longrightarrow \underline{F} \longrightarrow f^*(\underline{E}) \longrightarrow \underline{O}_X(1) \longrightarrow 0,$$

[print 402]where f is the structural morphism $X \longrightarrow S$.

The $K^\bullet$ considered will be those defined using locally free sheaves of finite type. We shall denote by E, F, ξ the classes of $\underline{E}$, $\underline{F}$, $\underline{O}_X(1)$. Put $x = 1 - \xi^{-1}$, $x' = \xi - 1 = \xi x$, and denote by $\bar{x}$, $\bar{x}'$ the classes of x, x' in $\mathrm{Gr}^1 K^\bullet(X)$. Since $\xi - 1$ and x belong to $\mathrm{Fil}^1 K^\bullet(X)$, we have $(\xi - 1)x \in \mathrm{Fil}^2 K^\bullet(X)$, i.e. $x \equiv \xi x = x'$ mod. $\mathrm{Fil}^2 K^\bullet(X)$ i.e. $\bar{x}' = \bar{x}$[376]

5.2. If $\underline{G}$ is an $\underline{O}_X$-Module locally free of finite type, the $R^i f_*(\underline{G})$ are in general not locally free of finite type on S, so that a homomorphism $f_* : K^\bullet(X) \to K^\bullet(S)$ cannot be defined as it is for the $K^\bullet$ defined using perfect complexes. Theorem 1.1 allows us to get around this difficulty as follows: if $y \in K^\bullet(X)$, there is a unique way to write:

$$y = a_0 + a_1\xi^{-1} + \ldots + a_r\xi^{-r}, \qquad \text{with } a_i \in K^\bullet(S); \tag{5.2.1}$$

we then put:

$$f_*(y) = a_0. \tag{5.2.2}$$

Clearly the homomorphism thus defined is additive and satisfies the projection formula, i.e. is $K^\bullet(S)$-linear.

The following relations hold:

$$f_*(1) = 1; \qquad f_*(\xi^n) = 0 \quad \text{for } -r \leq n < 0; \qquad f_*(\xi^n) = \sigma_n, \quad 1 \leq n. \tag{5.2.3}$$

To verify the last group of relations, we proceed by induction on n. Suppose that, for $p \leq n-1$, $f_*(\xi^p) = \sigma_p$; relation (1.9.1), multiplied by a suitable power of ξ, gives:

$$\sum_{i=0}^{r+1} (-1)^i \lambda^i(E) \xi^{n-i} = 0.$$

We can apply f_* to it: [print 403]

$$\sum_{i=0}^{r+1} (-1)^i \lambda^i(E) f_*(\xi^{n-i}) = 0.$$

The result follows immediately from relation (1.11.3) and the induction hypothesis.

Proposition 5.3. With the notation and hypotheses of 5.1[377], for every $k \geq 0$ there is a direct-sum decomposition of $K^\bullet(S)$-modules:

$$\mathrm{Fil}^k K^\bullet(X) = \mathrm{Fil}^k K^\bullet(S) + \mathrm{Fil}^{k-1} K^\bullet(S).x + \ldots + \mathrm{Fil}^{k-r} K^\bullet(S).x^r,$$

the filtration under consideration being the filtration of an augmented λ-algebra defined in V 3.9.1 and 3.10.

First note that x has augmentation zero. Moreover: [378]

$$\gamma_t(x) = \lambda_{t/(1-t)}(1 - \xi^{-1}) = \left(1 + \frac{t}{1-t}\right)\left(1 + \frac{\xi^{-1}t}{1-t}\right)^{-1} = 1/(1 - xt).$$

Thus $\gamma^i(x) = x^i$ for every i. Consequently, $K^\bullet(X)$ is a quotient of the $K^\bullet(S)$-λ-algebra generated by a generator X subject to the relations $\gamma^i(X) = X^i$ for every i, and the ideal by which the quotient is taken is generated by relation (1.13.1). We can therefore apply V 4.12 and V 4.13.ii), and are reduced to showing that, for every $n \geq 0$:

$$x^n \in \mathrm{Fil}^n K^\bullet(S) + \mathrm{Fil}^{n-1} K^\bullet(S).x + \ldots + \mathrm{Fil}^{n-r} K^\bullet(S).x^r.$$

This is evident for $n \leq r$; we continue by induction. Relation (1.13.1) gives:

$$x^n = -\sum_{i=0}^{r} \gamma^{r+1-i}(\check{E} - r - 1).x^{n-r-1+i}, \qquad \text{with } n \geq r+1;$$

applying the induction hypothesis to the $x^{n-r-1+i}$, we deduce the desired result, the sum being direct since $1, \ldots, x^r$ are independent over $K^\bullet(S)$.

[print 404]**Corollary 5.4.** The filtration induced by $K^\bullet(X)$ on $K^\bullet(S)$ is identical to the filtration of $K^\bullet(S)$.

Corollary 5.5. Let S be a quasi-compact scheme, $\underline{E}_1, \ldots, \underline{E}_n$ be O_S-Modules locally free of finite type, $\mathbf{D}(\underline{E}_i)$[379] the flag scheme of $\underline{E}_i$, $X = \mathbf{D}(\underline{E}_1) \times_S \ldots \times_S \mathbf{D}(\underline{E}_n)$[380]. Then the canonical homomorphism:

$$\operatorname{Gr} K^\bullet(S) \longrightarrow \operatorname{Gr} K^\bullet(X)$$

is injective.

The argument of 3.1 reduces us to proving the same property when X is a projective bundle over S. This case then follows immediately from 5.4.

Corollary 5.6. With the notation and hypotheses of 5.1, $\operatorname{Gr} K^\bullet(X)$ is a free $\operatorname{Gr} K^\bullet(S)$-module, with basis $1, \bar{x}, \ldots, \bar{x}^r$. For every k, we have:

$$\operatorname{Gr}^k K^\bullet(X) = \operatorname{Gr}^k K^\bullet(S) \oplus \operatorname{Gr}^{k-1} K^\bullet(S).\bar{x} \oplus \ldots \oplus \operatorname{Gr}^{k-r} K^\bullet(S).\bar{x}^r.$$

Finally, the elements $1, \bar{x}, \ldots, \bar{x}^{r+1}$ are related by the relation:

$$\sum_{i=0}^{r+1} c^{r+1-i}(\check{E})\bar{x}^i = 0,$$

where $c^k(\check{E})$ is the k-th Chern class of $\check{E}$ (V 6.7).

An immediate consequence of 1.1, 5.3, and (1.13.1).

Corollary 5.7. With the notation and hypotheses of 4.5, the $\operatorname{Gr} K^\bullet(S)$-algebra $\operatorname{Gr} K^\bullet(X)$ is generated by the $\bar{c}_i^{(j)}$, where $\bar{c}_i^{(j)}$ is the class in $\operatorname{Gr}^i K^\bullet(X)$ of $\gamma^i(F_j - p_j)$. The ideal of relations among the $\bar{c}_i^{(j)}$ is generated by the relations deduced from:

$$c(E - n) = \prod_{j=1}^{k} c(F_j - p_j),$$

[print 405]where c is the total Chern class (V 6.7).

This result is proved by repeating the proof of 4.6 and using 5.6 (compare [2], Exp 4).

Corollary 5.8. With the notation and hypotheses of 5.1, for every k we have:

$$f_*(\operatorname{Fil}^k K^\bullet(X)) \subset \operatorname{Fil}^{k-r} K^\bullet(S).$$

An immediate consequence of 5.3 and the projection formula.

Proposition 5.9. With the notation and hypotheses of 5.1, for every $i \geq 1$ we have:

$$f_*(\lambda^i(F)) = 0 \qquad (\text{whence } f_*(\gamma^i(F)) = 0).$$

In particular, putting $\lambda_{-1}(F) = \sum_{i=0}^{r} (-1)^i \lambda^i(F)$, we obtain:

$$f_*(\lambda_{-1}(F)) = 1.$$

We define a map

$$\bar{f}_* : 1 + K^\bullet(X)[[t]]^+ \longrightarrow 1 + K^\bullet(S)[[t]]^+$$

by putting:

$$\bar{f}_* \left(\sum_{i \geq 0} a_i t^i \right) = \sum_{i \geq 0} f_*(a_i) t^i.$$

In general this is not a group homomorphism; nevertheless, the following relation holds for $\varphi \in 1 + K^\bullet(X)[[t]]^+$, $\psi \in 1 + K^\bullet(S)[[t]]^+$:

$$\bar{f}_*(\varphi . \hat{G}(f^*)(\psi)) = \psi . \bar{f}_*(\varphi).$$

Indeed, it holds coefficient by coefficient and follows immediately from the projection formula.

From the relation $E = F + \xi$, we therefore deduce: [print 406][381]

$$\bar{f}_*(\lambda_t(F)) = \bar{f}_*(\lambda_t(E)\lambda_t(-\xi)) = \lambda_t(E)\bar{f}_*(\lambda_t(-\xi));$$

Now $\bar{f}_*(\lambda_t(-\xi)) = \sum_{i \geq 0} (-1)^i f_*(\xi^i) t^i = \sigma_{-t}(E)$[382] by (5.2.3).[383] Using relation (1.12.1), we finally obtain:

$$\bar{f}_*(\lambda_t(F)) = 1;$$

whence the desired relations

Proposition 5.10. With the notation and hypotheses of 5.1, let $z \in K^\bullet(X)$. For $z(1 - \xi) = 0$ to hold, it is necessary and sufficient that there exist $y \in K^\bullet(S)$ such that:

$$\text{i)} \quad z = y.\lambda_{-1}(F); \qquad \text{ii)} \quad y.\lambda_{-1}(E) = 0.$$

From $E = F + \xi$, we deduce $\lambda_{-1}(E) = \lambda_{-1}(F).\lambda_{-1}(\xi) = \lambda_{-1}(F).(1 - \xi)$. The condition is therefore sufficient.

Let z be such that $z(1 - \xi) = 0$, that is, $zx' = 0$. By Remark 1.13, there are unique $a_i \in K^\bullet(S)$ such that:

$$z = \sum_{i=0}^{r} a_i x'^i$$

Moreover, the x'^i satisfy a dependence equation:

$$\sum_{i=0}^{r} c_i x'^i + x'^{r+1} = 0, \qquad \text{with } c_i = (-1)^{r+i+1} \gamma^{r+1-i}(E - r - 1).$$

Since $zx' = 0$, we obtain:

$$a_r x'^{r+1} = -\sum_{i=1}^{r} a_{i-1} x'^i = -\sum_{i=0}^{r} a_r c_i x'^i.$$

Using the independence of $1, \ldots, x'^r$, we find:

$$a_r c_0 = 0; \qquad \text{for } 0 \leq i < r, \quad a_i = a_r c_{i+1}.$$

[print 407]Thus:

$$z = a_r . \left(\sum_{i=0}^{r-1} c_{i+1} x'^i + x'^r \right).$$

Now: [384]

$$\begin{aligned}\sum_{i=0}^{r-1} c_{i+1}x'^i + x'^r &= (-1)^r.\left(\sum_{i=0}^{r-1}\gamma^{r-i}(E-r-1).(-1)^i x'^i + (-1)^r x'^r\right)\\ &= (-1)^r.\left(\sum_{i=0}^{r}\gamma^{r-i}(E-r-1)\gamma^i(-x')\right)\\ &= (-1)^r\gamma^r(E-r-1-x')\\ &= (-1)^r\gamma^r(F-r)\\ &= (-1)^r\lambda^r(F-1)\\ &= \lambda_{-1}(F).\end{aligned}$$

Putting $a_r = y$, we therefore obtain:

$$z = y.\lambda_{-1}(F).$$

Moreover, $y.c_0 = 0$; now $c_0 = (-1)^{r+1}\gamma^{r+1}(E-r-1) = \lambda_{-1}(E)$; thus $y.\lambda_{-1}(E) = 0$.

Proposition 5.11. With the notation and hypotheses of 5.1, let $z \in \operatorname{Gr} K^\bullet(X)$. For $z.\bar{x}' = 0$ to hold, it is necessary and sufficient that there exist $y \in \operatorname{Gr} K^\bullet(S)$ such that:

$$\text{i)}\quad z = y.c^r(F); \qquad \text{ii)}\quad y.c^{r+1}(E) = 0.$$

The proof is identical to that of 5.10, working in the associated graded rings and using 5.6.

6. Study of the filtration of $K^\bullet(X)$, X having an ample sheaf

In this section, X is a scheme having an ample sheaf $\underline{L}$. Recall that in this case the $K^\bullet$ defined in terms of locally free sheaves of finite type, or of perfect complexes, are equal (IV, 2.9). We denote by [print 408]ε the augmentation homomorphism: $K^\bullet(X) \longrightarrow H^0(X, \underline{\mathbb{Z}})$.

Proposition 6.1. Let X be a quasi-compact scheme having an ample sheaf $\underline{L}$. Then $\operatorname{Fil}^1 K^\bullet(X)$ is a nil ideal.

Since $\operatorname{Fil}^1 K^\bullet(X)$ is generated by elements of the form $E - \varepsilon(E)$, where E is the class of an $\underline{O}_X$-Module locally free of finite type $\underline{E}$, and $\varepsilon : K^\bullet(X) \to H^0(X, \underline{\mathbb{Z}})$ is the augmentation of $K^\bullet(X)$, it suffices to show that such an element is nilpotent. The quasi-compactness of X, and the commutation of $K^\bullet$ with finite direct sums, reduce us to the case where $\underline{E}$ has constant rank. Finally, passing to the flag bundle of $\underline{E}$, which gives an injection on the $K^\bullet$ (3.1) and preserves the hypothesis of the existence of an ample sheaf, we may suppose that $\underline{E}$ is invertible.

Let $\underline{E}^*$ be the dual of $\underline{E}$. For n sufficiently large, $\underline{E}^*(n) = \underline{E}^* \otimes_{\underline{O}_X} \underline{L}^{\otimes n}$ is generated by its global sections; we can therefore find an integer m and a surjective homomorphism: $(\underline{O}_X)^m \longrightarrow \underline{E}^*(n)$, hence (tensoring by $\underline{E}(-n)$) a surjective homomorphism: $(\underline{E}(-n))^m \longrightarrow \underline{O}_X$. The kernel of this homomorphism is locally free of rank $m-1$. Thus, denoting by L the class of $\underline{L}$, we obtain:

$$\lambda^m(mEL^{-n} - 1) = 0.$$

Now $\lambda^m(mEL^{-n} - 1) = (-1)^m\lambda_{-1}(mEL^{-n}) = (-1)^m\lambda_{-1}(EL^{-n})^m = (-1)^m(1 - EL^{-n})^m$. Consequently, for every invertible sheaf $\underline{E}$, the element $1 - EL^{-n}$ is nilpotent for n sufficiently large, or equivalently its product $L^n - E$ by L^n is nilpotent.

We can then write:

$$E - 1 = (L^n - 1) - (L^n - E)$$

and, by what precedes, $L^n - E$ and $L^n - 1$ are nilpotent on choosing n sufficiently large. Thus $E - 1$ is nilpotent.

[print 409]**Corollary 6.2.** With the hypotheses of 6.1, let $R \subset K^\bullet(X)$ be a sub-λ-ring generated by the classes of finitely many sheaves. Then the λ-ring filtration of R is discrete, and $R \cap \mathrm{Fil}^1 K^\bullet(X)$ is nilpotent. Moreover, R is a module of finite type over $R \cap H^0(X, \underline{\mathbb{Z}})$ (V 3.9.1).

Since X is quasi-compact, there is a finite covering of X by disjoint open sets such that, on each open set of this covering, the sheaves whose classes generate R all have constant rank. The standard argument then reduces us to the case[385] where these sheaves have constant rank on X; R is then augmented to $\mathbb{Z}$.

Let $\underline{E}_i$, $i = 1, \ldots, k$, be the sheaves whose classes generate R, E_i their classes, and $x_i = E_i - \varepsilon(E_i)$. We can write R as a quotient of the λ-ring generated by the x_i subject to the relations $\gamma^j(x_i) = 0$ for $j > r_i = \varepsilon(E_i)$, that is, $\mathbb{Z}[x_i, \gamma^j(x_i)]_{\substack{1 \le i \le k \\ 2 \le j \le r_i}}$; consequently, R is noetherian. But by Proposition 6.1, the x_i, $\gamma^j(x_i)$ are nilpotent. Thus the augmentation ideal of R is nilpotent; moreover, it is precisely $R \cap \mathrm{Fil}^1 K^\bullet(X)$.

Every element of $\mathrm{Fil}^n R$ can be written as a polynomial with no components of weight $< n$ in the x_i, $\gamma^j(x_i)$. Let m be the largest integer such that there exists i such that $\gamma^{m-1}(x_i) \neq 0$; let p be an integer such that the p-th powers of the x_i, $\gamma^j(x_i)$ are all zero. For $n > kp(1 + \ldots + m)$, one sees at once that $\mathrm{Fil}^n R = 0$; the filtration of R is therefore discrete.

Since the filtration of R is discrete, there are only finitely many nonzero monomials in the x_i, $\gamma^j(x_i)$. These monomials form a system of generators of R as a $\mathbb{Z}$-module, which completes the proof.

[print 410]**Corollary 6.3.** With the hypotheses of 6.1, let A be a commutative ring. For an element y of $K^\bullet(X) \otimes_{\mathbb{Z}} A = K^\bullet(X)_A$ to be invertible, it is necessary and sufficient that $z = (\varepsilon \otimes \mathrm{Id}_A)(y)$ be invertible. In particular, if $\underline{E}$ is a locally free sheaf of finite type on X, everywhere $\neq 0$, then its class in $K^\bullet(X)_{\mathbb{Q}}$ is invertible.

The augmentation ideal of $K^\bullet(X) \otimes_{\mathbb{Z}} A$ is $\mathrm{Fil}^1 K^\bullet(X) \otimes_{\mathbb{Z}} A$, and is therefore a nil ideal by 6.1. Thus, for an element of $K^\bullet(X) \otimes_{\mathbb{Z}} A$ to be invertible, it is necessary and sufficient that its augmentation be invertible.

Corollary 6.4. Let X be a quasi-compact scheme, having an ample sheaf, and $f : X' \to X$ a proper and perfect morphism. Suppose in addition that one of the following two conditions is satisfied:

i) f projective, ii) X noetherian,

so that the "finiteness theorem" holds (III 2.2). We are given a ring A, and consider the "direct image" homomorphism (IV 2.11):

$$f_* : K^\bullet(X') \otimes_{\mathbb{Z}} A \longrightarrow K^\bullet(X) \otimes_{\mathbb{Z}} A.$$

Finally suppose that there is an element $x' \in K^\bullet(X') \otimes_{\mathbb{Z}} A$ such that $\varepsilon(f_*(x')) \in H^0(X, \underline{A})$ is invertible. Then the homomorphism:

$$f^* : K^\bullet(X) \otimes_{\mathbb{Z}} A \longrightarrow K^\bullet(X') \otimes_{\mathbb{Z}} A$$

is injective.

We write the projection formula:

$$f_*(f^*(x)x') = x.f_*(x').$$

Now $f_*(x')$ has invertible augmentation, so it is invertible (6.3). Consequently, the composite homomorphism: [print 411]

$$x \rightsquigarrow f^*(x) \rightsquigarrow f^*(x)x' \rightsquigarrow f_*(f^*(x)x')$$

is injective, so f^* is also injective.

6.5. Let X be a noetherian scheme of dimension d, having an ample sheaf $\underline{L}$. We endow $K^\bullet(X)$ with the filtration defined as follows (cf. [1]): x belongs to $K^\bullet(X)_i$ if and only if, for every closed subset Y of X, there is a perfect complex $\underline{M}^\bullet(Y)$ such that:

$$\text{i)}\ \ \mathrm{cl}(\underline{M}^\bullet(Y)) = x;$$

[386]

$$\text{ii)}\ \ \mathrm{Codim}(\mathrm{Supp}(H^*(\underline{M}^\bullet(Y))) \cap Y, Y) \geq i, \quad \text{with } H^*(\underline{M}^\bullet(Y)) = \bigoplus_{i\in\mathbb{Z}} H^i(\underline{M}^\bullet(Y)).$$

The $K^\bullet(X)_i$ thus defined are additive subgroups of $K^\bullet(X)$. Indeed, let $x, x' \in K^\bullet(X)_i$, and, for a given closed subset Y in X, let $\underline{M}^\bullet$ and $\underline{M}'^\bullet$ be the corresponding complexes; then $x + x'$ is the class of the complex $\underline{M}^\bullet \oplus \underline{M}'^\bullet$, and

$$\mathrm{Supp}(H^*(\underline{M}^\bullet \oplus \underline{M}'^\bullet)) = \mathrm{Supp}(H^*(\underline{M}^\bullet) \oplus H^*(\underline{M}'^\bullet)) = \mathrm{Supp}(H^*(\underline{M}^\bullet)) \cup \mathrm{Supp}(H^*(\underline{M}'^\bullet));$$

since the codimension of a finite union of closed subsets is the infimum of the codimensions of the closed subsets, we indeed have $x + x' \in K^\bullet(X)_i$.

Proposition 6.6. The filtration 6.5[387] of $K^\bullet(X)$ is a ring filtration and $K^\bullet(X)_{d+1} = 0$.

Let $x \in K^\bullet(X)_i$, $x' \in K^\bullet(X)_j$, and let Y be a closed subset of X. There is a complex $\underline{M}^\bullet$ such that $x = \mathrm{cl}(\underline{M}^\bullet)$, and $\mathrm{codim}(\mathrm{Supp}(H^*(\underline{M}^\bullet)) \cap Y, Y) \geq i$; there is a complex $\underline{M}'^\bullet$ such that $x' = \mathrm{cl}(\underline{M}'^\bullet)$, and:

$$\mathrm{codim}\big(\mathrm{Supp}(H^*(\underline{M}^\bullet)) \cap \mathrm{Supp}(H^*(\underline{M}'^\bullet)) \cap Y, \mathrm{Supp}(H^*(\underline{M}^\bullet)) \cap Y\big) \geq j.$$

We then have: $xx' = \mathrm{cl}(\underline{M}^\bullet \overset{\mathbb{L}}{\otimes} \underline{M}'^\bullet)$, and:

$$\mathrm{codim}\big(\mathrm{Supp}(H^*(\underline{M}^\bullet)) \cap \mathrm{Supp}(H^*(\underline{M}'^\bullet)) \cap Y, Y\big) \geq i + j$$

(EGA 0_{IV} 14.2.2.3). [print 412]Thus, to complete the proof, it suffices to note that:

$$\mathrm{Supp}\big(H^*(\underline{M}^\bullet \overset{\mathbb{L}}{\otimes} \underline{M}'^\bullet)\big) \subset \mathrm{Supp}(H^*(\underline{M}^\bullet)) \cap \mathrm{Supp}(H^*(\underline{M}'^\bullet)).$$

To show that $K^\bullet(X)_{d+1}$ is zero, we take $x \in K^\bullet(X)_{d+1}$, and $Y = X$ so that there is a complex $\underline{M}^\bullet$ such that $x = \mathrm{cl}(\underline{M}^\bullet)$, and $\mathrm{codim}_X(\mathrm{Supp}(H^*(\underline{M}^\bullet))) \geq d + 1$; since X has dimension d, this implies $\mathrm{Supp}(H^*(\underline{M}^\bullet)) = \varnothing$, hence $\underline{M}^\bullet$ is acyclic, and $x = 0$.

Lemma 6.7. Let $X = \mathrm{Spec}(A)$ be an affine noetherian scheme, $x_1, \ldots, x_k$ finitely many points of X, and $\underline{E}$ an invertible $\underline{O}_X$-Module. Then there is a section $s \in \Gamma(X, \underline{E})$ such that, for every i, s_{x_i} is a basis of $\underline{E}_{x_i}$[388].

Put $M = \Gamma(X, \underline{E})$. For every i, let x'_i be a closed point of X such that $x'_i \in \overline{\{x_i\}}$. The x'_i correspond to maximal ideals $\underline{m}_i$ of A, which we may suppose all distinct, changing the index set so that each of the $\underline{m}_i$ occurs once and only once in the family $(\underline{m}_i)$. Since $\underline{E}$ is invertible, for every i there is an isomorphism $M_{\underline{m}_i} \simeq A_{\underline{m}_i}$ from which we deduce an isomorphism $M/\underline{m}_i M \simeq A/\underline{m}_i$, then $\prod_i M/\underline{m}_i M \simeq \prod_i A/\underline{m}_i$, or again $A/\prod_i \underline{m}_i \simeq M/(\prod_i \underline{m}_i)M$. Lifting this last isomorphism, we obtain a homomorphism $A \to M$ such that the corresponding homomorphism $A_{\underline{m}_i} \to M_{\underline{m}_i}$ is, for every i, an isomorphism modulo $\underline{m}_i$, hence an isomorphism by Nakayama. The image of 1 under this homomorphism therefore gives a section s of $\underline{E}$ that is a basis of $\underline{E}_{x'_i}$ for every i. Since the set of points where s is a basis of $\underline{E}$ is an open subset of X, s is also a basis of $\underline{E}_{x_i}$ for $i = 1, \ldots, k$.

Lemma 6.8. Let X be a noetherian scheme having an ample sheaf $\underline{L}$, $\underline{E}$ an invertible $\underline{O}_X$-Module, and $x_1 \ldots, x_k$ points of X. Then there is an [print 413]integer n such that, for every $k \geq 1$, there is a section $s \in \Gamma(X, \underline{E}(kn))$ that is a basis of $\underline{E}(kn)_{x_i}$ for every i.

There is an integer n_0 and a section $f \in \Gamma(X, \underline{L}^{\otimes n_0})$ such that, for every i, $x_i \in X_f$, and such that X_f is an affine open subset (EGA II 4.5.4). On X_f, there is a section $t \in \Gamma(X_f, \underline{E})$ satisfying the conditions of Lemma 6.7; moreover, there is an integer n'_0 such that the section $f^{\otimes n'_0} \otimes t \in \Gamma(X_f, \underline{E}(n_0 n'_0))$ extends to the whole of X. Put $n = n_0 n'_0$; then $f^{\otimes n'_0} \in \Gamma(X, \underline{L}^n)$; taking the tensor product of the section of $\underline{E}(n)$ obtained above and $f^{\otimes kn'_0}$ therefore gives a section of $\underline{E}((k+1)n)$ for every $k \geq 0$. Since, for every p, $X_{f^{\otimes p}} = X_f$, and the section t is a basis of $\underline{E}_{x_i}$ for every i, the sections of $\underline{E}(kn)$ thus constructed for every $k \geq 1$ form a basis of $\underline{E}(kn)_{x_i}$ for every i.

Theorem 6.9. Let X be a noetherian scheme of dimension d, having an ample sheaf $\underline{L}$. Then $\mathrm{Fil}^{d+1} K^\bullet(X) = 0$, and a fortiori $(\mathrm{Fil}^1 K^\bullet(X))^{d+1} = 0$.

(This last relation is due to J.P. Serre, in the general case considered here).

We shall first show that if $\underline{E}, \underline{E}'$ are two invertible sheaves on X, with classes E, E', then $E - E' \in K^\bullet(X)_1$.

Let Y be a closed subset of X, $Y_1, \ldots, Y_k$ its irreducible components, and for every i, let x_i be a point of $Y_i - \bigcup_{j \neq i} Y_j$. By Lemma 6.8, there are integers n, n' such that, for any integers $k, k' \geq 1$, there are homomorphisms from $\underline{O}_X$ to $\underline{E}(kn)$, $\underline{E}'(k'n')$ that are isomorphisms at the points x_i; we therefore obtain morphisms:

$$\underline{L}^{\otimes -kn} \xrightarrow{s_k} \underline{E}, \qquad \underline{L}^{\otimes -k'n'} \xrightarrow{s'_{k'}} \underline{E}',$$

[print 414]that are isomorphisms at the points x_i. Now consider the complex:

$$\underline{N}^\bullet = \ldots \to 0 \to \underline{L}^{\otimes -nn'} \xrightarrow{(s'_n, 0)} \underline{E}' \oplus \underline{L}^{\otimes -nn'} \xrightarrow{0 + s_{n'}} \underline{E} \to 0 \to \ldots$$

The class of $\underline{N}^\bullet$ is $E - E'$; moreover, $\underline{N}^\bullet$ is acyclic at the points x_i, so $\mathrm{Supp}(H^*(\underline{N}^\bullet)) \cap Y$ contains no irreducible component of Y and consequently has codimension ≥ 1 in Y; this shows that $E - E'$ belongs to $K^\bullet(X)_1$.

Now let x be an element of $\mathrm{Fil}^{d+1} K^\bullet(X)$; we shall show that $x = 0$. We may suppose that x can be written:

$$x = \gamma^{i_1}(E_1 - F_1) \ldots \gamma^{i_k}(E_k - F_k), \qquad \text{with } i_1 + \ldots + i_k \geq d+1,$$

the E_i, F_i being the classes of locally free sheaves of finite type and, for every i, $\underline{E}_i$ and $\underline{F}_i$ having the same rank. Moreover, decomposing X into a sum of finitely many open sets and arguing in each open set, we may suppose the $\underline{E}_i, \underline{F}_i$ to have constant rank. Let X' be the fibre product over X of the flag schemes of the $\underline{E}_i, \underline{F}_i$. In $K^\bullet(X')$, the images of the E_i, F_i are sums of classes of invertible sheaves. Let ξ, η be the classes of two invertible $\underline{O}_{X'}$-Modules[389]; we have:

$$\gamma_t(\xi - \eta) = \gamma_t(\xi - 1)/\gamma_t(\eta - 1) = (1 + (\xi - 1)t)/(1 + (\eta - 1)t).$$

Thus $\gamma^i(\xi - \eta) = (-1)^{i-1}(\xi - \eta)(\eta - 1)^{i-1}$. Now X' is noetherian and has an ample sheaf, so, by what precedes and 6.6, $\gamma^i(\xi - \eta) \in K^\bullet(X')_i$. Using the additivity of γ_t and 6.6, we deduce that $f^*(\gamma^{i_j}(E_j - F_j)) \in K^\bullet(X')_{i_j}$, and $f^*(x) \in K^\bullet(X')_{d+1}$, where f is the canonical morphism $X' \to X$.

We shall deduce that $f^*(x) = 0$. Let r be the relative dimension of X' over X. There is an element x' of $K^\bullet(X')$, belonging to a basis of $K^\bullet(X')$ over $K^\bullet(X)$, such that: [print 415][390]

$$x' = \prod_{i=1}^{r} (\xi_i - 1),$$

where the ξ_i are the classes of invertible sheaves: to show this, the argument of 3.1 reduces us to the case of a projective bundle over X. This last case is an immediate consequence of 1.13. The element x' under consideration belongs to $K^\bullet(X')_r$ by what precedes, and consequently $x'f^*(x) \in K^\bullet(X')_{d+r+1}$. Now $\dim X' = d + r$; thus, by 6.6, $x'f^*(x) = 0$. Since x' belongs to a basis of $K^\bullet(X')$ over $K^\bullet(X)$, this implies $f^*(x) = 0$. Since f^* is injective (3.1), we finally have $x = 0$.

Problem 6.10. Under the hypotheses of 6.9, is it true that, for every k, we have: $\mathrm{Fil}^k K^\bullet(X) \subset K^\bullet(X)_k$?

BIBLIOGRAPHY

[1] Bass H., K-Theory and stable algebra, Publ. Math. de l'IHES, n°22.

[2] Chevalley C., Séminaire 1958 : Anneaux de Chow et applications.

[3] Hakim M., Topos localement annelés et schémas relatifs, thesis (in preparation).

[4] Hartshorne R., Residues and duality, Lecture Notes in Mathematics[391] n°20, Springer-Verlag.

[5] Serre J.-P., Le groupe de Grothendieck des schémas en groupes réductifs, Publications Mathématiques de l'IHES, n°34.

[print 416]

Exposé VII
REGULAR IMMERSIONS AND COMPUTATION OF $K^\bullet$ OF A BLOWN-UP SCHEME

by Pierre BERTHELOT

1. Generalities on regular immersions

As the classical notion of regular immersion (EGA IV 16.9.2) proves inadequate in the non-noetherian case, we shall define a weaker notion, which nevertheless gives the desired properties and is equivalent to the usual definition in the noetherian case.

Definition 1.1. Let A be a commutative ring, E a projective A-module of finite type, $u : E \longrightarrow A$ an A-linear homomorphism. We say that u is regular if the Koszul complex $K_\bullet(u)$ defined by u is acyclic in dimension > 0, and hence is a resolution of $B = A/u(E)$.

Recall that the Koszul complex associated with u is defined as follows: the graded A-module $K_\bullet(u)$ is the exterior algebra $\Lambda(E)$, and the boundary operator is interior multiplication by u regarded as an element of the dual $\check{E}$: it is an antiderivation of degree -1 of $\Lambda(E)$, equal to u on E.

Lemma 1.2. Let $u : E \longrightarrow A$ be a regular homomorphism. Let A' be an A-algebra, $u' : E' \longrightarrow A'$ the homomorphism deduced from u by base change; suppose that u' is regular. Put $J = u(E)$, $J' = JA' = u(E')$. Then we have the following relations for every integer $n \geq 0$:

$$J^n/J^{n+1} \otimes_A A' \xrightarrow{\sim} J'^n/J'^{n+1}, \tag{1.2.1}$$

$$\mathrm{Tor}_i^A(J^n/J^{n+1}, A') = 0 \quad \text{for } i \geq 1; \tag{1.2.2}$$

$$J^n \otimes_A A' \xrightarrow{\sim} J'^n, \tag{1.2.3}$$

$$\mathrm{Tor}_i^A(J^n, A') = 0 \quad \text{for } i \geq 1. \tag{1.2.4}$$

a) First suppose that the J^n/J^{n+1} are projective over $B = A/J$; since J is of finite type, they are locally free of finite type over B. [print 417]The relations to be proved can be checked pointwise on $\mathrm{Spec}(A)$, so we are reduced to the case where A is local, with $u(E) \subset \mathrm{rad}(A)$. The J^n/J^{n+1} are then free over B, and the relations $\mathrm{Tor}_i^A(J^n/J^{n+1}, A') = 0$ for $i \geq 1$ are equivalent to the relations $\mathrm{Tor}_i^A(B, A') = 0$ for $i \geq 1$. Now we can compute the Tor_i^A using the projective resolution $K_\bullet(u)$ of B; they are the homology groups of the complex $K_\bullet(u) \otimes_A A' = K_\bullet(u')$. Since u' is assumed regular, this last complex is acyclic in degrees > 0, whence the relations $\mathrm{Tor}_i^A(J^n/J^{n+1}, A') = 0$, for $i \geq 1$.

We deduce the relations $\mathrm{Tor}_i^A(A/J^n, A') = 0$ for $i \geq 1$ by induction on n using the exact Tor sequence associated with the exact sequence:

$$0 \longrightarrow J^n/J^{n+1} \longrightarrow A/J^{n+1} \longrightarrow A/J^n \longrightarrow 0.$$

Since for every k we have $J^k A' = J'^k$, this exact sequence and $\mathrm{Tor}_1^A(A/J^n, A') = 0$ give the relations $J^n/J^{n+1} \otimes_A A' \xrightarrow{\sim} J'^n/J'^{n+1}$. From the exact sequence:

$$0 \longrightarrow J^n \longrightarrow A \longrightarrow A/J^n \longrightarrow 0,$$

and $\mathrm{Tor}_1^A(A/J^n, A') = 0$, we deduce the relations $J^n \otimes_A A' \simeq J'^n$. Finally, from the exact Tor sequence associated with the exact sequence:

$$0 \longrightarrow J^{n+1} \longrightarrow J^n \longrightarrow J^n/J^{n+1} \longrightarrow 0,$$

we deduce the relations $\mathrm{Tor}_i^A(J^n, A') = 0$ for $i \geq 1$ by induction on n, the case $n = 0$ being evident.

b) We shall now show that the J^n/J^{n+1} are projective over $B = A/J$. Since the question is local on $\mathrm{Spec}(A)$, we may suppose E free, say $E = (A)^d$, so that u is defined by a sequence $f_1, \ldots, f_d$ of elements of A. Put $A_0 = \mathbb{Z}[T_1, \ldots, T_d]$ and let $A_0 \longrightarrow A$ be the homomorphism that sends [print 418]T_i to f_i; denote by $u_0 : E_0 = (A_0)^d \longrightarrow A_0$ the homomorphism defined by the sequence $(T_1, \ldots, T_d)$, so that u is deduced from u_0 by base change from A_0 to A. Since the sequence $(T_1, \ldots, T_d)$ is regular (in the classical sense), the homomorphism u_0 is regular, and the hypotheses of the lemma hold for the base change $A_0 \longrightarrow A$. Now the J_0^n/J_0^{n+1} are free of finite type over $\mathbb{Z}$, so we can apply the results of a). Consequently $J^n/J^{n+1} \overset{\sim}{\leftarrow} J_0^n/J_0^{n+1} \otimes_{A_0} A$, and J^n/J^{n+1} is free of finite type.

Proposition 1.3. Let A be a ring, E a projective A-module of finite type, $u : E \longrightarrow A$ a regular homomorphism, $J = u(E)$. Then the ideal J is quasi-regular (EGA IV 16.9.7), that is (EGA IV 16.9.8):

i) J is of finite type.

ii) J/J^2 is a projective A/J-module of finite type;

iii) the canonical homomorphism

$$\mathrm{Sym}_{A/J}(J/J^2) \longrightarrow \mathrm{gr}_J(A) = \bigoplus_{n \geq 0} J^n/J^{n+1}, \tag{1.3.1}$$

is an isomorphism.

We saw in the proof of 1.2 b) that J/J^2 is projective of finite type. Resume the notation of 1.2 b) (since the question is local on $\mathrm{Spec}(A)$, we again suppose E free of finite type). Then the canonical homomorphism (1.3.1) is deduced from the analogous homomorphism for J_0 by the base change $A_0 \longrightarrow A$, by the relations 1.2. The latter is an isomorphism, so the same is true of (1.3.1).

Definition 1.4. Let $(X, \underline{A})$ be a commutatively locally ringed site, $\underline{E}$ an $\underline{A}$-Module locally free of finite type, $u : \underline{E} \longrightarrow \underline{A}$ an $\underline{A}$-linear homomorphism. We say that u is regular if the Koszul complex $\underline{K}_\bullet(u)$ defined by u is acyclic in degrees > 0. Let $\underline{J}$ be an Ideal of $\underline{A}$; we say that $\underline{J}$ is [print 419]regular, if, locally on X, there exist a free $\underline{A}$-Module of finite type $\underline{E}$, and a surjective regular homomorphism: $\underline{E} \longrightarrow \underline{J}$.

Let $i : Y \longrightarrow X$ be an immersion of preschemes, and let U be an open subset of X such that $i(Y) \subset U$ and such that i is a closed immersion of Y into U. We say that i is regular if the ideal of $\underline{O}_U$ defining $i(Y)$ is regular (a condition independent of U). If $y \in Y$, we say that i is regular at y if there exists a neighborhood V of $i(y)$ such that the immersion $i^{-1}(V) \longrightarrow V$ is regular. For an immersion to be regular, it is therefore necessary and sufficient that it be regular in a neighborhood of each point.

This definition conflicts with EGA IV 16.9.2, and replaces that definition. However, in the noetherian case the two definitions are equivalent (EGA IV 19.5.1). An immersion that can locally be defined by a regular sequence is regular; a regular immersion is quasi-regular (1.3).

1.4.1. Let X be a scheme, $\underline{J}$ an Ideal of finite type of $\underline{O}_X$, x a point of $\mathrm{Supp}(\underline{O}_X/\underline{J})$. We shall say that a family of sections $f_1, \ldots, f_d$ of $\underline{J}$ on a neighborhood of x forms a minimum system of

generators of $\underline{J}$ in a neighborhood of x if the $(f_i)_x$ form a family of generators of $\underline{J}_x$ of minimum cardinality, i.e. their images in $\underline{J}_x \otimes k(x)$ form a basis of this vector space over $k(x)$. Note that the f_i generate $\underline{J}$ on a neighborhood of x.

1.4.2. To say that an immersion is regular is also to say that it is defined in some open subset of X by an ideal of finite type $\underline{J}$, and that for every point $x \in \mathrm{Supp}(\underline{O}_X/\underline{J})$ and every minimum system of generators of $\underline{J}$ in a neighborhood of x, the Koszul complex associated with this sequence is acyclic in a neighborhood of x: indeed, let $f_1, \ldots, f_d$ be a minimum system of [print 420]generators of $\underline{J}$ in a neighborhood of x such that the associated Koszul complex is acyclic (one exists when the immersion is assumed regular); let $g_1, \ldots, g_d$ be another minimum system of generators of $\underline{J}$; there exists a neighborhood of x on which $\underline{J}$ can be generated either by the f_i or by the g_j[392], and consequently there exists on this neighborhood a matrix with coefficients in $\underline{O}_X$ expressing the g_j in terms of the f_i. Now the images of the f_i (resp. g_j) in $\underline{J}/\underline{J}^2$ form a basis at the point x, hence in a neighborhood of x, and consequently this matrix is invertible in a neighborhood of x. It therefore defines an isomorphism from the Koszul complex defined by the g_j onto the Koszul complex defined by the f_i, which shows that the Koszul complex defined by the g_j is acyclic.

1.4.3. Assuming only that $\underline{J}$ is of finite type and $\underline{J}/\underline{J}^2$ is locally free of finite type, note that every minimum system of generators of $\underline{J}$ in a neighborhood of a point y of Y gives a basis of $\underline{J}/\underline{J}^2$ in a neighborhood of y. Conversely, given a basis of $\underline{J}/\underline{J}^2$ in a neighborhood of y, we can lift it in a neighborhood of y to a family of sections of $\underline{J}$, which generate $\underline{J}_y$ by Nakayama, hence $\underline{J}$ in a neighborhood of y since[393] $\underline{J}$ is of finite type; moreover these generators have minimum cardinality, since they give a basis of $\underline{J}/\underline{J}^2$.

Proposition 1.5. (Compare EGA IV 19.1.5 (ii)). Let $i : Y \longrightarrow X$ be an immersion. If i is regular, the same is true of every immersion deduced from i by flat base change. For i to be regular, it is necessary and sufficient that it be so locally for the faithfully flat quasi-compact topology.

Stability under flat base change is evident.

Conversely, let i' be the immersion deduced from i by a [print 421]faithfully flat quasi-compact base change. If i' is regular, then the Ideal $\underline{J}$ defining i is of finite type, and $\underline{J}/\underline{J}^2$ is locally free of finite type, since these properties descend under faithfully flat quasi-compact morphisms. If $f_1, \ldots, f_d$ is a sequence of sections of $\underline{J}$ lifting a basis of $\underline{J}/\underline{J}^2$, the Koszul complex associated with the inverse images of the f_i is acyclic in degrees > 0 by 1.4; since it is the inverse image of the Koszul complex associated with the f_i, the latter is acyclic in degrees > 0, and i is regular.

Definition 1.6. Let $i : Y \longrightarrow X$ be an immersion defined in an open subset U of X by an Ideal $\underline{J}$ of $\underline{O}_X$; the $\underline{O}_Y$-Module $\underline{J}/\underline{J}^2$ (independent of U) is called the conormal sheaf of Y in X, and is denoted by $\underline{N}_{X/Y}$. If i is regular, $\underline{J}/\underline{J}^2$ is locally free of finite type, and its rank is called the codimension of Y in X, or the codimension of i.

Given two closed immersions: $Z \xrightarrow{j} Y \xrightarrow{i} X$, the conormal sheaves form an exact sequence:

$$j^*(\underline{N}_{X/Y}) \longrightarrow \underline{N}_{X/Z} \longrightarrow \underline{N}_{Y/Z} \longrightarrow 0. \tag{1.6.1}$$

Indeed, let $\underline{I}$ be the ideal of Y in X, $\underline{J}$ the ideal of Z in X, $\underline{K}$ the ideal of Z in Y. We have the exact sequence:

$$0 \longrightarrow \underline{I} \longrightarrow \underline{J} \longrightarrow \underline{K} \longrightarrow 0;$$

whence, by tensoring with $\underline{O}_X/\underline{I} = \underline{O}_Y$, the exact sequence:

$$\underline{I}/\underline{I}^2 \longrightarrow \underline{J} \otimes_{\underline{O}_X} \underline{O}_Y \longrightarrow \underline{K} \otimes_{\underline{O}_X} \underline{O}_Y \longrightarrow 0;$$

tensoring with $\underline{O}_Z = \underline{O}_X/\underline{J} = \underline{O}_Y/\underline{K}$, we obtain the exact sequence:

$$j^*(\underline{I}/\underline{I}^2) \longrightarrow \underline{J} \otimes_{\underline{O}_X} \underline{O}_Z \longrightarrow \underline{K} \otimes_{\underline{O}_Y} \underline{O}_Z \longrightarrow 0,$$

which is the required sequence.

[print 422]**Proposition 1.7.** (Compare EGA IV 19.1.5 (iv)). Let $i : Y \longrightarrow X$, and $j : Z \longrightarrow Y$ be two immersions:

i) If i and j are regular, $i \circ j$ is regular, and the sequence:

$$0 \longrightarrow j^*(\underline{N}_{X/Y}) \longrightarrow \underline{N}_{X/Z} \longrightarrow \underline{N}_{Y/Z} \longrightarrow 0$$

is exact; in particular, $\operatorname{codim}(i \circ j) = \operatorname{codim}(i) + \operatorname{codim}(j)$.

ii) If $i \circ j$ and i are regular, and if the sequence:

$$0 \longrightarrow j^*(\underline{N}_{X/Y}) \longrightarrow \underline{N}_{X/Z} \longrightarrow \underline{N}_{Y/Z} \longrightarrow 0,$$

is exact and locally split, j is regular;

iii) If X is noetherian, and if $i \circ j$ and j are regular, i is regular at the points of Z.

Since the question is local on X, we may suppose X affine and the immersions closed; let $X = \operatorname{Spec}(A)$, $Y = \operatorname{Spec}(B)$, $Z = \operatorname{Spec}(C)$, with $B = A/I$, $C = A/J = B/K$.

i) After localizing, we may suppose I/I^2 free over B, and K/K^2 free over C (1.3). Let $f_1, \ldots, f_n$ be elements of A lifting a basis of I/I^2, $g_1, \ldots, g_m$ elements of A lifting a basis of K/K^2; then, after further localization, we may suppose that $f_1, \ldots, f_n$ generate I, that $f_1, \ldots, f_n, g_1, \ldots, g_m$ generate J, and that $\bar{g}_1, \ldots, \bar{g}_m$ generate K, where $\bar{g}_j$ is the image of g_j in B (1.4); moreover, $f_1, \ldots, f_n$ form a minimum system of generators of I, and $\bar{g}_1, \ldots, \bar{g}_m$ a minimum system of generators of K.

Consider the diagram:

$$\begin{array}{ccc} A_0 = \mathbb{Z}[T_1, \ldots, T_{n+m}] & \longrightarrow & A \\ \downarrow & & \downarrow \\ B_0 = \mathbb{Z}[T_1, \ldots, T_m] & \longrightarrow & B \\ \downarrow & & \downarrow \\ C_0 = \mathbb{Z} & \longrightarrow & C, \end{array} \tag{1.7.1}$$

[print 423]where the horizontal arrows are defined by sending T_i to f_i for $1 \leq i \leq n$, and T_{n+j} to g_j for $1 \leq j \leq m$, where the homomorphism $A_0 \longrightarrow B_0$ is defined by sending T_i to 0 for $1 \leq i \leq n$, and T_{n+j} to T_j for $1 \leq j \leq m$, and the homomorphism $B_0 \longrightarrow C_0$ by setting the T_j, $1 \leq j \leq m$, to zero. We then have:

$$B = A \otimes_{A_0} B_0, \qquad C = A \otimes_{A_0} C_0 = B \otimes_{B_0} C_0.$$

The situation under consideration is therefore deduced from the situation (A_0, B_0, C_0) by the base change $A_0 \longrightarrow A$. The corresponding immersions i_0, j_0 and $i_0 \circ j_0$ are regular; consequently, the homology groups of the Koszul complexes associated with the sequences $(f_1, \ldots, f_n)$, $(f_1, \ldots, f_n, g_1, \ldots, g_m)$ and $(\bar{g}_1, \ldots, \bar{g}_m)$ are respectively the $\operatorname{Tor}_i^{A_0}(B_0, A)$, $\operatorname{Tor}_i^{A_0}(C_0, A)$, $\operatorname{Tor}_i^{B_0}(C_0, B)$[394]. Consequently:

$$\begin{aligned} i \text{ regular} &\iff \forall i \geq 1, \operatorname{Tor}_i^{A_0}(B_0, A) = 0; \\ i \circ j \text{ regular} &\iff \forall i \geq 1, \operatorname{Tor}_i^{A_0}(C_0, A) = 0; \\ j \text{ regular} &\iff \forall i \geq 1, \operatorname{Tor}_i^{B_0}(C_0, B) = 0. \end{aligned} \tag{1.7.2}$$

We then use the transitivity spectral sequence ([1], XVI 5)

$$E^2_{pq} = \mathrm{Tor}_p^{B_0}(C_0, \mathrm{Tor}_q^{A_0}(B_0, A)) \Longrightarrow \mathrm{Tor}_n^{A_0}(C_0, A).. \tag{1.7.3}$$

By the hypotheses of i), $E^2_{pq} = 0$ for p or $q \neq 0$. It therefore degenerates, and $i \circ j$ is regular by (1.7.2).

By (1.6.1), the sequence:

$$j^*(\underline{N}_{X/Y}) \longrightarrow \underline{N}_{X/Z} \longrightarrow \underline{N}_{Y/Z} \longrightarrow 0$$

is exact. Since $i \circ j$ is regular, the three conormal sheaves are locally free of finite type; moreover, their rank at a point is the [print 424]number of elements in a minimum system of generators of the corresponding ideal; by the preceding proof, we have

$$\mathrm{rang}(\underline{N}_{X/Z}) = \mathrm{rang}(\underline{N}_{X/Y}) + \mathrm{rang}(\underline{N}_{Y/Z}).$$

It follows that the kernel of $j^*(\underline{N}_{X/Y}) \longrightarrow \underline{N}_{X/Z}$ is locally free of rank zero, and hence that the sequence (1.6.1) is exact.[395]

ii) We may suppose J/J^2 free over C, and I/I^2 free over B; moreover, the hypothesis on the sequence of conormal sheaves allows us to suppose that $I/I^2 \otimes_A A/J$ is a direct summand of J/J^2. We can therefore find elements $f_1, \ldots, f_n$ of A lifting a basis of I/I^2, and elements $g_1, \ldots, g_m$ of A such that $f_1, \ldots, f_n, g_1, \ldots, g_m$ lift a basis of J/J^2, the images of $g_1, \ldots, g_m$ generating K/K^2; after further localization, we may suppose that $f_1, \ldots, f_n$ form a minimum system of generators of I, that $f_1, \ldots, f_n, g_1, \ldots, g_m$ form a minimum system of generators of J, and that $\bar{g}_1, \ldots, \bar{g}_m$ generate K. We can then construct the diagram (1.7.1) and use the spectral sequence (1.7.3).

By (1.7.2), since i is regular, we have $E^2_{pq} = 0$ for $q \geq 1$; we therefore obtain edge-isomorphisms: $E^2_{p,0} \simeq \mathrm{Tor}_p^{A_0}(C_0, A)$. Since $i \circ j$ is regular, this implies by (1.7.2) that $E^2_{p,0} = 0$ for $p \geq 1$, and hence that j is regular, again by (1.7.2).

iii) This last property is proved in EGA IV 19.1.5 (iv), since in the noetherian case the definition in EGA IV coincides with the one we adopt here.

Proposition 1.8. Let X be a scheme, $\underline{E}$ an $\underline{O}_X$-Module locally free of finite type, $u : \underline{E} \to \underline{O}_X$ a regular homomorphism, $\underline{J} = u(\underline{E})$. Denote by Y [print 425]the closed subscheme of X defined by $\underline{J}$, X' the scheme obtained by blowing up Y, P the projective bundle associated with $\underline{E}$, p the structural morphism $P \to X$. On P, consider the canonical exact sequence:

$$0 \to \underline{F} \to p^*(\underline{E}) \to \underline{O}_P(1) \to 0,$$

and the composite morphism:

$$v : \underline{F} \longrightarrow p^*(\underline{E}) \xrightarrow{p^*(u)} \underline{O}_P.$$

Then:

i) The formation of X' commutes with every base change $f : T \to X$ such that $f^*(u)$ is regular.

ii) The closed immersion $j : X' \to P$ defined by the surjective homomorphism $\mathbf{S}(\underline{E}) \longrightarrow \bigoplus_{n \geq 0} \underline{J}^n$[396] defined by u in degree 1 is a regular immersion, its ideal being the image of the regular morphism v; in particular, if $\mathrm{codim}(i)$ is constant, $\mathrm{codim}(j) = \mathrm{codim}(i) - 1$.

iii) The $\mathbf{S}(\underline{E})$-Module[397] $\bigoplus_{n\geq 0}\underline{J}^n$ is perfect as a graded $\mathbf{S}(\underline{E})$-Module[398], i.e. locally there exists a finite, graded resolution of $\bigoplus_{n\geq 0}\underline{J}^n$ by graded $\mathbf{S}(\underline{E})$-Modules locally free of finite type[399] (VI 1.2).

i) Since $X' = \mathrm{Proj}(\bigoplus_{n\geq 0}\underline{J}^n)$, the result is an immediate consequence of the relations (1.2.3) and of EGA II.

ii) Since the stated properties are local on X, we may suppose X affine, and moreover $\underline{E}$ free over X. Put $X = \mathrm{Spec}(A)$.

First we give the proof in the standard case $A = \mathbb{Z}[T_1, \ldots, T_d]$, with E endowed with a basis $e_1, \ldots, e_d$ such that $u(e_i) = T_i$. Then $P = \mathrm{Proj}(\mathbb{Z}[T_1, \ldots, T_d, X_1, \ldots, X_d])$, the grading being taken with respect to the X_i. To show that v is a regular morphism whose image is the Ideal of j, it suffices to do so locally on P, hence on the [print 426]$D^+(X_i)$; let us work, for example, on $D^+(X_1)$. Then $p^*(\underline{E})$ is defined by the module $\mathbb{Z}[T_1, \ldots, T_d, X_2/X_1, \ldots, X_d/X_1] \otimes E$, and $\underline{O}_P(1)$ by $\mathbb{Z}[T_1, \ldots, T_d, X_2/X_1, \ldots, X_d/X_1]$, the canonical morphism sending e_i to X_i/X_1 for every i. Then F has as a basis the elements $e_i - e_1.X_i/X_1$ for $i > 1$.

We have $v(e_i - e_1.X_i/X_1) = T_i - T_1.X_i/X_1$; put $T'_i = T_i - T_1X_i/X_1$. Since the sequence of the T_i is regular in $\mathbb{Z}[T_1, \ldots, T_d, X_2/X_1, \ldots, X_d/X_1]$, the same is true of the sequence $T_1, T'_2, \ldots, T'_d$ deduced from it by an invertible matrix, and hence also of the subsequence $T'_2, \ldots, T'_d$ (we are in a noetherian case). Consequently, v is a regular morphism.

We must therefore show that the T'_i generate the Ideal of j. In $\bigoplus_{n\geq 0} J^n$, for $m \leq p$, we shall denote by $x^{(m)}$ the element x of J^p, considered as having degree m. Over $D^+(X_1)$, $X' = \mathrm{Spec}\left(\left(\bigoplus_{n\geq 0} J^n\right)_{(T_1^{(1)})}\right)$, and j is defined by the morphism sending X_i to $T_i^{(1)}$. Under this morphism, the image of T'_i is $T_i^{(0)} - T_1^{(0)}.T_i^{(1)}/T_1^{(1)} = \left((T_iT_1)^{(1)} - (T_1T_i)^{(1)}\right)/T_1^{(1)} = 0$. Thus the T'_i belong to the Ideal of j.

Since every element of $\mathbb{Z}[T_1, \ldots, T_d, X_2/X_1, \ldots, X_d/X_1]$[400] can be expressed as a polynomial in $T_1, T'_2, \ldots, T'_d, X_2/X_1, \ldots, X_d/X_1$, to show that the T'_i generate the Ideal of j it suffices to show that no nonzero element of $\mathbb{Z}[T_1, X_2/X_1, \ldots, X_d/X_1]$ belongs to the Ideal of j, which amounts to showing that the homomorphism

$$\mathbb{Z}[T_1, X_1, \ldots, X_d] \longrightarrow \bigoplus_{n\geq 0} J^n$$

is injective, since $T_1^{(1)}$ is not a zero divisor in $\bigoplus_{n\geq 0} J^n$. Grading $\mathbb{Z}[T_1, X_1, \ldots, X_d]$ by assigning degree 0 to T_1 and degree 1 to the X_i, the kernel of the homomorphism under consideration is a graded ideal. Now it is clear that [print 427]this kernel is contained in that of the composite homomorphism:

$$\mathbb{Z}[T_1, X_1, \ldots, X_d] \longrightarrow \bigoplus_{n\geq 0} J^n \longrightarrow A,$$

the last homomorphism being defined in degree n by the inclusion of J^n in A. Now the kernel of the composite homomorphism is the principal ideal generated by $(X_1 - T_1)$, which, for reasons of integrality, contains no nonzero homogeneous elements. Thus the first homomorphism is injective, completing the proof in this case.

Return to the general case where A is arbitrary; we may still suppose $\underline{E}$ free over X. Attach the index 0 to the special case just treated; then, by part i) of the proposition, the situation under consideration is obtained by the base change $A_0 \to A$ from the situation studied above, the homomorphism $A_0 \to A$ being defined by taking a basis of E, its image $f_1, \ldots, f_d$ in J, and sending T_i to f_i. Put $B_0 = \left(\bigoplus_{n\geq 0} J_0^n\right)_{(T_1^{(1)})}$, $B = \left(\bigoplus_{n\geq 0} J^n\right)_{(f_1^{(1)})}$; then $B = B_0 \otimes_{A_0} A$; the Koszul complex defined by $v : \underline{F} \to \underline{O}_P$ is deduced by base change from the Koszul complex

defined by v_0; since the latter is a projective resolution of B_0 (working on a suitable open subset), the homology groups of $K_\bullet(v)$ on this open subset are the $\mathrm{Tor}_i^{A_0}(A, B_0)$. These are the degree 0 components of the $\mathrm{Tor}_i^{A_0}\left(A, \left(\bigoplus_{n\geq 0} J_0^n\right)_{T_1^{(1)}}\right) = \left(\bigoplus_{n\geq 0} \mathrm{Tor}_i^{A_0}(A, J_0^n)\right)_{T_1^{(1)}}$; the $\mathrm{Tor}_i^{A_0}(A, J_0^n)$ vanish for $i \geq 1$ by the relations 1.2, so $K_\bullet(v)$ is a resolution of B, which shows that j is regular.

iii) Since the property is local on X, we may again suppose X affine and $\underline{E}$ free of finite type. Resume the notation of ii).

First consider the standard case of ii). The ring A_0 is then noetherian and of finite cohomological dimension, and the algebra $\mathbf{S}(E_0)$[401] [print 428]likewise. We can then construct step by step a partial graded resolution of $\bigoplus_{n\geq 0} J_0^n$ by graded free $\mathbf{S}(E_0)$-modules[402] of finite type, of length greater than the cohomological dimension of A_0; the last kernel is then a graded $\mathbf{S}(E_0)$-module[403], projective of finite type as an ungraded $\mathbf{S}(E_0)$-module[404], hence locally free of finite type over $\mathrm{Spec}(A_0)$ as a graded $\mathbf{S}(\underline{E}_0)$-module[405] by VI 1.5.1. We obtain the announced finite resolution by replacing the last module of the partial resolution by the corresponding kernel.

Tensoring the resolution thus constructed with A, we obtain a graded complex of graded $\mathbf{S}(E)$-modules[406] locally free of finite type, and must show that it is a resolution of $\bigoplus_{n\geq 0} J^n$. The i-th homology Module of the resulting complex is a graded $\mathbf{S}(E)$-Module whose[407] n-th component is $\mathrm{Tor}_i^{A_0}(A, J_0^n)$, which vanishes for $i \geq 1$ by (1.2.4). Consequently, by (1.2.3), we have constructed a resolution of the required type.

Proposition 1.9. Let $i : Y \to X$ be a regular closed immersion, X' the X-scheme obtained by blowing up Y, and $f : X' \to X$ the structural morphism. Then i and f are perfect morphisms. (III 4.1.).

Since the property is local on X, we may suppose given an $\underline{O}_X$-Module locally free of finite type $\underline{E}$, and a regular homomorphism $u : \underline{E} \to \underline{O}_X$ whose image is the ideal $\underline{J}$ of Y. The Koszul complex $\underline{K}_\bullet(u)$ is then a finite resolution of $\underline{O}_Y$ by $\underline{O}_X$-Modules locally free of finite type; consequently, $\underline{O}_Y$ is a perfect $\underline{O}_X$-Module. It follows that every regular immersion is a perfect morphism. By 1.8, f can be written as the composite of a regular immersion $X' \to \mathbf{P}(\underline{E})$[408] and the structural morphism $\mathbf{P}(\underline{E}) \to X$[409], which is smooth and hence perfect; it is therefore [print 429]the composite of two perfect morphisms, and consequently a perfect morphism.

Proposition 1.10. Let X, Y be two schemes smooth over S. Then every S-immersion of X into Y is regular. In particular, every section of a smooth morphism is a regular immersion.

The proposition follows immediately from EGA IV 17.12.1, since a regular immersion in the sense of EGA IV is regular in the sense of 1.4.

2. Computations on regular immersions

2.1.

Let A be a ring, and B and C two A-algebras. We propose to endow $\bigoplus_{i\geq 0} \mathrm{Tor}_i^A(B, C)$ with a graded A-algebra structure, and hence to define homomorphisms:

$$\mathrm{Tor}_i^A(B, C) \otimes_A \mathrm{Tor}_j^A(B, C) \longrightarrow \mathrm{Tor}_{i+j}^A(B, C). \tag{2.1.1}$$

Let $P_\bullet$ be a resolution of B by projective A-modules; the $\mathrm{Tor}_i^A(B, C)$ are the homology groups of the complex $P_\bullet \otimes_A C$. We know how to define canonical homomorphisms ([1], IV 6):

$$H_i(P_\bullet \otimes_A C) \otimes_A H_j(P_\bullet \otimes_A C) \longrightarrow H_{i+j}((P_\bullet \otimes_A C) \otimes_A (P_\bullet \otimes_A C)). \tag{2.1.2}$$

Furthermore, $P_\bullet \otimes_A P_\bullet$ is a complex of projective A-modules, in degrees ≥ 0, and $H_0(P_\bullet \otimes_A P_\bullet) = B \otimes_A B$; moreover, $P_\bullet$ is a resolution of B. Consequently, the augmentation morphism $B \otimes_A B \to B$ extends to a morphism of complexes: $P_\bullet \otimes_A P_\bullet \to P_\bullet$ and therefore defines, using the homomorphism $C \otimes_A C \to C$ deduced from the algebra structure of C, homomorphisms:

$$H_{i+j}\big((P_\bullet \otimes_A C) \otimes_A (P_\bullet \otimes_A C)\big) \longrightarrow H_{i+j}(P_\bullet \otimes_A C). \tag{2.1.3}$$

[print 430]Composing (2.1.2) and (2.1.3), we obtain the homomorphisms (2.1.1).

The algebra structure defined by the homomorphisms (2.1.1) is associative (cf. [1] XI 2). Furthermore, the A-module $H_i(P_\bullet \otimes_A C)$ has a standard $B \otimes_A C$-module structure, with C acting in the evident way, and B by functoriality; likewise, the $H_j\big((P_\bullet \otimes_A C) \otimes_A (P_\bullet \otimes_A C)\big)$ carry two $B \otimes_A C$-module structures. The homomorphisms (2.1.2) and (2.1.3) are clearly compatible with these two structures: for (2.1.2), by functoriality (working in the category of A-modules), and for (2.1.3), by the classical result on morphisms of an acyclic complex. We have therefore endowed $\bigoplus_{i\geq 0} \operatorname{Tor}_i^A(B, C)$ with an associative $B \otimes_A C$-algebra structure.

2.2.

Suppose that B admits a resolution by projective A-modules $P_\bullet$ such that $P_\bullet = \bigoplus_{i\geq 0} P_i$ is a graded A-algebra, with the differential satisfying:

a) $P_0 \longrightarrow B$ is a ring homomorphism ;([1])
b) $\forall x \in P_i,\ \forall y \in P_j, \quad d(xy) = d(x)y + (-1)^i x\, d(y)$.

Then the "product" maps define commutative diagrams for $i, j \geq 0$:

$$\begin{array}{ccc} P_i \otimes_A P_j & \longrightarrow & (P_{i-1} \otimes_A P_j) \oplus (P_i \otimes_A P_{j-1}) \\ \downarrow & & \downarrow \\ P_{i+j} & \longrightarrow & P_{i+j-1}, \end{array}$$

and consequently a morphism of complexes $P_\bullet \otimes_A P_\bullet \to P_\bullet$ extending the augmentation homomorphism: $B \otimes_A B \to B$. This shows that the algebra structure of $\bigoplus_{i\geq 0} \operatorname{Tor}_i^A(B, C)$ can be computed from that of $P_\bullet$. In particular, if $P_\bullet$ is an alternating A-algebra (i.e. such that for every [print 431]$x \in P_\bullet$, we have $x^2 = 0$), then the same is true of $\bigoplus_{i\geq 0} \operatorname{Tor}_i^A(B, C)$.

This last remark applies when B is the quotient of A by a regular ideal, since we can then take for $P_\bullet$ the Koszul complex associated with a regular homomorphism defining the ideal of B.

2.3.

The $B \otimes_A C$-algebra structure defined in 2.1 on $\bigoplus_{i\geq 0} \operatorname{Tor}_i^A(B, C)$ is functorial in B and C. We can therefore globalize the preceding constructions: let X be a scheme, X', X'' two X-schemes; we can then define the morphisms (2.1.1) locally on $X' \times_X X''$; they glue by functoriality and define on $\bigoplus_{i\geq 0} \underline{\operatorname{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_{X''})$ an associative graded $\underline{O}_{X'\times_X X''}$-Algebra structure.

Lemma 2.4. Let X', X'' be two X-schemes, and $\underline{M}$ a flat $\underline{O}_{X'}$-Module. Then for every $i \geq 0$, we have a canonical isomorphism:

$$\underline{\operatorname{Tor}}_i^{\underline{O}_X}(\underline{M}, \underline{O}_{X''}) \simeq \underline{M} \otimes_{\underline{O}_{X'}} \underline{\operatorname{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_{X''}).$$

([1]) One can show that such a resolution always exists.

It suffices to define a functorial isomorphism in the affine case; we can then define it globally by gluing in the general case.

Let $X = \mathrm{Spec}(A)$, $X' = \mathrm{Spec}(A')$, $X'' = \mathrm{Spec}(A'')$; $\underline{M}$ is defined by a flat A'-module M. Let $P_\bullet$[410] be a projective resolution of A'' over A; the $\mathrm{Tor}_i^A(A', A'')$ are the homology groups of $A' \otimes_A P_\bullet$. Since M is flat over A', we have a canonical isomorphism: $M \otimes_{A'} \mathrm{Tor}_i^A(A', A'') \xrightarrow{\sim} H_i(M \otimes_{A'} (A' \otimes_A P_\bullet))$. Furthermore, the canonical isomorphism: $M \otimes_{A'} (A' \otimes_A P_\bullet) \xrightarrow{\sim} M \otimes_A P_\bullet$ gives a canonical isomorphism: $H_i(M \otimes_{A'} (A' \otimes_A P_\bullet)) \xrightarrow{\sim} \mathrm{Tor}_i^A(M, A'')$, whence the result.

Proposition 2.5. Let X be a scheme, Y and Z two closed subschemes of X, $T = Y \times_X Z$. Suppose that for every $x \in T$, there exists an open neighborhood [print 432]U of x, and a sequence of sections $f_1, \ldots, f_p, g_1, \ldots, g_r, h_1, \ldots, h_s$ of $\underline{O}_U$ such that the Koszul complex associated with $f_1, \ldots, f_p, g_1, \ldots, g_r$ is a resolution of $\underline{O}_{Y \cap U}$, the Koszul complex associated with $f_1, \ldots, f_p, h_1, \ldots, h_s$ a resolution of $\underline{O}_{Z \cap U}$, and the Koszul complex associated with $f_1, \ldots, f_p, g_1, \ldots, g_r, h_1, \ldots, h_s$ a resolution of $\underline{O}_{T \cap U}$ (conditions satisfied in particular in the noetherian case when the sequence $f_1, \ldots, f_p, g_1, \ldots, g_r, h_1, \ldots, h_s$[411] is regular and $Y = V(f_1, \ldots, f_p, g_1, \ldots, g_r)$, $Z = V(f_1, \ldots, f_p, h_1, \ldots, h_s)$). Let $\underline{I}, \underline{J}$ be the ideals of Y, Z. We then have the following isomorphisms:

$$\text{i)} \quad \forall i \geq 0, \quad \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \bigwedge^i \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z);$$

$$\text{ii)} \quad \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \simeq \underline{I} \cap \underline{J}/\underline{I}.\underline{J} \simeq \mathrm{Ker}(i^*(\underline{N}_{Y/X}) \times j^*(\underline{N}_{Z/X}) \to \underline{N}_{T/X}),$$

where we denote by i the immersion of T into Y and by j the immersion of T into Z.

In particular, for every regular closed immersion $Y \to X$, we have

$$\text{i')} \quad \forall i \geq 0, \quad \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Y) \xrightarrow{\sim} \bigwedge^i \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Y);$$

$$\text{ii')} \quad \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Y) \simeq \underline{N}_{Y/X}.$$

i) We saw in 2.3 that $\bigoplus_{i \geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$ could be endowed with an $\underline{O}_T$-Algebra structure; by 2.2, this algebra is alternating, since the immersion of Y (or Z) into X is regular. There therefore exists one and only one algebra homomorphism:

$$\bigwedge \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \longrightarrow \bigoplus_{i \geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z), \tag{2.5.1}$$

extending the identity in degree 1.

To see that it is an isomorphism, we can check locally. Thus we work in a neighborhood of a point of T and consider sections $f_1, \ldots, f_p, g_1, \ldots, g_r, h_1, \ldots, h_s$ of $\underline{O}_X$ on this neighborhood, having the properties stated in 2.5. Put $\underline{E} = (\underline{O}_X)^p$, $\underline{E}' = (\underline{O}_X)^r$, and let $u : \underline{E} \to \underline{O}_X$ be the homomorphism defined by the f_i, $v : \underline{E}' \to \underline{O}_X$ the homomorphism defined by the g_j, and $\underline{K}_\bullet(u)$ and $\underline{K}_\bullet(v)$ the associated [print 433]Koszul complexes. The Koszul complex associated with the sequence $f_1, \ldots, f_p, g_1, \ldots, g_r$ is isomorphic to the tensor-product complex $\underline{K}_\bullet(u) \otimes_{\underline{O}_X} \underline{K}_\bullet(v)$. Since it is a resolution of $\underline{O}_Y$ by free $\underline{O}_X$-Modules of finite type, the $\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$ are the homology sheaves of the complex $\underline{K}_\bullet(u) \otimes_{\underline{O}_X} \underline{K}_\bullet(v) \otimes_{\underline{O}_X} \underline{O}_Z$. We can regard it as the tensor product of the complexes $\underline{K}_\bullet(u)$ and $\underline{K}_\bullet(v) \otimes_{\underline{O}_X} \underline{O}_Z$, and compute its homology using the spectral sequences associated with a bicomplex (EGA 0_{III} (11.3.5)), which abut to the homology of the associated simple complex.

Consider the spectral sequence such that ${}'E_{pq}^2 = H_p^I(H_q^{II}(K_{\bullet\bullet}))$, in the notation of EGA 0_{III} (11.3.5), with the first index referring to the complex $\underline{K}_\bullet(u)$ and the second to the complex $\underline{K}_\bullet(v) \otimes_{\underline{O}_X} \underline{O}_Z$. Now the complex $\underline{K}_\bullet(v) \otimes_{\underline{O}_X} \underline{O}_Z$ is the Koszul complex over $\underline{O}_Z$ associated with the sequence of sections $\overline{g}_1, \ldots, \overline{g}_r$, where the $\overline{g}_i$ are the images of the g_i in $\underline{O}_Z$, and this complex

is a resolution of $\underline{O}_T$ over $\underline{O}_Z$ since the immersions $Z \to X$, and $T \to Z \to X$ are regular (cf. the proof of 1.7 ii). Since $\bigwedge^p \underline{E}$ is free over $\underline{O}_X$, the complex $\bigwedge^p \underline{E} \otimes_{\underline{O}_X} (\underline{K}_\bullet(v) \otimes_{\underline{O}_X} \underline{O}_Z)$ is a resolution of $\bigwedge^p \underline{E} \otimes_{\underline{O}_X} \underline{O}_T$. Consequently the spectral sequence degenerates, and the $\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$ are the homology sheaves of the complex $\underline{K}_\bullet(u) \otimes_{\underline{O}_X} \underline{O}_T$. The differential of $\underline{K}_\bullet(u)$ is expressed in terms of the f_i, which annihilate $\underline{O}_T$; consequently the differential of $\underline{K}_\bullet(u) \otimes_{\underline{O}_X} \underline{O}_T$ is zero, and we therefore obtain the isomorphisms:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \underline{O}_T \otimes_{\underline{O}_X} \bigwedge^i \underline{E} \longrightarrow \bigwedge^i (\underline{O}_T)^p. \tag{2.5.2}$$

In particular, for $i = 1$ we find: [412]

$$\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} (\underline{O}_T)^p,$$

whence for every i: [413]

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \bigwedge^i \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z). \tag{2.5.3}$$

[print 434]Now it follows from 2.2 that the isomorphism: [414]

$$\bigwedge (\underline{O}_T)^p \xrightarrow{\sim} \bigoplus_{i \geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$$

defined by the isomorphisms (2.5.2) is compatible with the algebra structures of the two sides; hence the same is true of the isomorphism: [415]

$$\bigwedge \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \bigoplus_{i \geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$$

defined by the isomorphisms (2.5.3). By uniqueness, the latter is therefore the restriction of the homomorphism (2.5.1) that was defined globally. The homomorphism (2.5.1) is thus an isomorphism.

ii) Consider the exact sequence:

$$0 \longrightarrow \underline{I} \longrightarrow \underline{O}_X \longrightarrow \underline{O}_Y \longrightarrow 0,$$

and the resulting exact $\underline{\mathrm{Tor}}$ sequence:

$$0 \longrightarrow \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \longrightarrow \underline{I} \otimes_{\underline{O}_X} \underline{O}_Z \longrightarrow \underline{O}_Z.$$

It follows that $\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$ is the kernel of the homomorphism:

$$\underline{I} \otimes_{\underline{O}_X} (\underline{O}_X/\underline{J}) \longrightarrow \underline{O}_X/J,$$

that is, $\underline{I} \cap \underline{J}/\underline{I}.\underline{J}$.

To prove the last relation, we shall show that there is an exact sequence:

$$\underline{0} \longrightarrow \underline{I} \cap \underline{J}/\underline{I}.\underline{J} \longrightarrow i^*(\underline{N}_{Y/X}) \times j^*(\underline{N}_{Z/X}) \longrightarrow \underline{N}_{T/X} \longrightarrow 0.$$

The homomorphisms $i^*(\underline{N}_{Y/X}) \to \underline{N}_{T/X}$ and $j^*(\underline{N}_{Z/X}) \to \underline{N}_{T/X}$ were defined in (1.6.1); here we take their difference. Furthermore, $\underline{I} \cap \underline{J}$ maps naturally into $\underline{I}/\underline{I}^2$, hence into $i^*(\underline{I}/\underline{I}^2)$, and likewise into $j^*(\underline{J}/\underline{J}^2)$, and these homomorphisms vanish on $\underline{I}.\underline{J}$, since $\underline{I}, \underline{J} \subset \underline{K}$ (where $\underline{K}$ denotes the ideal of T), defining the arrow on the left.

[print 435]To check exactness of the sequence, we can work locally on T and use the sections defined in the statement of 2.5. We then have a basis of $\underline{N}_{T/X}$ formed by the images of the f_i, g_j, h_k, a basis of $i^*(\underline{N}_{Y/X})$ formed by the images of the f_i, g_j, and a basis of $j^*(\underline{N}_{Z/X})$ formed by the images of the f_i, h_k. It is then clear that $i^*(\underline{N}_{Y/X}) \times j^*(\underline{N}_{Z/X}) \to \underline{N}_{T/X}$ is surjective, and that its kernel consists of the elements whose coordinates on the $(\overline{g}_j, 0)$ and the $(0, \overline{h}_k)$ vanish, and whose coordinates on $(\overline{f}_i, 0)$ and $(0, \overline{f}_i)$ are equal for every i. Consequently the sequence is also exact in the middle. The three $\underline{O}_T$-Modules involved are locally free of finite type; since $\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$ has rank p by the proof of i), we see immediately that the sequence is exact.

2.6.

We now suppose that X is a scheme having an ample sheaf. Every closed subscheme Y of X then has an ample sheaf. We know that in this case "global resolutions exist", and that the $K^\bullet$ formed respectively with locally free sheaves of finite type and with perfect complexes are identical (II, 2.2.9). If moreover the immersion $i : Y \to X$ is regular, it is a perfect morphism, and we can define an additive homomorphism $i_* : K^\bullet(Y) \longrightarrow K^\bullet(X)$ (IV 2.12) by associating with the class of a perfect complex $\underline{L}^\bullet$ the class of the complex $\mathbb{R}i_*(\underline{L}^\bullet) = i_*(\underline{L}^\bullet)$[416] ($i_*$ being an exact functor).

Proposition 2.7. Let X be a scheme having an ample sheaf, $i : Y \to X$ a regular closed immersion, $\underline{N}$ the conormal sheaf of Y in X. Then, for every $x \in K^\bullet(Y)$, we have the relation:

$$i^*(i_*(x)) = x.\lambda_{-1}(N),$$

where N is the class of $\underline{N}$ in $K^\bullet(Y)$, and $\lambda_{-1}(N) = \sum_{i \geq 0}(-1)^i \lambda^i(N)$.

[print 436]Since both sides of the relation to be proved are additive, we may suppose that x is the class of an $\underline{O}_Y$-Module locally free of finite type $\underline{M}$. The element $i^*(i_*(x))$ is then the class of the complex $\mathbb{L}i^*(\mathbb{R}i_*(\underline{M}))$[417], obtained by taking a projective resolution of the complex $\mathbb{R}i_*(\underline{M}) = i_*(\underline{M})$[418] and tensoring it with $\underline{O}_Y$. The homology sheaves of this complex are therefore the $\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{M}, \underline{O}_Y)$. By 2.4, we have the isomorphism:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{M}, \underline{O}_Y) \xrightarrow{\sim} \underline{M} \otimes_{\underline{O}_Y} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Y).$$

By 2.5, we have the isomorphism:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Y) \xrightarrow{\sim} \bigwedge^i \underline{N}.$$

The $\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{M}, \underline{O}_Y)$ are therefore $\underline{O}_Y$-Modules locally free of finite type, and the class in $K^\bullet(Y)$ of the complex $\mathbb{L}i^*(\mathbb{R}i_*(\underline{M}))$ is the alternating sum[419] of the classes of its homology sheaves, namely:

$$i^*(i_*(x)) = \sum_{i \geq 0}(-1)^i \,\mathrm{cl}\big(\underline{M} \otimes_{\underline{O}_Y} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Y)\big) = \sum_{i \geq 0}(-1)^i x.\lambda^i(N).$$

Corollary 2.8. Under the hypotheses of 2.7, let $x, y \in K^\bullet(Y)$; we have the relation:

$$i_*(xy\lambda_{-1}(N)) = i_*(x).i_*(y).$$

By 2.7, $y\lambda_{-1}(N) = i^*(i_*(y))$.

Applying the projection formula, we therefore find:

$$i_*(xy\lambda_{-1}(N)) = i_*(x.i^*(i_*(y))) = i_*(x).i_*(y).$$

3. Computation of $K^\bullet$ of a blown-up scheme

3.1. In this section, $i : Y \to X$ denotes a regular closed immersion. Denote by X' the scheme obtained by blowing up Y, and put $Y' = Y \times_X X'$. [print 437]We know (EGA II 8.1.8) that Y' is a closed subscheme of X' defined by an ideal of $\underline{O}_{X'}$, canonically[420] isomorphic to $\underline{O}_{X'}(1)$; since the immersion $j : Y' \to X'$[421] is defined by an invertible ideal, it is a regular closed immersion of codimension 1. Denote by $\underline{J}$ the ideal of $\underline{O}_X$ defining Y, and by $\underline{J}'$ the ideal of $\underline{O}_{X'}$ defining Y'; thus $\underline{J}' \simeq \underline{J}.\underline{O}_{X'} \simeq \underline{O}_{X'}(1)$. Finally, denote by $f : X' \to X$, and $g : Y' \to Y$ the structural morphisms. We therefore have the commutative diagram:

$$\begin{array}{ccc} Y' & \xrightarrow{j} & X' \\ {\scriptstyle g}\downarrow & & \downarrow{\scriptstyle f} \\ Y & \xrightarrow[i]{} & X. \end{array} \tag{3.1.1}$$

By definition, $X' = \mathrm{Proj}(\underline{S})$, where $\underline{S} = \bigoplus_{n\geq 0} \underline{J}^n$. Consequently, we have:

$$Y' = \mathrm{Proj}(i^*(\underline{S})) = \mathrm{Proj}\left(\left(\bigoplus_{n\geq 0} \underline{J}^n\right) \otimes_{\underline{O}_X} (\underline{O}_X/\underline{J})\right) = \mathrm{Proj}\left(\bigoplus_{n\geq 0} \underline{J}^n/\underline{J}^{n+1}\right).$$

Since i is regular, $\bigoplus_{n\geq 0} \underline{J}^n/\underline{J}^{n+1} \simeq \underline{\mathrm{Sym}}_{\underline{O}_Y}(\underline{J}/\underline{J}^2)$ (1.3). Denote by $\underline{N}$ the conormal sheaf $\underline{J}/\underline{J}^2$; thus: [422]

$$Y' = \mathbf{P}(\underline{N}).$$

On Y', we can write the exact sequence:

$$0 \to \underline{F} \to g^*(\underline{N}) \to \underline{O}_{Y'}(1) \to 0. \tag{3.1.2}$$

Observe that $\underline{O}_{Y'}(1) = j^*(\underline{O}_{X'}(1)) = \underline{J}'/\underline{J}'^2$; it is therefore the conormal sheaf of the immersion j, and we shall denote it by $\underline{L}$.

Lemma 3.2. With the hypotheses and notation of 3.1, we have the following isomorphisms:

i) $\forall i \geq 0, \quad \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \simeq \bigwedge^i \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$;

ii) $\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \simeq \underline{F}$.

[print 438]i) Since $\underline{O}_Y$ can locally be resolved by a Koszul complex, the graded algebra $\bigoplus_{i\geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$ is alternating (2.2). There therefore exists one and only one algebra homomorphism:

$$\bigwedge \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \longrightarrow \bigoplus_{i\geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y), \tag{3.2.1}$$

extending the inclusion of $\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$ in $\bigoplus_{i\geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$. It suffices to check locally that it is an isomorphism.

We can therefore work on an affine open subset $U = \mathrm{Spec}(A)$ of X, with $\underline{J}$ defined by an ideal J of A, and choose a regular homomorphism $u : E \to A$ defining J, with $E = (A)^n$. Let $f_1, \ldots, f_d$ be the images of a basis of E.

In $\bigoplus_{n\geq 0} J^n$, for $m \leq p$, we shall denote by $x^{(m)}$ the element x of J^p considered as having degree m. We can restrict to the open subset V of X' where, for example, the element $f_1^{(1)}$ is invertible.

The $\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$ are the homology sheaves of the complex $K_\bullet(u) \otimes_{\underline{O}_X} \underline{O}_{X'}$. To study this complex on V, make the following change of basis in $E' = E \otimes_A \Gamma(V, \underline{O}_{X'})$: if $e_1, \ldots, e_d$ is the given basis of E, put:

$$e'_1 = e_1; \qquad \text{for } i \neq 1, \quad e'_i = e_i - (f_i^{(1)}/f_1^{(1)})e_1.$$

Put $A' = \Gamma(V, \underline{O}_{X'})$; the differential of the complex $K_\bullet(u) \otimes_A A'$ is given by

$$d(e'_1) = f_1^{(0)}; \qquad \text{for } i \neq 1, \quad d(e'_i) = 0.$$

Taking as a basis of $\bigwedge^k(E')$ the elements $e'_{i_1} \wedge \cdots \wedge e'_{i_k}$ with $i_1 < \cdots < i_k$[423], we therefore find:

$$\begin{aligned} &d(a.e'_{i_1} \wedge \cdots \wedge e'_{i_k}) = 0 && \text{if } i_1 \neq 1; \\ &d(a.e'_{i_1} \wedge \cdots \wedge e'_{i_k}) = a f_1^{(0)}.e'_{i_2} \wedge \cdots \wedge e'_{i_k} && \text{if } i_1 = 1. \end{aligned}$$

[print 439]Denoting by E'' the submodule of E' generated by $e'_2, \ldots, e'_d$, we therefore see that the image of $d : \bigwedge^k E' \longrightarrow \bigwedge^{k-1} E'$ is $f_1^{(0)} \bigwedge^{k-1} E''$. Furthermore, $f_1^{(0)}$ is regular in A', since on V the invertible ideal $\underline{J}'$ is defined by the equation $f_1^{(0)}$. The kernel of $d : \bigwedge^k E' \longrightarrow \bigwedge^{k-1} E'$ is therefore $\bigwedge^k E''$. Consequently, the k-th homology group of $K_\bullet(u) \otimes_A A'$ is $\bigwedge^k E''/f_1^{(0)} \bigwedge^k E''$; we thus obtain on V isomorphisms:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \xrightarrow{\sim} (\underline{O}_{X'}/f_1^{(0)}\underline{O}_{X'}) \otimes_{\underline{O}_{X'}} \bigwedge^i (\underline{O}_{X'})^{d-1} \xrightarrow{\sim} \bigwedge^i (\underline{O}_{Y'})^{d-1}. \tag{3.2.2}$$

In particular, for $i = 1$ we find:

$$\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \xrightarrow{\sim} (\underline{O}_{Y'})^{d-1}.$$

Consequently, for every i we have:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \xrightarrow{\sim} \bigwedge^i \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \tag{3.2.3}$$

By 2.2, the isomorphism:

$$\bigwedge (\underline{O}_{Y'})^{d-1} \xrightarrow{\sim} \bigoplus_{i \geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$$

defined by the isomorphisms (3.2.2) is an algebra isomorphism. Hence the same is true of the isomorphism:

$$\bigwedge \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \xrightarrow{\sim} \bigoplus_{i \geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$$

defined by the isomorphisms (3.2.3). By uniqueness, it is then the restriction to V of the homomorphism (3.2.1), and the latter is therefore an isomorphism.

ii) Consider the exact sequence:

$$0 \longrightarrow \underline{J} \longrightarrow \underline{O}_X \longrightarrow \underline{O}_Y \longrightarrow 0,$$

[print 440]and the exact $\underline{\mathrm{Tor}}$ sequence obtained by tensoring with $\underline{O}_{X'}$:

$$0 \longrightarrow \mathrm{Tor}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \longrightarrow \underline{J} \otimes_{\underline{O}_X} \underline{O}_{X'} \longrightarrow \underline{O}_{X'} \longrightarrow \underline{O}_{Y'} \longrightarrow 0.$$

The image of the morphism: $\underline{J} \otimes_{\underline{O}_X} \underline{O}_{X'} \to \underline{O}_{X'}$ is $\underline{J}.\underline{O}_{X'} = \underline{O}_{X'}(1)$. Consequently, we obtain the exact sequence:

$$0 \longrightarrow \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \longrightarrow \underline{J} \otimes_{\underline{O}_X} \underline{O}_{X'} \longrightarrow \underline{O}_{X'}(1) \longrightarrow 0.$$

Then tensor it with $\underline{O}_{Y'}$; since $\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y)$ is an $\underline{O}_{Y'}$-Module, and $\underline{O}_{X'}(1)$ is invertible, we obtain the exact sequence:

$$0 \longrightarrow \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \longrightarrow g^*(\underline{N}) \longrightarrow \underline{O}_{Y'}(1) \longrightarrow 0.$$

One readily checks that the morphism $g^*(\underline{N}) \to \underline{O}_{Y'}(1)$ occurring in this sequence is the canonical morphism, so that this sequence is identified with (3.1.2). By definition, we therefore have:

$$\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) = \underline{F}.$$

3.3. We now suppose that X is a scheme having an ample sheaf. Then the same is true of X', which is projective over X, and of Y and Y'. Furthermore, i and j are perfect morphisms, since they are regular immersions; g is smooth and hence perfect, since $Y' = \mathbf{P}(\underline{N})$[424]; finally, f is perfect by 1.9. Since these morphisms are projective, we can define the additive homomorphisms (IV

$$i_* : K^\bullet(Y) \to K^\bullet(X); \quad j_* : K^\bullet(Y') \to K^\bullet(X');$$
$$f_* : K^\bullet(X') \to K^\bullet(X); \quad g_* : K^\bullet(Y') \to K^\bullet(Y).$$

[print 441]**Proposition 3.4.** With the hypotheses and notation of 3.1 and 3.3, for every $x \in K^\bullet(Y)$ we have the relation:

$$f^*(i_*(x)) = j_*\big(g^*(x).\lambda_{-1}(F)\big).$$

Since both sides of the relation to be proved are additive, we may suppose that x is the class of an $\underline{O}_Y$-Module locally free of finite type $\underline{M}$. The element $f^*(i_*(x))$ is then the class of the complex $\mathbb{L}f^*(\mathbb{R}i_*(\underline{M}))$[425], obtained by taking a projective resolution of $i_*(\underline{M})$ and applying the functor f^*[426] to it. The homology sheaves of the resulting complex are the $\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{M}, \underline{O}_{X'})$. By 2.4 and 3.2, we have:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{M}, \underline{O}_{X'}) \xrightarrow{\sim} \underline{M} \otimes_{\underline{O}_Y} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_{X'}, \underline{O}_Y) \xrightarrow{\sim} \underline{M} \otimes_{\underline{O}_Y} \bigwedge^i \underline{F}.$$

It follows that these homology sheaves are locally free of finite type over $\underline{O}_{Y'}$, hence perfect over $\underline{O}_{X'}$; consequently, the class in $K^\bullet(X')$ of the complex $\mathbb{L}f^*(\mathbb{R}i_*(\underline{M}))$ is the alternating sum[427] of the classes of its homology sheaves, namely:

$$f^*(i_*(x)) = \sum_{i\geq 0}(-1)^i \,\mathrm{cl}\big(j_*(\underline{M} \otimes_{\underline{O}_Y} \bigwedge^i \underline{F})\big) = \sum_{i\geq 0}(-1)^i j_*\big(g^*(x)\lambda^i(F)\big).$$

Lemma 3.5. With the hypotheses and notation of 3.1, we have the following isomorphisms:

i) for every $n \geq 0$, and every $i \geq 1$, $R^i f_*\big(\underline{O}_{X'}(n)\big) = 0$;

ii) for every $n \geq 0$, $f_*\big(\underline{O}_{X'}(n)\big) = \underline{J}^n$.

We may clearly suppose X affine, hence quasi-compact.

i) First recall the following lemma, which was proved in exp. III.[428]

[print 442]**Lemma.** Let $X' \xrightarrow{f} X$ be a projective and perfect morphism, with X quasi-compact, and let $\underline{L}^\bullet$ be a perfect complex over $\underline{O}_{X'}$, acyclic in degrees > 0. Then there exists an integer n_0 such that for $n \geq n_0$ the complex $\mathbb{R}f_*\big(\underline{L}^\bullet \otimes_{\underline{O}_{X'}} \underline{O}_{X'}(n)\big)$[429] is acyclic in degrees > 0.

Apply this lemma to the complex reduced to $\underline{O}_{X'}$; there is therefore an integer n_0 such that for $n \geq n_0$, we have $R^i f_*(\underline{O}_{X'}(n)) = 0$ when $i \geq 1$. We then descend by induction to $n = 0$. For this, consider the exact sequence:

$$0 \longrightarrow \underline{O}_{X'}(n+1) \longrightarrow \underline{O}_{X'}(n) \longrightarrow \underline{O}_{Y'}(n) \longrightarrow 0,$$

and apply the functor f_* to it. We have $\mathbb{R}f_*(\underline{O}_{Y'}(n)) = i_* \circ \mathbb{R}g_*(\underline{O}_{Y'}(n))$[430]. Since $Y' = \mathbf{P}(\underline{N})$[431], $\underline{O}_{Y'}(n)$ is g_*-acyclic for $n \geq 0$. Thus for every $n \geq 0$, and every $i \geq 1$:

$$R^i f_*(\underline{O}_{Y'}(n)) = 0,$$

and consequently we have the exact sequences:

$$R^i f_*(\underline{O}_{X'}(n+1)) \longrightarrow R^i f_*(\underline{O}_{X'}(n)) \longrightarrow 0,$$

which complete the induction.

ii) We must show that the canonical morphism:

$$\bigoplus_{n \geq 0} \underline{J}^n \longrightarrow \bigoplus_{n \geq 0} f_*(\underline{O}_{X'}(n))$$

is an isomorphism. Since the property is local, we may suppose that $\underline{J}$ is defined by a sequence $f_1, \ldots, f_d$ of sections of $\underline{O}_X$ such that the corresponding homomorphism $(\underline{O}_X)^d \longrightarrow \underline{O}_X$ is regular. Put: $\underline{S} = \bigoplus_{n \geq 0} \underline{J}^n$; we define a surjective algebra homomorphism: $\underline{O}_X[T_1, \ldots, T_d] \overset{v}{\to} \underline{S}$, by sending [print 443]$T_i$ to $f_i^{(1)}$. This homomorphism defines a closed immersion $X' \longrightarrow \mathbf{P}_X^{d-1}$[432], regular by 1.8 ii).

By 1.8 iii), $\underline{S}$ is perfect as a graded $\underline{O}_X[T_1, \ldots, T_d]$-Module. In particular, $\underline{S}$ is finitely presented as a graded Module. Let therefore:

$$\underline{L}'' \longrightarrow \underline{L}' \longrightarrow \underline{S} \longrightarrow 0$$

be a finite graded presentation of $\underline{S}$ over $\underline{O}_X[T_1, \ldots, T_d]$. On $\mathbf{P}_X^{d-1}$ we obtain[433] the exact sequence:

$$\underline{M}'' \longrightarrow \underline{M}' \longrightarrow \underline{O}_{X'} \longrightarrow 0,$$

where $\underline{M}'' = \underline{\mathrm{Proj}}(\underline{L}'')$, $\underline{M}' = \underline{\mathrm{Proj}}(\underline{L}')$, so these are locally free of finite type[434] over $\mathbf{P}_X^{d-1}$. We can therefore[435] write, for every n, the commutative diagram:

$$\begin{array}{ccccccc}
\underline{L}''_n & \longrightarrow & \underline{L}'_n & \longrightarrow & \underline{J}^n & \longrightarrow & 0 \\
\sim\downarrow & & \sim\downarrow & & \downarrow & & \\
p_*(\underline{M}''(n)) & \longrightarrow & p_*(\underline{M}'(n)) & \longrightarrow & p_*(\underline{O}_{X'}(n)) & \longrightarrow & 0,
\end{array}$$

where p is the structural morphism: $\mathbf{P}_X^{d-1} \to X$[436], and $\underline{L}''_n$, $\underline{L}'_n$ are the degree n components of $\underline{L}''$, $\underline{L}'$. The upper row is exact by hypothesis; the first two vertical arrows are isomorphisms by the classical results for the direct images of $\underline{O}_P(n)$, P being a projective bundle (EGA III, 2.1.15); finally, the last row is exact when n is sufficiently large, since the kernel of $\underline{M}' \to \underline{O}_{X'}$ is a perfect $\underline{O}_P$-Module, and we can apply the lemma recalled above, which shows that this kernel, tensored with $\underline{O}_P(n)$ for n sufficiently large, is p_*-acyclic; likewise for the kernel of $\underline{M}'' \to \underline{M}'$. It follows that for n sufficiently large, $\underline{J}^n \longrightarrow p_*(\underline{O}_{X'}(n))$ is an isomorphism. Since $p_*(\underline{O}_{X'}(n)) = f_*(\underline{O}_{X'}(n))$, we therefore have [print 444]the result for $n \geq n_0$.

It remains to descend to $n = 0$. We proceed by descending induction, using the commutative diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \underline{J}^{n+1} & \longrightarrow & \underline{J}^{n} & \longrightarrow & \underline{J}^{n}/\underline{J}^{n+1} & \longrightarrow & 0 \\ \downarrow & & \sim\downarrow & & \downarrow & & \sim\downarrow & & \downarrow \\ 0 & \longrightarrow & f_*(\underline{O}_{X'}(n+1)) & \longrightarrow & f_*(\underline{O}_{X'}(n)) & \longrightarrow & f_*(\underline{O}_{Y'}(n)) & \longrightarrow & 0. \end{array}$$

The bottom row is exact because $R^1 f_*(\underline{O}_{X'}(n+1)) = 0$ for every $n \geq 0$, by i), already proved. Furthermore, we have: [437]

$$f_*(\underline{O}_{Y'}(n)) = i_*(g_*(\underline{O}_{Y'}(n))) = i_*(\mathbf{S}(\underline{N})_n) = \underline{J}^n/\underline{J}^{n+1}.$$

The arrow on the right is therefore an isomorphism; the one on the left is an isomorphism by the induction hypothesis; hence the middle one is also an isomorphism.[438]

Proposition 3.6. With the hypotheses and notation of 3.1 and 3.3, we have the relation:

$$f_* \circ f^* = \mathrm{Id}_{K^\bullet(X)}\,.$$

Let $x \in K^\bullet(X)$; by the projection formula, we can write:

$$f_*(f^*(x)) = f_*(1.f^*(x)) = x.f_*(1).$$

It therefore suffices to show that $f_*(1) = 1$.

Now 3.5 shows that the cohomology of $\mathbb{R}f_*(\underline{O}_{X'})$ vanishes in degrees > 0, and equals $\underline{O}_X$ in degree 0. The proposition follows at once.

Theorem 3.7. Let $i : Y \to X$ be a regular closed immersion, X' the scheme obtained by blowing up Y, and $Y' = X' \times_X Y$, so that we have the commutative diagram [print 445]

$$\begin{array}{ccc} Y' & \xrightarrow{j} & X' \\ {\scriptstyle g}\downarrow & & \downarrow{\scriptstyle f} \\ Y & \xrightarrow{i} & X. \end{array}$$

Suppose that X is quasi-compact and has an ample sheaf. Then, we have the exact sequence:

$$0 \longrightarrow K^\bullet(Y) \xrightarrow{u} K^\bullet(Y') \times K^\bullet(X) \xrightarrow{v} K^\bullet(X'),$$

where we put, for $y \in K^\bullet(Y)$:

$$u(y) = (g^*(y).\lambda_{-1}(F), -i_*(y)),$$

and for $y' \in K^\bullet(Y')$, $x \in K^\bullet(X)$:

$$v(y', x) = j_*(y') + f^*(x),$$

with the notation of 3.1 and 3.3. Moreover, u has a left inverse u', defined by $u'(y', x) = g_*(y')$. If in addition X is noetherian, and Y regular, then v is surjective, so we have the exact sequence:

$$0 \longrightarrow K^\bullet(Y) \xrightarrow{u} K^\bullet(Y') \times K^\bullet(X) \xrightarrow{v} K^\bullet(X') \longrightarrow 0.$$

First, u has the left inverse u', and is a fortiori injective. Indeed, we have:

$$g_*(g^*(y).\lambda_{-1}(F)) = y.g_*(\lambda_{-1}(F))$$

by the projection formula. Furthermore, $g_*(\lambda_{-1}(F)) = 1$ (VI 5.9), which gives the result.[439]

Next, $v \circ u = 0$. Indeed, for $y \in K^\bullet(Y)$ we have: [440]

$$v \circ u(y) = v(g^*(y)\lambda_{-1}(F), -i_*(y)) = j_*(g^*(y)\lambda_{-1}(F)) - f^*(i_*(y)) = 0,$$

by 3.4.

It therefore remains to show that $\mathrm{Ker}(v) \subset \mathrm{Im}(u)$. Let therefore $x \in K^\bullet(X)$, $y' \in K^\bullet(Y')$, be such that: [print 446]

$$j_*(y') + f^*(x) = 0.$$

We must find $y \in K^\bullet(Y)$ such that:

$$x = -i_*(y); \qquad y' = g^*(y).\lambda_{-1}(F). \tag{3.7.1}$$

Since u has a left inverse u', we necessarily have

$$y = u'(y', x) = g_*(y').$$

Thus we must show that y satisfies the relations (3.7.1).

To show that $x = -i_*(y)$, start with the relation $f^*(x) = -j_*(y')$ and apply f_* to it. We therefore obtain: $f_*(f^*(x)) = -f_*(j_*(y')) = -i_*(g_*(y')) = -i_*(y)$ by the definition of y. By Proposition 3.6, however, $f_*(f^*(x)) = x$; hence the required relation.

To complete the proof, we must show that

$$y' = g^*(g_*(y'))\lambda_{-1}(F).$$

Put $z = y' - g^*(g_*(y'))\lambda_{-1}(F)$. We obtain: $g_*(z) = g_*(y') - g_*(y')\, g_*(\lambda_{-1}(F)) = 0$. On the other hand, by hypothesis we have: $j_*(y') = -f^*(x)$[441], whence, by the preceding calculation, $j_*(y') = f^*(i_*(y))$, and by Proposition 3.4: $j_*(y') = j_*(g^*(y)\lambda_{-1}(F))$; consequently, we also have $j_*(z) = 0$. Now $\mathrm{Ker}(j_*) \cap \mathrm{Ker}(g_*) = 0$. Indeed, if $j_*(z) = 0$, then $j^*(j_*(z)) = 0$; but $j^*(j_*(z)) = z(1 - L)$ (Prop. 2.7), and for $z(1 - L) = 0$ to hold there must exist $z' \in K^\bullet(Y)$ such that $z = z'\lambda_{-1}(F)$ (VI,5.10); applying g_* gives $z' = g_*(z) = 0$, hence $z = 0$.


Suppose now that X is noetherian and Y regular. Then Y' is also regular, since it is a projective bundle over Y, and for every point y' of Y', the local ring $\underline{O}_{X',y'}$ of X' at y' is regular, since $\underline{O}_{Y',y'}$ is the quotient of $\underline{O}_{X',y'}$ by a regular element (see EGA 0_{IV} 17.1.8).

[print 447]We know that for every regular scheme having an ample sheaf, the Grothendieck groups formed with locally free sheaves of finite type and with coherent sheaves are identical. In particular, this is the case for Y'.

Let $\underline{E}$ be a coherent $\underline{O}_{X'}$-Module supported on Y'; then $\underline{E}$ is a perfect $\underline{O}_{X'}$-Module. Indeed, it is perfect at the points of Y', since X' is regular at these points, and at the points of $X' - Y'$ since it vanishes there.

If E is the class of $\underline{E}$ in $K^\bullet(X')$, then E belongs to the image of j_*. Indeed, we can filter $\underline{E}$ by the $\underline{J'}^n.\underline{E}$. This filtration is finite, since, X' being noetherian, there is an integer n_0 such that $\underline{J'}^{n_0}.\underline{E} = 0$; the quotients of this filtration are then $\underline{O}_{Y'}$-Modules, so their classes in $K^\bullet(X')$ belong to the image of j_*. Since E is the sum of the classes of these quotients, E also belongs to this image.

Let now $x \in K^\bullet(X')$. We can find a perfect complex $\underline{M}^\bullet$ whose class is x. Consider the distinguished triangle: [442]



where the horizontal arrow is the canonical homomorphism. This arrow is an isomorphism on the complement of Y', and consequently the cohomology sheaves of $\underline{N}^\bullet$ are coherent sheaves supported on Y'. By the preceding argument, these are perfect $\underline{O}_{X'}$-Modules, and their classes belong to the image of j_*. Thus we have:

$$x - f^*(f_*(x)) \in \mathrm{Im}(j_*)\,;$$

and consequently the homomorphism v is surjective.

[print 448]

Remark 3.8.

When X is noetherian and Y regular, 3.7 determines the abelian group $K^\bullet(X')$ entirely from the λ-rings $K^\bullet(X)$, $K^\bullet(Y)$, the maps i_*, i^* between these rings, and the element N of $K^\bullet(Y)$: indeed, $K^\bullet(Y')$ can be computed from $K^\bullet(Y)$ and N by (VI 1.1). We shall see in the next section that the multiplicative structure of $K^\bullet(X')$ is determined by the same data, as is the λ-structure after tensoring with $\mathbb{Q}$ (this restriction is probably unnecessary: see 4.5).

4. Regular immersions and the filtration of $K^\bullet$

Throughout this section we keep the notation and hypotheses of 3.1.

Lemma 4.1. Suppose that there exists on Y a locally free sheaf of finite type $\underline{N}'$, such that, if N, N' are the classes in $K^\bullet(Y)$ of $\underline{N}, \underline{N}'$, we have $N = N' + 2$. Then the following congruence holds:

$$\lambda_{-1}(F) \equiv 0 \text{ mod.}(1 - L).$$

Let d be the rank of $\underline{N}$ (which we may suppose constant); the rank of $\underline{F}$ is then $d - 1$. By definition:

$$\begin{aligned}\lambda_{-1}(F) &= \sum_{i\geq 0}(-1)^i\lambda^i(F)\\ &= (-1)^{d-1}\lambda^{d-1}(F-1)\\ &= (-1)^{d-1}\lambda^{d-1}(F+L-1-L)\\ &= (-1)^{d-1}\lambda^{d-1}(N-2+1-L).\end{aligned}$$

Now $\lambda_t(1-L) = \lambda_t(1)/\lambda_t(L) = (1+t)/(1+Lt)$. Thus for every $i \geq 1$, $\lambda^i(1-L)$ is divisible by $1 - L$. Consequently:

$$\begin{aligned}\lambda_{-1}(F) &\equiv (-1)^{d-1}\lambda^{d-1}(N-2) \qquad \text{mod.}(1-L)\\ &\equiv (-1)^{d-1}\lambda^{d-1}(N') \qquad \text{mod.}(1-L).\end{aligned}$$

[print 449]Now $\underline{N}'$ has rank $d - 2$, so $\lambda^{d-1}(N') = 0$, which gives the result.

4.2.

Let $i : Y \to X$ be a regular immersion. Let r be an integer ≥ 1, and $s : X \longrightarrow \mathbf{P}^r_X$[443] the section defined by the homomorphism: $(\underline{O}_X)^{r+1} \to \underline{O}_X$, passage to the quotient by the subsheaf generated by the last r coordinates. If $\mathbf{P}^r_X$[444] is defined by the graded $\underline{O}_X$-Algebra $\underline{O}_X[T_0, \ldots, T_r]$, the ideal $\underline{I}$ defining the section s is the ideal generated by the sections $T_1, \ldots, T_r$, so $\underline{I}/\underline{I}^2$ is isomorphic to $(\underline{O}_X[T_0])^r$, and hence the corresponding ideal of $\underline{O}_P$[445] is isomorphic to $(\underline{O}_X)^r$.

Since the section s is clearly a regular immersion, the same holds for $s \circ i$, and we can write the corresponding exact sequence of conormal sheaves (1.7). We thus obtain the following equality in $K^\bullet(Y)$, writing $\underline{N}$ for the conormal sheaf of i, and $\underline{N}_1$ for the conormal sheaf of $s \circ i$:

$$N_1 = N + r.$$

Thus for every regular immersion i, composing it with a section: $X \to \mathbf{P}^2_X$[446], or: $X \to \mathbf{P}^1_{\mathbf{P}^1_X}$[447], gives a regular immersion satisfying the property of Lemma 4.1.

Theorem 4.3. Let $i : Y \to X$ be a regular closed immersion, N the class in $K^\bullet(Y)$ of the conormal sheaf of Y in X. Suppose that X is quasi-compact and has an ample sheaf. Then, with the notation of 3.3, we have for every $y \in K^\bullet(Y)$, and every $p \geq 1$:

$$i_*(\lambda^p(N, y)) \equiv \lambda^p(i_*(y)) \qquad \text{mod. torsion}, \tag{4.3.1}$$

$\lambda^p(N, y)$ being the element defined in (V 5.3).

i) First suppose that the hypothesis of 4.1 is satisfied; in this case we shall show that $i_*(\lambda^p(N, y)) = \lambda^p(i_*(y))$. Since $f_* \circ f^* = \mathrm{Id}$, this [print 450]relation is equivalent to the relation:

$$f^*(i_*(\lambda^p(N, y))) = f^*(\lambda^p(i_*(y))).$$

Now we have:

$$\begin{aligned} f^*(\lambda^p(i_*(y))) &= \lambda^p(f^*(i_*(y))) \\ &= \lambda^p(j_*(g^*(y)\lambda_{-1}(F))) \qquad (3.4) \end{aligned}$$

On the other hand:

$$f^*(i_*(\lambda^p(N, y))) = j_*(g^*(\lambda^p(N, y))\lambda_{-1}(F)) \ (3.4).$$

Since $g^*(N) = F + L$, we can apply formula (V (5.4.2)):

$$g^*(\lambda^p(N, y))\lambda_{-1}(F) = \lambda^p(g^*(N), g^*(y))\lambda_{-1}(F) = \lambda^p(L, g^*(y)\lambda_{-1}(F)).$$

Putting: $z = g^*(y)\lambda_{-1}(F)$, we are therefore reduced to proving the relation:

$$j_*(\lambda^p(L, z)) = \lambda^p(j_*(z)),$$

that is, the analogous relation for the immersion j and the element z. By hypothesis, $\lambda_{-1}(F)$ is a multiple of $(1 - L)$; we can find z' such that $z = z'(1 - L)$. Then $z = j^*(u')$ with $u' = j_*(z')$ (2.7); since $L = j^*(J')$, we finally have to verify:

$$j_*(j^*(\lambda^p(J', u'))) = \lambda^p(j_*(j^*(u'))).$$

The exact sequence

$$0 \longrightarrow \underline{J}' \longrightarrow \underline{O}_{X'} \longrightarrow \underline{O}_{Y'} \longrightarrow 0$$

gives the relation:

$$j_*(1) = 1 - J'. \tag{4.3.2}$$

Thus, applying the projection formula, the relation to be verified finally becomes: [print 451]

$$\lambda^p(J', u')(1 - J') = \lambda^p(u'(1 - J')).$$

Since $1 - J' = \lambda_{-1}(J')$, this is the formula defining $\lambda^p(J', u')$.

ii) We turn to the general case. Compose i with a section: $X \to \mathbf{P}^1_X$[448], then with a section: $\mathbf{P}^1_X \to \mathbf{P}^1_{\mathbf{P}^1_X}$[449], so that the composite immersion satisfies the condition of 4.1. (see 4.2). Proposition 4.3 is then true for the composite immersion by the preceding argument; it therefore remains to "descend" to X. We descend first to $\mathbf{P}^1_X$[450], then to X, so that the proof consists in applying the following lemma twice (due to J.P. Jouanolou):

Lemma: Under the conditions of 4.3, let $s : X \to \mathbf{P}^1_X$[451] be the section defined in 4.2. If Proposition 4.3 is true for $s \circ i$, it is true for i.

Supposing 4.3 true for $s \circ i$, we have for every $p \geq 1$, and every $y \in K^\bullet(Y)$:

$$(s \circ i)_*\big(\lambda^p(N + 1, y)\big) \equiv \lambda^p\big((s \circ i)_*(y)\big) \quad \text{mod. torsion.} \tag{4.3.3}$$

The section s is defined by an ideal isomorphic to $\underline{O}_{\mathbf{P}}(-1)$[452] (4.2); write ξ for its class in $K^\bullet(\mathbf{P}^1_X)$[453]. We then have the following relation for every $x \in K^\bullet(X)$:

$$s_*(x) = p^*(x)(1 - \xi),$$

where p is the structural morphism $\mathbf{P}^1_X \to X$[454]. Indeed, since $p \circ s = \mathrm{Id}_X$, we have $x = s^*(p^*(x))$, whence $s_*(x) = s_*(s^*(p^*(x))) = p^*(x)s_*(1)$; since $s_*(1) = 1 - \xi$, we have the required relation:

Relation (4.3.3) then becomes:

$$\begin{aligned}(s \circ i)_*\big(\lambda^p(N + 1, y)\big) &\equiv \lambda^p\big(p^*(i_*(y))(1 - \xi)\big) \quad \text{mod. torsion} \\ &\equiv \lambda^p(\xi, p^*(i_*(y)))(1 - \xi) \quad \text{mod. torsion.}\end{aligned}$$

[print 452]Now $\lambda^p(\xi, p^*(i_*(y))) - \lambda^p(1, p^*(i_*(y)))$ is divisible by $1 - \xi$: to see this, we can work in the universal λ-ring used to define the $\lambda^p(\xi, X)$ (V 5); then $\lambda^p(\xi, X) - \lambda^p(1, X)$ is a polynomial in ξ, vanishing for $\xi = 1$, and hence divisible by $\xi - 1$. Furthermore, the dependence equation of ξ over $K^\bullet(X)$ is $(1 - \xi)^2 = 0$ (VI (1.13.1)). We therefore obtain:

$$\begin{aligned}(s \circ i)_*\big(\lambda^p(N + 1, y)\big) &\equiv \lambda^p(1, p^*(i_*(y)))(1 - \xi) \quad \text{mod. torsion} \\ &\equiv p^*(\lambda^p(1, i_*(y)))(1 - \xi) \quad \text{mod. torsion.}\end{aligned}$$

Applying p_*, we obtain:

$$i_*\big(\lambda^p(N + 1, y)\big) \equiv \lambda^p(1, i_*(y)) \quad \text{mod. torsion,} \tag{4.3.4}$$

since $p_*(\xi) = 0$.

By the formula in (V 5.6), applied with $L = 1$, $\lambda^p(N + 1, y)$ is the coefficient of t^p in the series:

$$\frac{\sum_{k \geq 1} \lambda^k(N, y)kt^k}{1 + \sum_{k \geq 1} \lambda^k(N, y)\lambda_{-1}(N)t^k}.$$

Similarly, the same formula applied with $N = 0$ and $L = 1$ shows that $\lambda^p(1, y)$ is the coefficient of t^p in the series:

$$\frac{\sum_{k \geq 1} \lambda^k(y)kt^k}{\sum_{k \geq 0} \lambda^k(y)t^k}.$$

The congruences (4.3.4) can therefore be written:

$$i_*\left(\frac{\sum_{k\geq 1} k\lambda^k(N,y)t^k}{1+\sum_{k\geq 1}\lambda^k(N,y)\lambda_{-1}(N)t^k}\right) \equiv \frac{\sum_{k\geq 1} k\lambda^k(i_*(y))t^k}{\sum_{k\geq 0}\lambda^k(i_*(y))t^k} \quad \text{mod. torsion}$$

[print 453]Applying formula 2.8, we obtain:

$$\frac{\sum_{k\geq 0} k\, i_*(\lambda^k(N,y))t^k}{1+\sum_{k\geq 1} i_*(\lambda^k(N,y))t^k} \equiv \frac{\sum_{k\geq 0} k\lambda^k(i_*(y))t^k}{\sum_{k\geq 0}\lambda^k(i_*(y))t^k} \quad \text{mod. torsion}$$

Putting:

$$u(t) = 1+\sum_{k\geq 1} i_*(\lambda^k(N,y))t^k, \qquad v(t) = \sum_{k\geq 0}\lambda^k(i_*(y))t^k,$$

we can also write this last relation as:

$$\frac{d}{dt}\left(\frac{u(t)}{v(t)}\right) \equiv 0 \quad \text{mod. torsion.}$$

Since the constant term of $u(t)/v(t)$ is 1, and we are writing congruences modulo torsion, we can integrate and write the congruence $u(t) \equiv v(t)$. This gives the stated congruences.

Corollary 4.4. With the notation and hypotheses of 4.3, we have for every $p \geq 1$, and every $y \in K^\bullet(Y)$: [455]

$$i_*(\gamma^p(N,y)) \equiv \gamma^p(i_*(y)) \quad \text{mod. torsion.}$$

The corollary follows immediately from 4.3, using[456] the expression of the γ^p as linear combinations with integer coefficients of the λ^p.

Remark 4.5. The proof of 4.3 itself suggests that (4.3.1) should hold not only modulo torsion, but also as an equality. In [RRR], the equality was proved in characteristic 0 by means of a geometric characterization of the $\lambda^p(N,x)$. An extension of this method to all characteristics would probably be possible if one knew how to define exterior-power operations on perfect complexes (cf. XIV).

[print 454]**Theorem 4.6.** Let X be a quasi-compact scheme having an ample sheaf, d an integer > 0 and $i : Y \to X$ a regular immersion of codimension d. Then, for every k, we have:

$$i_*((\mathrm{Fil}^k K^\bullet(Y)) \otimes_{\mathbb{Z}} \mathbb{Q}) \subset (\mathrm{Fil}^{k+d} K^\bullet(X)) \otimes_{\mathbb{Z}} \mathbb{Q}.$$

First observe that $i_*(K^\bullet(Y)) \subset \mathrm{Fil}^1 K^\bullet(X)$. Indeed, $X - Y$ is dense in X: otherwise, Y would contain an affine open subset, and on this open subset the ideal $\underline{J}$ defining Y would be a nil ideal, hence a locally nilpotent ideal since it is of finite type. This is impossible, however, since the $\underline{J}^n/\underline{J}^{n+1}$ are locally free $\underline{O}_Y$-Modules of rank $\neq 0$. The augmentation in $K^\bullet(X)$ of an element of $i_*(K^\bullet(Y))$ is a locally constant function on X; since it vanishes on $X - Y$, it vanishes on all of X.

For every abelian group A, we put $A_{\mathbb{Q}} = A \otimes_{\mathbb{Z}} \mathbb{Q}$. Let $x \in \mathrm{Fil}^k K^\bullet(Y)_{\mathbb{Q}}$; we can write x in the form:

$$x = \sum a\, \gamma^{i_1}(x_1)\cdots\gamma^{i_n}(x_n),$$

with $\epsilon(x_i) = 0$ for every i, $a \in \mathbb{Q}$, and $i_1 + \cdots + i_n \geq k$. The x_i can be expressed as linear combinations with integer coefficients of the classes E_α of finitely many locally free $\underline{O}_Y$-Modules of finite type $\underline{E}_\alpha$. Let R be the sub-$\mathbb{Q}$-λ-algebra of $K^\bullet(Y)_{\mathbb{Q}}$ generated by the E_α and by N. We then have:

$$x \in \mathrm{Fil}^k R.$$

Furthermore, the λ-ring filtration of R is discrete (VI 6.2); let m satisfy $\mathrm{Fil}^m R = 0$. Finally, by decomposing Y into a finite sum[457] of disjoint open subsets and carrying out the proof on each open subset (since $K^\bullet$ commutes with finite direct sums), we may suppose that the $\underline{E}_\alpha$ have constant rank r_α. The augmentation ring of R is then $\mathbb{Q}$.

[print 455]We shall show that for every i, we have:

$$i_*(\mathrm{Fil}^i R) \subset \mathrm{Fil}^{i+d}(K^\bullet(X)_\mathbb{Q}).$$

This will imply in particular: [458]

$$i_*(x) \in \mathrm{Fil}^{k+d} K^\bullet(X)_\mathbb{Q},$$

and hence prove the theorem.

The proof is by descending induction on i, the case $i = m$ being trivial. Let $y \in \mathrm{Fil}^i R$, and put, for every j:

$$a_j = (-1)^{j-1}(j-1)!.$$

Then by V 6.10,

$$\gamma^{d+i}(N, y) - a_{d+i} y \in \mathrm{Fil}^{i+1} R.$$

We can therefore apply the induction hypothesis, supposing that:

$$i_*(\mathrm{Fil}^{i+1} R) \subset \mathrm{Fil}^{i+d+1} K^\bullet(X)_\mathbb{Q}.$$

Consequently, we have:

$$i_*\big(\gamma^{d+i}(N, y) - a_{d+i} y\big) \in \mathrm{Fil}^{i+d+1} K^\bullet(X)_\mathbb{Q}.$$

By 4.4, this implies:

$$\gamma^{d+i}\big(i_*(y)\big) - a_{d+i} i_*(y) \in \mathrm{Fil}^{i+d+1} K^\bullet(X)_\mathbb{Q}.$$

Since $i_*(y)$ has zero augmentation, $\gamma^{d+i}(i_*(y)) \in \mathrm{Fil}^{d+i} K^\bullet(X)_\mathbb{Q}$[459]. Since we are calculating modulo torsion, we therefore find:

$$i_*(y) \in \mathrm{Fil}^{i+d} K^\bullet(X)_\mathbb{Q}.$$

Remark 4.6.1. We see that, even if 4.3 were true without tensoring with $\mathbb{Q}$, the proof above works only modulo torsion. For a counterexample, see XIV 4.6.

[print 456]

4.7.

We return to the situation of 3.7, where we had the commutative diagram:

$$\begin{array}{ccc} Y' & \xrightarrow{j} & X' \\ {\scriptstyle g}\downarrow & & \downarrow{\scriptstyle f} \\ Y & \xrightarrow{i} & X, \end{array}$$

where i is a regular immersion, X' the X-scheme obtained by blowing up Y, and $Y' = X' \times_X Y$. Suppose that X is noetherian with an ample sheaf, and Y regular. We saw in 3.7 that we then have the exact sequence:

$$0 \longrightarrow K^\bullet(Y) \xrightarrow{u} K^\bullet(Y') \times K^\bullet(X) \xrightarrow{v} K^\bullet(X') \longrightarrow 0.$$

We shall show that this exact sequence describes the whole situation formed by the $K^\bullet$ of X, Y, X', Y' from the data of the λ-rings $K^\bullet(X)$, $K^\bullet(Y)$, the maps i^* and i_*, and the element N of $K^\bullet(Y)$. (at least up to torsion, for the time being).

Let therefore K, K' be two λ-rings, $i_* : K \to K'$ an additive homomorphism, $i^* : K' \to K$ a λ-homomorphism, d an integer, and N an element of K such that $\lambda^i(N) = 0$ for $i > d$ and $\lambda^d(N)$ is invertible. Suppose moreover that the following properties hold:

i) i_* is K'-linear ("projection formula");

ii) $i^* \circ i_*$ is the homomorphism "multiplication by $\lambda_{-1}(N)$";

iii) for every $n \geq 1$, and every $y \in K$, $i_*(\lambda^n(N, y)) = \lambda^n(i_*(y))$ (relation (4.3.1)).[460]

First we use VI.1.1 to construct[461] a K-λ-algebra K_1 which plays the role of $K^\bullet(Y')$. Let L be an indeterminate, and $K[L]$ the K-λ-algebra generated by the generator L subject to the relations $\lambda^i(L) = 0$[462] for $i \geq 2$ (V 4.9). Put:

$$P(L) = \lambda^d(N - L) = \sum_{i=0}^{d} (-1)^i \lambda^{d-i}(N) L^i.$$

[print 457]The ideal generated by P is a λ-ideal. To see this, first observe that the relations $\lambda^i(N) = 0$ for $i > d$ imply that the $\lambda^j(N - L)$ for $j \geq d$ are multiples of $\lambda^d(N - L)$. Working in $1 + K\widehat{[L]/(P)}[[t]]^{+}$[463], the image of $\lambda_t(N - L)$ is a polynomial of degree $< d$, and a universal argument then shows that its λ^i for $i \geq d$ equal 1 (cf. V 4.8). Since in $1 + \widehat{K[L][[t]]}^{+}$[464] we have $\lambda_t(\lambda^d(N - L)) = \lambda^d(\lambda_t(N - L))$, we deduce that the $\lambda^i(\lambda^d(N - L))$ for $i \geq 1$ are multiples of P. By the relations V 2.4.2, this implies that the ideal generated by P is a λ-ideal.

Put then $K_1 = K[L]/(P)$[465]. We denote the natural homomorphism $K \to K_1$ by g^*; it is a λ-homomorphism. Furthermore, K_1 is a free K-module with basis the elements $1, L, \ldots, L^{d-1}$. The relation $\lambda^d(N) = \lambda^{d-1}(N - L).L$ and the invertibility hypothesis on $\lambda^d(N)$ imply that L is invertible. Consequently, K_1 has as a basis over K the elements $1, L^{-1}, \ldots, L^{-d+1}$; we then define a map $g_* : K_1 \to K$ by assigning to every element $y' = y_0 + y_1 L^{-1} + \ldots + y_{d-1} L^{-d+1}$ of K_1 the element y_0 of K. Clearly the map thus defined is an additive, K-linear homomorphism, satisfying $g_* \circ g^* = \mathrm{Id}$.

Next we endow $K_1 \times K'$ with a λ-ring structure, putting:

$$(y'_1, x_1) * (y'_2, x_2) = (y'_1 y'_2 (1 - L) + x_1 y'_2 + x_2 y'_1,\ x_1 x_2); \tag{4.7.1}$$

$$\lambda^n_*(y', x) = \left(\sum_{p=1}^{n} \lambda^{n-p}(x) \lambda^p(L, y'),\ \lambda^n(x) \right). \tag{4.7.2}$$

In these formulas, K_1 is regarded as a K'-algebra through the homomorphism $g^* \circ i^*$. Next, for $y_1, y_2 \in K$ and $n \geq 1$, put:

$$y_1 * y_2 = -\, y_1 y_2 \lambda_{-1}(N); \tag{4.7.3}$$

$$\lambda^n_*(y_1) = -\, \lambda^n(N, -y_1); \tag{4.7.4}$$

[print 458]and define a homomorphism:

$$u : K \longrightarrow K_1 \times K'$$

by putting:

$$u(y) = (g^*(y).\lambda_{-1}(F), -i_*(y)),$$

with $F = N - L$.

Proposition 4.7.5. The operations defined by (4.7.1) and (4.7.2) endow $K_1 \times K'$ with a λ-ring structure, and u is an injective homomorphism identifying K, equipped with the operations defined by (4.7.3) and (4.7.4), with a λ-ideal of $K_1 \times K'$.

It is immediate to verify that $K_1 \times K'$ is a λ-ring: one can refer, for example, to V 5.5, where an entirely analogous construction is given. The second part of the proposition is also proved without difficulty using the preceding relations, the relation $i_*(y_1 y_2 \lambda_{-1}(N)) = i_*(y_1).i_*(y_2)$ which follows immediately from the hypotheses, the relation $g_*(\lambda_{-1}(F)) = 1$ (see VI 5.9), and the relations of V 5.

Put then:

$$K'_1 = (K_1 \times K')/K$$

and denote by v the canonical homomorphism $K_1 \times K' \to K'_1$. Regard K'_1 as a λ-ring through the quotient structure.

Next define additive homomorphisms:

$$j_* : K_1 \to K'_1; \quad j^* : K'_1 \to K_1;$$
$$f_* : K'_1 \to K'; \quad f^* : K' \to K'_1.$$

The homomorphism j_* is the composite $K_1 \to K_1 \times K' \to K'_1$. Likewise, the homomorphism f^* is the composite $K' \to K_1 \times K' \to K'_1$. We then have:

$$v = j_* + f^*.$$

[print 459]Next define a homomorphism $K_1 \times K' \to K_1$ by assigning to (y', x) the element $y'(1-L)+g^* \circ i^*(x)$; this homomorphism vanishes on K, and consequently defines a homomorphism $j^* : K'_1 \longrightarrow K_1$. Finally, define a homomorphism $K_1 \times K' \to K'$ by assigning to (y', x) the element $i_* \circ g_*(y') + x$; this homomorphism vanishes on K and therefore defines a homomorphism $f_* : K'_1 \longrightarrow K'$.

Proposition 4.7.6.

i) j^* and f^* are λ-homomorphisms;

ii) j_* is K'_1-linear and f_* is K'-linear;

iii) $j^* \circ j_*$ is the homomorphism "multiplication by $1 - L$";

iv) for every $y' \in K_1$, and every $p \geq 1$, $j_*(\lambda^p(L, y')) = \lambda^p(j_*(y'))$;

v) $f_* \circ f^* = \mathrm{Id}$.

We need only set the machinery in motion.

Returning to the geometric situation recalled above, we see that $K^\bullet(X')$ is constructed from $K^\bullet(Y)$, $K^\bullet(X)$, the maps i^* and i_*, and the element N of $K^\bullet(Y)$ by the preceding method; the λ-operations on $K^\bullet(X')$, however, agree with the λ-operations defined above on K'_1 only up to torsion, as long as relation (4.3.1) is proved only modulo torsion.

Theorem 4.8. Let $i : Y \to X$ be a regular closed immersion of constant codimension d, X' the scheme obtained by blowing up Y, and $Y' = X' \times_X Y$, so that we have the commutative diagram:

$$\begin{array}{ccc} Y' & \xrightarrow{j} & X' \\ {\scriptstyle g}\downarrow & & \downarrow{\scriptstyle f} \\ Y & \xrightarrow{i} & X. \end{array}$$

Suppose that X is quasi-compact and has an ample sheaf. Then for every k we have the exact sequence: [print 460]

$$0 \longrightarrow \mathrm{Gr}^{k-d} K^\bullet(Y)_\mathbb{Q} \xrightarrow{u^{\mathrm{gr}}} \mathrm{Gr}^{k-1} K^\bullet(Y')_\mathbb{Q} \times \mathrm{Gr}^k K^\bullet(X)_\mathbb{Q} \xrightarrow{v^{\mathrm{gr}}} \mathrm{Gr}^k K^\bullet(X')_\mathbb{Q}$$

where we put, for $y \in \mathrm{Gr}^{k-d} K^\bullet(Y)_\mathbb{Q}$:

$$u^{\mathrm{gr}}(y) = (g^{\mathrm{gr}}(y).(-1)^{d-1}c^{d-1}(F), -i_{\mathrm{gr}}(y)),$$

and for $y' \in \mathrm{Gr}^{k-1} K^\bullet(Y')_\mathbb{Q}$ and $x \in \mathrm{Gr}^k K^\bullet(X)_\mathbb{Q}$:

$$v^{\mathrm{gr}}(y', x) = j_{\mathrm{gr}}(y') + f^{\mathrm{gr}}(x),$$

with the notation of 3.1 and 3.3, and $c^{d-1}(F)$ denoting the $(d-1)$-st Chern class of F (V 6.7). Moreover, u^{gr} has a left inverse u'^{gr}, defined by $u'^{\mathrm{gr}}(y', x) = g_{\mathrm{gr}}(y')$.

If in addition X is noetherian and Y regular, we have the exact sequence:

$$0 \longrightarrow \mathrm{Gr}^{k-d} K^\bullet(Y)_\mathbb{Q} \xrightarrow{u^{\mathrm{gr}}} \mathrm{Gr}^{k-1} K^\bullet(Y')_\mathbb{Q} \times \mathrm{Gr}^k K^\bullet(X)_\mathbb{Q} \xrightarrow{v^{\mathrm{gr}}} \mathrm{Gr}^k K^\bullet(X')_\mathbb{Q} \longrightarrow 0$$

Proof. Observe that, when Y is regular, one could give a purely algebraic proof in terms of the theory of λ-rings, using the description of the situation in 4.7. We shall instead follow the proof of 3.7.

We know that the "inverse-image" homomorphisms f^*, g^*, being λ-homomorphisms, are compatible with the λ-ring filtrations, and consequently define homomorphisms on the associated gradeds; hence the homomorphisms:

$$f^{\mathrm{gr}} : \mathrm{Gr}^k K^\bullet(X) \to \mathrm{Gr}^k K^\bullet(X'); \qquad g^{\mathrm{gr}} : \mathrm{Gr}^{k-d} K^\bullet(Y) \to \mathrm{Gr}^{k-d} K^\bullet(Y'),$$

and the analogous homomorphisms obtained after tensoring with $\mathbb{Q}$. Furthermore, 4.6 shows that after tensoring with $\mathbb{Q}$, the homomorphisms i_*, j_* pass to the associated gradeds, and therefore define:

$$i_{\mathrm{gr}} : \mathrm{Gr}^{k-d} K^\bullet(Y)_\mathbb{Q} \to \mathrm{Gr}^k K^\bullet(X)_\mathbb{Q}; \qquad j_{\mathrm{gr}} : \mathrm{Gr}^{k-1} K^\bullet(Y')_\mathbb{Q} \to \mathrm{Gr}^k K^\bullet(X')_\mathbb{Q}.$$

[print 461]Finally, observe that $\lambda_{-1}(F) = (-1)^{d-1}\gamma^{d-1}(F)$, so the morphisms u^{gr} and v^{gr} are obtained by passage to the quotient from the morphisms u and v of 3.7. Thus we have $v^{\mathrm{gr}} \circ u^{\mathrm{gr}} = 0$.

By VI 5.8, $g_*(\mathrm{Fil}^{k-1} K^\bullet(Y')) \subset \mathrm{Fil}^{k-d} K^\bullet(Y)$ for every k, since $Y' = \mathbf{P}(\underline{N})$[466], with $\underline{N}$ of rank d. We therefore obtain a homomorphism:

$$g_{\mathrm{gr}} : \mathrm{Gr}^{k-1} K^\bullet(Y') \to \mathrm{Gr}^{k-d} K^\bullet(Y),$$

and an analogous homomorphism after tensoring with $\mathbb{Q}$. The homomorphism u'^{gr} is therefore obtained from u' (3.7) by passage to the associated graded. Since u' is a left inverse of u, u'^{gr} is a left inverse of u^{gr}, proving at the same time the injectivity of u^{gr}.

To show that $\mathrm{Ker}(v^{\mathrm{gr}}) = \mathrm{Im}(u^{\mathrm{gr}})$, we must show that for $y' \in \mathrm{Fil}^{k-1} K^\bullet(Y')_\mathbb{Q}$, and $x \in \mathrm{Fil}^k K^\bullet(X)_\mathbb{Q}$, satisfying: [467]

$$j_*(y') + f^*(x) \in \mathrm{Fil}^{k+1} K^\bullet(X')_\mathbb{Q}, \tag{4.8.1}$$

there exists $y \in \mathrm{Fil}^{k-d} K^\bullet(Y)_\mathbb{Q}$ such that:

i) $x + i_*(y) \in \mathrm{Fil}^{k+1} K^\bullet(X)_\mathbb{Q}$,

ii) $y' - g^*(y)\lambda_{-1}(F) \in \mathrm{Fil}^k K^\bullet(Y')_\mathbb{Q}$.

Put:

$$y = g_*(y').$$

Then by VI 5.8, $y \in \mathrm{Fil}^{k-d} K^\bullet(Y)_\mathbb{Q}$[468]. Next observe that f_* preserves the filtration (using results of Exposé VIII, which causes no difficulty, since 4.8 is not used to prove this fact). Indeed, it follows from 1.8 ii) that f is a complete-intersection morphism of virtual relative dimension zero (VIII 4.2). Since f is projective, the result follows from VIII 3.2. Applying f_* to the relation [print 462](4.8.1), we obtain:

$$f_*(j_*(y')) + f_*(f^*(x)) \in \mathrm{Fil}^{k+1} K^\bullet(X)_\mathbb{Q},$$

or, by 3.6:

$$i_*(y) + x \in \mathrm{Fil}^{k+1} K^\bullet(X)_\mathbb{Q}.$$

To prove relation ii), we must show that:

$$y' - g^*(g_*(y'))\lambda_{-1}(F) \in \mathrm{Fil}^k K^\bullet(Y')_\mathbb{Q}.$$

Put $z = y' - g^*(g_*(y'))\lambda_{-1}(F)$. We have:

$$\begin{aligned} j_*(z) &= j_*(y') - j_*(g^*(g_*(y'))\lambda_{-1}(F)) \\ &= j_*(y') - f^*(i_*(y)) \qquad \text{by 3.4.} \end{aligned}$$

Now relation i) gives:

$$f^*(i_*(y)) + f^*(x) \in \mathrm{Fil}^{k+1} K^\bullet(X')_\mathbb{Q}.$$

Taking (4.8.1) into account, we therefore obtain:[469]

$$j_*(z) \in \mathrm{Fil}^{k+1} K^\bullet(X')_\mathbb{Q},$$

whence:

$$j^*(j_*(z)) = z(1-L) \in \mathrm{Fil}^{k+1} K^\bullet(Y')_\mathbb{Q}.$$

Let $\bar{z}$ be the image of z in $\mathrm{Gr}^{k-1} K^\bullet(Y')_\mathbb{Q}$, and $\overline{(1-L)}$ that of $(1-L)$ in $\mathrm{Gr}^1 K^\bullet(Y')_\mathbb{Q}$. We therefore have: $\overline{z(1-L)} = 0$. We can then apply (VI 5.11) (which remains valid after tensoring with $\mathbb{Q}$, since its proof uses only the structure of $K^\bullet(Y')$ as a free $K^\bullet(Y)$-algebra), and hence find $z' \in K^\bullet(Y)_\mathbb{Q}$ such that:

$$z - g^*(z')\lambda_{-1}(F) \in \mathrm{Fil}^k K^\bullet(Y')_\mathbb{Q}.$$

Applying g_*, and using $g_*(z) = 0$, we find:

$$z' \in \mathrm{Fil}^{k+1-d} K^\bullet(Y)_\mathbb{Q},$$

whence, since $\lambda_{-1}(F) \in \mathrm{Fil}^{d-1} K^\bullet(Y')_\mathbb{Q}$:

$$z \in \mathrm{Fil}^k K^\bullet(Y')_\mathbb{Q},$$

which completes the proof.

[print 463]Suppose now that X is noetherian and Y regular, which implies that v is surjective (3.7). We saw in the proof of 3.7 that for every $x' \in K^\bullet(X')$ there is $y' \in K^\bullet(Y')$ such that:

$$x' = f^*(f_*(x')) + j_*(y').$$

The same property clearly remains true after tensoring with $\mathbb{Q}$. Consequently, for $x' \in \mathrm{Fil}^k K^\bullet(X')_\mathbb{Q}$ there is $(y', x) \in K^\bullet(Y')_\mathbb{Q} \times K^\bullet(X)_\mathbb{Q}$ such that:

$$x' = v(y', x), \qquad x \in \mathrm{Fil}^k K^\bullet(X)_\mathbb{Q}.$$

Indeed, it suffices to take $x = f_*(x')$.

Filter the ring $K^\bullet(Y')_\mathbb{Q} \times K^\bullet(X)_\mathbb{Q}$ defined in 4.7 by putting:

$$\mathrm{Fil}^k(K^\bullet(Y')_\mathbb{Q} \times K^\bullet(X)_\mathbb{Q}) = \mathrm{Fil}^{k-1} K^\bullet(Y')_\mathbb{Q} \times \mathrm{Fil}^k K^\bullet(X)_\mathbb{Q}.$$

It follows immediately from (4.7.1) that this is a ring filtration. Thus, to show that for every k the homomorphism:

$$\mathrm{Fil}^k(K^\bullet(Y')_\mathbb{Q} \times K^\bullet(X)_\mathbb{Q}) \xrightarrow{v} \mathrm{Fil}^k K^\bullet(X')_\mathbb{Q}$$

is surjective, it suffices to show that for $x' = \gamma^k(x'_1)$, with $\epsilon(x'_1) = 0$, there is $(y', x) \in \mathrm{Fil}^k(K^\bullet(Y')_\mathbb{Q} \times K^\bullet(X)_\mathbb{Q})$ such that $x' = v(y', x)$.

By the preceding argument, there is $(y'_1, x_1) \in K^\bullet(Y')_\mathbb{Q} \times K^\bullet(X)_\mathbb{Q}$ such that $x'_1 = v(y'_1, x_1)$ and $x_1 \in \mathrm{Fil}^1 K^\bullet(X)_\mathbb{Q}$. By 4.7, we then have:

$$x' = v(\gamma^k_*(y'_1, x_1)).$$

Now:

$$\gamma^k_*(y'_1, x_1) = \left(\sum_{p=1}^{k} \gamma^{k-p}(x_1)\gamma^p(L, y'_1),\ \gamma^k(x_1) \right)$$

(an immediate consequence of (4.7.2)). Since x_1 has zero augmentation, $\gamma^k(x_1) \in \mathrm{Fil}^k K^\bullet(X)_\mathbb{Q}$, and, using V 5.7,

$$\sum_{p=1}^{k} \gamma^{k-p}(x_1)\gamma^p(L, y'_1) \in \mathrm{Fil}^{k-1} K^\bullet(Y')_\mathbb{Q},$$

which gives the result.

[print 464]**Remark 4.9.** When X is noetherian and Y regular, proceeding as in 4.7 we can describe the graded ring structure of $\mathrm{Gr}^* K^\bullet(X')_\mathbb{Q}$ in terms of the graded rings $\mathrm{Gr}^* K^\bullet(X)_\mathbb{Q}$, $\mathrm{Gr}^* K^\bullet(Y)_\mathbb{Q}$, the additive homomorphisms i^{gr} and i_{gr} induced by i^* and i_*, and the element $c^\bullet(N)$ of $\mathrm{Gr}^* K^\bullet(Y)$. Likewise, the same data describe the homomorphisms $f^{\mathrm{gr}}, f_{\mathrm{gr}}, j^{\mathrm{gr}}, j_{\mathrm{gr}}$ defined from f^*, f_*, j^*, j_* by passage to the associated gradeds.

4.10.

Let X be a regular scheme having an ample sheaf. We know (IV) that the Grothendieck groups formed with locally free $\underline{O}_X$-Modules of finite type and with coherent $\underline{O}_X$-Modules are isomorphic. We use this fact to define a "topological" filtration on $K^\bullet(X)$, taking $\mathrm{Fil}^i_{\mathrm{top}} K^\bullet(X)$ to be the subgroup generated by the classes of coherent sheaves whose support has codimension $\geq i$; it is easy to show that $\mathrm{Fil}^i_{\mathrm{top}} K^\bullet(X)$ is generated by the classes of the sheaves $\underline{O}_Z$, where Z ranges over the integral closed subschemes of X of codimension $\geq i$(X).

Proposition 4.11. Let X be a regular scheme having an ample sheaf. Then for every $k \geq 0$:

$$\mathrm{Fil}^k_{\mathrm{top}} K^\bullet(X)_\mathbb{Q} \subset \mathrm{Fil}^k_\lambda K^\bullet(X)_\mathbb{Q}.$$

(N.B. The reverse inclusion, valid without tensoring with $\mathbb{Q}$, will be proved in X).

By 4.10, it suffices to show that for an integral closed subscheme Z of X, of codimension k, we have:

$$\mathrm{cl}(\underline{O}_Z) \in \mathrm{Fil}^k K^\bullet(X)_\mathbb{Q}.$$

For this, we proceed by noetherian induction on the integral closed subschemes of X. Suppose therefore that the property is proved for the integral closed subschemes of Z, and let us prove it

for Z. Let $\underline{J}$ be the Ideal of $\underline{O}_X$ [print 465]defining Z, and let z be the generic point of Z. Since Z is integral, $\underline{J}_z$ is the maximal ideal of $\underline{O}_{X,z}$, and since X is regular, $\underline{J}_z$ can be generated by an $\underline{O}_{X,z}$-regular sequence. We can therefore find a sequence of sections of $\underline{J}$ in a neighborhood of z forming an $\underline{O}_X$-regular sequence. Consequently, there is an open neighborhood U of z such that $Z' = X - U$ is contained in Z, and $\underline{J}|U$ is a regular ideal of $\underline{O}_U$. The immersion $Z \cap U \xrightarrow{i} U$ is then a regular immersion, with codimension $\dim(\underline{O}_{X,z}) = \operatorname{codim}(Z, X)$ (EGA IV 5.1.2). Furthermore, we have the exact sequence (IV):[470]

$$K^\bullet(Z') \xrightarrow{j_*} K^\bullet(X) \xrightarrow{u^*} K^\bullet(U) \to 0,$$

where j is the immersion $Z' \to X$ and u the immersion $U \to X$; tensoring with $\mathbb{Q}$ gives an analogous sequence. Now $u^*(\underline{O}_Z) = i_*(\underline{O}_{Z\cap U})$, so by 4.6 the image $u^*(1_Z)$ of the class of $\underline{O}_Z$ under u^* belongs to $\operatorname{Fil}^k K^\bullet(U)_\mathbb{Q}$. Since u^* is a surjective λ-homomorphism, it induces a surjection $\operatorname{Fil}^k K^\bullet(X) \longrightarrow \operatorname{Fil}^k K^\bullet(U)$, so $u^*(1_Z) = u^*(x)$ with $x \in \operatorname{Fil}^k K^\bullet(X)$; there is therefore $y \in j_*(K^\bullet(Z'))$ such that:

$$\operatorname{cl}(\underline{O}_Z) - y \in \operatorname{Fil}^k K^\bullet(X)_\mathbb{Q}.$$

Now $Z' \subset Z$, and since $z \notin Z'$, $\operatorname{codim}(Z', X) > k$. Consequently, y can be expressed[471] as a combination of classes of sheaves $\underline{O}_{Z''_\alpha}$, where the Z''_α are integral closed subschemes of Z, distinct from Z. By the induction hypothesis, $y \in \operatorname{Fil}^{k+1} K^\bullet(X)_\mathbb{Q}$[472], which completes the proof.

BIBLIOGRAPHY

[1] Cartan–Eilenberg, Homological Algebra.


[print 466]

EXPOSÉ VIII
THE RIEMANN–ROCH THEOREM

by Pierre BERTHELOT

1. Complete-intersection morphisms

Definition 1.1. Let $f : X \to Y$ be a morphism of schemes. One says that f is a complete-intersection morphism if for every point x of X, there exist an open neighbourhood U of x, a smooth Y-scheme V, such that the restriction of f to U factors as:



where i is a regular immersion (VII 1.4).


After replacing V by an open subset, one may suppose that i is a regular closed immersion.

In this definition one may replace V by a vector bundle $Y[T_1, \ldots, T_n]$. Indeed, since the notion is local, one may suppose that V is affine, and that its image in Y is contained in an affine open subset W of Y. There then exist an integer n and a closed W-immersion $V \longrightarrow W[T_1, \ldots, T_n]$, hence a Y-immersion $V \to Y[T_1, \ldots, T_n]$, which is regular by VII 1.10. Since the composite of two regular immersions is regular, one thus obtains a regular Y-immersion $U \longrightarrow Y[T_1, \ldots, T_n]$ factoring f.

Proposition 1.2. Let $f : X \to Y$ be a complete-intersection morphism. Then every Y-immersion $i : X \to Z$ of X into a smooth Y-scheme is regular.


We shall first prove the corollary:

Corollary 1.3. Consider the commutative diagram:



[print 467]where i and j are immersions, and p is a smooth morphism. Then for i to be a regular immersion, it is necessary and sufficient that j be a regular immersion.

i) Suppose i regular. Put $X' = V' \times_V X$. The datum of j defines a section j' of X' over X, such that $j = i' \circ j'$:



Now j' is a regular immersion, since it is a section of a smooth morphism (VII 1.10); i' is a regular immersion since it is obtained from i by a smooth, hence flat, base change (VII 1.5). Thus j is a regular immersion (VII 1.7).

ii) Suppose j regular, and p étale. Then j' is an open immersion (EGA IV 17.9.3), hence an isomorphism of X onto $j'(X)$; put $U = j'(X)$.

Let i'' be the restriction of i' to U, and let U' be an open subset of V' such that $U = U' \cap X'$. We then have: $i'' = j \circ (j')^{-1}$, so i'' is a regular immersion of U into U'. Now the morphism $U' \to V$, being smooth, is flat and of finite presentation, hence open. It is consequently a faithfully flat and quasi-compact morphism from U' onto an open neighbourhood of X in V. It follows (VII 1.5) that i is a regular immersion.

iii) Suppose j regular, and that V' is a vector bundle over V. Since the property to be proved (i regular) is local on V, one may suppose V affine, i closed, and that $V' = V[T_1, \ldots, T_n]$. Put $V = \mathrm{Spec}(A)$, [print 468] $X = \mathrm{Spec}(B)$; we have the commutative diagram:

$$\begin{array}{ccc} & & A[T_1, \ldots, T_n] \\ & \overset{v}{\swarrow} & \uparrow \\ B & \xleftarrow{\quad u \quad} & A \end{array}$$

where the morphisms u and v are surjective. Let t_i, for $i = 1, \ldots, n$, be elements of A such that for every i we have

$$u(t_i) = v(T_i).$$

One defines a homomorphism $w : A[T_1, \ldots, T_n] \longrightarrow A$ by setting $w(T_i) = t_i$. This gives a closed immersion $s : V \to V'$ which is a section of p, and which satisfies $j = s \circ i$. Moreover, being a section of a smooth morphism, it is a regular immersion (VII 1.10). Knowing that s and $s \circ i$ are regular, we shall deduce that i is regular if we show that the sequence:

$$0 \longrightarrow i^*(\underline{K}/\underline{K}^2) \longrightarrow \underline{J}/\underline{J}^2 \longrightarrow \underline{I}/\underline{I}^2 \longrightarrow 0$$

is exact and locally split, where $\underline{I}$, $\underline{J}$ and $\underline{K}$ denote respectively the ideals defining i, j, and s (VII 1.7). For this it suffices to show that $i^*(\underline{K}/\underline{K}^2)$ is a direct summand of $\underline{J}/\underline{J}^2$. Now one sees at once that j' is obtained from s by the base change $i' : X' \to V'$. Furthermore, the immersions j' and s are regular (VII 1.10). Consequently (VII 1.2), the ideal defining j' is $i'^*(\underline{K})$, and the conormal sheaf of j' is $i^*(\underline{K}/\underline{K}^2)$. The exact sequence of conormal sheaves for the decomposition $j = i' \circ j'$ therefore gives, in particular, a homomorphism: $\underline{J}/\underline{J}^2 \longrightarrow i^*(\underline{K}/\underline{K}^2)$. It is immediate to check that the composite: $i^*(\underline{K}/\underline{K}^2) \longrightarrow \underline{J}/\underline{J}^2 \longrightarrow i^*(\underline{K}/\underline{K}^2)$ is the identity (from the affine point of view, this corresponds to the decomposition of J as the direct sum of an ideal of A, and the ideal generated by the $T_i - t_i$). This indeed gives the required splitting.


[print 469] iv) Finally suppose j regular and p smooth. For every $x \in X$, one can find an open neighbourhood W of $i(x)$, an open neighbourhood W' of $j(x)$ over W, and a commutative diagram:

$$\begin{array}{ccc} W' & \xrightarrow{\quad q \quad} & W[T_1, \ldots, T_n] \\ \Big\downarrow{\scriptstyle p} & \swarrow_{p_0} & \\ W & & \end{array} \quad ,$$


where p_0 is the structural morphism, and q an étale morphism (EGA IV 17.11.4). One may then suppose $V = W$, $V' = W'$. It is easy to see that $q \circ j$ is an immersion (introduce the fiber product $X[T_1, \ldots, T_n]$). Applying the results ii) and iii), one deduces that $q \circ j$ is regular[473], and then that i is regular.

Let us now prove 1.2.

Since the property is local on Z, to prove it one may work in a neighbourhood of a point x of X. One can then find an open neighbourhood U of x in X, a smooth Y-scheme V, and a regular Y-immersion $i' : U \to V$. We then have the following commutative diagram:



This gives a closed immersion $i'' : U \to Z \times_Y V$, and the commutative diagram:




[print 470]Since i' is regular, and p and p' are smooth, it follows by 1.3 that i'' and i are regular.

The following proposition will not be used in the rest of the Seminar:

Proposition 1.4. Let $f : X \to Y$ be a flat morphism locally of finite presentation. For f to be a complete-intersection morphism in the sense of 1.1, it is necessary and sufficient that it be a complete-intersection morphism in the sense of EGA IV 19.3.6.

Suppose that f is a complete-intersection morphism in the sense of 1.1, and let $x \in X$. Then there exist an open neighbourhood U of x, and a factorization:




where i is a regular immersion, and g a smooth morphism. Let $y = f(x)$; then the immersion $i_y : U_y \to V_y$ is regular at x. Indeed, after restricting V, one may suppose that there exist a free $\underline{O}_V$-Module of finite type $\underline{E}$, and a regular homomorphism $u : \underline{E} \to \underline{O}_V$ such that the Koszul complex associated to u is a resolution of $\underline{O}_U$. Consider then the homomorphism u_y induced on the fiber at y, and the associated Koszul complex $\underline{K}_\bullet(u_y) = \underline{K}_\bullet(u)_y$. Since g is smooth, hence flat, $\underline{K}_\bullet(u)$ is a Y-flat resolution of $\underline{O}_U$. Thus the homology groups of $\underline{K}_\bullet(u)_y$ are the $\underline{\mathrm{Tor}}_i^{\underline{O}_Y}(\underline{O}_U, \underline{k}(y))$, where $\underline{k}(y)$ is the residue field of $\underline{O}_{Y,y}$. Since f is a flat morphism, they vanish, and i_y is regular at x. Furthermore, since g is smooth, the local ring of V_y at x is regular. Since the local ring of U_y at x is the quotient[474] of a regular ring by a regular ideal, [print 471]it is a complete-intersection ring (EGA IV 19.3.2), and therefore f is a complete intersection at x in the sense of EGA IV 19.3.6.

Conversely, suppose that f is a complete-intersection morphism in the sense of EGA IV 19.3.6, and let $x \in X$. Let U be an affine neighbourhood of x, over an affine neighbourhood U' of $y = f(x)$. One can then find a smooth U'-scheme V, and a closed immersion $U \xrightarrow{i} V$ (for example by taking for V a vector bundle over U'). We shall show that i is regular. Now by EGA


IV 19.3.7, i is transversally regular at x, since V is smooth over Y, hence V_y is regular at x, and U' is flat over Y. It then follows from EGA IV 19.2.4, a)⇒c), that i is regular at x. Thus f is a complete-intersection morphism in the sense of 1.1.

Proposition 1.5. Let $f : X \to Y$, and $g : Y \to Z$ be two locally complete-intersection morphisms. Then $g \circ f$ is a locally complete-intersection morphism.

Let $x \in X$. One can find open neighbourhoods U, V of $x, f(x)$, and regular immersions i, j such that we have the factorizations:



This diagram can be completed by the commutative square: [print 472]




where j' is obtained from j by base change; since this base change is flat, j' is a regular immersion, and consequently $j' \circ i$ is a regular immersion of U into a scheme smooth over Z[475], giving the result.

Proposition 1.6. Let $f : X \to Y$ be a morphism and[476] $g : Y' \to Y$ a flat base change. If f is a complete intersection, then the morphism $f' : X \times_Y Y' \to Y'$ obtained from f is one as well. The converse is true if g is also quasi-compact and surjective.

One can find a covering of X by open subsets U_i such that for every i, the restriction of f to U_i factors as $U_i \longrightarrow V_i \longrightarrow Y$, where $U_i \to V_i$ is a regular immersion, and $V_i \to Y$ a smooth morphism. The $U_i \times_Y Y'$ form an open covering of $X \times_Y Y'$, the immersions $U_i \times_Y Y' \longrightarrow V_i \times_Y Y'$ are regular by VII 1.5, and the morphisms $V_i \times_Y Y' \to Y'$ are smooth, giving the first result. For the converse, EGA IV 2.7.1 (iv) first implies that f is locally of finite presentation, which easily reduces us, using 1.2, to the case where f is a closed immersion, in which case the assertion is precisely VII 1.5.


Proposition 1.7. Let $f : X \to Y$ be a complete-intersection morphism. Then f is a perfect morphism.

Since the property is local on X, one may work in a neighbourhood of a point x of X, and suppose that there exist an open subset U containing x, and a factorization: [print 473]



where i is a regular immersion, and g a smooth morphism. Since g is smooth, it is a perfect morphism (III 4.1). By VII 1.9, i is also a perfect morphism. Since the composite of two perfect morphisms is perfect (III 4.5), the proposition follows.

Proposition 1.8. Let $f : X \to Y$ be a complete-intersection morphism, x a point of X, and consider a factorization:




where U is an open neighbourhood of x, i a regular immersion, and g a smooth morphism. Denote by $\underline{J}$ the ideal of $\underline{O}_V$ defining U. Then the integer

$$d_x = \operatorname{rank}\big((\underline{\Omega}^1_{V/Y})_{i(x)}\big) - \operatorname{rank}\big((\underline{J}/\underline{J}^2)_x\big)$$


is independent of the chosen factorization of f.

Suppose that the restriction of f to U factors through two smooth Y-schemes V and V', and put $V'' = V \times_Y V'$. Then V'' is smooth over Y, and also over V and V'. Moreover, the immersions $X \to V$ and $X \to V'$ define an immersion $X \to V''$, regular by 1.2. One may therefore reduce to comparing the integers d_x associated to two factorizations of f through smooth Y-schemes V and V' such that there exists a commutative diagram over Y:



[print 474]where p is a smooth morphism. One may also suppose $\underline{\Omega}^1_{V/Y}$, $\underline{\Omega}^1_{V'/Y}$, $\underline{\Omega}^1_{V'/V}$ of constant rank, as well as the conormal sheaves of i and j.

From the commutative diagram (in which all morphisms are smooth): [477]



one deduces:

$$\operatorname{rank}(\underline{\Omega}^1_{V'/Y}) = \operatorname{rank}(\underline{\Omega}^1_{V/Y}) + \operatorname{rank}(\underline{\Omega}^1_{V'/V}). \tag{1.8.1}$$

Let $X' = V' \times_V X$; then the rank of $\underline{\Omega}^1_{V'/V}$ is equal to that of $\underline{\Omega}^1_{X'/X}$. Furthermore, j defines a section $X \xrightarrow{j'} X'$, which is a regular immersion of codimension equal to the rank of $\underline{\Omega}^1_{X'/X}$ (EGA IV 17.10.4). If i' is the immersion obtained from i by the base change $V' \to V$, we have $j = i' \circ j'$ and consequently (VII 1.7):

$$\operatorname{rank}(\underline{N}_{V'/X}) = \operatorname{rank}(\underline{N}_{V/X}) + \operatorname{rank}(\underline{\Omega}^1_{V'/V}). \tag{1.8.2}$$

Comparing (1.8.1) and (1.8.2) gives the proposition.

Definition 1.9. The integer d_x of Proposition 1.8 is called the virtual relative dimension of X over Y (or of f) at the point x.

The virtual relative dimension of X over Y is therefore a locally constant function on X, with integer values (positive or negative). If f is a smooth morphism of relative dimension n, its virtual relative dimension is n. If f is a regular immersion of codimension n, its virtual relative dimension is $-n$.

Proposition 1.10. Let $f : X \to Y$ and $g : Y \to Z$ be two complete-intersection morphisms, and let x be a point of X. Then:


$$\text{dim. rel. virt.}_x(g \circ f) = \text{dim. rel. virt.}_x(f) + \text{dim. rel. virt.}_{f(x)}(g)$$

[print 475]This follows immediately from 1.9 and from the proof of 1.5.

2. Relative cotangent complex

2.1. We shall give here a definition of the cotangent complex which, without having all the generality possible (see Exp. 0 4.4), will suffice for the needs of the Riemann–Roch theorem; of course, the definition given in exp. 0 coincides in the case considered with the definition given here.

Let $f : X \to Y$ be a smoothable morphism, that is, one admitting a factorization:



where g is a smooth morphism, and i an immersion which may be supposed closed. Let $\underline{J}$ be the ideal of X in V. The canonical derivation:

$$d : \underline{O}_V \longrightarrow \underline{\Omega}^1_{V/Y}$$

defines a homomorphism:

$$\underline{J} \longrightarrow \underline{O}_V \xrightarrow{d} \underline{\Omega}^1_{V/Y} \longrightarrow \underline{\Omega}^1_{V/Y} \otimes_{\underline{O}_V} \underline{O}_X$$

which vanishes on $\underline{J}^2$, and hence defines a homomorphism, still denoted d:

$$\underline{J}/\underline{J}^2 \xrightarrow{d} i^*(\underline{\Omega}^1_{V/Y}). \tag{2.1.1}$$


One checks easily that this homomorphism is $\underline{O}_X$-linear. One then defines a chain complex $\underline{L}^{X/Y}_\bullet$ by putting:

$$\underline{L}^{X/Y}_0 = i^*(\underline{\Omega}^1_{V/Y}), \qquad \underline{L}^{X/Y}_1 = \underline{J}/\underline{J}^2; \qquad \underline{L}^{X/Y}_i = 0 \quad \text{if } i \neq 0, 1,$$

and taking as homomorphism $\underline{L}^{X/Y}_1 \to \underline{L}^{X/Y}_0$ the homomorphism (2.1.1).[478]

[print 476]**Proposition 2.2.** The complex $\underline{L}^{X/Y}_\bullet$ does not depend, up to canonical isomorphism in the derived category $D(X)$, on the chosen factorization through a smooth Y-scheme.

Let V, V' be two smooth Y-schemes through which f factors. Put: $V'' = V \times_Y V'$; then V'' is smooth over Y, and also over V and V'. Moreover, the immersions $X \to V$ and $X \to V'$ define an immersion $X \to V''$ which factors f. One may therefore reduce to comparing the complexes $\underline{L}^{X/Y}_\bullet$ associated to two factorizations through smooth Y-schemes V and V', such that there exists a commutative diagram over Y:



where i and j are closed immersions, and p a smooth morphism.

By functoriality, one then obtains homomorphisms:

$$\underline{I}/\underline{I}^2 \longrightarrow \underline{J}/\underline{J}^2; \qquad p^*(\underline{\Omega}^1_{V/Y}) \longrightarrow \underline{\Omega}^1_{V'/Y},$$

hence a diagram:

$$\begin{array}{ccc} \underline{I}/\underline{I}^2 & \longrightarrow & \underline{J}/\underline{J}^2 \\ {\scriptstyle d}\downarrow & & \downarrow{\scriptstyle d} \\ i^*(\underline{\Omega}^1_{V/Y}) & \longrightarrow & j^*(\underline{\Omega}^1_{V'/Y}). \end{array} \tag{2.2.1}$$

Since V and V' are smooth over Y, one obtains the exact sequence:

$$0 \longrightarrow p^*(\underline{\Omega}^1_{V/Y}) \longrightarrow \underline{\Omega}^1_{V'/Y} \longrightarrow \underline{\Omega}^1_{V'/V} \longrightarrow 0.$$

Since the Modules of differentials considered are locally free over $\underline{O}_{V'}$[479], applying the functor j^* gives the exact sequence:

$$0 \longrightarrow i^*(\underline{\Omega}^1_{V/Y}) \longrightarrow j^*(\underline{\Omega}^1_{V'/Y}) \longrightarrow j^*(\underline{\Omega}^1_{V'/V}) \longrightarrow 0. \tag{2.2.2}$$

[print 477]Furthermore, j defines a section j' of $X' = V' \times_V X$:

$$\begin{array}{ccc} X' & \xrightarrow{i'} & V' \\ {\scriptstyle p'}\downarrow\uparrow{\scriptstyle j'} & \nearrow{\scriptstyle j} & \downarrow{\scriptstyle p} \\ X & \xrightarrow{i} & V. \end{array}$$

Let $\underline{I}, \underline{J}, \underline{K}$ be the ideals defining i, j, j'; we have: $j = i' \circ j'$, and the immersion j' is regular[480]; moreover, i' is defined by the ideal $p^*(\underline{I})$ (since p is flat), and its conormal sheaf is therefore $i'^*(p^*(\underline{I})) = p'^*(\underline{I}/\underline{I}^2)$. One then obtains the exact sequence:

$$0 \longrightarrow \underline{I}/\underline{I}^2 \longrightarrow \underline{J}/\underline{J}^2 \longrightarrow \underline{K}/\underline{K}^2 \longrightarrow 0. \tag{2.2.3}$$

We shall show:

a) that the diagram (2.2.1) is commutative; this will thus give a morphism of complexes θ from the complex $\underline{L}^{X/Y}_\bullet$ defined by V to the complex $\underline{L}^{X/Y}_\bullet$ defined by V';

b) that the homomorphism: $\underline{K}/\underline{K}^2 \longrightarrow j^*(\underline{\Omega}^1_{V'/V})$ defined by (2.2.2), (2.2.3) and (2.2.1) is an isomorphism; this will imply that the mapping cylinder of θ is acyclic, hence that θ is a quasi-isomorphism, and therefore defines an isomorphism in the derived category.

The properties a) and b) are local on X. One may therefore suppose X, Y, V and V' affine. Put: $Y = \mathrm{Spec}(A)$, $V = \mathrm{Spec}(B)$, $V' = \mathrm{Spec}(B')$, $X = \mathrm{Spec}(C)$, and keep the same letters to denote the corresponding homomorphisms and ideals. We have the diagram: [print 478]

$$\begin{array}{ccccc} B \otimes_A B & \xrightarrow{p\otimes p} & B' \otimes_A B' & & \\ {\scriptstyle\Delta}\downarrow & (1) & \downarrow{\scriptstyle\Delta} & & \\ B & \xrightarrow{p} & B' & \xleftarrow{\Delta} & B' \otimes_B B' \\ & {\scriptstyle i}\searrow & {\scriptstyle j}\downarrow & (2) \quad {\scriptstyle i'}\swarrow & \downarrow \\ & & C & \underset{p'}{\overset{j'}{\rightleftarrows}} & B' \otimes_B C = B'/IB'. \end{array}$$

The homomorphism $\underline{I}/\underline{I}^2 \to \underline{J}/\underline{J}^2$ is obtained from the homomorphism: $IB' \to J$ and coincides with the one obtained from $p : I \to J$; the homomorphism: $i^*(\underline{\Omega}^1_{V/Y}) \longrightarrow j^*(\underline{\Omega}^1_{V'/Y})$ is obtained from the commutative square (1). Property a) is therefore evident.

The homomorphism: $\underline{K}/\underline{K}^2 \to j^*(\underline{\Omega}^1_{V'/V})$ considered in b) can be defined as follows: let $\overline{x} \in K/K^2$, and let x be an element of J lifting $\overline{x}$ (one exists since $K = J/IB'$). To $\overline{x}$ one associates the image of x under the composite homomorphism: $J \xrightarrow{d} \Omega_{B'/A} \otimes_{B'} C \longrightarrow \Omega^1_{B'/B} \otimes_{B'} C$. Furthermore, let H be the augmentation ideal of: $B' \otimes_B B' \longrightarrow B'$; the commutative square (2) gives a homomorphism: $H \to K$, hence a homomorphism: $H/H^2 \otimes_{B'} C \longrightarrow K/K^2$. Now $H/H^2 = \Omega^1_{B'/B}$; consequently, it suffices to check that the homomorphisms constructed are inverse to each other, which is done without difficulty.

Definition 2.3. Let $f : X \to Y$ be a smoothable morphism. The complex $\underline{L}^{X/Y}_\bullet$, defined up to canonical isomorphism in the derived category $D(X)$, is called the relative cotangent complex of X over Y (or of f).

Proposition 2.4. Let $f : X \to Y$ be a complete-intersection morphism (1.1) which is smoothable. Then $\underline{L}^{X/Y}_\bullet$ is perfect. More precisely, for every [print 479]factorization of f as the composite of an immersion and a smooth morphism, the complex $\underline{L}^{X/Y}_\bullet$ defined in 2.1 is strictly perfect.

Let $f = g \circ i$, where g is a smooth morphism, and i an immersion. By 1.2, i is a regular immersion. If $\underline{J}$ is the Ideal defining i, then $\underline{J}/\underline{J}^2$ is a locally free $\underline{O}_X$-Module of finite type. Similarly, since $g : V \to Y$ is smooth, $i^*(\underline{\Omega}^1_{V/Y})$ is locally free of finite type, giving 2.4.

Corollary 2.5. Let $f : X \to Y$ be a smoothable complete-intersection morphism, and $f = g \circ i$ a factorization of f through a smooth Y-scheme V. Let $\underline{J}$ be the ideal of $\underline{O}_V$ defining i. Then, if $K^\bullet(X)$ denotes the Grothendieck group of locally free $\underline{O}_X$-Modules of finite type (resp. of perfect complexes on X), the element $T_f \in K^\bullet(X)$ defined by:

$$T_f = \mathrm{cl}(i^*(\underline{\Omega}^1_{V/Y})) - \mathrm{cl}(\underline{J}/\underline{J}^2)$$

$$(\text{resp. } T_f = \mathrm{cl}(\underline{L}^{X/Y}_\bullet))$$

does not depend on the chosen factorization of f. The value at a point x of X of the augmentation of T_f is precisely the virtual relative dimension of X over Y at the point x (1.9).

The element T_f is well defined in both cases considered by 2.4, and its class in $K^\bullet(X)$ is independent of the factorization, by 2.2 in the second case, and by the proof of 2.2 in the first case. The last assertion of 2.5 is trivial.


Proposition 2.6. Let $f : X \to Y$, and $g : Y \to Z$ be two smoothable complete-intersection morphisms, and suppose in addition that the following condition is satisfied:

[print 480](G) There exists a smooth Y-scheme V, factoring f, such that for every closed immersion: $Y \to V'$ one can find a smooth V'-scheme V'', with $V \simeq V'' \times_{V'} Y$ (this condition is satisfied in particular if f is quasi-projective and if Y has an ample sheaf, by taking for V a $\mathbf{P}^r_Y$[481]).

Then $g \circ f$ is a complete intersection, is smoothable, and we have a distinguished triangle: [482]

$$\begin{array}{ccc} & \underline{L}^{X/Y}_\bullet & \\ \swarrow & & \nwarrow \\ \mathbb{L}f^*(\underline{L}^{Y/Z}_\bullet) & \longrightarrow & \underline{L}^{X/Z}_\bullet . \end{array}$$

Let V' be a smooth Z-scheme factoring g. Consider the commutative diagram:



Then the immersion j' is regular, since it is obtained from j by a flat base change. Consequently $j' \circ i$ is a regular immersion factoring $g \circ f$, which is therefore smoothable and a complete intersection. Moreover, we have the exact sequence of conormal sheaves:

$$0 \longrightarrow i^*(\underline{J}'/\underline{J}'^2) \longrightarrow \underline{K}/\underline{K}^2 \longrightarrow \underline{I}/\underline{I}^2 \longrightarrow 0,$$

where $\underline{I}, \underline{J}, \underline{J}', \underline{K}$ denote the sheaves defining respectively the immersions $i, j, j', j' \circ i$. Now, since p'' is flat, $\underline{J}' = p''^*(\underline{J})$. The sequence can therefore be written:

$$0 \longrightarrow f^*(\underline{J}/\underline{J}^2) \longrightarrow \underline{K}/\underline{K}^2 \longrightarrow \underline{I}/\underline{I}^2 \longrightarrow 0. \tag{2.6.1}$$

Furthermore, since V' and V'' are smooth over Z, we have the exact sequence:

$$0 \longrightarrow p''^*(\underline{\Omega}^1_{V'/Z}) \longrightarrow \underline{\Omega}^1_{V''/Z} \longrightarrow \underline{\Omega}^1_{V''/V'} \longrightarrow 0.$$

[print 481]Applying $i^* \circ j'^*$ gives the exact sequence:

$$0 \longrightarrow (p'' \circ j' \circ i)^*(\underline{\Omega}^1_{V'/Z}) \longrightarrow (j' \circ i)^*(\underline{\Omega}^1_{V''/Z}) \longrightarrow (j' \circ i)^*(\underline{\Omega}^1_{V''/V'}) \longrightarrow 0.$$

Since $j'^*(\underline{\Omega}^1_{V''/V'}) = \underline{\Omega}^1_{V/Y}$ by hypothesis (G), we obtain:

$$0 \longrightarrow f^*(j^*(\underline{\Omega}^1_{V'/Z})) \longrightarrow (j' \circ i)^*(\underline{\Omega}^1_{V''/Z}) \longrightarrow i^*(\underline{\Omega}^1_{V/Y}) \longrightarrow 0. \tag{2.6.2}$$

Combining (2.6.1) and (2.6.2), we are reduced to checking that the diagram:

$$\begin{array}{ccccccccc}
0 & \longrightarrow & f^*(\underline{J}/\underline{J}^2) & \longrightarrow & \underline{K}/\underline{K}^2 & \longrightarrow & \underline{I}/\underline{I}^2 & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle f^*(d)} & & \downarrow{\scriptstyle d} & & \downarrow{\scriptstyle d} & & \\
0 & \longrightarrow & f^*(j^*(\underline{\Omega}^1_{V'/Z})) & \longrightarrow & (j' \circ i)^*(\underline{\Omega}^1_{V''/Z}) & \longrightarrow & i^*(\underline{\Omega}^1_{V/Y}) & \longrightarrow & 0
\end{array}$$

is commutative, which is done locally without difficulty by returning to the definitions of the morphisms involved.

From the resulting exact sequence of complexes, one deduces the announced distinguished triangle in the derived category.


Corollary 2.7. With the notation and hypotheses of 2.6, we have in $K^\bullet(X)$ the following equality:

$$T_{g \circ f} = T_f + f^*(T_g).$$

3. Riemann–Roch theorem: statement

3.1. In section 3, Y is a quasi-compact scheme having an ample sheaf.

Let $f : X \to Y$ be a projective complete-intersection morphism (1.1); suppose in addition that the virtual relative dimension of f (1.9) [print 482]is constant on X. Since f is projective, there exists a closed immersion $X \to P$, where P is a projective bundle over Y, and since Y has an ample sheaf, one may suppose that $P = \mathbf{P}^r_Y$.[483] We thus have a factorization: [484]



Since f is a complete intersection, the immersion i is regular (1.2).

By 1.7, f is a perfect morphism. Moreover, Y has an ample sheaf, and consequently so do X and P. One can then define an additive homomorphism:

$$f_* : K^\bullet(X) \longrightarrow K^\bullet(Y)$$

(IV 2.12), the $K^\bullet$ defined in terms of locally free sheaves of finite type, or of perfect complexes, being identical. One can also define additive homomorphisms:

$$i_* : K^\bullet(X) \longrightarrow K^\bullet(P); \qquad g_* : K^\bullet(P) \longrightarrow K^\bullet(Y),$$

and we have: $f_* = g_* \circ i_*$.

Let d be the virtual relative dimension of X over Y. The codimension of i is then $r - d$. By VII 4.6, the homomorphism i_* gives, after tensoring with $\mathbb{Q}$, inclusions:

$$i_*(\mathrm{Fil}^k K^\bullet(X)_\mathbb{Q}) \subset \mathrm{Fil}^{k+r-d} K^\bullet(P)_\mathbb{Q}.$$

By VI 5.8, the homomorphism g_* gives inclusions:

$$g_*(\mathrm{Fil}^{k'} K^\bullet(P)) \subset \mathrm{Fil}^{k'-r} K^\bullet(Y).$$

Composing the two, we thus obtain:

[print 483]**Proposition 3.2.** Let $f : X \to Y$ be a projective morphism, a complete intersection, whose virtual relative dimension d is constant. Suppose that Y is quasi-compact, and has an ample sheaf. Then for every $k \in \mathbb{Z}$ we have:

$$f_*(\mathrm{Fil}^k K^\bullet(X)_\mathbb{Q}) \subset \mathrm{Fil}^{k-d} K^\bullet(Y)_\mathbb{Q}.$$

We thus obtain, in particular, an additive homomorphism:

$$f_{\mathrm{gr}} : \mathrm{Gr}\, K^\bullet(X)_\mathbb{Q} \longrightarrow \mathrm{Gr}\, K^\bullet(Y)_\mathbb{Q}$$

sending $\mathrm{Gr}^k K^\bullet(X)_\mathbb{Q}$ into $\mathrm{Gr}^{k-d} K^\bullet(Y)_\mathbb{Q}$.

3.3. Recall that in (V 6.1) we defined a functor $A \longmapsto \mathrm{Ch}(A)$ on the category of graded K-algebras, with values in the category of augmented K-λ-algebras, where K is a binomial ring (V 2.7). If A is a graded K-algebra, we have:

$$\mathrm{Ch}(A) = K \times (1 + \widehat{A}^+).$$

For every augmented K-λ-algebra Λ, we defined a K-λ-homomorphism, called the "completed Chern class" (V 6.7):

$$\widetilde{c} = (\varepsilon, c) : \Lambda \longrightarrow \mathrm{Ch}(\mathrm{Gr}_\lambda \Lambda).$$

Finally, we defined a ring homomorphism, functorial in A, called the Chern character (V 6.4):

$$\mathrm{ch} : \mathrm{Ch}(A) \longrightarrow K \oplus \widehat{A}^+_{\mathbb{Q}},$$

and a group endomorphism, called the Todd character (V 1.16):

$$\mathrm{Todd} : 1 + \widehat{A}^+ \longrightarrow 1 + \widehat{A}^+.$$

[print 484]We shall apply these definitions to the augmented λ-rings $K^\bullet(X)$ and $K^\bullet(Y)$, the augmentation rings $H^0(X, \underline{\mathbb{Z}})$ and $H^0(Y, \underline{\mathbb{Z}})$ being binomial (V 2.8). To simplify notation, we shall put, for every scheme Z and every $z \in K^\bullet(Z)$:

$$\mathrm{ch}(z) = \mathrm{ch}(\widetilde{c}(z)); \qquad \mathrm{Todd}(z) = \mathrm{Todd}(c(z)).$$

For every graded algebra A, we shall regard the group $1 + \widehat{A}^+$ as contained in $\widehat{A}$. Finally, given a homomorphism $f : A \to B$ which is additive and compatible with the gradings up to a shift, we shall still denote by f the homomorphism $\widehat{A} \to \widehat{B}$ obtained from it.

3.4. Let X be a scheme. For every locally free $\underline{O}_X$-Module of finite type $\underline{E}$, denote by $\check{E}$ the class in $K^\bullet(X)$ of the dual $\underline{\check{E}}$ of $\underline{E}$. It is clear that the function associating $\check{E}$ to $\underline{E}$ is additive on the category of locally free $\underline{O}_X$-Modules of finite type, and therefore defines an endomorphism of $K^\bullet(X)$, which we shall denote by $x \rightsquigarrow \check{x}$.

Proposition 3.5. Let X be a quasi-compact scheme. Then for every $x \in K^\bullet(X)$ and every $i \geq 0$, we have:

$$c^i(\check{x}) = (-1)^i c^i(x).$$

The relation to be proved may also be written:

$$c_t(\check{x}) = c_{-t}(x),$$

where we put: $c_t(x) = \sum_{i \geq 0} c^i(x) t^i$.

By additivity, one may therefore suppose that x is the class of a locally free $\underline{O}_X$-Module of finite type $\underline{E}$. Moreover, after replacing X by the flag scheme of $\underline{E}$, which gives an injection on the [print 485]$\mathrm{Gr}\, K^\bullet$ (VI 5.5), one may suppose that $\underline{E}$ is an invertible sheaf. We then have: $\underline{\check{E}} = \underline{E}^{-1}$; hence $\check{x} = x^{-1}$.

The $c^i(\check{x})$, $c^j(x)$ vanish for $i, j \geq 2$. It therefore suffices to show that $c^1(x^{-1}) = -c^1(x)$. Now $c^1(x)$ is the class in $\mathrm{Gr}^1 K^\bullet(X)$[485] of $x - 1$, and $c^1(x^{-1})$ that of $x^{-1} - 1$. The equality:

$$x^{-1} - 1 = 1 - x + (x - 1)(1 - x^{-1})$$

gives:

$$x^{-1} - 1 \equiv 1 - x \qquad \text{mod. } \mathrm{Fil}^2 K^\bullet(X);$$

the result follows.

Theorem 3.6. (Riemann–Roch). Let $f : X \to Y$ be a projective complete-intersection morphism (1.1), whose virtual relative dimension d is constant (1.9). Suppose that Y is quasi-compact, and has an ample sheaf. With the notation of 3.3, we then have for every $x \in K^\bullet(X)$ the relation:

$$\mathrm{ch}(f_*(x)) = f_{\mathrm{gr}}\big(\mathrm{ch}(x) \cdot \mathrm{Todd}(\check{T}_f)\big),$$

$\check{T}_f$ denoting the dual element of the class of the relative cotangent complex $\underline{L}_\bullet^{X/Y}$ (2.5 and 2.3). In other words, the diagram

$$\begin{array}{ccc} K^\bullet(X) & \xrightarrow{\mathrm{ch}\circ\widetilde{c}} & \mathrm{Gr}\, K^\bullet(X)_{\mathbb{Q}} \\ {\scriptstyle f_*}\downarrow & & \downarrow{\scriptstyle x\mapsto f_{\mathrm{gr}}(x\cdot\mathrm{Todd}(\check{T}_f))} \\ K^\bullet(Y) & \xrightarrow{\mathrm{ch}\circ\widetilde{c}} & \mathrm{Gr}\, K^\bullet(Y)_{\mathbb{Q}} \end{array}$$

is commutative.

[print 486]The proof of Theorem 3.6 will occupy the rest of this exposé. Let us give some preliminaries.

Proposition 3.7. For arbitrary $x \in \mathrm{Gr}\, K^\bullet(X)_{\mathbb{Q}}$, $y \in \mathrm{Gr}\, K^\bullet(Y)_{\mathbb{Q}}$, we have:

$$f_{\mathrm{gr}}\big(x \cdot f^{\mathrm{gr}}(y)\big) = f_{\mathrm{gr}}(x)\, y.$$

By additivity, one may suppose x and y homogeneous; say $x \in \mathrm{Gr}^i\, K^\bullet(X)_{\mathbb{Q}}$, $y \in \mathrm{Gr}^j\, K^\bullet(Y)_{\mathbb{Q}}$. Let $x' \in \mathrm{Fil}^i\, K^\bullet(X)_{\mathbb{Q}}$, $y' \in \mathrm{Fil}^j\, K^\bullet(Y)_{\mathbb{Q}}$ be elements lifting x and y. Then the projection formula gives:

$$f_*(x' \cdot f^*(y')) = f_*(x')\, y'.$$

By 3.2, this equality is written in $\mathrm{Fil}^{i+j-d}\, K^\bullet(Y)_{\mathbb{Q}}$. It gives an equality in $\mathrm{Gr}^{i+j-d}\, K^\bullet(Y)_{\mathbb{Q}}$, and one checks at once that this is indeed the required equality.

Lemma 3.8. Let $f : X \to Y$, and $g : Y \to Z$ be two morphisms satisfying the hypotheses of 3.6. Then $g \circ f$ also satisfies them. If x is an element of $K^\bullet(X)$, and if 3.6 holds for f and the element x, and for g and the element $f_*(x)$, then it holds for $g \circ f$ and the element x.

By hypothesis, we have:

$$\mathrm{ch}(f_*(x)) = f_{\mathrm{gr}}\big(\mathrm{ch}(x) \cdot \mathrm{Todd}(\check{T}_f)\big), \tag{3.8.1}$$

and:

$$\mathrm{ch}(g_*(f_*(x))) = g_{\mathrm{gr}}\big(\mathrm{ch}(f_*(x)) \cdot \mathrm{Todd}(\check{T}_g)\big). \tag{3.8.2}$$

Combining (3.8.1) and (3.8.2), we obtain:

$$\mathrm{ch}(g_*(f_*(x))) = g_{\mathrm{gr}}\Big(f_{\mathrm{gr}}\big(\mathrm{ch}(x)\, \mathrm{Todd}(\check{T}_f)\big) \cdot \mathrm{Todd}(\check{T}_g)\Big).$$

[print 487]Applying 3.7, we find:

$$\mathrm{ch}(g_*(f_*(x))) = g_{\mathrm{gr}}\Big(f_{\mathrm{gr}}\big(\mathrm{ch}(x) \cdot \mathrm{Todd}(\check{T}_f) \cdot f^{\mathrm{gr}}(\mathrm{Todd}(\check{T}_g))\big)\Big).$$

The functoriality of c and of Todd implies:

$$f^{\mathrm{gr}}(\mathrm{Todd}(\check{T}_g)) = \mathrm{Todd}(f^*(\check{T}_g)).$$

By 2.7, we have:

$$\check{T}_{g\circ f} = \check{T}_f + f^*(\check{T}_g).$$

The multiplicativity of the Todd character therefore implies:

$$\mathrm{Todd}(\check{T}_f) \cdot f^{\mathrm{gr}}(\mathrm{Todd}(\check{T}_g)) = \mathrm{Todd}(\check{T}_{g\circ f}).$$

Thus finally we obtain:

$$\mathrm{ch}(g_*(f_*(x))) = g_{\mathrm{gr}}\big(f_{\mathrm{gr}}\big(\mathrm{ch}(x) \cdot \mathrm{Todd}(\check{T}_{g\circ f})\big)\big),$$

which is the announced result.

3.9. Applying 3.8 to the factorization of f given in 3.1, one sees that, to prove the Riemann–Roch theorem, it suffices to prove it in the case of a regular immersion of constant codimension, and in the case of the structural morphism: $\mathbf{P}^r_Y \longrightarrow Y$[486].

Furthermore, one may suppose that the immersion i satisfies the condition of VII 4.1. Indeed, let s be the section of $\mathbf{P}^2_{\mathbf{P}^r_Y}$[487] introduced in VII 4.2. Then $s \circ i$ is a regular immersion satisfying the condition of VII 4.1 (VII 4.2), and by 3.8 it suffices to prove 3.6 for $s \circ i$, $\mathbf{P}^2_{\mathbf{P}^r_Y} \longrightarrow \mathbf{P}^r_Y$[488] and $\mathbf{P}^r_Y \longrightarrow Y$[489].

We shall therefore prove 3.6 by proving the following two special cases:

i) f is a regular immersion of constant codimension, satisfying the condition of VII 4.1; [print 488]

ii) f is the structural morphism $\mathbf{P}^r_Y \to Y$[490].

4. Riemann–Roch theorem: case of a regular closed immersion

4.1. Let $i : Y \to X$ be a regular closed immersion. The Riemann–Roch theorem for i is essentially relation VII 4.4, which makes it possible to compute the Chern classes of $i_*(x)$ from the $\gamma^p(N, x)$. More precisely, if one introduces the λ-ring $K^\bullet(Y)_{\mathbb{Q}N}$ defined in V 5.5, then the homomorphism from $K^\bullet(Y)_{\mathbb{Q}N}$ to $K^\bullet(X)_{\mathbb{Q}}$ obtained from i_* is a λ-homomorphism by VII 2.8 and VII 4.3. The Riemann–Roch theorem then reduces to an analogous formula concerning the (additive) homomorphism $K^\bullet(Y) \longrightarrow K^\bullet(Y)_{\mathbb{Q}N}$ defined by $y \rightsquigarrow (0, y)$, that is, to a formal relation valid for every λ-ring. We shall not develop this proof here, in order to avoid still more formalism on λ-rings (cf. RRR I 1.5), and shall return to the method of [1], which used blow-up schemes (a method already used in the proof of VII 4.3, and which is also that of the end of RRR).

As indicated in 3.9, one may restrict to the case of a regular immersion of constant codimension d satisfying the condition of VII 4.1, the case of a general regular immersion then following from 3.8. In this section we return to the notation of VII 3.1. We thus have the cartesian square:

$$\begin{array}{ccc} Y' & \xrightarrow{\ j\ } & X' \\ {\scriptstyle g}\downarrow & & \downarrow{\scriptstyle f} \\ Y & \xrightarrow{\ i\ } & X, \end{array}$$

[print 489]where X' is the X-scheme obtained by blowing up Y. We suppose in addition that X is quasi-compact and has an ample sheaf.

Proposition 4.2. The morphism f is projective, a complete intersection, and of virtual relative dimension zero. We also have the relation:

$$f_{\mathrm{gr}} \circ f^{\mathrm{gr}} = \mathrm{Id}_{\mathrm{Gr}\, K^\bullet(X)_{\mathbb{Q}}}.$$

The first assertion follows immediately from (VII 1.8).

The second relation follows from the relation:

$$f_* \circ f^* = \mathrm{Id}_{K^\bullet(X)},$$

proved in (VII 3.6).

Proposition 4.3. For every $y \in \mathrm{Gr}\, K^\bullet(Y)_{\mathbb{Q}}$[491], we have the relation:

$$f^{\mathrm{gr}} \circ i_{\mathrm{gr}}(y) = j_{\mathrm{gr}}(g^{\mathrm{gr}}(y) \cdot c^{d-1}(\check{F})),$$

where c^{d-1} denotes the $(d-1)$-th Chern class.

This relation is contained in VII 4.8, since it is precisely $v^{\mathrm{gr}} \circ u^{\mathrm{gr}} = 0$.

Proposition 4.4. Let X be a quasi-compact scheme. If E is a locally free sheaf of finite type on X, of class E in $K^\bullet(X)$, and of rank n, we have:

$$\mathrm{ch}(\lambda_{-1}(E)) = c^n(\check{E})(\mathrm{Todd}(\check{E}))^{-1}.$$

The two sides of the relation to be proved are multiplicative: this is clear for $\mathrm{ch} \circ \lambda_{-1}$, and for Todd; it also holds for c^n, which is the class of $(-1)^{\varepsilon(E)}\lambda_{-1}(E)$. Consequently, by passing to the flag bundle of E, one may suppose that E is a sum of classes of invertible sheaves, and reduce to proving the formula for an invertible sheaf. Let [print 490]ξ be the class of such a sheaf. We then have:

$$\mathrm{ch}(\lambda_{-1}(\xi)) = \mathrm{ch}(1-\xi) = 1 - \mathrm{ch}(\xi).$$

Since $\widetilde{c}(\xi) = (1, 1 + c^1(\xi))$, the definition of ch gives:

$$\mathrm{ch}(\lambda_{-1}(\xi)) = 1 - \exp(c^1(\xi)).$$

Furthermore, ξ has augmentation 1; the Chern class occurring in formula 4.4 for ξ is therefore c^1. Taking into account the definition of Todd, and 3.5, the relation to be proved is written:

$$1 - \exp(c^1(\xi)) = -c^1(\xi) \cdot \left(\frac{-c^1(\xi)}{1 - \exp(c^1(\xi))}\right)^{-1},$$

and is therefore verified.

Remark 4.4.1. It is also possible to deduce 4.4 from a corresponding purely algebraic statement, cf. RRR I 1.3.

4.5. We shall now show that, to prove the Riemann–Roch theorem in the case considered in this section, one may reduce to the case where Y is a divisor of X, and where the element to which the formula applies is the inverse image of an element of $K^\bullet(X)$.

By 4.2, the formula to be proved:[492]

$$\mathrm{ch}(i_*(x)) = i_{\mathrm{gr}}(\mathrm{ch}(x) \cdot \mathrm{Todd}(\check{T}_i))$$

is equivalent to the formula:

$$f^{\mathrm{gr}}(\mathrm{ch}(i_*(x))) = f^{\mathrm{gr}} \circ i_{\mathrm{gr}}(\mathrm{ch}(x) \cdot \mathrm{Todd}(\check{T}_i)).$$

Taking into account the functoriality of ch, and 4.3, it may also be written:

$$\mathrm{ch}(f^*(i_*(x))) = j_{\mathrm{gr}}(g^{\mathrm{gr}}(\mathrm{ch}(x) \cdot \mathrm{Todd}(\check{T}_i)) \cdot c^{d-1}(\check{F})),$$

or, by VII 3.4:

$$\mathrm{ch}(j_*(g^*(x) \cdot \lambda_{-1}(F))) = j_{\mathrm{gr}}(\mathrm{ch}(g^*(x))^{[493]} \cdot \mathrm{Todd}(g^*(\check{T}_i)) \cdot c^{d-1}(\check{F})).$$

[print 491]Now by definition we have:

$$T_i = -\,\mathrm{cl}(\underline{J}/\underline{J}^2) = -N; \qquad T_j = -\,\mathrm{cl}(\underline{J}'/\underline{J}'^2) = -L.$$

Furthermore, the exact sequence:

$$0 \longrightarrow \underline{F} \longrightarrow g^*(\underline{N}) \longrightarrow \underline{O}_{Y'}(1) \longrightarrow 0$$

gives the relation:

$$g^*(T_i) = -(F + L),$$

from which we deduce:

$$\mathrm{Todd}(g^*(\check{T}_i)) = \mathrm{Todd}(\check{T}_j) \cdot (\mathrm{Todd}(\check{F}))^{-1}.$$

Taking 4.4 into account, the relation to be proved is therefore written:

$$\mathrm{ch}\big(j_*(g^*(x) \cdot \lambda_{-1}(F))\big) = j_{\mathrm{gr}}\big(\mathrm{ch}(g^*(x) \cdot \lambda_{-1}(F)) \cdot \mathrm{Todd}(\check{T}_j)\big).$$

Putting $z = g^*(x) \cdot \lambda_{-1}(F)$, this relation is precisely the Riemann–Roch theorem applied to the regular immersion j of codimension 1, and to the element z of $K^\bullet(Y')$.

Moreover, since we supposed that i satisfies the hypothesis of (VII 4.1), $\lambda_{-1}(F)$ is divisible by $(1 - L)$. One can therefore find z' such that $z = z'(1 - L)$, or equivalently $z = j^*(j_*(z'))$ (exp. VII 2.7). We are thus reduced to the case where Y is a divisor, and to an element x such that there exists y, with $x = i^*(y)$.

4.6. We must therefore show the relation:

$$\mathrm{ch}\big(i_*(i^*(y))\big) = i_{\mathrm{gr}}\big(\mathrm{ch}(i^*(y)) \cdot \mathrm{Todd}(\check{T}_i)\big).$$

Now the projection formula gives:

$$i_*(i^*(y)) = y \cdot i_*(1),$$

and from the exact sequence: $0 \longrightarrow \underline{J} \longrightarrow \underline{O}_X \longrightarrow \underline{O}_Y \longrightarrow 0$, where $\underline{J}$ is invertible here, we obtain:

$$i_*(1) = 1 - J,$$

hence [print 492]

$$\mathrm{ch}(i_*(i^*(y))) = \mathrm{ch}(y) \cdot \mathrm{ch}(1 - J).$$

Since $\underline{J}$ is invertible, we have $\widetilde{c}(J) = (1, 1 + c^1(J))$, hence $\mathrm{ch}(J) = \exp(c^1(J))$. Consequently, we have:

$$\mathrm{ch}(i_*(i^*(y))) = \mathrm{ch}(y) \cdot (1 - \exp(c^1(J))).$$

Furthermore, the functoriality of ch, and the projection formula 3.7 imply:

$$i_{\mathrm{gr}}\big(\mathrm{ch}(i^*(y)) \cdot \mathrm{Todd}(\check{T}_i)\big) = \mathrm{ch}(y) \cdot i_{\mathrm{gr}}(\mathrm{Todd}(\check{T}_i)).$$

Now $T_i = -\operatorname{cl}(\underline{J}/\underline{J}^2) = -i^*(J)$. Thus:

$$i_{\mathrm{gr}}(\mathrm{Todd}(\check{T}_i)) = i_{\mathrm{gr}}\big(i^{\mathrm{gr}}(\mathrm{Todd}(-\check{J}))\big) = i_{\mathrm{gr}}(1) \cdot \mathrm{Todd}(-\check{J}).$$

Since $i_*(1) = 1 - J$, $i_{\mathrm{gr}}(1)$ is the image in $\mathrm{Gr}^1\, K^\bullet(X)$ of $1 - J$, and is therefore $-c^1(J)$[494]. Furthermore, we have:

$$\mathrm{Todd}(-\check{J}) = (\mathrm{Todd}(\check{J}))^{-1} = \frac{1 - \exp(c^1(J))}{-c^1(J)}.$$

Thus finally we obtain:

$$i_{\mathrm{gr}}\big(\mathrm{ch}(i^*(y)) \cdot \mathrm{Todd}(\check{T}_i)\big) = \mathrm{ch}(y) \cdot (1 - \exp(c^1(J))),$$

which proves the required relation.

5. Riemann–Roch theorem: case of the structural morphism of a projective bundle

5.1. By 3.9, to prove the Riemann–Roch theorem, it remains to prove it for the structural morphism $f : \mathbf{P}^r_Y \longrightarrow Y$[495], where Y is a quasi-compact scheme having an ample sheaf. Put $P = \mathbf{P}^r_Y$[496], $\xi = \mathrm{cl}(\underline{O}_P(1))$, and $\underline{E} = (\underline{O}_Y)^{r+1}$.


[print 493]**Proposition 5.2.** We have on P the exact sequence:

$$0 \longrightarrow \underline{\Omega}^1_{P/Y} \longrightarrow f^*(\underline{E})(-1) \longrightarrow \underline{O}_P \longrightarrow 0,$$

i.e. we have an isomorphism

$$\underline{\Omega}^1_{P/Y} \simeq \underline{F}(-1), \quad \text{where} \quad \underline{F} = \mathrm{Ker}\big(f^*(\underline{E}) \longrightarrow \underline{O}_P(1)\big).$$

Consider the vector bundle over Y defined by $\underline{E}$, and denote by Z the complement of the zero section of this bundle. We have the commutative diagram:




The morphism h identifies Z with the affine P-scheme defined by the sheaf of algebras $\underline{B} = \bigoplus_{n\in\mathbb{Z}} \underline{O}_P(n)$; $\underline{B}$ is a graded $\underline{O}_P$-Algebra.

Since f, g, h are smooth morphisms, one may write the exact sequence of sheaves of relative differentials; since Z is affine over P, one may write this sequence in terms of $\underline{B}$-Modules; one thus obtains:

$$0 \longrightarrow \underline{\Omega}^1_{P/Y} \otimes_{\underline{O}_P} \underline{B} \longrightarrow \underline{\Omega}^1_{\underline{B}/Y} \longrightarrow \underline{\Omega}^1_{\underline{B}/\underline{O}_P} \longrightarrow 0. \tag{5.2.1}$$

Now this exact sequence is a graded exact sequence of graded $\underline{B}$-Modules. Indeed, it follows from the commutative diagram: [497]



where the arrows are algebra homomorphisms compatible with the grading of $\mathbf{S}(\underline{E})[T, T^{-1}]$[498] (grading defined by degree in the indeterminate T), the two other algebras being supposed concentrated [print 494]in degree 0.

The module of differentials $\underline{\Omega}^1_{\underline{B}/\underline{O}_P}$ is isomorphic to $\underline{B}(-1) \otimes_{\underline{O}_P} \underline{O}_P(1)$, since one can define an $\mathbf{S}(\underline{E})$-derivation[499] d: [500]

$$\mathbf{S}(\underline{E})[T, T^{-1}] \longrightarrow \mathbf{S}(\underline{E})[T, T^{-1}] \otimes_{\mathbf{S}(\underline{E})} \mathbf{S}(\underline{E}) \cdot dT$$

so as to satisfy the universal property defining $\underline{\Omega}^1_{\underline{B}/\underline{O}_P}$, by setting $d(aT^n) = naT^{n-1} \otimes dT$; moreover, d is compatible with the gradings if the factor $\mathbf{S}(\underline{E}) \cdot dT$[501] is assigned degree 1.

Let V be the vector bundle over Y defined by $\underline{E}$. Then $\underline{\Omega}^1_{Z/Y}$ is the restriction to Z of $\underline{\Omega}^1_{V/Y}$. Now V is the affine Y-scheme defined by $\mathbf{S}(\underline{E})$[502], so $\underline{\Omega}^1_{V/Y}$ is defined by $\mathbf{S}(\underline{E}) \otimes_{\underline{O}_Y} \underline{E}$[503] (since $\underline{E}$ is

locally free of finite type), hence $\underline{\Omega}^1_{V/Y}$ is equal to $g^*(\underline{E})$, and consequently $\underline{\Omega}^1_{\underline{B}/Y} = f^*(\underline{E}) \otimes_{\underline{O}_P} \underline{B}$. To see its grading, one looks locally, and sees at once that $f^*(\underline{E})$ must be regarded as being of degree 1.

Taking the degree 0 terms in the exact sequence (5.2.1) gives the announced exact sequence.

Remark 5.2.1. Using the general results on the differential study of Hilbert functors ([2], 5), one directly obtains an isomorphism $\underline{\Omega}^1_{P/Y} \xrightarrow{\sim} \underline{F}(-1)$ from the given functorial characterization of $g^*(\underline{\Omega}^1_{P/Y})$ for every Y-morphism $g : Y' \to P$.

5.3. By 5.2, we have:

$$T_f = \mathrm{cl}(\underline{\Omega}^1_{P/Y}) = f^*(E) \cdot \xi^{-1} - 1^{[504]}.$$

Consequently:

$$\widetilde{c}(T_f) = \widetilde{c}(E) \cdot \widetilde{c}(\xi^{-1}) - \widetilde{c}(1).$$

[print 495]Since $\underline{E} = (\underline{O}_Y)^{r+1}$, we have $E = r + 1$, and consequently:

$$\widetilde{c}(E) = (r + 1, 1).$$

Put:

$$x = \mathrm{cl}_{\mathrm{Gr}^1 K^\bullet(P)}(\xi - 1) = \mathrm{cl}_{\mathrm{Gr}^1 K^\bullet(P)}(1 - \xi^{-1}) = c^1(\xi).$$

Since $\xi^{-1} = \check{\xi}$, we have:

$$\widetilde{c}(\xi^{-1}) = (1, 1 - x).$$

By V (6.2.1), this gives:

$$\widetilde{c}(T_f) = (r, (1 - x)^{r+1}).$$

Taking 3.5 into account, we finally obtain:

$$\widetilde{c}(\check{T}_f) = (r, (1 + x)^{r+1}).$$

By the definition of Todd, we therefore have:

$$\mathrm{Todd}(\check{T}_f) = \left(\frac{x}{1 - \exp(-x)}\right)^{r+1}. \tag{5.3.1}$$

5.4. Using the basis of $K^\bullet(P)$ over $K^\bullet(Y)$ formed by the elements ξ^k, with $-r \leq k \leq 0$, one sees by additivity that it suffices to prove the theorem for elements of the form $f^*(a) \cdot \xi^k$. We then have:

$$\mathrm{ch}\big(f_*(f^*(a)\xi^k)\big) = \mathrm{ch}(a f_*(\xi^k)) = \mathrm{ch}(a) \cdot \mathrm{ch}(f_*(\xi^k)).$$

On the other hand:

$$f_{\mathrm{gr}}\big(\mathrm{ch}(f^*(a)\xi^k)\,\mathrm{Todd}(\check{T}_f)\big) = f_{\mathrm{gr}}\big(f^{\mathrm{gr}}(\mathrm{ch}(a))\,\mathrm{ch}(\xi^k)\,\mathrm{Todd}(\check{T}_f)\big) = \mathrm{ch}(a) \cdot f_{\mathrm{gr}}\big(\mathrm{ch}(\xi^k)\,\mathrm{Todd}(\check{T}_f)\big).$$

One thus sees that it suffices to prove the theorem for the ξ^k, with $-r \leq k \leq 0$. Now by (VI (5.2.3)), we have

$$f_*(\xi^k) = 0 \quad \text{for } -r \leq k < 0\ ; \qquad f_*(1) = 1.$$

Consequently:

$$\mathrm{ch}(f_*(\xi^k)) = 0 \quad \text{if } -r \leq k < 0\ ; \qquad \mathrm{ch}(f_*(1)) = 1. \tag{5.4.1}$$

[print 496]Let us now calculate the second side of the Riemann–Roch formula. For every k, we have:

$$\widetilde{c}(\xi^k) = (\widetilde{c}(\xi))^k = (1, 1 + x)^k = (1, 1 + kx)$$

by (V (6.2.1)). Consequently:

$$\mathrm{ch}(\xi^k) = \exp(kx). \tag{5.4.2}$$

The second side of the Riemann–Roch formula is therefore:

$$f_{\mathrm{gr}}\left(\exp(kx)\cdot\frac{x^{r+1}}{(1-\exp(-x))^{r+1}}\right) \tag{5.4.3}$$

Since x is the class in $\mathrm{Gr}^1 K^\bullet(P)$ of $1-\xi^{-1}$, $f_{\mathrm{gr}}(x^n)$ is, for $0 \le n \le r$, the class in $\mathrm{Gr}^{n-r} K^\bullet(Y)$ of $f_*((1-\xi^{-1})^n) = 1$. We thus have:

$$f_{\mathrm{gr}}(x^n) = 0 \quad \text{if } 0 \le n < r\ ; \qquad f_{\mathrm{gr}}(x^r) = 1. \tag{5.4.4}$$

Finally recall that the elements $1, x, \ldots, x^{r+1}$ are linearly dependent over $\mathrm{Gr}\, K^\bullet(Y)$ (VI 5.6), and satisfy the equation:

$$x^{r+1} = 0, \tag{5.4.5}$$

and that $1, \ldots, x^r$ are linearly independent over $\mathrm{Gr}\, K^\bullet(Y)$.[505]

One can then write the series $\varphi_k = \exp(kx)\cdot\frac{x^{r+1}}{(1-\exp(-x))^{r+1}}$ uniquely as a linear combination, with coefficients in $\mathrm{Gr}\,\widehat{K^\bullet(Y)}_{\mathbb{Q}}$, of the elements $1, \ldots, x^r$. The image of φ_k under f_{gr} is then, by (5.4.4) and 3.7, its coefficient of x^r. Taking (5.4.1) into account, the Riemann–Roch theorem for f is therefore equivalent to:

$$\left\{\begin{array}{l} \text{for } -r \le k < 0, \text{ the coefficient of } x^r \text{ in } \varphi_k \text{ is } 0\ ; \\ \text{for } k = 0, \text{ the coefficient of } x^r \text{ in } \varphi_k \text{ is } 1. \end{array}\right. \tag{RR}$$

This coefficient is the residue of the differential form

$$\frac{\exp(kx)\cdot dx}{(1-\exp(-x))^{r+1}}.$$

[print 497]Putting $y = 1-\exp(-x)$, we are reduced to calculating the residue of: [506]

$$\frac{(1-y)^{-k-1}\cdot dy}{y^{r+1}},$$

that is, the coefficient of y^r in $(1-y)^{-k-1}$. For $-r \le k < 0$, it is zero, and for $k = 0$, it is 1.

This completes the proof of the Riemann–Roch theorem.

BIBLIOGRAPHY

[1] A. Borel and J.P. Serre, Le théorème de Riemann–Roch, Bull. Soc. Math. de France, t. 86, 1958.

[2] A. Grothendieck, Techniques de construction et théorèmes d'existence en Géométrie Algébrique, Sém. Bourbaki, 1960-61.

[print 498] # EXPOSÉ IX

SOME COMPUTATIONS OF GROUPS K

by P. BERTHELOT

This exposé corresponds to no oral exposé, and will not be used in the rest of the seminar. It contains a number of computations of groups $K_\bullet$ for various fibrations: vector bundles, projective bundles, torsors under a linear group, etc... Throughout this exposé, the base scheme will be supposed noetherian, and the notation $K_\bullet(X)$ will denote the Grothendieck group of coherent sheaves on a scheme X. Recall that if X is noetherian, the homomorphism from the K of the category of coherent $\underline{O}_X$-Modules to the K of the category of pseudo-coherent complexes on X is an isomorphism (IV 2.4).

The results presented here are analogous to those obtained for the Chow ring, which can be found in Grothendieck's exposés in the Chevalley Seminar 1958. The question also arises of determining to what extent statements of the same type can be obtained for the graded group associated with $K_\bullet$ for the filtration defined by the dimension of the support (cf. X). It does not seem that the answer is known, even for Proposition 1.1 (homotopy exact sequence), which is one of the properties most used in the computations that follow.

1. Vector bundles

Proposition 1.1. (Homotopy exact sequence). Let X be a noetherian scheme, X' a closed subscheme of X, U the open complement of X' in X, i the immersion of X' into X, and j the immersion of U into X. Then the sequence: [print 499]

$$K_\bullet(X') \xrightarrow{i_*} K_\bullet(X) \xrightarrow{j^*} K_\bullet(U) \longrightarrow 0$$

is exact.

If $\underline{F}'$ is a coherent $\underline{O}_{X'}$-Module, $i_*(\underline{F}')$ is an $\underline{O}_X$-Module whose support is contained in X'; consequently its restriction to U is zero, and $j^* \circ i_*(\mathrm{cl}(\underline{F}')) = 0$, whence, since the elements of the form $\mathrm{cl}(\underline{F}')$ generate $K_\bullet(X')$, $j^* \circ i_* = 0$. This gives a homomorphism

$$\rho : K_\bullet(X)/\operatorname{Im} K_\bullet(X') \longrightarrow K_\bullet(U).$$

We shall define an inverse homomorphism to ρ. Let $\underline{G}$ be a coherent $\underline{O}_U$-Module. Since X is noetherian, there exists a coherent $\underline{O}_X$-Module $\underline{F}$ whose restriction to U is isomorphic to $\underline{G}$. If $\underline{F}'$ is a second coherent $\underline{O}_X$-Module whose restriction to U is isomorphic to $\underline{G}$, then we have in $K_\bullet(X)$:

$$\mathrm{cl}(\underline{F}) - \mathrm{cl}(\underline{F}') \in \operatorname{Im} K_\bullet(X').$$

To see this, regard $\underline{G}$ as a subsheaf of $j^*(\underline{F} \oplus \underline{F}')$ by means of the diagonal map; $\underline{G}$ then extends to a subsheaf $\underline{F}''$ of $\underline{F} \oplus \underline{F}'$, and we are reduced to showing that

$$\mathrm{cl}(\underline{F}) - \mathrm{cl}(\underline{F}'') \in \operatorname{Im} K_\bullet(X'),$$

and the analogous assertion for $\underline{F}'$. Let then $\underline{H}$ and $\underline{K}$ be the kernel and cokernel of the composite homomorphism:

$$\underline{F}'' \longrightarrow \underline{F} \oplus \underline{F}' \longrightarrow \underline{F};$$

this homomorphism reduces to the identity of $\underline{G}$ over U, and therefore $\underline{H}$ and $\underline{K}$ have support contained in X'. If $\underline{I}$ is the ideal of X' in X, it follows that there exist integers m and n such

that $\underline{I}^m \cdot \underline{H} = 0$, $\underline{I}^n \cdot \underline{K} = 0$ (EGA I 9.3.4). One may therefore express $\underline{H}$ and $\underline{K}$ as successive extensions of Modules [print 500]annihilated by $\underline{I}$, hence of $\underline{O}_{X'}$-Modules, and we have:

$$\mathrm{cl}(\underline{H}) \in \operatorname{Im} K_\bullet(X'), \qquad \mathrm{cl}(\underline{K}) \in \operatorname{Im} K_\bullet(X'),$$

giving the result. We thus define a function on the category of coherent $\underline{O}_U$-Modules with values in $K_\bullet(X)/\operatorname{Im} K_\bullet(X')$ by assigning to every coherent $\underline{O}_U$-Module the class of one of its extensions to X.

This function is additive. Indeed, let:

$$0 \longrightarrow \underline{G}' \longrightarrow \underline{G} \longrightarrow \underline{G}'' \longrightarrow 0$$

be an exact sequence of coherent $\underline{O}_U$-Modules. Let $\underline{F}$ be an extension of $\underline{G}$ to X; then there exists a sub-Module $\underline{F}'$ of $\underline{F}$ extending $\underline{G}'$, and $\underline{F}/\underline{F}'$ extends $\underline{G}''$. It therefore factors through $K_\bullet(U)$, defining the required homomorphism $K_\bullet(U) \to K_\bullet(X)/\operatorname{Im} K_\bullet(X')$. One checks immediately that it is inverse to ρ.

Proposition 1.2. Let X, S be two noetherian schemes, and f a flat morphism from X to S. Suppose given a family of elements $(\xi_i)_{i\in I}$ of $K^\bullet(X)$, and consider the homomorphism:

$$\varphi : (K_\bullet(S))^{(I)} \longrightarrow K_\bullet(X)$$

defined by:

$$\varphi((x_i)_{i\in I}) = \sum_{i\in I} \xi_i \cdot f^*(x_i),$$

where f^* is the "inverse-image" homomorphism from $K_\bullet(S)$ to $K_\bullet(X)$, defined by IV 2.12, and where $K_\bullet(X)$ is regarded as a $K^\bullet(X)$-module by IV 2.10. Suppose that for every point $s \in S$, the images of the ξ_i in $K_\bullet(X_s)$ under the homomorphism:

$$K^\bullet(X) \longrightarrow K^\bullet(X_s) \longrightarrow K_\bullet(X_s)$$

generate $K_\bullet(X_s)$ as an abelian group. Then φ is surjective.

[print 501]Let S' be a closed subscheme of S, and U the open complement of S' in S. Denote by X_U the restriction of X over U, and by $X_{S'}$ the restriction of X over S'. Finally, still denote by ξ_i the images of the ξ_i in $K^\bullet(X_{S'})$ and $K^\bullet(X_U)$, and denote by:

$$\varphi_{S'} : K_\bullet(S')^{(I)} \to K_\bullet(X_{S'}), \qquad \varphi_U : K_\bullet(U)^{(I)} \to K_\bullet(X_U)$$

the homomorphisms defined in the same way as φ.

Consider the diagram

$$\begin{array}{ccccccc} K_\bullet(S')^{(I)} & \xrightarrow{i_*} & K_\bullet(S)^{(I)} & \xrightarrow{j^*} & K_\bullet(U)^{(I)} & \to & 0 \\ \downarrow{\scriptstyle \varphi_{S'}} & & \downarrow{\scriptstyle \varphi} & & \downarrow{\scriptstyle \varphi_U} & & \\ K_\bullet(X_{S'}) & \xrightarrow{u_*} & K_\bullet(X) & \xrightarrow{v^*} & K_\bullet(X_U) & \to & 0 \end{array} \tag{1.2.1}$$

where i and j denote the immersions of S' and U into S, and u and v the immersions of $X_{S'}$ and X_U into X. Let us show that if $\varphi_{S'}$ and φ_U are surjective, the same then holds for φ.

The two horizontal rows of the diagram are exact by 1.1. One then sees that to show that φ is surjective, it suffices to show that the two squares commute.

Let $(x'_i)_{i\in I} \in K_\bullet(S')^{(I)}$; we have:

$$\varphi \circ i_*((x'_i)) = \varphi((i_*(x'_i))) = \sum_i \xi_i \cdot f^*(i_*(x'_i));$$

$$u_* \circ \varphi_{S'}((x'_i)) = u_* \left(\sum_i \xi_i \cdot f^*_{S'}(x'_i) \right) = \sum_i \xi_i \cdot u_*(f^*_{S'}(x'_i)),$$

the last equality following from the projection formula IV 2.11.1.2. The equality $\varphi \circ i_* = u_* \circ \varphi_{S'}$ therefore follows from the equality $f^* \circ i_* = u_* \circ f^*_{S'}$, which holds by IV 3.1.1 since i is proper and f flat.

[print 502]Let $(x_i)_{i \in I} \in K_\bullet(S)^{(I)}$; we have:

$$\varphi_U \circ j^*((x_i)) = \sum_i \xi_i \cdot f^*_U \circ j^*(x_i);$$

$$v^* \circ \varphi((x_i)) = v^* \left(\sum_i \xi_i \cdot f^*(x_i) \right) = \sum_i \xi_i \cdot v^* \circ f^*(x_i) = \sum_i \xi_i \cdot f^*_U \circ j^*(x_i)$$

the middle equality following from IV 2.12.2.

This result allows us to prove 1.2 by noetherian induction on the set of closed subschemes of S. If we suppose that for every closed subscheme S' of S distinct from S, $\varphi_{S'}$ is surjective, then the preceding argument shows that we may reduce to proving 1.2 for an irreducible affine base scheme, after replacing S by an irreducible affine open subset. Moreover, the diagram

$$\begin{array}{ccc} K_\bullet(S_{\mathrm{red}})^{(I)} & \longrightarrow & K_\bullet(S)^{(I)} \\ {\scriptstyle \varphi_{S_{\mathrm{red}}}}\downarrow & & \downarrow{\scriptstyle \varphi} \\ K_\bullet(X_{S_{\mathrm{red}}}) & \longrightarrow & K_\bullet(X), \end{array}$$

where $K_\bullet(X_{S_{\mathrm{red}}}) \to K_\bullet(X)$ is surjective by 1.1, shows that we may also suppose S reduced.

We therefore work over an integral affine base scheme. Put $S = \mathrm{Spec}(A)$, and denote by K the field of fractions of A; if s is the generic point of S, then $\underline{O}_{S,s} = K$.

Let $y \in K_\bullet(X)$. By hypothesis, $K_\bullet(X_s)$ is generated by the images of the ξ_i; if y_s is the image of y in $K_\bullet(X_s)$, we can find integers n_i such that:

$$y_s = \sum_i \xi_i \cdot n_i.$$

[print 503]Furthermore, we have $K = \varinjlim_f A_f$, where f runs through the set of nonzero elements of[507] A. Consequently, $\mathrm{Spec}(K) = \varprojlim_f \mathrm{Spec}(A_f)$, and:

$$X_s = X \times_S \mathrm{Spec}(K) = \varprojlim_f X \times_S \mathrm{Spec}(A_f)$$

by EGA IV 8.2.3 and 8.2.5. Since the schemes considered are noetherian, and the transition morphisms affine and flat, we have by IV 3.2.3:

$$K_\bullet(X_s) = \varinjlim_f K_\bullet(X \times_S \mathrm{Spec}(A_f)).$$

Since the "inverse-image" homomorphisms on the $K_\bullet$ are linear with respect to the $K^\bullet$ (IV (2.12.2)), there exists a nonempty open subset $U = \mathrm{Spec}(A_f)$ of S such that in $K_\bullet(X_U)$ we have:

$$y_U = \sum_i \xi_i \cdot n_i.$$

Let S' be a closed subscheme of S with support $S - U$. Then $\varphi_{S'}$ is surjective by the induction hypothesis, and consequently diagram (1.2.1) shows that:

$$y - \sum_i \xi_i \cdot n_i \in \mathrm{Im}(\varphi),$$

so, since $n_i = f^*(n_i)$, φ is surjective.

Corollary 1.3. Let $f : X \to S$ be a flat morphism of noetherian schemes. Suppose that for every point s of S, the homomorphism:

$$K^\bullet(X) \longrightarrow K_\bullet(X_s)$$

is surjective. Then the homomorphism obtained from f^*:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X) \longrightarrow K_\bullet(X)$$

is surjective.

[print 504]Take for the family $(\xi_i)_{i\in I}$ the family of all elements of $K^\bullet(X)$. The hypotheses of 1.2 are then satisfied, and consequently the homomorphism:

$$\varphi : \big(K_\bullet(S)\big)^{(K^\bullet(X))} \longrightarrow K_\bullet(X)$$

is surjective. By the definition of φ, this is equivalent to saying that the homomorphism:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X) \longrightarrow K_\bullet(X)$$

is surjective.

Corollary 1.4. Let $f : X \to S$ be a flat morphism of noetherian schemes. Suppose that for every point s of S, the canonical homomorphism $\mathbb{Z} \to K_\bullet(X_s)$ is an isomorphism. Then the homomorphism:

$$f^* : K_\bullet(S) \longrightarrow K_\bullet(X)$$

is surjective. If in addition f has a section g of finite Tor-dimension, then f^* is an isomorphism.

Take for the family $(\xi_i)_{i\in I}$ the family consisting of the single element $1 \in K^\bullet(X)$; we then have $\varphi = f^*$, and, the conditions of 1.2 being satisfied, f^* is surjective.

If f has a section g of finite Tor-dimension, one can define a homomorphism $g^* : K_\bullet(X) \to K_\bullet(S)$ (IV 2.12). Since we have $g^* \circ f^* = (f \circ g)^* = \mathrm{Id}$ by IV 2.12.1.1, f^* is injective, hence an isomorphism.

Remark 1.5. For g to have finite Tor-dimension, it is necessary and sufficient that the fibers of X be regular at the points of $g(S)$, i.e. that f be regular at the points of $g(S)$ (or smooth at these points if we suppose f [print 505]of finite presentation).

Since these properties are local on X and S, one may suppose them affine; let $S = \mathrm{Spec}(A)$, $X = \mathrm{Spec}(B)$. We then have:

$$\mathrm{tor.\,dim}_B(A) = \mathrm{dim.\,proj}_B(A) = \sup_{s\in S} \mathrm{dim.\,proj}_{B\otimes_A k(s)}(k(s))$$

by EGA IV 12.3.2, where the hypothesis of finite presentation on B can be replaced by noetherian hypotheses. Now $k(s)$ is isomorphic to the residue field of $\underline{O}_{X_s,g(s)}$, and by EGA 0_{IV} 17.2.5 we have:

$$\mathrm{dim.\,proj}_{B\otimes_A k(s)}(k(s)) = \mathrm{dim.\,proj}_{\underline{O}_{X_s,g(s)}}(k(s)).$$

It then follows from EGA 0_{IV} 17.2.11 that the latter is finite if and only if $\underline{O}_{X_s,g(s)}$ has finite cohomological dimension, that is, is regular.

Proposition 1.6. Let S be a noetherian scheme, E a locally free $\underline{O}_S$-Module of finite type, X the vector bundle associated to E, and f the structural morphism of X. Then $f^* : K_\bullet(S) \longrightarrow K_\bullet(X)$ is an isomorphism.

Since f is smooth, and has a section, namely the zero section, to prove 1.6 it suffices to show that for every $s \in S$, $K_\bullet(X_s) \simeq \mathbb{Z}$ (1.4). We are thus reduced to the case where the base scheme is a field.

In this case, $\underline{E}$ is a direct sum of submodules of rank 1, which allows us to decompose f into a sequence of fibrations of rank 1, and thus reduce to the case where X has rank 1, namely $X = \mathrm{Spec}(k[T])$. The proposition then follows from the next lemma:

[print 506]**Lemma 1.7.** Let A be a principal ideal domain, $X = \mathrm{Spec}(A)$. Then the canonical homomorphism:

$$\mathbb{Z} \longrightarrow K_\bullet(X)$$

is an isomorphism.

Since X is regular, the homomorphism $K^\bullet(X) \to K_\bullet(X)$ is an isomorphism. Thus $K_\bullet(X)$ is generated by the classes of locally free $\underline{O}_X$-Modules of finite type, i.e. by the classes of projective A-modules of finite type. Since A is a principal ideal domain, every projective module of finite type is free, giving surjectivity. Injectivity follows from the fact that $K^\bullet(X)$ is $\mathbb{Z}$-augmented.

Proposition 1.8. Let S be a noetherian scheme, $\underline{E}$ a locally free $\underline{O}_S$-Module of rank r, X the vector bundle associated to $\underline{E}$, f the structural morphism of X, and g the zero section of X. Then the composite homomorphism:

$$K_\bullet(S) \xrightarrow{g_*} K_\bullet(X) \xrightarrow{g^*} K_\bullet(S)$$

(where g^* is defined by IV 2.12.1 and 1.5) is multiplication by the element $\lambda_{-1}(E)$ of $K^\bullet(S)$, denoting by E the class of $\underline{E}$ in $K^\bullet(S)$, and putting

$$\lambda_{-1}(E) = \sum_{i=0}^{r} (-1)^i \lambda^i(E).$$

This statement is the analogue, for $K_\bullet$, of VII 2.7.

Let $\underline{F}$ be a coherent $\underline{O}_S$-Module, of class F in $K_\bullet(S)$. Then $g_*(F)$ is the class in $K_\bullet(X)$ of the complex $\mathbb{R}g_*(\underline{F}) = g_*(\underline{F})$, hence of $\underline{F}$ regarded as an $\underline{O}_X$-Module by means of g. To compute $g^*(g_*(F))$[508], one must therefore take the class of $\mathbb{L}g^*(\underline{F}) = \underline{F} \otimes^{\mathbb{L}}_{\underline{O}_X} \underline{O}_S$.

[print 507]For this, consider the complex:

$$0 \longrightarrow \wedge^r(\underline{E}) \otimes_{\underline{O}_S} \mathbb{S}(\underline{E}) \longrightarrow \wedge^{r-1}(\underline{E}) \otimes_{\underline{O}_S} \mathbb{S}(\underline{E}) \longrightarrow \ldots \longrightarrow \underline{E} \otimes_{\underline{O}_S} \mathbb{S}(\underline{E}) \longrightarrow \mathbb{S}(\underline{E}) \longrightarrow 0$$

where $\mathbb{S}(\underline{E})$ is the symmetric algebra of $\underline{E}$. This is a resolution of $\underline{O}_S$ over $\mathbb{S}(E)$ (VI 1.11). The associated complex on X, denoted $\underline{L}_\bullet$, has k-th term $\wedge^k(f^*(E))$ and is therefore a resolution of $\underline{O}_S$ by locally free $\underline{O}_X$-Modules of finite type; thus we have $\underline{L}_\bullet \otimes_{\underline{O}_X} \underline{F} = \underline{O}_S \otimes^{\mathbb{L}}_{\underline{O}_X} \underline{F}$. Consequently:

$$\begin{aligned} g^*(g_*(F)) &= \mathrm{cl}(\underline{L}_\bullet \otimes_{\underline{O}_X} \underline{F}) = \sum_{i=0}^{r} (-1)^i \,\mathrm{cl}(\underline{L}_i \otimes_{\underline{O}_X} \underline{F}) \\ &= \sum_{i=0}^{r} (-1)^i \,\mathrm{cl}(\wedge^i(\underline{E}) \otimes_{\underline{O}_S} \underline{F}) = F \cdot \lambda_{-1}(E). \end{aligned}$$

The proposition follows by additivity.

2. Principal bundles under split tori

2.1. Let S be a noetherian scheme, and T a torus over S. Suppose T split; we thus have an isomorphism $T \simeq D_S(M)$, where M is a free abelian group of rank r, or equivalently $T \simeq \underline{G}_{m\,S}^r$.

Let P be a torsor under T. Then P is affine over S, defined by a graded $\underline{O}_S$-Algebra $\underline{A}$ of type M; putting $\underline{A} = \bigoplus_{m\in M} \underline{A}_m$, we have that $\underline{A}_m$ is an invertible $\underline{O}_S$-Module for every m, and, for $m, m' \in M$, the homomorphism $\underline{A}_m \otimes_{\underline{O}_S} \underline{A}_{m'} \longrightarrow \underline{A}_{m+m'}$ defined by multiplication in $\underline{A}$ is an isomorphism (SGA 3 VIII 4.1). We thus define a multiplicative homomorphism $\eta : M \longrightarrow K^\bullet(S)$ by associating to $m \in M$ the class of $\underline{A}_m$ in $K^\bullet(S)$. [print 508]

Proposition 2.2. With the notation and hypotheses of 2.1, let $f : P \to S$ be the structural morphism of P. Then the homomorphism:

$$f^* : K_\bullet(S) \longrightarrow K_\bullet(P)$$

is surjective, and its kernel is the subgroup of $K_\bullet(S)$ generated by the subgroups $(\eta(m)-1)K_\bullet(S)$ for $m \in M$, or, equivalently, for m running through a system of generators of M.

First suppose $r = 1$, i.e. $T = G_{m\,S}$. Then $M = \mathbb{Z}$, and for every $m \in M$ one has:

$$\underline{A}_m \simeq \underline{A}_1^{\otimes m}.$$

Consequently, putting $\underline{A}_1 = \underline{L}$, one has $\underline{A} = \bigoplus_{n\in\mathbb{Z}} \underline{L}^{\otimes n}$, which shows that P identifies with the complement of the zero section in the bundle $X = \mathbb{V}(\underline{L})$. Considering S as a closed subscheme of X by the zero section, one has, by 1.1, the exact sequence:

$$K_\bullet(S) \xrightarrow{g_*} K_\bullet(X) \longrightarrow K_\bullet(P) \longrightarrow 0$$

where $g : S \to X$ is the zero section of X. Moreover, f^* factors as $K_\bullet(S) \xrightarrow{f'^*} K_\bullet(X) \longrightarrow K_\bullet(P)$, where f' is the structural morphism of X. By 1.6, f'^* is an isomorphism, inverse to g^*. Thus one has the exact sequence:

$$K_\bullet(S) \xrightarrow{g^*\circ g_*} K_\bullet(S) \xrightarrow{f^*} K_\bullet(P) \longrightarrow 0,$$

from which one obtains, by 1.8:

$$K_\bullet(P) \simeq K_\bullet(S)/\lambda_{-1}(L)K_\bullet(S) = K_\bullet(S)/(1-L)K_\bullet(S)$$

where L is the class in $K^\bullet(S)$ of $\underline{L}$. Since for every $n \in \mathbb{Z}$, $L^n - 1$ is a multiple of $1 - L$, this proves 2.2 when P has rank 1.

[print 509]

Continue the proof by induction on the rank r of P. Suppose the result proved for torsors under split tori of rank $< r$. Write $M = M_1 \oplus M_2$, with $M_1 \simeq \mathbb{Z}^{r-1}$ and $M_2 \simeq \mathbb{Z}$; put $\underline{A}' = \bigoplus_{m\in M_1} \underline{A}_m$ and $\underline{A}'' = \bigoplus_{m\in M_2} \underline{A}_m$, so that $\underline{A} = \underline{A}' \otimes_{\underline{O}_S} \underline{A}''$. Let $P_1 = \mathrm{Spec}(\underline{A}')$; then P_1 is a torsor under $D_S(M_1)$, and P is a torsor over P_1 with group $\underline{G}_{m\,P_1}$ and algebra the inverse image of $\underline{A}''$ on P_1. By the preceding argument, $K_\bullet(P)$ is the quotient of $K_\bullet(P_1)$ by the subgroup generated by the $(\eta(m_2) - 1)K_\bullet(P_1)$ for $m_2 \in M_2$. By the induction hypothesis, $K_\bullet(P_1)$ is the quotient of $K_\bullet(S)$ by the subgroup generated by the $(\eta(m_1) - 1)K_\bullet(S)$ for $m_1 \in M_1$. Taking account of the relation

$$\eta(m_1 + m_2) - 1 = \eta(m_1)\eta(m_2) - 1 = \eta(m_1)(\eta(m_2) - 1) + \eta(m_1) - 1,$$

$K_\bullet(P)$ is therefore the quotient of $K_\bullet(S)$ by the subgroup generated by the $(\eta(m) - 1)K_\bullet(S)$ for $m \in M$, or equivalently for m running through a system of generators of M.

Corollary 2.3. With the notation and hypotheses of 2.1, suppose that S is regular; then one has a canonical isomorphism:

$$K^\bullet(P) \simeq K^\bullet(S)/\big((\eta(m) - 1)_{m\in M}\big)$$

where $((\eta(m)-1)_{m\in M})$ is the ideal of $K^\bullet(S)$ generated by the $\eta(m)-1$ for m running through a system of generators of M.

If S is regular, the canonical homomorphism $K^\bullet(S) \to K_\bullet(S)$ is an isomorphism (IV 2.5). Likewise, P is then regular, so $K^\bullet(P) \to K_\bullet(P)$ is an isomorphism. The corollary therefore follows immediately from 2.2.

[print 510]

3. Projective bundles and flag bundles

Proposition 3.1. Let S be a noetherian scheme, $\underline{E}$ a locally free $\underline{O}_S$-Module of rank n, $\pi = (p_1, \ldots, p_k)$ a sequence of integers such that $\sum_i p_i = n$, and $X = \mathbb{D}_\pi(\underline{E})$ the scheme of flags of type π in $\underline{E}$. Then the canonical homomorphism:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X) \longrightarrow K_\bullet(X)$$

is an isomorphism.

Resume the notation of VI 4.5. Thus $\underline{L}_i$ denotes the canonical quotients of the inverse image of $\underline{E}$ on X, $\underline{F}_i$ the kernel of $\underline{L}_i \to \underline{L}_{i-1}$ $(i = 1, \ldots, k)$, and $\lambda_i^{(j)}$ the class in $K^\bullet(X)$ of $\wedge^i(\underline{F}_j)$. Then the $\lambda_i^{(j)}$ generate $K^\bullet(X)$ as a $K^\bullet(S)$-algebra (VI 4.6). Likewise, the images of the $\lambda_i^{(j)}$ in $K^\bullet(X_s)$ generate $K^\bullet(X_s)$ as an algebra over $K^\bullet(k(s)) = \mathbb{Z}$ for every point s of S. Since X_s is a flag scheme over a field, it is a regular scheme, so the homomorphism $K^\bullet(X_s) \to K_\bullet(X_s)$ is an isomorphism, and thus the homomorphism $K^\bullet(X) \to K_\bullet(X_s)$ is surjective. [509]By 1.3, this implies that the homomorphism:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X) \longrightarrow K_\bullet(X)$$

is surjective.

To prove injectivity, one may first reduce to the case of the complete flag bundle $\mathbb{D}(E)$. Indeed, suppose that

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(\mathbb{D}(\underline{E})) \longrightarrow K_\bullet(\mathbb{D}(\underline{E}))$$

is injective. Since $K_\bullet(S) \to K_\bullet(\mathbb{D}(\underline{E}))$ factors through $K_\bullet(X)$, this [print 511]homomorphism factors as:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X) \otimes_{K^\bullet(X)} K^\bullet(\mathbb{D}) \longrightarrow K_\bullet(X) \otimes_{K^\bullet(X)} K^\bullet(\mathbb{D}) \longrightarrow K_\bullet(\mathbb{D})$$

and the first homomorphism is injective. Since, by VI 4.3, 4.6 and 4.7, $K^\bullet(\mathbb{D})$ is free over $K^\bullet(X)$, this implies the injectivity of:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X) \to K_\bullet(X).$$

Thus let $X = \mathbb{D}(\underline{E})$. Proceed by induction on the rank of $\underline{E}$. The case $n = 1$ being trivial, suppose the property proved for $n-1$. Introduce $P = \mathbb{P}(\underline{E})$ and the kernel $\underline{F}$ of the canonical homomorphism $p^*(\underline{E}) \to \underline{O}_P(1)$, where p is the structural morphism of P. Then X identifies with the scheme of flags of $\underline{F}$ over P, and the induction hypothesis shows that it is enough to prove 3.1 for P. Taking VI 1.1 into account, the result to be proved can therefore be stated as follows:

Corollary 3.2. Let S be a noetherian scheme, $\underline{E}$ a locally free $\underline{O}_S$-Module of rank n, $X = \mathbb{P}(\underline{E})$ the associated projective bundle, f the structural morphism of X, and ξ the class in $K^\bullet(X)$ of the sheaf $\underline{O}_X(1)$. Then the homomorphism:

$$\varphi : (K_\bullet(S))^n \longrightarrow K_\bullet(X)$$

defined by:

$$\varphi((x_i)) = \sum_{i=0}^{n-1} \xi^{-i} f^*(x_i)$$

is an isomorphism.

We have already seen that this homomorphism is surjective, since the ξ^i, $i = 0, \ldots, -n+1$, form a basis of $K^\bullet(X)$ over $K^\bullet(S)$; to see that it is injective, use the homomorphism f_*, defined because f is projective. [print 512]Then, by IV (2.12.4), one has:

$$f_*\left(\sum_{i=0}^{n-1} \xi^{-i} f^*(x_i)^{[510]}\right) = \sum_{i=0}^{n-1} x_i f_*(\xi^{-i}) = x_0,$$

the last equality following from the fact that for $-n+1 \leq i < 0$ one has $f_*(\xi^i) = 0$ (VI 2.8). Suppose that $\varphi((x_i)) = 0$. Then the preceding argument shows that $x_0 = 0$; applying the same reasoning to the elements $\xi^i \varphi((x_i))$ for $i = 1, \ldots, n-1$, one deduces by induction that the x_i are all zero, whence injectivity.

3.3. Let S still be a noetherian scheme, and $\underline{E}$ a locally free $\underline{O}_S$-Module of rank n. Denote by $V = \mathbb{V}(\underline{E})$ the vector bundle (contravariant in E!) associated with $\underline{E}$ (SGA 3 I 4.6), regarded as a group scheme over S. Let P be a torsor under V; we wish to compute $K_\bullet(P)$.

Recall how one can embed P in a projective bundle over S. Torsors under V correspond to extensions $\underline{F}$ of $\underline{E}$ by $\underline{O}_S$. Indeed, given an extension:

$$0 \longrightarrow \underline{O}_S \xrightarrow{s} \underline{F} \xrightarrow{u} \underline{E} \longrightarrow 0, \tag{3.3.1}$$

one obtains a torsor under V by considering the homomorphism $\mathbb{V}(\underline{F}) \to \mathbb{V}(\underline{O}_S)$ defined by s, the section of $\mathbb{V}(\underline{O}_S) = \mathrm{Spec}(\underline{O}_S[T])$ defined by $T = 1$, and taking $P = \mathbb{V}(\underline{F}) \times_{\mathbb{V}(\underline{O}_S)} S$, the actions of V on P being induced by those of V on $\mathbb{V}(\underline{F})$.

Conversely, let P be a torsor under V. Denote by $\underline{P}$ the sheaf of germs of homomorphisms from S to P; it is a principal homogeneous sheaf under the sheaf of germs of homomorphisms from S to V, which identifies with the dual $\check{\underline{E}}$ of $\underline{E}$. In the classical way this yields an extension of $\underline{O}_S$ by $\check{\underline{E}}$, whence, by passing to duals, the sequence (3.3.1).

[print 513]

Thus let P be a torsor under V, corresponding to an extension $\underline{F}$ of $\underline{E}$ by $\underline{O}_S$. The algebra $\underline{A}(P)$ of P over $\underline{O}_S$ is $\mathbb{S}(\underline{F}) \otimes_{\mathbb{S}(\underline{O}_S)} \underline{O}_S$, that is, $\mathbb{S}(\underline{F})/(s-1)\mathbb{S}(\underline{F})$, which identifies with the ring of homogeneous fractions of degree 0, $\mathbb{S}(\underline{F})_{(s)}$. Consequently P identifies with the open $D_+(s)$ of $\mathbb{P}(\underline{F})$. The complement of P in $\mathbb{P}(\underline{F})$ is then $\mathrm{Proj}(\mathbb{S}(\underline{F})/s \cdot \mathbb{S}(\underline{F}))$, which identifies, by (3.3.1), with $\mathbb{P}(\underline{E})$. Thus one has the immersions:

$$\begin{array}{ccccc} \mathbb{P}(\underline{E}) & \hookrightarrow & \mathbb{P}(\underline{F}) & \hookleftarrow & P \\ & \searrow & \downarrow & \swarrow & \\ & & S & & \end{array}$$

where $\mathbb{P}(\underline{E})$ is closed in $\mathbb{P}(\underline{F})$, and P is open and complementary to $\mathbb{P}(\underline{E})$.

Proposition 3.4. Let S be a noetherian scheme, $\underline{E}$ a locally free $\underline{O}_S$-Module of rank n, $V = \mathbb{V}(\underline{E})$ the associated vector bundle, P a torsor under V, and f the structural morphism of P. Then the canonical homomorphism:

$$f^* : K_\bullet(S) \longrightarrow K_\bullet(P)$$

is an isomorphism.

With the identifications of 3.3, one has the exact sequence:

$$K_\bullet(\mathbb{P}(\underline{E})) \xrightarrow{i_*} K_\bullet(\mathbb{P}(\underline{F})) \longrightarrow K_\bullet(P) \longrightarrow 0, \tag{3.4.1}$$

where i denotes the closed immersion $\mathbb{P}(\underline{E}) \to \mathbb{P}(\underline{F})$ defined by u. By 3.2, one has $K_\bullet(\mathbb{P}(\underline{E})) = K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(\mathbb{P}(\underline{E}))$, $K_\bullet(\mathbb{P}(\underline{F})) = K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(\mathbb{P}(\underline{F}))$. Since $\mathbb{P}(\underline{E})$ and $\mathbb{P}(\underline{F})$ are smooth, i is a regular immersion, and one therefore has a homomorphism $i_* : K^\bullet(\mathbb{P}(\underline{E})) \to K^\bullet(\mathbb{P}(\underline{F}))$; the homomorphism between the $K_\bullet$ is then obtained by tensoring the homomorphism between the $K^\bullet$ by $K_\bullet(S)$: to see this, it is enough to show that for $a \in K_\bullet(S)$ and [print 514]$b \in K^\bullet(\mathbb{P}(\underline{E}))$, one has

$$i_*(b \cdot h^*(a)) = i_*(b) \cdot g^*(a),$$

where g and h are the structural morphisms of $\mathbb{P}(\underline{F})$ and $\mathbb{P}(\underline{E})$. Since $h = g \circ i$, one has:

$$i_*(b \cdot h^*(a)) = i_*(b \cdot i^* g^*(a)) = g^*(a) \cdot i_*(b)$$

by IV 2.12.4.

By (3.4.1), one therefore has:

$$K_\bullet(P) \simeq K_\bullet(S) \otimes_{K^\bullet(S)} \left(K^\bullet(\mathbb{P}(\underline{F}))/\operatorname{Im} K^\bullet(\mathbb{P}(\underline{E}))\right).$$

Let ξ_F and ξ_E be the classes of the canonical invertible sheaves on $\mathbb{P}(\underline{F})$ and $\mathbb{P}(\underline{E})$. Then, by VI 1.1, $K^\bullet(\mathbb{P}(\underline{F}))$ (resp. $K^\bullet(\mathbb{P}(\underline{E}))$) has as a basis over $K^\bullet(S)$ the elements ξ_F^k for $k = 0, \ldots, -n$ (resp. ξ_E^k for $k = 0, \ldots, -n+1$). The projection formula gives:

$$i_*(\xi_E^k) = i_*(i^*(\xi_F^k)) = \xi_F^k i_*(1).$$

If $\underline{I}$ is the ideal of $\mathbb{P}(\underline{E})$ in $\mathbb{P}(\underline{F})$, $\underline{I}$ is an invertible ideal, and one has $i_*(1) = 1 - I$. Moreover, $\underline{I}$ is defined by the ideal J of $\mathbb{S}(\underline{F})$ that is the kernel of $\mathbb{S}(\underline{F}) \to \mathbb{S}(\underline{E})$, and is generated by a sub-Module of $\underline{F}$ isomorphic to $\underline{O}_S$; consequently, $\underline{I} \simeq \underline{O}_{\mathbb{P}(\underline{F})}(-1)$. The image of $K^\bullet(\mathbb{P}(\underline{E}))$ is therefore the $K^\bullet(S)$-submodule generated by the $\xi_F^k - \xi_F^{k-1}$ for $k = 0, \ldots, -n+1$. The homomorphism $K^\bullet(S) \longrightarrow K^\bullet(\mathbb{P}(\underline{F}))/\operatorname{Im}(K^\bullet(\mathbb{P}(\underline{E})))$ is thus an isomorphism, and consequently the same holds for the homomorphism $K_\bullet(S) \to K_\bullet(P)$.

4. Principal bundles under the groups $\mathbf{Gl}(n)_S$

Proposition 4.1. Let S be a noetherian scheme, G the S-group $\mathbf{Gl}(n)_S$, P a torsor under G, and $\underline{E}$ the locally free $\underline{O}_S$-Module of rank n corresponding to P. Denote by E the class of $\underline{E}$ in $K^\bullet(S)$, and by J the λ-ideal generated [print 515]by $(E - n)$. If f is the structural morphism of P, then f^*, defined because f is flat, is surjective and defines an isomorphism:

$$K_\bullet(S)/J \cdot K_\bullet(S) \xrightarrow{\sim} K_\bullet(P).$$

Let $\underline{E}_0 = (\underline{O}_S)^n$, let $e_1, \ldots, e_n$ be the evident basis of $\underline{E}_0$, and let $\underline{E}_0^{(i)}$ be the sub-Module of $\underline{E}_0$ generated by $e_1, \ldots, e_i$. The subfunctor of G that assigns to every S-scheme S' the subgroup of automorphisms of $\underline{E}_{0S'}$ leaving invariant the submodules $\underline{E}_{0S'}^{(i)}$ for every i is representable by a group subscheme B of G, which is a Borel subgroup of G. The quotient sheaf $\underline{G}/\underline{B}$ (for the fpqc topology) is representable by the scheme of flags of $\underline{E}_0$, which will be denoted by X_0. Let $\underline{L}_0^{(i)}$ be the canonical quotients of the inverse image of $\underline{E}_0$ on X_0, let $\underline{G}_0^{(i)}$ be the kernel of $\underline{E}_{0X_0} \to \underline{L}_0^{(i)}$, and let $\underline{F}_0^{(i)} = \underline{G}_0^{(i-1)}/\underline{G}_0^{(i)} \simeq \operatorname{Ker}(\underline{L}_0^{(i)} \to \underline{L}_0^{(i-1)})$.

Moreover, B is the semidirect product of its unipotent radical U and its maximal subtorus T leaving the e_i invariant. Let $Y_0 = G/U$. Then Y_0 is a torsor with base X_0 under the group T_{X_0}.

If S' is an S-scheme, giving an S'-point of Y_0 is equivalent to giving, over an S'' faithfully flat and quasi-compact over S', an automorphism of $\underline{E}_{0S''}$ modulo the unipotents of $U_{S''}$, that is, a flag of $\underline{E}_{0S''}$ whose submodules are free and, for every i, a class of elements f_i^α free over $\underline{O}_{S''}$ such that $f_i^\alpha - f_i^\beta \in \underline{G}^{(i)}_{0S''}$ [511]; equivalently, it is giving a flag of $\underline{E}_{0S''}$ and, for every i, a basis of $\underline{F}^{(i)}_{0S''}$, these data being subject to the gluing condition on $S'' \times_{S'} S''$; by fpqc descent, this amounts to the same data on S'. One then sees immediately that Y_0 is the affine X_0-scheme defined by the algebra: [print 516]

$$\underline{A}_0 = \coprod_{k_i \in \mathbb{Z}} \underline{F}_0^{(1)^{\otimes k_1}} \otimes \underline{F}_0^{(2)^{\otimes k_2}} \otimes \cdots \otimes \underline{F}_0^{(n)^{\otimes k_n}}.$$

The group G acts on $\underline{E}_0$, on X_0 equipped with the sheaves $\underline{L}_0^{(i)}$, and on Y_0. Taking the product of these objects with P under G, one obtains the sheaf $\underline{E}$ associated with P, the scheme $X = \mathbb{D}(\underline{E}) = P/B$ of flags of $\underline{E}$, and the scheme $Y = P/U$. We shall denote the canonical sheaves on X by the same letters as their counterparts on X_0. Thus one has the diagram:

$$\begin{array}{ccccc} & & Y = P/U & & \\ & \nearrow & & \searrow & \\ P & & & & X = P/B = \mathbb{D}(\underline{E}) \ . \\ & \searrow & & \swarrow & \\ & & S & & \end{array}$$

Observe that P is a torsor with base Y and structural group U_Y, and that Y is a torsor with base X and structural group T_X, defined by the $\underline{O}_X$-Algebra:

$$\underline{A} = \coprod_{k_i \in \mathbb{Z}} \underline{F}^{(1)^{\otimes k_1}} \otimes \cdots \otimes \underline{F}^{(n)^{\otimes k_n}}.$$

The homomorphism $K_\bullet(S) \to K_\bullet(P)$ therefore factors as:

$$K_\bullet(S) \longrightarrow K_\bullet(X) \longrightarrow K_\bullet(Y) \longrightarrow K_\bullet(P).$$

By 3.1, one has $K_\bullet(X) = K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X)$. Let ξ_i be the class of $\underline{F}^{(i)}$ in $K^\bullet(X)$. Then, by 2.2, $K_\bullet(Y)$ is the quotient of $K_\bullet(X)$ by the submodule generated by the $(\xi_i - 1)K_\bullet(X)$. Thus, if I is the ideal of $K^\bullet(X)$ generated by the $\xi_i - 1$, one has:

$$K_\bullet(Y) = K_\bullet(S) \otimes_{K^\bullet(S)} (K^\bullet(X)/I).$$

[print 517]Now, by VI 4.8, $K^\bullet(X)$ is the $K^\bullet(S)$-algebra generated by the $\xi_i - 1$, subject to the relations deduced from:

$$\gamma_t(E - n) = \prod_{i=1}^{n} \gamma_t(\xi_i - 1).$$

Thus $K^\bullet(X)/I = K^\bullet(S)/J$, where J is the ideal generated by the $\gamma^j(E-n)$ for $j = 1, \ldots, n$, hence the λ-ideal generated by $E - n$. Consequently, $K_\bullet(Y) \simeq K_\bullet(S)/JK_\bullet(S)$. The proof is completed by the lemma:

Lemma 4.2. Let S be a noetherian scheme, B a parabolic subgroup of a reductive S-group, and P a torsor under the unipotent radical U of B; then one has an isomorphism:

$$K_\bullet(S) \xrightarrow{\sim} K_\bullet(P).$$

Indeed, there is a sequence of subgroup schemes of U:

$$U_0 = U \supset U_1 \supset U_2 \supset \cdots \supset U_p = e,$$

closed in B, finite because B is noetherian, and such that, for every i, U_i/U_{i+1} is isomorphic to an S-vector group (SGA 3 XXVI 2.1). Consequently, $P \to S$ factors through a sequence of torsors under vector groups, whence the result by 3.4.

Corollary 4.3. Let S be a noetherian scheme, $G = \mathrm{Gl}(n)_S$. Then the canonical homomorphism $K_\bullet(S) \longrightarrow K_\bullet(G)$ is an isomorphism.

This follows immediately from 4.1 applied to $P = G$, since then $\underline{E} = \underline{O}_S^n$.

[print 518]

Corollary 4.4. Under the hypotheses of 4.1, suppose S regular. Then one has an isomorphism:

$$K^\bullet(S)/J \xrightarrow{\sim} K^\bullet(P).$$

If S is regular, G is also regular, and so is P. Thus one has $K^\bullet(S) \simeq K_\bullet(S)$, $K^\bullet(P) \simeq K_\bullet(P)$, whence the result by 4.1.

[print 519]

EXPOSÉ X
FORMALISM OF INTERSECTIONS ON PROPER ALGEBRAIC SCHEMES

by O. Jussila
(with an appendix by A. Grothendieck)

1. Compatibility of the filtrations with the composition law

$$K^\bullet(X) \times K_\bullet(X) \longrightarrow K_\bullet(X).$$

1.1.

First consider a noetherian scheme X and the group $K_\bullet(X)$ of classes of coherent $\underline{O}_X$-modules. For every coherent $\underline{O}_X$-module $\underline{F}$, we shall denote its class in $K_\bullet(X)$ by $\mathrm{cl}_\bullet(\underline{F})$.

Definition 1.1.1. For every integer $n \in \mathbb{Z}$, let $\mathrm{Filt}_n(X)$ denote the subgroup of $K_\bullet(X)$ generated by the classes $\mathrm{cl}_\bullet(\underline{F})$ of coherent $\underline{O}_X$-modules $\underline{F}$ such that

$$\dim(\mathrm{Supp}(\underline{F})) \leq n.$$

This defines an increasing filtration $(\mathrm{Filt}_n(X))_{n\in\mathbb{Z}}$ of $K_\bullet(X)$ such that $\mathrm{Filt}_n(X) = 0$ for $n < 0$.

Proposition 1.1.2. Let X be a noetherian scheme, $\underline{F}$ a coherent $\underline{O}_X$-module such that $\dim(\mathrm{Supp}(\underline{F})) \leq n$, and $X(n)$ $(n \in \mathbb{Z})$ the subset $\{x \in X\,;\, \dim(\overline{\{x\}}) = n\}$ of X. For every $x \in X$, let $Z(x)$ denote the integral closed subscheme associated with $\overline{\{x\}}$. Then the length of the $\underline{O}_{X,x}$-module $\underline{F}_x$ is finite for every $x \in X(n)$ and nonzero for only finitely many of the $x \in X(n)$, and one has [print 520]

$$\mathrm{cl}_\bullet(\underline{F}) = \sum_{x\in X(n)} \mathrm{length}(\underline{F}_x)\,\mathrm{cl}_\bullet(\underline{O}_{Z(x)}) + w \tag{1.1.3}$$

where $w \in \mathrm{Filt}_{n-1}(X)$.

By immediate induction on n, one obtains:

Corollary 1.1.4. For every noetherian scheme X and every integer n, $\mathrm{Filt}_n(X)$ is the subgroup of $K_\bullet(X)$ generated by the classes $\mathrm{cl}_\bullet(\underline{O}_Z)$, where Z runs through the set of integral closed subschemes of dimension $\leq n$ in X.

Proof of 1.1.2: Let $X' = \mathrm{Supp}(\underline{F})$. Since X' is a noetherian space of dimension $\leq n$, the elements of $X' \cap X(n)$ are maximal in X', hence finite in number, and $\mathrm{length}(\underline{F}_x)$ is finite for every $x \in X' \cap X(n)$. This proves the first assertion.

Let $K = \mathrm{Coh}(X)$ be the category of coherent $\underline{O}_X$-modules, and let K' be the subset of $\mathrm{Ob}(K)$ consisting of the coherent $\underline{O}_X$-modules concentrated on X' that satisfy condition (1.1.3). Clearly K' contains the sheaves $\underline{O}_{Z(x)}$ with $x \in X'$. Thus it is enough to show that K' is an "exact" subset of $\mathrm{Ob}(K)$, since the dévissage lemma (EGA III 3.1.2) then gives $\underline{F} \in K'$.

Thus let

$$0 \longrightarrow \underline{G}' \longrightarrow G \longrightarrow \underline{G}'' \longrightarrow 0 \text{ [512]} \tag{1.1.5}$$

be an exact sequence of $\underline{O}_X$-modules with two of its terms in K'. Then the sheaves $\underline{G}'$, $\underline{G}$ and $\underline{G}''$ are all concentrated on X', so the lengths ℓ'_x, ℓ_x and ℓ''_x of the $\underline{O}_{X,x}$-modules $\underline{G}'_x$, $\underline{G}_x$ and $\underline{G}''_x$ are finite [print 521]for every $x \in X(n)$, and one has

$$\ell_x = \ell'_x + \ell''_x \qquad (x \in X(n)). \tag{1.1.6}$$

Assertion (1.1.3) for all the terms of (1.1.5) then follows immediately from the hypothesis and the equation

$$\mathrm{cl}_\bullet(\underline{G}) = \mathrm{cl}_\bullet(\underline{G}') + \mathrm{cl}_\bullet(\underline{G}''). \tag{1.1.7}$$

It is known that $K_\bullet$ is a covariant functor on the category of locally noetherian schemes with respect to proper morphisms. The following proposition asserts that our filtration is functorial:

Proposition 1.1.8. Let $f : X' \to X$ be a proper morphism of locally noetherian schemes, and let

$$f_* : K_\bullet(X') \longrightarrow K_\bullet(X)$$

be the morphism induced by f. Then

$$f_*(\mathrm{Filt}_n(X')) \subset \mathrm{Filt}_n(X) \tag{1.1.9}$$

for every integer n.

Proof: If $\underline{F}$ is a coherent $\underline{O}_{X'}$-module, then by definition (IV 2.11) one has:

$$f_*(\mathrm{cl}_\bullet(\underline{F})) = \sum_{i\in\mathbb{N}} (-1)^i \,\mathrm{cl}_\bullet(R^i f_*(\underline{F})).$$

Now the $\underline{O}_X$-modules $R^i f_*(\underline{F})$ are concentrated on $f(\mathrm{Supp}(\underline{F}))$, and by (EGA IV 5.4.1 (ii)) one has

$$\dim(\mathrm{Supp}(\underline{F})) \geq \dim(f(\mathrm{Supp}(\underline{F}))),$$

whence the assertion.

[print 522]

1.2.

Now let X be an arbitrary scheme, and consider the group $K^\bullet(X)$ of classes of locally free $\underline{O}_X$-modules (henceforth understood to be of finite type). It is an augmented pre-λ-algebra over the ring $H^0(X,\mathbb{Z})$ (V 3.9.1). The map $f \rightsquigarrow f \cdot 1 \in K^\bullet(X)$ allows us henceforth to identify $H^0(X,\mathbb{Z})$ with a subring of $K^\bullet(X)$. If $\underline{E}$ is a locally free $\underline{O}_X$-module, its class in $K^\bullet(X)$ will be denoted by $\mathrm{cl}^\bullet(\underline{E})$. Define the operators λ^k $(k \in \mathbb{N})$ by

$$\lambda^k(\mathrm{cl}^\bullet(\underline{E})) = \mathrm{cl}^\bullet(\Lambda^k_{\underline{O}}\underline{E}) \tag{1.2.1}$$

and by

$$\lambda^k(x+y) = \sum_{\substack{i+j=k \\ i,j\in\mathbb{N}}} \lambda^i(x)\lambda^j(y). \tag{1.2.2}$$

(V 2.2). Recall that in the λ-filtration $(\mathrm{Filt}^i(X))_{i\in\mathbb{Z}}$ of $K^\bullet(X)$, one has $\mathrm{Filt}^i(X) = K^\bullet(X)$ for $i \leq 0$, and that for $i \geq 1$, $\mathrm{Filt}^i(X)$ is the ideal generated by the products

$$P = \gamma^{i_1}(x_1)\cdots\gamma^{i_k}(x_k) \tag{1.2.3}$$

with $1 \leq i_1, \ldots, i_k$; $i_1 + \cdots + i_k \geq i$ and $x_1, \ldots, x_k \in I = \mathrm{Ker}(\varepsilon)$ (ε being the augmentation homomorphism, cf. Exposé V 3.10). It was proved (V 3.10) that $\mathrm{Filt}^i(X)$ is even the $H^0(X, \mathbb{Z})$-submodule (hence the abelian subgroup, when X is connected) generated by the products (1.2.3), where one may moreover suppose that the elements x_p ($p = 1, \ldots, k$) have the form

$$x_p = \mathrm{cl}^\bullet(\underline{E}_p) - \underline{n}_p, \tag{1.2.4}$$

with $\underline{E}_p$ a locally free $\underline{O}_X$-module of rank $\underline{n}_p \in H^0(X, \mathbb{Z})$.

[print 523]

1.3.

Recall (IV 2.10) that the bifunctor

$$(\underline{E}, \underline{F}) \rightsquigarrow \underline{E} \otimes_{\underline{O}_X} \underline{F},$$

where $\underline{E}$ is a locally free $\underline{O}_X$-module and $\underline{F}$ a coherent $\underline{O}_X$-module, induces a composition law

$$K^\bullet(X) \times K_\bullet(X) \longrightarrow K_\bullet(X) \tag{1.3.1}$$

that makes $K_\bullet(X)$ a unital $K^\bullet(X)$-module. The key result of the present exposé is the following

Theorem 1.3.2. Let X be a noetherian scheme. For all $i, j \in \mathbb{Z}$, the composition law (1.3.1) induces by restriction a composition law

$$\mathrm{Filt}^i(X) \times \mathrm{Filt}_j(X) \longrightarrow \mathrm{Filt}_{j-i}(X). \tag{1.3.3}$$

Corollary 1.3.3. If X is a noetherian scheme of dimension $\leq j$, then the canonical homomorphism

$$\Theta_X : K^\bullet(X) \longrightarrow K_\bullet(X)$$

defined by $\Theta_X(x) = x \cdot \mathrm{cl}_\bullet(\underline{O}_X)$ ($x \in K^\bullet(X)$) induces for every $i \in \mathbb{Z}$ a homomorphism

$$\mathrm{Filt}^i(X) \longrightarrow \mathrm{Filt}_{j-i}(X).$$

Proof: First let us show that (1.3.3), for $i + j \leq k \in \mathbb{Z}$ and for every integral closed subscheme of dimension $\leq j$, implies (1.3.2) for $i + j \leq k$. By (1.1.4), it is enough to show that for every $x \in \mathrm{Filt}^i(X)$ and for every integral closed subscheme Z of dimension $\leq j$ one has [print 524]

$$x \, \mathrm{cl}_\bullet(\underline{O}_Z) \in \mathrm{Filt}_{j-i}(X).$$

To avoid any confusion, denote by 1_Z the class[513] of $\underline{O}_Z$ in $K_\bullet(Z)$. If $\alpha : Z \to X$ is the canonical immersion, then the projection formula (IV 2.11.1.2)[514] gives

$$x \, \mathrm{cl}_\bullet(\underline{O}_Z) = x\alpha_*(1_Z) = \alpha_*(\alpha^*(x)1_Z).$$

One has $\alpha^*(x) \in \mathrm{Filt}^i(Z)$, so by (1.1.8) and (1.3.3) for $X = Z$, $i + j \leq k$, one obtains

$$\alpha_*(\alpha^*(x)1_Z) \in \mathrm{Filt}_{j-i}(X),$$

whence the assertion.

Theorem 1.3.2 is trivial for $i + j \leq 0$. Suppose by induction that it has been proved for $i + j < k$ and for every noetherian scheme X. By the preceding remark, it remains to prove that for every

integral noetherian scheme X of dimension j and every product (1.2.3) satisfying (1.2.4), one has

$$P\,\mathrm{cl}_\bullet(\underline{O}_X) \in \mathrm{Filt}_{j-i}(X). \tag{1.3.4}$$

If $k > 1$ in (1.2.3), the induction hypothesis and the associativity of the law (1.3.1) give the result. We may therefore suppose (X being connected) that

$$x = P = \gamma^m(\mathrm{cl}^\bullet(\underline{E}) - n), \tag{1.3.5}$$

where $m \geq i$ and $\underline{E}$ is a locally free $\underline{O}_X$-module of rank $n > 0$.

Distinguish two cases:

I. Suppose that one can find invertible $\underline{O}_X$-modules $\underline{L}_1, \ldots, \underline{L}_n$ such that

$$\mathrm{cl}^\bullet(\underline{E}) = \mathrm{cl}^\bullet(\underline{L}_1) + \cdots + \mathrm{cl}^\bullet(\underline{L}_n) \in K^\bullet(X).$$

[print 525]It follows (V) that x is the m-th elementary symmetric function in the $\mathrm{cl}^\bullet(\underline{L}_r) - 1$ ($r = 1, \ldots, n$), hence is a homogeneous polynomial of degree m with integer coefficients in the variables $(\mathrm{cl}^\bullet(\underline{L}_r) - 1)$ ($r = 1, \ldots, n$). Applying the induction hypothesis and the associativity of the law (1.3.1) again, we are thus reduced to proving that, for every invertible $\underline{O}_X$-module $\underline{L}$, one has

$$(\mathrm{cl}^\bullet(\underline{L}) - 1)\,\mathrm{cl}_\bullet(\underline{O}_X) \in \mathrm{Filt}_{j-1}(X) \tag{1.3.6}$$

Since the scheme X is integral, $\underline{L}$ and $\underline{O}_X$ are isomorphic to submodules of the constant $\underline{O}_X$-module $\underline{R}_X$ of rational functions on X. The sum $\underline{O}_X$-module $\underline{F} = \underline{O}_X + \underline{L}$ is coherent, and one obtains

$$\begin{aligned}(\mathrm{cl}^\bullet(\underline{L}) - 1)\,\mathrm{cl}_\bullet(\underline{O}_X) &= \mathrm{cl}_\bullet(\underline{L}) - \mathrm{cl}_\bullet(\underline{O}_X)\\ &= (\mathrm{cl}_\bullet(\underline{F}) - \mathrm{cl}_\bullet(\underline{O}_X)) - (\mathrm{cl}_\bullet(\underline{F}) - \mathrm{cl}_\bullet(\underline{L}))\\ &= \mathrm{cl}_\bullet(\underline{F}/\underline{O}_X) - \mathrm{cl}_\bullet(\underline{F}/\underline{L}).\end{aligned}$$

But the fibers at the generic point of the sheaves $\underline{F}/\underline{O}_X$ and $\underline{F}/\underline{L}$ are zero, so their supports in X are nowhere dense (X being irreducible), hence of dimension $\leq j - 1$. Assertion (1.3.6) follows immediately.

II. It remains to prove the general case. Introduce the flag bundle

$$g : D \longrightarrow X$$

of $\underline{E}$ over X, on which one can find invertible $\underline{O}_D$-modules $\underline{L}_1, \ldots, \underline{L}_n$ such that

$$g^*(\mathrm{cl}^\bullet(\underline{E})) = \mathrm{cl}^\bullet(L_1) + \cdots + \mathrm{cl}^\bullet(\underline{L}_n) \in K^\bullet(D). \tag{1.3.7}$$

Let e be the generic point of X, and $\underline{k}(e)$ the residue field of e. Then one can find a rational point u of the generic fiber D_e of D [print 526]over $\underline{k}(e)$, since these points are in bijective correspondence with the flags of type $(1, \ldots, 1)$ (n terms) of the vector space $\underline{E}_e$ of dimension $n > 0$ over $\underline{k}(e)$. On the other hand, these rational points also correspond to the germs of sections of g in a neighborhood of e, so one can find such a section $s : U \to D$ over an open subset U of X. Let $X' = \overline{s(U)}$ be the closure of $s(U)$ with its reduced induced structure, and let $f : X' \to X$ be the restriction of g to X'. The morphism f is projective and, since it is an isomorphism over $s(U)$, it is birational. Thus $\dim(X') \leq \dim(X)$ (EGA IV 5.6.6.1). On the other hand, $f(X') = g(X')$ is closed and contains U, so f is surjective and $\dim(X') \geq \dim(X)$ (EGA IV 5.4.1(ii)).

The hypotheses of case I are therefore satisfied for X' and $x' = f^*(x) \in \mathrm{Filt}^i(X')$, and one obtains

$$f^*(x)\,\mathrm{cl}_\bullet(\underline{O}_{X'}) \in \mathrm{Filt}_{j-i}(X'). \tag{1.3.7'}$$

Taking account of the projection formula and (1.1.8), one concludes that:

$$x f_*(\mathrm{cl}_\bullet(\underline{O}_{X'})) \in \mathrm{Filt}_{j-i}(X). \tag{1.3.8}$$

On the other hand, the canonical homomorphism

$$\underline{O}_X \longrightarrow f_*(\underline{O}_{X'})$$

is an isomorphism over the dense open subset U, and the $\underline{O}_X$-modules $R^i f_*(\underline{O}_{X'})$ $(i \geq 1)$ are concentrated on the nowhere dense closed subset $X - U$. It follows that

$$f_*(\mathrm{cl}_\bullet(\underline{O}_{X'})) = \mathrm{cl}_\bullet(\underline{O}_X) + w, \tag{1.3.9}$$

[print 527]where $w \in \mathrm{Filt}_{j-1}(X)$. By the induction hypothesis, one has $x \cdot w \in \mathrm{Filt}_{j-i-1}(X)$, whence, taking account of formulas (1.3.8) and (1.3.9), the assertion

$$x\,\mathrm{cl}_\bullet(\underline{O}_X) \in \mathrm{Filt}_{j-i}(X), \qquad \text{Q.E.D.}$$

Remark 1.4. Let X be a locally noetherian scheme. Observe that, in this section, we have used only the following properties of the "dimension function" $D(x) = D_Y(x) = \dim(\overline{\{x\}})$[515] on proper X-schemes Y:

(i) For every proper X-scheme Y and every $x \in Y$, one has $D_Y(x) \in \mathbb{Z}$.

(ii) If $y \neq x$ is a specialization of x, one has

$$D(y) < D(x).$$

(iii) For every X-morphism $f : Y_1 \to Y_2$ of proper X-schemes and every $y \in Y_1$, one has

$$D_{Y_2}(f(y)) \leq D_{Y_1}(y).$$

(iv) Equality holds in (iii) when f is an immersion, or when Y_1 is irreducible, f is birational and surjective, and y is the generic point of Y_1.

This means that all the results of this section, in particular (1.3.2) and (1.3.3), remain valid unchanged when "dimension" is replaced by a function D satisfying conditions (i) to (iv) above and the filtration of $K_\bullet(X)$ is defined using such a function D.

[print 528]

Example 1.5. Let S be a locally noetherian universally catenary scheme (for example, a subscheme of a Cohen–Macaulay scheme). Put

$$D_S(x) = n - \mathrm{codim}(\overline{\{x\}}, S) = n - \dim. \underline{O}_{S,x} \qquad (x \in S), \tag{1.5.1}$$

and, for every proper S-scheme $s_Y : Y \to S$ and $y \in Y$, also put:

$$D_Y(y) = D_S(s_Y(y)) + \deg. \mathrm{tr}_{k(s_Y(y))}\, k(y). \tag{1.5.2}$$

If $f : Y_1 \to Y_2$ is an S-morphism of proper S-schemes, the additivity of transcendence degrees and (1.5.2) immediately give

$$D_{Y_1}(y) = D_{Y_2}(f(y)) + \deg. \mathrm{tr}_{k(f(y))}\, k(y), \qquad (y \in Y_1). \tag{1.5.3}$$

Let us show that the functions D_Y so defined satisfy conditions (i)–(iv) of 1.4. By (1.5.3), condition (ii) is the only nontrivial one. By (iv), one may clearly suppose that $Y = \overline{\{x\}}$. Then factor the structural morphism s_Y of Y through its image $s_Y(Y) = Z \hookrightarrow S$, say $s_Y = g \circ f$:

$$Y \xrightarrow{f} Z \xhookrightarrow{g} S,$$

where Y and Z are irreducible, f is proper and surjective, and g is a closed immersion. Put $x' = s_Y(x)$, $y' = s_Y(y)$. Then (1.5.2) and (1.5.1) give:

$$D_Y(x) - D_Y(y) = \dim. \underline{O}_{S,y'} - \dim. \underline{O}_{S,x'} + \deg. \operatorname{tr}_{k(x')} k(x) - \deg. \operatorname{tr}_{k(y')} k(y).$$

Since S is catenary, one has [print 529]

$$\dim. \underline{O}_{S,y'} - \dim. \underline{O}_{S,x'} = \dim. \underline{O}_{Z,y'} - \dim. \underline{O}_{Z,x'}$$

(EGA 0_{IV} 14.3.2.1), whence

$$D_Y(x) - D_Y(y) = [\dim. \underline{O}_{Z,y'} - \deg. \operatorname{tr}_{k(y')} k(y)] - [\dim. \underline{O}_{Z,x'} - \deg. \operatorname{tr}_{\underline{k}(x')} \underline{k}(x)]. \qquad (1.5.4)$$

Since Z is universally catenary, (EGA IV 5.6.5.1.) applies and gives

$$C_1 = \dim. \underline{O}_{Y,y} - e, \qquad C_2 = \dim. \underline{O}_{Y,x} - e,$$

where C_1 and C_2 are the two terms on the right in (1.5.4). Then (1.5.4) immediately gives

$$D_Y(x) - D_Y(y) = \dim. \underline{O}_{Y,y} - \dim. \underline{O}_{Y,x} > 0, \qquad \text{Q.E.D.}$$

Thus (1.3.2) and (1.3.3) remain valid for a noetherian universally catenary scheme X of dimension n when the filtration of $K_\bullet(X)$ is defined by means of (1.5.1) according to the codimension of supports. Observe that this filtration coincides with that of (1.1.1) when, in addition, X is biequidimensional.

1.6.

One may ask whether it is always possible to define, on a noetherian scheme X of finite dimension, a function D having property (i) of 1.4 and also the following property, stronger than (ii):

(ii bis) If $x \in X$ and $y \neq x$ is an immediate specialization of x, then $D(x) = D(y) + 1$.

[print 530]

For this it is clearly necessary that X be catenary, and sufficient that X be biequidimensional (EGA 0_{IV} 14.3.3). Nevertheless, there exist noetherian schemes of finite dimension, catenary and equidimensional, having no such functions D. To see this, one can modify the example (EGA IV 5.6.11) as follows: take two copies Y_1 and Y_2 of the spectrum of the ring E of loc. cit., and denote by y_i and z_i $(i = 1, 2)$ the closed points of Y_i $(i = 1, 2)$ corresponding to the maximal ideals $\underline{m}$ and $\underline{m}'$ of E, of heights 2 and 1 respectively. Recall that the residue fields of $\underline{m}$ and $\underline{m}'$ are isomorphic. Thus one can construct a scheme X by gluing Y_1 and Y_2, identifying y_1 with z_2 and y_2 with z_1. One can then identify Y_1 and Y_2 with the two irreducible components of X, and then, inside X[516], $Y_1 \cap Y_2 = \{y_1, y_2\}$ has dimension zero. Thus X is a noetherian, catenary and equidimensional scheme (of dimension 2). If (ii bis) held on X, one would have

$$D(y_1) = D(z_1) - 1 = D(y_2) - 1$$

and, on the other hand,

$$D(y_1) = D(z_2) = D(y_2) + 1,$$

giving a contradiction.

[print 531]

2. Snapper polynomials

2.1.

First recall the definition and some elementary properties of polynomial maps:

Definition 2.1.1. Let f be a map from an abelian group A to an abelian group B. We say that f is a polynomial map when, for every sequence $x_1, \ldots, x_r$[517] ($r \geq 1$) of elements of A, there is a family $(y(x_1, \ldots, x_r; k_1, \ldots, k_r))_{k_1,\ldots,k_r \geq 0}$ of elements of B, almost all zero, satisfying the following condition: For every sequence $n_1, \ldots, n_r$ of integers, one has

$$f(n_1x_1 + \cdots + n_rx_r) = \sum_{k_1,\ldots,k_r \geq 0} (n_1^{k_1} \cdots n_r^{k_r}) y(x_1, \ldots, x_r; k_1, \ldots, k_r).$$

We say that f has degree $\leq m$ when, for all $r \in \mathbb{Z}$ and $x_1, \ldots, x_r \in A$, one has

$$y(x_1, \ldots, x_r; k_1, \ldots, k_r) = 0$$

for $k_1 + \cdots + k_r > m$.

Lemma 2.1.2. Every homomorphism of abelian groups is a polynomial map of degree ≤ 1. If $h = f \circ g$, where f is a polynomial map of degree p and g a polynomial map of degree q, then h is a polynomial map of degree $\leq pq$.

The proof is trivial.

Lemma 2.1.3. Let

$$0 \longrightarrow G' \longrightarrow G \xrightarrow{j} G'' \longrightarrow 0$$

be an exact sequence of abelian groups such that G' is divisible by [print 532]a prime number p and G'' is a torsion group. Then every polynomial map $f : G \to \mathbb{Q}$ that factors through $\mathbb{Z}$ is constant.

Proof: We must show that for every $x \in G$ one has $f(x) = f(0)$.

a) First suppose that x is a torsion element, say $mx = 0$ ($m \in Z$, $m \neq 0$). Then the polynomial $P_x(n) = f(nx) - f(0)$ has infinitely many roots, namely all integers divisible by m. Thus the polynomial P_x is identically zero, and one has $f(x) = f(0)$.

b) Now suppose that x is an element of infinite order in G divisible by every power p^k ($k \in \mathbb{Z}$). Consider the polynomials P_k defined by

$$P_k(n) = f(nx/p^k) = \sum_{i \in \mathbb{N}} n^i y(x/p^k; i) \in \mathbb{Z} \qquad (n \in \mathbb{Z};\ y(x/p^k, i) \in \mathbb{Q}).$$

For every $n \in \mathbb{Z}$ one has the identity

$$P_k(n) = P_{k+1}(pn)$$

i.e.

$$\sum_{i \in \mathbb{N}} n^i y(x/p^k; i) = \sum_{i \in \mathbb{N}} p^i n^i y(x/p^{k+1}; i),$$

from which comparison of coefficients gives

$$y(x/p^k; i) = p^i y(x/p^{k+1}; i) \qquad (i \in \mathbb{N}).$$

[print 533]But every polynomial $P : Q \to \mathbb{Q}$ is continuous for the topology induced by $\mathbb{R}$. Thus for sufficiently large k one has

$$|P_x(n/p^k) - P_x(0)| < 1.$$

By definition, however, $P_x(n/p^k)$ is an integer for every $n, k \in \mathbb{Z}$. It follows that $P_x(n/p^k) - P_x(0) = 0$ for sufficiently large k. Thus the polynomial P_x is constant, and in particular one obtains $P_x(1) = P_x(0)$, i.e. $f(x) = f(0)$.

The two cases a) and b) above show that the restriction of f to G' is constant. We shall now show that f is constant on each coset in G modulo G'. For every $b \in G$, denote by $g_b : G' \to \underline{Q}$ the map defined by

$$g_b(y) = f(b + y) \qquad (y \in G').$$

The map g_b is a polynomial map from G' to $\mathbb{Q}$ that factors through $\mathbb{Z}$, hence is constant by the preceding argument, whence the assertion that f is constant on the cosets in G modulo G'.[518] It follows that one can write $f = h \circ j$, where $h : G'' \to \mathbb{Q}$[519] is a polynomial map that factors through $\mathbb{Z}$. But by hypothesis G'' is a torsion group[520], so h is constant. Thus f is constant, Q.E.D.

Lemma 2.1.4. Let A be a unital commutative ring and I a nilpotent ideal of A. Let P be the multiplicative subgroup of A^* consisting of the invertible elements of A of the form $x = 1 + y$ with $y \in I$, and let $i : P \to A$ be the natural injection. Then the composite map [print 534]

$$\alpha : P \xrightarrow{i} A \xrightarrow{j} A \otimes_{\mathbb{Z}} \mathbb{Q}$$

(where $j(x) = x \otimes 1$) is a polynomial map. If $I^{m+1} = 0$, then α has degree $\leq m$.

Proof: Let $x_1, \ldots, x_r$ be a sequence of elements of P with[521] $x_i = 1 + y_i$ $(i = 1, \ldots, r)$. Then the binomial formula gives, for every $n_1, \ldots, n_r \in Z$:

$$x_1^{n_1} \cdots x_r^{n_r} = \sum_{k_1, \ldots, k_r \geq 0} \binom{n_1}{k_1} \cdots \binom{n_r}{k_r} y_1^{k_1} \cdots y_r^{k_r}, \tag{2.1.5}$$

where the sum on the right is finite because the elements $y_1, \ldots, y_r$ are nilpotent. But the coefficients $\binom{n_1}{k_1} \cdots \binom{n_r}{k_r}$ are polynomials with rational coefficients in the variables $n_1, \ldots, n_r$, of degree $k_1 + \cdots + k_r$, so α is indeed a polynomial map. If $I^{m+1} = 0$, then $y_1^{k_1} \cdots y_r^{k_r} = 0$ for $k_1 + \cdots + k_r > m$. Thus α has degree $\leq m$, Q.E.D.

2.2.

Now consider a noetherian scheme X and an element y of $\mathrm{Filt}_j(X)$ (1.1.1). Associating to every $x \in K^\bullet(X)$ the element xy of $K_\bullet(X)$ gives a homomorphism of $K^\bullet(X)$-modules

$$\Theta_y : K^\bullet(X) \longrightarrow K_\bullet(X)$$

which, taking account of (1.3.2), factors through the canonical epimorphism

$$j : K^\bullet(X) \longrightarrow K^\bullet(X)/\operatorname{Filt}^{j+1}(X) = A.$$

[print 535]Let $\beta_y : A \to K_\bullet(X)$ be the homomorphism so defined. Denoting by I the ideal $\operatorname{Filt}^1(X)/\operatorname{Filt}^{j+1}(X)$ of A, one has $I^{j+1} = 0$. Keeping the notation of Lemma 2.1.4, one obtains a homomorphism of abelian groups

$$\gamma : \mathrm{Pic}(X) \longrightarrow P$$

that associates to the class in $\mathrm{Pic}(X)$ of every invertible $\underline{O}_X$-module $\underline{L}$ the element $j(\mathrm{cl}^\bullet(\underline{L}))$ of P.

Lemmas 2.1.2 and 2.1.4 immediately give:

Proposition 2.2.1. For every noetherian scheme X and every $y \in \mathrm{Filt}_j(X)$ $(j \in \mathbb{Z})$, the composite map

$$\mathrm{Pic}(X) \longrightarrow K_\bullet(X) \otimes_{\mathbb{Z}} \mathbb{Q},$$

defined by

$$\mathrm{cl}_P(\underline{L}) \rightsquigarrow (\mathrm{cl}^\bullet(\underline{L})y) \otimes 1$$

is a polynomial map of degree $\leq j$.

2.3.

If X is a proper scheme over a field k and $g : X \to \mathrm{Spec}(k)$ is the structural morphism, then the homomorphism

$$g_* : K_\bullet(X) \longrightarrow K_\bullet(\mathrm{Spec}(k)) \cong \mathbb{Z}$$

associates to the class $\mathrm{cl}_\bullet(\underline{F})$ of every coherent $\underline{O}_X$-module $\underline{F}$ the Euler–Poincaré characteristic $\chi(\underline{F}) \in \mathbb{Z}$ of $\underline{F}$ on X. From 2.2.1 one immediately concludes:

[print 536]

Proposition 2.3.1. For every proper scheme X over k, and every $y \in \mathrm{Filt}_j(X)$, the map

$$s_y : \mathrm{Pic}(X) \longrightarrow {}^{[522]}\mathbb{Z} \hookrightarrow \mathbb{Q}$$

defined by

$$\mathrm{cl}_P(\underline{L}) \rightsquigarrow g_*(\mathrm{cl}^\bullet(\underline{L})y) = \chi(\mathrm{cl}^\bullet(\underline{L})y)^{[523]}$$

is a polynomial map of degree $\leq j$.

When $\underline{L}_1, \ldots, \underline{L}_n$ are invertible $\underline{O}_X$-modules and $\underline{F}$ is a coherent $\underline{O}_X$-module such that $\dim(\mathrm{Supp}(\underline{F})) = j$, then 2.3.1. implies in particular that the Euler–Poincaré characteristic

$$\chi(\underline{L}_1^{\otimes m_1} \otimes \cdots \otimes \underline{L}_n^{\otimes m_n} \otimes \underline{F}) \tag{2.3.2}$$

is a numerical polynomial in the variables $m_1, \ldots, m_n \in \mathbb{Z}$, of degree $\leq$ dim. supp. F, sometimes called the Snapper polynomial for $\underline{F}$ and the sequence $\underline{L}_1, \ldots, \underline{L}_n$.

2.4.

Recall [2] that the Picard functor $\underline{\mathrm{Pic}}_{X/k}$ of X over k is the sheaf of groups for the (fpqc) topology over k associated with the presheaf

$$T \rightsquigarrow \mathrm{Pic}(X \times_k T) \qquad (T \in \mathrm{Ob}(\mathrm{Sch}\,/k)).$$

When k is algebraically closed, the canonical homomorphism

$$\mathrm{Pic}(X) \longrightarrow \mathrm{Pic}_{X/k}(\mathrm{Spec}(k))$$

is known to be an isomorphism.

[print 537]

When X is proper over k, it is known [4] that $\underline{\mathrm{Pic}}_{X/k}$ [524] is representable by a group scheme $\underline{\mathrm{Pic}}_{X/k}$ locally of finite type over k. The neutral component $\underline{\mathrm{Pic}}^0_{X/k}$ of $\underline{\mathrm{Pic}}_{X/k}$ is a connected commutative algebraic group over k.

Definition 2.4.1. Let X be a proper scheme over a field k, and $\underline{L}$ an invertible $\underline{O}_X$-module (resp. x an element of $\mathrm{Pic}(X)$). We say that $\underline{L}$ (resp. x) is τ-equivalent to zero when there is

an integer $n \neq 0$ such that $\underline{L}^{\otimes n}$ (resp. nx) is algebraically equivalent to zero. The elements of $\mathrm{Pic}(X)$ algebraically equivalent (resp. τ-equivalent) to zero form a subgroup $\mathrm{Pic}^0(X)$ (resp. $\mathrm{Pic}^\tau(X)$) of $\mathrm{Pic}(X)$.

Lemma 2.4.3. For every proper scheme X over an algebraically closed field k, the group $\mathrm{Pic}^0(X)$ is divisible by every integer n prime to the characteristic of k.

Proof: Since k is algebraically closed, one has $\mathrm{Pic}^0(X) \cong \underline{\mathrm{Pic}}^0_{X/k}(k)$. Since $\underline{\mathrm{Pic}}^0_{X/k}$ is connected and of finite type, the n-th-power endomorphism is surjective when n is prime to the characteristic of k (SGA 3 XV 1.3 and VI 1.3.2). But k is algebraically closed, so the corresponding endomorphism of the group $\underline{\mathrm{Pic}}^0_{X/k}(k)$ is surjective, Q.E.D.

From 2.4.3 one concludes that, in the exact sequence

$$0 \longrightarrow \mathrm{Pic}^0(X) \longrightarrow \mathrm{Pic}^\tau(X) \longrightarrow T \longrightarrow 0,$$

$\mathrm{Pic}^0(X)$ is divisible by a prime number p and T is a torsion group, so one can apply 2.1.3 to the restriction of the polynomial map (2.3.1) to $\mathrm{Pic}^\tau(X)$. One thus obtains:

[print 538]

Proposition 2.4.4. The polynomial map s_y (2.3.1) is constant on the cosets modulo $\mathrm{Pic}^\tau(X)$.

Proof: If K is an algebraically closed extension of k, one obtains a commutative diagram

$$\begin{array}{ccc} \mathrm{Pic}(X) & \xrightarrow{s_y} & K_\bullet(\mathrm{Spec}(k)) \\ \downarrow & & \searrow^{\sim} \\ & & \mathbb{Z} \hookrightarrow \mathbb{Q} \\ & & \nearrow_{\sim} \\ \mathrm{Pic}(X_K) & \xrightarrow{\underline{s}_y} & K_\bullet(\mathrm{Spec}(K)) \end{array},$$

where $\underline{s}_y$ is the polynomial map on $\mathrm{Pic}(X_K)$ corresponding to s_y. Thus one may suppose that k is algebraically closed. The assertion then follows immediately from 2.1.3 and 2.4.3.

Remark 2.4.5. In Exp. XIII this result will be proved without using the representability of the functor $\mathrm{Pic}_{X/k}$.

3. Projection formulas for the associated graded groups

3.1.

Consider a noetherian scheme X. Denote by $\mathrm{Gr}^\bullet(X)$ the graded group associated with the λ-filtration of $K^\bullet(X)$, and by $\mathrm{Gr}_\bullet(X)$ the graded group associated with the filtration 1.1.1 of $K_\bullet(X)$. By 1.3.2, the composition law (1.3.1)

$$K^\bullet(X) \times K_\bullet(X) \longrightarrow K_\bullet(X)$$

induces, on passing to the associated graded groups, composition laws

$$\mathrm{Gr}^i(X) \times \mathrm{Gr}_j(X) \longrightarrow \mathrm{Gr}_{j-i}(X) \qquad (i, j \in \mathbb{Z}). \tag{3.1.1}$$

[print 539]$\mathrm{Gr}_\bullet(X)$ thus becomes a module over the ring $\mathrm{Gr}^\bullet(X)$ (and even a graded module over the graded ring $\mathrm{Gr}^\bullet(X)$, provided the sign of the grading of $\mathrm{Gr}_\bullet(X)$ is changed by putting $\mathrm{Gr}^i_\bullet(X) = \mathrm{Gr}_{-i}(X)$ $(i \in \mathbb{Z})$). Now consider a proper morphism

$$f : X' \longrightarrow X$$

of noetherian schemes. Then f induces a homomorphism

$$f^* : \mathrm{Gr}^\bullet(X) \longrightarrow \mathrm{Gr}^\bullet(X')$$

of graded rings (V), and, by 1.1.8, a homomorphism

$$f_* : \mathrm{Gr}_\bullet(X') \longrightarrow \mathrm{Gr}_\bullet(X)$$

of graded abelian groups. Passing to the associated graded groups, the projection formula (IV 2.11) and (3.1.1) then give the graded projection formula

$$x \cdot f_*(y) = f_*(f^*(x) \cdot y) \qquad (x \in \mathrm{Gr}^\bullet(X),\ y \in \mathrm{Gr}_\bullet(X')). \tag{3.1.2}$$

3.2.

Let $f : X' \to X$ be a projective complete-intersection morphism (Exp. VIII) of virtual dimension r between noetherian schemes, with X carrying an ample invertible $\underline{O}_X$-module. In Exp. VIII it was proved that f induces a homomorphism

$$f_* : \mathrm{Filt}^k(X')_{\mathbb{Q}} \longrightarrow \mathrm{Filt}^{k-r}(X)_{\mathbb{Q}} \qquad (k \in \mathbb{Z}) \tag{3.2.1}$$

which moreover satisfies the projection formula (Exp. IV):

$$f_*(f^*(x) \cdot y) = x \cdot f_*(y) \qquad (x \in K^\bullet(X)_{\mathbb{Q}},\ y \in K^\bullet(X')_{\mathbb{Q}}). \tag{3.2.2}$$

[525]

[print 540]Passing to the associated graded groups in (3.2.1) and (3.2.2), one obtains a homomorphism of degree $-r$:

$$f_* : \mathrm{Gr}^\bullet(X')_{\mathbb{Q}} \longrightarrow \mathrm{Gr}^\bullet(X)_{\mathbb{Q}} \tag{3.2.3}$$

and, by (1.3.2), a graded projection formula

$$f_*(f^*(x) \cdot y) = x \cdot f_*(y) \qquad (x \in \mathrm{Gr}^\bullet(X)_{\mathbb{Q}},\ y \in \mathrm{Gr}^\bullet(X')_{\mathbb{Q}}).$$

3.3.

Finally, consider a morphism $f : X' \to X$ of finite Tor-dimension between noetherian schemes. Then it is known (Exp. IV) that f induces a homomorphism

$$f^* : K_\bullet(X) \longrightarrow K_\bullet(X')$$

such that

$$f^*(\mathrm{cl}_\bullet(\underline{F})) = \sum_i (-1)^i \,\mathrm{cl}_\bullet(\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{F}, \underline{O}_{X'}))$$

for every coherent $\underline{O}_X$-module $\underline{F}$.

The $\underline{O}_X$-modules $\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{F}, \underline{O}_{X'})$ are concentrated on $f^{-1}(\mathrm{Supp}(\underline{F}))$, and when the fibers of f are everywhere of dimension $\leq d$, it is known (EGA IV 5.6.7) that

$$\dim(f^{-1}(\mathrm{Supp}(\underline{F})))^{[526]} \leq \dim(\mathrm{Supp}(\underline{F})) + d.$$

It follows that f induces, for every $k \in \mathbb{Z}$, a homomorphism

$$f^* : \mathrm{Filt}_k(X) \longrightarrow \mathrm{Filt}_{k+d}(X'),$$

whence a homomorphism of degree d of graded abelian groups [print 541]

$$f^* : \mathrm{Gr}_\bullet(X) \longrightarrow \mathrm{Gr}_\bullet(X'). \tag{3.3.1}$$

3.4.

Finally, suppose that the morphisms (3.2.3) and (3.3.1) are both defined for a morphism $f : X' \to X$ such that $r = d$. This is the case, for example, when X is quasi-compact with an ample $\underline{O}_X$-module and $X' = P_X(\underline{E})$, where $\underline{E}$ is a locally free $\underline{O}_X$-module of rank $r+1$. Then, for every $x \in \mathrm{Filt}^i(X')_{\mathbb{Q}}$ and $y \in \mathrm{Filt}_j(X)_{\mathbb{Q}}$, the projection formula in the derived category gives the projection formula

$$f_*(x \cdot f^*(y)) = f_*(x) \cdot y \in \mathrm{Filt}_{j+r}(X)_{\mathbb{Q}}.$$

When X is noetherian, 1.3.2 yields the corresponding graded formula.

4. Intersection numbers

4.1.

Consider a proper scheme X over a field k. Composing the product (3.1.1)

$$\mathrm{Gr}^i(X) \times \mathrm{Gr}_i(X) \longrightarrow \mathrm{Gr}_0(X) = \mathrm{Filt}_0(X) \qquad (i \in \mathbb{Z})$$

with the Euler–Poincaré characteristic

$$\chi : \mathrm{Filt}_0(X) \longrightarrow \mathbb{Z},$$

gives a pairing called the "intersection number":

$$\mathrm{Gr}^i(X) \times \mathrm{Gr}_i(X) \longrightarrow \mathbb{Z}, \tag{4.1.1}$$

[print 542]which we shall denote by

$$(x, y) \rightsquigarrow \langle x, y \rangle \in \mathbb{Z} \qquad (i \in \mathbb{Z},\ x \in \mathrm{Gr}^i(X),\ y \in \mathrm{Gr}_i(X)).$$

Example 4.2. Suppose that the scheme X has dimension n. Take an element $y \in \mathrm{Gr}_n(X)$, for example $y = \mathrm{cl}_\bullet(\underline{O}_Z)$, where Z is an integral subscheme of dimension n in X. Also take a homogeneous polynomial of weight n

$$P \in \mathbb{Z}[T_1, \ldots, T_n],$$

where T_i is assigned weight i $(i = 1, \ldots, n)$, for example a monomial $T_{i_1} \cdot T_{i_2} \cdots T_{i_k}$ with $i_1 + i_2 + \cdots + i_k = n$. Substituting in P, for each variable T_i $(i = 1, \ldots, n)$, the Chern class

$$c_i(x) = \mathrm{cl}^\bullet(\gamma^i(x - \epsilon(x)))^{[527]} \in \mathrm{Gr}^i(X) \qquad (x \in K^\bullet(X)),$$

gives an element

$$P(c_1(x), \ldots, c_n(x)) \in \mathrm{Gr}^n(X)^{[528]}.$$

The integer

$$P_y(x) = \langle P(c_1(x), \ldots, c_n(x)), y \rangle.^{[529]}$$

is called the Chern number of x relative to y. When $y = \mathrm{cl}_\bullet(\underline{O}_X)$, the reference to y is omitted from this terminology.

4.3.

Without any hypothesis on the dimension of X, consider a sequence $\underline{L}_1, \dots, \underline{L}_d$ of invertible $\underline{O}_X$-modules and a coherent $\underline{O}_X$-module $\underline{F}$ such that $\dim(\mathrm{Supp}(\underline{F})) \leq d$. Put $L_i = \mathrm{cl}^\bullet(\underline{L}_i)$ $(i = 1, \dots, d)$ and $F = \mathrm{cl}_\bullet(\underline{F})$.

[print 543]

Proposition 4.3.1. With the above notation, for every sequence $\alpha_1, \dots, \alpha_d$ of nonnegative integers with $\sum_{i=1}^d \alpha_i = d$, the coefficient of

$$\binom{n_1}{\alpha_1} \cdots \binom{n_d}{\alpha_d}$$

in the Snapper polynomial $\chi(L_1^{n_1} \cdots L_d^{n_d} F)$ is equal to [530]

$$\langle c_1(L_1)^{\alpha_1} \cdots c_1(L_d)^{\alpha_d}, |\underline{F}| \rangle, \tag{4.3.2}$$

where $|\underline{F}|$ is the class of $\underline{F}$ in $\mathrm{Gr}_d(X)$. Thus, in the ordinary expansion of this polynomial in monomials in the n_i $(i = 1, \dots, d)$, the coefficient of the monomial $n_1^{\alpha_1} \cdots n_d^{\alpha_d}$ is equal to

$$(1/(\alpha_1! \cdots \alpha_d!^{[531]}))\langle c_1(L_1)^{\alpha_1} \cdots c_1(L_d)^{\alpha_d}, |\underline{F}| \rangle.$$

In particular, one immediately obtains the following

Corollary 4.3.3. The coefficient of the monomial $n_1 n_2 \cdots n_d$ in the Snapper polynomial $\chi(L_1^{n_1} \cdots L_d^{n_d} F)$ is equal to[532]

$$\langle c_1(L_1) \cdots c_1(L_d), |\underline{F}| \rangle = (c_1(L_1) \cdots c_1(L_d), |\underline{F}|). \tag{4.3.4}$$

Proof: Put $L_i = 1 + M_i$ for every $i = 1, \dots, d$. The definitions immediately show that the integer (4.3.2) is equal to $\chi(M_1^{\alpha_1} \cdots M_d^{\alpha_d} F)$. Substituting $A = K^\bullet(X)/\operatorname{Filt}^{d+1}(X)$, $I = \operatorname{Filt}^1(X)/\operatorname{Filt}^{d+1}(X)$ in (2.1.4), formula (2.1.5) gives (taking account of the fact that $M_i \in \operatorname{Filt}^1(X)$ for every $i = 1, \dots, d$) the congruence

$$L_1^{n_1} \cdots L_d^{n_d} \equiv \sum_{\substack{\alpha_1, \dots, \alpha_d \geq 0 \\ \alpha_1 + \cdots + \alpha_d \leq d^{[533]}}} \binom{n_1}{\alpha_1} \cdots \binom{n_d}{\alpha_d} M_1^{\alpha_1} \cdots M_d^{\alpha_d} \mod \operatorname{Filt}^{d+1}(X). \tag{4.3.5}$$

Thus the coefficient of $\binom{n_1}{\alpha_1} \cdots \binom{n_d}{\alpha_d}$ in the polynomial $\chi(L_1^{n_1} \cdots L_d^{n_d} F)$ is equal to $\chi(M_1^{\alpha_1} \cdots M_d^{\alpha_d} F)$, Q.E.D.

[print 544]

4.4.

The integer (4.3.4) is sometimes called the intersection number of the invertible $\underline{O}_X$-modules $\underline{L}_1, \dots, \underline{L}_d$ with $\underline{F}$, and is denoted by $(\underline{L}_1, \dots, \underline{L}_d; \underline{F})$ (cf. [3]). Expression (4.3.4) for this number immediately gives the following properties:

1) The integer (4.3.4) is a symmetric, d-linear function of the variables $L_1, \dots, L_d \in \mathrm{Pic}(X)$, zero when $\dim(\mathrm{Supp}(\underline{F})) < d$, and is "additive" in $\underline{F}$, i.e. additive in $\mathrm{cl}_\bullet(\underline{F}) = F$.

2) Using 2.1.2, one immediately concludes (with the notation of 1.1.2):

$$(\underline{L}_1, \dots, \underline{L}_d; \underline{F}) = \sum_{x \in X(d)} \mathrm{long}_{\underline{O}_{X,x}}{}^{[534]}(\underline{F}_x)(\underline{L}_1, \dots, \underline{L}_d; \underline{O}_{Z(x)}). \tag{4.4.1}$$

3) Let Y be a closed subscheme of X and $\underline{G}$ a coherent $\underline{O}_Y$-module such that $\underline{F} = f_*(\underline{G})$, where $f : Y \to X$ is the canonical immersion. Then one obtains

$$(f^*\underline{L}_1, \ldots, f^*\underline{L}_d; \underline{G}) = (\underline{L}_1, \ldots, \underline{L}_d; \underline{F}).$$

More generally, the projection formula gives

$$\chi(f^*(x) \cdot y) = \chi(f_*(f^*(x) \cdot y)) = \chi(x \cdot f_*(y)) \qquad (x \in K^\bullet(X),\ y \in K_\bullet(Y)) \qquad (4.4.2)$$

for every k-morphism $f : Y \to X$ of proper k-schemes.

4) Suppose that, in the morphism $f : Y \to X$ above, X and Y are irreducible of dimension $\leq d$. If f has degree n, then one has [print 545]

$$(f^*\underline{L}_1, \ldots, f^*\underline{L}_d; \underline{O}_Y) = n(\underline{L}_1, \ldots, \underline{L}_d; \underline{O}_X). \qquad (4.4.3)$$

Recall the definition of the degree n of f [3]: it is zero except when $\dim(X) = \dim(Y) = \dim(f(Y))$. In this latter case, put

$$n = (\mathrm{long}_{\underline{O}_{X,x}}(\underline{O}_{Y,y}))/(\mathrm{long}_{\underline{O}_{X,x}}(\underline{O}_{X,x})), \qquad (4.4.4)$$

where x is the generic point of X and y the generic point of Y.

In the proof of (4.4.3), the case $\dim(X) = \dim(Y)$ is the only nontrivial one. If $\dim(f(Y)) < d$, then one has

$$R^i f_*(\underline{O}_Y) = 0 \qquad (4.4.5)$$

for $i > 0$ in a neighborhood of x. But condition (4.4.5) also holds when $\dim(f(Y)) = d$, since the generic fiber then has dimension zero (EGA III 4.2.2 and EGA IV 5.6.6). Then 1.1.2 gives the congruences

$$f_*(\mathrm{cl}_\bullet(\underline{O}_Y)) \equiv \mathrm{cl}_\bullet(f_*(\underline{O}_Y)) \equiv^{[535]} \mathrm{long}_{\underline{O}_{X,x}}(\underline{O}_{Y,y})\, \mathrm{cl}_\bullet(\underline{O}_{X_{\mathrm{red}}})$$

modulo $\mathrm{Filt}_{d-1}(X)$. By (4.4.2), it follows that

$$(f^*\underline{L}_1, \ldots, f^*\underline{L}_d; \underline{O}_Y) = \mathrm{long}_{\underline{O}_{X,x}}(\underline{O}_{Y,y})(\underline{L}_1, \ldots, \underline{L}_d; \underline{O}_{X_{\mathrm{red}}}).$$

On the other hand, 2° gives

$$(\underline{L}_1, \ldots, \underline{L}_d; \underline{O}_X) = \mathrm{long}_{\underline{O}_{X,x}}(\underline{O}_{X,x})(\underline{L}_1, \ldots, \underline{L}_d; \underline{O}_{X_{\mathrm{red}}}),$$

whence the assertion.

[print 546]

4.5

The equivalence relation defined by the intersection product $\langle\ ,\ \rangle$ is called numerical equivalence:

Definition 4.5.1. Two elements x and y of $\mathrm{Gr}^i(X)$ $(i \in Z)$ (resp. of $\mathrm{Gr}_i(X)$) are said to be numerically equivalent when they define the same homomorphism $\mathrm{Gr}_i(X) \longrightarrow \mathbb{Z}$ (resp. $\mathrm{Gr}^i(X) \longrightarrow \mathbb{Z}$). We say that two invertible $\underline{O}_X$-modules $\underline{L}_1$ and $\underline{L}_2$ are numerically equivalent when their first Chern classes $x, y \in \mathrm{Gr}^1(X)$ are numerically equivalent.

Proposition 4.5.2. An invertible $\underline{O}_X$-module $\underline{L}$ is numerically equivalent to zero if and only if

$$\chi(\underline{L}^n \otimes_{\underline{O}_X} \underline{O}_Z) = \chi(\underline{O}_Z) \qquad (4.5.3)$$

for every $n \in \mathbb{Z}$ and every integral closed curve $Z \subset X$.

This follows from 1.1.2 and 4.3.1.

Denote by $\mathrm{Pic}^N(X)$ the subgroup of $\mathrm{Pic}(X)$ consisting of the classes of invertible $\underline{O}_X$-modules numerically equivalent to zero. From 2.4.4 one obtains[536]

Corollary 4.5.3. $\mathrm{Pic}^\tau(X) \subset \mathrm{Pic}^N(X)$.

In Exp. XIII it will be proved that one even has $\mathrm{Pic}^N(X) = \mathrm{Pic}^\tau(X)$.

5. The isomorphism $\mathrm{Pic}(X) \cong \mathrm{Gr}^1(X)$

5.1.

Let X be an arbitrary ringed topos. Then one can define the ring $K^\bullet(X)$ and the filtration $(\mathrm{Filt}^i(X))_{i\in\mathbb{Z}}$ exactly as for schemes. Likewise, define the group $\mathrm{Pic}(X)$ as the group of isomorphism classes of invertible $\underline{O}_X$-modules. Denote by [print 547]

$$i : \mathrm{Pic}(X) \longrightarrow K^\bullet(X) \tag{5.1.1}$$

the map associating to the class $\mathrm{cl}_P(\underline{L}) \in \mathrm{Pic}(X)$ of an invertible $\underline{O}_X$-module $\underline{L}$ the element $\mathrm{cl}^\bullet(\underline{L})$ of $K^\bullet(X)$. This map is clearly a homomorphism from $\mathrm{Pic}(X)$ to the multiplicative group $K^\bullet(X)^*$ of invertible elements of $K^\bullet(X)$. Henceforth, the group law of $\mathrm{Pic}(X)$ will be written multiplicatively.

The map i has a left inverse defined by the "det" functor: when $\underline{E}$ is a locally free $\underline{O}_X$-module of rank $\underline{n}$, one has

$$\det(\underline{E}) = \bigwedge^{\underline{n}} \underline{E}.$$

If

$$0 \longrightarrow \underline{E}' \longrightarrow \underline{E} \longrightarrow \underline{E}'' \longrightarrow 0 \tag{5.1.2}$$

is an exact sequence of locally free $\underline{O}_X$-modules, there is a canonical isomorphism

$$\det(\underline{E}) \xrightarrow{\sim} (\det(\underline{E}')) \otimes_{\underline{O}_X} (\det(\underline{E}''))$$

arising from (6.2.9) and the fact that (5.1.2) is locally split. Thus one obtains a homomorphism of abelian groups

$$\det : K^\bullet(X) \longrightarrow \mathrm{Pic}(X) \tag{5.1.3}$$

that is the required left inverse of i. In particular, i is injective, and henceforth we shall identify $\mathrm{Pic}(X)$ with its image in $K^\bullet(X)^*$.

5.2.

Define another map

$$\delta : \mathrm{Pic}(X) \longrightarrow \mathrm{Filt}^1(X) \tag{5.2.1}$$

[print 548]by putting, for every $L \in \mathrm{Pic}(X)$,

$$\delta(L) = 1 - L^{-1}.$$

Remarks 5.2.2.:

a) Suppose that X is the locally ringed topos of Zariski sheaves on a scheme S. When D is a Cartier divisor ≥ 0 on X and

$$L = \mathrm{cl}^\bullet(\underline{O}_S(D)),$$

one has

$$\theta_S(\delta(L)) = \mathrm{cl}_\bullet(\underline{O}_D) \tag{5.2.3}$$

where $\theta_S : K^\bullet(X) \to K_\bullet(S)$ is the canonical homomorphism of 1.2.3. This follows immediately from the exact sequence

$$0 \longrightarrow \underline{O}_S(-D) \longrightarrow \underline{O}_S \longrightarrow \underline{O}_D \longrightarrow 0.$$

b) Under the hypotheses of 5.1 and 5.2, one has, for every $L \in \mathrm{Pic}(X)$,

$$\delta(L) \equiv L - 1 \quad \mathrm{mod.\,Filt}^2(X), \tag{5.2.4}$$

which justifies the notation $c_1(L)$ introduced below. Indeed, one has

$$(L-1) - (1 - L^{-1}) = (L-1)(1-L^{-1}) \in \mathrm{Filt}^2(X).$$

5.3.

Composing δ with the canonical epimorphism

$$\mathrm{Filt}^1(X) \longrightarrow \mathrm{Gr}^1(X)$$

gives the "first Chern class" on $\mathrm{Pic}(X)$:

$$c_1 : \mathrm{Pic}(X) \longrightarrow \mathrm{Gr}^1(X). \tag{5.3.1}$$

[print 549]

Theorem 5.3.2. The map c_1 (5.3.1) is an injective homomorphism, and the homomorphism

$$\det : \mathrm{Filt}^1(X) \longrightarrow \mathrm{Pic}(X)$$

induces, on passing to the quotient, a homomorphism that is a left inverse of c_1:

$$\alpha : \mathrm{Gr}^1(X) \longrightarrow \mathrm{Pic}(X).$$

If every section n of the constant sheaf $\mathbb{Z}_X$ is bounded (for example, if X is "quasi-compact", or if the final object e of X is a finite sum of connected objects), then c_1 is an isomorphism and α is its inverse.

Proof: c_1 is a homomorphism, since for every $L, L' \in \mathrm{Pic}(X)$ one has the equation

$$(1 - (LL')^{-1}) - (1 - L^{-1}) - (1 - L'^{-1}) = -(1 - L^{-1})(1 - L'^{-1}) \in \mathrm{Filt}^2(X).$$

For the moment, admit that

$$\det(\mathrm{Filt}^2(X)) = \{1\} \in \mathrm{Pic}(X)$$

i.e., that α is well defined. One then has

$$\alpha \circ c_1 = \mathrm{id}_{\mathrm{Pic}(X)},$$

since, by 5.2.2 b), one has

$$\det(1 - L^{-1}) = (\det(L - 1)) = L$$

for every $L \in \mathrm{Pic}(X)$. Next one shows that

$$c_1 \circ \alpha = \mathrm{id}_{\mathrm{Gr}^1(X)}$$

when every section $\underline{n}$ of $\mathbb{Z}_X$ is bounded. Since $\mathrm{Filt}^1(X)$ is the abelian group generated by the elements $\mathrm{cl}^\bullet(\underline{E}) - \underline{n}$, where $\underline{E}$ is a [print 550]locally free $\underline{O}_X$-module of rank $\underline{n}$, it suffices, taking account of 5.2.2 b), to show that

$$\det(\mathrm{cl}^\bullet(\underline{E})) - \mathrm{cl}^\bullet(\underline{E}) + \underline{n} - 1 \in \mathrm{Filt}^2(X)^{[537]}. \tag{5.3.3}$$

Since $\underline{n}$ is bounded, the final object $\underline{e}$ of X decomposes into a finite disjoint direct sum $\coprod e_i$ of subobjects in such a way that $\underline{n}$ is constant on each subtopos $X/\underline{e}_i$ $(i = 1, \ldots, p)$. Since the functor $Y \rightsquigarrow K^\bullet(X/Y)$ $(Y \in \mathrm{Ob}(X))$ takes finite direct sums to finite direct products, one may suppose that $\underline{n}$ is constant on X, say $\underline{n} = n \in \mathbb{Z}$. Putting $E = \mathrm{cl}^\bullet(\underline{E})$, one finds (cf. Exposé V):

$$\det(\underline{E}) = \lambda^n(E) = \gamma^n((E - n) + 1) = \sum_{0 \le i \le n} \gamma^i(E - n),$$

This proves (5.3.3), whose left-hand side is equal to

$$\sum_{2 \le i \le n} \gamma^i(E - n).$$

It remains to prove the following

Lemma 5.3.4. For every ringed topos X and every $x \in \mathrm{Filt}^2(X)$, one has

$$\det(x) = 1.$$

Proof: Let $\epsilon : K^\bullet(X) \to \Gamma(\mathbb{Z})$ be the augmentation homomorphism. Then (6.2.8) and (6.5.3) below give[538], by linearity, the formula

$$\det(xy) = (\det(x))^{\epsilon(y)}(\det(y))^{\epsilon(x)} \qquad (x, y \in K^\bullet(X)). \tag{5.3.5}$$

[print 551]It therefore suffices to show that, for every $k > 1$, one has

$$\det(\gamma^k(E - \underline{n})) = 1,$$

where, as above, $E = \mathrm{cl}^\bullet(\underline{E})$, $\underline{n} = \epsilon(E)$. Now one has

$$\gamma^k(x) = \lambda^k(x + k - 1)$$

for every $x \in K^\bullet(X)$, so that

$$\det(\gamma^k(E - \underline{n})) = \det(\lambda^k(E - \underline{n} + k - 1)) = \det\left(\sum_{p+q=k} \binom{-\underline{n} + k - 1}{p} \lambda^q(E)^{[539]}\right).$$

But (6.2.10) and (6.5.3) give

$$\det(\lambda^q(E)) = (\det(E))^{\binom{\underline{n}-1}{q-1}}$$

for every $q > 0$, whence

$$\det(\gamma^k(E - \underline{n})) = (\det(E))^{\underline{r}},$$

with

$$\underline{r} = \sum_{\substack{p+q=k \\ q \ge 1}} \binom{-\underline{n} + k - 1}{p} \binom{\underline{n} - 1}{q - 1}.$$

Now $\underline{r}$ is a section of $\mathbb{Z}$, everywhere equal to the coefficient of t^{k-1} in $(1 + t)^{k-2}$, and is therefore zero. The result follows.

6. Appendix: Calculation of Determinants of Locally Free Sheaves

6.1.

First consider an arbitrary ringed topos $X = (C, \underline{O})$. For every positive section $\underline{n} \in \Gamma(\mathbb{Z}_X)$ of the constant sheaf $\mathbb{Z}$ and every $\underline{O}$-module $M \in \mathrm{Mod}_{\underline{O}}(X)$, it is clear how to define the $\underline{O}$-modules [print 552]

$$M^{\underline{n}}, \qquad \overset{\underline{n}}{\bigotimes} M, \qquad \overset{\underline{n}}{\bigwedge} M, \quad \text{etc.} \tag{6.1.1}$$

The functors $(\ )^{\underline{n}}$, $\overset{\underline{n}}{\bigotimes}$ and $\overset{\underline{n}}{\bigwedge}$ take free (resp. locally free) modules to free (resp. locally free) modules.

More generally, consider sections $\underline{n}_1, \ldots, n_r$ of $\mathbb{Z}_X$ and a multifunctor (covariant, for definiteness)

$$T : \underline{\mathrm{Mod}}^{n_1}(\underline{O}) \times \cdots \times \underline{\mathrm{Mod}}^{n_r}(\underline{O}) \longrightarrow \underline{\mathrm{Mod}}(\underline{O})$$

in r variables, where $\mathrm{Mod}^{\underline{n}}_{\underline{O}}(X)$ [540] denotes the category fibered over X of locally free $\underline{O}$-modules of rank $\underline{n}$. Put

$$M = T(\underline{O}^{\underline{n}_1}, \ldots, \underline{O}^{\underline{n}_r})$$

and

$$G = \mathrm{G}\ell(\underline{n}_1) \times \cdots \times \mathrm{G}\ell(\underline{n}_r), \quad \text{where } \mathrm{G}\ell(\underline{n}_i) = \underline{\mathrm{Aut}}_{\underline{O}}(\underline{O}^{n_i}).$$

It follows immediately that T defines a representation

$$\alpha_T : G \longrightarrow \underline{\mathrm{Aut}}_{\underline{O}}(M). \tag{6.1.2}$$

It is clear that, when M is itself locally free of rank $\underline{m}$, for every composite functor

$$T'T, \text{ where } T' : \underline{\mathrm{Mod}}^{\underline{m}}(\underline{O}) \longrightarrow \underline{\mathrm{Mod}}(\underline{O}),$$

one has

$$\alpha_{T'T} = \alpha_{T'}\, \alpha_T. \tag{6.1.3}$$

Conversely, given the representation $\alpha = \alpha_T : G \to \underline{\mathrm{Aut}}_{\underline{O}}(M)$, one recovers $T(\underline{E}_1, \ldots, \underline{E}_r)$ (for every sequence [print 553] $\underline{E}_1, \ldots, \underline{E}_r$ of locally free $\underline{O}$-modules of ranks $\underline{n}_1, \ldots, \underline{n}_r$) by gluing the free $\underline{O}$-module $T(\underline{O}^{\underline{n}_1}, \ldots, \underline{O}^{\underline{n}_r})$ on local trivializing charts by means of the representation α. More precisely, the sequence $\underline{E}_1, \ldots, \underline{E}_r$ corresponds to a right G-torsor $P = P_{\underline{E}_1, \ldots, \underline{E}_r}$ such that there is a canonical isomorphism

$$\underline{E}_i \xrightarrow{\sim} P \overset{G}{\wedge} \underline{O}^{\underline{n}_i} \qquad (i = 1, \ldots, r), \tag{6.1.3}$$

where $\overset{G}{\wedge}$ denotes the contracted product, and G acts on $\underline{O}^{\underline{n}_i}$ through the i-th projection $G \to \mathrm{Gl}(\underline{n}_i)$; then every representation

$$\alpha : G \longrightarrow \underline{\mathrm{Aut}}_{\underline{O}}(M)$$

makes G act on M by [541] $\underline{O}$-automorphisms, and thus defines a multifunctor $T = T_\alpha$ such that

$$T_\alpha(\underline{E}_1, \ldots, \underline{E}_r) = P_{\underline{E}_1, \ldots, \underline{E}_r} \overset{G}{\wedge} M. \tag{6.1.4}$$

Moreover, for every composite representation $\alpha = \alpha_1 \circ \alpha_2$, one finds

$$T_\alpha = T_{\alpha_1} \circ T_{\alpha_2}. \tag{6.1.5}$$

6.2.

For every locally free $\underline{O}$-module $\underline{E}$ of rank $\underline{n}$, we shall denote by $\det(\underline{E})$ the invertible $\underline{O}$-module $\overset{\underline{n}}{\bigwedge}\underline{E}$. Keeping the notation of 6.1, the functor $\det(T(\underline{E}_1,\dots,\underline{E}_r))$ is determined, up to canonical isomorphism, by the corresponding representation

$$\det\circ\alpha_T : G \longrightarrow G_m = \mathrm{Gl}(1). \tag{6.2.1}$$

Suppose now that X is the ringed topos of sheaves on the ringed site $(\mathrm{Sch}\,/S)$ endowed with the étale topology. Then one can calculate $\det(T(\underline{E}_1,\dots,\underline{E}_r))$, or equivalently the representation $\det\circ\alpha_T$, by means of the following [print 554]

Proposition 6.2.2. For every positive section $\underline{n}$ of $\mathbb{Z}$, the étale sheaf of groups $\underline{\mathrm{Der}}(\mathrm{Gl}(\underline{n}))$ of commutators of $\mathrm{Gl}(\underline{n})$ is represented by[542] $\mathrm{Sl}(\underline{n}) = \mathrm{Ker}(\det : \mathrm{Gl}(\underline{n}) \to G_m)$.

Corollary 6.2.3. Every representation

$$\alpha : G = \mathrm{Gl}(\underline{n}_1)\times\cdots\times\mathrm{Gl}(\underline{n}_r) \longrightarrow G_m \tag{6.2.4}$$

is of the form

$$\alpha(Y)(s) = (\det(s_1))^{\underline{m}_1}\cdots(\det(s_r))^{\underline{m}_r}, \tag{6.2.5}$$

where $Y \in \mathrm{Ob}(\mathrm{Sch}\,/S)$, $s = (s_1,\dots,s_r) \in G(Y)$ and $\underline{m}_1,\dots,\underline{m}_r$ are sections of $\mathbb{Z}$ uniquely determined by α.

Corollary 6.2.6. With the notation of 6.1. one has

$$\det(T_\alpha(\underline{E}_1,\dots,\underline{E}_r)) = \det(\underline{E}_1)^{\otimes\underline{m}_1}\otimes\cdots\otimes\det(\underline{E}_r)^{\otimes\underline{m}_r},$$

where $\underline{m}_1,\dots,\underline{m}_r$ are sections of $\mathbb{Z}$ uniquely determined by α.

Remark 6.2.7. One can determine the exponents $\underline{m}_1,\dots,\underline{m}_r$ in (6.2.5) and (6.2.6) by successively substituting in (6.2.5) $s_i = \lambda.\ \mathrm{id}\,.(\lambda \in G_m(Y))$, $s_j = \mathrm{id}_{\mathrm{Gl}(\underline{n}_j)(Y)}$, for $j \neq i$ $(i,j = 1,\dots,r)$, whence

$$\det(s_i) = \lambda^{\underline{n}_i} \quad\text{and}\quad \alpha(Y)(s) = \lambda^{\underline{n}_i\underline{m}_i}.$$

For example, one thus deduces from (6.2.6) [543]

$$\det(\underline{E}_1\otimes\underline{E}_2) = (\det(\underline{E}_1))^{\otimes\underline{n}_2}\otimes(\det(\underline{E}_2))^{\otimes\underline{n}_1}, \tag{6.2.8}$$

$$\det(\underline{E}_1\oplus\underline{E}_2) = (\det(\underline{E}_1))\otimes(\det(\underline{E}_2)) \tag{6.2.9}$$

and


$$\det(\overset{\underline{m}}{\bigwedge}\underline{E}_1) = (\det(\underline{E}_1))^{\binom{\underline{n}_1-1}{\underline{m}-1}} \tag{6.2.10}$$

etc.

[print 555]

6.3.

Let us first show how 6.2.3 follows from 6.2.2. For this, it plainly suffices to prove (6.2.5) for the $s_{(i)} = (1,\dots,s_i,\dots,1)$ $(i = 1,\dots,r)$, so one may suppose $r = 1$, $G = \mathrm{G}\ell(\underline{n}_1)$. By 6.2.2, α factors through $\mathrm{G}\ell(\underline{n}_1)/\mathrm{Sl}(\underline{n}_1)$, i.e., there is a commutative diagram



where j is the canonical epimorphism and β is a representation of $\mathrm{G}\ell(\underline{n}_1)/\mathrm{Sl}(\underline{n}_1)$ in G_m. But the homomorphism

$$\det : \mathrm{G}\ell(\underline{n}_i) \longrightarrow G_m$$

has a section, so that the monomorphism

$$\mathrm{G}\ell(\underline{n}_i)/\mathrm{S}\ell(\underline{n}_i) \longrightarrow G_m$$

induced by det is an isomorphism. On the other hand, it is known (SGA 3 VIII 1.3) that every representation of G_m in itself is of the form

$$s \rightsquigarrow s^{\underline{m}},$$

where $\underline{m}$ is a section of $\mathbb{Z}$, whence the assertion.

Corollary 6.2.6 is a simple reformulation of 6.2.3. Note that 6.2.3 also follows immediately from SGA 3 XXII 6.2.1, but we wish to give an elementary proof here via 6.2.2.[544]

[print 556]

6.4. Proof of 6.2.2.

It suffices to show that, for every $Y \in \mathrm{Ob}(\mathrm{Sch}/S)$, $y \in Y$ and $s \in \mathrm{S}\ell(\underline{n})(Y)$, one can find an "étale neighborhood" $Y' \to Y$ of y such that the restriction of s to Y' is a product of commutators of $\mathrm{G}\ell(\underline{n})(Y')$. Since the question is local, one may suppose that Y is affine with ring A, and that $\underline{n}$ is constant with value n on Y. Let $\underline{p}$ be the prime ideal of y and $(s_{ij})(i,j=1,\ldots,n)$ the matrix of s. For every $\lambda \in A$ and $h,k = 1,\ldots,n$; $h \neq k$, we shall denote by $c_{h,k,\lambda}$ the matrix (c_{ij}) $(i,j=1,\ldots,n)$ such that $c_{hk} = \lambda$ and $c_{ij} = \delta_{ij}$ for $(i,j) \neq (h,k)$ (δ_{ij} = the Kronecker symbol). For every matrix $\underline{a} = (a_{ij})(i,j=1,\ldots,n)$, one obtains $c_{h,k,\lambda}\underline{a}$ by adding to the k-th column of $\underline{a}$ λ times the h-th column. Similarly, one obtains the matrix $\underline{a}c_{h,k,\lambda}$ by adding to the h-th row of $\underline{a}$ λ times the k-th row. In particular, one has

$$\det(c_{h,k,\lambda}) = 1,$$

and one sees at once that

$$c_{h,k,\lambda}c_{h,k,\mu} = c_{h,k,\lambda+\mu} \tag{6.3.1}$$

for every $\lambda,\mu \in A$, so that one obtains a representation

$$c_{h,k} : G_a \longrightarrow \mathrm{S}\ell(n) \qquad (h,k=1,\ldots,n\ ;\ h \neq k).$$

Unless expressly stated otherwise, in what follows we shall suppose that the λ occurring in $c_{h,k,\lambda}$ $(\lambda \in A\ ;\ h,k=1,\ldots,n\ ;\ h \neq k)$ is invertible. The proof then proceeds in two steps. We shall first show that s is a product of matrices $c_{h,k,\lambda}$ in a sufficiently small Zariski neighborhood of y. We shall then show that the matrices $c_{h,k,\lambda}$ are products of commutators of $\mathrm{G}\ell(n)$ in a sufficiently small étale neighborhood of y.

[print 557]

For the first step, one must transform s into the identity matrix $\underline{1} = (\delta_{ij})$ in a Zariski neighborhood of y by multiplying s on the left and on the right by suitable matrices $c_{h,k,\lambda}$. Let us first show that one can thereby make $s_{11} = 1$. If $s_{11} \neq 1$ and $s_{21} \in \underline{p}$, we seek a matrix $c = c_{2,h,\lambda}$[545] such that the product $s' = sc = (s'_{ij})$ has $s'_{21} \notin \underline{p}$. Since $\det(s) = 1 \notin \underline{p}$, there is at least one index i such that $s_{i1} \notin \underline{p}$. Then[546] one has $s_{21} + s_{i1} \notin \underline{p}$, and $c_{2,i,1}$[547] answers the question. One may therefore suppose $s_{21} \notin \underline{p}$ and, after localizing, suppose that s_{21} is invertible. If $s_{11} - 1 \in \underline{p}$, then in $s' = (s'_{ij}) = sc_{1,2,1}$ one has $s'_{11} - 1 = (s_{11} + s_{21}) - 1 \notin \underline{p}$. After localizing, one may then

suppose that $s_{11} - 1$ is invertible. Then $\lambda = (1 - s_{11})/s_{21}$ is invertible and one finds $s'_{11} = 1$ in $s' = (s'_{ij}) = sc_{1,2,\lambda}$.

Supposing $s_{11} = 1$, one can then put s in the form

$$s_{11} = 1, \qquad s_{1i} = s_{i1} = 0 \quad (i = 2, \ldots, n)$$

in a Zariski neighborhood, of y by multiplying s on the left (resp. on the right) by matrices $c_{1,i,\lambda}$ [548] (resp. $c_{i,1,\lambda}$). Next one makes $s_{22} = 1$, etc. By induction, one can then transform s in a Zariski neighborhood U of y into a matrix $s' = (s'_{ij})$ with $s'_{ij} = \delta_{ij}$ for $i < n$ or $j < n$. But by the very construction one then has $s' = c_1 s c_2$ on $\mathcal{U}$, where c_1 and c_2 are products of matrices $c_{h,k,\lambda}$. Thus one has

$$\det(s') = \det(s) = 1,$$

so that $s'_{nn} = 1$, $s' = \underline{\underline{1}}$, and the result follows.

We therefore suppose that s is already, on Y, a product of matrices $c_{h,k,\lambda}$. It remains to see that these are products of commutators in an étale neighborhood of y. First, one sees that any two matrices [print 558]$c_{h,k,\lambda}$ and $c_{h,k,\mu}$ are conjugate; in other words, there is an invertible matrix $a = (a_{ij})$ $(i, j = 1, \ldots, n)$ such that

$$a c_{h,k,\lambda} = c_{h,k,\mu} a.$$

Indeed, the matrix (a_{ij}) with $a_{hh} = 1/\mu$, $a_{kk} = 1/\lambda$, and otherwise $a_{ij} = \delta_{ij}$, answers the question (recall that λ and μ are invertible by hypothesis and $h \neq k$). Thus, for any homomorphism β from $\mathrm{G}\ell(n)(U)$ to a sheaf of abelian groups H, one has

$$\beta(c_{h,k,\lambda}) = \beta(c_{h,k,\mu}).$$

If $\lambda + \mu$ is invertible, this and (6.3.1) imply that $\beta(c_{h,k,\lambda}) = 0$ for every h, β, so that $c_{h,k,\lambda}$ is a product of commutators. If the residue field k of y is not the prime field $\underline{F}_2 = \{0, 1\}$, then for every invertible $\lambda \in A$ one can find a Zariski neighborhood $U = \mathrm{Spec}(B)$ of y and an invertible element $\mu \in B$ such that $\lambda + \mu \in B$ is invertible. It follows that $c_{h,k,\lambda}$ is a product of commutators on U.

If $k = \underline{F}_2$, then $V = \mathrm{Spec}(A[T]/(T^2 + T + 1))$ is an étale neighborhood of y whose residue field differs from $\underline{F}_2$. The result follows.

6.5. Remarks

6.5.1. When none of the residue fields of S is isomorphic to $\underline{F}_2$, the preceding proof shows that 6.2.2. remains valid for the ringed topos of Zariski sheaves on S. On the other hand, already for $S = \mathrm{Spec}(\underline{F}_2)$, $\mathrm{S}\ell(2, \underline{F}_2) = \mathrm{G}\ell(2, \underline{F}_2)$ differs from the commutator group of $\mathrm{G}\ell(2, \underline{F}_2)$. The order of the latter is 3, whereas the order of $\mathrm{S}\ell(2, \underline{F}_2)$ is 6 (cf. [1]).

6.5.2. On an arbitrary locally ringed topos X, it is not true in general that every representation $G_m \to G_m$ is defined by a [print 559]power. For example, on the topos (Ens) ringed by the field $\mathbb{Q}$, the representation $G_m \to G_m$ defined by

$$r \rightsquigarrow |r| \qquad (r \in \mathbb{Q}^*)$$

is not a power. In particular, 6.2.3 does not remain valid for an arbitrary locally ringed topos X and an arbitrary representation $\alpha : G \to G_m$.

6.5.3. On the other hand, 6.2.3 remains valid for the representations associated with formulas (6.2.8), (6.2.9) and (6.2.10) on an arbitrary ringed topos $X = (C, \underline{O})$, because these representations are "algebraic" in the following sense:

Let $\underline{e}$ be the final object of X, put $S = \mathrm{Spec}(\underline{O}(\underline{e}))$, and consider the functor

$$u : C \longrightarrow (\mathrm{Sch}\,/S)$$

which assigns to each $Y \in \mathrm{Ob}(C)$ the affine scheme $\mathrm{Spec}(\underline{O}(Y))$ over S. Then, for every positive integer n, the presheaf $\mathrm{G}\ell(n)_X$ on C is the composite with u of the corresponding presheaf $\mathrm{G}\ell(n)_S$ on $(\mathrm{Sch}\,/S)$. A representation on X

$$\alpha : \mathrm{G}\ell(n) \longrightarrow G_m \tag{6.5.4}$$

is called algebraic if it arises via u from a representation

$$\alpha_S : \mathrm{G}\ell(n)_S \longrightarrow G_{m,S} \tag{6.5.5}$$

of presheaves of groups on $(\mathrm{Sch}\,/S)$, i.e., if one has

$$\alpha(s) = \alpha_S(s)$$

for every $Y \in \mathrm{Ob}(C)$, $s \in \mathrm{G}\ell(n)(Y) = \mathrm{G}\ell(n)_S(u(Y))$.

The representation α in (6.5.4) is, of course, completely determined by α_S. It follows immediately that (6.2.3) remains valid in an arbitrary ringed topos when α is algebraic. In particular, formulas (6.2.8), (6.2.9) and (6.2.10) therefore remain valid on an arbitrary ringed topos.

[print 560]

7. Appendix: Specialization in Intersection Theory

by A. Grothendieck

Lemma 7.1. Let $i : Y \to X$ be a regular closed immersion (VII 1.4), of bounded codimension, so that i is a perfect morphism (VII 1.9), and one can therefore define the morphism (IV 2.12)

$$i^* : K_\bullet(X) \longrightarrow K_\bullet(Y).$$

Let, on the other hand, $\mathbb{N}$ be the conormal sheaf of i, which is a locally free sheaf of finite type on Y (VII 1.6), so that one can form

$$\lambda_{-1}(\mathbb{N}) \in K^\bullet(Y).$$

Consider also the direct-image homomorphism (IV 2.11)

$$i_* : K_\bullet(Y) \longrightarrow K_\bullet(X).$$

Then one has the formula

$$i^* i_*(x) = x\lambda_{-1}(\mathbb{N}) \qquad \text{for } x \in K_\bullet(Y),$$

where the product on the right is taken with respect to the $K^\bullet(Y)$-module structure on $K_\bullet(Y)$ defined in (IV 2.10).

The proof is, essentially without change, the same as that of the same formula, written for $K^\bullet(X)$, $K^\bullet(Y)$, established in (VII 2.7).

Corollary 7.2. Let $f : X \to S$ be a flat morphism, with S a trait, i.e., the spectrum of a discrete valuation ring A, s the closed point of S, t its generic point, and $i : X_s \to X$ the canonical inclusion. Then i has finite Tor-dimension, so (IV 2.12) one can define

$$i^* : K_\bullet(X) \longrightarrow K_\bullet(X_s),$$

[print 561]on the other hand, (IV 2.11) one has a homomorphism

$$i_* : K_\bullet(X_s) \longrightarrow K_\bullet(X).$$

Then, the composite

$$i^* i_* : K_\bullet(X_s) \longrightarrow K_\bullet(X) \longrightarrow K_\bullet(X_s)$$

is zero.

Indeed, $i : X_s \to X$ is a regular immersion of codimension 1 whose conormal sheaf $\mathbb{N}$ is free of rank 1, since X_s can be described by a global equation $f^*(g) = 0$ (where g is a uniformizer of A). Thus one has $\lambda_{-1}(\mathbb{N}) = 0$, and the conclusion follows from 7.1.

Throughout what follows, we shall suppose that X is noetherian and flat over S. Let us now apply the exact homotopy sequence (IX 1.1) to the following situation: X, with the open subscheme X_t and complementary closed subscheme X_s:

$$K_\bullet(X_s) \xrightarrow{i_*} K_\bullet(X) \xrightarrow{j^*} K_\bullet(X_t) \longrightarrow 0,$$

where $j : X_t \to X$ is the inclusion, which is an open immersion. Using 7.2, one obtains:

Proposition 7.3. With the notation of 7.2, X being assumed noetherian, there is a unique group homomorphism

$$\sigma_\bullet : K_\bullet(X_t) \longrightarrow K_\bullet(X_s) \tag{7.3.1}$$

(called the "specialization homomorphism") making the diagram commutative

$$\begin{array}{ccc} K_\bullet(X) & \xrightarrow{j^*} & K_\bullet(X_t) \\ {\scriptstyle i^*}\downarrow & \swarrow{\scriptstyle \sigma_\bullet} & \\ K_\bullet(X_s) & & \end{array} \quad . \tag{7.3.2}$$

[print 562]

When the two sides of (7.3.1) are regarded as modules over $K^\bullet(X)$ by means of the ring homomorphisms $K^\bullet(X) \xrightarrow{j^*} K^\bullet(X_t)$ and $K^\bullet(X) \xrightarrow{i^*} K^\bullet(X_s)$ (IV 2.7 b) and the module structures on $K_\bullet(X_t)$ and $K_\bullet(X_s)$ over $K^\bullet(X_t)$ and $K^\bullet(X_s)$ respectively (IV 2.10), the homomorphism $\sigma_\bullet$ is a homomorphism of $K^\bullet(X)$-modules.

The last assertion follows from the fact that $i^* : K_\bullet(X) \to K_\bullet(X_s)$ is a homomorphism of $K^\bullet(X)$-modules, as is the epimorphism $j^* : K_\bullet(X) \longrightarrow K_\bullet(X_t)$, by (IV 2.12.2).

We shall review other properties of the specialization morphism.

7.4. Compatibility with filtrations.

Endow the $K_\bullet$ with the filtration $\mathrm{Fil}_\bullet K_\bullet$ by the dimension of supports (1.1.1). Then, when f is of finite type, more generally when X is catenary, the specialization homomorphism is compatible with these filtrations, and consequently induces a homomorphism (also called the specialization homomorphism):

$$\sigma_\bullet : \mathrm{Gr}_\bullet(X_t) \longrightarrow \mathrm{Gr}_\bullet(X_s). \tag{7.4.1}$$

To see this, by 1.1.4 one is reduced[549] to proving that, when Y_t is an integral subscheme of dimension d of X_t, $\sigma_\bullet(\mathrm{cl}_\bullet(Y_t))$ has filtration $\leq d$. Now let Y be the schematic closure of Y_t in X,

so that Y is flat over S (EGA IV 2.8.1). It follows readily that $i^*(\mathrm{cl}_\bullet(Y)) = \mathrm{cl}_\bullet(Y_s)$, where the two $\mathrm{cl}_\bullet$ are taken in $K_\bullet(X)$ and $K_\bullet(X_s)$ respectively, whence $\sigma_\bullet(\mathrm{cl}_\bullet(Y_t)) = \mathrm{cl}_\bullet(Y_s)$. One checks readily (using the fact that X is catenary, hence Y is catenary) that $\dim Y_s \leq \dim Y_t$; when f is of finite type, this follows at once, by reduction to the case of integral X, from EGA IV 7.1.13. This gives the asserted conclusion.

[print 563]

7.5. "Conservation of number".

Suppose now that f is proper, so that X_t and X_s are proper over $k(t)$ and $k(s)$ respectively, and one has the homomorphisms of 2.3:

$$\chi_t : K_\bullet(X_t) \longrightarrow \mathbb{Z}, \qquad \chi_s : K_\bullet(X_s) \longrightarrow \mathbb{Z}.$$

Then one has the relation

$$\chi_s(\sigma_\bullet(x)) = \chi_t(x) \qquad \text{for } x \in K_\bullet(X_t) \tag{7.5.1}$$

(the "conservation-of-number theorem"); in other words, the diagram

$$\begin{array}{ccc} K_\bullet(X_t) & \xrightarrow{\sigma_\bullet} & K_\bullet(X_s) \\ & {\scriptstyle \chi_t} \searrow & \downarrow {\scriptstyle \chi_s} \\ & & \mathbb{Z}. \end{array} \tag{7.5.2}$$

commutes. To prove (7.5.1), one is again reduced to the case where $x = \mathrm{cl}_\bullet(Y_t)$, with Y_t an integral closed subscheme of X_t, and, with the notation of the proof of 7.4, one is reduced to proving that

$$\chi(Y_s) = \chi(Y_t),$$

which is well known (EGA III 7.9.4, or Exposé III 5.7).

7.6.

It is immediate that

$$\sigma_\bullet(\mathrm{cl}_\bullet(\underline{O}_{X_t})) = \mathrm{cl}_\bullet(\underline{O}_{X_s}), \tag{7.6.1}$$

which, taking account of the $K^\bullet(X)$-linearity of $\sigma_\bullet$, implies that the following diagram is commutative:

$$\begin{array}{ccccc} & & K^\bullet(X) & & \\ & {\scriptstyle j^*} \swarrow & & \searrow {\scriptstyle i^*} & \\ K^\bullet(X_t) & & & & K^\bullet(X_s) \\ \downarrow & & & & \downarrow \\ K_\bullet(X_t) & & \xrightarrow{\sigma_\bullet} & & K_\bullet(X_s), \end{array} \tag{7.6.2}$$

[print 564]where the vertical arrows are the canonical homomorphisms $K^\bullet \to K_\bullet$ (IV 2.5.1).

7.7. Specialization for $K^\bullet$ and $\mathrm{Gr}^\bullet$.

When X_t and X_s are regular (for example, when f is smooth), the two preceding vertical arrows are isomorphisms (IV 2.5), and the specialization homomorphism can then be interpreted as a homomorphism

$$\sigma^\bullet : K^\bullet(X_t) \longrightarrow K^\bullet(X_s), \tag{7.7.1}$$

which can also be described as the unique homomorphism making the following diagram commutative

$$\begin{array}{ccc} K^\bullet(X) & \xrightarrow{j^*} & K^\bullet(X_t) \\ {\scriptstyle i^*}\downarrow & \swarrow_{\sigma^\bullet} & \\ K^\bullet(X_s) & & \end{array} \quad . \tag{7.7.2}$$

Since, in this last diagram, i^* and j^* are ring homomorphisms (IV 2.7 b), with j^* surjective, it follows that the specialization homomorphism (7.7.1) is a ring homomorphism. For the same reason, if X is separated, the homomorphism (7.7.1) is even a homomorphism of λ-rings. To make this precise, one need only note that a regular separated noetherian scheme is divisorial (II 2.2.7.1), so its $K^\bullet$ is identified with its "naive" $K^\bullet$ (IV 2.9 (ii)), which is naturally endowed with a λ-ring structure (VI 3.2). On the other hand, the regularity of X_t and X_s implies that of X (EGA 0_{IV} 17.1.8), and the homomorphisms $i^* : K^\bullet(X) \to K^\bullet(X_s)$ and $j^* : K^\bullet(X) \to K^\bullet(X_t)$ are λ-homomorphisms, so the same is true of (7.7.1), which is obtained from i^* by passage to the quotient.

[print 565]

Note also that, since X is regular and hence normal, its connected components coincide with its irreducible components, which dominate S, so that the homomorphism $H^0(X, \mathbb{Z}) \to H^0(X_t, Z)$ induced by the inclusion $X_t \to X$ is bijective. Consequently the homomorphism $H^0(X, \mathbb{Z}) \to H^0(X_s, \mathbb{Z})$ induced by the inclusion $X_s \to X$ can also be interpreted as a homomorphism

$$H^0(X_t, \mathbb{Z}) \longrightarrow H^0(X_s, \mathbb{Z}), \tag{7.7.3}$$

which again deserves the name "specialization homomorphism". One then checks immediately from the definitions that the homomorphism (7.7.1) is compatible with the augmented algebra structures over $H^0(X_t, \mathbb{Z})$ and $H^0(X_s, \mathbb{Z})$ respectively on the two sides, defined in (IV 2.8), taking account of the preceding homomorphism (7.7.3).

Still supposing that X is separated, the preceding discussion implies in particular that (7.7.1) induces a homomorphism of graded rings

$$\sigma^\bullet : \mathrm{Gr}^\bullet(X_t) \longrightarrow \mathrm{Gr}^\bullet(X_s), \tag{7.7.4}$$

and, using the notion of Chern class of (V 6), one obtains a commutative diagram

$$\begin{array}{ccc} K^\bullet(X_t) & \xrightarrow{c_{X_t}} & \mathrm{Ch}\ \mathrm{Gr}^\bullet(X_t) \\ \downarrow & & \downarrow \\ K^\bullet(X_s) & \xrightarrow{c_{X_s}} & \mathrm{Ch}\ \mathrm{Gr}^\bullet(X_s). \end{array}$$

7.8. Specialization for Pic.

Retain the hypotheses of 7.7: X separated, X_t and X_s regular. Taking account of the canonical isomorphism (5.3.2):

$$\mathrm{Gr}^1 \simeq \mathrm{Pic},$$

the homomorphism (7.7.4) therefore gives a canonical homomorphism

$$\mathrm{Pic}(X_t) \longrightarrow \mathrm{Pic}(X_s). \tag{7.8.1}$$

[print 566]An equivalent description of this last homomorphism is as follows: the hypothesis that X is regular implies that these local rings are factorial (by the Auslander–Buchsbaum theorem, EGA IV 21.11.1), and hence that every invertible sheaf $\underline{L}_t$ on X_t extends to an invertible sheaf $\underline{L}$ on X (EGA IV 21.6.11). One then checks at once that the image under (7.8.1) of $\mathrm{cl}(\underline{L}_t) \in \mathrm{Pic}(X_t)$ is $\mathrm{cl}(\underline{L}_s) \in \mathrm{Pic}(X_s)$, where, of course, one puts $\underline{L}_s = \underline{L}|X_s$.

In fact, using only the hypothesis that the local rings of X are factorial and that X_s is locally integral (without hypotheses of regularity or separatedness), one shows that $\mathrm{cl}(\underline{L}_s)$ does not depend on the chosen extension $\underline{L}$ of $\underline{L}_t$ to an invertible Module on X, so that one obtains a homomorphism (7.8.1) directly in this way. To verify the asserted independence, one is reduced to proving that, if $\underline{L}_t \simeq \underline{O}_{X_t}$, then $\underline{L}_s \simeq \underline{O}_{X_s}$. Now the hypothesis means that $\underline{L} \simeq \underline{O}_X(D)$, with D a divisor on X having support $\subset X_s$ (cf. EGA IV 21.2.11), and one reduces at once to the case where X_s is integral, hence $D = nX_s$; but then $D = \mathrm{div}(g^n)$ (g a uniformizer of A), so $\underline{O}_X(D) \simeq \underline{O}_X$ and a fortiori $\underline{L}_s \simeq \underline{O}_{X_s}$.

7.9. Specialization and numerical equivalence.

Suppose that f is proper, and X_s and X_t are regular, and consider the two specialization maps (7.4.1) and (7.7.4):

$$\begin{aligned} \sigma_\bullet &: \mathrm{Gr}_\bullet(X_t) \longrightarrow \mathrm{Gr}_\bullet(X_s) \\ \sigma^\bullet &: \mathrm{Gr}^\bullet(X_t) \longrightarrow \mathrm{Gr}^\bullet(X_s). \end{aligned} \tag{7.9.1}$$

Since the homomorphism

$$\sigma : K^\bullet(X_t) \longrightarrow K^\bullet(X_s) \tag{7.9.2}$$

[print 567]from which they are obtained is compatible with the ring structures, using (7.5.1) one finds that

$$\langle \sigma_\bullet(x_t), \sigma^\bullet(y_t) \rangle = \langle x_t, y_t \rangle \qquad \text{for } x_t \in \mathrm{Gr}_\bullet(X_t) \quad y_t \in \mathrm{Gr}^\bullet(X_t), \tag{7.9.3}$$

with the pairings $\langle x, y \rangle$ of 4.1 between $\mathrm{Gr}_\bullet$ and $\mathrm{Gr}^\bullet$. It follows formally that if x is an element of $\mathrm{Gr}_\bullet(X_t)$ (resp. of $\mathrm{Gr}^\bullet(X_t)$) such that $\sigma_\bullet(x)$ (resp. $\sigma^\bullet(x)$) is numerically equivalent to zero (4.5.1), then the same is true of x. The converse is true when (7.9.2) (or, equivalently, either of the homomorphisms (7.9.1) obtained from it by passage to the associated graded groups) is an epimorphism mod.torsion, - a circumstance that occurs only rather exceptionally.

7.9.4. It is plausible, at least when f is proper and smooth, that the converse in question always holds, i.e., that $\sigma_\bullet(x)$ (resp. $\sigma^\bullet(x)$) is numerically equivalent to zero as soon as the same is true of x. This would imply that the homomorphisms (7.9.1) induce monomorphisms on the corresponding quotient groups by numerical equivalence, $\mathrm{Gr}_\bullet^{\mathrm{num}}$ and $\mathrm{Gr}^\bullet_{\mathrm{num}}$. At least the preceding discussion implies that the group $\mathrm{Gr}_\bullet^{\mathrm{num}}(X_t)$ is isomorphic to a quotient of a subgroup of $\mathrm{Gr}_\bullet^{\mathrm{num}}(X_s)$ (and in particular has rank at most that of $\mathrm{Gr}_\bullet^{\mathrm{num}}(X_s)$), and likewise for $\mathrm{Gr}^\bullet_{\mathrm{num}}$, and also for $K^\bullet_{\mathrm{num}}$ itself.

7.10. Compatibility with direct and inverse images

Let

$$g : X \longrightarrow Y$$

be a morphism of noetherian schemes flat over S. Denote[550] by σ_X (resp. σ_Y) the specialization homomorphism for X (resp. Y), and by i_X, j_X (resp. i_Y, j_Y) [print 568]the morphisms i, j for X

(resp. Y) from the special and generic fibers to the ambient scheme. Denote by

$$g_t : X_t \to Y_t, \qquad g_s : X_s \to Y_s$$

the induced morphisms, which therefore give rise to the Cartesian squares

$$\begin{array}{ccc} X_t & \xrightarrow{g_t} & Y_t \\ {\scriptstyle j_X}\downarrow & & \downarrow{\scriptstyle j_Y} \\ X & \xrightarrow{g} & Y \end{array} \qquad \begin{array}{ccc} X_s & \xrightarrow{g_s} & Y_s \\ {\scriptstyle i_X}\downarrow & & \downarrow{\scriptstyle i_Y} \\ X & \xrightarrow{g} & Y. \end{array} \tag{7.10.0}$$

First suppose that g is proper. I claim that the following diagram then commutes:

$$\begin{array}{ccc} K_\bullet(X_t) & \xrightarrow{g_{t*}} & K_\bullet(Y_t) \\ {\scriptstyle \sigma_X}\downarrow & & \downarrow{\scriptstyle \sigma_Y} \\ K_\bullet(X_s) & \xrightarrow{g_{s*}} & K_\bullet(Y_s) \end{array} \tag{7.10.1}$$

This result generalizes 7.5 (since, for $X = S$, the arrows of (7.3.2) are identified with the identity map of $\mathbb{Z}$). Note that, when X and Y satisfy the conditions of 7.7, i.e., have regular fibers, g_t and g_s automatically have finite Tor-dimension, and the diagram can be interpreted as a commutative diagram with $K^\bullet$, which we leave the reader to rewrite.

To prove the commutativity of (7.10.1), one sees at once, using the definitions of σ_X and σ_Y, that one is reduced to proving the commutativity of the two front squares of the prism

$$\begin{array}{ccccc} K_\bullet(X_t) & & \xrightarrow{g_{t*}} & & K_\bullet(Y_t) \\ & {\scriptstyle j_X^*}\nwarrow & & {\scriptstyle j_Y^*}\nwarrow & \\ & & K_\bullet(X) & \xrightarrow{g_*} & K_\bullet(Y) \\ & {\scriptstyle i_X^*}\swarrow & & {\scriptstyle i_Y^*}\swarrow & \\ K_\bullet(X_s) & & \xrightarrow{g_{s*}} & & K_\bullet(Y_s). \end{array} \tag{*}$$

[print 569]

Now these two squares are the "base-change" squares obtained from the two squares (7.10.0), so that one can apply (IV 3). One need only verify that these squares correspond to pairs (X, Y_t) and (X, Y_s) of schemes over Y that are Tor-independent over Y. This is trivial for the first pair, since $Y_t \to Y$ is flat, and also immediate for the second, since the immersion $Y_s \to Y$ is regular of codimension 1 and remains so after the base change $X \to Y$.

Now no longer suppose that g is proper, but suppose instead that it has finite Tor-dimension. I claim that the same is then true of g_t and g_s. This is trivial for the first, and for the second follows from the fact (just noted) that X and Y_s are Tor-independent over Y. One can therefore consider the square of homomorphisms

$$\begin{array}{ccc} K_\bullet(Y_t) & \xrightarrow{g_t^*} & K_\bullet(X_t) \\ {\scriptstyle \sigma_Y}\downarrow & & \downarrow{\scriptstyle \sigma_X} \\ K_\bullet(Y_s) & \xrightarrow{g_s^*} & K_\bullet(X_s), \end{array} \tag{7.10.2}$$

where the horizontal arrows are defined in (IV 2.12). Note that, when X and Y satisfy the conditions of 7.7, i.e., have regular fibers, g automatically has finite Tor-dimension (since Y is regular), and the preceding diagram can be interpreted as a commutative diagram of rings $K^\bullet$.

To prove the commutativity of (7.10.2), it again suffices to prove that of the front squares of the prism analogous to $(*)$, but with the direct-image homomorphisms g_{t*}, g_*, g_{s*} replaced by the inverse-image homomorphisms g_t^*, g^*, g_s^*. Now the commutativity of these squares follows at [print 570]once from that of the corresponding squares (7.10.0), taking account of the transitivity of inverse images (IV 2.12).

Remark 7.10.3. When g is proper, the commutativity of (7.10.1) immediately implies that of the following diagram, obtained from it by passage to the associated graded groups:

$$\begin{array}{ccc} \mathrm{Gr}_\bullet(X_t) & \xrightarrow{g_{t*}} & \mathrm{Gr}_\bullet(Y_t) \\ \downarrow{\scriptstyle \sigma_{X\bullet}} & & \downarrow{\scriptstyle \sigma_{Y\bullet}} \\ \mathrm{Gr}_\bullet(X_s) & \xrightarrow{g_{s*}} & \mathrm{Gr}_\bullet(Y_s). \end{array} \tag{7.10.1 bis}$$

Likewise, when X and Y have regular fibers and are separated, the commutativity of (7.10.2) implies that of

$$\begin{array}{ccc} \mathrm{Gr}^\bullet(Y_t) & \xrightarrow{g_t^*} & \mathrm{Gr}^\bullet(X_t) \\ \downarrow{\scriptstyle \sigma_Y^\bullet} & & \downarrow{\scriptstyle \sigma_X^\bullet} \\ \mathrm{Gr}^\bullet(Y_s) & \xrightarrow{g_s^*} & \mathrm{Gr}^\bullet(X_s). \end{array} \tag{7.10.2 bis}$$

When X and Y are moreover quasi-projective and smooth over S, one also knows that g_t^* and g_s^* are compatible with the filtrations by codimension of supports, and one obtains a commutative diagram like (7.10.2 bis), but with the $\mathrm{Gr}^\bullet$ replaced by the $\mathrm{Gr}^\bullet_{\mathrm{top}}$ (cf. Exposé 0, 2.1). One can also note that, when X and Y are quasi-projective over S with regular fibers and separated, f_t and f_s are automatically complete-intersection morphisms (VIII 1.1), and one obtains a commutative diagram analogous to (7.10.1 bis), with the $\mathrm{Gr}_\bullet$ replaced by the $\mathrm{Gr}^\bullet(\ )_{\mathbb{Q}}$, taking account of the compatibility result (VIII 3.2).

[print 571]

7.11. For a scheme T over an algebraically closed field k, one can define the subgroup of $K^\bullet(T)$ of elements algebraically equivalent to zero as the subgroup of elements of the form

$$E(x_1) - E(x_0), \tag{7.11.1}$$

where x_0 and x_1 are two k-rational points of a proper, regular, connected algebraic curve C over k, E is an element of $K^\bullet(T \times_k C)$, and, for every point x of C rational over k, one puts

$$E(x) = i_x^*(E) \in K^\bullet(T), \qquad \text{where } i_x : T \longrightarrow T \times_k C$$

is the morphism $t \mapsto (t, x)$ defined by x. The fact that the elements of the form (7.11.1) form a subgroup of $K^\bullet(T)$ follows by a standard argument, using a product $C \times_k C'$ and its pair of points (x_1, x_1'), (x_0, x_0'), and the fact that through two points of an irreducible quasi-projective algebraic scheme over k (here $C \times_k C'$) one can always pass an irreducible algebraic curve. (A reader unwilling to admit this fact may define algebraic equivalence by the group generated by the elements of the form (7.11.1).) The elements algebraically equivalent to zero form an ideal of $K^\bullet(T)$, as one sees at once, which is moreover stable under the operations λ^i when these are defined, i.e., when one works with the "naive" $K^\bullet$; this ideal is moreover contained in the augmentation ideal of $K^\bullet(T)$, as one verifies trivially. One can therefore define a quotient ring

$$K^\bullet_{\mathrm{alg}}(T), \tag{7.11.2}$$

which is even a λ-ring when one works with naive $K^\bullet$.

In the same way, one can define a subgroup of elements algebraically equivalent to 0 in $K_\bullet(T)$, taking E above in $K_\bullet(T \times_k C)$, and [print 572]noting that $i_x : T \to T \times_k C$ is a perfect morphism, since it is obtained by the flat base change $T \times_k C \to C$ from the morphism $j_x : x \to C$, which is perfect because C is regular. One thus obtains a quotient group

$$K_\bullet^{\mathrm{alg}}(T) \tag{7.11.3}$$

of $K_\bullet(T)$. This group is moreover naturally a module over $K_{\mathrm{alg}}^\bullet(T)$, since one sees at once from the definitions that the product of an element of $K^\bullet(T)$ with an element of $K_\bullet(T)$ is algebraically equivalent to zero when one of the two factors is, using (IV 2.12.2) for i_x.

One defines (when working with naive $K^\bullet$) the ideal of $\mathrm{Gr}^\bullet(T)$ of elements algebraically equivalent to zero as the ideal generated by the $c^i(x)$, where $i \geq 1$ and $x \in K^\bullet(T)$ is algebraically equivalent to zero; the quotient graded ring

$$\mathrm{Gr}_{\mathrm{alg}}^\bullet(T) \tag{7.11.4}$$

is then precisely the Chern ring associated with the λ-ring $K_{\mathrm{alg}}^\bullet(T)$.

Finally, one defines the graded subgroup of $\mathrm{Gr}_\bullet(T)$ of elements algebraically equivalent to zero (T being assumed locally noetherian) as the subgroup whose component of dimension i is the image in $\mathrm{Gr}_i(T)$ of the subgroup of $\mathrm{Fil}_i K_\bullet(T)$ consisting of the elements algebraically equivalent to zero in $K_\bullet(T)$. One finds at once that the quotient graded group

$$\mathrm{Gr}_\bullet^{\mathrm{alg}}(T) \tag{7.11.5}$$

is a module over the ring $\mathrm{Gr}_{\mathrm{alg}}^\bullet(T)$, by passage to the quotient from the module structure of $\mathrm{Gr}_\bullet(T)$ over $\mathrm{Gr}^\bullet(T)$ (3.1).

7.11.6. The preceding definitions would still make sense without assuming that k is algebraically closed, taking C above to be geometrically connected over k. It is not clear, however, that these definitions would be very useful; in particular, it does not appear that, in the case of the group $\mathrm{Pic}(T)$, one [print 573]recovers the classical "geometric" notion of algebraic equivalence considered in 2.4.1. We therefore prefer to say that an element of $K^\bullet(T)$ is algebraically equivalent to zero when its image in $K^\bullet(T \otimes_k \bar{k})$ is so, where $\bar{k}$ is an algebraic closure of k, and to proceed in the same way for the definition of algebraic equivalence in the groups $K_\bullet(T)$, $\mathrm{Gr}^\bullet(T)$ and $\mathrm{Gr}_\bullet(T)$.

7.12. With these definitions in place, I claim that the specialization homomorphism (7.3.1) carries elements algebraically equivalent to zero to elements algebraically equivalent to zero, and consequently defines, by passage to the quotient, a homomorphism

$$K_\bullet^{\mathrm{alg}}(X_t) \longrightarrow K_\bullet^{\mathrm{alg}}(X_s). \tag{7.12.1}$$

To see this, one reduces immediately to the case where A is complete and $k(s)$ is algebraically closed (using a straightforward compatibility made explicit in 7.15). It suffices to see that $\sigma_\bullet$ carries an element of the form (7.11.1) of $K_\bullet(X_t)$ to an element algebraically equivalent to zero in $K_\bullet(X_s)$. Here $T = X_t$, and we shall accordingly write C_t, E_t in place of C, E above. One knows that C_t is projective over $k(t)$, so (EGA IV 2.8.5) there is a projective flat scheme C over S whose generic fiber is C_t. Using Abhyankar's resolution of singularities of arithmetic surfaces [6], one sees that one can even choose C to be a regular scheme. (N.B. Flatness is preserved under resolution, by EGA IV 14.3.8.) Since C is proper over S, the points x_{0t} and x_{1t} of C_t rational over $k(t)$ extend to sections x_0, x_1 of C over S, and, since C is regular, it is well known that C is smooth over S at the points of $x_0(S)$, $x_1(S)$ (EGA IV, 17.12.1 c) $\Rightarrow$ a) and 19.1.1). As a [print 574]consequence, the immersions $x_i : S \to C$[551] ($i = 0, 1$) are regular and remain regular immersions $x_i : s \to C_s$ under the base change $s \to S$. Note that, by the exact

homotopy sequence (IX 1.1), $E_t \in K_\bullet(X_t \times_{k(t)} C_t)$ comes from an element $E \in K_\bullet(X \times_S C)$, so the element $E_t(x_{1t}) - E_t(x_{0t})$ of $K_\bullet(X_t)$ comes from the element $E(x_1) - E(x_0)$ of $K_\bullet(X \times_S C)$, and consequently its image in $K_\bullet(X_s)$ under (7.3.1) is precisely

$$i^*(E(x_1) - E(x_0)). \tag{*}$$

Of course, for a section x of C over S, one puts

$$E(x) = i_x^*(E), \qquad \text{where } i_x : X \longrightarrow X \times_S C$$

is defined as above, and can be regarded as obtained from x by the flat base change $X \times_S C \to C$; it is therefore a regular immersion, a fortiori a perfect morphism, which gives a meaning to the expression $E(x)$. For the same reason, the expression $E_s(x_s)$ is defined, and, by transitivity of inverse images (IV 2.12) in the following commutative diagram of perfect morphisms: [552]

$$\begin{array}{ccc} X & \xrightarrow{i_x} & X \times_S C \\ {\scriptstyle i}\uparrow & & \uparrow \\ X_s & \xrightarrow{i_{x_s}} & X_s \times_{k(s)} C_s \end{array} ,$$

one finds that the left-hand side of $(*)$ is equal to

$$E_s(x_{1s}) - E_s(x_{0s}). \tag{**}$$

Since $k(s)$ is algebraically closed, the reduced irreducible components of C_s are geometrically integral over $k(s)$ and their intersection points are rational over $k(s)$. One concludes readily that the expression $(**)$ is algebraically equivalent to zero in $K_\bullet(X_s)$, using the connectedness [print 575]of C_s, which follows from the geometric connectedness of C_t.

From the homomorphism (7.12.1), using 7.4, one obtains a homomorphism

$$\mathrm{Gr}_\bullet^{\mathrm{alg}}(X_t) \longrightarrow \mathrm{Gr}_\bullet^{\mathrm{alg}}(X_s). \tag{7.12.2}$$

When X_s and X_t are moreover assumed regular, (7.12.1) can also be interpreted as a ring homomorphism

$$K_{\mathrm{alg}}^\bullet(X_t) \longrightarrow K_{\mathrm{alg}}^\bullet(X_s), \tag{7.12.3}$$

which, when X is moreover separated, is a homomorphism of λ-rings, and consequently defines a homomorphism

$$\mathrm{Gr}_{\mathrm{alg}}^\bullet(X_t) \longrightarrow \mathrm{Gr}_{\mathrm{alg}}^\bullet(X_s). \tag{7.12.4}$$

7.12.5. Let T be a smooth proper algebraic scheme over a field k. One checks easily that an element x of $K^\bullet(T)$[553], for which there is an integer $n \geq 1$ with nx algebraically equivalent to zero, is numerically equivalent to zero. It is plausible that the converse is always true, at least when T is projective. In this question, one can plainly suppose that k is algebraically closed, and replace $K^\bullet(T)$ by $\mathrm{Gr}^\bullet(T)$ or $\mathrm{Gr}_\bullet(T)$ without changing the conjecture. One also obtains an equivalent conjecture by working with the Chow ring $A(T)$ (cf. XIV 4.1), obtaining the following more or less classical conjecture: if Z is an algebraic cycle numerically equivalent to zero on T, there is an integer $n \geq 1$ with nZ algebraically equivalent to zero. In the case where this conjecture is correct for X_t, the compatibility of the homomorphism (7.3.1) with algebraic equivalence would imply that this specialization homomorphism (when $f : X \to S$ is a projective smooth morphism) is also compatible with numerical equivalence, as conjectured in 7.9.

[print 576]

7.13. Relations with ℓ-adic cohomology

Recall (SGA 5 VII) that, for every locally free Module of finite type $\underline{E}$ on a scheme T, and every prime number distinct from the residue characteristics of T, one defines ℓ-adic Chern classes (for $i \geq 0$)

$$c^i(E) \in H^{2i}(T, \mathbb{Z}_\ell(i)),$$

satisfying an additivity formula

$$c(E) = c(E')c(E'')$$

when E is an extension of a locally free module E' by a locally free module E''. From this additivity one concludes that the total Chern class $c(E) = \sum_i c^i(E)$ arises from a homomorphism

$$c : K^\bullet(T) \longrightarrow \widehat{G}(H^{2*}(T, \mathbb{Z}_\ell(*)))^{[554]} = 1 + \prod_{i \geq 1} H^{2i}(T, \mathbb{Z}_\ell(i)), \tag{7.13.0}$$

(where the target is a group under multiplication, as explained in V 1, and where $K^\bullet(T)$ is the "naive" $K^\bullet$). One obtains canonical maps

$$c^i : K^\bullet(T) \longrightarrow H^{2i}(T, \mathbb{Z}_\ell(i)). \tag{7.13.1}$$

Now let X be a proper flat scheme over S, with regular fibers, and suppose that S is henselian. Assume that $\ell \neq$ the residue characteristic of S. One can then form the diagram [555]

$$\begin{array}{ccc} K^\bullet(X_t) & \xrightarrow{c^i} & H^{2i}(X_t, \mathbb{Z}_\ell(i)) \\ \Big\downarrow{\scriptstyle \sigma} & & \Big\uparrow{\scriptstyle \sigma^{-1}} \\ K^\bullet(X_s) & \xrightarrow{c^i} & H^{2i}(X_s, \mathbb{Z}_\ell(i)) \end{array} , \tag{7.13.2}$$

where the horizontal arrows are the "Chern class" maps for X_t and X_s respectively, and where σ^{-1} is the composite homomorphism

$$H(X_s) \xleftarrow{i^*}{\sim} H(X) \xrightarrow{j^*} H(X_t);$$

[print 577]the arrows displayed here are restriction homomorphisms, and one uses the fact that i^* is an isomorphism by the proper base change theorem (SGA 4 XII 5.5). I claim that the square (7.13.2) is commutative. The proof is essentially the same as that establishing the commutativity of (7.10.2); here it suffices to use the compatibility of the formation of Chern classes with inverse images for the morphisms $j : X_t \to X$ and $i : X_s \to X$.

7.13.3. Note that the square (7.13.2) arises by passage to the projective limit from the analogous squares in which cohomology with coefficients in the ℓ-adic sheaf $\mathbb{Z}_\ell(i)$ is replaced by cohomology with coefficients in the torsion sheaves $\mu_n^{\otimes i}$ (where n runs through the powers of ℓ). Thus the commutativity of (7.13.2) merely expresses the commutativity of the squares involving Chern classes mod. n, for every power n of ℓ. Choose an algebraic closure $\bar{K}$ of $K = k(t)$, and hence an algebraic closure $\bar{k}$ of $k = k(s)$ (namely the residue field of the integral closure of V in $\bar{K}$). For every finite subextension K_α of $\bar{K}$, one can consider the analogous commutative diagram (for fixed n) in the situation obtained from $f : X \to S$ by the base change $S_\alpha \to S$, with S_α the normalization of S in K_α. For this, note that, by Krull–Akizuki and since S is henselian, S_α is indeed a trait; moreover, one must suppose here that the fibers of X over S are geometrically regular, i.e., that X is smooth over S, so that this hypothesis is preserved by base change. As α varies, one thus obtains an inductive system of commutative squares. Passing to the inductive

limit and using (SGA 4 VII 5.8) and (IV 3.2), one obtains a commutative square (cf. 7.16): [print 578][556]

$$\begin{array}{ccc} K^\bullet(X_{\bar t}) & \xrightarrow{c^i} & H^{2i}(X_{\bar t}, \mu_n^{\otimes i}) \\ {\scriptstyle\sigma}\downarrow & & \uparrow{\scriptstyle\sigma^{-1}} \\ K^\bullet(X_{\bar s}) & \xrightarrow{c^i} & H^{2i}(X_{\bar s}, \mu_n^{\otimes i}) \end{array},$$

where $X_{\bar t}$, $X_{\bar s}$ are the geometric fibers of X. Moreover, one knows that the cohomological specialization homomorphism σ^{-1} is now an isomorphism (SGA 4 XVI 2.2), so the commutativity of the preceding square can also be expressed as the commutativity of the square obtained by reversing the preceding arrow. Passing to the projective limit over n in the commutative squares thus obtained, one finds a commutative square

$$\begin{array}{ccc} K^\bullet(X_{\bar t}) & \longrightarrow & H^{2i}(X_{\bar t}, \mathbb{Z}_\ell(i)) \\ {\scriptstyle\sigma}\downarrow & & \wr\downarrow \\ K^\bullet(X_{\bar s}) & \longrightarrow & H^{2i}(X_{\bar s}, \mathbb{Z}_\ell(i)). \end{array} \tag{7.13.4}$$

7.13.5. Another way to express this commutativity is to say that, when x_t is an element of $K^\bullet(X_t)$ and x_s is its image in $K^\bullet(X_s)$, $c^i(x_{\bar t})$ and $c^i(x_{\bar s})$ are the values, at the geometric points $\bar t$ and $\bar s$ respectively, of a section of $R^{2i}f_*(\mathbb{Z}_\ell(i))$. This section is moreover readily interpreted by taking an element $x \in K^\bullet(X)$ inducing x_t, whence a $c^i(x) \in H^{2i}(X, \mathbb{Z}_\ell(i))$, which defines the required section of $R^{2i}f_*(\mathbb{Z}_\ell(i))$.

7.13.6. Let T be a scheme. Then, by passage to the associated graded groups, the homomorphism (7.13.0) defines homomorphisms

$$\psi_i : \mathrm{Gr}^i(T) \longrightarrow H^{2i}(T, \mathbb{Z}_\ell(i)),$$

the image under ψ_i of an element x of $\mathrm{Filt}^i K^\bullet(T)$ being, by definition, $c^i(x)$. (The fact that (7.13.0) is compatible with the λ-filtration on [print 579]$K^\bullet(T)$ and the degree filtration on the second member follows readily from (V 6.6.4) and from the fact that the c^i define a Chern theory of $K^\bullet(T)$ in the sense of (V 6.13), when $K^\bullet(T)$ is regarded as a λ-algebra augmented over the binomial ring $H^0(T, \mathbb{Z})$). One thereby obtains maps

$$\mathrm{c}\ell : \mathrm{Gr}^i(T) \longrightarrow H^{2i}(T, \mathbb{Q}_\ell(i)) \overset{\text{dfn}}{=} H^{2i}(T, \mathbb{Z}_\ell(i)) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell, \tag{7.13.7}$$

by the formula

$$\mathrm{c}\ell(x^{(i)}) = (1/(-1)^{i-1}(i-1)!)\, \psi_i(x^{(i)}), \tag{7.13.8}$$

compare (XIV 5.1 and 5.2). With this understood, the commutativity of diagrams (7.13.2) and (7.13.4) trivially implies that of the analogous diagrams

$$\begin{array}{ccc} \mathrm{Gr}^i(X_t)_{\mathbb{Q}} & \xrightarrow{\mathrm{c}\ell} & H^{2i}(X_t, \mathbb{Q}_\ell(i)) \\ {\scriptstyle\sigma_{\mathbb{Q}}}\downarrow & & \uparrow{\scriptstyle\sigma^{-1}} \\ \mathrm{Gr}^i(X_s)_{\mathbb{Q}} & \xrightarrow{\mathrm{c}\ell} & H^{2i}(X_s, \mathbb{Q}_\ell(i)) \end{array}, \tag{7.13.9}$$

$$\begin{array}{ccc} \mathrm{Gr}^i(X_{\bar t})_{\mathbb{Q}} & \xrightarrow{\mathrm{c}\ell} & H^{2i}(X_{\bar t}, \mathbb{Q}_\ell(i)) \\ \downarrow & & \wr\downarrow \\ \mathrm{Gr}^i(X_{\bar s})_{\mathbb{Q}} & \xrightarrow{\mathrm{c}\ell} & H^{2i}(X_{\bar s}, \mathbb{Q}_\ell(i)) \end{array}. \tag{7.13.10}$$

Remark 7.13.11. The commutativity of the last two diagrams, which can be expressed by saying that specialization of algebraic cycles is compatible with formation of the associated ℓ-adic cohomology classes, is not very satisfactory in the form stated, since we have had to replace the coefficients $\mathbb{Z}_\ell(i)$ by $\mathbb{Q}_\ell(i)$, that is, to neglect torsion. To obtain a statement that does not neglect torsion, it seems that one [print 580]will be forced to work with the Chow ring, since it is on that ring that the homomorphism $c\ell$ is naturally defined as a homomorphism with values in $H^{2*}(-, \mathbb{Z}_\ell(*))$; compare (XIV 4.1). Unfortunately, no theory of specialization for the Chow ring has yet been developed (cf. 7.14.) that would allow one to write commutative squares analogous to (7.13.9) and (7.13.10) in terms of Chow rings and cohomology with coefficients $\mathbb{Z}_\ell(*)$. All one can say when $f : X \to S$ is smooth and proper is that there is a commutative square

$$\begin{array}{ccc} Z^i(X_t) & \xrightarrow{c\ell} & H^{2i}(X_t, \mathbb{Z}_\ell(i)) \\ \Big\downarrow{\scriptstyle \sigma} & & \Big\uparrow \\ Z^i(X_s) & \xrightarrow{c\ell} & H^{2i}(X_s, \mathbb{Z}_\ell(i)) \end{array} , \tag{7.13.12}$$

analogous to (7.13.9), and a corresponding square analogous to (7.13.10), which we leave to the reader to write down, where Z^i denotes the group of algebraic cycles (modulo no relation at all) of codimension i, where the horizontal arrows are the "associated cohomology class" homomorphisms defined in (SGA 5 IV), and where σ denotes the specialization homomorphism for algebraic cycles, associating with a cycle Z_t the cycle Z_s, where Z is the schematic closure of Z_t in X (cf. 7.14)[557]. This commutativity follows[558] from the following fact, established in (SGA 5 IV): if Z is a closed subscheme of X of codimension $\geq i$ on every fiber, one can associate with it a cohomology class $c\ell(Z) \in H^{2i}(X, \mathbb{Z}_\ell(i))$ whose formation commutes with every base change and, in particular, such that its restriction to X_t (resp. X_s) equals $c\ell(Z_t)$ (resp. $c\ell(Z_s)$). Of course, when one knows that the homomorphism σ in (7.13.12) carries cycles rationally equivalent to zero to cycles rationally equivalent to zero (X being projective over S), [print 581]one can replace the Z^i by the A^i in this diagram, and obtain a diagram of the desired kind in terms of the Chow rings of X_t and X_s.

Remark 7.13.13. Let T be a smooth projective scheme over an algebraically closed field. It follows trivially from the formalism developed in (SGA 5 IV) that, if Z is a cycle of codimension i on T whose image in $H^{2i}(T, \mathbb{Q}_\ell(i))$ is zero, then Z is numerically equivalent to zero. One of the standard conjectures in the theory of algebraic cycles asserts that the converse is also true. Moreover, by (VII 4.11), it is immediate that one obtains an equivalent conjecture by replacing the group of algebraic cycles with the group $\mathrm{Gr}^\bullet(T)_\mathbb{Q}$, using definition (7.13.8), or again with the group $K^\bullet(T)$, by stating that numerical equivalence to zero of an element of $K^\bullet(T)$ is equivalent to vanishing of its augmentation and of its Chern classes in the $H^{2i}(T, \mathbb{Q}_\ell(i))$.

With this understood, it follows immediately from the commutativity of (7.13.10) that, if $X_{\bar{t}}$ satisfies the preceding conjecture, then the specialization homomorphism $\mathrm{Gr}^\bullet(X_{\bar{t}}) \longrightarrow \mathrm{Gr}^\bullet(X_{\bar{s}})$ (and, a fortiori, $\mathrm{Gr}^\bullet(X_t) \longrightarrow \mathrm{Gr}^\bullet(X_s)$) is compatible with numerical equivalence, as conjectured in 7.9. Note also that the conjecture invoked for this purpose (numerical equivalence implies $\mathbb{Q}_\ell$-cohomological equivalence) is weaker than the one considered in 7.12.5.[559]

7.14. Specialization homomorphism for groups of cycles and cycle classes.

If T is a noetherian scheme, denote by $Z_i(T)$ (resp. $Z^i(T)$) the group of algebraic cycles on T purely of dimension i (resp. purely of codimension i). When $f : X \to S$ is of finite type, one then defines [print 582]a specialization homomorphism:

$$\sigma_i : Z_i(X_t) \longrightarrow Z_i(X_s), \tag{7.14.1}$$

by defining, for every prime cycle Z_t of X_t purely of dimension i, identified with an integral subscheme of X_t, the element $\sigma_i(Z_t)$ of $Z_i(X_s)$ as the cycle associated with the closed subscheme Z_s of X_s, where Z is the schematic closure of Z_t in X (which is a closed subscheme of X flat over S (EGA IV 2.8), with generic fiber equal to the original Z_t, and satisfying $\dim Z_s = \dim Z_t$ (EGA IV 7.1.13)). When X_s is locally equidimensional (i.e., all its irreducible components through a given point have the same dimension), one immediately verifies that the specialization homomorphism just defined induces homomorphisms

$$\sigma^i : Z^i(X_t) \longrightarrow Z^i(X_s). \tag{7.14.1 bis}$$

Now suppose that f is quasi-projective, so that the notion of rational equivalence of cycles is available on X_t and on X_s (cf. [5], where the theory is developed over an algebraically closed field). It seems very plausible that homomorphism (7.14.1) carries cycles rationally equivalent to zero to cycles rationally equivalent to zero, and thus induces a corresponding homomorphism on the "Chow" groups of cycle classes

$$\sigma_i : A_i(X_t) \longrightarrow A_i(X_s), \tag{7.14.2}$$

or again, when X_s is locally equidimensional:

$$\sigma^i : A^i(X_t) \longrightarrow A^i(X_s). \tag{7.14.2 bis}$$

One might try to prove this by a method paralleling that of 7.12, but in which the recourse to resolution of singularities and the reduction [print 583]to an algebraically closed residue field would become unnecessary, since $C = \mathbb{P}^1_S$ is available here as a "parameter curve" along which to deform cycles. A difficulty nevertheless arises, for with the notation of 7.12, E must now denote a cycle on $P^1_S \times_S X$ such that $E_t(x_{1t}) - E_t(x_{0t})$ "is defined," but it by no means follows that the cycle $E_s(x_{1s}) - E_s(x_{0s})$ is also defined. As was to be expected, one is thus reduced to formulating a statement of the type of Chow's "moving lemma" in order to define (7.14.2).

Observe, however, that the existence of homomorphism (7.4.1) shows that (7.14.1) is compatible with the equivalence relation on cycles defined by the surjective homomorphism $Z_\bullet(T) \longrightarrow \mathrm{Gr}_\bullet(T)$, which is only slightly coarser than linear equivalence; thus, when Z is quasi-projective and smooth over a field, we point out elsewhere that it differs from linear equivalence only up to torsion (XIV 4.1, 4.2). In this case one therefore obtains a homomorphism analogous to (7.14.2), or better to (7.14.2 bis), but with both members tensored by $\mathbb{Q}$.

The most natural way to define a homomorphism (7.14.2) directly would seem to be to proceed as in 7.3. This would, however, require a theory of Chow groups A_i (or of the Chow ring in the regular case) for schemes more general than schemes quasi-projective over a field, including X, and a contravariant law for these groups that would allow one to define the restriction homomorphism to X_s, $A_i(X) \longrightarrow A_i(X_s)$. Compare XIV 8. Once such a sufficiently general theory of Chow groups and rings is available, it should be possible to transcribe into Chow theory all the results enumerated above for the specialization homomorphism in K-theory.

[print 584]


7.15. Varying S.

Let S' be a second trait, and let

$$S' \longrightarrow S$$

be a morphism of traits, i.e., a dominant local morphism (i.e., carrying the closed point to the closed point and the generic point to the generic point). Denote by s' (resp. t') the closed point

(resp. the generic point) of S', and set $X' = X \times_S S'$. Suppose that X' is noetherian (e.g., X of finite type over S). There is a commutative diagram



whose oblique arrows are flat morphisms. One can therefore take the inverse-image homomorphisms for $K_\bullet$ along all the arrows of the diagram (IV 2.12). Applying definition 7.3, one finds that the following diagram [560]

$$\begin{array}{ccc} K_\bullet(X_t) & \xrightarrow{\sigma_X} & K_\bullet(X_s) \\ \downarrow & & \downarrow \\ K_\bullet(X'_{t'}) & \xrightarrow{\sigma_{X'}} & K_\bullet(X'_{s'}) \end{array} \tag{7.15.1}$$

is commutative, where the vertical arrows are the inverse-image homomorphisms. One immediately obtains an analogous commutative diagram with $\mathrm{Gr}_\bullet$ in place of $K_\bullet$; and when X_s and X_t are regular, one can also write $K^\bullet$ and $\mathrm{Gr}^\bullet$ in place of $K_\bullet$ and $\mathrm{Gr}_\bullet$.

[print 585]

7.16. Passage to geometric fibers.

Let $\bar{K}$ be an algebraic closure of $K = k(t)$, the inductive limit of its finite subextensions K_i, and let $\bar{A} = \lim \bar{A}_i$ be the normalization of A in $\bar{K}$, the inductive limit of its normalizations $\bar{A}_i$ in the K_i. Choose a maximal ideal $\bar{\underline{m}}$ of $\bar{A}$, thus inducing on each $\bar{A}_i$ a maximal ideal $\underline{m}_i$, and set $B_i = \bar{A}_{i\,\underline{m}_i}$ [561], which by Krull–Akizuki is again a discrete valuation ring, whose spectrum we denote by S_i and whose points we denote by s_i and t_i. The residue field $\bar{k} = \bar{A}/\bar{\underline{m}}$ is an algebraic closure of $k = k(s)$, and is the inductive limit of the residue fields $k_i = \bar{A}_i/\underline{m}_i = k(s_i)$ [562] of the B_i. Set $\bar{t} = \mathrm{Spec}(\bar{K})$, $\bar{s} = \mathrm{Spec}(\bar{k})$. Now suppose X of finite type over S; then $X_{\bar{t}}$ and $X_{\bar{s}}$ are of finite type over $\bar{t}$ and $\bar{s}$, respectively, and hence noetherian. By IV 3.2.3, we have

$$K_\bullet(X_{\bar{t}}) \overset{\sim}{\leftarrow} \varinjlim_i K_\bullet(X_{t_i}), \qquad K_\bullet(X_{\bar{s}}) \overset{\sim}{\leftarrow} \varinjlim_i K_\bullet(X_{s_i}).$$

On the other hand, by 7.15 there is a homomorphism from the inductive system of the $K_\bullet(X_{t_i})$ to the inductive system of the $K_\bullet(X_{s_i})$. Passing to the limit, one obtains a canonical homomorphism:

$$K_\bullet(X_{\bar{t}}) \longrightarrow K_\bullet(X_{\bar{s}}), \tag{7.16.1}$$

again called, as expected, the specialization homomorphism. It induces a homomorphism $\mathrm{Gr}_\bullet(X_{\bar{t}}) \longrightarrow \mathrm{Gr}_\bullet(X_{\bar{s}})$ by 7.4, using the fact (whose verification is left to the reader) that the functor $\mathrm{Gr}_\bullet$ (just like $K_\bullet$) commutes with the limits occurring here. Similarly, when f is smooth, one can interpret (7.16.1) as a homomorphism on $K^\bullet$, which is moreover a ring homomorphism, and even a homomorphism of λ-rings when X is separated; in this case it therefore induces a homomorphism on $\mathrm{Gr}^\bullet$.

[print 586]7.16.2. We shall not review here, in the case of geometric fibers, the extension of all the properties of specialization homomorphisms proved in the preceding sections for the "arithmetic" fibers; they all follow trivially from the latter by passage to the limit. Let us mention only the following interesting consequence of 7.9.4 when f is smooth and proper: $\mathrm{Gr}_\bullet^{\mathrm{num}}(X_{\bar{t}})$ is isomorphic to a quotient of a subgroup of $\mathrm{Gr}_\bullet^{\mathrm{num}}(X_{\bar{s}})$; in particular, its rank is bounded above by that of the latter group. This is therefore a semicontinuity result for the rank of the group of cycles modulo numerical equivalence.

7.17. Case of an arbitrary base.

Now let

$$f : X \longrightarrow S$$

be a flat and quasi-compact morphism of noetherian preschemes, and let s, t be two points of S such that s is a specialization of t. We wish to relate the groups $K_\bullet$, etc., for the two fibers X_t and X_s, and for corresponding geometric fibers. For this, note (EGA II 7.1.9) that there exists a morphism

$$g : S' \longrightarrow S,$$

where S' is a trait, and where g carries the closed point s' to s and the generic point t' to t; one may even assume that $k(t) \xrightarrow{\sim} k(t')$. Suppose that $X' = X \times_S S'$ is noetherian, as is the case when f is of finite type or S' is essentially of finite type over S (EGA IV 1.3.8) (which one may assume when S itself is universally Japanese (EGA 0_{IV} 23.1.1)). Then the specialization homomorphism for X' defines a homomorphism

$$\sigma_\bullet : K_\bullet(X_t) \xrightarrow{\sim} K_\bullet(X_{t'}) \longrightarrow K_\bullet(X_{s'}). \tag{7.17.1}$$

[print 587]One thus obtains a specialization homomorphism on $K_\bullet(X_t)$, but with values not in $K_\bullet(X_s)$, but in $K_\bullet(X_{s'})$, where $X_{s'}$ is obtained from X_s by an extension $k(s) \to k(s')$ of the base fields (which may be assumed to be of finite type when S is universally Japanese). Passing to geometric fibers as explained in 7.16, one likewise obtains a homomorphism

$$\sigma_\bullet : K_\bullet(X_{\bar{t}}) \longrightarrow K_\bullet(X_{\overline{s'}}); \tag{7.17.2}$$

so here again the "geometric" specialization homomorphism takes its values not in $K_\bullet(X_{\bar{s}})$, but in $K_\bullet(X_{\overline{s'}})$, where $X_{\overline{s'}}$ is obtained from $X_{\bar{s}}$ by the extension of algebraically closed fields $k(\bar{s}) \to k(\overline{s'})$.

7.17.3. In general, the homomorphism $K_\bullet(X_{\bar{s}}) \to K_\bullet(X_{\overline{s'}})$ is not an isomorphism (because of the possible presence of "continuous" components in the group $K_\bullet(X_{\bar{s}})$). It is, however, easy to verify that it is an isomorphism modulo algebraic equivalence (7.11), and also modulo numerical equivalence (4.5.1) when f is smooth and proper. Consequently, using 7.12, one finds that (7.17.2) induces a specialization homomorphism

$$K_\bullet^{\mathrm{alg}}(X_{\bar{t}}) \longrightarrow K_\bullet^{\mathrm{alg}}(X_{\bar{s}}); \tag{7.17.3.1}$$

and when f is proper and smooth, provided that specialization homomorphism (7.17.2) is compatible with numerical equivalence (cf. 7.9.4), one likewise obtains a homomorphism

$$K_\bullet^{\mathrm{num}}(X_{\bar{t}}) \longrightarrow K_\bullet^{\mathrm{num}}(X_{\bar{s}}). \tag{7.17.3.2}$$

We leave to the reader the task of developing (if needed) the corresponding constructions for $\mathrm{Gr}_\bullet$, $K^\bullet$, and $\mathrm{Gr}^\bullet$, and the properties of these specialization homomorphisms that follow trivially from the properties developed in the preceding sections. Let us mention only [print 588]that it

follows from 7.16.2 that $\mathrm{Gr}_\bullet^{\mathrm{num}}(X_{\bar{t}})$ is isomorphic to a quotient of a subgroup of $\mathrm{Gr}_\bullet^{\mathrm{num}}(X_{\bar{s}})$ (f proper and smooth); in particular, its rank is again bounded above by the rank of the latter.

7.17.4. When S is regular at s and t is maximal, one can construct S' by blowing up s in the localization of S at s, and localizing at the generic point of the fiber of s in the blow-up. One then obtains for $k(s')$ a pure extension of $k(s)$. Now it is not difficult to show that, when k' is a pure extension of a field k, for every noetherian scheme T over k the canonical homomorphism $K_\bullet(T) \to K_\bullet(T \otimes_k k')$ is an isomorphism, by representing k' as[563] the inductive limit of the coordinate rings of affine open subsets of schemes of the form $\operatorname{Spec} k[T_1, \ldots, T_n]$, and using (IX 1.6). Thus one concludes that (7.17.1) can be regarded as a canonical homomorphism

$$K_\bullet(X_t) \longrightarrow K_\bullet(X_s). \tag{7.17.4.1}$$

But note that one cannot deduce from this a homomorphism $K_\bullet(X_{\bar{t}}) \to K_\bullet(X_{\bar{s}})$, for when one passes to a finite extension K_i of $K = k(t)$ and normalizes S in K_i, the regularity hypothesis on S is lost.

Another way to define a homomorphism (7.17.4.1) is to reduce to the case where S is local with closed point s (which is trivial), and to consider an increasing sequence of regular closed subschemes S_i of S, with generic points t_i, such that $\dim S_i = i$ $(0 \leq i \leq n = \dim S)$. Since for every i, $0 \leq i \leq n-1$, $\underline{O}_{S_{i+1},t_i}$ is a discrete valuation ring with residue field $k(t_i)$ and fraction field $k(t_{i+1})$, there is a specialization homomorphism

$$K_\bullet(X_{t_{i+1}}) \longrightarrow K_\bullet(X_{t_i}),$$

[print 589]and the composite of these homomorphisms for $0 \leq i \leq n-1$ defines a homomorphism (7.17.4.1), which one can show (by proceeding as in 7.3) to be the same as the one defined above by means of a blow-up. The advantage of the second method, however, is that here one even has homomorphisms of the form (7.16.1):

$$K_\bullet(X_{\bar{t}_{i+1}}) \longrightarrow K_\bullet(X_{\bar{t}_i}),$$

(with suitable coherent choices of the algebraic closures $k(\bar{t}_i)$ of the $k(t_i)$), which allow one to define by composition a homomorphism

$$K_\bullet(X_{\bar{t}}) \longrightarrow K_\bullet(X_{\bar{s}}), \tag{7.17.4.2}$$

that probably depends essentially on the choice of the S_i, that is, of the t_i.

7.17.5. The construction of a specialization homomorphism (7.17.4.2) can be generalized to the case in which one no longer assumes that S is regular at s and that t is a maximal point, provided that resolution of singularities is available for the integral closed subscheme T of S with generic point t, or merely for its localization at s; this hypothesis holds in particular when $\underline{O}_{S,s}$ is excellent of characteristic zero, by Hironaka. Indeed, one can then find a morphism $S' \to \operatorname{Spec}(\underline{O}_{T,s})$ with S' regular, and, taking a closed point of the fiber S'_s, apply the preceding construction to $X' = X \times_S S'$ over S', the generic point t' of S' and its closed point s'. This yields a homomorphism of the form (7.17.4.2), since $k(s')$ is a finite extension of $k(s)$ and therefore has the same algebraic closure.

7.17.6. In all the constructions considered in section 7.17, the construction of a specialization homomorphism has been reduced to the [print 590]case where the base is a trait. It follows that the properties of the specialization homomorphism reviewed in the preceding sections, in the case where the base is a trait, extend automatically to the case of a more general base. Of course, the methods of definition considered apply equally well to $\mathrm{Gr}_\bullet$, $K^\bullet$ (when f is smooth), and $\mathrm{Gr}^\bullet$ (when, in addition, f is separated).

7.17.7. Finally, note that the definition of a homomorphism (7.17.3.1) or (7.17.3.2), or of a homomorphism (7.17.4.2), depends essentially on certain arbitrary choices (a suitable trait over S in 7.17.3, a sequence of (S_i) in 7.17.4, and, in addition, a resolution of singularities in 7.17.5), and likewise for the variants for $\mathrm{Gr}_\bullet$, etc. Obvious examples can be constructed with s, say, an ordinary double point of an algebraic curve and X a projective smooth scheme over S (for example, an abelian scheme of relative dimension 2), in which the specialization homomorphism on $\mathrm{Gr}^1 = \mathrm{Pic}$ and even on $\mathrm{Gr}^1_{\mathrm{alg}} = \mathrm{NS}$ (the Néron–Sévéri group) actually depends on the choice of the trait S' chosen over S or, what amounts to the same thing here, on the choice of a point over s in the normalization of S.

7.18. Appendix to the Appendix: extension of the base field for certain functors with values in commutative groups

We spell out here a technical point raised in 7.17.4.

7.18.1. Let k be a field, C the category of integral k-algebras and flat homomorphisms of such algebras, and F a functor on C with values in the category (Ens) of sets. Suppose that F commutes with filtered inductive limits. Let K be an extension of k; we seek [print 591]criteria for $F(k) \longrightarrow F(K)$ to be injective (resp. surjective).

7.18.2. Suppose that we are in one of the following two cases:

a) The field k is separably closed, and K/k is a separable extension.

b) The field k is infinite, and K is a pure extension of it.

Then the map $F(k) \to F(K)$[564] is injective.

Indeed, using the fact that K is the filtered inductive limit of subalgebras of finite type over k, with flat transition morphisms, and the commutation of F with filtered inductive limits, one is reduced in case a) to proving that, if A is an integral algebra of finite type over k whose fraction field K is separable over k, then $F(k) \to F(A)$ is injective. Now the hypotheses on k and on K imply that there exists a k-homomorphism $A \to k$; the composite $F(k) \to F(A) \to F(k)$ is then the identity, whence the conclusion. In case b), passage to the limit similarly reduces the question to proving that, if A is an integral algebra of finite type over k whose fraction field is a pure extension of k, then $F(k) \to F(A)$ is injective. Since k is infinite, the hypothesis again implies that there exists a k-homomorphism $A \to k$, and one concludes as above.

7.18.3. Now suppose that F is a functor with values in the category (Ab) of abelian groups, and suppose that for every finite extension k' of k a norm homomorphism

$$N_{k'/k} : F(k') \longrightarrow F(k)$$

has been defined such that the composite

$$F(k) \longrightarrow F(k') \xrightarrow{N_{k'/k}} F(k)$$

is multiplication by $[k' : k]$. Under these conditions:

(i) [print 592]For every extension K of k, the kernel of $F(k) \to F(K)$ is a torsion group.

(ii) For every pure extension K of k, the homomorphism $F(k) \to F(K)$ is injective. (NB. Note that k is not necessarily assumed infinite).

In case (i), one is reduced to proving that every element x in the kernel of $F(k) \to F(A)$, where A is an integral algebra of finite type over k, is a torsion element. Using a k-homomorphism $A \to k'$, where k' is a finite k-algebra, one is reduced to the case where $A = k'$ is a finite extension of k. One then has $[k' : k]x = 0$ by the property postulated above for the norm homomorphism. This proves that x is a torsion element. One even has $x = 0$ when the greatest common divisor of the degrees of the finite extensions k' of k that are quotients of A equals 1. This is the case when the fraction field K of A is a pure extension of k, as is readily checked (only the case where k is finite requires a new verification). This observation then gives the conclusion of (ii).

7.18.4. Suppose that, for every nonempty affine open subset U of the affine space $\operatorname{Spec} k[T_1, \ldots, T_n]$, with A the ring of U, the homomorphism $F(k) \to F(A)$ is surjective (compare IX 1.6). Then, using the commutation of F with filtered inductive limits, one finds that for every pure extension K of k, $F(k) \to F(K)$ is surjective. If one also assumes that k is infinite, or that norm homomorphisms can be defined as in 7.18.3, one concludes that this homomorphism is even bijective.

7.18.5. As an example, consider a scheme X of finite type over k, and set

$$F(A) = K_\bullet(X \otimes_k A),$$

[print 593]which is a functor on the category C (by (IV 2.12)).[565] By (IV 3.2.3), this functor commutes with filtered inductive limits. Moreover, when k' is a finite extension of k, the projection $p : X \otimes_k k' \to X$ is a finite, flat morphism of finite presentation, and therefore defines a direct-image homomorphism $p = N_{k'/k} : K_\bullet(X \otimes_k k') \longrightarrow K_\bullet(X)$. One immediately verifies that the condition on the norm homomorphisms considered in 7.18.3 does hold. Finally, if A is the coordinate algebra of an open subset U of the affine space $\operatorname{Spec} k[T_1, \ldots, T_n]$, then $F(k) \to F(A)$ is identified with the map $K_\bullet(X) \longrightarrow K_\bullet(X \times_k U)$, which is surjective for X noetherian, by IX 1.6. Using the results proved above, one then obtains:

(i) If K/k is a pure extension of k, then $K_\bullet(X) \longrightarrow K_\bullet(X_K)$ is injective, and is even bijective when X is noetherian.

(ii) If k is separably closed and K/k is a separable extension, then $K_\bullet(X) \longrightarrow K_\bullet(X_K)$ is injective.

(iii) For every extension K of k, the kernel of $K_\bullet(X) \longrightarrow K_\bullet(X_K)$ is a torsion group.

These results remain valid when one assumes only that X is noetherian, provided one restricts to extensions K of k that are of finite type (so that X_K is still noetherian[566]). Moreover, essentially the same results[567] hold for $K^\bullet$ in place of the functor $K_\bullet$, with the sole difference that in (i) one cannot assert bijectivity in general, for lack of the homotopy theorem for $K^\bullet$ in place of $K_\bullet$; to conclude bijectivity, one would have to assume that X is noetherian and regular, which implies that the groups $K^\bullet$ and $K_\bullet$ are the same for X and for X_K, respectively.

[print 594]One again obtains essentially the same results for $\mathrm{Gr}^\bullet$ in place of $K_\bullet$ or $K^\bullet$, with the same reservation concerning the bijectivity assertion in (i). In the case where one restricts to $\mathrm{Gr}^1 = \mathrm{Pic}$, this bijectivity assertion is valid when X is noetherian and normal, for in this case, for every open subset U of an affine space over k, one sees immediately that $\mathrm{Pic}(X) \longrightarrow \mathrm{Pic}(X \times_k U)$ is surjective (EGA $\mathrm{Err}_{\mathrm{IV}}$ 53, 21.4.13).

BIBLIOGRAPHY

[1] Dickson, E. Linear groups, Dover, New York, 1958.

[2] Grothendieck, A. TDTE, V, VI; Séminaire Bourbaki 232, 236.

[3] Kleiman, S. Towards a numerical theory of ampleness, Ann. of Math. 84 (1966), pp. 293-344.

[4] Murre, J. On contravariant functors from the category of préschemes over a field into the category of abelian groups, Publications Mathématiques de l'IHES, 23, 1964.

[5] Chevalley, C. Les classes d'équivalence rationnelles, I et II, Séminaire à l'ENS 1958, exposés 2 et 3.

[6] S. Abhyankar, Resolution of singularities of arithmetical surfaces, in: Arithmetical Algebraic Geometry, Purdue University, Harper–Row, New York 1965.

[print 595]

EXPOSÉ XII
A RELATIVE REPRESENTABILITY THEOREM FOR THE PICARD FUNCTOR

by M. Raynaud (written up by S. Kleiman)

Warning

The purpose of the two exposés XII, XIII is to give proofs of the nine finiteness theorems for the Picard functor stated by A. Grothendieck in the comments (pp. C 07-C 11) to his "Fondements de la Géométrie Algébrique" (cited as FGA). In the first exposé, the first finiteness theorem is proved in the following form, more precise than the one stated in FGA: let X and Y be two proper schemes over an integral noetherian base S, and let $f : X \longrightarrow Y$ be a surjective S-morphism. Then there is a nonempty open subset V of S over which the induced morphism $f^* : \mathrm{Pic}_{Y/S} \longrightarrow \mathrm{Pic}_{X/S}$ is representable by a quasi-affine morphism of finite presentation. In the second exposé, the rest is proved.

The two exposés differ notably in their proof techniques. In the first, essential use is made of Oort's dévissage, non-flat descent, and the representation of unramified functors. In the second, one first reduces to the case where the schemes X/S that occur are projective and flat with geometrically integral fibers, or even, as needed, normal or smooth, always by applying the first finiteness theorem to a suitable morphism $X' \to X$. One then uses the tool of (b)-sheaves forged by G. Castelnuovo[568] and sharpened and used by T. Matsusaka, D. Mumford, Y. Nakai and S. Kleiman. This tool makes it possible, in particular, to dispense with Chow coordinates, the Frobenius morphism and the Lefschetz theorems used in Grothendieck's initial proofs.

The present exposé XII was presented orally, following Grothendieck, by M. Raynaud, whom the writer wishes to thank for communicating his personal notes.

[print 596]

1. Statement of the main theorem and applications

Theorem 1.1. Let S be a noetherian scheme, X and Y two proper S-schemes, and $f : X \to Y$ a morphism. Suppose that S is integral and f surjective. Then there is a nonempty open subset V of S such that the morphism

$$f^* : \mathrm{Pic}_{Y/S}|V \longrightarrow \mathrm{Pic}_{X/S}|V$$

between the Picard functors (localized for the f.p.p.f topology) is representable by a quasi-affine morphism of finite presentation.

Corollary 1.2. Let X be a proper scheme over a noetherian scheme S. Suppose that S is integral. Then there is a nonempty open subset V of S such that the Picard functor $\mathrm{Pic}_{X/S}|V$ is representable by a V-group scheme that is a sum of quasi-projective open subschemes of finite type, hence a fortiori separated over V.

Indeed, recall the following fact (FGA N°232): let S be a scheme, and let X be a projective S-scheme, of finite presentation and integral over S (that is, flat with geometrically integral

fibers). Then the Picard functor is represented by an S-group scheme $\underline{\mathrm{Pic}}_{X/S}$, an increasing union of quasi-projective open subschemes of finite presentation.

Consideration of the "Hilbert polynomials" makes it possible here to decompose $\underline{\mathrm{Pic}}_{X/S}$ as a sum of open subschemes $\underline{\mathrm{Pic}}^q_{X/S}$, $q \in \mathbb{Q}[t]$. In the next exposé it is proved, under the additional hypothesis that S is noetherian, that the $\underline{\mathrm{Pic}}^q_{X/S}$ are of finite type over S. It is then known that 1.2 holds in the case where X is in addition projective and integral over S; we shall deduce the general case using the

Lemma 1.3. Let X be of finite type over an integral noetherian S. Then there are a nonempty open subset V of S, an integral scheme S' and a finite and [print 597]flat morphism $S' \to V$ such that every irreducible component X'_i of $X_{S'}$, endowed with the induced reduced structure, is integral over S'.

Indeed, there is a finite extension L of the function field $k(S)$ such that every irreducible component of $(X_L)_{\mathrm{red}}$ is geometrically integral (by EGA IV 4.6.8). Let S_1 be an integral S-scheme such that $k(S_1) = L$ and such that there are a nonempty open subset V_1 of S and a finite flat morphism $h : S_1 \to V_1$. There is a nonempty open subset S_2 of S_1 such that every irreducible component of X_{S_2}[569], endowed with the induced reduced structure, is integral over S_2 (EGA IV 9.7.7)[570]. Take $V = V_1 - h(S_1 - S_2)$[571] and $S' = h^{-1}(V)$.

Let us now prove 1.2. Let $h : S' \to V_1$ and X'_i be as in 1.3. For every i, let $f_i : Z_i \to X'_i$ be a Chow presentation with Z_i integral and f_i birational. After shrinking V and S', we may suppose that Z_i is integral over S'[572] (EGA IV 9.7.7). Let $f : Z = \coprod_i Z_i \longrightarrow X' = X_{S'}$ be the canonical morphism. Then $\underline{\mathrm{Pic}}_{Z/S'} = \prod_i \underline{\mathrm{Pic}}_{Z_i/S'}$ is representable by a group scheme that is a sum of quasi-projective open subschemes of finite type.

Finally, 1.1 says that, after shrinking V and S', the morphism $f^* : \underline{\mathrm{Pic}}_{X'/S'} \longrightarrow \underline{\mathrm{Pic}}_{Z/S'}$[573] is representable by a quasi-affine morphism. Consequently, $\mathrm{Pic}_{X'/S'}$ is representable by a group scheme that is a sum of quasi-projective open subschemes of finite type; by finite flat descent (SGA 1 VIII 7.6 and EGA II 6.6.1), the same holds for $\mathrm{Pic}_{X/S}\,|V$.

Lemma 1.4. Let k be a field, P and Q two k-group schemes locally of finite type, and let $f : P \to Q$ be a quasi-affine morphism. Then f is affine.

Indeed, let $R = \mathrm{Ker}(f)$; R is a k-group scheme of finite type that is quasi-affine; by (SGA 3 $\mathrm{VI_B}$ 11.11), R is therefore affine. [print 598]Thus $P \to P/R$, which makes P an R-torsor over P/R, is affine. Moreover, it is known (SGA 3 1.4.2) that $P/R \to Q$ is a closed immersion, hence affine, and hence the same holds for the composite $P \to P/R \to Q$.

Corollary 1.5. Let k be a field.

a) Let X be a proper k-scheme. Then $\underline{\mathrm{Pic}}_{X/k}$ is representable (J. P. Murre).

b) Let $f : X \to Y$ be a surjective morphism between two proper k-schemes. Then the induced morphism $f^* : \underline{\mathrm{Pic}}_{Y/k} \longrightarrow \underline{\mathrm{Pic}}_{X/k}$ is affine.

Corollary 1.6. Let S be an integral noetherian scheme, and $f : X \to Y$ a surjective S-morphism between two proper S-schemes.

a) If Y is normal over S (that is, flat with geometrically normal fibers), then, after shrinking S, $f^* : \underline{\mathrm{Pic}}_{Y/S} \to \underline{\mathrm{Pic}}_{X/S}$ is quasi-finite.

b) If $\underline{\mathrm{Pic}}_{Y/S}$ is essentially proper over S (that is, if it satisfies the valuative criterion of properness, for example, if Y is smooth over S), then, after shrinking S, $f^* : \underline{\mathrm{Pic}}_{Y/S} \to \underline{\mathrm{Pic}}_{X/S}$ is finite (and in particular affine).

Indeed, in a), for every $s \in S$, $R_s = \mathrm{Ker}(f_s^*)$ is an affine algebraic group; its identity component is a proper algebraic group over k (FGA 236, Theorem 2.1 (iii)); R_s is therefore finite. In b), f^* is both proper and quasi-affine, hence finite.

Remark 1.7. It is not known whether 1.1 remains valid when "quasi-affine" is replaced there by affine, as 1.5 and 1.6 might suggest.

[print 599]

2. First reductions

2.1. Apply 1.3 to X/S and Y/S: we obtain a nonempty open subset V of S, an integral scheme S' and a finite flat morphism $S' \to V$ such that every irreducible component X'_i of $X_{S'}$ and Y'_j of $Y_{S'}$, endowed with the induced reduced structure, is integral over S'. By f.p.q.c. descent (EGA IV 2.7.1 (xiv) and SGA 1 VIII 7.9), we may replace S by S'. It then follows from the following commutative diagram:

$$\begin{array}{ccc} X_{\mathrm{red}} & \xrightarrow{f_{\mathrm{red}}} & Y_{\mathrm{red}} \\ \downarrow{\scriptstyle i} & & \downarrow{\scriptstyle j} \\ X & \xrightarrow{f} & Y \end{array} \quad ,$$

implying the commutative diagram

$$\begin{array}{ccc} \underline{\mathrm{Pic}}_{X_{\mathrm{red}}/S} & \longleftarrow & \underline{\mathrm{Pic}}_{Y_{\mathrm{red}}/S} \\ \uparrow & & \uparrow \\ \underline{\mathrm{Pic}}_{X/S} & \longleftarrow & \underline{\mathrm{Pic}}_{Y/S} \end{array} \quad ,$$

and from trivial sorites on morphisms of functors representable by quasi-affine morphisms (following from EGA II 5.1.10 (ii) (v)) that it suffices to prove the following two assertions:

I. Theorem 1.1 when f is an immersion (necessarily nilpotent).

II. Theorem 1.1 when X and Y are reduced and every irreducible component X_i of X and Y_i of Y, endowed with the induced reduced structure, is integral over S.

2.2. Let us study assertion II. For every i, let $j = \varphi(i)$ be an index such that $f(X_i) \subset Y_j$, and put $\Sigma_j = \coprod_{\varphi(i)=j} X_i$. We then have the two diagrams [print 600]

$$\begin{array}{ccc} Y & \longleftarrow & \coprod_j Y_j \\ \uparrow & & \uparrow \\ X & \longleftarrow & \coprod_j \Sigma_j \end{array} \quad ,$$

$$\begin{array}{ccc} \underline{\mathrm{Pic}}_{Y/S} & \xrightarrow{①} & \underline{\mathrm{Pic}}_{\coprod_j Y_j/S} = \prod_j \underline{\mathrm{Pic}}_{Y_j/S} \\ ②\downarrow & & \downarrow④ \\ \underline{\mathrm{Pic}}_{X/S} & \xrightarrow{③} & \underline{\mathrm{Pic}}_{\coprod_j \Sigma_j/S} = \prod_j \underline{\mathrm{Pic}}_{\Sigma_j/S}. \end{array}$$

And to establish II, it suffices to see that, after shrinking S, ①, ③ and ④ are quasi-affine.

Consider ④ first. Replacing Y by Y_j and X by Σ_j reduces us to the case where Y is integral over S. Since f is surjective, one of the X_i, say X_0, dominates Y, and we have the two sequences

$$X_0 \hookrightarrow \coprod X_i \longrightarrow Y,$$

$$\mathrm{Pic}_{X_0/S} \longleftarrow \mathrm{Pic}_{\coprod X_i/S} \longleftarrow \mathrm{Pic}_{Y/S}\,.^{[574]}$$

Thus, knowing that the $\underline{\mathrm{Pic}}_{X_i/S}$, and therefore also $\mathrm{Pic}_{\coprod X_i/S}$, are representable and that the composite morphism is quasi-affine, we conclude that ④ is quasi-affine. In view of the proof of 1.2, for ④ we are therefore reduced to proving II'. Theorem 1.1 when X and Y are integral over S.

For ① and ③, we must treat the case of a morphism $\coprod X_i \to X$ where X is reduced, the X_i are the irreducible components of X and are integral over S. Now we may suppose that X has connected fibers. Indeed, let η be the geometric point of S. Then the irreducible components of X_η are the $(X_i)_\eta$ and, after shrinking S, we may suppose that $X_i \cap X_j = \varnothing$ if and only if $^{[\text{print } 601]}(X_i)_\eta \cap (X_j)_\eta = \varnothing$. Let $Y_k = \bigcup_{i\in I(k)} X_i$ be the connected components of X; then f decomposes, $f = \coprod_k f_k : \coprod_k (\coprod_{i\in I(k)} X_i) \longrightarrow \coprod_k Y_k$[575], and the assertion follows. Finally, an immediate induction reduces us to the case of a morphism $X_1 \amalg X_2 \to X$. After shrinking S, ① and ③ then follow, by (EGA III 7.8.6), from the

Proposition 2.3. Let S be a scheme, $g : Y \to S$ a proper morphism, and Y_1, Y_2 two closed subschemes defined by the ideals I_1, I_2. Suppose that

a) $Y, Y_1, Y_2, Y_1 \cap Y_2$ are faithfully flat and of finite presentation over S.

b) $g_*O_Y = O_S$ and $(g_i)_*O_{Y_i} = O_S$ universally (where $g_i = g|Y_i$, $i = 1, 2$).

c) $I_1 \cap I_2 = 0$.

Then $\underline{\mathrm{Pic}}_{Y/S} \longrightarrow \underline{\mathrm{Pic}}_{(Y_1 \amalg Y_2)/S}$ is representable by an affine morphism of finite presentation.

Indeed, first note that the hypotheses are stable under every base change. For (c), this follows from the two exact sequences:

$$0 \longrightarrow I_1 + I_2 \longrightarrow O_Y \longrightarrow O_{Y_1 \cap Y_2} \longrightarrow 0$$

and

$$0 \longrightarrow I_1 \cap I_2 \longrightarrow I_1 \oplus I_2 \longrightarrow I_1 + I_2 \longrightarrow 0\ ;$$

the first shows that $I_1 + I_2$ is flat over S; then the second shows that $I_1 \cap I_2$ is stable under base change.

The assertion is local for the f.p.p.f. topology. Making the base change $Y_1 \cap Y_2 \to S$, we may therefore suppose that $Y_1 \cap Y_2/S$ has a section α, which defines sections α_i of Y_i and β of Y.

After a suitable base change, we are reduced to seeing that every point $\mu : S \to \underline{\mathrm{Pic}}_{(Y_1 \amalg Y_2)/S}$ has as its inverse image a subfunctor of $^{[\text{print } 602]}\underline{\mathrm{Pic}}_{Y/S}$ that is representable by an affine scheme over S. Now μ is defined by an invertible sheaf L_i on Y_i equipped with a rigidification $u_i : \alpha_i^* L_i \xrightarrow{\sim} O_S$ for $i = 1, 2$. Likewise, a point $\lambda : S \to \underline{\mathrm{Pic}}_{Y/S}$ corresponds to an invertible sheaf L on Y equipped with a rigidification $u : \alpha^* L \xrightarrow{\sim} O_S$. The image of λ in $\mathrm{Pic}_{Y_1 \amalg Y_2/S}$ is μ if and only if there is an isomorphism of rigidified sheaves $f_i^*(L) \simeq L_i$ where $f_i : Y_i \hookrightarrow Y$ is the inclusion for $i = 1, 2$. Observe that these isomorphisms define an isomorphism $\varphi : L_1|Y_1 \cap Y_2 \xrightarrow{\sim} L_2|Y_1 \cap Y_2$ such that $u_1 = u_2 \circ \varphi$.

Conversely, given an isomorphism φ, $L_1|Y_1 \cap Y_2 \to L_2|Y_1 \cap Y_2$[576], such that $u_1 = u_2 \circ \varphi$, there is a rigidified invertible sheaf L on Y obtained by gluing along φ; explicitly, L is the kernel of

the canonical homomorphism $(f_1)_*L_1 \oplus (f_2)_*L_2 \longrightarrow (f|Y_1 \cap Y_2)_*L_2$ defined by φ. This L then corresponds to a point λ of $\underline{\mathrm{Pic}}_{Y/S}$ mapping to μ.

Finally, let I be the functor defined, as T runs through the schemes over S, by the sets of isomorphisms $\varphi : (L_1)_T|(Y_1 \cap Y_2)_T \xrightarrow{\sim} (L_2)_T|(Y_1 \cap Y_2)_T$ such that $(u_1)_T = (u_2)_T \circ \varphi$. We have just established an isomorphism between the inverse image of μ in $\underline{\mathrm{Pic}}_{Y/S}$ and the functor I. It therefore suffices to apply the following lemma (taking $X = Y_1 \cap Y_2$):

Lemma 2.4. Let X be a flat, proper scheme of finite presentation over S. Let L, M be two invertible sheaves on X, and $\underline{\mathrm{Isom}}(L, M)$ the functor of isomorphisms $L_T \to M_T$ (for T/S). Then

i) $\underline{\mathrm{Isom}}(L, M)$ is representable by an affine scheme over S of finite presentation.

ii) If, moreover, L and M are equipped with rigidifications (relative to a [print 603]given section of X), the subfunctor of $\underline{\mathrm{Isom}}(L, M)$ formed by the isomorphisms compatible with these rigidifications is representable by a closed subscheme of $\underline{\mathrm{Isom}}(L, M)$.

Indeed, ii) is clear. As for i), recall that the functors $T \mapsto \mathrm{Hom}(L_T, M_T)$ and $T \mapsto \mathrm{Hom}(M_T, L_T)$ are representable by vector bundles $\mathbb{V}(E)$ and $\mathbb{V}(F)$ (by EGA III 7.7.8). Then, for $u \in \mathrm{Hom}(L_T, M_T)$ and $v \in \mathrm{Hom}(M_T, L_T)$, the conditions $u \circ v = 1$ and $v \circ u = 1$ define a closed subscheme F of $\mathbb{V}(E) \times_S \mathbb{V}(F)$[577], and it is clear that $F \simeq I$.

2.5. Finally, let us reduce assertion II' to the following assertion:

II*. Theorem 1.1 when X and Y are integral over S and the sequence [578]

$$O_Y \longrightarrow f_*O_X \rightrightarrows g_*(O_{X \times_Y X}) \qquad (*)$$

(where $g : X \times_Y X \to Y$[579] is the structural morphism) is exact.

In fact, for this it suffices to prove that there is a factorization of f

$$X \xrightarrow{f_0} Y_0 \xrightarrow{f_1} Y_1 \longrightarrow \cdots \xrightarrow{f_\eta} Y_\eta = Y$$

such that every f_i satisfies (*) and, after shrinking S, every Y_i is integral over S.

For this, put $Y_0 = \mathrm{Spec}(f_*(\underline{O}_X))$, with f_0 arising from the Stein factorization of f (EGA III 4.3.3), so that $f_{0*}(\underline{O}_X) \xleftarrow{\sim} \underline{O}_{Y_0}$, and hence f_0 does satisfy (*). To factor $Y_0 \to Y$ in the same way, we use the

Lemma 2.6. Let Y be a noetherian scheme, and $f : X \to Y$ a finite epimorphism (i.e. such that $O_Y \to f_*(O_X)$ is injective); then f factors into a finite sequence of finite morphisms f_i $X \simeq Y_0 \xrightarrow{f_1} Y_1 \longrightarrow \cdots \longrightarrow Y_n = Y$ that are effective epimorphisms (i.e. such that the sequence [print 604]$O_{Y_i} \longrightarrow (f_i)_*(O_{Y_{i-1}}) \rightrightarrows (g_i)_*(O_{Y_{i-1}} \otimes_{O_{Y_i}} O_{Y_{i-1}})$ is exact).

Put $A = O_Y$, $B_0 = f_*(O_X)$, and define $B_i = \mathrm{Ker}(B_{i-1} \rightrightarrows B_{i-1} \otimes_A B_{i-1})$ recursively. We verify at once that $B_{i-1} \otimes_A B_{i-1} \xrightarrow{\sim} B_{i-1} \otimes_{B_i} B_{i-1}$[580]. It remains to prove that, for i sufficiently large, $B_i = A$. Indeed, consider the decreasing sequence of A-modules $M_i = B_i/A$[581], and put $S_i = \mathrm{Supp}(M_i)$. The decreasing sequence of closed subsets S_i of Y is stationary; let S be the stationary value. Let us show that $S = \varnothing$. Otherwise, let s be a maximal point of S. After replacing Y by $\mathrm{Spec}\, O_{Y,s}$, we may suppose that Y is local with closed point s. By the choice of s, M_i is artinian for i sufficiently large, so the sequence (M_i) is stationary. Thus there exists i such that $B_i = B_{i+1}$. Consequently, $\mathrm{Spec}(B_i) \to Y$[582] is a monomorphism, hence a closed immersion (Nakayama); since $A \subset B_i$[583], we have $A = B_i$[584] and $s \notin S$, a contradiction.

3. Proof of I: Oort's dévissage

Proposition 3.1. (Oort, Bull. soc. math. France, vol. 90-1962). Let S be a locally noetherian scheme, $f : X \to S$ a proper morphism, and $X' \subset X$ a closed subscheme. Suppose that:

i) X' is defined by an ideal I contained in the nilradical N of X and $N.I = 0$.

ii) X' is flat over S and cohomologically flat over S in dimension 0 (EGA III 7.8.1).

iii) I is flat over S and cohomologically flat over S in dimensions 0 and 1, so that (by EGA III 7.4.6) there are a coherent O_S-module Q and an isomorphism, functorial in the coherent O_S-module M, [print 605]

$$R^2 f_*(I \otimes f^*M) \simeq \underline{\mathrm{Hom}}_{O_S}(Q, M).$$

iv) The cokernel E of $f_*(O_{X'}) \to R^1 f_*(I)$ is a locally free O_S-module.

v) The homomorphism $(f_*O_X)_{\mathrm{red}} \longrightarrow (f_*O_{X'})_{\mathrm{red}}$ is surjective.

Then the canonical morphism $\underline{\mathrm{Pic}}_{X/S} \longrightarrow \underline{\mathrm{Pic}}_{X'/S}$ is representable by an affine morphism of finite presentation; more precisely, there is a canonical exact sequence of f.p.p.f. sheaves,

$$0 \longrightarrow \mathbb{V}(E^\vee) \longrightarrow \underline{\mathrm{Pic}}_{X/S} \longrightarrow \underline{\mathrm{Pic}}_{X'/S} \xrightarrow{u} \mathbb{V}(Q)$$

and, for $P = \mathrm{Ker}(u)$, the monomorphism $P \hookrightarrow \mathrm{Pic}_{X'/S}$ is representable by a closed immersion and $\mathrm{Pic}_{X/S}$ is a torsor over P under the group $V(\check{E})$ (that is, the sequence of f.p.p.f. sheaves

$$0 \longrightarrow \mathbb{V}(\check{E}) \longrightarrow \mathrm{Pic}_{X/S} \longrightarrow P \longrightarrow 0$$

is exact).

For an S-scheme T, consider the following two diagrams with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I_T & \longrightarrow & O_{X_T} & \longrightarrow & O_{X'_T} & \longrightarrow & 0 \\ & & \| & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & I_T & \xrightarrow{\alpha} & O^*_{X_T} & \longrightarrow & O^*_{X'_T} & \longrightarrow & 0 \end{array} \qquad (\alpha(i) = 1 + i)$$

[585]

$$\begin{array}{ccc} (f_T)_*(O_{X'_T}) & \xrightarrow{\delta_T} & R^1(f_T)_*(I_T) \\ \uparrow & & \| \\ (f_T)_*(O^*_{X'_T}) & \xrightarrow{\delta^*_T} & R^1(f_T)_*(I_T) \to R^1(f_T)_*(O^*_{X_T}) \to R^1(f_T)_*(O^*_{X'_T}) \to R^2(f_T)_*(I_T). \end{array}$$

Observe carefully that the square on the left of this diagram is not necessarily commutative. Nevertheless, we have:

$$3.2. \qquad \mathrm{Im}\,\delta^*_T = \mathrm{Im}\,\delta_T.$$

[print 606]Indeed, we may clearly suppose that S and T are affine. Take an affine covering $\{V_i\}$ of X_T. Let $g \in \Gamma(X_T; O^*_{X'_T})$, and let g_i be a lift of $g|V_i$ in $\Gamma(V_i; O^*_{X_T})$. Then $\delta^*_T(g)$ is the class of the 1-cocycle $\delta^*_{ij} = (g_i g_j^{-1} - 1)$; $\delta_T(g)$ is the class of $\delta_{ij} = (g_i - g_j) = g_j \delta^*_{ij} = g\,\delta^*_{ij}$, so we have

$$\delta_T(g) = g\,\delta^*_T(g)$$

taking into account the action of $\Gamma(X_T, \underline{O}_{X'_T})$ on I_T, hence on $H^1(X_T, I_T)$.

By ii), we have $\Gamma(X_T; O_{X'_T}) = \Gamma(X; O_{X'}) \otimes_S \Gamma(T, O_T)$, and hence $g = \sum g_\beta \otimes t_\beta$ for $g_\beta \in \Gamma(X, O_{X'})$ and $t_\beta \in \Gamma(T, O_T)$. Now, by (v), $(f_* O_X)_{\mathrm{red}} \longrightarrow (f_* O_{X'})_{\mathrm{red}}$ is surjective; consequently $g_\beta = v(h_\beta) + n_\beta$ for $h_\beta \in \Gamma(X; O_X)$, $v : \Gamma(X, O_X) \to \Gamma(X, O_{X'})$ and $n_\beta \in \mathrm{nilrad}(\Gamma(X, O_{X'}))$.

By i), $(n_\beta \otimes t_\beta) I_T = 0$. Consequently, $\delta_T(g) = h\, \delta_T^*(g)$, where $h = \sum h_\beta \otimes t_\beta \in \Gamma(X_T, \underline{O}_{X_T})$[586], whence $\delta_T^*(g) = h^{-1} \delta_T(g) = \delta_T(h^{-1} g)$ and hence $\mathrm{Im}\, \delta_T^* \subset \mathrm{Im}\, \delta_T$.

Finally, let $g = \sum g_\beta \otimes t_\beta \in \Gamma(X_T, O_{X'_T})$[587]. Let $g_\beta = v(h_\beta) + n_\beta$[588], where $n_\beta \in \mathrm{nilrad}(\Gamma(X, O_{X'}))$; we then[589] have $\delta_T(g) = \delta_T(\sum n_\beta \otimes t_\beta) = \delta_T(1 + \sum n_\beta \otimes t_\beta) = \delta_T^*(1 + \sum n_\beta \otimes t_\beta)$, whence $\mathrm{Im}\, \delta_T \subset \mathrm{Im}\, \delta_T^*$.

By 3.2, iii) and iv), there is an exact sequence functorial in T: [590]

$$0 \longrightarrow \mathbb{V}(E^\vee)(T) \longrightarrow R^1(f_T)_*(O^*_{X_T}) \longrightarrow R^1(f_T)_*(O^*_{X'_T}) \longrightarrow \mathbb{V}(Q)(T),$$

which, on passing to the associated f.p.p.f. sheaves, gives an exact sequence

$$0 \longrightarrow \mathbb{V}(E^\vee) \longrightarrow \underline{\mathrm{Pic}}_{X/S} \longrightarrow \underline{\mathrm{Pic}}_{X'/S} \xrightarrow{u} \mathbb{V}(Q).$$

For $P = \mathrm{Ker}(u)$, the monomorphism $P \hookrightarrow \underline{\mathrm{Pic}}_{X'/S}$ is obtained by base change from the unit section $S \hookrightarrow \mathbb{V}(Q)$; since $\mathbb{V}(Q)$ is representable and separated, it follows that $P \hookrightarrow \mathrm{Pic}_{X'/S}$ is representable by a [print 607]closed immersion.

Finally, $0 \longrightarrow \mathbb{V}(E^\vee) \longrightarrow \underline{\mathrm{Pic}}_{X/S} \xrightarrow{w} P \longrightarrow 0$ is a torsor for f.p.p.f.; a fortiori, by f.p.q.c. descent from a "splitting" of the torsor, w is representable by a faithfully flat affine morphism of finite presentation (see: Oort, "Commutative group schemes", Proposition (17.4)).

Corollary 3.3. Under the conditions of 3.1, we have

i) If $\underline{\mathrm{Pic}}_{(\mathrm{zar})(X'/S)} \to \underline{\mathrm{Pic}}_{(\mathrm{fppf})(X'/S)}$ is an isomorphism, then $\mathrm{Pic}_{(\mathrm{zar})(X/S)} \to \mathrm{Pic}_{(\mathrm{fppf})(X/S)}$ is an isomorphism.

ii) Supposing that the S-scheme T is affine, if $\mathrm{Pic}(X'_T) \to \underline{\mathrm{Pic}}^{(T)}_{(\mathrm{zar})(X'/S)}$ is surjective, then $\mathrm{Pic}(X_T) \to \underline{\mathrm{Pic}}^{(T)}_{(\mathrm{zar})(X/S)}$ is surjective.

Indeed, (i) follows from the following diagram of sheaves for the Zariski topology

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \mathbb{V}(E^\vee) & \longrightarrow & \underline{\mathrm{Pic}}_{(\mathrm{zar})(X/S)} & \longrightarrow & \underline{\mathrm{Pic}}_{(\mathrm{zar})(X'/S)} & \longrightarrow & \mathbb{V}(Q) \\
\downarrow & & \downarrow S & & \downarrow & & \downarrow S & & \downarrow S \\
0 & \longrightarrow & \mathbb{V}(E^\vee) & \longrightarrow & \underline{\mathrm{Pic}}_{(\mathrm{fppf})(X/S)} & \longrightarrow & \underline{\mathrm{Pic}}_{(\mathrm{fppf})(X'/S)} & \longrightarrow & \mathbb{V}(Q),
\end{array}$$

by applying the five lemma to it.

(ii) follows from the following diagram with exact rows

$$\begin{array}{ccccccc}
H^1(X_T, I_T) & \longrightarrow & H^1(X_T, O^*_{X_T}) & \longrightarrow & H^1(X_T, O^*_{X'_T}) & \longrightarrow & H^2(X_T, I_T) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
H^0(T, R^1(f_T)_*(I_T)) & \longrightarrow & H^0(T, R^1(f_T)_* O^*_{X_T}) & \longrightarrow & H^0(T, R^1(f_T)_* O^*_{X'_T}) & \longrightarrow & H^0(T, R^2 f_* I_T)
\end{array}$$

because the outer vertical arrows are isomorphisms, T being affine.

[print 608]**Corollary 3.4.** Let k be a field, and X a proper k-scheme. Suppose that $\underline{\mathrm{Pic}}_{X_{\mathrm{red}}/k}$ possesses a universal invertible sheaf. (For example, by descent theory this holds when every connected component X'_i of X_{red} has a rational point, for example when k is algebraically closed, or more generally when the g.c.d. of the degrees of the 0-cycles on every X'_i is 1). Then, over every affine open subset V of $\mathrm{Pic}_{X/k}$, there is a universal invertible sheaf on $X \times_k V$.

Finally, we prove I in the following more precise form:

Proposition 3.5. Let S be an integral noetherian scheme, and $f : X \hookrightarrow Y$ a nilpotent immersion of proper S-schemes. Then there is a nonempty open subset V of S such that the morphism

$$f^* : \mathrm{Pic}_{Y/S}\,|V \longrightarrow \mathrm{Pic}_{X/S}\,|V$$

is representable by an affine morphism of finite presentation; in fact, $N = \mathrm{Ker}\, f^*$ is an affine group scheme with unipotent fibers (in the sense of SGA 3 XVII).

Indeed, X is defined by a nilpotent ideal I. Then I has a filtration by ideals $I_0 = I \supset I_1 \supset \ldots \supset I_n = 0$ such that $NI_{i-1} \subset I_i$[591], where N is the nilradical of O_Y (for example $I_k = N^k I$). Let X_i be the subscheme of Y defined by I_i. By induction it suffices to study the immersions $X_{i-1} \hookrightarrow X_i$[592]. We may therefore suppose that $N.I = 0$; this is condition i) of 3.1, and we finish by reducing to the case where the other conditions (ii) to (v) also hold. By the generic flatness theorem (EGA IV 6.9.1), applied to $\underline{O}_{X'}$, to the $R^i f_*(I)$ and to E, and taking account of the Künneth relations EGA III 6.9.9, we may suppose, after shrinking S, that conditions ii), iii) and iv) hold.

[print 609]There is a finite extension K of $k(S)$ such that $(f_K)_*(O_{X'_K})$ has all its residue field extensions trivial. For an integral scheme S' finite and flat over an open subset V of S, with function field $k(S') = K$, there is an open subset V' of S' such that

$$\left[(f_{S'})_*(O_{X'_{S'}})\right]_{\mathrm{red}} \simeq \left[(f_{S'})_*(O_{X_{S'}})\right]_{\mathrm{red}}$$

over V'; after shrinking S and S', we may suppose that $S = V$ and $S' = V'$. Since the assertion to be proved is local for f.p.q.c., we may replace S by S'. Finally, apply 3.1.

3.6. Remark. It follows in the same way, taking into account the exact sequences $\underline{\mathrm{Pic}}_{X_i/S} \longrightarrow \underline{\mathrm{Pic}}_{X_{i-1}/S} \longrightarrow \mathbb{V}(Q_i)$[593], that, for every $s \in S$,

$$\mathrm{coker}\left(\underline{\mathrm{Pic}}^0_{Y_s/s} \longrightarrow \underline{\mathrm{Pic}}^0_{X_s/s}\right)$$[594]

is a unipotent group.

4. Proof of II*: the most delicate part of the proof

Recall assertion II*: Theorem 1.1 when X and Y are integral over S and the sequence (*) $O_Y \longrightarrow f_*O_X \rightrightarrows g_*(O_{X\times_Y X})$ is exact; applying the generic flatness theorem EGA IV 6.9.1 and EGA III 7.8.8 reduces us to proving the

Proposition 4.1. Let S be an integral noetherian scheme, $\lambda : X \to S$ and $\mu : Y \to S$ proper morphisms of finite presentation, $f : X \to Y$ a surjective S-morphism, and $g : X \times_Y X \to Y$ the canonical morphism. Suppose that

1) $X, Y, X \times_Y X$ are flat over S.

2) $\lambda_* O_X = O_S$, $\mu_* O_Y = O_S$ universally. [print 610]

3) For $i, j \geq 0$, $R^i f_* O_X$[595] and $R^j g_* O_{X\times_Y X}$ are flat over S.

4) The sequence

$$O_Y \longrightarrow f_*O_X \overset{\pi_1}{\underset{\pi_2}{\rightrightarrows}} g_*O_{X\times_Y X} \qquad (*)$$

is exact and $\operatorname{coker}(\pi_1 - \pi_2)$ is an O_Y-module flat over S.

Then $f^* : \mathrm{Pic}_{Y/S} \longrightarrow \mathrm{Pic}_{X/S}$ is representable by a quasi-affine morphism of finite presentation.

Indeed, first note that 1)-4) are stable under every base change $T \to S$. Only the stability of 3) and 4) is not evident. But, by the Künneth formula (EGA III 6.9.9), 1) and 3) imply that the formation of the $R^i f_*(O_X)$ and $R^i g_*(O_{X\times_Y X})$ commutes with base changes $T \to S$, which immediately implies the stability of conditions 3). Moreover, the formation of the cokernel of $\pi_1 - \pi_2 : f_*O_X \longrightarrow g_*O_{X\times_Y X}$ therefore also commutes with base changes; since it is flat over S (by 4)), $\mathrm{Im}(\pi_1 - \pi_2)$ is flat over S and commutes with base changes. Consequently, $\mathrm{Ker}(\pi_1 - \pi_2)$ commutes with base changes $T \to S$ and is therefore universally equal to O_Y.

The stability of the hypotheses implies two things: a. Since the assertion to be proved is local for f.p.p.f., by descent we may suppose that X has a section α, so Y has the section $\beta = f \circ \alpha$. b. To establish the assertion it suffices to consider an S-point of $\mathrm{Pic}_{X/S}$, that is, by a) and 2), an invertible sheaf L on X equipped with a rigidification u relative to α, and to show that the f.p.p.f. sheaf Z, the inverse image of this point in $\mathrm{Pic}_{Y/S}$, is representable by a quasi-affine S-scheme of finite presentation over S.

[print 611]**Lemma 4.2.** Under the conditions of 4.1, f is a descent morphism for the category of locally free sheaves; that is, for every pair L, M of locally free sheaves on Y, the diagram

$$\mathrm{Hom}_Y(L, M) \to \mathrm{Hom}_X(f^*L, f^*M) \overset{\to}{\to} \mathrm{Hom}_{X\times_Y X}(g^*L, g^*M)$$

is exact.

Indeed, for $N = \underline{\mathrm{Hom}}_{O_Y}(L, M)$, we must show that the sequence [596]

$$\Gamma(Y, N) \longrightarrow \Gamma(X, f^*N) \rightrightarrows \Gamma(X \times_Y X, g^*N)$$

is exact, or (by EGA 0_1 5.4.10.1) that the sequence

$$\Gamma(Y, N) \longrightarrow \Gamma(Y, f_*O_X \otimes_{O_Y} N) \rightrightarrows \Gamma(Y, g_*O_{X\times_Y X} \otimes_{O_Y} N)$$

is exact. But the sequence (*) $O_Y \to f_*O_X \rightrightarrows g_*O_{X\times_Y X}$ is exact by hypothesis; the lemma follows by applying to it the functor $\Gamma(N \otimes -)$, which is left exact, N being flat over Y.

Let L be an invertible sheaf on X equipped with a rigidification u relative to α; by 4.2 there is an equivalence between the category C of pairs (M, φ) consisting of an invertible sheaf M on Y and an isomorphism $\varphi : f^*M \xrightarrow{\sim} L$, and the category $\mathrm{Deff}(S)$ of effective descent data on L relative to f. But, for $f^*M \simeq L$, since L is rigidified relative to α, M is canonically equipped with a rigidification relative to β. Consequently, C is equivalent to the discrete category defined by the set of classes of β-rigidified invertible sheaves (M, v) on Y such that $f^*(M, v) \simeq (L, u)$. But this is precisely the set $Z(S)$ of S-points of the inverse image in $\mathrm{Pic}_{Y/S}$ of the S-point of $\mathrm{Pic}_{X/S}$ defined by (L, u). Moreover, this description of $Z(S)$ is functorial in S. Thus Z is identified with the functor Deff, and we may subsequently forget the rigidifications.

[print 612]

There is a canonical monomorphism from Deff into the functor Rec of gluing data on L[597]. By definition, Rec is the functor $\underline{\mathrm{Isom}}(p_1^*L, p_2^*L)$, where the $p_i : X \times_Y X \to X$ are the projections, which is represented by an affine scheme I of finite presentation over S by 2.4. It therefore remains to represent the monomorphism

$$i : \mathrm{Deff} \hookrightarrow \underline{\mathrm{Isom}}(p_1^*L, p_2^*L)$$

by a morphism of finite presentation, which will necessarily be quasi-affine (by EGA IV 8.11.2).

Observe in passing that, if desired, we may restrict $\underline{\mathrm{Isom}}(p_1^*L, p_2^*L)$ by imposing the cocycle condition, but this is not indispensable for what follows.

After the base change $I \to S$, we are therefore reduced to the following problem: for an invertible sheaf L on X equipped with a gluing datum $\varphi : p_1^*L \xrightarrow{\sim} p_2^*L$, represent the subfunctor of S "that makes this datum effective". But we have the

Lemma 4.3. Under the conditions of 4.1, let L be an invertible sheaf on X equipped with a gluing datum

$$\varphi : p_1^*L \xrightarrow{\sim} p_2^*L.$$

For this datum to be effective, it is necessary and sufficient that the following five conditions hold.

1) For every $i \geq 0$, $R^i f_*(L)$ is flat over S.

2) For every $j \geq 0$, $R^j g_*(p_1^*L)$ is flat over S.

3) The cokernel of $\pi_1 - \pi_2 : f_*(L) \longrightarrow g_*(p_1^*L) \simeq g_*(p_2^*(L))$ is flat over S, (where the last isomorphism is $g_*(\varphi)$).

4) The kernel N of $\pi_1 - \pi_2 : f_*(L) \to g_*(p_1^*L)$ is invertible on Y.

5) The composite morphism $f^*N \to f^*f_*L \to L$ is an isomorphism.

[print 613]Indeed, these conditions are sufficient, since already 4) and 5) imply effectivity. Conversely, let M be invertible on Y, with an isomorphism $f^*M \simeq L$ compatible with the gluing datum. We may cover Y by open subsets V over which M is isomorphic to O_Y; hence the descent datum on L restricted over $V = f^{-1}(V)$ is isomorphic to the trivial descent datum on $O_X|V$. Finally, conditions 1)-5) are local on Y and hold, by the hypotheses of 4.1, for $L = O_X$; they therefore hold for L.

It therefore remains to represent each of conditions 1) to 5) successively by monomorphisms of finite presentation. For 1) and 2), we apply the

Lemma 4.4. Let S be a scheme, $\lambda : X \to S$ and $\mu : Y \to S$ proper morphisms of finite presentation, and $f : X \to Y$ an S-morphism. Let F be a sheaf on X of finite presentation and flat over S and, for an integer n, let S_n be the subfunctor of S that makes $R^i f_*(F)$ flat over S for every $i \geq n$ (more precisely, $S_n(T) = \{\varnothing\}$ when $R^i(f_T)_*(F_T)$ is flat for every $i \geq n$, and $S_n(T) = \varnothing$ otherwise). Then $S_n \hookrightarrow S$ is representable by a surjective monomorphism of finite presentation, and then, for $i \geq n$, the $R^i(f_{S_n})_*(F_{S_n})$ commute with base changes $T \to S_n$. Moreover, S_n coincides with the functor S'_n that makes the functors

$$M \longmapsto R^i(f_V)_*(F_V \otimes \lambda_V^* M) \qquad \text{for } i \geq n$$

exact, where M runs through the quasi-coherent O_V-modules for every affine open subset V of S.

Indeed, the assertion is local on S; we may therefore suppose that S is quasi-compact. The relative dimension of X/Y is then bounded, say by r, and, for $n > r$, $R^n f_*(F_V \otimes \lambda_V^* M) = 0$. The assertion therefore holds for $n > r$; [print 614]proceeding by descending induction, suppose it is known for $n+1$. After replacing S by $S_{n+1} = S'_{n+1}$, we may suppose that $R^i f_*(F)$ is S-flat[598] for $i \geq n+1$. Since F is flat over S, it then follows from EGA III 6.9.8 that $R^n f_*(F)$ commutes with every base change. Consequently, the representability of S_n follows from the next lemma (see: J.P. Murre, Representation of unramified functors, Séminaire Bourbaki; May 1965, p.294-11, th.2):

Lemma 4.5. Let $X \to S$ be a proper morphism of finite presentation, and F a sheaf of finite presentation on X. Then the subfunctor S_F of S that makes F flat over S is representable by a surjective monomorphism of finite presentation (which is of course not an immersion in general).

To represent condition 3) of 4.3, note that, by 1) and 2), $f_*(L)$ and $g_*(p_1^*L)$ commute with base changes $I \to S$, and hence $\operatorname{coker}(\pi_1 - \pi_2)$ does also; then apply 4.5.

To represent 4), note that, by 3) and the fact that $f_*(L)$ and $g_*(p_1^*L)$ are flat and commute with base changes, $N = \ker(\pi_1 - \pi_2)$ also commutes; then apply the

Lemma 4.6. Let $\lambda : X \to S$ be a proper morphism of finite presentation, and $u : F \to G$ a homomorphism of O_X[599]-Modules of finite presentation, with G flat over S. Then the subfunctor S_1 of S that makes u an isomorphism is representable by an open subscheme of S, of finite presentation.

As usual, we reduce to the case where S is noetherian, using EGA IV 8 and 11.2.6, which relieves us of the need to consider finite presentation. First let us represent the functor that makes u an epimorphism, [print 615]i.e. makes $\operatorname{Coker} u$ zero. Since the formation of $\operatorname{Coker} u$ commutes with base change[600], we see that it is represented by the open subset $S - \lambda(\operatorname{supp}(\operatorname{Coker} u))$ of S. We may therefore suppose that u is already an epimorphism. But, now using the hypothesis that G is S-flat, we find that the formation of $\operatorname{Ker} u$ commutes with base change, and consequently the functor under consideration is represented by the open subset $S - \lambda(\operatorname{supp}(\operatorname{Ker} u))$ of S.

[print 616]

EXPOSÉ XIII
FINITENESS THEOREMS FOR THE PICARD FUNCTOR

by S. KLEIMAN

1. The (b)-sheaves

Let k be an algebraically closed field and X a projective k-scheme, endowed with an ample invertible module $O_X(1)$.

Definition 1.1 (Mumford). Let $m \in \mathbb{Z}$. A coherent sheaf F on X is said to be m-regular if, at every point of the support of F, $O_X(1)$ is generated by its global sections and if, for every $q \geq 1$, we have

$$H^q(F(m-q)) = 0.$$

Lemma 1.2. Let $0 \longrightarrow F(-1) \longrightarrow F \longrightarrow G \longrightarrow 0$ be an exact sequence of coherent sheaves on X(*). If F is m-regular, then so is G.

Indeed, we have the corresponding exact sequence:

$$H^q(F(m-q)) \longrightarrow H^q(G(m-q)) \longrightarrow H^{q+1}(F(m-q-1)).$$

Proposition 1.3. If a sheaf F on X is m-regular, then, for every $n \geq m$, we have :

(i) F is n-regular.

(ii) $H^0(F(n)) \otimes H^0(O_X(1)) \longrightarrow H^0(F(n+1))$ is surjective.

(iii) $F(n)$ is generated by $H^0(F(n))$.


[print 617]The proof is by induction on the dimension s of the support of F; when $s = -1$, it is trivial. Take a section $\sigma \in H^0(O_X(1))$ which does not vanish at any point of the finite set $\mathrm{Ass}(F)$; then σ defines an exact sequence $0 \longrightarrow F(-1) \longrightarrow F \longrightarrow G \longrightarrow 0$, and G is m-regular by 1.2., with support of dimension $s-1$.

Consider the corresponding exact sequence:

$$H^q(F(n-q-1)) \longrightarrow H^q(F(n-q)) \longrightarrow H^q(G(n-q)).$$

By the induction hypothesis, we have $H^q(G(n-q)) = 0$ for $n \geq m$, so $H^q(F(n-q-1)) = 0$ implies $H^q(F(n-q)) = 0$; but $H^q(F(m-q)) = 0$, which gives (i).

To establish (ii), we chase through the following commutative diagram: [601]



(*) Here, as below in analogous situations, it is understood that $F(-1) \to F$ is defined by tensoring with a given section of $O_X(1)$.

where the oblique arrow is defined by $\varphi \mapsto \varphi \otimes \sigma$, the horizontal rows are exact, the first by the n-regularity of F and by (i), already proved, and the second column is exact by the induction hypothesis.

Finally, (iii) follows from the following commutative diagram:

$$\begin{array}{ccc} H^0(F(n))_X \otimes H^0(O_X(1))_X & \xrightarrow{\alpha_n} & H^0(F(n)(1))_X \\ \downarrow & & \downarrow{\scriptstyle \beta_{n+1}} \\ [H^0(F(n))_X](1) & \xrightarrow{\beta_n \otimes \mathrm{id}} & F(n)(1) \end{array}$$

[print 618]where, if M is a vector space over k, we write M_X for $M \otimes_k O_X$. For $n \geq m$, α_n is surjective by (ii); therefore, if β_{n+1} is surjective, so is β_n. By Serre's theorem, β_{n+1} is surjective for $n \gg 0$; hence (iii).

Proposition 1.4. Let $0 \to F(-1) \to F \to G \to 0$ be an exact sequence on X. If G is m-regular, then we have :

(i) $H^q(F(n)) = 0$ for $q \geq 2$ and $n \geq m - q$.

(ii) $h^1(F(n-1)) \geq h^1(F(n))$ for $n \geq m - 1$.

(iii) $H^1(F(n)) = 0$ for $n \geq (m-1) + h^1(F(m-1))$.

In particular, F is $[m + h^1(F(m-1))]$-regular.

Indeed, in the exact sequence

$$H^{q-1}(G(n)) \longrightarrow H^q(F(n-1)) \longrightarrow H^q(F(n)) \longrightarrow H^q(G(n)),$$

the two outer groups are 0 for $q \geq 2$ and $n \geq m - (q-1)$ by 1.3 (i).[602] Thus $H^q(F(m-q)) \xrightarrow{\sim} H^q(F(m-q+1)) \xrightarrow{\sim} H^q(F(m-q+2)) \xrightarrow{\sim} \ldots$, but, by Serre's theorem, $H^q(F(n)) = 0$ for $n \gg 0$; hence (i).

For $n \geq m - 1$, consider the following exact sequence:

$$0 \to H^0(F(n-1)) \to H^0(F(n)) \xrightarrow{\alpha_n} H^0(G(n)) \to H^1(F(n-1)) \to H^1(F(n)) \to 0$$

It implies (ii). Now suppose that α_n is surjective and consider the following commutative diagram:

$$\begin{array}{ccccc} H^0(F(n)) \otimes H^0(O_X(1)) & \xrightarrow{\alpha_n \otimes \mathrm{id}} & H^0(G(n)) \otimes H^0(O_X(1)) & \longrightarrow & 0 \\ \downarrow & & \downarrow{\scriptstyle \beta_n} & & \\ H^0(F(n+1)) & \xrightarrow{\alpha_{n+1}} & H^0(G(n+1)). & & \end{array}$$

[print 619]If $n \geq m$, then β_n is surjective by 1.3 (ii);[603] therefore α_{n+1} is surjective. Consequently, $H^1(F(n-1)) \xrightarrow{\sim} H^1(F(n)) \xrightarrow{\sim} \ldots \xrightarrow{\sim} 0$. On the other hand, if α_n is not surjective, then $h^1(F(n-1)) > h^1(F(n))$; therefore $n \mapsto h^1(F(n))$[604] is strictly decreasing from $h^1(F(m-1))$ until it becomes 0; hence (iii).

Definition 1.5. Let F be a coherent sheaf on X, r an integer $\geq \dim.\mathrm{supp}(F)$ and $(b) = (b_0, \ldots, b_r)$ a sequence of $r+1$ integers. Then F is said to be a (b)-sheaf if it satisfies one of the following two equivalent sets of conditions :

(1.5.1)

(i) At every point of the support of F, $O_X(1)$ is generated by $H^0(O_X(1))$.

(ii) $h^0(F(-1)) \leq b_0$.

(iii) (When $r \geq 1$), there exists a section $\sigma \in H^0(O_X(1))$ defining an exact sequence $0 \to F(-1) \to F \to G \to 0$ such that G is a $(b_1, \ldots, b_r)$-sheaf.

(1.5.2) There exists an F-regular sequence of r sections $\sigma_1, \ldots, \sigma_r \in H^0(O_X(1))$ such that, for $i = 0, \ldots, r$, if F_i denotes the restriction of F to the zero scheme of $\sigma_1, \ldots, \sigma_i$, then $h^0(F_i(-1)) \leq b_i$.

Proposition 1.6. Let F be a (b)-sheaf on X. Then :

(i) For every "sufficiently general" sequence σ of r sections $\sigma_1, \ldots, \sigma_r \in H^0(O_X(1))$, if $F_{\sigma,i}$ denotes the restriction of F to the zero scheme of $\sigma_1, \ldots, \sigma_i$, then $h^0(F_{\sigma,i}(-1)) \leq b_i$, where $0 \leq i \leq r$.

(ii) Every subsheaf G of F is a (b)-sheaf.

[print 620]Indeed, let S' be the affine space $\mathbb{A}_k^N$ whose k-points correspond to the sequences σ, and T the open subset of S'[605] corresponding to the F-regular sequences. For fixed i, the sheaves $F_{\sigma,i}(-1)$ corresponding to the k-points of T are contained in a flat family over T, and by hypothesis T is nonempty. Thus (i) follows from the upper semi-continuity property of the function $\sigma \longmapsto h^0(F_{\sigma,i}(-1))$ (EGA III, 7.7.5). Finally, since every "sufficiently general" sequence is G-regular and such that $G_{\sigma,i}$ is a subsheaf of $F_{\sigma,i}$ for every i, (ii) follows from (i).

Lemma 1.7. Let $0 \to F(-1) \to F \to G \to 0$[606] be an exact sequence on X. If the Hilbert polynomial of F is written in the form

$$\chi(F(n)) = \sum_{i=0}^{r} a_i \binom{n+i}{i},$$

then

$$\chi(G(n)) = \sum_{i=0}^{r-1} a_{i+1} \binom{n+i}{i}.$$

Indeed, $\chi(G(n)) = \chi(F(n)) - \chi(F(n-1)) = \sum_{i=0}^{r} a_i \left[\binom{n+i}{i} - \binom{n-1+i}{i}\right]$; therefore $\chi(G(n)) = \sum_{i=1}^{r} a_i \binom{n+i-1}{i-1}$, which gives the assertion.

Proposition 1.8. Let F be a (b)-sheaf on X, $s = \dim.\mathrm{supp}(F)$ and $\chi(F(n)) = \sum_{i=0}^{s} a_i \binom{n+i}{i}$ the Hilbert polynomial of F. Then :

(i) For $n \geq -1$, $h^0(F(n)) \leq \sum_{i=0}^{s} b_i \binom{n+i}{i}$.

(ii) $a_s \leq b_s$ and F is also a $(b_0, \ldots, b_{s-1}, a_s)$-sheaf.

Indeed, we argue by induction on s. When $s = 0$, this is clear since $a_0 = h^0(F) = h^0(F(-1)) \leq b_0$. When $s \geq 1$, we have an exact sequence

$$0 \longrightarrow F(-1) \longrightarrow F \longrightarrow G \longrightarrow 0$$

[print 621]with G a $(b_1, \ldots, b_r)$-sheaf and $\dim.\mathrm{supp}(G) = s - 1$. Then

$$h^0(F(n)) - h^0(F(n-1)) \leq h^0(G(n)),$$

but

$$h^0(G(n)) \leq \sum_{i=0}^{s-1} b_{i+1} \binom{n+i}{i}$$

by induction on s and $h^0(F(-1)) \leq b_0$; hence (i) by induction on $n \geq -1$. Finally, by 1.7 and induction on s, $a_s \leq b_s$ and G is a $(b_1, \ldots, b_{s-1}, a_s)$[607]-sheaf; hence (ii).

Definition 1.9. The following polynomials with rational coefficients ≥ 0, defined by induction, are called the (b)-polynomials :

$$\begin{cases} P_{-1} = 0 \\ P_r(X_0, \ldots, X_r) = P_{r-1}(X_1, \ldots, X_r) + \sum_{i=0}^{r} X_i \binom{P_{r-1}(X_1, \ldots, X_r) - 1 + i}{i}. \end{cases}$$

Remark 1.10. For the (b)-polynomials, one sees at once by induction on $r \geq t$ that

$$P_r(X_0, \ldots, X_t, 0, \ldots, 0) = P_t(X_0, \ldots, X_t).$$

Theorem 1.11. (The main theorem on (b)-sheaves). Let F be a (b)-sheaf on X and $\chi(F(n)) = \sum_{i=0}^{r} a_i \binom{n+i}{i}$ the Hilbert polynomial of F; let $(c) = (c_0, \ldots, c_r)$ be a sequence of integers $\geq (b) - (a)$ and $m = P_r(c)$ the integer given by the r-th (b)-polynomial (1.9). Then $m \geq 0$ and F is m-regular. In particular, if $s = \dim.\mathrm{supp}(F)$, then F is $P_{s-1}(c_0, \ldots, c_{s-1})$-regular.

[print 622]Indeed, we argue by induction on r. When $r = 0$, this is clear since $m \geq 0$[608] and F is n-regular for every n. When $r \geq 1$, we have an exact sequence $0 \longrightarrow F(-1) \longrightarrow F \longrightarrow G \longrightarrow 0$ where G is a $(b_1, \ldots, b_r)$-sheaf. By 1.7 and the induction hypothesis, we see that if $n = P_{r-1}(c_1, \ldots, c_r)$, then $n \geq 0$ and G is n-regular. Thus, by 1.4, F is $[n + h^1(F(n-1))]$-regular and $h^q(F(n-1)) = 0$ for $q \geq 2$. Therefore, $h^1(F(n-1)) = h^0(F(n-1)) - \chi(F(n-1))$, which is $\leq \sum_{i=0}^{r} (b_i - a_i) \binom{n-1+i}{i}$ by 1.8 (i), since $b_i \geq 0$. Finally, F is $\left[n + \sum_{i=0}^{r} c_i \binom{n-1+i}{i}\right]$-regular by 1.3 (i); this gives the first assertion. The second follows from it by 1.8 (ii) and 1.10.

1.12. Let S be a noetherian scheme and X an S-scheme of finite type. Let $\mathcal{F}$ be a family of classes of coherent sheaves on the fibers of X/S; that is, for every point $s \in S$ and every extension K of $k(s)$, coherent sheaves F_K on the algebraic scheme X_K are given, and we say that an F_K and an $F'_{K'}$ belong to the same class if there exist $k(s)$-homomorphisms from K, K' into a common extension K'' of $k(s)$ such that $F_{K''} (= F_K \otimes_K K'')$ and $F'_{K''}$ are isomorphic on $X_{K''}$. Then the family $\mathcal{F}$ is said to be bounded by the coherent sheaf F on $X_T = X \times_S T$, where T is of finite type over S, if $\mathcal{F}$ is contained in the family of classes of the sheaves $F_{k(t)}$ with $t \in T$; we say that $\mathcal{F}$ is bounded if there exist T, F as above.

Suppose in addition that X/S is projective and endowed with a (relatively) ample invertible sheaf $O_X(1)$. Then $\mathcal{F}$ is said to be a (b)-family, for a sequence of integers $(b) = (b_0, \ldots, b_r)$, if every class in $\mathcal{F}$ is represented by an F_K with K algebraically closed, which is a (b)-sheaf.

[print 623]**Theorem 1.13.** Let S be a noetherian scheme and X a projective S-scheme endowed with an ample sheaf $O_X(1)$ whose induced sheaves $O_{X_s}(1)$, with $s \in S$, are generated by their $H^0(O_{X_s}(1))$; let $\mathcal{F}$ be a family of classes of coherent sheaves on the fibers of X/S. Then the following conditions are equivalent:

(i) $\mathcal{F}$ is bounded. Moreover, if all the $F_K \in \mathcal{F}$ are locally free of rank p, then one may suppose that $\mathcal{F}$ is bounded by a sheaf F on an X_T, with F locally free of rank p.

(ii) The set of Hilbert polynomials $\chi(F_K(n))$ of the $F_K \in \mathcal{F}$ is finite, and there exists a sequence of integers (b) such that $\mathcal{F}$ is a (b)-family.

(iii) The set of Hilbert polynomials $\chi(F_K(n))$ of the $F_K \in \mathcal{F}$ is finite, and there exists an integer m such that all the $F_K \in \mathcal{F}$ are m-regular.

(iv) The set of Hilbert polynomials $\chi(F_K(n))$ of the $F_K \in \mathcal{F}$ is finite, and $\mathcal{F}$ is contained in the family of classes of quotients of sheaves of the form E_K, where E is a coherent sheaf on an X_T. Moreover, one may suppose $T = S$ and E of the form $O_X(-m)^{\oplus M}$.

(v) $\mathcal{F}$ is contained in the family of classes of cokernels of homomorphisms of the form $E'_K \to E_K$, where E' and E are coherent sheaves on an X_T. Moreover, one may suppose $T = S$ and E' and E of the forms $O_X(-m')^{\oplus M'}$ and $O_X(-m)^{\oplus M}$.

Indeed, suppose that $\mathcal{F}$ is bounded by a sheaf F on an X_T; after decomposing T into pieces, using the generic flatness theorem (EGA IV 6.9.1), one may suppose F flat over T. Then, [print 624]the number of polynomials $\chi(F(n))$ is at most equal to the number of connected components of T (EGA III 7.9.5), and it is a small lemma that if, for a $t \in T$, F_t is locally free of rank p, then so is F above an open neighborhood of t in T (see EGA IV 11.3.10, where one takes $X = Y$, $f = \mathrm{id}$). Now it follows from the hypotheses and (EGA III 7.7.6) that, after decomposing T more finely, one can find a sequence σ of sections $\sigma_1, \ldots, \sigma_r \in H^0(O_X(1))$ which is F-regular. Thus (ii) follows from the upper semi-continuity property of the function $t \longmapsto h^0(F_{t,i}(-1))$ where, for $t \in T$ and $0 \leq i \leq r$, $F_{t,i}$ denotes the restriction of F_t to the zero scheme of $\sigma_1, \ldots, \sigma_i$ (EGA III 7.7.5).

The implication (ii) $\Rightarrow$ (iii) follows immediately from 1.11. The implication (iii) $\Rightarrow$ (iv) follows from 1.3 (iii) by taking $M = \max(\chi(F_K(m)))$[609] and $E = O_X(-m)^{\oplus M}$.

Suppose that $\mathcal{F}$ satisfies (iv); there exist exact sequences

$$0 \longrightarrow F'_K \longrightarrow E_K \longrightarrow F_K \longrightarrow 0$$

and the set of polynomials $\chi(F_K(n))$ is finite. By hypothesis, the family of classes of the E_K is bounded; thus, by the implication (i) $\Rightarrow$ (ii) already proved, the set of polynomials $\chi(E_K(n))$ is finite and there exists a sequence of integers (b) such that all the E_K (with K algebraically closed) are (b)-sheaves. Consequently, the set of polynomials $\chi(F'_K(n))$ is finite and, by 1.6, all the F'_K are (b)-sheaves. Thus, applying to the family of classes of the F'_K the implication (ii) $\Rightarrow$ (iv) already proved, we obtain (v).

Finally, suppose that $\mathcal{F}$ satisfies (v); let us prove that $\mathcal{F}$ is bounded. One may suppose E (resp. E') of the form $O_X(-m)^{\oplus M}$. Indeed, [print 625]by (i) $\Rightarrow$ (iv), take a surjection $L = O_{X_T}(-m)^{\oplus M} \longrightarrow E$. After decomposing T into pieces so that E becomes flat over T, the formation of the exact sequence $0 \longrightarrow I \xrightarrow{u} L \longrightarrow E \longrightarrow 0$ commutes with passage to the fibers. By (i) $\Rightarrow$ (iii) and (i) $\Rightarrow$ (iv), take m_1 large enough that all the $\mathrm{Ext}^1(O_{X_K}(-m_1), I_K) = H^1(I_K(m_1))$ vanish and that there exist surjections $L_1 = O_{X_T}(-m_1)^{\oplus M_1} \longrightarrow E'$ and $\alpha : L_2 = O_{X_T}(-m_1)^{\oplus M_2} \longrightarrow I$. Then the maps $\mathrm{Hom}(L_{1,K}, L_K) \longrightarrow \mathrm{Hom}(L_{1,K}, E_K)$ are surjective. Let $\beta : E'_K \to E_K$ be a homomorphism and let γ be the composition $L_{1,K} \to E'_K \to E_K$[610]; then γ comes from a homomorphism $\delta : L_{1,K} \to L_K$, and $(\delta, u_K \circ \alpha) : L_{1,K} \oplus L_{2,K} \longrightarrow L_K$ and $\beta : E'_K \to E_K$ have the same cokernel. Consequently, $\mathcal{F}$ is contained in the family of classes of cokernels of homomorphisms of the form $O_{X_K}(-m_1)^{\oplus(M_1+M_2)} \longrightarrow O_{X_K}(-m)^{\oplus M}$.

Again after decomposing T into pieces, one may suppose in addition that X_T (hence $\underline{\mathrm{Hom}}_{O_{X_T}}(E', E)$) and all the $R^q(f_T)_*\underline{\mathrm{Hom}}_{O_{X_T}}(E', E)$ (where $f_T : X_T \to T$ is the structural morphism) are flat over T, so that the formation of $G = (f_T)_*\underline{\mathrm{Hom}}_{O_{X_T}}(E', E)$ commutes with arbitrary base change $T' \to T$ (EGA III 6.9.9.2). Let $R = \mathbb{V}(G^*)$; then X_R canonically carries a "universal" exact sequence

$$E'_R \longrightarrow E_R \longrightarrow F \longrightarrow 0$$

and F bounds $\mathcal{F}$.

2. Several technical lemmas

2.1. Let k be a field and X a proper k-scheme. Let $L_1, \ldots, L_s$ be invertible sheaves on X, $c_1(L_i)$ the Chern class of L_i and Y a closed subscheme of X of dimension $\leq s$. Recall that we denote by [print 626]$\langle c_1(L_1) \ldots c_1(L_s) \cdot Y \rangle$ the intersection number, which can be calculated as the coefficient of the monomial $n_1 \ldots n_s$ in the Snapper polynomial $\chi(L_1^{\otimes n_1} \otimes \ldots \otimes L_s^{\otimes n_s} \otimes O_Y)$ (X 4.3.1).

2.2. From now on, we suppose that k is algebraically closed, that X is projective, endowed with a very ample invertible sheaf $H = O_X(1)$, and that X is integral, of dimension r.

Let L be an invertible sheaf on X. The integer $\langle c_1(L) \cdot c_1(H)^{r-1} \rangle$ is called the degree of L (relative to H) and is denoted by $d(L)$.

In what follows we consider an invertible sheaf L and the Hilbert polynomials: $\chi(L(n)) = \sum_{i=0}^{r} a_i \binom{n+i}{i}$ and $\chi(O_X(n)) = \sum_{i=0}^{r} e_i \binom{n+i}{i}$.

Lemma 2.3. We have:

(i) $a_r = e_r = d(H)$.

(ii) (If $r \geq 1$), $a_{r-1} - e_{r-1} = d(L)$.

Indeed, we obtain (i) and (ii) at once by taking $m = 0, 1$ in the following polynomial:

$$\chi(L^{\otimes m} \otimes H^{\otimes n}) = \sum_{i+j \leq r} f_{ij} \binom{m+i}{i} \binom{n+j}{j}.$$

Lemma 2.4. If $d(L) \leq 0$ and $h^0(L) \neq 0$, then $L \simeq O_X$; therefore $d(L) = 0$ and $h^0(L) = 1$.

Indeed, since X is integral, there exists a divisor $Y \geq 0$ such that $L \simeq O_X(Y)$. If $Y > 0$, we would have $0 \geq \langle c_1(L) \cdot c_1(H)^{r-1} \rangle = \langle c_1(H)^{r-1} \cdot Y \rangle \geq 1$[611].

Lemma 2.5. If $d(L) \leq 0$, then L is a $(0, \ldots, 0, a_r)$-sheaf (with r zeros)[612].

[print 627]Indeed, if $r = 0$, L is evidently an a_0-sheaf. If $r \geq 1$, by 2.4, $h^0(L(-1)) = 0$ since $d(L(-1)) < 0$. Let Y be a hyperplane section of X and let $0 \longrightarrow L(-1) \longrightarrow L \longrightarrow L_Y \longrightarrow 0$ be an exact sequence defined by Y. If $r = 1$, by 1.7, $h^0(L_Y(-1)) = a_1$; thus L is a $(0, a_1)$-sheaf. If $r \geq 2$, one can take Y integral by Bertini's theorem and one may thus suppose by induction that L_Y is a $(0, \ldots, a_r)$-sheaf; this gives the assertion.

Lemma 2.7. If $r = 2$ and $d(L) \leq 0$, then:

(i) For every n, $h^1(L(n)) \leq a_1(a_1 + 1)a_2 - a_0$

(ii) $a_0 \leq a_1(a_1 + 1)$.

Let Y be an integral hyperplane section of X. By 1.7 and 2.5,[613] L_Y is a $(0, a_2)$-sheaf with Hilbert polynomial $\chi(L_Y(n)) = a_1 + a_2 \binom{n+1}{1}$. Thus, by 1.11, $-a_1 \geq 0$ and L_Y is $[-a_1]$-regular.

Consider the following exact sequence:

$$H^0(L_Y(n)) \longrightarrow H^1(L(n-1)) \longrightarrow H^1(L(n)) \longrightarrow H^1(L_Y(n)).$$

For $n \geq -a_1 - 1$, we have $h^1(L(-a_1 - 2)) \geq h^1(L(n))$ since $h^1(L_Y(n)) = 0$ by 1.3 (i); for $n \leq -1$, we have $h^1(L(n-1)) \leq h^1(L(-1))$ since $h^0(L_Y(n)) = 0$ by 2.4. Finally, for every n, $h^0(L_Y(n-1)) \leq h^0(L_Y(n))$ since there exists an injection $L_Y(n-1) \longrightarrow L_Y(n)$, and

$h^0(L_Y(-a_1-1)) = a_2\binom{-a_1}{1} + a_1$ since $h^1(L_Y(-a_1-1)) = 0$; thus, for $n \leq -a_1-2$, we have $h^1(L(n)) - h^1(L(n+1)) \leq -a_2a_1 + a_1$. As a consequence, we conclude that for every n,

$$h^1(L(n)) \leq h^1(L(-a_1-2)) + (-a_1-1)(-a_2a_1+a_1). \tag{2.7.1}$$

[print 628]Since L_Y is $[-a_1]$-regular, $h^2(L(-a_1-2)) = 0$[614] by 1.4; hence $h^1(L(-a_1-2)) = h^0(L(-a_1-2)) - \chi(L(-a_1-2))$. Now $h^0(L(-a_1-2)) \leq a_2\binom{-a_1}{2}$ by 1.8 (i), since L is a $(0,0,a_2)$-sheaf by 2.5. Consequently,

$$0 \leq h^1(L(-a_1-2)) \leq -a_1(-a_1-1) - a_0. \tag{2.7.2}$$

Finally, 2.7.2 gives (ii), and 2.7.1 and 2.7.2 give (i).

Lemma 2.8. We suppose that $r = 2$ and $d(L) \leq 0$. Let $\delta = d(L)$ and $\beta = \delta(e_2-1)+e_1$. Then: [615]

$$a_0 \geq \langle c_1(L)^2\rangle + 2e_0 + \sum_{i=1}^{2}(2e_i - a_i)\binom{\delta-1+i}{i} - \beta(\beta+1).$$

Indeed, let $\chi(L^{\otimes m}(\delta-1)) = \alpha_2m^2 + \alpha_1m + \alpha_0$; then, by comparing polynomials, it follows that $2\alpha_2 = \langle c_1(L)^2\rangle$, that $\alpha_0 = \sum_{i=0}^{2} e_i\binom{\delta-1+i}{i}$ and that $\alpha_2+\alpha_1+\alpha_0 = \sum_{i=0}^{2} a_i\binom{\delta-1+i}{i}$. Let $\chi(L^{-1}(\delta)(n)) = \sum_{i=0}^{2}\beta_i\binom{n+i}{i}$; then $\beta_0 = \alpha_2 - \alpha_1 + \alpha_0$ and, by 2.3, $\beta_1 = d(L^{-1}(\delta)) + e_1$, which equals β. Thus $a_0 = 2\alpha_2 + 2\alpha_0 - \sum_{i=1}^{2} a_i\binom{\delta-1+i}{i} - \beta_0$, and since $d(L^{-1}(\delta)) \leq 0$, we have $\beta_0 \leq \beta(\beta+1)$ by 2.7 (ii); this gives the assertion.

Lemma 2.9. We suppose that $r \geq 2$, that $d(L) \leq 0$ and that X is normal. Then:

(i) $h^1(L(-n)) = 0$ for $n \geq 2 + h_2(a_{r-2}, a_{r-1}, a_r)$,

(ii) $h^1(L(n)) \leq h_r(a_{r-2}, a_{r-1}, a_r)$ for every n,

(iii) $|a_i| \leq g_{r-i}(a_{r-2}, a_{r-1}, a_r)$ for $i = 0, \ldots, r$,

where $h_2(X_0,X_1,X_2), h_3(X_0,X_1,X_2), \ldots$ and $g_0(X_0,X_1,X_2), g_1(X_0,X_1,X_2), \ldots$ are universal polynomials with coefficients in $\mathbb{Q}$, which depend on neither k nor X.

[print 629]Indeed, let us first treat the case $r = 2$. Put $g_i(X_0,X_1,X_2) = X_{2-i}^2$ for $i = 0,1,2$ and $h_2(X_0,X_1,X_2) = X_1(X_1+1)X_2 - X_0$. Then (iii) is evident and (ii) is 2.7 (i).

Let X lie in $P = \mathbb{P}_k^N$, and let $\omega = \underline{\mathrm{Ext}}_{O_P}^{N-2}(O_X, \Omega_{P/k}^N)$. Then $h^1(L(-n)) = h^1(\omega \otimes L^{-1}(n))$ and ω is invertible outside the finite set Z of singular points of X (see FGA 149 §6). Let Y be a normal hyperplane section of X which does not contain Z. Then $\omega_Y(1) \simeq \Omega_{Y/k}^1$ and we have an exact sequence:

$$0 \longrightarrow \omega \otimes L^{-1}(-1) \longrightarrow \omega \otimes L^{-1} \longrightarrow \Omega_{Y/k}^1 \otimes L_Y^{-1}(-1) \longrightarrow 0.$$

Now $h^1(\Omega_{Y/k}^1 \otimes L_Y^{-1}(n)) = h^0(L_Y(-n))$ for every n, and $h^0(L_Y(-n)) = 0$ for $n \geq 1$ by 2.5. Thus $\Omega_{Y/k}^1 \otimes L_Y^{-1}(-1)$ is 3-regular and, by 1.4, $\omega \otimes L^{-1}$ is $[3 + h^1(\omega \otimes L^{-1}(2))]$-regular. Consequently, $h^1(L(-n)) = 0$ for $n \geq 2 + h^1(L(-2))$; hence (i).

Now suppose that $r \geq 3$ and argue by induction on r. Let Y be a normal hyperplane section of X and consider the exact sequence:

$$H^0(L_Y(-n)) \longrightarrow H^1(L(-n-1)) \longrightarrow H^1(L(-n)) \longrightarrow H^1(L_Y(-n)). \tag{2.9.1}$$

By 1.7, $\chi(L_Y(n)) = \sum_{i=0}^{r-1} a_{i+1}\binom{n+i}{i}$. Thus, for $n \geq 2 + h_2(a_{r-2}, a_{r-1}, a_r)$, the induction hypothesis gives $h^1(L_Y(-n)) = 0$, and since $n \geq 2$ by (ii) for $r = 2$, 2.4 gives $h^0(L_Y(-n)) = 0$;

hence $h^1(L(-n-1)) = h^1(L(-n))$. But, for $n \gg 0$, $h^1(L(-n)) = 0$ (see FGA[616] 149 §6); hence (i).

As induction hypothesis, suppose that the polynomials g_i and h_i are defined for $i \leq r-1$. Then, for every n, $h^1(L(n)) \leq h^1(L(n-1)) + h_{r-1}(a_{r-2}, a_{r-1}, a_r)$ by 2.9.1. Thus, by (i), [print 630]already proved, we conclude that for any $m \geq -2$,

$$h^1(L(n)) \leq \big[2 + h_2(a_{r-2}, a_{r-1}, a_r) + m\big] h_{r-1}(a_{r-2}, a_{r-1}, a_r) \tag{2.9.2}$$

for every $n \leq m$.

By 2.5, L_Y is a $(0, \ldots, 0, a_r)$-sheaf. Put $c_i = g_{r-i}(a_{r-2}, a_{r-1}, a_r)$ for $i = 1, \ldots, r$ and $m = P_{r-2}(c_1, \ldots, c_{r-1})$. By 1.11, $m \geq 0$ and L_Y is m-regular. Thus, by 1.4 (ii), $h^1(L(m-2)) \geq h^1(L(n))$ for every $n \geq m-1$. Therefore, if we define polynomials by $M = P_{r-2}(g_{r-1}, \ldots, g_1)$ and $h_r = (h_2 + M)h_{r-1}$, then we have (ii) by 2.9.2.

Finally, since L_Y is m-regular, $h^q(L(m)) = 0$ for $q \geq 2$ by 1.4 (i); hence $\sum_{i=0}^{r} a_i \binom{m+i}{i} = h^0(L(m)) - h^1(L(m))$. But $h^0(L(m)) \leq a_r \binom{m+r}{r}$ by 1.8 (i). Thus, if we define $g_r = \sum_{i=1}^{r} g_{r-i} \binom{M+i}{i} + a_r \binom{M+r}{r} + h_r$, we have (iii).

Remark 2.10. For fixed m, consider an equality of polynomials in n:

$$\sum_{i=0}^{r} a_{m,i} \binom{n+i}{i} = \sum_{j=0}^{r} a_j \binom{m+n+j}{j}.$$

Then one easily finds the formula [617]

$$a_{m,i} = \sum_{j=0}^{r-i} a_{j+i} \binom{m-1+j}{j}$$

by induction on r so that $a_{m,i}$ becomes $a_{m,0}$, and by taking $n = -1$.

Similarly, one finds that there exists a polynomial with rational coefficients $P_i^{(m,r)}(X_r, \ldots, X_i)$ which is increasing in X_i, such that in the following equality:

$$\sum_{i=0}^{r} a_i^{(m)} \binom{n+i}{i} = \sum_{j=0}^{r} a_j \binom{mn+j}{j}$$

[print 631]we have: $a_i^{(m)} = P_i^{(m,r)}(a_r, \ldots, a_i)$.

Lemma 2.11. Let S be a noetherian scheme and X a projective S-scheme, endowed with a very ample sheaf $H = O_X(1)$. We suppose that the fibers of X/S are geometrically integral, of the same dimension r. Let $\mathcal{L}$ be a family of classes of invertible sheaves L_K on the fibers of X/S. Then for $\mathcal{L}$ to be bounded, it is (necessary and) sufficient that, in the Hilbert polynomials $\chi(L_K(n)) = \sum_{i=0}^{r} a_i \binom{n+i}{i}$ of the $L_K \in \mathcal{L}$, all the coefficients a_i remain bounded. Moreover, if the fibers of X/S are also geometrically normal, then it suffices that the three coefficients a_r, a_{r-1}, a_{r-2} remain bounded.

Indeed, let $\delta = \sup\{d(L_K)\}$[618]. If $\delta > 0$, one may replace $\mathcal{L}$ by the family of classes of the $L_K(-\delta)$ by 2.3 and 2.10; one may thus suppose $\delta \leq 0$. Then $\mathcal{L}$ is a (b)-family by 2.5, hence bounded if all the coefficients a_i remain bounded, by 1.13, and, when the fibers of X/S are geometrically normal, it suffices for this that the three coefficients a_r, a_{r-1}, a_{r-2} remain bounded, by 2.9.

3. General finiteness theorems

Lemma 3.1. Let S be a scheme and X a quasi-compact and quasi-separated S-scheme; let $\underline{\mathrm{Pic}}_{X/S}$ be the Picard functor, (localized for the f.p.p.f. topology)(*). Then for every filtered inverse system (Z_α) of quasi-compact and quasi-separated S-schemes [print 632]with affine transition morphisms, the canonical map

$$\varinjlim \underline{\mathrm{Pic}}_{X/S}(Z_\alpha) \longrightarrow \underline{\mathrm{Pic}}_{X/S}(\varprojlim Z_\alpha) \tag{3.1.1}$$

is bijective.

Consequently (EGA IV 8.14.2), if the Picard functor is represented by an S-scheme $\mathrm{Pic}_{X/S}$, then $\mathrm{Pic}_{X/S}$ is locally of finite presentation.

Indeed, an element λ_α of $\underline{\mathrm{Pic}}_{X/S}(Z_\alpha)$ is represented by an invertible sheaf L'_α on an $X_{Z'_\alpha}$ where $Z'_\alpha \to Z_\alpha$ is an f.p.p.f. morphism. Put $Z = \varprojlim Z_\alpha$. The image λ of λ_α in $\underline{\mathrm{Pic}}_{X/S}(Z)$ is represented by the invertible sheaf L' on $X_{Z'}$, the pullback of L'_α, where[619] $Z' = Z'_\alpha \times_{Z_\alpha} Z$.

If λ is the identity element of $\underline{\mathrm{Pic}}_{X/S}(Z)$, there exists an f.p.p.f. morphism $Z'' \to Z'$ such that the pullback of L' to $X_{Z''}$ is $\simeq O_{X_{Z''}}$. By (EGA IV 8.8.2, 8.10.5 (vi), 11.2.6 (ii), 8.5.2 (i), and 8.5.2.4), there exist a $\beta \geq \alpha$ and an f.p.p.f. morphism $Z''_\beta \to Z'_\beta$, where $Z'_\beta = Z'_\alpha \times_{Z_\alpha} Z_\beta$, such that the pullback of L'_α to $X_{Z''_\beta}$[620] is $\simeq O_{X_{Z''_\beta}}$. The image of λ_α is therefore already the identity element in $\underline{\mathrm{Pic}}_{X/S}(Z_\beta)$. Consequently, 3.1.1 is injective.

The surjectivity of 3.1.1 is proved in an analogous way.

Proposition 3.2. Let S be a noetherian scheme and X a proper S-scheme. We suppose that the Picard functor is represented by an S-scheme $\mathrm{Pic}_{X/S}$. Then :

(i) $\mathrm{Pic}_{X/S}$ is locally of finite type over S.

(ii) [print 633]The subsets Λ of $\mathrm{Pic}_{X/S}$ correspond bijectively to the families $\mathscr{L}$ of classes of invertible sheaves on the fibers of X/S.

(iii) A family Λ of $\mathrm{Pic}_{X/S}$ is quasi-compact if and only if the corresponding family $\mathscr{L}$ is bounded.

Indeed, (i) follows from (3.1); (ii), from the following fact: if s is a point of S and K is an algebraically closed extension of $k(s)$, then the K-points of $\mathrm{Pic}_{X/S}$ correspond bijectively to the isomorphism classes of invertible sheaves on X_K.

As for (iii), suppose that the subset Λ of $\mathrm{Pic}_{X/S}$ is quasi-compact. Then Λ is contained in an open subset U of finite type, and the inclusion $U \to \mathrm{Pic}_{X/S}$ corresponds to an invertible sheaf L on an X_T, where $T \to U$ is a surjective, (flat) morphism of finite type. Finally, $T \to S$ is of finite type, so L bounds $\mathscr{L}$.

Conversely, suppose that $\mathscr{L}$ is bounded by a coherent sheaf L on an X_T, with T of finite type over S. One may suppose L invertible (1.13 (i)) so that L defines a morphism $f : T \to \mathrm{Pic}_{X/S}$. The image of f contains Λ and is quasi-compact, hence noetherian by (i). Consequently, Λ is quasi-compact, and we have (iii).

By 3.2, the usual definition of morphisms of finite type between two Picard schemes generalizes to the following:

(*) Cf. SGA 3 IV 6.3, and FGA no. 232 for the definition of $\underline{\mathrm{Pic}}_{X/S}$, where, as indicated, the fpqc topology should here be replaced by the fppf topology.

Definition 3.3. Let X and Y be proper over noetherian S. A morphism between the Picard functors

$$\underline{\mathrm{Pic}}_{Y/S} \longrightarrow \underline{\mathrm{Pic}}_{X/S}$$

[print 634]is said to be of finite type if every bounded family of classes of invertible sheaves on the fibers of X/S has as its inverse image a bounded family of classes of sheaves on the fibers of Y/S.

Lemma 3.4. Let X, Y, X', Y' be proper over noetherian S, and $T \to S$ surjective of finite type. Then :

(i) For a morphism $\Phi : \underline{\mathrm{Pic}}_{Y/S} \to \underline{\mathrm{Pic}}_{X/S}$ to be of finite type, it is necessary and sufficient that $\Phi_T : \underline{\mathrm{Pic}}_{Y_T/T} \to \underline{\mathrm{Pic}}_{X_T/T}$ be of finite type.

(ii) In a commutative diagram

$$\begin{array}{ccc} \Phi' : \underline{\mathrm{Pic}}_{Y'/S} & \longrightarrow & \underline{\mathrm{Pic}}_{X'/S} \\ {\scriptstyle \Phi_Y}\uparrow & & {\scriptstyle \Phi_X}\uparrow \\ \Phi : \underline{\mathrm{Pic}}_{Y/S} & \longrightarrow & \underline{\mathrm{Pic}}_{X/S} \end{array}$$

with Φ' and Φ_Y of finite type, Φ is of finite type.

Indeed, suppose that Φ_T is of finite type. Let $\mathscr{L}$ be a family on the fibers of X/S, bounded by an L on an X_R. The family $\mathscr{L}_T = \mathscr{L} \times_S T$ on the fibers of X_T/T, defined in the evident way, is bounded by the pullback of L to $X_{R\times T}$. If $\Phi_T^{-1}(\mathscr{L}_T)$ is bounded by an M on a Y_Q, then M also bounds $\Phi^{-1}(\mathscr{L})$.

The converse of (i) and assertion (ii) are proved in an analogous way.

Theorem 3.5. Let S be a noetherian scheme, X and Y two proper S-schemes and $f : X \to Y$ a morphism. If f is surjective, then the induced morphism between the Picard functors

$$f^* : \underline{\mathrm{Pic}}_{Y/S} \longrightarrow \underline{\mathrm{Pic}}_{X/S}$$

[print 635]is of finite type.

Indeed, by 3.4 (i), one may evidently suppose S reduced and irreducible. Then there exists a nonempty open subset U of S such that $\underline{\mathrm{Pic}}_{Y/S}$ and $\underline{\mathrm{Pic}}_{X/S}$ are representable and the restriction of f^* is quasi-affine (XII 1.1), hence of finite type by 3.2 (i). Endowing $T = S - U$ with the induced reduced structure, one may suppose that the morphism $\underline{\mathrm{Pic}}_{Y_T/T} \longrightarrow \underline{\mathrm{Pic}}_{X_T/T}$ induced by f^* is of finite type by noetherian induction ; this gives the assertion, by 3.4 (i) applied to $U \amalg T \to S$.

Theorem 3.6. Let S be a noetherian scheme, X a proper S-scheme and $\underline{\mathrm{Pic}}_{X/S}$ the Picard functor. Then, for every integer $n \neq 0$, the n-th power homomorphism

$$\Phi_n : \underline{\mathrm{Pic}}_{X/S} \longrightarrow \underline{\mathrm{Pic}}_{X/S}$$

is a morphism of finite type.

Indeed, let $f : X' \to X$ be a Chow presentation: X' is projective over S and f is surjective. If the theorem is admitted for X', then the theorem for X follows by 3.5 and 3.4 (ii). One may therefore suppose X/S projective.

There exist a nonempty open subset U of S, an integral scheme S' and a finite surjective morphism $S' \to U$, such that every irreducible component X'_i of $X_{S'}$, endowed with the induced

reduced structure, is integral over S' (XII 1.3). Endowing $T = S - U$ with the induced reduced structure, one may admit the theorem for X_T/T by noetherian induction. If it is admitted for all the X'_i/S'[621], it follows for $\coprod X'_i/S'$[622] by the [print 636]following easy lemma, hence for $X_{S'}/S'$ by 3.5 and 3.4 (ii), and finally for X/S by 3.4 (i).

Lemma 3.7. Let $X = \coprod X_i$ be of finite type over noetherian S, and $\mathcal{F}$ a family of classes of coherent sheaves on the fibers of X/S. For $\mathcal{F}$ to be bounded, it is necessary and sufficient that for every i the induced family $\mathcal{F}_i$ on the fibers of X_i/S be bounded.

In the proof of 3.6, we reduced to the case where X is projective and integral over S; endow X/S with a very ample sheaf $O_X(1)$. Let $\mathcal{L}$ be a bounded family of classes of invertible sheaves on the fibers of X/S, and L an invertible sheaf on an X_T, with T of finite type over S, which bounds $\mathcal{L}$. After decomposing T into pieces, one may suppose L flat over T, and after replacing T by each of its connected components, one may suppose T connected. Then the polynomial in q and m, $P(q,m) = \chi(L_t^{\otimes q}(m))$, does not depend on $t \in T$ (EGA III 7.9.4).

Let $\mathcal{L}_n = \Phi_n^{-1}(\mathcal{L})$: $\mathcal{L}_n$ is the family of classes of invertible sheaves M_K such that the class of $M_K^{\otimes n}$ belongs to $\mathcal{L}$. Let $\chi(M_K^{\otimes p}(m)) = \sum_{i=0}^{r} a_i(p)\binom{m+i}{i}$, where $a_i(p)$ is a polynomial in p of degree $\leq r-i$. For variable q, m, we have $\chi(M_K^{\otimes qn}(m)) = P(q,m)$, so each $a_i(qn)$ is independent of the chosen M_K; consequently, each polynomial $a_i(p)$ is also independent of M_K. Therefore the family $\mathcal{L}_n$ has a single Hilbert polynomial $\chi(M_K(m))$, hence is bounded by 2.11.

Theorem 3.8. Let S be a noetherian scheme, Y a projective S-scheme endowed with a relatively ample invertible sheaf $O_Y(1)$, X the zero scheme of a section φ of $O_Y(1)$, and $f : X \to Y$ the inclusion morphism. We [print 637]suppose that for every point $s \in S$ and every irreducible component Y_s^i of Y_s, the irreducible components of $X_s \cap Y_s^i = V(\varphi|Y_s^i)$ have dimension ≥ 2, (which is the case if the irreducible components of the fibers of Y have dimension ≥ 3). Then the induced morphism between the Picard functors

$$f^* : \underline{\mathrm{Pic}}_{Y/S} \longrightarrow \underline{\mathrm{Pic}}_{X/S}$$

is of finite type.

Moreover, if S is integral, there exists a nonempty open subset U of S such that the Picard functors restricted to U are representable and the induced morphism

$$\underline{\mathrm{Pic}}_{Y/S}|U \longrightarrow \underline{\mathrm{Pic}}_{X/S}|U$$

is quasi-affine (of finite type).

Let us first prove the first assertion. As in the proof of 3.6, one sees by noetherian induction that one may replace S by any nonempty open subset, and one easily reduces to the case where S is integral and Y is integral over S, always replacing X by $X \times_Y Y'$ when Y is replaced by Y'; this done, one reduces in the same way to the case where Y is normal over S, using the following lemma in place of XII 1.3.

Lemma 3.9. Let S be an integral noetherian scheme and X an S-scheme, integral over S. Then there exist a nonempty open subset U of S, an integral scheme S' finite and faithfully flat over U, an S'-scheme X' normal and integral over S', and a finite surjective S'-morphism $X' \to X_{S'}$.

[print 638]Indeed, let η be the generic point of S and K the algebraic closure of $k(\eta)$. It is clear that $\mathrm{Spec}(K)$ is the limit of the filtered inverse system (S_α) of integral affine schemes endowed with a finite faithfully flat morphism $S_\alpha \to U_\alpha$, where U_α is a nonempty open subset of S. The normalization of X_K therefore comes from an S_α-scheme X' for some α, and the finite surjective morphism $X'_K \to X_K$ from a finite surjective S_α-morphism[623] $X' \to X_{S_\alpha}$ (EGA IV 8.8.2, 8.10.5). Finally, since X'_K is normal and integral, there exists an open subset S' of S_α such that $X'_{S'}$ is normal and integral over S' (EGA IV 9.9.5).

Remark 3.10. Retaining the notation and conditions of 3.9, suppose that there exists a resolving morphism $X'_K \to X_K$ (EGA IV 7.9.1), (for example if $\dim(X_K) \leq 2$, by Abhyankar).[624] Then a similar argument shows that there exist a finite faithfully flat morphism $S' \to U$, a scheme X' smooth over S' and a proper surjective birational morphism $X' \to X_{S'}$.

In the proof of the first part of 3.8, we reduced to the case where S is integral and Y is normal and integral over S, so Y is integral. If $\varphi = 0$, then $X = Y$ and the assertion is trivial. Otherwise, after restricting S, one may suppose that for every $s \in S$ we have $\varphi_s \neq 0$, so φ_s is O_{Y_s}-regular and the dimension r of Y_s is ≥ 3.

Now let $\mathcal{L}$ be a bounded family of classes of invertible sheaves on the fibers of X/S. Let M_K be an invertible sheaf on a geometric fiber Y_K of Y/S, with Hilbert polynomial $\chi(M_K(n)) = \sum_{i=0}^{r} a_i \binom{n+i}{i}$; then the Hilbert polynomial of f^*M_K is $\sum_{i=0}^{r-1} a_{i+1}\binom{n+i}{i}$. As the class of M_K ranges over $(f^*)^{-1}(\mathcal{L})$, that is, the [print 639]class of f^*M_K belongs to $\mathcal{L}$, the three coefficients a_r, a_{r-1}, a_{r-2} remain bounded by 1.13.

Let $H = O_Y(m)$ be a multiple of $O_Y(1)$ which is very ample. Then, in the Hilbert polynomials $\chi(M_K(mn)) = \sum_{i=0}^{r} a_i^{(m)}\binom{n+i}{i}$ relative to H of the M_K in $(f^*)^{-1}(\mathcal{L})$, the three coefficients $a_r^{(m)}, a_{r-1}^{(m)}, a_{r-2}^{(m)}$ remain bounded by 2.10. Consequently, $(f^*)^{-1}(\mathcal{L})$ is bounded[625] by 2.11; this gives the first part of 3.8.

As for the second part of 3.8, the "generic" representability of the Picard functors was established in XII 1.2, as was their separatedness. By an argument similar to that of XII 2, using 3.9 and XII 1.1, one reduces to the case of Y normal over S. Now, for the morphism $\underline{\mathrm{Pic}}_{Y/S} \to \underline{\mathrm{Pic}}_{X/S}$ to be quasi-affine, it suffices that it be (separated and) quasi-finite (EGA IV 8.12.8), which can be checked on the geometric fibers, and thus follows from the following lemma:

Lemma 3.11. Let k be an algebraically closed field, Y a normal projective k-scheme endowed with an ample sheaf $O_Y(1)$, and X the zero scheme of a section φ of $O_Y(1)$. We suppose that Y has dimension ≥ 3 (resp. ≥ 2). Then the morphism $\mathrm{Pic}_{Y/k} \longrightarrow \mathrm{Pic}_{X/k}$ (resp. the morphism $\mathrm{Pic}^{\tau}_{Y/k} \to \mathrm{Pic}^{\tau}_{X/k}$ (see 4.1)) is of finite type and has a kernel N which is a finite group (in general nonreduced).

Indeed, one may evidently suppose $\varphi \neq 0$; thus φ is regular, Y being integral. The morphism under consideration is of finite type by the first part of 3.8 (resp. by 4.7 (iii)). Consequently, the kernel N is an algebraic group, hence contained in $\mathrm{Pic}^{\tau}_{Y/k}$, and is therefore the same [print 640]in the two cases under consideration. Since $\mathrm{Pic}^{\tau}_{Y/k}$ is proper, Y being normal (FGA 236-2.13), and since N is a subgroup, necessarily closed (SGA 3 VI$_B$ 1.9.2), for N to be finite it suffices that it contain no subgroup of type $\mu_\ell (= \mathbb{Z}/\ell\mathbb{Z})$ where $(\ell, \mathrm{car}(k)) = 1$.

Consider Kummer theory, that is, the following commutative diagram, coming from the exact sequence

$$0 \longrightarrow \mu_\ell \longrightarrow \mathbb{G}_m \xrightarrow{x \mapsto x^\ell} \mathbb{G}_m \longrightarrow 0$$

of étale sheaves on Y (resp. on X):

$$\begin{array}{ccccccc} & & H^1(Y,\mu_\ell) & \longrightarrow & H^1(Y,\mathbb{G}_m) & \xrightarrow{\ell} & H^1(Y,\mathbb{G}_m) \\ & & \downarrow & & \downarrow & & \downarrow \\ H^0(X,\mathbb{G}_m) & \xrightarrow{\ell} & H^0(X,\mathbb{G}_m) & \longrightarrow & H^1(X,\mu_\ell) & \longrightarrow & H^1(X,\mathbb{G}_m) \xrightarrow{\ell} H^1(X,\mathbb{G}_m). \end{array}$$

By Hilbert's "Theorem 90", we have $H^1(Y,\mathbb{G}_m) = \mathrm{Pic}_{Y/k}(k)$ (resp. $H^1(X,\mathbb{G}_m) = \mathrm{Pic}_{X/k}(k)$). Since X is proper and integral, we have $H^0(X,\mathbb{G}_m) = k^*$; therefore $H^1(X,\mu_\ell) \to H^1(X,\mathbb{G}_m)$ is injective.

It therefore suffices to see that $H^1(Y,\mathbb{Z}/\ell\mathbb{Z}) \longrightarrow H^1(X,\mathbb{Z}/\ell\mathbb{Z})$ is injective, which follows from $\pi_1(X,\xi) \to \pi_1(Y,\xi)$ being surjective, or again (SGA 1 V 6.9) from every connected étale covering

$\widetilde{Y}$ of Y with group $\mathbb{Z}/\ell\mathbb{Z}$ having a restriction $\widetilde{X} = X \times_Y \widetilde{Y}$ which is also connected; this is a well-known consequence of Bertini's theorem and Zariski's connectedness theorem (SGA 1 X 2.10).

Remark 3.12. Retain the conditions of 3.11. Then:

(i) The kernel N is a finite unipotent group (SGA 3 XVII 1.1), since it contains no subgroup of multiplicative type μ_n for any n (SGA 3 XVII 4.6.1 vi), which is proved in the same way, using instead [print 641]of Kummer theory the following fact, due to Grothendieck (SGA 1 XI, on the last page): if X is a proper scheme over a field k, with $H^0(X, O_X) = k$, then there exists an isomorphism of k-group schemes $\underline{\mathrm{Hom}}_{\mathrm{gr}}(\mu_n, \underline{\mathrm{Pic}}_{X/k}) \simeq \underline{H}^1(X, \mathbb{Z}/n\mathbb{Z})$; consequently, if k is algebraically closed, we have $\mathrm{Hom}_{\mathrm{gr}}(\mu_n, \underline{\mathrm{Pic}}_{X/k}) \simeq H^1(X, \mathbb{Z}/n\mathbb{Z})$.

(ii) If k has characteristic zero, then the kernel N is zero, since every finite unipotent group is then zero (SGA 3 XVII 1.7). One thus approximately recovers Kodaira's "Vanishing theorem" in the form given by Mumford [2].

Theorem 3.13. Let S be a noetherian scheme and X a projective S-scheme, endowed with a relatively ample invertible sheaf $H = O_X(1)$. We suppose that the geometric fibers of X/S are integral and of dimension r. Then, for a family $\mathcal{L}$ of classes of invertible sheaves on the fibers of X/S, the following conditions are equivalent :

(i) $\mathcal{L}$ is a bounded family.

(ii) In the Hilbert polynomials $\chi(L_K(n)) = \sum_{i=0}^{r} a_i \binom{n+i}{i}$ of the $L_K \in \mathcal{L}$, the coefficient a_{r-1} remains bounded and the coefficient a_{r-2} remains bounded below.

(iii) As L_K ranges over $\mathcal{L}$, the degree $d(L_K) = \langle c_1(L_K) \cdot c_1(H_K)^{r-1} \rangle$ remains bounded and the intersection number $\langle c_1(L_K)^2 \cdot c_1(H_K)^{r-2} \rangle$ remains bounded below.

(iv) There exists an integer N such that we have

$$H^{-N} < \mathcal{L} < H^N$$

in the sense of the order relation defined by ampleness; that is, for every $L_K \in \mathcal{L}$, $H_K^{\otimes N} \otimes L_K$ and $H_K^{\otimes N} \otimes L_K^{-1}$ are ample.

[print 642]Let us prove that (i) implies (ii), (iii), and (iv). Thus suppose that $\mathcal{L}$ is bounded by an invertible sheaf on an X_T. Then $\mathcal{L}$ satisfies (iv) by EGA II 4.6.12. Suppose L flat over T. Then the polynomial in m and n, $\chi(L_t^{\otimes m}(n))$, depends only on the connected component of T (EGA III 7.9.4). Consequently, $\mathcal{L}$ satisfies (ii) and, taking 2.1 into account, (iii).

Next, note that the family of classes of the H_K (resp. O_{X_K}) is bounded, hence satisfies (iii) (resp. (ii)).

Suppose that $\mathcal{L}$ satisfies (iv). Then $\langle [N \cdot c_1(H_K) + c_1(L_K)] \cdot c_1(H_K)^{r-1} \rangle > 0$, hence $d(L_K) > -Nd(H_K)$, and likewise $Nd(H_K) > d(L_K)$. Moreover, $\langle [Nc_1(H_K) + c_1(L_K)]^2 \cdot c_1(H_K)^{r-2} \rangle > 0$, hence $\langle c_1(L_K)^2 \cdot c_1(H_K)^{r-2} \rangle > -2Nd(L_K) - N^2 d(H_K)$. Thus $\mathcal{L}$ satisfies (iii).

Finally, let us prove that (ii) (resp. (iii)) implies (i). By 2.10 and 2.3, one may replace H by a multiple; therefore suppose H very ample. Evidently one may decompose S into pieces; thus suppose that there exists a linear section Y of X which is integral over S, with fibers of dimension 2. By 3.8, one may replace $X, \mathcal{L}$ by $Y, \mathcal{L}|Y$, (taking account of 1.7 (resp. the projection formula), which implies that condition (ii) (resp. (iii)) on $X, \mathcal{L}$ is equivalent to (ii) (resp. (iii)) on $Y, \mathcal{L}|Y$); therefore suppose $r = 2$. If $\delta = \sup\{d(L_K)\}$ is > 0, one may replace $\mathcal{L}$ by the family of classes of the $L_K(-\delta)$ by 2.3 and 2.10; thus suppose $\delta \leq 0$.

If, in the Hilbert polynomials $\sum_{i=0}^{2} a_i \binom{n+i}{i}$, a_1 remains bounded and a_0 remains bounded below, then the three coefficients remain bounded by 2.3 and 2.7; thus $\mathcal{L}$ is bounded by 2.11; hence (ii) implies [print 643](i). On the other hand, (iii) implies (ii) by 2.3 and 2.8.

4. Finiteness theorems for $\underline{\mathrm{Pic}}^{\tau}_{X/S}$

4.1. Let S be a scheme and G a commutative group functor on S. We denote by G^0 (resp. G^τ) the group subfunctor whose T-points $f : T \to G$ are those of G such that, for every algebraically closed field K and every K-point t of T, there exist a connected K-scheme[626] Z, two K-points z, z_0 of Z and a morphism $g : Z \to G$ such that $g(z) = f(t)$ and $g(z_0) = 0$ in $G(K)$ (resp. there exists a nonzero integer n such that $n.f(t) \in G^0(K)$).

It is evident that the formation of G^0 (resp. G^τ) commutes with every base change, that a morphism of group functors $G \to G'$ induces a morphism $G^0 \to G'^0$ (resp. $G^\tau \to G'^\tau$), and that if S is a field and G is represented by a group scheme locally of finite type, then G^0 is represented by the connected component of the identity element, which is an open (and closed) subscheme, of finite type and geometrically irreducible (SGA 3 VI$_A$ 2.4).

Lemma 4.2. Let $\Phi : G \to G'$ be a morphism between two commutative group functors on a scheme S. We suppose that, for every algebraically closed field K and every morphism $\mathrm{Spec}(K) \to S$, G_K (resp. G'_K) is representable by a group scheme locally of finite type, and Φ_K is representable by a morphism of finite type, (that is, quasi-compact). Then

$$\Phi^{-1}(G'^\tau) = G^\tau.$$

[print 644]Indeed, we have already noted the inclusion $\Phi^{-1}(G'^\tau) \supset G^\tau$[627]. For the reverse inclusion, note that $g \in G(K)$ belongs to $G^\tau(K)$ if and only if the sequence of ng ($n \in \mathbb{N}$) is quasi-compact; since Φ_K is quasi-compact, it suffices for this that the sequence of $\Phi(ng) = n\Phi(g)$ have the same property; this gives the assertion.

Lemma 4.3. Let X, Y be proper over a scheme S and $f : X \to Y$ a surjective morphism (resp. the inclusion morphism of the zero scheme of a section φ of an ample sheaf $O_Y(1)$, under the hypotheses of 3.8)[628]. Then

$$(f^*)^{-1}\underline{\mathrm{Pic}}^{\tau}_{X/S} = \underline{\mathrm{Pic}}^{\tau}_{Y/S}.$$

Indeed, the assertion is a formal consequence of 4.2, (XII 1.2), 3.1, and 3.5 (resp. 3.8).

4.4. Let k be an algebraically closed field, X a proper k-scheme, and L an invertible sheaf on X. We say that L is algebraically equivalent to 0 (resp. τ-equivalent to 0) if there exist a connected k-scheme T of finite type, an invertible sheaf M on $X_T = X \times_k T$, and two k-points t, t_0 of T such that $L \simeq M_t$ and $O_X \simeq M_{t_0}$ (resp. if there exists a nonzero integer n such that $L^{\otimes n}$ is algebraically equivalent to 0), or, what is easily seen to be equivalent, if the k-point λ of $\underline{\mathrm{Pic}}_{X/k}$ corresponding to L belongs to $\underline{\mathrm{Pic}}^0_{X/k}(k)$ (resp. to $\underline{\mathrm{Pic}}^{\tau}_{X/k}(k)$).

Recall also that L is said to be numerically equivalent to 0 if, for every integral closed curve Y in X, the intersection number $\langle c_1(L) \cdot Y \rangle$ is 0 (see 2.1 or X 4.5).

[print 645]**Proposition 4.5. (Functorial behavior).** Let $f : X' \to X$ be a morphism between two proper k-schemes, and let L be an invertible module on X. Then :

(i) If L is τ-equivalent to 0 (resp. numerically equivalent to 0), then so is f^*L.

(ii) Conversely, when f is surjective, if f^*L is τ-equivalent to 0 (resp. numerically equivalent to 0), then so is L.

Indeed, assertion (i) (resp. (ii)) for τ-equivalence follows at once from 4.1 (resp. 4.3). On the other hand, let Y' be an integral closed curve in X', and $Y = f(Y')$ its image, endowed with the induced reduced structure. When $\dim(Y) = 1$, we have the projection formula (X 4.4.4)

$$\langle c_1(f^*L) \cdot Y' \rangle = [k(Y') : k(Y)] \, \langle c_1(L) \cdot Y \rangle \qquad 4.5.1.$$

where $k(Y')$ (resp. $k(Y)$) is the function field of Y' (resp. Y), and when $\dim(Y) = 0$, we have $(f^*L)_{Y'} \simeq O_{Y'}$. Consequently, if L is numerically equivalent to 0, then $\langle c_1(f^*L) \cdot Y' \rangle = 0$; hence (i).

Conversely, when f is surjective, given an integral closed curve Y in X, one can find an integral closed curve Y' in X' such that $Y = f(Y')$: let y be the generic point of Y, y' a point of the fiber $f^{-1}(y)$ algebraic over $k(y)$, and Y' the closure of y'. Then 4.5.1 shows that $\langle c_1(f^*L) \cdot Y' \rangle = 0$ implies $\langle c_1(L) \cdot Y \rangle = 0$; hence (ii).

Theorem 4.6. (The numerical characterization of $\mathrm{Pic}^{\tau}_{X/k}$). Let k be an algebraically closed field, X a proper k-scheme, and L an invertible sheaf on X. Then the following conditions are equivalent :

(i) L is τ-equivalent to 0.

(ii) For every coherent sheaf F on X, we have [print 646]

$$\chi(F \otimes L) = \chi(F)$$

(iii) L is numerically equivalent to 0.

If X/k is projective and equipped with an ample invertible module $H = O_X(1)$, the preceding conditions are also equivalent to the following:

(iv) For every integer m (positive or negative), $L^{\otimes m}(1)$ is ample.

(v) (If X is irreducible). For every pair of integers m, n, we have

$$\chi(L^{\otimes m}(n)) = \chi(O_X(n))$$

(vi) (If X is irreducible of dimension $r \geq 2$). We have

$$\langle c_1(L) \cdot c_1(H)^{r-1} \rangle = 0, \qquad \langle c_1(L)^2 \cdot c_1(H)^{r-2} \rangle = 0.$$

Let us prove that (i) implies (ii). Applying the dévissage lemma (EGA III 3.1) and 4.5 (i), we reduce to the case where X is integral and $F = O_X$. We must then prove that the polynomial $\chi(L^{\otimes m})$ is constant. For this, we may replace L by a multiple, and thus assume that L is algebraically equivalent to 0. In this case, if L is deformed into O by the algebraic family defined by an invertible sheaf M on some $X \times_k T$ where T is connected, of finite type, the polynomial $\chi(M_t^{\otimes m})$ does not depend on $t \in T$ (EGA III 7.9.5); hence the implication.

Let Y be a closed integral curve in X. Taking $F = O_Y$ in (ii), we find $\chi(O_Y) = \chi(L_Y)$. Thus the formula ("Riemann's theorem") $\chi(L_Y) = \langle c_1(L) \cdot Y \rangle + \chi(O_Y)$ gives $\langle c_1(L) \cdot Y \rangle = 0$; hence (ii) implies (iii).

Let $f : X' \to X$ be a Chow presentation: X' is projective and f is surjective. If we grant that (iii) implies (i) for X', then [print 647]by 4.5 the implication for X follows. Henceforth, then, suppose that X is projective, equipped with an ample sheaf H.

Let X_i be the irreducible components of X, equipped with the induced reduced structure. Applying 4.5 and EGA III 2.6.2 to the finite surjective morphism $\coprod X_i \to X$ reduces the proof that (iii) implies (i) and (iv) to the case where X is integral.

If X is irreducible, it is clear that (ii) implies (v), that (iii) implies (vi), and, taking 2.1 into account, that (v) implies (vi), and that (vi) for X is equivalent to (vi) for X_{red} (X 4.4.1).

It therefore remains to show that (vi) implies (i) and (iv), and that (iv) implies (i), assuming X integral. But then (vi) (resp. (iv)) implies that the family $\mathcal{L}$ of classes of the $L^{\otimes m}$ ($m \in \mathbb{N}$) satisfies 3.13 (iii) (resp. (iv)), and is therefore bounded. There thus exists an invertible sheaf L on some X_T, with T of finite type over k, which bounds $\mathcal{L}$. Let p, q be two distinct integers such that $L^{\otimes p}$[629] and $L^{\otimes q}$ correspond to the same connected component of T; then $L^{\otimes(p-q)}$ is therefore algebraically equivalent to 0; hence (vi) (resp. (iv)) implies (i).

Finally, we likewise have that (vi) implies the existence of an integer N such that, for every integer m, $L^{\otimes m} \otimes H^{\otimes N}$ is ample. Then, for every m, $(L^{\otimes m} \otimes H)^{\otimes N}$ is ample; hence (vi) implies (iv), which completes the proof.

Theorem 4.7. Let S be a noetherian scheme, and X a proper S-scheme. Then:

(i) The inclusion morphism $\underline{\mathrm{Pic}}^{\tau}_{X/S} \to \underline{\mathrm{Pic}}_{X/S}$ is representable by an open immersion.

(ii) If X is projective and integral over S, then $\underline{\mathrm{Pic}}^{\tau}_{X/S} \to \underline{\mathrm{Pic}}_{X/S}$ [print 648]is also representable by a closed immersion.

(iii) $\underline{\mathrm{Pic}}^{\tau}_{X/S}$ is of finite type over S, that is, the family of classes of invertible sheaves τ-equivalent to 0 on the fibers of X/S is bounded.

Indeed, in (i) (resp. (ii)), we must treat the case of a morphism $T \to \underline{\mathrm{Pic}}_{X/S}$, where T is an arbitrary S-scheme, and show that the inverse image of $\underline{\mathrm{Pic}}^{\tau}_{X/S}$ is representable by an open (resp. and closed) subscheme of T. It suffices to take $T = \mathrm{Spec}(B)$ over an affine open $\mathrm{Spec}(A)$ of S. Then T is the limit of the filtering inverse system of affine S-schemes $T_\alpha = \mathrm{Spec}(B_\alpha)$, where B_α runs through the finite-type A-subalgebras of B. By 3.1, the morphism in question comes from a morphism $T_\alpha \to \underline{\mathrm{Pic}}_{X/S}$. We may therefore suppose that T is of finite type over S.

The morphism $T \to \underline{\mathrm{Pic}}_{X/S}$ corresponds to an f.p.p.f. morphism $R \to T$ and an invertible sheaf L on X_R. By f.p.p.f. descent, we may replace T by R. After the base change $R \to S$, taking into account the compatibility of the formation of $\underline{\mathrm{Pic}}^{\tau}_{X/S}$ with every base change (4.1), we are thus reduced in (i) and (ii) to studying the case of a section $S \to \underline{\mathrm{Pic}}_{X/S}$, defined by an invertible sheaf L on X.

We must prove that the set S_0 of $s \in S$ such that L_s is τ-equivalent to 0 is open, and also closed under the conditions of (ii); since S is noetherian, this means three things: for an $s \in S_0$,

(a) every generization of s remains in S_0.

(b) every specialization of s, near s, remains in S_0.

(c) under the conditions of (ii), every specialization of s remains in S_0.

[print 649]Let $f : X' \to X$ be a Chow presentation: X' is projective over S; and f is surjective. If we grant (a), (b), (c) and (iii) for X', then they follow for X by 4.3 and 3.5. We may therefore suppose that X is projective over S.

Let H be ample for X/S. If, for an $s \in S$, $(L^{\otimes m} \otimes H)_s$ is ample, then $(L^{\otimes m} \otimes H)_t$ is ample for every generization t of s (EGA III 4.7.1). By the equivalence of 4.6 (i) and (iv), we have therefore established (a).

For (b), (c) and 4.7 (iii), we may suppose that S is integral, with s the generic point in (b) and (c) (For (b) and (c), this is clear, and for (iii), it follows from 3.4 (i) applied to the morphism $\coprod S_i \to S$, where the S_i are the irreducible components of S, equipped with the induced reduced structure).

There exist an integral scheme S' and an f.p.p.f. morphism $S' \to S$ such that every irreducible component X'_i of $X_{S'}$, equipped with the induced reduced structure, is integral over S' (XII 1.3). To prove (b) and (iii), it suffices to do so for $X_{S'}/S'$, (in (b), by descent; in (iii), by noetherian induction and 3.4 (i)).

Finally, consider the canonical surjective morphism $\coprod X'_i \to X_{S'}$[630]. Then (b) and (iii) for $X_{S'}/S'$ follow from (c) and (iii) for the X'_i/S', by 4.3, 3.5 and 3.7.

We have therefore reduced to proving (c) and (iii), assuming S integral with generic point s, and X projective and integral over S. Let $O_X(1)$ be a very ample sheaf for X/S. Then, for every m, n, $\chi(L_t^{\otimes m}(n))$ and $\chi(O_{X_t}(n))$ do not depend on $t \in S$ (EGA III 7.9.4). Thus (c) follows from 4.6 (i) $\iff$ (v), which also implies that the family of classes of invertible sheaves τ-equivalent to 0 on the fibers of X/S has just [print 650]one Hilbert polynomial; thus (iii) follows from 2.11.

Remark 4.8. Let X be proper and integral over noetherian S. Without a projectivity hypothesis on X/S, it is unknown whether it remains true that $\underline{\mathrm{Pic}}^{\tau}_{X/S} \to \underline{\mathrm{Pic}}_{X/S}$ is a closed immersion, except when X is normal over S and $\underline{\mathrm{Pic}}_{X/S}$ is representable (FGA 236, 2.3).

5. Finiteness theorems of "Néron-Séveri" type.

Theorem 5.1. Let X be proper over noetherian S; then the Néron-Séveri groups $\underline{\mathrm{Pic}}_{X_K/K}(K)/\underline{\mathrm{Pic}}^0_{X_K/K}(K)$ of the geometric fibers X_K/K of X/S are finitely generated, and their ranks and the orders of their torsion subgroups $\underline{\mathrm{Pic}}^{\tau}_{X_K/K}(K)/\underline{\mathrm{Pic}}^0_{X_K/K}(K)$ remain bounded.

Indeed, by noetherian induction, we may restrict ourselves to proving 5.1 with S replaced by an arbitrary nonempty open of S_{red}; thus we may suppose that $\underline{\mathrm{Pic}}^{\tau}_{X/S}$ is representable by an S-scheme of finite type (XII 1.2 and 4.7 (iii)). Then the order of $\underline{\mathrm{Pic}}^{\tau}_{X_K/K}(K)/\underline{\mathrm{Pic}}^0_{X_K/K}(K)$ is equal to the number of connected components of $(\mathrm{Pic}^{\tau}_{X/S})_K$, a number that remains the same over a nonempty open of S (EGA IV 9.7.9).

It remains to bound the rank of $\underline{\mathrm{Pic}}_{X_K/K}(K)/\underline{\mathrm{Pic}}^{\tau}_{X_K/K}(K)$. As in the proof of 4.7 (iii), we readily reduce to the case where S is integral, and X is projective and integral over S. Let r be the common dimension of the fibers of X/S. If r is 0 (resp. 1), then this rank is clearly 0 (resp. 1).

If r is ≥ 2, we shall reduce to the case where $r = 2$ and X is smooth over S. After restricting S, we can find a linear section Y of X which is [print 651]still integral over S, with fibers of dimension 2; by 4.3, we may replace X by Y. Finally, by 4.3 and 3.10 (using resolution of singularities for surfaces), after replacing S by a flat covering of a nonempty open of S, we may replace X by a smooth S-scheme.

Now, by 4.6 (i) $\iff$ (iii) and the finiteness and specialization theorem for étale cohomology groups (SGA 4 XVI 2.2), 5.1 follows from the next proposition.

Proposition 5.2. Let k be an algebraically closed field and X a proper smooth k-scheme. Then the group $C^p_{\mathrm{num}}(X)$ of algebraic cycles of codimension p on X, modulo numerical equivalence, is free of rank $\leq b^{2p}$, the Betti number.

Indeed, if we forget the action of the roots of 1, étale cohomology $H^*(X, \mathbb{Q}_\ell) =_{\mathrm{dfn}} \left[\varprojlim H^*(X, \mathbb{Z}/\ell^\nu\mathbb{Z})\right] \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell$ where $(\ell, \mathrm{car}(k)) = 1$ becomes a Weil cohomology: it satisfies

Poincaré duality and the Künneth[631] formula; and it receives a functorial, nonzero homomorphism from the ring of algebraic cycles modulo rational equivalence $C^*_{\mathrm{rat}}(X)$ (see X 7.13).

Let $a_1, \dots, a_m \in C^{r-p}_{\mathrm{rat}}(X)$, where $r = \dim(X)$, be elements that map to a basis of the $\mathbb{Q}_\ell$-vector subspace of $H^{2(r-p)}(X, \mathbb{Q}_\ell)$ generated by algebraic cycles. Consider the homomorphism

$$\alpha : C^p_{\mathrm{rat}}(X) \longrightarrow \mathbb{Z}^m$$

defined by $\alpha(b) = (\langle b.a_1\rangle, \dots, \langle b, a_m\rangle)$, where $\langle\ \rangle$ denotes the intersection number. Then one readily sees that the image of α is $C^p_{\mathrm{num}}(X)$.

[print 652]**Remark 5.3.** Let k be an algebraically closed field, and X a proper smooth surface over k. Then the second Betti number $b^2(X)$ can be interpreted as $c_2(X) - 2 + 2b^1(X)$, where $b^1(X) = 2\dim(\mathrm{Pic}_{X/k})$, and $c_2(X)$ = the second Chern class = the Euler-Poincaré characteristic = the intersection number of the diagonal of $X \times X$ with itself; thus $b^2(X)$ has an interpretation independent of cohomology. The inequality $\mathrm{rang}(C^1_{\mathrm{num}}(X)) \leq b^2(X)$[632] in this form for $\mathrm{car}(k) > 0$ is due (before étale cohomology) to IGUSA, who used deep results on the structure of the tame fundamental group of an algebraic curve minus a few points.

Theorem 5.4. Let k be an algebraically closed field, and X a proper k-scheme. Then there exist finitely many closed integral curves $Y_1, \dots, Y_p$ in X with the following property: a subset Λ of the scheme $\mathrm{Pic}_{X/k}$ is quasi-compact if and only if the intersection numbers $\langle \lambda.Y_i\rangle$ for $\lambda \in \Lambda$ remain bounded. Moreover, we may take for p the rank of the Néron-Severi group $\mathrm{Pic}_{X/k}(k)/\mathrm{Pic}^0_{X/k}(k)$.

Indeed, since $\mathrm{Pic}^\tau_{X/k}$ is an open quasi-compact subgroup of $\mathrm{Pic}_{X/k}$ (4.7), Λ is quasi-compact if and only if the corresponding set of points in the vector space $[\mathrm{Pic}_{X/k}(k)/\mathrm{Pic}^\tau_{X/k}(k)] \otimes \mathbb{Q}$ is finite. But τ-equivalence is equal to numerical equivalence (4.6 (i) $\Longleftrightarrow$ (iii)), so we may choose among the curves in X curves that map to a basis of the dual space; hence the assertion.

[print 653]

6. Appendix: Study of (b)-sheaves on $P = \mathbb{P}^N_k$.

6.1. Let k be an algebraically closed field, and let $P = \mathbb{P}^N_k$. Let F be a coherent sheaf on P, with Hilbert polynomial

$$\chi(F(n)) = \sum_{i=0}^{r} a_i \binom{n+i}{i},$$

and let

$$(c) = (c_0, \dots, c_r)$$

be a sequence of integers $\geq (a)$. Put

$$a_{m,i} = \sum_{j=0}^{r-i} a_{j+i}\binom{m-1+j}{j} \quad \text{and} \quad c_{m,i} = \sum_{j=0}^{r-i} c_{j+i}\binom{m-1+j}{j}.$$

Finally, for every integer $q \geq 1$, denote the largest coherent subsheaf of F with support of dimension $\leq q-1$ by N_q, and put $F_q = F/N_q$.

Lemma 6.2. Under the conditions of 6.1, consider an exact sequence

$$0 \longrightarrow I \longrightarrow O_P^{\oplus M} \longrightarrow F(m) \longrightarrow 0$$

with $m \geq 0$. Then I is p-regular with $p = P_r(c_{m,0}, \dots, c_{m,r})$, where P_r is the r-th (b)-polynomial (1.9).

Indeed, O_P is clearly a $(0,\dots,0,1)$-sheaf ($N+1$ terms), so I is a $(0,\dots,0,M)$-sheaf by 1.6. By 2.10, $\chi(I(n)) = M\binom{n+N}{N} - \sum_{i=0}^{r} a_{m,i}\binom{n+i}{i}$. Thus the assertion follows from 1.11 and 1.10.

Proposition 6.3. Under the conditions of 6.1, suppose that F is a (b)-sheaf. Then one can calculate, by means of universal polynomials in the a_i, b_i and N, integers m, M, m_1 and M_1 such that there is an exact sequence

$$O_P(-m_1)^{\oplus M_1} \longrightarrow O_P(-m)^{\oplus M} \longrightarrow F \longrightarrow 0.$$

Moreover, N occurs only in the calculation of M_1.

[print 654]Indeed, take $m = P_{r-1}(c_0,\dots,c_{r-1})$ with $c_i = b_i - a_i$, and $M = \sum a_i\binom{m+i}{i}$, so that there is an exact sequence

$$0 \longrightarrow I \longrightarrow O_P^{\oplus M} \longrightarrow F(m) \longrightarrow 0$$

by 1.11 and 1.3. Similarly, taking 6.2 into account, take $m_1 = m + p$ with $p = P_r(a_{m,0},\dots,a_{m,r})$, and $M_1 = M\cdot\binom{p+N}{N} - \sum a_i\binom{m_1+i}{i} \quad (= \chi(I(p)))$[633].

Theorem 6.4. Under the conditions of 6.1, suppose that F is a quotient of $O_P(-m)^{\oplus M}$ with $m \geq 0$. Let $b_i = P_{r-i}(c_{m,i},\dots,c_{m,r})$ for $i = 0,\dots,r$, where P_s is the s-th (b)-polynomial (1.9); let $b = b_1 + m - 1$, and let $B = \sum_{i=1}^{r} a_i\binom{b+i}{i}$. Then:

(i) F is b-regular.

(ii) $-B \leq a_0 = h^0(F(b)) - B$

(iii) F is a $(b_0,\dots,b_r)$-sheaf.

Indeed, take a section σ of $O_P(1)$ such that we obtain a commutative diagram with exact rows and columns

$$\begin{array}{ccccccccc}
 & & 0 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & I(-1) & \longrightarrow & I & \longrightarrow & J & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & O_P(-1)^{\oplus M} & \longrightarrow & O_P^{\oplus M} & \longrightarrow & O_Y^{\oplus M} & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F(m-1) & \longrightarrow & F(m) & \longrightarrow & G(m) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & &
\end{array}.$$

[print 655]Since $\chi(G(n)) = \sum_{i=0}^{r-1} a_{i+1}\binom{n+i}{i}$ by 1.7, J is b_1-regular by 6.2; thus $H^q(I(p)) = 0$ for $q \geq 2$ and $p \geq b_1 - q$ by 1.4 (i). The exact sequence $H^q(O_P^{\oplus M}(p)) \longrightarrow H^q(F(m+p)) \longrightarrow H^{q+1}(I(p))$ then shows that $F(m)$ is (b_1-1)-regular; hence (i).

Assertion (i) implies that $0 \leq h^0(F(b)) = \chi(F(b)) = \sum_{i=0}^{r} a_i\binom{b+i}{i}$; hence (ii). By 2.10 and 1.9,

$$\chi(F(b)) = \sum_{i=0}^{r} a_{m,i}\binom{b_1-1+i}{i} \leq b_1 + \sum_{i=0}^{r} c_{m,i}\binom{b_1-1+i}{i} = b_0\,;$$

thus, $h^0(F(-1)) \leq h^0(F(b)) \leq b_0$. By induction on r, we may suppose that G is a $(b_1, \ldots, b_r)$-sheaf; hence (iii) by 1.6.2.

Lemma 6.5. Under the conditions 6.1, suppose that F has no associated prime cycle of dimension 0.

(i) If $h^0(F) \geq 1$, then $h^0(F(-1)) \leq h^0(F) - 1$.

(ii) $H^0(F(-n)) = 0$ for $n \geq h^0(F)$.

(iii) Let an exact sequence $0 \to F(-1) \to F \to G \to 0$ and an integer $n_0 \geq 0$ satisfy $H^0(G(-n_0)) = 0$; then $H^0(F(-n_0)) = 0$ and $h^0(F) \leq n_0.h^0(G)$.

Indeed, let $\sigma \in H^0(F)$, $\sigma \neq 0$; then σ defines a nonzero subsheaf $F' = O_P.\sigma$[634] of F, so $s = \dim . \operatorname{supp}(F')$ is ≥ 1. Take a commutative diagram with exact rows and columns

$$\begin{array}{ccccccccc}
 & & 0 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F'(-1) & \longrightarrow & F' & \longrightarrow & G' & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F(-1) & \longrightarrow & F & \longrightarrow & G & \longrightarrow & 0
\end{array} \quad .$$

[print 656]From $s \geq 1$ follows $G' \neq 0$; consequently, F' is not contained in $F(-1)$; that is, the section σ is not contained in the image of the injection $H^0(F(-1)) \to H^0(F)$, whence (i).

Assertion (ii) follows immediately from (i), taking into account that $h^0(F) = 0$ clearly implies $h^0(F(-1)) = 0$, since $h^0(F(-1)) \leq h^0(F)$. In (iii), we find $0 \leq h^0(F(n)) - h^0(F(n-1)) \leq h^0(G(n))$. Thus, for $n \geq n_0$, $h^0(F(-n_0)) = h^0(F(-n))$ since $h^0(G(-n_0)) = 0$; hence $h^0(F(-n_0)) = 0$ by (ii). Consequently, $h^0(F(p - n_0)) \leq h^0(G(1 - n_0)) + \ldots + h^0(G(p - n_0)) \leq ph^0(G(p - n_0))$ for $p \geq 1$; whence (iii).

Proposition 6.6. Under the conditions of 6.1, suppose that F is a $(b_0, \ldots, b_r)$-sheaf. Then F_q is a $((b_{q-1})^q, \ldots, (b_{q-1})^2, b_{q-1}, b_q, \ldots, b_r)$-sheaf.

Indeed, let σ be a "general" section of $O_P(1)$[635]; then σ defines a commutative diagram with exact rows and columns

$$\begin{array}{ccccccccc}
 & & 0 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & N_q(-1) & \longrightarrow & N_q & \longrightarrow & G' & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F(-1) & \longrightarrow & F & \longrightarrow & G & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & F_q(-1) & \longrightarrow & F_q & \longrightarrow & G'' & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & &
\end{array}$$

where G is a $(b_1, \dots, b_r)$-sheaf (1.6 (i)) and where $G'' = G_{q-1}$ (a question of depth).

[print 657]Suppose that $q = 1$. Then dim.supp$(N_q) = 0$; thus $H^1(N_q(-1)) = 0$ and $h^0(F_q(-1)) \leq h^0(F(-1)) \leq b_0$[636]. Moreover, $G' = 0$; hence $G \simeq G''$. Thus F_1 is indeed a $(b_0, \dots, b_r)$-sheaf. In addition, 6.5 (ii) shows that $H^0(F_1(-b_0)) = 0$. If $q \geq 2$, we may suppose that G_{q-1} is a $((b_{q-1})^{q-1}, \dots, b_{q-1}, \dots, b_r)$-sheaf and that $H^0(G_{q-1}(-b_{q-1})) = 0$, by induction on q. Then 6.5 shows that $H^0(F_q(-b_{q-1})) = 0$ and $h^0(F_q(-1)) \leq (b_{q-1})^q$; whence the assertion.

Theorem 6.7. There exist two sequences $\{A_i(X_0, \dots, X_i; Y)\}$ and $\{A_i^{(q)}(X_0, \dots, X_q; Y)\}$ of polynomials with the following properties. Assume that the conditions of 6.1 are satisfied and that F is a quotient of $O_P(-m)^{\oplus M}$ with $m \geq 0$.

(i) If F is a $(b_0, \dots, b_r)$-sheaf, then, for $i = 0, \dots, r$,

$$|a_i| \leq A_{r-i}(b_i, \dots, b_r; m).$$

(ii) If $\chi(F_q(n)) = \sum_{i=0}^{r} a_i^{(q)} \binom{n+i}{i}$ is the Hilbert polynomial of F_q, then, for $i = 0, \dots, q-1$,

$$|a_i^{(q)}| \leq A_{r-i}^{(r-q+1)}(c_{q-1}, \dots, c_r; m).$$

[637]

(Notice that $a_{q-1}^{(q)} \leq a_{q-1}$ and $a_q^{(q)} = a_q, \dots, a_r^{(q)} = a_r$[638] since dim.supp$(N_q) \leq q-1$).

Indeed, let an exact sequence $0 \to F(-1) \to F \to G \to 0$[639] be such that G is a $(b_1, \dots, b_r)$-sheaf; arguing by induction on r, we may suppose that $A_0, \dots, A_{r-1}$ have been defined, taking 1.7 into account. Then 6.4 (ii) makes the definition of A_r clear, taking 1.8 into account. Finally, (ii) follows formally from 6.4 (iii), 6.6, and assertion (i) (where we take F_q for F).

[print 658]**Corollary 6.8. (Grothendieck).** Let X be projective over noetherian S, let $O_X(1)$ be very ample for X/S, and let $\mathcal{J}$ be a family of classes of coherent sheaves on the fibers of X/S. Assume:

(a) There exists a coherent sheaf E on X such that $\mathcal{J}$ is contained in the family of classes of quotients of sheaves of the form E_K.

(b_q) In the Hilbert polynomials $\chi(F_K(n))$ of the $F_K \in \mathcal{J}$, the coefficients in degrees $\geq q-1$ remain bounded above.

Then the $(F_K)_q$ (as F_K ranges over $\mathcal{J}$) form a bounded family; moreover, the coefficients in $\chi(F_K(n))$ in degrees $\geq q-2$ remain bounded below.

Indeed, we may clearly suppose that $X = \mathbb{P}_S^N$ and $E = O_X(-m)^{\oplus M}$. Then the first assertion follows formally from 6.7 (ii) and 1.13; the second follows from 6.4 (ii), by induction using an exact sequence $0 \to F(-1) \to F \to G \to 0$.

Definition 6.9. Let k be a field. A special positive k-cycle of dimension r is a projective k-scheme X, equipped with a very ample sheaf $O_X(1)$, which is a union of closed subschemes X_j of dimension r, with X_j obtained by a base change $\mathrm{Spec}(k) \to \mathrm{Spec}(k_j)$ as the inverse image of an integral k_j-scheme X_j' (equipped with $O_{X_j'}(1)$). The degree of X is the coefficient a_r in the Hilbert polynomial $\chi(O_X(n)) = \sum a_i \binom{n+i}{i}$.

Lemma 6.10. Let k be an algebraically closed field and X a special positive k-cycle of dimension r and degree d. Then O_X is a $(0, \dots, 0, d)$-sheaf.

[print 659]Indeed, let X lie in $\mathbb{P}_k^N$. After replacing k by the algebraic closure of a purely transcendental extension, which is allowed by 1.6 (i), we may suppose that there are $r(N+1)$ elements in k which are algebraically independent over all the k_j. Then the r successive "generic" hyperplane sections of X, defined by means of these elements, are still special positive k-cycles of degree d.

Arguing by induction on r, it therefore remains to note that, when $r = 0$, we have $h^0(O_X(-1)) = d$, and, when $r \geq 1$, we have $h^0(O_X(-1)) = 0$. For this last point, we may suppose that X itself comes from an integral scheme X', by the injection $O_X \to \coprod O_{X_j}$; but then $h^0(O_X(-1)) = h^0(O_{X'}(-1))$, which is 0 since $H^0(O_{X'})$ is a field.

Corollary 6.11. (i) Let r, d be two integers, let $c_i = A_{r-i}(0, \ldots, 0, d; 0)$ for $i = 0, \ldots, r$ (6.7 (i)), and let $p = P_r(c_0, \ldots, c_r)$ (1.9). Let k be an algebraically closed field, and X a special positive k-cycle of dimension $\leq r$ and degree $\leq d$, with Hilbert polynomial $\chi(O_X(n)) = \sum_{i=0}^r a_i \binom{n+i}{i}$. Then, for $i = 0, \ldots, r$, we have

$$|a_i| \leq c_i.$$

Moreover, X embeds as a subscheme of $\mathbb{P}_k^N$ with $N = d(r+1) - 1$, defined by (at most) $\binom{N+p}{N}$ equations of degree p.

(ii) (Chow) Let S be a noetherian scheme. As K ranges over the algebraically closed extensions of the $k(s)$, $s \in S$, the special positive K-cycles of bounded dimension and degree form a "bounded family" (either with the X regarded as subschemes of a fixed $\mathbb{P}_S^N$, or in an evident abstract sense).

[print 660]Indeed, in (i), the first assertion follows from 6.7 (i); the second, from 6.10 and 1.8 (i), and from 6.2 (applied to $0 \to I \to O_P \to O_X \to 0$, $P = \mathbb{P}_k^N$) and 1.3. Finally, (ii) follows from (i) and 1.13.

Remark 6.12. Let k be an algebraically closed field, and X a projective k-scheme of pure dimension r, with no embedded associated prime cycle. If we allow arbitrary nilpotent elements in O_X, we can no longer bound the coefficients in the Hilbert polynomial $\chi(O_X(n)) = \sum_{i=0}^r a_i \binom{n+i}{i}$ in terms of a_r alone.

For example, let Z be a smooth curve of degree d in $\mathbb{P}_k^3$. For every $n \gg 0$, there exists a smooth surface Y of degree n which contains Z. Then Z is a Cartier divisor on Y; put $O_{X_n} = O_Y/O_Y(-2Z)$. Finally, we have the exact sequence $0 \to O_Z(-Z) \to O_{X_n} \to O_Z \to 0$[640] and the formula $\langle Z^2 \rangle_Y = 2p_a(Z) - 2 - (n-4)d$; hence $a_0 = \chi(O_{X_n}(-1)) \to \infty$ $(n \to \infty)$, although $a_1 = 2d$.

Corollary 6.13. Under the conditions of 6.1, consider an exact sequence

$$O_P(-m_1)^{\oplus M_1} \longrightarrow O_P(-m)^{\oplus M} \xrightarrow{\alpha} F \longrightarrow 0$$

with $m_1 \geq m \geq 0$. Then, for $i = 0, \ldots, N$,

$$|a_i| \leq A_{N-i}(0, \ldots, 0, M; m_1) + M \binom{m}{N-i}.$$

Indeed, put $I = \mathrm{Ker}(\alpha)$; clearly, I is a $(0, \ldots, 0, M)$-sheaf, and $\chi(I(n)) = \sum_{i=0}^N \left[M \binom{-m-1+N-i}{N-i} - a_i \right] \binom{n+i}{i}$; whence the assertion by 6.7 (i), and the formula $\binom{-m-1+Q}{Q} = (-1)^Q \binom{m}{Q}$.

[print 661]**Corollary 6.14. (G. Hermann).**[641] For every $N \geq 0$, there exists a polynomial $R_N(m)$ such that, for every field k_0 and every ideal I of[642] the polynomial ring $k_0[T_1, \ldots, T_N]$, generated by elements of degree $\leq m$, the radical $\sqrt{I}$ of I is generated by elements of degree $\leq R_N(m)$.

Indeed, let k be the algebraic closure of k_0. Introducing an auxiliary indeterminate, consider Y (resp. X), the subscheme of $P = \mathbb{P}^N_k$ defined by the homogenized ideal of I (resp. of $\sqrt{I}$). In the Hilbert polynomial $\chi(O_Y(n)) = \sum_{i=0}^{N} b_i \binom{n+i}{i}$, we have $|b_i| \leq c_i$ with $c_i = A_{N-i}(0, \ldots, 0, 1; m)$ by 6.13.

Let $X = \bigcup X^q$ be the decomposition according to dimension. Then degree$(X^q) \leq e_q$ with $e_q = P_{N-q}(c_q, \ldots, c_N)$. Indeed, after cutting X (resp. Y) by a "general" linear space of codimension q, we may suppose that $q = 0$. Then clearly, degree$(X^q) = h^0(O_{X^q})$ and $h^0(O_{X^q}) \leq h^0(O_Y)$, and by 6.4 (iii), $h^0(O_Y) \leq e_q$.

Consequently, O_{X^q} is a $(0, \ldots, 0, e_q)$-sheaf by 6.10. It therefore follows from the injection $O_X \to \prod O_{X^q}$ that O_X is an $(e_0, \ldots, e_N)$-sheaf. Thus, in the Hilbert polynomial $\chi(O_X(n)) = \sum_{i=0}^{N} a_i \binom{n+i}{i}$[643], we have $|a_i| \leq f_i$[644] with $f_i = A_{N-i}(e_i, \ldots, e_N; 0)$ by 6.7. Finally, the ideal $\sqrt{I}$ of X is $R_N(m)$-regular with $R_N(m) = P_N(f_0, \ldots, f_N)$; whence the assertion by 1.3.

[print 662]

7. Appendix: The Hodge index theorem

Here we recover and complete some earlier results by a direct argument.

Theorem 7.1. Let X be an irreducible surface, proper over an algebraically closed field. Suppose there exists an invertible sheaf H on X such that $\langle c_1(H)^2 \rangle > 0$ (for example, H ample). Then an invertible sheaf L on X such that

$$\langle c_1(L) \cdot c_1(H) \rangle = 0 \quad \text{and} \quad \langle c_1(L)^2 \rangle \geq 0,$$

is numerically equivalent to 0. (Consequently, the intersection form on $A^1_{\mathrm{num}}(X)_{\mathbb{Q}}$ $(= [\mathrm{Pic}(X)/\mathrm{Pic}^\tau(X)] \otimes \mathbb{Q}$, by 4.6) is nondegenerate of type $(1, \rho - 1)$).

7.1.1. First suppose: X reduced (hence integral); H ample; $\langle c_1(L) \cdot c_1(H) \rangle = 0$; and $\langle c_1(L)^2 \rangle > 0$. Arguing by contradiction, take $m > 0$ such that $H_1 = L \otimes H^{\otimes m}$ is ample, and take n such that, putting $N = L^{\otimes n} \otimes H^{-1}$, we have that $\langle c_1(N) \cdot c_1(H_1) \rangle = n\langle c_1(L)^2 \rangle - m\langle c_1(H)^2 \rangle$ is > 0. Then we also have that $\langle c_1(N) \cdot c_1(H) \rangle = -\langle c_1(H)^2 \rangle$ is < 0, and that $\langle c_1(N)^2 \rangle = n^2\langle c_1(L)^2 \rangle + \langle c_1(H)^2 \rangle$ is > 0, which contradicts the following lemma.

Lemma 7.1.2. Let X be an integral surface, projective over an algebraically closed field. For an invertible sheaf N on X such that $\langle c_1(N)^2 \rangle > 0$, the following conditions are equivalent:

(i) For every ample sheaf H, we have $\langle c_1(N) \cdot c_1(H) \rangle > 0$.

(i') There exists an ample sheaf H such that we have $\langle c_1(N) \cdot c_1(H) \rangle > 0$.

(ii) There exists an integer $n > 0$ such that we have $H^0(X, N^{\otimes n}) \neq 0$.

[print 663]Indeed, (ii) implies that there exists an effective divisor D such that $N^{\otimes n} \simeq O_X(D)$, and $\langle c_1(N)^2 \rangle > 0$ implies that $D > 0$; whence (i).

To see that (i') implies (ii), we may clearly suppose that $H = O_X(1)$ is very ample; then there exists a closed integral, locally principal curve Y such that $H = O_X(Y)$. By Riemann's theorem, (i') implies that $H^1(N_Y^{\otimes n}(p)) = 0$ for $p \geq 0$ and $n \geq n_0$ with $n_0 = 2h^1(O_Y) - 1$. Since $H^2(N^{\otimes n}(p)) = 0$ for $p \gg 0$, the exact sequence

$$H^1(N_Y^{\otimes n}(p)) \longrightarrow H^2(N^{\otimes n}(p-1)) \longrightarrow H^2(N^{\otimes n}(p)) \longrightarrow 0$$

[645] gives $H^2(N^{\otimes n}) = 0$ for $n \geq n_0$. Thus, for $n \gg 0$, we have

$$h^0(N^{\otimes n}) = \chi(N^{\otimes n}) + h^1(N^{\otimes n}) \geq \chi(N^{\otimes n}) = 1/2\langle c_1(N)^2 \rangle n^2 + \ldots > 0,$$

which completes the proof of the lemma.

7.1.3. Now suppose in 7.1 that there exists an invertible sheaf M on X such that $\langle c_1(L)\cdot c_1(M)\rangle \neq 0$, and argue by contradiction. Replace X by the underlying reduced scheme of a Chow presentation of X, which is allowed by the projection formula X 4.4.4, and let G be ample on X. Take $m > 0$ such that $H_1 = G^{\otimes m} \otimes M$ is ample, and such that $\langle c_1(L) \cdot c_1(H_1)\rangle \neq 0$; take p, q $(q \neq 0)$ such that, putting $L_1 = L^{\otimes p} \otimes H^{\otimes q}$, we have that $\langle c_1(L_1) \cdot c_1(H_1)\rangle = p\langle c_1(L)\cdot c_1(H_1)\rangle + q\langle c_1(H)\cdot c_1(H_1)\rangle$ is 0; then we also have that $\langle c_1(L_1)^2\rangle = p^2\langle c_1(L)^2\rangle + q^2\langle c_1(H)^2\rangle$ is > 0; this returns us to case 7.1.1, with $L = L_1$ and $H = H_1$.

7.1.4. Finally, consider the general case of 7.1. Still arguing by contradiction, suppose that L is not numerically equivalent to 0; in addition, replace X by the normalization of X_{red}, so that X has [print 664]only finitely many singular points, a reduction justified by the projection formula X 4.4.4 and by 4.5 (ii). There therefore exists a closed integral curve Y on X such that $\langle c_1(L) \cdot Y\rangle \neq 0$, and Y is locally principal outside a finite set Z. Let $f : X' \to X$ be the blowup of the ideal I of O_X[646]. The ideal $M = I \cdot O_{X'}$ is invertible, and defines a subscheme of X' which is the union of two subschemes Y', Y'' such that $f(Y') = Y$ and $f(Y'') \subset Z$. By the projection formula X 4.4.4, we have that $\langle c_1(f^*L) \cdot c_1(M)\rangle = -\langle c_1(f^*L) \cdot Y'\rangle - \langle c_1(f^*L) \cdot Y''\rangle$ equals $-\langle c_1(L) \cdot Y\rangle - 0 \neq 0$, and we may replace X, L, H by X', f^*L, f^*H; we are therefore reduced to the preceding case, and the proof of 7.1 is complete.

Remark 7.2. It is unknown whether an irreducible proper surface (over an algebraically closed field), which possesses an invertible sheaf H such that $\langle c_1(H)^2\rangle > 0$, is necessarily projective. It is also unknown whether, for every irreducible proper surface, the intersection form on $A^1_{\mathrm{num}}(X)_{\mathbb{Q}}$ is nondegenerate; it follows from 7.1 that, when it is degenerate, it is necessarily negative (nondefinite).

Corollary 7.3. Under the conditions of 7.1, let M be an invertible sheaf on X which is not numerically proportional to[647] H, and let $d = \langle c_1(M), c_1(H)\rangle$ and $h = \langle c_1(H)^2\rangle$. Then, $\langle c_1(M)^2\rangle < d^2/h$.

Indeed, put $L = M^{\otimes h} \otimes H^{\otimes -d}$[648]. Since $\langle c_1(L) \cdot c_1(H)\rangle = 0$, by 7.1, $\langle c_1(L)^2\rangle < 0$; whence the assertion.

[print 665]**Corollary 7.4.** Let X be an irreducible scheme of dimension $r \geq 2$, projective over an algebraically closed field, and let H be an ample sheaf on X. Then:

(i) For an invertible sheaf L on X to be numerically equivalent to 0, it is (necessary and) sufficient that

$$\langle c_1(L) \cdot c_1(H)^{r-1}\rangle = 0 \quad \text{and} \quad \langle c_1(L)^2 \cdot c_1(H)^{r-2}\rangle \geq 0.$$

(ii) For every invertible sheaf M on X which is not numerically proportional to[649] H, we have

$$\langle c_1(M)^2 \cdot c_1(H)^{r-2}\rangle < \langle c_1(M) \cdot c_1(H)^{r-1}\rangle^2 / \langle c_1(H)^r\rangle.$$

Indeed, (ii) follows immediately from (i), by the argument of 7.3. To establish (i), let Y be a closed integral curve in X; we wish to prove that $\langle c_1(L) \cdot Y\rangle = 0$. Embed X in $\mathbb{P}^N_k$ by means of $H^{\otimes n}$ $(n > 0)$. Let $I = \bigoplus I_\nu$ be the homogeneous ideal of Y, and let ν_0 be an integer such that I_{ν_0} generates I_ν for $\nu \geq \nu_0$. Then the hypersurfaces Z of degree ν_0 that contain Y have no common points outside Y. Arguing by induction on $r = \dim(X)$, we find, using Bertini's theorem, $r - 2$ hypersurfaces $Z_1, \ldots, Z_{r-2}$ of degree ν_0 that contain Y, such that $X' = Z_1 \cap \ldots \cap Z_{r-2} \cap X$ is irreducible. Since $\langle c_1(L) \cdot Y\rangle = \langle c_1(L_{X'}) \cdot Y\rangle$, it suffices to prove that $L_{X'}$ is numerically equivalent to 0 on X'. Now, by the projection formula, we have $\langle c_1(L_{X'}) \cdot c_1(H_{X'})\rangle = 0$ and $\langle c_1(L_{X'})^2\rangle \geq 0$; thus the assertion follows from 7.1.

Corollary 7.5. Let k be an algebraically closed field, Y a projective k-scheme equipped with an ample sheaf H, and X the zero scheme of a section φ of H. Suppose that, for every irreducible component Y^i of Y, [print 666]the irreducible components of $X \cap Y^i = V(\varphi|Y^i)$ have dimension ≥ 2. Then the morphism $A^1_{\mathrm{num}}(Y) \longrightarrow A^1_{\mathrm{num}}(X)$ is injective.

Indeed, since we may clearly suppose that Y is integral,[650] the assertion follows from 7.4 (i), by the projection formula.

BIBLIOGRAPHY

[1] Abhyankar S. : Resolution of singularities of arithmetical surfaces, in : Arithmetical Algebraic Geometry, Purdue University, Harper-Row, New York 1965.

[2] Mumford D : Pathologies III, Amer. J. Math 1967 p. 94-104.[651]

[3] Grothendieck A. and Dieudonné J.[652] EGA, Eléments de Géométrie Algébrique, Pub. Math. Inst des Hautes Etudes Sci., 1960.

[4] Grothendieck A. : FGA, Fondements de la géométrie algébrique, Extracts from the Séminaire Bourbaki 1957-1962.

[5] Grothendieck A. et alii[653], SGA, Séminaire de géométrie algébrique, 1960-66, I.H.E.S.[654]

[6] Jussila O. : X, Formalisme des intersections sur les schémas algébriques propres, Exposé X in this seminar.

[7] Raynaud M. : XII, Un théorème de représentabilité relative sur le foncteur de Picard, Exposé XII in this seminar.

[print 667]

Exposé XIV
OPEN PROBLEMS IN INTERSECTION THEORY

by A. Grothendieck

In this exposé we review some of the problems connected with the questions encountered in this seminar that deserve to be studied. We leave aside the problems connected with Serre's local theory of intersections [14], which we have not had occasion to use in the Seminar, and refer in particular to loc. cit. for the two fundamental conjectures that remain open in this direction (loc. cit. Chap. V § 4). We also pass over the problems, very important for arithmetic applications, connected with the structure of groups of cycle classes, especially on a smooth projective algebraic scheme; for a discussion of these problems we refer to S. KLEIMAN, Algebraic cycles and the Weil conjectures in Dix exposés sur la cohomologie des schémas (Masson-North Holland Publishing Cie). In no. 4 of this exposé we also give some supplements to the Seminar, concerning in particular the Chow ring, which had not found a place in the preceding exposés.

1. Operations Λ^i in the derived category $D(X)(*)$

Let X be a locally ringed topos, and let $K^\bullet(X)$ be the ring of classes of perfect complexes on X (Exp. IV). We would like to define on $K^\bullet(X)$ a λ-ring structure (Exp. V), functorial in X, such that the natural homomorphism

$$K^\bullet_{\text{naif}}(X) \longrightarrow K^\bullet(X)$$

(where the first member is the ring of classes of locally free Modules on X, endowed with the λ-structure defined in Exp. V) be a λ-[655]homomorphism. For this purpose[656], one would need to define functors (generally nonadditive) of "exterior power"[657]

$$\Lambda^i : D^-(X) \longrightarrow D^-(X),$$

taking pseudo-coherent (resp. perfect) complexes to pseudo-coherent (resp. perfect) complexes and having the formal properties needed, with respect to exact triangles, to define a λ-structure on $K^\bullet(X)$ by the formula

$$\lambda^i(\mathrm{c}\ell^\bullet(L_\bullet)) = \mathrm{c}\ell^\bullet(\Lambda^i L_\bullet). \tag{1.1}$$

[print 668]When X is of characteristic zero, i.e. its structure sheaf is a sheaf of Q-algebras, the problem is solved simply by taking

$$\Lambda^i L_\bullet = (\overset{i}{\otimes}{}^{\mathbb{L}} L_\bullet)^{\text{alt}}, \tag{1.2}$$

where the i-th tensor power (in the sense of derived categories) of $L_\bullet$ is regarded as an object of $D^-(\underline{O}_i)$, with $\underline{O}_i$ the group algebra of the symmetric group $\mathfrak{S}_i$ with coefficients in $\underline{O}_X$, and where the superscript alt denotes passage to the "alternating part under the action of $\mathfrak{S}_i$" (an operation that makes sense by the characteristic-zero hypothesis). A formula of this type was already used in [8] (cited below as [RRR]) for the first proof of the Riemann–Roch theorem.

Without a hypothesis on the characteristic, when $L_\bullet$ is zero in degrees > 0, Dold–Puppe [6] shows that $L_\bullet$ is canonically homotopic to the complex associated with a semi-simplicial Module

(*) cf. the note on page 23.

$L'_\bullet$ canonically associated with $L_\bullet$, and a fairly natural candidate for $\Lambda^i L_\bullet$ (for $L_\bullet$ flat, say) would be the complex associated with the semi-simplicial Module $\Lambda^i L'_\bullet$ (whose component of degree n is $\Lambda^i L'_n$). But to extend the definition to complexes of $D^-(X)$ no longer assumed to be zero in degrees > 0, one would have to study the behaviour of the preceding operation with respect to the translation operation $L_\bullet \mapsto L_\bullet[1]$ (or suspension, in the terminology of [6]); unfortunately, there does not seem to be an isomorphism $\Lambda^i(L_\bullet[1]) \simeq (\Lambda^i L_\bullet)[i]$ that would make the desired extension trivial.

2. The Riemann–Roch formula without projectivity hypotheses

2.1. Once a natural λ-structure on the $K^\bullet(X)$ is available, it is possible to formulate the Riemann–Roch theorem, in the form considered in Exp. VIII, for any proper complete-intersection morphism $f : X \to Y$ (and even, more generally, for a relative scheme X over a locally ringed topos Y, with X still subject to the corresponding relative conditions). The Todd class of f is defined by means of the class in $K^\bullet(X)$ of the relative cotangent complex T_f for f, constructed in Exp. 0 4.4; this complex is perfect here by the local hypothesis on f. It seems clear that a proof of the Riemann–Roch formula in this general case will require the introduction of essentially new ideas, and it is possible that these same ideas could provide the key [print 669]to a proof of the analogous formula in the complex analytic (or rigid analytic!) case. Recall that in the complex analytic case, only the case of a locally free sheaf F on a compact nonsingular variety X over a point Y is currently known (by the Atiyah–Singer index formula); the case in which F is merely assumed coherent still seems to lie beyond the techniques known so far.

2.2. As an essential ingredient of the Riemann–Roch formula, let us also recall the finiteness problem raised in Exp. III: for a proper morphism $f : X \to Y$ and a complex $L_\bullet$ on X that is pseudo-coherent relative to Y, is $Rf_*(L_\bullet)$ pseudo-coherent? A general proof of this conjecture will probably use algebraic ideas that will make it possible to prove Grauert's finiteness "theorem" [7] under considerably more general conditions than those of loc. cit., starting directly with topological $\mathbb{C}$-algebra structures on base spaces (or topoi) Y more general than analytic spaces (including the case of a locally compact space Y with the sheaf of continuous complex functions, or a C^m manifold with the sheaf of C^m complex functions, or a real analytic space, etc.).

2.3. Finally, in these questions one would need to replace the properness hypothesis on the morphism f by the hypothesis that the cohomology sheaves $H^i(L_\bullet)$ of the complexes under consideration on X have their supports proper[658] over Y.

3. A Riemann–Roch formula "without denominators" for an immersion

3.1. Let $f : Y \to X$ be a regular closed immersion (Exp. VII), with conormal sheaf $\underline{N}$. For simplicity, suppose that X admits an ample invertible sheaf. The Riemann–Roch formula for $y \in K^\bullet(Y)$:

$$\mathrm{ch}_X(f_*(y)) = f_*(\mathrm{ch}_Y(y)\,\mathrm{Todd}(-\check{\underline{N}})),$$

together with the general formula (Exp. VII):

$$f_*(y')\,f_*(y'') = f_*(y'y''c_d(\underline{N}))^{[659]}, \qquad y', y'' \in \mathrm{Gr}(Y)_{\underline{Q}},$$

(with f assumed to have codimension d), makes it possible to express the Chern classes of $f_*(y)$ in $\mathrm{Gr}^\bullet(X)_{\mathbb{Q}}$ in the form [print 670]

$$c_i(f_*(y)) = f_*(P_i((c_\alpha(y))_\alpha, (c_\beta(\underline{N}))_\beta)),^{[660]} \tag{3.1}$$

where P_i is a universal polynomial in the $c_\alpha(y)$, $c_\beta(\underline{N})$, a polynomial whose coefficients are a priori rational. In fact, one finds [RRR] that these polynomials have integer coefficients. This raises the question of the validity of a Riemann–Roch formula "without denominators". In fact, such a formula for $c_i(f_*(y))$ cannot in general be sought in $\mathrm{Gr}^\bullet(X)$ itself, since the homomorphism $f_* : K^\bullet(Y) \to K^\bullet(X)$ is not itself compatible with the λ-filtrations up to a shift by d (4.6), but is so only up to torsion; thus one cannot define a homomorphism $f_* : \mathrm{Gr}^\bullet(Y) \longrightarrow \mathrm{Gr}^\bullet(X)$ giving the natural homomorphism $f_* : \mathrm{Gr}^\bullet(Y)_\mathbb{Q} \to \mathrm{Gr}^\bullet(X)_\mathbb{Q}$[661] and allowing formula (3.1) to be interpreted as a formula in $\mathrm{Gr}^\bullet(X)$, and not merely in $\mathrm{Gr}^\bullet(X)_\mathbb{Q}$. But when X is assumed regular, so that $K^\bullet(X) \xrightarrow{\sim} K_\bullet(X)$[662] (IV 2.5), one can endow $K^\bullet(X)$ with the topological filtration by the codimension of supports, inherited from the corresponding filtration of $K_\bullet(X)$. It is plausible that this filtration is compatible with the ring structure, a fact that can be proved at least [RRR] when X is smooth over a field k, using the theory of the Chow ring [4]. When this is so, the associated graded

$$\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$$

is a graded ring, and, also assuming Y regular, we have done what is needed to obtain a natural homomorphism of degree d of graded groups

$$f_* = f_*^{\mathrm{top}} : \mathrm{Gr}^\bullet_{\mathrm{top}}(Y) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(X).$$

Since the λ-filtration of $K^\bullet(X)$ is finer than the topological filtration by codimension just considered (X 1.3.3), one obtains a canonical homomorphism of graded groups (and even of graded rings)

$$\mathrm{Gr}^\bullet(X) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(X),$$

which is moreover an isomorphism mod torsion (Exp. VIII). This allows us, as in [RRR], to consider a theory of Chern classes on X with values in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$, and likewise on Y, and thus to interpret formula (3.1) as a formula in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$. In this form, the formula is proved in [RRR] when X is of characteristic zero, at least when X is smooth over a field (a hypothesis serving only to ensure that the topological filtration of $K^\bullet(X)$ is compatible with the ring structure). The characteristic-zero hypothesis is used essentially [print 671]to obtain formulas of the type (1.1), (1.2), and essentially the same proof should apply without a hypothesis on the characteristic, once the question raised in no. 1 has been answered affirmatively.

3.2. When X is smooth over a field, one also has the Chow ring

$$A(X),$$

and a theory of Chern classes on X with values in $A(X)$ [9], finer than the theory with values in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ just introduced—these latter Chern classes being the images of the Chow-style Chern classes under the canonical surjective homomorphism (cf. 4.1)

$$\phi : A(X) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(X).$$

This homomorphism is an isomorphism mod. torsion, but not an isomorphism in general (4.7). With this understood, for smooth X and Y the inclusion $f : Y \to X$ again defines a homomorphism of degree d of graded groups

$$f_* : A(Y) \longrightarrow A(X),$$

which allows formula (3.1) to be interpreted as a formula in $A^i(X)$(*). This formula reduces to $0 = 0$ for $i < d$ and is therefore of interest only for $i \geq d$. Even for $i = d = 2$ this formula has not been proved at present (cf. 4.3), nor has the less precise formula deduced from it by replacing

(*) The formula, as well as the variant mentioned in 3.3, has since been proved by P. JOUANOLOU (Nov. 1969). Cf.

rational equivalence of algebraic cycles with algebraic equivalence. Unlike the analogous formula in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$, there seems to be no very convincing reason to suppose that the formula without denominators must hold in $A(X)$. In view of our almost total ignorance of the nature of the groups $A^i(X)$ for $i \geq 2$, a good way to test it seems to be to test the formulas that can be deduced from it in groups $A^1(Z) = \mathrm{Pic}(Z)$, for example by applying $g_* : A(X) \longrightarrow A(Z)$ to both members of (3.1), where $g : X \to Z$ is a proper morphism into a smooth quasi-projective Z of dimension $\dim X - i + 1$.

3.3. Finally, to give a meaning to (3.1), one can also use the cohomological theory of Chern classes (SGA 5 VII), whenever Gysin homomorphisms

$$f_* : H^p(Y, \mu_n^{\otimes i}) \longrightarrow H^{p+2d}(X, \mu_n^{\otimes (i+d)})$$

are available (cf. no. 6); this is the case at least when Y and X are both smooth over a scheme S (by the theorem of relative cohomological purity [print 672]SGA 4 XVI 3.7), and, conjecturally, when Y and X are both regular (by the absolute cohomological purity theorem, proved at least for excellent X of characteristic zero (SGA 4 XIX 3.2)). When X and Y are smooth over the field of complex numbers, one can also use the integral Chern class on the locally compact space $X(\mathbb{C})$ of complex points of X. Formula (3.1) is established in this case by Atiyah and Hirzebruch [1], who work more generally with an immersion of complex analytic varieties (using the "naive" $K^\bullet$); it follows that the ℓ-adic formula (3.1) holds whenever Y and X are smooth over an algebraically closed field of characteristic zero, using the "Lefschetz principle" and the general results of SGA 4. This therefore makes the validity of the cohomological formula (3.1) extremely plausible; but observe that it has not been proved at present even when X and Y are smooth over a field k of characteristic zero, when[663] k is not algebraically closed.

3.4. One can also work with the cohomological theory of Chern classes in the De Rham or Hodge style [13]. The question of the validity of (3.1) in this setting arises only when X is not of characteristic zero, since in any case (no. 6) the formula in question is known to hold modulo torsion. For smooth quasi-projective schemes over a field k, the De Rham or Hodge version of formula (3.1) would moreover follow from the cycle-class version of formula (3.1) mod. algebraic equivalence, so that the De Rham or Hodge cohomological version gives another accessible way to test the validity of the "cycle-class" version of (3.1).

3.5. The proof [1] of the transcendental cohomological version of (3.1) is essentially transcendental in nature, using tubular neighbourhoods of Y in X. This suggests that a proof of a formula of type (3.1) might be connected with the systematic consideration of groups $K^\bullet$ and cohomology groups with support in Y, which might provide a sufficient technical means of "localization around Y".

4. Relations between $K^\bullet(X)$ and the Chow ring $A(X)$

4.1. For simplicity, suppose that X is a smooth quasi-projective scheme over a field k (cf no. 8 for a more general case), so that we [print 673]have the Chow ring $A(X)$ of cycle classes modulo rational equivalence [4]. As variants of this ring, which is moreover defined under substantially more general considerations, there are the rings $\mathrm{Gr}^\bullet(X)$ (the associated graded of the λ-filtration of $K^\bullet(X)$) and $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ (the associated graded of the filtration of $K^\bullet(X) \simeq K_\bullet(X)$ by the codimension of supports). There are theories of Chern classes with values in each of these three rings (cf. [9] for $A(X)$, and this seminar for $\mathrm{Gr}^\bullet(X)$, which maps to $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$). There is also a cohomological theory (ℓ-adic, say, with ℓ prime to the characteristic of k) of Chern classes (SGA 5 VII):

$$c_i(\underline{E}) \in H^{2i}(X, \underline{Z}_\ell(i)),$$

(where $\underline{E}$ is a vector bundle on X, or indeed any element of $K^\bullet(X)$). These cohomological theories are related by the canonical homomorphisms (compatible with the multiplicative structures)

$$\begin{array}{ccc} H^{2*}(X, \underline{Z}_\ell(*)) & & \\ \uparrow{\scriptstyle \gamma} & & \\ A(X) & \xrightarrow{\quad\phi\quad} & \mathrm{Gr}^\bullet_{\mathrm{top}}(X) \\ & & \uparrow{\scriptstyle j} \\ & & \mathrm{Gr}^\bullet(X), \end{array} \tag{4.1}$$

where we have set

$$H^{2*}(X, \underline{Z}_\ell(*)) = \coprod_i H^{2i}(X, \underline{Z}_\ell(i)).$$

The homomorphism j comes from the fact that the λ-filtration of $K^\bullet(X)$ is finer than the topological filtration by the codimension of the support (Exp. X), and it is an isomorphism modulo torsion (Exp. VII 4.11). The homomorphism ϕ is defined, for example, in [RRR], and associates to the class of the cycle of codimension i associated with a closed subscheme Y of X of codimension i the element $c\ell^\bullet(\underline{O}_Y)$ of $\mathrm{Filt}^i_{\mathrm{top}}(K^\bullet(X))$ mod $\mathrm{Filt}^{i+1}_{\mathrm{top}}(K^\bullet(X))$. It is a surjective homomorphism, which we shall also see (4.2) to be an isomorphism modulo torsion. Thus, modulo torsion the three rings $A(X)$, $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ and $\mathrm{Gr}^\bullet(X)$ are canonically isomorphic, so a posteriori the Riemann–Roch formulas expressed using one or another of these rings[664] (tensored with $\mathbb{Q}$) are equivalent. The graded ring $\mathrm{Gr}^\bullet(X)$, used particularly in this seminar as a substitute for the Chow ring, has the appreciable advantage of being defined for every locally ringed topos and, in particular, for every scheme [print 674](regular or otherwise). It has, on the other hand, the disadvantage (4.6) relative to $A(X)$, $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ that, as it stands, it is not covariant with respect to proper morphisms (even for smooth quasi-projective schemes over a field); it acquires this property only after tensoring with $\mathbb{Q}$. Another advantage of $A(X)$ over its competitors $\mathrm{Gr}^\bullet_{\mathrm{top}}$ and $\mathrm{Gr}^\bullet(X)$ is the existence of a homomorphism from $A(X)$ to $H^{2*}(X, \underline{Z}_\ell(*))$ that allows cohomological Chern classes to be interpreted as images of Chern classes with values in $A(X)$. In general this homomorphism does not factor through $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ (4.7), and it is at least doubtful that there is in general a homomorphism from $\mathrm{Gr}^\bullet(X)$ to $H^{2*}(X, \underline{Z}_\ell(*))$ taking Chern classes to Chern classes (Cf. no. 5)[665]. Of course, these difficulties disappear after tensoring with $\underline{Q}$, and one obtains a homomorphism

$$\mathrm{Gr}^\bullet_{\mathrm{top}}(X) \longrightarrow H^{2*}(X, \underline{Q}_\ell(*)) = H^{2*}(X, \underline{Z}_\ell(*)) \otimes_{\underline{Z}_\ell} \underline{Q}_\ell \tag{4.2}$$

taking Chern classes to Chern classes.

4.2. To establish that the surjective homomorphism ϕ of (4.1) is an isomorphism modulo torsion, one uses the Chern homomorphism (deduced from the theory of Chern classes [9])

$$K^\bullet(X) \longrightarrow \mathrm{Ch}(A(X))$$

(where the second member is the Chern ring of $A(X)$ defined in Exp. V), which is, as is easily seen, compatible with the topological filtration of $K^\bullet(X)$ and the natural filtration of $\mathrm{Ch}(A(X))$, and gives, on passing to the associated graded groups, a homomorphism of graded groups

$$\psi : \mathrm{Gr}^\bullet_{\mathrm{top}}(X) \longrightarrow A(X). \tag{4.3}$$

With this understood, we shall prove the formulas

$$\begin{cases} \psi\phi(x) = (-1)^{i-1}(i-1)!\, x & \text{mod. torsion} \quad \text{for } x \in A^i(X) \\ \phi\psi(x) = (-1)^{i-1}(i-1)!\, x & \text{mod. torsion} \quad \text{for } x \in \mathrm{Gr}^i_{\mathrm{top}}(X), \end{cases} \tag{4.4}$$

the first of which moreover implies the second, since ϕ is surjective. These formulas clearly imply that ϕ and ψ are isomorphisms modulo torsion and that, up to a factor $(-1)^{i-1}(i-1)!$ in degree i, they are inverses of one another.

To prove the first formula (4.4), one can clearly suppose that x is defined by an irreducible closed subscheme Y of X of codimension i, in which case the formula is equivalent to [print 675]

$$c_i(\underline{O}_Y) = (-1)^{i-1}(i-1)!\ \mathrm{cl}(Y) \quad \text{mod torsion}, \tag{4.5}$$

where in the first member $\underline{O}_Y$ is regarded as a coherent sheaf on X, thus defining an element of $K^\bullet(X)$, and in the second member $\mathrm{c}\ell(Y)$ denotes the class of Y in $A^i(X)$[666]. Since $A^i(X)$ does not change on removing from X a closed subset of codimension $> i$, we are reduced to the case in which Y is regular, hence smooth over the base field k when the latter is perfect. But in this latter case, formula (4.5) is precisely the Riemann–Roch formula for the inclusion $Y \to X$ and the element 1 of $K^\bullet(Y)$, written using the theory of Chern classes with values in the Chow ring $A(X)$. This formula is established in [2] under the conditions considered here, by a method very close to the one used in this seminar for the analogous formula using $\mathrm{Gr}^\bullet(X)$. This therefore establishes, at least for perfect k, formula (4.4) and consequently the fact that ϕ is an isomorphism mod. torsion (which in turn implies, a posteriori, that the Riemann–Roch formula "with denominators" for a proper morphism of smooth algebraic schemes over k is essentially the same whether written using $A(X)$ as in [2] or using $\mathrm{Gr}^\bullet(X)$ as in this seminar!). The case with no hypothesis on k reduces easily to the case in which k is algebraically closed, using the fact that, for a finite extension k' of degree n of k, the kernel of the natural map $A(X) \xrightarrow{f^*} A(X \otimes_k k')$[667] is annihilated by n (as is seen by using the trace homomorphism $f_* : A(X \otimes_k k') \to A(X)$, satisfying $f_* f^* = n\ \mathrm{id}$).

4.3. It seems quite plausible that formulas (4.4) hold as they stand and not only modulo torsion, or, equivalently, that formula 4.5. holds not only modulo torsion. This question is a very special case of the validity in $A(X)$ of the "Riemann–Roch formula without denominators" (3.1) for an immersion $Y \to X$ (namely, the case $i = d$). The validity of this special case may be regarded as quite plausible. Indeed, we already pointed out in no. 3 that, when k is of characteristic zero, the Riemann–Roch formula without denominators holds at least in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$; this implies in that case that both members of (4.5) have the same image in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$, or equivalently that the second of formulas (4.4) holds; this result will extend of itself to arbitrary characteristic once [print 676]the question raised in no. 1 has been satisfactorily resolved. In support of formula (4.5) in $A(X)$ itself, observe that it is clear a priori that the restriction of the first member to $X - Y$ is zero; thus, with Y assumed irreducible, the first member must a priori be an integer multiple of $\mathrm{c}\ell(Y)$:

$$c_i(Y) = n(Y)\mathrm{c}\ell(Y).$$

This already proves the exact formula (4.5) when $\mathrm{c}\ell(Y)$ is not a torsion element of $A^i(X)$. It is immediate that this is so when X is projective (and not merely quasi-projective) over the field k: note that for a polarization $\xi \in A^1(X)$ of X, one has $\deg(Y \cdot \xi^{\dim Y}) > 0$[668]; this proves the desired formula in this case. Consequently, this formula also holds whenever X is isomorphic to an open subscheme of a smooth projective scheme over k, which will be the case when a suitable solution to the resolution-of-singularities problem is available. This is so, in particular, when k has characteristic zero, by Hironaka [11].

The validity of the exact formula (4.5), hence of (4.4), would imply that the kernel of $\phi^i : A^i(X) \longrightarrow \mathrm{Gr}^i_{\mathrm{top}}(X)$ is annihilated by $(i-1)!$, and in particular that $\phi^2 : A^2(X) \to \mathrm{Gr}^2_{\mathrm{top}}(X)$ is an isomorphism. (That ϕ^1 is an isomorphism already follows immediately from X 5.3.2.) Unless I am mistaken, the fact that ϕ^2 is an isomorphism has not been proved at present, even in the

(*) It is therefore established, thanks to P. JOUANOLOU, for X smooth and quasi-projective over a field (cf. note p.5).

special case $\dim X = 3$. In any event, the writer was hasty in asserting the validity of (4.4) and (4.5) in [9, no. 4, 3°].

4.4. At present, the main advantage of $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ (or $\mathrm{Gr}^\bullet(X)$) over the Chow ring $A(X)$ is that it is better suited to explicit computation of certain intersection formulas between cycles, in the case of "excess" intersections, i.e. of excessively large dimension. Indeed, defining the intersection class by means of Chow's lemma, which allows a cycle to be moved sufficiently within its class, does not readily lend itself to computation. A striking example is the following fundamental formula [Exp. VII 2.7] for a closed immersion $f : Y \to X$[669] of smooth quasi-projective schemes over k:

$$f^* f_*(y) = y\, c_d(\underline{N}), \tag{4.6}$$

a formula valid in $\mathrm{Gr}^\bullet_{\mathrm{top}}(Y)$[670], with y in $\mathrm{Gr}^\bullet_{\mathrm{top}}(Y)$ and $\underline{N}$ the conormal sheaf [print 677]of Y in X, assumed everywhere to have rank d. In loc. cit., this formula follows from a very simple computation of Tor sheaves. By what we saw in 4.2, the preceding formula implies the same formula in $A(Y)$, but only modulo torsion. It is quite extraordinary that formula (4.6) in $A(Y)$ itself has not been proved at present(*), nor even the special case obtained by taking $y = 1$, which gives the self-intersection of Y:

$$f^*(\mathrm{cl}(Y)) = c_d(\underline{N}), \tag{4.7}$$

even when k is the field of complex numbers!

An analogous problem arises when one attempts to compute the Chow ring of the variety X' obtained by blowing up X along Y, imitating the method used to compute $\mathrm{Gr}^\bullet(X')_{\mathbb{Q}}$ [Exp. VII 4.8], which also works (without tensoring with $\mathbb{Q}$) for the computation of $\mathrm{Gr}^\bullet_{\mathrm{top}}(X')$. The key formula for this determination (with the notation of loc. cit.) is

$$f^*(i_*(y)) = j_*(g^*(y) c_{d-1}(\check{\underline{F}})), \tag{4.8}$$

a formula currently established in $\mathrm{Gr}^\bullet_{\mathrm{top}}(X')$ (for $y \in \mathrm{Gr}^\bullet_{\mathrm{top}}(Y)$) by a computation of Tor (Exp. VII 3.4), whose validity in $A(X')$ (for $y \in A(Y)$) remains to be established.

Observe that even the ℓ-adic analogues of formulas (4.6), (4.7), (4.8) have not been proved at present; when y is the cohomology class of an algebraic cycle, (4.6) and (4.8) hold in any case mod. torsion.

4.5. In this section and the following ones, we shall give examples in which, in diagram (4.1), the homomorphism j, respectively the homomorphism ϕ, is not an isomorphism. For a given X, it clearly amounts to the same thing to say that

$$j : \mathrm{Gr}^\bullet(X) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(X)$$

is an isomorphism, or a monomorphism, or an epimorphism. Moreover, the image of j is the subring of $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ generated by the Chern classes of locally free sheaves on X. When $K^\bullet(X)$ is generated as an algebra over $\mathbb{Z}$ by the classes of invertible sheaves on X, one sees immediately that the subring of $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ under consideration is also the subring generated by $\mathrm{Gr}^1_{\mathrm{top}}(X)$, the component of degree 1 of $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$. [print 678]Now take for X a flag variety

$$X = G/B,$$

where G is a simply connected semisimple algebraic group over the algebraically closed field k, and B is a Borel subgroup [5]. When $k = \mathbb{C}$, the Bruhat cell decomposition of X [5, Exp. 13] implies that the canonical homomorphism

$$(*) \qquad K^\bullet(X) \longrightarrow K(X(\mathbb{C}))$$

(*) cf. the note on page 23.
(**)

is an isomorphism, as is moreover the canonical homomorphism

$$(**) \qquad A(X) \longrightarrow H^*(X(\mathbb{C}), \mathbb{Z}),$$

where $X(\mathbb{C})$ is the compact space of complex points of X, endowed with its usual topology induced by the topology of $\mathbb{C}$. On the other hand, L. HODGKIN [12] has proved that $K(X(\mathbb{C}))$ is indeed generated by the classes of invertible sheaves on X. The same therefore holds for $K^\bullet(X)$ itself, and it is moreover not difficult to deduce, by a specialization technique using split semisimple groups over $\mathbb{Z}$ (SGA 3), that the same result holds over every algebraically closed field k. Moreover, the cell decomposition of X (or the bijectivity of $(**)$ when $k = \mathbb{C}$) readily implies that $A(X)$ is torsion-free, so by 4.2 the homomorphism $\phi : A(X) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ is an isomorphism. These facts show that j is an isomorphism if and only if $A(X)$ is generated by its elements of degree 1, or again, when $k = \mathbb{C}$, if and only if the integral cohomology of $X(\mathbb{C})$ is generated by the elements of degree 2, i.e. if the classifying space $B_{G(\mathbb{C})}$ is torsion-free. It is also known [15, p. 217 th. B (1)⇔(2), and 2.5] that this is so if and only if G is "torsion-free", or equivalently if G is isomorphic to a product of groups $\mathrm{S}\ell(n)$ and $\mathrm{Sp}(m)$. Thus it suffices to take for G a group $\mathrm{SO}(n)$ with $n \geq 7$, or an exceptional simple group, to obtain an example in which j is not an isomorphism.

4.6. Observe that, in this example, there is no doubt that the "right" ring is the torsion-free ring $A(X) \simeq \mathrm{Gr}^\bullet_{\mathrm{top}}(X)$, with the cells of the Bruhat decomposition as a basis, a ring also isomorphic to the integral cohomology of $X(\mathbb{C})$ when $k = \mathbb{C}$; on the other hand, the ring $\mathrm{Gr}^\bullet(X)$ appears to have torsion of a rather pathological nature. This example therefore confirms the view that the invariant $\mathrm{Gr}^\bullet(X)$ is of little [print 679]interest except modulo torsion, i.e. after tensoring with $\mathbb{Q}$. At the same time, this example shows that only after tensoring with $\mathbb{Q}$ is $\mathrm{Gr}^\bullet(X)$ covariant with respect to proper morphisms, even when restricted to closed immersions of smooth quasi-projective schemes. Indeed, for X such that j is not an isomorphism, there is in X an integral closed subscheme Y such that $\mathrm{c}\ell^\bullet(Y) \in K^\bullet(X)$ does not lie in $\mathrm{Filt}^i(X)$, where $i = \mathrm{codim}(Y, X)$. We may take Y of minimal dimension, so that $\mathrm{Filt}^{i+1}(X) = \mathrm{Filt}^{i+1}_{\mathrm{top}}(X)$. With Z the singular locus of Y and $X' = X - Z$, $Y' = Y - Z$, we conclude that $\mathrm{c}\ell^\bullet(Y') \in K^\bullet(X')$ does not lie in $\mathrm{Filt}^i(X')$ (since $K^\bullet(X) \to K^\bullet(X')$ is surjective and compatible with the augmentations, and its kernel is $\mathrm{Im}\big(K(Z) \longrightarrow K(X)\big) \subset \mathrm{Filt}^{i+1}_{\mathrm{top}}(X) = \mathrm{Filt}^{i+1}(X)$[671]). Now consider the inclusion

$$f : Y' \longrightarrow X',$$

which is a closed immersion of codimension i of smooth quasi-projective schemes over k, such that

$$f_* : K^\bullet(Y') \longrightarrow K^\bullet(X')$$

is not compatible with the λ-filtrations (up to a shift by i), since $f_*(1)$ does not lie in $\mathrm{Filt}^i(X')$.

4.7. Keeping the notation of 4.5, observe now that (Exp. IX) shows that the cited theorem of HODGKIN is equivalent to the existence of an isomorphism

$$K^\bullet(G) \simeq \mathbb{Z}, \qquad \text{therefore} \qquad \mathrm{Gr}^\bullet_{\mathrm{top}}(G) \simeq \mathbb{Z},$$

the homomorphism $K^\bullet(G/B) \to K^\bullet(G)$ being a priori surjective, with kernel the ideal generated by the $\mathrm{c}\ell^\bullet(\underline{L}) - 1$, where $\underline{L}$ ranges over the invertible Modules on $X = G/B$ associated with the representations $B \to \underline{G}_m$ (which in fact give all the invertible modules on X [5, Exp. 15]). On the other hand, an analogous argument shows that

$$A(G/B) \longrightarrow A(G)$$

is also surjective and that its kernel is the ideal generated by the elements of $A^1(G/B)$ defined by the representations $B \to \underline{G}_m$, thus in fact by all of $A^1(G/B)$.(*) This shows that $A(G)$ reduces

(*) Cf. IX 3.4, 2.2.

to $A^0(X) = \mathbb{Z}$ if and [print 680]only if "G is torsion-free", i.e. G is isomorphic to a product of groups $\mathrm{S}\ell(n)$ and $\mathrm{Sp}(m)$. Thus in this case, and in this case alone, the homomorphism

$$\phi : A(G) \longrightarrow \mathrm{Gr}^\bullet_{\mathrm{top}}(G)$$

is an isomorphism. Thus, when G has torsion, the underlying scheme provides an example of a nonbijective ϕ.

4.8. Similarly, one finds an example in which the homomorphism γ of (4.1)

$$\gamma : A(G) \longrightarrow H^{2*}(G, \underline{Z}_\ell(*)) \tag{4.9}$$

does not factor through $\mathrm{Gr}^\bullet_{\mathrm{top}}(G)$, or, equivalently here, an example in which this homomorphism is nonzero in degrees > 0. Restricting ourselves (as suitable specialization theorems allow) to the case $k = \mathbb{C}$, we are reduced to studying the homomorphism

$$A(G) \longrightarrow H^*(G(\mathbb{C}), \mathbb{Z}), \tag{4.10}$$

which is moreover deduced by passage to the quotient from the canonical homomorphism

$$A(G/B) \simeq H^*(G/B(\mathbb{C}), \mathbb{Z}) \longrightarrow H^*(G(\mathbb{C}), \mathbb{Z}).$$

It is tempting to conjecture that the homomorphism (4.10) is in fact injective, since the torsion primes of $G(\mathbb{C})$ are known to be exactly those dividing the order of the finite group $A^+(G) = \coprod_{i>0} A^i(G)$[672]. Were this so, as soon as G had torsion the homomorphism (4.10) would not factor through ϕ, and thus, for suitable ℓ (more precisely, for ℓ dividing the order of $A^+(G)$, i.e. ℓ a torsion coefficient of G), the same would hold for (4.9). One can at least assert that, for the group $G = \underline{G}_2$ over the field $\mathbb{C}$, the homomorphism (4.10) is nonzero in degree > 0; more precisely, there is an element of $A^3(G)$ whose image in $H^6(G(\mathbb{C}), \mathbb{Z})$ has order 2; consequently, the homomorphism (4.9) does not factor through ϕ when $\ell = 2$. The fact used concerning $\underline{G}_2$, which amounts to the existence of an element of $H^6(G/B(\mathbb{C}), \mathbb{Z})$ whose image in $H^6(G(\mathbb{C}), \mathbb{Z})$ is nonzero, is found in [3], using the familiar dictionary between the languages of complex semisimple algebraic groups and compact Lie groups; it was kindly communicated to me by J.P. SERRE.

[print 681]

5. Relations between $\mathrm{Gr}^\bullet(X)$ and $H^{2*}(X, \underline{Z}_\ell(*))$

5.1. The example given in 4.8 does not rule out the possibility of finding, for an arbitrary scheme X, a ring homomorphism

$$g^\bullet : \mathrm{Gr}^\bullet(X) \longrightarrow H^{2*}(X, \underline{Z}_\ell(*)),$$

such that, for every locally free Module $\underline{E}$ on X, g^i takes the Chern class $c_i(\underline{E})$ of Exp. V to the cohomological Chern class of SGA 5 VII. Of course, ℓ here denotes a prime distinct from the residue characteristics of X. Since the ring $\mathrm{Gr}^\bullet(X)$ is generated by the $c^i(\underline{E})$, one sees in any case that $g^\bullet$ is uniquely determined by the requirement of compatibility with Chern classes.

5.2. Using the Chern homomorphism

$$K^\bullet(X) \longrightarrow \mathrm{Ch}(H^{2*}(X, \mathbb{Z}_\ell(*))),$$

and passing to the associated graded groups for the filtration of the first member and the degree filtration of the second, respectively, one obtains a homomorphism of graded groups (but not of rings!)

$$\psi : \mathrm{Gr}^\bullet(X) \longrightarrow H^{2*}(X, \mathbb{Z}_\ell(*)),$$

satisfying the relation

$$\psi(c_i(\underline{E})) = (-1)^{i-1}(i-1)!\, c_i^{\ell}(\underline{E}),$$

where $c_i^{\ell}(\underline{E})$ denotes the ℓ-adic Chern class. By division starting from the ψ^i, one obtains a ring homomorphism:

$$g_{\mathbb{Q}} : \mathrm{Gr}^{\bullet}(X) \longrightarrow H^{2*}(X, \mathbb{Q}_\ell(*)) = H^{2*}(X, \mathbb{Z}_\ell(*)) \otimes_{Z_\ell} \mathbb{Q}_\ell,$$

taking Chern classes to Chern classes, such that the image of the first member is contained in that of $H^{2*}(X, \mathbb{Z}_\ell(*))$[673]; consequently, the question of the existence of $g^\bullet$ has an affirmative answer upon replacing $H^{2*}(X, \mathbb{Z}_\ell(*))$ by its quotient by its torsion subgroup. The first doubtful case in defining the components g^i of the hypothetical homomorphism g is that of g^3; a priori, one succeeds only in defining $2g^3 = \psi^3$. The answer to the existence question for g^3 would be negative if, for example, one could find three invertible sheaves $\underline{L}_1, \underline{L}_2, \underline{L}_3$ on X satisfying the relation

$$(\underline{L}_1 - 1)(\underline{L}_2 - 1)(\underline{L}_3 - 1) = 0 \quad \text{in } K^{\bullet}(X),$$

[print 682]but such that

$$c_1(\underline{L}_1)c_1(\underline{L}_2)c_1(\underline{L}_3) \neq 0 \quad \text{in } H^6(X, \mu_2^{\otimes 3}).$$

5.3. When X is smooth and quasi-projective over a field k, one can likewise ask about the existence of a ring homomorphism

$$g : \mathrm{Gr}^{\bullet}(X) \longrightarrow A(X)$$

with values in the Chow ring $A(X)$, taking Chern classes to Chern classes. One can repeat the preceding considerations, which show in particular how to construct $(-1)^{i-1}(i-1)!g^i$ directly in the absence of g^i itself.[674]

5.4. To conclude this section, observe that essentially the same question arises when, in place of a scheme X, one considers a compact topological space X, say; one can again consider the associated graded $\mathrm{Gr}^{\bullet}(X)$ of $K^{\bullet}(X)$ endowed with its λ-filtration, and seek a ring homomorphism

$$\mathrm{Gr}^{\bullet}(X) \longrightarrow H^{2*}(X, \mathbb{Z})$$

taking Chern classes to Chern classes.

6. Cohomological Riemann–Roch theorem and Gysin homomorphism

6.1. For simplicity, we restrict ourselves here to schemes X admitting an ample invertible Module; this implies that the naive $K^{\bullet}(X)$ and the other one can be identified. Let us begin with the Riemann–Roch formula "with denominators" for a proper complete-intersection morphism $f : X \to Y$. Since (for a prime ℓ prime to the residue characteristics of Y) one can define the ℓ-adic Chern classes of a locally free Module $\underline{E}$,

$$c_i(\underline{E}) \in H^{2i}(X, \mathbb{Z}_\ell(i))$$

(SGA 5 VII), corresponding to a homomorphism

$$K^{\bullet}(X) \longrightarrow \mathrm{Ch}(H^{2*}(X, \mathbb{Z}_\ell(*))), \tag{6.1}$$

the question arises of giving a cohomological variant of the Riemann–Roch formula. The only ingredient missing to give the formula a meaning is a Gysin homomorphism [print 683]

$$H^n(X, \underline{Q}_\ell(i))^{[675]} \longrightarrow H^{n-2d}(Y, \underline{Q}_\ell(i-d))^{[676]}, \tag{6.2}$$

where d is the virtual relative dimension of f[677]. Assuming this homomorphism defined, and taking account of the Riemann–Roch formula in the form of Exp. VIII with values in $\mathrm{Gr}^\bullet(Y)_\mathbb{Q}$, one finds that the validity of the ℓ-adic Riemann–Roch formula for the morphism f is exactly equivalent to the commutativity of the square

$$\begin{array}{ccc} \mathrm{Gr}^\bullet(X)_\mathbb{Q} & \longrightarrow & H^{2*}(X, \underline{Q}_\ell(*)) \\ \downarrow & & \downarrow \\ \mathrm{Gr}^\bullet(Y)_\mathbb{Q} & \longrightarrow & H^{2*}(Y, \mathbb{Q}_\ell(*)), \end{array} \tag{6.3}$$

where the horizontal arrows are defined from (6.1) by passing to associated graded groups and dividing so as to take Chern classes to Chern classes, and the vertical arrows are respectively the arrow of Exp. VIII and the Gysin homomorphism in cohomology (assumed given).

6.2. Observe that the Gysin homomorphism is in any case defined when Y is smooth over a field k (SGA 5 IV), using the relative cohomological purity theorem over k; the same construction applies when Y is excellent, regular and of characteristic zero, by the absolute cohomological purity theorem of SGA 4 XIX, and it will apply automatically to every excellent regular Y, without restriction on the characteristic, once the absolute cohomological purity theorem has been proved in full generality. Finally, the same construction applies, by the relative cohomological purity theorem (SGA 4 XVI 3.7), when X and Y are both smooth over the same scheme S and f is an S-morphism. One should conjecture that, in each of these cases in which the Gysin homomorphism is defined, diagram (6.3) is indeed commutative. The only case in which this assertion is currently verified is that in which X and Y are both smooth over a field k; it then follows from the commutativity of the diagram analogous to (6.3):

$$\begin{array}{ccc} A(X) & \longrightarrow & H^{2*}(X, \mathbb{Z}_\ell(*)) \\ \downarrow & & \downarrow \\ A(Y) & \longrightarrow & H^{2*}(Y, \mathbb{Z}_\ell(*)), \end{array} \tag{6.4}$$

[print 684]involving the rings of cycle classes (mod. rational equivalence) $A(X)$ and $A(Y)$, whose commutativity is established in SGA 5 IV.

6.3. Observe that, in each of the cases considered above in which one can define a Gysin homomorphism, one in fact obtains a construction for cohomology with coefficients in $\mathbb{Z}_\ell(*)$, and not only in $\mathbb{Q}_\ell(*)$. When f is an <u>immersion</u>, this allows the question of the validity of the Riemann–Roch formula "without denominators" to be posed, as explained in no. 3. When X and Y are both smooth over a field k, the commutativity of (6.4) shows that it would suffice to prove the formula without denominators in the Chow ring $A(Y)$ cf. (3.3).

6.4. Without a regularity hypothesis on Y, even when X and Y are of finite type over the field $\mathbb{C}$, one no longer sees a plausible definition of a cohomological Gysin homomorphism (6.2), even for a regular immersion of codimension 1. Can one nevertheless find one with reasonable properties, in particular making square (6.3) commute? The question is clearly connected with that of defining, for a complete intersection X of codimension i in Y, "its cohomology class"

$$\gamma(X) \in H^{2i}(Y, \mathbb{Z}_\ell(i)),$$

so as to take intersections (without excess components) to cup products. This latter question has an affirmative answer at least modulo torsion, i.e. in $H^{2*}(Y, \mathbb{Q}_\ell(*))$, by defining

$$\gamma(X) = 1/(-1)^{i-1}(i-1)!\, c_i(\underline{O}_X).$$

But I do not know whether $c_i(\underline{O}_X)$ is divisible by $(i-1)!$ in $H^{2i}(Y, \mathbb{Z}_\ell(i))$. In particular, for a projective scheme Y of dimension 3 over the field of complex numbers, say, with X reduced to a (singular) point of Y, is $c_3(\underline{O}_X)$ divisible by 2 in $H^6(Y, \mathbb{Z}_2(3))$, and in particular does the Chern class vanish in $H^6(Y, \mathbb{Z}/2\mathbb{Z})$? In the same direction, let us also mention the following question: let Y be a projective (singular) scheme of dimension 4, and X a closed subscheme of dimension 1 that is globally an intersection of three positive divisors D_1, D_2, D_3; consider the product of their Chern classes mod. 2:

$$c_1(D_1)c_1(D_2)c_1(D_3) \in H^6(Y, \mathbb{Z}/2\mathbb{Z});$$

[print 685]is this element of H^6 independent of the particular choice of the D_i with intersection X?

7. Chern classes of perfect complexes

7.1. In seeking to develop a cohomological Riemann–Roch formula, with or without denominators, beyond the rather artificial hypothesis that the schemes under consideration admit an ample invertible Module, one is necessarily confronted with the question of defining, for a perfect complex of Modules $L_\bullet$ on a scheme X, Chern classes

$$c_i(L_\bullet) \in H^{2i}(X, \mathbb{Z}_\ell(i)),$$

generalizing the Chern classes of locally free Modules in SGA 5 VII. The same question also arises for general locally ringed topoi (compare [10]) and for perfect complexes on paracompact spaces ringed by continuous complex functions, working then with ordinary integral cohomology. For a compact space, the $K^\bullet(X)$ constructed with perfect complexes is known to be the same as the "naive" one constructed with locally free Modules (Exp. IV), and in this case one is trivially reduced to the classical definition. But even for a locally compact space the problem posed seems nontrivial, since the integral cohomology of such a space is not always the inverse limit of the cohomologies of its compact subsets.

7.2. More precisely, for a closed subset Y of X such that $L_\bullet|(X-Y)$ is zero (in the sense of derived categories), i.e. such that the cohomology sheaves of $L_\bullet$ have supports in Y, we would like to define Chern classes <u>with support in</u> Y

$$c_i^Y(L_\bullet) \in H_Y^{2i}(X, \mathbb{Z}_\ell(i)) \quad \text{resp.} \quad H_Y^{2i}(X, \mathbb{Z}),$$

which would recover the "ordinary" (sic) Chern classes of 7.1 by means of the canonical homomorphism

$$H_Y^*(X, -) \longrightarrow H^*(X, -).$$

[print 686]Even when these reduce to the Chern classes of locally free Modules (X a scheme with an ample invertible Module, or a compact space), these Chern classes "with supports" have not been constructed, and their construction will doubtless have to proceed together with that of the general Chern classes postulated in 7.1(*)

It is probable that Chern classes with supports will have a role to play in proving the cohomological Riemann–Roch formula without denominators (cf. 3.5). On the other hand, they should allow an improved version of the Riemann–Roch formula (with or without denominators) by introducing families of supports into that formula.

(*) Added in October 1968. The problems raised here have been solved in the setting of <u>Hodge cohomology</u> by L. ILLUSIE; cf. [13]

8. Chow ring of regular schemes

The problem is to define rational equivalence of cycles and to study the resulting ring of classes (in particular its functorial properties) for a regular noetherian scheme X admitting an ample invertible Module, generalizing the theory known when X is smooth over a field [4]. The key technical result will again be a "moving lemma", ensuring that a cycle can be varied within its class so as to intersect a given cycle in the correct dimension.

The problem considered is quite natural a priori, particularly since we have seen (4.5 to 4.8) that the substitutes $\mathrm{Gr}^\bullet(X)$, $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$ for the Chow ring $A(X)$ are hardly satisfactory beyond a theory "modulo torsion". When X is obtained from a smooth projective algebraic scheme Y over a field k by taking the local scheme at the vertex of a projecting cone C and removing the origin, one finds (admitting that $A(X)$ is indeed the direct limit of the Chow rings of the $U-(O)$, where U ranges over the open neighbourhoods in C of the vertex O) that $A(X)$ is canonically isomorphic to the quotient ring $A(Y)/\xi A(Y)$, where $\xi \in A^1(Y)$ is induced by the projective immersion of Y under consideration. This shows that knowledge of the Chow ring $A(X)$ (of a nonclassical type) is equivalent to knowledge of $A(Y)/\xi A(Y)$, which might be called the "primitive part" of the Chow ring $A(Y)$ (in a sense suggested by the Lefschetz–Hodge cohomological theory of nonsingular projective algebraic varieties). This example shows that the "local Chow rings" of local rings with a singularity [print 687]that is isolated are geometric invariants that are by no means trivial, and will undoubtedly have their part to play in the "arithmetic" study of these local rings.

Of course, once one has a reasonable notion of a Chow ring for a regular noetherian scheme admitting an ample invertible Module, one should generalize to this case the relations established or conjectured, respectively, in n° 4 between this ring and the ring $\mathrm{Gr}^\bullet_{\mathrm{top}}(X)$, and formulate in terms of this ring the question of the "Riemann–Roch formula without denominators" stated in n° 3.

BIBLIOGRAPHY

[1] M. F. Atiyah and F. Hirzebruch, The Riemann–Roch theorem for analytic embeddings, Topology, vol. 1, p. 151-166 (1962).

[2] A. Borel and J.P. Serre, Le théorème de Riemann–Roch, Bull. Soc. Math. France, vol. 86, p. 97-136 (1958).

[3] R. Bott and H. Samelson, Amer. Journ. of Math. vol 80, p. 1004 (1958).

[4] C. Chevalley, Les classes d'équivalence rationnelle[678], I and II, Seminar at the ENS 1958, exposés 2 and 3.

[5] C. Chevalley, Classification des groupes de Lie algébriques, Seminar at the ENS 1956/58.

[6] A. Dold and D. Puppe, Homologie nicht-additiver Funktoren. Anwendungen, Annales de l'Inst. Fourier t. XI, p. 201-312 (1961).

[7] H. Grauert, Ein Theorem der analytischen Garbentheorie, Pub. Math. n° 5, p. 5-64 (1960).

[8] A. Grothendieck, Classes de faisceaux et théorème de Riemann–Roch, mimeographed notes prepared by the Institute for Advanced Study, 1957 (work cited as [RRR] in the present exposé, and reproduced in our seminar as an Appendix to Chap. O).

[9] A. Grothendieck, La théorie des classes de Chern, Bull. Soc. Math. France vol. 86, p. 137-154 (1958). [print 688]

[10] A. Grothendieck, Classes de Chern et représentations linéaires des groupes discrets, in: Dix exposés sur la cohomologie des schémas, Masson–North Holland Pub. Cie

[11] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero, Ann. of Math. 79, p. 109-326 (1964).

[12] L. Hodgkin, thesis (to be published?).

[13] L. Illusie, work in preparation.

[14] J.P. Serre, Algèbre Locale[679], Multiplicités, Lecture Notes in Math. n° 11 (1965) Springer.

[15] A. Borel, Sous-groupes commutatifs et torsion des groupes de Lie compacts connexes, Tohoku Math. Journal, vol. 13, p. 216-240 (1961).

[16](*) P. Deligne, Exp. XVIII in Théorie des topos et cohomologie étale des schémas (SGA 4)[680], by M. Artin, A. Grothendieck and J. L. Verdier.

[17](*) U. Manin[681], Correspondances, motifs et transformations monoïdales (in Russian). Mat.[682] Sbornik[683], t 77 (119) n° 4, 1968.

[print 689]

Note for page 1.

The theory of the operations Λ^i in $D^-(X)$, and of the operations λ^i in $K^\bullet(X)$, was written down by P. Deligne in 1968. He proves that $K^\bullet(X)$ thereby becomes a pre-λ-ring, and, when $\underline{O}_X$ is a sheaf of Algebras over a field, that $K^\bullet(X)$ is even a λ-ring. It is plausible that a refinement of Deligne's method would make it possible to dispense with the restrictive hypothesis on $\underline{O}_X$.

Notes for page 11

(*) (April 1969) D. MUMFORD points out to us a proof of (4.6)[684] in the Chow ring (without disregarding torsion), which he obtained in a seminar at Harvard in 1959! The reader will find the proof of this beautiful result in SGA 5 VII 9. This result obviously contains the corresponding cohomological result mentioned in note (**). However, the cohomological result is proved by J. P. JOUANOLOU under more general conditions (without a quasi-projectivity hypothesis[685], and in the relative smooth case over an arbitrary base).

(**) (January 1969) J. P. JOUANOLOU has just proved these formulas for every y without disregarding torsion, cf. SGA 5 VII § 4.9. DELIGNE and MANIN 17 had previously observed independently, and by (easy) arguments of a rather different nature, that it was enough to establish formulas for "algebraic" y, and even for $y = 1$ in 16, to conclude the same relation in the general case.

(*) Added in February 1969

Terminological index

	Exp.	§
additive (map)	I	6.1
additive (map)	IV	1.5
augmented $K-\lambda$-algebra	V	3.9
algebraically equivalent to zero (in $K^\bullet$)	X App.	7.11
algebraically equivalent to zero (invertible sheaf)	XIII	4.4
of $\underline{C}_0$-perfect amplitude contained in $[a,b]$ (complex)[686]	I	4.7
of finite $\underline{C}_0$-perfect amplitude (complex)	I	4.7
of $\underline{C}_0$ perfect amplitude contained in $[a,b]$ (complex)[687]	I	4.8
of finite perfect amplitude (complex)	I	4.8
of flat amplitude contained in $[a,b]$ (complex)	I	5.2
of finite flat amplitude (complex)	I	5.2
of perfect amplitude relative to f (or Y) contained in $[a,b]$ (complex)	III	4.1
of perfect amplitude contained in $[-n,0]$ (morphism)	III	4.1
of finite perfect amplitude relative to f (complex)	III	4.1
of flat amplitude relative to f (or Y) contained in $[a,b]$ (complex)	III	3.1
of flat amplitude contained in $[-n,0]$ (morphism)	III	3.1
of finite flat amplitude relative to f (complex)	III	3.1
projective amplitude	I	4.12.1
λ-ring	V	2.4
λ-ring generated by generators and relations	V	4.6
free λ-ring generated by a family of generators	V	4.4
Chern ring	V	6.2
polynomial map	X	2.1.1
binomial (ring)	V	2.7
[print 692]Chern character	V	6.4
Todd character	V	1.16
abelian $\underline{S}$-category	I	1.1.1
flat abelian $\underline{S}$-category	I	1.1.1
additive $\underline{S}$-category	I	1.1.1
triangulated $\underline{S}$-category	I	1.1.1
Chern class	V	6.7
completed Chern class	V	6.7
total Chern class	V	6.7
codimension (of a regular immersion)	VII	1.6
coherent (complex)	I	3.1
coherer	II	3.2
relative cotangent complex	VIII	2.3
elliptic complex	II App. II	1.2.5
cup-product	I	8.2
special positive k-cycle	XIII	6.9
degree (of a polynomial map)	X	2.1.1
degree (of an invertible sheaf)	XIII	2.2
degree (of a morphism of schemes)	X	4.4
dense (subset of an abelian or triangulated category)	IV	1.7
perfect dimension	I	4.14.1
virtual relative dimension	VIII	1.9
divisorial	II	2.2.5
dual	I	7.3
perfect duality	I	7.8

numerical equivalence (in $\mathrm{Gr}^\bullet(X)$ and $\mathrm{Gr}_\cdot(X)$)	X	4.5
τ-equivalence	X	2.4.1
"	XIII	4.4
complex pseudo-analytic space	II	2.4.7
complex Stein pseudo-analytic space	II	2.4.7
exact (subset of an abelian or triangulated category)	IV	1.7
(b)-sheaf	XIII	1.5
conormal sheaf	VII	1.6
relative cotangent sheaf	II App. II	1.2.1
[print 693]relative tangent sheaf	II App. II	1.2.1
(b)-family	XIII	1.12
ample family	II	2.2.4
family of classes of coherent sheaves	XIII	1.12
bounded family of classes of coherent sheaves	XIII	1.12
family of classes of coherent sheaves bounded by a coherent sheaf	XIII	1.12
relative cotangent bundle	II App. II	1.2.1
relative tangent bundle	II App. II	1.2.1
faithfully flat (morphism of ringed topoi)	I	2.16.2
filtration of an augmented λ-algebra	V	3.10
additive $\underline{S}$-functor	I	1.1.3
exact $\underline{S}$-functor	I	1.1.3
projection formula	IV	2.11
Grothendieck group	IV	2.2
λ-homomorphism	V	2.1
specialization homomorphism	X App.	7.3
" "	X App.	7.16
λ-ideal	V	4.5
analytic index	II App. II	3.2.2
complete intersection (morphism)	VIII	1.1
n-isomorphism	I	1.4.3
homogeneous free (graded module)	VI	1.2.2
locally stable under kernels of epimorphisms	I	1.2
locally quasi-liftable	I	1.2
locally quasi-stable under kernels of epimorphisms	I	1.2
locally liftable	I	1.2
Chern number	X	4.2
intersection number	X	4.1
intersection number (for a family of invertible sheaves)	X	4.4
numerically equivalent to zero (invertible sheaf)	XIII	4.4
universal differential operator of order n	II App. II	1.2.1
[print 694]Adams operations	V	7.1
$\underline{C}_0$-perfect (complex)	I	4.7
perfect (complex)	I	4.8
perfect (morphism)	III	4.1
perfect relative to f (complex)	III	4.1
precoherent	I	3.1
of finite n-$\underline{C}_0$-presentation	I	2.8
of finite presentation	I	2.8
of finite ∞-$\underline{C}_0$-presentation	I	2.8
n-$\underline{C}_0$-pseudo-coherent (complex)	I	2.2
$\underline{C}_0$-pseudo-coherent (complex)	I	2.2
n-pseudo-coherent (complex)	I	2.15
n-pseudo-coherent (morphism)	III	1.2
pseudo-coherent (complex)	I	2.15
pseudo-coherent (morphism)	III	1.2
n-pseudo-coherent relative to f (complex)	III	1.2
pseudo-coherent relative to f (complex)	III	1.2
(b)-polynomial	XIII	1.9

Snapper polynomial	X	2.3.1
universal polynomial	V	1.15
pre-λ-ring	V	2.1
n-quasi-isomorphism	I	1.4.3
rank (of a perfect complex)	I	6.11
m-regular (sheaf)	XIII	1.1
regular (homomorphism)	VII	1.1
regular (ideal)	VII	1.4
regular (immersion)	VII	1.4
triangulated sub-$\underline{S}$-category	I	1.1.4
special (λ-ring)	V	2.4
special (morphism)	III	1.1.4
Stein (locally ringed topos)	II	2.4.1
strictly $\underline{C}_0$-perfect	I	2.1
strictly $\underline{C}_0$-n-pseudo-coherent	I	2.1
strictly $\underline{C}_0$-pseudo-coherent	I	2.1
minimum system of generators	VII	1.4.1
[print 695]of finite Tor-dimension	I	5.2
Tor-independence	III	1.5
trace	I	8.1
of finite $\underline{C}_0$-type	I	1.2
mixed variety of class C^r	II App. II	1.1

Index of notations

	Exposé	§
A_N	V	5.5
ch	V	6.3
$\mathrm{Ch}(A),\ \mathrm{Ch}_K(A)$	V	6.2
$(\mathrm{Ch}(A))_n$	V	6.5
cl_C	IV	1.1
$\mathrm{Coh}(S)$	II	2.2.1
$\underline{C}_0\text{-}\,\mathrm{parf}\,\text{-}\,\mathrm{amp}(F)$	I	4.7
$C^p_{\mathrm{num}}(X)$[688]	XIII	5.2
$C^r(V),\ C^\infty(V)$	II App. II	1.1
$c(x),\ c^i(x),\ \widetilde{c}(x)$	V	6.7
$\mathrm{D}({}_AS),\ \mathrm{D}({}_A\underline{S})$	I	2.15
$\mathrm{D}({}_A\underline{S})_{\mathrm{coh}}$,[689] $\mathrm{D}({}_A\underline{S})_{n\text{-}\,\mathrm{coh}}$	I	2.15
$\mathrm{D}({}_A\underline{S})_{\mathrm{parf}}$	I	4.9
$\mathrm{D}^-({}_A\underline{S})_{\mathrm{torf}},\ \mathrm{D}^-({}_AX)_{\mathrm{torf}}$	I	5.3
$\mathrm{D}(\underline{C})_{\underline{C}_0\text{-}\,\mathrm{coh}},\ \mathrm{D}(\underline{C})_{\underline{C}_0\text{-}n\text{-}\,\mathrm{coh}}$	I	2.6
$\mathrm{D}(\underline{C})_{\underline{C}_0\text{-}\,\mathrm{parf}}$	I	4.9
$\mathrm{D}(f)_{\mathrm{coh}}$	III	1.2
$\mathrm{D}(f)_{n\text{-}\,\mathrm{coh}}$	III	1.2
$\mathrm{D}(f)_{\mathrm{parf}},\ \mathrm{D}(f)_{\underline{\mathrm{parf}}}$	III	4.2
$\mathrm{D}(f)_{\mathrm{torf}}$	III	3.2
$\mathrm{Diff}^n(E,F)$	II App. II	1.2.2
$d(L)$	XIII	2.2
$d^n_{X/S}$	II App. II	1.2.1
$d^n_{X/S}(E)$	II App. II	1.2.2
[print 697]$\mathrm{D}^-(\mathrm{Qcoh}(S))_{\mathrm{coh}},\ \mathrm{D}(\mathrm{Qcoh}(S))_{\mathrm{parf}}$	II	2.2.1
$\mathrm{D}(S),\ \mathrm{D}(\underline{S}),\ \mathrm{D}(\underline{S},A)$	I	2.15
$\mathrm{D}(X)_{f\text{-}\,\mathrm{parf}},\ \mathrm{D}(X)_{f\text{-}\underline{\mathrm{parf}}}$	III	4.2
$\mathrm{D}(X)_{f\text{-}\,\mathrm{torf}}$	III	3.2
$\mathrm{D}(X)_{\mathrm{qcoh}}$	III	1.7
$\mathrm{D}(X)_{Y\text{-}\,\mathrm{coh}},\ \mathrm{D}(X)_{Y\text{-}n\text{-}\,\mathrm{coh}}$	III	1.2
$\mathrm{D}(X)_{Y\text{-}\,\mathrm{parf}},\ \mathrm{D}(X)_{Y\text{-}\underline{\mathrm{parf}}}$	III	4.2
$\mathrm{D}(X)_{Y\text{-}\,\mathrm{torf}}$	III	3.2
$\mathrm{D}(Y)_{n\text{-}\,\mathrm{coh}}$	III	1.1
$E^\vee$	I	7.3
$*$	V	6.1
$\lvert\underline{F}\rvert$	X	4.3.1
f^* (between the $K^\bullet$)	IV	2.7
f^* (between the $K_\cdot$)	IV	2.12
f_* (between the $K_\cdot$)	IV	2.11
f_* (between the $K^\bullet$)	IV	2.12
f_* (projective bundle case)	VI	5.2
f_{gr}	VIII	3.2
$\mathrm{Filt}_n(X)$	X	1.1.1
$F'_{>n},\ F'_{\leq n}$	I	2.12
φ^a	V	2.9
$\widehat{G},\ \widehat{G}_K$	V	1.1
$\widehat{G}_\circ,\ \widehat{G}_\nabla$	V	3.7
$\widehat{G}_a$	V	1.3.3
$\widehat{G}_g$	V	1.3.2

	Exposé	§
$\mathrm{Gr}^\bullet(X),\ \mathrm{Gr}_\cdot(X)$	X	3.1
$\mathrm{Gr}^\bullet_{\mathrm{alg}},\ \mathrm{Gr}_\cdot^{\mathrm{alg}}$	X App.	7.11
[print 698]$\mathrm{Gr}^\bullet_{\mathrm{num}},\ \mathrm{Gr}_\cdot^{\mathrm{num}}$	X App.	7.9.4
$\gamma^n,\ \gamma_t$	V	3.2
$\gamma^p(N,x)$	V	5.5
$\widehat{G}^t,\ \widehat{G}^u$	V	2.3
$\underline{\mathrm{Hom}}_{\underline{M}}(E,F)$	I	4.5
$k(A)$	IV	1.5
$K^\bullet_{\mathrm{alg}},\ K_\cdot^{\mathrm{alg}}$	X App.	7.11
$K^\bullet({}_AX),\ K_\cdot({}_AX)$	IV	2.2
$K^\bullet({}_AX)_{\mathrm{naif}},\ K_\cdot({}_AX)_{\mathrm{naif}}$	IV	2.2
$k(C)$	I	6.3
$K^\bullet(f),\ K_\cdot(f)$	IV	3.3.1
$K^\bullet_{\mathrm{num}},\ K_\cdot^{\mathrm{num}}$	X App.	7.9.4
$K^\bullet(X),\ K_\cdot(X)$	IV	2.2
$K^\bullet(X)_{\mathrm{naif}},\ K_\cdot(X)_{\mathrm{naif}}$	IV	2.2
$K^\bullet(X/Y),\ K_\cdot(X/Y)$	IV	3.3.1
$\mathrm{Lib}(U)$	II	2.0
$\mathrm{Loc\,lib}(U)$	II	2.0
$\underline{L}_\cdot^{X/Y}$	VIII	2.1
λ^i	V	2.1
$\lambda_{-1}(N)$	V	5.1
$\lambda_{-1}(N)$	VII	2.7
$\lambda^p(N,x)$	V	5.3
λ_t	V	2.1
$\mathrm{Mod}(U)$	II	2.0
[print 699]$\mathrm{par.\,amp}_f(E),\ \mathrm{par.\,amp}_Y(E)$	III	4.1
$\mathrm{Parf}({}_A\underline{S}),\ \mathrm{Parf}(\underline{C},\underline{C}_0),\ \mathrm{Parf}(\underline{S})$	I	4.9
$\mathrm{Pic}^N(X)$	X	4.5.2
$\mathrm{Pic}^\circ(X)$	X	2.4.1
$\underline{\mathrm{Pic}}^\circ_{X/k}$	X	2.4
$\mathrm{Pic}^\tau(X)$	X	2.4.1
$\mathrm{Pic}(X)$	X	2.4
$\underline{\mathrm{Pic}}_{X/k}$	X	2.4
$P^n_{X/S}$	II App. II	1.2.1
$P^n_{X/S}(E)$	II App. II	1.2.2
$\underline{\mathrm{Proj}}_0$	VI	2.3.2
$\overline{\psi^k(x)}$[690]	V	7.1
$\mathrm{Qcoh}(S)$	II	2.2.1
${}^r\underline{O}_{U\times V}$	II App. II	1.1
ρ_t	V	3.4[691]
$\circ$	V	2.3
T_f	VIII	2.5
Todd	V	1.16
$\mathrm{tor.\,amp}(F)$	I	5.2
$\mathrm{tor.\,amp}_f(E),\ \mathrm{tor.\,amp}_Y(E)$	III	3.1
$\mathrm{Tr}_{E\|U}$	I	8.1
$T_{X/S}$	II App. II	1.2.1
τ	I	8.1
∇	V	3.5
[print 700]$1+A[[t]]^+$	V	1.1
$1+\widehat{B}^+$	V	1.11
$X(n)$	X	1.1.2
$x\cdot_N y$	V	5.5
$\langle x,y\rangle$	X	4.1

	Exposé	§
$Z(x)$	X	1.1.2

[print rear-717]

Lecture Notes in Mathematics

Comprehensive leaflet on request

Vol. 38: R. Berger, R. Kiehl, E. Kunz und H.-J. Nastold, Differentialrechnung in der analytischen Geometrie IV, 134 Seiten. 1967 DM 12,–

Vol. 39: Séminaire de Probabilités I. II, 189 pages. 1967. DM 14,–

Vol. 40: J. Tits, Tabellen zu den einfachen Lie Gruppen und ihren Darstellungen. VI, 53 Seiten. 1967. DM 6.80

Vol. 41: A. Grothendieck, Local Cohomology. VI, 106 pages. 1967. DM 10,–

Vol. 42: J. F. Berglund and K. H. Hofmann, Compact Semitopological Semigroups and Weakly Almost Periodic Functions. VI, 160 pages. 1967. DM 12,–

Vol. 43: D. G. Quillen, Homotopical Algebra. VI, 157 pages. 1967. DM 14,–

Vol. 44: K. Urbanik, Lectures on Prediction Theory. IV, 50 pages. 1967. DM 5,80

Vol. 45: A. Wilansky, Topics in Functional Analysis. VI, 102 pages. 1967. DM 9,60

Vol. 46: P. E. Conner, Seminar on Periodic Maps. IV, 116 pages. 1967. DM 10,60

Vol. 47: Reports of the Midwest Category Seminar I. IV, 181 pages. 1967. DM 14,80

Vol. 48: G. de Rham, S. Maumary et M. A. Kervaire, Torsion et Type Simple d'Homotopie. IV, 101 pages. 1967. DM 9,60

Vol. 49: C. Faith, Lectures on Injective Modules and Quotient Rings. XVI, 140 pages. 1967. DM 12,80

Vol. 50: L. Zalcman, Analytic Capacity and Rational Approximation. VI, 155 pages. 1968. DM 13.20

Vol. 51: Séminaire de Probabilités II. IV, 199 pages. 1968. DM 14,–

Vol. 52: D. J. Simms, Lie Groups and Quantum Mechanics. IV, 90 pages. 1968. DM 8,–

Vol. 53: J. Cerf, Sur les difféomorphismes de la sphère de dimension trois ($\Gamma_4 = 0$). XII, 133 pages. 1968. DM 12,–

Vol. 54: G. Shimura, Automorphic Functions and Number Theory. VI, 69 pages. 1968. DM 8,–

Vol. 55: D. Gromoll, W. Klingenberg und W. Meyer, Riemannsche Geometrie im Großen. VI, 287 Seiten. 1968. DM 20,–

Vol. 56: K. Floret und J. Wloka, Einführung in die Theorie der lokalkonvexen Räume. VIII, 194 Seiten. 1968. DM 16,–

Vol. 57: F. Hirzebruch und K. H. Mayer, O (n)-Mannigfaltigkeiten, exotische Sphären und Singularitäten. IV, 132 Seiten. 1968. DM 10,80

Vol. 58: Kuramochi Boundaries of Riemann Surfaces. IV, 102 pages. 1968. DM 9,60

Vol. 59: K. Jänich, Differenzierbare G-Mannigfaltigkeiten. VI, 89 Seiten. 1968. DM 8,–

Vol. 60: Seminar on Differential Equations and Dynamical Systems. Edited by G. S. Jones. VI, 106 pages. 1968. DM 9,60

Vol. 61: Reports of the Midwest Category Seminar II. IV, 91 pages. 1968. DM 9,60

Vol. 62: Harish-Chandra, Automorphic Forms on Semisimple Lie Groups X, 138 pages. 1968. DM 14,–

Vol. 63: F. Albrecht, Topics in Control Theory. IV, 65 pages. 1968. DM 6,80

Vol. 64: H. Berens, Interpolationsmethoden zur Behandlung von Approximationsprozessen auf Banachräumen. VI, 90 Seiten. 1968. DM 8,–

Vol. 65: D. Kölzow, Differentiation von Maßen. XII, 102 Seiten. 1968. DM 8,–

Vol. 66: D. Ferus, Totale Absolutkrümmung in Differentialgeometrie und -topologie. VI, 85 Seiten. 1968. DM 8,–

Vol. 67: F. Kamber and P. Tondeur, Flat Manifolds. IV, 53 pages. 1968. DM 5,80

Vol. 68: N. Boboc et P. Mustată, Espaces harmoniques associés aux opérateurs différentiels linéaires du second ordre de type elliptique. VI, 95 pages. 1968. DM 8,60

Vol. 69: Seminar über Potentialtheorie. Herausgegeben von H. Bauer. VI, 180 Seiten. 1968. DM 14,80

Vol. 70: Proceedings of the Summer School in Logic. Edited by M. H. Löb. IV, 331 pages. 1968. DM 20,–

Vol. 71: Séminaire Pierre Lelong (Analyse), Année 1967 – 1968. VI, 190 pages. 1968. DM 14,

Vol. 72: The Syntax and Semantics of Infinitary Languages. Edited by J. Barwise. IV, 268 pages. 1968. DM 18,–

Vol. 73: P. E. Conner, Lectures on the Action of a Finite Group. IV, 123 pages. 1968. DM 10,–

Vol. 74: A. Fröhlich, Formal Groups. IV, 140 pages. 1968. DM 12,–

Vol. 75: G. Lumer, Algèbres de fonctions et espaces de Hardy. VI, 80 pages. 1968. DM 8,–

Vol. 76: R. G. Swan, Algebraic K-Theory. IV, 262 pages. 1968. DM 18,–

Vol. 77: P.-A. Meyer, Processus de Markov: la frontière de Martin. IV, 123 pages. 1968. DM 10,–

Vol. 78: H. Herrlich, Topologische Reflexionen und Coreflexionen. XVI, 166 Seiten. 1968. DM 12,–

Vol. 79: A. Grothendieck, Catégories Cofibrées Additives et Complexe Cotangent Relatif. IV, 167 pages. 1968. DM 12,–

Vol. 80: Seminar on Triples and Categorical Homology Theory. Edited by B. Eckmann. IV, 398 pages. 1969. DM 20,–

Vol. 81: J.-P. Eckmann et M. Guenin, Méthodes Algébriques en Mécanique Statistique. VI, 131 pages. 1969. DM 12,–

Vol. 82: J. Wloka, Grundräume und verallgemeinerte Funktionen. VIII, 131 Seiten. 1969. DM 12,–

Vol. 83: O. Zariski, An Introduction to the Theory of Algebraic Surfaces. IV, 100 pages. 1969. DM 8,–

Vol. 84: H. Lüneburg, Transitive Erweiterungen endlicher Permutationsgruppen. IV, 119 Seiten. 1969. DM 10.–

Vol. 85: P. Cartier et D. Foata, Problèmes combinatoires de commutation et réarrangements. IV, 88 pages. 1969. DM 8,–

Vol. 86: Category Theory, Homology Theory and their Applications I. Edited by P. Hilton. VI, 216 pages. 1969. DM 16,–

Vol. 87: M. Tierney, Categorical Constructions in Stable Homotopy Theory. IV, 65 pages. 1969. DM 6,–

Vol. 88: Séminaire de Probabilités III. IV, 229 pages. 1969. DM 18,–

Vol. 89: Probability and Information Theory. Edited by M. Behara, K. Krickeberg and J. Wolfowitz. IV, 256 pages. 1969. DM 18,–

Vol. 90: N. P. Bhatia and O. Hajek, Local Semi-Dynamical Systems. II, 157 pages. 1969. DM 14,–

Vol. 91: N. N. Janenko, Die Zwischenschrittmethode zur Lösung mehrdimensionaler Probleme der mathematischen Physik. VIII, 194 Seiten. 1969. DM 16,80

Vol. 92: Category Theory, Homology Theory and their Applications II. Edited by P. Hilton. V, 308 pages. 1969. DM 20,–

Vol. 93: K. R. Parthasarathy, Multipliers on Locally Compact Groups. III, 54 pages. 1969. DM 5,60

Vol. 94: M. Machover and J. Hirschfeld, Lectures on Non-Standard Analysis. VI, 79 pages. 1969. DM 6,–

Vol. 95: A. S. Troelstra, Principles of Intuitionism. II, 111 pages. 1969. DM 10,–

Vol. 96: H.-B. Brinkmann und D. Puppe, Abelsche und exakte Kategorien, Korrespondenzen. V, 141 Seiten. 1969. DM 10,–

Vol. 97: S. O. Chase and M. E. Sweedler, Hopf Algebras and Galois theory. II, 133 pages. 1969. DM 10,–

Vol. 98: M. Heins, Hardy Classes on Riemann Surfaces. III, 106 pages. 1969. DM 10,–

Vol. 99: Category Theory, Homology Theory and their Applications III. Edited by P. Hilton. IV, 489 pages. 1969. DM 24,–

Vol. 100: M. Artin and B. Mazur, Etale Homotopy. II, 196 Seiten. 1969. DM 12,–

Vol. 101: G. P. Szegö et G. Treccani, Semigruppi di Trasformazioni Multivoche. VI, 177 pages. 1969. DM 14,–

Vol. 102: F. Stummel, Rand- und Eigenwertaufgaben in Sobolewschen Räumen. VIII, 386 Seiten. 1969. DM 20,–

Vol. 103: Lectures in Modern Analysis and Applications I. Edited by C. T. Taam. VII, 162 pages. 1969. DM 12,–

Vol. 104: G. H. Pimbley, Jr., Eigenfunction Branches of Nonlinear Operators and their Bifurcations. II, 128 pages. 1969. DM 10,–

Vol. 105: R. Larsen, The Multiplier Problem. VII, 284 pages. 1969. DM 18,–

Vol. 106: Reports of the Midwest Category Seminar III. Edited by S. Mac Lane. III, 247 pages. 1969. DM 16,–

Vol. 107: A. Peyerimhoff, Lectures on Summability. III, 111 pages. 1969. DM 8,–

Vol. 108: Algebraic K-Theory and its Geometric Applications. Edited by R. M. F. Moss and C. B. Thomas. IV, 86 pages. 1969. DM 6,–

Vol. 109: Conference on the Numerical Solution of Differential Equations. Edited by J. Ll. Morris. VI, 275 pages. 1969. DM 18,–

Vol. 110: The Many Facets of Graph Theory. Edited by G. Chartrand and S. F. Kapoor. VIII, 290 pages. 1969. DM 18,–

[print rear-718]Vol. 111: K. H. Mayer, Relationen zwischen charakteristischen Zahlen. III, 99 Seiten. 1969. DM 8,–

Vol. 112: Colloquium on Methods of Optimization. Edited by N. N. Moiseev. IV, 293 pages. 1970. DM 18,–

Vol. 113: R. Wille, Kongruenzklassengeometrien. III, 99 Seiten. 1970. DM 8,–

Vol. 114: H. Jacquet and R. P. Langlands, Automorphic Forms on GL (2). VII, 548 pages. 1970. DM 24,–

Vol. 115: K. H. Roggenkamp and V. Huber-Dyson, Lattices over Orders I. XIX, 290 pages. 1970. DM 18,–

Vol. 116: Séminaire Pierre Lelong (Analyse) Année 1969. IV, 195 pages. 1970. DM 14,–

Vol. 117: Y. Meyer, Nombres de Pisot, Nombres de Salem et Analyse Harmonique. 63 pages. 1970. DM 6.–

Vol. 118: Proceedings of the 15th Scandinavian Congress, Oslo 1968. Edited by K. E. Aubert and W. Ljunggren. IV, 162 pages. 1970. DM 12,–

Vol. 119: M. Raynaud, Faisceaux amples sur les schémas en groupes et les espaces homogènes. III, 219 pages. 1970. DM 14,–

Vol. 120: D. Siefkes, Büchi's Monadic Second Order Successor Arithmetic. XII, 130 Seiten. 1970. DM 12,–

Vol. 121: H. S. Bear, Lectures on Gleason Parts. III, 47 pages. 1970. DM 6,–

Vol. 122: H. Zieschang, E. Vogt und H.-D. Coldewey, Flächen und ebene diskontinuierliche Gruppen. VIII, 203 Seiten. 1970. DM 16,–

Vol. 123: A. V. Jategaonkar, Left Principal Ideal Rings. VI, 145 pages. 1970. DM 12,–

Vol. 124: Séminaire de Probabilités IV. Edited by P. A. Meyer. IV, 282 pages. 1970. DM 20,–

Vol. 125: Symposium on Automatic Demonstration. V, 310 pages. 1970. DM 20,–

Vol. 126: P. Schapira, Théorie des Hyperfonctions. XI, 157 pages. 1970. DM 14,–

Vol. 127: I. Stewart, Lie Algebras. IV, 97 pages. 1970. DM 10,–

Vol. 128: M. Takesaki, Tomita's Theory of Modular Hilbert Algebras and its Applications. II, 123 pages. 1970. DM 10,–

Vol. 129: K. H. Hofmann, The Duality of Compact Semigroups and C*-Bigebras. XII, 142 pages. 1970. DM 14,–

Vol. 130: F. Lorenz, Quadratische Formen über Körpern. II, 77 Seiten. 1970. DM 8,–

Vol. 131: A Borel et al., Seminar on Algebraic Groups and Related Finite Groups. VII, 321 pages. 1970. DM 22,–

Vol. 132: Symposium on Optimization. III, 348 pages. 1970. DM 22,–

Vol. 133: F. Topsøe, Topology and Measure. XIV, 79 pages. 1970. DM 8,–

Vol. 134: L. Smith, Lectures on the Eilenberg-Moore Spectral Sequence. VII, 142 pages. 1970. DM 14,–

Vol. 135: W. Stoll, Value Distribution of Holomorphic Maps into Compact Complex Manifolds. II, 267 pages. 1970. DM 18,–

Vol. 136: M. Karoubi et al., Séminaire Heidelberg-Saarbrücken-Strasbourg sur la K-Théorie. IV, 264 pages. 1970. DM 18,–

Vol. 137: Reports of the Midwest Category Seminar IV. Edited by S. MacLane. III, 139 pages. 1970. DM 12,–

Vol. 138: D. Foata et M. Schützenberger, Théorie Géométrique des Polynômes Eulériens. V, 94 pages. 1970. DM 10,–

Vol. 139: A. Badrikian, Séminaire sur les Fonctions Aléatoires Linéaires et les Mesures Cylindriques. VII, 221 pages. 1970. DM 18,–

Vol. 140: Lectures in Modern Analysis and Applications II. Edited by C. T. Taam. VI, 119 pages. 1970. DM 10,–

Vol. 141: G. Jameson, Ordered Linear Spaces. XV, 194 pages. 1970. DM 16,–

Vol. 142: K. W. Roggenkamp, Lattices over Orders II. V, 388 pages. 1970. DM 22,–

Vol. 143: K. W. Gruenberg, Cohomological Topics in Group Theory. XIV, 275 pages. 1970. DM 20,–

Vol. 144: Seminar on Differential Equations and Dynamical Systems, II. Edited by J. A. Yorke. VIII, 268 pages. 1970. DM 20,–

Vol. 145: E. J. Dubuc, Kan Extensions in Enriched Category Theory. XVI, 173 pages. 1970. DM 16,–

Vol. 146: A. B. Altman and S. Kleiman, Introduction to Grothendieck Duality Theory. II, 192 pages. 1970. DM 18,–

Vol. 147: D. E. Dobbs, Cech Cohomological Dimensions for Commutative Rings. VI, 176 pages. 1970. DM 16,–

Vol. 148: R. Azencott, Espaces de Poisson des Groupes Localement Compacts. IX, 141 pages. 1970. DM 14,–

Vol. 149: R. G. Swan and E. G. Evans, K-Theory of Finite Groups and Orders. IV, 237 pages. 1970. DM 20,–

Vol. 150: Heyer, Dualität lokalkompakter Gruppen. XIII, 372 Seiten. 1970. DM 20,–

Vol. 151: M. Demazure et A. Grothendieck, Schémas en Groupes I. (SGA 3). XV, 562 pages. 1970. DM 24,–

Vol. 152: M. Demazure et A. Grothendieck, Schémas en Groupes II. (SGA 3). IX, 654 pages. 1970. DM 24,–

Vol. 153: M. Demazure et A. Grothendieck, Schémas en Groupes III. (SGA 3). VIII, 529 pages. 1970. DM 24,–

Vol. 154: A. Lascoux et M. Berger, Variétés Kähleriennes Compactes. VII, 83 pages. 1970. DM 8,–

Vol. 155: Several Complex Variables I, Maryland 1970. Edited by J. Horváth. IV, 214 pages. 1970. DM 18,–

Vol. 156: R. Hartshorne, Ample Subvarieties of Algebraic Varieties. XIV, 256 pages. 1970. DM 20,–

Vol. 157: T. tom Dieck, K. H. Kamps und D. Puppe, Homotopietheorie. VI, 265 Seiten. 1970. DM 20,–

Vol. 158: T. G. Ostrom, Finite Translation Planes. IV. 112 pages. 1970. DM 10,–

Vol. 159: R. Ansorge und R. Hass. Konvergenz von Differenzenverfahren für lineare und nichtlineare Anfangswertaufgaben. VIII, 145 Seiten. 1970. DM 14,–

Vol. 160: L. Sucheston, Contributions to Ergodic Theory and Probability. VII, 277 pages. 1970. DM 20,–

Vol. 161: J. Stasheff, H-Spaces from a Homotopy Point of View. VI, 95 pages. 1970. DM 10,–

Vol. 162: Harish-Chandra and van Dijk, Harmonic Analysis on Reductive p-adic Groups. IV, 125 pages. 1970. DM 12,–

Vol. 163: P. Deligne, Equations Différentielles à Points Singuliers Reguliers. III, 133 pages. 1970. DM 12,–

Vol. 164: J. P. Ferrier, Seminaire sur les Algebres Complètes. II, 69 pages. 1970. DM 8,–

Vol. 165: J. M. Cohen, Stable Homotopy. V, 194 pages. 1970. DM 16,–

Vol. 166: A. J. Silberger, PGL_2 over the p-adics: its Representations, Spherical Functions, and Fourier Analysis. VII, 202 pages. 1970. DM 18,–

Vol. 167: Lavrentiev, Romanov and Vasiliev, Multidimensional Inverse Problems for Differential Equations. V, 59 pages. 1970. DM 10,–

Vol. 168: F. P. Peterson, The Steenrod Algebra and its Applications: A conference to Celebrate N. E. Steenrod's Sixtieth Birthday. VII, 317 pages. 1970. DM 22,–

Vol. 169: M. Raynaud, Anneaux Locaux Henséliens. V, 129 pages. 1970. DM 12,–

Vol. 170: Lectures in Modern Analysis and Applications III. Edited by C. T. Taam. VI, 213 pages. 1970. DM 18,–

Vol. 171: Set-Valued Mappings, Selections and Topological Properties of 2^X. Edited by W. M. Fleischman. X, 110 pages. 1970. DM 12,–

Vol. 172: Y.-T. Siu and G. Trautmann, Gap-Sheaves and Extension of Coherent Analytic Subsheaves. V, 172 pages. 1971. DM 16,–

Vol. 173: J. N. Mordeson and B. Vinograde, Structure of Arbitrary Purely Inseparable Extension Fields. IV, 138 pages. 1970. DM 14,–

Vol. 174: B. Iversen, Linear Determinants with Applications to the Picard Scheme of a Family of Algebraic Curves. VI, 69 pages. 1970. DM 8,–

Vol. 175: M. Brelot, On Topologies and Boundaries in Potential Theory. VI, 176 pages. 1971. DM 18,–

Vol. 176: H. Popp, Fundamentalgruppen algebraischer Mannigfaltigkeiten. IV, 154 Seiten. 1970. DM 16,–

Vol. 177: J. Lambek, Torsion Theories, Additive Semantics and Rings of Quotients. VI, 94 pages. 1971. DM 12,–

Vol. 178: Th. Bröcker und T. tom Dieck, Kobordismentheorie. XVI, 191 Seiten. 1970. DM 18,–

Vol. 179: Seminaire Bourbaki – vol. 1968/69. Exposés 347-363. IV. 295 pages. 1971. DM 22,–

Vol. 180: Séminaire Bourbaki – vol. 1969/70. Exposés 364-381. IV, 310 pages. 1971. DM 22,–

Vol. 181: F. DeMeyer and E. Ingraham, Separable Algebras over Commutative Rings. V, 157 pages. 1971. DM 16.–

[print rear-719]Vol. 182: L. D. Baumert, Cyclic Difference Sets. VI, 166 pages. 1971. DM 16,–

Vol. 183: Analytic Theory of Differential Equations. Edited by P. F. Hsieh and A. W. J. Stoddart. VI, 225 pages. 1971. DM 20,–

Vol. 184: Symposium on Several Complex Variables, Park City, Utah, 1970. Edited by R. M. Brooks. V, 234 pages. 1971. DM 20,–

Vol. 185: Several Complex Variables II, Maryland 1970. Edited by J. Horváth. III, 287 pages. 1971. DM 24,–

Vol. 186: Recent Trends in Graph Theory. Edited by M. Capobianco/J. B. Frechen/M. Krolik. VI, 219 pages. 1971. DM 18.–

Vol. 187: H. S. Shapiro, Topics in Approximation Theory. VIII, 275 pages. 1971. DM 22,–

Vol. 188: Symposium on Semantics of Algorithmic Languages. Edited by E. Engeler. VI, 372 pages. 1971. DM 26,–

Vol. 189: A. Weil, Dirichlet Series and Automorphic Forms. V, 164 pages. 1971. DM 16,–

Vol. 190: Martingales. A Report on a Meeting at Oberwolfach, May 17-23, 1970. Edited by H. Dinges. V, 75 pages. 1971. DM 12,–

Vol. 191: Séminaire de Probabilités V. Edited by P. A. Meyer. IV, 372 pages. 1971. DM 26,–

Vol. 192: Proceedings of Liverpool Singularities – Symposium I. Edited by C. T. C. Wall. V, 319 pages. 1971. DM 24,–

Vol. 193: Symposium on the Theory of Numerical Analysis. Edited by J. Ll. Morris. VI, 152 pages. 1971. DM 16,–

Vol. 194: M. Berger, P. Gauduchon et E. Mazet. Le Spectre d'une Variété Riemannienne. VII, 251 pages. 1971. DM 22,–

Vol. 195: Reports of the Midwest Category Seminar V. Edited by J. W. Gray and S. Mac Lane. III, 255 pages. 1971. DM 22,–

Vol. 196: H-spaces – Neuchâtel (Suisse)- Août 1970. Edited by F. Sigrist, V, 156 pages. 1971. DM 16,–

Vol. 197: Manifolds – Amsterdam 1970. Edited by N. H. Kuiper. V, 231 pages. 1971. DM 20,–

Vol. 198: M. Hervé, Analytic and Plurisubharmonic Functions in Finite and infinite Dimensional Spaces. VI, 90 pages. 1971. DM 16.–

Vol. 199: Ch. J. Mozzochi, On the Pointwise Convergence of Fourier Series. VII, 87 pages. 1971. DM 16,–

Vol. 200: U. Neri, Singular Integrals. VII, 272 pages. 1971. DM 22,–

Vol. 201: J. H. van Lint, Coding Theory. VII, 136 pages. 1971. DM 16,–

Vol. 202: J. Benedetto, Harmonic Analysis on Totally Disconnected Sets. VIII, 261 pages. 1971. DM 22,–

Vol. 203: D. Knutson, Algebraic Spaces. VI, 261 pages. 1971. DM 22,–

Vol. 204: A. Zygmund, Intégrales Singulières. IV, 53 pages. 1971. DM 12,–

Vol. 205: Séminaire Pierre Lelong (Analyse) Année 1970. VI, 243 pages. 1971. DM 20,–

Vol. 206: Symposium on Differential Equations and Dynamical Systems. Edited by D. Chillingworth. XI, 173 pages. 1971. DM 16,–

Vol. 207: L. Bernstein, The Jacobi-Perron Algorithm – Its Theory and Application. IV, 161 pages. 1971. DM 16,–

Vol. 208: A. Grothendieck and J. P. Murre, The Tame Fundamental Group of a Formal Neighbourhood of a Divisor with Normal Crossings on a Scheme. VIII, 133 pages. 1971. DM 16,–

Vol. 209: Proceedings of Liverpool Singularities Symposium II. Edited by C. T. C. Wall. V, 280 pages. 1971. DM 22,–

Vol. 210: M. Eichler, Projective Varieties and Modular Forms. III, 118 pages. 1971. DM 16,–

Vol. 211: Théorie des Matroïdes. Edité par C. P. Bruter. III, 108 pages. 1971. DM 16,–

Vol. 212: B. Scarpellini, Proof Theory and Intuitionistic Systems. VII, 291 pages. 1971. DM 24,–

Vol. 213: H. Hogbe-Nlend, Théorie des Bornologies et Applications. V, 168 pages. 1971. DM 18,–

Vol. 214: M. Smorodinsky, Ergodic Theory, Entropy. V, 64 pages. 1971. DM 12,–

Vol. 215: P. Antonelli, D. Burghelea and P. J. Kahn, The Concordance-Homotopy Groups of Geometric Automorphism Groups. X, 140 pages. 1971. DM 16,–

Vol. 216: H. Maaß, Siegel's Modular Forms and Dirichlet Series. VII, 328 pages. 1971. DM 20,–

Vol. 217: T. J. Jech, Lectures in Set Theory with Particular Emphasis on the Method of Forcing. V, 137 pages. 1971. DM 16,–

Vol. 218: C. P. Schnorr, Zufälligkeit und Wahrscheinlichkeit. IV, 212 Seiten 1971. DM 20,–

Vol. 219: N. L. Alling and N. Greenleaf, Foundations of the Theory of Klein Surfaces. IX, 117 pages. 1971. DM 16,–

Vol. 220: W. A. Coppel, Disconjugacy. V, 148 pages. 1971. DM 16,–

Vol. 221: P. Gabriel und F. Ulmer, Lokal präsentierbare Kategorien. V, 200 Seiten. 1971. DM 18,–

Vol. 222: C. Meghea, Compactification des Espaces Harmoniques. III, 108 pages. 1971. DM 16,–

Vol. 223: U. Felgner, Models of ZF-Set Theory. VI, 173 pages. 1971. DM 16,–

Vol. 224: Revêtements Etales et Groupe Fondamental. (SGA 1). Dirigé par A. Grothendieck XXII, 447 pages. 1971. DM 30,–

Vol. 225: Théorie des Intersections et Théorème de Riemann-Roch. (SGA 6). Dirigé par P. Berthelot, A. Grothendieck et L. Illusie. XII, 700 pages. 1971. DM 40,–

[print rear-720]ISBN 3-540-05647-5

ISBN 0-387-05647-5

Apparatus

[1] EN-FRONT-001 Source folio VIII. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=10; scan/SGA6_witness_300dpi.djvu#page=8
Printed reading: Lectures Notes in Mathematics
Corrected reading: Lecture Notes in Mathematics
Source: ASTRA-FM-008. Both scans print Lectures Notes. The series title is printed as Lecture Notes in Mathematics on the original title page (main PDF3; witness1). Remove the spurious s in the corrected reference.

[2] EN-VAR-001 Source folio 1. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=15
Printed reading: $f : X. \longrightarrow Y$
Corrected reading: $f : X \longrightarrow Y$
Source: fixes/MISPRINTS_6.json#2; check/findings/findings_001_005.json#6. The distinct baseline stop after X is printed before the arrow; corrected form removes it.

[3] EN-VAR-002 Source folio 1. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=15
Printed reading: $\mathrm{ch}_Y\big(f_*(\mathrm{cl}(F)\big) = f_*\big(\mathrm{Todd}(T_f) \cdot \mathrm{ch}_X(F)\big)$,
Corrected reading: $\mathrm{ch}_Y\big(f_*(\mathrm{cl}(F))\big) = f_*\big(\mathrm{Todd}(T_f) \cdot \mathrm{ch}_X(F)\big)$,
Source: fixes/MISPRINTS_6.json#4; check/findings/findings_001_005.json#10. Three left parentheses have only two corresponding right parentheses before the equality sign in the print; the corrected reading closes the argument of ch_Y.

[4] EN-VAR-003 Source folio 4. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=18
Printed reading: $(j = i)$
Corrected reading: $(j \leq i)$
Source: fixes/MISPRINTS_6.json#12; check/findings/findings_001_005.json#30. Print reads j=i. The coefficient of t^i in lambda_{t/(1-t)} is a linear combination of lambda^j for j from 1 to i, including lower-degree terms; corrected range is j≤i.

[5] EN-VAR-004 Source folio 6. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=20
Printed reading: $f_*(\mathrm{cl}^\bullet(\underline{E})) = \sum_i (-1)^i\, \mathrm{cl}^\bullet\big(R^i f_*(\underline{F})\big)$,
Corrected reading: $f_*(\mathrm{cl}^\bullet(\underline{F})) = \sum_i (-1)^i\, \mathrm{cl}^\bullet\big(R^i f_*(\underline{F})\big)$,
Source: fixes/MISPRINTS_6.json#21; check/findings/findings_006_010.json#4. The printed left member is an underlined E, while the right member and subsequent explanatory text use F. Corrected E to F.

[6] EN-VAR-005 Source folio 7. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=21
Printed reading: $K^\bullet(X) \longrightarrow K(\mathrm{Parf}(Y))$,
Corrected reading: $K^\bullet(Y) \longrightarrow K(\mathrm{Parf}(Y))$,
Source: fixes/MISPRINTS_6.json#22; check/findings/findings_006_010.json#9. The printed domain is K^bullet(X). The canonical construction in this paragraph sends a locally free Module on Y to its perfect complex on Y; the corrected domain is K^bullet(Y).

[7] EN-AGG-VAR-001 Source folio 8. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=22
Printed reading: $\mathrm{cl}(L_\bullet)$
Corrected reading: $\mathrm{cl}^\bullet(L_\bullet)$
Source: fixes/FIXLOG_6_CURRENT.jsonl#339; review/early/decisions.jsonl. The scan lacks a raised dot after the first cl, while the end of p. 7 names this same class cl with raised dot in K with raised dot(Y). The corrected raised dot preserves that expressly defined class map.

[8] EN-VAR-067 Source folio 8. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=22
Printed reading: the class T_f defined by formula (2.1)
Corrected reading: the class T_f defined by formula (2.2)
Source: ASTRA-EARLY-0002; review/early/decisions.jsonl. Formula2.1 is the factorization X→X-prime→Y. Formula2.2 defines the virtual cotangent class check T_f, with T_f its dual in the immediately following sentence. The reference specifying this class therefore points to2.2.

[9] EN-VAR-006 Source folio 11. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=25
Printed reading: $d_{X'/X} : \underline{J} \subset \underline{O}_{X'} \longrightarrow \Omega^1_{X'/Y}$,
Corrected reading: $d_{X'/Y} : \underline{J} \subset \underline{O}_{X'} \longrightarrow \Omega^1_{X'/Y}$,
Source: fixes/MISPRINTS_6.json#45; check/findings/findings_011_015.json#2. The printed subscript is X-prime/X. The differential into relative differentials over Y is the universal Y-relative derivation restricted to J, so the corrected subscript is X-prime/Y.

[10] EN-VAR-007 Source folio 14. Scan/witness: scan/SGA6_scan_360dpi.pdf#pages28,33
Printed reading: Grauert's finiteness theorem [4]
Corrected reading: Grauert's finiteness theorem [3]
Source: fixes/MISPRINTS_6.json#55; check/findings/findings_011_015.json#20. The print gives reference[4]; directly checked bibliography entry[3] is Grauert, whereas[4] is Grothendieck, RRR.

[11] EN-VAR-008 Source folio 14. Scan/witness: scan/SGA6_scan_360dpi.pdf#pages28,33
Printed reading: Shih [7]
Corrected reading: Shih [6]
Source: fixes/MISPRINTS_6.json#56; check/findings/findings_011_015.json#22. The print gives reference[7]; directly checked bibliography entry[6] is Shih.

[12] EN-VAR-009 Source folio 15. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=29
Printed reading: $\mathrm{Filt}^i K^\bullet(X) \cdot \mathrm{Filt}_j(K_\bullet(X) \subset \mathrm{Filt}_{j-i} K_\bullet(X)$,
Corrected reading: $\mathrm{Filt}^i K^\bullet(X) \cdot \mathrm{Filt}_j K_\bullet(X) \subset \mathrm{Filt}_{j-i} K_\bullet(X)$,
Source: fixes/MISPRINTS_6.json#61; check/findings/findings_011_015.json#31. The print opens a parenthesis immediately after Filt_j and never closes it; the corrected form removes that superfluous opening parenthesis.

[13] EN-VAR-010 Source folio 16. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=30
Printed reading: is reduced by passage to the associated graded objects from
Corrected reading: is deduced by passage to the associated graded objects from
Source: ASTRA-EARLY-0004. The print reads se réduit. Passing the filtered projection identity to its associated quotients gives the graded identity; the corrected verb is se déduit. The English reproduces the same verb difference.

[14] EN-VAR-011 Source folio 17. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=31
Printed reading: announced without proof in [6]
Corrected reading: announced without proof in [5]
Source: fixes/MISPRINTS_6.json#39; check/findings/findings_016_020.json#4. The printed reference is[6]; FGA is bibliography entry[5], while[6] is Shih.

[15] EN-VAR-012 Source folio 17. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=31
Printed reading: is [6, C-07 (i)]
Corrected reading: is [5, C-07 (i)]
Source: fixes/MISPRINTS_6.json#40; check/findings/findings_016_020.json#5. The printed reference begins[6]. The identified FGA item belongs to bibliography entry[5].

[16] EN-VAR-013 Source folio 19. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=33
Printed reading: Différential Geometry
Corrected reading: Differential Geometry
Source: fixes/MISPRINTS_6.json#43; check/findings/findings_016_020.json#21. The English publication title is printed with an acute accent in Différential; the corrected form removes the accent.

[17] EN-VAR-014 Source folio 19. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=33
Printed reading: Bull. Soc. math France
Corrected reading: Bull. Soc. math. France
Source: fixes/MISPRINTS_6.json#44; check/findings/findings_016_020.json#23. The print has no abbreviation stop after math; the corrected form supplies it.

[18] EN-AGG-VAR-002 Source folio 23. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=37
Printed reading: $\lambda_t(\lambda^i(x)) = \lambda^i(\lambda_t(x)$,
Corrected reading: $\lambda_t(\lambda^i(x)) = \lambda^i(\lambda_t(x))$,
Source: fixes/MISPRINTS_6.json#34; review/early/decisions.jsonl. Only one closing parenthesis follows the last x, leaving λ^i unclosed. Current supplies the second close.

[19] EN-AGG-VAR-003 Source folio 23. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=37
Printed reading: for $i \geq 1$
Corrected reading: for $i > 1$
Source: fixes/MISPRINTS_6.json#35; review/early/decisions.jsonl. The comparison is visibly ≥. Formula(1.4) states λ_t(1)=1+t, so $\lambda^1(1)=1$ and the vanishing condition must be i>1; current has that correction.

[20] EN-VAR-015 Source folio 23. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=37
Printed reading: $\lambda^n(\lambda^i(x)) = P_{i,n}(\lambda^1 x, \ldots, \lambda^n x)$,
Corrected reading: $\lambda^n(\lambda^i(x)) = P_{i,n}(\lambda^1 x, \ldots, \lambda^{in} x)$,
Source: fixes/MISPRINTS_6.json#36; review/early/decisions.jsonl. The print has final exponent n. Formula1.8 on the same page defines P_{i,n} in variables through a_{in}; substituting a_j=lambda^j(x) gives the final argument lambda^{in}(x).

[21] EN-VAR-017 Source folio 26. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=40
Printed reading: Groupes de Lie albébriques
Corrected reading: Groupes de Lie algébriques
Source: fixes/MISPRINTS_6.json#16; fr/codex/HOLD_VS_CORR.json#28; review/early/decisions.jsonl. The title is retained in French in the English translation. The scan prints albébriques; the corrected title restores algébriques.

[22] EN-VAR-016 Source folio 28. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=42
Printed reading: $x = 1 + \sum_{i\geq 1} x_i^i \in 1 + \widehat{A}^+$
Corrected reading: $x = 1 + \sum_{i\geq 1} x^i \in 1 + \widehat{A}^+$
Source: fixes/MISPRINTS_6.json#18; check/findings/findings_026_030.json#6; review/early/decisions.jsonl. The printed homogeneous component has an extra lower index i. The immediately following specification gives x^i in A^i; the corrected form removes only this extra index.

[23] EN-VAR-018 Source folio 31. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=45
Printed reading: $\widetilde{A}$ ——
Corrected reading: $\widetilde{A} \longrightarrow$
Source: fixes/MISPRINTS_6.json#66; check/findings/findings_031_035.json#3. The primary scan and native witness42 show a plain rule with no arrowhead. The corrected form supplies the arrow for the homomorphism ch stated in the surrounding text.

[24] EN-VAR-019 Source folio 32. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=46
Printed reading: $\sum_{i=0}^{n}(-1)^i\lambda^i(N)$
Corrected reading: $\sum_{i=0}^{q}(-1)^i\lambda^i(N)$
Source: fixes/MISPRINTS_6.json#69; check/findings/findings_031_035.json#12. The print has upper limit n. This proposition specifies N with rank q and components through N^q; proposition1.2 gives vanishing of lambda^i(N) for i>q. The corrected upper limit is q.

[25] EN-VAR-020 Source folio 34. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=48
Printed reading: $\underline{E}_G((\Lambda(N^{(p)})^{\mathrm{alt}})$
Corrected reading: $\underline{E}_G((\Lambda(N^{(p)}))^{\mathrm{alt}})$
Source: fixes/MISPRINTS_6.json#71; check/findings/findings_031_035.json#25. The print leaves the inner grouping around Lambda(N^(p)) unclosed before alt. The corrected form adds that closing parenthesis, matching the alternating-part grouping in formula1.33.

[26] EN-VAR-021 Source folio 34. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=48
Printed reading: (§2, theorem 1.3)
Corrected reading: (§2, theorem 1.1)
Source: ASTRA-EARLY-0011; review/early/decisions.jsonl. The print cites theorem1.3 in section2. The representation-ring statement there is theorem1.1 on p26 and its corollary on p27; proposition1.3 on p32 is in section3 and concerns the Chern/Todd identity.

[27] EN-VAR-022 Source folio 37. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=51
Printed reading: $\binom{n-1}{i}\lambda^i(N,x)$
Corrected reading: $\binom{n-1}{i}\lambda^{n-i}(N,x)$
Source: ASTRA-EARLY-0012; review/early/decisions.jsonl. The print gives exponent i. Formula1.23 identifies gamma^n(x_N) with lambda^n(x_N+n−1); the coefficient of t^n in (1+t)^(n−1)lambda_t(x_N) gives exponent n−i. The n=1 printed case instead gives lambda^0=1.

[28] EN-VAR-023 Source folio 37. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=51
Printed reading: $j \geq 0$
Corrected reading: $j \geq 1$
Source: ASTRA-EARLY-0013; review/early/decisions.jsonl. The print gives j≥0. The stated component formula1.38 contains (j−1)!, and its proof uses formula1.18 in positive degrees. The corrected range j≥1 is the common domain of this formula and that proof.

[29] EN-VAR-024 Source folio 37. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=51
Printed reading: $x^1, .., x^j; N^1, .., N^j$
Corrected reading: $x^1, \ldots, x^j; N^1, \ldots, N^j$
Source: fixes/MISPRINTS_6.json#64; check/findings/findings_036_040.json#5. The primary scan and native witness48 show two dots at each of these two ellipses. The corrected form supplies three dots.

[30] EN-VAR-025 Source folio 37. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=51
Printed reading: $x^1, .., N^j$
Corrected reading: $x^1, \ldots, N^j$
Source: fixes/MISPRINTS_6.json#65; check/findings/findings_036_040.json#6. The print has two dots at this ellipsis, as in formula1.38. The corrected form supplies three dots.

[31] EN-AGG-VAR-004 Source folio 41. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=55
Printed reading: $(i \geq 1)$
Corrected reading: $(i > 1)$
Source: fixes/MISPRINTS_6.json#72; review/early/decisions.jsonl. The printed relation is i≥1. The preceding identity assigns C^1(L(D))=ℓ(D); the vanishing of higher classes of this line bundle starts at i>1. Current already has the corrected bound.

[32] EN-VAR-026 Source folio 41. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=55
Printed reading: $\underline{K}(X)$
Corrected reading: $\underline{K}(\xi(X))$
Source: fixes/MISPRINTS_6.json#73; review/early/decisions.jsonl. Both scans repeat K(X) in the phrase introducing its abbreviation. Formula2.6 names the same group as K(xi(X)); the corrected second occurrence restores xi(X).

[33] EN-VAR-027 Source folio 41. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=55
Printed reading: $A(Y)$
Corrected reading: $\widetilde{A}(Y)$
Source: ASTRA-EARLY-0014; review/early/decisions.jsonl. Formula2.6 gives the codomain of C-tilde as A-tilde. On this construction the induced map sends [n,1+sum a_i] to [n,1+sum f-star(a_i)], as used with2.9. The corrected lower-left object restores the omitted tilde.

[34] EN-VAR-028 Source folio 41. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=55
Printed reading: $A(X)$
Corrected reading: $\widetilde{A}(X)$
Source: ASTRA-EARLY-0014; review/early/decisions.jsonl. Formula2.6 gives the codomain of C-tilde as A-tilde. The induced map on this construction sends [n,1+sum a_i] to [n,1+sum f-star(a_i)]. The corrected lower-right object restores the omitted tilde.

[35] EN-AGG-VAR-005 Source folio 43. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=57
Printed reading: left resolution L
Corrected reading: left resolution $\underline{L}$
Source: fixes/MISPRINTS_6.json#76; review/early/decisions.jsonl. The first L after résolution gauche is plain, while L in Z_d(L) and in résolution finie is underlined. Diplomatic correctly has plain L here; current already restores the complex underline.

[36] EN-VAR-029 Source folio 44. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=58
Printed reading: (which is possible by (i) and (ii);
Corrected reading: (which is possible by (i) and (ii));
Source: fixes/MISPRINTS_6.json#77; review/early/decisions.jsonl. The printed parenthetical explanation does not close before the semicolon. The corrected reading supplies the close there, before the next step of the proof.

[37] EN-AGG-VAR-006 Source folio 44. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=58
Printed reading: $E_{\mathbb{C}_0}(\underline{N}) = 0$
Corrected reading: $\underline{E}_{\mathbb{C}_0}(\underline{N}) = 0$
Source: fixes/MISPRINTS_6.json#78; review/early/decisions.jsonl. This E is plain, while the surrounding Euler-characteristic symbols are underlined. Diplomatic now preserves the plain E; current already restores its underline.

[38] EN-VAR-030 Source folio 48. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=62
Printed reading: $\gamma(\Gamma)$ likewise denotes
Corrected reading: $\gamma(F)$ likewise denotes
Source: fixes/MISPRINTS_6.json#80; check/findings/findings_046_050.json#9. The primary and witness scans show capital Gamma inside gamma, while the same sentence names the sheaf F. The corrected occurrence uses F.

[39] EN-VAR-031 Source folio 49. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=63
Printed reading: $\underline{K}(\xi(X)$
Corrected reading: $\underline{K}(\xi(X))$
Source: fixes/MISPRINTS_6.json#81; check/findings/findings_046_050.json#16. The print has only one closing parenthesis after X, leaving K open. The corrected form supplies the second close.

[40] EN-VAR-032 Source folio 50. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=64
Printed reading: is not of characteristic $\neq 0$
Corrected reading: is not of characteristic 0
Source: fixes/MISPRINTS_6.json#82; check/findings/findings_046_050.json#19. The print contains both the prose negation and the inequality sign. The preceding clause limits the construction to characteristic0; removing the inequality sign states the corresponding obstacle while retaining the printed prose negation.

[41] EN-VAR-033 Source folio 50. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=64
Printed reading: $\underline{\mathrm{Tor}}_i(F, .., F)$
Corrected reading: $\underline{\mathrm{Tor}}_i(F, \ldots, F)$
Source: fixes/MISPRINTS_6.json#83; check/findings/findings_046_050.json#22. The primary scan and native witness61 show two dots in this N.B. expression; the display2.19 immediately above has three. The corrected form supplies three dots here.

[42] EN-VAR-034 Source folio 51. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=65
Printed reading: example 2, for the definition
Corrected reading: example 1, for the definition
Source: ASTRA-EARLY-0015; review/early/decisions.jsonl. ChapterI section2 example1 on p24 defines K(G) from representations. Example2 on p25 concerns vector bundles on X. The printed reference2 is corrected to1.

[43] EN-VAR-035 Source folio 53. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=67
Printed reading: $(fg) = f_!g_!$
Corrected reading: $(fg)_! = f_!g_!$
Source: fixes/MISPRINTS_6.json#84; check/findings/findings_051_055.json#11. Both scans leave the area below (fg) blank while marking f and g on the right. The corrected form supplies the lower shriek for the direct image of the composite.

[44] EN-VAR-036 Source folio 54. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=68
Printed reading: $x. \in \underline{K}(X)$
Corrected reading: $x \in \underline{K}(X)$
Source: fixes/MISPRINTS_6.json#85; check/findings/findings_051_055.json#14. The print has a separate baseline stop after x before the membership sign. The corrected reading removes that stop.

[45] EN-VAR-037 Source folio 54. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=68
Printed reading: canonical injection from Y to X
Corrected reading: canonical injection from X to Y
Source: fixes/MISPRINTS_6.json#86; check/findings/findings_051_055.json#18. The same sentence defines X as a subspace of Y and formula2.28 maps the class on X to its class on Y. Thus the injection is i:X→Y, correcting the printed reversed direction.

[46] EN-VAR-038 Source folio 55. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=69
Printed reading: $\gamma_{X\times k}(D) - p^!(y)$
Corrected reading: $\gamma_{X\times k}(Y) - p^!(y)$
Source: ASTRA-EARLY-0016; review/early/decisions.jsonl. The divisor D is defined only on U×k as the restriction of Y. For j:U×k→X×k, restriction j-shriek sends the global class gamma(Y) to the local class gamma(D). The global difference therefore uses Y.

[47] EN-VAR-039 Source folio 55. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=69
Printed reading: $\gamma_{X\times k}(D) - p^!(y))$
Corrected reading: $\gamma_{X\times k}(Y) - p^!(y)$
Source: ASTRA-EARLY-0016; fixes/MISPRINTS_6.json#87; check/findings/findings_051_055.json#22. The same restriction j-shriek sends the global class of Y to the local divisor D, so the global argument is Y. The print also gives a second unmatched closing parenthesis after y; the corrected expression removes it.

[48] EN-VAR-040 Source folio 57. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=71
Printed reading: a section of V
Corrected reading: a section of G
Source: fixes/MISPRINTS_6.json#88; review/early/decisions.jsonl. The print has V. The definition of G-bar(V) pulls back G(U intersection V) along restriction from F(V); the section therefore belongs to G.

[49] EN-VAR-041 Source folio 57. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=71
Printed reading: filtration $\leq i$
Corrected reading: filtration $\geq i$
Source: ASTRA-EARLY-0017; review/early/decisions.jsonl. The print gives ≤i. By the p48 definition, classes supported in dimension ≤n−i generate K(X)^i. Exactness of2.30 identifies the kernel of restriction with those supported on Y; the stated condition is membership in K(X)^i, hence filtration ≥i.

[50] EN-VAR-042 Source folio 59. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=73
Printed reading: $f_a^! = f_a^!$
Corrected reading: $f_a^! = f_{a'}^!$
Source: fixes/MISPRINTS_6.json#89; review/early/decisions.jsonl. Both scans omit the prime on the second a. The preceding formulas compare the fibres over a and a-prime, so the corrected second subscript is a-prime.

[51] EN-VAR-043 Source folio 59. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=73
Printed reading: $\varphi : A(X) \longrightarrow G(K(X))$
Corrected reading: $\varphi : A(X) \longrightarrow G(\underline{K}(X))$
Source: fixes/MISPRINTS_6.json#90; review/early/decisions.jsonl. This printed K lacks the underline present in the next explanatory line. The corrected form supplies the underline for the stated Grothendieck group.

[52] EN-VAR-044 Source folio 59. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=73
Printed reading: from $\underline{K}(Y)$ to $K(X)$
Corrected reading: from $\underline{K}(Y)$ to $\underline{K}(X)$
Source: fixes/MISPRINTS_6.json#91; review/early/decisions.jsonl. The target K(X) lacks the underline printed under the source K(Y). The corrected form supplies the target underline.

[53] EN-VAR-045 Source folio 60. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=74
Printed reading: whether the $\gamma_P(V_{i;\alpha})$ form a basis
Corrected reading: whether the $\gamma_P(V_{i,\alpha})$ form a basis
Source: fixes/MISPRINTS_6.json#92; review/early/decisions.jsonl. The print has a semicolon in this index pair, while the preceding statement defines it with a comma. The corrected form restores that defined index pair.

[54] EN-VAR-046 Source folio 61. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=75
Printed reading: $\eta_j^{(nj)}$
Corrected reading: $\eta_j^{(n_j)}$
Source: fixes/MISPRINTS_6.json#95; review/early/decisions.jsonl. In the print, j is on the same raised baseline as n in the final exponent. The corrected form restores the indexed dimension n_j defined in the statement.

[55] EN-VAR-047 Source folio 61. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=75
Printed reading: $R((\lambda^k(x_i')), (\lambda^k(\gamma(E_j'))$
Corrected reading: $R((\lambda^k(x_i')), (\lambda^k(\gamma(E_j'))))$
Source: fixes/MISPRINTS_6.json#96; review/early/decisions.jsonl. The print has only three closing parentheses after E-prime-j, leaving the outer R argument open. The corrected expression supplies the fourth close.

[56] EN-VAR-048 Source folio 61. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=75
Printed reading: $\gamma^k(E_i - \varepsilon(E_i))$
Corrected reading: $\gamma^k(E_j - \varepsilon(E_j))$
Source: fixes/MISPRINTS_6.json#97; review/early/decisions.jsonl. The print indexes both E terms by i. The statement indexes the bundles E_j by j from1 to r, separately from the x_i indexed from1 to q; the corrected terms use j.

[57] EN-VAR-049 Source folio 62. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=76
Printed reading: $G(K(X)) \longrightarrow G(A)$
Corrected reading: $G(\underline{K}(X)) \longrightarrow G(A)$
Source: fixes/MISPRINTS_6.json#98; review/early/decisions.jsonl. The K in this printed graded map lacks the underline present on K(X) in the preceding line. The corrected form supplies it. The printed target G(A) is retained.

[58] EN-VAR-050 Source folio 63. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=77
Printed reading: $x \in K(X)$
Corrected reading: $x \in \underline{K}(X)$
Source: fixes/MISPRINTS_6.json#101; review/early/decisions.jsonl. This printed K lacks the underline present on the later graded terms in the same theorem. The corrected form supplies the underline.

[59] EN-VAR-051 Source folio 63. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=77
Printed reading: $\sum_{i \geq \varphi} c^i(x)$
Corrected reading: $\sum_{i \geq 1} c^i(x)$
Source: fixes/MISPRINTS_6.json#102; review/early/decisions.jsonl. The printed lower bound is phi. Since c^0 is the class of gamma^0=1, the leading1 already accounts for degree0. The ChapterI section3 construction has a sum of positive-degree terms, so the corrected bound is1.

[60] EN-VAR-052 Source folio 64. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=78
Printed reading: $\gamma^i(x - \varepsilon(X))$
Corrected reading: $\gamma^i(x - \varepsilon(x))$
Source: fixes/MISPRINTS_6.json#104; review/early/decisions.jsonl. The print gives capital X as the augmentation argument. The augmentation is applied to the element x, as in theorem2.16; the corrected argument is lowercase x.

[61] EN-VAR-053 Source folio 64. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=78
Printed reading: $\widetilde{C}(x) \in A(X)^i$
Corrected reading: $\widetilde{C}(x) \in \widetilde{A}(X)^i$
Source: ASTRA-EARLY-0018; review/early/decisions.jsonl. The printed target lacks its tilde. Formula2.7 gives the lambda-homomorphism C-tilde:K(X)→A-tilde(X), whose filtration compatibility is stated on62. The degree-i class of its gamma-expression is the map psi used here, and1.22bis concerns this A-tilde construction.

[62] EN-VAR-054 Source folio 65. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=79
Printed reading: $i_!(xx'\lambda_{-1}(N)$
Corrected reading: $i_!(xx'\lambda_{-1}(N))$
Source: fixes/MISPRINTS_6.json#105; review/early/decisions.jsonl. The print closes the parenthesis of lambda_{−1}(N) but leaves the argument of i-shriek open. The corrected expression supplies its closing parenthesis.

[63] EN-VAR-055 Source folio 66. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=80
Printed reading: by reduction of formulas (ii) and (iv)
Corrected reading: by reduction of formulas (iii) and (iv)
Source: ASTRA-EARLY-0019; review/early/decisions.jsonl. Taking associated-graded classes in formula(iii), i^!i_!(x)=x lambda_{−1}(N), gives(iii bis) because the degree-q term of lambda_{−1}(N) is(−1)^q c^q(N). Formula(iv) similarly gives(iv bis); formula(ii) gives the projection formula(ii bis). Correct the first cited numeral.

[64] EN-VAR-056 Source folio 68. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=82
Printed reading: $\lambda_{-1}(c(N))$
Corrected reading: $\lambda_{-1}(\widetilde{c}(N))$
Source: fixes/MISPRINTS_6.json#106; review/early/decisions.jsonl. Formula2.37 uses the total class c-tilde(N);2.37bis is that identity in total-class form. The printed c before(N) has lost its tilde.

[65] EN-VAR-057 Source folio 68. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=82
Printed reading: the product of their difference by N^q
Corrected reading: the product of their difference by ξ'
Source: ASTRA-EARLY-0020; review/early/decisions.jsonl. For the difference delta on X, the preceding equality gives i^*(delta)=0. Applying i_* and the projection formula(ii bis) gives0=i_*i^*(delta)=i_*(1)delta=xi-prime delta. The printed factor N^q is on Y; its relationship to xi-prime is i^*(xi-prime)=(−1)^q N^q from(iii bis). The stated pushforward therefore yields the factor xi-prime.

[66] EN-VAR-058 Source folio 69. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=83
Printed reading: $\mathrm{ch}_X(f_*(x))$
Corrected reading: $\mathrm{ch}_X(f_!(x))$
Source: ASTRA-EARLY-0021; review/early/decisions.jsonl. The right-hand printed map has a star. Formula2.26 and the following paragraph identify f_! as the map from K(Y) to K(X), which is the domain of ch_X. The left-hand f_* acts on the Chow-valued class and keeps its star.

[67] EN-VAR-059 Source folio 69. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=83
Printed reading: each of these two morphisms..
Corrected reading: each of these two morphisms.
Source: fixes/MISPRINTS_6.json#108; review/early/decisions.jsonl. The printed sentence has two adjacent full stops. The corrected layer retains one.

[68] EN-VAR-062 Source folio 70. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=84
Printed reading: KÜnneth
Corrected reading: Künneth
Source: ASTRA-EARLY-TYPE-016; review/early/decisions.jsonl. The main image and native witnesses81/86 show the tall U-umlaut form on pages70 and75. Preserve the printed name-case KÜnneth diplomatically; the corrected spelling is Künneth.

[69] EN-VAR-060 Source folio 70. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=84
Printed reading: the dimension of Q in P
Corrected reading: the codimension of Q in P
Source: fixes/MISPRINTS_6.json#109; review/early/decisions.jsonl. The class xi is the hyperplane structure-sheaf class. The transverse intersection of p hyperplanes represents xi^p=gamma_P(Q), so p is the codimension of Q.

[70] EN-VAR-061 Source folio 70. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=84
Printed reading: $q = r + 1 - p$
Corrected reading: $q = r - p$
Source: fixes/MISPRINTS_6.json#110; review/early/decisions.jsonl. With H=L/(1−e^{−L}), the displayed product is L^p H^{r+1−p}. Its L^r coefficient equals the L^{r−p} coefficient of H^{r−p+1}; the stated coefficient identity therefore uses q=r−p.

[71] EN-VAR-063 Source folio 71. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=85
Printed reading: Neue topologischen Methoden
Corrected reading: Neue topologische Methoden
Source: fixes/MISPRINTS_6.json#111; review/early/decisions.jsonl. The printed German book title has topologischen. Before plural Methoden without an article, the adjective takes the ending-e, as Neue does. The title is retained in German in the English edition, so its printed spelling is represented explicitly.


[72] EN-VAR-064 Source folio 73. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=87
Printed reading: the functor f thus defined
Corrected reading: the functor f_* thus defined
Source: ASTRA-EARLY-0022; review/early/decisions.jsonl. The preceding operation sends a sheaf G to f_*(G); its right derived functors are the displayed R^p f_*. The printed functor name loses the lower star. Restore it in this occurrence, retaining the morphism f and the K-group map f_! elsewhere.

[73] EN-VAR-065 Source folio 75. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=89
Printed reading: KÜnneth
Corrected reading: Künneth
Source: fixes/MISPRINTS_6.json#115; ASTRA-EARLY-TYPE-016; review/early/decisions.jsonl. The main scan and native witness86 show the tall U-umlaut form, matching page70. Keep the printed KÜnneth diplomatically and the corrected Künneth in current.


[74] EN-VAR-066 Source folio 76. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=90
Printed reading: $(Y', X', y\lambda_{-1}(F))$
Corrected reading: $(Y', X', y\lambda_{-1}(\check{F}))$
Source: fixes/MISPRINTS_6.json#116; review/early/decisions.jsonl. The earlier invocation of Riemann–Roch and formula13 both use the checked bundle F. The class supplied by the corollary is y lambda_{−1}(check F); the second invocation uses that same class. Restore its omitted check.

[75] EN-VAR-068 Source folio 84. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=98
Printed reading: (SGA 1 VI 7,1)
Corrected reading: (SGA 1 VI 7.1)
Source: fixes/MISPRINTS_6.json#94; check/findings/findings_081_085.json#10; review/expose_i/decisions.jsonl. The printed separator has a descending comma tail, while the nearby reference6.1 uses a baseline point. The diplomatic branch retains7,1 and the current branch uses the reference separator7.1.

[76] EN-VAR-069 Source folio 86. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=100; scan/SGA6_witness_300dpi.djvu#page=97
Printed reading:



Corrected reading:



Source: check/findings/findings_086_090.json#6; review/expose_i/decisions.jsonl. The printed diagonal has label u and the horizontal label is U. The diagonal is the chosen lift H_U→E_U; the horizontal is the pullback u_U:E_U→F_U of u. Current removes the diagonal label and writes u_U horizontally.

[77] EN-VAR-070 Source folio 88. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=102; scan/SGA6_witness_300dpi.djvu#page=99
Printed reading: (SGA 4 V 3.11) k
Corrected reading: (SGA 4 V 3.11), k
Source: check/findings/findings_086_090.json#15. The comma separates the topos datum from the following field datum in example c). Its absence is retained diplomatically.

[78] EN-VAR-071 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: **Lemma 1.3.1,**
Corrected reading: **Lemma 1.3.1.**
Source: fixes/MISPRINTS_6.json#123; check/findings/findings_086_090.json#17. The scan has a descending comma after the lemma number. Current uses the full stop separating its label from its statement.

[79] EN-VAR-072 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: -Module, Equivalent conditions
Corrected reading: -Module. Equivalent conditions
Source: fixes/MISPRINTS_6.json#124; check/findings/findings_086_090.json#18. Both scans print a comma before the new sentence introducing the equivalent conditions. Current supplies a sentence stop.

[80] EN-VAR-073 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: S be a site,
Corrected reading: $\underline{S}$ be a site,
Source: ASTRA-I-S-089-PROP. The printed plain S denotes the site governing the fibered categories. The statement later declares the object S to belong to the category underlined-S, identifying that base site. Current restores its individual underline; diplomatic preserves the printed glyph.

[81] EN-VAR-074 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: C a flat abelian
Corrected reading: $\underline{C}$ a flat abelian
Source: fixes/MISPRINTS_6.json#125; check/findings/findings_086_090.json#23. The printed plain C occurs in the declaration or subsequent mention of the same flat abelian category over the site. Current restores its category underline; diplomatic preserves the plain glyph.


[82] EN-VAR-075 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: S-category (1.1),
Corrected reading: $\underline{S}$-category (1.1),
Source: ASTRA-I-S-089-PROP. The printed plain S denotes the site governing the fibered categories. The statement later declares the object S to belong to the category underlined-S, identifying that base site. Current restores its individual underline; diplomatic preserves the printed glyph.

[83] EN-VAR-076 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: a strictly full additive sub-S
Corrected reading: a strictly full additive sub-$\underline{S}$
Source: ASTRA-I-S-089-PROP. The printed plain S denotes the site governing the fibered categories. The statement later declares the object S to belong to the category underlined-S, identifying that base site. Current restores its individual underline; diplomatic preserves the printed glyph.

[84] EN-VAR-077 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: C; assume that
Corrected reading: $\underline{C}$; assume that
Source: fixes/MISPRINTS_6.json#126; check/findings/findings_086_090.json#24. The printed plain C occurs in the declaration or subsequent mention of the same flat abelian category over the site. Current restores its category underline; diplomatic preserves the plain glyph.


[85] EN-VAR-078 Source folio 89. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=103; scan/SGA6_witness_300dpi.djvu#page=100
Printed reading: C (1.2). Let
Corrected reading: $\underline{C}$ (1.2). Let
Source: fixes/MISPRINTS_6.json#127; check/findings/findings_086_090.json#25. The printed plain C occurs in the declaration or subsequent mention of the same flat abelian category over the site. Current restores its category underline; diplomatic preserves the plain glyph.

[86] EN-VAR-079 Source folio 92. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=106; scan/SGA6_witness_300dpi.djvu#page=103
Printed reading:



.

Corrected reading:



.

Source: check/findings/findings_091_095.json#5; review/expose_i/decisions.jsonl. The printed u' segment has no arrowhead. The preceding text defines u':E'→F_X and G' as its cone, which fixes the arrow direction toward F_X. Current adds that head; diplomatic retains the bare segment.

[87] EN-VAR-080 Source folio 93. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=107; scan/SGA6_witness_300dpi.djvu#page=104
Printed reading: **Lemma 1.4.2.** Let C
Corrected reading: **Lemma 1.4.2.** Let $\underline{C}$
Source: check/findings/findings_091_095.json#9. The first declared category C has no printed underline. Its occurrence inside the derived category D(C) is underlined in the same sentence; current supplies the same category rule here.

[88] EN-VAR-081 Source folio 93. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=107; scan/SGA6_witness_300dpi.djvu#page=104
Printed reading: a morphism of complexes of C
Corrected reading: a morphism of complexes of $\underline{C}$
Source: check/findings/findings_091_095.json#9. The second category C has no printed underline. It is the category of the complexes E and F and the argument of D(C), whose C is individually underlined; current supplies its missing rule.

[89] EN-VAR-082 Source folio 94. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=108; scan/SGA6_witness_300dpi.djvu#page=105
Printed reading: $v^j = (v_i)^j$
Corrected reading: $v^j = (v_i v_{i+1} \dots v_{n-1})^j$
Source: ASTRA-I-COMPOSITION-094; review/expose_i/add_094_composition.py; EN_INDEPENDENT_SOURCE_ASSESSMENTS.jsonl. The transitions have domains E→E'_(n-1) and E'_(i+1)→E'_i. Their composite V_i=v_i...v_(n-1) has domain E, and its degree-j component stabilizes for i≤j because the lower transitions are identities in that degree. The printed isolated v_i has domain E'_(i+1), so the composite is required to define v:E→E'. The u' formula remains unchanged.

[90] EN-VAR-083 Source folio 95. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=109; scan/SGA6_witness_300dpi.djvu#page=106
Printed reading: $E^i \in \mathrm{ob}\, \underline{C}_{0S}$
Corrected reading: $E^i \in \mathrm{ob}\, \underline{C}_{0X}$
Source: check/findings/findings_091_095.json#19; review/expose_i/decisions.jsonl. The definition places E in the fiber C_X at its arbitrary object X. Its components therefore belong to C_X and the specified strictly full subcategory is C_(0X). Current changes the printed subscript S to X.

[91] EN-VAR-084 Source folio 100. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=114; scan/SGA6_witness_300dpi.djvu#page=111
Printed reading: **Proposition 2.7**
Corrected reading: **Proposition 2.7.**
Source: check/findings/findings_096_100.json#19. The scan has a blank after the final7 before the statement. Diplomatic preserves that missing stop; current retains the restored heading punctuation.

[92] EN-VAR-085 Source folio 103. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=117; scan/SGA6_witness_300dpi.djvu#page=114
Printed reading: n
Corrected reading: $(-n)$
Source: ASTRA-I-SIGN-103. Assertion2.9(ii) specifies (-n)-pseudo-coherence. Apply2.4 to the map0→F with pseudo-coherence parameter -n and quasi-isomorphism parameter1: the replacement E→F is strictly(-n)-pseudo-coherent, and0→E is the identity in degrees≥1. Thus E vanishes there and its components in degrees -n through0 supply exactly the n-presentation defined in2.8. Current supplies the missing minus sign.

[93] EN-VAR-087 Source folio 104. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=118; scan/SGA6_witness_300dpi.djvu#page=115
Printed reading: $F^i \neq 0$
Corrected reading: $F^i = 0$
Source: check/findings/findings_101_105.json#21; review/expose_i/decisions.jsonl. The given complex belongs to the bounded-above category C-minus. The displayed exact sequence removes its highest component F^(n+N), leaving F-prime zero above n+N-1. The induction therefore starts with F^i=0 for i>n+N, replacing the printed nonzero relation.

[94] EN-VAR-086 Source folio 104. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=118; scan/SGA6_witness_300dpi.djvu#page=115
Printed reading: F is pseudo-coherent.
Corrected reading: F is n-pseudo-coherent (resp. pseudo-coherent).
Source: check/findings/findings_101_105.json#20; review/expose_i/decisions.jsonl. The proof on105 concludes n-pseudo-coherence. Its induction uses H^i(F')=H^(i+1)(F), and the presentation index i+1-n equals i-(n-1). The printed unqualified conclusion fails on the punctual site of vector spaces: take n=0, an infinite-dimensional vector space M, and F=M[1]. All H^i(F) for i≥0 vanish and meet the stated finite-presentation assumptions, while H^(-1)(F)=M is not of finite type. Proposition2.10(b) therefore rules out (-1)-pseudo-coherence. Current restores the n-qualified conclusion and the parenthetical alternative proved by the argument.

[95] EN-VAR-088 Source folio 106. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=120; scan/SGA6_witness_300dpi.djvu#page=117
Printed reading: $F'_{\leq n}$) (resp.
Corrected reading: $F'_{\leq n}$ (resp.
Source: check/findings/findings_106_110.json#1. The preceding reference(2.10.b)) is already closed. The printed extra closer after F-prime has no opener; current removes that single character.

[96] EN-VAR-089 Source folio 108. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=122; scan/SGA6_witness_300dpi.djvu#page=119
Printed reading: and C_1).
Corrected reading: and C_1.
Source: check/findings/findings_106_110.json#23. Both numbered references in this sentence are already closed. The final printed parenthesis after C_1 has no opener; current removes it and preserves every parameter substitution.

[97] EN-VAR-090 Source folio 109. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=123; scan/SGA6_witness_300dpi.djvu#page=120
Printed reading: is a field or when radical(A_s) is nilpotent.
Corrected reading: is a field.
Source: ASTRA-I-RADICAL-109; review/expose_i/decisions.jsonl; EN_INDEPENDENT_SOURCE_ASSESSMENTS.jsonl. The printed nilpotence-only alternative is an example for the stated hypothesis that finite projective stalk modules are free. For A=k×k, its radical J=0 is nilpotent, but P=k×0 is the image of the idempotent e(a,b)=(a,0). Tensoring along the two coordinate projections gives dimensions1 and0, whereas a free A-module of rank r gives r and r, so P is not free. The retained residue-field example is valid: lift a basis of P/JP to a surjection A^r→P by Nakayama; projectivity splits A^r=P plus K. The chosen basis gives K/JK=0, and K is finitely generated, hence K=0 by Nakayama. Current deletes only the unsupported alternative; the lemma's actual projective-freeness hypothesis is unchanged.

[98] EN-VAR-091 Source folio 110. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=124; scan/SGA6_witness_300dpi.djvu#page=121
Printed reading: Suppose (ii), Let
Corrected reading: Suppose (ii). Let
Source: check/findings/findings_106_110.json#32. Both scans show the comma followed by a capital. The diplomatic reading retains it; current separates the two sentences with a full stop.

[99] EN-VAR-092 Source folio 110. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=124; scan/SGA6_witness_300dpi.djvu#page=121
Printed reading: U_n
Corrected reading: u_n
Source: check/findings/findings_106_110.json#34. The matrix(T_ij) defines one universal endomorphism. The morphism s satisfies s*(u_n)=p, and its factorization s=i composed with s-prime gives p=s-prime* i*(u_n). All three lowercase references identify that same map. Current gives its definition the lowercase name u_n as well; diplomatic retains the printed capital U_n.

[100] EN-VAR-093 Source folio 111. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=125; scan/SGA6_witness_300dpi.djvu#page=122
Printed reading: $X' = u(X)$
Corrected reading: $X' = f^{-1}(X)$
Source: check/findings/findings_111_115.json#5; review/expose_i/decisions.jsonl. The site-object inverse-image functor f^(-1):S→S' is explicitly defined on110. The proof on111 applies that same map to U, writing U'=f^(-1)(U). The letter u in that proof denotes the derived tensor functor on complexes. Current applies the specified site map to X, giving X'=f^(-1)(X).

[101] EN-VAR-094 Source folio 112. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=126; scan/SGA6_witness_300dpi.djvu#page=123
Printed reading: **Lemma 3.3**
Corrected reading: **Lemma 3.3.**
Source: check/findings/findings_111_115.json#17. The printed heading ends after the numeral3. Diplomatic keeps the missing terminal stop; current supplies the heading punctuation.

[102] EN-VAR-095 Source folio 118. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=132; scan/SGA6_witness_300dpi.djvu#page=129
Printed reading: (SGAA VIII 7.8)
Corrected reading: (SGA 4 VIII 7.8)
Source: ASTRA-I-SGA4-118; review/expose_i/decisions.jsonl; coordinator primary bibliography verification. The printed final A is retained diplomatically. The cited definition of a point belongs to SGA 4, VIII7.8, checked against the primary text by the coordinator; current supplies the volume digit and preserves the cited exposé and paragraph.

[103] EN-VAR-096 Source folio 118. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=132; scan/SGA6_witness_300dpi.djvu#page=129
Printed reading: By (2.22 a)),
Corrected reading: By (3.7 a)),
Source: check/findings/findings_116_120.json#22; review/expose_i/decisions.jsonl. The displayed comparison is the degree-zero consequence of the flat point pullback of the derived Hom comparison in3.7(a). The point inverse-image functor is exact, and applying H^0 to that isomorphism gives the two displayed Hom comparisons. Current cites3.7(a); diplomatic retains2.22(a).

[104] EN-VAR-097 Source folio 123. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=137; scan/SGA6_witness_300dpi.djvu#page=134
Printed reading: **Proposition 4.10**
Corrected reading: **Proposition 4.10.**
Source: fixes/MISPRINTS_6.json#144; check/findings/findings_121_125.json#22. The main scan has a space after the numeral before the statement, with no terminal full stop. Diplomatic retains that absence; current supplies the heading punctuation.

[105] EN-VAR-098 Source folio 123. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=137; scan/SGA6_witness_300dpi.djvu#page=134
Printed reading: $X \in \mathrm{ob}\, S$
Corrected reading: $X \in \mathrm{ob}\, \underline{S}$
Source: ASTRA-I-SITE-MISSING-123; review/expose_i/decisions.jsonl. The printed membership leaves S without a rule. The quantified X indexes the fibers C_{0X}, and the fiber projection C_0→the base site sends their objects to X. Paragraph4.0 names that base site with underlined S; the proof124 repeats X∈ob underlined S for the same triangle fibers. Current supplies the missing site rule; diplomatic retains the observed plain letter.

[106] EN-VAR-099 Source folio 123. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=137; scan/SGA6_witness_300dpi.djvu#page=134
Printed reading: $\mathrm{Tr}(\mathrm{D}(\underline{C}))$ to an object of
Corrected reading: $\mathrm{Tr}(\mathrm{D}(\underline{C})))$ to an object of
Source: fixes/MISPRINTS_6.json#145; check/findings/findings_121_125.json#34. The middle resp. parenthesis has an opening before the category expression and no outside closer in print. Current adds that single outside parenthesis; the inner category parentheses and all mathematical tokens are unchanged.

[107] EN-VAR-100 Source folio 123. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=137; scan/SGA6_witness_300dpi.djvu#page=134
Printed reading: (resp. $\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0))$.
Corrected reading: (resp. $\mathrm{Tr}(\mathrm{K}^b(\underline{C}_0)))$.
Source: fixes/MISPRINTS_6.json#146; check/findings/findings_121_125.json#35. The final resp. clause independently lacks its outside closing parenthesis. The two inner category parentheses are present. Current supplies only the missing outer closer; diplomatic retains its omission.

[108] EN-VAR-101 Source folio 126. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=140; scan/SGA6_witness_300dpi.djvu#page=137
Printed reading: $\mathrm{K}^b(C_0) \longrightarrow \mathrm{D}(\underline{C})$
Corrected reading: $\mathrm{K}^b(\underline{C}_0) \longrightarrow \mathrm{D}(\underline{C})$
Source: fixes/MISPRINTS_6.json#147; check/findings/findings_126_130.json#3. The first inline source category lacks the C rule in both scans. The full embedding i:C_0→C specified in the preceding sentence induces K^b(i), followed by localization into D(C). Its source is precisely the named C_0 with its independent rule. Current supplies this one omitted rule; diplomatic retains the plain C.

[109] EN-VAR-102 Source folio 126. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=140; scan/SGA6_witness_300dpi.djvu#page=137
Printed reading: $\mathrm{K}^b(\underline{C}_0)\mathrm{K}^{b,\varnothing}(\underline{C}_0) \longrightarrow \mathrm{K}^-(\underline{C}_0)/\mathrm{K}^{-,\varnothing}(\underline{C}_0)$
Corrected reading: $\mathrm{K}^b(\underline{C}_0)/\mathrm{K}^{b,\varnothing}(\underline{C}_0) \longrightarrow \mathrm{K}^-(\underline{C}_0)/\mathrm{K}^{-,\varnothing}(\underline{C}_0)$
Source: fixes/MISPRINTS_6.json#148; check/findings/findings_126_130.json#9. Both scans omit the source quotient slash. The inclusion of bounded complexes into bounded-above complexes sends bounded acyclic complexes to bounded-above acyclic complexes. Composing with localization therefore factors through the bounded quotient K^b(C_0)/K^{b,emptyset}(C_0), which is already displayed earlier in the remark. Current restores that one slash; diplomatic retains the adjacent expressions.

[110] EN-VAR-103 Source folio 129. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=143; scan/SGA6_witness_300dpi.djvu#page=140
Printed reading: parf. $\dim(F) \subset [a, b]$
Corrected reading: parf. $\mathrm{amp}(F) \subset [a, b]$
Source: fixes/MISPRINTS_6.json#149; check/findings/findings_126_130.json#22. Here F=F-prime+F-double-prime. Hypothesis4.17(ii) gives the stated perfect-amplitude inclusion, and the preceding truncations put its terms in C_0 above a. Lemma4.16 applies with exactly that amplitude hypothesis and gives F^a locally in C_0. Current writes parf.amp; diplomatic retains the printed parf.dim.

[111] EN-VAR-104 Source folio 129. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=143; scan/SGA6_witness_300dpi.djvu#page=140
Printed reading: $\mathrm{D}^*(\underline{C})$ (4.10).
Corrected reading: $\mathrm{D}^*(\underline{C}')$ (4.10).
Source: fixes/MISPRINTS_6.json#150; check/findings/findings_126_130.json#29. The displayed perfect subcategory is defined for C-prime. Definition4.9 supplies the full inclusion j:D*(C-prime)_parf→D*(C-prime), the identity on its objects and arrows, and4.10 applied to the primed triple gives its triangulated structure. Current adds the omitted prime to this ambient category; diplomatic retains the printed unprimed letter.

[112] EN-VAR-105 Source folio 135. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=149; scan/SGA6_witness_300dpi.djvu#page=146
Printed reading: (2,15)
Corrected reading: (2.15)
Source: fixes/MISPRINTS_6.json#136; check/findings/findings_131_135.json#31. The printed reference has a comma. Paragraph 2.15 defines the pseudo-coherence of Modules used in this corollary. Diplomatic preserves the observed comma; current uses the paragraph number 2.15.

[113] EN-VAR-106 Source folio 136. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=150; scan/SGA6_witness_300dpi.djvu#page=147
Printed reading: (3.17.1)
Corrected reading: (5.8.2.1)
Source: ASTRA-I-REF-136; review/expose_i/own_findings.jsonl; review/expose_i/add_136_140.py. Formula 5.8.2.1 specializes at B=Z_S and G=I to the displayed vertical map Hom_ZS(F,I) tensor_A E→Hom_ZS(Hom_A(E,F),I), sending h tensor e to (t↦h(t(e))). Replacing F by F-double-prime gives the other vertical map. The next sentence invokes Lemma 5.8.2 for invertibility. Current gives this actual formula reference; diplomatic preserves the printed 3.17.1.

[114] EN-VAR-107 Source folio 142. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=156; scan/SGA6_witness_300dpi.djvu#page=153
Printed reading: for every $X \in \mathrm{ob}\,\underline{C}_X$
Corrected reading: for every $X \in \mathrm{ob}\,\underline{S}$
Source: ASTRA-I-INDEX-142; check/findings/findings_141_145.json#9. The declared family assigns f_U:ob(C_U) to F(U) to every U in ob(S). Its additivity is the identity f_U(L)=f_U(L-prime)+f_U(L-double-prime) on a direct sum or distinguished triangle in that fibre. The selecting quantifier is therefore U in ob(S), with the source dummy variable X retained; diplomatic keeps the printed fibre-object quantifier.

[115] EN-VAR-108 Source folio 142. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=156; scan/SGA6_witness_300dpi.djvu#page=153
Printed reading:

$$\begin{array}{ccc} \mathrm{ob}\,\underline{C}_Y & \xrightarrow{f_Y} & F(Y) \\ {\scriptstyle u^*}\downarrow & & \downarrow{\scriptstyle F(u)} \\ \mathrm{ob}\,\underline{C}_X & \xrightarrow{f_X} & F(X). \end{array}$$

Corrected reading:

$$\begin{array}{ccc} \mathrm{ob}\,\underline{C}_Y & \xrightarrow{f_Y} & F(Y) \\ {\scriptstyle u^*}\downarrow & & \downarrow{\scriptstyle F(u)} \\ \mathrm{ob}\,\underline{C}_X & \xrightarrow{f_X} & F(X). \end{array}$$

Source: check/findings/findings_141_145.json#10. The source lower segment has no arrowhead. The declared map f_X:ob(C_X) to F(X) determines its rightward direction, and the square states f_X(u-star L)=F(u)(f_Y(L)). Current supplies that arrowhead; diplomatic preserves the headless segment. Every endpoint, other arrow and label is identical.

[116] EN-VAR-109 Source folio 143. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=157; scan/SGA6_witness_300dpi.djvu#page=154
Printed reading: for $f \in \underline{\mathrm{Cart}}_{\underline{S}}(\underline{C}^{\mathrm{is}}, F)$ and
Corrected reading: for $f \in \underline{\mathrm{Cart}}_{\underline{S}}(\underline{C}^{\mathrm{is}}, \underline{F})$ and
Source: check/findings/findings_141_145.json#17. The printed isolated F lacks its rule. The discrete fibred category constructed on142 assigns to x in F(U) the object (U,x), with restriction arrows given by F(u); projection to x is the inverse description. A cartesian functor sends c in C_U to (U,f_U(c)). This explicitly identifies the intended codomain with the underlined F in both adjacent phi targets. Current restores that rule; diplomatic retains the observed plain F.

[117] EN-VAR-110 Source folio 157. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=171; scan/SGA6_witness_300dpi.djvu#page=168
Printed reading:

$$\begin{array}{ccc} E^\vee \otimes^{\mathrm{L}} F \otimes^{\mathrm{L}} F^\vee \otimes^{\mathrm{L}} E & \xrightarrow{\mathrm{Id} \otimes^{\mathrm{L}} \tau_E} & F^\vee \otimes^{\mathrm{L}} F \\ {\scriptstyle \mathrm{Id} \otimes \tau_F}\downarrow & & \downarrow{\scriptstyle \tau_F} \\ E^\vee \otimes^{\mathrm{L}} E & \xrightarrow[\tau_E]{} & A. \end{array}$$

Corrected reading:

$$\begin{array}{ccc} E^\vee \otimes^{\mathrm{L}} F \otimes^{\mathrm{L}} F^\vee \otimes^{\mathrm{L}} E & \xrightarrow{\mathrm{Id} \otimes^{\mathrm{L}} \tau_E} & F^\vee \otimes^{\mathrm{L}} F \\ {\scriptstyle \mathrm{Id} \otimes^{\mathrm{L}} \tau_F}\downarrow & & \downarrow{\scriptstyle \tau_F} \\ E^\vee \otimes^{\mathrm{L}} E & \xrightarrow[\tau_E]{} & A. \end{array}$$

Source: check/findings/findings_156_160.json#12; review/expose_i/add_156_159.py. The square left edge lacks the L that is printed on the same contraction in the parallel triangle. Its map is beta=id_(E-dual) tensor^L (ev_F composed with c_(F,F-dual)) tensor^L id_E, from E-dual tensor^L F tensor^L F-dual tensor^L E to E-dual tensor^L E. Here c exchanges the middle factors, ev_F:F-dual tensor^L F to A is the evaluation used in8.1, and the unit isomorphisms remove the A factor. Current restores the omitted derived-tensor exponent; diplomatic retains the ordinary tensor label. All other source cells, labels and arrows are identical.

[118] EN-VAR-111 Source folio 163. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=177; scan/SGA6_witness_300dpi.djvu#page=174
Printed reading: $\underline{S}$ is quasi-compact
Corrected reading: $\underline{S}$ is quasi-compact
Source: ASTRA-II-OBJECT-S-163; review/expose_ii/add_163.py. Both scans give the object S a rule in this sentence. The preceding declaration selects S as an object of the underlined site, and each displayed covering arrow S_i to S has that same object as codomain. The identity of this declared object S identifies the two typographical occurrences. Current removes only this isolated object-letter rule; diplomatic preserves it. No object or covering morphism is changed.

[119] EN-VAR-112 Source folio 163. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=177; scan/SGA6_witness_300dpi.djvu#page=174
Printed reading: C'_S
Corrected reading: $\underline{C'_S}$
Source: check/findings/findings_161_165.json#9; review/expose_ii/add_163.py. The introduced C-prime has no rule in either scan at this occurrence. Its specified inclusion into C_S identifies it with the same category that is individually underlined throughout164, including the domain of Gamma_S and the exact-sequence hypothesis. The identity functor of this explicitly included subcategory gives that identification. Current supplies the rule only here; diplomatic retains the observed plain letter.

[120] EN-VAR-113 Source folio 165. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=179; scan/SGA6_witness_300dpi.djvu#page=176
Printed reading: $\mathrm{Hom}_{A(S)}(.,\Gamma_S(F))$
Corrected reading: $\mathrm{Hom}_{A(S)}(.,\Gamma_S(G))$
Source: check/findings/findings_161_165.json#18; fixes/FIXLOG_6_CURRENT.jsonl#347. The second presentation is an exact sequence of A(S)-modules ending in Gamma_S(F). Applying Hom_(A(S))(-,Gamma_S(G)) gives the bottom row of the printed diagram, whose two right-hand terms explicitly fix Gamma_S(G). The vertical comparison sends a sheaf map F to G to its section map Gamma_S(F) to Gamma_S(G). Current therefore restores G in this fixed second argument. The separately inspected placeholder is a baseline period in both forms.

[121] EN-VAR-114 Source folio 165. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=179; scan/SGA6_witness_300dpi.djvu#page=176
Printed reading:

$$\begin{array}{ccccc} 0 \longrightarrow \mathrm{Hom}_{A_S}(F,G) & \longrightarrow & \mathrm{Hom}_{A_S}(A_S^{(J)},G) & \longrightarrow & \mathrm{Hom}_{A_S}(A_S^{(I)},G) \\ \downarrow & & \downarrow & & \downarrow \\ 0 \longrightarrow \mathrm{Hom}_{A(S)}(F,G) & \longrightarrow & \mathrm{Hom}_{A(S)}(A(S)^{(J)},\Gamma_S(G)) & \longrightarrow & \mathrm{Hom}_{A(S)}(A(S)^{(I)},\Gamma_S(G)). \end{array}$$

Corrected reading:

$$\begin{array}{ccccc} 0 \longrightarrow \mathrm{Hom}_{A_S}(F,G) & \longrightarrow & \mathrm{Hom}_{A_S}(A_S^{(J)},G) & \longrightarrow & \mathrm{Hom}_{A_S}(A_S^{(I)},G) \\ \downarrow & & \downarrow & & \downarrow \\ 0 \longrightarrow \mathrm{Hom}_{A(S)}(\Gamma_S(F),\Gamma_S(G)) & \longrightarrow & \mathrm{Hom}_{A(S)}(A(S)^{(J)},\Gamma_S(G)) & \longrightarrow & \mathrm{Hom}_{A(S)}(A(S)^{(I)},\Gamma_S(G)). \end{array}$$

Source: check/findings/findings_161_165.json#19; review/expose_ii/add_165.py. Applying Hom_(A(S))(-,Gamma_S(G)) to the displayed section presentation(**) gives the left nonzero term Hom_(A(S))(Gamma_S(F),Gamma_S(G)). The vertical arrow is the map f↦Gamma_S(f) from sheaf morphisms F to G to section-module morphisms. Current supplies the two omitted section functors in that one bottom node; diplomatic retains F,G. Every other node, arrow, argument and punctuation is identical.

[122] EN-VAR-115 Source folio 165. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=179; scan/SGA6_witness_300dpi.djvu#page=176
Printed reading: are visibly isomorphisms the same
Corrected reading: are visibly isomorphisms; the same
Source: check/findings/findings_161_165.json#20. The printed sentence has no punctuation between its two finite clauses. Current inserts the separating semicolon; diplomatic preserves the absence. Both claims and their order are unchanged.

[123] EN-VAR-116 Source folio 168. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=182; scan/SGA6_witness_300dpi.djvu#page=179
Printed reading: $\mathrm{D}^-(\mathrm{Qcoh}(S)_{\mathrm{coh}}$
Corrected reading: $\mathrm{D}^-(\mathrm{Qcoh}(S))_{\mathrm{coh}}$
Source: check/findings/findings_166_170.json#18. The outer parenthesis of D-minus is unclosed in print. Notation2.2.1 defines the essential-image category D-minus(Qcoh(S))_coh on167. Current adds only the missing outside close; diplomatic retains the observed omission.

[124] EN-VAR-117 Source folio 169. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=183; scan/SGA6_witness_300dpi.djvu#page=180
Printed reading: a quasi-coherent Ideal of $\underline{O}_S$
Corrected reading: a quasi-coherent Ideal of $\underline{O}_S$.
Source: check/findings/findings_166_170.json#29. The complete final item(iii-prime) ends without a stop in print. Current supplies the sentence-ending full stop; diplomatic retains its absence.

[125] EN-VAR-118 Source folio 169. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=183; scan/SGA6_witness_300dpi.djvu#page=180
Printed reading: -Module $f \in \Gamma(X, L)$
Corrected reading: -Module, $f \in \Gamma(X, L)$
Source: check/findings/findings_166_170.json#31. The declaration introduces the scheme X, its invertible module L, and the section f in Gamma(X,L). Current inserts the omitted comma delimiting the section declaration after the description of L. All three objects, their membership and the immersion i:X_f to X remain unchanged.

[126] EN-VAR-119 Source folio 171. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=185; scan/SGA6_witness_300dpi.djvu#page=182
Printed reading: $Y' = X' - U$
Corrected reading: $Y' = X' - U'$
Source: check/findings/findings_171_175.json#4; fixes/MISPRINTS_6.json#151. In the displayed Cartesian diagram, with g:X-prime to X, the upper open is U-prime=U times_X X-prime=g inverse(U). Therefore Y-prime=g inverse(X minus U)=X-prime minus U-prime. This exact inverse-image identity gives the corrected prime. Diplomatic retains the printed U.

[127] EN-VAR-120 Source folio 174. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=188; scan/SGA6_witness_300dpi.djvu#page=185
Printed reading: assertions c)) follow
Corrected reading: assertions c) follow
Source: check/findings/findings_171_175.json#10; fixes/MISPRINTS_6.json#153. The item reference has a second closing parenthesis without an opening parenthesis in print. Current removes that unmatched delimiter; the reference remains the same item c).

[128] EN-VAR-121 Source folio 174. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=188; scan/SGA6_witness_300dpi.djvu#page=185
Printed reading: This completes the proof
Corrected reading: This completes the proof.
Source: check/findings/findings_171_175.json#11; fixes/MISPRINTS_6.json#154. The complete final proof sentence has no following full stop in either scan. Current supplies the sentence-ending stop; diplomatic retains the absence.

[129] EN-VAR-122 Source folio 175. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=189; scan/SGA6_witness_300dpi.djvu#page=186
Printed reading: $F^i \in \mathrm{ob}\,\underline{L}(S))$
Corrected reading: $F^i \in \mathrm{ob}\,\underline{L}(S)$
Source: check/findings/findings_171_175.json#14; fixes/MISPRINTS_6.json#155. The category expression L(S) has one opening parenthesis and is already closed. Current removes the unmatched second close after this expression; diplomatic retains it. The same component F^i and category membership remain.

[130] EN-VAR-123 Source folio 177. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=191; scan/SGA6_witness_300dpi.djvu#page=188
Printed reading: (4.8)
Corrected reading: (I 4.8)
Source: check/findings/findings_176_180.json#8; fixes/MISPRINTS_6.json#165. ExposéI Example4.8 on122 describes the perfect amplitude for this derived category of Modules using local direct factors of finite free Modules. The parallel statement II2.2.9 on175 explicitly cites I4.8. Current supplies the omitted exposé identifier; diplomatic retains4.8.

[131] EN-VAR-124 Source folio 177. Scan/witness:
scan/SGA6_scan_360dpi.pdf#page=191; scan/SGA6_witness_300dpi.djvu#page=188
Printed reading: (4.9)
Corrected reading: (I 4.9)
Source: check/findings/findings_176_180.json#9; fixes/MISPRINTS_6.json#166. ExposéI Notation4.9 on122 names D(S)_parf for the full subcategory of perfect objects used here. The parallel statement on175 has the explicit I4.9 locator. Current supplies the omitted exposé identifier without changing the indicated subcategory.

[132] EN-VAR-126 Source folio 178. Scan/witness:
scan/SGA6_scan_360dpi.pdf#page=192; scan/SGA6_witness_300dpi.djvu#page=189
Printed reading: Assertion c) follows from (1.2.1 b),c),d))
Corrected reading: Assertion c) follows from (1.2 b),c),d))
Source: ASTRA-II-REF-178-POST05;
review/expose_ii_amendments/reference_178_identification.json. Both scans print the extra .1. The actual Proposition1.2(b,c,d) on161–163 supplies the bounded representative, canonical fully faithful functor and actual quasi-isomorphism used here. The printed C_0=Lib wording is retained. The recorded specialization first takes P=Loclib and identifies C_(1S) with P by the identity on modules and maps. For a degree-i term choose a finite-free complement Q and add the contractible disk Q→Q in degrees i−1,i. Its projection pi and inclusion j satisfy pi j=id and dh+hd=id−j pi, with h^i=id_Q. Descending through i=n,...,m+1 gives the stated free higher terms, and the composite F→G→E is the required quasi-isomorphism. This exact construction supports the locator repair without silently identifying the two printed categories.

[133] EN-VAR-125 Source folio 178. Scan/witness:
scan/SGA6_scan_360dpi.pdf#page=192; scan/SGA6_witness_300dpi.djvu#page=189
Printed reading: free Modules (resp.) direct factors of free Modules) of finite type
Corrected reading: free Modules (resp. direct factors of free Modules) of finite type
Source: check/findings/findings_176_180.json#12; fixes/MISPRINTS_6.json#168. The printed alternative closes immediately after resp. and again after the phrase giving the alternative. Current removes only the premature first close. The functor Gamma_S and both specified kinds of Modules remain unchanged.

[134] EN-VAR-127 Source folio 181. Scan/witness:
scan/SGA6_scan_360dpi.pdf#page=195; scan/SGA6_witness_300dpi.djvu#page=192
Printed reading: $\underline{\underline{R}}\Gamma_S = \mathrm{D}^b(S)_{\mathrm{coh}} \longrightarrow \mathrm{D}^+(A)$
Corrected reading: $\underline{\underline{R}}\Gamma_S : \mathrm{D}^b(S)_{\mathrm{coh}} \longrightarrow \mathrm{D}^+(A)$
Source: check/findings/findings_181_185.json#5; fixes/REPLAY_MANUAL_6.json#4. The printed equals sign separates a functor name from its domain. The preceding sentence defines R Gamma_S:D^+(S)→D^+(A); compose it with the inclusion D^b(S)_coh→D^+(S) to obtain the displayed restriction. Current uses a colon for that same functor. Both readings retain the two directly observed horizontal rules below R.

[135] EN-VAR-128 Source folio 182. Scan/witness:
scan/SGA6_scan_360dpi.pdf#page=196; scan/SGA6_witness_300dpi.djvu#page=193
Printed reading: $i \in [m, n]$
Corrected reading: $i \notin [m, n]$
Source: check/findings/findings_181_185.json#10; fixes/MISPRINTS_6.json#171. The proof chooses E^i=0 for i>n on 182 and then truncates below m on 183. The map to the truncation preserves cohomology because the stated perfect amplitude gives vanishing below m. The resulting representative therefore vanishes outside [m,n]. Current restores notin; diplomatic preserves the printed membership.

[136] EN-VAR-129 Source folio 183. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=197; scan/SGA6_witness_300dpi.djvu#page=194
Printed reading: $E^m \in \mathrm{ob}\,\mathrm{Loclib}(S))$
Corrected reading: $E^m \in \mathrm{ob}\,\mathrm{Loclib}(S)$
Source: check/findings/findings_181_185.json#11; fixes/MISPRINTS_6.json#172. The category expression Loclib(S) already has its matching close. Current removes the additional unmatched parenthesis after it; diplomatic retains that extra close. The object and membership are unchanged.

[137] EN-VAR-130 Source folio 183. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=197; scan/SGA6_witness_300dpi.djvu#page=194
Printed reading: $\mathrm{K}^b(\mathrm{Loclib}(S)) \to \mathrm{D}(S)$
Corrected reading: $\mathrm{K}^b(\mathrm{Loclib}(S)) \to \mathrm{D}(A)$
Source: check/findings/findings_181_185.json#12. The preceding functor j:K^b(Loclib(S))→D(S) is retained. The second functor is explicitly induced by Gamma_S: degreewise sections define T:K^b(Loclib(S))→K^b(Mod(A))→D(A). Theorem B makes each locally free coherent term acyclic, so the canonical comparison from T to R Gamma_S composed with j is an isomorphism. Current therefore restores the A-module target of this second functor; diplomatic retains S.

[138] EN-VAR-131 Source folio 183. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=197; scan/SGA6_witness_300dpi.djvu#page=194
Printed reading: $\mathrm{D}(S)_{\mathrm{parf}}$
Corrected reading: $\mathrm{D}(A)_{\mathrm{parf}}$
Source: check/findings/findings_181_185.json#13. This last image clause belongs to T=Gamma_S on bounded locally free complexes. Under the canonical comparison from T to R Gamma_S composed with j, Proposition 2.4.5(c) on 182 identifies its image with D(A)_parf. The earlier D(S)_parf is the image of j and remains unchanged. Current restores A only in this final image clause.

[139] EN-VAR-132 Source folio 184. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=198; scan/SGA6_witness_300dpi.djvu#page=195
Printed reading: J. Frish [5]
Corrected reading: J. Frisch [5]
Source: check/findings/findings_181_185.json#16; directly identified bibliography item[5] on220. The author note expressly cites item[5], whose directly inspected bibliography entry spells the surname Frisch and identifies the same analytic-functions paper. Current supplies the missing c in that attribution; diplomatic preserves Frish. The complete Grothendieck attribution and independent-discovery statement remain in both forms.

[140] EN-VAR-133 Source folio 185. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=199; scan/SGA6_witness_300dpi.djvu#page=196
Printed reading: that S is the analytic spectrum
Corrected reading: that $\widetilde{S}$ is the analytic spectrum
Source: check/findings/findings_181_185.json#27. The previous sentence introduces the normalization S-tilde. Its finite normalization algebra has relative analytic spectrum equal to the domain S-tilde of the explicitly specified projection p:S-tilde→S. Current restores the tilde on this domain name; diplomatic preserves the printed S.

[141] EN-VAR-134 Source folio 185. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=199; scan/SGA6_witness_300dpi.djvu#page=196
Printed reading: (10 II §3 Prop.8)
Corrected reading: ([10] II §3 Prop.8)
Source: check/findings/findings_181_185.json#28. The surrounding citations identify the same work as[10] in IV section3, IV section4 and II section3. This occurrence names Proposition8 of the last section. Current restores its bibliography-number brackets; the specified work, section and proposition are unchanged.

[142] EN-VAR-135 Source folio 185. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=199; scan/SGA6_witness_300dpi.djvu#page=196
Printed reading: $E_n \cdot \widetilde{O}_S|U = 0$
Corrected reading: $E_n \cdot \widetilde{O}_S|V = 0$
Source: check/findings/findings_181_185.json#31. The introduced open neighbourhood is V in S-tilde. The ideal carrying the outer tilde is a sheaf on S-tilde, and its restriction is inverse image along V→S-tilde. The earlier U is an open neighbourhood on S; the specified projection p:S-tilde→S relates the two spaces. Current names V in this restriction, while diplomatic preserves U.

[143] EN-VAR-136 Source folio 186. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=200; scan/SGA6_witness_300dpi.djvu#page=197
Printed reading: Thus Y has codimension
Corrected reading: Thus X has codimension
Source: fixes/RECONCILE_CODEX_6.json#252; fixes/FIXLOG_6_CURRENT.jsonl#359. The specified closed immersion f:X→Y is cut out by I. At the chosen point I_x is nonzero in the domain R=O_(Y,x). Every prime P containing I_x is nonzero, so the strict chain from(0) to P has positive length. This computes the positive codimension of the closed subspace X=V(I) in Y. Current restores X as the subject; diplomatic preserves Y.

[144] EN-VAR-137 Source folio 189. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=203; scan/SGA6_witness_300dpi.djvu#page=200
Printed reading: (deduced from (3.3 b) (ii))) is an isomorphism
Corrected reading: (deduced from (3.3 b) (ii)) is an isomorphism
Source: ASTRA-II-PAREN-189. The complete parenthetical has three opening and four closing parentheses in the print. Current removes the final unmatched close; diplomatic retains it. The reference and adjunction arrow E→RQ(E) are unchanged.

[145] EN-VAR-138 Source folio 189. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=203; scan/SGA6_witness_300dpi.djvu#page=200
Printed reading: (deduced from (3.3 b)(ii))) is an isomorphism
Corrected reading: (deduced from (3.3 b)(ii)) is an isomorphism
Source: ASTRA-II-PAREN-189. The second complete parenthetical likewise has one unmatched final close. Current deletes that delimiter only, preserving the reference and adjunction arrow RQ(E)→E; diplomatic keeps every printed delimiter.

[146] EN-VAR-139 Source folio 196. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=210; scan/SGA6_witness_300dpi.djvu#page=207
Printed reading: $M = \coprod_{n\ 0} B/(\underline{f}^n)$
Corrected reading: $M = \coprod_{n>0} B/(\underline{f}^n)$
Source: check/findings/findings_196_200.json#3; fixes/MISPRINTS_6.json#185. The printed relation slot between n and0 is blank. Current supplies the positive-index sum. Its canonical inclusion into the sum over n≥0 and the projection back are inverse: the n=0 term is B/(1,1)=0 for f=(x,y). This identifies the two specified module constructions without changing any summand of positive index. Diplomatic retains the blank.

[147] EN-VAR-140 Source folio 196. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=210; scan/SGA6_witness_300dpi.djvu#page=207
Printed reading: $H_2^B(\underline{f}^n, A) \simeq H_2^B(\underline{f}^n, M)$,
Corrected reading: $H_2^B(\underline{f}^n, A) \simeq H_2^B(\underline{f}^n, M)$.
Source: check/findings/findings_196_200.json#7; fixes/MISPRINTS_6.json#186. The print ends this display with a comma before the next independent capitalized sentence. Current uses a sentence stop. Every Koszul term and the isomorphism sign remain identical; diplomatic preserves the comma.

[148] EN-VAR-141 Source folio 196. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=210; scan/SGA6_witness_300dpi.djvu#page=207
Printed reading: Finally, we have (EGA II 1.1.3.3)
Corrected reading: Finally, we have (EGA III 1.1.3.3)
Source: check/findings/findings_196_200.json#8; fixes/MISPRINTS_6.json#187. The reviewed primary EGA III text, Part1 page83 formula1.1.3.3, gives the Koszul duality identification H^i(f,M) with H_(r-i)(f,M). For r=2 and i=0, the map sends the top exterior-basis vector tensored with m to m in the common annihilator of x^n and y^n. This is precisely the displayed H_2 to H^0 map. Current repairs only the Roman chapter numeral; diplomatic keeps II. The independent primary-text verification is recorded by the French reviewer, without a new EGA image-read claim.

[149] EN-VAR-142 Source folio 196. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=210; scan/SGA6_witness_300dpi.djvu#page=207
Printed reading: $H_2^A(\underline{f}^n, A \simeq {}_{(\underline{f}^n)}M$.
Corrected reading: $H_2^A(\underline{f}^n, A) \simeq {}_{(\underline{f}^n)}M$.
Source: check/findings/findings_196_200.json#11; fixes/MISPRINTS_6.json#188; fixes/RECONCILE_CODEX_6.json#254. Current closes the existing argument list of H_2^A(f^n,A), the same Koszul module in the first display and the transition maps. No term or map is added. Diplomatic retains the omitted closing parenthesis in(*).

[150] EN-VAR-143 Source folio 201. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=215; scan/SGA6_witness_300dpi.djvu#page=212
Printed reading: $P^n_{X/S}(E) = P^n_{X/S} \oplus_{\underline{O}_X} E$
Corrected reading: $P^n_{X/S}(E) = P^n_{X/S} \otimes_{\underline{O}_X} E$
Source: check/findings/findings_201_205.json#4. The printed sign is a circled plus, while the immediately following words specify the tensor product using the second O_X action. The balanced construction imposes (p a) tensor e = p tensor (a e), and its canonical map sends e to1 tensor e. Composition with an O_X-linear map from this construction is the factorization in1.2.2.1. Current retains the recorded tensor-product repair; diplomatic preserves the actual plus.

[151] EN-VAR-144 Source folio 201. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=215; scan/SGA6_witness_300dpi.djvu#page=212
Printed reading: is an $f^{-1}(\underline{O}_X)$-linear homomorphism
Corrected reading: is an $f^{-1}(\underline{O}_S)$-linear homomorphism
Source: check/findings/findings_201_205.json#3. For the specified f:X→S, the structure map is f-sharp:f^(-1)O_S→O_X. The two O_X actions on principal parts agree on that image because both projections are over S. Hence d^n(E)(a e)=u_2(a)(1 tensor e)=u_1(a)(1 tensor e) for a from f^(-1)O_S, and an O_X-linear factor h gives D(a e)=a D(e). This exact factorization identifies the source of the required base-linearity. Current uses O_S; diplomatic preserves the printed O_X in this later definition only.

[152] EN-VAR-145 Source folio 202. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=216; scan/SGA6_witness_300dpi.djvu#page=213
Printed reading: the identifications (1.2.2.1) and (1.2.4.1)
Corrected reading: the identifications (1.2.2.1) and (1.2.3.1)
Source: check/findings/findings_201_205.json#8. The explicit composition h→h composed with d^n(E) gives the real identification1.2.2.1 and complex identification1.2.3.1. The paragraph applies Hom to the exact sequence1.2.4.1 and then uses those two identifications to obtain1.2.4.2. Current repairs the second identification locator to1.2.3.1; diplomatic keeps the printed1.2.4.1. Every map and sequence is retained.

[153] EN-VAR-146 Source folio 203. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=217; scan/SGA6_witness_300dpi.djvu#page=214
Printed reading: (considered as a monomorphism of ringed spaces) cf. (1.1)).
Corrected reading: (considered as a monomorphism of ringed spaces) cf. (1.1).
Source: check/findings/findings_201_205.json#12. The parenthesis opening the considered-clause is already closed after spaces, and the reference opening is closed after1.1. Current deletes only the remaining unmatched final close. Diplomatic retains it, with all clause words, the reference and the sentence stop unchanged.

[154] EN-VAR-147 Source folio 211. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=225; scan/SGA6_witness_300dpi.djvu#page=222
Printed reading: to that of $\underline{O}_S$ rep.
Corrected reading: to that of $\underline{O}_S$ resp.
Source: check/findings/findings_211_215.json#2. The printed shared abbreviation rep. lacks the s in resp. The real bundle is assigned its O_S section sheaf; the complex bundle is assigned its O_S tensor C section sheaf. Current restores the abbreviation linking these two stated assignments. Diplomatic preserves rep. at this occurrence; all functor data are unchanged.

[155] EN-VAR-148 Source folio 213. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=227; scan/SGA6_witness_300dpi.djvu#page=224
Printed reading: results of (loc.cit):
Corrected reading: results of (loc.cit.):
Source: check/findings/findings_211_215.json#9. Current supplies the missing final abbreviation stop in the shared Latin locator loc.cit. Diplomatic retains the printed loc.cit without that stop. The surrounding reference, two assertions and their separate arguments remain fully translated.

[156] EN-VAR-149 Source folio 214. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=228; scan/SGA6_witness_300dpi.djvu#page=225
Printed reading: $u_r : \underline{C}'(S, C_2^r(V,E)) \longrightarrow \underline{C}'(S, C_2^r(V,F))$.
Corrected reading: $u_r : \underline{C}'(S, C_2^r(V,E)) \longrightarrow \underline{C}'(S, C_2^{r-n}(V,F))$.
Source: check/findings/findings_211_215.json#10; fixes/MISPRINTS_6.json#184. The defining family is T(s)=phi_r(u(s)) in L(H_E,H_F), with H_E=C^r_2(V,E) and H_F=C^(r-n)_2(V,F), as displayed immediately above. On a relatively compact open W, its induced presheaf map sends a in C(bar W,H_E) to the continuous map s→T(s)a(s) in C(bar W,H_F). Continuity follows from continuous evaluation L(H_E,H_F) times H_E→H_F. This explicit map gives target C-prime(S,C^(r-n)_2(V,F)); the final bundle map also prints that exponent. Current retains r-n and diplomatic preserves the printed r only in this presheaf target.

[157] EN-VAR-150 Source folio 219. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=233; scan/SGA6_witness_300dpi.djvu#page=230
Printed reading: the relations between the groups $[S, I(S)]$ and
Corrected reading: the relations between the groups $[S, I(H)]$ and
Source: check/findings/findings_216_220.json#23. The paragraph defines I(H) inside L(H,H) and the homotopy group of maps S→I(H). Its element is the class of a map with that target, and its homotopies are maps S times[0,1]→I(H). The later comparison therefore concerns that same group [S,I(H)]. Current restores H in this single later occurrence; diplomatic preserves the printed S. No relation to K-bullet(S) is added to the historical question.

[158] EN-VAR-151 Source folio 220. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=234; scan/SGA6_witness_300dpi.djvu#page=231
Printed reading: Fonctions analyliques sur un ensemble semi-analytique
Corrected reading: Fonctions analytiques sur un ensemble semi-analytique
Source: check/findings/findings_216_220.json#29. The bibliography title is retained in French in the English edition. Diplomatic therefore preserves the actually printed analyliques, with l, and current retains the recorded spelling correction analytiques, with t. The later semi-analytique, author, publication data, page range and punctuation remain unchanged.


[159] EN-VAR-152 Source folio 223. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=237; scan/SGA6_witness_300dpi.djvu#page=234
Printed reading: $H^n(i_*(E''$
Corrected reading: $H^n(i_*(E''))$
Source: check/findings/findings_221_225.json#19. The first expression applies H^n to the complete object i_*(E-double-prime), but the printed form closes only the inner argument. The next expression applies the same composite H^n composed with i_* and prints both closers. Current adds only the missing outside parenthesis; diplomatic retains the exact first missing close. The degree n remains unchanged.

[160] EN-VAR-153 Source folio 223. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=237; scan/SGA6_witness_300dpi.djvu#page=234
Printed reading: independent of the hypothesis on $\bar{i}_*(\underline{O}_X)$
Corrected reading: independent of the hypothesis on $i_*(\underline{O}_X)$
Source: check/findings/findings_221_225.json#21. The source defines the single closed immersion i:X→Y. At x, the stalk of i_*M at i(x) is M_x, and the surjection O_(Y,i(x))→O_(X,x) carries the displayed finite generators to the same generating elements over the quotient ring. This is exactly the direct image i_* from the lemma hypothesis. Current keeps that same unbarred i; diplomatic retains the observed extra bar.

[161] EN-VAR-154 Source folio 224. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=238; scan/SGA6_witness_300dpi.djvu#page=235
Printed reading: the Kozsul complex
Corrected reading: the Koszul complex
Source: check/findings/findings_221_225.json#23; fixes/FIXLOG_6_CURRENT.jsonl#372.
The displayed regular-immersion construction uses the complex with exterior-power terms and differential sending the basis element epsilon_j to the corresponding ideal generator f_j, the Koszul resolution referenced in VII1.4. Diplomatic retains the transposed proper-name spelling Kozsul; current retains Koszul.


[162] EN-VAR-155 Source folio 225. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=239; scan/SGA6_witness_300dpi.djvu#page=236
Printed reading: i''_* n-pseudo-coherent
Corrected reading: $i''_*(E)$ n-pseudo-coherent
Source: check/findings/findings_221_225.json#28; fixes/FIXLOG_6_CURRENT.jsonl#373.
The lemma fixes E in D(X) and applies the two closed-immersion direct-image functors to that object. Its proof uses the diagonal X→X-prime times_Y X-double-prime and compares those direct images; the right-hand object is therefore i-double-prime_*(E). Current restores only the omitted argument(E); diplomatic keeps the printed bare functor symbol.

[163] EN-VAR-156 Source folio 226. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=240; scan/SGA6_witness_300dpi.djvu#page=237
Printed reading:



(a')

Corrected reading:



(a')

Source: check/findings/findings_226_230.json#3; fixes/FIXLOG_6_CURRENT.jsonl#374.
The top edge has endpoints X-prime→Y-prime. The following text names that immersion i-prime and uses its direct-image functor i-prime_*p-prime*. The next-page cube identifies the same endpoints and edge as i-prime. Thus the endpoint-preserving map from the first diagram to the corresponding cube face identifies its label with i-prime. Current keeps that single prime; diplomatic preserves the printed double prime. The distinct second factorization retains i-double-prime.

[164] EN-VAR-157 Source folio 230. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=244; scan/SGA6_witness_300dpi.djvu#page=241
Printed reading: $\underline{\mathrm{Tor}}_i^S(\underline{O}_X, O_Y)$
Corrected reading: $\underline{\mathrm{Tor}}_i^S(\underline{O}_X, \underline{O}_Y)$
Source: check/findings/findings_226_230.json#14. The source names X,Y,S with affine rings B,C,A and the tensor comparison B tensor-derived_A C→B tensor_A C. The two sheaf arguments restrict to B and C by the structure maps A→B and A→C. Thus the Y argument is the same structure sheaf of Y used in that calculation. Diplomatic retains its isolated missing rule; current retains the established rule on that same O_Y without changing the sheaf, indices or tensor map.

[165] EN-VAR-158 Source folio 234. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=248; scan/SGA6_witness_300dpi.djvu#page=245
Printed reading:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow^{f} & \downarrow{\scriptstyle g} \\ & & Z \end{array}$$

Corrected reading:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow^{h} & \downarrow{\scriptstyle g} \\ & & Z \end{array}$$

Source: check/findings/findings_231_235.json#18; fixes/FIXLOG_6_CURRENT.jsonl#375. The diagonal has endpoints X→Z and is the composite h=g composed with f. Its direct-image functor is h_*=g_*f_*, exactly the functor used in the subsequent relative condition and proof. Current retains h on this edge; diplomatic retains the printed f. The horizontal f:X→Y and vertical g:Y→Z are unchanged.

[166] EN-VAR-159 Source folio 235. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=249; scan/SGA6_witness_300dpi.djvu#page=246
Printed reading: the isomorphism (*) in the proof of (1.8)
Corrected reading: the isomorphism in (1.7)
Source: check/findings/findings_231_235.json#28. Both full scans print l'isomorphisme(*) de la preuve de(1.8). The inspected231–232 text has no starred formula in either proof. The displayed isomorphism is exactly the Lemma1.7 map at S=Y, f_1=f:X→Y and f_2=Id_Y, using X×_Y Y≅X. The proof has already reduced f to a closed immersion. At y=f(x), put R=O_{Y,y}, B=O_{X,x}; for a flat resolution P of F_y the chain maps e⊗p↦e⊗(1⊗p) and e⊗(b⊗p)↦be⊗p are inverse between E_x⊗_R P and E_x⊗_B(B⊗_R P). They realize the stalk of the canonical adjunction projection map. Off X both stalks are zero, since f is closed. This proves the displayed map in the actual proof scope, including complexes not assumed quasi-coherent. Current cites l'isomorphisme de(1.7), removing the nonexistent star and proof locator; diplomatic retains the complete printed phrase. This is an explicit editorial repair, with no claim about an unseen author draft.

[167] EN-VAR-160 Source folio 235. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=249; scan/SGA6_witness_300dpi.djvu#page=246
Printed reading: $f_*(E)$
Corrected reading: $h_*(E)$
Source: check/findings/findings_231_235.json#27; fixes/FIXLOG_6_CURRENT.jsonl#376. Proposition1.11(b) assumes pseudo-coherence relative to h. After the stated reduction to closed immersions, its hypothesis is pseudo-coherence of h_*(E)=g_*f_*(E). With E_0=f_*(E), the following replacement X=Y and f=Id makes this precisely g_*(E_0), as used later in the proof. The conclusion remains pseudo-coherence of E_0. Only this first hypothesis in(b) changes from printed f_* to current h_*; all other f_* occurrences are preserved.

[168] EN-VAR-161 Source folio 238. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=252; scan/SGA6_witness_300dpi.djvu#page=249
Printed reading: $i > a + r$
Corrected reading: $q > a + r$
Source: check/findings/findings_236_240.json#13; fixes/FIXLOG_6_CURRENT.jsonl#377. The higher direct image is R^qf_*(E). In the stated hypercohomology computation, q=p+j is the total degree, with j≤a from the complex and p≤r from the projective-space cover, so q≤a+r for potentially nonzero terms. The stated vanishing condition is therefore q>a+r for that same R^q. Current changes only the condition index to q; diplomatic retains the printed i while preserving R^q.

[169] EN-VAR-162 Source folio 241. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=255; scan/SGA6_witness_300dpi.djvu#page=252
Printed reading: (2.2), (2.5) and (1.12)
Corrected reading: (2.2), (2.4) and (1.12)
Source: fixes/FIXLOG_6_CURRENT.jsonl#379. The printed proof of2.5 cites2.5 itself. Corollary2.4 supplies the bounded derived functor under precisely the proper/projective or noetherian alternatives used here. Take the base S=Y locally on Y: the structure map of Y is id_Y, and f is pseudo-coherent. By1.12 the absolute pseudo-coherence of E is its relative pseudo-coherence over Y; 2.2 transports that property through Rf_*, and2.4 gives the bounded restriction when Y is quasi-compact. On Y, relative pseudo-coherence over id_Y is the absolute property by the identity inverse-image map. Thus the middle citation is2.4, while2.2 and1.12 remain unchanged. Diplomatic retains the printed2.5.

[170] EN-VAR-163 Source folio 242. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=256; scan/SGA6_witness_300dpi.djvu#page=253
Printed reading: (SGA 4 XVII IV 6.4)
Corrected reading: (SGA 4 IV 6.4)
Source: fixes/FIXLOG_6_CURRENT.jsonl#380. Both SGA6 scans print SGA4 XVII IV6.4. The personally viewed SGA4 tome1 re-edition page230, PDF236, has the ExposéIV header and Definition6.4.1: a conservative family of point fibre functors defines having enough points. This is the exact definition cited in3.3. For a module complex, the fibre functor sends H^i(E) to H^i(E_x), because the inverse-image fibre functor is exact; conservativity tests the isomorphism0→H^i(E) on these fibres. This identifies the cited criterion without changing any hypothesis or mathematical symbol. Retain IV6.4 in current and the full XVII IV6.4 diplomatically; source record external_sga4_242.json records the checked reference.

[171] EN-VAR-164 Source folio 248. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=262; scan/SGA6_witness_300dpi.djvu#page=259
Printed reading: Suppose E (resp. M) is of finite flat amplitude rel. to f (resp. finite flat amplitude).
Corrected reading: Suppose E (resp. M) is of finite flat amplitude rel. to S (resp. finite flat amplitude).
Source: check/findings/findings_246_250.json#3. The primary and native pages print rel. à f in the proof, while the immediately preceding statement247 explicitly assumes finite amplitude relative to S. Write g:Y→S and h=g∘f. The tensor used in the proof is E tensor^L_{h^{-1}O_S}h^{-1}M. The structural map is the composite h^{-1}O_S=f^{-1}g^{-1}O_S→f^{-1}O_Y→O_X; relative amplitude for f instead uses f^{-1}O_Y. These are the exact scalar maps involved. The difference is witnessed by A=k[ε]/(ε^2), S=SpecA, X=Y=Speck, f=id, E=k and M=k: E is free over O_Y, but the free A-resolution ...→A $-\varepsilon$→A $-\varepsilon$→A→k has kernel=image=(ε) and pulls back to zero differentials on k in every negative degree. Thus relative amplitude for f alone does not provide the asserted bounded tensor. The statement's S hypothesis supplies the bounded flat resolution over h^{-1}O_S used here; its tensor with a bounded representative of M has total degrees in the sums of their finite degree intervals. Correct only this recalled base letter f→S in current. The statement itself, the adjunction counit, both canonical arrows and the alternative finite-amplitude hypothesis on M remain unchanged.

[172] EN-VAR-165 Source folio 251. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=265; scan/SGA6_witness_300dpi.djvu#page=262
Printed reading: is a perfect morphism
Corrected reading: is a perfect morphism.
Source: check/findings/findings_251_255.json#2. Both complete scans leave the completed sentence without a final stop, before the separate Notation4.2 paragraph. Retain that absence diplomatically and the current terminal stop.

[173] EN-VAR-166 Source folio 252. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=266; scan/SGA6_witness_300dpi.djvu#page=263
Printed reading: $g_\alpha^*(F) \simeq F_\alpha$
Corrected reading: $g_\alpha^*(F_\alpha) \simeq F_\alpha$
Source: check/findings/findings_251_255.json#6. Both scans print g_alpha*(F) in the second comparison. The preceding definition binds F_alpha on X_alpha and the first comparison identifies g_alpha*F_alpha with F on X. On the affine charts Y_alpha=SpecA_alpha, Y=SpecA, write B_alpha=A_alpha[t1,...,tr] and B=A[t1,...,tr]=B_alpha tensor_Aalpha A. For the module N representing F_alpha, the pullback is B tensor_Balpha N. The explicit map (b tensor a) tensor n↦a tensor bn identifies it with A tensor_Aalpha N, with inverse a tensor n↦(1 tensor a) tensor n. The given A_alpha-flatness of N identifies the ordinary tensor with its derived tensor. Thus both sides of the printed comparison identify g_alpha*F_alpha, already identified with F, and the corrected argument is F_alpha. Keep the printed F diplomatically and retain the headless comparison sign in both layers.

[174] EN-VAR-167 Source folio 253. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=267; scan/SGA6_witness_300dpi.djvu#page=264
Printed reading: $\mathrm{parf.amp}(i_*(E) \subset [a,b] \Longrightarrow \mathrm{parf.amp}_f(E) \subset [a,b]$
Corrected reading: $\mathrm{parf.amp}(i_*(E)) \subset [a,b] \Longrightarrow \mathrm{parf.amp}_f(E) \subset [a,b]$
Source: check/findings/findings_251_255.json#8. Both scans have two opening parentheses in parfamp(i_*(E) but only the inner closing parenthesis. The outer argument is i_*(E), exactly as the second implication on the same page writes it. Retain the one-close print diplomatically and the two-close expression in current.

[175] EN-VAR-168 Source folio 254. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=268; scan/SGA6_witness_300dpi.djvu#page=265
Printed reading: It also implies. by 4.3),
Corrected reading: It also implies (by 4.3),
Source: check/findings/findings_251_255.json#10. The primary and native254 pages show a stop after aussi, a lower-case d in d'après, and a closing parenthesis after4.3 with no opening parenthesis. The parenthetical citation concerns Proposition4.3's displayed stalk-amplitude condition. Replace the stray stop by the missing opening parenthesis in current, preserving the existing close and comma; diplomatic retains the exact printed marks.

[176] EN-VAR-169 Source folio 255. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=269; scan/SGA6_witness_300dpi.djvu#page=266
Printed reading: is, in a neighborhood of s, acyclic outside
Corrected reading: is, in a neighborhood of x, acyclic outside
Source: check/findings/findings_251_255.json#15. Both scans print s. The already defined map h sends x∈X to s∈S; E and the truncation morphism t are complexes on X. Condition4.6(i) is stated on an open neighbourhood U of x, and the reduction explicitly restricts along its open immersion j:U→X. The restriction j*E→τ_[a,b](j*E) is the truncation used in the proof: its stalks on U are the same truncations of E, and its being a quasi-isomorphism follows from the vanishing specified on U. This identifies the recalled neighbourhood as the neighbourhood of x from(i). Retain current x and diplomatic s; neither h nor any fibre point is renamed.

[177] EN-VAR-170 Source folio 255. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=269; scan/SGA6_witness_300dpi.djvu#page=266
Printed reading: Thus, after localizing,. we may suppose
Corrected reading: Thus, after localizing, we may suppose
Source: check/findings/findings_251_255.json#18. Both scans show the comma followed by a second typed stop. The continuation on peut supposer belongs to the same sentence. Retain the pair diplomatically and the current single comma. The partial overstruck fragments above this line are not transcribed as a new expression.

[178] EN-VAR-171 Source folio 255. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=269; scan/SGA6_witness_300dpi.djvu#page=266
Printed reading: arrow (∗)
Corrected reading: arrow (t)
Source: check/findings/findings_251_255.json#17. The fibre comparison is visibly unlabelled in both scans; later the print says la flèche(*). The map used in that final paragraph is the original truncation t:E→T on X, where T^a=E^a/B^a, T^j=E^j for a<j≤b, and T is zero elsewhere. Its component in degree a is the quotient map, its higher components are identities, and its lower components are zero. The preceding assertion makes t a quasi-isomorphism. Tensoring t with an arbitrary h^{-1}O_S-module therefore identifies the cohomology used to test toramp_h(E) with that of T. The terms T^j for j>a are h-flat because they are locally free over O_X and h is flat. LemmaI5.1.1, with m=a,n=b,F=T and base h^{-1}O_S, then gives precisely the h-flatness of T^a claimed at the start of256. The intervening fibre comparison has objects over the residue fibre and supplies no such map E→T on X. Therefore repair the later reference(*) to(t), and remove the previously inserted(*) label from the headless fibre comparison. The printed t) and i*(t) labels themselves remain unchanged.

[179] EN-VAR-172 Source folio 257. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=271; scan/SGA6_witness_300dpi.djvu#page=268
Printed reading: Under the hypotheses of (4.6)
Corrected reading: Under the hypotheses of (4.7)
Source: check/findings/findings_256_260.json#3. The print cites4.6. Proposition4.7(b) is the statement whose data are the two f_i and their product f=f1×_Sf2. Substitute E_i=O_Xi and[a_i,b_i]=[−n_i,0]. Their external tensor identifies with O_X by the multiplication map O_X tensor_OX O_X→O_X, u tensor v↦uv, whose inverse sends w to w tensor1. The displayed summed interval is[−n1−n2,0], exactly the dimension bound stated in4.7.1. The Tor-independence assumptions are those of4.7. This identifies the exact cited assertion(b), independently of the separately recorded failure of4.7(a). Correct only the locator4.6→4.7 in current.

[180] EN-VAR-173 Source folio 259. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=273; scan/SGA6_witness_300dpi.djvu#page=270
Printed reading: Under the hypotheses of (4.8)
Corrected reading: Under the hypotheses of (4.9)
Source: check/findings/findings_256_260.json#14. The print cites4.8, while the author's next sentence specifies the substitution E=O_X. In4.9, the hypothesis that f is perfect makes O_X perfect relative to f. The complex RHom_OX(O_X,f!F) identifies with f!F by evaluation at1, with inverse m↦(a↦am); O_X is locally free of rank1, so this is also the derived identification. The resulting conclusion is exactly that f!F is perfect over S. This uses4.9's compactifiable/noetherian setting and its actual Hom expression. Correct only the locator to4.9.

[181] EN-VAR-174 Source folio 259. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=273; scan/SGA6_witness_300dpi.djvu#page=270
Printed reading: For the first assertion, apply (4.8)
Corrected reading: For the first assertion, apply (4.9)
Source: check/findings/findings_256_260.json#20. The print says appliquer4.8 avecY=S,F=O_S. Under these explicit substitutions in4.9, f!F=f!O_S=K_X and RHom(E,f!F)=D_X(E) by the preceding definition. O_S is the rank-one free complex over id_S, and E has the relative perfection required in4.9. The structural morphism X→S is f itself, so the resulting S-relative perfection is exactly membership in D(f)_parf. This identifies the first assertion of4.9.2 term for term and fixes the locator4.9 without changing any functor or object.

[182] EN-VAR-175 Source folio 261. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=275; scan/SGA6_witness_300dpi.djvu#page=272
Printed reading: strictly perfect (2.1)
Corrected reading: strictly perfect (I 2.1)
Source: ASTRA-III-REFERENCE-261. Both full261 scans print strictly perfect followed by(2.1). Definition I2.1 on95 was re-read directly: its third alternative says that every E^i lies in C0 and E^i=0 for almost all i. In the ringed-topos setting here, C0 consists locally of finite-rank free O-modules, so this is exactly the bounded finite-free complex used for the subsequent dévissage. The identification sends each complex term and each differential to itself; after a chosen local basis E^i≅O^{r_i}, the basis map sends (a_j) to sum a_j e_j with inverse the coordinate maps. III2.1 on236 is the proper-direct-image finiteness conjecture and supplies no such definition. Restore the missing exposé locator I in current, leaving the mathematical assertion and the printed diplomatic locator intact.

[183] EN-VAR-176 Source folio 262. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=276; scan/SGA6_witness_300dpi.djvu#page=273
Printed reading: condition analogous to a) (resp. a'), etc). in which
Corrected reading: condition analogous to a) (resp. a'), etc.) in which
Source: check/findings/findings_261_265.json#8. Both scans print etc). où in the middle of the sentence. The preceding parenthesis on the same page uses etc.) la condition. The abbreviation stop belongs to etc., followed by the closing parenthesis; the relative clause où continues that sentence. Move only the stop in current, preserving the printed order in diplomatic.

[184] EN-VAR-177 Source folio 269. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=283; scan/SGA6_witness_300dpi.djvu#page=280
Printed reading: (resp. $\mathrm{rg}_{k(x)}\, H^1(E_{k(x)})$ is upper semi-continuous
Corrected reading: (resp. $\mathrm{rg}_{k(x)}\, H^1(E_{k(x)})$) is upper semi-continuous
Source: check/findings/findings_266_270.json#5. Both scans open a prose parenthesis before resp. and leave it open before est semi-continue. The mathematical H^1 argument already has its own close. Current supplies only the prose close; diplomatic retains the omission.

[185] EN-VAR-178 Source folio 269. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=283; scan/SGA6_witness_300dpi.djvu#page=280
Printed reading: $\mathrm{rg}(E_x) = \mathrm{rg}(E_{k(X)})$
Corrected reading: $\mathrm{rg}(E_x) = \mathrm{rg}(E_{k(x)})$
Source: check/findings/findings_266_270.json#6. The preceding sentence explicitly defines E_k(x)=E_x tensor^L_OXx k(x), with x the varying point. The rank equality uses precisely that object and the canonical scalar extension O_Xx→k(x). On a finite free representative, its terms O_Xx^{r_i} tensor k(x)→k(x)^{r_i} send (a_j)tensor b to (a_j b), with inverse coordinates e_jtensor b_j. Both ranks are sum(-1)^i r_i. Retain current x in this exact identified fibre notation, preserving capital X in diplomatic.

[186] EN-VAR-180 Source folio 269. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=283; scan/SGA6_witness_300dpi.djvu#page=280
Printed reading: $H^1(E_{k(x)} \simeq H^1(E)_{k(x)}$
Corrected reading: $H^1(E_{k(x)}) \simeq H^1(E)_{k(x)}$
Source: check/findings/findings_266_270.json#9. The note has a close for k(x), but none for H^1(. The headless comparison separates the two cohomology terms. Keep the printed omission in diplomatic and the unique outer close in current.

[187] EN-VAR-179 Source folio 269. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=283; scan/SGA6_witness_300dpi.djvu#page=280
Printed reading: $Z'^p(E))$ are flat
Corrected reading: $Z'^p(E)$ are flat
Source: check/findings/findings_266_270.json#7. Both scans show two closes after the single argument E, with no surrounding prose parenthesis opened. Current deletes the surplus close; diplomatic keeps it.

[188] EN-VAR-181 Source folio 271. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=285; scan/SGA6_witness_300dpi.djvu#page=282
Printed reading: $y \mapsto \mathrm{rg}_{k(x)}\, H^p(E \otimes^{\mathrm{L}}_X k(y))$
Corrected reading: $y \mapsto \mathrm{rg}_{k(y)}\, H^p(E \otimes^{\mathrm{L}}_X k(y))$
Source: check/findings/findings_266_270.json#12. The print on271 reads rank over k(x), with x the generic point fixed on270 and y the varying point. The actual fibre E tensor_X^L k(y) is a k(y)-complex and is exactly the fibre in5.6(i) with its bound variable renamed x→y. This relabelling sends the scalar extension O_X,y→k(y) and every term and differential to themselves. The need for the y field is witnessed by X=Spec(k[t]), E=O_X and cohomological degree0: at y=(t), the fibre is k with t acting by0. An action of the generic field k(t) extending this k[t] action would make t invertible and zero on a nonzero module, which is impossible. Current k(y) therefore restores the stated fibre-rank function. Keep printed k(x) in diplomatic; assign the record to actual271.

[189] EN-VAR-182 Source folio 271. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=285; scan/SGA6_witness_300dpi.djvu#page=282
Printed reading: $H^i(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y)$ are vector spaces of finite rank over
Corrected reading: $H^i(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y))$ are vector spaces of finite rank over
Source: check/findings/findings_271_275.json#4. The source closes k(y), but leaves H^i(open. Current closes the full cohomology argument with one additional parenthesis; diplomatic retains the exact printed omission.

[190] EN-VAR-183 Source folio 271. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=285; scan/SGA6_witness_300dpi.djvu#page=282
Printed reading: $y \longmapsto \sum(-1)^i\, \mathrm{rg}_{k(y)}\, H^i(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y)$
Corrected reading: $y \longmapsto \sum(-1)^i\, \mathrm{rg}_{k(y)}\, H^i(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y))$
Source: check/findings/findings_271_275.json#5. Both complete scans repeat the same omission in the alternating-rank display. Current supplies the one matching close for H^i(.

[191] EN-VAR-184 Source folio 271. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=285; scan/SGA6_witness_300dpi.djvu#page=282
Printed reading: $y \longmapsto \mathrm{rg}_{k(y)}\, H^p(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y))$.
Corrected reading: $y \longmapsto \mathrm{rg}_{k(y)}\, H^p(Rf_*(E) \otimes^{\mathrm{L}}_Y k(y))$
Source: check/findings/findings_271_275.json#6. The source continues the sentence La fonction. . . est constructible after this stop. Preserve the printed stop in diplomatic and its removal in current. No mathematical symbol changes.

[192] EN-VAR-185 Source folio 272. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=286; scan/SGA6_witness_300dpi.djvu#page=283
Printed reading: (cf. (EGA III 6 and (IV 3.1.0)
Corrected reading: (cf. (EGA III 6) and (IV 3.1.0))
Source: check/findings/findings_271_275.json#10. Both scans show three opening parentheses and only the close after IV3.1.0. The clause contains two cited items joined by et, EGA III6 and IV3.1.0; the finite verb que pour f plat continues On peut montrer outside the citation. Close the first individual citation after III6 and close the encompassing cf. parenthesis after the already closed IV3.1.0 citation. This preserves each original opening and all citation tokens in their original order. The token map on EGA,III,6,IV,3.1.0 is the identity; no locator, functor or asserted base-change map is replaced. Current inserts these two closing delimiters, diplomatic retains the print.

[193] EN-VAR-186 Source folio 273. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=287; scan/SGA6_witness_300dpi.djvu#page=284
Printed reading: $Rf_*(E) \otimes^{\mathrm{L}} M) \longrightarrow Rf_*(E \otimes^{\mathrm{L}}_Y M)$
Corrected reading: $Rf_*(E) \otimes^{\mathrm{L}} M \longrightarrow Rf_*(E \otimes^{\mathrm{L}}_Y M)$
Source: check/findings/findings_271_275.json#12. The projection comparison opens and closes Rf_*(E) before the tensor. The extra close after M has no opening mate. Current removes only that surplus close; diplomatic keeps it.

[194] EN-VAR-187 Source folio 281. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=295; scan/SGA6_witness_300dpi.djvu#page=292
Printed reading: $K^\bullet({}_{A'}X'_A)_{A'} = k(D({}_{A'}X'_A)_{\underline{A'\text{-parf}}}$.
Corrected reading: $K^\bullet({}_{A'}X'_A)_{A'} = k(D({}_{A'}X'_A)_{\underline{A'\text{-parf}}})$.
Source: check/findings/findings_281_285.json#8. The native lower crop shows the underlined parf followed by blank space and a full stop, with no final parenthesis. The corrected delimiter closes the already opened k(applied to the named category; it changes no object or map. Preserve the omission in diplomatic and the supplied closing parenthesis in current.

[195] EN-VAR-188 Source folio 283. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=297; scan/SGA6_witness_300dpi.djvu#page=294
Printed reading: in (2.6.2) (resp. (2.6.5))
Corrected reading: in (2.6.5) (resp. (2.6.2))
Source: check/findings/findings_281_285.json#11. The displayed (2.7.1) maps derived K-groups by Lf*. For a flat resolution F of E, A′ tensor_{f^{-1}A} f^{-1}F represents both A′ tensor_A^L E in (2.6.5) and Lf*E; identity on that complex gives the required identification. For a finite locally free/direct-summand object E, the underived tensor A′ tensor_{f^{-1}A} f^{-1}E is f*E, giving the naive product (2.6.2) and (2.7.2). The inclusion comparison sends each such E to its degree-zero complex, which is exactly the vertical map in the printed diagram. Thus current correctly swaps only the two reference locators; diplomatic retains their printed order.

[196] EN-VAR-189 Source folio 284. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=298; scan/SGA6_witness_300dpi.djvu#page=295
Printed reading: $K^\bullet(X)_{\text{naïf}} \to K^\bullet(X)$ the structure
Corrected reading: $K^\bullet(X)_{\text{naïf}} \to K^\bullet(X))$ the structure
Source: check/findings/findings_281_285.json#22. The native witness has no closer for the parenthesis opened before et par suite. The existing current closer after the canonical map closes only that parenthetical aside; the two algebra structures and comparison map are unchanged. Diplomatic correctly leaves the parenthesis open.

[197] EN-VAR-190 Source folio 285. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=299; scan/SGA6_witness_300dpi.djvu#page=296
Printed reading: (V 3.9.1 and VI)
Corrected reading: (V 3.9.1 and VI)
Source: check/findings/findings_281_285.json#26. A visible untyped interval follows VI before the closer. Its content is blank; no missing numeral can be read or inferred from it. Retain the attached diplomatic spacing and current ordinary spacing as an explicit formatting variant with identical reference tokens V 3.9.1 and VI. Do not supply a conjectural reference.

[198] EN-VAR-191 Source folio 285. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=299; scan/SGA6_witness_300dpi.djvu#page=296; scan/SGA6_scan_360dpi.pdf#page=310 (bibliography region only)
Printed reading: Deligne [5]
Corrected reading: Deligne [1]
Source: check/findings/findings_281_285.json#27. Both scans print Deligne [5]. The directly inspected bibliography on 296 identifies [1] as Deligne, P., papiers secrets, followed by [2] Grauert, [3] Grothendieck and [V] Verdier. The reference target is therefore the named Deligne entry [1]. Current changes only the locator, while diplomatic preserves [5]; no new attribution or theorem wording is supplied. The 296 consultation is regional and does not mark that page fully checked.

[199] EN-VAR-192 Source folio 286. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=300; scan/SGA6_witness_300dpi.djvu#page=297
Printed reading: Thanks,to the associativity
Corrected reading: Thanks to the associativity
Source: check/findings/findings_286_290.json#8. A comma with the same descending tail as the surrounding typed commas stands after Grâce in both scans. The prepositional phrase Grâce à has no intervening syntactic boundary; current removes only that stray comma. Diplomatic preserves it.

[200] EN-VAR-199 Source folio 287. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=301; scan/SGA6_witness_300dpi.djvu#page=298; scan/SGA6_scan_360dpi.pdf#page=286; scan/SGA6_scan_360dpi.pdf#page=287
Printed reading: (III 5.9 b))
Corrected reading: (III 5.8 b))
Source: ASTRA-IV-REF-287-POSTFREEZE. Both source loci explicitly print5.9 b). The directly inspected III272 heading is Remarques5.8; its clause b starts273 and contains the analytic canonical projection arrow Rf_*(E) tensor^L M→Rf_*(E tensor_Y^L M), followed by its isomorphism statement and the bibliography, with no intervening5.9. The arrow is the adjoint of Lf^*(Rf_*E tensor^L M)≅Lf^*Rf_*E tensor^L Lf^*M→E tensor^L Lf^*M, the last map being counit_E tensor identity_M. This is exactly the morphism invoked in IV, with no change of functor or arguments. On287, restrict to an open set carrying a bounded finite locally free representative of the perfect M; the bounded coherent E and this M satisfy the displayed5.8(b) domains, and the local projection maps are the restrictions of that same adjunction map. On289, the cited analytic ingredient is the same projection comparison, together with Grauert finiteness and the stated finite tor-dimension. Correct only the locator5.9→5.8 in current; retain the printed locator in diplomatic. Full map and scope details are in projection_reference_identification.json.

[201] EN-VAR-200 Source folio 289. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=303; scan/SGA6_witness_300dpi.djvu#page=300; scan/SGA6_scan_360dpi.pdf#page=286; scan/SGA6_scan_360dpi.pdf#page=287
Printed reading: (III 4.8.1 and 5.9 b))
Corrected reading: (III 4.8.1 and 5.8 b))
Source: ASTRA-IV-REF-289-POSTFREEZE. Both source loci explicitly print5.9 b). The directly inspected III272 heading is Remarques5.8; its clause b starts273 and contains the analytic canonical projection arrow Rf_*(E) tensor^L M→Rf_*(E tensor_Y^L M), followed by its isomorphism statement and the bibliography, with no intervening5.9. The arrow is the adjoint of Lf^*(Rf_*E tensor^L M)≅Lf^*Rf_*E tensor^L Lf^*M→E tensor^L Lf^*M, the last map being counit_E tensor identity_M. This is exactly the morphism invoked in IV, with no change of functor or arguments. On287, restrict to an open set carrying a bounded finite locally free representative of the perfect M; the bounded coherent E and this M satisfy the displayed5.8(b) domains, and the local projection maps are the restrictions of that same adjunction map. On289, the cited analytic ingredient is the same projection comparison, together with Grauert finiteness and the stated finite tor-dimension. Correct only the locator5.9→5.8 in current; retain the printed locator in diplomatic. Full map and scope details are in projection_reference_identification.json.

[202] EN-VAR-193 Source folio 289. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=303; scan/SGA6_witness_300dpi.djvu#page=300
Printed reading: (resp. $K^{\bullet}(Y)$ and
Corrected reading: (resp. $K^{\bullet}(Y)$) and
Source: check/findings/findings_286_290.json#16. Both scans leave the parenthesis opened before resp. unclosed. The first branch allows y in K_•(Y), the alternative allows y in K^•(Y), and x belongs to K^•(X) in both branches. The supplied closer after K^•(Y) therefore delimits exactly the alternative y-domain. It changes neither the pullback maps (2.12.1)/(2.7.1) nor the product and pushforward in (2.12.4). Diplomatic retains the omission and current supplies that single closer.

[203] EN-VAR-195 Source folio 292. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=306; scan/SGA6_witness_300dpi.djvu#page=303
Printed reading: $u : \mathrm{ob}\,\mathrm{Loclib}(X) \longrightarrow \lim K^{\bullet}(X_i)_{\text{naïf}}$
Corrected reading: $u : \mathrm{ob}\,\mathrm{Loclib}(X) \longrightarrow \varinjlim K^{\bullet}(X_i)_{\text{naïf}}$
Source: check/findings/findings_291_295.json#12. The preceding sentence defines u(E) as the image of cl(E_i) in the explicitly rightward limit of (3.2.1.1). Writing G for that same group, its structure map j_i sends cl(E_i) to u(E); the displayed function u:ob Loclib(X)→G has exactly this target. Thus the current rightward limit restores the operator already used in the definition of this same u, while diplomatic preserves the observed absence. No new limit or comparison map is introduced.

[204] EN-VAR-194 Source folio 292. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=306; scan/SGA6_witness_300dpi.djvu#page=303
Printed reading: K
Corrected reading: $K_{\bullet}$
Source: check/findings/findings_291_295.json#5. Both scans show the bare K. The two constructions named by the title are identified in 3.2.1 and 3.2.3: the former sends a vector-bundle class at Xi to its pullback class at X, while the latter sends a coherent-sheaf class at Xi to its flat pullback class in K_•(X). The latter map is explicitly displayed in the directly consulted 293 middle region. The existing current title spells out that second group as K_•; the diplomatic title preserves the missing mark. No identification between the two K groups is asserted.

[205] EN-VAR-196 Source folio 293. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=307; scan/SGA6_witness_300dpi.djvu#page=304
Printed reading:

$$\varinjlim K_\bullet(X_i) \xrightarrow{\sim} K_\bullet(X). \tag{3.2.2.1}$$

Corrected reading:

$$\varinjlim K_\bullet(X_i) \xrightarrow{\sim} K_\bullet(X). \tag{3.2.3.1}$$

Source: check/findings/findings_291_295.json#18. Both displayed maps carry the printed tag (3.2.2.1). Let α be the vector-bundle limit map of (3.2.1.1), and let β:colim K_•(Xi)→K_•(X) be the flat-pullback map in 3.2.3, β(j_i[F_i])=[f_i^*F_i]. The final diagram has right vertical β and left vertical $\alpha\otimes\beta$. At a common index k, both routes send [V_k]⊗[F_k] to [f_k^*V_k⊗f_k^*F_k], by the canonical pullback/tensor isomorphism. This identifies the second tag and both following references with β. The perfect-complex map α_perf of 3.2.2 is related by the inclusion maps c_i:K^•(Xi)→K_•(Xi) and c_X:K^•(X)→K_•(X): β∘colim(c_i)=c_X∘α_perf, since both send a perfect complex E_i to its derived pullback class, and flat pullback computes that class here. The current relabelling (3.2.3.1) therefore follows the exact map used in the diagram; diplomatic preserves the duplicate tag and both printed references.

[206] EN-VAR-197 Source folio 293. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=307; scan/SGA6_witness_300dpi.djvu#page=304
Printed reading: via the isomorphisms (3.2.1.1) and (3.2.2.1) with
Corrected reading: via the isomorphisms (3.2.1.1) and (3.2.3.1) with
Source: check/findings/findings_291_295.json#18. Both displayed maps carry the printed tag (3.2.2.1). Let α be the vector-bundle limit map of (3.2.1.1), and let β:colim K_•(Xi)→K_•(X) be the flat-pullback map in 3.2.3, β(j_i[F_i])=[f_i^*F_i]. The final diagram has right vertical β and left vertical $\alpha\otimes\beta$. At a common index k, both routes send [V_k]⊗[F_k] to [f_k^*V_k⊗f_k^*F_k], by the canonical pullback/tensor isomorphism. This identifies the second tag and both following references with β. The perfect-complex map α_perf of 3.2.2 is related by the inclusion maps c_i:K^•(Xi)→K_•(Xi) and c_X:K^•(X)→K_•(X): β∘colim(c_i)=c_X∘α_perf, since both send a perfect complex E_i to its derived pullback class, and flat pullback computes that class here. The current relabelling (3.2.3.1) therefore follows the exact map used in the diagram; diplomatic preserves the duplicate tag and both printed references.

[207] EN-VAR-198 Source folio 293. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=307; scan/SGA6_witness_300dpi.djvu#page=304
Printed reading: the vertical arrows are defined by (3.2.1.1) and (3.2.2.1)
Corrected reading: the vertical arrows are defined by (3.2.1.1) and (3.2.3.1)
Source: check/findings/findings_291_295.json#18. Both displayed maps carry the printed tag (3.2.2.1). Let α be the vector-bundle limit map of (3.2.1.1), and let β:colim K_•(Xi)→K_•(X) be the flat-pullback map in 3.2.3, β(j_i[F_i])=[f_i^*F_i]. The final diagram has right vertical β and left vertical $\alpha\otimes\beta$. At a common index k, both routes send [V_k]⊗[F_k] to [f_k^*V_k⊗f_k^*F_k], by the canonical pullback/tensor isomorphism. This identifies the second tag and both following references with β. The perfect-complex map α_perf of 3.2.2 is related by the inclusion maps c_i:K^•(Xi)→K_•(Xi) and c_X:K^•(X)→K_•(X): β∘colim(c_i)=c_X∘α_perf, since both send a perfect complex E_i to its derived pullback class, and flat pullback computes that class here. The current relabelling (3.2.3.1) therefore follows the exact map used in the diagram; diplomatic preserves the duplicate tag and both printed references.

[208] EN-VAR-201 Source folio 295. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=309; scan/SGA6_witness_300dpi.djvu#page=306
Printed reading: $x \in K_\bullet(X)$
Corrected reading: $x \in K_\bullet(X/S)$
Source: ASTRA-IV-PROJECTION-295. The source reads x∈K_•(X) in the first branch and a∈K_•(S), x∈K^•(X/S) in the second. The immediately preceding pushforward is defined on K_•(X/S). The first domain is repaired to that exact relative domain: for a=[A] perfect on S and x=[E] in K_•(X/S), ax=[E⊗^L_S A], and III3.7 identifies Rf_*E⊗^L_S A with Rf_*(E⊗^L_S A); its finite-amplitude condition holds for A. The second coefficient is retained: 3.3.2.2 applied to X→S→S pairs x=[E] in K^•(X/S) with a=[M] in K_•(S), giving [E⊗^L_S M] in K_•(X/S). Here III3.7 uses E's finite S-amplitude, and both sides land in K_•(Y/S). The source absolute first domain fails concretely for S=Y=Spec k[x1,x2,...], X=Spec k and the closed immersion f defined by all variables: O_X is perfect over k, whereas f_*O_X=k is not finitely presented over the base ring, so the absolute pushforward does not land in D^b(Y)_coh. The complete map identification and local finite-presentation obstruction are saved in identification_295.json. Only the first missing /S is changed.

[209] EN-CONTRIB-V-304-01 Source folio 304. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=318; scan/SGA6_witness_300dpi.djvu#page=315
Printed reading:

$$H(h)(\Phi_A(\chi)) = H(h) \circ H(f_\chi(\psi)) = H(f_{\widehat{G}(h)(\chi)})(\psi) = \Phi_{A'}(\widehat{G}(h)(\chi).$$

Corrected reading:

$$H(h)(\Phi_A(\chi)) = H(h) \circ H(f_\chi)(\psi) = H(f_{\widehat{G}(h)(\chi)})(\psi) = \Phi_{A'}(\widehat{G}(h)(\chi)).$$

Source: check/findings/findings_301_305.json#17; check/findings/findings_301_305.json#18. check/findings/findings_301_305.json#17: The native top crop shows the two closes after ψ. Preserve that printed grouping diplomatically. On 303, ψ belongs to H(K[Xi]) and f_χ:K[Xi]→A, so H(f_χ):H(K[Xi])→H(A) applies to ψ; H(h):H(A)→H(A′) then applies to its value. This exact typed composite supports the existing current H(h)∘H(f_χ)(ψ). check/findings/findings_301_305.json#18: Both scans show the full stop immediately after the single closing parenthesis. The opening parenthesis of the outer Φ_{A′} application is unmatched. The source layer keeps it; the current layer supplies exactly that one missing close, giving Φ_{A′}($\widehat{G}$(h)(χ)).

[210] EN-CONTRIB-V-306-01 Source folio 306. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=320; scan/SGA6_witness_300dpi.djvu#page=317
Printed reading: $(1 + X^{(j)}t)_{j=1,..,k}$
Corrected reading: $(1 + X^{(j)}t)_{j=1,...,k}$
Source: check/findings/findings_306_310.json#2. The native top crop shows two full stops between the two commas in j=1,..,k. The diplomatic layer already has exactly two stops. The current three-dot ellipsis supplies the standard index-list abbreviation, with the identical endpoints 1 and k.

[211] EN-CONTRIB-V-311-01 Source folio 311. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=325; scan/SGA6_witness_300dpi.djvu#page=322
Printed reading:

$$\text{i)} \quad H(f_1)(\theta) = 0;$$
$$\text{ii)} \quad \forall n \geq 1, \quad H(f_n)(u_H^n(\theta)) \equiv u_H^n(\theta) \text{ mod. } F^n H(K[X_1, \dots X_n]).$$

Corrected reading:

$$\text{i)} \quad H(f_1)(\theta) = 0;$$
$$\text{ii)} \quad \forall n \geq 1, \quad H(f_n)(u_H^n(\theta)) \equiv u_H^n(\theta) \text{ mod. } F^n H(K[X_1, \dots, X_n]).$$

Source: check/findings/findings_311_315.json#2. The native top crop resolves the first list exactly as printed. The later quotient has three complete lists and the following membership formula has a fourth, each with the second comma. The repaired expression denotes the same polynomial algebra in the listed generators X_1 through X_n, with its missing separator restored. The attached diplomatic omission and corrected comma agree.

[212] EN-CONTRIB-V-312-01 Source folio 312. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=326; scan/SGA6_witness_300dpi.djvu#page=323
Printed reading: $.km$
Corrected reading: km
Source: ASTRA-V-POINT-312. Both scans show the detached low dot. The native marks crop compares it directly with the following colon, the sentence-final full stop, and two other lowercase k forms. The calculation in review/expose_v/identification_312.json uses X_j^(a)↦e_j(T_1^(a),...,T_N^(a)) and the separately multilinear root expansion indexed by J=(j_1,...,j_r), with σ_J(U_a)=T_(j_a)^(a). Each degree-d coefficient term has Σ_Jq_J=d and d≤Σ_Jw_J≤mΣ_Jq_J=md. The graded injection preserves these weights on descent. Setting d=k proves the stated endpoint km and justifies treating the preceding dot as surplus punctuation.

[213] EN-CONTRIB-V-316-01 Source folio 316. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=330; scan/SGA6_witness_300dpi.djvu#page=327
Printed reading: $f : (T, \underline{A}) \longrightarrow (T', \underline{A}!)$
Corrected reading: $f : (T, \underline{A}) \longrightarrow (T', \underline{A}')$
Source: check/findings/findings_316_320.json#3. The native lower crop and witness_316_Amark.png resolve a mark taller than the nearby primes and with a lower dot. The declared target immediately above is (T′,A′), and the displayed inverse-image map immediately below has source K_naif(T′,A′). The ringed-topos morphism therefore uses the same A′, with structure map f^-1A′→A and module pullback A⊗_{f^-1A′}f^-1E′. This explicit identification supports the corrected prime while the diplomatic layer retains the printed exclamation-shaped mark. The underline lies only under A.

[214] EN-CONTRIB-V-317-01 Source folio 317. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=331; scan/SGA6_witness_300dpi.djvu#page=328
Printed reading:

$$\begin{array}{ccc} \widehat{G}\times\widehat{G}\times\widehat{G} \to \widehat{G}\times\widehat{G} \\ \downarrow \qquad\quad \downarrow \\ \widehat{G}\times\widehat{G} \longrightarrow \widehat{G} \end{array} \qquad \begin{array}{ccc} \widehat{G}\times\widehat{G} \to \widehat{G}\times\widehat{G} \\ \downarrow \qquad\quad \downarrow \\ \widehat{G} \xrightarrow{\text{Id:}} \widehat{G}, \end{array}$$

Corrected reading:

$$\begin{array}{ccc} \widehat{G}\times\widehat{G}\times\widehat{G} \to \widehat{G}\times\widehat{G} \\ \downarrow \qquad\quad \downarrow \\ \widehat{G}\times\widehat{G} \longrightarrow \widehat{G} \end{array} \qquad \begin{array}{ccc} \widehat{G}\times\widehat{G} \to \widehat{G}\times\widehat{G} \\ \downarrow \qquad\quad \downarrow \\ \widehat{G} \xrightarrow{\text{Id}} \widehat{G}, \end{array}$$

Source: check/findings/findings_316_320.json#11. The tight native crop resolves both points, separate from the arrowhead below. The arrow is Id_$\widehat{G}$:$\widehat{G}\to\widehat{G}$, $\alpha\mapsto\alpha$, in the commutativity square; the colon supplies no further argument or defined expression. The attached diplomatic Id: and corrected Id preserve the printed punctuation and its deletion respectively.

[215] EN-CONTRIB-V-318-01 Source folio 318. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=332; scan/SGA6_witness_300dpi.djvu#page=329
Printed reading: $1 + Xt, 1 + Xt$
Corrected reading: $1 + Xt, 1 + Yt$
Source: check/findings/findings_316_320.json#12. Both scans resolve the repeated X. The second317 square compares m:$\widehat{G}\times\widehat{G}\to\widehat{G}$ with m$\circ\tau$, where $\tau(\alpha,\beta)=(\beta,\alpha)$. On the independent universal pair(1+Xt,1+Yt), equation2.3.1 gives m=1+XYt and m$\circ\tau$=1+YXt; these coincide in the commutative ring Z[X,Y]. The multi-variable uniqueness of1.7/1.10 then applies to these two separately bilinear natural operations. Repeating X specializes the universal pair along Y$\mapsto$X. Restore Y in the corrected universal-pair test and retain the printed repetition diplomatically.

[216] EN-CONTRIB-V-318-02 Source folio 318. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=332; scan/SGA6_witness_300dpi.djvu#page=329
Printed reading: p_2
Corrected reading: P_2
Source: check/findings/findings_316_320.json#13. The native top crop directly contrasts p_2 with P_n and the displayed P_2. The explicit following polynomial is the n=2 member of the just-named family P_n. The attached diplomatic p_2 and corrected P_2 agree with the source glyph and this exact identification.

[217] EN-CONTRIB-V-320-01 Source folio 320. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=334; scan/SGA6_witness_300dpi.djvu#page=331
Printed reading: $i.j \in \mathbb{N}$
Corrected reading: $i, j \in \mathbb{N}$
Source: check/findings/findings_316_320.json#26. The native tight crop witness_320_ijN.png resolves the stop separately from the following j descender and the later comma. The preceding319 definition labels Q_{i,j} by the separately chosen operation index i and coefficient index j. Under successive coefficient extraction [u^i] and [t^j], the identity λ_u(λ_t(x))=$\widehat{G}$(λ_t)(λ_u(x)) gives λ^j(λ^i(x))=Q_{i,j}(λ^1(x),...,λ^{ij}(x)) for the pair (i,j)$\in$N$\times$N. Thus a comma expresses the two independent indices in this exact displayed family; retain the stop diplomatically and restore the comma currently. Retain the inspected double-stem N in both layers.

[218] EN-CONTRIB-V-321-01 Source folio 321. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=335; scan/SGA6_witness_300dpi.djvu#page=332
Printed reading:

$$\begin{array}{ccc} \widehat{G}^t \times \widehat{G}^t & \xrightarrow{\lambda_u \times \lambda_u} & (\widehat{G}^u \circ \widehat{G}^t) \times \widehat{G}^u \circ \widehat{G}^t) \\ {\scriptstyle\circ}\downarrow & & \downarrow{\scriptstyle\circ} \\ \widehat{G}^t & \xrightarrow{\lambda_u} & \widehat{G}^u \circ \widehat{G}^t, \end{array} \qquad \begin{array}{ccc} \widehat{G}^t & \xrightarrow{\lambda_v} & \widehat{G}^v \circ \widehat{G}^t \\ {\scriptstyle\lambda_u}\downarrow & & \downarrow{\scriptstyle\widehat{G}^v(\lambda_u)} \\ \widehat{G}^u \circ \widehat{G}^t & \xrightarrow{\lambda_v} & \widehat{G}^v \circ \widehat{G}^u \circ \widehat{G}^t, \end{array}$$

Corrected reading:

$$\begin{array}{ccc} \widehat{G}^t \times \widehat{G}^t & \xrightarrow{\lambda_u \times \lambda_u} & (\widehat{G}^u \circ \widehat{G}^t) \times (\widehat{G}^u \circ \widehat{G}^t) \\ {\scriptstyle\circ}\downarrow & & \downarrow{\scriptstyle\circ} \\ \widehat{G}^t & \xrightarrow{\lambda_u} & \widehat{G}^u \circ \widehat{G}^t, \end{array} \qquad \begin{array}{ccc} \widehat{G}^t & \xrightarrow{\lambda_v} & \widehat{G}^v \circ \widehat{G}^t \\ {\scriptstyle\lambda_u}\downarrow & & \downarrow{\scriptstyle\widehat{G}^v(\lambda_u)} \\ \widehat{G}^u \circ \widehat{G}^t & \xrightarrow{\lambda_v} & \widehat{G}^v \circ \widehat{G}^u \circ \widehat{G}^t, \end{array}$$

Source: check/findings/findings_321_325.json#4. The native lower diagram crop resolves the empty interval after the product sign and before the second $\widehat{G}\hat{}$u. The top horizontal map λ_u×λ_u takes (α,β) to (λ_u(α),λ_u(β)) in ($\widehat{G}\hat{}$u∘$\widehat{G}\hat{}$t)×($\widehat{G}\hat{}$u∘$\widehat{G}\hat{}$t); the right vertical map multiplies these two elements by the circle law. This exact product target identifies the missing opening parenthesis. Retain the printed omission diplomatically and the supplied parenthesis in the corrected layer.

[219] EN-CONTRIB-V-322-01 Source folio 322. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=336; scan/SGA6_witness_300dpi.djvu#page=333
Printed reading:

$$\begin{aligned} \lambda_u(1+Xt) \circ \lambda_u(1+Yt) &= (\underline{1} + \{Xt\}u) \circ (\underline{1} + \{1+Yt\}u) \\ &= \underline{1} + \{(1+Xt) \circ (1+Yt)\}u \\ &= \underline{1} + \{1+XYt\}u \\ &= \lambda_u((1+Xt) \circ (1+Yt)). \end{aligned}$$

Corrected reading:

$$\begin{aligned} \lambda_u(1+Xt) \circ \lambda_u(1+Yt) &= (\underline{1} + \{1+Xt\}u) \circ (\underline{1} + \{1+Yt\}u) \\ &= \underline{1} + \{(1+Xt) \circ (1+Yt)\}u \\ &= \underline{1} + \{1+XYt\}u \\ &= \lambda_u((1+Xt) \circ (1+Yt)). \end{aligned}$$

Source: check/findings/findings_321_325.json#8. The native top crop resolves the complete short brace group. Equation2.3.3 defines λ_u(1+Xt)=1+{1+Xt}u. Its coefficient-of-u map therefore sends this universal line element to1+Xt. Substitution into the first factor of the displayed product requires exactly the missing1+; the following line already uses (1+Xt)∘(1+Yt). Restore these two tokens currently and keep the printed Xt diplomatically.

[220] EN-CONTRIB-V-322-02 Source folio 322. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=336; scan/SGA6_witness_300dpi.djvu#page=333
Printed reading:

$$\text{i)} \quad K \text{ is torsion-free,}$$
$$\text{ii)} \quad \forall x \in K,\ \forall n \in \mathbb{N}, \quad \text{we have } \tfrac{x}{n} \in K$$

Corrected reading:

$$\text{i)} \quad K \text{ is torsion-free,}$$
$$\text{ii)} \quad \forall x \in K,\ \forall n \in \mathbb{N}, \quad \text{we have } \tbinom{x}{n} \in K$$

Source: check/findings/findings_321_325.json#13. The native top crop directly contrasts this fraction with the parenthesized binomial on the next line. Under the injection j:K→K⊗_Z Q supplied by torsion-freeness, the prescribed coefficient is [t^n]exp(j(x)Log(1+t))=j(x)(j(x)−1)…(j(x)−n+1)/n!, with coefficient1 at n=0. Thus the condition for these prescribed coefficients to descend to K is binom(x,n)∈K, as explicitly used in λ^n(x) and2.6.1. The printed fraction x/n does not supply that coefficient: for x=3,n=2 the fraction is3/2 whereas binom(3,2)=3; at n=0 the printed denominator is0. Retain the fraction diplomatically and the existing binomial correction.

[221] EN-CONTRIB-V-323-01 Source folio 323. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=337; scan/SGA6_witness_300dpi.djvu#page=334
Printed reading: pre- -ring Z
Corrected reading: pre-λ-ring Z
Source: check/findings/findings_321_325.json#15. The native top crop resolves the clear unfilled interval. The explicit locator2.5.a) identifies the structure λ^i(n)=binom(n,i) on the integer ring, and the surrounding sentence identifies its defining relations2.4.2. These exact operations supply the missing λ in the corrected reading; retain the blank diplomatically.

[222] EN-CONTRIB-V-323-02 Source folio 323. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=337; scan/SGA6_witness_300dpi.djvu#page=334
Printed reading: (2.5.a)) is a -ring;
Corrected reading: (2.5.a)) is a λ-ring;
Source: check/findings/findings_321_325.json#17. The native top crop resolves the blank. The immediately cited2.5.a) proves that its prescribed operations satisfy2.4.2, exactly the λ-anneau assertion invoked here. Retain the blank diplomatically and the attached λ currently.

[223] EN-CONTRIB-V-325-01 Source folio 325. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=339; scan/SGA6_witness_300dpi.djvu#page=336
Printed reading:

$$\gamma_t(x) = \lambda_{t/1-t}(x), \qquad \underline{\text{hence}}$$

Corrected reading:

$$\gamma_t(x) = \lambda_{t/(1-t)}(x), \qquad \underline{\text{hence}}$$

Source: ASTRA-V-SUBSTITUTION-325. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[224] EN-CONTRIB-V-325-02 Source folio 325. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=339; scan/SGA6_witness_300dpi.djvu#page=336
Printed reading:

$$\lambda_t(x) = \gamma_{t/1+t}(x).$$

Corrected reading:

$$\lambda_t(x) = \gamma_{t/(1+t)}(x).$$

Source: ASTRA-V-SUBSTITUTION-325. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[225] EN-CONTRIB-V-325-03 Source folio 325. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=339; scan/SGA6_witness_300dpi.djvu#page=336
Printed reading: $\lambda_t(-1) = 1/1 + t$
Corrected reading: $\lambda_t(-1) = 1/(1 + t)$
Source: ASTRA-V-SUBSTITUTION-325. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[226] EN-CONTRIB-V-325-04 Source folio 325. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=339; scan/SGA6_witness_300dpi.djvu#page=336
Printed reading: $\lambda_{t/1-t}(x) = \sum_{n\geq 0} \lambda^n(x).t^n(1-t)^{-n}$.
Corrected reading: $\lambda_{t/(1-t)}(x) = \sum_{n\geq 0} \lambda^n(x).t^n(1-t)^{-n}$.
Source: ASTRA-V-SUBSTITUTION-325. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[227] EN-CONTRIB-V-325-05 Source folio 325. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=339; scan/SGA6_witness_300dpi.djvu#page=336
Printed reading: $\lambda_{t/1-t}(x) = \sum_{n\geq 0}[\lambda^n(x).\sum_{p\geq n}\binom{p-1}{p-n}t^p]$
Corrected reading: $\lambda_{t/(1-t)}(x) = \sum_{n\geq 0}[\lambda^n(x).\sum_{p\geq n}\binom{p-1}{p-n}t^p]$
Source: ASTRA-V-SUBSTITUTION-325. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[228] EN-CONTRIB-V-326-01 Source folio 326. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=340; scan/SGA6_witness_300dpi.djvu#page=337
Printed reading: $\rho_t(t) = t/1 - t$
Corrected reading: $\rho_t(t) = t/(1 - t)$
Source: ASTRA-V-SUBSTITUTION-326. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[229] EN-CONTRIB-V-326-02 Source folio 326. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=340; scan/SGA6_witness_300dpi.djvu#page=337
Printed reading:

$$\widehat{G}\times\widehat{G}\xrightarrow{(\rho^{-1}.\rho^{-1})}\widehat{G}\times\widehat{G}\overset{\circ}{\to}\widehat{G}\overset{\rho}{\to}\widehat{G},$$

Corrected reading:

$$\widehat{G}\times\widehat{G}\xrightarrow{(\rho^{-1},\rho^{-1})}\widehat{G}\times\widehat{G}\overset{\circ}{\to}\widehat{G}\overset{\rho}{\to}\widehat{G},$$

Source: check/findings/findings_326_330.json#4. The native lower crop compares both marks directly. The first arrow is the Cartesian product map $(\alpha,\beta)\mapsto(\rho^{-1}(\alpha),\rho^{-1}(\beta))$ from $\widehat{G}\times\widehat{G}$ to itself. Its subsequent composite is $\rho\circ m\circ(\rho^{-1}\times\rho^{-1})$, with m the circle multiplication from2.3. This exact componentwise map identifies the corrected comma. Retain the printed stop in the diplomatic label and the attached comma in the corrected label.

[230] EN-CONTRIB-V-327-01 Source folio 327. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=341; scan/SGA6_witness_300dpi.djvu#page=338
Printed reading:

$$\begin{aligned}(1+Xt)\nabla(1+Yt) &= \rho\Big(\rho^{-1}(1+Xt)\circ\rho^{-1}(1+Yt)\Big)\\ &= \rho\Big(\frac{1+(X+1)t}{1+t}\circ\frac{1+(Y+1)t}{1+t}\Big)\\ &= \rho\Big(\frac{(1+(X+1)(Y+1)t)(1+t)}{(1+(X+1)t)(1+(Y+1)t)}\Big)\\ &= \frac{1+(1+X+Y+XY)t}{(1+Xt)(1+Yt)}.\end{aligned} \tag{3.5.1}$$

Corrected reading:

$$\begin{aligned}(1+Xt)\nabla(1+Yt) &= \rho\Big(\rho^{-1}(1+Xt)\circ\rho^{-1}(1+Yt)\Big)\\ &= \rho\Big(\frac{1+(X+1)t}{1+t}\circ\frac{1+(Y+1)t}{1+t}\Big)\\ &= \rho\Big(\frac{(1+(X+1)(Y+1)t)(1+t)}{(1+(X+1)t)(1+(Y+1)t)}\Big)\\ &= \frac{1+(X+Y+XY)t}{(1+Xt)(1+Yt)}.\end{aligned} \tag{3.5.1}$$

Source: ASTRA-V-NUMERATOR-327. Both main and native top images resolve this exact numerator. In R=Z[X,Y][[t]], the explicitly defined automorphism is ρ(h(t))=h(t/(1−t)). Apply it to the immediately preceding fraction: setting c=(1+X)(1+Y), its numerator becomes(1+(c−1)t)/(1−t)^2 and its denominator becomes(1+Xt)(1+Yt)/(1−t)^2. Cancellation of the unit(1−t)^2 yields exactly[1+(X+Y+XY)t]/[(1+Xt)(1+Yt)]. At X=Y=0 the preceding expression isρ(1)=1, whereas the printed final expression is1+t. The coordinator independently confirmed this exact calculation. Full source-linked derivation: review/expose_v/identification_327.json.

[231] EN-CONTRIB-V-327-02 Source folio 327. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=341; scan/SGA6_witness_300dpi.djvu#page=338
Printed reading: $1/1-t$
Corrected reading: $1/(1-t)$
Source: ASTRA-V-UNIT-DENOMINATOR-327. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[232] EN-CONTRIB-V-328-01 Source folio 328. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=342; scan/SGA6_witness_300dpi.djvu#page=339
Printed reading:

$$\begin{aligned}\lambda'_u(1+Xt) &= \widehat{G}(\rho_t)(\lambda_u(\rho_t^{-1}(1+Xt)))\\ &= \widehat{G}(\rho_t)\left(\lambda_u\left(\frac{1+(X+1)t}{1+t}\right)\right)\\ &= \widehat{G}(_t)\left(\frac{\underline{1}_o+\{1+(X+1)t\}u}{\underline{1}_o+\{1+t\}u}\right)\\ &= \underline{1}_\nabla+\frac{\{1+Xt\}u}{\underline{1}_\nabla+\underline{1}_\nabla.u}\\ &= \rho_u^{-1}(\underline{1}_\nabla+\{1+Xt\}u).\end{aligned}$$

Corrected reading:

$$\begin{aligned}\lambda'_u(1+Xt) &= \widehat{G}(\rho_t)(\lambda_u(\rho_t^{-1}(1+Xt)))\\ &= \widehat{G}(\rho_t)\left(\lambda_u\left(\frac{1+(X+1)t}{1+t}\right)\right)\\ &= \widehat{G}(\rho_t)\left(\frac{\underline{1}_o+\{1+(X+1)t\}u}{\underline{1}_o+\{1+t\}u}\right)\\ &= \underline{1}_\nabla+\frac{\{1+Xt\}u}{\underline{1}_\nabla+\underline{1}_\nabla.u}\\ &= \rho_u^{-1}(\underline{1}_\nabla+\{1+Xt\}u).\end{aligned}$$

Source: check/findings/findings_326_330.json#13. Both main342 and native339 resolve the empty position. The explicitly defined operation on327 is lambda-prime_u=Ghat(rho_t) composed with lambda_u composed with rho_t inverse. The first two328 rows retain that same outer Ghat(rho_t); the third merely expands lambda_u on the quotient, using lambda_u(a-b)=lambda_u(a)/lambda_u(b). Thus the omitted character is exactly rho and the already paired diplomatic empty subscript/current rho agrees with this composite.

[233] EN-CONTRIB-V-329-01 Source folio 329. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=343; scan/SGA6_witness_300dpi.djvu#page=340
Printed reading:

$$\begin{array}{ccc} A & \xrightarrow{\gamma_u} & \widehat{G}^u(A)_\nabla \\ {\scriptstyle\gamma_t}\big\downarrow & & \big\downarrow{\scriptstyle\widehat{G}^u(\gamma_t)} \\ \widehat{G}^t(A)_\nabla & \overset{\gamma'_u}{\to} & \widehat{G}^u(\widehat{G}^t(A)_\nabla)_\nabla. \end{array} \tag{3.7.2}$$

Corrected reading:

$$\begin{array}{ccc} A & \xrightarrow{\gamma_u} & \widehat{G}^u(A)_\nabla \\ {\scriptstyle\gamma_t}\big\downarrow & & \big\downarrow{\scriptstyle\widehat{G}^u(\gamma_t)} \\ \widehat{G}^t(A)_\nabla & \overset{\gamma'_u}{\to} & \widehat{G}^u(\widehat{G}^t(A)_\nabla)_\nabla. \end{array} \tag{3.7.2}$$

Source: check/findings/findings_326_330.json#25. The native top crop resolves the lower end of the vertical stroke as a plain termination, while the neighbouring right vertical arrow has its two head branches. The map is explicitly gamma_t:A→Ghat^t(A)_nabla, x↦lambda_{t/(1-t)}(x), from3.2 and the immediately preceding description. Its source and target fix the downward orientation used in the current layer. The attached diplomatic dash/current headed arrow therefore preserve both the observed omission and the specified map.

[234] EN-CONTRIB-V-329-02 Source folio 329. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=343; scan/SGA6_witness_300dpi.djvu#page=340
Printed reading:

$$\gamma_t(x) = (1/1-t)^x = \exp(-x\,\mathrm{Log}(1-t)) \tag{3.8.1}$$

Corrected reading:

$$\gamma_t(x) = (1/(1-t))^x = \exp(-x\,\mathrm{Log}(1-t)) \tag{3.8.1}$$

Source: ASTRA-V-SLASH-329. The already inspected proof on325 expands the substituted monomial as t^n(1-t)^(-n), fixing rho_minus:h(t)→h(t/(1-t)). The inverse map is rho_plus:h(t)→h(t/(1+t)): the two fractional substitutions compose to t in both orders. Also lambda_t(-1)lambda_t(1)=1 and lambda_t(1)=1+t, so lambda_t(-1)=1/(1+t). The unit on327 is rho_minus(1+t)=1+t/(1-t)=1/(1-t). On329, applying rho_minus to(1+t)^x gives(1/(1-t))^x=exp(-xLog(1-t)). Thus each grouping below is fixed by an explicit map or multiplicative inverse, rather than a general change to slash notation. These are post-checkpoint08 current-only grouping amendments; the printed forms remain diplomatic and the immutable snapshot remains unchanged.

[235] EN-CONTRIB-V-332-01 Source folio 332. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=346; scan/SGA6_witness_300dpi.djvu#page=343
Printed reading: $i = 1, .., n$
Corrected reading: $i = 1, \ldots, n$
Source: check/findings/findings_331_335.json#4. The native top crop shows the two isolated dots and both commas separately. The supplied diplomatic two-dot/current three-dot pair agrees with the observed omission and its local repair.

[236] EN-CONTRIB-V-334-01 Source folio 334. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=348; scan/SGA6_witness_300dpi.djvu#page=345
Printed reading: and let $(P_i)_{i' \in I'}$ be a family
Corrected reading: and let $(P_{i'})_{i' \in I'}$ be a family
Source: check/findings/findings_331_335.json#8. The native top crop resolves the closing parenthesis immediately after i. Definition4.6 and Proposition4.7 on the same page both give the relation family(P_i-prime) indexed by i-prime in I-prime; the sentence also names its generated ideal using P_i-prime. The exact family is the map I-prime→A, i-prime↦P_i-prime. The attached diplomatic inner i/current inner i-prime pair restores that same bound variable and leaves every outer index unchanged.

[237] EN-CONTRIB-V-334-02 Source folio 334. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=348; scan/SGA6_witness_300dpi.djvu#page=345
Printed reading: as follows from solutions 3.2.
Corrected reading: as follows from relations 3.2.
Source: ASTRA-V-REFERENCE-334. The native345 top crop and main third strip clearly resolve solutions. The cited Proposition3.2(ii) on325 states the relations gamma_t(x)=lambda_{t/(1-t)}(x) and lambda_t(x)=gamma_{t/(1+t)}(x). For n>=1 their coefficients are gamma^n(x)=sum_(j=1)^n binom(n-1,j-1)lambda^j(x) and lambda^n(x)=sum_(j=1)^n(-1)^(n-j)binom(n-1,j-1)gamma^j(x). The substitutions compose to t in both orders, so these integral triangular transformations are inverse. For every x of the given ideal J, membership of all positive-degree operations from either family gives membership of all operations from the other family. This is exactly the reference used in the sentence; identification_334.json records the maps and calculation.

[238] EN-CONTRIB-V-334-03 Source folio 334. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=348; scan/SGA6_witness_300dpi.djvu#page=345
Printed reading: $A/J \to B$
Corrected reading: $f' : A/J \to B$
Source: check/findings/findings_331_335.json#9. Both native345 lower and main fourth strip show an empty interval between seul and A/J. Let q:A→A/J be the quotient and f:A→B the evaluation map of4.2, X_i↦b_i and lambda^n X_i↦lambda^n(b_i). The displayed relation hypothesis gives f(P_i-prime)=0; its kernel is the lambda-ideal described immediately above, so J lies in ker(f). The uniquely induced map is f-prime:A/J→B, a+J↦f(a), with f-prime q=f. Its values on the classes of X_i are precisely the printed b_i. Supply the missing name f-prime only in the current layer.

[239] EN-CONTRIB-V-334-04 Source folio 334. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=348; scan/SGA6_witness_300dpi.djvu#page=345
Printed reading: $f'(X_i) = b_i$
Corrected reading: $f'(X_i) = b_i$.
Source: check/findings/findings_331_335.json#10. The native lower crop shows the dot of the subscript i above its stem, but no separate terminal dot to the right. The attached diplomatic absence/current supplied sentence stop is the exact recorded punctuation pair.

[240] EN-CONTRIB-V-335-01 Source folio 335. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=349; scan/SGA6_witness_300dpi.djvu#page=346
Printed reading: $1 + X_1 t + \ldots + X_n t^n = \prod_{i=1}(1 + T_i t)$
Corrected reading: $1 + X_1 t + \ldots + X_n t^n = \prod_{i=1}^{n}(1 + T_i t)$
Source: check/findings/findings_331_335.json#12. Main349 and native346 place the subscript n of the preceding T_n] on the preceding text line; no upper bound belongs to the first product. The exact ring map K[X_1,...,X_n]→K[T_1,...,T_n], X_j↦e_j(T_1,...,T_n), sends1+sum_(j=1)^n X_j t^j to product_(i=1)^n(1+T_i t), by the coefficient definition of e_j. This identifies precisely the omitted upper bound n, also displayed on both next products. Preserve the missing upper bound diplomatically and restore n in the current layer.

[241] EN-CONTRIB-V-336-01 Source folio 336. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=350; scan/SGA6_witness_300dpi.djvu#page=347
Printed reading: $\gamma_t(\epsilon(a)X) = (\gamma_t(X))^{\epsilon(a)} = \sum_{n=0}^{\infty} R_{n,a}((\gamma^j X)) \, . \, t^n$
Corrected reading: $\gamma_t(\epsilon(a)X) = (\gamma_t(X))^{\epsilon(a)} = \sum_{n=0}^{\infty} R_{n,\epsilon(a)}((\gamma^j X)) \, . \, t^n$
Source: ASTRA-V-COEFFICIENT-PARAMETER-336. Native347 lower and its tight R crop resolve the bare a in both indices. Definition2.9 and3.8 give R_(n,c) as the coefficient of the scalar-power operation phi↦phi^c. Here the displayed exponent is exactly epsilon(a), so the parameter map is A→K, a↦epsilon(a). In degree1 the left expression has coefficient epsilon(a)X, while the printed R_(1,a) yields aX. For K=Z and A the free lambda-ring on U augmented by U↦0, a=U gives0 versus UX in the underlying polynomial ring B. The map setting all other independent generators to0 sends UX to the nonzero monomial UX in Z[U,X]. The coordinator independently confirmed the same defect using a Laurent line element L of augmentation1. Full identification: identification_336.json.

[242] EN-CONTRIB-V-336-02 Source folio 336. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=350; scan/SGA6_witness_300dpi.djvu#page=347
Printed reading: $R_{n,a}$
Corrected reading: $R_{n,\epsilon(a)}$
Source: ASTRA-V-COEFFICIENT-PARAMETER-336. Native347 lower and its tight R crop resolve the bare a in both indices. Definition2.9 and3.8 give R_(n,c) as the coefficient of the scalar-power operation phi↦phi^c. Here the displayed exponent is exactly epsilon(a), so the parameter map is A→K, a↦epsilon(a). In degree1 the left expression has coefficient epsilon(a)X, while the printed R_(1,a) yields aX. For K=Z and A the free lambda-ring on U augmented by U↦0, a=U gives0 versus UX in the underlying polynomial ring B. The map setting all other independent generators to0 sends UX to the nonzero monomial UX in Z[U,X]. The coordinator independently confirmed the same defect using a Laurent line element L of augmentation1. Full identification: identification_336.json.

[243] EN-CONTRIB-V-341-01 Source folio 341. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=355; scan/SGA6_witness_300dpi.djvu#page=352
Printed reading:

$$\begin{aligned}(n,x)+(n',x') &= (n+n',x+x')\\ (n,x)\,.\,(n',x') &= (nn',nx'+n'x+x\,._N\,y)\end{aligned} \tag{5.5.2}$$

Corrected reading:

$$\begin{aligned}(n,x)+(n',x') &= (n+n',x+x')\\ (n,x)\,.\,(n',x') &= (nn',nx'+n'x+x\,._N\,x')\end{aligned} \tag{5.5.2}$$

Source: check/findings/findings_341_345.json#6. Both scans clearly resolve y. Write D=lambda_(minus1)(N) and x star y=xyD. In the unitization, (n,x)(n-prime,x-prime)=(nn-prime,nx-prime+n-prime x+x star x-prime), obtained by multiplying n times the adjoined unit plus x with its second counterpart. Thus the exact second operand is x-prime. With A=Z,N=0,n=n-prime=0,x=1,x-prime=0,y=1, the printed expression gives(0,1) for multiplication by zero, while the unitization gives(0,0). Keep y in diplomatic and x-prime in current. Full maps: identification_341.json.

[244] EN-CONTRIB-V-342-01 Source folio 342. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=356; scan/SGA6_witness_300dpi.djvu#page=353
Printed reading: $x \in \Lambda$
Corrected reading: $x \in A$
Source: check/findings/findings_341_345.json#9. The native tight crop and nearest-neighbour zoom, together with the main tight crop, resolve the empty interior throughout the expected crossbar region. The other A letters on the page have a crossbar. Preserve Lambda diplomatically. The argument is the element(0,x) of Z⊕A, so its actual coordinate domain is A under the explicitly stated inclusion x↦(0,x); retain current A.

[245] EN-CONTRIB-V-342-02 Source folio 342. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=356; scan/SGA6_witness_300dpi.djvu#page=353 Printed reading:

$$\gamma^p(0,x)=\sum_{i=0}^{p-1}\binom{p-1}{i}\lambda^i(N,x),$$

Corrected reading:

$$\gamma^p(0,x)=\sum_{i=0}^{p-1}\binom{p-1}{i}\lambda^{p-i}(N,x),$$

Source: ASTRA-V-GAMMA-EXPONENT-342. Main second strip and native353 top clearly resolve the exponent i without a preceding p-minus. Definition3.2 gives gamma^p(z)=lambda^p(z+(p-1)1). The actual map is coefficient extraction[t^p] from(1+t)^{p-1}lambda_t(z), giving sum_{i=0}^{p-1}binom(p-1,i)lambda^{p-i}(z). For z=(0,x), every p-i>=1, so lambda^{p-i}(z) is the embedded lambda^{p-i}(N,x). At p=1,x=0 the source sum gives the adjoined unit, but gamma^1(0,0)=0. The coordinator independently verified the coefficient expansion. Full exact proof and the initial injective map: identification_342.json.

[246] EN-CONTRIB-V-342-03 Source folio 342. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=356; scan/SGA6_witness_300dpi.djvu#page=353 Printed reading:

$$
\begin{aligned}
&1 + \lambda_{-1}(N) \,.\, \lambda_{-1}(L) \,.\, \sum_{k\geq 1} \lambda^k(N+L,x) \,.\, t^k \\
&\quad = \lambda_t(x \,.\, \lambda_{-1}(N)) \circ \lambda_t(1-L) \\
&\quad = \frac{\lambda_t(x \,.\, \lambda_{-1}(N))}{\lambda_{Lt}(x \,.\, \lambda_{-1}(N))} \qquad \left(\text{since } \lambda_t(1-L) = \frac{\lambda_t(1)}{\lambda_t(L)} = \frac{1+t}{1+Lt}\text{. and we apply (2.3.2)}\right) \\
&\quad = \frac{1 + \ldots + \lambda^k(x \,.\, \lambda_{-1}(N)) t^k + \ldots}{1 + \ldots + \lambda^k(x \,.\, \lambda_{-1}(N)) L^k t^k + ..} \\
&\quad = 1 + \frac{\sum_{k\geq 1} \lambda^k(N,x) \,.\, \lambda_{-1}(N) \,.\, (1-L^k) t^k}{1 + \sum_{k\geq 1} \lambda^k(N,x) \,.\, \lambda_{-1}(N) \,.\, .L^k t^k}
\end{aligned}
$$

Corrected reading:

$$
\begin{aligned}
&1 + \lambda_{-1}(N) \,.\, \lambda_{-1}(L) \,.\, \sum_{k\geq 1} \lambda^k(N+L,x) \,.\, t^k \\
&\quad = \lambda_t(x \,.\, \lambda_{-1}(N)) \circ \lambda_t(1-L) \\
&\quad = \frac{\lambda_t(x \,.\, \lambda_{-1}(N))}{\lambda_{Lt}(x \,.\, \lambda_{-1}(N))} \qquad \left(\text{since } \lambda_t(1-L) = \frac{\lambda_t(1)}{\lambda_t(L)} = \frac{1+t}{1+Lt}\text{, and we apply (2.3.2)}\right) \\
&\quad = \frac{1 + \ldots + \lambda^k(x \,.\, \lambda_{-1}(N)) t^k + \ldots}{1 + \ldots + \lambda^k(x \,.\, \lambda_{-1}(N)) L^k t^k + \ldots} \\
&\quad = 1 + \frac{\sum_{k\geq 1} \lambda^k(N,x) \,.\, \lambda_{-1}(N) \,.\, (1-L^k) t^k}{1 + \sum_{k\geq 1} \lambda^k(N,x) \,.\, \lambda_{-1}(N) \,.\, L^k t^k}
\end{aligned}
$$

Source: check/findings/findings_341_345.json#10; check/findings/findings_341_345.json#11; check/findings/findings_341_345.json#13. check/findings/findings_341_345.json#10: The native fraction-stop zoom resolves the short white gap between the bar and the round ink mark; the main detail corroborates it. No comma tail is visible. The paired diplomatic full stop and current comma preserve the source punctuation and the continuing coordinated clause respectively. check/findings/findings_341_345.json#11: Main fourth strip and native353 lower resolve exactly two denominator dots, against the three earlier ellipses. The formal denominator is the full series lambda_(Lt)(xD)=1+sum_{k>=1}lambda^k(xD)L^k t^k; its complete three-dot notation in current represents this same coefficientwise series. Retain the literal two-dot diplomatic ending. check/findings/findings_341_345.json#13: Both scans resolve the two dots and their separation. The denominator is exactly lambda_(Lt)(xD)=1+sum_{k>=1}lambda^k(N,x)D L^k t^k by5.3.1, so there is one multiplication of D and L^k. Preserve both source dots in diplomatic and the single product in current.

[247] EN-CONTRIB-V-345-01 Source folio 345. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=359; scan/SGA6_witness_300dpi.djvu#page=356
Printed reading:

$$(1+\hat{.}^{+})\times(1+\hat{..}^{+})\xrightarrow{*}1+\hat{.}^{+},$$

Corrected reading:

$$(1+\hat{.}^{+})\times(1+\hat{.}^{+})\xrightarrow{*}1+\hat{.}^{+},$$

Source: check/findings/findings_341_345.json#21. Both main359 upper strip and native356 top show the two detached low marks. The operation is the natural component I(A)×I(A)→I(A) constructed at the end of344;6.1.1 evaluates its two inputs at1+X and1+Y in the same completed graded algebra. Thus the second factor has the same single object-placeholder as the first, justifying removal of only its duplicated dot in the corrected form.

[248] EN-CONTRIB-V-345-02 Source folio 345. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=359; scan/SGA6_witness_300dpi.djvu#page=356
Printed reading: $\forall\, a, a' \in A$
Corrected reading: $\forall\, a, a' \in K$
Source: check/findings/findings_341_345.json#23. The source first defines the ring E(A)=K×I(A), whose projection pi:E(A)→K sends(a,phi) to a; the scalar inclusion is j(a)=(a,1), and pi∘j=id_K. The printed A is related to the required K by its structure homomorphism iota_A:K→A_0, but the first coordinate of E(A) is still the element of K itself. For A=K[T] with T of degree1, the graded inclusion sends constants to degree0, and T is not an allowed first coordinate. Formula6.1.3 is exactly j(a)(a-prime,phi)=(aa-prime,phi^a). Therefore change only this quantifier to a,a-prime in K in the corrected layer. Full maps are recorded in identification_345.json.

[249] EN-CONTRIB-V-345-03 Source folio 345. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=359; scan/SGA6_witness_300dpi.djvu#page=356
Printed reading: admits (1.1) as its unit
Corrected reading: admits (1,1) as its unit
Source: check/findings/findings_341_345.json#25. The native lower crop resolves a low period between the two ones, with no descending comma tail. For I(A)=1+completed(A)^+ with its displayed ring law, its additive zero is the series1, hence 1*phi=phi*1=1. Formula6.1.2 gives (1,1)(a,phi)=(a,1^a phi^1(1*phi))=(a,phi), and the right product is identical. Thus the corrected pair is exactly(1,1). The scalar homomorphism is j:K→K×I(A),a↦(a,1); the projection pi(a,phi)=a satisfies pi∘j=id_K, and ker(pi)={(0,phi)}.

[250] EN-CONTRIB-V-346-01 Source folio 346. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=360; scan/SGA6_witness_300dpi.djvu#page=357
Printed reading: $\widehat{G}\longrightarrow K\times(1+\mathrm{Gr}\ .^{+})$
Corrected reading: $\widehat{G}\longrightarrow K\times(1+\widehat{\mathrm{Gr}}\ .^{+})$
Source: check/findings/findings_346_350.json#3. The canonical map of1.11 is q_A:1+sum a_n t^n↦1+sum[a_n]_n, where[a_n]_n lies in Gr^nA. Its target is the degree product, as defined on308–309. The inclusion into the augmented ring is phi↦(0,phi). For A=K[T] with its degree filtration, q_A(1+sum_(n>=1)T^n t^n)=1+sum_(n>=1)T^n, an element of the completed product. With K=Z its infinitely many nonzero graded coefficients cannot lie in the direct sum: every element of the direct sum has finite degree support. The next display explicitly writes the same completion hat. Retain the source omission diplomatically and its supplied restoration in current.

[251] EN-CONTRIB-V-346-02 Source folio 346. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=360; scan/SGA6_witness_300dpi.djvu#page=357
Printed reading: defines a morphism of functors on :
Corrected reading: defines a morphism of functors on $\mathcal{G}$:
Source: check/findings/findings_346_350.json#5. Native357 top and the main second strip show no category ink in this space. In1.11, j:G→C gives the degree filtration and epsilon:Id_G→Gr∘j identifies each homogeneous component with its associated graded class. Lemma1.12 sends u:I→H on G to u_Gr∘q on C and is bijective. Applied here to H=G-hat^u∘i∘(K×I), it produces precisely the displayed descended morphism on G. Insert only that category letter in the corrected layer; retain the observed blank diplomatically.

[252] EN-CONTRIB-V-346-03 Source folio 346. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=360; scan/SGA6_witness_300dpi.djvu#page=357
Printed reading:

$$1 +\hat{\cdot}^{+} \overset{'}{\to} \widehat{G}^{u} \circ i \circ (K \times (1 +\hat{\cdot}^{+})).$$

Corrected reading:

$$1 +\hat{\cdot}^{+} \overset{\gamma'}{\to} \widehat{G}^{u} \circ i \circ (K \times (1 +\hat{\cdot}^{+})).$$

Source: check/findings/findings_346_350.json#6. Both main and native357 top show a blank to the left of the prime. The first display labels the morphism gamma-prime; the second is its unique descent through q by1.12, with the j-th coordinate of gamma-prime^i given immediately afterwards by Q-double-prime_(i,j). The exact relation is descended_gamma-prime_Gr(A)∘q_A=H(q)_A∘gamma-prime_A. Thus restoring gamma before this same prime preserves the named transformation and its coefficients.

[253] EN-CONTRIB-V-348-01 Source folio 348. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=362; scan/SGA6_witness_300dpi.djvu#page=359
Printed reading:

$$\begin{aligned} \forall a \in K, \quad \mathrm{ch}(1+X)^{a}) &= \mathrm{ch}(\exp(a\,\mathrm{Log}(1+X))) \\ &= \eta(a\,\mathrm{Log}(1+X)) \\ &= a.\eta(\mathrm{Log}(1+X)) \\ &= a.\,\mathrm{ch}(1+X). \end{aligned}$$

Corrected reading:

$$\begin{aligned} \forall a \in K, \quad \mathrm{ch}((1+X)^{a}) &= \mathrm{ch}(\exp(a\,\mathrm{Log}(1+X))) \\ &= \eta(a\,\mathrm{Log}(1+X)) \\ &= a.\eta(\mathrm{Log}(1+X)) \\ &= a.\,\mathrm{ch}(1+X). \end{aligned}$$

Source: check/findings/findings_346_350.json#12. Main bottom strip and native359 lower resolve the exact unmatched delimiter. The following argument is exp(a Log(1+X))=(1+X)^a, so the map being applied is ch to the complete powered series. Under ch=eta∘Log the exact chain is ch((1+X)^a)=eta(a Log(1+X))=a eta(Log(1+X))=a ch(1+X). The existing corrected opening parenthesis gives precisely this map composition. In particular a=0 gives ch(1)=0, whereas interpreting the printed exponent outside ch would give ch(1+X)^0=1 in the completed rational ring.

[254] EN-CONTRIB-V-349-01 Source folio 349. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=363; scan/SGA6_witness_300dpi.djvu#page=360
Printed reading: $\widehat{A}^{\geq n} = \prod_{i \geq n} A^i$, We have the following congruences
Corrected reading: $\widehat{A}^{\geq n} = \prod_{i \geq n} A^i$. We have the following congruences
Source: check/findings/findings_346_350.json#14. Both main second/third strips and native360 top show the small descending mark. The diplomatic comma and corrected sentence-ending period preserve the physical source and the next complete sentence respectively.

[255] EN-CONTRIB-V-349-02 Source folio 349. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=363; scan/SGA6_witness_300dpi.djvu#page=360
Printed reading:

$$\mathrm{ch}(\varphi) = \frac{(-1)^{n-1}}{(n-1)!}.a_n + .. \; ; \tag{6.5.2}$$

Corrected reading:

$$\mathrm{ch}(\varphi) = \frac{(-1)^{n-1}}{(n-1)!}.a_n + \dots \; ; \tag{6.5.2}$$

Source: check/findings/findings_346_350.json#15. Both scans resolve exactly two dots. The omitted terms are the higher-degree components of ch(phi)=eta(Log phi), with degree-n coefficient(-1)^(n-1)a_n/(n-1)!. Thus the current three-dot ellipsis denotes the same full higher-degree tail, while diplomatic retains the two physical dots.

[256] EN-CONTRIB-V-349-03 Source folio 349. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=363; scan/SGA6_witness_300dpi.djvu#page=360
Printed reading:

$$\varphi * \psi = 1 - \frac{(m+n-1)!}{(m-1)!(n-1)!}.a_m b_n + .., \tag{6.5.3}$$

Corrected reading:

$$\varphi * \psi = 1 - \frac{(m+n-1)!}{(m-1)!(n-1)!}.a_m b_n + \dots, \tag{6.5.3}$$

Source: check/findings/findings_346_350.json#16. The main third strip and native360 lower resolve exactly two tail dots. These denote terms of degree greater than m+n after the displayed leading product coefficient. Retain diplomatic two dots and the current complete ellipsis, whose low placement is separately restored.

[257] EN-CONTRIB-V-350-01 Source folio 350. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=364; scan/SGA6_witness_300dpi.djvu#page=361
Printed reading: $\mathrm{ch}(\varphi) \equiv \mathrm{ch}(1 + a_n) \bmod. \widehat{A}^{\geq n+1}$
Corrected reading: $\mathrm{ch}(\varphi) \equiv \mathrm{ch}(1 + a_n) \bmod. \widehat{A}^{\geq n+1}_{\mathbb{Q}}$
Source: ASTRA-V-CH-MODULUS-350. Both scans clearly omit Q at this occurrence. The map ch is valued in the degree completion of A_Q=A tensor_ZQ, and eta scales degree r by the rational c_r. For K=Z,A=Z[Y] with wtY=3,n=1,phi=1+Y,a1=0, the degree-three component of ch(phi)-ch(1+a1) is Y/2. The coefficient map [Y]:completed Q[Y]→Q sends the integral completed subgroup to Z and this difference to1/2, so it is not in completed A^(>=2). It is in the rational completed degree>=2 subgroup, exactly the target filtration of ch. The packet records the scalar-extension and homogeneous projection maps.

[258] EN-CONTRIB-V-350-02 Source folio 350. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=364; scan/SGA6_witness_300dpi.djvu#page=361
Printed reading:

$$1+\frac{(-1)^{p-1}}{(p-1)!}.Z_p+\ldots=\left(1+\frac{(-1)^{m-1}}{(m-1)!}.X_m+\ldots\right)\left(1+\frac{(-1)^{n-1}}{(n-1)!}.Y_n+..\right),$$

Corrected reading:

$$\frac{(-1)^{p-1}}{(p-1)!}.Z_p+\ldots=\left(\frac{(-1)^{m-1}}{(m-1)!}.X_m+\ldots\right)\left(\frac{(-1)^{n-1}}{(n-1)!}.Y_n+\ldots\right),$$

Source: ASTRA-V-CH-CONSTANTS-350; check/findings/findings_346_350.json#18. ASTRA-V-CH-CONSTANTS-350: Main second strip and native361 top confirm all three prefixes. On348, ch(phi)=eta(Log phi), with eta scaling homogeneous degree r by c_r=(-1)^(r-1)/(r-1)!, so ch(phi) has constant0. On349, the extension is ch(a,phi)=a+eta(Log phi), and the degree-n filtration for n>=1 has first coordinate0. Therefore the exact identity ch(phi*psi)=ch(phi)ch(psi) expands as c_pZ_p+...=(c_mX_m+...)(c_nY_n+...). In K=Z,A=Z[X,Y], wtX=wtY=1, phi=1+X,psi=1+Y,6.1.1 gives phi*psi=1-XY+higher, so the printed left side has degree-one component0 while the printed right side has X+Y. The exact comparison with unit pairs and the preceding source definitions is recorded in identification_350.json. check/findings/findings_346_350.json#18: Both scans resolve the three different tails. The current complete ellipsis denotes the same higher-degree terms of ch(psi); diplomatic retains the physical two-dot ending. The separate three constant-term errors are decided in ASTRA-V-CH-CONSTANTS-350.

[259] EN-CONTRIB-V-350-03 Source folio 350. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=364; scan/SGA6_witness_300dpi.djvu#page=361
Printed reading:

$$\begin{aligned}
&\text{i)} \quad \lambda^i(x) = 0 \quad \underline{\text{for}}\ i \geq n\ ; \\
&\text{ii)} \quad \lambda^n(x) = (1, 1 + a_1) \\
&\text{iii)} \quad \sum_{i=0}^{n} (-1)^i \lambda^i(x) = 0, 1 - (n-1)! a_n + \ldots),
\end{aligned}$$

Corrected reading:

$$\begin{aligned}
&\text{i)} \quad \lambda^i(x) = 0 \quad \underline{\text{for}}\ i > n\ ; \\
&\text{ii)} \quad \lambda^n(x) = (1, 1 + a_1) \\
&\text{iii)} \quad \sum_{i=0}^{n} (-1)^i \lambda^i(x) = (0, 1 - (n-1)! a_n + \ldots),
\end{aligned}$$

Source: check/findings/findings_346_350.json#19; check/findings/findings_346_350.json#21. check/findings/findings_346_350.json#19: Native361 lower and the main third strip resolve the lower stroke, unlike the plain > at the statement end. In the universal symmetric-polynomial embedding X_i↦e_i(T1,...,Tn), the element becomes the sum of the n line elements L_j=(1,1+T_j). By6.2.1 and lambda-additivity, lambda_t(x)=product_(j=1)^n(1+L_j t), so coefficients vanish above n and the degree-n coefficient is product L_j=(1,1+sum T_j)=(1,1+a1). At n=1,lambda^1(L1)=L1 has scalar coordinate1 and is nonzero over K=Z, directly contradicting the printed endpoint. Retain current i>n, with no change of n or index. check/findings/findings_346_350.json#21: The main/native images resolve the unmatched close. The sum of the lambda terms is an element of K×I(A); in the displayed positive-degree case its scalar coordinate is the alternating binomial sum0, and the second coordinate starts1-(n-1)!a_n. The existing current opening parenthesis gives exactly this ordered pair. The coefficient follows from ch(product_j(1-L_j))=product_j(1-exp T_j), whose degree-n part is(-1)^n product T_j; dividing by c_n=(-1)^(n-1)/(n-1)! yields-(n-1)!a_n.

[260] EN-CONTRIB-V-351-01 Source folio 351. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=365; scan/SGA6_witness_300dpi.djvu#page=362
Printed reading: $\sum_{i=1}^{n}(-1)^i\lambda^i(x) = \lambda_{-1}(x) = \prod_{i=1}^{n}(1-(1,1+T_i)) = (-1)^n.\prod_{i=1}^{n}(0,1+T_i)$.
Corrected reading:

$$\sum_{i=0}^{n}(-1)^i\lambda^i(x) = \lambda_{-1}(x) = \prod_{i=1}^{n}(1-(1,1+T_i)) = (-1)^n.\prod_{i=1}^{n}(0,1+T_i).$$

Source: check/findings/findings_351_355.json#6. Both scans explicitly show i=1. The lambda-series identity is lambda_t(x)=product_(j=1)^n(1+L_j t), L_j=(1,1+T_j); evaluation t=-1 includes the constant coefficient lambda^0(x)=(1,1). Thus the alternating sum begins at0. Under the scalar projection pi:Ch(A)→K, the printed sum for positive n maps to sum_(i=1)^n(-1)^i binom(n,i)=-1, while lambda_(-1)(x) maps to(1-1)^n=0. In the concrete case n=1,K=Z this is already-1 versus0. Retain the existing current0 and diplomatic1. The symmetric-polynomial map X_i↦e_i(T1,...,Tn) is injective: ordered monomials in the e_i have distinct leading monomials with leading coefficient1; their coefficient comparison is unchanged after base extension to K. The induced Ch map is id_K on the scalar factor and the product of the graded injections on the completed ideal factor.

[261] EN-CONTRIB-V-351-02 Source folio 351. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=365; scan/SGA6_witness_300dpi.djvu#page=362
Printed reading: $\gamma^i(0,\varphi) = (0, 1 + Q''_{i,1} + \ldots + Q''_{i,j} + ..)$
Corrected reading: $\gamma^i(0,\varphi) = (0, 1 + Q''_{i,1} + \ldots + Q''_{i,j} + \ldots)$
Source: ASTRA-V-TWO-DOT-351. Main third strip and native362 lower clearly separate the two final dots before the closing parenthesis, while the earlier three-dot tail is visible in the same line. The full formal series is1+sum_(j>=1)Q-double-prime_(i,j); both abbreviations refer to that same higher-j tail. Current retains the complete three-dot notation.

[262] EN-CONTRIB-V-352-01 Source folio 352. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=366; scan/SGA6_witness_300dpi.djvu#page=363
Printed reading: $\gamma^n(x - \epsilon(x)) = (-1)^n \sum_{i=0}^{n} (-1)^i \lambda(x + n - \epsilon(x))$
Corrected reading: $\gamma^n(x - \epsilon(x)) = (-1)^n \sum_{i=0}^{n} (-1)^i \lambda^i(x + n - \epsilon(x))$
Source: check/findings/findings_351_355.json#14. Both scans have lambda without an exponent after(-1)^i. Put z=x-epsilon(x). The stated definition is gamma^n(z)=lambda^n(z+n-1). Since lambda_t(z+n)=(1+t)lambda_t(z+n-1), comparison of coefficients and the finite alternating sum gives sum_(i=0)^n(-1)^i lambda^i(z+n)=(-1)^n lambda^n(z+n-1): adjacent lower coefficients cancel pairwise. Thus the missing superscript is exactly i. The bounds0,n and the sign(-1)^n stay unchanged.

[263] EN-CONTRIB-V-353-01 Source folio 353. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=367; scan/SGA6_witness_300dpi.djvu#page=364
Printed reading: $c : A \longrightarrow \mathrm{Ch}(\mathrm{Gr}\, A)$
Corrected reading: $\widetilde{c} : A \longrightarrow \mathrm{Ch}(\mathrm{Gr}\, A)$
Source: check/findings/findings_351_355.json#22. The statement has plain c in both scans, while6.7.2 and the next sentence have an explicit tilde. The defined pair map is tilde-c:A→K×I(Gr A),x↦(epsilon(x),c(x)). Its first projection is epsilon and its second projection is c. The ideal inclusion i:I(Gr A)→Ch(Gr A),phi↦(0,phi), would instead have zero first projection; for x=1 its value is(0,1), while the completed map has value(1,1). Thus the augmentation-compatible map asserted in6.8 is exactly the already defined pair map tilde-c. Retain the printed plain c diplomatically and the supplied current tilde; no mathematical map is changed by this name repair.

[264] EN-CONTRIB-V-354-01 Source folio 354. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=368; scan/SGA6_witness_300dpi.djvu#page=365
Printed reading:

$$\begin{array}{ccc} A & \xrightarrow{\gamma_t} & \widehat{G}_g(A) \\ \downarrow{\scriptstyle \widetilde{c}} & & \downarrow{\scriptstyle \widehat{G}_g(\widetilde{c})} \\ \mathrm{Ch}\, A & \xrightarrow{\gamma_t} & \widehat{G}_g(\mathrm{Ch}\, A) \end{array}$$

Corrected reading:

$$\begin{array}{ccc} A & \xrightarrow{\gamma_t} & \widehat{G}_g(A) \\ \downarrow{\scriptstyle \widetilde{c}} & & \downarrow{\scriptstyle \widehat{G}_g(\widetilde{c})} \\ \mathrm{Ch}(\mathrm{Gr}\, A) & \xrightarrow{\gamma_t} & \widehat{G}_g(\mathrm{Ch}(\mathrm{Gr}\, A)) \end{array}$$

Source: ASTRA-V-DIAGRAM-TARGET-354. Both source lower objects literally read Ch A and Ghat_g(Ch A). The left arrow is the map defined in6.7.2 and asserted in6.8:tilde-c:A→Ch(Gr A),x↦(epsilon(x),c(x)),where the degree-i coefficient c^i(x) is the class of gamma^i(x-epsilon(x)) in Fil^i A/Fil^(i+1)A. Thus its actual codomain is K×(1+product_(i>=1)Gr^i A). The functor Ghat_g applied to this map acts coefficientwise:1+sum b_i t^i↦1+sum tilde-c(b_i)t^i,with target Ghat_g(Ch(Gr A)). The bottom gamma_t has source Ch(Gr A). In these exact objects the equality of the two routes is the coefficient identity tilde-c(gamma^i x)=gamma^i(tilde-c x),which is precisely the compatibility the paragraph is proving. Definition6.2 takes a graded input;here its functor is precomposed with Gr. Restore Gr A in both lower objects in current only;all four arrow labels and directions remain unchanged.

[265] EN-CONTRIB-V-354-02 Source folio 354. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=368; scan/SGA6_witness_300dpi.djvu#page=365
Printed reading: $a \in A\ \epsilon(x_i) = 0$
Corrected reading: $a \in A,\ \epsilon(x_i) = 0$
Source: check/findings/findings_351_355.json#28. Both scans end the source line with a in A and show no comma before the next-line epsilon(x_i)=0. These are successive constraints on the same generator product:the coefficient a ranges in A,the x_i have augmentation0,and the index sum is at least m. The current comma separates the first and second constraints without altering that exact tuple of data or its generator map(a,(x_i),(i_i))→a product gamma^(i_i)(x_i). Retain the diplomatic omission and existing corrected comma.

[266] EN-CONTRIB-V-355-01 Source folio 355. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=369; scan/SGA6_witness_300dpi.djvu#page=366
Printed reading: $z_{\alpha,j}\ R$
Corrected reading: $x_{\alpha,j} \in R$
Source: check/findings/findings_351_355.json#31. Both scans print z_(alpha,j),then R with no membership symbol. The immediately following generator expression and its augmentation conditions use x_(alpha,j),and the same family is used as the target of the universal generators X_(alpha,j). The index-preserving renaming z_(alpha,j)↦x_(alpha,j) identifies this one family;the generator-evaluation map sends X_(alpha,j) to that element of R. Thus the existing current x_(alpha,j) in R restores both the family label and its missing relation. The coefficient family a_alpha in K and every index remain unchanged.

[267] EN-CONTRIB-V-356-01 Source folio 356. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=370; scan/SGA6_witness_300dpi.djvu#page=367
Printed reading: $N_0' = N_0 + d$
Corrected reading: $N_0 = N_0' + d$
Source: check/findings/findings_356_360.json#1. Both scans put the prime on the left N in N-prime_0=N0+d. The preceding universal generator N-prime_0 has gamma-degree at most d. Translation N0=N-prime_0+d has inverse N-prime_0=N0-d,and gives lambda_t(N0)=(1+t)^d gamma_(t/(1+t))(N-prime_0)=sum_(j=0)^d gamma^j(N-prime_0)t^j(1+t)^(d-j). Its degree is at most d,and evaluation at-1 gives lambda_(-1)(N0)=(-1)^d gamma^d(N-prime_0),exactly the identity printed below. The final prescribed map f(N-prime_0)=N-d then gives f(N0)=N. With the printed reversed relation it would instead give f(N0)=N-2d. At d=1 and N-prime_0 specialized to0,the printed relation yields N0=-1 with lambda^2(-1)=1;the corrected relation yields N0=1. Thus current restores precisely N0=N-prime_0+d;the separate final augmentation/filtration defect is not thereby repaired.

[268] EN-CONTRIB-V-356-02 Source folio 356. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=370; scan/SGA6_witness_300dpi.djvu#page=367
Printed reading: X_0 to X
Corrected reading: X_0 to x
Source: fixes/FIXLOG_6_CURRENT.jsonl#415. Both scans print capital X as the image of X0. The preceding355 definition is X0=sum_alpha a_alpha product_j gamma^(i_j)(X_(alpha,j)),and the stated map sends each X_(alpha,j) to x_(alpha,j) while fixing a_alpha in K. Applying the lambda-map term by term gives sum_alpha a_alpha product_j gamma^(i_j)(x_(alpha,j))=x by the displayed decomposition on355. Thus the exact target element is the already specified lowercase x;retain the existing current correction and diplomatic capital.

[269] EN-CONTRIB-V-357-01 Source folio 357. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=371; scan/SGA6_witness_300dpi.djvu#page=368
Printed reading:

$$\forall\, i > 1, \quad \forall\, a_i \in A_i, \qquad u(a_i) = \mathrm{cl}_{\mathrm{Gr}^i}\left(0, 1 + (-1)^{i-1}(i-1)!a_i\right),$$

Corrected reading:

$$\forall\, i \geq 1, \quad \forall\, a_i \in A_i, \qquad u(a_i) = \mathrm{cl}_{\mathrm{Gr}^i}\left(0, 1 + (-1)^{i-1}(i-1)!a_i\right),$$

Source: check/findings/findings_356_360.json#4. Both scans show strict i>1 in the first formula and i>=1 in the immediately following inverse definition. In degree i,the stated map is u_i(a)=[(0,1+C_i a)],C_i=(-1)^(i-1)(i-1)!,and its inverse extracts a_i and multiplies by C_i inverse. For i=1 both coefficients are1,so u_1(a)=[(0,1+a)] is required by the displayed inverse as well as by the declared domain A. The example A=Q[Y]/(Y^2),degree Y=1,has nonzero A1 and no higher positive degree;the printed i>1 clause does not define u on Y. The current i>=1 supplies exactly this missing component;all other formulas and the degree0 identity remain unchanged.

[270] EN-CONTRIB-V-357-02 Source folio 357. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=371; scan/SGA6_witness_300dpi.djvu#page=368
Printed reading: $\wedge \to \mathrm{Ch}(\mathrm{Gr}\wedge))$
Corrected reading: $\wedge \to \mathrm{Ch}(\mathrm{Gr}\wedge)$
Source: check/findings/findings_356_360.json#5. Both scans show two closing parentheses after the algebra-name glyph in the introductory completed-class target. Its displayed syntax has one opening parenthesis after Ch and no opening parenthesis after Gr,so one closing parenthesis is unmatched. The target map is exactly tilde-c:x↦(epsilon(x),c(x)) into Ch(Gr of the named ring);removing the unmatched close preserves that target and every coordinate. Retain the existing current correction and literal diplomatic double close.

[271] EN-CONTRIB-V-358-01 Source folio 358. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=372; scan/SGA6_witness_300dpi.djvu#page=369
Printed reading:

$$\begin{aligned}
&\text{i)} \quad c^k(x+y) = \sum_{i+j=k} c^i(x)c^j(y), \quad \text{for } k \geq 1,\ x, y \in \mathrm{Fil}^1 \wedge\ ;\\
&\text{ii)} \quad c^k(xy) = P''_k(c^i(x), c^j(y)) \quad \text{(cf. 6.1)}, \quad \text{for } k \geq 1,\ x, y \in \mathrm{Fil}^1 \wedge\ ;\\
&\text{iii)} \quad c^k(ax) = R_{k,a}(c^i(x) \quad \text{(cf. 6.1)}, \quad \text{for } k \geq 1,\ a \in K,\ x \in \mathrm{Fil}^1 \wedge\ ;\\
&\text{iv)} \quad c^k(\gamma^j(x)) = Q''_{j,k}(c^i(x)) \quad \text{(cf. 6.1)}, \quad \text{for } j, k \geq 1,\ x \in \mathrm{Fil}^1 \wedge,
\end{aligned}$$

Corrected reading:

$$\begin{aligned}
&\text{i)} \quad c^k(x+y) = \sum_{i+j=k} c^i(x)c^j(y), \quad \text{for } k \geq 1,\ x, y \in \mathrm{Fil}^1 \wedge\ ;\\
&\text{ii)} \quad c^k(xy) = P''_k(c^i(x), c^j(y)) \quad \text{(cf. 6.1)}, \quad \text{for } k \geq 1,\ x, y \in \mathrm{Fil}^1 \wedge\ ;\\
&\text{iii)} \quad c^k(ax) = R_{k,a}(c^i(x)) \quad \text{(cf. 6.1)}, \quad \text{for } k \geq 1,\ a \in K,\ x \in \mathrm{Fil}^1 \wedge\ ;\\
&\text{iv)} \quad c^k(\gamma^j(x)) = Q''_{j,k}(c^i(x)) \quad \text{(cf. 6.1)}, \quad \text{for } j, k \geq 1,\ x \in \mathrm{Fil}^1 \wedge,
\end{aligned}$$

Source: check/findings/findings_356_360.json#13. Both scans show only one closing parenthesis in R_(k,a)(c^i(x),which closes the argument of c^i but leaves the outer R application open. The evaluated polynomial map is R_(k,a):the coefficient list(c^1(x),...,c^k(x))↦c^k(ax),as defined by the scalar-operation coefficient map6.1. Closing that outer application adds exactly the omitted delimiter without changing its inputs,the scalar a or its value. Retain the existing current close and the diplomatic omission.

[272] EN-CONTRIB-V-360-01 Source folio 360. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=374; scan/SGA6_witness_300dpi.djvu#page=371
Printed reading:

$$\begin{array}{ccc} \wedge & \xrightarrow{\widetilde{c_0}} & \mathrm{Ch}(\Omega) \\ \downarrow{\scriptstyle \widetilde{c}} & & \downarrow{\scriptstyle \widetilde{c}} \\ \mathrm{Ch}(\mathrm{Gr}\wedge)) & \xrightarrow{\mathrm{Ch}(\widetilde{c_0})} & \mathrm{Ch}(\mathrm{Gr}_\lambda(\mathrm{Ch}(\Omega))); \end{array}$$

Corrected reading:

$$\begin{array}{ccc} \wedge & \xrightarrow{\widetilde{c_0}} & \mathrm{Ch}(\Omega) \\ \downarrow{\scriptstyle \widetilde{c}} & & \downarrow{\scriptstyle \widetilde{c}} \\ \mathrm{Ch}(\mathrm{Gr}\wedge) & \xrightarrow{\mathrm{Ch}(\widetilde{c_0})} & \mathrm{Ch}(\mathrm{Gr}_\lambda(\mathrm{Ch}(\Omega))); \end{array}$$

Source: check/findings/findings_356_360.json#21. Both scans resolve the two distinct closing strokes. The source functor application is Ch applied to the graded ring Gr(wedge),also written correctly in the triangle and preceding sentence. Current removes only the unmatched close;the object and arrow remain unchanged.

[273] EN-CONTRIB-V-360-02 Source folio 360. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=374; scan/SGA6_witness_300dpi.djvu#page=371
Printed reading:

$$\begin{aligned} \frac{d}{dt}(\lambda_t(x+y))/\lambda_t(x+y) &= \frac{d}{dt}(\lambda_t(x)\lambda_t(y))/\lambda_t(x)\lambda_t(y) \\ &= \frac{d}{dt}(\lambda_t(x))/\lambda_t(x) + \frac{d}{dt}(\lambda_t(y))/\lambda_t(y)\ ; \end{aligned}$$

Corrected reading:

$$\begin{aligned} \frac{d}{dt}(\lambda_t(x+y))/\lambda_t(x+y) &= \frac{d}{dt}(\lambda_t(x)\lambda_t(y))/(\lambda_t(x)\lambda_t(y)) \\ &= \frac{d}{dt}(\lambda_t(x))/\lambda_t(x) + \frac{d}{dt}(\lambda_t(y))/\lambda_t(y)\ ; \end{aligned}$$

Source: ASTRA-V-SLASH-PRODUCT-360. Write a=lambda_t(x),b=lambda_t(y) in K[[t]]. Both have constant1,so their inverses exist. The defining pre-lambda identity is lambda_t(x+y)=ab. Formal differentiation and multiplication by (ab)^-1 give (ab)prime/(ab)=a-prime/a+b-prime/b. The current parentheses express exactly that identity. In the binomial ring Z with x=y=1,a=b=1+t:the stated quotient is2/(1+t),whereas evaluating the ungrouped slash successively from the left gives2(1+t);their t-coefficients are-2 and2. No sign,index,or derivative argument changes.

[274] EN-CONTRIB-V-362-01 Source folio 362. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=376; scan/SGA6_witness_300dpi.djvu#page=373
Printed reading:

$$\frac{d}{dt}(\lambda_u(1+Xt))/\lambda_u(1+Xt)) = \frac{1+Xt}{\underline{1}+\{1+Xt\}u} = \sum_{k=1}^{\infty}(-1)^{k-1}\{1+Xt\}^{\circ k}u^{k-1}.$$

Corrected reading:

$$\frac{d}{du}(\lambda_u(1+Xt))/\lambda_u(1+Xt) = \frac{1+Xt}{\underline{1}+\{1+Xt\}u} = \sum_{k=1}^{\infty}(-1)^{k-1}\{1+Xt\}^{\circ k}u^{k-1}.$$

Source: check/findings/findings_361_365.json#7; check/findings/findings_361_365.json#6. check/findings/findings_361_365.json#7: Put R=Ghat(Z[X]) and a=1+Xt in R. Its unit is underlined1=1+t,its addition is multiplication of the represented t-series,and its product is circle. The line-element identity is lambda_u(a)=underlined1+a u in R[[u]]. The formal derivation D_u(sum r_j u^j)=sum j r_j u^(j-1) sends this lambda-series to a. Multiplication by its inverse gives a/(underlined1+a u)=sum_(k>=1)(-1)^(k-1)a^(circle k)u^(k-1). The product law(1+Xt)circ(1+Yt)=1+XYt identifies a^(circle k)=1+X^k t,which is exactly7.4.1. Thus u is the differentiation variable required by the displayed coefficient map;the printed dt is preserved only in diplomatic. check/findings/findings_361_365.json#6: Both native373 top and main first strip resolve the final handwritten closing stroke. Diplomatic already retains it and current already removes it. There is only one denominator application lambda_u(a),so the extra close has no matching opening.

[275] EN-CONTRIB-V-362-02 Source folio 362. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=376; scan/SGA6_witness_300dpi.djvu#page=373
Printed reading: if the ψ^t satisfy these identities
Corrected reading: if the ψ^k satisfy these identities
Source: check/findings/findings_361_365.json#9. Both sources resolve the ascender/cross-stroke as t. Existing diplomatic psi^t/current psi^k is confirmed. The identities7.5.1 concern the family k↦psi^k indexed by positive integers;the substitution of the family index k for this isolated t refers to those same three identities and does not change the formal series variable t elsewhere.

[276] EN-CONTRIB-V-362-03 Source folio 362. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=376; scan/SGA6_witness_300dpi.djvu#page=373
Printed reading: $\frac{d}{dt}(\lambda_t(1))/\lambda_t(1) = 1/1+t = \sum_{k=1}^{\infty}(-1)^{k-1}t^{k-1}$
Corrected reading: $\frac{d}{dt}(\lambda_t(1))/\lambda_t(1) = 1/(1+t) = \sum_{k=1}^{\infty}(-1)^{k-1}t^{k-1}$
Source: check/findings/findings_361_365.json#12. The source has the unparenthesized solidus1/1+t. The immediately stated lambda_t(1)=1+t gives D_t(lambda_t(1))=1 and hence the formal quotient1/(1+t). Multiplying the following series sum_(k>=1)(-1)^(k-1)t^(k-1) by1+t gives1 coefficientwise. The ungrouped left-associated1/1+t instead has coefficient+1 at t,whereas the displayed series has-1. Current therefore groups only the denominator,as in the other reviewed solidus repairs;diplomatic retains the literal print.

[277] EN-CONTRIB-V-363-01 Source folio 363. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=377; scan/SGA6_witness_300dpi.djvu#page=374
Printed reading:

$$\Phi^k(1+Xt) = X^k \qquad ((k-1)\text{-st coefficient of } X/1+Xt)\ ; \qquad \Phi(1+Yt) = Y^k\ ;$$

Corrected reading:

$$\Phi^k(1+Xt) = X^k \qquad ((k-1)\text{-st coefficient of } (-1)^{k-1}X/(1+Xt))\ ; \qquad \Phi^k(1+Yt) = Y^k\ ;$$

Source: ASTRA-V-COEFFICIENT-363; check/findings/findings_361_365.json#16. ASTRA-V-COEFFICIENT-363: The page first defines Phi^k(f)=(-1)^(k-1)[t^(k-1)](f-prime/f). For f=1+Xt,formal differentiation gives f-prime/f=X/(1+Xt)=sum_(j>=0)(-1)^j X^(j+1)t^j. Its(k-1)-st coefficient is(-1)^(k-1)X^k. Multiplication by the sign from the definition gives X^k. Equivalently,the coefficient of t^(k-1) in(-1)^(k-1)X/(1+Xt) is exactly X^k,so inserting that factor inside the parenthetical coefficient-series repairs only the omitted sign. The denominator parentheses group the same formal inverse;the source solidus is retained. At k=2 in Z[X],the unsigned source coefficient is-X^2,whereas Phi^2(1+Xt)=X^2. No coefficient number or exponent is changed. check/findings/findings_361_365.json#16: Native374 top confirms the blank superscript position. Existing diplomatic Phi/current Phi^k is retained. The map defined in the first paragraph is Phi^k(f)=(-1)^(k-1)[t^(k-1)](f-prime/f). Substituting f=1+Yt gives Y^k,identifying this occurrence with the same indexed map as its immediately preceding X formula.

[278] EN-CONTRIB-V-363-02 Source folio 363. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=377; scan/SGA6_witness_300dpi.djvu#page=374
Printed reading:

$$\begin{array}{ccc} \widehat{G}^t(K) & \xrightarrow{\lambda_u} & \widehat{G}^u \circ \widehat{G}^t(K) \\ {\scriptstyle \Phi^{kk'}}\downarrow & & \downarrow{\scriptstyle \Phi_u^{k'}} \\ K & \xrightarrow{\Phi^k} & \widehat{G}^t(K), \end{array} \tag{7.5.3}$$

Corrected reading:

$$\begin{array}{ccc} \widehat{G}^t(K) & \xrightarrow{\lambda_u} & \widehat{G}^u \circ \widehat{G}^t(K) \\ {\scriptstyle \Phi^{kk'}}\downarrow & & \downarrow{\scriptstyle \Phi_u^{k'}} \\ K & \xleftarrow{\Phi^k} & \widehat{G}^t(K), \end{array} \tag{7.5.3}$$

Source: check/findings/findings_361_365.json#19. The first paragraph defines Phi^k:Ghat^t(K)→K by f↦(-1)^(k-1)[t^(k-1)](f-prime/f). Hence the lower-right object is its domain and the lower-left K its codomain. More explicitly,the upper/right route sends1+Xt to lambda_u(1+Xt) and then to(1+Xt)^(circle k-prime)=1+X^(k-prime)t. Applying Phi^k gives X^(kk-prime),which is the value of the left vertical Phi^(kk-prime) on1+Xt. This is exactly the two displayed calculations immediately below the square,and fixes the lower arrow right-to-left without changing any object or label.

[279] EN-CONTRIB-VI-365-01 Source folio 365. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=379; scan/SGA6_witness_300dpi.djvu#page=376
Printed reading: $X = \underline{\underline{P}}(\underline{E})$
Corrected reading: $X = \mathbf{P}(\underline{E})$
Source: ASTRA-VI-PROJECTIVE-P-365. Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all1 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.

[280] EN-CONTRIB-VI-365-02 Source folio 365. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=379; scan/SGA6_witness_300dpi.djvu#page=376
Printed reading: $\mathbf{S}(\underline{E})$-Module module $\underline{M}$ of finite presentation
Corrected reading: $\mathbf{S}(\underline{E})$-Module $\underline{M}$ of finite presentation
Source: check/findings/findings_361_365.json#36. Both scans repeat the noun. Current removes the second, lower-case copy; diplomatic keeps both words. The graded module, ring and all qualification words remain identical.

[281] EN-CONTRIB-VI-365-03 Source folio 365. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=379; scan/SGA6_witness_300dpi.djvu#page=376
Printed reading: see EGA II 8.12.1.
Corrected reading: see EGA II 8.12.1).
Source: check/findings/findings_361_365.json#37. Both scans end the EGA locator with a stop alone. Current adds one close before that stop, matching the opening before the symmetric-algebra explanation.

[282] EN-CONTRIB-VI-366-01 Source folio 366. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=380; scan/SGA6_witness_300dpi.djvu#page=377
Printed reading:

$$\underline{M}_n \otimes_{\underline{O}_S} \underline{B}(-n) \longrightarrow M, \tag{1.2.1}$$

Corrected reading:

$$\underline{M}_n \otimes_{\underline{O}_S} \underline{B}(-n) \longrightarrow \underline{M}, \tag{1.2.1}$$

Source: ASTRA-VI-TARGET-RULE-366. Both images, including the native300dpi enlarged display, leave the target M unruled although the source M_n and the later M_k carry their own rules. The map is explicitly x tensor a↦a.x, with x∈M_n and a∈B_{k-n}, landing in M_k. These component maps assemble in the very same graded sheaf M already fixed in1.2. Current restores its letter rule; diplomatic retains the exact isolated omission. No module, grading or morphism is replaced.

[283] EN-CONTRIB-VI-368-01 Source folio 368. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=382; scan/SGA6_witness_300dpi.djvu#page=379
Printed reading: do not depend on the decomposition
Corrected reading: do not depend on the decomposition.
Source: check/findings/findings_366_370.json#7. Both images end the first paragraph without a stop. Diplomatic keeps the omission; current supplies the sentence-ending full stop already present.

[284] EN-CONTRIB-VI-369-01 Source folio 369. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=383; scan/SGA6_witness_300dpi.djvu#page=380
Printed reading: $X_0 = \underline{\underline{P}}(\underline{E}_0)$
Corrected reading: $X_0 = \mathbf{P}(\underline{E}_0)$
Source: ASTRA-VI-PROJECTIVE-P-369. Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all2 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.

[285] EN-CONTRIB-VI-369-02 Source folio 369. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=383; scan/SGA6_witness_300dpi.djvu#page=380
Printed reading: $\underline{O}_X$
Corrected reading: $\underline{O}_{X_0}$
Source: ASTRA-VI-BASE-SHEAF-369. The primary and native pages both print O_X with no zero. The same sentence defines X_0=P(E_0), the diagram gives g:X→X_0, and F=g^*(F_0). The printed component operation is g^*F_0=O_X tensor over g^{-1}O_{X_0} with g^{-1}F_0, using the ring map g-sharp:g^{-1}O_{X_0}→O_X. Thus the module supplied before this pullback is over O_{X_0}; adding only the zero restores the domain sheaf already specified by the displayed g. Diplomatic keeps O_X; current uses O_{X_0}.

[286] EN-CONTRIB-VI-369-03 Source folio 369. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=383; scan/SGA6_witness_300dpi.djvu#page=380
Printed reading: $X_\mu = \underline{\underline{P}}(\underline{E}_\mu)$
Corrected reading: $X_\mu = \mathbf{P}(\underline{E}_\mu)$
Source: ASTRA-VI-PROJECTIVE-P-369. Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all2 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.

[287] EN-CONTRIB-VI-371-01 Source folio 371. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=385; scan/SGA6_witness_300dpi.djvu#page=382
Printed reading: it is clear that it surjective.
Corrected reading: it is clear that it is surjective.
Source: check/findings/findings_371_375.json#1. Both full scans omit est with no insertion above the line. Current supplies that verb only; diplomatic keeps the printed omission.

[288] EN-CONTRIB-VI-372-01 Source folio 372. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=386; scan/SGA6_witness_300dpi.djvu#page=383
Printed reading: $\underline{Z}_{r+1}$
Corrected reading: $\underline{Z}_r$
Source: ASTRA-VI-SYZYGY-372. The complete main and native pages define Z_i=ker(d_i:L_i→L_{i−1}), then print remplacer L_{r+1} par Z_{r+1}. Write Z_{−1}=M. The exact sequences0→Z_i→L_i→Z_{i−1}→0 and the stated affine projective-dimension bound pd(M)≤r+1 give Ext^1(Z_r,N)=Ext^{r+2}(M,N)=0. Hence the finite-presentation Z_r is projective on each affine open and1.5.1 gives its graded local freeness. The exact truncation is0→Z_r→L_r→…→L_0→M→0, with the kernel inclusion as first differential. The old differential factors L_{r+1}→Z_r→L_r; this factorization and the identities on L_r,…,L_0 give its comparison map. An injective top term with this same exact tail maps isomorphically onto ker(d_r)=Z_r, uniquely by the kernel universal property. The printed Z_{r+1} instead has zero composite with d_{r+1}. Concrete source instance:r=0,S=Spec k,B=k[t],M=B/(t),L_1=B(−1)→L_0=B by multiplication by t. Then Z_0=(t),Z_1=0; using Z_1 leaves the nonzero kernel(t) unresolved. Current changes only the final subscript to r. The earlier i≥r+1 sentence is true and remains intact. Independent text/math audit:independent_syzygy_audit_372.json; it makes no additional image-reading claim.

[289] EN-CONTRIB-VI-372-02 Source folio 372. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=386; scan/SGA6_witness_300dpi.djvu#page=383
Printed reading:

i) $\underline{L}^{(m)} = \underline{L}$ for $m \leq m_0$;
ii) $\underline{L}^{(m)} = 0$ for $m \geq m_1$;
iii) for every m, $\mathrm{gr}^m(\underline{L}) = \underline{L}^{(m)}/\underline{L}^{(m+1)}$ is isomorphic as a graded B-Module to a graded B-Module of the form $\underline{N}_m \otimes_{\underline{O}_S} \underline{B}(m)$, where $\underline{N}_m$ is an $\underline{O}_S$-Module locally free of finite type.

Corrected reading:

i) $\underline{L}^{(m)} = \underline{L}$ for $m \leq m_0$;
ii) $\underline{L}^{(m)} = 0$ for $m \geq m_1$;
iii) for every m, $\mathrm{gr}^m(\underline{L}) = \underline{L}^{(m)}/\underline{L}^{(m+1)}$ is isomorphic as a graded $\underline{B}$-Module to a graded $\underline{B}$-Module of the form $\underline{N}_m \otimes_{\underline{O}_S} \underline{B}(m)$, where $\underline{N}_m$ is an $\underline{O}_S$-Module locally free of finite type.

Source: ASTRA-VI-ALGEBRA-RULE-372. Both images leave the first B after comme unruled, while the next B and B(m) have their own rules. The proposition explicitly sets B=Sym(E). Its filtered submodules, quotient L^(m)/L^(m+1), and comparison N_m tensor B(m) use this same algebra and its original action. The algebra identification is id_B, with quotient action b·[l]=[bl]; no algebra is replaced. Current restores the one missing B rule, diplomatic keeps its absence.

[290] EN-CONTRIB-VI-373-01 Source folio 373. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=387; scan/SGA6_witness_300dpi.djvu#page=384
Printed reading:

$$\underline{M}_{n_0} \otimes_{\underline{O}_S} \underline{B}(-n_0) \longrightarrow M.$$

Corrected reading:

$$\underline{M}_{n_0} \otimes_{\underline{O}_S} \underline{B}(-n_0) \longrightarrow \underline{M}.$$

Source: ASTRA-VI-TARGET-RULE-373. The first displayed comparison ends in ruled M; the second ends in plain M in both images. Both are explicitly the map(1.2.1), x tensor a↦a.x, whose degree-k target is M_k. The second uses its n_0 component and lands in the same fixed graded sheaf M. Current restores this one target rule; diplomatic preserves the observed omission.

[291] EN-CONTRIB-VI-374-01 Source folio 374. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=388; scan/SGA6_witness_300dpi.djvu#page=385
Printed reading: $\lambda_t(-\xi) = 1/1 + \xi t$
Corrected reading: $\lambda_t(-\xi) = 1/(1 + \xi t)$
Source: ASTRA-VI-RECIPROCAL-374. Both complete images print the inline slash expression1/1+xi t without denominator parentheses. The immediately preceding identity is lambda_t(xi)=1+xi t. By the stated pre-lambda addition law, lambda_t(−xi)lambda_t(xi)=lambda_t(0)=1 in K^bullet(X)[[t]]. The series1+xi t has constant coefficient1, and its uniquely determined inverse is sum_{j≥0}(−xi)^j t^j. This identifies the full denominator1+xi t without changing its letters or their order. For the source instance S=Spec k,r=0,X=P^0_k,xi=1, the ordinary ungrouped expression1/1+t gives1+t, whose product with1+t is1+2t+t^2, while the required product is1. Current inserts the two denominator parentheses; diplomatic retains the source slash expression.

[292] EN-CONTRIB-VI-375-01 Source folio 375. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=389; scan/SGA6_witness_300dpi.djvu#page=386
Printed reading:

$$(\underline{F}(n)) - (\underline{G}(n)) = \sum_i ((\underline{L}_i(n)) - (\underline{M}_i(n)) + (\underline{N}_i(n))),$$

Corrected reading:

$$(\underline{F}(n)) - (\underline{G}(n)) = \sum_i n_i ((\underline{L}_i(n)) - (\underline{M}_i(n)) + (\underline{N}_i(n))),$$

Source: check/findings/findings_371_375.json#12. Both375 scans lack the coefficients present in the immediately preceding free-group equality. The automorphism T_n of A sends each basis generator(E) to(E tensor O_X(n)); its inverse is T_{−n}. Z-linearity gives T_n(sum_i n_i R_i)=sum_i n_i T_n(R_i) with R_i=(L_i)−(M_i)+(N_i). Thus every displayed coefficient is preserved by the printed tensorization map. Current restores n_i; diplomatic keeps its omission.

[293] EN-CONTRIB-VI-376-01 Source folio 376. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=390; scan/SGA6_witness_300dpi.djvu#page=387
Printed reading:

$$(f_*(\underline{F}(n))) - (f_*(\underline{G}(n))) = \sum_i n_i((f_*(\underline{L}_i(n)) - (f_*(\underline{M}_i(n))) + (f_*(\underline{N}_i(n)))).$$

Corrected reading:

$$(f_*(\underline{F}(n))) - (f_*(\underline{G}(n))) = \sum_i n_i((f_*(\underline{L}_i(n))) - (f_*(\underline{M}_i(n))) + (f_*(\underline{N}_i(n)))).$$

Source: check/findings/findings_376_380.json#1. Both full images have two closing parentheses after L_i(n), three after M_i(n), and four after N_i(n). The free-group expression is the image of (L_i(n))−(M_i(n))+(N_i(n)) under the generator map (H)↦(f_*H); each generator has its own outer pair. Current restores precisely the missing close after the first generator.

[294] EN-CONTRIB-VI-376-02 Source folio 376. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=390; scan/SGA6_witness_300dpi.djvu#page=387
Printed reading: By 1.2, for sufficiently large
Corrected reading: By 1.3, for sufficiently large
Source: ASTRA-VI-REFERENCE-376. Both full images print D'après1.2. Section1.2 defines the graded-module functors; the stated simultaneous vanishing and local freeness are the conclusions of1.3 on368, already personally read. The application map substitutes each of the finite family L_i,M_i,N_i,F,G for its variable F and keeps the same projective-bundle morphism f:X→S. Each is a locally free finite-type O_X-module and is flat over S because f is flat. Taking the maximum of their finitely many twist bounds gives every local-freeness assertion and R^1 f_*L_i(n)=0. The long exact sequence then yields0→f_*L_i(n)→f_*M_i(n)→f_*N_i(n)→0. Current changes only the cited final digit from2 to3.

[295] EN-CONTRIB-VI-377-01 Source folio 377. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=391; scan/SGA6_witness_300dpi.djvu#page=388
Printed reading:

$$\sum_{i=0}^{r+1}(-1)^i\lambda^i(E)\sigma_{\rho-i} = 0 \qquad \text{for every } p \geq 1. \tag{1.11.3}$$

Corrected reading:

$$\sum_{i=0}^{r+1}(-1)^i\lambda^i(E)\sigma_{p-i} = 0 \qquad \text{for every } p \geq 1. \tag{1.11.3}$$

Source: check/findings/findings_376_380.json#13. The enlarged native crop native377_rho.png resolves the round rho with its continuous descending stroke; the p in pour tout p has its separate serif head. The homogeneous degree-p term of the printed Koszul complex is wedge^i(E) tensor Sym^(p−i)(E), whose class is lambda^i([E]) sigma_(p−i). Taking the alternating sum gives the displayed relation with p−i. Current retains this uniquely indexed reading; diplomatic keeps rho−i.

[296] EN-CONTRIB-VI-378-01 Source folio 378. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=392; scan/SGA6_witness_300dpi.djvu#page=389
Printed reading:

$$\sum_{j=0}^{r} x_j \left(\sum_{i=0}^{r+1} (-1)^i \lambda^i(E) \sigma_{p(j)-i} \right) = 0,$$

Corrected reading:

$$\sum_{j=0}^{r} x_j \left(\sum_{i=0}^{r+1} (-1)^i \lambda^i(E) \sigma_{p(j)-i} \right) = 0, \tag{1.11.4}$$

Source: check/findings/findings_376_380.json#15. Both full378 images have an empty label area. The later two references explicitly name1.11.4. Substitution p(j)=n_0+r+j in this first equation and interchange of the two finite sums give the next equality; the i≤r summands vanish by1.11.2, leaving (−1)^(r+1)lambda^(r+1)(E) sum_j x_j sigma_(n_0−1+j)=0. Multiplication by (−1)^(r+1) gives the printed reduction. This identifies the first zero equation as the referenced relation. Keep it unnumbered diplomatically and restore label1.11.4 in current.

[297] EN-CONTRIB-VI-379-01 Source folio 379. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=393; scan/SGA6_witness_300dpi.djvu#page=390
Printed reading: $\xi^k, \ldots, \xi^{k+r+1}$
Corrected reading: $\xi^k, \ldots, \xi^{k+r}$
Source: check/findings/findings_376_380.json#21. Both images clearly include the final +1. Multiplication m_(xi^k):K^bullet(X)→K^bullet(X), a↦xi^k a, is a K^bullet(S)-module automorphism with inverse m_(xi^(−k)). It sends the already proved basis1,xi,...,xi^r exactly to xi^k,...,xi^(k+r). The next power satisfies1.9.1 multiplied by xi^k. For S=Spec k,r=0,X=P^0_k,xi=1, the printed two-element list is(1,1), and the nonzero coefficient vector(1,−1) maps to0. Current removes only the final +1.

[298] EN-CONTRIB-VI-379-02 Source folio 379. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=393; scan/SGA6_witness_300dpi.djvu#page=390
Printed reading: $X = \underline{\underline{P}}^r_S$
Corrected reading: $X = \mathbf{P}^r_S$
Source: ASTRA-VI-PROJECTIVE-P-379. Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all1 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.

[299] EN-CONTRIB-VI-380-01 Source folio 380. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=394; scan/SGA6_witness_300dpi.djvu#page=391
Printed reading: $X = \underline{\underline{P}}(\underline{E})$
Corrected reading: $X = \mathbf{P}(\underline{E})$
Source: ASTRA-VI-PROJECTIVE-P-380. Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all1 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.

[300] EN-CONTRIB-VI-382-01 Source folio 382. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=396; scan/SGA6_witness_300dpi.djvu#page=393
Printed reading:

$$\bar{f}^*(\underline{M}) \longrightarrow \underline{M}$$

Corrected reading:

$$\bar{f}^*(\underline{M}) \longrightarrow \underline{\widetilde{M}}$$

Source: ASTRA-VI-ADJOINT-TARGET-382. Both images and native382_top.png show a bare M at the target, while the preceding line twice has M-tilde. The preceding isomorphism is phi:M→bar-f_*M-tilde. The stated adjunction Hom_Y(bar-f^*M,N)=Hom_(S,A)(M,bar-f_*N) sends phi to epsilon_(M-tilde)∘bar-f^*phi:bar-f^*M→M-tilde. On an affine stalk A_p this is m tensor a↦a m in the localized module. Thus the displayed adjoint has exactly the previously defined target M-tilde. Current restores its tilde alone.

[301] EN-CONTRIB-VI-382-02 Source folio 382. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=396; scan/SGA6_witness_300dpi.djvu#page=393
Printed reading: $e > o$
Corrected reading: $e > 0$
Source: check/findings/findings_381_385.json#14. Both complete images show the small o, compared with the full-height zero in d>0 above. NOTE6 handles index zero only, so retain this baseline o diplomatically. The construction uses a homogeneous section of positive integral degree, and applies the same D_+ chart construction with substitution(s,d)↦(t,e); its lower bound is the integer0. Current restores0 in this one baseline position.

[302] EN-CONTRIB-VI-383-01 Source folio 383. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=397; scan/SGA6_witness_300dpi.djvu#page=394
Printed reading:

$$\begin{aligned} i^* \circ \bar{f}^*_{U,s}(\underline{M}_{(s),U}) &\simeq i^*\Big(f^{-1}_{U,s}(\underline{M}_{(s),U}) \otimes_{f^{-1}_{U,s}((B|U)_{(s)})} \underline{O}_{D_+(s)}\Big) \\ &\simeq i^{-1}\Big(i^{-1}_{U,s}(\underline{M}_{(s),U}) \otimes_{f^{-1}_{U,s}((B|U)_{(s)})} \underline{O}_{D_+(s)}\Big) \otimes_{i^{-1}(\underline{O}_{D_+(s)})} \underline{O}_{D_+(st)} \\ &\simeq f^{-1}_{U,st}(\underline{M}_{(s),U}) \otimes_{f^{-1}_{U,st}((\underline{B}|U)_{(s)})} \underline{O}_{D_+(st)}. \end{aligned}$$

Corrected reading:

$$\begin{aligned} i^* \circ \bar{f}^*_{U,s}(\underline{M}_{(s),U}) &\simeq i^*\Big(f^{-1}_{U,s}(\underline{M}_{(s),U}) \otimes_{f^{-1}_{U,s}((\underline{B}|U)_{(s)})} \underline{O}_{D_+(s)}\Big) \\ &\simeq i^{-1}\Big(f^{-1}_{U,s}(\underline{M}_{(s),U}) \otimes_{f^{-1}_{U,s}((\underline{B}|U)_{(s)})} \underline{O}_{D_+(s)}\Big) \otimes_{i^{-1}(\underline{O}_{D_+(s)})} \underline{O}_{D_+(st)} \\ &\simeq f^{-1}_{U,st}(\underline{M}_{(s),U}) \otimes_{f^{-1}_{U,st}((\underline{B}|U)_{(s)})} \underline{O}_{D_+(st)}. \end{aligned}$$

Source: check/findings/findings_381_385.json#17; ASTRA-VI-ALGEBRA-RULES-383. check/findings/findings_381_385.json#17: Native383_inner.png resolves the low i stem and separate upper ink; the preceding f has an ascender and hook. The diagram gives f_(U,st)=f_(U,s)∘i. Its inverse-image composition is i^−1∘f_(U,s)^−1=f_(U,st)^−1, with the equality on a stalk over y given by the identity on M_(f(y)). Applying i^* to the preceding tensor expression therefore yields f_(U,s)^−1 inside i^−1. Current changes only this inner i to f; diplomatic retains i. ASTRA-VI-ALGEBRA-RULES-383: Both images leave these two B letters plain, while the next four B occurrences are ruled. The first and second lines expand the same pullback tensor product over R=(B|U)_(s). The coefficient-ring comparison is id_R, induced by id_B under restriction and homogeneous localization b/s^m↦b/s^m. Current restores the two omitted rules without replacing any coefficient object.

[303] EN-CONTRIB-VI-384-01 Source folio 384. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=398; scan/SGA6_witness_300dpi.djvu#page=395
Printed reading:

$$F_k(L) = \bigoplus_{n \geq k} f_*(\underline{L}(n)). \tag{2.4.1}$$

Corrected reading:

$$F_k(\underline{L}) = \bigoplus_{n \geq k} f_*(\underline{L}(n)). \tag{2.4.1}$$

Source: ASTRA-VI-FUNCTOR-ARG-RULE-384. Both images and native384_LR.png leave the argument L in F_k(L) plain, while the immediately preceding L and the right-side L(n) have rules. The definition F_k(L)=direct-sum_(n≥k) f_*(L tensor O_X(n)) takes the identical O_X-module L as argument. Current restores this rule; the underlying object comparison is id_L.

[304] EN-CONTRIB-VI-384-02 Source folio 384. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=398; scan/SGA6_witness_300dpi.djvu#page=395
Printed reading:

$$\underline{\underline{\mathrm{R}}}F_k : D^+(\mathcal{A}) \longrightarrow D^+(\mathcal{C}). \tag{2.4.2}$$

Corrected reading:

$$\mathbb{R}F_k : D^+(\mathcal{A}) \longrightarrow D^+(\mathcal{C}). \tag{2.4.2}$$

Source: ASTRA-VI-DERIVED-R-384. Both complete images show two separate horizontal rules under the derived-functor R, resolved on384 in native384_LR.png and on385 beside2.5.3. Diplomatic renders both rules literally. Current uses the double-struck R for these specific occurrences. The notation map denotes the same functor: an injective resolution L→I^bullet is sent to F_k(I^bullet), with the identity chain map giving the comparison between the two renderings. No unviewed occurrence is changed.

[305] EN-CONTRIB-VI-384-03 Source folio 384. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=398; scan/SGA6_witness_300dpi.djvu#page=395
Printed reading: $L \rightsquigarrow f_*(\underline{L}(n))$
Corrected reading: $\underline{L} \rightsquigarrow f_*(\underline{L}(n))$
Source: ASTRA-VI-MAP-ARG-RULE-384. Both complete images leave the source L of L⇝f_*L(n) plain and rule its target occurrence. This is exactly the functor T_n:L↦f_*(L tensor O_X(n)) just used in the acyclicity statement, with source comparison id_L and identity on the tensor factor O_X(n). Current restores the omitted source rule.

[306] EN-CONTRIB-VI-385-01 Source folio 385. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=399; scan/SGA6_witness_300dpi.djvu#page=396
Printed reading: $z.\alpha_d(s^n)^{-1}$
Corrected reading: $z.\alpha_{nd}(s^n)^{-1}$
Source: ASTRA-VI-DEGREE-385. Both full images and native385_alpha.png print alpha_d(s^n). The immediately following definition has alpha_d:Gamma(V,B_d)→Gamma(f^−1(V),O_X(d)), while s^n has degree nd. The graded algebra map alpha sends a homogeneous b of degree q to its section of O_X(q); its multiplicativity gives alpha_(nd)(s^n)=alpha_d(s)^(tensor n). Multiplication by this section on D_+(V,s) is an isomorphism O→O(nd), whose inverse sends the restricted z of degree nd to a section of F. For B=k[x_0,x_1],s=x_0,d=1,n=2, x_0^2 belongs to B_2 and the printed alpha_1 has domain B_1; alpha_2(x_0^2)=alpha_1(x_0)^2 is the exact typed value. Current changes only this first alpha subscript to nd; the general definition of alpha_d stays intact.

[307] EN-CONTRIB-VI-385-02 Source folio 385. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=399; scan/SGA6_witness_300dpi.djvu#page=396
Printed reading:

$$\underline{\mathrm{Proj}}_0 \circ \underline{\underline{\mathrm{R}}}F_k \longrightarrow \mathrm{Id}_{D^+(\mathcal{A})}\,. \tag{2.5.3}$$

Corrected reading:

$$\underline{\mathrm{Proj}}_0 \circ \mathbb{R}F_k \longrightarrow \mathrm{Id}_{D^+(\mathcal{A})}\,. \tag{2.5.3}$$

Source: ASTRA-VI-DERIVED-R-385. Both complete images show two separate horizontal rules under the derived-functor R, resolved on384 in native384_LR.png and on385 beside2.5.3. Diplomatic renders both rules literally. Current uses the double-struck R for these specific occurrences. The notation map denotes the same functor: an injective resolution L→I^bullet is sent to F_k(I^bullet), with the identity chain map giving the comparison between the two renderings. No unviewed occurrence is changed.

[308] EN-CONTRIB-VI-385-03 Source folio 385. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=399; scan/SGA6_witness_300dpi.djvu#page=396
Printed reading: $X = \underline{\underline{P}}(E)$
Corrected reading: $X = \mathbf{P}(\underline{E})$
Source: ASTRA-VI-PROJECTIVE-P-385; ASTRA-VI-PROJECTIVE-ARG-RULE-385. ASTRA-VI-PROJECTIVE-P-385: Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all1 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.
ASTRA-VI-PROJECTIVE-ARG-RULE-385: Both complete images and native385_prop.png leave E inside P(E) plain; the same proposition introduces the ruled sheaf E immediately before it. The construction is X=Proj(Sym(E)), using exactly this sheaf. Its comparison is induced by id_E on each symmetric power, hence id_Sym(E) and id_Proj(Sym(E)). Current restores this one omitted rule; diplomatic retains plain E.

[309] EN-CONTRIB-VI-385-04 Source folio 385. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=399; scan/SGA6_witness_300dpi.djvu#page=396
Printed reading: $\underline{L}$.
Corrected reading: $\underline{L}^{\bullet}$
Source: check/findings/findings_381_385.json#32. The complete images and native385_prop.png show a baseline full stop immediately after the ruled L; native385_rule_X.png compares it with the nearby sentence stop. The attached subscript bullet places the dot below that baseline. Diplomatic now renders L followed by an ordinary math stop; current keeps the raised complex dot. The same complex is the argument of RF_k and Proj_0 RF_k in the personally read386 top. Its component indexing and differential are unchanged: the comparison is identity in every degree.

[310] EN-CONTRIB-VI-386-01 Source folio 386. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=400; scan/SGA6_witness_300dpi.djvu#page=397
Printed reading:

i) $\underline{\underline{\mathrm{R}}F_k(\underline{L}^{\bullet})}$ is $\mathcal{C}_0$-perfect;

ii) The canonical morphism (2.5.3) : $\mathrm{Proj}_0 \circ \underline{\underline{\mathrm{R}}F_k(\underline{L}^{\bullet})} \longrightarrow \underline{L}^{\bullet}$ is an isomorphism in $D^+(\mathcal{A})$.

Corrected reading:

i) $\mathbb{R}F_k(\underline{L}^{\bullet})$ is $\mathcal{C}_0$-perfect;

ii) The canonical morphism (2.5.3) : $\mathrm{Proj}_0 \circ \mathbb{R}F_k(\underline{L}^{\bullet}) \longrightarrow \underline{L}^{\bullet}$ is an isomorphism in $D^+(\mathcal{A})$.

Source: ASTRA-VI-DERIVED-R-386. Both complete images show two lower rules on each total derived-functor R counted here. Diplomatic reproduces those two rules; current uses double-struck R for the identical functor represented by F_k applied to an injective resolution, with identity comparison on that representative. On389 the two separate R^i symbols are visibly plain and are left unchanged.

[311] EN-CONTRIB-VI-386-02 Source folio 386. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=400; scan/SGA6_witness_300dpi.djvu#page=397
Printed reading:

$$\underline{\underline{\mathrm{R}}}F_k(\underline{L}^\bullet) = F_k(\underline{L}^\bullet)$$

Corrected reading:

$$\mathbb{R}F_k(\underline{L}^\bullet) = F_k(\underline{L}^\bullet)$$

Source: ASTRA-VI-DERIVED-R-386. Both complete images show two lower rules on each total derived-functor R counted here. Diplomatic reproduces those two rules; current uses double-struck R for the identical functor represented by F_k applied to an injective resolution, with identity comparison on that representative. On389 the two separate R^i symbols are visibly plain and are left unchanged.

[312] EN-CONTRIB-VI-387-01 Source folio 387. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=401; scan/SGA6_witness_300dpi.djvu#page=398
Printed reading:

$$\underline{\mathrm{Proj}_0}(F_k(\underline{L}^\bullet)) \longrightarrow \underline{L}^\bullet, \qquad ,$$

Corrected reading:

$$\underline{\mathrm{Proj}_0}(F_k(\underline{L}^\bullet)) \longrightarrow \underline{L}^\bullet,$$

Source: check/findings/findings_386_390.json#5. Both complete387 images have a comma after L and a second separated comma farther right. Diplomatic retains both; current keeps one, with the same sentence continuation et c'est un isomorphisme.

[313] EN-CONTRIB-VI-389-01 Source folio 389. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=403; scan/SGA6_witness_300dpi.djvu#page=400
Printed reading: $\underline{M}^\bullet = \underline{\underline{\mathrm{R}}}F_k(\underline{L}^\bullet)$
Corrected reading: $\underline{M}^\bullet = \mathbb{R}F_k(\underline{L}^\bullet)$
Source: ASTRA-VI-DERIVED-R-389. Both complete images show two lower rules on each total derived-functor R counted here. Diplomatic reproduces those two rules; current uses double-struck R for the identical functor represented by F_k applied to an injective resolution, with identity comparison on that representative. On389 the two separate R^i symbols are visibly plain and are left unchanged.

[314] EN-CONTRIB-VI-389-02 Source folio 389. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=403; scan/SGA6_witness_300dpi.djvu#page=400
Printed reading: $\underline{\mathrm{Proj}_0}(\underline{\underline{\mathrm{R}}}F_k(\underline{L}^\bullet))$
Corrected reading: $\underline{\mathrm{Proj}_0}(\mathbb{R}F_k(\underline{L}^\bullet))$
Source: ASTRA-VI-DERIVED-R-389. Both complete images show two lower rules on each total derived-functor R counted here. Diplomatic reproduces those two rules; current uses double-struck R for the identical functor represented by F_k applied to an injective resolution, with identity comparison on that representative. On389 the two separate R^i symbols are visibly plain and are left unchanged.

[315] EN-CONTRIB-VI-389-03 Source folio 389. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=403; scan/SGA6_witness_300dpi.djvu#page=400
Printed reading: $f^* = \underline{\mathrm{L}}f^*$
Corrected reading: $f^* = \mathbb{L}f^*$
Source: ASTRA-VI-DERIVED-L-389. Both images show two distinct rules below L in Lf^*. Diplomatic renders those rules literally, current uses the double-struck L. The identity comparison of the two notations is the same derived pullback functor; flatness makes O_X tensor_(f^−1O_S) f^−1(−) exact, and the resolution comparison maps induce the printed identification with f^*.

[316] EN-CONTRIB-VI-389-04 Source folio 389. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=403; scan/SGA6_witness_300dpi.djvu#page=400
Printed reading: where the $\underline{K}_i$ are the perfect complexes
Corrected reading: where the $\underline{K}_i^\bullet$ are the perfect complexes
Source: check/findings/findings_386_390.json#11. Both images show ruled K_i without the raised complex dot; the preceding display has it on both K_i occurrences. The prose explicitly identifies these with the complexes of2.7. The comparison is degreewise identity on each K_i^bullet with its existing differential. Diplomatic keeps the missing dot; current retains its supplied dot.

[317] EN-CONTRIB-VI-390-01 Source folio 390. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=404; scan/SGA6_witness_300dpi.djvu#page=401
Printed reading: $\underline{\underline{D}}(\underline{E}_i)$
Corrected reading: $\mathbf{D}(\underline{E}_i)$
Source: ASTRA-VI-FLAG-D-390. All three flag-scheme D occurrences have two distinct lower rules in both complete images. Diplomatic reproduces both rules; current retains the existing bold D. The notation map is identity on the functor of complete flags of locally free quotients of the specified E: it sends each flag and each pullback flag to itself, inducing the identity of its representing scheme. No scheme or flag type is changed.

[318] EN-CONTRIB-VI-390-03 Source folio 390. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=404; scan/SGA6_witness_300dpi.djvu#page=401
Printed reading: $\underline{E}_i$
Corrected reading: $\underline{E}_i$,
Source: check/findings/findings_386_390.json#18. Both complete images end that source line at E_i, with X= beginning the next. Diplomatic retains the omission. Current retains the supplied comma separating the sheaf/flag definitions from the following definition of X in the same hypothesis list.

[319] EN-CONTRIB-VI-390-02 Source folio 390. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=404; scan/SGA6_witness_300dpi.djvu#page=401
Printed reading: $X = \underline{\underline{D}}(\underline{E}_1) \times_S \ldots \times_S \underline{\underline{D}}(\underline{E}_n)$
Corrected reading: $X = \mathbf{D}(\underline{E}_1) \times_S \ldots \times_S \mathbf{D}(\underline{E}_n)$
Source: ASTRA-VI-FLAG-D-390. All three flag-scheme D occurrences have two distinct lower rules in both complete images. Diplomatic reproduces both rules; current retains the existing bold D. The notation map is identity on the functor of complete flags of locally free quotients of the specified E: it sends each flag and each pullback flag to itself, inducing the identity of its representing scheme. No scheme or flag type is changed.

[320] EN-CONTRIB-VI-391-01 Source folio 391. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=405; scan/SGA6_witness_300dpi.djvu#page=402
Printed reading:

$$X_k = \underline{\underline{P}}(\underline{F}_{k-1}); \qquad \underline{F}_k = \mathrm{Ker}\big(f_k^*(\underline{F}_{k-1}) \longrightarrow \underline{O}_{X_k}(1)\big),$$

Corrected reading:

$$X_k = \mathbf{P}(\underline{F}_{k-1}); \qquad \underline{F}_k = \mathrm{Ker}\big(f_k^*(\underline{F}_{k-1}) \longrightarrow \underline{O}_{X_k}(1)\big),$$

Source: ASTRA-VI-PROJECTIVE-P-391. Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all1 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.

[321] EN-CONTRIB-VI-392-01 Source folio 392. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=406; scan/SGA6_witness_300dpi.djvu#page=403
Printed reading: $X = \underline{\underline{D}}(\underline{E}) \times_S \underline{\underline{D}}(\underline{F})$
Corrected reading: $X = \mathbf{D}(\underline{E}) \times_S \mathbf{D}(\underline{F})$
Source: ASTRA-VI-FLAG-OPERATORS-392. Both complete images show two separate lower rules on each named operator counted here. Diplomatic restores those two rules, and current retains the existing bold operators. Each notation transport is identity on the represented flag or rank-r quotient functor, and on all its pullback maps; the representing scheme comparison is therefore the induced identity.

[322] EN-CONTRIB-VI-392-02 Source folio 392. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=406; scan/SGA6_witness_300dpi.djvu#page=403
Printed reading: **Theorem 3.3**
Corrected reading: **Theorem 3.3.**
Source: check/findings/findings_391_395.json#9. Both full images have no stop after3.3. Diplomatic keeps the omission; current retains the supplied label stop, paired against the source.

[323] EN-CONTRIB-VI-392-03 Source folio 392. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=406; scan/SGA6_witness_300dpi.djvu#page=403
Printed reading: A-Modules locally free
Corrected reading: $\underline{A}$-Modules locally free
Source: ASTRA-VI-TOPOS-COEFFICIENT-RULE-392. Both full source images leave A in des A-Modules plain, while the theorem introduces the ruled A in (T,A), and the two proof occurrences have the rule. These modules carry exactly the action of that same ring object: the comparison is id_A:A→A, with restriction of scalars along id_A equal to the identical module/action. Current restores this one omitted rule; diplomatic keeps the plain glyph.

[324] EN-CONTRIB-VI-393-01 Source folio 393. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=407; scan/SGA6_witness_300dpi.djvu#page=404
Printed reading: Let $\underline{E}$ be a locally free $\underline{A}$-Module of finite rank n. There exists a torsor $\underline{P}$ under $\underline{\mathrm{Gl}}(n, \underline{A})$ such that $\underline{E} = \underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A})} (\underline{A})^n$, where $(\underline{A})^n$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A})$-Module by the standard representation. We can then define a λ-homomorphism from $R_{\mathbb{Z}}(\underline{\underline{\mathrm{Gl}}}(n))$ to K by associating to every representation ρ of $\underline{\underline{\mathrm{Gl}}}(n)$ in a free $\mathbb{Z}$-module of rank p the class of the locally free $\underline{A}$-Module $\underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A})} (\underline{A})^p$, where $(\underline{A})^p$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A})$-module by the representation ρ. Under this homomorphism, the class of $\underline{E}$ is therefore the image of the standard representation. To check a polynomial relation among the $\lambda^i(E)$, it therefore suffices that this relation hold for the standard representation in $R_{\mathbb{Z}}(\underline{\underline{\mathrm{Gl}}}(n))$.
Corrected reading: Let $\underline{E}$ be a locally free $\underline{A}$-Module of finite rank n. There exists a torsor $\underline{P}$ under $\underline{\mathrm{Gl}}(n, \underline{A})$ such that $\underline{E} = \underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A})} (\underline{A})^n$, where $(\underline{A})^n$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A})$-Module by the standard representation. We can then define a λ-homomorphism from $R_{\mathbb{Z}}(\mathbf{Gl}(n))$ to K by associating to every representation ρ of $\mathbf{Gl}(n)$ in a free $\mathbb{Z}$-module of rank p the class of the locally free $\underline{A}$-Module $\underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A})} (\underline{A})^p$, where $(\underline{A})^p$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A})$-module by the representation ρ. Under this homomorphism, the class of $\underline{E}$ is therefore the image of the standard representation. To check a polynomial relation among the $\lambda^i(E)$, it therefore suffices that this relation hold for the standard representation in $R_{\mathbb{Z}}(\mathbf{Gl}(n))$.
Source: ASTRA-VI-GL-OPERATORS-393. Both full images show seven double-rule group-scheme Gl names and thirteen single-rule Gl(n,A)/Gl(p,A) names. For each object U the group-scheme evaluation is Gl_n(A(U))=GL_n(A(U)), with identical matrix multiplication and restriction maps. A frame p and vector v map to p(v), giving the contracted-product comparison [p,v]↦p(v) used by the source. Diplomatic retains the observed one/two rules; current retains bold group-scheme Gl and singly ruled group-sheaf Gl. No object or action is replaced.

[325] EN-CONTRIB-VI-393-02 Source folio 393. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=407; scan/SGA6_witness_300dpi.djvu#page=404
Printed reading: We can argue similarly with two locally free $\underline{A}$-Modules $\underline{E}, \underline{F}$ of ranks n, p; we can then find a torsor $\underline{P}$ under $\underline{\mathrm{Gl}}(n, \underline{A}) \times \underline{\mathrm{Gl}}(p, \underline{A})$ such that $\underline{E} = \underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A}) \times \underline{\mathrm{Gl}}(p,\underline{A})} (\underline{A})^n$, where $(\underline{A})^n$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A}) \times \underline{\mathrm{Gl}}(p, \underline{A})$-Module by letting $\underline{\mathrm{Gl}}(n, \underline{A})$ act by the standard representation and $\underline{\mathrm{Gl}}(p, \underline{A})$ by the trivial representation, and conversely for $\underline{F}$. We then define a λ-homomorphism from $R_{\mathbb{Z}}(\underline{\underline{\mathrm{Gl}}}(n) \times \underline{\underline{\mathrm{Gl}}}(p))$ to K in the same way as above, and are thus reduced to checking the relations among the $\lambda^i(E)$, $\lambda^j(F)$ for the corresponding elements of $R_{\mathbb{Z}}(\underline{\underline{\mathrm{Gl}}}(n) \times \underline{\underline{\mathrm{Gl}}}(p))$.
Corrected reading: We can argue similarly with two locally free $\underline{A}$-Modules $\underline{E}, \underline{F}$ of ranks n, p; we can then find a torsor $\underline{P}$ under $\underline{\mathrm{Gl}}(n, \underline{A}) \times \underline{\mathrm{Gl}}(p, \underline{A})$ such that $\underline{E} = \underline{P} \times_{\underline{\mathrm{Gl}}(n,\underline{A}) \times \underline{\mathrm{Gl}}(p,\underline{A})} (\underline{A})^n$, where $(\underline{A})^n$ is regarded as a $\underline{\mathrm{Gl}}(n, \underline{A}) \times \underline{\mathrm{Gl}}(p, \underline{A})$-Module by letting $\underline{\mathrm{Gl}}(n, \underline{A})$ act by the standard representation and $\underline{\mathrm{Gl}}(p, \underline{A})$ by the trivial representation, and conversely for $\underline{F}$. We then define a λ-homomorphism from $R_{\mathbb{Z}}(\mathbf{Gl}(n) \times \mathbf{Gl}(p))$ to K in the same way as above, and are thus reduced to checking the relations among the $\lambda^i(E)$, $\lambda^j(F)$ for the corresponding elements of $R_{\mathbb{Z}}(\mathbf{Gl}(n) \times \mathbf{Gl}(p))$.
Source: ASTRA-VI-GL-OPERATORS-393. Both full images show seven double-rule group-scheme Gl names and thirteen single-rule Gl(n,A)/Gl(p,A) names. For each object U the group-scheme evaluation is Gl_n(A(U))=GL_n(A(U)), with identical matrix multiplication and restriction maps. A frame p and vector v map to p(v), giving the contracted-product comparison [p,v]↦p(v) used by the source. Diplomatic retains the observed one/two rules; current retains bold group-scheme Gl and singly ruled group-sheaf Gl. No object or action is replaced.

[326] EN-CONTRIB-VI394-001 Source folio 394. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=408; scan/SGA6_witness_300dpi.djvu#page=405
Printed reading: -scheme $\underline{\underline{D}}_{\pi}(\underline{E})$
Corrected reading: -scheme $\mathbf{D}_{\pi}(\underline{E})$
Source: ASTRA-VI-FLAG-OPERATORS-394. Both complete images show two separate lower rules on each named operator counted here. Diplomatic restores those two rules, and current retains the existing bold operators. Each notation transport is identity on the represented flag or rank-r quotient functor, and on all its pullback maps; the representing scheme comparison is therefore the induced identity.

[327] EN-CONTRIB-VI394-002 Source folio 394. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=408; scan/SGA6_witness_300dpi.djvu#page=405
Printed reading: $\underline{\underline{D}}_{\pi}(\underline{E}) = \underline{\underline{P}}(\underline{E})$
Corrected reading: $\mathbf{D}_{\pi}(\underline{E}) = \mathbf{P}(\underline{E})$
Source: ASTRA-VI-FLAG-OPERATORS-394; ASTRA-VI-PROJECTIVE-P-394. ASTRA-VI-FLAG-OPERATORS-394: Both complete images show two separate lower rules on each named operator counted here. Diplomatic restores those two rules, and current retains the existing bold operators. Each notation transport is identity on the represented flag or rank-r quotient functor, and on all its pullback maps; the representing scheme comparison is therefore the induced identity. ASTRA-VI-PROJECTIVE-P-394: Both complete source images were compared during this folio's review. The native image was re-opened at closure of VI (native385_prop.png for385); all1 projective P occurrences show two distinct lower rules. NOTE6 explicitly accepts the bold transport for the symmetric-algebra S, but supplies no projective-P exception. For each base map T→S, the projective notation represents the set of surjections E_T→L with L invertible, modulo isomorphism of the quotient. The comparison between the two rendered P names sends every quotient to itself and has itself as inverse; it commutes with pullback. On Proj(Sym E), it is induced by id_E, hence by id on every symmetric power and every degree-zero localization chart. Thus the corrected bold P preserves precisely the represented functor while the diplomatic form reproduces the two observed lower rules.

[328] EN-CONTRIB-VI394-003 Source folio 394. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=408; scan/SGA6_witness_300dpi.djvu#page=405
Printed reading: $\underline{\underline{D}}_{\pi}(\underline{E}) = \underline{\underline{G}}_{r,n}(\underline{E})$
Corrected reading: $\mathbf{D}_{\pi}(\underline{E}) = \mathbf{G}_{r,n}(\underline{E})$
Source: ASTRA-VI-FLAG-OPERATORS-394. Both complete images show two separate lower rules on each named operator counted here. Diplomatic restores those two rules, and current retains the existing bold operators. Each notation transport is identity on the represented flag or rank-r quotient functor, and on all its pullback maps; the representing scheme comparison is therefore the induced identity.

[329] EN-CONTRIB-VI394-004 Source folio 394. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=408; scan/SGA6_witness_300dpi.djvu#page=405
Printed reading: $\underline{\underline{D}}(\underline{E})$
Corrected reading: $\mathbf{D}(\underline{E})$
Source: ASTRA-VI-FLAG-OPERATORS-394. Both complete images show two separate lower rules on each named operator counted here. Diplomatic restores those two rules, and current retains the existing bold operators. Each notation transport is identity on the represented flag or rank-r quotient functor, and on all its pullback maps; the representing scheme comparison is therefore the induced identity.

[330] EN-CONTRIB-VI394-005 Source folio 394. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=408; scan/SGA6_witness_300dpi.djvu#page=405
Printed reading: on $\underline{\underline{D}}_{\pi}(\underline{E})$
Corrected reading: on $\mathbf{D}_{\pi}(\underline{E})$
Source: ASTRA-VI-FLAG-OPERATORS-394. Both complete images show two separate lower rules on each named operator counted here. Diplomatic restores those two rules, and current retains the existing bold operators. Each notation transport is identity on the represented flag or rank-r quotient functor, and on all its pullback maps; the representing scheme comparison is therefore the induced identity.

[331] EN-CONTRIB-VI394-006 Source folio 395. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=409; scan/SGA6_witness_300dpi.djvu#page=406
Printed reading:

$$\underline{\underline{D}}_{\pi'}(\underline{E}) \simeq \underline{\underline{D}}_{\pi''}(\underline{F}).$$

Corrected reading:

$$\mathbf{D}_{\pi'}(\underline{E}) \simeq \mathbf{D}_{\pi''}(\underline{F}).$$

Source: ASTRA-VI-FLAG-OPERATORS-395. Both complete images show two separate lower rules on each named operator counted here. Diplomatic restores those two rules, and current retains the existing bold operators. Each notation transport is identity on the represented flag or rank-r quotient functor, and on all its pullback maps; the representing scheme comparison is therefore the induced identity.

[332] EN-CONTRIB-VI394-007 Source folio 395. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=409; scan/SGA6_witness_300dpi.djvu#page=406
Printed reading: isobaric relations deduced from the relation::
Corrected reading: isobaric relations deduced from the relation:
Source: check/findings/findings_391_395.json#39. Both full395 images show two distinct colon pairs. Diplomatic retains both; current removes only the duplicate colon.

[333] EN-CONTRIB-VI394-008 Source folio 395. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=409; scan/SGA6_witness_300dpi.djvu#page=406
Printed reading:

$$\sum_{i=1}^{n} c_i = \prod_{j=1}^{k} \left(\sum_{i=1}^{p_j} c_i^{(j)} \right)$$

Corrected reading:

$$1 + \sum_{i=1}^{n} c_i = \prod_{j=1}^{k} \left(1 + \sum_{i=1}^{p_j} c_i^{(j)} \right)$$

Source: ASTRA-VI-BLOCK-SYMMETRIC-CONSTANTS-395. Both complete primary/native395 images print the displayed equality without either constant1, and the personally read396 top assigns weight i to c_i and c_i^(j). In Z[xi_1,...,xi_n][t], the coefficient of t^r in product_a(1+xi_a t) is the sum of monomials indexed by r-element subsets. Intersection with the declared disjoint blocks gives the exact bijection I↦(I∩B_1,...,I∩B_k), preserving each monomial and total cardinality. The empty subset contributes1 in every factor and in the total product. Consequently product_a(1+xi_a t)=product_j(1+sum_i sigma_i(block_j)t^i)=1+sum_r sigma_r(xi_1,...,xi_n)t^r. Apply the coefficientwise ring homomorphism to C=A[xi]/(sigma_r−c_r); its images are the stated c_r and c_i^(j). The n positive-degree coefficients give exactly c_r=sum_(a_1+...+a_k=r) product_j c_(a_j)^(j), with the empty-block coefficient1. The isobaric instruction selects these coefficient identities; its weights are formal bookkeeping and assert no grading on arbitrary A or specialized C. For A=Z,n=k=2,p_1=p_2=1,c_1=3,c_2=2, evaluation xi_1↦1,xi_2↦2 kills both defining relations and gives a unital C→Z. It sends the printed equality to5=2, while the corrected total equality gives6=6 and its coefficients give3=1+2,2=1·2. Adding1+ before the total sum and inside each block factor preserves every printed index and the isobaric instruction, and restores exactly the constants forced by the elementary-symmetric definitions. Diplomatic retains the original; current accepts this minimal part of the earlier Codex proposal. The formerly proposed extra terminal comma remains rejected because both scans omit it. Independent coefficient/context audit:independent_block_symmetric_audit_395.json, a text/math audit carrying no additional image claim; its separate source-context checks do not claim a proof of the lemma's basis or ideal-generation assertions.

[334] EN-CONTRIB-VI394-009 Source folio 396. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=410; scan/SGA6_witness_300dpi.djvu#page=407
Printed reading: being effected with weight
Corrected reading: being assigned weight
Source: check/findings/findings_396_400.json#1. Both complete396 images print effectués. The operation specified is the assignment w(c_i)=w(c_i^(j))=i used to select the isobaric coefficient equations; the corresponding French verb is affecter. Current corrects this word to affectés, preserving every variable and the assigned weight; diplomatic retains effectués.

[335] EN-CONTRIB-VI394-010 Source folio 396. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=410; scan/SGA6_witness_300dpi.djvu#page=407
Printed reading: $X = \underline{\underline{D}}_{\pi}(\underline{E})$
Corrected reading: $X = \mathbf{D}_{\pi}(\underline{E})$
Source: ASTRA-VI-DOUBLE-OPERATORS-396. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[336] EN-CONTRIB-VI394-011 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: -Module locally of rank
Corrected reading: -Module locally free of rank
Source: check/findings/findings_396_400.json#7. Both full397 images omit libre. This corollary is explicitly the case p_i=1 of4.5, whose hypothesis is the same locally free sheaf E of rank n. The specialization keeps id_E and changes only the partition to n blocks of size1; it therefore carries the printed4.5 local-freeness hypothesis. Retain the existing current libre and the diplomatic omission.

[337] EN-CONTRIB-VI394-012 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: $\underline{F}_1, \ldots, \underline{F}_m$
Corrected reading: $\underline{F}_1, \ldots, \underline{F}_n$
Source: ASTRA-VI-FLAG-LAST-INDEX-397. The complete source images show the last index with three m strokes; native397_Fn.png resolves them at native resolution. The attached transcription has n. Section4.5 defines F_i=ker(L_i→L_(i−1)) for1≤i≤k, and this corollary sets every p_i=1 with sum p_i=n, hence k=n. The exact indexed family is therefore (F_i)_(1≤i≤n), whose ordered class map gives xi_i=[F_i]. Current retains that final n; diplomatic restores the printed m. No m is bound in this passage.

[338] EN-CONTRIB-VI394-013 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: (notation of 3.5)
Corrected reading: (notation of 4.5)
Source: ASTRA-VI-FLAG-NOTATION-REFERENCE-397. Both complete397 images print3.5. The personally read4.5 on396 defines exactly the canonical quotients L_i and their kernels F_i. Under the stated specialization p_i=1,k=n, the comparison keeps each kernel F_i and sends it to its class xi_i. The reference is therefore4.5; current changes only its first digit.

[339] EN-CONTRIB-VI394-014 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: we then consider $\underline{\underline{P}}(\underline{E})$
Corrected reading: we then consider $\mathbf{P}(\underline{E})$
Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[340] EN-CONTRIB-VI394-015 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: and on $\underline{\underline{P}}(\underline{E})$
Corrected reading: and on $\mathbf{P}(\underline{E})$
Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[341] EN-CONTRIB-VI394-016 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading:

$$0 \longrightarrow \underline{F} \longrightarrow g^*(\underline{E}) \longrightarrow \underline{O}_{\underline{\underline{P}}(\underline{E})}(1) \longrightarrow 0,$$

Corrected reading:

$$0 \longrightarrow \underline{F} \longrightarrow g^*(\underline{E}) \longrightarrow \underline{O}_{\mathbf{P}(\underline{E})}(1) \longrightarrow 0,$$


Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[342] EN-CONTRIB-VI394-017 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: structural morphism of $\underline{\underline{P}}(\underline{E})$
Corrected reading: structural morphism of $\mathbf{P}(\underline{E})$
Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.


[343] EN-CONTRIB-VI394-018 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: usual flag bundle $\underline{\underline{D}}(\underline{F})$
Corrected reading: usual flag bundle $\mathbf{D}(\underline{F})$
Source: ASTRA-VI-DOUBLE-OPERATORS-397. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[344] EN-CONTRIB-VI394-019 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: over $\underline{\underline{P}}(\underline{E})$
Corrected reading: over $\mathbf{P}(\underline{E})$
Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[345] EN-CONTRIB-VI394-020 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading:



Corrected reading:



Source: ASTRA-VI-DOUBLE-OPERATORS-397; ASTRA-VI-PROJECTIVE-OPERATORS-397. ASTRA-VI-DOUBLE-OPERATORS-397: Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks. ASTRA-VI-PROJECTIVE-OPERATORS-397: Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[346] EN-CONTRIB-VI394-021 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: factors through $K^{\bullet}(\underline{\underline{P}}(\underline{E}))$
Corrected reading: factors through $K^{\bullet}(\mathbf{P}(\underline{E}))$
Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[347] EN-CONTRIB-VI394-022 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: induction hypothesis applied to $\underline{\underline{D}}(\underline{F})$
Corrected reading: induction hypothesis applied to $\mathbf{D}(\underline{F})$
Source: ASTRA-VI-DOUBLE-OPERATORS-397. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[348] EN-CONTRIB-VI394-023 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: and Corollary 3.4
Corrected reading: and Corollary 4.4
Source: ASTRA-VI-FLAG-BASIS-REFERENCE-397. Both complete397 images print corollaire3.4. The basis statement used is the personally read Corollaire4.4 on396. Here A=K^bullet(P(E)), the rank-(n−1) kernel is F, and the inductive presentation maps A[T_2,...,T_n]/(sigma_i−lambda^i(F)) isomorphically to K^bullet(D(F)) by T_i↦xi_i. Applying4.4 to these n−1 ordered variables gives exactly xi_2^alpha_2...xi_(n−1)^alpha_(n−1),0≤alpha_i≤n−i. Current changes only3.4 to4.4.

[349] EN-CONTRIB-VI394-024 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: has as a basis over $K^\bullet(\underline{\underline{P}}(\underline{E}))$
Corrected reading: has as a basis over $K^\bullet(\mathbf{P}(\underline{E}))$
Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[350] EN-CONTRIB-VI394-025 Source folio 397. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=411; scan/SGA6_witness_300dpi.djvu#page=408
Printed reading: moreover, $K^\bullet(\underline{\underline{P}}(\underline{E}))$
Corrected reading: moreover, $K^\bullet(\mathbf{P}(\underline{E}))$
Source: ASTRA-VI-PROJECTIVE-OPERATORS-397. Both complete397 images show the same pair of rules on all nine projective P operators, including the one in the O subscript and the diagram. Diplomatic renders both rules literally; current uses bold P, preserving the functor of invertible quotients by its identity natural transformation.

[351] EN-CONTRIB-VI394-026 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: usual flag bundle $\underline{\underline{D}}(\underline{E})$
Corrected reading: usual flag bundle $\mathbf{D}(\underline{E})$
Source: ASTRA-VI-DOUBLE-OPERATORS-398. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[352] EN-CONTRIB-VI394-027 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: $\underline{\underline{D}}_\pi(\underline{E})$
Corrected reading: $\mathbf{D}_\pi(\underline{E})$
Source: ASTRA-VI-DOUBLE-OPERATORS-398. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[353] EN-CONTRIB-VI394-028 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: It follows that $K^\bullet(\underline{\underline{D}}(\underline{E}))$
Corrected reading: It follows that $K^\bullet(\mathbf{D}(\underline{E}))$
Source: ASTRA-VI-DOUBLE-OPERATORS-398. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[354] EN-CONTRIB-VI394-029 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: on $\underline{\underline{D}}(\underline{E})$
Corrected reading: on $\mathbf{D}(\underline{E})$
Source: ASTRA-VI-DOUBLE-OPERATORS-398. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[355] EN-CONTRIB-VI394-030 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: (notation of 3.5 and 3.7)
Corrected reading: (notation of 4.5 and 4.7)
Source: ASTRA-VI-FLAG-SPLITTING-REFERENCES-398. Both complete398 images print3.5 and3.7. The personally read4.5 defines the block sheaves F_j, and4.7 defines the classes of the invertible factors on each full-flag bundle. Pull back F_j to its full-flag bundle, take its successive rank-one kernels Q_i^(j), and apply [Q_i^(j)]=xi_i^(j); these exact operations give the variables of this paragraph. Current changes the two leading reference digits to4, leaving the source readings diplomatic.

[356] EN-CONTRIB-VI394-031 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: is a subring of $K^\bullet(\underline{\underline{D}}(\underline{E}))$
Corrected reading: is a subring of $K^\bullet(\mathbf{D}(\underline{E}))$
Source: ASTRA-VI-DOUBLE-OPERATORS-398. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[357] EN-CONTRIB-VI394-032 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: -algebra of $K^\bullet(\underline{\underline{D}}(\underline{E}))$
Corrected reading: -algebra of $K^\bullet(\mathbf{D}(\underline{E}))$
Source: ASTRA-VI-DOUBLE-OPERATORS-398. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[358] EN-CONTRIB-VI394-033 Source folio 398. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=412; scan/SGA6_witness_300dpi.djvu#page=409
Printed reading: there is in $K^\bullet(\underline{\underline{D}}(\underline{E}))$
Corrected reading: there is in $K^\bullet(\mathbf{D}(\underline{E}))$
Source: ASTRA-VI-DOUBLE-OPERATORS-398. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[359] EN-CONTRIB-VI394-034 Source folio 399. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=413; scan/SGA6_witness_300dpi.djvu#page=410
Printed reading: $K^\bullet(\underline{\underline{D}}(\underline{E}))$
Corrected reading: $K^\bullet(\mathbf{D}(\underline{E}))$
Source: ASTRA-VI-DOUBLE-OPERATORS-399. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[360] EN-CONTRIB-VI394-035 Source folio 399. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=413; scan/SGA6_witness_300dpi.djvu#page=410
Printed reading: $\underline{\underline{D}}(\underline{E})$
Corrected reading: $\mathbf{D}(\underline{E})$
Source: ASTRA-VI-DOUBLE-OPERATORS-399. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[361] EN-CONTRIB-VI394-036 Source folio 399. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=413; scan/SGA6_witness_300dpi.djvu#page=410
Printed reading: denote by $\underline{\underline{G}}_S^{n,r}$
Corrected reading: denote by $\mathbf{G}_S^{n,r}$
Source: ASTRA-VI-DOUBLE-OPERATORS-399. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[362] EN-CONTRIB-VI394-037 Source folio 399. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=413; scan/SGA6_witness_300dpi.djvu#page=410
Printed reading: the $\underline{\underline{G}}_S^{n,r}$ form an inductive system
Corrected reading: the $\mathbf{G}_S^{n,r}$ form an inductive system
Source: ASTRA-VI-DOUBLE-OPERATORS-399. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[363] EN-CONTRIB-VI394-038 Source folio 399. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=413; scan/SGA6_witness_300dpi.djvu#page=410
Printed reading: $\underline{\underline{G}}_S^{n,r} \longrightarrow \underline{\underline{G}}_S^{n,r+1}$
Corrected reading: $\mathbf{G}_S^{n,r} \longrightarrow \mathbf{G}_S^{n,r+1}$
Source: ASTRA-VI-DOUBLE-OPERATORS-399. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[364] EN-CONTRIB-VI394-039 Source folio 399. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=413; scan/SGA6_witness_300dpi.djvu#page=410
Printed reading: $(\underline{O}_{\underline{\underline{G}}_S^{n,r}})^r$
Corrected reading: $(\underline{O}_{\mathbf{G}_S^{n,r}})^r$
Source: ASTRA-VI-DOUBLE-OPERATORS-399. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[365] EN-CONTRIB-VI394-040 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading: $(\underline{O}_{\underline{\underline{G}}_S^{n,r}})^{r+1}$
Corrected reading: $(\underline{O}_{\mathbf{G}_S^{n,r}})^{r+1}$
Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[366] EN-CONTRIB-VI394-041 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading: $\underline{\underline{G}}_S^{n,\infty}$
Corrected reading: $\mathbf{G}_S^{n,\infty}$
Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[367] EN-CONTRIB-VI394-042 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading:

$$K^\bullet(\underline{\underline{G}}_S^{n,\infty}) = \varprojlim_r (K^\bullet(\underline{\underline{G}}_S^{n,r})). \tag{4.9.1}$$

Corrected reading:

$$K^\bullet(\mathbf{G}_S^{n,\infty}) = \varprojlim_r (K^\bullet(\mathbf{G}_S^{n,r})). \tag{4.9.1}$$

Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[368] EN-CONTRIB-VI394-043 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading: $\underline{\underline{G}}_S^{n,r}$
Corrected reading: $\mathbf{G}_S^{n,r}$
Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[369] EN-CONTRIB-VI394-044 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading: their class in $K^\bullet(\underline{\underline{G}}_S^{n,r})$
Corrected reading: their class in $K^\bullet(\mathbf{G}_S^{n,r})$
Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[370] EN-CONTRIB-VI394-045 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading: $K^\bullet(\underline{\underline{G}}_S^{n,\infty})$
Corrected reading: $K^\bullet(\mathbf{G}_S^{n,\infty})$
Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[371] EN-CONTRIB-VI394-046 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading: there is an isomorphism.:
Corrected reading: there is an isomorphism:
Source: check/findings/findings_396_400.json#21. Both complete400 images show a separate period before the colon. Diplomatic keeps it; current retains the removal of the stray period.

[372] EN-CONTRIB-VI394-047 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading:

$$K^\bullet(\underline{\underline{G}}_S^{n,\infty}) \xrightarrow{\sim} K^\bullet(S)[[c_1,\dots,c_n]].$$

Corrected reading:

$$K^\bullet(\mathbf{G}_S^{n,\infty}) \xrightarrow{\sim} K^\bullet(S)[[c_1,\dots,c_n]].$$

Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[373] EN-CONTRIB-VI394-048 Source folio 400. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=414; scan/SGA6_witness_300dpi.djvu#page=411
Printed reading: By 4.8, $K^\bullet(\underline{\underline{G}}_S^{n,r})$
Corrected reading: By 4.8, $K^\bullet(\mathbf{G}_S^{n,r})$
Source: ASTRA-VI-DOUBLE-OPERATORS-400. Every listed operator was directly inspected in both complete pages: D counts1,4,7,2 on396–399; G counts5,9 on399–400. Diplomatic restores literal double rules and current uses bold operators. Each rendering denotes the identical represented flag/quotient functor, with identity comparison on each quotient and its pullbacks.

[374] EN-CONTRIB-VI394-049 Source folio 401. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=415; scan/SGA6_witness_300dpi.djvu#page=412
Printed reading:

$$K^\bullet(\underline{\underline{G}}_S^{n,r}) = K^\bullet(S)[c_1,\dots,c_n]/I_r,$$

Corrected reading:

$$K^\bullet(\mathbf{G}_S^{n,r}) = K^\bullet(S)[c_1,\dots,c_n]/I_r,$$

Source: ASTRA-VI-DOUBLE-OPERATORS-401. Both complete401 images show two rules under the first G. Diplomatic restores those rules; current retains bold G. The two renderings represent the same rank-n quotient functor by the identity on every quotient and every base-change map.

[375] EN-CONTRIB-VI394-050 Source folio 401. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=415; scan/SGA6_witness_300dpi.djvu#page=412
Printed reading: $K^\bullet(\underline{\underline{G}}_S^{n,\infty})$
Corrected reading: $K^\bullet(\mathbf{G}_S^{n,\infty})$
Source: ASTRA-VI-DOUBLE-OPERATORS-401. Both complete401 images show two rules under the first G. Diplomatic restores those rules; current retains bold G. The two renderings represent the same rank-n quotient functor by the identity on every quotient and every base-change map.

[376] EN-CONTRIB-VI394-051 Source folio 402. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=416; scan/SGA6_witness_300dpi.djvu#page=413
Printed reading: i.e.. $\bar{x}' = -\bar{x}$
Corrected reading: i.e. $\bar{x}' = \bar{x}$
Source: check/findings/findings_401_405.json#9; check/findings/findings_401_405.json#8. check/findings/findings_401_405.json#9: Both complete402 images show the minus. Under the quotient map q:Fil^1K(X)→Fil^1K(X)/Fil^2K(X), x-prime−x=(xi−1)x belongs to Fil^2, so q(x-prime)=q(x). This supplies the precise comparison behind the existing current removal of the minus. For S=Spec(k),X=P^1, x=1−[O(−1)] has Euler characteristic1 and x^2=0, so its degree-one class has infinite order; the printed negative equality fails there. check/findings/findings_401_405.json#8: Both complete402 images show two consecutive full stops after the e. The abbreviation is already complete at i.e.; current removes only the immediately duplicated final stop, and diplomatic retains both.

[377] EN-CONTRIB-VI394-052 Source folio 403. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=417; scan/SGA6_witness_300dpi.djvu#page=414
Printed reading: With the notation and hypotheses of 4.1
Corrected reading: With the notation and hypotheses of 5.1
Source: check/findings/findings_401_405.json#22. Both complete403 images print4.1. The exact data used in the statement are f:P(E)→S, rank(E)=r+1 and x=1−[O(−1)], defined in5.1–5.2. The specialization of the general flag construction to its rank-one quotient gives this same projective bundle and class map, so the displayed decomposition is on precisely the object of5.1. Current retains5.1; diplomatic retains4.1.

[378] EN-CONTRIB-VI394-053 Source folio 403. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=417; scan/SGA6_witness_300dpi.djvu#page=414
Printed reading:

$$\gamma_t(x) = \lambda_{t/1-t}(1-\xi^{-1}) = \left(1+\frac{t}{1-t}\right)\left(1+\frac{\xi^{-1}t}{1-t}\right)^{-1} = 1/(1-xt).$$

Corrected reading:

$$\gamma_t(x) = \lambda_{t/(1-t)}(1-\xi^{-1}) = \left(1+\frac{t}{1-t}\right)\left(1+\frac{\xi^{-1}t}{1-t}\right)^{-1} = 1/(1-xt).$$

Source: ASTRA-VI-GAMMA-SUBSTITUTION-403. Both complete403 images print lambda_(t/1−t), without denominator parentheses. The next two explicit fractions have denominator1−t. In A[[t]], the continuous substitution rho(t)=t/(1−t) is inverse to eta(t)=t/(1+t): direct substitution gives rho(eta(t))=eta(rho(t))=t. Applying rho coefficientwise to lambda_u(1−xi^-1)=(1+u)/(1+xi^-1 u) gives1/(1−xt), x=1−xi^-1, exactly the rest of the displayed equation. Current adds only the denominator parentheses; diplomatic keeps the source slash string. The literal arithmetic parse t/1−t=0 would give1; on P^1 the printed final expression is1+xt with chi(x)=1, so that parse fails. This is a specified formal substitution, not an unsupported rewriting of notation.

[379] EN-CONTRIB-VI394-054 Source folio 404. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=418; scan/SGA6_witness_300dpi.djvu#page=415
Printed reading: $\underline{\underline{D}}(\underline{E}_i)$
Corrected reading: $\mathbf{D}(\underline{E}_i)$
Source: ASTRA-VI-DOUBLE-OPERATORS-404. Both complete404 images show two rules on D(E_i). Diplomatic restores both rules, current retains bold D. Both denote the identical complete-flag functor, with identity comparison on each flag and its pullbacks.

[380] EN-CONTRIB-VI394-055 Source folio 404. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=418; scan/SGA6_witness_300dpi.djvu#page=415
Printed reading: $X = \underline{\underline{D}}(\underline{E}_1) \times_S \ldots \times_S \underline{\underline{D}}(\underline{E}_n)$
Corrected reading: $X = \mathbf{D}(\underline{E}_1) \times_S \ldots \times_S \mathbf{D}(\underline{E}_n)$
Source: ASTRA-VI-DOUBLE-OPERATORS-404. Both complete404 images show two rules on D(E_i). Diplomatic restores both rules, current retains bold D. Both denote the identical complete-flag functor, with identity comparison on each flag and its pullbacks.

[381] EN-CONTRIB-VI394-056 Source folio 406. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=420; scan/SGA6_witness_300dpi.djvu#page=417
Printed reading:

$$\bar{f}(\lambda_t(F)) = \bar{f}_*(\lambda_t(E)\lambda_t(-\xi)) = \lambda_t(E)\bar{f}_*(\lambda_t(-\xi));$$

Corrected reading:

$$\bar{f}_*(\lambda_t(F)) = \bar{f}_*(\lambda_t(E)\lambda_t(-\xi)) = \lambda_t(E)\bar{f}_*(\lambda_t(-\xi));$$

Source: check/findings/findings_406_410.json#1; check/findings/findings_401_405.json#37. check/findings/findings_406_410.json#1: Both complete406 images show an empty subscript position under the first barred f. The map defined on405 is the coefficientwise extension sum a_it^i↦sum f_*(a_i)t^i; applying it to lambda_t(F) produces exactly the next two equal terms by the projection formula. Current restores its lower star; diplomatic retains the omitted star. check/findings/findings_401_405.json#37: Both complete406 images show an empty subscript position under the first barred f. The map defined on405 is the coefficientwise extension sum a_it^i↦sum f_*(a_i)t^i; applying it to lambda_t(F) produces exactly the next two equal terms by the projection formula. Current restores its lower star; diplomatic retains the omitted star.

[382] EN-CONTRIB-VI394-057 Source folio 406. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=420; scan/SGA6_witness_300dpi.djvu#page=417
Printed reading: $\bar{f}(\lambda_t(-\xi)) = \sum_{i\geq 0}(-1)^i f_*(\xi^i)t^i = \sigma_{-t}(E)$
Corrected reading: $\bar{f}_*(\lambda_t(-\xi)) = \sum_{i\geq 0}(-1)^i f_*(\xi^i)t^i = \sigma_{-t}(E)$
Source: check/findings/findings_406_410.json#2; check/findings/findings_401_405.json#38. check/findings/findings_406_410.json#2: Both complete406 images show this second omission. The explicitly displayed coefficient formula sum(−1)^i f_*(xi^i)t^i is the value of the same coefficientwise map bar f_* on lambda_t(−xi). Current restores the star; diplomatic retains the printed omission. check/findings/findings_401_405.json#38: Both complete406 images show this second omission. The explicitly displayed coefficient formula sum(−1)^i f_*(xi^i)t^i is the value of the same coefficientwise map bar f_* on lambda_t(−xi). Current restores the star; diplomatic retains the printed omission.

[383] EN-CONTRIB-VI394-058 Source folio 406. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=420; scan/SGA6_witness_300dpi.djvu#page=417
Printed reading: by (4.2.3).
Corrected reading: by (5.2.3).
Source: check/findings/findings_406_410.json#3; check/findings/findings_401_405.json#39. check/findings/findings_406_410.json#3: Both complete406 images print4.2.3. Formula5.2.3 on the personally read402 gives f_*(xi^i)=sigma_i for i≥1 and f_*(1)=1. Applying these equalities coefficient by coefficient sends sum(−1)^i f_*(xi^i)t^i to sigma_-t(E), exactly the equality cited here. Current retains the reference5.2.3; diplomatic keeps4.2.3. check/findings/findings_401_405.json#39: Both complete406 images print4.2.3. Formula5.2.3 on the personally read402 gives f_*(xi^i)=sigma_i for i≥1 and f_*(1)=1. Applying these equalities coefficient by coefficient sends sum(−1)^i f_*(xi^i)t^i to sigma_-t(E), exactly the equality cited here. Current retains the reference5.2.3; diplomatic keeps4.2.3.

[384] EN-CONTRIB-VI394-059 Source folio 407. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=421; scan/SGA6_witness_300dpi.djvu#page=418
Printed reading:

$$\begin{aligned}\sum_{i=0}^{r-1} c_{i+1}x'^{i} + x'^{r} &= (-1)^r.\left(\sum_{i=0}^{r-1}\gamma^{r-i}(E-r-1).(-1)^i x'^{i} + (-1)^r x'^{r}\right)\\ &= (-1)^r.\left(\sum_{i=0}^{r}\gamma^{r-i}(E-r-1)\gamma^{i}(-x')\right)\\ &= (-1)^r\gamma^r(E-r-1-x')\\ &= (-1)^r\gamma^r(F-x')\\ &= (-1)^r\lambda^r(F-1)\\ &= \lambda_{-1}(F).\end{aligned}$$

Corrected reading:

$$\begin{aligned}\sum_{i=0}^{r-1} c_{i+1}x'^{i} + x'^{r} &= (-1)^r.\left(\sum_{i=0}^{r-1}\gamma^{r-i}(E-r-1).(-1)^i x'^{i} + (-1)^r x'^{r}\right)\\ &= (-1)^r.\left(\sum_{i=0}^{r}\gamma^{r-i}(E-r-1)\gamma^{i}(-x')\right)\\ &= (-1)^r\gamma^r(E-r-1-x')\\ &= (-1)^r\gamma^r(F-r)\\ &= (-1)^r\lambda^r(F-1)\\ &= \lambda_{-1}(F).\end{aligned}$$

Source: check/findings/findings_406_410.json#8. Both complete407 images print F−x-prime. The preceding expression is E−r−1−x-prime. In the displayed K ring, substitution E=F+xi and x-prime=xi−1 gives E−r−1−x-prime=F−r exactly. Thus current retains F−r. The next equality is verified by [t^r]((1−t)^r lambda_(t/(1−t))(F))=sum_(j=0)^r(−1)^(r−j)lambda^j(F)=lambda^r(F−1); multiplication by(−1)^r gives lambda_-1(F). Diplomatic retains the printed x-prime.

[385] EN-CONTRIB-VI394-060 Source folio 409. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=423; scan/SGA6_witness_300dpi.djvu#page=420
Printed reading: The standard argument then reduces. us to the case
Corrected reading: The standard argument then reduces us to the case
Source: ASTRA-VI-STRAY-STOP-409. Both complete409 images and native409_permet.png show a separate stop after permet. The source verb governs the immediately following de se ramener, with alors between them; the stop cuts that same verbal construction. Diplomatic restores the observed period, while current retains the uninterrupted permet alors de se ramener. The reduction itself is the displayed restriction to the finite clopen rank partition, with inverse extension by zero.

[386] EN-CONTRIB-VI394-061 Source folio 411. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=425; scan/SGA6_witness_300dpi.djvu#page=422
Printed reading:

$$\text{ii)}\ \ \mathrm{Codim}(\mathrm{Supp}(H^*(\underline{M}^\bullet(Y))) \cap Y, Y) \geq i, \quad \text{with } H^*(\underline{M}^\bullet(Y)) = \bigoplus_{i \geq 0} H^i(\underline{M}^\bullet(Y)).$$

Corrected reading:

$$\text{ii)}\ \ \mathrm{Codim}(\mathrm{Supp}(H^*(\underline{M}^\bullet(Y))) \cap Y, Y) \geq i, \quad \text{with } H^*(\underline{M}^\bullet(Y)) = \bigoplus_{i \in \mathbb{Z}} H^i(\underline{M}^\bullet(Y)).$$

Source: ASTRA-VI-TOTAL-COHOMOLOGY-411. Both complete411 images print i≥0. The source allows arbitrary integer shifts of perfect complexes(I4.7–4.11,IV1.2 and2.2). On Spec(k),M=k[2] has its only cohomology in degree−2, but class1; the printed sum is zero and puts1 in K-dot(X)_1, contradicting6.6 at d=0. The all-degree sum projects to the printed nonnegative sum by identity on nonnegative summands and zero on negative ones, with kernel the sum of the negative cohomology sheaves. The proof on412 requires empty support to mean acyclicity and requires the tensor-support inclusion; k[−2] tensor k[2]→k,a⊗b↦ab, explicitly disproves that inclusion for the printed sum. Every integer q occurs as the sole degree of the allowed k[−q], so preserving the stated perfect category and the proof forces inclusion of every integer in this index. Current uses i∈Z; diplomatic keeps i≥0. An even shift[−2N] with inverse[2N] gives the exact same-class, same-total-support comparison with representatives concentrated in nonnegative cohomological degrees. Full convention checks, maps and counterexamples are in cohomology_index_411.json.

[387] EN-CONTRIB-VI394-062 Source folio 411. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=425; scan/SGA6_witness_300dpi.djvu#page=422
Printed reading: The filtration 5.5
Corrected reading: The filtration 6.5
Source: check/findings/findings_411_415.json#5. Both complete411 images print5.5. The immediately preceding6.5 defines the filtration K-dot(X)_i used in6.6;5.5 concerns the flag-scheme lambda presentation and supplies no such filtration. The locator map sends the displayed definition to6.5. Retain5.5 diplomatically and6.5 currently.

[388] EN-CONTRIB-VI394-063 Source folio 412. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=426; scan/SGA6_witness_300dpi.djvu#page=423
Printed reading: is a basis of $\underline{E}_{x'_i}$
Corrected reading: is a basis of $\underline{E}_{x_i}$
Source: check/findings/findings_411_415.json#12. Both complete412 images and native412_lemma_stalk_detail.png show a vertical prime at the index x; the proof prime in native412_proof_stalk_detail.png supplies a direct glyph comparison. Diplomatic retains the observed prime, without asserting the typist's intended stroke length. The statement names only x_i and its section germ s_(x_i), whose module is E_(x_i). The proof chooses a closed specialization x-prime_i later. For primes p_i⊂m_i the localization A_(m_i)→A_(p_i) induces E_(x-prime_i) tensor_A_(m_i) A_(p_i)→E_(x_i), e⊗a↦ae, with inverse e/a↦e⊗a^(-1); it carries the constructed basis to s_(x_i). Thus current E_(x_i) matches both the bound point and the actual conclusion; the primes inside the proof remain.

[389] EN-CONTRIB-VI394-064 Source folio 414. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=428; scan/SGA6_witness_300dpi.djvu#page=425
Printed reading: $\underline{O}_X$,-Modules
Corrected reading: $\underline{O}_{X'}$-Modules
Source: check/findings/findings_411_415.json#22. Both complete414 images show the low comma and its tail, unlike the raised straight primes in the nearby X-prime occurrences. Diplomatic retains that comma. The immediately defined flag scheme is X-prime, the two classes xi,eta belong to K-dot(X-prime), and their invertible sheaves enter the displayed gamma identity in that same group. Current O_(X-prime) restores this stated base. Pullback f^*:K-dot(X)→K-dot(X-prime) is the explicit map used in the following line; no base is silently identified.

[390] EN-CONTRIB-VI394-065 Source folio 415. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=429; scan/SGA6_witness_300dpi.djvu#page=426
Printed reading:

$$x' = \prod_{i=0}^{r} (\xi_i - 1),$$

Corrected reading:

$$x' = \prod_{i=1}^{r} (\xi_i - 1),$$

Source: ASTRA-VI-BASIS-PRODUCT-415. The complete primary and native415 images both print the lower index0 and upper indexr. The preceding414 sentence defines r as the relative dimension of the actual flag tower and asserts that x-prime is a K(X)-basis element. The reviewed projective-bundle basis of1.13 transforms from1,xi,...,xi^q to1,(xi−1),...,(xi−1)^q by the triangular binomial matrix; its inverse is xi^j=sum_(m=0)^j binom(j,m)(xi−1)^m. Composing these exact module-basis maps through the3.1 projective tower produces the top basis monomial with sum of the stage relative dimensions=r factors. The printed r+1 assertion fails already for the identity flag tower over Spec(k):r=0,every line class is1,and the printed single factor is0,whereas the corrected empty product is the basis element1. On P^1 the explicit ring Z[delta]/(delta^2),delta=xi−1,also makes every two-factor product of line-class differences zero. Current changes the lower endpoint0→1 and retains the printed upper endpoint. The alternative endpoint edit0≤i≤r−1 is related exactly by j=i+1,xi_j=eta_(j−1),including r=0; the choice here claims no unique unseen author indexing intention. The source fixes the factor count. The top-coordinate map pi_top(x-prime·f^*(a))=a proves the following cancellation without assuming multiplication by x-prime injective on all K(X-prime). Full maps and counterexamples are in basis_product_415.json.

[391] EN-CONTRIB-VI394-066 Source folio 415. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=429; scan/SGA6_witness_300dpi.djvu#page=426
Printed reading: Lectures notes in Mathematics
Corrected reading: Lecture Notes in Mathematics
Source: check/findings/findings_411_415.json#30. Both complete415 images print Lectures notes. The volume's own title matter gives the series name Lecture Notes in Mathematics. The bibliographic locator(R.Hartshorne,Residues and duality,Springer,series volume20) is preserved by the spelling correction; current retains Lecture Notes, diplomatic the printed wording.

[392] EN-CONTRIB-VII-001 Source folio 420. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=434
Printed reading: can be generated either the f_i or by the g_j
Corrected reading: can be generated either by the f_i or by the g_j
Source: check/findings/findings_416_420.json#16. The source omits par before the first family. The parallel soit par les g_j and the passive construction engendré par identify the missing preposition. The existing current insertion is exact.

[393] EN-CONTRIB-VII-002 Source folio 420. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=434
Printed reading: in a neighborhood of $\ddot{y}$ since
Corrected reading: in a neighborhood of y since
Source: check/findings/findings_416_420.json#18. The witness431 clearly separates the two dots, while the main makes a short horizontal blob. The point in this sentence is the same y introduced earlier; the statement concerns the stalk J_y and its neighborhood under the identity point map. No new decorated point is defined. Retain diplomatic ddot y and the existing current y.

[394] EN-CONTRIB-VII-003 Source folio 423. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=437
Printed reading: $\mathrm{Tor}_i^{A_0}(C_0, B)$
Corrected reading: $\mathrm{Tor}_i^{B_0}(C_0, B)$
Source: check/findings/findings_421_425.json#11. Both main and witness434 clearly print A_0. The third complex is K_{B_0}(T_1,...,T_m)⊗_{B_0}B, with T_j acting as g-bar_j. The first factor is the free Koszul resolution of C_0=B_0/(T_1,...,T_m), so its homology is exactly Tor_i^{B_0}(C_0,B), as the third line of1.7.2 independently states. This cannot be replaced by Tor over A_0: in the admissible case n=1,m=0,A=A_0=Z[T],B=B_0=C=C_0=Z, the third Koszul complex is the empty complex Z in degree0, whereas Tor_1^{Z[T]}(Z,Z)=Z from0→Z[T] –T→ Z[T]→Z→0. Tor_1^Z(Z,Z)=0 agrees with the actual complex. Correct only the current superscript to B_0.

[395] EN-CONTRIB-VII-004 Source folio 424. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=438
Printed reading: the sequence (1.8.1) is exact.
Corrected reading: the sequence (1.6.1) is exact.
Source: check/findings/findings_421_425.json#16. The main source clearly prints1.8.1. The displayed sequence immediately preceding the argument is the conormal sequence1.6.1, with maps induced by I⊂J and J→J/I. The proof shows the left kernel is locally free of rank0 and hence0, extending that sequence by0 on the left as in1.7(i). The following425 canonical projective-bundle sequence has different terms and is unnumbered in the source. The existing current1.6.1 therefore identifies the actual sequence proved exact.

[396] EN-CONTRIB-VII-005 Source folio 425. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=439
Printed reading: $\underline{\underline{S}}(\underline{E}) \longrightarrow \bigoplus_{n\geq 0} \underline{J}^n$
Corrected reading: $\mathbf{S}(\underline{E}) \longrightarrow \bigoplus_{n\geq 0} \underline{J}^n$
Source: ASTRA-VII-0023. The four main-scan glyphs each have two distinct lower strokes. The witness436 lower strip confirms the last three; the first is clear in the main upper strip. Restore both rules literally in diplomatic. Current retains the explicitly declared NOTE_6 bold-S representation of this same double-rule symmetric-algebra notation. The exact mathematical object is S(E)=⊕_{n≥0}Sym^n(E), with multiplication induced by symmetric tensor product. This occurrence-scoped transport preserves the distinction from the plain typed word Sym on418 and from the separately single-underlined E. It is not propagated to other exposés.

[397] EN-CONTRIB-VII-006 Source folio 425. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=439
Printed reading: The $\underline{S}(\underline{E})$-Module
Corrected reading: The $\mathbf{S}(\underline{E})$-Module
Source: ASTRA-VII-0023. The four main-scan glyphs each have two distinct lower strokes. The witness436 lower strip confirms the last three; the first is clear in the main upper strip. Restore both rules literally in diplomatic. Current retains the explicitly declared NOTE_6 bold-S representation of this same double-rule symmetric-algebra notation. The exact mathematical object is S(E)=⊕_{n≥0}Sym^n(E), with multiplication induced by symmetric tensor product. This occurrence-scoped transport preserves the distinction from the plain typed word Sym on418 and from the separately single-underlined E. It is not propagated to other exposés.

[398] EN-CONTRIB-VII-007 Source folio 425. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=439
Printed reading: is perfect as a graded $\underline{S}(\underline{E})$-Module
Corrected reading: is perfect as a graded $\mathbf{S}(\underline{E})$-Module
Source: ASTRA-VII-0023. The four main-scan glyphs each have two distinct lower strokes. The witness436 lower strip confirms the last three; the first is clear in the main upper strip. Restore both rules literally in diplomatic. Current retains the explicitly declared NOTE_6 bold-S representation of this same double-rule symmetric-algebra notation. The exact mathematical object is S(E)=⊕_{n≥0}Sym^n(E), with multiplication induced by symmetric tensor product. This occurrence-scoped transport preserves the distinction from the plain typed word Sym on418 and from the separately single-underlined E. It is not propagated to other exposés.

[399] EN-CONTRIB-VII-008 Source folio 425. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=439
Printed reading: by graded $\underline{S}(\underline{E})$-Modules locally free of finite type
Corrected reading: by graded $\mathbf{S}(\underline{E})$-Modules locally free of finite type
Source: ASTRA-VII-0023. The four main-scan glyphs each have two distinct lower strokes. The witness436 lower strip confirms the last three; the first is clear in the main upper strip. Restore both rules literally in diplomatic. Current retains the explicitly declared NOTE_6 bold-S representation of this same double-rule symmetric-algebra notation. The exact mathematical object is S(E)=⊕_{n≥0}Sym^n(E), with multiplication induced by symmetric tensor product. This occurrence-scoped transport preserves the distinction from the plain typed word Sym on418 and from the separately single-underlined E. It is not propagated to other exposés.

[400] EN-CONTRIB-VII-009 Source folio 426. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=440
Printed reading: Since every element of $\mathbb{Z}[T_1, \ldots, T_d, X_2/X_1, \ldots, X_d, X_1]$
Corrected reading: Since every element of $\mathbb{Z}[T_1, \ldots, T_d, X_2/X_1, \ldots, X_d/X_1]$
Source: check/findings/findings_426_430.json#9. The main and witness437 both clearly print X_d,X_1. Set z_i=X_i/X_1; the chart ring previously defined on this page is S=Z[T_1,...,T_d,z_2,...,z_d]. The substitutions T′_i=T_i−T_1z_i have the polynomial inverse T_i=T′_i+T_1z_i, so the claimed coordinate expression is exact for S. It fails for the printed ring with the additional X_1: for d=2 in Z[T_1,T_2,X_1^{±1},X_2], specialize T_1=T_2=X_2=0 and X_1=w. All the claimed new generators T_1,T′_2,z_2 map to0, but X_1 maps to the nonconstant invertible indeterminate w. Thus X_1 cannot be a polynomial in those generators. The existing current slash restores precisely the chart ring; diplomatic preserves the source comma.

[401] EN-CONTRIB-VII-010 Source folio 427. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=441
Printed reading: $\underline{\underline{S}}(E_0)$
Corrected reading: $\mathbf{S}(E_0)$
Source: ASTRA-VII-0028. The directly viewed occurrences have two separate rules below S. Keep those literally in diplomatic, and use the explicitly documented bold-S notation transport in current, as for the inspected425 occurrences. The object in each occurrence is the same symmetric algebra ⊕ Sym^n(E), with its symmetric-product multiplication. E and E_0 receive only their separately observed individual rules. This decision applies only to these counted VII occurrences.

[402] EN-CONTRIB-VII-011 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: by graded free $\underline{\underline{S}}(E_0)$-modules
Corrected reading: by graded free $\mathbf{S}(E_0)$-modules
Source: ASTRA-VII-0029. The directly viewed occurrences have two separate rules below S. Keep those literally in diplomatic, and use the explicitly documented bold-S notation transport in current, as for the inspected425 occurrences. The object in each occurrence is the same symmetric algebra ⊕ Sym^n(E), with its symmetric-product multiplication. E and E_0 receive only their separately observed individual rules. This decision applies only to these counted VII occurrences.

[403] EN-CONTRIB-VII-012 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: the last kernel is then a graded $\underline{\underline{S}}(E_0)$-module
Corrected reading: the last kernel is then a graded $\mathbf{S}(E_0)$-module
Source: ASTRA-VII-0029. The directly viewed occurrences have two separate rules below S. Keep those literally in diplomatic, and use the explicitly documented bold-S notation transport in current, as for the inspected425 occurrences. The object in each occurrence is the same symmetric algebra ⊕ Sym^n(E), with its symmetric-product multiplication. E and E_0 receive only their separately observed individual rules. This decision applies only to these counted VII occurrences.

[404] EN-CONTRIB-VII-013 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: as an ungraded $\underline{\underline{S}}(E_0)$-module
Corrected reading: as an ungraded $\mathbf{S}(E_0)$-module
Source: ASTRA-VII-0029. The directly viewed occurrences have two separate rules below S. Keep those literally in diplomatic, and use the explicitly documented bold-S notation transport in current, as for the inspected425 occurrences. The object in each occurrence is the same symmetric algebra ⊕ Sym^n(E), with its symmetric-product multiplication. E and E_0 receive only their separately observed individual rules. This decision applies only to these counted VII occurrences.

[405] EN-CONTRIB-VII-014 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: as a graded $\underline{\underline{S}}(\underline{E}_0)$-module
Corrected reading: as a graded $\mathbf{S}(\underline{E}_0)$-module
Source: ASTRA-VII-0029. The directly viewed occurrences have two separate rules below S. Keep those literally in diplomatic, and use the explicitly documented bold-S notation transport in current, as for the inspected425 occurrences. The object in each occurrence is the same symmetric algebra ⊕ Sym^n(E), with its symmetric-product multiplication. E and E_0 receive only their separately observed individual rules. This decision applies only to these counted VII occurrences.

[406] EN-CONTRIB-VII-015 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: a graded complex of graded $\underline{\underline{S}}(E)$-modules
Corrected reading: a graded complex of graded $\mathbf{S}(E)$-modules
Source: ASTRA-VII-0029. The directly viewed occurrences have two separate rules below S. Keep those literally in diplomatic, and use the explicitly documented bold-S notation transport in current, as for the inspected425 occurrences. The object in each occurrence is the same symmetric algebra $\oplus$ Sym^n(E), with its symmetric-product multiplication. E and E_0 receive only their separately observed individual rules. This decision applies only to these counted VII occurrences.

[407] EN-CONTRIB-VII-016 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: is a graded $\underline{\underline{S}}(E)$-Module whose
Corrected reading: is a graded $\mathbf{S}(E)$-Module whose
Source: ASTRA-VII-0029. The directly viewed occurrences have two separate rules below S. Keep those literally in diplomatic, and use the explicitly documented bold-S notation transport in current, as for the inspected425 occurrences. The object in each occurrence is the same symmetric algebra $\oplus$ Sym^n(E), with its symmetric-product multiplication. E and E_0 receive only their separately observed individual rules. This decision applies only to these counted VII occurrences.

[408] EN-CONTRIB-VII-017 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: $X' \to \underline{\underline{P}}(\underline{E})$
Corrected reading: $X' \to \mathbf{P}(\underline{E})$
Source: ASTRA-VII-0032. Both printed projective-bundle P letters have two distinct rules, independently of the single E rule. Diplomatic preserves both P strokes literally. Current uses the already declared bold-P notation for the same Proj(Sym E) construction: the identity of the graded symmetric algebra induces the identity on every degree-zero standard affine chart and hence on the glued projective bundle. This is an occurrence-scoped rendering choice, with no change of object.

[409] EN-CONTRIB-VII-018 Source folio 428. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=442
Printed reading: $\underline{\underline{P}}(\underline{E}) \to X$
Corrected reading: $\mathbf{P}(\underline{E}) \to X$
Source: ASTRA-VII-0032. Both printed projective-bundle P letters have two distinct rules, independently of the single E rule. Diplomatic preserves both P strokes literally. Current uses the already declared bold-P notation for the same Proj(Sym E) construction: the identity of the graded symmetric algebra induces the identity on every degree-zero standard affine chart and hence on the glued projective bundle. This is an occurrence-scoped rendering choice, with no change of object.

[410] EN-CONTRIB-VII-019 Source folio 431. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=445
Printed reading: M, Let $P_{\bullet}$
Corrected reading: M. Let $P_{\bullet}$
Source: check/findings/findings_431_435.json#10. The enlarged main lower strip shows the descending comma beside the overstruck M. The following capital Soit introduces a new independent sentence, so current full stop is the established punctuation correction; diplomatic comma is retained.

[411] EN-CONTRIB-VII-020 Source folio 432. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=446
Printed reading: conditions satisfied in particular in the noetherian case when the sequence $f_1, \dots, f_p, g_1, \dots g_r, h_1, \dots, h_s$
Corrected reading: conditions satisfied in particular in the noetherian case when the sequence $f_1, \dots, f_p, g_1, \dots, g_r, h_1, \dots, h_s$
Source: check/findings/findings_431_435.json#13. The main upper strip visibly lacks a comma between the last ellipsis dot and g_r in this one list; the other corresponding lists include it. The source lists the same successive entries g_1 through g_r, and current restores the missing punctuation without altering the sequence.

[412] EN-CONTRIB-VII-021 Source folio 433. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=447
Printed reading:

$$\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} (\underline{O}_T)^n,$$

Corrected reading:

$$\underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} (\underline{O}_T)^p,$$

Source: check/findings/findings_431_435.json#32. Both scans clearly print n. The preceding formula identifies Tor_i with O_T⊗∧^iE, with E=O_X^p by the explicit definition on432. At i=1 the basis classes[e_j] for1≤j≤p map bijectively to the p standard coordinates of O_T^p. Thus the rank is precisely p, also stated independently on435. No n is defined in this proof. Retain the existing current p and diplomatic n.

[413] EN-CONTRIB-VII-022 Source folio 433. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=447
Printed reading:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \bigwedge^i \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z). \tag{2.5.3}$$

Corrected reading:

$$\underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \bigwedge^i \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z). \tag{2.5.3}$$

Source: check/findings/findings_431_435.json#35. Main and witness444 plainly print the subscript i. The degree1 identification ψ:Tor_1→O_T^p from the previous display induces ∧^iψ; composing the2.5.2 isomorphism Tor_i→∧^i(O_T)^p with its inverse gives exactly Tor_i→∧^iTor_1. The printed Tor_i cannot supply this: take A=k[t_1,t_2], Y=Z=Spec(k), p=2 and r=s=0. Tensoring the Koszul resolution with k gives zero differentials, so Tor_1=k^2 and Tor_2=k. At i=2 the printed right member $\wedge^2$Tor_2=$\wedge^2$k=0, while the left member is k. The corrected $\wedge^2$Tor_1=$\wedge^2 k^2 \cong$k agrees. Keep the existing current1.

[414] EN-CONTRIB-VII-023 Source folio 434. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=448
Printed reading:

$$\bigwedge(\underline{O}_T)^n \xrightarrow{\sim} \bigoplus_{i\geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$$

Corrected reading:

$$\bigwedge(\underline{O}_T)^p \xrightarrow{\sim} \bigoplus_{i\geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$$

Source: check/findings/findings_431_435.json#37. Both main and witness445 print n. The source explicitly says this graded isomorphism is defined by2.5.2, whose degreei source is ∧^i(O_T^p). The direct sum of those degreei maps has source∧(O_T^p), with degree1 basis indexed1,. . . ,p. This is the same established rank correction as433; current had not yet propagated it to this independently printed occurrence. Correct only this current n to p.

[415] EN-CONTRIB-VII-024 Source folio 434. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=448
Printed reading:

$$\bigwedge \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \bigoplus_{i\geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$$

Corrected reading:

$$\bigwedge \underline{\mathrm{Tor}}_1^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z) \xrightarrow{\sim} \bigoplus_{i\geq 0} \underline{\mathrm{Tor}}_i^{\underline{O}_X}(\underline{O}_Y, \underline{O}_Z)$$

Source: check/findings/findings_431_435.json#41. Both scans print the subscript i. This map is explicitly the direct sum of2.5.3 and the restriction of2.5.1, which is defined to extend the identity in degree1. Its generator is therefore the degree1 sheaf Tor_1 under that identity. In the admissible p=2,r=s=0 example A=k[t_1,t_2], Y=Z=Spec(k), the displayed target has dimensions1,2,1 in degrees0,1,2. Substituting Tor_2=k as the printed generator would instead give exterior dimensions1,1,0, so the displayed graded isomorphism fails. The existing current Tor_1 gives exactly1,2,1 and the stipulated identity in degree1.

[416] EN-CONTRIB-VII-025 Source folio 435. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=449
Printed reading: $\underline{\underline{R}}i_*(\underline{L}^\bullet) = i_*(\underline{L}^\bullet)$
Corrected reading: $\mathbb{R}i_*(\underline{L}^\bullet) = i_*(\underline{L}^\bullet)$
Source: check/findings/findings_431_435.json#48. Main and witness446 show a typed R with two independent rules below it. Diplomatic now preserves those two strokes literally; current retains the documented double-struck R notation for the same derived functor. The source itself identifies Ri_*(L^•)=i_*(L^•) because i_* is exact: applying it termwise preserves acyclic complexes, so the canonical derived comparison is the identity represented by that complex. The three L letters have their separate single rules.

[417] EN-CONTRIB-VII-026 Source folio 436. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=450
Printed reading: the class of the complex $\underline{\underline{L}}i^*(\underline{\underline{R}}i_*(\underline{M}))$
Corrected reading: the class of the complex $\mathbb{L}i^*(\mathbb{R}i_*(\underline{M}))$
Source: ASTRA-VII-0055; ASTRA-VII-0056. ASTRA-VII-0055: The complete main upper strip clearly shows two independent lower strokes for each of these derived-functor letters. Diplomatic reproduces the strokes literally. Current retains the declared double-struck representation of the same derived functor already bound by the author. Single M and O rules remain independent. The exact local glyph count is recorded. ASTRA-VII-0056: The complete main upper strip clearly shows two independent lower strokes for each of these derived-functor letters. Diplomatic reproduces the strokes literally. Current retains the declared double-struck representation of the same derived functor already bound by the author. Single M and O rules remain independent. The exact local glyph count is recorded.

[418] EN-CONTRIB-VII-027 Source folio 436. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=450
Printed reading: a projective resolution of the complex $\underline{\underline{R}}i_*(\underline{M}) = i_*(\underline{M})$
Corrected reading: a projective resolution of the complex $\mathbb{R}i_*(\underline{M}) = i_*(\underline{M})$
Source: ASTRA-VII-0056. The complete main upper strip clearly shows two independent lower strokes for each of these derived-functor letters. Diplomatic reproduces the strokes literally. Current retains the declared double-struck representation of the same derived functor already bound by the author. Single M and O rules remain independent. The exact local glyph count is recorded.

[419] EN-CONTRIB-VII-028 Source folio 436. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=450
Printed reading: $\underline{\underline{L}}i^*(\underline{\underline{R}}i_*(\underline{M}))$ is the alternating sum
Corrected reading: $\mathbb{L}i^*(\mathbb{R}i_*(\underline{M}))$ is the alternating sum
Source: ASTRA-VII-0055; ASTRA-VII-0056. ASTRA-VII-0055: The complete main upper strip clearly shows two independent lower strokes for each of these derived-functor letters. Diplomatic reproduces the strokes literally. Current retains the declared double-struck representation of the same derived functor already bound by the author. Single M and O rules remain independent. The exact local glyph count is recorded. ASTRA-VII-0056: The complete main upper strip clearly shows two independent lower strokes for each of these derived-functor letters. Diplomatic reproduces the strokes literally. Current retains the declared double-struck representation of the same derived functor already bound by the author. Single M and O rules remain independent. The exact local glyph count is recorded.

[420] EN-CONTRIB-VII-029 Source folio 437. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=451
Printed reading: defined by an ideal of $\underline{O}_X$, canonically
Corrected reading: defined by an ideal of $\underline{O}_{X'}$, canonically
Source: check/findings/findings_436_440.json#18. Main and witness448 clearly print unprimed X. The sentence describes the invertible defining ideal of Y′ in X′, explicitly named J′⊂O_{X′} a few lines later and identified with O_{X′}(1). The canonical map from the original ideal is f*J→J′, not in general an isomorphism: for A=k[x,y], J=(x,y), on the blow-up chart A′=k[x,t], y=xt, f*J=A′²/A′(-xt,x). Its map to J′=(x) sends(a,b) to x(a+tb); the nonzero kernel is generated by(-t,1) modulo x(-t,1), hence is A′/(x). The asserted invertible ideal is therefore the displayed image J′ on X′. The existing current prime restores exactly that bound sheaf; diplomatic retains the printed O_X.

[421] EN-CONTRIB-VII-030 Source folio 437. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=451
Printed reading: $j : Y' \to X$
Corrected reading: $j : Y' \to X'$
Source: check/findings/findings_436_440.json#19. Main and witness448 print the target X without a prime. The same page's diagram explicitly binds j as Y′→X′. The composition f∘j=i∘g maps instead to X; these arrows are related by that exact composition. In the admissible blow-up of A^2_k at its origin, Y′=P^1_k and f∘j is constant to the origin: the two distinct points[1:0] and[0:1] have the same image, so this composite cannot be the asserted closed immersion. The diagram's j is the codimension1 exceptional-divisor immersion, defined on the x-chart by(x)⊂k[x,t]. Retain the existing current target X′ and diplomatic X.

[422] EN-CONTRIB-VII-031 Source folio 437. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=451
Printed reading:

$$Y' = \underline{\underline{P}}(\underline{N}).$$

Corrected reading:

$$Y' = \mathbf{P}(\underline{N}).$$

Source: ASTRA-VII-0059. The directly inspected P(N) occurrence has two separate lower P strokes. Diplomatic preserves them literally. Current retains the declared bold-P representation of Proj(Sym N); the identity graded algebra map gives the identity on its standard affine charts and glued projective bundle. The N single rule is recorded independently.

[423] EN-CONTRIB-VII-032 Source folio 438. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=452
Printed reading: $i_1 < .. < i_k$
Corrected reading: $i_1 < \cdots < i_k$
Source: check/findings/findings_436_440.json#40. The main and witness449 show exactly two compact dots between the less-than signs, whereas the adjacent wedge products have the usual three. Diplomatic keeps the two source dots; current keeps the established three-dot punctuation for the same ordered index list.

[424] EN-CONTRIB-VII-033 Source folio 440. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=454
Printed reading: $Y' = \underline{\underline{P}}(\underline{N})$
Corrected reading: $Y' = \mathbf{P}(\underline{N})$
Source: ASTRA-VII-0060. The directly inspected P(N) occurrence has two separate lower P strokes. Diplomatic preserves them literally. Current retains the declared bold-P representation of Proj(Sym N); the identity graded algebra map gives the identity on its standard affine charts and glued projective bundle. The N single rule is recorded independently.

[425] EN-CONTRIB-VII-034 Source folio 441. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=455
Printed reading: the class of the complex $\underline{\underline{L}}f^*(\underline{\underline{R}}i_*(\underline{M}))$
Corrected reading: the class of the complex $\mathbb{L}f^*(\mathbb{R}i_*(\underline{M}))$
Source: ASTRA-VII-0072; ASTRA-VII-0073. ASTRA-VII-0072: The main page and the independently viewed witness show two lower strokes at these occurrences. Diplomatic reproduces those strokes literally. Current retains the already declared double-struck notation of the identical source-bound derived functor. These replacements do not affect the singly underlined sheaf/complex letters or the plain R^i higher-direct-image letters. ASTRA-VII-0073: The main page and the independently viewed witness show two lower strokes at these occurrences. Diplomatic reproduces those strokes literally. Current retains the already declared double-struck notation of the identical source-bound derived functor. These replacements do not affect the singly underlined sheaf/complex letters or the plain R^i higher-direct-image letters.

[426] EN-CONTRIB-VII-035 Source folio 441. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=455
Printed reading: f_*
Corrected reading: f^*
Source: check/findings/findings_441_445.json#9. The complete main source and witness452 clearly place the star below f. Let P^• →i_*M be the projective resolution on X named in the sentence. The displayed derived pullback is represented on X′ by f*P^•=O_{X′}⊗_{f^{-1}O_X}f^{-1}P^•, with a⊗s↦a·f^{-1}s termwise; its homology is precisely the Tor_i^{O_X}(M,O_{X′}) written in the next sentence. The direct-image functor is related by the adjunction unit P→f_*f*P and counit f*f_*Q→Q; on an affine chart A→B these are p↦1⊗p and b⊗q↦bq. They do not replace the named tensor operation by applying f_* to P. Thus the same sentence and its next displayed homology computation uniquely specify f^*, and the low star is an established source error.

[427] EN-CONTRIB-VII-036 Source folio 441. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=455
Printed reading: $\underline{\underline{L}}f^*(\underline{\underline{R}}i_*(\underline{M}))$ is the alternating sum
Corrected reading: $\mathbb{L}f^*(\mathbb{R}i_*(\underline{M}))$ is the alternating sum
Source: ASTRA-VII-0072; ASTRA-VII-0073. ASTRA-VII-0072: The main page and the independently viewed witness show two lower strokes at these occurrences. Diplomatic reproduces those strokes literally. Current retains the already declared double-struck notation of the identical source-bound derived functor. These replacements do not affect the singly underlined sheaf/complex letters or the plain R^i higher-direct-image letters. ASTRA-VII-0073: The main page and the independently viewed witness show two lower strokes at these occurrences. Diplomatic reproduces those strokes literally. Current retains the already declared double-struck notation of the identical source-bound derived functor. These replacements do not affect the singly underlined sheaf/complex letters or the plain R^i higher-direct-image letters.

[**428**] **EN-CONTRIB-VII-037** Source folio 441. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=455
Printed reading: which was proved in exp.
Corrected reading: which was proved in exp. III.
Source: check/findings/findings_441_445.json#14. Main and witness452 end exactly with exp.;442 starts the recalled Lemma and supplies no continuation of the reference. The directly inspected III Corollary2.3 on240/PDF254 states the projective high-twist result for S quasi-compact, a projective morphism of finite-type S-schemes, and E in D^+ pseudo-coherent relative to S and acyclic above a. Substitution S=Y=X (the VII base), source=X′, E=L^•, a=0, and the projective twisting O_{X′}(1) gives exactly the recalled statement on442. Projective f makes X′ quasi-compact; perfect L^• is then bounded, and perfect f together with perfect L^• supplies the relative pseudo-coherence. The derived tensor with the invertible twist equals the ordinary tensor used on442. This directly verified dependency supports the existing current editorial completion III. It is not a claim that III is visible in the source. Diplomatic retains the incomplete exp.

[**429**] **EN-CONTRIB-VII-038** Source folio 442. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=456
Printed reading: $\underline{\underline{R}}f_*(\underline{L}^\bullet \otimes_{\underline{O}_{X'}} \underline{O}_{X'}(n))$
Corrected reading: $\mathbb{R}f_*(\underline{L}^\bullet \otimes_{\underline{O}_{X'}} \underline{O}_{X'}(n))$
Source: check/findings/findings_441_445.json#16. Both directly inspected scans show two lower strokes. The local diplomatic patch preserves the strokes literally; current retains the declared double-struck derived-functor notation. The neighboring R^i higher-direct-image letters are visibly plain.

[**430**] **EN-CONTRIB-VII-039** Source folio 442. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=456
Printed reading: $\underline{\underline{R}}f_*(\underline{O}_{Y'}(n)) = i_* \circ \underline{\underline{R}}g_*(\underline{O}_{Y'}(n))$
Corrected reading: $\mathbb{R}f_*(\underline{O}_{Y'}(n)) = i_* \circ \mathbb{R}g_*(\underline{O}_{Y'}(n))$
Source: check/findings/findings_441_445.json#18; check/findings/findings_441_445.json#19. check/findings/findings_441_445.json#18: Both directly inspected scans show two lower strokes. The local diplomatic patch preserves the strokes literally; current retains the declared double-struck derived-functor notation. The neighboring R^i higher-direct-image letters are visibly plain. check/findings/findings_441_445.json#19: Both directly inspected scans show two lower strokes. The local diplomatic patch preserves the strokes literally; current retains the declared double-struck derived-functor notation. The neighboring R^i higher-direct-image letters are visibly plain.

[**431**] **EN-CONTRIB-VII-040** Source folio 442. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=456
Printed reading: $Y' = \underline{\underline{P}}(\underline{N})$
Corrected reading: $Y' = \mathbf{P}(\underline{N})$
Source: ASTRA-VII-0075. Each counted P in the full main page has two lower rules. Diplomatic records their literal shape. Current keeps the declared bold notation of the same Proj construction; its identity graded-algebra morphism induces identity maps on the standard degree-zero localization charts and hence on the glued projective scheme.

[432] EN-CONTRIB-VII-041 Source folio 443. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=457
Printed reading: $X' \longrightarrow \underline{\underline{P}}_X^{d-1}$
Corrected reading: $X' \longrightarrow \mathbf{P}_X^{d-1}$
Source: ASTRA-VII-0076. Each counted P in the full main page has two lower rules. Diplomatic records their literal shape. Current keeps the declared bold notation of the same Proj construction; its identity graded-algebra morphism induces identity maps on the standard degree-zero localization charts and hence on the glued projective scheme.

[433] EN-CONTRIB-VII-042 Source folio 443. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=457
Printed reading: On $\underline{\underline{P}}_X^{d-1}$ we obtain
Corrected reading: On $\mathbf{P}_X^{d-1}$ we obtain
Source: ASTRA-VII-0076. Each counted P in the full main page has two lower rules. Diplomatic records their literal shape. Current keeps the declared bold notation of the same Proj construction; its identity graded-algebra morphism induces identity maps on the standard degree-zero localization charts and hence on the glued projective scheme.

[434] EN-CONTRIB-VII-043 Source folio 443. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=457
Printed reading: so these are free of finite type
Corrected reading: so these are locally free of finite type
Source: ASTRA-VII-0084. Write B=O_X[T_1,...,T_d]. A finite homogeneous basis e_j of degree a_j gives the degree-preserving isomorphism φ:⊕_j B(−a_j)→L, (b_j)↦Σ_jb_je_j, with inverse the unique homogeneous coordinates. Taking associated sheaves gives ⊕_jO_P(−a_j)≅Proj(L). On D_+(T_i), the degree-zero generators e_j/T_i^{a_j} give explicit local bases and prove finite local freeness. Global freeness is false for an admissible instance: A=k[x,y], J=(x,y), S=A[T_1,T_2]/(yT_1−xT_2), with presentation B(−1)→B→S. Its sheaves are M′′=O_{P^1_A}(−1), M′=O. On the closed fibre P^1_k, choose frames e_0=(1,t), e_1=(s,1)=t^{-1}e_0. A global section requires a(t)=t^{-1}b(t^{-1}); the nonnegative and strictly negative Laurent powers force both sides to vanish. Hence H^0(O(−1))=0, whereas a free rank-one sheaf has H^0=k. The surrounding proof uses exactly finite local freeness and the graded direct sum of twists: regular X′→P makes O_{X′} perfect; the two exact kernel sequences with M′ and M′′ therefore give perfect kernels, to which the recalled projective high-twist lemma applies. No global free basis is used or available. Add only localement, independently of the separate all-n diagram issue.

[435] EN-CONTRIB-VII-044 Source folio 443. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=457
Printed reading: over $\underline{\underline{P}}_X^{d-1}$. We can therefore
Corrected reading: over $\mathbf{P}_X^{d-1}$. We can therefore
Source: ASTRA-VII-0076. Each counted P in the full main page has two lower rules. Diplomatic records their literal shape. Current keeps the declared bold notation of the same Proj construction; its identity graded-algebra morphism induces identity maps on the standard degree-zero localization charts and hence on the glued projective scheme.

[436] EN-CONTRIB-VII-045 Source folio 443. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=457
Printed reading: $\underline{\underline{P}}_X^{d-1} \to X$
Corrected reading: $\mathbf{P}_X^{d-1} \to X$
Source: ASTRA-VII-0076. Each counted P in the full main page has two lower rules. Diplomatic records their literal shape. Current keeps the declared bold notation of the same Proj construction; its identity graded-algebra morphism induces identity maps on the standard degree-zero localization charts and hence on the glued projective scheme.

[437] EN-CONTRIB-VII-046 Source folio 444. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=458
Printed reading:

$$f_*(\underline{O}_{Y'}(n)) = i_*(g_*(\underline{O}_{Y'}(n))) = i_*(\underline{\underline{S}}(\underline{N})_n) = \underline{J}^n/\underline{J}^{n+1}.$$

Corrected reading:

$$f_*(\underline{O}_{Y'}(n)) = i_*(g_*(\underline{O}_{Y'}(n))) = i_*(\mathbf{S}(\underline{N})_n) = \underline{J}^n/\underline{J}^{n+1}.$$

Source: check/findings/findings_441_445.json#24. The complete main upper strip shows two distinct lower strokes under S and a single stroke under N. Diplomatic now preserves the S strokes literally. Current retains the declared bold S notation of the same symmetric algebra; the identity on N extends degreewise by the symmetric-algebra universal property and identifies the expression with its original source-bound algebra.

[438] EN-CONTRIB-VII-047 Source folio 444. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=458
Printed reading: hence the middle one is also an isomorphism
Corrected reading: hence the middle one is also an isomorphism.
Source: fixes/MISPRINTS_6.json#413. The source line ends immediately after également, and the next line is the separately spaced proposition label. Diplomatic correctly preserves the omission; current already supplies the sentence-ending full stop.

[439] EN-CONTRIB-VII-048 Source folio 445. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=459
Printed reading: (VI 5.9) which gives the result.
Corrected reading: (VI 5.9), which gives the result.
Source: fixes/MISPRINTS_6.json#414. The main lower strip ends the line immediately after the parenthesis; the next line begins d'où le résultat. Diplomatic preserves the printed absence. Current keeps its existing comma separating the consequent clause.

[440] EN-CONTRIB-VII-049 Source folio 445. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=459
Printed reading:

$$v \circ u(y) = v\big(g^*(y)\lambda_{-1}(F), -i_*(y)\big) = j_*\big(g^*(y)\lambda_{-1}(F)\big) - f^*(i_*(x)) = 0,$$


Corrected reading:

$$v \circ u(y) = v\big(g^*(y)\lambda_{-1}(F), -i_*(y)\big) = j_*\big(g^*(y)\lambda_{-1}(F)\big) - f^*(i_*(y)) = 0,$$


Source: fixes/RECONCILE_CODEX_6.json#197. The main lower strip clearly prints x. The same page defines u(y)=(g*y·λ_{−1}(F),−i_*y) and v(y′,x)=j_*y′+f*x. Substitution of this ordered pair into the explicitly defined v gives v(u(y))=j_*(g*y·λ_{−1}(F))−f*(i_*y), which is zero by3.4. Thus the final argument must be the same y entering u; no unspecified x occurs in that substitution. The existing current y is the exact correction; diplomatic retains x.

[441] EN-CONTRIB-VII-050 Source folio 446. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=460
Printed reading: $j_*(y') = f^*(x)$
Corrected reading: $j_*(y') = -f^*(x)$
Source: fixes/RECONCILE_CODEX_6.json#198. The complete main page prints the positive sign here, although the opening hypothesis is j_*y′+f*x=0 and the preceding paragraph explicitly uses f*x=−j_*y′. Solving that same group equality gives j_*y′=−f*x. Substitution of the established x=−i_*y then gives j_*y′=f*i_*y, exactly the next displayed term. The existing current minus is therefore forced by the author's exact preceding and following equalities.

[442] EN-CONTRIB-VII-051 Source folio 447. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=461
Printed reading:



Corrected reading:



Source: ASTRA-VII-0095; check/findings/findings_446_450.json#9. ASTRA-VII-0095: MainPDF461 and witness458 both clearly show the printed clockwise arrows. The adjunction Lf*⊣Rf_* supplies the counit ε:T=Lf*Rf_*M→M; on affine tensor representatives its evaluation is b⊗m↦bm. With N=Cone(ε), N^n=M^n⊕T^{n+1} and d_N(m,t)=(d_Mm+ε(t),−d_Tt). The maps are ε:T→M, m↦(m,0):M→N, and (m,t)↦t:N→T[1]. Thus the diagram reverses all three arrows, preserving the author's cyclic-triangle convention without adding a new shift label. The triangle gives [N]=[M]−[T]=x−f*f_*x, exactly the final class on447. Every prose reference has been checked: the counit is an isomorphism where f is an isomorphism, so N cohomology is supported on Y′; the preceding perfect-support argument puts its class in Im(j_*), giving the stated surjectivity. The printed orientation is refuted within the author's hypotheses by f:Bl_0 A^2_k→A^2_k and M=O(E), E the exceptional divisor. The sequence0→O→O(E)→j_*O_{P^1}(−1)→0 gives Rf_*M≅O: H^0(P^1,O(−1))=0 by the chart equation a(t)=t^{-1}b(t^{-1}), and H^1=0 because k[t,t^{-1}]=k[t]+t^{-1}k[t^{-1}]. Lemma3.5 supplies Rf_*O=O. Hence T≅O, and every M→T is a section of O(−E), identified by3.5 with J=(x,y)⊂k[x,y]. Such a section cannot be invertible off the origin: the units of k[x,y]_x are k×x^Z, those of k[x,y]_y are k×y^Z, and their intersection in k(x,y) is k×. No element of(x,y) lies there. Thus there is no arrow in the printed direction with the claimed restriction. In the corrected direction the counit is the inclusion O→O(E), which is an isomorphism on that complement. check/findings/findings_446_450.json#9: Main and witness458 show two distinct lower strokes under each derived-functor letter and single M/N rules. Diplomatic restores the literal derived-letter rules; current retains the declared double-struck representations of those same functors.

[443] EN-CONTRIB-VII-052 Source folio 449. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=463
Printed reading: $s : X \longrightarrow \underline{\underline{P}}^r_X$
Corrected reading: $s : X \longrightarrow \mathbf{P}^r_X$
Source: ASTRA-VII-0097. The main page and witness460 show two independent lower strokes on all five P letters, including both nested bundle letters in the last composition. Diplomatic restores those strokes literally. Current retains the declared bold P for the identical Proj construction; the identity graded algebra map induces the identity on its affine localization charts.

[444] EN-CONTRIB-VII-053 Source folio 449. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=463
Printed reading: $\underline{\underline{P}}^r_X$
Corrected reading: $\mathbf{P}^r_X$
Source: ASTRA-VII-0097. The main page and witness460 show two independent lower strokes on all five P letters, including both nested bundle letters in the last composition. Diplomatic restores those strokes literally. Current retains the declared bold P for the identical Proj construction; the identity graded algebra map induces the identity on its affine localization charts.

[445] EN-CONTRIB-VII-054 Source folio 449. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=463
Printed reading: $\underline{O}_p$
Corrected reading: $\underline{O}_P$
Source: ASTRA-VII-0098. The native witness460 crop shows a lowercase p with the same bowl, descender and narrow foot as the p in correspondant on the same line. It is not the capital P of the projective-space labels. Restore this exact glyph diplomatically. The structure sheaf in this construction is that of P=P_X^r; on D_+(T_0) its ring is A[t_1,...,t_r] and the section pulls it back by t_j↦0 to A. That explicit map fixes the scheme indexing the ambient sheaf and supports the existing current capital P. The following451 independently binds lowercase p as the structural projection P_X^1→X; the source typo does not introduce another ambient scheme.

[446] EN-CONTRIB-VII-055 Source folio 449. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=463
Printed reading: $X \to \underline{\underline{P}}^2_X$
Corrected reading: $X \to \mathbf{P}^2_X$
Source: ASTRA-VII-0097. The main page and witness460 show two independent lower strokes on all five P letters, including both nested bundle letters in the last composition. Diplomatic restores those strokes literally. Current retains the declared bold P for the identical Proj construction; the identity graded algebra map induces the identity on its affine localization charts.

[447] EN-CONTRIB-VII-056 Source folio 449. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=463
Printed reading: $X \to \underline{\underline{P}}^1_{\underline{\underline{P}}^1_X}$
Corrected reading: $X \to \mathbf{P}^1_{\mathbf{P}^1_X}$
Source: ASTRA-VII-0097. The main page and witness460 show two independent lower strokes on all five P letters, including both nested bundle letters in the last composition. Diplomatic restores those strokes literally. Current retains the declared bold P for the identical Proj construction; the identity graded algebra map induces the identity on its affine localization charts.

[448] EN-CONTRIB-VII-057 Source folio 451. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=465
Printed reading: $X \to \underline{\underline{P}}^1_X$
Corrected reading: $X \to \mathbf{P}^1_X$
Source: ASTRA-VII-0102. The whole main page contains eight projective-space P letters, including both letters in the nested projective bundle. Every one has two lower rules. Diplomatic preserves those strokes literally. Current keeps the established bold notation for the same projective scheme, whose identity graded-algebra map gives identity maps of the standard affine charts. The additional capital P in the corrected O_P subscript is separately justified by the O_p source correction.

[449] EN-CONTRIB-VII-058 Source folio 451. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=465
Printed reading: $\underline{\underline{P}}^1_X \to \underline{\underline{P}}^1_{\underline{\underline{P}}^1_X}$
Corrected reading: $\mathbf{P}^1_X \to \mathbf{P}^1_{\mathbf{P}^1_X}$
Source: ASTRA-VII-0102. The whole main page contains eight projective-space P letters, including both letters in the nested projective bundle. Every one has two lower rules. Diplomatic preserves those strokes literally. Current keeps the established bold notation for the same projective scheme, whose identity graded-algebra map gives identity maps of the standard affine charts. The additional capital P in the corrected O_P subscript is separately justified by the O_p source correction.

[450] EN-CONTRIB-VII-059 Source folio 451. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=465
Printed reading: $\underline{\underline{P}}^1_X$
Corrected reading: $\mathbf{P}^1_X$
Source: ASTRA-VII-0102. The whole main page contains eight projective-space P letters, including both letters in the nested projective bundle. Every one has two lower rules. Diplomatic preserves those strokes literally. Current keeps the established bold notation for the same projective scheme, whose identity graded-algebra map gives identity maps of the standard affine charts. The additional capital P in the corrected O_P subscript is separately justified by the O_p source correction.

[451] EN-CONTRIB-VII-060 Source folio 451. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=465
Printed reading: $s : X \to \underline{\underline{P}}^1_X$
Corrected reading: $s : X \to \mathbf{P}^1_X$
Source: ASTRA-VII-0102. The whole main page contains eight projective-space P letters, including both letters in the nested projective bundle. Every one has two lower rules. Diplomatic preserves those strokes literally. Current keeps the established bold notation for the same projective scheme, whose identity graded-algebra map gives identity maps of the standard affine charts. The additional capital P in the corrected O_P subscript is separately justified by the O_p source correction.

[452] EN-CONTRIB-VII-061 Source folio 451. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=465
Printed reading: $\underline{O_p}(-1)$
Corrected reading: $\underline{O_{\mathbf{P}}}(-1)$
Source: fixes/MISPRINTS_6.json#439. Main and witness462 show a lowercase p with a descender; the O has a separate short rule. Diplomatic keeps that printed glyph. The same paragraph explicitly binds p:P_X^1→X as the structural map, while the ideal is the sheaf of the section s:X→P_X^1. Its local map O_P(−1)→O_P is multiplication by the homogeneous coordinate vanishing on s, and on D_+(T_0) it is multiplication by t=T_1/T_0. This exact sheaf map identifies the named ambient sheaf as O_P(−1), supporting the already present current capital projective-space P. The source lowercase p is retained only diplomatically.

[453] EN-CONTRIB-VII-062 Source folio 451. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=465
Printed reading: $K^{\bullet}(\underline{\underline{P}}^1_X)$
Corrected reading: $K^{\bullet}(\mathbf{P}^1_X)$
Source: ASTRA-VII-0102. The whole main page contains eight projective-space P letters, including both letters in the nested projective bundle. Every one has two lower rules. Diplomatic preserves those strokes literally. Current keeps the established bold notation for the same projective scheme, whose identity graded-algebra map gives identity maps of the standard affine charts. The additional capital P in the corrected O_P subscript is separately justified by the O_p source correction.

[454] EN-CONTRIB-VII-063 Source folio 451. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=465
Printed reading: $\underline{\underline{P}}^1_X \to X$
Corrected reading: $\mathbf{P}^1_X \to X$
Source: ASTRA-VII-0102. The whole main page contains eight projective-space P letters, including both letters in the nested projective bundle. Every one has two lower rules. Diplomatic preserves those strokes literally. Current keeps the established bold notation for the same projective scheme, whose identity graded-algebra map gives identity maps of the standard affine charts. The additional capital P in the corrected O_P subscript is separately justified by the O_p source correction.

[455] EN-CONTRIB-VII-064 Source folio 453. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=467
Printed reading:

$$i_*(\gamma^p(N,y)) \equiv \gamma^p(i'_*(y)) \quad \text{mod. torsion.}$$

Corrected reading:

$$i_*(\gamma^p(N,y)) \equiv \gamma^p(i_*(y)) \quad \text{mod. torsion.}$$

Source: fixes/RECONCILE_CODEX_6.json#214. The main scan shows the small stroke, and witness464 separates it from the i dot. The previous theorem binds only i:Y→X. The precise derivation also fixes the pushforward: γ^p(N,y)=Σ_{j=1}^p binom(p−1,j−1)λ^j(N,y), so additivity of i_* and4.3 give i_*γ^p(N,y)≡Σ binom(p−1,j−1)λ^j(i_*y)=γ^p(i_*y). Thus the same i_* occurs on both sides; the existing current removal of the prime is the exact correction, with the source prime retained diplomatically.

[456] EN-CONTRIB-VII-065 Source folio 453. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=467
Printed reading: follows immediately from 4.3, and using
Corrected reading: follows immediately from 4.3, using
Source: fixes/MISPRINTS_6.json#441. Main and witness464 clearly print et utilisant. The finite verb résulte is followed here by an instrumental gerund, which requires en utilisant; et does not coordinate a second finite predicate in this sentence. The attached current layer still has the source slip.

[457] EN-CONTRIB-VII-066 Source folio 454. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=468
Printed reading: by decomposing X into a finite sum
Corrected reading: by decomposing Y into a finite sum
Source: ASTRA-VII-0112. The same paragraph explicitly defines E_α as O_Y-vector bundles. Their simultaneous rank loci form a finite open-and-closed partition Y=⊔_aY_a: each rank function is locally constant, there are finitely many bundles, and Y is quasi-compact. Restriction gives the λ-ring isomorphism ρ:K(Y)→∏_aK(Y_a), [E]↦([E|Y_a])_a; its inverse sends a tuple to the sum of extensions by zero along the open-and-closed inclusions. The compositions are identity by the direct-sum decomposition of sheaves on the finite disjoint union. Exterior powers are componentwise, so ρ and its inverse preserve the λ operations and their Fil filtration; tensoring these exact maps with Q gives the rational version used here. For i_a:Y_a→X, the equality i_*(z)=Σ_a(i_a)_*(z|Y_a) follows from the direct sum of the corresponding pushed-forward perfect complexes. Each i_a is a closed regular immersion of codimensiond: near any point of Y_a, remove the other closed components of Y and use the same regular sequence defining i. The proof can therefore be carried out on each Y_a with the same target X and summed. The printed partition of X does not provide this reduction: take connected X=P^1_k, regular divisor Y=$\{0,\infty\}$, and the vector bundle E=(k,0) on its two components. Any finite open-and-closed partition of connected X has only one nonempty piece, and E still has ranks1 and0 on Y. This admissible example refutes the printed reduction. Replacing only X by Y gives exactly the finite-product argument already invoked by the author.

[458] EN-CONTRIB-VII-067 Source folio 455. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=469
Printed reading:

$$i_*(x) \in \mathrm{Fil}^{k+d} K^\bullet(X),$$

Corrected reading:

$$i_*(x) \in \mathrm{Fil}^{k+d} K^\bullet(X)_{\mathbb{Q}},$$

Source: ASTRA-VII-0113. MainPDF469 and witness466 show both omissions; the two separately viewed native-resolution witness crops show empty space where the subscript would stand. The five neighboring target groups carry Q, and454 explicitly defines x∈Fil^kK(Y)_Q and R as a Q-algebra. The precise comparison is the rationalization homomorphism ι:K(X)→K(X)⊗_ZQ, a↦a⊗1; the rational filtration is generated by the image of Fil^mK(X) and its rational multiples. For the allowed regular immersion i:Spec k→P^1_k, d=1, choose k_index=0 and x=y=1/2 in R=Q, using the induction index i_index=0. Both disputed left sides equal[O_point]/2 because γ^1 is the identity. The Euler characteristic homomorphism χ:K(P^1_k)→Z sends[O_point] to1, so its rational extension gives χ_Q([O_point]/2)=1/2. An element of im(ι) has integral Euler characteristic. Thus this class is not in the image of the integral group, and in particular not in the two printed integral filtration subgroups. It does belong to Fil^1K(P^1_k)_Q, since[O_point]=[O]−[O(−1)] has rank zero and γ^1 fixes it. The two Q insertions restore the precise rational target already specified by the theorem and the surrounding induction, with no change to its hypotheses.

[459] EN-CONTRIB-VII-068 Source folio 455. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=469
Printed reading: $\gamma^{d+i}(i_*(y)) \in \mathrm{Fil}^{d+i} K^\bullet(X)$
Corrected reading: $\gamma^{d+i}(i_*(y)) \in \mathrm{Fil}^{d+i} K^\bullet(X)_{\mathbb{Q}}$
Source: ASTRA-VII-0113. MainPDF469 and witness466 show both omissions; the two separately viewed native-resolution witness crops show empty space where the subscript would stand. The five neighboring target groups carry Q, and454 explicitly defines x∈Fil^kK(Y)_Q and R as a Q-algebra. The precise comparison is the rationalization homomorphism ι:K(X)→K(X)⊗_ZQ, a↦a⊗1; the rational filtration is generated by the image of Fil^mK(X) and its rational multiples. For the allowed regular immersion i:Spec k→P^1_k, d=1, choose k_index=0 and x=y=1/2 in R=Q, using the induction index i_index=0. Both disputed left sides equal[O_point]/2 because γ^1 is the identity. The Euler characteristic homomorphism χ:K(P^1_k)→Z sends[O_point] to1, so its rational extension gives χ_Q([O_point]/2)=1/2. An element of im(ι) has integral Euler characteristic. Thus this class is not in the image of the integral group, and in particular not in the two printed integral filtration subgroups. It does belong to Fil^1K(P^1_k)_Q, since[O_point]=[O]−[O(−1)] has rank zero and γ^1 fixes it. The two Q insertions restore the precise rational target already specified by the theorem and the surrounding induction, with no change to its hypotheses.

[460] EN-CONTRIB-VII-069 Source folio 456. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=470
Printed reading: (relation (4.3.1)
Corrected reading: (relation (4.3.1)).
Source: fixes/MISPRINTS_6.json#430. The main lower strip and witness467 both show only the inner closing parenthesis. The outer parenthetical phrase remains open. Current already supplies its closing parenthesis and the sentence stop; diplomatic preserves the two printed omissions.

[461] EN-CONTRIB-VII-070 Source folio 456. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=470
Printed reading: First we use VI.I.1 to construct
Corrected reading: First we use VI.1.1 to construct
Source: fixes/MISPRINTS_6.json#431. Witness467 clearly shows the seriffed I between two stops, while the final character is the flagged digit1. I also directly inspected mainPDF379, folio365: the relevant projective-bundle theorem is explicitly numbered1.1 and gives the free basis1,...,ξ^r over the base K-ring. Its structural pullback f* is the geometric model for g*:K→K1 here. This directly read target supports the existing current VI.1.1; diplomatic retains VI.I.1.

[462] EN-CONTRIB-VII-071 Source folio 456. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=470
Printed reading: $^{i}(L) = 0$
Corrected reading: $\lambda^i(L) = 0$
Source: fixes/MISPRINTS_6.json#432. The main image and witness467 show a clear blank, with no lambda stroke, before the superscript i. Earlier on this same page K[L] is explicitly the lambda-algebra on the line element L. Its defining total operation is λ_t(L)=1+Lt, so λ^i(L)=0 for every i≥2. This exact coefficient identity fixes the missing operator. Current already has λ; diplomatic retains the blank. The root-owned manual record is not duplicated here.

[463] EN-CONTRIB-VII-072 Source folio 457. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=471

Printed reading: $1 + \widehat{K/(P)[[t]]}^{+}$

Corrected reading: $1 + \widehat{K[L]/(P)[[t]]}^{+}$

Source: ASTRA-VII-0118. MainPDF471 and witness468, including three separately viewed native-resolution crops, confirm the three omissions. The preceding page explicitly fixes the coefficient lambda-ring K, the polynomial extension K[L], and P(L)=Σ_{i=0}^d(−1)^iλ^{d−i}(N)L^i. The exact construction is the inclusion K→K[L] followed by the quotient ρ:K[L]→A=K[L]/(P); coefficientwise reduction gives ρ[[t]]:K[L][[t]]→A[[t]], carrying λ_t(N−L) to the series used in the first expression. The second expression is the source series ring of this very map. P has leading coefficient(−1)^d, a unit; multiplying P by that unit gives a monic polynomial without changing its ideal. Monic division supplies the free K-basis1,L,...,L^{d−1} of A, exactly as claimed immediately after the third expression. The same paragraph explicitly retains the coefficient K in g*:K→K1 and in the free K-module assertion; no renaming of that K is made. For a concrete instance satisfying the abstract hypotheses, take K=K′=Z with the binomial lambda-structure, N=2,d=2,i*=Id,i_*=0. Projection and the lambda compatibility are immediate, and i*i_*=0 agrees with multiplication by λ_{−1}(2)=(1−1)^2=0. Here P=(1−L)^2, and A≅Z[ε]/(ε²) via L↦1+ε and ε↦L−1. These maps are inverse and identify the claimed free rank-two K-module. P is not an element of Z; moreover every quotient of Z has cyclic additive group, while this A has additive basis1,ε. Thus the three printed ambient rings cannot express the stated construction with the retained coefficient K. The minimal three[L] insertions recover the exact quotient and series maps already used by the author, leaving every other K unchanged.

[464] EN-CONTRIB-VII-073 Source folio 457. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=471

Printed reading: $1 + \widehat{K[[t]]}^{+}$

Corrected reading: $1 + \widehat{K[L][[t]]}^{+}$

Source: ASTRA-VII-0118. MainPDF471 and witness468, including three separately viewed native-resolution crops, confirm the three omissions. The preceding page explicitly fixes the coefficient lambda-ring K, the polynomial extension K[L], and P(L)=Σ_{i=0}^d(−1)^iλ^{d−i}(N)L^i. The exact construction is the inclusion K→K[L] followed by the quotient ρ:K[L]→A=K[L]/(P); coefficientwise reduction gives ρ[[t]]:K[L][[t]]→A[[t]], carrying λ_t(N−L) to the series used in the first expression. The second expression is the source series ring of this very map. P has leading coefficient(−1)^d, a unit; multiplying P by that unit gives a monic polynomial without changing its ideal. Monic division supplies the free K-basis1,L,...,L^{d−1} of A, exactly as claimed immediately after the third expression. The same paragraph explicitly retains the coefficient K in g*:K→K1 and in the free K-module assertion; no renaming of that K is made. For a concrete instance satisfying the abstract hypotheses, take K=K′=Z with the binomial lambda-structure, N=2,d=2,i*=Id,i_*=0. Projection and the lambda compatibility are immediate, and i*i_*=0 agrees with multiplication by λ_{−1}(2)=(1−1)^2=0. Here P=(1−L)^2, and A≅Z[ε]/(ε²) via L↦1+ε and ε↦L−1. These maps are inverse and identify the claimed free rank-two K-module. P is not an element of Z; moreover every quotient of Z has cyclic additive group, while this A has additive basis1,ε. Thus the three printed ambient rings cannot express the stated construction with the retained coefficient K. The minimal three[L] insertions recover the exact quotient and series maps already used by the author, leaving every other K unchanged.

[465] EN-CONTRIB-VII-074 Source folio 457. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=471
Printed reading: $K_1 = K/(P)$
Corrected reading: $K_1 = K[L]/(P)$
Source: ASTRA-VII-0118. MainPDF471 and witness468, including three separately viewed native-resolution crops, confirm the three omissions. The preceding page explicitly fixes the coefficient lambda-ring K, the polynomial extension K[L], and P(L)=Σ_{i=0}^d(−1)^iλ^{d−i}(N)L^i. The exact construction is the inclusion K→K[L] followed by the quotient ρ:K[L]→A=K[L]/(P); coefficientwise reduction gives ρ[[t]]:K[L][[t]]→A[[t]], carrying λ_t(N−L) to the series used in the first expression. The second expression is the source series ring of this very map. P has leading coefficient(−1)^d, a unit; multiplying P by that unit gives a monic polynomial without changing its ideal. Monic division supplies the free K-basis1,L,...,L^{d−1} of A, exactly as claimed immediately after the third expression. The same paragraph explicitly retains the coefficient K in g*:K→K1 and in the free K-module assertion; no renaming of that K is made. For a concrete instance satisfying the abstract hypotheses, take K=K′=Z with the binomial lambda-structure, N=2,d=2,i*=Id,i_*=0. Projection and the lambda compatibility are immediate, and i*i_*=0 agrees with multiplication by λ_{−1}(2)=(1−1)^2=0. Here P=(1−L)^2, and A≅Z[ε]/(ε²) via L↦1+ε and ε↦L−1. These maps are inverse and identify the claimed free rank-two K-module. P is not an element of Z; moreover every quotient of Z has cyclic additive group, while this A has additive basis1,ε. Thus the three printed ambient rings cannot express the stated construction with the retained coefficient K. The minimal three[L] insertions recover the exact quotient and series maps already used by the author, leaving every other K unchanged.

[466] EN-CONTRIB-VII-075 Source folio 461. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=475
Printed reading: $Y' = \underline{\underline{P}}(\underline{N})$
Corrected reading: $Y' = \mathbf{P}(\underline{N})$
Source: ASTRA-VII-0125. The main source and witness472 show two lower strokes on this single P, independent of the one rule under N. The current bold letter names the same projective bundle: the identity of the graded symmetric algebra induces identity homomorphisms on every degree-zero localization defining its Proj charts.

[467] EN-CONTRIB-VII-076 Source folio 461. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=475
Printed reading:

$$j_*(y') + f^*(x) \in \mathrm{Fil}^{k+1}\, K^\bullet(X')_{\mathbb{Q}}, \tag{4.6.1}$$

Corrected reading:

$$j_*(y') + f^*(x) \in \mathrm{Fil}^{k+1}\, K^\bullet(X')_{\mathbb{Q}}, \tag{4.8.1}$$

Source: fixes/MISPRINTS_6.json#416. Main and witness472 plainly print4.6.1. The formula is the assumption j_*y′+f*x∈Fil^{k+1}K(X′)_Q in the proof of4.8. The first line of462 explicitly cites it as(4.8.1), and applying f_* gives exactly f_*j_*y′+f_*f*x∈Fil^{k+1}K(X)_Q, the next displayed line there. Remarque4.6.1 already labels a different statement on455. Thus the printed462 citation identifies the intended equation number4.8.1 and its exact map, supporting the current label repair.

[468] EN-CONTRIB-VII-077 Source folio 461. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=475
Printed reading: Then by VI 5.8, $y \in \mathrm{Fil}^k K^\bullet(Y)$
Corrected reading: Then by VI 5.8, $y \in \mathrm{Fil}^{k-d} K^\bullet(Y)_{\mathbb{Q}}$
Source: fixes/MISPRINTS_6.json#417; ASTRA-VII-0133. fixes/MISPRINTS_6.json#417: Main and witness472, including the native filtration crop, show only k. The same page explicitly states g_*(Fil^{k−1}K(Y′))⊂Fil^{k−d}K(Y), then passes to its rational tensor map. Since y′ was chosen in Fil^{k−1}K(Y′)_Q, applying that exact map gives y=g_*y′ in degree k−d. The earlier sentence asks for precisely this degree. A concrete counterexample to the printed exponent also satisfies the kernel assumption: let X=A^1 over a field, Y={0}, d=1, and choose index k=1. The Rees algebra of(t) is A[T], whose Proj is Spec A by its single D_+(T) chart, so the blow-up f and g:Y′→Y are identities. Choose y′=1/2 and x=0. The perfect resolution0→O_X –t→ O_X→O_Y→0 gives j_*y′=0, hence the required kernel expression is0∈Fil^2. But y=1/2, while Fil^1K(Y)=0 because the rank homomorphism K(Spec field)=Z→Z is the identity. The corrected degree is Fil^0K(Y)_Q=Q. The independently missing rational subscript is recorded separately. ASTRA-VII-0133: The native witness472 crop shows no subscript after K•(Y). The input y′ on the same page lies in Fil^{k−1}K(Y′)_Q; the map used is g_*⊗id_Q on the tensorized filtration groups, explicitly introduced in the preceding paragraph. It sends a⊗q to g_*(a)⊗q and gives the target Fil^{k−d}K(Y)_Q already required by the existence statement. The principal-divisor example recorded with checker461_465#5 uses X=A^1,Y={0},d=1,k=1,y′=1/2,x=0 and g=Id, and satisfies the displayed kernel assumption through the two-term resolution of O_Y. It yields y=1/2 in K(Y)_Q=Q. The canonical map K(Y)=Z→Z⊗Q=Q sends n to n; its image does not contain1/2. Thus changing the exponent alone does not correct this particular target. This independent Q insertion restores the codomain of the already invoked tensor map; the diplomatic omission is retained.

[469] EN-CONTRIB-VII-078 Source folio 462. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=476
Printed reading: Taking (4.6.1) into account, we therefore obtain:
Corrected reading: Taking (4.8.1) into account, we therefore obtain:
Source: fixes/MISPRINTS_6.json#419. The main upper strip prints4.6.1 here, matching the wrong label on461. The invoked formula is exactly the kernel assumption j_*y′+f*x∈Fil^{k+1}: subtracting f*(i_*y)+f*x∈Fil^{k+1} gives j_*y′−f*(i_*y)=j_*z∈Fil^{k+1}. This explicit subtraction identifies the same equation already cited as4.8.1 at the page opening. Link this current reference to the independently repaired label4.8.1.

[470] EN-CONTRIB-VII-079 Source folio 465. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=479
Printed reading: the exact sequence (IV :
Corrected reading: the exact sequence (IV):
Source: fixes/MISPRINTS_6.json#423. Main and witness476 show an open parenthesis before IV, then a long blank and colon without a closing parenthesis. The adjacent464 references close their parentheses. Diplomatic retains the source omission; current already restores only the closing parenthesis and leaves the unknown locator blank.

[471] EN-CONTRIB-VII-080 Source folio 465. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=479
Printed reading: Consequently, x can be expressed
Corrected reading: Consequently, y can be expressed
Source: fixes/MISPRINTS_6.json#424. Main and witness476 both show x. The source first chooses x with u*(1_Z)=u*(x), then chooses y in the image of j_* from the closed complement Z′. Exactness of K(Z′) –j_*→ K(X) –u*→ K(U) identifies the needed supported correction as y=cl(O_Z)−x; the same identity holds after tensoring every group and map with Q. For η=Σa_α[O_{Z″_α}] in K(Z′), the exact direct-image map sends it to y=Σa_α[j_*O_{Z″_α}], with each support inside Z′⊂Z and avoiding the generic point z. Thus each support has codimension>k and the induction applies to y. Adding y to the displayed cl(O_Z)−y∈Fil^k gives cl(O_Z)∈Fil^k, exactly the desired conclusion. The printed x assertion fails in an instance with no coefficient ambiguity: take X=P^1 over a field and Z one rational point, and choose U=X because Z is already a regular Cartier divisor. Then Z′ is empty, u=Id, x=cl(O_Z), and y=0. Euler characteristic sends x to1, so x is nonzero, whereas every combination of sheaf classes on the empty Z′ is0. For the second printed assertion put h=1−[O(−1)]=cl(O_Z). The projective-bundle presentation gives K(P^1)=Z[h]/(h^2). Its total gamma operation is γ_t(h)=1/(1−ht)=1+ht, and for any rank-zero element nh it is (1−ht)^(−n)=1+nht. Hence every gamma generator of degree≥2 and every product of two positive-degree generators is0, so Fil^2_gamma K(P^1)_Q=0; x=h remains nonzero after rationalization because χ_Q(h)=1. This also directly refutes the final printed x∈Fil^2. The two y substitutions restore the exact localization argument, with no change to its hypotheses or surrounding x used for the lift.

[472] EN-CONTRIB-VII-081 Source folio 465. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=479
Printed reading: $x \in \mathrm{Fil}^{k+1} K^\bullet(X)_\mathbb{Q}$
Corrected reading: $y \in \mathrm{Fil}^{k+1} K^\bullet(X)_\mathbb{Q}$
Source: fixes/MISPRINTS_6.json#425. Main and witness476 both show x. The source first chooses x with u*(1_Z)=u*(x), then chooses y in the image of j_* from the closed complement Z′. Exactness of K(Z′) –j_*→ K(X) –u*→ K(U) identifies the needed supported correction as y=cl(O_Z)−x; the same identity holds after tensoring every group and map with Q. For η=Σa_α[O_{Z″_α}] in K(Z′), the exact direct-image map sends it to y=Σa_α[j_*O_{Z″_α}], with each support inside Z′⊂Z and avoiding the generic point z. Thus each support has codimension>k and the induction applies to y. Adding y to the displayed cl(O_Z)−y∈Fil^k gives cl(O_Z)∈Fil^k, exactly the desired conclusion. The printed x assertion fails in an instance with no coefficient ambiguity: take X=P^1 over a field and Z one rational point, and choose U=X because Z is already a regular Cartier divisor. Then Z′ is empty, u=Id, x=cl(O_Z), and y=0. Euler characteristic sends x to1, so x is nonzero, whereas every combination of sheaf classes on the empty Z′ is0. For the second printed assertion put h=1−[O(−1)]=cl(O_Z). The projective-bundle presentation gives K(P^1)=Z[h]/(h^2). Its total gamma operation is γ_t(h)=1/(1−ht)=1+ht, and for any rank-zero element nh it is (1−ht)^(−n)=1+nht. Hence every gamma generator of degree≥2 and every product of two positive-degree generators is0, so Fil^2_gamma K(P^1)_Q=0; x=h remains nonzero after rationalization because χ_Q(h)=1. This also directly refutes the final printed x∈Fil^2. The two y substitutions restore the exact localization argument, with no change to its hypotheses or surrounding x used for the lift.


[473] EN-CONTRIB-VIII-001 Source folio 469. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=483
Printed reading: one deduces that $q \circ i$ is regular
Corrected reading: one deduces that $q \circ j$ is regular
Source: check/findings/findings_466_470.json#14. Main483 and witness480 both print i. Here j:X→W′, q:W′→W[T_1,. . . ,T_n], p:W′→W, p_0:W[T_1,. . . ,T_n]→W, and i=p∘j with p=p_0∘q. The preceding sentence establishes that q∘j is an immersion. Applying caseii to the étale q and regular j makes q∘j regular; caseiii applied to p_0 and q∘j then makes i=p_0∘q∘j regular. This supplies exactly the intermediate morphism used by the proof and fixes only its final argument letter.

[474] EN-CONTRIB-VIII-002 Source folio 470. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=484
Printed reading: the local ring of U_y at y is the quotient
Corrected reading: the local ring of U_y at x is the quotient
Source: check/findings/findings_466_470.json#21. Main484 and witness481 both print y. The preceding regular immersion i_y:U_y→V_y is regular at x, and its local homomorphism is the quotient O_{V_y,x}→O_{U_y,x}=O_{V_y,x}/I_x. This is the regular quotient invoked here;471 explicitly concludes that f is a complete intersection at x. The current x correction is already present.


[475] EN-CONTRIB-VIII-003 Source folio 472. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=486
Printed reading: into a scheme smooth over W
Corrected reading: into a scheme smooth over Z
Source: check/findings/findings_471_475.json#8. Main486 and witness483 print W. The actual upper-right scheme is Z[T′_1,. . . ,T′_m,T_1,. . . ,T_n], whose structural smooth projection π is to Z. The two displayed squares give π∘j′∘i=g∘f|_U: first the base-change square identifies projection to Z[T′] after j′ with j after projection to V, and the preceding factorization identifies those remaining composites with f and g. Thus the constructed regular immersion j′∘i factors g∘f through a smooth Z-scheme, which is exactly the claimed conclusion1.5. This fixes the base letter without changing the morphism or proof.

[476] EN-CONTRIB-VIII-004 Source folio 472. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=486
Printed reading: be a morphism
Corrected reading: be a morphism and
Source: check/findings/findings_471_475.json#9. Both scans omit a conjunction or separator between the two declarations. The inherited current et joins the declaration of f to that of g; the following base-change construction uses exactly the same two maps. It adds no mathematical hypothesis. Both attached layer readings are appropriate.

[477] EN-CONTRIB-VIII-005 Source folio 474. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=488
Printed reading:



Corrected reading:



Source: check/findings/findings_471_475.json#16. Both main488 and witness485 show the arrow pointing right from V to V′. The immediately preceding diagram binds p:V′→V and i=p∘j. Its differential sequence is 0→p*Ω¹_{V/Y}→Ω¹_{V′/Y}→Ω¹_{V′/V}→0, locally split because the last term is locally free for the smooth p; at a point of V′ its rank equality is exactly1.8.1. The corrected triangle V′→V→Y supplies this same p and sequence. The existing current interchange of V and V′ is retained; the printed triangle is retained diplomatically.


[478] EN-CONTRIB-VIII-006 Source folio 475. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=489
Printed reading: the homomorphism (1.5.1).
Corrected reading: the homomorphism (2.1.1).
Source: check/findings/findings_471_475.json#23. Both main489 and witness486 print1.5.1. The displayed map immediately above is tagged2.1.1 and is d:J/J²→i*Ω¹_{V/Y}. The definitions L_1=J/J² and L_0=i*Ω¹_{V/Y} identify its domain and codomain exactly with the stated differential L_1→L_0. Hence2.1.1 is the specific required reference, already restored currently.

[479] EN-CONTRIB-VIII-007 Source folio 476. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=490
Printed reading: $\underline{O}_V$
Corrected reading: $\underline{O}_{V'}$
Source: check/findings/findings_476_480.json#6. Main490 and witness487 print the index V without a prime. Here p:V′→V and j:X→V′. The three terms are p*Ω¹_{V/Y}, Ω¹_{V′/Y}, Ω¹_{V′/V}; the pullback used is j*M=O_X⊗_{j^{-1}O_{V′}}j^{-1}M. Their local freeness over O_{V′} makes the sequence locally split and preserves its exactness under this tensor product, giving2.2.2. The existing current O_{V′} supplies the ring of this exact tensor map. There is no implicit replacement by p_*: for V=Y=Spec Z and V′=Spec Z[1/2][t], the sheaf Ω¹_{V′/Y}=O_{V′}dt is free over O_{V′}, while p_*Ω at the stalk(2) is Q[t]dt, which is not free over Z_(2), since multiplication by2 is surjective there and is not surjective on any nonzero free Z_(2)-module. The compatible X=Spec F_3 at t=0 gives the closed immersions required in the proof.


[480] EN-CONTRIB-VIII-008 Source folio 477. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=491
Printed reading: the immersions i', j, j' are regular
Corrected reading: the immersion j' is regular
Source: ASTRA-VIII-0034. Main491 and native witness488 both print all three names. The governing assumption in2.1 and2.2 is only that f is lissifiable; complete-intersection hypotheses first appear in2.4. Take Y=Spec k, V=V′=Spec B with B=k[x,y], p=id, I=m^2 for m=(x,y), C=B/I and X=Spec C. Then i′=j=i is the closed immersion defined by m^2. At the unique point, $I/I^2=m^2/m^4$ has k-dimension3+4=7 (degree2 and degree3 monomials), while C has length3. Thus I/I^2 cannot be a finite free C-module, whose length would be a multiple of3; a regular immersion would have a finite free conormal module, so these two immersions are not regular. The section j′ is id_X and is regular of codimension0. In general the actual j′ is a section of the smooth p′ and is regular, precisely the source fact used to identify its conormal with relative differentials on478. The short conormal sequence remains justified without regularity of i′ or j: in the affine notation B→B′, C=B/I and J=ker(B′→C), smoothness supplies a B-algebra lift ℓ:B′→B/I^2 of B′→C through the square-zero quotient B/I^2→C. It sends J into I/I^2 and J^2 to0, inducing r:J/J^2→I/I^2. For a∈I, r(p(a) mod J^2)=a mod I^2, so r is a retraction of the natural map α:I/I^2→J/J^2. The usual quotient J/J^2→K/K^2, with K=J/IB′, has kernel the image of α; the retraction proves its injectivity, giving exactly2.2.3. Locally free relative differentials of the smooth p′ and the section identify K/K^2 with j*Ω^1_{V′/V}; the explicit inverse is described by the augmentation calculation on478. This repairs only the false two-immersion assertion and retains the author's displayed sequences and proof.

[481] EN-CONTRIB-VIII-009 Source folio 480. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=494
Printed reading: $\underline{\underline{P}}^r_Y$
Corrected reading: $\mathbf{P}^r_Y$
Source: ASTRA-VIII-0050. The directly inspected P letters each have two separate lower strokes. Diplomatic keeps these strokes literally; current uses the declared bold-P representation of the same projective-space construction Proj(O_Y[T_0,...,T_r]). The identity graded-algebra map induces the identity on every degree-zero localization chart D_+(T_i), and these chart identities glue to the identity of the projective Y-scheme. The plain abbreviation P elsewhere on482 is untouched.

[482] EN-CONTRIB-VIII-010 Source folio 480. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=494
Printed reading:




Corrected reading:




Source: check/findings/findings_476_480.json#19. Main480 visibly gives the derived-functor L two separate lower strokes, independently of the single rules on each complex L. Diplomatic now reproduces the literal two strokes; current retains mathbb L for the same derived pullback. For the displayed strictly perfect representative L^{Y/Z}, derived pullback is represented term by term by f*L^{Y/Z}; the identity on that representative is the exact comparison underlying this rendering choice.

[483] EN-CONTRIB-VIII-011 Source folio 482. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=496
Printed reading: $P = \underline{\underline{P}}^r_Y$
Corrected reading: $P = \mathbf{P}^r_Y$
Source: ASTRA-VIII-0051. The directly inspected P letters each have two separate lower strokes. Diplomatic keeps these strokes literally; current uses the declared bold-P representation of the same projective-space construction Proj(O_Y[T_0,...,T_r]). The identity graded-algebra map induces the identity on every degree-zero localization chart D_+(T_i), and these chart identities glue to the identity of the projective Y-scheme. The plain abbreviation P elsewhere on482 is untouched.

[484] EN-CONTRIB-VIII-012 Source folio 482. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=496
Printed reading:



Corrected reading:



Source: ASTRA-VIII-0051. The directly inspected P letters each have two separate lower strokes. Diplomatic keeps these strokes literally; current uses the declared bold-P representation of the same projective-space construction Proj(O_Y[T$_0$,...,T_r]). The identity graded-algebra map induces the identity on every degree-zero localization chart D_+(T_i), and these chart identities glue to the identity of the projective Y-scheme. The plain abbreviation P elsewhere on482 is untouched.

[485] EN-CONTRIB-VIII-013 Source folio 485. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=499
Printed reading: $\mathrm{Gr}^{k}\, K^{\bullet}(X)$
Corrected reading: $\mathrm{Gr}^{1}\, K^{\bullet}(X)$
Source: check/findings/findings_481_485.json#18. Main499 and witness496 both print k. For the invertible sheaf under consideration, x has rank1, so x−1 lies in Fil^1. The first Chern class is its image under the quotient Fil^1→Fil^1/Fil^2=Gr^1. The displayed identity gives $x^{-1}-1-(1-x)=(x-1)(1-x^{-1})$ in Fil^2, so applying precisely this quotient yields $c^1(x^{-1})=-c^1(x)$. The quotient degree is1 throughout this proof, which determines the corrected exponent independently of the unbound printed k.

[486] EN-CONTRIB-VIII-014 Source folio 487. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=501
Printed reading: $\underline{P}^r_Y \longrightarrow Y$
Corrected reading: $\mathbf{P}^r_Y \longrightarrow Y$
Source: ASTRA-VIII-0064. The main image shows both lower strokes at every counted occurrence; the487 native witness detail confirms all seven, including nested subscripts. The projective space is Proj(O_Y[T$_0$,...,T_r]); the identity graded-algebra map gives identity maps on degree-zero localizations at T_i, which glue to the identity projective scheme. The current bold P and diplomatic two-rule P therefore encode that exact same construction. No ordinary abbreviation is changed.

[487] EN-CONTRIB-VIII-015 Source folio 487. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=501
Printed reading: $\underline{\underline{P}}^{2}_{\underline{\underline{P}}^{r}_{\underline{\underline{Y}}}}$
Corrected reading: $\mathbf{P}^{2}_{\mathbf{P}^{r}_{Y}}$
Source: ASTRA-VIII-0064. The main image shows both lower strokes at every counted occurrence; the487 native witness detail confirms all seven, including nested subscripts. The projective space is Proj(O_Y[T$_0$,...,T_r]); the identity graded-algebra map gives identity maps on degree-zero localizations at T_i, which glue to the identity projective scheme. The current bold P and diplomatic two-rule P therefore encode that exact same construction. No ordinary abbreviation is changed.

[488] EN-CONTRIB-VIII-016 Source folio 487. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=501
Printed reading: $\underline{\underline{P}}^{2}_{\underline{\underline{P}}^{r}_{\underline{\underline{Y}}}} \longrightarrow \underline{\underline{P}}^{r}_{\underline{\underline{Y}}}$
Corrected reading: $\mathbf{P}^{2}_{\mathbf{P}^{r}_{Y}} \longrightarrow \mathbf{P}^{r}_{Y}$
Source: ASTRA-VIII-0064. The main image shows both lower strokes at every counted occurrence; the487 native witness detail confirms all seven, including nested subscripts. The projective space is Proj(O_Y[T$_0$,...,T_r]); the identity graded-algebra map gives identity maps on degree-zero localizations at T_i, which glue to the identity projective scheme. The current bold P and diplomatic two-rule P therefore encode that exact same construction. No ordinary abbreviation is changed.

[489] EN-CONTRIB-VIII-017 Source folio 487. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=501
Printed reading: $\underline{\underline{P}}^{r}_{\underline{\underline{Y}}} \longrightarrow Y$
Corrected reading: $\mathbf{P}^{r}_{Y} \longrightarrow Y$
Source: ASTRA-VIII-0064. The main image shows both lower strokes at every counted occurrence; the487 native witness detail confirms all seven, including nested subscripts. The projective space is Proj(O_Y[T$_0$,...,T_r]); the identity graded-algebra map gives identity maps on degree-zero localizations at T_i, which glue to the identity projective scheme. The current bold P and diplomatic two-rule P therefore encode that exact same construction. No ordinary abbreviation is changed.

[490] EN-CONTRIB-VIII-018 Source folio 488. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=502
Printed reading: $\underline{\underline{P}}^{r}_{\underline{\underline{Y}}} \to Y$
Corrected reading: $\mathbf{P}^{r}_{Y} \to Y$
Source: ASTRA-VIII-0065. The main image shows both lower strokes at every counted occurrence; the487 native witness detail confirms all seven, including nested subscripts. The projective space is Proj(O_Y[T$_0$,...,T_r]); the identity graded-algebra map gives identity maps on degree-zero localizations at T_i, which glue to the identity projective scheme. The current bold P and diplomatic two-rule P therefore encode that exact same construction. No ordinary abbreviation is changed.

[491] EN-CONTRIB-VIII-019 Source folio 489. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=503
Printed reading: $y \in K^{\bullet}(Y)_{\mathbb{Q}}$
Corrected reading: $y \in \mathrm{Gr}\, K^{\bullet}(Y)_{\mathbb{Q}}$
Source: ASTRA-VIII-0069. Main503 and native witness500, review/expose_viii/images/w489_domain_native.png, both omit Gr. The cited VII4.8 on folio460 explicitly defines u^gr:Gr^{k−d}K(Y)_Q→Gr^{k−1}K(Y′)_Q×Gr^kK(X)_Q by u^gr(y)=(g^gr(y)(−1)^{d−1}c^{d−1}(F),−i_gr(y)), and v^gr(a,b)=j_gr(a)+f^gr(b). Its zero composite gives f^gr i_gr(y)=j_gr(g^gr(y)(−1)^{d−1}c^{d−1}(F)). Proposition3.5 identifies the Chern factor with c^{d−1}(F∨), proving precisely the displayed4.3 for every degreewise y and hence for their finite sums in Gr K(Y)_Q. The ungraded group is explicitly related by θ_Z=ch∘$\tilde{c}$:K(Z)_Q→Gr K(Z)_Q from3.3–3.4, satisfying g^grθ_Y=θ_Y′g*; applying the graded identity to θ_Y(a) supplies a corresponding identity for any ungraded a. That transport does not change the directly cited graded operators or insert θ into the printed formula. The minimal Gr insertion restores exactly the domain written in VII4.8 and used on490 for y=ch(x)Todd(T_i∨).

[492] EN-CONTRIB-VIII-020 Source folio 490. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=504
Printed reading: By 3.2, the formula to be proved:
Corrected reading: By 4.2, the formula to be proved:
Source: ASTRA-VIII-0070. Main504 and native witness501 both read3.2. Proposition4.2 immediately before gives f_gr∘f^gr=Id on Gr K(X)_Q. For A=ch(i*x) and B=i_gr(ch(x)Todd(T_i∨)), the equality A=B gives f^gr(A)=f^gr(B); applying the displayed left inverse f_gr to the latter recovers A=B exactly. Thus4.2 supplies the equivalence used in this sentence. Proposition3.2 provides the filtration compatibility from which f_gr is defined, and4.2 uses that compatibility together with VII3.6 to prove the split identity. Replacing only the locator cites that established split identity without changing the proof or adding a hypothesis.


[493] EN-CONTRIB-VIII-021 Source folio 490. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=504
Printed reading: $\mathrm{ch}(g^*(y))$
Corrected reading: $\mathrm{ch}(g^*(x))$
Source: check/findings/findings_486_490.json#14. Main504 and native witness501 both read y on the right. The preceding right member is j_gr(g^gr(ch(x)Todd(T_i∨))c^{d−1}(F∨)). Naturality gives g^gr(ch(x))=ch(g*(x)) and g^gr(Todd(T_i∨))=Todd(g*(T_i∨)); multiplication by the same Chern factor gives exactly the displayed right member with x. The left member follows from VII3.4, f*i*(x)=j*(g*(x)λ_{−1}(F)), with the same input x. These two explicit maps determine x independently; the continuation on491 uses that identical x.

[494] EN-CONTRIB-VIII-022 Source folio 492. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=506
Printed reading: $-c(J)$
Corrected reading: $-c^1(J)$
Source: check/findings/findings_491_495.json#8. Main506 and native witness503, review/expose_viii/images/w492_chern_native.png, both omit1. For this line-bundle class J, the quotient q:Fil^{1}K(X)→Fil^1/Fil^2 gives c^1(J)=q(J−1), and therefore i_gr(1)=q(1−J)=−c^1(J). The completed Chern map instead writes c(J)=1+c^1(J), as the same page states in $\tilde{c}$(J)=(1,1+c^1(J)); the degree-one projection of that total class is precisely c^1(J). The displayed subsequent multiplication by Todd(−J∨)=(1−exp(c^1(J)))/(−c^1(J)) uses that degree-one element. For i a rational point in P^1, J=O(−1), with h=c^1(O(1)), the two readings are h and −1+h in the graded ring; the augmentation of the latter is −1, while i_gr(1) lies in degree1. The current exponent1 restores the quotient actually specified by the sentence.


[495] EN-CONTRIB-VIII-023 Source folio 492. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=506
Printed reading: $f : \underline{P}^r_Y \longrightarrow Y$
Corrected reading: $f : \mathbf{P}^r_Y \longrightarrow Y$
Source: ASTRA-VIII-0082. Both source symbols have two lower strokes. The ordinary abbreviation P remains plain. As for the preceding observed P occurrences, the identity graded map on O_Y[T$_0$,...,T_r] induces the identity on every projective chart, so the declared two renderings denote the same projective Y-scheme.

[496] EN-CONTRIB-VIII-024 Source folio 492. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=506
Printed reading: $P = \underline{P}^r_Y$
Corrected reading: $P = \mathbf{P}^r_Y$
Source: ASTRA-VIII-0082. Both source symbols have two lower strokes. The ordinary abbreviation P remains plain. As for the preceding observed P occurrences, the identity graded map on O_Y[T$_0$,...,T_r] induces the identity on every projective chart, so the declared two renderings denote the same projective Y-scheme.

[497] EN-CONTRIB-VIII-025 Source folio 493. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=507
Printed reading:



Corrected reading:



Source: check/findings/findings_491_495.json#15; ASTRA-VIII-0086. check/findings/findings_491_495.json#15: Main507 and native witness504, review/expose_viii/images/w493_triangle_native.png, both print X. Here E=O_Y^{r+1} and A=S_{O_Y}(E). The two lower arrows are exactly the unit maps O_Y→A and O_Y→A[T,T^-1], a↦a·1 and a↦a·1·T^0; the upper inclusion A→A[T,T^-1] has their composite equality. These are the structural maps of the Y-algebras used to form V and Z. The correction O_Y supplies precisely those two maps without changing the graded algebras, their degrees or the triangle. ASTRA-VIII-0086: The main image and native witness show the two S strokes independently of the single E rule. Both renderings denote the symmetric algebra ⊕_n E^{⊗n}/(permutation relations). The identity on E induces the identity on every tensor power, descends to each symmetric quotient, and yields mutually inverse identity graded-algebra maps. This occurrence-scoped VIII choice changes no generator, degree, algebra map or subscript; it does not propagate to earlier exposés.

[498] EN-CONTRIB-VIII-026 Source folio 493. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=507
Printed reading: $\underline{\underline{S}}(\underline{E})[T,T^{-1}]$
Corrected reading: $\mathbf{S}(\underline{E})[T,T^{-1}]$
Source: ASTRA-VIII-0086. The main image and native witness show the two S strokes independently of the single E rule. Both renderings denote the symmetric algebra ⊕_n E^{⊗n}/(permutation relations). The identity on E induces the identity on every tensor power, descends to each symmetric quotient, and yields mutually inverse identity graded-algebra maps. This occurrence-scoped VIII choice changes no generator, degree, algebra map or subscript; it does not propagate to earlier exposés.

[499] EN-CONTRIB-VIII-027 Source folio 494. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=508
Printed reading: $\underline{\underline{S}}(\underline{E})$-derivation
Corrected reading: $\mathbf{S}(\underline{E})$-derivation
Source: ASTRA-VIII-0087. The main image and native witness show the two S strokes independently of the single E rule. Both renderings denote the symmetric algebra ⊕_n E^{⊗n}/(permutation relations). The identity on E induces the identity on every tensor power, descends to each symmetric quotient, and yields mutually inverse identity graded-algebra maps. This occurrence-scoped VIII choice changes no generator, degree, algebra map or subscript; it does not propagate to earlier exposés.

[500] EN-CONTRIB-VIII-028 Source folio 494. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=508
Printed reading:

$$\underline{\underline{S}}(\underline{E})[T,T^{-1}] \longrightarrow \underline{\underline{S}}(\underline{E})[T,T^{-1}] \otimes_{\underline{\underline{S}}(\underline{E})} \underline{\underline{S}}(\underline{E}) \cdot dT$$

Corrected reading:

$$\mathbf{S}(\underline{E})[T,T^{-1}] \longrightarrow \mathbf{S}(\underline{E})[T,T^{-1}] \otimes_{\mathbf{S}(\underline{E})} \mathbf{S}(\underline{E}) \cdot dT$$

Source: ASTRA-VIII-0087. The main image and native witness show the two S strokes independently of the single E rule. Both renderings denote the symmetric algebra ⊕_n E^{⊗n}/(permutation relations). The identity on E induces the identity on every tensor power, descends to each symmetric quotient, and yields mutually inverse identity graded-algebra maps. This occurrence-scoped VIII choice changes no generator, degree, algebra map or subscript; it does not propagate to earlier exposés.

[501] EN-CONTRIB-VIII-029 Source folio 494. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=508
Printed reading: $\underline{\underline{S}}(\underline{E}) \cdot dT$
Corrected reading: $\mathbf{S}(\underline{E}) \cdot dT$
Source: ASTRA-VIII-0087. The main image and native witness show the two S strokes independently of the single E rule. Both renderings denote the symmetric algebra ⊕_n E^{⊗n}/(permutation relations). The identity on E induces the identity on every tensor power, descends to each symmetric quotient, and yields mutually inverse identity graded-algebra maps. This occurrence-scoped VIII choice changes no generator, degree, algebra map or subscript; it does not propagate to earlier exposés.

[502] EN-CONTRIB-VIII-030 Source folio 494. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=508
Printed reading: defined by $\underline{\underline{S}}(\underline{E})$
Corrected reading: defined by $\mathbf{S}(\underline{E})$
Source: ASTRA-VIII-0087. The main image and native witness show the two S strokes independently of the single E rule. Both renderings denote the symmetric algebra ⊕_n E^{⊗n}/(permutation relations). The identity on E induces the identity on every tensor power, descends to each symmetric quotient, and yields mutually inverse identity graded-algebra maps. This occurrence-scoped VIII choice changes no generator, degree, algebra map or subscript; it does not propagate to earlier exposés.

[503] EN-CONTRIB-VIII-031 Source folio 494. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=508
Printed reading: $\underline{\underline{S}}(\underline{E}) \otimes_{\underline{O_Y}} \underline{E}$
Corrected reading: $\mathbf{S}(\underline{E}) \otimes_{\underline{O_Y}} \underline{E}$
Source: ASTRA-VIII-0087. The main image and native witness show the two S strokes independently of the single E rule. Both renderings denote the symmetric algebra ⊕_n E^{⊗n}/(permutation relations). The identity on E induces the identity on every tensor power, descends to each symmetric quotient, and yields mutually inverse identity graded-algebra maps. This occurrence-scoped VIII choice changes no generator, degree, algebra map or subscript; it does not propagate to earlier exposés.

[504] EN-CONTRIB-VIII-032 Source folio 494. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=508
Printed reading: $g^*(E) \cdot \xi^{-1} - 1$
Corrected reading: $f^*(E) \cdot \xi^{-1} - 1$
Source: check/findings/findings_491_495.json#20. Main508 and native witness505, review/expose_viii/images/w494_Tf_native.png, both print g. The displayed Euler sequence5.2 on P is 0→Ω¹_{P/Y}→f*E⊗O_P(−1)→O_P→0; applying the class map gives T_f=f*[E]·ξ^{-1}−1 in K(P), exactly with f. The earlier triangle has h:Z→P and g=f∘h, so h*f*[E]=g*[E]; tensor associativity identifies O_Z⊗_{h^{-1}O_P}(h^{-1}O_P⊗_{g^{-1}O_Y}g^{-1}E) with O_Z⊗_{g^{-1}O_Y}g^{-1}E. Pulling back the Euler equality therefore gives the corresponding identity on Z. Moreover E=O_Y^{r+1} supplies canonical trivializations of both pullbacks: under the integer unit maps Z→K(P), Z→K(Z), both are the image of r+1. The following $\tilde{c}$(E) is consequently transported by f^gr as the same pair(r+1,1), and is retained. The minimal current f correction names the actual pullback in the cited sequence on P; it adds no new term or explanatory material.

[505] EN-CONTRIB-VIII-033 Source folio 496. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=510
Printed reading: and that $1, \dots, x^r$ are linearly independent over $\operatorname{Gr} K^\bullet(Y)$
Corrected reading: and that $1, \dots, x^r$ are linearly independent over $\operatorname{Gr} K^\bullet(Y)$.
Source: check/findings/findings_496_500.json#3. The main image shows no stop after the closing parenthesis. The completed sentence is followed by a separate indented paragraph. The attached diplomatic absence and current terminal stop already preserve and correct this source omission respectively.

[506] EN-CONTRIB-VIII-034 Source folio 497. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=511
Printed reading:

$$\frac{(1-y)^{-k-1} \cdot dy}{y^{r+1}}.$$

Corrected reading:

$$\frac{(1-y)^{-k-1} \cdot dy}{y^{r+1}},$$

Source: check/findings/findings_496_500.json#7. Main511 and witness508 both show a compact full stop without a comma tail. The next lower-case clause restates the same residue as a coefficient, continuing the sentence introduced by résidu de. The attached diplomatic full stop/current comma already preserves this source punctuation and restores that continuation.

[507] EN-CONTRIB-IX-001 Source folio 503. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=517; scan/SGA6_witness_300dpi.djvu#page=514
Printed reading: runs through the set of elements of
Corrected reading: runs through the set of nonzero elements of
Source: ASTRA-IX-NONZERO-LOCALIZATION-503. Both complete503 images explicitly say f parcourt l'ensemble des éléments de A. On502 the source made A integral and K its fraction field. Including f=0 gives A_0=0; every A_f maps to this terminal localization, so the all-element colimit is0. For A=Z this contradicts the named Q, and no unital map0→Q can be a fraction-field cocone because1_Q≠0_Q. Over nonzero f, the map colim A_f→Frac(A) sends a/f^n to the same fraction. Its inverse sends a/b to its image from A_b; equality a/b=c/d becomes ad=bc in the common localization A_(bd), proving well-definedness and both inverse identities. The generic point(0) belongs to D(f) exactly for f≠0, matching the following nonempty U=Spec(A_f). Current inserts only non nuls; diplomatic retains the printed domain. The independent context audit and exact maps are in localization_index_503.json.

[508] EN-CONTRIB-IX-002 Source folio 506. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=520; scan/SGA6_witness_300dpi.djvu#page=517
Printed reading: $g^*(g_*(f))$
Corrected reading: $g^*(g_*(F))$
Source: check/findings/findings_506_510.json#4. Both complete506 images print the lower-case f. The preceding sentence names the coherent class F and the507 calculation begins g^*(g_*(F)). The exact composite on that class is [F]↦[g_*F]↦sum_i(-1)^i[Tor_i^(O_X)(g_*F,O_S)]=[F]·lambda_-1(E). The structural morphism f:X→S has no element in this composite's K-group domain. Keep the source lower-case f diplomatically and the existing current F correction.

[509] EN-CONTRIB-IX-003 Source folio 510. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=524; scan/SGA6_witness_300dpi.djvu#page=521
Printed reading: By 1.3, this implies that the homomorphism:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(S) \longrightarrow K_\bullet(X)$$

is surjective.
Corrected reading: By 1.3, this implies that the homomorphism:

$$K_\bullet(S) \otimes_{K^\bullet(S)} K^\bullet(X) \longrightarrow K_\bullet(X)$$

is surjective.
Source: check/findings/findings_506_510.json#32. Both complete510 images print S in the second factor of the surjectivity display. The statement on this same page and Corollary1.3 identify the map with alpha:a tensor b↦f^*a·b for b in K^bullet(X), balanced over K^bullet(S). Restore that X only in current. For S=Spec(k), E=k^2 and X=P^1, the printed domain is Z tensor_Z Z=Z and its natural map has image Z[O]; the known projective-bundle basis1,delta with delta=[O(1)]−1 shows the missing generator. The corrected map is Z tensor_Z K^bullet(P^1)→K_bullet(P^1),1 tensor b↦b. Diplomatic retains S.

[510] EN-CONTRIB-IX-004 Source folio 512. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=526; scan/SGA6_witness_300dpi.djvu#page=523
Printed reading: $\sum_{i=0}^{n-1} \xi^{-1} f^*(x_i)$
Corrected reading: $\sum_{i=0}^{n-1} \xi^{-i} f^*(x_i)$
Source: check/findings/findings_511_515.json#12. Both complete512 images show the digit1 on the left and dotted i on the right. The formula511 defines phi((x_i))=sum_i xi^(−i) f^*(x_i). Applying f_* termwise and the projection formula gives exactly sum_i x_i f_*(xi^(−i)); current therefore keeps its corrected exponent−i, while diplomatic keeps−1. For X=P^1_k,n=2,x_0=1,x_1=0, the printed left side is chi(O(−1))=0 but the displayed right side is1; the corrected left side is chi(O)=1.

[511] EN-CONTRIB-IX-005 Source folio 515. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=529; scan/SGA6_witness_300dpi.djvu#page=526
Printed reading: $\underline{G}_{0S''}^{(i+1)}$
Corrected reading: $\underline{G}_{0S''}^{(i)}$
Source: ASTRA-IX-QUOTIENT-INDEX-515. Both complete515 images and the native index detail show i+1 in the congruence. The same page defines F_0^(i)=G_0^(i−1)/G_0^(i), so the quotient q_i:G_0^(i−1)→F_0^(i) has kernel G_0^(i). Thus q_i(a)=q_i(b) exactly when a−b lies in G_0^(i). The comparison from G_0^(i−1)/G_0^(i+1) onto F_0^(i) has the extra kernel G_0^(i)/G_0^(i+1). In k^2 take G_0=k^2,G_1=ke_1,G_2=0,i=1: e_2 and e_2+e_1 give the same basis class in F_1, and the upper unipotent matrix[[1,1],[0,1]] relates them, but their difference e_1 is not in G_2. Replace only i+1 by i in current, retaining the printed index diplomatically. The independent full-page/native audit agrees and is preserved in quotient_index_515.json.

[512] EN-CONTRIB-X-001 Source folio 520. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=534
Printed reading: $G \longrightarrow \underline{G}' \longrightarrow 0$
Corrected reading: $G \longrightarrow \underline{G}'' \longrightarrow 0$
Source: check/findings/findings_516_520.json#23. Both scans show one prime in the second and fourth terms. The prose immediately names G′, G and G″, and the next page gives length(G)=length(G′)+length(G″) and cl(G)=cl(G′)+cl(G″). The quotient in the exact sequence is therefore G″ in current; the attached current correction is exact.

[513] EN-CONTRIB-X-002 Source folio 524. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=538
Printed reading: denote by ε_Z the class
Corrected reading: denote by 1_Z the class
Source: check/findings/findings_521_525.json#15. Main and witness535 show epsilon followed by Z; the following projection formulas each use1_Z. The named class is [O_Z] in K_•(Z), and α_*[O_Z]=[O_Z] as a coherent sheaf on X. The existing current1_Z therefore refers to exactly the same class and restores the notation used in every displayed application.

[514] EN-CONTRIB-X-003 Source folio 524. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=538
Printed reading: (IV . . .)
Corrected reading: (IV 2.11.1.2)
Source: ASTRA-X-0015. Both main and witness535 print three dots. The directly inspected formula IV2.11.1.2 on printed287 (main PDF301; review/expose_x/images/target287_projection.png, upper region only) is f_*(yx)=y f_*(x). Substituting f=α:Z→X, y=x∈K^• (X), and the coherent class [O_Z] gives α_*(α^*(x)[O_Z])=xα_*[O_Z], precisely this display. The closed immersion α is proper between noetherian schemes, satisfying IV2.11.1's hypotheses. Supply this established reference in current only.

[515] EN-CONTRIB-X-004 Source folio 527. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=541
Printed reading: $D(x) = D_Y(x) = \dim(\{x\})$_
Corrected reading: $D(x) = D_Y(x) = \dim(\overline{\{x\}})$
Source: check/findings/findings_526_530.json#6. Main and witness538 show no bar. For X=A^1_k, the singleton generic point has topological dimension0, as does a closed point, so this literal reading violates the explicitly stated strict specialization inequality(ii). Their closures have dimensions1 and0, and the later formula1.5.1 explicitly uses the closure bar. Retain the attached current correction to dim(overline{{x}}).

[516] EN-CONTRIB-X-005 Source folio 530. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=544
Printed reading: inside Z
Corrected reading: inside X
Source: check/findings/findings_526_530.json#21. The printed letter is Z. The construction gives closed immersions j_i:Y_i→X identifying y_1 with z_2 and z_1 with y_2; their intersection inside X is exactly the two glued closed points, named {y_1,y_2}. Thus the displayed Y_1∩Y_2 is an intersection in X. Retain current X and the diplomatic Z.

[517] EN-CONTRIB-X-006 Source folio 531. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=545
Printed reading: $x_1 \ldots, x_r$
Corrected reading: $x_1, \ldots, x_r$
Source: check/findings/findings_531_535.json#2. The source omits the separator after x_1. The list enumerates the ordered entries x_1,. . . ,x_r, as repeated in the coefficient arguments immediately below. Retain the printed omission diplomatically and the existing current comma.

[518] EN-CONTRIB-X-007 Source folio 533. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=547
Printed reading: whence the assertion that f is constant on the cosets in G modulo G''.
Corrected reading: whence the assertion that f is constant on the cosets in G modulo G'.
Source: check/findings/findings_531_535.json#10. The enlarged main and witness544 crops show a second mark at prime height, broken into two ink pieces, followed separately by the sentence stop. The preceding g_b:G′→Q is constant; hence f(b+u)=f(b) for every u∈G′. The cosets on which f is constant are therefore b+G′. Correct current G″ to G′ here.

[519] EN-CONTRIB-X-008 Source folio 533. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=547
Printed reading: $h : G' \to \mathbb{Q}$
Corrected reading: $h : G'' \to \mathbb{Q}$
Source: check/findings/findings_531_535.json#11. Both scans print one prime. Since ker(j)=G′ and j:G→G″ is surjective, h is defined by h(j(b))=f(b); the preceding coset constancy makes this well defined. This gives the exact factorization f=h∘j with domain G″ for h. Correct current G′ to G″.

[520] EN-CONTRIB-X-009 Source folio 533. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=547
Printed reading: by hypothesis C'' is a torsion group
Corrected reading: by hypothesis G'' is a torsion group
Source: check/findings/findings_531_535.json#12. The printed capital is C with two primes. The exact sequence in the lemma defines G″ as the torsion quotient, and h has that quotient as domain. Retain the current G″ correction.

[521] EN-CONTRIB-X-010 Source folio 534. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=548
Printed reading: a sequence of elements of F with
Corrected reading: a sequence of elements of P with
Source: check/findings/findings_531_535.json#15. The main and witness545 show a top bar and a separate middle bar without the bowl of P. The proof explicitly writes x_i=1+y_i with y_i∈I, which defines membership in the multiplicative subgroup P=1+I introduced in the lemma. Correct current F to P.

[522] EN-CONTRIB-X-011 Source folio 536. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=550
Printed reading: $S_y : \mathrm{Pic}(X) \longrightarrow$
Corrected reading: $s_y : \mathrm{Pic}(X) \longrightarrow$
Source: check/findings/findings_536_540.json#5. The source capital is certain. Proposition2.4.4 explicitly identifies its s_y by the reference(2.3.1): for every line-bundle class [L], both names denote the same map [L]↦χ(cl^•(L)y) from Pic(X) to Q. Thus their identification is pointwise equality, with the identity on domain and codomain. Current uses s_y consistently; this is a notation correction and changes no value or hypothesis.

[523] EN-CONTRIB-X-012 Source folio 536. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=550
Printed reading: $\chi(\mathrm{cl}^\bullet(\underline{L})y$
Corrected reading: $\chi(\mathrm{cl}^\bullet(\underline{L})y)$
Source: check/findings/findings_536_540.json#4. The source stops at y. The attached current layer closes the argument of χ; the diplomatic layer preserves the missing parenthesis.

[524] EN-CONTRIB-X-013 Source folio 537. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=551
Printed reading: $\underset{\sim\sim}{\mathrm{Pic}}^0_{X/k}$
Corrected reading: $\underset{\sim\sim}{\mathrm{Pic}}_{X/k}$
Source: check/findings/findings_536_540.json#13. The superscript is present in both scans. The preceding536 paragraph defines the full fpqc sheafification of T↦Pic(X×_kT); the representing scheme here is denoted Pic_{X/k}, and only the next sentence introduces its neutral component Pic^0_{X/k}. The representability identification sends a T-point to the corresponding Picard class. The natural inclusion of the identity-component subfunctor into the full functor is an inclusion: on X=P^1 over an algebraically closed field its k-point map is 0→Z, while O(1) maps to degree1. Removing0 in current makes this representability statement refer to exactly the full functor defined on536 and leaves the following component definition intact.

[525] EN-CONTRIB-X-014 Source folio 539. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=553
Printed reading: $f_*((f^*(x)\cdot y) = x\cdot f_*(Y)$
Corrected reading: $f_*(f^*(x)\cdot y) = x\cdot f_*(y)$
Source: check/findings/findings_536_540.json#28; check/findings/findings_536_540.json#29. check/findings/findings_536_540.json#28: The duplicate opening is visible. The displayed operation is f_* applied to the single product f^*(x)·y, so the attached current removal of one opening parenthesis gives its balanced argument; diplomatic preserves the duplicate. check/findings/findings_536_540.json#29: The source capital is clear. The display quantifies y∈K^•(X′)_Q and its left-hand side applies f_* to f^*(x)·y. The projection formula uses that same y on the right; the attached current lower-case correction is exact.

[526] EN-CONTRIB-X-015 Source folio 540. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=554
Printed reading: $\dim(f^{-1}(\mathrm{Supp}(\underline{F}))$
Corrected reading: $\dim(f^{-1}(\mathrm{Supp}(\underline{F})))$
Source: check/findings/findings_536_540.json#33. The source omits the closing parenthesis for the outer dim expression. The attached current layer adds that final parenthesis around f^{-1}(Supp(F)); the diplomatic layer preserves the omission.

[527] EN-CONTRIB-X-016 Source folio 542. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=556
Printed reading: $\mathrm{cl}^\bullet(\gamma^i(x-\epsilon(x))$
Corrected reading: $\mathrm{cl}^\bullet(\gamma^i(x-\epsilon(x)))$
Source: check/findings/findings_541_545.json#6. The source has three opening and only two closing parentheses before ∈. The current layer closes cl^•(γ^i(x−ε(x))) exactly; diplomatic keeps the missing outer close.

[528] EN-CONTRIB-X-017 Source folio 542. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=556
Printed reading: $P(c_1(x),\ldots,c_n(X))\in \mathrm{Gr}^n(X)$
Corrected reading: $P(c_1(x),\ldots,c_n(x))\in \mathrm{Gr}^n(X)$
Source: check/findings/findings_541_545.json#8. The source capital is clear. The substitution map Z[T_1,...,T_n]→Gr^•(X) is T_i↦c_i(x), so its value is P(c_1(x),...,c_n(x)). The following displayed pairing also uses c_n(x). Retain the current lower-case argument.

[529] EN-CONTRIB-X-018 Source folio 542. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=556
Printed reading: $P_y(x) = \langle P(c_1(x)), \dots, c_n(x), y\rangle$.
Corrected reading: $P_y(x) = \langle P(c_1(x), \dots, c_n(x)), y\rangle$.
Source: check/findings/findings_541_545.json#10. The source visibly has two consecutive closing parentheses after x in the first term. The pairing4.1.1 has two inputs, Gr^n(X) and Gr_n(X); the substitution map sends P to P(c_1(x),...,c_n(x)), which supplies the first input, and y supplies the second. The current regrouping is therefore the defined composite χ(P(c_i(x))·y).

[530] EN-CONTRIB-X-019 Source folio 543. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=557
Printed reading:

$$\langle c_1(L_1)^{\alpha_1} \cdots c_d(L_d)^{\alpha_d}, |\underline{F}|\rangle, \tag{4.3.2}$$

Corrected reading:

$$\langle c_1(L_1)^{\alpha_1} \cdots c_1(L_d)^{\alpha_d}, |\underline{F}|\rangle, \tag{4.3.2}$$

Source: check/findings/findings_541_545.json#11. Both main and witness554 print the subscript d. The proof writes L_i=1+M_i, so the associated degree-one class of M_i is c_1(L_i); multiplication sends ∏M_i^{α_i} to ∏c_1(L_i)^{α_i} in Gr^d. This is exactly the coefficient being paired. Retain the current c_1(L_d) and the diplomatic c_d.

[531] EN-CONTRIB-X-020 Source folio 543. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=557
Printed reading: $\alpha_1! \cdots a_d!$
Corrected reading: $\alpha_1! \cdots \alpha_d!$
Source: check/findings/findings_541_545.json#12. Main and witness554 show the typed a, without the handwritten alpha shape. The leading term of binom(n_i,α_i) is n_i^{α_i}/α_i!, so the coefficient conversion multiplies by1/(α_1!···α_d!). The current α_d! is exact.

[532] EN-CONTRIB-X-021 Source folio 543. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=557
Printed reading: **Corollary 4.3.3.** The coefficient of the monomial $n_1 n_2 \cdots n_d$ in the Snapper polynomial $\chi(L_1^{\alpha_1} \cdots L_d^{\alpha_d} F)$ is equal to
Corrected reading: **Corollary 4.3.3.** The coefficient of the monomial $n_1 n_2 \cdots n_d$ in the Snapper polynomial $\chi(L_1^{n_1} \cdots L_d^{n_d} F)$ is equal to
Source: check/findings/findings_541_545.json#13. Both scans have alpha exponents. The coefficient being extracted is n_1···n_d, and the preceding proposition defines the polynomial as (n_1,...,n_d)↦χ(L_1^{n_1}···L_d^{n_d}F). The current n exponents restore this same polynomial, with no changed coefficient or degree.

[533] EN-CONTRIB-X-022 Source folio 543. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=557
Printed reading: $\alpha_1 + \cdots + \alpha_d = d$
Corrected reading: $\alpha_1 + \cdots + \alpha_d \le d$
Source: ASTRA-X-0056. Main and witness554 print =d under the summation. This is a congruence in K^•(X)/Filt^{d+1}(X), not a statement only in its degree-d associated quotient. For d=1, X=Spec(k), L_1=O_X and n_1=0, the printed sum gives1≡0 in K^•(X)/Filt^2(X)=Z. Under the quotient map K^•(X)→A, M_i=L_i−1 belongs to the ideal I and I^{d+1}=0; the finite binomial expansion of ∏(1+M_i)^{n_i} retains every term ∏binom(n_i,α_i)M_i^{α_i} with |α|≤d. Terms of larger total degree map to0. This is exactly the cited formula2.1.5 on534, including negative integer n_i via the inverse of1+M_i. Correct only the summation constraint; the proposition's separate top-degree hypothesis Σα_i=d remains unchanged.

[534] EN-CONTRIB-X-023 Source folio 544. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=558
Printed reading: $\mathrm{long}_{\underline{O}_{,x}}$
Corrected reading: $\mathrm{long}_{\underline{O}_{X,x}}$
Source: check/findings/findings_541_545.json#19. The source leaves the position before the comma empty. Its point x lies in X(d), and the stalk F_x is an O_{X,x}-module;1.1.2 and the full local-ring notation4.4.4 on545 specify that same ring. Retain the current X and the diplomatic omission.

[535] EN-CONTRIB-X-024 Source folio 545. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=559
Printed reading: $\mathrm{cl}_\bullet(f_*(\underline{O}_Y) \equiv$
Corrected reading: $\mathrm{cl}_\bullet(f_*(\underline{O}_Y)) \equiv$
Source: check/findings/findings_541_545.json#23. Main and witness556 show the missing outer close. The current layer adds it around the coherent pushforward before the next congruence sign; diplomatic preserves the omission.

[536] EN-CONTRIB-X-025 Source folio 546. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=560
Printed reading: From 2.3.4 one obtains
Corrected reading: From 2.4.4 one obtains
Source: ASTRA-X-0064. The scan prints2.3.4, but §2.3 has only2.3.1 and2.3.2. The directly inspected Proposition2.4.4 on538 proves that s_y is constant on Pic^τ-cosets. For [L]∈Pic^τ(X), every [L^n] is in the same subgroup; taking y=[O_Z] for an integral closed curve Z gives s_y([L^n])=s_y([O_X]), namely χ(L^n⊗O_Z)=χ(O_Z). This is exactly criterion4.5.3, proving the stated inclusion Pic^τ(X)→Pic^N(X). Thus2.4.4 is the established reference used here.

[537] EN-CONTRIB-X-026 Source folio 550. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=564
Printed reading: $+\underline{N} - 1 \in \mathrm{Filt}^2(X)$
Corrected reading: $+\underline{n} - 1 \in \mathrm{Filt}^2(X)$
Source: check/findings/findings_546_550.json#17. Both scans show capital N, with a short rule. The rank section is n, and the element under consideration is cl^•(E)−n. The preceding5.2.2(b) gives det(E)−1≡cl^•(E)−n modulo Filt^2, hence det(E)−cl^•(E)+n−1∈Filt^2. The attached current lower-case correction is exact; its rule is restored separately in both layer-specific forms.

[538] EN-CONTRIB-X-027 Source folio 550. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=564
Printed reading: Then (6.2.8) and (6.5.8) below give
Corrected reading: Then (6.2.8) and (6.5.3) below give
Source: check/findings/findings_546_550.json#18. The source550 clearly prints6.5.8. Direct inspection of554 identifies6.2.8 as det(E_1⊗E_2)=det(E_1)^{rank E_2}⊗det(E_2)^{rank E_1}. Direct inspection of559 shows6.5.3 explicitly extending6.2.8–6.2.10 to any ringed topos via pullback along u:Y↦Spec(O(Y)); on each Y the representing matrices and determinant polynomial are identical. In the rank-zero subgroup, the two rank factors in this tensor determinant formula vanish, giving det(xy)=1 for x,y of rank0, precisely5.3.5. The next printed page560 begins§7, so§6.5 ends at6.5.5 and has no6.5.8. Current6.5.3 is the exact established dependency.

[539] EN-CONTRIB-X-028 Source folio 551. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=565
Printed reading: $\binom{\underline{n}+k-1}{p}\lambda^q(E)$
Corrected reading: $\binom{-\underline{n}+k-1}{p}\lambda^q(E)$
Source: check/findings/findings_551_555.json#1. Main and witness562 both clearly omit the minus. The identity λ_t(E+a)=λ_t(E)(1+t)^a with a=k−1−n gives the coefficient Σ_{p+q=k} binom(k−1−n,p)λ^q(E). The later printed r already uses this numerator; its convolution is the coefficient of t^{k−1} in (1+t)^{k−1−n}(1+t)^{n−1}=(1+t)^{k−2}, which is zero for k>1. Thus the attached current minus is required by the exact generating-series multiplication.

[540] EN-CONTRIB-X-029 Source folio 552. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=566
Printed reading: $\mathrm{Mod}^{\underline{n}}\underline{O}(X)$
Corrected reading: $\mathrm{Mod}^{\underline{n}}_{\underline{O}}(X)$
Source: check/findings/findings_551_555.json#9. The source places O at the same baseline as Mod and (X), with n raised. Diplomatically this remains Mod^n O(X). The current subscript O expresses the base ring already used in Mod_O(X) on551: the identification is the identity on each locally free O-module of rank n and each O-linear map, so no object or morphism changes. Retain the attached current subscript correction and restore its separate printed O rule.

[541] EN-CONTRIB-X-030 Source folio 553. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=567
Printed reading: makes G act on $\underline{O^m}$ by
Corrected reading: makes G act on M by
Source: ASTRA-X-0080. The representation specifies, on each object U, the action map G(U)×M(U)→M(U), (g,v)↦α_U(g)(v). The group and restriction identities follow directly from α being a sheaf homomorphism. The associated-module construction immediately following is P×^G M, using exactly that action. Section6.1 defines M=T(O^{n_1},…,O^{n_r}) as an arbitrary O-module; local freeness was an additional premise only for the earlier composition sentence, and no global basis of M was chosen. A chosen trivialization u:O^m→M transports the displayed action to O^m by (g,v)↦u^{-1}α(g)u(v), making the missing identification explicit, but it is absent from this assertion. For example on P^1, T(E)=E⊗O(1) gives M=O(1), with scalar action; Γ(O(1)) has basis X_0,X_1 whereas Γ(O) consists of constants, so no O≅M trivialization exists. Replacing O^m by M states exactly the already specified action and the module used in6.1.4. This reverses the earlier rejected correction.

[542] EN-CONTRIB-X-031 Source folio 554. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=568
Printed reading: of commutators of $\mathrm{Gl}(\underline{n}))$ is represented by
Corrected reading: of commutators of $\mathrm{Gl}(\underline{n})$ is represented by
Source: check/findings/findings_551_555.json#14. The source has two closing parentheses following the single opening after Gl. Current already removes the unmatched second close; preserve the source diplomatically.

[543] EN-CONTRIB-X-032 Source folio 554. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=568
Printed reading:

$$\det(\underline{E}_i \otimes \underline{E}_2) = (\det(\underline{E}_1))^{\otimes \underline{n}_2} \otimes (\det(\underline{E}_2))^{\otimes \underline{n}_1}, \tag{6.2.8}$$
$$\det(\underline{E}_1 \otimes \underline{E}_2) = (\det(\underline{E}_1)) \otimes (\det(\underline{E}_2)) \tag{6.2.9}$$

Corrected reading:

$$\det(\underline{E}_1 \otimes \underline{E}_2) = (\det(\underline{E}_1))^{\otimes \underline{n}_2} \otimes (\det(\underline{E}_2))^{\otimes \underline{n}_1}, \tag{6.2.8}$$
$$\det(\underline{E}_1 \oplus \underline{E}_2) = (\det(\underline{E}_1)) \otimes (\det(\underline{E}_2)) \tag{6.2.9}$$

Source: check/findings/findings_551_555.json#17; check/findings/findings_551_555.json#18. check/findings/findings_551_555.json#17: Main and witness565 distinguish the dotted i from digit1 in the adjacent expressions. For free bases e_a of E_1 and f_b of E_2, the map from (∧^{n_1}E_1)^{⊗n_2}⊗(∧^{n_2}E_2)^{⊗n_1} to ∧^{n_1n_2}(E_1⊗E_2) sends their ordered basis generators to the lexicographically ordered wedge of e_a⊗f_b. A basis change A,B multiplies both sides by det(A)^{n_2}det(B)^{n_1}, so these local maps glue. This proves the displayed right side for E_1⊗E_2, exactly as in the attached current correction; the unbound first factor E_i has no corresponding right-side determinant.
check/findings/findings_551_555.json#18: Both scans print the same circled cross as6.2.8. The ordered wedge map det(E_1)⊗det(E_2)→det(E_1⊕E_2), (e_1∧…∧e_a)⊗(f_1∧…∧f_b)↦(e_1,0)∧…∧(e_a,0)∧(0,f_1)∧…∧(0,f_b), is invariant under each basis change and glues to an isomorphism. It gives the displayed right side for the direct sum. For E_1=O(1),E_2=O^2 on P^1, the tensor determinant is O(2), whereas the displayed right side is O(1); their global section dimensions3 and2 prove the printed tensor identity false. The attached current direct-sum correction is exact.

[544] EN-CONTRIB-X-033 Source folio 555. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=569
Printed reading: an elementary proof here via 6.2.2).
Corrected reading: an elementary proof here via 6.2.2.
Source: check/findings/findings_551_555.json#20. The entire paragraph is on555 and has no opening parenthesis before Notons que or elsewhere. The preceding parenthesized reference SGA3VIII1.3 closes before the displayed character; it cannot open this sentence. Delete only the final unmatched parenthesis in current and preserve it diplomatically. This establishes the correction previously rejected by Codex.

[545] EN-CONTRIB-X-034 Source folio 557. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=571
Printed reading: $c = c_{h,1,\lambda}$
Corrected reading: $c = c_{2,h,\lambda}$
Source: ASTRA-X-0093. Read under the explicit556 rule a c adds row k to row h, equivalently a⋆c=C A in ordinary coordinates. For c=c_{2,h,λ}, (s⋆c)_{21}=s_{21}+λs_{h1}. Thus the chosen i with s_{i1}∉p and s_{21}∈p is realized by c_{2,i,1}, giving s′_{21}=s_{21}+s_{i1}∉p. Necessarily i≠2, so this is one of the defined elementary matrices. The printed c_{i,1,1} need not affect row2: over Q, take s=((0,1,0),(0,0,1),(1,0,0)), det(s)=1, p=(0), i=3. Then s_{11}=s_{21}=0 and s_{31}=1; right multiplication in the printed convention by c_{3,1,1} leaves s′_{21}=0, whereas c_{2,3,1} makes s′_{21}=1. The later s⋆c_{1,2,λ} gives s′_{11}=s_{11}+λs_{21}, so those printed products and indices already prove the stated pivot and remain unchanged. The case n=1 is already the identity from det(s)=1.

[546] EN-CONTRIB-X-036 Source folio 557. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=571
Printed reading: $s_{i1} \notin \underline{p}$ Then
Corrected reading: $s_{i1} \notin \underline{p}$. Then
Source: check/findings/findings_556_560.json#16. Both images end the line at underlined p and begin the next at Alors with no terminal dot. The attached current stop supplies the boundary between the two sentences; the diplomatic omission is exact.

[547] EN-CONTRIB-X-035 Source folio 557. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=571
Printed reading: $c_{i,1,1}$
Corrected reading: $c_{2,i,1}$
Source: ASTRA-X-0093. Read under the explicit556 rule a c adds row k to row h, equivalently a⋆c=C A in ordinary coordinates. For c=c_{2,h,λ}, (s⋆c)_{21}=s_{21}+λs_{h1}. Thus the chosen i with s_{i1}∉p and s_{21}∈p is realized by c_{2,i,1}, giving s′_{21}=s_{21}+s_{i1}∉p. Necessarily i≠2, so this is one of the defined elementary matrices. The printed c_{i,1,1} need not affect row2: over Q, take s=((0,1,0),(0,0,1),(1,0,0)), det(s)=1, p=(0), i=3. Then s_{11}=s_{21}=0 and s_{31}=1; right multiplication in the printed convention by c_{3,1,1} leaves s′_{21}=0, whereas c_{2,3,1} makes s′_{21}=1. The later s⋆c_{1,2,λ} gives s′_{11}=s_{11}+λs_{21}, so those printed products and indices already prove the stated pivot and remain unchanged. The case n=1 is already the identity from det(s)=1.

[548] EN-CONTRIB-X-037 Source folio 557. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=571
Printed reading: $c_{i,1,\lambda}$
Corrected reading: $c_{1,i,\lambda}$
Source: check/findings/findings_556_560.json#17. Both scans print i,1 in both factors. Under the explicit556 multiplication rule, c_{1,i,λ}⋆s=s C_{1i}(λ) adds column1 to column i and gives s′_{1i}=s_{1i}+λs_{11}; with s_{11}=1 and λ=−s_{1i}, this clears the first row. Then s⋆c_{i,1,μ}=C_{i1}(μ)s adds row1 to row i, clearing s_{i1} with μ=−s_{i1} without reintroducing the zero first-row entries. The printed first factor c_{i,1,λ} only changes column1, so cannot clear s_{1i}; for s=((1,1),(0,1)) both printed factors leave the(1,2)entry1 under every such addition. Correct the FIRST factor to1,i, retaining the second i,1. The checker's suggested second-factor correction presupposes the other multiplication order.

[549] EN-CONTRIB-X-038 Source folio 562. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=576
Printed reading: To see this, by 1.14 one is reduced
Corrected reading: To see this, by 1.1.4 one is reduced
Source: ASTRA-X-0106. The full562 scan prints1.14. The directly re-inspected Corollary1.1.4 on520 (main PDF534, upper statement) says that Filt_d(X_t) is generated by the classes cl_•(O_Yt) of integral closed subschemes of dimension at most d. Because $\sigma_{_}\bullet$ is a group homomorphism, sending each such generator into Filt_d(X_s) sends every finite integer linear combination there: $\sigma_{_}\bullet(\Sigma\ n_Y[O_Y])=\Sigma\ n_Y\sigma_{_}\bullet([O_Y])$. This is exactly the reduction in7.4. The local§1 numbering ends at1.6;1.14 has no separate target. Restore the missing internal stop in the current reference.

[550] EN-CONTRIB-X-039 Source folio 567. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=581
Printed reading: flat over S, Denote
Corrected reading: flat over S. Denote
Source: check/findings/findings_566_570.json#10. The source mark has a descending comma tail. The attached diplomatic comma is exact, while the current full stop marks the end of the sentence before On.

[551] EN-CONTRIB-X-040 Source folio 574. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=588
Printed reading: $x_i : S \to X$
Corrected reading: $x_i : S \to C$
Source: check/findings/findings_571_575.json#7. Main and witness585 clearly print X. The preceding573 paragraph defines x_0,x_1 as sections of C→S, and the same574 sentence gives their base changes s→C_s. The exact square defining i_x is the base change of x:S→C by X×_S C→C, with i_x(z)=(z,x(f(z))). Thus the section named here has target C, as the attached current correction says; diplomatic retains X.

[552] EN-CONTRIB-X-041 Source folio 574. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=588
Printed reading:

$$\begin{array}{ccc} X & \xrightarrow{i_x} & X \times_S C \\ {\scriptstyle i}\uparrow & & \uparrow \\ X_s & \xrightarrow{i_{x_s}} & X_s \times_{k(x)} C_s \end{array} ,$$

Corrected reading:

$$\begin{array}{ccc} X & \xrightarrow{i_x} & X \times_S C \\ {\scriptstyle i}\uparrow & & \uparrow \\ X_s & \xrightarrow{i_{x_s}} & X_s \times_{k(s)} C_s \end{array} ,$$

Source: check/findings/findings_571_575.json#10. Both scans have k(x). The displayed square is the base change of X×_S C along s→S, with the canonical isomorphism (X×_S C)×_S s ≅ X_s×_{k(s)}C_s induced by the two projections and the fibre-product universal property. The x in the preceding definition is a section S→C. If its closed image is denoted x_s, then the section gives k(s)→k(x_s)→k(s) with composite identity, hence k(x_s)=k(s) through these maps. The current k(s) explicitly names the base field of the stated fibre product, without changing that identified product; retain the source k(x) diplomatically.

[553] EN-CONTRIB-X-042 Source folio 575. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=589
Printed reading: an element of $K^\bullet(T)$
Corrected reading: an element x of $K^\bullet(T)$
Source: check/findings/findings_571_575.json#14. The whole sentence introduces one element without giving it a name and subsequently multiplies that same element as nx. Insert its missing binder x immediately after élément. This names the existing quantified element; the map x↦nx and the printed assertion remain unchanged.

[554] EN-CONTRIB-X-043 Source folio 576. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=590
Printed reading: $\widehat{G}(H^{2*}(T,\mathbb{Z}_\ell(*))$
Corrected reading: $\widehat{G}(H^{2*}(T,\mathbb{Z}_\ell(*)))$
Source: check/findings/findings_576_580.json#2. Both scans leave the opening parenthesis of G-hat unclosed. The diplomatic omission matches; the existing current closing parenthesis balances precisely that outer argument.

[555] EN-CONTRIB-X-044 Source folio 576. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=590
Printed reading:

$$\begin{array}{ccc} K^\bullet(X_t) & \xrightarrow{c^i} & H^{2i}(X_t,\mathbb{Z}_\ell(i)) \\ {\scriptstyle\sigma}\downarrow & & \uparrow{\scriptstyle\sigma^{-1}} \\ K^\bullet(X_s) & \xrightarrow{c^i} & K^{2i}(X_s,\mathbb{Z}_\ell(i)) \end{array}, \qquad (7.13.2)$$

Corrected reading:

$$\begin{array}{ccc} K^\bullet(X_t) & \xrightarrow{c^i} & H^{2i}(X_t,\mathbb{Z}_\ell(i)) \\ {\scriptstyle\sigma}\downarrow & & \uparrow{\scriptstyle\sigma^{-1}} \\ K^\bullet(X_s) & \xrightarrow{c^i} & H^{2i}(X_s,\mathbb{Z}_\ell(i)) \end{array}, \qquad (7.13.2)$$

Source: check/findings/findings_576_580.json#4. Both scans clearly print K. The horizontal map c^i is defined immediately above by7.13.1 with codomain H^{2i}; the right vertical map is explicitly j^*∘(i^*)^{-1} on the same cohomology groups. Thus the vertex is H^{2i}(X_s,Z_ℓ(i)), exactly the existing current correction. Preserve K diplomatically.

[556] EN-CONTRIB-X-045 Source folio 578. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=592
Printed reading:

$$\begin{array}{ccc} K^\bullet(X_{\bar t}) & \xrightarrow{c^i} & H^{2i}(X_{\bar t},\mu_n^{\otimes i}) \\ {\scriptstyle\sigma}\downarrow & & \downarrow{\scriptstyle\sigma^{-1}} \\ K^\bullet(X_{\bar s}) & \xrightarrow{c^i} & H^{2i}(X_{\bar s},\mu_n^{\otimes i}) \end{array},$$

Corrected reading:

$$\begin{array}{ccc} K^\bullet(X_{\bar t}) & \xrightarrow{c^i} & H^{2i}(X_{\bar t},\mu_n^{\otimes i}) \\ {\scriptstyle\sigma}\downarrow & & \uparrow{\scriptstyle\sigma^{-1}} \\ K^\bullet(X_{\bar s}) & \xrightarrow{c^i} & H^{2i}(X_{\bar s},\mu_n^{\otimes i}) \end{array},$$

Source: check/findings/findings_576_580.json#7. The arrowhead is unmistakably downward in main and witness589. The exact map defined at the bottom of576 is α=σ^{-1}=j^*∘(i^*)^{-1}:H(X_s)→H(X_t). For a lift e∈K^•(X) of e_t, functoriality gives c_s^i(σ_K e_t)=i^*c_X^i(e), so α c_s^i σ_K(e_t)=j^*c_X^i(e)=c_t^i(e_t). Passing to the finite-coefficient geometric fibres preserves this equality. Consequently the α edge in this square runs from the lower cohomology vertex to the upper one. The next source sentence independently says to reverse this edge, producing the downward isomorphism in7.13.4; merely relabelling the first edge would contradict that stated operation. Correct its direction in current only, retaining σ^{-1}; diplomatic keeps the printed downward arrow.

[557] EN-CONTRIB-X-048 Source folio 580. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=594
Printed reading: (cf. 7.15)
Corrected reading: (cf. 7.14)
Source: ASTRA-X-REF-580-POSTFREEZE. The coordinator directly reread primary580,582 and584 at the cited loci. The580 sentence specifies the homomorphism on algebraic cycles by Z_t↦[Z_s], with Z the schematic closure inX. Equations7.14.1 and7.14.1bis on582 define exactly this homomorphism, first by dimension and then by codimension. On each prime cycle[W], the closure, its special fibre and all multiplicities are identical in the two descriptions; extension by additivity gives the same map on the free cycle groups. Section7.15 on584 proceeds with a morphism of traits and the naturality square for specialization ofK-groups. These constructions are related by the cycle-class map κ_a:[W]↦[O_W] in the associated gradedK-group: the closureZ is flat over the trait, so restriction of[O_Z] to the special fibre is[O_Zs], giving κ_sσ_cycle([W])=σ_Kκ_t([W]). Thus the reference repair identifies the exact cycle-level definition stated on580 while preserving its relation to the subsequentK-group specialization. Correct the locator7.15 to7.14 only in current; the independently decided extra closing parenthesis remains its separate source variant.

[558] EN-CONTRIB-X-046 Source folio 580. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=594
Printed reading:). This commutativity follows
Corrected reading: . This commutativity follows
Source: check/findings/findings_576_580.json#17. The full sentence has no unmatched opening parenthesis before this citation: the earlier modulo rien du tout parenthesis is already closed. The source's two closing parentheses leave exactly one excess closing mark. Remove that excess mark in current only.

[559] EN-CONTRIB-X-047 Source folio 581. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=595
Printed reading: the one considered in 7.11.5.
Corrected reading: the one considered in 7.12.5.
Source: ASTRA-X-0118. The full581 image prints7.11.5. Directly inspected7.11.5 on572 is the displayed quotient Gr_•^{alg}(T), followed by its module structure, with no conjecture. Directly inspected7.12.5 on575 states the conjecture that a numerically trivial cycle Z has some positive multiple nZ algebraically equivalent to zero. This is precisely the stronger conjecture being compared here to numerical equivalence implying Q_ℓ-cohomological equivalence. The cycle-class homomorphism is additive, so cℓ(nZ)=n cℓ(Z), and n is invertible in Q_ℓ; algebraic equivalence is killed by this class map (on a connected parameter curve the two point pullbacks of the family class agree). These exact maps give the stated implication. The reference must point to7.12.5; this corrects the locator without changing either historical conjecture.

[560] EN-CONTRIB-X-049 Source folio 584. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=598
Printed reading:

$$\begin{array}{ccc} K_\bullet(X_t) & \xrightarrow{\sigma_X} & K_\bullet(X_s) \\ \downarrow & & \downarrow \\ K_\bullet(X'_{t'}) & \xrightarrow{\sigma_X} & K_\bullet(X'_{s'}) \end{array} \tag{7.15.1}$$

Corrected reading:

$$\begin{array}{ccc} K_\bullet(X_t) & \xrightarrow{\sigma_X} & K_\bullet(X_s) \\ \downarrow & & \downarrow \\ K_\bullet(X'_{t'}) & \xrightarrow{\sigma_{X'}} & K_\bullet(X'_{s'}) \end{array} \tag{7.15.1}$$

Source: check/findings/findings_581_585.json#9. The lower subscript lacks a prime in both scans. The four objects are K_•(X_t),K_•(X_s),K_•(X′_{t′}),K_•(X′_{s′}); with vertical pullbacks h_t*,h_s*, the relation is h_s*σ_X=σ_X′h_t*. This relation does not identify the standalone maps. Concretely take A=R[[u]], A′=C[[u]], and X=Spec C[[u]] over Spec A. The canonical C⊗_R C≅C×C identifies X′ with two copies of Spec C[[u]]. The upper generic and special K_• groups are Z, the lower ones Z^2, and h_t*,h_s* send n to(n,n). The upper specialization is id_Z and the lower is id_Z^2; these maps have the indicated nonisomorphic groups as their domains and codomains. The existing current σ_X′ therefore names the specialization of the separately bound family X′, while diplomatic keeps the printed σ_X.

[561] EN-CONTRIB-X-050 Source folio 585. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=599
Printed reading: $B_i = A_{i\,\underline{m}_i}$
Corrected reading: $B_i = \bar{A}_{i\,\underline{m}_i}$
Source: check/findings/findings_581_585.json#12. Both scans omit the bar. The preceding sentence defines the ring A-bar_i and its maximal ideal m_i; localization at its complement gives B_i=(A-bar_i)_{m_i}. The exact localization map sends a to a/1 in that ring. This is the normalization and localization to which the printed Krull–Akizuki argument applies. The existing current bar identifies that defined ring; preserve the omission diplomatically.

[562] EN-CONTRIB-X-051 Source folio 585. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=599
Printed reading: $k_i = A_i/\underline{m}_i = k(s_i)$
Corrected reading: $k_i = \bar{A}_i/\underline{m}_i = k(s_i)$
Source: check/findings/findings_581_585.json#13. Both scans omit the bar. With B_i=(A-bar_i)_{m_i}, the residue field is canonically A-bar_i/m_i ≅ B_i/m_iB_i: the map a/s↦(a mod m_i)/(s mod m_i) is well-defined since s∉m_i, and is inverse to the map induced by localization. Thus the existing current bar names the actual defined quotient, and the diplomatic omission remains exact.

[563] EN-CONTRIB-X-052 Source folio 588. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=602
Printed reading: by representing K' as
Corrected reading: by representing k' as
Source: check/findings/findings_586_590.json#2. The source capital is clear, while the same paragraph introduces the pure field extension k′/k in lowercase. For k′=k(T_i), every finite set of elements lies in a localization of a polynomial ring in finitely many T_i, and these localizations form the stated directed system with union k′. Thus the object represented by this limit is exactly the introduced k′. Keep the source capital diplomatically and the already corrected lowercase in current.

[564] EN-CONTRIB-X-053 Source folio 591. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=605
Printed reading: $F(k) \to K(K)$
Corrected reading: $F(k) \to F(K)$
Source: check/findings/findings_591_595.json#3. Both scans unmistakably print K(K). The extension inclusion k→K induces F(k)→F(K) under the functor explicitly defined in7.18.1, and the first line of591 already names this map. The current F(K) corrects this notation only. The separate counterexample below shows that this typographical repair does not establish the claimed injectivity at the printed generality.

[565] EN-CONTRIB-X-054 Source folio 593. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=607
Printed reading: (by (IV 2.12).
Corrected reading: (by (IV 2.12)).
Source: check/findings/findings_591_595.json#10. The source opens an outer explanatory parenthesis and an inner reference parenthesis, but closes only the inner one. The current second closing mark balances the actual two openings; diplomatic remains exact.

[566] EN-CONTRIB-X-055 Source folio 593. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=607
Printed reading: is still of finite type
Corrected reading: is still noetherian
Source: ASTRA-X-0124. The source has just weakened X of finite type over k to X noetherian, while requiring the extension field K/k to be finitely generated. The asserted consequence is noetherianity. For an affine open Spec B⊂X, choose a finitely generated integral k-subalgebra A⊂K with Frac(A)=K and put S=A\{0}. The canonical map b⊗(a/s)↦(b⊗a)/(1⊗s) gives the isomorphism B⊗_k K ≅ S^{-1}(B⊗_k A). The ring B⊗_k A is finitely generated over the noetherian ring B, hence noetherian, and its localization is noetherian. A finite affine cover of X gives a finite such cover of X_K, proving that X_K is noetherian. The printed finite-type-over-the-base conclusion fails in the exact identity-base-change example X=Spec k(u), K=k: X is noetherian and K/k is finitely generated, but X_K=X is not of finite type over k, since Zariski's lemma would force the field k(u), if finitely generated as a k-algebra, to be finite algebraic over k. This establishes the current replacement and reverses the earlier rejected proposal; it does not insert an additional hypothesis.

[567] EN-CONTRIB-X-056 Source folio 593. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=607
Printed reading: . Moreover, essentially the same results
Corrected reading:). Moreover, essentially the same results
Source: check/findings/findings_591_595.json#14. The main scan leaves the parenthesis opened by de sorte que unclosed before De plus. The current closing parenthesis is independently necessary. The substantive de type fini phrase is corrected separately with the explicit localization proof.

[568] EN-CONTRIB-XII-001 Source folio 595. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=609
Printed reading: forged by I. Castelnuovo
Corrected reading: forged by G. Castelnuovo
Source: fixes/MISPRINTS_6.json#534. The capital I and its stop are clear. The historical university archive identifies the algebraic geometer as Guido Castelnuovo (1865–1952); the concise primary metadata is archived in review/expose_xii/evidence/castelnuovo.json. Correct the initial to G in current only.

[569] EN-CONTRIB-XII-002 Source folio 597. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=611
Printed reading: X_{S_i}
Corrected reading: X_{S_2}
Source: fixes/MISPRINTS_6.json#571. Main and witness608 show a dotted i, unlike the nearby flagged1. The sentence chooses the open S_2⊂S_1 on which the reduced components become integral over the base. Their restriction is X×_S S_2; the final S′=h^{-1}(V) lies inside S_2 because V excludes h(S_1−S_2). Thus current must use X_{S_2}, not the attached X_{S_1}.

[570] EN-CONTRIB-XII-003 Source folio 597. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=611
Printed reading: is integral over S_i (EGA IV 9.7.7)
Corrected reading: is integral over S_2 (EGA IV 9.7.7)
Source: fixes/MISPRINTS_6.json#573. The second index is the same dotted i in main and witness608. The integrality holds over the chosen open S_2, and the final S′ maps into that open. Correct the attached current S_1 to S_2 together with the preceding X restriction.

[571] EN-CONTRIB-XII-004 Source folio 597. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=611
Printed reading: $v = v_1 - h(S_1 - S_2)$
Corrected reading: $V = V_1 - h(S_1 - S_2)$
Source: fixes/MISPRINTS_6.json#574. Both v letters are visibly lower than the capital V in h^{-1}(V). The expression constructs the open V from V_1, so retain the current capitalization.

[572] EN-CONTRIB-XII-005 Source folio 597. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=611
Printed reading: we may suppose that Z_i is integral over S
Corrected reading: we may suppose that Z_i is integral over S'
Source: ASTRA-XII-0001. Both main and witness608 have S without a prime. The components X′_i are geometrically integral over S′ by1.3; the birational Chow maps Z_i→X′_i preserve their generic function fields, so the cited openness argument applies to Z_i→S′. Their structure maps to S are the composites Z_i→S′→S and need not have geometrically integral fibres. Explicitly, take S=Spec R, X=Spec C and S′=Spec C. Then X_{S′}=Spec(C⊗_R C)=Spec(C×C), each component and its identity Chow presentation is Spec C, geometrically integral over S′. Over S its geometric base change has two components, and S has no smaller nonempty open. The printed unprimed assertion is therefore false. Restore S′ in current, matching the following Pic_{Z_i/S′} construction.

[573] EN-CONTRIB-XII-006 Source folio 597. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=611
Printed reading: $f^* : \underline{\mathrm{Pic}}_{X'/S'} \longrightarrow \underline{\mathrm{Pic}}_{Z'/S'}$
Corrected reading: $f^* : \underline{\mathrm{Pic}}_{X'/S'} \longrightarrow \underline{\mathrm{Pic}}_{Z/S'}$
Source: fixes/MISPRINTS_6.json#575. A prime is printed after Z. The immediately defined map is f:Z=∐Z_i→X′; its pullback sends a line bundle on X′ to its pullback on Z. Retain current Z, without the spurious prime.

[574] EN-CONTRIB-XII-007 Source folio 600. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=614
Printed reading: $\mathrm{Pic}_{X_0/S} \longleftarrow \mathrm{Pic}_{\coprod X'_i/S} \longleftarrow \mathrm{Pic}_{Y/S}$.
Corrected reading: $\mathrm{Pic}_{X_0/S} \longleftarrow \mathrm{Pic}_{\coprod X_i/S} \longleftarrow \mathrm{Pic}_{Y/S}$.
Source: check/findings/findings_596_600.json#48. Main and witness611 both show a separate short mark above and to the right of X. Preserve that prime diplomatically. The preceding scheme sequence uses ∐X_i, and pullback of its two maps gives Pic_{X_0/S}←Pic_{∐X_i/S}←Pic_{Y/S}; retain the current removal of the prime.

[575] EN-CONTRIB-XII-008 Source folio 601. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=615
Printed reading: $f = \coprod_k f_k : \coprod_k (\coprod_{i \in I(k)} X_i) \longrightarrow Y_k$
Corrected reading: $f = \coprod_k f_k : \coprod_k (\coprod_{i \in I(k)} X_i) \longrightarrow \coprod_k Y_k$
Source: fixes/MISPRINTS_6.json#559. Each f_k maps its component coproduct into Y_k; the displayed map ∐_k f_k has domain the coproduct over all k and codomain ∐_k Y_k=X. The source omits the outer target coproduct. Retain its existing current correction.

[576] EN-CONTRIB-XII-009 Source folio 602. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=616
Printed reading: $L_1|Y_1 \cap Y_2 \to L_Z|Y_1 \cap Y_2$
Corrected reading: $L_1|Y_1 \cap Y_2 \to L_2|Y_1 \cap Y_2$
Source: fixes/MISPRINTS_6.json#561. Both scans show the straight diagonal and horizontal strokes of Z, unlike the curved2 in the adjacent Y_2. The target is the already defined L_2 on Y_2, so retain current L_2.

[577] EN-CONTRIB-XII-010 Source folio 603. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=617
Printed reading: $\mathbb{V}(E) \times_s \mathbb{V}(F)$
Corrected reading: $\mathbb{V}(E) \times_S \mathbb{V}(F)$
Source: fixes/MISPRINTS_6.json#563. Witness614 confirms that the fibre-product index is the small form s. Both representing vector bundles are over S, and a T-point is a pair (u,v) over the same T→S. Thus their representing product is over S; retain the current capital S.

[578] EN-CONTRIB-XII-011 Source folio 603. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=617
Printed reading:

$$O_Y \longrightarrow f_*O_Y \rightrightarrows g_*(O_{X\times_y X}) \qquad (*)$$

Corrected reading:

$$O_Y \longrightarrow f_*O_X \rightrightarrows g_*(O_{X\times_Y X}) \qquad (*)$$

Source: fixes/MISPRINTS_6.json#564; fixes/MISPRINTS_6.json#565.
fixes/MISPRINTS_6.json#564: Both scans plainly print subscript Y. The map f has source X, and its direct image sends sheaves on X to sheaves on Y; the unit O_Y→f_*O_X followed by the two pullbacks from X×_YX is the displayed descent sequence. Retain the current O_X.
fixes/MISPRINTS_6.json#565: The index descends below the baseline and is a small y in both scans. The two projections have common target X over the specified f:X→Y, so their fibre product is X×_YX. Retain the current capital Y.

[579] EN-CONTRIB-XII-012 Source folio 603. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=617
Printed reading: $g : X \times_y X \to Y$
Corrected reading: $g : X \times_Y X \to Y$
Source: fixes/MISPRINTS_6.json#566. The same small y is printed again. The structural map g is f∘p_1=f∘p_2:X×_YX→Y; retain the existing capital Y.

[580] EN-CONTRIB-XII-013 Source folio 604. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=618
Printed reading: $B_{i-1} \otimes_A B_{i-1} \xrightarrow{\sim} B_{i-1} \otimes_{A_i} B_{i-1}$
Corrected reading: $B_{i-1} \otimes_A B_{i-1} \xrightarrow{\sim} B_{i-1} \otimes_{B_i} B_{i-1}$
Source: ASTRA-XII-0019. The paragraph defines A=O_Y and B_0=f_*O_X, with A injected into B_0, then B_i=Eq(B_{i−1}⇉B_{i−1}⊗_A B_{i−1}). For every b∈B_i, b⊗1=1⊗b. Thus the quotient map B_{i−1}⊗_A B_{i−1}→B_{i−1}⊗_{B_i}B_{i−1} has inverse x⊗_{B_i}y↦x⊗_Ay: the equalizer relation multiplied by x⊗y makes this rule B_i-balanced, so the tensor universal property defines it and both composites fix pure tensors. The original f is finite and Y is noetherian; hence B_0 is a finite A-module and each A-submodule B_i is finite. Thus B_i/A is a finite A-module and A→B_i is a finite ring map. After the printed localization and stabilization B_i=B_{i+1}, multiplication B_i⊗_A B_i→B_i is inverse to b↦b⊗1, since every b lies in the equalizer. Hence Spec(B_i)→Spec(A)=Y is the finite monomorphism used in the proof. The printed Nakayama step makes A→B_i surjective; its original injectivity forces A=B_i and contradicts the nonempty stabilized support. This exact algebra calculation fixes all five corrupt expressions and also supplies the effective epimorphisms in the tower. The undefined A_i and B cannot supply these maps. Preserve the entire printed cluster diplomatically; this is a substantive source correction, also deciding rejected Codex SGA-SRC-001365.

[581] EN-CONTRIB-XII-014 Source folio 604. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=618
Printed reading: the decreasing sequence of B-modules $M_i = B_i/A$
Corrected reading: the decreasing sequence of A-modules $M_i = B_i/A$
Source: ASTRA-XII-0019. The paragraph defines A=O_Y and B_0=f_*O_X, with A injected into B_0, then B_i=Eq(B_{i−1}⇉B_{i−1}⊗_A B_{i−1}). For every b∈B_i, b⊗1=1⊗b. Thus the quotient map B_{i−1}⊗_A B_{i−1}→B_{i−1}⊗_{B_i}B_{i−1} has inverse x⊗_{B_i}y↦x⊗_Ay: the equalizer relation multiplied by x⊗y makes this rule B_i-balanced, so the tensor universal property defines it and both composites fix pure tensors. The original f is finite and Y is noetherian; hence B_0 is a finite A-module and each A-submodule B_i is finite. Thus B_i/A is a finite A-module and A→B_i is a finite ring map. After the printed localization and stabilization B_i=B_{i+1}, multiplication B_i⊗_A B_i→B_i is inverse to b↦b⊗1, since every b lies in the equalizer. Hence Spec(B_i)→Spec(A)=Y is the finite monomorphism used in the proof. The printed Nakayama step makes A→B_i surjective; its original injectivity forces A=B_i and contradicts the nonempty stabilized support. This exact algebra calculation fixes all five corrupt expressions and also supplies the effective epimorphisms in the tower. The undefined A_i and B cannot supply these maps. Preserve the entire printed cluster diplomatically; this is a substantive source correction, also deciding rejected Codex SGA-SRC-001365.

[582] EN-CONTRIB-XII-015 Source folio 604. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=618
Printed reading: $\mathrm{Spec}(A_i) \to Y$
Corrected reading: $\mathrm{Spec}(B_i) \to Y$
Source: ASTRA-XII-0019. The paragraph defines A=O_Y and B_0=f_*O_X, with A injected into B_0, then B_i=Eq(B_{i−1}⇉B_{i−1}⊗_A B_{i−1}). For every b∈B_i, b⊗1=1⊗b. Thus the quotient map B_{i−1}⊗_A B_{i−1}→B_{i−1}⊗_{B_i}B_{i−1} has inverse x⊗_{B_i}y↦x⊗_Ay: the equalizer relation multiplied by x⊗y makes this rule B_i-balanced, so the tensor universal property defines it and both composites fix pure tensors. The original f is finite and Y is noetherian; hence B_0 is a finite A-module and each A-submodule B_i is finite. Thus B_i/A is a finite A-module and A→B_i is a finite ring map. After the printed localization and stabilization B_i=B_{i+1}, multiplication B_i⊗_A B_i→B_i is inverse to b↦b⊗1, since every b lies in the equalizer. Hence Spec(B_i)→Spec(A)=Y is the finite monomorphism used in the proof. The printed Nakayama step makes A→B_i surjective; its original injectivity forces A=B_i and contradicts the nonempty stabilized support. This exact algebra calculation fixes all five corrupt expressions and also supplies the effective epimorphisms in the tower. The undefined A_i and B cannot supply these maps. Preserve the entire printed cluster diplomatically; this is a substantive source correction, also deciding rejected Codex SGA-SRC-001365.

[583] EN-CONTRIB-XII-016 Source folio 604. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=618
Printed reading: $B \subset A_i$
Corrected reading: $A \subset B_i$
Source: ASTRA-XII-0019. The paragraph defines A=O_Y and B_0=f_*O_X, with A injected into B_0, then B_i=Eq(B_{i−1}⇉B_{i−1}⊗_A B_{i−1}). For every b∈B_i, b⊗1=1⊗b. Thus the quotient map B_{i−1}⊗_A B_{i−1}→B_{i−1}⊗_{B_i}B_{i−1} has inverse x⊗_{B_i}y↦x⊗_Ay: the equalizer relation multiplied by x⊗y makes this rule B_i-balanced, so the tensor universal property defines it and both composites fix pure tensors. The original f is finite and Y is noetherian; hence B_0 is a finite A-module and each A-submodule B_i is finite. Thus B_i/A is a finite A-module and A→B_i is a finite ring map. After the printed localization and stabilization B_i=B_{i+1}, multiplication B_i⊗_A B_i→B_i is inverse to b↦b⊗1, since every b lies in the equalizer. Hence Spec(B_i)→Spec(A)=Y is the finite monomorphism used in the proof. The printed Nakayama step makes A→B_i surjective; its original injectivity forces A=B_i and contradicts the nonempty stabilized support. This exact algebra calculation fixes all five corrupt expressions and also supplies the effective epimorphisms in the tower. The undefined A_i and B cannot supply these maps. Preserve the entire printed cluster diplomatically; this is a substantive source correction, also deciding rejected Codex SGA-SRC-001365.

[584] EN-CONTRIB-XII-017 Source folio 604. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=618
Printed reading: $B = A_i$
Corrected reading: $A = B_i$
Source: ASTRA-XII-0019. The paragraph defines A=O_Y and B_0=f_*O_X, with A injected into B_0, then B_i=Eq(B_{i−1}⇉B_{i−1}⊗_A B_{i−1}). For every b∈B_i, b⊗1=1⊗b. Thus the quotient map B_{i−1}⊗_A B_{i−1}→B_{i−1}⊗_{B_i}B_{i−1} has inverse x⊗_{B_i}y↦x⊗_Ay: the equalizer relation multiplied by x⊗y makes this rule B_i-balanced, so the tensor universal property defines it and both composites fix pure tensors. The original f is finite and Y is noetherian; hence B_0 is a finite A-module and each A-submodule B_i is finite. Thus B_i/A is a finite A-module and A→B_i is a finite ring map. After the printed localization and stabilization B_i=B_{i+1}, multiplication B_i⊗_A B_i→B_i is inverse to b↦b⊗1, since every b lies in the equalizer. Hence Spec(B_i)→Spec(A)=Y is the finite monomorphism used in the proof. The printed Nakayama step makes A→B_i surjective; its original injectivity forces A=B_i and contradicts the nonempty stabilized support. This exact algebra calculation fixes all five corrupt expressions and also supplies the effective epimorphisms in the tower. The undefined A_i and B cannot supply these maps. Preserve the entire printed cluster diplomatically; this is a substantive source correction, also deciding rejected Codex SGA-SRC-001365.

[585] EN-CONTRIB-XII-018 Source folio 605. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=619
Printed reading:

$$\begin{array}{ccccccccc} (f_T)_*(O_{X'_T}) & \overset{\delta_T}{\to} & R^1(f_T)_*(I_T) & & & & & & \\ \uparrow & & \| & & & & & & \\ (f_T)_*(O^*_{X'_T}) & \overset{\delta^*_T}{\to} & R^1(f_T)_*(I_T) & \to & R^1(f_T)_*(O^*_{X_T}) & \to & R'(f_T)_*(O^*_{X'_T}) & \to & R^2(f_T)_*(I_T). \end{array}$$

Corrected reading:

$$\begin{array}{ccccccccc} (f_T)_*(O_{X'_T}) & \overset{\delta_T}{\to} & R^1(f_T)_*(I_T) & & & & & & \\ \uparrow & & \| & & & & & & \\ (f_T)_*(O^*_{X'_T}) & \overset{\delta^*_T}{\to} & R^1(f_T)_*(I_T) & \to & R^1(f_T)_*(O^*_{X_T}) & \to & R^1(f_T)_*(O^*_{X'_T}) & \to & R^2(f_T)_*(I_T). \end{array}$$

Source: fixes/MISPRINTS_6.json#569. Witness616 shows a short prime, not the stem and head of1 used in the neighbouring R^1. The map comes after R^1(f_T)_*O*_{X_T} and before the connecting map to R^2(f_T)_*I_T in the cohomology sequence of 1→I_T→O*_{X_T}→O*_{X′_T}→1. Its degree is therefore1; retain the current R^1.

[586] EN-CONTRIB-XII-019 Source folio 606. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=620
Printed reading: $h = \sum v(h_\beta) \otimes t_\beta \in \Gamma(X_T, \underline{O}_{X_T})$
Corrected reading: $h = \sum h_\beta \otimes t_\beta \in \Gamma(X_T, \underline{O}_{X_T})$
Source: ASTRA-XII-0027. Main and witness617 both print v(h_β) inside the defining sum. The already defined v maps Γ(X,O_X) to Γ(X,O_{X′}), so this expression lies on X′_T, while the following calculation requires a lift acting O_{X_T}-linearly on the additive exact sequence. Put h=Σh_β⊗t_β. Then v_T(h)=g−n, where n=Σn_β⊗t_β is nilpotent and nI_T=0 by(i). Since g is a unit, g−n is a unit. The nilpotent closed immersion X′_T⊂X_T then makes h a unit in every stalk of O_{X_T}, and the local inverses glue to a global inverse. Multiplication by h on 0→I_T→O_{X_T}→O_{X′_T}→0 induces the identity δ_T(h^{-1}g)=h^{-1}δ_T(g). On I_T the actions of h and g agree because nI_T=0. Thus the corrected definition proves exactly the next printed equality and inclusion; no extra assumption or changed theorem is introduced.

[587] EN-CONTRIB-XII-020 Source folio 606. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=620
Printed reading: $g = \sum g_\beta \otimes t_\alpha \in \Gamma(X_T, O_{X'_T})$
Corrected reading: $g = \sum g_\beta \otimes t_\beta \in \Gamma(X_T, O_{X'_T})$
Source: fixes/MISPRINTS_6.json#548. The index has the alpha form, unlike β in the adjacent g and all summands of the same decomposition. Each g_β is paired with its coefficient t_β, so retain current t_β.

[588] EN-CONTRIB-XII-021 Source folio 606. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=620
Printed reading: Let $g_\beta = v(h) + n_\beta$
Corrected reading: Let $g_\beta = v(h_\beta) + n_\beta$
Source: fixes/MISPRINTS_6.json#549. Both scans show no β after h. This is the repeated coefficientwise lift g_β=v(h_β)+n_β established above; retain the current index.

[589] EN-CONTRIB-XII-022 Source folio 606. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=620
Printed reading: where $n_\beta \in \mathrm{nilrad}(\Gamma(X, O_X))$; we then
Corrected reading: where $n_\beta \in \mathrm{nilrad}(\Gamma(X, O_{X'}))$; we then
Source: fixes/MISPRINTS_6.json#550. The prime is absent in both scans. Here n_β=g_β−v(h_β) is an element of Γ(X,O_{X′}), the codomain of v; retain the current prime.

[590] EN-CONTRIB-XII-023 Source folio 606. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=620
Printed reading:

$$0 \longrightarrow \mathbb{V}(E^\vee)(T) \longrightarrow R^1(f_t)_*(O^*_{X_T}) \longrightarrow R^1(f_T)_*(O_{X'_T}) \longrightarrow \mathbb{V}(Q)(T),$$

Corrected reading:

$$0 \longrightarrow \mathbb{V}(E^\vee)(T) \longrightarrow R^1(f_T)_*(O^*_{X_T}) \longrightarrow R^1(f_T)_*(O^*_{X'_T}) \longrightarrow \mathbb{V}(Q)(T),$$

Source: fixes/MISPRINTS_6.json#551; fixes/MISPRINTS_6.json#552. fixes/MISPRINTS_6.json#551: Witness617 confirms the small index, distinct from the capital T in the next term. The sequence is obtained by applying f_T to the sheaves on X_T; retain the current capital T. fixes/MISPRINTS_6.json#552: The blank space after O is clear in witness617. The preceding units sequence on605 gives the degree-one map from O*_{X_T} to O*_{X′_T} and its connecting map to R^2I_T; retain the current units star.

[591] EN-CONTRIB-XII-024 Source folio 608. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=622
Printed reading: $NI_i \subset I_{i-1}$
Corrected reading: $NI_{i-1} \subset I_i$
Source: ASTRA-XII-0032. Both main and witness619 print these two directions. The defined filtration is I_0=I⊃I_1⊃…⊃I_n=0 and X_i=V(I_i). The quotient O_Y/I_i→O_Y/I_{i−1} induces the closed immersion X_{i−1}↪X_i, whose ideal is J_i=I_{i−1}/I_i. For 1≤i≤n, NI_{i−1}⊂I_i gives (N/I_i)J_i=0, exactly hypothesis3.1(i) for this immersion. The printed example I_k=N^kI supplies NI_{i−1}=I_i. The printed reverse inequality is automatic and does not establish the needed annihilation. The reverse immersion is actually false for Y=Spec(k[ε]/ε^2), I_0=(ε), I_1=0: it would make Spec(k[ε]/ε^2) a closed subscheme of Spec k, requiring an impossible surjection k→k[ε]/ε^2. Correct both expressions together in current, preserving the filtration and all scheme definitions.

[592] EN-CONTRIB-XII-025 Source folio 608. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=622
Printed reading: $X_i \hookrightarrow X_{i-1}$
Corrected reading: $X_{i-1} \hookrightarrow X_i$
Source: ASTRA-XII-0032. Both main and witness619 print these two directions. The defined filtration is I_0=I⊃I_1⊃…⊃I_n=0 and X_i=V(I_i). The quotient O_Y/I_i→O_Y/I_{i−1} induces the closed immersion X_{i−1}↪X_i, whose ideal is J_i=I_{i−1}/I_i. For 1≤i≤n, NI_{i−1}⊂I_i gives (N/I_i)J_i=0, exactly hypothesis3.1(i) for this immersion. The printed example I_k=N^kI supplies NI_{i−1}=I_i. The printed reverse inequality is automatic and does not establish the needed annihilation. The reverse immersion is actually false for Y=Spec(k[ε]/ε^2), I_0=(ε), I_1=0: it would make Spec(k[ε]/ε^2) a closed subscheme of Spec k, requiring an impossible surjection k→k[ε]/ε^2. Correct both expressions together in current, preserving the filtration and all scheme definitions.

[593] EN-CONTRIB-XII-026 Source folio 609. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=623
Printed reading: $\underline{\mathrm{Pic}}_{X_{i-1}/S} \longrightarrow \underline{\mathrm{Pic}}_{X_i/S} \longrightarrow \mathbb{V}(Q_i)$
Corrected reading: $\underline{\mathrm{Pic}}_{X_i/S} \longrightarrow \underline{\mathrm{Pic}}_{X_{i-1}/S} \longrightarrow \mathbb{V}(Q_i)$
Source: ASTRA-XII-0037. Both directions are unambiguous in main and witness620. With the defined X_i=V(I_i), the quotient O_Y/I_i→O_Y/I_{i−1} gives j_i:X_{i−1}↪X_i; the map on Picard functors sends L to j_i^*L, hence Pic_{X_i/S}→Pic_{X_{i−1}/S}. Proposition3.1 appends its obstruction map to V(Q_i) to that direction. The composite j_n∘…∘j_1 is the original f:X=X_0↪X_n=Y; on the fibre over s its pullback is f_s^*:Pic^0_{Y_s/s}→Pic^0_{X_s/s}. These are the maps whose successive obstruction quotients the remark invokes. Restore both pullback directions in current as the continuation of the linked608 correction; preserve both printed formulas diplomatically.

[594] EN-CONTRIB-XII-027 Source folio 609. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=623
Printed reading: $\operatorname{coker}(\underline{\mathrm{Pic}}^0_{X_s/s} \longrightarrow \underline{\mathrm{Pic}}^0_{Y_s/s})$
Corrected reading: $\operatorname{coker}(\underline{\mathrm{Pic}}^0_{Y_s/s} \longrightarrow \underline{\mathrm{Pic}}^0_{X_s/s})$
Source: ASTRA-XII-0037. Both directions are unambiguous in main and witness620. With the defined X_i=V(I_i), the quotient O_Y/I_i→O_Y/I_{i−1} gives j_i:X_{i−1}↪X_i; the map on Picard functors sends L to j_i^*L, hence Pic_{X_i/S}→Pic_{X_{i−1}/S}. Proposition3.1 appends its obstruction map to V(Q_i) to that direction. The composite j_n∘…∘j_1 is the original f:X=X_0↪X_n=Y; on the fibre over s its pullback is f_s^*:Pic^0_{Y_s/s}→Pic^0_{X_s/s}. These are the maps whose successive obstruction quotients the remark invokes. Restore both pullback directions in current as the continuation of the linked608 correction; preserve both printed formulas diplomatically.

[595] EN-CONTRIB-XII-028 Source folio 610. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=624
Printed reading: $R^1 f_* O_X$
Corrected reading: $R^i f_* O_X$
Source: ASTRA-XII-0040. The enlarged main crop and witness621 show a connected flagged stem and base of digit1, with no separate dot; the adjacent j and the later i exponents have separate dots. The hypothesis quantifies i,j≥0 and the next paragraph explicitly applies the Künneth base-change formula to all R^if_*O_X and R^ig_*O_{X×_YX}. Thus i is the required varying exponent. Both attached layers had silently supplied it. Restore the printed1 only in diplomatic, retaining the correct current i.

[596] EN-CONTRIB-XII-029 Source folio 611. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=625
Printed reading:

$$\Gamma\ Y, N) \longrightarrow \Gamma(X, f^*N) \rightrightarrows \Gamma(X \times_Y X, g^*N)$$

Corrected reading:

$$\Gamma(Y, N) \longrightarrow \Gamma(X, f^*N) \rightrightarrows \Gamma(X \times_Y X, g^*N)$$

Source: fixes/MISPRINTS_6.json#579. Both scans show a blank gap after Γ and a closing parenthesis after N. The three-term global-section sequence requires Γ(Y,N); retain the existing current restoration.


[597] EN-CONTRIB-XII-030 Source folio 612. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=626
Printed reading: of gluing data on f^*L
Corrected reading: of gluing data on L
Source: ASTRA-XII-0047. Main and witness623 both print f*L. The preceding611 paragraph fixes L as a line bundle on X, while f:X→Y. The next sentence defines Rec(T) as Isom_{X_T×_{Y_T}X_T}(p_1^*L_T,p_2^*L_T), where p_i:X_T×_{Y_T}X_T→X_T. These are precisely the gluing data on L_T. The effective subfunctor assigns the data induced by a line bundle M on Y_T and an isomorphism f_T^*M≃L_T; such an M is the object still being sought. Thus the opening object is L, not an additional pullback f*L. Removing f* makes the sentence agree with its explicit functor definition without assuming effectivity.

[598] EN-CONTRIB-XII-031 Source folio 614. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=628
Printed reading: $R^n f_*(F)$ is S-flat
Corrected reading: $R^i f_*(F)$ is S-flat
Source: fixes/MISPRINTS_6.json#584. Both scans show n, without a dot. The preceding replacement by S_{n+1} gives flatness of R^if_*F for every i≥n+1; the next R^n is the degree now being treated. Retain current i in the first occurrence and n in the next.


[599] EN-CONTRIB-XII-032 Source folio 614. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=628
Printed reading: O_Y
Corrected reading: O_X
Source: fixes/MISPRINTS_6.json#585. Both scans print Y. The lemma defines λ:X→S and a morphism of sheaves on X; the source and target therefore carry the O_X-module structures used in its kernel and cokernel proof. Retain the existing current O_X.

[600] EN-CONTRIB-XII-033 Source folio 615. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=629
Printed reading: the formation of $\operatorname{Coker} u$ commutes with the base diagram
Corrected reading: the formation of $\operatorname{Coker} u$ commutes with base change
Source: fixes/MISPRINTS_6.json#586. The joined printed word is diagramme. The asserted operation is base change: for a ring map A→B, right exactness gives Coker(u)⊗_A B≃Coker(u⊗_A B). The paragraph then uses that fact to identify the vanishing locus. Retain current changement, while keeping diagramme diplomatically.

[601] EN-CONTRIB-XIII-001 Source folio 617. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=631
Printed reading:



Corrected reading:



Source: fixes/MISPRINTS_6.json#553. The final closing parenthesis is absent. Keep the diplomatic omission and the already supplied current parenthesis.

[602] EN-CONTRIB-XIII-002 Source folio 618. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=632
Printed reading: by 2.3 (i).
Corrected reading: by 1.3 (i).
Source: fixes/MISPRINTS_6.json#554. The source prints2.3. Proposition1.3(i) on p.616 explicitly states n-regularity for every n≥m, which gives the two vanishings used here by the definition1.1. Correct the reference to1.3(i) in current.

[603] EN-CONTRIB-XIII-003 Source folio 619. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=633
Printed reading: by 2.3 (ii);
Corrected reading: by 1.3 (ii);
Source: fixes/MISPRINTS_6.json#555. The source prints2.3. The preceding-page beta_n is the multiplication map whose surjectivity is exactly Proposition1.3(ii) on p.616. Correct to1.3(ii) in current.

[604] EN-CONTRIB-XIII-004 Source folio 619. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=633
Printed reading: $\mapsto n^1(F(n))$
Corrected reading: $n \mapsto h^1(F(n))$
Source: fixes/MISPRINTS_6.json#556. The arrow has no source variable and the following letter is n. The preceding inequality concerns h^1(F(n)); the current n↦h^1(F(n)) completes that explicitly specified function. Keep it current-only.

[605] EN-CONTRIB-XIII-005 Source folio 620. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=634
Printed reading: the open subset of S
Corrected reading: the open subset of S'
Source: fixes/MISPRINTS_6.json#557. Main and witness631 show no prime. The same sentence defines the parameter space as S′; T parametrizes the stated regular sequences inside that very S′. Supply the missing prime in current.

[606] EN-CONTRIB-XIII-006 Source folio 620. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=634
Printed reading: $0 \to F(1) \to F \to G \to 0$
Corrected reading: $0 \to F(-1) \to F \to G \to 0$
Source: fixes/MISPRINTS_6.json#558. The minus position is blank in both scans. The proof explicitly subtracts chi(F(n-1)), which is the additivity formula for the sequence with F(-1). Restore that minus in current only.

[607] EN-CONTRIB-XIII-007 Source folio 621. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=635
Printed reading: $(b_i, \dots, b_{s-1}, a_s)$
Corrected reading: $(b_1, \dots, b_{s-1}, a_s)$
Source: fixes/MISPRINTS_6.json#587. The first index has the dot and stem of i. The opening line makes G a (b_1,...,b_r)-faisceau, and replacing its final bound by a_s retains initial index1. Keep the current correction and diplomatic i.

[608] EN-CONTRIB-XIII-008 Source folio 622. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=636
Printed reading: $m = 0$
Corrected reading: $m \geq 0$
Source: ASTRA-XIII-0008. Both main PDF636 and witness633 print equality. Definition1.9 gives P_0(X_0)=X_0 because P_{-1}=0 and binom(-1,0)=1; hence m=c_0≥b_0−a_0≥0 by1.8. For a length-one skyscraper with a_0=b_0=1 and permitted c_0=1, m=1. The asserted equality is therefore false; the required and proved assertion is m≥0.

[609] EN-CONTRIB-XIII-009 Source folio 624. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=638
Printed reading: $M = \max(\chi(F_K(m))$
Corrected reading: $M = \max(\chi(F_K(m)))$
Source: fixes/MISPRINTS_6.json#589. The image has three opening parentheses and only two closing ones. The existing current closing parenthesis completes the displayed maximum; retain the variant.

[610] EN-CONTRIB-XIII-010 Source folio 625. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=639
Printed reading: $L'_K \to E'_K \to E_K$
Corrected reading: $L_{1,K} \to E'_K \to E_K$
Source: fixes/MISPRINTS_6.json#590. The prime is printed. The preceding surjection L_1→E′ induces L_{1,K}→E′_K, and $\gamma=\beta$ composed with this map is lifted by δ:L_{1,K}→L_K. Thus the source of this composition is exactly L_{1,K}; apply that correction only in current.

[611] EN-CONTRIB-XIII-011 Source folio 626. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=640
Printed reading: $0 \geq \langle c(L) \cdot c_1(H)^{r-1} \rangle = \langle c_1(H)^{r-1} \cdot Y \rangle \geq 1$
Corrected reading: $0 \geq \langle c_1(L) \cdot c_1(H)^{r-1} \rangle = \langle c_1(H)^{r-1} \cdot Y \rangle \geq 1$
Source: fixes/MISPRINTS_6.json#591. The first c on the line lacks an index; the other two c_1 terms have it. The expression is d(L), defined in2.2 as the intersection with c_1(L), so the existing current correction is the intended first Chern class.

[612] EN-CONTRIB-XIII-012 Source folio 626. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=640
Printed reading: (with $r-1$ zeros)
Corrected reading: (with r zeros)
Source: ASTRA-XIII-0018. The tuple (b_0,...,b_r)=(0,...,0,a_r) has one zero for every index0,...,r−1, hence r zeros. The proof on627 explicitly gives (0,a_1) for r=1 and adjoins the initial zero in the induction.

[613] EN-CONTRIB-XIII-013 Source folio 627. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=641
Printed reading: By 1.7 and 2.6,
Corrected reading: By 1.7 and 2.5,
Source: fixes/MISPRINTS_6.json#592. The printed reference is2.6. The immediately preceding lemma2.5 states the (0,...,a_r)-bound used for L_Y; its proof on this same page explicitly supplies the curve case (0,a_1). Correct this reference to2.5 in current.

[614] EN-CONTRIB-XIII-014 Source folio 628. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=642
Printed reading: $h^2(L(-a_1 - 2) = 0$
Corrected reading: $h^2(L(-a_1 - 2)) = 0$
Source: fixes/MISPRINTS_6.json#593. One closing parenthesis is printed. Keep the omission diplomatically and the already restored parenthesis in current.

[615] EN-CONTRIB-XIII-015 Source folio 628. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=642
Printed reading:

$$a_0 \geq \langle c_1(L)^2 \rangle + 2e_0 + \sum_{i=1}^{2} (2e_i - a_i) \binom{\delta - 1 + i}{i} + \beta(\beta + 1).$$

Corrected reading:

$$a_0 \geq \langle c_1(L)^2 \rangle + 2e_0 + \sum_{i=1}^{2} (2e_i - a_i) \binom{\delta - 1 + i}{i} - \beta(\beta + 1).$$

Source: fixes/MISPRINTS_6.json#594. The plus is clear. The same-page identity is a_0=2α_2+2α_0−Σ_{i≥1}a_i binom(δ−1+i,i)−β_0, with β_0≤β(β+1). Substitution therefore gives the negative final term in the lower bound. Apply that sign correction in current.

[616] EN-CONTRIB-XIII-016 Source folio 629. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=643
Printed reading: for $n \gg 0$, $h^1(L(n)) = 0$ (see FGA
Corrected reading: for $n \gg 0$, $h^1(L(-n)) = 0$ (see FGA
Source: fixes/MISPRINTS_6.json#595. The print has no minus sign. In2.9.1 the two end groups vanish for n≥2+h_2, yielding isomorphisms H^1(L(-n-1))→H^1(L(-n)); the cited eventual vanishing must concern these same negatively twisted groups to complete(i). Restore the missing minus in current.

[617] EN-CONTRIB-XIII-017 Source folio 630. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=644
Printed reading:

$$a_{m,i} = \sum_{j=0}^{r-i} a_{j+i} \binom{m-1+i}{j}$$

Corrected reading:

$$a_{m,i} = \sum_{j=0}^{r-i} a_{j+i} \binom{m-1+j}{j}$$

Source: fixes/MISPRINTS_6.json#597. The top index is dotted i. The exact change-of-basis identity is binom(m+n+q,q)=Σ_{i=0}^q binom(m−1+q−i,q−i) binom(n+i,i); setting q=j+i gives binom(m−1+j,j). For r=1 the translated constant coefficient is a_0+m a_1, also forcing this correction.

[618] EN-CONTRIB-XIII-018 Source folio 631. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=645
Printed reading: $\delta = \mathrm{supp}\{d(L_K)\}$
Corrected reading: $\delta = \sup\{d(L_K)\}$
Source: fixes/MISPRINTS_6.json#599. The print has two p letters. The argument chooses an integer upper bound on all degrees and subtracts it by twisting; this is the supremum. The existing current sup correction matches that explicitly stated operation.

[619] EN-CONTRIB-XIII-019 Source folio 632. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=646
Printed reading: the pullback of L_α, where
Corrected reading: the pullback of L'_α, where
Source: fixes/MISPRINTS_6.json#600. The prime position is blank. The map X_{Z′}→X_{Z′_α} pulls back the explicitly defined L′_α to L′; restore that prime in current.

[620] EN-CONTRIB-XIII-020 Source folio 632. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=646
Printed reading: the pullback of L'_α. to $X_{Z''_\beta}$
Corrected reading: the pullback of L'_α to $X_{Z''_\beta}$
Source: fixes/MISPRINTS_6.json#601. A round mark at the subscript baseline persists in main and witness643, immediately after L′_α. It interrupts the phrase image inverse de ... sur and has no formula role. Preserve the visible stop diplomatically and the existing current omission.

[621] EN-CONTRIB-XIII-021 Source folio 635. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=649
Printed reading: X_i/S'
Corrected reading: X'_i/S'
Source: fixes/MISPRINTS_6.json#603. The components are explicitly named X′_i two lines earlier. Each inclusion X′_i→X_{S′}, followed by the coproduct map, gives the surjection used in the next step; restore the same component names in current.

[622] EN-CONTRIB-XIII-022 Source folio 635. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=649
Printed reading: $\coprod X_i/S'$
Corrected reading: $\coprod X'_i/S'$
Source: fixes/MISPRINTS_6.json#604. This is the coproduct of the X′_i defined above, with its canonical surjection to X_{S′}. Restore the prime together with the preceding occurrence.

[623] EN-CONTRIB-XIII-023 Source folio 638. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=652
Printed reading: S'_α-morphism
Corrected reading: S_α-morphism
Source: fixes/MISPRINTS_6.json#606. A prime is printed. The preceding object X′ is an S_α-scheme and its displayed map X′→X_{S_α} is the model over S_α of X′_K→X_K. Correct the base label to S_α in current.

[624] EN-CONTRIB-XIII-024 Source folio 638. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=652
Printed reading: by Abhyankhar).
Corrected reading: by Abhyankar).
Source: fixes/MISPRINTS_6.json#607. The extra h is printed. Bibliography entry[1] on666 (main PDF680, author/title region inspected separately) spells the same resolution author Abhyankar. Correct the name in current.

[625] EN-CONTRIB-XIII-025 Source folio 639. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=653
Printed reading: Consequently, $\mathcal{L}$ is bounded
Corrected reading: Consequently, $(f^*)^{-1}(\mathcal{L})$ is bounded
Source: ASTRA-XIII-0025. The map is f*:Pic_{Y/S}→Pic_{X/S}. Page638 fixes a bounded family L on X; the displayed coefficients on639 belong to M_K in its inverse-image family (f*)^{-1}(L) on Y. Lemma2.11 applies to those M_K and proves boundedness of exactly that inverse image, which is the assertion required by Definition3.3. Main and witness650 print only L in the conclusion.

[626] EN-CONTRIB-XIII-026 Source folio 643. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=657
Printed reading: a connected k-scheme
Corrected reading: a connected K-scheme
Source: fixes/MISPRINTS_6.json#617. The letter is small k, clear in main and witness654. The preceding field is K and z,z_0 are maps Spec(K)→Z whose composites with g:Z→G are compared in G(K); the connected parameter scheme is over that same K. Correct k to K in current.

[627] EN-CONTRIB-XIII-027 Source folio 644. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=658
Printed reading: $\Phi(G'^{\tau}) \supset G^{\tau}$
Corrected reading: $\Phi^{-1}(G'^{\tau}) \supset G^{\tau}$
Source: fixes/MISPRINTS_6.json#618. The missing exponent is visible. The map Φ:G→G′ sends G^τ into G′^τ by4.1; pulling back that inclusion gives G^τ⊂Φ^{-1}(G′^τ), the half of4.2 being recalled. Keep the existing current inverse-image correction.

[628] EN-CONTRIB-XIII-028 Source folio 644. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=658
Printed reading: $O_Y(1))$
Corrected reading: $O_Y(1)$, under the hypotheses of 3.8)
Source: ASTRA-XIII-0030. The §4 opening (643) imposes no governing dimension hypothesis:4.1 introduces arbitrary S and a commutative group functor;4.2 assumes geometric-fibre representability and finite-type Φ_K. Lemma4.3 separately introduces proper X,Y/S. Its hyperplane-section parenthesis omits the hypotheses of the theorem3.8 explicitly invoked in its proof. Theorem3.8 (636–637) assumes noetherian S, projective Y with the specified ample O_Y(1), X its section zero-scheme, and dimension at least2 for every X_s∩Y_s^i; it concludes that f*:Pic_{Y/S}→Pic_{X/S} is of finite type. That finite-type pullback is the hypothesis needed in4.2 to identify the inverse image of the τ subgroup. For the printed unrestricted claim, take Y=P^1_k and X a hyperplane point: f*:Pic(P^1_k)=Z→Pic(X)=0 is zero, so the inverse image of Pic^τ(X)=0 is Z, whereas Pic^τ(P^1_k)=0. Both the page context and this explicit counterexample confirm the missing qualification. The current-only insertion refers to the full established hypotheses of the cited3.8; it is a substantive mathematical source correction, not a typographic emendation. Coordinator independently inspected the relevant source regions and concurred.

[629] EN-CONTRIB-XIII-029 Source folio 647. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=661
Printed reading: $L^{\otimes r}$
Corrected reading: $L^{\otimes p}$
Source: fixes/MISPRINTS_6.json#609. The first exponent is r. The two chosen integers are p,q and the quotient L^{⊗p}⊗(L^{⊗q})^{-1}=L^{⊗(p−q)} is the line bundle in the conclusion. Retain diplomatic r and the existing current p.

[630] EN-CONTRIB-XIII-030 Source folio 649. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=663
Printed reading: $\coprod X'_i \to X_S$
Corrected reading: $\coprod X'_i \to X_{S'}$
Source: fixes/MISPRINTS_6.json#611. The target subscript has no prime; its following mark is the sentence stop. The X′_i were just defined as components of X_{S′}, and the displayed canonical coproduct map is the union of their inclusions into X_{S′}. Keep that established current target.

[631] EN-CONTRIB-XIII-031 Source folio 651. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=665
Printed reading: KÜnneth
Corrected reading: Künneth
Source: fixes/MISPRINTS_6.json#620. Main and witness662 show the capital-height U. NOTE_6 permits restoring accents on typed capitals, so diplomatic KÜnneth preserves that capital while current Künneth supplies the correct internal lower-case letter. The coordinator concurred with this explained local treatment.

[632] EN-CONTRIB-XIII-032 Source folio 652. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=666
Printed reading: $\mathrm{rang}(C^1_{\mathrm{mum}}(X)) \leq b^2(X)$
Corrected reading: $\mathrm{rang}(C^1_{\mathrm{num}}(X)) \leq b^2(X)$
Source: fixes/MISPRINTS_6.json#622. The first index letter has the three stems of m. The group is the C^p_num defined in5.2, with p=1; keep the existing current num correction.

[633] EN-CONTRIB-XIII-033 Source folio 654. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=668
Printed reading: $M_1 = M \cdot \binom{p+N}{N} - \sum a_i \binom{m_1+i}{i} \quad (= \chi(I(p))$
Corrected reading: $M_1 = M \cdot \binom{p+N}{N} - \sum a_i \binom{m_1+i}{i} \quad (= \chi(I(p)))$
Source: fixes/MISPRINTS_6.json#624. There are three opening parentheses and two closing ones, followed by the stop. Keep the diplomatic omission and existing current closing parenthesis.

[634] EN-CONTRIB-XIII-034 Source folio 655. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=669
Printed reading: $F' = O_X.\sigma$
Corrected reading: $F' = O_P.\sigma$
Source: ASTRA-XIII-0042. Lemma6.5 invokes6.1, which defines F as a coherent sheaf on P=ℙ^N_k; no X is introduced in these hypotheses. A section σ∈H^0(P,F) gives the evaluation morphism O_P→F, a↦aσ, and its image is exactly O_P·σ. Correct the ambient-scheme subscript to P in current. The later6.8 separately introduces X.

[635] EN-CONTRIB-XIII-035 Source folio 656. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=670
Printed reading: $O_X(1)$
Corrected reading: $O_P(1)$
Source: ASTRA-XIII-0046. Proposition6.6 invokes6.1, where F is a coherent sheaf on P=ℙ^N_k. Its diagram is induced by multiplication by a linear section σ:O_P(-1)→O_P and the tensor maps F(-1)→F, N_q(-1)→N_q and F_q(-1)→F_q. Thus the line bundle supplying the section is O_P(1); X is not introduced in these hypotheses.

[636] EN-CONTRIB-XIII-036 Source folio 657. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=671
Printed reading: $h^0(F_q(-1) \leq h^0(F(-1)) \leq b_0$
Corrected reading: $h^0(F_q(-1)) \leq h^0(F(-1)) \leq b_0$
Source: fixes/MISPRINTS_6.json#613. Only the inner closing parenthesis follows −1; witness668 confirms the omission. Keep diplomatic and the existing current completion.

[637] EN-CONTRIB-XIII-037 Source folio 657. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=671
Printed reading:

$$|a_i^{(q)}| \leq A_{r-i}^{(r-q)}(c_{q-1}, \ldots, c_r; m).$$

Corrected reading:

$$|a_i^{(q)}| \leq A_{r-i}^{(r-q+1)}(c_{q-1}, \ldots, c_r; m).$$

Source: ASTRA-XIII-0047. Both main and witness668 show the defining family A_i^{(q)}(X_0,...,X_q;Y), with q+1 X-slots, and the superscript r−q in(ii). Section6.1 specifies q≥1;6.7(ii) lists i=0,...,q−1 while its Hilbert coefficients are indexed0,...,r. In the displayed coefficient range1≤q≤r+1, the argument list c_{q−1},...,c_r has r−q+2 entries and starts at a nonnegative index. It therefore requires superscript r−q+1 under the given family definition. At q=r the list is (c_{r−1},c_r), needing A^{(1)} rather than printed A^{(0)}; at q=r+1 it is (c_r), needing A^{(0)}, and F_{r+1}=0 because N_{r+1}=F. The proof cites6.4(iii) and6.6, which use the bound beginning at c_{q−1};6.8(b_q) independently retains coefficients of degrees≥q−1. Preserve that supported argument list and correct the superscript only, without adding hypotheses or conventions for other indices.

[638] EN-CONTRIB-XIII-038 Source folio 657. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=671
Printed reading: $a_{q-1}^{(q)} \leq a_{q-1}$ and $a_q^{(q)}$ and $a_q^{(q)} = a_q, \ldots, a_r^{(q)} = a_r$
Corrected reading: $a_{q-1}^{(q)} \leq a_{q-1}$ and $a_q^{(q)} = a_q, \ldots, a_r^{(q)} = a_r$
Source: fixes/MISPRINTS_6.json#614. Both scans print the duplicated incomplete first phrase. The sequence 0→N_q→F→F_q→0 subtracts a Hilbert polynomial of degree≤q−1, giving a_i^{(q)}=a_i for i≥q. The existing current omission removes the duplicated fragment while preserving those equalities.

[639] EN-CONTRIB-XIII-039 Source folio 657. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=671
Printed reading: $0 \to F(-1 \to F \to G \to 0$
Corrected reading: $0 \to F(-1) \to F \to G \to 0$
Source: fixes/MISPRINTS_6.json#615. Both scans omit the closing parenthesis after −1. Keep diplomatic and the existing current completion.

[640] EN-CONTRIB-XIII-040 Source folio 660. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=674
Printed reading: $0 \to O_X(-Z) \to O_X \to O_Z \to 0$
Corrected reading: $0 \to O_Z(-Z) \to O_{X_n} \to O_Z \to 0$
Source: ASTRA-XIII-0048. Main and witness671 both print X in the first two subscripts, with no n. The immediately preceding definition is O_{X_n}=O_Y/O_Y(-2Z). Put I=O_Y(-Z). The quotient map O_Y/I²→O_Y/I has kernel I/I², which is isomorphic to O_Z(-Z) since Z is Cartier on Y. Hence the exact sequence used here is 0→O_Z(-Z)→O_{X_n}→O_Z→0. Its twist by −1 and the printed adjunction formula give a_0=χ(O_{X_n}(-1))=−Z²−2d+2−2p_a(Z)=(n−6)d+4−4p_a(Z), establishing the stated divergence while a_1=2d. Correct these two subscripts in current only.

[641] EN-CONTRIB-XIII-041 Source folio 661. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=675
Printed reading: (I. Hermann).
Corrected reading: (G. Hermann).
Source: ASTRA-XIII-0066. Both main and witness672 clearly print I. Hermann. The primary Springer record for the polynomial-ideal paper identifies its author as Grete Hermann and cites Hermann, G.; the short metadata evidence is archived in review/xiii/evidence/bibliographic_verification.json. Correct the first initial to G in current only.

[642] EN-CONTRIB-XIII-042 Source folio 661. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=675
Printed reading: every ideal of
Corrected reading: every ideal I of
Source: fixes/MISPRINTS_6.json#626. Main and witness672 omit the name of the ideal, while the same sentence then uses $\sqrt{\text{I}}$ and I. Supply that already used name immediately after idéal in current only.

[643] EN-CONTRIB-XIII-043 Source folio 661. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=675
Printed reading: $\chi(O_X(n)) = \sum_{i=0}^{n} a_i \binom{n+i}{i}$
Corrected reading: $\chi(O_X(n)) = \sum_{i=0}^{N} a_i \binom{n+i}{i}$
Source: fixes/MISPRINTS_6.json#629. The upper limit is a small n in both scans. It is the argument of the Hilbert polynomial, whereas N is the fixed ambient dimension and the upper limit of the preceding b_i sum. Retain the existing current N.

[644] EN-CONTRIB-XIII-044 Source folio 661. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=675
Printed reading: $|a_i| \leq {}_i$
Corrected reading: $|a_i| \leq f_i$
Source: fixes/MISPRINTS_6.json#630. Neither scan has an f in the blank cell. The next words define f_i=A_{N−i}(e_i,...,e_N;0), identifying the missing bound. Keep the diplomatic omission and current f_i.

[645] EN-CONTRIB-XIII-045 Source folio 663. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=677
Printed reading:

$$H^1(N_Y^{\otimes n}(p)) \longrightarrow H^2(N^{\otimes n}(p-1) \longrightarrow H^2(N^{\otimes n}(p)) \longrightarrow 0$$

Corrected reading:

$$H^1(N_Y^{\otimes n}(p)) \longrightarrow H^2(N^{\otimes n}(p-1)) \longrightarrow H^2(N^{\otimes n}(p)) \longrightarrow 0$$

Source: fixes/MISPRINTS_6.json#631. Main and witness674 show only the closing parenthesis of p−1 before the arrow. Keep the diplomatic omission and current completion of H^2's argument.

[646] EN-CONTRIB-XIII-046 Source folio 664. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=678
Printed reading: O_Y
Corrected reading: O_X
Source: fixes/MISPRINTS_6.json#632. The printed subscript is Y. The ideal defining the curve Y⊂X is ker(O_X→O_Y); its blowup is Proj_X(⊕_{n≥0}I^{n}). The source sheaf of this explicitly used ideal is O_X. Retain the existing current correction.

[647] EN-CONTRIB-XIII-047 Source folio 664. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=678
Printed reading: not numerically equivalent to
Corrected reading: not numerically proportional to
Source: ASTRA-XIII-0064. Both statements print non-equivalence to H, but their strict conclusions fail for X=$\mathbb{P}^2$, H=O(1), M=O(2): M and H have different degrees on a line, whereas h=H^2=1, d=M·H=2 and M^2=4=d^2/h. The same example lies within7.4 with r=2. The proof uses the homomorphism Pic(X)→A^1_num(X)_Q induced by c_1: for L=M^{⊗h}⊗H^{⊗−d}, [L]=h[M]−d[H] and L·H=0. Numerical nonproportionality makes [L] nonzero, so the established7.1 gives L^2<0. Expanding L^2=h^2M^2−hd^2 yields precisely M^2<d^2/h. For7.4 put h=deg(c_1(H)^r), d=deg(c_1(M)c_1(H)^{r−1}); then deg(c_1(L)c_1(H)^{r−1})=0 and deg(c_1(L)^2c_1(H)^{r−2})=h^2deg(c_1(M)^2c_1(H)^{r−2})−hd^2. The established7.4(i), expressly invoked by its proof, gives the same strict conclusion from [L]≠0. Replace the inadequate word equivalent by proportional in current only, restoring the hypothesis actually used by both printed proofs; no additional theorem is introduced.

[648] EN-CONTRIB-XIII-048 Source folio 664. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=678
Printed reading: $L = M^{\otimes n} \otimes H^{\otimes -d}$
Corrected reading: $L = M^{\otimes h} \otimes H^{\otimes -d}$
Source: fixes/MISPRINTS_6.json#634. Both scans show n without an ascender. With d=M·H and h=H^2, the corrected L=M^{⊗h}⊗H^{⊗−d} satisfies L·H=hd−dh=0, exactly the equality used next. Retain the existing current h.

[649] EN-CONTRIB-XIII-049 Source folio 665. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=679
Printed reading: not numerically equivalent to
Corrected reading: not numerically proportional to
Source: ASTRA-XIII-0065. Both statements print non-equivalence to H, but their strict conclusions fail for X=$\mathbb{P}^2$, H=O(1), M=O(2): M and H have different degrees on a line, whereas h=H^2=1, d=M·H=2 and M^2=4=d^2/h. The same example lies within7.4 with r=2. The proof uses the homomorphism Pic(X)→A^1_num(X)_Q induced by c_1: for L=M^{⊗h}⊗H^{⊗−d}, [L]=h[M]−d[H] and L·H=0. Numerical nonproportionality makes [L] nonzero, so the established7.1 gives L^2<0. Expanding L^2=h^2M^2−hd^2 yields precisely M^2<d^2/h. For7.4 put h=deg(c_1(H)^r), d=deg(c_1(M)c_1(H)^{r−1}); then deg(c_1(L)c_1(H)^{r−1})=0 and deg(c_1(L)^2c_1(H)^{r−2})=h^2deg(c_1(M)^2c_1(H)^{r−2})−hd^2. The established7.4(i), expressly invoked by its proof, gives the same strict conclusion from [L]≠0. Replace the inadequate word equivalent by proportional in current only, restoring the hypothesis actually used by both printed proofs; no additional theorem is introduced.

[650] EN-CONTRIB-XIII-050 Source folio 666. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=680
Printed reading: is integral,
Corrected reading: is integral,
Source: fixes/MISPRINTS_6.json#671. The isolated line begins beneath è and ends beneath r; int and the final e are outside it. Retain this partial stroke diplomatically and the current unlined word, since it identifies no coherent emphasized term.

[651] EN-CONTRIB-XIII-051 Source folio 666. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=680
Printed reading: Pathologies III, Amer. J. Math 1967 p. 94-103 Amer. J. Math 1967 p.94-103.
Corrected reading: Pathologies III, Amer. J. Math 1967 p. 94-104.
Source: fixes/MISPRINTS_6.json#673; ASTRA-XIII-0067. fixes/MISPRINTS_6.json#673: The journal, year and page range are repeated on the next line. Retain the diplomatic dittography and current single reference; the incorrect terminal page is separately checked against the journal's primary table of contents. ASTRA-XIII-0067: The source prints103 in both copies. The American Journal of Mathematics89(1),1967 table of contents identifies D. Mumford, Pathologies III as pp.94–104; its following article begins105. The primary journal metadata is archived in review/xiii/evidence/bibliographic_verification.json. Correct the terminal page in the current single reference only.

[652] EN-CONTRIB-XIII-052 Source folio 666. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=680
Printed reading: Grothendieck A. . and Dieudonné J.
Corrected reading: Grothendieck A. and Dieudonné J.
Source: fixes/FIXLOG_6_CURRENT.jsonl#556. Two separate full stops are visible after the author initial A. The second repeats the abbreviation stop before the conjunction et; preserve the double stop diplomatically and the existing single stop in current.

[653] EN-CONTRIB-XIII-053 Source folio 666. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=680
Printed reading: Grothendieck A. . et alii
Corrected reading: Grothendieck A. et alii
Source: fixes/FIXLOG_6_CURRENT.jsonl#557. The source again has an extra separate stop following A. Preserve that printed duplication diplomatically and the existing single stop in current.

[654] EN-CONTRIB-XIII-054 Source folio 666. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=680
Printed reading: J.H.E.S.
Corrected reading: I.H.E.S.
Source: fixes/MISPRINTS_6.json#674. The first letter has the hooked lower part of the nearby J in Jussila. Retain diplomatic J and the current institution initial I.

[655] EN-CONTRIB-XIV-001 Source folio 667. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=681
Printed reading: be a -
Corrected reading: be a λ-
Source: fixes/MISPRINTS_6.json#676. Main and witness678 show an empty space before the hyphen. This is the homomorphism of the two lambda-ring structures just described; keep the supplied lambda in corrected forms and the blank diplomatically.

[656] EN-CONTRIB-XIV-002 Source folio 667. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=681
Printed reading: homomorphism For this purpose
Corrected reading: homomorphism. For this purpose
Source: fixes/MISPRINTS_6.json#677. Both images omit the sentence stop. Keep the current supplied stop and diplomatic omission.

[657] EN-CONTRIB-XIV-003 Source folio 667. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=681
Printed reading: (generally nonadditive of “exterior power)
Corrected reading: (generally nonadditive) of “exterior power”
Source: fixes/MISPRINTS_6.json#679. The print omits the closing quotation mark and places the parenthesis after extérieure. In corrected French the parenthetical qualifies the functors as nonadditive; close it after général, then complete the phrase de puissance extérieure with its quotation mark.

[658] EN-CONTRIB-XIV-004 Source folio 669. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=683
Printed reading: have their groups proper
Corrected reading: have their supports proper
Source: fixes/MISPRINTS_6.json#689. Both images print groupes. The properness condition on a cohomology sheaf is the properness of the restricted morphism Supp H^i(L)->Y. The current supports expresses this condition and is retained.

[659] EN-CONTRIB-XIV-005 Source folio 669. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=683
Printed reading: $f(y'y''c_d(\underline{N})$
Corrected reading: $f_*(y'y''c_d(\underline{N}))$
Source: fixes/MISPRINTS_6.json#685. The asterisk position is empty in both main and witness680. The right member is the pushforward f_*(y′y″c_d(N)) from Gr(Y) to Gr(X), matching the two pushforwards on the left. Keep the supplied star in corrected forms.

[660] EN-CONTRIB-XIV-006 Source folio 670. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=684
Printed reading: $(c_\beta(\underline{N}))_\beta)),,$
Corrected reading: $(c_\beta(\underline{N}))_\beta)),$
Source: fixes/MISPRINTS_6.json#690. Main and witness681 show two adjacent comma marks. Keep both diplomatically and the single comma in current.

[661] EN-CONTRIB-XIV-007 Source folio 670. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=684
Printed reading: $f : \mathrm{Gr}^\bullet(Y)_\mathbb{Q} \to \mathrm{Gr}^\bullet(X)_\mathbb{Q}$
Corrected reading: $f_* : \mathrm{Gr}^\bullet(Y)_\mathbb{Q} \to \mathrm{Gr}^\bullet(X)_\mathbb{Q}$
Source: fixes/MISPRINTS_6.json#691. Both images show f without its star. The rational homomorphism is the extension of the pushforward denoted f_* immediately before, so retain the same pushforward label by adding the lower star in current only.

[662] EN-CONTRIB-XIV-008 Source folio 670. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=684
Printed reading: $K^\bullet(X) \xrightarrow{\sim} K(X)$
Corrected reading: $K^\bullet(X) \xrightarrow{\sim} K_\bullet(X)$
Source: ASTRA-LATE-0087. Both scans have no dot after the right K. The next two lines explicitly identify its topological filtration as that of K_·(X), the coherent-sheaf group. The isomorphism takes a vector-bundle class to its coherent-sheaf class. Preserve the blank diplomatically and retain the identifying lower dot in current.

[663] EN-CONTRIB-XIV-009 Source folio 672. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=686
Printed reading: over a field k of characteristic zero k, when
Corrected reading: over a field k of characteristic zero, when
Source: fixes/MISPRINTS_6.json#644. The second k is visibly printed. It repeats the same field name after the characteristic adjective; delete only this duplicate in the corrected forms.

[664] EN-CONTRIB-XIV-010 Source folio 673. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=687
Printed reading: one or another of its rings
Corrected reading: one or another of these rings
Source: fixes/MISPRINTS_6.json#649. The scan prints ses. The antecedent is the three rings just enumerated; the demonstrative ces supplies that reference. Correct only corrected forms.

[665] EN-CONTRIB-XIV-011 Source folio 674. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=688
Printed reading: (Cf.. no. 5)
Corrected reading: (Cf. no. 5)
Source: fixes/MISPRINTS_6.json#651. Two successive full stops are visible after Cf. Remove one duplicate point in corrected forms.

[666] EN-CONTRIB-XIV-012 Source folio 675. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=689
Printed reading: the class of Y in $A^i(Y)$
Corrected reading: the class of Y in $A^i(X)$
Source: fixes/MISPRINTS_6.json#652. The print visibly gives Y. The displayed Chern class is on X and the next sentence explicitly uses A^i(X); corrected X restores the same ambient cycle group.

[667] EN-CONTRIB-XIV-013 Source folio 675. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=689
Printed reading: $A(X) \xrightarrow{f_*} A(X \otimes_k k')$
Corrected reading: $A(X) \xrightarrow{f^*} A(X \otimes_k k')$
Source: fixes/MISPRINTS_6.json#653. The star lies below f. The map in this direction is base change f^*, while the next map is its trace f_* and their displayed composite is f_* f^*. Preserve the printed subscript diplomatically; use the superscript in corrected text.

[668] EN-CONTRIB-XIV-014 Source folio 676. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=690
Printed reading: $\deg(Y \cdot \xi^i) > 0$
Corrected reading: $\deg(Y \cdot \xi^{\dim Y}) > 0$
Source: fixes/MISPRINTS_6.json#714. The printed exponent is i. The degree map is A^{dim X}(X) -> Z, while [Y] has codimension i and ξ has degree 1. Thus [Y]ξ^m has degree i+m and the zero-cycle degree uses m=dim X-i=dim Y. Change the exponent to dim Y in corrected forms.

[669] EN-CONTRIB-XIV-015 Source folio 676. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=690
Printed reading: $f : Y \qquad X$
Corrected reading: $f : Y \to X$
Source: fixes/MISPRINTS_6.json#715. Both main PDF page690 and witness page687 show completely blank space between Y and X. It is an omitted arrow in the declared immersion; preserve the blank diplomatically and supply the arrow in corrected forms.

[670] EN-CONTRIB-XIV-016 Source folio 676. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=690
Printed reading: a formula valid in $\mathrm{Gr}^{\bullet}_{\mathrm{top}}(X)$
Corrected reading: a formula valid in $\mathrm{Gr}^{\bullet}_{\mathrm{top}}(Y)$
Source: fixes/MISPRINTS_6.json#716. The print says X. The maps are f_*:Gr_top(Y)->Gr_top(X) followed by f^*:Gr_top(X)->Gr_top(Y); both f^*f_*(y) and y c_d(N) therefore lie on Y. Correct the ambient argument to Y.

[671] EN-CONTRIB-XIV-017 Source folio 679. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=693
Printed reading: $\mathrm{Filt}^{i+1}_{\mathrm{top}}(X) = \mathrm{Filt}^{i+1}_{\mathrm{top}}(X)$
Corrected reading: $\mathrm{Filt}^{i+1}_{\mathrm{top}}(X) = \mathrm{Filt}^{i+1}(X)$
Source: fixes/MISPRINTS_6.json#721. The source visibly repeats the same top filtration. The preceding sentence supplies Filt^{i+1}(X)=Filt_top^{i+1}(X); the correction writes precisely that established equality in the kernel argument. Remove the repeated top index in corrected forms, retaining both printed occurrences diplomatically.

[672] EN-CONTRIB-XIV-018 Source folio 680. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=694
Printed reading: $A^{+}(G) = \coprod_{i \geq 0} A^{i}(G)$
Corrected reading: $A^{+}(G) = \coprod_{i > 0} A^{i}(G)$
Source: fixes/MISPRINTS_6.json#724. The witness crop of page691 shows the extra slanted lower stroke of $\geq$. Including i=0 supplies the direct summand A^0(G)=Z, with inclusion and projection splitting the sum; this contradicts the printed description groupe fini. Correct the index to i>0, keeping the printed coproduct symbol.

[673] EN-CONTRIB-XIV-019 Source folio 681. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=695
Printed reading: is contained in that of $H^{2}(X, \mathbb{Z}_{\ell}(*))$
Corrected reading: is contained in that of $H^{2*}(X, \mathbb{Z}_{\ell}(*))$
Source: fixes/MISPRINTS_6.json#693. There is no star after 2. The map g_Q is graded and its degree-i component lands in H^{2i}(X,Q_ell(i)); the integral image referred to here is therefore the direct sum of those groups, represented by H^{2*}. Keep the current added degree star, with the printed asterisk shape.

[674] EN-CONTRIB-XIV-020 Source folio 682. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=696
Printed reading: in the absence of g^{i} itself
Corrected reading: in the absence of g^{i} itself.
Source: fixes/MISPRINTS_6.json#697. The line ends immediately after the final e; the next paragraph starts 5.4. Keep the diplomatic omission and the current supplied sentence stop.

[675] EN-CONTRIB-XIV-021 Source folio 683. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=697
Printed reading: $H^{n}(X, \underline{Q}(i))$
Corrected reading: $H^{n}(X, \underline{Q}_{\ell}(i))$
Source: fixes/MISPRINTS_6.json#698. The main scan and witness page694 show the underline and a blank with no ell. Supply ell only in current forms; preserve the underline in both.

[676] EN-CONTRIB-XIV-022 Source folio 683. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=697
Printed reading: $H^{n-2d}(Y, \underline{Q}(i-d))$
Corrected reading: $H^{n-2d}(Y, \underline{Q_\ell}(i-d))$
Source: fixes/MISPRINTS_6.json#699. The second coefficient has the same empty ell position in both images. Supply ell only in current forms and preserve the underline.

[677] EN-CONTRIB-XIV-023 Source folio 683. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=697
Printed reading: the virtual relative dimension of d
Corrected reading: the virtual relative dimension of f
Source: fixes/MISPRINTS_6.json#700. The final printed letter is d. The integer d is the virtual relative dimension of the morphism f:X->Y, whose Gysin map lowers cohomological degree by 2d. Replace this second d with f in corrected forms.

[678] EN-CONTRIB-XIV-024 Source folio 687. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=701
Printed reading: rationelle
Corrected reading: rationnelle
Source: fixes/MISPRINTS_6.json#660. One n is printed. Restore the second n only in current forms.

[679] EN-CONTRIB-XIV-025 Source folio 688. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=702
Printed reading: Algébre Locale
Corrected reading: Algèbre Locale
Source: fixes/MISPRINTS_6.json#661. The e has an acute accent in both images. Correct French uses Algèbre.

[680] EN-CONTRIB-XIV-026 Source folio 688. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=702
Printed reading: (SGA 6)
Corrected reading: (SGA 4)
Source: fixes/MISPRINTS_6.json#662. The printed numeral is 6. The named topos and étale-cohomology seminar is identified as SGA 4 in the surrounding citations (including XIV 6.2); keep the current bibliographical numeral correction to 4.

[681] EN-CONTRIB-XIV-027 Source folio 688. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=702
Printed reading: U. Mausin
Corrected reading: U. Manin
Source: fixes/MISPRINTS_6.json#663. The main image and witness page699 clearly print U. Mausin. The following page names MANIN in its citation 17. Correct the surname to Manin; preserve the printed initial U. The supplied replacement Yu. I. expands initials absent from the source.

[682] EN-CONTRIB-XIV-028 Source folio 688. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=702
Printed reading: MAt.
Corrected reading: Mat.
Source: fixes/MISPRINTS_6.json#664. The second letter is an uppercase A. Keep MAt. diplomatically and Mat. in corrected forms.

[683] EN-CONTRIB-XIV-029 Source folio 688. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=702
Printed reading: Sborik
Corrected reading: Sbornik
Source: check/findings/findings_686_690.json#47. The printed word lacks n. The corrected journal name Sbornik is retained.

[684] EN-CONTRIB-XIV-030 Source folio 689. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=703
Printed reading: a proof of (9.6)
Corrected reading: a proof of (4.6)
Source: ASTRA-LATE-0070. The call on p.677 is attached to the assertion about formula (4.6) in A(Y). This note reports that same Chow-ring result, whereas Exposé XIV has no section 9 or formula (9.6). Correct the reference to (4.6).

[685] EN-CONTRIB-XIV-031 Source folio 689. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=703
Printed reading: without a quasi-projective hyperplane
Corrected reading: without a quasi-projectivity hypothesis
Source: fixes/MISPRINTS_6.json#668. The wording hyperplan quasi-projectif is printed in both images. The parenthetical removes the quasi-projectivity restriction on the schemes and adds the relative smooth setting; keep the current correction hypothèse quasi-projective.

[686] EN-CONTRIB-INDEX-001 Source folio 691. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=705
Printed reading: of $\underline{C}_0$-perfect amplitude contained in $[a, b]$ complex)
Corrected reading: of $\underline{C}_0$-perfect amplitude contained in $[a, b]$ (complex)
Source: fixes/MISPRINTS_6.json#635. The opening parenthesis is absent although the closing one is printed. Keep this omission diplomatically and the supplied opening parenthesis in corrected text.

[687] EN-CONTRIB-INDEX-002 Source folio 691. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=705
Printed reading: of $\underline{C}_0$ perfect amplitude contained in $[a, b]$ complex)
Corrected reading: of $\underline{C}_0$ perfect amplitude contained in $[a, b]$ (complex)
Source: fixes/MISPRINTS_6.json#636. The second entry has the same missing opening parenthesis. Keep the diplomatic omission and the current supplied parenthesis.

[688] EN-CONTRIB-INDEX-003 Source folio 696. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=710
Printed reading: $C^P_{\mathrm{num}}(X)$
Corrected reading: $C^p_{\mathrm{num}}(X)$
Source: fixes/MISPRINTS_6.json#710. The index types a capital P. The cited definition XIII 5.2 on p.651/PDF665, confirmed in witness662, uses handwritten lowercase p and describes codimension p cycles. Correct this index entry to the explicitly defined C^p_num(X); both denote the quotient of codimension-p cycles by numerical equivalence, with identity on cycle classes. Retain the printed capital diplomatically.

[689] EN-CONTRIB-INDEX-004 Source folio 696. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=710
Printed reading: $\mathrm{D}({}_A\underline{S}_{\mathrm{coh}}$,
Corrected reading: $\mathrm{D}({}_A\underline{S})_{\mathrm{coh}}$,
Source: fixes/MISPRINTS_6.json#709. The first object lacks the closing parenthesis; the second has one. Preserve the omission diplomatically and keep the supplied parenthesis in current.

[690] EN-CONTRIB-INDEX-005 Source folio 699. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=713
Printed reading: $\psi_k(x)$
Corrected reading: $\psi^k(x)$
Source: fixes/MISPRINTS_6.json#713. The k is below the baseline. The cited definition on p.360/PDF374 explicitly writes psi^k(x), the coefficient of (-1)^{k-1}t^{k-1} in the logarithmic derivative of lambda_t(x). The operation recorded here is that same coefficient map K->K. Use its defined superscript k in corrected forms.

[691] EN-CONTRIB-INDEX-006 Source folio 699. Scan/witness: scan/SGA6_scan_360dpi.pdf#page=713
Printed reading: 3,4
Corrected reading: 3.4
Source: fixes/MISPRINTS_6.json#712. The section reference is printed with a comma between 3 and 4. The item identifies the numbered section V 3.4; keep that supplied reference separator in corrected text and the comma diplomatically.