

Heat transport: multiplication, flags and actual evaluation

$$\mathcal{E} = \{ \sum a_n z^n : \sum |a_n| \sqrt{n!} \, h^n < \infty \text{ for every } h > 0 \}$$

$$T_t = e^{-t\partial_z^2}, \quad f \star_t g = T_t((T_{-t}f)(T_{-t}g))$$

$$f \star_t g = \sum_{n \geq 0} \frac{(-2t)^n}{n!} f^{(n)} g^{(n)}, \quad z \star_t z = z^2 - 2t$$

The complete coefficient algebra keeps two descending support chains:

$$\nu_0 \geq \nu_2 \geq \cdots, \quad \nu_1 \geq \nu_3 \geq \cdots, \quad a_n \neq 0 \Rightarrow \nu_n = 1_L$$

Multiplication includes the mixed odd-odd contribution

$$(\nu \otimes \mu)_0 = (\nu_0 \wedge \mu_0) \vee (\nu_1 \wedge \mu_1)$$

$$(\nu \otimes \mu)_n = \bigvee_{r=0}^n \nu_r \wedge \mu_{n-r} \quad (n \geq 1)$$

$$\Phi(f, \nu) = (f, (1_K \otimes \nu_n)_n) \in \mathcal{E} \times \mathcal{F}(K_L)$$

The exact character identifies which divisor is transported

$$\chi_{t,z}(f) = (T_{-t}f)(z), \quad \chi_{t,z}(H_a) = H_{a-t}(z)$$

$$\chi_{t,z}(H_t) = H_0(z), \quad H_t(z) = \chi_{t,z}(H_{2t})$$

Every entire function, time parameter and support label remains in these maps.

Proofs: HTP1-24, SHP1-28, HDT1-6. All series converge in the displayed coefficient space.

Kernel: Brad Rodgers & Terence Tao, arXiv:1801.05914v5. Full carrier: The Clankers, v11, def:lattice-split.