

Supported time-zero is a projection

$e = (0, 1_L), \quad \tau = (0, 0_L), \quad \Phi(c, \lambda) = (c, 1_K \otimes \lambda)$

Exact finite example: the polynomial z^4

$q(U_{(t, 1_L)} z^4) = z^4 - 12tz^2 + 12t^2$



The full map retains the changed support matrix

$U_{(1, 1_L)} U_{(-1, 1_L)} = U_e \neq U_\tau = I_5$

$\Phi(U_e) = (I_5, B_{1_L})$

Fixed-image labels keep the two parity chains:

$\nu_0 \geq \nu_2 \geq \nu_4, \quad \nu_1 \geq \nu_3$

B_{1_L}				
1	0	1	0	1
0	1	0	1	0
0	0	1	0	1
0	0	0	1	0
0	0	0	0	1

$1 = 1_{K_L}, \quad 0 = 0_{K_L}$

The actual theta family already lies in this fixed image

$\mathbf{a}(t) = ((h_0(t), 1_L), \tau, (h_1(t), 1_L), \tau, (h_2(t), 1_L), \dots)$

$(-1)^j h_j(t) > 0 \quad (t \in \mathbb{R}), \quad U_{(0, \lambda)} \mathbf{a}(t) = \mathbf{a}(t) \quad (\lambda \in L)$

Every label is retained. The kernel positivity and convergent heat action prove this identity.

Here q is the amplitude map. Proofs: THS3, THS9, THS16–25; full independent-label operator: HSL1–14.

Original heat kernel: Brad Rodgers & Terence Tao, arXiv:1801.05914v5. Full carrier: The Clankers, v11, def:lattice-split.