

Exact arithmetic receivers: endpoint jets and the full primary block

$$\mathcal{M}f_b(s) = 2b^s \zeta(s) \left[\frac{\pi^2 \Gamma((s+4)/2)}{2\pi^{(s+4)/2}} - \frac{3\pi \Gamma((s+2)/2)}{4\pi^{(s+2)/2}} \right]$$

Endpoints $s = 0$ and $s = 1$

$$\mathcal{M}f_b(0) = 1/4, \quad \mathcal{M}f_b(1) = b/4$$

Finite combinations at distinct widths adjust every finite endpoint jet.
The actual restriction-image class and its numerical trace stay fixed.

At an original nontrivial zero ρ of multiplicity m

$$\mathcal{O}/h^{\min(m, N)}\mathcal{O}, \quad h = s - \rho, \quad N[v] = [hv]$$

$$[1] \mapsto [h] \mapsto [h^2] \mapsto 0 \quad (m = N = 3)$$

$$(N^\uparrow)^3(v, \lambda) = (0, \lambda) = e \cdot (v, \lambda)$$

WJ17-18 prove the general block and all scaling factors. The three-step chain is the explicitly stated $m = N = 3$ example.
Trivial-zero and pole contributions require the retained original multiplier and embedded lattices; no divisor is discarded.