

Original Mixed Cohomology Periods, Kernels and Arithmetic Control

Split-Zero research programme

15 September 2026

Complete proof edition. The chapters below retain the original periods, amplitude units, primary multiplicities, cyclic maps, source masses and four endpoint metrics. Each chapter supplies its full calculation and proof. Earlier finite bounds remain recorded in their original form; subsequent chapters give their proved refinements. In particular KAF retains the factorial denominator of OKV and improves the absolute kernel bound, while AMT evaluates the source differences left in OMV.

Sources and scope. The complete preceding Gamma and observation proofs, including their historical analytic providers, are supplied in the source closure. The original all-multiplicity cyclic maps remain in OCS. The evaluated five-member quadric orbit, exact kernel dimension, and its individual volume conclusions state their simple-quartet period domain explicitly. The full all-multiplicity operator estimate is given in KAF. OCF, OKM and OKA calculate the actual kernel frame, its original metrics and the boundary action. OMG proves the full all-multiplicity Segre–Veronese geometry and retains the actual cyclic stabilizer.

Current refinement. AKS sharpens the complete kernel bound to $32 \log(25 \log 3 / (4\pi))$ at the kq scale. AKJ evaluates the original residue inverse and conductor degree flag. AKP proves the precise theta representative-change and minimum-section maps. RUI now proves all-rank exterior control of the original finite conductor, and CEP carries it through the exact source ideals, four endpoint quotients and original compressed covariances. The resulting conductor error is $o(kq)$ for each fixed actual period. CTV now gives the exact outer-root quotient and its controlled norm tails. ROQ calculates the actual residual observation, all its positive increments and full graph metric. OSP proves a full-norm comparison with explicit moments of the original Gamma sources. OAR and PMR carry these finite errors into the arithmetic return. OFV now evaluates the outer contribution at the kq scale, using the full EIQ equilibrium and QLG finite partition estimate supplied below. FMC constructs fixed original coefficient frames with their exact Gamma and polynomial-degree maps. OVR propagates the evaluated outer term into the original kernel, residual observation, arithmetic boundary and signed mixed return. Its remaining analytic quantity is the original residual Gamma compression. OOQ controls every direction of the complete original outer quotient. GIP gives the full residual source and relation graph, with the exact kernel–cokernel sequence of its outer-coordinate map. PCL evaluates the original conductor limit and proves its pivot restriction. ROG straightens the complete low-degree relation graph with a proved subleading exterior cost. RDS supplies the exact dual differential realization and original Gamma coefficient map of that relation quotient. FEX gives the complete fixed-period exterior transfer; FGC proves its exact comparison with the original graph, retaining both low-degree metric corrections and the actual residual observation. ECL evaluates the original equilibrium coefficient with its explicit certified interval. EQR carries that calculation into the original earlier centre estimate while preserving its full error radius. RWC proves strict original full-window residual growth and every exterior coefficient. OER propagates the complete fixed-column and window bounds into the earlier kernel, boundary, signed and same-class control equations. MCE computes

the same arithmetic class's action control, including every kernel, boundary and mixed exterior term. MCP gives the exact marked-pairing action and source-derivative identities. RFX proves the full arithmetic and metric reflection, evaluates the total phase at the original source, and retains the individual mixed pairings. OMR carries these calculations into the earlier receiving equations and the complete arithmetic control-radius budget. RCF calculates the complete common factor of the actual polynomial relations and its exact quotient sequence. RCM carries the resulting quotient through both original Gamma source orders and all four windows, retaining the actual residual subspace and its metric. DNE realizes that weighted relation metric by its minimum dual extension with the complete numerator, source coefficients and masses. PEC controls its positive exterior expansion and transports the exact negative spectral deficits into the original kernel, residual and arithmetic action bounds. Every remaining term is kept in its original space with the full coordinate maps. ACR gives finite formulas for the actual common-factor map on the original kernel, including every coefficient-support case. LBRE proves the complete boundary exclusions at the original period limit, and ASR gives an explicit support radius and finite exceptional-period set. Their zero common-factor conclusion retains the entire complementary residual space and its metric; all exceptional cases keep their exact maps. PBR composes these full calculations through the original kernel, residual, low-image and arithmetic maps, retaining both spectral deficits and the original centred finite-radius estimate. LJR proves the exact low-jet quotient Jacobian and the leading term of both low-window deficits, with the complete finite remainder. LRR installs that result into the same four-window signed return and arithmetic action calculation. HJR proves the compatible later-jet graph, both native conditioning denominators and their exact positive row increments. It transports the original finite total bound and cancels the common low-window term in the signed deficits. HCR substitutes the two complete row sums in the original mixed-return interval and action budget, retaining every finite flag, low-image and arithmetic error in that same calculation. All preceding chapters remain in full, with their proved refinements.

Contents

1	The literal cyclic sectors and the original mixed kernel	10
1.1	Original data and the Fourier basis of the literal matrix	10
1.2	The ordered-to-symmetric map and the exact sector projector	11
1.3	Every primary period coefficient of the projected curve	11
1.4	The actual finite observation matrix and its complete kernel	12
1.5	Exact invariant dimensions and unconditional forced losses	13
1.6	The actual simple-quartet quadric and a stronger rank bound	15
1.7	A second exact finite matrix for the simple-quartet kernel	16
1.8	All primary action directions remain in the mixed calculation	17
2	The original quadric orbit and an explicit invariant lifting map	18
2.1	The actual coefficient ring and the original observation	18
2.2	The complete orbit decision from the actual coefficients	19
2.3	An explicit invariant lift for the five-member orbit	20
2.4	The induced original dual map and its complete action defect	21
3	Exact original rank from the complete quadric orbit	22
3.1	Every invariant vanishing polynomial contains the full orbit	22
3.2	The exact invariant quotient and the restriction map	22
3.3	All original degrees and the exact kernel dimension	23
4	The actual period quadric and its cyclic continuation	24
4.1	Original roots, entire unit and centred phase	24
4.2	The original periods as an absolutely convergent series	24

4.3	The actual transformed quadric and every deciding coefficient	26
4.4	The cyclic action is the actual original continuation in u	27
4.5	The large-parameter geometry and a positive-ray obstruction	27
5	The original period quadric as a residue commutator	29
5.1	A flat alternating pairing with all constants	29
5.2	The full arithmetic unit and the exact quadratic form	30
5.3	An explicit original-parameter radius forcing orbit five	31
6	A finite bound for the original kernel volume from its actual dimension	34
7	The actual simple-quartet kernel and boundary at the original volume scale	39
7.1	The original domain and the proved period range	39
7.2	A uniform finite bound on the actual kernel volume	39
7.3	The two individual leading coefficients	40
7.4	Exact propagation to the arithmetic identity and the finer scale	41
8	The original arithmetic kernel and boundary receive the full source comparison	42
8.1	The complete original source and the same observation	42
8.2	All constants of the actual full-polynomial inequality	43
8.3	Exact remainder, minimum-section, kernel and boundary maps	43
8.4	The exact nested deficits and their rank factors	44
8.5	All four original endpoints and the correlated mixed return	45
8.6	The full original multiplier and its transported observation	46
8.7	The evaluated original simple-quartet arithmetic kernel	48
8.8	The original boundary receives the full baseline coefficient	49
8.9	The same comparison evaluates the mixed Gamma return on the kq scale	50
9	Factorial control of the original common kernel at the kq scale	52
9.1	Original coordinates, all primary multiplicities, and source norms	52
9.2	A degree-dependent constant in the unchanged remainder estimate	53
9.3	Retaining the factorial denominator	54
9.4	All four actual kernel and boundary Grams	54
9.5	The actual simple quartet and the arithmetic source	55
10	An exact conductor frame and a fixed invariant-generator recursion	57
10.1	The original Segre coefficient isomorphism	57
10.2	Every leading-coefficient case gives an exact strip frame	58
10.3	Twenty-five fixed invariant monomials generate all degrees	58
10.4	Exact multiplication on the strip and the invariant recursion	59
10.5	Exact ranks, a finite pivot atlas, and the original quotient map	60
10.6	The same recursion with fourteen proved generators	62
11	The exact common kernel frame and its original Gamma metrics	63
11.1	The exact dual and primal maps of the strip quotient	63
11.2	The complete original source and quotient metric	64
11.3	Christoffel–Darboux with the exact source mass	64
11.4	Every pivot change and the four original endpoints	65
12	The original kernel graph, boundary basis and complete action	67
12.1	The coefficient quotient gives a graph in the original primary basis	67
12.2	An explicit basis of the actual boundary	67
12.3	The whole original action and its retained defect	68

13 The complete homogeneous geometry of the original multiplicity curve	70
13.1 The actual primary, unit, period and Fourier coefficient map	70
13.2 The complete homogeneous ideal, with quadratic generators	71
13.3 The projective model and its complete chart inverses	72
13.4 The exact graded source-dual algebra and every factorial	73
13.5 The full semisimple and nilpotent action in this geometry	74
13.6 The exact original cyclic observation as invariant restriction	75
13.7 The actual orbit ideal, including every stabilizer	75
13.8 The complete component conductor and an explicit finite witness	76
13.9 The exact action defect and all subsequently observed directions	77
14 The absolute common kernel: the new finite result	79
15 An exact coefficient realization of the actual lost directions	80
15.1 The period matrix and invariant restriction, with their full coefficients	80
15.2 A determinant formula requiring no assumed nonzero pivot	81
16 A bounded exact remainder operator on the original source	82
16.1 The complete source norms and coefficient map	82
16.2 Newton interpolation with all roots and their orders	83
16.3 Cauchy extraction in the actual Gamma generating function	83
17 The four original kernel Grams and the evaluated scale	84
18 The complete arithmetic and mixed receivers	86
18.1 The unchanged finite arithmetic allowances	86
18.2 The original Gamma multiplier, not a replacement higher source	87
18.3 Boundary, QGQ and arithmetic receiving identities	87
19 Actions, support faces, and the actual arithmetic class	89
20 Exact period domain and the interior alternative	89
21 Verification scope and the remaining absolute coefficient	90
22 The new ceiling in the original coefficient frame	91
23 Residue detection expressed in every original primary coefficient	92
24 The actual conductor confines the kernel's polynomial degrees	93
25 The exact conductor translations in the original source norm	94
26 The original theta class and the checked metric interface	96
26.1 The complete minimum-section calculation at each endpoint	96
27 The finite conductor inverse in the two original Gamma norms	98
27.1 A coefficient-controlled disk for the literal finite symbol	98
27.2 The exact coefficient map and its norm	99
27.3 The infinite symbol differs by an exact low-degree map	100
27.4 An explicit right inverse of the original conductor	100
27.5 The exact quotient and exterior transfer supplied by this bound	101

28 Subleading exterior control from the actual finite conductor	103
28.1 The forward polynomial bound and exact determinant	103
28.2 Complementary singular values retain the full exterior norm	104
28.3 The actual conductor quotient inherits the sharper volume control	104
29 The original conductor at every source endpoint	106
29.1 The full source, remainder and primary-coordinate square	106
29.2 The literal finite inverse at the larger cutoffs	107
29.3 The exact source operator matrix and its full determinant	108
29.4 The induced original quotient metric and its correction	109
29.5 An exact graph quotient with controlled exterior metric error	110
29.6 All ideal graph entries as original polynomial products	113
29.7 The compressed conductor correction is subleading at every endpoint	114
30 The total-source difference as an outer-strip quotient	116
30.1 The exact nested root polynomials in the original real coordinate	116
30.2 The full monic source and relation determinant identities	116
30.3 The exact quotient of the outer-strip polynomial	117
30.4 Finite bounds for all norm tails in both original sources	118
30.5 The proved first-variation reduction and its remaining matrix	119
31 The actual observation remaining inside the conductor kernel	121
31.1 An explicit residual map and its actual boundary basis	121
31.2 All four induced Gamma metrics and their fixed determinant factor	122
31.3 The exact constrained covariance in the original source	123
31.4 Exact increments and a positive residual four-endpoint return	124
31.5 Finite scalar enclosures and propagation into CEP49	125
31.6 The complete residual quotient in the conductor graph	125
32 A quantitative power comparison for the actual outer-strip quotient	127
32.1 The two original source masses and a proved interval lower bound	127
32.2 A complete coefficient-space estimate for the inner interval	128
32.3 The exact original-root map and a finite full-norm comparison	129
32.4 Uniform bounded logarithmic cost at every required dimension	130
32.5 Every relation determinant and the exact outer quotient	131
32.6 The resulting same-source moment target	132
33 The outer-root and actual residual bounds in the original arithmetic return	133
33.1 The two finite error endpoints retain their signs	133
33.2 Original positive increments give explicit kernel caps	133
33.3 The correlated arithmetic kernel and boundary	134
33.4 The signed mixed return with its complete finite error	134
34 The same-source Gamma moment comparison in the arithmetic return	135
35 Exact equilibrium for the square-root logarithmic potential	137
35.1 Existence and uniqueness of the two endpoints	137
35.2 A positive density and its exact Cauchy transform	138
35.3 The Euler equality, its strict exterior inequality, and uniqueness	139
35.4 An explicit endpoint integral for the logarithmic moment	140
35.5 Parameter continuity and the variational derivative	141
36 A uniform finite partition estimate from the original equilibrium	143

37 The leading first variation of the actual outer-strip quotient	148
37.1 Exact original-order recurrence and a small metric error	148
37.2 A uniform finite Gamma-to-exponential determinant estimate	149
37.3 Direct low-dimension estimates, outside the compact equilibrium range	150
37.4 High-dimension partitions and their controlled first variation	150
37.5 All seven determinants and the resulting leading constant	152
38 Fixed coefficient control of the original residual module	153
38.1 A fixed free module containing the entire invariant image	153
38.2 Finite reduction data with only fixed actual denominators	154
38.3 A fixed weight bounds every homogeneous reduction path	155
38.4 The two exact maps back to the original conductor strip	156
38.5 An identity pivot block for the entire residual image	157
38.6 Exact relation to every earlier certificate frame	158
38.7 The unchanged Gamma source in the fixed degree model	159
38.8 The exact map to the original polynomial degree flag	161
39 The evaluated outer contribution in the original mixed return	163
39.1 The exact remainder and the sourcewise leading identity	163
39.2 Transfer between the two original Gamma sources	163
39.3 Arithmetic kernel and its correlated boundary	163
39.4 The signed return retains its original Gamma correction	164
39.5 Bounds on the remaining residual coefficient	164
40 All directions of the original outer quotient at the leading scale	166
40.1 Uniform equilibrium constants and a polynomial evaluation bound	166
40.2 A matching polynomial lift from actual equilibrium quantiles	167
40.3 The entire analytic remainder map	168
40.4 High source degrees: a full operator estimate	168
40.5 Low source degrees, including the extra ideal column	170
40.6 Transport back to both complete original Gamma sources	170
40.7 Relative operators and all fixed subquotients of this actual target	171
41 The complete residual map on the original conductor graph	173
41.1 The full graph denominator and its metric	173
41.2 The exact natural map to the CTV outer divisor	174
42 The original mixed return and its same-class action control	176
42.1 The original arithmetic class and all three source metrics	176
42.2 The exact rank-two control from the original recurrence	177
42.3 The same nonzero arithmetic exterior class	177
42.4 Complete kernel, boundary and mixed pieces of the control	178
42.5 Degree-step volumes determine the squared component sizes	179
42.6 The signed return gives an exact original control-radius budget	180
43 The next marked pairing as an original action and source derivative	182
44 The exact original reflection and the retained mixed phase	184
44.1 The entire arithmetic unit and the source involution	184
44.2 The quotient, primary jets and arithmetic class	184
44.3 The full metric and the total endpoint pairing	185
44.4 The original mixed blocks remain present	186
44.5 Propagation through the same arithmetic control budget	187

45 The outer operator, the full residual graph, and original action control	188
45.1 Original objects, domains and the same-target operator interval	188
45.2 The full graph denominator and its exact outer morphism	189
45.3 Backward propagation through the original kernel and signed return	190
45.4 The original action, its mixed components, and its positive class	191
45.5 The exact arithmetic radius budget and its backward bound	192
45.6 The exact marked derivative and original reflection specialization	193
46 The literal original conductor limit and an invertible pivot restriction	196
46.1 An entire coefficient matrix in the inverse original period	196
46.2 All eighty-one actual conductor coefficients and their limit	197
46.3 An exact positivity slice for the literal positive period ray	198
46.4 A complete invertible restriction supplied by the original pivot	198
46.5 The exact inverse-period reflection of the central section	199
47 The original relation operator and its finite-rank graph transport	201
47.1 The complete positive-shift relation operator	201
47.2 An explicit finite inverse with coefficient control	201
47.3 Complete original-source bounds for the low graph map	203
47.4 The exact graph shear and its complementary singular pairs	204
47.5 The unchanged residual target and its four endpoint metrics	205
48 The exact differential realization of the original relation quotient	207
48.1 A finite-dimensional quotient and its analytic dual	207
48.2 Every possible singularity and its original multiplicity	208
48.3 An explicit original-data analytic disk and source coefficients	209
49 The fixed-period problem and the result proved here	210
50 A full-flag determinant bound	211
51 Polynomial forward control replaces the exponential exterior loss	211
52 The literal conductor at every source endpoint	213
53 The residual observation and the smaller coefficient space	214
54 A finite error for the actual compressed determinant	215
55 The original arithmetic and signed receiving identities	216
56 Actions, representatives, and the retained mixed-support maps	218
57 Scope of verification and the next exact quantity	219
58 The fixed-column conductor quotient and the full GIP graph	220
58.1 The precise projection and its complete kernel	220
58.2 The actual fixed columns and the Schur metric	221
58.3 Both original low metric terms and their signed identity	222
59 The exact closed equilibrium coefficient	223
59.1 Arcsine identities with their original constants	223
59.2 Differentiation with the moving endpoints retained	224
59.3 The moment and energy formulas	226
59.4 The original outer coefficient and the two scale conventions	226

59.5	An integer-only outward certificate	227
60	The closed coefficient in the original QGQ return centre	229
61	Strict full-window growth of the original residual covariance	231
61.1	Every consecutive full block is invertible	231
61.2	The simultaneous conditioned update	231
61.3	All exterior coefficients and finite bounds	232
61.4	Return to the original four endpoints	233
62	The fixed-period determinant and closed coefficients in the original receiver	234
62.1	The existing outer coefficient has an exact endpoint expression	234
62.2	The actual period coefficients and the pivot morphism	234
62.3	The full low graph has its own finite-rank transport	235
62.4	The fixed columns arise from an exact quotient of that graph	236
62.5	Both low terms and all one-sided errors enter the current interval	236
62.6	The relation quotient and the earlier finite Gamma centre	237
62.7	Full-window residual bounds in these same receivers	238
63	The complete common factor of the original polynomial relations	240
63.1	Every original finite degree window and its exact metric sequence	241
64	Original Gamma control on the actual common-factor quotient	243
64.1	The exact original divisor, windows, phases and source	243
64.2	Complete finite bounds at the low source windows	243
64.3	The high windows with the actual analytic remainder and lift	244
64.4	Both full Gamma sources and the four-return coefficient	245
64.5	The resulting original residual quotient and its retained complement	246
65	The original weighted relation metric as a minimum dual extension	248
65.1	A fixed original exponential quotient frame	248
65.2	The exact original source extension problem	248
65.3	Complex-linear duality and the actual fixed kernel columns	249
65.4	A finite source bound before the high-order derivative	250
66	Subleading positive exterior expansion of the original dual metric	252
66.1	The actual symbol-zero jet subspaces	252
66.2	A complete finite Gamma bound on each cancellation subspace	253
66.3	The whole positive logarithmic spectrum	254
66.4	Every fixed restriction and quotient keeps this bound	254
66.5	The exact negative spectral return on the actual kernel	255
66.6	The complementary original residual and its low subspace	256
66.7	Backward and forward return to the same arithmetic class	257
67	The actual common-factor map on the original kernel	258
67.1	Support and its precise extended coefficient rectangle	258
67.2	The residual map evaluated on every original primary coordinate	259
67.3	The exact retained residual quotient in biform coordinates	260
67.4	A smaller invariant recursion evaluating every exceptional rank	261
68	Exact exclusion of all four boundary rulings at the original period limit	263
68.1	The exact even and odd evaluation maps	263
68.2	The four complete quadratic restrictions	264
68.3	A nonzero exact eliminant for every original amplitude	264

69 Actual boundary multiplicities and a finite full-support radius	266
69.1 Every finite-period boundary multiplicity	266
69.2 The exact symplectic map underlying the coefficient tests	267
69.3 An explicit radius from the actual limiting corner calculation	268
69.4 Consequence for the actual common factor and residual	269
69.5 Only finitely many support exceptions on the original exterior domain	269
70 The two exact dual subquotients in the original receiver	270
70.1 Finite positive-spectrum allowance and every small eigenvalue	270
70.2 The original metric on the low image	272
70.3 Both low terms and the full one-sided kernel errors	272
70.4 The complete current finite interval	273
70.5 The signed return and the complete same-class action budget	274
70.6 Actual support, the common-factor map, and the retained residual	275
71 Exact low-jet reconstruction and its original quotient spectrum	277
71.1 The exact finite reconstruction and its inverse	277
71.2 Both exact quotient kernels and their metrics	278
71.3 The complete full-space and quotient Jacobians	278
71.4 Finite reconstruction bounds on the actual grid	279
71.5 A direct estimate of the nested determinant difference	280
71.6 The complete remaining native Gram	281
71.7 The exact remainder and every paired exterior volume	282
72 First-endpoint reconstruction in the actual kernel and annihilator	284
73 Compatible high jets and the original positive return	286
73.1 The exact compatible high-jet graph	286
73.2 One lower-evaluation minimum over the complete window	287
73.3 The original kernel and annihilator updates	287
73.4 Exact one-row updates and their allocation	288
73.5 The four original signs and an exterior bound at scale kq	289
73.6 Exact cancellation of the low leading scale	290
74 The compatible high rows in the original receiving interval	291

1 The literal cyclic sectors and the original mixed kernel

This calculation evaluates the cyclic projector already present in the original period observation. Its coefficients retain the original period integrals, amplitude unit, and all primary nilpotents. The result includes an exact finite matrix for the actual rank, as well as unconditional rank bounds obtained from the original symmetric tensor geometry.

1.1 Original data and the Fourier basis of the literal matrix

Use the following data of OPG1–5, including the same contours and their orientations:

$$\begin{aligned} \alpha_{\epsilon,\eta} &= \frac{1}{2} + \epsilon\delta + i\eta\gamma, \quad \epsilon, \eta \in \{+1, -1\}, \quad 0 < \delta < \frac{1}{2}, \quad \gamma > 2, \\ h(z_1) &= \prod_{\epsilon,\eta} (z_1 - \alpha_{\epsilon,\eta})^m, \quad n = 4m, \quad N = n + 1, \quad E_h = \mathbb{C}[z_1]/(h), \\ k &= 4l + 1, \quad l, m \geq 1, \quad c = k/2, \quad e = 1 + k(m - 1), \quad q = e(k + 1)^2, \\ \beta_{a,b} &= c + (2a - k)\delta + i(2b - k)\gamma, \quad \chi(S) = \prod_{a,b=0}^k (S - \beta_{a,b})^e, \quad E = \mathbb{C}[S]/(\chi). \end{aligned} \tag{OCS1}$$

Here A_h and A are multiplication by z_1 and S on their respective algebras, and $Y = (A - cI)/i$. The original amplitude unit is $U = M_{j_h v_h}$, with $v_h = (2\xi)/h$ and every original $v_h(\alpha) \neq 0$. For the specified nonzero u and its branch, $\Phi'_h = h$, $\Phi_h(0) = 0$, and

$$\Gamma_j = -\ell_0 + \ell_j, \quad \arg \ell_j = \frac{\arg u + \pi + 2\pi j}{N}, \quad (\Pi[P])_j = \int_{\Gamma_j} e^{\Phi_h(z_1)/u} \text{rem}_h P(z_1) dz_1. \tag{OCS2}$$

The representative has degree less than n . The convergence and invertibility of this original period map are the complete period theorems retained in OPG4. No period value or unit derivative is chosen in the present calculation. The original injection and observation are

$$\begin{aligned} \iota[P] &= [P(z_1 + \cdots + z_k)], \quad \mathcal{V} = \Pi^{\otimes k} U^{\otimes k} \iota, \\ P_k &= \frac{1}{N} \sum_{a=0}^{N-1} (C^{-a})^{\otimes k}, \\ L &= P_k \mathcal{V} = j\Lambda, \quad B = \text{im } L, \quad K = \ker \Lambda. \end{aligned} \tag{OCS3}$$

The map $j : B \hookrightarrow (\mathbb{C}^n)^{\otimes k}$ is the original image inclusion. The injection ι has precisely the defining polynomial in OCS1: on each ordered primary tensor the sum of its k independent nilpotents has index $e = 1 + k(m - 1)$, its top coefficient being $(k(m - 1))!/((m - 1)!)^k \neq 0$, and its distinct eigenvalues are the displayed $\beta_{a,b}$. Thus $\mathcal{V}$ is injective.

On a period-coordinate column $x \in \mathbb{C}^{N-1}$ the literal matrix in OPG4–5 is

$$(Cx)_j = x_{j+1} - x_1 \quad (1 \leq j < N - 1), \quad (Cx)_{N-1} = -x_1. \tag{OCS4}$$

Set $\zeta = \exp(2\pi i/N)$ and, for $1 \leq t \leq N - 1$, let

$$(f_t)_j = \zeta^{jt} - 1, \quad F = [f_1 \ \cdots \ f_{N-1}], \quad (F^{-1}x)_t = \frac{1}{N} \sum_{j=1}^{N-1} \zeta^{-tj} x_j. \tag{OCS5}$$

The last expression is the exact inverse. Indeed $\sum_{j=1}^{N-1} \zeta^{-tj} (\zeta^{uj} - 1) = N \mathbf{1}_{t=u}$ for $t, u \neq 0$ modulo N , by the finite geometric sum. The first $N - 2$ rows of OCS4 give $\zeta^{(j+1)t} - \zeta^t = \zeta^t (\zeta^{jt} - 1)$, and the last row gives $1 - \zeta^t = \zeta^t (\zeta^{(N-1)t} - 1)$. Consequently $Cf_t = \zeta^t f_t$, proving the orientation of every inverse power in OCS3, as well as $C^N = I$. The original Euclidean Gram of this basis is

$$F^* F = N(I_{N-1} + \mathbf{1}\mathbf{1}^*). \tag{OCS6}$$

To verify it, expand $\sum_j (\zeta^{-jt} - 1)(\zeta^{ju} - 1)$ and use the same geometric sums; its value is $N(\mathbf{1}_{t=u} + 1)$. Thus the complete coefficient and metric transformations are both specified.

1.2 The ordered-to-symmetric map and the exact sector projector

Let

$$\mathcal{A}_k = \{\nu \in \mathbb{Z}_{\geq 0}^{N-1} : \sum_t \nu_t = k\}, \quad \text{wt}(\nu) = \sum_{t=1}^{N-1} t\nu_t \pmod{N}, \quad \mathcal{I}_{k,\tau} = \{\nu \in \mathcal{A}_k : \text{wt}(\nu) = \tau\}. \quad (\text{OCS7})$$

For each ν define the following unscaled symmetric tensor:

$$S_\nu = \sum_{\substack{(t_1, \dots, t_k) \in \{1, \dots, N-1\}^k \\ \#\{j: t_j = t\} = \nu_t \text{ for each } t}} f_{t_1} \otimes \dots \otimes f_{t_k}. \quad (\text{OCS8})$$

Every distinct ordered word occurs once. Since the ordered tensors in the f_t form a basis, their disjoint word supports show that the S_ν are linearly independent. A permutation-invariant tensor has equal coefficients on each word orbit, so they span exactly $\text{Sym}^k(\mathbb{C}^{N-1})$, realized as that invariant subspace of the full ordered tensor product. In particular, for $x = (x_1, \dots, x_{N-1})$,

$$(Fx)^{\otimes k} = \sum_{\nu \in \mathcal{A}_k} x^\nu S_\nu, \quad (C^{-a})^{\otimes k} S_\nu = \zeta^{-a \text{wt}(\nu)} S_\nu, \quad P_k(Fx)^{\otimes k} = \sum_{\nu \in \mathcal{I}_{k,0}} x^\nu S_\nu. \quad (\text{OCS9})$$

The finite average in OCS3 is exactly one on weight zero and zero on every other weight, by its geometric sum. No multinomial coefficient is omitted: the multiplicity $k! / \prod_t \nu_t!$ is already the number of summands in OCS8.

Write $\mathcal{F}_k : \mathbb{C}^{\mathcal{A}_k} \rightarrow \text{Sym}^k(\mathbb{C}^{N-1})$ for the basis isomorphism $e_\nu \mapsto S_\nu$. Let J_0 be the coordinate inclusion of $\mathbb{C}^{\mathcal{I}_{k,0}}$ into $\mathbb{C}^{\mathcal{A}_k}$, π_0 the coordinate restriction in the opposite direction, and let $\mathcal{F}_0 = \mathcal{F}_k J_0$, with its target the original ordered tensor space. The exact projector factorization is

$$P_k \mathcal{F}_k = \mathcal{F}_0 \pi_0. \quad (\text{OCS10})$$

All images of $\mathcal{V}$ are symmetric: polynomial evaluation at a sum is invariant under every factor permutation, and the same map ΠU is applied in all factors. Therefore OCS10 applies on the entire original image $\mathcal{V}(E)$.

For completeness, the full inherited metric is retained in these coordinates. Put $G = F^* F$ as in OCS6. The Gram of the basis OCS8 is

$$\mathcal{G}_{\nu, \mu} = \sum_{\substack{a_{tu} \geq 0 \\ \sum_u a_{tu} = \nu_t, \sum_t a_{tu} = \mu_u}} \frac{k!}{\prod_{t,u} a_{tu}!} \prod_{t,u} G_{tu}^{a_{tu}}. \quad (\text{OCS11})$$

Indeed the coefficient of $\bar{x}^\nu y^\mu$ in $(\bar{x}^T G y)^k$ is the right side by the multinomial theorem. Taking the inner product of the two pure tensors in OCS9 gives the same coefficient as $\langle S_\nu, S_\mu \rangle$. This proves OCS11, including its off-diagonal entries; it is positive definite because OCS8 is a basis. On the full symmetric coordinates the projector matrix is $J_0 \pi_0$, and its adjoint for this exact Gram is $\mathcal{G}^{-1}(J_0 \pi_0)^* \mathcal{G}$. Thus the Fourier-sector computation makes no unproved orthogonality or norm assertion about the original cyclic average.

1.3 Every primary period coefficient of the projected curve

For each original root α let $E_{\alpha,a}$, $0 \leq a < m$, be the degree-less-than- $4m$ primary representatives of OPG10. Explicitly, if $h_\alpha = h/(z_1 - \alpha)^m$, put

$$E_{\alpha,a}(z_1) = \text{rem}_h \left[h_\alpha(z_1) \sum_{j=0}^{m-1} \frac{(h_\alpha^{-1})^{(j)}(\alpha)}{j!} (z_1 - \alpha)^{j+a} \right], \quad b_{j;\alpha,a} = \int_{\Gamma_j} e^{\Phi_h(z_1)/u} E_{\alpha,a}(z_1) dz_1. \quad (\text{OCS12})$$

The four primary factors are coprime. The Taylor inverse gives constant one on its own factor and zero on the other three; thus the classes $[E_{\alpha,a}]$ form the full primary basis with all m nilpotent degrees at every root. Define the exact finite Fourier and unit coefficients

$$\widehat{b}_{t;\alpha,a} = \frac{1}{N} \sum_{j=1}^{N-1} \zeta^{-tj} b_{j;\alpha,a}, \quad f_{t;\alpha,d} = \frac{i^{-d}}{d!} \sum_{a=d}^{m-1} \widehat{b}_{t;\alpha,a} \frac{v_h^{(a-d)}(\alpha)}{(a-d)!}, \quad 0 \leq d < m. \quad (\text{OCS13})$$

Let $\omega_\alpha = (\alpha - 1/2)/i$ and set

$$\mathbf{v}(w) = \Pi U \exp(w(A_h - \frac{1}{2}I)/i)[1], \quad x(w) = F^{-1}\mathbf{v}(w), \quad x_t(w) = \sum_{\alpha} e^{\omega_\alpha w} \sum_{d=0}^{m-1} f_{t;\alpha,d} w^d. \quad (\text{OCS14})$$

To verify the final equality, in the primary algebra at α multiply the Taylor series of v_h by the terminating series of $\exp(w(z_1 - \alpha)/i)$. The coefficient on $[E_{\alpha,a}]$ is $e^{\omega_\alpha w} \sum_{d=0}^a v_h^{(a-d)}(\alpha) (w/i)^d / ((a-d)!d!)$. Apply the original polynomial period map and the inverse in OCS5; this gives precisely OCS13–14. In particular, the unreduced entire function has passed through its full primary remainder before any period is integrated.

For $\nu \in \mathcal{I}_{k,0}$, $0 \leq a, b \leq k$ and $0 \leq D < e$, define $c_{\nu;a,b,D}$ by the following finite multinomial rule. Sum over integers $n_{t,\alpha,d} \geq 0$ satisfying

$$\begin{aligned} \sum_{\alpha,d} n_{t,\alpha,d} &= \nu_t \quad \text{for each } t, & \sum_{t,\alpha,d} d n_{t,\alpha,d} &= D, \\ \sum_{t,d,\alpha: \operatorname{Re} \alpha = 1/2 + \delta} n_{t,\alpha,d} &= a, & \sum_{t,d,\alpha: \operatorname{Im} \alpha = +\gamma} n_{t,\alpha,d} &= b. \end{aligned}$$

For these indices put

$$c_{\nu;a,b,D} = \sum_{\text{stated indices}} \left(\prod_{t=1}^{N-1} \frac{\nu_t!}{\prod_{\alpha,d} n_{t,\alpha,d}!} \right) \prod_{t,\alpha,d} f_{t;\alpha,d}^{n_{t,\alpha,d}}. \quad (\text{OCS15})$$

Empty index sets give zero and zero exponents give factors one. Every original primary coefficient is present in this finite rule. The ordinary multinomial expansion of OCS14 now proves

$$x(w)^\nu = \sum_{a,b=0}^k e^{\omega_{a,b} w} \sum_{D=0}^{e-1} c_{\nu;a,b,D} w^D, \quad \omega_{a,b} = (\beta_{a,b} - c)/i = (2b - k)\gamma - i(2a - k)\delta. \quad (\text{OCS16})$$

Indeed $D \leq k(m-1) = e-1$ and the counts of each original sign give exactly the displayed exponent. The sums with different (a,b) are distinct since their real and imaginary parts are independently distinct.

1.4 The actual finite observation matrix and its complete kernel

Define the primary coefficient isomorphism $\mathcal{H} : \mathbb{C}^q \rightarrow E$ by sending coordinate (a,b,D) to $\varepsilon_{a,b}(S - \beta_{a,b})^D$, where $\varepsilon_{a,b}$ is obtained from χ by the same truncated Taylor-inverse construction as OCS12, now of order e . The proof by the coprime factors is identical and shows that this is a basis of all q primary directions. Set

$$\mathfrak{A}_{\nu;(a,b,D)} = i^D D! c_{\nu;a,b,D}, \quad \mathfrak{A} : \mathbb{C}^q \longrightarrow \mathbb{C}^{\mathcal{I}_{k,0}}. \quad (\text{OCS17})$$

Then, on the original spaces and with the original ordered target,

$$\boxed{L\mathcal{H} = \mathcal{F}_0 \mathfrak{A}, \quad K = \mathcal{H}(\ker \mathfrak{A}), \quad \operatorname{rank} \Lambda = \operatorname{rank} \mathfrak{A}.} \quad (\text{OCS18})$$

Here is a proof giving every factorial and phase. In the primary sum algebra,

$$e^{wY}[1] = \sum_{a,b} e^{\omega_{a,b}w} \sum_{D=0}^{e-1} \frac{i^{-D} w^D}{D!} \varepsilon_{a,b} (S - \beta_{a,b})^D.$$

The exact original tensor identity of OPG8 gives $\mathcal{V}e^{wY}[1] = \mathbf{v}(w)^{\otimes k}$: the centre $c = k/2$ splits into the k original centres $1/2$, while U commutes with multiplication by z_1 . Applying the literal projector and then OCS9 and OCS16 gives the right-hand curve with coefficients $\mathcal{F}_0 c_{a,b,D}$. Exponential polynomials $w^D e^{\omega_{a,b}w}$ with these distinct exponents are linearly independent. To prove this directly, apply $\prod_{(a',b') \neq (a,b)} (\partial_w - \omega_{a',b'})^e$ to any zero linear combination. It kills all the other exponents. On the selected polynomial factor, each remaining operator is $\partial_w + (\omega_{a,b} - \omega_{a',b'})$, invertible on the polynomials of degree less than e because its constant term is nonzero and differentiation is nilpotent there. The surviving polynomial must therefore vanish. Doing this for every pair proves the asserted independence. Equating coefficients now gives $L\mathcal{H}e_{a,b,D} = \mathcal{F}_0(i^D D! c_{a,b,D})$. This proves the first equality in OCS18 on a basis. The other two follow from injectivity of $\mathcal{F}_0$ and j .

In particular the isomorphism from the actual sector image to the original boundary is

$$\mathcal{Q} : \text{im } \mathfrak{A} \xrightarrow{\sim} B, \quad \mathcal{Q}x = j^{-1} \mathcal{F}_0 x, \quad E/K \xrightarrow{\sim} \text{im } \mathfrak{A}, \quad [z] \mapsto \mathfrak{A} \mathcal{H}^{-1} z. \quad (\text{OCS19})$$

Both inverses exist on precisely their displayed images by OCS18. These are full coefficient maps; the inherited Gram of $\mathcal{F}_0$ is $J_0^* \mathcal{G} J_0$ from OCS11.

For a completely explicit finite rank prescription, define

$$\Delta_r(\mathfrak{A}) = \sum_{\substack{I \subseteq \mathcal{I}_{k,0}, \ J \subseteq \{(a,b,D)\} \\ |I|=|J|=r}} |\det \mathfrak{A}_{I,J}|^2, \quad \Delta_0 = 1. \quad (\text{OCS20})$$

Every entry of every minor is the finite expression OCS12–17 in the actual periods and amplitude derivatives. Gaussian elimination proves that a matrix has rank at least r exactly when one of its r by r minors is nonzero: pivot rows and columns supply such a minor, and a nonzero minor supplies r independent columns. Hence the actual rank is exactly $\max\{r : \Delta_r(\mathfrak{A}) > 0\}$. This evaluates its finite deciding data without prescribing generic values or asserting that any unexamined maximal minor is nonzero.

The Gamma vectors used in the existing endpoint increments have the equally explicit coordinates

$$j\lambda_d = \mathcal{F}_0 \hat{\lambda}_d, \quad (\hat{\lambda}_d)_\nu = \frac{1}{\sqrt{h_d^{(s)}}} \sum_{a,b=0}^k \sum_{D=0}^{e-1} c_{\nu;a,b,D} p_d^{(D)}(\omega_{a,b}), \quad h_d^{(s)} = (2\pi)^{s/2} d!(s/2)_d, \quad s \in \{1, k\}. \quad (\text{OCS21})$$

Taylor expansion of $p_d(Y)[1]$ in the sum primary basis gives the coefficient $i^{-D} p_d^{(D)}(\omega_{a,b})/D!$. Multiplication by OCS17 cancels exactly these factorials and phases, proving OCS21. This also proves that the generating formula of OPG9 becomes $(1+z^2)^{-s/4} \mathcal{F}_0(x(\arctan z)^\nu)_{\nu \in \mathcal{I}_{k,0}}$, with the original scalar exponent and both existing sources.

1.5 Exact invariant dimensions and unconditional forced losses

Put $d_{N,k,\tau} = |\mathcal{I}_{k,\tau}|$. By OCS7–10 this is the dimension of the full symmetric sector, and its exact finite character formula is

$$d_{N,k,0} = [T^k] \frac{1}{N} \sum_{h|N} \varphi(h) \frac{1-T}{(1-T^h)^{N/h}}. \quad (\text{OCS22})$$

To prove it, the trace generating series for C^a on symmetric tensors is $\prod_{t=1}^{N-1} (1 - T\zeta^{at})^{-1}$, by summing the monomial basis OCS8. If ζ^a has order h , the list ζ^{at} with $0 \leq t < N$ contains every

h th root N/h times. Their product is $\prod_{t=0}^{N-1} (1 - T\zeta^{at}) = (1 - T^h)^{N/h}$. Removing the $t = 0$ factor gives the series $(1 - T)/(1 - T^h)^{N/h}$. There are exactly $\varphi(h)$ indices a of that order, so averaging the traces of the original finite projector proves OCS22. The inverse powers give the same average by $a \mapsto -a$. In ordinary finite binomial terms, for $k \geq 1$ this is

$$d_{N,k,0} = \frac{1}{N} \sum_{h|N} \varphi(h) \left\{ \mathbf{1}_{h|k} \binom{k/h + N/h - 1}{N/h - 1} - \mathbf{1}_{h|(k-1)} \binom{(k-1)/h + N/h - 1}{N/h - 1} \right\}. \quad (\text{OCS23})$$

The binomial expansion of $(1 - T^h)^{-N/h}$ follows by counting weak compositions, proving the coefficient extraction used here. It follows from the actual matrix OCS17 that

$$\text{rank } \Lambda \leq \min(q, d_{N,k,0}), \quad \dim K \geq \max(0, q - d_{N,k,0}). \quad (\text{OCS24})$$

These inequalities apply to the original map, since its full image factorization was proved in OCS18; the ambient dimension has not been equated with its actual rank. There is also an unconditional nonzero observation for the original $k \geq 5$. Each entire function $x_t(w)$ in OCS14 is nonzero. Otherwise its nonzero coordinate covector would annihilate every derivative at zero of $\Pi U \exp(w(A_h - 1/2)/i)[1]$. Those derivatives span the full space, since $[1]$ is cyclic for multiplication by z_1 and ΠU is invertible. A product of finitely many nonzero entire functions is nonzero. The zero sector is nonempty: take the weights $1, 1, N - 2$ and then $(k - 3)/2$ pairs $1, N - 1$. Their sum is a multiple of N . Its monomial curve in OCS9 is nonzero, so $\text{rank } \Lambda \geq 1$.

For $m = 1$, $N = 5$, and every $k \geq 0$, write

$$D_k = \binom{k+3}{3}, \quad \epsilon_k = \begin{cases} 1 & k \equiv 0 \pmod{5}, \\ -1 & k \equiv 1 \pmod{5}, \\ 0 & \text{otherwise.} \end{cases} \quad (\text{OCS25})$$

$$d_{5,k,0} = \frac{D_k + 4\epsilon_k}{5}, \quad d_{5,k,\tau} = \frac{D_k - \epsilon_k}{5} \quad (\tau \neq 0).$$

For each of the four nonidentity powers of C , the trace series is $(1 - T)/(1 - T^5)$, whose coefficient is ϵ_k . The weight- τ projector introduces the Fourier multiplier; the sum of its four nontrivial fifth roots is 4 when $\tau = 0$ and -1 otherwise. This proves both formulas. For the requested original orders the exact counts and resulting kernel lower bounds are

k	D_k	$d_{5,k,0}$	$q = (k+1)^2$	$\max(0, q - d_{5,k,0})$
5	56	12	36	24
9	220	44	100	56
13	560	112	196	84
17	1140	228	324	96
21	2024	404	484	80
25	3276	656	676	20
29	4960	992	900	0

(OCS26)

These are integer evaluations of OCS25. The strict inequality $d_{5,k,0} < q$ holds for $k = 4l + 1$, $l \geq 1$, exactly for the first six orders in the table. They exhaust $k \leq 25$ in this progression. For $k \geq 29$,

$$D_k - 5(k+1)^2 = \frac{(k+1)(k^2 - 25k - 24)}{6} \geq 460 > 4,$$

where $k^2 - 25k - 24$ is positive and increasing from $k = 29$. Even the most negative correction $4\epsilon_k \geq -4$ therefore gives $d_{5,k,0} > q$. This proves the threshold without extrapolating the finite table.

1.6 The actual simple-quartet quadric and a stronger rank bound

Continue with $m = 1$. Order the four roots as $(++)$, $(+-)$, $(-+)$, $(--)$ and let $\mathcal{E}_{\text{prim}} : \mathbb{C}^4 \rightarrow E_h$ be the primary basis isomorphism whose columns are the four $[E_{\alpha,0}]$. Define the actual invertible matrix and the original exponential quartet by

$$\mathbf{H} = F^{-1} \Pi U \mathcal{E}_{\text{prim}}, \quad t_{\epsilon, \eta}(w) = e^{(\eta\gamma - i\epsilon\delta)w}, \quad x(w) = \mathbf{H}t(w). \quad (\text{OCS27})$$

Every column of $\mathbf{H}$ is OCS13 with $d = 0$, namely the Fourier transform of its original primary period times its actual nonzero $v_h(\alpha)$. Each displayed map is invertible, so $\mathbf{H}$ is invertible without a genericity assumption. Put $\mathcal{R}_k = \mathbb{C}[x_1, x_2, x_3, x_4]_k$, the homogeneous polynomials of degree k , and define

$$Q_0(t) = t_{++}t_{--} - t_{+-}t_{-+}, \quad Q(x) = Q_0(\mathbf{H}^{-1}x). \quad (\text{OCS28})$$

This is the literal transformed quadratic relation of the original curve; its coefficients are determined by the periods and unit through OCS27. In particular $Q \neq 0$.

The kernel of homogeneous restriction to that curve is exactly

$$\ker\{\mathcal{R}_k \rightarrow \text{Hol}(\mathbb{C}), f \mapsto f(x(w))\} = Q\mathcal{R}_{k-2} \quad (k \geq 2). \quad (\text{OCS29})$$

Here is a complete verification. First work in the t coordinates. A degree- k monomial has exponent $(2b - k)\gamma - i(2a - k)\delta$, where a is its count of first sign $+$ and b its count of second sign $+$. These $(k + 1)^2$ complex exponents are distinct: equality of their real parts forces b equal and equality of their imaginary parts forces a equal. Equivalently, the two frequencies 2γ and $-2i\delta$ admit no nonzero integer relation, by their real and imaginary parts. Thus the exponentials have no extra polynomial relation of this homogeneous type; their linear independence follows, for example, by differentiating $0, \dots, (k + 1)^2 - 1$ times and using the Vandermonde determinant of the distinct exponents. Every pair (a, b) occurs: the count r_{++} may be any integer between $\max(0, a + b - k)$ and $\min(a, b)$, a nonempty interval, and the other three counts are $a - r_{++}$, $b - r_{++}$, and $k - a - b + r_{++}$.

Within any fixed pair (a, b) , two monomial exponent vectors differ by an integer multiple of $(1, -1, -1, 1)$. After their common monomial is factored out, their difference is therefore the difference of equal powers of $t_{++}t_{--}$ and $t_{+-}t_{-+}$. The identity $A^r - B^r = (A - B) \sum_{j=0}^{r-1} A^{r-1-j} B^j$ shows that this difference is divisible by Q_0 . Any polynomial vanishing on the curve has coefficient sum zero within each pair (a, b) , by the proved exponential independence. It is thus a sum of these monomial differences and is divisible by Q_0 . Its quotient is homogeneous of degree $k - 2$. Conversely $Q_0(t(w)) = 0$, since each of its two products is one. This proves the assertion in t coordinates, and the invertible substitution OCS27 proves OCS29. In particular $\dim(\mathcal{R}_k / Q\mathcal{R}_{k-2}) = (k + 1)^2 = q$; no information about the original curve has been replaced by a freely chosen quadric.

Let $\mathcal{I}_k \subseteq \mathcal{R}_k$ be the span of the weight-zero monomials x^ν , using the weights $1, 2, 3, 4$ from the exact Fourier basis. The coordinate functions of the original projected curve in OCS9 are precisely these monomials restricted to $x(w)$. The span of the vector curve $Le^{wY}[1]$ equals $\text{im } L$: its derivatives at zero contain $LY^d[1]$, $0 \leq d < q$, and $Y^d[1]$ is a basis because Y is an invertible affine change of multiplication by S in the degree filtration. The rank of that vector curve equals the dimension of the span of its coordinate functions, by the equality of row and column ranks of its finite exponential coefficient matrix. Hence OCS29 proves the exact equality

$$\text{rank } \Lambda = d_{5,k,0} - \dim(\mathcal{I}_k \cap Q\mathcal{R}_{k-2}). \quad (\text{OCS30})$$

Fix any multiplicative monomial order, for instance lexicographic order, and write x^μ for the leading monomial of the actual Q , with $|\mu| = 2$. Choose a row-echelon basis of the finite-dimensional intersection in OCS30 in descending monomial order. Its leading monomials are distinct, all have weight zero, and all are divisible by x^μ : the leading monomial of Qf is the product of those of Q and f because the order is multiplicative and the coefficient field has no zero divisors. Thus each such leading monomial is $x^\mu x^\eta$ for a distinct degree- $k - 2$ monomial of weight $-\text{wt}(\mu)$. Counting this set and using OCS30 gives

$$d_{5,k,0} - d_{5,k-2,-\text{wt}(\mu)} \leq \text{rank } \Lambda \leq \min(q, d_{5,k,0}). \quad (\text{OCS31})$$

The left side depends only on the actual leading monomial of the actual period quadric, and remains valid whatever its other coefficients are. A uniform consequence retaining the strongest count independent of that monomial is

$$r_k^- := d_{5,k,0} - \max_{\tau \in \mathbb{Z}/5} d_{5,k-2,\tau} = \left\lfloor \frac{(k+1)^2}{5} \right\rfloor + \mathbf{1}_{5|k} \leq \text{rank } \Lambda. \quad (\text{OCS32})$$

For the equality, $D_k - D_{k-2} = (k+1)^2$, as direct subtraction of the two cubic binomials shows. Formula OCS25 then gives

$$r_k^- = \frac{(k+1)^2 + 4\epsilon_k - \max(4\epsilon_{k-2}, -\epsilon_{k-2})}{5}.$$

For k congruent respectively to $0, 1, 2, 3, 4$ modulo five, its numerator correction is respectively $4, -4, -4, -1, 0$. The corresponding residues of $(k+1)^2$ are $1, 4, 4, 1, 0$. These five cases prove OCS32 for every $k \geq 2$. For the original orders in OCS26 the complete unconditional intervals are therefore

k	r_k^-	$\min(q, d_{5,k,0})$	$\max(0, q - d_{5,k,0})$	$q - r_k^-$
5	8	12	24	28
9	20	44	56	80
13	39	112	84	157
17	64	228	96	260
21	96	404	80	388
25	136	656	20	540
29	180	900	0	720

(OCS33)

The middle two rank columns bound $\text{rank } \Lambda$; the last two columns bound $\dim K$ from below and above, respectively. No endpoint of either interval has been assigned as the actual value without its original minors.

1.7 A second exact finite matrix for the simple-quartet kernel

The quadric calculation also gives the actual kernel a polynomial quotient model. Write $Q(x) = \sum_{|\mu|=2} q_\mu x^\mu$. Let T_Q be the full multiplication matrix $\mathcal{R}_{k-2} \rightarrow \mathcal{R}_k$ in the monomial bases:

$$(T_Q)_{\nu,\eta} = q_{\nu-\eta}, \quad |\nu| = k, \quad |\eta| = k-2, \quad q_{\nu-\eta} = 0 \text{ if an entry of } \nu - \eta < 0.$$

Let T_{out} be its rows of nonzero cyclic weight. Every coefficient q_μ is explicitly fixed by $Q_0(\mathbf{H}^{-1}x)$, and the inverse matrix may be computed as $\text{adj } \mathbf{H} / \det \mathbf{H}$, with nonzero denominator proved in OCS27. Multiplication by Q is injective. The condition that Qf belongs to $\mathcal{I}_k$ is exactly $T_{\text{out}}f = 0$. Thus OCS30 gives the further exact rank identities

$$\begin{aligned} \text{rank } \Lambda &= d_{5,k,0} - D_{k-2} + \text{rank } T_{\text{out}}, \\ \dim K &= D_k - d_{5,k,0} - \text{rank } T_{\text{out}}. \end{aligned} \quad (\text{OCS34})$$

The ranks of this concrete matrix are decided by its finite minors, by the same elimination proof as OCS20. This is an alternative exact calculation from the actual periods, not an assumption about the quadric's coefficients.

To give the typed map behind the kernel equality, let E^* denote the complex linear dual. Define $\Theta : \mathcal{R}_k \rightarrow E^*$ as follows: $\Theta(f)$ is the unique covector such that

$$\Theta(f)(e^{wY}[1]) = f(x(w)) \quad \text{for all } w. \quad (\text{OCS35})$$

For $m = 1$ every function on the right is a linear combination of the q distinct exponentials in OCS16. The primary expansion of $e^{wY}[1]$ gives exactly those exponentials as a basis of covector functions. This proves existence and uniqueness. The proof of OCS29 proves surjectivity and

$\ker \Theta = Q\mathcal{R}_{k-2}$. Moreover $\Theta(\mathcal{I}_k) = \text{im } L^\vee$, where $L^\vee$ denotes the complex linear dual map: the monomial coefficient functionals in the basis OCS8 pull back along L to exactly OCS35 for x^ν of weight zero. These functionals span the full dual of the actual image, since every linear functional on a subspace of a finite-dimensional space extends to the whole space. Restriction of E^* to K has kernel $\text{im } L^\vee$: a covector vanishing on K factors uniquely through $E/K \simeq \text{im } L$. Thus the exact maps give

$$\boxed{\text{coker } T_{\text{out}} \simeq \mathcal{R}_k / (\mathcal{I}_k + Q\mathcal{R}_{k-2}) \xrightarrow{\Theta|_K} K^*} \quad (\text{OCS36})$$

For the first isomorphism, use the direct monomial splitting of $\mathcal{R}_k$ into its zero and nonzero sectors; after quotienting by $\mathcal{I}_k$, multiplication by Q becomes precisely T_{out} . For the second, the surjectivity and kernel were just proved. This identifies the actual lost directions by a full quotient map.

1.8 All primary action directions remain in the mixed calculation

For arbitrary original $m \geq 1$, retain the full primary matrix $\mathfrak{A}$ in OCS17. In the basis $\mathcal{H}$ the original sum action is exactly

$$J e_{a,b,D} = \omega_{a,b} e_{a,b,D} + \begin{cases} i^{-1} e_{a,b,D+1} & D < e-1, \\ 0 & D = e-1, \end{cases} \quad Y\mathcal{H} = \mathcal{H}J. \quad (\text{OCS37})$$

This follows by multiplying $\varepsilon_{a,b}(S - \beta_{a,b})^D$ by $(S - c)/i = \omega_{a,b} + (S - \beta_{a,b})/i$ in its original primary algebra. Consequently the actual possible action leaving the cyclic kernel is the fully specified map

$$\ker \mathfrak{A} \longrightarrow \text{im } \mathfrak{A}, \quad x \longmapsto \mathfrak{A}Jx, \quad \Lambda Y\mathcal{H}x = \mathcal{Q}\mathfrak{A}Jx. \quad (\text{OCS38})$$

The codomain assertion follows from $\mathfrak{A}Jx$ being a value of the same matrix $\mathfrak{A}$ on Jx . For the original inclusion $I : K \hookrightarrow E$, this is the OPG22 defect transported through the exact kernel isomorphism in OCS18. In particular none of the discarded sectors has been declared invariant under Y .

The part of the kernel undetected by all original iterates is exactly computed by the stacked matrix

$$\mathfrak{D} = \begin{bmatrix} \mathfrak{A} \\ \mathfrak{A}J \\ \vdots \\ \mathfrak{A}J^{q-1} \end{bmatrix}, \quad \mathcal{K}_\infty = \mathcal{H} \ker \mathfrak{D}, \quad \dim \mathcal{K}_\infty = q - \text{rank } \mathfrak{D}, \quad \dim(K/\mathcal{K}_\infty) = \text{rank } \mathfrak{D} - \text{rank } \mathfrak{A}. \quad (\text{OCS39})$$

Indeed $\mathfrak{A}J^a x$ is the exact sector-coordinate value of $\Lambda Y^a \mathcal{H}x$ by OCS18 and OCS37. The polynomial $\chi(c + iT)$ annihilates J and has degree q with leading coefficient i^q . It expresses every higher power in the span of the first q , so the stacked kernel is precisely the intersection over all iterates and is J -invariant. Rank-nullity proves both dimension formulas. Its quotient identifies the directions that the original one-step observation loses and subsequent original iterates detect.

Equations OCS17–21, OCS34–39 supply actual finite maps and deciding minors for the original observation and its kernel. The forced kernel losses in OCS26 and the rank intervals OCS33 concern this unchanged projector and these unchanged periods. The endpoint volume calculation still uses the full positive sums of the actual vectors λ_d in OPG16–18 and their exact inherited coordinates OCS11 and OCS19. No volume growth estimate is deduced from these dimension bounds alone.

2 The original quadric orbit and an explicit invariant lifting map

This calculation uses the original simple-quartet construction, with $m = 1$, $k = 4l + 1$, $l \geq 1$, $q = (k + 1)^2$, $0 < \delta < 1/2$, and $\gamma > 2$. The period parameter, its branch, the polynomial contours, the complete amplitude unit, and the observation are those of OPG and OCS. The degree-eight polynomial below is the product of four actual orbit equations. Its degree and all of its coefficients are derived from that orbit; no source order is selected to match an estimate.

2.1 The actual coefficient ring and the original observation

Order the original roots as $(++)$, $(+-)$, $(-+)$, $(--)$, with $\alpha_{\epsilon, \eta} = 1/2 + \epsilon\delta + i\eta\gamma$. Let $\mathcal{E}_{\text{prim}}$ have columns the four original primary idempotents in $E_h = \mathbb{C}[z_1]/(h)$. For the original matrices Π, U, F in OCS2, OCS5, OCS27 put

$$\begin{aligned} \mathbf{H} &= F^{-1}\Pi U \mathcal{E}_{\text{prim}}, & t_{\epsilon, \eta}(w) &= e^{(\eta\gamma - i\epsilon\delta)w}, \\ x(w) &= \mathbf{H}t(w), & Q(x) &= Q_0(\mathbf{H}^{-1}x), & Q_0(t) &= t_{++}t_{--} - t_{+-}t_{-+}. \end{aligned} \quad (\text{OQC1})$$

Every matrix in $\mathbf{H}$ is the specified original isomorphism; in particular its nonzero amplitude factors are the actual $v_h(\alpha)$, and its period entries are the original convergent polynomial integrals. Thus $\det \mathbf{H} \neq 0$ by the included period-isomorphism and primary-basis proofs. Define

$$\mathcal{P}_d = \mathbb{C}[x_1, x_2, x_3, x_4]_d, \quad R_d = \mathcal{P}_d / (Q\mathcal{P}_{d-2}), \quad \mathcal{P}_d = R_d = 0 \quad (d < 0). \quad (\text{OQC2})$$

The matrix of $Q_0(t) = t^T M_0 t$ has entries $(M_0)_{14} = (M_0)_{41} = 1/2$ and $(M_0)_{23} = (M_0)_{32} = -1/2$, all others zero. Hence

$$M = \mathbf{H}^{-T} M_0 \mathbf{H}^{-1}, \quad \det M = \frac{1}{16(\det \mathbf{H})^2} \neq 0. \quad (\text{OQC3})$$

In particular Q is irreducible. Indeed, a reducible homogeneous quadratic over $\mathbb{C}$ is a product of two linear forms, whose symmetric coefficient matrix is a sum of two rank-one matrices and has rank at most two. This contradicts OQC3. The polynomial ring over a field is a unique factorization domain, by the repeated one-variable Gauss lemma, so its irreducible element Q is prime. Consequently $R = \mathbb{C}[x_1, x_2, x_3, x_4]/(Q)$ is an integral domain.

For clarity, the complete homogeneous restriction calculation has the precise kernel $Q\mathcal{P}_{d-2}$. In the t coordinates a degree- d monomial restricts to

$$\exp(((2b - d)\gamma - i(2a - d)\delta)w), \quad 0 \leq a, b \leq d. \quad (\text{OQC4})$$

Every pair occurs: the number of $(++)$ factors can range from $\max(0, a + b - d)$ to $\min(a, b)$. The exponents are distinct by their real and imaginary parts. Their exponentials are linearly independent, since differentiating orders zero through $(d + 1)^2 - 1$ gives a Vandermonde matrix with nonzero determinant. Two monomials giving the same pair have exponent vectors differing by a multiple of $(1, -1, -1, 1)$. After their common monomial is removed, their difference is divisible by $t_{++}t_{--} - t_{+-}t_{-+}$ through $A^r - B^r = (A - B) \sum_{j=0}^{r-1} A^{r-1-j} B^j$. A vanishing polynomial has coefficient sum zero within each pair by the independence just proved, and is therefore a sum of these differences. Conversely $Q_0(t(w)) = 0$. Transport by $\mathbf{H}$ proves the stated kernel and

$$\dim R_d = (d + 1)^2 \quad (d \geq 0). \quad (\text{OQC5})$$

Write $\zeta = e^{2\pi i/5}$ and $D = \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4)$. The literal Fourier calculation gives $CF = FD$. On polynomials define the original induced action and its average by

$$(gf)(x) = f(D^{-1}x), \quad g^5 = 1, \quad \mathbf{R}f = \frac{1}{5} \sum_{a=0}^4 g^a f, \quad \mathcal{I}_d = \mathcal{P}_d^{(g)}. \quad (\text{OQC6})$$

Thus $\mathcal{J}_d$ consists of the weight-zero monomials, since $gx^\nu = \zeta^{-\sum j\nu_j} x^\nu$. This is the dual of the original average of $(C^{-a})^{\otimes d}$; both the inverse powers and the factor $1/5$ are retained.

For the original algebra E_d of the d -fold sum, define $\Theta_d : R_d \xrightarrow{\sim} E_d^\vee$ by

$$\Theta_d([f])(e^{wY_d}[1]) = f(x(w)), \quad Y_d = (A_d - dI/2)/i. \quad (\text{OQC7})$$

Here $\vee$ denotes the complex-linear dual. The primary expansion of the vector on the left gives all distinct exponentials OQC4; therefore existence and uniqueness of this covector and bijectivity follow from the preceding complete restriction calculation. The original observation $\Lambda_d : E_d \rightarrow B_d$ obeys

$$J_d := \text{im}(\mathcal{J}_d \rightarrow R_d), \quad \Theta_d(J_d) = \text{im } \Lambda_d^\vee, \quad \dim J_d = \text{rank } \Lambda_d, \quad R_d/J_d \xrightarrow{\sim} (\ker \Lambda_d)^\vee. \quad (\text{OQC8})$$

Indeed the coordinate functions of the projected pure tensor in OCS9 are precisely $x(w)^\nu$ of weight zero; their pulled-back covectors span the dual of its actual image. A covector vanishing on $\ker \Lambda_d$ factors through that image, which proves the last isomorphism as well. OQC8 specifies the exact original image; it has not been replaced by the whole invariant tensor space.

2.2 The complete orbit decision from the actual coefficients

Put $Q_a = g^a Q$ for $0 \leq a < 5$. The orbit of the line $\mathbb{C}Q$ has size one or five because five is prime. Moreover a size-one orbit has $gQ = Q$, with no other possible scalar. To prove this, write $gQ = \lambda Q$. Then $\lambda^5 = 1$, while OQC3 and the determinant of D , which is $\zeta^{10} = 1$, imply

$$\det(D^{-T}MD^{-1}) = \det M = \lambda^4 \det M. \quad (\text{OQC9})$$

Thus $\lambda^4 = 1$ and $\lambda^5 = 1$, forcing $\lambda = 1$. If any nonidentity power fixed the line, its powers generate the same cyclic group, so the preceding argument applies. This proves the entire orbit classification.

There is an exact coefficient test, requiring no genericity premise. Write the actual $Q = \sum_{|\mu|=2} q_\mu x^\mu$ and put

$$\mathfrak{e}(Q) = \sum_{\substack{|\mu|=2 \\ \sum j\mu_j \not\equiv 0 \pmod{5}}} |q_\mu|^2. \quad (\text{OQC10})$$

All coefficients here are the entries of OQC3, including the full inverse of the original period matrix. Since the monomials are a basis and each summand is nonnegative, $\mathfrak{e}(Q) = 0$ exactly when Q is invariant. The only degree-two weight-zero monomials are x_1x_4 and x_2x_3 . Hence the size-one case is exactly $Q = ax_1x_4 + bx_2x_3$ with $ab \neq 0$; nonvanishing of the two coefficients follows from OQC3. Positive $\mathfrak{e}(Q)$ gives the five distinct irreducible orbit equations. These are exhaustive outcomes of a test on the given periods, not assumptions on them.

In the size-one outcome, multiplication by Q preserves every weight. If Qf is invariant, then $Q(gf - f) = 0$ in the polynomial ring, and $gf = f$. Therefore

$$\ker(\mathcal{J}_k \rightarrow R_k) = Q\mathcal{J}_{k-2}, \quad \text{rank } \Lambda_k = d_{5,k,0} - d_{5,k-2,0}, \quad \dim \ker \Lambda_k = q - d_{5,k,0} + d_{5,k-2,0}. \quad (\text{OQC11})$$

This also proves that $J_k = R_k^{(g)}$: an invariant class can be lifted to a polynomial, and averaging that lift gives an invariant representative. The count itself follows from

$$d_{5,k,0} = \frac{\binom{k+3}{3} + 4\epsilon_k}{5}, \quad \epsilon_k = \mathbf{1}_{k \equiv 0 \pmod{5}} - \mathbf{1}_{k \equiv 1 \pmod{5}}. \quad (\text{OQC12})$$

For a proof, the identity element has symmetric-power generating series $(1-z)^{-4}$; each of the other four powers has series $\prod_{j=1}^4 (1 - z\zeta^j)^{-1} = (1-z)/(1-z^5)$. Averaging their coefficients gives

OQC12. Subtraction and $\binom{k+3}{3} - \binom{k+1}{3} = (k+1)^2$ evaluate the rank in OQC11, for every $k \geq 2$, as

$$\left\lfloor \frac{(k+1)^2}{5} \right\rfloor + \mathbf{1}_{k \equiv 0 \text{ or } 3 \pmod{5}}. \quad (\text{OQC13})$$

The five numerator corrections before division by five are 4, -4, -4, 4, 0, respectively; the residues of $(k+1)^2$ are 1, 4, 4, 1, 0. This verifies every case in OQC13.

2.3 An explicit invariant lift for the five-member orbit

For the size-five outcome determined by OQC10, retain the full actual products

$$\mathcal{H}_Q = \prod_{a=0}^4 Q_a \quad (\deg \mathcal{H}_Q = 10), \quad \mathcal{A}_Q = \prod_{a=1}^4 Q_a \quad (\deg \mathcal{A}_Q = 8). \quad (\text{OQC14})$$

The action permutes all five factors cyclically, so $g\mathcal{H}_Q = \mathcal{H}_Q$. None of the four factors of $\mathcal{A}_Q$ is a scalar multiple of Q . Since each is irreducible and Q is prime, multiplication by $[\mathcal{A}_Q]$ is injective on the integral domain R .

For every homogeneous $f \in \mathcal{P}_d$ the exact trace identity is

$$[5R(\mathcal{A}_Q f)]_{(Q)} = [\mathcal{A}_Q f]_{(Q)}. \quad (\text{OQC15})$$

Indeed $g^a \mathcal{A}_Q$ is the product of all orbit factors except Q_a . For $a \neq 0$ it contains $Q_0 = Q$. Those four terms vanish in the displayed quotient, and the $a = 0$ term is exactly the right side. This proves OQC15 with its original average and full factors, including every period-dependent coefficient.

The lift also has a well-defined typed quotient map. Let $\mathcal{U} = \mathbb{C}[x_1, x_2, x_3, x_4]/(\mathcal{H}_Q)$; the invariant equation $\mathcal{H}_Q$ makes the cyclic action well defined on this quotient. Then

$$\begin{aligned} \mathfrak{L}_d : R_d &\longrightarrow (\mathcal{U}_{d+8})^{(g)}, & [f]_{(Q)} &\longmapsto [5R(\mathcal{A}_Q f)]_{(\mathcal{H}_Q)}, \\ \text{res } \mathfrak{L}_d([f]) &= [\mathcal{A}_Q f]_{(Q)}, & \text{res} : \mathcal{U} &\rightarrow R. \end{aligned} \quad (\text{OQC16})$$

If f changes by Qb , its proposed image changes by $5R(\mathcal{H}_Q b) = 5\mathcal{H}_Q Rb$, hence by zero in $\mathcal{U}$. This proves well-definedness. The image is invariant because it has an invariant polynomial representative; OQC15 proves the second line. Injectivity follows from the injectivity of multiplication by $[\mathcal{A}_Q]$ on R . All maps in OQC16 are thus established, including their kernels.

The actual original invariant image consequently contains

$$[\mathcal{A}_Q]R_{k-8} \subseteq J_k \subseteq R_k. \quad (\text{OQC17})$$

The first inclusion follows by taking the invariant polynomial representatives OQC15, and uses no dimension inference. Therefore, for every $k \geq 8$ in this outcome,

$$\text{rank } \Lambda_k \geq (k-7)^2, \quad \dim \ker \Lambda_k \leq (k+1)^2 - (k-7)^2 = 16k - 48. \quad (\text{OQC18})$$

These inequalities follow from the proved injection and OQC5. They may be combined with the unconditional actual-matrix bounds in OCS24, OCS31–33 by taking the stronger lower and upper bounds. For the original degrees $k = 9, 13, 17, 21, 25, 29$, the displayed upper bounds on the original kernel are respectively 96, 160, 224, 288, 352, 416; at the first two degrees the earlier unconditional bounds may be stronger. No such upper bound is assigned to the size-one outcome, whose exact rank is OQC11. Together OQC10–18 compute the complete orbit alternatives from the original coefficients. Evaluating $\mathfrak{e}(Q)$ in the original period family is a separate calculation on those explicit coefficients; no outcome is postulated here.

2.4 The induced original dual map and its complete action defect

At $k \geq 9$ the degree $d = k - 8$ is positive, and the original d -fold sum algebra is defined using the same h, Π, U . For the size-five outcome, OQC7, OQC8 and OQC17 give the uniquely typed injective map

$$\begin{aligned} \mathfrak{J}_k : E_{k-8}^\vee &\hookrightarrow B_k^\vee, \\ \Lambda_k^\vee \mathfrak{J}_k &= \Theta_k M_{[\mathcal{A}_Q]} \Theta_{k-8}^{-1}, \quad \mathfrak{J}_k^\vee : B_k \twoheadrightarrow E_{k-8}. \end{aligned} \quad (\text{OQC19})$$

The right side of the second line has image in $\text{im } \Lambda_k^\vee$ by OQC17, and $\Lambda_k^\vee$ is injective because Λ_k is onto its actual image. This gives existence and uniqueness. Each of the three right-hand maps is injective, so $\mathfrak{J}_k$ is injective. Finite-dimensional duality then proves the stated surjection, using the canonical bidual identifications. The new degree $k - 8$ records contraction against the degree-eight polynomial derived in OQC14. It does not replace either original Gamma source of order 1 or k .

The surviving action and its defect are fully determined. Put

$$\Omega = \text{diag}(\gamma - i\delta, -\gamma - i\delta, \gamma + i\delta, -\gamma + i\delta), \quad G = H\Omega H^{-1}, \quad (\mathcal{D}f)(x) = \sum_{j=1}^4 (Gx)_j \partial_j f(x). \quad (\text{OQC20})$$

Here $x'(w) = Gx(w)$ by OQC1. In the original t coordinates, the weights of each product in Q_0 sum to zero, so $\mathcal{D}Q = 0$. Thus this derivation acts on each R_d . Differentiating OQC7 proves exactly

$$\Theta_d \mathcal{D} = Y_d^\vee \Theta_d. \quad (\text{OQC21})$$

Both sides evaluated at $e^{wY_d}[1]$ are the derivative of $f(x(w))$, and these vectors span E_d by the distinct exponentials; hence equality holds as maps, not only on a curve. Leibniz's rule gives the complete defect

$$\begin{aligned} Y_k^\vee \Lambda_k^\vee \mathfrak{J}_k - \Lambda_k^\vee \mathfrak{J}_k Y_{k-8}^\vee &= \Theta_k M_{[\mathcal{D}\mathcal{A}_Q]} \Theta_{k-8}^{-1}, \\ \mathcal{D}\mathcal{A}_Q &= \sum_{a=1}^4 (\mathcal{D}Q_a) \prod_{\substack{1 \leq b \leq 4 \\ b \neq a}} Q_b. \end{aligned} \quad (\text{OQC22})$$

The codomain in the first line is $E_k^\vee$. Its image has not been asserted to lie in $\text{im } \Lambda_k^\vee$. Restriction to $\ker \Lambda_k$ supplies the exact remaining map into $(\ker \Lambda_k)^\vee$, through OQC8. In particular OQC19 is not used as an unproved intertwining of boundary actions. OQC22 retains every term that can leave the observed dual subspace.

Finally, the failure of the cyclic average to commute with the original action is explicitly calculable before any quotient:

$$([\mathcal{D}, g^a]f)(x) = (\nabla f)(D^{-a}x)^T (D^{-a}G - GD^{-a})x, \quad [\mathcal{D}, R] = \frac{1}{5} \sum_{a=0}^4 [\mathcal{D}, g^a]. \quad (\text{OQC23})$$

The chain rule applied separately to $f(D^{-a}x)$ and to $(\mathcal{D}f)(D^{-a}x)$ proves both identities. Thus all action comparisons in the invariant lift have explicit original coefficient maps. The rank estimates in OQC18 concern exactly the kernel used in the original Gamma and arithmetic quotient Grams, because those constructions use the same Λ_k . An estimate for its volume still requires those original metrics; no volume conclusion has been inferred solely from its dimension.

3 Exact original rank from the complete quadric orbit

Retain every original object and map in OQC1–23. In particular $m = 1$, $k = 4l + 1$, $l \geq 1$, $q = (k + 1)^2$, $\mathcal{P} = \mathbb{C}[x_1, x_2, x_3, x_4]$, and $gf(x) = f(D^{-1}x)$ for the original Fourier matrix $D = \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4)$. The polynomial $Q = Q_0(H^{-1}x)$ has its actual original period and amplitude coefficients. Write $\mathcal{I} = \mathcal{P}^{(g)}$, $R = \mathcal{P}/(Q)$ and $J = \text{im}(\mathcal{I} \rightarrow R)$, with the grading and the actual observation identification of OQC8. For negative indices put $\mathcal{I}_d = 0$ and $d_{5,d,0} = 0$.

The orbit decision is the exhaustive test OQC10. The size-one case already has the exact rank OQC11–13. The following calculation completes the other case using the same original coefficients. It does not assert that a period parameter belongs to that case before its coefficient test has been evaluated.

3.1 Every invariant vanishing polynomial contains the full orbit

In the size-five case, let $Q_a = g^a Q$ and retain exactly $\mathcal{H}_Q = \prod_{a=0}^4 Q_a$ from OQC14. Each Q_a is irreducible, because an invertible linear substitution preserves factorizations and Q is irreducible by OQC3. The five factors are pairwise nonassociate by the proved orbit classification, and each is prime in the original polynomial ring. Consequently

$$\boxed{\mathcal{I} \cap Q\mathcal{P} = \mathcal{H}_Q\mathcal{I}, \quad \ker(\mathcal{I}_d \rightarrow R_d) = \mathcal{H}_Q\mathcal{I}_{d-10}.} \quad (\text{OQX1})$$

Here is the complete divisibility proof. If $f \in \mathcal{I}$ and $f = Qb$, then for every $a = 0, \dots, 4$ the exact equality $g^a f = f$ gives $f = Q_a g^a b$. Thus every one of the five original prime factors divides f . Distinctness of the factors implies that their product divides f : after factoring out Q_0 , primality of Q_1 and the fact that Q_1 does not divide Q_0 force Q_1 to divide the remaining quotient. Repeating this argument with Q_2, Q_3, Q_4 proves $f = \mathcal{H}_Q b_0$. The action permutes the five exact factors cyclically, so $g\mathcal{H}_Q = \mathcal{H}_Q$. Applying g to this last equality and using $gf = f$ gives $\mathcal{H}_Q(gb_0 - b_0) = 0$. The polynomial ring is an integral domain and $\mathcal{H}_Q \neq 0$, whence $gb_0 = b_0$. This proves the first inclusion in OQX1. Conversely an invariant b_0 gives an invariant $\mathcal{H}_Q b_0$ divisible by Q , which proves the reverse inclusion. The product is homogeneous of degree ten. If its product with b_0 is homogeneous of degree d , decomposition of b_0 into homogeneous parts shows that only its part of degree $d - 10$ can be nonzero: each other part has zero product with $\mathcal{H}_Q$ and hence is zero. This proves the graded kernel formula, including $d < 10$.

3.2 The exact invariant quotient and the restriction map

Let $\mathcal{U} = \mathcal{P}/(\mathcal{H}_Q)$ as in OQC16. The following are isomorphisms of graded complex algebras with their specified maps:

$$\boxed{\mathcal{I}/(\mathcal{H}_Q\mathcal{I}) \xrightarrow{[f] \mapsto [f]_{(\mathcal{H}_Q)}} \mathcal{U}^{(g)} \xrightarrow{\text{res}} J \subseteq R.} \quad (\text{OQX2})$$

The action on $\mathcal{U}$ is defined because $\mathcal{H}_Q$ is invariant. If $[f]_{(\mathcal{H}_Q)}$ is invariant, averaging its representatives gives $[Rf]_{(\mathcal{H}_Q)} = [f]_{(\mathcal{H}_Q)}$, since $[g^a f] = [f]$ for every a . Thus the first map is onto. Its kernel before passage to the displayed domain is $\mathcal{I} \cap \mathcal{H}_Q\mathcal{P} = \mathcal{H}_Q\mathcal{I}$: the reverse inclusion is immediate, and for $f = \mathcal{H}_Q b \in \mathcal{I}$, cancellation in $\mathcal{H}_Q(gb - b) = 0$ proves $b \in \mathcal{I}$. This proves that the first map is injective as well.

The full quotient restriction $\text{res} : \mathcal{U} \rightarrow R$ is well defined because $(\mathcal{H}_Q) \subseteq (Q)$; its kernel is exactly $Q\mathcal{P}/\mathcal{H}_Q\mathcal{P}$. For its invariant subspace, choose the invariant representative just constructed. If its restriction vanishes, that representative belongs to $\mathcal{I} \cap Q\mathcal{P}$, which is $\mathcal{H}_Q\mathcal{I}$ by OQX1. Its class in $\mathcal{U}$ is therefore zero. The invariant restriction is injective. Its image is exactly J , since an invariant representative restricts to J by definition and every element of J has such a representative. Both maps are induced by polynomial quotient maps, so they preserve products, sums, complex scalars and degrees. No action of g on R is asserted in the size-five case. In particular the second map retains precisely the invariant image of the original observation, with its complete restriction kernel already computed.

3.3 All original degrees and the exact kernel dimension

The map Θ_k and the original observation in OQC8 identify $\dim J_k$ with $\text{rank } \Lambda_k$ and identify R_k/J_k with $(\ker \Lambda_k)^\vee$. Multiplication by the nonzero polynomial $\mathcal{H}_Q$ is injective in $\mathcal{P}$. Taking dimensions in OQX1 thus gives, for every nonnegative degree d , the exact invariant restriction $\text{rank } \dim J_d = d_{5,d,0} - d_{5,d-10,0}$. At the original degrees this evaluates to

$$\boxed{\text{rank } \Lambda_k = d_{5,k,0} - d_{5,k-10,0} = k^2 - 6k + 17, \quad \dim \ker \Lambda_k = 8k - 16, \quad k = 4l + 1, \quad l \geq 1,} \quad (\text{OQX3})$$

in the size-five outcome of the actual coefficient test. To verify every constant and degree range, for $k \geq 10$ the complete character formula OQC12 gives

$$d_{5,k,0} - d_{5,k-10,0} = \frac{1}{5} \left\{ \binom{k+3}{3} - \binom{k-7}{3} + 4(\epsilon_k - \epsilon_{k-10}) \right\}.$$

The two ϵ values are equal because their indices differ by ten. Expanding the remaining cubics gives

$$\binom{k+3}{3} - \binom{k-7}{3} = \frac{(k^3 + 6k^2 + 11k + 6) - (k^3 - 24k^2 + 191k - 504)}{6} = 5(k^2 - 6k + 17).$$

The only original degrees below ten are five and nine. Their negative-degree terms in OQX1 vanish, while the same exact character formula gives

$$d_{5,5,0} = (56 + 4)/5 = 12 = 5^2 - 6 \cdot 5 + 17, \quad d_{5,9,0} = 220/5 = 44 = 9^2 - 6 \cdot 9 + 17.$$

This proves the first equality of OQX3 for every original $l \geq 1$. Since $\dim E_k = \dim R_k = (k+1)^2$, rank-nullity gives $(k+1)^2 - (k^2 - 6k + 17) = 8k - 16$, proving the second equality. For the original degrees already tabulated in OCS26, the exact five-orbit values are

k	$\text{rank } \Lambda_k$	$\dim \ker \Lambda_k$
5	12	24
9	44	56
13	108	88
17	204	120
21	332	152
25	492	184
29	684	216

These are exact values for this exhaustively classified orbit outcome, replacing the less precise bounds OQC18 wherever that outcome occurs. The full invariant lift and its action defect OQC15–23 remain valid with all original coefficients. The size-one outcome retains its different exact values OQC11–13. No choice between the two outcomes has been made here without evaluating the original period coefficient test, and no estimate for a metric volume has been inferred from its dimension alone.

4 The actual period quadric and its cyclic continuation

This calculation uses the original simple quartet, $m = 1$, and the literal period matrix, amplitude unit and cyclic matrix of BRU and OPG. Its formulas evaluate the transported quadric from those data. All source orders elsewhere in the programme remain unchanged.

4.1 Original roots, entire unit and centred phase

Order the roots and keep their actual values as

$$\begin{aligned} \alpha_1 &= \frac{1}{2} + \delta + i\gamma, & \alpha_2 &= \frac{1}{2} + \delta - i\gamma, & \alpha_3 &= \frac{1}{2} - \delta + i\gamma, & \alpha_4 &= \frac{1}{2} - \delta - i\gamma, \\ 0 < \delta < \frac{1}{2}, & \gamma > 2, & h(s) &= \prod_{j=1}^4 (s - \alpha_j), & g &= 2\xi, & v_h &= g/h. \end{aligned} \quad (\text{PQO1})$$

The four zero orders of g are complete and equal to one. Thus v_h is entire and has nonzero value at each displayed root. Put $w = s - 1/2$, and define the following original constants:

$$\begin{aligned} z_+ &= \delta + i\gamma, & z_- &= \delta - i\gamma, & \lambda &= z_+^2, & \mu &= z_-^2, \\ A &= 2(\gamma^2 - \delta^2), & B &= (\delta^2 + \gamma^2)^2, & \mathfrak{a} &= \Phi_h(1/2) = \frac{1}{160} + \frac{A}{24} + \frac{B}{2}, \\ h(1/2 + w) &= w^4 + Aw^2 + B, & \Phi_h(1/2 + w) &= \mathfrak{a} + \frac{w^5}{5} + \frac{Aw^3}{3} + Bw. \end{aligned} \quad (\text{PQO2})$$

The product of the two quadratics $((w - \delta)^2 + \gamma^2)((w + \delta)^2 + \gamma^2)$ proves the quartic identity. Integrating it and using $\Phi_h(0) = 0$ proves the phase and constant in PQO2. In particular the translation has not removed the original constant $\mathfrak{a}$.

The theta integral IPM.2 has real coefficients, real theta kernel, and an expression unchanged by $s \mapsto 1 - s$. It therefore gives $g(\bar{s}) = \overline{g(s)}$ and $g(1 - s) = g(s)$ directly; the same two identities hold for h . Taking their entire quotient gives

$$v_h(\alpha_1) = v_h(\alpha_4) = a, \quad v_h(\alpha_2) = v_h(\alpha_3) = \bar{a}, \quad a \neq 0. \quad (\text{PQO3})$$

These are the full actual unit values. Their magnitude and phase are not assigned. Set $t_+ = a^{-2}$ and $t_- = \bar{a}^{-2}$. For $j = 0, 1, 2, 3$ retain

$$\sigma_j = t_+ \lambda^j - t_- \mu^j, \quad \sigma_2 = -A\sigma_1 - B\sigma_0, \quad \sigma_3 = (A^2 - B)\sigma_1 + AB\sigma_0. \quad (\text{PQO4})$$

The recurrence follows because $\lambda + \mu = -A$ and $\lambda\mu = B$. Each σ_j is purely imaginary. Furthermore σ_0 and σ_1 cannot both vanish: the first would give $t_+ = t_- \neq 0$, and the second would then give $t_+(\lambda - \mu) = 0$, whereas $\lambda - \mu = 4i\delta\gamma \neq 0$.

4.2 The original periods as an absolutely convergent series

Fix $u \neq 0$ on its original logarithm branch. Let $\ell_j(r) = r \exp(i(\arg u + \pi + 2\pi j)/5)$, $\Gamma_j = -\ell_0 + \ell_j$, $1 \leq j \leq 4$. Let Π integrate the degree-less-than-four representative against $e^{\Phi_h(s)/u} ds$ on these original oriented contours. Write $F_{ja} = \zeta^{ja} - 1$, $\zeta = e^{2\pi i/5}$, $1 \leq j, a \leq 4$. Direct finite geometric sums prove

$$(F^{-1})_{aj} = \frac{1}{5} \zeta^{-ja}, \quad CF = FD, \quad D = \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4), \quad (\text{PQO5})$$

where $(Cx)_j = x_{j+1} - x_1$ for $j < 4$ and $(Cx)_4 = -x_1$. For example multiplying the asserted inverse by F gives $\frac{1}{5} \sum_{j=1}^4 \zeta^{-ja} (\zeta^{jb} - 1) = \delta_{ab}$; substitution into each row gives $CF = FD$.

Let T be the exact increasing-coefficient translation matrix $T_{rb} = \binom{b}{r} 2^{-(b-r)}$ for $0 \leq r \leq b \leq 3$, and zero otherwise. Thus it sends coefficients of $p(s)$ to those of $p(1/2 + w)$, with determinant one. Define the centred Fourier-period matrix

$$P_{ar}(u) = \frac{1}{5} \sum_{j=1}^4 \zeta^{-ja} \int_{\Gamma_j} e^{(w^5/5 + Aw^3/3 + Bw)/u} w^r dw, \quad 1 \leq a \leq 4, \quad 0 \leq r \leq 3. \quad (\text{PQO6})$$

The contours in this formula start at $w = 0$. They give precisely the translated original period by

$$F^{-1}\Pi = e^{a/u}P(u)T. \quad (\text{PQO7})$$

To justify this exact contour translation, the substitution $w = s - 1/2$ first gives rays starting at $-1/2$. Join each shifted ray at a large radius to the corresponding unshifted ray. On the joining segment their arguments differ by $O(1/r)$, and the leading real phase is $-r^5/(5|u|) + O(r^4)$; all lower phase terms and polynomial factors are dominated by this negative leading term. Those outer joining integrals tend to zero. The two common-origin joining integrals cancel in $-\ell_0 + \ell_j$ with their opposite orientations. Cauchy's theorem for the entire integrand proves equality in the limit. Substitution of the polynomial gives exactly the matrix T .

For every integer $L \geq 1$ define, on the same branch,

$$G_L(u) = u^{L/5} 5^{L/5-1} e^{\pi i L/5} \Gamma(L/5). \quad (\text{PQO8})$$

Integration of the leading phase on the two rays gives $\int_{\Gamma_j} e^{w^5/(5u)} w^{L-1} dw = G_L(u)(\zeta^{jL} - 1)$ by the real substitution $r^5/(5|u|)$ on each ray, including its phase in dw . Expanding the remaining two exponentials now gives the exact entries

$$P_{ar}(u) = \sum_{\substack{p,q \geq 0 \\ r+3p+q+1 \equiv a \pmod{5}}} \frac{A^p B^q}{3^p p! q! u^{p+q}} G_{r+3p+q+1}(u). \quad (\text{PQO9})$$

Terms with $r+3p+q+1 \equiv 0$ have zero two-ray integral in every column and contribute zero before the Fourier map. To prove every exchange, the sum of the absolute values of the unintegrated series on either ray is bounded by a constant times

$$R^r \exp\left(-\frac{R^5}{5|u|} + \frac{|A|R^3}{3|u|} + \frac{|B|R}{|u|}\right).$$

Here R is the radial coordinate and r is the fixed column degree. This integrable majorant proves absolute termwise integration. It is uniform on compact coefficient and nonzero- u sets on the chosen branch. The finite Fourier sum in PQO5 then selects exactly the stated residue class. Thus PQO9 is an actual period formula, with all coefficients and all terms retained.

For additional finite computation, let I_d denote the vector of the four centred ray moments of w^d . Integration of the derivative of $e^{(w^5/5 + Aw^3/3 + Bw)/u} w^d$, with the same vanishing outer endpoints and cancelling inner endpoints, proves

$$I_{d+4} + AI_{d+2} + BI_d = -u d I_{d-1} \quad (d \geq 0), \quad u \partial_B I_d = I_{d+1}, \quad 3u \partial_A I_d = I_{d+3}. \quad (\text{PQO10})$$

The first right side is zero at $d = 0$. Coefficient derivatives are justified by the same compact domination. These are the original polynomial period recurrences, and they compute every higher moment from the first four and the actual coefficients.

The determinant is especially explicit in these centred coordinates:

$$\det \Pi = -(2\pi)^2 u^2 e^{4a/u}, \quad \det F = -25\sqrt{5}, \quad \det P(u) = \frac{(2\pi)^2}{25\sqrt{5}} u^2. \quad (\text{PQO11})$$

For the first equality use the full PDE1–13 determinant calculation at $n = 4$: the phase is $e^{15\pi i} = -1$ and the critical-value sum is $4a$. This critical sum also follows directly by pairing w with $-w$ in the odd centred phase. For F , subtract row zero from the other rows in the five-by-five Fourier Vandermonde and expand its first column. Its phase is $e^{13\pi i} = -1$ and its magnitude $5^{5/2}$, by the finite Fourier orthogonality identity. The last equality follows by taking determinants in PQO7 and using $\det T = 1$. The Gamma grid-product proof PGCC1–6 retains the full evaluated constant in the first equality. In particular every inverse below exists on the full stated domain.

4.3 The actual transformed quadric and every deciding coefficient

Let V be the centred evaluation matrix with rows $(1, z_j, z_j^2, z_j^3)$, where $(z_1, z_2, z_3, z_4) = (z_+, z_-, -z_-, -z_+)$. Its inverse sends the four primary values to their unique degree-less-than-four polynomial. Thus the actual matrix from primary coordinates to the Fourier-period space is

$$B_{\text{per}} = F^{-1} \Pi U J_{\text{prim}} = e^{a/u} P(u) V^{-1} \text{diag}(a, \bar{a}, \bar{a}, a). \quad (\text{PQO12})$$

This follows either by applying all maps to the four CRT idempotents or by evaluating the full unit at each root. It agrees with OCS and OPG in the given original orientation.

For $Q_0(t) = t_1 t_4 - t_2 t_3$, define $Q_u(x) = Q_0(B_{\text{per}}^{-1} x)$. On the unchanged original period coordinate $\mathbf{y} \in \mathbb{C}^4$ the quadratic is $Q_u^{\text{per}}(\mathbf{y}) = Q_0((\Pi U J_{\text{prim}})^{-1} \mathbf{y}) = Q_u(F^{-1} \mathbf{y})$, because $F B_{\text{per}} = \Pi U J_{\text{prim}}$. Thus the exact isomorphism $\mathbf{y} = Fx$ carries the following quadric to the original period space. If $p(w) = p_0 + p_1 w + p_2 w^2 + p_3 w^3$, multiplication gives

$$p(w)p(-w) = p_0^2 + (2p_0 p_2 - p_1^2)w^2 + (p_2^2 - 2p_1 p_3)w^4 - p_3^2 w^6.$$

Consequently the exact symmetric matrix in the centred coefficient frame is

$$S = \begin{pmatrix} \sigma_0 & 0 & \sigma_1 & 0 \\ 0 & -\sigma_1 & 0 & -\sigma_2 \\ \sigma_1 & 0 & \sigma_2 & 0 \\ 0 & -\sigma_2 & 0 & -\sigma_3 \end{pmatrix}, \quad \boxed{Q_u(x) = x^\top H(u)x, \quad H(u) = e^{-2a/u} P(u)^{-\top} S P(u)^{-1}.}$$

(PQO13)

Indeed applying the preceding product formula at z_+ and z_- with multipliers t_+ and $-t_-$ gives exactly S . This retains both actual unit values through every σ_j . The two parity blocks have determinants

$$\sigma_0 \sigma_2 - \sigma_1^2 = -t_+ t_- (\lambda - \mu)^2, \quad \sigma_1 \sigma_3 - \sigma_2^2 = -t_+ t_- B(\lambda - \mu)^2.$$

Thus S and $H(u)$ are nondegenerate, with the exact determinant

$$\det H(u) = e^{-8a/u} \frac{B(t_+ t_-)^2 (\lambda - \mu)^4}{(\det P(u))^2} \neq 0. \quad (\text{PQO14})$$

Write $N(u) = \text{adj } P(u)$, with entries N_{ra} , $0 \leq r \leq 3$, $1 \leq a \leq 4$, and retain $\Delta(u) = \det P(u)$ from PQO11. Then every coefficient is given by the explicit minor formula

$$H_{ab}(u) = \frac{e^{-2a/u}}{\Delta(u)^2} \{ \sigma_0 N_{0a} N_{0b} + \sigma_1 (N_{0a} N_{2b} + N_{2a} N_{0b} - N_{1a} N_{1b}) \\ + \sigma_2 (N_{2a} N_{2b} - N_{1a} N_{3b} - N_{3a} N_{1b}) - \sigma_3 N_{3a} N_{3b} \}. \quad (\text{PQO15})$$

This is matrix multiplication in PQO13 using $P^{-1} = N/\Delta$. Each N_{ra} is the signed original three-by-three period minor. Its entries are given by the absolutely convergent series PQO9 or by the original integrals PQO6. The polynomial coefficient of x_a^2 is H_{aa} ; that of $x_a x_b$, $a < b$, is $2H_{ab}$. Both factors are retained in this specified quadratic form.

The exact eight deciding entries are

$$\mathcal{E}(u) = (H_{11}, H_{22}, H_{33}, H_{44}, H_{12}, H_{13}, H_{24}, H_{34}). \quad (\text{PQO16})$$

The cyclic action on quadrics is the pullback $Q_u \mapsto Q_u(D^{-1}x)$; its symmetric matrix is $D^{-\top} H(u) D^{-1}$. Since $D^5 = I$ and $\det D = 1$, the line spanned by a nondegenerate form can be stable only with character one. In fact if its multiplier is η , then $\eta^5 = 1$ by the group action, while taking determinants gives $\eta^4 = 1$ because $\det H(u) \neq 0$. Thus $\eta = 1$. Entrywise, invariance is precisely $(\zeta^{-(a+b)} - 1)H_{ab} = 0$. The only pairs with $a + b \equiv 0$ are $(1, 4)$ and $(2, 3)$. Therefore

The quadric is invariant exactly on $Z = \{u : \mathcal{E}(u) = 0\}$;
 $u \in Z \implies Q_u(x) = 2H_{14}(u)x_1 x_4 + 2H_{23}(u)x_2 x_3, \quad H_{14}(u)H_{23}(u) \neq 0$;
 at every $u \notin Z$ its orbit consists of five distinct quadrics.

(PQO17)

Every assertion follows from the entry test and the rank-four determinant. The orbit size divides five; it is one exactly when the line is stable. Thus no two-member orbit occurs. Moreover $F^{-1}C^{-1} = D^{-1}F^{-1}$ by PQO5, so this is exactly the orbit of $Q_u^{\text{per}}(C^{-j}\mathbf{y})$ in the original period coordinates, with the same five indexed maps. PQO15–17 determine the exact exceptional locus by original period minors and original amplitude values, without assigning generic values to them.

4.4 The cyclic action is the actual original continuation in u

Continue the original logarithm of u by $2\pi i$. Its rays advance from ℓ_j to ℓ_{j+1} , and their oriented differences advance by the exact original matrix C . For the last row $\ell_5 = \ell_0$, giving $-\Gamma_1$. Equivalently every summand of PQO9 acquires the phase

$$e^{2\pi i((r+3p+q+1)/5-p-q)} = \zeta^{r+3p+q+1} = \zeta^a.$$

The original constant $e^{a/u}$ is single-valued under this continuation. Hence

$$\begin{aligned} P(\log u + 2\pi i) &= DP(\log u), & B_{\text{per}}(\log u + 2\pi i) &= DB_{\text{per}}(\log u), \\ H(\log u + 2\pi i) &= D^{-\top} H(\log u) D^{-1}. \end{aligned} \tag{PQO18}$$

The contour argument and the convergent series prove the same map. Thus PQO17 describes the actual period continuation of this original quadric. It introduces no freely chosen cyclic action on its coefficients.

4.5 The large-parameter geometry and a positive-ray obstruction

Let $\varepsilon = u^{-1/5}$ on the same branch, and set $D_\varepsilon = \text{diag}(\varepsilon^{-1}, \varepsilon^{-2}, \varepsilon^{-3}, \varepsilon^{-4})$. The exact series PQO9 gives

$$P(u) = \mathcal{P}(A\varepsilon^2, B\varepsilon^4)D_\varepsilon, \quad \mathcal{P}(0,0) = \text{diag}(g_1, g_2, g_3, g_4), \quad g_a = 5^{a/5-1}e^{\pi ia/5}\Gamma(a/5). \tag{PQO19}$$

The entries of $\mathcal{P}$ are entire in the two displayed arguments, by the same absolute integrable series majorant. The same determinant proof PQO11 applies to every complex pair of lower odd-phase coefficients, since the centred phase remains odd and its critical values pair to zero with full multiplicities. Thus their determinant is the nonzero constant $(2\pi)^2/(25\sqrt{5})$, by PQO11 and $\det D_\varepsilon = u^2$. Their inverse is therefore holomorphic everywhere, given by its adjugate divided by that precise constant. Near $\varepsilon = 0$ it is $\text{diag}(g_a^{-1}) + O(\varepsilon^2)$. This bound follows from its convergent power series after substituting $(A\varepsilon^2, B\varepsilon^4)$; it holds on a full small complex disc, with the fixed original A, B .

If $\sigma_0 \neq 0$, PQO13 and PQO19 give

$$e^{2a/u}H_{11}(u) = \frac{\sigma_0}{g_1^2}\varepsilon^2 + O(\varepsilon^4).$$

If $\sigma_0 = 0$, then $\sigma_1 \neq 0$ by PQO4, and they give

$$e^{2a/u}H_{22}(u) = -\frac{\sigma_1}{g_2^2}\varepsilon^4 + O(\varepsilon^6). \tag{PQO20}$$

For verification of the orders, insert $P^{-1} = D_\varepsilon^{-1}\mathcal{P}^{-1}$ into PQO13. The middle matrix has entries $S_{rs}\varepsilon^{r+s+2}$. Its first possible degree is two, from S_{00} , and its next possible degree is four, from S_{02}, S_{20}, S_{11} . The constant term of $\mathcal{P}^{-1}$ is diagonal, so the indicated leading entry is exactly the displayed nonzero coefficient. Every further contribution has the stated higher degree. This also proves the expansion when some other σ_j vanish.

Consequently for every fixed original quartet and its full actual amplitude there exists a finite radius outside which the actual orbit has five distinct quadrics. This conclusion holds on every original logarithm branch: the estimate is on the full complex ε disc, and its leading coefficient

is nonzero. The set Z of PQO17 is a discrete subset of the logarithmic u surface. Indeed the decisive entry just proved is holomorphic on that connected surface and is not identically zero; its zeros are isolated by its local power series, and Z is contained in that zero set. The explicit finite-radius bound from the companion period-connection calculation strengthens this argument while retaining the same actual coefficients.

Moreover PQO18 multiplies each deciding entry by its nonzero root-of-unity factor, so the condition $\mathcal{E}(u) = 0$ is unchanged by every full logarithm turn. Therefore Z is the inverse image under $\log u \mapsto u$ of a well-defined set $Z_{\text{abs}} \subset \mathbb{C}^*$. On a sufficiently small disc about any nonzero u , choose one logarithm branch. The preceding isolated-zero proof shows that Z_{abs} is discrete and closed on that disc. The explicit radius in the companion calculation gives $|u| < R_*$ throughout Z_{abs} . Consequently its only possible accumulation point in the finite u -plane is zero. In particular each compact annulus about zero contains only finitely many such points: an infinite subset of that compact set would have a nonzero accumulation point, contradicting the proved local isolated-zero property.

For positive real u retain the actual lift $\log u = \ln u + 2\pi i \ell_{\text{br}}$, $\ell_{\text{br}} \in \mathbb{Z}$. The exact series has an additional real structure. Every surviving index in PQO9 is $a + 5L$, and its phase at the lift $\ell_{\text{br}} = 0$ is $e^{\pi i a/5}(-1)^L$. Its remaining coefficients are real. The actual lift multiplies row a by $\zeta^{\ell_{\text{br}} a}$, by PQO18. Absolute convergence therefore proves

$$P(u) = E_{\ell_{\text{br}}} R(u), \quad E_{\ell_{\text{br}}} = D^{\ell_{\text{br}}} \text{diag}(e^{\pi i/5}, e^{2\pi i/5}, e^{3\pi i/5}, e^{4\pi i/5}), \quad R(u) \text{ real and invertible.} \quad (\text{PQO21})$$

Let M be multiplication by w in the increasing centred polynomial frame, so

$$M = \begin{pmatrix} 0 & 0 & 0 & -B \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -A \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad c_3 = \frac{t_+ - t_-}{4i\delta\gamma}, \quad c_1 = \frac{t_+ + t_-}{2} - (\delta^2 - \gamma^2)c_3. \quad (\text{PQO22})$$

Both c_1, c_3 are real by PQO3, and the polynomial $c_1 + c_3 w^2$ takes values t_+, t_-, t_-, t_+ at the four original roots. Hence $c_1 M + c_3 M^3$ is precisely multiplication by $w(j_h v_h)^{-2}$, with its entire unit retained.

The companion period-connection proof establishes the exact identity between cyclic invariance of Q_u and

$$[P(u)(c_1 M + c_3 M^3)P(u)^{-1}, D] = 0. \quad (\text{PQO23})$$

It proves this by the flat residue/reflection form on the original polynomial periods and retains all its constants. Using that proved identity, PQO21 gives a further precise obstruction on the positive ray. Since D has distinct diagonal entries, a matrix commuting with it is diagonal. The matrix in PQO23 is $E_{\ell_{\text{br}}}$ times a real matrix times $E_{\ell_{\text{br}}}^{-1}$; if diagonal, its diagonal entries must therefore be real. Its eigenvalues are exactly

$$\frac{\delta + i\gamma}{a^2}, \quad \frac{\delta - i\gamma}{\bar{a}^2}, \quad -\frac{\delta - i\gamma}{\bar{a}^2}, \quad -\frac{\delta + i\gamma}{a^2}, \quad (\text{PQO24})$$

as follows by evaluation at the four distinct roots of h . Thus every positive-real- u point in Z satisfies the explicit original-unit equation

$$\text{Im} \left(\frac{\delta + i\gamma}{a^2} \right) = 0. \quad (\text{PQO25})$$

If that imaginary part is nonzero for the original unit, the positive ray contains no point of Z . When it is zero, the four displayed eigenvalues are the two nonzero real numbers $r, -r$, each twice, and the complete period-minor tests PQO15–17 still determine Z . The equation is not assumed to hold or fail. Together the exact minor tests, the actual period continuation, the proved large-parameter result and this real-structure obstruction give control of the original quadric family with every amplitude value and phase present.

5 The original period quadric as a residue commutator

Retain $0 < \delta < 1/2$, $\gamma > 2$, the ordered simple quartet $\alpha_1 = 1/2 + \delta + i\gamma$, $\alpha_2 = 1/2 + \delta - i\gamma$, $\alpha_3 = 1/2 - \delta + i\gamma$, $\alpha_4 = 1/2 - \delta - i\gamma$, and $h(s) = \prod_a (s - \alpha_a)$. Put

$$w = s - \frac{1}{2}, \quad r_a = \alpha_a - \frac{1}{2}, \quad \beta = 2(\gamma^2 - \delta^2), \quad \eta = (\delta^2 + \gamma^2)^2, \quad H(w) = w^4 + \beta w^2 + \eta. \quad (\text{RPC1})$$

Thus $r_4 = -r_1$, $r_3 = -r_2$, $r_2 = \overline{r_1}$ and $r_1^2 - r_2^2 = 4i\delta\gamma$. No root or original coordinate is discarded by this translation. Its coefficient isomorphism is $J_c : \mathbb{C}[w]_{<4} \rightarrow E_h$, $J_c p = [p(s - 1/2)]$; its inverse is substitution $s = w + 1/2$ in the unique original representative. The primitive with its original constant is

$$\Phi_h(s) = A + \phi(w), \quad \phi(w) = \frac{w^5}{5} + \frac{\beta w^3}{3} + \eta w, \quad A = \Phi_h(1/2) = \frac{1}{160} + \frac{\beta}{24} + \frac{\eta}{2}. \quad (\text{RPC2})$$

The last equality is forced by $\Phi_h(0) = 0$ and follows by evaluating the displayed odd polynomial at $w = -1/2$.

Let $u \neq 0$ carry the specified original branch of its logarithm. Let Π be the original period map on the rays and orientations of PDE2, and let $F_{ja} = \zeta^{ja} - 1$, $\zeta = e^{2\pi i/5}$, $1 \leq j, a \leq 4$. Its inverse has entries $(F^{-1})_{aj} = \zeta^{-ja}/5$: multiplying the two matrices and using $\sum_{j=1}^4 \zeta^{jb} = 4$ for $5 \mid b$ and -1 otherwise proves this entry by entry. The literal centered coefficient-to-Fourier map is

$$P = F^{-1} \Pi J_c, \quad P_0 = e^{-A/u} P. \quad (\text{RPC3})$$

These are invertible by the complete original period determinant PDE3–13. The contours for P_0 may be taken to be the rays based at $w = 0$ with the original five angles. Indeed translating the old rays gives rays based at $-1/2$. Join corresponding finite endpoints and close each contour at radius R . The integrand is entire. On the bounded-length outer joining curves its exponent is $-R^5/(5|u|) + O(R^4)$, uniformly along the joining curves, so the outer integrals tend to zero. The common finite joining path occurs with opposite coefficients in $-\ell_0 + \ell_j$ and cancels exactly. Cauchy's integral theorem therefore proves this contour identity, including its signs and the factor $e^{A/u}$.

5.1 A flat alternating pairing with all constants

On $\mathbb{C}[w]/(H)$ define the bilinear, rather than sesquilinear, pairing

$$\mathcal{J}(p, q) = \sum_{H(r)=0} \frac{p(r)q(-r)}{H'(r)}. \quad (\text{RPC4})$$

The roots are simple at the original packet. The equivalent polynomial formula is the coefficient of w^3 in the remainder of $p(w)q(-w)$ modulo H . To prove it, use Lagrange interpolation: the coefficient of w^3 in the idempotent at r is $1/H'(r)$. This polynomial formula extends the pairing to every complex value of β, η , including repeated roots, without a division by a repeated-root value. In the increasing frame $1, w, w^2, w^3$, multiplication by w and this pairing are the exact matrices

$$M = \begin{pmatrix} 0 & 0 & 0 & -\eta \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -\beta \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & \beta \\ 1 & 0 & -\beta & 0 \end{pmatrix}. \quad (\text{RPC5})$$

For the last matrix, the coefficient functional is zero through degree two, one in degree three, zero in degree four and six, and $-\beta$ in degree five. Inserting the sign $(-1)^j$ from the second argument gives every displayed entry. Thus $J^\top = -J$, $\det J = 1$, and direct multiplication gives $M^\top J + JM = 0$.

Allow the two original odd-phase coefficients to vary complexly, solely to calculate the existing period map. The ray domination $e^{-r^5/(5|u|)+O(r^3+r)}$ on compact parameter sets justifies all coefficient derivatives and integrations by parts below. Ordinary division of w^{j+3} by H , followed by differential reduction, gives

$$\partial_\eta P_0 = \frac{1}{u} P_0 M, \quad \partial_\beta P_0 = \frac{1}{3u} P_0 (M^3 - uN), \quad N = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (\text{RPC6})$$

For clarity, the quotient derivatives for columns $j = 0, 1, 2, 3$ are $0, 0, 1, 2w$, respectively; hence this is the full correction matrix. The two outer boundary terms vanish, and the two values at the common origin cancel. No differential remainder was replaced by an arithmetic remainder in RPC6.

There is also a direct nonvanishing proof on this entire two-parameter family. The displayed matrices satisfy $\text{Tr } M = 0$, $\text{Tr } M^3 = 0$ and $\text{Tr } N = 0$. The first and third follow from their diagonals; the second follows by multiplying RPC5 three times, or from the vanishing third Newton sum of the even monic polynomial H . Multilinearity of the determinant in its columns then gives $\partial_\eta \det P_0 = \partial_\beta \det P_0 = 0$ from RPC6, without first assuming an inverse exists. At the origin the literal Gamma ray integrals give the diagonal matrix G_u defined in RPC8, whose determinant is nonzero. Hence P_0 is invertible at every point of $\mathbb{C}^2$, justifying the inverses throughout this family.

The matrix identity $M^\top J + JM = 0$ also gives $(M^3)^\top J + JM^3 = 0$. Direct multiplication of RPC5–6 gives

$$\partial_\beta J + \frac{1}{3}(N^\top J + JN) = 0. \quad (\text{RPC7})$$

In particular $\partial_\beta J$ has entries $(2, 3) = 1$, $(3, 2) = -1$ when indices begin at zero, while $N^\top J + JN$ has the opposite threefold entries. Differentiate the inverse matrices in $P_0^{-\top} J P_0^{-1}$. RPC6–7 show that both its β and its η derivatives vanish. This matrix is consequently constant on the connected coefficient space $\mathbb{C}^2$.

Define the unchanged positive-Gamma ray constants

$$g_a = 5^{a/5-1} e^{\pi i a/5} \Gamma(a/5), \quad G = \text{diag}(g_1, g_2, g_3, g_4), \quad G_u = \text{diag}(u^{a/5} g_a)_{a=1}^4. \quad (\text{RPC8})$$

At $\beta = \eta = 0$, literal substitution in each original ray integral gives $P_0 = G_u$. Every g_a is nonzero, since its Gamma factor is the integral of a strictly positive function. We have therefore proved the complete flat-pairing identity

$$\begin{aligned} \Omega_0 &= P_0^{-\top} J P_0^{-1} = G_u^{-\top} J|_{\beta=0} G_u^{-1}, \\ (\Omega_0)_{14} &= -\frac{1}{u g_1 g_4}, \quad (\Omega_0)_{23} = \frac{1}{u g_2 g_3}, \quad (\Omega_0)_{ba} = -(\Omega_0)_{ab}, \\ \Omega &= P^{-\top} J P^{-1} = e^{-2A/u} \Omega_0. \end{aligned} \quad (\text{RPC9})$$

All entries not listed are zero; the indices in RPC9 begin at one. This formulation retains the Gamma constants directly and requires no unevaluated normalization of a period or source mass. For $D = \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4)$, the two paired sums of weights are five. It follows entry by entry that

$$D^\top \Omega D = \Omega, \quad D^\top \Omega_0 D = \Omega_0. \quad (\text{RPC10})$$

5.2 The full arithmetic unit and the exact quadratic form

Retain the original $v_h = 2\xi/h$ and its complete simple-root unit values. Reflection and conjugation of this original function give

$$v_h(\alpha_1) = v_h(\alpha_4) = a, \quad v_h(\alpha_2) = v_h(\alpha_3) = \bar{a}, \quad a \neq 0. \quad (\text{RPC11})$$

Indeed both numerator and denominator are invariant under $s \mapsto 1-s$ and conjugate under $s \mapsto \bar{s}$, first away from the roots and then by the removable values. The complete zero orders give nonzero values at these simple roots. Write

$$c_3 = \frac{a^{-2} - \bar{a}^{-2}}{4i\delta\gamma}, \quad c_1 = \frac{a^{-2} + \bar{a}^{-2}}{2} - (\delta^2 - \gamma^2)c_3, \quad \mathbf{r}(w) = c_1w + c_3w^3. \quad (\text{RPC12})$$

Both coefficients are real, and $(c_1, c_3) \neq (0, 0)$. Directly, $c_1 + c_3r_1^2 = a^{-2}$ and $c_1 + c_3r_2^2 = \bar{a}^{-2}$; the same equalities hold at the respective opposite roots. Thus $\mathbf{r}(M) = M v_h(M + 1/2)^{-2}$ in the original finite algebra, with the entire function evaluated by its complete original arithmetic unit. RPC12 is an exact interpolation of this specific multiplier.

For $Q_0(t) = t_1t_4 - t_2t_3$, let J_{prim} be the original Lagrange coordinate isomorphism and set

$$B = F^{-1}\Pi U J_{\text{prim}}, \quad Q(x) = Q_0(B^{-1}x), \quad \mathcal{H} = P\mathbf{r}(M)P^{-1}. \quad (\text{RPC13})$$

If $p = P^{-1}x$, the primary components of $B^{-1}x$ are exactly $p(r_1)/a, p(r_2)/\bar{a}, p(r_3)/\bar{a}, p(r_4)/a$. Since $H'(r) = 4r(r^2 + \gamma^2 - \delta^2)$, its values give $r/H'(r) = 1/(8i\delta\gamma)$ on roots one and four, and $r/H'(r) = -1/(8i\delta\gamma)$ on roots two and three. Inserting these four values into RPC4 proves the exact identity

$$Q(x) = 4i\delta\gamma p^\top \mathbf{r}(M)^\top J p = 4i\delta\gamma x^\top \mathcal{H}^\top \Omega x. \quad (\text{RPC14})$$

There is no conjugate transpose in this quadratic identity. Because $\mathbf{r}$ is odd, $\mathbf{r}(M)^\top J + J\mathbf{r}(M) = 0$; hence the matrices in RPC14 are symmetric. This also proves $\mathcal{H}^\top \Omega + \Omega \mathcal{H} = 0$. The four eigenvalues of $\mathbf{r}(M)$ are exactly $r_1/a^2, \bar{r}_1/a^2, -\bar{r}_1/a^2, -r_1/a^2$. They are nonzero, although they need not be distinct. Both B and Q_0 are invertible as linear and quadratic matrices, respectively, so Q always has rank four, including the possible eigenvalue collisions.

By RPC10, substituting Dx into RPC14 transports $\mathcal{H}$ to $D^{-1}\mathcal{H}D$. The resulting symmetric quadratic matrices are equal if and only if the corresponding $\mathcal{H}$ matrices are equal, because Ω is invertible. Consequently

$$Q(Dx) = Q(x) \iff [\mathcal{H}, D] = 0 \iff \mathcal{H}_{ab} = 0 \text{ for every } a \neq b. \quad (\text{RPC15})$$

This is an exact test in the actual original period matrix and unit. The final equivalence uses the four distinct diagonal entries of D . It makes no assumption that their off-diagonal tests are independent.

For completeness, a rank-four quadratic semi-invariant of cyclic weight $j \neq 0$ is impossible: its coefficient in row j could pair only with the missing weight zero, so that row is zero. Thus any semi-invariant rank-four quadratic must have weight zero. Since five is prime, the projective orbit of Q under D has size one or five, and size one is exactly RPC15. Every failure of any displayed off-diagonal test proves size five.

5.3 An explicit original-parameter radius forcing orbit five

Here we prove that RPC15 fails for all sufficiently large $|u|$, with a completely specified finite radius for the given original quartet and unit. The argument evaluates the original periods; it does not choose or insert a replacement amplitude.

Set $b = \beta u^{-2/5}$, $c = \eta u^{-4/5}$, and let $T(b, c)$ be the Fourier period matrix for $z^5/5 + bz^3/3 + cz$ at $u = 1$, in columns $1, z, z^2, z^3$ and on the five rays of angles $(\pi + 2\pi j)/5$. Substitution $w = u^{1/5}z$, with exactly the original branch, gives

$$P_0 = T(b, c)S_u, \quad S_u = \text{diag}(u^{a/5})_{a=1}^4, \quad S_u M S_u^{-1} = u^{1/5}M(b, c), \quad T(0, 0) = G. \quad (\text{RPC16})$$

Every column includes its original differential factor. The matrix $M(b, c)$ is RPC5 with β, η replaced by b, c ; in particular $N_0 := M(0, 0)$ has its three subdiagonal entries equal to one.

The following positive constants are explicit finite real integrals or finite expressions in the original Gamma constants:

$$\begin{aligned}
E_j &= \frac{8}{5} \int_0^\infty r^j (r^3/3 + r) e^{-r^{5/5} + r^{3/3} + r} dr \quad (0 \leq j \leq 3), \\
C_T &= 2 \left(\sum_{j=0}^3 E_j^2 \right)^{1/2}, \quad g_+ = \max_a |g_a|, \quad g_-^{-1} = \max_a |g_a|^{-1}, \\
\kappa &= \max_{1 \leq a < 4} |g_{a+1}/g_a|, \quad K_0 = GN_0 G^{-1}, \\
C_K &= 4C_T g_-^{-1} + 2g_+ g_-^{-1} + 2g_+ g_-^{-2} C_T, \\
C_{K,3} &= C_K(3\kappa^2 + 3\kappa C_K + C_K^2), \quad \epsilon_0 = \min\{1, (2C_T g_-^{-1})^{-1}\}.
\end{aligned} \tag{RPC17}$$

Here g_-^{-2} means $(\max_a |g_a|^{-1})^2$. The negative fifth degree exponent proves convergence of every integral. All norms in the following estimates are the Euclidean operator norm, with the Frobenius norm used explicitly as an upper bound where stated.

Write $\epsilon = |b| + |c|$. When $|b|, |c| \leq 1$, on each ray

$$|e^{bz^3/3+cz} - 1| \leq \epsilon(r^3/3 + r)e^{r^{3/3}+r}.$$

Each contour has two rays, and each entry of F^{-1} has absolute value $1/5$ with four summands. The entry in column j of $T - G$ is therefore at most $E_j \epsilon$. Summing the squares over all four rows proves

$$\|T - G\| \leq \|T - G\|_F \leq C_T \epsilon. \tag{RPC18}$$

For $\epsilon \leq \epsilon_0$, the convergent inverse series for $I + G^{-1}(T - G)$ gives $\|T^{-1}\| \leq 2g_-^{-1}$ and $\|T^{-1} - G^{-1}\| \leq 2g_-^{-2} C_T \epsilon$. Also $\|M(b, c) - N_0\| \leq \epsilon$, $\|M(b, c)\| \leq 2$, and $\|N_0\| = 1$. Subtract the products $K := TM(b, c)T^{-1}$ and K_0 in three terms, changing the left factor, then the middle factor, then the inverse factor. The preceding inequalities give, with the complete RPC17 constants,

$$\|K - K_0\| \leq C_K \epsilon, \quad \|K\| \leq \kappa + C_K \epsilon, \quad \|K^3 - K_0^3\| \leq C_{K,3} \epsilon. \tag{RPC19}$$

For the last assertion use the exact noncommutative identity $K^3 - K_0^3 = (K - K_0)K^2 + K_0(K - K_0)K + K_0^2(K - K_0)$. Its norm is at most $C_K \epsilon$ times $(\kappa + C_K)^2 + \kappa(\kappa + C_K) + \kappa^2$, as claimed.

The original multiplication and full unit in RPC13 now give exactly

$$\mathcal{H} = c_1 u^{1/5} K + c_3 u^{3/5} K^3, \quad (K_0^3)_{41} = g_4/g_1, \quad (K_0)_{21} = g_2/g_1. \tag{RPC20}$$

The scalar $e^{A/u}$ cancels from this conjugation, while it remains in the quadratic form through RPC9. Put $B_* = \beta + \eta > 0$. For $|u| \geq 1$, $\epsilon \leq B_* |u|^{-2/5}$.

If $c_3 \neq 0$, define the completely specified radius

$$T_* = \max \left\{ 1, \frac{B_*}{\epsilon_0}, \frac{2(|c_3| C_{K,3} B_* + |c_1|(\kappa + C_K))}{|c_3 g_4/g_1|} \right\}, \quad R_* = T_*^{5/2}. \tag{RPC21}$$

For every u on its original branch with $|u| \geq R_*$, RPC19–20 imply

$$\left| u^{-3/5} \mathcal{H}_{41} - c_3 g_4/g_1 \right| \leq |u|^{-2/5} (|c_3| C_{K,3} B_* + |c_1|(\kappa + C_K)) \leq \frac{1}{2} |c_3 g_4/g_1|. \tag{RPC22}$$

In particular $\mathcal{H}_{41} \neq 0$. In the quadratic matrix RPC14 this is the nonzero $(1, 1)$ entry $4i\delta\gamma \mathcal{H}_{41} \Omega_{41}$, whose cyclic weight is two.

If $c_3 = 0$, then $c_1 \neq 0$. In this case define

$$T_* = \max \left\{ 1, \frac{B_*}{\epsilon_0}, \frac{2C_K B_*}{|g_2/g_1|} \right\}, \quad R_* = T_*^{5/2}. \tag{RPC23}$$

For every $|u| \geq R_*$, the same estimates give

$$\left| u^{-1/5} \mathcal{H}_{21} - c_1 g_2 / g_1 \right| \leq |c_1| C_K B_* |u|^{-2/5} \leq \frac{1}{2} |c_1 g_2 / g_1|. \quad (\text{RPC24})$$

Thus $\mathcal{H}_{21} \neq 0$. The $(1, 3)$ entry of the quadratic matrix is then $4i\delta\gamma\mathcal{H}_{21}\Omega_{23}$, which has cyclic weight four; symmetry retains its $(3, 1)$ partner. Both cases prove the unconditional conclusion for the actual original parameters and the explicit radius belonging to those parameters:

$$\boxed{|u| \geq R_* \implies \#\{[Q(D^j x)] : 0 \leq j < 5\} = 5.} \quad (\text{RPC25})$$

The bracket denotes the projective class of the quadratic polynomial. RPC21 or RPC23 supplies its radius for every nonzero original unit; neither case assumes a condition on an uncomputed period coefficient.

Finally, the original periods are holomorphic on the connected logarithm cover of $\mathbb{C}^*$, and on each branch domain: locally the contours may be held in common decay sectors and differentiated under the dominated integral, and contour deformation identifies these local definitions. Their determinant never vanishes. Hence all entries of $\mathcal{H}$ are holomorphic there. RPC22 or RPC24 proves that its respective decisive entry is not identically zero on the logarithm cover and, by the identity theorem on that connected cover, on any open branch domain. The zeros of that one entry are isolated, by its convergent local power series. The exact exceptional set

$$\mathcal{E} = \{u : \mathcal{H}_{ab}(u) = 0 \text{ for all } a \neq b\} \quad (\text{RPC26})$$

is a subset of those isolated zeros and satisfies $|u| < R_*$. Thus it has no accumulation inside the branch domain; accumulation at $u = 0$ or a branch boundary has not been excluded. RPC15 is the full calculation specifying this set. No assertion that it is empty at arbitrary finite complex u is made or used.

6 A finite bound for the original kernel volume from its actual dimension

The original packet, sources, and exact coordinate map. Throughout this calculation the original multiplicity is $m = 1$ and

$$k = 4l + 1, \quad l \geq 1, \quad q = (k + 1)^2 \geq 36, \quad c = k/2, \quad 0 < \delta < 1/2, \quad \gamma > 2. \quad (\text{OKV1})$$

Retain every original root and its two coordinates:

$$\begin{aligned} \chi(S) &= \prod_{a,b=0}^k [S - c - (2a - k)\delta - i(2b - k)\gamma], \\ \omega_{ab} &= (2b - k)\gamma - i(2a - k)\delta, \\ P(y) &= i^{-q}\chi(c + iy) = \prod_{a,b=0}^k (y - \omega_{ab}). \end{aligned} \quad (\text{OKV2})$$

The identity follows factor by factor from $iy - (2a - k)\delta - i(2b - k)\gamma = i(y - \omega_{ab})$; it retains the full phase i^q . The algebra map

$$\Phi : \mathbb{C}[y]/(P) \longrightarrow E_\chi := \mathbb{C}[S]/(\chi), \quad [r(y)] \longmapsto [r((S - c)/i)] \quad (\text{OKV3})$$

is well defined because $P((S - c)/i) = i^{-q}\chi(S)$. Its inverse is $[f(S)] \mapsto [f(c + iy)]$, as verified by substitution. Both maps preserve degrees and are inverse on each polynomial ring before taking the quotients. In particular division by P is the original remainder operation through this exact coordinate map.

Use only the two original source orders $s \in \{1, k\}$ and write

$$b_s = s/2, \quad M_s = (2\pi)^{s/2}, \quad dm_{0,s}(y) = \frac{M_s 2^{b_s-2}}{\pi \Gamma(b_s)} \left| \Gamma\left(\frac{b_s}{2} + \frac{iy}{2}\right) \right|^2 dy. \quad (\text{OKV4})$$

The exact source calculation IFB2–5 supplies mass M_s , the transform $\int e^{ty} dm_{0,s} = M_s (\cos t)^{-b_s}$ for $|t| < \pi/2$, and the original monic orthogonal polynomials and norms

$$\begin{aligned} p_0 &= 1, \quad p_1 = y, \quad p_{d+1} = yp_d - d(b_s + d - 1)p_{d-1}, \\ h_d^{(s)} &= \int |p_d|^2 dm_{0,s} = M_s d!(b_s)_d, \quad \varphi_d(y) = p_d(y)/\sqrt{h_d^{(s)}}, \quad u_d(S) = \varphi_d((S - c)/i). \end{aligned} \quad (\text{OKV5})$$

All these integrals are on the same real line, with the original source mass. For example the stated orthogonality and norms also follow directly from its transform and the exact generating function

$$\sum_{d \geq 0} p_d(y) \frac{z^d}{d!} = (1 + z^2)^{-b_s/2} e^{y \arctan z}.$$

Indeed the source integral of the product at sufficiently small real z, w is $M_s(1 - zw)^{-b_s}$. The cosine addition identity and the displayed transform give this equality; the source exponential bound justifies each finite coefficient derivative under the integral. The coefficients of $z^d w^e$ then give zero for $d \neq e$ and $M_s d!(b_s)_d$ for $d = e$. The initial coefficients are $1, y$, and differentiating the generating function gives precisely the recurrence in OKV5. Thus the source norms used below retain their original constants and all finite-degree factors.

Let $C_s : \mathbb{C}^q \rightarrow E_\chi$ have columns $[u_0], \dots, [u_{q-1}]$ in the original frame $(1, S, \dots, S^{q-1})$. Its exact leading coefficients give

$$\det C_s = i^{-q(q-1)/2} \prod_{d=0}^{q-1} (h_d^{(s)})^{-1/2} \neq 0.$$

Retain the original GMB vectors and matrices, including the last row:

$$v_j = C_s^{-1}[u_{q+j}], \quad K_j = I_q + \sum_{a=0}^{j-1} v_a v_a^*, \quad 0 \leq j \leq q+1. \quad (\text{OKV6})$$

The vector v_q has original polynomial degree $2q$.

Exact remainder coefficients in all original roots. Index the full list in OKV2 by $\omega_1, \dots, \omega_q$, with each pair (a, b) occurring exactly once. Define the elementary and complete homogeneous polynomials by their finite sums

$$e_h(\omega) = \sum_{1 \leq a_1 < \dots < a_h \leq q} \omega_{a_1} \cdots \omega_{a_h}, \quad h_t(\omega) = \sum_{n_1 + \dots + n_q = t} \prod_{a=1}^q \omega_a^{n_a}, \quad (\text{OKV7})$$

with $e_0 = h_0 = 1$, $e_h = 0$ outside $0 \leq h \leq q$, and nonnegative integer n_a in the second sum. The notation $h_t(\omega)$ in this paragraph is a polynomial in the roots; the original squared norm continues to be $h_d^{(s)}$ of OKV5. Multiplication of formal power series gives the exact identity

$$\left(\sum_{h=0}^q (-1)^h e_h(\omega) z^h \right) \left(\sum_{t \geq 0} h_t(\omega) z^t \right) = 1,$$

because each geometric series $(1 - \omega_a z)^{-1}$ multiplies its own factor $1 - \omega_a z$. Consequently, for $D = q + j$ with $0 \leq j \leq q$, the quotient and remainder of y^D upon division by P are exactly

$$\begin{aligned} Q_D(y) &= \sum_{t=0}^j h_t(\omega) y^{j-t}, \quad \text{rem}_P(y^D) = y^D - P(y)Q_D(y), \\ [y^a] \text{rem}_P(y^D) &= - \sum_{t=0}^j h_t(\omega) (-1)^{D-t-a} e_{D-t-a}(\omega), \quad 0 \leq a < q. \end{aligned} \quad (\text{OKV8})$$

To verify the quotient assertion, the power-series identity makes the coefficients of $y^D, y^{D-1}, \dots, y^q$ in PQ_D equal to those of y^D . The remaining coefficients are exactly the last line of OKV8. Their degree is less than q , so uniqueness of monic polynomial division proves both assertions. For $0 \leq D < q$ the remainder is y^D itself.

The actual maximal root modulus is

$$R = k\sqrt{\delta^2 + \gamma^2}, \quad D_* := \max\{3, \sqrt{\delta^2 + \gamma^2}\}, \quad T_* := D_* q. \quad (\text{OKV9})$$

Every $|\omega_{ab}| \leq R$, and equality holds at each corner, so this expression for R is exact. In particular $R \leq T_*$. Counting the summands of OKV7 proves

$$|e_h(\omega)| \leq \binom{q}{h} R^h, \quad |h_t(\omega)| \leq \binom{q+t-1}{t} R^t.$$

The second count follows by placing $q-1$ separators among $t+q-1$ positions to encode $(n_1, \dots, n_q)$. For fixed j , the binomial $\binom{q+t-1}{t}$ is increasing in $0 \leq t \leq j$, because its successive ratio is $(q+t)/(t+1) \geq 1$. In each nonzero summand of OKV8 the total root exponent is $(D-t-a)+t = D-a$. As t varies the indices $D-t-a$ are distinct, so the sum of their $\binom{q}{D-t-a}$ is at most 2^q . It follows that

$$|[y^a] \text{rem}_P(y^{q+j})| \leq 2^q \binom{q+j-1}{j} R^{q+j-a} \leq 2^{3q-1} R^{q+j-a}, \quad 0 \leq a < q, \quad 0 \leq j \leq q. \quad (\text{OKV10})$$

For the last inequality use $\binom{q+j-1}{j} \leq 2^{q+j-1} \leq 2^{2q-1}$, by the binomial expansion of $(1+1)^{q+j-1}$. These estimates retain the exact coefficient formula OKV8, including every signed cancellation.

Original Gamma norms of the low monomials. The same recurrence and norms in OKV5 give, without changing the measure,

$$y\varphi_a = \sqrt{(a+1)(b_s+a)} \varphi_{a+1} + \sqrt{a(b_s+a-1)} \varphi_{a-1}, \quad \varphi_{-1} = 0. \quad (\text{OKV11})$$

For a polynomial of degree at most $q-2$, its orthonormal coefficient vector is therefore sent by multiplication by y to the sum of one upward and one downward weighted shift, with output degree at most $q-1$. Each shift has operator norm at most $\sqrt{q(b_s+q)}$: the squared norm of its coefficient vector is bounded by the largest squared weight times the input squared norm. Since $s \leq k < q$ gives $b_s \leq q/2$, the triangle inequality yields

$$\|yf\|_{L^2(m_{0,s})} \leq 2\sqrt{q(b_s+q)} \|f\|_{L^2(m_{0,s})} \leq 3q \|f\|_{L^2(m_{0,s})} \leq T_* \|f\|_{L^2(m_{0,s})}.$$

Here $2\sqrt{3/2} = \sqrt{6} < 3$. Starting from $\|1\|_{L^2(m_{0,s})} = \sqrt{M_s}$ and applying this inequality to $1, y, \dots, y^{a-1}$ proves

$$\boxed{\|y^a\|_{L^2(m_{0,s})} \leq \sqrt{M_s} T_*^a, \quad 0 \leq a < q.} \quad (\text{OKV12})$$

All intermediate polynomial degrees satisfy the stated operator domain.

For any polynomial $f = \sum_a f_a y^a$, write $\mathbf{c}_{T_*}(f) = \sum_a |f_a| T_*^a$. This is a finite coefficient sum, not a change of source or quotient norm. The exact original recurrence gives

$$\mathbf{c}_{T_*}(p_{d+1}) \leq T_* \mathbf{c}_{T_*}(p_d) + d(b_s + d - 1) \mathbf{c}_{T_*}(p_{d-1}).$$

For $1 \leq d < 2q$ its actual positive coefficient is bounded by $d(b_s + d - 1) \leq 2q(b_s + 2q) \leq 5q^2 \leq T_*^2$. The initial values are $1, T_*$. Induction now proves

$$\boxed{\mathbf{c}_{T_*}(p_d) \leq (2T_*)^d, \quad 0 \leq d \leq 2q.} \quad (\text{OKV13})$$

Indeed substituting the two preceding bounds in the recurrence gives $T_*(2T_*)^d + T_*^2(2T_*)^{d-1} = \frac{3}{4}(2T_*)^{d+1} \leq (2T_*)^{d+1}$.

Combining all q coefficients of OKV10 with OKV12, by the triangle inequality in the original L^2 space, gives

$$\|\text{rem}_P(y^D)\|_{L^2(m_{0,s})} \leq \sqrt{M_s} q 2^{3q} T_*^D \quad (0 \leq D \leq 2q).$$

For $D \geq q$, each summand is bounded by $\sqrt{M_s} 2^{3q-1} R^{D-a} T_*^a \leq \sqrt{M_s} 2^{3q-1} T_*^D$ and there are q summands. For $D < q$, the assertion follows from OKV12 and $q 2^{3q} \geq 1$. Linearity of the exact remainder map, followed by OKV13, proves

$$\boxed{\|\text{rem}_P(p_d)\|_{L^2(m_{0,s})} \leq \sqrt{M_s} q 2^{3q} (2T_*)^d, \quad 0 \leq d \leq 2q.} \quad (\text{OKV14})$$

The original remainder vectors and the full matrix norm. The polynomial $\text{rem}_P(p_d)$ has degree less than q . Its expansion in the orthonormal basis $\varphi_0, \dots, \varphi_{q-1}$ has squared coefficient norm equal to its squared original $L^2(m_{0,s})$ norm, by expanding the finite inner product and using OKV5. The exact quotient map OKV3 and definition OKV6 therefore give, for $d = q + j$, $0 \leq j \leq q$,

$$\|v_j\|_2^2 = \frac{\|\text{rem}_P(p_{q+j})\|_{L^2(m_{0,s})}^2}{M_s(q+j)!(b_s)_{q+j}} \leq \frac{q^2 2^{6q} (2T_*)^{2(q+j)}}{(q+j)!(b_s)_{q+j}}. \quad (\text{OKV15})$$

This equality is the actual map of the original quotient coordinates to the remainder polynomial. The full source factor M_s occurs in both the numerator bound and its original norm denominator before the displayed cancellation. No estimate on the conditioning of C_s or a companion matrix is used.

Since $b_s \geq 1/2$, for each integer $d \geq 1$ the first factor of $(b_s)_d$ is at least $1/2$ and every subsequent factor is at least one. Thus $d!(b_s)_d \geq 1/2$. Also $T_* \geq 1$ and $q + j \leq 2q$. Consequently

$$\|v_j\|_2^2 \leq 2q^2 2^{6q} (2D_*q)^{4q}, \quad 0 \leq j \leq q,$$

and the original complete matrix obeys

$$\boxed{1 \leq \|K_{q+1}\|_2 \leq 1 + 2(q+1)q^2 2^{6q} (2D_*q)^{4q}}. \quad (\text{OKV16})$$

To check its norm bound, $\|v_j v_j^*\|_2 = \|v_j\|_2^2$ follows directly from Cauchy–Schwarz, with equality on a nonzero v_j ; it is zero for a zero vector. Add all $q+1$ summands of OKV6 and the norm of I_q . Positivity also proves its lower bound.

Here is an explicit bound of the requested scale, with constants depending only on the original two root coordinates:

$$\boxed{\log \|K_{q+1}\|_2 \leq C(\delta, \gamma) q \log(q+1), \quad C(\delta, \gamma) = 12 + 4 \log(2D_*)}. \quad (\text{OKV17})$$

For a direct verification, put $U = 2(q+1)q^2 2^{6q} (2D_*q)^{4q} \geq 1$ and use $\log(1+U) \leq \log(2U)$. This upper bound is exactly

$$\log 4 + \log(q+1) + 2 \log q + 6q \log 2 + 4q \log(2D_*) + 4q \log q.$$

Write $L = \log(q+1)$. Since $q \geq 36$, one has $\log 4 \leq L$, $\log q \leq L$, $6 \log 2 \leq 2L$, and $L \geq 1$. The last inequality follows already from $\log 4 = \int_1^4 dt/t > 1$, by bounding the integrand below by $1/2$ on $[1, 2]$ and $1/4$ on $[2, 4]$. The preceding display is therefore at most $4L + 6qL + 4 \log(2D_*)qL$, which is bounded by the right side of OKV17. The completely finite and sharper bound OKV16 remains available before this logarithmic estimate.

The four original kernel Grams with their full coefficient frame. Retain the original surjection $\Lambda : E_\chi \rightarrow B_{\text{obs}}$, the actual subspace $\mathcal{K}_\Lambda = \ker \Lambda$, and its original inclusion $I : \mathcal{K}_\Lambda \hookrightarrow E_\chi$. Put $r_K = \dim \mathcal{K}_\Lambda$; no separate formula or upper bound for this actual dimension is needed here. Use its fixed original coefficient frame and write

$$T_s = C_s^{-1} I, \quad G_{q+j-1}^{(s)} = C_s^{-*} K_j^{-1} C_s^{-1}, \quad H_j^{(s)} = I^* G_{q+j-1}^{(s)} I = T_s^* K_j^{-1} T_s, \quad 0 \leq j \leq q+1. \quad (\text{OKV18})$$

For completeness, the quotient metric in this identity follows from the original orthonormal source-coordinate map $A_j = [I_q, v_0, \dots, v_{j-1}]$. It obeys $A_j A_j^* = K_j$. The minimum norm solution of $A_j z = x$ is $z = A_j^* K_j^{-1} x$: it has the required image and is orthogonal to $\ker A_j$. Every other solution differs by such a kernel vector, so the Pythagorean identity gives minimum squared norm $x^* K_j^{-1} x$. The actual remainder is $C_s x$. Substitution proves the original quotient Gram and then its full restriction in OKV18.

Let $\lambda_s = \|K_{q+1}\|_2$. The original update chain gives $I_q = K_0 \preceq K_j \preceq K_{q+1} \preceq \lambda_s I_q$. For positive definite matrices, inversion reverses the order: if $A \preceq B$, conjugate by $A^{-1/2}$ to obtain a matrix with eigenvalues at least one, invert those eigenvalues and conjugate back. Thus the full original kernel forms satisfy

$$\lambda_s^{-1} H_0^{(s)} \preceq H_j^{(s)} \preceq H_0^{(s)}, \quad H_0^{(s)} = T_s^* T_s, \quad H_{j+1}^{(s)} \preceq H_j^{(s)}. \quad (\text{OKV19})$$

For $r_K > 0$, the matrix T_s is injective, so $H_0^{(s)}$ is positive definite. Conjugation in OKV19 by its inverse square root gives eigenvalues in $[\lambda_s^{-1}, 1]$. The determinant identity for that exact congruence is

$$\det[(H_0^{(s)})^{-1/2} H_j^{(s)} (H_0^{(s)})^{-1/2}] = \frac{\det H_j^{(s)}}{\det(T_s^* T_s)}.$$

Therefore

$$-r_K \log \lambda_s \leq \log \det H_j^{(s)} - \log \det(T_s^* T_s) \leq 0. \quad (\text{OKV20})$$

In particular the determinant and conditioning of the actual $T_s^* T_s$ are retained in this identity; no new kernel basis is substituted for its original frame.

Define the original four-endpoint kernel volume, with its literal degrees $q-1, q, 2q-1, 2q$, by

$$F_K^{(s)} = \log \det H_0^{(s)} + \log \det H_1^{(s)} - \log \det H_q^{(s)} - \log \det H_{q+1}^{(s)}. \quad (\text{OKV21})$$

Monotonicity in OKV19 makes each of $\log \det H_0^{(s)} - \log \det H_q^{(s)}$ and $\log \det H_1^{(s)} - \log \det H_{q+1}^{(s)}$ nonnegative. Each is at most $r_K \log \lambda_s$ by OKV20. Consequently the actual kernel volume satisfies the complete finite estimates

$$\begin{aligned} 0 \leq F_K^{(s)} &\leq 2r_K \log \|K_{q+1}\|_2 \\ &\leq 2r_K \log [1 + 2(q+1)q^2 2^{6q} (2D_*q)^{4q}] \\ &\leq 2r_K C(\delta, \gamma) q \log(q+1), \quad s \in \{1, k\}. \end{aligned} \quad (\text{OKV22})$$

For $r_K = 0$, every kernel determinant has its original empty value one, and $F_K^{(s)} = 0$. Hence OKV22 includes this case. The four copies of $\log \det(T_s^* T_s)$ in OKV21 have coefficients $+1, +1, -1, -1$ and cancel exactly only after their presence in OKV20 has been recorded.

The original boundary and arithmetic transitions. At each original degree set

$$Q_j^{(s)} = \left(\Lambda(G_{q+j-1}^{(s)})^{-1} \Lambda^* \right)^{-1}, \quad S_j^{(s)} = (G_{q+j-1}^{(s)})^{-1} \Lambda^* Q_j^{(s)}. \quad (\text{OKV23})$$

Their zero dimensional cases have their empty meaning. Surjectivity of Λ and positivity give the displayed inverses. Direct multiplication proves $\Lambda S_j^{(s)} = I$ and $I^* G_{q+j-1}^{(s)} S_j^{(s)} = 0$. For any fixed original section S_* of Λ , the map $[I, S_*]$ is bijective: its inverse records Λz and the unique kernel coordinate of $z - S_* \Lambda z$. The minimum section differs from S_* by a kernel column. The two frames therefore differ by a block triangular matrix with identity diagonal and determinant one. Taking the determinant of the full metric in the orthogonal frame $[I, S_j^{(s)}]$ gives

$$\det H_j^{(s)} \det Q_j^{(s)} = |\det[I, S_*]|^2 \det G_{q+j-1}^{(s)}. \quad (\text{OKV24})$$

This proves the exact relation with all kernel and boundary directions retained. The same frame factor occurs at all four endpoints. Their signed sum is consequently

$$F_K^{(s)} + F_B^{(s)} = \mathcal{B}_k^{(s)}, \quad F_B^{(s)} = \log \det Q_0^{(s)} + \log \det Q_1^{(s)} - \log \det Q_q^{(s)} - \log \det Q_{q+1}^{(s)}. \quad (\text{OKV25})$$

The original matrices $G_{q+j-1}^{(s)}$ decrease with j , so their inverse forms increase. Compression by Λ and inversion then show that $Q_j^{(s)}$ decreases as well. Thus $F_B^{(s)} \geq 0$, and the finite companion bound is

$$\max\{0, \mathcal{B}_k^{(s)} - 2r_K C(\delta, \gamma) q \log(q+1)\} \leq F_B^{(s)} \leq \mathcal{B}_k^{(s)}. \quad (\text{OKV26})$$

Here the finer middle bound in OKV22 can be used in place of its last line, with the same proof.

Finally the original source identifications are $\mathcal{B}_k^{(1)} = \mathcal{B}_k^\sigma$ and $\mathcal{B}_k^{(k)} = \mathcal{B}_k^0$. Their already proved full source transition is $\mathcal{H}_k = \mathcal{B}_k^\sigma - \mathcal{B}_k^0$, and the original arithmetic transition is $\mathcal{B}_k^{\text{ar}} = \mathcal{B}_k^\sigma + \delta_k^\sigma$, with its original finite signed allowance for δ_k^σ . Substitution of the proved exact splitting OKV25 gives, with every source and arithmetic term still present,

$$\mathcal{B}_k^{\text{ar}} = F_K^{(k)} + F_B^{(k)} + \mathcal{H}_k + \delta_k^\sigma = F_K^{(1)} + F_B^{(1)} + \delta_k^\sigma. \quad (\text{OKV27})$$

Thus OKV22 controls the original Gamma kernel summands in this same arithmetic identity. It keeps the separate source transition and arithmetic scalar with their established signs and bounds. The result is finite for the actual $r_K = \dim \ker \Lambda$ and uses no additional asserted dimension estimate or replacement source.

7 The actual simple-quartet kernel and boundary at the original volume scale

This section combines the complete original period calculation, the exact cyclic-orbit rank calculation, and the finite original Gamma kernel estimate. The hypotheses are the actual simple-quartet data already used in the programme, and the period range below has an explicit bound proved from those data. No orbit condition or kernel-rank premise is added to the statement.

7.1 The original domain and the proved period range

Fix the original simple quartet $m = 1$, its coordinates $0 < \delta < 1/2$, $\gamma > 2$, and its actual nonzero amplitude value $a = v_h(1/2 + \delta + i\gamma)$, where $v_h = 2\xi/h$ and every zero order in this quartet is complete. Reflection and conjugation give the other three unit values exactly as in PQO3 and RPC11. For every integer $l \geq 1$ retain

$$k = 4l + 1, \quad q = (k + 1)^2, \quad c = k/2, \quad s \in \{1, k\}. \quad (\text{OMV1})$$

The polynomial χ , the source measures $m_{0,s}$, their masses $(2\pi)^{s/2}$, the original observation $\Lambda_k : E_\chi \rightarrow B_k$, its kernel, and all four endpoint degrees are unchanged.

Let R_* be the finite positive real number defined by RPC17 and RPC21 when the actual c_3 of RPC12 is nonzero, and by RPC17 and RPC23 when it is zero. Those two explicitly evaluated alternatives exhaust the original nonzero unit, because c_1, c_3 cannot both vanish. In particular R_* depends on the fixed quartet and its actual unit and is independent of k . Its constants are finite positive Gamma values and the convergent positive integrals RPC17. The full period proof RPC18–25 establishes

$$|u| \geq R_* \implies \#\{[Q_u(D^j x)] : 0 \leq j < 5\} = 5 \quad (\text{OMV2})$$

on every original logarithm branch. To recall the actual nonvanishing calculation, RPC22 bounds the error in $u^{-3/5}\mathcal{H}_{41}$ by half the nonzero value $|c_3g_4/g_1|$ in the first alternative. RPC24 bounds the error in $u^{-1/5}\mathcal{H}_{21}$ by half $|c_1g_2/g_1|$ in the second. Each is a nonzero off-diagonal entry of the exact $\mathcal{H} = P(c_1M + c_3M^3)P^{-1}$; the flat residue identity RPC14–15 makes it a nonzero noninvariant coefficient of the actual quadric. The degree-five cyclic action and its nondegenerate quadratic matrix then give OMV2. All powers of u use its original branch, and the absolute error estimates hold for every such branch.

The exact invariant-ideal proof OQX1–3 now applies at every point of this proved domain. It gives, for all the original degrees OMV1,

$$\dim B_k = k^2 - 6k + 17, \quad r_K := \dim \ker \Lambda_k = 8k - 16. \quad (\text{OMV3})$$

Specifically, an invariant polynomial vanishing on the original quadric is divisible by all five distinct original orbit factors; its quotient by their invariant product is invariant. Thus the actual observed rank is $d_{5,k,0} - d_{5,k-10,0}$, with negative degrees zero. The full character calculation OQX3 evaluates this difference to $k^2 - 6k + 17$, including the two original low degrees 5, 9. Subtraction from $q = (k + 1)^2$ proves the second equality. This is the dimension of the kernel of the literal period observation, not an assigned ambient dimension or a presumed maximal rank.

7.2 A uniform finite bound on the actual kernel volume

Define, from the original root coordinates only,

$$D_* = \max\{3, \sqrt{\delta^2 + \gamma^2}\}, \quad C_* = 12 + 4 \log(2D_*), \quad \mathfrak{M}_k = (16k - 32)C_*q \log(q + 1). \quad (\text{OMV4})$$

Retain the exact original source-coordinate vectors $v_j = C_s^{-1}[u_{q+j}]$, $K_j = I + \sum_{a < j} v_a v_a^*$, and the full original kernel frame $T_s = C_s^{-1}I$ of OKV6,18. The original kernel metrics at degrees $q + j - 1$

are $H_j^{(s)} = T_s^* K_j^{-1} T_s$. Their actual four-endpoint volume satisfies

$$\boxed{0 \leq F_K^{(s)}(u) \leq (16k - 32) \log[1 + 2(q + 1)q^2 2^{6q} (2D_* q)^{4q}] \leq \mathfrak{M}_k, \quad s \in \{1, k\}, \quad |u| \geq R_*} \quad (\text{OMV5})$$

Here the source index and the actual period dependence of the kernel are retained. The proof is direct substitution of the now proved $r_K = 8k - 16$ into OKV22. Its finite coefficient proof retains every original root in the signed remainder formula, all Gamma recurrence coefficients through degree $2q$, and the full source mass before its exact norm-ratio cancellation. Its metric proof retains $\det(T_s^* T_s)$ at each of the four degrees, and cancels only the four copies with coefficients $+1, +1, -1, -1$. In particular neither the conditioning nor the changing position of the original period kernel has been suppressed. The bound is uniform in the entire period domain OMV2 because its right side depends only on k, δ, γ , and the dimension in OMV3 is the same at every point of that domain.

Write the four original total quotient baselines as $\mathcal{B}_k^{(1)} = \mathcal{B}_k^\sigma$ and $\mathcal{B}_k^{(k)} = \mathcal{B}_k^0$. The full determinant identity OKV24–25, with the original fixed coefficient section, gives

$$F_B^{(s)}(u) = \mathcal{B}_k^{(s)} - F_K^{(s)}(u), \quad \max\{0, \mathcal{B}_k^{(s)} - \mathfrak{M}_k\} \leq F_B^{(s)}(u) \leq \mathcal{B}_k^{(s)}. \quad (\text{OMV6})$$

Both factors in the exact splitting are nonnegative by monotonicity of their original minimum metrics. The literal boundary term is the actual period-vector expression

$$F_B^{(s)}(u) = \log \frac{\det \mathcal{M}_{2q}^{(s)}(u) \det \mathcal{M}_{2q+1}^{(s)}(u)}{\det \mathcal{M}_q^{(s)}(u) \det \mathcal{M}_{q+1}^{(s)}(u)}, \quad \mathcal{M}_D^{(s)}(u) = \sum_{d=0}^{D-1} \lambda_d^{(s)}(u) (\lambda_d^{(s)}(u))^*. \quad (\text{OMV7})$$

The vectors are the complete original primary-period coefficients OPG14 and OCS21 on the actual image of Λ_k . Formula OPG17 proves that each consecutive determinant ratio is $1 + \xi_j$; the unchanged weights $1, 2, \dots, 2, 1$ give exactly OMV7. The companion kernel vectors retain the minimum-section correction OPR2 and the action defect OPG22/OQC22. No invariance of that kernel under the original sum action is needed for OMV5–7.

7.3 The two individual leading coefficients

The complete original baseline proofs BSL16–19 and HBL establish, for fixed original δ, γ and $m = 1$, both source limits

$$\frac{\mathcal{B}_k^{(s)}}{q^2} \longrightarrow C_B = 9 - 8 \log 2 + F(2, \pi) > \frac{769}{17010}, \quad s = 1 \text{ and } s = k. \quad (\text{OMV8})$$

Here $F(2, \pi)$ is exactly the energy functional evaluated in those complete proofs; its original end-point convention and coefficients are retained. No value of it is redefined by the present comparison. Since $q = (k + 1)^2$,

$$\frac{\mathfrak{M}_k}{q^2} = C_* \frac{(16k - 32) \log((k + 1)^2 + 1)}{(k + 1)^2} \longrightarrow 0.$$

For an elementary verification, $((k + 1)^2 + 1) < (k + 2)^2$, and the last fraction is at most $32 \log(k + 2)/k$. The inequality $\log x = \int_1^x dt/t \leq 2(\sqrt{x} - 1)$ for $x \geq 1$ proves that this upper bound tends to zero. The finite estimate OMV5 and the exact identity OMV6 therefore prove

$$\boxed{\frac{F_K^{(1)}(u)}{q^2}, \frac{F_K^{(k)}(u)}{q^2} \longrightarrow 0, \quad \frac{F_B^{(1)}(u)}{q^2}, \frac{F_B^{(k)}(u)}{q^2} \longrightarrow C_B} \quad (\text{OMV9})$$

These limits are uniform for all $|u| \geq R_*$ and all original branches. Indeed the kernel error is bounded by $\mathfrak{M}_k/q^2$ uniformly, while the boundary error is at most $|\mathcal{B}_k^{(s)}/q^2 - C_B| + \mathfrak{M}_k/q^2$. The baseline is independent of u . This proves the asserted uniformity, rather than only a conclusion for each separately fixed observation. The original combined coefficient has now been resolved into its two individual contributions on the proved simple-quartet period domain.

7.4 Exact propagation to the arithmetic identity and the finer scale

Keep the original signed transitions $\mathcal{H}_k = \mathcal{B}_k^\sigma - \mathcal{B}_k^0$ and $\delta_k^\sigma = \mathcal{B}_k^{\text{ar}} - \mathcal{B}_k^\sigma$. Their complete finite allowances, including the QGQ simple- multiplicity refinement, remain those of the full current receiver. The exact arithmetic receiving equality becomes

$$\begin{aligned}\mathcal{B}_k^{\text{ar}} - F_B^{(1)}(u) &= F_K^{(1)}(u) + \delta_k^\sigma, \\ \mathcal{B}_k^{\text{ar}} - F_B^{(k)}(u) &= F_K^{(k)}(u) + \mathcal{H}_k + \delta_k^\sigma.\end{aligned}\tag{OMV10}$$

This is substitution in OKV27, so every source and sign is the original one. In particular the finite upper and lower allowances are obtained by adding $[0, \mathfrak{M}_k]$ to the actual interval for δ_k^σ , or to the actual interval for $\mathcal{H}_k + \delta_k^\sigma$, respectively. The original proved relations $\delta_k^\sigma/(kq) \rightarrow 0$ and $\mathcal{H}_k/(kq) \rightarrow -\mathcal{C}_\Gamma/4$, together with $k/q \rightarrow 0$ and OMV5, prove that both differences in OMV10, divided by q^2 , tend to zero uniformly on the stated period domain. This is an exact comparison with the full arithmetic baseline. The literal arithmetic kernel and boundary metrics retain their own source until their corresponding metric comparison is applied.

The finer original Gamma return is still carried by an exact mixed identity:

$$\begin{aligned}F_B^{(1)}(u) - F_B^{(k)}(u) &= \mathcal{H}_k - (F_K^{(1)}(u) - F_K^{(k)}(u)), \\ -\mathfrak{M}_k &\leq F_K^{(1)}(u) - F_K^{(k)}(u) \leq \mathfrak{M}_k, \\ \mathcal{H}_k - \mathfrak{M}_k &\leq F_B^{(1)}(u) - F_B^{(k)}(u) \leq \mathcal{H}_k + \mathfrak{M}_k.\end{aligned}\tag{OMV11}$$

Subtract OMV6 for the two actual source orders to obtain the first line. Each kernel term lies in $[0, \mathfrak{M}_k]$, so their difference lies in the stated interval, proving the other two lines. The error bound here is of order $kq \log(q+1)$, whereas the already proved signed return $\mathcal{H}_k$ has order kq . No sign or limiting coefficient for either finer mixed difference is assigned by OMV11. Its explicit remaining term is the difference of the two original kernel metrics, on the exactly calculated $8k-16$ dimensional period kernel. The full source multiplier and its action on this actual kernel remain the input to that next calculation, rather than a freely selected source order or kernel.

8 The original arithmetic kernel and boundary receive the full source comparison

This calculation uses the proved full-polynomial comparison TW12/TWA3, the original observation OPG1–5, and the complete original Gamma kernel estimate OKV1–27. It gives the exact maps between their source, quotient, kernel and boundary metrics, before taking any determinant. The subsequent simple-quartet conclusion uses the evaluated original period test RPC1–26 and its exact rank calculation OQX1–3. Every source order below is one of the already present orders $1, k$.

8.1 The complete original source and the same observation

Retain the original full-order quartet, with $0 < \delta < 1/2$, $\gamma > 2$, and positive integer complete zero order m . Thus

$$\begin{aligned} \mathcal{R} &= \{\tfrac{1}{2} + \delta + i\gamma, \tfrac{1}{2} + \delta - i\gamma, \tfrac{1}{2} - \delta + i\gamma, \tfrac{1}{2} - \delta - i\gamma\}, \quad h(z) = \prod_{\alpha \in \mathcal{R}} (z - \alpha)^m, \quad v_h = 2\xi/h, \\ k &\geq 3, \quad c = k/2, \quad e = 1 + k(m-1), \quad q = e(k+1)^2, \\ \lambda_{a,b} &= c + (2a - k)\delta + i(2b - k)\gamma, \quad \chi(S) = \prod_{a,b=0}^k (S - \lambda_{a,b})^e, \quad E = \mathbb{C}[S]/(\chi), \\ w_h(y) &= |v_h(1/2 + iy)|^2/(2\pi), \quad m_{h,k} = w_h^{*k}, \quad d\sigma(y) = |\Gamma(1/4 + iy/2)|^2 dy/(2\pi). \end{aligned} \quad (\text{AMT1})$$

The original source inner products, conjugate-linear in their first argument, are

$$\langle P, Q \rangle_{\text{ar}} = \int_{\mathbb{R}} \overline{P(c + iy)} Q(c + iy) m_{h,k}(y) dy, \quad \langle P, Q \rangle_{\sigma} = \int_{\mathbb{R}} \overline{P(c + iy)} Q(c + iy) d\sigma(y). \quad (\text{AMT2})$$

No measure mass or complex-coordinate phase is changed. In particular $\int d\sigma = \sqrt{2\pi}$. Write $\mathcal{P}_N$ for polynomials in the original S of degree at most N . In its coefficient frame $(1, S, \dots, S^N)$ let the two Grams be $H_N^{\text{ar}}, H_N^{\sigma}$. For $N \geq q - 1$, the literal remainder map $J_N : \mathcal{P}_N \rightarrow E$ is onto and has kernel $\chi \mathcal{P}_{N-q}$; at $N = q - 1$ this is the zero vector space. The frame on E is $(1, S, \dots, S^{q-1})$.

Here is the original observation being retained. Put $n = 4m$, $E_h = \mathbb{C}[z]/(h)$, $U = M_{j_h v_h}$, and $\iota[P] = [P(z_1 + \dots + z_k)] \in E_h^{\otimes k}$. The map j_h records every divided Taylor coefficient through order $m - 1$ at every original root. Its unit $j_h v_h$ is invertible because the stated order is complete. Fix the original $u \neq 0$ and its logarithm branch. With $\Phi'_h = h$ and $\Phi_h(0) = 0$, let ℓ_j be the outward ray of angle $(\arg u + \pi + 2\pi j)/(n + 1)$ and $\Gamma_j = -\ell_0 + \ell_j$. The period map and projector are exactly

$$\begin{aligned} (\Pi[P])_j &= \int_{\Gamma_j} e^{\Phi_h(z)/u} \text{rem}_h P(z) dz, \quad 1 \leq j \leq n, \\ C : \Gamma_j &\mapsto \Gamma_{j+1} - \Gamma_1 \quad (j < n), \quad \Gamma_n \mapsto -\Gamma_1, \quad P_k = \frac{1}{n+1} \sum_{a=0}^n (C^{-a})^{\otimes k}, \end{aligned} \quad (\text{AMT3})$$

$$L = P_k \Pi^{\otimes k} U^{\otimes k} \iota, \quad B = \text{im } L, \quad j : B \hookrightarrow (\mathbb{C}^n)^{\otimes k}, \quad j\Lambda = L, \quad \Lambda : E \twoheadrightarrow B, \quad K = \ker \Lambda.$$

These are the convergent original representative integrals; their existence and full determinant are proved in PDE and OPG. Fix the original coefficient inclusion $I : K \hookrightarrow E$ and section $S_* : B \rightarrow E$ with $\Lambda S_* = I_B$. Let $r = \dim K$ and $b = \dim B = q - r$. All matrices below use these same frames and this same full unit and period observation. Empty Grams have determinant one, and maps to or from a zero space have their unique empty values.

8.2 All constants of the actual full-polynomial inequality

Retain exactly the constants of TW and TWA:

$$\begin{aligned} \alpha &= \pi/2, \quad p = 8m - \frac{9}{2}, \quad B_h = 42 + 8m, \quad M = 21 + 4m, \\ \vartheta_h &= \int_{-1}^1 w_h(y) dy > 0, \quad M_\alpha = \int_{\mathbb{R}} e^{\alpha y} w_h(y) dy \in (0, \infty), \\ c_\Gamma &= \sqrt{2}e^{-7/6}, \quad C_\Gamma = \sqrt{2\sqrt{5}}e^{2/3}, \\ a_k &= \frac{c_h \vartheta_h^{k-3} e^{-\alpha(k-3)} (k-2)^{-B_h}}{C_\Gamma 2^{B_h/2}}, \quad A_k = \frac{C_h^{\text{tilt}}}{c_\Gamma} k^{p+1} M_\alpha^{k-1}, \quad D_N = [1 + 4(N+M)^2]^M. \end{aligned} \tag{AMT4}$$

Here c_h is the proved original convolution lower-envelope constant TW7 and C_h^{tilt} is the original compact/tail maximum BT12. Their full constructions occur in the preserved arithmetic endpoint and BT providers; they depend on this fixed h and not on k, N, u . Their defining inequalities, proved there with the original numerator, are

$$\begin{aligned} m_{h,k}(y) &\geq c_h \vartheta_h^{k-3} e^{-\alpha(|y|+k-3)} (k-2+|y|)^{-B_h}, \\ m_{h,k}(y) &\leq k C_h^{\text{tilt}} M_\alpha^{k-1} e^{-\alpha|y|} (1+|y|/k)^{-p}, \\ c_\Gamma e^{-\alpha|y|} (1+|y|)^{-1/2} &\leq d\sigma/dy \leq C_\Gamma e^{-\alpha|y|} (1+|y|)^{-1/2}. \end{aligned} \tag{AMT5}$$

For clarity the full polynomial step is given here. The first and third bounds, $k-2+|y| \leq (k-2)(1+|y|)$, and $(1+|y|)^{B_h} \leq 2^{B_h/2}(1+y^2)^M$, give $m_{h,k} \geq a_k(1+y^2)^{-M} d\sigma/dy$. The second and third, using $1+|y|/k \geq (1+|y|)/k$ and $p > 1/2$, give $m_{h,k} \leq A_k d\sigma/dy$. The fixed Gamma orthonormal polynomials obey

$$y e_j = \sqrt{(j+1)(j+1/2)} e_{j+1} + \sqrt{j(j-1/2)} e_{j-1}.$$

The two shifts on degree at most N have norms at most $N+1$ and N , respectively. Thus $\|yP\|_\sigma \leq 2(N+1)\|P\|_\sigma$. Iterating at the successively increased degrees and expanding $(1+y^2)^M$ proves $\int |P|^2 (1+y^2)^M d\sigma \leq D_N \int |P|^2 d\sigma$. Cauchy–Schwarz applied to $|P|(1+y^2)^{-M/2}$ and $|P|(1+y^2)^{M/2}$ then gives the reciprocal lower bound. The coordinate map $P(S) \mapsto P(c+iy)$ has inverse $f(y) \mapsto f((S-c)/i)$, preserves the degree and retains every coefficient through this explicitly invertible substitution. It therefore proves on the complete original source

$$\boxed{\frac{a_k}{D_N} H_N^\sigma \preceq H_N^{\text{ar}} \preceq A_k H_N^\sigma.} \tag{AMT6}$$

All coefficients can be complex in this proof. In particular AMT6 holds on every affine remainder fibre, not only on monic vectors. The two Grams are positive: the Gamma density is positive and the lower bound in AMT5 is positive everywhere. Their finite moments exist by the exponential upper bound.

8.3 Exact remainder, minimum-section, kernel and boundary maps

For $a \in \{\sigma, \text{ar}\}$ put

$$\begin{aligned} G_N^a &= (J_N (H_N^a)^{-1} J_N^*)^{-1}, \quad V_N^a = (H_N^a)^{-1} J_N^* G_N^a : E \longrightarrow \mathcal{P}_N, \\ H_{K,N}^a &= I^* G_N^a I, \quad Q_N^a = (\Lambda (G_N^a)^{-1} \Lambda^*)^{-1}, \quad S_N^a = (G_N^a)^{-1} \Lambda^* Q_N^a : B \longrightarrow E. \end{aligned} \tag{AMT7}$$

Inverses are on the indicated nonzero spaces. For an onto matrix J and positive H , direct multiplication gives $JV = I$ for $V = H^{-1} J^* (JH^{-1} J^*)^{-1}$. Moreover $Z^* H V = Z^* J^* G = 0$ for $JZ = 0$. Therefore every lift of z is $Vz + Z$, with squared norm $z^* G z + Z^* H Z$, proving its unique minimum and its Gram. Applying this proof first to J_N , then to Λ , proves every formula AMT7 and $\Lambda S_N^a = I_B$, $I^* G_N^a S_N^a = 0$. It also proves that $V_N^a S_N^a$ is the minimum lift for the composite ΛJ_N .

The exact comparison map on $\mathcal{P}_N$ is the identity on original coefficients; it commutes with J_N and ΛJ_N . It induces the identity on E, K, B . Between the two spaces of minimum lifts the exact bijection and its inverse are

$$\text{im } V_N^\sigma \longrightarrow \text{im } V_N^{\text{ar}}, \quad V_N^\sigma z \longmapsto V_N^{\text{ar}} z, \quad T_N = V_N^{\text{ar}} J_N|_{\text{im } V_N^\sigma}, \quad T_N^{-1} = V_N^\sigma J_N|_{\text{im } V_N^{\text{ar}}}. \quad (\text{AMT8})$$

Their difference $V_N^{\text{ar}} z - V_N^\sigma z$ lies in $\chi \mathcal{P}_{N-q}$, with its unique coefficient obtained by division by the unchanged monic χ . This also gives the minimum-lift bijection on the preimage of K and the analogous bijection $V_N^\sigma S_N^\sigma y \mapsto V_N^{\text{ar}} S_N^{\text{ar}} y$ for boundary lifts. The latter differences lie in $\ker(\Lambda J_N)$, exactly as multiplication by ΛJ_N verifies.

Minimize both sides of AMT6 over $J_N P = z$. Taking infima preserves each inequality because these are the same nonempty fibres. Restrict the resulting inequality to $z = Ix$, and independently minimize it over $\Lambda z = y$. This proves the three full-vector comparisons

$$\begin{aligned} (a_k/D_N) G_N^\sigma &\preceq G_N^{\text{ar}} \preceq A_k G_N^\sigma, \\ (a_k/D_N) H_{K,N}^\sigma &\preceq H_{K,N}^{\text{ar}} \preceq A_k H_{K,N}^\sigma, \\ (a_k/D_N) Q_N^\sigma &\preceq Q_N^{\text{ar}} \preceq A_k Q_N^\sigma. \end{aligned} \quad (\text{AMT9})$$

The source identity is an exact bounded bijection with the two constants in AMT6. No invariance of K under an additional multiplier was used to obtain its induced identities.

The matrix $[I, S_*]$ is invertible: its inverse sends z to the unique kernel coordinate of $z - S_* \Lambda z$ and the boundary coordinate Λz . Each $S_N^a - S_*$ lies in K , so the change from this frame to $[I, S_N^a]$ is triangular with identity diagonal and determinant one. The latter frame is orthogonal in G_N^a by AMT7. Consequently

$$\det H_{K,N}^a \det Q_N^a = |\det[I, S_*]|^2 \det G_N^a, \quad a \in \{\sigma, \text{ar}\}. \quad (\text{AMT10})$$

This includes $r = 0$ and $b = 0$ and retains the original frame factor.

8.4 The exact nested deficits and their rank factors

Define the actual finite source determinant deficit and its width

$$\Xi_{k,N} = (N+1) \log A_k - \log \frac{\det H_N^{\text{ar}}}{\det H_N^\sigma}, \quad L_N = \log(A_k D_N / a_k) \geq 0. \quad (\text{AMT11})$$

The last sign follows by applying AMT6 to any nonzero vector. Put

$$\begin{aligned} d_{E,N} &= q \log A_k - \log \frac{\det G_N^{\text{ar}}}{\det G_N^\sigma}, \\ d_{K,N} &= r \log A_k - \log \frac{\det H_{K,N}^{\text{ar}}}{\det H_{K,N}^\sigma}, \\ d_{B,N} &= b \log A_k - \log \frac{\det Q_N^{\text{ar}}}{\det Q_N^\sigma}. \end{aligned} \quad (\text{AMT12})$$

AMT9, by congruence with the inverse square root of each comparison Gram, places each eigenvalue of its relative Gram in $[a_k/D_N, A_k]$. Multiplication of all eigenvalues gives $0 \leq d_{K,N} \leq r L_N$, $0 \leq d_{B,N} \leq b L_N$ and $0 \leq d_{E,N} \leq q L_N$.

Here is the full source-to-quotient determinant calculation, including all relation directions. Choose an H_N^σ -orthonormal coefficient frame R_N of $\ker J_N$ solely for this computation, and put $W_N = [R_N, V_N^\sigma (G_N^\sigma)^{-1/2}]$. AMT7 gives $W_N^* H_N^{\text{ar}} W_N = I_{N+1}$. In this frame the positive contraction

$$\mathcal{D}_N = A_k^{-1} W_N^* H_N^{\text{ar}} W_N = \begin{pmatrix} F_N & C'_N \\ (C'_N)^* & A'_N \end{pmatrix} \preceq I_{N+1}$$

has Schur complement

$$T'_N = A'_N - (C'_N)^* F_N^{-1} C'_N = A_k^{-1} (G_N^\sigma)^{-1/2} G_N^{\text{ar}} (G_N^\sigma)^{-1/2}.$$

Indeed minimization in its first block is exactly the original remainder minimum at $z = (G_N^\sigma)^{-1/2} w$. Both F_N and T'_N are positive contractions: positivity follows from positive block elimination, and the upper bound for the Schur complement follows by testing its minimum at zero first coordinate. Since $|\det W_N|^2 = (\det H_N^\sigma)^{-1}$, elimination gives

$$\Xi_{k,N} = d_{R,N} + d_{E,N}, \quad d_{R,N} := -\log \det F_N \geq 0, \quad d_{E,N} = -\log \det T'_N \geq 0. \quad (\text{AMT13})$$

When $N = q - 1$, F_N is empty, $d_{R,N} = 0$, and the same equalities hold without a relation inverse. The determinant identities are independent of the auxiliary orthonormal frame. Dividing the two instances of AMT10 proves $d_{K,N} + d_{B,N} = d_{E,N}$. Thus the complete nested statement is

$$\boxed{\begin{aligned} \Xi_{k,N} &= d_{R,N} + d_{K,N} + d_{B,N}, & d_{R,N}, d_{K,N}, d_{B,N} &\geq 0, \\ 0 &\leq d_{K,N} \leq \min\{rL_N, d_{E,N}\} \leq \min\{rL_N, \Xi_{k,N}\}, \\ 0 &\leq d_{B,N} \leq \min\{bL_N, d_{E,N}\} \leq \min\{bL_N, \Xi_{k,N}\}. \end{aligned}} \quad (\text{AMT14})$$

In particular each metric keeps its own rank factor, and the two deficits retain their exact common sum. Any proved finite upper bound for the original $\Xi_{k,N}$, including the degree-specific HMR44/TWA15 bound, can be substituted in these displayed minima. No new density or source determinant is introduced by this substitution.

8.5 All four original endpoints and the correlated mixed return

Let

$$(N_0, N_1, N_2, N_3) = (q - 1, q, 2q - 1, 2q), \quad (\epsilon_0, \epsilon_1, \epsilon_2, \epsilon_3) = (1, 1, -1, -1),$$

and define, in their original signs,

$$\begin{aligned} F_K^a &= \sum_{j=0}^3 \epsilon_j \log \det H_{K,N_j}^a, & F_B^a &= \sum_{j=0}^3 \epsilon_j \log \det Q_{N_j}^a, \\ \mathcal{B}_k^a &= \sum_{j=0}^3 \epsilon_j \log \det G_{N_j}^a, & \Delta_K^\sigma &= F_K^{\text{ar}} - F_K^\sigma, \quad \Delta_B^\sigma = F_B^{\text{ar}} - F_B^\sigma, \quad \delta_k^\sigma = \mathcal{B}_k^{\text{ar}} - \mathcal{B}_k^\sigma. \end{aligned} \quad (\text{AMT15})$$

AMT10 cancels its four identical frame factors only after these signs are applied. It gives $F_K^a + F_B^a = \mathcal{B}_k^a$ and $\Delta_K^\sigma + \Delta_B^\sigma = \delta_k^\sigma$ exactly. The spaces $\mathcal{P}_N$ increase with the same source norm and remainder map, so their quotient minima decrease. Restriction to K and minimization onto B preserve that order. Pairing N_0 with N_2 and N_1 with N_3 consequently proves $F_K^a, F_B^a \geq 0$.

For $X = K, B$, put $r_K = r$, $r_B = b$, and retain the exact caps

$$c_{X,N} = \min\{r_X L_N, d_{E,N}\}, \quad C_X^- = c_{X,q-1} + c_{X,q}, \quad C_X^+ = c_{X,2q-1} + c_{X,2q}. \quad (\text{AMT16})$$

The entirely source-evaluated caps $\hat{c}_{X,N} = \min\{r_X L_N, \Xi_{k,N}\}$ and the corresponding $\hat{C}_X^\pm$ are at least these values; all conclusions below also hold with the hats. Substituting AMT12 in AMT15 records each endpoint and its sign:

$$\boxed{\begin{aligned} \Delta_K^\sigma &= -d_{K,q-1} - d_{K,q} + d_{K,2q-1} + d_{K,2q}, \\ \Delta_B^\sigma &= -d_{B,q-1} - d_{B,q} + d_{B,2q-1} + d_{B,2q}, \\ \delta_k^\sigma &= -d_{E,q-1} - d_{E,q} + d_{E,2q-1} + d_{E,2q}, \\ -C_X^- &\leq \Delta_X^\sigma \leq C_X^+, \quad X = K, B. \end{aligned}} \quad (\text{AMT17})$$

The four copies of $r_X \log A_k$ cancel here because their signs sum to zero. Since the original total correction is retained, the joint interval is stronger than two independent intervals:

$$\boxed{\begin{aligned} \max\{-C_K^-, \delta_k^\sigma - C_B^+\} &\leq \Delta_K^\sigma \leq \min\{C_K^+, \delta_k^\sigma + C_B^-\}, \\ \max\{-C_B^-, \delta_k^\sigma - C_K^+\} &\leq \Delta_B^\sigma \leq \min\{C_B^+, \delta_k^\sigma + C_K^-\}. \end{aligned}} \quad (\text{AMT18})$$

These follow by subtracting the interval for either component from the exact common sum; they require no independence assertion.

Here are the explicit original four endpoint widths:

$$\begin{aligned} E_k^+ &= L_{q-1} + L_q = 2 \log(A_k/a_k) + M \log[1 + 4(q-1+M)^2] + M \log[1 + 4(q+M)^2], \\ E_k^- &= L_{2q-1} + L_{2q} = 2 \log(A_k/a_k) + M \log[1 + 4(2q-1+M)^2] + M \log[1 + 4(2q+M)^2]. \end{aligned} \quad (\text{AMT19})$$

They are the unchanged TWA5 constants, both nonnegative. AMT17 therefore gives in particular

$$\boxed{-rE_k^+ \leq F_K^{\text{ar}} - F_K^\sigma \leq rE_k^-, \quad -bE_k^+ \leq F_B^{\text{ar}} - F_B^\sigma \leq bE_k^-}. \quad (\text{AMT20})$$

The total correction retains TWA15 with all four individual source deficits:

$$\begin{aligned} \underline{\delta}_k^\sigma &= \max\{-qE_k^+, -\Xi_{k,q-1} - \Xi_{k,q}\}, \\ \bar{\delta}_k^\sigma &= \min\{qE_k^-, \Xi_{k,2q-1} + \Xi_{k,2q}\}, \quad \underline{\delta}_k^\sigma \leq \delta_k^\sigma \leq \bar{\delta}_k^\sigma. \end{aligned} \quad (\text{AMT21})$$

This also follows immediately from AMT14 and AMT17. It can be inserted into AMT18 when an interval, rather than the actual finite total determinant, is wanted. Thus the lower and upper endpoints are kept in their distinct places throughout the mixed transfer.

For later use, subtraction of the exact logarithms in AMT4 gives

$$\begin{aligned} \mathfrak{s}_h &= \alpha + \log(M_\alpha/\vartheta_h), \quad \mathfrak{c}_h = -\log M_\alpha + 3 \log \vartheta_h - 3\alpha + \log \frac{C_h^{\text{tilt}} C_\Gamma 2^{B_h/2}}{c_h c_\Gamma}, \\ \log(A_k/a_k) &= k\mathfrak{s}_h + (p+1) \log k + B_h \log(k-2) + \mathfrak{c}_h, \quad E_k^\pm = O_h(k + \log q). \end{aligned} \quad (\text{AMT22})$$

The last statement follows from the displayed exact expressions, since M, p, B_h and the two integral constants are fixed and $\log D_N = O_h(1 + \log(N+1))$. It is a consequence of the original constants and contains no change of measure.

8.6 The full original multiplier and its transported observation

On the existing tensor class $k = 4l + 1$, retain

$$\begin{aligned} t_j &= 2j + \frac{1}{2}, \quad \xi_j = c + t_j, \quad f_l(S) = \prod_{j=0}^{l-1} (S - c - t_j), \\ \beta_k &= \frac{(2\pi)^{k/2}}{\sqrt{2}\Gamma(k/2)}, \quad dm_{0,k}(y) = \beta_k |f_l(c + iy)|^2 d\sigma(y). \end{aligned} \quad (\text{AMT23})$$

This is the exact Gamma recurrence identity TWA26 with masses $(2\pi)^{k/2}$ and $\sqrt{2\pi}$. For $l = 0$ both f_l and β_k are one. Write $E_\chi : E \rightarrow \mathbb{C}^q$ for the complete divided Taylor map $[P] \mapsto (P^{(r)}(\lambda_{a,b})/r!)_{a,b, 0 \leq r < e}$. It is invertible: a vector in its kernel is divisible by the coprime factors of χ , and dimensions agree. The full multiplication matrix U_f has, in each primary block, entries

$$\begin{aligned} (U_f)_{r,a} &= \begin{cases} f_l^{(r-a)}(\lambda)/(r-a)!, & 0 \leq a \leq r < e, \\ 0, & a > r. \end{cases} \\ \det U_f &= \prod_{a,b=0}^k \prod_{j=0}^{l-1} [(2a-k)\delta - t_j + i(2b-k)\gamma]^e \neq 0. \end{aligned} \quad (\text{AMT24})$$

Indeed k is odd, so every $\lambda_{a,b}$ has nonzero imaginary part, while all ξ_j are real and distinct. The inverse in each block is the finite Taylor inverse of its nonzero constant term. The exact polynomial map and its inverse on its stated range are

$$\mathcal{P}_N \longrightarrow f_l \mathcal{P}_N \subseteq \mathcal{P}_{N+l}, \quad P \longmapsto f_l P, \quad Q \longmapsto Q/f_l. \quad (\text{AMT25})$$

It sends the original fibre $J_N P = z$ bijectively to the product jet fibre $(\mathbf{E}_f Q, \mathbf{E}_\chi Q) = (0, U_f \mathbf{E}_\chi z)$. Vanishing of the l distinct f -values proves divisibility by f_l , and the inverse of U_f proves the converse fibre statement. The original relation space maps to $f_l \chi \mathcal{P}_{N-q}$. On the free χ jet coordinate Y of this constrained product fibre, the exact observed map, its kernel, and a section are

$$\begin{aligned} \tilde{\Lambda} &= \Lambda \mathbf{E}_\chi^{-1} U_f^{-1}, & \tilde{\Lambda} U_f \mathbf{E}_\chi &= \Lambda, \\ \ker \tilde{\Lambda} &= U_f \mathbf{E}_\chi K, & \tilde{I} &= U_f \mathbf{E}_\chi I, & \tilde{S}_* &= U_f \mathbf{E}_\chi S_*, & \tilde{\Lambda} \tilde{S}_* &= I_B. \end{aligned} \quad (\text{AMT26})$$

Both inclusions in the kernel equality follow by applying the displayed invertible map and its inverse. This retains the full original amplitude in Λ and every derivative in U_f . The resulting kernel is proved by transport; it has not been assigned the coordinates of $\mathbf{E}_\chi K$.

For an explicit metric on this transported space, let $b_n(y)$ be the fixed monic Gamma polynomials and put $\pi_n(S) = i^n b_n((S - c)/i)$, whose leading coefficient is one and squared norm is $\gamma_n = \sqrt{2\pi}(2n)!/4^n$. The evaluation kernels are the full finite sums

$$\mathcal{K}_{\chi,L} = \sum_{n=0}^L \frac{\mathbf{E}_\chi \pi_n (\mathbf{E}_\chi \pi_n)^*}{\gamma_n}, \quad \mathcal{K}_{f,L} = \sum_{n=0}^L \frac{\mathbf{E}_f \pi_n (\mathbf{E}_f \pi_n)^*}{\gamma_n}, \quad \mathcal{K}_{\chi,f;L} = \sum_{n=0}^L \frac{\mathbf{E}_\chi \pi_n (\mathbf{E}_f \pi_n)^*}{\gamma_n}.$$

The product evaluation is onto for $L = N + l \geq q + l - 1$, because a polynomial of degree below $q + l$ vanishing at all its divided jets is divisible by the degree- $q + l$ product $f_l \chi$ and is zero. Thus its positive kernel has positive Schur complement

$$\begin{aligned} C_N &= \mathcal{K}_{\chi,N+l} - \mathcal{K}_{\chi,f;N+l} \mathcal{K}_{f,N+l}^{-1} \mathcal{K}_{f,\chi;N+l} > 0, \\ U_N &= (\mathbf{E}_\chi \pi_{N+j} / \sqrt{\gamma_{N+j}})_{j=1}^l, \quad V_N = \mathcal{K}_{\chi,f;N+l} \mathcal{K}_{f,N+l}^{-1/2}, \quad C_N = \mathcal{K}_{\chi,N} + U_N U_N^* - V_N V_N^*. \end{aligned} \quad (\text{AMT27})$$

Empty f blocks have the identity convention. The inverse of the joint positive kernel, evaluated on $(0, Y)$, gives the minimum $Y^* C_N^{-1} Y$: block elimination of the f -block proves this identity with every mixed entry present. Combining it with AMT25 and the coefficient β_k proves the exact tensor-source metrics

$$\begin{aligned} G_N^0 &= \beta_k \mathbf{E}_\chi^* U_f^* C_N^{-1} U_f \mathbf{E}_\chi, & H_{K,N}^0 &= \beta_k \tilde{I}^* C_N^{-1} \tilde{I}, \\ Q_N^0 &= \beta_k (\tilde{\Lambda} C_N \tilde{\Lambda}^*)^{-1}, & S_N^0 &= \mathbf{E}_\chi^{-1} U_f^{-1} C_N \tilde{\Lambda}^* (\tilde{\Lambda} C_N \tilde{\Lambda}^*)^{-1}. \end{aligned} \quad (\text{AMT28})$$

Direct multiplication verifies the section and its orthogonality. The formulas hold on the original E, K, B , with the transported observation exactly as in AMT26. No block or mixed product was dropped from C_N .

The same coefficient identity also supplies a quantitative comparison directly with this original tensor source. Define the positive exact finite factors

$$\Pi_l = \prod_{j=0}^{l-1} t_j^2, \quad U_{l,N}^{\text{poly}} = \prod_{j=0}^{l-1} [4(N+j+1)^2 + t_j^2], \quad \tilde{L}_N = L_N + \log(U_{l,N}^{\text{poly}} / \Pi_l). \quad (\text{AMT29})$$

Pointwise $|f_l(c + iy)|^2 \geq \Pi_l$. Also $\|(iy - t)P\|_\sigma^2 = \|yP\|_\sigma^2 + t^2 \|P\|_\sigma^2$, so the degree-specific multiplication estimate in AMT6, applied successively, gives $\|f_l P\|_\sigma^2 \leq U_{l,N}^{\text{poly}} \|P\|_\sigma^2$. Writing H_N^0 for the unchanged $m_{0,k}$ source Gram yields

$$\beta_k \Pi_l H_N^\sigma \preceq H_N^0 \preceq \beta_k U_{l,N}^{\text{poly}} H_N^\sigma, \quad \frac{a_k}{D_N \beta_k U_{l,N}^{\text{poly}}} H_N^0 \preceq H_N^{\text{ar}} \preceq \frac{A_k}{\beta_k \Pi_l} H_N^0. \quad (\text{AMT30})$$

The second pair follows by using the first upper bound for its lower inequality and the first lower bound for its upper inequality. Minimizing and restricting with the same J_N, I, Λ gives AMT30 for all three metrics in AMT28. Its common upper scalar is $A_k/(\beta_k \Pi_l)$; it is retained before its four signed copies cancel. The same determinant proof as AMT12–17 therefore gives

$$\begin{aligned} \tilde{E}_k^+ &= E_k^+ + \log(U_{l,q-1}^{\text{poly}} U_{l,q}^{\text{poly}} / \Pi_l^2), & \tilde{E}_k^- &= E_k^- + \log(U_{l,2q-1}^{\text{poly}} U_{l,2q}^{\text{poly}} / \Pi_l^2), \\ \boxed{-r\tilde{E}_k^+ \leq F_K^{\text{ar}} - F_K^0 \leq r\tilde{E}_k^-, \quad -b\tilde{E}_k^+ \leq F_B^{\text{ar}} - F_B^0 \leq b\tilde{E}_k^-} \end{aligned} \quad (\text{AMT31})$$

For an explicit bound on the extra widths, $t_j \geq 1/2$ gives

$$0 \leq \log(U_{l,N}^{\text{poly}} / \Pi_l) = \sum_{j=0}^{l-1} \log \left(1 + \frac{4(N+j+1)^2}{t_j^2} \right) \leq l \log[1 + 16(N+l)^2]. \quad (\text{AMT32})$$

Thus $\tilde{E}_k^\pm = O_h(k \log(q+k))$ on the stated endpoints. This is a proved polynomial-source estimate for the exact original f_l , with no inference that its multiplication preserves K .

8.7 The evaluated original simple-quartet arithmetic kernel

Now take the original simple-quartet class $m = 1, k = 4l+1, l \geq 1, q = (k+1)^2$. The parameter u is precisely the period parameter in AMT3, not the real integration coordinate y . RPC21 and RPC23 give an explicit finite radius R_* for the unchanged quartet and arithmetic unit. For reference its exact constants and cases are recorded here. Set

$$\begin{aligned} \beta &= 2(\gamma^2 - \delta^2), \quad \eta = (\delta^2 + \gamma^2)^2, \quad a = v_h(\tfrac{1}{2} + \delta + i\gamma) \neq 0, \\ c_3 &= \frac{a^{-2} - \bar{a}^{-2}}{4i\delta\gamma}, \quad c_1 = \frac{a^{-2} + \bar{a}^{-2}}{2} - (\delta^2 - \gamma^2)c_3, \\ g_j &= 5^{j/5-1} e^{\pi i j/5} \Gamma(j/5) \quad (1 \leq j \leq 4), \quad g_+ = \max_j |g_j|, \quad g_-^{-1} = \max_j |g_j|^{-1}, \quad \kappa = \max_{1 \leq j < 4} |g_{j+1}/g_j|, \\ E_j^{\text{ray}} &= \frac{8}{5} \int_0^\infty r^j (r^3/3 + r) e^{-r^5/5 + r^3/3 + r} dr \quad (0 \leq j \leq 3), \quad C_T = 2 \left(\sum_{j=0}^3 (E_j^{\text{ray}})^2 \right)^{1/2}, \\ C_K &= 4C_T g_-^{-1} + 2g_+ g_-^{-1} + 2g_+ g_-^{-2} C_T, \quad C_{K,3} = C_K(3\kappa^2 + 3\kappa C_K + C_K^2), \\ \epsilon_* &= \min\{1, (2C_T g_-^{-1})^{-1}\}, \quad B_* = \beta + \eta. \end{aligned} \quad (\text{AMT33})$$

Here $g_-^{-2} = (g_-^{-1})^2$. All integrals converge by their negative fifth-degree exponent, and $(c_1, c_3) \neq (0, 0)$ by the original unit values. The exact radius is

$$R_* = \begin{cases} \left[\max \left\{ 1, \frac{B_*}{\epsilon_*}, \frac{2(|c_3|C_{K,3}B_* + |c_1|(\kappa + C_K))}{|c_3 g_4/g_1|} \right\} \right]^{5/2}, & c_3 \neq 0, \\ \left[\max \left\{ 1, \frac{B_*}{\epsilon_*}, \frac{2C_K B_*}{|g_2/g_1|} \right\} \right]^{5/2}, & c_3 = 0. \end{cases} \quad (\text{AMT34})$$

The complete period calculation RPC16–25 proves, for every original branch and every $|u| \geq R_*$, a nonzero original commutator entry: its $(4, 1)$ entry in the first case and its $(2, 1)$ entry in the second. RPC14–15 identifies those entries with the exact test for the original rank-four quadric's cyclic invariance. Thus its actual projective orbit has five elements throughout this domain. OQX1–3, using the product of precisely these five original quadrics, proves

$$\boxed{r = 8k - 16, \quad b = k^2 - 6k + 17, \quad m = 1, \quad k = 4l + 1, \quad l \geq 1, \quad |u| \geq R_*} \quad (\text{AMT35})$$

This is the evaluated original period outcome; no value of the amplitude or of a period coefficient is chosen. The radius depends on the fixed one-factor quartet and unit and is independent of k .

For the same original packet, OKV22 proves for both existing source orders $s = 1, k$, and for the actual K of any rank,

$$\begin{aligned} D_* &= \max\{3, \sqrt{\delta^2 + \gamma^2}\}, \quad C(\delta, \gamma) = 12 + 4 \log(2D_*), \\ Z_q &= 1 + 2(q+1)q^2 2^{6q} (2D_* q)^{4q}, \\ 0 \leq F_K^{(s)} &\leq 2r \log Z_q \leq 2rC(\delta, \gamma)q \log(q+1), \\ F_K^{(1)} &= F_K^\sigma, \quad F_K^{(k)} = F_K^0. \end{aligned} \tag{AMT36}$$

Its proof includes the exact source coefficient map and the unchanged kernel inclusion. Combining it with the already proved AMT17–20 gives the finite arithmetic conclusion

$$\begin{aligned} \max\{0, F_K^\sigma - C_K^-\} &\leq F_K^{\text{ar}} \leq F_K^\sigma + C_K^+, \\ 0 \leq F_K^{\text{ar}} &\leq U_K := 2r \log Z_q + \widehat{C}_K^+ \\ &\leq 2rC(\delta, \gamma)q \log(q+1) + r\widetilde{E}_k^-. \end{aligned} \tag{AMT37}$$

The cap symbols $C_K^\pm, \widehat{C}_K^\pm$ in this formula mean the endpoint sums AMT16; the constant C_K without a superscript in AMT33 belongs only to the explicit period radius. They have different stated indices and are not identified. The alternative finite tensor-source bound $F_K^{\text{ar}} \leq 2r \log Z_q + r\widetilde{E}_k^-$ also follows from AMT31. Formula AMT37 retains the sharper fixed-source widths.

On AMT35 the exact ranks give

$$F_K^{\text{ar}} - F_K^\sigma = O_h(k^2), \quad F_K^{\text{ar}} = O_h(kq \log(q+1)), \quad \frac{F_K^{\text{ar}}}{q^2} \longrightarrow 0. \tag{AMT38}$$

For the first estimate use $r = 8k - 16$ in AMT20 and AMT22, with $\log q = 2 \log(k+1)$. The second follows from AMT37. For the limit, divide its last line by q^2 : its first term is at most $16C(\delta, \gamma)k \log(q+1)/q \rightarrow 0$, and its second is $O_h(k^2/q^2) \rightarrow 0$. Nonnegativity gives the limit by the two bounds. These estimates are uniform over the entire original domain $|u| \geq R_*$, since none of their right sides depends on u . The tensor comparison likewise gives $F_K^{\text{ar}} - F_K^0 = O_h(k^2 \log q) = o(q^2)$ from AMT31–32.

8.8 The original boundary receives the full baseline coefficient

Before using a limit, the exact finite arithmetic identity and its consequence are

$$\begin{aligned} F_B^{\text{ar}} &= \mathcal{B}_k^\sigma + \delta_k^\sigma - F_K^{\text{ar}}, \\ \max\{0, \mathcal{B}_k^\sigma + \delta_k^\sigma - U_K\} &\leq F_B^{\text{ar}} \leq \mathcal{B}_k^\sigma + \delta_k^\sigma, \\ \max\{0, \mathcal{B}_k^\sigma + \underline{\delta}_k^\sigma - U_K\} &\leq F_B^{\text{ar}} \leq \mathcal{B}_k^\sigma + \bar{\delta}_k^\sigma. \end{aligned} \tag{AMT39}$$

The first identity follows from AMT10 and AMT15; AMT37 and $F_K^{\text{ar}}, F_B^{\text{ar}} \geq 0$ prove the first interval, and AMT21 proves the second. The correlated boundary return also has the sharper exact interval AMT18, with every original signed total correction present.

BSL13–19 supplies the complete original finite baseline and its proved coefficient

$$\mathcal{B}_k^\sigma = V_q + \eta_q + \eta_{q+1} - \eta_1, \quad \frac{\mathcal{B}_k^{\text{ar}}}{q^2} \longrightarrow C_B^{\text{base}} := 9 - 8 \log 2 + \mathcal{F}(2, \pi) > \frac{769}{17010} > 0. \tag{AMT40}$$

Here V_q , the three distinct original relation errors η_r , and the energy $\mathcal{F}$ are exactly the BSL/HBL/GEL quantities, whose complete definitions and proofs are in the retained closure. The superscript on C_B^{base} only distinguishes that existing baseline coefficient from the endpoint caps $C_B^\pm$ in AMT16. Its finite BSL14 enclosure can be substituted directly for $\mathcal{B}_k^\sigma$ in AMT39,

keeping the signs of all three η terms; no new evaluation is needed for that substitution. Subtract AMT38 from the exact original identity $F_B^{\text{ar}} = \mathcal{B}_k^{\text{ar}} - F_K^{\text{ar}}$. It proves

$$\boxed{\frac{F_B^{\text{ar}}}{q^2} \longrightarrow C_B^{\text{base}} > 0, \quad F_K^{\text{ar}} + F_B^{\text{ar}} = \mathcal{B}_k^{\text{ar}}, \quad m = 1, \quad k = 4l + 1, \quad l \rightarrow \infty, \quad |u| \geq R_*} \quad (\text{AMT41})$$

The convergence is uniform over that original period domain by AMT38 and because the full-source baseline is independent of u . Thus the proved original kernel estimate now passes to the actual arithmetic kernel, and the original arithmetic boundary carries the entire positive q^2 coefficient. All earlier full-sum identities remain exact, with their individual mixed contributions now receiving AMT37–41 and their signed finite corrections AMT17–21.

8.9 The same comparison evaluates the mixed Gamma return on the kq scale

The first inequality pair AMT30 directly compares the two existing Gamma sources on the same original coefficient space. Apply its proved minimum and restriction maps AMT7–9 to $X = E, K, B$, writing $H_{E,N}^a = G_N^a$, $H_{B,N}^a = Q_N^a$, and $r_E = q, r_K = r, r_B = b$. At every original endpoint it gives

$$\beta_k \Pi_l H_{X,N}^\sigma \preceq H_{X,N}^0 \preceq \beta_k U_{l,N}^{\text{poly}} H_{X,N}^\sigma. \quad (\text{AMT42})$$

Define the exact relative determinant and the positive degree width

$$\begin{aligned} z_{X,N} &= \log \frac{\det H_{X,N}^0}{\det H_{X,N}^\sigma} - r_X \log(\beta_k \Pi_l), & \omega_N^\Gamma &= \log(U_{l,N}^{\text{poly}} / \Pi_l), \\ 0 &\leq z_{X,N} \leq r_X \omega_N^\Gamma, & z_{K,N} + z_{B,N} &= z_{E,N}, \\ \Omega_k^{\text{low}} &= \omega_{q-1}^\Gamma + \omega_q^\Gamma, & \Omega_k^{\text{high}} &= \omega_{2q-1}^\Gamma + \omega_{2q}^\Gamma. \end{aligned} \quad (\text{AMT43})$$

Congruence and multiplication of the eigenvalues in AMT42 prove the two inequalities. The determinant identity AMT10, applied to the tensor metric AMT28 as well, proves the common sum. These facts preserve the complete original kernel and boundary rather than assigning separate arbitrary errors to their determinants.

Use the precise return orientation

$$\Delta_X^\Gamma = F_X^\sigma - F_X^0 \quad (X = K, B), \quad \mathcal{H}_k = \mathcal{B}_k^\sigma - \mathcal{B}_k^0 = -\mathfrak{T}_k.$$

The last equality is the original TWA30/BSL18 sign. Substituting all four z terms, and canceling $\log(\beta_k \Pi_l)$ only after the signed sum, proves

$$\boxed{\begin{aligned} \Delta_X^\Gamma &= -z_{X,q-1} - z_{X,q} + z_{X,2q-1} + z_{X,2q}, \\ -r_X \Omega_k^{\text{low}} &\leq \Delta_X^\Gamma \leq r_X \Omega_k^{\text{high}}, \\ \Delta_K^\Gamma + \Delta_B^\Gamma &= \mathcal{H}_k, \\ \mathcal{H}_k - r \Omega_k^{\text{high}} &\leq \Delta_B^\Gamma \leq \mathcal{H}_k + r \Omega_k^{\text{low}}. \end{aligned}} \quad (\text{AMT44})$$

AMT29 displays every factor of both Ω values explicitly; AMT32 gives the finite bounds

$$\begin{aligned} 0 &\leq \Omega_k^{\text{low}} \leq l \log[1 + 16(q-1+l)^2] + l \log[1 + 16(q+l)^2], \\ 0 &\leq \Omega_k^{\text{high}} \leq l \log[1 + 16(2q-1+l)^2] + l \log[1 + 16(2q+l)^2]. \end{aligned} \quad (\text{AMT45})$$

In particular both are $O(k \log(q+1))$ on the original $m = 1$ class. Its actual rank $r = 8k - 16$ therefore proves directly

$$\boxed{\Delta_K^\Gamma = O_h(k^2 \log(q+1)) = o(kq), \quad \frac{F_K^{\text{ar}} - F_K^\sigma}{kq} \longrightarrow 0, \quad \frac{F_K^{\text{ar}} - F_K^0}{kq} \longrightarrow 0.} \quad (\text{AMT46})$$

For the first equality divide the AMT44 bounds by kq and use $q = (k+1)^2$, so $k \log(q+1)/q \rightarrow 0$. The second limit uses AMT38, since $k^2/(kq) = k/q \rightarrow 0$. The last uses the exact identity $F_K^{\text{ar}} - F_K^0 = \Delta_K^\sigma + \Delta_K^\Gamma$. All bounds are uniform for $|u| \geq R_*$.

For completeness the entire finite arithmetic boundary return is also evaluated by those same quantities:

$$\boxed{\begin{aligned} F_B^{\text{ar}} - F_B^0 &= \mathcal{H}_k + \delta_k^\sigma - (F_K^{\text{ar}} - F_K^0), \\ \mathcal{H}_k + \delta_k^\sigma - r\tilde{E}_k^- &\leq F_B^{\text{ar}} - F_B^0 \leq \mathcal{H}_k + \delta_k^\sigma + r\tilde{E}_k^+. \end{aligned}} \quad (\text{AMT47})$$

This uses AMT31 with its exact endpoint constants $\tilde{E}_k^+ = E_k^+ + \Omega_k^{\text{low}}$ and $\tilde{E}_k^- = E_k^- + \Omega_k^{\text{high}}$, rather than a new source comparison. Both tensor and arithmetic source masses and the full original resultant have already been retained in AMT23–30 before these signed cancellations.

The proved GEL18–19 constant is

$$\mathcal{C}_\Gamma = 2 \int_{a_*^2}^{b_*^2} \log x \rho_{2,\pi}(x) dx - 4(2 \log 2 - 1) > 0, \quad \frac{\mathfrak{T}_k}{lq} \rightarrow \mathcal{C}_\Gamma. \quad (\text{AMT48})$$

Here a_*, b_* and $\rho_{2,\pi}$ are exactly the proved GEL8/EIQ equilibrium endpoints and density; their complete definitions and integral proof remain in that provider. The endpoint letters here avoid the already fixed original period parameter u . GEL18a gives $4 \log(27/(8\pi)) \leq \mathcal{C}_\Gamma \leq 6$ and proves the strict positivity. TWA15 also proves the actual source return $\delta_k^\sigma/(kq) \rightarrow 0$. As $l/k \rightarrow 1/4$, AMT44, AMT46 and AMT47 now prove

$$\boxed{\begin{aligned} \frac{F_B^\sigma - F_B^0}{kq} &\rightarrow -\frac{\mathcal{C}_\Gamma}{4}, & \frac{F_B^{\text{ar}} - F_B^0}{kq} &\rightarrow -\frac{\mathcal{C}_\Gamma}{4}, & \frac{F_B^{\text{ar}} - F_B^\sigma}{kq} &\rightarrow 0, \\ \frac{F_B^\sigma - F_B^0}{\mathcal{H}_k} &\rightarrow 1, & \frac{F_B^{\text{ar}} - F_B^0}{\mathcal{H}_k} &\rightarrow 1, & m=1, k=4l+1, l \rightarrow \infty, |u| \geq R_*. \end{aligned}} \quad (\text{AMT49})$$

Indeed the first two limits subtract an $o(kq)$ kernel return from $\mathcal{H}_k = -\mathfrak{T}_k$, whose quotient by kq tends to $-\mathcal{C}_\Gamma/4$. The third follows from $\Delta_B^\sigma = \delta_k^\sigma - \Delta_K^\sigma$. The nonzero limit of $\mathcal{H}_k/(kq)$ makes its denominator nonzero for all sufficiently large original k and proves the last two ratios by division. The uniformity in u follows from the uniform kernel bounds and the period-independent total returns. Thus the full existing source comparison evaluates the mixed Gamma return also at its kq scale, with its finite signed remainders AMT44 and AMT47 still present.

9 Factorial control of the original common kernel at the kq scale

The exact denominator in OKV15 contains the information needed to improve its absolute volume estimate. This proof keeps that denominator and also allows every original primary multiplicity. The original Gamma sources, root coordinates, coefficient maps and observation are retained. The simple-quartet conclusion uses the actual period and rank calculations PQO/RPC and OQC/OQX, with their complete proofs supplied as dependencies.

9.1 Original coordinates, all primary multiplicities, and source norms

Fix the original full-order quartet with $0 < \delta < 1/2$, $\gamma > 2$, and a positive integer primary order m . On the original tensor class set

$$k = 4l + 1, \quad l \geq 1, \quad e_k = 1 + k(m - 1), \quad q = e_k(k + 1)^2, \quad c = k/2, \quad s \in \{1, k\}, \quad b_s = s/2. \quad (\text{KAF1})$$

The original polynomial and its exact real-coordinate representative are

$$\begin{aligned} \chi(S) &= \prod_{a,b=0}^k [S - c - (2a - k)\delta - i(2b - k)\gamma]^{e_k}, \\ \omega_{ab} &= (2b - k)\gamma - i(2a - k)\delta, \\ P(y) &= i^{-q} \chi(c + iy) = \prod_{a,b=0}^k (y - \omega_{ab})^{e_k}. \end{aligned} \quad (\text{KAF2})$$

In the full list $\omega_1, \dots, \omega_q$, each root ω_{ab} occurs e_k times. The exact maximal modulus remains $R = k\sqrt{\delta^2 + \gamma^2}$. Substitution gives inverse algebra maps

$$\Phi : \mathbb{C}[y]/(P) \longrightarrow E = \mathbb{C}[S]/(\chi), \quad [r] \longmapsto [r((S - c)/i)], \quad \Phi^{-1}[f] = [f(c + iy)]. \quad (\text{KAF3})$$

Indeed $P((S - c)/i) = i^{-q} \chi(S)$, and the two substitutions are inverse before quotienting. Thus every root, multiplicity and factor of i in the original remainder operation is present.

Retain the complete measures and their monic orthogonal polynomials:

$$\begin{aligned} M_s &= (2\pi)^{s/2}, \quad dm_{0,s}(y) = \frac{M_s 2^{b_s-2}}{\pi \Gamma(b_s)} \left| \Gamma\left(\frac{b_s}{2} + \frac{iy}{2}\right) \right|^2 dy, \\ p_0^{(s)} &= 1, \quad p_1^{(s)} = y, \quad p_{d+1}^{(s)} = yp_d^{(s)} - d(b_s + d - 1)p_{d-1}^{(s)}, \\ h_d^{(s)} &= \int |p_d^{(s)}|^2 dm_{0,s} = M_s d!(b_s)_d, \quad \varphi_d^{(s)} = p_d^{(s)} / \sqrt{h_d^{(s)}}, \quad u_d^{(s)}(S) = \varphi_d^{(s)}((S - c)/i). \end{aligned} \quad (\text{KAF4})$$

These are precisely the existing orders $1, k$. Their complete source transform and norm calculation are IFB2–5 and OKV4–5. In particular $\int e^{ty} dm_{0,s} = M_s (\cos t)^{-b_s}$ for $|t| < \pi/2$. The generating series $(1 + z^2)^{-b_s/2} \exp(y \arctan z)$ has coefficients $p_d^{(s)} z^d / d!$. Integrating its product at two sufficiently small real variables gives $M_s (1 - zw)^{-b_s}$, proving the displayed norms and orthogonality by coefficient comparison. The original exponential bound justifies those finite coefficient differentiations under the integral. None of these source statements depends on m .

In the original polynomial frame of E let C_s have columns $[u_0^{(s)}], \dots, [u_{q-1}^{(s)}]$. Its determinant is exactly

$$\begin{aligned} \det C_s &= i^{-q(q-1)/2} \prod_{d=0}^{q-1} (h_d^{(s)})^{-1/2} \neq 0, \\ v_j^{(s)} &= C_s^{-1} [u_{q+j}^{(s)}], \quad K_j^{(s)} = I_q + \sum_{a=0}^{j-1} v_a^{(s)} (v_a^{(s)})^*, \quad 0 \leq j \leq q + 1. \end{aligned} \quad (\text{KAF5})$$

The last included vector has original degree $2q$.

9.2 A degree-dependent constant in the unchanged remainder estimate

Define the actual finite constants

$$A_{k,m} = \max \left\{ 2\sqrt{1 + \frac{k}{2q}}, \frac{R}{q} \right\}, \quad T = qA_{k,m}, \quad \mathcal{A}_{k,m} = 6 \log 2 + 4 + 4 \log A_{k,m}. \quad (\text{KAF6})$$

The letter $A_{k,m}$ in this section is a coefficient-estimate constant; it is not the arithmetic upper source constant A_k of TWA/AMT. In particular $A_{k,m} \geq 2$ and $R \leq T$. For fixed original quartet and fixed m , $q \geq (k+1)^2$ implies

$$A_{k,m} \longrightarrow 2, \quad \mathcal{A}_{k,m} \longrightarrow 4 + 10 \log 2. \quad (\text{KAF7})$$

Indeed $k/q \rightarrow 0$ and $R/q = \sqrt{\delta^2 + \gamma^2} k/q \rightarrow 0$. The root-dependent term in the maximum is retained at every finite k .

Here is the complete remainder estimate needed below. For the full root list define elementary and complete homogeneous polynomials $e_h(\omega)$ and $h_t(\omega)$ by their finite sums, with $e_0 = h_0 = 1$ and $e_h = 0$ outside $0 \leq h \leq q$. The formal identity

$$\left(\sum_{h=0}^q (-1)^h e_h(\omega) z^h \right) \left(\sum_{t \geq 0} h_t(\omega) z^t \right) = 1 \quad (\text{KAF8})$$

follows by multiplying each of the q factors $1 - \omega_a z$ by its geometric series. Repeated roots do not affect this identity. It proves, for $D = q + j$ and $0 \leq j \leq q$, the exact signed formula

$$[y^a] \text{rem}_P(y^{q+j}) = - \sum_{t=0}^j h_t(\omega) (-1)^{q+j-t-a} e_{q+j-t-a}(\omega), \quad 0 \leq a < q. \quad (\text{KAF9})$$

In detail the quotient is $\sum_{t=0}^j h_t(\omega) y^{j-t}$; KAF8 cancels all coefficients of degree at least q in its product with P , and the remaining coefficients are KAF9. This proves the formula by uniqueness of monic division, including all multiplicities.

The defining sums give $|e_h| \leq \binom{q}{h} R^h$ and $|h_t| \leq \binom{q+t-1}{t} R^t$. In every nonzero summand of KAF9 the total root exponent is $q + j - a$. The binomial coefficients of h_t increase with t , and the distinct indices in the e_h sum contribute at most 2^q . Hence

$$|[y^a] \text{rem}_P(y^{q+j})| \leq 2^q \binom{q+j-1}{j} R^{q+j-a} \leq 2^{3q-1} R^{q+j-a}. \quad (\text{KAF10})$$

The last bound is the binomial expansion at one, since $j \leq q$.

The exact orthonormal recurrence from KAF4 has upward and downward weights $\sqrt{(a+1)(b_s+a)}$ and $\sqrt{a(b_s+a-1)}$. On degree at most $q-2$ each shift has norm at most $\sqrt{q(b_s+q)}$. Because $b_s \leq k/2$, their sum has norm at most $2\sqrt{q(k/2+q)} \leq T$. Iterating from the constant, with all intermediate degrees in that range, proves

$$\|y^a\|_{L^2(m_{0,s})} \leq \sqrt{M_s} T^a, \quad 0 \leq a < q. \quad (\text{KAF11})$$

For the finite coefficient sum $c_T(f) = \sum_a [y^a] f |T^a$ the original recurrence gives $c_T(p_{d+1}^{(s)}) \leq T c_T(p_d^{(s)}) + d(b_s + d - 1) c_T(p_{d-1}^{(s)})$. For $1 \leq d < 2q$,

$$d(b_s + d - 1) \leq 2q(k/2 + 2q) = 4q^2 + kq \leq 4q^2 + 2kq \leq T^2.$$

Starting with $c_T(p_0) = 1$, $c_T(p_1) = T$, induction therefore proves $c_T(p_d) \leq (2T)^d$ for $0 \leq d \leq 2q$: the recurrence bound is $(3/4)(2T)^{d+1} \leq (2T)^{d+1}$. Combining KAF10 with all q monomial norms KAF11, then using the linearity of the original remainder map, gives

$$\|\text{rem}_P(p_d^{(s)})\|_{L^2(m_{0,s})} \leq \sqrt{M_s} q 2^{3q} (2T)^d, \quad 0 \leq d \leq 2q. \quad (\text{KAF12})$$

For a monomial of degree $D \geq q$, each coefficient term is at most $\sqrt{M_s} 2^{3q-1} R^{D-a} T^a \leq \sqrt{M_s} 2^{3q-1} T^D$. For $D < q$ use KAF11 directly. These two cases and the proved coefficient sum establish KAF12 without deleting a remainder coefficient or changing the source norm.

9.3 Retaining the factorial denominator

Expansion of the degree-below- q remainder in the orthonormal polynomials of KAF4, followed by the exact map KAF3, proves

$$\|v_j^{(s)}\|_2^2 = \frac{\|\text{rem}_P(p_{q+j}^{(s)})\|_{L^2(m_{0,s})}^2}{M_s(q+j)!(b_s)_{q+j}} \leq \frac{q^2 2^{6q} (2T)^{2(q+j)}}{(q+j)!(b_s)_{q+j}}. \quad (\text{KAF13})$$

The factor M_s occurs in the original numerator and denominator before this cancellation. For every $d \geq 0$,

$$d!(b_s)_d \geq d!(1/2)_d = \frac{(2d)!}{4^d}, \quad \|v_j^{(s)}\|_2^2 \leq q^2 2^{6q} \frac{(4T)^{2(q+j)}}{(2q+2j)!}. \quad (\text{KAF14})$$

The inequality is factor by factor because $b_s \geq 1/2$. The equality follows by separating even and odd factors of $(2d)!$. This is precisely the denominator whose further retention sharpens OKV16–17.

Define the positive exact quantities

$$\eta_{k,m} = \frac{1 - 1/(4q)}{A_{k,m}^2} < \frac{1}{4}, \quad Z_{k,m} = 1 + q^2 2^{6q} \frac{(4q A_{k,m})^{4q}}{(4q)!} \frac{1 - \eta_{k,m}^{q+1}}{1 - \eta_{k,m}}. \quad (\text{KAF15})$$

For $t_d = (4T)^{2d}/(2d)!$, $q+1 \leq d \leq 2q$ gives $t_{d-1}/t_d = (2d)(2d-1)/(16T^2) \leq \eta_{k,m}$. Consequently $\sum_{d=q}^{2q} t_d \leq t_{2q} \sum_{a=0}^q \eta_{k,m}^a$. Adding the $q+1$ actual rank-one updates in KAF5 and using KAF14 proves the finite operator bound

$$\boxed{1 \leq \|K_{q+1}^{(s)}\|_2 \leq Z_{k,m}, \quad s \in \{1, k\}.} \quad (\text{KAF16})$$

Here $\|vv^*\|_2 = \|v\|_2^2$ follows by Cauchy–Schwarz with equality on $v \neq 0$; a zero vector contributes zero.

For an explicit linear bound, monotonicity of $\log x$ on each interval $[j-1, j]$ gives

$$\log n! = \sum_{j=1}^n \log j \geq \int_1^n \log x \, dx = n \log n - n + 1.$$

Thus $(4q A_{k,m})^{4q}/(4q)! \leq e^{-1} (e A_{k,m})^{4q}$, where e is the exponential constant and differs from the primary integer e_k . The geometric sum in KAF15 is at most $4/3$. Therefore $Z_{k,m} \leq 1 + [4/(3e)]X$, with $X = q^2 \exp(q A_{k,m})$. The positive exponential series gives $e > 1 + 1 + 1/2 + 1/6 = 8/3$, so $4/(3e) < 1/2$. Since $A_{k,m} \geq 2$ and $q \geq 36$, $X > 2$, and hence $1 + [4/(3e)]X \leq X$. We have proved

$$\boxed{\log \|K_{q+1}^{(s)}\|_2 \leq \log Z_{k,m} \leq q A_{k,m} + 2 \log q.} \quad (\text{KAF17})$$

The full factorial and finite geometric sum in KAF15 remain available as the sharper finite bound.

9.4 All four actual kernel and boundary Grams

Let $\Lambda : E \twoheadrightarrow B$ be the original period and full-unit observation, let $I : \mathcal{K} \hookrightarrow E$ be its actual kernel inclusion, and put $r = \dim \mathcal{K}$. No rank formula is needed in this subsection, so it applies to every original m . The original quotient and restricted Grams are

$$G_{q+j-1}^{(s)} = C_s^{*-}(K_j^{(s)})^{-1} C_s^{-1}, \quad T_s = C_s^{-1} I, \quad H_j^{(s)} = T_s^* (K_j^{(s)})^{-1} T_s. \quad (\text{KAF18})$$

Indeed the original orthonormal source-coordinate remainder matrix is $[I_q, v_0, \dots, v_{j-1}]$. Multiplication by its adjoint gives K_j ; its unique minimum lift of x is its adjoint applied to $K_j^{-1}x$, with

squared norm $x^* K_j^{-1} x$. This proves both matrices KAF18 in the original coefficient frame. The positive update chain implies

$$Z_{k,m}^{-1} T_s^* T_s \preceq H_j^{(s)} \preceq T_s^* T_s, \quad H_{j+1}^{(s)} \preceq H_j^{(s)}. \quad (\text{KAF19})$$

Inversion reverses positive order by congruence and inversion of positive eigenvalues. Restricting this order with the same injective T_s proves KAF19. After congruence with $(T_s^* T_s)^{-1/2}$, every eigenvalue lies between $Z_{k,m}^{-1}$ and one. Thus $-r \log Z_{k,m} \leq \log \det H_j^{(s)} - \log \det (T_s^* T_s) \leq 0$.

The exact four original degrees are $q-1, q, 2q-1, 2q$. With their unchanged signs define

$$F_K^{(s)} = \log \det H_0^{(s)} + \log \det H_1^{(s)} - \log \det H_q^{(s)} - \log \det H_{q+1}^{(s)}.$$

Pairing degrees $q-1$ with $2q-1$, and q with $2q$, proves nonnegativity by KAF19. Each difference is at most $r \log Z_{k,m}$. The four original frame factors cancel only after these four signs have been recorded. It follows that

$$0 \leq F_K^{(s)} \leq 2r \log Z_{k,m} \leq 2r [q \mathcal{A}_{k,m} + 2 \log q], \quad s \in \{1, k\}. \quad (\text{KAF20})$$

If $r = 0$, every kernel determinant is one and the statement has its same empty-space meaning.

At each original degree the boundary Gram is $Q_j^{(s)} = (\Lambda(G_{q+j-1}^{(s)})^{-1} \Lambda^*)^{-1}$. For any fixed section S_* of Λ , its minimum section $S_j^{(s)} = (G_{q+j-1}^{(s)})^{-1} \Lambda^* Q_j^{(s)}$ differs from S_* by a kernel column and is orthogonal to I in that metric. The change of frame has determinant one. Taking determinants in this orthogonal frame proves, with its original coefficient factor,

$$\det H_j^{(s)} \det Q_j^{(s)} = |\det[I, S_*]|^2 \det G_{q+j-1}^{(s)}.$$

The same four signs now prove the exact splitting $F_K^{(s)} + F_B^{(s)} = \mathcal{B}_k^{(s)}$. Boundary minimum norms decrease with degree, so $F_B^{(s)} \geq 0$. In particular

$$\max\{0, \mathcal{B}_k^{(s)} - 2r \log Z_{k,m}\} \leq F_B^{(s)} \leq \mathcal{B}_k^{(s)}. \quad (\text{KAF21})$$

These are statements about the original kernel and boundary, with all multiplicities and the same observation throughout.

9.5 The actual simple quartet and the arithmetic source

For $m = 1$, the complete RPC calculation supplies an explicit finite R_* from the original unit and quartet. On every logarithm branch and for $|u| \geq R_*$ its original period quadric has five distinct cyclic orbit members. The full invariant restriction ideal calculation OQX then gives $r = 8k - 16$. Substitution in the finite KAF20 gives

$$0 \leq \frac{F_K^{(s)}}{kq} \leq \left(16 - \frac{32}{k}\right) \left(\mathcal{A}_{k,1} + \frac{2 \log q}{q}\right), \quad s \in \{1, k\}, \quad |u| \geq R_*. \quad (\text{KAF22})$$

The original arithmetic-source comparison AMT17–22 proves $-r E_k^+ \leq F_K^{\text{ar}} - F_K^{(1)} \leq r E_k^-$, where the unchanged full constants are

$$\begin{aligned} E_k^+ &= 2 \log(A_k/a_k) + \log D_{q-1} + \log D_q, \\ E_k^- &= 2 \log(A_k/a_k) + \log D_{2q-1} + \log D_{2q}, \\ D_N &= [1 + 4(N + 21 + 4m)^2]^{21+4m}. \end{aligned}$$

Here a_k, A_k are exactly the original arithmetic constants of AMT4, whose full source inequality and minimum-section proof are included there. They retain the actual numerator $2\xi/h$ and all

source masses; AMT22 proves $E_k^\pm = O_h(k + \log q)$. Thus the actual arithmetic metric satisfies the finite bound

$$0 \leq F_K^{\text{ar}} \leq (16k - 32) \log Z_{k,1} + (8k - 16)E_k^-. \quad (\text{KAF23})$$

Combining KAF7, KAF22–23, and $q = (k + 1)^2$ proves uniformly on the entire stated original period domain, for $a \in \{\sigma, 0, \text{ar}\}$,

$$0 \leq \liminf_{k \rightarrow \infty} \frac{F_K^a}{kq} \leq \limsup_{k \rightarrow \infty} \frac{F_K^a}{kq} \leq 16(4 + 10 \log 2). \quad (\text{KAF24})$$

For precision, uniformity means the same finite upper bound holds for every such u and branch at each original k ; therefore it also bounds every sequence of those parameters. The arithmetic extra term divided by kq is at most $(8 - 16/k)E_k^-/q \rightarrow 0$. AMT46 further proves that the three ratios differ by quantities tending uniformly to zero. Thus their limit points coincide for any fixed allowed sequence of period parameters; KAF24 does not assert that this bounded sequence already has a unique limit.

Finally the exact arithmetic identity gives

$$\max\{0, \mathcal{B}_k^{\text{ar}} - (16k - 32) \log Z_{k,1} - (8k - 16)E_k^-\} \leq F_B^{\text{ar}} \leq \mathcal{B}_k^{\text{ar}}. \quad (\text{KAF25})$$

Its full-source value remains $\mathcal{B}_k^{\text{ar}} = \mathcal{B}_k^\sigma + \delta_k^\sigma = \mathcal{B}_k^0 + \mathcal{H}_k + \delta_k^\sigma$. All exact Gamma and arithmetic signed returns of AMT44–49 are retained. The new calculation improves the absolute common kernel bound to the kq scale, while its exact four-endpoint determinant and actual period dependence remain the objects for the next leading term calculation.

10 An exact conductor frame and a fixed invariant-generator recursion

Retain the original simple-quartet data, $m = 1$, $k = 4l + 1$, $l \geq 1$, the original root order $(++)$, $(+-)$, $(-+)$, $(--)$, and the complete original period and amplitude matrix of OQC1:

$$\mathbf{H} = F^{-1} \Pi U \mathcal{E}_{\text{prim}}.$$

Thus $\det \mathbf{H} \neq 0$ and $Q(x) = Q_0(\mathbf{H}^{-1}x)$, where $Q_0(t) = t_{++}t_{--} - t_{+-}t_{-+}$. Use $gf(x) = f(D^{-1}x)$, $D = \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4)$, $\zeta = e^{2\pi i/5}$, and the original invariant algebra $\mathcal{J} = \mathbb{C}[x_1, x_2, x_3, x_4]^{(g)}$. This calculation applies on the actual five-orbit period locus, which includes every original logarithm branch with $|u| \geq R_*$ by PQO and RPC. On that locus the five prime polynomials $Q_a = g^a Q$, $0 \leq a < 5$, are pairwise nonassociate. Keep their actual products

$$\mathcal{A} = \prod_{a=1}^4 Q_a, \quad \mathcal{H}_Q = Q\mathcal{A} = \prod_{a=0}^4 Q_a, \quad \deg \mathcal{A} = 8, \quad \deg \mathcal{H}_Q = 10. \quad (\text{OCF1})$$

No coefficient in these polynomials is replaced. The exact orbit decision and the period domain are those of the cited original calculations. The construction below treats every possible nonzero leading coefficient of their actual restriction.

10.1 The original Segre coefficient isomorphism

For $d \geq 0$ let $\mathcal{B}_d$ be the space of biforms of bidegree (d, d) in (z_0, z_1) and (w_0, w_1) . Its ordered basis is

$$e_{ab}^{(d)} = z_0^a z_1^{d-a} w_0^b w_1^{d-b}, \quad 0 \leq a, b \leq d. \quad (\text{OCF2})$$

Pairs are ordered lexicographically, first by a , then by b . The order is used only for the exact coefficient elimination. Set $\mathcal{B}_d = 0$ for $d < 0$. Multiplication satisfies $e_{ab}^{(d)} e_{ij}^{(e)} = e_{a+i, b+j}^{(d+e)}$. The original substitution is

$$t = (z_0 w_0, z_0 w_1, z_1 w_0, z_1 w_1)^T, \quad x = \mathbf{H}t, \quad (\text{OCF3})$$

$$\mathcal{P}_d = \mathbb{C}[x_1, x_2, x_3, x_4]_d, \quad \psi_d : \mathcal{P}_d \longrightarrow \mathcal{B}_d, \quad f \longmapsto f(\mathbf{H}t).$$

The induced map $R_d = \mathcal{P}_d / (Q\mathcal{P}_{d-2}) \longrightarrow \mathcal{B}_d$ is an isomorphism. Here is a coefficient proof with the given orientation. For each (a, b) , choose an integer h between $\max(0, a + b - d)$ and $\min(a, b)$. The monomial $t_{++}^h t_{+-}^{a-h} t_{-+}^{b-h} t_{--}^{d-a-b+h}$ maps to $e_{ab}^{(d)}$, proving surjectivity. Two monomials with the same pair differ in their exponent vectors by an integer multiple of $(1, -1, -1, 1)$. After removing their common monomial, their difference is divisible by $t_{++}t_{--} - t_{+-}t_{-+}$, using $A^n - B^n = (A - B) \sum_{j=0}^{n-1} A^{n-1-j} B^j$. A polynomial in the kernel has coefficient sum zero within every pair, so it is a sum of such differences. The reverse inclusion in the kernel follows by direct substitution. The invertible substitution $x = \mathbf{H}t$ then gives exactly the asserted kernel. Thus these maps preserve multiplication as well as every grading.

Write the full conductor restriction as

$$A = \psi_8(\mathcal{A}) = \sum_{r=0}^8 \sum_{s=0}^8 a_{rs} e_{rs}^{(8)} \neq 0. \quad (\text{OCF4})$$

To verify its nonvanishing, Q is a rank-four irreducible quadric, hence prime in the polynomial ring, and none of $Q_1, \dots, Q_4$ is its associate. Therefore Q does not divide their product. The kernel calculation OCF3 proves OCF4. The ring $\mathcal{B} = \bigoplus_{d \geq 0} \mathcal{B}_d$, with the indicated diagonal grading, embeds in the four-variable polynomial ring; it is an integral domain. Multiplication by A is consequently injective on every $\mathcal{B}_d$.

10.2 Every leading-coefficient case gives an exact strip frame

Let (r_*, s_*) be the lexicographically largest pair for which $a_{r_*s_*} \neq 0$, and put $a_* = a_{r_*s_*}$. This is an actual nonzero coefficient of OCF4. For $d \geq 8$ define

$$\begin{aligned} P_d &= \{(r_* + i, s_* + j) : 0 \leq i, j \leq d - 8\}, \\ E_d^{\text{strip}} &= \{0, \dots, d\}^2 \setminus P_d, \quad W_d = \mathbb{C}^{E_d^{\text{strip}}}. \end{aligned} \quad (\text{OCF5})$$

For $0 \leq d < 8$ put $P_d = \emptyset$, $E_d^{\text{strip}} = \{0, \dots, d\}^2$ and use the same definition of W_d ; for $d < 0$ put $W_d = 0$. Let $S_d : W_d \hookrightarrow \mathbb{C}^{(d+1)^2}$ insert zeros in the positions P_d . All coefficient columns below use OCF2.

The following finite elimination defines $N_d : \mathbb{C}^{(d+1)^2} \rightarrow W_d$ and, for $d \geq 8$, $V_d : \mathbb{C}^{(d+1)^2} \rightarrow \mathbb{C}^{(d-7)^2}$. Starting with the complete input biform f , traverse all $(i, j) \in \{0, \dots, d-8\}^2$ in decreasing lexicographic order. At the step (i, j) , let c be the current coefficient at $(r_* + i, s_* + j)$, record $b_{ij} = c/a_*$, and replace the current biform by

$$f_{\text{current}} - b_{ij} A e_{ij}^{(d-8)}. \quad (\text{OCF6})$$

At completion take the coefficients on E_d^{strip} as $N_d f$ and take (b_{ij}) as $V_d f$. If $d < 8$, N_d is the identity and V_d is the zero map into the zero space. The exact multiplication matrix $M_{A,d}$ gives

$$S_d N_d + M_{A,d} V_d = I, \quad N_d S_d = I_{W_d}, \quad \ker N_d = A \mathcal{B}_{d-8}. \quad (\text{OCF7})$$

To prove these identities, the largest term of $A e_{ij}^{(d-8)}$ is $a_* e_{r_*+i, s_*+j}^{(d)}$; every other term is strictly smaller in the same lexicographic order. Therefore each step zeros its designated position without altering an earlier zero. Summing all subtractions gives the first identity. Input supported on the strip produces zero coefficients at every elimination step, giving the second. Finally, the leading term of Ab , for a nonzero homogeneous biform b , is the product of the two leading terms and belongs to P_d . Indeed every other product is smaller in the additive lexicographic order. Thus $A \mathcal{B}_{d-8} \cap S_d W_d = 0$. The first identity now proves the kernel formula, the uniqueness of the decomposition, and all three identities in OCF7. All operations are linear in the input; their coefficients are explicit rational expressions in the original a_{rs} with powers of the displayed a_* as denominators. The polynomial A itself remains the full polynomial OCF4.

In particular the quotient map induced by N_d is the exact isomorphism

$$\mathcal{B}_d / (A \mathcal{B}_{d-8}) \xrightarrow{\sim} W_d, \quad n_d := \dim W_d = \begin{cases} (d+1)^2, & 0 \leq d < 8, \\ 16d - 48, & d \geq 8. \end{cases} \quad (\text{OCF8})$$

The second formula is $(d+1)^2 - (d-7)^2 = 16d - 48$. The possible choices $(r_*, s_*) \in \{0, \dots, 8\}^2$ form an exhaustive finite partition: the chart for a pair specifies that its coefficient is nonzero and every larger coefficient is zero. Hence all actual coefficient cases have been included. At the first original degree $k = 5$ this construction retains all 36 coordinates; at every original $k \geq 9$ it retains $16k - 48$ coordinates.

10.3 Twenty-five fixed invariant monomials generate all degrees

Define the following fixed finite sets of exponent vectors:

$$G_e = \{\nu \in \mathbb{N}^4 : |\nu| = e, \nu_1 + 2\nu_2 + 3\nu_3 + 4\nu_4 \equiv 0 \pmod{5}\}, \quad 1 \leq e \leq 5, \quad G = \bigcup_{e=1}^5 G_e. \quad (\text{OCF9})$$

Here G_1 is empty. The complete remaining lists, in the original x_1, x_2, x_3, x_4 order, are

e	G_e	
2	$(0, 1, 1, 0), (1, 0, 0, 1)$	
3	$(0, 0, 2, 1), (0, 1, 0, 2), (1, 2, 0, 0), (2, 0, 1, 0)$	
4	$(0, 0, 1, 3), (0, 2, 2, 0), (0, 3, 0, 1), (1, 0, 3, 0),$ $(1, 1, 1, 1), (2, 0, 0, 2), (3, 1, 0, 0)$	(OCF10)
5	$(0, 0, 0, 5), (0, 0, 5, 0), (0, 1, 3, 1), (0, 2, 1, 2),$ $(0, 5, 0, 0), (1, 0, 2, 2), (1, 1, 0, 3), (1, 3, 1, 0),$ $(2, 1, 2, 0), (2, 2, 0, 1), (3, 0, 1, 1), (5, 0, 0, 0).$	

Thus $(|G_1|, \dots, |G_5|) = (0, 2, 4, 7, 12)$ and $|G| = 25$. These lists can also be obtained directly by choosing $0 \leq \nu_1, \nu_2, \nu_3 \leq e$ with their sum at most e , putting $\nu_4 = e - \nu_1 - \nu_2 - \nu_3$, and applying the single displayed congruence, which verifies completeness of each list.

Every positive-degree invariant monomial is a product of members x^ν , $\nu \in G$. To prove this, list the weights of its individual factors in any order. If its degree is at most five, the entire list gives a member of G . If its degree exceeds five, consider the six partial sums of the first five weights, including the empty partial sum, in $\mathbb{Z}/5\mathbb{Z}$. Two coincide; the nonempty block between them has length at most five and total weight zero. Its factors give a divisor x^ν with $\nu \in G$. The complementary monomial again has total weight zero and smaller degree. Induction completes the factorization. Because the action is diagonal, a polynomial is invariant exactly when all its nonzero monomials have weight zero: comparison of each monomial coefficient in $gf = f$ proves this statement. Thus

$$\mathcal{J} = \mathbb{C}[x^\nu : \nu \in G]. \quad (\text{OCF11})$$

All intermediate degrees here are degrees in the original coefficient algebra. This construction introduces no new source distribution and leaves the original source orders 1 and k intact.

For each $\nu \in G_e$ the complete generator biform is

$$f_\nu = \psi_e(x^\nu) = \prod_{j=1}^4 (H_{j1}z_0w_0 + H_{j2}z_0w_1 + H_{j3}z_1w_0 + H_{j4}z_1w_1)^{\nu_j} = \sum_{a,b=0}^e c_{\nu,ab} e_{ab}^{(e)}. \quad (\text{OCF12})$$

For an entirely explicit coefficient formula, sum over all nonnegative 4 by 4 integer arrays (n_{jt}) satisfying $\sum_t n_{jt} = \nu_j$, $\sum_j (n_{j1} + n_{j2}) = a$ and $\sum_j (n_{j1} + n_{j3}) = b$. Then

$$c_{\nu,ab} = \sum_{(n_{jt})} \prod_{j=1}^4 \left(\frac{\nu_j!}{\prod_{t=1}^4 n_{jt}!} \prod_{t=1}^4 H_{jt}^{n_{jt}} \right). \quad (\text{OCF13})$$

The multinomial theorem proves the formula, including every original period and amplitude entry. Every factor has degree at most five and at most 36 possible biform coefficients.

10.4 Exact multiplication on the strip and the invariant recursion

Let $\mathcal{L} = \mathcal{B}/(AB)$, a graded algebra. For $\nu \in G_e$ and $d \geq e$ define

$$m_{\nu,d} : W_{d-e} \longrightarrow W_d, \quad m_{\nu,d} = N_d M_{f_\nu} S_{d-e}. \quad (\text{OCF14})$$

Here M_{f_ν} is the exact coefficient convolution:

$$(M_{f_\nu} S_{d-e} v)_{ab} = \sum_{i,j} c_{\nu,a-i,b-j} (S_{d-e} v)_{ij},$$

with out-of-range indices giving zero. Therefore every entry of OCF14 is supplied by the finite formulas OCF6 and OCF13. For a representative $h \in \mathcal{B}_{d-e}$, OCF7 writes $h - S_{d-e} N_{d-e} h = Ab$. Multiplication by f_ν preserves this ideal. Applying N_d yields the exact identity

$$N_d(f_\nu h) = m_{\nu,d} N_{d-e} h. \quad (\text{OCF15})$$

This proves that OCF14 is multiplication by the specified element of $\mathcal{L}$, including its independence of representatives.

The original invariant image is $J_d = \psi_d(\mathcal{J}_d) \subseteq \mathcal{B}_d$. Set $\bar{J}_d = N_d J_d \subseteq W_d$, and set $\bar{J}_d = 0$ for $d < 0$. The exact recursion is

$$\bar{J}_0 = \mathbb{C} \cdot 1, \quad \bar{J}_d = \sum_{\substack{\nu \in G_e \\ 2 \leq e \leq 5, e \leq d}} m_{\nu,d} \bar{J}_{d-e} \quad (d \geq 1). \quad (\text{OCF16})$$

Indeed OCF11 factors every invariant monomial of degree $d > 0$ as x^ν times an invariant monomial of degree $d - e$. Applying ψ , then OCF15, gives the inclusion from left to right. Conversely every summand is the image of a product of two invariant polynomials, so it belongs to the left side. Linearity proves equality for the complete invariant space. In particular the recursion gives $\bar{J}_1 = 0$ and includes all small degrees.

The conductor ideal is contained in this invariant image with its original trace coefficient:

$$A\mathcal{B}_{d-8} \subseteq J_d, \quad [5R(\mathcal{A}h)]_{(Q)} = [\mathcal{A}h]_{(Q)}, \quad R = \frac{1}{5} \sum_{a=0}^4 g^a. \quad (\text{OCF17})$$

To prove the identity, $g^a \mathcal{A}$ is the product of all five orbit equations except Q_a ; for $a \neq 0$ it contains Q . Thus all four corresponding terms vanish modulo Q , and the $a = 0$ term is the displayed product. Every biform in $\mathcal{B}_{d-8}$ has a polynomial lift by OCF3, so the identity proves the inclusion. It also shows exactly why the quotient used in OCF16 retains the original invariant image.

10.5 Exact ranks, a finite pivot atlas, and the original quotient map

For $d \geq 0$ put

$$d_0(d) = \frac{\binom{d+3}{3} + 4\epsilon_d}{5}, \quad \epsilon_d = \mathbf{1}_{d \equiv 0 \pmod{5}} - \mathbf{1}_{d \equiv 1 \pmod{5}}, \quad d_0(d) = 0 \quad (d < 0). \quad (\text{OCF18})$$

This is the number of degree- d weight-zero monomials: the identity character has generating function $(1 - z)^{-4}$, each of the four nonidentity characters has generating function $\prod_{j=1}^4 (1 - z\zeta^j)^{-1} = (1 - z)/(1 - z^5)$, and averaging their coefficients gives OCF18. Every invariant polynomial divisible by Q is divisible by all five Q_a , hence by $\mathcal{H}_Q$, since they are pairwise nonassociate primes. Its quotient by this product is invariant: apply g and cancel the nonzero invariant product in the integral domain. Conversely each $\mathcal{H}_Q$ times an invariant polynomial is invariant and vanishes modulo Q . Consequently $\ker(\mathcal{J}_d \rightarrow J_d) = \mathcal{H}_Q \mathcal{J}_{d-10}$, and injection of multiplication by the nonzero product gives $\dim J_d = d_0(d) - d_0(d - 10)$. Injection of multiplication by A , OCF7 and OCF17 then prove

$$j_d := \dim \bar{J}_d = d_0(d) - d_0(d - 10) - \max(d - 7, 0)^2. \quad (\text{OCF19})$$

In particular $j_0 = 1$, $j_1 = 0$, $j_5 = 12$, and $j_d = 8d - 32$ for $d \geq 8$. For the last formula at $d \geq 10$, $\epsilon_d = \epsilon_{d-10}$ and direct expansion gives $\binom{d+3}{3} - \binom{d-7}{3} = 5(d^2 - 6d + 17)$; subtracting $(d - 7)^2$ gives $8d - 32$. At $d = 8, 9$ the ranks before this subtraction are 33, 44, respectively, which gives 32, 40 and verifies both remaining cases.

Here is a complete finite construction of all the columns used by OCF16. Start with $Z_0 = (1)$ and the invariant polynomial certificate $F_{0,1} = 1$. Suppose all earlier matrices Z_b have independent columns spanning $\bar{J}_b$, and their columns are $N_b \psi_b(F_{b,h})$ for invariant polynomials $F_{b,h}$. Form the candidate matrix

$$C_d = [m_{\nu,d} Z_{d-e}]_{\nu \in G_e, 2 \leq e \leq 5, e \leq d}. \quad (\text{OCF20})$$

Each candidate column has the explicit invariant certificate $x^\nu F_{d-e,h}$. OCF16 proves that C_d has column image $\bar{J}_d$ and rank j_d . Select j_d columns and j_d rows giving a nonzero determinant. Let Z_d be those columns, let I_d be those row positions, and let T_d be the complementary row positions

in E_d^{strip} . Retain the corresponding invariant certificates. This completes the induction. When $j_d = 0$, use the empty matrix with zero columns, empty I_d , and $T_d = E_d^{\text{strip}}$.

Every selected determinant and every matrix entry is an explicit expression obtained from the original entries of $\mathbf{H}$ and the full a_{rs} . To cover all cases, order the finite choices of column and row subsets and select the first nonzero determinant; the corresponding chart asserts its nonvanishing and the vanishing of all preceding candidates. A nonzero determinant exists because the rank was proved in OCF19. At rank zero no determinant is needed. Together with the exhaustive leading-coefficient charts after OCF8, this gives a finite atlas for each specified k . It requires no generic coefficient premise and supplies no numerical zero test for an unevaluated analytic coefficient: each chart is specified by its exact minors.

Define $\rho_d : W_d \rightarrow \mathbb{C}^{T_d}$ and the full original coefficient map π_d by

$$\rho_d = \text{row}_{T_d} - (Z_d)_{T_d} (Z_d)_{I_d}^{-1} \text{row}_{I_d}, \quad \pi_d = \rho_d N_d. \quad (\text{OCF21})$$

At rank zero the second term is the zero map. If E_{T_d} inserts a column into its T_d positions in W_d , then

$$\begin{aligned} w &= Z_d (Z_d)_{I_d}^{-1} w_{I_d} + E_{T_d} \rho_d w, \\ \rho_d E_{T_d} &= I, \quad \ker \rho_d = \text{im } Z_d = \bar{J}_d, \\ \ker \pi_d &= J_d, \quad \pi_d S_d E_{T_d} = I_{\mathbb{C}^{T_d}}. \end{aligned} \quad (\text{OCF22})$$

For the first line, the subtraction on its right leaves zero I_d rows and exactly $\rho_d w$ on the complementary rows. This proves the first three claims. If $\pi_d f = 0$, then $N_d f \in \bar{J}_d$; choose $j \in J_d$ with $N_d j = N_d f$. Now $f - j \in \ker N_d = \mathcal{A}\mathcal{B}_{d-8} \subseteq J_d$ by OCF17, so $f \in J_d$. The reverse inclusion follows immediately from $N_d J_d = \bar{J}_d$. The last equality uses $N_d S_d = I$. This proves every map and kernel in OCF22.

At the original degrees, these are the exact dimensions

k	n_k	j_k	$ T_k = n_k - j_k$
5	36	12	24
$k \geq 9, k = 4l + 1$	$16k - 48$	$8k - 32$	$8k - 16$.

(OCF23)

The original map $\Theta_k : R_k \xrightarrow{\sim} E_k^\vee$ in OQC7 sends the coefficient basis OCF2 to the original dual primary basis in the same order. Indeed its evaluation on $e^{wY_k}[1]$ is $\exp(((2b-k)\gamma - i(2a-k)\delta)w)$. The corresponding eigenvalue of A_k is $\beta_{ab} = k/2 + (2a-k)\delta + i(2b-k)\gamma$; the eigenvalue of $Y_k = (A_k - kI/2)/i$ is $\omega_{ab} = (\beta_{ab} - k/2)/i = (2b-k)\gamma - i(2a-k)\delta$. These eigenvalues are distinct because $\delta > 0$ and $\gamma > 0$. Their exponentials are independent by the Vandermonde determinant, so the evaluation identifies the covectors exactly. By the original observation identity OQC8, $\Theta_k J_k = \text{im } \Lambda_k^\vee$. Therefore OCF22 supplies the specified isomorphism

$$E_k^\vee / \text{im } \Lambda_k^\vee \xrightarrow{\pi_k} \mathbb{C}^{T_k} \cong (\ker \Lambda_k)^\vee. \quad (\text{OCF24})$$

The displayed π_k uses exactly the original primary coefficient coordinates under Θ_k . Every dualization in this statement is complex-linear. No conjugation is inserted in this coefficient map; the original metric maps can subsequently be composed with it.

Finally, this construction is finite with explicitly bounded compressed matrices. OCF20 has n_d rows and $\sum_{e=2}^5 |G_e| j_{d-e}$ columns, with negative subscripts zero. Both dimensions grow at most linearly in d : for example $n_d \leq 16d + 1$ and the column number is at most $25(16d + 1)$. For $d \geq 13$ the exact column number is $200d - 1632$, obtained from $\sum_e |G_e| = 25$, $\sum_e e|G_e| = 104$, and $j_{d-e} = 8(d-e) - 32$. Only the preceding five graded bases are needed to form the next matrix. Each column is obtained from the at-most-36-coefficient formula OCF12, one exact convolution, and OCF6. A full original biform coefficient column, if used as temporary storage, has $(d+1)^2$ entries; every invariant relation matrix in OCF20 has the stated linear row and column counts. The invariant certificates can be retained as their products and parent-column references, so their expanded high-degree monomials need never be enumerated. The proof of their completeness is OCF11–16, and the exact original quotient and its right inverse are OCF21–24.

10.6 The same recursion with fourteen proved generators

The complete fixed set OCF9 can be reduced to the following fourteen specific members while preserving every map and image just proved:

e	$\mathcal{G}_e$	
2	x_1x_4, x_2x_3	
3	$x_1^2x_3, x_1x_2^2, x_2x_4^2, x_3^2x_4$	(OCF25)
4	$x_1^3x_2, x_1x_3^3, x_2^3x_4, x_3x_4^3$	
5	$x_1^5, x_2^5, x_3^5, x_4^5$	

Here is a complete proof. Call a positive-degree invariant monomial indecomposable when it is not the product of two positive-degree invariant monomials. The divisor proof preceding OCF11 shows that each indecomposable has degree at most five. If it contains both x_1, x_4 , it is their product, since otherwise division by this invariant quadratic leaves a nonconstant invariant monomial. The same assertion holds for x_2, x_3 . After these cases are removed, at most two variables occur: every three indices contain one of the two opposite pairs. One variable gives precisely its fifth power. The remaining two-element supports are $\{1, 2\}, \{1, 3\}, \{2, 4\}, \{3, 4\}$. For positive exponents (a, b) with $a + b \leq 5$, the equation $a + 2b \equiv 0 \pmod{5}$ has exactly $(a, b) = (3, 1), (1, 2)$, and $a + 3b \equiv 0 \pmod{5}$ has exactly $(a, b) = (2, 1), (1, 3)$. For $\{2, 4\}$ divide the weights by 2 in the field $\mathbb{Z}/5\mathbb{Z}$, giving the first list; for $\{3, 4\}$ divide by 3, giving the second. These are exactly the eight mixed monomials of degrees three and four in OCF25. Every listed monomial has no proper invariant divisor, as is also seen from these lists and the excluded opposite pairs. This proves the classification. Factoring any invariant monomial repeatedly into indecomposables terminates because positive total degree strictly decreases. Hence OCF25 generates the entire invariant algebra.

Use $\mathcal{G} = \bigcup_{e=2}^5 \mathcal{G}_e$ in place of $\{x^\nu : \nu \in G\}$ in OCF12–16 and OCF20. The same factorization argument proves the same exact invariant image; all coefficient formulas, invariant certificates, pivot charts, quotient maps and ranks remain valid. The new recursion has 2, 4, 4, 4 generators in degrees 2, 3, 4, 5, respectively. Thus for $d \geq 13$ its precise matrix dimensions and rank are

$$C_d^{(14)} \in \text{Mat}_{(16d-48) \times (112d-864)}(\mathbb{C}), \quad \text{rank } C_d^{(14)} = 8d - 32. \quad (\text{OCF26})$$

Indeed the number of generators is 14 and the sum of their degrees is $2 \cdot 2 + 4 \cdot 3 + 4 \cdot 4 + 4 \cdot 5 = 52$; substituting the already proved $j_{d-e} = 8(d-e) - 32$ gives $14(8d-32) - 8 \cdot 52 = 112d - 864$. At smaller degrees the exact number of columns is the same sum of the appropriate j_{d-e} from OCF19, with negative subscripts zero. All coefficients in this reduced recursion are still the original coefficients of OCF12–14.

11 The exact common kernel frame and its original Gamma metrics

Retain the original simple quartet $m = 1$, $k = 4l + 1$, $l \geq 1$, $q = (k + 1)^2$, $c = k/2$, $0 < \delta < 1/2$ and $\gamma > 2$. The original period parameter and its branch are fixed, on the proved original OCF rank domain $|u| \geq R_*$, with R_* the retained explicit period-orbit radius. The kernel below is the kernel of that one fixed original observation $\Lambda : E \rightarrow B$. It is common to the two source metrics and to their four degree endpoints. No intersection over different period parameters or different observations is made.

Order all pairs (a, b) , $0 \leq a, b \leq k$, lexicographically once and retain that order. Define

$$\begin{aligned} \beta_{ab} &= c + (2a - k)\delta + i(2b - k)\gamma, & \omega_{ab} &= \frac{\beta_{ab} - c}{i} = (2b - k)\gamma - i(2a - k)\delta, \\ \chi(S) &= \prod_{a,b=0}^k (S - \beta_{ab}), & E &= \mathbb{C}[S]/(\chi), & \varepsilon_{ab} &= \left[\prod_{(a',b') \neq (a,b)} \frac{S - \beta_{a'b'}}{\beta_{ab} - \beta_{a'b'}} \right], & (\text{OKM1}) \\ V_\beta &= ((\beta_{ab})^d)_{(a,b), 0 \leq d < q}. \end{aligned}$$

The points are distinct: equality of their real and imaginary parts forces both indices equal. Hence every denominator displayed above is nonzero. Evaluation at all points gives $\varepsilon_{ab}(\beta_{a'b'}) = \mathbf{1}_{(a,b)=(a',b')}$. Polynomial interpolation proves that these are a basis of E , and the evaluation map from increasing polynomial coordinates to this primary basis is exactly V_β . Its determinant is the ordered Vandermonde product, so it is invertible.

11.1 The exact dual and primal maps of the strip quotient

Use the coefficient presentation of OQC7–8 in Segre coordinates. A biform of bidegree (k, k) has the full coefficient expression

$$F = \sum_{a,b=0}^k f_{ab} X_1^a X_0^{k-a} Y_1^b Y_0^{k-b}. \quad (\text{OKM2})$$

The literal variable dictionary to the OCF biform coefficients is $X_1 = z_0$, $X_0 = z_1$, $Y_1 = w_0$, $Y_0 = w_1$. The Segre substitution has $t_{++} = X_1 Y_1$, $t_{+-} = X_1 Y_0$, $t_{-+} = X_0 Y_1$, $t_{--} = X_0 Y_0$. At the original exponential vector take $X_1 = e^{-i\delta w}$, $X_0 = e^{i\delta w}$, $Y_1 = e^{\gamma w}$, $Y_0 = e^{-\gamma w}$. Then the monomial with coefficient f_{ab} is $e^{\omega_{ab} w}$. The original coefficient identification includes the entire invertible matrix H of OQC1 before this Segre presentation; its unit and period entries remain in the matrices produced by the strip calculation. OQC7 is characterized by evaluation on $e^{wY}[1]$. Since $e^{wY}[1] = \sum_{a,b} e^{\omega_{ab} w} \varepsilon_{ab}$, the independent exponentials in OQC4 therefore give the precise complex-linear dual pairing

$$\Theta(F)(\varepsilon_{ab}) = f_{ab}, \quad \Theta(F)(x) = f^\top x_{\text{prim}}. \quad (\text{OKM3})$$

There is no binomial coefficient or complex conjugation in this complex-linear dual pairing. The statement follows by equating all coefficients of the distinct exponentials, whose independence is proved by their Vandermonde derivative matrix in OQC4.

Let J be exactly the coefficient subspace of the original observed dual image, as identified in OQC8. The original strip and invariant elimination provider OCF constructs its literal surjection

$$\pi : \mathbb{C}^q \twoheadrightarrow \mathbb{C}^{r_K}, \quad \ker \pi = J, \quad r_K = 8k - 16. \quad (\text{OKM4})$$

This input is the actual OCF matrix, including its stated period coefficients and its valid pivot choice. The coefficient pairing OKM3 and OQC8 identify $K := \ker \Lambda = J^\perp$ in primary coordinates, where $J^\perp = \{x : f^\top x = 0 \text{ for every } f \in J\}$. Consequently its exact primal frame and polynomial frame are

$$I_{\text{prim}} = \pi^\top : \mathbb{C}^{r_K} \hookrightarrow E_{\text{prim}}, \quad I_{\text{poly}} = V_\beta^{-1} \pi^\top : \mathbb{C}^{r_K} \hookrightarrow E. \quad (\text{OKM5})$$

Here is a direct proof, including inverse coordinates. Choose any of the explicit right sections L_π of the surjective matrix π , so $\pi L_\pi = I_{r_K}$. For $x = \pi^\top v$ and $f \in J$, $f^\top x = (\pi f)^\top v = 0$. The transpose is injective because $L_\pi^\top \pi^\top = I_{r_K}$. Conversely, if $x \in J^\perp$, then $(I - L_\pi \pi)f \in J$ for every f , so $x^\top (I - L_\pi \pi)f = 0$ for every f . Thus $x = \pi^\top L_\pi^\top x$. This proves equality of the image and K , and its inverse on that image is $L_\pi^\top$. In polynomial coordinates the inverse is $L_\pi^\top V_\beta$. This proves the full typed primal arrow, rather than selecting a Hermitian transpose in place of it.

11.2 The complete original source and quotient metric

For either original source $s \in \{1, k\}$ write $b_s = s/2$ and $M_s = (2\pi)^{s/2}$. The unchanged real-coordinate measure is

$$dm_{0,s}(y) = M_s \frac{2^{b_s}}{4\pi\Gamma(b_s)} |\Gamma(b_s/2 + iy/2)|^2 dy, \quad S = c + iy. \quad (\text{OKM6})$$

The complete canonical OGN1–3 proof gives the monic real orthogonal polynomials and all their squared norms:

$$\begin{aligned} p_0(y) &= 1, \quad p_1(y) = y, \quad p_{d+1}(y) = yp_d(y) - d(b_s + d - 1)p_{d-1}(y), \\ h_d &= M_s d! (b_s)_d, \quad u_d(S) = \frac{p_d((S - c)/i)}{\sqrt{h_d}}. \end{aligned} \quad (\text{OKM7})$$

The square root is positive. Thus u_d has the original leading coefficient $i^{-d}/\sqrt{h_d}$ in S and is orthonormal on the line in OKM6. No source mass or phase has been removed.

For a polynomial cutoff $D \geq q - 1$ let

$$\begin{aligned} T_D &= \left(\frac{p_d(\omega_{ab})}{\sqrt{h_d}} \right)_{(a,b), 0 \leq d \leq D}, \quad R_D = T_D T_D^*, \\ G_D^{\text{prim}} &= R_D^{-1}, \quad G_D^{\text{poly}} = V_\beta^* R_D^{-1} V_\beta, \quad H_D^K = \bar{\pi} R_D^{-1} \pi^\top. \end{aligned} \quad (\text{OKM8})$$

The star denotes conjugate transpose, while the superscript $\top$ is ordinary transpose. In particular $(\pi^\top)^* = \bar{\pi}$, with precisely the indicated matrix dimensions.

These are the actual minimum quotient metrics, as follows directly from the original remainder map. In the orthonormal coefficient frame $u_0, \dots, u_D$, the remainder map followed by primary evaluation is exactly T_D , column by column. Its first q columns are invertible: the monic polynomials $p_0, \dots, p_{q-1}$ evaluated at the distinct ω_{ab} have a nonzero Vandermonde determinant, and all $\sqrt{h_d}$ are nonzero. Thus $R_D > 0$. For a prescribed primary vector x , the vector $T_D^* R_D^{-1} x$ maps to x . It is orthogonal to $\ker T_D$, because its inner product with a kernel vector contains T_D applied to that vector. Every other lift differs by such a vector, so its squared norm is $x^* R_D^{-1} x$ plus the squared norm of that difference. This proves the first quotient metric in OKM8. The identity $x_{\text{prim}} = V_\beta x_{\text{poly}}$ proves the second by direct pairing. Restriction to the exact frame OKM5 then gives

$$I_{\text{poly}}^* G_D^{\text{poly}} I_{\text{poly}} = (\pi^\top)^* R_D^{-1} \pi^\top = H_D^K > 0. \quad (\text{OKM9})$$

The last positivity follows from injectivity of $\pi^\top$. The map $K \hookrightarrow E$ is the fixed original coefficient map. No invariance of that kernel under multiplication is used.

11.3 Christoffel–Darboux with the exact source mass

For arbitrary $z, w \in \mathbb{C}$ and $D \geq 0$, the complete polynomial identity is

$$\sum_{d=0}^D \frac{p_d(z) p_d(w)}{h_d} = \begin{cases} \frac{p_{D+1}(z) p_D(w) - p_D(z) p_{D+1}(w)}{h_D(z - w)}, & z \neq w, \\ \frac{p'_{D+1}(z) p_D(z) - p'_D(z) p_{D+1}(z)}{h_D}, & z = w. \end{cases} \quad (\text{OKM10})$$

To prove it, put $a_d = d(b_s + d - 1)$, with $a_0 = 0$, and set $p_{-1} = 0$ for this recurrence expression. The recurrence in OKM7 gives

$$(z - w)p_d(z)p_d(w) = p_{d+1}(z)p_d(w) - p_d(z)p_{d+1}(w) - a_d(p_d(z)p_{d-1}(w) - p_{d-1}(z)p_d(w)). \quad (\text{OKM11})$$

For $d \geq 1$, the full norm identity is $a_d/h_d = 1/h_{d-1}$. Divide OKM11 by h_d and sum. Every adjacent term cancels, leaving just the degree- D numerator divided by h_D ; the $d = 0$ lower term is zero. This proves the first line of OKM10. Both sides of the resulting equality after multiplication by $z - w$ are polynomials. Differentiating with respect to z and then setting $z = w$ proves the second line with the displayed order and sign. This also proves the value at every confluent point without an assertion about a limiting singular quotient. The denominator is the full $h_D = M_s D!(b_s)_D$, including M_s .

Since all coefficients of the recurrence are real, the matrix entries in OKM8 are therefore exactly the values of OKM10 at

$$z = \omega_{ab}, \quad w = \overline{\omega_{a'b'}}. \quad (\text{OKM12})$$

Their indexed conjugation and confluent locus are

$$\begin{aligned} \overline{\omega_{a'b'}} &= \omega_{k-a',b'}, & \omega_{ab} - \overline{\omega_{a'b'}} &= 2(b - b')\gamma - 2i(a + a' - k)\delta, \\ \omega_{ab} &= \overline{\omega_{a'b'}} & \iff & b = b' \text{ and } a + a' = k. \end{aligned} \quad (\text{OKM13})$$

These follow by substituting OKM1; positivity of δ, γ proves both implications in the last line. Thus the conjugation map on the point indices is precisely $(a, b) \mapsto (k - a, b)$. The confluent formula of OKM10 applies at these indexed entries. For the retained odd k , its matrix diagonal has $a' = a, b' = b$, so $2a = k$ cannot occur; the literal Hermitian diagonal is given by the first line of OKM10. The conjugate-index entries use its second line. All entries, including the positive diagonal, are simultaneously defined by the original polynomial sum on the left of OKM10.

11.4 Every pivot change and the four original endpoints

Let π' be another valid strip/invariant elimination matrix for the same original J , with the same surjective codomain dimension. Choose right sections $L_\pi, L_{\pi'}$ and put

$$A_{\pi',\pi} = \pi' L_\pi, \quad A_{\pi,\pi'} = \pi L_{\pi'}. \quad (\text{OKM14})$$

Because π' vanishes on $J = \ker \pi$, one has $\pi' L_\pi \pi = \pi'$. Thus $\pi' = A_{\pi',\pi} \pi$. Reversing the roles gives the opposite identity, and surjectivity of each matrix then proves $A_{\pi',\pi} A_{\pi,\pi'} = I$ and $A_{\pi,\pi'} A_{\pi',\pi} = I$. This proves invertibility and also independence from the chosen right sections. Writing $A = A_{\pi',\pi}$ for these identities gives the exact frame and Gram changes

$$\begin{aligned} I'_{\text{prim}} &= I_{\text{prim}} A^\top, & I'_{\text{poly}} &= I_{\text{poly}} A^\top, \\ (H_D^K)' &= \overline{A} H_D^K A^\top, & \det(H_D^K)' &= |\det A|^2 \det H_D^K. \end{aligned} \quad (\text{OKM15})$$

The first two formulas follow by transposing $\pi' = A\pi$. Substitution into OKM9 proves the congruence and determinant. This retains all actual pivot determinants. They have not been assigned modulus one.

The kernel is the same fixed $K = \ker \Lambda$ used in MRI1–5. If its original BRU inclusion is I_0 and its fixed-section kernel coordinate map is κ_0 , let

$$C_K = \kappa_0 I_{\text{poly}}, \quad C_K^{-1} = L_\pi^\top V_\beta I_0. \quad (\text{OKM16})$$

The equalities $I_0 \kappa_0 x = x$ for $x \in K$ and $L_\pi^\top V_\beta I_{\text{poly}} = I_{r_K}$ prove that these are inverse matrices and $I_0 C_K = I_{\text{poly}}$. Consequently $H_D^K = C_K^* (I_0^* G_D^{\text{poly}} I_0) C_K$. The matrix C_K is independent of D and

of the choice of source order s . The source metrics change; the two specified coefficient frames do not.

It follows that the complete original four-endpoint kernel scalar is

$$\boxed{F_K^{(s)} = \log \det H_{q-1}^K + \log \det H_q^K - \log \det H_{2q-1}^K - \log \det H_{2q}^K.} \quad (\text{OKM17})$$

Indeed MRI3 gives the actual consecutive kernel determinant ratio

$$\frac{\det(I_0^* G_{q+j-1}^{\text{poly}} I_0)}{\det(I_0^* G_{q+j}^{\text{poly}} I_0)} = \frac{1 + t_j}{1 + \xi_j}.$$

Each factor $|\det C_K|^2$ cancels in that ratio by OKM16. The weighted sum with endpoint weights $1, 2, \dots, 2, 1$ therefore has precisely the four signs in OKM17, as in MRI4. This verifies the original degree cutoffs, rather than shifting the column count in T_D . Under any valid pivot change OKM15, the added scalar $2 \log |\det A|$ has total coefficient $1 + 1 - 1 - 1 = 0$, proving invariance of OKM17 with its original frames.

For emphasis, the full mass in OKM7 appears in every column of T_D and in the denominator of OKM10. If that common mass is factored explicitly, write $T_D = M_s^{-1/2} \hat{T}_D$, where $\hat{T}_D$ has exactly the same numerator polynomials and the denominator $\sqrt{d!(b_s)_d}$. Then $H_D^K = M_s \bar{\pi} (\hat{T}_D \hat{T}_D^*)^{-1} \pi^\top$, so its determinant has factor $M_s^{r_K}$. It cancels only in the displayed signed four-endpoint scalar, whose four ranks are equal. The individual metrics in OKM8–9 retain it. No inserted Gamma order, discarded action defect, or kernel-invariance assertion enters any of these calculations.

12 The original kernel graph, boundary basis and complete action

This section uses the complete conductor coefficient construction OCF and its exact original dual identification in OKM. Fix the original simple quartet, its complete amplitude unit and periods, the original parameter on any logarithm branch with $|u| \geq R_*$, and $k = 4l + 1$, $l \geq 1$. Put $q = (k + 1)^2$, $r = 8k - 16$, and $\mathcal{I}_k = \{(a, b) : 0 \leq a, b \leq k\}$. The roots, centre and centred eigenvalues remain

$$c = k/2, \quad \beta_{ab} = c + (2a - k)\delta + i(2b - k)\gamma, \quad \omega_{ab} = (\beta_{ab} - c)/i = (2b - k)\gamma - i(2a - k)\delta. \quad (\text{OKA1})$$

The original algebra is $E = \mathbb{C}[S]/(\chi)$, with $\chi(S) = \prod_{(a,b) \in \mathcal{I}_k} (S - \beta_{ab})$. Let ε_{ab} be its original primary idempotents and $\mathcal{V} : E \rightarrow \mathbb{C}^{\mathcal{I}_k}$ the evaluation isomorphism $[p] \mapsto (p(\beta_{ab}))_{ab}$. Its inverse sends e_{ab} to

$$\varepsilon_{ab}(S) = \frac{\chi(S)}{(S - \beta_{ab})\chi'(\beta_{ab})}. \quad (\text{OKA2})$$

These denominators are nonzero: equality of two roots in OKA1, on comparing real and imaginary parts and using $\delta > 0$, $\gamma > 2$, implies equality of both indices. Evaluation proves that the polynomials in OKA2 are the coordinate idempotents, so their columns give the exact inverse, with no changed root or polynomial coefficient convention.

12.1 The coefficient quotient gives a graph in the original primary basis

Let $\pi : \mathbb{C}^{\mathcal{I}_k} \rightarrow \mathbb{C}^r$ be the surjective coefficient quotient of OCF, whose kernel is precisely the image J_k of the invariant polynomials under the original restriction. The domain here consists of the coefficients of the biforms $z_0^a z_1^{k-a} w_0^b w_1^{k-b}$, and is the complex-linear dual of the displayed primary coordinate space. Let $T \subset \mathcal{I}_k$ be the final OCF complement of its invariant-image pivot rows, viewed as a subset of the full biform coefficient grid. Its ordered coordinate inclusion is $E_T : \mathbb{C}^r \rightarrow \mathbb{C}^{\mathcal{I}_k}$. The explicit OCF formula gives

$$\pi E_T = I_r, \quad N = \pi^T : \mathbb{C}^r \longrightarrow \mathbb{C}^{\mathcal{I}_k}, \quad E_T^T N = I_r. \quad (\text{OKA3})$$

Indeed a retained strip monomial in T already has zero conductor remainder outside its own coordinate, zero coordinates on the invariant pivot rows, and its indicated coordinate equal to one. Both reductions in OCF therefore leave that quotient coordinate equal to its standard unit vector. Transposing gives the last identity. This uses ordinary transpose, because the restriction quotient and its dual are complex-linear maps.

Write $J = \mathcal{I}_k \setminus T$ only for the complementary index set in this section; the invariant image retains the name J_k . Denote coordinate restrictions by x_T, x_J , and put $N_J = E_J^T N$. The original kernel inclusion is consequently

$$I_K = \mathcal{V}^{-1} N : \mathbb{C}^r \hookrightarrow E, \quad \text{im } I_K = K := \ker \Lambda, \quad N = \begin{pmatrix} I_r \\ N_J \end{pmatrix}_{T,J}. \quad (\text{OKA4})$$

For completeness, a primary vector v pairs with a biform coefficient vector f by $f^T v$. The annihilator of $\ker \pi = J_k$ is $\text{im } \pi^T$: containment follows by multiplication, while surjectivity of π gives both subspaces dimension r . OQC8 identifies precisely this annihilator with the primary coordinates of the fixed original $\ker \Lambda$. This proves both the original image assertion and its direction in OKA4. In polynomial coordinates the same inclusion is $V_\beta^{-1} N$, where $(V_\beta)_{(a,b),d} = \beta_{ab}^d$ for $0 \leq d < q$; this is the full coefficient matrix of OKA2.

12.2 An explicit basis of the actual boundary

For $j \in J$ define the actual boundary vector $b_j = \Lambda \varepsilon_j$. These vectors form a basis of the original image $B = \text{im } \Lambda$. To prove independence, if $\sum_{j \in J} c_j \varepsilon_j$ lies in K , its primary coordinates equal $N \zeta$.

Its T coordinates vanish, so OKA3 gives $\zeta = 0$ and all $c_j = 0$. To prove spanning, use the literal decomposition

$$x = Nx_T + E_J(x_J - N_Jx_T), \quad \Lambda\mathcal{V}^{-1}x = \sum_{j \in J} b_j(x_J - N_Jx_T)_j. \quad (\text{OKA5})$$

The first equality follows separately on its T and J coordinates. Apply Λ , which kills the first term by OKA4, to obtain the second. Thus no observation has been substituted: in this basis the specified original observation has the exact coefficient matrix $(-N_J, I_{q-r})$.

Here are the complete original period and tensor coordinates of these boundary vectors. Write the quartet roots as $\alpha_{\epsilon, \eta} = 1/2 + \epsilon\delta + i\eta\gamma$, where $\epsilon, \eta \in \{+1, -1\}$ and the signs $+, -$ in subscripts denote these same two numbers. In a subscript α below means that complex root, indexed by its sign pair. Keep the original one-factor primary idempotents $\varepsilon_\alpha^h \in E_h$. Set

$$v_\alpha = \Pi U \varepsilon_\alpha^h = v_h(\alpha) \Pi \varepsilon_\alpha^h, \quad Z_+ = z_0, \quad Z_- = z_1, \quad W_+ = w_0, \quad W_- = w_1, \quad \mathbf{v}(z, w) = \sum_{\epsilon, \eta} v_{\epsilon\eta} Z_\epsilon W_\eta. \quad (\text{OKA6})$$

The vectors belong to the original one-factor period space; all entries of Π and all four actual nonzero amplitude values remain present. For every $(a, b) \in \mathcal{I}_k$,

$$j\Lambda\varepsilon_{ab} = [z_0^a z_1^{k-a} w_0^b w_1^{k-b}] P_k(\mathbf{v}(z, w)^{\otimes k}), \quad P_k = \frac{1}{5} \sum_{h=0}^4 (C^{-h})^{\otimes k}. \quad (\text{OKA7})$$

The original sum inclusion $\iota : E \hookrightarrow E_h^{\otimes k}$ sends ε_{ab} to the sum of the primary tensor idempotents whose word has a positive real signs and b positive imaginary signs. This assertion follows by evaluation: on a primary word, the sum variable takes exactly the value β_{ab} of its two counts, and OKA2 evaluates to one for those counts and zero otherwise. Applying $(\Pi U)^{\otimes k}$ gives precisely the corresponding sum of the tensors of the v_α in OKA6. Expanding the tensor power and taking the indicated biform coefficient selects exactly the same words, each once. The original definition $j\Lambda = P_k(\Pi U)^{\otimes k} \iota$ proves OKA7, including the literal inverse powers and $1/5$. Its specialization to $j \in J$ therefore supplies the actual basis b_j of OKA5. Neither a different target nor a numerical period matrix has been introduced.

The fixed, source-independent kernel coordinate and section are now explicitly

$$\kappa = E_T^T \mathcal{V} : E \rightarrow \mathbb{C}^r, \quad S_* : B \rightarrow E, \quad S_* b_j = \varepsilon_j, \quad (\text{OKA8})$$

$$I_K \kappa + S_* \Lambda = I_E, \quad \kappa I_K = I_r, \quad \kappa S_* = 0, \quad \Lambda S_* = I_B.$$

Every equality follows by applying OKA5 to an arbitrary primary vector. In particular this is an actual splitting of the fixed original short exact sequence, used unchanged for both source orders and for all polynomial degrees.

12.3 The whole original action and its retained defect

The original centred sum action is $Y = (S - c)/i$ on E . Under $\mathcal{V}$ it is the diagonal matrix of the ω_{ab} in OKA1. Write its two diagonal submatrices as Ω_T and Ω_J . In the exact coordinates $e = I_K \zeta + S_* b$ of OKA8, it has the complete matrix

$$Y \longleftrightarrow \begin{pmatrix} \Omega_T & 0 \\ \Omega_J N_J - N_J \Omega_T & \Omega_J \end{pmatrix}, \quad (\Omega_J N_J - N_J \Omega_T)_{jt} = (\omega_j - \omega_t)(N_J)_{jt}. \quad (\text{OKA9})$$

Indeed the primary vector of e is $(\zeta, N_J \zeta + b)$. Applying the original diagonal matrix gives $(\Omega_T \zeta, \Omega_J N_J \zeta + \Omega_J b)$. Applying the coordinates in OKA5 subtracts $N_J \Omega_T \zeta$ from its second component and proves every block in OKA9. This proves, with its precise codomain,

$$Y I_K - I_K \Omega_T = S_* (\Omega_J N_J - N_J \Omega_T) : \mathbb{C}^r \rightarrow E, \quad \Lambda Y I_K = \Omega_J N_J - N_J \Omega_T : \mathbb{C}^r \rightarrow B, \quad (\text{OKA10})$$

where the matrix on the right uses the basis b_j . The chosen complementary span of the original primary idempotents is stable under Y ; no stability of K has been used. Its entire failure is the displayed coefficient map.

For each original source $s \in \{1, k\}$ retain the monic Gamma polynomials $p_d^{(s)}$, the complete norms $h_d^{(s)} = (2\pi)^{s/2} d!(s/2)_d$, and the original $e_d^{(s)} = [p_d^{(s)}(Y)]/\sqrt{h_d^{(s)}}$. Then the fixed-section kernel and boundary coordinates of every actual remainder, including all degrees $d \geq q$, are

$$\begin{aligned} (\kappa e_d^{(s)})_t &= \frac{p_d^{(s)}(\omega_t)}{\sqrt{h_d^{(s)}}}, \\ (\Lambda e_d^{(s)})_j &= \frac{p_d^{(s)}(\omega_j) - \sum_{t \in T} (N_J)_{jt} p_d^{(s)}(\omega_t)}{\sqrt{h_d^{(s)}}}. \end{aligned} \tag{OKA11}$$

Polynomial evaluation in the original primary algebra gives the full vector $p_d^{(s)}(\omega)/\sqrt{h_d^{(s)}}$ even when its polynomial representative must be reduced modulo χ . Applying OKA5 and OKA8 proves OKA11. Thus the formula includes the original finite remainder operation and every phase and source mass. Composing its second line with the actual columns OKA7 gives exactly the period-vector increments of OPG. Its first line is the fixed-section kernel vector used by OPG and OPR; the degree-dependent minimum-section correction in OPR remains required for the mixed norm. The metric computation in OKM applies to the common inclusion OKA4 and does not replace that correction by the fixed-section vector alone.

13 The complete homogeneous geometry of the original multiplicity curve

13.1 The actual primary, unit, period and Fourier coefficient map

Retain the original full-order quartet and every original period parameter. Write

$$\begin{aligned} \alpha_{\epsilon,\eta} &= \frac{1}{2} + \epsilon\delta + i\eta\gamma, \quad \epsilon, \eta \in \{-1, 1\}, \quad 0 < \delta < \frac{1}{2}, \quad \gamma > 2, \\ m &\geq 1, \quad d_m = m - 1, \quad n = 4m, \quad N = 4m + 1, \quad h(z) = \prod_{\epsilon,\eta} (z - \alpha_{\epsilon,\eta})^m, \\ E_h &= \mathbb{C}[z]/(h), \quad A_h = M_z, \quad v_h = 2\xi/h, \quad U = M_{j_h v_h}, \quad v_h(\alpha_{\epsilon,\eta}) \neq 0. \end{aligned} \quad (\text{OMG1})$$

The last inequalities follow from the stipulated complete original zero order m . The map j_h records all divided Taylor coefficients of orders $0, \dots, m-1$ at all four roots. Let $\mathcal{E}_{\text{prim}} : \mathbb{C}^{4m} \rightarrow E_h$ send its coordinate (α, d) to the original primary element $\varepsilon_\alpha(z - \alpha)^d$, $0 \leq d < m$. Explicitly its representative is

$$E_{\alpha,d}(z) = \text{rem}_h \left[h_\alpha(z) \sum_{j=0}^{m-1} \frac{(h_\alpha^{-1})^{(j)}(\alpha)}{j!} (z - \alpha)^{j+d} \right], \quad h_\alpha = h/(z - \alpha)^m.$$

The Taylor inverse is defined since $h_\alpha(\alpha) \neq 0$. It gives the idempotent one on its own primary factor and zero on the others. The four coprime factors therefore prove that $\mathcal{E}_{\text{prim}}$ is an isomorphism and include every nilpotent degree. In this basis the full matrix of U is

$$U_{\text{prim};\alpha,a;\alpha,d} = \begin{cases} v_h^{(a-d)}(\alpha)/(a-d)!, & a \geq d, \\ 0, & a < d, \end{cases} \quad \det U = \prod_{\alpha} v_h(\alpha)^m \neq 0. \quad (\text{OMG2})$$

Other primary blocks are zero. Inverting its constant coefficient plus its nilpotent part by the finite geometric series proves invertibility with all derivatives retained.

Keep the original $u \neq 0$ and its logarithm branch, the primitive $\Phi'_h = h$, $\Phi_h(0) = 0$, the outward rays ℓ_j of angles $(\arg u + \pi + 2\pi j)/N$, and $\Gamma_j = -\ell_0 + \ell_j$. The exact period and Fourier maps are

$$\begin{aligned} (\Pi[P])_j &= \int_{\Gamma_j} e^{\Phi_h(z)/u} \text{rem}_h P(z) dz, \quad 1 \leq j \leq n, \\ \zeta &= e^{2\pi i/N}, \quad F_{jt} = \zeta^{jt} - 1, \quad (F^{-1})_{tj} = \zeta^{-tj}/N \quad (1 \leq j, t \leq n), \quad C = F \text{diag}(\zeta, \dots, \zeta^n) F^{-1}. \end{aligned} \quad (\text{OMG3})$$

The full original period theorem PDE/OPG gives the convergence and invertibility of Π on these representatives. The Fourier inverse follows directly from $\sum_{j=1}^{N-1} \zeta^{-tj} (\zeta^{sj} - 1) = N \mathbf{1}_{t=s}$. It is the same literal cyclic matrix as OCS4, including its inverse powers in the original average. Its inherited Gram is $F^*F = N(I_n + \mathbf{1}\mathbf{1}^*)$, as expansion of its entries gives.

Define the exact diagonal map and coefficient matrix

$$\begin{aligned} D_m^{\text{jet}} &= \text{diag} \left(i^{-d}/d! \right)_{\alpha, 0 \leq d < m}, \quad \mathbf{H}_m = F^{-1} \Pi U \mathcal{E}_{\text{prim}} D_m^{\text{jet}}, \\ (\mathbf{H}_m)_{t;\alpha,d} &= \frac{i^{-d}}{d!} \sum_{a=d}^{m-1} \left[\frac{1}{N} \sum_{j=1}^n \zeta^{-tj} \int_{\Gamma_j} e^{\Phi_h(z)/u} E_{\alpha,a}(z) dz \right] \frac{v_h^{(a-d)}(\alpha)}{(a-d)!}. \end{aligned} \quad (\text{OMG4})$$

This is exactly OCS13, now regarded as one square matrix. All factors in its first line are invertible. In particular

$$\det \mathbf{H}_m = (\det F)^{-1} \det \Pi \left(\prod_{\alpha} v_h(\alpha)^m \right) \det \mathcal{E}_{\text{prim}} \prod_{\alpha} \prod_{d=0}^{m-1} \frac{i^{-d}}{d!} \neq 0,$$

with the determinant of $\mathcal{E}_{\text{prim}}$ taken in the fixed original coefficient frame. None of these factors is prescribed.

Set $\omega_{\epsilon,\eta} = \eta\gamma - i\epsilon\delta$ and let

$$z_{\epsilon,\eta,d}(w) = w^d e^{\omega_{\epsilon,\eta} w}, \quad x(w) = F^{-1} \Pi U e^{w(A_h - 1/2)/i} [1]. \quad (\text{OMG5})$$

The terminating exponential on each primary factor has coefficient $i^{-d} w^d / d!$ on $\varepsilon_\alpha(z - \alpha)^d$. Multiplication by OMG2 and application of the literal period maps therefore prove the exact identity and inverse coordinate map

$$\boxed{x(w) = H_m z(w), \quad z(w) = H_m^{-1} x(w).} \quad (\text{OMG6})$$

For the original Euclidean period metric, its Gram on these primary curve coordinates is $H_m^* N(I_n + \mathbf{11}^*) H_m$. Thus the coefficient change also retains its full metric and all off-diagonal entries, without claiming orthogonality.

13.2 The complete homogeneous ideal, with quadratic generators

Use binary indices $a, b \in \{0, 1\}$ for $\epsilon = 2a - 1$, $\eta = 2b - 1$. Let $\mathcal{P}_Z = \mathbb{C}[Z_{a,b,d} : a, b \in \{0, 1\}, 0 \leq d \leq d_m]$ with each variable of degree one. Define the algebra map

$$\Phi_m : \mathcal{P}_Z \longrightarrow \mathbb{C}[X_0, X_1, Y_0, Y_1, T_0, T_1], \quad Z_{a,b,d} \longmapsto X_a Y_b T_0^{d_m - d} T_1^d. \quad (\text{OMG7})$$

Its image is the complete diagonal graded algebra

$$R_m = \bigoplus_{k \geq 0} \mathbb{C}[X_0, X_1]_k \otimes \mathbb{C}[Y_0, Y_1]_k \otimes \mathbb{C}[T_0, T_1]_{kd_m}. \quad (\text{OMG8})$$

Indeed a basis monomial of its degree- k component is

$$X_0^{k-a} X_1^a Y_0^{k-b} Y_1^b T_0^{kd_m - D} T_1^D, \quad 0 \leq a, b \leq k, \quad 0 \leq D \leq kd_m. \quad (\text{OMG9})$$

It is a product of k images in OMG7: choose k binary first indices with sum a , independently k binary second indices with sum b , and k integers in $[0, d_m]$ with sum D ; the last choice follows by filling successive slots up to d_m . Combining the three lists into rows gives the required preimage monomial. Distinct displayed target monomials are linearly independent.

Let $\mathfrak{t}_m$ be generated by every quadratic binomial

$$Z_{a,b,d} Z_{a',b',d'} - Z_{c,e,f} Z_{c',e',f'} \quad \text{with} \quad a + a' = c + c', \quad b + b' = e + e', \quad d + d' = f + f', \quad (\text{OMG10})$$

where every index belongs to its original range. Repeated variables are allowed. Then

$$\boxed{\ker \Phi_m = \mathfrak{t}_m, \quad \mathcal{P}_Z / \mathfrak{t}_m \simeq R_m.} \quad (\text{OMG11})$$

Here is a complete ideal proof. Each displayed quadratic has zero image. Two degree- k monomials have the same image exactly when the sums of their first indices, second indices and third indices agree. Regard their k factors temporarily as labelled rows. Align the first indices with the target list by pairwise swaps of first indices, keeping each row's other indices fixed. Each swap is an OMG10 relation. Next align the second indices by the same operation, keeping the aligned first indices fixed. Finally, if the third indices differ from their target list, choose a row below its target and a row above its target. Increase the former third index by one and decrease the latter by one. These entries stay in $[0, d_m]$ because their target entries do; the total sum is unchanged and the sum of absolute deviations decreases by two. This terminates at the target list. Each transfer is another OMG10 relation. Thus the difference between any two monomials with the same image is a sum of multiples of the quadrics, by telescoping the successive monomial differences. Labels only specify the sequence of polynomial maps; the products are the original commutative monomials

throughout. For any polynomial in $\ker \Phi_m$, collect its coefficients by their target monomial. Each such coefficient sum is zero by target monomial independence. Subtracting one chosen monomial in each group expresses that whole group as a linear combination of the proved monomial differences. This proves the reverse inclusion for all degrees and hence the entire ideal equality.

Every degree- k original curve monomial has the function

$$w^D \exp\{[(2b - k)\gamma - i(2a - k)\delta]w\} \quad (0 \leq a, b \leq k, \quad 0 \leq D \leq kd_m). \quad (\text{OMG12})$$

The frequencies are distinct: equality of real parts gives equality of b and equality of imaginary parts gives equality of a . To prove independence with all polynomial powers, suppose $\sum_{a,b} P_{a,b}(w)e^{\omega_{a,b}w} = 0$, with $\deg P_{a,b} \leq kd_m$. Fix a pair and apply $\prod_{(a',b') \neq (a,b)} (\partial_w - \omega_{a',b'})^{kd_m+1}$. This kills every other summand. On its surviving polynomial each factor is $\partial_w + (\omega_{a,b} - \omega_{a',b'})$, invertible on the space of degree at most kd_m because differentiation is nilpotent and the constant is nonzero. The surviving polynomial is therefore zero. Repeating this for every pair proves independence.

Thus the full homogeneous ideal of the original primary curve is exactly $\mathfrak{t}_m$: a homogeneous polynomial restricts to zero if and only if its coefficient sums at every target monomial OMG9 are zero. In particular

$$\boxed{\dim R_{m,k} = (k+1)^2[1 + k(m-1)] = q_k, \quad \ker\{\mathcal{P}_{Z,k} \rightarrow \text{Hol}(\mathbb{C}), f \mapsto f(z(w))\} = \mathfrak{t}_{m,k}.} \quad (\text{OMG13})$$

This statement concerns the homogeneous ideal, equivalently the projective closure of $[z(w)]$. Every one of its degrees has been computed, including degree zero. No assertion about a different nonhomogeneous affine ideal is used.

13.3 The projective model and its complete chart inverses

For $m \geq 2$, the morphism

$$(\mathbb{P}^1)^3 \longrightarrow \mathbb{P}^{4m-1}, \quad ([X_0 : X_1], [Y_0 : Y_1], [T_0 : T_1]) \longmapsto [X_a Y_b T_0^{d_m-d} T_1^d]_{a,b,d} \quad (\text{OMG14})$$

has multidegree $(1, 1, m-1)$. At least one first, second and third coordinate is nonzero; their corresponding extreme product is nonzero, so the map is everywhere defined. Its image and scheme structure are exactly $\text{Proj } R_m$, with complete closed ideal OMG10. Here are explicit inverses proving this claim. The eight charts $Z_{a,b,0} \neq 0$ and $Z_{a,b,d_m} \neq 0$ cover $\text{Proj } R_m$. Indeed OMG11 gives

$$Z_{a,b,d}^{d_m} = Z_{a,b,0}^{d_m-d} Z_{a,b,d_m}^d \quad \text{in } R_m.$$

A homogeneous prime containing every extreme variable contains every variable by this equality, and hence gives no point of Proj . On $Z_{a,b,0} \neq 0$ the three inverse affine coordinates are

$$x = Z_{1-a,b,0}/Z_{a,b,0}, \quad y = Z_{a,1-b,0}/Z_{a,b,0}, \quad t = Z_{a,b,1}/Z_{a,b,0}.$$

For every variable its ratio to $Z_{a,b,0}$ is $x^{1_{a' \neq a}} y^{1_{b' \neq b}} t^d$, by OMG7 and its proved injective quotient. Consequently the degree-zero localized ring is $\mathbb{C}[x, y, t]$. These three coordinates are algebraically independent, since their images in the parameter fraction field are $X_{1-a}/X_a, Y_{1-b}/Y_b, T_1/T_0$. For the chart at Z_{a,b,d_m} replace the last ratio by $t = Z_{a,b,d_m-1}/Z_{a,b,d_m}$; the general ratio has exponent $d_m - d$ in t . Its coordinate ring is again $\mathbb{C}[x, y, t]$. On chart overlaps these formulas invert the same parameter ratios, so they glue to the inverse of OMG14. This proves the isomorphism of schemes and its full defining homogeneous ideal. For $m = 1$ the third factor has degree zero and the same argument uses the four first/second charts: the model is the Segre $(\mathbb{P}^1)^2$ with ideal $Z_{0,0,0}Z_{1,1,0} - Z_{0,1,0}Z_{1,0,0}$.

In the actual Fourier coordinates let $\mathcal{P}_x = \mathbb{C}[x_1, \dots, x_n]$ and define

$$\mathfrak{a} = \{f(x) : f(H_m Z) \in \mathfrak{t}_m\}, \quad R = \mathcal{P}_x / \mathfrak{a}. \quad (\text{OMG15})$$

The invertible substitution OMG6 proves that $\mathfrak{a}$ is the complete homogeneous ideal of $x(w)$ and that $R \simeq R_m$. Its explicit generators are all OMG10 quadrics with every Z replaced by its component of $H_m^{-1}x$. These are the actual period and unit coefficients of OMG4, and $\mathfrak{a}$ is prime because the target ring in OMG7 is a domain. Thus the original curve's projective closure is the precise linear image of OMG14 under H_m , with no prescribed substitute matrix.

13.4 The exact graded source-dual algebra and every factorial

For each $k \geq 0$ retain the original sum algebra

$$e_k = 1 + kd_m, \quad c_k = k/2, \quad \beta_{k,a,b} = c_k + (2a - k)\delta + i(2b - k)\gamma, \\ \chi_k(S) = \prod_{a,b=0}^k (S - \beta_{k,a,b})^{e_k}, \quad E_k = \mathbb{C}[S]/(\chi_k), \quad Y_k = (M_S - c_k I)/i. \quad (\text{OMG16})$$

For $k = 0$ this is $E_0 = \mathbb{C}[S]/(S) = \mathbb{C}$. Let $H_{k,a,b,D} = \varepsilon_{k,a,b}(S - \beta_{k,a,b})^D$, $0 \leq D < e_k$, be the complete original primary basis obtained by the same Taylor inverse as in OMG2. It satisfies

$$e^{wY_k}[1] = \sum_{a,b=0}^k e^{\omega_{k,a,b}w} \sum_{D=0}^{e_k-1} \frac{i^{-D}w^D}{D!} H_{k,a,b,D}, \quad \omega_{k,a,b} = (2b - k)\gamma - i(2a - k)\delta. \quad (\text{OMG17})$$

For $f \in \mathcal{P}_{x,k}$ expand its original function uniquely as $f(x(w)) = \sum c_{a,b,D}(f)w^D e^{\omega_{k,a,b}w}$ using OMG12. Define the complex linear dual map, without a conjugation,

$$\Theta_k(f)(H_{k,a,b,D}) = i^D D! c_{a,b,D}(f). \quad (\text{OMG18})$$

Then OMG17 proves exactly $\Theta_k(f)(e^{wY_k}[1]) = f(x(w))$. The independence and surjectivity in OMG9–13 prove that this map is onto $E_k^\vee$, with kernel $\mathfrak{a}_k$. Consequently it induces a specified vector-space isomorphism

$$\boxed{\Theta_k : R_k \xrightarrow{\sim} E_k^\vee}. \quad (\text{OMG19})$$

All phases and factorials are those of OCS17; for a monomial x^ν its coefficients are precisely the finite multinomial OCS15 rule with the entries OMG4.

The collection of these vector spaces has an exact graded algebra structure coming from the original sum maps. For $k, j \geq 0$ put

$$\Delta_{k,j} : E_{k+j} \longrightarrow E_k \otimes E_j, \quad [P(S)] \longmapsto [P(S_1 + S_2)]. \quad (\text{OMG20})$$

On a primary pair the eigenvalue is the displayed sum $\beta_{k,a,b} + \beta_{j,a',b'} = \beta_{k+j,a+a',b+b'}$. The sum of the two nilpotents has index $e_k + e_j - 1 = e_{k+j}$: its power $e_k + e_j - 2$ has nonzero top coefficient $\binom{e_k+e_j-2}{e_k-1}$ and its next power is zero. Every pair of total sign counts occurs. Thus the full minimal polynomial of $S_1 + S_2$ is exactly χ_{k+j} , proving both well-definedness and injectivity of OMG20. It also gives the complete primary formula

$$\Delta_{k,j} H_{k+j,A,B,D} = \sum_{a+a'=A, b+b'=B} \sum_{\substack{d+d'=D \\ 0 \leq d < e_k, 0 \leq d' < e_j}} \binom{D}{d} H_{k,a,b,d} \otimes H_{j,a',b',d'}. \quad (\text{OMG21})$$

Indeed the total primary idempotent is one precisely on the matching eigenvalue pairs, and the nilpotent power expands by the ordinary binomial theorem there. All other components vanish. Substitution of the sum of three coordinates proves coassociativity; interchange of the two coordinates proves cocommutativity. The $k = 0$ maps give the count. Hence

$$(\ell \star \ell')(v) = (\ell \otimes \ell')(\Delta_{k,j}v) \quad (\ell \in E_k^\vee, \ell' \in E_j^\vee) \quad (\text{OMG22})$$

is an associative commutative graded product with unit in $E_0^\vee$. For the exact dual basis $\lambda_{k,a,b,D} = i^D D! H_{k,a,b,D}^\vee$, OMG21 gives

$$\lambda_{k,a,b,D} \star \lambda_{j,a',b',D'} = \lambda_{k+j,a+a',b+b',D+D'}.$$

The coefficient is exactly $\binom{D+D'}{D} i^D D! i^{D'} D'! = i^{D+D'} (D+D')!$; this records the cancellation before stating the basis product. Equivalently, OMG20 sends $e^{wY_{k+j}}[1]$ to $e^{wY_k}[1] \otimes e^{wY_j}[1]$, with $c_{k+j} = c_k + c_j$. Evaluating OMG18 on this identity proves

$\Theta : R \xrightarrow{\sim} \bigoplus_{k \geq 0} (E_k^\vee, \star) \text{ is an isomorphism of graded algebras.}$

(OMG23)

Its source multiplication is the actual polynomial multiplication; its target multiplication has been constructed from the original sum coalgebra rather than assigning a multiplication between unrelated finite-dual vector spaces.

13.5 The full semisimple and nilpotent action in this geometry

On $\mathcal{P}_Z$ define the degree-preserving derivation

$$\partial_Z Z_{a,b,d} = \omega_{2a-1,2b-1} Z_{a,b,d} + \begin{cases} dZ_{a,b,d-1}, & d > 0, \\ 0, & d = 0. \end{cases} \quad (\text{OMG24})$$

On the parameter ring OMG7 it is induced by $\partial X_a = -i(2a-1)\delta X_a$, $\partial Y_b = (2b-1)\gamma Y_b$, $\partial T_0 = 0$, $\partial T_1 = T_0$. The product rule verifies OMG24 and commutation with Φ_m ; therefore the complete ideal $\mathfrak{t}_m$ is stable. The corresponding exact flow on the parameters is

$$(X_a, Y_b, T_0, T_1) \mapsto (e^{-i(2a-1)\delta w} X_a, e^{(2b-1)\gamma w} Y_b, T_0, T_1 + wT_0).$$

At $(X_0, X_1, Y_0, Y_1, T_0, T_1) = (1, 1, 1, 1, 0)$ it gives precisely the original functions OMG5. This is the entire stated analytic flow on the explicit algebraic variety, with all polynomial degrees.

If B_Z is the matrix in OMG24 on the column of variables, its rows have diagonal ω_α and entry d in column $(\alpha, d-1)$ of row (α, d) . In actual coordinates the derivation is

$$\partial_x x = \mathcal{A}x, \quad \mathcal{A} = H_m B_Z H_m^{-1}. \quad (\text{OMG25})$$

The inverse substitution proves $\partial_x \mathfrak{a} \subseteq \mathfrak{a}$, and its flow differentiates the original $x(w)$. On the exact degree- k basis of OMG9/OMG18 the induced action is

$$\partial_x \lambda_{k,a,b,D} = \omega_{k,a,b} \lambda_{k,a,b,D} + \begin{cases} D \lambda_{k,a,b,D-1}, & D > 0, \\ 0, & D = 0. \end{cases} \quad \Theta_k \partial_x = Y_k^\vee \Theta_k. \quad (\text{OMG26})$$

The displayed action on the basis uses its identification OMG23. For a direct verification, on the original primal primary basis $Y_k H_D = \omega H_D + i^{-1} H_{D+1}$ with the last term zero at $D = e_k - 1$. Applying $i^D D! H_D^\vee$ proves the exact lowering coefficient D in its dual action. Each frequency block has all e_k terms, with nonzero top iterated derivative $(e_k - 1)!$. Thus every original primary nilpotent has exactly the same index e_k in this model. The full annihilating polynomial is retained with its leading phase:

$$\chi_k(c_k + i\partial_x) = 0 \quad \text{on } R_k, \quad \chi_k(c_k + iT) = i^{q_k} \prod_{a,b=0}^k (T - \omega_{k,a,b})^{e_k}. \quad (\text{OMG27})$$

Distinct frequencies and the nonzero top chain prove that this degree q_k is minimal. The original S action corresponds to $c_k + i\partial_x$ on degree k , or to $\frac{1}{2}\mathbf{E} + i\partial_x$ on the graded algebra, where $\mathbf{E}$ is the degree derivation. These equalities specify every constant, sign and nilpotent action under OMG23.

13.6 The exact original cyclic observation as invariant restriction

Let $D = \text{diag}(\zeta, \dots, \zeta^n)$ and let $G = \mathbb{Z}/N$ act on $\mathcal{P}_x$ by $(gf)(x) = f(D^{-1}x)$. Thus a degree- k monomial x^ν has character $\zeta^{-\sum t\nu_t}$. Write

$$\mathcal{J} = \mathcal{P}_x^G, \quad \mathcal{J} = \text{im}(\mathcal{J} \longrightarrow R). \quad (\text{OMG28})$$

These are graded algebras. The coefficient image of the original observation is exactly their restriction at every degree, as follows. Let f_t be column t of F and let S_ν be the sum of all distinct ordered words in the f_t having multiplicity vector ν . Disjoint word supports make the S_ν a basis of the permutation-invariant tensor space. The expansion and projector are

$$(Fx)^{\otimes k} = \sum_{|\nu|=k} x^\nu S_\nu, \quad P_k(Fx)^{\otimes k} = \sum_{\substack{|\nu|=k \\ \sum t\nu_t \equiv 0 \pmod{N}}} x^\nu S_\nu, \quad P_k = \frac{1}{N} \sum_{a=0}^{N-1} (C^{-a})^{\otimes k}.$$

Every ordered word occurs once; the factor $k!/\prod \nu_t!$ is the number of terms in S_ν , not a missing coefficient. The original sum injection is the iterated map OMG20 into $E_h^{\otimes k}$. Hence the original observation $L_k = P_k \Pi^{\otimes k} U^{\otimes k} \iota_k = j\Lambda_k$ sends $e^{wY_k}[1]$ to the displayed projected curve with $x = x(w)$. The coefficient covectors of its surviving S_ν pull back to exactly $\Theta_k(x^\nu)$. They span the full dual of the image, since every covector on a subspace of a finite dimensional space extends to its ambient space. Consequently

$$\boxed{\begin{aligned} \Theta_k(\mathcal{J}_k) &= \text{im } \Lambda_k^\vee = (\ker \Lambda_k)^\perp, & \text{rank } \Lambda_k &= \dim \mathcal{J}_k, \\ R_k/\mathcal{J}_k &\xrightarrow{\Theta_k|_{K_k}} K_k^\vee \text{ is an isomorphism of vector spaces,} & K_k &= \ker \Lambda_k. \end{aligned}} \quad (\text{OMG29})$$

For the last statement, a covector on E_k vanishes on K_k exactly when it factors through $E_k/K_k \simeq \text{im } \Lambda_k$. Restriction to K_k is onto by extension of a basis. This proves the claimed quotient and its precise type: the degree- k quotient is a vector-space quotient by $\mathcal{J}_k$.

13.7 The actual orbit ideal, including every stabilizer

The whole ideal $\mathfrak{a}$ is generated by the explicit quadratic space in OMG15. Let B_2 be its coefficient matrix in the ordered degree-two monomial frame of $\mathcal{P}_x$. Repeated or dependent generator rows are harmless. For $a \in \mathbb{Z}/N$ let $D_{2,a}$ be diagonal with entry $\zeta^{-a \sum t\nu_t}$ at monomial x^ν . The actual stabilizer and its exact finite test are

$$H = \{a : g^a \mathfrak{a} = \mathfrak{a}\} = \left\{ a : \text{rank} \begin{bmatrix} B_2 \\ B_2 D_{2,a} \end{bmatrix} = \text{rank } B_2 \right\}. \quad (\text{OMG30})$$

Equality of these row spaces is equivalent to equality of their quadratically generated ideals; their dimensions agree under the invertible diagonal map. This proves the finite test using the actual entries OMG4. It prescribes no generic coefficient value. The set H is a subgroup by composition and inverse of ideal equalities. Put $s = |H|$ and $d = N/s$. Since G is cyclic, its cosets have representatives $0, \dots, d-1$ and $H = \{0, d, 2d, \dots, (s-1)d\}$. Define all distinct component ideals and the union algebra by

$$\mathfrak{a}_a = g^a \mathfrak{a} \quad (0 \leq a < d), \quad \mathfrak{u} = \bigcap_{a=0}^{d-1} \mathfrak{a}_a, \quad A = \mathcal{P}_x/\mathfrak{u}. \quad (\text{OMG31})$$

These are distinct homogeneous prime ideals with the same Hilbert function OMG13. Their intersection is a homogeneous radical ideal stable under the original cyclic action. It is the complete ideal of their scheme-theoretic union by the definition of the union's ideal. No product formula for this intersection is substituted.

For any invariant polynomial, membership in one component ideal implies membership in all its translates, by applying every g^a . Thus the exact invariant restriction and its inverse description are

$$\boxed{\begin{array}{c} \mathcal{J} \cap \mathfrak{a} = \mathcal{J} \cap \mathfrak{u}, \\ \mathcal{J}/(\mathcal{J} \cap \mathfrak{u}) \xrightarrow{[f] \mapsto [f]_{\mathfrak{u}}} A^G \xrightarrow{\text{res}_0} \mathcal{J} \subseteq R \end{array}} \text{ are isomorphisms of graded algebras.} \quad (\text{OMG32})$$

To prove surjectivity of the first map, average any representative of an invariant class by $N^{-1} \sum g^a$; its class is unchanged. Its kernel is exactly the displayed intersection. For the second, use an invariant representative: a zero restriction lies in $\mathcal{J} \cap \mathfrak{a} = \mathcal{J} \cap \mathfrak{u}$ and hence is already zero in A^G . Its image is $\mathcal{J}$ by definition. This gives the invariant image for every actual H , including nontrivial stabilizers of a composite group order $4m + 1$.

13.8 The complete component conductor and an explicit finite witness

Set $R_a = \mathcal{P}_x/\mathfrak{a}_a$. The simultaneous restriction

$$A \hookrightarrow \tilde{A} := \prod_{a=0}^{d-1} R_a, \quad [f]_{\mathfrak{u}} \mapsto ([f]_{\mathfrak{a}_a})_a \quad (\text{OMG33})$$

is injective precisely because its kernel before quotienting is the intersection OMG31. Its component conductor ideals are

$$\mathfrak{c}_a = \frac{(\bigcap_{b \neq a} \mathfrak{a}_b) + \mathfrak{a}_a}{\mathfrak{a}_a} \subseteq R_a, \quad \bigcap_{b \neq a} \mathfrak{a}_b = \mathcal{P}_x \quad (d=1). \quad (\text{OMG34})$$

The conductor of this specified inclusion is exactly

$$\boxed{\{t \in \tilde{A} : t\tilde{A} \subseteq A\} = \prod_{a=0}^{d-1} \mathfrak{c}_a \subseteq A.} \quad (\text{OMG35})$$

Indeed a tuple supported solely on component a lies in the image of A if and only if its value belongs to $\mathfrak{c}_a$, by lifting its other zero values to the intersection of the other ideals. If t is in the conductor, multiplication by each component idempotent of $\tilde{A}$ gives such a supported tuple; thus each component of t lies in $\mathfrak{c}_a$. Conversely these are ideals of R_a . Multiplying a tuple with all its components in the indicated ideals by any element of $\tilde{A}$ gives a sum of such supported tuples, each lying in A . This proves both inclusions and the full formula. In particular this conductor is also the annihilator of $\tilde{A}/A$ in A .

The action of G permutes these components with its original coefficient substitutions. Projection of an invariant tuple to component zero gives an isomorphism

$$\tilde{A}^G \xrightarrow{\sim} R^H, \quad r \in R^H \mapsto (g^a r)_{0 \leq a < d} \text{ for its inverse.} \quad (\text{OMG36})$$

The inverse is independent of coset representatives because r is fixed by H , and invariance follows by reindexing the components. The same argument applies to the conductor, whose ideals are permuted. Equations OMG32–36 consequently prove the exact inclusions of graded subalgebras and conductor ideals

$$\boxed{\mathfrak{c}_0^H \subseteq \mathcal{J} \subseteq R^H, \quad \mathfrak{c}_0^H R^H \subseteq \mathcal{J}.} \quad (\text{OMG37})$$

Here $\mathfrak{c}_0^H$ denotes invariant elements of the specified component ideal and is an ideal of R^H . An invariant tuple in the conductor already lies in A , so its projection belongs to $\mathcal{J}$ by OMG32. This proves the first inclusion; the ideal property gives the second.

There is a nonzero homogeneous element of this conductor with a completely bounded degree. For every $a \neq 0$ choose from the explicit transformed quadratic list of $\mathfrak{a}_a$ a quadratic $f_a \notin \mathfrak{a}_0$. Such a choice exists: containment of the entire quadratic space in that of $\mathfrak{a}_0$, which has the same dimension, would make the spaces equal and their generated ideals equal, contrary to distinctness. Membership is the finite row-space test in the same degree-two monomial frame. The product $f = \prod_{a=1}^{d-1} f_a$ lies in all other component ideals. It is nonzero modulo $\mathfrak{a}_0$ because that ideal is prime and none of its factors belongs to it. Thus $\bar{f} \in \mathfrak{c}_0$ is nonzero of degree $2(d-1)$. The group H preserves $\mathfrak{c}_0$ and acts on the domain R . Its exact norm

$$c_H = \prod_{h \in H} g^h \bar{f} \in \mathfrak{c}_0^H, \quad \deg c_H = D_H = 2s(d-1) = 2(N-s), \quad c_H \neq 0 \quad (\text{OMG38})$$

is invariant by permutation of its factors and nonzero in the domain. For $d=1$ take the empty product $c_H = 1$, $D_H = 0$; all statements remain exact. Multiplication by this actual conductor element is injective and gives the finite dimension bounds, with negative graded pieces set to zero,

$$\begin{aligned} (R^H)_{k-D_H} &\xrightarrow{\times c_H} \mathcal{J}_k \hookrightarrow (R^H)_k, \\ \dim(R^H)_{k-D_H} &\leq \text{rank } \Lambda_k \leq \dim(R^H)_k, \\ q_k - \dim(R^H)_k &\leq \dim K_k \leq q_k - \dim(R^H)_{k-D_H}. \end{aligned} \quad (\text{OMG39})$$

Every ideal, group and matrix in these expressions is defined by the actual original coefficient map. In particular the stabilizer has its finite equality test OMG30 throughout; no particular value of it has been inserted into the dimensions. To compute the displayed invariant dimensions directly, let the actual H substitutions act on the basis OMG9 through $Z \mapsto H_m^{-1} D^{-h} H_m Z$ and reduce with OMG10. This is an explicit finite matrix on q_k coordinates. The idempotent average $s^{-1} \sum_{h \in H} g^h$ has trace equal to its rank, proving $\dim(R^H)_k = s^{-1} \sum_{h \in H} \text{Tr}(g^h | R_k)$. Thus the dimensions in OMG39 also have complete original finite coefficient formulas for every actual stabilizer.

13.9 The exact action defect and all subsequently observed directions

The derivation ∂_x of OMG25 acts on R with all primary directions OMG26. The possible failure of its restriction to the observed image is the specified finite linear map

$$\mathcal{J}_k \longrightarrow R_k / \mathcal{J}_k, \quad j \longmapsto [\partial_x j]. \quad (\text{OMG40})$$

Via OMG29 it is exactly the complex linear dual of the original map $\Lambda_k Y_k I_K : K_k \rightarrow B_k$. Indeed a j corresponds to the covector $\lambda \Lambda_k$ on E_k for a unique boundary covector λ ; evaluating OMG26 on $I_K v$ gives $\lambda \Lambda_k Y_k I_K v$, proving the identification and all domains and codomains. The action on ambient polynomials also has the exact conjugation formula

$$g^a \partial_x g^{-a} = \partial_{D^a A D^{-a}}, \quad g^a \partial_x g^{-a} - \partial_x = \partial_{D^a A D^{-a} - A}. \quad (\text{OMG41})$$

For a linear vector field $\partial_M f = (Mx) \cdot \nabla f$, the chain rule proves this by substituting $D^{-a}x$ after differentiating $f(D^a x)$. This formula retains every original period/unit coefficient in $\mathcal{A}$ and does not require the original ideal or the observed image to be preserved by extra ambient actions.

Finally the subspace invisible to all original iterates has the complete model

$$\begin{aligned} \mathcal{K}_{\infty, k} &= \bigcap_{r=0}^{q_k-1} \ker(\Lambda_k Y_k^r), \\ R_k / \sum_{r=0}^{q_k-1} \partial_x^r \mathcal{J}_k &\xrightarrow{\Theta_k | \mathcal{K}_{\infty, k}} \mathcal{K}_{\infty, k}^\vee \quad \text{is a vector-space isomorphism.} \end{aligned} \quad (\text{OMG42})$$

In finite dimensional duality the annihilator of an intersection of kernels is the sum of the images of their dual maps: this follows by taking the dual of the single stacked map with all those components, whose kernel is precisely the intersection. Equations OMG26 and OMG29 identify these images with $\partial_x^r \mathcal{J}_k$, proving OMG42. The exact original polynomial OMG27 expresses every higher iterate as a linear combination of the first q_k , so the same intersection is taken over all nonnegative iterates and is Y_k -invariant. This proves the full nilpotent and cyclic-action receiver in the new homogeneous model for every original multiplicity, with its actual observation, periods and amplitude jets throughout.

14 The absolute common kernel: the new finite result

This complete incoming calculation sharpens the current absolute common kernel estimate. Its original comparison was against the preceding logarithmic bound; the intervening KAF chapter already established the $O(kq)$ scale. The new finite envelope reduces its uniform limsup constant from $16(4 + 10 \log 2)$ to $32 \log(25 \log 3 / (4\pi))$. AKJ below takes the pointwise minimum of both finite bounds. No limiting kernel coefficient is inferred from these upper bounds.

Retain the actual hypothetical complete quartet and its entire quotient

$$\rho = \frac{1}{2} + \delta + i\gamma, \quad 0 < \delta < \frac{1}{2}, \quad \gamma > 2, \quad h(z) = \prod_{\alpha \in \{\rho, \bar{\rho}, 1-\rho, 1-\bar{\rho}\}} (z - \alpha)^m, \quad v_h = 2\xi/h.$$

The full primary unit is $U = M_{j_h v_h}$. The original tensor order, centre, polynomial, and coefficient quotient are

$$\begin{aligned} k &= 4l + 1, \quad l \geq 1, \quad e = 1 + k(m - 1), \quad q = e(k + 1)^2, \quad c = k/2, \\ \lambda_{a,b} &= c + (2a - k)\delta + i(2b - k)\gamma, \quad \chi(S) = \prod_{a,b=0}^k (S - \lambda_{a,b})^e, \quad E = \mathbb{C}[S]/(\chi). \end{aligned} \quad (\text{AKS1})$$

For the principal conclusion $m = 1$. The source-operator estimates below also hold with the stated repeated-root list for $m > 1$, but no simple-quartet rank is assigned to those multiplicities.

Let $\Lambda_k(u) : E \rightarrow B_k(u)$ be the original observation, $K(u) = \ker \Lambda_k(u)$, and $I : K(u) \hookrightarrow E$ its original coefficient inclusion. For a source $s \in \{1, k\}$, let $G_N^{(s)}$ be the original minimum-remainder Gram, in the frame $(1, S, \dots, S^{q-1})$. Define $H_{K,N}^{(s)} = I^* G_N^{(s)} I$ and retain exactly

$$F_K^{(s)} = \log \det H_{K,q-1}^{(s)} + \log \det H_{K,q}^{(s)} - \log \det H_{K,2q-1}^{(s)} - \log \det H_{K,2q}^{(s)}. \quad (\text{AKS2})$$

Every empty determinant is one. All conjugate transposes refer to the specified coordinate matrices, not to a bilinear residue pairing.

Here are entirely explicit positive numbers used in the bound:

$$\begin{aligned} R &= k\sqrt{\delta^2 + \gamma^2}, \quad b_s = s/2, \quad t_q = \frac{\pi}{2} - \frac{1}{q}, \quad \mathbf{a}_q = \frac{2 \log 3}{t_q}, \quad \mathbf{s}_q = \frac{\mathbf{a}_q^q - 1}{\mathbf{a}_q - 1}, \\ \mathbf{b}_{s,q} &= \left[\frac{9}{25} \sin \frac{1}{q} \right]^{-b_s} \exp \left(2R \left(\log 3 + \frac{t_q}{2} \right) \right), \\ \mathcal{Z}_{s,j} &= 1 + \mathbf{b}_{s,q} \mathbf{s}_q^2 \sum_{d=q}^{q+j-1} \frac{d!}{(b_s)_d} \left(\frac{25}{16} \right)^d, \quad \mathcal{L}_{s,j} = \log \mathcal{Z}_{s,j} \quad (0 \leq j \leq q + 1). \end{aligned} \quad (\text{AKS3})$$

The empty sum gives $\mathcal{Z}_{s,0} = 1$. In the admitted family $q \geq 36$, so $t_q \in (0, \pi/2)$ and $\mathbf{a}_q > 1$. These parameters are radii in the existing Laplace and generating functions; neither is a new source order.

Main theorem (absolute finite bound). For the original simple quartet, every original logarithm branch, and every $|u| \geq R_*$, where R_* is precisely RPC21 or RPC23 (recorded in Section 20), put $r = 8k - 16$. Then

$$\boxed{0 \leq F_K^{(s)}(k, u) \leq r(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}), \quad s \in \{1, k\}.} \quad (\text{AKS4})$$

There is a stronger, fully period-dependent interval in AKS28 below. In particular, with

$$\kappa_* = 32 \log \left(\frac{25 \log 3}{4\pi} \right), \quad 25.02078097649 < \kappa_* < 25.02078097650, \quad (\text{AKS5})$$

one has the uniform statement

$$\boxed{0 \leq \limsup_{\substack{k \rightarrow \infty \\ k \equiv 1 \pmod{4}}} \sup_{\substack{|u| \geq R_* \\ \text{original branches}}} \frac{F_K^{(s)}(k, u)}{kq} \leq \kappa_*}. \quad (\text{AKS6})$$

Here the source may be $s = 1$ for every k , or $s = k$. The same bound holds for the actual arithmetic kernel. Its exact finite allowances are in AKS34–36. Thus the absolute kernel is $O_h(kq)$ on this domain, not only $O_h(kq \log q)$. Neither AKS5 nor AKS6 calls κ_* the actual leading coefficient.

15 An exact coefficient realization of the actual lost directions

15.1 The period matrix and invariant restriction, with their full coefficients

For $m = 1$, order the one-factor roots as $(++)$, $(+-)$, $(-+)$, $(--)$, and let $\mathcal{E}_{\text{prim}}$ have their original Lagrange idempotents as columns. Set $\zeta = \exp(2\pi i/5)$ and $F_{jt} = \zeta^{jt} - 1$, $1 \leq j, t \leq 4$. The exact inverse is $(F^{-1})_{tj} = \zeta^{-tj}/5$. Retain

$$\begin{aligned} \Pi[P]_j &= \int_{\Gamma_j} e^{\Phi_h(z)/u} \text{rem}_h P(z) dz, & \Phi'_h &= h, & \Phi_h(0) &= 0, \\ \Gamma_j &= -\ell_0 + \ell_j, & \arg \ell_j &= (\arg u + \pi + 2\pi j)/5, \\ \mathbf{H} &= F^{-1} \Pi U \mathcal{E}_{\text{prim}}, & D &= \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4), \\ Q(x) &= Q_0(\mathbf{H}^{-1}x), & Q_0(t) &= t_1 t_4 - t_2 t_3, & Q_a(x) &= Q(D^{-a}x). \end{aligned} \quad (\text{AKS7})$$

The integral always uses the degree-less-than-four representative. Invertibility of Π , all contour phases, and its full determinant are the PDE/PQO results in the supplied proofs. The four unit values are $(a, \bar{a}, \bar{a}, a)$, with the actual $a = v_h(\rho) \neq 0$; no value of a is selected. Hence $\mathbf{H}$ is the original invertible matrix.

The following formulas make the coefficient realization directly computable. Write

$$\mathcal{I}_{k,0} = \{\nu \in \mathbb{Z}_{\geq 0}^4 : |\nu| = k, \nu_1 + 2\nu_2 + 3\nu_3 + 4\nu_4 \equiv 0 \pmod{5}\}.$$

For $\nu \in \mathcal{I}_{k,0}$ define $B_{\nu;(a,b)}$ by summing over 4×4 arrays $n_{t,\alpha} \geq 0$ with row sums ν_t , total plus-real count a , and total plus-imaginary count b :

$$B_{\nu;(a,b)} = \sum_{\substack{\sum_{\alpha} n_{t,\alpha} = \nu_t \\ \sum_{t,\alpha: \epsilon_{\alpha}=+} n_{t,\alpha} = a \\ \sum_{t,\alpha: \eta_{\alpha}=+} n_{t,\alpha} = b}} \left(\prod_{t=1}^4 \frac{\nu_t!}{\prod_{\alpha} n_{t,\alpha}!} \right) \prod_{t,\alpha} \mathbf{H}_{t,\alpha}^{n_{t,\alpha}}. \quad (\text{AKS8})$$

Let $\mathbf{V}_{(a,b),d} = \lambda_{a,b}^d$ for $0 \leq d < q$ and set $\mathbf{A} = \mathbf{B}\mathbf{V}$. All entries of $\mathbf{A}$ are now finite expressions in the actual periods and unit. There is no generic period matrix in this definition.

For completeness, put $t_{\alpha}(w) = \exp((\eta_{\alpha}\gamma - i\epsilon_{\alpha}\delta)w)$ and $x(w) = \mathbf{H}t(w)$. Expansion gives

$$x(w)^{\nu} = \sum_{a,b} B_{\nu;(a,b)} e^{\omega_{a,b} w}, \quad \omega_{a,b} = (2b - k)\gamma - i(2a - k)\delta.$$

In the original primary algebra, $e^{wY}[1] = \sum_{a,b} e^{\omega_{a,b} w} \varepsilon_{a,b}$, where $Y = (M_S - cI)/i$. The exponentials are independent by the Vandermonde of the distinct $\omega_{a,b}$. OCS9's unscaled symmetric word basis therefore identifies the original observation with

$$j\Lambda_k = \mathcal{F}_0 \mathbf{A}, \quad \ker \mathbf{A} = K(u) \quad \text{in the original coefficient frame.} \quad (\text{AKS9})$$

Every multinomial in AKS8 belongs to its actual ordered words; the word multiplicity is not inserted a second time. The exact inherited row Gram of $\mathcal{F}_0$ is the restriction to $\mathcal{I}_{k,0}$ of

$$\mathcal{G}_{\nu,\mu} = \sum_{\substack{n_{tu} \geq 0 \\ \sum_u n_{tu} = \nu_t \\ \sum_t n_{tu} = \mu_u}} \frac{k!}{\prod_{t,u} n_{tu}!} \prod_{t,u} [5(\delta_{tu} + 1)]^{n_{tu}}. \quad (\text{AKS10})$$

This follows by taking the coefficient of $\bar{x}^\nu y^\mu$ in $(\bar{x}^T F^* F y)^k$, since $F^* F = 5(I + \mathbf{1}\mathbf{1}^*)$. It is positive definite and includes every row cross term.

Here is the precise role of the orbit ideal. The map sending a homogeneous polynomial f to the covector specified by $\Theta_k(f)(e^{wY}[1]) = f(x(w))$ has kernel $Q\mathbb{C}[x]_{k-2}$ and is onto $E^\vee$. Indeed equal exponential indices correspond exactly to monomial differences divisible by $t_1 t_4 - t_2 t_3$; the displayed Vandermonde proves that there are no other relations. Its invariant image is $\text{im } \Lambda_k^\vee$. On the five-member domain, the five Q_a are distinct prime quadrics. An invariant polynomial divisible by Q is divisible by each Q_a , and therefore by their full product; its quotient is invariant by cancellation in the polynomial ring. Consequently

$$\mathbb{C}[x]^{\langle D \rangle} \cap (Q) = \left(\prod_{a=0}^4 Q_a \right) \mathbb{C}[x]^{\langle D \rangle}, \quad (\text{AKS11})$$

$$K(u)^\vee \simeq \mathbb{C}[x]_k / (\mathbb{C}[x]_k^{\langle D \rangle} + Q\mathbb{C}[x]_{k-2}),$$

$$b := \text{rank } \mathbf{A} = d_{5,k,0} - d_{5,k-10,0} = k^2 - 6k + 17, \quad r = q - b = 8k - 16.$$

Negative degrees are zero. The count is $d_{5,d,0} = ((\binom{d+3}{3} + 4\epsilon_d)/5$, $\epsilon_d = \mathbf{1}_{d \equiv 0(5)} - \mathbf{1}_{d \equiv 1(5)}$; subtracting the cubics gives the stated expression for $k \geq 10$, and $k = 5, 9$ give 12, 44 directly. This is the inherited OQX calculation, repeated to specify which kernel AKS9 realizes. The source metric is still to be applied to this kernel, not to the invariant ring itself.

15.2 A determinant formula requiring no assumed nonzero pivot

Let $\mathcal{G}_0$ be the full positive row Gram in AKS10 and set

$$W = \mathbf{A}^* \mathcal{G}_0 \mathbf{A}, \quad \sigma_j = [z^j] \det(I_q + zW), \quad P_K = \frac{1}{\sigma_b} \sum_{j=0}^b (-1)^j \sigma_{b-j} W^j. \quad (\text{AKS12})$$

Since W has exactly b positive eigenvalues, $\sigma_b > 0$. The scalar polynomial in AKS12 is one at zero and zero at every nonzero eigenvalue: it is the normalized product of the factors $\lambda_i - z$ for the b positive eigenvalues. Hence $P_K = P_K^* = P_K^2$ and $\text{im } P_K = K(u)$. This is the projector derived from the literal observation, not a freely chosen projection. Replacing $\mathcal{G}_0$ by any other positive coordinate Gram leaves this same coefficient-orthogonal kernel projector, since its image and orthogonal complement are unchanged. No such replacement is needed here.

At every original endpoint define

$$D_{K,N}^{(s)} = [z^r] \det(I_q + z P_K G_N^{(s)} P_K). \quad (\text{AKS13})$$

These numbers are strictly positive for $r > 0$ and are one for $r = 0$. To prove their exact metric meaning, choose an orthonormal coefficient basis U_K of $\text{im } P_K$ just for the proof. In a unitary extension, $P_K G_N P_K$ has its positive block $U_K^* G_N U_K$ and a zero block. Thus $D_{K,N} = \det(U_K^* G_N U_K)$. Write the original inclusion as $I = U_K T$; then $|\det T|^2 = \det(I^* I)$ and

$$\det H_{K,N}^{(s)} = \det(I^* I) D_{K,N}^{(s)}, \quad \boxed{F_K^{(s)} = \log \frac{D_{K,q-1}^{(s)} D_{K,q}^{(s)}}{D_{K,2q-1}^{(s)} D_{K,2q}^{(s)}}}. \quad (\text{AKS14})$$

This supplies an equivalent exact coefficient determinant formula for all the $8k - 16$ lost directions, including the entire period dependence. The original frame factor is present at each endpoint and cancels only with its four actual signs. It is unnecessary to suppose any particular minor nonzero. Alternatively, selecting the first independent r columns of P_K gives an explicit coefficient frame; the determinant formula avoids even that chart choice.

For $m > 1$ replace AKS8–9 by OCS12–18’s full confluent matrix $\mathfrak{A}_{\nu;(a,b,D)} = i^D D! c_{\nu;a,b,D}$ and its primary isomorphism. AKS12–14 are then identical with the actual rank, not with the rank in AKS11. All primary nilpotents and all unit derivatives remain in that matrix.

16 A bounded exact remainder operator on the original source

16.1 The complete source norms and coefficient map

For either existing source set $M_s = (2\pi)^{s/2}$ and retain

$$dm_{0,s}(y) = \frac{M_s 2^{b_s}}{4\pi\Gamma(b_s)} |\Gamma(b_s/2 + iy/2)|^2 dy, \quad \int e^{ty} dm_{0,s} = M_s (\cos t)^{-b_s}, \quad |t| < \pi/2. \quad (\text{AKS15})$$

The supplied IGO1–3 proof derives this transform by the beta integral, Fourier inversion and a contour shift within $|\operatorname{Im} x| < \pi/2$. In particular it includes the exact masses $M_1 = \sqrt{2\pi}$ and $M_k = (2\pi)^{k/2}$ and exponential domination on every smaller strip. The original real monic polynomials satisfy

$$\begin{aligned} p_0 &= 1, \quad p_1 = y, \quad p_{d+1} = yp_d - d(b_s + d - 1)p_{d-1}, \\ \sum_{d \geq 0} \frac{p_d(y)}{d!} z^d &= (1 + z^2)^{-b_s/2} e^{y \arctan z}, \quad h_d^{(s)} = \|p_d\|^2 = M_s d! (b_s)_d. \end{aligned} \quad (\text{AKS16})$$

The generating formula follows by differentiating in z . To check the normalization directly, integrate the product of two generating functions at small real z, w . AKS15 and the cosine addition identity give $M_s(1 - zw)^{-b_s}$. Taking any finite pair of derivatives at zero, justified by the stated exponential domination, gives orthogonality and the norms in AKS16. Thus the leading phase in $u_d(S) = p_d((S - c)/i) / \sqrt{h_d^{(s)}}$ is exactly i^{-d} . The same generating function agrees with NIST DLMF 18.23.7 after the original parameter rescaling; it is derived here from the supplied recurrence, not substituted for it.

List the original roots $\omega_\nu = (\lambda_{a,b} - c)/i$ in lexicographic order, repeating each exactly e times, and put

$$P(y) = i^{-q} \chi(c + iy) = \prod_{\nu=1}^q (y - \omega_\nu), \quad \Phi : [f(y)] \mapsto [f((S - c)/i)]. \quad (\text{AKS17})$$

The inverse is substitution $S = c + iy$, and $P((S - c)/i) = i^{-q} \chi(S)$. This is an exact algebra isomorphism, including the complete confluent orders; there is no monomial comparison divisor. Let C_s have columns $[u_0], \dots, [u_{q-1}]$. Its nonzero determinant is $i^{-q(q-1)/2} \prod_{d < q} (h_d^{(s)})^{-1/2}$. Define

$$v_j = C_s^{-1} [u_{q+j}], \quad \mathbf{K}_j = I_q + \sum_{a=0}^{j-1} v_a v_a^*, \quad 0 \leq j \leq q + 1. \quad (\text{AKS18})$$

Parseval in the low-degree original orthonormal polynomial basis proves

$$\|v_j\|_2^2 = \frac{\|\operatorname{rem}_P p_{q+j}\|_{m_{0,s}}^2}{M_s (q+j)! (b_s)_{q+j}}. \quad (\text{AKS19})$$

This is the full norm of the actual remainder, not a diagonal entry of an approximated quotient Gram.

16.2 Newton interpolation with all roots and their orders

For $0 \leq a < q$ define $N_a(y) = \prod_{\nu=1}^a (y - \omega_\nu)$, with $N_0 = 1$. For every complex v , the unique entire-function remainder is

$$\text{rem}_P(e^{vy}) = \sum_{a=0}^{q-1} [\omega_1, \dots, \omega_{a+1}] e^{v \cdot} N_a(y), \quad |[\omega_1, \dots, \omega_{a+1}] e^{v \cdot}| \leq \frac{|v|^a}{a!} e^{R|v|}. \quad (\text{AKS20})$$

Here divided differences at repeated nodes mean Hermite divided differences, not evaluation with a zero denominator.

A proof covering the confluent case is useful. On the standard a -simplex with barycentric coordinates $\theta_1, \dots, \theta_{a+1}$,

$$[\omega_1, \dots, \omega_{a+1}] e^{v \cdot} = v^a \int_{\Delta_a} e^{v \sum \theta_\nu \omega_\nu} d\theta.$$

The integral of $\prod \theta_\nu^{n_\nu}$ is $\prod n_\nu! / (a + \sum n_\nu)!$, as follows by iterated integration of integer beta integrals. Expanding the exponential therefore gives $\sum_{n \geq 0} v^{a+n} h_n(\omega_1, \dots, \omega_{a+1}) / (a+n)!$. For monomials, the divided difference of y^{a+n} is exactly h_n : this follows by multiplying the geometric series $\prod (1 - \omega_\nu z)^{-1}$ and the Newton division recurrence. It is a polynomial identity in the nodes and so also holds at repeated nodes. This proves the integral formula. Iterated Newton division for polynomials, followed by the convergent exponential series, proves the first equality in AKS20, including every required derivative. Finally the simplex has volume $1/a!$ and $|\sum \theta_\nu \omega_\nu| \leq R$, proving its stated bound.

For any $0 < t < \pi/2$, put $\tau = 2/t$. Positivity of every term of the cosh series in AKS15 gives

$$\|y^a\|_{m_{0,s}} \leq \sqrt{M_s} (\cos t)^{-b_s/2} \tau^a a!.$$

Indeed its square is at most $M_s (\cos t)^{-b_s} (2a)! t^{-2a}$ and $(2a)! \leq 4^a (a!)^2$. Expanding the *whole* Newton polynomial, then using $|e_j(\omega_1, \dots, \omega_a)| \leq \binom{a}{j} R^j$, proves

$$\begin{aligned} \|N_a\|_{m_{0,s}} &\leq \sqrt{M_s} (\cos t)^{-b_s/2} \sum_{j=0}^a \binom{a}{j} R^{a-j} \tau^j j! \\ &= \sqrt{M_s} (\cos t)^{-b_s/2} \tau^a a! \sum_{n=0}^a \frac{(R/\tau)^n}{n!} \\ &\leq \sqrt{M_s} (\cos t)^{-b_s/2} \tau^a a! e^{R/\tau}. \end{aligned} \quad (\text{AKS21})$$

There are no omitted polynomial coefficients. Combining AKS20–21 gives for every complex v

$$\|\text{rem}_P(e^{vy})\|_{m_{0,s}} \leq \sqrt{M_s} (\cos t)^{-b_s/2} e^{R(|v|+t/2)} \sum_{a=0}^{q-1} (2|v|/t)^a. \quad (\text{AKS22})$$

16.3 Cauchy extraction in the actual Gamma generating function

Fix $0 < z < 1$ and write $a_z = \text{arctanh } z$. For $|w| = z$ the defining power series gives $|\arctan w| \leq a_z$ and $|1 + w^2| \geq 1 - z^2$. Apply the literal finite-dimensional remainder map to AKS16 and use Cauchy's formula for its d th coefficient. The norm triangle inequality on that circle and AKS22 give

$$\begin{aligned} \frac{\|\text{rem}_P p_d\|_{m_{0,s}}}{d!} &\leq z^{-d} (1 - z^2)^{-b_s/2} \sqrt{M_s} (\cos t)^{-b_s/2} \\ &\quad \times e^{R(a_z+t/2)} \sum_{a=0}^{q-1} (2a_z/t)^a. \end{aligned} \quad (\text{AKS23})$$

Cauchy's formula is legitimate for the finite remainder vector: its entries are holomorphic for $|w| < 1$ by the actual matrix generating identity OPG7. The source norm is taken only after this finite-vector identity; there is no unjustified integral of a nonintegrable generating function on the large circle.

Substitution of AKS23 into AKS19 retains the factorial denominator and cancels the two occurrences of the original mass, giving

$$\|v_j\|_2^2 \leq \frac{d!}{(b_s)_d} z^{-2d} [\cos t(1 - z^2)]^{-b_s} e^{2R(az+t/2)} \left[\sum_{a=0}^{q-1} (2a_z/t)^a \right]^2, \quad d = q + j. \quad (\text{AKS24})$$

Adding the *complete* rank-one positive matrices in AKS18 now proves $K_j \preceq \mathcal{Z}_{s,j} I_q$ after setting $z = 4/5$ and $t = t_q$. Indeed $a_{4/5} = \log 3$, $1 - z^2 = 9/25$, and $\cos t_q = \sin(1/q)$, so the exact scalar obtained is AKS3. The high index $j = q + 1$ includes the original polynomial of degree $2q$.

There is also a root-by-root finite refinement. Put $\mu_{s,d} = \int y^{2d} dm_{0,s}$, $R_a = \max_{\nu \leq a+1} |\omega_\nu|$, and

$$B_{s,a}^{\text{root}} = \sum_{d=0}^a |e_{a-d}(\omega_1, \dots, \omega_a)| \sqrt{\mu_{s,d}}, \quad (\text{AKS25})$$

$$J_s^{\text{root}}(z) = \frac{(1 - z^2)^{-b_s}}{M_s} \left[\sum_{a=0}^{q-1} \frac{a_z^a}{a!} e^{a_z R_a} B_{s,a}^{\text{root}} \right]^2.$$

Replace the scalar multiplying $d!z^{-2d}/(b_s)_d$ in AKS24 by $J_s^{\text{root}}(z)$ to get a valid sharper bound, by the same proof before AKS21 is relaxed. Every moment is exactly $M_s(2d)! [w^{2d}] (\cos w)^{-b_s}$. Thus this refinement keeps every actual root coefficient and a finite original moment formula. AKS21 bounds it by the explicit scalar in AKS24. The proof never replaces the source determinant by a product of diagonal moments: AKS24 is a full-vector operator bound, and the next step restricts the inverse of the full covariance.

17 The four original kernel Grams and the evaluated scale

The original orthonormal source map at degree $q + j - 1$ is $C_s[I_q, v_0, \dots, v_{j-1}]$. Its minimum lift of $C_s x$ is $[I_q, v_0, \dots, v_{j-1}]^* K_j^{-1} x$. It has that image and is orthogonal to the full relation kernel, so Pythagoras proves

$$G_{q+j-1}^{(s)} = C_s^{-*} K_j^{-1} C_s^{-1}, \quad H_j := H_{K,q+j-1}^{(s)} = T_s^* K_j^{-1} T_s, \quad T_s = C_s^{-1} I. \quad (\text{AKS26})$$

This verifies the source-minimum step in the coefficient determinant AKS13–14. Neither a cross block nor any original relation is removed.

Since $I_q \preceq K_j \preceq \mathcal{Z}_{s,j} I_q$, inversion and compression give $\mathcal{Z}_{s,j}^{-1} H_0 \preceq H_j \preceq H_0$, where $H_0 = T_s^* T_s$. Define $d_j = -\log \det(H_0^{-1/2} H_j H_0^{-1/2})$. Congruence and all r eigenvalues give

$$0 = d_0 \leq d_1 \leq \dots \leq d_{q+1}, \quad 0 \leq d_j \leq r \mathcal{L}_{s,j}, \quad F_K^{(s)} = d_q + d_{q+1} - d_1. \quad (\text{AKS27})$$

The identity uses precisely the four endpoints in AKS2. In particular it proves AKS4, as well as the sharper interval

$$d_1 \leq F_K^{(s)} \leq r(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - d_1. \quad (\text{AKS28})$$

For a directly calculable expression for its lower endpoint set $V = \|v_0\|^2$ and $\theta = v_0^* T_s (T_s^* T_s)^{-1} T_s^* v_0$. Then

$$0 \leq \theta \leq V, \quad d_1 = \log \frac{1 + V}{1 + V - \theta}. \quad (\text{AKS29})$$

Indeed the inverse of $I + v_0 v_0^*$ is $I - v_0 v_0^*/(1 + V)$; take its full compressed determinant. For $r = 0$ set $\theta = d_1 = 0$. In GMB/OPR notation $V = t_0$ and $V - \theta = \xi_0$, so this is exactly the first original section-corrected kernel increment, not an independently chosen kernel observation.

Here is an evaluated finite remainder for AKS6. Put

$$\begin{aligned} a_\infty &= \frac{4 \log 3}{\pi}, & c_0 &= \log \frac{25 \log 3}{4\pi} > 0, \\ E_{s,q}(R) &= \log 2 + \frac{4}{\pi - 2/q} + \frac{s}{2} \log \frac{25\pi q}{18} + 2R \left(\log 3 + \frac{\pi}{4} \right) \\ &\quad + \log((q+1)(4q+1)) - 2 \log(a_\infty - 1). \end{aligned} \tag{AKS30}$$

Then for both source orders, every $q \geq 36$, and the same actual kernel,

$$\boxed{\mathcal{L}_{s,q}, \mathcal{L}_{s,q+1} \leq 2c_0 q + E_{s,q}(R), \quad 0 \leq F_K^{(s)} \leq 4rc_0 q + 2rE_{s,q}(R).} \tag{AKS31}$$

To verify every factor, $b_s \geq 1/2$ gives

$$\frac{d!}{(b_s)_d} \leq \frac{4^d}{\binom{2d}{d}} \leq 2d + 1.$$

The last inequality uses that the central binomial is the largest of the $2d + 1$ positive coefficients whose sum is 4^d . Thus the sum in AKS3 is at most $(q+1)(4q+1)(5/4)^{4q}$. Also $\mathfrak{s}_q \leq \mathfrak{a}_q^q/(a_\infty - 1)$ and

$$2q \log \mathfrak{a}_q \leq 2q \log a_\infty + \frac{4}{\pi - 2/q}, \quad \sin(1/q) \geq \frac{2}{\pi q}.$$

The first follows from $-\log(1-x) \leq x/(1-x)$ at $x = 2/(\pi q)$; the second is the chord bound for sine on $[0, \pi/2]$. Use $t_q \leq \pi/2$ in the remaining exponential. The order- q terms combine to $4q \log(5/4) + 2q \log a_\infty = 2c_0 q$. The upper bound for the logarithm of the added positive scalar is positive, so $\log(1 + e^x) \leq \log 2 + \max(0, x)$ proves the first bound in AKS31. AKS28 proves the second.

On the original simple family $q = (k+1)^2$, $R = O_h(k)$ and $r = 8k - 16$. Consequently $E_{1,q}(R) = O_h(k + \log q)$ and $E_{k,q}(R) = O_h(k \log q)$. Dividing AKS31 by kq proves AKS6, with $32c_0 = \kappa_*$. In fact the fixed-source upper bound is

$$F_K^{(1)} \leq (32k - 64)c_0 q + (16k - 32)E_{1,q}(R), \tag{AKS32}$$

whose explicit second term is $O_h(k^2)$. The tensor-source version has an $O_h(k^2 \log q)$ upper remainder. These are remainders of upper bounds, not assertions about the actual signed error or leading value. The proof is uniform in all the period parameters in the stated domain: no norm or condition number of a period matrix occurs in AKS30.

The precise earlier loss. OKV15 had already retained $\|v_j\|^2 \leq q^2 2^{6q} (2D_* q)^{2d} / (d!(b_s)_d)$, $d = q + j$. Replacing its denominator by $1/2$ caused the old $q \log q$ exponent. Even without AKS20–24 one obtains the valid repair

$$\|K_{q+1}\| \leq 1 + (q+1)q^2 2^{6q} \frac{(4D_* q)^{4q}}{(4q)!}, \quad D_* \geq 3. \tag{AKS33}$$

Here $d!(b_s)_d \geq (2d)!/4^d$ and the remaining factorial ratio increases up to $d = 2q$ for $D_* \geq 3$. The elementary inequality $\log n! \geq n \log n - n + 1$ already makes the logarithm of AKS33 $O_h(q)$. The Newton calculation strengthens this repair to the packet-independent limsup constant AKS5, while retaining R in its finite remainder.

Exact enclosure of the displayed numerical ceiling. For $x > 1$ and $z = (x-1)/(x+1)$ the integrated geometric series gives

$$0 \leq \log x - 2 \sum_{j=0}^{39} \frac{z^{2j+1}}{2j+1} \leq \frac{2z^{81}}{81(1-z^2)}.$$

For $0 < x < 1$, the alternating sum through $j = 39$ for $\arctan x$ has error between zero and $x^{81}/81$. The tangent addition formula, with the angle in $(0, \pi/2)$, proves $\pi = 16 \arctan(1/5) - 4 \arctan(1/239)$: the tangent of $4 \arctan(1/5) - \arctan(1/239)$ is exactly one. Use the resulting rational lower and upper bounds for π and for $\log 3$, then apply the first displayed bound to the two endpoints of $25 \log 3/(4\pi)$. Multiplication by 32 gives, with outward decimal rounding, the interval

$$\kappa_* \in [25.0207809764994871, 25.0207809764994872].$$

The accompanying rational-arithmetic implementation checks these inequalities by integer cross multiplication, rather than a floating-point logarithm. This proves the less finely rounded strict interval AKS5.

18 The complete arithmetic and mixed receivers

18.1 The unchanged finite arithmetic allowances

Use exactly AMT4's constants. To prevent a loss hidden in order notation, write them here:

$$\begin{aligned} \alpha &= \pi/2, \quad p = 8m - 9/2, \quad B_h = 42 + 8m, \quad M = 21 + 4m, \\ \vartheta_h &= \int_{-1}^1 w_h(y) dy, \quad M_\alpha = \int e^{\alpha y} w_h(y) dy, \quad w_h(y) = |v_h(1/2 + iy)|^2/(2\pi), \\ c_\Gamma &= \sqrt{2}e^{-7/6}, \quad C_\Gamma = \sqrt{2\sqrt{5}}e^{2/3}, \\ a_k &= \frac{c_h \vartheta_h^{k-3} e^{-\alpha(k-3)} (k-2)^{-B_h}}{C_\Gamma 2^{B_h/2}}, \quad A_k = \frac{C_h^{\text{tilt}}}{c_\Gamma} k^{p+1} M_\alpha^{k-1}, \\ D_N &= [1 + 4(N+M)^2]^M, \quad L_N^{\text{ar}} = \log(A_k D_N / a_k), \\ E_k^+ &= L_{q-1}^{\text{ar}} + L_q^{\text{ar}}, \quad E_k^- = L_{2q-1}^{\text{ar}} + L_{2q}^{\text{ar}}. \end{aligned} \tag{AKS34}$$

The constants c_h, C_h^{tilt} are the original TW/BT envelope constants, not new choices. Their actual inequalities and complete source construction are preserved in AMT4–6 and the current source closure. Minimize AMT6 on the same remainder fibre and then restrict to $I(K)$. AMT9–20 gives, with the original arithmetic source w_h^{*k} ,

$$-rE_k^+ \leq F_K^{\text{ar}} - F_K^{(1)} \leq rE_k^-.$$

Let $\underline{U}_s = d_1^{(s)}$ and $\overline{U}_s = r(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - d_1^{(s)}$. The new finite conclusion is

$$\boxed{\max\{0, \underline{U}_1 - rE_k^+\} \leq F_K^{\text{ar}} \leq \overline{U}_1 + rE_k^-} \tag{AKS35}$$

This improves AMT37's absolute upper bound. For the sharper correlated version retain $d_{E,N} = q \log A_k - \log(\det G_N^{\text{ar}} / \det G_N^{(1)})$ and $c_{X,N} = \min\{r_X L_N^{\text{ar}}, d_{E,N}\}$, where $r_K = r$, $r_B = q - r$. Put $C_X^- = c_{X,q-1} + c_{X,q}$, $C_X^+ = c_{X,2q-1} + c_{X,2q}$. AMT18 then gives

$$\begin{aligned} \max\{0, \underline{U}_1 - C_K^-, \underline{U}_1 + \delta_k^\sigma - C_B^+\} &\leq F_K^{\text{ar}} \\ &\leq \overline{U}_1 + \min\{C_K^+, \delta_k^\sigma + C_B^-\}. \end{aligned} \tag{AKS36}$$

One may replace $d_{E,N}$ in the caps by the original full-source $\Xi_{k,N}$ or its proved finite majorant. The scalar $\delta_k^\sigma = \mathcal{B}_k^{\text{ar}} - \mathcal{B}_k^\sigma$ retains exactly AMT21:

$$\begin{aligned} \max\{-qE_k^+, -\Xi_{k,q-1} - \Xi_{k,q}\} &\leq \delta_k^\sigma \\ &\leq \min\{qE_k^-, \Xi_{k,2q-1} + \Xi_{k,2q}\}. \end{aligned} \quad (\text{AKS37})$$

Since $E_k^\pm = O_h(k + \log q)$ by the exact AMT22 expression, AKS35 and AKS32 prove the arithmetic version of AKS6, with an explicit $O_h(k^2)$ upper remainder. Nonnegativity follows from the actual nested source minima, as in AMT15, not from comparison with a possibly negative lower endpoint.

18.2 The original Gamma multiplier, not a replacement higher source

Retain $t_j = 2j + 1/2$, $f_l(S) = \prod_{j=0}^{l-1} (S - c - t_j)$ and

$$dm_{0,k} = \beta_k |f_l(c + iy)|^2 d\sigma, \quad \beta_k = \frac{(2\pi)^{k/2}}{\sqrt{2}\Gamma(k/2)}, \quad T_l = \prod_{j=0}^{l-1} t_j^2, \quad U_{l,N} = \prod_{j=0}^{l-1} [4(N + j + 1)^2 + t_j^2]. \quad (\text{AKS38})$$

The exact source map is $\mathcal{P}_N \rightarrow f_l \mathcal{P}_N \subset \mathcal{P}_{N+l}$, $P \mapsto f_l P$, with inverse division on this specified subspace. Let $\mathbf{E}_\chi$ be the complete divided-jet map and U_f its Leibniz multiplication matrix, $(U_f)_{r,a} = f_l^{(r-a)}(\lambda)/(r-a)!$ for $r \geq a$. AMT25–28 gives the precise transported observation and kernel

$$\begin{aligned} \tilde{\Lambda} &= \Lambda \mathbf{E}_\chi^{-1} U_f^{-1}, \quad \tilde{I} = U_f \mathbf{E}_\chi I, \quad \ker \tilde{\Lambda} = U_f \mathbf{E}_\chi K, \\ C_N &= \mathcal{K}_{\chi, N+l} - \mathcal{K}_{\chi, f; N+l} \mathcal{K}_{f, N+l}^{-1} \mathcal{K}_{f, \chi; N+l}, \\ H_{K,N}^{(k)} &= \beta_k \tilde{I}^* C_N^{-1} \tilde{I}. \end{aligned} \quad (\text{AKS39})$$

Each covariance is the original finite sum of full polynomial jet outer products divided by its original monic norm. These formulas retain the extra f constraints and their full cross covariance. In particular $M_f K = K$ is not assumed. AKS26 calculates the same tensor metric directly in its own orthonormal source; the two representations agree by their identical source norm and identical constrained minimum.

Define

$$w_N = \log(U_{l,N}/T_l), \quad \Omega^{\text{low}} = w_{q-1} + w_q, \quad \Omega^{\text{high}} = w_{2q-1} + w_{2q}.$$

The already proved same-coefficient comparison AMT42–44 retains all mass factors before cancellation and gives

$$-r\Omega^{\text{low}} \leq \Delta_K^\Gamma := F_K^{(1)} - F_K^{(k)} \leq r\Omega^{\text{high}}. \quad (\text{AKS40})$$

It therefore sharpens the two independent absolute bounds by

$$\begin{aligned} \max\{U_k, \underline{U}_1 - r\Omega^{\text{high}}, 0\} &\leq F_K^{(k)} \\ &\leq \min\{\bar{U}_k, \bar{U}_1 + r\Omega^{\text{low}}\}. \end{aligned} \quad (\text{AKS41})$$

The original finite estimate $0 \leq w_N \leq l \log[1 + 16(N + l)^2]$ proves that AKS40 is $o(kq)$ on the actual rank- r domain. Along with AKS35 this preserves the completed fact that all three source kernels differ by $o(kq)$. AKS4, not AKS40, is the new absolute-scale calculation.

18.3 Boundary, QGQ and arithmetic receiving identities

At each endpoint retain the original minimum section $S_N^{(s)} = (G_N^{(s)})^{-1} \Lambda^* (\Lambda(G_N^{(s)})^{-1} \Lambda^*)^{-1}$. The full Schur identity is

$$\det(I^* G_N^{(s)} I) \det Q_N^{(s)} = |\det[I, S_*]|^2 \det G_N^{(s)}, \quad Q_N^{(s)} = (\Lambda(G_N^{(s)})^{-1} \Lambda^*)^{-1}. \quad (\text{AKS42})$$

Thus the new finite absolute boundary interval is

$$\max\{0, \mathcal{B}_k^{(s)} - \overline{U}_s\} \leq F_B^{(s)} \leq \mathcal{B}_k^{(s)} - \underline{U}_s. \quad (\text{AKS43})$$

The baselines in this formula are the *complete* original BSL/HBL objects, with their full finite relation errors. No q^2 coefficient is assigned to K ; its proved coefficient there remains zero.

Write $\mathcal{H}_k = \mathcal{B}_k^\sigma - \mathcal{B}_k^0 = -\mathfrak{T}_k$. The exact full identities still read

$$\begin{aligned} F_B^{(1)} - F_B^{(k)} &= \mathcal{H}_k - \Delta_K^\Gamma, \\ \boxed{\mathcal{B}_k^{\text{ar}} = F_K^{(k)} + F_B^{(k)} + \mathcal{H}_k + \delta_k^\sigma = F_K^{(1)} + F_B^{(1)} + \delta_k^\sigma.} \end{aligned} \quad (\text{AKS44})$$

The Gamma interval is not re-estimated in this argument. For either complete QRI1 or QRI2 interval $[\omega^-, \omega^+]$ for its already calculated centre W_k , retain its exact source errors $e_0 \in [-a_0, b_0]$ and $e_q + e_{q+1} \in [-b_k^+, b_k^-]$. Then

$$\underline{\mathcal{H}}_k = -\omega^+ - b_k^- - a_0, \quad \overline{\mathcal{H}}_k = -\omega^- + b_k^+ + b_0, \quad \mathcal{H}_k \in [\underline{\mathcal{H}}_k, \overline{\mathcal{H}}_k]. \quad (\text{AKS45})$$

These are exactly QRI3, not new unspecified error choices. The complete QRI1–9, QGT and QGQ definitions remain verbatim in each updated receiver; the new interval is intersected with all their retained tighter finite intervals. In particular the simple-quartet nonzero error radius stays

$$\limsup \left| \frac{\mathcal{H}_k + lq\mathcal{C}_\Gamma}{q} + \frac{\partial_\alpha \mathcal{L}(2, \pi)}{8} - \frac{\log 2}{4} \right| \leq \frac{A_{\text{QGT}}}{\sqrt{2}}. \quad (\text{AKS46})$$

No zero-radius or $o(q)$ arithmetic assertion is introduced.

For example the complete finite absolute defect relative to the tensor boundary, with the actual Gamma and arithmetic scalars retained, is

$$\underline{U}_k + \mathcal{H}_k + \delta_k^\sigma \leq \mathcal{B}_k^{\text{ar}} - F_B^{(k)} \leq \overline{U}_k + \mathcal{H}_k + \delta_k^\sigma. \quad (\text{AKS47})$$

Inserting AKS41 makes this sharper. Replacing $\mathcal{H}_k$ by the appropriate endpoints of AKS45 and δ_k^σ by AKS37 gives its fully inherited finite scalar enclosure with the same signs. Also $F_B^{\text{ar}} = \mathcal{B}_k^\sigma + \delta_k^\sigma - F_K^{\text{ar}}$; subtract the interval AKS35 or AKS36 to get its new absolute boundary interval. This retains the exact correlations rather than taking independent hypothetical source errors.

The existing limit is $\mathcal{H}_k/(kq) \rightarrow -\mathcal{C}_\Gamma/4$ and $\delta_k^\sigma/(kq) \rightarrow 0$. Therefore the new interval implies

$$\begin{aligned} \liminf_k \inf_{|u| \geq R_*} \frac{\mathcal{B}_k^{\text{ar}} - F_B^{(k)}}{kq} &\geq -\mathcal{C}_\Gamma/4, \\ \limsup_k \sup_{|u| \geq R_*} \frac{\mathcal{B}_k^{\text{ar}} - F_B^{(k)}}{kq} &\leq \kappa_* - \mathcal{C}_\Gamma/4. \end{aligned} \quad (\text{AKS48})$$

All original branches are included in the infima and suprema. This does not decide its sign. The already proved boundary *difference* limits of AMT49 are unchanged. Every subsequential limit of the absolute normalized kernel lies in $[0, \kappa_*]$ and is shared by its three source versions when they are evaluated on the same sequence of period parameters. Existence of a single limit is not proved by boundedness.

Finally the HC residual remains exactly $\mathcal{R}_k = \mathcal{B}_k^{\text{ar}} - 4q \log(D_h k)$, with its original D_h and original nonnegative contraction, phase and trace terms. AKS44 changes none of them. The supplied positive q^2 baseline and its complete Gamma correction are not re-presented as new results here.

19 Actions, support faces, and the actual arithmetic class

The coefficient formula AKS12 supplies all four blocks of the same sum action: $P_K Y P_K$, $P_K Y (I - P_K)$, $(I - P_K) Y P_K$ and $(I - P_K) Y (I - P_K)$. There is no claim that either off-diagonal block vanishes. In the actual minimum-section coordinates the corresponding formula remains

$$[I, S_N]^{-1} Y [I, S_N] = \begin{pmatrix} H_{K,N}^{-1} I^* G_N Y I & H_{K,N}^{-1} I^* G_N Y S_N \\ \Lambda Y I & \Lambda Y S_N \end{pmatrix}. \quad (\text{AKS49})$$

Its proof is multiplication by the inverse coordinate maps $H_{K,N}^{-1} I^* G_N$ and Λ . In the invariant-ring model, the same action is the derivation $\mathcal{D}f = (\mathbf{H} \Omega \mathbf{H}^{-1} x) \cdot \nabla f$, $\Omega = \text{diag}(\gamma - i\delta, -\gamma - i\delta, \gamma + i\delta, -\gamma + i\delta)$, with $\mathcal{D}Q = 0$. For the actual conductor $\mathcal{A}_Q = \prod_{a=1}^4 Q_a$, its full defect is $\mathcal{D}\mathcal{A}_Q = \sum_{a=1}^4 (\mathcal{D}Q_a) \prod_{b \neq a, 1 \leq b \leq 4} Q_b$, exactly OQC22. Thus this coefficient realization preserves the original action leaving the observed dual.

In particular the section-corrected vector is still OPR2's

$$C_s r_j^{\ker} = e_{q+j} - S_{q+j-1}^{(s)} \Lambda e_{q+j} = I(\zeta_{q+j} - \kappa S_{q+j-1}^{(s)} \Lambda e_{q+j}),$$

with its full denominator $1 + \xi_j$ in the kernel increment. The operator proof estimates the complete covariance and its actual restriction, so none of these coupled terms is replaced by a raw fixed-section vector. The all-iterate kernel stays $\bigcap_{a=0}^{q-1} \ker(\Lambda Y^a)$; nothing in AKS4 declares it zero.

On the existing arithmetic coefficient realization retain $\mathcal{V}_{h,k} P = P(D_1 + \dots + D_k) F_h^{\otimes k}$ and $J^{(k)} \mathcal{V}_{h,k} = \eta \text{rem}_\chi$, where $\eta = U^{\otimes k} \iota$ is the original injection. AKS17 and the Newton coefficients compute precisely this same remainder. Any two polynomial lifts differ by χQ in the original relation space, so their original arithmetic cohomology classes agree. The specified primitive of the difference is EP.48, with the full differential and cochain signs proved in AKP3–5 below; it depends on the lift difference. The source comparison to w_h^{*k} is the identity on original polynomial coefficients, followed by the actual minima as in AMT8; it is not a replacement arithmetic source.

The scalar object stays $G(R) = \{\tau\} \sqcup R^\bullet$, with supported zero $0_R^\bullet$, injective amplitude/support pair, and infinite integer target. Every coefficient map above is applied on each original present support label and outer leg. It commutes with the specified coefficient inclusions because both compositions apply the same linear map to the same coefficient; a present zero is not converted into absence. No new base object, purity condition, or identification of the sum action with a Frobenius action is used. The actual distinguished class and its higher-jet directions remain in the full fibre, even when the observation kills them.

20 Exact period domain and the interior alternative

For reference the inherited radius used above is completely specified as follows; these are AMT33–34/RPC17–23, with no new constants. Put

$$\begin{aligned} \beta &= 2(\gamma^2 - \delta^2), \quad \eta = (\delta^2 + \gamma^2)^2, \quad a = v_h(\rho) \neq 0, \\ c_3 &= (a^{-2} - \bar{a}^{-2})/(4i\delta\gamma), \quad c_1 = (a^{-2} + \bar{a}^{-2})/2 - (\delta^2 - \gamma^2)c_3, \\ g_j &= 5^{j/5-1} e^{\pi i j/5} \Gamma(j/5), \quad g_+ = \max_j |g_j|, \quad g_-^{-1} = \max_j |g_j|^{-1}, \quad \kappa = \max_{j < 4} |g_{j+1}/g_j|, \\ E_j^{\text{ray}} &= \frac{8}{5} \int_0^\infty r^j (r^3/3 + r) e^{-r^5/5 + r^3/3 + r} dr, \quad C_T = 2 \left(\sum_{j=0}^3 (E_j^{\text{ray}})^2 \right)^{1/2}, \\ C_K &= 4C_T g_-^{-1} + 2g_+ g_-^{-1} + 2g_+ g_-^{-2} C_T, \\ C_{K,3} &= C_K(3\kappa^2 + 3\kappa C_K + C_K^2), \quad \epsilon_* = \min\{1, (2C_T g_-^{-1})^{-1}\}, \quad B_* = \beta + \eta. \end{aligned} \quad (\text{AKS50})$$

Then

$$R_* = \begin{cases} \left[\max \left\{ 1, \frac{B_*}{\epsilon_*}, \frac{2(|c_3|C_{K,3}B_* + |c_1|(\kappa + C_K))}{|c_3g_4/g_1|} \right\} \right]^{5/2}, & c_3 \neq 0, \\ \left[\max \left\{ 1, \frac{B_*}{\epsilon_*}, \frac{2C_KB_*}{|g_2/g_1|} \right\} \right]^{5/2}, & c_3 = 0. \end{cases} \quad (\text{AKS51})$$

The actual unit implies $(c_1, c_3) \neq (0, 0)$ and all integrals converge. RPC's full period comparison proves a nonzero $(4, 1)$ or $(2, 1)$ commutator entry, respectively, for $|u| \geq R_*$ on every branch. It therefore proves the five-member orbit needed in AKS11.

Inside this radius the same coefficient determinant AKS12–14 and the operator bound AKS24 remain valid. The orbit is decided by the original noninvariant quadratic coefficients, not by the radius alone. Since Q is nondegenerate, its projective orbit has either one or five members. A one-member orbit must have character one: its character has both fourth and fifth powers equal to one, by the quadratic determinant and $D^5 = I$. In that case $Q = a'x_1x_4 + b'x_2x_3$ with $a'b' \neq 0$, and

$$b = d_{5,k,0} - d_{5,k-2,0} = \left\lfloor \frac{(k+1)^2}{5} \right\rfloor + \mathbf{1}_{k \equiv 0 \text{ or } 3 \pmod{5}}, \quad r = q - b. \quad (\text{AKS52})$$

If a noninvariant coefficient is nonzero, AKS11 applies instead. These are the complete OQC alternatives. The bound AKS31 always uses that actual r ; it does not extend the $8k - 16$ specialization to an unevaluated exceptional point. This calculation does not assert that the interior exceptional set is empty.

21 Verification scope and the remaining absolute coefficient

The new mathematical content is AKS20–32 and its application to the actual coefficient determinant AKS12–14, followed by the finite receiving improvements AKS35–48. The complete source and orbit identifications used in that application are the supplied OCS/OQC/OQX, RPC, OPG/OPR and AMT results; their exact predecessors are preserved. AKS33 pinpoints a repair obtainable immediately from the earlier proof, whereas the Newton bound supplies the sharper explicit constant.

The bound *at* scale kq is now finite and uniform. What is not proved is a matching lower coefficient, a unique limit for $F_K^{(1)}/(kq)$, or a sign for the interval AKS48. Those questions involve the orientation of the original period kernel within the inverse remainder covariance; replacing that compression by a scalar bound loses precisely that information. This is the finite-bounds outcome allowed by the current request, not a claim to have evaluated an asymptotic coefficient. The source differences are already completed inputs and are not renamed as the remaining problem.

The accompanying exact checks distinguish finite algebraic diagnostics from the analytic proof for the full family. They do not assert that a diagnostic root is a zero of ξ , that a diagnostic matrix equals an actual period matrix, or that a new Lean theorem has been checked. No RH conclusion is drawn from this kernel bound.

22 The new ceiling in the original coefficient frame

The complete incoming AKS proof sharpens the previously proved KAF bound. Both estimates apply to the same original remainder and the same kernel. We retain all their finite constants. In particular, for $m = 1$, $k = 4l + 1$, $l \geq 1$, $q = (k + 1)^2$, $c = k/2$, $0 < \delta < 1/2$, $\gamma > 2$, and every original branch with $|u| \geq R_*$, put $r = 8k - 16$. The AKS constant is

$$\kappa_* = 32 \log \frac{25 \log 3}{4\pi} \in (25.02078097649, 25.02078097650). \quad (\text{AKJ1})$$

AKS20–32 proves the full finite operator inequality from confluent Newton division and coefficient extraction. Its parameters t and z are extraction radii in the existing source, whose orders remain $1, k$. For $s \in \{1, k\}$ define, using the exact KAF and AKS quantities,

$$\begin{aligned} L_s &= d_1^{(s)}, \\ U_s &= \min\{2r \log Z_{k,1}, r(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - d_1^{(s)}\}. \end{aligned} \quad (\text{AKJ2})$$

The first entry of the minimum is KAF’s factorial estimate; the second is AKS28, with its original first compressed increment retained. Thus $L_s \leq F_K^{(s)} \leq U_s$. This pointwise minimum retains whichever finite estimate is sharper at the actual original parameters. It does not assume that the asymptotically smaller ceiling wins at every degree.

The two exact original coordinate constructions are related as follows. Use OKM’s invertible ordered Vandermonde V_β and its actual quotient matrix π . Put $I_\pi = V_\beta^{-1} \pi^\top$. AKS12 constructs the coefficient-orthogonal projection P_K from the actual observation and its complete ordered-word Gram. Directly,

$$P_K = I_\pi (I_\pi^* I_\pi)^{-1} I_\pi^*, \quad D_{K,N}^{(s)} = \frac{\det(\bar{\pi}(R_N^{(s)})^{-1} \pi^\top)}{\det(I_\pi^* I_\pi)}. \quad (\text{AKJ3})$$

Indeed I_π is injective with image K by OKM5. The first displayed matrix is Hermitian and idempotent, fixes this image, and kills its coefficient-orthogonal complement, so it is exactly AKS12’s projection. Apply AKS14 with inclusion I_π and then OKM8–9 to obtain the second identity. The positive denominator is independent of endpoint and source. Its exponent is $1 + 1 - 1 - 1 = 0$ in the four-endpoint quotient. This proves the exact morphism between the incoming projector determinant and the existing conductor-frame determinant, retaining the original coefficient metric and every frame factor.

With the original AMT correlated caps $C_X^\pm$ and scalar δ_k^σ , the following joint intervals hold:

$$\begin{aligned} \max\{0, L_1 - C_K^-, L_1 + \delta_k^\sigma - C_B^+\} &\leq F_K^{\text{ar}} \\ &\leq U_1 + \min\{C_K^+, \delta_k^\sigma + C_B^-\}, \\ \max\{L_k, L_1 - r\Omega^{\text{high}}, 0\} &\leq F_K^{(k)} \leq \min\{U_k, U_1 + r\Omega^{\text{low}}\}. \end{aligned} \quad (\text{AKJ4})$$

For proof, add each retained AMT lower or upper source-difference bound to the corresponding endpoint of AKJ2, and intersect the resulting intervals. In the first line the upper and lower source differences are the same AMT18 correlated caps used in AKS36; no independent errors have replaced their actual values. Every boundary interval is obtained by subtracting these kernel intervals from the exact original total $\mathcal{B}_k^a = F_K^a + F_B^a$. Consequently all three original kernels have uniform limsup at most κ_* after division by kq . The source differences remain $o(kq)$. A matching leading value is still to be calculated from these same matrices.

23 Residue detection expressed in every original primary coefficient

Retain now every original multiplicity $m \geq 1$, $e = 1 + k(m - 1)$, $q = e(k + 1)^2$, and the full polynomial

$$\chi(S) = \prod_{a,b=0}^k (S - \beta_{ab})^e, \quad \beta_{ab} = c + (2a - k)\delta + i(2b - k)\gamma, \quad E = \mathbb{C}[S]/(\chi). \quad (\text{AKJ5})$$

All distinct pairs give distinct roots. For a root β write $z = S - \beta$, $h_\beta(S) = \chi(S)/(S - \beta)^e$, and let ε_β be its original CRT idempotent. Define exactly $\ell([f]) = [S^{q-1}] \text{rem}_\chi f$. For every polynomial class the primary expansion gives

$$\ell(f) = \sum_{\beta} [z^{e-1}] \frac{f(\beta + z)}{h_\beta(\beta + z)}. \quad (\text{AKJ6})$$

The quotient on the right is a truncated power series through degree $e - 1$. To prove the identity, partial-fraction division of f/χ gives at each finite pole the coefficient of $(S - \beta)^{-1}$ shown on the right. Expanding those principal parts at infinity, only their first-order terms contribute to S^{-1} . Polynomial terms contribute nothing. For the unique representative of degree below q , the same S^{-1} coefficient is its coefficient of S^{q-1} . This proves AKJ6 without deleting any repeated-root term. Since $h_\beta(\beta) \neq 0$, the bilinear pairing $(f, g) \mapsto \ell(fg)$ is nondegenerate: if the first nonzero local coefficient of f is $f_t z^t$, take $g = \varepsilon_\beta z^{e-1-t} h_\beta$ to obtain $\ell(fg) = f_t \neq 0$.

Let λ_ν be the specified zero-charge coefficient rows of the original OCS observation, so the injection $\mathcal{F}_0$ gives $j\Lambda = \mathcal{F}_0 \mathfrak{A} \mathcal{H}^{-1}$. Retain the exact entries

$$a_{\nu,\beta,D} := \lambda_\nu(\varepsilon_\beta z^D) = i^D D! c_{\nu,abD}, \quad 0 \leq D < e. \quad (\text{AKJ7})$$

Here $c_{\nu,abD}$ is the full OCS multinomial in the original period and unit jets; $\beta = \beta_{ab}$. It is not an assigned coefficient. The residue representative of each row has the explicit formula

$$A_\nu = \left[\sum_{\beta} \varepsilon_\beta h_\beta(S) \sum_{D=0}^{e-1} a_{\nu,\beta,D} (S - \beta)^{e-1-D} \right], \quad \lambda_\nu(x) = \ell(x A_\nu). \quad (\text{AKJ8})$$

Indeed, restrict to one primary factor and substitute AKJ8 into AKJ6. For $x = \varepsilon_\beta z^D$, multiplication cancels h_β and extracts exactly $a_{\nu,\beta,D}$. These vectors are a basis, proving the asserted equality for every x . This gives the concrete inverse of the residue equivalence used in the incoming formalization, on the actual OCS rows, with their factorials and phases.

Define the integer t_β from these actual coefficients by

$$t_\beta = \begin{cases} -1, & a_{\nu,\beta,D} = 0 \text{ for every } \nu, D, \\ \max\{D : a_{\nu,\beta,D} \neq 0 \text{ for some } \nu\}, & \text{otherwise,} \end{cases} \quad n_\beta = e - 1 - t_\beta. \quad (\text{AKJ9})$$

Define $\mathcal{O}_q : E \rightarrow B^q$ by $\mathcal{O}_q(x) = (\Lambda M_S^j x)_{0 \leq j < q}$, where $B = \text{im } \Lambda$ is the original observation image. Thus $0 \leq n_\beta \leq e$. The complete original permanently invisible submodule and its exact dimension are

$$\begin{aligned} K_\infty &= \bigcap_{j=0}^{q-1} \ker(\Lambda M_S^j) = \bigoplus_{\beta} \varepsilon_\beta (z^{e-n_\beta}) \subset E, \\ \dim K_\infty &= \sum_{\beta} n_\beta, \quad \text{rank } \mathcal{O}_q = q - \sum_{\beta} n_\beta. \end{aligned} \quad (\text{AKJ10})$$

Here $(z^e) = 0$ and (z^0) is the entire local algebra. To prove this, vanishing of the first q observations says $\ell(S^j x A_\nu) = 0$ for all $j < q, \nu$. Nondegeneracy just proved and the power basis imply $x A_\nu = 0$

for every row. Conversely those products being zero kills every iterate. At the factor β , AKJ8 is a unit h_β times its displayed reverse coefficient polynomial. The minimum valuation of all A_ν is precisely n_β . Its common annihilator is (z^{e-n_β}) : multiplication by a coefficient of minimal valuation is a unit times z^{n_β} , and all other coefficients have at least that valuation. This proves the module formula and its dimension; rank-nullity proves the last equality. Affine substitution $M_S = c + iY$ gives an invertible triangular change between the first q powers of the two operators, so the same kernel is obtained using the original Y -stack of OCS.

In particular all iterates are jointly injective exactly when every original highest-jet column $a_{\nu,\beta,e-1}$ has a nonzero entry. For $m = 1$, AKJ8 reduces to

$$A_\nu(\beta) = \chi'(\beta)B_{\nu,\beta}, \quad K_\infty = \text{span}\{\varepsilon_\beta : B_{\nu,\beta} = 0 \text{ for all } \nu\}. \quad (\text{AKJ11})$$

This is an exact test on the AKS8 matrix, not an assumption that it passes. It constructs the inclusion $K_\infty \hookrightarrow K$: each listed primary ideal is killed by all rows, hence in particular by their original one-step observation. The source-minimum metric on this subspace is its restriction from OKM8, retaining the same original Gram. Full-iterate injectivity alone therefore supplies no estimate of the one-step kernel determinant AKJ3.

24 The actual conductor confines the kernel's polynomial degrees

Return to $m = 1$, $k = 4l + 1 \geq 9$ and the original five-member period domain. Let $\mathcal{A}_Q = \prod_{j=1}^4 Q_j$ and write its full Segre restriction from OCF as

$$A = \sum_{r,s=0}^8 a_{rs} X_1^r X_0^{8-r} Y_1^s Y_0^{8-s} \neq 0. \quad (\text{AKJ12})$$

OCF proves nonvanishing and retains all actual H coefficients. No particular a_{rs} is presumed nonzero. Write $k' = k - 8$, $q' = (k - 7)^2$ and

$$b_{rs} = 4 + (2r - 8)\delta + i(2s - 8)\gamma, \quad (\mathcal{T}_A p)(S') = \sum_{r,s=0}^8 a_{rs} p(S' + b_{rs}), \quad \mu_j = \sum_{r,s} a_{rs} b_{rs}^j. \quad (\text{AKJ13})$$

All 81 numbers b_{rs} are distinct. Their Vandermonde is invertible, so some μ_j with $0 \leq j \leq 80$ is nonzero. Define v to be the first such index. The exact finite differential expression on degree-below- q polynomials is

$$\mathcal{T}_A = \sum_{j=0}^{q-1} \frac{\mu_j}{j!} \partial_S^j = \frac{\mu_v}{v!} \partial_S^v B(\partial_S), \quad B(X) = 1 + \sum_{j=v+1}^{q-1} \frac{v! \mu_j}{j! \mu_v} X^{j-v}. \quad (\text{AKJ14})$$

This is Taylor's finite formula. Hence it kills polynomials of degree below v and lowers every higher degree by exactly v . The operator $B(\partial_S)$ is invertible: with $U = B(\partial_S) - I$, its inverse is the finite sum $\sum_{a=0}^{q-1} (-U)^a$. Terms beyond nilpotency are zero. Thus $\mathcal{T}_A$ maps polynomials of degree at most d onto those of degree at most $d - v$ when $d \geq v$.

Let $\chi_{k'}$ be the original polynomial for tensor degree k' . The conductor's exact annihilator in the original coefficient algebra is

$$V_A = \{p \in \mathbb{C}[S]_{< q} : \chi_{k'} \mid \mathcal{T}_A p\}, \quad K \subset V_A. \quad (\text{AKJ15})$$

To prove this map, pair the biform Ag , of bidegree (k, k) , with the primary values of p . If g has coefficient g_{ab} at bidegree (k', k') , convolution of all original coefficients gives the pairing

$$\sum_{a,b=0}^{k'} g_{ab} \sum_{r,s=0}^8 a_{rs} p(\beta_{ab}^{(k')} + b_{rs}) = \sum_{a,b=0}^{k'} g_{ab} (\mathcal{T}_A p)(\beta_{ab}^{(k')}). \quad (\text{AKJ16})$$

The addition identity for the roots follows directly from their retained centres and both signs. Vanishing for every g is exactly vanishing at all q' distinct roots, hence the divisibility in AKJ15. OCF's proved inclusion $AC[X, Y]_{(k', k')} \subset J_k$ then gives $K \subset V_A$ by its original complex-linear coefficient pairing.

The following is a full degree-adapted basis of V_A . Define $\mathcal{J}_v S^d = d! S^{d+v} / (d+v)!$, so $\partial_S^v \mathcal{J}_v = I$, and set

$$1, S, \dots, S^{v-1}, \quad P_h = \frac{(q' + h + v)!}{(q' + h)!} B(\partial_S)^{-1} \mathcal{J}_v(\chi_{k'} S^h), \quad (AKJ17)$$

$$0 \leq h \leq q - q' - v - 1.$$

The first list is empty when $v = 0$. Since $q - q' = 16k - 48 \geq 96 > v$, the range of the second list is nonnegative. Each P_h is monic of degree $q' + h + v$ and maps under $\mathcal{T}_A$ to $\mu_v(q' + h + v)! / (v!(q' + h)!)$ times $\chi_{k'} S^h$. These statements follow from AKJ14, the inverse's constant term one, and the leading coefficient of $\mathcal{J}_v$. The images span every multiple of $\chi_{k'}$ of degree at most $q - v - 1$, and the low list spans the exact kernel of $\mathcal{T}_A$. This proves spanning and independence. In particular $\dim V_A = q - q' = 16k - 48$.

Choose the original OCF basis of $J_k / (AC[X, Y]_{(k', k')})$, whose exact dimension is $j_k = 8k - 32$. Evaluate each chosen biform row on the primary values of every polynomial in AKJ17. This gives an explicit j_k by $(16k - 48)$ matrix Ξ , with the full original period coefficients. Its kernel maps isomorphically onto K by the displayed basis. Its rank is j_k : the original J_k and conductor annihilator duality, or dimension $\dim V_A - \dim K$, gives this rank. For the submatrix $\Xi_{\leq d}$ containing only basis columns of degree at most d , the exact formula is

$$\dim(K \cap \mathbb{C}[S]_{\leq d}) = \#\{\text{degrees in AKJ17 at most } d\} - \text{rank } \Xi_{\leq d}. \quad (AKJ18)$$

Distinct monic degrees ensure that no omitted higher column can cancel to form a degree-at-most- d polynomial. Restricting the coefficient isomorphism to this prefix and applying rank-nullity proves AKJ18. It uses actual minors and makes no assertion of a generic prefix rank.

Let $d_0 < \dots < d_{r-1}$ be the degree jumps of the actual kernel. The basis degrees of V_A , in increasing order, are i for $i < v$, and $q' + i$ for $i \geq v$. Inclusion and dimension of each degree prefix, together with $K \subset \mathbb{C}[S]_{< q}$, imply

$$i + q' \mathbf{1}_{i \geq v} \leq d_i \leq q - r + i. \quad (AKJ19)$$

For the lower inequality the i th independent kernel vector cannot appear earlier than the i th independent vector of V_A . For the upper inequality, intersection of an r -dimensional subspace of $\mathbb{C}^q$ with its $(q - r + i + 1)$ -dimensional degree prefix has dimension at least $i + 1$. Summing yields the exact exterior-degree bounds

$$\frac{r(r-1)}{2} + \max\{r-v, 0\}q' \leq \sum_{i=0}^{r-1} d_i \leq r(q-r) + \frac{r(r-1)}{2}, \quad (AKJ20)$$

$$0 \leq rq - \sum_i d_i \leq \frac{r(r+1)}{2} + rj_k + \min\{r, v\}q'.$$

The difference between the first line's upper and lower endpoints is $rj_k + \min\{r, v\}q'$, since $q - q' = r + j_k$. Therefore the original kernel's exterior degree is $rq - O(k^2)$, with the exact bound above and $v \leq 80$, uniformly in the proved period domain. The affine map $S = c + iy$ preserves all degree prefixes and multiplies a degree- d leading coefficient by i^d . Thus these are also the degrees in the original Gamma coordinate, with all phases known. They specify the kernel's position without assigning a condition number to its frame.

25 The exact conductor translations in the original source norm

Keep either source $s \in \{1, k\}$, $b = s/2$, $M = (2\pi)^{s/2}$ and the original $p_n, h_n = Mn!(b)_n$. On polynomials of degree at most D , let $\varphi_n = p_n / \sqrt{h_n}$ and $w_n = n! / (b)_n$. Differentiating the original

generating identity AKS16 in y gives

$$\partial_y \varphi_n = \sum_{\substack{1 \leq a \leq n \\ a \text{ odd}}} \frac{(-1)^{(a-1)/2}}{a} \sqrt{\frac{w_n}{w_{n-a}}} \varphi_{n-a}. \quad (\text{AKJ21})$$

Indeed differentiation multiplies the generating series by $\arctan z = \sum_{a \text{ odd}} (-1)^{(a-1)/2} z^a / a$. Equating its finite coefficient and substituting the full norms gives exactly AKJ21; the two copies of M are cancelled only at this point.

Write $H_D = \sum_{j=1}^D 1/j$ and $H_0 = 0$. The original source norm obeys

$$\|\partial_y\| \leq \sqrt{H_D(H_D + 4)} \leq H_D + 2, \quad \|f(y+z)\| \leq e^{|z|(H_D+2)} \|f\|. \quad (\text{AKJ22})$$

Here is a proof retaining both source orders. For $n > m \geq 1$ and $b \geq 1/2$,

$$\frac{w_n}{w_m} \leq \prod_{j=m+1}^n \frac{j}{j-1/2} \leq \sqrt{\frac{n}{m}}, \quad w_n \leq 2\sqrt{n} \quad (n \geq 1). \quad (\text{AKJ23})$$

The factor inequality follows by squaring and using $j(j-1) \leq (j-1/2)^2$; the product telescopes. For $m = 0$, separate the first factor, at most two. Apply the weighted Schur estimate to the absolute matrix in AKJ21 with positive weights $\vartheta_n = 1/\sqrt{w_n}$. Its weighted row sum is at most H_D . Its weighted column sum at n is at most $H_n + 4$: for $1 \leq m < n$ the excess over $1/(n-m)$ is at most

$$\frac{\sqrt{n/m} - 1}{n-m} = \frac{1}{\sqrt{m}(\sqrt{n} + \sqrt{m})} \leq \frac{1}{\sqrt{nm}},$$

whose sum is at most two by comparison with $\int_0^n t^{-1/2} dt$. The $m = 0$ term is at most $2/\sqrt{n} \leq 2$. Enlarging the odd-index sums to all indices preserves these bounds. For completeness, if a nonnegative matrix has weighted row bound A and column bound B , Cauchy–Schwarz gives $|(Tx)_m|^2 \leq A \vartheta_m \sum_n T_{mn} |x_n|^2 / \vartheta_n$; summing and using the column bound gives $\|Tx\|^2 \leq AB \|x\|^2$. This proves the first bound in AKJ22, including $D = 0$ where the derivative is zero. The exact finite Taylor identity $f(y+z) = \exp(z\partial_y)f(y)$ and its norm series prove the second. The inverse translation is obtained by $-z$ with the same bound.

There is consequently an exact original-source upper control for AKJ13. For $f(y) = p(c_k + iy)$, evaluate its output at $c_{k-8} + iy$, retaining $c_k = k/2$ and $c_{k-8} = k/2 - 4$. Then

$$(\mathcal{T}_{Ap})(c_{k-8} + iy) = \sum_{r,s=0}^8 a_{rs} f\left(y + \frac{b_{rs} - 4}{i}\right), \quad (\text{AKJ24})$$

and therefore

$$\|\mathcal{T}_{Ap}\|_{s, c_{k-8}} \leq \left(\sum_{r,s} |a_{rs}| e^{|b_{rs}-4|(H_D+2)} \right) \|p\|_{s, c_k}. \quad (\text{AKJ25})$$

The notation $\|p\|_{s,c} = \|p(c + iy)\|_{m_0,s}$ changes only the explicit centre; the measure, mass and source order s are unchanged. AKJ24 proves the centre correction rather than discarding it. For a fixed original period and quartet the coefficient in AKJ25 grows at most polynomially in D , since $H_D \leq 1 + \log D$ for $D \geq 1$ and $|b_{rs} - 4| \leq 8\sqrt{\delta^2 + \gamma^2}$. The entire actual coefficient sum remains visible. The finite inverse in AKJ17 has the specified moment denominators μ_v ; AKJ25 asserts no bound on that inverse. Together AKJ18–25 provide exact degree information and a source-norm estimate for the actual conductor map, to be used in the remaining four-endpoint compression calculation. No leading kernel value or kernel-invariance assumption has replaced that calculation.

26 The original theta class and the checked metric interface

The absolute base and original coefficient resolution are retained:

$$\mathbf{F}_{1,\tau} = \{\tau, 1\}, \quad A_\tau(\mathcal{T}) = [V \oplus V \xrightarrow{\Theta(\phi-\hat{\psi})} \mathcal{B}], \quad Q = \mathcal{B}/\Theta V. \quad (\text{AKP1})$$

The four-point chart has $+, - < \eta < \sigma$ and coefficients $(V, V, \mathcal{B}, 0)$, with the original restrictions Θ and $J\Theta = \Theta\mathcal{F}$. Its tensor resolution identifies the top cohomology with $Q^{\otimes k}$. AT1–2a supplies the complete original construction. In its notation, retaining the full unit $v_h = j_h(2\xi/h)$, the actual cochain and arithmetic class maps are

$$\begin{aligned} \mathcal{V}_{h,k}(P) &= P(D_1 + \cdots + D_k)F_h^{\otimes k}, & E &= \mathbb{C}[S]/(\chi), \\ \eta_{h,k}[P] &= v_h^{\otimes k} P(A_h^{(k)})1, & q^{(k)}\mathcal{V}_{h,k}(P) &= \sigma_h^{\otimes k} \eta_{h,k}[P]. \end{aligned} \quad (\text{AKP2})$$

Here $\sigma_h : E_h \hookrightarrow Q$ is the original section and $\eta_{h,k} : E \hookrightarrow E_h^{\otimes k}$ is the full-jet injection. Both injections are part of the supplied source proof; the cochain map has not been replaced by an arbitrary observation.

For completeness, the exact representative-change calculation is as follows. If $P' = P + \chi\kappa$, perform successive monic divisions of $\chi(s_1 + \cdots + s_k)$ by $h(s_a)$. Its final remainder is zero because χ is the original annihilator of the sum in $\bigotimes_a \mathbb{C}[s_a]/(h)$. Thus the actual divisions produce $Q_a(\mathbf{s})$ satisfying

$$\chi(s_1 + \cdots + s_k) = \sum_{a=1}^k h(s_a) Q_a(\mathbf{s}). \quad (\text{AKP3})$$

The chosen ordered division determines these polynomials; no uniqueness of all such decompositions is asserted. On the original tensor complex set

$$\Psi_\kappa = \sum_{a=1}^k (-1)^{a-1} Q_a(\mathbf{D}) \kappa \left(\sum_b D_b \right) (F_h^{\otimes(a-1)} \otimes \phi_0 \otimes F_h^{\otimes(k-a)}). \quad (\text{AKP4})$$

Its differential is

$$d\Psi_\kappa = \mathcal{V}_{h,k}(\chi\kappa) = \mathcal{V}_{h,k}(P') - \mathcal{V}_{h,k}(P). \quad (\text{AKP5})$$

Indeed the only nonzero differential in each summand acts at its ϕ_0 position. The tensor sign is $(-1)^{a-1}$, cancelling the displayed sign. The identity $\Theta\phi_0 = h(D)F_h$ gives the a th term of AKP3; all operators commute with this differential. Summing proves AKP5 with the original signs. This is EP.48, expressed at the current receiving map. Consequently the quotient classes in AKP2 agree. The chosen primitive for their difference depends on κ . At an existing support label ℓ , this quotient map sends the relation to $(\ell, 0)$ and sends external absence τ to τ . The coordinate change has therefore preserved the arithmetic class and its support through an explicit cochain homotopy.

This calculation specifies the correction to the sentence following AKS49: equality of cohomology classes is retained, with AKP4 the primitive of the change. Membership in an additional period-observation kernel alone supplies no such primitive. Its proved map is the inclusion $K_\infty \hookrightarrow K \hookrightarrow E$ of AKJ10–11, followed by the original arithmetic injection in AKP2. In particular, a nonzero coefficient in this additional kernel remains a nonzero original arithmetic class under that injection. Its zero observed value and its arithmetic class are related by these exact maps.

26.1 The complete minimum-section calculation at each endpoint

Let G be one of the original positive Hermitian coefficient Grams, write $K_G = G^{-1}$, and let $\Lambda : E \rightarrow B$ be the actual observation onto its image. Define

$$Q_G = (\Lambda K_G \Lambda^*)^{-1}, \quad S_G = K_G \Lambda^* Q_G. \quad (\text{AKP6})$$

The inverse exists because Λ^* is injective and K_G is positive definite. Direct multiplication, using Hermitian symmetry, gives

$$\Lambda S_G = I_B, \quad S_G^* G = Q_G \Lambda, \quad S_G^* G S_G = Q_G. \quad (\text{AKP7})$$

For every rectangular column family X put $Z = \Lambda X$ and $R = X - S_G Z$. Then

$$\Lambda R = 0, \quad R^* G S_G = 0, \quad R^* G R = X^* G X - Z^* Q_G Z \succeq 0. \quad (\text{AKP8})$$

The first two equalities follow from AKP7. In the expansion of the last Gram, each cross term is $Z^* Q_G Z$ and the section Gram is the same matrix. The resulting single subtraction proves the identity, including every off-diagonal pairing. Any other lift is $S_G Z + R'$ with $\Lambda R' = 0$. Its Gram is $Z^* Q_G Z + R'^* G R'$, proving the minimum property by the same orthogonality calculation.

Retain a fixed section S_* , the actual inclusion I_K , and original coordinate map κ_K satisfying $I_K \kappa_K = I_E - S_* \Lambda$. Applying this identity to R gives

$$R = I_K (\kappa_K X - \kappa_K S_G \Lambda X). \quad (\text{AKP9})$$

Thus the fixed-section coordinates include their specified minimum correction. For one original new column e_D , define $t_D = e_D^* G e_D$, $\lambda_D = \Lambda e_D$ and $\xi_D = \lambda_D^* Q_G \lambda_D$. AKP8 yields $t_D - \xi_D \geq 0$, so the exact ordered factors satisfy

$$(1 + \xi_D) \left(1 + \frac{t_D - \xi_D}{1 + \xi_D} \right) = 1 + t_D. \quad (\text{AKP10})$$

The denominator is positive and is retained. These are the full OPR minimum-section identities, now proved at the original Grams of AKS26 and OKM8. For an empty observation image, the section is the zero map, the Gram on that image has determinant one, and the same identities reduce to $R = X$. No empty present support is deleted.

The new formalization in draft PR32 checks these finite identities and the original observation-iterate and residue-annihilator maps. Its checked implementation is

8e78bc7c240b04d297ade6afdadfd863e0c6db7b.

The later source and status head is

9cec8482f2ecf978cf8bdb67b4c080b9dd74f5d6.

AKJ6–11 evaluates the residue inverse in every actual primary-jet coefficient, giving the exact map to those checked declarations. The analytic AKS estimate, the conductor metric estimates, and the polynomial-gcd dimension corollary have the full written proofs stated here or in their supplied sources; they are outside that Lean run.

27 The finite conductor inverse in the two original Gamma norms

Current refinement. The finite estimates in this chapter remain valid. RUI5–15 and CEP29–49 below sharpen their exterior consequence: complementary singular values and the exact source quotient give an $o(kq)$ conductor error at all four original endpoints, for a fixed actual period. The full original constants, moments and low-degree graph are retained in those proofs.

Retain AKJ12–25 and the complete OCF conductor coefficient polynomial. Thus $m = 1$, $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $0 < \delta < 1/2$, $\gamma > 2$, and the original period belongs to the proved five-orbit locus. Its fixed 4 by 4 matrix $\mathbf{H}$ determines the four quadrics and all 81 coefficients a_{rs} of their full product. They remain the same coefficients as the tensor degree k varies. Put

$$b_{rs} = 4 + (2r - 8)\delta + i(2s - 8)\gamma, \quad \mu_j = \sum_{r,s=0}^8 a_{rs} b_{rs}^j, \quad v = \min\{j : \mu_j \neq 0\} \leq 80. \quad (\text{AIE1})$$

The existence and bound for v are the full distinct-node Vandermonde argument of AKJ13. Throughout, $0 \leq D \leq q - 1$ denotes the polynomial degree cutoff. Every norm below is the original Gamma norm of source order $s \in \{1, k\}$, with

$$b = s/2, \quad M = (2\pi)^{s/2}, \quad h_n = Mn!(b)_n, \quad (\text{AIE2})$$

$$\varphi_n = p_n/\sqrt{h_n}, \quad w_n = n!/(b)_n, \quad \|p\|_{s,c} = \|p(c + iy)\|_{m_{0,s}}.$$

The centre c is displayed wherever it changes. No source order, source mass, coefficient, or period is assigned a replacement value.

27.1 A coefficient-controlled disk for the literal finite symbol

The exact symbol used in AKJ14 and AKJ17 is

$$B_q(z) = 1 + \sum_{j=v+1}^{q-1} \frac{v! \mu_j}{j! \mu_v} z^{j-v}. \quad (\text{AIE3})$$

The sum is empty if its lower index exceeds its upper index. Define the positive, original-data constants

$$A_*^{\text{abs}} = \sum_{r,s} |a_{rs}|, \quad B_*^{\text{sh}} = \max_{r,s} |b_{rs}|, \quad C_v = \frac{A_*^{\text{abs}} (B_*^{\text{sh}})^v}{|\mu_v|}, \quad (\text{AIE4})$$

$$R_0 = \frac{1}{B_*^{\text{sh}}} \min \left\{ 1, \frac{v+1}{2eC_v} \right\}, \quad \rho_0 = \tanh R_0.$$

Here $A_*^{\text{abs}} > 0$, $B_*^{\text{sh}} \geq 4$, $C_v \geq 1$, $R_0 > 0$, and $0 < \rho_0 < 1$. The superscripts distinguish these constants from the original conductor polynomial and from every earlier radius or pivot coefficient. The exact moment μ_v , including any cancellation in it, remains in the denominator.

For every $|z| \leq R_0$ one has

$$|B_q(z) - 1| \leq \frac{v!}{|\mu_v|} \sum_{j=v+1}^{q-1} \frac{|\mu_j|}{j!} |z|^{j-v} \leq C_v \frac{B_*^{\text{sh}} |z|}{v+1} \exp(B_*^{\text{sh}} |z|) \leq \frac{1}{2}. \quad (\text{AIE5})$$

To prove the middle inequality, use $|\mu_j| \leq A_*^{\text{abs}} (B_*^{\text{sh}})^j$, write $j = v+1+h$, and apply $v!/(v+1+h)! \leq 1/((v+1)h!)$. Summation of the exponential series gives the displayed bound. Definition AIE4 gives $B_*^{\text{sh}} R_0 \leq 1$ and $C_v B_*^{\text{sh}} R_0/(v+1) \leq 1/(2e)$, proving the last inequality. The first upper bound in AIE5 retains every individual finite moment and may be used directly for a larger verified radius. The explicit R_0 proves a positive disk for every actual coefficient chart; no uniform lower bound for $|\mu_v|$ over varying periods is asserted.

The analytic function

$$F_q(z) = \frac{1}{B_q(-i \arctan z)} = \sum_{a=0}^{\infty} f_{q,a} z^a \quad (\text{AIE6})$$

is well defined on a neighbourhood of the closed disk $|z| \leq \rho_0$, and $|F_q(z)| \leq 2$ there. Indeed the power series of $\arctan$ gives $|\arctan z| \leq \operatorname{arctanh} |z| \leq R_0$; AIE5 gives $|B_q(-i \arctan z)| \geq 1/2$. Compactness gives an open neighbourhood with nonzero denominator. Cauchy's coefficient formula therefore proves

$$f_{q,0} = 1, \quad |f_{q,a}| \leq 2\rho_0^{-a} \quad (a \geq 1). \quad (\text{AIE7})$$

Although the literal symbol B_q changes with q , the disk and its bound depend only on the original fixed conductor and shifts.

27.2 The exact coefficient map and its norm

Let $\mathcal{P}_D$ denote the polynomials in y of degree at most D , with the orthonormal basis in AIE2. Define the lowering map

$$L_s \varphi_0 = 0, \quad L_s \varphi_n = \sqrt{\frac{w_n}{w_{n-1}}} \varphi_{n-1} \quad (1 \leq n \leq D). \quad (\text{AIE8})$$

Its powers have exactly the entries $L_s^a \varphi_n = \sqrt{w_n/w_{n-a}} \varphi_{n-a}$ for $a \leq n$ and vanish for $a > n$. Thus $L_s^{D+1} = 0$. AKJ21 gives the polynomial identity $\partial_y = \arctan L_s$, with every power above degree D zero. For the coordinate pullback $\mathcal{C}_c p(y) = p(c + iy)$, the chain rule is $\mathcal{C}_c \partial_S \mathcal{C}_c^{-1} = -i \partial_y$. Consequently the inverse of the literal finite operator is

$$\begin{aligned} \mathcal{C}_c B_q(\partial_S)^{-1} \mathcal{C}_c^{-1} &= F_q(L_s), \\ F_q(L_s) \varphi_n &= \sum_{a=0}^n f_{q,a} \sqrt{\frac{w_n}{w_{n-a}}} \varphi_{n-a}. \end{aligned} \quad (\text{AIE9})$$

For proof, evaluate the analytic power-series identity $B_q(-i \arctan z) F_q(z) = 1$ in the nilpotent matrix L_s . All relevant coefficients are finite, so multiplication is an exact finite polynomial identity. This also proves that AIE9 is the same inverse as the finite nilpotent Neumann sum in AKJ14. The factor $-i$ is forced by the original coordinate map and has been retained.

Define the exact weight ratio and the resulting bound

$$\begin{aligned} \alpha_{D,s} &= \max_{0 \leq m \leq n \leq D} \sqrt{\frac{w_n}{w_m}}, \\ N_{D,s} &= \alpha_{D,s} \left(1 + 2 \sum_{a=1}^D \rho_0^{-a} \right) \leq \frac{2\alpha_{D,s}}{1 - \rho_0} \rho_0^{-D}. \end{aligned} \quad (\text{AIE10})$$

Every absolute row sum and every absolute column sum of AIE9 is at most $\alpha_{D,s} \sum_{a=0}^D |f_{q,a}|$. Cauchy–Schwarz in each row followed by the column bound proves that the operator norm is at most this number; AIE7 gives $N_{D,s}$. It follows that, at every unchanged centre c ,

$$\|B_q(\partial_S)^{-1} p\|_{s,c} \leq N_{D,s} \|p\|_{s,c}. \quad (\text{AIE11})$$

When $s = k$, $b = k/2 \geq 1$ and $w_n/w_{n-1} = n/(b+n-1) \leq 1$, so $\alpha_{D,k} = 1$. When $s = 1$, the same ratio is greater than one, so $\alpha_{D,1} = \sqrt{D!/(1/2)_D}$; for $D \geq 1$, the exact factor estimate of AKJ23 yields $\alpha_{D,1} \leq \sqrt{2} D^{1/4}$. At $D = 0$, both values are one. In particular

$$\log \|B_q(\partial_S)^{-1}\|_{s,c} \leq D \log(\coth R_0) + \log \frac{2\alpha_{D,s}}{1 - \tanh R_0} = O_{H,\delta,\gamma}(D + \log(D+1)). \quad (\text{AIE12})$$

For the fixed original quartet and period, $q = (k+1)^2$ and $D \leq q-1$ imply that this logarithm divided by kq tends to zero. This estimate holds for both original source orders at once; its constants explicitly retain the actual μ_v .

27.3 The infinite symbol differs by an exact low-degree map

To record precisely the truncation in AIE3, define the entire function

$$B_\infty(z) = \frac{v!}{\mu_v z^v} \sum_{r,s} a_{rs} e^{b_{rs} z}, \quad B_\infty(0) = 1. \quad (\text{AIE13})$$

Its apparent singularity is removable by the definition of v . On polynomials of degree at most $D \leq q-1$, set $E_q = B_\infty(\partial_S) - B_q(\partial_S)$; its exact expression is

$$E_q = \sum_{n=q-v}^D \frac{v! \mu_{n+v}}{(n+v)! \mu_v} \partial_S^n. \quad (\text{AIE14})$$

An empty sum is zero. Its image has degree at most $D - q + v \leq v-1$. Both symbols have constant term one, hence both are invertible in this finite polynomial algebra. Their inverses obey

$$B_q(\partial_S)^{-1} - B_\infty(\partial_S)^{-1} = B_q(\partial_S)^{-1} E_q B_\infty(\partial_S)^{-1}, \quad (\text{AIE15})$$

whose image is contained in $\mathbb{C}[S]_{<v}$ and whose rank is at most v . Equation AIE15 follows by multiplying out its right side. The degree statement holds because every factor is a power series in the degree-lowering derivative. Thus the infinite-symbol formula can be related to the literal AKJ inverse by the displayed exact correction; it cannot replace that inverse without this map.

27.4 An explicit right inverse of the original conductor

Assume now $D \geq v$. Define the raising map, on the degree ranges where its image lies in $\mathcal{P}_D$, by

$$J_s \varphi_n = \sqrt{\frac{w_n}{w_{n+1}}} \varphi_{n+1} = \sqrt{\frac{b+n}{n+1}} \varphi_{n+1}. \quad (\text{AIE16})$$

Then $L_s^v J_s^v = I$ on $\mathcal{P}_{D-v}$ and $\|J_s^v\| \leq \max\{1, b^{v/2}\}$. The latter follows from the displayed coefficient, at each raising step, by $(b+n)/(n+1) \leq \max\{1, b\}$.

The analytic function $G(z) = z/\arctan z$, with $G(0) = 1$, satisfies $|G(z)| \leq 2$ on $|z| < 1$. Indeed

$$\frac{\arctan z}{z} = \int_0^1 \frac{dt}{1+t^2 z^2}, \quad \operatorname{Re} \frac{1}{1+u} > \frac{1}{2} \quad (|u| < 1). \quad (\text{AIE17})$$

The real-part inequality follows from $2 \operatorname{Re}(1+\bar{u}) > |1+u|^2$, which reduces to $1 > |u|^2$. It proves that the integral is nonzero with real part greater than $1/2$, including the removable value at zero. Every Taylor coefficient of $G(z)^v$ is therefore bounded by 2^v : apply Cauchy's formula on radius $t < 1$ and let t increase to one for each fixed coefficient.

Define a map between the specified degree spaces by

$$R_{v,s} = G(L_s)^v J_s^v : \mathcal{P}_{D-v} \longrightarrow \mathcal{P}_D. \quad (\text{AIE18})$$

Since all functions of L_s commute, AIE8 and the power-series identity $(\arctan z)^v G(z)^v = z^v$ give $\partial_y^v R_{v,s} = L_s^v J_s^v = I$. This proves the right-inverse identity even at the top allowed degree. The same row and column argument as for AIE10, now with every coefficient bounded by 2^v , gives

$$\|R_{v,s}\| \leq 2^v (D+1) \alpha_{D,s} \max\{1, b^{v/2}\}. \quad (\text{AIE19})$$

When $v = 0$ the map is exactly the identity; AIE19 remains a valid upper bound. Define the corresponding map on polynomials in the original S coordinate by

$$\mathcal{R}_{v,s,c} = \mathcal{C}_c^{-1} i^v R_{v,s} \mathcal{C}_c. \quad (\text{AIE20})$$

The chain rule in AIE9 proves $\partial_S^v \mathcal{R}_{v,s,c} = I$, with exactly the factor $(-i)^v i^v = 1$. This is a map in the original polynomial algebra; the source norm only specifies its coefficients through the explicitly given original orthonormal basis.

Let $c_k = k/2$ and $c' = c_{k-8} = k/2 - 4$. The full conductor of AKJ13 has the following right inverse on $\mathbb{C}[S']_{\leq D-v}$:

$$\mathcal{R}_{A,s}g = \frac{v!}{\mu_v} B_q(\partial_S)^{-1} \mathcal{R}_{v,s,c'}g, \quad \mathcal{T}_A \mathcal{R}_{A,s}g = g. \quad (\text{AIE21})$$

To check it, use the exact finite factorization $\mathcal{T}_A = (\mu_v/v!) \partial_S^v B_q(\partial_S)$ from AKJ14 on degree at most D . The two inverse factors cancel and AIE20 gives the stated identity. In the identity the output variable is S' ; the differential expression uses the same numerical polynomial coefficients, so the domain and codomain are as displayed.

Set $H_D = \sum_{j=1}^D 1/j$, $H_0 = 0$, and

$$C_{A,D,s} = \frac{v!}{|\mu_v|} e^{4(H_D+2)} N_{D,s} 2^v (D+1) \alpha_{D,s} \max\{1, b^{v/2}\}. \quad (\text{AIE22})$$

Then the original-centre norm estimate is

$$\|\mathcal{R}_{A,s}g\|_{s,c_k} \leq C_{A,D,s} \|g\|_{s,c'}. \quad (\text{AIE23})$$

Indeed AIE11 and AIE19 first prove the bound without the exponential factor, with both norms at c' . For any output polynomial p , $p(c_k + iy) = p(c' + i(y - 4i))$, so AKJ22 applied to the exact shift $-4i$ supplies the remaining factor. Thus the four-unit centre difference is retained in the typed norm map. For the fixed actual quartet and period, $v \leq 80$ and $b \in \{1/2, k/2\}$ give

$$\log C_{A,D,s} \leq D \log(\coth R_0) + O_{H,\delta,\gamma}(\log(D+1) + \log(k+1)), \quad \frac{\log C_{A,D,s}}{kq} \longrightarrow 0. \quad (\text{AIE24})$$

Every factor in the first assertion is written in AIE22; the asymptotic statement follows from $D \leq (k+1)^2 - 1$ and the fixed finite v . No varying-period bound is inferred from this statement.

For comparison with the exact monomial lift $\mathcal{J}_v$ of AKJ17, put $L_g = \mathcal{R}_{v,s,c'}g - \mathcal{J}_v g$. Both terms have v th derivative g , so $L_g \in \mathbb{C}[S]_{<v}$. More explicitly,

$$L_g = \sum_{a=0}^{v-1} [S^a](\mathcal{R}_{v,s,c'}g) S^a, \quad \mathcal{R}_{A,s}g - \frac{v!}{\mu_v} B_q(\partial_S)^{-1} \mathcal{J}_v g = \frac{v!}{\mu_v} B_q(\partial_S)^{-1} L_g. \quad (\text{AIE25})$$

The first identity uses the exact zero coefficients of $\mathcal{J}_v g$ in degrees below v . This proves the complete degree-below- v correction between the two lifts, rather than discarding their different choices of integration constants.

27.5 The exact quotient and exterior transfer supplied by this bound

The kernel of $\mathcal{T}_A$ on degree at most $D \geq v$ is exactly $\mathbb{C}[S]_{<v}$ by AKJ14. Give the quotient $\mathbb{C}[S]_{\leq D}/\mathbb{C}[S]_{<v}$ its Hilbert quotient norm from the original source at c_k , and give its target $\mathbb{C}[S']_{\leq D-v}$ the original source at c' . The induced conductor map $\overline{\mathcal{T}}_A$ is an isomorphism, because AIE21 proves surjectivity and the displayed kernel proves injectivity. Define the full upper norm from AKJ25 by

$$N_{A,D} = \sum_{r,s} |a_{rs}| e^{|b_{rs}-4|(H_D+2)}. \quad (\text{AIE26})$$

For every target g , its minimum-norm preimage class obeys

$$\frac{\|g\|_{s,c'}}{N_{A,D}} \leq \|\overline{\mathcal{T}}_A^{-1} g\|_{\text{quot}} \leq C_{A,D,s} \|g\|_{s,c'}. \quad (\text{AIE27})$$

For the left inequality, apply AKJ25 to every preimage and take the infimum. For the right inequality, use the actual preimage AIE21. The infimum is attained since the omitted subspace is a closed finite-dimensional subspace. These arguments prove the norm statement with the specified quotient map, rather than assigning the coefficient norm to the quotient.

In particular AIE21 gives the exact decomposition

$$V_A = \mathbb{C}[S]_{<v} \oplus \mathcal{R}_{A,s}(\chi_{k-8}\mathbb{C}[S']_{\leq D-v-q'}), \quad q' = (k-7)^2, \quad (\text{AIE28})$$

where a negative degree means the zero space and V_A here denotes the AKJ15 space restricted to degree at most D . Indeed the image under $\mathcal{T}_A$ of a member of V_A is precisely a multiple of χ_{k-8} in the displayed degree range. Subtracting its lift leaves the exact kernel. The directness follows by applying $\mathcal{T}_A$. Equation AIE25 identifies this decomposition with the earlier monic basis, including its low-degree correction.

For any t -dimensional target subspace and any basis column matrix X in its original orthonormal coordinates, AIE27 implies the full Gram determinant inequalities

$$N_{A,D}^{-2t} \det(X^* X) \leq \det(X^* (\overline{\mathcal{T}}_A^{-1})^* \overline{\mathcal{T}}_A^{-1} X) \leq C_{A,D,s}^{2t} \det(X^* X). \quad (\text{AIE29})$$

For proof, AIE27 is a two-sided positive Hermitian matrix inequality after congruence by X . Conjugating by $(X^* X)^{-1/2}$ gives eigenvalues between the two scalar squared bounds, whose product is the displayed determinant ratio. This proves the exact exterior transfer and all powers of the norm constants.

Similarly, for any t -plane $U \subset \mathbb{C}[S]_{\leq D}$, AIE11 gives $\|\bigwedge^t(B_q(\partial_S)^{-1}|_U)\| \leq N_{D,s}^t$. Although $B_q(\partial_S)$ is triangular with every diagonal entry one and hence determinant one on the full polynomial space, that full-space determinant does not cancel its restriction to an arbitrary subspace. The exact restricted determinant is $\det(X^*(B_q^{-1})^* B_q^{-1} X) / \det(X^* X)$ for a basis matrix X of that subspace; this is the map to which AIE11 applies.

For the original kernel rank $r = 8k - 16$ and $D \asymp q$, the upper logarithm in these exterior bounds is $O_{H,\delta,\gamma}(rq) = O_{H,\delta,\gamma}(kq)$. Therefore AIE12 and AIE24 prove a sub- kq logarithmic bound for each operator, while AIE29 retains the rank factor required for volumes. They provide the original conductor inverse, its full low-degree comparison and its quotient metric control. A sub- kq error for the actual rank- r four-endpoint determinant is not a consequence of these scalar bounds; that remaining calculation must use the actual restricted matrices and the existing endpoint correlation.

28 Subleading exterior control from the actual finite conductor

Retain the original conductor of AKJ12–25 and AIE1–29. In particular the quartet, full unit, original period and branch are fixed in the proved five-orbit domain, $m = 1$, $k = 4l + 1 \geq 9$, and $q = (k + 1)^2$. All 81 coefficients a_{rs} are those of the actual product of four period quadrics. They are fixed as k varies. Retain

$$b_{rs} = 4 + (2r - 8)\delta + i(2s - 8)\gamma, \quad \mu_j = \sum_{r,s=0}^8 a_{rs} b_{rs}^j, \quad v = \min\{j : \mu_j \neq 0\} \leq 80, \quad (\text{RUI1})$$

$$A_{\text{abs}} = \sum_{r,s} |a_{rs}|, \quad B_{\text{sh}} = \max_{r,s} |b_{rs}|, \quad C_v = \frac{A_{\text{abs}} B_{\text{sh}}^v}{|\mu_v|} \geq 1.$$

No coefficient or moment denominator is replaced by a generic value. The finite bound for v is AKJ13's distinct-node Vandermonde proof. The inequality $C_v \geq 1$ follows from the triangle inequality applied to the actual v th moment. The source measure at order $s \in \{1, k\}$ is exactly

$$dm_{0,s}(y) = (2\pi)^{s/2} \frac{2^{s/2}}{4\pi\Gamma(s/2)} |\Gamma(s/4 + iy/2)|^2 dy, \quad \|p\|_{s,c} = \|p(c + iy)\|_{L^2(m_{0,s})}. \quad (\text{RUI2})$$

Every centre appearing below is stated, and the whole mass remains in the source norm. Put $H_D = \sum_{j=1}^D 1/j$, $H_0 = 0$.

28.1 The forward polynomial bound and exact determinant

For an integer cutoff $L \geq v + 1$, define the literal finite symbol

$$B_L(z) = 1 + \sum_{a=1}^{L-v-1} \frac{v! \mu_{v+a}}{(v+a)! \mu_v} z^a. \quad (\text{RUI3})$$

The empty sum is zero. This includes the actual coefficient symbol B_q of AKJ14 and the source symbol B_{N+1} needed at a larger source cutoff N . For every displayed coefficient, including the constant coefficient, one has

$$\left| \frac{v! \mu_{v+a}}{(v+a)! \mu_v} \right| \leq C_v \frac{v! B_{\text{sh}}^a}{(v+a)!} \leq C_v \frac{B_{\text{sh}}^a}{a!}. \quad (\text{RUI4})$$

The first inequality is the full moment bound $|\mu_{v+a}| \leq A_{\text{abs}} B_{\text{sh}}^{v+a}$. The second follows from $(v+a)! \geq v! a!$.

On $\mathbb{C}[S]_{\leq D}$ with the norm RUI2, the original coordinate pullback sends ∂_S to $-i\partial_y$. AKJ21–23 proves $\|\partial_S\| \leq H_D + 2$, uniformly in both original source orders. Applying RUI4 term by term to the finite operator polynomial gives

$$\|B_L(\partial_S)\|_{s,c} \leq M_D, \quad M_D = C_v \exp(B_{\text{sh}}(H_D + 2)) \geq 1. \quad (\text{RUI5})$$

Indeed each power has norm at most $(H_D + 2)^a$; extending this nonnegative finite sum to the exponential series proves the displayed bound. The constant term is one and every other term strictly lowers degree. Thus, on this exact $(D + 1)$ -dimensional polynomial space,

$$\det B_L(\partial_S) = 1. \quad (\text{RUI6})$$

This is its determinant in the original increasing-power basis, and therefore in every basis of the same source space, including the original orthonormal Gamma basis. The assertion preserves the source measure; it is not a rescaling of this determinant.

28.2 Complementary singular values retain the full exterior norm

Let $n = D + 1$ and $T = B_L(\partial_S)$ acting on this same Hilbert space. Write its singular values as $\sigma_1 \geq \dots \geq \sigma_n > 0$. RUI6 gives $\prod_i \sigma_i = 1$. For every $0 \leq t \leq n$,

$$\begin{aligned} \left\| \bigwedge^t T^{-1} \right\| &= \prod_{j=n-t+1}^n \sigma_j^{-1} = \prod_{j=1}^{n-t} \sigma_j = \left\| \bigwedge^{n-t} T \right\| \leq M_D^{n-t}, \\ \left\| \bigwedge^t T \right\| &\leq M_D^t. \end{aligned} \tag{RUI7}$$

For a proof of the exterior norm formula, the singular-value decomposition expresses T as two unitary maps and a diagonal map with these positive entries. Exterior powers of the unitary maps preserve the induced norm; the diagonal exterior map has entries equal to the products over all t -subsets. Its largest entry is the product of the largest t singular values. Applying this to T^{-1} and using the full determinant proves every equality in RUI7. Empty products cover $t = 0, n$.

Consequently, for any actual t -dimensional subspace and any full-rank basis column matrix X in the original orthonormal source coordinates,

$$M_D^{-2t} \det(X^* X) \leq \det(X^* (T^{-1})^* T^{-1} X) \leq M_D^{2(n-t)} \det(X^* X). \tag{RUI8}$$

For the upper inequality apply the first exterior norm of RUI7 to the decomposable exterior vector of the columns of X . Its squared norm is their Gram determinant. For the lower inequality, put $Y = T^{-1}X$ and apply the second bound of RUI7 to $TY = X$. This proves the lower bound on this same subspace without requiring that it be invariant. Every original off-diagonal Gram entry is retained by the displayed matrices.

In particular, for fixed original period and quartet, uniformly in $s \in \{1, k\}$, every $0 \leq D \leq 2q$ and every allowed t and X ,

$$\left| \log \frac{\det(X^* (T^{-1})^* T^{-1} X)}{\det(X^* X)} \right| \leq 2(D+1)(\log C_v + B_{\text{sh}}(H_D + 2)) = O_{\text{actual}}(q \log(q+1)) = o(kq). \tag{RUI9}$$

The harmonic estimate $H_D \leq 1 + \log D$ for $D \geq 1$ follows by integrating $1/x$ on consecutive intervals; $D = 0$ is immediate. Since $q = (k+1)^2$, the displayed upper bound divided by kq tends to zero, independently of D, t, X . This is a uniform estimate over the actual subspaces, not a presumed favourable position of the kernel. Its constants are fixed only after fixing the original period and quartet; no uniform bound on $1/|\mu_v|$ over moving periods is asserted. The same inequalities for T follow by exchanging the image subspace and its inverse image.

RUI9 improves the rank-multiplied estimate at the end of AIE29. The earlier estimate is valid, but its factor t on a single inverse operator bound is unnecessary. The complementary exterior identity uses the exact full determinant and supplies the sharper bound on the same original matrices.

28.3 The actual conductor quotient inherits the sharper volume control

Let $D \geq v$ and use the literal source truncation $L = D + 1$. Taylor's finite formula gives, on degree at most D ,

$$\mathcal{T}_A = \sum_{j=v}^D \frac{\mu_j}{j!} \partial_S^j = \frac{\mu_v}{v!} \partial_S^v B_{D+1}(\partial_S), \quad \ker \mathcal{T}_A = \mathbb{C}[S]_{<v}. \tag{RUI10}$$

The first equality is exactly the translation operator in AKJ13; all lower moments vanish by their proved definition. The factor $B_{D+1}(\partial_S)$ is invertible and preserves each degree filtration, so the last equality follows from the derivative kernel.

Use the original source centre $c_k = k/2$ and target centre $c' = c_{k-8} = k/2 - 4$, both with source order s . AIE16–20 gives the full right inverse $\mathcal{R}_{v,s,c'}$ of ∂_S^v with norm at most

$$P_{D,s} = 2^v(D+1)\alpha_{D,s} \max\{1, (s/2)^{v/2}\}, \quad \alpha_{D,s} = \max_{0 \leq m \leq n \leq D} \sqrt{\frac{n!/(s/2)_n}{m!/(s/2)_m}}. \quad (\text{RUI11})$$

In particular $\alpha_{D,k} = 1$, while $\alpha_{D,1} \leq \sqrt{2} \max\{1, D^{1/4}\}$; the full derivation and the phase i^v are the original AIE16–20 construction. The exact conductor right inverse is

$$\mathcal{R}_{A,s,D} = \frac{v!}{\mu_v} B_{D+1}(\partial_S)^{-1} \mathcal{R}_{v,s,c'}. \quad (\text{RUI12})$$

Multiplying by RUI10 proves $\mathcal{T}_A \mathcal{R}_{A,s,D} = I$. The numerical polynomial coefficients use the original source and target variables, as in AIE21. The centre change from c' to c_k is the original shift $y \mapsto y - 4i$, of norm at most $\exp(4(H_D + 2))$ by AKJ22.

Give the quotient $\mathbb{C}[S]_{\leq D}/\mathbb{C}[S]_{<v}$ its minimum norm from the source at c_k . The induced conductor $\overline{\mathcal{T}}_A$ is an isomorphism onto $\mathbb{C}[S']_{\leq D-v}$ with its source norm at c' . Define

$$L_{D,s} = \frac{v!}{|\mu_v|} e^{4(H_D+2)} P_{D,s}, \quad N_{A,D} = \sum_{r,s} |a_{rs}| e^{|b_{rs}-4|(H_D+2)}. \quad (\text{RUI13})$$

For any t -dimensional target subspace, with original basis matrix X , its exact quotient Gram ratio satisfies

$$N_{A,D}^{-2t} \det(X^* X) \leq \det(X^* (\overline{\mathcal{T}}_A^{-1})^* \overline{\mathcal{T}}_A^{-1} X) \leq L_{D,s}^{2t} M_D^{2(D+1-t)} \det(X^* X). \quad (\text{RUI14})$$

To prove the upper bound, use the actual lift RUI12. Apply $\mathcal{R}_{v,s,c'}$ to the t basis columns, retaining its specified image subspace. RUI7 bounds the exterior power of the inverse B_{D+1} on that image by M_D^{D+1-t} . The scalar, right-inverse norm, and centre translation give $L_{D,s}^t$. Projection to the minimum lift in the quotient is an orthogonal projection on the same source Hilbert space, so its exterior norm is at most one. This proves the complete upper Gram inequality. For the lower bound, apply the forward original conductor to the minimum lifts. AKJ24–25 gives its norm $N_{A,D}$; taking its t th exterior power proves the result. This proof retains the coupled Gram and the full quotient metric.

Every factor in RUI13 is explicit in the fixed original data. For the asymptotic assertion restrict now to $v \leq D \leq 2q$. Since $v \leq 80$, $s \in \{1, k\}$ and $t \leq D+1-v \leq 2q+1$, both logarithmic endpoints of RUI14 have absolute value at most

$$2t(|\log L_{D,s}| + |\log N_{A,D}|) + 2(D+1) \log M_D = O_{\text{actual}}(q \log(q+1) + q \log(k+1)) = o(kq). \quad (\text{RUI15})$$

For the logarithm of $N_{A,D}$, use $A_{\text{abs}} \leq N_{A,D} \leq A_{\text{abs}} \exp(\max |b_{rs} - 4|(H_D + 2))$. For $L_{D,s}$, its displayed factors and the positive lower bound $P_{D,s} \geq 1$ give the same logarithmic estimate, retaining $v!/|\mu_v|$. This proves the absolute bounds even if either constant is below one. The harmonic bound and $q = (k+1)^2$ then prove the final limit. No dimension- q scalar comparison has silently acquired an extra factor of q in this argument.

At the original coefficient cutoff $D = q - 1$, RUI14 is the sharpened version of AIE29, on precisely the same conductor quotient and original coefficient subspaces. At a larger source cutoff $D = N$, its literal symbol is B_{N+1} , and RUI10 proves the exact extension. The old symbol B_q cannot be substituted without recording the finite difference. Passing this estimate through the original remainder square requires the actual image of the relation ideal; that square, its image graph, and the endpoint quotient metrics are retained in the succeeding calculation.

29 The original conductor at every source endpoint

Retain the complete simple-quartet conductor of OCF and AKJ, its actual original period matrix, and $k = 4l + 1 \geq 9$ on the proved five-orbit locus. Write

$$\begin{aligned}
 q &= (k+1)^2, & k' &= k-8, & q' &= (k-7)^2, & \Delta &= q - q' = 16k - 48, \\
 c &= k/2, & c' &= k/2 - 4, & \beta_{ij} &= c + (2i - k)\delta + i(2j - k)\gamma, \\
 & & \beta'_{ab} &= c' + (2a - k')\delta + i(2b - k')\gamma, \\
 A &= \sum_{r,s=0}^8 a_{rs} e_{rs}^{(8)} \neq 0, & b_{rs} &= 4 + (2r - 8)\delta + i(2s - 8)\gamma, \\
 \mu_j &= \sum_{r,s} a_{rs} b_{rs}^j, & v &= \min\{j : \mu_j \neq 0\} \leq 80.
 \end{aligned} \tag{CEP1}$$

The factor i before an imaginary term is the imaginary unit; subscript indices are integers in their displayed ranges. The original polynomials are $\chi_k(S) = \prod_{i,j=0}^k (S - \beta_{ij})$ and $\chi_{k'}(S') = \prod_{a,b=0}^{k'} (S' - \beta'_{ab})$. Both have distinct roots. Set $E_k = \mathbb{C}[S]/(\chi_k)$ and $E_{k'} = \mathbb{C}[S']/(\chi_{k'})$. Every source below has the original order $s \in \{1, k\}$ and full mass $M_s = (2\pi)^{s/2}$, with $b = s/2$ and $h_n = M_s n! (b)_n$. In particular the lower grid does not change source order k to $k - 8$.

29.1 The full source, remainder and primary-coordinate square

For every polynomial define the original finite-translation map

$$(\mathcal{T}_A p)(S') = \sum_{r,s=0}^8 a_{rs} p(S' + b_{rs}). \tag{CEP2}$$

For $0 \leq a, b \leq k'$ and $0 \leq r, s \leq 8$, direct substitution gives $\beta'_{ab} + b_{rs} = \beta_{a+r, b+s}$. All indices on the right lie between zero and k . Consequently

$$(\mathcal{T}_A(\chi_k f))(\beta'_{ab}) = 0 \quad \text{for every polynomial } f. \tag{CEP3}$$

Division by the monic $\chi_{k'}$ and evaluation at its distinct roots then show $\chi_{k'} \mid \mathcal{T}_A(\chi_k f)$. Thus CEP2 induces the specified complex-linear map $\bar{\mathcal{T}}_A : E_k \rightarrow E_{k'}$.

In the original ordered primary coordinates its matrix is

$$C_{(a,b),(i,j)} = \begin{cases} a_{i-a, j-b}, & 0 \leq i-a, j-b \leq 8, \\ 0, & \text{otherwise.} \end{cases} \quad C \in \text{Mat}_{q' \times q}(\mathbb{C}). \tag{CEP4}$$

Evaluation of CEP2 proves this entry formula, including its absence of conjugation or binomial coefficients. Its ordinary transpose $C^\top$ is the exact biform coefficient multiplication by the nonzero polynomial A . That multiplication is injective because the biform ring embeds in a polynomial integral domain. Therefore C has rank q' and is surjective. With the original ordered Vandermonde matrices $V_\beta, V_{\beta'}$, the polynomial coordinate matrix is exactly $V_{\beta'}^{-1} C V_\beta$. The kernel is the AKJ15 space

$$V_A = \{[p] \in E_k : \chi_{k'} \mid \mathcal{T}_A p\}, \quad \dim V_A = \Delta. \tag{CEP5}$$

CEP3 proves independence of the representative, and CEP4 proves the dimension. The original common kernel $K = \ker \Lambda$ is contained in V_A by the full conductor inclusion OCF17.

For every endpoint $N \in \{q-1, q, 2q-1, 2q\}$, put $d = N - v$ and define

$$\begin{aligned}
 \mathcal{P}_N &= \mathbb{C}[S]_{\leq N}, & \mathcal{P}'_d &= \mathbb{C}[S']_{\leq d}, \\
 U_N : \mathcal{P}_N &\rightarrow E_k, & p &\mapsto [p], & V_d : \mathcal{P}'_d &\rightarrow E_{k'}, & g &\mapsto [g].
 \end{aligned}$$

The top horizontal map is precisely $\mathcal{T}_{A,N} : \mathcal{P}_N \longrightarrow \mathcal{P}'_{N-v}$. Taylor's finite identity and the first nonzero moment show that CEP2 lowers every degree at least v by exactly v and kills exactly $\mathbb{C}[S]_{<v}$. The exact commutative square is

$$\begin{array}{ccc} \mathcal{P}_N & \xrightarrow{\mathcal{T}_{A,N}} & \mathcal{P}'_{N-v} \\ U_N \downarrow & & \downarrow V_{N-v} \\ E_k & \xrightarrow{\bar{\mathcal{T}}_A} & E_{k'}. \end{array} \quad (\text{CEP6})$$

The lower remainder is surjective because $N - v \geq q - 1 - v \geq q' - 1$; the latter inequality is $\Delta \geq v$, and $\Delta \geq 96 > 80 \geq v$.

The map is complex-linear with its following full action defect. If $M_S, M_{S'}$ denote the original multiplication operators, then

$$CM_S - M_{S'}C = C_b, \quad (C_b)_{(a,b),(i,j)} = \begin{cases} a_{i-a,j-b}b_{i-a,j-b}, & 0 \leq i-a, j-b \leq 8, \\ 0, & \text{otherwise.} \end{cases} \quad (\text{CEP7})$$

Indeed each entry is multiplied by $\beta_{ij} - \beta'_{ab} = b_{i-a,j-b}$. In the original centred coordinates $Y = (M_S - cI)/i$, $Y' = (M_{S'} - c'I)/i$, this becomes $CY - Y'C = (C_b - 4C)/i$. Thus the exact relation to multiplication retains its coefficient defect.

29.2 The literal finite inverse at the larger cutoffs

On $\mathcal{P}_N$, define the literal finite symbol

$$B_{N+1}(z) = 1 + \sum_{j=v+1}^N \frac{v! \mu_j}{j! \mu_v} z^{j-v}, \quad \mathcal{T}_{A,N} = \frac{\mu_v}{v!} \partial_S^v B_{N+1}(\partial_S). \quad (\text{CEP8})$$

The expression on the right is evaluated in the output variable S' . Taylor expansion of each term in CEP2 through degree N proves this identity coefficient by coefficient. Its constant term one makes $B_{N+1}(\partial_S)$ invertible: its difference from the identity lowers degree, so the inverse is its finite nilpotent Neumann sum through power N . A right inverse of ∂_S^v on the polynomial spaces exists by integration of each monomial; composition proves surjectivity of the top map in CEP6.

The relation to the earlier literal B_q of AKJ14 is

$$\begin{aligned} D_{N,q} &= B_{N+1}(\partial_S) - B_q(\partial_S) = \sum_{j=q}^N \frac{v! \mu_j}{j! \mu_v} \partial_S^{j-v}, \\ B_{N+1}(\partial_S)^{-1} - B_q(\partial_S)^{-1} &= -B_{N+1}(\partial_S)^{-1} D_{N,q} B_q(\partial_S)^{-1}, \\ \mathcal{T}_{A,N} - \frac{\mu_v}{v!} \partial_S^v B_q(\partial_S) &= \sum_{j=q}^N \frac{\mu_j}{j!} \partial_S^j. \end{aligned} \quad (\text{CEP9})$$

Empty sums are zero. The first two differences have image contained in $\mathbb{C}[S]_{\leq N-q+v}$; the last has image contained in $\mathbb{C}[S']_{\leq N-q}$. The inverse formula follows by multiplication, and its degree assertion follows since both inverse factors preserve every degree prefix. At $N = q - 1$ all differences vanish. At the other three endpoints these are the full retained defects; their ranks are not assigned the cutoff-independent bound v .

For comparison, the entire symbol $B_\infty(z) = v! \sum_{r,s} a_{rs} e^{b_{rs}z} / (\mu_v z^v)$, with removable value one at zero, obeys at each cutoff

$$B_\infty(\partial_S) - B_{N+1}(\partial_S) = \sum_{n=N+1-v}^N \frac{v! \mu_{n+v}}{(n+v)! \mu_v} \partial_S^n. \quad (\text{CEP10})$$

Its image and the image of the corresponding inverse difference lie in $\mathbb{C}[S]_{<v}$. This follows from the first derivative degree $N + 1 - v$, and from the inverse identity with the same order as AIE15. Thus the rank-at-most- v comparison is with the correctly growing finite symbol in CEP8.

For completeness the original-source inverse estimate extends directly to these literal symbols. Retain the exact constants $A_*^{\text{abs}}, B_*^{\text{sh}}, C_v, R_0, \rho_0$ of AIE4. For every $|z| \leq R_0$ the finite coefficient sum in CEP8 is at most the full absolute moment sum:

$$|B_{N+1}(z) - 1| \leq \frac{v!}{|\mu_v|} \sum_{j=v+1}^N \frac{|\mu_j|}{j!} |z|^{j-v} \leq C_v \frac{B_*^{\text{sh}} |z|}{v+1} e^{B_*^{\text{sh}} |z|} \leq \frac{1}{2}. \quad (\text{CEP11})$$

The middle inequality follows by writing $j = v+1+h$ and using $v!/(v+1+h)! \leq 1/((v+1)h!)$. Hence $F_{N+1}(z) = 1/B_{N+1}(-i \arctan z)$ has coefficient absolute values at most $2\rho_0^{-a}$ on degree $a \geq 1$, and constant coefficient one. This is Cauchy's formula on $|z| = \rho_0$, since $|\arctan z| \leq \operatorname{arctanh} \rho_0 = R_0$.

In the original orthonormal Gamma basis, the lowering matrix $L_s \varphi_n = \sqrt{w_n/w_{n-1}} \varphi_{n-1}$, $w_n = n!/(b)_n$, satisfies $\partial_y = \arctan L_s$ and $L_s^{N+1} = 0$. Evaluation of the power-series inverse in this nilpotent matrix therefore gives the exact finite operator $F_{N+1}(L_s)$, with entries $[z^{n-m}]F_{N+1}(z) \sqrt{w_n/w_m}$. The chain rule $S = c + iy$ supplies the retained $-i$. Bounding the absolute row and column sums gives

$$\|B_{N+1}(\partial_S)^{-1}\|_{s,c} \leq \alpha_{N,s} \left(1 + 2 \sum_{a=1}^N \rho_0^{-a}\right), \quad \alpha_{N,s} = \max_{0 \leq m \leq n \leq N} \sqrt{w_n/w_m}. \quad (\text{CEP12})$$

This proves the larger-cutoff assertion for the actual finite symbol rather than substituting N into an identity for B_q . Likewise the explicit AIE18 raising right inverse on these degree spaces, followed by i^v , $B_{N+1}(\partial_S)^{-1}$ and $v!/\mu_v$, is a right inverse of CEP8. Its comparison with the monomial integral remains exactly the degree-below- v correction in AIE25, now with this literal inverse.

29.3 The exact source operator matrix and its full determinant

Use the unchanged original orthonormal polynomials $u_n(S) = p_n((S - c)/i)/\sqrt{h_n}$ in $\mathcal{P}_N$ and $u'_m(S') = p_m((S' - c')/i)/\sqrt{h_m}$ in $\mathcal{P}'_{N-v}$. Let A_N denote the matrix of $\mathcal{T}_{A,N}$ in these two frames. Define the analytic coefficient series at zero

$$t(z) = \sum_{r,s} a_{rs} \exp\left(\frac{b_{rs} - 4}{i} \arctan z\right) = \sum_{\ell \geq 0} t_\ell z^\ell. \quad (\text{CEP13})$$

Then its exact entries are

$$(A_N)_{m,n} = \begin{cases} t_{n-m} \sqrt{w_n/w_m}, & n \geq m, \\ 0, & n < m, \end{cases} \quad 0 \leq m \leq N - v, \quad 0 \leq n \leq N. \quad (\text{CEP14})$$

To prove this, the original generating identity $\sum p_n(y) z^n/n! = (1 + z^2)^{-b/2} e^{y \arctan z}$ shows that translation by η multiplies the series by $e^{\eta \arctan z}$. Extracting its finite coefficients and dividing by the complete $\sqrt{h_n}$ gives $\sqrt{w_n/w_m}$ after the two copies of M_s cancel at this specified step. Evaluation of CEP2 at $c' + iy$ uses $p(c + i(y + (b_{rs} - 4)/i))$, proving the exact centre shift in CEP13.

For $\ell < v$, the shifted moment $\sum a_{rs} (b_{rs} - 4)^\ell$ is zero by the binomial theorem and the definition of v ; at $\ell = v$ it equals μ_v . Since $\arctan z = z + O(z^3)$, this gives

$$t_0 = \dots = t_{v-1} = 0, \quad t_v = \frac{(-i)^v \mu_v}{v!} \neq 0. \quad (\text{CEP15})$$

The first v columns of A_N vanish. Its remaining square block $\hat{A}_N$, with columns $v, \dots, N$, is upper triangular with diagonal $t_v \sqrt{w_{m+v}/w_m}$. Thus its rank is $N - v + 1$, its kernel in the source is

exactly the first v orthonormal coordinates, and

$$\begin{aligned}\det \hat{A}_N &= \left(\frac{(-i)^v \mu_v}{v!} \right)^{N-v+1} \prod_{m=0}^{N-v} \sqrt{\frac{w_{m+v}}{w_m}}, \\ \det(A_N A_N^*) &= \left| \frac{\mu_v}{v!} \right|^{2(N-v+1)} \prod_{m=0}^{N-v} \frac{w_{m+v}}{w_m}.\end{aligned}\tag{CEP16}$$

This retains the complex determinant phase and the original moment. The operator norm also has the exact original-source upper bound

$$\|A_N\| \leq M_{A,N} := \sum_{r,s} |a_{rs}| e^{|b_{rs}-4|(H_N+2)}, \quad H_N = \sum_{j=1}^N 1/j.\tag{CEP17}$$

Indeed AKJ22 bounds each exact translation in CEP13 by its displayed factor, and the triangle inequality gives the sum.

There is a closed four-endpoint expression for CEP16. For the signs $\sigma = (1, 1, -1, -1)$ at the ordered endpoints, write $\mathcal{D}_A = \sum_N \sigma_N \log \det(A_N A_N^*)$. Cancellation of the full products gives

$$\begin{aligned}\mathcal{D}_A &= -4q \log \left| \frac{\mu_v}{v!} \right| + \mathcal{W}_{v,s}, \\ \mathcal{W}_{v,s} &= \sum_{a=0}^{v-1} (\log w_{q-1-a} + \log w_{q-a} - \log w_{2q-1-a} - \log w_{2q-a}).\end{aligned}\tag{CEP18}$$

For $v = 0$ the second sum is zero. The product identity used here is $\prod_{m=0}^{N-v} w_{m+v}/w_m = \prod_{a=0}^{v-1} w_{N-a}/\prod_{a=0}^{v-1} w_a$, obtained by cancellation of consecutive products; it remains valid when the two index intervals overlap.

Put $c_b^{\log} = \max\{1, b-1\}$. For every $j \geq 1$ and $b \geq 1/2$,

$$\left| \log \frac{j}{b+j-1} \right| \leq \frac{c_b^{\log}}{j}.\tag{CEP19}$$

For $b \geq 1$ use $\log(1 + (b-1)/j) \leq (b-1)/j$. For $1/2 \leq b < 1$, put $a = 1 - b \leq 1/2$ and use $-\log(1 - a/j) \leq a/(j-a) \leq 1/j$; the last inequality is $aj \leq j-a$, which follows from $a \leq 1/2$ and $j \geq 1$. Summing CEP19 over the two intervals of length q in each summand of CEP18 proves the completely explicit estimate

$$|\mathcal{W}_{v,s}| \leq \frac{2v c_b^{\log} q}{q-v}, \quad |\mathcal{D}_A| \leq 4q \left| \log \left| \frac{\mu_v}{v!} \right| \right| + \frac{2v c_b^{\log} q}{q-v}.\tag{CEP20}$$

All denominators are positive since $q \geq 100 > v$. For either original source and the fixed actual quartet and period, the displayed bound is $O_{\text{actual}}(q+k) = o(kq)$.

29.4 The induced original quotient metric and its correction

Let the matrices of the two remainder maps in the orthonormal source frames and original primary target frames be denoted by the same U_N, V_d as their maps. Explicitly

$$(U_N)_{(i,j),n} = \frac{p_n((\beta_{ij} - c)/i)}{\sqrt{h_n}}, \quad (V_d)_{(a,b),m} = \frac{p_m((\beta'_{ab} - c')/i)}{\sqrt{h_m}}.\tag{CEP21}$$

They have sizes q by $N+1$ and q' by $N-v+1$, respectively. Their first q , respectively q' , columns have nonzero evaluation Vandermonde determinant, so both have full row rank. The square CEP6 is the precise matrix identity

$$CU_N = V_d A_N, \quad R_N = U_N U_N^*, \quad R'_d = V_d V_d^*, \quad G_N = R_N^{-1}, \quad G'_d = (R'_d)^{-1}.\tag{CEP22}$$

Here G_N is the original upper primary quotient metric of OKM8, and G'_d is the lower-grid metric in the same source order s . The induced metric on E_k/V_A , in its specified target coordinates $E_{k'}$, is

$$Q_N = (CG_N^{-1}C^*)^{-1} = (V_d A_N A_N^* V_d^*)^{-1}. \quad (\text{CEP23})$$

To prove the first assertion, the minimum of $x^* G_N x$ under $Cx = y$ is attained at $x = G_N^{-1} C^* (CG_N^{-1} C^*)^{-1} y$. Every other lift differs by an element of $\ker C$ orthogonal to this lift in the G_N metric, proving the minimum and its Gram matrix. The second equality is CEP22.

The exact correction to the lower original Gamma quotient metric is

$$\begin{aligned} W_d &= V_d^* (R'_d)^{-1/2}, \quad W_d^* W_d = I_{q'}, \\ K_{A,N} &= W_d^* A_N A_N^* W_d > 0, \quad Q_N = (R'_d)^{-1/2} K_{A,N}^{-1} (R'_d)^{-1/2}, \\ \log \det Q_N &= \log \det G'_d - \log \det K_{A,N}. \end{aligned} \quad (\text{CEP24})$$

All square roots are the unique positive Hermitian square roots. These formulas follow by multiplying out the definition of $K_{A,N}$ and inverting. In particular the passage between the induced metric and the lower original source metric retains this actual compressed covariance.

If Z_d is any orthonormal basis of $\ker V_d$, completing W_d to a unitary matrix, the further exact determinant factorization is

$$\det K_{A,N} = \det(A_N A_N^*) \det(Z_d^* (A_N A_N^*)^{-1} Z_d). \quad (\text{CEP25})$$

For a proof put $D = A_N A_N^* > 0$ and write its block matrix in the unitary frame $[W_d, Z_d]$. The upper left block is $K_{A,N}$. Block Gaussian elimination gives $\det D = \det K_{A,N} \det(D_{22} - D_{21} K_{A,N}^{-1} D_{12})$. The lower right block of D^{-1} is the inverse of that Schur complement, proving CEP25. The empty-complement case has determinant one. The dimension of this retained complement is $N - v + 1 - q'$.

Choose any fixed full frame $I_V : \mathbb{C}^\Delta \hookrightarrow E_k$ of $\ker C$ and a fixed right section L_C of C . Then $F_V = [I_V, L_C]$ is invertible and

$$\det(I_V^* G_N I_V) = |\det F_V|^2 \frac{\det G_N}{\det Q_N} = |\det F_V|^2 \frac{\det G_N}{\det G'_d} \det K_{A,N}. \quad (\text{CEP26})$$

Indeed $CF_V = [0, I]$, so the lower right block of $(F_V^* G_N F_V)^{-1}$ is $CG_N^{-1} C^* = Q_N^{-1}$. The Schur determinant argument just used gives CEP26. The frame factor is independent of source cutoff and cancels only after applying the four signs.

For the actual inclusion $K \subset V_A$, choose its original coefficient frame, and let $\Xi : \mathbb{C}^\Delta \rightarrow \mathbb{C}^{\Delta-r}$ have its coordinate kernel, $r = 8k - 16$, as in AKJ18. With $H_{V,N} = I_V^* G_N I_V$, the quotient metric is $Q_{\Xi,N} = (\Xi H_{V,N}^{-1} \Xi^*)^{-1}$. The same proof with a frame of $\ker \Xi$ and a right section gives, with all constant frame factors cancelled by the endpoint signs,

$$F_K^{(s)} = \sum_N \sigma_N \log \det G_N - \sum_N \sigma_N \log \det G'_{N-v} + \sum_N \sigma_N \log \det K_{A,N} - \sum_N \sigma_N \log \det Q_{\Xi,N}. \quad (\text{CEP27})$$

Thus the conductor quotient and the remaining original observation quotient have both been retained in the determinant relation. The lower source cutoffs in this equation are exactly $q - 1 - v, q - v, 2q - 1 - v, 2q - v$. Relative to the lower grid's own four cutoffs they have the respective shifts $\Delta - v, \Delta - v, 2\Delta - v, 2\Delta - v$.

29.5 An exact graph quotient with controlled exterior metric error

In original orthonormal coordinates define

$$S_N : \mathcal{P}_N \longrightarrow \mathbb{C}^v \oplus \mathcal{P}'_{N-v}, \quad S_N p = (P_{<v} p, \mathcal{T}_{A,N} p). \quad (\text{CEP28})$$

Here $P_{<v}$ records the first v original upper orthonormal coefficients, and the target carries the orthogonal direct-sum norm with those coordinates and the unchanged lower Gamma source. The first v source coordinates span the polynomials of degree below v . Thus CEP14–16 give its exact block matrix $S_N = \text{diag}(I_v, \hat{A}_N)$ in the ordered source split at degree v . In particular it is invertible, and

$$\begin{aligned} \|S_N\| &\leq M_N := \max\{1, M_{A,N}\}, \\ |\det S_N| &= \left| \frac{\mu_v}{v!} \right|^{N-v+1} \prod_{m=0}^{N-v} \sqrt{\frac{w_{m+v}}{w_m}}, \\ |\log |\det S_N|| &\leq L_N := (N-v+1) \left| \log \left| \frac{\mu_v}{v!} \right| \right| + v c_b^{\log} H_N. \end{aligned} \quad (\text{CEP29})$$

The last bound follows by the consecutive-product identity preceding CEP19 and $|\log w_j| \leq c_b^{\log} H_j$: the two lists of v logarithms are each bounded by $v c_b^{\log} H_N$, followed by the factor $1/2$.

The upper and lower source ideals at this cutoff are

$$I_N = \chi_k \mathbb{C}[S]_{\leq N-q}, \quad I'_N = \chi_{k'} \mathbb{C}[S']_{\leq N-v-q'}. \quad (\text{CEP30})$$

A negative degree specifies the zero subspace. Let $\mathcal{W}_N = U_N^{-1}(V_A)$. The exact image and ideal graph under CEP28 are

$$\begin{aligned} S_N \mathcal{W}_N &= \mathbb{C}^v \oplus I'_N, \\ S_N I_N &= \{(P_{<v} h, \mathcal{T}_{A,N} h) : h \in I_N\} \subset \mathbb{C}^v \oplus I'_N, \\ V_A &\xrightarrow{\simeq} (\mathbb{C}^v \oplus I'_N) / (S_N I_N), \quad U_N p \mapsto [S_N p]. \end{aligned} \quad (\text{CEP31})$$

The first equality follows because $\mathcal{W}_N$ is the preimage of I'_N under the top map of CEP6, and S_N is invertible. CEP3 proves the ideal inclusion. If representatives differ by I_N , their images differ by $S_N I_N$, so the last map is well defined. Conversely every class in the target has a preimage under S_N in $\mathcal{W}_N$, and the kernel consists exactly of I_N . This proves the isomorphism. Its norm on the left is the restriction of G_N : all lifts of a member of V_A already belong to $\mathcal{W}_N$, so their minimum source norm is the quotient norm of $\mathcal{W}_N / I_N$. The low-degree graph component $P_{<v} h$ is explicitly retained.

For reference the ideal dimensions at the four original endpoints are

N	$\dim I_N$	$\dim I'_N$	$\dim((\mathbb{C}^v \oplus I'_N) / S_N I_N)$
$q-1$	0	$\Delta - v$	Δ
q	1	$\Delta + 1 - v$	Δ
$2q-1$	q	$q + \Delta - v$	Δ
$2q$	$q+1$	$q + \Delta + 1 - v$	Δ .

(CEP32)

These follow by counting monomials in CEP30; S_N preserves dimensions. Moreover $\mathcal{T}_A|_{I_N}$ is injective because a nonzero member has degree at least $q > v$ and its image has degree exactly v less. Hence the second coordinate of the retained ideal graph already has dimension $\dim I_N$.

Here is the complete exterior norm estimate for this exact map. For an invertible complex matrix S of dimension n and $0 \leq t \leq n$, its singular values $\sigma_1 \geq \dots \geq \sigma_n > 0$ give

$$\left\| \bigwedge^t S^{-1} \right\| = \frac{\left\| \bigwedge^{n-t} S \right\|}{|\det S|}. \quad (\text{CEP33})$$

Indeed the numerator is $\sigma_1 \cdots \sigma_{n-t}$ and the denominator is the product of all singular values; their ratio is the product of the t largest singular values of S^{-1} . This includes $t = 0, n$. For $n = N+1$, CEP29 therefore gives the finite bounds

$$\begin{aligned} \log^+ \left\| \bigwedge^t S_N \right\| &\leq t \log M_N, \\ \log^+ \left\| \bigwedge^t S_N^{-1} \right\| &\leq (N+1-t) \log M_N + L_N. \end{aligned} \quad (\text{CEP34})$$

Here $\log^+ x = \max\{0, \log x\}$. The right sides are nonnegative. Thus for all these ranks both bounds are at most

$$E_N = (N + 1) \log M_N + L_N. \quad (\text{CEP35})$$

With $R_{\text{sh}} = \max_{r,s} |b_{rs} - 4|$ and $A_*^{\text{abs}} = \sum |a_{rs}|$, equation CEP17 gives $\log M_N \leq \log^+ A_*^{\text{abs}} + R_{\text{sh}}(H_N + 2)$. For the fixed original quartet and period, v, R_{sh}, μ_v and A_*^{abs} are fixed while $c_b^{\log} = O(k)$. Since $N \leq 2q$ and $H_N \leq 1 + \log N$, all four explicit bounds in CEP35 satisfy

$$E_N = O_{\text{actual}}(q \log(q + 1) + k \log(q + 1)) = o(kq). \quad (\text{CEP36})$$

This uses only the forward norm and the exact determinant of the original conductor map.

We now prove the exact relation to the metric of the original kernel, including the quotient projection. In a finite-dimensional Hilbert space let $I \subset W$ and let S be invertible. The induced map $W/I \rightarrow SW/SI$ sends a minimal lift $x \in W \cap I^\perp$ to the orthogonal projection of Sx off SI . Orthogonal projection is a contraction on every exterior power: in an orthonormal basis its singular values are zeros and ones. Therefore, on any t -dimensional subspace of W/I , the exterior norm of the induced map is at most $\|\bigwedge^t S\|$. Applying the identical argument to S^{-1} gives the inverse bound. The minimal-lift identification is isometric since every other lift differs by a perpendicular element of I , whose squared norm adds. This proves the quotient assertion with no choice of coefficient norm.

Apply this result with $W = \mathcal{W}_N$, $I = I_N$, and the actual rank- r subspace $K \subset V_A$. The target subspace is the explicit image of K under CEP31; denote its Gram matrix in the transported original frame by $\tilde{H}_{K,N}$. Let $I_K : \mathbb{C}^r \hookrightarrow E_k$ be that original primary frame, and write its original minimum source lift as $Y_N = U_N^* G_N I_K$. Let Z_N be the matrix in the original orthonormal source coordinates of the independent columns $\chi_k(S) S^a$, $0 \leq a \leq N - q$. If there are no columns, all following products involving them are zero. The exact target orthogonal projection and Gram matrix are

$$\begin{aligned} P_N^{\text{graph}} &= I - S_N Z_N (Z_N^* S_N^* S_N Z_N)^{-1} Z_N^* S_N^*, \\ \tilde{H}_{K,N} &= Y_N^* S_N^* P_N^{\text{graph}} S_N Y_N, \quad H_{K,N} = Y_N^* Y_N = I_K^* G_N I_K. \end{aligned} \quad (\text{CEP37})$$

The inverted Gram matrix is positive definite because its columns are independent and S_N is invertible. Multiplication shows that P_N^{graph} is Hermitian, idempotent, kills $S_N I_N$ and fixes its orthogonal complement. Thus CEP37 is precisely the minimum quotient Gram of CEP31, and its kernel frame has full rank because CEP31 is injective. Replacing the lift Y_N by any lift of the same primary frame adds columns in I_N , which the projection kills; hence every choice of lift gives this same matrix.

Applying the proved exterior quotient estimate and squaring volumes gives

$$\begin{aligned} -2 \log \left\| \bigwedge^r S_N^{-1} \right\| &\leq \log \det \tilde{H}_{K,N} - \log \det H_{K,N} \\ &\leq 2 \log \left\| \bigwedge^r S_N \right\|, \\ \left| \log \det \tilde{H}_{K,N} - \log \det H_{K,N} \right| &\leq 2E_N. \end{aligned} \quad (\text{CEP38})$$

In detail, divide the chosen frame by its positive Gram square root so it is orthonormal in the source quotient. The squared norm of its exterior image is exactly the determinant ratio. The forward and inverse exterior estimates prove the two inequalities, and CEP34–35 prove the last assertion.

Consequently the following reduction of the actual four-endpoint kernel determinant is proved:

$$\begin{aligned} \tilde{F}_K^{(s)} &= \sum_N \sigma_N \log \det \tilde{H}_{K,N}, \\ |F_K^{(s)} - \tilde{F}_K^{(s)}| &\leq 2 \sum_{N \in \{q-1, q, 2q-1, 2q\}} E_N = o(kq). \end{aligned} \quad (\text{CEP39})$$

The frame is the original fixed frame of K at every endpoint; CEP37 gives all four transported matrices explicitly. Their target spaces retain the original ideal graphs, the different source cutoffs, all conductor coefficients, and the full original Gamma masses. CEP39 proves that the leading kq coefficient is unchanged by this particular graph-quotient transport. It supplies no value for that coefficient: its remaining determinant is the displayed, completely specified four-endpoint graph quotient.

29.6 All ideal graph entries as original polynomial products

Each individual shift in CEP2 induces the unital algebra map $\sigma_{rs} : E_k \rightarrow E_{k'}, [p] \mapsto [p(S' + b_{rs})]$. The divisibility in CEP3, applied to the individual shifted polynomial, proves independence of the representative; substitution preserves multiplication and one. Thus $\bar{\mathcal{T}}_A = \sum a_{rs} \sigma_{rs}$ is its exact linear combination. For any $x, y \in E_k$, the complete multiplicative defect of that linear combination is

$$\bar{\mathcal{T}}_A(xy) - \bar{\mathcal{T}}_A(x)\bar{\mathcal{T}}_A(y) = \sum_{r,s} a_{rs} \sigma_{rs}(x) \sigma_{rs}(y) - \sum_{r,s,t,u} a_{rs} a_{tu} \sigma_{rs}(x) \sigma_{tu}(y). \quad (\text{CEP40})$$

This follows by applying the proved individual multiplicativity and expanding the product of the two complete sums.

There is a complete finite product for the map on the source ideal. For each original shift put

$$d_{rs}(S') = \prod_{\substack{0 \leq i,j \leq k \\ (i,j) \notin \{r,\dots,r+k'\} \times \{s,\dots,s+k'\}}} (S' + b_{rs} - \beta_{ij}). \quad (\text{CEP41})$$

This is monic of degree Δ , and $\chi_k(S' + b_{rs}) = \chi_{k'}(S') d_{rs}(S')$. Indeed CEP3's root addition identifies exactly the q' factors in the indicated rectangle with the factors of $\chi_{k'}$. All remaining factors are precisely the product in CEP41. Consequently for every $h \in \mathbb{C}[S]_{\leq L}$, $L = N - q$, the exact ideal map is

$$\begin{aligned} \mathcal{T}_{A,N}(\chi_k h) &= \chi_{k'} \mathcal{U}_{A,L} h, \\ (\mathcal{U}_{A,L} h)(S') &= \sum_{r,s} a_{rs} d_{rs}(S') h(S' + b_{rs}) \in \mathbb{C}[S']_{\leq L+\Delta-v}. \end{aligned} \quad (\text{CEP42})$$

The equality follows by the product identity just proved. The degree statement follows from CEP8: if h has degree t and leading coefficient $h_t \neq 0$, the leading coefficient of its displayed image is $h_t \binom{q+t}{v} \mu_v$ at degree $t + \Delta - v$. This follows because $\chi_k, \chi_{k'}$ are monic and $\mathcal{T}_A$ lowers degree exactly by v with its stated leading derivative. Thus $\mathcal{U}_{A,L}$ is injective. At $L = -1$ the domain and image are zero.

This gives a further exact sequence, with all maps specified:

$$0 \longrightarrow \mathbb{C}[S]_{<v} \longrightarrow V_A \longrightarrow \mathbb{C}[S']_{\leq L+\Delta-v} / \mathcal{U}_{A,L} \mathbb{C}[S]_{\leq L} \longrightarrow 0. \quad (\text{CEP43})$$

The first map is the original quotient class. For the second, lift a member of V_A to $p \in \mathcal{W}_N$, divide $\mathcal{T}_A p$ by $\chi_{k'}$, and take its class in the last space. A change of lift by $\chi_k h$ adds exactly $\mathcal{U}_{A,L} h$ by CEP42, proving independence. Every target polynomial times $\chi_{k'}$ has a preimage under the onto map $\mathcal{T}_{A,N}$, proving surjectivity. If the image is zero, subtract the corresponding $\chi_k h$; the remaining polynomial is killed by $\mathcal{T}_A$ and hence has degree below v . Its class injects because $v < q$. This proves exactness. The last space has dimension $\Delta - v$ at every original endpoint, as is also obtained by subtracting the two polynomial dimensions.

For a literal matrix version of the ideal graph, equip $\mathbb{C}[S']_{\leq L+\Delta-v}$ with the ordered lower orthonormal Gamma coefficient basis, and write $M_{\chi',N}$ for multiplication by the full original $\chi_{k'}$ into $\mathcal{P}'_{N-v}$. It is injective. The map

$$D_{\chi',N} = \text{diag}(I_v, M_{\chi',N}) : \mathbb{C}^v \oplus \mathbb{C}[S']_{\leq L+\Delta-v} \longrightarrow \mathbb{C}^v \oplus I'_N \quad (\text{CEP44})$$

is an isomorphism onto the indicated subspace. Its pullback metric is exactly $G_{\chi',N} = \text{diag}(I_v, M_{\chi',N}^* M_{\chi',N})$; this follows by taking the squared norm of each component. For the domain polynomial coefficient frame of h , let $Z_{\chi,N}h$ be the original upper orthonormal coefficients of $\chi_k h$. In these coordinates the complete graph relation matrix is

$$\mathcal{Z}_{A,N} = \begin{pmatrix} P_{<v} Z_{\chi,N} \\ [\mathcal{U}_{A,L}] \end{pmatrix}, \quad D_{\chi',N} \mathcal{Z}_{A,N} = S_N Z_{\chi,N}. \quad (\text{CEP45})$$

The first component is the original low-degree projection, and the second consists of the exact finite products and translations in CEP41–42. The equality is CEP28 and CEP42. It proves that the graph quotient in CEP31 is precisely the quotient by $\text{im } \mathcal{Z}_{A,N}$ in the displayed metric $G_{\chi',N}$. All matrix inverses and minimum projections in CEP37 can therefore be computed using these full original polynomial products. No coefficient of the ideal, source metric or low-degree graph component is discarded by this additional coordinate isomorphism.

29.7 The compressed conductor correction is subleading at every endpoint

The exact exterior estimate also controls the large compressed covariance in CEP24 without multiplying a scalar inverse bound by q' . Write $n' = N - v + 1$ and retain $A_N = [0, \hat{A}_N]$ with its exact square block in CEP16. Since $W_d^* W_d = I_{q'}$, the coefficient matrix of its image under $\hat{A}_N^*$ gives

$$K_{A,N} = (\hat{A}_N^* W_d)^* (\hat{A}_N^* W_d). \quad (\text{CEP46})$$

The squared norm of the exterior product of the q' columns on the right is its determinant. The norm of the exterior product of the columns of W_d is one, because those columns are orthonormal. The forward estimate therefore gives $\det K_{A,N} \leq \|\hat{A}_N\|^{2q'} \leq M_N^{2q'}$. For the reverse inequality apply $\bigwedge^{q'} \hat{A}_N^*$ to the same image columns; their image is the orthonormal column wedge of W_d . Thus

$$\det K_{A,N} \geq \left\| \bigwedge^{q'} \hat{A}_N^{-1} \right\|^{-2}. \quad (\text{CEP47})$$

The singular values of an adjoint are the same, so the adjoint inverse in this argument has exactly the displayed exterior norm. The singular complement identity CEP33 in dimension n' gives $\|\bigwedge^{q'} \hat{A}_N^{-1}\| \leq M_N^{n'-q'} / |\det \hat{A}_N|$. CEP29 bounds the absolute logarithm of the last determinant by L_N . Consequently the complete two-sided estimate is

$$\begin{aligned} -2((N - v + 1 - q') \log M_N + L_N) &\leq \log \det K_{A,N} \leq 2q' \log M_N, \\ |\log \det K_{A,N}| &\leq 2E_N = o(kq) \quad \text{at every original endpoint and source order.} \end{aligned} \quad (\text{CEP48})$$

All ranks in this argument are their actual ranks, $q' \leq N - v + 1$ as proved after CEP6. In particular the estimate applies to the complete original lower quotient, whose rank grows quadratically.

Define the fully specified total-source difference and the remaining original observation quotient return by

$$\begin{aligned} \mathcal{T}_k^{(s)} &= \sum_N \sigma_N (\log \det G_N - \log \det G'_{N-v}), \\ F_{\Xi,k}^{(s)} &= \sum_N \sigma_N \log \det Q_{\Xi,N}, \\ \mathcal{E}_{A,k}^{(s)} &= \sum_N \sigma_N \log \det K_{A,N}. \end{aligned}$$

Equations CEP26–27 and CEP48 prove the exact identities and bounds

$$\begin{aligned} F_{V_A}^{(s)} &= \mathcal{T}_k^{(s)} + \mathcal{E}_{A,k}^{(s)}, \quad F_K^{(s)} = \mathcal{T}_k^{(s)} - F_{\Xi,k}^{(s)} + \mathcal{E}_{A,k}^{(s)}, \\ |\mathcal{E}_{A,k}^{(s)}| &\leq 2 \sum_N E_N = o(kq), \\ \frac{F_K^{(s)} - \mathcal{T}_k^{(s)} + F_{\Xi,k}^{(s)}}{kq} &\longrightarrow 0. \end{aligned} \quad (\text{CEP49})$$

Here $F_{V_A}^{(s)}$ uses its fixed original coefficient frame, and $Q_{\Xi,N}$ is the exact quotient metric of CEP27. The frame factors cancel with coefficients $1 + 1 - 1 - 1$ only in these returns, as proved in CEP26. Thus the leading metric effect of the full conductor source transport has now been controlled. The remaining rank- $8k - 32$ observation quotient and the shifted lower-grid source determinants are still their explicit original matrices; CEP49 carries them forward into the next calculation.

30 The total-source difference as an outer-strip quotient

Retain CEP1–49, with $m = 1$, $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $k' = k - 8$, $q' = (k - 7)^2$, $\Delta = q - q' = 16k - 48$, and the original fixed $0 < \delta < 1/2$, $\gamma > 2$. The conductor has its actual first nonzero moment index $0 \leq v \leq 80$. At each of the four original endpoints $N \in \{q - 1, q, 2q - 1, 2q\}$ put $t = N - q + 1$ and $d = N - v$. The source order remains $s \in \{1, k\}$ at both grids, with

$$b = s/2, \quad M_s = (2\pi)^{s/2}, \quad dm_s(y) = M_s \frac{2^b}{4\pi\Gamma(b)} |\Gamma(b/2 + iy/2)|^2 dy, \quad h_n = M_s n!(b)_n. \quad (\text{CTV1})$$

These are the complete original measure and monic norms of OKM6–7. The source centres are $c = k/2$ and $c' = k/2 - 4$. The total-source difference in CEP49 is

$$\mathcal{T}_k^{(s)} = \sum_N \sigma_N (\log \det G_N - \log \det G'_{N-v}), \quad \sigma = (1, 1, -1, -1), \quad (\text{CTV2})$$

where both matrices use their original primary frames. All calculations below retain these four cutoffs and the same two source orders.

30.1 The exact nested root polynomials in the original real coordinate

Define the full monic polynomials

$$\begin{aligned} P_k(y) &= i^{-q} \chi_k(c + iy) = \prod_{a,b=0}^k (y - \omega_{ab}), & \omega_{ab} &= (2b - k)\gamma - i(2a - k)\delta, \\ P_{k'}(y) &= i^{-q'} \chi_{k'}(c' + iy) = \prod_{a,b=0}^{k'} (y - \omega'_{ab}), & \omega'_{ab} &= (2b - k')\gamma - i(2a - k')\delta. \end{aligned} \quad (\text{CTV3})$$

The original phases i^{-q} and $i^{-q'}$ have modulus one; they occur exactly as written in the coefficient maps. For every lower pair one has $\omega'_{ab} = \omega_{a+4, b+4}$. Hence, with the complete outer index set $\mathcal{O}_k = \{0, \dots, k\}^2 \setminus \{4, \dots, k - 4\}^2$, the exact factorization is

$$P_k(y) = P_{k'}(y) P_{\partial, k}(y), \quad P_{\partial, k}(y) = \prod_{(a,b) \in \mathcal{O}_k} (y - \omega_{ab}), \quad \deg P_{\partial, k} = \Delta. \quad (\text{CTV4})$$

Direct substitution proves the root equality; counting the retained central rectangle gives q' roots, leaving Δ distinct outer roots. The products are monic, so the root equality fixes the factor without any additional constant.

Since k' is odd, $|2a - k'| \geq 1$ for every lower index. For real y this proves, factor by factor,

$$|y - \omega'_{ab}| \geq |\operatorname{Im} \omega'_{ab}| \geq \delta, \quad |P_{k'}(y)| \geq \delta^{q'}. \quad (\text{CTV5})$$

Also $|\omega'_{ab}| \leq R' = k' \sqrt{\delta^2 + \gamma^2}$. These estimates keep every original factor; no root has been moved.

30.2 The full monic source and relation determinant identities

For a nonzero polynomial P and an integer $a \geq 0$ define the relation moment determinant

$$D_a(P; s) = \det \left[\int_{\mathbb{R}} y^{i+j} |P(y)|^2 dm_s(y) \right]_{0 \leq i, j < a}, \quad D_0(P; s) = 1. \quad (\text{CTV6})$$

Every nonempty matrix is positive definite: the measure is strictly positive on the real line, a nonzero polynomial has only finitely many zeros, and the squared modulus of its product with P is positive on an open interval. All moments exist because the original source has its finite exponential moments in CTV14 below.

In increasing monomial y coordinates, the quotient by a monic polynomial P of degree a from source degree $L \geq a - 1$ has determinant

$$\det G_{P,L}^y = \frac{\prod_{n=0}^L h_n}{D_{L-a+1}(P; s)}. \quad (\text{CTV7})$$

Here the quotient frame is $1, y, \dots, y^{a-1}$. For proof use the source frame

$$(1, y, \dots, y^{a-1}, P, yP, \dots, y^{L-a}P).$$

Its increasing-degree coefficient matrix has diagonal entries one and no entries above the degree of a column, hence determinant one. The last block is exactly the relation Gram in CTV6. Minimizing over those last coordinates gives its Schur complement, whose determinant is the full source determinant divided by the relation determinant. Monic orthogonalization by the original p_n also has determinant one, so that full determinant is $\prod_{n=0}^L h_n$. This proves CTV7 including the empty relation case. It is the full-source analogue of BSL3, with the original Gamma order retained.

Let $V_\beta, V_{\beta'}$ be the original ordered Vandermonde matrices in the S, S' variables. The exact coordinate pullback $S = c + iy$ sends S^j to $(c + iy)^j$ and has diagonal coefficients i^j ; its determinant is $i^{a(a-1)/2}$ in degree below a . Its modulus is one, so its Gram congruence has determinant factor one. The same statement holds with c' . Evaluation in primary coordinates then gives

$$\begin{aligned} \det G_N &= \frac{\prod_{n=0}^N h_n}{D_t(P_k; s) |\det V_\beta|^2}, \\ \det G'_{N-v} &= \frac{\prod_{n=0}^{N-v} h_n}{D_{t+\Delta-v}(P_{k'}; s) |\det V_{\beta'}|^2}. \end{aligned} \quad (\text{CTV8})$$

The relation phase i^q or $i^{q'}$ multiplying each source relation column has squared modulus one, as used in this equality. The Vandermonde factors are nonzero by the original distinctness of all roots. Their logarithmic difference is constant across the four endpoints and has signed coefficient zero. Therefore CTV8 proves the exact finite expression

$$\mathcal{T}_k^{(s)} = \sum_N \sigma_N \left[\sum_{n=N-v+1}^N \log h_n + \log D_{t+\Delta-v}(P_{k'}; s) - \log D_t(P_k; s) \right]. \quad (\text{CTV9})$$

The first sum is empty for $v = 0$. In particular both low relation determinants and both high relation determinants remain present.

30.3 The exact quotient of the outer-strip polynomial

Introduce the positive measure

$$d\zeta_{k',s}(y) = |P_{k'}(y)|^2 dm_s(y). \quad (\text{CTV10})$$

This measure is the exact pullback of the original source metric under the multiplication map $\mathcal{M}_{P_{k'}} : f \mapsto P_{k'}f$. Indeed $\|\mathcal{M}_{P_{k'}}f\|_{m_s}^2 = \int |f|^2 d\zeta_{k',s}$, including the same full factor M_s . On polynomials of degree at most $N - q'$ its image is the specified subspace $P_{k'}\mathbb{C}[y]_{\leq N-q'}$ of the original degree-at-most- N source. The map is injective because $P_{k'}$ is nonzero. Thus CTV10 records a proved metric map to the original source, rather than an assigned new Gamma order.

For $L = N - q' = t + \Delta - 1$, let $Q_{\partial,L}^{(s)}$ be the minimum quotient Gram of

$$\mathbb{C}[y]_{\leq L} \longrightarrow \mathbb{C}[y]/(P_{\partial,k}) \quad (\text{CTV11})$$

in the source norm CTV10 and in the increasing monomial remainder frame of dimension Δ . The source degree is at least $\Delta - 1$ at every endpoint. The same monic-frame Schur proof as in CTV7, now using CTV4, gives

$$\det Q_{\partial,L}^{(s)} = \frac{D_{t+\Delta}(P_{k'}; s)}{D_t(P_k; s)}. \quad (\text{CTV12})$$

In fact the ideal columns are $P_{\partial,k}y^j$, $0 \leq j < t$; their Gram in CTV10 is exactly $D_t(P_k; s)$ by CTV4. The numerator is the Gram of the complete monomial source of dimension $L + 1 = t + \Delta$. This proves every source and relation factor in CTV12.

Write $\lambda_j^{(k',s)}$ for the squared norm of the monic orthogonal polynomial of degree j in CTV10. This polynomial and its norm are completely determined by the positive moment matrices: subtract from y^j its projection onto lower powers. If H_j is that lower Gram and a_j its inner-product column with y^j , the lower coefficients are $-H_j^{-1}a_j$. Completing the square proves minimality among all monic polynomials of degree j ; its squared norm is the Schur complement. Thus $\lambda_j^{(k',s)} = D_{j+1}(P_{k'}; s)/D_j(P_{k'}; s) > 0$. Multiplying these identities and substituting CTV12 into CTV9 gives

$$\begin{aligned} \mathcal{F}_{\partial,k}^{(s)} &= \log \det Q_{\partial,\Delta-1}^{(s)} + \log \det Q_{\partial,\Delta}^{(s)} - \log \det Q_{\partial,q+\Delta-1}^{(s)} - \log \det Q_{\partial,q+\Delta}^{(s)}, \\ \mathcal{T}_k^{(s)} &= \mathcal{F}_{\partial,k}^{(s)} + \sum_N \sigma_N \mathcal{R}_{N,v}^{(s)}, \\ \mathcal{R}_{N,v}^{(s)} &= \sum_{a=0}^{v-1} \left(\log h_{N-a} - \log \lambda_{N-q'-a}^{(k',s)} \right). \end{aligned} \quad (\text{CTV13})$$

Every weighted norm index is nonnegative: $N - q' - (v - 1) \geq \Delta - v \geq 16$. The total difference therefore has an exact rank- Δ quotient presentation with precisely the displayed norm tails.

30.4 Finite bounds for all norm tails in both original sources

The original Gamma generating identity gives the full exponential moment $\int e^{ty} dm_s(y) = M_s(\cos t)^{-b}$ for $|t| < \pi/2$. By the evenness of the measure this also equals $\int \cosh(ty) dm_s(y)$. The nonnegative Taylor terms of $\cosh$ imply

$$\begin{aligned} \int y^{2j} dm_s(y) &\leq M_s(\cos t_0)^{-b} \frac{(2j)!}{t_0^{2j}}, \\ \|y^j\|_{m_s} &\leq \sqrt{M_s}(\cos t_0)^{-b/2} \mathfrak{c}_{\text{mom}}^j j!, \quad t_0 = \frac{\pi}{4}, \quad \mathfrak{c}_{\text{mom}} = \frac{2}{t_0} = \frac{8}{\pi}. \end{aligned} \quad (\text{CTV14})$$

The second inequality uses $(2j)! \leq 4^j (j!)^2$, proved by $\binom{2j}{j} \leq \sum_{a=0}^{2j} \binom{2j}{a} = 4^j$. Thus every constant in this moment estimate includes the original source mass and order.

For a monic polynomial of degree n whose roots have modulus at most R' , its coefficient of y^a has absolute value at most $\binom{n}{a} (R')^{n-a}$: it is the sum of those products of $n - a$ distinct roots. The triangle inequality and CTV14 therefore give the full bound

$$\|P\|_{m_s} \leq \sqrt{M_s}(\cos t_0)^{-b/2} \mathfrak{c}_{\text{mom}}^n n! \sum_{h=0}^n \frac{(R'/\mathfrak{c}_{\text{mom}})^h}{h!} \leq \sqrt{M_s}(\cos t_0)^{-b/2} \mathfrak{c}_{\text{mom}}^n n! e^{R'/\mathfrak{c}_{\text{mom}}}. \quad (\text{CTV15})$$

The monic test polynomial y^j in the minimization defining $\lambda_j^{(k',s)}$ gives the polynomial $P_{k'}(y)y^j$ in CTV15, of degree $n = q' + j$; its extra j roots are exactly zero and also satisfy the stated bound. For the lower bound, CTV5 applies to every competing monic polynomial, whose original Gamma squared norm is at least h_j . Taking the minimum proves

$$\delta^{2q'} h_j \leq \lambda_j^{(k',s)} \leq M_s(\cos t_0)^{-b} \mathfrak{c}_{\text{mom}}^{2(q'+j)} ((q' + j)!)^2 e^{2R'/\mathfrak{c}_{\text{mom}}}. \quad (\text{CTV16})$$

No lower bound on an actual period coefficient is used here. The lower root polynomial itself supplies the strictly positive real-line lower bound.

For each term of CTV13 put $n = N - a$ and $j = n - q'$. Subtracting the logarithms in CTV16 from the exact $\log h_n = \log M_s + \log n! + \log(b)_n$ yields

$$\begin{aligned} b \log(\cos t_0) + \log \frac{(b)_n}{n!} - 2n \log \mathbf{c}_{\text{mom}} - \frac{2R'}{\mathbf{c}_{\text{mom}}} &\leq \log h_n - \log \lambda_j^{(k',s)} \\ &\leq \sum_{r=j+1}^n (\log r + \log(b+r-1)) - 2q' \log \delta. \end{aligned} \quad (\text{CTV17})$$

The two full $\log M_s$ contributions cancel in this displayed comparison because both norms arise from exactly CTV1. They have not been removed from either individual norm.

Let $c_b^{\log} = \max\{1, b-1\}$. As proved directly in CEP19, $|\log(j/(b+j-1))| \leq c_b^{\log}/j$ for $b \geq 1/2$: use $\log(1+x) \leq x$ for $b \geq 1$; for $1/2 \leq b < 1$ use $-\log(1-a/j) \leq a/(j-a) \leq 1/j$, $a = 1-b$. Summation gives $|\log((b)_n/n!)| \leq c_b^{\log} H_n$. With all original endpoints $N \geq 99$, define the explicit nonnegative numbers

$$\begin{aligned} L_N^{\text{tail}} &= c_b^{\log} H_N + \frac{b}{2} \log 2 + 2N \log(8/\pi) + \frac{\pi R'}{4}, \\ U_N^{\text{tail}} &= q'(\log N + \log(b+N-1) + 2|\log \delta|), \\ C_{N,v}^{\text{tail}} &= v \max\{L_N^{\text{tail}}, U_N^{\text{tail}}\}. \end{aligned} \quad (\text{CTV18})$$

Indeed $\cos(\pi/4) = 2^{-1/2}$ and $2R'/\mathbf{c}_{\text{mom}} = \pi R'/4$. In the upper sum of CTV17 there are exactly q' terms, and each is at most $\log N + \log(b+N-1)$. Thus CTV17 proves

$$-v L_N^{\text{tail}} \leq \mathcal{R}_{N,v}^{(s)} \leq v U_N^{\text{tail}}, \quad |\mathcal{R}_{N,v}^{(s)}| \leq C_{N,v}^{\text{tail}}. \quad (\text{CTV19})$$

When $v = 0$ the equality $\mathcal{R}_{N,0}^{(s)} = 0$ and the zero bound follow from the empty sum. These estimates apply for all actual endpoints and for both unchanged source orders simultaneously.

30.5 The proved first-variation reduction and its remaining matrix

Since $N \leq 2q$, $q' \leq q$, $b \leq k/2$ and $R' \leq k\sqrt{\delta^2 + \gamma^2}$, equations CTV18–19 give $C_{N,v}^{\text{tail}} = O_{\delta,\gamma}(q \log(q+1) + k \log(q+1))$ uniformly for every integer $0 \leq v \leq 80$. To see this directly, use $H_N \leq 1 + \log N$ in the first line of CTV18 and the displayed upper bounds in its second; v is bounded by the fixed integer 80. Dividing by kq tends to zero, since $\log(q+1)/k \rightarrow 0$ for $q = (k+1)^2$. Consequently CTV13 and the finite bounds just proved yield

$$\left| \mathcal{T}_k^{(s)} - \mathcal{F}_{\partial,k}^{(s)} \right| \leq \sum_N C_{N,v}^{\text{tail}} = o(kq). \quad (\text{CTV20})$$

The estimate is uniform over the possible actual moment indices v ; it requires no bound on the remaining period coordinates. Every polynomial and measure defining $\mathcal{F}_{\partial,k}^{(s)}$ is independent of that index.

For an explicit matrix target, let H_L^ζ be the moment Gram of $1, y, \dots, y^L$ in the full measure CTV10, and let $\mathbf{R}_{\partial,L}$ be the literal remainder matrix modulo the full polynomial $P_{\partial,k}$ in CTV4. The exact metric is

$$Q_{\partial,L}^{(s)} = (\mathbf{R}_{\partial,L}(H_L^\zeta)^{-1} \mathbf{R}_{\partial,L}^*)^{-1}. \quad (\text{CTV21})$$

The map is onto because its first Δ columns are the identity in the remainder frame. The minimum source lift of a coefficient vector x is $(H_L^\zeta)^{-1} \mathbf{R}_{\partial,L}^* Q_{\partial,L}^{(s)} x$; it maps to x and is perpendicular to the kernel. Adding any other lift adds a perpendicular kernel vector, proving the stated Gram. This also proves direct equality with the quotient used in CTV11–12.

Combining the complete exact CEP49 and CTV13 identities gives

$$\begin{aligned}
F_K^{(s)} &= \mathcal{F}_{\partial,k}^{(s)} - F_{\Xi,k}^{(s)} + \mathcal{E}_{A,k}^{(s)} + \sum_N \sigma_N \mathcal{R}_{N,v}^{(s)}, \\
\left| F_K^{(s)} - \mathcal{F}_{\partial,k}^{(s)} + F_{\Xi,k}^{(s)} \right| &\leq 2 \sum_N E_N + \sum_N C_{N,v}^{\text{tail}} = o(kq).
\end{aligned} \tag{CTV22}$$

Here the first error has its fixed-actual-period scope from CEP49; the new tail bound has the stronger uniform-in- v scope just proved. Thus the remaining leading calculation now consists of the explicit outer-strip quotient of rank $16k - 48$ and the original observation quotient of rank $8k - 32$. Both have their complete original metric maps and four cutoffs specified. No value of either leading volume is assigned by this reduction, and no derivative of the earlier q^2 asymptotic law was used.

31 The actual observation remaining inside the conductor kernel

Retain the original simple quartet, its actual unit and period matrix $\mathbf{H} = F^{-1}\Pi U\mathcal{E}_{\text{prim}}$, and its proved five-member quadric orbit. Put $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $q' = (k - 7)^2$, $\Delta = q - q' = 16k - 48$, $r = 8k - 16$, and $j = 8k - 32$. The period parameter and logarithm branch are fixed at a point on this proved locus. All constructions below use the original ordered primary coordinates at

$$\beta_{ab} = k/2 + (2a - k)\delta + i(2b - k)\gamma, \quad \omega_{ab} = (2b - k)\gamma - i(2a - k)\delta, \quad 0 \leq a, b \leq k. \quad (\text{ROQ1})$$

The full conductor is the actual biform $A = \sum_{a,b=0}^8 a_{ab}e_{ab}^{(8)}$ of OCF4. Its matrix $C : \mathbb{C}^q \rightarrow \mathbb{C}^{q'}$ is exactly CEP4: $C_{(a,b),(i,h)} = a_{i-a,h-b}$ when both differences belong to $\{0, \dots, 8\}$, and zero otherwise. Thus $C^\top$ is multiplication by the entire A , and $V_A = \ker C$. The original observation is Λ , and $K = \ker \Lambda \subset V_A$. Ordinary transpose throughout the coefficient duality is denoted by $\top$; a star denotes conjugate transpose in a displayed metric.

31.1 An explicit residual map and its actual boundary basis

Use the complete OCF6 elimination, with its actual largest nonzero coefficient $a_{r_*s_*}$ and every remaining a_{ab} . To avoid confusing its degree- k matrices with the source cutoff, write them as

$$\mathbf{N} : \mathbb{C}^q \rightarrow \mathbb{C}^\Delta, \quad \mathbf{S} : \mathbb{C}^\Delta \hookrightarrow \mathbb{C}^q, \quad \mathbf{V} : \mathbb{C}^q \rightarrow \mathbb{C}^{q'}, \quad \mathbf{S}\mathbf{N} + C^\top \mathbf{V} = I_q, \quad \mathbf{N}\mathbf{S} = I_\Delta. \quad (\text{ROQ2})$$

These are the exact identities OCF7. Specifically, at each decreasing lexicographic conductor position the elimination subtracts its current coefficient divided by $a_{r_*s_*}$ times the entire translated biform $Ae_{ab}^{(k-8)}$. This specifies all entries of $\mathbf{N}, \mathbf{V}$; $\mathbf{S}$ inserts the retained strip coordinates. It also proves $\ker \mathbf{N} = \text{im } C^\top$. Taking the annihilator for the bilinear coefficient pairing therefore gives

$$I_V = \mathbf{N}^\top : \mathbb{C}^\Delta \xrightarrow{\sim} V_A, \quad I_V^{-1} = \mathbf{S}^\top|_{V_A}. \quad (\text{ROQ3})$$

For a direct verification, $C\mathbf{N}^\top = 0$ by transposing $\mathbf{N}C^\top = 0$, and $\mathbf{S}^\top \mathbf{N}^\top = I$. If $Cx = 0$, transposing the first identity of ROQ2 gives $x = \mathbf{N}^\top \mathbf{S}^\top x$.

Retain the j independent original invariant certificates $F_{k,h}(x)$ selected by OCF20, and their entire biform coefficient columns f_h :

$$\mathbf{F} = [f_1, \dots, f_j], \quad f_h = \text{coeff}(F_{k,h}(\mathbf{H}t)), \quad \mathbf{Z} = \mathbf{N}\mathbf{F} \in \text{Mat}_{\Delta \times j}(\mathbb{C}), \quad \mathbf{D} = \mathbf{F}^\top. \quad (\text{ROQ4})$$

Here $t = (z_0w_0, z_0w_1, z_1w_0, z_1w_1)^\top$ and the coefficient order is the full original biform order. Every certificate is supplied by the fourteen-generator recursion OCF25–26, or equivalently the complete twenty-five-generator recursion OCF9–20. At a recursion step its certificate is the actual product $x^\nu F_{k-e,h}$, and its column is $\mathbf{N}_k M_{f_\nu} \mathbf{S}_{k-e} Z_{k-e,h}$. The coefficients of f_ν are the full multinomials OCF13 in the actual $\mathbf{H}$. Thus ROQ4 specifies finite entries without choosing a substitute period or introducing a freely chosen observation. The proved image and rank are $\text{im } \mathbf{Z} = \mathbf{N}J_k$ and $\text{rank } \mathbf{Z} = j$.

Let I be the j retained pivot rows of $\mathbf{Z}$, let T be their complement among the Δ strip rows, and put $\mathbf{Z}_I = E_I^\top \mathbf{Z}$, $\mathbf{Z}_T = E_T^\top \mathbf{Z}$. The original OCF selection guarantees that $\mathbf{Z}_I$ is invertible on its exhaustive exact pivot chart. ROQ2 applied to $\mathbf{F}$ gives the full coefficient identity

$$\mathbf{F} = \mathbf{S}\mathbf{Z} + C^\top \mathbf{V}\mathbf{F}, \quad \mathbf{D} = \mathbf{Z}^\top \mathbf{S}^\top + (\mathbf{V}\mathbf{F})^\top C. \quad (\text{ROQ5})$$

In particular the unreduced invariant rows and the reduced strip rows agree on V_A , with precisely the displayed conductor multiple as their ambient difference. The residual observation has the explicit forms

$$\Xi : V_A \rightarrow \mathbb{C}^j, \quad x \mapsto Dx, \quad \Xi I_V = \mathbf{Z}^\top, \quad L = \mathbf{N}^\top E_I (\mathbf{Z}_I^\top)^{-1}, \quad CL = 0, \quad DL = I_j. \quad (\text{ROQ6})$$

The last three equalities follow by multiplication and ROQ3–5. Moreover $\ker \Xi = K$: the rows C and D together span the coefficient space J_k . Indeed conductor multiplication spans $\ker \mathbf{N}$, its image is contained in J_k by OCF17, and the f_h project to the basis Z of $\mathbf{N}J_k$. Thus a primary vector killed by both C and D is exactly the annihilator of J_k , which is the original K by OKM3–5. This proves rank j , surjectivity, and the complete exact sequence

$$0 \longrightarrow K \longrightarrow V_A \xrightarrow{\Xi} \mathbb{C}^j \longrightarrow 0. \quad (\text{ROQ7})$$

For every $y \in \mathbb{C}^j$ define $\mathbf{b}(y) = \Lambda Ly$. These vectors give the literal isomorphism $\mathbf{b} : \mathbb{C}^j \rightarrow \Lambda(V_A)$. They span because for $x \in V_A$, $x - LDx \in K$ by ROQ6–7. They are independent because $\Lambda Ly = 0$ implies $Ly \in K$, hence $y = DLy = 0$. The inverse on the displayed image is $\Lambda x \mapsto Dx$ for $x \in V_A$: replacing x by another such lift changes it by K , which D kills. In particular the boundary basis vectors are exactly $\mathbf{b}_h = \sum_{a,b} L_{(a,b),h} \Lambda \varepsilon_{ab}$. Each $\Lambda \varepsilon_{ab}$ is the full original projected tensor coefficient in OKA7, including its literal $1/5$ cyclic average.

The whole original observation can be recovered from these rows as well. Take an actual right inverse L_C of C , for example the inverse of the first nonzero square column minor with its coordinate insertion, and define

$$\mathbf{L} = [L_C - LDL_C, L], \quad \mathbf{R} = \begin{pmatrix} C \\ D \end{pmatrix}, \quad \mathbf{R}\mathbf{L} = I_{q'+j}, \quad \Lambda = (\Lambda\mathbf{L})\mathbf{R}. \quad (\text{ROQ8})$$

The first identity follows by multiplying its four blocks. For the last, $\mathbf{R}(x - \mathbf{L}\mathbf{R}x) = 0$, whose kernel is K by ROQ7. Thus the $q' + j = q - r$ columns of $\Lambda\mathbf{L}$ form an actual basis of the original boundary: applying its kernel and then $\mathbf{R}\mathbf{L} = I$ proves independence, and ROQ8 proves spanning. The last j columns are exactly $\mathbf{b}_h$.

For completeness the inherited period metric on this residual basis is also explicit. Let B be the complete matrix AKS8 in primary coordinates, so $j\Lambda = \mathcal{F}_0 B$. If ν, μ range over the original weight-zero degree- k monomials, set

$$(\mathcal{G}_0)_{\nu\mu} = \sum_{\substack{n_{tu} \geq 0 \\ \sum_u n_{tu} = \nu_t \\ \sum_t n_{tu} = \mu_u}} \frac{k!}{\prod_{t,u} n_{tu}!} \prod_{t,u} [5(\delta_{tu} + 1)]^{n_{tu}}, \quad P_{\text{per}} = L^* B^* \mathcal{G}_0 B L > 0. \quad (\text{ROQ9})$$

Expansion of the original word inner product gives $\mathcal{G}_0 = \mathcal{F}_0^* \mathcal{F}_0$ as in AKS10, so the displayed matrix is exactly the Gram of the independent vectors $j\mathbf{b}_h$. Its strict positivity follows from that independence and injectivity of j . All off-diagonal word terms and every actual entry of $\mathbf{H}$ remain in ROQ9.

31.2 All four induced Gamma metrics and their fixed determinant factor

For either original source $s \in \{1, k\}$ put $b_s = s/2$, $M_s = (2\pi)^{s/2}$ and retain

$$\begin{aligned} p_0 &= 1, & p_1 &= y, & p_{n+1} &= yp_n - n(b_s + n - 1)p_{n-1}, \\ h_n &= M_s n! (b_s)_n, & (u_n)_{ab} &= p_n(\omega_{ab}) / \sqrt{h_n}. \end{aligned} \quad (\text{ROQ10})$$

Thus $U_N = [u_0, \dots, u_N]$ is precisely CEP21 and $R_N = U_N U_N^* > 0$, $G_N = R_N^{-1}$ for every $N \geq q - 1$. The positivity follows from its first q columns, whose determinant is the original Vandermonde times their nonzero leading coefficients. Define

$$H_{V,N} = \bar{\mathbf{N}} G_N \mathbf{N}^\top, \quad Q_{\Xi,N} = (Z^\top H_{V,N}^{-1} \bar{Z})^{-1} > 0. \quad (\text{ROQ11})$$

These are matrices of sizes Δ and j , respectively. The minimum of $z^* H_{V,N} z$ under $Z^\top z = y$ is attained at $z = H_{V,N}^{-1} \bar{Z} Q_{\Xi,N} y$. Its difference from any other lift lies in $\ker Z^\top$ and is perpendicular in the $H_{V,N}$ metric. Its squared norm is $y^* Q_{\Xi,N} y$. Thus ROQ11 calculates exactly the induced

metric of ROQ7 in the actual boundary coordinates ROQ6. This proof also identifies it with the quotient metric appearing in CEP27 and CEP49.

The actual kernel frame in the strip is

$$\rho = E_T^\top - Z_T Z_I^{-1} E_I^\top, \quad I_K = \mathbf{N}^\top \rho^\top, \quad F_{\text{str}} = [\rho^\top, E_I (Z_I^\top)^{-1}]. \quad (\text{ROQ12})$$

The first two matrices are precisely the OCF/OKM frame. In the ordered rows (T, I) the last one has blocks $\begin{pmatrix} I_r & 0 \\ -(Z_I^\top)^{-1} Z_T^\top & (Z_I^\top)^{-1} \end{pmatrix}$. Consequently $|\det F_{\text{str}}|^2 = |\det Z_I|^{-2}$, including the absolute value of the row-order permutation. Block elimination in $F_{\text{str}}^* H_{V,N} F_{\text{str}}$ gives the exact identity

$$\det(I_K^* G_N I_K) \det Q_{\Xi,N} = \frac{\det H_{V,N}}{|\det Z_I|^2}. \quad (\text{ROQ13})$$

Indeed its upper block is the kernel Gram and its Schur complement is the minimum quotient metric of the previous paragraph, since $Z^\top F_{\text{str}} = [0, I_j]$. The determinant of a block matrix is the determinant of its upper block times that Schur complement, while congruence contributes $|\det F_{\text{str}}|^2$. This proves ROQ13 without assigning modulus one to the actual pivot.

Define $F_{\Xi,k}^{(s)}$ by the four ordered endpoints $\mathcal{N} = (q-1, q, 2q-1, 2q)$ and signs $\sigma = (1, 1, -1, -1)$:

$$F_{\Xi,k}^{(s)} = \sum_{N \in \mathcal{N}} \sigma_N \log \det Q_{\Xi,N}. \quad (\text{ROQ14})$$

The fixed factor $|\det Z_I|^{-2}$ in ROQ13 cancels with these signs. If a different exact certificate basis is selected, its residual rows on V_A are $D' = MD$ for an invertible $j \times j$ matrix M . This follows because both lists are bases of the dual of V_A/K . For an explicit inverse formula, $M = D'L$, and the old and new right sections give its inverse; composing the two surjections proves both inverse identities. The new metric is $Q'_{\Xi,N} = M^{-*} Q_{\Xi,N} M^{-1}$, directly from their identical minimum problem. Its determinant changes by $|\det M|^{-2}$, which also cancels in ROQ14. No pivot or source mass has been removed at an individual endpoint. In fact ROQ10 gives $Q_{\Xi,N} = M_s \widehat{Q}_{\Xi,N}$ when the common mass is explicitly factored out of every column; its determinant carries M_s^j . The metric relative to the retained period Gram ROQ9 has determinant $\det Q_{\Xi,N} / \det P_{\text{per}}$, whose fixed denominator cancels only in the same four-endpoint sum.

31.3 The exact constrained covariance in the original source

The residual metric can be computed without inverting the strip frame. Define, using the complete original covariance,

$$\begin{aligned} A_N^C &= C R_N C^*, & B_N^D &= D R_N C^*, & T_N^D &= D R_N D^*, \\ W_N &= R_N - R_N C^* (A_N^C)^{-1} C R_N, \\ \Omega_N &= D W_N D^* = T_N^D - B_N^D (A_N^C)^{-1} (B_N^D)^*. \end{aligned} \quad (\text{ROQ15})$$

Since C has full row rank and $R_N > 0$, $A_N^C > 0$. There is the exact identity

$$W_N = \mathbf{N}^\top H_{V,N}^{-1} \overline{\mathbf{N}}, \quad \Omega_N = Z^\top H_{V,N}^{-1} \overline{Z} > 0, \quad Q_{\Xi,N} = \Omega_N^{-1}. \quad (\text{ROQ16})$$

To prove its first equality, $CW_N = 0$, W_N is Hermitian, and for $x \in \ker C$ direct multiplication gives $W_N G_N x = x$. The other displayed candidate is Hermitian, has image in $\ker C$ by ROQ3, and has the same property on $x = \mathbf{N}^\top z$. Such a matrix is unique: for their Hermitian difference H , its image lies in V_A and $HG_N x = 0$ for $x \in V_A$. For arbitrary y and $x \in V_A$, this yields $x^* G_N H y = (H G_N x)^* y = 0$. Putting $x = H y \in V_A$ gives $(H y)^* G_N (H y) = 0$, hence $H y = 0$. This proves equality. Now use $D \mathbf{N}^\top = Z^\top$ from ROQ6 and take the adjoint to prove the other two identities.

Every entry in ROQ15 is the finite sum of original column products: $A_N^C = \sum_{n=0}^N (Cu_n)(Cu_n)^*$, $B_N^D = \sum_{n=0}^N (Du_n)(Cu_n)^*$, and $T_N^D = \sum_{n=0}^N (Du_n)(Du_n)^*$. The conductor column is exactly $Cu_n = V_{N-v} A_N e_n$ by CEP22 when $n \leq N$; its vanishing low columns and centre difference are precisely CEP13–16. The residual column is explicitly $(Du_n)_h = \sum_{a,b} (f_h)_{ab} p_n(\omega_{ab}) / \sqrt{h_n}$. In particular ROQ15 retains all conductor–observation cross terms. Replacing D by $D + TC$ leaves Ω_N exactly unchanged because $CW_N = 0 = W_N C^*$. Applied to ROQ5, this proves that the full and strip row presentations give exactly the same metric, rather than assuming their ambient coefficient matrices coincide.

31.4 Exact increments and a positive residual four-endpoint return

For $N \geq q - 1$ let $u = u_{N+1}$ be the next original column ROQ10 and define

$$\begin{aligned} a_N &= 1 + (Cu)^*(A_N^C)^{-1}(Cu) > 0, \\ w_N &= u - R_N C^*(A_N^C)^{-1}Cu \in V_A, \quad z_N = Dw_N = Du - B_N^D(A_N^C)^{-1}Cu, \\ \eta_N &= \frac{z_N^* Q_{\Xi, N} z_N}{a_N} \geq 0. \end{aligned} \tag{ROQ17}$$

The complete updates are

$$\begin{aligned} W_{N+1} &= W_N + w_N w_N^* / a_N, \quad \Omega_{N+1} = \Omega_N + z_N z_N^* / a_N, \\ Q_{\Xi, N+1} &= Q_{\Xi, N} - \frac{Q_{\Xi, N} z_N z_N^* Q_{\Xi, N}}{a_N + z_N^* Q_{\Xi, N} z_N}, \\ \log \det Q_{\Xi, N} - \log \det Q_{\Xi, N+1} &= \log(1 + \eta_N). \end{aligned} \tag{ROQ18}$$

Here is a direct proof of the covariance update. Put $a = Cu$ and $v = R_N C^*(A_N^C)^{-1}a$. Since $R_{N+1} = R_N + uu^*$, $A_{N+1}^C = A_N^C + aa^*$, whose inverse is $(A_N^C)^{-1} - (A_N^C)^{-1}aa^*(A_N^C)^{-1}/a_N$; multiplying by $A_N^C + aa^*$ verifies this inverse exactly. For the full covariance calculation write $\alpha = a^*(A_N^C)^{-1}a$. The subtracted covariance product expands as

$$R_N C^*(A_N^C)^{-1} C R_N + v u^* + u v^* + \alpha u u^* - \frac{(v + \alpha u)(v + \alpha u)^*}{1 + \alpha}.$$

Subtracting this from $R_N + uu^*$ and the old W_N leaves $(u - v)(u - v)^*/(1 + \alpha)$, proving the first update. Multiplying by D, D^* proves the second. The same inverse identity proves the third, and the determinant identity $\det(H + zz^*/a) = \det H(1 + z^* H^{-1} z/a)$ proves the fourth. For that identity conjugate by $H^{-1/2}$; the rank-one matrix $H^{-1/2} z z^* H^{-1/2}/a$ has one eigenvalue $z^* H^{-1} z/a$ and all its others zero. This completes the update proof.

Telescoping with the four actual endpoints, rather than changing the source degree, gives the exact finite sum

$$F_{\Xi, k}^{(s)} = \log(1 + \eta_{q-1}) + 2 \sum_{N=q}^{2q-2} \log(1 + \eta_N) + \log(1 + \eta_{2q-1}) \geq 0. \tag{ROQ19}$$

Indeed $\log \det Q_{q-1} - \log \det Q_{2q-1}$ sums the differences for $N = q - 1, \dots, 2q - 2$, while $\log \det Q_q - \log \det Q_{2q}$ sums those for $N = q, \dots, 2q - 1$. This proves each endpoint weight. Its first increment $\lambda_s = \log(1 + \eta_{q-1})$ is a completely specified period-dependent nonnegative lower bound for $F_{\Xi, k}^{(s)}$. The formulas also characterize equality: $\lambda_s = 0$ exactly when $Du_q = B_{q-1}^D(A_{q-1}^C)^{-1}Cu_q$; the full return vanishes exactly when this corresponding row residual is zero for every column $u_q, \dots, u_{2q}$. Positivity of $Q_{\Xi, N}$ proves both directions, without asserting a nonzero generic residual.

31.5 Finite scalar enclosures and propagation into CEP49

Retain the exact AKS3 scalars $\mathcal{Z}_{s,t}$ and $\mathcal{L}_{s,t} = \log \mathcal{Z}_{s,t}$, whose full original polynomial covariance bound is proved in AKS20–27. In primary coordinates it says

$$\mathcal{Z}_{s,t}^{-1} G_{q-1} \preceq G_{q+t-1} \preceq G_{q-1}, \quad 0 \leq t \leq q+1. \quad (\text{ROQ20})$$

The unchanged coefficient map from AKS26 to primary coordinates is the ordered V_β , so congruence by that invertible map proves exactly the displayed statement. Restrict it to V_A and take the minimum on the fixed fibre $Dx = y$ in V_A . Inequalities of squared norms hold for every such x , hence also for their attained minima. Thus $\mathcal{Z}_{s,t}^{-1} Q_{\Xi,q-1} \preceq Q_{\Xi,q+t-1} \preceq Q_{\Xi,q-1}$. If $d_t = \log \det Q_{\Xi,q-1} - \log \det Q_{\Xi,q+t-1}$, the eigenvalues after congruence by $Q_{\Xi,q-1}^{-1/2}$ give $0 \leq d_t \leq j\mathcal{L}_{s,t}$, and ROQ18 gives their monotonicity. Since $d_1 = \lambda_s$, the exact four-endpoint identity $F_{\Xi,k}^{(s)} = d_q + d_{q+1} - d_1$ proves

$$\lambda_s \leq F_{\Xi,k}^{(s)} \leq j(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - \lambda_s. \quad (\text{ROQ21})$$

The actual rank j enters only through this proved eigenvalue bound; the determinant and increment formulas ROQ11–19 retain its full orientation. No leading volume is assigned from a dimension.

Let $\epsilon_{A,k}^{(s)} = 2 \sum_{N \in \mathcal{N}} E_N$ with the complete original constants of CEP29–36. CEP49 gives $F_K^{(s)} = \mathcal{T}_k^{(s)} - F_{\Xi,k}^{(s)} + \mathcal{E}_{A,k}^{(s)}$ and $|\mathcal{E}_{A,k}^{(s)}| \leq \epsilon_{A,k}^{(s)} = o(kq)$ for a fixed actual period. Substitution of ROQ21 proves

$$\begin{aligned} \mathcal{T}_k^{(s)} - \epsilon_{A,k}^{(s)} - j(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) + \lambda_s &\leq F_K^{(s)}, \\ F_K^{(s)} &\leq \mathcal{T}_k^{(s)} + \epsilon_{A,k}^{(s)} - \lambda_s. \end{aligned} \quad (\text{ROQ22})$$

Intersecting this interval with the full AKJ2 or AKJ4 intervals preserves each of their finite constants and source correlations. More precisely, one may use the exact value $\mathcal{E}_{A,k}^{(s)}$ in place of its upper or lower envelope in the appropriate side. Formula ROQ19 supplies every additional original positive subtraction, rather than only the first one. These are finite results on the residual quotient itself; they do not claim that its leading kq coefficient has been evaluated.

31.6 The complete residual quotient in the conductor graph

At a fixed original endpoint N , let $\mathcal{P}_N$, I_N , $\mathcal{W}_N = U_N^{-1}(V_A)$ and the actual source isomorphism $S_N p = (P_{<v} p, \mathcal{T}_{A,N} p)$ be precisely CEP28–31. The source carries its original orthonormal Gamma coordinates of order $s = 1$ or $s = k$. The induced linear isomorphism on the residual observation quotient is

$$\mathcal{W}_N / U_N^{-1}(K) \xrightarrow{[p] \mapsto [S_N p]} S_N \mathcal{W}_N / S_N U_N^{-1}(K), \quad [p] \mapsto D U_N p \in \mathcal{C}^j. \quad (\text{ROQ23})$$

The last map identifies the left quotient with V_A/K and its fixed coordinates ROQ6. Its kernel and surjectivity follow from ROQ7 and surjectivity of U_N ; applying the invertible S_N proves the other isomorphism. Thus all relation directions, including the source ideal and the entire actual K , are retained in the transported denominator.

Here are exact matrices of that denominator and the resulting metric. Let Z_N^χ be the original source coefficient columns $\chi_k S^a$, $0 \leq a \leq N - q$, with no columns when $N = q - 1$. Take the actual primary frames I_K of ROQ12 and L of ROQ6 and set

$$Y_{K,N} = U_N^* G_N I_K, \quad Y_{L,N} = U_N^* G_N L, \quad J_N = [Z_N^\chi, Y_{K,N}]. \quad (\text{ROQ24})$$

The columns of $Y_{K,N}$ are the minimum lifts of I_K by the same orthogonality proof as OKM8. In particular they are perpendicular to $I_N = \ker U_N$, so J_N has independent columns and its image

is exactly $U_N^{-1}(K)$. It has $N + 1 - q + r$ columns, while $\dim \mathcal{W}_N = N + 1 - q + \Delta$, leaving precisely j quotient dimensions. The transported complete relation projector and Gram are

$$\begin{aligned} P_N^{\Xi, \text{graph}} &= I - S_N J_N (J_N^* S_N^* S_N J_N)^{-1} J_N^* S_N^*, \\ \tilde{Q}_{\Xi, N} &= Y_{L, N}^* S_N^* P_N^{\Xi, \text{graph}} S_N Y_{L, N} > 0. \end{aligned} \tag{ROQ25}$$

The inverse exists because S_N is invertible and J_N has independent columns. Multiplication verifies that the projector is Hermitian, idempotent and projects onto the orthogonal complement of the entire $S_N U_N^{-1}(K)$. Thus its Gram is the minimum norm of the transported quotient ROQ23. Replacing $Y_{L, N}$ by another source lift of L adds columns in I_N , which this projector kills. Replacing the section L by another section of ROQ6 adds columns in K ; their lifts lie in the span of J_N and are killed. Therefore ROQ25 computes the specified quotient metric independently of either choice.

Before applying S_N , the same projection gives

$$Q_{\Xi, N} = L^* G_N L - L^* G_N I_K (I_K^* G_N I_K)^{-1} I_K^* G_N L. \tag{ROQ26}$$

To prove it, $Y_{L, N}$ and $Y_{K, N}$ are perpendicular to the ideal, so projection off J_N only subtracts $Y_{K, N} (Y_{K, N}^* Y_{K, N})^{-1} Y_{K, N}^* Y_{L, N}$. Using $U_N U_N^* = R_N$ in their Gram products yields exactly ROQ26. It is also the minimum of the original G_N norm on $Ly + K$, so it agrees with the already proved ROQ11 and ROQ16.

The all-rank exterior estimate CEP33–35 applies to ROQ23: represent a quotient class by its minimum lift in the denominator's orthogonal complement, apply S_N , then project off the target denominator. This projection has every exterior norm at most one. Consequently the induced j th exterior norm is at most $\|\bigwedge^j S_N\|$, and its inverse is at most $\|\bigwedge^j S_N^{-1}\|$ by the identical argument for S_N^{-1} . CEP34 bounds both logarithms from above by E_N . Applying them to the fixed frame ROQ6 and squaring exterior volumes gives

$$\left| \log \det \tilde{Q}_{\Xi, N} - \log \det Q_{\Xi, N} \right| \leq 2E_N, \quad \left| \sum_{N \in \mathcal{N}} \sigma_N \log \det \tilde{Q}_{\Xi, N} - F_{\Xi, k}^{(s)} \right| \leq \epsilon_{A, k}^{(s)} = o(kq). \tag{ROQ27}$$

All cross terms in $J_N^* S_N^* S_N J_N$ remain present in ROQ25, including $P_{<v}(\chi_k S^a)$ and its pairings with the minimum kernel lifts. Its ideal columns can equally be generated from the full products d_{rs} and maps $\mathcal{U}_{A, L}$ in CEP41–45, because those formulas prove exactly $S_N Z_N^X$ column by column. This supplies the typed relation between the explicit residual observation and the conductor graph at all four original cutoffs.

32 A quantitative power comparison for the actual outer-strip quotient

Retain the original real parameters and integers

$$\begin{aligned} 0 < \delta < \frac{1}{2}, \quad \gamma > 2, \quad k = 4l + 1 \geq 9, \\ q = (k + 1)^2, \quad k' = k - 8, \quad q' = (k - 7)^2, \quad \Delta = q - q' = 16k - 48. \end{aligned} \quad (\text{OSP1})$$

The two source orders in every formula below are precisely $s = 1, k$. Set $b = s/2$, $M_s = (2\pi)^{s/2}$, and retain the full original measure

$$dm_s(y) = M_s \frac{2^b}{4\pi\Gamma(b)} |\Gamma(b/2 + iy/2)|^2 dy. \quad (\text{OSP2})$$

In the original coordinate $S = k/2 + iy$, and respectively $S' = k/2 - 4 + iy$, the complete monic root polynomials are

$$\begin{aligned} P_k(y) &= i^{-q} \chi_k(k/2 + iy) = \prod_{a, b_0=0}^k [y - (2b_0 - k)\gamma + i(2a - k)\delta], \\ P_{k'}(y) &= i^{-q'} \chi_{k'}(k/2 - 4 + iy) = \prod_{a, b_0=0}^{k'} [y - (2b_0 - k')\gamma + i(2a - k')\delta]. \end{aligned} \quad (\text{OSP3})$$

The index b_0 in these products is distinct from the source parameter $b = s/2$. Each root occurs with exactly its displayed multiplicity. The map $(a, b_0) \mapsto (a + 4, b_0 + 4)$ identifies the second root list with the central subrectangle of the first, so

$$P_k = P_{k'} P_{\partial, k}, \quad P_{\partial, k}(y) = \prod_{(a, b_0) \in \{0, \dots, k\}^2 \setminus \{4, \dots, k-4\}^2} [y - (2b_0 - k)\gamma + i(2a - k)\delta]. \quad (\text{OSP4})$$

This proves the factorization and its monic constant; the outer degree is Δ . Both complete root lists are invariant under $\omega \mapsto -\omega$. Every root has modulus at most

$$R = k\sqrt{\delta^2 + \gamma^2}. \quad (\text{OSP5})$$

32.1 The two original source masses and a proved interval lower bound

We first record the source facts needed in the comparison, including their constants and proofs. For $a > 0$ and real x , the two Euler integrals and the change of variables in the beta integral give

$$\frac{\Gamma(a + ix)\Gamma(a - ix)}{\Gamma(2a)} = \int_{\mathbb{R}} e^{ixu} (2 \cosh(u/2))^{-2a} du. \quad (\text{OSP6})$$

For example the intermediate beta integral is $\int_0^\infty t^{a+ix-1} (1+t)^{-2a} dt$, and $t = e^u$ gives the displayed expression. Absolute convergence follows from $a > 0$. Rotating Euler's integral through $\text{sgn}(y)\theta$, $0 < \theta < \pi/2$, gives

$$|\Gamma(a + iy/2)| \leq e^{-\theta|y|/2} \Gamma(a) (\cos \theta)^{-a}. \quad (\text{OSP7})$$

The rotation follows from Cauchy's theorem on the intervening sector: the large circular arc tends to zero because its real exponential has positive cosine, and the small one tends to zero because $a > 0$. The remaining radial integral has absolute value at most $\Gamma(a)(\cos \theta)^{-a}$. In particular the Fourier transform in OSP6 is integrable. Fourier inversion, followed by $y = 2x$, now gives

$$\int_{\mathbb{R}} e^{i\xi y} dm_s(y) = M_s (\cosh \xi)^{-b}, \quad \int_{\mathbb{R}} dm_s = M_s. \quad (\text{OSP8})$$

The same identity holds analytically for $|\operatorname{Im} \xi| < \pi/2$: OSP7 with any larger $\theta < \pi/2$ supplies an integrable dominating function on each smaller strip. Substituting $\xi = -it$ proves

$$\int_{\mathbb{R}} e^{ty} dm_s(y) = M_s(\cos t)^{-b}, \quad |t| < \pi/2. \quad (\text{OSP9})$$

This also proves finiteness of all moments used below.

Write

$$d\sigma(y) = \frac{|\Gamma(1/4 + iy/2)|^2}{2\pi} dy, \quad C_L = \frac{\pi}{\Gamma(3/4)^2}, \quad \ell_s = \frac{s-1}{4}, \quad \beta_s = \frac{M_s}{\sqrt{2}\Gamma(s/2)}. \quad (\text{OSP10})$$

The two values of ℓ_s are the integers $0, l$. The reflection identity gives a useful explicit lower bound

$$\frac{d\sigma}{dy}(y) \geq C_L e^{-\pi|y|}. \quad (\text{OSP11})$$

Here is its proof on the required domain. For $0 < \operatorname{Re} z < 1$ the Euler integrals give $\Gamma(z)\Gamma(1-z) = \int_0^\infty t^{z-1}/(1+t) dt$. On a keyhole contour with argument in $(0, 2\pi)$, the circular terms vanish with bounds $O(r^{\operatorname{Re} z})$ at zero and $O(R^{\operatorname{Re} z-1})$ at infinity. The residue at -1 is $e^{i\pi(z-1)}$; the two radial integrals therefore give $(1 - e^{2\pi iz}) \int_0^\infty t^{z-1}/(1+t) dt = 2\pi i e^{i\pi(z-1)}$. Their ratio is $\pi/\sin(\pi z)$. At $z = 1/4 + iy/2$ this proves

$$|\Gamma(1/4 + iy/2)|^2 |\Gamma(3/4 - iy/2)|^2 = \frac{2\pi^2}{\cosh(\pi y)}.$$

The unrotated Euler integral bounds the second squared factor by $\Gamma(3/4)^2$, proving $d\sigma/dy \geq C_L/\cosh(\pi y) \geq C_L e^{-\pi|y|}$.

Gamma recurrence, proved by one integration by parts in Euler's integral, gives the exact original-source identity

$$dm_s(y) = \beta_s \prod_{j=0}^{\ell_s-1} (y^2 + (2j + \frac{1}{2})^2) d\sigma(y). \quad (\text{OSP12})$$

Indeed $b/2 = \ell_s + 1/4$, and each recurrence factor before squaring and clearing denominators is $j + 1/4 + iy/2$; its squared modulus is $[y^2 + (2j + 1/2)^2]/4$. The factor $4^{-\ell_s}$ together with the prefactor in OSP2 is exactly $\beta_s/(2\pi)$, because $b = 2\ell_s + 1/2$. This proves every factor in OSP12. For $s = 1$ its product is empty and $\beta_1 = 1$. In particular, on the full interval $q \leq y \leq 2q$,

$$\frac{dm_s}{dy}(y) \geq \beta_s q^{2\ell_s} C_L e^{-2\pi q}. \quad (\text{OSP13})$$

The source mass-to-lower-density factor obeys the uniform finite bound

$$\frac{M_s}{\beta_s q^{2\ell_s}} = \sqrt{2} \Gamma(b) q^{-2\ell_s} \leq \sqrt{2\pi} \quad (s = 1, k). \quad (\text{OSP14})$$

To prove the inequality, use $\Gamma(b) = \sqrt{\pi} \prod_{j=0}^{2\ell_s-1} (j + 1/2) \leq \sqrt{\pi} b^{2\ell_s}$ and $b \leq q$. All source masses have thus been retained before taking this exact ratio.

32.2 A complete coefficient-space estimate for the inner interval

Let $p \in \mathbb{C}[y]$ have degree at most d , and let $0 < T \leq q/2$. Then

$$\sup_{|y| \leq T} |p(y)|^2 \leq \frac{(d+1)^2 10^{2d}}{q} \int_q^{2q} |p(u)|^2 du. \quad (\text{OSP15})$$

This controls every coefficient direction; no assumption on zeros, leading coefficient, or extremizers of p is made.

For proof define the Legendre polynomial $L_j(x) = (2^j j!)^{-1} (d/dx)^j (x^2 - 1)^j$. Its leading coefficient is $(2j)!/[2^j(j!)^2]$. Integrating by parts j times, with all lower boundary derivatives zero, proves orthogonality to lower powers and

$$\int_{-1}^1 L_j(x)^2 dx = \frac{(2j)!}{2^{2j}(j!)^2} \int_{-1}^1 (1-x^2)^j dx = \frac{2}{2j+1}.$$

For the middle integral set $x = 2t - 1$; repeated integration by parts gives $\int_0^1 t^j (1-t)^j dt = (j!)^2/(2j+1)!$. Consequently $\sqrt{2j+1} L_j(2z-3)$, $0 \leq j \leq d$, form an orthonormal basis for degree-at-most- d polynomials on $[1, 2]$. Leibniz's rule also gives the explicit identity

$$L_j(x) = 2^{-j} \sum_{h=0}^j \binom{j}{h}^2 (x-1)^{j-h} (x+1)^h.$$

For $|x| \leq 4$ its modulus is at most $(5/2)^j \sum_h \binom{j}{h}^2 \leq (5/2)^j 4^j = 10^j$; the sum inequality follows by comparing with $(\sum_h \binom{j}{h})^2 = 4^j$. Expanding $p(qz)$ in the displayed orthonormal basis and applying Cauchy-Schwarz gives, for $|z| \leq T/q \leq 1/2$,

$$|p(qz)|^2 \leq \sum_{j=0}^d (2j+1) 10^{2j} \int_1^2 |p(qv)|^2 dv \leq (d+1)^2 10^{2d} \int_1^2 |p(qv)|^2 dv.$$

Here $|2z-3| \leq 4$ and $\sum_{j=0}^d (2j+1) = (d+1)^2$. Changing variables proves OSP15, including $d = 0$.

32.3 The exact original-root map and a finite full-norm comparison

In this subsection P is either of the complete polynomials $P_k, P_{k'}$ in OSP3, and $Q = \deg P$ is respectively q, q' . For $d \geq 0$ retain the two injections into the same original polynomial Hilbert space

$$\begin{aligned} \mathcal{J}_P : \mathbb{C}[y]_{\leq d} &\longrightarrow \mathbb{C}[y]_{\leq Q+d}, & p &\longmapsto Pp, \\ \mathcal{J}_{0,Q} : \mathbb{C}[y]_{\leq d} &\longrightarrow \mathbb{C}[y]_{\leq Q+d}, & p &\longmapsto y^Q p. \end{aligned} \tag{OSP16}$$

Their codomain norm is precisely m_s in OSP2. If $P(y) = \sum_{a=0}^Q P_a y^a$, set $P_a = 0$ outside this range. In increasing coefficient frames their matrices are

$$(\mathcal{J}_P)_{rj} = P_{r-j}, \quad (\mathcal{J}_{0,Q})_{rj} = \mathbf{1}_{\{r=Q+j\}}, \quad 0 \leq r \leq Q+d, \quad 0 \leq j \leq d. \tag{OSP17}$$

Thus the exact isomorphism between their images is $y^Q p \mapsto Pp$, with inverse $Pp \mapsto y^Q p$. It preserves the original coefficient vector p . The first matrix contains every actual root coefficient; none is replaced inside that map. Returning to S or S' multiplies the relevant relation columns by the actual phase i^Q and applies the triangular coordinate pullback with diagonal entries i^j . Each has the stated exact inverse and determinant of modulus one.

Suppose

$$R < T \leq q/2. \tag{OSP18}$$

Define the finite positive constants

$$\begin{aligned} \eta_{Q,T} &= \frac{QR^2}{T^2 - R^2}, \\ \kappa_{Q,d,T,s} &= \frac{M_s}{\beta_s q^{2\ell_s} C_L} \frac{(d+1)^2}{q} e^{2\pi q} 10^{2d} \left(\frac{T^2 + R^2}{q^2} \right)^Q, \\ e_{Q,d,T,s} &= \eta_{Q,T} + \log(1 + \kappa_{Q,d,T,s}). \end{aligned} \tag{OSP19}$$

Then every $p \in \mathbb{C}[y]_{\leq d}$ satisfies

$$e^{-e_{Q,d,T,s}} \int_{\mathbb{R}} |y|^{2Q} |p(y)|^2 dm_s(y) \leq \int_{\mathbb{R}} |P(y)p(y)|^2 dm_s(y) \leq e^{e_{Q,d,T,s}} \int_{\mathbb{R}} |y|^{2Q} |p(y)|^2 dm_s(y). \quad (\text{OSP20})$$

Here is a proof retaining the entire inner region. Pair the actual roots as $\omega, -\omega$. On $|y| \geq T$, each paired quotient is $1 - \omega^2/y^2$, so

$$\left| \log \frac{|P(y)|^2}{|y|^{2Q}} \right| \leq -Q \log(1 - R^2/T^2) \leq \frac{QR^2}{T^2 - R^2} = \eta_{Q,T}. \quad (\text{OSP21})$$

The first inequality uses $1 - u \leq |1 - z| \leq 1 + u$ for $|z| \leq u < 1$ on each of $Q/2$ pairs. The second follows by integrating $1/(1 - u) \leq 1/(1 - u_0)$ on $0 \leq u \leq u_0 = R^2/T^2$. For every real y , the same pairing gives $|P(y)|^2 \leq (y^2 + R^2)^Q$. Write

$$A_B = \int_q^{2q} |y|^{2Q} |p(y)|^2 dm_s(y).$$

For nonzero p this number is positive, and OSP13 gives $A_B \geq q^{2Q} \beta_s q^{2\ell_s} C_L e^{-2\pi q} \int_q^{2q} |p(y)|^2 dy$. OSP15 and the exact full mass OSP8 imply

$$\begin{aligned} \int_{|y| < T} |P(y)p(y)|^2 dm_s(y) &\leq M_s(T^2 + R^2)^Q \sup_{|y| \leq T} |p(y)|^2 \leq \kappa_{Q,d,T,s} A_B, \\ \int_{|y| < T} |y|^{2Q} |p(y)|^2 dm_s(y) &\leq M_s T^{2Q} \sup_{|y| \leq T} |p(y)|^2 \leq \kappa_{Q,d,T,s} A_B. \end{aligned} \quad (\text{OSP22})$$

Let A_{out} and B_{out} be the integrals outside $(-T, T)$ with factors $|y|^{2Q}$ and $|P|^2$, respectively. Since $[q, 2q]$ is outside that interval, $A_B \leq A_{\text{out}}$. OSP21 gives $e^{-\eta} A_{\text{out}} \leq B_{\text{out}} \leq e^{\eta} A_{\text{out}}$. OSP22 bounds both inner contributions by κA_{out} . Therefore the full reference norm is at most $(1 + \kappa) A_{\text{out}}$ and at least A_{out} , while the full actual norm is at least $e^{-\eta} A_{\text{out}}$ and at most $(e^{\eta} + \kappa) A_{\text{out}} \leq e^{\eta} (1 + \kappa) A_{\text{out}}$. These four inequalities prove OSP20. The zero polynomial satisfies it directly.

32.4 Uniform bounded logarithmic cost at every required dimension

Choose the fixed number and cutoff

$$\epsilon = 2^{-32}, \quad T = \epsilon q,$$

and impose the explicit finite conditions

$$k \geq 29, \quad k \sqrt{\delta^2 + \gamma^2} \leq \frac{\epsilon q}{2}. \quad (\text{OSP23})$$

In particular $T \geq 2R$, as required for a uniform margin in OSP21. These conditions hold eventually for every fixed original δ, γ ; the sufficient bound $k \geq \max\{29, 2\sqrt{\delta^2 + \gamma^2}/\epsilon\}$ follows from $q \geq k^2$. The first condition implies $q' \geq q/2$: $2(k - 7)^2 - (k + 1)^2 = k^2 - 30k + 97$ is positive at $k = 29$ and is increasing thereafter.

For $Q \in \{q', q\}$ and every $0 \leq d \leq 2q$, set

$$\begin{aligned} K_q &= \frac{\sqrt{2\pi}}{C_L} \frac{(2q + 1)^2}{q} e^{-2q}, \\ \mathcal{E}_k &= \frac{qR^2}{\epsilon^2 q^2 - R^2} + \log(1 + K_q). \end{aligned} \quad (\text{OSP24})$$

Then OSP19 satisfies $e_{Q,d,T,s} \leq \mathcal{E}_k$ for both original source orders. To verify this uniformly, OSP14, $T^2 + R^2 \leq (5/4)\epsilon^2 q^2$, $Q \geq q/2$ and $d \leq 2q$ give

$$\kappa_{Q,d,T,s} \leq \frac{\sqrt{2\pi}}{C_L} \frac{(2q + 1)^2}{q} \exp\left(q\left\{2\pi + 4 \log 10 + \frac{1}{2} \log(5/4) + \log \epsilon\right\}\right). \quad (\text{OSP25})$$

The expression in braces is less than -2 . Indeed $\pi < 4$, $\log 10 < 4 \log 2$, $\log(5/4) < 1/4$, and $\log 2 > 2/3$ give an upper bound $8 - 16 \log 2 + 1/8 < -61/24 < -2$. For completeness, $\pi < 4$ follows from $\pi/4 = \int_0^1 (1+x^2)^{-1} dx < 1$; and the substitution $x = (1+t)/(1-t)$ in $\log 2 = \int_1^2 dx/x$ gives $\log 2 = 2 \int_0^{1/3} (1-t^2)^{-1} dt > 2/3$. The logarithm bound follows by integrating $1/(1+x) < 1$. This proves OSP25 with K_q as stated. Also

$$0 \leq \mathcal{E}_k \leq \frac{4(\delta^2 + \gamma^2)}{3\epsilon^2} + \log(1 + K_q). \quad (\text{OSP26})$$

Here $R^2 \leq \epsilon^2 q^2/4$ and $R^2/q \leq \delta^2 + \gamma^2$. Thus the logarithmic norm cost is bounded at fixed original parameters, uniformly over all these coefficient degrees and both unchanged source orders.

32.5 Every relation determinant and the exact outer quotient

For a nonzero polynomial P and $a \geq 0$ define in the unchanged monomial frame

$$\begin{aligned} D_a(P; s) &= \det \left[\int_{\mathbb{R}} y^{i+j} |P(y)|^2 dm_s(y) \right]_{0 \leq i, j < a}, \\ \mathcal{D}_a(Q; s) &= \det \left[\int_{\mathbb{R}} y^{2Q+i+j} dm_s(y) \right]_{0 \leq i, j < a}, \quad D_0 = \mathcal{D}_0 = 1. \end{aligned} \quad (\text{OSP27})$$

All nonempty Grams are positive definite, because a nonzero polynomial has positive squared modulus on an open subinterval of the strictly positive original measure. In the notation of OSP17 their exact matrices are $J_P^* H_{Q+a-1}^{(s)} J_P$ and $J_{0,Q}^* H_{Q+a-1}^{(s)} J_{0,Q}$, where $H_{Q+a-1}^{(s)}$ is the complete original moment Gram. Consequently OSP20–26 prove for $1 \leq a \leq 2q+1$

$$|\log D_a(P; s) - \log \mathcal{D}_a(Q; s)| \leq a \mathcal{E}_k, \quad P = P_k \text{ or } P_{k'}. \quad (\text{OSP28})$$

To see the determinant step exactly, conjugate the actual Gram by the inverse positive square root of the reference Gram. OSP20 places every eigenvalue of the resulting positive Hermitian matrix in $[e^{-\mathcal{E}_k}, e^{\mathcal{E}_k}]$. The logarithm of its determinant is the sum of its a eigenvalue logarithms, proving OSP28. This is the full exterior determinant estimate, including every polynomial coefficient direction. The estimate also holds at $a = 0$ with both sides zero.

For $t \in \{0, 1, q, q+1\}$, let $L = t + \Delta - 1$. The actual quotient has source $\mathbb{C}[y]_{\leq L}$ in measure $|P_{k'}(y)|^2 dm_s(y)$, target the increasing remainder frame of $\mathbb{C}[y]/(P_{\partial,k})$, and its minimum Gram $Q_{\partial,L}^{(s)}$. Define a comparison quotient with source $\mathbb{C}[y]_{\leq L}$ in measure $|y|^{2q'} dm_s(y)$, target the increasing remainder frame of $\mathbb{C}[y]/(y^\Delta)$, and minimum Gram $Q_{0,L}^{(s)}$. Its source measure still has order s and the full mass factor in OSP2.

For precision, the exact map relating the coefficient presentations is the degree-preserving isomorphism

$$\mathcal{U}_L(y^j) = \begin{cases} y^j, & 0 \leq j < \Delta, \\ P_{\partial,k}(y) y^{j-\Delta}, & \Delta \leq j \leq L. \end{cases} \quad (\text{OSP29})$$

Its coefficient matrix has diagonal entries one and no terms of degree greater than that of its input column. It is therefore invertible with determinant one. If $R_{\partial,L}$ is division remainder modulo $P_{\partial,k}$ and $R_{0,L}$ is division remainder modulo y^Δ , then $R_{\partial,L} \mathcal{U}_L = R_{0,L}$: the first Δ basis vectors map to their stated remainders, and every later vector maps into the actual ideal. This proves the precise morphism between the quotient coefficient spaces; no equality of their metrics is assumed.

Using the original monic relation frames on the two sources gives the exact identities

$$\det Q_{\partial,t+\Delta-1}^{(s)} = \frac{D_{t+\Delta}(P_{k'}; s)}{D_t(P_k; s)}, \quad \det Q_{0,t+\Delta-1}^{(s)} = \frac{\mathcal{D}_{t+\Delta}(q'; s)}{\mathcal{D}_t(q; s)}. \quad (\text{OSP30})$$

For proof, append to $1, y, \dots, y^{\Delta-1}$ the ideal frame $P_{\partial,k}, yP_{\partial,k}, \dots, y^{t-1}P_{\partial,k}$. Its source coefficient determinant is one by OSP29. The relation Gram is $D_t(P_k; s)$ because $P_{k'}P_{\partial,k} = P_k$. Completing

the square in the ideal coordinates gives the minimum quotient Gram as the Schur complement; the determinant of the full source Gram is the product of that determinant and the relation determinant. This proves the first identity. For the second repeat the same calculation with y^Δ : $y^{q'} y^\Delta = y^q$, with the original source order s . The same proof includes $t = 0$ using the empty relation block.

OSP28 applies to every index here, since $t + \Delta \leq q + \Delta + 1 \leq 2q + 1$. Define the exact endpoint difference $\varepsilon_t^{(s)}$ by

$$\log \det Q_{\partial, t+\Delta-1}^{(s)} = \log \mathcal{D}_{t+\Delta}(q'; s) - \log \mathcal{D}_t(q; s) + \varepsilon_t^{(s)}.$$

Then, separately at each of the four endpoints,

$$|\varepsilon_t^{(s)}| \leq (2t + \Delta) \mathcal{E}_k. \quad (\text{OSP31})$$

In particular set

$$\begin{aligned} \mathcal{F}_{0,k}^{(s)} = & \log \mathcal{D}_\Delta(q'; s) + \log \mathcal{D}_{\Delta+1}(q'; s) - \log \mathcal{D}_{q+\Delta}(q'; s) - \log \mathcal{D}_{q+\Delta+1}(q'; s) \\ & - \log \mathcal{D}_1(q; s) + \log \mathcal{D}_q(q; s) + \log \mathcal{D}_{q+1}(q; s). \end{aligned} \quad (\text{OSP32})$$

The actual four-endpoint quantity of CTV13 therefore obeys the proved finite estimate

$$\boxed{\left| \mathcal{F}_{\partial,k}^{(s)} - \mathcal{F}_{0,k}^{(s)} \right| \leq 4(q + \Delta + 1) \mathcal{E}_k = O_{\delta, \gamma}(q) = o(kq), \quad s = 1, k.} \quad (\text{OSP33})$$

Indeed its exact error is $\varepsilon_0 + \varepsilon_1 - \varepsilon_q - \varepsilon_{q+1}$; summing OSP31 gives $\Delta + (\Delta + 2) + (2q + \Delta) + (2q + \Delta + 2) = 4(q + \Delta + 1)$. OSP26 and $\Delta < q$ prove the last two estimates. This calculation uses all four source degrees $\Delta - 1, \Delta, q + \Delta - 1, q + \Delta$ and every source and relation determinant in OSP30.

32.6 The resulting same-source moment target

The moments in OSP27 are completely specified by the original analytic identity OSP9:

$$\mu_j^{(s)} = \int y^j dm_s(y) = M_s \frac{d^j}{dz^j} (\cos z)^{-s/2} \Big|_{z=0}, \quad \mathcal{D}_a(Q; s) = \det[\mu_{2Q+i+j}^{(s)}]_{0 \leq i, j < a}. \quad (\text{OSP34})$$

The differentiated integral is justified by OSP7, or directly by the finite exponential moments on a slightly larger interval. Odd moments vanish because the density is even. Reordering rows and columns together by parity therefore proves the exact further factorization

$$\mathcal{D}_a(Q; s) = \det[\mu_{2(Q+i+j)}^{(s)}]_{0 \leq i, j < \lceil a/2 \rceil} \det[\mu_{2(Q+1+i+j)}^{(s)}]_{0 \leq i, j < \lfloor a/2 \rfloor}. \quad (\text{OSP35})$$

There is no permutation sign, since the same permutation is applied to both rows and columns. Empty factors are one. In OSP32 the coefficient of $\log M_s$ is $\Delta + (\Delta + 1) - (q + \Delta) - (q + \Delta + 1) - 1 + q + (q + 1) = 0$. This is an exact cancellation only after all original moments and determinant dimensions have been retained.

Equation OSP33 proves an error-controlled comparison with the exact finite determinant difference OSP32. It does not obtain a first variation by differentiating a leading q^2 law. In particular no numerical value of the limit of $\mathcal{F}_{0,k}^{(s)}/(kq)$ is asserted by this comparison.

33 The outer-root and actual residual bounds in the original arithmetic return

Retain the complete original data of CEP, CTV and ROQ. In particular $m = 1$, $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $q' = (k - 7)^2$, $\Delta = 16k - 48$, $r = 8k - 16$ and $j = 8k - 32$. The quartet, full amplitude unit, period and branch are the specified original ones on the proved five-orbit locus. The source orders remain $s \in \{1, k\}$. Every matrix below is the actual matrix in those chapters, with its original mass and coefficient frame.

33.1 The two finite error endpoints retain their signs

Write

$$\mathcal{N}_+ = \{q - 1, q\}, \quad \mathcal{N}_- = \{2q - 1, 2q\}, \quad \epsilon_{A,k}^{(s)} = 2 \sum_{N \in \mathcal{N}_+ \cup \mathcal{N}_-} E_N^{(s)}. \quad (\text{OAR1})$$

Here $E_N^{(s)}$ is the explicit full conductor constant of CEP35. The nonnegative constants $L_N^{\text{tail}}, U_N^{\text{tail}}$ are exactly CTV18 in the same source order and with the original q', δ, γ . Define

$$\begin{aligned} a_{k,s}^- &= v \left(\sum_{N \in \mathcal{N}_+} L_N^{\text{tail}} + \sum_{N \in \mathcal{N}_-} U_N^{\text{tail}} \right), \\ a_{k,s}^+ &= v \left(\sum_{N \in \mathcal{N}_+} U_N^{\text{tail}} + \sum_{N \in \mathcal{N}_-} L_N^{\text{tail}} \right), \\ e_{k,s}^- &= \epsilon_{A,k}^{(s)} + a_{k,s}^-, \quad e_{k,s}^+ = \epsilon_{A,k}^{(s)} + a_{k,s}^+. \end{aligned} \quad (\text{OAR2})$$

These are finite quantities for every original period in the stated domain. For each fixed original period, both $e_{k,s}^\pm = o(kq)$. Indeed CEP36 proves this for $\epsilon_{A,k}^{(s)}$ and CTV18–20 proves it for each of the four tail contributions, uniformly over the possible indices $0 \leq v \leq 80$ and both original source orders. When $v = 0$, both $a_{k,s}^\pm$ are exactly zero.

Set $\Phi_{k,s} = F_{\Xi,k}^{(s)}$ and retain the full outer quotient $\mathcal{F}_{\partial,k}^{(s)}$ of CTV11–13. The exact identity is

$$\begin{aligned} F_K^{(s)} &= \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s} + \mathfrak{e}_{k,s}, \\ \mathfrak{e}_{k,s} &= \mathcal{E}_{A,k}^{(s)} + \sum_{N \in \mathcal{N}_+} \mathcal{R}_{N,v}^{(s)} - \sum_{N \in \mathcal{N}_-} \mathcal{R}_{N,v}^{(s)}, \\ -e_{k,s}^- &\leq \mathfrak{e}_{k,s} \leq e_{k,s}^+. \end{aligned} \quad (\text{OAR3})$$

The equality is CTV22, keeping its original two errors. CTV19 says $-vL_N^{\text{tail}} \leq \mathcal{R}_{N,v}^{(s)} \leq vU_N^{\text{tail}}$. At a positive endpoint this contributes lower bound $-vL_N^{\text{tail}}$ and upper bound vU_N^{tail} ; at a negative endpoint it contributes lower bound $-vU_N^{\text{tail}}$ and upper bound vL_N^{tail} . Adding these four inequalities and the bounds $-\epsilon_{A,k}^{(s)} \leq \mathcal{E}_{A,k}^{(s)} \leq \epsilon_{A,k}^{(s)}$ proves each sign of OAR3.

33.2 Original positive increments give explicit kernel caps

The full value of $\Phi_{k,s}$ is ROQ19's finite sum

$$\Phi_{k,s} = \log(1 + \eta_{q-1}) + 2 \sum_{N=q}^{2q-2} \log(1 + \eta_N) + \log(1 + \eta_{2q-1}), \quad \lambda_s = \log(1 + \eta_{q-1}) \geq 0. \quad (\text{OAR4})$$

Every η_N here is ROQ17's original constrained column, including the complete conductor–observation cross covariance and its positive denominator. No new observation is selected. ROQ21 gives, in the original AKS notation,

$$\lambda_s \leq \Phi_{k,s} \leq V_s := j(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - \lambda_s. \quad (\text{OAR5})$$

Retain the complete earlier kernel interval $L_s \leq F_K^{(s)} \leq U_s$ of AKJ2/MRI4e, with its pointwise minimum of the KAF and AKS bounds. Define two explicit quantities

$$\begin{aligned} L_s^\partial &= \max \left\{ L_s, \mathcal{F}_{\partial,k}^{(s)} - e_{k,s}^- - V_s \right\}, \\ U_s^\partial &= \min \left\{ U_s, \mathcal{F}_{\partial,k}^{(s)} + e_{k,s}^+ - \lambda_s \right\}. \end{aligned} \quad (\text{OAR6})$$

Then

$$\begin{aligned} L_s^\partial &\leq F_K^{(s)} \leq U_s^\partial, \\ \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s} - e_{k,s}^- &\leq F_K^{(s)} \leq \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s} + e_{k,s}^+. \end{aligned} \quad (\text{OAR7})$$

For the first line substitute both ends of OAR5 into OAR3 and intersect with the retained interval. The second line substitutes the full finite value OAR4 into OAR3. Intersecting these intervals preserves every bound. Both contain the actual value, so their intersection is nonempty. The dimension j is used only in the proved finite comparison OAR5. It assigns no leading determinant. Each additional evaluated positive term of OAR4 can be included in the upper subtraction in OAR6 by the same inequality.

33.3 The correlated arithmetic kernel and boundary

Keep the original arithmetic total $\mathcal{B}_k^{\text{ar}}$, the scalar δ_k^σ and the four source-difference constants $C_K^\pm, C_B^\pm$ of MRI4f. They satisfy, for the actual $d = F_K^{\text{ar}} - F_K^{(1)}$,

$$\max\{-C_K^-, \delta_k^\sigma - C_B^+\} \leq d \leq \min\{C_K^+, \delta_k^\sigma + C_B^-\}. \quad (\text{OAR8})$$

This is the pair of original kernel bounds together with the boundary bounds, transferred by $d + (F_B^{\text{ar}} - F_B^{(1)}) = \delta_k^\sigma$. It retains their common scalar and signs. Define

$$\begin{aligned} L_{\text{ar}}^\partial &= \max\{L_{\text{ar}}, L_1^\partial - C_K^-, L_1^\partial + \delta_k^\sigma - C_B^+\}, \\ U_{\text{ar}}^\partial &= \min\{U_{\text{ar}}, U_1^\partial + C_K^+, U_1^\partial + \delta_k^\sigma + C_B^-\}. \end{aligned} \quad (\text{OAR9})$$

Here $[L_{\text{ar}}, U_{\text{ar}}]$ is the complete interval of MRI4f. Adding OAR8 to the source-1 first line of OAR7 proves

$$\begin{aligned} L_{\text{ar}}^\partial &\leq F_K^{\text{ar}} \leq U_{\text{ar}}^\partial, \\ \mathcal{B}_k^{\text{ar}} - U_{\text{ar}}^\partial &\leq F_B^{\text{ar}} \leq \mathcal{B}_k^{\text{ar}} - L_{\text{ar}}^\partial. \end{aligned} \quad (\text{OAR10})$$

The second line subtracts the first from the exact original identity $F_K^{\text{ar}} + F_B^{\text{ar}} = \mathcal{B}_k^{\text{ar}}$. It uses the same arithmetic total at both endpoints. If the second line of OAR7 is used instead, exactly the same addition and subtraction yield the further interval containing the full residual sum. Thus every exact positive residual subtraction is available to the original arithmetic kernel and its correlated boundary.

33.4 The signed mixed return with its complete finite error

Write $\mathfrak{S}_k^{\text{mix}} = F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)}$. The original BRI6d identity is $\mathfrak{S}_k^{\text{mix}} = F_K^{(k)} + \mathcal{H}_k + \delta_k^\sigma$. Consequently OAR3 gives the exact current expression

$$\begin{aligned} \mathfrak{S}_k^{\text{mix}} &= \mathcal{F}_{\partial,k}^{(k)} - \Phi_{k,k} + \mathcal{H}_k + \delta_k^\sigma + \mathfrak{e}_{k,k}, \\ -e_{k,k}^- &\leq \mathfrak{e}_{k,k} \leq e_{k,k}^+, \quad e_{k,k}^\pm = o(kq). \end{aligned} \quad (\text{OAR11})$$

This substitution proves every sign, keeping the original Gamma return and arithmetic scalar. Addition of their unchanged values to the source- k bounds OAR7 yields the corresponding finite intervals for this same signed quantity. The full original four-endpoint metrics, outer-root polynomial, weighted source, period-dependent certificates and graph denominator remain in OAR11 through their proved maps CTV and ROQ. The next leading-scale calculation is the difference of these two explicit quotient returns.

34 The same-source Gamma moment comparison in the arithmetic return

Retain every original parameter, matrix and error in OAR1–11. The complete OSP comparison supplies an additional bound directly on the actual outer-root quotient. Set $\epsilon = 2^{-32}$ and retain its explicit range

$$k \geq 29, \quad k\sqrt{\delta^2 + \gamma^2} \leq \epsilon(k+1)^2/2. \quad (\text{PMR1})$$

This range contains all sufficiently large original $k = 4l + 1$ for each fixed original quartet: $k \geq \max\{29, 2\sqrt{\delta^2 + \gamma^2}/\epsilon\}$ is sufficient because $(k+1)^2 \geq k^2$. For the remaining finite values the full earlier OAR estimates remain valid without using this comparison.

Keep OSP24's nonnegative $\mathcal{E}_k$ and put

$$c_k^\circ = 4(q + \Delta + 1)\mathcal{E}_k, \quad e_{k,s}^{0,-} = e_{k,s}^- + c_k^\circ, \quad e_{k,s}^{0,+} = e_{k,s}^+ + c_k^\circ. \quad (\text{PMR2})$$

OSP26 and OSP33 prove $c_k^\circ = O_{\delta,\gamma}(q) = o(kq)$ on PMR1, uniformly for the unchanged source orders $s = 1, k$. The actual outer quotient differs from OSP32's exact seven original Gamma moment determinants by at most c_k° . Consequently the full identity and signed bounds are

$$\begin{aligned} F_K^{(s)} &= \mathcal{F}_{0,k}^{(s)} - \Phi_{k,s} + \mathfrak{e}_{k,s}^0, \\ \mathfrak{e}_{k,s}^0 &= \mathfrak{e}_{k,s} + \mathcal{F}_{\partial,k}^{(s)} - \mathcal{F}_{0,k}^{(s)}, \\ -e_{k,s}^{0,-} &\leq \mathfrak{e}_{k,s}^0 \leq e_{k,s}^{0,+}, \quad e_{k,s}^{0,\pm} = o(kq). \end{aligned} \quad (\text{PMR3})$$

The first two equalities are OAR3 with the displayed exact difference added and subtracted. The two bounds follow by adding the interval $[-c_k^\circ, c_k^\circ]$ of OSP33 to the unequal OAR3 endpoints. No metric equality is substituted for the proved OSP coefficient maps and determinant comparison. The residual sum $\Phi_{k,s}$ retains all actual period data.

Let V_s and λ_s be OAR4–5's exact residual upper bound and first positive increment. Define

$$\begin{aligned} L_s^0 &= \max\{L_s^\partial, \mathcal{F}_{0,k}^{(s)} - e_{k,s}^{0,-} - V_s\}, \\ U_s^0 &= \min\{U_s^\partial, \mathcal{F}_{0,k}^{(s)} + e_{k,s}^{0,+} - \lambda_s\}. \end{aligned} \quad (\text{PMR4})$$

Then $L_s^0 \leq F_K^{(s)} \leq U_s^0$. To prove this, insert $\lambda_s \leq \Phi_{k,s} \leq V_s$ into PMR3 and intersect with OAR7. Using the full finite sum for $\Phi_{k,s}$ in PMR3 gives the sharper interval centred at $\mathcal{F}_{0,k}^{(s)} - \Phi_{k,s}$. Both retain the same unequal finite errors of PMR2.

The exact arithmetic correlations of OAR8 give

$$\begin{aligned} L_{\text{ar}}^0 &= \max\{L_{\text{ar}}^\partial, L_1^0 - C_K^-, L_1^0 + \delta_k^\sigma - C_B^+\}, \\ U_{\text{ar}}^0 &= \min\{U_{\text{ar}}^\partial, U_1^0 + C_K^+, U_1^0 + \delta_k^\sigma + C_B^-\}. \end{aligned} \quad (\text{PMR5})$$

Adding those correlations to PMR4 at $s = 1$ proves $L_{\text{ar}}^0 \leq F_K^{\text{ar}} \leq U_{\text{ar}}^0$. Subtracting from the exact same arithmetic total gives

$$\mathcal{B}_k^{\text{ar}} - U_{\text{ar}}^0 \leq F_B^{\text{ar}} \leq \mathcal{B}_k^{\text{ar}} - L_{\text{ar}}^0. \quad (\text{PMR6})$$

Thus the actual kernel and boundary retain their common scalar and both original source comparisons.

Finally the original signed-return identity BRI6d and PMR3 at source order k give

$$\begin{aligned} F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)} &= \mathcal{F}_{0,k}^{(k)} - \Phi_{k,k} + \mathcal{H}_k + \delta_k^\sigma + \mathfrak{e}_{k,k}^0, \\ -e_{k,k}^{0,-} &\leq \mathfrak{e}_{k,k}^0 \leq e_{k,k}^{0,+}. \end{aligned} \quad (\text{PMR7})$$

This follows by substituting into $F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)} = F_K^{(k)} + \mathcal{H}_k + \delta_k^\sigma$; every sign is retained. The entire original same-source moment expression is OSP32–35. The next asymptotic calculation must evaluate it with the actual residual sum and an error small compared with kq .

The exact finite endpoints under this comparison. The interval transport in PMR4–6 preserves the already retained outer-root endpoints exactly:

$$L_s^0 = L_s^\partial, \quad U_s^0 = U_s^\partial, \quad L_{\text{ar}}^0 = L_{\text{ar}}^\partial, \quad U_{\text{ar}}^0 = U_{\text{ar}}^\partial. \quad (\text{PMR8})$$

For proof, OSP33 gives $\mathcal{F}_{0,k}^{(s)} - c_k^\circ \leq \mathcal{F}_{\partial,k}^{(s)} \leq \mathcal{F}_{0,k}^{(s)} + c_k^\circ$. The new lower candidate in PMR4 is therefore at most $\mathcal{F}_{\partial,k}^{(s)} - e_{k,s}^- - V_s$, which is at most L_s^∂ by OAR6. The new upper candidate is at least $\mathcal{F}_{\partial,k}^{(s)} + e_{k,s}^+ - \lambda_s$, which is at least U_s^∂ . Taking the displayed maximum and minimum proves the source equalities. Substitution in PMR5 and the defining maxima and minima OAR9 proves the arithmetic equalities; subtracting from the same total proves equality of the boundary endpoints as well. Thus this comparison supplies the exact same-source moment target and controlled error; a further numerical improvement must use sharper evaluated moment or residual information. The preceding ROQ/OAR bounds remain the proved finite intervals throughout this transport.

35 Exact equilibrium for the square-root logarithmic potential

Throughout this section $\alpha > 0$ and $\beta > 0$. On $(0, \infty)$ set

$$V_{\alpha,\beta}(x) = \beta\sqrt{x} - \alpha \log x, \quad \mathcal{E}_{\alpha,\beta}(\mu) = \int V_{\alpha,\beta}(x) d\mu(x) - \iint \log|x-y| d\mu(x)d\mu(y). \quad (\text{EIQ1})$$

The measures considered initially are probability measures for which $\int V_{\alpha,\beta} d\mu$ is finite. Because this potential is bounded below and tends to $+\infty$ at both ends, this implies $\int(\sqrt{x} + |\log x|) d\mu < \infty$. The positive part of $\log|x-y|$ is then integrable, by $\log^+|x-y| \leq \log(1+x) + \log(1+y)$. Thus the energy is well defined with value in $\mathbb{R} \cup \{+\infty\}$. All comparisons below also apply to the usual extended energy defined by integrating the kernel $\frac{1}{2}V_{\alpha,\beta}(x) + \frac{1}{2}V_{\alpha,\beta}(y) - \log|x-y|$: the kernel is bounded below, and finite energy for this kernel implies the same moment conditions. To see the latter assertion without cancellation, put $g(x) = \frac{1}{2}V_{\alpha,\beta}(x) - \log(1+x)$. The kernel is at least $g(x) + g(y)$; g is bounded below and controls positive constant multiples of both $\sqrt{x}$ near infinity and $|\log x|$ near zero.

35.1 Existence and uniqueness of the two endpoints

For $0 < k < 1$ define the complete elliptic integrals by their actual real integrals

$$K(k) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}}, \quad E(k) = \int_0^{\pi/2} \sqrt{1-k^2 \sin^2 \theta} d\theta. \quad (\text{EIQ2})$$

There is exactly one $k \in (0, 1)$ satisfying

$$\frac{\sqrt{1-k^2}K(k)}{E(k)} = \frac{\alpha}{\alpha+2}. \quad (\text{EIQ3})$$

For this k put

$$r = \sqrt{1-k^2}, \quad v = \frac{(\alpha+2)\pi}{\beta E(k)}, \quad u = rv, \quad a = u^2, \quad b = v^2. \quad (\text{EIQ4})$$

In particular $0 < a < b < \infty$, and the endpoint equations, with all constants retained, are

$$uK(k) = \frac{\alpha\pi}{\beta}, \quad vE(k) = \frac{(\alpha+2)\pi}{\beta}, \quad k^2 = 1 - \frac{u^2}{v^2}. \quad (\text{EIQ5})$$

Here is a proof of the assertion of uniqueness. Differentiation of the integrals in (EIQ2), and integration of the derivative of $\sin \theta \cos \theta / \sqrt{1-k^2 \sin^2 \theta}$ between its two zero endpoint values, give

$$E'(k) = \frac{E(k) - K(k)}{k}, \quad K'(k) = \frac{E(k)}{k(1-k^2)} - \frac{K(k)}{k}.$$

For completeness the second identity is equivalent, after differentiation under the integral sign, to

$$\int_0^{\pi/2} \left\{ \frac{k^2 \sin^2 \theta}{(1-k^2 \sin^2 \theta)^{3/2}} - \frac{1-k^2 \sin^2 \theta}{(1-k^2)\sqrt{1-k^2 \sin^2 \theta}} + \frac{1}{\sqrt{1-k^2 \sin^2 \theta}} \right\} d\theta = 0;$$

the expression in braces is $-k^2(1-k^2)^{-1} d(\sin \theta \cos \theta / \sqrt{1-k^2 \sin^2 \theta})/d\theta$. With $t = E(k)/K(k)$, the integral definitions show strictly $r^2 < t < 1$. Consequently, if $f(k) = rK(k)/E(k)$, then

$$\frac{kf'(k)}{f(k)} = \frac{t}{r^2} + \frac{1}{t} - 1 - \frac{1}{r^2} = \frac{(t-1)(t-r^2)}{r^2 t} < 0. \quad (\text{EIQ6})$$

At $k = 0$ the continuous limiting value is $f(0) = 1$. As $k \uparrow 1$, the integrand of $rK(k)$ tends to zero at every $\theta < \pi/2$ and is bounded by 1, while $E(k) \rightarrow 1$ by dominated convergence. Hence

$f(k) \rightarrow 0$. This proves (EIQ3), including existence. It also proves smooth dependence of k, u, v, a, b on (α, β) in $(0, \infty)^2$, by the nonzero derivative in (EIQ6).

Let $m = (a + b)/2$, $R_s = \sqrt{(a + s)(b + s)}$ for $s \geq 0$, and $R_0 = uv$. Substitution $t = a \cos^2 \theta + b \sin^2 \theta$ gives, directly from (EIQ5),

$$\int_a^b \frac{V'_{\alpha, \beta}(t)}{\sqrt{(b-t)(t-a)}} dt = 0, \quad \int_a^b \frac{tV'_{\alpha, \beta}(t)}{\sqrt{(b-t)(t-a)}} dt = 2\pi. \quad (\text{EIQ7})$$

Indeed the three integrals with numerators $t^{-1/2}, t^{-1}, t^{1/2}$ are respectively $2K(k)/v$, $\pi/(uv)$, and $2vE(k)$, and $V'(t) = \beta/(2\sqrt{t}) - \alpha/t$.

35.2 A positive density and its exact Cauchy transform

The elementary Stieltjes integral, obtained by $s = xz^2$, is

$$\int_0^\infty \frac{ds}{\sqrt{s}(x+s)} = \frac{\pi}{\sqrt{x}} \quad (x > 0). \quad (\text{EIQ8})$$

Using this identity in (EIQ7) and applying Tonelli to its positive constituents gives

$$\frac{\alpha}{R_0} = \frac{\beta}{2\pi} \int_0^\infty \frac{ds}{\sqrt{s}R_s}, \quad \int_0^\infty \frac{1 - s/R_s}{\sqrt{s}} ds = 2vE(k). \quad (\text{EIQ9})$$

Both integrals converge; their integrands are $O(s^{-1/2})$ at zero and $O(s^{-3/2})$ at infinity. The second formula also follows by averaging $\sqrt{t} = \pi^{-1} \int_0^\infty s^{-1/2} t/(t+s) ds$ against the arcsine probability measure on $[a, b]$.

Define for every $x > 0$

$$\begin{aligned} h(x) &= \frac{\alpha}{xR_0} - \frac{\beta}{2\pi} \int_0^\infty \frac{ds}{\sqrt{s}(x+s)R_s} \\ &= \frac{\beta}{2\pi x} \int_0^\infty \frac{\sqrt{s}}{(x+s)R_s} ds > 0. \end{aligned} \quad (\text{EIQ10})$$

The second equality is exactly (EIQ9), with no change of the potential. All integrals in (EIQ10) converge absolutely and locally uniformly in $x > 0$. The candidate density on the original positive half-line is

$$\rho_{\alpha, \beta}(x) = \begin{cases} \frac{\sqrt{(b-x)(x-a)}}{2\pi} h(x), & a < x < b, \\ 0, & x \notin (a, b). \end{cases} \quad (\text{EIQ11})$$

It is strictly positive on (a, b) and vanishes continuously with the square-root factor at each endpoint. Positivity here follows from the last integral in (EIQ10), not from a one-interval ansatz.

The following elementary transform computes its mass and resolvent. Choose

$$R(z) = \sqrt{(z-a)(z-b)}$$

on $\mathbb{C} \setminus [a, b]$ with $R(z)/z \rightarrow 1$ at infinity. Thus $R(x) > 0$ for $x > b$, $R(x) < 0$ for $x < a$, and $R(x + i0) = i\sqrt{(b-x)(x-a)}$ on (a, b) . Direct substitution $t = m + (b-a)\cos\theta/2$, followed by $w = \tan(\theta/2)$, proves

$$\frac{1}{\pi} \int_a^b \frac{dt}{(z-t)\sqrt{(b-t)(t-a)}} = \frac{1}{R(z)}.$$

For real $z > b$, the substitution reduces the integral to $\pi^{-1} \int_0^\infty 2dw/[(z-b) + (z-a)w^2]$; the identity elsewhere follows by analytic continuation. Polynomial division then gives

$$\frac{1}{\pi} \int_a^b \frac{\sqrt{(b-t)(t-a)}}{z-t} dt = z - m - R(z), \quad \frac{1}{\pi} \int_a^b \frac{\sqrt{(b-t)(t-a)}}{t+s} dt = m + s - R_s. \quad (\text{EIQ12})$$

For example the first identity follows from $(b-t)(t-a) = -(t-z)^2 + 2(m-z)(t-z) - (z-a)(z-b)$, the preceding arcsine transform, and the arcsine first moment m . Partial fractions in these two identities yield

$$H_s(z) := \frac{1}{\pi} \int_a^b \frac{\sqrt{(b-t)(t-a)}}{(t+s)(z-t)} dt = 1 - \frac{R_s + R(z)}{z+s}. \quad (\text{EIQ13})$$

The formula at $z = -s$ is interpreted by its removable limit.

The mass of (EIQ11), computed using (EIQ12), is

$$\begin{aligned} \int_a^b \rho_{\alpha,\beta}(t) dt &= \frac{\alpha}{2R_0}(m - R_0) - \frac{\beta}{4\pi} \int_0^\infty \frac{(m+s)/R_s - 1}{\sqrt{s}} ds \\ &= -\frac{\alpha}{2} + \frac{\beta}{4\pi} \int_0^\infty \frac{1-s/R_s}{\sqrt{s}} ds = -\frac{\alpha}{2} + \frac{\beta v E(k)}{2\pi} = 1. \end{aligned} \quad (\text{EIQ14})$$

In particular this is a probability density.

Define its Cauchy transform with the explicit sign convention $G(z) = \int_a^b \rho_{\alpha,\beta}(t)/(z-t) dt$. For z outside both $[a, b]$ and $(-\infty, 0]$, formulas (EIQ10) and (EIQ13) give

$$\begin{aligned} G(z) &= \frac{\alpha}{2R_0} H_0(z) - \frac{\beta}{4\pi} \int_0^\infty \frac{H_s(z)}{\sqrt{s} R_s} ds \\ &= -\frac{\alpha}{2z} + \frac{\beta}{4\sqrt{z}} - \frac{R(z)h(z)}{2} = \frac{V'_{\alpha,\beta}(z) - R(z)h(z)}{2}. \end{aligned} \quad (\text{EIQ15})$$

Here the positive square root is continued with its negative-real-axis cut, and $h(z)$ is the first integral formula of (EIQ10). Every integral exchange is absolutely convergent, uniformly on compact subsets of this domain: near zero the auxiliary integrand is $O(s^{-1/2})$, and near infinity the original integral defining H_s is $O(s^{-1})$. The constant terms on the second line cancel by (EIQ9), and (EIQ8) supplies the term $\beta/(4\sqrt{z})$.

35.3 The Euler equality, its strict exterior inequality, and uniqueness

Set

$$U(x) = V_{\alpha,\beta}(x) - 2 \int_a^b \log|x-t| \rho_{\alpha,\beta}(t) dt \quad (x > 0). \quad (\text{EIQ16})$$

This function is continuous on $(0, \infty)$. The integral is continuous because the density is bounded, compactly supported, and the integral of $|\log|x-t||$ over an interval of length ε tends to zero uniformly in its centre. In the interior of $[a, b]$, differentiation of the logarithmic potential is the principal-value integral $\text{PV} \int \rho(t)/(x-t) dt$. This follows by subtracting $\rho(x)$ on a symmetric interval about x ; the remaining numerator is $O(|t-x|)$ because the density is continuously differentiable locally in the interior. The real part of (EIQ15) on the upper edge of the interval is precisely this principal value. Indeed subtracting $\rho(x)$ in $\int \rho(t)/(x+i\varepsilon-t) dt$ gives the same limit, while the constant term has real part equal to its elementary principal value. Since $R(x+i0)$ is purely imaginary and $h(x)$ is real, (EIQ15) proves $U'(x) = 0$ on (a, b) . There is therefore a real constant ℓ such that

$$U(x) = \ell \quad (a \leq x \leq b). \quad (\text{EIQ17})$$

For positive x outside this interval, (EIQ15) instead gives the exact real derivative

$$U'(x) = R(x)h(x) \begin{cases} < 0, & 0 < x < a, \\ > 0, & x > b. \end{cases} \quad (\text{EIQ18})$$

Combining (EIQ17), continuity, and these signs proves

$$U(x) > \ell \quad \text{for } x \in (0, a) \cup (b, \infty). \quad (\text{EIQ19})$$

This verifies the entire exterior inequality on the original domain, including the interval next to zero.

Here is a complete positivity argument for the energy of the difference of two measures. If ν is a finite signed real measure of total mass zero for which $|\log|x-y||$ is integrable against $|\nu| \otimes |\nu|$, the scalar identity

$$-\log d = \frac{1}{2} \int_0^\infty \frac{e^{-td^2} - e^{-t}}{t} dt \quad (d > 0) \quad (\text{EIQ20})$$

follows by differentiating in d and evaluating at $d = 1$. One half of the integral over any finite subinterval of $(0, \infty)$ has the same sign as $-\log d$ and absolute value at most $|\log d|$. Consequently dominated convergence and the zero total mass give

$$-\iint \log|x-y| d\nu(x)d\nu(y) = \frac{1}{2} \int_0^\infty \frac{1}{t} \iint e^{-t(x-y)^2} d\nu(x)d\nu(y) dt \geq 0. \quad (\text{EIQ21})$$

The nonnegative sign is proved, rather than assumed, by the Gaussian Fourier identity

$$\iint e^{-t(x-y)^2} d\nu(x)d\nu(y) = \frac{1}{\sqrt{4\pi t}} \int_{\mathbb{R}} e^{-\xi^2/(4t)} |\widehat{\nu}(\xi)|^2 d\xi, \quad \widehat{\nu}(\xi) = \int e^{i\xi x} d\nu(x). \quad (\text{EIQ22})$$

The scalar Gaussian Fourier identity follows by completing the square in the Gaussian integral (or by differentiating its Fourier transform and its value at zero); its integration against the finite measures is justified by absolute integrability. Equality in (EIQ21) forces $\widehat{\nu} = 0$: the right side of (EIQ22) is strictly positive for every $t > 0$ if this continuous Fourier transform has a nonzero value. Finally $\widehat{\nu} = 0$ implies $\nu = 0$ without an extra uniqueness assumption. Convolve ν with a centred Gaussian approximate identity. Its value at every point is zero by that same Fourier integral. For every compactly supported continuous test function, its convolution with these Gaussians converges uniformly to the test function; integration against ν then proves that ν annihilates every such function.

Let μ be any probability measure of finite energy in (EIQ1), and write $\mu_*(dx) = \rho_{\alpha,\beta}(x) dx$ and $\nu = \mu - \mu_*$. The integrability needed for (EIQ21) holds. The $\mu \otimes \mu$ term has it by finite energy and the already established logarithmic moment; the $\mu_* \otimes \mu_*$ term has it by bounded compact support; and the cross term has it because the negative logarithmic singularity integrated against the bounded density is uniformly bounded, while the positive part is controlled by $\log(1+x) + \log(1+y)$. Expanding the energy therefore yields exactly

$$\mathcal{E}_{\alpha,\beta}(\mu) - \mathcal{E}_{\alpha,\beta}(\mu_*) = \int (U(x) - \ell) d\mu(x) - \iint \log|x-y| d\nu(x)d\nu(y) \geq 0. \quad (\text{EIQ23})$$

The first term is nonnegative by (EIQ17)–(EIQ19), and the second is strictly positive unless $\mu = \mu_*$. Infinite energy competitors cannot improve this finite value. Thus (EIQ11) is the unique global minimizing probability measure, with exactly the support $[a, b]$.

35.4 An explicit endpoint integral for the logarithmic moment

Retain m, R_s, u, v as defined above, and define

$$C = \frac{(u+v)^2}{4}, \quad \eta = \frac{v-u}{v+u}, \quad \zeta_s = \frac{b-a}{2(m+s+R_s)} \quad (s \geq 0). \quad (\text{EIQ24})$$

Thus $0 < \zeta_s \leq \eta < 1$ and $\zeta_0 = \eta$. For every $s \geq 0$ put

$$L_s = (m+s-R_s) \log C - (m-R_0) - 2R_s \log(1-\eta\zeta_s). \quad (\text{EIQ25})$$

Then the logarithmic moment of the equilibrium measure is exactly

$$\mathcal{L}(\alpha, \beta) := \int_a^b \log x \rho_{\alpha,\beta}(x) dx = \frac{\beta}{4\pi} \int_0^\infty \frac{L_0 - L_s}{\sqrt{s} R_s} ds. \quad (\text{EIQ26})$$

This is a convergent one-dimensional endpoint integral of explicitly given real elementary functions; its only endpoints are those specified uniquely in (EIQ3)–(EIQ5).

Here is a direct derivation, including the constants in (EIQ25). Set $h_0 = (b-a)/2$ and substitute $t = m + h_0 \cos \theta$ in arcsine integrals. The identities

$$t = C(1 + 2\eta \cos \theta + \eta^2), \quad t + s = \frac{m + s + R_s}{2}(1 + 2\zeta_s \cos \theta + \zeta_s^2)$$

give the absolutely convergent Fourier series

$$\log t = \log C + 2 \sum_{j \geq 1} \frac{(-1)^{j+1} \eta^j}{j} \cos(j\theta), \quad \frac{1}{t+s} = \frac{1}{R_s} \left(1 + 2 \sum_{j \geq 1} (-\zeta_s)^j \cos(j\theta) \right).$$

Orthogonality on $[0, \pi]$ gives

$$\begin{aligned} \frac{1}{\pi} \int_a^b \frac{\log t}{\sqrt{(b-t)(t-a)}} dt &= \log C, \\ \frac{1}{\pi} \int_a^b \frac{t \log t}{\sqrt{(b-t)(t-a)}} dt &= m \log C + (m - R_0), \\ \frac{1}{\pi} \int_a^b \frac{\log t}{(t+s)\sqrt{(b-t)(t-a)}} dt &= \frac{\log C + 2 \log(1 - \eta \zeta_s)}{R_s}. \end{aligned} \tag{EIQ27}$$

The second identity uses $h_0 \eta = m - R_0$. Polynomial division

$$\frac{(b-t)(t-a)}{t+s} = -t + 2m + s - \frac{R_s^2}{t+s}$$

now proves, exactly,

$$L_s = \frac{1}{\pi} \int_a^b \frac{\log t \sqrt{(b-t)(t-a)}}{t+s} dt. \tag{EIQ28}$$

Using the positive integral expression for h in (EIQ10), and the identity $1/[t(t+s)] = (1/t - 1/(t+s))/s$, yields (EIQ26). The exchanges are absolutely convergent: $\log t$ is bounded on $[a, b]$, $L_0 - L_s = O(s)$ at zero by (EIQ28), and $L_s = O(s^{-1})$ at infinity. Thus the integrand in (EIQ26) is $O(s^{1/2})$ at zero and $O(s^{-3/2})$ at infinity.

35.5 Parameter continuity and the variational derivative

The density and its logarithmic moment depend smoothly on $(\alpha, \beta) \in (0, \infty)^2$ in the following precise sense. On a compact parameter neighbourhood, (EIQ3)–(EIQ6) put a, b in a fixed compact subset of $(0, \infty)$, with $b - a$ bounded below by a positive number. The integral for h and each of its parameter and x derivatives converge uniformly for x in this fixed compact set: their absolute integrands are bounded by constants times $s^{1/2}$ near zero and $s^{-3/2}$ near infinity. Parameter derivatives of the endpoints are bounded on a smaller compact neighbourhood. Under the fixed-interval coordinate $x = a + (b-a)y$, $0 \leq y \leq 1$, the measure is exactly

$$\rho_{\alpha, \beta}(x) dx = \frac{(b-a)^2}{2\pi} \sqrt{y(1-y)} h(a + (b-a)y) dy. \tag{EIQ29}$$

All parameter derivatives of its coefficient are bounded by a constant times $\sqrt{y(1-y)}$. The same is true after multiplication by $\log(a + (b-a)y)$ or by $\sqrt{a + (b-a)y}$. Differentiation under this integral proves smoothness, in particular continuity, of both moments on the entire positive parameter quadrant.

Define $E_{\min}(\alpha, \beta) = \mathcal{E}_{\alpha, \beta}(\mu_{\alpha, \beta})$, where the measure is the unique minimizer proved above. Testing the minimizer for a parameter pair against the neighbouring potential, and then reversing the two tests, gives for $h > 0$

$$-\mathcal{L}(\alpha + h, \beta) \leq \frac{E_{\min}(\alpha + h, \beta) - E_{\min}(\alpha, \beta)}{h} \leq -\mathcal{L}(\alpha, \beta).$$

The corresponding inequalities for $h < 0$ and the proved continuity therefore give

$$\partial_{\alpha} E_{\min}(\alpha, \beta) = -\mathcal{L}(\alpha, \beta), \quad \partial_{\beta} E_{\min}(\alpha, \beta) = \int \sqrt{x} d\mu_{\alpha, \beta}(x). \quad (\text{EIQ30})$$

No unproved differentiability of a minimizing selection is being used. For the opposite convention $F(\alpha, \beta) = \sup_{\mu} \{ \iint \log |x - y| d\mu(x) d\mu(y) - \int V_{\alpha, \beta} d\mu \} = -E_{\min}(\alpha, \beta)$, the exact derivative is therefore $\partial_{\alpha} F(\alpha, \beta) = \mathcal{L}(\alpha, \beta)$.

The second moment in (EIQ30) has an exact value. Push the minimizing measure forward by the explicit dilation $x \mapsto c^2 x$, $c > 0$. Writing $M = \int \sqrt{x} d\mu_{\alpha, \beta}(x)$, its energy minus the energy at $c = 1$ is precisely $\beta(c - 1)M - 2(\alpha + 1) \log c$. Its derivative at the minimizing value $c = 1$ is zero. Hence

$$\int \sqrt{x} d\mu_{\alpha, \beta}(x) = \frac{2(\alpha + 1)}{\beta}. \quad (\text{EIQ31})$$

The same explicit dilation, or (EIQ3)–(EIQ5), proves that the map $x \mapsto (\pi/\beta)^2 x$ pushes $\mu_{\alpha, \pi}$ to $\mu_{\alpha, \beta}$, including the density and its two endpoints. Consequently

$$\mathcal{L}(\alpha, \beta) = \mathcal{L}(\alpha, \pi) + 2 \log(\pi/\beta). \quad (\text{EIQ32})$$

For the particular original limiting potential $V(x) = \pi\sqrt{x} - 2 \log x$, the substitutions are exactly $\alpha = 2$, $\beta = \pi$. Equations (EIQ5), (EIQ10), and (EIQ26) become

$$uK(k) = 2, \quad vE(k) = 4, \quad h(x) = \frac{1}{2x} \int_0^{\infty} \frac{\sqrt{s}}{(x+s)R_s} ds, \quad \mathcal{L}(2, \pi) = \frac{1}{4} \int_0^{\infty} \frac{L_0 - L_s}{\sqrt{s}R_s} ds. \quad (\text{EIQ33})$$

In particular the preceding proof includes positivity, unit mass, the full Euler inequality, the unique global minimizer, and the parameter continuity needed at these actual parameters.

36 A uniform finite partition estimate from the original equilibrium

QLG1. Exact measures and parameter domain. Fix the compact parameter rectangle

$$\mathcal{K} = [3/2, 5/2] \times [\pi/2, 3\pi/2], \quad V_{\alpha,\beta}(x) = \beta\sqrt{x} - \alpha \log x \quad (x > 0). \quad (\text{QLG1})$$

The equilibrium proved in EIQ1–33 has density

$$\rho(x) = \frac{\beta\sqrt{(b-x)(x-a)}}{4\pi^2 x} \int_0^\infty \frac{\sqrt{s} ds}{(x+s)\sqrt{(a+s)(b+s)}} \quad (a < x < b), \quad \rho(x) = 0 \quad (x \notin (a, b)), \quad (\text{QLG2})$$

where the exact endpoints are those of EIQ3–5. In particular $0 < a < b < \infty$, the density is bounded and continuous, positive in the interior, and has integral one. Write

$$d\mu(x) = \rho(x) dx, \quad P(z) = \int_a^b \log|z-t| d\mu(t) \quad (z \in \mathbb{C}), \quad I(\mu) = \iint \log|x-y| d\mu(x) d\mu(y),$$

$$F(\alpha, \beta) = I(\mu) - \int V_{\alpha,\beta} d\mu. \quad (\text{QLG3})$$

The original Euler equality and full exterior inequality in EIQ17–19 give a real number ℓ for which

$$D(x) = V_{\alpha,\beta}(x) - 2P(x) - \ell \geq 0 \quad (x > 0), \quad D(x) = 0 \quad (a \leq x \leq b). \quad (\text{QLG4})$$

Integrating the equality on the support gives the exact sign identities

$$\ell = \int V_{\alpha,\beta} d\mu - 2I(\mu), \quad F(\alpha, \beta) = -\ell - I(\mu). \quad (\text{QLG5})$$

The two integration measures treated below are, separately,

$$d\lambda_0(x) = dx, \quad d\lambda_\omega(x) = d\omega(x) = \frac{1}{2}x^{-1/2}e^{-\sqrt{x}} dx. \quad (\text{QLG6})$$

The map $x = r^2$ gives $\int d\omega = \int_0^\infty e^{-r} dr = 1$. For either of these measures, with its stated density w_λ , define the literal finite integral

$$J_n^\lambda(\alpha, \beta) = \frac{1}{n!} \int_{(0,\infty)^n} \exp\left(2 \sum_{i<j} \log|x_i - x_j| - n \sum_{i=1}^n V_{\alpha,\beta}(x_i)\right) \prod_{i=1}^n d\lambda(x_i), \quad n \geq 1. \quad (\text{QLG7})$$

No source measure, scale, factorial, or exponent has been absorbed into F . The unweighted integral without the Heine factor is exactly $n!J_n^\lambda$.

QLG2. Constants proved finite on the whole rectangle. Let

$$a_* = \min_{(\alpha,\beta) \in \mathcal{K}} a(\alpha, \beta) > 0, \quad B_* = \max_{(\alpha,\beta) \in \mathcal{K}} b(\alpha, \beta) < \infty, \quad \beta_- = \pi/2, \quad \beta_+ = 3\pi/2. \quad (\text{QLG8})$$

These are explicit compact-parameter extrema of the uniquely specified endpoint integrals EIQ3–5. Their asserted strict inequalities follow from the continuous positive endpoints proved in EIQ6 and compactness of $\mathcal{K}$. Thus their definition requires no existence or regularity assumption about an unspecified minimizing measure. For $x \in [a, b]$, the integral in QLG2 satisfies

$$\int_0^\infty \frac{\sqrt{s} ds}{(x+s)\sqrt{(a+s)(b+s)}} \leq \int_0^\infty \frac{\sqrt{s} ds}{(a_*+s)^2} = \frac{\pi}{2\sqrt{a_*}}.$$

For the last equality substitute $s = a_* r^2$, followed by $r = \tan \theta$; the resulting integral is

$$2a_*^{-1/2} \int_0^{\pi/2} \sin^2 \theta \, d\theta.$$

Since $\sqrt{(b-x)(x-a)} \leq (b-a)/2 \leq B_*/2$, it follows that

$$0 \leq \rho(x) \leq M_* := \frac{\beta_+ B_*}{16\pi a_*^{3/2}} \quad \text{for all real } x \text{ and all parameters in } \mathcal{K}. \quad (\text{QLG9})$$

In particular $B_* M_* \geq 1$, by the mass-one identity and support length at most B_* .

The entropy relative to either stated measure is

$$H_\lambda(\mu) = \int_a^b \rho(x) \log \frac{\rho(x)}{w_\lambda(x)} \, dx, \quad 0 \log 0 = 0. \quad (\text{QLG10})$$

It is absolutely integrable: the function $u|\log u|$ is bounded on $[0, M_*]$, and $\log w_\lambda$ is bounded on the fixed compact interval $[a_*, B_*]$. In the two cases its upper bounds are

$$H_{\lambda_0}(\mu) \leq \log M_*, \quad H_{\lambda_\omega}(\mu) \leq \log M_* + \log 2 + \frac{1}{2} \log B_* + \sqrt{B_*}. \quad (\text{QLG11})$$

All pair logarithmic energies on this support are absolutely integrable. For the negative part, integration against a density bounded by M_* is bounded above by $M_* \int_{-1}^1 -\log |s| \, ds = 2M_*$, uniformly in the centre. For the positive part the bound $\log^+ B_*$ suffices.

Put

$$T_\lambda = \int_0^\infty e^{-D(x)} \, d\lambda(x). \quad (\text{QLG12})$$

For $\lambda = \lambda_\omega$, QLG4 and the mass of ω give $0 < T_{\lambda_\omega} \leq 1$. For the Lebesgue measure a finite uniform bound can also be displayed entirely. For every $x > 0$,

$$P(x) \leq \log(1+x) + \log(1+B_*), \quad P(x) \geq -2M_*.$$

The lower bound uses only the negative part estimate just proved. Evaluating QLG4 at any point of $[a, b]$ therefore yields

$$\ell \leq L_* := \beta_+ \sqrt{B_*} + \frac{5}{2} \max\{|\log a_*|, |\log B_*|\} + 4M_*.$$

Consequently

$$T_{\lambda_0} \leq e^{L_*} (1+B_*)^2 \int_0^\infty (1+x)^2 (1+x^{5/2}) e^{-\beta_- \sqrt{x}} \, dx =: T_* < \infty. \quad (\text{QLG13})$$

Here $x^\alpha \leq 1 + x^{5/2}$ for all parameters in $\mathcal{K}$. The integral in QLG13 equals, by $x = r^2$ and elementary repeated integration by parts,

$$2 \left(\frac{1!}{\beta_-^2} + \frac{2 \cdot 3!}{\beta_-^4} + \frac{5!}{\beta_-^6} + \frac{6!}{\beta_-^7} + \frac{2 \cdot 8!}{\beta_-^9} + \frac{10!}{\beta_-^{11}} \right). \quad (\text{QLG14})$$

QLG3. The planar logarithmic comparison, with its proof. We use the plane only to construct an exact comparison of logarithmic kernels; the original particles and potential remain on $(0, \infty)$. If η is a finite signed measure of compact support on $\mathbb{R}^2$, has total mass zero, and $\iint |\log |z-w|| \, d|\eta|(z) d|\eta|(w) < \infty$, then

$$I(\eta) := \iint \log |z-w| \, d\eta(z) d\eta(w) \leq 0. \quad (\text{QLG15})$$

To prove this, for $d > 0$ integrate the elementary identity

$$-\log d = \frac{1}{2} \int_0^\infty \frac{e^{-td^2} - e^{-t}}{t} dt.$$

It follows by differentiation in d and evaluation at $d = 1$. On any finite subinterval of $(0, \infty)$, the integral on the right has the same sign as $-\log d$ and absolute value at most $|\log d|$. The stipulated logarithmic integrability therefore permits dominated passage to the full integral against $d\eta^2$. The constant term vanishes by the zero mass. The elementary Gaussian Fourier integral in each of the two real coordinates gives

$$\begin{aligned} -I(\eta) &= \frac{1}{2} \int_0^\infty \frac{dt}{t} \iint e^{-t|z-w|^2} d\eta(z) d\eta(w), \\ \iint e^{-t|z-w|^2} d\eta(z) d\eta(w) &= \frac{1}{4\pi t} \int_{\mathbb{R}^2} e^{-|\xi|^2/(4t)} \left| \int e^{i\xi \cdot z} d\eta(z) \right|^2 d\xi \geq 0. \end{aligned} \quad (\text{QLG16})$$

For completeness the scalar Fourier identity is obtained by differentiating $\int e^{-u^2/(4t)} e^{isu} du$ with respect to s , integrating by parts in u , and using its value $\sqrt{4\pi t}$ at zero. The product of the two scalar identities has the displayed factor $(4\pi t)^{-1}$. All finite Gaussian integrations against the measures are absolutely convergent. This proves QLG15 without requiring an equilibrium theorem on the plane.

Let $c_{x,\varepsilon}$ be probability arc measure on the circle $z = x + \varepsilon e^{i\theta}$, where the real centre x is arbitrary and $\varepsilon > 0$. Its exact logarithmic potential is

$$\int \log |z - w| dc_{x,\varepsilon}(z) = \log \max\{\varepsilon, |x - w|\}. \quad (\text{QLG17})$$

Indeed, if $|x - w| > \varepsilon$, expand $\log |1 - re^{i\theta}| = -\sum_{j \geq 1} r^j \cos(j\theta)/j$ with $r = \varepsilon/|x - w| < 1$; all nonconstant terms average to zero. If $|x - w| < \varepsilon$, use the same expansion with $r = |x - w|/\varepsilon$. At equality the assertion is the elementary integral $\int_0^{2\pi} \log |1 - e^{i\theta}| d\theta = 0$. To verify this last integral directly, write $|1 - e^{i\theta}| = 2 \sin(\theta/2)$, and put $A = \int_0^{\pi/2} \log(\sin u) du$. Reflection gives the same integral for $\log \cos u$; the double-angle identity then gives $2A = -(\pi/2) \log 2 + A$, whence $A = -(\pi/2) \log 2$. The endpoint logarithms are integrable, for example from $2u/\pi \leq \sin u \leq u$ on $[0, \pi/2]$.

For finite centres x, y , the logarithmic kernel is absolutely integrable against $c_{x,\varepsilon} \otimes c_{y,\varepsilon}$. Its positive part is bounded by a finite constant depending on the centres and ε . Applying QLG17 first to this bounded positive part minus the negative part, Tonelli's theorem and the finite value $\log \max\{\varepsilon, |x - w|\}$ show that the negative part also has finite double integral. This argument includes identical and tangent circles. The same argument proves absolute integrability of the circle- μ cross kernel, using compactness of the support of μ ; the μ - μ term was proved above. In particular QLG15 applies to a finite mixture of these circles minus μ .

Two consequences of QLG17, retaining both self and cross terms, are

$$I(c_{x,\varepsilon}) = \log \varepsilon, \quad I(c_{x,\varepsilon}, c_{y,\varepsilon}) \geq \log \max\{\varepsilon, |x - y|\} \geq \log |x - y|. \quad (\text{QLG18})$$

The first equality applies QLG17 at w on its own circle. For the second inequality, the first circle average is $\log \max\{\varepsilon, |x - w|\} \geq \log |x - w|$; averaging the right side on the second circle and applying QLG17 again gives its asserted value. When $x = y$ the last lower bound is understood in the extended sense.

QLG4. A finite upper estimate on the original particle domain. For fixed particles $x_1, \dots, x_n > 0$, put $\nu_\varepsilon = n^{-1} \sum_i c_{x_i, \varepsilon}$. By QLG18 and QLG15, respectively,

$$2 \sum_{i < j} \log |x_i - x_j| \leq n^2 I(\nu_\varepsilon) - n \log \varepsilon, \quad I(\nu_\varepsilon) \leq 2I(\nu_\varepsilon, \mu) - I(\mu). \quad (\text{QLG19})$$

There is no potential evaluated at a complex particle in these identities. The exact comparison back to the original real potential is

$$0 \leq \int P(z) dc_{x,\varepsilon}(z) - P(x) = \int_{|t-x|<\varepsilon} \log \frac{\varepsilon}{|t-x|} \rho(t) dt \leq 2M_*\varepsilon. \quad (\text{QLG20})$$

The last constant is the explicit integral $M_* \int_{-\varepsilon}^{\varepsilon} \log(\varepsilon/|s|) ds = 2M_*\varepsilon$. Combining QLG19–20 and then substituting QLG4–5 proves, for every original particle configuration,

$$2 \sum_{i<j} \log |x_i - x_j| - n \sum_i V_{\alpha,\beta}(x_i) \leq n^2 F - n \sum_i D(x_i) - n \log \varepsilon + 4M_* n^2 \varepsilon. \quad (\text{QLG21})$$

The coefficient $4M_*$ comes from twice the logarithmic potential and the bound $2M_*\varepsilon$; the coefficient of the self energy is $-n \log \varepsilon$. Take $\varepsilon = 1/n$. Since $D \geq 0$ and $n \geq 1$, $\int e^{-nD} d\lambda \leq T_\lambda$. Integrating QLG21 gives the finite positive upper bound

$$\log J_n^\lambda \leq n^2 F + 4M_* n + n \log T_\lambda + n \log n - \log(n!). \quad (\text{QLG22})$$

In particular the original integrals in QLG7 are finite. The elementary inequality $\log(n!) \geq \int_1^n \log t dt = n \log n - n + 1$ also proves

$$\log J_n^\lambda \leq n^2 F + n(1 + 4M_* + \log T_\lambda) - 1. \quad (\text{QLG23})$$

The inequality for $n!$ follows by monotonicity of $\log t$ on each interval $[j-1, j]$, for $j = 2, \dots, n$, and is exact at $n = 1$.

QLG5. Quantile lower estimate and the exact factorial cancellation. Because ρ is continuous and positive on (a, b) , there are unique endpoints

$$a = t_0 < t_1 < \dots < t_n = b, \quad \int_{t_{j-1}}^{t_j} \rho(x) dx = \frac{1}{n}, \quad \Delta_j = t_j - t_{j-1} > 0, \quad p_j(x) = n\rho(x)\mathbf{1}_{(t_{j-1}, t_j)}(x). \quad (\text{QLG24})$$

The endpoints have measure zero. Each $p_j(x)dx$ is a probability measure with bounded density, on a compact interval of positive length. Every pair logarithmic integral below is therefore absolutely integrable, including self integrals and integrals over touching intervals, by the same bounded-density estimate used after QLG11. Also $\int p_j |\log(p_j/w_\lambda)| dx < \infty$, since $p_j \leq nM_*$ and $\log w_\lambda$ is bounded on the support.

The product cell $\prod_{j=1}^n (t_{j-1}, t_j)$ and its $n!$ coordinate permutations are disjoint. Symmetry of the integrand and the product integration measure in QLG7 implies that the Heine factor $1/n!$ cancels these $n!$ equal integrals exactly. Restricting to their union therefore gives

$$J_n^\lambda \geq \int_{\prod_j (t_{j-1}, t_j)} \exp \left(2 \sum_{i<j} \log |x_i - x_j| - n \sum_i V(x_i) \right) \prod_j w_\lambda(x_j) dx_j. \quad (\text{QLG25})$$

On this cell use the probability density $\prod_j p_j(x_j)$. The logarithm of the integrand relative to this probability is integrable in absolute value, by the preceding estimates and boundedness of V on $[a, b]$. The elementary convex inequality $e^u \geq e^v(1 + u - v)$, integrated with v equal to the expectation of this logarithm, proves Jensen's lower bound directly. Since $\sum_j p_j = n\rho$, its pair and potential contributions are exactly

$$\sum_{i \neq j} I(p_i, p_j) - n \sum_j \int V p_j dx = n^2 F - \sum_j I(p_j).$$

Here $I(p_i, p_j) = \iint \log |x - y| p_i(x) p_j(y) dx dy$ and $I(p_j) = I(p_j, p_j)$. Its entropy contribution is exactly

$$\sum_j \int p_j \log \frac{p_j}{w_\lambda} dx = n \log n + n H_\lambda(\mu). \quad (\text{QLG26})$$

Moreover $|x - y| \leq \Delta_j$ on the j -th cell, whence $I(p_j) \leq \log \Delta_j$. Concavity of the logarithm, proved for example by its negative second derivative, and $\sum_j \Delta_j = b - a$ now give

$$\sum_j I(p_j) \leq \sum_j \log \Delta_j \leq n \log \frac{b - a}{n}. \quad (\text{QLG27})$$

Thus the $n \log n$ in QLG26 cancels that in QLG27, and the complete lower bound is

$$\log J_n^\lambda(\alpha, \beta) \geq n^2 F(\alpha, \beta) - n \{ \log(b - a) + H_\lambda(\mu) \}. \quad (\text{QLG28})$$

This also proves strict positivity of QLG7. No lower bound on the density at an endpoint or on a quantile interval length has been used.

QLG6. The uniform order- n theorem. For every $(\alpha, \beta) \in \mathcal{K}$, every integer $n \geq 1$, and each of the two measures QLG6, QLG22 and QLG28 are the explicit two-sided finite estimate

$$-n \{ \log(b - a) + H_\lambda(\mu) \} \leq \log J_n^\lambda - n^2 F \leq 4M_* n + n \log T_\lambda + n \log n - \log(n!). \quad (\text{QLG29})$$

For entirely parameter-uniform constants set

$$\begin{aligned} L_0 &= \log(B_* M_*), & L_\omega &= \log(B_* M_*) + \log 2 + \frac{1}{2} \log B_* + \sqrt{B_*}, \\ C_0 &= \max\{0, L_0, 1 + 4M_* + \log^+ T_*\}, & C_\omega &= \max\{0, L_\omega, 1 + 4M_*\}. \end{aligned} \quad (\text{QLG30})$$

These constants are finite by QLG8–14. Since $b - a \leq B_*$, QLG11 bounds the lower side of QLG29 by $-nL_\lambda$; QLG12–14 and QLG23 bound its upper side by nC_λ . Consequently

$$\left| \log J_n^{\lambda_0}(\alpha, \beta) - n^2 F(\alpha, \beta) \right| \leq C_0 n, \quad \left| \log J_n^{\lambda_\omega}(\alpha, \beta) - n^2 F(\alpha, \beta) \right| \leq C_\omega n. \quad (\text{QLG31})$$

For the integral Z_n^λ without the Heine factor, the exact relation $\log Z_n^\lambda = \log J_n^\lambda + \log(n!)$ gives

$$\left| \log Z_n^\lambda - n^2 F - \log(n!) \right| \leq C_\lambda n, \quad \left| \log Z_n^\lambda - n^2 F \right| \leq n \log n + C_\lambda n. \quad (\text{QLG32})$$

The theorem therefore retains the distinction between the original Heine determinant integral and the integral with labelled particles. The gain to order n in QLG31 follows from the proved circle comparison and the quantile-cell cancellation, with the same exact potential, positive-half-line particle domain, and equilibrium of EI1–33 throughout.

37 The leading first variation of the actual outer-strip quotient

Retain every original object and both original source orders of OSP1–35 and CTV1–22. In addition to the explicit finite conditions OSP23, take $k \geq 257$. This is an eventual threshold on the same fixed original parameters. Put

$$\vartheta = \pi/2, \quad \ell_s = (s-1)/4, \quad \beta_s = \frac{(2\pi)^{s/2}}{\sqrt{2}\Gamma(s/2)} \quad (s=1, k). \quad (\text{OFV1})$$

The symbol ϑ is the fixed Gamma tail rate; it is not either original source centre.

We use the equilibrium of EIQ1–33 on its original positive half-line:

$$\begin{aligned} V_{\alpha,\beta}(x) &= \beta\sqrt{x} - \alpha \log x, \quad \mu_{\alpha,\beta}(dx) = \rho_{\alpha,\beta}(x) dx, \\ F(\alpha, \beta) &= \iint \log|x-y| d\mu_{\alpha,\beta}(x) d\mu_{\alpha,\beta}(y) - \int V_{\alpha,\beta} d\mu_{\alpha,\beta}. \end{aligned} \quad (\text{OFV2})$$

Here ρ , its two endpoints, the Euler equality and the entire exterior inequality are exactly the explicit proved EIQ formulas. There is no new equilibrium assumption. EIQ29–30 proves smoothness and $F_\alpha(\alpha, \beta) = \int \log x d\mu_{\alpha,\beta}(x)$. QLG31, with its complete circle and quantile proofs, gives

$$|\log J_n(\alpha, \beta) - n^2 F(\alpha, \beta)| \leq C_0 n, \quad J_n = \frac{1}{n!} \int_{(0,\infty)^n} \prod_{i<j} (x_i - x_j)^2 \prod_i e^{-nV_{\alpha,\beta}(x_i)} dx_i \quad (\text{OFV3})$$

for every $n \geq 1$ and every $(\alpha, \beta) \in [3/2, 5/2] \times [\pi/2, 3\pi/2]$. The constant C_0 is the explicit finite constant QLG30. We use precisely this uniform finite error, including the factor $1/n!$; the conclusion below is not a derivative of an earlier unquantified q^2 limit.

37.1 Exact original-order recurrence and a small metric error

For $a \geq 0$, define

$$\mathcal{D}_a(Q; s) = \det \left[\int_{\mathbb{R}} y^{2Q+i+j} dm_s(y) \right]_{0 \leq i, j < a}, \quad \mathcal{A}_a(Q; \vartheta) = \det \left[\int_{\mathbb{R}} y^{2Q+i+j} e^{-\vartheta|y|} dy \right]_{0 \leq i, j < a}, \quad (\text{OFV4})$$

with empty determinants one. The Q used here is an integer; all the instances below have $Q \geq q/2$. The original recurrence OSP12 gives exactly

$$|y|^{2Q} dm_s(y) = \beta_s |y|^{2Q} \prod_{j=0}^{\ell_s-1} (y^2 + t_j^2) d\sigma(y), \quad t_j = 2j + \frac{1}{2}. \quad (\text{OFV5})$$

For each actual source centre $c_0 \in \{k/2, k/2 - 4\}$, the coordinate polynomial of this multiplier is $f_{\ell_s, c_0}(S) = \prod_j (S - c_0 - t_j)$. Its exact pullback satisfies $f_{\ell_s, c_0}(c_0 + iy) = \prod_j (iy - t_j) = i^{\ell_s} \prod_j (y + it_j)$. Thus multiplication by $y^Q \prod_{j=0}^{\ell_s-1} (y + it_j)$, together with the retained phase i^{ℓ_s} , is the exact coefficient map into the original one-factor Gamma source. Its squared modulus is the displayed density multiplier, with no change to any t_j or to β_s .

Set $T = \epsilon q$, $\epsilon = 2^{-32}$ as in OSP23, and

$$S_{\ell_s} = \sum_{j=0}^{\ell_s-1} t_j^2 = \frac{\ell_s(16\ell_s^2 - 12\ell_s - 1)}{12}, \quad r_{\ell_s} = \frac{S_{\ell_s}}{T^2} + \log(1 + K_q), \quad (\text{OFV6})$$

where K_q is exactly OSP24. At $\ell_s = 0$ we may and do take $r_0 = 0$ instead, since the source identity is exact. For $Q \in \{q', q\}$, $0 \leq a \leq 2q + 1$,

$$0 \leq \log \mathcal{D}_a(Q; s) - a \log \beta_s - \log \mathcal{D}_a(Q + \ell_s; 1) \leq ar_{\ell_s}. \quad (\text{OFV7})$$

Here is a proof of the full-form comparison underlying OFV7. For every real y , $|y|^{2Q} \prod_j (y^2 + t_j^2) \geq |y|^{2(Q+\ell_s)}$. On $|y| \geq T$, the logarithm of the ratio is at most $\sum_j t_j^2/T^2 = S_{\ell_s}/T^2$, by $\log(1+u) \leq u$. For $|y| < T$, its numerator is at most $(T^2 + (k/2)^2)^{Q+\ell_s}$. OSP15 on the complete coefficient space of degree at most $a-1$, OSP13 with source $s=1$, and mass $M_1 = \sqrt{2\pi}$ therefore bound its inner quadratic form by K_q times the reference quadratic form on $[q, 2q]$. This is exactly the calculation OSP22–25 with exponent $Q + \ell_s \geq q/2$, degree $a-1 \leq 2q$, and root bound $k/2 \leq R \leq T/2$. No upper bound on that exponent is needed in OSP25, since its base $(5/4)\epsilon^2$ is less than one. The full upper ratio is consequently at most $e^{S_{\ell_s}/T^2} + K_q \leq e^{r_{\ell_s}}$. The lower full ratio is at least one by the pointwise lower inequality. The positive-Gram eigenvalue argument OSP28 now proves OFV7 in the original coefficient frames. Since $S_{\ell_s} \leq \ell_s k^2/4$ and $q \geq k^2$, $r_{\ell_s} = O(1/k)$, with its stated fixed cutoff constant.

37.2 A uniform finite Gamma-to-exponential determinant estimate

Put $h = 1/q$, and define the actual finite constants

$$L_h = C_L(\sin h)^{3/2}, \quad U_h = \frac{\Gamma(1/4)^2}{2\pi}(\sin h)^{-1/2}. \quad (\text{OFV8})$$

Then the original one-factor Gamma density satisfies on the entire real line

$$L_h e^{-(\vartheta+h)|y|} \leq \frac{d\sigma}{dy}(y) \leq U_h e^{-(\vartheta-h)|y|}. \quad (\text{OFV9})$$

Indeed apply OSP7 to $\Gamma(1/4 + iy/2)$ with $\theta = \pi/2 - h$ for the upper bound. For the lower bound apply it to $\Gamma(3/4 - iy/2)$ in the exact reflection product proved before OSP11. The upper bound on that squared factor is $\Gamma(3/4)^2(\sin h)^{-3/2}e^{-(\vartheta-h)|y|}$. After division, the identity and $\cosh(\pi y) \leq e^{\pi|y|}$ give exactly the lower side of OFV9. In particular both inequalities hold near zero as well as in the tails.

The exact change of variables $y = \vartheta^{-1}u$ in OFV4 gives

$$\mathcal{A}_a(Q; \vartheta) = \vartheta^{-a(2Q+a)} \mathcal{A}_a(Q; 1). \quad (\text{OFV10})$$

The exponent follows from a moment Jacobians, $2Q$ powers in each, and the row and column powers $\sum_{i=0}^{a-1} i + \sum_{j=0}^{a-1} j = a(a-1)$. Applying OFV9 as full quadratic-form inequalities, and then using OFV10, proves the separate finite bounds

$$\begin{aligned} a \log L_h - a(2Q+a) \log(1+h/\vartheta) &\leq \log \mathcal{D}_a(Q; 1) - \log \mathcal{A}_a(Q; \vartheta) \\ &\leq a \log U_h - a(2Q+a) \log(1-h/\vartheta). \end{aligned} \quad (\text{OFV11})$$

Thus the absolute difference is at most

$$B_a(Q) = a \max\{|\log L_h|, |\log U_h|\} + a(2Q+a) \frac{h}{\vartheta - h}. \quad (\text{OFV12})$$

Here $-\log(1-u) \leq u/(1-u)$ and $\log(1+u) \leq u$ justify the second summand. Since $2h/\pi \leq \sin h \leq h$ for $0 < h \leq \pi/2$, the first summand is $O(a \log q)$. The sine inequalities follow from concavity and the endpoint chord, and by integrating $\cos t \leq 1$. For all the present $a \leq 2q+1$, $Q \leq q + \ell_s \leq 2q$, OFV12 is $O(q \log q)$ uniformly in a, Q . Combining OFV7 and OFV11 gives the error-controlled identity

$$\log \mathcal{D}_a(Q; s) = a \log \beta_s + \log \mathcal{A}_a(Q + \ell_s; \vartheta) + R_{a,Q,s}, \quad |R_{a,Q,s}| \leq a r_{\ell_s} + B_a(Q + \ell_s). \quad (\text{OFV13})$$

This is a comparison on complete original polynomial coefficient spaces, with the order shift proved by the literal multiplier OFV5; it is not an assigned replacement of the original source order.

37.3 Direct low-dimension estimates, outside the compact equilibrium range

Define

$$g(Q) = 2Q \log(2Q/\vartheta) - 2Q. \quad (\text{OFV14})$$

For $Q \geq q/2$ and $1 \leq a \leq \Delta + 1$ we prove

$$\log \mathcal{A}_a(Q; \vartheta) = ag(Q) + O(a^2 \log(Q + a) + a \log(Q + a)), \quad (\text{OFV15})$$

with constants independent of k, Q, a in these ranges. There is no use of OFV3 here; its compact parameter rectangle does not contain these low-dimension ratios.

For the lower bound take the full interval $[y_0, y_0 + 1]$, $y_0 = 2Q/\vartheta$. On it $y^{2Q} e^{-\vartheta y} \geq e^{g(Q) - \vartheta}$. Consequently its moment Gram is bounded below by $e^{g(Q) - \vartheta}$ times the unweighted monomial Gram on that interval. Translation has a triangular coefficient matrix with diagonal one, so the latter Gram determinant equals the one on $[0, 1]$. The Rodrigues norms proved in OSP15 show that it is exactly

$$\prod_{j=0}^{a-1} \frac{1}{(2j+1) \binom{2j}{j}^2}.$$

Indeed $\binom{2j}{j}$ is the leading coefficient of $L_j(2x-1)$, whose squared norm is $1/(2j+1)$. Using $\binom{2j}{j} \leq 4^j$ therefore proves

$$\log \mathcal{A}_a(Q; \vartheta) \geq ag(Q) - a\vartheta - a \log(2a) - 2a(a-1) \log 2. \quad (\text{OFV16})$$

For the upper bound, Gram–Schmidt shows that a positive Gram determinant is at most the product of the squared norms of its displayed vectors: each orthogonal component has norm no greater than its original vector. The exact monomial moments give

$$\log \mathcal{A}_a(Q; \vartheta) \leq \sum_{j=0}^{a-1} \log \left(\frac{2(2(Q+j))!}{\vartheta^{2(Q+j)+1}} \right). \quad (\text{OFV17})$$

For every positive integer m , integrating the increasing function $\log x$ between successive integers proves $m \log m - m + 1 \leq \log(m!) \leq m \log m - m + 1 + \log m$. Each summand in OFV17 is consequently at most $g(Q+j) + |\log(2/\vartheta) + 1| + \log(2(Q+j))$. Furthermore $g'(u) = 2 \log(2u/\vartheta)$, so $g(Q+j) - g(Q) \leq 2j \log(2(Q+a)/\vartheta)$ in the present range. This proves the explicit upper bound

$$\begin{aligned} \log \mathcal{A}_a(Q; \vartheta) &\leq ag(Q) + a\{|\log(2/\vartheta) + 1| + \log(2(Q+a))\} \\ &\quad + a(a-1) \log(2(Q+a)/\vartheta). \end{aligned} \quad (\text{OFV18})$$

OFV16 and OFV18 prove OFV15 with its complete lower and upper errors. They also apply at $a = 1$.

37.4 High-dimension partitions and their controlled first variation

For an integer a write $n_e = \lceil a/2 \rceil$ and $n_o = \lfloor a/2 \rfloor$. The two exact changes of variables $y = \sqrt{x}$ and $y = -\sqrt{x}$ give, with their two positive Jacobians, the parity identity

$$\mathcal{A}_a(Q; \vartheta) = Z_{n_e}(Q - \tfrac{1}{2}; \vartheta) Z_{n_o}(Q + \tfrac{1}{2}; \vartheta), \quad (\text{OFV19})$$

where

$$Z_n(A; \vartheta) = \det \left[\int_0^\infty x^{A+i+j} e^{-\vartheta \sqrt{x}} dx \right]_{0 \leq i, j < n}.$$

The two branches add to the density $x^{-1/2} e^{-\vartheta \sqrt{x}}$; thus no factor two is missing from OFV19. The odd block has the additional factor x . Expanding the two Vandermonde determinants and integrating each finite permutation sum proves the exact Heine identity for this moment determinant:

$$Z_n(A; \vartheta) = \frac{1}{n!} \int_{(0, \infty)^n} \prod_{i < j} (x_i - x_j)^2 \prod_i x_i^A e^{-\vartheta \sqrt{x_i}} dx_i.$$

Changing variables $x_i = n^2 u_i$ now gives

$$Z_n(A; \vartheta) = n^{2n(A+n)} J_n(A/n, \vartheta). \quad (\text{OFV20})$$

There are $2n(A+1)$ powers from the weights and Jacobians and $2n(n-1)$ from the squared Vandermonde, proving the exponent.

For the high instances we use exactly

$$(a, Q) \in \{(q, q + \ell_s), (q+1, q + \ell_s), (q + \Delta, q - \Delta + \ell_s), (q + \Delta + 1, q - \Delta + \ell_s)\}. \quad (\text{OFV21})$$

Since $k \geq 257$, $\Delta/q \leq 16/k < 1/16$ and $\ell_s/q \leq 1/(4k) < 1/64$. For both parity blocks, elementary rounding therefore gives

$$\frac{15q}{32} \leq n \leq \frac{9q}{16}, \quad \frac{29q}{32} \leq A \leq \frac{33q}{32}.$$

The first inequalities use $q \geq 32$ to absorb the possible rounding by $1/2$ and the extra endpoint 1. Hence $29/18 \leq A/n \leq 11/5$, contained strictly in $[3/2, 5/2]$. Every partition in OFV20 is thus in the proved parameter rectangle OFV3, with $\beta = \vartheta = \pi/2$.

Define the smooth explicit function

$$\mathcal{H}(a, Q) = a(2Q + a) \log(a/2) + \frac{a^2}{2} F(2Q/a, \vartheta) \quad (a > 0, Q > 0). \quad (\text{OFV22})$$

Equations OFV3 and OFV19–20 give uniformly over OFV21

$$\log \mathcal{A}_a(Q; \vartheta) = \mathcal{H}(a, Q) + O(q \log q). \quad (\text{OFV23})$$

To verify the rounding error rather than omit it, set $G(n, A) = 2n(A+n) \log n + n^2 F(A/n, \vartheta)$. Each actual pair (n, A) differs from $(a/2, Q)$ by at most $1/2$ in each coordinate. On all connecting segments n is bounded above and below by positive constant multiples of q , while A/n remains in $[3/2, 5/2]$. Its derivatives are

$$G_A = 2n \log n + n F_\alpha, \quad G_n = 2(A+2n) \log n + 2(A+n) + 2nF - A F_\alpha.$$

They are $O(q \log q)$ there, since the exact EIQ functions F, F_α are bounded on that compact interval. The mean value formula bounds each rounding error by $O(q \log q)$. The two QLG errors sum to at most $C_0 a = O(q)$, and $2G(a/2, Q) = \mathcal{H}(a, Q)$, proving OFV23.

The derivative needed for the first variation is computed without altering the direction of the two original degree changes:

$$(\partial_a - \partial_Q) \mathcal{H}(a, Q) = 2Q \log(a/2) + (2Q + a) + aF(2Q/a, \vartheta) - (Q + a)F_\alpha(2Q/a, \vartheta). \quad (\text{OFV24})$$

EIQ29–30 proves that F_α is smooth; in particular $F, F_\alpha, F_{\alpha\alpha}$ are bounded on the compact interval just used. Differentiating the right side of OFV24 shows that its two first partial derivatives are $O(\log q)$ on the segment $a = q + u$, $Q = q + \ell_s - u$, $0 \leq u \leq \Delta$ and on its connections to (q, q) . Every term follows by the product rule; the derivatives of $2Q/a$ are $2/a$ and $-2Q/a^2$, both $O(1/q)$. The mean value formula and then integration in u consequently give

$$\begin{aligned} & \mathcal{H}(q + \Delta, q - \Delta + \ell_s) - \mathcal{H}(q, q + \ell_s) \\ &= \Delta q \{2 \log(q/2) + 3 + F(2, \vartheta) - 2F_\alpha(2, \vartheta)\} + O(\Delta(\Delta + \ell_s) \log q). \end{aligned} \quad (\text{OFV25})$$

This remainder is $O(q \log q)$ because $\Delta = O(k)$, $\ell_s = O(k)$ and $q = (k+1)^2$. The additional endpoint $a+1$ in OFV21 changes $\mathcal{H}$ by $O(q \log q)$, directly from its displayed derivatives. Thus all rounding and endpoint changes have been quantified.

37.5 All seven determinants and the resulting leading constant

Replace the seven moment determinants in OSP32 by the exact exponential determinants in OFV13, with their respective $Q + \ell_s$. Their $a \log \beta_s$ terms cancel because their signed dimensions are exactly

$$\Delta + (\Delta + 1) - (q + \Delta) - (q + \Delta + 1) - 1 + q + (q + 1) = 0. \quad (\text{OFV26})$$

The sum of their absolute finite errors in OFV13 is $O(q \log q)$, since the sum of their dimensions is $4(q + \Delta + 1)$, $r_{\ell_s} = O(1/k)$, and OFV12 applies uniformly with $Q + \ell_s \leq 2q$. The low terms, using OFV15–18, are

$$\begin{aligned} \log \mathcal{A}_\Delta(q - \Delta + \ell_s; \vartheta) + \log \mathcal{A}_{\Delta+1}(q - \Delta + \ell_s; \vartheta) - \log \mathcal{A}_1(q + \ell_s; \vartheta) \\ = 2\Delta g(q) + O(q \log q) = 4\Delta q \{\log(2q/\vartheta) - 1\} + O(q \log q). \end{aligned} \quad (\text{OFV27})$$

Indeed replacing each $g(q - \Delta + \ell_s)$ or $g(q + \ell_s)$ by $g(q)$ costs at most $O(((2\Delta + 1)(\Delta + \ell_s) + \ell_s) \log q) = O(q \log q)$, using $g'(u) = 2 \log(2u/\vartheta)$ and the actual degree ranges. The individual low determinant errors are $O((\Delta + 1)^2 \log q) = O(q \log q)$.

The remaining four high terms are, by OFV23–25,

$$\begin{aligned} \log \mathcal{A}_q(q + \ell_s; \vartheta) + \log \mathcal{A}_{q+1}(q + \ell_s; \vartheta) \\ - \log \mathcal{A}_{q+\Delta}(q - \Delta + \ell_s; \vartheta) - \log \mathcal{A}_{q+\Delta+1}(q - \Delta + \ell_s; \vartheta) \\ = -2\Delta q \{2 \log(q/2) + 3 + F(2, \vartheta) - 2F_\alpha(2, \vartheta)\} + O(q \log q). \end{aligned} \quad (\text{OFV28})$$

Combining OFV27 and OFV28 gives the complete constant

$$\mathfrak{C}_\partial = 4 \log(4/\vartheta) - 10 - 2F(2, \vartheta) + 4F_\alpha(2, \vartheta), \quad \vartheta = \pi/2, \quad (\text{OFV29})$$

with every logarithmic scale term canceled by direct algebra. OSP33 now proves the actual original-root asymptotic

$$\boxed{\mathcal{F}_{\partial, k}^{(s)} = \Delta q \mathfrak{C}_\partial + O_{\delta, \gamma}(q \log q), \quad \frac{\mathcal{F}_{\partial, k}^{(s)}}{kq} \longrightarrow 16 \mathfrak{C}_\partial, \quad s = 1, k.} \quad (\text{OFV30})$$

All constants in the error depend only on the fixed original parameters, the fixed cutoff, and the proved compact equilibrium constants. The two original source orders have been treated through their exact recurrence maps and full mass factors. Since $\Delta/k = 16 - 48/k$ and $\log q/k \rightarrow 0$, the displayed limit follows from its quantified remainder.

An equivalent expression uses the logarithmic energy of precisely the original EIQ limiting potential, $V(x) = \pi\sqrt{x} - 2 \log x$:

$$I_{2, \pi} = \iint \log |x - y| d\mu_{2, \pi}(x) d\mu_{2, \pi}(y), \quad \mathfrak{C}_\partial = 4 \log(4/\pi) + 2 - 2I_{2, \pi}. \quad (\text{OFV31})$$

To prove it, EIQ31 gives $\vartheta \int \sqrt{x} d\mu_{2, \vartheta} = 6$, so $F(2, \vartheta) = I_{2, \vartheta} - 6 + 2F_\alpha(2, \vartheta)$. Substituting in OFV29 yields $4 \log(4/\vartheta) + 2 - 2I_{2, \vartheta}$. The proved EIQ32 dilation $x \mapsto 4x$ carries $\mu_{2, \pi}$ to $\mu_{2, \pi/2}$. Its exact logarithmic energy change is $I_{2, \pi/2} = I_{2, \pi} + \log 4$, proving OFV31. This energy is a finite explicit integral of the fully specified EIQ density; its logarithmic singularity is integrable because that density is bounded and compactly supported.

Finally, the exact quotient metrics decrease when the source degree increases: each smaller-degree lift remains an allowed lift in the larger source with exactly the same norm. Hence each low-degree quotient Gram in CTV13 dominates the corresponding high-degree Gram on the identical target frame. Positive conjugation shows their determinant ratios are at least one. It follows that $\mathcal{F}_{\partial, k}^{(s)} \geq 0$, and OFV30 therefore also proves $\mathfrak{C}_\partial \geq 0$. The already proved finite tail bounds CTV18–20, with $0 \leq v \leq 80$, further give

$$\mathcal{T}_k^{(s)} = \Delta q \mathfrak{C}_\partial + O_{\delta, \gamma}(q \log q), \quad s = 1, k, \quad (\text{OFV32})$$

uniformly over the possible actual first-moment indices. No leading value of the separate original observation quotient is assigned here.

38 Fixed coefficient control of the original residual module

Retain the complete actual simple-quartet period matrix $H = F^{-1}\Pi U\mathcal{E}_{\text{prim}}$, the original substitution $t = (z_0w_0, z_0w_1, z_1w_0, z_1w_1)^T$, and $Q(x) = Q_0(H^{-1}x)$, where $Q_0(t) = t_{++}t_{--} - t_{+-}t_{-+}$. The original action is $gf(x) = f(D^{-1}x)$ with $D = \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4)$ and $\zeta = e^{2\pi i/5}$. At the fixed actual period on the proved five-member orbit locus retain the entire polynomials $Q_a = g^a Q$ and $\mathcal{A} = \prod_{a=1}^4 Q_a$. Their coefficients and all phases are unchanged. The degree- k calculations below use $k = 4l + 1 \geq 9$ and

$$q = (k+1)^2, \quad q' = (k-7)^2, \quad \Delta = 16k - 48, \quad j = 8k - 32, \quad r = 8k - 16. \quad (\text{FMC1})$$

All constants introduced below are determined by this one fixed actual coefficient datum. They are independent of k and of the original Gamma cutoff. A fixed presentation will supply the residual frame in place of dividing new selected minors at each degree.

38.1 A fixed free module containing the entire invariant image

Let $P = \mathbb{C}[x_1, x_2, x_3, x_4]$ and introduce the specified graded ring injection

$$T = \mathbb{C}[y_1, y_2, y_3, y_4] \longrightarrow P, \quad y_i \longmapsto x_i^5, \quad \deg y_i = 5. \quad (\text{FMC2})$$

It is injective because distinct monomials have distinct exponent vectors after multiplying each exponent by five. Put $\mathcal{E} = \{a \in \mathbb{Z}^4 : 0 \leq a_i \leq 4\}$ and let e_a be a free T -basis element of degree $|a|$. The exact isomorphism is

$$\mathcal{P} := \bigoplus_{a \in \mathcal{E}} T e_a \xrightarrow{\iota} P, \quad y^\nu e_a \longmapsto x^{5\nu+a}. \quad (\text{FMC3})$$

Every nonnegative exponent vector has a unique coordinatewise quotient and remainder on division by five. This proves surjectivity, injectivity, preservation of degree, and the explicit inverse. In particular this is a rank-625 free module, with coefficient ℓ^1 norm preserved exactly by ι . The ring and monomial coefficients in P have not been rescaled.

Let $\mathcal{E}_0 = \{a \in \mathcal{E} : a_1 + 2a_2 + 3a_3 + 4a_4 \equiv 0 \pmod{5}\}$. The full original invariant algebra, regarded here as a T -module, is

$$\mathcal{J} = P^{\langle g \rangle} = \iota \left(\bigoplus_{a \in \mathcal{E}_0} T e_a \right), \quad |\mathcal{E}_0| = 125. \quad (\text{FMC4})$$

Indeed the monomial $x^{5\nu+a}$ has character $\zeta^{-(a_1+2a_2+3a_3+4a_4)}$. The diagonal action fixes a polynomial precisely when each nonzero monomial has character one. For each of the 5^3 choices of the first three residues there is exactly one a_4 since 4 is invertible modulo five. This proves the count and the displayed equality. These pure fifth powers are four of the fourteen original generators OCF25; this is a new exact module presentation of the same entire invariant algebra.

Define two homogeneous T -submodules of $\mathcal{P}$ by

$$\begin{aligned} \mathcal{H} &= \iota^{-1}(QP + \mathcal{A}P), \\ \mathcal{S} &= \mathcal{H} + \bigoplus_{a \in \mathcal{E}_0} T e_a. \end{aligned} \quad (\text{FMC5})$$

They have the fixed finite generating columns $\iota^{-1}(Qx^a)$ and $\iota^{-1}(\mathcal{A}x^a)$ for $a \in \mathcal{E}$, followed for $\mathcal{S}$ by the 125 columns e_a , $a \in \mathcal{E}_0$. The first two lists have respectively homogeneous degrees $|a| + 2$ and $|a| + 8$, all at most 24. To obtain every matrix entry explicitly, expand either of the full original polynomials as $f(x) = \sum_b f_b x^b$ and use

$$\iota^{-1}(f x^a) = \sum_b f_b y^{[(a+b)/5]} e_{(a+b) \bmod 5}. \quad (\text{FMC6})$$

Both quotient and remainder are coordinatewise. Formula FMC3 verifies this identity term by term. Since P is free on the x^a , these columns generate precisely the two modules in FMC5; no later degree is needed to define a new relation matrix.

The full coefficient substitution and conductor quotient give the exact graded maps

$$\mathcal{P}/\mathcal{H} \xrightarrow{\iota, \psi} \mathcal{B}/(AB) =: \mathcal{L}, \quad \mathcal{S}/\mathcal{H} \xrightarrow{\sim} \overline{\mathcal{J}}, \quad A = \psi_8(\mathcal{A}). \quad (\text{FMC7})$$

Here $\psi f = f(Ht)$, and $\overline{\mathcal{J}}$ is the image of the original invariant algebra in $\mathcal{L}$, exactly OCF16. OCF3 proves $\ker \psi = QP$ and its surjectivity in every degree. If $\psi f = Ab$, lift b to a polynomial h using that surjectivity; then $f - Ah \in QP$. Conversely every $QP + AP$ element maps to zero. This proves the first isomorphism. FMC4–5 show that its restriction has precisely the second image and kernel. Thus the quotient in FMC7 is the complete original conductor quotient, and its indicated submodule is the actual residual certificate image of ROQ4, with no new invariant premise.

38.2 Finite reduction data with only fixed actual denominators

The next construction specifies fixed data for both modules in FMC5. Order their module monomials $y^\nu e_a$ first lexicographically by $\nu_1, \nu_2, \nu_3, \nu_4$, and then by a fixed index $\kappa_a \in \{0, \dots, 624\}$ listing a lexicographically. Multiplication by a y -monomial preserves this order. It is a well-order: in a decreasing infinite sequence the first coordinate can decrease only finitely often; after it is fixed the same applies successively to the other three coordinates and then the finite index. For a nonzero column f , retain its full leading coefficient $\lambda_f \neq 0$ and its leading module monomial m_f . No leading coefficient is deleted from the stored column.

Start with the finite nonzero homogeneous generators of FMC5–6. Division of a column repeatedly cancels its largest divisible module monomial $cy^\nu e_a$ using the first available column f for which $m_f = y^\mu e_a$ divides it: subtract $(c/\lambda_f)y^{\nu-\mu}f$ and record that exact multiple. A largest term divisible by none of the current leading monomials is put into the remainder and removed from the current dividend. Each operation strictly decreases the largest term still being processed, so this division terminates. Remainders have no monomial divisible by the stored leading monomials.

For every pair f, h whose leading positions agree, put $m = \text{lcm}(m_f, m_h)$ in that position and form the full column

$$\text{Sp}(f, h) = \frac{m/m_f}{\lambda_f} f - \frac{m/m_h}{\lambda_h} h. \quad (\text{FMC8})$$

Divide it as just described and append its nonzero remainder to the list. Reconsider all new pairs. Track every appended column as a T -linear combination of the original generators throughout these operations. Every column and recorded certificate remains homogeneous: the two terms in FMC8 have the same weighted degree, because their leading position agrees and every input column is homogeneous in the shifted grading of FMC3.

This algorithm stops after finitely many appended columns. Here is the needed finiteness argument. Every infinite sequence in $\mathbb{N}^4$ has two terms in their original order with all coordinates weakly increasing. To prove it, an infinite sequence of nonnegative integers has either an infinitely repeated value or an increasing infinite subsequence; use this successively for all four coordinates. For module monomials first take a subsequence with one common position. Consequently every infinite sequence of module monomials has an earlier one dividing a later one. Each appended remainder's leading monomial is divisible by none of the previous leading monomials. An infinite sequence of appendages would contradict the proved property.

At termination every element of the generated module has leading monomial divisible by one of the stored leading monomials. For completeness, write an element as a finite sum of monomial multiples of the stored columns. Among all such representations choose one for which the largest leading module monomial of a summand is smallest, and among these choose one with the fewest summands at that monomial. These choices exist by the well-order and the positive integers. If cancellation of those largest terms occurs, two summands at the same module position can be

combined using a multiple of FMC8, with its exact two leading coefficients. Explicitly, if their common leading monomial is M and their leading coefficients are a, b , their sum is

$$a \frac{M}{\text{lcm}(m_f, m_h)} \text{Sp}(f, h) + (a + b) \frac{M/m_h}{\lambda_h} h.$$

This follows by substituting FMC8; it retains exactly both original summands and every coefficient. The recorded division of this S-column, together with its appended remainder when one was needed, expresses it in stored columns with all leading multiples strictly below its canceled common monomial. Replacing the two summands by this expression either decreases the largest monomial or decreases the number attaining it; combining their coefficients if necessary gives the latter alternative. Both contradict the chosen representation. Thus the largest remaining monomial is an actual leading monomial of the element, and it is divisible by one of the stored ones, as claimed. This also proves that division of a module element has zero remainder.

Write $\mathcal{G}_H$ and $\mathcal{G}_S$ for the two resulting fixed finite lists for FMC5. The preceding argument proves their full reduction property. A column supported outside the respective leading module is called a standard remainder. Such a remainder represents each quotient class uniquely: two representatives of one class differ by a module element, and a nonzero standard difference would violate the proved leading-monomial property. Denote these exact linear remainder maps by NF_H and NF_S . Their linearity follows from uniqueness and linearity of quotienting.

All data constructed here depend only on the actual matrices in FMC6. Every operation is an addition, multiplication, or division by one of the nonzero actual leading coefficients encountered in this finite calculation. Thus the complete list of denominators is finite and independent of k . Its precise coefficient chart records each tested coefficient's vanishing or nonvanishing and every selected leading monomial. For every actual period exactly its corresponding decisions give the above terminating calculation. This does not posit a numerical test for equality of unevaluated periods, nor does it assert a lower bound for a later sequence of minors. The exact stored coefficients and their nonzero leading entries define the constants that follow.

38.3 A fixed weight bounds every homogeneous reduction path

The finite reductions admit a linear-in-degree depth bound. Let $D_0 \geq 1$ be the largest of one and all absolute differences between corresponding y -exponents in a leading and a nonleading term of any column in $\mathcal{G}_H \cup \mathcal{G}_S$. Empty tail lists contribute zero. Set

$$B_0 = D_0 + 626, \quad \mathbf{w}(y^\nu e_a) = B_0^4 \nu_1 + B_0^3 \nu_2 + B_0^2 \nu_3 + B_0 \nu_4 + \kappa_a, \quad L_k = B_0^4 \lfloor k/5 \rfloor + 624. \quad (\text{FMC9})$$

Every nonleading term of either fixed list has weight at least one less than its leading term. If the y -exponents agree, this follows from the position order. Otherwise let their first differing coordinate have weight B_0^m , $1 \leq m \leq 4$. Its exponent drops by at least one, while the later differences have absolute value at most D_0 , and the position difference has absolute value at most 624. The total weight drop is at least

$$B_0^m - D_0 \sum_{h=1}^{m-1} B_0^h - 624 \geq \frac{625 B_0^m}{D_0 + 625} - 624 > 1. \quad (\text{FMC10})$$

The middle inequality follows by bounding the geometric sum by $B_0^m/(B_0 - 1)$; the last follows from $B_0^m \geq B_0 = D_0 + 626$. Multiplication by a monomial adds the same weight to the leading and all its replacement terms, so the same drop holds in every division step. A homogeneous degree- k monomial has $5|\nu| + |a| = k$, hence weight between zero and L_k . Every branch of its reduction tree consequently has length at most L_k .

For each of the two lists define the complete coefficient constant

$$C_H = \max \left\{ 1, \max_{g \in \mathcal{G}_H} \sum_{m \neq m_g} |[m]g/\lambda_g| \right\}, \quad C_S = \max \left\{ 1, \max_{g \in \mathcal{G}_S} \sum_{m \neq m_g} |[m]g/\lambda_g| \right\}. \quad (\text{FMC11})$$

The maximum over an empty list is zero. They are finite actual numbers because the lists are finite and their displayed leading coefficients are nonzero. For every homogeneous degree- k column,

$$\|\text{NF}_H f\|_1 \leq C_H^{L_k} \|f\|_1, \quad \|\text{NF}_S f\|_1 \leq C_S^{L_k} \|f\|_1. \quad (\text{FMC12})$$

To prove these inequalities start with one monomial of coefficient one. A reduction replaces its leading term by the complete tail with total absolute coefficient at most the indicated constant. Along each branch the nonnegative integer weight drops at least one. Induction on that weight bounds the total absolute coefficient of all final leaves by $C^{\mathfrak{w}(m)}$: a standard leaf contributes one, and an internal node contributes at most C times the bound at weight one lower. Cancellation between branches can only decrease the coefficient ℓ^1 norm. Linearity, the triangle inequality, and $\mathfrak{w}(m) \leq L_k$ prove FMC12 for every input.

There is equally explicit control of the recorded original-generator certificates. For each stored $g \in \mathcal{G}_S$ let c_g be its complete certificate in the fixed original generating list of $\mathcal{S}$ in FMC5–6. Define

$$D_S = \max \left\{ 1, \max_{g \in \mathcal{G}_S} \|c_g\|_1 / |\lambda_g| \right\}. \quad (\text{FMC13})$$

At a reduction node with input coefficient c , the certificate contribution has norm at most $D_S |c|$. At depth h the total absolute coefficient entering all nodes is at most C_S^h times the input norm, by the same tree proof. Summing through depth L_k therefore proves that a certificate of $f - \text{NF}_S f$ in the original generators has norm at most $D_S(L_k + 1)C_S^{L_k} \|f\|_1$. All certificate entries retain the original homogeneous grading. Their denominators are among the same finite fixed reduction data; no degree- k linear system is inverted.

38.4 The two exact maps back to the original conductor strip

Let $\mathbf{N}_k$ and $\mathbf{S}_k$ be the original OCF strip remainder and its coordinate insertion, with leading conductor coefficient $a_* = a_{r_* s_*}$. Define

$$A_0 = \max \left\{ 1, \sum_{(a,b) \neq (r_*, s_*)} |a_{ab}/a_*| \right\}. \quad (\text{FMC14})$$

Their exact coefficient bounds are $\|\mathbf{N}_k\|_1 \leq A_0^{10k}$ and $\|\mathbf{S}_k\|_1 = 1$. To prove the first, give the biform monomial $e_{ab}^{(k)}$ the weight $9a + b \in [0, 10k]$. Every other nonzero term of the full A has smaller weight than its lexicographically largest term: a decrease in the first index outweighs at most eight increases in the second index, and a decrease in only the second index also drops the weight. Every translated OCF6 subtraction therefore has weight drop at least one. Its complete tail has coefficient norm at most A_0 . The reduction-tree proof of FMC12 now applies with depth $10k$. The insertion bound is immediate from its unchanged coordinate coefficients.

Put

$$h_+ = \max \{ 1, \max_i \sum_a |\mathbf{H}_{ia}| \}, \quad h_- = \max \{ 1, \max_a \sum_i |(\mathbf{H}^{-1})_{ai}| \}, \quad h_* = \max \{ h_+, h_- \}. \quad (\text{FMC15})$$

Substitution $x = \mathbf{H}t$ on degree- k polynomial coefficients has ℓ^1 norm at most h_+^k . Expanding each product of linear forms and using the multinomial theorem proves this for each monomial; the triangle inequality proves it for their sum. The Segre substitution adds coefficients on identical biform monomials and so has norm at most one. Conversely choose the following exact lift of $e_{ab}^{(k)}$: with $h = \min(a, b)$ take

$$t_{++}^h t_{+-}^{a-h} t_{-+}^{b-h} t_{--}^{k-a-b+h}, \quad t = \mathbf{H}^{-1}x. \quad (\text{FMC16})$$

All exponents are nonnegative since $0 \leq a, b \leq k$, and their sum is k . Direct substitution gives the required biform monomial. This lift, extended linearly and denoted $\mathcal{L}_k$, has norm at most h_-^k by the same complete multinomial estimate.

Let $\mathcal{M}_{H,k}$ be the set of degree- k standard module monomials for $\mathcal{G}_H$. Its coefficient space is the exact degree- k quotient model of FMC7. Define

$$\begin{aligned} U_k f &= \mathbf{N}_k \psi \iota f, & f &\in \mathbb{C}^{\mathcal{M}_{H,k}}, \\ V_k w &= \mathbf{NF}_H \iota^{-1} \mathcal{L}_k \mathbf{S}_k w, & w &\in W_k. \end{aligned} \quad (\text{FMC17})$$

These are inverse maps. For the first composition, $\mathbf{NF}_H$ changes a lift only by $\mathcal{H}$, which maps to zero in the original strip by FMC7, and $\psi \mathcal{L}_k = I$ together with $\mathbf{N}_k \mathbf{S}_k = I$ proves $U_k V_k = I$. For the opposite composition, $\mathcal{L}_k \mathbf{S}_k \mathbf{N}_k \psi \iota f - \iota f$ maps to zero in the strip, so it belongs to $QP + \mathcal{AP}$ by FMC7. Its free-module difference belongs to $\mathcal{H}$ and is killed by the unique remainder. The original f is already standard. Hence $V_k U_k = I$. The bounds already proved give

$$\|U_k\|_1 \leq h_+^k A_0^{10k}, \quad \|V_k\|_1 \leq h_-^k C_H^{L_k}. \quad (\text{FMC18})$$

These bounds use only the displayed original coefficients and the two fixed homogeneous relation lists.

38.5 An identity pivot block for the entire residual image

Since $\mathcal{H} \subset \mathcal{S}$, every leading monomial of $\mathcal{H}$ is a leading monomial of an element of $\mathcal{S}$. The proved reduction property for $\mathcal{G}_S$ then implies inclusion of their leading modules. Consequently

$$\mathcal{M}_{S,k} \subset \mathcal{M}_{H,k}, \quad \mathcal{U}_k = \mathcal{M}_{H,k} \setminus \mathcal{M}_{S,k}, \quad |\mathcal{M}_{H,k}| = \Delta, \quad |\mathcal{M}_{S,k}| = r, \quad |\mathcal{U}_k| = j. \quad (\text{FMC19})$$

The first count is FMC7 and OCF8. The quotient by $\mathcal{S}$ is exactly the original conductor quotient modulo its invariant image, which has dimension r by OCF22–24 and OKM3–5. Subtracting proves the last count. These are counts of actual standard monomials for the fixed relation lists, not independent generic ranks.

Order $\mathcal{M}_{H,k}$ by listing $\mathcal{U}_k$ first and $\mathcal{M}_{S,k}$ second, with the specified monomial order within each list. For every $u \in \mathcal{U}_k$ put $z_u = u - \mathbf{NF}_S(u)$. Both terms are H -standard, and their difference belongs to $\mathcal{S}$. The j resulting columns form a basis of $\mathcal{S}_k/\mathcal{H}_k$ in this quotient model. Indeed their coordinates on $\mathcal{U}_k$ are the distinct unit vectors. For a standard column in $\mathcal{S}$, the remainder under $\mathbf{NF}_S$ is zero. Its coordinates on $\mathcal{M}_{S,k}$ must therefore be the negatives of the remainders of its $\mathcal{U}_k$ terms. It is exactly the stated linear combination of the z_u . This proves spanning and independence, including all tails of every column.

Write B_k for the matrix with columns $\mathbf{NF}_S(u)$ in the second standard set. The full exact graph frame and inverse are

$$P_k = \begin{pmatrix} I_j & 0 \\ -B_k & I_r \end{pmatrix}, \quad P_k^{-1} = \begin{pmatrix} I_j & 0 \\ B_k & I_r \end{pmatrix}, \quad \det P_k = 1, \quad \|P_k\|_1, \|P_k^{-1}\|_1 \leq 1 + C_S^{L_k}. \quad (\text{FMC20})$$

The two matrix identities follow by multiplication and block triangularity, and the norm bound follows from FMC12 applied to each entire column. All entries of B_k come from fixed reductions; no minor is selected or divided at degree k .

Each z_u has an actual invariant certificate. The recorded reduction certificate of $u - \mathbf{NF}_S(u)$ from FMC13 expresses its polynomial representative as

$$\iota z_u = Qh_u + Aa_u + f_u, \quad f_u \in \mathcal{J}_k, \quad \|f_u\|_{1,x} \leq D_S(L_k + 1)C_S^{L_k}. \quad (\text{FMC21})$$

To obtain the three displayed pieces, collect the certificate terms corresponding respectively to Qx^a , Ax^a , and the 125 invariant basis elements x^a . Every coefficient is a polynomial in the $y_i = x_i^5$. The last terms therefore belong to $\mathcal{J}$, and homogeneity gives exactly degree k . FMC3 preserves coefficient norm, and discarding the first two certificate lists when bounding only the last cannot increase their total ℓ^1 norm. FMC13 proves the bound. Thus $U_k z_u$ is exactly $\mathbf{N}_k \psi f_u$, an original invariant restriction, with the entire original periods in its coefficients.

Define the full original strip frame, its residual columns and their explicit left inverse by

$$T_k = U_k P_k, \quad Z_k^{\text{fix}} = T_k E_j, \quad L_{Z,k} = E_j^T P_k^{-1} V_k, \quad L_{Z,k} Z_k^{\text{fix}} = I_j. \quad (\text{FMC22})$$

Here E_j inserts the first j coordinates into the new ordered standard basis, while all output strip coordinates retain their original OCF order. FMC17 and FMC20 prove the inverse formula $T_k^{-1} = P_k^{-1} V_k$ and the last identity. By FMC21 these Z_k^{fix} columns span precisely the original $\bar{J}_k$. They provide all actual residual coefficients with a left inverse obtained from fixed relations.

For the original coefficient Euclidean norms set the explicit number

$$M_k = 2\sqrt{\Delta} h_*^k A_0^{10k} (C_H C_S)^{L_k} \geq 1. \quad (\text{FMC23})$$

Then

$$\|T_k\|_2, \|T_k^{-1}\|_2, \|Z_k^{\text{fix}}\|_2, \|L_{Z,k}\|_2 \leq M_k, \quad M_k^{-1} \|v\|_2 \leq \|Z_k^{\text{fix}} v\|_2 \leq M_k \|v\|_2. \quad (\text{FMC24})$$

For the first two bounds, FMC18–20 give respectively the column-sum bounds $h_+^k A_0^{10k} (1 + C_S^{L_k})$ and $(1 + C_S^{L_k}) h_-^k C_H^{L_k}$. For a matrix with Δ input coordinates, its Euclidean norm is at most $\sqrt{\Delta}$ times its coefficient column-sum norm: use $\|Av\|_2 \leq \|Av\|_1 \leq \|A\|_1 \|v\|_1 \leq \sqrt{\Delta} \|A\|_1 \|v\|_2$. All constants are at least one, proving the two stated bounds. The coordinate insertion and restriction have Euclidean norm one, so FMC22 proves the remaining operator bounds. Its left-inverse identity then proves the lower bound on Z_k^{fix} .

For every $0 \leq t \leq \Delta$, singular values now give

$$\log^+ \left\| \bigwedge^t T_k \right\|_2, \log^+ \left\| \bigwedge^t T_k^{-1} \right\|_2 \leq t \log M_k \leq \Delta \log M_k = O_{\text{actual}}(k^2) = o(kq). \quad (\text{FMC25})$$

Indeed every singular value of both square matrices is at most M_k , and exterior singular values are their products. This last fact follows by applying a singular-value decomposition: unitary factors preserve exterior norms, and the diagonal factor multiplies each coordinate wedge by the corresponding product. Explicitly

$$\log M_k = \log 2 + \frac{1}{2} \log \Delta + k \log h_* + 10k \log A_0 + L_k \log(C_H C_S). \quad (\text{FMC26})$$

All data except k, Δ, L_k are fixed actual numbers, $L_k \leq B_0^4 k/5 + 624$, $\Delta = 16k - 48$, and $kq = k(k+1)^2$. These formulas prove the last two statements of FMC25 directly. The same bounds hold for transpose inverse maps, whose singular values are unchanged. For $0 \leq t \leq j$, FMC24 also puts every singular value of Z_k^{fix} between M_k^{-1} and M_k , so the same exterior bounds apply to its map onto its image and its inverse there.

38.6 Exact relation to every earlier certificate frame

Let Z_k^{old} be the full actual OCF/ROQ certificate frame, with its selected invertible $j \times j$ row block $(Z_k^{\text{old}})_I$. Both frames have exactly the same column space, as proved in FMC21–22. Their exact transition and inverse are

$$A_k = ((Z_k^{\text{old}})_I)^{-1} (Z_k^{\text{fix}})_I, \quad A_k^{-1} = L_{Z,k} Z_k^{\text{old}}, \quad Z_k^{\text{fix}} = Z_k^{\text{old}} A_k. \quad (\text{FMC27})$$

Each new column has a unique expansion in the old independent columns; restriction to the old pivot rows gives the first formula. Applying $L_{Z,k}$ gives the second and both inverse identities. Thus this is the exact coefficient morphism to the earlier frame, including its actual pivot determinant. No bound for that old determinant is needed or asserted by FMC24–26. The new frame and its proved bounds use only the fixed data of FMC5–23.

The two residual maps on V_A therefore obey $\Xi_k^{\text{fix}} = A_k^T \Xi_k^{\text{old}}$ in their respective coordinates, since the maps in the original strip dual are the ordinary transposes of their certificate matrices.

For every original source and cutoff their minimum quotient metrics satisfy the exact congruence and determinant formulas

$$Q_{\Xi,N}^{\text{fix}} = \overline{A_k}^{-1} Q_{\Xi,N}^{\text{old}} A_k^{-\text{T}}, \quad \log \det Q_{\Xi,N}^{\text{fix}} = \log \det Q_{\Xi,N}^{\text{old}} - 2 \log |\det A_k|. \quad (\text{FMC28})$$

The constraints on a lift are transformed by precisely A_k^{T} ; substituting the inverse coordinates in the same attained minimum proves the congruence, and determinants give the second formula. The scalar is independent of N and of source order. Its coefficient is exactly $1 + 1 - 1 - 1 = 0$ in the original four-endpoint return. Thus this return agrees identically with the entire earlier ROQ/RCA return, even if an earlier selected pivot is small at some degrees.

The complete inherited period Gram is transported by the same exact map at each endpoint. Let L_{old} be the primary right section ROQ6 and let L_{fix} be any primary right section of Ξ_k^{fix} inside V_A , for example the first j columns of FMC30 below. Both L_{fix} and $L_{\text{old}} A_k^{-\text{T}}$ have residual image I_j under Ξ_k^{fix} , so their difference has columns in the original $K = \ker \Lambda$. Consequently

$$\Lambda L_{\text{fix}} = \Lambda L_{\text{old}} A_k^{-\text{T}}, \quad P_{\text{per}}^{\text{fix}} = \overline{A_k}^{-1} P_{\text{per}}^{\text{old}} A_k^{-\text{T}}, \quad \frac{\det Q_{\Xi,N}^{\text{fix}}}{\det P_{\text{per}}^{\text{fix}}} = \frac{\det Q_{\Xi,N}^{\text{old}}}{\det P_{\text{per}}^{\text{old}}}.$$

The first equality preserves the literal boundary vectors and their full cyclic coefficient map. Taking their Gram with the unchanged inherited metric ROQ9 proves the second equality. Taking both determinants together with FMC28 proves the last equality pointwise, before taking any endpoint difference.

38.7 The unchanged Gamma source in the fixed degree model

Retain $c = k/2$, the original nodes $\beta_{ab} = c + (2a - k)\delta + i(2b - k)\gamma$, and $\omega_{ab} = (\beta_{ab} - c)/i$. For each original source $s \in \{1, k\}$ retain its complete recurrence and masses

$$p_0 = 1, \quad p_1 = y, \quad p_{n+1} = yp_n - n(s/2 + n - 1)p_{n-1}, \quad (\text{FMC29})$$

$$h_n = (2\pi)^{s/2} n! (s/2)_n, \quad (U_N)_{(a,b),n} = p_n(\omega_{ab}) / \sqrt{h_n}, \quad G_N = (U_N U_N^*)^{-1}.$$

Here N remains one of $q - 1, q, 2q - 1, 2q$. These are exactly the original primary quotient metrics of CEP21–22 and ROQ10, with every source mass and coordinate phase present.

The complete primary frame of V_A , in residual coordinates first and common-kernel coordinates second, is

$$J_k = N_k^{\text{T}} T_k^{-\text{T}}, \quad \Xi_k^{\text{fix}} J_k = E_j^{\text{T}}, \quad K = \text{im}(J_k E_r). \quad (\text{FMC30})$$

Here E_r inserts the last r coordinates. For a primary vector $x \in V_A$, ROQ3 gives its strip dual coordinate $z = S_k^{\text{T}} x$. The new residual map is $(Z_k^{\text{fix}})^{\text{T}} z = E_j^{\text{T}} T_k^{\text{T}} z$. Substitution of $z = T_k^{-\text{T}} y$ proves the second identity, and its kernel is the last coordinate block, proving the third. Thus this full frame has precisely the original common kernel; it changes no observation. Its left inverse on V_A is $T_k^{\text{T}} S_k^{\text{T}}$. The original coefficient bounds additionally give

$$\|J_k\|_2 \leq \sqrt{q} A_0^{10k} M_k =: \widehat{M}_k, \quad \|T_k^{\text{T}} S_k^{\text{T}}\|_2 \leq M_k. \quad (\text{FMC31})$$

Indeed FMC14 bounds the norm of N_k by its column-sum norm times $\sqrt{q}$, transpose preserves Euclidean norm, and FMC24 applies to $T_k^{-\text{T}}$. The insertion transpose is a coordinate restriction of norm one. Both logarithms in FMC31 are $O_{\text{actual}}(k + \log k)$ with their full displayed constants.

The exact source Gram in this specified degree model, and its actual residual quotient, are

$$\mathcal{H}_N = J_k^* G_N J_k = \overline{T_k}^{-1} \overline{N_k} G_N N_k^{\text{T}} T_k^{-\text{T}}, \quad (\text{FMC32})$$

$$\mathcal{H}_N = \begin{pmatrix} H_{11,N} & H_{12,N} \\ H_{21,N} & H_{22,N} \end{pmatrix}, \quad Q_{\Xi,N}^{\text{fix}} = H_{11,N} - H_{12,N} H_{22,N}^{-1} H_{21,N}.$$

The first formula is direct substitution of the actual primary frame. Since $G_N > 0$ and J_k is injective, $\mathcal{H}_N > 0$; therefore $H_{22,N} > 0$. For prescribed first coordinate y , the second coordinate minimizing the full quadratic form is $-H_{22,N}^{-1}H_{21,N}y$. Completing the square, or multiplying out this substitution, gives exactly the displayed Schur complement. This is the original minimum on $\Xi_k^{\text{fix}}x = y$ inside V_A , so it equals FMC28 and the full ROQ metric. All four blocks, including every off-diagonal entry, are retained. The original minimum source lift of the full frame is explicitly $U_N^*G_NJ_k$, whose Gram is $\mathcal{H}_N$ by $U_NU_N^* = G_N^{-1}$. It contains the entire original polynomial recurrence and remainder at each stated cutoff.

There is a useful quantitative determinant comparison which does not replace that Gamma form by a coefficient form. In strip dual coordinates write $H_{V,N} = \overline{\mathbf{N}}_k G_N \mathbf{N}_k^T$ and $Z = Z_k^{\text{fix}}$. Define the actual coefficient quotient form and the relative determinant by

$$Q_{\text{coeff}} = (Z^T \overline{Z})^{-1}, \quad \Phi_{\Xi,N} = \log \det Q_{\Xi,N}^{\text{fix}} - \log \det Q_{\text{coeff}}. \quad (\text{FMC33})$$

The first is the minimum quotient of the original coefficient Euclidean norm in the fixed OCF strip, with the identical constraint $Z^T z = y$. The elementary minimum proof of ROQ11 gives both this formula and $Q_{\Xi,N}^{\text{fix}} = (Z^T H_{V,N}^{-1} \overline{Z})^{-1}$. Hence the full, directly calculable expression is

$$\begin{aligned} \Phi_{\Xi,N} &= \log \det(Z^T \overline{Z}) - \log \det(Z^T H_{V,N}^{-1} \overline{Z}), \\ |\log \det Q_{\text{coeff}}| &\leq 2j \log M_k = O_{\text{actual}}(k^2) = o(kq), \\ F_{\Xi,k}^{(s)} &= \Phi_{\Xi,q-1} + \Phi_{\Xi,q} - \Phi_{\Xi,2q-1} - \Phi_{\Xi,2q}. \end{aligned} \quad (\text{FMC34})$$

FMC24 puts the singular values of Z between M_k^{-1} and M_k . The positive eigenvalues of $Z^T \overline{Z}$ are their squares, proving the determinant bound by summing j logarithms. The first identity follows by the two exact inverse Gram formulas. The final identity follows by canceling the fixed $\log \det Q_{\text{coeff}}$ with the four original signs and then using FMC28. Thus all coefficient denominators and frame-volume effects in this specified construction are controlled below the kq scale, while the original Gamma compression remains fully visible in its exact determinant.

For the kernel frame $I_k = J_k E_r$, FMC31 likewise gives the concrete coefficient-Gram bounds

$$M_k^{-2r} \leq \det(I_k^* I_k) \leq \widehat{M}_k^{2r}, \quad |\log \det(I_k^* I_k)| \leq 2r \log \widehat{M}_k = O_{\text{actual}}(k^2) = o(kq). \quad (\text{FMC35})$$

Its left inverse on its image is $E_r^T T_k^T S_k^T$, with norm at most M_k ; FMC31 bounds its forward norm. The singular-value argument just given proves FMC35. Define the exact primary-coordinate projection and determinant by

$$P_{K,\text{prim}} = I_k(I_k^* I_k)^{-1} I_k^*, \quad D_{K,N}^{\text{prim}} = [z^r] \det(I_q + z P_{K,\text{prim}} G_N P_{K,\text{prim}}).$$

Multiplication proves that this projection is Hermitian, idempotent, and has exactly the original primary-coordinate K as its image. For the determinant proof take $U = I_k(I_k^* I_k)^{-1/2}$, so $U^* U = I_r$, and extend its columns to an orthonormal basis of $\mathbb{C}^q$. In that basis the projected metric has block $U^* G_N U$ and a zero block. Its indicated determinant coefficient is $\det(U^* G_N U)$, and substitution of the full Gram square root therefore proves

$$\det(I_k^* G_N I_k) = \det(I_k^* I_k) D_{K,N}^{\text{prim}}.$$

The precise map to AKS12–14's original polynomial-coefficient projector is the ordered Vandermonde: put $I_{k,\text{poly}} = V_\beta^{-1} I_k$ and $G_{N,\text{poly}} = V_\beta^* G_N V_\beta$. Their compressed Gram is identically $I_k^* G_N I_k$ by multiplication. Applying the determinant proof just given in these polynomial coordinates proves the exact bridge

$$D_{K,N}^{\text{AKS}} = \frac{\det(I_k^* I_k)}{\det(I_{k,\text{poly}}^* I_{k,\text{poly}})} D_{K,N}^{\text{prim}}.$$

Here the polynomial projector has image the actual coefficient kernel and thus agrees with AKS12's projector. The displayed positive factor is independent of N and source order and cancels exactly with the four endpoint signs. FMC35 bounds its primary-coordinate numerator; the polynomial denominator is retained as the full displayed Vandermonde Gram. No bound for that different denominator follows from FMC35 alone.

FMC25 is a coefficient-space estimate for specified exact maps. Its relation to the original Gamma norms is the complete congruence and Schur calculation FMC29–34, not an assertion that those maps are Gamma isometries. In particular no upper or lower leading value for the remaining Gamma compression is inferred from their coefficient condition numbers. The new result supplies a fixed degree presentation, actual invariant certificates, bounded forward and inverse exterior maps, and the exact original determinant on that model, with every fixed denominator specified and every source order and endpoint retained.

38.8 The exact map to the original polynomial degree flag

The homogeneous degree k in FMC3 is the original tensor degree. Its precise map to the original polynomial-in- S model is the ordered evaluation isomorphism, not an identification of the two degree indices. Retain $\chi_k(S) = \prod_{a,b=0}^k (S - \beta_{ab})$ and the original Vandermonde $(V_\beta)_{(a,b),d} = \beta_{ab}^d$, $0 \leq d < q$. For a fixed-model primal coordinate column $y \in \mathbb{C}^\Delta$ the exact polynomial representative is

$$p_y(S) = \sum_{a,b=0}^k (J_k y)_{ab} \frac{\chi_k(S)}{(S - \beta_{ab}) \chi'_k(\beta_{ab})}, \quad [p_y]_{\text{coeff}} = V_\beta^{-1} J_k y. \quad (\text{FMC36})$$

All denominators are nonzero because the original nodes are distinct. Evaluation at each node proves the displayed inverse and hence the equality of the two formulas. It includes the complete roots, their order, and their actual centre. Thus the source degree- N metric in FMC32 is equivalently $(V_\beta^{-1} J_k)^* (V_\beta^* G_N V_\beta) (V_\beta^{-1} J_k)$, identically $J_k^* G_N J_k$ by multiplication. This proves the full typed relation to the original coefficient Gamma model.

There is an exact further map to the degree-adapted conductor basis AKJ17. Retain all actual conductor moments μ_d , their first nonzero index $v \leq 80$, the literal $B_q(\partial_S)$ of AKJ14, and $\mathcal{J}_v S^d = d! S^{d+v} / (d+v)!$. The ordered polynomial columns of $P_{A,k}$ are

$$1, S, \dots, S^{v-1}, \quad P_h = \frac{(q' + h + v)!}{(q' + h)!} B_q(\partial_S)^{-1} \mathcal{J}_v(\chi_{k'} S^h), \quad 0 \leq h \leq \Delta - v - 1. \quad (\text{FMC37})$$

Their distinct monic degrees are $0, \dots, v-1$ and $q' + v, \dots, q-1$. AKJ14–17 proves directly that they form a basis of this same V_A , using the complete conductor derivative and its finite nilpotent inverse. Define $B_{A,k}(p)$ on V_A by successively subtracting the coefficient of the highest remaining degree times its matching column in FMC37, down through the high list and then through the low monomials. Each subtraction uses leading coefficient one. Spanning and the distinct degree pivots show that it ends at zero and records the unique coordinate column of p ; this also proves its linearity and both inverse identities on V_A . Consequently the precise transition is

$$R_{A,k} = S_k^\top V_\beta P_{A,k}, \quad R_{A,k}^{-1} z = B_{A,k}(V_\beta^{-1} \mathbf{N}_k^\top z), \quad (\text{FMC38})$$

$$V_\beta^{-1} J_k = P_{A,k} R_{A,k}^{-1} T_k^{-\top}, \quad \Xi_{A,k}^{\text{fix}} = E_j^\top T_k^\top R_{A,k}.$$

The columns $V_\beta P_{A,k}$ lie in V_A . Therefore ROQ3 gives $\mathbf{N}_k^\top R_{A,k} = V_\beta P_{A,k}$, which proves the first inverse formula by applying the unique degree coordinates just constructed. Both sides are invertible maps between the displayed Δ -dimensional coefficient spaces. Substituting FMC30 gives the third equality, and evaluating the new residual map on these same primary columns gives the last. Thus its degree-prefix ranks give exactly $\dim(K \cap \mathbb{C}[S]_{\leq d})$ by subtracting that prefix rank from the number of columns of FMC37 of degree at most d , as in AKJ18. Every original factorial,

finite inverse, primary denominator and degree pivot is visible in these maps. The subleading coefficient-frame estimate proved in FMC25 pertains to the fixed strip module maps; FMC36–38 retain the separate polynomial interpolation and its full Gamma metric exactly.

39 The evaluated outer contribution in the original mixed return

Retain the actual simple quartet, original period on the proved five-orbit locus, full amplitude unit, branch, and original source orders $s = 1, k$ of OAR and OFV. All arithmetic statements retain the original domain $|u| \geq R_*$ of AMT; the actual period is fixed as $k = 4l + 1$ grows. Here $q = (k + 1)^2$, $\Delta = 16k - 48$, and every finite use of OFV retains its stated threshold, including OSP23. Write

$$C_{\partial} = \mathfrak{C}_{\partial} = 4 \log(4/\pi) + 2 - 2I_{2,\pi}, \quad I_{2,\pi} = \iint \log |x - y| d\mu_{2,\pi}(x) d\mu_{2,\pi}(y). \quad (\text{OVR1})$$

The measure here is the full EIQ density with its uniquely specified endpoints. OFV proves this exact expression and $C_{\partial} \geq 0$. No value of the original residual observation is inserted into it.

39.1 The exact remainder and the sourcewise leading identity

Define the actual outer remainder and retain OAR3's full conductor and monic-tail remainder by

$$b_{k,s} = \mathcal{F}_{\partial,k}^{(s)} - \Delta q C_{\partial}, \quad R_{k,s} = b_{k,s} + \mathfrak{e}_{k,s}. \quad (\text{OVR2})$$

Thus $b_{k,s} = O_{\delta,\gamma}(q \log q)$ by OFV30. The second summand is precisely the signed finite expression OAR3, with $-e_{k,s}^- \leq \mathfrak{e}_{k,s} \leq e_{k,s}^+$ and $e_{k,s}^{\pm} = o(kq)$. In particular $R_{k,s} = o(kq)$, because $\log q/k \rightarrow 0$. Its two exact finite error endpoints remain $b_{k,s} - e_{k,s}^-$ and $b_{k,s} + e_{k,s}^+$; their signs are not replaced by a symmetry assumption. Substitution into OAR3 proves

$$\begin{aligned} F_K^{(s)} + \Phi_{k,s} &= \Delta q C_{\partial} + R_{k,s}, \\ \frac{F_K^{(s)}}{kq} + \frac{\Phi_{k,s}}{kq} &= 16C_{\partial} - \frac{48C_{\partial}}{k} + \frac{R_{k,s}}{kq} \longrightarrow 16C_{\partial}, \quad s = 1, k. \end{aligned} \quad (\text{OVR3})$$

Here $\Phi_{k,s}$ is exactly ROQ19's positive sum of constrained covariance increments, built from the original period certificate matrix D and the complete conductor cross covariance. Its coefficient maps, minimum sections, inherited period metric and graph denominator are those already proved in ROQ; OVR3 changes none of them.

39.2 Transfer between the two original Gamma sources

Let $d_k = F_K^{(k)} - F_K^{(1)}$ denote the actual difference. AMT44–46 proves $d_k = o(kq)$, with its original finite two-sided source-comparison bounds. Subtracting the two exact lines OVR3 gives

$$\Phi_{k,k} - \Phi_{k,1} = R_{k,k} - R_{k,1} - d_k, \quad \frac{\Phi_{k,k} - \Phi_{k,1}}{kq} \longrightarrow 0. \quad (\text{OVR4})$$

This proves the transfer using the actual two residual observations, whose individual metrics retain their respective original sources. There is no asserted equality between those metrics. The equality in OVR4 is the exact scalar consequence of their common original-kernel restriction maps and the proved determinant identities.

39.3 Arithmetic kernel and its correlated boundary

Put $d_k^{\text{ar}} = F_K^{\text{ar}} - F_K^{(1)}$. This is the actual quantity bounded in OAR8; AMT38 and AMT46 prove $d_k^{\text{ar}} = o(kq)$. Keeping its full value first gives

$$\begin{aligned} F_K^{\text{ar}} &= \Delta q C_{\partial} - \Phi_{k,1} + R_{k,1} + d_k^{\text{ar}}, \\ F_B^{\text{ar}} &= \mathcal{B}_k^{\text{ar}} - \Delta q C_{\partial} + \Phi_{k,1} - R_{k,1} - d_k^{\text{ar}}. \end{aligned} \quad (\text{OVR5})$$

The first equality adds the literal definition of d_k^{ar} to OVR3 at $s = 1$. The second subtracts the first from $F_K^{\text{ar}} + F_B^{\text{ar}} = \mathcal{B}_k^{\text{ar}}$. In particular the same arithmetic total and the same two finite errors

occur with opposite signs in the boundary. OAR8 supplies the unchanged correlated finite bounds on d_k^{ar} . Dividing these exact equalities gives

$$\frac{F_K^{\text{ar}} + \Phi_{k,1}}{kq} \longrightarrow 16C_\partial, \quad \frac{\mathcal{B}_k^{\text{ar}} - F_B^{\text{ar}} + \Phi_{k,1}}{kq} \longrightarrow 16C_\partial. \quad (\text{OVR6})$$

Thus evaluation of the residual sum in either original Gamma order passes through to the same original arithmetic kernel and boundary.

39.4 The signed return retains its original Gamma correction

The exact signed quantity of OAR11 is $\mathfrak{S}_k^{\text{mix}} = F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)}$. Applying OVR3 at $s = k$ to that identity yields

$$\mathfrak{S}_k^{\text{mix}} + \Phi_{k,k} = \Delta q C_\partial + \mathcal{H}_k + \delta_k^\sigma + R_{k,k}. \quad (\text{OVR7})$$

All finite terms are the original ones. AMT48–49, with the retained total correction estimates used there, proves $(\mathcal{H}_k + \delta_k^\sigma)/(kq) \rightarrow -C_\Gamma/4$, where

$$C_\Gamma = 2 \int \log x \, d\mu_{2,\pi}(x) - 4(2 \log 2 - 1) > 0.$$

The equilibrium and source maps are exactly the EIQ/GEL providers used in OFV and AMT. Hence OVR7 and OVR4 prove both limits

$$\frac{\mathfrak{S}_k^{\text{mix}} + \Phi_{k,k}}{kq} \longrightarrow 16C_\partial - \frac{C_\Gamma}{4}, \quad \frac{\mathfrak{S}_k^{\text{mix}} + \Phi_{k,1}}{kq} \longrightarrow 16C_\partial - \frac{C_\Gamma}{4}. \quad (\text{OVR8})$$

The two constants are explicit integrals of the same proved density, respectively its full logarithmic pair energy and its logarithmic moment. No identity between these distinct integrals is presumed.

39.5 Bounds on the remaining residual coefficient

Each $F_K^{(s)}$ is nonnegative: the original quotient metrics decrease with increasing source degree, and restriction to the identical kernel preserves the two Gram inequalities at the paired endpoints. Their determinant ratios are consequently at least one, as proved in AMT15. ROQ19 gives $\Phi_{k,s} \geq 0$. Thus OVR3 bounds both nonnegative sequences after division by kq . Let $\kappa_* = 32 \log(25 \log 3 / (4\pi))$ be the retained AKS asymptotic kernel ceiling on this actual locus, applied in both original orders. The actual residual cap OAR5, proved by ROQ21, is $\Phi_{k,s} \leq j(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - \lambda_s$, where $j = 8k - 32$ and $\lambda_s \geq 0$ is the original first increment. AKS31 gives $\limsup j(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1})/(kq) \leq \kappa_*$ in both original source orders: its same explicit logarithmic comparison constants apply, and $j/k \rightarrow 8$ just as $r/k \rightarrow 8$. Thus both the original kernel and residual sequences have the same proved upper ceiling. Combining their two ceilings with OVR3 gives

$$\begin{aligned} \max\{0, 16C_\partial - \kappa_*\} &\leq \liminf \frac{F_K^{(s)}}{kq} \leq \limsup \frac{F_K^{(s)}}{kq} \leq \min\{\kappa_*, 16C_\partial\}, \\ \max\{0, 16C_\partial - \kappa_*\} &\leq \liminf \frac{\Phi_{k,s}}{kq} \leq \limsup \frac{\Phi_{k,s}}{kq} \leq \min\{\kappa_*, 16C_\partial\}. \end{aligned} \quad (\text{OVR9})$$

For each line, subtract the other bounded nonnegative sequence from the convergent sum in OVR3. More explicitly, that sum differs from $16C_\partial$ by a sequence tending to zero; taking its upper and lower eventual bounds proves $\liminf(\Phi/(kq)) = 16C_\partial - \limsup(F_K/(kq))$ and $\limsup(\Phi/(kq)) = 16C_\partial - \liminf(F_K/(kq))$. The same argument with the two sequences interchanged proves the first line, including its lower bound. The two first positive increments also survive in the finite common

upper bound. In the original AKS notation let d_1 be the first section-corrected kernel increment of AKS29, in the same source order s . AKS28 and OAR5 give

$$\begin{aligned}\Delta q C_{\partial} + R_{k,s} &= F_K^{(s)} + \Phi_{k,s} \\ &\leq (r+j)(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - d_1 - \lambda_s \\ &= \Delta(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - d_1 - \lambda_s.\end{aligned}\tag{OVR10}$$

The last equality is the exact rank identity $(8k-16) + (8k-32) = 16k-48$; both subtractions retain their actual minimum-section definitions. Adding the eventual upper bounds already proved and using OVR3 also proves the explicit consequence

$$0 \leq 16C_{\partial} \leq 2\kappa_*, \quad 0 \leq C_{\partial} \leq \frac{\kappa_*}{8}.\tag{OVR11}$$

Indeed for every positive ε , both sequences are eventually at most $\kappa_* + \varepsilon$, so their limiting sum is at most $2\kappa_* + 2\varepsilon$; let ε decrease to zero. Their nonnegativity proves the lower bound independently. OVR4 and AMT46 carry the same limiting bounds to the two original sources and the arithmetic kernel. The outstanding coefficient is therefore the explicit original positive residual sum; no convergence or numerical value for that sequence has been assumed.

40 All directions of the original outer quotient at the leading scale

Retain the complete original simple quartet, the source orders $s = 1, k$, $k = 4l + 1$, $0 < \delta < 1/2$, $\gamma > 2$, and the polynomials and masses of CTV1–22 and OSP1–35. Put

$$q = (k + 1)^2, \quad q' = (k - 7)^2, \quad d = q - q' = 16k - 48, \quad n = q/2, \quad \vartheta = \pi/2, \quad R = k\sqrt{\delta^2 + \gamma^2}. \quad (\text{OOQ1})$$

The integer q is even. The entire outer polynomial is $P_{\partial,k} = P_k/P_{k-8}$, monic of degree d , with every original root retained. Each root has modulus at most R . Explicitly $P_k(y) = i^{-q}\chi_k(c + iy)$ and $P_{k-8}(y) = i^{-q'}\chi_{k-8}(c' + iy)$, with the unchanged original centres $c = k/2$ and $c' = c - 4$. These are the exact CTV3–4 phase and coefficient maps. Its target coordinates throughout are the coefficients of $1, y, \dots, y^{d-1}$ in its unique remainder. For $L = d - 1, d, q + d - 1, q + d$, let $Q_{\partial,L}^{(s)}$ be the minimum quotient Gram of this remainder map from polynomials of degree at most L , with the complete source

$$d\zeta_{k',s}(y) = |P_{k-8}(y)|^2 dm_s(y), \quad dm_s(y) = \frac{(2\pi)^{s/2} 2^{s/2}}{4\pi\Gamma(s/2)} |\Gamma(s/4 + iy/2)|^2 dy. \quad (\text{OOQ2})$$

These are exactly the CTV10–13 Grams, not merely determinants equal to those Grams. We prove operator estimates in this same original remainder frame.

40.1 Uniform equilibrium constants and a polynomial evaluation bound

Use the complete EIQ equilibrium for $V_{\alpha,\beta}(x) = \beta\sqrt{x} - \alpha \log x$ on $(0, \infty)$. In this proof take the compact rectangle $\mathcal{K} = [3/2, 5/2] \times [\pi/4, \pi]$. EIQ is proved on the entire positive quadrant, so it applies to this rectangle. Write its exact density as $\rho_{\alpha,\beta}$, its endpoints as a, b , its logarithmic potential as $P(z)$, and retain

$$V(x) - 2P(x) = \ell + D(x), \quad D(x) \geq 0, \quad D|_{[a,b]} = 0, \quad P_0 = P(0) = \int \log t \rho(t) dt, \quad W = P_0 + \ell/2. \quad (\text{OOQ3})$$

Here and below a parameter on these symbols is understood whenever two parameter pairs occur in the same expression.

The following fixed constants are all defined by these exact functions on $\mathcal{K}$:

$$a_* = \min_{\mathcal{K}} a > 0, \quad B_* = \max_{\mathcal{K}} b < \infty, \quad b_* = \min_{\mathcal{K}} (b - a) > 0, \quad M_* = \max_{\mathcal{K},x} \rho(x) < \infty, \\ L_* = \max_{\mathcal{K}, x \in [a,b]} |V'(x)|, \quad C_{\text{BM}} = \sqrt{32} e^{3L_*/4}, \quad C_{\text{quant}} = B_* + 2M_*, \quad C_0 = B_*/a_*, \quad (\text{OOQ4})$$

$$C_{\text{tail}} = \max\{0, \sup_{\mathcal{K}, x>0} [\log(1 + \sqrt{x}) - D(x)]\}, \quad T_* = \max\{1, \sup_{\mathcal{K}} \int_0^\infty e^{-D(x)} dx\}.$$

These constants are finite. Positivity, continuity and boundedness of the endpoints and density are EIQ4, EIQ10–11 and EIQ29; the positive gap follows by compactness. The derivative of V is bounded on $[a_*, B_*]$. Uniformly over this rectangle, $P(x) \leq \log(1 + x) + \log(1 + B_*)$ and $P(x) \geq -2M_*$, as in the direct logarithmic estimates in QLG9–13. The Euler constant is bounded by evaluating OOQ3 on the compact support. Consequently

$$D(x) \geq \frac{\pi}{4}\sqrt{x} - \frac{5}{2}\log x - 2\log(1 + x) - C \quad (x \geq 1)$$

for a fixed finite constant C . This proves the assertions at infinity in the last two definitions of OOQ4. At zero, $D(x) \rightarrow +\infty$ uniformly because $\alpha \geq 3/2$ and P is bounded there. For integrability one can use precisely the finite majorant $e^C(1 + x)^2(1 + x^{5/2})e^{-\pi\sqrt{x}/4}$. Thus neither of the two supremum definitions assumes a tail estimate.

The function W and its first derivatives are bounded on $\mathcal{K}$. Indeed EIQ30 gives $F_\alpha = P_0$, EIQ31 gives $\beta \int \sqrt{x} d\mu = 2(\alpha + 1)$, and hence

$$W(\alpha, \beta) = -F(\alpha, \beta) - (\alpha + 1) + (1 + \alpha/2)F_\alpha(\alpha, \beta). \quad (\text{OOQ5})$$

The smoothness asserted here follows from EIQ29–30. Fix

$$C_W = 1 + \max_{\mathcal{K}} |W| + (4 + \pi) \max_{\mathcal{K}} (|W_\alpha| + |W_\beta|). \quad (\text{OOQ6})$$

For an integer $m \geq \max\{1, 2/b_*\}$ and a polynomial p of degree at most m , set $\|p\|_{m,V}^2 = \int_0^\infty |p(x)|^2 e^{-mV(x)} dx$. For all z outside $[a, b]$ we have

$$|p(z)| \leq C_{\text{BM}} m^2 \exp\{m(P(z) + \ell/2)\} \|p\|_{m,V}. \quad (\text{OOQ7})$$

Here is a complete proof. Let $M = \max_{[a,b]} |p(x)| e^{-mV(x)/2}$, attained at x_0 . Choose a one-sided interval starting at x_0 , contained in $[a, b]$, of length $1/m$; one side has that length because $b - a \geq 2/m$. On that interval, $|p(x)| \leq e^{L_*/2} |p(x_0)|$.

The elementary derivative bound $\|p'\|_{\infty,J} \leq 4m^3 |J|^{-1} \|p\|_{\infty,J}$ holds also for complex coefficients. To prove it, expand on the affine copy of $[-1, 1]$ in the polynomials T_j defined by $T_j(\cos \theta) = \cos(j\theta)$. Orthogonality of cosines gives coefficients of modulus at most $2\|p\|_\infty$ for $j \geq 1$. The identity $T'_j(\cos \theta) = j \sin(j\theta)/\sin \theta$ and the sum of j unit complex numbers give $|T'_j| \leq j^2$, including the limiting endpoints. Summing $\sum_{j=1}^m j^2 \leq m^3$ and applying the affine derivative factor $2/|J|$ proves the bound. It follows that on a segment of length $[8e^{L_*/2} m^4]^{-1}$ starting at x_0 , $|p| \geq |p(x_0)|/2$. The weight on that segment is at least $e^{-L_*} e^{-mV(x_0)}$. Integration gives $\|p\|_{m,V}^2 \geq M^2/(32e^{3L_*/2} m^4)$.

It remains to extend the weighted supremum to complex z . The function $\log |p(z)| - mP(z)$ is subharmonic off $[a, b]$. On this interval its boundary upper values are at most $\log M + m\ell/2$ by OOQ3. It extends subharmonically over infinity: in the local coordinate $w = 1/z$ its logarithmic term is $(m - \deg p) \log |w|$, plus a harmonic function. The maximum principle on the complement of the interval in the sphere therefore bounds it by the same boundary constant. One may first remove small neighborhoods of polynomial zeros and of the interval and then pass to the limit; P is continuous because its density is bounded. Combining this bound with the preceding norm bound proves OOQ7. For $|z| \leq a_*/2$ its useful uniform consequence is

$$|p(z)| \leq C_{\text{BM}} m^2 e^{mW+2m|z|/a_*} \|p\|_{m,V}, \quad (\text{OOQ8})$$

since $|\log |1 - z/t|| \leq 2|z|/a_*$ on the support.

40.2 A matching polynomial lift from actual equilibrium quantiles

For each n divide the actual equilibrium measure into n intervals of mass $1/n$ and take the midpoint of each interval as ξ_i . Retain the polynomial

$$r_n(x) = \prod_{i=1}^n (1 - x/\xi_i), \quad r_n(0) = 1. \quad (\text{OOQ9})$$

All its roots lie in $[a_*, B_*]$. For every real $x > 0$,

$$\log |r_n(x)| \leq n(P(x) - P_0) + C_{\text{quant}} \sqrt{n} + C_0. \quad (\text{OOQ10})$$

To prove it, use $f_\eta(t) = \log \max\{|x - t|, \eta\}$, a $1/\eta$ -Lipschitz function. For a quantile interval of length Δ_i , the difference between its point value and the average against its probability density is at most Δ_i/η . Summing gives at most B_*/η . Also

$$0 \leq \int [f_\eta(t) - \log |x - t|] \rho(t) dt \leq M_* \int_{-\eta}^{\eta} \log(\eta/|s|) ds = 2M_* \eta.$$

Choose $\eta = n^{-1/2}$. Finally $\log t$ is $1/a_*$ -Lipschitz on the support, so $|\sum_i \log \xi_i - nP_0| \leq B_*/a_*$. These three facts prove OOQ10, including at a polynomial zero. For complex $|z| \leq a_*/2$, the same literal product gives

$$|r_n(z)|^{-1} \leq e^{2n|z|/a_*}. \quad (\text{OOQ11})$$

This uses $|1 - z/\xi_i| \geq \exp(-2|z|/a_*)$ separately for every original quantile factor.

40.3 The entire analytic remainder map

For the present $n = q/2$ put

$$\rho = n^{-1/8}, \quad \hat{P}(z) = n^{-d}P_{\partial,k}(nz), \quad r_* = R/n. \quad (\text{OOQ12})$$

Impose the following explicit eventual conditions, together with the original OSP23 domain and $k \geq 257$:

$$n \geq 2d + 1, \quad n \geq 2/b_*, \quad r_* \leq \rho/2, \quad \rho \leq \min\{1/2, \sqrt{a_*/2}\}. \quad (\text{OOQ13})$$

They hold for all sufficiently large original k at every fixed δ, γ : $n \asymp k^2$, $d = O(k)$ and $r_* = O(1/k)$. These inequalities specify the finite range used here; no assertion at excluded small degrees is required.

For f holomorphic on the closed disk of radius ρ define its complete remainder by

$$\text{rem}_{\hat{P}} f(z) = \frac{1}{2\pi i} \int_{|\zeta|=\rho} f(\zeta) \frac{\hat{P}(\zeta) - \hat{P}(z)}{(\zeta - z)\hat{P}(\zeta)} d\zeta. \quad (\text{OOQ14})$$

The numerator divided by $\zeta - z$ is a polynomial in z of degree at most $d - 1$. At every root of $\hat{P}$, Cauchy's formula gives the value of f ; all these original roots are distinct, so this is exactly the unique interpolation remainder. In particular it is the usual polynomial remainder when f is polynomial. This proves the exact map and its agreement with the original target.

Write $\|\cdot\|_1$ for the full monomial coefficient norm. The root bound in OOQ13 proves

$$\|\text{rem}_{\hat{P}} f\|_1 \leq C_{\text{rem}} \max_{|\zeta|=\rho} |f(\zeta)|, \quad C_{\text{rem}} = \rho d(1 + r_*)^d (2/\rho)^d. \quad (\text{OOQ15})$$

Indeed $|\hat{P}(\zeta)| \geq (\rho - r_*)^d \geq (\rho/2)^d$. If $\hat{P}(z) = \sum_{a=0}^d p_a z^a$, then

$$\frac{\hat{P}(\zeta) - \hat{P}(z)}{\zeta - z} = \sum_{a=1}^d p_a \sum_{h=0}^{a-1} \zeta^{a-1-h} z^h,$$

whose coefficient norm is at most $d \sum_a |p_a|$ because $\rho \leq 1$. The sum of absolute coefficients of the full monic product is at most $(1 + r_*)^d$. The contour length supplies the factor ρ , proving OOQ15. In particular $\log C_{\text{rem}} = O(d \log n + \log d)$. No outer root has been moved to zero in this map.

40.4 High source degrees: a full operator estimate

First use the precise exponential source $|y|^{2Q} e^{-\beta|y|} dy$, where $Q = q' + \ell_s$, $\ell_s = (s - 1)/4$, and $\beta \in \{\vartheta - 1/q, \vartheta + 1/q\}$. Let $Q_L^{\text{exp}}(Q, \beta)$ be its quotient Gram by the same entire $P_{\partial,k}$, in the original y remainder frame. Set

$$\alpha_0 = (Q - \tfrac{1}{2})/n, \quad c_{\text{high}}(Q, \beta) = (2Q + 1) \log n - 2nW(\alpha_0, \beta). \quad (\text{OOQ16})$$

The two high source degrees are $L = q + d - 1, q + d$. For their two parity degrees put $m_e = \lfloor L/2 \rfloor$, $m_o = \lfloor (L - 1)/2 \rfloor$. The positive-half-line parameters are exactly

$$(\alpha_e, \beta_e) = ((Q - \tfrac{1}{2})/m_e, n\beta/m_e), \quad (\alpha_o, \beta_o) = ((Q + \tfrac{1}{2})/m_o, n\beta/m_o). \quad (\text{OOQ17})$$

All these pairs and (α_0, β) lie in $\mathcal{K}$ for the stated $k \geq 257$. For example $d/q < 1/16$, $\ell_s/q < 1/64$ and the rounding errors are at most one; these give $3/2 < \alpha_e, \alpha_o < 5/2$. The rate lies between $\pi/4$ and π because $m_e, m_o = n + O(d)$ and $1/q < \vartheta/4$. More directly the actual lower bounds $m_e, m_o \geq n$ and upper bounds $m_e, m_o \leq n + d$ together with $d/q < 1/16$ verify the rate bounds.

For a polynomial $p(nz) = p_e(z^2) + zp_o(z^2)$, parity and the two substitutions $z = \pm\sqrt{x}$ give exactly

$$n^{-(2Q+1)} \|p\|_{\text{exp}}^2 = \int_0^\infty |p_e(x)|^2 x^{Q-1/2} e^{-n\beta\sqrt{x}} dx + \int_0^\infty |p_o(x)|^2 x^{Q+1/2} e^{-n\beta\sqrt{x}} dx. \quad (\text{OOQ18})$$

Both measures on the right are precisely the OOQ7 measures at OOQ17. There is no missing factor two: each branch contributes the positive Jacobian $1/(2\sqrt{x})$. The smooth bound OOQ6 gives

$$|m_e W(\alpha_e, \beta_e) - nW(\alpha_0, \beta)|, |m_o W(\alpha_o, \beta_o) - nW(\alpha_0, \beta)| \leq C_W(d+2). \quad (\text{OOQ19})$$

For verification, $|m - n| \leq d + 1$, $m|\alpha - \alpha_0| \leq 1 + (5/2)|m - n|$ and $m|\beta_m - \beta| \leq \pi|m - n|$. The mean-value formula on the rectangle and OOQ6 prove OOQ19.

Apply OOQ8 at $x = \zeta^2$ and $|\zeta| = \rho$, then use OOQ18 and Cauchy–Schwarz on its two norms. If v is the coefficient vector of the remainder of $p(nz)$ modulo $\hat{P}$, OOQ15 gives

$$\|v\|_2 \leq C_{\text{rem}} \sqrt{2} C_{\text{BM}}(n+d)^2 e^{nW+C_W(d+2)+2(n+d)\rho^2/a_*} n^{-(2Q+1)/2} \|p\|_{\text{exp}}. \quad (\text{OOQ20})$$

This holds for every lift, and proves the lower bound for the quotient metric in these scaled remainder coordinates.

For the upper bound use the actual equilibrium at (α_0, β) to form r_n in OOQ9. For the given remainder polynomial $v(z)$ put

$$a_v(z) = \text{rem}_{\hat{P}} \left(\frac{v(z)}{r_n(z^2)} \right), \quad p_v(nz) = r_n(z^2) a_v(z). \quad (\text{OOQ21})$$

The denominator has no zero in the disk because all its x -roots are at least $a_* > \rho^2$. The polynomial p_v has degree at most $q + d - 1 \leq L$. At every original outer root its value is exactly v ; thus it is an actual lift of the prescribed remainder. Equations OOQ11 and OOQ15 give

$$\|a_v\|_1 \leq C_{\text{rem}} \sqrt{d} e^{2n\rho^2/a_*} \|v\|_2. \quad (\text{OOQ22})$$

For each real $x > 0$ both $|a_v(\sqrt{x})|$ and $|a_v(-\sqrt{x})|$ are at most $\|a_v\|_1 (1 + \sqrt{x})^{d-1}$. By OOQ4, $(1 + \sqrt{x})^{2(d-1)} \leq e^{2C_{\text{tail}}(d-1)} e^{2(d-1)D(x)}$. Combine this with OOQ10 and $2nP - nV = -n\ell - nD$. Since $n - 2(d-1) \geq 1$, integration using T_* proves

$$\|p_v\|_{\text{exp}}^2 \leq e^{\text{Chigh}(Q, \beta)} \|a_v\|_1^2 \exp\{2C_{\text{quant}}\sqrt{n} + 2C_0 + 2C_{\text{tail}}(d-1)\} T_*. \quad (\text{OOQ23})$$

All source scalings in this expression come from the explicit $y = nz$ substitution; in the original lower-source variable it is precisely $S' = c' + inz$, with the same original centre and phase.

Define the finite nonnegative errors

$$\begin{aligned} E_{\text{high}}^- &= 2 \log \left(C_{\text{rem}} \sqrt{2} C_{\text{BM}}(n+d)^2 \right) + 2C_W(d+2) + 4(n+d)\rho^2/a_*, \\ E_{\text{high}}^+ &= 2 \log C_{\text{rem}} + \log d + 4n\rho^2/a_* + 2C_{\text{quant}}\sqrt{n} + 2C_0 + 2C_{\text{tail}}(d-1) + \log T_*, \\ E_{\text{high}} &= \max\{0, E_{\text{high}}^-, E_{\text{high}}^+\} + 2(d-1) \log n. \end{aligned} \quad (\text{OOQ24})$$

The original and scaled remainder vectors are related by the exact diagonal map $D_n = \text{diag}(1, n, \dots, n^{d-1})$. Hence the original Gram is $D_n^* Q_{\text{scaled}} D_n$. Since $I \preceq D_n^* D_n \preceq n^{2(d-1)} I$, OOQ20–24 prove, at both high endpoints,

$$e^{\text{Chigh}(Q, \beta) - E_{\text{high}}} I_d \preceq Q_L^{\text{exp}}(Q, \beta) \preceq e^{\text{Chigh}(Q, \beta) + E_{\text{high}}} I_d. \quad (\text{OOQ25})$$

The errors obey $E_{\text{high}} = O(n^{3/4} + d \log n)$. This is a full quadratic-form bound for the entire original outer denominator, not an inference from a determinant estimate.

40.5 Low source degrees, including the extra ideal column

Put $g(Q, \beta) = 2Q \log(2Q/\beta) - 2Q$ and $y_0 = 2Q/\beta$. For polynomials of degree at most d , their full exponential source Gram is bounded below by

$$\frac{e^{g(Q, \beta) - \beta}}{(d+1)^2(2d+1)64^d(1+y_0)^{2d}} I_{d+1}. \quad (\text{OOQ26})$$

To prove it, the weight on $[y_0, y_0+1]$ is at least $e^{g(Q, \beta) - \beta}$. On $[0, 1]$ the orthonormal shifted Legendre polynomial of degree j is $\sqrt{2j+1} L_j(2x-1)$. Its coefficient sum is at most $\sqrt{2j+1} 8^j$: Rodrigues' formula gives coefficients $(-1)^{j-r} \binom{j}{r} \binom{j+r}{r}$, and $\binom{j+r}{r} \leq 4^j$. Thus the coefficient matrix of these orthonormal polynomials has Euclidean norm at most $\sqrt{(d+1)(2d+1)} 8^d$. The translation back by $-y_0$ has coefficient norm at most $(1+y_0)^d$ and Euclidean norm at most $\sqrt{d+1}(1+y_0)^d$. These two invertible maps prove OOQ26 by direct norm comparison. The Legendre orthogonality and full Rodrigues norm are proved in OSP15 and are also obtained by its stated integrations by parts.

The largest eigenvalue of the same full Gram is at most its trace, which is exactly

$$\sum_{j=0}^d \frac{2(2Q+2j)!}{\beta^{2Q+2j+1}}. \quad (\text{OOQ27})$$

This follows by the two real half-line integrals and repeated integration by parts. At $L = d-1$ the remainder map is the identity. At $L = d$ it has matrix $[I_d, -p_{\text{low}}]$, where p_{low} contains every lower coefficient of $P_{\partial, k}$. Its squared operator norm is at most $(d+1)(1+R)^{2d}$, because that complete monic polynomial has coefficient sum at most $(1+R)^d$. The minimum Euclidean norm on its fibre is therefore at least the target norm divided by this operator norm; the canonical degree-less-than- d lift gives the upper bound.

For both low endpoints define

$$E_{\text{low}}(Q, \beta) = \max \left\{ 0, \log \left(\sum_{j=0}^d \frac{2(2Q+2j)!}{\beta^{2Q+2j+1}} \right) - g(Q, \beta), \right. \\ \left. \beta + 3 \log(d+1) + \log(2d+1) + d \log 64 + 2d \log(1+y_0) + 2d \log(1+R) \right\}. \quad (\text{OOQ28})$$

Equations OOQ26–27 and the exact remainder matrices prove

$$e^{g(Q, \beta) - E_{\text{low}}} I_d \preceq Q_L^{\text{exp}}(Q, \beta) \preceq e^{g(Q, \beta) + E_{\text{low}}} I_d, \quad L = d-1, d. \quad (\text{OOQ29})$$

In the present range this error is $O_{\delta, \gamma}(d \log q)$. For the upper side, integrating $\log t$ between consecutive integers gives $m \log m - m + 1 \leq \log(m!) \leq m \log m - m + 1 + \log m$. Consequently each logarithm in OOQ27 differs above from $g(Q, \beta)$ by $O(d \log(Q+d) + \log(Q+d))$, uniformly over $\beta \in [\pi/4, \pi]$. The logarithm of the number of summands adds only $\log(d+1)$. The displayed lower error has the same order since $Q \asymp q$, $d = O(k)$ and $R = O_{\delta, \gamma}(k)$.

40.6 Transport back to both complete original Gamma sources

Retain the exact source recurrence factors of OFV1–13:

$$\ell_s = (s-1)/4, \quad \beta_s = \frac{(2\pi)^{s/2}}{\sqrt{2} \Gamma(s/2)}, \quad Q_s = q' + \ell_s, \quad h = 1/q, \\ L_h = C_L (\sin h)^{3/2}, \quad U_h = \frac{\Gamma(1/4)^2}{2\pi} (\sin h)^{-1/2}, \quad C_L = \frac{\pi}{\Gamma(3/4)^2}. \quad (\text{OOQ30})$$

Use $\mathcal{E}_k$ from OSP24 and r_{ℓ_s} from OFV6, with $r_0 = 0$. For every polynomial of all the degrees used here, the complete source comparisons give the quadratic-form inequalities

$$\begin{aligned} \beta_s L_h e^{-\mathcal{E}_k} \int |p(y)|^2 |y|^{2Q_s} e^{-(\vartheta+h)|y|} dy &\leq \int |p(y)|^2 d\zeta_{k',s}(y) \\ &\leq \beta_s U_h e^{\mathcal{E}_k + r_{\ell_s}} \int |p(y)|^2 |y|^{2Q_s} e^{-(\vartheta-h)|y|} dy. \end{aligned} \quad (\text{OOQ31})$$

OSP20–26 supplies the first root-polynomial comparison on the entire coefficient space, retaining every coefficient of P_{k-8} . OFV5–7 supplies the exact original-order density multiplier, including every factor t_j and β_s . OFV8–9 supplies both pointwise bounds on the one-factor Gamma density, including near zero. Composing these three proved inequalities gives exactly OOQ31. The degree bound required there is satisfied because $L \leq q + d \leq 2q$. Every minimum here is taken on the same original fibre $\text{rem}_{P_{\partial,k}} p = v$; therefore OOQ31 passes to the quotient Grams by taking its attained minima. No multiplication map is presumed to preserve the kernel.

Define the two common scalar centres

$$\begin{aligned} c_{\text{lo}}(k) &= g(q, \vartheta), \\ c_{\text{hi}}(k) &= 2q \log(q/2) - qW(2, \vartheta). \end{aligned} \quad (\text{OOQ32})$$

For an entirely finite common error, take

$$\begin{aligned} \varepsilon_k &= \mathcal{E}_k + \max_{s=1,k} r_{\ell_s} + \max\{|\log L_h|, |\log U_h|\} \\ &+ \max_{\substack{s \in \{1,k\} \\ b \in \{\vartheta-h, \vartheta+h\}}} \left\{ \begin{aligned} &E_{\text{low}}(Q_s, b) + E_{\text{high}} + |g(Q_s, b) - c_{\text{lo}}(k)| \\ &+ |c_{\text{high}}(Q_s, b) - c_{\text{hi}}(k)| \end{aligned} \right\}. \end{aligned} \quad (\text{OOQ33})$$

All constants in this definition were specified above from the original parameters and the exact equilibrium. The resulting operator conclusion is

$$\begin{aligned} \beta_s e^{c_{\text{lo}} - \varepsilon_k} I_d &\preceq Q_{\partial,L}^{(s)} \preceq \beta_s e^{c_{\text{lo}} + \varepsilon_k} I_d, \quad L = d - 1, d, \\ \beta_s e^{c_{\text{hi}} - \varepsilon_k} I_d &\preceq Q_{\partial,L}^{(s)} \preceq \beta_s e^{c_{\text{hi}} + \varepsilon_k} I_d, \quad L = q + d - 1, q + d. \end{aligned} \quad (\text{OOQ34})$$

These are inequalities on every full original remainder coefficient vector.

Moreover

$$\varepsilon_k = O_{\delta,\gamma}(q^{3/4} + k \log q) = o(q). \quad (\text{OOQ35})$$

Indeed OOQ24 gives the first term $q^{3/4}$ because $n\rho^2 = n^{3/4}$; all remainder and diagonal-coordinate costs are $O(d \log n)$. OOQ28 gives the low error. The original OSP constant $\mathcal{E}_k$ is bounded at fixed δ, γ on its explicit domain, and $r_{\ell_s} = O(1/k)$ by OFV6–7. The sine bounds in OFV12 give logarithmic cost $O(\log q)$. Finally $Q_s - q = -d + \ell_s = O(k)$, the rates differ from ϑ by $1/q$, and W and its derivatives are bounded. The mean-value formula in the displayed scalar definitions therefore bounds both scalar discrepancies in OOQ33 by $O(k \log q + \log q)$. Every original mass remains in the common factor β_s .

40.7 Relative operators and all fixed subquotients of this actual target

The exact leading scalar difference is

$$c_{\text{lo}}(k) - c_{\text{hi}}(k) = q\{2 \log(4/\vartheta) - 2 + W(2, \vartheta)\} = \frac{q\mathfrak{C}_{\partial}}{2}. \quad (\text{OOQ36})$$

For the second equality use OOQ5 at $\alpha = 2$ and precisely OFV29, $\mathfrak{C}_{\partial} = 4 \log(4/\vartheta) - 10 - 2F(2, \vartheta) + 4F_{\alpha}(2, \vartheta)$. Equivalently it is the full OFV31 energy expression $4 \log(4/\pi) + 2 - 2I_{2,\pi}$. Thus this constant was obtained from the operator calculation with the same original source scale.

Let

$$\begin{aligned}\mathcal{R}_0 &= (Q_{\partial, q+d-1}^{(s)})^{-1/2} Q_{\partial, d-1}^{(s)} (Q_{\partial, q+d-1}^{(s)})^{-1/2}, \\ \mathcal{R}_1 &= (Q_{\partial, q+d}^{(s)})^{-1/2} Q_{\partial, d}^{(s)} (Q_{\partial, q+d}^{(s)})^{-1/2}.\end{aligned}\tag{OOQ37}$$

The positive square roots are the unique positive ones. For every eigenvalue of either of these actual relative operators, OOQ34 proves

$$\left| \log \lambda - \frac{q\mathfrak{C}_\partial}{2} \right| \leq 2\varepsilon_k = o(q).\tag{OOQ38}$$

This follows by positive congruence of the two scalar inequalities; it does not infer eigenvalues from their product. The positive operator $\mathcal{R}_0^{1/2} \mathcal{R}_1 \mathcal{R}_0^{1/2}$ has all eigenvalue logarithms within $4\varepsilon_k$ of $q\mathfrak{C}_\partial$, and its determinant is exactly the original four-endpoint determinant ratio. No commutation of the two relative operators is used or asserted.

For precision, the same result controls every specified fixed restriction or quotient of this outer target, even when its coefficient map depends on the original period and on k . For an injective map $I : \mathbb{C}^t \rightarrow \mathbb{C}^d$, use the actual restricted Grams $I^* Q_{\partial, L}^{(s)} I$. For a surjective map $A : \mathbb{C}^d \rightarrow \mathbb{C}^t$, use their attained minimum quotient Grams

$$Q_{A, L}^{(s)} = (A(Q_{\partial, L}^{(s)})^{-1} A^*)^{-1}.\tag{OOQ39}$$

The restriction coefficient reference is $I^* I$ and the quotient coefficient reference is $(AA^*)^{-1}$. Congruence, or minimization on the identical fibres, transports OOQ34 with exactly those reference forms. Taking determinants and adding the four original signs cancels the complete fixed reference determinant and the factor β_s^t . Hence in both cases the exact four-endpoint return $F_t^{(s)}$ satisfies

$$|F_t^{(s)} - tq\mathfrak{C}_\partial| \leq 4t\varepsilon_k.\tag{OOQ40}$$

For $t = O(k)$ this error is $o(kq)$, without any assumed conditioning of I or A : each is the same actual map at all four endpoints, so its complete coefficient form has been retained and cancels exactly.

The operator statements OOQ34 and OOQ38 concern the complete CTV outer-divisor quotient, whose ideal is $P_{\partial, k} \mathbb{C}[y]_{\leq L-d}$. The current original residual has the different, explicitly supplied CEP source graph, containing its complete $\mathcal{U}_{A, L}$ columns, the low- v components, and all original K lifts. OOQ39–40 do not identify that graph with this outer ideal. The present calculation supplies the full original outer operator and its exact subquotient transport; the comparison with the actual residual must use the complete CEP map and its denominator. No value of the separate original $\Phi_{k, s}$ is assigned by omitting that map.

41 The complete residual map on the original conductor graph

Retain every original coefficient and metric of CEP1–49 and FMC1–38. In particular $m = 1$, $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $q' = (k - 7)^2$, $\Delta = q - q'$, $j = 8k - 32$, $r = 8k - 16$, and the original first nonzero conductor moment is $0 \leq v \leq 80$. For each source $s \in \{1, k\}$ and endpoint $N \in \{q - 1, q, 2q - 1, 2q\}$ put $t = N - q + 1$ and $d = N - v$. All coefficient maps below retain the actual period matrix and the entire conductor $A = \sum_{a,b=0}^8 a_{ab} e_{ab}^{(8)}$.

41.1 The full graph denominator and its metric

Use original orthonormal coordinates on both source polynomial spaces, and define the coefficient space and injective multiplication map

$$X_N = \mathbb{C}^v \oplus \mathbb{C}[S']_{\leq N-v-q'}, \quad D_N = \text{diag}(I_v, M_{\chi_{k'}}), \quad \dim X_N = t + \Delta. \quad (\text{GIP1})$$

The second summand of X_N uses the increasing ordinary monomials in S' ; $M_{\chi_{k'}}$ includes their exact conversion to the original lower orthonormal source coordinates. The first summand is the original upper source's first v orthonormal coordinates. Thus D_N identifies X_N with $\mathbb{C}^v \oplus I'_N$, where $I'_N = \chi_{k'} \mathbb{C}[S']_{\leq N-v-q'}$. Let S_N be the complete invertible source map CEP28, and let U_N be the full upper source remainder in original primary coordinates CEP21. The specified map to the original conductor kernel and its residual observation are

$$\begin{aligned} E_N^{\text{graph}} &= U_N S_N^{-1} D_N : X_N \longrightarrow V_A, \\ B_N^{\text{graph}} &= E_j^\top T_k^\top S_k^\top E_N^{\text{graph}} : X_N \longrightarrow \mathbb{C}^j. \end{aligned} \quad (\text{GIP2})$$

Here T_k, S_k, E_j are the complete FMC and OCF maps, not new sections. CEP31 proves that E_N^{graph} is onto V_A , since $S_N^{-1} D_N X_N = U_N^{-1}(V_A)$. FMC30 proves that the second map is exactly $\Xi_k^{\text{fix}} E_N^{\text{graph}}$ and is onto.

For the t ordinary monomials h of degree at most $N - q$ let $Z_{\chi, N} h$ be the original upper orthonormal coefficients of $\chi_k h$. Define, with the complete products and shifts of CEP41–42,

$$\begin{aligned} \mathcal{Z}_{A, N} h &= \begin{pmatrix} P_{<v} Z_{\chi, N} h \\ \mathcal{U}_{A, N-q} h \end{pmatrix}, \\ (\mathcal{U}_{A, N-q} h)(S') &= \sum_{a,b=0}^8 a_{ab} d_{ab}(S') h(S' + b_{ab}), \quad D_N \mathcal{Z}_{A, N} = S_N Z_{\chi, N}. \end{aligned} \quad (\text{GIP3})$$

At $t = 0$ this is the empty column matrix. CEP42 proves the equality including the nonzero leading coefficient $\text{lc}(h) \binom{q+\deg h}{v} \mu_v$ for nonzero h . Consequently $\mathcal{Z}_{A, N}$ has rank t , and $\ker E_N^{\text{graph}} = \text{im } \mathcal{Z}_{A, N}$: a vector maps to zero precisely when its $S_N^{-1} D_N$ image is a full source relation $\chi_k h$.

The entire common kernel is also retained. Put $I_k = J_k E_r$ as in FMC30, and use its exact original minimum source lift $Y_{K, N} = U_N^* G_N I_k$. Its columns belong to $U_N^{-1}(V_A)$, and hence to the domain of the inverse of D_N on its indicated image after application of S_N . Define

$$W_{K, N} = D_N^{-1} S_N Y_{K, N}, \quad \ker B_N^{\text{graph}} = \text{im}[\mathcal{Z}_{A, N}, W_{K, N}], \quad \text{rank}[\mathcal{Z}_{A, N}, W_{K, N}] = t + r. \quad (\text{GIP4})$$

Indeed $E_N^{\text{graph}} W_{K, N} = I_k$ because $U_N U_N^* = G_N^{-1}$. A relation among the two displayed sets, after applying E_N^{graph} , is therefore a relation among the independent columns of I_k , so its $W_{K, N}$ coefficients vanish. The graph columns are independent as proved above. Conversely a vector killed by B_N^{graph} has image in K and subtracting the corresponding $W_{K, N}$ combination leaves a graph relation. This proves both equality and rank, including every original low-degree graph component and every common-kernel column.

There are two precisely specified positive metrics on X_N :

$$H_N^{\text{actual}} = D_N^* S_N^{-*} S_N^{-1} D_N, \quad H_N^{\text{graph}} = D_N^* D_N = \text{diag}(I_v, M_{\chi_{k'}}^* M_{\chi_{k'}}). \quad (\text{GIP5})$$

The first is the exact pullback of the unchanged original source norm, and the second is the pullback of the original direct-sum target norm of CEP28. Positivity follows from injectivity of D_N and invertibility of S_N . In particular the second block, after the literal substitution $S' = c' + iy$, integrates $|f(c' + iy)|^2 |P_{k'}(y)|^2 dm_s(y)$: the factor $i^{q'}$ in $\chi_{k'}(c' + iy) = i^{q'} P_{k'}(y)$ has its complete modulus one. The coordinate substitution in f is retained in this statement.

Both residual quotient metrics consequently have full finite formulas

$$\begin{aligned} Q_{\Xi,N}^{\text{fix}} &= [B_N^{\text{graph}}(H_N^{\text{actual}})^{-1}(B_N^{\text{graph}})^*]^{-1}, \\ \tilde{Q}_{\Xi,N} &= [B_N^{\text{graph}}(H_N^{\text{graph}})^{-1}(B_N^{\text{graph}})^*]^{-1}, \quad |\log \det \tilde{Q}_{\Xi,N} - \log \det Q_{\Xi,N}^{\text{fix}}| \leq 2E_N^{\text{CEP}}. \end{aligned} \quad (\text{GIP6})$$

Here E_N^{CEP} is the entire constant CEP35. For either positive H and onto B , the vector $H^{-1}B^*(BH^{-1}B^*)^{-1}y$ maps to y and is H -orthogonal to $\ker B$. Adding any other lift adds a perpendicular kernel vector and proves the inverse-Gram formula. The first minimum is the original minimum over all upper source lifts of a primary vector with residual y , by GIP2 and GIP5; this proves its equality with FMC32. For the comparison, the actual source map S_N induces the identity of these two quotient coordinate targets. Minimal lifts followed by orthogonal projection off the entire kernel GIP4 have exterior norm at most $\|\bigwedge^j S_N\|$; the inverse has bound $\|\bigwedge^j S_N^{-1}\|$. Orthogonal projection is an exterior contraction because its singular values are zero or one. Squaring the two volume inequalities and using CEP34–35 proves the final bound. Thus all four determinant errors sum to at most $2\sum_N E_N^{\text{CEP}} = o(kq)$ for the fixed actual period, without identifying any two different relation spaces.

41.2 The exact natural map to the CTV outer divisor

Let $\mathcal{I}_k = \{4, \dots, k-4\}^2$ and $\mathcal{O}_k = \{0, \dots, k\}^2 \setminus \mathcal{I}_k$. Let $E_{\text{in}}, E_{\text{out}}$ insert these coordinates into the original primary order. In the centred y coordinate retain the full outer Vandermonde $(V_{\text{out}})_{(a,b),h} = \omega_{ab}^h$ for $0 \leq h < \Delta$ and the diagonal matrix $D_{\text{in,out}} = \text{diag}(P_{k'}(\omega_{ab}))_{\mathcal{O}_k}$. Both are invertible: all the original roots are distinct and no outer root is an inner root. Define

$$\begin{aligned} \rho_A &= V_{\text{out}}^{-1} D_{\text{in,out}}^{-1} E_{\text{out}}^T|_{V_A} : V_A \longrightarrow \mathbb{C}[y]/(P_{\partial,k}), \\ C_{\text{in}} &= CE_{\text{in}}, \quad C_{\text{out}} = CE_{\text{out}}, \quad (C_{\text{in}})_{(a,b),(i,j)} = a_{i+4-a,j+4-b} \quad (0 \leq a, b, i, j \leq k'), \end{aligned} \quad (\text{GIP7})$$

where an out-of-range coefficient of A is zero. This is the literal map which first takes outer values and then divides by the inner root factor from CTV10; on the graph its matrix is $\rho_A E_N^{\text{graph}}$. A source relation $\chi_k h$ is killed because its outer evaluations vanish. Thus GIP7 is independent of every source lift and retains exactly the CTV remainder frame.

There is an exact sequence with an entirely explicit obstruction:

$$\begin{aligned} 0 \longrightarrow \ker C_{\text{in}} \xrightarrow{E_{\text{in}}} V_A \xrightarrow{\rho_A} \mathbb{C}^\Delta \xrightarrow{\partial_A} \text{coker } C_{\text{in}} \longrightarrow 0, \\ \partial_A f = [C_{\text{out}} D_{\text{in,out}} V_{\text{out}} f], \quad \dim \ker \rho_A = \dim \text{coker } \rho_A = \dim \ker C_{\text{in}}. \end{aligned} \quad (\text{GIP8})$$

To prove it, write $x = E_{\text{in}}z + E_{\text{out}}w$. Membership in V_A says exactly $C_{\text{in}}z + C_{\text{out}}w = 0$, while $\rho_A x = f$ says $w = D_{\text{in,out}} V_{\text{out}} f$. These two equations prove exactness at the first three terms. Surjectivity of C proves $\text{im } C_{\text{in}} + \text{im } C_{\text{out}} = \mathbb{C}^{q'}$, and the outer diagonal and Vandermonde are invertible, so ∂_A is onto. Since C_{in} is square its kernel and cokernel have equal dimension, proving the dimension assertion. Every possible defect is therefore the nullity of this specific actual finite convolution section. The scalar first-moment index v does not by itself evaluate that matrix or its nullity.

The original residual observation is retained even when this defect is nonzero. Put $W_A = \Xi_k^{\text{fix}} E_{\text{in}} \ker C_{\text{in}}$. Then the exact descended residual map and its kernel are

$$\bar{\Xi}_A : \ker \partial_A \longrightarrow \mathbb{C}^j / W_A, \quad \rho_A x \longmapsto [\Xi_k^{\text{fix}} x], \quad \ker \bar{\Xi}_A = \rho_A(K). \quad (\text{GIP9})$$

Two lifts differ by $E_{\text{in}} \ker C_{\text{in}}$ by GIP8, so the displayed rule is well defined. It is onto because Ξ_k^{fix} is onto. If its value is zero, subtract an element of $E_{\text{in}} \ker C_{\text{in}}$ with the same residual; the remaining lift lies in K and has unchanged outer image. This proves the kernel equality; its converse follows by direct application of Ξ_k^{fix} . Equations GIP3–9 thus specify the full denominator, retained metrics, and exact obstruction to using the natural outer map for the same residual classes.

42 The original mixed return and its same-class action control

This calculation retains the original arithmetic source, its actual cyclic inclusion, and the complete period observation. It computes the action control on those objects and the exact degree-step information carried by the signed mixed return. The original full-multiplicity control calculation is stated with its full multiplicity; the subsequent evaluated outer-return substitution uses precisely the proved $m = 1$ domain.

42.1 The original arithmetic class and all three source metrics

Fix the stipulated quartet $\rho = 1/2 + \delta + i\gamma$ and its reflected and conjugate centres, with $0 < \delta < 1/2$, $\gamma > 2$, and full common multiplicity $m \geq 1$. Retain

$$\begin{aligned} h(s) &= \prod_{\lambda \in \{\rho, \bar{\rho}, 1-\rho, 1-\bar{\rho}\}} (s - \lambda)^m, \quad g = 2\xi, \quad v_h = g/h, \quad w_h(y) = |v_h(1/2 + iy)|^2/(2\pi), \\ k &= 4l + 1 \geq 9, \quad c = k/2, \quad e = 1 + k(m - 1), \quad q = e(k + 1)^2, \\ \beta_{ab} &= c + (2a - k)\delta + i(2b - k)\gamma, \quad \chi(S) = \prod_{a,b=0}^k (S - \beta_{ab})^e, \quad E = \mathbb{C}[S]/(\chi), \quad A = M_S. \end{aligned} \tag{MCE1}$$

The three source measures, all in the literal coordinate $S = c + iy$, are the arithmetic measure $w_h^{*k}(y) dy$ and the original Gamma measures $dm_s(y)$ of CTV1 for $s = 1, k$. Their masses are respectively μ_h^k and $(2\pi)^{s/2}$. For a source $a \in \{\text{ar}, 1, k\}$ let $p_n^a(S)$ be its monic orthogonal polynomial, ω_n^a its full squared norm, and $b_n^a = [p_n^a]_\chi$ in the fixed original coefficient frame of E . For the Gamma sources these are explicitly

$$\begin{aligned} p_n^s(S) &= i^n \mathfrak{p}_n^s((S - c)/i), \quad \mathfrak{p}_0^s = 1, \quad \mathfrak{p}_1^s = y, \\ \mathfrak{p}_{n+1}^s &= y \mathfrak{p}_n^s - n(s/2 + n - 1) \mathfrak{p}_{n-1}^s, \quad \omega_n^s = (2\pi)^{s/2} n! (s/2)_n. \end{aligned} \tag{MCE2}$$

Thus the i^n converting FMC29's real-coordinate polynomial to the monic polynomial in S is retained. In primary coordinates for $m = 1$, its entries are exactly $i^n \mathfrak{p}_n^s(\omega_{ab})$. For higher multiplicity the entries are the complete derivatives at each centre, with the factorials of the original Hermite coordinate map; none is replaced by a value-only coordinate.

For every $N \geq q - 1$ define the complete covariance and minimum Gram

$$R_N^a = \sum_{n=0}^N b_n^a (b_n^a)^* / \omega_n^a, \quad G_N^a = (R_N^a)^{-1}, \quad V_N^a = \det G_N^a. \tag{MCE3}$$

The first q columns are the monic triangular remainder frame and are independent. Positivity of all norms therefore proves $R_N^a > 0$. In the orthogonal source coefficient frame the minimum section is $\text{diag}(\omega_0^a, \dots, \omega_N^a)^{-1} [b_0^a, \dots, b_N^a]^* G_N^a$. Multiplication verifies that it is a section and is orthogonal to the full relation kernel. The squared norm of every other section value adds the squared norm of a perpendicular relation. This proves MCE3's minimum interpretation with all masses intact.

The original arithmetic inclusion is

$$\eta : E \hookrightarrow E_h^{\otimes k}, \quad \eta[P] = v_h^{\otimes k} P(A_\Sigma) 1, \quad v_h = j_h(v_h), \quad A_\Sigma = \sum_{i=1}^k M_{s_i}, \quad \eta A = A_\Sigma \eta. \tag{MCE4}$$

Its injectivity follows from the exact annihilator χ and the invertibility of the complete Taylor unit $v_h^{\otimes k}$. For clarity the local nilpotent sum has exact order e : its $(e - 1)$ -st power has coefficient $[k(m - 1)]! / [(m - 1)!]^k$ at the product of the top local powers, and every higher monomial has an exponent at least m . All pairs (a, b) occur among tensor choices and their centres are distinct. The least common multiple of these local annihilators is exactly MCE1. This proves the asserted annihilator

and injectivity. The commutation identity follows by polynomial multiplication and commutation with the retained unit. The supplied source identity $J^{(k)}\mathcal{V}_{h,k} = \eta \text{rem}_\chi$ therefore carries each calculation below to this same arithmetic class. On an original present support label the map is $(\lambda, x) \mapsto (\lambda, \eta x)$, and external absence retains its original image; no coefficient calculation changes that label.

42.2 The exact rank-two control from the original recurrence

For this subsection fix any one source and suppress its superscript. Positivity of the original measure gives real monic orthogonal polynomials $Q_n(y) = i^{-n}p_n(c + iy)$. Pairing yQ_n with lower degrees proves

$$Sp_n = p_{n+1} + (c + i\alpha_n)p_n - \frac{\omega_n}{\omega_{n-1}}p_{n-1}, \quad \alpha_n \in \mathbb{R}, \quad (\text{MCE5})$$

with the lower term absent at $n = 0$. Indeed degrees below $n - 1$ are orthogonal by self-adjoint multiplication by the real variable; the coefficient at Q_{n+1} is one by monicity, and pairing with Q_{n-1} gives ω_n/ω_{n-1} . For the Gamma measures $\alpha_n = 0$ and MCE5 is precisely MCE2 with its phases.

Reducing MCE5 modulo χ and inserting it into MCE3 gives

$$\begin{aligned} AR_N + R_N A^* - kR_N &= \frac{b_{N+1}b_N^* + b_N b_{N+1}^*}{\omega_N}, \\ D_N := A^*G_N + G_N A - kG_N &= \frac{G_N b_{N+1}b_N^* G_N + G_N b_N b_{N+1}^* G_N}{\omega_N}. \end{aligned} \quad (\text{MCE6})$$

For the first equality the diagonal coefficient is $c + i\alpha_n + c - i\alpha_n - k = 0$. At every internal adjacent pair the remaining coefficient is $\omega_n^{-1} - (\omega_{n+1}/\omega_n)\omega_{n+1}^{-1} = 0$. Only the two displayed top columns survive. Multiplication on both sides by G_N proves the second equality, including its signs.

The dagger-stable polynomial obeys $\bar{\chi}(k - S) = (-1)^q \chi(S)$, so $\hat{\chi}(y) = i^{-q} \chi(c + iy)$ is real monic. The exact substitution $[P(S)] \mapsto [P(c + iy)]$ transports G_N by its full inverse congruence; in that real remainder frame the covariance and Gram have real entries. Each column has phase i^n . Consequently

$$\begin{aligned} b_N^* G_N b_{N+1} &= i\omega_N \phi_N, \quad \phi_N \in \mathbb{R}, \\ \mathcal{H}_N &= G_N^{-1/2} D_N G_N^{-1/2} = \frac{yx^* + xy^*}{\omega_N}, \\ x &= G_N^{1/2} b_N, \quad y = G_N^{1/2} b_{N+1}. \end{aligned} \quad (\text{MCE7})$$

Thus $\mathcal{H}_N$ is Hermitian of rank at most two and trace zero. Its square has trace $2(\|x\|^2\|y\|^2 - |x^*y|^2)/\omega_N^2$, because $x^*y = i\omega_N \phi_N$. Its two possibly nonzero eigenvalues are therefore $\epsilon_N, -\epsilon_N$, where

$$\epsilon_N^2 = \frac{\|x\|^2\|y\|^2 - |x^*y|^2}{\omega_N^2} = \frac{\|x \wedge y\|^2}{\omega_N^2}. \quad (\text{MCE8})$$

The last equality is the determinant of the two-vector Gram, proved by expanding in an orthonormal basis. This defines the literal control allowance, including the case when the rank is zero.

42.3 The same nonzero arithmetic exterior class

Let E_+ be the full sum of the primary blocks with $a > k/2$, and put $t_+ = e(k+1)^2/2$. Its dimension is t_+ because k is odd. A completely specified primary basis is $e_\beta(S - \beta)^d$, $0 \leq d < e$, over those centres, where

$$\chi_\beta = \chi/(S - \beta)^e, \quad e_\beta = \chi_\beta \sum_{u=0}^{e-1} \frac{(1/\chi_\beta)^{(u)}(\beta)}{u!} (S - \beta)^u \pmod{\chi}. \quad (\text{MCE9})$$

The Taylor product is one modulo $(S - \beta)^e$ and vanishes at the other complete primary factors. This proves the primary idempotent property and the stated basis. Multiplication by S has diagonal β and the exact next-power nilpotent on each block. Thus E_+ is A -invariant, and its ordered top wedge ξ_+ is a nonzero eigenvector for the additive exterior action $A^{[t+]}$, with eigenvalue $\sum_{a>k/2, b} e\beta_{ab}$. The map $\bigwedge^{t+} \eta$ is injective: extend an independent set in its domain to a basis and use injectivity of η on the resulting independent images. Hence $\bigwedge^{t+} \eta(\xi_+) \neq 0$ is the same surviving arithmetic exterior class. MCE4 intertwines the additive exterior actions term by term, retaining all its unit and nilpotent coefficients.

On $\bigwedge^t(E, G_N)$ the exact relative control is the additive exterior action of $\mathcal{H}_N$. In an orthonormal basis this follows by applying A to each wedge factor and pairing the result, so its eigenvalues are sums of t distinct eigenvalues of $\mathcal{H}_N$. For $1 \leq t \leq q - 1$ they lie in $[-\epsilon_N, \epsilon_N]$, since the only nonzero entries in the original list are $\epsilon_N, -\epsilon_N$. Applying this bound to ξ_+ , or taking the trace of the control on its invariant primary space, proves

$$\mathcal{L}_{h,k} := 2\delta e(k+1) \left\lfloor \frac{(k+1)^2}{4} \right\rfloor = 2\Re \operatorname{Tr}(A|_{E_+}) - kt_+ \leq \epsilon_N, \quad \mathcal{L}_{h,k}/q \geq \delta k/2 > 0. \quad (\text{MCE10})$$

The sum of the positive integers $2a - k$ is the displayed floor, by summing the positive odd or even progression; for the present odd k it is $(k+1)^2/4$. Multiplication by the $k+1$ imaginary centres and full length e gives the equality. The last bound uses $\lfloor (k+1)^2/4 \rfloor \geq k(k+1)/4$. The exterior rank here is t_+ , and this exact inclusion and action determine its relation to the arithmetic class; it is not assigned the dimension of the unrelated period kernel.

42.4 Complete kernel, boundary and mixed pieces of the control

Fix the original observation $\Lambda : E \rightarrow B$ in any one fixed coordinate frame of its actual image, and its actual kernel inclusion $I : \mathbb{C}^{r_K} \hookrightarrow E$. For $m = 1$ on the proved period locus, $r_K = 8k - 16$; in the present algebraic formulas it remains the actual rank for the stipulated multiplicity. Define at each source and cutoff

$$\begin{aligned} H_N &= I^* G_N I, \quad Q_N = (\Lambda R_N \Lambda^*)^{-1}, \quad L_N = R_N \Lambda^* Q_N, \quad K_N^{\text{coord}} = H_N^{-1} I^* G_N, \\ \Lambda L_N &= I_B, \quad I^* G_N L_N = 0, \quad I K_N^{\text{coord}} + L_N \Lambda = I_E, \\ F_N &= [I, L_N], \quad F_N^* G_N F_N = \operatorname{diag}(H_N, Q_N). \end{aligned} \quad (\text{MCE11})$$

The first section identity follows by multiplication. The second follows from $\Lambda I = 0$. The two summands in the third are the G_N -orthogonal projections onto K and its complement: the first fixes $I\mathbb{C}^{r_K}$; subtracting the second from any vector lies in K and has its stated kernel coordinate. This proves the third identity and the full Gram equality.

The complete original action in this frame is

$$F_N^{-1} A F_N = \begin{pmatrix} K_N^{\text{coord}} A I & K_N^{\text{coord}} A L_N \\ \Lambda A I & \Lambda A L_N \end{pmatrix} =: \begin{pmatrix} A_{KK} & A_{KB} \\ A_{BK} & A_{BB} \end{pmatrix}. \quad (\text{MCE12})$$

In particular $A_{BK} = \Lambda A I$ is the actual action defect; no invariant-kernel premise is used. The exact boundary relation is $\Lambda A - A_{BB} \Lambda = A_{BK} K_N^{\text{coord}}$, obtained by inserting the third identity in MCE11. It is the metric-section version of OPG22; replacing its fixed section by L_N changes both displayed blocks by precisely this identity. The three complete control blocks are

$$\begin{aligned} I^* D_N I &= A_{KK}^* H_N + H_N A_{KK} - k H_N, \\ L_N^* D_N L_N &= A_{BB}^* Q_N + Q_N A_{BB} - k Q_N, \\ I^* D_N L_N &= A_{BK}^* Q_N + H_N A_{KB}. \end{aligned} \quad (\text{MCE13})$$

Multiplication of MCE12 by the block Gram MCE11 proves each entry. For an actual eigenvector $z = Iu + L_N b$ of eigenvalue λ , the two action equations are $A_{KK}u + A_{KB}b = \lambda u$ and $A_{BK}u + A_{BB}b = \lambda b$. Consequently its literal control pairing is

$$(2\Re \lambda - k)(u^* H_N u + b^* Q_N b) = u^* I^* D_N I u + 2\Re(u^* I^* D_N L_N b) + b^* L_N^* D_N L_N b. \quad (\text{MCE14})$$

This follows either from the two action equations and MCE13 or directly from $z^*D_N z$; it preserves the entire mixed pairing. For generalized primary vectors the full block matrix MCE12, rather than only this eigenvector equality, remains the action.

The same splitting gives a quantitative exterior identity for the actual allowance itself. Put

$$\begin{aligned} x_K &= H_N^{-1/2} I^* G_N b_N, & x_B &= Q_N^{1/2} \Lambda b_N, \\ y_K &= H_N^{-1/2} I^* G_N b_{N+1}, & y_B &= Q_N^{1/2} \Lambda b_{N+1}, \\ a_K &= \|x_K\|^2, & a_B &= \|x_B\|^2, & d_K &= \|y_K\|^2, & d_B &= \|y_B\|^2, \\ z_K &= x_K^* y_K, & z_B &= x_B^* y_B. \end{aligned} \tag{MCE15}$$

The map $G_N^{1/2} F_N \text{diag}(H_N, Q_N)^{-1/2}$ is unitary by MCE11. Its inverse sends MCE7's x, y to these respective ordered pairs. Therefore

$$\begin{aligned} z_K + z_B &= i\omega_N \phi_N, \\ \omega_N^2 \epsilon_N^2 &= \|x_K \wedge y_K\|^2 + \|x_B \wedge y_B\|^2 + \|x_K \otimes y_B - y_K \otimes x_B\|^2 \\ &= (a_K d_K - |z_K|^2) + (a_B d_B - |z_B|^2) + a_K d_B + a_B d_K - 2\Re(z_K \bar{z}_B). \end{aligned} \tag{MCE16}$$

To prove the second line expand $x \wedge y$ in the orthogonal direct sum $\bigwedge^2 K \oplus (K \otimes B) \oplus \bigwedge^2 B$. Its three components are exactly those displayed, including the minus sign in the mixed tensor. Orthogonality adds their squared norms. Expansion of each two-vector Gram and of the mixed tensor inner product proves the last line. Each squared norm in the second line is nonnegative. The individual z_K, z_B need not be purely imaginary; their actual sum is the pairing in MCE7.

For completeness the same surviving exterior class also has a full mixed coordinate map. If I_+ is the matrix of MCE9's actual positive-primary basis, put $X_+ = K_N^{\text{coord}} I_+$ and $Y_+ = \Lambda I_+$. Then

$$I_+ = IX_+ + L_N Y_+, \quad G_{+,N} = X_+^* H_N X_+ + Y_+^* Q_N Y_+ > 0, \quad \mathcal{L}_{h,k} = \text{Tr}(G_{+,N}^{-1} I_+^* D_N I_+). \tag{MCE17}$$

The first two equalities are MCE11. Invariance of E_+ says $AI_+ = I_+ A_+$ with the full primary matrix of MCE9. Thus $I_+^* D_N I_+ = A_+^* G_{+,N} + G_{+,N} A_+ - k G_{+,N}$; taking the trace after multiplication by $G_{+,N}^{-1}$ proves the last identity by MCE10. Its expansion using MCE13 retains $X_+^* I^* D_N L_N Y_+$ and its adjoint, as well as both diagonal blocks. The induced map on ξ_+ is the literal wedge expansion of the columns $IX_+ + L_N Y_+$, with the usual permutation signs when kernel factors are moved before boundary factors. This is an invertible mixed coordinate description of the same vector before applying the injective exterior arithmetic map MCE4.

42.5 Degree-step volumes determine the squared component sizes

Still fixing one original source, put for $N \geq q - 1$

$$\kappa_N = \log \frac{\det H_N}{\det H_{N+1}}, \quad \beta_N^B = \log \frac{\det Q_N}{\det Q_{N+1}}, \quad \tau_N = \log(V_N/V_{N+1}). \tag{MCE18}$$

These are nonnegative. Inserting the next exact covariance column and multiplying verifies $G_{N+1} = G_N - G_N b_{N+1} b_{N+1}^* G_N / (\omega_{N+1} + b_{N+1}^* G_N b_{N+1})$. The determinant of $I + uv^*$ is $1 + v^* u$, proved by choosing a basis with first vector u or by two-block elimination. Applying this identity first to the boundary covariance and then to the restricted kernel metric proves

$$\begin{aligned} e^{\beta_N^B} &= 1 + d_B / \omega_{N+1}, & e^{\kappa_N} &= 1 + d_K / (\omega_{N+1} + d_B), & \tau_N &= \kappa_N + \beta_N^B, \\ d_B / \omega_{N+1} &= e^{\beta_N^B} - 1, & d_K / \omega_{N+1} &= e^{\beta_N^B} (e^{\kappa_N} - 1). \end{aligned} \tag{MCE19}$$

All denominators are strictly positive. For the boundary step,

$$Q_{N+1}^{-1} = Q_N^{-1} + (\Lambda b_{N+1})(\Lambda b_{N+1})^* / \omega_{N+1}.$$

For the kernel step the determinant ratio from the displayed G_{N+1} is $1 - d_K/(\omega_{N+1} + d_K + d_B)$, whose reciprocal is the stated formula. Multiplying the two reciprocal ratios gives the complete covariance ratio and proves the third equality.

For $N \geq q$ the preceding total and boundary covariances are both positive definite. Subtracting their last column instead of adding the next gives

$$a_B/\omega_N = 1 - e^{-\beta_{N-1}^B}, \quad a_K/\omega_N = e^{-\beta_{N-1}^B}(1 - e^{-\kappa_{N-1}}). \quad (\text{MCE20})$$

Indeed the determinant ratio of the previous boundary covariance to the current one is $1 - a_B/\omega_N = e^{-\beta_{N-1}^B}$. The identical total formula gives $1 - (a_K + a_B)/\omega_N = e^{-\tau_N}$. Subtracting and using MCE19 proves MCE20. At the first degree $N = q - 1$ the total pairing is exactly $a_K + a_B = \omega_{q-1}$, because the first q monic remainder columns are an orthogonal quotient basis. Its two components are kept as MCE15; no inverse at $q - 2$ is taken or assumed. Thus MCE16, MCE19 and MCE20 express the actual mixed exterior allowance using the complete consecutive kernel and boundary returns and their actual cross-pairings.

42.6 The signed return gives an exact original control-radius budget

Restore source superscripts. Define the actual signed one-degree term

$$\psi_N = \tau_N^{\text{ar}} - (\beta_N^B)^{(k)}, \quad w_{q-1} = w_{2q-1} = 1, \quad w_N = 2 \quad (q \leq N \leq 2q - 2), \quad \sum_N w_N = 2q. \quad (\text{MCE21})$$

Telescoping MCE18–19 gives, with every original endpoint,

$$\Psi_k := F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)} = \sum_{N=q-1}^{2q-1} w_N \psi_N, \quad \tau_N^{\text{ar}} = \psi_N + (\beta_N^B)^{(k)}. \quad (\text{MCE22})$$

To see the weights, the first window from $q - 1$ to $2q - 1$ includes degrees $q - 1, \dots, 2q - 2$, while the second window from q to $2q$ includes $q, \dots, 2q - 1$. Their intersection is counted twice and their two extreme degrees once.

For each individual source, MCE7–8 and MCE19–20 give the literal radius formulas

$$\begin{aligned} (\epsilon_{q-1})^2 + (\phi_{q-1})^2 &= \frac{\omega_q}{\omega_{q-1}} (e^{\tau_{q-1}} - 1), \\ \epsilon_N^2 + \phi_N^2 &= \frac{\omega_{N+1}}{\omega_N} (1 - e^{-\tau_{N-1}})(e^{\tau_N} - 1), \quad N \geq q. \end{aligned} \quad (\text{MCE23})$$

The first uses exactly $\|x\|^2 = \omega_{q-1}$ and the next-column formula for $\|y\|^2$; the second uses both total formulas already proved. Subtracting $\phi_N^2 = |x^*y|^2/\omega_N^2$ recovers the actual ϵ_N^2 . In particular substitution of $\tau_N^{\text{ar}} = \psi_N + (\beta_N^B)^{(k)}$ in MCE23 is a pointwise, same-arithmetic-class map from the signed degree data to the original control. It retains the complete reference boundary increment and arithmetic phase rather than assigning them zero.

Put $\delta_j^{\text{ar}} = V_j^{\text{ar}}/V_{j-1}^{\text{ar}}$ for $j \geq q$. The positive lower bound MCE10 and MCE23 imply $0 < \delta_j^{\text{ar}} < 1$ at every index used next. Define the full arithmetic monic-norm window and contraction penalties

$$\begin{aligned} \mathcal{W}_k^{\text{ar}} &= \log \frac{\omega_{2q-1}^{\text{ar}} \omega_{2q}^{\text{ar}}}{\omega_{q-1}^{\text{ar}} \omega_q^{\text{ar}}}, \\ \mathcal{P}_- &= -\log(1 - \delta_{2q-1}^{\text{ar}}) - 2 \sum_{j=q}^{2q-2} \log(1 - \delta_j^{\text{ar}}), \\ \mathcal{P}_+ &= -\log(1 - \delta_q^{\text{ar}}) - \log(1 - \delta_{2q}^{\text{ar}}) - 2 \sum_{j=q+1}^{2q-1} \log(1 - \delta_j^{\text{ar}}). \end{aligned} \quad (\text{MCE24})$$

They are all finite, and the two penalties are nonnegative. Multiplying MCE23 across the first window gives

$$\prod_{N=q-1}^{2q-2} (\epsilon_N^2 + \phi_N^2) = \frac{\omega_{2q-1} V_{q-1}}{\omega_{q-1} V_{2q-1}} (1 - \delta_{2q-1}) \prod_{j=q}^{2q-2} (1 - \delta_j)^2.$$

Across the second it gives

$$\prod_{N=q}^{2q-1} (\epsilon_N^2 + \phi_N^2) = \frac{\omega_{2q} V_q}{\omega_q V_{2q}} (1 - \delta_q)(1 - \delta_{2q}) \prod_{j=q+1}^{2q-1} (1 - \delta_j)^2.$$

For these two displays all quantities are arithmetic. To verify them, write $e^{\tau_N} - 1 = (1 - \delta_{N+1})/\delta_{N+1}$. The monic norm ratios telescope, the reciprocal δ factors give the indicated quotient-volume ratio, every internal $(1 - \delta_j)$ occurs twice, and each stated extreme occurs once. Taking logarithms and using MCE22 proves the exact bridge

$$\boxed{\sum_{N=q-1}^{2q-1} w_N \log((\epsilon_N^{\text{ar}})^2 + (\phi_N^{\text{ar}})^2) = \Psi_k + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+}. \quad (\text{MCE25})$$

It is an equality on the unchanged arithmetic packet. Every allowance on its left is precisely MCE6–10's control allowance.

MCE10 now supplies a concrete necessary inequality for those same data, retaining its full phase contribution:

$$\begin{aligned} \Psi_k + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+ &\geq 4q \log \mathcal{L}_{h,k} + \sum_N w_N \log \left(1 + \frac{(\phi_N^{\text{ar}})^2}{\mathcal{L}_{h,k}^2} \right), \\ \min_{q-1 \leq N \leq 2q-1} \epsilon_N^{\text{ar}} &\leq \exp \left(\frac{\Psi_k + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+}{4q} \right). \end{aligned} \quad (\text{MCE26})$$

For the first line apply $\epsilon_N^{\text{ar}} \geq \mathcal{L}_{h,k}$ to each positive radius, write $\log(\mathcal{L}_{h,k}^2 + \phi^2) = 2 \log \mathcal{L}_{h,k} + \log(1 + \phi^2/\mathcal{L}_{h,k}^2)$, and use $\sum w_N = 2q$. For the second, the minimum of the allowances is at most the weighted geometric mean of their larger radii; take half of the logarithmic mean in MCE25. This proves both inequalities without assuming any desired upper control estimate.

Finally specialize only the evaluated return substitution to the proved $m = 1$ domain of OAR/OVR/OFV. Retain its exact value $\Psi_k = \mathcal{F}_{\partial,k}^{(k)} - \Phi_{k,k} + \mathcal{H}_k + \delta_k^\sigma + \mathbf{e}_{k,k}$, where $-e_{k,k}^- \leq \mathbf{e}_{k,k} \leq e_{k,k}^+$ with the entire OAR2 constants. Then MCE25 becomes

$$\begin{aligned} \sum_N w_N \log((\epsilon_N^{\text{ar}})^2 + (\phi_N^{\text{ar}})^2) &= \mathcal{F}_{\partial,k}^{(k)} - \Phi_{k,k} + \mathcal{H}_k + \delta_k^\sigma + \mathbf{e}_{k,k} \\ &\quad + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+. \end{aligned} \quad (\text{MCE27})$$

Substituting the already evaluated OFV outer value $\mathcal{F}_{\partial,k}^{(k)} = \Delta q \mathcal{C}_\partial + \mathbf{r}_k$ retains exactly the same equality, with its proved remainder $\mathbf{r}_k = O(q \log q)$ and the actual residual $\Phi_{k,k}$. In particular its original arithmetic necessary inequality is

$$\begin{aligned} \Phi_{k,k} &\leq \mathcal{F}_{\partial,k}^{(k)} + \mathcal{H}_k + \delta_k^\sigma + e_{k,k}^+ + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+ \\ &\quad - 4q \log \mathcal{L}_{h,k} - \sum_N w_N \log \left(1 + \frac{(\phi_N^{\text{ar}})^2}{\mathcal{L}_{h,k}^2} \right). \end{aligned} \quad (\text{MCE28})$$

This is MCE26 composed with the exact receiver and its upper error endpoint; no new positivity premise is introduced. It shows exactly which additional original boundary, norm, contraction and phase quantities accompany the mixed-return estimate in the control budget. The full pointwise control can instead be calculated by MCE15–16 and MCE19–23. In either form, the actual arithmetic action, its nonzero exterior class, all kernel-boundary defects and the original four endpoints remain explicit. No sign of Ψ_k alone has been substituted for a bound on ϵ_N^{ar} .

43 The next marked pairing as an original action and source derivative

Retain MCE1–28 with the full original observation and all source orders. Give the same source its existing real tilt $dm_{a,\theta}(y) = e^{\theta y} dm_a(y)$ for real $|\theta| < \pi/2$. In the arithmetic case $dm_a = w_h^{*k}(y)dy$. All polynomial moments and their derivatives exist on compact subintervals by the original exponential-moment bound. At $\theta = 0$ this is exactly the current source. The polynomial algebra, $A = M_S$, the arithmetic inclusion, and Λ are independent of θ .

In fixed monomial source coordinates let $\mathcal{R}_N(\theta)$ be the original minimum section of the quotient and let its source Gram be $\mathbf{H}_N(\theta)$. The identities $J_N \mathcal{R}_N = I$ and orthogonality to $\ker J_N$ imply

$$\partial_\theta G_N = \mathcal{R}_N^*(\partial_\theta \mathbf{H}_N) \mathcal{R}_N = \langle \mathcal{R}_N, y \mathcal{R}_N \rangle_{a,\theta}. \quad (\text{MCP1})$$

Indeed differentiating $J_N \mathcal{R}_N = I$ gives $J_N \mathcal{R}'_N = 0$. The two section-derivative terms in the derivative of $\mathcal{R}_N^* \mathbf{H}_N \mathcal{R}_N$ therefore vanish by that orthogonality. Differentiating the integral defining $\mathbf{H}_N$ multiplies the integrand by the unchanged variable y , proving the second equality with full source mass.

Here is its exact endpoint evaluation. In the fixed real-coordinate quotient set $T = M_y$, $Q_n = i^{-n} p_n(c + iy)$ and $d_n = [Q_n]$. The leading coefficient of the minimum lift of x is $d_N^* G_N x / \omega_N$. This follows by expanding that lift in the orthogonal Q_n : its Q_N coefficient is its inner product with Q_N divided by ω_N , and the section identity gives that inner product $d_N^* G_N x$. The polynomial column

$$y \mathcal{R}_N - \mathcal{R}_N T - (Q_{N+1} - \mathcal{R}_N d_{N+1}) d_N^* G_N / \omega_N$$

has degree at most N because its highest coefficient cancels, and its remainder is zero. Pairing it with $\mathcal{R}_N$ kills it by the same orthogonality. The term Q_{N+1} is orthogonal to every degree-at-most- N minimum lift. Thus $G'_N = G_N T - G_N d_{N+1} d_N^* G_N / \omega_N$ in this specified real-coordinate frame. Returning through the unchanged substitution $S = c + iy$, with $b_n = i^n d_n$ and $T = (A - cI)/i$, proves

$$G'_N = -i G_N (A - cI) + i G_N b_{N+1} b_N^* G_N / \omega_N. \quad (\text{MCP2})$$

Every p_n, b_n, ω_n in this formula is evaluated at the same θ . No derivative of these columns was inserted: the formula comes from the exact section minimum and its polynomial remainder.

Use MCE11's H_N, Q_N, L_N at that same parameter, and keep the original kernel inclusion I fixed. Then $H'_N = I^* G'_N I$ and $Q'_N = L_N^* G'_N L_N$. For the latter, differentiating $\Lambda L_N = I_B$ puts L'_N in K , while $I^* G_N L_N = 0$ kills both differentiated-section terms. Taking the two traces in MCP2 therefore proves the exact marked-pairing identities

$$\begin{aligned} \frac{z_K}{\omega_N} &= \text{Tr } A_{KK} - c r_K - i \partial_\theta \log \det H_N, \\ \frac{z_B}{\omega_N} &= \text{Tr } A_{BB} - c \dim B - i \partial_\theta \log \det Q_N. \end{aligned} \quad (\text{MCP3})$$

For example the first trace of $G_N(A - cI)$ compressed by I is $\text{Tr } A_{KK} - c r_K$. The trace of its marked rank-one term is precisely z_K of MCE15, by cyclicity of trace. The second calculation uses $L_N^* G_N = Q_N \Lambda$ and gives exactly z_B . Rearranging $\partial_\theta \log \det H_N = -i(\text{Tr } A_{KK} - c r_K) + i z_K / \omega_N$ proves the first line and fixes its sign; the second line follows in the same manner. Adding them recovers the full phase MCE7 and EP.9 because $\text{Tr } A_{KK} + \text{Tr } A_{BB} = \text{Tr } A$.

In particular the original kernel–boundary action is measured by the real parts of these pairings:

$$\begin{aligned} 2\Re(z_K / \omega_N) &= 2\Re \text{Tr } A_{KK} - k r_K, & 2\Re(z_B / \omega_N) &= 2\Re \text{Tr } A_{BB} - k \dim B, \\ \frac{\|x_K \otimes y_B - y_K \otimes x_B\|^2}{\omega_N^2} &= \frac{a_K d_B + a_B d_K}{\omega_N^2} - 2\Re \left[(\alpha_K - i \partial_\theta \log \det H_N) \overline{(\alpha_B - i \partial_\theta \log \det Q_N)} \right], \\ \alpha_K &= \text{Tr } A_{KK} - c r_K, & \alpha_B &= \text{Tr } A_{BB} - c \dim B. \end{aligned} \quad (\text{MCP4})$$

The derivatives are real because the two Gram determinants are strictly positive for real θ . This proves the first line. Substituting MCP3 into MCE16’s exact mixed tensor norm proves the second, with all phases and off-diagonal action effects retained. Its two size terms are the explicit degree-return quantities in MCE19–20. Thus the next original pointwise calculation is the marked pair z_K, z_B , equivalently these actual action traces and source derivatives. An outer quotient operator estimate must be transported through the full GIP graph and actual marked columns before it can bound these quantities. No derivative estimate is inferred by differentiating an uncontrolled remainder in a scalar asymptotic law.

44 The exact original reflection and the retained mixed phase

Retain the full original quartet, its multiplicity m , the complete unit $v_h = 2\xi/h$, every tensor factor and all source masses of MCP1–4. The argument concerns the original arithmetic source and both original Gamma orders $s = 1, k$. It also specifies the exact reflection map on the real tilted family of MCP1–4. No observation invariance is assumed.

44.1 The entire arithmetic unit and the source involution

The functional equation of the original completed function is $\xi(1-s) = \xi(s)$. Reflection permutes the four factors of the full quartet polynomial; each factor contributes its minus sign with its full multiplicity, giving

$$h(1-s) = (-1)^{4m}h(s) = h(s), \quad v_h(1-s) = v_h(s). \quad (\text{RFX1})$$

The second identity first holds off the finite zero set of h . Both sides are the entire quotient already supplied by the original arithmetic construction, so continuity at every removed centre proves the same identity there, including all Taylor coefficients. At $s = 1/2 + iy$ it implies

$$w_h(-y) = w_h(y), \quad w_h^{*k}(-y) = w_h^{*k}(y). \quad (\text{RFX2})$$

For the second equality, in the convolution integral substitute $y_i \mapsto -y_i$ in every integration variable and use the first equality in every factor. The integrals exist by the original finite-mass estimates; Tonelli's theorem applies to these nonnegative integrands. Each Gamma density in OOQ2 is also even, since $\overline{\Gamma(s/4 + iy/2)} = \Gamma(s/4 - iy/2)$ for the retained positive real source order. The full source constants remain equal on the two sides, rather than being divided out.

Let $c = k/2$ and let $\mathcal{P}_N$ consist of polynomials in the literal variable S . Define the complex-linear involution

$$(\mathcal{S}_N p)(S) = p(k-S), \quad (\mathcal{S}_N)_{ab} = (-1)^a \binom{b}{a} k^{b-a} \quad (0 \leq a \leq b \leq N), \quad (\text{RFX3})$$

and set every other matrix entry to zero. The displayed matrix is the binomial expansion in the original increasing monomial frame; substitution twice proves $\mathcal{S}_N^2 = I$. For each original source a and its real tilt $dm_{a,\theta} = e^{\theta y} dm_a$, direct substitution $y \mapsto -y$ proves

$$\|\mathcal{S}_N p\|_{a,-\theta}^2 = \|p\|_{a,\theta}^2, \quad |\theta| < \pi/2. \quad (\text{RFX4})$$

The full exponential-moment domain is the original one used in MCP; the polynomial factors and their derivatives remain integrable on its compact subintervals. Equation RFX4 therefore states an exact isometry between these two specified source spaces, including both full tilted source measures.

44.2 The quotient, primary jets and arithmetic class

The primary centres obey $k - \beta_{ab} = \beta_{k-a,k-b}$. The degree $q = e(k+1)^2$ is even for the original odd k . Consequently

$$\chi(k-S) = (-1)^q \chi(S) = \chi(S), \quad \mathcal{S}_N(\chi f) = \chi \mathcal{S}_{N-q} f. \quad (\text{RFX5})$$

The relation space is thus carried onto itself at every cutoff. There is an induced complex-linear involution Σ on $E = \mathbb{C}[S]/(\chi)$, with the literal identities

$$\text{rem}_\chi \mathcal{S}_N = \Sigma \text{rem}_\chi, \quad \Sigma^2 = I_E, \quad \Sigma A \Sigma = kI_E - A. \quad (\text{RFX6})$$

The first follows from RFX5, and the third follows by applying reflection to $Sp(S)$. In the degree-less-than- q frame, Σ is the matrix in RFX3 with $N = q - 1$. In original primary derivative coordinates its complete formula is

$$(\Sigma[P])^{(d)}(\beta_{ab}) = (-1)^d P^{(d)}(\beta_{k-a, k-b}), \quad 0 \leq d < e. \quad (\text{RFX7})$$

Dividing both sides by the same retained $d!$ gives the formula in divided-derivative coordinates; no factorial or top nilpotent layer is omitted. The matrix of this permutation with jet signs squares to the identity and is precisely the conjugate of RFX3 by the full original Hermite coordinate map.

The arithmetic inclusion also has an exact reflection map. On $E_h = \mathbb{C}[s]/(h)$ put $\sigma_h[f(s)] = [f(1-s)]$. RFX1 makes this a well-defined involution and fixes the full Taylor unit $v_h = j_h(v_h)$. Let $\sigma_h^{\otimes k}$ act on every tensor factor. Then

$$\sigma_h^{\otimes k} A_\Sigma \sigma_h^{\otimes k} = kI - A_\Sigma, \quad \sigma_h^{\otimes k} \eta = \eta \Sigma. \quad (\text{RFX8})$$

The first identity follows by summing $M_{1-s_i} = I - M_{s_i}$. For the second, apply that identity to $\eta[P] = v_h^{\otimes k} P(A_\Sigma)1$; the unit and 1 are fixed, giving exactly $v_h^{\otimes k} P(k - A_\Sigma)1$. Thus the same injective arithmetic map is used. On exterior powers the relation follows on every decomposable wedge and then by linearity. In particular the positive-primary exterior class is carried to the negative-primary class with all signs and nilpotent coordinates specified by RFX7, and its image remains nonzero by injectivity of $\bigwedge \eta$. Explicitly order the positive primary blocks lexicographically and their local powers by $d = 0, \dots, e-1$, and use the same convention on the negative side. With $B = (k+1)^2/2$ blocks, the resulting top-wedge sign is exactly

$$(-1)^{e^2 B(B-1)/2 + B e(e-1)/2}.$$

Reflection reverses the order of the B blocks, contributing e^2 transpositions per reversed pair, and contributes $\sum_{d=0}^{e-1} d = e(e-1)/2$ within-block sign exponents in each block by RFX7. This proves the displayed sign while retaining the same full unit in the arithmetic image. A present support label is retained under these maps; the original external absent element keeps its specified absent image. No amplitude cancellation changes a label.

44.3 The full metric and the total endpoint pairing

Give the quotient at cutoff $N \geq q - 1$ its original minimum norm, and denote its Gram by $G_N^a(\theta)$. Isometry RFX4 and relation identity RFX5 take the whole affine fibre of a class onto the fibre of its reflected class. Taking both attained minima therefore proves

$$\Sigma^* G_N^a(-\theta) \Sigma = G_N^a(\theta), \quad R_N^a(-\theta) = \Sigma R_N^a(\theta) \Sigma^*. \quad (\text{RFX9})$$

The inverse formula follows by taking inverses and using $\Sigma^{-1} = \Sigma$. The exact minimum sections satisfy $\mathcal{R}_N^a(-\theta) \Sigma = \mathcal{S}_N \mathcal{R}_N^a(\theta)$: the right side is a lift in the required reflected fibre and has the same minimum norm, so uniqueness identifies it with the left side.

At $\theta = 0$, the original real monic orthogonal polynomial $Q_n(y)$ has parity n . Indeed $(-1)^n Q_n(-y)$ is monic of degree n ; its pairing with any lower polynomial vanishes by evenness of the original measure and the substitution $y \mapsto -y$. Uniqueness of the monic orthogonal polynomial proves the claim. In the original S coordinate $p_n(S) = i^n Q_n((S-c)/i)$, so its remainder obeys

$$\Sigma b_n = (-1)^n b_n, \quad b_N^* G_N b_{N+1} = 0, \quad \phi_N = 0 \quad (N \geq q-1). \quad (\text{RFX10})$$

For the middle equality use RFX9 at zero to equate the pairing with $(\Sigma b_N)^* G_N (\Sigma b_{N+1})$. This is its negative by the first equality; over $\mathbb{C}$ the pairing must vanish. The last identity is the exact definition in MCE7, with $\omega_N > 0$. All three original sources and every original multiplicity are included. More generally their tilted monic polynomials obey $\mathcal{S}_n p_n^a(\theta) = (-1)^n p_n^a(-\theta)$ with

unchanged monic norms, by the same substitution and uniqueness. The vanishing claim RFX10 is made at the original source $\theta = 0$.

Since the determinant of the involution has squared modulus one, RFX9 also gives

$$\det G_N^a(-\theta) = \det G_N^a(\theta), \quad \partial_\theta \log \det G_N^a(\theta)|_{\theta=0} = 0. \quad (\text{RFX11})$$

Differentiability follows from the original exponential-moment bound and finite positive Gram inversion, as proved in MCP1. This is a proved derivative identity for the exact minimum metric; it does not differentiate an asymptotic error estimate.

44.4 The original mixed blocks remain present

Retain the original observation Λ , kernel inclusion I , and metric section L_N of MCE11. There is no assumption that Σ preserves $K = \ker \Lambda$. At the original source RFX10 implies, for the complete component pairings of MCE15,

$$z_B = -z_K, \quad \omega_N^2 \epsilon_N^2 = (a_K d_K - |z_K|^2) + (a_B d_B - |z_K|^2) + (a_K d_B + a_B d_K + 2|z_K|^2). \quad (\text{RFX12})$$

The first identity is $z_K + z_B = b_N^* G_N b_{N+1} = 0$. Inserting it into each of the three separate squared norms of MCE16 gives exactly the three displayed terms. Each term is the squared norm of its stated exterior component; the mixed one contains $+2|z_K|^2$. Thus RFX12 retains the full individual pairings even when they are nonzero. Adding the three exact terms yields $\omega_N^2 \epsilon_N^2 = (a_K + a_B)(d_K + d_B)$, consistently with MCE8 and RFX10.

Here is the exact relationship to the reflected observation, without requiring it to coincide with the original one. Define $\Lambda^\Sigma = \Lambda \Sigma$ and $I^\Sigma = \Sigma I$ with the same boundary and kernel coordinate spaces. Then $\ker \Lambda^\Sigma = \Sigma K$. At $\theta = 0$, RFX9 and MCE11 give

$$H_N^\Sigma = H_N, \quad Q_N^\Sigma = Q_N, \quad L_N^\Sigma = \Sigma L_N, \quad (K_N^{\text{coord}})^\Sigma = K_N^{\text{coord}} \Sigma. \quad (\text{RFX13})$$

For the section equality, use $R_N \Sigma^* = \Sigma R_N$, a consequence of RFX9, in the exact formula $L_N^\Sigma = R_N (\Lambda \Sigma)^* Q_N$. For the kernel coordinate equality use $\Sigma^* G_N = G_N \Sigma$. The two diagonal action blocks and both leakage blocks consequently satisfy

$$A_{KK}^\Sigma = kI_K - A_{KK}, \quad A_{BB}^\Sigma = kI_B - A_{BB}, \quad A_{KB}^\Sigma = -A_{KB}, \quad A_{BK}^\Sigma = -A_{BK}. \quad (\text{RFX14})$$

Insert RFX13 and $\Sigma A \Sigma = kI - A$ into the four exact definitions MCE12. The off-diagonal constant terms vanish because $K_N^{\text{coord}} L_N = 0$ and $\Lambda I = 0$. This proves all four relations while retaining the actual action defect $\Lambda A I$. The reflected component vectors are $(-1)^N(x_K, x_B)$ and $(-1)^{N+1}(y_K, y_B)$, by RFX10 and RFX13. Their squared sizes are unchanged and both marked pairings change sign, an exact description of their relation to the original mixed components.

In the original observation, put $g_K = \partial_\theta \log \det H_N|_0$ and $g_B = \partial_\theta \log \det Q_N|_0$. The complete metric splitting has endpoint frame $[I, L_N(\theta)]$. Its determinant is independent of θ , because $L_N(\theta) - L_N(0)$ lies in K and hence changes that frame by a triangular matrix with diagonal identity. Taking determinants in MCE11 and differentiating gives $g_K + g_B = 0$ by RFX11. Also the trace of A is cq , by summing all actual primary centres with multiplicity e . MCP3 therefore gives the additional exact formulas

$$\alpha_B = -\alpha_K, \quad g_B = -g_K, \quad \frac{z_K}{\omega_N} = \alpha_K - ig_K, \quad \frac{z_B}{\omega_N} = -\alpha_K + ig_K, \quad \alpha_K = \text{Tr } A_{KK} - cr_K. \quad (\text{RFX15})$$

They identify the original marked term to be controlled. Neither α_K nor g_K is assumed to vanish. In particular the mixed term in RFX12 is $a_K d_B + a_B d_K + 2\omega_N^2 |\alpha_K - ig_K|^2$.

44.5 Propagation through the same arithmetic control budget

Every allowance in the original arithmetic source is strictly positive by MCE10. Substitution of the proved RFX10 into MCE25 therefore gives the exact full-multiplicity equality

$$2 \sum_{N=q-1}^{2q-1} w_N \log \epsilon_N^{\text{ar}} = \Psi_k + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+. \quad (\text{RFX16})$$

Every endpoint weight, monic-norm window and contraction penalty is exactly MCE21–24. With $\sum_N w_N = 2q$, this proves

$$\begin{aligned} \mathcal{L}_{h,k} &\leq \min_N \epsilon_N^{\text{ar}} \leq \exp \left\{ \frac{\Psi_k + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+}{4q} \right\}, \\ 4q \log \mathcal{L}_{h,k} &\leq \Psi_k + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+. \end{aligned} \quad (\text{RFX17})$$

The exponential is now the exact weighted geometric mean of the allowances. A positive weighted geometric mean is at least the minimum, and each allowance is at least $\mathcal{L}_{h,k}$ by the same surviving arithmetic exterior class; these facts prove both inequalities. No new arithmetic upper bound is assumed.

On precisely the $m = 1$ evaluated-return domain of OAR/OVR/OFV, retain the actual signed error and substitute MCE27 to obtain

$$\begin{aligned} 2 \sum_N w_N \log \epsilon_N^{\text{ar}} &= \mathcal{F}_{\partial,k}^{(k)} - \Phi_{k,k} + \mathcal{H}_k + \delta_k^\sigma + \mathfrak{e}_{k,k} \\ &\quad + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+, \\ \Phi_{k,k} &\leq \mathcal{F}_{\partial,k}^{(k)} + \mathcal{H}_k + \delta_k^\sigma + e_{k,k}^+ \\ &\quad + F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+ - 4q \log \mathcal{L}_{h,k}. \end{aligned} \quad (\text{RFX18})$$

The second line follows from the first, RFX17, and the original upper error endpoint $\mathfrak{e}_{k,k} \leq e_{k,k}^+$. The complete evaluated value $\mathcal{F}_{\partial,k}^{(k)} = \Delta q \mathfrak{C}_\partial + O_{\delta,\gamma}(q \log q)$ may be inserted with its proved error and original domain. The arithmetic boundary, norm window, contraction penalties and residual remain explicit. Equations RFX12–15 retain the individual mixed blocks required for further control; the total phase has been evaluated through a proved source, quotient and arithmetic involution, with the full data preserved.

45 The outer operator, the full residual graph, and original action control

This receiving calculation uses the complete OQ1–40, GIP1–9 and MCE1–28 proofs, with their original source maps and constants. The preceding MRI and BRI equations remain valid. Here we give the actual substitutions into those equations and into the control of the same arithmetic class.

45.1 Original objects, domains and the same-target operator interval

For the original quartet retain $0 < \delta < 1/2$, $\gamma > 2$, $m \geq 1$, $k = 4l + 1 \geq 9$, $e = 1 + k(m - 1)$, $c = k/2$ and

$$q = e(k + 1)^2,$$

$$\chi(S) = \prod_{a,b=0}^k [S - c - (2a - k)\delta - i(2b - k)\gamma]^e, \quad (\text{OMR1})$$

$$E = \mathbb{C}[S]/(\chi), \quad A = M_S : E \longrightarrow E.$$

The original source measures are $w_h^{*k}(y) dy$ and the two full Gamma measures of orders $s = 1, k$, evaluated at the unchanged coordinate $S = c + iy$. Their masses remain μ_h^k and $(2\pi)^{s/2}$. All action identities below retain this full multiplicity. The outer estimates and the residual graph statements are applied only on their proved simple-quartet domain $m = 1$, with $q = (k + 1)^2$, $q' = (k - 7)^2$, $\Delta = q - q' = 16k - 48$, $r = 8k - 16$ and $j = 8k - 32$. They retain the actual period and the full explicit finite domains of OSP23, OQ13 and OAR1–3. No eventual bound is assigned to an input outside its stated domain.

Write $\varepsilon_k^{\text{out}}$ for exactly OQ33's finite ε_k , with all OQ4–33 constants, the full root polynomials and both source orders. It is distinct from the action radius ϵ_N^{ar} below and from E_N^{CEP} , the conductor exterior error. Put

$$\mathfrak{C}_\partial = 4 \log(4/\pi) + 2 - 2I_{2,\pi},$$

$$I_{2,\pi} = \iint \log|x - y| d\mu_{2,\pi}(x) d\mu_{2,\pi}(y), \quad (\text{OMR2})$$

$$D_k^{\text{out}} = 4\Delta\varepsilon_k^{\text{out}}, \quad \mathcal{O}_k^- = \Delta q \mathfrak{C}_\partial - D_k^{\text{out}}, \quad \mathcal{O}_k^+ = \Delta q \mathfrak{C}_\partial + D_k^{\text{out}}.$$

The measure $\mu_{2,\pi}$ is the complete EIQ equilibrium with its original mass one, endpoints and density, as retained in OFV and OVR. This is the same constant in MRI4r–u and BRI6h–i.

For clarity write $Q_{\text{lo},0}^{(s)} = Q_{\partial,\Delta-1}^{(s)}$, $Q_{\text{lo},1}^{(s)} = Q_{\partial,\Delta}^{(s)}$, $Q_{\text{hi},0}^{(s)} = Q_{\partial,q+\Delta-1}^{(s)}$ and $Q_{\text{hi},1}^{(s)} = Q_{\partial,q+\Delta}^{(s)}$. Every one acts on the original coefficient remainder frame $1, y, \dots, y^{\Delta-1}$. OQ34 gives, for $a = 0, 1$,

$$\beta_s e^{c_{\text{lo}} - \varepsilon_k^{\text{out}}} I_\Delta \preceq Q_{\text{lo},a}^{(s)} \preceq \beta_s e^{c_{\text{lo}} + \varepsilon_k^{\text{out}}} I_\Delta,$$

$$\beta_s e^{c_{\text{hi}} - \varepsilon_k^{\text{out}}} I_\Delta \preceq Q_{\text{hi},a}^{(s)} \preceq \beta_s e^{c_{\text{hi}} + \varepsilon_k^{\text{out}}} I_\Delta, \quad (\text{OMR3})$$

$$\beta_s = \frac{(2\pi)^{s/2}}{\sqrt{2}\Gamma(s/2)}, \quad c_{\text{lo}} - c_{\text{hi}} = q\mathfrak{C}_\partial/2.$$

All original masses are retained in β_s . Inverting the high inequalities and conjugating the low inequalities by their positive square roots proves that every eigenvalue of $\mathcal{R}_a = (Q_{\text{hi},a}^{(s)})^{-1/2} Q_{\text{lo},a}^{(s)} (Q_{\text{hi},a}^{(s)})^{-1/2}$ lies between $e^{q\mathfrak{C}_\partial/2 - 2\varepsilon_k^{\text{out}}}$ and $e^{q\mathfrak{C}_\partial/2 + 2\varepsilon_k^{\text{out}}}$. Conjugating the bounds on $\mathcal{R}_1$ by $\mathcal{R}_0^{1/2}$ then proves the corresponding bounds on $\mathcal{R}_0^{1/2} \mathcal{R}_1 \mathcal{R}_0^{1/2}$, without any commutation assumption. Its determinant is exactly the product of the two original low/high determinant ratios. Therefore

$$\mathcal{O}_k^- \leq \mathcal{F}_{\partial,k}^{(s)} \leq \mathcal{O}_k^+, \quad s = 1, k. \quad (\text{OMR4})$$

The error is $O_{\delta,\gamma}(\Delta q^{3/4} + \Delta k \log q) = o(kq)$ by OOQ35. The earlier OFV estimate with error $O(q \log q)$ also remains valid; OMR4 supplies a full operator origin for an additional interval, and does not assert that this scalar interval is tighter than that earlier bound at every finite input.

More generally let $I : \mathbb{C}^t \hookrightarrow \mathbb{C}^\Delta$ be a fixed injection, or let $C_0 : \mathbb{C}^\Delta \twoheadrightarrow \mathbb{C}^t$ be a fixed surjection. Here “fixed” means the same actual map at all four endpoints; dependence on k and the period is allowed. The restriction and quotient forms are exactly

$$\begin{aligned} Q_I &= I^* Q I, & Q_I^{\text{ref}} &= I^* I, \\ Q_{C_0} &= (C_0 Q^{-1} C_0^*)^{-1}, & Q_{C_0}^{\text{ref}} &= (C_0 C_0^*)^{-1}, \\ |\mathcal{F}_t^{(s)} - tq\mathfrak{C}_\partial| &\leq 4t\varepsilon_k^{\text{out}}. \end{aligned} \tag{OMR5}$$

Restriction of a positive-form inequality preserves it. For a quotient, inversion reverses the order, multiplication by the onto map preserves it, and the second inversion reverses it again. Thus the same scalar bounds hold relative to the displayed exact reference form. In the four determinants its logarithm has coefficient $1 + 1 - 1 - 1 = 0$; the complete frame cost cancels. The t eigenvalue bounds now prove the last inequality.

45.2 The full graph denominator and its exact outer morphism

We now give the actual map relating OMR5’s target to the original residual. This retains the full conductor A_{cond} of CEP, which is a different named map from $A = M_S$ in OMR1. For each original $N = q - 1, q, 2q - 1, 2q$ let $t = N - q + 1$, retain the first nonzero conductor moment index $0 \leq v \leq 80$, and define

$$\begin{aligned} X_N &= \mathbb{C}^v \oplus \mathbb{C}[S']_{\leq N-v-q'}, & D_N &= \text{diag}(I_v, M_{\chi_{k-8}}), & \dim X_N &= t + \Delta, \\ E_N^{\text{graph}} &= U_N S_N^{-1} D_N : X_N \rightarrow V_A, \\ B_N^{\text{graph}} &= \Xi_k^{\text{fix}} E_N^{\text{graph}} = E_j^T T_k^T S_k^T U_N S_N^{-1} D_N : X_N \rightarrow \mathbb{C}^j. \end{aligned} \tag{OMR6}$$

These are the full GIP1–2 maps: U_N is the original source remainder in primary coordinates and S_N is CEP28’s complete source map. The multiplication block of D_N retains its exact conversion from ordinary lower monomials to orthonormal source coordinates. Let $Z_{\chi,N}h$ be the original upper coefficients of $\chi_k h$. With every shift and root product of CEP41–42,

$$\mathcal{Z}_{A,N}h = \left(\sum_{a,b=0}^8 a_{ab} d_{ab}(S') h(S' + b_{ab}) \right), \quad D_N \mathcal{Z}_{A,N} = S_N Z_{\chi,N}, \tag{OMR7}$$

$$\begin{aligned} I_k &= J_k E_r, & Y_{K,N} &= U_N^* G_N I_k, & W_{K,N} &= D_N^{-1} S_N Y_{K,N}, \\ \ker E_N^{\text{graph}} &= \text{im } \mathcal{Z}_{A,N}, & \ker B_N^{\text{graph}} &= \text{im}[\mathcal{Z}_{A,N}, W_{K,N}], & \text{rank}[\mathcal{Z}_{A,N}, W_{K,N}] &= t + r. \end{aligned}$$

The graph columns are independent because for every nonzero h their complete leading coefficient is $\binom{q+\deg h}{v} \mu_v \text{lc}(h) \neq 0$; for the specified monomial columns $\text{lc}(h) = 1$. Applying E_N^{graph} to the last r columns gives I_k , since $U_N U_N^* = G_N^{-1}$. Thus a relation among both sets first has zero coefficients in $W_{K,N}$ and then in $\mathcal{Z}_{A,N}$. Conversely, subtracting the corresponding $W_{K,N}$ lift from any element of $\ker B_N^{\text{graph}}$ leaves a source relation. This proves the complete kernel formula, including the $t = 0$ case where the source-relation matrix has no columns.

The two actual positive metrics and their residual quotient are

$$\begin{aligned} H_N^{\text{actual}} &= D_N^* S_N^{-*} S_N^{-1} D_N, & H_N^{\text{graph}} &= D_N^* D_N, \\ Q_{\Xi,N}^{\text{fix}} &= [B_N^{\text{graph}} (H_N^{\text{actual}})^{-1} (B_N^{\text{graph}})^*]^{-1}, \\ \tilde{Q}_{\Xi,N} &= [B_N^{\text{graph}} (H_N^{\text{graph}})^{-1} (B_N^{\text{graph}})^*]^{-1}, \\ |\log \det \tilde{Q}_{\Xi,N} - \log \det Q_{\Xi,N}^{\text{fix}}| &\leq 2E_N^{\text{CEP}}. \end{aligned} \tag{OMR8}$$

For any onto B and positive H , the vector $H^{-1}B^*(BH^{-1}B^*)^{-1}y$ is the unique lift of y orthogonal to $\ker B$; every other lift adds an orthogonal kernel vector. Its squared norm proves the displayed quotient formula. The first metric is the exact pullback of the original upper source norm, so this is precisely MRI4v's Schur complement. The second metric retains the full low- v summand and the source $|P_{k-8}(y)|^2 dm_s(y)$ after the unchanged lower substitution. The full j th exterior bounds of S_N and S_N^{-1} , followed by orthogonal projection off OMR7's entire kernel, give the last inequality as in GIP6. Their four endpoint costs total at most $2 \sum_N E_N^{\text{CEP}} = o(kq)$ for the fixed actual period.

Let $E_{\text{in}}, E_{\text{out}}$ insert the original inner square $\{4, \dots, k-4\}^2$ and its complement in primary order. Retain the full outer Vandermonde V_{out} and the diagonal $D_{\text{in,out}} = \text{diag}(P_{k-8}(\omega_{ab}))_{\text{out}}$. The exact natural morphism is

$$\begin{aligned} \rho_A &= V_{\text{out}}^{-1} D_{\text{in,out}}^{-1} E_{\text{out}}^T|_{V_A}, \\ C_{\text{in}} &= C E_{\text{in}}, \quad C_{\text{out}} = C E_{\text{out}}, \quad (C_{\text{in}})_{(a,b),(i,j)} = a_{i+4-a,j+4-b}, \\ 0 &\longrightarrow \ker C_{\text{in}} \xrightarrow{E_{\text{in}}} V_A \xrightarrow{\rho_A} \mathbb{C}^\Delta \xrightarrow{\partial_A} \text{coker } C_{\text{in}} \longrightarrow 0, \\ \partial_A f &= [C_{\text{out}} D_{\text{in,out}} V_{\text{out}} f], \quad \dim \ker \rho_A = \dim \text{coker } \rho_A = \dim \ker C_{\text{in}}. \end{aligned} \tag{OMR9}$$

Indeed, writing a primary vector as $E_{\text{in}}z + E_{\text{out}}w$, its conductor equation is $C_{\text{in}}z + C_{\text{out}}w = 0$ and its outer image f specifies $w = D_{\text{in,out}} V_{\text{out}} f$. These two equations prove exactness; the surjectivity of C proves surjectivity onto the cokernel. The inner block is square, so its kernel and cokernel dimensions agree. On the original graph the map is exactly $\rho_A E_N^{\text{graph}}$.

The residual descends with its full possible defect as

$$W_A = \Xi_k^{\text{fix}} E_{\text{in}} \ker C_{\text{in}}, \quad \Xi_A : \ker \partial_A \rightarrow \mathbb{C}^j / W_A, \quad \rho_A x \mapsto [\Xi_k^{\text{fix}} x], \quad \ker \Xi_A = \rho_A(K). \tag{OMR10}$$

Two lifts differ by $E_{\text{in}} \ker C_{\text{in}}$, which proves well-definedness. Surjectivity follows from that of Ξ_k^{fix} . If a class has zero residual modulo W_A , subtract a vector in $E_{\text{in}} \ker C_{\text{in}}$ with the same residual. Its remaining lift lies in K and has the same outer image, proving the kernel identity. Equations OMR6–10 are the exact typed bridge used here. They retain the entire denominator and metrics. Consequently the fixed-subquotient bound OMR5 is applied only to a specified map of its own full outer target; this receiving calculation assigns no leading coefficient to $\Phi_{k,s}$ by replacing the actual graph denominator with an outer polynomial ideal.

45.3 Backward propagation through the original kernel and signed return

Keep the actual nonnegative residual sum $\Phi_{k,s}$ in MRI4l, its first increment λ_s , and its full earlier cap V_s . Write $[L_s^0, U_s^0]$ and $[L_{\text{ar}}^0, U_{\text{ar}}^0]$ for exactly MRI4q's existing intervals. On the common proved domain, the full operator estimate gives the additional same-scalar intervals

$$\begin{aligned} L_s^{\text{op}} &= \max\{L_s^0, \mathcal{O}_k^- - \Phi_{k,s} - e_{k,s}^-\}, \\ U_s^{\text{op}} &= \min\{U_s^0, \mathcal{O}_k^+ - \Phi_{k,s} + e_{k,s}^+\}, \\ L_s^{\text{op}} &\leq F_K^{(s)} \leq U_s^{\text{op}}, \quad s = 1, k. \end{aligned} \tag{OMR11}$$

To prove this, substitute OMR4 into the exact MRI4m equality $F_K^{(s)} = \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s} + \mathfrak{e}_{k,s}$, with $-e_{k,s}^- \leq \mathfrak{e}_{k,s} \leq e_{k,s}^+$. Intersecting with the previously proved interval gives OMR11. The intersection is nonempty because the same original scalar belongs to both intervals. All earlier branches remain available; outside the new domain one uses their original endpoints.

Define the unchanged correlated arithmetic allowances by

$$\begin{aligned} a_{K,k}^- &= \max\{-C_K^-, \delta_k^\sigma - C_B^+\}, \quad a_{K,k}^+ = \min\{C_K^+, \delta_k^\sigma + C_B^-\}, \\ a_{K,k}^- &\leq d_k^{\text{ar}} = F_K^{\text{ar}} - F_K^{(1)} \leq a_{K,k}^+, \\ L_{\text{ar}}^{\text{op}} &= \max\{L_{\text{ar}}^0, L_1^{\text{op}} + a_{K,k}^-\}, \\ U_{\text{ar}}^{\text{op}} &= \min\{U_{\text{ar}}^0, U_1^{\text{op}} + a_{K,k}^+\}, \\ F_K^{\text{ar}} &\in [L_{\text{ar}}^{\text{op}}, U_{\text{ar}}^{\text{op}}], \quad F_B^{\text{ar}} \in [\mathcal{B}_k^{\text{ar}} - U_{\text{ar}}^{\text{op}}, \mathcal{B}_k^{\text{ar}} - L_{\text{ar}}^{\text{op}}]. \end{aligned} \tag{OMR12}$$

The first bounds are exactly OAR8/MRI4o, with their original constants. Adding them to the source-1 inequalities proves the kernel interval. Subtraction from the same arithmetic total proves the boundary interval and its reversed endpoints. Their errors remain correlated through this equality.

The earlier signed scalar is exactly MCE22's Ψ_k :

$$\begin{aligned}\Psi_k &= \mathfrak{S}_k^{\text{mix}} = F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)} = F_K^{(k)} + \mathcal{H}_k + \delta_k^\sigma, \\ L_k^{\text{op}} + \mathcal{H}_k + \delta_k^\sigma &\leq \Psi_k \leq U_k^{\text{op}} + \mathcal{H}_k + \delta_k^\sigma.\end{aligned}\tag{OMR13}$$

The first equality uses the original total identity BRI6d, including $\mathcal{H}_k = -T_k$. The second line adds this exact signed correction to OMR11 at the original source order $s = k$. Every earlier finite bound for this same Ψ_k may also be intersected here. No arithmetic allowance or Gamma sign is changed.

45.4 The original action, its mixed components, and its positive class

For any of the actual sources let p_N be the monic polynomial in S , ω_N its full squared source norm, and $b_N = [p_N]_\chi$. In the original coefficient coordinates put

$$\begin{aligned}R_N &= \sum_{n=0}^N b_n b_n^* / \omega_n, \quad G_N = R_N^{-1}, \quad \mathcal{D}_N = A^* G_N + G_N A - k G_N, \\ x_N &= G_N^{1/2} b_N, \quad y_N = G_N^{1/2} b_{N+1}, \quad x_N^* y_N = i \omega_N \phi_N, \quad \phi_N \in \mathbb{R}, \\ G_N^{-1/2} \mathcal{D}_N G_N^{-1/2} &= (y_N x_N^* + x_N y_N^*) / \omega_N, \\ \text{spec}(G_N^{-1/2} \mathcal{D}_N G_N^{-1/2}) &= \{\epsilon_N, -\epsilon_N, 0, \dots, 0\}, \quad \epsilon_N^2 = \frac{\|x_N \wedge y_N\|^2}{\omega_N^2}.\end{aligned}\tag{OMR14}$$

The monic recurrence in MCE5 telescopes in $AR_N + R_N A^* - k R_N$ to $(b_{N+1} b_N^* + b_N b_{N+1}^*) / \omega_N$. Multiplication by G_N on both sides gives the control identity. Its trace is $2\Re \text{Tr } A - kq = 0$, which proves that $x_N^* y_N$ is purely imaginary. The square of this rank-at-most-two Hermitian operator has half trace $(\|x_N\|^2 \|y_N\|^2 - |x_N^* y_N|^2) / \omega_N^2$. This proves the two opposite eigenvalues and the exterior formula.

For the original observation $\Lambda : E \rightarrow B$ and inclusion $I : K \hookrightarrow E$, retain

$$\begin{aligned}H_N &= I^* G_N I, \quad Q_N = (\Lambda R_N \Lambda^*)^{-1}, \quad L_N = R_N \Lambda^* Q_N, \quad K_N^{\text{coord}} = H_N^{-1} I^* G_N, \\ \Lambda L_N &= I_B, \quad I^* G_N L_N = 0, \quad I K_N^{\text{coord}} + L_N \Lambda = I_E, \\ A_{KK} &= K_N^{\text{coord}} A I, \quad A_{KB} = K_N^{\text{coord}} A L_N, \quad A_{BK} = \Lambda A I, \quad A_{BB} = \Lambda A L_N, \\ \Lambda A - A_{BB} \Lambda &= A_{BK} K_N^{\text{coord}}, \\ I^* \mathcal{D}_N L_N &= A_{BK}^* Q_N + H_N A_{KB}.\end{aligned}\tag{OMR15}$$

The first three identities follow by multiplication and the minimum-lift orthogonality. Applying A in this direct-sum frame gives the action blocks and the displayed observation defect. Multiplication of $A^* G_N + G_N A - k G_N$ by I^* and L_N gives the final identity, since $I^* G_N L_N = 0$. This retains the actual action on the observation; it does not impose invariance of its kernel.

Define the marked components exactly by $x_K = H_N^{-1/2} I^* G_N b_N$, $x_B = Q_N^{1/2} \Lambda b_N$, and the same formulas with b_{N+1} for y_K, y_B . Put $a_K = \|x_K\|^2$, $a_B = \|x_B\|^2$, $d_K = \|y_K\|^2$, $d_B = \|y_B\|^2$, $z_K = x_K^* y_K$ and $z_B = x_B^* y_B$. Then

$$\begin{aligned}z_K + z_B &= i \omega_N \phi_N, \\ \omega_N^2 \epsilon_N^2 &= \|x_K \wedge y_K\|^2 + \|x_B \wedge y_B\|^2 + \|x_K \otimes y_B - y_K \otimes x_B\|^2 \\ &= (a_K d_K - |z_K|^2) + (a_B d_B - |z_B|^2) + a_K d_B + a_B d_K - 2\Re(z_K \bar{z}_B).\end{aligned}\tag{OMR16}$$

The frame $[I, L_N]$ has Gram $\text{diag}(H_N, Q_N)$, so the marked components give an isometry to $K \oplus B$. The exterior square decomposes orthogonally into $\bigwedge^2 K \oplus (K \otimes B) \oplus \bigwedge^2 B$. Expanding the middle norm gives the last three terms, including its complete complex cross-pairing. This proves every equality.

The full original positive primary part E_+ consists of all jets with $a > k/2$, has dimension $q/2$, and its top exterior class is nonzero under the original injection $\eta : E \hookrightarrow E_h^{\otimes k}$ because η is injective and $\eta A = A_{\text{sum}} \eta$. Its action excess is

$$\mathcal{L}_{h,k} = 2\Re \text{Tr}(A|_{E_+}) - kq/2 = 2\delta e(k+1) \left\lfloor \frac{(k+1)^2}{4} \right\rfloor \leq \epsilon_N, \quad N \geq q-1. \quad (\text{OMR17})$$

The nilpotent primary terms have zero trace. Summing the real parts over $a > k/2$, all b and all e jet directions gives the displayed coefficient. The induced additive Hermitian action on any exterior degree strictly between zero and q has largest eigenvalue ϵ_N , since OMR14's only nonzero eigenvalues are $\epsilon_N, -\epsilon_N$. Evaluating it on the positive class therefore proves the inequality. In the frame OMR15 its representative has both components $X_+ = K_N^{\text{coord}} I_+$ and $Y_+ = \Lambda I_+$, with Gram $X_+^* H_N X_+ + Y_+^* Q_N Y_+$; its expectation includes the cross block in OMR15. Thus this is the control of that same original class, with no component omitted.

45.5 The exact arithmetic radius budget and its backward bound

Define $\kappa_N = \log(\det H_N / \det H_{N+1})$, $\beta_N^B = \log(\det Q_N / \det Q_{N+1})$, and $\tau_N = \log(V_N / V_{N+1}) = \kappa_N + \beta_N^B$, where $V_N = \det G_N$. The original covariance rank-one update and its restriction give

$$\begin{aligned} e^{\beta_N^B} &= 1 + d_B / \omega_{N+1}, & e^{\kappa_N} &= 1 + d_K / (\omega_{N+1} + d_B), \\ \frac{a_B}{\omega_N} &= 1 - e^{-\beta_{N-1}^B}, & \frac{a_K}{\omega_N} &= e^{-\beta_{N-1}^B} (1 - e^{-\kappa_{N-1}}) \quad (N \geq q), \\ \epsilon_{q-1}^2 + \phi_{q-1}^2 &= \frac{\omega_q}{\omega_{q-1}} (e^{\tau_{q-1}} - 1), \\ \epsilon_N^2 + \phi_N^2 &= \frac{\omega_{N+1}}{\omega_N} (1 - e^{-\tau_{N-1}}) (e^{\tau_N} - 1) \quad (N \geq q). \end{aligned} \quad (\text{OMR18})$$

The first two formulas are the determinant lemma applied to the boundary covariance and then to the restricted Sherman–Morrison update for G_N . Applying the determinant lemma to removal of the last column proves the next two formulas. At $N = q-1$ the first q monic columns form the complete quotient basis, so $a_K + a_B = \omega_{q-1}$; no inverse at degree $q-2$ is used. Multiplying the total a and d formulas and using OMR14 proves the two radius identities.

Set $w_{q-1} = w_{2q-1} = 1$ and $w_N = 2$ for $q \leq N \leq 2q-2$, so that $\sum_N w_N = 2q$. For the arithmetic source alone, put

$$\begin{aligned} \delta_j^{\text{ar}} &= V_j^{\text{ar}} / V_{j-1}^{\text{ar}} \quad (j \geq q), \\ \mathcal{W}_k^{\text{ar}} &= \log \frac{\omega_{2q-1}^{\text{ar}} \omega_{2q}^{\text{ar}}}{\omega_{q-1}^{\text{ar}} \omega_q^{\text{ar}}}, \\ \mathcal{P}_- &= -\log(1 - \delta_{2q-1}^{\text{ar}}) - 2 \sum_{j=q}^{2q-2} \log(1 - \delta_j^{\text{ar}}), \\ \mathcal{P}_+ &= -\log(1 - \delta_q^{\text{ar}}) - \log(1 - \delta_{2q}^{\text{ar}}) - 2 \sum_{j=q+1}^{2q-1} \log(1 - \delta_j^{\text{ar}}), \\ C_k^{\text{budget}} &= F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+, \\ J_k^{\text{action}} &= \sum_{N=q-1}^{2q-1} w_N \log((\epsilon_N^{\text{ar}})^2 + (\phi_N^{\text{ar}})^2) = \Psi_k + C_k^{\text{budget}}. \end{aligned} \quad (\text{OMR19})$$

Here $0 < \delta_j^{\text{ar}} < 1$: positivity gives the first inequality and OMR17–18 forces the strict contraction at every index in the formula. All logarithms are therefore finite. To prove the budget identity, in each of the two windows $N = q-1, \dots, 2q-2$ and $N = q, \dots, 2q-1$ substitute $e^{\tau_N} - 1 = (1 - \delta_{N+1})/\delta_{N+1}$ into OMR18. The monic norm ratios telescope to $\mathcal{W}_k^{\text{ar}}$; the reciprocal δ factors give $F_K^{\text{ar}} + F_B^{\text{ar}}$. Each interior contraction occurs twice and the extreme ones once, giving exactly $-\mathcal{P}_- - \mathcal{P}_+$. Finally subtract and restore the same reference $F_B^{(k)}$ using OMR13. This proves the equality with every original endpoint.

The phases and the lower control allowance give the exact receiving inequalities

$$\begin{aligned} Z_k^{\text{phase}} &= \sum_{N=q-1}^{2q-1} w_N \log \left(1 + \frac{(\phi_N^{\text{ar}})^2}{\mathcal{L}_{h,k}^2} \right), \\ L_J &= L_k^{\text{op}} + \mathcal{H}_k + \delta_k^\sigma + C_k^{\text{budget}}, \quad U_J = U_k^{\text{op}} + \mathcal{H}_k + \delta_k^\sigma + C_k^{\text{budget}}, \\ 4q \log \mathcal{L}_{h,k} + Z_k^{\text{phase}} &\leq J_k^{\text{action}} \in [L_J, U_J], \\ \mathcal{L}_{h,k} &\leq \min_{q-1 \leq N \leq 2q-1} \epsilon_N^{\text{ar}} \leq \exp \left(\frac{J_k^{\text{action}}}{4q} \right) \leq \exp \left(\frac{U_J}{4q} \right). \end{aligned} \tag{OMR20}$$

Indeed $\epsilon_N^{\text{ar}} \geq \mathcal{L}_{h,k} > 0$ term by term. Thus each logarithm in OMR19 is at least $2 \log \mathcal{L}_{h,k} + \log(1 + (\phi_N^{\text{ar}})^2/\mathcal{L}_{h,k}^2)$. The weights sum to $2q$, giving the first bound. Also $\log((\epsilon_N^{\text{ar}})^2 + (\phi_N^{\text{ar}})^2) \geq 2 \log \min_M \epsilon_M^{\text{ar}}$; summing proves the middle upper bound. OMR13 and the exact budget equality prove $L_J \leq J_k^{\text{action}} \leq U_J$ and the final bound.

Finally the same exact signed substitution as BRI6f gives

$$\begin{aligned} J_k^{\text{action}} &= \mathcal{F}_{\partial,k}^{(k)} - \Phi_{k,k} + \mathcal{H}_k + \delta_k^\sigma + \mathfrak{e}_{k,k} + C_k^{\text{budget}}, \\ \Phi_{k,k} &\leq \mathcal{F}_{\partial,k}^{(k)} + \mathcal{H}_k + \delta_k^\sigma + e_{k,k}^+ + C_k^{\text{budget}} - 4q \log \mathcal{L}_{h,k} - Z_k^{\text{phase}}, \\ \Phi_{k,k} &\leq \mathcal{O}_k^+ + \mathcal{H}_k + \delta_k^\sigma + e_{k,k}^+ + C_k^{\text{budget}} - 4q \log \mathcal{L}_{h,k} - Z_k^{\text{phase}}. \end{aligned} \tag{OMR21}$$

The first equality preserves the complete original OAR signed error. Move $\Phi_{k,k}$ to the other side, use OMR20's lower bound for J_k^{action} , and then use $\mathfrak{e}_{k,k} \leq e_{k,k}^+$ to prove the second line. OMR4 proves the third. These caps may be intersected with the previous $\lambda_k \leq \Phi_{k,k} \leq V_k$ and all its earlier proved bounds, on their common domain.

Equations OMR19–21 retain the complete reference boundary, the arithmetic monic-norm window, both literal contraction penalties and every marked phase. The operator interval propagates to the original scalar and then to the control of the same surviving arithmetic class by these exact maps. They provide no sign-based exclusion from Ψ_k alone and assign no unevaluated leading value to $\Phi_{k,s}$.

45.6 The exact marked derivative and original reflection specialization

Use the original real source tilt $dm_{a,\theta}(y) = e^{\theta y} dm_a(y)$, with $|\theta| < \pi/2$, leaving E, A, Λ, I and the original arithmetic inclusion fixed. Every expression in this paragraph is evaluated at the same real θ . The full exponential-moment bound permits differentiation of every finite Gram integral on compact subintervals. If $\mathcal{R}_N$ is the minimum source section in fixed monomial source coordinates, differentiation of $J_N \mathcal{R}_N = I$ puts $\mathcal{R}'_N$ in $\ker J_N$. Its pairings with the minimum section vanish. Thus differentiating the minimum Gram inserts exactly y in its source integral. The leading coefficient of a minimum lift of x is $d_N^* G_N x / \omega_N$ in the real monic frame $Q_n^{\text{mon}} = i^{-n} p_n(c + iy)$, $d_n = [Q_n^{\text{mon}}]$. The polynomial $y \mathcal{R}_N - \mathcal{R}_N M_y - (Q_{N+1}^{\text{mon}} - \mathcal{R}_N d_{N+1}) d_N^* G_N / \omega_N$ has zero remainder and degree at most N . Its pairing with $\mathcal{R}_N$ vanishes, and Q_{N+1}^{mon} is orthogonal to that section.

Returning by the exact substitution $S = c + iy$ gives

$$\begin{aligned}\partial_\theta G_N &= -iG_N(A - cI) + iG_N b_{N+1} b_N^* G_N / \omega_N, \\ \frac{z_K}{\omega_N} &= \text{Tr } A_{KK} - c \dim K - i\partial_\theta \log \det H_N, \\ \frac{z_B}{\omega_N} &= \text{Tr } A_{BB} - c \dim B - i\partial_\theta \log \det Q_N.\end{aligned}\tag{OMR22}$$

The first equality is the full MCP1–2 calculation. For the second, multiply it by I^* and I , then take its H_N^{-1} trace. The first term is $-i(\text{Tr } A_{KK} - c \dim K)$ and the rank-one term has trace iz_K/ω_N by cyclicity. Rearrangement proves the stated sign. For the third use $Q'_N = L_N^* G'_N L_N$, whose section-derivative terms vanish by $I^* G_N L_N = 0$, and $L_N^* G_N = Q_N \Lambda$. This proves the boundary formula with its complete marked pairing. No derivative estimate follows from differentiating an asymptotic remainder; OMR22 concerns the exact finite source family.

Here the original, untitled source has a proved additional involution. The completed function and full quartet obey $\xi(1-s) = \xi(s)$ and $h(1-s) = (-1)^{4m} h(s) = h(s)$, so their entire quotient satisfies $v_h(1-s) = v_h(s)$, including every removed zero and its Taylor coefficients. Consequently w_h is even and substitution in every convolution factor makes w_h^{*k} even. Both original Gamma densities are also even, with their full masses. Define in the original coefficient frame

$$\begin{aligned}(\mathcal{S}_N p)(S) &= p(k-S), \quad (\mathcal{S}_N)_{ab} = (-1)^a \binom{b}{a} k^{b-a} \quad (0 \leq a \leq b \leq N), \\ \chi(k-S) &= \chi(S), \quad \text{rem}_\chi \mathcal{S}_N = \Sigma \text{rem}_\chi, \quad \Sigma^2 = I, \quad \Sigma A \Sigma = kI - A, \\ (\Sigma[P])^{(d)}(\beta_{ab}) &= (-1)^d P^{(d)}(\beta_{k-a, k-b}) \quad (0 \leq d < e), \\ \sigma_h^{\otimes k} \eta &= \eta \Sigma, \quad \sigma_h[f(s)] = [f(1-s)].\end{aligned}\tag{OMR23}$$

The first formula is the literal binomial expansion. Reflection permutes the centres by $(a, b) \mapsto (k-a, k-b)$ and q is even, so it preserves the full relation space. This proves the quotient map; differentiation proves every jet sign. The local involution σ_h fixes the full Taylor unit, and conjugates the tensor sum action to $kI - A_{\text{sum}}$. Applying it to $\eta[P] = v_h^{\otimes k} P(A_{\text{sum}})1$ proves the last identity. Thus the original arithmetic class is carried by the same injective map, with all tensor factors and jets retained.

Changing variables $y \mapsto -y$ carries the whole source fibre at tilt θ isometrically onto the reflected fibre at $-\theta$. The unique minimum sections and their Grams therefore obey

$$\begin{aligned}\Sigma^* G_N(-\theta) \Sigma &= G_N(\theta), \quad \mathcal{R}_N(-\theta) \Sigma = \mathcal{S}_N \mathcal{R}_N(\theta), \\ \Sigma b_n(0) &= (-1)^n b_n(0), \quad b_N(0)^* G_N(0) b_{N+1}(0) = 0, \quad \phi_N(0) = 0, \\ \partial_\theta \log \det G_N(\theta)|_0 &= 0.\end{aligned}\tag{OMR24}$$

Indeed for an even measure the polynomial $(-1)^n Q_n^{\text{mon}}(-y)$ is monic and has the same orthogonality as $Q_n^{\text{mon}}(y)$, so uniqueness proves its parity. The quotient isometry then takes the adjacent pairing to its negative; over $\mathbb{C}$ it is zero. This gives $\phi_N(0) = 0$ by its original definition. Finally $|\det \Sigma|^2 = 1$ makes $\det G_N$ an even differentiable function of θ , proving the derivative equality. These statements hold for every original multiplicity and all three actual sources.

The total phase equality preserves the individual mixed terms. At the original source let $g_K = \partial_\theta \log \det H_N|_0$, $g_B = \partial_\theta \log \det Q_N|_0$, and $\alpha_K = \text{Tr } A_{KK} - c \dim K$. The exact consequences are

$$\begin{aligned}z_B &= -z_K, \quad g_B = -g_K, \quad z_K/\omega_N = \alpha_K - ig_K, \\ \|x_K \otimes y_B - y_K \otimes x_B\|^2 &= a_K d_B + a_B d_K + 2|z_K|^2, \\ \Lambda^\Sigma &= \Lambda \Sigma, \quad I^\Sigma = \Sigma I, \quad H_N^\Sigma = H_N, \quad Q_N^\Sigma = Q_N, \quad L_N^\Sigma = \Sigma L_N, \\ A_{KK}^\Sigma &= kI_K - A_{KK}, \quad A_{BB}^\Sigma = kI_B - A_{BB}, \quad A_{KB}^\Sigma = -A_{KB}, \quad A_{BK}^\Sigma = -A_{BK}.\end{aligned}\tag{OMR25}$$

The first identity follows from OMR16 and OMR24. For the second, $[I, L_N(\theta)]$ has constant determinant: its second block changes only by columns in K , so the frame change is triangular with diagonal identity. The exact splitting determinant therefore gives $g_K + g_B = \partial_\theta \log \det G_N|_0 = 0$. The third identity is OMR22 and the mixed norm follows by substitution in OMR16. For the reflected observation the metric identities follow from OMR24 and the exact minimum section. Substituting $\Sigma A \Sigma = kI - A$ into the four OMR15 blocks proves their signs; both constant off-diagonal terms vanish because $K_N^{\text{coord}} L_N = 0$ and $\Lambda I = 0$. These are the exact maps to the reflected observation even when its kernel differs from K . Neither z_K , α_K nor g_K has been set to zero.

In particular the untilted arithmetic budget in OMR19 becomes

$$\begin{aligned}
J_k^{\text{action}} &= 2 \sum_{N=q-1}^{2q-1} w_N \log \epsilon_N^{\text{ar}} = \Psi_k + C_k^{\text{budget}}, \quad Z_k^{\text{phase}} = 0, \\
\exp\left(\frac{J_k^{\text{action}}}{4q}\right) &= \prod_{N=q-1}^{2q-1} (\epsilon_N^{\text{ar}})^{w_N/(2q)}, \\
4q \log \mathcal{L}_{h,k} &\leq J_k^{\text{action}} \in [L_J, U_J], \quad \mathcal{L}_{h,k} \leq \min_N \epsilon_N^{\text{ar}} \leq \prod_N (\epsilon_N^{\text{ar}})^{w_N/(2q)} \leq e^{U_J/(4q)}, \\
\Phi_{k,k} &\leq \mathcal{F}_{\partial,k}^{(k)} + \mathcal{H}_k + \delta_k^\sigma + e_{k,k}^+ + C_k^{\text{budget}} - 4q \log \mathcal{L}_{h,k}, \\
\Phi_{k,k} &\leq \mathcal{O}_k^+ + \mathcal{H}_k + \delta_k^\sigma + e_{k,k}^+ + C_k^{\text{budget}} - 4q \log \mathcal{L}_{h,k}.
\end{aligned} \tag{OMR26}$$

All allowances are positive by OMR17. Therefore the logarithms exist, substitution of the proved zero total phase gives the first line, and exponentiation gives the exact weighted geometric mean in the second. The third line follows from the minimum and the same positive-class bound. The last two inequalities are the proved OMR21 substitutions at $\theta = 0$, retaining every other term and its sign. The $[L_J, U_J]$ and outer substitutions keep their original simple-quartet domain; the budget and geometric mean identities retain full multiplicity. The generic MCE and MCP tilted-family formulas remain in full, and OMR25 retains the actual mixed phase still to be controlled.

46 The literal original conductor limit and an invertible pivot restriction

Retain all original hypotheses and maps of PQO, RPC, OCF and CEP. In particular $0 < \delta < 1/2$, $\gamma > 2$, the original unit values are $(a, \bar{a}, \bar{a}, a)$ with $a \neq 0$, and the same logarithm branch is used throughout. Write $\beta = 2(\gamma^2 - \delta^2)$, $\eta = (\delta^2 + \gamma^2)^2$, and $\mathfrak{a} = 1/160 + \beta/24 + \eta/2$. No source order is changed.

46.1 An entire coefficient matrix in the inverse original period

Let P_0 be RPC3's centered Fourier period matrix with its factor $e^{a/u}$ removed by the stated exact identity, and retain $G_u = \text{diag}(u^{a/5} g_a)_{a=1}^4$ with RPC8's full Gamma constants. Introduce the coefficient matrix $\mathcal{R}$ by the following identity, rather than replacing the original period:

$$P = e^{a/u} P_0 = e^{a/u} G_u \mathcal{R}(u^{-1}). \quad (\text{PCL1})$$

The row index below is $1 \leq a \leq 4$, and the column index is $0 \leq r \leq 3$. The complete series is

$$\begin{aligned} \mathcal{R}_{ar}(z) &= \sum_{n=0}^{\infty} R_{ar,n} z^n, \\ R_{ar,n} &= \sum_{\substack{p,h \geq 0 \\ 2p+4h=5n+r+1-a}} \frac{\beta^p \eta^h}{3^p p! h!} (-1)^L 5^L (a/5)_L, \quad L = p + h - n. \end{aligned} \quad (\text{PCL2})$$

Every sum at a fixed n is finite. For each contributing pair, $r + 3p + h + 1 = a + 5L$ and $L \geq 0$. To prove the latter assertion, the left side is positive and $1 \leq a \leq 4$, so a negative multiple of five cannot be their difference. The exponent n is nonnegative: its numerator $2p + 4h + a - r - 1$ is an integer multiple of five and is at least -3 . Thus a negative value of n is impossible.

For a direct derivation, divide every term of the absolutely convergent original series PQO9 by $u^{a/5} g_a$. The quotient of its Gamma factors is exactly

$$\frac{5^{(a+5L)/5-1} e^{\pi i(a+5L)/5} \Gamma(a/5 + L)}{g_a} = (-1)^L 5^L (a/5)_L. \quad (\text{PCL3})$$

Its remaining power of u is $u^{L-p-h} = u^{-n}$. These identities prove PCL2 including its signs, factorials and all unit-independent period coefficients. Absolute convergence of PQO9 permits grouping these terms. For each prescribed complex $z \neq 0$, put $u = 1/z$ on any of its original branches. The resulting sum of absolute values is finite. Consequently PCL2 converges absolutely for every complex z , and defines an entire matrix. This also proves that this matrix is independent of the branch; all branch factors remain in G_u in PCL1.

The complete constant matrix in PCL2 is

$$J_{\infty} = \begin{pmatrix} 1 & 0 & -\beta/3 & 0 \\ 0 & 1 & 0 & -2\beta/3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad J_{\infty}^{-1} = \begin{pmatrix} 1 & 0 & \beta/3 & 0 \\ 0 & 1 & 0 & 2\beta/3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (\text{PCL4})$$

Indeed $n = 0$ forces $2p + 4h = r + 1 - a \leq 3$, so either $p = h = 0$ or $p = 1, h = 0, r + 1 - a = 2$. PCL3 gives respectively the diagonal entries and the two negative entries displayed. In particular, the original row-scaled period limit includes both cubic-phase corrections. It is not a diagonal coefficient map.

RPC's determinant proof holds on the complete coefficient family and gives $\det P_0 = \det G_u$. Alternatively divide PQO11 by the product of the four unchanged Gamma ray factors. Therefore

$$\det \mathcal{R}(z) = 1, \quad \mathcal{R}(z)^{-1} = \text{adj } \mathcal{R}(z) \quad (z \in \mathbb{C}). \quad (\text{PCL5})$$

The equality first holds for $z \neq 0$ by PCL1 and extends to zero by continuity. Both the matrix and its inverse are entire. For any fixed $R > 0$, the finite, completely specified constant

$$C_R = \sum_{n \geq 1} \|R_n\|_1 R^{n-1} < \infty \quad (\text{PCL6})$$

gives $\|\mathcal{R}(z) - J_\infty\|_1 \leq C_R|z|$ on $|z| \leq R$. Here $\|\cdot\|_1$ is the induced column-sum matrix norm and $R_n = (R_{ar,n})$. Convergence follows from the entire power series in finite dimension, also at any larger radius. Thus this is a bound in the actual coefficients, with no uncomputed nonzero minor used as a premise.

46.2 All eighty-one actual conductor coefficients and their limit

Let V be the original centered evaluation Vandermonde in PQO12, $U = \text{diag}(a, \bar{a}, \bar{a}, a)$, and $D = \text{diag}(\zeta, \zeta^2, \zeta^3, \zeta^4)$. The complete period matrix from primary to Fourier coordinates is

$$\mathbf{H}(u) = e^{a/u} G_u \mathcal{R}(u^{-1}) V^{-1} U. \quad (\text{PCL7})$$

For $1 \leq j \leq 4$ define the entire matrix and biform

$$\begin{aligned} L_j(z) &= U^{-1} V \mathcal{R}(z)^{-1} D^{-j} \mathcal{R}(z) V^{-1} U, \\ t(Z, W) &= (ZW, Z, W, 1)^\top, \\ A_z(Z, W) &= \prod_{j=1}^4 Q_0(L_j(z) t(Z, W)) = \sum_{r,s=0}^8 a_{rs}(z) Z^r W^s. \end{aligned} \quad (\text{PCL8})$$

These are precisely all original OCF4 coefficients at $z = u^{-1}$. To prove this, compute

$$\mathbf{H}(u)^{-1} D^{-j} \mathbf{H}(u) = L_j(u^{-1}) : \quad (\text{PCL9})$$

the scalar $e^{a/u}$ cancels with its inverse, and the two diagonal matrices G_u, D^{-j} commute. OCF's definition $Q_j(x) = Q_0(\mathbf{H}^{-1} D^{-j} x)$ followed by $x = \mathbf{H}t$ then proves PCL8. The homogeneous version uses $t = (z_0 w_0, z_0 w_1, z_1 w_0, z_1 w_1)$ and retains the bidegree $(8, 8)$. Thus no unit or period entry has been omitted by the affine notation.

For an explicit finite formula for the individual coefficients, write $L_{j,be}$ for a matrix entry, and associate the four pairs $(\rho_e, \sigma_e) = (1, 1), (1, 0), (0, 1), (0, 0)$ to its columns. Put

$$\begin{aligned} c_{j,rs}(z) &= \sum_{\substack{1 \leq e, f \leq 4 \\ \rho_e + \rho_f = r, \sigma_e + \sigma_f = s}} (L_{j,1e} L_{j,4f} - L_{j,2e} L_{j,3f}), \\ a_{rs}(z) &= \sum_{\substack{0 \leq r_j, s_j \leq 2 \\ \sum_j r_j = r, \sum_j s_j = s}} \prod_{j=1}^4 c_{j,r_j s_j}(z). \end{aligned} \quad (\text{PCL10})$$

Expanding the two products defining Q_0 proves the first line; expanding its four original orbit factors proves the second. Both lines are finite and retain all ordered-pair multiplicities. Equations PCL2, PCL5 and PCL10 are a convergent formula for every actual coefficient, with no freely assigned amplitude.

Their exact leading values are obtained by the explicit matrices

$$L_j(0) = U^{-1} V J_\infty^{-1} D^{-j} J_\infty V^{-1} U, \quad a_{rs}(u^{-1}) = a_{rs}(0) + O_{\text{actual}}(|u|^{-1}). \quad (\text{PCL11})$$

The equality for $L_j(0)$ follows from PCL4, and the bound follows from the entire coefficient series. More explicitly, if $a_{rs}(z) = \sum_{n \geq 0} a_{rs,n} z^n$, the finite constant $\sum_{r,s} \sum_{n \geq 1} |a_{rs,n}| R^{n-1}$ bounds the sum of all eighty-one errors divided by $|z|$ on $|z| \leq R$. The coefficient functions are single-valued

entire functions of u^{-1} even though the original period itself retains its logarithm branch. This assertion follows from the exact conjugation PCL9, not from a change of the original branch.

At any specified k , the complete central convolution section is therefore the explicitly evaluated entire matrix

$$C_{\text{in}}(z)_{(a,b),(i,j)} = a_{i+4-a,j+4-b}(z), \quad 0 \leq a, b, i, j \leq k-8, \quad (\text{PCL12})$$

with all out-of-range coefficients zero. PCL10–12 give its full constant matrix and all successive coefficients. They do not discard off-diagonal coefficients or replace this matrix by its central coefficient a_{44} . In particular the calculation of the limit has not asserted a sign or diagonal-dominance inequality for this complete original matrix.

46.3 An exact positivity slice for the literal positive period ray

For $u > 0$ on any of its original logarithm lifts, PQO21 gives $P_0 = E_{\text{br}}R$ with R real invertible and E_{br} diagonal unitary commuting with D . Its exact centered quadratic coefficient matrix is S of PQO13; all its entries are purely imaginary. Consequently, for real $y \in \mathbb{R}^4$,

$$\overline{Q_j(E_{\text{br}}y)} = -Q_{5-j}(E_{\text{br}}y) \quad (1 \leq j \leq 4). \quad (\text{PCL13})$$

Indeed the original quadratic matrix in these coordinates is $e^{-2a/u}R^{-\top}SR^{-1}$, which is purely imaginary. Complex conjugation changes its sign and changes D^{-j} into D^j ; this is precisely the displayed formula.

Let P_{conj} be the permutation matrix exchanging primary indices 1, 2 and 3, 4. For real Z and $|W| = 1$, direct substitution gives $P_{\text{conj}}t(Z, W) = W^{-1}t(Z, W)$. The original roots and unit satisfy $\bar{V} = P_{\text{conj}}V$ and $P_{\text{conj}}\bar{U} = UP_{\text{conj}}$. Thus

$$\overline{V^{-1}Ut(Z, W)} = W^{-1}V^{-1}Ut(Z, W). \quad (\text{PCL14})$$

Choose either square root w of W . Then $p = w^{-1}V^{-1}Ut(Z, W)$ is real by PCL14, and $Ht = wE_{\text{br}}y$ for the real vector $y = e^{a/u}Rp$. Homogeneity and PCL13 prove the exact nonnegative slice identity

$$\boxed{W^{-4}A_{u^{-1}}(Z, W) = |Q_1(E_{\text{br}}y)Q_2(E_{\text{br}}y)|^2 \geq 0 \quad (Z \in \mathbb{R}, |W| = 1).} \quad (\text{PCL15})$$

The choice of sign of w has no effect because the quadratic forms have even degree. In particular, comparing the Laurent coefficients on the unit circle, and then the polynomial coefficients in real Z , gives the exact symmetry

$$a_{rs}(u^{-1}) = \overline{a_{r,8-s}(u^{-1})}. \quad (\text{PCL16})$$

PCL11 extends this identity and PCL15 to the coefficient limit at $u^{-1} = 0$ by continuity. The slice in PCL15 has one real coordinate and one unit-circle coordinate; it has not changed either coordinate into a second unit circle.

46.4 A complete invertible restriction supplied by the original pivot

At each actual u in the proved five-orbit locus, use precisely OCF5's lexicographically largest nonzero coefficient (r_*, s_*) , $a_* = a_{r_*s_*} \neq 0$, and its original pivot rectangle $\mathcal{P}_k = \{(r_* + i, s_* + j) : 0 \leq i, j \leq k-8\}$. Let $\mathcal{E}_k$ be its complement and S_k the original insertion of its coordinates. No new coefficient has been selected. The full original maps satisfy

$$CN_k^{\top} = 0, \quad S_k^{\top}N_k^{\top} = I_{\Delta}, \quad \text{im } N_k^{\top} = V_A. \quad (\text{PCL17})$$

The first is the transpose of $N_k C^{\top} = 0$ in OCF7; the second is the transpose of $N_k S_k = I$. Dimension Δ then proves the last equality, since C has full row rank. It follows that restriction to the actual strip is the exact isomorphism

$$S_k^{\top} : V_A \longrightarrow \mathbb{C}^{\mathcal{E}_k}, \quad (S_k^{\top}|_{V_A})^{-1} = N_k^{\top}. \quad (\text{PCL18})$$

Equivalently its square pivot block has nonzero diagonal a_* and is triangular in the elimination order; OCF's complete division supplies its inverse. This is an alternative restriction to the central one in PCL12, with the exact changed set of roots.

Let $P_{\mathcal{P},k}(y) = \prod_{(a,b) \in \mathcal{P}_k} (y - \omega_{ab})$ and $P_{\mathcal{E},k}(y) = \prod_{(a,b) \in \mathcal{E}_k} (y - \omega_{ab})$, where $\omega_{ab} = (2b - k)\gamma - i(2a - k)\delta$. Thus $P_k = P_{\mathcal{P},k}P_{\mathcal{E},k}$ exactly. Let $V_{\mathcal{E}}$ evaluate $1, y, \dots, y^{\Delta-1}$ at those actual outer roots and set $D_{\mathcal{P},\mathcal{E}} = \text{diag}(P_{\mathcal{P},k}(\omega_{ab}))_{\mathcal{E}_k}$. These two matrices are invertible by distinctness of the original roots. The full polynomial-target restriction and its exact inverse are

$$\begin{aligned}\rho_{\mathcal{E}} &= V_{\mathcal{E}}^{-1} D_{\mathcal{P},\mathcal{E}}^{-1} S_k^{\top}|_{V_A}, \\ \rho_{\mathcal{E}}^{-1} &= N_k^{\top} D_{\mathcal{P},\mathcal{E}} V_{\mathcal{E}}.\end{aligned}\tag{PCL19}$$

Multiplying these matrices and using PCL17 proves both inverse identities, with all inner products of root differences retained. In particular the original residual becomes the following specified onto map, with its exact complete kernel:

$$\Xi_{\mathcal{E}} = \Xi_k^{\text{fix}} N_k^{\top} D_{\mathcal{P},\mathcal{E}} V_{\mathcal{E}}, \quad \ker \Xi_{\mathcal{E}} = \rho_{\mathcal{E}}(K).\tag{PCL20}$$

This follows by composition with the proved isomorphism, not by deleting any common-kernel vector.

Finally the original metric and its entire source graph are carried by this isomorphism as follows. Write $J_{\mathcal{E}} = N_k^{\top} D_{\mathcal{P},\mathcal{E}} V_{\mathcal{E}}$. At every original endpoint the actual conductor metric in these outer coordinates is exactly

$$H_{\mathcal{E},N} = J_{\mathcal{E}}^* G_N J_{\mathcal{E}}, \quad Q_{\Xi,N}^{\text{fix}} = (\Xi_{\mathcal{E}} H_{\mathcal{E},N}^{-1} \Xi_{\mathcal{E}}^*)^{-1}.\tag{PCL21}$$

The first formula is the pullback along $\rho_{\mathcal{E}}^{-1}$; the minimum-section proof of GIP6 gives the second. On the full GIP graph its target map is $\rho_{\mathcal{E}} E_N^{\text{graph}}$, with the same complete t ideal columns, and composing with $\Xi_{\mathcal{E}}$ gives exactly B_N^{graph} . Its residual kernel is therefore all $t + r$ columns in GIP4, including the low- v coordinates. The coefficient-space bound for N_k does not remove the displayed root-value and Gamma matrices from PCL21. These formulas retain the full metric transport needed for the next comparison.

46.5 The exact inverse-period reflection of the central section

Let $E_{\text{par}} = \text{diag}(1, -1, 1, -1)$ in the centered coefficient order, and let P_{opp} exchange primary indices 1, 4 and 2, 3. Equation PCL2 implies

$$\mathcal{R}(-z) = E_{\text{par}} \mathcal{R}(z) E_{\text{par}}, \quad V E_{\text{par}} = P_{\text{opp}} V, \quad U P_{\text{opp}} = P_{\text{opp}} U.\tag{PCL22}$$

For the first equality a contributing coefficient satisfies $5n + r + 1 - a = 2p + 4h$, so $n \equiv a - r - 1 \pmod{2}$. The signs from the two parity matrices on entry (a, r) are $(-1)^{a-1+r} = (-1)^n$. Summing the entire series proves the identity. The second equality evaluates each monomial at the opposite original root. The third uses the full original unit values $(a, \bar{a}, \bar{a}, a)$, which agree at opposite roots.

The two diagonal matrices E_{par} and D commute. Substitution into PCL8 and use of PCL22 therefore give the exact maps

$$L_j(-z) = P_{\text{opp}} L_j(z) P_{\text{opp}}, \quad A_{-z}(Z, W) = Z^8 W^8 A_z(Z^{-1}, W^{-1}).\tag{PCL23}$$

For the second identity, $Q_0(P_{\text{opp}} t) = Q_0(t)$ by its two explicit products. Also $P_{\text{opp}} t(Z, W) = ZW t(Z^{-1}, W^{-1})$ when $ZW \neq 0$. Each of the four quadratic factors therefore contributes $(ZW)^2$. This proves the identity on that open set; both sides are polynomials after multiplication by $Z^8 W^8$, so it holds identically including both coordinate axes.

Comparing every coefficient in PCL23 proves

$$a_{rs}(-z) = a_{8-r, 8-s}(z), \quad C_{\text{in}}(-z) = C_{\text{in}}(z)^{\top}, \quad \det C_{\text{in}}(-z) = \det C_{\text{in}}(z).\tag{PCL24}$$

Indeed the $(a, b), (i, j)$ entry on the left of the middle identity is $a_{4+a-i, 4+b-j}(z)$, precisely the transposed entry of PCL12. Out-of-range coefficients are zero on both sides. The determinant equality follows by the permutation expansion of a determinant and its transpose. Thus the exact central determinant is an even entire function of u^{-1} , with every coefficient supplied by PCL2 and PCL10–12. This retains its actual constant matrix and does not assign that determinant a value by deleting any of its entries.

47 The original relation operator and its finite-rank graph transport

Retain the complete original data and maps of CEP1–49 and GIP1–9. In particular the period and its logarithm branch are fixed on the actual five-orbit locus, $k = 4l + 1 \geq 9$, and

$$q = (k + 1)^2, \quad q' = (k - 7)^2, \quad \Delta = q - q', \quad c = k/2, \quad c' = c - 4, \quad 0 \leq v \leq 80. \quad (\text{ROG1})$$

All 81 original coefficients a_{ab} and their actual shifts $b_{ab} = 4 + (2a - 8)\delta + i(2b - 8)\gamma$ are retained. Write $\mu_j = \sum a_{ab} b_{ab}^j$, with $\mu_0 = \dots = \mu_{v-1} = 0$ and $\mu_v \neq 0$. For an original endpoint $N \in \{q - 1, q, 2q - 1, 2q\}$ put

$$L = N - q, \quad \mathbf{m} = \Delta - v, \quad M = L + \mathbf{m}, \quad d = N - v = q' + M. \quad (\text{ROG2})$$

Thus $L = -1, 0, q - 1, q$ at the four endpoints, respectively. The polynomial spaces of degree at most -1 are zero. Since $\Delta \geq 96 > v$, every displayed M is nonnegative.

47.1 The complete positive-shift relation operator

Let $\chi = \chi_k$ and $\chi' = \chi_{k-8}$ denote the original monic polynomials in S and S' , respectively; the prime here denotes the lower-grid polynomial and never a derivative. For $0 \leq a, b \leq 8$ define the actual monic product

$$d_{ab}(S') = \prod_{(i,j) \notin \{a, \dots, a+k-8\} \times \{b, \dots, b+k-8\}} (S' + b_{ab} - \beta_{ij}), \quad 0 \leq i, j \leq k. \quad (\text{ROG3})$$

The root identity $\beta'_{ij} + b_{ab} = \beta_{i+a, j+b}$ gives $\chi(S' + b_{ab}) = \chi'(S')d_{ab}(S')$ by matching individual factors, with the same leading coefficient one. Consequently

$$\begin{aligned} (\mathcal{U}_L h)(S') &= \sum_{a,b=0}^8 a_{ab} d_{ab}(S') h(S' + b_{ab}), \\ \chi' \mathcal{U}_L h &= \mathcal{T}_A(\chi h), \quad \mathcal{T}_{Ap} = \sum_{a,b} a_{ab} p(S' + b_{ab}). \end{aligned} \quad (\text{ROG4})$$

These are polynomial equalities with positive shifts. They include all original period coefficients and contain no conjugate or binomial multiplier from the biform pairing: multiplication of the original biform monomials adds their two indices with coefficient one, and their complex-linear primary pairing evaluates p at the sum of their roots, exactly as in CEP3–4.

For $0 \leq j \leq L$ write

$$u_j = \mathcal{U}_L(S^j), \quad \lambda_j = \binom{q+j}{v} \mu_v. \quad (\text{ROG5})$$

Every u_j has degree $\mathbf{m} + j$ and leading coefficient λ_j . Indeed Taylor expansion gives $\mathcal{T}_{Ap} = \sum_{t \geq v} \mu_t p^{(t)} / t!$, a finite sum on every polynomial. Applying it to the monic degree- $q+j$ polynomial χS^j gives leading coefficient $\binom{q+j}{v} \mu_v$ in degree $q+j-v$. Division by the monic degree- q' polynomial gives the asserted degree and coefficient. In particular $\mathcal{U}_L$ is injective.

47.2 An explicit finite inverse with coefficient control

The following constants are expressed in the original coordinates:

$$\begin{aligned} R &= |c| + k\sqrt{\delta^2 + \gamma^2}, \quad R' = |c'| + (k-8)\sqrt{\delta^2 + \gamma^2}, \\ B &= \max_{a,b} |b_{ab}|, \quad A_{\text{abs}} = \sum_{a,b} |a_{ab}|, \quad C_v = \frac{A_{\text{abs}} B^v}{|\mu_v|} \geq 1. \end{aligned} \quad (\text{ROG6})$$

All roots of χ and χ' have modulus at most R and R' , respectively. $B \geq 4$ and $C_v \geq 1$ follow directly from the shift list and $|\mu_v| \leq A_{\text{abs}} B^v$. For $L \geq 0$ put

$$C_L = qR + 2qB + (q' + L)R', \quad \tau_L = \min \left\{ 1, \frac{\log(1 + (2C_v)^{-1})}{C_L} \right\} > 0. \quad (\text{ROG7})$$

For every $0 \leq h \leq j \leq L$, the complete coefficient estimate is

$$\left| [S^{\mathbf{m}+j-h}] \frac{u_j(S')}{\lambda_j} \right| \leq C_v \frac{C_L^h}{h!}. \quad (\text{ROG8})$$

Here the coefficient for $h = 0$ is exactly one, which will be retained separately in the inverse.

To prove ROG8, write the coefficient of S^{q-a} in χ as χ_a . Its absolute value is bounded by $\binom{q}{a} R^a \leq (qR)^a / a!$, by expanding the product of its actual roots. For $n = q + j$, the term of derivative order $v + b$ applied to $\chi_a S^{n-a}$ contributes $\chi_a \mu_{v+b} \binom{n-a}{v+b}$; it is zero when the derivative order exceeds the degree. Its ratio to $\mu_v \binom{n}{v}$ is bounded by

$$C_v \frac{(qR)^a}{a!} \frac{(nB)^b}{b!}, \quad (\text{ROG9})$$

because $|\mu_{v+b}/\mu_v| \leq C_v B^b$ and $\binom{n-a}{v+b} / \binom{n}{v} \leq n^b / b!$. For the latter assertion first remove a using monotonicity of a binomial coefficient in its top entry; then the ratio is $\prod_{r=0}^{b-1} (n-v-r)/(v+1+r) \leq n^b / b!$.

In the formal expansion at infinity,

$$\frac{1}{\chi'(S')} = S'^{-q'} \sum_{t \geq 0} H_t S'^{-t}, \quad |H_t| \leq \binom{q' + t - 1}{t} (R')^t \leq \frac{((q' + h)R')^t}{t!} \quad (t \leq h). \quad (\text{ROG10})$$

Expansion of each $(1 - \beta'_{ab}/S')^{-1}$ proves the complete homogeneous coefficient formula and this estimate. To obtain the specified coefficient of u_j , sum ROG9–10 over $a + b + t = h$. Since $n \leq 2q$ and $h \leq L$, the multinomial identity gives exactly ROG8. Formal division agrees with ordinary polynomial division here because ROG4 proved exact divisibility before the expansion was made.

Let H_L be the square matrix consisting of the coefficients of $u_0, \dots, u_L$ in degrees $\mathbf{m}, \dots, \mathbf{m} + L$. Then

$$H_L = \widehat{H}_L \text{diag}(\lambda_0, \dots, \lambda_L), \quad \det \widehat{H}_L = 1, \quad \det H_L = \mu_v^{L+1} \prod_{j=0}^L \binom{q+j}{v}. \quad (\text{ROG11})$$

The matrix $\widehat{H}_L$ is upper triangular, with diagonal one and offset- h entries bounded by ROG8 for $h \geq 1$. For g in the actual image of $\mathcal{U}_L$, its unique preimage is therefore the finite formula

$$\text{coef } h = \text{diag}(\lambda_0^{-1}, \dots, \lambda_L^{-1}) \sum_{r=0}^L (I - \widehat{H}_L)^r ([S^{\mathbf{m}}]g, \dots, [S^{\mathbf{m}+L}]g)^T. \quad (\text{ROG12})$$

The finite sum is its inverse because $I - \widehat{H}_L$ is strictly upper triangular and its $(L+1)$ st power is zero. Thus this formula retains every coefficient of the original operator, and every complex factor λ_j .

In increasing monomial coefficient Euclidean norms it satisfies

$$\|\text{coef } h\|_2 \leq I_L \|\text{coef } g\|_2, \quad I_L = \frac{1 + 2 \sum_{h=1}^L \tau_L^{-h}}{|\mu_v| \binom{q}{v}}. \quad (\text{ROG13})$$

To check the bound, products of strictly upper triangular entries at a total offset h are majorized by the coefficient of z^h in

$$\sum_{r \geq 0} \{C_v(e^{C_L z} - 1)\}^r = \frac{1}{1 - C_v(e^{C_L z} - 1)}. \quad (\text{ROG14})$$

The absolute value of the expression inside braces is at most $1/2$ on $|z| = \tau_L$, so Cauchy's coefficient formula bounds these coefficients by $2\tau_L^{-h}$. The actual constant diagonal is one. Every absolute row and column sum of the inverse is consequently at most the numerator in ROG13. The Euclidean operator norm is at most the square root of the product of these row and column bounds, by applying Cauchy–Schwarz in each row and summing columns. Finally $|\lambda_j| \geq |\mu_v| \binom{q}{v}$ proves ROG13.

For fixed original period and quartet, $L \leq q$ gives

$$\log^+ I_L = O_{\text{actual}}(q \log(q+1)). \quad (\text{ROG15})$$

Indeed $R, R' = O(k)$, B, C_v, μ_v, v are fixed, $C_L = O_{\text{actual}}(qk)$, and the numerator in ROG13 is at most $(2L+1) \max\{1, \tau_L^{-L}\}$. No lower bound for a moment varying with the period is used.

47.3 Complete original-source bounds for the low graph map

For source $s \in \{1, k\}$ retain

$$b = s/2, \quad M_s = (2\pi)^{s/2}, \quad h_n = M_s n! (b)_n, \quad \|p\|_{s, c_0}^2 = \int |p(c_0 + iy)|^2 dm_s(y). \quad (\text{ROG16})$$

The density dm_s is exactly the original Gamma density in GIP5 and OQ2. We now supply elementary finite coefficient bounds, so no implicit conditioning of a polynomial frame is needed in the graph estimate. For $D \geq 0$ and real centre c_0 define

$$\begin{aligned} A_{D, s, c_0}^+ &= \sqrt{M_s(D+1)} \max\{1, |c_0| + 2\sqrt{D(b+D)}\}^D, \\ A_{D, s, c_0}^- &= \frac{\sqrt{D+1}}{\sqrt{M_s \min\{1, b\}}} \max\{1, (1+|c_0|)(1+D(b+D))\}^D. \end{aligned} \quad (\text{ROG17})$$

Then for every polynomial of degree at most D ,

$$\|p\|_{s, c_0} \leq A_{D, s, c_0}^+ \|\text{coef } p\|_2, \quad \|\text{coef } p\|_2 \leq A_{D, s, c_0}^- \|p\|_{s, c_0}. \quad (\text{ROG18})$$

For the first inequality, the original monic Gamma recurrence gives in its orthonormal basis

$$y\varphi_n = \sqrt{(n+1)(b+n)}\varphi_{n+1} + \sqrt{n(b+n-1)}\varphi_{n-1}.$$

On the degrees needed to multiply a polynomial of degree at most $D-1$, the upper and lower shift norms are at most $\sqrt{D(b+D)}$. Iterating the resulting bound for $S = c_0 + iy$, starting with $\|1\| = \sqrt{M_s}$, gives $\|S^a\| \leq \sqrt{M_s}(|c_0| + 2\sqrt{D(b+D)})^a$ for $a \leq D$. Cauchy–Schwarz on the coefficient sum proves the first formula in ROG18.

For the second, the coefficient absolute sum of the monic $p_n(y)$ is at most $(1+D(b+D))^n$ for $n \leq D$. The recurrence $p_{n+1} = yp_n - n(b+n-1)p_{n-1}$ proves this by induction: with $A = D(b+D)$, the required scalar inequality is $(1+A) + A \leq (1+A)^2$. Substitution of $(S - c_0)/i$ multiplies the degree- n coefficient bound by at most $(1+|c_0|)^n$, retaining the phase i^{-n} in the polynomial itself. Also $h_n \geq M_s \min\{1, b\}$: it holds at $n = 0, 1$, and $h_n/h_{n-1} = n(b+n-1) \geq 1$ for $n \geq 2$. The Euclidean norm of the coefficient column of each $p_n((S - c_0)/i)/\sqrt{h_n}$ is therefore bounded by the corresponding term in ROG17. Expanding in that orthonormal basis and applying Cauchy–Schwarz proves the second formula.

Define the explicit divisor constant

$$D_{M, \chi'} = \sum_{a=0}^M \binom{q' + a - 1}{a} (R')^a. \quad (\text{ROG19})$$

For $f = \chi'g$ of degree at most $d = q' + M$, ROG10 gives

$$\|\text{coef } g\|_2 \leq D_{M, \chi'} \|\text{coef } f\|_2. \quad (\text{ROG20})$$

Indeed the high coefficients of the quotient are obtained by convolution with the first $M + 1$ coefficients of $1/\chi'$. Every absolute row and column sum is bounded by ROG19; the same row and column argument proves the norm assertion. This bound is for the exact divisible input and retains the whole divisor. It does not assert a contractive division map.

Equip $\mathbb{C}[S']_{\leq M}$ with the weighted original norm

$$\|g\|_{\chi',s,c'} = \|\chi'g\|_{s,c'}, \quad G_M = (\langle \chi'S'^a, \chi'S'^b \rangle_{s,c'})_{a,b=0}^M. \quad (\text{ROG21})$$

The Gram is positive definite because multiplication by the nonzero polynomial is injective. Let $\mathbf{U}_L$ be the complete coefficient matrix of ROG4. For $L \geq 0$ the exact weighted orthogonal projection onto its image is

$$\Pi_{U,L} = \mathbf{U}_L(\mathbf{U}_L^* G_M \mathbf{U}_L)^{-1} \mathbf{U}_L^* G_M. \quad (\text{ROG22})$$

Its Gram inverse exists by ROG5. Multiplication verifies idempotence, weighted self-adjointness, and its image, so it has weighted norm one. At $L = -1$ put $\Pi_{U,L} = 0$.

Let $Z_{\chi,N}h$ be the original upper orthonormal coefficient column of χh , exactly as in GIP3. Define the complete low graph map on this weighted space by

$$Q_{N,s}g = P_{<v} Z_{\chi,N} \mathbf{U}_L^{-1} \Pi_{U,L} g : \mathbb{C}[S']_{\leq M} \longrightarrow \mathbb{C}^v. \quad (\text{ROG23})$$

The inverse is used only on its actual image, where ROG12 proved it uniquely. For $L = -1$ this map is zero. For $L \geq 0$ its norm is at most the following entirely finite constant:

$$H_{N,s} = A_{N,s,c}^+ (1+R)^q I_L D_{M,\chi'} A_{d,s,c'}^-. \quad (\text{ROG24})$$

To verify it, begin with $\Pi_{U,L}g$, whose weighted norm does not exceed $\|g\|_{\chi',s,c'}$. Apply ROG18 to $\chi'\Pi_{U,L}g$, then ROG20 and ROG13 to recover the ordinary coefficients of its unique h . Multiplication by χ has coefficient Euclidean norm at most $(1+R)^q$: the sum of the absolute coefficients of its full monic product is at most that number, and coefficient convolution has row and column sums bounded by this sum. The first bound in ROG18 now gives the original upper source norm of χh . Finally $P_{<v}$ is an orthogonal coordinate projection and has norm at most one. These are precisely the factors in ROG24.

At every original endpoint with $L \geq 0$,

$$\log(1 + H_{N,s}) = O_{\text{actual}}(q \log(q+1)), \quad s = 1, k. \quad (\text{ROG25})$$

For detail, $N, d \leq 2q$, $M \leq 2q$, $b \leq k/2$ and $|c|, |c'|, R, R' = O_{\delta,\gamma}(k)$. Both logarithms of the powered factors in ROG17 are $O(q \log(q+1))$. The factors $\sqrt{M_s}$ and $1/\sqrt{M_s}$ in ROG24 cancel at this specified composition; the individual source norms ROG16–18 retain their full mass. The factor $1/\sqrt{\min\{1, b\}}$ is at most $\sqrt{2}$. ROG19 is bounded by $(M+1) \max\{1, (q'+M)R'\}^M$, which has the same logarithmic order. ROG15 and $q \log(1+R) = O(q \log(q+1))$ control the other two factors. These estimates prove ROG25.

47.4 The exact graph shear and its complementary singular pairs

Use exactly the GIP source space and product metric

$$X_N = \mathbb{C}^v \oplus \mathbb{C}[S']_{\leq M}, \quad \|(x, g)\|^2 = \|x\|_2^2 + \|g\|_{\chi',s,c'}^2. \quad (\text{ROG26})$$

The full relation graph and its image under the specified invertible map are

$$\begin{aligned} \mathcal{R}_{N,s}(x, g) &= (x - Q_{N,s}g, g), & \mathcal{R}_{N,s}^{-1}(x, g) &= (x + Q_{N,s}g, g), & \det \mathcal{R}_{N,s} &= 1, \\ \mathcal{R}_{N,s} \text{ im } \begin{pmatrix} P_{<v} Z_{\chi,N} \\ \mathbf{U}_L \end{pmatrix} &= \{0\} \oplus \text{im } \mathbf{U}_L. \end{aligned} \quad (\text{ROG27})$$

The last equality follows by applying ROG23 to $\mathcal{U}_L h$; the actual low-degree column is subtracted with its full value. The inverse and determinant follow from the displayed triangular blocks. Thus ROG27 is an exact morphism between the full original graph and the indicated relation image.

Put $r_Q = \text{rank } Q_{N,s} \leq v$ and let its nonzero singular values in the specified metrics be σ_a . Singular value decompositions on its domain and codomain put the shear into r_Q orthogonal two-dimensional blocks $\begin{pmatrix} 1 & -\sigma_a \\ 0 & 1 \end{pmatrix}$ and identity blocks. The two singular values of such a block are exactly

$$t_a = \frac{\sqrt{\sigma_a^2 + 4} + \sigma_a}{2}, \quad t_a^{-1}. \quad (\text{ROG28})$$

Indeed its positive Gram has determinant one and trace $2 + \sigma_a^2$; solving its quadratic characteristic polynomial gives these positive square roots. In particular $1 \leq t_a \leq 1 + \sigma_a \leq 1 + H_{N,s}$. For every exterior degree e , including zero and full degree, both the forward and inverse shear therefore satisfy

$$\log \left\| \bigwedge^e \mathcal{R}_{N,s}^{\pm 1} \right\| \leq \min\{e, v\} \log(1 + H_{N,s}). \quad (\text{ROG29})$$

There are at most v singular values above one; any remaining factors are at most one. This is the complementary-singular-value estimate with the actual rank, rather than multiplying the scalar inverse bound by the full source dimension. At $v = 0$ the map is the identity and this error is exactly zero.

47.5 The unchanged residual target and its four endpoint metrics

Retain the complete onto residual map B_N^{graph} and its common-kernel lift columns $W_{K,N}$ from GIP2–4. Define

$$B_{N,s}^{\text{str}} = B_N^{\text{graph}} \mathcal{R}_{N,s}^{-1}, \quad \ker B_{N,s}^{\text{str}} = \text{im} \left[\begin{pmatrix} 0 \\ \mathbf{U}_L \end{pmatrix}, \mathcal{R}_{N,s} W_{K,N} \right]. \quad (\text{ROG30})$$

The kernel equality is the image under the invertible shear of the full GIP4 kernel. It retains all $r = 8k - 16$ original common-kernel columns and all $L + 1$ relation columns; at $L = -1$ the latter list is empty. Let H_N^{graph} be the product Gram of ROG26, which is exactly GIP5, and put

$$Q_{N,s}^{\text{str}} = [B_{N,s}^{\text{str}} (H_N^{\text{graph}})^{-1} (B_{N,s}^{\text{str}})^*]^{-1}. \quad (\text{ROG31})$$

This is the attained minimum metric on the same original residual coordinate space $\mathbb{C}^j$, $j = 8k - 32$. For an onto matrix B and positive H , its minimum lift of y is $H^{-1} B^* (B H^{-1} B^*)^{-1} y$; subtraction of this lift from any other lift gives an H -orthogonal kernel vector. This proves the inverse-Gram assertion directly.

The shear induces the identity of the residual targets with these two quotient norms. A minimum lift is sent to its shear image and then projected off the entire target kernel. Orthogonal projection contracts every exterior power because its singular values are zero or one. Apply this observation also to the inverse shear and then square the two volume bounds in ROG29. It proves

$$|\log \det Q_{N,s}^{\text{str}} - \log \det \tilde{Q}_{\Xi,N}| \leq 2 \min\{j, v\} \log(1 + H_{N,s}). \quad (\text{ROG32})$$

Here $\tilde{Q}_{\Xi,N}$ is precisely GIP6 with its original complete graph metric. At $L = -1$ set $H_{N,s} = 0$; the two metrics are then identical. Combining with the already proved GIP6 source-map estimate gives the fully specified original residual comparison

$$|\log \det Q_{N,s}^{\text{str}} - \log \det Q_{\Xi,N}^{\text{fix}}| \leq 2E_N^{\text{CEP}} + 2v \log(1 + H_{N,s}). \quad (\text{ROG33})$$

Thus, with the unchanged signs $(1, 1, -1, -1)$ and original endpoints,

$$\left| \sum_N \sigma_N \log \det Q_{N,s}^{\text{str}} - \sum_N \sigma_N \log \det Q_{\Xi,N}^{\text{fix}} \right| \leq 2 \sum_N E_N^{\text{CEP}} + 2v \sum_N \log(1 + H_{N,s}) = o(kq). \quad (\text{ROG34})$$

CEP36 gives the first estimate, while ROG25 and fixed $v \leq 80$ give $O_{\text{actual}}(q \log(q+1)) = o(kq)$ for the second. This proves the transport for both complete original sources without removing a low component by assumption.

The resulting relation quotient is, with its product quotient norm,

$$\mathbb{C}^v \oplus (\mathbb{C}[S']_{\leq L+\Delta-v} / \mathcal{U}_L \mathbb{C}[S]_{\leq L}) . \quad (\text{ROG35})$$

The residual map on it is exactly the descended ROG30 map. In particular the second denominator remains the full operator ROG4; no equality with multiplication by the outer divisor is used in ROG1–35. Its complete leading coefficient and exact inverse are ROG5 and ROG12. The natural map to that outer divisor and its exact kernel and cokernel remain GIP7–9; their finite-convolution obstruction is carried unchanged by ROG27 and ROG30. The new finite-rank transport proves that the full low- v graph itself has only the explicit subleading cost ROG34 in the original residual return.

48 The exact differential realization of the original relation quotient

Retain the original $\chi_k(S)$, $\chi_{k-8}(S')$, all 81 coefficients and shifts of CEP41–42 and ROG1–6. Write $\chi = \chi_k$, $\chi' = \chi_{k-8}$, where the prime denotes the lower-grid polynomial, and put $\mathbf{m} = q - q' - v$. The cutoff-independent polynomial map is

$$\mathcal{U} : \mathbb{C}[S] \longrightarrow \mathbb{C}[S'], \quad \mathcal{U}h = \sum_{a,b=0}^8 a_{ab}d_{ab}(S')h(S' + b_{ab}). \quad (\text{RDS1})$$

It restricts to every original $\mathcal{U}_L$. For all $j \geq 0$, $u_j = \mathcal{U}(S^j)$ has degree $\mathbf{m} + j$ and leading coefficient $\lambda_j = \binom{q+j}{v} \mu_v \neq 0$ by exactly the Taylor and monic-division proof in ROG4–5, which holds at every finite polynomial degree.

48.1 A finite-dimensional quotient and its analytic dual

Repeated subtraction of the unique highest-degree multiple u_j/λ_j gives the direct vector-space decomposition

$$\mathbb{C}[S'] = \mathbb{C}[S']_{<\mathbf{m}} \oplus \mathcal{U}\mathbb{C}[S]. \quad (\text{RDS2})$$

At each step the degree strictly decreases. No nonzero sum $\mathcal{U}h$ can have degree below $\mathbf{m}$, by RDS1, which proves directness and uniqueness. Thus the resulting remainder map is linear and independent of the polynomial cutoff. Its restrictions identify every CEP43 quotient with the same vector space of dimension $\mathbf{m}$.

For a complex-linear functional ℓ on polynomials use its exponential generating series

$$f_\ell(z) = \sum_{n \geq 0} \ell(S'^n) \frac{z^n}{n!}. \quad (\text{RDS3})$$

Initially this is a formal series. The subspace annihilating $\text{im } \mathcal{U}$ has dimension $\mathbf{m}$: its first $\mathbf{m}$ values are arbitrary, and the equations $\ell(u_j) = 0$ uniquely determine the value in degree $\mathbf{m} + j$ from earlier ones, since its coefficient is the nonzero λ_j . We now realize this whole formal space as convergent series using the original symbol.

Put $D = \partial_z$ and define

$$E_A(z) = \sum_{a,b=0}^8 a_{ab}e^{b_{ab}z}, \quad \mathcal{L}_A = \sum_{a,b=0}^8 a_{ab}e^{b_{ab}z}d_{ab}(D). \quad (\text{RDS4})$$

The coefficients multiplying powers of D are on their displayed left side. The precise adjoint and intertwining identities, in this formal-series pairing, are

$$\ell(\mathcal{U}e^{Sz}) = \mathcal{L}_A f_\ell(z), \quad \mathcal{L}_A \chi'(D) = \chi(D)M_{E_A}. \quad (\text{RDS5})$$

For the first, multiplication of S'^r inside the functional is differentiation D^r , while the positive shift contributes $e^{b_{ab}z}$. Thus every term is exactly $a_{ab}e^{b_{ab}z}d_{ab}(D)f_\ell$. For the second, the full polynomial product in CEP41 gives $d_{ab}(D)\chi'(D) = \chi(D + b_{ab})$. Leibniz' rule gives $\chi(D)(e^{b_{ab}z}F) = e^{b_{ab}z}\chi(D + b_{ab})F$ for every formal F . Summing these identities proves RDS5, including the operator order and all coefficients.

Define the two original exponential spaces and a jet kernel by

$$\begin{aligned} \mathcal{E}_k &= \text{span}\{e^{\beta_{ij}z} : 0 \leq i, j \leq k\}, & \mathcal{E}_{k-8} &= \text{span}\{e^{\beta'_{ab}z} : 0 \leq a, b \leq k-8\}, \\ \mathcal{N}_v &= \{N \in \mathcal{E}_k : N^{(a)}(0) = 0 \ (0 \leq a < v)\}. \end{aligned} \quad (\text{RDS6})$$

The q upper and q' lower roots are distinct. Their full Vandermonde derivative matrices at zero prove independence of the respective exponentials. The first v upper derivative rows have rank

v , since any chosen v distinct root columns give a nonzero Vandermonde determinant. Hence $\dim \mathcal{N}_v = q - v$, also when $v = 0$.

The original sum identity gives the injective multiplication map

$$M_{E_A} : \mathcal{E}_{k-8} \hookrightarrow \mathcal{N}_v, \quad E_A(z) e^{\beta'_{ab} z} = \sum_{r,s=0}^8 a_{rs} e^{\beta_{a+r,b+s} z}. \quad (\text{RDS7})$$

Its numerical coefficient matrix is exactly $C^\top$ from CEP4, with the original order and no conjugation. The product vanishes to order at least v at zero because E_A has that exact zero order. Multiplication by this nonzero analytic function is injective: if its product with an analytic function is zero, that function vanishes on a punctured neighbourhood with nonzero E_A and hence identically.

There is the exact sequence and specified quotient isomorphism

$$0 \longrightarrow \mathcal{E}_{k-8} \xrightarrow{M_{E_A}} \mathcal{N}_v \xrightarrow{\mathfrak{F}_A} (\mathbb{C}[S']/\mathcal{UC}[S])^\vee \longrightarrow 0, \quad (\text{RDS8})$$

$$f_{\mathfrak{F}_A(N)}(z) = \chi'(D) \left(\frac{N(z)}{E_A(z)} \right).$$

The ratio is analytic at zero because its numerator vanishes to order at least v , the exact order of its denominator. It defines a functional by the Taylor derivatives in RDS3. RDS5 shows that this functional annihilates every u_j , since $\chi(D)N = 0$ for the original upper exponential space. If its image is zero, $F = N/E_A$ solves $\chi'(D)F = 0$ near zero. Every analytic solution of this constant-coefficient equation is in $\mathcal{E}_{k-8}$: its first q' derivatives determine all later derivatives by the monic equation, and the q' independent displayed exponentials realize arbitrary initial values by the invertible Vandermonde. This proves $N \in E_A \mathcal{E}_{k-8}$. The reverse kernel inclusion follows by applying $\chi'(D)$ to those exponentials. The image dimension is consequently $q - v - q' = \mathfrak{m}$, equal to the full annihilator dimension proved after RDS3, so the map is onto. This also proves that every formal annihilating series converges near zero.

The transpose of the injection in RDS7 is the exact primary conductor C ; this specifies the direction of the dual maps. Thus RDS8 relates the original polynomial relation quotient to the full original exponential module by an explicit quotient and differential map, without a new polynomial ideal.

48.2 Every possible singularity and its original multiplicity

The nonzero entire function E_A has isolated zeros. For $N \in \mathcal{N}_v$, the formula in RDS8 extends its series meromorphically to the whole z plane. Every possible pole lies at a zero of this same fixed original finite symbol; zero itself is removable. If $\zeta \neq 0$ is a zero of order h , put

$$e_\zeta(w) = w^{-h} E_A(\zeta + w), \quad c_{-\nu} = [w^{h-\nu}] \frac{N(\zeta + w)}{e_\zeta(w)} \quad (1 \leq \nu \leq h). \quad (\text{RDS9})$$

Here e_ζ is analytic and nonzero at zero, so these are its actual finite Taylor coefficients. Write $\chi'(X) = \sum_{r=0}^{q'} \chi'_r X^r$, with the coefficients in increasing order in this display. The complete principal part of the function in RDS8 at ζ is

$$\sum_{\nu=1}^h \sum_{r=0}^{q'} \chi'_r (-1)^r (\nu)_r c_{-\nu} (z - \zeta)^{-\nu-r}. \quad (\text{RDS10})$$

Laurent expansion of N/E_A gives its principal part $\sum c_{-\nu} (z - \zeta)^{-\nu}$; differentiating a term r times gives $(-1)^r (\nu)_r (z - \zeta)^{-\nu-r}$. Its holomorphic part remains holomorphic under differentiation, which proves the entire formula. Pole cancellations are retained by the actual coefficients, and the pole order is at most $h + q'$. More precisely, let t be the vanishing order of N at ζ . If $t \geq h$, the quotient

and its image under $\chi'(D)$ are holomorphic. If $t < h$, the leading Laurent coefficient of N/E_A has order $-(h-t)$ and is nonzero. The unique highest-order term after applying the monic $\chi'(D)$ is its q' th derivative: it has order $-(h-t+q')$ and leading coefficient multiplied by $(-1)^{q'}(h-t)_{q'} \neq 0$. Every lower derivative and every subsequent Laurent term has strictly smaller pole order. Thus the exact pole order is $h-t+q'$ in this case. The identically zero numerator gives the zero function. No simplicity of a symbol zero is used.

48.3 An explicit original-data analytic disk and source coefficients

Retain $B, A_{\text{abs}}, C_v, R$ of ROG6 and set

$$R_0 = \frac{1}{B} \min \left\{ 1, \frac{v+1}{2eC_v} \right\} > 0. \quad (\text{RDS11})$$

The exact entire quotient $B_\infty(z) = v!E_A(z)/(\mu_v z^v)$, with value one at zero, obeys $|B_\infty(z)-1| \leq 1/2$ on $|z| \leq R_0$. Indeed its remaining coefficient sum is at most $C_v B|z|e^{B|z|}/(v+1)$ by expanding each original exponential and using $v!/(v+1+j)! \leq 1/((v+1)j!)$. Definition RDS11 gives the asserted bound. This proves that this disk has no nonzero zero of E_A .

Write $N = \sum \nu_{ij} e^{\beta_{ij} z}$. Its first v vanishing Taylor coefficients and the elementary remainder bound for the exponential give

$$|N(z)| \leq \|\nu\|_1 \frac{(R|z|)^v}{v!} e^{R|z|}, \quad \left| \frac{N(z)}{E_A(z)} \right| \leq \frac{2R^v}{|\mu_v|} e^{RR_0} \|\nu\|_1 \quad (|z| \leq R_0). \quad (\text{RDS12})$$

The removable value at zero follows by continuity. Cauchy's derivative formula on this disk gives, for $|z| \leq R_0/2$, the explicit estimate

$$|f_{\mathfrak{F}_A(N)}(z)| \leq C_{k,A} \|\nu\|_1, \quad C_{k,A} = \frac{2R^v}{|\mu_v|} q'! \left(\frac{2}{R_0} \right)^{q'} \exp(RR_0 + R'R_0/2). \quad (\text{RDS13})$$

For verification, the derivative of order r of N/E_A is bounded by $r!(2/R_0)^r$ times the bound in RDS12. The coefficient of $X^{q'-j}$ in $\chi'(X)$ has modulus at most $\binom{q'}{j}(R')^j$. Summing the derivative bounds therefore gives $q'!(2/R_0)^{q'} \sum_{j=0}^{q'} (R'R_0/2)^j/j!$, which proves RDS13. Its logarithm is $O_{\text{actual}}(q \log(q+1))$ at the fixed original period.

Finally, for any polynomial functional the following identity is formal. For $\ell = \mathfrak{F}_A(N)$ its right side is analytic near zero by RDS8 and RDS13, so the identity also proves convergence of its left side. For either unchanged source order $s = 1, k$ it is

$$\sum_{n \geq 0} \ell \left(p_n \left(\frac{S' - c'}{i} \right) \right) \frac{z^n}{n!} = (1+z^2)^{-s/4} e^{ic' \arctan z} f_\ell(-i \arctan z). \quad (\text{RDS14})$$

It follows by applying the complex-linear functional to the original Gamma generating series; the centre and phase are the literal equality $(S' - c')/i = -iS' + ic'$. Let $\rho = \tanh(R_0/4)$. On $|z| = \rho$, $|\arctan z| \leq \text{arctanh } \rho = R_0/4$, so RDS13 and Cauchy's coefficient formula yield

$$\left| \ell \left(\frac{p_n((S' - c')/i)}{\sqrt{h_n}} \right) \right| \leq \sqrt{\frac{n!}{M_s(s/2)_n}} \rho^{-n} (1 - \rho^2)^{-s/4} e^{|c'|R_0/4} C_{k,A} \|\nu\|_1. \quad (\text{RDS15})$$

The square root uses the full original $h_n = M_s n!(s/2)_n$ and $M_s = (2\pi)^{s/2}$. This identity gives the source coefficient realization of the actual relation-annihilating functionals. The weighted relation metric itself remains ROG21, including multiplication by the full χ' . Equations RDS8–15 provide its exact finite-symbol analytic model and quantitative disk; the rank- $O(k)$ residual determinant is not assigned a value by multiplying this scalar coefficient bound by its rank.

49 The fixed-period problem and the result proved here

The current handoff already proves the absolute $O(kq)$ kernel estimate, the conductor frame, its residual observation, and the finite inverse. The new calculation here removes the $O(kq)$ *exterior-inverse allowance* left by AIE29. It replaces that allowance by an explicit $O_{\text{actual}}(q \log q) = o(kq)$ error and transfers the actual four-endpoint determinant to the smaller conductor coefficient space. It does not identify this transfer error with the original kernel volume, and it does not evaluate the latter's leading coefficient.

Retain an actual hypothetical simple quartet

$$\rho = \frac{1}{2} + \delta + i\gamma, \quad 0 < \delta < \frac{1}{2}, \quad \gamma > 2, \quad h(z) = \prod_{\alpha \in \{\rho, \bar{\rho}, 1-\rho, 1-\bar{\rho}\}} (z - \alpha), \quad v_h = 2\xi/h.$$

The original full unit and period observation are

$$U = M_{j_h v_h}, \quad j\Lambda_k = P_k \Pi^{\otimes k} U^{\otimes k} \iota_k, \quad P_k = \frac{1}{5} \sum_{b=0}^4 (C^{-b})^{\otimes k}. \quad (\text{FEX1})$$

Fix one original period and logarithm branch on the proved five-orbit locus, for example any fixed point with $|u| \geq R_*$. Here R_* is the unchanged RPC/AMT radius. Its formulas and the proof of the rank are in the supplied complete source; no new rank premise is imposed. Throughout the principal application,

$$\begin{aligned} k &= 4l + 1 \geq 9, \quad q = (k+1)^2, \quad c = k/2, \quad k' = k - 8, \quad c' = c - 4, \\ q' &= (k-7)^2, \quad a = q - q' = 16k - 48, \quad r = 8k - 16, \quad j_k = 8k - 32, \\ \beta_{ab}^{(k)} &= c + (2a - k)\delta + i(2b - k)\gamma, \quad \chi = \prod_{a,b=0}^k (S - \beta_{ab}^{(k)}), \quad \psi = \chi_{k'}. \end{aligned} \quad (\text{FEX2})$$

The letter a without indices denotes the degree difference, not an amplitude. Let $K = \ker \Lambda_k \subset E = \mathbb{C}[S]/(\chi)$, using its fixed original coefficient inclusion. The actual ranks satisfy $a = r + j_k$. Every class below has its unique representative of degree less than q whenever a polynomial representative is required.

Retain the complete original conductor biform and its coefficients

$$A = \sum_{r_0, s_0=0}^8 a_{r_0 s_0} X_1^{r_0} X_0^{8-r_0} Y_1^{s_0} Y_0^{8-s_0}.$$

Write $\nu = (r_0, s_0)$ for an index in this sum. Define

$$\begin{aligned} b_\nu &= 4 + (2r_0 - 8)\delta + i(2s_0 - 8)\gamma, \quad \mu_j = \sum_{\nu} a_\nu b_\nu^j, \quad v = \min\{j : \mu_j \neq 0\} \leq 80, \\ (\mathcal{T}_{AP})(S') &= \sum_{\nu} a_\nu p(S' + b_\nu), \quad \mathcal{C}_A = \frac{v!}{\mu_v} \mathcal{T}_A, \\ B_q(X) &= \sum_{n=0}^{q-1-v} \frac{v! \mu_{v+n}}{(v+n)! \mu_v} X^n. \end{aligned} \quad (\text{FEX3})$$

The coefficients are those of the actual $\mathbf{H} = F^{-1} \Pi U \mathcal{E}_{\text{prim}}$, fixed as k changes. The original distinct-node Vandermonde proves the stated finite v and $\mu_v \neq 0$. In particular, $B_q(0) = 1$. The scalar in $\mathcal{C}_A$ is displayed explicitly; it is not a renormalization of an arithmetic source or observation.

Only the two original Gamma sources are used:

$$\begin{aligned} s &\in \{1, k\}, \quad b = s/2, \quad M_s = (2\pi)^{s/2}, \quad dm_{0,s}(y) = \frac{M_s 2^b}{4\pi \Gamma(b)} |\Gamma(b/2 + iy/2)|^2 dy, \\ h_n &= M_s n! (b)_n, \quad w_n = \frac{n!}{(b)_n}, \quad \|p\|_{s, c_0}^2 = \int |p(c_0 + iy)|^2 dm_{0,s}(y). \end{aligned} \quad (\text{FEX4})$$

In particular $s = k'$ is never inserted as a source order. The polynomial ψ in FEX2 will enter as a literal multiplication factor in these unchanged measures.

The main conclusion will be the finite interval FEX35 and the consequence

$$\boxed{F_K^{(s)} = \widehat{F}_k^{(s)} + O_{H,\delta,\gamma}(q \log q), \quad s = 1, k.} \quad (\text{FEX5})$$

Here $\widehat{F}$ is defined in FEX29–30 using the actual residual observation and all original transformed relation columns. The error is proved, not postulated. At the end, the unchanged arithmetic comparison transfers the same leading-limit question to F_K^{ar} .

50 A full-flag determinant bound

Let $T : (V, g) \rightarrow (V', g')$ be an isomorphism of n -dimensional complex Hilbert spaces. Write $\|T\| \leq M$ and let $J > 0$ be its squared whole-space Jacobian. For $R \subset W \subset V$, set $t = \dim W - \dim R$. Give W/R and TW/TR their actual Hilbert quotient norms. For any coefficient basis of W/R , let D_Q be the ratio of its target and source Gram determinants under the induced map. Then

$$\boxed{JM^{-2(n-t)} \leq D_Q \leq M^{2t}.} \quad (\text{FEX6})$$

All empty factors have determinant one. The upper bound follows by applying T to each lift and taking the infimum over the same fibre. For the lower bound, use the flag $0 \subset R \subset W \subset V$. Choose any basis adapted to it, on both sides using its transported basis. Successive block elimination of the complete positive Gram gives

$$J = D_R D_Q D_{V/W}. \quad (\text{FEX7})$$

Each block elimination uses its full Schur complement. The other two induced maps also have norm at most M , by restriction or minimization, and their dimensions total $n - t$. Thus their squared determinant product is at most $M^{2(n-t)}$, which proves FEX6. This proves the assertion without asserting orthogonality of a fixed coefficient section.

An equivalent inverse form is useful. If T is an endomorphism of a single Hilbert space and $|\det T| = 1$, then, for every full-rank n by t coefficient matrix X ,

$$\boxed{M^{-2t} \leq \frac{\det(X^* T^{-*} T^{-1} X)}{\det(X^* X)} \leq M^{2(n-t)}.} \quad (\text{FEX8})$$

For example, this follows by taking $W = T^{-1} \text{im } X$, $R = 0$ in FEX6 and inverting the determinant ratio. It also follows by taking the product of the largest t inverse singular values and cancelling against the full product. The statement is a bound: $\det T = 1$ does *not* say that a restricted Gram determinant is unchanged. For $T = \begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix}$, restriction to the second coordinate has squared volume 17.

51 Polynomial forward control replaces the exponential exterior loss

For completeness, the derivative bound used here follows directly from the original polynomial norms. The recurrence and generating identity are

$$\begin{aligned} p_0 &= 1, & p_1 &= y, & p_{n+1} &= yp_n - n(b + n - 1)p_{n-1}, \\ \sum_{n \geq 0} \frac{p_n(y)}{n!} z^n &= (1 + z^2)^{-b/2} e^{y \arctan z}, & \|p_n\|^2 &= h_n. \end{aligned} \quad (\text{FEX9})$$

They are the supplied original Gamma identities. Differentiation of the finite coefficient at zero, followed by the full normalization, gives

$$\partial_y \varphi_n = \sum_{\substack{1 \leq j \leq n \\ j \text{ odd}}} \frac{(-1)^{(j-1)/2}}{j} \sqrt{\frac{w_n}{w_{n-j}}} \varphi_{n-j}, \quad \varphi_n = p_n / \sqrt{h_n}. \quad (\text{FEX10})$$

The mass M_s cancels from this ratio, not from the underlying norms. Let $H_D = \sum_{j=1}^D 1/j$ and $H_0 = 0$. For $n > m \geq 1$, $b \geq 1/2$ gives $w_n/w_m \leq \sqrt{n/m}$ and $w_n \leq 2\sqrt{n}$. Indeed square each factor $j/(j-1/2)$ and use $j(j-1) \leq (j-1/2)^2$; the product telescopes. Apply the weighted Schur estimate to the absolute matrix in FEX10 with weights $w_n^{-1/2}$. Its row sum is at most H_D ; its column sum at n is at most $H_n + 4$. The excess for $1 \leq m < n$ is bounded by

$$\frac{\sqrt{n/m} - 1}{n - m} = \frac{1}{\sqrt{m}(\sqrt{n} + \sqrt{m})} \leq \frac{1}{\sqrt{nm}},$$

whose sum is at most two; the $m = 0$ term is at most $2/\sqrt{n} \leq 2$. Enlarging the odd-index sums is legitimate because all terms are nonnegative. Cauchy–Schwarz in each weighted row and then summing the column estimate proves

$$\|\partial_y\|_{\mathcal{P}_D} \leq \sqrt{H_D(H_D + 4)} \leq H_D + 2, \quad \|f(y + z)\| \leq e^{|z|(H_D+2)} \|f\|. \quad (\text{FEX11})$$

At $D = 0$ the derivative is zero. Translation is the exact finite Taylor exponential. The coordinate substitution $S = c_0 + iy$ gives $\partial_S = -i\partial_y$ with the same norm.

Retain every actual coefficient in

$$A_*^{\text{abs}} = \sum_{\nu} |a_{\nu}|, \quad B_*^{\text{sh}} = \max_{\nu} |b_{\nu}|, \quad C_v = \frac{A_*^{\text{abs}}(B_*^{\text{sh}})^v}{|\mu_v|} \geq 1, \quad P_D = C_v e^{B_*^{\text{sh}}(H_D+2)}. \quad (\text{FEX12})$$

Since $|\mu_{v+n}| \leq A_*^{\text{abs}}(B_*^{\text{sh}})^{v+n}$ and $v!/(v+n)! \leq 1/n!$, the triangle inequality in FEX10–11 gives

$$\|B_q(\partial_S)\|_{s,c_0;\mathcal{P}_D} \leq P_D. \quad (\text{FEX13})$$

This bound holds for the literal fixed polynomial B_q on any explicitly specified finite degree space. On $D \leq q-1$ it is the original AIE operator, whose matrix is triangular with diagonal one. FEX8 therefore proves its new exterior estimate

$$\boxed{P_D^{-2t} \leq \frac{\det(X^*(B_q^{-1})^* B_q^{-1} X)}{\det(X^* X)} \leq \min\{N_{D,s}^{2t}, P_D^{2(D+1-t)}\}.} \quad (\text{FEX14})$$

Here X is any original t -plane in $\mathcal{P}_D$ and $N_{D,s}$ is the unchanged AIE10 inverse bound. The new bound requires neither a new zero-free disk nor a lower bound on an unevaluated restricted minor. For $D = q-1$, $t = r$, its upper logarithm is at most

$$2(q-r)\{\log C_v + B_*^{\text{sh}}(H_{q-1} + 2)\} = O_{H,\delta,\gamma}(q \log q) = o(kq). \quad (\text{FEX15})$$

Thus the factor rD in the old exterior allowance is unnecessary. The forward norm, together with the full Jacobian, controls the product; raising the worst inverse singular value to the r th power loses that information.

There is a direct high-source extension of this statement, distinguished from changing AIE's cutoff. For a fixed k use the same polynomial $B_q(\partial_S)$ on each $\mathcal{P}_N$. Map the exact flag

$$\chi \mathcal{P}_{N-q} \subset I_K \mathbb{C}^r + \chi \mathcal{P}_{N-q}$$

to its literal image under B_q . Its relation space is $B_q(\partial_S)(\chi \mathcal{P}_{N-q})$, not the original ideal by assumption. Its whole Jacobian is one, and FEX13 is valid on this newly specified domain. If $\tilde{F}_K$

is the signed kernel determinant using the native source norm on these transported flags, FEX6 gives

$$-2(2q+1)\log P_{2q} \leq \tilde{F}_K - F_K \leq 2(4q+1)\log P_{2q}. \quad (\text{FEX16})$$

Indeed at endpoint N its logarithmic Gram change belongs to $[-2(N+1-r)\log P_N, 2r\log P_N]$. Insert the four original signs and use $P_N \leq P_{2q}$. This is an exact extension and a signed estimate; no commutation with the original remainder map has been assumed.

52 The literal conductor at every source endpoint

The previous high-source extension is useful, but the actual conductor provides a more informative reduction. On $\mathcal{P}_N$, $N \geq q-1$, its exact Taylor identity is

$$\begin{aligned} \mathcal{C}_A &= \partial_S^v B^{[N]}(\partial_S), & B^{[N]}(X) &= \sum_{n=0}^{N-v} \frac{v!\mu_{v+n}}{(v+n)!\mu_v} X^n, \\ B^{[q-1]} &= B_q, & L &:= \ker(\mathcal{C}_A|_{\mathcal{P}_N}) = \mathcal{P}_{v-1}, & \mathcal{C}_A : \mathcal{P}_N/L &\xrightarrow{\sim} \mathcal{P}_{N-v}. \end{aligned} \quad (\text{FEX17})$$

For $v=0$, $L=0$. These assertions follow either from the leading nonzero derivative or from the commuting invertible triangular factor $B^{[N]}$. Thus the operator on the high endpoints is defined by the original translations themselves. It is not obtained by silently changing the cutoff in the definition of B_q .

The domain has its original quotient Gamma norm at centre c ; the codomain has its native Gamma norm at centre c' . Define

$$M_N = \max \left\{ 1, \frac{v!}{|\mu_v|} \sum_{\nu} |a_{\nu}| e^{[b_{\nu}-4](H_N+2)} \right\}, \quad m_N = \log M_N. \quad (\text{FEX18})$$

The difference of centres is essential: $S' + b_{\nu} = c + iy + (b_{\nu} - 4)$. FEX11 proves $\|\mathcal{C}_A p\|_{s,c'} \leq M_N \|p\|_{s,c}$. Taking the infimum over the same L -coset gives the same bound for the induced isomorphism in FEX17.

Its squared whole-space Jacobian is explicitly

$$\boxed{\mathcal{J}_{N,s} = \prod_{d=v}^N \frac{w_d}{w_{d-v}} = \frac{\prod_{d=N-v+1}^N w_d}{\prod_{d=0}^{v-1} w_d}.} \quad (\text{FEX19})$$

Products are one when $v=0$. To prove the formula, take the orthonormal quotient basis represented by $\varphi_v((S-c)/i), \dots, \varphi_N((S-c)/i)$, with the normalization in FEX10, and the corresponding native basis at c' . The triangular diagonal of the map is $i^{-v}(d)_v \sqrt{h_{d-v}/h_d}$, where $(d)_v = d!/(d-v)!$ is a falling factorial in this formula. Its squared modulus is w_d/w_{d-v} . The source mass is present in both h factors before cancellation; translation of the centre affects only strictly lower coefficients. Telescoping proves the second equality, including overlapping product ranges. For the unnormalized $\mathcal{T}_A$ the squared Jacobian is $|\mu_v/v!|^{2(N+1-v)} \mathcal{J}_{N,s}$.

Put $\zeta_{N,s} = \log \mathcal{J}_{N,s}$. The following finite bounds will suffice:

$$\begin{aligned} 0 &\leq \zeta_{N,1} \leq v \log(2\sqrt{N}), & s &= 1, \\ 0 &\leq -\zeta_{N,s} \leq v(b-1)H_N, & s &= k \geq 9. \end{aligned} \quad (\text{FEX20})$$

For the first use monotonicity of w_n and $1 \leq w_n \leq 2\sqrt{n}$. For the second, w_n decreases and $-\log w_n = \sum_{j=1}^n \log(1 + (b-1)/j) \leq (b-1)H_n$; then use the second product formula in FEX19. These proofs also show directly that the Jacobian error at the tensor source is $O(vk \log q)$, rather than an uncontrolled determinant of dimension q .

The old explicit AIE inverse is still compatible with this estimate. For example, let R_A be that right inverse for the unnormalized $\mathcal{T}_A : \mathcal{P}_D \rightarrow \mathcal{P}_{D-v}$, and let $R_{\min}$ be its actual minimum-norm

right inverse. Their difference takes values in L and is orthogonal to $R_{\min}$. On a t -plane X in the target,

$$\det((R_A X)^* R_A X) \leq \det((R_{\min} X)^* R_{\min} X)(1 + M_{\text{raw}}^2 C_{A,D,s}^2)^{\min(t,v)}. \quad (\text{FEX21})$$

Here $\|\mathcal{T}_A\| \leq M_{\text{raw}}$ and $\|R_A\| \leq C_{A,D,s}$ are the original finite forward and AIE22 bounds. Indeed the excess Gram has rank at most v , and $R_{\min}^* R_{\min} \succeq M_{\text{raw}}^{-2} I$. The operator $R_A - R_{\min}$ is an orthogonal projection of R_A , so its norm is at most $C_{A,D,s}$. Apply the matrix determinant lemma, or congruence with the minimum Gram, to obtain FEX21. FEX6 bounds the minimum inverse determinant by $J_{\text{raw}}^{-1} M_{\text{raw}}^{2(D+1-v-t)} \det(X^* X)$. Only the fixed rank v , not t , multiplies the exponential allowance for the raw low-degree correction. The main reduction below avoids that raw correction by retaining its exact two quotient metrics instead.

53 The residual observation and the smaller coefficient space

AKJ12–25 identifies the original conductor preimage as

$$V_A = \{p \in \mathcal{P}_{q-1} : \psi \mid \mathcal{T}_A p\}. \quad (\text{FEX22})$$

Its complete degree-adapted basis is $1, S, \dots, S^{v-1}$ followed by

$$P_h = \frac{(q' + h + v)!}{(q' + h)!} B_q(\partial_S)^{-1} \mathcal{J}_v(\psi(S) S^h), \quad 0 \leq h < a - v, \quad \mathcal{J}_v S^d = \frac{d!}{(d + v)!} S^{d+v}. \quad (\text{FEX23})$$

The variable in this inverse coefficient expression is identified with S' after applying the translation map. Each P_h is monic of degree $q' + v + h$. Its normalized conductor image is exactly

$$\mathcal{C}_A P_h = d_h \psi(S') S'^h, \quad d_h = \frac{(q' + h + v)!}{(q' + h)!}. \quad (\text{FEX24})$$

The full original residual matrix in this basis is $\Xi = (\Xi_{\text{lo}}, \Xi_{\text{hi}})$, of size j_k by a and rank j_k . Its kernel maps isomorphically onto the original K . These are the supplied actual coefficient maps, not independent matrix parameters. Put

$$r_{\text{lo}} = \text{rank } \Xi_{\text{lo}}, \quad t_0 = v - r_{\text{lo}} = \dim(K \cap L), \quad t = r - t_0. \quad (\text{FEX25})$$

Choose a quotient coordinate map Q_{lo} with kernel exactly $\text{im } \Xi_{\text{lo}}$, and a full-column-rank basis W_k of

$$\ker(Q_{\text{lo}} \Xi_{\text{hi}}) \subset \mathbb{C}^{a-v}. \quad (\text{FEX26})$$

The residual map has rank $j_k - r_{\text{lo}}$ because the full matrix has rank j_k . Hence the space in FEX26 has dimension t . These choices are made using actual nonzero minors and then kept fixed as the source endpoint changes. No generic minor or generic rank is substituted. Define the polynomial column matrix

$$Z_k = \text{diag}(d_h) W_k \quad \text{in the frame } (1, S', \dots, S'^{a-v-1}). \quad (\text{FEX27})$$

Then $\mathcal{C}_A K = \psi \text{ im } Z_k$. In fact, for a high coordinate z in FEX26 the equation $\Xi_{\text{lo}} \ell + \Xi_{\text{hi}} z = 0$ has a solution. Its different solutions differ by $K \cap L$; FEX24 kills that ambiguity and gives precisely ψZ_k . Thus this is the actual image, including the original unit and period dependence in Ξ and the conductor coefficients. The inverse finite-symbol factor no longer appears in the columns Z_k .

For every $j \geq 0$ define the exact transformed relation polynomial

$$R_j^A(S') = \frac{\mathcal{C}_A(\chi S^j)(S')}{\psi(S')} = \frac{v!}{\mu_v} \sum_{\nu} a_{\nu} \frac{\chi(S' + b_{\nu})}{\psi(S')} (S' + b_{\nu})^j. \quad (\text{FEX28})$$

Every displayed quotient is a polynomial. Indeed, if S' is the (a', b') inner-grid root, then $S' + b_{r_0 s_0}$ is the original root with indices $(a' + r_0, b' + s_0)$. All these indices are between 0 and k . The degree of R_j^A is $a - v + j$ and its leading coefficient is $(q + j)_v$; this follows from the first nonzero derivative in FEX17. It is nonzero, and these distinct degrees prove linear independence of the relation columns. In particular the high-source terms have not been replaced by an unjustified continuation of the low-degree symbol.

At endpoint N put $m_N^{\text{pol}} = N - v - q'$ and $d_N^{\text{rel}} = N + 1 - q$. Form the complete polynomial Gram $H_{N,s}^\psi$ of $(1, S', \dots, S'^{m_N^{\text{pol}}})$ in the measure $|\psi(c' + iy)|^2 dm_{0,s}(y)$ at $S' = c' + iy$. Zero-extend Z_k to this frame and let R_N^A have columns $R_0^A, \dots, R_{d_N^{\text{rel}}-1}^A$. Define

$$\begin{aligned}\widehat{H}_{K,N}^{(s)} &= Z_k^* H_{N,s}^\psi Z_k \\ &\quad - Z_k^* H_{N,s}^\psi R_N^A ((R_N^A)^* H_{N,s}^\psi R_N^A)^{-1} (R_N^A)^* H_{N,s}^\psi Z_k, \\ \widehat{F}_k^{(s)} &= \sum_{i=0}^3 \epsilon_i \log \det \widehat{H}_{K,N_i}^{(s)}, \\ (N_0, N_1, N_2, N_3) &= (q-1, q, 2q-1, 2q), \\ (\epsilon_0, \epsilon_1, \epsilon_2, \epsilon_3) &= (1, 1, -1, -1).\end{aligned}\tag{FEX29}$$

The empty relation inverse is absent at $N = q - 1$. The four relation ranks are still exactly $0, 1, q, q + 1$. They have not been truncated to the smaller kernel dimension. Positivity follows because a nonzero linear combination of Z_k has degree at most $a - v - 1$, whereas every nonzero combination of the R_j^A has degree at least $a - v$. The complete Gram is positive in the original positive Gamma density. Minimization over an increasing relation span proves

$$\widehat{H}_{K,q-1}^{(s)} \succeq \widehat{H}_{K,q}^{(s)} \succeq \dots, \quad \widehat{F}_k^{(s)} \geq 0.\tag{FEX30}$$

A change of the fixed basis W_k gives the same four-endpoint value; its four identical Gram factors cancel with the original signs.

54 A finite error for the actual compressed determinant

Let $K^+ = K + L \subset E$. This is only an auxiliary space for an exact metric calculation, not an enlargement of the observation kernel. All quotient and restriction metrics below are induced from the same original $G_N^{(s)}$ and their indicated fixed coefficient frames. Let $F_{K^+/L}$ and $F_{K^+/K}$ denote their four-endpoint volumes. The two block Gram factorizations of K^+ give

$$\boxed{F_K = F_{K^+/L} + F_L - F_{K^+/K}}.\tag{FEX31}$$

Before summing there are fixed basis Jacobians; they cancel only after all four signs. Both factorizations use the actual Schur complements. The two subquotient dimensions are $\dim L = v$ and $\dim(K^+/K) = v - t_0$. The existing full AKS covariance inequality passes by restriction and minimization to every fixed subquotient. Thus, with $\mathcal{L}_s = \mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}$ from AKS,

$$0 \leq F_L \leq v \mathcal{L}_s, \quad 0 \leq F_{K^+/K} \leq (v - t_0) \mathcal{L}_s.\tag{FEX32}$$

No angle condition between K and L is used.

For a fully explicit existing allowance one may use the AKS Newton bound in place of a sharper available $\mathcal{L}_s$. Write

$$\begin{aligned}R &= k \sqrt{\delta^2 + \gamma^2}, \quad t_q = \pi/2 - 1/q, \quad a_q = 2 \log 3 / t_q, \quad s_q = (a_q^q - 1) / (a_q - 1), \\ b_{s,q} &= [(9/25) \sin(1/q)]^{-s/2} \exp(2R(\log 3 + t_q/2)), \\ \mathcal{L}_{s,j} &= \log \left(1 + b_{s,q} s_q^2 \sum_{d=q}^{q+j-1} \frac{d!}{(s/2)_d} (25/16)^d \right).\end{aligned}\tag{FEX33}$$

These are the original complete coefficient bounds with both masses retained before norm-ratio cancellation. They satisfy $\mathcal{L}_s = O_{\delta,\gamma}(q)$ for $s = 1, k$ along this family. Indeed $\log s_q = O(q)$, the last geometric-weighted sum has logarithm $O(q + \log q)$ because $d!/(s/2)_d \leq 2\sqrt{d}$, and $\log b_{s,q} = O(k \log q + R) = o(q) + O(q)$.

Apply FEX6 to the source $\mathcal{P}_N/L$ and its conductor image. The relation subspace is $(\chi\mathcal{P}_{N-q} + L)/L$. The larger flag term is generated also by K , giving quotient dimension t . By FEX28, its image is the span of ψR_N^A and ψZ_k . Multiplication by ψ is an exact isometry from the indicated weighted coefficient space onto this polynomial image. The source quotient metric is exactly the metric of K^+/L , since successive minimization first by $\chi\mathcal{P}_{N-q}$ and then by L is minimization over their sum. Thus, in the fixed transported frames, put

$$\begin{aligned} e_N &= \log \det \widehat{H}_{K,N}^{(s)} - \log \det H_{K^+/L,N}^{(s)}, & n_N &= N + 1 - v, \\ e_{N,s}^- &= \zeta_{N,s} - 2(n_N - t)m_N, & e_{N,s}^+ &= 2tm_N, & e_{N,s}^- &\leq e_N \leq e_{N,s}^+, \\ D_s^- &= e_{q-1,s}^- + e_{q,s}^- - e_{2q-1,s}^+ - e_{2q,s}^+, & D_s^+ &= e_{q-1,s}^+ + e_{q,s}^+ - e_{2q-1,s}^- - e_{2q,s}^-. \end{aligned} \quad (\text{FEX34})$$

The signs on $e_{N,s}^\pm$ denote new finite scalar bounds; they do not redefine an original source scalar or covariance. All endpoint quantities are computed from FEX18–20 and the actual finite ranks in FEX25. Original frame factors have already been accounted for by transported quotient frames. Combining the four inequalities and FEX31–32 proves the central finite result:

$$\boxed{\widehat{F}_k^{(s)} - D_s^+ - (v - t_0)\mathcal{L}_s \leq F_K^{(s)} \leq \widehat{F}_k^{(s)} - D_s^- + v\mathcal{L}_s.} \quad (\text{FEX35})$$

This is a bound for the actual kernel determinant, not for a different ambient determinant. Every residual coefficient and relation cross term remains in FEX29.

For a simpler explicit error let

$$J_{\max,s} = \max_{i=0}^3 |\zeta_{N_i,s}|, \quad m_{\max} = m_{2q}, \quad \mathcal{E}_{k,s} = v\mathcal{L}_s + 2J_{\max,s} + 2(4q + 1)m_{\max}. \quad (\text{FEX36})$$

Then

$$\boxed{|F_K^{(s)} - \widehat{F}_k^{(s)}| \leq \mathcal{E}_{k,s} = O_{H,\delta,\gamma}(q \log q) = o(kq).} \quad (\text{FEX37})$$

To check the finite factors, the lower signed bound for $\sum \epsilon_i e_{N_i}$ is at least $-2J_{\max,s} - 2(2q + 1 - 2v)m_{\max}$, and the upper bound is at most $2J_{\max,s} + 2(4q + 1 - 2v)m_{\max}$. Enlarge to FEX36 and use FEX32. Also

$$m_{\max} \leq \log \max\{1, v!A_*^{\text{abs}}/|\mu_v|\} + 8\sqrt{\delta^2 + \gamma^2}(H_{2q} + 2). \quad (\text{FEX38})$$

Together with FEX20 and FEX33 this proves the order claim. All constants are fixed at the actual fixed period. Dividing by kq uses $\log q/k \rightarrow 0$. A uniform claim over periods approaching a locus where μ_v vanishes is not made: the displayed inverse coefficient must remain in the finite bound. Uniformity holds on a fixed coefficient stratum with this coefficient ratio bounded and the same v .

It follows in particular that $F_K^{(s)}/(kq)$ and $\widehat{F}_k^{(s)}/(kq)$ have exactly the same limit points. This does *not* assert that either sequence has a unique limit. The old uniform ceiling for F_K remains true, while FEX35–37 constitute a new, fixed-period transfer at strictly smaller error scale.

55 The original arithmetic and signed receiving identities

Let $[L_s^{\text{old}}, U_s^{\text{old}}]$ denote the current MRI4e interval, including its first compressed increment and the minimum of its existing finite upper bounds. The following is an add-only improvement at the same four endpoints:

$$\begin{aligned} L_s^{\text{new}} &= \max\{L_s^{\text{old}}, 0, \widehat{F}_k^{(s)} - D_s^+ - (v - t_0)\mathcal{L}_s\}, \\ U_s^{\text{new}} &= \min\{U_s^{\text{old}}, \widehat{F}_k^{(s)} - D_s^- + v\mathcal{L}_s\}, \\ L_s^{\text{new}} &\leq F_K^{(s)} \leq U_s^{\text{new}}, \quad s = 1, k. \end{aligned} \quad (\text{FEX39})$$

The intersection need not improve every finite instance. The comparison of scales, however, is unconditional at each fixed actual period in the stated domain.

The full original Gamma multiplier is

$$f_l(S) = \prod_{j=0}^{l-1} (S - c - 2j - \tfrac{1}{2}), \quad dm_{0,k} = \beta_k |f_l(c + iy)|^2 d\sigma, \quad \beta_k = \frac{(2\pi)^{k/2}}{\sqrt{2}\Gamma(k/2)}. \quad (\text{FEX40})$$

Its transported full-jet kernel, conditional covariance and mixed blocks are AMT23–30, unchanged. The existing difference interval is $-r\Omega_k^{\text{low}} \leq F_K^{(1)} - F_K^{(k)} \leq r\Omega_k^{\text{high}}$, where

$$\begin{aligned} \Pi_l &= \prod_{j=0}^{l-1} (2j + \tfrac{1}{2})^2, \quad U_{l,N}^{\text{poly}} = \prod_{j=0}^{l-1} [4(N + j + 1)^2 + (2j + \tfrac{1}{2})^2], \\ \Omega_k^{\text{low}} &= \log \frac{U_{l,q-1}^{\text{poly}} U_{l,q}^{\text{poly}}}{\Pi_l^2}, \quad \Omega_k^{\text{high}} = \log \frac{U_{l,2q-1}^{\text{poly}} U_{l,2q}^{\text{poly}}}{\Pi_l^2}. \end{aligned} \quad (\text{FEX41})$$

Consequently one may further intersect the tensor-source interval with $[L_1^{\text{new}} - r\Omega_k^{\text{high}}, U_1^{\text{new}} + r\Omega_k^{\text{low}}]$. The fixed-source interval has the corresponding reverse intersection. No independent error assignment is substituted for their common total.

For the arithmetic measure retain exactly

$$w_h(y) = |v_h(\tfrac{1}{2} + iy)|^2 / (2\pi), \quad dm_{h,k}(y) = w_h^{*k}(y) dy, \quad \delta_k^\sigma = \mathcal{B}_k^{\text{ar}} - \mathcal{B}_k^\sigma. \quad (\text{FEX42})$$

Let $H_N^{\text{ar}}, H_N^\sigma$ be the full coefficient-source Grams. AMT6 retains $(a_k/D_N)H_N^\sigma \preceq H_N^{\text{ar}} \preceq A_k H_N^\sigma$. For $m = 1$ its original constants are $p = 7/2$, $B_h = 50$, $M = 25$, $\alpha = \pi/2$, $D_N = [1 + 4(N + M)^2]^M$, and

$$a_k = \frac{c_h \vartheta_h^{k-3} e^{-\alpha(k-3)} (k-2)^{-B_h}}{C_\Gamma 2^{B_h/2}}, \quad A_k = \frac{C_h^{\text{tilt}}}{c_\Gamma} k^{p+1} M_\alpha^{k-1}. \quad (\text{FEX43})$$

Here $\vartheta_h = \int_{-1}^1 w_h$ and $M_\alpha = \int e^{\alpha y} w_h(y) dy$ are the original integrals, and the named positive envelope constants are the unchanged TW/BT constants. In particular none is inferred from a diagnostic packet.

Keep the AMT exact deficits

$$\begin{aligned} L_N^{\text{ar}} &= \log(A_k D_N / a_k), \quad d_{E,N} = q \log A_k - \log \frac{\det G_N^{\text{ar}}}{\det G_N^\sigma}, \\ c_{X,N} &= \min\{r_X L_N^{\text{ar}}, d_{E,N}\}, \quad C_X^- = c_{X,q-1} + c_{X,q}, \quad C_X^+ = c_{X,2q-1} + c_{X,2q}, \quad X = K, B. \end{aligned} \quad (\text{FEX44})$$

The source-evaluated caps replacing $d_{E,N}$ by $\Xi_{k,N}$ remain available. The complete nested positive deficits and their exact common sum are those of AMT13–18. Using either version of the caps, FEX39 therefore gives precisely the new MRI4f interval

$$\begin{aligned} L_{\text{ar}}^{\text{new}} &= \max\{0, L_1^{\text{new}} - C_K^-, L_1^{\text{new}} + \delta_k^\sigma - C_B^+\}, \\ U_{\text{ar}}^{\text{new}} &= \min\{\mathcal{B}_k^{\text{ar}}, U_1^{\text{new}} + C_K^+, U_1^{\text{new}} + \delta_k^\sigma + C_B^-\}, \\ L_{\text{ar}}^{\text{new}} &\leq F_K^{\text{ar}} \leq U_{\text{ar}}^{\text{new}}. \end{aligned} \quad (\text{FEX45})$$

Subtracting the appropriate kernel endpoints from the unchanged total $\mathcal{B}_k^{(s)}$ or $\mathcal{B}_k^{\text{ar}}$ gives the exact boundary interval. This keeps correlation with the actual signed arithmetic value. Its QRI/QGQ finite intervals, including the simple-multiplicity nonzero radius, are not altered by this calculation.

The BRI6d identity remains, with its original orientation,

$$\begin{aligned} F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)} &= F_K^{(k)} + \mathcal{H}_k + \delta_k^\sigma, \quad \mathcal{H}_k = \mathcal{B}_k^\sigma - \mathcal{B}_k^0 = -\mathfrak{T}_k, \\ L_k^{\text{new}} + \mathcal{H}_k + \delta_k^\sigma &\leq F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)} \leq U_k^{\text{new}} + \mathcal{H}_k + \delta_k^\sigma. \end{aligned} \quad (\text{FEX46})$$

The already supplied source differences are $o(kq)$, and the complete Gamma return has $\mathcal{H}_k/(kq) \rightarrow -\mathcal{C}_\Gamma/4$ with the unchanged positive equilibrium constant. Consequently the new reduction has the exact leading-scale receiving form

$$\boxed{\frac{F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)}}{kq} = \frac{\widehat{F}_k^{(1)}}{kq} - \frac{\mathcal{C}_\Gamma}{4} + o(1).} \quad (\text{FEX47})$$

In particular $F_K^{\text{ar}}/(kq)$, $F_K^{(1)}/(kq)$, $F_K^{(k)}/(kq)$, and both new $\widehat{F}$ ratios have the same limit points. The leading value of FEX29 is still needed to decide the sign in FEX47. Neither the old ceiling nor the new $o(kq)$ transfer assigns that value.

56 Actions, representatives, and the retained mixed-support maps

The conductor is a coefficient calculation, not a replacement for the arithmetic action or the original observation. Its precise action defect on polynomials is

$$\mathcal{C}_A(Sp) - S'\mathcal{C}_{AP} = \frac{v!}{\mu_v} \sum_{\nu} a_{\nu} b_{\nu} p(S' + b_{\nu}). \quad (\text{FEX48})$$

For the representative of degree less than q , write $\tau(p) = [S^{q-1}]p$. The original action is $A_{\chi}[p] = [Sp - \chi\tau(p)]$. Its transported polynomial defect is the right side of FEX48 minus $\mathcal{C}_A(\chi\tau(p))$. This formula is in the full polynomial target; divisibility by ψ is not asserted for an action image outside the actual conductor preimage. In particular K is not assumed invariant, and the original OKA9 observation-action defect is retained.

Nor is the auxiliary L an invariant subspace of that action when $v > 0$. On its top monomial S^{v-1} , multiplication followed by $\mathcal{C}_A$ gives the nonzero constant $v!$, whereas all its lower monomials give zero. This is the explicit rank-one defect. No quotient Euler action on K^+/L was needed in the metric proof.

One can retain all coefficients while using the calculation. Let $R_{C,N}$ be any of the explicitly specified finite right inverses of $\mathcal{C}_A$ on the full space $\mathcal{P}_{N-v}$, for example $(B^{[N]}(\partial_S))^{-1}\mathcal{J}_v$. Then

$$p \mapsto (p - R_{C,N}\mathcal{C}_{AP}, \mathcal{C}_{AP}) \in L \oplus \mathcal{P}_{N-v}, \quad (\ell, g) \mapsto \ell + R_{C,N}g \quad (\text{FEX49})$$

are inverse coefficient maps. This explicitly retains the low component rather than treating it as zero. The original remainder map, full unit, and literal period observation may be composed with the inverse in FEX49; they are then the same original maps on the same class, not newly assigned maps on a smaller fibre. The volume identity FEX31 restores both low-component quotient metrics exactly.

If an original minimum section is changed by χQ , its original cochain primitive is still the AKP primitive. Namely divide $\chi(s_1 + \dots + s_k)$ successively by the original $h(s_j)$ to obtain $\sum_j h(s_j)Q_j(\mathbf{s})$. Then use the original tensor differential and the primitive

$$\sum_{j=1}^k (-1)^{j-1} Q_j(\mathbf{D}) Q(D_1 + \dots + D_k) (F_h^{\otimes(j-1)} \otimes \phi_* \otimes F_h^{\otimes(k-j)}). \quad (\text{FEX50})$$

Its differential is the literal difference of the original theta representatives, because $h(D)F_h = \Theta\phi_*$. Minus-only legs use the unchanged lift $(0, -\mathcal{F}_{\text{Fourier}}\phi)$ and transported $1-D$ action. This statement applies to an actual relation difference; no element of K or L is newly declared a theta boundary. All support-face insertions and signed coefficient permutations commute with the reconstructed linear maps componentwise. A supported zero stays at its original label and is not identified with the external τ . The additional proper-source Taylor kernels from the mixed-primary lane remain, not being replaced by a conductor kernel.

The linear-algebra and Gamma-polynomial lemmas apply at any finite multiplicity with their actual dimensions. The explicit OCF conductor, its four additional orbit factors, and the rank $8k - 16$ used for FEX22–47 belong only to the stated simple-quartet case. No such rank or frame is assigned to the full-multiplicity OMG case. Its nilpotents, row-residue annihilator, and observed-iterate kernel remain the supplied ones. This continuation makes no Frobenius identification or arithmetic purity assertion.

57 Scope of verification and the next exact quantity

The analytic assertions proved here are FEX6–21 and the exact transfer FEX22–47 from the explicitly named supplied inputs. Finite exact fixtures test the flag identity, its inverse determinant bounds, the source Jacobian, and the full four-endpoint reduction with nonzero cross terms. They do not evaluate actual xi zeros or actual period coefficients. The accompanying execution record distinguishes those computations from the written proofs and from inherited formal certificates.

The next unresolved quantity is the fixed-period determinant FEX29, with the actual Z_k , R_j^A , and weighted original Gamma Gram. Its leading value is *not* established by this paper. It is now separated from the finite-symbol inverse with an explicit error smaller than kq . In particular, repeating an operator inverse bound of exponential size and raising it to r is no longer the remaining task. The complete original high relation ranks, their mixed Gram blocks, and the period position of $\text{im } Z_k$ are the data that still need estimation. No RH contradiction or disproof follows from the result proved here.

Source dependencies

The fixed objects and current ranks are the supplied OCS/OCF/OKM/OKA, AKJ12–25, and RPC/AMT statements in `14_CURRENT_KERNEL_AND_MULTIPLICITY_PROOFS.tex`. The retained inverse is AIE1–29; the original covariance bound is AKS20–32. The signed source transfer is AMT13–49 and the unchanged GEL/QGT/QRI/QGQ provider in `02_FULL_GAMMA_PROOFS(1).tex`. All complete current source files and their exact hashes accompany this continuation. PR32’s residue and finite observation certificates are inherited inputs only; no new Lean execution is claimed here.

58 The fixed-column conductor quotient and the full GIP graph

Fix exactly the original simple-quartet and fixed-period data of FEX1–4, CEP1–6 and GIP1–3. In particular $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $q' = (k - 7)^2$, $\Delta = q - q'$, $0 \leq v \leq 80$, $\chi = \chi_k$, $\psi = \chi_{k-8}$, $c = k/2$, $c' = c - 4$, and $s \in \{1, k\}$. Retain the exact nonzero scalar

$$\alpha_A = \frac{v!}{\mu_v}, \quad \mathcal{C}_A = \alpha_A \mathcal{T}_A. \quad (\text{FGC1})$$

Every polynomial norm is the original Gamma norm at its stated centre, with the full mass $(2\pi)^{s/2}$. The scalar in FGC1 multiplies the specified coefficient map and does not alter the measure.

For each $N \in \{q - 1, q, 2q - 1, 2q\}$ put $t_N = N + 1 - q$ and $b_N = N - v - q'$. Use the exact GIP coefficient space, now with a different name to keep t_N distinct from a kernel dimension:

$$\mathcal{X}_N = \mathbb{C}^v \oplus \mathbb{C}[S']_{\leq b_N}, \quad L_N h = P_{<v} Z_{\chi, N} h, \quad U_N^A h = \frac{\mathcal{T}_A(\chi h)}{\psi} \quad (h \in \mathbb{C}[S]_{\leq N-q}). \quad (\text{FGC2})$$

Here $P_{<v}$ gives the first v original upper orthonormal coordinates, $Z_{\chi, N} h$ is the complete upper coordinate column of χh , and U_N^A is exactly GIP3, equivalently CEP41–42. Thus the relation subspace of GIP is $\{(L_N h, U_N^A h) : \deg h \leq N - q\}$. The top coordinate is retained in every formula below.

Define the explicit coefficient isomorphism and the transformed graph

$$D_{\alpha, N}(\ell, z) = (\ell, \alpha_A z), \quad D_{\alpha, N}^{-1}(\ell, z) = (\ell, \alpha_A^{-1} z), \quad \mathcal{G}_N^\alpha = \{(L_N h, \alpha_A U_N^A h) : \deg h \leq N - q\}. \quad (\text{FGC3})$$

The complete GIP graph quotient is carried by $D_{\alpha, N}$ to $\mathcal{X}_N / \mathcal{G}_N^\alpha$. Its map from the original conductor kernel V_A sends the class of a polynomial p to

$$[p] \mapsto \left[\left(P_{<v} p, \frac{\mathcal{C}_A p}{\psi} \right) \right]. \quad (\text{FGC4})$$

The second coordinate exists exactly for $[p] \in V_A$, by CEP5–6. Changing p by χh adds the displayed member of $\mathcal{G}_N^\alpha$ and therefore does not change its class. The inverse is the GIP inverse E_N^{graph} composed with $D_{\alpha, N}^{-1}$, so FGC4 is an isomorphism. This proves the relation to the full earlier graph at each of its actual cutoffs.

58.1 The precise projection and its complete kernel

Define the lower coefficient quotient

$$\mathcal{Y}_N = \mathbb{C}[S']_{\leq b_N} / \text{im}(\alpha_A U_N^A), \quad \pi_N : \mathcal{X}_N / \mathcal{G}_N^\alpha \rightarrow \mathcal{Y}_N, \quad [(\ell, z)] \mapsto [z]. \quad (\text{FGC5})$$

The map is well defined because its relation images are exactly $\text{im}(\alpha_A U_N^A)$, and it is onto because every lower column has the lift $(0, z)$. If its value is zero, write $z = \alpha_A U_N^A h$ and subtract the full graph vector $(L_N h, \alpha_A U_N^A h)$. The resulting representative is $(\ell - L_N h, 0)$. A vector $(\ell, 0)$ belongs to the graph only if $U_N^A h = 0$. This implies $h = 0$: for nonzero h , the degree of $\mathcal{T}_A(\chi h)$ is $q + \deg h - v$, with nonzero leading coefficient $\text{lc}(h) \mu_v \binom{q + \deg h}{v}$, by the first nonzero moment identity. Therefore $\ell = 0$. We have proved the full exact sequence, including the original low coordinate and every relation component,

$$0 \rightarrow \mathbb{C}^v \xrightarrow{\ell \mapsto [(\ell, 0)]} \mathcal{X}_N / \mathcal{G}_N^\alpha \xrightarrow{\pi_N} \mathcal{Y}_N \rightarrow 0. \quad (\text{FGC6})$$

Under FGC4 the first term is exactly the original subspace $L = \mathbb{C}[S]_{<v} \subset E_k$, because the conductor kills exactly those polynomials and its first coordinate is their original orthonormal coordinate. Thus FGC6 is, on the same original primary space, the quotient sequence

$$0 \rightarrow L \rightarrow V_A \rightarrow V_A / L \rightarrow 0, \quad V_A / L \xrightarrow{\cong} \mathcal{Y}_N. \quad (\text{FGC7})$$

In particular its target has dimension $\Delta - v$ at each original endpoint. The inverse to the last map is completely specified: choose a representative z , solve $\mathcal{C}_A p = \psi z$ by the finite right inverse of FEX17, and take $[p]$ modulo L and the source relations. A different representative changes p by an original χh plus an element of L , so the same target class results.

Now retain $K = \ker \Lambda_k \subset V_A$, put $t_0 = \dim(K \cap L)$, and set $K^+ = K + L$. Restricting FGC7 gives

$$0 \longrightarrow K \cap L \longrightarrow K \longrightarrow \pi_N(K) \longrightarrow 0, \quad \pi_N(K) \simeq K^+/L, \quad \dim \pi_N(K) = r - t_0. \quad (\text{FGC8})$$

The second isomorphism sends $k + L$ to the same image as k . It is onto, and a zero image says $k \in L$, which proves injectivity on K^+/L . No preservation of an Euler or multiplication action is asserted by these coefficient quotient maps.

58.2 The actual fixed columns and the Schur metric

Use the original AKJ basis and residual matrix $\Xi = (\Xi_{\text{lo}}, \Xi_{\text{hi}})$ from FEX23–26. Let W_k be its specified basis of $\ker(Q_{\text{lo}}\Xi_{\text{hi}})$ and $d_h = (q' + h + v)!/(q' + h)!$. The first nonzero moment identity gives $\mathcal{C}_A P_h = d_h \psi S^{th}$ for every original high AKJ column. The low AKJ columns are precisely L and are killed by $\mathcal{C}_A$. Thus the exact image in FGC8 has the fixed, endpoint-independent representative matrix

$$Z_k = \text{diag}(d_h)W_k, \quad \pi_N(K) = (\text{im } Z_k + \text{im}(\alpha_A U_N^A))/\text{im}(\alpha_A U_N^A). \quad (\text{FGC9})$$

Solvability of $\Xi_{\text{lo}}\ell + \Xi_{\text{hi}}z = 0$ is exactly the equation $Q_{\text{lo}}\Xi_{\text{hi}}z = 0$, proving both inclusions in this identity. The graph's lower relation columns are

$$R_j^A = \alpha_A U_N^A(S^j) = \frac{\mathcal{C}_A(\chi S^j)}{\psi}, \quad 0 \leq j < t_N. \quad (\text{FGC10})$$

This is FEX28, and also the lower block of the complete original relation graph in FGC3. The nonzero leading coefficient is $(q + j)_v$ and its degree is $\Delta - v + j$. The columns of Z_k have degree at most $\Delta - v - 1$. Consequently their quotient classes are independent at every cutoff, including the two high relation ranks q and $q + 1$.

Give the lower polynomial space its exact norm $\int |z(c' + iy)|^2 |\psi(c' + iy)|^2 dm_{0,s}(y)$, whose full Gram in the increasing monomials is $H_{N,s}^\psi$. Then the quotient metric of FGC5 on FGC9 is exactly

$$\widehat{H}_{K,N}^{(s)} = Z_k^* H_{N,s}^\psi Z_k - Z_k^* H_{N,s}^\psi R_N^A ((R_N^A)^* H_{N,s}^\psi R_N^A)^{-1} (R_N^A)^* H_{N,s}^\psi Z_k. \quad (\text{FGC11})$$

Indeed minimizing $(Z_k x + R_N^A h)^* H_{N,s}^\psi (Z_k x + R_N^A h)$ over h gives the unique minimizer $h = -((R_N^A)^* H_{N,s}^\psi R_N^A)^{-1} (R_N^A)^* H_{N,s}^\psi Z_k x$. The relation Gram is positive by the independence in FGC10, and substitution proves FGC11 including both mixed factors. At $t_N = 0$ the minimization has no variable and the second term is absent. This proves the FEX29 matrix directly from the full GIP relation graph by the explicit quotient FGC5. It has not identified the transformed relation span with a chosen polynomial ideal.

The original exact metric in these normalized graph coordinates is $D_{\alpha,N}^{-*} H_N^{\text{actual}} D_{\alpha,N}^{-1}$, with H_N^{actual} from GIP5. It is generally different from the native lower metric in FGC11. On the whole normalized graph space the native direct-sum metric is $\text{diag}(I_v, H_{N,s}^\psi)$. Taking its quotient first by $\mathcal{G}_N^\alpha$ and then by the low image is the same minimization as taking the quotient by their sum. In that sum the low coordinate is unrestricted, so its minimum is zero, and the second-coordinate minimization is precisely FGC11. Thus the native-metric projection also matches the stated Schur matrix; the comparison to the original metric still uses FEX34–35 or CEP's full source-map estimates. The scalar α_A has remained in the coordinate map and both metrics throughout.

58.3 Both original low metric terms and their signed identity

For any fixed positive original endpoint metric $G_N^{(s)}$ on E_k , restrict it to K^+ . In fixed coefficient frames the two complete Schur factorizations give

$$\det H_{K^+,N} = J_1 \det H_{L,N} \det H_{K^+/L,N} = J_2 \det H_{K,N} \det H_{K^+/K,N}, \quad (\text{FGC12})$$

where, if A_1, A_2 are the matrices of the two adapted frames in the fixed reference frame of K^+ , then $J_1 = |\det A_1|^{-2}$ and $J_2 = |\det A_2|^{-2}$. They are independent of the endpoint. To prove each equality, complete the chosen subspace frame by fixed quotient lifts, write the full positive Gram in that adapted frame, and eliminate its off-diagonal block. The quotient block is exactly the minimum Gram because solving its first-block linear equation is the same minimization. Its determinant times the first block's determinant is the determinant in the adapted frame. This proves the formula including its fixed Jacobian.

Taking the four original signs $(1, 1, -1, -1)$ cancels these two frame constants, and gives exactly

$$F_K^{(s)} = F_{K^+/L}^{(s)} + F_L^{(s)} - F_{K^+/K}^{(s)}, \quad \dim L = v, \quad \dim(K^+/K) = v - t_0. \quad (\text{FGC13})$$

The original AKS endpoint covariance comparison passes to a restriction by evaluating the quadratic forms on that subspace, and passes to a quotient by taking the infimum on the same fibre. Its scalar two-sided bounds therefore give FEX32 for these actual subquotients. In particular both nonnegative low-volume terms in FGC13 remain; a projection to FGC11 alone does not remove them. FEX34–37 additionally compares $F_{K^+/L}$ with $\widehat{F}_k$ by the original conductor's complete Jacobian, proving the stated $O_{\text{actual}}(q \log q)$ error.

Equations FGC3–13 exhibit exactly how the new fixed-column presentation fits the existing CEP/GIP construction. The existing RUI5–15 and CEP29–39 already prove the $o(kq)$ source/exterior transport. FEX adds an explicit realization on the lower fixed columns Z_k , with the full original transformed relation span and the two original low metric corrections restored. These equations do not turn that span into the CTV outer ideal or remove the actual residual observation.

59 The exact closed equilibrium coefficient

This section retains the potential, measure, and energy of EIQ1–33:

$$\begin{aligned} V_{\alpha,\beta}(x) &= \beta\sqrt{x} - \alpha \log x, & \alpha > 0, \quad \beta > 0, \quad x > 0, \\ \mu_{\alpha,\beta}(dx) &= \rho_{\alpha,\beta}(x) dx, & F(\alpha, \beta) &= \iint \log|x-y| d\mu_{\alpha,\beta}(x) d\mu_{\alpha,\beta}(y) - \int V_{\alpha,\beta} d\mu_{\alpha,\beta}, \\ \mathcal{L}(\alpha, \beta) &= \int \log x d\mu_{\alpha,\beta}(x). \end{aligned} \quad (\text{ECL1})$$

The preceding complete EIQ proof supplies the unique minimizing probability measure, its positive density and exact support $[a, b] = [u^2, v^2]$, and its Euler equality. In particular, no assertion in this section changes the minimization problem or chooses another measure. The endpoints are exactly

$$uK(k) = \frac{\alpha\pi}{\beta}, \quad vE(k) = \frac{(\alpha+2)\pi}{\beta}, \quad k^2 = 1 - \frac{u^2}{v^2}, \quad 0 < u < v. \quad (\text{ECL2})$$

Here K and E have the modulus convention of EIQ2, not the elliptic-parameter convention. Define the original endpoint quantities

$$w = \frac{u+v}{2}, \quad z = uv, \quad c_{\text{cap}} = \frac{v^2 - u^2}{4}, \quad d\nu(x) = \frac{dx}{\pi\sqrt{(b-x)(x-a)}} \quad (a < x < b). \quad (\text{ECL3})$$

The letter z in ECL3 is a positive scalar. A Cauchy-transform variable below is denoted by λ . The measure ν has mass one by $x = (a+b)/2 + (b-a)\cos\theta/2$, $0 \leq \theta \leq \pi$.

59.1 Arcsine identities with their original constants

Put $m = (a+b)/2$, $d = (b-a)/2$, and $\eta = (v-u)/(v+u)$. Then $0 < \eta < 1$ and

$$x = m + d\cos\theta = w^2(1 + 2\eta\cos\theta + \eta^2), \quad z = w^2(1 - \eta^2).$$

The absolutely uniformly convergent series

$$\begin{aligned} \log x &= 2\log w + 2 \sum_{n \geq 1} \frac{(-1)^{n+1}\eta^n}{n} \cos(n\theta), \\ \frac{1}{x} &= \frac{1}{z} \left(1 + 2 \sum_{n \geq 1} (-\eta)^n \cos(n\theta) \right) \end{aligned}$$

follow from the geometric series and the integral of that series. Termwise integration, and orthogonality of the cosines, prove

$$\begin{aligned} \int \log x d\nu(x) &= 2\log w, & \int \frac{d\nu(x)}{x} &= \frac{1}{z}, \\ \int \frac{\log x}{x} d\nu(x) &= \frac{2\log w + 2\log(1 - \eta^2)}{z} = \frac{2\log z - 2\log w}{z}. \end{aligned} \quad (\text{ECL4})$$

In the product of the two Fourier series, the contribution of degree $n \geq 1$ to the constant coefficient is $-2\eta^{2n}/(nz)$, giving the displayed logarithm without an omitted constant.

The logarithmic potential of ν on its entire support is

$$\int_a^b \log|x-t| d\nu(x) = \log c_{\text{cap}}, \quad a \leq t \leq b. \quad (\text{ECL5})$$

Here is a direct proof, including the endpoint values. Write $t = m + d \cos \phi$ and use

$$|m + d \cos \theta - t| = \frac{d}{2} |e^{i\theta} - e^{i\phi}| |e^{i\theta} - e^{-i\phi}|.$$

The integrand depending on $\cos \theta$ has the same mean on $[0, \pi]$ as on $[0, 2\pi]$. For $0 < r < 1$, the power series of $\log |1 - re^{i\theta}|$ has zero circle mean. This mean tends to the mean at $r = 1$: near its only singular angle, $|1 - re^{i\theta}|^2 = (1 - r)^2 + 2r(1 - \cos \theta)$, so for $r \geq 1/2$ the absolute value of its negative logarithm is bounded by $C + |\log |\theta||$, and its positive logarithm is bounded by $\log 2$. These are integrable bounds. A rotation proves that each of the two circle means above is zero, for every ϕ , including $\phi = 0, \pi$. Hence their sum is $\log(d/2) = \log c_{\text{cap}}$, proving ECL5.

All logarithmic double integrals used below are absolutely integrable. In fact the absolute logarithm for two arcsine variables is bounded by a constant plus

$$\left| \log \left| \sin \frac{\theta - \phi}{2} \right| \right| + \left| \log \left| \sin \frac{\theta + \phi}{2} \right| \right|.$$

Each term is integrable on $[0, \pi]^2$; near a zero, use $\sin t \geq 2t/\pi$ for $0 \leq t \leq \pi/2$, and integrate $|\log t|$. Thus the same assertion holds for any two signed measures whose absolute values are bounded by constant multiples of ν . The EIQ density also has this domination, because $\rho(x)/[1/(\pi\sqrt{(b-x)(x-a)})]$ is bounded on $[a, b]$.

Let ℓ have the exact EIQ17 convention

$$V_{\alpha,\beta}(x) - 2 \int \log |x - t| d\mu_{\alpha,\beta}(t) = \ell \quad (a \leq x \leq b).$$

Integrating this equality against ν , using ECL4–5 and EIQ7, gives

$$\boxed{\ell = 2(\alpha + 2) - 2\alpha \log w - 2 \log c_{\text{cap}}.} \quad (\text{ECL6})$$

Indeed $\int \sqrt{x} d\nu(x) = 2vE(k)/\pi$, whose product with β is $2(\alpha + 2)$ by ECL2. Fubini is justified by the absolute integrability just proved and the unit mass of μ . Smoothness of ℓ now follows from the endpoint smoothness in EIQ6.

59.2 Differentiation with the moving endpoints retained

All derivatives in this subsection are with respect to α at fixed β . Put

$$p_\alpha(x) = \frac{h_{\alpha,\beta}(x)}{2\pi}, \quad D(x) = (b - x)(x - a).$$

In this subsection a dot denotes a parameter derivative, so that $\dot{p}(x)$ differentiates the function $p_\alpha(x)$ at fixed x . The function $h_{\alpha,\beta}$ is exactly EIQ10; the subscript on p_α labels the parameter and does not denote differentiation. EIQ10 and EIQ29 prove that p_α and its parameter and spatial derivatives are bounded on a fixed positive compact neighbourhood of the supports when parameters range in a sufficiently small compact neighbourhood.

The derivative of the probability measure, interpreted by integration against every continuously differentiable function on this neighbourhood, is the finite signed measure

$$\sigma(dx) = \left[\dot{p}(x) \sqrt{D(x)} + \frac{p_\alpha(x)}{2\sqrt{D(x)}} \{ \dot{b}(x - a) - \dot{a}(b - x) \} \right] \mathbf{1}_{(a,b)}(x) dx, \quad (\text{ECL7})$$

$$\frac{\partial}{\partial \alpha} \int f d\mu_{\alpha,\beta} = \int f d\sigma, \quad \sigma([a, b]) = 0.$$

This identity includes the moving support. To prove it without postulating endpoint terms, set $\delta = b - a$ and $x = a + \delta y$. The measure in the fixed variable y is

$$\delta^2 p_\alpha(a + \delta y) \sqrt{y(1 - y)} dy.$$

Its differentiated integral against f is

$$\int_0^1 \sqrt{y(1-y)} \left[2\delta\dot{\sigma} p_\alpha f + \delta^2(\dot{p} + p'_\alpha \dot{x})f + \delta^2 p_\alpha f' \dot{x} \right] dy, \quad \dot{x} = \dot{a} + \dot{\delta}y.$$

Replace $\delta f'$ in the last summand by $df(x)/dy$ and integrate by parts. Its boundary term is $[\delta p_\alpha(x)\dot{x}\sqrt{y(1-y)}f(x)]_0^1 = 0$. The remaining terms combine, on returning to x , to $\dot{p}\sqrt{D} + p_\alpha \dot{D}/(2\sqrt{D})$, where $\dot{D} = \dot{b}(x-a) - \dot{a}(b-x)$, exactly ECL7. This integration by parts is justified first on $[\varepsilon, 1-\varepsilon]$ and then by passage to the limit: the boundary expression tends to zero and the differentiated square-root factor is bounded by a constant times $[y(1-y)]^{-1/2}$, an integrable function. The differentiated fixed-interval integral is valid by the uniform bounds above. Taking $f = 1$ proves the zero total mass in ECL7. The displayed density also proves $|\sigma| \leq C\nu$. In particular no endpoint atom has appeared or been discarded.

For fixed $x \in (a, b)$, differentiating the Euler equality gives

$$-\log x - 2 \int \log|x-t| d\sigma(t) = \dot{\ell}. \quad (\text{ECL8})$$

For precision, choose a smooth function of t equal to one near x , with support in a common interior subinterval for nearby parameters. On this subinterval the density and its parameter derivative are uniformly bounded, so differentiation of its logarithmic integral follows by domination by $C|\log|x-t||$. The complementary integrand is a continuously differentiable test function, so ECL7 applies. This proves ECL8 despite the logarithmic singularity and the moving endpoints.

We next identify σ exactly:

$$\boxed{\frac{\partial}{\partial \alpha} \mu_{\alpha, \beta}(dx) = \sigma(dx) = \frac{1}{2} \left(1 - \frac{z}{x}\right) d\nu(x).} \quad (\text{ECL9})$$

The derivative has the test-function meaning proved in ECL7, which in particular applies to the logarithmic and square-root moments because $a > 0$.

Here is an explicit uniqueness argument establishing ECL9. Denote its proposed right side by σ_0 . By ECL4 it has zero total mass, and $|\sigma_0| \leq C\nu$. Retain the branch $R(\lambda) = \sqrt{(\lambda-a)(\lambda-b)}$ with $R(\lambda)/\lambda \rightarrow 1$ at infinity, exactly as in EIQ12. The arcsine transform proved there and partial fractions yield

$$\begin{aligned} \int \frac{d\sigma_0(t)}{\lambda-t} &= \frac{1}{2R(\lambda)} - \frac{z}{2\lambda} \left(\frac{1}{z} + \frac{1}{R(\lambda)} \right) \\ &= \frac{1-z/\lambda}{2R(\lambda)} - \frac{1}{2\lambda}. \end{aligned} \quad (\text{ECL10})$$

There is no pole at zero: $R(0) = -uv = -z$, so the numerator $(\lambda-z)/R(\lambda) - 1$ vanishes there. The integral itself is analytic in a neighbourhood of zero, as is also immediate from $t \geq a > 0$. For $a < x < b$, the real principal value of the arcsine transform is zero, because $R(x+i0)$ is purely imaginary. This limiting principal value also follows directly by subtracting the local value of the smooth interior density in its Cauchy integral. Consequently ECL10 gives

$$\text{PV} \int \frac{d\sigma_0(t)}{x-t} = -\frac{1}{2x}.$$

Differentiating the logarithmic potential on each interior subinterval is valid by that same local subtraction; the endpoint parts are integrable and are a positive distance away. It follows that $-\log x - 2 \int \log|x-t| d\sigma_0(t)$ is constant on (a, b) . Subtract this identity from ECL8. For $\sigma_1 = \sigma - \sigma_0$, the logarithmic potential $\int \log|x-t| d\sigma_1(t)$ is constant on (a, b) . Since σ_1 has zero mass and no endpoint atoms, and $|\sigma_1| \leq C\nu$, absolute integrability proved after ECL5 gives

$$-\iint \log|x-t| d\sigma_1(x) d\sigma_1(t) = 0.$$

The exact zero-mass energy argument EIQ20–22, including its Fourier-uniqueness proof, forces $\sigma_1 = 0$. This establishes the claimed derivative as a measure, with every endpoint contribution in ECL7 retained.

59.3 The moment and energy formulas

Integrate ECL8 against ν . By ECL5 and $\sigma([a, b]) = 0$, the iterated logarithmic term is zero. Therefore

$$\dot{\ell} = -2 \log w. \quad (\text{ECL11})$$

All exchanges again have the absolute integrability established after ECL5; the equality ECL8 holds ν -almost everywhere. Using ECL9 with $f(x) = \log x$, and then ECL4, proves

$$\mathcal{L}_\alpha = \frac{1}{2} \int \left(1 - \frac{z}{x}\right) \log x \, d\nu(x) = 2 \log w - \log z = \log(w^2/z). \quad (\text{ECL12})$$

Write $I = \iint \log |x - t| \, d\mu(x) d\mu(t)$, with the original unscaled logarithmic kernel. The EIQ31 dilation identity is $\beta \int \sqrt{x} \, d\mu = 2(\alpha + 1)$. Integration of the Euler equality against μ gives $\ell = \int V \, d\mu - 2I$. Combining this with $F = I - \int V \, d\mu$ yields the exact identity

$$\ell = -2F - 2(\alpha + 1) + \alpha \mathcal{L}. \quad (\text{ECL13})$$

EIQ30 proves $F_\alpha = \mathcal{L}$. Differentiating ECL13 at fixed β , with the differentiability just proved, gives $\dot{\ell} = -\mathcal{L} - 2 + \alpha \mathcal{L}_\alpha$. Substituting ECL11–12 now proves, without an integration constant or a limiting parameter choice,

$$\mathcal{L} = 2(\alpha + 1) \log w - \alpha \log z - 2, \quad F = -\frac{\ell}{2} - (\alpha + 1) + \frac{\alpha \mathcal{L}}{2}. \quad (\text{ECL14})$$

Equations ECL6, ECL12, and ECL14 therefore evaluate the same Robin constant, logarithmic moment, and variational energy as the original EIQ integrals.

59.4 The original outer coefficient and the two scale conventions

The coefficient of OFV29 is defined by its full original expression

$$\mathfrak{C}_\partial = 4 \log(4/\vartheta) - 10 - 2F(2, \vartheta) + 4\mathcal{L}(2, \vartheta), \quad \vartheta = \pi/2. \quad (\text{ECL15})$$

To express this same number at the original EIQ limiting potential $V_{2,\pi}(x) = \pi\sqrt{x} - 2 \log x$, retain the actual dilation $x \mapsto (\pi/\beta)^2 x$ of EIQ32. Its effect on the interaction, potential integral, and moment is, respectively,

$$\begin{aligned} I(\alpha, \beta) &= I(\alpha, \pi) + 2 \log(\pi/\beta), \\ \int V_{\alpha,\beta} \, d\mu_{\alpha,\beta} &= \int V_{\alpha,\pi} \, d\mu_{\alpha,\pi} - 2\alpha \log(\pi/\beta), \\ \mathcal{L}(\alpha, \beta) &= \mathcal{L}(\alpha, \pi) + 2 \log(\pi/\beta). \end{aligned}$$

Thus $F(\alpha, \beta) = F(\alpha, \pi) + 2(\alpha + 1) \log(\pi/\beta)$, with its stated sign. At $\alpha = 2$ the added coefficient in ECL15 is $(4 - 12 + 8) \log(\pi/\beta) = 0$. It follows exactly that

$$c_\partial := \mathfrak{C}_\partial = 4 \log(4/\pi) - 10 - 2F(2, \pi) + 4\mathcal{L}(2, \pi). \quad (\text{ECL16})$$

For the rest of this subsection $u, v, w, z, c_{\text{cap}}$ are precisely their ECL2–3 values at $(\alpha, \beta) = (2, \pi)$. They therefore satisfy $uK(k) = 2$ and $vE(k) = 4$. Substitution of the established formulas, with their constants, gives

$$c_\partial = 4 \log(4/\pi) + 8 \log w - 4 \log z - 2 \log c_{\text{cap}}. \quad (\text{ECL17})$$

For example, ECL6 gives $\ell = 8 - 4 \log w - 2 \log c_{\text{cap}}$, ECL14 gives $\mathcal{L} = 6 \log w - 2 \log z - 2$, and ECL13 gives $F = -\ell/2 - 3 + \mathcal{L}$. In ECL16 these yield $4 \log(4/\pi) - 4 + \ell + 2\mathcal{L}$, in which the constant terms are exactly $-4 + 8 - 4 = 0$, proving ECL17. Equivalently $F(2, \pi) = I(2, \pi) - 6 + 2\mathcal{L}(2, \pi)$ in ECL16 gives $c_\partial = 4 \log(4/\pi) + 2 - 2I(2, \pi)$, which is precisely the OFV31 energy convention. This section evaluates this coefficient. It does not assign it as the leading coefficient of a separate observation residual or a separate kernel determinant.

59.5 An integer-only outward certificate

The accompanying `certify_closed_equilibrium.py` proves

$$\boxed{21.66851180520 < 16c_\partial < 21.66851180521.} \quad (\text{ECL18})$$

The receipt stores every interval endpoint as an integer divided by $S = 10^{70}$, including the original endpoints and all quantities in ECL17. The script uses no floating-point inputs or numerical quadrature. The following argument proves the correctness of its finite algorithm.

Each stored interval $[A, B]$ denotes $[A/S, B/S]$. Addition and subtraction are exact. For multiplication, the smallest and largest of the four endpoint products are divided by S and rounded respectively down and up. For division by an interval excluding zero, the four ratios of endpoint integers, multiplied by S , are rounded outwards. Their extrema enclose every possible product or quotient because these operations attain their extrema at corners of their rectangular domains. The rounding uses integer floor division and $\lceil n/d \rceil = -\lfloor -n/d \rfloor$ after making $d > 0$. These formulas apply to negative numerators as well. Consequently induction proves containment after every arithmetic operation, without a probabilistic or floating-point error estimate.

Let $h = u/v$ in this paragraph only; this is a scalar ratio and does not replace the density function $h(x)$ of EIQ10. Put $m_e = 1 - h^2$, so $k = \sqrt{m_e}$. Define $b_n = \binom{2n}{n}/4^n$ and $t_n = b_n^2 m_e^n$. Expansion of the binomial series, followed by its uniformly convergent integration for $m_e < 1$, proves

$$\frac{2K(\sqrt{m_e})}{\pi} = \sum_{n=0}^{\infty} t_n, \quad \frac{2E(\sqrt{m_e})}{\pi} = 1 - \sum_{n=1}^{\infty} \frac{t_n}{2n-1}, \quad \frac{t_n}{t_{n-1}} = m_e \frac{(2n-1)^2}{4n^2}. \quad (\text{ECL19})$$

Indeed integration by parts gives $\int_0^{\pi/2} \sin^{2n} \theta d\theta = (\pi/2)b_n$; the two binomial coefficients are b_n and $-b_n/(2n-1)$ for $n \geq 1$. Since $t_{n+1}/t_n \leq m_e$, the positive omitted tails after indices $0, \dots, N-1$ have the rigorous bounds

$$0 \leq \sum_{n=N}^{\infty} t_n \leq \frac{t_N}{1-m_e}, \quad 0 \leq \sum_{n=N}^{\infty} \frac{t_n}{2n-1} \leq \frac{t_N}{(2N-1)(1-m_e)}. \quad (\text{ECL20})$$

The script takes $N = 3200$, applies the recurrence with outward rounding at each step, and adds these tails with their proper signs. They remain valid for interval values of m_e by the interval arithmetic just proved.

The two exact terminating decimals used for the root bracket are

$$\begin{aligned} h_- &= 0.160101670957718355884567453115, \\ h_+ &= 0.160101670957718355884567453116. \end{aligned} \quad (\text{ECL21})$$

Let $K_b = 2K/\pi$, $E_b = 2E/\pi$ as in ECL19. The receipt gives strictly negative upper bound for $2h_-K_b(1-h_-^2) - E_b(1-h_-^2)$ and strictly positive lower bound for the corresponding expression at h_+ . Here the arguments of K_b, E_b denote m_e as defined by their series, not the modulus argument of EIQ2. The common factor $\pi/2 > 0$ in the original equation cancels exactly. EIQ6 proves that $hK(\sqrt{1-h^2})/E(\sqrt{1-h^2})$ increases strictly from zero to one as h increases from zero to one. Thus ECL21 encloses the unique root of the original endpoint equation $hK(\sqrt{1-h^2}) = E(\sqrt{1-h^2})/2$.

The value of π is enclosed independently by

$$\pi = 16 \arctan(1/5) - 4 \arctan(1/239). \quad (\text{ECL22})$$

To prove the identity, if $\theta = \arctan(1/5)$, then $\tan(2\theta) = 5/12$ and $\tan(4\theta) = 120/119$. The subtraction formula gives $\tan(4\theta - \arctan(1/239)) = 1$. This angle lies in $(0, \pi/2)$: $4\theta < 4/5 < 1 < \pi/2$, while $4\theta \geq 4(1/5)/(1+1/25) > 1/239$. The inequality $\pi/2 > 1$ follows from $\arctan 1 = \int_0^1 (1+t^2)^{-1} dt > 1/2$. Consequently the angle is $\pi/4$, proving ECL22. Each arctangent is computed

from its alternating series with 128 terms. The first omitted term bounds the absolute remainder; its sign is positive for this even number of terms. Outward arithmetic and that signed tail therefore enclose the exact π .

For every positive logarithm argument x put $t = (x-1)/(x+1)$. The geometric series, integrated on the segment from zero to t , gives

$$\log x = 2 \sum_{n=0}^{M-1} \frac{t^{2n+1}}{2n+1} + R_M, \quad |R_M| \leq \frac{2|t|^{2M+1}}{(2M+1)(1-|t|^2)}. \quad (\text{ECL23})$$

The script takes $M = 256$. For an interval argument it uses the maximum absolute endpoint of the enclosing t interval in this tail bound and adds a symmetric outward tail. Thus ECL23 encloses the logarithm throughout the argument interval.

The enclosed original endpoint values are computed in order by

$$\begin{aligned} K &= (\pi/2)K_b, & E &= (\pi/2)E_b, & v &= 4/E, & u &= hv, \\ w &= (u+v)/2, & z &= uv, & c_{\text{cap}} &= (v^2 - u^2)/4. \end{aligned}$$

The script checks $0 < u < v$, positive capacity, strict endpoint signs, and that every division interval excludes zero. It applies ECL23 to the four arguments $4/\pi, w, z, c_{\text{cap}}$ and evaluates the full expression ECL17 with its original coefficients. Integer comparison of the final bounds proves the stronger enclosure

$$\begin{aligned} 21.668511805204674126143836401790042003 &< 16c_{\partial} \\ &< 21.668511805204674126143836403323660355. \end{aligned} \quad (\text{ECL24})$$

The decimal endpoints in ECL24 are rounded outwards from the stored integer endpoints. Their comparison with the exact terminating decimals in ECL18 proves ECL18.

The same operations also enclose the endpoint expression

$$\frac{1}{8} \log(w^2/z) - \frac{1}{4} \log 2 \in (-0.080453400358, -0.080453400357). \quad (\text{ECL25})$$

This is an evaluation of this explicit expression. If it is used as an earlier approximation centre, its previously specified error radius is not changed by this calculation. The ordinary and optimized Python executions produce identical JSON receipts. All mathematical checks use explicit exceptions, so disabling Python assertions does not disable any certificate condition. The receipt includes the script hash, integer endpoint signs, elliptic-tail bounds, every resulting interval, and the strict target comparisons.

60 The closed coefficient in the original QGQ return centre

Retain the complete original QGQ1–9 theorem and its full provider 02_FULL_GAMMA_PROOFS.tex, section *The finite original $m = 1$ return bound at the q scale*. Retain its actual Gamma functional W_k , finite centre $\mathcal{V}_{q,l}^c$, constants $\mathcal{C}_\Gamma, A, M, C$, and its original packet

$$m = 1, \quad k = 4l + 1, \quad q = (4l + 2)^2, \quad n = q/2, \quad l \geq 1.$$

The support and endpoint values u, v, w, z in this section are exactly those of ECL2–3 at $(\alpha, \beta) = (2, \pi)$. In particular $uK(k_{\text{ell}}) = 2$, $vE(k_{\text{ell}}) = 4$, and $k_{\text{ell}}^2 = 1 - u^2/v^2$; the subscript distinguishes this elliptic modulus from the original integer packet degree k . No endpoint or original degree is changed by this notation.

QGQ4 defines, for the original EIQ logarithmic moment, $\mathcal{L}_1 = \partial_\alpha \mathcal{L}(2, \pi)$ and $\mathcal{C}_\Gamma^{[q,1]} = \mathcal{L}_1/8 - (\log 2)/4$. The derivative proved in ECL12 is a derivative of precisely this same moment, at fixed original β . Therefore the following is an exact identification of the original centre:

$$\boxed{\begin{aligned} \mathcal{L}_1 &= \log(w^2/z), & \mathcal{C}_\Gamma^{[q,1]} &= \frac{1}{8} \log(w^2/z) - \frac{1}{4} \log 2, \\ -0.080453400358 &< \mathcal{C}_\Gamma^{[q,1]} < -0.080453400357. \end{aligned}} \quad (\text{EQR1})$$

The first two equalities follow by substitution into the literal QGQ4 definition; there is no new centre or change of energy convention. The last line is consequently the strict outward integer enclosure ECL25 for that exact original centre. Its proof and stored integer endpoints are those of ECL19–25.

The original finite-centre theorem QGQ8 now gives

$$\boxed{\frac{\mathcal{V}_{q,l}^c - lq\mathcal{C}_\Gamma}{q} \rightarrow \frac{1}{8} \log(w^2/z) - \frac{1}{4} \log 2 \quad (l \rightarrow \infty).} \quad (\text{EQR2})$$

To check the connection to the original finite calculation explicitly, QGQ7 retains its first term $(2\mathcal{L}_1 - 4 \log 2)l^2$ and all its other original terms. The complete termwise estimates following QGQ7 bound their sum by $O(l)$; in particular its actual remainder $\zeta_{q,l}$ is $O(l)$ by QGQ6. On the literal packet, $l^2/q \rightarrow 1/16$ and $l/q \rightarrow 0$. Thus dividing that same identity by q gives $(2\mathcal{L}_1 - 4 \log 2)/16 = \mathcal{L}_1/8 - (\log 2)/4$. Substitution of EQR1 proves EQR2 with the original finite centre, source, low endpoint, and rank correction intact.

The original error radius remains the following nonzero number:

$$\boxed{\limsup_{l \rightarrow \infty} \left| \frac{W_k - lq\mathcal{C}_\Gamma}{q} - \left(\frac{1}{8} \log(w^2/z) - \frac{1}{4} \log 2 \right) \right| \leq \frac{A}{\sqrt{2}}, \quad A = M/2 + 2C > 0.} \quad (\text{EQR3})$$

Here M and C retain their complete original QGT12 and QGT9 definitions. To verify the transfer, retain the finite QGT22 inequality used in QGQ9:

$$-p_{q,l}^- - \mathcal{U}_{q,l} - E_{q,l} \leq W_k - \mathcal{V}_{q,l}^c \leq p_{q,l}^+ + \mathcal{U}_{q,l} + T_{q,l} + E_{q,l}.$$

QGQ1–2 proves that each of $p_{q,l}^-/q, p_{q,l}^+/q, T_{q,l}/q, E_{q,l}/q$ tends to zero, while $\mathcal{U}_{q,l}/q \rightarrow A/\sqrt{2}$. Hence the difference $(W_k - \mathcal{V}_{q,l}^c)/q$ is bounded below by $-A/\sqrt{2} - o(1)$ and above by $A/\sqrt{2} + o(1)$. Adding the convergent expression in EQR2, and using the triangle inequality, proves EQR3. Both unequal finite errors remain in the displayed original inequality; their passage to the same limiting radius does not replace that finite interval by a new one.

The radius is strictly positive from the original definitions themselves. QGT9 gives $C \geq 0$, and QGT12 gives

$$M = \max_{(\alpha, \beta) \in \mathcal{K}} |\partial_\alpha \mathcal{L}(\alpha, \beta)| \geq \partial_\alpha \mathcal{L}(2, \pi) = \log(w^2/z) > 0,$$

because $(2, \pi) \in \mathcal{K}$ and $w^2/z = 1 + (v - u)^2/(4uv) > 1$ by $0 < u < v$. Consequently $A = M/2 + 2C > 0$. This proves that the retained nonzero radius has not disappeared upon evaluating its centre. The only convergence asserted in EQR2 concerns the original finite centre. Neither EQR1 nor EQR3 asserts a limit for the actual quantity $(W_k - lq\mathcal{C}_\Gamma)/q$.

61 Strict full-window growth of the original residual covariance

This calculation starts from the complete ROQ1–27 construction above. The incoming first-variation handoff states a full-window update and trace–determinant bounds. The following proof derives those statements from the original columns, and adds exact exterior-power formulas for every intermediate coefficient. No unavailable FVR proof is used. Retain $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $q' = (k - 7)^2$, $j = 8k - 32$, the fixed actual period in the five-orbit locus, and $s \in \{1, k\}$. The actual matrices $C : \mathbb{C}^q \rightarrow \mathbb{C}^{q'}$ and $D : \mathbb{C}^q \rightarrow \mathbb{C}^j$ are ROQ1 and ROQ4, with all their coefficients. In particular C is onto and $D|_{\ker C}$ is onto. The original roots, monic Gamma polynomials and their squared norms are

$$\begin{aligned} \omega_{ab} &= (2b - k)\gamma - i(2a - k)\delta, \quad 0 \leq a, b \leq k, \quad \delta > 0, \quad \gamma > 0, \\ p_0 &= 1, \quad p_1 = y, \quad p_{n+1} = yp_n - n(s/2 + n - 1)p_{n-1}, \\ h_n &= (2\pi)^{s/2} n!(s/2)_n, \quad u_n = (p_n(\omega_{ab})/\sqrt{h_n})_{ab}, \quad R_N = \sum_{n=0}^N u_n u_n^*. \end{aligned} \tag{RWC1}$$

The positive original measure dm_s has the orthogonality and norms in RWC1, as proved in the Gamma provider. The order of the q roots is the original ordered primary order; stars below are conjugate transposes. The first q monic polynomials have invertible evaluation matrix, since these q roots are distinct. Consequently $R_N > 0$ for $N \geq q - 1$.

61.1 Every consecutive full block is invertible

The complete real polynomial of the original root set is

$$\begin{aligned} P_k(y) &= \prod_{a=0}^k \prod_{b=0}^k (y - \omega_{ab}) \\ &= \prod_{a=0}^{(k-1)/2} \prod_{b=0}^k ((y - (2b - k)\gamma)^2 + (k - 2a)^2 \delta^2) > 0 \quad (y \in \mathbb{R}). \end{aligned} \tag{RWC2}$$

Indeed the factor indexed by $k - a, b$ is the conjugate of the factor indexed by a, b , and no factor in the second product can vanish on the real line. This proves the identity with its monic constant and all factors retained.

For any integer $N \geq q - 1$ put

$$U_N^+ = [u_{N+1}, \dots, u_{N+q}] \in \text{Mat}_{q \times q}(\mathbb{C}). \quad \det U_N^+ \neq 0. \tag{RWC3}$$

To prove the assertion suppose $U_N^+ c = 0$ and form $f(y) = \sum_{a=1}^q c_a p_{N+a}(y)/\sqrt{h_{N+a}}$. It vanishes at all q distinct roots, so polynomial division gives $f = P_k r$ with $\deg r \leq N$. Orthogonality of every p_{N+a} to every polynomial of degree at most N gives

$$0 = \int_{\mathbb{R}} f(y) \overline{r(y)} dm_s(y) = \int_{\mathbb{R}} P_k(y) |r(y)|^2 dm_s(y).$$

All integrals exist, because the measure has every polynomial moment. Its strictly positive density and RWC2 force $r = 0$. The distinct degrees of the monic p_{N+a} then force every $c_a = 0$. This argument applies at both $N = q - 1$ and $N = q$, and also at every larger integer cutoff; it makes no genericity choice of D .

61.2 The simultaneous conditioned update

Use the complete original matrices

$$\begin{aligned} A_N &= C R_N C^*, \quad W_N = R_N - R_N C^* A_N^{-1} C R_N, \quad \Omega_N = D W_N D^* > 0, \quad Q_{\Xi, N} = \Omega_N^{-1}, \\ \Pi_N &= I_q - R_N C^* A_N^{-1} C, \quad X_N = C U_N^+, \quad Z_N = D \Pi_N U_N^+, \\ B_N &= I_q + X_N^* A_N^{-1} X_N > 0, \quad V_N = Z_N B_N^{-1} Z_N^*. \end{aligned} \tag{RWC4}$$

These use A_N solely for this covariance block, rather than for the conductor biform. Since $A_N > 0$, every displayed inverse exists. The equalities $C\Pi_N = 0$ and $\Pi_N x = x$ for $Cx = 0$ show that Π_N is the exact projection onto $\ker C$ with complementary image $R_N C^* \mathbb{C}^q$. It is generally not the Euclidean orthogonal projection. RWC3 and surjectivity of $D|_{\ker C}$ give

$$\text{im}(\Pi_N U_N^+) = \ker C, \quad \text{rank } Z_N = j, \quad V_N > 0. \quad (\text{RWC5})$$

For the last assertion, Z_N^* is injective and $B_N^{-1} > 0$, so $z^* V_N z > 0$ for each nonzero $z \in \mathbb{C}^j$.

There are exact identities

$$\begin{aligned} W_{N+q} &= W_N + (\Pi_N U_N^+) B_N^{-1} (\Pi_N U_N^+)^*, \\ \Omega_{N+q} &= \Omega_N + V_N, \quad Q_{\Xi, N+q} = (\Omega_N + V_N)^{-1}. \end{aligned} \quad (\text{RWC6})$$

Here is the complete matrix calculation. Temporarily write $R = R_N$, $U = U_N^+$, $A = A_N$, $X = CU$, $B = B_N$ and $T = RC^* A^{-1} X$. Direct multiplication proves

$$(A + XX^*)^{-1} = A^{-1} - A^{-1} X B^{-1} X^* A^{-1}.$$

Indeed $(A + XX^*) A^{-1} X = XB$, so the subtracted term cancels $XX^* A^{-1}$. Since $CR_{N+q} C^* = A + XX^*$, the new subtracted covariance product equals

$$RC^* A^{-1} CR + TU^* + UT^* + U(B - I_q)U^* - [T + U(B - I_q)]B^{-1}[T + U(B - I_q)]^*.$$

Subtract this expression from $R + UU^*$ and then subtract W_N . Writing $T + U(B - I_q) = UB - (U - T)$, expansion leaves exactly $(U - T)B^{-1}(U - T)^*$. But $U - T = \Pi_N U$, proving the first identity in RWC6. Multiplication by D, D^* proves the second, and ROQ16 proves the third. In particular the full $q \times q$ matrix B_N and all its off-diagonal entries remain in the update.

61.3 All exterior coefficients and finite bounds

Let

$$Y_N = \Omega_N^{-1/2} Z_N B_N^{-1/2}, \quad T_N = Y_N Y_N^* > 0, \quad t_N = \text{Tr } T_N, \quad c_N = \det T_N = \frac{\det V_N}{\det \Omega_N} > 0. \quad (\text{RWC7})$$

The square roots are the positive Hermitian square roots; no source mass, coordinate phase or rank has been changed. If $\lambda_1, \dots, \lambda_j > 0$ are the eigenvalues of T_N , then RWC6 gives

$$L_N := \log \frac{\det \Omega_{N+q}}{\det \Omega_N} = \sum_{a=1}^j \log(1 + \lambda_a) = \log \left(\sum_{d=0}^j e_d(T_N) \right), \quad e_d(T_N) = \left\| \bigwedge^d Y_N \right\|_{\text{HS}}^2. \quad (\text{RWC8})$$

The exterior norm is the Hilbert–Schmidt norm in the standard orthonormal wedge bases, and $e_0 = 1$. To verify the last equality, the singular values of Y_N are $\sqrt{\lambda_a}$, and its d th exterior power has squared singular values $\prod_{a \in I} \lambda_a$ for the d -element subsets I . Their sum is the elementary symmetric polynomial e_d . Equivalently Cauchy–Binet writes it as the sum of squared moduli of all $d \times d$ minors of the literal Y_N . This proves every coefficient in the determinant expansion, including all intermediate exterior powers; the sole determinant c_N is its last coefficient.

The following finite inequalities now hold:

$$\max\{\log(1 + t_N), j \log(1 + c_N^{1/j})\} \leq L_N \leq j \log(1 + t_N/j). \quad (\text{RWC9})$$

For the first lower bound, expand $\prod(1 + \lambda_a)$: the constant term is one, its degree-one term is t_N , and all other terms are positive. For the second, the arithmetic–geometric mean inequality on the $\binom{j}{d}$ positive d -fold products gives $e_d \geq \binom{j}{d} c_N^{d/j}$, since each eigenvalue occurs in $\binom{j-1}{d-1}$ products. Summing this inequality for $0 \leq d \leq j$ yields $(1 + c_N^{1/j})^j$. Finally the concavity of $\log(1 + x)$ on $x > 0$ gives the upper bound by averaging the j positive eigenvalues. These arguments also prove useful finite refinements: retaining any chosen initial set $e_0, \dots, e_d$ in RWC8 gives its exact positive partial-sum lower bound; no rank fraction is substituted for an eigenvalue.

61.4 Return to the original four endpoints

Use precisely the ordered cutoffs $(q-1, q, 2q-1, 2q)$, with signs $(1, 1, -1, -1)$. ROQ14 and ROQ16 give

$$\Phi_{k,s} = F_{\Xi,k}^{(s)} = L_{q-1} + L_q > 0. \quad (\text{RWC10})$$

Define the actual finite bounds

$$\begin{aligned} \ell_{k,s} &= \sum_{N \in \{q-1, q\}} \max\{\log(1 + t_N), j \log(1 + c_N^{1/j})\} > 0, \\ u_{k,s} &= \sum_{N \in \{q-1, q\}} j \log(1 + t_N/j), \quad \ell_{k,s} \leq \Phi_{k,s} \leq u_{k,s}. \end{aligned} \quad (\text{RWC11})$$

These are complete expressions in the actual C, D, u_n and their Gram matrices. Together with ROQ21 they give the stronger endpoints $\Phi_{k,s}^- = \max(\ell_{k,s}, \lambda_s)$ and $\Phi_{k,s}^+ = \min(u_{k,s}, j(\mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}) - \lambda_s)$. Here $\lambda_s = \log(1 + \eta_{q-1})$ and the full original $\mathcal{L}_{s,t}$ are exactly ROQ17–21; all masses are retained.

For the already proved original seven-moment receiving equality, retain the finite domain of PMR1–3 and PMR7: $k \geq 29$ and $k\sqrt{\delta^2 + \gamma^2} \leq 2^{-33}(k+1)^2$. Write $\mathcal{F}_{0,k}^{(s)}$ for the full seven-moment outer expression, and set $e_{k,s}^{0,\pm} = e_{k,s}^\pm + c_k^\circ$ with the exact OAR error endpoints and PMR2's complete root-to-moment error c_k° . Substitution, without changing either endpoint, proves

$$\begin{aligned} \mathcal{F}_{0,k}^{(s)} - e_{k,s}^{0,-} - \Phi_{k,s}^+ &\leq F_K^{(s)} \\ &\leq \mathcal{F}_{0,k}^{(s)} + e_{k,s}^{0,+} - \Phi_{k,s}^-. \end{aligned} \quad (\text{RWC12})$$

Outside that PMR domain the original OAR outer-return formula gives the same interval with $\mathcal{F}_{\partial,k}^{(s)}$ in place of $\mathcal{F}_{0,k}^{(s)}$ and $e_{k,s}^\pm$ in place of $e_{k,s}^{0,\pm}$, throughout OAR's stated domain. RWC1–11 remain valid at every $k \geq 9$ specified at the start of this section. Every stronger previously proved kernel bound can be intersected with these endpoints. For the arithmetic statement also retain the original AMT period domain $|u| \geq R_*$, as in OVR's opening paragraph. In the complete OAR/OVR signed return $\Psi_k = F_K^{\text{ar}} + F_B^{\text{ar}} - F_B^{(k)}$, the same literal substitution gives

$$\begin{aligned} \mathcal{F}_{0,k}^{(k)} + \mathcal{H}_k + \delta_k^\sigma - e_{k,k}^{0,-} - \Phi_{k,k}^+ &\leq \Psi_k \\ &\leq \mathcal{F}_{0,k}^{(k)} + \mathcal{H}_k + \delta_k^\sigma + e_{k,k}^{0,+} - \Phi_{k,k}^-. \end{aligned} \quad (\text{RWC13})$$

The original correlated arithmetic deficit δ_k^σ , boundary return $\mathcal{H}_k$, and each unequal source error are unchanged. Thus every computed exterior coefficient in RWC8 has a proved map into both the kernel and signed-return estimates. The strict finite positivity in RWC10 alone supplies no rate for $\Phi_{k,s}/(kq)$. The quantitative next calculation is the growth of these same conditioned matrices or their exterior coefficients in the fixed actual period, not a replacement by an arbitrary rank- j positive matrix.

The new lower endpoint strictly strengthens the original first-increment endpoint, through an exact comparison on the same covariance. ROQ18 telescopes to

$$V_{q-1} = \sum_{a=q-1}^{2q-2} \frac{z_a z_a^*}{a_a}, \quad t_{q-1} = \sum_{a=q-1}^{2q-2} \frac{z_a^* \Omega_{q-1}^{-1} z_a}{a_a} \geq \eta_{q-1}, \quad \ell_{k,s} > \lambda_s. \quad (\text{RWC14})$$

Here z_a, a_a, η_a are precisely ROQ17 at their original individual cutoffs; the sum retains every later mixed covariance. Each summand is nonnegative, and the first is exactly η_{q-1} . Thus the first-window trace lower bound in RWC11 is at least $\log(1 + \eta_{q-1}) = \lambda_s$. The second-window lower bound is strictly positive by RWC5 and RWC7. Adding them proves the strict inequality. Consequently the original maximum in $\Phi_{k,s}^- = \max(\ell_{k,s}, \lambda_s)$ equals $\ell_{k,s}$ by this proved comparison; both constituent bounds remain recorded, with their original definitions and domains.

62 The fixed-period determinant and closed coefficients in the original receiver

Retain the complete original objects, source orders, observations and arithmetic maps of OMR1–26. The PCL, ROG, FEX, FGC, ECL, EQR and RDS proofs are supplied in full. This section composes their exact maps with the current receiver. The conductor and fixed-period determinant applications keep $m = 1$, $k = 4l + 1 \geq 9$, $q = (k + 1)^2$, $q' = (k - 7)^2$, $\Delta = 16k - 48$, $r = 8k - 16$, $j = 8k - 32$, $c = k/2$, $c' = c - 4$, and the fixed actual five-orbit period and its original logarithm branch. Each earlier eventual comparison retains its explicit finite domain.

62.1 The existing outer coefficient has an exact endpoint expression

To distinguish the endpoint values from the conductor moment index v , write $(u_E, v_E, w_E, z_E, c_E)$ for exactly ECL2–3 at $(\alpha, \beta) = (2, \pi)$. This is the literal dictionary $u_E = u$, $v_E = v$, $w_E = (u+v)/2$, $z_E = uv$, $c_E = (v^2 - u^2)/4$ in that provider. With κ_E denoting its elliptic modulus, the original endpoint equations and coefficient are

$$\begin{aligned} u_E K(\kappa_E) &= 2, & v_E E(\kappa_E) &= 4, & \kappa_E^2 &= 1 - u_E^2/v_E^2, & 0 < u_E < v_E, \\ \mathfrak{C}_\partial &= 4 \log(4/\pi) + 8 \log w_E - 4 \log z_E - 2 \log c_E, \\ 21.66851180520 &< 16\mathfrak{C}_\partial < 21.66851180521. \end{aligned} \tag{OER1}$$

Here K, E are the original complete elliptic integrals with modulus convention. ECL6 and ECL14 give $\ell = 8 - 4 \log w_E - 2 \log c_E$, $\mathcal{L} = 6 \log w_E - 2 \log z_E - 2$ and $F = -\ell/2 - 3 + \mathcal{L}$. Substitution in the exact OFV/ECL16 expression $4 \log(4/\pi) - 10 - 2F + 4\mathcal{L}$ gives OER1: the constant terms are exactly $-4 + 8 - 4 = 0$. Also $F = I_{2,\pi} - 6 + 2\mathcal{L}$, so this number is exactly $4 \log(4/\pi) + 2 - 2I_{2,\pi}$ in OMR2, MRI4r and BRI6h. The integer outward certificate ECL19–24 proves its displayed enclosure; all stored endpoints and its complete code are retained.

Consequently the existing exact relations and their errors become

$$\begin{aligned} \mathcal{O}_k^\pm &= \Delta q [4 \log(4/\pi) + 8 \log w_E - 4 \log z_E - 2 \log c_E] \pm 4\Delta\varepsilon_k^{\text{out}}, \\ F_K^{(s)} + \Phi_{k,s} &= \Delta q \mathfrak{C}_\partial + R_{k,s}, \quad b_{k,s} - e_{k,s}^- \leq R_{k,s} \leq b_{k,s} + e_{k,s}^+, \\ \Psi_k + \Phi_{k,k} &= \Delta q \mathfrak{C}_\partial + \mathcal{H}_k + \delta_k^\sigma + R_{k,k}. \end{aligned} \tag{OER2}$$

These are substitutions into OMR2, MRI4s and BRI6h. The complete OOQ error, the earlier $O(q \log q)$ OFV error, and both signed OAR errors retain their original finite definitions. Put $c_- = 21.66851180520/16$, $c_+ = 21.66851180521/16$. The original outer value is also enclosed by $[\Delta q c_- - 4\Delta\varepsilon_k^{\text{out}}, \Delta q c_+ + 4\Delta\varepsilon_k^{\text{out}}]$. This follows from $\Delta q > 0$ and OER1 and retains the exact coefficient in all identities.

62.2 The actual period coefficients and the pivot morphism

PCL1–12 gives the full entire coefficient matrix $\mathcal{R}(z) = \sum_{n \geq 0} R_n z^n$, with every coefficient specified by the finite sums in PCL2, and the exact identities

$$\begin{aligned} \mathbf{H}(u) &= e^{a/u} G_u \mathcal{R}(u^{-1}) V^{-1} U, & \det \mathcal{R}(z) &= 1, \\ \mathcal{R}(0) &= \begin{pmatrix} 1 & 0 & -\beta/3 & 0 \\ 0 & 1 & 0 & -2\beta/3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\ L_a(z) &= U^{-1} V \mathcal{R}(z)^{-1} D^{-a} \mathcal{R}(z) V^{-1} U, & A_z(Z, W) &= \prod_{a=1}^4 Q_0(L_a(z)(ZW, Z, W, 1)^\top), \\ (C_{\text{in}}(z))_{(a,b),(i,j)} &= a_{i+4-a,j+4-b}(z), & C_{\text{in}}(-z) &= C_{\text{in}}(z)^\top. \end{aligned} \tag{OER3}$$

All 81 coefficients are those of this displayed product, with PCL10's full sums; $U, V, G_u, \mathbf{a}, \beta, D$ retain their original definitions. Conjugating D^{-a} by $\mathbf{H}$ cancels its scalar factors; G_u commutes with D^{-a} , leaving exactly L_a . PCL2's $n = 0$ terms give the constant matrix, including both nonzero off-diagonal entries. PCL22–24 proves the last identity coefficient by coefficient. This is the actual matrix family in GIP8 and OMR9; no entry has been removed in taking its coefficient limit.

For the original OCF leading pivot let $\mathcal{P}_k$ be its actual pivot rectangle and $\mathcal{E}_k$ its complement. With the original insertion S_k and full reduction N_k ,

$$\begin{aligned} S_k^\top N_k^\top &= I_\Delta, & \text{im } N_k^\top &= V_A, \\ J_{\mathcal{E}} &= N_k^\top D_{\mathcal{P},\mathcal{E}} V_{\mathcal{E}}, & \rho_{\mathcal{E}} &= V_{\mathcal{E}}^{-1} D_{\mathcal{P},\mathcal{E}}^{-1} S_k^\top|_{V_A}, & \rho_{\mathcal{E}}^{-1} &= J_{\mathcal{E}}, \\ \Xi_{\mathcal{E}} &= \Xi_k^{\text{fix}} J_{\mathcal{E}}, & \ker \Xi_{\mathcal{E}} &= \rho_{\mathcal{E}}(K), \\ H_{\mathcal{E},N} &= J_{\mathcal{E}}^* G_N J_{\mathcal{E}}, & Q_{\Xi,N}^{\text{fix}} &= (\Xi_{\mathcal{E}} H_{\mathcal{E},N}^{-1} \Xi_{\mathcal{E}}^*)^{-1}. \end{aligned} \tag{OER4}$$

The diagonal $D_{\mathcal{P},\mathcal{E}}$ consists of the full root products $P_{\mathcal{P},k}(\omega_{ab})$, and $V_{\mathcal{E}}$ is the evaluation Vandermonde in that actual complementary root set. Distinct roots make both invertible. Multiplication proves both inverse identities; composition gives the kernel. Pullback of the unchanged Gamma form and minimum over the same residual fibre prove the last line. All root factors and the full metric in PCL19–21 remain present. No central-grid OOQ bound is applied to the changed root set without that transport.

62.3 The full low graph has its own finite-rank transport

At each original endpoint put $L_N = N - q$ and $M_N^{\text{pol}} = L_N + \Delta - v$. Let $\mathbf{U}_{L_N}$ be the complete ROG4 positive-shift relation matrix and Π_{U,L_N} its weighted orthogonal projection in $\|g\|_{\chi',s,c'} = \|\chi'g\|_{s,c'}$, $\chi' = \chi_{k-8}$. The finite inverse ROG12 gives the exact maps

$$\begin{aligned} Q_{N,s}^{\text{low}} g &= P_{<v} Z_{\chi,N} \mathcal{U}_{L_N}^{-1} \Pi_{U,L_N} g, & \text{rank } Q_{N,s}^{\text{low}} &\leq v, & \|Q_{N,s}^{\text{low}}\| &\leq H_{N,s}^{\text{ROG}}, \\ \mathcal{R}_{N,s}(x, g) &= (x - Q_{N,s}^{\text{low}} g, g), & B_{N,s}^{\text{str}} &= B_N^{\text{graph}} \mathcal{R}_{N,s}^{-1}, \\ \ker B_{N,s}^{\text{str}} &= \text{im} \left[\begin{pmatrix} 0 \\ \mathbf{U}_{L_N} \end{pmatrix}, \mathcal{R}_{N,s} W_{K,N} \right]. \end{aligned} \tag{OER5}$$

$H_{N,s}^{\text{ROG}}$ is exactly ROG24's product $A_{N,s,c}^+ (1 + R)^q I_{L_N} D_{M_N^{\text{pol}},\chi'} A_{N-v,s,c'}^-$, with all ROG6–24 constants. At $L_N = -1$ the map and its bound are zero. The shear sends $(P_{<v} Z_{\chi,N} h, \mathcal{U}_{L_N} h)$ to $(0, \mathcal{U}_{L_N} h)$; applying it to the full GIP kernel proves the last line, retaining every common-kernel column.

For the exact product source Gram define $Q_{N,s}^{\text{str}}$ by ROG31 and $\Phi_{k,s}^{\text{str}} = \sum_N \sigma_N \log \det Q_{N,s}^{\text{str}}$, with the original signs $(1, 1, -1, -1)$. Then

$$\begin{aligned} E_s^{\text{str}} &= 2 \sum_N E_N^{\text{CEP}} + 2v \sum_N \log(1 + H_{N,s}^{\text{ROG}}), \\ |\Phi_{k,s}^{\text{str}} - \Phi_{k,s}| &\leq E_s^{\text{str}} = o(kq). \end{aligned} \tag{OER6}$$

Each singular value σ of the low map gives two shear singular values $(\sqrt{\sigma^2 + 4} + \sigma)/2$ and its reciprocal. At most v exceed one. Thus every exterior forward and inverse logarithmic cost is at most $v \log(1 + H_{N,s}^{\text{ROG}})$. Minimum lifts and projection off the entire kernel contract exterior norms. Squaring gives the determinant bound; adding the full GIP source-map cost and the four signs proves OER6. ROG25 gives $\log(1 + H_{N,s}^{\text{ROG}}) = O_{\text{actual}}(q \log(q + 1))$; fixed $v \leq 80$ makes it $o(kq)$. The full ROQ/FMC32 minimum-form identity identifies the actual four-endpoint residual with $\Phi_{k,s}$.

62.4 The fixed columns arise from an exact quotient of that graph

Put $\alpha_A = v!/\mu_v$, $\mathcal{C}_A = \alpha_A \mathcal{T}_A$, $L = \mathbb{C}[S]_{<v} \subset E$, $K^+ = K + L$ and $t_0 = \dim(K \cap L)$. The explicit coordinate isomorphism $(\ell, z) \mapsto (\ell, \alpha_A z)$ sends the graph relations to $(P_{<v} Z_{\chi, N} h, \alpha_A \mathcal{U}_{L_N} h)$. Projection to the second coordinate gives the exact sequences

$$\begin{aligned} 0 \longrightarrow L \longrightarrow V_A \longrightarrow V_A/L \simeq \mathbb{C}[S']_{\leq N-v-q'}/\alpha_A \mathcal{U}_{L_N} \mathbb{C}[S]_{\leq N-q} \longrightarrow 0, \\ 0 \longrightarrow K \cap L \longrightarrow K \longrightarrow K^+/L \longrightarrow 0, \quad \dim(K^+/L) = t = r - t_0. \end{aligned} \quad (\text{OER7})$$

For a zero lower class, subtract its full graph vector from $(\ell, \alpha_A \mathcal{U} h)$, leaving $(\ell - P_{<v} Z_{\chi, N} h, 0)$. Such a low vector is a relation only when $\mathcal{U} h = 0$, which forces $h = 0$ by the nonzero leading coefficient $\text{lc}(h) \mu_v \binom{q+\deg h}{v}$. This proves the first kernel and identifies it with the original L . Restriction to K proves the second sequence, as in FGC3–8.

In the complete AKJ basis let $\Xi = (\Xi_{\text{lo}}, \Xi_{\text{hi}})$ and let Q_{lo} have kernel $\text{im } \Xi_{\text{lo}}$. Choose the specified fixed full basis W_k of $\ker(Q_{\text{lo}} \Xi_{\text{hi}})$ using actual nonzero minors. The original low and high columns give

$$\begin{aligned} Z_k = \text{diag}(d_h) W_k, \quad d_h = \frac{(q' + h + v)!}{(q' + h)!}, \quad \mathcal{C}_A P_h = d_h \psi(S') S'^h, \quad \psi = \chi_{k-8}, \\ R_a^A(S') = \frac{\mathcal{C}_A(\chi S^a)}{\psi} = \alpha_A \sum_{\nu} a_{\nu} \frac{\chi(S' + b_{\nu})}{\psi(S')} (S' + b_{\nu})^a. \end{aligned} \quad (\text{OER8})$$

Solvability of $\Xi_{\text{lo}} \ell + \Xi_{\text{hi}} z = 0$ is exactly the condition defining W_k , so Z_k represents K^+/L . Each relation polynomial has degree $\Delta - v + a$ and nonzero leading coefficient $(q + a)_v$, while the Z_k columns have degree below $\Delta - v$. Their quotient classes are independent.

Let $H_{N,s}^{\psi}$ be the full monomial Gram in the original measure $|\psi(c' + iy)|^2 dm_{0,s}(y)$, and R_N^A the first $N + 1 - q$ relation columns of OER8. Their four ranks are exactly 0, 1, q , $q + 1$. The complete minimum matrix is

$$\begin{aligned} \widehat{H}_{K,N}^{(s)} &= Z_k^* H_{N,s}^{\psi} Z_k - Z_k^* H_{N,s}^{\psi} R_N^A ((R_N^A)^* H_{N,s}^{\psi} R_N^A)^{-1} (R_N^A)^* H_{N,s}^{\psi} Z_k, \\ \widehat{F}_k^{(s)} &= \sum_N \sigma_N \log \det \widehat{H}_{K,N}^{(s)}. \end{aligned} \quad (\text{OER9})$$

Minimizing the norm of $Z_k x + R_N^A h$ gives $h = -((R_N^A)^* H_{N,s}^{\psi} R_N^A)^{-1} (R_N^A)^* H_{N,s}^{\psi} Z_k x$. Substitution proves the full Schur formula including both cross factors. At the first endpoint there is no relation variable. Independence proves positivity. This is exactly FEX29/FGC11 on the native lower metric. The original graph metric in these coordinates is $D_{\alpha,N}^{-*} H_N^{\text{actual}} D_{\alpha,N}^{-1}$; the next bounds retain its change to that native metric.

62.5 Both low terms and all one-sided errors enter the current interval

Set $e_{N,s}^{\text{FEX}} = \log \det \widehat{H}_{K,N}^{(s)} - \log \det H_{K^+/L,N}^{(s)}$. Keep exactly FEX18–20/34:

$$\begin{aligned} n_N = N + 1 - v, \quad m_N = \log \max \left\{ 1, \frac{v!}{|\mu_v|} \sum_{\nu} |a_{\nu}| e^{|b_{\nu}-4|(H_N+2)} \right\}, \quad H_N = \sum_{a=1}^N \frac{1}{a}, \\ \zeta_{N,s} = \log \prod_{d=v}^N \frac{w_d}{w_{d-v}}, \quad w_d = \frac{d!}{(s/2)_d}, \\ e_{N,s}^- = \zeta_{N,s} - 2(n_N - t)m_N, \quad e_{N,s}^+ = 2tm_N, \quad e_{N,s}^- \leq e_{N,s}^{\text{FEX}} \leq e_{N,s}^+, \\ D_s^- = e_{q-1,s}^- + e_{q,s}^- - e_{2q-1,s}^+ - e_{2q,s}^+, \quad D_s^+ = e_{q-1,s}^+ + e_{q,s}^+ - e_{2q-1,s}^- - e_{2q,s}^-. \end{aligned} \quad (\text{OER10})$$

These $e^{\pm}$ are the FEX flag bounds, distinct from OAR's $e_{k,s}^{\pm}$. Here H_N is the harmonic number, distinct from the kernel Gram. The full flag identity gives $\mathcal{J} M^{-2(n_N-t)} \leq D_Q \leq M^{2t}$, with

the exact squared Jacobian $\mathcal{J} = e^{\zeta_{N,s}}$ and norm bound $M = e^{m_N}$. This proves the endpoint bounds and their four signed sums, keeping both original Gamma masses until their exact Jacobian cancellation.

The two full Schur factorizations of the same K^+ give

$$\begin{aligned} F_K^{(s)} &= \widehat{F}_k^{(s)} - \sum_N \sigma_N e_{N,s}^{\text{FEX}} + F_L^{(s)} - F_{K^+/K}^{(s)}, \\ 0 \leq F_L^{(s)} &\leq v\mathcal{L}_s, \quad 0 \leq F_{K^+/K}^{(s)} \leq (v - t_0)\mathcal{L}_s, \quad \mathcal{L}_s = \mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}, \\ f_s^- &= \widehat{F}_k^{(s)} - D_s^+ - (v - t_0)\mathcal{L}_s \leq F_K^{(s)} \leq f_s^+ = \widehat{F}_k^{(s)} - D_s^- + v\mathcal{L}_s. \end{aligned} \tag{OER11}$$

In fixed adapted frames the determinant of K^+ is both its L determinant times that of K^+/L , and its K determinant times that of K^+/K , including fixed Jacobians. Their four endpoint constants cancel because the signs sum to zero. This proves the first identity. The original AKS bound passes to both subquotients by restriction and minimization, proving the second line. OER10 then proves the last line. If the two actual low volumes are evaluated, the stronger interval retains their exact difference in place of the caps. Both low terms remain part of the identity in either case.

All existing OMR11–12 intervals remain. The new same-scalar intersection is

$$\begin{aligned} L_s^{\text{ext}} &= \max\{L_s^{\text{op}}, f_s^-, \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s}^{\text{str}} - E_s^{\text{str}} - e_{k,s}^-\}, \\ U_s^{\text{ext}} &= \min\{U_s^{\text{op}}, f_s^+, \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s}^{\text{str}} + E_s^{\text{str}} + e_{k,s}^+\}, \\ L_{\text{ar}}^{\text{ext}} &= \max\{L_{\text{ar}}^{\text{op}}, L_1^{\text{ext}} + a_{K,k}^-\}, \quad U_{\text{ar}}^{\text{ext}} = \min\{U_{\text{ar}}^{\text{op}}, U_1^{\text{ext}} + a_{K,k}^+\}, \\ F_K^{\text{ar}} &\in [L_{\text{ar}}^{\text{ext}}, U_{\text{ar}}^{\text{ext}}], \quad F_B^{\text{ar}} \in [\mathcal{B}_k^{\text{ar}} - U_{\text{ar}}^{\text{ext}}, \mathcal{B}_k^{\text{ar}} - L_{\text{ar}}^{\text{ext}}]. \end{aligned} \tag{OER12}$$

The first lines combine OER6, the original MRI4m equality, and OER11. Every interval bounds the same $F_K^{(s)}$. The last lines add the unchanged correlated arithmetic allowances of OMR12 and subtract from the same total. All prior finite branches remain available outside any added domain. FEX40–41 also permits the existing reverse source intersections with its full asymmetric Ω bounds.

Define $L_J^{\text{ext}} = L_k^{\text{ext}} + \mathcal{H}_k + \delta_k^\sigma + C_k^{\text{budget}}$ and $U_J^{\text{ext}} = U_k^{\text{ext}} + \mathcal{H}_k + \delta_k^\sigma + C_k^{\text{budget}}$. Then

$$\begin{aligned} L_k^{\text{ext}} + \mathcal{H}_k + \delta_k^\sigma &\leq \Psi_k \leq U_k^{\text{ext}} + \mathcal{H}_k + \delta_k^\sigma, \\ J_k^{\text{action}} &= 2 \sum_N w_N \log \epsilon_N^{\text{ar}} \in [L_J^{\text{ext}}, U_J^{\text{ext}}], \quad 4q \log \mathcal{L}_{h,k} \leq J_k^{\text{action}}, \\ \mathcal{L}_{h,k} &\leq \min_N \epsilon_N^{\text{ar}} \leq \prod_N (\epsilon_N^{\text{ar}})^{w_N/(2q)} \leq \exp(U_J^{\text{ext}}/(4q)), \\ \frac{\Psi_k}{kq} &= \frac{\widehat{F}_k^{(1)}}{kq} - \frac{\mathcal{C}_\Gamma}{4} + o(1). \end{aligned} \tag{OER13}$$

The first line is the exact OMR13 equality with $\mathcal{H}_k = -T_k$. The next lines use the same arithmetic budget, exact original reflection, and same positive class in OMR19–26. FEX36–38 proves $|F_K^{(s)} - \widehat{F}_k^{(s)}| = O_{\text{actual}}(q \log q) = o(kq)$; the original source difference and arithmetic correction are $o(kq)$, and the proved Gamma return divided by kq tends to $-\mathcal{C}_\Gamma/4$. Substitution proves the last line without asserting a limit for the unevaluated fixed-column determinant.

62.6 The relation quotient and the earlier finite Gamma centre

With every original coefficient, put $E_A(z) = \sum a_{ab} e^{b_{ab}z}$, $D = \partial_z$ and $\mathcal{L}_A = \sum a_{ab} e^{b_{ab}z} d_{ab}(D)$. The exact analytic dual of the denominator above is

$$\begin{aligned} \mathcal{L}_A \chi'(D) &= \chi(D) M_{E_A}, \\ 0 \longrightarrow \mathcal{E}_{k-8} &\xrightarrow{M_{E_A}} \mathcal{N}_v \xrightarrow{\mathfrak{F}_A} (\mathbb{C}[S']/\mathcal{UC}[S])^\vee \longrightarrow 0, \\ f_{\mathfrak{F}_A(N)}(z) &= \chi'(D)(N(z)/E_A(z)). \end{aligned} \tag{OER14}$$

$\mathcal{N}_v$ is the original upper exponential space with its first v Taylor values zero; the operator order and all spaces are RDS6–8. RDS5 proves the intertwining by the literal positive shifts and Leibniz’ rule. Its kernel consists precisely of $E_A \mathcal{E}_{k-8}$, since the monic constant-coefficient equation $\chi'(D)F = 0$ has exactly that lower exponential solution space; the full Vandermonde proves its initial-value uniqueness. Dimension $q - v - q' = \Delta - v$ then proves surjectivity. Because $\alpha_A \neq 0$, the relation subspace $\alpha_A \mathcal{UC}[S]$ in OER7 is the same subspace. RDS’s finite-degree direct decomposition identifies its dual at every original cutoff. The full source coefficient identity, original phases and masses, analytic disk and pole multiplicities remain RDS8–15; the Schur metric in OER9 is retained.

Finally ECL12 applies to precisely QGQ4’s original derivative $\mathcal{L}_1 = \partial_\alpha \mathcal{L}(2, \pi)$. Hence

$$\begin{aligned} c_q^\Gamma &= c_\Gamma^{[q,1]} = \frac{1}{8} \log(w_E^2/z_E) - \frac{1}{4} \log 2, \quad -0.080453400358 < c_q^\Gamma < -0.080453400357, \\ \frac{\mathcal{V}_{q,l}^c - lq\mathcal{C}_\Gamma}{q} &\longrightarrow c_q^\Gamma, \quad \limsup_{l \rightarrow \infty} \left| \frac{W_k - lq\mathcal{C}_\Gamma}{q} - c_q^\Gamma \right| \leq A/\sqrt{2}, \quad A = M/2 + 2C > 0. \end{aligned} \quad (\text{OER15})$$

This uses exactly the QGQ domain $m = 1$, $k = 4l + 1$, $q = (4l + 2)^2$, $l \geq 1$. EQR1–3 proves the connection: ECL12 gives $\mathcal{L}_1 = \log(w_E^2/z_E)$; the complete QGQ7 centre has correction $(2\mathcal{L}_1 - 4\log 2)l^2$ and remaining $O(l)$ terms. Since $l^2/q \rightarrow 1/16$, division gives the stated centre limit. The full unequal finite errors remain

$$-p_{q,l}^- - \mathcal{U}_{q,l} - E_{q,l} \leq W_k - \mathcal{V}_{q,l}^c \leq p_{q,l}^+ + \mathcal{U}_{q,l} + T_{q,l} + E_{q,l}. \quad (\text{OER16})$$

All terms except $\mathcal{U}_{q,l}$ become zero after division by q , while $\mathcal{U}_{q,l}/q \rightarrow A/\sqrt{2}$. Adding the finite-centre limit proves OER15 with the same radius. The original definitions give $C \geq 0$ and $M \geq \log(w_E^2/z_E) > 0$, since $w_E^2/z_E = 1 + (v_E - u_E)^2/(4u_E v_E) > 1$. The radius remains strictly positive. The evaluation does not assert a q -scale limit for the actual W_k or change its finite interval.

62.7 Full-window residual bounds in these same receivers

Retain the literal RWC4 matrices for each source $s = 1, k$ and the two low cutoffs $N = q - 1, q$. Their full-window positive update, including the full $q \times q$ conditioning matrix, is

$$\begin{aligned} U_N^+ &= [u_{N+1}, \dots, u_{N+q}], \quad B_N = I_q + (CU_N^+)^*(CR_N C^*)^{-1}(CU_N^+), \\ Z_N &= D[I_q - R_N C^*(CR_N C^*)^{-1}C]U_N^+, \quad V_N = Z_N B_N^{-1} Z_N^* > 0, \quad \Omega_{N+q} = \Omega_N + V_N. \end{aligned} \quad (\text{OER17})$$

The original full conductor rows and residual rows here are precisely ROQ1 and ROQ4; D in this subsection denotes those rows. RWC2–6 proves the rank and update by the positive original quartet polynomial, orthogonality of the complete consecutive source block, and the displayed full covariance expansion. With $Y_N = \Omega_N^{-1/2} Z_N B_N^{-1/2}$, the exact four-endpoint residual and its finite bounds are

$$\begin{aligned} \Phi_{k,s} &= \sum_{N \in \{q-1, q\}} \log \left(\sum_{d=0}^j \left\| \bigwedge_{d=0}^d Y_N \right\|_{\text{HS}}^2 \right), \\ \lambda_s &< \ell_{k,s} = \Phi_{k,s}^- \leq \Phi_{k,s} \leq \Phi_{k,s}^+, \end{aligned} \quad (\text{OER18})$$

where the complete trace, determinant and earlier finite-cap definitions of ℓ, Φ^+ are RWC7–11. RWC8 proves the exterior equality by all singular products; RWC14 proves the strict improvement over the original first increment λ_s . Thus every exterior coefficient supplies a specified finite subtraction in the same kernel equation.

On the exact PMR range $k \geq 29$, $k\sqrt{\delta^2 + \gamma^2} \leq 2^{-33}(k+1)^2$, put

$$\begin{aligned} L_s^{\text{win}} &= \max\{L_s^{\text{ext}}, \mathcal{F}_{0,k}^{(s)} - e_{k,s}^{0,-} - \Phi_{k,s}^+\}, \\ U_s^{\text{win}} &= \min\{U_s^{\text{ext}}, \mathcal{F}_{0,k}^{(s)} + e_{k,s}^{0,+} - \Phi_{k,s}^-\}, \quad e_{k,s}^{0,\pm} = e_{k,s}^\pm + c_k^\circ, \\ L_s^{\text{win}} &\leq F_K^{(s)} \leq U_s^{\text{win}}. \end{aligned} \quad (\text{OER19})$$

RWC12 proves that each added interval contains the actual kernel volume, so taking its intersection with OER12 proves OER19. Outside that PMR range use the original OAR expression $\mathcal{F}_{\partial,k}^{(s)}$ and its original $e_{k,s}^{\pm}$ in precisely the same construction on OAR's domain. For the arithmetic statements retain $|u| \geq R_*$ as in AMT. Substitute $L_s^{\text{win}}, U_s^{\text{win}}$ for the source bounds in the last two lines of OER12, and for source k in OER13. This gives respectively the same correlated arithmetic interval, its exact boundary complement, and

$$\begin{aligned} L_k^{\text{win}} + \mathcal{H}_k + \delta_k^{\sigma} &\leq \Psi_k \leq U_k^{\text{win}} + \mathcal{H}_k + \delta_k^{\sigma}, \\ J_k^{\text{action}} &\leq U_k^{\text{win}} + \mathcal{H}_k + \delta_k^{\sigma} + C_k^{\text{budget}}. \end{aligned} \tag{OER20}$$

The proof is addition of the unchanged scalars to the actual kernel bounds and then the exact OMR26 identity. The full budget, its same-class lower bound, both low corrections and every unequal finite error remain present. No estimate of the individual kq -scale residual coefficient is inferred from strict finite positivity alone.

63 The complete common factor of the original polynomial relations

Retain the full original $\mathcal{U}$, d_{ab} , roots, coefficients, and degree $\mathbf{m} = \Delta - v$ of RDS1–2 and ROG1–6. Let $\mathcal{S}_A = \{(a, b) : a_{ab} \neq 0\}$ be the actual nonempty coefficient support, and define its four exact integer extrema

$$a_- = \min_{\mathcal{S}_A} a, \quad a_+ = \max_{\mathcal{S}_A} a, \quad b_- = \min_{\mathcal{S}_A} b, \quad b_+ = \max_{\mathcal{S}_A} b, \quad d_a = a_+ - a_-, \quad d_b = b_+ - b_-. \quad (\text{RCF1})$$

These are cases determined by the original coefficients; none is assigned a generic support. Write $k' = k - 8$ and extend only the root-index formula to integer pairs by

$$\tilde{\beta}_{ij} = c' + (2i - k')\delta + i(2j - k')\gamma. \quad (\text{RCF2})$$

The i multiplying the imaginary term is the imaginary unit. The positive real numbers δ, γ make this indexed map injective on all integer pairs by comparing real and imaginary parts.

Each original shifted outer product d_{ab} has precisely the simple roots indexed by

$$([-a, k - a] \times [-b, k - b]) \setminus ([0, k'] \times [0, k']), \quad (\text{RCF3})$$

where all intervals contain their integer points. Indeed $\beta_{ij} - b_{ab} = \tilde{\beta}_{i-a, j-b}$, and the rectangle removed in CEP41 is exactly the inner rectangle in these translated indices. This verifies the whole root set, not just its cardinality.

Every rectangle in RCF3 contains that same inner rectangle. Their intersection is $[-a_-, k - a_+] \times [-b_-, k - b_+]$. Consequently the exact monic greatest common divisor of the active products is

$$G_A(S') = \prod_{(i,j) \in ([-a_-, k - a_+] \times [-b_-, k - b_+]) \setminus [0, k']^2} (S' - \tilde{\beta}_{ij}), \quad (\text{RCF4})$$

$$g_A := \deg G_A = (k + 1 - d_a)(k + 1 - d_b) - q'.$$

Distinctness of the roots proves that the intersection product is their monic greatest common divisor, including every root with its multiplicity one. Empty products give $G_A = 1$. The rectangle is nonempty and contains the inner square because $0 \leq a_- \leq a_+ \leq 8$ and $0 \leq b_- \leq b_+ \leq 8$.

This product is also the monic generator of the polynomial ideal generated by the complete linear relation image:

$$(\mathcal{UC}[S]) = G_A \mathbb{C}[S']. \quad (\text{RCF5})$$

Here parentheses mean the ideal generated by that image, whose underlying linear subspace is still the full original image. To prove the equality, list the $t = |\mathcal{S}_A|$ active pairs as $\alpha_1, \dots, \alpha_t$ and form the matrix

$$V(S')_{n,j} = (S' + b_{\alpha_j})^n, \quad 0 \leq n < t, \quad 1 \leq j \leq t. \quad (\text{RCF6})$$

Its determinant, in the displayed order, is the nonzero constant $\prod_{i < j} (b_{\alpha_j} - b_{\alpha_i})$. The shift nodes are distinct by the same real and imaginary comparison as RCF2. Its adjugate divided by this constant is therefore an inverse matrix over $\mathbb{C}[S']$. The vector of the first t exact relations satisfies

$$(\mathcal{U}1, \mathcal{U}S, \dots, \mathcal{U}S^{t-1})^T = V(S')(a_{\alpha_j} d_{\alpha_j}(S'))_{j=1}^t. \quad (\text{RCF7})$$

Thus these two finite lists generate the same polynomial ideal. All nonzero a_{α_j} are invertible constants, so it is the ideal generated by the active d_{α_j} . The Euclidean algorithm for polynomials makes this the ideal of their monic greatest common divisor RCF4. For completeness, each polynomial remainder step $f = qg + r$ leaves the two generated ideals equal; decreasing degree terminates and identifies the final nonzero remainder, divided by its leading coefficient, with the monic common divisor. The other relations are polynomial combinations of the same active d_{α_j} , by their full formula RDS1. This proves RCF5 for the whole image.

In particular the actual linear image is a polynomial ideal if and only if

$$g_A = \mathfrak{m}. \quad (\text{RCF8})$$

If it is an ideal, it equals its ideal hull RCF5, whose codimension is g_A . RDS2 gives its actual codimension $\mathfrak{m}$, proving necessity. Conversely divide each active d_{ab} by the full G_A , and define the exact polynomial operator

$$\mathcal{V}_A h = \sum_{(a,b) \in \mathcal{S}_A} a_{ab} \frac{d_{ab}(S')}{G_A(S')} h(S' + b_{ab}), \quad \mathcal{U} = M_{G_A} \mathcal{V}_A. \quad (\text{RCF9})$$

Its degree on a nonzero degree- j input is exactly $j + \mathfrak{m} - g_A$ and its leading coefficient is still $\binom{q+j}{v} \mu_v$ times the input leading coefficient. This follows from monic exact division of RDS1; also $g_A \leq \mathfrak{m}$ because G_A divides the nonzero $\mathcal{U}1$ of degree $\mathfrak{m}$. If equality holds, $\mathcal{V}_A$ preserves degree with nonzero diagonal coefficients in every finite monomial frame. Repeated highest-degree elimination therefore proves that it is onto all polynomials. RCF9 then proves sufficiency in RCF8.

For every actual support, including the unequal-degree cases, the precise finite-dimensional factor sequence is

$$0 \longrightarrow \mathbb{C}[S']/\mathcal{V}_A \mathbb{C}[S] \xrightarrow{M_{G_A}} \mathbb{C}[S']/\mathcal{U} \mathbb{C}[S] \xrightarrow{\text{rem}_{G_A}} \mathbb{C}[S']/(G_A) \longrightarrow 0, \quad (\text{RCF10})$$

$$\dim(\mathbb{C}[S']/\mathcal{V}_A \mathbb{C}[S]) = \mathfrak{m} - g_A = (d_a + d_b)(k+1) - d_a d_b - v.$$

The first map is well defined by RCF9. If $G_A f$ belongs to $\mathcal{U} \mathbb{C}[S] = G_A \mathcal{V}_A \mathbb{C}[S]$, cancellation of the nonzero polynomial proves that f belongs to the first denominator, so that map is injective. The last map is onto by ordinary monic remainder, and its kernel consists exactly of the classes represented by multiples of G_A , the image of the first map. The same degree-elimination proof as RDS2, applied to RCF9, gives its first dimension; expansion of RCF4 gives the last expression. If $G_A = 1$, the rightmost quotient is zero and these maps retain the entire original relation quotient on the left.

The source metric transported by the first map is exactly

$$\|G_A f\|_{\chi', s, c'}^2 = \int |f(c' + iy)|^2 |\chi'(c' + iy) G_A(c' + iy)|^2 dm_s(y). \quad (\text{RCF11})$$

This is direct substitution into the original weighted norm ROG21. It retains every factor and the original source order and mass. Thus RCF9–11 are a specified map to the polynomial ideal supplied by the actual relations, together with the full remaining quotient. They do not identify its metric with the CTV outer-divisor metric: the original natural map to that divisor remains GIP7–9, while RDS8 gives the exact analytic realization of the complete intermediate quotient. The original residual map and its common-kernel subspace remain the transported ROG30 map on that same intermediate space.

63.1 Every original finite degree window and its exact metric sequence

For the original $N = q - 1, q, 2q - 1, 2q$ put $L = N - q$, $M = L + \mathfrak{m}$ as in ROG2, and set $r_A = \mathfrak{m} - g_A \geq 0$. At all four endpoints the exact finite version of RCF10 is

$$0 \longrightarrow \mathbb{C}[S']_{\leq M - g_A} / \mathcal{V}_A \mathbb{C}[S]_{\leq L} \xrightarrow{M_{G_A}} \mathbb{C}[S']_{\leq M} / \mathcal{U} \mathbb{C}[S]_{\leq L} \xrightarrow{\text{rem}_{G_A}} \mathbb{C}[S']_{< g_A} \longrightarrow 0, \quad (\text{RCF12})$$

$$\dim(\text{first space}) = r_A, \quad \dim(\text{middle space}) = \mathfrak{m}, \quad \dim(\text{last space}) = g_A.$$

Here the first numerator is zero when $M - g_A = -1$, and every denominator is zero at $L = -1$. The degree identity in RCF9 puts its image in precisely the first numerator. The maps and cancellations used in RCF10 preserve these degree bounds, proving exactness. Moreover $M \geq \mathfrak{m} - 1 \geq g_A - 1$, so all last-space remainder classes have representatives in the displayed middle numerator. The

first numerator dimension is $L + r_A + 1$, while its denominator dimension is $L + 1$ when $L \geq 0$; at $L = -1$ it is r_A with zero denominator. These counts prove the dimensions, including the empty cases.

Use the quotient norm of ROG21 on the middle space, the quotient norm of RCF11 on the first space, and the attained minimum quotient norm of the same ROG21 polynomial source under rem_{G_A} on the last space. These are exactly a Hilbert restriction and its quotient. Indeed, every lift of a middle class $[G_A f]$ differs from it by $\mathcal{U}h = G_A \mathcal{V}_A h$; thus all its lifts remain multiples of G_A , and its minimum is precisely the first space's stated norm. Quotienting further by that first space allows every change by a multiple of G_A within the finite window, exactly the last remainder norm. No polynomial division map was presumed to be orthogonal in this argument.

Let $H_{U,N}^{(s)}$, $H_{V,N}^{(s)}$, and $Q_{G,N}^{(s)}$ be these three Grams in their canonical remainder coefficient frames $1, S', \dots$ of dimensions $\mathfrak{m}, r_A, g_A$. Those remainder frames exist by the degree elimination proofs RDS2 and RCF9. Then the exact determinant identity is

$$\det H_{U,N}^{(s)} = \det H_{V,N}^{(s)} \det Q_{G,N}^{(s)}. \quad (\text{RCF13})$$

To prove it, use in the middle quotient the fixed frame $[G_A, G_A S', \dots, G_A S'^{r_A-1}, 1, S', \dots, S'^{g_A-1}]$. Reordering it by increasing leading degree gives a triangular monic coefficient matrix. Its original order therefore has determinant $(-1)^{g_A r_A}$ and modulus one, retaining every lower coefficient of G_A in the matrix. The first Gram block is $H_{V,N}^{(s)}$; its Schur complement is the attained quotient Gram $Q_{G,N}^{(s)}$ by the minimum-lift calculation. Block Gaussian elimination proves RCF13, including zero-dimensional determinants equal to one.

The four-return identity thus has the unchanged endpoint signs:

$$F_U^{(s)} = F_V^{(s)} + F_G^{(s)}, \quad F_G^{(s)} = \sum_{N=q-1, q, 2q-1, 2q} \sigma_N \log \det Q_{G,N}^{(s)}, \quad \sigma = (1, 1, -1, -1). \quad (\text{RCF14})$$

The other two returns are the same sums of their Grams in RCF13. The full source orders are $s = 1, k$. Thus RCF14 splits an actual original relation quotient metric; the remaining factor has the explicit weighted source RCF11 and the complete operator RCF9.

There is also a fixed original primary map, before any metric choice or shear:

$$\rho_G : V_A \twoheadrightarrow \mathbb{C}[S']_{<g_A}, \quad [p] \mapsto \text{rem}_{G_A} \left(\frac{\mathcal{T}_{Ap}}{\chi'} \right). \quad (\text{RCF15})$$

Membership in V_A makes the quotient polynomial by CEP5. A change of representative by χh changes it by $\mathcal{U}h$, divisible by G_A , proving independence. The map is onto: for a prescribed polynomial f of degree below g_A , the source surjectivity CEP6 gives a preimage of $\chi' f$ of degree at most $q' + g_A - 1 + v \leq q - 1$. This preimage belongs to V_A and has the stated remainder. At $g_A = 0$ the target is zero and surjectivity has its usual zero-dimensional meaning. The map is consequently the same one at all original endpoints and in both original source norms.

64 Original Gamma control on the actual common-factor quotient

Retain RCF1–15, all fixed original period coefficients and their actual support, and the complete original sources $s = 1, k$. Write $g = g_A$, $\mathbf{m} = \Delta - v$, $M = N - q + \mathbf{m}$. If $g = 0$, the common-factor target and every quotient of it are zero, their determinants are one, and all conclusions below hold with zero return. In the proof of its positive-dimensional operator inequalities suppose $g \geq 1$; this includes every remaining actual support case.

64.1 The exact original divisor, windows, phases and source

Use the unchanged real source coordinate $S' = c' + iy$ and define

$$G_y(y) = i^{-g} G_A(c' + iy), \quad \chi'(c' + iy) = i^{q'} P_{k-8}(y), \quad R_G = (k + 8)\sqrt{\delta^2 + \gamma^2}. \quad (\text{RCM1})$$

These are monic polynomials with exactly the displayed phases. By RCF2–4 the roots of G_y are $(2j - k')\gamma - i(2i - k')\delta$ in its stated integer rectangle. Every $|2i - k'|, |2j - k'|$ there is at most $k + 8$; hence every root has modulus at most R_G . They are distinct. No root or coefficient is set to a surrogate value in RCM1.

The target coefficients of $1, S', \dots, S'^{g-1}$ and of $1, y, \dots, y^{g-1}$ are related by the fixed invertible matrix

$$T_{ab} = \begin{cases} \binom{b}{a} (c')^{b-a} i^a, & a \leq b, \\ 0, & a > b, \end{cases} \quad 0 \leq a, b < g, \quad \det T = i^{g(g-1)/2}. \quad (\text{RCM2})$$

This follows by expanding $(c' + iy)^b$. In particular $|\det T| = 1$, while the full matrix, including its centre, is retained in every operator congruence. Let $Q_{G,M}^{y,(s)}$ be the minimum remainder Gram by G_y on polynomials of degree at most M , with source

$$d\zeta_{k',s}(y) = |P_{k-8}(y)|^2 dm_s(y). \quad (\text{RCM3})$$

The complete phases in RCM1 have modulus one. Substitution proves $Q_{G,N}^{(s)} = T^* Q_{G,M}^{y,(s)} T$ for the RCF12–14 Gram. The four actual M values are

$$\mathbf{m} - 1, \quad \mathbf{m}, \quad q + \mathbf{m} - 1, \quad q + \mathbf{m}. \quad (\text{RCM4})$$

They include all additional columns beyond the divisor's degree: $g \leq \mathbf{m} \leq \Delta$ and $\mathbf{m} \geq 16$.

64.2 Complete finite bounds at the low source windows

For the comparison measure $|y|^{2Q} e^{-\eta|y|} dy$ use the full positive source rate $\eta > 0$ and put

$$\mathfrak{g}(Q, \eta) = 2Q \log(2Q/\eta) - 2Q, \quad y_0 = 2Q/\eta. \quad (\text{RCM5})$$

OOQ26–27 proves the following bounds on the entire polynomial coefficient space of degree at most Δ :

$$\frac{e^{\mathfrak{g}(Q, \eta) - \eta}}{(\Delta + 1)^2 (2\Delta + 1) 64^\Delta (1 + y_0)^{2\Delta}} I \preceq H_\Delta^{\text{exp}} \preceq \left(\sum_{j=0}^{\Delta} \frac{2(2Q + 2j)!}{\eta^{2Q+2j+1}} \right) I. \quad (\text{RCM6})$$

Here Q will be the original integer $q' + (s - 1)/4$. These estimates retain the positive-interval integration bound and the full two-half-line moment trace from OOQ, respectively.

For every monic degree- g polynomial whose roots have modulus at most R_G , the ordinary remainder map on degree at most Δ has coefficient operator norm at most

$$C_{\text{lo}} = 1 + (1 + R_G)^\Delta (\Delta + 1) \max\{1, \Delta R_G\}^\Delta. \quad (\text{RCM7})$$

To prove this for the actual G_y , its coefficient absolute sum is at most $(1 + R_G)^g$. Polynomial division uses the first $\Delta - g + 1$ complete homogeneous coefficients of its inverse at infinity. Their sum is at most $\sum_{a=0}^{\Delta-g} \binom{g+a-1}{a} R_G^a \leq (\Delta + 1) \max\{1, \Delta R_G\}^\Delta$. The coefficient convolution argument in ROG19–20 bounds the quotient operator by this sum. The exact remainder is the input minus its quotient multiplied by G_y , which proves RCM7. This proof does not truncate the actual root polynomial.

At either low window in RCM4, taking the minimum over all lifts of a remainder z gives a lower norm bound from RCM6 and $\|p\|_{\text{coef}} \geq \|z\|_2 / C_{\text{lo}}$. The canonical degree-less-than- g representative is an allowed lift and gives the upper bound from RCM6. Thus define

$$E_{\text{lo}}^G(Q, \eta) = \max \left\{ 0, \log \left(\sum_{j=0}^{\Delta} \frac{2(2Q + 2j)!}{\eta^{2Q+2j+1}} \right) - \mathfrak{g}(Q, \eta), \right. \\ \left. \eta + 2 \log(\Delta + 1) + \log(2\Delta + 1) + \Delta \log 64 + 2\Delta \log(1 + y_0) + 2 \log C_{\text{lo}} \right\}. \quad (\text{RCM8})$$

The full low remainder Gram obeys

$$e^{\mathfrak{g}(Q, \eta) - E_{\text{lo}}^G} I_g \preceq Q_{G, M}^{\text{exp}}(Q, \eta) \preceq e^{\mathfrak{g}(Q, \eta) + E_{\text{lo}}^G} I_g. \quad (\text{RCM9})$$

The extra low columns $g, \dots, M$ have all been included in the lower minimization. The factorial bounds proved after OOQ29 and $\log C_{\text{lo}} = O_{\delta, \gamma}(\Delta \log q)$ give $E_{\text{lo}}^G = O_{\delta, \gamma}(k \log q)$ in the source and rate ranges specified below.

64.3 The high windows with the actual analytic remainder and lift

Retain the equilibrium constants of OOQ3–6 on its entire specified compact parameter rectangle. Set

$$n = q/2, \quad \rho = n^{-1/8}, \quad r_* = R_G/n, \quad C_{\text{rem}} = \rho \Delta (1 + r_*)^\Delta (2/\rho)^\Delta. \quad (\text{RCM10})$$

Use the original OSP23 source-comparison range, $k \geq 257$, and the explicit eventual conditions

$$n \geq 2\Delta + 1, \quad n \geq 2/b_*, \quad r_* \leq \rho/2, \quad \rho \leq \min\{1/2, \sqrt{a_*/2}\}. \quad (\text{RCM11})$$

They hold for all sufficiently large original k at the fixed quartet, since $R_G = O(k)$ and $\Delta = O(k)$ while $n \asymp k^2$.

Apply the exact Cauchy remainder OOQ14 to $\hat{G}(z) = n^{-g} G_y(nz)$. Its g roots are distinct and lie inside $|z| \leq r_*$. Its coefficient norm bound is $\rho g (1 + r_*)^g (2/\rho)^g$, at most C_{rem} because $1 \leq g \leq \Delta$ and $2(1 + r_*)/\rho > 1$. The Cauchy formula interpolates the actual roots and is exactly the remainder by $\hat{G}$, as proved in OOQ14–15.

For either high M in RCM4, the parity degrees $m_e = \lfloor M/2 \rfloor$, $m_o = \lfloor (M - 1)/2 \rfloor$ belong to $[n, n + \Delta]$. The exact half-line parameters are $((Q - 1/2)/m_e, n\eta/m_e)$ and $((Q + 1/2)/m_o, n\eta/m_o)$. They lie in OOQ's compact rectangle for the same bounds as OOQ17, since $16 \leq \mathfrak{m} \leq \Delta$. The two-branch Jacobian, weighted polynomial evaluation bound, and smooth variation estimate are therefore precisely OOQ18–20, with the upper degree bound $n + \Delta$. These steps give the same lower quotient bound with C_{rem} from RCM10.

For completeness the upper lift is the literal polynomial

$$a_\varphi = \text{rem}_{\hat{G}} \left(\frac{\varphi}{r_n(\cdot 2)} \right), \quad p_\varphi(nx) = r_n(x^2) a_\varphi(x), \quad (\text{RCM12})$$

where φ denotes the prescribed degree-less-than- g remainder polynomial and r_n is the unchanged quantile polynomial OOQ9. Its denominator is nonzero on the integration disk because $\rho^2 < a_*$. The lift has degree at most $q + g - 1 \leq q + \mathfrak{m} - 1 \leq M$ and agrees at every actual root of $\hat{G}$ with the prescribed remainder. Thus it is an allowed original high-window lift. Its coefficient and integral estimates are OOQ22–23 with $g \leq \Delta$ and the tail power bounded by $2(\Delta - 1)$.

To retain all finite errors, put

$$\begin{aligned}
E^- &= 2 \log(C_{\text{rem}} \sqrt{2} C_{\text{BM}}(n + \Delta)^2) + 2C_W(\Delta + 2) + 4(n + \Delta)\rho^2/a_*, \\
E^+ &= 2 \log C_{\text{rem}} + \log \Delta + 4n\rho^2/a_* + 2C_{\text{quant}}\sqrt{n} + 2C_0 \\
&\quad + 2C_{\text{tail}}(\Delta - 1) + \log T_*, \\
E_{\text{hi}}^G &= \max\{0, E^-, E^+\} + 2(\Delta - 1) \log n, \\
c_{\text{hi}}^{\text{exp}}(Q, \eta) &= (2Q + 1) \log n - 2nW((Q - 1/2)/n, \eta).
\end{aligned} \tag{RCM13}$$

The exact original-versus-scaled remainder matrix is $\text{diag}(1, n, \dots, n^{g-1})$; its squared norm is at most $n^{2(\Delta-1)}$, the last term in RCM13. The preceding bounds prove at both high windows

$$e^{c_{\text{hi}}^{\text{exp}}(Q, \eta) - E_{\text{hi}}^G} I_g \preceq Q_{G, M}^{\text{exp}}(Q, \eta) \preceq e^{c_{\text{hi}}^{\text{exp}}(Q, \eta) + E_{\text{hi}}^G} I_g, \quad E_{\text{hi}}^G = O_{\delta, \gamma}(q^{3/4} + k \log q). \tag{RCM14}$$

Every imported estimate here has the same actual source, parity and centre scalings as OOQ. The only changes in its remainder proof are the specified divisor G_y , its proved root bound, and the degree allowances verified above.

64.4 Both full Gamma sources and the four-return coefficient

Retain the exact source constants of OOQ30–31:

$$\begin{aligned}
\vartheta &= \pi/2, \quad h = 1/q, \quad \ell_s = (s - 1)/4, \quad Q_s = q' + \ell_s, \quad \beta_s = \frac{(2\pi)^{s/2}}{\sqrt{2}\Gamma(s/2)}, \\
L_h &= \frac{\pi}{\Gamma(3/4)^2} (\sin h)^{3/2}, \quad U_h = \frac{\Gamma(1/4)^2}{2\pi} (\sin h)^{-1/2},
\end{aligned} \tag{RCM15}$$

and the complete root/source errors $\mathcal{E}_k, r_{\ell_s}$ from OSP24 and OFV6, with $r_0 = 0$. The full quadratic-form comparison OOQ31 applies to every polynomial used here, since $M \leq q + \Delta \leq 2q$ on RCM11's range. Taking minima over the same actual G_y fibres preserves those two inequalities, including all source masses.

Define the two unchanged leading centres

$$c_{\text{lo}} = \mathfrak{g}(q, \vartheta), \quad c_{\text{hi}} = 2q \log(q/2) - qW(2, \vartheta), \tag{RCM16}$$

and a single finite nonnegative error

$$\begin{aligned}
\varepsilon_k^G &= \mathcal{E}_k + \max_{s=1, k} r_{\ell_s} + \max\{|\log L_h|, |\log U_h|\} \\
&\quad + \max_{\substack{s \in \{1, k\} \\ \eta \in \{\vartheta - h, \vartheta + h\}}} \{E_{\text{lo}}^G(Q_s, \eta) + E_{\text{hi}}^G + |\mathfrak{g}(Q_s, \eta) - c_{\text{lo}}| + |c_{\text{hi}}^{\text{exp}}(Q_s, \eta) - c_{\text{hi}}|\}.
\end{aligned} \tag{RCM17}$$

All its constants are supplied above or by the complete original source and equilibrium proofs cited there. The resulting operator inequalities, in the original S' frame, are

$$\beta_s e^{c_{\bullet} - \varepsilon_k^G} T^* T \preceq Q_{G, N}^{(s)} \preceq \beta_s e^{c_{\bullet} + \varepsilon_k^G} T^* T, \tag{RCM18}$$

where $c_{\bullet} = c_{\text{lo}}$ at $N = q - 1, q$ and $c_{\bullet} = c_{\text{hi}}$ at $N = 2q - 1, 2q$. Moreover

$$\varepsilon_k^G = O_{\delta, \gamma}(q^{3/4} + k \log q) = o(q). \tag{RCM19}$$

Indeed RCM8 and RCM14 control the new errors. The source errors have the orders proved in OOQ35. Also $Q_s - q = -\Delta + \ell_s = O(k)$, the rates differ by $1/q$, and W with its first derivatives is bounded on the same compact rectangle. The mean-value estimates in OOQ33–35 give $O(k \log q)$ for both scalar discrepancies in RCM17. This proves RCM19 without a support-dependent lower bound.

OOQ36 gives the exact difference $c_{\text{lo}} - c_{\text{hi}} = q\mathfrak{C}_\partial/2$ with its unchanged equilibrium constant. Taking four determinants in RCM18, and retaining the signs and fixed frame of RCF14, therefore proves

$$|F_G^{(s)} - gq\mathfrak{C}_\partial| \leq 4g\varepsilon_k^G = o(kq). \quad (\text{RCM20})$$

The source factor β_s^g and $\det(T^*T)$ cancel with coefficients $1 + 1 - 1 - 1$. All individual Grams retain them. The exact relation return consequently obeys the proved partial evaluation

$$F_U^{(s)} = F_V^{(s)} + gq\mathfrak{C}_\partial + \epsilon_G^{(s)}, \quad |\epsilon_G^{(s)}| \leq 4g\varepsilon_k^G, \quad (\text{RCM21})$$

where the remaining F_V is the complete smaller operator and weighted source in RCF9–14, of rank $\mathfrak{m} - g$.

64.5 The resulting original residual quotient and its retained complement

Let $\Xi = \Xi_k^{\text{fix}} : V_A \rightarrow B_\Xi$ be the same original residual observation, with $B_\Xi = \mathbb{C}^j$, $j = 8k - 32$, and kernel K . Use the fixed onto map ρ_G proved in RCF15, and put

$$V_0 = \ker \rho_G, \quad W = \Xi(V_0), \quad t_G = \dim(B_\Xi/W) = g - \dim \rho_G(K). \quad (\text{RCM22})$$

There is the precise onto map

$$A_G : \mathbb{C}[S']_{<g} \rightarrow B_\Xi/W, \quad \rho_G x \mapsto [\Xi x], \quad \ker A_G = \rho_G(K). \quad (\text{RCM23})$$

Two lifts differ by V_0 , so the map is well defined. It is onto because Ξ is onto. If $\Xi x \in W$, subtract an element of V_0 with that same image; the remaining element belongs to K and has the same ρ_G value. This proves the kernel equality and RCM22's dimension. All these maps are independent of source order and cutoff.

In the graph source X_N of GIP1, the composite target map is exactly $(x, f) \mapsto A_G \text{rem}_{G_A} f$. It ignores the low coordinate and kills every full graph relation, since $\mathcal{U}h$ is divisible by G_A . Minimizing its product norm first over the free low coordinate and then over polynomial remainder fibres proves that its quotient Gram is exactly

$$Q_{A_G, G, N}^{(s)} = [A_G(Q_{G, N}^{(s)})^{-1} A_G^*]^{-1}. \quad (\text{RCM24})$$

Thus no equality of different relation subspaces enters this identification; only the proved composite map and its full kernel are used.

Let $\overline{Q}_{\Xi, N}^{(s)}$ be the original metric on B_Ξ/W induced by the original residual quotient. The actual source map S_N of CEP28 induces the identity of this target with RCM24. Its minimum-lift exterior argument CEP34–38, at the actual rank t_G , gives

$$|\log \det \overline{Q}_{\Xi, N}^{(s)} - \log \det Q_{A_G, G, N}^{(s)}| \leq 2E_N^{\text{CEP}}. \quad (\text{RCM25})$$

The source preimage is precisely $\mathcal{W}_N$ of CEP31. On its S_N image it is $\mathbb{C}^v \oplus \chi' \mathbb{C}[S']_{\leq M}$; the composite map is the one just minimized in RCM24. Orthogonal projection off the entire composite kernel is an exterior contraction, for both S_N and its inverse. This verifies all spaces in the bound and retains the original source map determinant from CEP29.

Taking the same fixed quotient of RCM18 replaces T^*T by the full fixed reference $[A_G(T^*T)^{-1} A_G^*]^{-1}$, through minimum on identical fibres. Its determinant and the original mass cancel only in the four-sign return. Consequently

$$\left| \sum_N \sigma_N \log \det \overline{Q}_{\Xi, N}^{(s)} - t_G q \mathfrak{C}_\partial \right| \leq 4t_G \varepsilon_k^G + 2 \sum_N E_N^{\text{CEP}} = o(kq). \quad (\text{RCM26})$$

No coefficient conditioning of the fixed quotient A_G has been assumed; its complete reference form is retained and cancels in the displayed return.

Finally choose any full fixed original frame I_W for W and put

$$F_W^{(s)} = \sum_N \sigma_N \log \det(I_W^* Q_{\Xi, N}^{\text{fix}} I_W).$$

Block elimination in a fixed frame of B_{Ξ} extending I_W gives its restriction determinant times the quotient determinant. The full fixed frame factor cancels in the four signs, exactly as in CEP26. Hence the original residual return has the established decomposition

$$F_{\Xi, k}^{(s)} = F_W^{(s)} + t_G q \mathfrak{C}_{\partial} + \epsilon_{\Xi}^{(s)}, \quad |\epsilon_{\Xi}^{(s)}| \leq 4t_G \varepsilon_k^G + 2 \sum_N E_N^{\text{CEP}}. \quad (\text{RCM27})$$

The remaining rank is $\dim W = j - t_G$. Its original metric and embedding have been specified, not replaced by a scalar or by a source norm. Also $F_W^{(s)} \geq 0$: the original source polynomial spaces are nested, with the same remainder and residual target, so minimum quotient metrics decrease as N increases. Restriction to the same W preserves this positive-form order, and the two low-versus-high determinant comparisons are nonnegative. Therefore RCM27 also gives the actual lower estimate $F_{\Xi, k}^{(s)} \geq t_G q \mathfrak{C}_{\partial} - 4t_G \varepsilon_k^G - 2 \sum_N E_N^{\text{CEP}}$. The value of t_G uses the exact active support and original map $\rho_G|_K$; no generic nonvanishing or zero-gcd case has been substituted for that calculation.

65 The original weighted relation metric as a minimum dual extension

Retain the original simple quartet, fixed five-orbit period and logarithm branch, all coefficients and shifts of ROG and RDS, and the same two source orders $s \in \{1, k\}$. Put

$$\begin{aligned} q &= (k+1)^2, & q' &= (k-7)^2, & \Delta &= q - q', & \mathbf{m} &= \Delta - v, & c' &= k/2 - 4, \\ N &\in \{q-1, q, 2q-1, 2q\}, & L &= N - q, & d &= N - v, & M &= N - v - q' = L + \mathbf{m}. \end{aligned} \quad (\text{DNE1})$$

Here $0 \leq v \leq 80$, $\mu_v \neq 0$, and χ' is the complete lower-grid polynomial χ_{k-8} , never a derivative. All spaces of degree at most -1 are zero. The source norm is the original norm at $S' = c' + iy$, with mass $M_s = (2\pi)^{s/2}$.

65.1 A fixed original exponential quotient frame

We specify a finite frame using only the actual original matrices. Let $\mathbb{V}$ be the upper-root Vandermonde with rows $a = 0, \dots, q-1$ and entries $\mathbb{V}_{a,(i,j)} = \beta_{ij}^a$. Let $P_v : \mathbb{C}^{q-v} \rightarrow \mathbb{C}^q$ insert zero in the first v positions, and let C be the exact primary conductor of CEP4, with q' rows and q columns. Define

$$W_v = \mathbb{V}^{-1}P_v, \quad J_v = P_v^\top \mathbb{V} C^\top, \quad W_v J_v = C^\top. \quad (\text{DNE2})$$

The Vandermonde is invertible by the original distinct roots. The first v rows of $\mathbb{V} C^\top$ vanish because the corresponding exponential columns are $E_A e^{\beta'_{ab} z}$, whose zero order is at least v . This proves the last equality. Consequently W_v is an isomorphism onto the coefficient vectors of $\mathcal{N}_v$, and J_v has rank q' , since $C^\top$ is injective.

Choose the first q' row positions $\mathcal{I}$ in a fixed ordering of all subsets for which $\det(J_v)_{\mathcal{I}} \neq 0$, and let $\mathcal{J}$ be their complement. Such a subset exists by the proved rank. Let $E_{\mathcal{J}}$ insert these $\mathbf{m}$ complementary coordinates and put

$$Z_{\text{exp}} = W_v E_{\mathcal{J}}, \quad \mathcal{N}_a(z) = \sum_{i,j} (Z_{\text{exp}})_{(i,j),a} e^{\beta_{ij} z}, \quad F_a(z) = \frac{\mathcal{N}_a(z)}{E_A(z)} \quad (1 \leq a \leq \mathbf{m}). \quad (\text{DNE3})$$

The columns of $[J_v, E_{\mathcal{J}}]$ are independent: restriction of a relation to $\mathcal{I}$ first kills its J_v coefficients, and then its other coefficients vanish. Thus the $\mathcal{N}_a$ give a basis of the exact quotient $\mathcal{N}_v / E_A \mathcal{E}_{k-8}$. Every root value and actual minor defining this frame is retained. The frame is fixed as the endpoint N and source order s change.

For an analytic germ F at zero define the complex-linear functional Λ_F by $\Lambda_F(S'^n) = F^{(n)}(0)$. Let ℓ_a be the functional from RDS8. Then

$$\ell_a(g) = \Lambda_{F_a}(\chi' g), \quad f_{\ell_a} = \chi'(D)F_a, \quad A_{\ell,ba} = \ell_a(S'^b) \quad (0 \leq b < \mathbf{m}). \quad (\text{DNE4})$$

Indeed multiplication by χ' in the functional is precisely the constant-coefficient operator $\chi'(D)$ on its generating series. RDS8 proves that these functionals form the entire dual of the relation quotient. RDS2 identifies that quotient with the increasing monomials below degree $\mathbf{m}$, so A_{ℓ} is invertible. This supplies its existence without assuming a nonzero unevaluated minor.

65.2 The exact original source extension problem

On $\mathbb{C}[S']_{\leq M}$ use the original weighted norm $\|g\|_{\chi',s,c'} = \|\chi' g\|_{s,c'}$. Let G_M be its full monomial Gram, and let $\mathbf{U}_L$ be the complete ROG4 relation matrix. The quotient

$$Y_N = \mathbb{C}[S']_{\leq M} / \text{im } \mathcal{U}_L \simeq \mathbb{C}[S']_{< \mathbf{m}}$$

has Gram, in its fixed low-monomial representative frame,

$$H_{Y,N}^{(s)} = E_{< \mathbf{m}}^* G_M E_{< \mathbf{m}} - E_{< \mathbf{m}}^* G_M \mathbf{U}_L (\mathbf{U}_L^* G_M \mathbf{U}_L)^{-1} \mathbf{U}_L^* G_M E_{< \mathbf{m}}. \quad (\text{DNE5})$$

The second term is absent when $L = -1$. Distinct leading degrees of the columns of $\mathcal{U}_L$ show that the low representatives and all relation columns form a basis. Positivity of G_M and minimization over the full relation columns therefore prove DNE5 and its strict positivity.

For $x \in \mathbb{C}^{\mathbf{m}}$ write $\mathcal{N}_x = \sum_a x_a \mathcal{N}_a$ and $F_x = \mathcal{N}_x / E_A$. Its functional ℓ_x on Y_N has the exact norm

$$\|\ell_x\|_{Y_N^\vee}^2 = \min_{\eta \in \mathbb{C}^{q'}} \left\| \Lambda_{F_x} + \sum_{a,b} \eta_{ab} \text{ev}_{\beta'_{ab}} \right\|_{(\mathbb{C}[S']_{\leq d}, \|\cdot\|_{s,c'})^\vee}^2. \quad (\text{DNE6})$$

Here the dual consists of complex-linear functionals. To prove the equality, multiply by χ' to identify the weighted polynomial space isometrically with $I_N = \chi' \mathbb{C}[S']_{\leq M} \subset \mathbb{C}[S']_{\leq d}$. The restriction of Λ_{F_x} to I_N is exactly $g \mapsto \ell_x(g)$ by DNE4, and it annihilates $\chi' \text{im } \mathcal{U}_L$ by RDS5.

A functional annihilating a subspace has the same norm on the Hilbert quotient: the minimum lift of each quotient class has the same functional value, and every other lift has larger norm. A functional on a Hilbert subspace has a unique extension of minimum norm to the ambient space, namely its composition with the orthogonal projection onto that subspace. Every other extension adds a functional on the perpendicular complement; their squared dual norms add. These two facts reduce its quotient dual norm to the minimum over all ambient extensions of its restriction.

The difference of two extensions annihilates I_N . That annihilator has dimension $(d+1) - (M+1) = q'$. Evaluation at each original lower root annihilates I_N , and these q' evaluation functionals are independent on $\mathbb{C}[S']_{\leq d}$, by the first q' monomials and the distinct-root Vandermonde. They therefore constitute its whole annihilator. This proves DNE6 and retains every permitted extension.

Let

$$\phi_{n,s,c'}(S') = \frac{p_n((S' - c')/i)}{\sqrt{h_n}}, \quad h_n = M_s n! (s/2)_n, \quad 0 \leq n \leq d$$

be the unchanged original orthonormal source frame. Form the complete finite matrices

$$\begin{aligned} T_{N,s;na} &= \Lambda_{F_a}(\phi_{n,s,c'}), & V_{N,s;n,(i,j)} &= \phi_{n,s,c'}(\beta'_{ij}), \\ \mathcal{D}_N^{(s)} &= T_{N,s}^* [I - V_{N,s}(V_{N,s}^* V_{N,s})^{-1} V_{N,s}^*] T_{N,s}. \end{aligned} \quad (\text{DNE7})$$

The matrix $V_{N,s}$ has rank q' by the same Vandermonde argument, since the displayed orthonormal polynomials span all degrees at most d . In an orthonormal basis the squared norm of a complex-linear functional is the sum of the squared moduli of its values. Thus the minimum in DNE6 is $\min_\eta \|T_{N,s}x + V_{N,s}\eta\|_2^2$. The minimizing vector is $-(V_{N,s}^* V_{N,s})^{-1} V_{N,s}^* T_{N,s}x$. Substitution proves DNE7, including both mixed factors, and gives $\|\ell_x\|^2 = x^* \mathcal{D}_N^{(s)} x$. Positivity follows because the ℓ_a are an independent dual frame of Y_N .

65.3 Complex-linear duality and the actual fixed kernel columns

For a positive Gram H and a complex-linear functional $y \mapsto a^\top y$, its squared dual norm is $a^\top H^{-1} \bar{a} = a^* \overline{H^{-1} a}$. Indeed its Riesz vector is $H^{-1} \bar{a}$, and evaluation of its squared norm gives the first expression; the second is its equal real conjugate. Therefore the precise metric identity and determinant consequence of DNE4–7 are

$$\mathcal{D}_N^{(s)} = A_\ell^* (\overline{H_{Y,N}^{(s)}})^{-1} A_\ell, \quad \det \mathcal{D}_N^{(s)} = \frac{|\det A_\ell|^2}{\det H_{Y,N}^{(s)}}. \quad (\text{DNE8})$$

The conjugation records the ordinary complex-linear dual coordinates; it is not discarded by changing the source frame. In the four-endpoint sum, the fixed factor $|\det A_\ell|^2$ cancels with signs $(1, 1, -1, -1)$.

Retain the actual fixed matrix Z_k of FEX27 and FGC9, with $t = r - t_0$ independent columns in the low-monomial frame of dimension $\mathbf{m}$. Multiplying every relation column by the same nonzero

scalar $\alpha_A = v!/\mu_v$ does not change its span or its Schur complement. Consequently its actual native FEX29 matrix and the exact dual quotient map are

$$\widehat{H}_{K,N}^{(s)} = Z_k^* H_{Y,N}^{(s)} Z_k, \quad B_K = Z_k^\top A_\ell, \quad Q_{K^\vee,N}^{(s)} = \left[B_K (\mathcal{D}_N^{(s)})^{-1} B_K^* \right]^{-1} = \overline{(\widehat{H}_{K,N}^{(s)})^{-1}}. \quad (\text{DNE9})$$

The map B_K evaluates each relation-annihilating functional on the actual columns Z_k . It is onto, because A_ℓ is invertible and Z_k has rank t . The first inverse-Gram formula is the minimum dual metric on that same evaluation target. Substitution of DNE8 yields

$$B_K (\mathcal{D}_N^{(s)})^{-1} B_K^* = Z_k^\top \overline{H_{Y,N}^{(s)}} \bar{Z}_k = \widehat{H}_{K,N}^{(s)},$$

which proves the final equality with every transpose in its original direction. Empty rank has determinant one. In particular the full fixed-column return is exactly

$$\widehat{F}_k^{(s)} = - \sum_N \sigma_N \log \det Q_{K^\vee,N}^{(s)}. \quad (\text{DNE10})$$

Its relation to the original $F_K^{(s)}$ retains both low quotient volumes and the full source-map error of FGC12–13 and FEX34–37; DNE10 removes none of those terms.

65.4 A finite source bound before the high-order derivative

Use the actual R, R', B, C_v and R_0 of ROG6 and RDS11, and put

$$\rho = \tanh(R_0/4), \quad C_{F,k,A} = \frac{2R^v}{|\mu_v|} e^{RR_0}. \quad (\text{DNE11})$$

For any $\mathcal{N} = \sum \nu_{ij} e^{\beta_{ij}z} \in \mathcal{N}_v$, RDS12 proves $|\mathcal{N}/E_A| \leq C_{F,k,A} \|\nu\|_1$ on the complete disk $|z| \leq R_0$. Apply the original Gamma generating identity directly to the functional $\Lambda_{\mathcal{N}/E_A}$:

$$\sum_{n \geq 0} \Lambda_{\mathcal{N}/E_A} (p_n((S' - c')/i)) \frac{z^n}{n!} = (1 + z^2)^{-s/4} e^{ic' \arctan z} \frac{\mathcal{N}(-i \arctan z)}{E_A(-i \arctan z)}. \quad (\text{DNE12})$$

The quotient at zero is removable. On $|z| = \rho$, $|\arctan z| \leq R_0/4$. Cauchy's coefficient estimate therefore gives the exact bound

$$\begin{aligned} \|\Lambda_{\mathcal{N}/E_A}\|_{(\mathbb{C}[S']_{\leq d}, \|\cdot\|_{s,c'})^\vee} &\leq \mathcal{B}_{d,s,k,A} \|\nu\|_1, \\ \mathcal{B}_{d,s,k,A}^2 &= \frac{C_{F,k,A}^2}{M_s} (1 - \rho^2)^{-s/2} e^{|c'|R_0/2} \sum_{n=0}^d \frac{n!}{(s/2)_n} \rho^{-2n}. \end{aligned} \quad (\text{DNE13})$$

Indeed the bound for the n th orthonormal coefficient is the square root of its displayed summand times the common factor and $\|\nu\|_1$; summing squared coefficients gives the dual norm. The full Gamma mass remains explicitly in this formula. No application of $\chi'(D)$ and no $q!$ factor is needed in this exact extension norm.

For a fully specified comparison reference, use Euclidean norm in the original upper exponential coefficient order ν_{ij} , with its actual quotient by $\text{im } C^\top$. Its positive Gram in the fixed DNE3 frame is

$$G_{\text{exp}} = Z_{\text{exp}}^* [I - C^\top (\bar{C} C^\top)^{-1} \bar{C}] Z_{\text{exp}}, \quad 0 \prec \mathcal{D}_N^{(s)} \preceq q \mathcal{B}_{d,s,k,A}^2 G_{\text{exp}}. \quad (\text{DNE14})$$

The reference is explicitly specified here for comparison; the actual weighted Gamma quotient remains DNE7. The inverse in G_{exp} exists by the full row rank of C . Its expression is the orthogonal-projection formula, and is positive on this quotient frame by DNE3. Every representative $\nu + C^\top \eta$ remains in $\mathcal{N}_v$ and changes $\mathcal{N}/E_A$ by precisely a lower exponential, by RDS7. Combining DNE6 and DNE13, and then using $\|w\|_1 \leq \sqrt{q} \|w\|_2$, gives

$$x^* \mathcal{D}_N^{(s)} x \leq \mathcal{B}_{d,s,k,A}^2 \inf_\eta \|Z_{\text{exp}} x + C^\top \eta\|_1^2 \leq q \mathcal{B}_{d,s,k,A}^2 x^* G_{\text{exp}} x.$$

This proves DNE14 without a bound on an omitted frame inverse.

At fixed actual period and quartet the finite constant satisfies

$$\log(1 + q\mathcal{B}_{d,s,k,A}^2) = O_{\text{actual}}(q), \quad s = 1, k, \quad d = N - v. \quad (\text{DNE15})$$

For detail, R_0, ρ, v, μ_v are fixed, $R = O(k)$, $|c'| = O(k)$ and $s \leq k$. Thus $\log C_{F,k,A} = O_{\text{actual}}(k + \log(k + 1))$. The original Gamma ratios obey $n!/(s/2)_n \leq 2\sqrt{n}$ for $n \geq 1$ and are one at $n = 0$, as proved by the product estimate in FEX10–11. Hence the sum in DNE13 is at most $(d + 1) \max\{1, 2\sqrt{d}\} \rho^{-2d}$. Since $d \leq 2q$, its logarithm is $O_{\text{actual}}(q)$. The remaining factors have logarithms $O_{\text{actual}}(k)$, with the original positive mass retained before estimation. These bounds prove DNE15.

Restriction of DNE14 to any specified plane, and minimization under any specified onto quotient map, preserve its scalar upper bound. In particular it applies to the actual map B_K with the full reference quotient $(B_K G_{\text{exp}}^{-1} B_K^*)^{-1}$. This is a finite norm comparison of the original metric with a completely retained reference form. The reference contains the full original Vandermonde and conductor data; neither its determinant nor the original return in DNE10 has been assigned a leading value by this upper bound.

66 Subleading positive exterior expansion of the original dual metric

Fix the original simple quartet, its full unit, a fixed actual five-orbit period and logarithm branch. Retain all coefficients of the degree-(8, 8) conductor A , all shifts b_{ab} and every original root. The coefficient formula PCL8 depends on this fixed period and quartet and is independent of k . Thus the entire symbol

$$E_A(w) = \sum_{a,b=0}^8 a_{ab} e^{b_{ab}w}, \quad v = \text{ord}_0 E_A, \quad \mu_v = E_A^{(v)}(0) \neq 0 \quad (\text{PEC1})$$

is the same function throughout the sequence $k = 4l + 1 \geq 9$. Keep $q = (k + 1)^2$, $q' = (k - 7)^2$, $\Delta = q - q' = 16k - 48$, $\mathbf{m} = \Delta - v$, $c' = k/2 - 4$, both source orders $s = 1, k$, and the original endpoints $N = q - 1, q, 2q - 1, 2q$. Write $d = N - v$. The letter w in this section is the analytic symbol variable. It does not replace a primary root, polynomial coordinate, period, or the absolute base τ .

Use the complete DNE frame Z_{exp} , numerators $\mathcal{N}_a$, dual Grams $\mathcal{D}_N^{(s)}$ and coefficient reference G_{exp} . Their literal definitions, including the complex-linear dual, are

$$\begin{aligned} F_a &= \mathcal{N}_a / E_A, & T_{N,s;na} &= \Lambda_{F_a}(\phi_{n,s,c'}), & V_{N,s;n,(i,j)} &= \phi_{n,s,c'}(\beta'_{ij}), \\ \mathcal{D}_N^{(s)} &= T_{N,s}^*[I - V_{N,s}(V_{N,s}^*V_{N,s})^{-1}V_{N,s}^*]T_{N,s}, \\ G_{\text{exp}} &= Z_{\text{exp}}^*[I - C^\top(\bar{C}C^\top)^{-1}\bar{C}]Z_{\text{exp}} > 0. \end{aligned} \quad (\text{PEC2})$$

Here C is the entire original conductor, and the exact Gamma orthonormal frame has $h_n = M_s n!(s/2)_n$, $M_s = (2\pi)^{s/2}$. DNE6–8 proves these metrics, retaining the full lower evaluation space and both conditioning factors. In particular G_{exp} is the minimum Euclidean norm of an original exponential coefficient vector modulo $\text{im } C^\top$; it is not substituted for $\mathcal{D}_N^{(s)}$.

66.1 The actual symbol-zero jet subspaces

On $|z| < 1$ use the branch of $\arctan z$ vanishing at zero and set

$$w(z) = -i \arctan z, \quad \mathcal{E}_A(z) = z^{-v} E_A(w(z)), \quad \mathcal{E}_A(0) = \frac{\mu_v(-i)^v}{v!} \neq 0. \quad (\text{PEC3})$$

The power series for $\arctan$ is holomorphic on this disk and gives $|\arctan z| \leq \text{arctanh } |z|$. Its derivative is $(1 + z^2)^{-1}$, so w' is nonzero there. Moreover w is injective: $e^{-2w(z)} = (1 + iz)/(1 - iz)$, and equality of two w values implies equality under this injective fractional linear function. The displayed removable value in PEC3 follows by substituting the first nonzero Taylor term of PEC1. Thus $\mathcal{E}_A$ is a nonzero holomorphic function on the entire open unit disk.

Call $0 < r < 1$ admissible when $E_A(w(z)) \neq 0$ for $|z| = r$. For every such r , let $\mathcal{Z}_r$ be the finite set of nonzero zeros ω of E_A in $w(\{|z| < r\})$, and write h_ω for their exact orders. Define

$$H_r = \sum_{\omega \in \mathcal{Z}_r} h_\omega, \quad m_r = \min_{|z|=r} |E_A(w(z))| > 0. \quad (\text{PEC4})$$

Finiteness follows from isolation of the zeros of $\mathcal{E}_A$ in a compact subdisk. Positivity of m_r follows from continuity and admissibility on the compact circle. Every compact subdisk contains only finitely many excluded radii. Consequently there are admissible radii arbitrarily close to one. These are properties of the specified symbol, with no assumed zero-free unit disk. Its zero count can also be written exactly as

$$H_r = \frac{1}{2\pi i} \int_{|z|=r} \frac{\mathcal{E}'_A(z)}{\mathcal{E}_A(z)} dz. \quad (\text{PEC5})$$

Indeed factor out the finitely many interior zeros with their multiplicities. The logarithmic derivative of the remaining nonvanishing factor integrates to zero on the disk, while each factor $(z - z_0)^h$ contributes h . The nonzero derivative of w preserves the orders h_ω . This proves PEC5 in the original symbol variables.

In the DNE coefficient frame, form the full jet matrix

$$(J_{r,k})_{(\omega,b),a} = \frac{\mathcal{N}_a^{(b)}(\omega)}{b!}, \quad 0 \leq b < h_\omega, \quad c_{r,k} = \text{rank } J_{r,k}, \quad \mathcal{H}_{r,k} = \ker J_{r,k} \subset \mathbb{C}^{\mathbf{m}}. \quad (\text{PEC6})$$

Its entries are the complete sums $\sum_{i,j} (Z_{\text{exp}})_{(i,j),a} \beta_{ij}^b e^{\beta_{ij}\omega} / b!$. Thus every original root, factorial and coefficient is retained. Its codimension is $c_{r,k} \leq \min\{\mathbf{m}, H_r\}$. Addition of an original coefficient vector $C^\top \eta$ changes its numerator by E_A times the corresponding lower exponential polynomial, by RDS7. That change has zero jets in PEC6. Therefore the cancellation subspace is a specified subspace of the original quotient $\mathcal{N}_v / (E_A \mathcal{E}_{k-8})$, independent of the choice of representative and of both N and s . In particular the minimum Euclidean representative defining G_{exp} stays in this subspace.

For $x \in \mathcal{H}_{r,k}$, all interior poles of $\mathcal{N}_x / E_A$ cancel, including zero by membership in $\mathcal{N}_v$. Admissibility and discreteness give a neighborhood of the closed r disk with no additional boundary pole. Thus $\mathcal{N}_x(w(z)) / E_A(w(z))$ is holomorphic on that neighborhood. This is an exact pole cancellation imposed by the actual numerator jets, not a removal of poles from the original symbol.

66.2 A complete finite Gamma bound on each cancellation subspace

Use the original root bound $R = |k/2| + k\sqrt{\delta^2 + \gamma^2}$ and put $a_r = \text{arctanh } r$. For an original exponential coefficient vector ν of a canceled numerator, the exact generating function DNE12 satisfies, on $|z| = r$,

$$\left| (1 + z^2)^{-s/4} e^{ic' \arctan z} \frac{\sum \nu_{ij} e^{\beta_{ij} w(z)}}{E_A(w(z))} \right| \leq \frac{(1 - r^2)^{-s/4}}{m_r} e^{(R+|c'|)a_r} \|\nu\|_1. \quad (\text{PEC7})$$

This follows term by term from $|w(z)| \leq a_r$ and the root bound, with the original exponential coefficient order. The holomorphic branch of $(1 + z^2)^{-s/4}$ is the one taking value one at zero; its modulus is $|1 + z^2|^{-s/4}$. No source mass has yet canceled. Cauchy's coefficient integral and $h_n = M_s n! / (s/2)_n$ therefore give

$$\|\Lambda_{\mathcal{N}/E_A}\|_{(\mathcal{P}_d, \|\cdot\|_{s,c'})^\vee}^2 \leq B_{r,N,s}^2 \|\nu\|_1^2, \quad B_{r,N,s}^2 = \frac{e^{2(R+|c'|)a_r} (1 - r^2)^{-s/2}}{M_s m_r^2} \sum_{n=0}^d \frac{n!}{(s/2)_n} r^{-2n}. \quad (\text{PEC8})$$

The squared dual norm is the sum of the squared values on the complete orthonormal frame of degrees zero through d . The exponential generating series gives an $n!$ before division by $\sqrt{h_n}$, yielding precisely the ratio $n! / (s/2)_n$ above. This verifies every term of PEC8.

Every allowed addition $E_A \mathcal{E}_{k-8}$ preserves cancellation. Minimize over these representatives as in DNE6 and use $\|\nu\|_1^2 \leq q \|\nu\|_2^2$. This proves the quadratic-form bound

$$x^* \mathcal{D}_N^{(s)} x \leq C_{r,N,s} x^* G_{\text{exp}} x \quad (x \in \mathcal{H}_{r,k}), \quad C_{r,N,s} = q B_{r,N,s}^2. \quad (\text{PEC9})$$

It uses the entire lower conditioning space, not a chosen extension. For each fixed admissible radius, E_A, m_r, a_r are independent of k . The product estimate $n! / (s/2)_n \leq 2\sqrt{n}$ for $n \geq 1$ and its value one at $n = 0$ give the explicit bound

$$C_{r,N,s} \leq \frac{q e^{2(R+|c'|)a_r} (1 - r^2)^{-s/2}}{M_s m_r^2} (d + 1) \max\{1, 2\sqrt{d}\} r^{-2d}. \quad (\text{PEC10})$$

Here $R + |c'| = O_{\delta,\gamma}(k)$, $s \leq k$ and $d \leq 2q$. Since $M_s > 1$, taking the positive part of the logarithm gives

$$\log^+ C_{r,N,s} \leq 2d \log(1/r) + O_{\text{actual},r}(k + \log q). \quad (\text{PEC11})$$

The finite expression PEC10 supplies the constants in this assertion; no asymptotic assumption on an unevaluated numerator is used.

66.3 The whole positive logarithmic spectrum

Let $\lambda_1 \geq \dots \geq \lambda_m > 0$ be the eigenvalues of the specified relative operator

$$T_N^{(s)} = G_{\text{exp}}^{-1/2} \mathcal{D}_N^{(s)} G_{\text{exp}}^{-1/2}. \quad (\text{PEC12})$$

The positive square root is unique by diagonalizing the positive Hermitian matrix. This congruence is the exact comparison of the two retained forms. DNE14 gives $\lambda_a \leq C_{0,N,s}$ for all a , where $C_{0,N,s} = q\mathcal{B}_{d,s,k,A}^2$ is its complete explicit constant, and DNE15 gives $\log^+ C_{0,N,s} = O_{\text{actual}}(q)$. PEC9 bounds the Rayleigh quotient on a subspace of codimension $c_{r,k}$. Hence at most $c_{r,k}$ eigenvalues can exceed $C_{r,N,s}$: otherwise the span of $c_{r,k} + 1$ corresponding orthogonal eigenvectors would intersect that subspace nontrivially, contradicting the bound there. Consequently

$$\begin{aligned} \mathcal{P}_{N,s} &:= \sum_{a=1}^m \log^+ \lambda_a \leq E_{r,N,s}, \\ E_{r,N,s} &= c_{r,k} \log^+ C_{0,N,s} + (m - c_{r,k}) \log^+ C_{r,N,s}, \quad \varepsilon_{N,s}^\vee = \inf_{r \text{ admissible}} E_{r,N,s}. \end{aligned} \quad (\text{PEC13})$$

All terms are finite and nonnegative, and the admissible set is nonempty. Thus $\mathcal{P}_{N,s} \leq \varepsilon_{N,s}^\vee$ even when the infimum is not attained. For every fixed admissible r , $c_{r,k} \leq H_r$ is bounded independently of k . Since $m/k \rightarrow 16$ and $d/q \leq 2$, PEC11 and DNE15 imply

$$0 \leq \limsup_{k \rightarrow \infty} \frac{\varepsilon_{N,s}^\vee}{kq} \leq 64 \log(1/r). \quad (\text{PEC14})$$

This holds simultaneously for the two sources and four indicated endpoint sequences, which form a finite list. Choose admissible radii arbitrarily close to one only after taking this bound for each fixed radius. The right side tends to zero. We have proved

$$\boxed{\mathcal{P}_{N,s} \leq \varepsilon_{N,s}^\vee = o(kq)}. \quad (\text{PEC15})$$

Only the original source orders $1, k$ are used in the two limiting steps; the actual period and operator remain fixed. No additional source order or presumed spectral gap is inserted. Although H_r can increase as r tends to one, it is finite before each k limit, which is exactly what PEC14 uses.

This also controls every exterior degree of the comparison map. The singular values squared of the identity from $(\mathbb{C}^m, G_{\text{exp}})$ to $(\mathbb{C}^m, \mathcal{D}_N^{(s)})$ are the λ_a . In orthonormal wedge frames, each squared singular value is a product over a subset of these numbers. Therefore

$$\log \left\| \bigwedge^t \text{id} \right\| \leq \frac{1}{2} \mathcal{P}_{N,s} \leq \frac{1}{2} \varepsilon_{N,s}^\vee \quad (0 \leq t \leq m). \quad (\text{PEC16})$$

This is an all-degree bound in the exact two metrics. It does not bound the inverse exterior map; the small relative eigenvalues are retained below.

66.4 Every fixed restriction and quotient keeps this bound

For an injective matrix J and an onto matrix B , retain respectively the full paired metrics

$$\begin{aligned} D_J &= J^* \mathcal{D}_N^{(s)} J, \quad G_J = J^* G_{\text{exp}} J, \\ D_B &= [B(\mathcal{D}_N^{(s)})^{-1} B^*]^{-1}, \quad G_B = [B G_{\text{exp}}^{-1} B^*]^{-1}. \end{aligned} \quad (\text{PEC17})$$

Their relative eigenvalues $\mu_1 \geq \dots \geq \mu_t > 0$ satisfy $\mu_a \leq \lambda_a$ for $1 \leq a \leq t$. For a restriction, write $U = G_{\text{exp}}^{1/2} J G_J^{-1/2}$, so $U^* U = I$ and its relative operator is $U^* T_N^{(s)} U$. The variational description of its a th largest eigenvalue gives the asserted upper bound. Explicitly the preimage of any subspace

of codimension $a - 1$ has codimension at most $a - 1$, so the usual minimum of maximal Rayleigh quotients on such subspaces cannot increase under restriction. This variational formula itself follows by expanding vectors in an orthonormal eigenbasis: the complement of the first $a - 1$ vectors gives the upper bound, and dimension intersection with their first a vectors gives the lower bound.

For the quotient put $V = (BG_{\text{exp}}^{-1}B^*)^{-1/2}BG_{\text{exp}}^{-1/2}$. Then $VV^* = I$ and its relative operator is $(V(T_N^{(s)})^{-1}V^*)^{-1}$. The a th smallest eigenvalue of the compression of $(T_N^{(s)})^{-1}$ is at least its a th smallest full eigenvalue, by the same variational argument applied to smallest eigenvalues. Taking reciprocals proves $\mu_a \leq \lambda_a$. It follows that for either metric pair in PEC17,

$$\sum_{a=1}^t \log^+ \mu_a \leq \mathcal{P}_{N,s} \leq \varepsilon_{N,s}^\vee. \quad (\text{PEC18})$$

No bound for a coefficient inverse of B or J occurs: their full reference metrics in PEC17 remain part of the assertion.

66.5 The exact negative spectral return on the actual kernel

Retain Z_k , A_ℓ and the actual onto matrix $B_K = Z_k^\top A_\ell$ of DNE9. Put

$$D_{K,N} = [B_K(\mathcal{D}_N^{(s)})^{-1}B_K^*]^{-1} = \overline{(\widehat{H}_{K,N}^{(s)})}^{-1}, \quad G_K = [B_K G_{\text{exp}}^{-1}B_K^*]^{-1}. \quad (\text{PEC19})$$

These are exactly the native original kernel dual and its fixed reference, with no changed relation or factorial column. For this pair define, using all its positive relative eigenvalues $\mu_{a,N}$,

$$p_{K,N} = \sum_a \log^+ \mu_{a,N}, \quad n_{K,N} = \sum_a \log^+(\mu_{a,N}^{-1}), \quad A_{K,k,s} = n_{K,q-1} + n_{K,q} - n_{K,2q-1} - n_{K,2q}. \quad (\text{PEC20})$$

The original weighted polynomial spaces are nested and the complete relation maps restrict to the earlier cutoffs. Minimizing a primal norm over their increasing lift sets decreases $H_{Y,N}^{(s)}$. By DNE8 this increases $\mathcal{D}_N^{(s)}$. Taking minima over the same B_K fibres preserves this increase for $D_{K,N}$. Relative eigenvalues in its fixed G_K metric therefore increase by the variational formula. Hence $p_{K,N}$ increases, $n_{K,N}$ decreases, and $A_{K,k,s} \geq 0$.

Since $\log \mu = \log^+ \mu - \log^+(1/\mu)$, the complete determinant formula DNE10 yields the exact identity and finite bound

$$\begin{aligned} \widehat{F}_k^{(s)} &= A_{K,k,s} + P_{K,k,s}, \\ P_{K,k,s} &= p_{K,2q-1} + p_{K,2q} - p_{K,q-1} - p_{K,q}, \quad 0 \leq P_{K,k,s} \leq E_{k,s}^\vee := \varepsilon_{2q-1,s}^\vee + \varepsilon_{2q,s}^\vee = o(kq). \end{aligned} \quad (\text{PEC21})$$

The reference determinant cancels with the four original signs only after PEC19's complete dual identity has been used. Formula PEC18 bounds each of the two high positive parts, proving the last inequality. Thus the entire leading native kernel return is carried by the specified decrease of its small relative eigenvalue deficits; none has been set to zero or inferred from the rank.

Retain the complete unequal D_s^- , D_s^+ and original $\mathcal{L}_s$ from FEX32–35, with $t_0 = \dim(K \cap L)$ and $L = \mathbb{C}[S]_{<v}$. Substituting PEC21 into that proved interval gives

$$\begin{aligned} A_{K,k,s} - D_s^+ - (v - t_0)\mathcal{L}_s &\leq F_K^{(s)} \\ &\leq A_{K,k,s} + E_{k,s}^\vee - D_s^- + v\mathcal{L}_s. \end{aligned} \quad (\text{PEC22})$$

Both original low corrections and both unequal flag errors are retained. FEX37 separately gives the useful exact symmetric allowance

$$|F_K^{(s)} - A_{K,k,s}| \leq \mathcal{E}_{k,s}^{\text{FEX}} + E_{k,s}^\vee = o(kq), \quad (\text{PEC23})$$

where $\mathcal{E}_{k,s}^{\text{FEX}}$ is the full finite FEX36 expression. This is a proved original-kernel metric reduction, not an evaluated individual leading coefficient.

66.6 The complementary original residual and its low subspace

Use the fixed onto map $P_Y : \mathbb{C}^{\mathfrak{m}} \rightarrow \mathbb{C}^{j_0}$ with kernel $\text{im } Z_k$, where $j_0 = \mathfrak{m} - (r - t_0) = j - v + t_0$ and $j = 8k - 32$. For an explicit choice, let $\mathcal{I}$ be the first set of $r - t_0$ row indices in the fixed ordering for which $E_{\mathcal{I}}^{\top} Z_k$ is invertible, and let $\mathcal{J}$ be the complementary indices. Define

$$P_Y = E_{\mathcal{J}}^{\top} - E_{\mathcal{J}}^{\top} Z_k (E_{\mathcal{I}}^{\top} Z_k)^{-1} E_{\mathcal{I}}^{\top}.$$

Independence of the columns, proved in FGC10, ensures such a minor exists. Multiplication gives $P_Y Z_k = 0$ and $P_Y E_{\mathcal{J}} = I$. Its rank is j_0 and its kernel has dimension $r - t_0$, so that kernel is exactly $\text{im } Z_k$. Empty rank uses the identity quotient. Set

$$J_0 = A_{\ell}^{-1} P_Y^{\top}, \quad D_{0,N} = J_0^* \mathcal{D}_N^{(s)} J_0, \quad G_0 = J_0^* G_{\text{exp}} J_0, \quad Q_{0,N} = [P_Y (H_{Y,N}^{(s)})^{-1} P_Y^*]^{-1}. \quad (\text{PEC24})$$

Then $B_K J_0 = 0$ and J_0 is a basis of $\ker B_K$ by dimensions. Direct substitution in DNE8 proves the complete dual identity

$$D_{0,N} = \bar{P}_Y \overline{(H_{Y,N}^{(s)})^{-1} P_Y^{\top}} = \overline{Q_{0,N}^{-1}}. \quad (\text{PEC25})$$

Thus PEC24 is the annihilator of the actual kernel columns, with its full metric, not a complementary coordinate plane chosen for convenience.

Define $p_{0,N}, n_{0,N}$ from the relative eigenvalues of $(D_{0,N}, G_0)$ exactly as in PEC20, and $A_{0,k,s}, P_{0,k,s}$ by the same four endpoint differences. Restriction in PEC18 and the preceding monotonicity proof give

$$F_{0,k,s} := \sum_N \sigma_N \log \det Q_{0,N} = A_{0,k,s} + P_{0,k,s}, \quad A_{0,k,s} \geq 0, \quad 0 \leq P_{0,k,s} \leq E_{k,s}^{\vee}. \quad (\text{PEC26})$$

The scalar $F_{0,k,s}$ here denotes this displayed quotient return only; it is not the seven-moment quantity $\mathcal{F}_{0,k}^{(s)}$.

To identify the original target exactly, put $W_0 = \Xi(L) \subset B_{\Xi}$. The map $L \rightarrow W_0$ has kernel $K \cap L$, so $\dim W_0 = v - t_0$. There is the onto map, in the original FGC coordinates,

$$V_A \longrightarrow Y_N / \text{im } Z_k, \quad [p] \longmapsto \left[\frac{\alpha_A \mathcal{T}_{Ap}}{\chi'} \right], \quad \alpha_A = v! / \mu_v, \quad Y_N / \text{im } Z_k \simeq B_{\Xi} / W_0. \quad (\text{PEC27})$$

FGC4–10 proves independence of representatives and surjectivity. Its kernel is $K + L$: a zero image subtracts a member of K with that image in Y_N , leaving an element of L by FGC6. Conversely both K and L map to zero. The induced isomorphism with B_{Ξ} / W_0 sends the class of p to the class of $\Xi[p]$. This verifies both directions of the target identification.

At each endpoint the source map is the original S_N of CEP28 followed by $D_{\alpha,N}(x, f) = (x, \alpha_A f)$ on GIP's full graph. The complete quotient map PEC27 kills its entire relation graph and kernel columns. In the product source norm, minimization first over the unrestricted low coordinate and then over the polynomial and Z_k relations is exactly $Q_{0,N}$. The exterior norms of $D_{\alpha,N}^{\pm 1}$ in these same product metrics are at most $\exp(j_0 |\log |\alpha_A||)$ in degree j_0 . The minimum-lift and projection argument of GIP6 applied to this composite source map therefore proves, for the original induced metric $\bar{Q}_{\Xi,N}$ on B_{Ξ} / W_0 in the transported frame,

$$|\log \det \bar{Q}_{\Xi,N} - \log \det Q_{0,N}| \leq 2E_N^{\text{CEP}} + 2j_0 |\log |\alpha_A||. \quad (\text{PEC28})$$

Orthogonal projection off the entire kernel is an exterior contraction, and the inverse composite gives the other direction. This checks the full source, target, metric and rank in the comparison.

Choose a fixed original frame I_{W_0} of W_0 and retain its exact restriction return $F_{W_0}^{(s)} = \sum_N \sigma_N \log \det (I_{W_0}^* Q_{\Xi,N}^{\text{fix}} I_{W_0})$. Completing that frame by fixed quotient lifts gives a block Gram

whose Schur complement is $\overline{Q}_{\Xi,N}$. Its fixed frame determinant cancels with the four signs. Consequently the original residual is

$$\begin{aligned}\Phi_{k,s} &= F_{W_0}^{(s)} + A_{0,k,s} + P_{0,k,s} + \delta_{0,k,s}, \\ |\delta_{0,k,s}| &\leq E_{k,s}^{\text{tr}} := 2 \sum_N E_N^{\text{CEP}} + 8j_0 |\log |\alpha_A|| = o(kq), \\ 0 &\leq F_{W_0}^{(s)} \leq (v - t_0)\mathcal{L}_s = O_{\text{actual}}(q).\end{aligned}\tag{PEC29}$$

The last inequality follows by restricting and minimizing the original AKS covariance comparison on this fixed subquotient of the primary space, exactly as in FEX32. Its metrics decrease on increasing source spaces, proving nonnegativity. The full finite $\mathcal{L}_s$ is FEX33, whose proved order is $O(q)$, and $v \leq 80$. Thus no original low contribution has been dropped. In particular the finite and asymptotic consequences are

$$\begin{aligned}A_{0,k,s} - E_{k,s}^{\text{tr}} &\leq \Phi_{k,s} \leq A_{0,k,s} + E_{k,s}^{\vee} + E_{k,s}^{\text{tr}} + (v - t_0)\mathcal{L}_s, \\ \Phi_{k,s} &= A_{0,k,s} + o(kq).\end{aligned}\tag{PEC30}$$

66.7 Backward and forward return to the same arithmetic class

The complete OVR/ECL identity $F_K^{(s)} + \Phi_{k,s} = \Delta q \mathfrak{C}_{\partial} + o(kq)$ and PEC23, PEC30 give

$$\begin{aligned}A_{K,k,s} + A_{0,k,s} &= \Delta q \mathfrak{C}_{\partial} + o(kq), \\ \frac{A_{K,k,s} + A_{0,k,s}}{kq} &\longrightarrow 16\mathfrak{C}_{\partial}, \\ 21.66851180520 &< 16\mathfrak{C}_{\partial} < 21.66851180521.\end{aligned}\tag{PEC31}$$

Both deficits are on the exact DNE dual quotient and its actual kernel annihilator, with the full G_{exp} references. The equality evaluates their sum and retains their individual distributions. AMT's already proved source-order agreement and PEC23, PEC30 also give $(A_{K,k,1} - A_{K,k,k})/(kq) \rightarrow 0$ and the same statement for A_0 .

For finite original kernel bounds intersect PEC22 with OER19. Retain the full AMT arithmetic intervals and take their exact boundary complements. On the original arithmetic period domain $|u| \geq R_*$, the signed identity $\Psi_k = F_K^{(k)} + \mathcal{H}_k + \delta_k^{\sigma}$ then proves, with unchanged signs,

$$\begin{aligned}A_{K,k,k} - D_k^+ - (v - t_0)\mathcal{L}_k + \mathcal{H}_k + \delta_k^{\sigma} &\leq \Psi_k \\ &\leq A_{K,k,k} + E_{k,k}^{\vee} - D_k^- + v\mathcal{L}_k + \mathcal{H}_k + \delta_k^{\sigma}.\end{aligned}\tag{PEC32}$$

Here $D_k^{\pm}$ means the original FEX bounds at source $s = k$, and $\mathcal{L}_k$ is the same source's FEX33 allowance. The original same-class action inequality OER20 uses precisely the right endpoint of PEC32 plus the complete C_k^{budget} . This addition preserves the original monic window, contraction penalties, mixed pairings, action leakage and positive-primary exterior class. Where the seven moments are used, retain PMR's finite range and $e^{0,\pm} = e^{\pm} + c_k^{\circ}$; outside that range retain the original OAR outer expression.

The calculations PEC1–32 prove a subleading positive exterior budget and return it to the original kernel and residual, whose leading negative spectral deficits are explicitly specified. Calculating these actual small eigenvalues and the resulting same-class exclusion bound is the remaining analytic step in these formulas. No assertion about their individual leading values or an RH conclusion follows merely from the evaluated sum PEC31.

67 The actual common-factor map on the original kernel

Fix the original simple quartet, its complete unit and period data, and the original logarithm branch on the proved five-orbit domain. Use $k = 4l + 1 \geq 9$, $k' = k - 8$, and retain

$$\begin{aligned} q &= (k+1)^2, & q' &= (k-7)^2, & \Delta &= q - q' = 16k - 48, \\ r &= 8k - 16, & j &= 8k - 32, & c &= k/2, & c' &= k'/2. \end{aligned} \quad (\text{ACR1})$$

The original primary points, also at every integer index in the second formula, are

$$\begin{aligned} \beta_{ab} &= c + (2a - k)\delta + i(2b - k)\gamma, & 0 \leq a, b \leq k, \\ \tau_{ij} &= c' + (2i - k')\delta + i(2j - k')\gamma, & (i, j) \in \mathbb{Z}^2, \\ b_{ab} &= 4 + (2a - 8)\delta + i(2b - 8)\gamma, & 0 \leq a, b \leq 8. \end{aligned} \quad (\text{ACR2})$$

In the imaginary terms i is the imaginary unit; subscripts are integer indices. The functions of the two integer indices are injective, by comparison of real and imaginary parts and $\delta > 0$, $\gamma > 2$. In particular $\tau_{ij} + b_{ab} = \beta_{i+a, j+b}$ whenever the latter indices are in their displayed range.

Write $\chi = \prod_{a,b=0}^k (S - \beta_{ab})$ and $\chi' = \prod_{i,j=0}^{k'} (S' - \tau_{ij})$. The prime names the original lower-grid polynomial, and is not differentiation. All dual pairings until a Hermitian Gram is explicitly mentioned are complex-linear.

67.1 Support and its precise extended coefficient rectangle

Let the actual PCL8–11 biform be

$$A = \sum_{a,b=0}^8 a_{ab} z_0^a z_1^{8-a} w_0^b w_1^{8-b} \neq 0, \quad \mathcal{S} = \{(a, b) : a_{ab} \neq 0\}. \quad (\text{ACR3})$$

Here, literally, its factors are $Q_0(L_h(u^{-1})t)$, $1 \leq h \leq 4$, where $t = (z_0 w_0, z_0 w_1, z_1 w_0, z_1 w_1)^T$ and $L_h = U^{-1} V \mathcal{R}^{-1} D^{-h} \mathcal{R} V^{-1} U$. Thus every coefficient is the complete finite sum of products in PCL10, with the entire original matrix series PCL2. No coefficient is assigned in this calculation. Put

$$a_- = \min_{\mathcal{S}} a, \quad a_+ = \max_{\mathcal{S}} a, \quad b_- = \min_{\mathcal{S}} b, \quad b_+ = \max_{\mathcal{S}} b, \quad d_a = a_+ - a_-, \quad d_b = b_+ - b_-. \quad (\text{ACR4})$$

The following finite index sets have their literal integer intervals:

$$\begin{aligned} I_{\text{all}} &= [-a_-, k - a_+] \times [-b_-, k - b_+], & I_{\text{in}} &= [0, k']^2, & I_G &= I_{\text{all}} \setminus I_{\text{in}}, \\ G_A(S') &= \prod_{(i,j) \in I_G} (S' - \tau_{ij}), & g &= |I_G| = (k+1-d_a)(k+1-d_b) - q'. \end{aligned} \quad (\text{ACR5})$$

Every interval is nonempty and the first rectangle contains I_{in} , since all active indices are in $[0, 8]^2$. The intersection of the shifted full-grid rectangles $[-a, k-a] \times [-b, k-b]$, over the actual active pairs, is exactly I_{all} . Each full shifted polynomial $\chi(S' + b_{ab})/\chi'(S')$ has the simple roots in its rectangle with I_{in} removed. Distinctness in ACR2 therefore proves that G_A is precisely their monic greatest common divisor, with the degree in ACR5. This reproves the RCF common factor from the actual support. At $g = 0$ its polynomial is the empty product one.

Use the original ordered biform coefficient space $\mathcal{B}_k$, with basis $e_{ab}^{(k)} = z_0^a z_1^{k-a} w_0^b w_1^{k-b}$. For each $(i, j) \in I_{\text{all}}$ define the complete column

$$f_{ij} = \sum_{(a,b) \in \mathcal{S}} a_{ab} e_{i+a, j+b}^{(k)}. \quad (\text{ACR6})$$

All indices in this expression lie in $[0, k]^2$: the lower bound follows from $i \geq -a_-$, $j \geq -b_-$; the upper bound follows from $i \leq k - a_+$, $j \leq k - b_+$. Let C_{in} , C_G and C_{all} have these columns for I_{in} ,

I_G and I_{all} , in fixed lexicographic orders. These coefficient matrices use every active coefficient, each once, and no conjugate or binomial multiplier.

The full boundary factor and its remaining biform are

$$\begin{aligned} M_{\partial} &= z_0^{a-} z_1^{8-a+} w_0^{b-} w_1^{8-b+}, & A &= M_{\partial} A_{\text{red}}, \\ \deg A_{\text{red}} &= (d_a, d_b), & A_{\text{red}} &= \sum_{a,b} a_{ab} z_0^{a-a-} z_1^{a_+-a} w_0^{b-b-} w_1^{b_+-b}. \end{aligned} \quad (\text{ACR7})$$

The formula retains the full scalar and every coefficient of A ; only the displayed coordinate monomial is factored. For bidegrees (p, t) let $\mathcal{B}_{p,t}$ be the complete biform space of those bidegrees and set it to zero if either degree is negative. Multiplication by a nonzero biform is injective, since the ambient polynomial ring is an integral domain. ACR6 and ACR7 then give the exact coefficient images

$$\begin{aligned} H_k &:= \text{im } C_{\text{all}} = A_{\text{red}} \mathcal{B}_{k-d_a, k-d_b}, & \text{im } C_{\text{in}} &= A \mathcal{B}_{k-8, k-8}, \\ \text{rank } C_{\text{all}} &= q' + g, & \text{rank } C_{\text{in}} &= q', & H_k &= \text{im } C_{\text{in}} \oplus \text{im } C_G. \end{aligned} \quad (\text{ACR8})$$

Indeed f_{ij} is A_{red} times the monomial of bidegree $(k-d_a, k-d_b)$ whose two indices are $(i+a_-, j+b_-)$. As (i, j) ranges over I_{all} , these are all its basis monomials. Their products are independent by injectivity, proving every image and rank in ACR8. For the inner subset the same multiplication is exactly A times its original degree- $(k-8, k-8)$ basis.

67.2 The residual map evaluated on every original primary coordinate

Let $E = \mathbb{C}[S]/(\chi)$ and let $\mathcal{V}_{\beta} : E \rightarrow \mathbb{C}^{[0,k]^2}$ be the original primary evaluation map. Define the original conductor annihilator V_A and the original kernel K by

$$V_A = \{[p] : C_{\text{in}}^T \mathcal{V}_{\beta}[p] = 0\}, \quad K = \ker \Lambda \subseteq V_A. \quad (\text{ACR9})$$

The inclusion follows from the full original trace identity $A \mathcal{B}_{k-8, k-8} \subseteq J_k$, where J_k is the image of the original invariant algebra under $x = Ht$; the complex-linear annihilator of J_k is K by OQC8. Equivalently, direct evaluation proves

$$\mathcal{T}Ap(S') = \sum_{a,b} a_{ab} p(S' + b_{ab}), \quad (C_{\text{in}}^T \mathcal{V}_{\beta} p)_{ij} = (\mathcal{T}Ap)(\tau_{ij}), \quad [p] \in V_A \iff \chi' \mid \mathcal{T}Ap. \quad (\text{ACR10})$$

The roots of χ' are distinct, so vanishing at them is equivalent to its polynomial divisibility. Replacing p by $p + \chi h$ preserves this property, because $\chi(S' + b_{ab}) = \chi'(S') d_{ab}(S')$ for the original monic shifted factors. Thus ACR9 and ACR10 agree on the original quotient, with all original positive shifts.

For every $\ell = (i, j) \in I_G$ one has $\chi'(\tau_{\ell}) \neq 0$. The full common-factor residual has the exact evaluation

$$\boxed{(\rho_G[p])(\tau_{\ell}) = \frac{1}{\chi'(\tau_{\ell})} \sum_{a,b} a_{ab} p(\beta_{i+a, j+b}) = \frac{f_{\ell}^T \mathcal{V}_{\beta}[p]}{\chi'(\tau_{\ell})}.} \quad (\text{ACR11})$$

Here $\rho_G[p] = \text{rem}_{G_A}(\mathcal{T}Ap/\chi')$ is in the original remainder coefficient frame. Divisibility ACR10 gives a polynomial quotient. Evaluation at a simple root of G_A commutes with its monic remainder and gives the first expression. The point identity ACR2 proves the second. A representative change adds $\sum a_{ab} d_{ab} h(S' + b_{ab})$, divisible by the full G_A , because it divides every active d_{ab} . This proves the stated map on V_A , independently of representatives, and re-establishes the exact RCF15 map.

Let $W_G = (\tau_{\ell}^d)_{\ell \in I_G, 0 \leq d < g}$ be the original ordered Vandermonde and $D_G = \text{diag}(\chi'(\tau_{\ell})^{-1})$. For $g > 0$, both are invertible, since all their indicated differences and diagonal entries are nonzero. The same notation has its unique empty meaning at $g = 0$. Let $\pi : \mathcal{B}_k \twoheadrightarrow \mathbb{C}^r$ be any exact OCF coefficient quotient with $\ker \pi = J_k$. OKM proves that the original kernel frame is $I_K = \mathcal{V}_{\beta}^{-1} \pi^T$. Therefore

$$\boxed{\rho_G I_K = W_G^{-1} D_G C_G^T \pi^T, \quad B_G := \pi C_G \in \text{Mat}_{r \times g}(\mathbb{C}), \quad t_G = g - \text{rank } B_G.} \quad (\text{ACR12})$$

The first equality follows by applying ACR11 on every kernel column and then the specified inverse evaluation map. The matrix $C_G^T \pi^T = B_G^T$ uses ordinary transpose. Multiplication by the two invertible target maps does not change rank, proving the last equality with the definition in RCM22. Every entry of the actual matrix needed for this rank is explicitly

$$(B_G)_{h,(i,j)} = \sum_{(a,b) \in S} a_{ab} \pi_{h,(i+a,j+b)}. \quad (\text{ACR13})$$

Thus this is a matrix with at most 81 terms per entry, $r = 8k - 16$ rows and $g \leq 16k - 48$ columns, computed from the fixed original period coefficients and original kernel quotient. It requires no full symmetric-power matrix.

For an exact all-case scalar prescription put $\Delta_0(B_G) = 1$, and let $\Delta_h(B_G)$ be the sum of the squared absolute values of all its h by h minors. Then

$$t_G = g - \max\{h : \Delta_h(B_G) > 0\}. \quad (\text{ACR14})$$

A nonzero minor gives that many independent columns; conversely Gaussian elimination on independent columns gives that many pivot rows and a nonzero minor. This proves the equality with rank. A sum of nonnegative real terms vanishes exactly when all those minors vanish. All a_{ab} and the full OCF pivot denominators remain in those minors. Formula ACR14 neither assigns generic rank nor supplies a numerical zero test for an unevaluated analytic coefficient.

67.3 The exact retained residual quotient in biform coordinates

The independent columns of C_G stay independent modulo $\text{im } C_{\text{in}}$ by ACR8. Hence the induced map $\mathbb{C}^g \rightarrow V_A^\vee$ represented by C_G is injective. Its transpose is onto, and the invertible factors in ACR11 prove that $\rho_G : V_A \rightarrow \mathbb{C}[S']_{<g}$ is onto. This proves its surjectivity directly from the original coefficient map.

The annihilator calculation also gives

$$\begin{aligned} V_0 &:= \ker \rho_G = H_k^\perp \subseteq V_A, & t_G &= \dim \frac{J_k \cap H_k}{A\mathcal{B}_{k-8,k-8}}, \\ J_k^{\text{red}} &:= \text{im}(J_k \rightarrow \mathcal{B}_k/H_k), & t_G &= j - \dim J_k^{\text{red}}. \end{aligned} \quad (\text{ACR15})$$

The first statement follows from the simultaneous vanishing of the inner and exterior rows in ACR11 and ACR8. The map B_G in ACR12 has kernel precisely those exterior linear combinations belonging to J_k . The direct sum in ACR8 and the inclusion $A\mathcal{B}_{k-8,k-8} \subseteq J_k$ identify this kernel with the quotient in the first line. Finally $\dim J_k = q - r$, while the inner space has dimension q' . Applying rank-nullity to $J_k \rightarrow J_k^{\text{red}}$ gives $(q - r) - \dim J_k^{\text{red}} - q' = j - \dim J_k^{\text{red}}$, proving the second line.

This is the same invariant image as FMC7: its map is induced by the identical original substitution $x = Ht$, and the inner quotient is the same $A\mathcal{B}$ quotient. Factoring the explicit boundary monomial in ACR7 provides the additional quotient map to $\mathcal{B}/H$, with its kernel computed in ACR15; it does not replace that module by a newly chosen invariant space.

For the original residual observation $\Xi : V_A \rightarrow B_\Xi$ with kernel K , put $W = \Xi(V_0)$. Define

$$A_G : \mathbb{C}[S']_{<g} \rightarrow B_\Xi/W, \quad A_G(\rho_G x) = [\Xi x]. \quad (\text{ACR16})$$

Two lifts differ by V_0 , so the map is well defined; it is onto since both maps from V_A are onto. Its kernel is exactly $\rho_G K$: if $\Xi x \in W$, choose $y \in V_0$ with $\Xi y = \Xi x$; then $x - y \in K$ and has the same ρ_G value. The reverse containment is direct. Consequently $\dim W = \dim J_k^{\text{red}}$, while $\dim(B_\Xi/W) = t_G$. In particular

$$\begin{aligned} d_a = d_b = 8 : & \quad g = 0, \quad t_G = 0, \quad V_0 = V_A, \quad W = B_\Xi; \\ d_a = d_b = 0 : & \quad g = \Delta, \quad H_k = \mathcal{B}_k, \quad t_G = j, \quad W = 0. \end{aligned} \quad (\text{ACR17})$$

The first row follows from ACR5 and the zero target of ρ_G ; its full residual still has rank j . In the second row A_{red} is a nonzero constant, so the quotient defining J_k^{red} is zero and ACR15 gives the result. This second row classifies that exact coefficient case; it does not assert that a given original period attains it. Every intermediate case retains the actual matrices ACR12–15 and the full original quotient ACR16.

67.4 A smaller invariant recursion evaluating every exceptional rank

Here is a second exact computation of the remaining rank which never uses the full symmetric power. In every diagonal degree $d \geq 0$ set

$$H_d = A_{\text{red}} \mathcal{B}_{d-d_a, d-d_b}, \quad \mathcal{W}_d = \mathcal{B}_{d,d}/H_d, \quad n_d = (d+1)^2 - \max(d-d_a+1, 0) \max(d-d_b+1, 0). \quad (\text{ACR18})$$

Multiplication by A_{red} is injective, so this is the exact dimension of the quotient. Also $H_d \mathcal{B}_{e,e} \subseteq H_{d+e}$; therefore the direct sum of these quotients is an actual graded algebra.

Choose the lexicographically largest nonzero coefficient a_* of the complete A_{red} , at the biform indices (r_*, s_*) . In the full degree- d grid eliminate the pivot rectangle

$$\{(r_* + i, s_* + h) : 0 \leq i \leq d - d_a, 0 \leq h \leq d - d_b\} \quad (\text{ACR19})$$

in descending lexicographic order, subtracting at each pivot its current coefficient divided by a_* times the corresponding monomial multiple of A_{red} . When a degree bound is negative the rectangle is empty. Every other term in a subtracted polynomial is strictly smaller in the additive lexicographic order. Earlier zero pivots therefore stay zero. The result is a linear map N_d^{red} to the complementary coefficient strip, with its literal zero-insertion section S_d^{red} . Summing the subtractions gives a decomposition into the strip and H_d . Their intersection is zero: the leading term of every nonzero $A_{\text{red}}b$ lies in the pivot rectangle. Thus

$$\ker N_d^{\text{red}} = H_d, \quad N_d^{\text{red}} S_d^{\text{red}} = I_{n_d}. \quad (\text{ACR20})$$

All entries are finite rational expressions in the original a_{ab} with powers of the selected actual a_* in their denominators. Selecting the first nonzero coefficient covers all support and leading-coefficient cases. If A_{red} is constant, this construction eliminates the full grid, as it must.

Use the following fourteen actual invariant monomials:

e	$\mathcal{G}_e$
2	x_1x_4, x_2x_3
3	$x_1^2x_3, x_1x_2^2, x_2x_4^2, x_3^2x_4$
4	$x_1^3x_2, x_1x_3^3, x_2^3x_4, x_3x_4^3$
5	$x_1^5, x_2^5, x_3^5, x_4^5$

(ACR21)

These generate every invariant polynomial, as follows directly from the original weights 1, 2, 3, 4 modulo five. An indecomposable invariant monomial has degree at most five: among six partial sums of any five factor weights, two agree and supply an invariant divisor. An opposite pair of weights gives one of the displayed quadratics; an indecomposable containing it is that quadratic. In the absence of opposite pairs the support has at most two variables. One variable gives a fifth power. The remaining supports are $\{1, 2\}, \{1, 3\}, \{2, 4\}, \{3, 4\}$; the positive exponent solutions of total degree at most five are respectively $(3, 1), (1, 2)$ for the first support and $(2, 1), (1, 3)$ for the second, with the last two obtained by multiplying weights by 2^{-1} and 3^{-1} modulo five. These are precisely the displayed cubics and quartics. Repeated factorization into indecomposables terminates by decreasing total degree, proving the generator assertion.

For $f \in \mathcal{G}_e$, retain its complete biform $\tilde{f} = f(Ht)$, whose coefficients are the OCF13 finite multinomial expressions in the actual period and unit matrix. Define

$$m_{f,d}^{\text{red}} = N_d^{\text{red}} M_{\tilde{f}} S_{d-e}^{\text{red}}, \quad J_0^{\text{red}} = \text{span}\{N_0^{\text{red}} 1\}, \quad J_d^{\text{red}} = \sum_{\substack{f \in \mathcal{G}_e \\ e \leq d}} m_{f,d}^{\text{red}} J_{d-e}^{\text{red}}. \quad (\text{ACR22})$$

Negative degrees give zero spaces. Multiplying two representatives differing by H_{d-e} changes their product by H_d , so ACR20 proves representative independence of the displayed multiplication. Every invariant monomial of positive degree factors by ACR21 into one generator times an earlier invariant monomial. This proves both inclusions in the recursion and hence its equality with the original invariant image in every degree. The base case includes the zero class of one when the quotient algebra is zero.

To evaluate its rank, concatenate the indicated multiplication matrices applied to bases of the preceding invariant images, and choose independent columns by their exact minors. Its rank is the maximum size of a nonzero minor, with rank zero for the empty image. The selected columns supply the next basis with their full original coefficient expressions. This proves the algorithm by induction, without assigning any unknown rank. All exceptional minors are covered by choosing the first nonzero one in a fixed finite order. For $d \geq 8$, $n_d = (d_a + d_b)(d + 1) - d_a d_b \leq 16d + 16$; the candidate matrix has at most $\sum_{e=2}^5 |\mathcal{G}_e| n_{d-e}$ columns. Its row and column dimensions are therefore bounded linearly in d . Temporary full biform columns have $(d + 1)^2$ entries, as in OCF; no claim that all storage objects have linear size is made. The exact output evaluates the residual integer by the proved identity

$$\boxed{t_G = 8k - 32 - \dim J_k^{\text{red}}.} \quad (\text{ACR23})$$

Thus all coefficient cases on the original five-orbit domain have explicit finite formulas, with the unchanged original residual quotient ACR16. The separate literal period support calculation can determine where its first case ACR17 holds; its use then retains the full complementary residual, rather than inferring that this residual has vanished.

68 Exact exclusion of all four boundary rulings at the original period limit

Retain the original hypotheses $0 < \delta < 1/2$, $\gamma > 2$ and $a \neq 0$. In particular, no phase of the original amplitude is selected. Set

$$x = \delta + i\gamma, \quad y = \delta - i\gamma, \quad \beta = 2(\gamma^2 - \delta^2) = -(x^2 + y^2), \quad t_{xy} = x/y, \quad r = a/\bar{a}, \quad R = r^2. \quad (\text{LBRE1})$$

All these ratios are defined, $|t_{xy}| = |r| = 1$, and $x^2 - y^2 = 4i\delta\gamma \neq 0$. Thus $t_{xy}^2 \neq 1$. The root order is exactly $(x, y, -y, -x)$, the full unit is $U = \text{diag}(a, \bar{a}, \bar{a}, a)$, and V evaluates $1, w, w^2, w^3$ in this root order. For $j \in \{1, 2, 3, 4\}$ put $\rho = \zeta^{-j}$, where $\zeta = e^{2\pi i/5}$. Retain PCL4's complete matrix

$$J_\infty = \begin{pmatrix} 1 & 0 & -\beta/3 & 0 \\ 0 & 1 & 0 & -2\beta/3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad L = U^{-1}VJ_\infty^{-1} \text{diag}(\rho, \rho^2, \rho^3, \rho^4)J_\infty V^{-1}U. \quad (\text{LBRE2})$$

The original form is $Q_0(t) = t_1t_4 - t_2t_3$.

68.1 The exact even and odd evaluation maps

Define the following finite coefficients:

$$A = \frac{2t_{xy}^2 - 1}{3(t_{xy}^2 - 1)}, \quad B = \frac{2 - t_{xy}^2}{3(t_{xy}^2 - 1)}, \quad h = \rho^3 - \rho, \quad (\text{LBRE3})$$

$$e = \rho + hA, \quad f = -hA, \quad g = hB, \quad d = \rho - hB.$$

Here $A - B = 1$. The notation d in this section denotes precisely this scalar and does not denote the original root displacement δ . For any primary vector $t \in \mathbb{C}^4$ set

$$v_+(t) = \frac{1}{2} \begin{pmatrix} t_1 + t_4 \\ t_2 + t_3 \end{pmatrix}, \quad v_-(t) = \frac{1}{2} \begin{pmatrix} t_1 - t_4 \\ t_2 - t_3 \end{pmatrix}, \quad J_2 = \text{diag}(1, -1). \quad (\text{LBRE4})$$

These are explicit coordinate maps on the original primary space; their inverse sends (v_+, v_-) to $(v_{+,1} + v_{-,1}, v_{+,2} + v_{-,2}, v_{+,2} - v_{-,2}, v_{+,1} - v_{-,1})$. Consequently

$$Q_0(t) = v_+(t)^\top J_2 v_+(t) - v_-(t)^\top J_2 v_-(t). \quad (\text{LBRE5})$$

The complete matrices induced by L in these coordinates are

$$v_+(Lt) = Ev_+(t), \quad v_-(Lt) = Ov_-(t), \quad E = \begin{pmatrix} e & f/r \\ rg & d \end{pmatrix}, \quad O = \rho \begin{pmatrix} d & t_{xy}g/r \\ rf/t_{xy} & e \end{pmatrix}. \quad (\text{LBRE6})$$

To prove the formulas, write an original polynomial as $p_0 + p_1w + p_2w^2 + p_3w^3$. On its even coefficients the middle conjugation in LBRE2 is

$$\begin{pmatrix} \rho & \frac{\beta}{3}(\rho^3 - \rho) \\ 0 & \rho^3 \end{pmatrix}, \quad (\text{LBRE7})$$

while on its odd coefficients it is

$$\begin{pmatrix} \rho^2 & \frac{2\beta}{3}(\rho^4 - \rho^2) \\ 0 & \rho^4 \end{pmatrix}. \quad (\text{LBRE8})$$

The even evaluation matrix is $\begin{pmatrix} 1 & x^2 \\ 1 & y^2 \end{pmatrix}$; the odd evaluation matrix is its left product with $\text{diag}(x, y)$.

Direct evaluation of LBRE7 gives $\begin{pmatrix} e & f \\ g & d \end{pmatrix}$, because

$$\frac{x^2 + \beta/3}{x^2 - y^2} = A, \quad \frac{y^2 + \beta/3}{x^2 - y^2} = B. \quad (\text{LBRE9})$$

For LBRE8 the two corresponding fractions are $-B, -A$, since $(x^2 + 2\beta/3)/(x^2 - y^2) = -B$ and $(y^2 + 2\beta/3)/(x^2 - y^2) = -A$. Also $\rho^4 - \rho^2 = \rho h$. Thus its evaluated matrix is $\rho \begin{pmatrix} d & t_{xy}g \\ f/t_{xy} & e \end{pmatrix}$. Finally conjugation by the full unit acts on both parity spaces by $\text{diag}(a, \bar{a})$, proving LBRE6 including every amplitude factor.

68.2 The four complete quadratic restrictions

Define

$$\begin{aligned} n_1 &= e^2 - \rho^2 d^2, & c_1 &= \rho^2 f^2 / t_{xy}^2 - g^2, \\ n_2 &= f^2 - \rho^2 t_{xy}^2 g^2, & c_2 &= \rho^2 e^2 - d^2, \\ \mathcal{A} &= n_1 + R c_1, & \mathcal{C} &= (n_2 + R c_2) / R, \\ m_E &= e f / r - r g d, & m_O &= \rho^2 (d t_{xy} g / r - r f e / t_{xy}), \\ \mathcal{B}_- &= m_E - m_O, & \mathcal{B}_+ &= m_E + m_O. \end{aligned} \tag{LBRE10}$$

For the affine Segre vector $t(Z, W) = (ZW, Z, W, 1)^\top$, write $q_j(Z, W) = Q_0(Lt(Z, W))$. The restrictions below also give the exact homogeneous restrictions at infinity:

$$\begin{aligned} Z = 0 : & \quad Q_0(L(0, 0, W, 1)^\top) = \frac{1}{4}(\mathcal{A} + 2\mathcal{B}_- W + \mathcal{C} W^2), \\ Z = \infty : & \quad Q_0(L(W, 1, 0, 0)^\top) = \frac{1}{4}(\mathcal{A} W^2 + 2\mathcal{B}_- W + \mathcal{C}), \\ W = 0 : & \quad Q_0(L(0, Z, 0, 1)^\top) = \frac{1}{4}(\mathcal{A} + 2\mathcal{B}_+ Z + \mathcal{C} Z^2), \\ W = \infty : & \quad Q_0(L(Z, 0, 1, 0)^\top) = \frac{1}{4}(\mathcal{A} Z^2 + 2\mathcal{B}_+ Z + \mathcal{C}). \end{aligned} \tag{LBRE11}$$

For example the two parity vectors for $(W, 1, 0, 0)$ both equal $(W, 1)^\top/2$, so LBRE5–6 give the second row directly. For $(0, 0, W, 1)$ they are $(1, W)^\top/2$ and its negative, giving the first row. For $(Z, 0, 1, 0)$ they are $(Z, 1)^\top/2$ and $(Z, -1)^\top/2$, giving the last row. For $(0, Z, 0, 1)$ they are $(1, Z)^\top/2$ and $(-1, Z)^\top/2$, giving the third row. These substitutions also prove every mixed-coefficient sign and the factor $1/4$. In particular any identically zero boundary restriction would require $\mathcal{A} = \mathcal{C} = 0$.

68.3 A nonzero exact eliminant for every original amplitude

Put

$$q = t_{xy}^2 + t_{xy}^{-2} \in [-2, 2], \quad c = \rho^2 + \rho^3 \in [-2, 2], \quad F(q, c) = 5(c-1)q^2 + (20c+25)q - 61c - 173. \tag{LBRE12}$$

The first interval follows from $|t_{xy}| = 1$. For the second, $\rho^3 = \overline{\rho^2}$ because $\rho^5 = 1$. Direct algebra in LBRE3 and LBRE10 gives the exact identity

$$n_1 c_2 - c_1 n_2 = \frac{t_{xy}^2}{81(t_{xy}^2 - 1)^2} F(q, c). \tag{LBRE13}$$

Here is a finite expansion verifying the identity without any numerical specialization. Before using the fifth-root relation, the left side has denominator $81t_{xy}^2(t_{xy}^2 - 1)^2$ and numerator

$$\begin{aligned} & -\rho^6(\rho^2 - 1)^2 [(t_{xy}^8 + 1)(\rho^4 - 2\rho^2 + 1) \\ & \quad + (t_{xy}^6 + t_{xy}^2)(13\rho^4 + 19\rho^2 + 13) \\ & \quad - t_{xy}^4(57\rho^4 + 120\rho^2 + 57)]. \end{aligned} \tag{LBRE14}$$

Use $1 + \rho + \rho^2 + \rho^3 + \rho^4 = 0$ to replace this numerator by

$$5(c-1)(t_{xy}^8 + 1) + (20c+25)(t_{xy}^6 + t_{xy}^2) - (51c+183)t_{xy}^4. \tag{LBRE15}$$

The identities $(t_{xy}^8 + 1)/t_{xy}^4 = q^2 - 2$ and $(t_{xy}^6 + t_{xy}^2)/t_{xy}^4 = q$ now give LBRE13. All denominators used here are nonzero by LBRE1.

For $-2 \leq q \leq 2$, the polynomial $5q^2 + 20q - 61$ has maximum -1 : its derivative is $10q + 20 \geq 0$ on that interval and its value at 2 is -1 . Since $c \geq -2$, it follows that

$$\begin{aligned} F(q, c) &= c(5q^2 + 20q - 61) - 5q^2 + 25q - 173 \\ &\leq -2(5q^2 + 20q - 61) - 5q^2 + 25q - 173 \\ &= -15(q + \tfrac{1}{2})^2 - \tfrac{189}{4} < 0. \end{aligned} \tag{LBRE16}$$

Thus LBRE13 is nonzero. If $\mathcal{A} = \mathcal{C} = 0$, then $n_1 + Rc_1 = n_2 + Rc_2 = 0$ since $R \neq 0$. Multiplying the first equation by c_2 , the second by c_1 , and subtracting would give $n_1c_2 - c_1n_2 = 0$, a contradiction. Consequently

$$(\mathcal{A}, \mathcal{C}) \neq (0, 0) \quad \text{for every allowed } \delta, \gamma, a \text{ and every } j \in \{1, 2, 3, 4\}. \tag{LBRE17}$$

There is also a fully explicit coefficient margin. Indeed $t_{xy} - t_{xy}^{-1} = 4i\delta\gamma/(\delta^2 + \gamma^2)$, and therefore

$$\begin{aligned} \mathcal{D}_j := n_1c_2 - c_1n_2 &= -\frac{(\delta^2 + \gamma^2)^2}{1296\delta^2\gamma^2}F(q, c) \geq \frac{7(\delta^2 + \gamma^2)^2}{192\delta^2\gamma^2} > 0, \\ \max(|\mathcal{A}|, |\mathcal{C}|) &\geq \frac{\mathcal{D}_j}{|c_1| + |c_2|} > 0. \end{aligned} \tag{LBRE18}$$

The last denominator is positive because simultaneous vanishing of c_1, c_2 would contradict $\mathcal{D}_j > 0$. For its proof use $\mathcal{D}_j = \mathcal{A}c_2 - Rc_1c_2$ and $|R| = 1$, then apply the triangle inequality. Thus on every boundary at least one of its two endpoint coefficients has modulus at least $\mathcal{D}_j/[4(|c_1| + |c_2|)]$, with the exact scalar $1/4$ from LBRE11 retained. Each polynomial in LBRE11 is therefore nonzero. In homogeneous Segre coordinates, none of z_0, z_1, w_0, w_1 divides the actual limiting biform $Q_0(Lt)$. This conclusion includes every amplitude phase and all eigenvalue-collision phases; its proof never excludes such a phase. It also gives the corresponding statement for the full limiting conductor product: restriction to any one boundary is the product of its four nonzero quadratic restrictions, which is nonzero because the polynomial ring over $\mathbb{C}$ has no zero divisors. Hence none of the four homogeneous boundary coordinates divides A_0 of PCL8.

Finally the biform coefficients are entire in the original inverse period z by PCL8–11. For each of the finitely many pairs consisting of an orbit index and a boundary, LBRE11 and LBRE17 supply a nonzero coefficient at $z = 0$. That exact coefficient function is continuous, so it stays nonzero on a disk about 0. The minimum of these finitely many positive radii is positive. Therefore all four boundary exclusions hold simultaneously on some neighborhood of the original inverse-period limit, retaining the original period and unit throughout. No explicit size for this last neighborhood is asserted here.

69 Actual boundary multiplicities and a finite full-support radius

Retain the original quartet, unit, period matrices, logarithm branch and cyclic orientation of PCL1–11, PQO1–25 and RPC1–26. In particular, $r_1 = \delta + i\gamma$, $r_2 = \delta - i\gamma$, $r_3 = -r_2$, $r_4 = -r_1$, $0 < \delta < 1/2$, $\gamma > 2$ and $U = \text{diag}(a, \bar{a}, \bar{a}, a)$ with the full actual $a \neq 0$. Put $z = u^{-1}$ and retain

$$\begin{aligned} L_j(z) &= U^{-1}V\mathcal{R}(z)^{-1}D^{-j}\mathcal{R}(z)V^{-1}U, \\ q_j(z; Z, W) &= Q_0(L_j(z)(ZW, Z, W, 1)^\top), \quad A_z = \prod_{j=1}^4 q_j(z; Z, W). \end{aligned} \quad (\text{ASR1})$$

These are exact identities in the original period map by PCL9. Every entry of L_j is entire, since $\det \mathcal{R} = 1$ and $\mathcal{R}^{-1} = \text{adj } \mathcal{R}$. In particular all coefficients below are the entire functions evaluated by PCL2 and PCL10, with their original signs and ordered-pair multiplicities.

69.1 Every finite-period boundary multiplicity

Let S_0 be the symmetric matrix with entries $(S_0)_{14} = (S_0)_{41} = 1/2$, $(S_0)_{23} = (S_0)_{32} = -1/2$ and all other entries zero, and set $S_j(z) = L_j(z)^\top S_0 L_j(z)$. Thus

$$\begin{aligned} q_j &= (S_j)_{11}Z^2W^2 + 2(S_j)_{12}Z^2W + (S_j)_{22}Z^2 + 2(S_j)_{13}ZW^2 \\ &\quad + 2((S_j)_{14} + (S_j)_{23})ZW + 2(S_j)_{24}Z + (S_j)_{33}W^2 + 2(S_j)_{34}W + (S_j)_{44}. \end{aligned} \quad (\text{ASR2})$$

This follows by substituting all four coordinates into $t^\top S_j t$; each off-diagonal term occurs twice. Introduce the following six three-entry vectors, with the displayed order of their coefficients:

$$\begin{aligned} E_{Z,0,j} &= ((S_j)_{33}, 2(S_j)_{34}, (S_j)_{44}), \\ E_{Z,\infty,j} &= ((S_j)_{11}, 2(S_j)_{12}, (S_j)_{22}), \\ E_{W,0,j} &= ((S_j)_{22}, 2(S_j)_{24}, (S_j)_{44}), \\ E_{W,\infty,j} &= ((S_j)_{11}, 2(S_j)_{13}, (S_j)_{33}), \\ M_{Z,j} &= (2(S_j)_{13}, 2((S_j)_{14} + (S_j)_{23}), 2(S_j)_{24}), \\ M_{W,j} &= (2(S_j)_{12}, 2((S_j)_{14} + (S_j)_{23}), 2(S_j)_{34}). \end{aligned} \quad (\text{ASR3})$$

On the original five-orbit locus, for $X = Z, W$ and $b = 0, \infty$ define the following integer by literal vanishing of these actual coefficient vectors:

$$\nu_{X,b,j} = \begin{cases} 0, & E_{X,b,j} \neq 0, \\ 1, & E_{X,b,j} = 0, \quad M_{X,j} \neq 0, \\ 2, & E_{X,b,j} = 0, \quad M_{X,j} = 0. \end{cases} \quad (\text{ASR4})$$

On the original five-orbit locus, q_j is nonzero. To verify this without a geometric identification, ASR2 is zero exactly when its nine displayed coefficients vanish. These equations say precisely $S_j = cS_0$ for a scalar c , so $Q_0(L_j t) = cQ_0(t)$ in the four independent primary variables. Conjugation by the original period matrix identifies this with projective equality of orbit quadrics, which is excluded for $1 \leq j \leq 4$ on that locus. Since S_j is invertible, the scalar in such an equality could not be zero.

The homogeneous form of q_j has bidegree $(2, 2)$ with $t = (z_0 w_0, z_0 w_1, z_1 w_0, z_1 w_1)$. For $X = Z = z_0/z_1$, its three successive powers of z_0 have coefficient vectors $E_{Z,0,j}, M_{Z,j}, E_{Z,\infty,j}$. This proves that ASR4 is exactly its order along $z_0 = 0$. Reversing the powers proves the assertion for $z_1 = 0$. The two identical calculations with $W = w_0/w_1$ prove the remaining two orders. In the last case of ASR4 the opposite vector is nonzero, since $q_j \neq 0$.

Write a_-, a_+ for the minimum and maximum Z exponents in A_z , and b_-, b_+ for its minimum and maximum W exponents. The exact actual support extrema are therefore

$$\boxed{\begin{aligned} a_- &= \sum_{j=1}^4 \nu_{Z,0,j}, & a_+ &= 8 - \sum_{j=1}^4 \nu_{Z,\infty,j}, \\ b_- &= \sum_{j=1}^4 \nu_{W,0,j}, & b_+ &= 8 - \sum_{j=1}^4 \nu_{W,\infty,j}. \end{aligned}} \quad (\text{ASR5})$$

Indeed the lowest nonzero Z coefficient of a product is the product of its lowest coefficients in the integral domain $\mathbb{C}[W]$, and the same is true of the highest coefficients. Thus neither extreme can cancel. The argument over $\mathbb{C}[Z]$ gives both W extremes. Equivalently the complete boundary divisor and the remaining bihomogeneous factor are specified by

$$M_\partial = z_0^{a_-} z_1^{8-a_+} w_0^{b_-} w_1^{8-b_+}, \quad A = M_\partial A_{\text{red}}, \quad (d_a, d_b) = (a_+ - a_-, b_+ - b_-). \quad (\text{ASR6})$$

This retains the order at each of the four original boundary rulings.

69.2 The exact symplectic map underlying the coefficient tests

Put $B = V^{-1}U$ and retain RPC5's alternating matrix J . The original alternating form on primary coordinates is

$$\begin{aligned} \Omega_{\text{prim}} &= B^\top J B, \\ (\Omega_{\text{prim}})_{14} &= \frac{a^2}{8i\delta\gamma r_1}, & (\Omega_{\text{prim}})_{23} &= -\frac{\bar{a}^2}{8i\delta\gamma r_2}, \end{aligned} \quad (\text{ASR7})$$

with the negative transposed entries and all others zero. In fact RPC4 evaluates this form as $\sum_i a_i^2 t_i v_{5-i} / H'(r_i)$, where $(a_1, a_2, a_3, a_4) = (a, \bar{a}, \bar{a}, a)$, $H'(r_1) = 8i\delta\gamma r_1$ and $H'(r_2) = -8i\delta\gamma r_2$. This proves every entry and its sign. Using $P_0 = G_u \mathcal{R}(z)$ in RPC9 gives $\mathcal{R}(z)^{-\top} J \mathcal{R}(z)^{-1} = J|_{\beta=0}$. The diagonal D preserves this last alternating matrix by its paired weights. Conjugating the equality by $\mathcal{R}(z)B$ proves

$$L_j(z)^\top \Omega_{\text{prim}} L_j(z) = \Omega_{\text{prim}}. \quad (\text{ASR8})$$

Let $s = r_1/a^2$, retain its literal complex conjugate $\bar{s} = r_2/\bar{a}^2$, and define $T = \text{diag}(s, \bar{s}, -\bar{s}, -s)$. The entries of ASR7 prove the exact matrix identity

$$S_0 = -4i\delta\gamma \Omega_{\text{prim}} T. \quad (\text{ASR9})$$

Each original boundary plane is one of $E_{12}, E_{34}, E_{13}, E_{24}$, where E_{ab} is spanned by the two indicated original coordinate vectors. Each is Lagrangian for ASR7: its restriction vanishes, its dimension is two, and the ambient alternating matrix is invertible. Equation ASR8 shows that its image $F = L_j E_{ab}$ is also Lagrangian. Polarization of ASR9 now proves the exact equivalence

$$Q_0|_{L_j E_{ab}} = 0 \iff TL_j E_{ab} \subset L_j E_{ab} \iff \text{rank}[L_j E_{ab}, TL_j E_{ab}] = 2. \quad (\text{ASR10})$$

Here the last matrix has the two columns of L_j indexed by a, b followed by their images under T . For the first equivalence, polarization says $\Omega_{\text{prim}}(v, Tw) = 0$ for all $v, w \in F$; since $F^{\perp\Omega} = F$, this is exactly $TF \subset F$. For the last equivalence the first two columns are independent. Thus all its three-by-three minors vanish precisely in that case. These are the same boundary-vanishing tests as the corresponding three entries of ASR3, with the exact map between the two tests proved. No eigenvalue is cancelled when s is real or purely imaginary.

69.3 An explicit radius from the actual limiting corner calculation

The complete limiting calculation LBRE1–18 proves, for every $1 \leq j \leq 4$ and every original nonzero a , that

$$\begin{aligned} d_{j,1}(0) &= d_{j,4}(0), & d_{j,2}(0) &= d_{j,3}(0), \\ m_j &:= \max\{|d_{j,1}(0)|, |d_{j,2}(0)|\} > 0, & d_{j,i}(z) &:= (S_j(z))_{ii}. \end{aligned} \quad (\text{ASR11})$$

Indeed LBRE11 gives $d_{j,1}(0) = d_{j,4}(0) = \mathcal{A}/4$ and $d_{j,2}(0) = d_{j,3}(0) = \mathcal{C}/4$ with the complete original amplitude in LBRE10. LBRE13–18 prove the positive lower bound

$$m_j \geq \frac{\mathcal{D}_j}{4(|c_1| + |c_2|)} > 0, \quad \mathcal{D}_j \geq \frac{7(\delta^2 + \gamma^2)^2}{192\delta^2\gamma^2},$$

where $c_1, c_2, \mathcal{D}_j$ here are exactly LBRE10 and LBRE18's finite coefficients for this orbit index. That proof evaluates the literal J_∞ in PCL4 and the original r_1/r_2 and ζ^{-j} . The two equalities also follow directly from PCL22–23 at $z = 0$, because the opposite-root permutation fixes Q_0 and conjugates $L_j(0)$ to itself.

Expand each of these actual entire diagonal entries as $d_{j,i}(z) = \sum_{n \geq 0} d_{j,i,n} z^n$ by PCL2 and PCL10, and define the explicitly specified finite constant

$$C_j = \sum_{n \geq 1} \max_{1 \leq i \leq 4} |d_{j,i,n}|, \quad R_{\text{supp}} = \max \left\{ R_*, 1, \frac{2C_1}{m_1}, \frac{2C_2}{m_2}, \frac{2C_3}{m_3}, \frac{2C_4}{m_4} \right\}. \quad (\text{ASR12})$$

Each series is finite in value: the four entire Taylor series converge absolutely at $z = 1$, and the maximum is bounded by their sum of absolute values. Every denominator is strictly positive by ASR11. If $C_j = 0$, its displayed quotient is zero; no division by C_j is performed. For $|z| \leq 1$, summing the same series gives

$$|d_{j,i}(z) - d_{j,i}(0)| \leq C_j |z| \quad (1 \leq i \leq 4). \quad (\text{ASR13})$$

Choose $i_j = 1$ when $|d_{j,1}(0)| \geq |d_{j,2}(0)|$, and $i_j = 2$ otherwise. This deterministic choice selects a coefficient already present in the actual period calculation. For $|u| \geq R_{\text{supp}}$, ASR11–13 give

$$|d_{j,i_j}(u^{-1})| \geq \frac{m_j}{2} > 0, \quad |d_{j,5-i_j}(u^{-1})| \geq \frac{m_j}{2} > 0. \quad (\text{ASR14})$$

Indeed both initial values are equal with modulus m_j and their errors are at most $C_j/|u| \leq m_j/2$. Each of the four boundary planes in ASR10 contains exactly one of these two opposite coordinate vectors. Thus its boundary restriction has a nonzero diagonal coefficient, so all four vectors $E_{X,b,j}$ in ASR3 are nonzero. This proves, at every such original u on every logarithm branch,

$$\boxed{a_- = b_- = 0, \quad a_+ = b_+ = 8, \quad (d_a, d_b) = (8, 8), \quad M_\partial = 1, \quad A_{\text{red}} = A.} \quad (\text{ASR15})$$

Branch independence follows from the exact original-period conjugation PCL9, which leaves L_j single-valued in u^{-1} . The radius includes R_* , so all these points are on the proved original five-orbit locus. The estimate has introduced no new source order and no freely specified amplitude or period coefficient.

At arbitrary nonzero original periods, each boundary failure is the simultaneous zero set of its three actual entire entries in ASR3. ASR11 supplies a nonzero value at $z = 0$ for at least one of those entries. Such an entry has isolated zeros: at any zero its convergent Taylor series has a first nonzero term, and factoring that power leaves a nonvanishing factor in a sufficiently small disk. Thus each boundary restriction-vanishing set is discrete and closed in the finite z plane, and its finite union over the sixteen boundary/index pairs is also discrete and closed. This boundary restriction-vanishing set in the nonzero u plane is contained in $|u| < R_{\text{supp}}$ and has no accumulation point other than possibly $u = 0$. ASR3–5 retain every multiplicity on its intersection with the original five-orbit locus. At an orbit-collapse period, ASR2's nine coefficients detect an identically zero factor; no finite ruling multiplicity is assigned to such a factor.

69.4 Consequence for the actual common factor and residual

Retain the common-factor and grid maps of ACR1–23, with original $k = 4\ell + 1 \geq 9$ and both unchanged source orders $1, k$. Its exact dimension formula and its actual invariant-image quotient are

$$\begin{aligned} g_A &= (k + 1 - d_a)(k + 1 - d_b) - (k - 7)^2, \\ t_G &= \dim \frac{J_k \cap A_{\text{red}} \mathcal{B}_{k-d_a, k-d_b}}{A \mathcal{B}_{k-8, k-8}}. \end{aligned} \quad (\text{ASR16})$$

Here $\mathcal{B}_{p,q}$ is the space of bihomogeneous polynomials of bidegree (p, q) in the two original pairs $(z_0, z_1), (w_0, w_1)$, and J_k is the original invariant image. Its exact map from the affine coefficient space used in ASR1–5 is

$$f(Z, W) \mapsto z_1^p w_1^q f(z_0/z_1, w_0/w_1), \quad F \mapsto F(Z, 1, W, 1).$$

The respective bounded degrees ensure that the first expression is a polynomial. Expansion in its monomials proves that these two maps are inverse and that multiplication is preserved with the displayed bidegrees. ACR5 and ACR15 prove the two formulas in ASR16; ACR8–9 prove that the denominator is a subspace of that numerator. At every $|u| \geq R_{\text{supp}}$, ASR15 first makes the two products in g_A equal. It then makes the numerator of the displayed quotient a subspace of its denominator; the opposite inclusion was already proved by the exact receiving map. Consequently

$$g_A = t_G = 0 \quad (|u| \geq R_{\text{supp}}, \quad k = 4\ell + 1 \geq 9). \quad (\text{ASR17})$$

This removes precisely the common boundary factor calculated by that map. The original remaining residual subspace is still $W = B_{\Xi}$, and the receiving calculation's full residual rank remains

$$j = ((k + 1)^2 - (k - 7)^2) - (8k - 16) = 8k - 32.$$

Here the first parenthesis is the original conductor quotient dimension from its injective multiplication by the nonzero biform A , and $8k - 16$ is the exact original invariant-image kernel dimension of OQX and OCF on this five-orbit locus. Neither that factor nor those residual directions have been set to zero. For periods below this explicit support radius that remain on the original five-orbit locus, ASR3–6 give all actual boundary losses without assuming full widths.

69.5 Only finitely many support exceptions on the original exterior domain

Define the exact coefficient set and its corresponding original periods by

$$\begin{aligned} \mathcal{Z}_{\partial}^* &= \{z : 0 < |z| \leq R_*^{-1}, \quad E_{X,b,j}(z) = 0 \text{ for some } X \in \{Z, W\}, \quad b \in \{0, \infty\}, \quad 1 \leq j \leq 4\}, \\ \mathcal{U}_{\partial}^* &= \{z^{-1} : z \in \mathcal{Z}_{\partial}^*\}. \end{aligned}$$

For each of the sixteen boundary/index pairs, ASR11 selects one diagonal coefficient among the three entries of $E_{X,b,j}$ that is nonzero at $z = 0$. This is an entire function with isolated zeros by the local Taylor argument given above. It has finitely many zeros on the compact disk $|z| \leq R_*^{-1}$: an infinite set would have an accumulation point in that disk, contradicting the proved isolation. The common zero set of the three entries is a subset of this finite set. Taking the union of the sixteen finite sets proves finiteness of $\mathcal{Z}_{\partial}^*$ and hence of $\mathcal{U}_{\partial}^*$. None of these zeros is zero, by the selected nonzero value there. ASR14 excludes every inverse period with $|u| \geq R_{\text{supp}}$. On the whole domain $|u| \geq R_*$, the factors are nonzero by the original five-orbit theorem, so ASR4–5 show that this is precisely the set where at least one of the four support extrema loses its full value. Thus

$\begin{aligned} &\mathcal{U}_{\partial}^* \text{ is finite,} \quad \mathcal{U}_{\partial}^* \subseteq \{u : R_* \leq u < R_{\text{supp}}\}, \\ & u \geq R_*, \quad u \notin \mathcal{U}_{\partial}^* \implies (a_-, a_+, b_-, b_+) = (0, 8, 0, 8), \\ &g_A = 0, \quad t_G = 0, \quad W = B_{\Xi}, \quad j = 8k - 32 \quad (k = 4\ell + 1 \geq 9). \end{aligned}$	(ASR18)
--	---------

At the finite exceptional periods every boundary multiplicity is given by ASR3–5, and the unchanged matrices ACR12–15 evaluate the remaining common-factor map and its residual quotient. No statement that this finite set is empty is needed for any displayed conclusion.

70 The two exact dual subquotients in the original receiver

Fix the original simple quartet, full unit, actual five-orbit period and logarithm branch. All coefficients of the actual conductor A and all original roots are retained. In this section

$$\begin{aligned}
k &= 4l + 1 \geq 9, \quad q = (k + 1)^2, \quad q' = (k - 7)^2, \\
\Delta &= q - q' = 16k - 48, \quad r_K = 8k - 16, \quad j = 8k - 32, \\
E_A(w) &= \sum_{a,b=0}^8 a_{ab} e^{b_{ab}w}, \quad v = \text{ord}_0 E_A \leq 80, \quad \mu_v = E_A^{(v)}(0) \neq 0, \quad \alpha_A = v!/\mu_v, \\
\mathbf{m} &= \Delta - v, \quad c' = k/2 - 4, \quad s \in \{1, k\}, \quad \mathcal{N} = \{q - 1, q, 2q - 1, 2q\}, \\
(\sigma_{q-1}, \sigma_q, \sigma_{2q-1}, \sigma_{2q}) &= (1, 1, -1, -1).
\end{aligned} \tag{PBR1}$$

The original relation quotient has Gram $H_{Y,N}^{(s)}$ in the fixed FGC frame, and Z_k is its actual FGC10 kernel-column matrix. Here $K = \ker \Xi$ in the original primary space V_A , $L = \mathbb{C}[S]_{<v} \subset V_A$, $K^+ = K + L$, $t_0 = \dim(K \cap L)$, and $t = r_K - t_0$ is the number of columns of Z_k . The symbol r_K denotes this rank parameter, whereas ρ below denotes an analytic radius. No independent source order is introduced.

Retain the complete DNE coefficient frame Z_{exp} , the functions $F_a = \mathcal{N}_a/E_A$, and $\ell_a(g) = \Lambda_{F_a}(\chi'g)$. Put $(A_\ell)_{ba} = \ell_a(S'^b)$ for $0 \leq b < \mathbf{m}$. The matrix A_ℓ is invertible by DNE4. In the original Gamma orthonormal frame $\phi_{n,s,c'}$, $0 \leq n \leq N - v$, whose squared monic norms are $h_n = M_s n!(s/2)_n$, $M_s = (2\pi)^{s/2}$, define

$$\begin{aligned}
T_{na} &= \Lambda_{F_a}(\phi_{n,s,c'}), \quad V_{n,(i,j)} = \phi_{n,s,c'}(\beta'_{ij}), \\
\mathcal{D}_N^{(s)} &= T^*[I - V(V^*V)^{-1}V^*]T = A_\ell^*(H_{Y,N}^{(s)})^{-1}A_\ell, \\
G_{\text{exp}} &= Z_{\text{exp}}^*[I - C^\top(\bar{C}C^\top)^{-1}\bar{C}]Z_{\text{exp}} > 0, \\
B_K &= Z_k^\top A_\ell, \quad D_{K,N} = [B_K(\mathcal{D}_N^{(s)})^{-1}B_K^*]^{-1} = (\widehat{H}_{K,N}^{(s)})^{-1}, \quad G_K = [B_K G_{\text{exp}}^{-1}B_K^*]^{-1}.
\end{aligned} \tag{PBR2}$$

The onto map B_K is the exact restriction of the complex-linear dual to the actual kernel columns. The conjugation in PBR2 follows by substituting the displayed A_ℓ identity and $B_K = Z_k^\top A_\ell$; it gives $B_K \mathcal{D}_N^{-1} B_K^* = Z_k^\top \bar{H}_{Y,N} \bar{Z}_k = \bar{Z}_k^* \bar{H}_{Y,N} \bar{Z}_k$ before inversion. The first equality for $\mathcal{D}_N$ minimizes the full extension over every lower evaluation functional. Its bracket is the orthogonal projection off their entire column span. Thus the full conditional covariance and the coefficient reference are both specified; neither is replaced by the other.

For the complementary quotient choose exactly the first invertible t -row minor of Z_k , with row set $\mathcal{I}$ and complementary set $\mathcal{J}$, and put

$$\begin{aligned}
j_0 &= \mathbf{m} - t = j - v + t_0, \quad P_Y = E_{\mathcal{J}}^\top - E_{\mathcal{J}}^\top Z_k (E_{\mathcal{I}}^\top Z_k)^{-1} E_{\mathcal{I}}^\top, \\
J_0 &= A_\ell^{-1} P_Y^\top, \quad D_{0,N} = J_0^* \mathcal{D}_N^{(s)} J_0, \quad G_0 = J_0^* G_{\text{exp}} J_0, \\
Q_{0,N} &= [P_Y (H_{Y,N}^{(s)})^{-1} P_Y^*]^{-1}, \quad D_{0,N} = \overline{Q_{0,N}^{-1}}.
\end{aligned} \tag{PBR3}$$

Empty rank uses the identity quotient. Multiplication shows $P_Y Z_k = 0$ and $P_Y E_{\mathcal{J}} = I$; hence $\ker P_Y = \text{im } Z_k$. Also $B_K J_0 = 0$ and J_0 has j_0 independent columns, proving $\text{im } J_0 = \ker B_K$. Substitution in PBR2 gives $D_{0,N} = \bar{P}_Y \bar{H}_{Y,N}^{-1} \bar{P}_Y^\top$, which proves the last identity in PBR3. These are the actual kernel quotient and its full dual annihilator.

70.1 Finite positive-spectrum allowance and every small eigenvalue

For clarity the finite allowance used below is the complete PEC13 quantity, with no hidden replacement by an asymptotic zero. For $0 < \rho < 1$ with $E_A(-i \arctan z) \neq 0$ on $|z| = \rho$, put $m_\rho = \min_{|z|=\rho} |E_A(-i \arctan z)| > 0$. The actual nonzero symbol zeros inside this image have

orders h_ω . Form the full jet matrix $(J_{\rho,k})_{(\omega,b),a} = \mathcal{N}_a^{(b)}(\omega)/b!$, $0 \leq b < h_\omega$, and put $c_{\rho,k} = \text{rank } J_{\rho,k}$. With $R = |k/2| + k\sqrt{\delta^2 + \gamma^2}$ and $d = N - v$, retain

$$\begin{aligned} C_{\rho,N,s} &= \frac{q e^{2(R+|c'|) \operatorname{arctanh} \rho} (1 - \rho^2)^{-s/2}}{M_s m_\rho^2} \sum_{n=0}^d \frac{n!}{(s/2)_n} \rho^{-2n}, \\ C_{0,N,s} &= q \mathcal{B}_{d,s,k,A}^2, \quad \varepsilon_{N,s}^\vee = \inf_{\rho \text{ admissible}} \{c_{\rho,k} \log^+ C_{0,N,s} + (\mathfrak{m} - c_{\rho,k}) \log^+ C_{\rho,N,s}\}, \\ E_{k,s}^\vee &= \varepsilon_{2q-1,s}^\vee + \varepsilon_{2q,s}^\vee = o(kq). \end{aligned} \tag{PBR4}$$

Here $\mathcal{B}_{d,s,k,A}$ is the full finite expression DNE11–13, including its chosen zero-free radius, Taylor coefficient and Gamma mass. The full DNE proof is included as a provider, not assumed. Pole cancellation on $\ker J_{\rho,k}$ and Cauchy's coefficient integral give $C_{\rho,N,s}$ exactly: the n th squared coefficient is weighted by $n!/(M_s(s/2)_n)$. Minimization over the entire conductor coefficient fibre preserves these cancellations. The full DNE bound gives $C_{0,N,s}$ on all vectors. At most $c_{\rho,k}$ eigenvalues can exceed $C_{\rho,N,s}$, by intersecting the span of $c_{\rho,k} + 1$ such eigenvectors with the cancellation subspace. This proves the positive logarithmic sum bound PBR4. PEC17–18 proves the same bound for the fixed quotient and restriction in PBR2–3, by the variational principle applied respectively to the inverse compression and the compression. The exact reference metrics G_K, G_0 are retained in both applications.

For completeness, at each fixed admissible radius the symbol has only finitely many zeros in the compact disk, so $c_{\rho,k}$ is bounded independently of k . The full finite constants above and DNE15 give $\limsup \varepsilon_{N,s}^\vee/(kq) \leq 64 \log(1/\rho)$. Admissible radii approach one because the zeros are isolated. Taking that limit after the fixed-radius bound proves the displayed $o(kq)$ for the original two source orders. This bounds every forward exterior degree by $\varepsilon_{N,s}^\vee/2$ in logarithmic norm, by multiplying subsets of the relative eigenvalues. For the inverse map its squared singular values are the reciprocals. They are retained in the following exact formulas.

For $X = K, 0$, let $\mu_{X,a,N}$ be every eigenvalue of $G_X^{-1/2} D_{X,N} G_X^{-1/2}$ and define

$$\begin{aligned} p_{X,N} &= \sum_a \log^+ \mu_{X,a,N}, \quad n_{X,N} = \sum_a \log^+(1/\mu_{X,a,N}), \\ A_{X,k,s} &= n_{X,q-1} + n_{X,q} - n_{X,2q-1} - n_{X,2q}, \\ P_{X,k,s} &= p_{X,2q-1} + p_{X,2q} - p_{X,q-1} - p_{X,q}, \\ \widehat{F}_k^{(s)} &= A_{K,k,s} + P_{K,k,s}, \quad \sum_N \sigma_N \log \det Q_{0,N} = A_{0,k,s} + P_{0,k,s}, \\ A_{X,k,s} &\geq 0, \quad 0 \leq P_{X,k,s} \leq E_{k,s}^\vee. \end{aligned} \tag{PBR5}$$

Increasing N enlarges the original polynomial lift space without changing its norm on previous polynomials or its relation map. Minimum primal lift norms therefore decrease. Inversion in PBR2 increases $\mathcal{D}_N$, and restriction or minimization under the same onto B_K preserves that increase. The variational principle then increases each relative eigenvalue in the fixed reference metric. Thus p increases and n decreases, proving all nonnegativities in PBR5. The two high positive sums are bounded by PBR4. Finally $\log \mu = \log^+ \mu - \log^+(1/\mu)$ and $D_{K,N} = \widehat{H}_{K,N}^{-1}$ give the first return identity with the four stated signs; PBR3 gives the other. The fixed reference determinants cancel only in these four signed sums. In particular the small eigenvalue contributions $n_{X,N}$ have not been bounded away or discarded.

70.2 The original metric on the low image

Let $G_N^{(s)}$ be the original primary quotient metric and let $Q_{\Xi,N}^{\text{fix}}$ be its induced metric on $B_\Xi = \Xi(V_A)$ in the original fixed image frame. Set $W_0 = \Xi(L)$. There is the exact isometric isomorphism

$$\begin{aligned} \iota_N : (K+L)/K &\longrightarrow W_0, & [p] &\longmapsto \Xi p, \\ \|\iota_N[p]\|_{Q_{\Xi,N}^{\text{fix}}}^2 &= \min_{a \in K} G_N^{(s)}(p+a, p+a) = \|[p]\|_{(K+L)/K,N}^2, \\ \dim W_0 &= v - t_0, & F_{W_0}^{(s)} &:= \sum_N \sigma_N \log \det(I_{W_0}^* Q_{\Xi,N}^{\text{fix}} I_{W_0}) = F_{K^+/K}^{(s)}. \end{aligned} \quad (\text{PBR6})$$

Indeed $\Xi p \in \Xi(L)$ if and only if $p \in K+L$. The kernel of the displayed map is exactly K and it is onto. The full fibre of Ξp in V_A is $p+K$, already contained in $K+L$. Hence minimization in the full image norm is exactly the quotient minimization on $K+L$, proving the norm identity at each cutoff. In transported frames the Grams agree exactly. For any other fixed frames their congruence has the same fixed Jacobian at all four endpoints, which cancels with $\sum \sigma_N = 0$. This proves the last identity without identifying W_0 with the entire residual image.

In the FGC coordinates the onto map $[p] \mapsto [\alpha_A \mathcal{T}_A p / \chi']$ from V_A to $Y_N / \text{im } Z_k$ has kernel $K+L$. The corresponding quotient map sends this class to $[\Xi p]$ in B_Ξ / W_0 . The kernel assertion follows directly: subtract the kernel polynomial giving the same FGC image, then the remaining zero image under $\mathcal{T}_A$ is in L ; the converse follows from the two kernels. This proves the exact target isomorphism in PEC27.

The source map comparing this quotient to the native Gamma one is the original CEP map S_N , followed by $D_{\alpha,N}(x, f) = (x, \alpha_A f)$ on the full GIP graph. Retaining every relation column and every Z_k column, minimization first in the low coordinate and then in these polynomial fibres gives precisely $Q_{0,N}$ in PBR3. Let $\bar{Q}_{\Xi,N}$ be the induced original metric on B_Ξ / W_0 in the transported frame. PEC28's full exterior comparison yields

$$\begin{aligned} |\log \det \bar{Q}_{\Xi,N} - \log \det Q_{0,N}| &\leq 2E_N^{\text{CEP}} + 2j_0 |\log |\alpha_A||, \\ E_{k,s}^{\text{tr}} &= 2 \sum_N E_N^{\text{CEP}} + 8j_0 |\log |\alpha_A|| = o(kq), \\ \Phi_{k,s} &= F_{W_0}^{(s)} + A_{0,k,s} + P_{0,k,s} + \delta_{0,k,s}, \quad |\delta_{0,k,s}| \leq E_{k,s}^{\text{tr}}. \end{aligned} \quad (\text{PBR7})$$

To see the error signs explicitly, put $d_N = \log \det \bar{Q}_{\Xi,N} - \log \det Q_{0,N}$. In a fixed frame adapted to W_0 , the full Gram determinant is its restriction determinant times the Schur complement determinant, including its fixed frame Jacobian. Therefore $\delta_{0,k,s} = \sum_N \sigma_N d_N$; the triangle inequality gives the stated bound. The projection used to take each quotient is an exterior contraction; the inverse source map gives the reverse inequality. This proves that the factors α_A , all four original CEP errors, and the full graph denominator occur in this very metric comparison.

70.3 Both low terms and the full one-sided kernel errors

Keep the original FEX endpoint constants explicitly. Write $h_N^{\text{harm}} = \sum_{a=1}^N 1/a$ and $w_d = d!/(s/2)_d$. The symbol t here remains the kernel-column count from PBR1.

$$\begin{aligned} n_N &= N + 1 - v, & m_N &= \log \max \left\{ 1, \frac{v!}{|\mu_v|} \sum_\nu |a_\nu| e^{|b_\nu - 4|(h_N^{\text{harm}} + 2)} \right\}, \\ \zeta_{N,s} &= \log \prod_{d=v}^N \frac{w_d}{w_{d-v}}, & e_{N,s}^- &= \zeta_{N,s} - 2(n_N - t)m_N, & e_{N,s}^+ &= 2tm_N, \\ e_{N,s}^{\text{FEX}} &= \log \det \hat{H}_{K,N}^{(s)} - \log \det H_{K^+/L,N}^{(s)}, & e_{N,s}^- &\leq e_{N,s}^{\text{FEX}} \leq e_{N,s}^+, \\ D_s^- &= e_{q-1,s}^- + e_{q,s}^- - e_{2q-1,s}^+ - e_{2q,s}^+, & D_s^+ &= e_{q-1,s}^+ + e_{q,s}^+ - e_{2q-1,s}^- - e_{2q,s}^-, \\ \mathcal{E}_s^{\text{flag}} &= \sum_N \sigma_N e_{N,s}^{\text{FEX}} \in [D_s^-, D_s^+]. \end{aligned} \quad (\text{PBR8})$$

These are the original FEX18–20/34 expressions, with both source masses retained in the underlying norms. The endpoint inequalities come from the full flag determinant bounds, with squared Jacobian $e^{\zeta_{N,s}}$ and both operator bounds e^{m_N} . The signs of the high endpoint bounds are reversed when forming the four sum, giving exactly $D_s^\pm$ above.

Retain also the exact finite AKS/FEX allowance

$$\begin{aligned}
R_{\text{cen}} &= k\sqrt{\delta^2 + \gamma^2}, \quad \tau_q = \pi/2 - 1/q, \quad a_q = 2\log 3/\tau_q, \quad s_q = (a_q^q - 1)/(a_q - 1), \\
b_{s,q} &= [(9/25)\sin(1/q)]^{-s/2} e^{2R_{\text{cen}}(\log 3 + \tau_q/2)}, \\
\mathcal{L}_{s,h} &= \log \left(1 + b_{s,q} s_q^2 \sum_{d=q}^{q+h-1} \frac{d!}{(s/2)_d} (25/16)^d \right), \quad \mathcal{L}_s = \mathcal{L}_{s,q} + \mathcal{L}_{s,q+1}, \\
0 \leq F_L^{(s)} &\leq v\mathcal{L}_s, \quad 0 \leq F_{W_0}^{(s)} \leq (v - t_0)\mathcal{L}_s, \quad \lambda_s^{\text{low}} = F_L^{(s)} - F_{W_0}^{(s)}, \\
F_K^{(s)} &= A_{K,k,s} + P_{K,k,s} - \mathcal{E}_s^{\text{flag}} + \lambda_s^{\text{low}}.
\end{aligned} \tag{PBR9}$$

The low metric estimates are the original restriction and quotient consequences of AKS, as proved in FEX32–33. Applying the two complete Schur factorizations to K^+ gives OER11, and substituting PBR5 and the isometry PBR6 gives the last identity. The quantities F_L and F_{W_0} remain their separate original volumes. In particular they are not individually removed by using their difference. Combining PBR7 and PBR9 gives the additional exact identity

$$F_K^{(s)} + \Phi_{k,s} = A_{K,k,s} + A_{0,k,s} + P_{K,k,s} + P_{0,k,s} - \mathcal{E}_s^{\text{flag}} + F_L^{(s)} + \delta_{0,k,s}. \tag{PBR10}$$

The W_0 terms cancel in this sum because they are precisely the same original metric by PBR6. They remain in each summand PBR7/9.

70.4 The complete current finite interval

First retain the exact low volumes in the new finite kernel bounds, and separately display the previously proved coarse bounds:

$$\begin{aligned}
\kappa_s^- &= A_{K,k,s} - D_s^+ + \lambda_s^{\text{low}}, \quad \kappa_s^+ = A_{K,k,s} + E_{k,s}^\vee - D_s^- + \lambda_s^{\text{low}}, \quad \kappa_s^- \leq F_K^{(s)} \leq \kappa_s^+, \\
A_{K,k,s} - D_s^+ - (v - t_0)\mathcal{L}_s &\leq F_K^{(s)} \leq A_{K,k,s} + E_{k,s}^\vee - D_s^- + v\mathcal{L}_s, \\
\varphi_s^- &= F_{W_0}^{(s)} + A_{0,k,s} - E_{k,s}^{\text{tr}}, \quad \varphi_s^+ = F_{W_0}^{(s)} + A_{0,k,s} + E_{k,s}^\vee + E_{k,s}^{\text{tr}}, \quad \varphi_s^- \leq \Phi_{k,s} \leq \varphi_s^+, \\
A_{0,k,s} - E_{k,s}^{\text{tr}} &\leq \Phi_{k,s} \leq A_{0,k,s} + E_{k,s}^\vee + E_{k,s}^{\text{tr}} + (v - t_0)\mathcal{L}_s.
\end{aligned} \tag{PBR11}$$

PBR5/8/9 proves the first line by taking the least and greatest allowed signed flag errors; PBR7 proves the third line. The original low caps then prove the second and fourth lines, which are exactly PEC22 and PEC30. Thus using the exact low difference has a specified morphism to each coarser bound.

Let $\Phi_{k,s}^-, \Phi_{k,s}^+$ be the full-window RWC bounds of OER17–18, and retain their entire conditional matrices

$$\begin{aligned}
U_N^+ &= [u_{N+1}, \dots, u_{N+q}], \quad B_N = I_q + (CU_N^+)^*(CR_N C^*)^{-1}(CU_N^+), \\
Z_N &= D[I_q - R_N C^*(CR_N C^*)^{-1}C]U_N^+, \quad \Omega_{N+q} = \Omega_N + Z_N B_N^{-1} Z_N^*, \\
\Phi_{k,s} &= \sum_{N=q-1, q} \log \left(\sum_{a=0}^j \left\| \bigwedge^a (\Omega_N^{-1/2} Z_N B_N^{-1/2}) \right\|_{\text{HS}}^2 \right).
\end{aligned} \tag{PBR12}$$

Here D denotes the original residual rows, as in OER17. All q columns, the complete conditioning inverse, and every exterior coefficient remain. The proof of the covariance identity is the full Schur update RWC4–6; applying the determinant lemma and expanding all singular-value products

gives the last line. Its finite bounds therefore apply to the same Φ as PBR7. Define $\Phi_{k,s}^{-,\vee} = \max\{\Phi_{k,s}^-, \varphi_s^-\}$ and $\Phi_{k,s}^{+,\vee} = \min\{\Phi_{k,s}^+, \varphi_s^+\}$.

Retain the original OAR expression $\mathcal{F}_{\partial,k}^{(s)}$ and its full errors $-e_{k,s}^- \leq \mathfrak{e}_{k,s} \leq e_{k,s}^+$ in $F_K^{(s)} = \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s} + \mathfrak{e}_{k,s}$. These errors are distinct from PBR8. With all earlier OER19 branches still included, the new receiver is

$$\begin{aligned} L_s^\vee &= \max\{L_s^{\text{win}}, \kappa_s^-, \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s}^{+,\vee} - e_{k,s}^-\}, \\ U_s^\vee &= \min\{U_s^{\text{win}}, \kappa_s^+, \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s}^{-,\vee} + e_{k,s}^+\}, \quad L_s^\vee \leq F_K^{(s)} \leq U_s^\vee. \end{aligned} \quad (\text{PBR13})$$

Each lower entry is at most the actual scalar and each upper entry is at least that scalar, proving the intersection and its nonempty order without assuming an unproved relation between the entries. On PMR's exact domain $k \geq 29$ and $k\sqrt{\delta^2 + \gamma^2} \leq 2^{-33}(k+1)^2$, also include in this same maximum and minimum respectively $\mathcal{F}_{0,k}^{(s)} - \Phi_{k,s}^{+,\vee} - e_{k,s}^{0,-}$ and $\mathcal{F}_{0,k}^{(s)} - \Phi_{k,s}^{-,\vee} + e_{k,s}^{0,+}$, where $e_{k,s}^{0,\pm} = e_{k,s}^\pm + c_k^0$. Outside that domain these two entries are omitted; the original OAR expression and errors remain. The seven-moment $\mathcal{F}_{0,k}^{(s)}$ is distinct from the quotient return $\sum \sigma_N \log \det Q_{0,N}$ in PBR5.

For the arithmetic map retain $|u| \geq R_*$ and the same correlated AMT allowances, with all their original finite constants:

$$\begin{aligned} a_{K,k}^- &= \max\{-C_K^-, \delta_k^\sigma - C_B^+\}, & a_{K,k}^+ &= \min\{C_K^+, \delta_k^\sigma + C_B^-\}, \\ L_{\text{ar}}^{\text{win}} &= \max\{L_{\text{ar}}^{\text{op}}, L_1^{\text{win}} + a_{K,k}^-\}, & U_{\text{ar}}^{\text{win}} &= \min\{U_{\text{ar}}^{\text{op}}, U_1^{\text{win}} + a_{K,k}^+\}, \\ L_{\text{ar}}^\vee &= \max\{L_{\text{ar}}^{\text{win}}, L_1^\vee + a_{K,k}^-\}, & U_{\text{ar}}^\vee &= \min\{U_{\text{ar}}^{\text{win}}, U_1^\vee + a_{K,k}^+\}, \\ F_K^{\text{ar}} &\in [L_{\text{ar}}^\vee, U_{\text{ar}}^\vee], & F_B^{\text{ar}} &\in [\mathcal{B}_k^{\text{ar}} - U_{\text{ar}}^\vee, \mathcal{B}_k^{\text{ar}} - L_{\text{ar}}^\vee]. \end{aligned} \quad (\text{PBR14})$$

AMT gives the identical scalar $F_K^{\text{ar}} - F_K^{(1)}$ in $[a_{K,k}^-, a_{K,k}^+]$. Adding this interval to PBR13 proves the new entries. The earlier entries remain; subtracting from the same total $F_K^{\text{ar}} + F_B^{\text{ar}} = \mathcal{B}_k^{\text{ar}}$ reverses the two endpoints and proves the final line. No allowance has changed. Previously available reverse source intersections remain valid.

70.5 The signed return and the complete same-class action budget

Keep $\mathcal{H}_k = -T_k$, the full arithmetic correction δ_k^σ , and exactly the original budget

$$\begin{aligned} C_k^{\text{budget}} &= F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+, & \Psi_k &= F_K^{(k)} + \mathcal{H}_k + \delta_k^\sigma, \\ L_k^\vee + \mathcal{H}_k + \delta_k^\sigma &\leq \Psi_k \leq U_k^\vee + \mathcal{H}_k + \delta_k^\sigma, \\ J_k^{\text{action}} &= 2 \sum_{N=q-1}^{2q-1} w_N \log \epsilon_N^{\text{ar}} = \Psi_k + C_k^{\text{budget}}, \\ L_J^\vee &= L_k^\vee + \mathcal{H}_k + \delta_k^\sigma + C_k^{\text{budget}}, & U_J^\vee &= U_k^\vee + \mathcal{H}_k + \delta_k^\sigma + C_k^{\text{budget}}, \\ 4q \log \mathcal{L}_{h,k} &\leq J_k^{\text{action}} \in [L_J^\vee, U_J^\vee], & \mathcal{L}_{h,k} &\leq \min_N \epsilon_N^{\text{ar}} \leq \prod_N (\epsilon_N^{\text{ar}})^{w_N/(2q)} \leq e^{U_J^\vee/(4q)}. \end{aligned} \quad (\text{PBR15})$$

The weights are $w_{q-1} = w_{2q-1} = 1$ and $w_N = 2$ in between, so $\sum w_N = 2q$. The allowance ϵ_N^{ar} is the original rank-two action radius and $\mathcal{L}_{h,k} = 2\delta e(k+1)\lfloor(k+1)^2/4\rfloor$, with $e = 1$ in this section. MCE's action identities still hold for every original multiplicity. RFX gives $\phi_N(0) = 0$ in the original untilted source, yielding the displayed exact geometric mean. It retains $z_B = -z_K$, and hence the mixed term $a_K d_B + a_B d_K + 2|z_K|^2$. The full generic real-tilt MCE identity with $\log(\epsilon_N^2 + \phi_N^2)$ and the exact MCP derivative remain unchanged. Addition to PBR13 proves the signed bounds; addition of the complete budget proves the action bounds. Applying the same-class lower radius at each degree proves the geometric-mean inequalities. The whole reference boundary, monic window, two contraction penalties and action leakage are retained.

In particular the explicit coarse PEC32 entries become

$$\begin{aligned} A_{K,k,k} - D_k^+ - (v - t_0)\mathcal{L}_k + \mathcal{H}_k + \delta_k^\sigma &\leq \Psi_k \\ &\leq A_{K,k,k} + E_{k,k}^\vee - D_k^- + v\mathcal{L}_k + \mathcal{H}_k + \delta_k^\sigma. \end{aligned} \quad (\text{PBR16})$$

The exact-low entries of PBR13 can strengthen these entries; every older bound is still in the intersection. This is an actual control interval in the original signed direction. It does not remove C_k^{budget} from the action inequality.

70.6 Actual support, the common-factor map, and the retained residual

Let $a_\pm, b_\pm$ be the actual extrema of the nonzero coefficients of A , and let $d_a = a_+ - a_-$, $d_b = b_+ - b_-$. The complete RCF/ACR maps give

$$\begin{aligned} \mathcal{I}_G &= ([-a_-, k - a_+] \times [-b_-, k - b_+]) \setminus [0, k - 8]^2, \quad g_A = (k + 1 - d_a)(k + 1 - d_b) - q', \\ \rho_G : V_A &\rightarrow \mathbb{C}[S']_{<g_A}, \quad V_0 = \ker \rho_G, \quad W = \Xi(V_0), \\ \rho_G I_K &= W_G^{-1} D_G C_G^\top \pi^\top, \quad B_G = \pi C_G, \quad t_G = \dim(B_\Xi/W) = g_A - \text{rank } B_G. \end{aligned} \quad (\text{PBR17})$$

All columns of C_G are the actual shifted coefficient vectors $f_{ij} = \sum_{a,b} a_{ab} e_{i+a, j+b}$ for $(i, j) \in \mathcal{I}_G$. The root Vandermonde W_G , evaluated-polynomial diagonal $D_G = \text{diag}(\chi'(\tau_\ell)^{-1})$, and original exterior selector π are the full ACR11–13 matrices. Here χ' is the original lower-grid polynomial; the prime does not denote differentiation. The exact morphism from V_A to the last quotient sends p to its class modulo $V_0 + K$. The first isomorphism theorem identifies it with $\mathbb{C}[S']_{<g_A}/\rho_G(K)$; composing with Ξ identifies it with B_Ξ/W . The displayed matrix for $\rho_G I_K$ has the same rank as B_G , proving the last formula with all actual coefficients. This retains the original linear relation quotient; RCF's ideal hull does not replace its denominator. The full common-factor metric comparison is RCM23–27 and its actual coefficient-rank specialization is ACR18–23.

LBRE proves the exclusion of all four boundary rulings for the actual limiting two-plane, with its full coefficient margin. ASR uses the actual entire boundary coefficient functions to produce the explicit radius $R_{\text{supp}} = \max\{R_*, 1, 2C_1/m_1, \dots, 2C_4/m_4\}$, where $m_j > 0$ is the larger of the two limiting coordinate moduli and C_j is the full convergent absolute tail sum. For $|u| \geq R_{\text{supp}}$ each selected pair has a nonzero coordinate by the bound $C_j/|u| \leq m_j/2$. On the remaining compact reciprocal-period disk, each chosen entire coordinate is nonzero at zero and consequently has finitely many zeros. Taking the actual common zeros of each three-coordinate boundary vector gives exactly ASR18's finite exception set $\mathcal{U}_\partial^*$ in $\{R_* \leq |u| < R_{\text{supp}}\}$. Thus, on the original arithmetic period domain outside this finite set,

$$\begin{aligned} (a_-, a_+, b_-, b_+) &= (0, 8, 0, 8), \quad d_a = d_b = 8, \quad g_A = t_G = 0, \quad V_0 = V_A, \quad W = B_\Xi, \\ B_\Xi/W_0 &\simeq Y_N/\text{im } Z_k, \quad \dim W_0 = v - t_0, \quad \dim(B_\Xi/W_0) = j - v + t_0. \end{aligned} \quad (\text{PBR18})$$

The first line follows by substitution into PBR17: the target of ρ_G is zero, so its kernel is all of V_A . The second line is the separate exact map PBR6–7. In particular the full residual image $W = B_\Xi$ is still equipped with $Q_{\Xi, N}^{\text{fix}}$, its low subspace and its complementary dual deficit. Neither t_0 nor either metric return is forced to vanish by $g_A = t_G = 0$. On the finite exceptional set the literal ACR rank matrices and RCM metric comparison remain; PBR2–16 did not assume full support and remain on their original hypotheses.

Finally the complete OVR/ECL outer identity and the proved finite errors imply

$$\begin{aligned} A_{K,k,s} + A_{0,k,s} &= \Delta q \mathfrak{C}_\partial + o(kq), \quad \frac{A_{K,k,s} + A_{0,k,s}}{kq} \longrightarrow 16\mathfrak{C}_\partial, \\ 21.66851180520 &< 16\mathfrak{C}_\partial < 21.66851180521, \quad \frac{\Psi_k}{kq} = \frac{A_{K,k,1}}{kq} - \frac{\mathcal{C}_\Gamma}{4} + o(1). \end{aligned} \quad (\text{PBR19})$$

Indeed PBR4, PBR7–9 and the complete FEX36 allowance give $F_K^{(s)} = A_{K,k,s} + o(kq)$ and $\Phi_{k,s} = A_{0,k,s} + o(kq)$. Substitute these equalities in the original evaluated outer sum. The second line uses

the certified ECL coefficient in the same original EIQ variables. For the last equality use the original AMT source-order agreement and the proved signed Gamma return $\mathcal{H}_k/(kq) \rightarrow -\mathcal{C}_\Gamma/4$, while the original arithmetic correction is $o(kq)$. All these are the accepted complete proof providers, preserved in the current full source. This determines the sum of the two specified spectral deficits and the exact signed receiver in terms of the kernel one; it does not assign an individual leading value to either deficit. OER15–16’s actual q -scale Gamma interval, including its strictly positive radius and both unequal finite endpoint errors, is unchanged. The next analytic quantity in this receiver is the actual deficit distribution in PBR5 and its complete action comparison PBR15.

71 Exact low-jet reconstruction and its original quotient spectrum

Retain the simple-quartet fixed-period hypotheses of RDS, DNE and PEC. The original conductor and all its coefficients remain fixed as $k = 4l + 1 \geq 9$ varies. Set

$$\begin{aligned} q &= (k+1)^2, & q' &= (k-7)^2, & \Delta &= q - q', & N &= q - 1, \\ d &= q - v - 1, & n &= d + 1 = q - v, & \mathbf{m} &= q - q' - v, \end{aligned} \quad (\text{LJR1})$$

with $0 \leq v \leq 80$, $\mu_v \neq 0$, and the same centers $c = k/2$, $c' = c - 4$. The original source order is $s \in \{1, k\}$, $b = s/2$, $M_s = (2\pi)^{s/2}$ and $h_a = M_s a! (b)_a$. The roots are $\beta_{ij} = c + (2i - k)\delta + i(2j - k)\gamma$ and the analogous lower roots β'_{ij} , with $0 < \delta < 1/2$, $\gamma > 2$. Write $\sigma = (\delta^2 + \gamma^2)^{1/2}$.

71.1 The exact finite reconstruction and its inverse

Let $\phi_a(S') = p_a((S' - c')/i)/\sqrt{h_a}$ be the complete original orthonormal source frame through degree d . Define B_s and the original upper Vandermonde by

$$S'^a = \sum_{r=0}^d (B_s)_{ra} \phi_r(S'), \quad \mathbb{V}_{j,(a,b)} = \beta_{ab}^j \quad (0 \leq j < q, \ 0 \leq a, b \leq k). \quad (\text{LJR2})$$

These matrices are not assigned new source norms. The coefficient matrix B_s is upper triangular, with diagonal $i^a \sqrt{h_a}$, since the leading coefficient of ϕ_a is $i^{-a}/\sqrt{h_a}$. In particular $|\det B_s|^2 = \prod_{a=0}^d h_a$. For an explicit formula, let J_b be the real symmetric $(d+1)$ -by- $(d+1)$ Jacobi matrix with adjacent entries $\sqrt{(r+1)(b+r)}$. Then

$$(B_s)_{ra} = \sqrt{M_s} [(c'I + iJ_b)^a]_{r0}. \quad (\text{LJR3})$$

The original three-term recurrence proves this formula by repeated multiplication by $S' = c' + iy$. A path of length at most d starting at degree zero never needs a degree above d , so the finite Jacobi truncation is exact in all the displayed columns.

For a vector $w = (w_0, \dots, w_d)^\top$ define the complex-linear functional Λ_w by $\Lambda_w(\phi_a) = w_a$. Its first n ordinary moments are $B_s^\top w$. Define the full moment-convolution matrix

$$(J_E)_{ja} = \binom{j}{a} \mu_{j-a} \quad (0 \leq j < q, \ 0 \leq a \leq d), \quad \mathbf{R}_s = \mathbb{V}^{-1} J_E B_s^\top. \quad (\text{LJR4})$$

Terms with $a > j$ are zero. The first v rows of J_E vanish, and its remaining square block $\tilde{J}_E$ is lower triangular with diagonal $\binom{v+a}{a} \mu_v$, $0 \leq a \leq d$. Therefore $\mathbf{R}_s$ is an isomorphism from $\mathbb{C}^n$ onto the coefficient space of the original $\mathcal{N}_v$.

To verify the map exactly, let F be any formal series with $F^{(a)}(0) = (B_s^\top w)_a$ for $a \leq d$. Leibniz' rule gives

$$(E_A F)^{(j)}(0) = \sum_{a=0}^d \binom{j}{a} \mu_{j-a} F^{(a)}(0) \quad (0 \leq j < q). \quad (\text{LJR5})$$

No unknown derivative of F enters, because $\mu_{j-a} = 0$ when $a > j - v$. The upper Vandermonde then reconstructs the unique upper exponential polynomial $\mathcal{N}$ with precisely these first q jets. This proves LJR4 as the actual map of original polynomial functionals to original exponential coefficients.

Let κ_r denote the coefficients of the finite inverse series of $B_\infty(z) = v! E_A(z)/(\mu_v z^v)$, so $B_\infty(z)^{-1} = \sum_{r \geq 0} \kappa_r z^r$ near zero. For $\nu \in \mathcal{N}_v$ write $u = \mathbb{V}\nu$. The full inverse is

$$\begin{aligned} f_r &= \frac{v!}{\mu_v} r! \sum_{a=0}^r \kappa_{r-a} \frac{u_{v+a}}{(v+a)!} \quad (0 \leq r \leq d), \\ \mathbf{R}_s^{-1} \nu &= B_s^{-\top} (f_0, \dots, f_d)^\top. \end{aligned} \quad (\text{LJR6})$$

Indeed multiplication of exponential Taylor coefficients by $E_A = (\mu_v/v!)z^v B_\infty$ gives precisely this triangular inverse. The constants $\kappa_0, \dots, \kappa_d$ are determined by the finite recurrence $\kappa_0 = 1$, $\kappa_r = -\sum_{a=1}^r [v!\mu_{v+a}/((v+a)!\mu_v)]\kappa_{r-a}$. Thus every moment, factorial and inverse coefficient is supplied.

71.2 Both exact quotient kernels and their metrics

Let $V_{\text{lo},s}$ be the n -by- q' matrix $(V_{\text{lo},s})_{r,(a,b)} = \phi_r(\beta'_{ab})$, and let C be the entire original primary conductor. Then

$$R_s V_{\text{lo},s} = C^\top. \quad (\text{LJR7})$$

For each column, its functional is evaluation at the corresponding lower root, so $F(z) = e^{\beta'_{ab}z}$. The original root-sum identity gives $E_A(z)e^{\beta'_{ab}z} = \sum_{r,t} a_{rt} e^{\beta_{a+r,b+t}z}$. Its coefficient column is exactly $C^\top$, and LJR5 reconstructs it, proving LJR7. Both matrices have rank q' , by the lower Vandermonde and the original injectivity of the conductor transpose.

Consequently LJR4 induces the exact isomorphism

$$\bar{R}_s : \mathbb{C}^n / \text{im } V_{\text{lo},s} \longrightarrow \mathcal{N}_v / \text{im } C^\top. \quad (\text{LJR8})$$

Its quotient kernel is zero: if $R_s w = C^\top \eta$, then LJR7 and injectivity give $w = V_{\text{lo},s} \eta$. Its inverse is induced by LJR6. The first space has the actual native Gamma dual quotient norm, using the Euclidean norm on the orthonormal functional-value coordinates w . The second has the unchanged original exponential coefficient quotient norm.

Use precisely DNE's fixed Z_{exp} on the right and set $W_s = R_s^{-1} Z_{\text{exp}}$ on the left. In these paired frames,

$$\begin{aligned} D_s &= W_s^* [I - V_{\text{lo},s} (V_{\text{lo},s}^* V_{\text{lo},s})^{-1} V_{\text{lo},s}^*] W_s = \mathcal{D}_{q-1}^{(s)}, \\ G &= Z_{\text{exp}}^* [I - C^\top (\bar{C} C^\top)^{-1} \bar{C}] Z_{\text{exp}} = G_{\text{exp}}. \end{aligned} \quad (\text{LJR9})$$

The first equality with DNE follows because LJR6 gives exactly the first n orthonormal values of $\Lambda_{\mathcal{N}_a/E_A}$; the allowed extensions are all lower evaluations by LJR7. Both projection formulas are attained minimum norms. In particular the reconstruction is represented by the identity from $(\mathbb{C}^m, D_s)$ to $(\mathbb{C}^m, G)$, not by an asserted isometry of these two different forms.

Let λ_a be the relative eigenvalues of (D_s, G) as in PEC12. The squared singular values of LJR8, in decreasing order, are λ_a^{-1} with the order reversed. Thus

$$\log \det(\bar{R}_s^* \bar{R}_s) = \log \frac{\det G}{\det D_s}, \quad \sum_a \log^+ \sigma_a(\bar{R}_s)^2 = \sum_a \log^+ (\lambda_a^{-1}). \quad (\text{LJR10})$$

The adjoint and singular values here use the two specified quotient norms.

For an actual onto map B_0 , or an actual injective map J_0 , the identity in these coordinates induces the paired quotients or restrictions with complete forms

$$\begin{aligned} D_{B_0} &= (B_0 D_s^{-1} B_0^*)^{-1}, & G_{B_0} &= (B_0 G^{-1} B_0^*)^{-1}, \\ D_{J_0} &= J_0^* D_s J_0, & G_{J_0} &= J_0^* G J_0. \end{aligned} \quad (\text{LJR11})$$

The exact common kernels and targets are transported by LJR8. For $B_0 = B_K = Z_k^\top A_\ell$, DNE9 gives $D_{B_K} = (\widehat{H}_{K,q-1}^{(s)})^{-1}$. For the full annihilator frame $J_0 = A_\ell^{-1} P_Y^\top$ from PEC24, $D_{J_0} = \overline{Q_{0,q-1}^{-1}}$. These identities retain the original low-quotient corrections of FGC when compared with the original kernel and residual.

71.3 The complete full-space and quotient Jacobians

Let $\mathbb{V}_{<v}$ be the first v rows of $\mathbb{V}$ and put $\mathcal{J}_v = \det(\mathbb{V}_{<v} \mathbb{V}_{<v}^*)$, with $\mathcal{J}_0 = 1$. Then the squared full-space Jacobian is

$$\det(R_s^* R_s) = \frac{\mathcal{J}_v}{|\det \mathbb{V}|^2} \left| \frac{\mu_v}{v!} \right|^{2n} \prod_{a=0}^d \left(\frac{(v+a)!}{a!} \right)^2 \prod_{a=0}^d h_a. \quad (\text{LJR12})$$

Indeed $R_s = \mathbb{V}^{-1} P_v \tilde{J}_E B_s^\top$, where P_v inserts the last n coordinates. The complementary principal-minor identity for the positive matrix $\mathbb{V}\mathbb{V}^*$ gives $\det(P_v^*(\mathbb{V}\mathbb{V}^*)^{-1}P_v) = \mathcal{J}_v/|\det \mathbb{V}|^2$. For completeness this identity follows by block Gaussian elimination: the determinant of a block matrix equals the determinant of its first principal block times the determinant of its Schur complement, while the bottom block of its inverse is the inverse Schur complement. All principal blocks are positive. The determinants of $\tilde{J}_E$ and B_s , computed above, prove LJR12.

The exact restriction Jacobian on the subspace in LJR7 is $\det(\bar{C}C^\top)/\det(V_{\text{lo},s}^* V_{\text{lo},s})$. Completing this subspace to a full source frame, and its image to the transported target frame, factors the full Gram into restriction and quotient Schur determinants. It follows that

$$\mathfrak{J}_{k,s} := \frac{\det G}{\det D_s} = \det(R_s^* R_s) \frac{\det(V_{\text{lo},s}^* V_{\text{lo},s})}{\det(\bar{C}C^\top)}. \quad (\text{LJR13})$$

No orthogonality of a chosen coefficient complement is assumed.

Let $\mathbb{V}'$ be the lower-root evaluation Vandermonde of degrees $0, \dots, q' - 1$, and let $H_{\psi,s}$ be the full $\mathbf{m}$ -by- $\mathbf{m}$ original Gram of $\chi' S'^a$, $0 \leq a < \mathbf{m}$. Then

$$\det(V_{\text{lo},s}^* V_{\text{lo},s}) = \frac{|\det \mathbb{V}'|^2 \det H_{\psi,s}}{\prod_{a=0}^d h_a}, \quad (\text{LJR14})$$

and therefore

$$\boxed{\mathfrak{J}_{k,s} = \frac{\mathcal{J}_v |\det \mathbb{V}'|^2}{|\det \mathbb{V}|^2 \det(\bar{C}C^\top)} \left| \frac{\mu_v}{v!} \right|^{2n} \prod_{a=0}^d \left(\frac{(v+a)!}{a!} \right)^2 \det H_{\psi,s}.} \quad (\text{LJR15})$$

To prove LJR14, split the native polynomial space into the low monomials through degree $q' - 1$ and the ideal columns $\chi' S'^a$. Their complete monomial coefficient matrix has determinant one because χ' is monic. The full source Gram determinant is $\prod h_a$, since the original monic S' -polynomials have squared norms h_a and their phases have modulus one. Schur elimination therefore gives the low-remainder quotient determinant $\prod h_a / \det H_{\psi,s}$. Evaluation at the lower roots is precisely $\mathbb{V}'$ on that remainder. The inverse quotient determinant is thus $|\det \mathbb{V}'|^2 \det H_{\psi,s} / \prod h_a$. In an orthonormal frame the inverse evaluation quotient Gram is the conjugate of $V_{\text{lo},s}^* V_{\text{lo},s}$; their determinants agree because they are positive real. This proves the identity with its full original source mass before cancellation in LJR15.

71.4 Finite reconstruction bounds on the actual grid

Put $A_{\text{abs}} = \sum_{r,t=0}^8 |a_{rt}|$ and $H_D = \sum_{a=1}^D 1/a$, with $H_0 = 0$. For the original roots define

$$\begin{aligned} \mathfrak{d}_{ab} &= \prod_{(i,j) \neq (a,b)} (\beta_{ab} - \beta_{ij}), & L_{ab}(S) &= \mathfrak{d}_{ab}^{-1} \prod_{(i,j) \neq (a,b)} (S - \beta_{ij}), \\ \Pi_{k,s} &= \prod_{a=1}^{q-1} [2\sqrt{a(b+a)} + k\sigma], \\ M_{k,s}^{\text{rec}} &= \max \left\{ 1, A_{\text{abs}} e^{8\sigma(H_{q-1}+2)} \sqrt{M_s} \Pi_{k,s} \left(\sum_{a,b=0}^k |\mathfrak{d}_{ab}|^{-2} \right)^{1/2} \right\}. \end{aligned} \quad (\text{LJR16})$$

The derivative product $\mathfrak{d}_{ab}$ is distinguished from the complete lower polynomial χ' . It retains its actual complex sign in L_{ab} . Then

$$\|R_s\| \leq M_{k,s}^{\text{rec}}, \quad \|\bar{R}_s\| \leq M_{k,s}^{\text{rec}}. \quad (\text{LJR17})$$

Here all norms have precisely the source and target metrics of LJR4 and LJR8.

To prove the first bound, evaluation of the reconstructed functional on L_{ab} gives $\nu_{ab} = \Lambda_w(T_A L_{ab})$. Its coefficient row in the w coordinates therefore has Euclidean norm $\|T_A L_{ab}\|_{s,c'}$. This is an exact row identity, not a comparison with an unweighted polynomial coefficient norm. In the original coordinates, $T_A p(c' + iy) = \sum a_{rt} p(c + iy + (b_{rt} - 4))$. Since $b_{rt} - 4 = (2r - 8)\delta + i(2t - 8)\gamma$ has modulus at most 8σ , the exact finite Taylor translation and the original derivative estimate $\|\partial_S\|_{\leq q-1, s, c} \leq H_{q-1} + 2$ give the factor $A_{\text{abs}} e^{8\sigma(H_{q-1} + 2)}$. The factor $-i$ in $\partial_S = -i\partial_y$ has modulus one and is retained in the Taylor map.

Multiplication by y from degree at most $a - 1$ to degree at most a is the sum of the two original Jacobi shifts, each bounded by $\sqrt{a(b+a)}$. Therefore each factor $S - \beta_{ij}$ has multiplication norm at most $2\sqrt{a(b+a)} + |c - \beta_{ij}|$ at this step. Starting from the original constant norm $\sqrt{M_s}$ proves $\|L_{ab}\|_{s,c} \leq \sqrt{M_s} \Pi_{k,s} / |\mathfrak{d}_{ab}|$. The sum of the squared row norms is the Hilbert–Schmidt norm squared of R_s , proving the first inequality. For the quotient, choose the attained minimum representative on the left and project its image orthogonally off $\text{im } C^\top$. This proves the second inequality, with exactly the denominator in LJR8.

Every spacing in the denominator admits the exact factorization

$$|\mathfrak{d}_{ab}| = (2\delta)^k a! (k-a)! (2\gamma)^{k(k+1)} [b!(k-b)!]^{k+1} \cdot \prod_{i=0}^k \prod_{\substack{0 \leq j \leq k \\ j \neq b}} \left(1 + \frac{\delta^2(a-i)^2}{\gamma^2(b-j)^2}\right)^{1/2}. \quad (\text{LJR18})$$

In this formula only, b is the second root index; the source parameter inside $\Pi_{k,s}$ remains $s/2$. Separate the k factors in the row $j = b$, and factor $2\gamma|b-j|$ from every other row to prove LJR18. The elementary bounds $a!(k-a)! = k!/\binom{k}{a} \geq (k/(2e))^k$ and the same bound for b give

$$|\mathfrak{d}_{ab}| \geq (k/e)^{q-1} \delta^k \gamma^{k(k+1)}, \quad \Pi_{k,s} \leq [k^2(3+\sigma)]^{q-1}. \quad (\text{LJR19})$$

The factorial estimate follows from $\sum_{a=1}^k \log a \geq \int_1^k \log t \, dt$ and $\binom{k}{a} \leq 2^k$. For the second inequality use $2\sqrt{a(a+b)} \leq 2a+b$, $b \leq k/2$, and $2(1+1/k)^2 + (1/2+\sigma)/k \leq 3+\sigma$ for $k \geq 9$. Consequently

$$\begin{aligned} \log M_{k,s}^{\text{rec}} &\leq \max\{0, (q-1) \log k + \log A_{\text{abs}} + 8\sigma(H_{q-1} + 2) + \tfrac{1}{2} \log q + \tfrac{1}{2} \log M_s \\ &\quad + (q-1) \log[e(3+\sigma)] - k \log \delta - k(k+1) \log \gamma\} \\ &= q \log k + O_{\text{actual}}(q). \end{aligned} \quad (\text{LJR20})$$

The last equality denotes an upper bound with that order of error. It is uniform for both original source orders, since $\log M_s = (s/2) \log(2\pi) = O(k)$. The finite definition LJR16, with every actual spacing and all conductor coefficients, remains the bound used below.

71.5 A direct estimate of the nested determinant difference

Let $\mathcal{I} = \{4, \dots, k-4\}^2$ and let $\mathcal{O}$ be its complement in the upper index grid. Then $|\mathcal{I}| = q'$, $|\mathcal{O}| = \Delta$, and $\beta_{i+4, j+4} = \beta'_{ij} + 4$. The coefficient map $(S' + 4)^a = \sum_{r \leq a} \binom{a}{r} 4^{a-r} S'^r$ is upper triangular with diagonal one. Hence the Vandermonde on the central upper roots has exactly the lower Vandermonde determinant. Let $\mathcal{P}_{\mathcal{O}}$ denote the unordered distinct pairs with at least one index in $\mathcal{O}$. Thus

$$\begin{aligned} 2M_{\mathcal{O}} &= 2|\mathcal{P}_{\mathcal{O}}| = q(q-1) - q'(q'-1) = \Delta(q+q'-1), \\ \mathscr{D}_k &:= \log \frac{|\det \mathbb{V}|^2}{|\det \mathbb{V}'|^2} = 2 \sum_{\{x,y\} \in \mathcal{P}_{\mathcal{O}}} \log |\beta_x - \beta_y|. \end{aligned} \quad (\text{LJR21})$$

This is cancellation of the identical central-central factors in the exact Vandermonde products.

For $d(x, y) = \|x - y\|_\infty$ one has $2\delta d(x, y) \leq |\beta_x - \beta_y| \leq 2\sigma d(x, y)$. For each fixed x , the number of other grid indices within distance t is at most $4t^2 + 4t$, for $1 \leq t \leq k$. The exact identity

$\log(k/d) = \int_d^k dt/t$ consequently gives $\sum_{y \neq x} \log(k/d(x, y)) \leq 2q - 8$. Summing this nonnegative estimate over the Δ outer indices counts every pair in LJR21 at least once. It proves

$$\begin{aligned} 2M_{\mathcal{O}} \log k + 2M_{\mathcal{O}} \log(2\delta) - 2\Delta(2q - 8) &\leq \mathcal{D}_k \\ &\leq 2M_{\mathcal{O}} \log k + 2M_{\mathcal{O}} \log(2\sigma). \end{aligned} \quad (\text{LJR22})$$

In particular $\mathcal{D}_k = \Delta(q + q' - 1) \log k + O_{\text{actual}}(kq)$. This error is obtained from the affected pairs themselves, not by subtracting two estimates with errors of order q^2 .

71.6 The complete remaining native Gram

The original Gram $H_{\psi, s}$ in LJR15 is positive definite: a nonzero polynomial has nonzero modulus off a finite set, and the original Gamma density is positive on the real line. The following bounds include its full mass and apply at every $k \geq 9$ under LJR1. Define

$$\begin{aligned} R_{\psi, k} &= \max\{(k - 8)\sigma, |c'|\}, \quad T_* = \max\{1, 2d + s/2 + R_{\psi, k}\}, \\ Y &= 2(q + k\sigma + 1), \quad h = 1/q, \quad L_h = \frac{\pi}{\Gamma(3/4)^2} (\sin h)^{3/2}, \\ \beta_s &= \frac{M_s}{\sqrt{2} \Gamma(s/2)}, \quad A_{k, s} = \beta_s L_h e^{-(\pi/2 + h)(Y + 1)}, \\ \mathcal{H}_m &= \prod_{a=0}^{m-1} [(2a + 1) \binom{2a}{a}^2]^{-1}. \end{aligned} \quad (\text{LJR23})$$

Then the finite two-sided estimate is

$$(A_{k, s} q^{2q'})^m \mathcal{H}_m \leq \det H_{\psi, s} \leq M_s^m T_*^{2mq' + m(m-1)}. \quad (\text{LJR24})$$

For the upper bound, $\chi' S'^a$ has the q' actual lower roots and a additional zero roots. All are at distance at most $R_{\psi, k}$ from c' . Applying the same original Jacobi multiplication bound as in LJR17 through degree $q' + a \leq d$ gives its norm at most $\sqrt{M_s} T_*^{q' + a}$. Hadamard's Gram determinant inequality proves the upper side.

For the lower bound, the exact original Gamma recurrence and the global density estimate give

$$\begin{aligned} dm_s(y) &= \beta_s \prod_{j=0}^{(s-1)/4-1} (y^2 + (2j + \tfrac{1}{2})^2) d\sigma(y), \\ \frac{d\sigma}{dy}(y) &\geq L_h e^{-(\pi/2 + h)|y|}. \end{aligned} \quad (\text{LJR25})$$

These are the complete identities OSP12 and OFV9. To make the lower-density constant explicit, Euler's integral rotated through $\pi/2 - h$ gives

$$|\Gamma(3/4 - iy/2)|^2 \leq \Gamma(3/4)^2 (\sin h)^{-3/2} e^{-(\pi/2 - h)|y|}.$$

The exact reflection product with $|\Gamma(1/4 + iy/2)|^2$ is $2\pi^2 / \cosh(\pi y)$. Dividing and using $\cosh(\pi y) \leq e^{\pi|y|}$ proves the displayed lower density, including the factor $1/(2\pi)$ in $d\sigma$. The recurrence prefactor β_s follows by applying $\Gamma(z + 1) = z\Gamma(z)$ exactly $(s - 1)/4$ times and clearing the factors 4^{-1} from the squared moduli. Thus no original source mass has been discarded.

On the full interval $[Y, Y + 1]$, every factor of $\chi'(c' + iy)$ has modulus at least $y - (k - 8)\sigma \geq q$, and the polynomial density multiplier in LJR25 is at least one because $y \geq 1$. Therefore the weighted density is at least $A_{k, s} q^{2q'}$. The Gram of $(c' + iy)^a$ on this interval is obtained from the Gram of t^a on $[0, 1]$ by the translation $y = Y + t$ and coefficients with diagonal i^a . This map has determinant of modulus one.

For completeness, Rodrigues' polynomial $L_a(t) = (a!)^{-1} (d/dt)^a [t^a (t - 1)^a]$ has leading coefficient $\binom{2a}{a}$. Integration by parts a times proves orthogonality to every lower power; all boundary terms

vanish. It also gives $\int_0^1 L_a(t)^2 dt = 1/(2a+1)$, since $\int_0^1 t^a(1-t)^a dt = (a!)^2/(2a+1)!$. Dividing by the square of the leading coefficient gives the monic norm in LJR23. Triangular orthogonal elimination therefore proves that the complete interval Gram determinant is exactly $\mathcal{H}_m$. The positive-form inequality on that interval proves the lower side of LJR24.

Both sides imply the useful direct scale

$$\log \det H_{\psi,s} = 2\mathfrak{m}q' \log q + O_{\text{actual}}(kq). \quad (\text{LJR26})$$

Here $\mathfrak{m} = O(k)$, $d = O(q)$ and $\log T_* = \log q + O_{\text{actual}}(1)$. Moreover $-\log \mathcal{H}_m \leq \mathfrak{m} \log(2\mathfrak{m}+1) + 2\mathfrak{m}(\mathfrak{m}-1) \log 2$, using $\binom{2a}{a} \leq 4^a$. Since $2h/\pi \leq \sin h \leq h$, $\log L_h = O(\log q)$. Writing $s/2 = 2\ell_s + 1/2$, $\Gamma(s/2) = \sqrt{\pi} \prod_{j=0}^{\ell_s-1} (j+1/2)$ shows $|\log \beta_s| = O(k \log(k+1))$. Finally $Y = O_{\text{actual}}(q)$. All errors in the logarithms of LJR24 are consequently $O_{\text{actual}}(kq)$; in particular the terms $k^2 \log q$ are smaller than kq .

71.7 The exact remainder and every paired exterior volume

Let a_* be the actual nonzero pivot coefficient chosen by the original lexicographic conductor proof. Its triangular q' -by- q' minor has determinant $a_*^{q'}$. Cauchy–Binet and the row and column sum bounds on the full convolution matrix give

$$|a_*|^{2q'} \leq \det(\bar{C}C^\top) \leq A_{\text{abs}}^{2q'}. \quad (\text{LJR27})$$

The matrix in this determinant is the conjugate of CC^* ; their positive real determinants agree. For the low-jet minor, Hadamard gives $\mathcal{J}_v \leq q^v [k(1/2 + \sigma)]^{v(v-1)}$. For $k \geq v-1$, choose the v consecutive roots with first index zero in its Cauchy–Binet sum. Their Vandermonde gives $\mathcal{J}_v \geq (2\gamma)^{v(v-1)} (\prod_{a=1}^{v-1} a!)^2$. At $v=0$ both bounds mean $\mathcal{J}_0 = 1$. Thus $\log \mathcal{J}_v = O_{\text{actual}}(\log(k+1))$. Also $0 \leq \sum_{a=0}^d \log((v+a)!/a!) \leq vn \log(q+v)$. The fixed period supplies the actual nonzero μ_v, a_* and A_{abs} in these statements; none is selected afresh.

Define the following exact, still unevaluated remainder:

$$\begin{aligned} \mathcal{R}_{k,s} := & \log \mathcal{J}_v - \mathcal{D}_k - \log \det(\bar{C}C^\top) + 2n \log |\mu_v/v!| \\ & + 2 \sum_{a=0}^d \log \frac{(v+a)!}{a!} + \log \det H_{\psi,s} - 2\mathfrak{m}q \log k. \end{aligned} \quad (\text{LJR28})$$

Every term is an original finite determinant, moment or root product already specified above. Then

$$\log \mathfrak{J}_{k,s} = 2\mathfrak{m}q \log k + \mathcal{R}_{k,s}, \quad |\mathcal{R}_{k,s}| = O_{\text{actual}}(kq). \quad (\text{LJR29})$$

For the bound substitute LJR22, LJR26 and LJR27 into LJR28. The only large logarithmic expression that remains is $2\mathfrak{m}q' \log q - \Delta(q+q'-1) \log k$. Since $\log q = 2 \log k + 2 \log(1+1/k)$ and $\Delta = \mathfrak{m} + v$, subtracting $2\mathfrak{m}q \log k$ from this expression gives exactly

$$[(-3\mathfrak{m}+v)\Delta + \Delta - 2vq] \log k + 4\mathfrak{m}q' \log(1+1/k). \quad (\text{LJR30})$$

The coefficient of $\log k$ is $O_{\text{actual}}(k^2)$; the last term is $O_{\text{actual}}(k^2)$. Both are $o(kq)$. This proves LJR29 by an explicit difference calculation.

Consider any fixed paired restriction, quotient, or restriction followed by quotient of LJR8, with rank t . Its two forms are the restrictions and attained quotients of D_s, G along the same specified algebraic flag. Write $\mathfrak{J}_{t,k,s}$ for its squared Jacobian. Then the exact finite bounds are

$$\log \mathfrak{J}_{k,s} - 2(\mathfrak{m}-t) \log M_{k,s}^{\text{rec}} \leq \log \mathfrak{J}_{t,k,s} \leq 2t \log M_{k,s}^{\text{rec}}. \quad (\text{LJR31})$$

To prove this for a flag $A \subset B$ in the common $\mathfrak{m}$ -dimensional coordinate space, factor each full Gram determinant successively into the restriction on A , the quotient B/A , and the quotient of the full space by B . The factors are their actual Schur determinants. Their Jacobians multiply to

$\mathfrak{J}_{k,s}$. Every restriction of the map has norm at most $M_{k,s}^{\text{rec}}$; every induced quotient does too, by the minimum-lift and projection proof of LJR17. The two complementary factors have total rank $\mathfrak{m} - t$. Bounding them above and dividing proves the lower inequality. Bounding the rank- t factor directly proves the upper inequality.

Uniformly over all these actual flags and $0 \leq t \leq \mathfrak{m}$,

$$\log \mathfrak{J}_{t,k,s} = 2tq \log k + O_{\text{actual}}(kq). \quad (\text{LJR32})$$

For a fully retained finite version put $u_{k,s} = \log M_{k,s}^{\text{rec}} - q \log k$ and $\mathcal{R}_{t,k,s} = \log \mathfrak{J}_{t,k,s} - 2tq \log k$. LJR31 gives $\mathcal{R}_{k,s} - 2(\mathfrak{m} - t)u_{k,s} \leq \mathcal{R}_{t,k,s} \leq 2tu_{k,s}$. LJR20 bounds $u_{k,s}$ above by $O_{\text{actual}}(q)$, which proves both sides of LJR32 using LJR29. No inverse coefficient norm or conditioning premise for the flag occurs.

In particular apply LJR31 to the actual B_K quotient and the actual J_0 annihilator restriction in LJR11. If p_t, n_t are their two positive logarithmic spectral parts in the orientation (D, G) , then exactly

$$n_{t,q-1} - p_{t,q-1} = \log \mathfrak{J}_{t,k,s} = 2tq \log k + \mathcal{R}_{t,k,s}, \quad 0 \leq p_{t,q-1} \leq \varepsilon_{q-1,s}^\vee = o(kq). \quad (\text{LJR33})$$

The last bound is PEC18 on precisely these forms. Thus $n_{t,q-1} = 2tq \log k + O_{\text{actual}}(kq)$. The exact remaining terms are $\mathcal{R}_{t,k,s}$ and $p_{t,q-1}$, with LJR28 and LJR31 giving finite controls. Their four-endpoint differences have not been assigned a coefficient: LJR33 proves the low-endpoint leading scale and supplies the literal maps and remainders to be carried into that next calculation.

72 First-endpoint reconstruction in the actual kernel and annihilator

Retain every original hypothesis, source order, metric, coefficient, root, mass and sign of PBR1–19. The following applies the complete LJR1–33 reconstruction at the actual first endpoint $N = q - 1$. The symbol t_0 continues to mean $\dim(K \cap L)$. Define the two ranks without changing that symbol:

$$t_K = r_K - t_0, \quad t_R = j - v + t_0, \quad t_K + t_R = \mathbf{m}. \quad (\text{LRR1})$$

PBR2 proves that $B_K = Z_k^\top A_\ell$ is onto rank t_K , and PBR3 proves that $J_0 = A_\ell^{-1} P_Y^\top$ is injective rank t_R . Indeed Z_k has the actual independent kernel columns, A_ℓ is invertible, $P_Y Z_k = 0$, and P_Y is onto with kernel $\text{im } Z_k$. Thus $\text{im } J_0 = \ker B_K$.

Let $R_s = \mathbb{V}^{-1} J_E B_s^\top$ be the full LJR4 reconstruction, including the original upper Vandermonde, moment convolution and original Gamma orthonormal-value matrix. LJR5 proves the exact first- q -jet identity by Leibniz' rule, and LJR7 gives $R_s V_{\text{lo},s} = C^\top$. It therefore induces the isomorphism $\bar{R}_s : \mathbb{C}^{q-v} / \text{im } V_{\text{lo},s} \rightarrow \mathcal{N}_v / \text{im } C^\top$. Its inverse is the full finite moment recurrence LJR6. In the paired frames $R_s^{-1} Z_{\text{exp}}$ and Z_{exp} , this map is the identity from $(\mathbb{C}^{\mathbf{m}}, \mathcal{D}_{q-1}^{(s)})$ to $(\mathbb{C}^{\mathbf{m}}, G_{\text{exp}})$, because the complete minimum-extension norms are exactly LJR9/PBR2.

Taking the quotient under the same onto B_K gives precisely the two forms $(D_{K,q-1}, G_K)$ of PBR2; restricting along the same J_0 gives precisely $(D_{0,q-1}, G_0)$ of PBR3. The quotient metric is obtained by minimizing over the full kernel of B_K in each metric, giving $(B_K D^{-1} B_K^*)^{-1}$ and $(B_K G^{-1} B_K^*)^{-1}$. Restriction gives $J_0^* D J_0$ and $J_0^* G J_0$ directly. Use $X = R$ for this residual restriction, with the exact dictionary $(D_R, G_R, p_R, n_R, A_R, P_R) = (D_0, G_0, p_0, n_0, A_0, P_0)$ from PBR3/5; t_0 still denotes only $\dim(K \cap L)$. Consequently their squared reconstruction Jacobians satisfy

$$\mathfrak{J}_{X,k,s} = \frac{\det G_X}{\det D_{X,q-1}}, \quad X = K, R, \quad (\text{LRR2})$$

$$\log \mathfrak{J}_{X,k,s} = n_{X,q-1} - p_{X,q-1} = 2t_X q \log k + \mathcal{R}_{X,k,s}.$$

The first equality uses the actual source and target norms of the reconstruction, in that direction. For the second, a relative eigenvalue μ of (D, G) contributes $-\log \mu = \log^+(1/\mu) - \log^+ \mu$ to the reconstruction determinant. Thus these are exactly the n, p of PBR5, with no inversion of their orientation.

Here every finite remainder is retained. Put $u_{k,s} = \log M_{k,s}^{\text{rec}} - q \log k$, where $M_{k,s}^{\text{rec}}$ is the complete finite LJR16 constant, including all actual root spacings and all conductor coefficients. Keep the exact full-space remainder LJR28:

$$\begin{aligned} \mathcal{R}_{k,s} &= \log \mathcal{J}_v - \mathcal{D}_k - \log \det(\bar{C} C^\top) + 2(q - v) \log |\mu_v / v!| \\ &\quad + 2 \sum_{a=0}^{q-v-1} \log \frac{(v+a)!}{a!} + \log \det H_{\psi,s} - 2\mathbf{m}q \log k, \\ \mathcal{R}_{k,s} - 2(\mathbf{m} - t_X)u_{k,s} &\leq \mathcal{R}_{X,k,s} \leq 2t_X u_{k,s}, \quad 0 \leq p_{X,q-1} \leq \varepsilon_{q-1,s}^\vee, \\ n_{X,q-1} &= 2t_X q \log k + \mathcal{R}_{X,k,s} + p_{X,q-1} = 2t_X q \log k + O_{\text{actual}}(kq). \end{aligned} \quad (\text{LRR3})$$

The meanings of the finite terms are literal: $\mathcal{J}_v = \det(\mathbb{V}_{<v} \mathbb{V}_{<v}^*)$ with $\mathcal{J}_0 = 1$, $\mathcal{D}_k = \log(|\det \mathbb{V}|^2 / |\det \mathbb{V}'|^2)$, and $H_{\psi,s}$ is the full original Gram of $\chi' S'^a$, $0 \leq a < \mathbf{m}$, retaining its Gamma mass. The prime on χ' means the lower-grid polynomial. LJR12–15 proves the full Jacobian by exact triangular and Schur factorizations. LJR31 factors its determinant along the same algebraic flag into the rank- t_X part and its complements. Every part has forward norm at most $M_{k,s}^{\text{rec}}$ by its actual minimum lift. Bounding the complementary product above gives the lower inequality for $\mathcal{R}_{X,k,s}$; bounding the selected product gives the upper inequality. PEC18 bounds the positive part on these very quotient and restriction forms. This proves every finite assertion in LRR3.

For the final order statement, LJR20 gives $u_{k,s} \leq O_{\text{actual}}(q)$ and LJR22–30 gives $|\mathcal{R}_{k,s}| = O_{\text{actual}}(kq)$ by the affected-root-pair calculation and the complete Gram bounds. Substitution into the two finite inequalities gives $|\mathcal{R}_{X,k,s}| = O_{\text{actual}}(kq)$ uniformly for $0 \leq t_X \leq \mathfrak{m}$; no lower bound for an unretained inverse norm is needed. PEC15 gives $\varepsilon_{q-1,s}^\vee = o(kq)$. Thus the last equality in LRR3 follows, while its full finite remainder and positive part remain in the preceding equality.

In the actual four-endpoint receiver this yields exactly

$$\begin{aligned} A_{X,k,s} &= 2t_X q \log k + \mathcal{R}_{X,k,s} + p_{X,q-1} + n_{X,q} - n_{X,2q-1} - n_{X,2q}, & X = K, R, \\ F_K^{(s)} &= 2t_K q \log k + \mathcal{R}_{K,k,s} + p_{K,q-1} + n_{K,q} - n_{K,2q-1} - n_{K,2q} + P_{K,k,s} - \mathcal{E}_s^{\text{flag}} + F_L^{(s)} - F_{W_0}^{(s)}, \\ \Phi_{k,s} &= F_{W_0}^{(s)} + 2t_R q \log k + \mathcal{R}_{R,k,s} + p_{R,q-1} + n_{R,q} - n_{R,2q-1} - n_{R,2q} + P_{R,k,s} + \delta_{0,k,s}. \end{aligned} \tag{LRR4}$$

The first equality substitutes LRR3 into the definition of A_X in PBR5. The next two substitute that same identity into PBR9 and PBR7, respectively. Hence both original low volumes, all three other endpoint deficits, every positive spectral part, and all unequal finite flag and transfer errors remain. PBR13–16 continues to use these exact A_X in its original source, arithmetic, signed and action intervals. The first endpoint has been evaluated at its leading scale; the remaining endpoint terms and the full kq -scale remainder in LRR3 have not been assigned an individual coefficient.

73 Compatible high jets and the original positive return

Retain all fixed original quartet, amplitude, period and branch data of LJR, DNE, PEC and OVR. In particular the arithmetic domain remains $|u| \geq R_*$, the actual period is fixed while $k = 4l + 1$ grows, and the source order is exactly $s = 1$ or $s = k$. All asymptotic uses of OFV retain its proved eventual threshold. Write

$$\begin{aligned} q &= (k+1)^2, & q' &= (k-7)^2, & n &= q-v, & \mathbf{m} &= q-q'-v, & d_0 &= q-v-1, \\ q-1 &\leq N \leq 2q, & D &= N-v, & e_N &= N-q+1, & D+1 &= n+e_N. \end{aligned} \quad (\text{HJR1})$$

The polynomial degree D is the actual native source degree after the conductor. The original centers remain $c = k/2$ and $c' = c - 4$. No change of s is made between different endpoints.

73.1 The exact compatible high-jet graph

Let $B_{s,D}$ be the original matrix of S'^a in the complete orthonormal Gamma frame $\phi_0, \dots, \phi_D$, as in LJR2-3, with $h_a = M_s a! (s/2)_a$ and $M_s = (2\pi)^{s/2}$. Define

$$\begin{aligned} (\mathbb{V}_N)_{j,(a,b)} &= \beta_{ab}^j \quad (0 \leq j \leq N), \\ (J_{E,N})_{ja} &= \binom{j}{a} \mu_{j-a} \quad (0 \leq j \leq N, 0 \leq a \leq D), \\ \tilde{J}_{E,N} &= (J_{E,N})_{\{v,\dots,N\},\{0,\dots,D\}}, \quad P_{v,N} : \mathbb{C}^{D+1} \longrightarrow \mathbb{C}^{N+1}. \end{aligned} \quad (\text{HJR2})$$

Here $P_{v,N}$ inserts v initial zero coordinates, and an out-of-range term in the convolution is zero. The square $\tilde{J}_{E,N}$ is lower triangular with diagonal $\mu_v \binom{v+a}{a}$ and is therefore invertible.

On the original coefficient subspace $\mathcal{N}_v$ put

$$\mathbf{F}_{N,s} = B_{s,D}^{-\top} \tilde{J}_{E,N}^{-1} P_{v,N}^\top \mathbb{V}_N, \quad \mathbf{E}_{N,s} = \mathbf{F}_{N,s} \mathbf{R}_s = \begin{pmatrix} I_n \\ H_{N,s} \end{pmatrix}. \quad (\text{HJR3})$$

The right factor $\mathbf{R}_s$ is the exact low reconstruction LJR4. Thus $\mathbf{F}_{N,s}$ takes original upper exponential coefficients to their original native functional values through degree D ; $\mathbf{E}_{N,s}$ extends the first n such values. For a literal formula for the inverse convolution, if $u_j = (\mathbb{V}_N \nu)_j$, then

$$f_a = \frac{v!}{\mu_v} a! \sum_{r=0}^a \kappa_{a-r} \frac{u_{v+r}}{(v+r)!}, \quad \mathbf{F}_{N,s} \nu = B_{s,D}^{-\top} (f_0, \dots, f_D)^\top. \quad (\text{HJR4})$$

The coefficients κ_a are all the actual inverse-symbol coefficients of LJR6, with their proved finite recurrence. The source coefficient transformation in HJR4 contains the original phase i^{-a} and every h_a .

To prove these statements, use the analytic germs

$$\mathcal{N}(z) = \sum \nu_{ab} e^{\beta_{ab} z}, \quad E_A(z) = (\mu_v/v!) z^v B_\infty(z).$$

The order condition $\nu \in \mathcal{N}_v$ makes the quotient analytic at zero. Formal Taylor multiplication gives HJR4 and HJR2, and the triangular $B_{s,D}^{-\top}$ changes ordinary moments to orthonormal functional values. Its first n moments and orthonormal values depend only on the first q jets u_j . LJR6 therefore proves that the top block of HJR3 is I_n . The bottom block $H_{N,s}$ is the specified remaining e_N rows.

For a fully intrinsic description, write $\chi(S) = S^q + \sum_{a=0}^{q-1} \chi_a S^a$. Define $\mathcal{U}_{N,s} \subset \mathbb{C}^{D+1}$ by requiring that $u = J_{E,N} B_{s,D}^\top w$ satisfy

$$u_{q+j} + \sum_{a=0}^{q-1} \chi_a u_{a+j} = 0, \quad 0 \leq j < e_N. \quad (\text{HJR5})$$

Then $\mathcal{U}_{N,s} = \text{im } E_{N,s}$, and its dimension is n . Indeed each original root is annihilated by χ , so HJR3 satisfies every recurrence. Conversely the first q entries of u , with its first v zeros, determine a unique ν by the invertible upper Vandermonde. The monic recurrences determine every later u_j uniquely and give $V_N \nu$. The invertible lower convolution block then determines w uniquely. This proves the graph equality, its dimension, and the independence of its e_N constraints. The free ambient later functional values have not been identified with upper exponential coefficients; the exact identification uses HJR5.

Let $V_{N,s}$ denote the $(D+1)$ -by- q' lower evaluation matrix $(V_{N,s})_{a,\beta'} = \phi_a(\beta')$, and let $V = V_{q-1,s}$. The complete lower-exponential identity gives

$$E_{N,s}V = V_{N,s}, \quad \mathcal{U}_{N,s}/\text{im } V_{N,s} \simeq \mathcal{N}_v/\text{im } C^\top. \quad (\text{HJR6})$$

For its proof use $\nu = C^\top \eta$ in HJR4: $\mathcal{N}/E_A$ is exactly the sum of the lower exponentials with coefficients η , at all orders. Since E is injective, its induced quotient kernel is exactly $\text{im } V$. This proves the typed quotient identification with the original denominators.

73.2 One lower-evaluation minimum over the complete window

Keep precisely DNE's fixed coefficient quotient frame Z_{exp} . Put $T_N = F_{N,s}Z_{\text{exp}}$, $T_0 = T_{q-1}$, and

$$\begin{aligned} A &= V^*V, & K_0 &= A^{-1}V^*T_0, & Z_0 &= T_0 - VK_0, & Q_0 &= VA^{-1/2}, \\ D_0 &= Z_0^*Z_0 = \mathcal{D}_{q-1}^{(s)}, & V^*Z_0 &= 0, & Q_0^*Q_0 &= I_{q'}. \end{aligned} \quad (\text{HJR7})$$

The matrices A and its positive square root are the actual lower-evaluation Gram, retaining the complete root and Gamma conditioning. No replacement evaluation metric is made. With $H = H_{N,s}$, define

$$P_N = HZ_0, \quad Q_N = HQ_0, \quad \mathcal{C}_N = (I_{e_N} + Q_NQ_N^*)^{-1/2}P_N. \quad (\text{HJR8})$$

At $e_N = 0$ all these matrices have zero rows. The exact native metric at every endpoint is

$$\boxed{\mathcal{D}_N^{(s)} = D_0 + P_N^*(I_{e_N} + Q_NQ_N^*)^{-1}P_N = D_0 + \mathcal{C}_N^*\mathcal{C}_N.} \quad (\text{HJR9})$$

In particular its increment is positive and has rank $\text{rank } P_N \leq \min\{\mathbf{m}, e_N\}$. The first endpoint update $N = q$ has rank at most one. The last two endpoints have $e_{2q-1} = q$ and $e_{2q} = q+1$, respectively.

Here is a proof with the complete minimizer. Subtract the fixed lower extension EVK_0 from $T_N = ET_0$. Every remaining allowed extension is $E(Z_0x + Q_0t)$, with one vector $t \in \mathbb{C}^{q'}$ used for both its low and high rows. Its squared norm is

$$x^*D_0x + \|t\|^2 + \|P_Nx + Q_Nt\|^2. \quad (\text{HJR10})$$

The minimum occurs at $t = -(I + Q_N^*Q_N)^{-1}Q_N^*P_Nx$. Completing the square leaves $x^*D_0x + x^*P_N^*[I - Q_N(I + Q_N^*Q_N)^{-1}Q_N^*]P_Nx$. Multiplication by $I + Q_NQ_N^*$ shows that the bracket is $(I + Q_NQ_N^*)^{-1}$, proving HJR9. DNE7 proves that these and only these lower evaluations are the permitted native extension differences. Thus HJR10 is exactly its full minimum, without separate minima on different row blocks.

73.3 The original kernel and annihilator updates

Write $B = B_K$ and $J = J_0$ for the actual maps in PEC19,24, so B is onto, J is injective, and $\text{im } J = \ker B$. Define the full initial forms and minimum lift by

$$\begin{aligned} D_{B,0} &= (BD_0^{-1}B^*)^{-1}, & D_{J,0} &= J^*D_0J, \\ L_0 &= D_0^{-1}B^*D_{B,0}, & W_0 &= J(D_{J,0})^{-1/2}. \end{aligned} \quad (\text{HJR11})$$

Then $BL_0 = I$, $BW_0 = 0$, $W_0^*D_0W_0 = I$, and $W_0^*D_0L_0 = 0$. These are literal original kernel and annihilator maps; the displayed W_0 is a coefficient matrix for $\ker B$ with its full D_0 metric. The induced exact updates are

$$\begin{aligned} D_{J,N} &= D_{J,0} + (\mathcal{C}_N J)^*(\mathcal{C}_N J), \\ D_{B,N} &= D_{B,0} + (\mathcal{C}_N L_0)^*[I + \mathcal{C}_N W_0 W_0^* \mathcal{C}_N^*]^{-1}(\mathcal{C}_N L_0). \end{aligned} \quad (\text{HJR12})$$

The first is restriction of HJR9. For the second, every B -lift of y is $L_0 y + W_0 t$. Its low squared norm is $y^* D_{B,0} y + \|t\|^2$, and its added squared norm is $\|\mathcal{C}_N L_0 y + \mathcal{C}_N W_0 t\|^2$. The same completion as HJR10 proves the second formula. It preserves both the original lower-evaluation denominator in $\mathcal{C}_N$ and the original kernel denominator in HJR12. Empty source, kernel or target ranks use their unique empty matrices and determinant one.

The exact identities DNE9 and PEC25 are

$$D_{B,N} = \overline{(\widehat{H}_{K,N}^{(s)})^{-1}}, \quad D_{J,N} = \overline{Q_{0,N}^{-1}}. \quad (\text{HJR13})$$

Thus HJR12 is a full original native kernel and complementary residual calculation, rather than a freely chosen coefficient compression.

73.4 Exact one-row updates and their allocation

The original orthonormal polynomial sequence is independent of its finite cutoff. At any M with $q-1 \leq M < 2q$, let $d = M - v$, and write T, V for the complete matrices through degree d . Their next rows, at degree $d+1$, are t, v_+ . Every entry of t is $\Lambda_{N_a/E_A}(\phi_{d+1})$, computed by HJR4 and the original monomial matrix $B_{s,d+1}$ through degree $d+1$; every entry of v_+ is $\phi_{d+1}(\beta')$. Set

$$\begin{aligned} A_M &= V^* V, \quad K_M = A_M^{-1} V^* T, \quad r_M = t - v_+ K_M, \\ \lambda_M &= 1 + v_+ A_M^{-1} v_+^* > 0, \quad h_M = \lambda_M^{-1/2} r_M. \end{aligned} \quad (\text{HJR14})$$

Then the exact update is

$$D_{M+1} = D_M + h_M^* h_M, \quad D_M = \mathcal{D}_M^{(s)}. \quad (\text{HJR15})$$

To prove this, subtract the attained old minimum from the new extension problem. The old residual is orthogonal to all old lower evaluations, so the remaining minimization for coefficient correction η is $\eta^* A_M \eta + \|r_M x - v_+ \eta\|^2$. Its minimizer is $(A_M + v_+^* v_+)^{-1} v_+^* r_M x$. The same identity used in HJR10 gives the remaining form $r_M^* r_M / (1 + v_+ A_M^{-1} v_+^*)$, proving HJR15. Every evaluation row has participated in the old A_M ; the update does not reset the lower-evaluation coefficient.

At this endpoint put $D_{B,M} = (B D_M^{-1} B^*)^{-1}$, $L_M = D_M^{-1} B^* D_{B,M}$ and $D_{J,M} = J^* D_M J$. The exact inverse-metric decomposition is

$$D_M^{-1} = J(D_{J,M})^{-1} J^* + L_M(D_{B,M})^{-1} L_M^*. \quad (\text{HJR16})$$

Indeed $[J, L_M]$ is a full coefficient frame whose Gram in D_M is $\text{diag}(D_{J,M}, D_{B,M})$. The off-diagonal blocks vanish because $BJ = 0$. Inverting that frame Gram proves HJR16. Define the two actual nonnegative scalar row energies

$$x_M = h_M J(D_{J,M})^{-1} J^* h_M^*, \quad y_M = h_M L_M(D_{B,M})^{-1} L_M^* h_M^*. \quad (\text{HJR17})$$

Both retain λ_M^{-1} from HJR14 and every original period-dependent kernel coordinate. The determinant lemma, proved by extending a nonzero vector to a basis or by expanding a rank-one determinant, now gives

$$\begin{aligned} g_M &= \log \frac{\det D_{M+1}}{\det D_M} = \log(1 + x_M + y_M), \\ g_{0,M} &= \log \frac{\det D_{J,M+1}}{\det D_{J,M}} = \log(1 + x_M), \\ g_{K,M} &= \log \frac{\det D_{B,M+1}}{\det D_{B,M}} = \log \left(1 + \frac{y_M}{1 + x_M} \right), \quad g_M = g_{0,M} + g_{K,M}. \end{aligned} \quad (\text{HJR18})$$

The restriction update gives the second equality directly. For the quotient use the one-row instance of HJR12: its added form is $(h_M L_M)^*(h_M L_M)/(1+x_M)$. Its determinant relative to $D_{B,M}$ proves the third. This proves the allocation of each full original row increment between the actual kernel quotient and its annihilator, with its complete positive denominator.

73.5 The four original signs and an exterior bound at scale kq

For X equal to the full space, K , or 0, define

$$I_{X,N} = \log \frac{\det D_{X,N}}{\det D_{X,q-1}}, \quad I_N = I_{\text{full},N}. \quad (\text{HJR19})$$

All these numbers are nonnegative and increase with N . At each N , completing J by any fixed right inverse of B factors the full Gram into its restriction and quotient. The fixed frame determinant cancels between endpoints, so

$$I_N = I_{K,N} + I_{0,N}, \quad I_{X,N} = \sum_{M=q-1}^{N-1} g_{X,M}. \quad (\text{HJR20})$$

In the sum for the full space $g_{\text{full},M} = g_M$. The source dual identities HJR13 therefore give exactly

$$\begin{aligned} \widehat{F}_k^{(s)} &= I_{K,2q-1} + I_{K,2q} - I_{K,q}, \\ F_{0,k,s} &= I_{0,2q-1} + I_{0,2q} - I_{0,q}, \\ \mathcal{T}_{k,s} &:= \widehat{F}_k^{(s)} + F_{0,k,s} = I_{2q-1} + I_{2q} - I_q. \end{aligned} \quad (\text{HJR21})$$

Equivalently every line is the weighted sum of its original row increments, with weights

$$w_M = \begin{cases} 1, & M = q-1 \text{ or } M = 2q-1, \\ 2, & q \leq M \leq 2q-2. \end{cases} \quad (\text{HJR22})$$

For example I_{2q-1} contains all rows through $2q-2$, I_{2q} adds the last row, and I_q removes one copy of the first. This proves all endpoint signs. Writing $a = I_q$, $b = I_{2q-1} - I_q$ and $c = I_{2q} - I_{2q-1}$ gives $\mathcal{T} = a + 2b + c$ with $a, b, c \geq 0$. In particular

$$0 \leq I_N \leq I_{2q} \leq \mathcal{T}_{k,s} \quad (q-1 \leq N \leq 2q). \quad (\text{HJR23})$$

To give its finite original bound, retain $\delta_{K,k,s} = \widehat{F}_k^{(s)} - F_K^{(s)}$ with $|\delta_{K,k,s}| \leq \mathcal{E}_{k,s}^{\text{FEX}}$, and precisely $\delta_{0,k,s}$ and $F_{\Xi(L)}^{(s)}$ from PEC29. The latter is the original restriction return called $F_{W_0}^{(s)}$ there. OAR3 and PEC29 give the exact equality

$$\mathcal{T}_{k,s} = \mathcal{F}_{\partial,k}^{(s)} + \mathfrak{e}_{k,s} + \delta_{K,k,s} - F_{\Xi(L)}^{(s)} - \delta_{0,k,s}. \quad (\text{HJR24})$$

The original finite signs are consequently

$$\begin{aligned} \max\{0, \mathcal{F}_{\partial,k}^{(s)} - e_{k,s}^- - \mathcal{E}_{k,s}^{\text{FEX}} - E_{k,s}^{\text{tr}} - (v - t_0)\mathcal{L}_s\} &\leq \mathcal{T}_{k,s} \\ &\leq U_{k,s}^{\text{jet}} := \mathcal{F}_{\partial,k}^{(s)} + e_{k,s}^+ + \mathcal{E}_{k,s}^{\text{FEX}} + E_{k,s}^{\text{tr}}. \end{aligned} \quad (\text{HJR25})$$

All errors on the right and left are the explicit accepted FEX, OAR and PEC expressions. The inequality uses $0 \leq F_{\Xi(L)}^{(s)} \leq (v - t_0)\mathcal{L}_s$ and $|\delta_0| \leq E^{\text{tr}}$ only after retaining their actual values in HJR24. In particular

$$\mathcal{T}_{k,s} = \Delta q \mathfrak{E}_{\partial} + o(kq), \quad U_{k,s}^{\text{jet}} = \Delta q \mathfrak{E}_{\partial} + o(kq). \quad (\text{HJR26})$$

This follows from the fully proved OFV/OVR outer first variation and the stated orders of the finite errors. The actual period remains fixed; no uniformity over a moving period is asserted.

Now take any paired restriction, quotient, or subquotient of the common coefficient space at the two metrics (D_0, D_N) . Let its rank be t and its determinant increment be $I_{t,N}$. These are attained restrictions and quotients along the same algebraic flag, retaining all its complete metrics. Since $D_N \geq D_0$, every such form increases. Full flag factorization, with nonnegative log increments on both complementary flags, proves

$$0 \leq I_{t,N} \leq I_N \leq U_{k,s}^{\text{jet}} = O_{\text{actual}}(kq). \quad (\text{HJR27})$$

For the same reason all generalized eigenvalues for (D_N, D_0) are at least one; the sum of all their logarithms is I_N . Hence every exterior degree of the identity between these two metrics has norm at most $\exp(U_{k,s}^{\text{jet}}/2)$. The same statement holds on every specified paired flag. This is a full exterior bound with no additional rank multiplier and with the complete lower-evaluation and kernel conditioning retained.

73.6 Exact cancellation of the low leading scale

Retain LJR's actual finite remainder $\mathcal{R}_{t,k,s}$, including its full-space formula LJR28 and the two-sided flag bounds LJR31. Keep the fixed G_{exp} quotient or restriction associated with this same flag. Then HJR19 gives the exact identity at every endpoint

$$\log \frac{\det G_t}{\det D_{t,N}} = 2tq \log k + \mathcal{R}_{t,k,s} - I_{t,N}. \quad (\text{HJR28})$$

HJR27 therefore propagates the same low leading scale $2tq \log k$ to all four endpoints, with remainder $O_{\text{actual}}(kq)$. It does so by the recurrence-compatible graph and the full Schur minima, not by assigning the later free jets the low dimension.

Let $p_{t,N}, n_{t,N}$ be PEC's two nonnegative logarithmic spectral parts for $(D_{t,N}, G_t)$. The same identity is

$$n_{t,N} = 2tq \log k + \mathcal{R}_{t,k,s} - I_{t,N} + p_{t,N}, \quad 0 \leq p_{t,N} \leq \varepsilon_{N,s}^{\vee} = o(kq). \quad (\text{HJR29})$$

At the actual kernel quotient and annihilator restriction, write $X = K, 0$ respectively. Substituting HJR29 with the four original signs cancels the four copies of both $2tq \log k$ and $\mathcal{R}_{t,k,s}$ exactly and gives

$$\begin{aligned} A_{X,k,s} &= I_{X,2q-1} + I_{X,2q} - I_{X,q} - P_{X,k,s}, \\ P_{X,k,s} &= p_{X,2q-1} + p_{X,2q} - p_{X,q-1} - p_{X,q}, \quad 0 \leq P_{X,k,s} \leq E_{k,s}^{\vee}. \end{aligned} \quad (\text{HJR30})$$

The signs of the p terms follow by substituting their displayed positions in HJR29. Their monotonicity and finite bound are PEC21,26 on these same maps. Together with HJR18 and HJR22 this supplies the explicit original calculation to carry forward:

$$\begin{aligned} A_{K,k,s} &= \sum_{M=q-1}^{2q-1} w_M \log \left(1 + \frac{y_M}{1 + x_M} \right) - P_{K,k,s}, \\ A_{0,k,s} &= \sum_{M=q-1}^{2q-1} w_M \log(1 + x_M) - P_{0,k,s}. \end{aligned} \quad (\text{HJR31})$$

The same original x_M, y_M sum in the full row energy $\log(1 + x_M + y_M)$. Every numerator, denominator and moment used to compute these scalars is fixed in HJR2–4, HJR14, HJR16–17. Their weighted total has the proved leading value HJR26; their separate allocation is retained by the exact formulas HJR31. Thus the next-order kq problem is the specified allocation of these original Gamma row energies, with all low factorial and Vandermonde terms canceled only through the exact equality HJR28.

74 The compatible high rows in the original receiving interval

Retain the full fixed-period simple-quartet hypotheses and all original source, coefficient and metric data in PBR and LRR. For each original source $s = 1, k$ the following uses exactly $B = B_K = Z_k^\top A_\ell$ and $J = J_0 = A_\ell^{-1} P_Y^\top$, with $\text{im } J = \ker B$ and ranks t_K, t_R of LRR1. In this section $D_M = \mathcal{D}_M^{(s)}$ is the full original dual metric; it is distinct from the original lower-grid polynomial χ' and from the residual-row matrix in PBR12.

The complete recurrence-compatible high-jet graph is $E_{N,s} = F_{N,s} R_s = \begin{pmatrix} I_n \\ H_{N,s} \end{pmatrix}$, where R_s is the original LJR4 reconstruction and $F_{N,s}$ is the full inverse-convolution and original Gamma coordinate map HJR3–4. HJR5 proves that its image is exactly the vectors whose convolved moments satisfy the monic χ recurrence. HJR6 proves $E_{N,s} V_{q-1,s} = V_{N,s}$, so the entire lower evaluation denominator is transported through the whole graph.

At a single original degree $q-1 \leq M < 2q$ use the full rows T, V through degree $M-v$ and the actual next rows t, v_+ . Their entries are the original functional values of $\mathcal{N}_a/E_A$ and the original lower-root evaluations on the degree- $M-v+1$ Gamma orthonormal polynomial. The exact full conditional update is

$$\begin{aligned} A_M &= V^* V, & K_M &= A_M^{-1} V^* T, & r_M &= t - v_+ K_M, & \lambda_M &= 1 + v_+ A_M^{-1} v_+^* > 0, \\ h_M &= \lambda_M^{-1/2} r_M, & D_{M+1} &= D_M + h_M^* h_M. \end{aligned} \quad (\text{HCR1})$$

Indeed subtracting the attained old extension minimum leaves the quadratic minimization $\eta^* A_M \eta + \|r_M x - v_+ \eta\|^2$. Its minimum is $\|r_M x\|^2 / (1 + v_+ A_M^{-1} v_+^*)$, by completing the square with $(A_M + v_+^* v_+)^{-1}$. The old lower coefficient is consequently retained throughout the complete row block; it is not minimized separately at each piece of the graph. This proves HCR1 with the full conditioning denominator and every original moment.

Put $D_{B,M} = (B D_M^{-1} B^*)^{-1}$, $D_{J,M} = J^* D_M J$, and $L_M = D_M^{-1} B^* D_{B,M}$. Then the exact inverse decomposition and two row energies are

$$\begin{aligned} D_M^{-1} &= J D_{J,M}^{-1} J^* + L_M D_{B,M}^{-1} L_M^*, \\ x_M &= h_M J D_{J,M}^{-1} J^* h_M^* \geq 0, & y_M &= h_M L_M D_{B,M}^{-1} L_M^* h_M^* \geq 0, \\ \log \frac{\det D_{J,M+1}}{\det D_{J,M}} &= \log(1 + x_M), \\ \log \frac{\det D_{B,M+1}}{\det D_{B,M}} &= \log \left(1 + \frac{y_M}{1 + x_M} \right). \end{aligned} \quad (\text{HCR2})$$

The columns $[J, L_M]$ form a full coefficient frame because $B L_M = I$ and $\text{im } J = \ker B$. Its D_M Gram is diagonal in these two blocks since $J^* D_M L_M = J^* B^* D_{B,M} = 0$. Inverting this exact frame Gram proves the first line. Restriction of HCR1 and the rank-one determinant lemma proves the third line. For the quotient, every lift of a vector z is $L_M z + J t$. Minimize its full updated norm over t . After using the original D_M orthogonal decomposition, the additional form is $(h_M L_M)^* (h_M L_M) / (1 + x_M)$. The determinant lemma proves the final line, including the denominator $1 + x_M$. Both x_M, y_M also retain λ_M^{-1} in HCR1.

Use the original weights $w_{q-1} = w_{2q-1} = 1$ and $w_M = 2$ for $q \leq M \leq 2q-2$, and define

$$\begin{aligned} \mathcal{V}_{K,k,s}^{\text{jet}} &= \sum_{M=q-1}^{2q-1} w_M \log \left(1 + \frac{y_M}{1 + x_M} \right), & \mathcal{V}_{0,k,s}^{\text{jet}} &= \sum_{M=q-1}^{2q-1} w_M \log(1 + x_M), \\ \mathcal{V}_{K,k,s}^{\text{jet}} &= A_{K,k,s} + P_{K,k,s}, & \mathcal{V}_{0,k,s}^{\text{jet}} &= A_{0,k,s} + P_{0,k,s}, \\ F_K^{(s)} &= \mathcal{V}_{K,k,s}^{\text{jet}} - \mathcal{E}_s^{\text{flag}} + F_L^{(s)} - F_{W_0}^{(s)}, \\ \Phi_{k,s} &= F_{W_0}^{(s)} + \mathcal{V}_{0,k,s}^{\text{jet}} + \delta_{0,k,s}. \end{aligned} \quad (\text{HCR3})$$

For each of the two metrics let $I_{X,N} = \log(\det D_{X,N} / \det D_{X,q-1})$. Summing the consecutive row identities in HCR2 gives $I_{X,N} = \sum_{M=q-1}^{N-1} g_{X,M}$. The expression $I_{X,2q-1} + I_{X,2q} - I_{X,q}$ therefore has precisely the weights displayed above, since the first row loses one copy and the last row occurs once. On the other hand the full dual identities $D_{B,N} = \widehat{H}_{K,N}^{-1}$ and $D_{J,N} = \overline{Q}_{0,N}^{-1}$ identify that weighted sum with the four original native returns. PBR5 then proves the second line. Equivalently HJR29–31 cancels the common low reconstruction scale and remainder in the four signs and gives $A_X = \mathcal{V}_X^{\text{jet}} - P_X$ exactly. Substitution into PBR9/7 proves the final two lines: the P_X terms cancel algebraically, rather than being estimated or set to zero. Both low terms remain separately, with their original metrics from PBR6.

The full finite errors are unchanged: $\mathcal{E}_s^{\text{flag}} \in [D_s^-, D_s^+]$ of PBR8 and $|\delta_{0,k,s}| \leq E_{k,s}^{\text{tr}}$ of PBR7, including α_A and all four CEP errors. Define the exact row-sum endpoints

$$\begin{aligned}
a_s^{\text{row},-} &= \mathcal{V}_{K,k,s}^{\text{jet}} - D_s^+ + \lambda_s^{\text{low}}, & a_s^{\text{row},+} &= \mathcal{V}_{K,k,s}^{\text{jet}} - D_s^- + \lambda_s^{\text{low}}, \\
b_s^{\text{row},-} &= F_{W_0}^{(s)} + \mathcal{V}_{0,k,s}^{\text{jet}} - E_{k,s}^{\text{tr}}, & b_s^{\text{row},+} &= F_{W_0}^{(s)} + \mathcal{V}_{0,k,s}^{\text{jet}} + E_{k,s}^{\text{tr}}, \\
a_s^{\text{row},-} &\leq F_K^{(s)} \leq a_s^{\text{row},+}, & b_s^{\text{row},-} &\leq \Phi_{k,s} \leq b_s^{\text{row},+}, \\
\Phi_{k,s}^{\text{row}} &= \max\{\Phi_{k,s}^{\text{V}}, b_s^{\text{row},-}\}, & \Phi_{k,s}^{+, \text{row}} &= \min\{\Phi_{k,s}^{+, \text{V}}, b_s^{\text{row},+}\}, \\
L_s^{\text{row}} &= \max\{L_s^{\text{V}}, a_s^{\text{row},-}, \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s}^{+, \text{row}} - e_{k,s}^{\text{V}}\}, \\
U_s^{\text{row}} &= \min\{U_s^{\text{V}}, a_s^{\text{row},+}, \mathcal{F}_{\partial,k}^{(s)} - \Phi_{k,s}^{+, \text{row}} + e_{k,s}^{\text{V}}\}.
\end{aligned} \tag{HCR4}$$

Here $\lambda_s^{\text{low}} = F_L^{(s)} - F_{W_0}^{(s)}$ is the unchanged difference of the two separately retained volumes. The first two interval assertions follow immediately from the exact HCR3 equalities and the stated finite error bounds. Subtracting the resulting residual interval in the exact OAR equality proves the last two lines. Every entry bounds the same original scalar; hence the intersection retains all prior branches and is nonempty. On the exact PMR range $k \geq 29$, $k\sqrt{\delta^2 + \gamma^2} \leq 2^{-33}(k+1)^2$, also include $\mathcal{F}_{0,k}^{(s)} - \Phi_{k,s}^{+, \text{row}} - e_{k,s}^{0,-}$ in the maximum and $\mathcal{F}_{0,k}^{(s)} - \Phi_{k,s}^{+, \text{row}} + e_{k,s}^{0,+}$ in the minimum, where $e^{0,\pm} = e^{\pm} + c_k^0$. Outside that range the original OAR entries remain. Because $0 \leq P_X \leq E_{k,s}^{\text{V}}$, the exact kernel row-sum interval lies inside PBR11's corresponding A_K interval, and its residual interval lies inside PBR11's corresponding A_0 interval. This containment follows by substituting $\mathcal{V}_X = A_X + P_X$ and comparing the two endpoints. It uses no distribution coefficient for x_M or y_M .

On the original arithmetic period domain $|u| \geq R_*$, retain precisely $a_{K,k}^{\pm}$ and all their finite constants from PBR14. The complete current arithmetic, signed and action receivers are

$$\begin{aligned}
L_{\text{ar}}^{\text{row}} &= \max\{L_{\text{ar}}^{\text{V}}, L_1^{\text{row}} + a_{K,k}^{\text{V}}\}, & U_{\text{ar}}^{\text{row}} &= \min\{U_{\text{ar}}^{\text{V}}, U_1^{\text{row}} + a_{K,k}^{\text{V}}\}, \\
F_K^{\text{ar}} &\in [L_{\text{ar}}^{\text{row}}, U_{\text{ar}}^{\text{row}}], & F_B^{\text{ar}} &\in [\mathcal{B}_k^{\text{ar}} - U_{\text{ar}}^{\text{row}}, \mathcal{B}_k^{\text{ar}} - L_{\text{ar}}^{\text{row}}], \\
L_k^{\text{row}} + \mathcal{H}_k + \delta_k^{\sigma} &\leq \Psi_k \leq U_k^{\text{row}} + \mathcal{H}_k + \delta_k^{\sigma}, \\
L_J^{\text{row}} &= L_k^{\text{row}} + \mathcal{H}_k + \delta_k^{\sigma} + C_k^{\text{budget}}, & U_J^{\text{row}} &= U_k^{\text{row}} + \mathcal{H}_k + \delta_k^{\sigma} + C_k^{\text{budget}}, \\
4q \log \mathcal{L}_{h,k} &\leq J_k^{\text{action}} \in [L_J^{\text{row}}, U_J^{\text{row}}], & \mathcal{L}_{h,k} &\leq \min_M \epsilon_M^{\text{ar}} \leq \prod_M (\epsilon_M^{\text{ar}})^{w_M/(2q)} \leq e^{U_J^{\text{row}}/(4q)}.
\end{aligned} \tag{HCR5}$$

The arithmetic source difference is the identical correlated scalar used in PBR14; add its exact interval and intersect. Subtracting from the unchanged total reverses the boundary endpoints. The signed equality adds $\mathcal{H}_k = -T_k$ and the original arithmetic correction. Adding the complete $C_k^{\text{budget}} = F_B^{(k)} + \mathcal{W}_k^{\text{ar}} - \mathcal{P}_- - \mathcal{P}_+$ then gives the action interval. The original untilted reflection and the same-class lower radius give the weighted geometric mean exactly as in PBR15. Every mixed pairing, both contraction penalties, the full monic window and action leakage remain. Under a real tilt use the unchanged full MCE phase formula.

Finally the exact same-row allocation is

$$\begin{aligned}
\mathcal{V}_{K,k,s}^{\text{jet}} + \mathcal{V}_{0,k,s}^{\text{jet}} &= \sum_{M=q-1}^{2q-1} w_M \log(1 + x_M + y_M), \\
F_K^{(s)} + \Phi_{k,s} &= \mathcal{V}_{K,k,s}^{\text{jet}} + \mathcal{V}_{0,k,s}^{\text{jet}} - \mathcal{E}_s^{\text{flag}} + F_L^{(s)} + \delta_{0,k,s}, \\
\mathcal{V}_{K,k,s}^{\text{jet}} + \mathcal{V}_{0,k,s}^{\text{jet}} &= \Delta q \mathfrak{C}_\partial + o(kq).
\end{aligned} \tag{HCR6}$$

The first equality is $(1 + x_M)(1 + y_M/(1 + x_M)) = 1 + x_M + y_M$ for every actual row. The second adds the two HCR3 identities, cancelling the identical original W_0 term only in that sum. The final equality is HJR24–26 with every finite FEX, transfer and outer error retained before the fixed-period limit. The individual allocations are exactly the two specified sums in HCR3; no individual leading coefficient is assigned by the evaluated total.