KOMOREBI MATHEMATICS / PAPER III

Typed paths, exact coordinates, and obstruction packets

The visual guide

Exact models for transport, holonomy, gauge, and nonlinear operations

Komorebi P.

Final companion edition / 5 October 2026

$$H - I = \Gamma_{0\leftarrow 2}\big(\Gamma_{2\leftarrow 1}\Gamma_{1\leftarrow 0} - \Gamma_{2\leftarrow 0}\big).$$

A reversible loop reveals its transported defect.

Twenty-four worked pages and twenty-two exact model figures accompany the traditional paper and its Lean guide. Integer, rational and symbolic calculations determine the figures. The preserved Python model sources and their explicit finite verification ranges are included in the source package.

A route through the examples

The labels V–1 through V–24 identify worked pages independently of PDF pagination. Each page names its corresponding statement in the final traditional edition. The Lean guide provides the separate formal contract.

V–1 Three objects, three different questions
V–2 An atlas of the objects
V–3 Read the product in execution order
V–4 All sixteen words, only five walks
V–5 Coefficients need not commute
V–6 A reachable packet can receive zero
V–7 Reachability is only an upper bound
V–8 Paths through changing packet lists
V–9 One vector, three exact frames
V–10 Nonlinear charts still cancel exactly
V–11 Reversibility is a real hypothesis
V–12 Measure the mismatch at its base
V–13 The identity can hide in the spectrum
V–14 Gauge changes the picture, not flatness
V–15 Flat is not the same as calibrated
V–16 Resolve the entire obstruction
V–17 Why diagonal checks are insufficient
V–18 A gauge can move every packet label
V–19 An obstruction rejects one candidate
V–20 A tree reads its named leaves
V–21 Where a nonlinear defect can enter
V–22 Mixed terms are not optional
V–23 From loop data to a packet witness
V–24 Read the evidence with the pictures

Three reading routes

Typed transfer: V–3 to V–8. **Coordinates, connection, and gauge:** V–9 to V–18, then V–23. **Candidate witnesses and trees:** V–19 to V–22, then V–24.

From the finite models to the broader paper

The traditional paper continues with rooted tree expansions, nonlinear refinement defects, obstruction objects, filtered lifting, elimination, and smooth projector transport. Read the finite examples as concrete preparation. The broader ordinary proofs keep their own hypotheses and are distinct from the finite Lean contract. Rooted trees: Section 18 (p. 37). Nonlinear defects: Section 20 (p. 42). Obstruction objects: Section 21 (p. 44). Filtered lifting: Section 23 (p. 51). Smooth transport: Section 27 (p. 61).

Read products in traversal order


A path $e_1, e_2, \ldots, e_n$ acts as $L_{e_n} \cdots L_{e_2} L_{e_1}$: the first traversed edge is the rightmost operator. Row indices are targets and column indices are sources throughout the guide.

V1 / PAPER III / A WORKED VISUAL COMPANION

Three objects, three different questions

Final traditional paper: Section 1 (p. 4); Section 4 (p. 10); Section 5 (p. 12)

An edge operator moves data between named packets. An exact coordinate change describes the same data in another complete chart. A connection supplies an additional choice of transport. The three can use similar arrows while making different mathematical assertions.



Three actual models used below: a single supported matrix entry, one ambient vector in three coordinate systems, and a shear obtained as chosen loop transport. The coordinates in the middle are not three different ambient vectors.

One paper, three tests

For an edge word, ask whether the target of each step matches the source of the next. For exact coordinates, ask whether analysis and synthesis are inverses. For a connection, ask whether its two-step transports agree with its declared direct transports. None of these tests can replace another.

This guide follows explicit integer and rational matrices through these tests. Each example has an inspectable input, calculation, and conclusion. Vector drawings display those calculations rather than replace them. A graph with two paths can have zero net transfer; a loop with trace two and determinant one can still differ from the identity.

The edition used here

The worked models originate in the finite algebraic v1.0.1 release. The final traditional paper also develops rooted tree expansions, nonlinear defects, obstruction objects, filtered lifting, and analytic transport under their stated hypotheses. Those ordinary mathematical developments have a wider scope than the finite Lean contract. The present models explain the finite mechanisms and provide a route into the broader paper.

> **Read this as:** Follow the object, not the arrow shape. A typed transfer can vanish, a complete coordinate loop must telescope, and an independently chosen connection can have genuine algebraic holonomy.

V2 / ORIENTATION / KEEP THE TYPES VISIBLE

An atlas of the objects

Final traditional paper: Definition 2.1 (p. 5); Definition 5.5 (p. 14); Definition 7.4 (p. 19)

Rows are targets and columns are sources throughout. If E_{ji} is a matrix unit, then $E_{ji}e_i = e_j$. The notation e_i means a basis vector; a named graph edge is written a, b, c, d or e, f when its role is explicit.

Object	What it records	First worked page
$\mathrm{src}, \mathrm{tgt} : E \to V$	The ordered endpoints of each distinct edge, including parallel edges.	V – 3–V – 4
$L_e = c(e)E_{\mathrm{tgt}(e),\mathrm{src}(e)}$	One packet-supported operator.	V – 3
$L_w = L_{e_n} \cdots L_{e_1}$	Execute the written word from left to right.	V – 4–V – 5
$T = \sum_e L_e$	Add all edge operators before taking powers.	V – 4–V – 7
F_a	A synthesis frame from chart a into the ambient space.	V – 9
$\mathcal{A}_a, \mathcal{S}_a$	Inverse analysis and synthesis maps; they may be maps of sets.	V – 10
$J_{b\leftarrow a}$	Canonical coordinate transport, forced by the exact charts.	V – 9–V – 10
$\Gamma_{b\leftarrow a}$	Additional connection data; reversibility must be stated.	V – 11–V – 12
$\Delta_{c,b,a}$	Two-step minus direct transport, a map from C_a to C_c.	V – 12
$H_{c,b,a}$	Loop transport based at a, an automorphism of C_a.	V – 13
G_a	An invertible gauge change at each chart.	V – 14–V – 15
$\Pi_\mu \,\mathrm{Ob}\, \Pi_\nu$	Output–input block of $\mathrm{Ob} = H - I$.	V – 16–V – 18
$r(x, p)$	A residual at packet p for the named candidate x.	V – 19
$t, \mathrm{inputs}(t)$	A finite operation tree and its syntactically used input labels.	V – 20

Reading routes

For transfer, read V–3–8. For geometry without a differential-geometric claim, read V–9–18. For candidate certificates and nonlinear scope, read V–19–22. V–23 puts the matrix mechanisms together; V–24 gives the formal boundary and replay guide.

> **Read this as:** A $C_a \to C_c$ defect and a $C_a \to C_a$ holonomy are not the same typed map. Returning the defect to the base fiber requires a specified inverse transport.

V3 / MODEL A / ONE EDGE IS ONE MATRIX ENTRY

Read the product in execution order

Final traditional paper: Lemma 2.2 (p. 5); Theorem 2.3 (p. 5); Theorem 2.4 (p. 6)

Work over $\mathbb{Z}$ with packets $0, 1, 2$. Let $e : 0 \to 1$ have coefficient 2, and let $f : 1 \to 2$ have coefficient 3. The elementary operators are

$$L_e = 2E_{10}, \qquad L_f = 3E_{21}.$$

The location of the nonzero entry records the type of the transfer, not an arbitrary matrix convention.

L_e


	0	1	2
0	0	0	0
1	2	0	0
2	0	0	0

L_f

	0	1	2
0	0	0	0
1	0	0	0
2	0	3	0

$L_f L_e$

	0	1	2
0	0	0	0
1	0	0	0
2	6	0	0



Each grid is an actual 3×3 matrix. The rightmost matrix factor acts first. In $L_f L_e$, packet 1 is the shared intermediate index; in $L_e L_f$ the interface is broken.

On a vector $(x_0, x_1, x_2)^{\mathsf{T}}$, the first step produces $(0, 2x_0, 0)^{\mathsf{T}}$, and the second produces $(0, 0, 6x_0)^{\mathsf{T}}$. Thus

$$L_f L_e = 6E_{20}, \qquad L_e L_f = 0.$$

For general edges, a summand in the entrywise product can survive only when the same intermediate packet is both $\mathrm{tgt}(e)$ and $\mathrm{src}(f)$:

$$(L_f L_e)_{yx} = \sum_z (L_f)_{yz} (L_e)_{zx}.$$

This is why a broken interface annihilates the entire product, even if both edge coefficients are nonzero.

What the zero says

It says that this ordered composition does not pass data through the declared packet types. It does not say that either edge is individually zero. For a longer word, finding one broken adjacent interface is sufficient to prove that the whole word vanishes.


> **Read this as:** Write the traversed edges first. Then reverse their order when writing the operator product: e followed by f is $L_f L_e$.

V4 / MODEL A / PARALLEL EDGES REMAIN DISTINCT

All sixteen words, only five walks

Final traditional paper: Theorem 3.8 (p. 8); Corollary 3.9 (p. 9)

Add four edges on three packets: $a : 0 \to 1$ with coefficient 2, $b : 0 \to 1$ with coefficient -1, $c : 1 \to 2$ with coefficient 3, and $d : 2 \to 0$ with coefficient 1. The parallel edges a and b are separate labels even though their source and target agree.



$$(T^2)_{20} = 6 - 3 = 3$$

first executed edge = column

second executed edge = row	a	b	c	d
a	×	×	×	2
b	×	×	×	-1
c	6	-3	×	×
d	×	×	3	×

×: broken packet interface

The board tests all ordered pairs. Columns are the first executed edge and rows are the second. A shaded cell gives the ordered coefficient of a valid walk; a cross means the product is zero by type mismatch.

The surviving words are ac, bc, cd, da, db. Their coefficients are $6, -3, 3, 2, -1$, respectively. Their endpoints determine where these numbers enter the matrix:

$$T = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 3 & 0 \end{pmatrix}, \qquad T^2 = \begin{pmatrix} 0 & 3 & 0 \\ 0 & 0 & 1 \\ 3 & 0 & 0 \end{pmatrix}.$$

At row 2, column 0, the two parallel choices give $6 - 3 = 3$. Deduplicating the edge labels before summing would alter the operator. On the other hand, adding the parallel coefficients in the already assembled matrix is legitimate: both act in the same supported matrix position.

The same calculation extends from pairs to all finite words by distributivity. The included check enumerates every word through length six and independently accumulates the endpoint coefficients. The empty word is treated separately as the identity.

> **Read this as:** The matrix power is a signed, ordered walk sum. It is not simply the number of paths, and a graph edge is not identified solely by its endpoints.

V5 / MODEL A / A MATRIX RING AS THE COEFFICIENT SEMIRING

Coefficients need not commute

Final traditional paper: Lemma 3.7 (p. 8); Section 9 (p. 22)

The edge theorem allows a noncommutative coefficient semiring. Use $R = M_2(\mathbb{Q})$ and let the two coefficients themselves be matrices:

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \qquad B = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

Attach A to $0 \to 1$ and B to $1 \to 2$. The single surviving endpoint coefficient is BA, not AB.

A		B		BA		AB	
1	1	1	0	1	1	2	1
0	1	1	1	1	2	1	1

$$e : 0 \to 1 \text{ carries } A; \quad f : 1 \to 2 \text{ carries } B$$

$$BA(1,0)^{\mathsf{T}} = (1,1)^{\mathsf{T}}, \qquad AB(1,0)^{\mathsf{T}} = (2,1)^{\mathsf{T}}$$

All entries are exact integers. The two products differ on the diagonal and act differently on the same input vector. This example is within the semiring-valued scalar model by taking the scalar ring to be $M_2(\mathbb{Q})$.

The distinction is visible before any abstraction:

$$BA \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \qquad AB \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$

For a written walk $(e_1, \ldots, e_n)$, its coefficient must therefore be

$$c^{\leftarrow}(w) = c(e_n)c(e_{n-1}) \cdots c(e_1).$$

A commutative example can hide an orientation mistake because reversing the coefficients changes nothing. This one cannot.

One ring is not a dependent family of fibers

The 3×3 matrix over $M_2(\mathbb{Q})$ can be displayed as an ordinary 6×6 matrix with 2×2 blocks. This is a concrete representation of the one-ring example. It does not establish an arbitrary family of differently typed edge fibers, which is outside the finite formal release's scalar edge-system interface.


> **Read this as:** The order rule carries mathematical information. It survives exact tests over a noncommutative ring and must not be inferred from pictures drawn over scalar real numbers alone.

V6 / MODEL A / CANCELLATION IS NOT A BROKEN INTERFACE

A reachable packet can receive zero

Final traditional paper: Corollary 3.9 (p. 9); Corollary 3.10 (p. 9)

Consider a diamond: two length-two routes from packet 0 to packet 3. Three edge coefficients are 1 and the lower final coefficient is −1. Both routes are correctly typed. Neither is annihilated by an interface mismatch.



The dashed edge carries the coefficient −1; it is an actual edge, not an uncertain or missing one. Both paths exist. The unsigned count is two, while their exact operator contributions sum to zero.

The endpoint identity gives

$$(T^2)_{30} = (1)(1) + (-1)(1) = 0.$$

Thus the paper's reachability implication has a definite direction:

$$\text{no compatible walk} \implies (T^n)_{ji} = 0.$$

Its converse is false. A zero entry may come from absent paths, zero coefficients, zero divisors in the coefficient ring, or cancellation among nonzero terms. The diamond exhibits the last possibility over $\mathbb{Z}$, where no zero divisor is needed.

Three different support records

The declared graph records permitted routes. The support of an individual word records the endpoints of that word when its coefficient is nonzero. The support of the sum records what remains after all contributions have been added. Passing from either of the first two records to the last requires the coefficients, not just adjacency.

A deliberately false test in the companion script asserts that reachability forces nonzero transfer. This diamond makes the test fail. That is a check against overreading the theorem, not a defect in the theorem.

> **Read this as:** Do not interpret a visible route as an occupied output. Graph support is a useful upper bound; signed operator support can be strictly smaller.

V7 / MODEL A / AN ORBIT THAT STAYS TWO-DIMENSIONAL

Reachability is only an upper bound

Final traditional paper: Corollary 3.11 (p. 9)

Keep the diamond and add an isolated packet 4. Start with the single seed $e_0 \in \mathbb{Q}^5$. Direct multiplication gives

$$Te_0 = e_1 + e_2, \qquad T^2 e_0 = 0.$$

Every later orbit vector is zero. The generated T-orbit hull is therefore exactly the span of the first two displayed columns.

$[e_0 \;\; Te_0 \;\; T^2 e_0]$ a basis of the orbit hull

0	1	0	0
1	0	1	0
2	0	1	0
3	0	0	0
4	0	0	0

1	0
0	1
0	1
0	0
0	0

$\text{Reach}(0) = \{0, 1, 2, 3\}$

$\text{Hull}_T(e_0) = \{(a, b, b, 0, 0)\}$

e_3 : reachable but absent

e_4 : unreachable and absent

rank $= 2$

The first matrix contains the actual orbit columns. The second supplies a basis of their span. The orbit hull lies inside the four-coordinate reachable subspace, but it has dimension two, not four.

A vector belongs to this hull precisely when it has the form

$$(a, b, b, 0, 0)^{\mathsf{T}}, \qquad a, b \in \mathbb{Q}.$$

Packet 4 is excluded because it is unreachable. Packet 3 is also absent, but for a different reason: the two contributions cancel. Even the reachable packets 1 and 2 remain linked; powers of the single total operator do not separate them.

The admissible operator grammar matters

Allowing each individual edge operator is a different generation problem. For instance, applying the edge $0 \to 1$ and then $1 \to 3$ individually produces e_3. It does not follow that e_3 belongs to the orbit of the summed operator T. The paper's orbit-support theorem concerns the explicitly stated operator action.

> **Read this as:** Unreachable means impossible under the declared action. Reachable means only that graph geometry has not ruled it out. Linear dependence and cancellation still have to be checked.

V8 / MODEL B / RECTANGULAR TWO- AND THREE-STEP PRODUCTS

Paths through changing packet lists

Final traditional paper: Proposition 8.1 (p. 21)

Here the packet lists have sizes $2, 3, 2, 1$ in succession. Let the actual maps be

$$\mathbb{Q}^2 \xrightarrow{A} \mathbb{Q}^3 \xrightarrow{B} \mathbb{Q}^2 \xrightarrow{C} \mathbb{Q}.$$

Unlike a power of one matrix, each step has its own entries and its own source and target list.

$A : \mathbb{Q}^2 \to \mathbb{Q}^3$

1	0
2	1
0	3

$B : \mathbb{Q}^3 \to \mathbb{Q}^2$

1	−1	2
0	1	1

$C : \mathbb{Q}^2 \to \mathbb{Q}$

2	−1

CBA

−4	6

	$(j,k)=(0,0)$	$(0,1)$	$(0,2)$	$(1,0)$	$(1,1)$	$(1,2)$	
$i=0$	2	−4	0	0	−2	0	$=-4$
$i=1$	0	−2	12	0	−1	−3	$=6$

The lower board expands one final row into all six intermediate-index choices for each input packet. The entries are $C_{0j}B_{jk}A_{ki}$, with $k \in \{0,1,2\}$ and $j \in \{0,1\}$. Zero terms remain visible.

The intermediate and total products are

$$BA = \begin{pmatrix} -1 & 5 \\ 2 & 4 \end{pmatrix}, \qquad CBA = \begin{pmatrix} -4 & 6 \end{pmatrix}.$$

For input packet 0, the six contributions sum to $2 - 4 + 0 + 0 - 2 + 0 = -4$. For input packet 1, they sum to $0 - 2 + 12 + 0 - 1 - 3 = 6$. These are literal matrix entries, not probabilities or path counts.

The formal scope is deliberately precise

The finite formal release's two- and three-step forms allow changing packet types. Its arbitrary-length heterogeneous product theorem allows changing *factors* over one common finite packet index. That formal statement does not assert a fully dependent arbitrary-length chain of changing index types. The finite rectangular illustration here uses exactly three factors.

For a single factor, the intermediate-index list is empty and contributes that matrix entry once. For zero factors, the result is the identity on the common space. These conventions must not be replaced by an empty sum equal to zero.

> **Read this as:** Heterogeneous can mean different matrices or different spaces. State which one is allowed before using a path formula.

V9 / MODEL C / CANONICAL TRANSPORT HAS NO HOLONOMY


One vector, three exact frames

Final traditional paper: Theorem 4.1 (p. 10); Theorem 4.2 (p. 10)

Let F_a synthesize chart coordinates into one ambient $\mathbb{Q}^2$. Choose

$$F_0 = I, \qquad F_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \qquad F_2 = \begin{pmatrix} 2 & 0 \\ 1 & 1 \end{pmatrix}.$$

The same ambient vector $x = (3,2)^{\mathsf{T}}$ has chart coordinates $u_a = F_a^{-1}x$.



Each panel is a coordinate plane, not a different physical location. The numerical arrows differ because the frames differ. Multiplying each coordinate vector by its corresponding F_a recovers the same x.

Transport from chart a to chart b is forced:

$$J_{b\leftarrow a} = F_b^{-1}F_a.$$

The route $0 \to 1 \to 2$ agrees with the direct route because the middle frame cancels:

$$(F_2^{-1}F_1)(F_1^{-1}F_0) = F_2^{-1}F_0.$$

Returning to chart 0 cancels every remaining frame. The final coordinate vector is $(3,2)^{\mathsf{T}}$, not a sheared or rotated variant.

No metric is required

The frames need not be orthogonal. The second one is a shear, and the third has determinant two. Exact invertibility, rather than preservation of angles or lengths, gives the cocycle and loop identities. The same telescoping argument works for every finite chart path.

> **Read this as:** A canonical coordinate loop is a sequence of changes in description of one object. Nonidentity holonomy requires additional transport data, not merely nonorthogonal charts.

V10 / MODEL C / NATURALITY DOES NOT REQUIRE DERIVATIVES


Nonlinear charts still cancel exactly

Final traditional paper: Proposition 10.1 (p. 23)

Exact coordinate equivalences can be nonlinear. On $\mathbb{Q}^2$, use

$$\mathcal{A}(x, y) = (x, y + x^2), \qquad \mathcal{S}(u, v) = (u, v - u^2).$$

Both composites are the identity by direct substitution. The real drawing below is a geometric view of these same polynomial formulas.



Horizontal segments become parabolas, but no information is discarded. The drawing samples the displayed polynomial boundaries for rendering; inverse identities and route equality are checked symbolically, not inferred from the curves.

Now represent the nonlinear operation $F(x, y) = (x + y, xy)$ in the new chart. Set $w = u + v - u^2$. Its correct coordinate formula is

$$\widehat{F}_b(u, v) = \mathcal{A}F\mathcal{S}(u, v) = (w,\ u(v - u^2) + w^2).$$

For $(x, y) = (1, 2)$, applying F first gives $(3, 2)$ and then $(3, 11)$ in the new chart. Changing the input first gives $(u, v) = (1, 3)$, and $\widehat{F}_b(1, 3) = (3, 11)$ as well.

The equality holds as a polynomial identity for all inputs. It is the set-level naturality theorem: only the inverse laws for the coordinate maps are needed. No Taylor expansion, derivative, or higher chain rule is involved.

> **Read this as:** Nonlinearity alone does not create a coordinate defect. Reusing the old formula after changing coordinates would be an extra modeling choice, not exact conjugation.

V11 / MODEL D / THE MINIMAL AND REVERSIBLE INTERFACES

Reversibility is a real hypothesis

Final traditional paper: Definition 5.1 (p. 12); Definition 5.5 (p. 14)

Choose the forward scalar transports $\Gamma_{1\leftarrow 0} = 2$, $\Gamma_{2\leftarrow 1} = 3$, and $\Gamma_{2\leftarrow 0} = 5$. Their two-step/direct multiplicative comparison is $D^{\times} = 6/5$. A loop also needs a declared return map $\Gamma_{0\leftarrow 2}$.

connection data	$\Gamma_{0\leftarrow 2}$	H	$D^{\times}$	$\Gamma_{0\leftarrow 2}\Gamma_{2\leftarrow 0}$
self-normalized only	$\frac{1}{6}$	1	$6/5$	$\frac{5}{6}$
reversible	$\frac{1}{5}$	$\frac{6}{5}$	$6/5$	1

$$\Gamma_{1\leftarrow 0} = 2, \quad \Gamma_{2\leftarrow 1} = 3, \quad \Gamma_{2\leftarrow 0} = 5$$

The first row has H = identity at this triangle, but fails its cocycle.

All transports are invertible rational scalars and self-transports are one. The first row is permitted by the minimal interface. The second uses the inverse of the direct forward edge and therefore satisfies that part of reversibility.

In the first row, choose the return scalar $1/6$. Then

$$H = (1/6) \cdot 3 \cdot 2 = 1, \qquad \Gamma_{2\leftarrow 1}\Gamma_{1\leftarrow 0} - \Gamma_{2\leftarrow 0} = 1.$$

This triangle loop is the identity even though its cocycle equation fails. The return is not the inverse of the direct transport: $(1/6)5 \neq 1$. Thus one cannot transfer the reversible defect–holonomy equivalence to the minimal interface.

In the reversible row, the return is $1/5$. Now the loop is $6/5$, and the additive defect of one is converted into the loop obstruction $1/5$ by that inverse return map.

Two safeguards

Global flatness in the minimal interface implies trivial loops by using its cocycle on the whole route, including the return. Separately, in the reversible interface, triviality of each triangle loop is equivalent to that triangle's cocycle. Neither assertion says that one isolated loop certifies all triangles.


> **Read this as:** Self-normalized means $\Gamma_{a\leftarrow a} = I$. Reversible additionally means $\Gamma_{a\leftarrow b} = \Gamma_{b\leftarrow a}^{-1}$. These are different assumptions.

V12 / MODEL D / AN EXACT REVERSIBLE TRIANGLE

Measure the mismatch at its base

Final traditional paper: Theorem 5.10 (p. 15); Section 9 (p. 22)

Retain the reversible scalar model with forward values $2, 3, 5$ and all reverse values given by inverses. Two routes from chart 0 to chart 2 multiply the input by 6 and 5, respectively. The additive discrepancy is one times the original input.



loop input 1 **returns** $6/5$

loops	returned value
0	1
1	$\frac{6}{5}$
2	$\frac{36}{25}$
3	$\frac{216}{125}$
4	$\frac{1296}{625}$

The dashed direct edge is a specified comparison map, not a missing edge. The table shows the exact effect of traversing the same loop repeatedly. These are rational values of this example, not an analytic convergence claim.

The defect lives at the target of the forward routes:

$$\Delta_{2,1,0} = 3 \cdot 2 - 5 = 1.$$

The holonomy lives back at chart 0:

$$H_{2,1,0} = 5^{-1} \cdot 3 \cdot 2 = \frac{6}{5}, \qquad H - I = \frac{1}{5} = \Gamma_{0\leftarrow 2}\Delta.$$

For input 5, the direct and two-step outputs are 25 and 30. Returning by $1/5$ gives 5 and 6. The target discrepancy of 5 has become a base discrepancy of 1. The types and locations of these differences matter even in a one-dimensional example.

Because the return map is invertible, it cannot turn a nonzero defect into zero. This is the mechanism behind $\Delta = 0 \iff H = I$ in the reversible interface.

> **Read this as:** Holonomy compares paths after both have been returned to the same base fiber. It is the transported mismatch, not an untyped difference between outputs in different charts.

V13 / MODEL E / A SHEAR WITH NONTRIVIAL HOLONOMY

The identity can hide in the spectrum


Final traditional paper: Definition 5.9 (p. 15); Definition 7.4 (p. 19)

Put $\Gamma_{1\leftarrow 0} = H$, $\Gamma_{2\leftarrow 1} = I$, and $\Gamma_{2\leftarrow 0} = I$, with reverse maps given by inverses, where

$$H = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

The loop $0 \to 1 \to 2 \to 0$ then has exactly this shear as its holonomy.

The unit square under H



H

1	1
0	1

$H - I$

0	1
0	0

$$\operatorname{tr} H = 2, \quad \det H = 1, \quad (H - I)e_1 = e_0$$

The dashed square is the original set. The solid parallelogram is its exact image. The second coordinate basis vector moves to $(1, 1)^{\mathsf{T}}$; its nonzero displacement is the first basis vector.

The usual scalar invariants can be misleading:

$$\operatorname{tr} H = 2, \qquad \det H = 1, \qquad \chi_H(t) = (t-1)^2.$$

These are also the invariants of the 2×2 identity matrix. Nevertheless,

$$\mathrm{Ob} = H - I = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \neq 0, \qquad \mathrm{Ob}\, e_1 = e_0.$$

A complete obstruction test must therefore retain the operator or a certificate strong enough to detect its nonzero action. Trace, determinant, and eigenvalue lists alone do not distinguish this loop from the identity.

What the picture certifies, and what it does not

The plotted corners are the images of the four unit-square corners under the displayed matrix. The equation $\mathrm{Ob}\, e_1 = e_0$ is the exact witness. No smooth connection, curvature tensor, or continuum limit is used in this model; its holonomy is the finite algebraic loop product. The traditional paper treats smooth transport separately under additional hypotheses.

> **Read this as:** Flatness is the equality $H = I$, not agreement of a short list of spectral summaries. A single explicit moved vector can disprove that equality.

V14 / MODEL E / CONJUGATION AND ITS LIMITS


Gauge changes the picture, not flatness

Final traditional paper: Theorem 6.2 (p. 17)

At the base chart, change coordinates by $D = \mathrm{diag}(2, 1)$. Transform every connection edge by the prescribed endpoint gauges. The loop transforms by conjugation, giving

$$H^D = DHD^{-1} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}.$$

The shear looks stronger in these fixed numerical axes, but it is not a different gauge-invariant flatness verdict.



Both figures are exact images of a unit square under their displayed matrices. The Frobenius norms are computed in the indicated fixed coordinate bases. A nonorthogonal gauge need not preserve such a norm.

The rank of the obstruction is one on both sides. Trace, determinant, and characteristic polynomial are preserved under this conjugation. Most importantly,

$$D(H - I)D^{-1} = 0 \quad \Longleftrightarrow \quad H - I = 0.$$

An invertible coordinate change cannot turn a nonidentity loop into the identity. The fact that a particular residual norm changes does not contradict the invariance of exact vanishing.

The base chart determines the outer gauge

In a triangle product, adjacent intermediate gauges cancel:

$$(G_a \Gamma_{a \leftarrow c} G_c^{-1})(G_c \Gamma_{c \leftarrow b} G_b^{-1})(G_b \Gamma_{b \leftarrow a} G_a^{-1}) = G_a H G_a^{-1}.$$

Only the base gauge survives around the loop. For the additive defect, the surviving outer gauges instead belong to its different source and target fibers.

> **Read this as:** Exact nonvanishing is gauge invariant. The numerical size of a matrix entry or a chosen norm generally is not.

V15 / MODEL D / A FLAT CONNECTION WITH NONIDENTITY EDGES


Flat is not the same as calibrated

Final traditional paper: Theorem 6.3 (p. 17)

Make all three canonical charts literally identical, so $J_{b\leftarrow a} = 1$ over $\mathbb{Q}$. Choose scalar gauges $g_0 = 1$, $g_1 = 2$, and $g_2 = 3$, and define

$$\Gamma_{b\leftarrow a} = g_b g_a^{-1}.$$

The edge $0 \to 1$ is multiplication by two, although its canonical counterpart is the identity.



$$\Gamma_{1\leftarrow 0} = 2, \quad \Gamma_{2\leftarrow 1} = 3/2, \quad \Gamma_{2\leftarrow 0} = 3$$

The rulers display the chosen gauge scales: the marks are $g_a n$ for integer n. They visualize the supplied scalar factors, not an intrinsic metric. Relative transport is the ratio of the target and source scales.

The two-step route $0 \to 1 \to 2$ has multiplier $(3/2)2 = 3$, equal to the direct route $0 \to 2$. More generally,

$$(g_c g_b^{-1})(g_b g_a^{-1}) = g_c g_a^{-1}$$

for every triple. Thus this connection is flat and reversible, but it is not calibrated to the original identity-valued J.

Recovering the gauge from one root

Choose a base chart o. The theorem constructs

$$G_a = \Gamma_{a\leftarrow o} J_{o\leftarrow a}, \qquad \Gamma_{b\leftarrow a} = G_b J_{b\leftarrow a} G_a^{-1}.$$

With $o = 0$ and $J = 1$, the recovered G_a are exactly $1, 2, 3$. The statement concerns the complete pairwise interface and its cocycle. It is not a topological claim about an arbitrary graph with missing direct comparisons.

> **Read this as:** Flatness makes the chosen transports path-independent. Calibration identifies them with one particular canonical transport before any gauge change.

V16 / MODEL F / INPUT–OUTPUT PACKET BLOCKS

Resolve the entire obstruction

Final traditional paper: Theorem 7.5 (p. 19); Corollary 7.6 (p. 20)

Use three coordinate packets with projectors P_0, P_1, P_2, and set

$$H = I + 2E_{01} + 3E_{12}, \qquad \mathrm{Ob} = H - I.$$

Here H is invertible, so it can be the loop product of a reversible three-chart connection. Its off-diagonal entries encode two packet-mixing effects.

H

1	2	0
0	1	3
0	0	1

$\mathrm{Ob} = H - I$

0	2	0
0	0	3
0	0	0

	$\nu = 0$	$\nu = 1$	$\nu = 2$
$\mu = 0$	0	$2E_{01}$	0
$\mu = 1$	0	0	$3E_{12}$
$\mu = 2$	0	0	0

$$\mathrm{Ob}(4,5,6)^{\mathsf{T}} = (10,18,0)^{\mathsf{T}}$$

sum all nine blocks

The board on the right contains all nine actual embedded blocks $P_\mu \,\mathrm{Ob}\, P_\nu$. Row labels are output packets and column labels are input packets. Only the addresses $(0,1)$ and $(1,2)$ carry nonzero maps.

Completeness on both sides supplies the reconstruction:

$$\mathrm{Ob} = (P_0 + P_1 + P_2)\,\mathrm{Ob}\,(P_0 + P_1 + P_2) = \sum_{\mu,\nu} P_\mu \,\mathrm{Ob}\, P_\nu.$$

For $x = (4,5,6)^{\mathsf{T}}$, the input in packet 1 contributes $10e_0$ and the input in packet 2 contributes $18e_1$. Their sum is the full displacement $\mathrm{Ob}\, x$.

A block becomes a witness when it acts

The nonzero block $2E_{01}$ has the explicit source vector e_1, supported entirely in packet 1, and $P_0 \,\mathrm{Ob}\, e_1 = 2e_0 \neq 0$. Similarly, $P_1 \,\mathrm{Ob}\, e_2 = 3e_1 \neq 0$. The general theorem supplies such a witness classically when a nonzero block is known; the example displays it concretely.

> **Read this as:** An obstruction packet is not just a colored matrix cell. It consists of a specific packet-pair label and an exact nonvanishing statement about the corresponding map.

V17 / MODEL E / COMPLETENESS, TWO SIDES, AND OBLIQUE PACKETS

Why diagonal checks are insufficient

Final traditional paper: Theorem 7.5 (p. 19)

Return to $\mathrm{Ob} = E_{01}$. With the two standard coordinate projectors, both diagonal blocks vanish:

$$P_0 \,\mathrm{Ob}\, P_0 = P_1 \,\mathrm{Ob}\, P_1 = 0.$$

The obstruction nevertheless survives in the off-diagonal block $P_0 \,\mathrm{Ob}\, P_1 = E_{01}$.

Ob

0	1
0	0

$\sum_i P_i \,\mathrm{Ob}\, P_i$

0	0
0	0

nonzero diagonal test misses it

Q_0

1	0
1	0

Q_1

0	0
−1	1

$$Q_0 + Q_1 = I, \quad Q_i^2 = Q_i$$

$$\sum_{i,j} Q_i \,\mathrm{Ob}\, Q_j = \mathrm{Ob}; \qquad Q_0^{\mathsf{T}} \neq Q_0$$

The diagonal-only test on the left misses the whole obstruction. On the right, the displayed oblique projectors still form a complete exact resolution; they need not be orthogonal projections for the Euclidean inner product.

There is a second failure mode. Supplying only P_0 gives a family of idempotents but not a complete one. The single tested block is zero, while the omitted packet carries the missing input. The implication "all tested blocks vanish, therefore the operator vanishes" requires the completeness hypothesis.

Exact reconstruction does not require right angles

For the displayed Q_0, Q_1,

$$Q_0 + Q_1 = I, \qquad Q_i^2 = Q_i, \qquad Q_0 Q_1 = Q_1 Q_0 = 0,$$

but $Q_0^{\mathsf{T}} \neq Q_0$. Expanding both copies of I still proves

$$\sum_{i,j} Q_i \,\mathrm{Ob}\, Q_j = \mathrm{Ob}\,.$$

The block reconstruction identity itself uses the sum-to-identity condition. Idempotence is additionally used when constructing a witness that remains supported in one packet after projection. The release does not require pairwise annihilation for its general reconstruction statement.

> **Read this as:** Test every input–output block of a complete resolution. Testing only the diagonal, or only the packets supplied so far, answers a weaker question.

V18 / MODEL E / FIXED VERSUS TRANSPORTED PROJECTORS

A gauge can move every packet label

Final traditional paper: Proposition 7.7 (p. 21)

The diagonal gauge on V–14 commuted with each standard projector. Now use the packet-mixing gauge

$$G = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \qquad \mathrm{Ob} = E_{01}.$$

The total obstruction transforms by conjugation, but its pattern in the *fixed* standard packets changes.

Ob		G		$G\,\mathrm{Ob}\,G^{-1}$	
0	1	1	0	−1	1
0	0	1	1	−1	1
one nonzero entry		$GP_0 \neq P_0G$		four nonzero entries	

fixed axes: different packet pattern / moving axes: conjugate labeled blocks

The first obstruction has one nonzero entry. Its conjugate has four. Both have rank one and are nonzero. The change in fixed block pattern is not a change in whether the connection is flat.

Direct multiplication gives

$$G\,\mathrm{Ob}\,G^{-1} = \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix}.$$

For example, the original $(0,0)$ block vanishes, while the new $(0,0)$ block relative to the old coordinate projectors does not. This is why the theorem about preservation of each block's zero status requires $GP_i = P_iG$.

Transporting the packet frame as well

Define $Q_i = GP_iG^{-1}$. Then the identities cancel exactly:

$$Q_i(G\,\mathrm{Ob}\,G^{-1})Q_j = G(P_i\,\mathrm{Ob}\,P_j)G^{-1}.$$

Relative to these transported packets, the corresponding labeled blocks again have matching vanishing status. These are the same oblique projectors displayed on V–17. One must distinguish moving the operator alone in a fixed resolution from moving the entire resolved description.

> **Read this as:** The total obstruction is intrinsic under gauge. Its location in a fixed packet grid is not, unless the gauge preserves that grid.

V19 / MODEL G / THE QUANTIFIERS ARE THE CONTENT

An obstruction rejects one candidate

Final traditional paper: Theorem 7.2 (p. 18)

Let the candidates be rational numbers and the packet labels be $0, 1$. Define

$$r(x, 0) = x^2 - 1, \qquad r(x, 1) = x - 1.$$

A candidate is a solution when both residuals vanish. The five rows of data below are a display of this explicit residual function, not a search over an unspecified class.

candidate x	-2	-1	0	1	2
$r(x, 0) = x^2 - 1$	3	0	-1	0	3
$r(x, 1) = x - 1$	-3	-2	-1	0	1
	rejected	rejected	rejected	solution	rejected

$$x = -1 : \quad r(-1, 0) = 0, \quad r(-1, 1) = -2$$

A shaded cell is a nonzero exact residual. At $x = -1$, the first equation passes but the second supplies an obstruction. At $x = 1$, both pass. The rejected candidate does not obstruct the successful one.

The statement proved by the generic certificate has the form

$$\neg(\forall p,\ r(x, p) = 0) \quad \Longleftrightarrow \quad \exists p,\ r(x, p) \neq 0.$$

The same candidate x appears on both sides. A certificate consists of the packet label and its nonvanishing proof. The general converse uses classical reasoning and does not itself provide an algorithm to find the label.

What would be required to rule out all solutions?

One would need the stronger assertion $\forall x\, \exists p,\ r(x, p) \neq 0$. The value $x = 1$ disproves that assertion in this example. A witness for $x = -1$ supplies no evidence for it. Likewise, finding an obstruction block for one triangle of one chosen connection does not classify every possible connection.

The term *obstruction* in this generic interface does not automatically mean a cokernel class, a cohomology group, a norm bound, or a minimal certificate. Those require additional constructions and theorems.

> **Read this as:** Keep the candidate attached to the witness. The theorem's strength is its exact logical scope, not a universal existence or nonexistence claim.

V20 / MODEL H / ACTUAL SYNTAX, ARBITRARY SEMANTICS


A tree reads its named leaves

Final traditional paper: Definition 11.1 (p. 24); Theorem 11.2 (p. 24)

Use the grammar's constant, input, unary, and binary constructors to build

$$t = (x^2 + 3) - y.$$

Here "square," "add," and "subtract" are explicitly supplied meanings of operation symbols. At $x = 2, y = 5$, every node value can be evaluated exactly.



$$\text{inputs}(t) = \{x, y\}$$

unused = -100 or 100

same value: 2

$x - x$: $\{x\}$ syntactically

but always zero

The rooted tree records input-slot order. Arrows indicate evaluation from children to parent, not chart transport. An environment may contain additional variables, but this tree does not read them.

The input set is $\text{inputs}(t) = \{x, y\}$. If two environments agree at x and y, structural induction proves that the two evaluations agree: input leaves agree, constants agree, and every parent applies the same supplied operation to equal child values.

Syntactic locality is not minimal dependence

The tree $x - x$ has input set $\{x\}$, although its value is always zero under ordinary subtraction. Locality provides a sufficient agreement condition; it does not compute the smallest set of semantically relevant variables. Nor does it prove that two different trees are equivalent.

The semantics may act on arbitrary values. For example, a binary concatenation symbol applied to strings "ab" and "cd" gives "abcd." No linearity or differentiability is required. Conversely, swapping the ordered inputs of the subtraction root above changes the value from 2 to -2.

> **Read this as:** The proved tree theorem controls which input labels are read. It does not supply a nonlinear chain rule, a stability estimate, or a solution of a tree equation.

V21 / MODEL H / EXACT COORDINATES VERSUS A CHOSEN RULE


Where a nonlinear defect can enter

Final traditional paper: Proposition 10.1 (p. 23); Roadmap 11.7 (p. 26); Theorem 20.2 (p. 42)

Take the unary operation $f(x) = x^2$ over $\mathbb{Q}$. Suppose both input and output transport are chosen as multiplication by two, while the same numerical formula f is reused at the target. The two routes give

$$2f(x) = 2x^2, \qquad f(2x) = 4x^2.$$

Their difference is $-2x^2$, which is generally nonzero.



x	0	1	2
$2f(x)$	0	2	8
$f(2x)$	0	4	16
$2f(x) - f(2x)$	0	-2	-8

correct chart formula: $f_b(v) = v^2/2$

Solid: $2x^2$. Dashed: $4x^2$. The curves are a real visualization of exact polynomial formulas; the table contains exact rational-domain evaluations. The mismatch is caused by the specified transport/formula combination.

For a canonical coordinate change with input and output chart $v = 2x$, the target formula is not still v^2. It is

$$\widehat{f}_b(v) = 2f(v/2) = \frac{v^2}{2}.$$

Consequently $\widehat{f}_b(2x) = 2x^2$, and the exact coordinate defect vanishes. This is the same distinction illustrated in the two-dimensional example on V–10.

An example is not a general propagation theorem

The polynomial calculation here is exact and checked symbolically. It explains a chosen node defect, while the traditional paper develops composition identities, iterated defects, fixed point stability, and derivative defects under explicit hypotheses. Those broader ordinary proofs must be read on their own terms. This one example and the finite Lean locality theorem do not certify them.

> **Read this as:** A nonzero nonlinear route difference must come from identifiable extra data: a changed transport, an approximation, a restricted model, or a reused formula that was not conjugated correctly.

V22 / FINITE EXAMPLES / WHY MULTILINEARITY MATTERS

Mixed terms are not optional


Final traditional paper: Roadmap 11.6 (p. 25); Theorem 18.3 (p. 38)

A bilinear node distributes over two independently resolved inputs:

$$(a+b)(c+d) = ac + ad + bc + bd.$$

The four terms correspond to all ordered choices of one input packet from each slot. Keeping only matching packet labels loses the mixed terms.

$$(2+3)(5+7) = 60$$



Diagonal pieces are not the whole product

$$ac + bd = 10 + 21 = 31$$

$$ad + bc = 14 + 15 = 29$$

$$(2+3)^2 = 25 \neq 2^2 + 3^2 = 13$$

missing square term: $2 \cdot 2 \cdot 3 = 12$

The rectangle has width $2+3$ and height $5+7$, with differently scaled horizontal and vertical display units. Each cell is labeled by its actual product. The lighter diagonal pair contributes only 31 of the total 60.

For a unary square, inserting $a+b$ does not give a^2+b^2. The extra $2ab$ arises because the operation is nonlinear in its single argument. It can be analyzed by representing the square using a binary multiplication node with a repeated input, but that is an additional choice of syntax and semantics.

The broader tree theorem needs its hypotheses

General packet tree expansion needs correctly typed input slots, finite packet-label sets at each node, and multilinearity in every slot used for distributive expansion. Empty inputs and unary nodes need explicit conventions. Slot order must survive even when values or operations do not commute. The traditional paper supplies rooted tree reconstruction and pruning results in its ordinary mathematical development.

This page supplies elementary exact examples of the issue. The finite v1.0.1 Lean contract remains narrower: it contains syntactic input locality, not the general dependent tree expansion or the traditional paper's analytic convergence results.

> **Read this as:** A branching picture does not justify throwing away mixed interactions. Exact packet expansion requires the appropriate multilinearity and typing, not merely a finite-looking diagram.

V23 / CAPSTONE / AN END-TO-END FINITE DIAGNOSIS

From loop data to a packet witness

Final traditional paper: Theorem 5.10 (p. 15); Theorem 6.2 (p. 17); Corollary 7.6 (p. 20)

Supply a reversible connection on three copies of $\mathbb{Q}^2$ by choosing

$$\Gamma_{1\leftarrow 0} = A, \qquad \Gamma_{2\leftarrow 1} = B, \qquad \Gamma_{2\leftarrow 0} = C = I,$$

and declaring every reverse edge to be the inverse. This time both nontrivial forward matrices are retained.

A

1	1
0	1

B

1	0
1	1

$H = BA$

1	1
1	2

$H - I$

0	1
1	1

$$e_0 = (1,0) \;\mapsto\; (1,0) \;\mapsto\; (1,1) \;\mapsto\; (1,1)$$

$$P_1(H - I)e_0 = e_1 \neq 0$$

---

replace direct edge $C = I$ by $C = BA$: then $C^{-1}BA = I$

The vector row follows e_0 through A, then B, then C^{-1}. Returning to the base produces $(1,1)^{\mathsf{T}}$, whose difference from the input is the source-supported obstruction witness e_1.

The calculation gives

$$\Delta = BA - C = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \qquad H = C^{-1}BA = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}.$$

Choose the standard packet projectors. The $(1,0)$ block is nonzero, and the explicit candidate vector satisfies $P_1(H - I)e_0 = e_1$. This proves nonflatness of the supplied connection at the supplied triangle.

Repairing data is not the same as changing gauge

Replacing the direct map $C = I$ by $C = BA$, and updating its reverse, makes the new loop equal to I. That changes the connection. It cannot be a gauge of the original nonflat connection, because gauge preserves whether holonomy is the identity. For this complete three-chart example all cocycles then hold; changing one edge in a larger network would require checking the other affected loops as well.

> **Read this as:** Declare the data, multiply in traversal order, form the defect, resolve all blocks, and exhibit an input on which one block acts nontrivially. Each step has a checkable mathematical object.

V24 / EVIDENCE / EXAMPLES, PROOFS, AND FORMAL SCOPE

Read the evidence with the pictures

Final traditional paper: Section 12 (p. 26)

This guide preserves the twenty-four worked pages and twenty-two exact figures from the finite visual appendix. It updates their references to the final traditional paper and explains the relation to that paper's broader nonlinear and analytic material.

Read each evidence record in its scope

Evidence	What it supports
Finite Lean release	The frozen declarations under their original hypotheses and pinned foundations. The Lean guide records the precise theorem map, source identity checks, and inherited build evidence. No fresh kernel compilation is claimed here.
Ordinary mathematics	The traditional paper's expanded proofs, including nonlinear, obstruction, lifting, and smooth transport results. Inclusion in that paper does not add these results to the finite Lean contract.
Exact model checks	The inherited 12 September 2026 report records 124 passed checks and four rejected false controls. Current model and checker hashes match that record. Fresh replay was blocked by missing SymPy. The scope is the displayed finite matrices, paths, polynomials, candidate tables, and trees.
Figure regeneration	The twenty-two shipped TikZ sources match their preserved original bytes. Their regeneration was not rerun because SymPy was unavailable. Byte identity is source provenance, not a fresh execution result.

Separately, eleven mixed exact integer and numerical verification groups passed in a fresh run of the preserved strengthened source and checker, not the newly edited final TeX. Three canonical schematic PNGs also regenerated byte-for-byte. Neither count is a replay of this guide's twenty-two exact plates.

Boundaries that remain in force

The formal edge system uses a single scalar coefficient semiring. The arbitrary-length heterogeneous path theorem uses a common packet index, with changing-type forms available for two and three factors. General dependent tree expansion and analytic convergence lie outside this finite formal contract. The traditional paper states and proves its broader results with their own hypotheses.

The generic obstruction theorem rejects its named candidate. A nonzero holonomy block rejects flatness for its named loop data. Neither conclusion supplies a universal classification, a new arithmetic realization, or an algorithmic search guarantee.

Replay the worked layer

From the editable visual-guide directory, run `python build.py` to compile this PDF. Run `python checks/verify_examples.py` for the exact examples and `python source/generate_graphics.py` for the figures. Use a temporary copy to retain the delivered evidence unchanged. The full package preserves the finite release and its original formal sources separately.

> **Read this as:** Use the figures to understand the mechanisms, the traditional paper to read the mathematical proofs, and the Lean guide to identify the exact formal contract. These are related records with distinct scopes.