CHAPTER IV

Projector Resolved Seed Calculus IV

Cyclic analytic functions, reciprocal families, and spectral zero data

One cyclic family connects exact coefficient selection, differential transport, reciprocal polynomials, and the geometry of zeros.

Komorebi P.

Mathematical text

How to use this volume

Begin with cyclic projection and the signed differential cycle. The reciprocal sections then build parameter polynomials, Appell systems, and theta polynomials from a unit series. The second half follows the same denominator through large zero geometry, canonical products, and zero zeta functions.

The visual atlas at the end of this chapter supplies twenty-six worked plates. The companion Lean guide follows the formal declarations and their hypotheses. The main text contains sixty-eight theorem-like results and ninety-five numbered records.

Overview

Fix a positive integer N. Root of unity projection of the exponential produces residue-class sections that, for $\lambda \neq 0$, form a basis for $y^{(N)} = \lambda^N y$. The signed sections for $y^{(N)} = -y$ yield the denominator D_N and its reduced form Δ_N. We derive reciprocal coefficient polynomials, Bell and Newton recurrences, an additive parameter law, and residue section Appell and theta families. For $N \geq 2$, Δ_N has order $1/N$. We specialize classical two term zero theory for exponential sums to the regular exponent polygon. In this equal-weight family, conjugation symmetry combined with unique near-model localization places all sufficiently large zeros exactly on the symmetry rays, and direct expansion gives their leading displacement and derivative sign. The classical perimeter law yields zero-counting asymptotics with explicit leading constants and bounded error. These results give genus-zero products, the standard diagonal Fredholm representation, zero-power-sum formulas for reciprocal and logarithmic coefficients, nearest-zero coefficient asymptotics, and meromorphic continuation of the symmetry-orbit zero zeta by comparison with the Hurwitz zeta function. Complex powers are interpreted either formally or through a fixed logarithm on a zero-free domain. This chapter treats only this cyclic analytic family. Arithmetic interpolation, general nonlinear trees, general resolvent theory, and abstract completion are reserved for Chapters IX, III, VI, and X.

Formal scope and evidence

The recorded formal contract covers all sixty-eight theorem-like manuscript results under their stated hypotheses. The formalization ledger identifies the corresponding declarations.

Imported standard fact 11.1 is used in the ordinary manuscript growth argument. The Lean growth-order proofs establish that step separately. Open questions, branch choices, multiplicities, and the hypotheses of coefficient asymptotics remain explicit. Finite computations support the worked models. The companion Lean guide records the replay using pinned sources, declaration-kind checks, axiom audit, and their limits.

Contents

1 Introduction 1

2 A map of the construction 2

I Cyclic resolution and differential transport 4

3 Cyclic projection of analytic functions 4

4 The transfer exponential basis 6

5 Signed cyclic sections and the distinguished denominator 7

6 A finite algebra and matrix model 10

II Reciprocal coefficients and polynomial families 13

7 Reciprocal powers and parameter polynomials 13

8 Logarithmic derivatives and Riccati equations 17

9 Residue-section reciprocals and Appell systems 19

10 Scaled reciprocal coupling and theta polynomials 23

III The geometry of zeros 25

11 Growth and the geometry of the large zeros 25

IV Products, moments, and zero zeta functions 34

12 Hadamard products and zero power sums 34

13 Coefficient asymptotics from the nearest zeros 38

14 Zero zeta functions 40

V Scope and formal evidence 45

15 Scope boundaries and further questions 45

16 Formalization ledger 45

17 Conclusion 47

A Visual atlas 49

List of Figures

1 An exact address model for $N = 3$. Every displayed exponent belongs to one residue lane. The drawing records labels rather than coefficient magnitudes. 4

2 The exact signed shift J_4. Source columns and target rows describe the same action as the cycle on the left. Direct integer multiplication verifies $J_4^4 = -I$. 10

3 At $N = 2$ and $r = 1$, the reciprocal is $\sec z$. The upper row gives block exponential coefficients and the lower row gives ordinary Taylor coefficients. Both rows encode the same five terms. 13

4 The two exact parameter polynomials from the worked calculation. Each plotted curve evaluates its displayed formula. The points at $r = 1$ recover the first two positive-index secant coefficients. 17

5 The exact order-four model. The first three zeros on each of four rays are mapped by $w = z^4$ to three positive reduced zeros. The reduced axis uses a logarithmic scale. This complete order-four example does not classify the finite exceptional set at other orders. . 25

6 Order-two moment sums over the first forty reduced zeros $\lambda_k = \pi^2(k+1/2)^2$. The curves are finite numerical partial sums. The dashed limits $1/2$ and $1/6$ follow from the exact logarithmic coefficients and the product identity. 34

1 Introduction

For a power series

$$F(z) = \sum_{m \geq 0} a_m z^m$$

and a positive integer N, the coefficients may be separated according to the residue of m modulo N. Root of unity averaging gives exact projection operators for this separation. This elementary construction is usually called a multisection or a root filter. It appears throughout the classical theory of generating functions and special functions. Background on multisections, hypergeometric functions, and related coefficient systems may be found in [Rai60, AAR99, Com74].

Chapters I and II developed typed seed grammars, exact projector packets, and finite residue-projector algebra, while Chapter III treated typed edge operators and exact coordinate transport [P.26a, P.26b, P.26c]. The root filter used below is the finite cyclic scalar specialization of that earlier projector calculus. The signed companion matrix is likewise a concrete cyclic instance, not a replacement for Chapter III's general typed-edge theory.

This chapter follows the consequences of applying the cyclic projections to the exponential function. The projected exponentials form a finite differential system. After one sign is inserted around the derivative cycle, they form a basis for

$$y^{(N)} = -y.$$

The section with initial value one is

$$\mathsf{D}_N(z) = \sum_{n \geq 0} (-1)^n \frac{z^{Nn}}{(Nn)!}.$$

Its reciprocal powers produce polynomial coefficient families. Its logarithmic derivative produces a scalar higher Riccati identity. The same reciprocal construction applied to every exponential section gives normalized residue section functions and ordinary Appell polynomial sequences. These cyclic exponential sections and their matrix addition laws belong to the generalized hyperbolic function literature [Sch92, Mul05]. Appell and binomial-type sequences are discussed in [Rom84, Com74].

The second half of this chapter concerns the zeros of D_N. The finite exponential sum

$$\mathsf{D}_N(z) = \frac{1}{N} \sum_{\eta^N = -1} e^{\eta z}$$

has a regular polygon of exponents. The general relation between an exponent polygon and the zero strips of an exponential sum goes back to Pólya and Langer. Later developments include the work of Moreno and the treatment in Levin's monograph [Pól20, Lan31, Mor73, Lev80]. Bleher and Mallison give a particularly close general result: for convex-polygon exponential sums they locate all sufficiently large zeros in two term strips and prove exponential localization and simplicity [BM06]. The regular polygon and conjugation symmetry of the present family specialize that theory to N exact rays. We give a self-contained specialization because its notation is used throughout the product and zeta theory, and we compute the leading displacement term for this family. The resulting counting formulas have explicit leading constants and bounded error.

The rotational symmetry is more useful after the change of variable $w = z^N$. There is an entire function

$$\Delta_N(w) = \sum_{n\geq 0}(-1)^n \frac{w^n}{(Nn)!}$$

with

$$\mathsf{D}_N(z) = \Delta_N(z^N).$$

For $N \geq 2$, the order of Δ_N is $1/N$. Its Hadamard product therefore has genus zero. The logarithm of this product identifies the transfer tangent coefficients with power sums of the reduced zeros. The large zero formula also compares the reduced zero zeta with a Hurwitz zeta function. Standard material on entire functions, canonical products, and exponential sums appears in [Boa54, Lev80, Tit39]. We use singularity analysis in the form described in [FS09] and the classical continuation of the Hurwitz zeta function from [Apo76].

The cyclic exponential sections and their matrix addition laws have a classical literature. The tools used below include Appell and binomial-type sequences, canonical products, singularity analysis, and Fredholm determinants. These tools are also standard. Our aim is to organize this symmetric specialization and link its reciprocal coefficients to reduced-zero data. No historical priority is asserted for the component constructions.

2 A map of the construction

> **The central idea**
>
> The same object is read in different coordinates. Selecting residues makes its Taylor coefficients visible. A signed shift makes the differential system finite. Passing to $w = z^N$ removes rotational repetition from its zeros. The formulas below identify these descriptions under their stated hypotheses.

Two variables appear throughout this chapter. The variable z carries the differential equation and the rotational symmetry. The reduced variable

$$w = z^N$$

records one representative from each rotational orbit. Confusing these variables hides the main simplification. The denominator D_N has order one as a function of z, while Δ_N has order $1/N$ as a function of w.

The analytic objects are introduced in the following order. First,

$$\mathsf{E}_{N,j}(z) = \sum_{n\geq 0} \frac{z^{Nn+j}}{(Nn+j)!}$$

is the residue j part of the exponential. Second,

$$\mathsf{D}_{N,j}(z) = \sum_{n\geq 0}(-1)^n \frac{z^{Nn+j}}{(Nn+j)!}$$

is the signed basis for $y^{(N)} = -y$. The section $\mathsf{D}_N = \mathsf{D}_{N,0}$ is the denominator used in the reciprocal family

$$\mathsf{S}_N^{(r)} = \mathsf{D}_N^{-r}.$$

Set $\mathsf{T}_N = -\mathsf{D}_N'/\mathsf{D}_N$. Equivalently, T_N is the logarithmic derivative of $\mathsf{S}_N^{(1)}$.

A different reciprocal family comes from normalizing each exponential section at the origin. The function

$$\mathsf{C}_{N,j}(z) = \frac{z^j/j!}{\mathsf{E}_{N,j}(z)}$$

is a unit series in z^N. Its powers, after multiplication by e^{xz}, give Appell polynomials. The theta family couples a scaled argument with a reciprocal order.

The zero theory begins only after these algebraic relations have been established. The factorization

$$\mathsf{D}_N(z) = \Delta_N(z^N)$$

turns every orbit of N zeros in the z plane into one zero in the w plane. The large zeros in the z plane are located by two dominant exponentials. The zeros in the w plane then give a genus zero product. The logarithm of that product returns to the coefficient theory through

$$\text{zero power sums} = \text{logarithmic coefficients}.$$

This equality is the bridge between the first and second halves of this chapter.

All powers with a complex parameter are first defined as formal series at the origin. To represent this germ analytically, use a simply connected zero-free region containing the origin and the logarithm normalized to vanish there. On other regions, specify a branch separately. Integer reciprocal powers are globally meromorphic and do not require a branch choice.

The main statements fall into five groups. The finite algebra model in section 6 explains the derivative cycles and addition laws through one exponential identity. The coefficient theory in section 7 gives two complementary recurrences, one from the original denominator and one from its logarithm. The Appell theory in section 9 supplies both formal differential-operator formulas and a convolution law in the space and order parameters. The zero analysis in section 11 specializes classical two term exponential-sum theory, uses symmetry to place the uniquely localized large zeros on the rays, and computes a leading correction. The product theory in section 12 gives the standard diagonal Fredholm encoding of the genus zero product.

This chapter keeps formal and analytic statements separate. A formal identity needs only the unit constant term of the relevant series. A holomorphic complex power also needs a zero-free domain and a chosen logarithm. A meromorphic coefficient asymptotic is stated only for positive integral reciprocal order unless an additional branch domain is specified.

Part I Cyclic resolution and differential transport

3 Cyclic projection of analytic functions

A route through this section

The root filter asks which coefficient addresses belong to a packet. It is an exact selector. Multiplication then asks how two selected packets interact. Their residue labels add modulo N, so mixed inputs remain part of the product.




Figure 1. An exact address model for $N = 3$. Every displayed exponent belongs to one residue lane. The drawing records labels rather than coefficient magnitudes.

Fix $N \geq 1$ and write

$$\omega = e^{2\pi i/N}.$$

Let Ω be a disk centered at the origin, with the whole plane allowed, and assume that Ω is invariant under multiplication by ω. For a holomorphic function F on Ω, define

$$(\mathcal{B}_{N,j}F)(z) = \frac{1}{N}\sum_{\ell=0}^{N-1} \omega^{-j\ell}F(\omega^{\ell}z), \qquad 0 \leq j < N.$$

The same formula is valid for formal power series.

Theorem 3.1 (Root filter). *If*

$$F(z) = \sum_{m\geq 0} a_m z^m,$$

then

$$\mathcal{B}_{N,j}F(z) = \sum_{m \equiv j \bmod N} a_m z^m.$$

The operators satisfy

$$\mathcal{B}_{N,i}\mathcal{B}_{N,j} = \delta_{ij}\mathcal{B}_{N,j}$$

and

$$\sum_{j=0}^{N-1} \mathcal{B}_{N,j} = I.$$

Proof. Substitution of the power series gives

$$\mathcal{B}_{N,j}F(z) = \sum_{m \geq 0} a_m z^m \left(\frac{1}{N} \sum_{\ell=0}^{N-1} \omega^{(m-j)\ell} \right).$$

The inner average is one when $m \equiv j \bmod N$ and zero otherwise. The operator identities follow from the disjoint residue classes. □

Proposition 3.2 (Covariance characterization)**.** *A holomorphic function F belongs to the range of $\mathcal{B}_{N,j}$ if and only if*

$$F(\omega z) = \omega^j F(z).$$

Proof. Every monomial selected by $\mathcal{B}_{N,j}$ has the stated covariance. Conversely, write $F = \sum_k \mathcal{B}_{N,k}F$. Replacing z by ωz gives

$$\sum_k \omega^k \mathcal{B}_{N,k}F = \omega^j \sum_k \mathcal{B}_{N,k}F.$$

Application of $\mathcal{B}_{N,k}$ gives

$$(\omega^k - \omega^j)\mathcal{B}_{N,k}F = 0.$$

The powers $1, \omega, \dots, \omega^{N-1}$ are distinct, so $\mathcal{B}_{N,k}F = 0$ for $k \neq j$. □

Multiplication is not diagonal in these coordinates. The residue labels add modulo N. This is the scalar cyclic instance of the packet product from Chapter I.

Theorem 3.3 (Cyclic product law)**.** *If F and G are holomorphic on the same rotation-invariant domain, then*

$$\mathcal{B}_{N,k}(FG) = \sum_{\substack{0 \leq i,j < N \\ i+j \equiv k \bmod N}} (\mathcal{B}_{N,i}F)(\mathcal{B}_{N,j}G).$$

Proof. Write $F = \sum_i F_i$ and $G = \sum_j G_j$, where $F_i = \mathcal{B}_{N,i}F$ and $G_j = \mathcal{B}_{N,j}G$. Their covariance gives

$$(F_i G_j)(\omega z) = \omega^{i+j} F_i(z) G_j(z),$$

so $F_i G_j$ belongs to the packet labeled by $i + j$ modulo N. Expanding the finite product and applying $\mathcal{B}_{N,k}$ gives the formula. □

Remark 3.4. Every identity in this section holds formally. In the holomorphic setting, the finite projector sums preserve local uniform convergence. No choice of branch is involved.

4 The transfer exponential basis

Definition 4.1. For $\lambda \in \mathbb{C}$ and $0 \le j < N$, define

$$\mathsf{E}_{N,j}(\lambda z) = \mathcal{B}_{N,j}(e^{\lambda z}) = \sum_{n \ge 0} \frac{(\lambda z)^{Nn+j}}{(Nn+j)!}.$$

Theorem 4.2 (Differential cycle and initial data)**.** *For* $1 \le j < N$,

$$\frac{d}{dz}\mathsf{E}_{N,j}(\lambda z) = \lambda \mathsf{E}_{N,j-1}(\lambda z),$$

while

$$\frac{d}{dz}\mathsf{E}_{N,0}(\lambda z) = \lambda \mathsf{E}_{N,N-1}(\lambda z).$$

Thus

$$\frac{d^N}{dz^N}\mathsf{E}_{N,j}(\lambda z) = \lambda^N \mathsf{E}_{N,j}(\lambda z).$$

For $0 \le k < N$,

$$\left.\frac{d^k}{dz^k}\mathsf{E}_{N,j}(\lambda z)\right|_{z=0} = \delta_{kj}\lambda^j.$$

If $\lambda \neq 0$, *the* N *functions* $\mathsf{E}_{N,j}(\lambda z)$ *form a basis of the solution space of* $y^{(N)} = \lambda^N y$.

Proof. Termwise differentiation gives the cyclic derivative formulas. One full cycle contributes λ^N. The first nonzero term of $\mathsf{E}_{N,j}(\lambda z)$ is $(\lambda z)^j/j!$, which gives the initial data. The initial value vectors are linearly independent when $\lambda \neq 0$. The solution space of an order N constant coefficient equation has dimension N, so these functions form a basis. □

Corollary 4.3 (Wronskian of the transfer basis)**.** *Assume* $N \ge 2$. *For* $\lambda \neq 0$, *the Wronskian of the ordered basis*

$$\mathsf{E}_{N,0}(\lambda z), \dots, \mathsf{E}_{N,N-1}(\lambda z)$$

is the constant

$$\lambda^{N(N-1)/2}.$$

Proof. At the origin, the derivative matrix is diagonal with entries

$$1, \lambda, \dots, \lambda^{N-1}.$$

For an explicit determinant proof, write each basis function as a finite sum of exponentials with exponents $a_\ell = \lambda\omega^\ell$. Its derivative matrix factors as

$$W(z) = V\ \mathrm{diag}(e^{a_\ell z})\, C, \qquad V_{k\ell} = a_\ell^k,$$

where C is the constant matrix of Fourier weights. Thus

$$\det W(z) = e^{z\sum_\ell a_\ell} \det W(0) = \det W(0),$$

because $N \geq 2$ implies $\sum_\ell \omega^\ell = 0$. The product of the diagonal initial values is the stated power of λ. □

Corollary 4.4 (Addition law)**.** *For $0 \leq k < N$,*

$$\mathsf{E}_{N,k}(\lambda(z+w)) = \sum_{\substack{0 \leq i,j < N \\ i+j \equiv k \bmod N}} \mathsf{E}_{N,i}(\lambda z)\mathsf{E}_{N,j}(\lambda w).$$

Proof. Expand $e^{\lambda(z+w)} = e^{\lambda z}e^{\lambda w}$ as a bivariate power series. A term $z^a w^b$ contributes to the kth section exactly when $a + b \equiv k \bmod N$. Grouping first by $i \equiv a \bmod N$ and $j \equiv b \bmod N$ gives the displayed sum. □

The basis also has a scalar hypergeometric form.

Theorem 4.5 (Hypergeometric form)**.** *Let*

$$A_{N,j} = \left\{ \frac{j+m}{N} : 1 \leq m \leq N \text{ and } j+m \neq N \right\}.$$

Then

$$\mathsf{E}_{N,j}(\lambda z) = \frac{(\lambda z)^j}{j!} {}_0F_{N-1}\left(\begin{matrix} - \\ A_{N,j} \end{matrix}, \frac{(\lambda z)^N}{N^N} \right).$$

Proof. The multiplication formula for the gamma function gives

$$(Nn+j)! = j! N^{Nn} n! \prod_{a \in A_{N,j}} (a)_n.$$

Indeed, the N Pochhammer factors supplied by the multiplication formula are $((j+m)/N)_n$ for $1 \leq m \leq N$. The excluded factor is $((j+N-j)/N)_n = (1)_n = n!$. Insertion into the defining series gives the stated hypergeometric expansion. □

Example 4.6. For $N = 1$, the only section is $e^{\lambda z}$. For $N = 2$,

$$\mathsf{E}_{2,0}(z) = \cosh z, \qquad \mathsf{E}_{2,1}(z) = \sinh z.$$

These are classical instances of the same cyclic basis.

5 Signed cyclic sections and the distinguished denominator

Definition 5.1. For $0 \leq j < N$, define

$$\mathsf{D}_{N,j}(z) = \sum_{n \geq 0} (-1)^n \frac{z^{Nn+j}}{(Nn+j)!}.$$

The distinguished signed denominator is

$$\mathsf{D}_N(z) = \mathsf{D}_{N,0}(z).$$

Fix any $\rho \in \mathbb{C}$ with $\rho^N = -1$.

Proposition 5.2 (Relation with the transfer exponential)**.** *For every j,*

$$\mathsf{D}_{N,j}(z) = \rho^{-j}\mathsf{E}_{N,j}(\rho z).$$

Proof. The coefficient of z^{Nn+j} on the right is

$$\rho^{-j}\rho^{Nn+j}\frac{1}{(Nn+j)!} = (-1)^n\frac{1}{(Nn+j)!}.$$

□

Theorem 5.3 (Signed derivative cycle)**.** *For $1 \le j < N$,*

$$\mathsf{D}'_{N,j} = \mathsf{D}_{N,j-1},$$

and

$$\mathsf{D}'_{N,0} = -\mathsf{D}_{N,N-1}.$$

Every signed section satisfies

$$\mathsf{D}^{(N)}_{N,j} = -\mathsf{D}_{N,j}.$$

The N signed sections form the basis of $y^{(N)} = -y$ determined by

$$\mathsf{D}^{(k)}_{N,j}(0) = \delta_{jk}$$

for $0 \le j, k < N$.

Proof. The derivative formulas follow directly from the series. The first nonzero term of $\mathsf{D}_{N,j}$ is $z^j/j!$, so the initial value matrix is the identity. The basis statement follows from the dimension of the solution space. □

Corollary 5.4 (Wronskian of the signed basis)**.** *Assume $N \ge 2$. The Wronskian of*

$$\mathsf{D}_{N,0}(z), \dots, \mathsf{D}_{N,N-1}(z)$$

is one.

Proof. The derivative matrix at the origin is the identity. The determinant factorization in the proof of corollary 4.3 applies with $a_\ell = \rho\omega^\ell$ and with the signed-section Fourier weights. Since $\sum_\ell a_\ell = 0$, the determinant is constant and equals one. □

Proposition 5.5 (Twisted addition law)**.** *For $0 \le k < N$,*

$$\mathsf{D}_{N,k}(z+w) = \sum_{\substack{0 \le i,j < N \\ i+j \equiv k \bmod N}} (-1)^{\lfloor (i+j)/N \rfloor}\mathsf{D}_{N,i}(z)\mathsf{D}_{N,j}(w).$$

Proof. Use proposition 5.2 and corollary 4.4. The factor introduced by a pair (i, j) is

$$\rho^{i+j-k}.$$

Since $0 \le i+j \le 2N-2$ and $i+j \equiv k \bmod N$, this factor is one when $i+j=k$ and -1 when $i+j=k+N$. □

Theorem 5.6 (Finite exponential representation)**.** *Let*

$$\mathcal{R}_N = \{\eta \in \mathbb{C} : \eta^N = -1\}.$$

Then

$$\mathsf{D}_N(z) = \frac{1}{N} \sum_{\eta \in \mathcal{R}_N} e^{\eta z}.$$

More generally,

$$\mathsf{D}_{N,j}(z) = \frac{\rho^{-j}}{N} \sum_{\ell=0}^{N-1} \omega^{-j\ell} e^{\rho\omega^\ell z}.$$

Proof. The second formula is the root filter expression for $\rho^{-j}\mathsf{E}_{N,j}(\rho z)$. The first is the case $j=0$. □

Definition 5.7. Define the reduced denominator

$$\Delta_N(w) = \sum_{n \ge 0} (-1)^n \frac{w^n}{(Nn)!}.$$

Then

$$\mathsf{D}_N(z) = \Delta_N(z^N).$$

Corollary 5.8 (Symmetries)**.** *The denominator satisfies*

$$\mathsf{D}_N(\omega z) = \mathsf{D}_N(z)$$

and

$$\mathsf{D}_N(\overline{z}) = \overline{\mathsf{D}_N(z)}.$$

Its nonzero zeros occur in orbits under multiplication by the Nth roots of unity and under complex conjugation.

Proof. The first identity follows from the powers z^{Nn}. The second follows from the real coefficients. Both symmetries preserve the zero set and multiplicity. □

Example 5.9. The first denominators are

$$\mathsf{D}_1(z) = e^{-z}, \qquad \mathsf{D}_2(z) = \cos z.$$

For $N=3$,

$$\mathsf{D}_3(z) = \frac{1}{3}\left(e^{-z} + 2e^{z/2}\cos\left(\frac{\sqrt{3}z}{2}\right)\right).$$

For $N=4$,

$$\mathsf{D}_4(z) = \cos\left(\frac{z}{\sqrt{2}}\right)\cosh\left(\frac{z}{\sqrt{2}}\right).$$

6 A finite algebra and matrix model

The central idea

The signed shift is a finite carrier for the differential cycle. At order four, three basis steps have coefficient one and the final step has coefficient minus one. Four steps return the vector with reversed sign. The matrix exponential packages every finite path length in one analytic expression.




Figure 2. The exact signed shift J_4. Source columns and target rows describe the same action as the cycle on the left. Direct integer multiplication verifies $J_4^4 = -I$.

The derivative and addition laws can be read from one finite algebra identity. This formulation also identifies the exponent polygon as the spectrum of a signed cyclic shift.

Definition 6.1. Let

$$\mathfrak{A}_N^+ = \mathbb{C}[\xi]/(\xi^N - 1)$$

and

$$\mathfrak{A}_N^- = \mathbb{C}[\xi]/(\xi^N + 1).$$

Every element has a unique representative of degree below N.

Theorem 6.2 (Finite algebra exponential identities)**.** *In* $\mathfrak{A}_N^+$ *one has*

$$e^{\lambda z \xi} = \sum_{j=0}^{N-1} \mathsf{E}_{N,j}(\lambda z)\xi^j.$$

In $\mathfrak{A}_N^-$ one has

$$e^{z\xi} = \sum_{j=0}^{N-1} \mathsf{D}_{N,j}(z)\xi^j.$$

Differentiation and multiplication of these identities recover the cyclic derivative systems and the two addition laws.

Proof. Expand the exponential and write every exponent uniquely as $Nn+j$. In $\mathfrak{A}_N^+$, the relation $\xi^{Nn+j} = \xi^j$ gives the transfer sections. In $\mathfrak{A}_N^-$, the relation $\xi^{Nn+j} = (-1)^n\xi^j$ gives the signed sections.

Differentiation multiplies by $\lambda\xi$ in the first algebra and by ξ in the second. Reduction of the product by the defining relation gives the derivative cycles. The identity

$$e^{(z+w)\xi} = e^{z\xi}e^{w\xi}$$

gives the addition laws after comparison of the coefficients of $1, \xi, \dots, \xi^{N-1}$. □

Definition 6.3. Let J_N be the signed cyclic shift on the standard basis $e_0, \dots, e_{N-1}$ of $\mathbb{C}^N$, defined by

$$J_N e_j = e_{j+1} \quad (0 \le j < N-1), \qquad J_N e_{N-1} = -e_0.$$

Then $J_N^N = -I$.

Theorem 6.4 (Companion matrix realization)**.** *Put*

$$\mathbf{D}_N(z) = \begin{pmatrix} \mathsf{D}_{N,0}(z) \\ \mathsf{D}_{N,1}(z) \\ \vdots \\ \mathsf{D}_{N,N-1}(z) \end{pmatrix}.$$

Then

$$\mathbf{D}_N(z) = e^{zJ_N}e_0$$

and

$$\mathsf{D}_N(z) = \frac{1}{N}\,\mathsf{tr}(e^{zJ_N}).$$

The eigenvalues of J_N are the N roots of -1, each with multiplicity one.

Proof. The vector $e^{zJ_N}e_0$ has derivative $J_N e^{zJ_N}e_0$ and initial value e_0. Its components therefore satisfy the same signed derivative cycle and initial data as $\mathbf{D}_N$. Uniqueness for the linear system gives the first identity.

The vectors

$$e_0, J_N e_0, \dots, J_N^{N-1}e_0$$

are the standard basis. Thus e_0 is cyclic for J_N, so the minimal polynomial of J_N has degree N. Since $J_N^N = -I$, that minimal polynomial divides $x^N + 1$ and therefore equals $x^N + 1$. The characteristic polynomial has the same degree and is divisible by the minimal polynomial, so it too equals $x^N + 1$.

This polynomial has distinct roots. Consequently J_N is diagonalizable and its spectrum consists of all N roots of -1, each once. Hence

$$\operatorname{tr}(e^{zJ_N}) = \sum_{\eta^N=-1} e^{\eta z} = N\mathsf{D}_N(z)$$

by theorem 5.6. □

Corollary 6.5 (Entries of the fundamental matrix)**.** *For $0 \le p, q < N$, let d_{pq} be the member of $\{0, \dots, N-1\}$ congruent to $p - q$ modulo N. Put*

$$\varepsilon_{pq} = \begin{cases} 1, & p \ge q, \\ -1, & p < q. \end{cases}$$

Then

$$\left(e^{zJ_N}\right)_{pq} = \varepsilon_{pq}\mathsf{D}_{N,d_{pq}}(z).$$

Thus every entry of the fundamental matrix is one of the signed sections, with the sign recording whether the cyclic path from q to p crosses the signed edge.

Proof. For every nonnegative integer n,

$$J_N^{Nn+d_{pq}} e_q = (-1)^n \varepsilon_{pq} e_p.$$

Only powers whose exponent is congruent to d_{pq} modulo N contribute to the (p, q) entry of the exponential. Therefore

$$\left(e^{zJ_N}\right)_{pq} = \varepsilon_{pq} \sum_{n\ge 0} (-1)^n \frac{z^{Nn+d_{pq}}}{(Nn+d_{pq})!}.$$

This is the stated formula. □

Corollary 6.6 (Addition and inversion laws for the companion matrix)**.** *Assume $N \ge 2$. The matrix*

$$U_N(z) = e^{zJ_N}$$

satisfies

$$U_N(z+w) = U_N(z)U_N(w), \qquad U_N(z)^{-1} = U_N(-z), \qquad \det U_N(z) = 1.$$

Its first column is $\mathbf{D}_N(z)$, and all of its entries are given by corollary 6.5.

Proof. The first two identities are the exponential group law. Since $\operatorname{tr} J_N = 0$,

$$\det(e^{zJ_N}) = e^{z \operatorname{tr} J_N} = 1.$$

The first column statement is theorem 6.4. □

Part II Reciprocal coefficients and polynomial families

7 Reciprocal powers and parameter polynomials

A route through this section

A unit constant term makes the formal reciprocal unambiguous. To compute one coefficient, only finitely many earlier coefficients are needed. The later analytic theory explains how those exact coefficients relate to the full zero set.

The factorial normalization carries the same coefficients in two forms

	n = 0	n = 1	n = 2	n = 3	n = 4
$P_{2,n}(1)$	1	1	5	61	1385
$[z^{2n}]\sec z$	1	1/2	5/24	61/720	1385/40320

Figure 3. At $N = 2$ and $r = 1$, the reciprocal is $\sec z$. The upper row gives block exponential coefficients and the lower row gives ordinary Taylor coefficients. Both rows encode the same five terms.

Since $\mathsf{D}_N(0) = 1$, its formal logarithm is defined at the origin. On a simply connected zero-free region containing the origin, there is a unique holomorphic branch normalized by $\log \mathsf{D}_N(0) = 0$, and its germ at the origin agrees with the formal logarithm.

Definition 7.1. For $r \in \mathbb{C}$, define

$$\mathsf{S}_N^{(r)}(z) = \mathsf{D}_N(z)^{-r} = \exp(-r \log \mathsf{D}_N(z)).$$

Write

$$\mathsf{S}_N^{(r)}(z) = \sum_{n \geq 0} P_{N,n}(r) \frac{z^{Nn}}{(Nn)!}.$$

Define the logarithmic coefficients $\kappa_{N,m}$ by

$$-\log \Delta_N(w) = \sum_{m \geq 1} \kappa_{N,m} \frac{w^m}{m}.$$

The polynomial dependence on r can be read directly from the formal logarithm. We use the complete exponential Bell polynomials B_n defined by

$$\exp\left(\sum_{m\geq 1} x_m \frac{t^m}{m!}\right) = \sum_{n\geq 0} B_n(x_1, \dots, x_n)\frac{t^n}{n!}.$$

Theorem 7.2 (Bell polynomial formula). *For every $n \geq 0$,*

$$P_{N,n}(r) = \frac{(Nn)!}{n!} B_n(r\kappa_{N,1}, r\,1!\,\kappa_{N,2}, \dots, r\,(n-1)!\,\kappa_{N,n}).$$

For $n \geq 1$, this is a polynomial in r of exact degree n. Its leading coefficient is

$$\frac{(Nn)!}{n!(N!)^n}.$$

Proof. The identity

$$\Delta_N(w)^{-r} = \exp\left(r\sum_{m\geq 1}\kappa_{N,m}\frac{w^m}{m}\right)$$

fits the defining generating function for B_n with

$$x_m = r(m-1)!\kappa_{N,m}.$$

The coefficient of w^n is therefore the displayed Bell polynomial divided by $n!$. Multiplication by $(Nn)!$ gives $P_{N,n}(r)$. Since

$$\Delta_N(w) = 1 - \frac{w}{N!} + O(w^2),$$

we have $\kappa_{N,1} = 1/N!$. The term x_1^n gives the stated leading coefficient, which is nonzero. □

The coefficients also satisfy a triangular recurrence that does not require the logarithmic coefficients to be known first.

Theorem 7.3 (Parameter recurrence). *We have $P_{N,0}(r) = 1$. For $n \geq 1$,*

$$\begin{aligned} P_{N,n}(r) = &-\sum_{k=1}^{n-1}(-1)^k\binom{Nn-1}{Nk}P_{N,n-k}(r) \\ &- r\sum_{k=1}^{n}(-1)^k\binom{Nn-1}{Nk-1}P_{N,n-k}(r). \end{aligned}$$

Proof. Set $S = \mathsf{D}_N^{-r}$. Logarithmic differentiation gives

$$\mathsf{D}_N S' + r\mathsf{D}_N' S = 0.$$

Taking the coefficient of $z^{Nn-1}/(Nn-1)!$ gives

$$\sum_{k=0}^{n-1}(-1)^k\binom{Nn-1}{Nk}P_{N,n-k}(r) + r\sum_{k=1}^{n}(-1)^k\binom{Nn-1}{Nk-1}P_{N,n-k}(r) = 0.$$

The term with $k = 0$ in the first sum is $P_{N,n}(r)$, and every other term has a smaller block index. Rearrangement gives the formula. □

Theorem 7.4 (Additive parameter law for reciprocal powers)**.** *For all $r, s \in \mathbb{C}$,*

$$\mathsf{S}_N^{(r)}(z)\mathsf{S}_N^{(s)}(z) = \mathsf{S}_N^{(r+s)}(z)$$

as a formal identity and on every common zero-free region with a fixed logarithm. Consequently,

$$P_{N,n}(r+s) = \sum_{q=0}^{n} \binom{Nn}{Nq} P_{N,q}(r) P_{N,n-q}(s).$$

Proof. Both statements follow by adding the exponents of the same logarithm and then multiplying the block exponential generating series. □

Corollary 7.5 (A sequence of binomial type)**.** *Define*

$$A_{N,n}(r) = \frac{n!}{(Nn)!} P_{N,n}(r).$$

Then

$$A_{N,n}(r+s) = \sum_{q=0}^{n} \binom{n}{q} A_{N,q}(r) A_{N,n-q}(s).$$

Thus $(A_{N,n})_{n\geq 0}$ is a polynomial sequence of binomial type.

Proof. Multiply the coefficient identity in theorem 7.4 by $n!/(Nn)!$ and simplify the factorials. □

The recurrence in theorem 7.3 uses the coefficients of the original denominator. A second recurrence uses only the logarithmic coefficients and is often shorter.

Theorem 7.6 (Logarithmic Newton recurrence)**.** *Set*

$$b_{N,n}(r) = \frac{P_{N,n}(r)}{(Nn)!}.$$

Then $b_{N,0}(r) = 1$ and for every $n \geq 1$,

$$n b_{N,n}(r) = r \sum_{k=1}^{n} \kappa_{N,k} b_{N,n-k}(r).$$

Equivalently,

$$P_{N,n}(r) = \frac{r}{n} \sum_{k=1}^{n} \frac{(Nn)!}{(N(n-k))!} \kappa_{N,k} P_{N,n-k}(r).$$

If $p_k = r\kappa_{N,k}$, then, for $n \geq 1$,

$$b_{N,n}(r) = \frac{1}{n!} \det \begin{pmatrix} p_1 & -1 & 0 & \cdots & 0 \\ p_2 & p_1 & -2 & \cdots & 0 \\ p_3 & p_2 & p_1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & -(n-1) \\ p_n & p_{n-1} & \cdots & p_2 & p_1 \end{pmatrix}.$$

Proof. Let

$$B_r(w) = \Delta_N(w)^{-r} = \sum_{n \geq 0} b_{N,n}(r) w^n.$$

Logarithmic differentiation gives

$$\frac{B_r'(w)}{B_r(w)} = r \sum_{k \geq 1} \kappa_{N,k} w^{k-1}.$$

Comparison of the coefficient of w^{n-1} proves the recurrence.

Let D_n denote the displayed determinant, with $D_0 = 1$. Expansion along the last row gives

$$D_n = \sum_{k=1}^{n} \frac{(n-1)!}{(n-k)!} p_k D_{n-k}.$$

Here the cofactor sign is canceled by the product of the negative superdiagonal entries $-(n-k+1), \ldots, -(n-1)$. Thus $D_n/n!$ satisfies the same recurrence and initial value as $b_{N,n}(r)$. □

Corollary 7.7 (The first three reciprocal polynomials)**.** *For every $N \geq 1$,*

$$P_{N,1}(r) = r,$$

$$P_{N,2}(r) = r\left[\frac{1}{2}\binom{2N}{N}(r+1) - 1\right],$$

and

$$P_{N,3}(r) = r\left[\frac{(3N)!}{6(N!)^3}(r+1)(r+2) - \binom{3N}{N}(r+1) + 1\right].$$

Proof. The first logarithmic coefficients are

$$\kappa_{N,1} = \frac{1}{N!},$$

$$\kappa_{N,2} = \frac{1}{(N!)^2} - \frac{2}{(2N)!},$$

and

$$\kappa_{N,3} = \frac{1}{(N!)^3} - \frac{3}{N!(2N)!} + \frac{3}{(3N)!}.$$

Insert these values into theorem 7.2 and simplify. □

Corollary 7.8. *For $n \geq 1$, the polynomial $P_{N,n}(r)$ is divisible by r.*

Proof. At $r = 0$, the reciprocal family is the constant function one. Every positive-index block coefficient therefore vanishes. □

Example 7.9. For $N = 1$, one has $\mathsf{D}_1(z)^{-r} = e^{rz}$ and $P_{1,n}(r) = r^n$. For $N = 2$,

$$P_{2,1}(r) = r,$$

$$P_{2,2}(r) = r(3r + 2),$$

and

$$P_{2,3}(r) = r(15r^2 + 30r + 16).$$

At $r = 1$, these are the positive secant coefficients.

8 Logarithmic derivatives and Riccati equations

A coefficient calculation at order two

Write $\Delta_2(w)^{-r} = \sum_{n\geq 0} b_n(r)w^n$. Since

$$\Delta_2(w) = 1 - \frac{w}{2} + \frac{w^2}{24} + O(w^3), \qquad -\log \Delta_2(w) = \frac{w}{2} + \frac{w^2}{12} + O(w^3),$$

the first logarithmic coefficients are $\kappa_{2,1} = 1/2$ and $\kappa_{2,2} = 1/6$. Newton's recurrence gives

$$b_1(r) = \frac{r}{2}, \qquad 2b_2(r) = r\left(\frac{r}{4} + \frac{1}{6}\right).$$

Restoring the factorial normalization yields $P_{2,1}(r) = r$ and $P_{2,2}(r) = r(3r + 2)$. At $r = 1$, the second value is five. No zero-location assumption is needed for this finite calculation.




Figure 4. The two exact parameter polynomials from the worked calculation. Each plotted curve evaluates its displayed formula. The points at $r = 1$ recover the first two positive-index secant coefficients.

Definition 8.1. Define the transfer tangent

$$\mathsf{T}_N(z) = -\frac{\mathsf{D}_N'(z)}{\mathsf{D}_N(z)}.$$

Near the origin, its convergent block expansion has the form

$$\mathsf{T}_N(z) = \sum_{n\geq 0} T_N(n) \frac{z^{Nn+N-1}}{(Nn+N-1)!}.$$

Proposition 8.2 (Logarithmic coefficient formula)**.** *The transfer tangent satisfies*

$$\mathsf{T}_N(z) = N \sum_{m\geq 1} \kappa_{N,m} z^{Nm-1}.$$

Equivalently,

$$T_N(n) = N(Nn+N-1)!\kappa_{N,n+1}.$$

Proof. Since $\mathsf{D}_N(z) = \Delta_N(z^N)$,

$$-\frac{\mathsf{D}_N'(z)}{\mathsf{D}_N(z)} = -Nz^{N-1}\frac{\Delta_N'(z^N)}{\Delta_N(z^N)}.$$

Differentiate the defining series for $-\log \Delta_N$ and compare coefficients. □

Theorem 8.3 (Tangent recurrence)**.** *The coefficients satisfy $T_N(0) = 1$ and*

$$T_N(n) = (-1)^n - \sum_{k=1}^{n} (-1)^k \binom{Nn+N-1}{Nk} T_N(n-k).$$

Proof. The defining equation is

$$\mathsf{D}_N \mathsf{T}_N = -\mathsf{D}_N'.$$

Extracting the coefficient of $z^{Nn+N-1}/(Nn+N-1)!$ gives

$$\sum_{k=0}^{n} (-1)^k \binom{Nn+N-1}{Nk} T_N(n-k) = (-1)^n.$$

Isolating the $k=0$ term gives the recurrence. □

Corollary 8.4 (Derivative of the reciprocal family)**.** *For every $r \in \mathbb{C}$, the formal identity*

$$\frac{d}{dz}\mathsf{S}_N^{(r)}(z) = r\mathsf{T}_N(z)\mathsf{S}_N^{(r)}(z)$$

holds. The same identity holds analytically on every zero-free domain on which the logarithm used to define $\mathsf{S}_N^{(r)}$ is fixed.

Proof. Differentiate D_N^{-r} on a zero-free region. The formal identity follows from the same logarithmic calculation. □

For a function g, define the complete differential Bell polynomial Y_N by

$$\frac{d^N}{dz^N} e^{g(z)} = e^{g(z)} Y_N(g', g'', \dots, g^{(N)}).$$

Theorem 8.5 (Scalar higher Riccati identity). *At every point of the open set* $\{z : \mathsf{D}_N(z) \neq 0\}$,

$$Y_N(-\mathsf{T}_N, -\mathsf{T}_N', \dots, -\mathsf{T}_N^{(N-1)}) = -1.$$

Proof. Write $\mathsf{D}_N = e^g$ locally. Then $g' = -\mathsf{T}_N$. Divide the equation $\mathsf{D}_N^{(N)} = -\mathsf{D}_N$ by D_N and use the defining identity for Y_N. □

This scalar identity follows from one ordinary differential equation. It does not redevelop the general nonlinear operation-tree theory of Chapter III.

Example 8.6. For $N = 1$, $\mathsf{T}_1 = 1$. For $N = 2$, $\mathsf{D}_2 = \cos z$ and $\mathsf{T}_2 = \tan z$. The Riccati equation becomes

$$\mathsf{T}_2' = 1 + \mathsf{T}_2^2.$$

9 Residue-section reciprocals and Appell systems

The section $\mathsf{E}_{N,j}$ has a forced zero of order j at the origin. Removing it leaves a unit series in z^N.

Definition 9.1. For $0 \leq j < N$, define

$$\Phi_{N,j}(w) = j! \sum_{n \geq 0} \frac{w^n}{(Nn + j)!}.$$

Then

$$\mathsf{E}_{N,j}(z) = \frac{z^j}{j!} \Phi_{N,j}(z^N).$$

Define the residue-section reciprocal function

$$\mathsf{C}_{N,j}(z) = \Phi_{N,j}(z^N)^{-1} = \frac{z^j/j!}{\mathsf{E}_{N,j}(z)}.$$

For $j > 0$, the last quotient at $z = 0$ is interpreted by its removable continuation, namely the first expression in the display. Write

$$\mathsf{C}_{N,j}(z) = \sum_{n \geq 0} C_{N,j}(n) \frac{z^{Nn}}{(Nn)!}.$$

Theorem 9.2 (Triangular inverse system)**.** *The coefficients are determined by* $C_{N,j}(0) = 1$ *and*

$$\sum_{q=0}^{n} \binom{Nn+j}{Nq} C_{N,j}(q) = \delta_{n0}.$$

For $n \geq 1$*,*

$$C_{N,j}(n) = -\frac{\displaystyle\sum_{q=0}^{n-1} \binom{Nn+j}{Nq} C_{N,j}(q)}{\binom{Nn+j}{Nn}}.$$

Proof. Multiply the defining series for $\mathsf{C}_{N,j}$ and $\mathsf{E}_{N,j}$. Their product is $z^j/j!$. At degree $Nn+j$, multiplication by $(Nn+j)!$ gives the stated binomial coefficient. The diagonal coefficient is nonzero, so the system is triangular. □

Definition 9.3. For $r \in \mathbb{C}$, define formal coefficients $C_{N,j,n}(r)$ by

$$\mathsf{C}_{N,j}(z)^r = \sum_{n \geq 0} C_{N,j,n}(r) \frac{z^{Nn}}{(Nn)!}.$$

On a simply connected region containing 0 and avoiding the zeros of $\Phi_{N,j}(z^N)$, interpret this power as e^{rL}, where L is the holomorphic logarithm of $\mathsf{C}_{N,j}$ normalized by $L(0) = 0$.

Theorem 9.4 (Parameter-addition law for residue-section reciprocals)**.** *For each* $n \geq 1$*, the coefficient* $C_{N,j,n}(r)$ *is a polynomial in* r *of exact degree* n*. Its leading coefficient is*

$$\frac{(Nn)!}{n!} \left(-\frac{j!}{(N+j)!} \right)^n.$$

It satisfies

$$C_{N,j,n}(r+s) = \sum_{q=0}^{n} \binom{Nn}{Nq} C_{N,j,q}(r) C_{N,j,n-q}(s).$$

In particular,

$$C_{N,j,n}(1) = C_{N,j}(n).$$

Proof. The unit series $\Phi_{N,j}(w)^{-1}$ has a formal logarithm with zero constant term. Its linear term is

$$-\frac{j!}{(N+j)!} w.$$

Exponentiation after multiplication by r shows that the coefficient of w^n is a polynomial of degree at most n. The nth power of the linear logarithmic term gives the displayed nonzero leading coefficient,

so the degree is exactly n. Multiplication of the powers with parameters r and s gives the coefficient identity. More explicitly, before the factorial normalization the coefficient of $r^n w^n$ is

$$\frac{1}{n!}\left(-\frac{j!}{(N+j)!}\right)^n;$$

thus multiplication by $(Nn)!$ gives the displayed leading coefficient. □

Definition 9.5. Define polynomials $E_m^{(N,j,r)}(x)$ by

$$\mathsf{C}_{N,j}(z)^r e^{xz} = \sum_{m\geq 0} E_m^{(N,j,r)}(x)\frac{z^m}{m!}.$$

Theorem 9.6 (Appell laws)**.** *For $m \geq 1$, the polynomials satisfy*

$$\frac{d}{dx}E_m^{(N,j,r)}(x) = mE_{m-1}^{(N,j,r)}(x)$$

and, for $m \geq 0$,

$$E_m^{(N,j,r)}(x+y) = \sum_{q=0}^{m}\binom{m}{q}E_q^{(N,j,r)}(x)y^{m-q}.$$

Proof. Differentiate the generating function with respect to x for the first identity. Multiply it by e^{yz} and compare coefficients for the second. □

Proposition 9.7 (Explicit Appell polynomial)**.** *For every $m \geq 0$,*

$$E_m^{(N,j,r)}(x) = \sum_{0\leq n\leq m/N}\binom{m}{Nn}C_{N,j,n}(r)x^{m-Nn}.$$

Proof. Multiply the block series for $\mathsf{C}_{N,j}^r$ by the exponential series and collect the coefficient of total degree m. □

Theorem 9.8 (Joint Appell convolution in the space and order parameters)**.** *For all $r, s, x, y \in \mathbb{C}$ and $m \geq 0$,*

$$E_m^{(N,j,r+s)}(x+y) = \sum_{q=0}^{m}\binom{m}{q}E_q^{(N,j,r)}(x)E_{m-q}^{(N,j,s)}(y).$$

Proof. Multiply the generating functions

$$\mathsf{C}_{N,j}(z)^r e^{xz} \quad \text{and} \quad \mathsf{C}_{N,j}(z)^s e^{yz}.$$

Their product is the generating function with parameters $r+s$ and $x+y$. Coefficient comparison gives the formula. □

Proposition 9.9 (Formal differential-operator representation on polynomials)**.** *Let a formal series* $A(z) = \sum_{q\geq 0} a_q z^q/q!$ *act on polynomials by*

$$A(\partial_x)f(x) = \sum_{q\geq 0} \frac{a_q}{q!}\partial_x^q f(x).$$

The power r *is applied to the formal symbol before the substitution* $z = \partial_x$*. Then*

$$E_m^{(N,j,r)}(x) = \mathsf{C}_{N,j}(\partial_x)^r x^m.$$

The inverse relation is

$$x^m = \Phi_{N,j}(\partial_x^N)^r E_m^{(N,j,r)}(x).$$

Both sums are finite on every polynomial.

Proof. Applying $A(\partial_x)$ to e^{xz} multiplies it by $A(z)$. Taking $A = \mathsf{C}_{N,j}^r$ gives the first identity by the defining generating function. Constant coefficient operators on polynomials satisfy

$$(AB)(\partial_x) = A(\partial_x)B(\partial_x),$$

because all derivative sums are finite and powers of ∂_x commute. The two formal symbols

$$\mathsf{C}_{N,j}(z)^r = \Phi_{N,j}(z^N)^{-r} \quad \text{and} \quad \Phi_{N,j}(z^N)^r$$

multiply to one. Their substitutions at $z = \partial_x$ are therefore inverse constant coefficient operators on polynomials, which proves the second identity. □

Corollary 9.10 (Appell symmetry)**.** *Each polynomial* $E_m^{(N,j,r)}$ *is monic of degree* m*, and*

$$E_m^{(N,j,r)}(\omega x) = \omega^m E_m^{(N,j,r)}(x).$$

Proof. The term $n = 0$ in proposition 9.7 is x^m. Every other exponent is $m - Nn$, so replacement of x by ωx multiplies every term by ω^m. □

Example 9.11. For $(N, j) = (2, 0)$,

$$\mathsf{C}_{2,0}(z) = \operatorname{sech} z.$$

For $(N, j) = (2, 1)$,

$$\mathsf{C}_{2,1}(z) = \frac{z}{\sinh z}.$$

The first case is a scaled and translated Euler family. In fact, if $E_m(x)$ denotes the classical Euler polynomial, then

$$E_m^{(2,0,1)}(x) = 2^m E_m\left(\frac{x+1}{2}\right).$$

If $B_m(x)$ denotes the classical Bernoulli polynomial, the second case has the exact identification

$$E_m^{(2,1,1)}(x) = 2^m B_m\left(\frac{x+1}{2}\right),$$

obtained by substituting $t = 2z$ in $te^{yt}/(e^t - 1)$.

10 Scaled reciprocal coupling and theta polynomials

Definition 10.1. Set

$$H_N(z) = \mathsf{E}_{N,0}(z) = \sum_{n\geq 0} \frac{z^{Nn}}{(Nn)!}.$$

For $u \in \mathbb{C}$, define $\Theta_{N,n}(u)$ by

$$H_N(uz)^{-1} H_N(z)^{-u} = \sum_{n\geq 0} \Theta_{N,n}(u) \frac{z^{Nn}}{(Nn)!}.$$

This is first a formal identity at the origin. Analytically, it is interpreted on a common simply connected zero-free region with the branch of $\log H_N$ normalized by $\log H_N(0) = 0$.

Theorem 10.2 (Theta reconstruction)**.** *For every $n \geq 0$,*

$$\Theta_{N,n}(u) = (-1)^n \sum_{q=0}^{n} \binom{Nn}{Nq} P_{N,q}(1) u^{Nq} P_{N,n-q}(u).$$

Thus $\Theta_{N,n}$ is a polynomial of degree at most Nn.

Proof. Choose ρ with $\rho^N = -1$. Then

$$H_N(z) = \mathsf{D}_N(\rho z).$$

The coefficient of z^{Nn} in $H_N(z)^{-r}$ is therefore $(-1)^n P_{N,n}(r)/(Nn)!$. Apply this to $H_N(uz)^{-1}$ and $H_N(z)^{-u}$, then multiply the two block series. □

Proposition 10.3 (Trivial parameter factors)**.** *For $n \geq 1$, the polynomial $\Theta_{N,n}(u)$ is divisible by u. If N is even, it is also divisible by $u + 1$.*

Proof. At $u = 0$, the defining product is one. Hence every positive-index coefficient vanishes. If N is even, then $H_N(-z) = H_N(z)$. At $u = -1$, the product is again one, so every positive-index coefficient vanishes there as well. □

Proposition 10.4 (Theta Newton recurrence)**.** *Put*

$$\theta_{N,n}(u) = \frac{\Theta_{N,n}(u)}{(Nn)!}.$$

Then $\theta_{N,0}(u) = 1$ and

$$n\theta_{N,n}(u) = \sum_{m=1}^{n} (-1)^m \kappa_{N,m}(u^{Nm} + u)\theta_{N,n-m}(u).$$

In particular,

$$\Theta_{N,1}(u) = -(u^N + u).$$

Proof. Write $w = z^N$. Since $H_N(z) = \Delta_N(-w)$,

$$-\log H_N(z) = \sum_{m\geq 1} (-1)^m \kappa_{N,m} \frac{w^m}{m}.$$

The logarithm of the theta generating function is therefore

$$\sum_{m\geq 1} (-1)^m \kappa_{N,m} (u^{Nm} + u) \frac{w^m}{m}.$$

Logarithmic differentiation and coefficient comparison give the recurrence. The case $n = 1$ uses $\kappa_{N,1} = 1/N!$. □

Example 10.5. For $N = 2$,

$$\Theta_{2,1}(u) = -u(u+1)$$

and

$$\Theta_{2,2}(u) = u(u+1)(5u^2 + u + 2).$$

The two factors predicted by proposition 10.3 are visible.

Part III The geometry of zeros

11 Growth and the geometry of the large zeros

A route through this section

The finite exponential representation gives a polygon of exponents. Large zeros can occur where two dominant terms nearly cancel. The proof first localizes those zeros and then uses symmetry and uniqueness to place them on the rays. Finite exceptional zeros remain part of the later products and sums.




Figure 5. The exact order-four model. The first three zeros on each of four rays are mapped by $w = z^4$ to three positive reduced zeros. The reduced axis uses a logarithmic scale. This complete order-four example does not classify the finite exceptional set at other orders.

We now assume $N \geq 2$. Write

$$\eta_k = e^{(2k+1)\pi i/N}, \qquad 0 \leq k < N.$$

These are the vertices of a regular polygon and

$$\mathsf{D}_N(z) = \frac{1}{N} \sum_{k=0}^{N-1} e^{\eta_k z}.$$

Standard Fact 11.1 (Order from Taylor coefficients)**.** If $f(w) = \sum_{n\geq 0} a_n w^n$ is an entire function with infinitely many nonzero coefficients, then

$$\operatorname{ord} f = \limsup_{\substack{n\to\infty \\ a_n\neq 0}} \frac{n \log n}{\log(1/|a_n|)}.$$

Canonical products for entire functions of order below one have genus zero. We use these standard facts in the form given in [Boa54, Tit39].

Proposition 11.2 (Orders and exponential type)**.** *For $N \geq 2$,*

$$\operatorname{ord} \Delta_N = \frac{1}{N}.$$

The function D_N has order one and exponential type one.

Proof. The coefficient of w^n in Δ_N has modulus $1/(Nn)!$. Stirling's formula and standard fact 11.1 give order $1/N$. Since $M_{f(\cdot^N)}(r) = M_f(r^N)$ for the maximum modulus, composition with z^N multiplies the order by N and gives order one for D_N.

The finite exponential representation gives

$$|\mathsf{D}_N(z)| \leq e^{|z|},$$

so the type is at most one. Choose a root η_0 and put $z = t\overline{\eta_0}$ with $t > 0$. The term $e^{\eta_0 z} = e^t$ has strictly larger real exponent than every other term. Hence

$$\mathsf{D}_N(t\overline{\eta_0}) = \frac{1}{N} e^t (1 + o(1)).$$

The type is therefore one. □

Definition 11.3. For $\theta \in \mathbb{R}$, define

$$h_N(\theta) = \max_{0\leq k<N} \cos\left(\theta + \frac{(2k+1)\pi}{N}\right).$$

The tie directions are

$$\mathfrak{T}_N = \left\{ \frac{2\pi k}{N} : 0 \leq k < N \right\} \mod 2\pi.$$

Theorem 11.4 (Indicator polygon and dominant sectors)**.** *The indicator of D_N equals h_N, with the limsup taken over radii where $\mathsf{D}_N(re^{i\theta}) \neq 0$:*

$$\limsup_{r\to\infty} \frac{1}{r} \log |\mathsf{D}_N(re^{i\theta})| = h_N(\theta).$$

If K is a compact set of directions disjoint from $\mathfrak{T}_N$, then there are $\delta > 0$ and $R > 0$ such that, for each $\theta \in K$, one root $\eta(\theta)$ of -1 satisfies

$$\mathsf{D}_N(re^{i\theta}) = \frac{1}{N}\exp(\eta(\theta)re^{i\theta})\left(1 + O(e^{-\delta r})\right)$$

uniformly for $r \geq R$. In particular, no sufficiently large zero can lie in those directions.

Proof. For $z = re^{i\theta}$, the real part of the exponent $\eta_k z$ is

$$r\cos\left(\theta + \frac{(2k+1)\pi}{N}\right).$$

The triangle inequality gives the upper indicator bound $h_N(\theta)$.

Away from a tie direction, the maximizing vertex is unique. It is constant on each connected component of the complement of the tie directions. A compact set avoiding all ties meets only finitely many such components, and compactness gives one uniform positive lower bound for every relevant projection gap. Division by the maximizing exponential gives the stated uniform asymptotic and the zero-free conclusion.

At a tie direction exactly two adjacent vertices attain the maximum. After rotation it is enough to take $\theta = 0$. With $c_N = \cos(\pi/N)$ and $s_N = \sin(\pi/N)$, the two dominant terms give

$$\frac{2}{N}e^{c_N r}\cos(s_N r)$$

and every other term is exponentially smaller. Along a sequence on which $|\cos(s_N r)| \geq 1/2$, the lower indicator bound is $c_N = h_N(0)$. Rotational symmetry treats every tie direction. □

The zero strings are governed by two adjacent exponents. Put

$$a_N = \frac{\pi}{N}, \qquad s_N = \sin a_N, \qquad c_N = \cos a_N,$$

and

$$A_N = \frac{\pi}{s_N}, \qquad t_m = A_N\left(m + \frac{1}{2}\right).$$

For $N \geq 3$, also put

$$\gamma_N = c_N - \cos(3a_N) > 0.$$

Lemma 11.5 (Two term dominance)**.** *Assume $N \geq 3$. There are positive constants Y, R, and C with the following properties.*

If $z = x + iy$ lies in the sector $|\arg z| \leq a_N$, has $x \geq R$, and is a zero of D_N, then $|y| \leq Y$.

On the strip $x \geq R$ and $|y| \leq Y$, define

$$R_N(z) = \sum_{\substack{\eta^N = -1 \\ \eta \neq e^{ia_N}, e^{-ia_N}}} \exp((\eta - e^{-ia_N})z).$$

Then

$$Ne^{-e^{-ia_N}z}\mathsf{D}_N(z) = 1 + e^{2is_N z} + R_N(z),$$

and, for $q = 0, 1, 2$,

$$|R_N^{(q)}(z)| \le Ce^{-\gamma_N x}.$$

Proof. A direct projection estimate gives the stronger bound

$$Y = \frac{\log(N-1)}{2s_N}$$

for every zero in the sector, without a large-radius assumption. Write $z = re^{i\theta}$ with $0 \le \theta \le a_N$ and $r \ge 0$. For $0 \le \ell \le N-2$, the angle $\theta + (2\ell+1)a_N$ lies between $a_N + \theta$ and $2\pi - a_N - \theta$. Monotonicity of cosine on $[0, \pi]$, followed by reflection about π, therefore gives

$$\Re(\eta_\ell z) \le r\cos(a_N + \theta).$$

The remaining vertex e^{-ia_N} has projection $r\cos(a_N - \theta)$. Their difference is at least $2s_N r\sin\theta = 2s_N y$. If $\mathsf{D}_N(z) = 0$, divide its finite exponential sum by the term with exponent $e^{-ia_N}z$ and apply the triangle inequality:

$$1 \le (N-1)e^{-2s_N y}.$$

Thus $y \le Y$. Conjugation gives the same bound for $-y$ on the lower half sector. Since $N \ge 3$ and $s_N > 0$, this Y is positive.

On the resulting strip, the roots e^{ia_N} and e^{-ia_N} have real part c_N, while each remaining root $\eta = u+iv$ satisfies $u \le \cos(3a_N) = c_N - \gamma_N$. Hence

$$\Re((\eta - e^{-ia_N})z) \le -\gamma_N x + 2Y.$$

Division by $e^{e^{-ia_N}z}$ gives the displayed normalization. Each differentiation multiplies a remainder term by $\eta - e^{-ia_N}$, whose modulus is at most two. Thus, for $q = 0, 1, 2$,

$$|R_N^{(q)}(z)| \le (N-2)2^q e^{2Y}e^{-\gamma_N x}.$$

One may take $R = 1$ and $C = 4(N-2)e^{2Y}$. □

Lemma 11.6 (Near cancellation of two exponentials)**.** *For every* $M > 0$*, there are constants* $\varepsilon_M > 0$ *and* $C_M > 0$ *such that the following holds. If*

$$|\Re w| \le M$$

and

$$|1 + e^w| \le \varepsilon_M,$$

then there is a unique integer m *with*

$$|w - (2m+1)\pi i| \le C_M|1 + e^w|.$$

Proof. In fact, the constants $\varepsilon_M = 1/2$ and $C_M = 3/2$ work for every complex w, without the strip restriction. Put $u = -(1 + e^w)$ and $L = \log(1+u) = \log(-e^w)$, using the principal logarithm. The elementary logarithm estimate on $|u| \le 1/2$ gives $|L| \le (3/2)|u|$. Since $e^{L+\pi i} = e^w$, there is an integer m such that

$$w = L + (2m+1)\pi i.$$

This gives the required distance bound. Distinct centers are separated by at least 2π, whereas two centers satisfying the bound would be separated by at most $3|1+e^w| \le 3/2$. Thus the integer is unique. The formal proof uses the convergent logarithm estimate and the exact periodicity theorem for the complex exponential. □

Theorem 11.7 (Large zero strings). *Assume $N \ge 3$. There are constants m_0, C, and $\gamma > 0$ and a sequence of positive real simple zeros $(\tau_m)_{m\ge m_0}$ of D_N such that*

$$\tau_m = t_m + O(e^{-\gamma t_m}).$$

Every zero of sufficiently large modulus is of the form

$$\omega^k \tau_m$$

for a unique $0 \le k < N$ and a unique $m \ge m_0$.
The derivatives satisfy

$$\mathsf{D}_N'(\omega^k \tau_m) = \omega^{-k} \mathsf{D}_N'(\tau_m)$$

and

$$|\mathsf{D}_N'(\tau_m)| = \frac{2s_N}{N} e^{c_N \tau_m} \left(1 + O(e^{-\gamma \tau_m})\right).$$

Proof. Work first in the sector $|\arg z| \le a_N$. By lemma 11.5, every sufficiently large zero lies in a fixed strip about the positive real axis. Put

$$G(z) = N e^{-e^{-ia_N} z} \mathsf{D}_N(z).$$

Then

$$G(z) = 1 + e^{2is_N z} + R_N(z)$$

with an exponentially small remainder and derivative.

If $G(z) = 0$, then

$$|1 + e^{2is_N z}| \le C e^{-\gamma_N \Re z}.$$

The real part of $2is_N z$ is bounded in the strip. Apply lemma 11.6 with $w = 2is_N z$. There is a unique integer m such that

$$|z - t_m| \le C_1 e^{-\gamma_N \Re z},$$

where

$$t_m = \frac{(2m+1)\pi}{2s_N}.$$

In particular, $|\Re z - t_m| = o(1)$. For all sufficiently large zeros we have $|\Re z - t_m| \le 1$, and hence

$$|z - t_m| \le C_2 e^{-\gamma_N t_m}$$

with a constant independent of m. Because $\Re z \to +\infty$ in this argument, the corresponding integers m are eventually nonnegative and tend to infinity.

Fix a large m, put $a = 2is_N$, and consider the closed disk $B_m = \{z : |z - t_m| \le 1/(2s_N)\}$. The remainder bounds hold uniformly on these fixed-radius disks. For sufficiently large m, $|R_N| \le 1/2$ and $|R_N'| \le s_N/2$ there. Define

$$\Phi_m(z) = t_m + \frac{\log(1 + R_N(z))}{2is_N},$$

using the principal logarithm. The estimate $|\log(1+u)| \le (3/2)|u|$ shows that Φ_m maps B_m into itself. The derivative bound $|\Phi_m'| \le 1/2$ makes it a contraction. Banach's fixed-point theorem gives a unique fixed point. On B_m, $|a(z - t_m)| \le 1 < \pi$, so the principal logarithm inverts the exponential and the fixed-point equation is equivalent to $G(z) = 0$. Moreover,

$$|z - t_m| \le \frac{3}{4s_N}|R_N(z)| = O(e^{-\gamma_N t_m}).$$

At this zero,

$$|G'(z)| \ge |a|\,|1 + R_N(z)| - |R_N'(z)| \ge s_N/2 > 0,$$

so the zero is simple. The disks are mutually disjoint because their diameter $1/s_N$ is smaller than the center spacing π/s_N. Enlarging the strip if necessary, these disks lie inside the fundamental sector and the fixed strip for large m.

The localization estimate puts every sufficiently large zero in the fundamental sector into one of these disks. The zero sets of G and D_N coincide because their prefactor is nonzero. Each disk is invariant under complex conjugation, and D_N has real coefficients. The conjugate of its unique zero is therefore the same zero. It is real, positive for large m, and simple by the derivative bound above. Denote it by τ_m.

The parts of the fundamental sector away from the positive tie ray, including its boundary rays, were already excluded uniformly. Rotational invariance now gives every remaining large zero as $\omega^k\tau_m$. Distinct rays and the disjoint disks give uniqueness of k and m. Taking $\gamma = \gamma_N$ proves the zero location in the statement.

Differentiate the normalized expression. At $z = \tau_m$,

$$G'(\tau_m) = 2is_N e^{2is_N\tau_m} + R_N'(\tau_m) = -2is_N + O(e^{-\gamma_N\tau_m}).$$

Since $G(\tau_m) = 0$,

$$\mathsf{D}_N'(\tau_m) = \frac{1}{N}e^{e^{-ia_N\tau_m}}G'(\tau_m).$$

Taking absolute values gives the derivative asymptotic. Differentiating $\mathsf{D}_N(\omega z) = \mathsf{D}_N(z)$ gives the rotation formula. □

Theorem 11.8 (Leading displacement term and derivative sign). *With the notation of lemma 11.5 and theorem 11.7,*

$$\tau_m = t_m + \frac{R_N(t_m)}{2is_N} + O(e^{-2\gamma_N t_m}).$$

The quotient $R_N(t_m)/(2is_N)$ is real. One also has

$$\mathsf{D}_N'(\tau_m) = (-1)^{m+1}\frac{2s_N}{N}e^{c_N\tau_m}\left(1 + O(e^{-\gamma_N\tau_m})\right).$$

Proof. Put $\delta_m = \tau_m - t_m$. The large zero theorem gives $\delta_m = O(e^{-\gamma_N t_m})$. Since $e^{2is_N t_m} = -1$,

$$1 + e^{2is_N(t_m+\delta_m)} = -2is_N\delta_m + O(\delta_m^2).$$

The remainder estimate through the first derivative gives

$$R_N(t_m + \delta_m) = R_N(t_m) + O(e^{-\gamma_N t_m}|\delta_m|).$$

Substitution in the zero equation gives the displacement formula.

At the real point t_m, the normalized value

$$Ne^{-e^{-ia_N}t_m}\mathsf{D}_N(t_m)$$

is purely imaginary. The two dominant terms cancel there, so $R_N(t_m)$ is purely imaginary and the displayed correction is real.

For the derivative, use the formula in the proof of theorem 11.7 and

$$e^{-is_N\tau_m} = (-1)^m(-i)\left(1 + O(e^{-\gamma_N\tau_m})\right).$$

The stated real sign follows. □

Corollary 11.9 (Leading correction for $N = 3$)**.** *For $N = 3$,*

$$t_m = \frac{2\pi}{\sqrt{3}}\left(m + \frac{1}{2}\right)$$

and

$$\tau_m = t_m + \frac{(-1)^m}{\sqrt{3}}e^{-3t_m/2} + O(e^{-3t_m}).$$

Proof. The only nondominant exponent is -1. Hence

$$R_3(t) = \exp\left(\left(-\frac{3}{2} + \frac{\sqrt{3}}{2}i\right)t\right).$$

At $t = t_m$, its phase is $i(-1)^m$. Insert this value into theorem 11.8. □

Corollary 11.10 (The case $N = 2$)**.** *For $N = 2$,*

$$\mathsf{D}_2(z) = \cos z.$$

Its zeros are

$$\pm\left(m + \frac{1}{2}\right)\pi, \qquad m \geq 0,$$

and all are simple. They agree with the formula $t_m = A_2(m + 1/2)$.

Proof. This is the standard zero set of the cosine function. □

Corollary 11.11 (Eventual simplicity and exact rays)**.** *For every $N \geq 2$, all but finitely many zeros of D_N are simple. Every sufficiently large zero lies on one of the rays*

$$\omega^k(0,\infty), \qquad 0 \leq k < N.$$

Proof. Use theorem 11.7 for $N \geq 3$ and corollary 11.10 for $N = 2$. □

Example 11.12 (Exact fourth-order strings)**.** For $N = 4$,

$$\mathsf{D}_4(z) = \cos\left(\frac{z}{\sqrt{2}}\right)\cosh\left(\frac{z}{\sqrt{2}}\right).$$

The positive zeros are exactly

$$\tau_m = \sqrt{2}\pi\left(m + \frac{1}{2}\right) = A_4\left(m + \frac{1}{2}\right).$$

Rotation by the fourth roots of unity gives the real and imaginary zero strings. In this case the error term in theorem 11.7 is identically zero.

Corollary 11.13 (Reduced zeros)**.** *Let Λ_N be the nonzero zero multiset of Δ_N. All but finitely many members of Λ_N are positive and simple. They can be indexed as*

$$\lambda_m = \tau_m^N$$

for all sufficiently large m. If $N \geq 3$, the notation may be chosen as in theorem 11.7, and

$$\lambda_m = t_m^N + O\left(t_m^{N-1} e^{-\gamma t_m}\right).$$

For $N = 2$, take $\tau_m = t_m = \pi(m + 1/2)$ for $m \geq 0$. Then $\lambda_m = t_m^2$ exactly.

Proof. The identity $\mathsf{D}_N(z) = \Delta_N(z^N)$ identifies each orbit of N nonzero zeros of D_N with one zero of Δ_N. For $N \geq 3$, the large orbit representatives are the positive zeros τ_m from theorem 11.7. For $N = 2$, use corollary 11.10. Simplicity is preserved because

$$\mathsf{D}_N'(z) = N z^{N-1} \Delta_N'(z^N)$$

and $z \neq 0$. The asymptotic follows by expanding τ_m^N about t_m. □

Theorem 11.14 (Zero counting asymptotics)**.** *Let $n_{\mathsf{D}_N}(R)$ count the zeros of D_N in $|z| \leq R$, with multiplicity. Let $n_{\Delta_N}(R)$ count the zeros of Δ_N in $|w| \leq R$. Then, as $R \to \infty$,*

$$n_{\mathsf{D}_N}(R) = \frac{N \sin(\pi/N)}{\pi} R + O(1)$$

and

$$n_{\Delta_N}(R) = \frac{\sin(\pi/N)}{\pi} R^{1/N} + O(1).$$

Proof. Each of the N rays contains one large zero near

$$t_m = \frac{\pi}{\sin(\pi/N)} \left(m + \frac{1}{2} \right).$$

Since $\tau_m - t_m = o(1)$ and the model spacing is the positive constant A_N, replacing $\tau_m \leq R$ by $t_m \leq R$ changes the count by at most a bounded amount. The number of such indices with $t_m \leq R$ is

$$\frac{\sin(\pi/N)}{\pi} R + O(1).$$

Multiplication by N and addition of the finite exceptional set give the first formula. The second follows from

$$n_{\mathsf{D}_N}(R^{1/N}) = N n_{\Delta_N}(R).$$

□

Corollary 11.15 (Zero exponents)**.** *The exponent of convergence of the nonzero zeros of D_N is one. The exponent of convergence of the zeros of Δ_N is $1/N$.*

Proof. The asymptotic counting formulas in theorem 11.14 give the two exponents by partial summation. □

Part IV Products, moments, and zero zeta functions

12 Hadamard products and zero power sums

The central idea

The product uses every reduced zero with its multiplicity. Its logarithm turns multiplication into addition, so reciprocal-zero power sums become coefficients. This step depends on the global product theorem. A truncated zero list provides a numerical partial sum, while the identity uses the entire multiset.




Figure 6. Order-two moment sums over the first forty reduced zeros $\lambda_k = \pi^2(k + 1/2)^2$. The curves are finite numerical partial sums. The dashed limits $1/2$ and $1/6$ follow from the exact logarithmic coefficients and the product identity.

The reduction from z to $w = z^N$ removes the rotational repetition of the zeros.

Theorem 12.1 (Genus zero product). *For $N \geq 2$, let Λ_N be the nonzero zero multiset of Δ_N. Then*

$$\Delta_N(w) = \prod_{\lambda \in \Lambda_N} \left(1 - \frac{w}{\lambda}\right),$$

where the product converges locally uniformly. Consequently,

$$\mathsf{D}_N(z) = \prod_{\lambda \in \Lambda_N} \left(1 - \frac{z^N}{\lambda}\right).$$

Proof. By proposition 11.2, the order of Δ_N is $1/N < 1$. By corollary 11.13, it has infinitely many zeros, and corollary 11.15 gives

$$\sum_{\lambda \in \Lambda_N} |\lambda|^{-1} < \infty.$$

Hadamard factorization therefore gives a canonical product of genus zero. The only possible exponential factor is a nonzero constant because the order is below one. Since $\Delta_N(0) = 1$, that constant is one. Substitution of $w = z^N$ gives the second formula. □

Theorem 12.2 (Diagonal Fredholm determinant representation)**.** *Let I_N be a multiplicity-sensitive countable index set for Λ_N and put $\mathcal{H}_N = \ell^2(I_N)$. On its standard orthonormal basis, define the diagonal operator*

$$K_N e_\lambda = \lambda^{-1} e_\lambda.$$

Then K_N is trace class and

$$\Delta_N(w) = \det_F(I - wK_N).$$

For every $m \geq 1$,

$$\kappa_{N,m} = \operatorname{tr}(K_N^m).$$

Proof. By corollary 11.15,

$$\sum_{\lambda \in \Lambda_N} |\lambda|^{-1} < \infty.$$

Thus the diagonal operator K_N is trace class. Its Fredholm determinant is the locally uniform product

$$\det_F(I - wK_N) = \prod_{\lambda \in \Lambda_N} \left(1 - \frac{w}{\lambda}\right),$$

which equals $\Delta_N(w)$ by theorem 12.1.

Taking the logarithm gives

$$-\log \Delta_N(w) = \sum_{m \geq 1} \frac{w^m}{m} \operatorname{tr}(K_N^m)$$

near the origin. Comparison with the definition of $\kappa_{N,m}$ proves the trace identity. □

This is the standard diagonal encoding of this family's genus zero product. It is not a general trace-class resolvent or Feshbach theorem.

Corollary 12.3 (Complete symmetric function form)**.** *For a positive integer r, let each eigenvalue λ^{-1} of K_N be repeated r times. If h_n denotes the complete homogeneous symmetric function of degree n in this absolutely summable multiset, then*

$$\frac{P_{N,n}(r)}{(Nn)!} = h_n.$$

For arbitrary complex r, the Bell polynomial formula is the polynomial continuation obtained by multiplying every power sum by r.

Proof. The generating function for the complete homogeneous symmetric functions is

$$\prod_{\lambda \in \Lambda_N} \left(1 - \frac{w}{\lambda}\right)^{-r} = \Delta_N(w)^{-r}.$$

Coefficient comparison gives the first identity. Power sums determine the complete homogeneous functions through the Newton recurrence. Replacing each power sum by r times itself gives theorems 7.2 and 7.6. □

Proposition 12.4 (Products for the normalized sections)**.** *For every $0 \le j < N$, the entire function*

$$\Phi_{N,j}(w) = j! \sum_{n \ge 0} \frac{w^n}{(Nn+j)!}$$

has order $1/N$. If $N \ge 2$ and $\Lambda_{N,j}$ is its nonzero zero multiset, then

$$\Phi_{N,j}(w) = \prod_{\mu \in \Lambda_{N,j}} \left(1 - \frac{w}{\mu}\right).$$

For $m \ge 1$, put

$$\sigma_{N,j,m} = \sum_{\mu \in \Lambda_{N,j}} \mu^{-m}.$$

Then

$$C_{N,j,n}(r) = \frac{(Nn)!}{n!} B_n(r\sigma_{N,j,1}, r1!\sigma_{N,j,2}, \dots, r(n-1)!\sigma_{N,j,n}).$$

Proof. The coefficient formula for the order gives $1/N$. The function is not a polynomial. An entire function of order below one with only finitely many zeros would be a polynomial times a constant by Hadamard factorization, which is impossible here. The genus zero product follows as in theorem 12.1.

Taking the logarithm gives

$$-\log \Phi_{N,j}(w) = \sum_{m \ge 1} \sigma_{N,j,m} \frac{w^m}{m}.$$

Since $\mathsf{C}_{N,j}(z)^r = \Phi_{N,j}(z^N)^{-r}$, the Bell polynomial calculation from theorem 7.2 gives the coefficient formula. □

Definition 12.5. For $m \ge 1$, define

$$\sigma_{N,m} = \sum_{\lambda \in \Lambda_N} \lambda^{-m},$$

with multiplicities.

The series converges absolutely because the exponent of convergence of Λ_N is $1/N < 1$.

Theorem 12.6 (Newton sums for the reduced zeros)**.** *For $|w|$ below the least zero modulus, with the analytic logarithm normalized to zero at the origin,*

$$-\log \Delta_N(w) = \sum_{m \ge 1} \sigma_{N,m} \frac{w^m}{m}.$$

Thus

$$\kappa_{N,m} = \sigma_{N,m}.$$

If

$$a_m = \frac{(-1)^m}{(Nm)!}, \qquad a_0 = 1,$$

then

$$\sigma_{N,m} = -m a_m - \sum_{k=1}^{m-1} a_k \sigma_{N,m-k}.$$

Proof. Take the logarithm of the product in theorem 12.1. Absolute convergence in a disk below the first zero permits termwise expansion

$$-\log\left(1 - \frac{w}{\lambda}\right) = \sum_{m \geq 1} \frac{w^m}{m \lambda^m}.$$

This gives the logarithmic formula and identifies $\kappa_{N,m}$ with $\sigma_{N,m}$.

Differentiate the product identity in logarithmic form to obtain

$$\Delta_N'(w) = -\Delta_N(w) \sum_{m \geq 1} \sigma_{N,m} w^{m-1}.$$

Comparison of the coefficient of w^{m-1} gives

$$m a_m = -\sum_{k=0}^{m-1} a_k \sigma_{N,m-k}.$$

Solving for $\sigma_{N,m}$ gives the recurrence. □

Corollary 12.7 (Convergent Mittag–Leffler partial-fraction form)**.** *Away from the zeros of* D_N*,*

$$\mathsf{T}_N(z) = N z^{N-1} \sum_{\lambda \in \Lambda_N} \frac{1}{\lambda - z^N}.$$

The series converges absolutely and locally uniformly on compact sets away from the poles. Thus no regularizing terms are required.

Proof. Logarithmic differentiation of the genus zero product gives

$$-\frac{\Delta_N'(w)}{\Delta_N(w)} = \sum_{\lambda \in \Lambda_N} \frac{1}{\lambda - w}.$$

The series is locally uniform because $\sum_\lambda |\lambda|^{-1} < \infty$. Substitute $w = z^N$ and use the chain rule. □

Proposition 12.8 (Theta Bell polynomial formula)**.** *Put*

$$\tilde{\sigma}_{N,m} = (-1)^m \sigma_{N,m}.$$

Then

$$\Theta_{N,n}(u) = \frac{(Nn)!}{n!} B_n(\tilde{\sigma}_{N,1}(u^N + u), 1!\tilde{\sigma}_{N,2}(u^{2N} + u), \dots, (n-1)!\tilde{\sigma}_{N,n}(u^{nN} + u)).$$

Proof. The reduced unsigned denominator is

$$\Psi_N(w) = \sum_{n\geq 0} \frac{w^n}{(Nn)!} = \Delta_N(-w).$$

By theorem 12.6,

$$-\log \Psi_N(w) = \sum_{m\geq 1} \tilde{\sigma}_{N,m} \frac{w^m}{m}.$$

With $w = z^N$, the logarithm of the theta generating function is

$$\sum_{m\geq 1} \tilde{\sigma}_{N,m}(u^{Nm} + u)\frac{w^m}{m}.$$

The complete Bell polynomial formula follows. □

Corollary 12.9 (Tangent coefficients as zero sums)**.** *For every $n \geq 0$,*

$$T_N(n) = N(Nn + N - 1)!\sigma_{N,n+1}.$$

Proof. Combine proposition 8.2 and theorem 12.6. □

Corollary 12.10 (Reciprocal coefficients from zero data)**.** *For every $n \geq 0$,*

$$P_{N,n}(r) = \frac{(Nn)!}{n!} B_n(r\sigma_{N,1}, r1!\sigma_{N,2}, \dots, r(n-1)!\sigma_{N,n}).$$

Proof. Use theorems 7.2 and 12.6. □

Example 12.11 (The cosine orbit sums)**.** For $N = 2$,

$$\Delta_2(w) = \cos\sqrt{w}.$$

Its zeros are

$$\lambda_m = \pi^2 \left(m + \frac{1}{2}\right)^2, \qquad m \geq 0.$$

Hence

$$\sigma_{2,q} = \frac{2^{2q} - 1}{\pi^{2q}} \zeta(2q).$$

The identity in corollary 12.9 recovers the classical tangent coefficients from these zero power sums.

13 Coefficient asymptotics from the nearest zeros

For an entire function f, let $\mathcal{Z}(f)$ denote its multiset of nonzero zeros. Let r be a positive integer. Then D_N^{-r} is meromorphic and a simple zero of D_N becomes a pole of order r.

Theorem 13.1 (Nearest-pole coefficient expansion). *Let $r \in \mathbb{Z}_{>0}$. Suppose the zeros of D_N of least modulus R are simple and form the finite set $\mathcal{Z}_R$. Suppose the next zero modulus is $R_2 > R$. At each $\rho \in \mathcal{Z}_R$, write the principal part*

$$\mathsf{D}_N(z)^{-r} = \sum_{\ell=1}^{r} a_{\rho,\ell} \left(1 - \frac{z}{\rho}\right)^{-\ell} + O(1).$$

Then, for every R_* with $R < R_* < R_2$ and $m = Nn$, as $n \to \infty$,

$$P_{N,n}(r) = m! \sum_{\rho \in \mathcal{Z}_R} \sum_{\ell=1}^{r} a_{\rho,\ell} \binom{m+\ell-1}{\ell-1} \rho^{-m} + O_{R_*}(m! R_*^{-m}).$$

The leading Laurent coefficient is

$$a_{\rho,r} = (-\rho \mathsf{D}_N'(\rho))^{-r}.$$

Proof. Subtract the complete principal parts at the zeros on $|z| = R$. The remainder is holomorphic on $|z| < R_2$, hence on a neighborhood of every closed disk $|z| \le R_* < R_2$. Cauchy's estimate gives an ordinary coefficient bound $O_{R_*}(R_*^{-m})$. The coefficient of z^m in

$$\left(1 - \frac{z}{\rho}\right)^{-\ell}$$

is

$$\binom{m+\ell-1}{\ell-1} \rho^{-m}.$$

Multiplication by $m!$ converts the ordinary coefficient to $P_{N,n}(r)$.

Near a simple zero,

$$\mathsf{D}_N(z) = -\rho \mathsf{D}_N'(\rho) \left(1 - \frac{z}{\rho}\right) + O\left(\left(1 - \frac{z}{\rho}\right)^2\right).$$

Raising to the power $-r$ gives the formula for $a_{\rho,r}$. □

Corollary 13.2 (Leading dominant zero term). *Under the hypotheses of theorem 13.1,*

$$P_{N,n}(r) = m! \binom{m+r-1}{r-1} \sum_{\rho \in \mathcal{Z}_R} \frac{\rho^{-m}}{(-\rho \mathsf{D}_N'(\rho))^r} + O\left(m! m^{r-2} R^{-m}\right) + O_{R_*}\left(m! R_*^{-m}\right).$$

For $r = 1$, the first error term is absent.

Proof. The terms with $\ell < r$ in theorem 13.1 are bounded by a constant times $m^{r-2} R^{-m}$. The term with $\ell = r$ gives the displayed main term. □

Corollary 13.3 (Orbit reduced nearest circle term)**.** *Assume the hypotheses of theorem 13.1. Choose one representative from each orbit of the nearest zeros under multiplication by ω, and call the resulting set $\mathcal{Z}_R^{\mathrm{orb}}$. With $m = Nn$,*

$$P_{N,n}(r) = Nm!\binom{m+r-1}{r-1}\sum_{\tau\in\mathcal{Z}_R^{\mathrm{orb}}}\frac{\tau^{-m}}{(-\tau\mathsf{D}_N'(\tau))^r} + O\left(m!m^{r-2}R^{-m}\right) + O_{R_*}\left(m!R_*^{-m}\right).$$

For $r = 1$, the first error term is absent.

Proof. If $\rho = \omega^k\tau$, then

$$\mathsf{D}_N'(\rho) = \omega^{-k}\mathsf{D}_N'(\tau).$$

Therefore

$$-\rho\mathsf{D}_N'(\rho) = -\tau\mathsf{D}_N'(\tau).$$

Since $m = Nn$, one also has $\rho^{-m} = \tau^{-m}$. Every orbit therefore contributes N equal terms in corollary 13.2. □

Corollary 13.4 (Secant asymptotic)**.** *For $N = 2$ and $r = 1$,*

$$P_{2,n}(1) \sim 2(2n)!\left(\frac{2}{\pi}\right)^{2n+1}.$$

Proof. The nearest zeros of $\cos z$ are $\rho = \pm\pi/2$, both simple. Their contributions in corollary 13.2 are equal. □

14 Zero zeta functions

The assumption $N \geq 2$ remains in force throughout this section. The zero counting theorem gives the exact abscissae of absolute convergence. The exponentially accurate zero locations give more. They compare the zero zeta with a Hurwitz zeta function.

Definition 14.1. Define

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = \sum_{\rho\in\mathcal{Z}(\mathsf{D}_N)}|\rho|^{-s}$$

and

$$\mathfrak{Z}_N^{\mathrm{red}}(s) = \sum_{\lambda\in\Lambda_N}|\lambda|^{-s},$$

with multiplicities. Each term means $e^{-s\log|\rho|}$ or $e^{-s\log|\lambda|}$.

Theorem 14.2 (Abscissae of absolute convergence)**.** *The series $\mathfrak{Z}_N^{\mathrm{abs}}$ converges absolutely and locally uniformly when*

$$\Re s > 1.$$

It fails to converge absolutely when $\Re s \le 1$. The series $\mathfrak{Z}_N^{\mathrm{red}}$ converges absolutely and locally uniformly when

$$\Re s > \frac{1}{N}.$$

It fails to converge absolutely when $\Re s \le 1/N$. Both functions are holomorphic on their open half-planes of absolute convergence.

Proof. By theorem 11.14, the zeros of D_N have linear counting density, while the reduced zeros have counting order $R^{1/N}$. Partial summation gives absolute convergence in the stated open half-planes. The positive terms obtained after taking absolute values diverge on the boundary and to its left. Local uniform convergence follows by the Weierstrass test on compact subsets. □

Theorem 14.3 (Meromorphic continuation of the symmetry-orbit zero zeta)**.** *Put*

$$A_N = \frac{\pi}{\sin(\pi/N)}.$$

There is an entire function Q_N^{red} such that

$$\mathfrak{Z}_N^{\mathrm{red}}(s) = A_N^{-Ns}\zeta\left(Ns, \frac{1}{2}\right) + Q_N^{\mathrm{red}}(s)$$

on $\Re s > 1/N$. The right-hand side gives a meromorphic continuation to $\mathbb{C}$ with one simple pole at $s = 1/N$. Its residue is

$$\frac{\sin(\pi/N)}{N\pi}.$$

There is an entire function Q_N^{abs} such that, initially on $\Re s > 1$,

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = NA_N^{-s}\zeta\left(s, \frac{1}{2}\right) + Q_N^{\mathrm{abs}}(s).$$

Thus $\mathfrak{Z}_N^{\mathrm{abs}}$ continues meromorphically to $\mathbb{C}$ with one simple pole at $s = 1$, of residue

$$\frac{N\sin(\pi/N)}{\pi}.$$

Proof. For $N = 2$, corollary 11.10 and $\mathsf{D}_2(z) = \Delta_2(z^2)$ give

$$\lambda_m = \pi^2\left(m + \frac{1}{2}\right)^2, \qquad m \ge 0.$$

Thus the first formula holds with $Q_2^{\mathrm{red}} = 0$.

For $N \ge 3$, corollary 11.13 writes all but finitely many reduced zeros as

$$\lambda_m = \left(A_N\left(m + \frac{1}{2}\right)\right)^N (1 + \varepsilon_m)$$

where $\lambda_m > 0$, $1 + \varepsilon_m > 0$, and

$$\varepsilon_m = O\left(t_m^{-1}e^{-\gamma t_m}\right).$$

On every compact set of s values,

$$(1+\varepsilon_m)^{-s}-1=O(|\varepsilon_m|)$$

uniformly for large m. The difference

$$\lambda_m^{-s}-A_N^{-Ns}\left(m+\frac{1}{2}\right)^{-Ns}$$

therefore has a normally convergent sum on every compact subset of $\mathbb{C}$. Indeed, the model power has at most polynomial growth in m on a compact set, while $|\varepsilon_m|$ decays exponentially. The finitely many exceptional terms $|\lambda|^{-s}$ and the finitely many omitted model terms are entire functions of s. Their addition to the normally convergent tail defines Q_N^{red} and proves the first formula. Thus that formula is established for every $N\geq 2$.

In either case, the Hurwitz zeta function has a meromorphic continuation with one simple pole at its first argument equal to one and residue one. Composition with Ns gives the stated pole and residue.

For every $N\geq 2$, each nonzero zero λ of Δ_N gives N zeros of D_N, each of modulus $|\lambda|^{1/N}$. Hence

$$\mathfrak{Z}_N^{\mathrm{abs}}(s)=N\mathfrak{Z}_N^{\mathrm{red}}(s/N).$$

Substitution gives the second formula with

$$Q_N^{\mathrm{abs}}(s)=NQ_N^{\mathrm{red}}(s/N),$$

which is entire. □

Corollary 14.4 (Exact zeta models in orders two and four)**.** *For $N=2$,*

$$\mathfrak{Z}_2^{\mathrm{red}}(s)=\pi^{-2s}\zeta\left(2s,\frac{1}{2}\right)$$

and

$$\mathfrak{Z}_2^{\mathrm{abs}}(s)=2\pi^{-s}\zeta\left(s,\frac{1}{2}\right).$$

For $N=4$, put $A_4=\sqrt{2}\pi$. Then

$$\mathfrak{Z}_4^{\mathrm{red}}(s)=A_4^{-4s}\zeta\left(4s,\frac{1}{2}\right)$$

and

$$\mathfrak{Z}_4^{\mathrm{abs}}(s)=4A_4^{-s}\zeta\left(s,\frac{1}{2}\right).$$

In these two cases the entire correction terms in theorem 14.3 vanish identically.

Proof. For $N=2$, the reduced zeros are

$$\pi^2\left(m+\frac{1}{2}\right)^2,\qquad m\geq 0.$$

For $N=4$, example 11.12 gives the reduced zeros

$$A_4^4\left(m+\frac{1}{2}\right)^4,\qquad m\geq 0.$$

Summing their negative powers proves the reduced formulas. The absolute formulas follow from

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = N\mathfrak{Z}_N^{\mathrm{red}}(s/N).$$

□

Definition 14.5. Choose a logarithm for each member of the finite exceptional part of Λ_N. Use the real logarithm for the eventual positive sequence. Define

$$\mathfrak{Z}_N^{\mathrm{orb}}(s) = \sum_{\lambda \in \Lambda_N} e^{-s \operatorname{Log} \lambda}.$$

This series is initially defined on $\Re s > 1/N$.

Corollary 14.6 (Branch-dependent symmetry-orbit continuation)**.** *The branch-dependent orbit zeta $\mathfrak{Z}_N^{\mathrm{orb}}$ continues meromorphically to $\mathbb{C}$. It has the same single pole and residue as $\mathfrak{Z}_N^{\mathrm{red}}$. Changing the logarithms of the finite exceptional zeros changes the continuation by an entire function.*

Proof. The tail consists of positive reduced zeros, so it has the same comparison with the Hurwitz zeta. Only finitely many terms depend on the branch choices. □

Definition 14.7. Let q be a positive integer. Choose logarithms of the simple zeros of D_N with arguments in a fixed bounded interval. Define

$$\mathfrak{Z}_{N,q}^{\mathrm{der}}(s) = \sum_{\substack{\mathsf{D}_N(\rho)=0 \\ \mathsf{D}_N'(\rho)\neq 0}} \frac{e^{-s \operatorname{Log} \rho}}{\mathsf{D}_N'(\rho)^q}.$$

Theorem 14.8 (Derivative-weighted convergence)**.** *If $N \geq 3$ and $q \geq 1$, the series $\mathfrak{Z}_{N,q}^{\mathrm{der}}(s)$ converges locally uniformly for every $s \in \mathbb{C}$. It therefore defines an entire function.*
For $N = 2$, the series converges absolutely and locally uniformly on $\Re s > 1$, where it defines a holomorphic function.

Proof. For $N \geq 3$, theorem 11.7 gives

$$|\mathsf{D}_N'(\rho)|^{-1} \leq Ce^{-c_N|\rho|}$$

for every sufficiently large simple zero. Let L be a compact subset of the s plane. Because the chosen arguments of the zeros lie in a fixed bounded interval, there are constants C_L and M_L such that

$$\sup_{s \in L} |e^{-s \operatorname{Log} \rho}| \leq C_L |\rho|^{M_L}$$

for every sufficiently large zero. Hence the absolute value of the corresponding summand is at most

$$C_L' |\rho|^{M_L} e^{-qc_N|\rho|}.$$

The linear zero counting estimate makes this majorant summable. The Weierstrass test gives absolute local uniform convergence of the series of entire summands, so its sum is entire. The finite exceptional set causes no difficulty.

For $N = 2$, the derivative of the cosine has modulus one at every zero. The assertion then follows from the linear zero spacing. □

Remark 14.9. The derivative weighted sum omits the finitely many zeros that might be multiple. A treatment of those zeros requires the complete principal parts of D_N^{-q} rather than the single derivative weight.

Part V Scope and formal evidence

15 Scope boundaries and further questions

The large-zero theorem settles eventual simplicity but does not classify the finite exceptional set. A remaining question is whether every zero of D_N is simple and every reduced zero of Δ_N is positive. Both statements hold for $N = 2$ and $N = 4$. The argument here proves them only outside a compact set for general N.

The leading correction in theorem 11.8 may admit iteration to a full exponential expansion for τ_m. Such an expansion is an open problem, not a result of this chapter. It could refine the entire correction in the zero-zeta comparison and lead to explicit special values.

The symmetry-orbit zero zeta has a meromorphic continuation with one pole. No functional equation is obtained. A branch dependent sum over the N roots of each reduced zero must specify logarithms on the rays.

The coefficient families in sections 7, 9 and 10 have exact formal parameter-addition laws. Their arithmetic behavior, including odd-prime congruences, dyadic weighting, and p-adic or Amice–Mahler interpolation, belongs to Chapter IX. General bounded-operator semigroup, resolvent, and Feshbach theory belongs to Chapter VI. Chapter III covers general nonlinear operation trees, while abstract completion and occupancy questions belong to Chapter X. None of those theories is inferred from the scalar complex-analytic results proved here.

16 Formalization ledger

The recorded formal verification used Lean 4 v4.34.0-rc2 and Mathlib revision `aeb72f1e9981b6d35b426b16dca7759be3e7ba06` [dMU21, The20]. Its 151 theorem-map rows resolve to 1164 unique declarations in the namespace `Komorebi.PaperIV`: 953 theorem declarations and 211 constructions. The recorded namespace audit scanned 2234 declarations, including 1999 theorem declarations. The recorded build with warnings treated as errors and supplied kernel audit allowed only `propext`, `Classical.choice`, and `Quot.sound` as standard transitive axioms.

Manuscript records	Compiled Chapter IV scope
3.1–3.3	Actual holomorphic root filters, convergent coefficient selection, orthogonal resolution, the exact range–covariance characterization, and the complete pointwise cyclic product law. The formal versions hold as well.
4.1–4.5 and 5.1–5.8	Entire sections, finite exponential formulas, derivative cycles, both Wronskians, and unique entire ODE basis expansions. Both pointwise addition laws and denominator symmetries are checked. The exact ordinary hypergeometric formula follows from the omitted-parameter list, Gamma normalization, and Mathlib's everywhere convergent series.

Manuscript records	Compiled Chapter IV scope (continued)
6.1–6.6	Quotient-algebra power bases and actual exponential-series convergence in the finite-dimensional coefficient topology. Both pointwise section-coordinate identities. Signed-shift powers, all matrix-exponential entries, group laws and trace are checked. The characteristic polynomial $X^N + 1$, its distinct roots, exact spectrum and determinant one are proved.
7.1–7.8	Named reciprocal families, exact parameter recurrences, Bell formula, parameter degree and leading coefficient, and polynomial divisibility. Parameter addition holds with any single fixed holomorphic logarithm on the stated domain. Newton's recurrence gives the literal Hessenberg determinant. All three compact reciprocal-polynomial formulas are checked.
8.1–8.5	The full tangent coefficient series has the exact kappa and factorial normalization, including $N = 1$. Reciprocal differentiation holds on every stated fixed logarithm domain. The actual higher Riccati identity holds at every point where the denominator is nonzero.
9.1–9.10	Arbitrary-parameter residue reciprocals and their analytic germs, polynomial degree and leading term, sparse Appell laws, joint convolution, monicity and covariance. The Hasse and derivative-truncation encodings are proved equal. The named Appell polynomial satisfies both the forward action and the exact $\Phi_{N,j}(\partial_x^N)^r$ inverse formula.
10.1–10.4	Exact formal theta reconstruction, polynomial lift and degree, literal divisibility by u and, for even N, by $u + 1$, and the theta Newton recurrence with its first coefficient.
11.2, 11.4–11.11	Two-sided maximum-modulus bounds prove exact order and type. The indicator limsup formula holds in every direction, including ties. Dominant-exponential errors and zero freeness are uniform on compact sets avoiding ties. Near cancellation and the order-two zero set are checked. For $N \geq 3$, gap and remainder bounds, sector confinement, global exhaustion and multiplicity identities give the full large-zero-string theorem. The real leading correction, quadratic exponential error, signed derivative asymptotic, eventual alternating sign and cubic correction are checked.
11.13–11.15	The reduced series is entire, and $\mathsf{D}_N(z) = \Delta_N(z^N)$ preserves multiplicity away from zero. Finite exceptional sets, eventual positive simple reduced zeros, and their stated displacement are checked, including $N = 2$. Both full disk-counting laws retain all exceptional multiplicities. The full-zero reciprocal sums have exact thresholds $s > 1$ and $s > 1/N$, including boundary divergence.
12.1	For every $N \geq 2$, the complete multiplicity-indexed product converges uniformly on compact sets and equals Δ_N. Jensen and Poisson estimates prove rigidity of the actual quotient from derived growth bounds. No growth certificate remains. The original-variable product for D_N is also checked.
12.2–12.4	The actual diagonal operator is trace class, with a basis-independent trace and determinant given by limits of actual finite compressions. Infinite complete homogeneous coefficients are limits of finite monomial sums. Every normalized section has exact order $1/N$. For $N \geq 2$ its full zero product and zero-moment Bell formulas are proved. The actual complex-power germ at zero has the named residue Taylor coefficients.

Manuscript records	Compiled Chapter IV scope (continued)
12.6–12.10	The normalized logarithm on each zero-free disk has the convergent zero-moment series. Its coefficients equal $\kappa_{N,m}$ and satisfy Newton's recurrence. The Mittag–Leffler tangent expansion converges absolutely and locally uniformly away from poles. Tangent coefficients and the named reciprocal and theta families have their exact zero-moment Bell formulas. The full tangent series is identified with the analytic function.
13.1–13.4	Analytic division constructs each simple-zero principal part. Its leading coefficient is uniquely determined. Removing the principal parts gives a holomorphic remainder and the exact factorial-scaled Cauchy estimate. Both leading-term errors are bounded, with the first absent for $r = 1$. Rotation gives the orbit-reduced main term. The secant formula holds as a ratio limit, with an exponentially vanishing relative error.
14.2–14.4, 14.6	The exact all-zero sums retain analytic multiplicity. Their sharp absolute-convergence domains, boundary divergence and local uniform convergence are checked. Both Hurwitz comparisons, entire corrections, global meromorphic continuations, unique simple poles and exact residues are proved. The order-two/four models have identically zero corrections. Branch choices on the finite exceptional part change the continuation by an entire function and preserve its pole and residue.
14.8	The derivative-weighted series uses every actual simple zero and genuine logarithms with bounded arguments. Absolute and local uniform convergence and an entire sum are proved for $N \geq 3$. For $N = 2$, absolute/local uniform convergence and holomorphy hold on $\Re s > 1$.

The declaration-level theorem map is authoritative for the formal contract. It records full scoped coverage of all 68 theorem-like records: 68 full, zero partial, and zero unmapped. Rows 143–151 close the final fifteen partial records without altering their statements or assumptions. Imported standard fact 11.1 remains imported. Formal identities, normalized analytic germs, and fixed logarithm domain identities retain their respective stated meanings. Finite computations do not change these proof boundaries.

17 Conclusion

Cyclic projection turns the exponential into a finite differential basis, and the signed cycle produces the distinguished denominator D_N. Its reciprocal powers yield Bell and Newton recurrences, an additive parameter law, Appell systems, and theta polynomials. Reduction by $w = z^N$ then connects the reciprocal coefficients to a genus zero product and reduced-zero power sums.

Classical two term localization, specialized to the regular exponent polygon and combined with reflection symmetry, places all sufficiently large zeros on symmetry rays. The displacement, derivative and counting estimates yield the family-specific Hurwitz comparison for the orbit zero zeta.

References

[AAR99] George E. Andrews, Richard Askey, and Ranjan Roy. *Special Functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1999.

[Apo76] Tom M. Apostol. *Introduction to Analytic Number Theory*. Undergraduate Texts in Mathematics. Springer, New York, 1976.

[BM06] Pavel Bleher and Robert Mallison, Jr. Zeros of sections of exponential sums. *International Mathematics Research Notices*, 2006:Article ID 38937, 49 pages, 2006. doi: 10.1155/IMRN/2006/38937.

[Boa54] Ralph P. Boas. *Entire Functions*. Academic Press, New York, 1954.

[Com74] Louis Comtet. *Advanced Combinatorics: The Art of Finite and Infinite Expansions*. D. Reidel Publishing Company, Dordrecht, 1974.

[dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021.

[FS09] Philippe Flajolet and Robert Sedgewick. *Analytic Combinatorics*. Cambridge University Press, Cambridge, 2009.

[Lan31] Rudolph E. Langer. On the zeros of exponential sums and integrals. *Bulletin of the American Mathematical Society*, 37(4):213–239, 1931.

[Lev80] Boris Ya. Levin. *Distribution of Zeros of Entire Functions*, volume 5 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, revised edition, 1980.

[Mor73] Carlos J. Moreno. The zeros of exponential polynomials. I. *Compositio Mathematica*, 26(1):69–78, 1973.

[Mul05] Martin E. Muldoon. Generalized hyperbolic functions, circulant matrices and functional equations. *Linear Algebra and its Applications*, 406:272–284, 2005. doi: 10.1016/j.laa.2005.04.011.

[P.26a] Komorebi P. Projector-resolved seed calculus i: Typed seeds, operator grammars, and exact reconstruction, 2026. Version 1.2.1.

[P.26b] Komorebi P. Projector-resolved seed calculus ii: Polyphase refinement, weighted transfer, and dyadic–odd resolution, 2026. Version 1.2.1.

[P.26c] Komorebi P. Projector-resolved seed calculus iii: Typed edge operators, connection defects, and obstruction packets, 2026. Version 1.0.1.

[Pól20] George Pólya. Geometrisches über die verteilung der nullstellen gewisser ganzer transzendenter funktionen. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften, Mathematisch Physikalische Klasse*, 50:285–290, 1920.

[Rai60] Earl D. Rainville. *Special Functions*. The Macmillan Company, New York, 1960.

[Rom84] Steven Roman. *The Umbral Calculus*, volume 111 of *Pure and Applied Mathematics*. Academic Press, New York, 1984.

[Sch92] Jens Schwaiger. On generalized hyperbolic functions and their characterization by functional equations. *Aequationes Mathematicae*, 43:198–210, 1992. doi: 10.1007/BF01835702.

[The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs*, pages 367–381. Association for Computing Machinery, 2020.

[Tit39] Edward C. Titchmarsh. *The Theory of Functions*. Oxford University Press, Oxford, second edition, 1939.

A Visual atlas

The twenty-six plates retain the worked models of the separate visual guide. The stable labels for Plates IV.1–IV.26 name mathematical mechanisms and point back to the relevant statements. The plates distinguish exact identities, rational certificates, sampled curves, and inherited formal evidence.

Three routes through the atlas

For algebra, read Plates IV.3–IV.16. For zero geometry and its consequences, read Plates IV.17–IV.24. Plate IV.25 follows one coefficient through the complete calculation. Plate IV.26 explains the evidence attached to the models.

A coefficient can become zero data

Chapter IV: Section 2, Section 7, Section 11, Section 12.

This guide follows a concrete analytic family through three descriptions: selected Taylor coefficients, a finite differential system, and the zeros of an entire function. The formulas are linked, but the evidence needed for each description is different.

COEFFICIENT LANES

z^0	z^3	z^6	z^9
z^1	z^4	z^7	z^{10}
z^2	z^5	z^8	z^{11}


$m \equiv j \pmod 3$

RECIPROCAL COEFFICIENTS

1	0	0
$-\frac{1}{2}$	1	0
$\frac{1}{24}$	$-\frac{1}{2}$	1

$1, \quad \frac{1}{2}, \quad \frac{5}{24}, \ldots$

ZERO ORBITS



$\{\rho, i\rho, -\rho, -i\rho\}$

Keep three small models in view

For $N = 2$, the signed denominator is $D_2(z) = \cos z$. Its reciprocal is the familiar secant series. For $N = 3$, a third exponential perturbs the zeros of a dominant cosine. For $N = 4$, an exact factorization gives a complete zero picture:

$$D_4(z) = \cos(z/\sqrt{2})\cosh(z/\sqrt{2}).$$

The examples move between these orders deliberately. Order two makes the coefficient arithmetic transparent. Order three shows a genuine displacement. Order four makes orbit reduction completely explicit.

Three routes through the guide

Route	Pages and question
Algebra first	Plate IV.3 to Plate IV.16: how do selection, signs, reciprocals, and finite differential operators work?
Zeros first	Plate IV.17 to Plate IV.24: where can cancellation happen, and how does it determine coefficients?
One full calculation	Plate IV.25: derive the same number, 69, from denominator coefficients and from zero moments.

Read this as: The pictures carry computable objects. They do not replace the global analytic arguments or extend the inherited Lean contract.

Exact identities and rational certificates are distinguished from sampled curves. All plates are reproducible vector drawings. The source package preserves the original proofread manuscript.

Two variables, several normalizations

Chapter IV: Section 2, Definition 5.7, Definition 7.1, Definition 9.1.

The most important coordinate change is $w = z^N$. A term of degree n in w has degree Nn in z. Its coefficient does not change, but an exponential-generating-function label does.

$$\underbrace{b_{N,n}(r)\,w^n}_{\text{ordinary coefficient}} \quad = \quad \underbrace{P_{N,n}(r)\frac{z^{Nn}}{(Nn)!}}_{\text{block exponential coefficient}} \quad , \qquad P_{N,n}(r) = (Nn)!b_{N,n}(r).$$

Object	Definition and role
$E_{N,j}(z)$	$\displaystyle\sum_{n\geq 0}\frac{z^{Nn+j}}{(Nn+j)!}$ selects one residue class of the exponential.
$D_{N,j}(z)$	The same series with $(-1)^n$. One full derivative cycle contributes a minus sign.
$D_N(z)$	$D_{N,0}(z)$, the denominator whose zeros control the signed reciprocal family.
$\Delta_N(w)$	$\displaystyle\sum_{n\geq 0}\frac{(-1)^n w^n}{(Nn)!}$. Thus $D_N(z) = \Delta_N(z^N)$.
$P_{N,n}(r)$	Block-EGF coefficients of D_N^{-r}. They are polynomials in r.
$\kappa_{N,m}$	$-\log\Delta_N(w) = \displaystyle\sum_{m\geq 1}\kappa_{N,m}w^m/m$.
$T_N(z)$	$-D_N'(z)/D_N(z)$, a function, not the coefficient $T_N(n)$.
$\Phi_{N,j}(w)$	$j!\displaystyle\sum_{n\geq 0}\frac{w^n}{(Nn+j)!}$ removes the forced zero of $E_{N,j}$.
$C_{N,j}(z)$	$1/\Phi_{N,j}(z^N)$. This is generally not $1/D_N(z)$.
Λ_N	Zeros of Δ_N, counted with multiplicity, one per rotational orbit of zeros of D_N.

A normalization test

The coefficient $5/24$ of w^2 in $\Delta_2(w)^{-1}$ is the same term as $5z^4/4!$. It is not $5w^2/2!$. Keeping the two factorials separate prevents errors in Bell formulas, convolution weights, and zero-moment reconstruction.

> **Read this as:** Use z for differentiation and rotational geometry. Use w for reciprocal recurrences, canonical products, and orbit-zero sums.


A root filter is an exact vote

Chapter IV: Theorem 3.1, Proposition 3.2.

For $N = 4$, write $\omega = i$. To extract residue j, evaluate the function at four rotated arguments and attach the phase $i^{-j\ell}$:

$$(\mathcal{B}_{4,j}F)(z) = \frac{1}{4}\sum_{\ell=0}^{3} i^{-j\ell}F(i^{\ell}z).$$

A monomial $a_m z^m$ is multiplied by the finite average of $i^{(m-j)\ell}$. Matching phases all vote 1. Nonmatching phases cancel exactly.



An example that can be checked coefficient by coefficient

Take $F(z) = \sum_{m=0}^{12}(m+1)z^m$. Then

$$\mathcal{B}_{4,2}F = 3z^2 + 7z^6 + 11z^{10}.$$

Applying the same selector again changes nothing. Applying a different selector gives zero. Summing the four selected polynomials returns the original F.

Selection is also a symmetry condition

For the selected polynomial F_2, rotation gives $F_2(iz) = -F_2(z)$. More generally, a residue-j section satisfies $F_j(\omega z) = \omega^j F_j(z)$. The phase label therefore describes both the coefficient addresses and the response to rotation.

> **Read this as:** This is not an approximate Fourier fit. The finite root sum is exactly 0 or 1 on every coefficient.

The checker compares the actual complex average with independent coefficient selection for this polynomial. The drawing shows the four complex phases, not a numerical convergence argument.

Products mix the residue lanes

Chapter IV: Theorem 3.3.

Selection separates a function additively. It does not make multiplication diagonal. A residue-i term multiplied by a residue-j term lands in residue $i + j \pmod N$. For $N = 3$, use

$$F = 1 + 2z + 3z^2 + z^3, \qquad G = 2 + z + z^2.$$

The table displays every product of their three selected sections.

	2	z	z^2
$z^3 + 1$	$2z^3 + 2$	$z^4 + z$	$z^5 + z^2$
$2z$	$4z$	$2z^2$	$2z^3$
$3z^2$	$6z^2$	$3z^3$	$3z^4$

$$\mathcal{B}_{3,0}(FG) = 2 + 7z^3$$

shaded cells: $(0,0), (1,2), (2,1)$ all land in residue 0

The shaded cells contribute to residue zero:

$$2(1 + z^3) + (2z)(z^2) + (3z^2)z = 2 + 7z^3.$$

Keeping only the product of the residue-zero inputs would give $2 + 2z^3$ and lose five units of the z^3 coefficient. The mixed cells are part of the exact result.

The general rule

$$\mathcal{B}_{N,k}(FG) = \sum_{i+j \equiv k \ (N)} (\mathcal{B}_{N,i}F)(\mathcal{B}_{N,j}G).$$

Every pair belongs to exactly one output residue. No term is discarded and no coefficient is estimated. This finite cyclic convolution is the algebra behind the later addition formulas and block-coefficient recurrences.

> **Read this as:** A packet label tells you where an interaction lands, not that interactions with other packets vanish.

All nine table entries and all three output sections are checked as exact polynomial identities. This is the scalar cyclic specialization used in this chapter.

One signed edge determines the cycle

Chapter IV: Theorem 5.3, Theorem 6.4, Corollary 6.5, Corollary 6.6.

Let $J_3e_0 = e_1$, $J_3e_1 = e_2$, and $J_3e_2 = -e_0$. A complete turn changes sign, so $J_3^3 = -I$. The signed sections are the coordinates of $e^{zJ_3}e_0$.

J_3

0	0	-1
1	0	0
0	1	0

e^{zJ_3}

D_0	$-D_2$	$-D_1$
D_1	D_0	$-D_2$
D_2	D_1	D_0

$$J_3^3 = -I$$

$$\mathbf{D}_3(0) = e_0$$

$$\det e^{zJ_3} = 1$$

$$D_0 \mapsto -D_2, \qquad D_1 \mapsto D_0, \qquad D_2 \mapsto D_1$$

Here D_j inside the matrix means $D_{3,j}(z)$. For example,

$$D_{3,0} = 1 - \frac{z^3}{3!} + \frac{z^6}{6!} - \cdots, \qquad D_{3,1} = z - \frac{z^4}{4!} + \cdots.$$

Rows are output coordinates and columns are input coordinates. The upper-triangular minus signs record crossing the signed edge while moving from the column to the row. They are not arbitrary orientation choices.

Two checks connect the matrix and the scalar equation

The initial derivative matrix is the identity. Each initial jet $(y(0), y'(0), y''(0))$ therefore determines a unique linear combination. Also,

$$\frac{1}{3}\operatorname{Tr}(e^{zJ_3}) = D_3(z), \qquad \det(XI - J_3) = X^3 + 1.$$

The roots of -1 appearing later in the exponential sum are already the spectrum of this concrete matrix.

> **Read this as:** The functions, their initial data, the addition law, and the exponent polygon come from one signed shift.

The exact tests check every matrix-entry Taylor coefficient through degree 14 for $N = 1, \dots, 5$. The displayed determinant-one law is for $N \geq 2$. $N = 1$ is a separate boundary case.

Addition is quotient multiplication

Chapter IV: Proposition 5.5, Theorem 6.2.

Work in $\mathbb{C}[\xi]/(\xi^3+1)$. Every element has coordinates in $1, \xi, \xi^2$. The entire exponential has the finite-coordinate form

$$e^{z\xi} = D_{3,0}(z) + D_{3,1}(z)\xi + D_{3,2}(z)\xi^2.$$

The multiplication table makes the carry sign unavoidable.

	1	ξ	ξ^2
1	1	ξ	ξ^2
ξ	ξ	ξ^2	-1
ξ^2	ξ^2	-1	$-\xi$

$$\xi^3 = -1$$

wrap once: change sign

$$\xi^2\xi^2 = -\xi$$

$$D_{3,0}(z+w) = D_{3,0}(z)D_{3,0}(w) - D_{3,1}(z)D_{3,2}(w) - D_{3,2}(z)D_{3,1}(w)$$

Multiply the coordinate forms of $e^{z\xi}$ and $e^{w\xi}$. Their product is $e^{(z+w)\xi}$. The coefficient of 1 receives one contribution without a wrap and two contributions with a wrap. Reading that coefficient gives the displayed addition law.

Compare the unsigned model

In $\mathbb{C}[\xi]/(\xi^3 - 1)$, the same table has no minus signs. Its exponential coordinates are the unsigned sections $E_{3,j}$. Thus the difference between the two differential systems is a single algebraic relation, not a change in how coefficients are grouped. For general N, the carry factor is

$$(-1)^{\lfloor (i+j)/N \rfloor}, \qquad 0 \le i, j < N.$$

Only zero or one wrap is possible in a product of two coordinate basis elements.

> **Read this as:** The twisted addition law is the ordinary exponential group law expressed in a signed cyclic algebra.

Bivariate Taylor identities are checked through total degree ten in every section for $N = 2, 3, 4$. Omitting the carry sign is one of the deliberately rejected controls.

Why a hypergeometric name appears

Chapter IV: Theorem 4.5.

The hypergeometric form is another way to organize the same coefficients. It does not introduce an extra function into the construction. Split a factorial into its residue classes and divide each factor by N:

$$(Nn+j)! = j!N^{Nn}n! \prod_{a\in A_{N,j}} (a)_n.$$

The parameter equal to 1 is omitted from $A_{N,j}$ because the hypergeometric convention already supplies $(1)_n = n!$.

$$N=3,\ j=1,\ n=2: \quad 7! = 1!\,3^6\,2!\,(2/3)_2(4/3)_2$$

$\frac{2}{3}$	$\frac{5}{3}$
1	2
$\frac{4}{3}$	$\frac{7}{3}$

$(2/3)_2$

$(1)_2 = 2!$

$(4/3)_2$

$$A_{3,1} = \{2/3, 4/3\}$$

$$E_{3,1}(z) = z\,{}_0F_2\left(;\frac{2}{3},\frac{4}{3};\frac{z^3}{27}\right)$$

The omitted unit parameter is already the hypergeometric $n!$.

Check the displayed case

For $N=3, j=1, n=2$, the six factors after the initial $1!$ are $2,3,4,5,6,7$. Dividing by three and grouping them gives the three rows of the plate. The middle row supplies $2!$. The others supply $(2/3)_2$ and $(4/3)_2$. The right-hand side is exactly 5040. Consequently,

$$E_{3,1}(z) = z\left(1+\frac{z^3}{4!}+\frac{z^6}{7!}+\cdots\right).$$

The prefactor z retains the forced zero at the origin. The hypergeometric argument $z^3/27$ retains the scaling N^N.

> **Read this as:** Parameters, the omitted unit parameter, the prefactor, and the argument scaling all come from factorial bookkeeping.

Five choices of (N, j) are checked against the exact factorial identity for $n = 0, \dots, 7$. These finite checks support the example. The general analytic identification belongs to the inherited manuscript.


Four zeros become one orbit zero

Chapter IV: Definition 5.7, Example 11.12, Corollary 11.13.

For $N = 4$ the factorization is exact:

$$D_4(z) = \cos(z/\sqrt{2})\cosh(z/\sqrt{2}), \qquad D_4(z) = \Delta_4(z^4).$$

Set $A_4 = \sqrt{2}\pi$. All the zeros are $A_4(m + 1/2)i^k$, with $m \geq 0$ and $0 \leq k < 4$. Each zero is simple.



The drawing uses $u = z/A_4$ and $v = w/A_4^4$, so its marked coordinates are rational. The first four-point orbit has fourth power $1/16$. The second has fourth power $81/16$.

Multiplicity is not multiplied in the reduced function

At a nonzero point, $z \mapsto z^4$ has nonzero derivative. It preserves the local multiplicity of a zero. Four distinct simple zeros of D_4 therefore correspond to one simple zero of Δ_4, not one zero of multiplicity four. The reduced zeros are

$$\lambda_m = A_4^4(m + 1/2)^4 = 4\pi^4(m + 1/2)^4.$$

This is why an orbit factor in the z plane is $1 - z^4/\lambda_m$. It is the product of the four corresponding linear factors, with their phases included.

> **Read this as:** Orbit reduction removes repetition. It changes growth and counting exponents, while preserving each nonzero zero's local multiplicity.

The quartic factorization, its differential equation, and the displayed orbit powers are checked independently. Here the complete zero set is known exactly. A finite numerical root search is not being used to establish it.

A reciprocal is a triangular solve

Chapter IV: Theorem 7.3, Theorem 7.6.

Start in the reduced variable with

$$\Delta_2(w) = 1 - \frac{w}{2} + \frac{w^2}{24} - \frac{w^3}{720} + \cdots.$$

Write $1/\Delta_2(w) = \sum b_n w^n$. Matching the product with 1 gives a unit lower-triangular system. At each row there is exactly one new unknown.

denominator convolution				b		unit series
1	0	0	0	1	$=$	1
$-\frac{1}{2}$	1	0	0	$\frac{1}{2}$		0
$\frac{1}{24}$	$-\frac{1}{2}$	1	0	$\frac{5}{24}$		0
$-\frac{1}{720}$	$\frac{1}{24}$	$-\frac{1}{2}$	1	$\frac{61}{720}$		0

$$b_1 = \tfrac{1}{2}$$

$$b_2 = \tfrac{1}{4} - \tfrac{1}{24}$$

$$b_3 = \tfrac{61}{720}$$

$$P_{2,n}(1) = (2n)!b_n : \qquad 1,\ 1,\ 5,\ 61$$

The first three equations are

$$b_1 - \tfrac{1}{2} = 0, \qquad b_2 - \tfrac{1}{2}b_1 + \tfrac{1}{24} = 0, \qquad b_3 - \tfrac{1}{2}b_2 + \tfrac{1}{24}b_1 - \tfrac{1}{720} = 0.$$

They yield $b_1 = 1/2$, $b_2 = 5/24$, and $b_3 = 61/720$. Multiplication by $(2n)!$ produces the secant coefficients $1, 5, 61$.

Why this is an exact finite calculation

To compute the coefficient of w^n, only denominator coefficients through w^n are needed. Later terms of the entire series cannot affect it. A finite series inverse is therefore exact at its declared truncation order, even though evaluating that truncated series is only an approximation to the full analytic function away from the origin.

> **Read this as:** A unit constant term makes coefficient recovery triangular. It does not by itself specify the radius of analytic convergence.

The checks compare an independent binomial-series expansion, the parameter recurrence, and the logarithmic Newton recurrence through block index six. The table uses ordinary coefficients, with factorial conversion shown separately.


The reciprocal order is a polynomial

Chapter IV: Theorem 7.2, Corollary 7.7.

Let r vary instead of fixing the reciprocal power. Since

$$\Delta_N(w)^{-r} = \exp\left(r \sum_{m \geq 1} \kappa_{N,m} \frac{w^m}{m} \right),$$

the coefficient of w^n is a polynomial of degree n in r. Bell polynomials collect the possible ways logarithmic terms contribute total degree n.



$$r^3\kappa_1^3 \qquad 3r^2\kappa_1\kappa_2 \qquad 2r\kappa_3$$

$$15r^3 \qquad 30r^2 \qquad 16r$$

$$P_{2,3}(r) = \tfrac{6!}{3!}\, B_3(r\kappa_1, r\kappa_2, 2r\kappa_3)$$

For $N = 2$, the first three logarithmic coefficients are

$$\kappa_1 = \frac{1}{2}, \qquad \kappa_2 = \frac{1}{6}, \qquad \kappa_3 = \frac{1}{15}.$$

The third Bell polynomial is $B_3(x_1, x_2, x_3) = x_1^3 + 3x_1x_2 + x_3$. Substituting $x_1 = r\kappa_1$, $x_2 = r\kappa_2$, $x_3 = 2r\kappa_3$ and multiplying by $6!/3!$ gives

$$P_{2,3}(r) = 15r^3 + 30r^2 + 16r.$$

At $r = 1$ this is 61. At $r = 2$ it is 272. At $r = 0$ it vanishes, as every positive-index reciprocal coefficient must.

The leading term has a simple source

Only n copies of the linear logarithmic term can contribute $r^n w^n$. Therefore

$$[r^n]P_{N,n}(r) = \frac{(Nn)!}{n!(N!)^n}.$$

> **Read this as:** The polynomial parameter records how many logarithmic pieces are combined. It is not a numerical interpolation in the reciprocal order.

The partition plate is a finite combinatorial expansion. The exact tests verify the leading coefficient and parameter polynomial through block index six for five orders.

Two recurrences reach the same answer

Chapter IV: Theorem 7.3, Theorem 7.4, Theorem 7.6.

One recurrence uses the denominator coefficients. Another uses its logarithm. With $b_n(r) = P_{N,n}(r)/(Nn)!$, the shorter logarithmic form is

$$nb_n(r) = r\sum_{k=1}^{n} \kappa_{N,k} b_{n-k}(r), \qquad b_0 = 1.$$


The negative superdiagonal in the Newton determinant is essential.

$n!b_n(r)$ for $n = 3$

$\frac{r}{2}$	-1	0
$\frac{r}{6}$	$\frac{r}{2}$	-2
$\frac{r}{15}$	$\frac{r}{6}$	$\frac{r}{2}$

$\det /6 = P_{2,3}(r)/720$

$P_{2,3}(2) = 61 + 75 + 75 + 61 = 272$



block weights: $1, 15, 15, 1$

The additive parameter law has block weights

Using the same logarithm, $D_N^{-r} D_N^{-s} = D_N^{-(r+s)}$. Thus

$$P_{N,n}(r+s) = \sum_{q=0}^{n} \binom{Nn}{Nq} P_{N,q}(r) P_{N,n-q}(s).$$

For $N = 2, n = 3, r = s = 1$, the four weights are $1, 15, 15, 1$. The two middle terms are $15 \cdot 1 \cdot 5$ and $15 \cdot 5 \cdot 1$. Together with the endpoints, they give 272. Using $\binom{n}{q}$ instead would be incorrect for these coefficients. That ordinary binomial law belongs to the separately normalized sequence $n!P_{N,n}/(Nn)!$.

> **Read this as:** Agreement of independent recurrences is a useful check precisely because their bookkeeping is different. Neither permits silently changing the factorial convention.

The checker compares both recurrence routes and the determinant, then checks the parameter-addition identity symbolically. A deliberately incorrect ordinary-binomial substitution is rejected.

Compress the ODE into a tangent

Chapter IV: Proposition 8.2, Corollary 8.4, Theorem 8.5.

The transfer tangent is $T_N = -D'_N/D_N$. It compresses the linear differential equation into a nonlinear scalar identity. This division is allowed only on the zero-free set of the denominator.

LINEAR EQUATION	LOGARITHMIC DERIVATIVE
$D_2'' = -D_2$	$T_2' = 1 + T_2^2$
$D_3''' = -D_3$	$T_3'' - 3T_3T_3' + T_3^3 = 1$

$$T_N = -D'_N/D_N \quad \text{only where } D_N \neq 0$$

$$T_4(z) = \frac{z^3}{3!} + 34\frac{z^7}{7!} + \cdots$$

The coefficients retain the reduced logarithmic information:

$$T_N(z) = N\sum_{m\geq 1} \kappa_{N,m} z^{Nm-1}, \qquad T_N(n) = N(Nn+N-1)!\kappa_{N,n+1}.$$

For order four, $\kappa_{4,1} = 1/24$ and $\kappa_{4,2} = 17/10080$. These give $T_4(0) = 1$ and $T_4(1) = 34$, explaining the last line of the plate.

A nonlinear equation with a precise origin

If $D = e^g$ locally, then $g' = -T$. Differentiating e^g three times gives $D'''/D = -T'' + 3TT' - T^3$. Setting $D'''/D = -1$ yields the cubic Riccati identity shown above. No additional nonlinear model has been assumed. The reciprocal family also satisfies $(D_N^{-r})' = rT_N D_N^{-r}$ on a fixed-log domain. Behavior at zeros of D_N depends on the reciprocal order and the branch conventions stated on Plate IV.13.

> **Read this as:** The nonlinear tangent equation is exact where division is legal. It is not a global regular differential equation across the zeros.


The tests verify the tangent coefficient recurrence, the exact cubic ODE, and the symbolic Riccati reduction after clearing denominators. The order-one tangent is the separate constant function $T_1 = 1$.

A complex power needs a domain

Chapter IV: Section 2, Definition 7.1, Corollary 8.4.

A formal unit series has a formal logarithm at the origin. To interpret that germ analytically, choose a simply connected zero-free region containing 0 and normalize the logarithm to vanish there.

$$D_N(z)^{-r} = \exp(-rL(z)), \qquad e^{L(z)} = D_N(z), \quad L(0) = 0.$$



$$D(z) \sim D'(\rho)(z - \rho)$$

$$\log D \longmapsto \log D + 2\pi i$$

$$D^{-r} \longmapsto e^{-2\pi i r} D^{-r}$$

$$r = 1/2 : \ -1 \qquad r \in \mathbb{Z} : \ 1$$

For $D_2 = \cos z$, the open disk $|z| < \pi/2$ is an admissible domain. A small loop around the simple zero $\rho = \pi/2$ is not contained in that disk. Analytic continuation around the zero changes the local logarithm by $2\pi i$, and the power by $e^{-2\pi i r}$.

Three different kinds of statement

A *formal* identity compares coefficients at the origin. A *holomorphic* identity for complex powers uses a common specified logarithm. A positive *integer* reciprocal power is globally meromorphic and needs no branch choice. These scopes cannot be exchanged without a further argument. For example, $D^{-r}D^{-s} = D^{-(r+s)}$ is immediate using one fixed logarithm. Combining values built from unrelated branches can introduce additional phase factors.

> **Read this as:** The branch choice is mathematical data. It is not a plotting convention that can be removed by drawing a cleaner curve.

The domain picture is schematic. The phase change comes from the local simple-zero model and winding number, not from a sampled loop. Later nearest-pole asymptotics retain positive integral reciprocal order.

Three similar reciprocals differ

Chapter IV: Definition 9.1, Theorem 9.2, Example 9.11.

The signed denominator and the normalized unsigned sections produce different reciprocal families. In order two,

$$D_2^{-1} = \sec z, \qquad C_{2,0} = \operatorname{sech} z, \qquad C_{2,1} = z/\sinh z.$$

Their first block-EGF coefficients make the difference visible.

function	1	$z^2/2!$	$z^4/4!$	$z^6/6!$
$\sec z$	1	1	5	61
$\operatorname{sech} z$	1	-1	5	-61
$z/\sinh z$	1	$-\frac{1}{3}$	$\frac{7}{15}$	$-\frac{31}{21}$

$$E_{2,1}(z) = \sinh z = z\Phi_{2,1}(z^2), \qquad C_{2,1}(0) = 1$$

Remove the forced zero before taking a reciprocal

The section $E_{N,j}$ begins with $z^j/j!$. Define

$$E_{N,j}(z) = \frac{z^j}{j!}\Phi_{N,j}(z^N), \qquad \Phi_{N,j}(0) = 1.$$

Then $C_{N,j}(z) = 1/\Phi_{N,j}(z^N)$ has value 1 at the origin. For $j > 0$, the notation $(z^j/j!)/E_{N,j}(z)$ means this removable continuation. It does not mean evaluating an undefined $0/0$ expression. For $N = 2, j = 1$, the first nontrivial coefficient is $-1/3$ in front of $z^2/2!$, equivalently $-z^2/6$. Dropping the numerator z would instead produce the pole of $1/\sinh z$.

> **Read this as:** A minus sign in the denominator series and removal of an initial zero are different operations. Their reciprocal coefficients should not be interchanged.

All entries in the table are exact rationals computed from independent triangular inverses. The table explicitly uses block-EGF columns. Ordinary coefficients are obtained by dividing by the displayed factorials.


An infinite symbol acts finitely

Chapter IV: Theorem 9.6, Theorem 9.8, Proposition 9.9.

For $C_{2,0}(z) = \operatorname{sech} z$, define $E_m(x)$ by $\operatorname{sech} z\, e^{xz} = \sum E_m(x) z^m/m!$. The formal differential-operator formula is

$$E_m(x) = \operatorname{sech}(\partial_x) x^m.$$

Although the symbol is an infinite series, derivatives above degree m vanish.



For $m = 4$, only three terms survive:

$$\left(1 - \frac{\partial_x^2}{2!} + 5\frac{\partial_x^4}{4!}\right) x^4 = x^4 - 6x^2 + 5.$$

The four real roots of this polynomial are $\pm 1, \pm\sqrt{5}$. They are not the zeros of the original analytic denominator. The plot distinguishes the input polynomial and its transformed output.

An inverse operation, not a truncation error

Applying $\cosh(\partial_x)$ returns x^4 exactly. The general inverse uses the reciprocal formal symbol. The lowering and translation identities are

$$E_m' = m E_{m-1}, \qquad E_m(x+y) = \sum_{q=0}^{m} \binom{m}{q} E_q(x) y^{m-q}.$$

Both follow from the generating function. No operator-series convergence assumption on an infinite-dimensional function space is needed for this polynomial calculation.

> **Read this as:** Formal substitution into the derivative is safe here because every polynomial sees only finitely many derivatives. That is a domain statement.

The polynomial, its inverse image, and the Appell identities are checked symbolically through degree eight in four normalized-section families. The displayed curves are samples of exact polynomials.


Scale and order enter differently

Chapter IV: Theorem 10.2, Proposition 10.3, Proposition 10.4.

Let $H_N(z) = E_{N,0}(z)$. The theta family uses the same parameter u twice, but in two different roles:

$$H_N(uz)^{-1} H_N(z)^{-u} = \sum_{n \geq 0} \Theta_{N,n}(u) \frac{z^{Nn}}{(Nn)!}.$$

Argument scaling contributes u^{Nm} at logarithmic block m. Reciprocal order contributes u.



Their combined logarithm has coefficient

$$(-1)^m \kappa_{N,m} \frac{u^{Nm} + u}{m}.$$

At the first block this gives $\Theta_{N,1}(u) = -(u^N + u)$. Every positive-index theta polynomial vanishes at $u = 0$. For even N, $H_N(-z) = H_N(z)$, so the defining product is also 1 at $u = -1$.

The odd-order boundary is visible

For $N = 3$, $\Theta_{3,1}(-1) = 2$, not zero. The factor $u + 1$ is not a uniform law across orders. In contrast,

$$\Theta_{2,2}(u) = u(u+1)(5u^2 + u + 2).$$

The two guaranteed factors follow from the whole generating function, not from guessing the first few polynomial roots.

> **Read this as:** A common parameter does not make argument scaling and reciprocal order interchangeable. Their powers must be tracked separately.

The exact checker compares theta convolution with its logarithmic Newton recurrence through block index four. The odd-order factor claim is explicitly rejected as an invalid control.


Growth comes from a polygon

Chapter IV: Definition 11.3, Theorem 11.4.

Write the signed denominator as a finite exponential sum:

$$D_N(z) = \frac{1}{N} \sum_{\eta^N=-1} e^{\eta z}.$$

For $z = re^{i\theta}$, the size of one exponential is $e^{r\Re(\eta e^{i\theta})}$. The maximal projection determines the directional exponential growth.



The indicator is

$$h_N(\theta) = \max_{0\le k<N} \cos\left(\theta + \frac{(2k+1)\pi}{N}\right).$$

Away from a tie direction, one exponential dominates all the others by a uniform positive projection gap on compact angular sets. For sufficiently large radius, smaller terms cannot cancel it. Those sectors are eventually zero-free.

Why zeros remain possible on a growing ray

At a tie, two adjacent exponents have the same real projection. Their complex phases can be opposite. Their sum can therefore vanish even though each term is exponentially large. The minima in the order-four indicator plot occur precisely at the tie directions $\theta = k\pi/2$. The indicator uses a limsup in radius. It does not assert that $r^{-1}\log|D_N(re^{i\theta})|$ has an ordinary limit through the zeros on a tie ray.

> **Read this as:** The polygon predicts where large cancellation is possible. It does not, by itself, enumerate zeros or prove their simplicity.

The order-four polygon and support-function formula are explicit. The curve is a sampled display of that formula. Eventual localization and uniqueness require the separate zero-analysis arguments.

Two large terms nearly cancel


Chapter IV: Lemma 11.5, Theorem 11.7, Corollary 11.9.

For real t, order three has the exact formula

$$D_3(t) = \frac{e^{-t} + 2e^{t/2}\cos(\sqrt{3}t/2)}{3}.$$

Divide by the nonzero factor $2e^{t/2}/3$. The resulting equation separates the dominant cosine from the small third exponential.

$$\tfrac{3}{2}e^{-t/2}D_3(t) = \cos(\sqrt{3}t/2) + \tfrac{1}{2}e^{-3t/2}$$



solid: full expression; dashed: two-term model; circles: model zeros

Ignoring the last term gives model zeros

$$t_m = \frac{2\pi}{\sqrt{3}}(m + 1/2).$$

At t_m, the full normalized expression equals $\frac{1}{2}e^{-3t_m/2} > 0$, so t_m is not an exact zero. Linearizing the cosine at that point predicts

$$\delta_m^{(1)} = \frac{(-1)^m}{\sqrt{3}}e^{-3t_m/2}.$$

The alternating slope explains why the true zero is to the right of the first model zero, then to the left of the second, and so on in the eventual regime.

Symmetry needs uniqueness

Real coefficients alone allow nonreal conjugate pairs. The manuscript first proves that a near-model disk contains exactly one zero. Conjugation preserves that disk, so the unique zero must equal its conjugate. This additional uniqueness step, not the symmetry alone, places it exactly on the ray.

> **Read this as:** The correction is caused by a real, nonzero remainder. A good cosine approximation does not make its model zeros exact.

The curves are sampled analytic expressions. The next page independently certifies four narrow real-zero intervals rather than trusting the plotted crossing or the floating-point root finder.

A root estimate becomes a certificate

Chapter IV: Theorem 11.7, Theorem 11.8, Corollary 11.9.

The root finder is used only to suggest intervals. Each displayed interval is accepted by exact rational arithmetic: opposite endpoint signs prove existence, and an interval bound for D_3' excludes zero throughout the interval, proving uniqueness inside it.

	certified lower endpoint	$\tau_m - t_m$	$\delta_m/\delta_m^{(1)} - 1$
$m = 0$	1.84981279919014347629	$+3.60134 \times 10^{-2}$	-5.243×10^{-2}
$m = 1$	5.44123335502366335540	-1.64738×10^{-4}	$+2.471 \times 10^{-4}$
$m = 2$	9.06899753487157765805	$+7.13700 \times 10^{-7}$	-1.071×10^{-6}
$m = 3$	12.69659554654675729440	-3.09277×10^{-9}	$+4.639 \times 10^{-9}$

Each upper endpoint is the displayed lower endpoint plus 10^{-20}.

Root intervals: exact rational certificates. Displacement columns: numerical diagnostics.

How the certificate works

Evaluate the Taylor polynomial

$$S_{90}(t) = \sum_{n=0}^{90} \frac{(-1)^n t^{3n}}{(3n)!}$$

at the two rational endpoints. From the next term onward, magnitudes decrease, because their successive ratio is

$$\frac{t^3}{(3n+1)(3n+2)(3n+3)} < 1$$

there. The alternating-series remainder is bounded by the first omitted term. A separate outward interval evaluation of the differentiated polynomial, plus its tail bound, certifies the derivative sign. The first bracket is

$$1.84981279919014347629 < \tau_0 < 1.84981279919014347630.$$

Its displacement differs from the leading asymptotic prediction by about five percent. The later examples have much smaller relative discrepancies, as the plate records.

> **Read this as:** A small residual is not a root certificate. Endpoint signs plus a derivative bound give a checkable local existence-and-uniqueness statement.


Four certified intervals do not establish that no other real or complex zeros exist. The displacement columns are high-precision diagnostics, not a finite verification of a global big-O theorem. All rational enclosures are included in the checks file.

Counting changes under the power map

Chapter IV: Example 11.12, Theorem 11.14.

Order four gives a case with no asymptotic displacement at all. In normalized coordinates the positive radial locations are $u = m + 1/2$, while the reduced-zero locations are $v = (m + 1/2)^4$.

$$u = |z|/(\sqrt{2}\pi)$$



$$v = |w|/(4\pi^4)$$



$$n_{D_4}(\sqrt{2}\pi u) = 4\lfloor u + 1/2 \rfloor, \quad n_{\Delta_4}(4\pi^4 v) = \lfloor v^{1/4} + 1/2 \rfloor$$

The two rows describe the same zero orbits. The upper row counts four zeros each time a new radius is reached. The lower row counts one reduced zero each time its fourth-power location is reached. Both formulas include zeros on the boundary of the disk.

Separate radius conversion from orbit multiplicity

In unscaled variables the general relation is

$$n_{D_N}(R^{1/N}) = N n_{\Delta_N}(R).$$

There are two operations here: replace the radius by its Nth root and divide the count by N. Forgetting either operation changes the leading constant. For this family the asymptotics are

$$n_{D_N}(R) = \frac{N \sin(\pi/N)}{\pi} R + O(1), \qquad n_{\Delta_N}(R) = \frac{\sin(\pi/N)}{\pi} R^{1/N} + O(1).$$

The different powers agree with the entire-function orders 1 and $1/N$.

> **Read this as:** The reduced function has fewer repeated zeros and a different radius scale. Its counting law is not obtained by dividing a same-radius count alone.

The exact order-four staircase formulas are tested at rational radii including jump points. The large-zero theorem for general N permits finitely many exceptions. These exact order-four identities do not justify discarding that boundary elsewhere.

A product and an operator share zeros


Chapter IV: Theorem 12.1, Theorem 12.2.

For $N \geq 2$, the reduced entire function has order $1/N < 1$. Its multiplicity-sensitive genus-zero product is

$$\Delta_N(w) = \prod_{\lambda \in \Lambda_N} (1 - w/\lambda).$$

Putting $K_N e_\lambda = \lambda^{-1} e_\lambda$ on $\ell^2(\Lambda_N)$ gives the corresponding diagonal Fredholm determinant. Repeated zeros require repeated diagonal entries.

$$\Delta_2(w) = \cos\sqrt{w}, \qquad P_L(w) = \textstyle\prod_{m=0}^{L-1}(1 - w/\lambda_m)$$



thick: entire function; dotted $L = 1$; dashed $L = 3$; dash-dot $L = 10$

Finite compression is an approximation

For $N = 2$, $\lambda_m = \pi^2(m + 1/2)^2$. The finite determinant P_L is exactly the displayed finite product, but it is not exactly Δ_2. For $0 \leq w \leq 1$ and $L \geq 1$, an elementary tail estimate gives

$$0 \leq 1 - \frac{\Delta_2(w)}{P_L(w)} \leq w \sum_{m \geq L} \lambda_m^{-1} \leq \frac{1}{9(L - 1/2)}.$$

The last bound uses $\pi > 3$ and integral comparison. Thus truncation has an explicit error mechanism.

> **Read this as:** The infinite product is justified by the analytic theorem. Finite determinants provide controlled approximations, not substitutes for that theorem.

The exact finite determinant test includes a repeated eigenvalue and rejects deduplication. The curves use known cosine zeros. Order one is excluded: $\Delta_1(w) = e^{-w}$ has no zeros, so its empty product would be 1, not the function.


Finite coefficients give infinite moments

Chapter IV: Theorem 12.6, Corollary 12.7, Corollary 12.9, Corollary 12.10.

Taking the logarithm of the canonical product identifies

$$\kappa_{N,q} = \sigma_{N,q} := \sum_{\lambda \in \Lambda_N} \lambda^{-q} = \operatorname{Tr}(K_N^q), \qquad q \geq 1.$$

The same exact numbers can be computed from finitely many denominator coefficients by Newton's recurrence. No numerical zero list is needed for that calculation.



five finite contributions plus the remaining tail; higher moments emphasize the nearest zero

For order two,

$$\sigma_{2,1} = \frac{1}{2}, \quad \sigma_{2,2} = \frac{1}{6}, \quad \sigma_{2,3} = \frac{1}{15}, \quad \sigma_{2,4} = \frac{17}{630}.$$

The plate shows how successive inverse powers emphasize the nearest zero. Its shaded pieces are contributions from the first five positive reduced zeros, followed by the remaining tail. The total in each row is the exact rational moment.

The tangent is a sum of poles

Logarithmic differentiation also gives

$$T_N(z) = N z^{N-1} \sum_{\lambda \in \Lambda_N} \frac{1}{\lambda - z^N}$$

away from poles. Summability of $|\lambda|^{-1}$ permits the locally uniform partial-fraction expansion without added regularizing terms.

> **Read this as:** A global analytic product theorem connects the infinite zero sum to a finite algebraic recurrence. The recurrence alone does not prove the product theorem.

Displayed moments are checked against formal logarithm coefficients and exact order-two/order-four zeta values. Bar widths are numerical shares. General power sums use λ^{-q}, not $|\lambda|^{-q}$ when exceptional nonpositive zeros are present.

Nearest poles set coefficient growth


Chapter IV: Theorem 13.1, Corollary 13.2, Corollary 13.3, Corollary 13.4.

For $1/\cos z$, the nearest poles are $\rho = \pm\pi/2$. They contribute equally to even-degree coefficients. Including both poles gives

$$P_{2,n}(1) \sim 2(2n)! \left(\frac{2}{\pi}\right)^{2n+1}.$$

The factor two is orbit multiplicity, not an adjustable fitted constant.

$$\log_{10}(1 - R_n), \quad R_n = \frac{P_{2,n}(1)}{2(2n)!(2/\pi)^{2n+1}}$$



Each ratio is enclosed with a rational Machin-formula bound for π.

A finite quantitative check

For $n = 1, \ldots, 8$, the exact integer coefficients and a rational enclosure for π bound the ratio R_n strictly between zero and one. In this tested range, $1 - R_n < 3^{-(2n+1)}$. The plot displays the logarithm of the enclosed error, not an independently asserted all-n bound.

Higher reciprocal powers need more Laurent data

For positive integral order r, a simple denominator zero becomes a pole of order r. The highest Laurent term contributes a factor $\binom{m+r-1}{r-1}$, but lower Laurent terms still contribute on the same exponential scale. Subtracting *all* principal parts is what leaves a holomorphic remainder controlled by the next zero radius.


> **Read this as:** Nearest-zero geometry explains factorial-times-exponential growth. Multiplicity, reciprocal order, and lower principal parts determine its precise normalization.

The ratio certificates use a rational Machin-formula enclosure for π. The inherited general theorem retains simple nearest zeros, a gap to the next modulus, and positive integral reciprocal order. The plot does not remove those hypotheses.

Convergence is not continuation

Chapter IV: Theorem 14.2, Theorem 14.3, Corollary 14.4.

Reduced zeros grow like m^N. Their absolute zero-zeta series therefore has half-plane of absolute convergence $\Re s > 1/N$. The absolute zero zeta in the original variable instead requires $\Re s > 1$.



For $N = 2$ and $N = 4$, the lattices are exact:

$$\mathfrak{Z}_2^{\mathrm{red}}(s) = \pi^{-2s}\zeta(2s, 1/2), \qquad \mathfrak{Z}_4^{\mathrm{red}}(s) = (\sqrt{2}\pi)^{-4s}\zeta(4s, 1/2).$$

Each identity first holds where its defining series converges. The right side supplies its meromorphic continuation. There is a single pole, at $1/2$ or $1/4$, with residue $1/(2\pi)$ or $1/(4\sqrt{2}\pi)$ respectively.

The factor N matters twice

For every order in this chapter,

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = N\mathfrak{Z}_N^{\mathrm{red}}(s/N).$$

For general N, exponentially accurate zero locations leave an entire correction to the Hurwitz model. A finite exceptional zero set also contributes entire terms. This reasoning establishes continuation, not a zeta functional equation.

> **Read this as:** An analytic continuation can exist at points where the original infinite sum diverges. It must not be evaluated there by naively summing that series.

The shaded regions show convergence, and the open circles mark poles of the continued functions. They are not zeros of the zeta functions. Branch-dependent orbit zetas and derivative-weighted zetas retain their separately stated conventions.

One closed calculation: the number 69

Chapter IV: Corollary 7.7, Example 11.12, Theorem 12.6.

Use $N = 4$ throughout this page. Its denominator, its exact zero lattice, and the reciprocal recurrences provide two routes to the same coefficient. All symbols in the plate refer to this one order.

denominator	$1 - w/24 + w^2/40320 + \cdots$
negative logarithm	$w/24 + (17/20160)w^2 + \cdots$
zero moments	$\sigma_1 = 1/24, \quad \sigma_2 = 17/10080$
reciprocal	$1 + w/24 + 69w^2/40320 + \cdots$
tangent	$z^3/3! + 34z^7/7! + \cdots$

$$69 = 8!\,\tfrac{\sigma_1^2+\sigma_2}{2}, \qquad 34 = 4 \cdot 7!\,\sigma_2$$

Route one: invert the denominator

The w^2 coefficient of Δ_4^{-1} is

$$\frac{1}{24^2} - \frac{1}{8!} = \frac{69}{8!}.$$

Thus $P_{4,2}(1) = 69$. Keeping r variable gives $P_{4,2}(r) = 35r^2 + 34r$.

Route two: use the zero moments

The exact reduced zeros are $4\pi^4(m+1/2)^4$. Their first two power sums are $\sigma_1 = 1/24$ and $\sigma_2 = 17/10080$. The Bell/Newton identity gives the same reciprocal coefficient $(\sigma_1^2 + \sigma_2)/2$. The tangent independently reads off the second moment as $4 \cdot 7!\sigma_2 = 34$. This links coefficients, geometry, and derivatives without a fitted parameter or an inferred zero list.

> **Read this as:** The content is the agreement of precisely normalized descriptions. A matching small example illustrates the general bridges. It is not a proof of all their analytic hypotheses.

Every rational value and polynomial identity on this page is checked exactly. The infinite zero-sum interpretation uses the inherited canonical-product theorem and the exact order-four zero formula.

What is evidence, and for what?

Chapter IV: Section 15, Section 16.

These twenty-six worked plates connect explicit mathematical models with the main text and its formal companion. Each evidence record states the scope of the model, computation, or formal result.

Evidence layer	What it supports
Formal record	The sources, theorem map, dependency pins, saved axiom reports, and mathematical review retain their own scope. The 68-result contract and its replay using pinned sources are described in the Lean guide.
Exact and symbolic suite	The recorded isolated execution passed all 147 finite, symbolic, and enclosure checks and rejected five deliberately false controls. The unchanged checker and models were run with SymPy 1.14.0 and mpmath 1.3.0.
Certified zero brackets	Four rational intervals of width 10^{-20} establish local existence and uniqueness only. The full suite reproduced all four rational brackets, opposite endpoint signs, and derivative exclusion throughout each interval. These are local certificates, not a global zero census.
Numerical visualization	Sampled analytic curves, finite products, moment shares, and displacement diagnostics have the roles stated on their pages. They are not exact global zero censuses.
Figure provenance	Regeneration from the models reproduced all twenty-four editable TikZ files. The mathematical labels and objects are preserved.
Editorial preservation	The original worked formulas, model code, checker, and vector figure sources are preserved. Locators refer to the main text.

Boundaries that remain in force

The general family has eventual positivity and simplicity of its reduced zeros, not an assertion that every reduced zero is positive and simple. This chapter leaves universal simplicity, positivity of every reduced zero, full displacement expansions, and a zeta functional equation open. The exact order-two and order-four models do not settle those questions for every order. Imported standard fact 11.1 stays outside the 68-result contract. Fixed logarithms, zero-free domains, multiplicities, and the hypotheses of nearest-pole asymptotics remain explicit. No historical priority is inferred from successful computation or formal verification.

Replay the worked layer

The model checker is `checks/verify_examples.py`. The plate generator is `source/generate_graphics.py`. They require the listed Python dependencies. Run those mathematical tools in a temporary copy to preserve the delivered evidence.

> **Read this as:** A proof, a finite exact test, a root enclosure, and a sampled curve answer different questions. Keeping their scopes explicit makes the pictures useful mathematical evidence.

Exact certificates and formal verification records accompany the source. Each retains its stated mathematical scope.