CORE SERIES XI

Tame and Wild Packet Geometry

For local Langlands

Visual guide

Komorebi P.

Revised edition 1.1.0 / 18 October 2026

Reading scope

This guide follows the paper's mathematical route from finite Frobenius packets to finite wild comparison, then explains the actual arithmetic constructions developed in Blocks 166–167.

The general arithmetic realization endpoint is stated as **Conjecture 66**. The comparison with the established local Langlands correspondence remains a theorem conditional on an actual realization and H5 for that same candidate.

All mathematical artwork is deterministic Python/Matplotlib vector output. The complete editable source and its figure generator accompany this guide.

How to use this companion

Read the first two parts as an atlas of finite mathematical objects. The third part explains where finite objects acquire an actual arithmetic carrier. The final part separates the remaining realization conjecture from the conditional comparison theorem.

Each spread develops one construction or one distinction. Displays preserve the normalizations used in the accompanying paper. The source line identifies an original manuscript section or a retained proof note. These are retained source anchors, rather than a claim that every argument has been newly formalized. The separate research atlas provides the later conductor, occurrence and native-marking reconciliation.

The symbol K denotes the original local field in the finite-theory discussion. In the explicit arithmetic constructions, $F = \mathbb{F}_p((t))$ is the base field and $K = F[\beta]$ is its totally ramified parameter field. A symplectic quotient dimension and a Heisenberg representation dimension are different quantities.

Theorem numbering. The historical proof obligation called "66" corresponds to the endpoint printed as Theorem 16.2 in the original manuscript. Conjecture 66 now isolates its general arithmetic realization requirement. It is not a claim that row 66 of the historical theorem register is this endpoint.

Supplementary materials. The accompanying release includes Markdown (.md) files containing detailed proofs, construction notes, proof audits, and research records. These documents distinguish established results, conditional results, and the remaining obligations associated with Conjecture 66.

Evidence. Blocks 166–167 contain ordinary mathematical arguments using the stated classical type-theory results. Their bounded exact computations support specified identities. They are not fresh Lean certificates for the arbitrary-length arithmetic theorems. The separate Lean guide states the actual formalization scope.

Two visual routes, one claim map

This is the core visual guide. It explains the mathematical objects and the restricted arithmetic constructions in the monograph. The separate 36-page research atlas explains the subsequent comparison tests, failed routes and remaining certificate. The two guides are complementary, not competing versions of the same document.

Start with the question	Continue to the linked reader
What is being realized?	Monograph realization contract and research plate V01. Finite comparison and arithmetic existence are different assertions.
How does recovery preserve the candidate?	Monograph restricted constructions and research plate V10. A changed assembly carrier does not mean a changed inducing representation.
Why do numerical tests not settle attachment?	Progress Sections 22–25 and research plates V11–V19. The full and skipped rank-eight profiles have capacities six and four despite the same three numerical invariants.
What does a dyadic marking retain?	Progress Sections 26–29 and research plates V20–V29. Squares, section products, native choices and phase data must not be collapsed into one commutator form.
What is formally checked?	Lean guide: the existence input and the conditional signature. A supplied field in a structure is not constructed by typesetting its declaration.

Keep the five reader PDFs together. These links use the actual companion pages, while the original lessons keep their source-note and section anchors. No diagram in either guide certifies arithmetic inputs that its source argument leaves open.

Reading corrections without changing the contract

One candidate throughout. A source-defined induction candidate and a directly constructed type candidate are not identified merely by notation. Occurrence, full-character normalization and the noncompact extension must concern the same representation. The Part II occurrence analysis retains the excluded original route and the corrected route's remaining obligation.

A transported mark is not a prescribed native mark. The positive scalar-pushout attachment carries the supplied finite marking through a center-fixing homomorphism. It does not identify that output with an independently fixed host subgroup. Conversely, the stronger native problem must not be inserted as a new premise into the different transported-marking result.

The equal-capacity example is conditional. At rank eight the full profile has exactly six quotient dimensions. The finite Arf test applies when the actual native refinement is independently prescribed. The abstract six-dimensional forms in the research atlas are examples, not a computation of that candidate's native square bits.

A successful finite repair has inputs. The transvection starts with an allowed symplectic map and an isotropic discrepancy vector. The odd-action correction starts with genuine supplied finite actions. Neither constructs the arithmetic normalizer or chooses every honest phase. See plates V25–V29.

Evidence and publication status

This revision rebuilds the core guide from its editable source and preserves its Python-generated vector figures. The historical matrix calculations and ordinary construction arguments keep their dates and scopes. The Lean and evidence guide records the separate current compiler replay; the PDF build itself does not certify arithmetic constructors.

The versioned ledger distinguishes publication and formal verification from the open arithmetic endpoints. The selected Henniart–Lemaire source interface has now been checked against the full official text, including its normalization conventions. The complete Block 2 editable archive remains unavailable. Its research-reader pages are preserved. No external peer review is claimed.

For the final same-candidate comparison, read the monograph comparison proposition alongside research plate V32. The conjecture itself does not assert the independent H5 step.

Contents

How to use this companion 1

Two visual routes, one claim map 2

Reading corrections without changing the contract 3

1 Three mathematical objects 6

2 Monodromy is a finite transfer chain 7

3 Fourier projectors isolate tame inertia 8

4 One orbit gives one tame atom 9

5 Regular orbits have exact counts 10

6 A primitive projector selects a cuspidal block 11

7 Tensor refinement is a diagonal orbit problem 12

8 Exponent coordinates test inertia triviality 13

9 The closing scalar carries noncompact information 14

10 Local polynomials and fixed vectors 15

11 Finite Gauss phases are unit scalars 16

12 Translations and multipliers create Heisenberg packets 17

13 An isotropic average selects one line 18

14 Projective choices leave a commutator invariant 19

15 Clifford multiplicity is the correct action space 20

16 The type-side quotient is an actual group quotient 21

17 A supplied bridge reduces comparison to a scalar 22

18 Odd phases have a determinant certificate 23

19 Two dyadic refinements can have opposite signs 24

20 Refinements form an affine family 25

21 A spanning tree fixes the vertex gauges 26

22 Cycle defects measure edge residuals exactly 27

23 Orthogonal breaks become tensor factors 28

24 A recursive matrix gives a literal parameter 29

25 The approximation degrees and critical levels 30

26 Odd towers retain native trace zero 31

27 Dyadic trace one is deep enough 32

28 Native trace and ambient trace are different 33

29 Full characters require shallow compatibility 34

30 The same arithmetic candidate has exact profiles 35

31 Dyadic capacity has two distinct dimensions 36

32 A sector can retain the whole source packet 37

33 Retain the whole marked finite extension 38

34 A native quadratic marking adds an embedding condition 39

35 Low-rank repair enlarges an actual compact group 40

36 Rank four gains a four-dimensional carrier 41

37 Induction in stages recovers the same candidate 42

38 Compact recovery must remember the noncompact root 43

39 Central completion is a construction choice 44

40 Why nontrivial stabilizer actions need another step 45

41 Genuine scalar phases can descend from a finite cover 46

42 Numerical matching does not construct realization 47

43 Conjecture 66 isolates the missing existence claim 48

44 H5 compares this same actual candidate 49


45 What the restricted realizations establish 50

46 Exact computation checks a bounded identity 51

47 Where the complete arguments are retained 52

1 Three mathematical objects

Source: Original manuscript, Sections 2, 11, 16; contract audit 162.



The finite comparison does not create the arithmetic witness.

An atom, its finite packet data, and its arithmetic realization have different roles.

A Weil atom is an irreducible Weil representation. A finite packet records selected quotient spaces, projectors, central extensions, phases, and transports. An arithmetic realization is an actual admissible representation of the required general linear group together with a comparison and recovery of that packet.


Monodromy organizes an atom into a finite Weil–Deligne chain. It does not supply an atom that is missing. Likewise a finite Heisenberg comparison identifies already supplied finite representations, rather than automatically constructing a representation of $\mathrm{GL}_n(K)$.

Reading the diagram. The final existence arrow is the content of Conjecture 66. Finite comparison, arithmetic construction, and local-factor characterization are separate steps.

2 Monodromy is a finite transfer chain

Source: Original Section 2.

$$F = \mathrm{diag}(1, q, q^2, q^3, q^4), \qquad FNF^{-1} = q^{-1}N$$



$$N(V_j) \subseteq V_{j-1}; \qquad N^5 = 0$$

The arrows point from V_j to V_{j-1}, the direction of N.

Put $V = \bigoplus_{j=0}^{m-1} V_j$ and let P_j be the coordinate projector. If $N(V_j) \subseteq V_{j-1}$ and $N(V_0) = 0$, then

$$N = \sum_{j=1}^{m-1} P_{j-1} N P_j, \qquad N^m = 0.$$

Every term in N^r follows a path of length r down the chain. There is no path of length m. With $F = \mathrm{diag}(1, q, \ldots, q^{m-1})$, each nonzero transfer block gains the conjugation factor $q^{j-1}q^{-j} = q^{-1}$.

Reading the diagram. Nilpotence and covariance follow from the displayed block structure. Assigning the initial supercuspidal atom is a separate arithmetic input.

3 Fourier projectors isolate tame inertia

Source: Original Section 3; geometric Frobenius convention.

Let t generate a finite tame quotient of order M, with $(q, M) = 1$. For $j \in \mathbb{Z}/M\mathbb{Z}$ define

$$P_j = \frac{1}{M} \sum_{a=0}^{M-1} \zeta_M^{-ja} r(t)^a, \qquad \zeta_M = e^{2\pi i/M}.$$

The finite Fourier identities give $P_j P_k = \delta_{jk} P_j$ and $\sum_j P_j = I$. Thus $P_j V$ is the ζ_M^j character space for $r(t)$. The guide uses geometric Frobenius with

$$\Phi^{-1} r(t) \Phi = r(t)^q.$$

Equivalently $\Phi r(t) \Phi^{-1} = r(t)^{q^{-1}}$, where the inverse is taken modulo M. Substituting in the Fourier sum gives


$$\Phi P_j \Phi^{-1} = P_{qj}.$$

Scope. If the forward relation $\Phi r(t) \Phi^{-1} = r(t)^q$ is used instead, the projector label is $q^{-1}j$. The two directions have the same orbit sets but different conjugation identities. The release corrects the historical mixed-convention sentence.

Reading the diagram. A packet is an actual eigenspace selected by an idempotent. Frobenius carries it to the next character packet.

4 One orbit gives one tame atom

Source: Original Theorem 3.2.



For $q = 2$ modulo 15, the displayed geometric projector orbit is $(1, 2, 4, 8)$.

For an irreducible tame Weil representation, a union of Frobenius orbits of nonzero inertia packets would give a stable subspace. Irreducibility therefore forces one orbit, of some length f.


The stabilizer of a packet is $W_{K_f} = \langle I_K, \Phi^f \rangle$, where K_f/K is the unramified extension of degree f. Inertia acts on the packet by one scalar character. Its remaining stabilizer action is generated by Φ^f; an irreducible complex action of this cyclic quotient is one dimensional. Hence

$$\tau \simeq \operatorname{Ind}_{W_{K_f}}^{W_K} \xi, \qquad \dim \tau = f.$$

Reading the diagram. Orbit transitivity supplies both the dimension and the induced model. It is a theorem about the tame case.

5 Regular orbits have exact counts

Source: Original Proposition 3.6.



Exact integer counts; the vertical scale is logarithmic.

Characters of $\mathbb{F}_{q^n}^{\times}$ are indexed modulo $q^n - 1$. Characters fixed by a Frobenius power of degree $r \mid n$ number $q^r - 1$. Möbius inversion removes those belonging to smaller orbits:

$$c_n(q) = \frac{1}{n} \sum_{d \mid n} \mu(d)(q^{n/d} - 1).$$

For $q = 2, n = 4$, the regular exponent orbits modulo 15 are

$$(1, 2, 4, 8), \quad (3, 6, 12, 9), \quad (7, 14, 13, 11).$$

Their number is $\frac{1}{4}(15 - 3) = 3$. The plotted values are computed by the exact divisor sum and retained in `figure_data.json`.

Reading the diagram. Regularity is an exact orbit property. The plot illustrates an integer formula rather than supplying its proof.

6 A primitive projector selects a cuspidal block

Source: Original Section 4.

Let $G_n = \mathrm{GL}_n(\mathbb{F}_q)$ and $T_n \simeq \mathbb{F}_{q^n}^{\times}$ be an elliptic torus. A regular torus character determines an irreducible cuspidal representation κ_θ by the classical finite-field construction. If its character is χ_θ and degree is d_θ, then

$$e_\theta = \frac{d_\theta}{|G_n|} \sum_{g \in G_n} \chi_\theta(g^{-1}) g$$

is its primitive central idempotent in $\mathbb{C}[G_n]$.

On an irreducible representation with character ψ, this central element acts by a scalar. Taking traces and using character orthogonality makes the scalar 1 on κ_θ and 0 on every other irreducible block.

For a proper parabolic $P = LU$, averaging over its unipotent radical gives


$$e_U = |U|^{-1} \sum_{u \in U} u, \qquad e_U e_\theta = e_\theta e_U = 0.$$

The average selects U-fixed vectors; the cuspidal block has none.

Scope. The existence and classification of κ_θ are classical inputs. The idempotent calculation explains its exact finite selector.

7 Tensor refinement is a diagonal orbit problem

Source: Original Theorem 5.1.



The 4×6 grid separates into two diagonal translation orbits, labelled 0 and 1.

The simultaneous translation action on $\mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ has

$$d = \gcd(n, m) \quad \text{orbits, each of length} \quad L = \operatorname{lcm}(n, m).$$

Mackey decomposition then gives

$$\operatorname{Ind}_{W_{K_n}}^{W_K} \xi \otimes \operatorname{Ind}_{W_{K_m}}^{W_K} \eta \simeq \bigoplus_{i=0}^{d-1} \operatorname{Ind}_{W_{K_L}}^{W_K} \left(\xi_{L,n} \eta_{L,m}^{\Phi^i}\right).$$

Here the subscripts denote norm lifts and the Frobenius superscript follows the fixed source convention. The dimension identity is exact: $dL = nm$.

Scope. These are interaction packets before combining isomorphic summands. A refined induced summand need not remain irreducible.

8 Exponent coordinates test inertia triviality

Source: Original Proposition 5.3.

Write the residue character exponents as $a \bmod (q^n - 1)$ and $b \bmod (q^m - 1)$. The ith refined packet has exponent

$$c_i = a\frac{q^L - 1}{q^n - 1} + bq^i\frac{q^L - 1}{q^m - 1} \pmod{q^L - 1}.$$

Norm lifting multiplies an exponent by the quotient of group orders. Shifting the second character by Frobenius multiplies its exponent by q^i. Multiplication of characters adds the two terms.

For $q = 2, n = 4, m = 6$, the common modulus is $2^{12} - 1 = 4095$. Taking $a = b = 1$, the two packets have


$$c_0 = 273 + 65 = 338, \qquad c_1 = 273 + 130 = 403.$$

Both are nonzero modulo 4095, so neither packet contributes inertia invariants.

Reading the diagram. The test is $c_i = 0$ in the common modulus. A numerical-looking residue formula is an exact character calculation.

9 The closing scalar carries noncompact information

Source: Original Propositions 3.5 and 4.4.



Closing edge a: characteristic polynomial $X^4 - a$

The closing scalar is basis-invariant after multiplication around the cycle.

In an orbit basis let $\Phi e_j = e_{j+1}$ for $j < n-1$ and $\Phi e_{n-1} = a e_0$. Then

$$\det(XI - \Phi) = X^n - a, \qquad \det \Phi = (-1)^{n-1} a.$$

Rescaling the individual basis vectors changes edge coefficients, but their product remains a. An unramified twist ν does not change the inertia orbit and sends the closing scalar to $a\nu(\Phi)^n$.

Reading the diagram. Compact inertia information alone does not determine the unramified completion. The cyclic scalar is additional data.

10 Local polynomials and fixed vectors

Source: Original Propositions 5.4–5.8.

For a packet of orbit length r and closing scalar a,

$$\det(I - u\Phi \mid V^{I_K}) = \begin{cases} 1 - au^r, & \text{if inertia acts trivially,} \\ 1, & \text{otherwise.} \end{cases}$$

Thus tame tensor local factors multiply $(1 - a_i u^{r_i})^{-1}$ over precisely the refined packets whose inertia exponent vanishes.

An inertia-trivial companion cycle has a Weil-fixed vector exactly when $a_i = 1$, and then its fixed space is one dimensional. Therefore

$$\dim(\tau_{\xi} \otimes \tau_{\eta})^{W_K} = \#\{i : c_i = 0,\ a_i = 1\}.$$

For an irreducible nontrivial tame atom of dimension $n > 1$, wild inertia acts trivially and inertia has no fixed vectors. Hence

$$\mathrm{Sw}(\tau) = 0, \qquad \mathrm{Art}(\tau) = n.$$

Scope. These exact tame formulas do not establish the full wild pair-factor identities required by H5.

11 Finite Gauss phases are unit scalars

Source: Original Section 6.

For a nontrivial additive character ψ of $\mathbb{F}_q$ and a nontrivial multiplicative character χ of $\mathbb{F}_q^\times$, define

$$G(\chi,\psi)=\sum_{x\in\mathbb{F}_q^\times}\chi(x)\psi(x),\qquad \gamma(\chi,\psi)=q^{-1/2}G(\chi,\psi).$$

The exact identities are

$$|G(\chi,\psi)|^2=q,\qquad \gamma(\chi,\psi)\gamma(\chi^{-1},\psi)=\chi(-1).$$


The magnitude calculation substitutes $x=ty$ in the double sum. The inner sum is $q-1$ for $t=1$ and -1 otherwise; multiplicative-character orthogonality supplies the result.

The normalized scalar therefore lies on the unit circle. Its value depends on the declared additive character and lifting normalization. Finite Gauss phases capture a residue-field part of normalization.

Scope. A full wild local factor also contains ramification depth, additive-character conductor, measure normalization, and pair interaction data. One finite phase is not that full factor.

12 Translations and multipliers create Heisenberg packets

Source: Original Section 7.



Noncommuting operators generate the full matrix algebra.

The commutator is central and nontrivial when the pairing is nonzero.

For odd p let $V = X \oplus X^*$ and

$$b((x,\xi),(y,\eta)) = \eta(x) - \xi(y).$$

The group $H(V) = V \times \mathbb{F}_p$ has product

$$(v,z)(w,t) = (v+w, z+t+\tfrac{1}{2}b(v,w)).$$

On $\mathcal{H} = \mathbb{C}^X$ the Schrödinger operator is

$$(\rho(x,\xi,z)f)(u) = \psi(z + \xi(u) + \tfrac{1}{2}\xi(x)) f(u+x).$$

Translations and character multipliers generate every matrix unit by finite Fourier inversion. Thus the representation is irreducible and has dimension $|X| = \sqrt{|V|}$.

13 An isotropic average selects one line

Source: Original Propositions 7.3–7.6.

The Schrödinger character is supported on the center:

$$\operatorname{Tr}\rho(v,z)=\begin{cases}|X|\psi(z), & v=0,\\ 0, & v\neq 0.\end{cases}$$

A nonzero translation has no fixed coordinate. A nontrivial multiplier has vanishing character sum.

For a maximal isotropic subspace L and a subgroup lift $\widetilde{L}$ with trivial central coordinate, the average

$$P_L=\frac{1}{|L|}\sum_{\ell\in\widetilde{L}}\rho(\ell)$$

is a self-adjoint idempotent. Its trace is $|X|/|L|=1$, because only the identity contributes. Its range is therefore one dimensional.


For $L=X$ this range consists of constant functions:

$$(P_Lf)(u)=|X|^{-1}\sum_{x\in X}f(u+x).$$

Reading the diagram. The complementary-line projector later used in arithmetic attachment is an exact operator in the Heisenberg image algebra. Its existence is finite representation theory.

14 Projective choices leave a commutator invariant

Source: Original Section 8.



Radical directions commute with every projective generator.

A cocycle contains choices; its alternating commutator forgets scalar rescaling.

For $T_aT_b = c(a,b)T_{a+b}$ define $b_c(a,b) = c(a,b)c(b,a)^{-1}$. Replacing T_a by $\lambda_a T_a$ multiplies c by the symmetric coboundary $\lambda_a\lambda_b\lambda_{a+b}^{-1}$, so b_c is unchanged.

The radical consists of a commuting projectively with every b. Its quotient carries a nondegenerate alternating pairing. In the nondegenerate case the twisted group algebra has scalar center and is a full matrix algebra:

$$\mathbb{C}_c[A] \simeq M_{\sqrt{|A|}}(\mathbb{C}).$$

Scope. This constructs the finite envelope for a supplied projective action. It does not produce a canonical action for every arbitrary wild inertia atom.

15 Clifford multiplicity is the correct action space

Source: Original Section 9.

Supply finite groups $N \triangleleft G$, elementary abelian quotient $A = G/N$, and irreducibles Π and ρ, with ρ fixed up to isomorphism by G and $\Pi|_N$ isotypic. The multiplicity space is

$$M_\rho = \mathrm{Hom}_N(W, X), \qquad W \otimes M_\rho \simeq X.$$

For a section $s : A \to G$ and intertwiners I_a of ρ, write

$$\Pi(s(a)) = I_a \otimes T_a.$$

The operators T_a give a projective action on M_ρ. Section and intertwiner changes alter it only by scalar changes and conjugation. Its commutator, radical, and symplectic quotient are intrinsic to this supplied datum.

If the commutator is nondegenerate, $\dim M_\rho = \sqrt{|A|}$. A tame stabilizer acting by automorphisms of the datum preserves the induced symplectic form.

Scope. The normal group, isotypic constituent, and one-break interpretation are assumptions of the restricted construction. They have not been constructed for every wild atom.

16 The type-side quotient is an actual group quotient

Source: Original Section 10.

For an actual maximal simple character θ of $\mathrm{GL}_n(K)$, classical simple-type theory supplies

$$H^1_\theta \subseteq J^1_\theta \subseteq J_\theta, \qquad V_\theta = J^1_\theta / H^1_\theta.$$

The commutator pairing is encoded by

$$\psi_0(h_\theta(\bar{x}, \bar{y})) = \theta([x, y]).$$

It is nondegenerate and alternating. The corresponding irreducible Heisenberg representation η_θ of J^1_θ contains θ and satisfies


$$\dim \eta_\theta = \sqrt{|J^1_\theta / H^1_\theta|}.$$

The canonical beta extension supplies a normalized extension of this representation using the declared type-theory data and finite corrections.

Scope. An arbitrary abstract symplectic space is not automatically a quotient of an actual simple character. An equivalent inducing carrier need not have this original quotient or a newly invented canonical beta extension.

17 A supplied bridge reduces comparison to a scalar

Source: Original Section 11.



A cyclic generator phase removes the discrepancy character.

The symplectic isometry is an input. Stone–von Neumann supplies the lower intertwiner.

Supply an equivariant symplectic isometry $\iota : V_\rho \to V_\theta$, a common nontrivial central character, and, in characteristic two, the compatible refinements and lift of central extensions.

The two pulled-back irreducible Heisenberg representations then have the same central character. Stone–von Neumann gives an invertible U with

$$U\omega_\rho(h) = \omega_\theta(\iota_H(h))U.$$

Schur's lemma makes U unique up to a nonzero scalar. After honest cyclic stabilizer linearizations, their discrepancy is a character $\delta : C \to \mathbb{C}^\times$. Equality at one generator kills this character.

Scope. Finite linearization does not identify the stabilizer action with an arithmetic action on a newly constructed common candidate.

18 Odd phases have a determinant certificate

Source: Original Section 12.

For an odd prime p, a symplectic space V of dimension $2r$, and fixed-point-free $g \in \mathrm{Sp}(V)$, the declared finite Weil normalization gives

$$\Phi_p(g, V) = \left(\frac{-1}{p}\right)^r \left(\frac{\det(g - I)}{p}\right).$$

With $P_g(X) = \det(XI - g)$, even dimension gives $\det(g - I) = P_g(1)$. The phase is therefore read from a characteristic polynomial and a square class.

For a hyperbolic space $U \oplus U^*$ with $g = (a, (a^{-1})^*)$ and $a - I$ invertible,

$$\Phi_p(g, U \oplus U^*) = \left(\frac{\det a}{p}\right).$$

If g has order prime to p, then


$$V = \ker(g - I) \perp \mathrm{im}(g - I).$$

The moving summand is nondegenerate and fixed-point-free. Orthogonal block phases multiply.

Scope. The fixed-point-free formula applies to the moving part. The full trace with a fixed part requires its declared normalization data.

19 Two dyadic refinements can have opposite signs

Source: Original Section 13.



The entries are exact values in $\mathbb{F}_2$; both polar forms are $xy' + x'y$.

A quadratic refinement satisfies $q(x+y) + q(x) + q(y) = b(x,y)$. On a plane,

$$q_0(x,y) = xy, \qquad q_1(x,y) = xy + x + y.$$


For q_0 there are three zeros and one one, so its Gauss sum is 2. For q_1 there is one zero and three ones, so its sum is -2. The alternating forms agree but the Arf invariants differ.

$$\sum_{x \in V} (-1)^{q(x)} = (-1)^{\mathrm{Arf}(q)} 2^r, \qquad \dim V = 2r.$$

Reading the diagram. Keeping the alternating form alone loses dyadic phase information. The refinement and the marked central extension must travel with the packet.

20 Refinements form an affine family

Source: Original Propositions 13.2 and 13.8–13.9.



Exact counts of refinements of a fixed nondegenerate alternating space.

The difference between two refinements is linear. Nondegeneracy identifies every linear form with $b(v,\cdot)$, so all refinements are

$$q_v(x) = q(x) + b(v,x), \qquad \mathrm{Arf}(q_v) = \mathrm{Arf}(q) + q(v).$$

The numbers of the two Arf types are


$$2^{r-1}(2^r+1), \qquad 2^{r-1}(2^r-1).$$

For $r = 1$ these are 3 and 1. For $r = 3$ they are 36 and 28. Orthogonal sums add Arf invariants and multiply normalized Gauss signs.

Reading the diagram. Equal dimension and equal alternating form do not force equal refinement. An embedding fixed in advance must preserve q pointwise.

21 A spanning tree fixes the vertex gauges

Source: Original Theorem 14.2.



Four vertices and five edges leave two fundamental cycle tests.

Let $A_e : V_u \to V_v$ and $B_e : W_u \to W_v$ be invertible transports, with reverse-edge inverses. Fix a root comparison U_r.

Choose a spanning tree. Along its unique root path p_v define

$$U_v = B_{p_v} U_r A_{p_v}^{-1}.$$

This forces every tree-edge equation $U_v A_e = B_e U_u$. Each non-tree edge closes one fundamental cycle. Its edge equation holds exactly when the root comparison intertwines that cycle's holonomy.

Reading the diagram. All edge equations are equivalent to the fundamental cycle identities $U_r A_\gamma = B_\gamma U_r$. Holonomy need not be individually trivial; the two holonomy representations must agree through the root map.

22 Cycle defects measure edge residuals exactly

Source: Original Proposition 14.6.

For a non-tree edge $e : u \to v$, let

$$D_e = U_r A_{p_v}^{-1} A_e A_{p_u} - B_{p_v}^{-1} B_e B_{p_u} U_r.$$

Substituting the tree-defined gauges gives the exact identity

$$U_v A_e - B_e U_u = B_{p_v} D_e A_{p_u}^{-1}.$$

Consequently


$$\|U_v A_e - B_e U_u\| \le \|B_{p_v}\| \, \|D_e\| \, \|A_{p_u}^{-1}\|.$$

Small cycle defects give small edge residuals when the tree transports are conditioned. Zero defects give exact finite gluing. In the scalar case there are $|E| - |V| + 1$ independent tests on a connected graph.

Scope. The theorem assumes both sets of fibers and transports already exist. Gauge propagation is not an arithmetic existence theorem.

23 Orthogonal breaks become tensor factors

Source: Original Section 15.



$$V = V_{s_1} \perp V_{s_2} \perp V_{s_3}$$

$$\omega_V \simeq \omega_{s_1} \otimes \omega_{s_2} \otimes \omega_{s_3}$$

Symplectic direct sum and Heisenberg tensor product are compatible.

For $V = \perp_s V_s$ and compatible central characters,


$$\omega_V \simeq \bigotimes_s \omega_{V_s}, \qquad \prod_s \sqrt{|V_s|} = \sqrt{|V|}.$$

The groups combine by a central product. Tensoring the individual intertwiners gives the full finite bridge, once the stabilizer actions and phases agree. Transport-graph compatibility is then the matrix holonomy condition.

Scope. Separate actual type quotients at each break do not, merely by tensoring, become one actual maximal simple type in the source atom's rank. Constructing a common arithmetic carrier and recovering a candidate are the extra realization requirements.

24 A recursive matrix gives a literal parameter

Source: Blocks 166–167, tower notes.



The derived field and the final field are not identified.

The recursive lift is an actual filtered right inverse of normalized corestriction.

Start over a coefficient field L with uniformizer t_L and form

$$E = L(\alpha), \qquad \alpha^p - \alpha = t_L^{-q}.$$

At a recursive stage, C is an actual matrix over E. The outer matrix is


$$B = \alpha I_m + \mathrm{Lift}_E(C).$$

The lift has prescribed corestriction and preserves its exact filtration valuation. The field and simple-stratum proof uses the actual displayed matrix, together with the cited type-theory derived-stratum criterion.

Scope. The derived field $E[C]$ is not identified with the final field $L[B]$. The recursion proves a literal defining sequence without silently declaring intermediate fields to be nested inside the final field.

25 The approximation degrees and critical levels

Source: Blocks 166–167, combined arithmetic statements.



These examples were recomputed exactly; the all-length assertions are proved separately.

For increasing permitted $h_1 < \cdots < h_d$, put $q_i = (p+1)h_i$ and $N = p^d$. The approximation-field degrees are $p, p^2, \ldots, p^d$, with

$$n_j = p^{d-j} q_{d-j+1}.$$

For odd p, require $p \nmid h_i$. For $p = 2$, require all h_i odd and positive. The leading ambient level is n_1; $n_2, \ldots, n_d$ are the successive critical approximation levels.

Scope. Every amplified ambient approximation still has leading level n_1. The declining values plotted here are critical levels, not new leading valuations for those ambient matrices.

26 Odd towers retain native trace zero

Source: Block 166, recursive tower and full-character proofs.

For odd p, $\mathrm{Tr}_{E/L}(\alpha) = 0$. Choose b with $qb \equiv -1 \pmod{p}$, let $a = (qb+1)/p$, and set $z = t_L^a \alpha^b$. Then z is a uniformizer.

For its Eisenstein polynomial f and constant-coordinate projection P_0, the normalized corestriction uses $\kappa = -f'(z)/z^{p-1}$. Writing $u_0 = s(P_0)$,


$$\mathrm{Lift}_E(C) = u_0^{-1} C P_0, \qquad \mathrm{tr}_L \,\mathrm{Lift}_E(C) = \mathrm{Tr}_{E/L}(\kappa^{-1}\,\mathrm{tr}_E\, C).$$

If $\mathrm{tr}_E\, C = 0$, the lift has trace zero. Adding αI_m preserves trace zero. Induction therefore gives trace zero at every native stage and every native correction.

Reading the diagram. This is stronger than the automatic ambient trace cancellation from a multiplicity divisible by p. Native trace controls the scalar restriction used in full-character construction.

27 Dyadic trace one is deep enough

Source: Block 167, depth invariant.



The scalar correction disappears when its argument lies in $t\mathcal{O}_F$.

Fix ψ_F of conductor $t\mathcal{O}_F$. On a scalar unit $u \in U_F^s$, the additive correction sees $(u-1)\,\mathrm{tr}\,\delta$. Hence


$$v_F(\mathrm{tr}\,\delta) + s \geq 1 \quad \Longrightarrow \quad \psi_F((u-1)\,\mathrm{tr}\,\delta) = 1.$$

The dyadic native corrections have trace 1. Its valuation is 0, so $s = 1$ already satisfies the inequality. Thus trace zero is sufficient but unnecessary.

Scope. Euler's weighted-visibility viewpoint motivated this step. No Euler coefficient theorem is an arithmetic proof input here. In characteristic two, the weights are uniformizer depths, not integer powers of two in the coefficient field.

28 Native trace and ambient trace are different

Source: Block 167, exact trace transport.



Native trace and amplified ambient trace have different roles.

A trace-one native matrix amplified by an even multiplicity has ambient trace zero.

For odd q, let $a = (q+1)/2$ and $z = t_L^a \alpha$. Then

$$z^2 + t_L^a z + t_L = 0, \quad \mathrm{Tr}_{E/L}\,\alpha = 1, \quad \mathrm{Lift}_E(C) = (z^2/t_L)CP_0.$$

The trace identity is

$$\mathrm{tr}_L\,\mathrm{Lift}_E(C) = \mathrm{Tr}_{E/L}(\alpha\,\mathrm{tr}_E\,C).$$

It sends inner trace 1 to outer trace 1. At native rank 2^j, every constructor has trace 1; its next native correction has trace $1 - 2 \cdot 1 = 1$. In rank 8, the earlier amplified approximations instead have traces $0, 0$, and the final matrix has trace 1.

Reading the diagram. Only the appropriate native trace establishes the scalar-character compatibility at each recursive stage.

29 Full characters require shallow compatibility

Source: Blocks 166–167, central character induction.

The full-character induction has an essential order. First construct the full native character, then transfer it to a higher multiplicity, add the actual correction on its forced domain, and extend at the next maximal native parameter field.

On a scalar $u \in U_F^1$, multiplicity p transfer gives

$$\Theta_j(uI) = \theta_j(u^p) = 1.$$

The correction is trivial on the scalar intersection by trace zero in the odd case, or integral trace and the depth inequality in the dyadic case. The forced deep character therefore agrees with the imposed trivial scalar character on their intersection. A finite abelian unit-character extension supplies the full next character.

For an odd dyadic critical level r, put $m = \lfloor r/2 \rfloor$. The endpoint satisfies $2(m+1) - r = 1$, making the additive cross term lie in the character's radical.

Scope. Checking a leading trace alone does not check shallow forced values. The earlier fixed-field central argument failed on an actual scalar unit; the recursive native-field proof repairs that problem.

30 The same arithmetic candidate has exact profiles

Source: Combined statements 166–167.

Let $N = p^d$ and retain the constructed full character, canonical beta extension, and compatible central completion. Both families give an actual irreducible admissible supercuspidal candidate of $\mathrm{GL}_N(F)$ with torsion number 1.

For odd p,

$$\begin{aligned} \mathrm{Sw}(\pi) &= p^{d-1}(p+1)h_d, \\ \dim_{\mathbb{F}_p}(J^1/H^1) &= N-1, \quad \dim \eta = p^{(N-1)/2}, \\ a(\pi^\vee \times \pi) &= p^{2d} - 1 + \sum_{i=1}^{d} (p^{2i} - p^{2i-2})h_i. \end{aligned}$$


For $p = 2$,

$$\begin{aligned} \mathrm{Sw}(\pi) &= 3\,2^{d-1}h_d, \\ \dim_{\mathbb{F}_2}(J^1/H^1) &= N-2, \quad \dim \eta = 2^{N/2-1}, \\ a(\pi^\vee \times \pi) &= N^2 - 1 + \sum_{i=1}^{d} 3\,4^{i-1}h_i. \end{aligned}$$

Scope. These identities belong to the same constructed candidate. Matching numerical profiles does not identify a source Weil atom, an endo-class, or H5 pair factors.

31 Dyadic capacity has two distinct dimensions

Source: Block 167, capacity proof.



Compare symplectic dimensions first. The original low-rank quotient remains too small.

The host symplectic quotient has dimension $N-2$ over $\mathbb{F}_2$; the supplied whole source of d planes has dimension $2d$. Its original-type embedding is possible by dimension exactly when $d \geq 3$.

N	host quotient	source quotient	host carrier	source carrier
2	0	2	1	2
4	2	4	2	4
8	6	6	8	8
16	14	8	128	16

Scope. The low-rank obstruction concerns the original maximal-simple-character carrier. Enlarged actual arithmetic carriers repair ranks 2 and 4 without changing this quotient theorem.

32 A sector can retain the whole source packet

Source: Blocks 166–167, scalar-pushout attachment.

Suppose the supplied full break characters have labelled plane quotients and satisfy the one-vertex scalar-pushout assembly contract, initially with $C = 1$. Their source representation has dimension $p^d = N$.

For odd p, the actual leading subspace has dimension $(p-1)p^{d-1} \geq 2d$. For dyadic $d \geq 3$, use the host quotient dimension $N - 2 \geq 2d$. Choose the declared symplectic embedding into the actual host.

An orthogonal splitting $W = V_{\mathrm{src}} \perp V_{\mathrm{src}}^{\perp}$ gives


$$\eta_W \simeq \eta_{\mathrm{src}} \otimes \eta_{\perp}.$$

A maximal-isotropic average on the complementary Heisenberg factor selects a one-dimensional representation line. Tensoring its projector with the identity gives a rank-N image-algebra projector. The retained induction/recovery theorem recovers the same Λ and then the same π.

Scope. For $p > 3$, genuinely supplied plane data are required: a native minimal rank-p quotient has dimension $p - 1$ and is not automatically a plane. The construction retains source labels but does not prove their old numerical depth placement.

33 Retain the whole marked finite extension

Source: Block 167, markings proof.



The whole finite subgroup travels through the map.

A separately fixed native marking needs its own comparison.

Squares are preserved by the actual map of marked extensions.

For a full character θ_i, its scalar pushout is

$$\mathcal{P}_i = (\mathbb{C}^\times \times J_i^1)/\{(\theta_i(h)^{-1}, h) : h \in H_i^1\}.$$

A center-fixing comparison retains every supplied character value. In the dyadic case retain the image of the entire marked finite subgroup, its central embedding, lifts, squares, refinements, and phases. An actual extension map F_0 obeys $F_0(a)^2 = F_0(a^2)$.

Scope. The unmarked complex pushouts of D_8 and Q_8 coincide, while their marked finite μ_2 extensions differ. Retaining only the central $\{\pm 1\}$ loses that distinction. Invisible kernel elements already removed by the central character cannot be recovered by pushout.

34 A native quadratic marking adds an embedding condition

Source: Block 167, markings theorem.

Transported source markings and an independently prescribed native host marking are different inputs. If the native host refinement q_W is independently supplied, a prescribed embedding f must satisfy

$$q_W(f(v)) = q_{\mathrm{src}}(v) \quad \text{for every } v.$$

With a free embedding and equal dimensions, quadratic classification requires equal Arf invariants. This is the rank-eight case: both quotient spaces have dimension 6.


For $d \geq 4$, the dyadic host has at least one spare symplectic plane. The complement's Arf type can absorb the difference, so either source Arf type admits an embedding into the supplied native quadratic host.

Reading the diagram. Dimension controls symplectic room. Arf controls a quadratic isometry class. A pointwise equation controls an already chosen embedding.

Scope. Capacity does not manufacture a canonical native finite marking or identify it with the transported marking. That comparison remains outside the general realization proof.

35 Low-rank repair enlarges an actual compact group

Source: Block 167, low-rank constructions.



The new carrier occurs in the original candidate; it is not an abstract extra tensor factor.

At rank two, write $q = 2m + 1$, $m \geq 1$, and use

$$J = U_K^1 U_A^{m+1}, \qquad P = U_K^1 U_A^m.$$

Then $J \triangleleft P$, $[P : J] = 2$, and $P \cap (K^\times J) = J$. The conjugate character ratio $\lambda = \theta^g/\theta$ is nontrivial and sign-valued. With $H_0 = \ker \lambda$,

$$P/H_0 \simeq \mathbb{F}_2^2, \qquad E = \mathrm{Ind}_J^P \theta, \quad \dim E = 2.$$

The represented diagonal sign and off-diagonal generators anticommute, proving the nondegenerate Heisenberg structure.

Scope. The original quotient J^1/H^1 is still zero-dimensional. The new pair is a separate arithmetic assembly carrier.

36 Rank four gains a four-dimensional carrier

Source: Block 167, rank-four enlarged carrier proof.

Let E_0/F be the actual outer quadratic approximation field and $B = \mathrm{End}_{E_0}(F^4)$. With lower odd critical level $r = 3h_1 = 2m + 1$ and half leading level $a = 3h_2 > r$, define

$$P = J\,U_B^m.$$

The filtered-centralizer estimate proves this is a compact group with J normal of index two. It does not assume $E_0 \subset K$.

For a new coset representative u, the ratio $\lambda = \theta^u/\theta$ on the actual H^1 is nontrivial and sign-valued. Setting $H_0 = \ker \lambda$ gives


$$P/H_0 \simeq \mathbb{F}_2^4, \qquad E = \mathrm{Ind}_J^P \eta, \quad \dim E = 4.$$

Its represented commutators are scalar; the quotient has exponent two; irreducibility and the exact scalar preimage prove nondegeneracy. The finite scalar center can exceed μ_2.

Scope. No literal extraspecial quotient is claimed. Full character values and marked attachment are retained through the actual finite extension.

37 Induction in stages recovers the same candidate

Source: Block 167, low-rank recovery proof.



The noncompact extension Λ is fixed before forming Σ.

At both low ranks, $K^\times$ normalizes P. Put

$$\mathcal{N} = K^\times J, \quad \mathcal{M} = K^\times P, \qquad \Sigma = \mathrm{Ind}_{\mathcal{N}}^{\mathcal{M}} \Lambda.$$

The subgroup $\mathcal{N}$ is normal of index two, $\Sigma|_P \simeq E$, and Σ is irreducible. Induction in stages proves

$$\pi \simeq \mathrm{Ind}_{\mathcal{M}}^{G} \Sigma.$$

Reading the diagram. The parameter, full character, original canonical beta-extension data, central character, and noncompact scalar remain those of the original candidate. The inducing carrier changes.

38 Compact recovery must remember the noncompact root

Source: Block 167, enlarged-carrier contract.

Let $X = \mathrm{Ind}_J^G \eta$ and retain $A_0 = \Lambda(\varpi_K)$. Define the normalized shift

$$(Tf)(g) = A_0^{-1} f(\varpi_K g).$$

The original candidate is the quotient

$$X/(T-1)X \simeq \pi.$$

This does not split before central normalization: a vector of X has finite support modulo compact J, whereas a scalar uniformizer shifts determinant valuations. It has no nonzero eigenvector with the required central eigenvalue.

After imposing the already fixed central character, the relevant quotient is cyclic of order N. Then $T^N = 1$ and the finite average


$$e_\Lambda = \frac{1}{N} \sum_{i=0}^{N-1} T^i$$

selects the same extension. The operator acts on a complex representation space, where division by N is valid.

Scope. This is a different recovery mechanism from a finite sector inside the original η. The root is retained, not reconstructed from the compact packet alone.

39 Central completion is a construction choice

Source: Blocks 166–167, full-character completion.



This diagram starts with an already constructed compatible arithmetic host.

The full character is trivial on U_F^1. Residue data and the noncompact scalar can then match a prescribed smooth central quasicharacter ω trivial on U_F^1, when completion choices are free or already compatible. For $F = \mathbb{F}_2((t))$, the residue unit group is trivial, so these covered central characters are unramified. An Nth root in $\mathbb{C}^\times$ fixes the action required by $\omega(t)$. All choices are recorded before invoking H5.

Scope. An incompatible immutably supplied completion is not silently replaced. Nor is an arbitrary one-factor wild determinant assumed trivial. The source central character must belong to the covered class.

Reading the diagram. A root choice completes an existing candidate. It does not prove the general realization conjecture or the same candidate's pair-factor equalities.

40 Why nontrivial stabilizer actions need another step

Source: Block 166, stabilizer contract.

For the current totally ramified degree-p^r maximal-type host, character-preserving normalizers have

$$N/(F^\times J^1) \simeq \mathbb{Z}/p^r\mathbb{Z}.$$

A native action of a finite prime-to-p group C on the quotient therefore factors through a p-group and is trivial.

The kernel of the action of center-fixing scalar-extension automorphisms on W is $\mathrm{Hom}(W, \mathbb{C}^\times)$, an elementary abelian p-group. Hence the full native scalar-extension action is trivial as well. Honest implementers on an irreducible Heisenberg carrier are scalar by Schur's lemma.

Scope. This is an obstruction to the native inner-normalizer route for this host. It does not refute the original finite contract or Conjecture 66. Original breakwise H2 may supply a genuinely nontrivial symplectic action on separate actual type targets. Transporting abstract operators does not by itself identify a common arithmetic action.

41 Genuine scalar phases can descend from a finite cover

Source: Block 166, scalar native attachment.

There is a proved extension beyond $C = 1$ when the prime-to-p stabilizer acts trivially on the source extension and its honest operators on the entire rank-N source carrier are $\chi(c)I_N$ in the actual inertia image.

Under the same-source determinant central-character condition, the required scalar phase can be implemented by actual central matrices after a finite cover $\widetilde{C} \twoheadrightarrow C$. The kernel acts trivially on the whole recovered candidate, so the represented action descends to C.

Reading the diagram. A cover may be needed even when the represented operator is a scalar. A direct matrix homomorphism from C need not exist.

Scope. The theorem requires a genuine inertia phase on the whole carrier, the fixed reciprocity convention, and compatible completion. It does not apply merely because an arbitrary abstract phase on a multiplicity subspace is scalar. It does not change a fixed canonical beta extension to force a desired action.

42 Numerical matching does not construct realization

Source: Blocks 162, 166–167, profile and capacity audits.

For compact $J \subset \mathrm{GL}_n(F)$, the homomorphism $v_F \circ \det$ has finite image in $\mathbb{Z}$, hence image zero. An unramified character

$$\nu_a(g) = a^{v_F(\det g)}$$


is therefore trivial on J. The compact packet restrictions of π and $\pi \otimes \nu_a$ agree, while the central character changes by a^n on a scalar uniformizer. Choosing $a^n \neq 1$ gives a different atom.

Block 167 also records a rank-eight skipped-degree $8 \to 2$ host with the same torsion, Swan, and self-pair conductor as the full tower, but quotient dimension 4 rather than 6.

Reading the diagram. The finite packet, numerical profiles, and noncompact normalization contain different information. No one of these alone is a substitute for an arithmetic witness preserving the full declared datum.

43 Conjecture 66 isolates the missing existence claim

Source: Historical Theorem 16.2; current realization contract.



The upper existence arrow is conjectural. The lower comparison follows conditionally.

Conjecture 66 (arithmetic packet realization). Fix the declared class $\mathcal{A}$ of wild atoms and the paper's normalizations. For every $\rho \in \mathcal{A}$ with supplied H1–H4 assembled finite datum d_ρ, there is one actual admissible supercuspidal representation π of $\mathrm{GL}_{\dim \rho}(F)$ and an arithmetic realization witness $R(d_\rho, \pi)$ recovering the packet and preserving its declared stabilizer action, central-extension and phase data, and compatible transports.

Here R includes actual arithmetic inducing data and explicit comparison and recovery. It is not merely the existence of abstract operators with the right commutators.

Scope. The conjecture is restricted to the declared admitted class. It does not assert that arbitrary coefficient targets, arbitrary partial hosts, or independently conflicting completion choices can all be extended.

44 H5 compares this same actual candidate

Source: Contract audit 162; conditional comparison proposition.

Suppose π is an actual admissible supercuspidal representation of the correct rank and $R(d_\rho, \pi)$ is supplied. If H5 gives equality of its required pair local factors with those of $\mathrm{LLC}(\rho)$ on the stipulated characterizing twist family, with the fixed normalizations, then

$$\pi \simeq \mathrm{LLC}(\rho).$$

The proof applies the supplied local-factor characterization to these two actual representations. It does not need existence for other diagrams or H5 for every alternative completion.

There is a logical difference between

$$\forall \pi\, [R(d, \pi) \Rightarrow \mathrm{H5}(\pi)] \quad \text{and} \quad \exists \pi\, [R(d, \pi) \wedge \mathrm{H5}(\pi)].$$

The first statement can be vacuous. The second includes an actual witness. Choosing the established reference atom only supplies a profile-matching object; it leaves its recovery of d unproved.

Reading the diagram. H5 remains a conditional premise about the same independently constructed candidate. It is not renamed as an unconditional factor-comparison theorem.

45 What the restricted realizations establish

Source: Blocks 166–167, final composition reviews.

- Actual totally ramified parameter fields, literal defining sequences, full simple characters, and covered central completions exist throughout the stated odd-prime and dyadic families.
- Under the one-vertex, $C = 1$, supplied full-character scalar-pushout contract, the whole marked packet has an actual arithmetic assembly carrier and same-candidate recovery.
- In the dyadic low ranks, the actual enlarged carriers restore capacity without changing the original maximal-simple quotient.
- Genuine scalar inertia phases have the separately stated finite-cover descent result.

The remaining general obligations include nontrivial prescribed symplectic stabilizer actions on a common arithmetic candidate, general sources and transport diagrams, comparison with independently fixed canonical native markings, and numerical layer placement when the declared contract requires it.

Scope. The constructions are substantial restricted realization theorems. Their scopes do not close universal Conjecture 66, and their profile identities do not prove H5.

46 Exact computation checks a bounded identity

Source: Block 167, computation contract; final verification records.

The rank-eight run for $h = (1, 3, 5)$ produces an actual matrix with 25 nonzero Laurent-polynomial entries. Its checked leading and adjacent-correction grades are

$$-60, \quad -18, \quad -3.$$

It checks normalized lifts, actual matrix identities, native traces 1, and amplified ambient traces $0, 0, 1$. The odd $p = 3$ example $h = (1, 2, 4)$ has levels $144, 24, 4$ and native trace zero.

These finite outputs have contracts, scripts, input pins, results, and execution records in the release. The critical exponents, simplicity, full-character existence, and arbitrary-length statements use the separate ordinary mathematical arguments.

Rebuilding the synchronized readers. From the checkpoint root, run `python tools/build_readers.py --output PATH`, with PATH outside the checkpoint. The builder stages every style, include and figure dependency, compiles the readers, and resolves the cross-document destinations. The vector plates are preserved source assets. Their Python generator remains available, but figure regeneration is not claimed by this reader build.

Scope. A bounded matrix run cannot establish the universal realization claim. No fresh Lean certificate for Blocks 166–167 is asserted by this guide.

47 Where the complete arguments are retained

Source: Release source collection.

Topic	Retained source anchor
Finite tame and wild route	Original manuscript, Sections 2–16. The release traditional paper preserves the route with the Frobenius correction and explicit realization boundary.
Original endpoint contract	`ORIGINAL_CONTRACT_162.md`: common-type versus separate-type readings, same-candidate H5, and exact witness requirement.
Odd arithmetic constructor	`PROOF166.md`; `TRACE_ZERO_AS_TOWER166.md`; `RECURSIVE_TOWER166.md`.
Full-character induction	`TRACE_ZERO_CHARACTER_INDUCTION166.md`; `DYADIC_FULL_CHARACTER167.md`.
Dyadic construction	`PROOF167.md`; `DYADIC_AS_TOWER167.md`; `DYADIC_CAPACITY167.md`.
Marked finite extensions	`DYADIC_MARKINGS167.md`, including native-refinement conditions and whole-subgroup transport.
Enlarged low ranks	`ENLARGED_CARRIER_CONTRACT167.md`; the rank-two, rank-four, and induction-in-stages notes.
Stabilizer boundary	`STABILIZER_CONTRACT166.md`; `SCALAR_NATIVE_ATTACHMENT166.md`.
Reviews and computation	The accompanying audit directories, exact computation contracts, source pins, and execution manifests.

The .md materials accompany the PDFs. Historical optimistic filenames or superseded reviews are preserved as history and do not override the final statements and scope records.