Projector-Resolved Seed Calculus IV

Cyclic Analytic Functions, Reciprocal Families, and Spectral Zero Data

Komorebi P.

Version 1.0.0 — proofread draft — September 5, 2026

Abstract

Fix a positive integer N. Root-of-unity projection of the exponential produces residue-class sections that, for $\lambda \neq 0$, form a basis for $y^{(N)} = \lambda^N y$. The signed sections for $y^{(N)} = -y$ yield the denominator D_N and its reduced form Δ_N. We derive reciprocal coefficient polynomials, Bell and Newton recurrences, an additive parameter law, and residue-section Appell and theta families. For $N \geq 2$, Δ_N has order $1/N$. We specialize classical two-term zero theory for exponential sums to the regular exponent polygon. In this equal-weight family, conjugation symmetry combined with unique near-model localization places all sufficiently large zeros exactly on the symmetry rays, and direct expansion gives their leading displacement and derivative sign. The classical perimeter law yields zero-counting asymptotics with explicit leading constants and bounded error. These results give genus-zero products, the standard diagonal Fredholm representation, zero-power-sum formulas for reciprocal and logarithmic coefficients, nearest-zero coefficient asymptotics, and meromorphic continuation of the symmetry-orbit zero zeta by comparison with the Hurwitz zeta function. Complex powers are interpreted either formally or through a fixed logarithm on a zero-free domain. The paper treats only this cyclic analytic family. Arithmetic interpolation, general nonlinear trees, general resolvent theory, and abstract completion are reserved for Papers IX, III, VI, and X.

Formalization and evidence boundary. The released Paper IV Lean contract covers all 68 theorem-like manuscript results under their stated hypotheses. The accompanying v1.0.0 theorem map and formalization ledger identify the declarations and their scopes. Imported standard fact 11.1 remains explicitly imported and is not needed by the checked growth-order proofs. Open problems remain open. Complex powers retain their specified logarithms and domains; conditional asymptotic statements retain their assumptions. Finite computations are regression evidence. Formal verification checks the encoded specifications and does not establish historical priority or replace independent mathematical review.

Contents

1 Introduction **3**

2 A map of the construction **4**

3 Cyclic projection of analytic functions **5**

4 The transfer exponential basis **6**

5 Signed cyclic sections and the distinguished denominator **8**

6 A finite algebra and matrix model **10**

7 Reciprocal powers and parameter polynomials **12**

8 Logarithmic derivatives and Riccati equations **15**

9 Residue-section reciprocals and Appell systems **16**

10 Scaled reciprocal coupling and theta polynomials **19**

11 Growth and the geometry of the large zeros **21**

12 Hadamard products and zero power sums **27**

13 Coefficient asymptotics from the nearest zeros **31**

14 Zero zeta functions **33**

15 Scope boundaries and further questions **36**

16 Formalization ledger for version 1.0.0 **36**

17 Conclusion **38**

1 Introduction

For a power series

$$F(z) = \sum_{m \geq 0} a_m z^m$$

and a positive integer N, the coefficients may be separated according to the residue of m modulo N. Root-of-unity averaging gives exact projection operators for this separation. This elementary construction is usually called a multisection or a root filter. It appears throughout the classical theory of generating functions and special functions. Background on multisections, hypergeometric functions, and related coefficient systems may be found in [Rai60, AAR99, Com74].

Papers I and II developed typed seed grammars, exact projector packets, and finite residue-projector algebra, while Paper III treated typed edge operators and exact coordinate transport [P.26a, P.26b, P.26c]. The root filter used below is the finite cyclic scalar specialization of that earlier projector calculus. The signed companion matrix is likewise a concrete cyclic instance, not a replacement for Paper III's general typed-edge theory.

This article follows the consequences of applying the cyclic projections to the exponential function. The projected exponentials form a finite differential system. After one sign is inserted around the derivative cycle, they form a basis for

$$y^{(N)} = -y.$$

The section with initial value one is

$$\mathsf{D}_N(z) = \sum_{n \geq 0} (-1)^n \frac{z^{Nn}}{(Nn)!}.$$

Its reciprocal powers produce polynomial coefficient families. Its logarithmic derivative produces a scalar higher Riccati identity. The same reciprocal construction applied to every exponential section gives normalized residue-section functions and ordinary Appell polynomial sequences. These cyclic exponential sections and their matrix addition laws belong to the generalized-hyperbolic-function literature [Sch92, Mul05]. Appell and binomial-type sequences are discussed in [Rom84, Com74].

The second half of the paper concerns the zeros of D_N. The finite exponential sum

$$\mathsf{D}_N(z) = \frac{1}{N} \sum_{\eta^N = -1} e^{\eta z}$$

has a regular polygon of exponents. The general relation between an exponent polygon and the zero strips of an exponential sum goes back to Pólya and Langer. Later developments include the work of Moreno and the treatment in Levin's monograph [Pól20, Lan31, Mor73, Lev80]. Bleher and Mallison give a particularly close general result: for convex-polygon exponential sums they locate all sufficiently large zeros in two-term strips and prove exponential localization and simplicity [BM06]. The regular polygon and conjugation symmetry of the present family specialize that theory to N exact rays. We give a self-contained specialization because its notation is used throughout the product and zeta theory, and we compute the leading displacement term for this family. The resulting counting formulas have explicit leading constants and bounded error.

The rotational symmetry is more useful after the change of variable $w = z^N$. There is an entire function

$$\Delta_N(w) = \sum_{n \geq 0} (-1)^n \frac{w^n}{(Nn)!}$$

with

$$\mathsf{D}_N(z) = \Delta_N(z^N).$$

For $N \geq 2$, the order of Δ_N is $1/N$. Its Hadamard product therefore has genus zero. The logarithm of this product identifies the transfer tangent coefficients with power sums of the reduced zeros. The large zero formula also compares the reduced zero zeta with a Hurwitz zeta function. Standard material on entire functions, canonical products, and exponential sums appears in [Boa54, Lev80, Tit39]. We use singularity analysis in the form described in [FS09] and the classical continuation of the Hurwitz zeta function from [Apo76].

The cyclic exponential sections and their matrix addition laws have a classical literature. The tools used below—Appell and binomial-type sequences, canonical products, singularity analysis, and Fredholm determinants—are also standard. Our aim is to organize this symmetric specialization and link its reciprocal coefficients to reduced-zero data. No historical priority is asserted for the component constructions.

2 A map of the construction

Two variables appear throughout the paper. The variable z carries the differential equation and the rotational symmetry. The reduced variable

$$w = z^N$$

records one representative from each rotational orbit. Confusing these variables hides the main simplification. The denominator D_N has order one as a function of z, while Δ_N has order $1/N$ as a function of w.

The analytic objects are introduced in the following order. First,

$$\mathsf{E}_{N,j}(z) = \sum_{n\geq 0} \frac{z^{Nn+j}}{(Nn+j)!}$$

is the residue j part of the exponential. Second,

$$\mathsf{D}_{N,j}(z) = \sum_{n\geq 0} (-1)^n \frac{z^{Nn+j}}{(Nn+j)!}$$

is the signed basis for $y^{(N)} = -y$. The section $\mathsf{D}_N = \mathsf{D}_{N,0}$ is the denominator used in the reciprocal family

$$\mathsf{S}_N^{(r)} = \mathsf{D}_N^{-r}.$$

Set $\mathsf{T}_N = -\mathsf{D}_N'/\mathsf{D}_N$; equivalently, T_N is the logarithmic derivative of $\mathsf{S}_N^{(1)}$.

A different reciprocal family comes from normalizing each exponential section at the origin. The function

$$\mathsf{C}_{N,j}(z) = \frac{z^j/j!}{\mathsf{E}_{N,j}(z)}$$

is a unit series in z^N. Its powers, after multiplication by e^{xz}, give Appell polynomials. The theta family couples a scaled argument with a reciprocal order.

The zero theory begins only after these algebraic relations have been established. The factorization

$$\mathsf{D}_N(z) = \Delta_N(z^N)$$

turns every orbit of N zeros in the z plane into one zero in the w plane. The large zeros in the z plane are located by two dominant exponentials. The zeros in the w plane then give a genus zero product. The logarithm of that product returns to the coefficient theory through

$$\text{zero power sums} = \text{logarithmic coefficients}.$$

This equality is the bridge between the first and second halves of the paper.

All powers with a complex parameter are first defined as formal series at the origin. To represent this germ analytically, use a simply connected zero-free region containing the origin and the logarithm normalized to vanish there. On other regions, specify a branch separately. Integer reciprocal powers are globally meromorphic and do not require a branch choice.

The main statements fall into five groups. The finite algebra model in section 6 explains the derivative cycles and addition laws through one exponential identity. The coefficient theory in section 7 gives two complementary recurrences, one from the original denominator and one from its logarithm. The Appell theory in section 9 supplies both formal differential-operator formulas and a convolution law in the space and order parameters. The zero analysis in section 11 specializes classical two-term exponential-sum theory, uses symmetry to place the uniquely localized large zeros on the rays, and computes a leading correction. The product theory in section 12 gives the standard diagonal Fredholm encoding of the genus-zero product.

The article keeps formal and analytic statements separate. A formal identity needs only the unit constant term of the relevant series. A holomorphic complex power also needs a zero-free domain and a chosen logarithm. A meromorphic coefficient asymptotic is stated only for positive integral reciprocal order unless an additional branch domain is specified.

3 Cyclic projection of analytic functions

Fix $N \geq 1$ and write

$$\omega = e^{2\pi i/N}.$$

Let Ω be a disk centered at the origin, with the whole plane allowed, and assume that Ω is invariant under multiplication by ω. For a holomorphic function F on Ω, define

$$(\mathcal{B}_{N,j}F)(z) = \frac{1}{N}\sum_{\ell=0}^{N-1} \omega^{-j\ell} F(\omega^{\ell} z), \qquad 0 \leq j < N.$$

The same formula is valid for formal power series.

Theorem 3.1 (Root filter)**.** *If*

$$F(z) = \sum_{m\geq 0} a_m z^m,$$

then

$$\mathcal{B}_{N,j}F(z) = \sum_{m\equiv j \bmod N} a_m z^m.$$

The operators satisfy

$$\mathcal{B}_{N,i}\mathcal{B}_{N,j} = \delta_{ij}\mathcal{B}_{N,j}$$

and

$$\sum_{j=0}^{N-1} \mathcal{B}_{N,j} = I.$$

Proof. Substitution of the power series gives

$$\mathcal{B}_{N,j}F(z) = \sum_{m\geq 0} a_m z^m \left(\frac{1}{N}\sum_{\ell=0}^{N-1} \omega^{(m-j)\ell}\right).$$

The inner average is one when $m \equiv j \bmod N$ and zero otherwise. The operator identities follow from the disjoint residue classes. □

Proposition 3.2 (Covariance characterization)**.** *A holomorphic function F belongs to the range of $\mathcal{B}_{N,j}$ if and only if*

$$F(\omega z) = \omega^j F(z).$$

Proof. Every monomial selected by $\mathcal{B}_{N,j}$ has the stated covariance. Conversely, write $F = \sum_k \mathcal{B}_{N,k}F$. Replacing z by ωz gives

$$\sum_k \omega^k \mathcal{B}_{N,k}F = \omega^j \sum_k \mathcal{B}_{N,k}F.$$

Application of $\mathcal{B}_{N,k}$ gives

$$(\omega^k - \omega^j)\mathcal{B}_{N,k}F = 0.$$

The powers $1, \omega, \ldots, \omega^{N-1}$ are distinct, so $\mathcal{B}_{N,k}F = 0$ for $k \neq j$. □

Multiplication is not diagonal in these coordinates. The residue labels add modulo N; this is the scalar cyclic instance of the packet product from Paper I.

Theorem 3.3 (Cyclic product law)**.** *If F and G are holomorphic on the same rotation-invariant domain, then*

$$\mathcal{B}_{N,k}(FG) = \sum_{\substack{0 \le i,j < N \\ i+j \equiv k \bmod N}} (\mathcal{B}_{N,i}F)(\mathcal{B}_{N,j}G).$$

Proof. Write $F = \sum_i F_i$ and $G = \sum_j G_j$, where $F_i = \mathcal{B}_{N,i}F$ and $G_j = \mathcal{B}_{N,j}G$. Their covariance gives

$$(F_i G_j)(\omega z) = \omega^{i+j} F_i(z) G_j(z),$$

so $F_i G_j$ belongs to the packet labelled by $i + j$ modulo N. Expanding the finite product and applying $\mathcal{B}_{N,k}$ gives the formula. □

Remark 3.4. Every identity in this section holds formally. In the holomorphic setting, the finite projector sums preserve local uniform convergence. No choice of branch is involved.

4 The transfer exponential basis

Definition 4.1. For $\lambda \in \mathbb{C}$ and $0 \le j < N$, define

$$\mathsf{E}_{N,j}(\lambda z) = \mathcal{B}_{N,j}(e^{\lambda z}) = \sum_{n \ge 0} \frac{(\lambda z)^{Nn+j}}{(Nn+j)!}.$$

Theorem 4.2 (Differential cycle and initial data)**.** *For $1 \le j < N$,*

$$\frac{d}{dz}\mathsf{E}_{N,j}(\lambda z) = \lambda \mathsf{E}_{N,j-1}(\lambda z),$$

while

$$\frac{d}{dz}\mathsf{E}_{N,0}(\lambda z) = \lambda \mathsf{E}_{N,N-1}(\lambda z).$$

Thus

$$\frac{d^N}{dz^N}\mathsf{E}_{N,j}(\lambda z) = \lambda^N \mathsf{E}_{N,j}(\lambda z).$$

For $0 \le k < N$,

$$\left.\frac{d^k}{dz^k}\mathsf{E}_{N,j}(\lambda z)\right|_{z=0} = \delta_{kj}\lambda^j.$$

If $\lambda \neq 0$, the N functions $\mathsf{E}_{N,j}(\lambda z)$ form a basis of the solution space of $y^{(N)} = \lambda^N y$.

Proof. Termwise differentiation gives the cyclic derivative formulas. One full cycle contributes λ^N. The first nonzero term of $\mathsf{E}_{N,j}(\lambda z)$ is $(\lambda z)^j/j!$, which gives the initial data. The initial value vectors are linearly independent when $\lambda \neq 0$. The solution space of an order N constant coefficient equation has dimension N, so these functions form a basis. □

Corollary 4.3 (Wronskian of the transfer basis)**.** *Assume $N \geq 2$. For $\lambda \neq 0$, the Wronskian of the ordered basis*

$$\mathsf{E}_{N,0}(\lambda z), \ldots, \mathsf{E}_{N,N-1}(\lambda z)$$

is the constant

$$\lambda^{N(N-1)/2}.$$

Proof. At the origin, the derivative matrix is diagonal with entries

$$1, \lambda, \ldots, \lambda^{N-1}.$$

For an explicit determinant proof, write each basis function as a finite sum of exponentials with exponents $a_\ell = \lambda\omega^\ell$. Its derivative matrix factors as

$$W(z) = V\ \mathrm{diag}(e^{a_\ell z})\, C, \qquad V_{k\ell} = a_\ell^k,$$

where C is the constant matrix of Fourier weights. Thus

$$\det W(z) = e^{z\sum_\ell a_\ell} \det W(0) = \det W(0),$$

because $N \geq 2$ implies $\sum_\ell \omega^\ell = 0$. The product of the diagonal initial values is the stated power of λ. □

Corollary 4.4 (Addition law)**.** *For $0 \leq k < N$,*

$$\mathsf{E}_{N,k}(\lambda(z+w)) = \sum_{\substack{0\leq i,j<N \\ i+j\equiv k \bmod N}} \mathsf{E}_{N,i}(\lambda z)\mathsf{E}_{N,j}(\lambda w).$$

Proof. Expand $e^{\lambda(z+w)} = e^{\lambda z}e^{\lambda w}$ as a bivariate power series. A term $z^a w^b$ contributes to the kth section exactly when $a + b \equiv k \bmod N$. Grouping first by $i \equiv a \bmod N$ and $j \equiv b \bmod N$ gives the displayed sum. □

The basis also has a scalar hypergeometric form.

Theorem 4.5 (Hypergeometric form)**.** *Let*

$$A_{N,j} = \left\{ \frac{j+m}{N} : 1 \leq m \leq N \text{ and } j+m \neq N \right\}.$$

Then

$$\mathsf{E}_{N,j}(\lambda z) = \frac{(\lambda z)^j}{j!} {}_0F_{N-1}\left(\begin{matrix} - \\ A_{N,j} \end{matrix}, \frac{(\lambda z)^N}{N^N} \right).$$

Proof. The multiplication formula for the gamma function gives

$$(Nn+j)! = j!N^{Nn}n! \prod_{a\in A_{N,j}} (a)_n.$$

Indeed, the N Pochhammer factors supplied by the multiplication formula are $((j+m)/N)_n$ for $1 \leq m \leq N$; the excluded factor is $((j+N-j)/N)_n = (1)_n = n!$. Insertion into the defining series gives the stated hypergeometric expansion. □

Example 4.6. For $N = 1$, the only section is $e^{\lambda z}$. For $N = 2$,

$$\mathsf{E}_{2,0}(z) = \cosh z, \qquad \mathsf{E}_{2,1}(z) = \sinh z.$$

These are classical instances of the same cyclic basis.

5 Signed cyclic sections and the distinguished denominator

Definition 5.1. For $0 \le j < N$, define

$$\mathsf{D}_{N,j}(z) = \sum_{n\ge 0} (-1)^n \frac{z^{Nn+j}}{(Nn+j)!}.$$

The distinguished signed denominator is

$$\mathsf{D}_N(z) = \mathsf{D}_{N,0}(z).$$

Fix any $\rho \in \mathbb{C}$ with $\rho^N = -1$.

Proposition 5.2 (Relation with the transfer exponential)**.** *For every j,*

$$\mathsf{D}_{N,j}(z) = \rho^{-j} \mathsf{E}_{N,j}(\rho z).$$

Proof. The coefficient of z^{Nn+j} on the right is

$$\rho^{-j}\rho^{Nn+j}\frac{1}{(Nn+j)!} = (-1)^n \frac{1}{(Nn+j)!}.$$

□

Theorem 5.3 (Signed derivative cycle)**.** *For $1 \le j < N$,*

$$\mathsf{D}'_{N,j} = \mathsf{D}_{N,j-1},$$

and

$$\mathsf{D}'_{N,0} = -\mathsf{D}_{N,N-1}.$$

Every signed section satisfies

$$\mathsf{D}^{(N)}_{N,j} = -\mathsf{D}_{N,j}.$$

The N signed sections form the basis of $y^{(N)} = -y$ determined by

$$\mathsf{D}^{(k)}_{N,j}(0) = \delta_{jk}$$

for $0 \le j, k < N$.

Proof. The derivative formulas follow directly from the series. The first nonzero term of $\mathsf{D}_{N,j}$ is $z^j/j!$, so the initial value matrix is the identity. The basis statement follows from the dimension of the solution space. □

Corollary 5.4 (Wronskian of the signed basis)**.** *Assume $N \ge 2$. The Wronskian of*

$$\mathsf{D}_{N,0}(z), \dots, \mathsf{D}_{N,N-1}(z)$$

is one.

Proof. The derivative matrix at the origin is the identity. The determinant factorization in the proof of corollary 4.3 applies with $a_\ell = \rho\omega^\ell$ and with the signed-section Fourier weights. Since $\sum_\ell a_\ell = 0$, the determinant is constant and equals one. □

Proposition 5.5 (Twisted addition law)**.** *For $0 \le k < N$,*

$$\mathsf{D}_{N,k}(z+w) = \sum_{\substack{0\le i,j<N \\ i+j\equiv k \bmod N}} (-1)^{\lfloor (i+j)/N \rfloor} \mathsf{D}_{N,i}(z)\mathsf{D}_{N,j}(w).$$

Proof. Use proposition 5.2 and corollary 4.4. The factor introduced by a pair (i,j) is

$$\rho^{i+j-k}.$$

Since $0 \le i+j \le 2N-2$ and $i+j \equiv k \bmod N$, this factor is one when $i+j=k$ and -1 when $i+j=k+N$. □

Theorem 5.6 (Finite exponential representation)**.** *Let*

$$\mathcal{R}_N = \{\eta \in \mathbb{C} : \eta^N = -1\}.$$

Then

$$\mathsf{D}_N(z) = \frac{1}{N} \sum_{\eta\in\mathcal{R}_N} e^{\eta z}.$$

More generally,

$$\mathsf{D}_{N,j}(z) = \frac{\rho^{-j}}{N} \sum_{\ell=0}^{N-1} \omega^{-j\ell} e^{\rho\omega^\ell z}.$$

Proof. The second formula is the root filter expression for $\rho^{-j}\mathsf{E}_{N,j}(\rho z)$. The first is the case $j=0$. □

Definition 5.7. Define the reduced denominator

$$\Delta_N(w) = \sum_{n\ge 0} (-1)^n \frac{w^n}{(Nn)!}.$$

Then

$$\mathsf{D}_N(z) = \Delta_N(z^N).$$

Corollary 5.8 (Symmetries)**.** *The denominator satisfies*

$$\mathsf{D}_N(\omega z) = \mathsf{D}_N(z)$$

and

$$\mathsf{D}_N(\overline{z}) = \overline{\mathsf{D}_N(z)}.$$

Its nonzero zeros occur in orbits under multiplication by the Nth roots of unity and under complex conjugation.

Proof. The first identity follows from the powers z^{Nn}. The second follows from the real coefficients. Both symmetries preserve the zero set and multiplicity. □

Example 5.9. The first denominators are

$$\mathsf{D}_1(z) = e^{-z}, \qquad \mathsf{D}_2(z) = \cos z.$$

For $N=3$,

$$\mathsf{D}_3(z) = \frac{1}{3}\left(e^{-z} + 2e^{z/2}\cos\left(\frac{\sqrt{3}z}{2}\right)\right).$$

For $N=4$,

$$\mathsf{D}_4(z) = \cos\left(\frac{z}{\sqrt{2}}\right)\cosh\left(\frac{z}{\sqrt{2}}\right).$$

6 A finite algebra and matrix model

The derivative and addition laws can be read from one finite algebra identity. This formulation also identifies the exponent polygon as the spectrum of a signed cyclic shift.

Definition 6.1. Let

$$\mathfrak{A}_N^+ = \mathbb{C}[\xi]/(\xi^N - 1)$$

and

$$\mathfrak{A}_N^- = \mathbb{C}[\xi]/(\xi^N + 1).$$

Every element has a unique representative of degree below N.

Theorem 6.2 (Finite algebra exponential identities)**.** *In $\mathfrak{A}_N^+$ one has*

$$e^{\lambda z \xi} = \sum_{j=0}^{N-1} \mathsf{E}_{N,j}(\lambda z)\xi^j.$$

In $\mathfrak{A}_N^-$ one has

$$e^{z\xi} = \sum_{j=0}^{N-1} \mathsf{D}_{N,j}(z)\xi^j.$$

Differentiation and multiplication of these identities recover the cyclic derivative systems and the two addition laws.

Proof. Expand the exponential and write every exponent uniquely as $Nn + j$. In $\mathfrak{A}_N^+$, the relation $\xi^{Nn+j} = \xi^j$ gives the transfer sections. In $\mathfrak{A}_N^-$, the relation $\xi^{Nn+j} = (-1)^n \xi^j$ gives the signed sections.

Differentiation multiplies by $\lambda\xi$ in the first algebra and by ξ in the second. Reduction of the product by the defining relation gives the derivative cycles. The identity

$$e^{(z+w)\xi} = e^{z\xi} e^{w\xi}$$

gives the addition laws after comparison of the coefficients of $1, \xi, \ldots, \xi^{N-1}$. □

Definition 6.3. Let J_N be the signed cyclic shift on the standard basis $e_0, \ldots, e_{N-1}$ of $\mathbb{C}^N$, defined by

$$J_N e_j = e_{j+1} \quad (0 \le j < N-1), \qquad J_N e_{N-1} = -e_0.$$

Then $J_N^N = -I$.

Theorem 6.4 (Companion matrix realization)**.** *Put*

$$\mathbf{D}_N(z) = \begin{pmatrix} \mathsf{D}_{N,0}(z) \\ \mathsf{D}_{N,1}(z) \\ \vdots \\ \mathsf{D}_{N,N-1}(z) \end{pmatrix}.$$

Then

$$\mathbf{D}_N(z) = e^{zJ_N} e_0$$

and

$$\mathsf{D}_N(z) = \frac{1}{N} \operatorname{tr}(e^{zJ_N}).$$

The eigenvalues of J_N are the N roots of -1, each with multiplicity one.

Proof. The vector $e^{zJ_N}e_0$ has derivative $J_Ne^{zJ_N}e_0$ and initial value e_0. Its components therefore satisfy the same signed derivative cycle and initial data as $\mathbf{D}_N$. Uniqueness for the linear system gives the first identity.

The vectors

$$e_0, J_Ne_0, \dots, J_N^{N-1}e_0$$

are the standard basis. Thus e_0 is cyclic for J_N, so the minimal polynomial of J_N has degree N. Since $J_N^N = -I$, that minimal polynomial divides $x^N + 1$ and therefore equals $x^N + 1$. The characteristic polynomial has the same degree and is divisible by the minimal polynomial, so it too equals $x^N + 1$. This polynomial has distinct roots. Consequently J_N is diagonalizable and its spectrum consists of all N roots of -1, each once. Hence

$$\operatorname{tr}(e^{zJ_N}) = \sum_{\eta^N=-1} e^{\eta z} = N\mathsf{D}_N(z)$$

by theorem 5.6. □

Corollary 6.5 (Entries of the fundamental matrix)**.** *For $0 \le p, q < N$, let d_{pq} be the member of $\{0, \dots, N-1\}$ congruent to $p - q$ modulo N. Put*

$$\varepsilon_{pq} = \begin{cases} 1, & p \ge q, \\ -1, & p < q. \end{cases}$$

Then

$$\left(e^{zJ_N}\right)_{pq} = \varepsilon_{pq}\mathsf{D}_{N,d_{pq}}(z).$$

Thus every entry of the fundamental matrix is one of the signed sections, with the sign recording whether the cyclic path from q to p crosses the signed edge.

Proof. For every nonnegative integer n,

$$J_N^{Nn+d_{pq}}e_q = (-1)^n\varepsilon_{pq}e_p.$$

Only powers whose exponent is congruent to d_{pq} modulo N contribute to the (p, q) entry of the exponential. Therefore

$$\left(e^{zJ_N}\right)_{pq} = \varepsilon_{pq}\sum_{n\ge 0}(-1)^n\frac{z^{Nn+d_{pq}}}{(Nn+d_{pq})!}.$$

This is the stated formula. □

Corollary 6.6 (Addition and inversion laws for the companion matrix)**.** *Assume $N \ge 2$. The matrix*

$$U_N(z) = e^{zJ_N}$$

satisfies

$$U_N(z+w) = U_N(z)U_N(w), \qquad U_N(z)^{-1} = U_N(-z), \qquad \det U_N(z) = 1.$$

Its first column is $\mathbf{D}_N(z)$, and all of its entries are given by corollary 6.5.

Proof. The first two identities are the exponential group law. Since $\operatorname{tr} J_N = 0$,

$$\det(e^{zJ_N}) = e^{z\operatorname{tr}J_N} = 1.$$

The first column statement is theorem 6.4. □

7 Reciprocal powers and parameter polynomials

Since $\mathsf{D}_N(0) = 1$, its formal logarithm is defined at the origin. On a simply connected zero-free region containing the origin, there is a unique holomorphic branch normalized by $\log \mathsf{D}_N(0) = 0$, and its germ at the origin agrees with the formal logarithm.

Definition 7.1. For $r \in \mathbb{C}$, define

$$\mathsf{S}_N^{(r)}(z) = \mathsf{D}_N(z)^{-r} = \exp\bigl(-r \log \mathsf{D}_N(z)\bigr).$$

Write

$$\mathsf{S}_N^{(r)}(z) = \sum_{n\geq 0} P_{N,n}(r) \frac{z^{Nn}}{(Nn)!}.$$

Define the logarithmic coefficients $\kappa_{N,m}$ by

$$-\log \Delta_N(w) = \sum_{m\geq 1} \kappa_{N,m} \frac{w^m}{m}.$$

The polynomial dependence on r can be read directly from the formal logarithm. We use the complete exponential Bell polynomials B_n defined by

$$\exp\left(\sum_{m\geq 1} x_m \frac{t^m}{m!}\right) = \sum_{n\geq 0} B_n(x_1, \ldots, x_n) \frac{t^n}{n!}.$$

Theorem 7.2 (Bell polynomial formula)**.** *For every $n \geq 0$,*

$$P_{N,n}(r) = \frac{(Nn)!}{n!} B_n\bigl(r\kappa_{N,1}, r\,1!\,\kappa_{N,2}, \ldots, r\,(n-1)!\,\kappa_{N,n}\bigr).$$

For $n \geq 1$, this is a polynomial in r of exact degree n. Its leading coefficient is

$$\frac{(Nn)!}{n!(N!)^n}.$$

Proof. The identity

$$\Delta_N(w)^{-r} = \exp\left(r \sum_{m\geq 1} \kappa_{N,m} \frac{w^m}{m}\right)$$

fits the defining generating function for B_n with

$$x_m = r(m-1)!\kappa_{N,m}.$$

The coefficient of w^n is therefore the displayed Bell polynomial divided by $n!$. Multiplication by $(Nn)!$ gives $P_{N,n}(r)$. Since

$$\Delta_N(w) = 1 - \frac{w}{N!} + O(w^2),$$

we have $\kappa_{N,1} = 1/N!$. The term x_1^n gives the stated leading coefficient, which is nonzero. □

The coefficients also satisfy a triangular recurrence that does not require the logarithmic coefficients to be known first.

Theorem 7.3 (Parameter recurrence)**.** *We have $P_{N,0}(r) = 1$. For $n \geq 1$,*

$$\begin{aligned} P_{N,n}(r) = & -\sum_{k=1}^{n-1}(-1)^k \binom{Nn-1}{Nk} P_{N,n-k}(r) \\ & - r\sum_{k=1}^{n}(-1)^k \binom{Nn-1}{Nk-1} P_{N,n-k}(r). \end{aligned}$$

Proof. Set $S = \mathsf{D}_N^{-r}$. Logarithmic differentiation gives

$$\mathsf{D}_N S' + r\mathsf{D}'_N S = 0.$$

Taking the coefficient of $z^{Nn-1}/(Nn-1)!$ gives

$$\sum_{k=0}^{n-1}(-1)^k \binom{Nn-1}{Nk} P_{N,n-k}(r) + r\sum_{k=1}^{n}(-1)^k \binom{Nn-1}{Nk-1} P_{N,n-k}(r) = 0.$$

The term with $k = 0$ in the first sum is $P_{N,n}(r)$, and every other term has a smaller block index. Rearrangement gives the formula. □

Theorem 7.4 (Additive parameter law for reciprocal powers)**.** *For all $r, s \in \mathbb{C}$,*

$$\mathsf{S}_N^{(r)}(z)\mathsf{S}_N^{(s)}(z) = \mathsf{S}_N^{(r+s)}(z)$$

as a formal identity and on every common zero-free region with a fixed logarithm. Consequently,

$$P_{N,n}(r+s) = \sum_{q=0}^{n} \binom{Nn}{Nq} P_{N,q}(r) P_{N,n-q}(s).$$

Proof. Both statements follow by adding the exponents of the same logarithm and then multiplying the block exponential generating series. □

Corollary 7.5 (A sequence of binomial type)**.** *Define*

$$A_{N,n}(r) = \frac{n!}{(Nn)!} P_{N,n}(r).$$

Then

$$A_{N,n}(r+s) = \sum_{q=0}^{n} \binom{n}{q} A_{N,q}(r) A_{N,n-q}(s).$$

Thus $(A_{N,n})_{n\geq 0}$ is a polynomial sequence of binomial type.

Proof. Multiply the coefficient identity in theorem 7.4 by $n!/(Nn)!$ and simplify the factorials. □

The recurrence in theorem 7.3 uses the coefficients of the original denominator. A second recurrence uses only the logarithmic coefficients and is often shorter.

Theorem 7.6 (Logarithmic Newton recurrence)**.** *Set*

$$b_{N,n}(r) = \frac{P_{N,n}(r)}{(Nn)!}.$$

Then $b_{N,0}(r) = 1$ and for every $n \geq 1$,

$$nb_{N,n}(r) = r \sum_{k=1}^{n} \kappa_{N,k} b_{N,n-k}(r).$$

Equivalently,

$$P_{N,n}(r) = \frac{r}{n} \sum_{k=1}^{n} \frac{(Nn)!}{(N(n-k))!} \kappa_{N,k} P_{N,n-k}(r).$$

If $p_k = r\kappa_{N,k}$, then, for $n \geq 1$,

$$b_{N,n}(r) = \frac{1}{n!} \det \begin{pmatrix} p_1 & -1 & 0 & \cdots & 0 \\ p_2 & p_1 & -2 & \cdots & 0 \\ p_3 & p_2 & p_1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & -(n-1) \\ p_n & p_{n-1} & \cdots & p_2 & p_1 \end{pmatrix}.$$

Proof. Let

$$B_r(w) = \Delta_N(w)^{-r} = \sum_{n \geq 0} b_{N,n}(r) w^n.$$

Logarithmic differentiation gives

$$\frac{B_r'(w)}{B_r(w)} = r \sum_{k \geq 1} \kappa_{N,k} w^{k-1}.$$

Comparison of the coefficient of w^{n-1} proves the recurrence.

Let D_n denote the displayed determinant, with $D_0 = 1$. Expansion along the last row gives

$$D_n = \sum_{k=1}^{n} \frac{(n-1)!}{(n-k)!} p_k D_{n-k}.$$

Here the cofactor sign is cancelled by the product of the negative superdiagonal entries $-(n-k+1), \ldots, -(n-1)$. Thus $D_n/n!$ satisfies the same recurrence and initial value as $b_{N,n}(r)$. □

Corollary 7.7 (The first three reciprocal polynomials)**.** *For every $N \geq 1$,*

$$P_{N,1}(r) = r,$$

$$P_{N,2}(r) = r \left[\frac{1}{2} \binom{2N}{N} (r+1) - 1 \right],$$

and

$$P_{N,3}(r) = r \left[\frac{(3N)!}{6(N!)^3} (r+1)(r+2) - \binom{3N}{N} (r+1) + 1 \right].$$

Proof. The first logarithmic coefficients are

$$\kappa_{N,1} = \frac{1}{N!},$$

$$\kappa_{N,2} = \frac{1}{(N!)^2} - \frac{2}{(2N)!},$$

and

$$\kappa_{N,3} = \frac{1}{(N!)^3} - \frac{3}{N!(2N)!} + \frac{3}{(3N)!}.$$

Insert these values into theorem 7.2 and simplify. □

Corollary 7.8. *For $n \geq 1$, the polynomial $P_{N,n}(r)$ is divisible by r.*

Proof. At $r = 0$, the reciprocal family is the constant function one. Every positive-index block coefficient therefore vanishes. □

Example 7.9. For $N = 1$, one has $\mathsf{D}_1(z)^{-r} = e^{rz}$ and $P_{1,n}(r) = r^n$. For $N = 2$,

$$P_{2,1}(r) = r,$$

$$P_{2,2}(r) = r(3r + 2),$$

and

$$P_{2,3}(r) = r(15r^2 + 30r + 16).$$

At $r = 1$, these are the positive secant coefficients.

8 Logarithmic derivatives and Riccati equations

Definition 8.1. Define the transfer tangent

$$\mathsf{T}_N(z) = -\frac{\mathsf{D}_N'(z)}{\mathsf{D}_N(z)}.$$

Near the origin, its convergent block expansion has the form

$$\mathsf{T}_N(z) = \sum_{n\geq 0} T_N(n) \frac{z^{Nn+N-1}}{(Nn+N-1)!}.$$

Proposition 8.2 (Logarithmic coefficient formula)**.** *The transfer tangent satisfies*

$$\mathsf{T}_N(z) = N \sum_{m\geq 1} \kappa_{N,m} z^{Nm-1}.$$

Equivalently,

$$T_N(n) = N(Nn + N - 1)!\kappa_{N,n+1}.$$

Proof. Since $\mathsf{D}_N(z) = \Delta_N(z^N)$,

$$-\frac{\mathsf{D}_N'(z)}{\mathsf{D}_N(z)} = -Nz^{N-1}\frac{\Delta_N'(z^N)}{\Delta_N(z^N)}.$$

Differentiate the defining series for $-\log \Delta_N$ and compare coefficients. □

Theorem 8.3 (Tangent recurrence)**.** *The coefficients satisfy $T_N(0) = 1$ and*

$$T_N(n) = (-1)^n - \sum_{k=1}^{n} (-1)^k \binom{Nn+N-1}{Nk} T_N(n-k).$$

Proof. The defining equation is

$$\mathsf{D}_N \mathsf{T}_N = -\mathsf{D}_N'.$$

Extracting the coefficient of $z^{Nn+N-1}/(Nn+N-1)!$ gives

$$\sum_{k=0}^{n} (-1)^k \binom{Nn+N-1}{Nk} T_N(n-k) = (-1)^n.$$

Isolating the $k = 0$ term gives the recurrence. □

Corollary 8.4 (Derivative of the reciprocal family)**.** *For every $r \in \mathbb{C}$, the formal identity*

$$\frac{d}{dz}\mathsf{S}_N^{(r)}(z) = r\mathsf{T}_N(z)\mathsf{S}_N^{(r)}(z)$$

holds. The same identity holds analytically on every zero-free domain on which the logarithm used to define $\mathsf{S}_N^{(r)}$ is fixed.

Proof. Differentiate D_N^{-r} on a zero-free region. The formal identity follows from the same logarithmic calculation. □

For a function g, define the complete differential Bell polynomial Y_N by

$$\frac{d^N}{dz^N}e^{g(z)} = e^{g(z)}Y_N(g', g'', \dots, g^{(N)}).$$

Theorem 8.5 (Scalar higher Riccati identity)**.** *At every point of the open set $\{z : \mathsf{D}_N(z) \neq 0\}$,*

$$Y_N(-\mathsf{T}_N, -\mathsf{T}_N', \dots, -\mathsf{T}_N^{(N-1)}) = -1.$$

Proof. Write $\mathsf{D}_N = e^g$ locally. Then $g' = -\mathsf{T}_N$. Divide the equation $\mathsf{D}_N^{(N)} = -\mathsf{D}_N$ by D_N and use the defining identity for Y_N. □

This scalar identity follows from one ordinary differential equation. It does not redevelop the general nonlinear operation-tree theory of Paper III.

Example 8.6. For $N = 1$, $\mathsf{T}_1 = 1$. For $N = 2$, $\mathsf{D}_2 = \cos z$ and $\mathsf{T}_2 = \tan z$. The Riccati equation becomes

$$\mathsf{T}_2' = 1 + \mathsf{T}_2^2.$$

9 Residue-section reciprocals and Appell systems

The section $\mathsf{E}_{N,j}$ has a forced zero of order j at the origin. Removing it leaves a unit series in z^N.

Definition 9.1. For $0 \leq j < N$, define

$$\Phi_{N,j}(w) = j! \sum_{n\geq 0} \frac{w^n}{(Nn+j)!}.$$

Then

$$\mathsf{E}_{N,j}(z) = \frac{z^j}{j!}\Phi_{N,j}(z^N).$$

Define the residue-section reciprocal function

$$\mathsf{C}_{N,j}(z) = \Phi_{N,j}(z^N)^{-1} = \frac{z^j/j!}{\mathsf{E}_{N,j}(z)}.$$

For $j > 0$, the last quotient at $z = 0$ is interpreted by its removable continuation, namely the first expression in the display. Write

$$\mathsf{C}_{N,j}(z) = \sum_{n\geq 0} C_{N,j}(n)\frac{z^{Nn}}{(Nn)!}.$$

Theorem 9.2 (Triangular inverse system)**.** *The coefficients are determined by* $C_{N,j}(0) = 1$ *and*

$$\sum_{q=0}^{n} \binom{Nn+j}{Nq} C_{N,j}(q) = \delta_{n0}.$$

For $n \geq 1$*,*

$$C_{N,j}(n) = -\frac{\displaystyle\sum_{q=0}^{n-1} \binom{Nn+j}{Nq} C_{N,j}(q)}{\displaystyle\binom{Nn+j}{Nn}}.$$

Proof. Multiply the defining series for $\mathsf{C}_{N,j}$ and $\mathsf{E}_{N,j}$. Their product is $z^j/j!$. At degree $Nn+j$, multiplication by $(Nn+j)!$ gives the stated binomial coefficient. The diagonal coefficient is nonzero, so the system is triangular. □

Definition 9.3. For $r \in \mathbb{C}$, define formal coefficients $C_{N,j,n}(r)$ by

$$\mathsf{C}_{N,j}(z)^r = \sum_{n\geq 0} C_{N,j,n}(r) \frac{z^{Nn}}{(Nn)!}.$$

On a simply connected region containing 0 and avoiding the zeros of $\Phi_{N,j}(z^N)$, interpret this power as e^{rL}, where L is the holomorphic logarithm of $\mathsf{C}_{N,j}$ normalized by $L(0) = 0$.

Theorem 9.4 (Parameter-addition law for residue-section reciprocals)**.** *For each* $n \geq 1$*, the coefficient* $C_{N,j,n}(r)$ *is a polynomial in* r *of exact degree* n*. Its leading coefficient is*

$$\frac{(Nn)!}{n!} \left(-\frac{j!}{(N+j)!}\right)^n.$$

It satisfies

$$C_{N,j,n}(r+s) = \sum_{q=0}^{n} \binom{Nn}{Nq} C_{N,j,q}(r) C_{N,j,n-q}(s).$$

In particular,

$$C_{N,j,n}(1) = C_{N,j}(n).$$

Proof. The unit series $\Phi_{N,j}(w)^{-1}$ has a formal logarithm with zero constant term. Its linear term is

$$-\frac{j!}{(N+j)!} w.$$

Exponentiation after multiplication by r shows that the coefficient of w^n is a polynomial of degree at most n. The nth power of the linear logarithmic term gives the displayed nonzero leading coefficient, so the degree is exactly n. Multiplication of the powers with parameters r and s gives the coefficient identity. More explicitly, before the factorial normalization the coefficient of $r^n w^n$ is

$$\frac{1}{n!} \left(-\frac{j!}{(N+j)!}\right)^n;$$

thus multiplication by $(Nn)!$ gives the displayed leading coefficient. □

Definition 9.5. Define polynomials $E_m^{(N,j,r)}(x)$ by

$$\mathsf{C}_{N,j}(z)^r e^{xz} = \sum_{m\geq 0} E_m^{(N,j,r)}(x)\frac{z^m}{m!}.$$

Theorem 9.6 (Appell laws)**.** *For $m \geq 1$, the polynomials satisfy*

$$\frac{d}{dx}E_m^{(N,j,r)}(x) = mE_{m-1}^{(N,j,r)}(x)$$

and, for $m \geq 0$,

$$E_m^{(N,j,r)}(x+y) = \sum_{q=0}^{m}\binom{m}{q}E_q^{(N,j,r)}(x)y^{m-q}.$$

Proof. Differentiate the generating function with respect to x for the first identity. Multiply it by e^{yz} and compare coefficients for the second. □

Proposition 9.7 (Explicit Appell polynomial)**.** *For every $m \geq 0$,*

$$E_m^{(N,j,r)}(x) = \sum_{0\leq n\leq m/N}\binom{m}{Nn}C_{N,j,n}(r)x^{m-Nn}.$$

Proof. Multiply the block series for $\mathsf{C}_{N,j}^r$ by the exponential series and collect the coefficient of total degree m. □

Theorem 9.8 (Joint Appell convolution in the space and order parameters)**.** *For all $r, s, x, y \in \mathbb{C}$ and $m \geq 0$,*

$$E_m^{(N,j,r+s)}(x+y) = \sum_{q=0}^{m}\binom{m}{q}E_q^{(N,j,r)}(x)E_{m-q}^{(N,j,s)}(y).$$

Proof. Multiply the generating functions

$$\mathsf{C}_{N,j}(z)^r e^{xz} \quad \text{and} \quad \mathsf{C}_{N,j}(z)^s e^{yz}.$$

Their product is the generating function with parameters $r + s$ and $x + y$. Coefficient comparison gives the formula. □

Proposition 9.9 (Formal differential-operator representation on polynomials)**.** *Let a formal series $A(z) = \sum_{q\geq 0} a_q z^q/q!$ act on polynomials by*

$$A(\partial_x)f(x) = \sum_{q\geq 0}\frac{a_q}{q!}\partial_x^q f(x).$$

The power r is applied to the formal symbol before the substitution $z = \partial_x$. Then

$$E_m^{(N,j,r)}(x) = \mathsf{C}_{N,j}(\partial_x)^r x^m.$$

The inverse relation is

$$x^m = \Phi_{N,j}(\partial_x^N)^r E_m^{(N,j,r)}(x).$$

Both sums are finite on every polynomial.

Proof. Applying $A(\partial_x)$ to e^{xz} multiplies it by $A(z)$. Taking $A = \mathsf{C}^r_{N,j}$ gives the first identity by the defining generating function. Constant coefficient operators on polynomials satisfy

$$(AB)(\partial_x) = A(\partial_x)B(\partial_x),$$

because all derivative sums are finite and powers of ∂_x commute. The two formal symbols

$$\mathsf{C}_{N,j}(z)^r = \Phi_{N,j}(z^N)^{-r} \quad \text{and} \quad \Phi_{N,j}(z^N)^r$$

multiply to one. Their substitutions at $z = \partial_x$ are therefore inverse constant coefficient operators on polynomials, which proves the second identity. □

Corollary 9.10 (Appell symmetry)**.** *Each polynomial $E_m^{(N,j,r)}$ is monic of degree m, and*

$$E_m^{(N,j,r)}(\omega x) = \omega^m E_m^{(N,j,r)}(x).$$

Proof. The term $n = 0$ in proposition 9.7 is x^m. Every other exponent is $m - Nn$, so replacement of x by ωx multiplies every term by ω^m. □

Example 9.11. For $(N, j) = (2, 0)$,

$$\mathsf{C}_{2,0}(z) = \operatorname{sech} z.$$

For $(N, j) = (2, 1)$,

$$\mathsf{C}_{2,1}(z) = \frac{z}{\sinh z}.$$

The first case is a scaled and translated Euler family. In fact, if $E_m(x)$ denotes the classical Euler polynomial, then

$$E_m^{(2,0,1)}(x) = 2^m E_m\left(\frac{x+1}{2}\right).$$

If $B_m(x)$ denotes the classical Bernoulli polynomial, the second case has the exact identification

$$E_m^{(2,1,1)}(x) = 2^m B_m\left(\frac{x+1}{2}\right),$$

obtained by substituting $t = 2z$ in $te^{yt}/(e^t - 1)$.

10 Scaled reciprocal coupling and theta polynomials

Definition 10.1. Set

$$H_N(z) = \mathsf{E}_{N,0}(z) = \sum_{n\geq 0} \frac{z^{Nn}}{(Nn)!}.$$

For $u \in \mathbb{C}$, define $\Theta_{N,n}(u)$ by

$$H_N(uz)^{-1} H_N(z)^{-u} = \sum_{n\geq 0} \Theta_{N,n}(u) \frac{z^{Nn}}{(Nn)!}.$$

This is first a formal identity at the origin. Analytically, it is interpreted on a common simply connected zero-free region with the branch of $\log H_N$ normalized by $\log H_N(0) = 0$.

Theorem 10.2 (Theta reconstruction)**.** *For every $n \geq 0$,*

$$\Theta_{N,n}(u) = (-1)^n \sum_{q=0}^{n} \binom{Nn}{Nq} P_{N,q}(1) u^{Nq} P_{N,n-q}(u).$$

Thus $\Theta_{N,n}$ is a polynomial of degree at most Nn.

Proof. Choose ρ with $\rho^N = -1$. Then

$$H_N(z) = \mathsf{D}_N(\rho z).$$

The coefficient of z^{Nn} in $H_N(z)^{-r}$ is therefore $(-1)^n P_{N,n}(r)/(Nn)!$. Apply this to $H_N(uz)^{-1}$ and $H_N(z)^{-u}$, then multiply the two block series. □

Proposition 10.3 (Trivial parameter factors)**.** *For $n \geq 1$, the polynomial $\Theta_{N,n}(u)$ is divisible by u. If N is even, it is also divisible by $u + 1$.*

Proof. At $u = 0$, the defining product is one. Hence every positive-index coefficient vanishes. If N is even, then $H_N(-z) = H_N(z)$. At $u = -1$, the product is again one, so every positive-index coefficient vanishes there as well. □

Proposition 10.4 (Theta Newton recurrence)**.** *Put*

$$\theta_{N,n}(u) = \frac{\Theta_{N,n}(u)}{(Nn)!}.$$

Then $\theta_{N,0}(u) = 1$ and

$$n\theta_{N,n}(u) = \sum_{m=1}^{n} (-1)^m \kappa_{N,m} (u^{Nm} + u) \theta_{N,n-m}(u).$$

In particular,

$$\Theta_{N,1}(u) = -(u^N + u).$$

Proof. Write $w = z^N$. Since $H_N(z) = \Delta_N(-w)$,

$$-\log H_N(z) = \sum_{m \geq 1} (-1)^m \kappa_{N,m} \frac{w^m}{m}.$$

The logarithm of the theta generating function is therefore

$$\sum_{m \geq 1} (-1)^m \kappa_{N,m} (u^{Nm} + u) \frac{w^m}{m}.$$

Logarithmic differentiation and coefficient comparison give the recurrence. The case $n = 1$ uses $\kappa_{N,1} = 1/N!$. □

Example 10.5. For $N = 2$,

$$\Theta_{2,1}(u) = -u(u + 1)$$

and

$$\Theta_{2,2}(u) = u(u + 1)(5u^2 + u + 2).$$

The two factors predicted by proposition 10.3 are visible.

11 Growth and the geometry of the large zeros

We now assume $N \geq 2$. Write

$$\eta_k = e^{(2k+1)\pi i/N}, \qquad 0 \leq k < N.$$

These are the vertices of a regular polygon and

$$\mathsf{D}_N(z) = \frac{1}{N} \sum_{k=0}^{N-1} e^{\eta_k z}.$$

Standard Fact 11.1 (Order from Taylor coefficients)**.** If $f(w) = \sum_{n\geq 0} a_n w^n$ is an entire function with infinitely many nonzero coefficients, then

$$\operatorname{ord} f = \limsup_{\substack{n\to\infty \\ a_n \neq 0}} \frac{n \log n}{\log(1/|a_n|)}.$$

Canonical products for entire functions of order below one have genus zero. We use these standard facts in the form given in [Boa54, Tit39].

Proposition 11.2 (Orders and exponential type)**.** *For $N \geq 2$,*

$$\operatorname{ord} \Delta_N = \frac{1}{N}.$$

The function D_N has order one and exponential type one.

Proof. The coefficient of w^n in Δ_N has modulus $1/(Nn)!$. Stirling's formula and standard fact 11.1 give order $1/N$. Since $M_{f(\,\cdot^N)}(r) = M_f(r^N)$ for the maximum modulus, composition with z^N multiplies the order by N and gives order one for D_N.

The finite exponential representation gives

$$|\mathsf{D}_N(z)| \leq e^{|z|},$$

so the type is at most one. Choose a root η_0 and put $z = t\overline{\eta_0}$ with $t > 0$. The term $e^{\eta_0 z} = e^t$ has strictly larger real exponent than every other term. Hence

$$\mathsf{D}_N(t\overline{\eta_0}) = \frac{1}{N} e^t (1 + o(1)).$$

The type is therefore one. □

Definition 11.3. For $\theta \in \mathbb{R}$, define

$$h_N(\theta) = \max_{0\leq k<N} \cos\left(\theta + \frac{(2k+1)\pi}{N}\right).$$

The tie directions are

$$\mathfrak{T}_N = \left\{ \frac{2\pi k}{N} : 0 \leq k < N \right\} \mod 2\pi.$$

Theorem 11.4 (Indicator polygon and dominant sectors)**.** *The indicator of D_N equals h_N, with the limsup taken over radii where $\mathsf{D}_N(re^{i\theta}) \neq 0$:*

$$\limsup_{r\to\infty} \frac{1}{r} \log |\mathsf{D}_N(re^{i\theta})| = h_N(\theta).$$

If K is a compact set of directions disjoint from $\mathfrak{T}_N$, then there are $\delta > 0$ and $R > 0$ such that, for each $\theta \in K$, one root $\eta(\theta)$ of -1 satisfies

$$\mathsf{D}_N(re^{i\theta}) = \frac{1}{N}\exp(\eta(\theta)re^{i\theta})\left(1 + O(e^{-\delta r})\right)$$

uniformly for $r \geq R$. In particular, no sufficiently large zero can lie in those directions.

Proof. For $z = re^{i\theta}$, the real part of the exponent $\eta_k z$ is

$$r\cos\left(\theta + \frac{(2k+1)\pi}{N}\right).$$

The triangle inequality gives the upper indicator bound $h_N(\theta)$.

Away from a tie direction, the maximizing vertex is unique. It is constant on each connected component of the complement of the tie directions. A compact set avoiding all ties meets only finitely many such components, and compactness gives one uniform positive lower bound for every relevant projection gap. Division by the maximizing exponential gives the stated uniform asymptotic and the zero-free conclusion.

At a tie direction exactly two adjacent vertices attain the maximum. After rotation it is enough to take $\theta = 0$. With $c_N = \cos(\pi/N)$ and $s_N = \sin(\pi/N)$, the two dominant terms give

$$\frac{2}{N}e^{c_N r}\cos(s_N r)$$

and every other term is exponentially smaller. Along a sequence on which $|\cos(s_N r)| \geq 1/2$, the lower indicator bound is $c_N = h_N(0)$. Rotational symmetry treats every tie direction. □

The zero strings are governed by two adjacent exponents. Put

$$a_N = \frac{\pi}{N}, \qquad s_N = \sin a_N, \qquad c_N = \cos a_N,$$

and

$$A_N = \frac{\pi}{s_N}, \qquad t_m = A_N\left(m + \frac{1}{2}\right).$$

For $N \geq 3$, also put

$$\gamma_N = c_N - \cos(3a_N) > 0.$$

Lemma 11.5 (Two term dominance)**.** *Assume $N \geq 3$. There are positive constants Y, R, and C with the following properties.*

If $z = x + iy$ lies in the sector $|\arg z| \leq a_N$, has $x \geq R$, and is a zero of D_N, then $|y| \leq Y$.

On the strip $x \geq R$ and $|y| \leq Y$, define

$$R_N(z) = \sum_{\substack{\eta^N = -1 \\ \eta \neq e^{ia_N}, e^{-ia_N}}} \exp\big((\eta - e^{-ia_N})z\big).$$

Then

$$Ne^{-e^{-ia_N}z}\mathsf{D}_N(z) = 1 + e^{2is_N z} + R_N(z),$$

and, for $q = 0, 1, 2$,

$$|R_N^{(q)}(z)| \leq Ce^{-\gamma_N x}.$$

Proof. A direct projection estimate gives the stronger bound

$$Y = \frac{\log(N-1)}{2s_N}$$

for every zero in the sector, without a large-radius assumption. Write $z = re^{i\theta}$ with $0 \le \theta \le a_N$ and $r \ge 0$. For $0 \le \ell \le N-2$, the angle $\theta + (2\ell+1)a_N$ lies between $a_N + \theta$ and $2\pi - a_N - \theta$. Monotonicity of cosine on $[0, \pi]$, followed by reflection about π, therefore gives

$$\Re(\eta_\ell z) \le r\cos(a_N + \theta).$$

The remaining vertex e^{-ia_N} has projection $r\cos(a_N - \theta)$. Their difference is at least $2s_N r\sin\theta = 2s_N y$. If $\mathsf{D}_N(z) = 0$, divide its finite exponential sum by the term with exponent $e^{-ia_N}z$ and apply the triangle inequality:

$$1 \le (N-1)e^{-2s_N y}.$$

Thus $y \le Y$. Conjugation gives the same bound for $-y$ on the lower half sector. Since $N \ge 3$ and $s_N > 0$, this Y is positive.

On the resulting strip, the roots e^{ia_N} and e^{-ia_N} have real part c_N, while each remaining root $\eta = u + iv$ satisfies $u \le \cos(3a_N) = c_N - \gamma_N$. Hence

$$\Re\big((\eta - e^{-ia_N})z\big) \le -\gamma_N x + 2Y.$$

Division by $e^{e^{-ia_N}z}$ gives the displayed normalization. Each differentiation multiplies a remainder term by $\eta - e^{-ia_N}$, whose modulus is at most two. Thus, for $q = 0, 1, 2$,

$$|R_N^{(q)}(z)| \le (N-2)2^q e^{2Y} e^{-\gamma_N x}.$$

One may take $R = 1$ and $C = 4(N-2)e^{2Y}$. □

Lemma 11.6 (Near cancellation of two exponentials)**.** *For every $M > 0$, there are constants $\varepsilon_M > 0$ and $C_M > 0$ such that the following holds. If*

$$|\Re w| \le M$$

and

$$|1 + e^w| \le \varepsilon_M,$$

then there is a unique integer m with

$$|w - (2m+1)\pi i| \le C_M|1 + e^w|.$$

Proof. In fact, the constants $\varepsilon_M = 1/2$ and $C_M = 3/2$ work for every complex w, without the strip restriction. Put $u = -(1 + e^w)$ and $L = \log(1+u) = \log(-e^w)$, using the principal logarithm. The elementary logarithm estimate on $|u| \le 1/2$ gives $|L| \le (3/2)|u|$. Since $e^{L+\pi i} = e^w$, there is an integer m such that

$$w = L + (2m+1)\pi i.$$

This gives the required distance bound. Distinct centers are separated by at least 2π, whereas two centers satisfying the bound would be separated by at most $3|1 + e^w| \le 3/2$. Thus the integer is unique. The formal proof uses the convergent logarithm estimate and the exact periodicity theorem for the complex exponential. □

Theorem 11.7 (Large zero strings)**.** *Assume $N \geq 3$. There are constants m_0, C, and $\gamma > 0$ and a sequence of positive real simple zeros $(\tau_m)_{m \geq m_0}$ of D_N such that*

$$\tau_m = t_m + O(e^{-\gamma t_m}).$$

Every zero of sufficiently large modulus is of the form

$$\omega^k \tau_m$$

for a unique $0 \leq k < N$ and a unique $m \geq m_0$.

The derivatives satisfy

$$\mathsf{D}_N'(\omega^k \tau_m) = \omega^{-k} \mathsf{D}_N'(\tau_m)$$

and

$$|\mathsf{D}_N'(\tau_m)| = \frac{2s_N}{N} e^{c_N \tau_m} \left(1 + O(e^{-\gamma \tau_m})\right).$$

Proof. Work first in the sector $|\arg z| \leq a_N$. By lemma 11.5, every sufficiently large zero lies in a fixed strip about the positive real axis. Put

$$G(z) = N e^{-e^{-ia_N} z} \mathsf{D}_N(z).$$

Then

$$G(z) = 1 + e^{2is_N z} + R_N(z)$$

with an exponentially small remainder and derivative.

If $G(z) = 0$, then

$$|1 + e^{2is_N z}| \leq C e^{-\gamma_N \Re z}.$$

The real part of $2is_N z$ is bounded in the strip. Apply lemma 11.6 with $w = 2is_N z$. There is a unique integer m such that

$$|z - t_m| \leq C_1 e^{-\gamma_N \Re z},$$

where

$$t_m = \frac{(2m+1)\pi}{2s_N}.$$

In particular, $|\Re z - t_m| = o(1)$. For all sufficiently large zeros we have $|\Re z - t_m| \leq 1$, and hence

$$|z - t_m| \leq C_2 e^{-\gamma_N t_m}$$

with a constant independent of m. Because $\Re z \to +\infty$ in this argument, the corresponding integers m are eventually nonnegative and tend to infinity.

Fix a large m, put $a = 2is_N$, and consider the closed disk $B_m = \{z : |z - t_m| \leq 1/(2s_N)\}$. The remainder bounds hold uniformly on these fixed-radius disks. For sufficiently large m, $|R_N| \leq 1/2$ and $|R_N'| \leq s_N/2$ there. Define

$$\Phi_m(z) = t_m + \frac{\log(1 + R_N(z))}{2is_N},$$

using the principal logarithm. The estimate $|\log(1+u)| \leq (3/2)|u|$ shows that Φ_m maps B_m into itself. The derivative bound $|\Phi_m'| \leq 1/2$ makes it a contraction. Banach's fixed-point theorem gives a unique fixed point. On B_m, $|a(z - t_m)| \leq 1 < \pi$, so the principal logarithm inverts the exponential and the fixed-point equation is equivalent to $G(z) = 0$. Moreover,

$$|z - t_m| \leq \frac{3}{4s_N} |R_N(z)| = O(e^{-\gamma_N t_m}).$$

At this zero,

$$|G'(z)| \geq |a|\,|1 + R_N(z)| - |R'_N(z)| \geq s_N/2 > 0,$$

so the zero is simple. The disks are mutually disjoint because their diameter $1/s_N$ is smaller than the center spacing π/s_N. Enlarging the strip if necessary, these disks lie inside the fundamental sector and the fixed strip for large m.

The localization estimate puts every sufficiently large zero in the fundamental sector into one of these disks. The zero sets of G and D_N coincide because their prefactor is nonzero. Each disk is invariant under complex conjugation, and D_N has real coefficients. The conjugate of its unique zero is therefore the same zero. It is real, positive for large m, and simple by the derivative bound above. Denote it by τ_m.

The parts of the fundamental sector away from the positive tie ray, including its boundary rays, were already excluded uniformly. Rotational invariance now gives every remaining large zero as $\omega^k\tau_m$. Distinct rays and the disjoint disks give uniqueness of k and m. Taking $\gamma = \gamma_N$ proves the zero location in the statement.

Differentiate the normalized expression. At $z = \tau_m$,

$$G'(\tau_m) = 2is_N e^{2is_N\tau_m} + R'_N(\tau_m) = -2is_N + O(e^{-\gamma_N\tau_m}).$$

Since $G(\tau_m) = 0$,

$$\mathsf{D}'_N(\tau_m) = \frac{1}{N} e^{e^{-ia_N\tau_m}} G'(\tau_m).$$

Taking absolute values gives the derivative asymptotic. Differentiating $\mathsf{D}_N(\omega z) = \mathsf{D}_N(z)$ gives the rotation formula. □

Theorem 11.8 (Leading displacement term and derivative sign)**.** *With the notation of lemma 11.5 and theorem 11.7,*

$$\tau_m = t_m + \frac{R_N(t_m)}{2is_N} + O(e^{-2\gamma_N t_m}).$$

The quotient $R_N(t_m)/(2is_N)$ is real. One also has

$$\mathsf{D}'_N(\tau_m) = (-1)^{m+1}\frac{2s_N}{N} e^{c_N\tau_m}\left(1 + O(e^{-\gamma_N\tau_m})\right).$$

Proof. Put $\delta_m = \tau_m - t_m$. The large zero theorem gives $\delta_m = O(e^{-\gamma_N t_m})$. Since $e^{2is_N t_m} = -1$,

$$1 + e^{2is_N(t_m+\delta_m)} = -2is_N\delta_m + O(\delta_m^2).$$

The remainder estimate through the first derivative gives

$$R_N(t_m + \delta_m) = R_N(t_m) + O(e^{-\gamma_N t_m}|\delta_m|).$$

Substitution in the zero equation gives the displacement formula.

At the real point t_m, the normalized value

$$Ne^{-e^{-ia_N t_m}}\mathsf{D}_N(t_m)$$

is purely imaginary. The two dominant terms cancel there, so $R_N(t_m)$ is purely imaginary and the displayed correction is real.

For the derivative, use the formula in the proof of theorem 11.7 and

$$e^{-is_N\tau_m} = (-1)^m(-i)\left(1 + O(e^{-\gamma_N\tau_m})\right).$$

The stated real sign follows. □

Corollary 11.9 (Leading correction for $N = 3$)**.** *For $N = 3$,*

$$t_m = \frac{2\pi}{\sqrt{3}}\left(m + \frac{1}{2}\right)$$

and

$$\tau_m = t_m + \frac{(-1)^m}{\sqrt{3}} e^{-3t_m/2} + O(e^{-3t_m}).$$

Proof. The only nondominant exponent is -1. Hence

$$R_3(t) = \exp\left(\left(-\frac{3}{2} + \frac{\sqrt{3}}{2}i\right) t\right).$$

At $t = t_m$, its phase is $i(-1)^m$. Insert this value into theorem 11.8. □

Corollary 11.10 (The case $N = 2$)**.** *For $N = 2$,*

$$\mathsf{D}_2(z) = \cos z.$$

Its zeros are

$$\pm\left(m + \frac{1}{2}\right)\pi, \qquad m \geq 0,$$

and all are simple. They agree with the formula $t_m = A_2(m + 1/2)$.

Proof. This is the standard zero set of the cosine function. □

Corollary 11.11 (Eventual simplicity and exact rays)**.** *For every $N \geq 2$, all but finitely many zeros of D_N are simple. Every sufficiently large zero lies on one of the rays*

$$\omega^k(0, \infty), \qquad 0 \leq k < N.$$

Proof. Use theorem 11.7 for $N \geq 3$ and corollary 11.10 for $N = 2$. □

Example 11.12 (Exact fourth-order strings)**.** For $N = 4$,

$$\mathsf{D}_4(z) = \cos\left(\frac{z}{\sqrt{2}}\right)\cosh\left(\frac{z}{\sqrt{2}}\right).$$

The positive zeros are exactly

$$\tau_m = \sqrt{2}\pi\left(m + \frac{1}{2}\right) = A_4\left(m + \frac{1}{2}\right).$$

Rotation by the fourth roots of unity gives the real and imaginary zero strings. In this case the error term in theorem 11.7 is identically zero.

Corollary 11.13 (Reduced zeros)**.** *Let Λ_N be the nonzero zero multiset of Δ_N. All but finitely many members of Λ_N are positive and simple. They can be indexed as*

$$\lambda_m = \tau_m^N$$

for all sufficiently large m. If $N \geq 3$, the notation may be chosen as in theorem 11.7, and

$$\lambda_m = t_m^N + O\left(t_m^{N-1} e^{-\gamma t_m}\right).$$

For $N = 2$, take $\tau_m = t_m = \pi(m + 1/2)$ for $m \geq 0$; then $\lambda_m = t_m^2$ exactly.

Proof. The identity $\mathsf{D}_N(z) = \Delta_N(z^N)$ identifies each orbit of N nonzero zeros of D_N with one zero of Δ_N. For $N \geq 3$, the large orbit representatives are the positive zeros τ_m from theorem 11.7; for $N = 2$, use corollary 11.10. Simplicity is preserved because

$$\mathsf{D}'_N(z) = Nz^{N-1}\Delta'_N(z^N)$$

and $z \neq 0$. The asymptotic follows by expanding τ_m^N about t_m. □

Theorem 11.14 (Zero counting asymptotics)**.** *Let $n_{\mathsf{D}_N}(R)$ count the zeros of D_N in $|z| \leq R$, with multiplicity. Let $n_{\Delta_N}(R)$ count the zeros of Δ_N in $|w| \leq R$. Then, as $R \to \infty$,*

$$n_{\mathsf{D}_N}(R) = \frac{N \sin(\pi/N)}{\pi} R + O(1)$$

and

$$n_{\Delta_N}(R) = \frac{\sin(\pi/N)}{\pi} R^{1/N} + O(1).$$

Proof. Each of the N rays contains one large zero near

$$t_m = \frac{\pi}{\sin(\pi/N)} \left(m + \frac{1}{2} \right).$$

Since $\tau_m - t_m = o(1)$ and the model spacing is the positive constant A_N, replacing $\tau_m \leq R$ by $t_m \leq R$ changes the count by at most a bounded amount. The number of such indices with $t_m \leq R$ is

$$\frac{\sin(\pi/N)}{\pi} R + O(1).$$

Multiplication by N and addition of the finite exceptional set give the first formula. The second follows from

$$n_{\mathsf{D}_N}(R^{1/N}) = N n_{\Delta_N}(R).$$

□

Corollary 11.15 (Zero exponents)**.** *The exponent of convergence of the nonzero zeros of D_N is one. The exponent of convergence of the zeros of Δ_N is $1/N$.*

Proof. The asymptotic counting formulas in theorem 11.14 give the two exponents by partial summation. □

12 Hadamard products and zero power sums

The reduction from z to $w = z^N$ removes the rotational repetition of the zeros.

Theorem 12.1 (Genus zero product)**.** *For $N \geq 2$, let Λ_N be the nonzero zero multiset of Δ_N. Then*

$$\Delta_N(w) = \prod_{\lambda \in \Lambda_N} \left(1 - \frac{w}{\lambda} \right),$$

where the product converges locally uniformly. Consequently,

$$\mathsf{D}_N(z) = \prod_{\lambda \in \Lambda_N} \left(1 - \frac{z^N}{\lambda} \right).$$

Proof. By proposition 11.2, the order of Δ_N is $1/N < 1$. By corollary 11.13, it has infinitely many zeros, and corollary 11.15 gives

$$\sum_{\lambda \in \Lambda_N} |\lambda|^{-1} < \infty.$$

Hadamard factorization therefore gives a canonical product of genus zero. The only possible exponential factor is a nonzero constant because the order is below one. Since $\Delta_N(0) = 1$, that constant is one. Substitution of $w = z^N$ gives the second formula. □

Theorem 12.2 (Diagonal Fredholm determinant representation)**.** *Let I_N be a multiplicity-sensitive countable index set for Λ_N and put $\mathcal{H}_N = \ell^2(I_N)$. On its standard orthonormal basis, define the diagonal operator*

$$K_N e_\lambda = \lambda^{-1} e_\lambda.$$

Then K_N is trace class and

$$\Delta_N(w) = \det_F(I - wK_N).$$

For every $m \geq 1$,

$$\kappa_{N,m} = \operatorname{tr}(K_N^m).$$

Proof. By corollary 11.15,

$$\sum_{\lambda \in \Lambda_N} |\lambda|^{-1} < \infty.$$

Thus the diagonal operator K_N is trace class. Its Fredholm determinant is the locally uniform product

$$\det_F(I - wK_N) = \prod_{\lambda \in \Lambda_N} \left(1 - \frac{w}{\lambda}\right),$$

which equals $\Delta_N(w)$ by theorem 12.1.

Taking the logarithm gives

$$-\log \Delta_N(w) = \sum_{m \geq 1} \frac{w^m}{m} \operatorname{tr}(K_N^m)$$

near the origin. Comparison with the definition of $\kappa_{N,m}$ proves the trace identity. □

This is the standard diagonal encoding of this family's genus-zero product. It is not a general trace-class resolvent or Feshbach theorem.

Corollary 12.3 (Complete symmetric function form)**.** *For a positive integer r, let each eigenvalue λ^{-1} of K_N be repeated r times. If h_n denotes the complete homogeneous symmetric function of degree n in this absolutely summable multiset, then*

$$\frac{P_{N,n}(r)}{(Nn)!} = h_n.$$

For arbitrary complex r, the Bell polynomial formula is the polynomial continuation obtained by multiplying every power sum by r.

Proof. The generating function for the complete homogeneous symmetric functions is

$$\prod_{\lambda \in \Lambda_N} \left(1 - \frac{w}{\lambda}\right)^{-r} = \Delta_N(w)^{-r}.$$

Coefficient comparison gives the first identity. Power sums determine the complete homogeneous functions through the Newton recurrence. Replacing each power sum by r times itself gives theorems 7.2 and 7.6. □

Proposition 12.4 (Products for the normalized sections)**.** *For every $0 \le j < N$, the entire function*

$$\Phi_{N,j}(w) = j! \sum_{n \ge 0} \frac{w^n}{(Nn+j)!}$$

has order $1/N$. If $N \ge 2$ and $\Lambda_{N,j}$ is its nonzero zero multiset, then

$$\Phi_{N,j}(w) = \prod_{\mu \in \Lambda_{N,j}} \left(1 - \frac{w}{\mu}\right).$$

For $m \ge 1$, put

$$\sigma_{N,j,m} = \sum_{\mu \in \Lambda_{N,j}} \mu^{-m}.$$

Then

$$C_{N,j,n}(r) = \frac{(Nn)!}{n!} B_n(r\sigma_{N,j,1}, r1!\sigma_{N,j,2}, \dots, r(n-1)!\sigma_{N,j,n}).$$

Proof. The coefficient formula for the order gives $1/N$. The function is not a polynomial. An entire function of order below one with only finitely many zeros would be a polynomial times a constant by Hadamard factorization, which is impossible here. The genus zero product follows as in theorem 12.1.

Taking the logarithm gives

$$-\log \Phi_{N,j}(w) = \sum_{m \ge 1} \sigma_{N,j,m} \frac{w^m}{m}.$$

Since $\mathsf{C}_{N,j}(z)^r = \Phi_{N,j}(z^N)^{-r}$, the Bell polynomial calculation from theorem 7.2 gives the coefficient formula. □

Definition 12.5. For $m \ge 1$, define

$$\sigma_{N,m} = \sum_{\lambda \in \Lambda_N} \lambda^{-m},$$

with multiplicities.

The series converges absolutely because the exponent of convergence of Λ_N is $1/N < 1$.

Theorem 12.6 (Newton sums for the reduced zeros)**.** *For $|w|$ below the least zero modulus, with the analytic logarithm normalized to zero at the origin,*

$$-\log \Delta_N(w) = \sum_{m \ge 1} \sigma_{N,m} \frac{w^m}{m}.$$

Thus

$$\kappa_{N,m} = \sigma_{N,m}.$$

If

$$a_m = \frac{(-1)^m}{(Nm)!}, \qquad a_0 = 1,$$

then

$$\sigma_{N,m} = -m a_m - \sum_{k=1}^{m-1} a_k \sigma_{N,m-k}.$$

Proof. Take the logarithm of the product in theorem 12.1. Absolute convergence in a disk below the first zero permits termwise expansion

$$-\log\left(1-\frac{w}{\lambda}\right)=\sum_{m\geq 1}\frac{w^m}{m\lambda^m}.$$

This gives the logarithmic formula and identifies $\kappa_{N,m}$ with $\sigma_{N,m}$.

Differentiate the product identity in logarithmic form to obtain

$$\Delta_N'(w)=-\Delta_N(w)\sum_{m\geq 1}\sigma_{N,m}w^{m-1}.$$

Comparison of the coefficient of w^{m-1} gives

$$ma_m=-\sum_{k=0}^{m-1}a_k\sigma_{N,m-k}.$$

Solving for $\sigma_{N,m}$ gives the recurrence. □

Corollary 12.7 (Convergent Mittag–Leffler partial-fraction form)**.** *Away from the zeros of* D_N*,*

$$\mathsf{T}_N(z)=Nz^{N-1}\sum_{\lambda\in\Lambda_N}\frac{1}{\lambda-z^N}.$$

The series converges absolutely and locally uniformly on compact sets away from the poles. Thus no regularizing terms are required.

Proof. Logarithmic differentiation of the genus zero product gives

$$-\frac{\Delta_N'(w)}{\Delta_N(w)}=\sum_{\lambda\in\Lambda_N}\frac{1}{\lambda-w}.$$

The series is locally uniform because $\sum_\lambda|\lambda|^{-1}<\infty$. Substitute $w=z^N$ and use the chain rule. □

Proposition 12.8 (Theta Bell polynomial formula)**.** *Put*

$$\widetilde{\sigma}_{N,m}=(-1)^m\sigma_{N,m}.$$

Then

$$\Theta_{N,n}(u)=\frac{(Nn)!}{n!}B_n(\widetilde{\sigma}_{N,1}(u^N+u),1!\widetilde{\sigma}_{N,2}(u^{2N}+u),\dots,(n-1)!\widetilde{\sigma}_{N,n}(u^{nN}+u)).$$

Proof. The reduced unsigned denominator is

$$\Psi_N(w)=\sum_{n\geq 0}\frac{w^n}{(Nn)!}=\Delta_N(-w).$$

By theorem 12.6,

$$-\log\Psi_N(w)=\sum_{m\geq 1}\widetilde{\sigma}_{N,m}\frac{w^m}{m}.$$

With $w=z^N$, the logarithm of the theta generating function is

$$\sum_{m\geq 1}\widetilde{\sigma}_{N,m}(u^{Nm}+u)\frac{w^m}{m}.$$

The complete Bell polynomial formula follows. □

Corollary 12.9 (Tangent coefficients as zero sums)**.** *For every $n \geq 0$,*

$$T_N(n) = N(Nn + N - 1)!\sigma_{N,n+1}.$$

Proof. Combine proposition 8.2 and theorem 12.6. □

Corollary 12.10 (Reciprocal coefficients from zero data)**.** *For every $n \geq 0$,*

$$P_{N,n}(r) = \frac{(Nn)!}{n!} B_n(r\sigma_{N,1}, r1!\sigma_{N,2}, \ldots, r(n-1)!\sigma_{N,n}).$$

Proof. Use theorems 7.2 and 12.6. □

Example 12.11 (The cosine orbit sums)**.** For $N = 2$,

$$\Delta_2(w) = \cos\sqrt{w}.$$

Its zeros are

$$\lambda_m = \pi^2 \left(m + \frac{1}{2}\right)^2, \qquad m \geq 0.$$

Hence

$$\sigma_{2,q} = \frac{2^{2q} - 1}{\pi^{2q}} \zeta(2q).$$

The identity in corollary 12.9 recovers the classical tangent coefficients from these zero power sums.

13 Coefficient asymptotics from the nearest zeros

For an entire function f, let $\mathcal{Z}(f)$ denote its multiset of nonzero zeros. Let r be a positive integer. Then D_N^{-r} is meromorphic and a simple zero of D_N becomes a pole of order r.

Theorem 13.1 (Nearest-pole coefficient expansion)**.** *Let $r \in \mathbb{Z}_{>0}$. Suppose the zeros of D_N of least modulus R are simple and form the finite set $\mathcal{Z}_R$. Suppose the next zero modulus is $R_2 > R$. At each $\rho \in \mathcal{Z}_R$, write the principal part*

$$\mathsf{D}_N(z)^{-r} = \sum_{\ell=1}^{r} a_{\rho,\ell} \left(1 - \frac{z}{\rho}\right)^{-\ell} + O(1).$$

Then, for every R_* with $R < R_* < R_2$ and $m = Nn$, as $n \to \infty$,

$$P_{N,n}(r) = m! \sum_{\rho \in \mathcal{Z}_R} \sum_{\ell=1}^{r} a_{\rho,\ell} \binom{m + \ell - 1}{\ell - 1} \rho^{-m} + O_{R_*}(m! R_*^{-m}).$$

The leading Laurent coefficient is

$$a_{\rho,r} = (-\rho \mathsf{D}_N'(\rho))^{-r}.$$

Proof. Subtract the complete principal parts at the zeros on $|z| = R$. The remainder is holomorphic on $|z| < R_2$, hence on a neighborhood of every closed disk $|z| \leq R_* < R_2$. Cauchy's estimate gives an ordinary coefficient bound $O_{R_*}(R_*^{-m})$. The coefficient of z^m in

$$\left(1 - \frac{z}{\rho}\right)^{-\ell}$$

is

$$\binom{m+\ell-1}{\ell-1}\rho^{-m}.$$

Multiplication by $m!$ converts the ordinary coefficient to $P_{N,n}(r)$.

Near a simple zero,

$$\mathsf{D}_N(z) = -\rho\mathsf{D}'_N(\rho)\left(1-\frac{z}{\rho}\right) + O\left(\left(1-\frac{z}{\rho}\right)^2\right).$$

Raising to the power $-r$ gives the formula for $a_{\rho,r}$. □

Corollary 13.2 (Leading dominant zero term)**.** *Under the hypotheses of theorem 13.1,*

$$P_{N,n}(r) = m!\binom{m+r-1}{r-1}\sum_{\rho\in\mathcal{Z}_R}\frac{\rho^{-m}}{(-\rho\mathsf{D}'_N(\rho))^r} + O\left(m!m^{r-2}R^{-m}\right) + O_{R_*}\left(m!R_*^{-m}\right).$$

For $r = 1$, the first error term is absent.

Proof. The terms with $\ell < r$ in theorem 13.1 are bounded by a constant times $m^{r-2}R^{-m}$. The term with $\ell = r$ gives the displayed main term. □

Corollary 13.3 (Orbit reduced nearest circle term)**.** *Assume the hypotheses of theorem 13.1. Choose one representative from each orbit of the nearest zeros under multiplication by ω, and call the resulting set $\mathcal{Z}_R^{\mathrm{orb}}$. With $m = Nn$,*

$$P_{N,n}(r) = Nm!\binom{m+r-1}{r-1}\sum_{\tau\in\mathcal{Z}_R^{\mathrm{orb}}}\frac{\tau^{-m}}{(-\tau\mathsf{D}'_N(\tau))^r} + O\left(m!m^{r-2}R^{-m}\right) + O_{R_*}\left(m!R_*^{-m}\right).$$

For $r = 1$, the first error term is absent.

Proof. If $\rho = \omega^k\tau$, then

$$\mathsf{D}'_N(\rho) = \omega^{-k}\mathsf{D}'_N(\tau).$$

Therefore

$$-\rho\mathsf{D}'_N(\rho) = -\tau\mathsf{D}'_N(\tau).$$

Since $m = Nn$, one also has $\rho^{-m} = \tau^{-m}$. Every orbit therefore contributes N equal terms in corollary 13.2. □

Corollary 13.4 (Secant asymptotic)**.** *For $N = 2$ and $r = 1$,*

$$P_{2,n}(1) \sim 2(2n)!\left(\frac{2}{\pi}\right)^{2n+1}.$$

Proof. The nearest zeros of $\cos z$ are $\rho = \pm\pi/2$, both simple. Their contributions in corollary 13.2 are equal. □

14 Zero zeta functions

The assumption $N \geq 2$ remains in force throughout this section. The zero counting theorem gives the exact abscissae of absolute convergence. The exponentially accurate zero locations give more. They compare the zero zeta with a Hurwitz zeta function.

Definition 14.1. Define

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = \sum_{\rho \in \mathcal{Z}(\mathsf{D}_N)} |\rho|^{-s}$$

and

$$\mathfrak{Z}_N^{\mathrm{red}}(s) = \sum_{\lambda \in \Lambda_N} |\lambda|^{-s},$$

with multiplicities. Each term means $e^{-s \log |\rho|}$ or $e^{-s \log |\lambda|}$.

Theorem 14.2 (Abscissae of absolute convergence)**.** *The series $\mathfrak{Z}_N^{\mathrm{abs}}$ converges absolutely and locally uniformly when*

$$\Re s > 1.$$

It fails to converge absolutely when $\Re s \leq 1$. The series $\mathfrak{Z}_N^{\mathrm{red}}$ converges absolutely and locally uniformly when

$$\Re s > \frac{1}{N}.$$

It fails to converge absolutely when $\Re s \leq 1/N$. Both functions are holomorphic on their open half planes of absolute convergence.

Proof. By theorem 11.14, the zeros of D_N have linear counting density, while the reduced zeros have counting order $R^{1/N}$. Partial summation gives absolute convergence in the stated open half planes. The positive terms obtained after taking absolute values diverge on the boundary and to its left. Local uniform convergence follows by the Weierstrass test on compact subsets. □

Theorem 14.3 (Meromorphic continuation of the symmetry-orbit zero zeta)**.** *Put*

$$A_N = \frac{\pi}{\sin(\pi/N)}.$$

There is an entire function Q_N^{red} such that

$$\mathfrak{Z}_N^{\mathrm{red}}(s) = A_N^{-Ns} \zeta\left(Ns, \frac{1}{2}\right) + Q_N^{\mathrm{red}}(s)$$

on $\Re s > 1/N$. The right hand side gives a meromorphic continuation to $\mathbb{C}$ with one simple pole at $s = 1/N$. Its residue is

$$\frac{\sin(\pi/N)}{N\pi}.$$

There is an entire function Q_N^{abs} such that, initially on $\Re s > 1$,

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = N A_N^{-s} \zeta\left(s, \frac{1}{2}\right) + Q_N^{\mathrm{abs}}(s).$$

Thus $\mathfrak{Z}_N^{\mathrm{abs}}$ continues meromorphically to $\mathbb{C}$ with one simple pole at $s = 1$, of residue

$$\frac{N \sin(\pi/N)}{\pi}.$$

Proof. For $N = 2$, corollary 11.10 and $\mathsf{D}_2(z) = \Delta_2(z^2)$ give

$$\lambda_m = \pi^2 \left(m + \frac{1}{2}\right)^2, \qquad m \geq 0.$$

Thus the first formula holds with $Q_2^{\mathrm{red}} = 0$.

For $N \geq 3$, corollary 11.13 writes all but finitely many reduced zeros as

$$\lambda_m = \left(A_N \left(m + \frac{1}{2}\right)\right)^N (1 + \varepsilon_m)$$

where $\lambda_m > 0$, $1 + \varepsilon_m > 0$, and

$$\varepsilon_m = O\left(t_m^{-1} e^{-\gamma t_m}\right).$$

On every compact set of s values,

$$(1 + \varepsilon_m)^{-s} - 1 = O(|\varepsilon_m|)$$

uniformly for large m. The difference

$$\lambda_m^{-s} - A_N^{-Ns} \left(m + \frac{1}{2}\right)^{-Ns}$$

therefore has a normally convergent sum on every compact subset of $\mathbb{C}$. Indeed, the model power has at most polynomial growth in m on a compact set, while $|\varepsilon_m|$ decays exponentially. The finitely many exceptional terms $|\lambda|^{-s}$ and the finitely many omitted model terms are entire functions of s. Their addition to the normally convergent tail defines Q_N^{red} and proves the first formula. Thus that formula is established for every $N \geq 2$.

In either case, the Hurwitz zeta function has a meromorphic continuation with one simple pole at its first argument equal to one and residue one. Composition with Ns gives the stated pole and residue.

For every $N \geq 2$, each nonzero zero λ of Δ_N gives N zeros of D_N, each of modulus $|\lambda|^{1/N}$. Hence

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = N \mathfrak{Z}_N^{\mathrm{red}}(s/N).$$

Substitution gives the second formula with

$$Q_N^{\mathrm{abs}}(s) = N Q_N^{\mathrm{red}}(s/N),$$

which is entire. □

Corollary 14.4 (Exact zeta models in orders two and four)**.** *For $N = 2$,*

$$\mathfrak{Z}_2^{\mathrm{red}}(s) = \pi^{-2s} \zeta\left(2s, \frac{1}{2}\right)$$

and

$$\mathfrak{Z}_2^{\mathrm{abs}}(s) = 2\pi^{-s} \zeta\left(s, \frac{1}{2}\right).$$

For $N = 4$, put $A_4 = \sqrt{2}\pi$. Then

$$\mathfrak{Z}_4^{\mathrm{red}}(s) = A_4^{-4s} \zeta\left(4s, \frac{1}{2}\right)$$

and

$$\mathfrak{Z}_4^{\mathrm{abs}}(s) = 4A_4^{-s} \zeta\left(s, \frac{1}{2}\right).$$

In these two cases the entire correction terms in theorem 14.3 vanish identically.

Proof. For $N = 2$, the reduced zeros are

$$\pi^2 \left(m + \frac{1}{2}\right)^2, \qquad m \geq 0.$$

For $N = 4$, example 11.12 gives the reduced zeros

$$A_4^4 \left(m + \frac{1}{2}\right)^4, \qquad m \geq 0.$$

Summing their negative powers proves the reduced formulas. The absolute formulas follow from

$$\mathfrak{Z}_N^{\mathrm{abs}}(s) = N \mathfrak{Z}_N^{\mathrm{red}}(s/N).$$

□

Definition 14.5. Choose a logarithm for each member of the finite exceptional part of Λ_N. Use the real logarithm for the eventual positive sequence. Define

$$\mathfrak{Z}_N^{\mathrm{orb}}(s) = \sum_{\lambda \in \Lambda_N} e^{-s \operatorname{Log} \lambda}.$$

This series is initially defined on $\Re s > 1/N$.

Corollary 14.6 (Branch-dependent symmetry-orbit continuation)**.** *The branch-dependent orbit zeta $\mathfrak{Z}_N^{\mathrm{orb}}$ continues meromorphically to $\mathbb{C}$. It has the same single pole and residue as $\mathfrak{Z}_N^{\mathrm{red}}$. Changing the logarithms of the finite exceptional zeros changes the continuation by an entire function.*

Proof. The tail consists of positive reduced zeros, so it has the same comparison with the Hurwitz zeta. Only finitely many terms depend on the branch choices. □

Definition 14.7. Let q be a positive integer. Choose logarithms of the simple zeros of D_N with arguments in a fixed bounded interval. Define

$$\mathfrak{Z}_{N,q}^{\mathrm{der}}(s) = \sum_{\substack{\mathsf{D}_N(\rho)=0 \\ \mathsf{D}_N'(\rho)\neq 0}} \frac{e^{-s \operatorname{Log} \rho}}{\mathsf{D}_N'(\rho)^q}.$$

Theorem 14.8 (Derivative-weighted convergence)**.** *If $N \geq 3$ and $q \geq 1$, the series $\mathfrak{Z}_{N,q}^{\mathrm{der}}(s)$ converges locally uniformly for every $s \in \mathbb{C}$. It therefore defines an entire function.*

For $N = 2$, the series converges absolutely and locally uniformly on $\Re s > 1$, where it defines a holomorphic function.

Proof. For $N \geq 3$, theorem 11.7 gives

$$|\mathsf{D}_N'(\rho)|^{-1} \leq C e^{-c_N |\rho|}$$

for every sufficiently large simple zero. Let L be a compact subset of the s plane. Because the chosen arguments of the zeros lie in a fixed bounded interval, there are constants C_L and M_L such that

$$\sup_{s \in L} |e^{-s \operatorname{Log} \rho}| \leq C_L |\rho|^{M_L}$$

for every sufficiently large zero. Hence the absolute value of the corresponding summand is at most

$$C_L' |\rho|^{M_L} e^{-q c_N |\rho|}.$$

The linear zero counting estimate makes this majorant summable. The Weierstrass test gives absolute local uniform convergence of the series of entire summands, so its sum is entire. The finite exceptional set causes no difficulty.

For $N = 2$, the derivative of the cosine has modulus one at every zero. The assertion then follows from the linear zero spacing. □

Remark 14.9. The derivative weighted sum omits the finitely many zeros that might be multiple. A treatment of those zeros requires the complete principal parts of D_N^{-q} rather than the single derivative weight.

15 Scope boundaries and further questions

The large-zero theorem settles eventual simplicity but does not classify the finite exceptional set. A remaining question is whether every zero of D_N is simple and every reduced zero of Δ_N is positive. Both statements hold for $N = 2$ and $N = 4$. The argument here proves them only outside a compact set for general N.

The leading correction in theorem 11.8 may admit iteration to a full exponential expansion for τ_m. Such an expansion is an open problem, not a result of this paper. It could refine the entire correction in the zero-zeta comparison and lead to explicit special values.

The symmetry-orbit zero zeta has a meromorphic continuation with one pole. No functional equation is obtained. A branch-dependent sum over the N roots of each reduced zero must specify logarithms on the rays.

The coefficient families in sections 7, 9 and 10 have exact formal parameter-addition laws. Their arithmetic behavior, including odd-prime congruences, dyadic weighting, and p-adic or Amice–Mahler interpolation, belongs to Paper IX. General bounded-operator semigroup, resolvent, and Feshbach theory belongs to Paper VI. Paper III owns general nonlinear operation trees, while abstract completion and occupancy questions belong to Paper X. None of those theories is inferred from the scalar complex-analytic results proved here.

16 Formalization ledger for version 1.0.0

The released Lean project uses Lean 4 v4.34.0-rc2 and Mathlib revision `aeb72f1e9981b6d35b426b16dca7759be3e7ba06` [dMU21, The20]. Its 151 theorem-map rows resolve to 1164 unique declarations in the namespace `Komorebi.PaperIV`: 953 theorem declarations and 211 constructions. The full namespace audit scans 2234 declarations, including 1999 theorem declarations. A warning-fatal build and the supplied kernel audit allow only `propext`, `Classical.choice`, and `Quot.sound` as standard transitive axioms.

Manuscript records	Compiled Paper IV scope
3.1–3.3	Actual holomorphic root filters, convergent coefficient selection, orthogonal resolution, the exact range–covariance characterization, and the complete pointwise cyclic product law. The formal versions hold as well.
4.1–4.5 and 5.1–5.8	Entire sections, finite exponential formulas, derivative cycles, both Wronskians, and unique entire ODE basis expansions. Both pointwise addition laws and denominator symmetries are checked. The exact ordinary hypergeometric formula follows from the omitted-parameter list, Gamma normalization, and Mathlib's everywhere convergent series.

Manuscript records	Compiled Paper IV scope (continued)
6.1–6.6	Quotient-algebra power bases and actual exponential-series convergence in the finite-dimensional coefficient topology; both pointwise section-coordinate identities. Signed-shift powers, all matrix-exponential entries, group laws and trace are checked. The characteristic polynomial $X^N + 1$, its distinct roots, exact spectrum and determinant one are proved.
7.1–7.8	Named reciprocal families, exact parameter recurrences, Bell formula, parameter degree and leading coefficient, and polynomial divisibility. Parameter addition holds with any single fixed holomorphic logarithm on the stated domain. Newton's recurrence gives the literal Hessenberg determinant; all three compact reciprocal-polynomial formulas are checked.
8.1–8.5	The full tangent coefficient series has the exact kappa and factorial normalization, including $N = 1$. Reciprocal differentiation holds on every stated fixed-logarithm domain. The actual higher Riccati identity holds at every point where the denominator is nonzero.
9.1–9.10	Arbitrary-parameter residue reciprocals and their analytic germs, polynomial degree and leading term, sparse Appell laws, joint convolution, monicity and covariance. The Hasse and derivative-truncation encodings are proved equal; the named Appell polynomial satisfies both the forward action and the exact $\Phi_{N,j}(\partial_x^N)^r$ inverse formula.
10.1–10.4	Exact formal theta reconstruction, polynomial lift and degree, literal divisibility by u and, for even N, by $u + 1$, and the theta Newton recurrence with its first coefficient.
11.2, 11.4–11.11	Two-sided maximum-modulus bounds prove exact order and type. The indicator limsup formula holds in every direction, including ties; dominant-exponential errors and zero-freeness are uniform on compact sets avoiding ties. Near cancellation and the order-two zero set are checked. For $N \geq 3$, gap and remainder bounds, sector confinement, global exhaustion and multiplicity identities give the full large-zero-string theorem. The real leading correction, quadratic exponential error, signed derivative asymptotic, eventual alternating sign and cubic correction are checked.
11.13–11.15	The reduced series is entire, and $\mathsf{D}_N(z) = \Delta_N(z^N)$ preserves multiplicity away from zero. Finite exceptional sets, eventual positive simple reduced zeros, and their stated displacement are checked, including $N = 2$. Both full disk-counting laws retain all exceptional multiplicities. The full-zero reciprocal sums have exact thresholds $s > 1$ and $s > 1/N$, including boundary divergence.
12.1	For every $N \geq 2$, the complete multiplicity-indexed product converges uniformly on compact sets and equals Δ_N. Jensen and Poisson estimates prove rigidity of the actual quotient from derived growth bounds; no growth certificate remains. The original-variable product for D_N is also checked.
12.2–12.4	The actual diagonal operator is trace class, with a basis-independent trace and determinant given by limits of actual finite compressions. Infinite complete homogeneous coefficients are limits of finite monomial sums. Every normalized section has exact order $1/N$; for $N \geq 2$ its full zero product and zero-moment Bell formulas are proved. The actual complex-power germ at zero has the named residue Taylor coefficients.
12.6–12.10	The normalized logarithm on each zero-free disk has the convergent zero-moment series. Its coefficients equal $\kappa_{N,m}$ and satisfy Newton's recurrence. The Mittag–Leffler tangent expansion converges absolutely and locally uniformly away from poles. Tangent coefficients and the named reciprocal and theta families have their exact zero-moment Bell formulas. The full tangent series is identified with the analytic function.

Manuscript records	Compiled Paper IV scope (continued)
13.1–13.4	Analytic division constructs each simple-zero principal part; its leading coefficient is uniquely determined. Removing the principal parts gives a holomorphic remainder and the exact factorial-scaled Cauchy estimate. Both leading-term errors are bounded, with the first absent for $r = 1$. Rotation gives the orbit-reduced main term. The secant formula holds as a ratio limit, with an exponentially vanishing relative error.
14.2–14.4, 14.6	The exact all-zero sums retain analytic multiplicity. Their sharp absolute-convergence domains, boundary divergence and local uniform convergence are checked. Both Hurwitz comparisons, entire corrections, global meromorphic continuations, unique simple poles and exact residues are proved. The order-two/four models have identically zero corrections. Branch choices on the finite exceptional part change the continuation by an entire function and preserve its pole and residue.
14.8	The derivative-weighted series uses every actual simple zero and genuine logarithms with bounded arguments. Absolute and local uniform convergence and an entire sum are proved for $N \geq 3$; for $N = 2$, absolute/local uniform convergence and holomorphy hold on $\Re s > 1$.

The declaration-level theorem map is authoritative. All 68 theorem-like records have their full scoped statements covered: 68 full, zero partial, and zero unmapped. Rows 143–151 close the final fifteen partial records without altering their statements or assumptions. Imported standard fact 11.1 remains imported. Formal identities, normalized analytic germs, and fixed-logarithm domain identities retain their respective stated meanings. Finite computations do not change these proof boundaries.

17 Conclusion

Cyclic projection turns the exponential into a finite differential basis, and the signed cycle produces the distinguished denominator D_N. Its reciprocal powers yield Bell and Newton recurrences, an additive parameter law, Appell systems, and theta polynomials. Reduction by $w = z^N$ then connects the reciprocal coefficients to a genus-zero product and reduced-zero power sums.

Classical two-term localization, specialized to the regular exponent polygon and combined with reflection symmetry, places all sufficiently large zeros on symmetry rays. The displacement, derivative and counting estimates yield the family-specific Hurwitz comparison for the orbit zero zeta.

References

[AAR99] George E. Andrews, Richard Askey, and Ranjan Roy. *Special Functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1999.

[Apo76] Tom M. Apostol. *Introduction to Analytic Number Theory*. Undergraduate Texts in Mathematics. Springer, New York, 1976.

[BM06] Pavel Bleher and Robert Mallison, Jr. Zeros of sections of exponential sums. *International Mathematics Research Notices*, 2006:Article ID 38937, 49 pages, 2006. doi: 10.1155/IMRN/2006/38937.

[Boa54] Ralph P. Boas. *Entire Functions*. Academic Press, New York, 1954.

[Com74] Louis Comtet. *Advanced Combinatorics: The Art of Finite and Infinite Expansions*. D. Reidel Publishing Company, Dordrecht, 1974.

[dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021.

[FS09] Philippe Flajolet and Robert Sedgewick. *Analytic Combinatorics*. Cambridge University Press, Cambridge, 2009.

[Lan31] Rudolph E. Langer. On the zeros of exponential sums and integrals. *Bulletin of the American Mathematical Society*, 37(4):213–239, 1931.

[Lev80] Boris Ya. Levin. *Distribution of Zeros of Entire Functions*, volume 5 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, revised edition, 1980.

[Mor73] Carlos J. Moreno. The zeros of exponential polynomials. I. *Compositio Mathematica*, 26(1):69–78, 1973.

[Mul05] Martin E. Muldoon. Generalized hyperbolic functions, circulant matrices and functional equations. *Linear Algebra and its Applications*, 406:272–284, 2005. doi: 10.1016/j.laa.2005.04.011.

[P.26a] Komorebi P. Projector-resolved seed calculus i: Typed seeds, operator grammars, and exact reconstruction, 2026. Version 1.2.1.

[P.26b] Komorebi P. Projector-resolved seed calculus ii: Polyphase refinement, weighted transfer, and dyadic–odd resolution, 2026. Version 1.2.1.

[P.26c] Komorebi P. Projector-resolved seed calculus iii: Typed edge operators, connection defects, and obstruction packets, 2026. Version 1.0.1.

[Pól20] George Pólya. Geometrisches über die verteilung der nullstellen gewisser ganzer transzendenter funktionen. *Sitzungsberichte der Bayerischen Akademie der Wissenschaften, Mathematisch Physikalische Klasse*, 50:285–290, 1920.

[Rai60] Earl D. Rainville. *Special Functions*. The Macmillan Company, New York, 1960.

[Rom84] Steven Roman. *The Umbral Calculus*, volume 111 of *Pure and Applied Mathematics*. Academic Press, New York, 1984.

[Sch92] Jens Schwaiger. On generalized hyperbolic functions and their characterization by functional equations. *Aequationes Mathematicae*, 43:198–210, 1992. doi: 10.1007/BF01835702.

[The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs*, pages 367–381. Association for Computing Machinery, 2020.

[Tit39] Edward C. Titchmarsh. *The Theory of Functions*. Oxford University Press, Oxford, second edition, 1939.