Nonlinear Packet Calculus, Obstruction Complexes, and Holonomy

By Komorebi P.

Abstract

Let X and Y carry finite complete families of pairwise annihilating idempotents. Every map from X to Y then has exact packet coordinates. We develop the resulting nonlinear calculus. Fréchet derivatives split into packet jets, and composition satisfies a packet form of the Faà di Bruno formula. Polynomial fixed point equations admit convergent rooted tree expansions under an explicit majorant condition. Graded products and Maurer Cartan equations become exact convolutions of packet labels. We define nonlinear refinement defects and prove estimates for iterates, fixed points, and derivatives. A correction map gives a canonical obstruction class in its cokernel. We prove naturality, packet descent, a relative obstruction exact sequence, mapping cone formulations, quantitative nonlinear correction, Lyapunov Schmidt reduction, and complete filtered lifting. For a filtered correction complex, a spectral sequence distinguishes graded obstruction candidates from the classes that survive in the total problem. Exact elimination gives Schur formulas through second order. Fundamental cycle holonomy controls finite transport systems. Smooth complete packet families have a canonical unitary transport, and products of direct packet rotations converge to it.

1. Introduction

A finite family of pairwise annihilating idempotents gives exact coordinates. If

$$P_\alpha P_\beta = 0 \quad \text{for } \alpha \neq \beta, \qquad \sum_\alpha P_\alpha = I,$$

then every vector is the sum of its packet components. Linear operators become block matrices in these coordinates. A nonlinear map cannot be reconstructed from its diagonal restrictions, since mixed interactions between

Keywords: nonlinear packet coordinates, Faà di Bruno formula, obstruction complex, filtered lifting, holonomy

AMS Classification: Primary: 47H10, Secondary: 46T20, 18G35, 05C25, 17B55.

packet components are usually present. It can still be represented exactly if its input is the complete packet tuple.

This article develops the nonlinear representation and the obstruction theory that follows from it. Exact coordinates preserve a nonlinear problem, but they do not solve it. Once the equation is written in packet coordinates, its derivatives, compositions, correction maps, refinement failures, and gluing defects can be separated without discarding mixed terms.

The first part treats the local nonlinear calculus. Packet analysis and synthesis conjugate arbitrary maps. Fréchet derivatives split into multilinear packet jets. The higher chain rule then becomes a Faà di Bruno formula whose terms carry intermediate packet labels. Standard background on Fréchet calculus and the higher chain rule may be found in [1], [2], and [3].

The second part studies nonlinear iteration. Repeated substitution is indexed by rooted trees rather than operator words. A scalar majorant gives an absolute convergence theorem for polynomial fixed point equations. The quadratic case recovers the Catalan series. Rooted tree combinatorics are discussed in [13], [10], and [5].

The third part concerns correction and obstruction. A linear correction map

$$D : U \longrightarrow V$$

places the obstruction to solving $Du = -r$ in the canonical quotient $\operatorname{coker} D$. This quotient has a universal property and behaves naturally under commuting squares. Mapping cones extend the same construction to cochain complexes. In a Hilbert space with closed range, the quotient is represented by the orthogonal projection onto $\operatorname{Ker} D^*$. The Moore Penrose inverse then gives the least norm correction. Relevant background appears in [14] and [12].

A vanishing linear obstruction is only the first step. We give a contraction theorem for nonlinear correction, followed by a local Lyapunov Schmidt reduction. In a complete filtered problem, every lifting step has its own cokernel class. Vanishing at all depths and compatible choices of corrections produce an exact inverse limit solution. A filtered correction complex carries a spectral sequence. Its first page records graded obstruction candidates, while later differentials record interactions that can cancel them. We use the standard spectral sequence of a filtered complex from [14] and [11].

The final part studies elimination and gluing. The implicit function theorem gives an exact Schur derivative after hidden variables are removed. Finite transport systems reduce, after a spanning tree gauge, to one holonomy element for each fundamental cycle. The same idea extends to smooth projector families through Kato transport. We use standard material from [3], [4], [6], [7], and [9].

The article does not claim that every nonlinear problem has a finite obstruction space. The cokernel is canonical only after a correction map has

been specified. It may be infinite dimensional. A first order obstruction may vanish while a higher order obstruction remains. The purpose of the calculus is to identify the exact object in which each failure lives and to show how those objects change under packets, refinement, filtration, and transport.

2. **A map of the nonlinear calculus**

Four different questions recur throughout the paper. They should not be merged into one vague idea of nonlinear structure.

The first question is representation. Given a nonlinear map F, can its packet form be written without approximation? The answer is yes. Analysis and synthesis are inverse coordinate maps, so conjugation preserves the full equation.

The second question is interaction. Which input packets influence a given output packet, and at what derivative order? Packet jets answer this question. They form a directed hypergraph rather than an ordinary block matrix because one nonlinear interaction can have several inputs.

The third question is correction. Suppose an approximate solution leaves a defect r. A declared linearized correction map D removes the part of r lying in $\operatorname{im} D$. The quotient class

$$[r] \in \operatorname{coker} D$$

is the part that no allowed first order correction can remove. This is the obstruction. A nonlinear theorem is still needed after that class vanishes.

The fourth question is gluing. Local coordinate changes can solve every edge problem and still fail around a loop. The remaining cycle product is holonomy. It is an obstruction to one global frame, not an obstruction to solving a single local equation.

These four layers fit into the sequence

$$\begin{aligned}\text{exact coordinates} \longrightarrow \text{packet interactions} \longrightarrow \text{correction classes}\\ \longrightarrow \text{global gluing.}\end{aligned}$$

Each arrow needs hypotheses. Exact coordinates need a complete packet system. Differential interactions need regularity. A cokernel obstruction needs a specified correction map. Holonomy needs invertible local transports.

A second distinction concerns scale. A refinement defect compares a fine nonlinear model with a coarse one. It is a point in the coarse output space. An obstruction class is the image of such a defect, or of another residual, in a quotient by allowed corrections. The two are related only after the correction map has been declared. A small defect can have a nonzero obstruction, and a large defect can have zero obstruction if it lies entirely in the correction range.

The paper uses finite packet systems in the coordinate and graph sections. Banach space hypotheses enter only where norms, contractions, or Fréchet

derivatives are used. Module and complex language is used for cokernels, filtered lifting, and cohomological obstructions. This separation keeps each result at the level of generality supported by its proof.

Standing conventions. All packet systems are finite unless a statement says otherwise. In a normed setting every packet projector and every coarsening operator is bounded. Differentiability of operator valued maps is taken in operator norm. A Banach space isomorphism is a bounded linear bijection with bounded inverse. For a decreasing filtration $(F^m X)$, completeness means that the canonical map

$$X \longrightarrow \varprojlim_m X/F^m X$$

is an isomorphism and separatedness means $\bigcap_m F^m X = 0$. Statements about congruence classes in a filtered nonlinear problem also assume that the nonlinear map preserves filtration congruences at the stated depth. These assumptions are repeated where they are essential to a proof.

3. **Exact nonlinear packet coordinates**

Let X be a module over a unital ring. A finite packet system on X is a family

$$\mathcal{P} = (P_\alpha)_{\alpha \in I} \subseteq \mathrm{End}(X)$$

of nonzero idempotents such that

$$P_\alpha P_\beta = 0 \quad \text{when } \alpha \neq \beta, \qquad \sum_{\alpha \in I} P_\alpha = I_X.$$

Write

$$X_\alpha = P_\alpha X, \qquad X^{\mathcal{P}} = \bigoplus_{\alpha \in I} X_\alpha.$$

The analysis and synthesis maps are

$$\mathcal{A}_{\mathcal{P}} x = (P_\alpha x)_\alpha, \qquad \mathcal{S}_{\mathcal{P}}(x_\alpha)_\alpha = \sum_\alpha x_\alpha.$$

They are inverse isomorphisms.

Definition 3.1 (Nonlinear packet coordinate map). Let $\mathcal{P}$ be a packet system on X and let $\mathcal{Q}$ be a packet system on Y. For any map $F : X \to Y$, define

$$F_{\mathcal{Q} \leftarrow \mathcal{P}} = \mathcal{A}_{\mathcal{Q}} F \mathcal{S}_{\mathcal{P}}.$$

Thus

$$F_{\mathcal{Q} \leftarrow \mathcal{P}}((x_\alpha)_\alpha) = \left(Q_\gamma F \left(\sum_\alpha x_\alpha \right) \right)_\gamma .$$

THEOREM 3.2 (Exact nonlinear coordinate theorem). *For every map* $F : X \to Y$,

$$\mathcal{S}_{\mathcal{Q}} F_{\mathcal{Q}\leftarrow\mathcal{P}} \mathcal{A}_{\mathcal{P}} = F.$$

If $G : Y \to Z$ *and* $\mathcal{R}$ *is a packet system on* Z, *then*

$$(G \circ F)_{\mathcal{R}\leftarrow\mathcal{P}} = G_{\mathcal{R}\leftarrow\mathcal{Q}} \circ F_{\mathcal{Q}\leftarrow\mathcal{P}}.$$

No linearity, continuity, or differentiability assumption is needed.

Proof. The identities $\mathcal{S}_{\mathcal{P}}\mathcal{A}_{\mathcal{P}} = I_X$ and $\mathcal{S}_{\mathcal{Q}}\mathcal{A}_{\mathcal{Q}} = I_Y$ give

$$\mathcal{S}_{\mathcal{Q}} F_{\mathcal{Q}\leftarrow\mathcal{P}} \mathcal{A}_{\mathcal{P}} = \mathcal{S}_{\mathcal{Q}}\mathcal{A}_{\mathcal{Q}} F \mathcal{S}_{\mathcal{P}}\mathcal{A}_{\mathcal{P}} = F.$$

For composition,

$$G_{\mathcal{R}\leftarrow\mathcal{Q}} F_{\mathcal{Q}\leftarrow\mathcal{P}} = \mathcal{A}_{\mathcal{R}} G \mathcal{S}_{\mathcal{Q}}\mathcal{A}_{\mathcal{Q}} F \mathcal{S}_{\mathcal{P}} = \mathcal{A}_{\mathcal{R}} (G \circ F) \mathcal{S}_{\mathcal{P}}.$$

□

COROLLARY 3.3 (Dynamical equivalence). *Suppose* $X = Y$ *and the same packet system is used on both sides. Packet coordinates preserve fixed points, periodic points, injectivity, surjectivity, and invertibility. In particular,*

$$F(x) = x \quad \Longleftrightarrow \quad F_{\mathcal{P}\leftarrow\mathcal{P}}(\mathcal{A}_{\mathcal{P}} x) = \mathcal{A}_{\mathcal{P}} x.$$

Proof. The packet map is conjugate to F through the isomorphism $\mathcal{A}_{\mathcal{P}}$. Every listed property is invariant under conjugacy. □

PROPOSITION 3.4 (Change of packet coordinates). *Let* $\mathcal{P}$ *and* $\mathcal{P}'$ *be packet systems on* X. *Let* $\mathcal{Q}$ *and* $\mathcal{Q}'$ *be packet systems on* Y. *For any two packet systems* $\mathcal{U}$ *and* $\mathcal{V}$ *on the same space, write*

$$J_{\mathcal{U}\leftarrow\mathcal{V}} = \mathcal{A}_{\mathcal{U}} \mathcal{S}_{\mathcal{V}}.$$

Then

$$F_{\mathcal{Q}'\leftarrow\mathcal{P}'} = J_{\mathcal{Q}'\leftarrow\mathcal{Q}} F_{\mathcal{Q}\leftarrow\mathcal{P}} J_{\mathcal{P}\leftarrow\mathcal{P}'}.$$

Proof. Insert $\mathcal{S}_{\mathcal{Q}}\mathcal{A}_{\mathcal{Q}} = I_Y$ and $\mathcal{S}_{\mathcal{P}}\mathcal{A}_{\mathcal{P}} = I_X$ into the definition of $F_{\mathcal{Q}'\leftarrow\mathcal{P}'}$. □

Example 3.5 (A mixed quadratic map). Let $X = Y = \Bbbk^2$ with coordinate packets. For

$$F(x_1, x_2) = (x_1 x_2, x_1^2 + x_2),$$

the first output packet depends on both input packets. The second output packet contains a quadratic contribution from the first input packet and a linear contribution from the second. Packet coordinates record these interactions exactly. Replacing F by the two diagonal restrictions would lose the product $x_1 x_2$.

Remark 3.6. Exact packet coordinates do not imply packet independence. They retain the complete input tuple. Every output packet may depend on every input packet.

4. **Packet jets and higher composition**

Let X and Y be Banach spaces and let the packet projectors on both spaces be bounded. Assume that $F : X \to Y$ is m times continuously Fréchet differentiable. The derivative

$$D^r F(x) : X^r \longrightarrow Y$$

is a continuous symmetric multilinear map.

Definition 4.1 (Packet jet). For $1 \le r \le m$, define

$$J^r_{\gamma|\alpha_1,\dots,\alpha_r} F(x) = Q_\gamma D^r F(x)\big|_{X_{\alpha_1}\times\cdots\times X_{\alpha_r}}.$$

This is a continuous multilinear map

$$X_{\alpha_1} \times \cdots \times X_{\alpha_r} \longrightarrow Y_\gamma.$$

Theorem 4.2 (Exact reconstruction of derivatives). *For $h_1, \dots, h_r \in X$,*

$$D^r F(x)[h_1, \dots, h_r] = \sum_\gamma \sum_{\alpha_1,\dots,\alpha_r} J^r_{\gamma|\alpha_1,\dots,\alpha_r} F(x)[P_{\alpha_1} h_1, \dots, P_{\alpha_r} h_r].$$

Proof. Insert $I_Y = \sum_\gamma Q_\gamma$ on the output and $I_X = \sum_\alpha P_\alpha$ in every input slot. Multilinearity expands the finite sums. □

Proposition 4.3 (Jet norm bound). *The operator norm of each packet jet satisfies*

$$\left\| J^r_{\gamma|\alpha_1,\dots,\alpha_r} F(x) \right\| \le \|Q_\gamma\| \, \|D^r F(x)\| \prod_{j=1}^{r} \|P_{\alpha_j}\|.$$

When all packets are orthogonal projections on Hilbert spaces, every nonzero projector has norm one and the bound reduces to the norm of the full derivative.

Proof. Apply the definition of the operator norm to the composition of $D^r F(x)$ with the input and output projectors. □

Corollary 4.4 (Packet Taylor formula). *Assume F is $m+1$ times continuously Fréchet differentiable on an open set containing the segment $x + th$ for $0 \le t \le 1$. Then*

$$\begin{aligned} Q_\gamma F(x+h) = Q_\gamma F(x) &+ \sum_{r=1}^{m} \frac{1}{r!} \sum_{\alpha_1,\dots,\alpha_r} J^r_{\gamma|\alpha_1,\dots,\alpha_r} F(x)[P_{\alpha_1} h, \dots, P_{\alpha_r} h] \\ &+ Q_\gamma R_{m+1}(x, h), \end{aligned}$$

where the Bochner integral remainder is

$$R_{m+1}(x,h) = \frac{1}{m!}\int_0^1 (1-t)^m D^{m+1}F(x+th)[h,\dots,h]\,dt.$$

Proof. The Banach space Taylor formula with integral remainder gives the unprojected identity. Apply Q_γ to every term and use theorem 4.2 on each derivative. Boundedness of Q_γ allows it to pass through the Bochner integral. □

For a fixed base point, the nonzero packet jets form a directed hypergraph. An order r interaction has r input packet labels and one output packet label. This support depends on the chosen packet system.

For a block $B = \{i_1 < \cdots < i_s\}$ and an intermediate packet γ, write

$$G_{\gamma,B}(x \mid h) = J^s_{\gamma|\alpha_{i_1},\dots,\alpha_{i_s}} G(x)[h_{i_1},\dots,h_{i_s}].$$

THEOREM 4.5 (Packet Faà di Bruno formula). *Let $G : X \to Y$ and $F : Y \to Z$ be n times continuously Fréchet differentiable. Let $\mathcal{P}$, $\mathcal{Q}$, and $\mathcal{R}$ be packet systems on X, Y, and Z. Fix input packets $\alpha_1,\dots,\alpha_n$, an output packet δ, and directions $h_i \in X_{\alpha_i}$. Then*

$$\begin{aligned} &J^n_{\delta|\alpha_1,\dots,\alpha_n}(F\circ G)(x)[h_1,\dots,h_n] \\ &= \sum_{\pi=\{B_1,\dots,B_k\}} \sum_{\gamma_1,\dots,\gamma_k} J^k_{\delta|\gamma_1,\dots,\gamma_k} F(Gx)\big[G_{\gamma_1,B_1}(x \mid h),\dots,G_{\gamma_k,B_k}(x \mid h)\big]. \end{aligned}$$

The first sum runs over all set partitions of $\{1,\dots,n\}$.

Proof. The higher chain rule writes $D^n(F\circ G)(x)$ as a sum over set partitions. Insert $I_Y = \sum_\gamma Q_\gamma$ in every argument of the derivative of F. Insert the input and output packet identities into each derivative of G and into the final output. Group the terms by the partition and by the intermediate packet labels. □

PROPOSITION 4.6 (Number of labeled composition terms). *If Y has M packets, then the number of packet labeled terms at order n before zero jets are removed is*

$$N_{n,M} = \sum_{k=1}^{n} S(n,k) M^k,$$

where $S(n,k)$ is a Stirling number of the second kind. Thus $N_{n,M}$ is the nth Touchard polynomial evaluated at M.

Proof. A set partition with k blocks receives one independent intermediate packet label on each block. It therefore gives M^k labeled terms. There are $S(n,k)$ such partitions. □

Corollary 4.7 (Structural pruning). *A term in theorem* 4.5 *vanishes whenever one of its constituent packet jets vanishes. The support of the composed jet is therefore contained in the set of decorated partitions built from the supports of the jets of* F *and* G*.*

Proof. Every summand is a composition of the displayed packet jets. A zero factor makes the entire summand zero. □

Example 4.8 (The second order composition law). At order two, the partition formula has two shapes. For packet directions h_1 and h_2,

$$Q_\delta D^2(F\circ G)(x)[h_1,h_2] = \sum_{\gamma_1,\gamma_2} J^2_{\delta|\gamma_1,\gamma_2}F(Gx)[J^1_{\gamma_1|\alpha_1}G(x)h_1, J^1_{\gamma_2|\alpha_2}G(x)h_2] + \sum_{\gamma} J^1_{\delta|\gamma}F(Gx)[J^2_{\gamma|\alpha_1,\alpha_2}G(x)[h_1,h_2]].$$

The first line records two first order changes entering a quadratic interaction of F. The second records a genuine second order interaction created inside G before F is applied.

Remark 4.9. Jet sparsity is not invariant under an arbitrary change of coordinates. A useful packet system normally comes from symmetry, scale, support, grading, valuation, or another structure that is already present in the problem. Every sparsity claim should therefore name the packet system with respect to which it is made.

5. **Rooted tree expansions**

Repeated nonlinear substitution is indexed by rooted trees. We begin with a polynomial fixed point equation on a Banach space.

Let X be a Banach space. Fix $s \in X$ and continuous multilinear maps

$$B_r : X^r \longrightarrow X, \qquad 2 \le r \le d.$$

Set

$$a = \|s\|, \qquad b_r = \|B_r\|, \qquad \phi(t) = a + \sum_{r=2}^{d} b_r t^r.$$

An ordered rooted tree has leaves labeled by s and every internal vertex of arity r labeled by B_r. Its value is obtained by evaluating from the leaves toward the root.

Theorem 5.1 (Polynomial tree expansion). *Suppose there is* $R > 0$ *such that*

$$\phi(R) \le R$$

and

$$q = \sum_{r=2}^{d} r b_r R^{r-1} < 1.$$

Then the map

$$\Phi(x) = s + \sum_{r=2}^{d} B_r(x, \dots, x)$$

has a unique fixed point in the closed ball $\overline{B}_R(0)$*. The sum of the values of all finite ordered rooted trees converges absolutely to that fixed point.*

Proof. For $x, y \in \overline{B}_R(0)$, multilinearity and telescoping in the r input positions give

$$\|B_r(x, \dots, x) - B_r(y, \dots, y)\| \le r b_r R^{r-1} \|x - y\|.$$

Hence Φ is q Lipschitz on the ball. The inequality $\phi(R) \le R$ shows that Φ preserves the ball. The contraction mapping theorem gives a unique fixed point u in that ball.

Let $\mathfrak{T}_m$ be the finite set of ordered rooted trees of height less than m. A leaf has value s. If the root of τ has arity r and ordered subtrees $\tau_1, \dots, \tau_r$, define

$$B_\tau(s) = B_r(B_{\tau_1}(s), \dots, B_{\tau_r}(s)).$$

Induction on m shows that the Picard iterate $u_0 = 0$ and $u_{m+1} = \Phi(u_m)$ satisfies

$$u_m = \sum_{\tau \in \mathfrak{T}_m} B_\tau(s).$$

Define a nonnegative scalar weight by assigning weight a to a leaf and applying the scalar map $(t_1, \dots, t_r) \mapsto b_r t_1 \cdots t_r$ at a vertex of arity r. A second induction gives

$$\sum_{\tau \in \mathfrak{T}_m} \|B_\tau(s)\| \le \sum_{\tau \in \mathfrak{T}_m} w(\tau) = t_m,$$

where $t_0 = 0$ and $t_{m+1} = \phi(t_m)$. The sequence (t_m) is increasing because ϕ has nonnegative coefficients. It is bounded by R because $t_0 \le R$ and $\phi(R) \le R$. The sum of the norms of all tree values is therefore finite. The tree series converges absolutely. Its partial sums are the Picard iterates, so its sum is u. □

Corollary 5.2 (Quadratic Catalan expansion). *Let* $B : X \times X \to X$ *be bounded and consider*

$$u = s + B(u, u).$$

Put $a = \|s\|$ *and* $b = \|B\|$*. If* $4ab < 1$*, then*

$$u = \sum_{n \ge 1} \sum_{\substack{\tau \text{ full ordered binary tree} \\ \text{with } n \text{ leaves}}} B_\tau(s)$$

converges absolutely. It also satisfies

$$\|u\| \leq \frac{1-\sqrt{1-4ab}}{2b}$$

when $b > 0$.

Proof. If $b = 0$, the equation is $u = s$ and the tree series contains only its one leaf. If $a = 0$, every tree value is zero. Choose any R with $0 < R < 1/(2b)$ and apply theorem 5.1 to obtain uniqueness of the zero fixed point in that ball.

Assume now that $a > 0$ and $b > 0$. Take

$$R = \frac{1-\sqrt{1-4ab}}{2b}.$$

Then $a + bR^2 = R$ and $2bR < 1$. Apply theorem 5.1. The number of full ordered binary trees with n leaves is the Catalan number C_{n-1}, which gives the displayed majorant. □

THEOREM 5.3 (Packet labeled tree reconstruction). *Let X carry a packet system $\mathcal{P}$. At each internal vertex of a tree, insert the output identity and one input identity for every child. Summing over all internal packet labels reconstructs the original tree value exactly.*

Proof. At a vertex labeled by an r multilinear map B_r, insert

$$I_X = \sum_{\beta} P_\beta$$

on the output and in every input slot. Multilinearity gives an exact finite sum of packet blocks at that vertex. Apply this identity successively from the leaves toward the root. Since the tree and the packet set are finite, all sums are finite and their total reconstructs the original tree value. □

COROLLARY 5.4 (Tree pruning). *A packet labeled tree vanishes if one internal vertex carries a zero packet block of the corresponding multilinear map. Removing all such trees leaves the exact series unchanged.*

Proof. A zero vertex map makes the value of the full tree zero. □

The tree theorem separates combinatorial growth from analytic convergence. The number of trees may grow rapidly, while the norm of each tree decreases through the multilinear bounds. The scalar map ϕ is the exact majorant that balances these two effects. Packet pruning changes the combinatorics but not the proof. It removes trees whose values are already zero before any norm estimate is used.

Example 5.5 (Catalan parity packets). Over a field of characteristic zero, the formal equation

$$u = z + u^2$$

has the branch

$$u(z) = \frac{1 - \sqrt{1 - 4z}}{2}.$$

Write $u = u_0 + u_1$, where u_0 contains the even powers of z and u_1 contains the odd powers. Exact packet projection gives

$$u_0 = u_0^2 + u_1^2, \qquad u_1 = z + 2u_0u_1.$$

The two equations remain coupled. Their sum is the original fixed point equation.

6. **Graded nonlinear operations**

Let G be a finite abelian group and let

$$A = \bigoplus_{g \in G} A_g$$

be a G graded module.

Definition 6.1 (Grade respecting operation). An r multilinear map $\mu : A^r \to A$ respects the grading when

$$\mu(A_{g_1}, \dots, A_{g_r}) \subseteq A_{g_1 + \cdots + g_r}.$$

THEOREM 6.2 (Graded packet convolution). *If $a_j = \sum_{g \in G} a_{j,g}$ and μ respects the grading, then*

$$P_k \mu(a_1, \dots, a_r) = \sum_{g_1 + \cdots + g_r = k} \mu(a_{1,g_1}, \dots, a_{r,g_r}).$$

Proof. Expand every input into its homogeneous components. Each term lands in the component whose label is the sum of its input labels. Projection onto A_k retains exactly the displayed terms. □

COROLLARY 6.3 (Cyclic product formula). *For a cyclically graded algebra*

$$A = \bigoplus_{j \in \mathbb{Z}/N\mathbb{Z}} A_j, \qquad A_i A_j \subseteq A_{i+j},$$

one has

$$P_k(ab) = \sum_{i+j=k} a_i b_j.$$

Proof. Apply theorem 6.2 to multiplication. □

THEOREM 6.4 (Packet Maurer Cartan equations). *Let*

$$\mathcal{L} = \bigoplus_{g \in G} \mathcal{L}_g$$

be a differential graded Lie algebra over a ring in which two is invertible. Assume

$$d\mathcal{L}_g \subseteq \mathcal{L}_g, \qquad [\mathcal{L}_g, \mathcal{L}_h] \subseteq \mathcal{L}_{g+h}.$$

For a degree one element $\eta = \sum_g \eta_g$, the Maurer Cartan equation

$$d\eta + \frac{1}{2}[\eta, \eta] = 0$$

is equivalent to

$$d\eta_k + \frac{1}{2} \sum_{g+h=k} [\eta_g, \eta_h] = 0$$

for every $k \in G$.

Proof. The differential preserves the group label and the bracket adds labels. Apply the projector onto $\mathcal{L}_k$ and use exact reconstruction. □

The same argument applies to higher brackets.

PROPOSITION 6.5 (Packet equations for L_∞ operations). *Let $\mathcal{L}$ be complete and carry operations ℓ_r that respect a finite abelian group grading. Assume the Maurer Cartan series*

$$\sum_{r \geq 1} \frac{1}{r!} \ell_r(\eta, \ldots, \eta)$$

converges and the required factorials are invertible. Then its packet of label k is

$$\sum_{r \geq 1} \frac{1}{r!} \sum_{g_1 + \cdots + g_r = k} \ell_r(\eta_{g_1}, \ldots, \eta_{g_r}).$$

The full Maurer Cartan equation holds if and only if every packet equation vanishes.

Proof. Apply theorem 6.2 to each operation ℓ_r. Completeness permits the convergent sums to be projected term by term. □

Definition 6.6 (Divided square). A divided square on degree one elements is a map q satisfying

$$q(x + y) = q(x) + q(y) + [x, y], \qquad q(ax) = a^2 q(x),$$

whenever the expressions are defined.

PROPOSITION 6.7 (Divided square packet equations). *Assume $q(\mathcal{L}_g) \subseteq \mathcal{L}_{2g}$. Choose an ordering of the finite label set. The equation*

$$d\eta + q(\eta) = 0$$

is equivalent to

$$d\eta_k + \sum_{2g=k} q(\eta_g) + \sum_{\substack{g<h \\ g+h=k}} [\eta_g, \eta_h] = 0$$

for every k.

Proof. Repeated use of the polarization identity for q gives

$$q\left(\sum_g \eta_g\right) = \sum_g q(\eta_g) + \sum_{g<h} [\eta_g, \eta_h].$$

Project onto the component of label k. □

Example 6.8 (Matrix difference grading). Let $A = M_N(\Bbbk)$ and assign the matrix unit E_{rs} the label $r - s$ modulo N. Since

$$E_{rs}E_{uv} = \delta_{su}E_{rv},$$

the output label is

$$(r - s) + (u - v) = r - v$$

whenever the product is nonzero. Matrix multiplication therefore obeys the cyclic convolution law exactly. The same grading resolves commutators and polynomial matrix equations.

Remark 6.9. These results begin with a graded differential or higher algebra. They do not assert that every deformation problem has such a controlling object. Differential graded and infinity algebra methods are developed in [10].

7. **Nonlinear refinement defects**

Let

$$F_f : X_f \longrightarrow Y_f, \qquad F_c : X_c \longrightarrow Y_c$$

be fine and coarse maps. Let

$$C_X : X_f \longrightarrow X_c, \qquad C_Y : Y_f \longrightarrow Y_c$$

be linear coarsening maps. Whenever a norm, a Lipschitz constant, or a Fréchet derivative is used, the spaces are Banach spaces and the coarsening maps are bounded.

Definition 7.1 (Nonlinear refinement defect). The pointwise defect of F is

$$\Delta_F(x) = C_Y F_f(x) - F_c(C_X x).$$

The two models are exactly compatible when Δ_F vanishes everywhere.

The defect is an ordinary vector in Y_c. It becomes an obstruction only after a correction map into Y_c is chosen and the defect is passed to the corresponding cokernel. This distinction matters in applications. Norm estimates measure the size of the defect, while a quotient determines whether the defect is removable.

THEOREM 7.2 (Exact defect under composition). *Let $G_f : Y_f \to Z_f$ and $G_c : Y_c \to Z_c$, with coarsening $C_Z : Z_f \to Z_c$. Then*

$$\Delta_{G\circ F}(x) = \Delta_G(F_f(x)) + G_c\big(F_c(C_X x) + \Delta_F(x)\big) - G_c\big(F_c(C_X x)\big).$$

If G_c is L Lipschitz on the relevant set, then

$$\|\Delta_{G\circ F}(x)\| \le \|\Delta_G(F_f(x))\| + L\|\Delta_F(x)\|.$$

Proof. Add and subtract $G_c(C_Y F_f(x))$. Use

$$C_Y F_f(x) = F_c(C_X x) + \Delta_F(x).$$

The norm estimate follows from the Lipschitz bound. □

COROLLARY 7.3 (Linear product rule). *If all maps in theorem 7.2 are linear, then*

$$\Delta_{GF} = \Delta_G F_f + G_c \Delta_F.$$

Proof. Linearity turns the last difference in theorem 7.2 into $G_c\Delta_F(x)$. □

THEOREM 7.4 (Iterated orbit defect). *Let $F_f : X_f \to X_f$ and $F_c : X_c \to X_c$, with coarsening $C : X_f \to X_c$. Suppose F_c is q Lipschitz. For a fine orbit starting at x, assume*

$$\|\Delta_F(F_f^j x)\| \le \varepsilon_j.$$

Then

$$\Big\|CF_f^n(x) - F_c^n(Cx)\Big\| \le \sum_{j=0}^{n-1} q^{n-1-j}\varepsilon_j.$$

If $0 \le q < 1$ and every $\varepsilon_j \le \varepsilon$, then

$$\Big\|CF_f^n(x) - F_c^n(Cx)\Big\| \le \varepsilon\frac{1-q^n}{1-q}.$$

Proof. Let

$$e_n = \Big\|CF_f^n(x) - F_c^n(Cx)\Big\|.$$

The definition of the defect and the Lipschitz property give

$$e_{n+1} \le \varepsilon_n + q e_n.$$

Iterating this recurrence proves the first estimate. The second is the geometric sum. □

THEOREM 7.5 (Fixed point stability). *Assume F_c is a contraction with constant $q < 1$. Let x_f and x_c be fixed points of F_f and F_c. If*

$$\|\Delta_F(x_f)\| \le \varepsilon,$$

then

$$\|Cx_f - x_c\| \le \frac{\varepsilon}{1-q}.$$

When the defect at x_f is zero, the fixed points are compatible.

Proof. Using the fixed point equations,

$$\begin{aligned}\|Cx_f - x_c\| &\le \|CF_f(x_f) - F_c(Cx_f)\| + \|F_c(Cx_f) - F_c(x_c)\| \\ &\le \varepsilon + q\|Cx_f - x_c\|.\end{aligned}$$

Rearrange. □

COROLLARY 7.6 (Several refinement levels). *Let $C_\ell : X_{\ell+1} \to X_\ell$ for $0 \le \ell < m$. Suppose x_ℓ is a fixed point at level ℓ, the coarse map at level ℓ has contraction constant $q_\ell < 1$, and the adjacent defect at $x_{\ell+1}$ is at most ε_ℓ. Write*

$$C_{a,b} = C_a C_{a+1} \cdots C_{b-1} \quad \text{for } a < b, \qquad C_{a,a} = I.$$

Then

$$\|C_{0,m}x_m - x_0\| \le \sum_{\ell=0}^{m-1} \|C_{0,\ell}\| \frac{\varepsilon_\ell}{1-q_\ell}.$$

Proof. Insert the intermediate vectors $C_{0,\ell}x_\ell$ and telescope. Apply theorem 7.5 to every adjacent pair. □

PROPOSITION 7.7 (Derivative defects). *Assume F_f and F_c are r times Fréchet differentiable and the coarsening maps are bounded and linear. Then*

$$D\Delta_F(x) = C_Y DF_f(x) - DF_c(C_X x)C_X.$$

For $r \ge 2$,

$$D^r\Delta_F(x)[h_1, \dots, h_r] = C_Y D^r F_f(x)[h_1, \dots, h_r] - D^r F_c(C_X x)[C_X h_1, \dots, C_X h_r].$$

Proof. Differentiate the defining identity for Δ_F. The coarsening maps are linear, so their higher derivatives vanish. □

Example 7.8 (Sharp affine bound). Let $F_c(y) = qy + b$ with $0 < q < 1$. Choose a fine fixed point whose coarsening has constant defect ε in the direction of the scalar state. The fixed point mismatch is exactly $\varepsilon/(1-q)$. Thus the constant in theorem 7.5 cannot be improved without more information.

8. **Canonical obstruction objects**

We begin by isolating the part of a defect that cannot be removed by an allowed linear correction.

Definition 8.1 (Linear correction datum). A linear correction datum consists of a homomorphism

$$D : U \longrightarrow V$$

and a defect $r \in V$. A correction is an element $u \in U$ satisfying

$$Du = -r.$$

The obstruction class is

$$\mathrm{Ob}_D(r) = [r] \in \operatorname{coker} D = V/\operatorname{im} D.$$

Theorem 8.2 (Cokernel obstruction criterion). *A linear correction exists if and only if*

$$\mathrm{Ob}_D(r) = 0.$$

The quotient map

$$q_D : V \longrightarrow \operatorname{coker} D$$

has the following universal property. If $\varphi : V \to W$ is linear and $\varphi D = 0$, then there is a unique linear map

$$\overline{\varphi} : \operatorname{coker} D \longrightarrow W$$

with $\varphi = \overline{\varphi} q_D$.

Proof. The class of r is zero exactly when $r \in \operatorname{im} D$, which is equivalent to solvability of $Du = -r$. If $\varphi D = 0$, then φ vanishes on $\operatorname{im} D$. It therefore descends uniquely to the quotient. □

The universal property explains why the cokernel is the canonical obstruction object relative to D. Any linear obstruction test that vanishes on correctable defects factors through it.

Theorem 8.3 (Naturality of obstruction classes). *Suppose*

$$\begin{array}{ccc} U & \xrightarrow{D} & V \\ a \downarrow & & \downarrow b \\ U' & \xrightarrow{D'} & V' \end{array}$$

is a commuting square. Then there is a unique induced map

$$\overline{b} : \operatorname{coker} D \longrightarrow \operatorname{coker} D'$$

satisfying

$$\overline{b}\ \mathrm{Ob}_D(r) = \mathrm{Ob}_{D'}(br).$$

Proof. The relation $bD = D'a$ shows that the quotient map $q_{D'}b$ vanishes on $\operatorname{im} D$. Apply the universal property in theorem 8.2. □

THEOREM 8.4 (Packet decomposition of an obstruction). *Suppose*

$$U = \bigoplus_{\alpha \in I} U_\alpha, \qquad V = \bigoplus_{\alpha \in I} V_\alpha,$$

and $D(U_\alpha) \subseteq V_\alpha$ *for every* α. *Let*

$$D_\alpha : U_\alpha \longrightarrow V_\alpha$$

be the restriction. Then

$$\operatorname{coker} D \cong \bigoplus_{\alpha \in I} \operatorname{coker} D_\alpha.$$

Under this isomorphism, the obstruction class of $r = \sum_\alpha r_\alpha$ *is the tuple*

$$\big([r_\alpha]\big)_\alpha.$$

Hence r *is correctable if and only if every packet obstruction vanishes.*

Proof. The grading assumption gives

$$\operatorname{im} D = \bigoplus_{\alpha} \operatorname{im} D_\alpha.$$

Taking the quotient of the direct sum by this direct sum of images gives the displayed isomorphism. □

Remark 8.5. The conclusion of theorem 8.4 requires the correction map to preserve the packets. Off diagonal correction blocks can couple packet defects. In that case the full cokernel remains canonical, but it need not split into the separate packet cokernels.

PROPOSITION 8.6 (Projectors on the obstruction quotient). *Let* $(Q_\alpha)_{\alpha \in I}$ *be a complete family of pairwise annihilating idempotents on* V. *Assume*

$$Q_\alpha(\operatorname{im} D) \subseteq \operatorname{im} D$$

for every α. *Then each* Q_α *induces an idempotent* $\overline{Q}_\alpha$ *on* $\operatorname{coker} D$, *and*

$$\operatorname{coker} D = \bigoplus_{\alpha \in I} \overline{Q}_\alpha(\operatorname{coker} D).$$

For every defect $r \in V$,

$$\overline{Q}_\alpha \operatorname{Ob}_D(r) = [Q_\alpha r].$$

No packet decomposition of the correction space U *is required.*

Proof. If $v - v' \in \operatorname{im} D$, then

$$Q_\alpha(v - v') \in \operatorname{im} D.$$

Hence $[v] \mapsto [Q_\alpha v]$ is well defined on the quotient. The idempotent, annihilation, and reconstruction identities descend from V to $V/\operatorname{im} D$. The formula for an obstruction class follows from the definition. □

8.1. *Relative obstruction under coarsening.* A coarse problem can forget an obstruction that is still present in the fine problem. The next exact sequence identifies the part that is lost.

THEOREM 8.7 (Relative obstruction exact sequence). *Suppose the following diagram of modules commutes and has exact rows*:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U_0 & \xrightarrow{i_U} & U_f & \xrightarrow{p_U} & U_c & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle D_0} & & \downarrow{\scriptstyle D_f} & & \downarrow{\scriptstyle D_c} & & \\ 0 & \longrightarrow & V_0 & \xrightarrow{i_V} & V_f & \xrightarrow{p_V} & V_c & \longrightarrow & 0. \end{array}$$

There is a natural exact sequence

$$\begin{aligned} 0 \longrightarrow \operatorname{Ker} D_0 \longrightarrow \operatorname{Ker} D_f \longrightarrow \operatorname{Ker} D_c \xrightarrow{\delta} \operatorname{coker} D_0 \\ \longrightarrow \operatorname{coker} D_f \longrightarrow \operatorname{coker} D_c \longrightarrow 0. \end{aligned}$$

For $u_c \in \operatorname{Ker} D_c$, choose $u_f \in U_f$ with $p_U u_f = u_c$. There is a unique $v_0 \in V_0$ such that

$$D_f u_f = i_V v_0,$$

and the connecting map is

$$\delta(u_c) = [v_0].$$

Proof. Since $D_c u_c = 0$, commutativity gives

$$p_V D_f u_f = D_c p_U u_f = 0.$$

Exactness of the lower row gives a unique v_0 with $D_f u_f = i_V v_0$. A different lift has the form $u_f + i_U u_0$. The corresponding element of V_0 changes by $D_0 u_0$, so $[v_0]$ is independent of the lift.

The map $\operatorname{Ker} D_0 \to \operatorname{Ker} D_f$ is injective because i_U is injective. If $u_f \in \operatorname{Ker} D_f$ maps to zero in U_c, then $u_f = i_U u_0$ for a unique u_0. The identity

$$i_V D_0 u_0 = D_f i_U u_0 = 0$$

and injectivity of i_V show that $u_0 \in \operatorname{Ker} D_0$. This proves exactness at $\operatorname{Ker} D_f$.

If u_c comes from an element of $\operatorname{Ker} D_f$, one may choose that element as its lift and obtain $\delta(u_c) = 0$. Conversely, if $\delta(u_c) = 0$, then $v_0 = D_0 u_0$ for some u_0. Replacing u_f by $u_f - i_U u_0$ gives an element of $\operatorname{Ker} D_f$ that maps to u_c. This proves exactness at $\operatorname{Ker} D_c$.

The class $[v_0] \in \operatorname{coker} D_0$ maps to zero in $\operatorname{coker} D_f$ whenever it equals $\delta(u_c)$, since the chosen lift satisfies $i_V v_0 = D_f u_f$. Conversely, if $i_V v_0 = D_f u_f$, then $u_c = p_U u_f$ lies in $\operatorname{Ker} D_c$ and $\delta(u_c) = [v_0]$. This proves exactness at $\operatorname{coker} D_0$.

If a class in $\operatorname{coker} D_f$ comes from $\operatorname{coker} D_0$, its image in $\operatorname{coker} D_c$ is zero. Conversely, suppose $[v_f]$ maps to zero in $\operatorname{coker} D_c$. Choose u_c with $D_c u_c = p_V v_f$ and lift it to $u_f \in U_f$. Then

$$p_V(v_f - D_f u_f) = 0,$$

so $v_f - D_f u_f = i_V v_0$ for some $v_0 \in V_0$. Thus $[v_f]$ comes from $[v_0]$. Surjectivity at the last term follows from surjectivity of p_V. □

Corollary 8.8 (Coarse vanishing and fine vanishing). *Let $[r_f] \in \operatorname{coker} D_f$. If its image in $\operatorname{coker} D_c$ is zero, then $[r_f]$ comes from $\operatorname{coker} D_0$. Coarse vanishing therefore means that the fine obstruction is supported in information removed by coarsening. It does not imply $[r_f] = 0$.*

Proof. This is exactness of the sequence in theorem 8.7 at $\operatorname{coker} D_f$. □

Example 8.9 (A fine obstruction erased by coarsening). Take all three correction spaces to be zero. Let

$$V_f = \Bbbk^2, \qquad V_c = \Bbbk, \qquad p_V(x, y) = x + y,$$

and let

$$V_0 = \operatorname{Ker} p_V = \operatorname{span}\{(1, -1)\}.$$

The fine defect $(1, -1)$ is a nonzero class in $\operatorname{coker} D_f = V_f$, while its coarse image is zero. The class comes from $\operatorname{coker} D_0 = V_0$, exactly as corollary 8.8 predicts.

The same construction has a cohomological form.

Definition 8.10 (Mapping cone). Let $D : C^\bullet \to E^\bullet$ be a cochain map. Its mapping cone is

$$\operatorname{Cone}(D)^n = E^n \oplus C^{n+1}$$

with differential

$$d_{\operatorname{Cone}}(e, c) = (d_E e + Dc, -d_C c).$$

Theorem 8.11 (Cohomological obstruction through the cone). *Let $e \in E^n$ satisfy $d_E e = 0$. The class of $(e, 0)$ in*

$$H^n(\operatorname{Cone}(D))$$

vanishes if and only if the class $[e] \in H^n(E)$ lies in the image of

$$H^n(D) : H^n(C) \longrightarrow H^n(E).$$

When C and E are concentrated in degree zero,

$$H^0(\mathrm{Cone}(D)) \cong \operatorname{coker} D.$$

Proof. The class of $(e, 0)$ vanishes exactly when there are $e' \in E^{n-1}$ and $c \in C^n$ with

$$(e, 0) = d_{\mathrm{Cone}}(e', c).$$

This means

$$e = d_E e' + Dc, \qquad d_C c = 0.$$

Equivalently, $[e] = H^n(D)[c]$. The degree zero statement follows directly from the definition of the cone differential. □

Corollary 8.12 (Packet obstruction complexes). *If $C^\bullet$ and $E^\bullet$ split into finite packet subcomplexes and D preserves those packets, then* $\mathrm{Cone}(D)$ *splits into packet subcomplexes and*

$$H^n(\mathrm{Cone}(D)) \cong \bigoplus_\alpha H^n(\mathrm{Cone}(D_\alpha)).$$

Proof. Both the cone and its cohomology commute with finite direct sums. □

Example 8.13 (A two component obstruction). Let

$$D : \Bbbk \longrightarrow \Bbbk^2, \qquad D(t) = (t, t).$$

The cokernel is one dimensional. The defect (a, b) is correctable exactly when $a = b$. The obstruction can be represented by $a - b$, but this scalar representative depends on a choice of isomorphism from the cokernel to $\Bbbk$. The quotient class itself is canonical.

9. **Split Hilbert obstructions and nonlinear correction**

The quotient obstruction has a concrete orthogonal representative when the correction map acts between Hilbert spaces and has closed range.

Theorem 9.1 (Orthogonal obstruction projector). *Let $D : H_1 \to H_2$ be a bounded operator between Hilbert spaces with closed range. Let $D^\dagger$ be its Moore Penrose inverse. Then*

$$P_{\mathrm{ob}} = I_{H_2} - DD^\dagger$$

is the orthogonal projector onto

$$(\operatorname{im} D)^\perp = \operatorname{Ker} D^*.$$

For $r \in H_2$, the following are equivalent:

(1) *$Du = -r$ is solvable*
(2) *$[r] = 0$ in* $\operatorname{coker} D$

(3) $P_{\mathrm{ob}} r = 0$.

When these conditions hold, $-D^{\dagger} r$ is the least norm correction. For every r,

$$\operatorname{dist}(r, \operatorname{im} D) = \|P_{\mathrm{ob}} r\|.$$

Proof. Closed range gives the orthogonal decomposition

$$H_2 = \operatorname{im} D \oplus \operatorname{Ker} D^{*}.$$

The Moore Penrose identities show that $DD^{\dagger}$ is the orthogonal projector onto $\operatorname{im} D$. The stated equivalences and distance formula follow from this decomposition. The least norm property is the standard characterization of $D^{\dagger} r$. □

A vanishing linear obstruction does not by itself solve a nonlinear equation. The next theorem gives a quantitative condition that does.

THEOREM 9.2 (Right inverse correction theorem). *Let $D : X \to Y$ be a bounded linear map between Banach spaces. Assume $Z = \operatorname{im} D$ is closed and there is a bounded right inverse*

$$R : Z \longrightarrow X, \qquad DR = I_Z.$$

Put $X_R = R(Z)$. Let

$$N : \overline{B}_{\rho}(0) \cap X_R \longrightarrow Z$$

be L Lipschitz. Let $y \in Z$ and put

$$q = \|R\| L.$$

Assume $q < 1$ and

$$\|R(y - N(0))\| \leq (1 - q)\rho.$$

Then the equation

$$Dx + N(x) = y$$

has a unique solution in $\overline{B}_{\rho}(0) \cap X_R$.

Proof. The space X_R is closed. Indeed, if $Rz_n \to x$, then $z_n = DRz_n \to Dx$, so $x = R(Dx)$. Define

$$\Psi(x) = R(y - N(x)).$$

This map takes X_R to itself. For $x \in \overline{B}_{\rho}(0) \cap X_R$,

$$\begin{aligned} \|\Psi(x)\| &\leq \|R(y - N(0))\| + \|R\| \, \|N(x) - N(0)\| \\ &\leq (1 - q)\rho + q\rho = \rho. \end{aligned}$$

Also,

$$\|\Psi(x) - \Psi(x')\| \leq q \|x - x'\|.$$

The contraction mapping theorem gives a unique fixed point in the stated set. Applying D to the fixed point identity gives the equation. Conversely, a solution in X_R equals $R(Dx)$ and is therefore a fixed point of Ψ. □

COROLLARY 9.3 (Stability with respect to the target). *Under the hypotheses of theorem* 9.2, *let* x *and* x' *be the fixed points associated with targets* y *and* y'. *If both fixed point maps preserve the same ball, then*

$$\|x - x'\| \leq \frac{\|R\|}{1-q}\|y - y'\|.$$

Proof. Subtract the two fixed point equations and use the Lipschitz bound for N:

$$\|x - x'\| \leq \|R\|\|y - y'\| + q\|x - x'\|.$$

□

The range equation can be separated from the remaining nonlinear obstruction. Standard forms of this reduction appear in [8] and [15].

THEOREM 9.4 (Local Lyapunov Schmidt reduction). *Let* $\mathcal{U} \subseteq X$ *be an open neighborhood of zero and let* $F : \mathcal{U} \to Y$ *be a* C^r *map between Banach spaces with* $r \geq 1$, $F(0) = 0$, *and* $L = DF(0)$. *Assume there are closed decompositions*

$$X = K \oplus X_1, \qquad Y = R \oplus O,$$

where $K = \operatorname{Ker} L$, $R = \operatorname{im} L$, *and*

$$L_1 = L|_{X_1} : X_1 \longrightarrow R$$

is a Banach space isomorphism. Let π_R *and* π_O *be the two projections on* Y. *Then there are neighborhoods* $U_K \subseteq K$ *and* $U_1 \subseteq X_1$ *of zero and a unique* C^r *map*

$$\psi : U_K \longrightarrow U_1$$

with $\psi(0) = 0$ *such that*

$$\pi_R F(k + \psi(k)) = 0$$

for every $k \in U_K$. *After the neighborhoods are made smaller if needed, the solutions of* $F(x) = 0$ *near zero are exactly the points*

$$x = k + \psi(k)$$

for which the reduced obstruction map

$$\kappa(k) = \pi_O F(k + \psi(k))$$

vanishes. At the base point,

$$D\psi(0) = 0, \qquad D\kappa(0) = 0.$$

Proof. Define

$$G(k, x_1) = \pi_R F(k + x_1).$$

The derivative of G in the X_1 variable at $(0, 0)$ is L_1, which is invertible. The implicit function theorem gives the unique map ψ. Every nearby x has a unique

decomposition $x = k + x_1$. The range equation forces $x_1 = \psi(k)$, and the remaining equation is precisely $\kappa(k) = 0$.

Differentiate $G(k, \psi(k)) = 0$ at zero. Since L vanishes on K, one obtains $L_1 D\psi(0) = 0$, hence $D\psi(0) = 0$. Finally,

$$D\kappa(0) = \pi_O L|_K = 0.$$

□

COROLLARY 9.5 (Quadratic reduced obstruction). *Assume the hypotheses of theorem* 9.4 *with* $r \geq 2$. *For* $k_1, k_2 \in K$,

$$D^2\kappa(0)[k_1, k_2] = \pi_O D^2 F(0)[k_1, k_2].$$

Proof. Differentiate $\kappa(k) = \pi_O F(k + \psi(k))$ twice. The terms containing $D\psi(0)$ vanish. The term containing $D^2\psi(0)$ is multiplied by $\pi_O L$, which is zero because $L(X) = R$. □

Example 9.6 (A purely quadratic obstruction). Let

$$F(x, y) = (y, x^2)$$

on $\mathbb{R}^2$. At the origin,

$$L(h, k) = (k, 0).$$

The kernel is the x axis, the range is the first output axis, and the cokernel is the second output axis. The range equation forces $y = 0$. The reduced obstruction map is

$$\kappa(x) = x^2.$$

The linear obstruction vanishes for every kernel direction, but the nonlinear equation still has only the solution $x = 0$ near the origin.

Remark 9.7. The reduced map κ begins at quadratic order. This is why vanishing of the first order cokernel class need not settle a nonlinear problem. Higher terms can produce a genuine obstruction on the kernel directions.

10. **Filtered lifting**

Let X and Y be modules with decreasing filtrations

$$X = F^0X \supseteq F^1X \supseteq \cdots, \qquad Y = F^0Y \supseteq F^1Y \supseteq \cdots.$$

Assume both are separated and complete for their filtration topologies. Let $\Phi : X \to Y$ be continuous and filtration congruence preserving in the sense that

$$h \in F^mX \quad \Longrightarrow \quad \Phi(x + h) - \Phi(x) \in F^mY$$

for every $x \in X$ and every m. This condition makes the statement $\Phi(x) \in F^mY$ depend only on the class of x modulo F^mX.

Suppose x_m satisfies

$$\Phi(x_m) \in F^m Y.$$

Assume that corrections $h \in F^m X$ obey a first order congruence

$$\Phi(x_m + h) \equiv \Phi(x_m) + L_m h \pmod{F^{m+1}Y},$$

where $L_m : F^m X \to F^m Y$ is a module homomorphism satisfying

$$L_m(F^{m+1}X) \subseteq F^{m+1}Y$$

and therefore induces

$$\overline{L}_m : F^m X/F^{m+1}X \longrightarrow F^m Y/F^{m+1}Y.$$

At depth m, only the associated graded correction is visible. The class below is therefore finite whenever the two graded pieces have finite length, even if the completed problem is infinite. Its vanishing answers one precise question: whether the current approximation can be improved by one level.

Definition 10.1 (Depth obstruction). The depth m obstruction of x_m is

$$\mathrm{Ob}_m(x_m) = [\Phi(x_m)] \in \operatorname{coker} \overline{L}_m,$$

where $\Phi(x_m)$ is first viewed in $F^m Y/F^{m+1}Y$.

Theorem 10.2 (One step lifting criterion). *The approximation x_m has a correction $h_m \in F^m X$ such that*

$$\Phi(x_m + h_m) \in F^{m+1}Y$$

if and only if

$$\mathrm{Ob}_m(x_m) = 0.$$

When a correction exists, the set of its classes in

$$F^m X/F^{m+1}X$$

is a coset of $\operatorname{Ker} \overline{L}_m$. *If the graded pieces are vector spaces, this coset is an affine space.*

Proof. The next depth condition is equivalent to

$$\overline{L}_m[h_m] = -[\Phi(x_m)]$$

in $F^m Y/F^{m+1}Y$. This is solvable exactly when the displayed defect has zero cokernel class. If h_m and h'_m are two solutions, their difference lies in $\operatorname{Ker} \overline{L}_m$. Adding an element of the kernel to one solution gives another, so the solution set is a coset of the kernel. □

COROLLARY 10.3 (Finite branching count). *Suppose the graded pieces are finite vector spaces over a finite field. If the depth obstruction vanishes, then the number of correction classes at depth m is*

$$|\operatorname{Ker} \overline{L}_m|.$$

If the obstruction does not vanish, the number is zero.

Proof. A nonempty affine space over a finite vector space has the same number of elements as that vector space. □

LEMMA 10.4 (Well defined lift classes). *Under the filtration congruence assumption, the condition*

$$\Phi(x) \in F^m Y$$

depends only on the class of x in $X/F^m X$. Fix a representative x_m of a depth m class. The map

$$[h] \longmapsto [x_m + h]_{m+1}$$

identifies correction classes $[h] \in F^m X/F^{m+1}X$ with the classes at depth $m+1$ that reduce to $[x_m]_m$. Under this identification, the solutions of the associated graded correction equation in theorem 10.2 are exactly the children whose defect lies in $F^{m+1}Y$. A different representative of the same parent gives a translated parametrization of the same set of children.

Proof. If $x' - x \in F^m X$, then filtration congruence preservation gives

$$\Phi(x') - \Phi(x) \in F^m Y.$$

Thus one defect lies in $F^m Y$ exactly when the other does. If $h - h' \in F^{m+1}X$, then $x_m + h$ and $x_m + h'$ define the same depth $m+1$ class. Conversely, two depth $m+1$ classes reducing to $[x_m]_m$ differ from x_m by elements of $F^m X$, and they are equal precisely when those elements agree modulo $F^{m+1}X$. The first order congruence shows that such a child has defect in $F^{m+1}Y$ exactly when its correction class solves the associated graded equation. Replacing x_m by another representative translates the correction coordinate by an element of $F^m X/F^{m+1}X$ and leaves the child classes themselves unchanged. □

Definition 10.5 (Lift tree). Fix an initial approximation. A vertex at depth m is a class $[x]_m \in X/F^m X$ with $\Phi(x) \in F^m Y$ that reduces to the fixed initial class. A depth $m+1$ vertex is joined to its reduction in $X/F^m X$. By lemma 10.4, this rooted graph is independent of the chosen representatives. It is called the lift tree.

THEOREM 10.6 (Finitely branching lift tree). *Assume every level of the lift tree is nonempty and every vertex has only finitely many children. Then the tree contains an infinite branch. If X and Y are separated and complete and Φ is continuous, that branch determines an exact solution of $\Phi(x) = 0$.*

Proof. The first assertion is König's infinity argument. The root has finitely many children. Since vertices occur at arbitrarily large depths, at least one child has descendants at arbitrarily large depths. Choose such a child and repeat the argument at every chosen vertex. This gives one compatible vertex at each depth.

Choose representatives recursively. Once x_m represents the selected depth m vertex, choose a representative x_{m+1} of its selected child with

$$x_{m+1} - x_m \in F^m X.$$

The resulting sequence is Cauchy for the filtration topology. Each representative satisfies $\Phi(x_m) \in F^m Y$. The conclusion now follows from theorem 10.8. □

Corollary 10.7 (Unique lift branch). *Suppose an initial approximation is fixed, every depth obstruction vanishes along its descendants, and every map $\overline{L}_m$ has zero kernel. Then at most one compatible lift branch extends the initial approximation.*

Proof. By theorem 10.2, the correction classes at each depth form an affine space under $\operatorname{Ker} \overline{L}_m$. A zero kernel leaves at most one correction class, so induction gives at most one branch. □

Theorem 10.8 (Complete filtered lifting). *Assume there is a compatible sequence*

$$x_{m+1} \equiv x_m \pmod{F^m X}$$

with

$$\Phi(x_m) \in F^m Y$$

for every m. Then x_m converges to an element $x_\infty \in X$ and

$$\Phi(x_\infty) = 0.$$

In particular, if the obstruction vanishes at every depth and compatible corrections are chosen, an exact solution exists.

Proof. For every fixed r, the congruences imply

$$x_n - x_m \in F^r X$$

whenever $n \geq m \geq r$. Thus (x_m) is Cauchy for the filtration topology. Completeness gives a limit x_∞.

Fix r. For every $m \geq r$, one has $\Phi(x_m) \in F^m Y \subseteq F^r Y$. The submodule $F^r Y$ is open and therefore closed in the filtration topology. Continuity of Φ gives $\Phi(x_m) \to \Phi(x_\infty)$, so $\Phi(x_\infty) \in F^r Y$. Since this holds for every r and Y is separated,

$$\Phi(x_\infty) \in \bigcap_r F^r Y = 0.$$

□

THEOREM 10.9 (Naturality of filtered obstructions). *Suppose a morphism of filtered lifting problems sends x_m to x'_m, intertwines L_m with L'_m, and sends $\Phi(x_m)$ to $\Phi'(x'_m)$ on the associated graded pieces. Then the induced cokernel map sends*

$$\mathrm{Ob}_m(x_m) \quad \textit{to} \quad \mathrm{Ob}'_m(x'_m).$$

Proof. Apply theorem 8.3 to the induced commuting square on the depth m graded pieces. □

Example 10.10 (Three lifting outcomes). The scalar equation $x^2 - 2 = 0$ at $x = 3$ modulo 7 has invertible derivative. It has one correction modulo 49 and gives $x = 10$ modulo 49.

For $x^2 - 3 = 0$ at $x = 0$ modulo 3, the derivative is zero and the defect class is nonzero. No correction exists.

For $x^2 = 0$ at $x = 0$ modulo 3, the derivative and defect both vanish. Every correction digit lies in the kernel, so the lift branches.

PROPOSITION 10.11 (Maurer Cartan lifting obstruction). *Let $\mathcal{L}$ be a complete filtered differential graded Lie algebra over a ring in which two is invertible. Assume*

$$dF^i\mathcal{L} \subseteq F^i\mathcal{L}, \qquad [F^i\mathcal{L}, F^j\mathcal{L}] \subseteq F^{i+j}\mathcal{L}.$$

For a degree one element η, put

$$\Omega(\eta) = d\eta + \frac{1}{2}[\eta, \eta].$$

Suppose $\Omega(\eta) \in F^m\mathcal{L}^2$ with $m \geq 1$. For every $h \in F^m\mathcal{L}^1$,

$$\Omega(\eta + h) \equiv \Omega(\eta) + d_\eta h \pmod{F^{m+1}\mathcal{L}^2}, \qquad d_\eta = d + [\eta, \cdot].$$

The obstruction to improving η by one depth is therefore the class of $\Omega(\eta)$ in the cokernel of

$$\mathrm{gr}^m(d_\eta) : \mathrm{gr}^m \mathcal{L}^1 \longrightarrow \mathrm{gr}^m \mathcal{L}^2.$$

Proof. Direct expansion gives

$$\Omega(\eta + h) = \Omega(\eta) + dh + [\eta, h] + \frac{1}{2}[h, h].$$

The filtration rule places $[h, h]$ in $F^{2m}\mathcal{L}^2$, which is contained in $F^{m+1}\mathcal{L}^2$ because $m \geq 1$. The congruence follows. The last statement is theorem 10.2 applied to the induced graded correction map. □

11. **Filtered obstruction complexes**

A direct packet decomposition and a filtration behave differently. In a direct sum of subcomplexes, cohomology splits packet by packet. In a filtration, a class visible at one graded depth can be killed by a later differential.

There are two different mechanisms for a coarse obstruction to vanish. The first is packetwise vanishing. Every fine class is already zero. The second is cancellation through compatibility. Fine classes are nonzero, but later differentials pair them and remove them before the total cohomology is reached. A direct sum decomposition allows only the first mechanism. A filtration can allow both.

Standard Fact 11.1 (Spectral sequence of a finite filtered complex). Let $C^\bullet$ be a cochain complex with a finite decreasing filtration by subcomplexes

$$C^\bullet = F^0C^\bullet \supseteq F^1C^\bullet \supseteq \cdots \supseteq F^{N+1}C^\bullet = 0.$$

There is a natural spectral sequence with

$$E_0^{p,q} = \mathrm{gr}^p\, C^{p+q}, \qquad E_1^{p,q} = H^{p+q}(\mathrm{gr}^p\, C^\bullet),$$

which converges to the associated graded object of $H^{p+q}(C^\bullet)$ with its induced filtration. A filtered cochain map induces a morphism of spectral sequences.

This is the standard filtered complex construction. We use it in the form given in [14] and [11]. When $C^\bullet$ is a correction or deformation complex, the groups on its pages can be read as graded obstruction candidates. For an arbitrary filtered complex they are simply cohomology classes of the associated graded problem.

Proposition 11.2 (Fine obstruction survival). *A class on the first page contributes to the total obstruction only if it is represented by a permanent cycle, so it is neither killed nor identified as a boundary on any later page. If the spectral sequence degenerates at the first page, then $H^n(C^\bullet) = 0$ exactly when $E_1^{p,n-p} = 0$ for every p.*

Proof. Convergence identifies the last page with the associated graded object of the total cohomology. A first page class contributes to that object exactly when it determines a nonzero class on the last page. If the sequence degenerates at the first page, the associated graded pieces of $H^n(C^\bullet)$ are the groups $E_1^{p,n-p}$. A finite filtered module is zero exactly when all its associated graded pieces are zero. □

Corollary 11.3 (Direct packet complexes). *Suppose*

$$C^\bullet = \bigoplus_{\alpha \in I} C_\alpha^\bullet$$

is a finite direct sum of subcomplexes. Then

$$H^n(C^\bullet) \cong \bigoplus_{\alpha \in I} H^n(C_\alpha^\bullet).$$

A total cohomology class vanishes if and only if every packet class vanishes.

Proof. The differential acts independently on the summands. Kernels and images therefore split as finite direct sums, and so does their quotient. □

Example 11.4 (A fine class killed at the next page). Let $C^0 = \Bbbk x$ and $C^1 = \Bbbk y$, with differential $d(x) = y$. Give C^0 the filtration

$$F^0C^0 = \Bbbk x, \qquad F^1C^0 = 0,$$

and give C^1 the filtration

$$F^0C^1 = F^1C^1 = \Bbbk y, \qquad F^2C^1 = 0.$$

The associated graded differential on the first page is zero in the relevant positions, so both x and y appear as nonzero fine classes. The next differential sends the class of x to the class of y. The total complex is acyclic. Thus nonzero fine obstruction classes can cancel through a later refinement differential.

Remark 11.5. The spectral sequence does not guarantee that a total obstruction vanishes. It records which fine classes survive and which are removed by compatibility across depths.

12. **Nonlinear elimination**

Split the variables and equations as

$$x = (u, v), \qquad F(u, v) = \begin{pmatrix} F_P(u, v) \\ F_Q(u, v) \end{pmatrix}.$$

Assume F is twice continuously Fréchet differentiable near (u_0, v_0) and

$$A = D_v F_Q(u_0, v_0)$$

is invertible. The implicit function theorem gives a local map $v = \varphi(u)$ with

$$F_Q(u, \varphi(u)) = 0.$$

Define the reduced equation

$$R(u) = F_P(u, \varphi(u)).$$

Theorem 12.1 (Exact local elimination). *After the neighborhoods are chosen small enough, the equation*

$$F(u, v) = 0$$

is equivalent to

$$v = \varphi(u) \qquad \text{and} \qquad R(u) = 0.$$

Thus eliminating the hidden variable loses no local solutions.

Proof. The hidden equation $F_Q(u, v) = 0$ has the unique local solution $v = \varphi(u)$ by the implicit function theorem. After this identity is imposed, the remaining component of F is exactly $R(u)$. □

THEOREM 12.2 (Nonlinear Schur derivative). *At* (u_0, v_0),

$$DR = D_u F_P - D_v F_P A^{-1} D_u F_Q.$$

Proof. Differentiate $F_Q(u, \varphi(u)) = 0$:

$$D_u F_Q + AD\varphi = 0.$$

Thus

$$D\varphi = -A^{-1} D_u F_Q.$$

Substitute this into

$$DR = D_u F_P + D_v F_P D\varphi.$$

□

The second derivative records the first nonlinear feedback from the hidden variables.

THEOREM 12.3 (Second Schur derivative). *For retained directions* h, k, *set*

$$\widehat{h} = (h, D\varphi h), \qquad \widehat{k} = (k, D\varphi k).$$

Then

$$D^2\varphi[h, k] = -A^{-1} D^2 F_Q[\widehat{h}, \widehat{k}]$$

and

$$D^2 R[h, k] = D^2 F_P[\widehat{h}, \widehat{k}] - D_v F_P A^{-1} D^2 F_Q[\widehat{h}, \widehat{k}].$$

All derivatives on the right are evaluated at (u_0, v_0).

Proof. Differentiate the identity $F_Q(u, \varphi(u)) = 0$ twice. The chain rule gives

$$D^2 F_Q[\widehat{h}, \widehat{k}] + AD^2\varphi[h, k] = 0.$$

This proves the first formula. Differentiate $R(u) = F_P(u, \varphi(u))$ twice and substitute the first formula. □

COROLLARY 12.4 (Packet jets of the reduced equation). *If the retained and hidden spaces carry packet systems, then the first and second packet jets of* R *are obtained by projecting the formulas in theorems* 12.2 *and* 12.3. *Hidden packet feedback is carried by the factors involving* A^{-1}.

Proof. Apply the packet projectors to the input and output slots of the two derivative formulas. □

Example 12.5 (Two retained variables and one hidden variable). Suppose the derivative of F at a solution has block form

$$DF = \begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

where D is invertible. Eliminating the hidden variable replaces the retained derivative A by

$$A - BD^{-1}C.$$

The feedback term is exact. Keeping only A would ignore the path from retained variables into the hidden variable and back.

Remark 12.6. Elimination and obstruction reduction address different situations. The Schur formula assumes that the hidden derivative is invertible. Lyapunov Schmidt reduction allows a nontrivial kernel and cokernel, then leaves a reduced obstruction equation on the kernel directions.

13. **Finite transport and holonomy**

Let $\Gamma = (V, E)$ be a finite connected graph and fix a vector space X. At every vertex take one labeled copy of X. Every oriented edge $u \to v$ carries an invertible transport

$$T_{vu} \in \mathrm{GL}(X), \qquad T_{uv} = T_{vu}^{-1}.$$

Equivalently, one may begin with separate but isomorphic vertex fibers and choose one frame in each fiber. The matrices below are then the transports in those frames. For a path $\gamma = (v_0, \dots, v_r)$, set

$$T_\gamma = T_{v_r v_{r-1}} \cdots T_{v_1 v_0}.$$

For a loop, this product is its holonomy. Standard graph terminology may be found in [4].

Local solvability appears here as the existence of every edge map. Global solvability asks whether all local frames can be chosen at once. A spanning tree has no cycles, so its edge data can always be removed by a gauge. Every remaining edge closes a loop and carries the full global obstruction for that loop.

Definition 13.1 (Gauge change). A gauge is a family $G_v \in \mathrm{GL}(X)$. It changes the edge transports to

$$T'_{vu} = G_v T_{vu} G_u^{-1}.$$

Proposition 13.2 (Gauge covariance). *For a loop γ based at v_0,*

$$\mathrm{Hol}'(\gamma) = G_{v_0} \,\mathrm{Hol}(\gamma) G_{v_0}^{-1}.$$

Thus the conjugacy class of each loop holonomy is gauge invariant.

Proof. Every intermediate gauge factor cancels with its inverse. The initial and terminal factors remain at the base vertex. □

THEOREM 13.3 (Spanning tree normalization). *Choose a root and a spanning tree $\mathcal{T}$ of Γ. There is a gauge in which every tree edge transport is the identity. In that gauge, the transport on each edge outside the tree is the holonomy of its fundamental cycle, with the orientation used to define that cycle.*

Proof. Set the gauge at the root to the identity. Propagate gauges along the unique tree paths so that every tree edge becomes the identity. An edge outside the tree closes one fundamental cycle. Since every tree factor is now the identity, the remaining edge transport is the cycle product. □

THEOREM 13.4 (Cycle gluing criterion). *There are vertex coordinates in which every edge transport is the identity if and only if every fundamental cycle holonomy is the identity.*

Proof. If all edge transports are identity maps, every loop product is the identity. Conversely, use theorem 13.3. Identity fundamental holonomy forces every edge outside the tree to become the identity as well. □

COROLLARY 13.5 (Number of independent cycle tests). *For a connected graph with $|V|$ vertices and $|E|$ unoriented edges, a spanning tree leaves*

$$|E| - |V| + 1$$

fundamental cycle holonomies.

Proof. A spanning tree has $|V| - 1$ edges. Every remaining edge creates one fundamental cycle. □

13.1. *Intertwining two transport systems.* Let A be a transport system on copies of a fixed vector space V_0, and let B be a transport system on copies of a fixed vector space W_0. Fix a root r and an isomorphism

$$U_r : V_0 \longrightarrow W_0.$$

THEOREM 13.6 (Fundamental holonomy intertwining criterion). *The following conditions are equivalent.*

(1) *There is a family of isomorphisms $U_v : V_0 \to W_0$ with the prescribed root value U_r and satisfying*

$$U_v A_{vu} = B_{vu} U_u$$

for every oriented edge $u \to v$.

(2) *For every fundamental loop γ based at r,*

$$U_r \operatorname{Hol}_A(\gamma) = \operatorname{Hol}_B(\gamma) U_r.$$

(3) *The same identity holds for every loop based at r.*

When these conditions hold, the family (U_v) is uniquely determined by U_r.

Proof. If the edge intertwiners exist, multiply their equations around a loop. Every intermediate vertex map cancels. This proves that the first condition implies the third, and the third implies the second.

Conversely, choose a spanning tree. Starting from U_r, propagate U_v along the unique tree path by the rule

$$U_v = B_{vu} U_u A_{vu}^{-1}$$

on every tree edge. This is the only possible extension satisfying the tree edge equations. An edge outside the tree closes one fundamental loop. Its intertwining equation holds exactly when U_r intertwines the two corresponding fundamental holonomies. The second condition therefore supplies every remaining edge equation. Uniqueness follows from propagation along the tree. □

COROLLARY 13.7 (Scalar holonomy bridge). *Suppose every vertex space is one dimensional. Two transport systems are gauge equivalent if and only if their holonomies agree on a fundamental cycle basis.*

Proof. Choose a nonzero root isomorphism. Scalar loop maps commute with it, so theorem 13.6 reduces to equality of the cycle scalars. □

THEOREM 13.8 (Gauge classification of transports). *Let $G \subseteq \mathrm{GL}(X)$ be the chosen structure group and fix a root r. Restrict both edge transports and gauge values to G. Gauge equivalence classes of G valued transport systems are naturally identified with*

$$\mathrm{Hom}(\pi_1(\Gamma, r), G)/G,$$

where G acts by conjugation. After a spanning tree and orientations of the remaining edges are chosen, this set is identified with simultaneous conjugacy classes of

$$b_1 = |E| - |V| + 1$$

tuples in G.

Proof. A transport system assigns a group element to every oriented edge, with inversion under reversal. The product along a based loop gives a homomorphism from the free group $\pi_1(\Gamma, r)$ to G. A gauge change conjugates this homomorphism by the gauge value at the root. Conversely, normalize a transport system on a spanning tree. The remaining edge values prescribe the images of a free basis of $\pi_1(\Gamma, r)$, and every homomorphism arises in this way. A further gauge preserving the tree normalization is constant at all vertices and acts by simultaneous conjugation. The number of non tree edges is b_1. □

COROLLARY 13.9 (Abelian cohomology class). *When the structure group A is abelian, gauge classes of transports are naturally*

$$H^1(\Gamma, A) \cong \mathrm{Hom}(H_1(\Gamma, \mathbb{Z}), A).$$

After a basis of $H_1(\Gamma, \mathbb{Z})$ is chosen, this group is identified with A^{b_1}.

Proof. An edge assignment is an A valued one cochain. A gauge change multiplies it by a coboundary. A graph has no two cells, so every one cochain is a cocycle. The usual computation of the first cohomology of a connected graph gives A^{b_1}. □

Corollary 13.10 (Projective transport). *The same normalization and gluing statements hold for transports valued in a projective linear group. A projective transport is globally trivial exactly when all fundamental holonomies are identity elements in that projective group.*

Proof. Apply theorems 13.3 and 13.4 with the projective linear group as the structure group. □

Example 13.11 (A square loop). A square graph has four vertices and four edges, so it has one fundamental cycle. After a tree gauge, three edges are identities and the fourth is the full loop holonomy. The system glues to one global frame exactly when that remaining matrix is the identity.

14. **Smooth projector transport**

Finite graph transport has a continuous counterpart. Let H be a Hilbert space and let

$$\mathcal{P}(t) = (P_\alpha(t))_{\alpha \in I}$$

be a finite family of complete orthogonal packet systems that is continuously differentiable in operator norm. Thus

$$P_\alpha(t)P_\beta(t) = 0 \quad \text{for } \alpha \neq \beta, \qquad \sum_\alpha P_\alpha(t) = I.$$

Define the canonical generator

$$K(t) = \sum_\alpha \dot{P}_\alpha(t)P_\alpha(t).$$

Lemma 14.1 (Complete packet commutator identity). *For every packet label β,*

$$\dot{P}_\beta(t) = [K(t), P_\beta(t)].$$

The generator is also skew adjoint:

$$K(t)^* = -K(t).$$

Proof. Differentiating $P_\alpha^2 = P_\alpha$ gives

$$P_\alpha \dot{P}_\alpha P_\alpha = 0.$$

Differentiating $P_\beta P_\alpha = 0$ for $\alpha \neq \beta$ gives

$$P_\beta \dot{P}_\alpha = -\dot{P}_\beta P_\alpha.$$

Hence

$$KP_\beta = \dot{P}_\beta P_\beta$$

and

$$P_\beta K = -\dot{P}_\beta (I - P_\beta).$$

Their difference is $\dot{P}_\beta$. Also,

$$K + K^* = \sum_\alpha (\dot{P}_\alpha P_\alpha + P_\alpha \dot{P}_\alpha) = \sum_\alpha \dot{P}_\alpha = 0.$$

□

THEOREM 14.2 (Canonical packet transport). *Let $U(t)$ solve*

$$\dot{U}(t) = K(t)U(t), \qquad U(0) = I.$$

Then $U(t)$ is unitary and transports every packet exactly:

$$P_\alpha(t)U(t) = U(t)P_\alpha(0)$$

for all α.

Proof. Skew adjointness of $K(t)$ gives

$$\frac{d}{dt}(U(t)^*U(t)) = 0,$$

so $U(t)$ is unitary. By lemma 14.1,

$$\frac{d}{dt}\big(U(t)^{-1}P_\alpha(t)U(t)\big) = U(t)^{-1}\big(\dot{P}_\alpha - [K, P_\alpha]\big)U(t) = 0.$$

The expression equals $P_\alpha(0)$ at time zero. □

COROLLARY 14.3 (One moving subspace). *For one orthogonal projection $P(t)$ and its complement, the canonical generator is*

$$K(t) = [\dot{P}(t), P(t)].$$

The transport in theorem 14.2 *maps* $\operatorname{im} P(0)$ *unitarily onto* $\operatorname{im} P(t)$.

Proof. Apply the complete packet formula to $P(t)$ and $I - P(t)$, then simplify the sum. □

Definition 14.4 (Direct rotation of nearby packet systems). Let $(P_\alpha)_{\alpha\in I}$ and $(Q_\alpha)_{\alpha\in I}$ be complete orthogonal packet systems on the same Hilbert space. Put

$$S(\mathcal{Q}, \mathcal{P}) = \sum_{\alpha\in I} Q_\alpha P_\alpha.$$

When $S(\mathcal{Q}, \mathcal{P})$ is invertible, define

$$\mathcal{U}(\mathcal{Q}, \mathcal{P}) = S(\mathcal{Q}, \mathcal{P})\big(S(\mathcal{Q}, \mathcal{P})^* S(\mathcal{Q}, \mathcal{P})\big)^{-1/2}.$$

PROPOSITION 14.5 (Exact direct packet rotation). *If*

$$\max_{\alpha\in I}\|P_\alpha - Q_\alpha\| < 1,$$

then $S(\mathcal{Q},\mathcal{P})$ is invertible. The operator $\mathcal{U}(\mathcal{Q},\mathcal{P})$ is unitary and satisfies

$$Q_\alpha \mathcal{U}(\mathcal{Q},\mathcal{P}) = \mathcal{U}(\mathcal{Q},\mathcal{P})P_\alpha$$

for every α.

Proof. The identities

$$Q_\alpha S(\mathcal{Q},\mathcal{P}) = S(\mathcal{Q},\mathcal{P})P_\alpha = Q_\alpha P_\alpha$$

follow from pairwise orthogonality. Also

$$S^*S = \sum_\alpha P_\alpha Q_\alpha P_\alpha, \qquad SS^* = \sum_\alpha Q_\alpha P_\alpha Q_\alpha.$$

Put $\delta = \max_\alpha \|P_\alpha - Q_\alpha\|$. On $P_\alpha H$,

$$P_\alpha Q_\alpha P_\alpha = P_\alpha - P_\alpha(P_\alpha - Q_\alpha)P_\alpha,$$

so this block is bounded below by $1-\delta > 0$. The same argument on $Q_\alpha H$ applies to $Q_\alpha P_\alpha Q_\alpha$. Thus both S^*S and SS^* are invertible. It follows that S is injective and surjective. The polar factor $\mathcal{U} = S(S^*S)^{-1/2}$ is unitary. Since S^*S commutes with every P_α, the intertwining identity for S passes to $\mathcal{U}$. □

LEMMA 14.6 (One step direct rotation estimate). *Assume $\mathcal{P}(t)$ is twice continuously differentiable in operator norm on $[0,T]$. There are constants $h_0 > 0$ and $C > 0$ such that, whenever $t, t+h \in [0,T]$ and $|h| \le h_0$,*

$$\|\mathcal{U}(\mathcal{P}(t+h),\mathcal{P}(t)) - I - hK(t)\| \le Ch^2.$$

The constants can be chosen uniformly in t.

Proof. Taylor's theorem in the Banach space of bounded operators gives

$$P_\alpha(t+h) = P_\alpha(t) + h\dot{P}_\alpha(t) + E_\alpha(t,h), \qquad \|E_\alpha(t,h)\| \le C_1h^2,$$

uniformly in t and α. Hence

$$S(\mathcal{P}(t+h),\mathcal{P}(t)) = I + hK(t) + E(t,h), \qquad \|E(t,h)\| \le C_2h^2.$$

Since $K(t)^* = -K(t)$,

$$S^*S = I + E_2(t,h), \qquad \|E_2(t,h)\| \le C_3h^2.$$

For small h, the spectrum of S^*S lies in a fixed compact subinterval of $(0,\infty)$. The functional calculus for $z^{-1/2}$ gives

$$(S^*S)^{-1/2} = I + E_3(t,h), \qquad \|E_3(t,h)\| \le C_4h^2.$$

Multiplying the last two expansions proves the estimate. □

THEOREM 14.7 (Direct rotations converge to packet transport). *Assume the complete packet family $\mathcal{P}(t)$ is twice continuously differentiable in operator norm on $[0,T]$. For a partition*

$$0 = t_0 < t_1 < \cdots < t_N = T$$

with sufficiently small mesh, define

$$V_\pi = \mathcal{U}(\mathcal{P}(t_N), \mathcal{P}(t_{N-1})) \cdots \mathcal{U}(\mathcal{P}(t_1), \mathcal{P}(t_0)).$$

Then

$$V_\pi \longrightarrow U(T)$$

in operator norm, where U is the canonical transport from theorem 14.2. On a fixed compact interval the error is bounded by a constant times the mesh.

Proof. Let $U(t,s)$ be the exact propagator generated by K. Since K is continuously differentiable, the integral equation for $U(t+h,t)$ gives a uniform estimate

$$\|U(t+h,t) - I - hK(t)\| \le C_1 h^2.$$

By lemma 14.6, the direct rotation V_j from t_j to t_{j+1} therefore satisfies

$$\|V_j - U(t_{j+1}, t_j)\| \le C_2 h_j^2, \qquad h_j = t_{j+1} - t_j.$$

Both V_j and the exact propagator are unitary. A telescoping product comparison gives

$$\|V_\pi - U(T,0)\| \le \sum_j \|V_j - U(t_{j+1}, t_j)\| \le C_2 \sum_j h_j^2 \le C_2 T |\pi|.$$

This proves norm convergence and the stated first order global bound. □

THEOREM 14.8 (Convergence of sampled transports). *Assume K is Lipschitz on $[0,T]$. For a partition*

$$0 = t_0 < t_1 < \cdots < t_N = T,$$

define

$$U_\pi = \exp\big((t_N - t_{N-1})K(t_{N-1})\big) \cdots \exp\big((t_1 - t_0)K(t_0)\big).$$

As the mesh $|\pi|$ tends to zero,

$$U_\pi \longrightarrow U(T)$$

in operator norm. There is a constant C, depending only on T, the supremum norm of K, and its Lipschitz constant, such that

$$\|U_\pi - U(T)\| \le C|\pi|.$$

Proof. On each interval, compare the exact propagator with the frozen exponential. Variation of constants and the Lipschitz bound give a local error bounded by a constant times the square of the interval length. Every factor is unitary, so multiplication does not amplify norms. Summing the local errors gives a bound by a constant times

$$\sum_j (t_{j+1} - t_j)^2 \leq T|\pi|.$$

□

Corollary 14.9 (Continuous holonomy from graph products). *If $P_\alpha(0) = P_\alpha(T)$ for every α, then $U(T)$ preserves every initial packet. Its restriction to each packet is the holonomy of the closed packet path. The products in theorems* 14.7 *and* 14.8 *converge to these holonomies. The direct rotation product has the added property that every factor maps each sampled packet exactly to its successor.*

Proof. The transport identity in theorem 14.2 gives packet preservation at the endpoint. The two convergence statements follow from theorems 14.7 and 14.8. □

Example 14.10 (A rotating line). In $\mathbb{R}^2$, let

$$v(t) = (\cos t, \sin t)$$

and let $P(t)$ be the orthogonal projection onto the line spanned by $v(t)$. The canonical transport is the ordinary rotation matrix

$$U(t) = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}.$$

It carries the initial line and its orthogonal complement to the two moving packets. Products of short rotations converge to the same transport.

Standard Fact 14.11 (Curvature of a projected connection). Let P be a smooth orthogonal projector valued map on a parameter manifold. The connection

$$\nabla = P\, d$$

on the range bundle has curvature

$$\mathcal{F} = P(dP \wedge dP)P.$$

Proof. For a section s satisfying $Ps = s$,

$$\nabla^2 s = P\, d(P\, ds) = P(dP) \wedge ds.$$

Since $ds = (dP)s + P ds$ and $P(dP)P = 0$, the surviving term is

$$P(dP \wedge dP)Ps.$$

□

15. **Realization data and limits of the calculus**

The continuous theorem transports a declared projector family. It does not construct the family, prove a continuum limit for a physical model, or show that its holonomy is trivial. The curvature formula in standard fact 14.11 is classical. Relating it to global holonomy still requires geometric information as in [6], [7], and [9].

To apply the results in another setting, one must identify the following objects:

(1) an ambient nonlinear equation or continuation problem
(2) the allowed correction space
(3) a linear correction map or a correction complex
(4) the defect whose class is to be tested
(5) a packet decomposition or filtration compatible with the correction data
(6) transition maps between refinement levels
(7) transport maps when local solutions must be glued.

Once these objects are fixed, the cokernel and cone constructions are functorial by theorems 8.3 and 8.11. Packet splitting and descent follow from theorem 8.4, proposition 8.6, and corollary 8.12. Relative loss under coarsening is measured by theorem 8.7. Filtered lifting follows from theorems 10.2, 10.6 and 10.8. Gluing and comparison of transport systems follow from theorems 13.4 and 13.6. Smooth packet transport follows from theorems 14.2 and 14.7.

Several limits remain explicit. Exact packet coordinates are a representation theorem, not a general solution theorem. The cokernel is canonical relative to the correction map, but the correction map itself belongs to the problem data. The cokernel may be infinite dimensional. A split Hilbert projector requires closed range. Coarse vanishing can erase a nonzero fine obstruction. A vanishing linear obstruction does not remove higher nonlinear terms. A spectral sequence records possible cancellation, but it does not force cancellation. Finite graph holonomy and smooth projector transport do not establish a continuum field theory.

These results do not assert that all obstructions vanish. Nonlinear continuation failures can be placed in exact quotient or cohomology objects whenever a correction problem is specified. Those objects admit packet decomposition, filtered refinement, natural transport, and holonomy tests under explicit hypotheses.

Use of language model tools

The author used an OpenAI language model to help reorganize and draft parts of this article. The model suggested formulations and proof approaches

for the polynomial tree theorem in section 5, the cokernel, relative obstruction, and mapping cone results in section 8, the Lyapunov Schmidt and filtered lifting arguments, the holonomy intertwining theorem in section 13, and the direct packet rotation theorem in section 14. It was also used during a later consistency audit to locate missing regularity and filtration hypotheses. The author checked the statements, proofs, and citations and accepts full responsibility for them. The model is not an author.

References

[1] R. Abraham, J. E. Marsden, and T. Ratiu, *Manifolds*, *Tensor Analysis*, *and Applications*, second ed., *Applied Mathematical Sciences* **75**, Springer, New York, 1988.

[2] G. M. Constantine and T. H. Savits, A multivariate Faà di Bruno formula with applications, *Transactions of the American Mathematical Society* **348** no. 2 (1996), 503–520.

[3] K. Deimling, *Nonlinear Functional Analysis*, Springer, Berlin, 1985.

[4] J. L. Gross and T. W. Tucker, *Topological Graph Theory*, John Wiley and Sons, New York, 1987.

[5] E. Hairer, C. Lubich, and G. Wanner, *Geometric Numerical Integration*: *Structure Preserving Algorithms for Ordinary Differential Equations*, second ed., Springer, Berlin, 2006.

[6] T. Kato, On the adiabatic theorem of quantum mechanics, *Journal of the Physical Society of Japan* **5** (1950), 435–439.

[7] T. Kato, *Perturbation Theory for Linear Operators*, Springer, Berlin, 1995.

[8] H. Kielhöfer, *Bifurcation Theory*: *An Introduction with Applications to Partial Differential Equations*, second ed., Springer, New York, 2012.

[9] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry. Volume I*, John Wiley and Sons, New York, 1963.

[10] J. L. Loday and B. Vallette, *Algebraic Operads*, Springer, Berlin, 2012.

[11] J. McCleary, *A User's Guide to Spectral Sequences*, second ed., Cambridge University Press, Cambridge, 2001.

[12] R. Penrose, A generalized inverse for matrices, *Proceedings of the Cambridge Philosophical Society* **51** (1955), 406–413.

[13] R. P. Stanley, *Enumerative Combinatorics*. *Volume* 1, second ed., *Cambridge Studies in Advanced Mathematics* **49**, Cambridge University Press, Cambridge, 2012.

[14] C. A. Weibel, *An Introduction to Homological Algebra*, Cambridge University Press, Cambridge, 1994.

[15] E. Zeidler, *Nonlinear Functional Analysis and Its Applications*. *I*: *Fixed Point Theorems*, Springer, New York, 1986.

Komorebi P.