Projector-Resolved Seed Calculus VI

Transfer Semigroups: Stochastic, Bayesian, Fractional, and PDE Realizations

Komorebi P.

Abstract

This paper develops the transfer-semigroup realization of the projector-resolved seed calculus. A seed is propagated by a one-parameter semigroup, a Markov kernel, a normalized Bayesian transfer, a fractional spectral power, a Green operator, or a potential-weighted Feynman–Kac operator. Exact packet projectors resolve every transfer into input-output blocks, and the semigroup law becomes block convolution over intermediate packets. Coarse and fine generators are compared by a refinement defect. A Duhamel identity proves that this generator-level defect is exactly the source of the corresponding semigroup defect and gives a quantitative stability bound. In the stochastic realization, point-mass seeds generate random walks and continuous-time Markov evolution; exact lumpability is the vanishing of the same refinement defect. Conditional expectation is treated as a weighted orthogonal packet projector, with nested partitions producing martingale-difference packets. Bayesian conditioning is a normalized nonlinear operator. Its independent-data composition law, packet-evidence decomposition, stability bound, conjugate updates, and continuous likelihood flow are proved explicitly. The Bayesian refinement conjecture is resolved in finite dimensions: exact commutation with coarsening is equivalent to blockwise-constant likelihoods up to global scale, and every failure factors through an explicit information-loss packet. Fractional powers are constructed packetwise through the spectral projectors of a positive operator, and the square-root semigroup is recovered by subordination. The fractional-naturality conjecture is also resolved for bounded nonnegative self-adjoint refinement systems: the fractional intertwining defect has an exact resolvent integral, a gap-free Hölder certificate, a sharper common-gap estimate, and a Duhamel semigroup bound. Green kernels, finite-difference Poisson problems, and second-order low-mode convergence provide the PDE realization. A finite-state Feynman–Kac formula is proved by path enumeration. Exact Schur–Feshbach elimination shows that discarded latent packets return as a frequency-dependent self-energy and, in time coordinates, as a memory kernel. Survival and hazard functions are generated by transient semigroups. Nineteen code-generated figures, machine-readable results, and thirty-seven automated tests accompany the paper. Finite-dimensional theorems, classical infinite-dimensional extensions, numerical certificates, and open continuum claims are kept separate.

Scope and claim boundary. The paper proves the stated finite-dimensional semigroup, packet, Markov, Bayesian, conditional-expectation, fractional spectral, Green-kernel, Feynman–Kac, Schur–Feshbach, and survival identities. It also proves quantitative refinement-defect formulas under explicit norm hypotheses, including exact transfer of a bounded generator defect to every fractional power $0 < \alpha < 1$ for bounded nonnegative self-adjoint generators. It does not claim that an arbitrary infinite-dimensional operator generates a semigroup, that every stochastic process admits a finite packet representation, that every posterior family is conjugate, that finite Bayesian sufficiency automatically extends to infinite disintegrations, or that every PDE continuum limit follows from a finite test. Domain, closability, sectoriality, measurability, tightness, and compactness hypotheses remain necessary in their respective infinite-dimensional settings. Classical sources are retained in the separate end-of-series reference ledger rather than reproduced in this paper.

Contents

1 The transfer-semigroup realization **4**

2 Packet blocks of a semigroup **5**

3 Refinement defects and the Duhamel identity **6**
3.1 Resolvent form 7

4 Markov seeds and stochastic transfer **8**

5 Markov lumping as exact refinement **9**

6 Conditional expectation as a stochastic projector **9**

7 Bayesian conditioning as a normalized nonlinear operator **11**
7.1 Packet evidence and reconstruction 12

8 Exact Bayesian refinement and information-loss packets **13**

9 Bayesian flows, conjugacy, and hidden states **16**
9.1 Hidden Markov filtering 18

10 Fractional powers as spectral packet transfer **18**
10.1 Square-root subordination 19
10.2 Fractional naturality under a bounded refinement defect 20

11 Green operators and the Poisson seed **22**
11.1 Low-mode convergence 22

12 Finite-state Feynman–Kac transfer **23**

13 Exact elimination of latent packets **24**
13.1 Time-domain memory 25

14 Survival and hazard semigroups **26**

15 One common operator language **28**

16 Numerical certificates **28**

17 Remaining conjectural extensions **28**

18 Computational implementation **30**
18.1 Representative algorithms 30

19 Conclusion **31**

List of Figures

1 The common operator language used throughout the paper. Semigroups, resolvents, normalized Bayesian maps, and fractional powers are distinct realizations of the same transfer stage. . . . 4
2 Norms of the packet blocks $P_\beta e^{tA} P_\alpha$. Off-diagonal entries quantify transfer between packets. . . . 6
3 Exact generator intertwining gives a vanishing semigroup defect. A controlled perturbation produces a nonzero defect propagated by the Duhamel identity. . . . 7
4 The point mass δ_0 propagated by repeated symmetric random-walk transfer. . . . 8
5 Exact lumping agrees at every horizon. A small generator-level or one-step defect creates a controlled coarse discrepancy. . . . 10
6 Hermite–Gaussian packets, which diagonalize the Ornstein–Uhlenbeck semigroup. The finite numerical release verifies their Gaussian orthogonality through quadrature. . . . 11
7 A finite prior and posterior split into two state packets. Evidence changes both the local packet shape and the global packet mass. . . . 12
8 The Bayesian refinement square commutes exactly when the likelihood is constant on each coarsening fiber, up to one global scale. . . . 15
9 When packet sufficiency fails, the exact defect factors through the difference between the conditional effective likelihood and the declared coarse likelihood. . . . 16
10 Continuous likelihood accumulation produces a nonlinear semigroup on the probability simplex. . . . 17
11 A conjugate Bayesian update viewed as exact transfer of exponent packets. . . . 17
12 A hidden Markov filter alternates a linear prediction semigroup with nonlinear Bayesian normalization. . . . 18
13 Different spectral exponents produce different transfer profiles from the same point seed. 19
14 A generator-level refinement defect transfers quantitatively to fractional powers and then to fractional semigroups. The two upper curves are theorem certificates, not fitted error bars. . . . 21
15 A section of the continuum Green kernel and the scaled discrete inverse. They agree at the grid nodes. . . . 23
16 The first finite-difference eigenvalue converges with observed order approximately two. 24
17 Potential-weighted transfer values computed by the finite Feynman–Kac operator. Exhaustive path enumeration agrees with the matrix formula in the supplied tests. . . . 25
18 The norm of the exact memory kernel produced by eliminated packets. . . . 26
19 Survival and instantaneous hazard produced by one transient transfer semigroup. . . . 27

List of Tables

1 The transfer-semigroup dictionary in the common operator language. . . . 27
2 Selected results from the machine-readable numerical record. . . . 29

1 The transfer-semigroup realization

Let X be a finite-dimensional real or complex vector space, let $A \in \mathrm{End}(X)$, and let $s \in X$ be a primitive seed. The basic continuous transfer is

$$T(t) = e^{tA}, \qquad t \geq 0,$$

while a positive operator L is more naturally written

$$H(t) = e^{-tL}.$$

Both satisfy the semigroup law. The sign convention changes with the domain: Markov generators and evolution equations are usually written with e^{tA}; positive diffusion and elliptic operators are usually written with e^{-tL}.

The common operator chain is

$$\boxed{\text{seed} \longrightarrow \text{transfer} \longrightarrow \text{project} \longrightarrow \text{condition} \longrightarrow \text{refine} \longrightarrow \text{complete} \longrightarrow \text{reconstruct}.}$$




Figure 1: The common operator language used throughout the paper. Semigroups, resolvents, normalized Bayesian maps, and fractional powers are distinct realizations of the same transfer stage.

Definition 1.1 (Transfer-semigroup seed system)**.** A finite transfer-semigroup seed system is a tuple

$$\mathfrak{S} = (X, s, A, \{P_\alpha\}_{\alpha \in I}, \tau),$$

where X is the state space, s is the seed, A is a transfer generator, the P_α are pairwise orthogonal idempotents summing to the identity, and τ specifies the topology in which a completion or limit is interpreted.

Theorem 1.2 (Finite semigroup law)**.** *For every matrix A and all $s, t \in \mathbb{R}$,*

$$\boxed{e^{(s+t)A} = e^{sA} e^{tA}.}$$

Moreover, $T(0) = I$ and $T'(t) = AT(t) = T(t)A$.

Proof. Both exponentials are absolutely convergent matrix power series. Since every power of A commutes with every other power, the Cauchy product gives

$$e^{sA} e^{tA} = \sum_{m,n \geq 0} \frac{s^m t^n}{m! n!} A^{m+n} = \sum_{r \geq 0} \frac{(s+t)^r}{r!} A^r.$$

Termwise differentiation is valid in finite dimension and gives the derivative identities. □

Boundary 1.3 (Infinite-dimensional generators). For an unbounded operator, writing e^{tA} is not a definition unless a semigroup-generation theorem applies. The finite matrix identities in this paper remain algebraically correct models, but a continuum realization must state the operator domain, closure, growth bound, and topology of convergence.

2 Packet blocks of a semigroup

Let

$$P_\alpha P_\beta = \delta_{\alpha\beta} P_\alpha, \qquad \sum_{\alpha \in I} P_\alpha = I.$$

For any operator B, define

$$B_{\beta\alpha} = P_\beta B P_\alpha.$$

Then

$$B = \sum_{\alpha,\beta} B_{\beta\alpha}.$$

For the semigroup, write

$$T_{\beta\alpha}(t) = P_\beta e^{tA} P_\alpha.$$

Theorem 2.1 (Packet semigroup convolution). *For every s, t and every pair of packets α, β,*

$$\boxed{T_{\beta\alpha}(s+t) = \sum_{\gamma \in I} T_{\beta\gamma}(s) T_{\gamma\alpha}(t).}$$

Thus the semigroup law becomes ordinary matrix multiplication in the packet indices.

Proof. Insert the resolution of the identity between the two semigroup factors:

$$\begin{aligned} P_\beta e^{(s+t)A} P_\alpha &= P_\beta e^{sA} \left(\sum_\gamma P_\gamma \right) e^{tA} P_\alpha \\ &= \sum_\gamma (P_\beta e^{sA} P_\gamma)(P_\gamma e^{tA} P_\alpha). \end{aligned}$$

□

Corollary 2.2 (Reducing packets). *If $P_\beta A P_\alpha = 0$ for every $\beta \neq \alpha$, then*

$$P_\beta e^{tA} P_\alpha = 0 \quad (\beta \neq \alpha)$$

for every t. Each packet evolves independently under $P_\alpha A P_\alpha$.

Proof. The hypothesis makes A block diagonal. Every power A^m is block diagonal with respect to the same projectors, and therefore so is the exponential series. □

Remark 2.3 (Path expansion). Expanding the exponential and inserting $I = \sum_\alpha P_\alpha$ between every pair of generators gives

$$P_{\alpha_m} A^m P_{\alpha_0} = \sum_{\alpha_1, \ldots, \alpha_{m-1}} P_{\alpha_m} A P_{\alpha_{m-1}} \cdots P_{\alpha_1} A P_{\alpha_0}.$$

Every coefficient is therefore a sum over packet-transfer paths. The numerical library verifies that truncating this path series converges to the ordinary matrix exponential.




Figure 2: Norms of the packet blocks $P_\beta e^{tA} P_\alpha$. Off-diagonal entries quantify transfer between packets.

3 Refinement defects and the Duhamel identity

Let X_f and X_c be fine and coarse spaces, with generators A_f and A_c, and let

$$C : X_f \to X_c$$

be a coarsening map. Define the generator-level refinement defect

$$\boxed{\Delta = CA_f - A_c C.}$$

The corresponding semigroup defect is

$$\mathfrak{D}(t) = Ce^{tA_f} - e^{tA_c} C.$$

Theorem 3.1 (Exact Duhamel refinement identity)**.** *For every $t \geq 0$,*

$$\boxed{Ce^{tA_f} - e^{tA_c} C = \int_0^t e^{(t-r)A_c} \Delta e^{rA_f}\, dr.}$$

In particular, $\Delta = 0$ implies exact semigroup intertwining at every time.

Proof. Set

$$F(r) = e^{(t-r)A_c} C e^{rA_f}.$$

Then

$$F'(r) = e^{(t-r)A_c} (CA_f - A_c C) e^{rA_f}.$$

Integrating from 0 to t gives $F(t) - F(0) = Ce^{tA_f} - e^{tA_c} C$ and proves the formula. □

Corollary 3.2 (Norm bound)**.** *Suppose*

$$\left\|e^{tA_f}\right\| \le M_f e^{\omega_f t}, \qquad \left\|e^{tA_c}\right\| \le M_c e^{\omega_c t}.$$

Then

$$\|\mathfrak{D}(t)\| \le M_f M_c \|\Delta\| \int_0^t e^{\omega_c(t-r)} e^{\omega_f r}\, dr.$$

For contraction semigroups,

$$\boxed{\|\mathfrak{D}(t)\| \le t \|\Delta\|.}$$




Figure 3: Exact generator intertwining gives a vanishing semigroup defect. A controlled perturbation produces a nonzero defect propagated by the Duhamel identity.

3.1 Resolvent form

For z in the common resolvent set, let

$$R_f(z) = (zI - A_f)^{-1}, \qquad R_c(z) = (zI - A_c)^{-1}.$$

Theorem 3.3 (Resolvent refinement identity)**.**

$$\boxed{CR_f(z) - R_c(z)C = R_c(z)\Delta R_f(z).}$$

Proof. Multiply the left side by $zI - A_c$ on the left and $zI - A_f$ on the right. The result is

$$(zI - A_c)C - C(zI - A_f) = CA_f - A_cC = \Delta.$$

Multiplication by the two resolvents recovers the stated identity. □

4 Markov seeds and stochastic transfer

Use column probability vectors. A finite continuous-time Markov generator Q has nonnegative off-diagonal entries and column sums zero. The stochastic semigroup is

$$K(t) = e^{tQ}.$$

Theorem 4.1 (Finite Markov semigroup)**.** *If Q has nonnegative off-diagonal entries and column sums zero, then e^{tQ} has nonnegative entries and column sums one for every $t \geq 0$.*

Proof. Choose $\lambda \geq \max_i(-Q_{ii})$. Then $B = Q + \lambda I$ has nonnegative entries, so e^{tB} is entrywise nonnegative by its power series. Hence

$$e^{tQ} = e^{-\lambda t} e^{tB}$$

is nonnegative. Since $\mathbf{1}^T Q = 0$, differentiation gives

$$\frac{d}{dt}\mathbf{1}^T e^{tQ} = \mathbf{1}^T Q e^{tQ} = 0,$$

so $\mathbf{1}^T e^{tQ} = \mathbf{1}^T$. □




Figure 4: The point mass δ_0 propagated by repeated symmetric random-walk transfer.

Example 4.2 (Symmetric random walk)**.** Starting from δ_0 and applying the transfer

$$T = \frac{1}{2}(S + S^{-1})$$

for n steps gives

$$\mathbb{P}(X_n = k) = 2^{-n}\binom{n}{(n+k)/2}$$

when $k \equiv n \pmod 2$, and zero otherwise. Hence

$$\mathbb{E}[X_n] = 0, \qquad \mathrm{Var}(X_n) = n.$$

The parity packet is exact: all mass remains in the one residue class permitted by n.

5 Markov lumping as exact refinement

Let K be a column-stochastic fine transition matrix, let C aggregate fine states into coarse packets, and let $\bar{K}$ be a coarse transition matrix. Define

$$\Delta = CK - \bar{K}C.$$

Theorem 5.1 (Exact lumpability)**.** *If $\Delta = 0$, then*

$$\boxed{CK^n = \bar{K}^n C}$$

for every $n \geq 0$. Thus fine evolution followed by coarse reconstruction equals coarse evolution from the beginning.

Proof. The result is immediate for $n = 0$. If $CK^n = \bar{K}^n C$, then

$$CK^{n+1} = \bar{K}^n CK = \bar{K}^{n+1} C.$$

□

Theorem 5.2 (Finite-horizon defect formula)**.** *For arbitrary $\Delta = CK - \bar{K}C$,*

$$\boxed{CK^n - \bar{K}^n C = \sum_{j=0}^{n-1} \bar{K}^{n-1-j} \Delta K^j.}$$

If K and $\bar{K}$ are contractions in the chosen norm, then

$$\left\| CK^n - \bar{K}^n C \right\| \leq n \left\| \Delta \right\|.$$

Proof. The identity follows by telescoping:

$$\bar{K}^{n-1-j} CK^{j+1} - \bar{K}^{n-j} CK^j = \bar{K}^{n-1-j} (CK - \bar{K}C) K^j.$$

Summing from $j = 0$ to $n - 1$ leaves $CK^n - \bar{K}^n C$. The norm bound follows termwise. □

6 Conditional expectation as a stochastic projector

Let $\Omega = \{1, \ldots, n\}$ carry strictly positive probabilities π_i, and let $\mathcal{Q} = \{B_1, \ldots, B_m\}$ be a partition. Define

$$(P_{\mathcal{Q}} f)_i = \frac{\sum_{j \in B(i)} \pi_j f_j}{\sum_{j \in B(i)} \pi_j},$$

where $B(i)$ is the block containing i.

Theorem 6.1 (Finite conditional-expectation projector)**.** *With the weighted inner product*

$$\langle f, g \rangle_\pi = \sum_i \pi_i f_i \overline{g_i},$$

the operator $P_{\mathcal{Q}}$ satisfies

$$P_{\mathcal{Q}}^2 = P_{\mathcal{Q}}, \qquad \langle P_{\mathcal{Q}} f, g \rangle_\pi = \langle f, P_{\mathcal{Q}} g \rangle_\pi.$$

Its range is the space of functions constant on each partition block.




Figure 5: Exact lumping agrees at every horizon. A small generator-level or one-step defect creates a controlled coarse discrepancy.

Proof. After one application, the result is constant on every block, so applying the same block average again changes nothing. Weighted self-adjointness follows by expanding both inner products block by block; both equal

$$\sum_{B\in\mathcal{Q}} \frac{1}{\pi(B)} \left(\sum_{i\in B} \pi_i f_i\right) \overline{\left(\sum_{j\in B} \pi_j g_j\right)}.$$

□

Theorem 6.2 (Tower law and martingale packets)**.** *If $\mathcal{Q}_c$ is coarser than $\mathcal{Q}_f$, then*

$$\boxed{P_c P_f = P_f P_c = P_c.}$$

For a nested tower $P_0 \le P_1 \le \cdots \le P_r$, define

$$D_0 = P_0, \qquad D_j = P_j - P_{j-1}.$$

Then

$$D_i D_j = \delta_{ij} D_j, \qquad P_r = \sum_{j=0}^{r} D_j.$$

Proof. A coarse-block average of a fine-block average is the coarse-block average, and a fine-block average of a coarse-constant function leaves it unchanged. The difference identities follow from $P_i P_j = P_{\min(i,j)}$. □

Classical Input 6.3 (Gaussian chaos realization). For the standard Gaussian measure, the normalized probabilists' Hermite polynomials form orthonormal chaos packets. The Ornstein–Uhlenbeck semigroup acts diagonally:

$$T_t H_n = e^{-nt} H_n.$$




Figure 6: Hermite–Gaussian packets, which diagonalize the Ornstein–Uhlenbeck semigroup. The finite numerical release verifies their Gaussian orthogonality through quadrature.

Thus conditional expectation, martingale differences, and Gaussian chaos are all projector realizations of stochastic refinement.

7 Bayesian conditioning as a normalized nonlinear operator

Let $p = (p_i)$ be a probability vector and let $\lambda = (\lambda_i)$ be a nonnegative likelihood with positive evidence

$$Z_\lambda(p) = \sum_i \lambda_i p_i.$$

Define

$$\boxed{\mathcal{B}_\lambda(p) = \frac{\lambda \odot p}{Z_\lambda(p)}.}$$

This map is nonlinear solely because of its scalar reconstruction factor.

Theorem 7.1 (Independent-likelihood composition)**.** *For likelihoods λ and μ with positive evidences,*

$$\boxed{\mathcal{B}_\mu(\mathcal{B}_\lambda(p)) = \mathcal{B}_{\lambda\odot\mu}(p).}$$

Proof. The first normalization appears in both numerator and denominator of the second update and cancels:

$$\frac{\mu_i(\lambda_i p_i/Z_\lambda)}{\sum_j \mu_j\lambda_j p_j/Z_\lambda} = \frac{\mu_i\lambda_i p_i}{\sum_j \mu_j\lambda_j p_j}.$$

□

7.1 Packet evidence and reconstruction

Let the state indices be partitioned into packets I_α. Define

$$Z_\alpha = \sum_{i \in I_\alpha} \lambda_i p_i, \qquad m_\alpha = \frac{Z_\alpha}{\sum_\beta Z_\beta}, \qquad q_{\alpha,i} = \frac{\lambda_i p_i}{Z_\alpha} \quad (i \in I_\alpha).$$

Theorem 7.2 (Bayesian packet reconstruction)**.**

$$\boxed{\mathcal{B}_\lambda(p) = \sum_\alpha m_\alpha q_\alpha.}$$

The nonlinear coupling between packets occurs only through the total evidence used to normalize the packet evidences.

Proof. For $i \in I_\alpha$,

$$m_\alpha q_{\alpha,i} = \frac{Z_\alpha}{Z} \frac{\lambda_i p_i}{Z_\alpha} = \frac{\lambda_i p_i}{Z}.$$

The packet supports are disjoint, so summing reconstructs the full posterior. □




Figure 7: A finite prior and posterior split into two state packets. Evidence changes both the local packet shape and the global packet mass.

Theorem 7.3 (Likelihood perturbation bound)**.** *Let p be fixed and let $\lambda, \mu \geq 0$ have positive evidences Z_λ, Z_μ. Then*

$$\boxed{\|\mathcal{B}_\lambda(p) - \mathcal{B}_\mu(p)\|_1 \leq \frac{2\sum_i |\lambda_i - \mu_i| p_i}{\min(Z_\lambda, Z_\mu)}.}$$

Proof. Add and subtract $\mu \odot p / Z_\lambda$ and use the triangle inequality:

$$\left\| \frac{\lambda \odot p}{Z_\lambda} - \frac{\mu \odot p}{Z_\mu} \right\|_1 \leq \frac{\|(\lambda - \mu) \odot p\|_1}{Z_\lambda} + \|\mu \odot p\|_1 \left| \frac{1}{Z_\lambda} - \frac{1}{Z_\mu} \right|.$$

The second term equals $|Z_\lambda - Z_\mu|/Z_\lambda$ and is bounded by the same weighted L^1 difference divided by Z_λ. Interchanging λ and μ gives the minimum-evidence form. □

8 Exact Bayesian refinement and information-loss packets

Let the fine state set I be partitioned into nonempty coarse packets I_a, $a \in A$. The coarsening map on probability vectors is

$$(C_\pi p)_a = P_a = \sum_{i \in I_a} p_i.$$

For a fine likelihood $\lambda = (\lambda_i)_{i \in I}$ and a prior p, define the effective coarse likelihood

$$r_a(p) = \frac{\sum_{i \in I_a} \lambda_i p_i}{P_a} \qquad (P_a > 0).$$

Its value on a zero-mass packet is irrelevant and may be set to zero.

Theorem 8.1 (Exact effective-likelihood factorization)**.** *Whenever the fine evidence is positive,*

$$\boxed{C_\pi \mathcal{B}_\lambda(p) = \mathcal{B}_{r(p)}(C_\pi p).}$$

Thus the only fine information needed for the coarse posterior, beyond the coarse prior masses, is the vector of conditional packet evidences $r(p)$.

Proof. For $P = C_\pi p$ and $Z = \sum_i \lambda_i p_i = \sum_a P_a r_a(p)$,

$$\big(C_\pi \mathcal{B}_\lambda(p)\big)_a = \frac{\sum_{i \in I_a} \lambda_i p_i}{Z} = \frac{P_a r_a(p)}{\sum_b P_b r_b(p)} = \big(\mathcal{B}_{r(p)}(P)\big)_a.$$

□

The next result gives the exact necessary and sufficient condition for a coarse likelihood to work for every prior.

Theorem 8.2 (Bayesian refinement criterion)**.** *Assume that the partition contains at least two coarse packets and that the fine and coarse likelihoods are strictly positive. The identity*

$$\boxed{C_\pi \mathcal{B}_\lambda(p) = \mathcal{B}_{\bar\lambda}(C_\pi p)}$$

holds for every fine prior p if and only if there exists one scalar $c > 0$ such that

$$\boxed{\lambda_i = c\,\bar\lambda_a \qquad (i \in I_a).}$$

Equivalently, the likelihood is constant on each coarse packet, with the coarse values determined up to the globally invisible Bayesian scaling c.

Proof. If $\lambda_i = c\bar\lambda_a$ on I_a, then $r_a(p) = c\bar\lambda_a$ for every prior, and the preceding theorem gives the desired identity because Bayesian normalization removes the common scalar c.

Conversely, fix two fine states $i, i' \in I_a$ and one state $j \in I_b$ in another coarse packet. Vary the distribution of a fixed coarse mass inside I_a between i and i' while leaving every coarse packet mass unchanged. The right side depends only on those coarse masses and is therefore unchanged. The left-side posterior mass of packet a depends on the conditional average of λ_i and $\lambda_{i'}$. It can be

independent of that variation only if $\lambda_i = \lambda_{i'}$. Hence λ is constant on every packet; write the value on I_a as λ_a.

Now use priors supported on one state of packet a and one state of packet b. Equality of the two coarse posterior odds gives

$$\frac{\lambda_a}{\lambda_b} = \frac{\bar{\lambda}_a}{\bar{\lambda}_b}.$$

This holds for every pair of coarse packets, so $\lambda_a = c\bar{\lambda}_a$ for one common $c > 0$. □

Remark 8.3 (The one-packet boundary). If the coarse partition has only one packet, every posterior coarsens to the scalar 1, so commutation is automatic and no within-packet likelihood information can be identified. The nontrivial criterion begins with at least two coarse packets.

When sufficiency fails, the defect still has an exact finite factorization. Fix a declared coarse likelihood $\bar{\lambda}$, put

$$P = C_\pi p, \qquad \delta(p) = r(p) - \bar{\lambda}, \qquad \bar{b} = \mathcal{B}_{\bar{\lambda}}(P), \qquad Z_r = \langle r(p), P\rangle.$$

Theorem 8.4 (Bayesian information-loss packet)**.** *The coarsening defect is*

$$\boxed{C_\pi \mathcal{B}_\lambda(p) - \mathcal{B}_{\bar{\lambda}}(P) = \frac{1}{Z_r}\left(\operatorname{diag}(P) - \bar{b}P^{\mathsf{T}}\right)\delta(p).}$$

The factor matrix annihilates $\bar{\lambda}$. This records the common-scale null direction at the declared coarse likelihood; the full exact defect remains homogeneous under simultaneous likelihood rescaling because the numerator and evidence denominator scale together. Moreover,

$$\boxed{\left\|C_\pi \mathcal{B}_\lambda(p) - \mathcal{B}_{\bar{\lambda}}(P)\right\|_1 \le \frac{2\sum_a P_a|\delta_a(p)|}{\min\{Z_r, Z_{\bar{\lambda}}\}}}$$

and

$$\sum_a P_a|\delta_a(p)| \le \sum_i p_i \left|\lambda_i - \bar{\lambda}_{\pi(i)}\right|.$$

Proof. The effective-likelihood theorem reduces the defect to

$$\mathcal{B}_{\bar{\lambda}+\delta}(P) - \mathcal{B}_{\bar{\lambda}}(P).$$

Since $Z_r = Z_{\bar{\lambda}} + P^{\mathsf{T}}\delta$, direct subtraction gives

$$\frac{1}{Z_r}\left(\operatorname{diag}(P) - \bar{b}P^{\mathsf{T}}\right)\delta.$$

Also,

$$\left(\operatorname{diag}(P) - \bar{b}P^{\mathsf{T}}\right)\bar{\lambda} = P \odot \bar{\lambda} - \bar{b}Z_{\bar{\lambda}} = 0.$$

The first norm bound is the likelihood perturbation theorem applied on the coarse simplex. Finally,

$$P_a|\delta_a| = P_a\left|\sum_{i\in I_a}\frac{p_i}{P_a}(\lambda_i - \bar{\lambda}_a)\right| \le \sum_{i\in I_a} p_i|\lambda_i - \bar{\lambda}_a|,$$

and summing over a proves the second inequality. □

Corollary 8.5 (Nested Bayesian packet tower)**.** *For a nested finite packet tower, Bayesian conditioning commutes with every coarsening map for every prior if and only if, at each level, the fine likelihood is blockwise proportional to the declared coarse likelihood on every coarsening fiber. Under this condition, all composite coarsening squares commute.*

Proof. Apply the refinement criterion at each adjacent pair of levels. Composition of the commuting squares gives the full tower statement. □

Corollary 8.6 (Canonical coarsest Bayesian-sufficient partition)**.** *For a finite family of likelihoods* $\{\lambda^{(y)}\}_{y\in Y}$*, define*

$$i \sim j \quad \Longleftrightarrow \quad \lambda_i^{(y)} = \lambda_j^{(y)} \text{ for every } y \in Y.$$

The equivalence classes form the unique coarsest partition on which every Bayesian update in the family closes exactly for every prior.

Proof. Every likelihood is constant on each equivalence class, so the refinement theorem gives exact closure. Conversely, any partition sufficient for every likelihood must place only states with equal likelihood values in the same packet. It therefore refines the displayed equivalence-class partition. □




Figure 8: The Bayesian refinement square commutes exactly when the likelihood is constant on each coarsening fiber, up to one global scale.

Numerical Certificate 8.7 (Bayesian refinement). For the finite test partition $\{0,1\} \sqcup \{2,3\}$, a blockwise-constant likelihood gave commutation residual 1.11×10^{-16}. A nonconstant likelihood produced an exact defect-factorization residual 7.29×10^{-17}. Its measured L^1 defect was 4.8787×10^{-2}, below the certified bound 1.1016×10^{-1}. Two priors with identical coarse masses but different hidden within-packet shapes produced coarse posteriors differing by 1.2400×10^{-1}, which is a direct necessity witness.

Boundary 8.8 (Finite exact theorem). The conjectured Bayesian refinement law is therefore resolved for finite packet systems and arbitrary priors. Infinite-dimensional disintegration, posterior consistency, and stability of data-dependent partitions require their own measure-theoretic and asymptotic hypotheses.




Figure 9: When packet sufficiency fails, the exact defect factors through the difference between the conditional effective likelihood and the declared coarse likelihood.

9 Bayesian flows, conjugacy, and hidden states

Fix a score vector g. The likelihood flow is

$$\lambda_i(t) = e^{tg_i},$$

and the Bayesian flow is

$$\Phi_t(p)_i = \frac{e^{tg_i} p_i}{\sum_j e^{tg_j} p_j}.$$

Theorem 9.1 (Bayesian nonlinear semigroup)**.**

$$\boxed{\Phi_{s+t} = \Phi_s \circ \Phi_t.}$$

Its generator is the centered multiplicative field

$$\boxed{\frac{d}{dt} p_i(t) = (g_i - \langle g, p(t) \rangle) p_i(t).}$$

Proof. The semigroup law is the likelihood-composition theorem with $e^{sg} e^{tg} = e^{(s+t)g}$. Differentiating the normalized exponential formula gives

$$\dot{p}_i = g_i p_i - p_i \frac{\sum_j g_j e^{tg_j} p_j(0)}{\sum_j e^{tg_j} p_j(0)}.$$

The quotient is $\langle g, p(t) \rangle$. □

Example 9.2 (Beta–Binomial conjugate packet)**.** A Beta(a, b) prior and s successes in n trials produce

$$\text{Beta}(a + s, b + n - s).$$

The prior exponents are additive packets: the likelihood increments them by $(s, n - s)$. In the numerical example,

$$(a, b) = (2, 3), \quad (s, n) = (7, 10),$$

so the posterior is Beta$(9, 6)$ with mean 0.6.




Figure 10: Continuous likelihood accumulation produces a nonlinear semigroup on the probability simplex.




Figure 11: A conjugate Bayesian update viewed as exact transfer of exponent packets.

9.1 Hidden Markov filtering

Let K be column stochastic and let $\lambda^{(r)}$ be the likelihood associated with the rth observation. The filter recursion is

$$\widehat{p}_r = Kp_{r-1}, \qquad p_r = \mathcal{B}_{\lambda^{(r)}}(\widehat{p}_r).$$

Proposition 9.3 (Prediction-update reconstruction)**.** *Every hidden Markov filtering step is the composition of a linear Markov transfer and a normalized nonlinear likelihood transfer. If both preserve nonnegative mass and the evidence is positive, then every p_r is a probability vector.*

Proof. Column stochasticity gives $\widehat{p}_r \geq 0$ and $\mathbf{1}^T\widehat{p}_r = 1$. Bayesian normalization then gives $p_r \geq 0$ and $\mathbf{1}^T p_r = 1$. □




Figure 12: A hidden Markov filter alternates a linear prediction semigroup with nonlinear Bayesian normalization.

10 Fractional powers as spectral packet transfer

Let $L = L^* \geq 0$ on a finite-dimensional Hilbert space. Its spectral resolution is

$$L = \sum_j \lambda_j P_j.$$

For $\alpha > 0$, define

$$\boxed{L^\alpha = \sum_j \lambda_j^\alpha P_j.}$$

The associated fractional semigroup is

$$T_\alpha(t) = e^{-tL^\alpha} = \sum_j e^{-t\lambda_j^\alpha} P_j.$$

Theorem 10.1 (Fractional packet calculus)**.** *For $\alpha, \beta > 0$,*

$$L^{\alpha} L^{\beta} = L^{\alpha+\beta},$$

and for $s, t \geq 0$,

$$\boxed{T_{\alpha}(s) T_{\alpha}(t) = T_{\alpha}(s+t).}$$

Every spectral packet P_j reduces L^{α} and $T_{\alpha}(t)$.

Proof. All statements hold coefficientwise on the mutually orthogonal spectral projectors. For example,

$$L^{\alpha} L^{\beta} = \sum_{j,k} \lambda_j^{\alpha} \lambda_k^{\beta} P_j P_k = \sum_{j} \lambda_j^{\alpha+\beta} P_j.$$

□




Figure 13: Different spectral exponents produce different transfer profiles from the same point seed.

10.1 Square-root subordination

Theorem 10.2 (Poisson subordination)**.** *For $t > 0$ and $L \geq 0$,*

$$\boxed{e^{-t\sqrt{L}} = \frac{t}{2\sqrt{\pi}} \int_0^{\infty} s^{-3/2} e^{-t^2/(4s)} e^{-sL}\, ds.}$$

The integral converges in operator norm in finite dimension.

Proof. By the spectral theorem it suffices to prove the scalar identity

$$e^{-t\sqrt{\lambda}} = \frac{t}{2\sqrt{\pi}} \int_0^{\infty} s^{-3/2} e^{-t^2/(4s) - \lambda s}\, ds$$

for $\lambda \geq 0$. This is the classical Gaussian integral after the substitution $u = t/(2\sqrt{s})$. Applying the identity to every eigenvalue and recombining the spectral projectors proves the operator formula. □

10.2 Fractional naturality under a bounded refinement defect

Let $A \geq 0$ act on a fine Hilbert space, let $B \geq 0$ act on a coarse Hilbert space, and let C be a bounded coarsening map. In the finite-dimensional statement below, set

$$D = CA - BC, \qquad D_\alpha = CA^\alpha - B^\alpha C, \qquad 0 < \alpha < 1.$$

Theorem 10.3 (Fractional intertwining theorem)**.** *For bounded nonnegative self-adjoint A and B,*

$$\boxed{D_\alpha = \frac{\sin(\pi\alpha)}{\pi} \int_0^\infty t^\alpha (tI + B)^{-1} D (tI + A)^{-1}\, dt.}$$

The improper integral converges in operator norm. Moreover,

$$\boxed{\|D_\alpha\| \leq K_\alpha \|C\|^{1-\alpha} \|D\|^\alpha, \qquad K_\alpha = \frac{2^{1-\alpha} \sin(\pi\alpha)}{\pi\alpha(1-\alpha)}.}$$

If $A, B \geq mI$ for some $m > 0$, then the stronger linear estimate holds:

$$\boxed{\|D_\alpha\| \leq \alpha m^{\alpha-1} \|D\|.}$$

In particular, exact generator intertwining $CA = BC$ implies exact fractional intertwining $CA^\alpha = B^\alpha C$.

Proof. For $0 < \alpha < 1$, the Balakrishnan formula gives

$$A^\alpha = \frac{\sin(\pi\alpha)}{\pi} \int_0^\infty t^{\alpha-1} A (tI + A)^{-1}\, dt,$$

and likewise for B. Using

$$A(tI + A)^{-1} = I - t(tI + A)^{-1}$$

and the rectangular resolvent identity

$$(tI + B)^{-1} C - C(tI + A)^{-1} = (tI + B)^{-1}(CA - BC)(tI + A)^{-1},$$

the difference of the two Balakrishnan integrals becomes the displayed resolvent-defect formula.

For small t,

$$\left\| CA(tI + A)^{-1} - B(tI + B)^{-1} C \right\| \leq 2 \|C\|,$$

because the two positive resolvent factors are contractions. For large t, the resolvent identity gives the bound $\|D\|/t$. Splitting the integral at

$$r = \frac{\|D\|}{2\|C\|}$$

when C and D are nonzero yields

$$\begin{aligned} \|D_\alpha\| &\leq \frac{\sin(\pi\alpha)}{\pi} \left(2\|C\| \int_0^r t^{\alpha-1}\, dt + \|D\| \int_r^\infty t^{\alpha-2}\, dt \right) \\ &= K_\alpha \|C\|^{1-\alpha} \|D\|^\alpha. \end{aligned}$$

The zero cases are immediate.

If $A, B \geq mI$, then both resolvents have norm at most $(t + m)^{-1}$, so

$$\|D_\alpha\| \leq \frac{\sin(\pi\alpha)}{\pi} \|D\| \int_0^\infty \frac{t^\alpha}{(t + m)^2}\, dt.$$

After $t = mu$, the beta integral equals

$$m^{\alpha-1} \frac{\alpha\pi}{\sin(\pi\alpha)},$$

which gives the linear estimate. □

Corollary 10.4 (Fractional semigroup refinement)**.** *For $s \geq 0$,*

$$\boxed{Ce^{-sA^\alpha} - e^{-sB^\alpha}C = -\int_0^s e^{-(s-r)B^\alpha} D_\alpha e^{-rA^\alpha}\, dr.}$$

Hence

$$\left\| Ce^{-sA^\alpha} - e^{-sB^\alpha}C \right\| \leq s \left\| D_\alpha \right\|.$$

Under the common gap $A, B \geq mI$,

$$\boxed{\left\| Ce^{-sA^\alpha} - e^{-sB^\alpha}C \right\| \leq se^{-sm^\alpha} \alpha m^{\alpha-1} \left\| D \right\|.}$$

Proof. Differentiate

$$r \longmapsto e^{-(s-r)B^\alpha} Ce^{-rA^\alpha}$$

and integrate from 0 to s. Positive fractional semigroups are contractions; under the gap, the product of their two norm bounds is e^{-sm^α}. □

Corollary 10.5 (Fractional refinement towers)**.** *Let $D_\ell = C_\ell A_{\ell+1} - A_\ell C_\ell$ in a bounded nonnegative self-adjoint tower, with $\sup_\ell \|C_\ell\| \leq M$. If*

$$\sum_\ell \|D_\ell\|^\alpha < \infty,$$

then the fractional defects

$$C_\ell A_{\ell+1}^\alpha - A_\ell^\alpha C_\ell$$

are summable. If the tower has a uniform gap $m > 0$, it is enough that $\sum_\ell \|D_\ell\| < \infty$. On every bounded time interval, the corresponding fractional-semigroup defects are summable as well.




Figure 14: A generator-level refinement defect transfers quantitatively to fractional powers and then to fractional semigroups. The two upper curves are theorem certificates, not fitted error bars.

Boundary 10.6 (Unbounded and nonnormal generators). The theorem resolves fractional naturality for bounded nonnegative self-adjoint refinement systems and gives the exact estimate needed for finite discretization towers. For unbounded positive generators, the same resolvent argument requires a declared common-domain condition and a bounded extension of $CA - BC$. Sectorial or nonnormal generators require their own contour calculus and resolvent bounds. Those domain questions are not inferred from the finite certificates.

Boundary 10.7 (Non-self-adjoint fractional powers). The spectral definition above is complete for finite self-adjoint positive operators. Sectorial and nonnormal operators require a resolvent or contour functional calculus; the same packet language can be used only after those analytic hypotheses are established.

11 Green operators and the Poisson seed

On the interval $(0,1)$ with Dirichlet boundary conditions, the Green kernel for $-u'' = f$ is

$$\boxed{G(x,y) = \min(x,y) - xy.}$$

Thus the point source δ_y generates the solution kernel, and a general source is reconstructed by

$$u(x) = \int_0^1 G(x,y) f(y)\, dy.$$

For n interior points with spacing $h = 1/(n+1)$, define

$$L_h = \frac{1}{h^2} \begin{pmatrix} 2 & -1 & & \\ -1 & 2 & \ddots & \\ & \ddots & \ddots & -1 \\ & & -1 & 2 \end{pmatrix}.$$

Theorem 11.1 (Exact discrete Green reconstruction at grid nodes). *Let $x_i = ih$. Then*

$$\boxed{\frac{1}{h}(L_h^{-1})_{ij} = G(x_i, x_j).}$$

Thus the scaled discrete inverse reproduces the continuum Dirichlet Green kernel exactly at the grid nodes.

Proof. The inverse of the unscaled tridiagonal matrix has entries

$$\frac{\min(i,j)(n+1-\max(i,j))}{n+1}.$$

Multiplying by h^2 and then dividing by h gives

$$h \min(i,j) \left(1 - \frac{\max(i,j)}{n+1}\right) = \min(x_i, x_j) - x_i x_j.$$

□

11.1 Low-mode convergence

The kth discrete eigenvalue is

$$\lambda_{k,h} = \frac{4}{h^2} \sin^2\left(\frac{k\pi h}{2}\right),$$

whereas the continuum eigenvalue is $(k\pi)^2$.




Figure 15: A section of the continuum Green kernel and the scaled discrete inverse. They agree at the grid nodes.

Theorem 11.2 (Second-order low-mode convergence)**.** *For fixed k,*

$$\lambda_{k,h} = (k\pi)^2 + O(h^2).$$

More precisely, for sufficiently small h,

$$0 \le (k\pi)^2 - \lambda_{k,h} \le \frac{(k\pi)^4}{12} h^2.$$

Proof. For $x \ge 0$, $0 \le x^2 - \sin^2 x \le x^4/3$. Set $x = k\pi h/2$ and multiply by $4/h^2$. □

12 Finite-state Feynman–Kac transfer

Let K be a column-stochastic matrix, let V be a potential on the states, and let f be terminal data. Define

$$M_V = \mathrm{diag}(e^{-V})K^T.$$

The backward Feynman–Kac value after n steps is

$$u_n = M_V^n f.$$

Theorem 12.1 (Discrete Feynman–Kac path formula)**.** *For an initial state i,*

$$\boxed{(u_n)_i = \mathbb{E}_i\left[\exp\left(-\sum_{r=0}^{n-1} V(X_r)\right) f(X_n)\right].}$$




Figure 16: The first finite-difference eigenvalue converges with observed order approximately two.

Proof. For $n = 1$,

$$(M_V f)_i = e^{-V(i)} \sum_j K_{ji} f_j,$$

which is the stated expectation. Iterating and expanding the matrix product sums over every path $(i = i_0, i_1, \dots, i_n)$ with probability $\prod_r K_{i_{r+1},i_r}$ and weight $\prod_r e^{-V(i_r)}$. □

13 Exact elimination of latent packets

Let $X = PX \oplus QX$ and write

$$A = \begin{pmatrix} A_{PP} & A_{PQ} \\ A_{QP} & A_{QQ} \end{pmatrix}.$$

Theorem 13.1 (Schur–Feshbach projected resolvent)**.** *Whenever $zI - A_{QQ}$ is invertible, define*

$$\boxed{F_P(z) = zI - A_{PP} - A_{PQ}(zI - A_{QQ})^{-1} A_{QP}.}$$

Then

$$\boxed{P(zI - A)^{-1} P = F_P(z)^{-1}.}$$

Proof. Solve the hidden block equation

$$(zI - A_{QQ})x_Q = A_{QP} x_P + y_Q$$

for x_Q and substitute it into the retained equation. The resulting retained coefficient is $F_P(z)$. Equivalently, this is the block inverse formula. □

The self-energy

$$\Sigma_P(z) = A_{PQ}(zI - A_{QQ})^{-1} A_{QP}$$

contains the exact effect of the eliminated packets. A bare truncation A_{PP} discards this term and is generally not exact.




Figure 17: Potential-weighted transfer values computed by the finite Feynman–Kac operator. Exhaustive path enumeration agrees with the matrix formula in the supplied tests.

13.1 Time-domain memory

For the homogeneous system $\dot{x} = Ax$, variation of constants gives

$$x_Q(t) = e^{tA_{QQ}}x_Q(0) + \int_0^t e^{(t-r)A_{QQ}}A_{QP}x_P(r)\,dr.$$

Substitution yields

$$\dot{x}_P(t) = A_{PP}x_P(t) + A_{PQ}e^{tA_{QQ}}x_Q(0) + \int_0^t \mathcal{K}(t-r)x_P(r)\,dr,$$

where

$$\boxed{\mathcal{K}(t) = A_{PQ}e^{tA_{QQ}}A_{QP}.}$$

Theorem 13.2 (Memory/self-energy correspondence)**.** *For z to the right of the spectrum of A_{QQ},*

$$\boxed{\int_0^\infty e^{-zt}\mathcal{K}(t)\,dt = A_{PQ}(zI - A_{QQ})^{-1}A_{QP} = \Sigma_P(z).}$$

Proof. Use

$$\int_0^\infty e^{-zt}e^{tA_{QQ}}\,dt = (zI - A_{QQ})^{-1}$$

and multiply on the left and right by the two transfer blocks. □




Figure 18: The norm of the exact memory kernel produced by eliminated packets.

14 Survival and hazard semigroups

Let A be a transient-state subgenerator: its off-diagonal entries are nonnegative and its column sums are nonpositive. Let p_0 be an initial transient-state distribution. Define

$$p(t) = e^{tA}p_0, \qquad S(t) = \mathbf{1}^T p(t).$$

The exit-rate row vector is

$$a^T = -\mathbf{1}^T A \geq 0.$$

Theorem 14.1 (Semigroup survival identities)**.** *The event-time density and hazard are*

$$\boxed{f(t) = a^T e^{tA}p_0 = -S'(t), \qquad h(t) = \frac{f(t)}{S(t)}.}$$

Consequently, S is nonincreasing and

$$S(t) = \exp\left(-\int_0^t h(r)\,dr\right)$$

whenever $S(t) > 0$.

Proof. Differentiate:

$$S'(t) = \mathbf{1}^T A e^{tA}p_0 = -a^T e^{tA}p_0.$$

The hazard identity is the definition of conditional instantaneous exit rate. Integrating $(\log S)' = -h$ gives the final formula. □




Figure 19: Survival and instantaneous hazard produced by one transient transfer semigroup.

Realization	Seed	Transfer	Packet/reconstruction law
Markov	point mass δ_i	K^n or e^{tQ}	state lumping and $CK^n = \bar{K}^n C$
Conditional	random variable	$P_{\mathcal{Q}}$	nested partitions and tower law
Bayesian	prior p	$\mathcal{B}_\lambda$	packet evidences and normalized synthesis
Fractional	spectral source	e^{-tL^α}	spectral projectors P_j
PDE	source f or δ_y	resolvent/Green operator	eigenmode or spatial packet synthesis
Feynman–Kac	terminal data f	$(\mathrm{diag}(e^{-V})K^T)^n$	weighted path reconstruction
Latent system	retained state	Feshbach/self-energy	hidden packets return as memory
Survival	initial transient law	e^{tA}	sum over transient packets

Table 1: The transfer-semigroup dictionary in the common operator language.

15 One common operator language

The preceding constructions are not identical, but they use the same packet operations. The domain dictionary is summarized in table 1.

Theorem 15.1 (Naturality under exact intertwining)**.** *Let F be any analytic function on a neighborhood of the spectra of A_f and A_c. If*

$$CA_f = A_c C,$$

then

$$\boxed{CF(A_f) = F(A_c)C.}$$

In particular, exact generator refinement propagates simultaneously to semigroups, resolvents, integer powers, and analytic transfer functions.

Proof. The identity $CA_f^m = A_c^m C$ follows by induction. Apply it termwise to the convergent power series of F. Resolvents follow either from the resolvent identity or from multiplication by the two invertible factors. □

Remark 15.2 (Where nonlinearity enters)*.* Bayesian conditioning does not fit the preceding linear functional calculus because normalization depends on the current state. It nevertheless obeys an exact composition law and has a closed packet reconstruction. Paper III supplies the general nonlinear coordinate theorem; the Bayesian map is a particularly transparent probabilistic realization.

16 Numerical certificates

All calculations in this release were run on CPU. No GPU timing or performance claim is made. Floating-point residuals are implementation checks and are not substitutes for the symbolic proofs.

Numerical Certificate 16.1 (Test suite)*.* The complete release passes

$$\boxed{37/37 \text{ automated tests.}}$$

The tests cover projector reconstruction, packet-block semigroup composition, exact and approximate Markov lumping, Bayesian composition and flow, the exact Bayesian refinement criterion, information-loss factorization, canonical sufficient partitions, conditional-expectation towers, Hermite orthogonality, fractional semigroups, square-root subordination, fractional intertwining and semigroup-defect bounds, discrete Green kernels, PDE convergence, Feynman–Kac enumeration, Schur–Feshbach elimination, memory kernels, refinement defects, hidden Markov filtering, and survival curves.

17 Remaining conjectural extensions

The finite theory closes several operator identities but does not settle the corresponding universal continuum questions.

Conjecture 17.1 (Transfer-semigroup realization)**.** Every sufficiently regular realization of the seed calculus whose transfer stage is local in time admits a closed generator on a completed state space, and the packet analysis maps form a compatible system for its semigroup or evolution family.

Conjecture 17.2 (Uniform refinement control)**.** For a nested family of physically or analytically meaningful discretizations, there exist norms and constants independent of resolution such that

$$\left\| C_\ell e^{tA_{\ell+1}} - e^{tA_\ell} C_\ell \right\| \le M\, t\, \|C_\ell A_{\ell+1} - A_\ell C_\ell\|$$

Certificate	Value
Packet-block semigroup composition residual	2.554×10^{-15}
Operator reconstruction residual	0
Random-walk mass error	0
Exact Markov generator-lumping defect	5.551×10^{-17}
Exact finite-horizon lumping error	2.220×10^{-16}
Perturbed ten-step lumping error	6.523×10^{-3}
Linear defect bound for that example	2.000×10^{-1}
Bayesian composition residual	5.551×10^{-17}
Bayesian packet reconstruction residual	0
Bayesian refinement commutation residual	1.110×10^{-16}
Bayesian defect-factorization residual	7.286×10^{-17}
Bayesian flow generator residual	1.959×10^{-11}
Conditional-expectation tower residual	0
Hermite Gram residual	9.992×10^{-15}
Fractional semigroup residual	2.082×10^{-17}
Maximum square-root subordination error	1.665×10^{-16}
Exact fractional-naturality residual	0
Fractional-power defect / gap certificate	0.51059/0.95079
Fractional-semigroup defect / certificate	0.12969/0.44480
Observed low-mode PDE convergence order	1.999854
Feynman–Kac enumeration residual	1.665×10^{-16}
Feshbach projected-resolvent residual	5.551×10^{-17}
Memory Laplace-transform residual	6.939×10^{-18}
Duhamel refinement identity residual	2.108×10^{-16}
Survival monotonicity violation	0
Hidden-Markov normalization error	1.110×10^{-16}

Table 2: Selected results from the machine-readable numerical record.

on every bounded time interval. Summable generator defects would then produce a coherent inverse-limit evolution.

Conjecture 17.3 (Latent-packet closure)**.** For a stable infinite packet system, exact elimination of unresolved packets produces a causal memory kernel whose Laplace transform is the limiting packet self-energy. A Markovian closure exists precisely when this self-energy is constant, or can be represented by finitely many additional state packets.

Boundary 17.4 (What remains outside Paper VI)*.* The following issues are not proved here:

1. existence and uniqueness for arbitrary stochastic differential equations;
2. posterior consistency or asymptotic normality in infinite statistical models;
3. domain invariance for unbounded fractional operators;
4. tightness and compactness required for stochastic continuum limits;
5. nonlinear PDE well-posedness beyond the stated finite and linear models;
6. an optimal packet choice for a general Markov, Bayesian, or PDE problem.

These belong to later realization-specific work or to external analytic theory.

18 Computational implementation

The accompanying Python package implements the formulas and finite models used in the paper.

`projector_transfer_semigroups`

The computational organization mirrors the paper:

`packets.py`	projectors, blocks, reconstruction, block composition
`semigroup.py`	matrix semigroups, defects, Feshbach operators, memory
`markov.py`	random walks, exact lumping, survival
`bayes.py`	normalized updates, packet evidence, conjugacy, flows, HMM
`conditional.py`	conditional expectation and Gaussian chaos
`fractional.py`	spectral powers, subordination, and fractional refinement certificates
`pde.py`	Dirichlet Laplacians, Green kernels, eigenvalue error
`feynman_kac.py`	weighted transfer and exhaustive path enumeration.

18.1 Representative algorithms

Listing 1: Normalized finite Bayesian update.

```
def bayes_update(prior, likelihood):
    p = normalize(prior)
    evidence = likelihood @ p
    if evidence <= 0:
        raise ValueError("positive evidence required")
    return likelihood * p / evidence
```

Listing 2: Packet block composition.

```
def block_composition(left, right):
    result = []
    for beta in range(len(left)):
        row = []
        for alpha in range(len(left)):
            value = sum(
                left[beta][gamma] @ right[gamma][alpha]
                for gamma in range(len(left))
            )
            row.append(value)
        result.append(row)
    return result
```

Listing 3: Finite Schur-Feshbach effective operator.

```
def feshbach_effective_operator(A, retained, z):
    App, Apq, Aqp, Aqq = split_blocks(A, retained)
    return z*I - App - Apq @ inv(z*I - Aqq) @ Aqp
```

The full implementation, tests, tables, figure-generation code, and exact numerical record are included with the release rather than embedded in the typeset paper.

19 Conclusion

The transfer-semigroup realization gives one common operator language to several subjects that are often presented separately. Markov propagation is linear transfer of a probability seed. Conditional expectation is stochastic projection. Bayesian conditioning is normalized nonlinear transfer, and its exact finite coarsening law is now characterized by packetwise likelihood sufficiency. Fractional calculus is spectral packet reweighting, and bounded positive refinement defects now transfer to fractional powers and fractional semigroups by explicit resolvent and Duhamel estimates. A Green operator reconstructs a field from source packets. Feynman–Kac propagation adds path weights to stochastic transfer. Schur–Feshbach elimination shows that hidden packets are not destroyed; they return through self-energy and memory. Survival and hazard are scalar reconstructions of a transient semigroup.

The central structural identity is

$$\boxed{\text{generator defect} \longrightarrow \text{Duhamel semigroup defect} \longrightarrow \text{controlled refinement error.}}$$

Together with exact packet reconstruction, this makes the semigroup paper the natural bridge from the foundational calculus to the quantum, computational, and arithmetic realizations that follow.