

The Milky Way Joins the Extragalactic World: I. PHANGS

NEAL J. EVANS II,¹ DAVIDE ELIA,^{2,3} JIAYI SUN,⁴ SOPHIA STUBER,^{5,6} AND ADAM K. LEROY⁷

¹*Department of Astronomy, The University of Texas at Austin, 2515 Speedway, Stop C1400, Austin, Texas 78712-1205, USA*

²*INAF-IAPS, Via del Fosso del Cavaliere 100, I-00133 Roma, Italy*

³*INAF, Sezione di Lecce, Via per Arnesano, I-73100 Lecce, Italy*

⁴*Department of Physics and Astronomy, University of Kentucky, 506 Library Drive, Lexington, KY 40506, USA*

⁵*National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan*

⁶*Max Planck Institute for Radio Astronomy, Auf dem Hügel 69, 53121 Bonn, Germany*

⁷*Department of Astronomy, The Ohio State University, 140 West 18th Ave, Columbus, OH 43210, USA*

ABSTRACT

Complete catalogs of molecular clouds in the Milky Way allow analysis of the molecular medium and the star formation properties of the Milky Way that closely follows the method used for nearby galaxies, in particular in the PHANGS project. The dependencies of the depletion time on other properties of molecular gas in the Milky Way are similar to those of other galaxies when analyzed in an analogous way. They exhibit a large scatter and relatively weak correlations. The strongest correlation is a decrease in depletion time with increasing velocity dispersion. We explore the effects of spatial resolution, sensitivity, cloud identification method, averaging method, and tracer choice on our results. Metallicity effects in converting observations to mass can have a substantial impact. Somewhat fortuitously, differences in conversion methods between PHANGS and the Milky Way do not affect the current comparison. Inadequate spatial resolution, resulting in unresolved cloud structure, has the most important effect on interpretation of both extragalactic observations and existing catalogs of Milky Way clouds.

Keywords: ISM: molecules, ISM: clouds, star formation(1569)

1. INTRODUCTION

Our goal is to place the Milky Way in the context of recent comprehensive surveys of the molecular gas and star formation studies of large samples of nearby galaxies. Using similar analysis techniques provides insight into the limitations of the extragalactic data. This paper focuses on CO observations and the PHANGS-ALMA survey in particular. Another paper will focus on surveys of tracers of denser gas (Bigiel et al. 2016; Jiménez-Donaire et al. 2019; Neumann et al. 2025). A spin-off of the current paper on the morphology of the Milky Way in comparison to barred galaxies was also published (Evans et al. 2026).

The PHANGS-ALMA survey (Leroy et al. 2021) has provided data on the molecular gas and star formation rate (SFR) on ~ 100 pc scales for 90 galaxies. The sample targeted nearby ($d < 20$ Mpc), low to moderately

inclined ($i \lesssim 75^\circ$), relatively massive ($M_\star/M_\odot \gtrsim 10^{9.75}$) star forming ($\text{SFR}/M_\star > 10^{-11} \text{ yr}^{-1}$) galaxies.

The PHANGS-ALMA team have explored many aspects of the data (Rosolowsky et al. 2021; Sun et al. 2022; Zakardjian et al. 2023; Leroy et al. 2022; Brunetti et al. 2024). The PHANGS-JWST project is providing infrared images of the sample (Williams et al. 2024) and information on star formation, feedback, and dust physics (Lee et al. 2023). They have explored such issues as the duration of the embedded (not seen in visible light tracers) star formation (Ramambason et al. 2026). PHANGS-HST (Lee et al. 2022) has obtained images of some galaxies with very high angular resolution, which Faustino Vieira et al. (2025) have used to create cloud catalogs based on extinction. These studies provide a comprehensive view of molecular gas and star formation, but the distance-limited spatial resolution may affect the conclusions. Comparison to studies in the Milky Way can provide valuable checks on that, so the goal is to add the Milky Way to the PHANGS data set. The Milky Way satisfies all the criteria for

the PHANGS sample, except of course the inclination: $M_*/M_\odot = 10^{10.72}$ and $\text{SFR}/M_* = 3 \times 10^{-11} \text{ yr}^{-1}$ (Imig et al. 2025; Licquia & Newman 2015).

Studies of other galaxies often make comparisons to the Milky Way, but they are hampered by the incomplete picture of our Milky Way caused by our location. Vast differences in spatial resolution across the Milky Way complicate analysis, and cloud properties are often biased toward nearby clouds. We now have quite complete maps of CO $J = 1 \rightarrow 0$ in the Milky Way (Dame et al. 2001) and a cloud catalog that accounts for all the CO emission (Miville-Deschênes et al. 2017). Elia et al. (2022) derived a face-on view of star formation in the Milky Way, using a model of star formation rate as a function of dense clump mass. More recently, Elia et al. (2025a, hereafter E25) have shown that three methods of predicting the total star formation rate and its distribution over Galactocentric radius (R_{Gal}) provide consistent information.

Additional surveys of the Milky Way are underway in different tracers with improved resolution, such as MWISP in CO, ^{13}CO , and $\text{C}^{18}\text{O } J = 1 \rightarrow 0$ (Su et al. 2019; Sun et al. 2020; Zhou et al. 2025), SEDIGISM in $^{13}\text{CO } J = 2 \rightarrow 1$ (Duarte-Cabral et al. 2021; Duarte-Cabral & The Sedigism Consortium 2022; Neralwar et al. 2022) and OGHReS in CO, ^{13}CO , and $\text{C}^{18}\text{O } J = 2 \rightarrow 1$ (Urquhart et al. 2025, and references therein). The time is right to place the Milky Way in the extragalactic context, at least in terms of its star formation and molecular gas properties.

Our primary comparison is to the paper by Leroy et al. (2025, hereafter L25), who have published a comprehensive study of cloud-scale gas properties, depletion times, and star formation efficiencies in a sub-sample of 67 galaxies from PHANGS-ALMA. They further provided a detailed, step-by-step description of their methods for computing the various properties of the clouds and star formation processes. As much as is possible, we follow the same steps for the Milky Way. Some details differ because of the different nature of the datasets, but we specify these differences and consider the likely consequences. One difference is deliberate. Because our spatial resolution is much higher, we can examine the effects of spatial resolution on the outcomes, providing insights into the sub-grid assumptions used in the analysis of extragalactic data. To simulate the effects of the resolution of the PHANGS data, we assign each cloud in the Milky Way to a cell in a rectangular grid with a size equal to the PHANGS resolution.

Because many quantities are used, and the notation has to distinguish different kinds of averages, we provide for reference a table of symbols (Table 1). For

PHANGS, we generally follow their notation, but remind the reader that “cloud” means 150 pc cloud-scale. In contrast, our cloud scale is much smaller and varies across the Milky Way. We use the term “cell” for the 150 pc scale. Most of the quantities need no definition, but we clarify that $\bar{n}$ is the mean density of all colliding partners; in a molecular cloud, this is $n(\text{H}_2) + n(\text{He}) + \dots$, calculated from

$$\bar{n} = M/(V \mu m_{\text{amu}}), \quad (1)$$

where M is the mass, V is the volume, m_{amu} is the atomic mass unit. A standard assumption for the mean molecular weight, $\mu = 2.37$ (Kauffmann et al. 2008). Shirley et al. (2026) has recently improved this value to $\mu = 2.3514 \pm 0.0006$, for solar abundances, which we adopt. Variations with metallicity will be less than 1%.

2. DATASETS

2.1. Extragalactic Data

PHANGS-ALMA has produced a number of products that we use. Machine-readable tables recording high-level measurements from the various PHANGS datasets, described in Sun et al. (2022), were downloaded from CANFAR⁸. Most of our comparisons are to plots and fits in L25 using tables published online with the paper. However, we use the tables published in the Erratum (Leroy et al. 2026), which correct some issues with the original tables.

In addition, the extinction-based cloud catalogs of Faustino Vieira et al. (2025) provide a complementary view of structures, which is agnostic as to the composition, as both atomic and molecular gas is counted. In particular, we use their catalog for NGC4321 to gauge the effect of using different tracers. It was one of the molecular morphological analogs for the Milky Way identified by Evans et al. (2026). While NGC4548 was even more similar to the Milky Way, it was not included in Faustino Vieira et al. (2025). For NGC4321, the spatial resolution allows detection of much smaller (radii defined by $\sqrt{A/\pi}$ down to 8 pc, where A is the area) and less massive (minimum mass of $1.5 \times 10^3 M_\odot$) clouds than those identified from the ALMA data. From the directory, /AstroDataCitationDOI/CISTI.CANFAR/25.0036/data, we downloaded the catalog for NGC4321 (NGC4321_cloud.catalogue.ecsv) and the ho-

⁸ The data repository is <https://www.canfar.net/storage/list/AstroDataCitationDOI/CISTI.CANFAR/22.0072/data> and we used NGC4321_hexagon_1p5kpc.ecsv from the v4p0_public_release for example.

Table 1. Symbol Definitions

Quantity	PHANGS	MW	PHANGS	MW	PHANGS	MW	Note
	Clouds		Cells		Hexagons		
Size	...	0.1-70 pc	150 pc	150 pc	1.5 kpc	1.5 kpc	
Mass	...	M_{cloud}	M_{cl}	M_{cell}	M_{hex}	M_{hex}	
Free-fall Time	...	t_{ff}	$\tau_{\text{ff}}^{\text{cloud}}$	$t_{\text{ff,cell}}$	$\langle \tau_{\text{ff}}^{\text{cloud}} \rangle$	$\langle t_{\text{ff}} \rangle_{\text{hex}}$	1
Surface Density	...	Σ_{cloud}	$\Sigma_{\text{mol}}^{\text{cloud}}$	Σ_{cell}	$\langle \Sigma_{\text{mol}}^{\text{cloud}} \rangle$	$\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$	2
Surface Density	...	Σ_{cloud}	$\Sigma_{\text{mol}}^{\text{cloud}}$	$\langle \Sigma_{\text{cloud}} \rangle$	$\langle \Sigma_{\text{mol}}^{\text{cloud}} \rangle$	$\langle \Sigma_{\text{cloud}} \rangle_{\text{hex}}$	3
Velocity Dispersion (σ_v)	...	σ_v^{cloud}	$\sigma_{\text{mol}}^{\text{cloud}}$	σ_v^{cell}	$\langle \sigma_{\text{mol}}^{\text{cloud}} \rangle$	$\langle \sigma_v^{\text{cell}} \rangle_{\text{hex}}$	4
Virial Parameter	...	α_{vir}	$\alpha_{\text{vir}}^{\text{cloud}}$	$\alpha_{\text{vir,cell}}$	$\langle \alpha_{\text{vir}}^{\text{cloud}} \rangle$	$\langle \alpha_{\text{vir}} \rangle_{\text{hex}}$	5
Average Density ($\bar{n}$)	...	$\bar{n}$	...	$\bar{n}_{\text{cell}}$	...	$\langle \bar{n}_{\text{cell}} \rangle_{\text{hex}}$	6

NOTE—Notes apply to the column immediately to the left. 1. Averages are the inverses of mass-weighted averages over speeds, as defined by equation 1. 2. Mass-weighted mean of $\Sigma_{\text{cell}} = M_{\text{cell}}/A_{\text{cell}}$. 3. Mass-weighted mean of Σ_{cloud} for all clouds within the cell. 4. Mass-weighted mean of σ_v^{cloud} for all clouds within the cell. 5. Mass-weighted mean of $\alpha_{\text{vir,cell}}$ based on total size, mass, and velocity dispersion in a cell. 6. Mass-weighted mean of $\bar{n}_{\text{cell}}$ based on the mass and volume of a cell.

mogenized catalog of all four galaxies (homogenised_cloud_catalogue.ecsv).

2.2. Milky Way data

The data on molecular gas in the Milky Way, for comparison to PHANGS, comes from the catalog of clouds defined by CO $J = 1 \rightarrow 0$ emission (Miville-Deschênes et al. 2017, hereafter MD17). E25 adjusted this catalog to use the same rotation curve as was used by the Hi-GAL survey, adjusted masses for a value of α_{CO} that varied with metallicity, Z , and provided star formation rates predicted by the model of Evans et al. (2022, hereafter EKO), using a star formation efficiency per free-fall time (ϵ_{ff}) that varied with the virial parameter (α_{vir}), as found by Kim et al. (2021).

For a few comparisons, we also use the catalog of Rice et al. (2016, hereafter R16), which is less complete (includes about 25% of the total CO $J = 1 \rightarrow 0$ emission of the Milky Way) and discriminates against low mass clouds compared to the catalog of MD17. It is useful to study the effects of different cloud identification methods.

We use the catalogs of sources rather than going back to the input data. This avoids repeating past work. An exception is the measure of SFR. Because the SFRs in the catalog of Elia et al. (2025b) are based on the model of EKO, which assume a relation between SFR and cloud mass, free-fall time, and virial parameter, tests of the dependence of t_{dep} on cloud properties would be circular. However, E25 showed that the SFR based on an alternative approach, consisting in converting the *Herschel* emission at 70 μm into a luminosity at that wave-

length (by using the heliocentric distance of the nearest compact source), and hence into SFR according to the prescription of Li et al. (2010), gave a total SFR and dependence on R_{Gal} that is very similar to that of SFR based on EKO. Consequently, we use the SFR based on the whole 70 μm emission, after converting pixel intensities into luminosities through distances assigned to the closest clumps, as described in E25. The resulting surface densities of SFR and molecular gas satisfy a Kennicutt-Schmidt relation with a power-law exponent of 1.10 ± 0.06 , identical within uncertainty to that found for the PHANGS galaxies (Sun et al. 2023).

3. PROCEDURES

Here we outline the procedures used by L25 and how we depart from them. “PA” indicates the PHANGS-ALMA step and “CP” indicates the current paper. The issue of the luminosity to mass conversion factor (α_{CO}) and its dependence on metallicity are discussed in §4.1.

1. (PA) Choose a threshold of CO $J = 2 \rightarrow 1$ emission and apply a local, best-estimate mass conversion factor (α_{CO}) to estimate the mass on the scale of 150 pc. The threshold value of integrated intensity is 0.6 K km s $^{-1}$ to achieve a uniform sensitivity limit.

(CP) We have CO $J = 1 \rightarrow 0$ emission, rather than $J = 2 \rightarrow 1$ data. We have cloud catalogs rather than images. The threshold for the Milky Way data is lower. For our standard procedure, we apply no criterion on integrated intensity, but we adopt the requirements for clouds used by

E25 as follows: $R_{\text{Gal}} \leq 20$ kpc, $M_{\text{mol}} \geq 1 M_{\odot}$, $\alpha_{\text{vir}} < 100$. Restrictions on CO integrated intensity are then added to test the effects of sensitivity limitations (§4.3). The “best-estimate” value of α_{CO} will differ because of the different transition, but this difference turns out to be small (§4.1).

2. (PA) Convolve the CO to a resolution of 150 pc (FWHM). Estimate the surface density and effective line width, correct for inclination to calculate face-on values, and calculate virial parameter, α_{vir} , and free-fall time, t_{ff} . Following Sun et al. (2022), they use the following definition of average free-fall time:

$$\langle t_{\text{ff}} \rangle = \left(\frac{\sum_i M_i (t_{\text{ff}i})^{-1}}{\sum_i M_i} \right)^{-1}, \quad (2)$$

where M_i and $t_{\text{ff}i}$ are the mass and free-fall time of object i . This method actually averages speeds of formation and weights by mass, then converts to t_{ff} . The value of $t_{\text{ff}i}$ is calculated from

$$t_{\text{ff}i} = \sqrt{3\pi/(32G\rho)}, \quad (3)$$

and

$$\rho = 3M/(4\pi R_{\text{pix}}^3). \quad (4)$$

Finally,

$$R_{\text{pix}} = \min[l/2, (l^2 H/(8 \cos i))^{1/3}], \quad (5)$$

where l is the side of the cell, H is the scale height of molecular gas (taken to be 100 pc) and i is the inclination angle of the galaxy.

(CP) We have catalogs of these properties already and higher resolution. Each cloud is assigned to a rectangular cell, 150 pc on a side, based on the Galactic coordinates of its centroid and its distance. Clouds are not split between cells. The cloud distance is the primary source of uncertainty, especially in regions where kinematic distances are questionable, such as the region of the bar (Baba 2026). The procedure is described in more detail in E25, where it was used for 500 pc cells.

While L25 defines $t_{\text{ff}i}$ as the free-fall time of “cloud” i , that definition is for 150 pc resolution; for comparison, we use $t_{\text{ff,cell}}$, calculated from the mean density of a cell (mass of cell divided by volume). For the Milky Way, $\cos i = 1$, and $R_{\text{pix}} = 65.5$ pc. We calculate both the average of the cloud level properties from our catalog and the smoothed value based on resolution of the 150 pc cell, labeled “cell.” For example, the average surface density of all the clouds in a cell is $\langle \Sigma_{\text{cloud}} \rangle$,

while the smoothed value is $\Sigma_{\text{cell}} = M_{\text{cell}}/A_{\text{cell}}$, where M_{cell} is the total mass of all the clouds in the cell and A_{cell} is the area of the cell (2.25×10^4 pc²). The latter is the equivalent of $\Sigma_{\text{mol}}^{\text{cloud}}$ of L25 while the former is not available in PHANGS because of resolution. Note that the superscript “cloud” in the L25 notation denotes cloud-scale rather than cloud, but we will use cell because many clouds exist in a typical 150 pc cell. One issue with using the catalog, rather than the original CO data arises in the linewidth. We use only the linewidth of clouds, averaged (unweighted) over the cell, which does not take into account velocity differences of clouds within the cell, so we are less likely to overestimate the velocity dispersion within a cell. The virial parameter of the cell is computed from the total molecular mass M_{cell} and radius $r_{\text{cell}} = 65.5$ pc to match the L25 method, along with the velocity dispersion $\sigma_{\text{v}}^{\text{cell}}$ (so it could be underestimated as well):

$$\alpha_{\text{vir,cell}} = 5r_{\text{cell}}\sigma_{\text{v}}^{\text{cell}2}/(GM_{\text{cell}}). \quad (6)$$

3. (PA) Divide the galaxy into hexagonal apertures with diameter 1.5 kpc. Within each aperture, compute the mass-weighted expectation value of the surface density, velocity dispersion and virial parameter from the “cloud” level values (really the 150 pc cell values).

(CP) We also divide the Milky Way into 1.5 kpc hexagons and compute average values of properties within those hexagons. Our standard procedure also uses mass-weighting, where mass is the total mass of the cell, M_{cell} , to best replicate the procedure in L25. The results are labeled with “hex,” as in $\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$ for the mass surface density. Because we have information on individual clouds, we also compute the average of cloud surface densities, again mass weighted by the mass in the cell: $\langle \Sigma_{\text{cloud}} \rangle_{\text{hex}}$. For consistency with our limitation of $R_{\text{Gal}} < 20$ kpc, we restrict the centers of the hexagons. At this stage, we add information on the 70 μm SFR, averaged over the hexagon. Figure 1 shows the hexagons with the surface densities of molecular gas and star formation rate from E25 in color code. For further analysis, each hexagon must have non-zero molecular mass. This requirement cuts the number of hexagons to 316. The total SFR of the Milky Way decreased only slightly from 1.41 to 1.21 $M_{\odot} \text{ yr}^{-1}$, indicating that, according to distance assignments done in E25, little 70 μm emission resides outside

hexagons with molecular clouds. Comparison of the two panels of Figure 1 shows that most of the hexagons without molecular mass are on the far side of the Milky Way.

4. (PA) Calculate the molecular gas depletion time within the 1.5 kpc hexagons by convolving Σ_{SFR} and Σ_{mol} to 1.5 kpc resolution and dividing them. (CP) We used the prediction of star formation rate from the 70 μm values for each hexagon. Then, we computed the depletion time in each hexagon, $t_{\text{dep,hex}}$, from the total star formation and mass in each hexagon. This seems the closest approach to recreating the L25 procedure, although we do not exactly match the convolution process.
5. (PA) Divide the average cloud t_{ff} by t_{dep} to estimate the efficiency per free-fall time for each 1.5 kpc region. (CP) We do the same with the definition of t_{ff} in equation 2. We explore the effects of using different definitions in Appendix C. Using our notation, $\epsilon_{\text{ff}}(\text{hex}) = \langle t_{\text{ff}} \rangle_{\text{hex}} / t_{\text{dep,hex}}$.
6. (PA) Limit the analysis to regions with surface densities at the hex-scale ($\langle \Sigma_{\text{mol}}^{\text{cloud}} \rangle$, see Table 1) greater than $20 \text{ M}_{\odot} \text{ pc}^{-2}$. (CP) We apply the same cut to our hexagons, using $\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$, which is our equivalent. At this stage, we also require a non-zero value for the 70 μm SFR in the hexagon. This step cuts the number of hexagons to 195, eliminating many of the outer Milky Way hexagons (Figure 1). The total 70 μm SFR drops only to $1.10 \text{ M}_{\odot} \text{ yr}^{-1}$, indicating that there is little star formation at lower surface densities.

We create three products for each property: the average of the cloud level properties from the catalog, grouped into 150 pc cells, labeled “cloud”; the value based on resolution of the 150 pc cell, labeled “cell”; and the mass-weighted average over each 1.5 kpc hexagon, labeled “hex.” The last two are most directly comparable to the PHANGS-ALMA results, while the first tests the effects of the limited resolution and sensitivity. See Table 1 for definitions of symbols used by L25 and here.

As a result of these procedures, we have values for the Milky Way that have been derived in a way that mimics the procedures used for other galaxies. We will also have information on the effect of spatial resolution on the extragalactic analysis.

Appendix A addresses relatively simple issues of how to compare the two datasets. In §4, we discuss more complex or consequential issues.

4. PRELIMINARY CONSIDERATIONS

4.1. Luminosity-to-Mass Conversion

The mass of molecular gas is related to the luminosity of a CO line by the conversion factor, α_{CO} with units $\text{M}_{\odot}(\text{K km s}^{-1} \text{ pc}^2)^{-1}$. Until recently, studies of the Milky Way have used a constant value for α_{CO} , but recent studies covering a wider range of R_{Gal} have corrected for metallicity effects (Evans et al. 2022; Elia et al. 2025a). In this regard, the extragalactic community has been ahead, faced with galaxies of widely different metallicities (Bolatto et al. 2013) and excitation conditions (Downes & Solomon 1998). In the most recent authoritative formulation, Schinnerer & Leroy (2024) favor the following formula:

$$\alpha_{\text{CO}} = \alpha_{\text{CO},\odot} f(Z) g(\Sigma_{\star}), \quad (7)$$

where Z is the metallicity, $f(Z)$ corrects for metallicity effects, $\Sigma_{\star}$ is the surface density of stars, and $g(\Sigma_{\star})$ corrects for the enhancement of CO emission in regions of high stellar density, like centers of galaxies.⁹ We refer to this model as SL.

The factor $f(Z)$, which they refer to as the CO-dark factor, has been much debated. Schinnerer & Leroy (2024) recommend

$$f(Z) = (Z/Z_{\odot})^{-a} \quad (8)$$

and favor $a = 1.5$ in the range of 0.2 to 2.0 solar. We define Z to be relative to solar, so here we use $f(Z) = Z^{-a}$.

The factor $g(\Sigma_{\star})$, called the starburst factor, accounts for the fact that clouds in galactic centers are more luminous per unit mass (Schinnerer & Leroy 2024).

$$g(\Sigma_{\star}) = \left(\frac{\max(\Sigma_{\star}, 100 \text{ M}_{\odot} \text{ pc}^{-2})}{100 \text{ M}_{\odot} \text{ pc}^{-2}} \right)^{-0.25}. \quad (9)$$

It is only relevant for $\Sigma_{\star} > 100 \text{ M}_{\odot} \text{ pc}^{-2}$. We estimate its effects in Appendix B. It only affects the inner 2 kpc of the Milky Way and by less than a factor of 2.

In addition, recent papers including L25 use a variable R_{21} to convert from $J = 2 \rightarrow 1$ to $J = 1 \rightarrow 0$ intensity that depends on the surface density of star formation (Sun et al. 2025):

$$R_{21} = 0.65(\Sigma_{\text{SFR}}/0.018)^{0.125}, \quad (10)$$

with Σ_{SFR} measured in $\text{M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$. This is not relevant to the Milky Way, for which the surveys are in the $J = 1 \rightarrow 0$ transition.

⁹ They use $\alpha_{\text{CO,MW}}^{1-0}$, but we replace “MW” with the Sun symbol to mean the value determined in the solar neighborhood because there is no single value for the Milky Way, and we drop the superscript because our data are all for $J = 1 \rightarrow 0$.




Figure 1. Layout of the hexagonal bins showing the molecular gas surface density (left) and the 70 μm SFR surface density (right). The right panel presents the same data as Figure 5 (left) of E25, but re-binned from the original rectangular grid into hexagons. The white circle and X mark the positions of the Sun and the Galactic center, respectively, further highlighted by the intersections of the displayed vertical and horizontal dotted lines.

While many references still use a constant α_{CO} for the Milky Way, we have adopted for our recent papers (E25) a metallicity-dependent conversion factor,

$$\alpha_{\text{CO}}(Z) = 4.50Z^{-0.8} \quad (11)$$

with the exponent based on simulations (Gong et al. 2020; Hu et al. 2022). Previous studies in the MW show that $a = 0.8$ works best when using the Milky Way gradient in oxygen abundance (Evans et al. 2022; Elia et al. 2025a). The common choice for $\alpha_{\text{CO},\odot}$ is $4.35 \text{ M}_{\odot} (\text{K km s}^{-1} \text{ pc}^2)^{-1}$, based on adding helium, when converting the local “standard” X_{CO} of $2 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$. If all the elements are added, the correct value is $\alpha_{\text{CO},\odot} = 4.50$ (Evans et al. 2022), which we have adopted for the Milky Way.

While the SL formulae for $\alpha_{\text{CO}}(Z)$ depend much more strongly on Z than those we have used for the Milky Way, the effect is muted by the weaker dependence of Z on R_{Gal} used in PHANGS papers. For the Milky Way we used a measured oxygen abundance gradient of $-0.044 \pm 0.009 \text{ dex/kpc}$ (Méndez-Delgado et al. 2022). Other elements show even larger gradients. Lacking a consistent set of metallicity measurements for the full sample, the PHANGS team adopted metallicities from scaling relations, as described in detail by Sun et al.

(2022). In particular, their gradient is normalized to the effective (half-light) radius of the galaxy (r_e):

$$\log Z(R_{\text{Gal}}) = \log Z(r_e) - 0.1R_{\text{Gal}}/r_e \quad (12)$$

with $\log Z(r_e)$ a function of total stellar mass (Sánchez et al. 2014, 2019).

$$\begin{aligned} \log Z(r_e) = & 0.04 + 0.01(\log(M_{\star}/M_{\odot}) - 11.5) \\ & \times \exp(-\log(M_{\star}/M_{\odot}) + 11.5). \end{aligned} \quad (13)$$

We refer to the combination of $\alpha_{\text{CO}}(Z)$ and $Z(R_{\text{Gal}})$ models as SL in Figure 2.

To apply the PHANGS formulae to the Milky Way, we need to know $M_{\star}$ and r_e . Our knowledge of the stellar properties of the Milky Way is in a state of rapid change. Values for the total stellar mass differing by a factor of two have been published recently (Imig et al. 2025; Lian et al. 2025), along with estimates of r_e ranging from 4.12 kpc (Lian et al. 2025) to 4.75 kpc (Imig et al. 2025) to 5.75 kpc (Lian et al. 2024). We adopt $r_e = 4.75 \text{ kpc}$ and $\log M_{\star} = 10.72$ (Imig et al. 2025) for this purpose, yielding a gradient of -0.021 dex/kpc , less than half the measured value, as can be seen in the top panel of Figure 2.¹⁰ For comparison to ex-

¹⁰ For discussion of r_e for the Milky Way, see Appendix B.

tragalactic gradients, the measured metallicity gradient in the Milky Way of -0.044 dex/kpc and $r_e = 4.75$ kpc implies -0.209 dex/ r_e , larger than in any of the 19 PHANGS galaxies studied by Groves et al. (2023). Those are characterized by a mean of -0.059 ± 0.056 dex/ r_e , and the most negative value is -0.188 ± 0.005 for NGC1365. These formulae used for extragalactic work predict $Z(r_e) = 1.05$ and $Z(R_{\text{Gal},\odot}) = 0.70$, where we take $R_{\text{Gal},\odot} = 8.34$ kpc, for consistency with E25. The SL formula would be in serious disagreement with the metallicity gradient for the Milky Way, measured in nebular lines (Méndez-Delgado et al. 2022). Studies of stars also support a steeper metallicity gradient in the Milky Way (Anders et al. 2017), than what would be inferred from the SL formulae.

The various formulations for $\alpha_{\text{CO}}(R_{\text{Gal}})$ are plotted in the lower panel of Figure 2. The different formulations for $Z(R_{\text{Gal}})$ result in rather similar $\alpha_{\text{CO}}(R_{\text{Gal}})$ at large R_{Gal} if we use the SL models of $Z(R_{\text{Gal}})$; they would be very different if we used the SL formula for $\alpha_{\text{CO}}(Z)$ and the actual $Z(R_{\text{Gal}})$ measured for the Milky Way (green curve). The sharp-eyed reader may notice a downturn for the innermost point on the SL plots, caused by the $g(\Sigma_*)$ factor.

When comparing Milky Way data to the PHANGS data, we will use the same conversion methods (Schinnerer & Leroy 2024) that were used in the PHANGS data for consistency. As Figure 2 shows, the difference between these conversion methods and the one we believe applies to the Milky Way is very small. However, this seems a rather fortuitous cancellation of two effects.

4.2. Effects of Resolution

We consider the effects of the limited spatial resolution for extragalactic observations by sorting the Milky Way data into cells of 150 pc on a side, simulating the resolution of the PHANGS-ALMA data. We compare various properties when we average over the resolved clouds in the catalog compared to aggregating them into a single mass.

The surface density (Σ_{mol}) is often used to characterize molecular clouds, the volume density directly predicts the free-fall time, and the virial parameter summarizes the dynamical state. We compare in Table 2 the distribution of these three quantities for the MD17 sample, sorted into cells if we average the Σ_{mol} of the clouds (denoted Σ_{cloud}) versus dividing the total mass in the cell by the area of the cell (denoted Σ_{cell}), roughly approximating the extragalactic resolution. The low resolution clearly lowers the values of Σ_{mol} by about a factor of three (cf. entries 1 and 2 in Table 2). The tail to very low Σ_{cell} in Figure 3 reflects cells with very




Figure 2. Two models for $\alpha_{\text{CO}}(Z)$ are plotted versus R_{Gal} (bottom), along with two models for $Z(R_{\text{Gal}})$ (top). The Milky Way model for $Z(R_{\text{Gal}})$ assumes a gradient of -0.044 dex/kpc extending throughout the Milky Way (labeled E25). The model labeled SL uses the extragalactic model with a gradient of $-0.10/r_e$ dex. The model labeled SL-MWZ uses those models for $\alpha_{\text{CO}}(Z)$ but the measured $[\text{O}/\text{H}]$ gradient in the Milky Way (Méndez-Delgado et al. 2022).

few or only low mass clouds; these would escape detection by PHANGS-ALMA altogether, since their limit is $4 M_{\odot} \text{ pc}^{-2}$, as shown by the vertical line. In contrast, mass-weighted averaging over hexagons increases the surface density as shown by line 3 in Table 2, where $\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$ is somewhat larger than the actual average cloud surface density ($\langle \Sigma_{\text{cloud}} \rangle$), but the distribution of $\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$ is much closer to the actual distribution of Σ_{cloud} (Fig. 3). An extragalactic observer using 150 pc resolution to observe the Milky Way would underestimate the mean cloud-level surface density in a 150 pc beam, but slightly overestimate the mean surface density by mass-weighting the data into hexagons.

4.3. Effects of Sensitivity

We next ask how the Milky Way properties change if we institute cuts that simulate the sensitivity limits of PHANGS-ALMA. We compare basic results with the full catalog and with cuts at the cloud level where we require an intensity of CO $J = 1 \rightarrow 0$ emission of 1.0 K km s^{-1} , equivalent to their $J = 2 \rightarrow 1$ sensitivity of 0.65 K km s^{-1} (see Appendix A for details). Only 814 clouds fail the CO intensity criterion, with a total mass of $6 \times 10^7 M_{\odot}$. Table 2 in lines labeled “MW-L25” provides comparisons of the median, mean, and standard deviation of the average densities ($\bar{n}$), surface densities

Table 2. Effects of Resolution and Sensitivity

Sample	Number	$\log(\bar{n}/\text{cm}^{-3})$			$\Sigma_{\text{mol}} (\text{M}_{\odot} \text{ pc}^{-2})$			α_{vir}			$\log(M_{\text{tot}}/\text{M}_{\odot})$	Note
		Med	Mean	Std.	Med	Mean	Std.	Med	Mean	Std.		
MW-E25-cloud	7389	1.05	1.07	0.65	17.9	28.7	29.4	8.03	15.41	18.40	9.13	1
MW-E25-cell	4558	0.82	0.86	0.57	4.3	13.1	27.1	10.88	36.11	117.44	9.13	2
MW-E25-hex	316	0.70	0.74	0.55	20.1	34.0	44.0	6.89	11.19	20.50	9.11	3
MW-L25-cloud	6575	1.14	1.14	0.64	21.2	31.7	29.8	7.50	14.52	17.71	9.12	4
MW-L25-cell	3871	0.94	0.94	0.57	6.1	15.2	28.9	9.40	28.51	81.10	9.12	5
MW-L25-hex	307	0.72	0.76	0.55	20.8	35.1	44.3	6.56	10.88	20.80	9.11	6

NOTE— 1. Data from catalog in [Elia et al. \(2025a\)](#) using their criteria: $M_{\text{mol}} > 1 \text{ M}_{\odot}$, $R_{\text{gal}} < 20 \text{ kpc}$, and $\alpha_{\text{vir}} < 100$.

2. Same as 1 but assuming resolution of a 150 pc cell.

3. Same as 1 but based on mass-weighted averages in 1.5 kpc hexagons.

4. Data from catalog in [Elia et al. \(2025a\)](#) adding criterion on $I_{\text{CO}} > 1.0 \text{ K km s}^{-1}$.

5. Same as 4 but assuming resolution of a 150 pc cell.

6. Same as 4 but based on mass-weighted averages in 1.5 kpc hexagons.




Figure 3. (a) The logarithmic mass surface density distribution for CO clouds calculated two ways, both normalized to the peak value. The cyan histogram is the distribution of Σ_{cloud} of clouds averaged over the cell ($\langle \Sigma_{\text{cloud}} \rangle$ in Table 1). The red histogram is the ratio of the total mass in the cell to the area of the cell (Σ_{cell} in Table 1). (b) The distribution of the logarithm of the (mass) surface density distribution for CO clouds calculated three ways, all normalized to the peak value. The red histogram is the ratio of the total mass of the cell to the area of the cell (Σ_{cell}), repeated from panel a. The green histogram is the mass-weighted average of the cloud level surface density (Σ_{mc}) over the hexagons ($\langle \Sigma_{\text{cloud}} \rangle_{\text{hex}}$ in Table 1). The magenta histogram is the mass-weighted average of Σ_{cell} over the hexagons ($\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$ in Table 1). No sensitivity limit is used for the sample. The vertical line is $\log(4 \text{ M}_{\odot} \text{ pc}^{-2})$, the sensitivity limit of PHANGS-ALMA.

(Σ_{mol}), and virial parameters (α_{vir}). Adding the criterion on I_{CO} corresponding to the PHANGS-ALMA sensitivity limits raises the values of $\bar{n}$ and Σ_{mol} and lowers

the values for α_{vir} for the Milky Way clouds because less populated cells are left out of the averages, but the effect is very small.

We use only the criteria for clouds used by E25 for the next set of comparisons.

4.4. Tracers and Cloud Identification Methods

Because different tracers (CO $J = 1 \rightarrow 0$, CO $J = 2 \rightarrow 1$, extinction) and different cloud separation methods have been used to define clouds, we compare the properties from some characteristic references. These both affect the definition of a cloud. For example, Wang et al. (2025) have produced a catalog of Milky Way clouds based on extinction, but favoring the denser, smaller leaves on a dendrogram analysis. The comparison to properties from CO and other dust-based catalogs emphasizes the inherent ambiguities in cloud definition.

We explore first the properties of the samples of clouds in the Milky Way, as compared to a similar PHANGS galaxy. We choose NGC4321. This is a barred galaxy, with data in both PHANGS, as well as Faustino Vieira et al. (2025), who derive cloud properties from extinction. For the properties measured from CO $J = 2 \rightarrow 1$, we restrict the entries in L25 Table 2 (Leroy et al. 2026) to this galaxy. Because density and mass are not given, we calculate the total mass and $\bar{n}$, from the surface density, assuming a volume of the hexagon with a total thickness of 100 pc. For NGC4321 measured in extinction (Faustino Vieira et al. 2025), we get mass from “Mass”, radius from “R_eq”, and area from “Area_exact”. As explained in Faustino Vieira et al. (2023, equation 7), the surface density estimates are effectively scaled to an assumed opacity at 250 μm of 21.6 $\text{cm}^2 \text{g}^{-1}$. That opacity model assumed grain growth and ice accumulation for 10^6 yr at a density, $n = 10^6 \text{ cm}^{-3}$ (Ossenkopf & Henning 1994), implausible conditions for most molecular cloud material. The most recent evaluation of this opacity for molecular clouds (KP5) is 8.12 $\text{cm}^2 \text{g}^{-1}$ (Pontoppidan et al. 2024), constructed to reproduce opacity laws in molecular clouds by Chapman et al. (2009) and quite consistent with recent JWST observations (e.g., Rubinstein et al. 2024). We use that opacity instead of the one used by Faustino Vieira et al. (2025).

For the Milky Way, we use the catalog in E25, but also the catalog from Rice et al. (2016), labeled Rice in figures and tables; this catalog was made from the same CO $J = 1 \rightarrow 0$ data but a different cloud identification method. We are using here the original cloud catalogs, not values averaged into hexagons. In particular we compare surface density (Σ_{mol} , Figure 4), mean density ($\bar{n}$, Figure 4), and virial parameter (α_{vir} , Figure 4). Statistics of the distributions are presented in Table 3.

Tracer choice can have a substantial effect on derived surface density (panel a of Figure 4). The distribution from extinction mapping (Faustino Vieira et al. 2025) finds surface densities lower by about a factor of two compared to the CO $J = 2 \rightarrow 1$ distribution from the same galaxy. (Before adjusting the opacities, the factor was 5.2.) This difference is probably mostly due to the lower surface density sensitivity limit for the extinction method and the mass-weighting for the CO data, as well as limitations on the maximum surface densities measurable with the extinction method (Faustino Vieira et al. 2023, 2025). The extinction values are however similar to Milky Way values.

The two methods of cloud identification for Milky Way clouds have quite similar distributions and statistical properties, though the Rice values are 1.2-1.3 times higher. Tracer choice seems to have a larger effect on surface density estimates than does cloud identification method, especially with uncertainties in opacity laws.

For the volume density ($\bar{n}$), the results are quite different (panel b of Figure 4). While the four distributions differ in width, their medians and means are striking in their similarity. The smaller cloud sizes for the extinction method result in somewhat higher $\bar{n}$ values, but still only different by a factor of three.

The distributions of virial parameter present yet a different story (panel c of Figure 4). Extinction measurements provide no measure of virial parameter, so we have only three distributions. The CO $J = 2 \rightarrow 1$ distribution is centered at lower values of α_{vir} than are the Milky Way distributions, though the Rice distributions are much more similar than those of E25. This difference has been discussed by Evans et al. (2021).

To summarize, the choice of tracer has a direct effect on estimates of surface density, but little apparent effect on volume density for the three considered here. In contrast, the cloud identification method has a major effect on virial parameter, but modest effects on surface density and volume density.

5. MOLECULAR GAS DEPLETION TIME AS A FUNCTION OF OTHER PROPERTIES

In this section, we compare the Milky Way to results from PHANGS with plots like those in §3.1 to §3.3 of L25. Table 4 provides the statistical properties of the variables for the Milky Way, while Table 5 contains the equivalent statistics for the data in the Erratum, (Leroy et al. 2026). We follow the procedures in L25 as closely as possible. In particular, we do not apply any criteria on I_{CO} or Σ_{mol} at the cloud level; the criterion on Σ_{mol} is applied only to analysis of data at the hexagon level,

Table 3. Effects of Tracers and Definitions

Sample	Number	$\log \bar{n}$ (cm^{-3})			Σ_{mol} ($\text{M}_{\odot} \text{pc}^{-2}$)			α_{vir}			$\log M_{\text{tot}}$ ($\text{M}_{\odot}$)	Note
		Med	Mean	Std.	Med	Mean	Std.	Med	Mean	Std.		
NGC4321 A_V	24196	1.46	1.43	0.24	27.3	31.4	16.3	...	...	...	9.36	1
NGC4321 $J = 2 \rightarrow 1$	124	0.92	0.96	0.24	48.3	61.5	41.5	1.51	1.95	1.24	9.64	2
MW Rice	1037	1.04	1.04	0.50	21.2	38.3	43.4	2.44	3.45	3.34	8.39	3
MW E25	7389	1.05	1.07	0.65	17.9	28.7	29.4	8.03	15.41	18.40	9.13	4

NOTE— 1. Data from catalog in [Faustino Vieira et al. \(2025\)](#) using extinction to define clouds.

2. Data from catalog in [Leroy et al. \(2025\)](#) using CO $J = 2 \rightarrow 1$ to define clouds.

3. Data from catalog in [Rice et al. \(2016\)](#), using higher threshold to define clouds.

4. Data from catalog in [Elia et al. \(2025a\)](#) using criteria of E25: $M_{\text{mol}} > 1 \text{ M}_{\odot}$, $R_{\text{gal}} < 20 \text{ kpc}$, and $\alpha_{\text{vir}} < 100$.




Figure 4. The histograms of (a) surface density (Σ_{mol}), (b) average density ($\bar{n}$), and (c) virial parameter (α_{vir}) for the Milky Way (from E25 in green and from R16 in light green, respectively) and for NGC4321, both from CO $J = 2 \rightarrow 1$ (L25) in blue and extinction ([Faustino Vieira et al. 2025](#)) in deep sky blue, respectively. The last distribution has been scaled to account for lower opacities as discussed in §4.4.

consistent with step 6 in §3. Therefore the statistical properties of the variables in Table 4 differ from those in Table 2. As explained in [Leroy et al. \(2026\)](#), some points in the figures published by L25 were incorrectly placed. We show the equivalent to Table 3 in L25 with correlation and fit information in Table 6.

The statistics in Table 4 would apply to the Milky Way, as analyzed as much as possible as a galaxy in the PHANGS sample. Comparing the mean values to the entries in Table 5 for the L25 sample indicates that the depletion and free-fall times for the Milky Way are longer, the velocity dispersion is lower, the virial parameter is higher, and the surface densities are very similar. All these properties have large dispersions, so the Milky Way clearly fits within the distribution of properties of the L25 sample. The efficiency per free-fall time, averaged over all hexagons, has a mean value of 0.91% in the Milky Way and 0.55% for L25. The “theoretical”

Table 4. Statistics of Key Parameters For Milky Way

Quantity	Median	Mean	Std Dev	Units
$t_{\text{dep,hex}}$	1.42	3.68	10.12	Gyr
$\langle t_{\text{ff}} \rangle_{\text{hex}}$	9.37	9.21	2.49	Myr
$\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$	42.53	59.53	50.17	$\text{M}_{\odot} \text{pc}^{-2}$
$\langle \sigma_{\text{v}}^{\text{cell}} \rangle_{\text{hex}}$	5.91	6.01	1.80	km s^{-1}
$\langle \alpha_{\text{vir}} \rangle_{\text{hex}}$	4.11	4.94	3.14	...
$\epsilon_{\text{ff}}(\text{hex})$	0.58	0.91	0.98	%
$\langle \epsilon_{\text{ff}}(\text{EKO}) \rangle_{\text{hex}}$	0.50	0.89	1.15	%

NOTE— 1. $\epsilon_{\text{ff}}(\text{hex})$ is $\langle t_{\text{ff}} \rangle_{\text{hex}} / t_{\text{dep,hex}}$ for each hexagon.

2. $\langle \epsilon_{\text{ff}}(\text{EKO}) \rangle_{\text{hex}}$ is calculated for each hexagon from $\langle \alpha_{\text{vir}} \rangle_{\text{hex}}$ and the EKO formula.

Table 5. Statistics of Key Parameters For L25 Data

Quantity	Median	Mean	Std Dev	Units
t_{dep}	2.13	2.50	1.76	Gyr
$\langle \tau_{\text{ff}}^{\text{cloud}} \rangle$	8.68	8.62	2.40	Myr
$\langle \Sigma_{\text{mol}}^{\text{cloud}} \rangle$	38.14	56.52	54.71	$\text{M}_{\odot} \text{ pc}^{-2}$
$\langle \sigma_{\text{mol}}^{\text{cloud}} \rangle$	5.43	7.12	4.95	km s^{-1}
$\langle \alpha_{\text{vir}}^{\text{cloud}} \rangle$	2.01	3.08	3.95	...
$\epsilon_{\text{ff}}(\text{hex})$	0.39	0.55	0.49	%
$\langle \epsilon_{\text{ff}}(\text{EKO}) \rangle_{\text{hex}}$	1.72	1.90	1.31	%

NOTE— 1. $\epsilon_{\text{ff}}(\text{hex})$ is $t_{\text{ff}}/t_{\text{dep}}$ for each hexagon.

2. $\langle \epsilon_{\text{ff}}(\text{EKO}) \rangle_{\text{hex}}$ is calculated for each hexagon from $\langle \alpha_{\text{vir}} \rangle_{\text{hex}}$ and the EKO formula.

version, $\langle \epsilon_{\text{ff}}(\text{EKO}) \rangle_{\text{hex}}$, calculated from the virial parameters of each hexagon, agrees with the observed value in the Milky Way, but is nearly 4 times larger than the observed value in the L25 sample.

Following L25, we test correlations with the Spearman test, with r_s the correlation coefficient and p the probability that the assumption of no correlation can be rejected. Values of $p \leq 0.05$ are needed for rejection of the null hypothesis. We also fit the data in each plot to the formula

$$y = m(x - x_0) + b, \quad (14)$$

where y is $\log_{10} t_{\text{dep}}$ and x is the decimal logarithm of the variable being tested, while x_0 is the median of x . We use a different fitter (LevMarlsq from astropy modeling) but our results differ from those in Leroy et al. (2026) by only insignificant amounts.

First, we reproduced Figure 4 through 7 in L25 (1 through 4 in Leroy et al. (2026)). After checking that the corrected plots were reproduced, we retained only a fit to the L25 data as a blue line for comparison in the plots of Milky Way data. The data points for the Milky Way are shown as filled circles, color-coded by R_{Gal} . The fit to the Milky Way data is shown as a green line. For each plot, we also fitted the data restricted to the region occupied by the data in L25, indicated by the orange box, and those fits are represented by orange lines. The right axis shows the data normalized to the median value of depletion time for comparison to those figures in L25.

Figure 5 (the equivalent of Figure 4 of L25, Figure 1 of Leroy et al. (2026)) shows no significant correlation between the depletion time and the free-fall time, using our hex-level data. Including only the points lying within the boundaries of the L25 data removes the hexagons

with large t_{dep} , but the fit is scarcely affected. The Milky Way lacks hexagons with very low values of free-fall time found in the PHANGS sample, albeit rarely, as shown by the empty area to the left of the orange box.

Next we make plots analogous to their figures 6 and 7 of L25 (Figure 3 of the Erratum), corresponding to the fits in their Table 3. The right panel of figure 5 shows the depletion time versus mass surface density. For the Milky Way, we use the surface density, smoothed over the cell and mass-weighted averaged into hexagons, $\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$. This is closest in spirit to what L25 does. We see a barely significant correlation.

The average σ_v over the clouds in a cell, mass averaged into hexagons (panel a of Figure 6, equivalent to Figure 6 of L25, the lower panel of Figure 3 of Leroy et al. (2026)) shows, like the extragalactic data, a negative correlation, but the Milky Way plot lacks the large σ_v points. At least in part, this is because the Milky Way values are average σ_v at the cloud level, which do not include velocity differences between clouds that are combined into σ_v with the spatial resolution of 150 pc.

The dependence of depletion time on virial parameter (panel b of Figure 6, equivalent to Figure 7 of L25, and 4 of (Leroy et al. 2026)) indicates a slight negative correlation, very similar to that of the PHANGS data. Recall that our star formation rate is based on the 70 μm emission, SFR_{70} . If we use the CO mass and the formula from EKO that is based on a relation with α_{vir} , we get a strong positive correlation, but this would be circular.

This section has shown that the Milky Way, analyzed similarly to the PHANGS sample, has similar dependences of t_{dep} on various parameters. There are relatively strong anti-correlations between t_{dep} and line width or virial parameter. Correlations between t_{dep} and $\langle \tau_{\text{ff}} \rangle_{\text{hex}}$ or $\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$ are very weak and probably not significant; these were also weak but more significant in the PHANGS data. A plot of the slopes of the fitted lines with uncertainties for both the PHANGS galaxies and the Milky Way (Figure 7) shows that these differ but only at the 2σ level. The errorbars on the Milky Way are larger because there are many fewer data points.

As L25 note, the results of these correlations change substantially if a fixed value of α_{CO} is used instead of the standard procedure. We confirm this fact (Appendix E, Table 7) and find that the differences in the fits between the Milky Way and the PHANGS galaxies are smaller with this choice (Figure 13). While we agree with L25 that the evidence for a variable α_{CO} is strong, remaining uncertainties in this function are problematic.

Table 6. Correlation coefficients and fit parameters of $\log t_{\text{dep}}$ as function of x

Sample	x	r_s	p	m	σ_m	x_0	b	σ_b
PHANGS	$\log \langle t_{\text{ff}} \rangle_{\text{hex}}$	0.18	2.81e-07	0.51	0.08	6.94	9.31	1.06e-02
MW	$\log \langle t_{\text{ff}} \rangle_{\text{hex}}$	-0.05	5.68e-01	-0.13	0.29	6.97	9.21	3.84e-02
MW (PHANGS)	$\log \langle t_{\text{ff}} \rangle_{\text{hex}}$	-0.05	5.68e-01	-0.25	0.25	6.97	9.15	3.84e-02
PHANGS	$\log \langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$	-0.22	1.40e-10	-0.31	0.04	1.58	9.32	1.07e-02
MW	$\log \langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$	0.06	4.93e-01	0.08	0.14	1.63	9.21	3.85e-02
MW (PHANGS)	$\log \langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$	0.06	4.93e-01	0.13	0.13	1.63	9.15	3.85e-02
PHANGS	$\log \langle \sigma_v^{\text{cell}} \rangle_{\text{hex}}$	-0.31	6.28e-20	-0.51	0.05	0.74	9.33	1.05e-02
MW	$\log \langle \sigma_v^{\text{cell}} \rangle_{\text{hex}}$	-0.25	1.39e-03	-1.14	0.28	0.77	9.20	3.60e-02
MW (PHANGS)	$\log \langle \sigma_v^{\text{cell}} \rangle_{\text{hex}}$	-0.25	1.39e-03	-0.61	0.27	0.77	9.14	3.60e-02
PHANGS	$\log \langle \alpha_{\text{vir}} \rangle_{\text{hex}}$	-0.27	3.67e-15	-0.32	0.03	0.30	9.31	3.41e-02
MW	$\log \langle \alpha_{\text{vir}} \rangle_{\text{hex}}$	-0.25	1.38e-03	-0.37	0.13	0.61	9.21	3.67e-02
MW (PHANGS)	$\log \langle \alpha_{\text{vir}} \rangle_{\text{hex}}$	-0.25	1.38e-03	-0.27	0.11	0.61	9.16	3.67e-02

NOTE— 1. MW (PHANGS) denotes MW data restricted to range of PHANGS data.




Figure 5. In panel a, the hex-level depletion time is plotted versus cloud-average free-fall time, mass-weighted averaged into hexagons, for the Milky Way. The covariance of the uncertainties in these parameters is shown in Figure 4 of L25. Panel b has the equivalent plot for $\langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$. In both panels, the points are color-coded to show R_{Gal} . The fit to the Milky Way data is shown in green, and the fit to Milky Way values restricted to the range of those in L25 is shown in orange. The orange box is the region including all the points in the PHANGS data. The blue line is the fit to the equivalent data on the PHANGS galaxies. The right axis shows the data normalized to the median value of depletion time for comparison to those figures in L25.

6. RADIAL DISTRIBUTIONS

Many of the properties we are studying vary systematically with R_{Gal} , as can be seen from the color coding, so we also plotted them versus R_{Gal} , following the procedures in L25. We first reproduced the plots of properties in the PHANGS sample versus galactocentric radius in L25. Then we plotted the same properties versus R_{Gal}

for the Milky Way. Figure 8 is the equivalent of Figure 8 of L25 (Figure 5 in Leroy et al. (2026)) for the Milky Way. Patterns are similar. Rather than their running median method, we use scipy binned statistics to compute median, mean, and standard deviation in 20 bins. The Milky Way shows very different values in the innermost hexagon (henceforth the CMZ) in most




Figure 6. In panel a, the mean depletion time is plotted versus the cloud-average velocity dispersion, mass-weighted averaged into hexagons, for the Milky Way. In panel b, the plot is versus $\langle\alpha_{\text{vir}}\rangle_{\text{hex}}$. For both panels, the points are color-coded to show R_{Gal} . The fit to the Milky Way data is shown in green, and the fit to Milky Way values restricted to the range of those in L25 is shown in orange. The orange box is the region including all the points in the PHANGS data. The blue line is the fit to the equivalent data on the PHANGS galaxies. The right axis shows the data normalized to the median value of depletion time for comparison to those figures in L25.




Figure 7. The slopes of the fits in Table 6 are plotted with uncertainties for both the PHANGS galaxies and the Milky Way.

plots, similar to many PHANGS galaxies. As is true for the PHANGS data, the only variables that show a clear trend with R_{Gal} outside the CMZ are $\langle\sigma_{\text{v}}^{\text{cell}}\rangle_{\text{hex}}$ and $\langle\alpha_{\text{vir}}\rangle_{\text{hex}}$. Zhou et al. (2025) also see a clear decrease of velocity dispersion with R_{Gal} for the clouds in their survey area.

The Milky Way and the galaxies in L25 generally exhibit the same trends with R_{Gal} . However, in the region of $1 < R_{\text{Gal}} < 8$ kpc, the Milky Way tends to have higher $\langle\sigma_{\text{v}}^{\text{cell}}\rangle_{\text{hex}}$ and $\langle\alpha_{\text{vir}}\rangle_{\text{hex}}$. These differences may be related to the fact that the Milky Way has a decrease

in the molecular surface density and star formation rate (Elia et al. 2022, 2025a), associated with a bar in the inner Milky Way (Benjamin et al. 2005; Lucey et al. 2023). While other barred galaxies support the interpretation of the Milky Way data as a real dip in the molecular gas (Evans et al. 2026), it is still true that distances in that region may be incorrect. A recent analysis of the Milky Way data using simulations of barred galaxies also found a dip of about the same depth, but perhaps a smaller radial extent (Baba 2026).

7. DISCUSSION

When analyzed in a similar way, the Milky Way data occupies areas in the plots of t_{dep} versus other parameters that are similar to the PHANGS galaxies with some differences. In particular, both datasets show tremendous scatter about any relationships. On the scale of 1.5 kpc hexagons, mass-weighted properties of clouds/cells of diameter 150 pc produce little correlation between depletion time and either free-fall time or mass surface density. The strongest correlation is a negative one between depletion time and velocity dispersion, seen in both the PHANGS sample and the Milky Way, suggesting faster star formation in more turbulent gas. Velocity dispersion correlates with mass, and more massive clouds are more likely to be bound (Evans et al. 2021), possibly explaining the correlation. On the other hand, t_{dep} anticorrelates with virial parameter, suggesting faster star formation in *less* bound gas. This fact, seen in




Figure 8. Milky Way version of Figure 8 of L25. The magenta line shows the running median with 20 bins, while the green line with error bars shows the mean and standard deviation. The blue line is the running median of the L25 data. The quantities on the y-axis follow the definitions in Table 1.

L25, is reproduced in the Milky Way. We first address the (small) differences between the L25 and Milky Way results. Then we discuss expected relations based on theory or empirical arguments. After a digression to consider further the effects of resolution, we discuss the comparison of the observations with expectations.

7.1. Differences between PHANGS galaxies and the Milky Way

As seen in Figure 7, the analysis of the PHANGS sample and Milky Way differ significantly about the slope in the $t_{\text{dep,hex}} - \langle t_{\text{ff}} \rangle_{\text{hex}}$ diagram. Figure 1 of Leroy et al. (2026) contains a number of data points with $\langle t_{\text{ff}} \rangle_{\text{hex}}$ significantly lower than any hexagons in the Milky Way. If the lower limit on $\langle t_{\text{ff}} \rangle_{\text{hex}}$ included in the fit to the L25 data is increased gradually, the correlation disap-

pears and the fitted slope becomes close to zero. The difference between L25 and the Milky Way is largely driven by the lack of hexagons with small $\langle t_{\text{ff}} \rangle_{\text{hex}}$ in the Milky Way. These low values occur most commonly at small galactocentric radii (Figure 8). Many of these exist in the PHANGS sample, but the Milky Way has only one central region. Similar considerations may explain the difference in slopes in the $t_{\text{dep}} - \langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$ diagram (Figure 5).

The Milky Way shows a steeper slope in the $t_{\text{dep}} - \langle \sigma_{\text{v}}^{\text{cell}} \rangle_{\text{hex}}$ diagram, perhaps caused by the hexagons with very large t_{dep} and low $\langle \sigma_{\text{v}}^{\text{cell}} \rangle_{\text{hex}}$. When the fit is restricted to the area in the L25 plots, the fit flattens to a slope of -0.61 , more similar to the L25 fit. As noted above the velocity dispersions may be underestimated because they ignore cloud-to-cloud variations that are unresolved in the extragalactic data. We tested the effects of spatial resolution by producing the same plot using the data with 60, 90, and 120 pc resolution in the tables in L25. The best-fitting slopes hardly differed; for example, for the 60 pc data, the slope was -0.56 ± 0.10 , not significantly different from that for 150 pc (-0.51 ± 0.05). Ongoing observations with better spatial resolution will allow further study of this issue.

In summary, the Milky Way differs from the PHANGS galaxies in relatively small ways, and some of the difference may be caused by selection effects.

7.2. Expected Relations

As discussed by L25, simple considerations and relations found for whole galaxies lead to some expectations for the relations plotted in §5. If the free-fall time is the pacing timescale for star formation, we would expect that $t_{\text{dep}} \propto t_{\text{ff}}$ if the efficiency per free-fall time (ϵ_{ff}) is constant (Krumholz et al. 2012; Pokhrel et al. 2021).

On the scale of whole galaxies, Σ_{SFR} correlates strongly with Σ_{mol} . As discussed by L25, clouds with similar ϵ_{ff} should show a dependence of

$$t_{\text{dep}} \propto \langle \Sigma_{\text{mol}} \rangle^{-a} \quad (15)$$

with $a = 0.5$. This is equivalent to a Kennicutt-Schmidt (KS) relation

$$\Sigma_{\text{SFR}} \propto \langle \Sigma_{\text{mol}} \rangle^{1+a}. \quad (16)$$

The KS relation for total gas is fitted well with $a = 0.4$ (Kennicutt 1998; Kennicutt & Evans 2012), while that including only the molecular component favors $a = 0$ (Bigiel et al. 2011). Studies of nearby molecular clouds where star formation and surface density can be traced to small scales find mixed results for a . Some studies found a linear dependence of Σ_{SFR} on Σ_{mol} ($a = 0$) above a threshold around $120 \text{ M}_{\odot} \text{ pc}^{-2}$ (e.g. Heiderman et al. 2010; Lada et al. 2010, 2012). More recent

studies that trace the structure within clouds, both in young stars and molecular gas, primarily at $\Sigma_{\text{mol}} > 100 \text{ M}_{\odot} \text{ pc}^{-2}$ find a stronger dependence of Σ_{SFR} on Σ_{mol} , with $a = 1.0$, but a possible decrease in a at very high Σ_{mol} (Pokhrel et al. 2021).

If, instead of t_{ff} , the crossing time is the pacing time for star formation (Elmegreen 2000), one might expect for fixed size that $t_{\text{dep}} \propto \sigma_v^{-1}$; L25 provide a more detailed rationale for this dependence based on the ‘‘Heyer-Keto’’ relation (Heyer et al. 2009) and a fixed virial parameter. In reality, neither size nor virial parameter is fixed at the cloud scale. However, size is fixed at 150 pc for the cell-based values and their hexagonal averages. Thus the crossing time for a region on the scale of 150 pc may be a relevant timescale.

Simulations predict that the star formation rate decreases rapidly with increasing virial parameter because the efficiency per free-fall time depends exponentially on the square root of α_{vir} (Padoan et al. 2012; Kim et al. 2021). Adopting values from Evans et al. (2022),

$$\epsilon_{\text{ff}}(\text{EKO}) = 0.30 \exp(-2.018 \alpha_{\text{vir}}^{1/2}), \quad (17)$$

so t_{dep} should increase exponentially with $\alpha_{\text{vir}}^{-0.5}$ because $t_{\text{dep}} \propto 1/\epsilon_{\text{ff}}$.

The fact is that none of these expected relations are seen in the data, with the exception of the $t_{\text{dep}} - \sigma_v$ relation in the Milky Way, and we noted earlier that this relation is quite sensitive to a few data points. We will explore possible explanations of each relation after further consideration of the effect of spatial resolution on the volume density and thus free-fall time.

7.3. Effect of Resolution Redux

The similarity of the average volume density ($\bar{n}$) across samples and tracers (Table 3 and Figure 4) is quite remarkable. The average values of the log of $\bar{n}$, translated to actual values are essentially all $10\text{--}30 \text{ cm}^{-3}$. This density is much lower than usually assumed for molecular clouds. The average densities are far below the critical density, which is often erroneously attributed to any emission from a molecule, but more consistent with effective densities, which will produce a 1 K km s^{-1} emission line for an extinction of 1 mag (Evans 1999; Shirley 2015). The effective densities are 15 cm^{-3} for CO $J = 1 \rightarrow 0$ and 60 cm^{-3} for $J = 2 \rightarrow 1$ (Evans et al. 2021). The mean densities of the clouds are smaller than the effective densities, but not greatly so.

However, molecular gas with a density of 10 cm^{-3} and the usually assumed $T_K = 10 \text{ K}$ would have very low pressure. To explore this issue further, we utilize the wide range of distances present in the MD17 catalog to study how $\bar{n}$ varies with distance and thus effective

spatial resolution on scales smaller than 150 pc. MD17 already discussed the effects of finite resolution on the radius, mass, and virial parameter in their Appendix C. Because at least $N_{\text{min}} = 5$ pixels were required to define a cloud, the effective angular resolution is larger than the beam size by $\sqrt{N_{\text{min}}}$. We discuss the effects on parameters of interest in terms of this effective spatial resolution given by the effective beam size times the distance. To avoid including the clouds in the CMZ, we limited the sample to $R_{\text{Gal}} \geq 0.5 \text{ kpc}$.




Figure 9. The median, mean, and standard deviation of $\log \bar{n}$ in bins of 1 kpc of distance from the Sun (upper axis). The effective resolution is indicated on the lower axis. The right axis shows the corresponding values of $\log t_{\text{ff}}$.

Figure 9 shows the density versus resolution at distances from 0 to 10 kpc. The corresponding t_{ff} from equation 3 is indicated on the right. While the median and mean $\log(\bar{n})$ over the whole sample correspond to about $\bar{n} = 10 \text{ cm}^{-3}$, this number clearly varies substantially with distance (resolution). While clouds within 1 kpc have mean $\bar{n}$ close to 100 cm^{-3} , the estimated $\bar{n}$ drops with distance. We interpret this to mean that many clouds are unresolved and the fraction of empty space within the identified cloud grows with distance. Beyond 35 pc resolution, the mean $\bar{n}$ levels off, suggesting that the observations are essentially in the cloud counting regime, and larger resolutions add clouds just as fast as individual clouds are beam-diluted. To test this idea, we plot the number of Gaussian components combined to define a cloud by MD17. In other definition schemes, these might be clouds being combined into a complex. The number is 2 for nearby clouds, rises to a

plateau around 4 for 3-6 kpc, then starts rising, reaching 6 at 9 kpc (Figure 10).




Figure 10. The median, mean, and standard deviation of the number of Gaussian components per cataloged cloud in MD17 in bins of 1 kpc of distance from the Sun (upper axis). The effective resolution is indicated on the lower axis.

This effect explains why the mean $\bar{n}$ over the whole sample is so low and why it agrees with the L25 results: for most of the Milky Way and all of the PHANGS galaxies, the average density refers to a region with many smaller clouds and substantial empty space. By empty space, we mean gas not emitting in the CO $J = 1 \rightarrow 0$ line. This could be CO-dark gas, Cold Neutral Medium (CNM), Warm Neutral Medium (WNM), or even ionized gas.

This pattern persists to smaller scales. The maps with the highest spatial dynamic range are those of nearby clouds, like the Taurus molecular cloud, with 3×10^6 pixels in CO (e.g., Goldsmith et al. 2008). They confirm that “clouds” are wispy, highly inhomogeneous, and likely associated with neutral atomic gas. Interestingly, Soler et al. (2023) calculate a mean volume density ($\bar{n}$) of about 10 cm^{-3} (after converting from density of nuclei), for the Taurus cloud using dust extinction. The volume density measured on smaller scales within the Taurus cloud varies from 75 cm^{-3} to 1200 cm^{-3} as one goes from the weakest CO emission to the strongest (Goldsmith et al. 2008).

The nearby clouds are not representative of the more massive regions of star formation, but studies of those

also find highly filamentary and fragmented structures on scales much smaller than available to either MD17 or PHANGS (e.g., Sanhueza et al. 2019; Xu et al. 2024; Schisano et al. 2026; Evans et al. 2020; Patra et al. 2025; Oakes et al. 2025, and references therein). Such a “Swiss cheese with many holes and little cheese” model was originally suggested by Zuckerman & Evans (1974).

One might hope to use the nearby clouds in the MD17 sample to define the actual $\bar{n}$ to use for calculating free-fall times of unresolved clouds. However, a blow up of the closest 1 kpc shows that $\bar{n}$ continues to rise with smaller resolution, reaching 300 cm^{-3} for the bin from 0 to 0.2 kpc (Figure 11). The density derived from observations of a cloud is always a function of the resolution and the tracer observed.




Figure 11. The median, mean, and standard deviation of $\log \bar{n}$ in bins of 0.2 kpc of distance from the Sun (upper axis). The effective resolution is indicated on the lower axis. The right axis shows the corresponding values of $\log t_{\text{ff}}$.

The virial parameter shows little change with resolution from 1-50 pc (Figure 12). Similarly, Oakes et al. (2025) find no emergent scale for bound objects between 3 and 300 pc in an analysis of molecular structure in NGC253. In the Milky Way, clearly bound objects emerge at scales of about 1 pc, where they depart from Larson’s laws (Plume et al. 1997; Peretto et al. 2023; Evans et al. 2021). Infall is well established at about 0.1 pc (Evans et al. 2015; Yang et al. 2020). These two scales are associated with the terms dense clumps and protostellar cores, respectively (Bergin & Tafalla 2007; McKee & Ostriker 2007). The velocity dispersion is also relatively insensitive to distance (resolution) with mean

values around $4 \pm 2 \text{ km s}^{-1}$. There is a rise at about 8 kpc to values closer to 6 km s^{-1} (Figure 12).

7.4. Observations versus Expectations

The fact that almost no clouds in the PHANGS samples or in the MD17 catalog are fully resolved could help explain the lack of correlation of t_{dep} and t_{ff} . The value of t_{ff} calculated from a collection of clouds will be overestimated and not really relevant to the time scale for depletion due to star formation. Highly structured clouds with low average densities can also explain the fact that t_{dep} depends weakly on surface density (Figure 5 and Table 6).

The L25 data show a negative correlation between t_{ff} and Σ_{mol} , while the Milky Way data show no correlation at all. The majority of the data points for both the PHANGS galaxies and the Milky Way lie at $\Sigma_{\text{mol}} < 120 \text{ M}_{\odot} \text{ pc}^{-2}$, where no strong correlation exists between surface density and SFR in Milky Way clouds (e.g. Heiderman et al. 2010; Lada et al. 2010, 2012). If the L25 data are restricted to the (only 59) points with $\Sigma_{\text{mol}} > 120 \text{ M}_{\odot} \text{ pc}^{-2}$, the values of a (Equation 15) become 0.63 or 0.77 for the normalized data. The data points that meet that criterion are mostly near galaxy centers, with median galactocentric radius of 1.5 kpc.

If the observations are primarily in the cloud-counting regime, and star formation is primarily dependent on the total mass of molecular gas, a simple prediction is that $\Sigma_{\text{SFR}} \propto \Sigma_{\text{mol}}$, and thus $t_{\text{dep}} \propto \Sigma_{\text{mol}}/\Sigma_{\text{SFR}}$ is constant with Σ_{mol} , as seen in the Milky Way data. Relations like KS work well on full galaxy scales and in some cases on structures within molecular clouds, but not when averaged over regions with a mix of star-forming gas and more diffuse gas. In addition, the measures of star formation rate are often displaced in time and space from the molecular gas, as first noted by Schruba et al. (2010), and emphasized by Kruijssen & Longmore (2014).

The lack of correlation between t_{dep} and α_{vir} suggests that virial parameters on 150 pc scales are not useful predictors of star formation rate. In contrast, including virial parameters on a cloud-by-cloud basis in the MD17 catalog produced good agreement on the overall star formation rate of the Milky Way, as well as the radial distribution (Evans et al. 2022). L25 suggest that the high velocity dispersion, high virial parameter regions are in the inner regions of galaxies, where higher densities and contributions from stellar surface density overcome the higher dispersions. This idea is supported by the trend to higher surface densities and shorter t_{ff} in the very central regions of both PHANGS galaxies and the Milky Way in Figure 8 and by Figure 6 showing

that the hexagons with high $\langle \alpha_{\text{vir}} \rangle_{\text{hex}}$ are in the inner Milky Way.

The one predicted relation that does appear in the data is that $t_{\text{dep}} \propto \sigma_v^{-1}$. The fit to the PHANGS data finds an exponent of about -0.5 but the MW data find -1.14 ± 0.28 , reasonably consistent with the simple prediction. (Recall, however, that we simply averaged cloud velocity dispersions to construct the cell-level value, as noted in §3.) In the crossing time picture, the velocity dispersion over 150 pc reflects the time for structural change: clouds merging, moving in or out of the cell, etc. In the Keto-Heyer picture, it would reflect more of an equilibrium state, but only if α_{vir} is fixed. In fact, α_{vir} varies widely in either the MD17 or R16 cloud catalogs.

7.5. Relation to Discussion in L25

A substantial discussion of how to interpret the PHANGS results is found in §4 of L25. All the interpretation in the previous section should be seen as “in addition to”, rather than as “instead of”, that discussion. The fact that the individual locations of star formation are not fully resolved even in the Milky Way adds emphasis to the points made there. The failure of expected relations to emerge in the data on kpc scales applies to the Milky Way as well. We agree with L25, who argue that the utility of these measurements lies in comparison to simulations that predict observables on similar scales. The large range of distances, and thus spatial resolution, available in the Milky Way data allows more stringent tests of simulations.

The Milky Way adds assurance that the trends seen in the PHANGS galaxies are not an artifact of observing at a large distance. We agree with the conclusion in L25 that “there is no clear evidence for a specific scale where the surface density traces physical density,” and Figures 9 and 11 extend that conclusion to scales as small as 1 pc.

8. SUMMARY

Analysis of cloud catalog data for the Milky Way with a method similar to that used for PHANGS galaxies produces broadly similar results. If the depletion time is plotted versus other parameters for which a relationship might be expected, the result is a very large scatter and relatively weak relationships. Simulations of star formation in galaxies should be able to reproduce these observations. The data in the Milky Way can be used to test simulations on still smaller scales by using the range of distances, hence resolutions, available.

Many of the trends that do exist and the differences between the Milky Way and the PHANGS galaxies can




Figure 12. The median, mean, and standard deviation of α_{vir} and σ_v in bins of 1 kpc of distance from the Sun (upper axis). The effective resolution is indicated on the lower axis.

be traced to different conditions that exist near the centers of galaxies, including the Milky Way. The radial distributions of many variables in the Milky Way are similar to those found in the PHANGS galaxies, with some differences, perhaps related to the effects of the bar in the Milky Way (Evans et al. 2026; Baba 2026). More specific results are listed next.

1. The formula used by L25 for how the luminosity to mass conversion factor (α_{CO}) depends on metallicity (Z) differs substantially from that used in studies in the Milky Way. However, the formula for how Z depends on galactocentric radius also differs substantially from the measured dependence in the Milky Way. The two effects cancel nearly exactly, so our analysis does not depend strongly on which we use. This cancellation is fortuitous but probably masks deeper issues. See §4.1 and Figure 2 in particular.
2. The coarse spatial resolution of the L25 data set (150 pc) and the surface density limitation ($\Sigma_{\text{mol}} \geq 4 \text{ M}_{\odot} \text{ pc}^{-2}$) mean that observers from another galaxy would miss many clouds in the Milky Way catalog and underestimate the surface density of those they detect. However the mass-weighting scheme when averaging into hexagons of L25 increases the resulting measures of surface

density to be more similar to the underlying cloud level distribution. See §4.2 and Figure 3.

3. The sensitivity limit ($I_{\text{CO}} \geq 0.6 \text{ K km s}^{-1}$) of the L25 analysis has a minimal effect on the inferred properties. See §4.3. Better sensitivity in surveys like PHANGS would have few benefits.
4. The tracer used ($\text{CO } J = 1 \rightarrow 0$, $\text{CO } J = 2 \rightarrow 1$, extinction) matters for the inferred surface density and average volume density, but only at the factor of two level. See §4.4 and Figure 4.
5. The method of cloud definition (MD17 versus R16) affects measures of the virial parameter substantially. See §4.4 and panel c of Figure 4.
6. The relations between t_{dep} and other variables expected from simple considerations (§7.2) are not generally seen, neither in the L25 data nor in the Milky Way. There are a number of explanations for this, as detailed in L25 and §7.4. Two of the more important explanations are that gas and star formation behave differently near galactic centers and that the simple considerations do not account for hierarchical cloud structure (§7.3).
7. The method of averaging free-fall times has a moderate effect. Mass-weighting the speeds ($1/t_{\text{ff}}$) as adopted by L25 produces an effective free-fall time

about 30% less than averaging t_{ff} . See Appendix C.

8. Averaging virial parameters because of coarse resolution causes formulae using α_{vir} to underestimate the efficiency per free-fall time and thus star formation rate, though mass-weighting mitigates this problem. See Appendix D.

ACKNOWLEDGMENTS

We are grateful to C. McKee and D. Hollenbach for interesting comments on the mean cloud densities in

Table 3. This research used the Canadian Advanced Network For Astronomy Research (CANFAR) operated in partnership by the Canadian Astronomy Data Centre and The Digital Research Alliance of Canada with support from the National Research Council of Canada the Canadian Space Agency, CANARIE and the Canadian Foundation for Innovation. NJE is grateful to the Astronomy Department of the University of Texas for research support, INAF-IAPS (Roma, Italy) for hospitality during a visit, and to H. Khan-Farooki, along with two anonymous cornea donors, for the gift of sight.

Software: astropy (Astropy Collaboration et al. 2013, 2018, 2022), scipy (Virtanen et al. 2020)

APPENDIX

A. SOME DETAILS OF COMPARISONS

We note here some relevant facts about the comparison of the Milky Way survey and the PHANGS-ALMA survey. The threshold value for PHANGS-ALMA is 0.6 K km s^{-1} (L25). Using $R_{21} = 0.65$ (Schinnerer & Leroy 2024), this corresponds to 1.0 K km s^{-1} for the $J = 1 \rightarrow 0$ line. For the Dame et al. (2001) survey, the sensitivity varies but is always less than 0.25 K (Fig. 22 in MD17). With a velocity resolution of 1.3 km s^{-1} , this corresponds to 0.33 K km s^{-1} . Thus the sensitivity of the Milky Way survey is at least 3 times better than the threshold for PHANGS-ALMA. For the great majority of the data, the intensity sensitivity is less than 0.13 K km s^{-1} , 8 times better than the PHANGS-ALMA threshold. We consider the effects of this difference in §4.3.

The spatial resolution of PHANGS-ALMA depends on the distance to the galaxy, but they have been convolved to a common spatial scale of 150 pc for the primary analysis. Some analysis with higher resolution (as small as 60 pc for 9 galaxies) for the more nearby galaxies was also done. For the $8''.5$ beam of the Dame survey, the resolution varies dramatically with heliocentric distance. At 20 kpc, it is 50 pc, compared to 0.3 pc for the closest molecular clouds. Especially with subsets located closer to the Sun, we can examine the effects of spatial resolution (§4.2 and §7.3).

For the Milky Way clouds, we use $\Sigma_{\text{cloud}} = M_{\text{cloud}}/A_{\text{cloud}}$, where A_{cloud} is the area of the cloud; for average density, we use $\bar{n}_{\text{cloud}} = M_{\text{cloud}}/V_{\text{cloud}}$, where V_{cloud} is the volume of the ellipsoid that defines the cloud in MD17. For the Milky Way, our cells are squares of 150 pc on a side. To compute the volume, we assume a scale height of 100 pc for consistency with L25. Then the smoothed surface density of a cell (Σ_{cell}) is the mass of the cell over the area of the cell. The smoothed volume density ($\bar{n}_{\text{cell}}$) is the mass of the cell divided by the volume of the cell. For the hexagons, we use the same formulae for PHANGS and the Milky Way, mass-weighting the averages of values in the cells.

L25 define $\sigma_{\text{mol}}^{\text{cloud}}$ as the effective (1D) velocity dispersion within their 150 pc resolution, corrected for galaxy inclination by multiplying by $(\cos i)^{0.5}$. Heyer et al. (2001) defines “effective dispersion” in Equation A7 as $\sigma_v = \delta v / (8 \ln 2)^{0.5}$, where δv is the FWHM of summed spectrum of all pixels. Because we work with catalogs, we simply average the MD17 catalog values of velocity dispersions (σ_v^{cloud}) for all clouds in a cell to get σ_v^{cell} . As noted above, this may underestimate the full dispersion measured by L25, but it more accurately describes the actual cloud-level dispersion.

L25 define $\alpha_{\text{vir}}^{\text{cloud}} = 5\sigma_{\text{mol}}^{\text{cloud}2} R_{\text{pix}} / (fGM_{\text{mol}}^{\text{beam}})$; f is a factor that depends in the cloud density structure (Bertoldi & McKee 1992), but it was assumed to be unity in L25 and we assume the same. We define $\alpha_{\text{vir,cell}} = 5\sigma_v^{\text{cell}2} R / (GM_{\text{cell}})$. L25 use $R_{\text{pix}} = (l^2 H / 8 \cos i)^{1/3}$, where $l = 150 \text{ pc}$ and $H = 100 \text{ pc}$. For $\cos i = 1$, this yields 65.5 pc. Our $\alpha_{\text{vir,cell}}$ is computed from the same formula, with $R = 65.5 \text{ pc}$, the averaged velocity dispersion (σ_v^{cell}), and the total mass in a cell (M_{cell}). For the hexagons, we calculate a mass-weighted average of $\alpha_{\text{vir,cell}}$, consistent with L25.

For all quantities labeled with $\langle x \rangle_{\text{hex}}$, the values are mass-weighted averages over the values in cells:

$$\langle x \rangle_{\text{hex}} = \frac{\sum_i M_i x_i}{\sum_i M_i} \quad (\text{A1})$$

B. STELLAR SURFACE DENSITY

As noted in §4.1, the CO luminosity to mass conversion factor can depend on stellar surface density through the starburst factor (Schinnerer & Leroy 2024). This becomes important only where $\Sigma_\star > 100 \text{ M}_\odot \text{ pc}^{-2}$. The standard exponential disk model for stellar density (Binney & Piffl 2015) does not predict values of $\Sigma_\star$ above this value, but Lian et al. (2024) have used APOGEE and GAIA data to derive nonparametric $\Sigma_\star$ distributions of bolometric luminosity in age bins from $R_{\text{Gal}} = 0$ to 17 kpc. Their Figure 1 shows the total luminosity surface density versus R_{Gal} . Assuming mass-to-light of unity, it exceeds $100 \text{ M}_\odot \text{ pc}^{-2}$ only in the innermost ring, $R_{\text{Gal}} = 0$ to 2 kpc. Much of that is due to very old stars which have large heights and may not contribute to the gravity acting on the molecular gas, as shown by the data including only ages less than 8 Gyr. The data for their youngest bin (0-2 Gyr) shows a very strong dip, similar to that seen in the molecular and star formation distributions.

The starburst factor in the Milky Way would be unity everywhere but perhaps in the central 2 kpc. If we use the Lian et al. (2024) data, assume mass-to-light of unity, and take the implied total stellar density of $\Sigma_\star = 1.2 \times 10^3 \text{ M}_\odot \text{ pc}^{-2}$, the starburst factor would be 0.53. If we use only the stars younger than 8 Gyr, the factor would be 0.74. We use the latter.

The stellar distribution also affects simulation of the extragalactic form for $\alpha_{\text{CO}}(R_{\text{Gal}})$ through the method of determining $Z(R_{\text{Gal}})$, which uses a form that depends on R_{Gal}/r_e , where r_e is the effective radius, containing half the light of the galaxy. To simulate this for the Milky Way, we used $r_e = 4.75 \text{ kpc}$ (Imig et al. 2025).

C. DEFINITIONS OF t_{ff}

The definition of t_{ff} in Equation 2 emphasizes the regions most active in star formation by mass-weighting and averaging speeds, rather than t_{ff} . To test the effects, we compared results with two other averaging methods: a straight average of free-fall times, and an unweighted average of speeds ($1/t_{\text{ff}}$). The test was performed in the step of averaging cells into hexagons for the Milky Way data, using the L25 criteria. After computing the three quantities for each hexagon, we computed the mean values over all hexagons for each of the three methods. The straight average of t_{ff} leads to the longest average (11.1 Myr); the average of speeds yields 9.2 Myr; and the mass-weighted average of speeds is 7.4 Myr. Using mass-weighted speeds does not have a major effect, but it decreases the timescale by about 30%. Because of the non-linearity, it is certainly better to average speeds than free-fall times.

D. EFFECTS OF AVERAGING α_{vir}

The virial parameter is also mass-weighted in the L25 method. To test the effect of this, we computed it with and without mass-weighting, using the same method described in Appendix C. The mean value with mass weighting is 44% smaller than that without mass weighting.

Another question is whether averaging α_{vir} misleads us about star formation. To test this, we calculated the star formation rate of the Milky Way three ways, all using the EKO formula (Equation 17). First, we redid the EKO method cloud by cloud with the various cuts above as a check and got $\text{SFR} = 1.39 \text{ M}_\odot \text{ yr}^{-1}$, only slightly lower than the $1.47 \text{ M}_\odot \text{ yr}^{-1}$ in Elia et al. (2025a). Second, we calculated with the total mass, the average ϵ_{ff} using each hexagon's mass-weighted α_{vir} , and the mass-weighted inverse free-fall time, with the result of $1.06 \text{ M}_\odot \text{ yr}^{-1}$, two-thirds the correct value. Third, we computed with the total mass, the ϵ_{ff} calculated from the mean mass-weighted α_{vir} , and the mass-weighted inverse free-fall time, using the values in Table 4, with the result of $0.40 \text{ M}_\odot \text{ yr}^{-1}$, about one-third the correct value.

One needs to average ϵ_{ff} , not α_{vir} because of the non-linearity. The most bound clouds have a high weight in the star formation process. For that reason, any averaging (or lower spatial resolution) will decrease the predicted star formation rate because the active clouds are diluted by the inactive clouds.

On the other hand, the mass-weighting has decreased α_{vir} on average (compare values in Table 2 and 4). Many of the clouds with high α_{vir} in the MD17 catalog have low mass and are down-weighted in the L25 method. The value of $\langle \alpha_{\text{vir}} \rangle_{\text{hex}}$ in Table 4 is however still larger than that for the PHANGS sample. The MD17 method of cloud identification also produces values of α_{vir} that are larger than those in a less complete catalog of Milky Way clouds compiled by R16, as can be seen in the entries for each in Table 3.


E. EFFECT OF ASSUMPTIONS ABOUT α_{CO}

L25 noted that the choice of assumptions about α_{CO} had a substantial impact on their results. We confirm this fact using the data in their table. Making the same assumption of a constant $\alpha_{\text{CO}} = 4.35$, we find weaker correlations in

Table 7. Correlations and Fits for fixed α_{CO}

Sample	x	r_s	p	m	σ_m	x_0	b	σ_b
PHANGS	$\log \langle t_{\text{ff}} \rangle_{\text{hex}}$	-0.18	3.11e-07	-0.22	0.05	6.91	9.37	9.95e-03
MW	$\log \langle t_{\text{ff}} \rangle_{\text{hex}}$	-0.03	7.22e-01	0.01	0.25	6.97	9.19	3.60e-02
PHANGS	$\log \langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$	0.13	1.28e-04	0.09	0.03	1.66	9.37	1.00e-02
MW	$\log \langle \Sigma_{\text{cell}} \rangle_{\text{hex}}$	0.03	6.64e-01	0.02	0.13	1.65	9.19	3.60e-02
PHANGS	$\log \langle \sigma_v^{\text{cell}} \rangle_{\text{hex}}$	0.14	7.32e-05	0.21	0.05	0.74	9.37	1.02e-02
MW	$\log \langle \sigma_v^{\text{cell}} \rangle_{\text{hex}}$	-0.14	7.73e-02	-0.68	0.27	0.77	9.18	3.54e-02
PHANGS	$\log \langle \alpha_{\text{vir}} \rangle_{\text{hex}}$	-0.02	5.36e-01	-0.00	0.05	0.26	9.38	4.58e-02
MW	$\log \langle \alpha_{\text{vir}} \rangle_{\text{hex}}$	-0.08	2.99e-01	-0.14	0.14	0.65	9.19	3.59e-02

general in both the PHANGS and Milky Way data, as shown in Table 7. The differences in the fits for the Milky Way and the PHANGS galaxies are even less, except for the velocity dispersion, on which t_{dep} still depends more strongly in the Milky Way.

**Figure 13.** The slopes of the fits in Table 7 are plotted with uncertainties for both the PHANGS galaxies and the Milky Way, with the assumption of fixed $\alpha_{\text{CO}} = 4.35$.

REFERENCES

- Anders, F., Chiappini, C., Minchev, I., et al. 2017, A&A, 600, A70, doi: [10.1051/0004-6361/201629363](https://doi.org/10.1051/0004-6361/201629363)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, ApJ, 935, 167, doi: [10.3847/1538-4357/ac7c74](https://doi.org/10.3847/1538-4357/ac7c74)
- Baba, J. 2026, PASJ, 78, 1093, doi: [10.1093/pasj/psag045](https://doi.org/10.1093/pasj/psag045)
- Benjamin, R. A., Churchwell, E., Babler, B. L., et al. 2005, ApJL, 630, L149, doi: [10.1086/491785](https://doi.org/10.1086/491785)
- Bergin, E. A., & Tafalla, M. 2007, ARA&A, 45, 339, doi: [10.1146/annurev.astro.45.071206.100404](https://doi.org/10.1146/annurev.astro.45.071206.100404)
- Bertoldi, F., & McKee, C. F. 1992, ApJ, 395, 140, doi: [10.1086/171638](https://doi.org/10.1086/171638)
- Bigiel, F., Leroy, A. K., Walter, F., et al. 2011, ApJL, 730, L13, doi: [10.1088/2041-8205/730/2/L13](https://doi.org/10.1088/2041-8205/730/2/L13)
- Bigiel, F., Leroy, A. K., Jiménez-Donaire, M. J., et al. 2016, ApJL, 822, L26, doi: [10.3847/2041-8205/822/2/L26](https://doi.org/10.3847/2041-8205/822/2/L26)
- Binney, J., & Piffl, T. 2015, MNRAS, 454, 3653, doi: [10.1093/mnras/stv2225](https://doi.org/10.1093/mnras/stv2225)
- Bolatto, A. D., Wolfire, M., & Leroy, A. K. 2013, ARA&A, 51, 207, doi: [10.1146/annurev-astro-082812-140944](https://doi.org/10.1146/annurev-astro-082812-140944)

- Brunetti, N., Wilson, C. D., He, H., et al. 2024, MNRAS, 530, 597, doi: [10.1093/mnras/stae890](https://doi.org/10.1093/mnras/stae890)
- Chapman, N. L., Mundy, L. G., Lai, S.-P., & Evans, II, N. J. 2009, ApJ, 690, 496, doi: [10.1088/0004-637X/690/1/496](https://doi.org/10.1088/0004-637X/690/1/496)
- Dame, T. M., Hartmann, D., & Thaddeus, P. 2001, ApJ, 547, 792, doi: [10.1086/318388](https://doi.org/10.1086/318388)
- Downes, D., & Solomon, P. M. 1998, ApJ, 507, 615, doi: [10.1086/306339](https://doi.org/10.1086/306339)
- Duarte-Cabral, A., & The Sedigism Consortium. 2022, MNRAS, 513, 296, doi: [10.1093/mnras/stac909](https://doi.org/10.1093/mnras/stac909)
- Duarte-Cabral, A., Colombo, D., Urquhart, J. S., et al. 2021, The SEDIGISM survey: molecular clouds in the inner Galaxy, Monthly Notices of the Royal Astronomical Society, Volume 500, Issue 3, pp.3027-3049, doi: [10.1093/mnras/staa2480](https://doi.org/10.1093/mnras/staa2480)
- Elia, D., Evans, N. J., Soler, J. D., et al. 2025a, ApJ, 980, 216, doi: [10.3847/1538-4357/adaeb2](https://doi.org/10.3847/1538-4357/adaeb2)
- Elia, D., Evans, II, N. J., Soler, J. D., et al. 2025b, ApJ, 992, 161, doi: [10.3847/1538-4357/ae0723](https://doi.org/10.3847/1538-4357/ae0723)
- Elia, D., Molinari, S., Schisano, E., et al. 2022, ApJ, 941, 162, doi: [10.3847/1538-4357/aca27d](https://doi.org/10.3847/1538-4357/aca27d)
- Elmegreen, B. G. 2000, ApJ, 530, 277, doi: [10.1086/308361](https://doi.org/10.1086/308361)
- Evans, Neal J., I., Heyer, M., Miville-Deschênes, M.-A., Nguyen-Luong, Q., & Merello, M. 2021, ApJ, 920, 126, doi: [10.3847/1538-4357/ac1425](https://doi.org/10.3847/1538-4357/ac1425)
- Evans, Neal J., I., Kim, K.-T., Wu, J., et al. 2020, ApJ, 894, 103, doi: [10.3847/1538-4357/ab8938](https://doi.org/10.3847/1538-4357/ab8938)
- Evans, N. J., Kim, J.-G., & Ostriker, E. C. 2022, ApJL, 929, L18, doi: [10.3847/2041-8213/ac6427](https://doi.org/10.3847/2041-8213/ac6427)
- Evans, II, N. J. 1999, ARA&A, 37, 311, doi: [10.1146/annurev.astro.37.1.311](https://doi.org/10.1146/annurev.astro.37.1.311)
- Evans, II, N. J., Di Francesco, J., Lee, J.-E., et al. 2015, ApJ, 814, 22, doi: [10.1088/0004-637X/814/1/22](https://doi.org/10.1088/0004-637X/814/1/22)
- Evans, II, N. J., Elia, D., Hawkins, K., Stuber, S., & Sun, J. 2026, ApJL, 998, L23, doi: [10.3847/2041-8213/ae3e80](https://doi.org/10.3847/2041-8213/ae3e80)
- Faustino Vieira, H., Duarte-Cabral, A., Davis, T. A., et al. 2023, MNRAS, 524, 161, doi: [10.1093/mnras/stad1876](https://doi.org/10.1093/mnras/stad1876)
- Faustino Vieira, H., Duarte-Cabral, A., Smith, M. W. L., et al. 2025, MNRAS, 538, 2744, doi: [10.1093/mnras/staf411](https://doi.org/10.1093/mnras/staf411)
- Goldsmith, P. F., Heyer, M., Narayanan, G., et al. 2008, ApJ, 680, 428, doi: [10.1086/587166](https://doi.org/10.1086/587166)
- Gong, M., Ostriker, E. C., Kim, C.-G., & Kim, J.-G. 2020, ApJ, 903, 142, doi: [10.3847/1538-4357/abbdab](https://doi.org/10.3847/1538-4357/abbdab)
- Groves, B., Kreckel, K., Santoro, F., et al. 2023, MNRAS, 520, 4902, doi: [10.1093/mnras/stad114](https://doi.org/10.1093/mnras/stad114)
- Heiderman, A., Evans, Neal J., I., Allen, L. E., Huard, T., & Heyer, M. 2010, ApJ, 723, 1019, doi: [10.1088/0004-637X/723/2/1019](https://doi.org/10.1088/0004-637X/723/2/1019)
- Heyer, M., Krawczyk, C., Duval, J., & Jackson, J. M. 2009, ApJ, 699, 1092, doi: [10.1088/0004-637X/699/2/1092](https://doi.org/10.1088/0004-637X/699/2/1092)
- Heyer, M. H., Carpenter, J. M., & Snell, R. L. 2001, ApJ, 551, 852, doi: [10.1086/320218](https://doi.org/10.1086/320218)
- Hu, C.-Y., Schrubba, A., Sternberg, A., & van Dishoeck, E. F. 2022, ApJ, 931, 28, doi: [10.3847/1538-4357/ac65fd](https://doi.org/10.3847/1538-4357/ac65fd)
- Imig, J., Holtzman, J. A., Zasowski, G., et al. 2025, ApJ, 990, 203, doi: [10.3847/1538-4357/adf723](https://doi.org/10.3847/1538-4357/adf723)
- Jiménez-Donaire, M. J., Bigiel, F., Leroy, A. K., et al. 2019, ApJ, 880, 127, doi: [10.3847/1538-4357/ab2b95](https://doi.org/10.3847/1538-4357/ab2b95)
- Kauffmann, J., Bertoldi, F., Bourke, T. L., Evans, II, N. J., & Lee, C. W. 2008, A&A, 487, 993, doi: [10.1051/0004-6361:200809481](https://doi.org/10.1051/0004-6361:200809481)
- Kennicutt, Robert C., J. 1998, ApJ, 498, 541, doi: [10.1086/305588](https://doi.org/10.1086/305588)
- Kennicutt, R. C., & Evans, N. J. 2012, ARA&A, 50, 531, doi: [10.1146/annurev-astro-081811-125610](https://doi.org/10.1146/annurev-astro-081811-125610)
- Kim, J.-G., Ostriker, E. C., & Filippova, N. 2021, ApJ, 911, 128, doi: [10.3847/1538-4357/abe934](https://doi.org/10.3847/1538-4357/abe934)
- Kruijssen, J. M. D., & Longmore, S. N. 2014, MNRAS, 439, 3239, doi: [10.1093/mnras/stu098](https://doi.org/10.1093/mnras/stu098)
- Krumholz, M. R., Dekel, A., & McKee, C. F. 2012, ApJ, 745, 69, doi: [10.1088/0004-637X/745/1/69](https://doi.org/10.1088/0004-637X/745/1/69)
- Lada, C. J., Forbrich, J., Lombardi, M., & Alves, J. F. 2012, ApJ, 745, 190, doi: [10.1088/0004-637X/745/2/190](https://doi.org/10.1088/0004-637X/745/2/190)
- Lada, C. J., Lombardi, M., & Alves, J. F. 2010, ApJ, 724, 687, doi: [10.1088/0004-637X/724/1/687](https://doi.org/10.1088/0004-637X/724/1/687)
- Lee, J. C., Whitmore, B. C., Thilker, D. A., et al. 2022, ApJS, 258, 10, doi: [10.3847/1538-4365/ac1fe5](https://doi.org/10.3847/1538-4365/ac1fe5)
- Lee, J. C., Sandstrom, K. M., Leroy, A. K., et al. 2023, ApJL, 944, L17, doi: [10.3847/2041-8213/acaaae](https://doi.org/10.3847/2041-8213/acaaae)
- Leroy, A. K., Schinnerer, E., Hughes, A., et al. 2021, ApJS, 257, 43, doi: [10.3847/1538-4365/ac17f3](https://doi.org/10.3847/1538-4365/ac17f3)
- Leroy, A. K., Rosolowsky, E., Usero, A., et al. 2022, ApJ, 927, 149, doi: [10.3847/1538-4357/ac3490](https://doi.org/10.3847/1538-4357/ac3490)
- Leroy, A. K., Sun, J., Meidt, S., et al. 2025, ApJ, 985, 14, doi: [10.3847/1538-4357/adbcab](https://doi.org/10.3847/1538-4357/adbcab)
- . 2026, ApJ, 1003, 248, doi: [10.3847/1538-4357/ae5e59](https://doi.org/10.3847/1538-4357/ae5e59)
- Li, Y., Calzetti, D., Kennicutt, R. C., et al. 2010, ApJ, 725, 677, doi: [10.1088/0004-637X/725/1/677](https://doi.org/10.1088/0004-637X/725/1/677)
- Lian, J., Wang, T., Feng, Q., Huang, Y., & Guo, H. 2025, ApJL, 990, L37, doi: [10.3847/2041-8213/adfc73](https://doi.org/10.3847/2041-8213/adfc73)
- Lian, J., Zasowski, G., Chen, B., et al. 2024, Nature Astronomy, 8, 1302, doi: [10.1038/s41550-024-02315-7](https://doi.org/10.1038/s41550-024-02315-7)
- Licquia, T. C., & Newman, J. A. 2015, ApJ, 806, 96, doi: [10.1088/0004-637X/806/1/96](https://doi.org/10.1088/0004-637X/806/1/96)
- Lucey, M., Pearson, S., Hunt, J. A. S., et al. 2023, MNRAS, 520, 4779, doi: [10.1093/mnras/stad406](https://doi.org/10.1093/mnras/stad406)
- McKee, C. F., & Ostriker, E. C. 2007, ARA&A, 45, 565, doi: [10.1146/annurev.astro.45.051806.110602](https://doi.org/10.1146/annurev.astro.45.051806.110602)

- Méndez-Delgado, J. E., Amayo, A., Arellano-Córdova, K. Z., et al. 2022, *MNRAS*, 510, 4436, doi: [10.1093/mnras/stab3782](https://doi.org/10.1093/mnras/stab3782)
- Miville-Deschênes, M.-A., Murray, N., & Lee, E. J. 2017, *ApJ*, 834, 57, doi: [10.3847/1538-4357/834/1/57](https://doi.org/10.3847/1538-4357/834/1/57)
- Neralwar, K. R., Colombo, D., Duarte-Cabral, A., et al. 2022, *A&A*, 664, A84, doi: [10.1051/0004-6361/202142513](https://doi.org/10.1051/0004-6361/202142513)
- Neumann, L., Jiménez-Donaire, M. J., Leroy, A. K., et al. 2025, *A&A*, 693, L13, doi: [10.1051/0004-6361/202453208](https://doi.org/10.1051/0004-6361/202453208)
- Oakes, E. K., Faesi, C. M., Rosolowsky, E., et al. 2025, *ApJ*, 993, 193, doi: [10.3847/1538-4357/ae0642](https://doi.org/10.3847/1538-4357/ae0642)
- Ossenkopf, V., & Henning, T. 1994, *A&A*, 291, 943
- Padoan, P., Haugbølle, T., & Nordlund, Å. 2012, *ApJL*, 759, L27, doi: [10.1088/2041-8205/759/2/L27](https://doi.org/10.1088/2041-8205/759/2/L27)
- Patra, S., Evans, N. J., Kim, K.-T., et al. 2025, *ApJ*, 983, 133, doi: [10.3847/1538-4357/adbf8d](https://doi.org/10.3847/1538-4357/adbf8d)
- Peretto, N., Rigby, A. J., Louvet, F., et al. 2023, *MNRAS*, 525, 2935, doi: [10.1093/mnras/stad2453](https://doi.org/10.1093/mnras/stad2453)
- Plume, R., Jaffe, D. T., Evans, II, N. J., Martín-Pintado, J., & Gómez-González, J. 1997, *ApJ*, 476, 730, doi: [10.1086/303654](https://doi.org/10.1086/303654)
- Pokhrel, R., Gutermuth, R. A., Krumholz, M. R., et al. 2021, *ApJL*, 912, L19, doi: [10.3847/2041-8213/abf564](https://doi.org/10.3847/2041-8213/abf564)
- Pontoppidan, K. M., Evans, N., Bergner, J., & Yang, Y.-L. 2024, *Research Notes of the American Astronomical Society*, 8, 68, doi: [10.3847/2515-5172/ad303f](https://doi.org/10.3847/2515-5172/ad303f)
- Ramambason, L., Chevance, M., Kim, J., et al. 2026, *A&A*, 706, A186, doi: [10.1051/0004-6361/202556206](https://doi.org/10.1051/0004-6361/202556206)
- Rice, T. S., Goodman, A. A., Bergin, E. A., Beaumont, C., & Dame, T. M. 2016, *ApJ*, 822, 52, doi: [10.3847/0004-637X/822/1/52](https://doi.org/10.3847/0004-637X/822/1/52)
- Rosolowsky, E., Hughes, A., Leroy, A. K., et al. 2021, *MNRAS*, 502, 1218, doi: [10.1093/mnras/stab085](https://doi.org/10.1093/mnras/stab085)
- Rubinstein, A. E., Evans, N. J., Tyagi, H., et al. 2024, *ApJ*, 974, 112, doi: [10.3847/1538-4357/ad6b92](https://doi.org/10.3847/1538-4357/ad6b92)
- Sánchez, S. F., Rosales-Ortega, F. F., Iglesias-Páramo, J., et al. 2014, *A&A*, 563, A49, doi: [10.1051/0004-6361/201322343](https://doi.org/10.1051/0004-6361/201322343)
- Sánchez, S. F., Barrera-Ballesteros, J. K., López-Cobá, C., et al. 2019, *MNRAS*, 484, 3042, doi: [10.1093/mnras/stz019](https://doi.org/10.1093/mnras/stz019)
- Sanhueza, P., Contreras, Y., Wu, B., et al. 2019, *ApJ*, 886, 102, doi: [10.3847/1538-4357/ab45e9](https://doi.org/10.3847/1538-4357/ab45e9)
- Schinnerer, E., & Leroy, A. K. 2024, *ARA&A*, 62, 369, doi: [10.1146/annurev-astro-071221-052651](https://doi.org/10.1146/annurev-astro-071221-052651)
- Schisano, E., Molinari, S., Coletta, A., et al. 2026, *A&A*, 707, A221, doi: [10.1051/0004-6361/202555619](https://doi.org/10.1051/0004-6361/202555619)
- Schruba, A., Leroy, A. K., Walter, F., Sandstrom, K., & Rosolowsky, E. 2010, *ApJ*, 722, 1699, doi: [10.1088/0004-637X/722/2/1699](https://doi.org/10.1088/0004-637X/722/2/1699)
- Shirley, Y. L. 2015, *PASP*, 127, 299, doi: [10.1086/680342](https://doi.org/10.1086/680342)
- Shirley, Y. L., Mangum, J. G., Narayanan, D., & Di Francesco, J. 2026, *PASP*, 138, 043001, doi: [10.1088/1538-3873/ae4270](https://doi.org/10.1088/1538-3873/ae4270)
- Soler, J. D., Zucker, C., Peek, J. E. G., et al. 2023, *A&A*, 675, A206, doi: [10.1051/0004-6361/202346241](https://doi.org/10.1051/0004-6361/202346241)
- Su, Y., Yang, J., Zhang, S., et al. 2019, *ApJS*, 240, 9, doi: [10.3847/1538-4365/aaf1c8](https://doi.org/10.3847/1538-4365/aaf1c8)
- Sun, J., Leroy, A. K., Rosolowsky, E., et al. 2022, *AJ*, 164, 43, doi: [10.3847/1538-3881/ac74bd](https://doi.org/10.3847/1538-3881/ac74bd)
- Sun, J., Leroy, A. K., Ostriker, E. C., et al. 2023, *ApJL*, 945, L19, doi: [10.3847/2041-8213/acbd9c](https://doi.org/10.3847/2041-8213/acbd9c)
- Sun, J., Teng, Y.-H., Chiang, I.-D., et al. 2025, *ApJ*, 994, 263, doi: [10.3847/1538-4357/ae10be](https://doi.org/10.3847/1538-4357/ae10be)
- Sun, Y., Yang, J., Xu, Y., et al. 2020, *ApJS*, 246, 7, doi: [10.3847/1538-4365/ab5b97](https://doi.org/10.3847/1538-4365/ab5b97)
- Urquhart, J. S., König, C., Colombo, D., et al. 2025, *MNRAS*, 539, 3105, doi: [10.1093/mnras/staf665](https://doi.org/10.1093/mnras/staf665)
- Virtanen, P., others, & SciPy 1.0 Contributors. 2020, *Nature Methods*, 17, 261, doi: [10.1038/s41592-019-0686-2](https://doi.org/10.1038/s41592-019-0686-2)
- Wang, T., Yuan, H., Chen, B., et al. 2025, *ApJS*, 280, 16, doi: [10.3847/1538-4365/adad89](https://doi.org/10.3847/1538-4365/adad89)
- Williams, T. G., Lee, J. C., Larson, K. L., et al. 2024, *ApJS*, 273, 13, doi: [10.3847/1538-4365/ad4be5](https://doi.org/10.3847/1538-4365/ad4be5)
- Xu, F., Wang, K., Liu, T., et al. 2024, *ApJS*, 270, 9, doi: [10.3847/1538-4365/acfee5](https://doi.org/10.3847/1538-4365/acfee5)
- Yang, Y.-L., Evans, II, N. J., Smith, A., et al. 2020, *ApJ*, 891, 61, doi: [10.3847/1538-4357/ab7201](https://doi.org/10.3847/1538-4357/ab7201)
- Zakardjian, A., Pety, J., Herrera, C. N., et al. 2023, *A&A*, 678, A171, doi: [10.1051/0004-6361/202244520](https://doi.org/10.1051/0004-6361/202244520)
- Zhou, X., Yang, J., Sun, Y., et al. 2025, *AJ*, 170, 239, doi: [10.3847/1538-3881/adfe68](https://doi.org/10.3847/1538-3881/adfe68)
- Zuckerman, B., & Evans, II, N. J. 1974, *ApJL*, 192, L149, doi: [10.1086/181613](https://doi.org/10.1086/181613)