

Mathematical Methods of Forecasting

Yakovlev Konstantin

Contents

1	Lab work 10	2
1.1	Motivation	2
1.2	Problem statement	2
1.3	Problem solution	3
1.4	Code analysis	3
1.5	Experiment on synthetic dataset	3

1 Lab work 10

1.1 Motivation

The work investigates the problem of predicting a complex structured target variable. The problem is supposed to be solved by higher-order partial least squares [2]. This model predicts tensor $\underline{\mathbf{Y}}$ from tensor $\underline{\mathbf{X}}$ using projection into the latent space and solving regression problem on latent variables. The aim of computational experiment is to predict gyroscope data using accelerometer data. The experiment is held on a WISDM dataset [1].

1.2 Problem statement

Given a dataset $\mathfrak{D} = (\underline{\mathbf{X}}, \underline{\mathbf{Y}})$, where tensors $\underline{\mathbf{X}} \in \mathbb{R}^{I_1 \times \dots \times I_N}$ and $\underline{\mathbf{Y}} \in \mathbb{R}^{J_1 \times \dots \times J_M}$, $I_1 = J_1$. We also require that $N \geq 3$, $M \geq 3$. Assume that $\underline{\mathbf{X}}$ is decomposed as a sum of rank- $(1, L_2, \dots, L_N)$ Tucker-blocks, and $\underline{\mathbf{Y}}$ is decomposed as a sum of rank- $(1, K_2, \dots, K_M)$ Tucker-blocks. This can be written in a following form:

$$\underline{\mathbf{X}} \approx \sum_{r=1}^R \underline{\mathbf{G}}_r \times_1 \mathbf{t}_r \times_2 \mathbf{P}_r^{(1)} \times_3 \dots \times_N \mathbf{P}_r^{(N-1)}, \quad (1.1)$$

$$\underline{\mathbf{Y}} \approx \sum_{r=1}^R \underline{\mathbf{D}}_r \times_1 \mathbf{t}_r \times_2 \mathbf{Q}_r^{(1)} \times_3 \dots \times_M \mathbf{Q}_r^{(M-1)}, \quad (1.2)$$

where R is a number of latent vectors, $\mathbf{t}_r \in \mathbb{R}^{I_1}$ is a r -th latent vector, $\mathbf{P}_r^{(n)} \in \mathbb{R}^{I_{n+1} \times L_{n+1}}$, $\mathbf{P}_r^{(n)\top} \mathbf{P}_r^{(n)} = \mathbf{I}$, $\mathbf{Q}_r^{(m)} \in \mathbb{R}^{J_{m+1} \times K_{m+1}}$, $\mathbf{Q}_r^{(m)\top} \mathbf{Q}_r^{(m)} = \mathbf{I}$, and $\underline{\mathbf{G}} \in \mathbb{R}^{1 \times L_1 \times \dots \times L_N}$, $\underline{\mathbf{D}} \in \mathbb{R}^{1 \times J_1 \times K_1 \times \dots \times K_M}$ are core tensors.

For simplicity, let us define the following notation of Tucker decomposition:

$$\sum_{r=1}^R \underline{\mathbf{G}}_r \times_1 \mathbf{t}_r \times_2 \mathbf{P}_r^{(1)} \times_3 \dots \times_N \mathbf{P}_r^{(N-1)} = [\underline{\mathbf{G}}; \mathbf{t}, \mathbf{P}^{(1)}, \dots, \mathbf{P}^{(N-1)}].$$

Let $\underline{\mathbf{C}} \in \mathbb{R}^{I_2 \times \dots \times I_N \times J_2 \times \dots \times J_M}$, where:

$$c_{i_2, \dots, i_N, j_1, \dots, j_M} = \sum_{i_1=1}^{I_1} x_{i_1, i_2, \dots, i_N} y_{j_1, \dots, j_M}.$$

The problem is to find matrices $\underline{\mathbf{P}}_r^{(n)}$, $\underline{\mathbf{Q}}_r^{(m)}$ and latent vectors \mathbf{t}_r from the following optimization problem:

$$\max_{\mathbf{P}^{(n)}, \mathbf{Q}^{(m)}} \|[\underline{\mathbf{C}}; \mathbf{P}^{(1)\top}, \dots, \mathbf{P}^{(N-1)\top}, \mathbf{Q}^{(1)\top}, \dots, \mathbf{Q}^{(M-1)\top}]\|_F^2 \quad (1.3)$$

$$\text{s.t., } \mathbf{P}^{(n)\top} \mathbf{P}^{(n)} = \mathbf{I}, \mathbf{Q}^{(m)\top} \mathbf{Q}^{(m)} = \mathbf{I}. \quad (1.4)$$

Based on the found matrices $\underline{\mathbf{P}}^{(n)}$, $\underline{\mathbf{Q}}^{(m)}$ from (1.3), we find a latent vector \mathbf{t} from the following optimization problem:

$$\mathbf{t} = \arg \min_{\mathbf{t}} \|\underline{\mathbf{X}} - [\underline{\mathbf{G}}; \mathbf{t}, \mathbf{P}^{(1)}, \dots, \mathbf{P}^{(N-1)}]\|_F^2. \quad (1.5)$$

The following procedure should be carried R times. The next step will be performed with a residual tensor:

$$\underline{\mathbf{X}} - [\underline{\mathbf{G}}; \mathbf{t}, \mathbf{P}^{(1)}, \dots, \mathbf{P}^{(N-1)}].$$

1.3 Problem solution

The optimization algorithm is described in [2]. On each of R steps it computes orthogonal Tucker decomposition of $\underline{\mathbf{C}}_r$ and performs SVD decomposition in order to find \mathbf{t}_r . The algorithm may stop earlier when frobenius norm of both residuals is less than ε .

Prediction from a new observation $\underline{\mathbf{X}}^{\text{test}}$ can be written in matricized form:

$$\underline{\mathbf{Y}}_{(1)}^{\text{test}} = \underline{\mathbf{X}}_{(1)}^{\text{test}} \mathbf{W} \mathbf{Q}^{*\top}, \quad (1.6)$$

where matrices \mathbf{W} and \mathbf{Q} have R columns, which are the following:

$$\begin{aligned} \mathbf{w}_r &= (\mathbf{P}_r^{(N-1)} \otimes \dots \otimes \mathbf{P}_r^{(1)}) \underline{\mathbf{G}}_{r(1)}^+, \\ \mathbf{q}_r^* &= \underline{\mathbf{D}}_{r(1)} (\mathbf{Q}_r^{(M-1)} \otimes \dots \otimes \mathbf{Q}_r^{(1)})^\top. \end{aligned}$$

1.4 Code analysis

The implementation of HOPLS was taken from GitHub repository¹. The computational experiment can be found on GitHub repository².

1.5 Experiment on synthetic dataset

Consider an accelerometer tensor $\underline{\mathbf{X}} \in \mathbb{R}^{P \times T \times C}$, where $P = 10$ is a number of multivariate time series, $T = 40$ is a number of time steps, $C = 10$ is the number of channels. The task is to predict gyroscope data $\underline{\mathbf{Y}} \in \mathbb{R}^{P \times T \times C}$. Tensor $\underline{\mathbf{X}}$ was generated as:

$$x_{p,t,c} = \sin(\omega_{p,t,c}(t/T) + \varphi_{p,t,c}), \quad \omega_{p,t,c} \sim \mathcal{U}(20, 30), \quad \varphi_{p,t,c} \sim \mathcal{U}(0, \pi). \quad (1.7)$$

Tensor $\underline{\mathbf{Y}}$ was generated as:

$$y_{p,t,c} = x_{p,t,c'} w_{c',c} + \varepsilon_{p,t,c}, \quad (1.8)$$

where $\mathbf{W} \in \mathbb{R}^{C \times C}$, $w_{c',c} \sim \mathcal{U}(0, 1)$, $\varepsilon_{p,t,c} \sim \mathcal{N}(0, 0.01)$.

The task is to predict $\underline{\mathbf{Y}}$ by $\underline{\mathbf{X}}$, using method described above. Define a similarity between real $\underline{\mathbf{Y}}$ and $\underline{\mathbf{Y}}^{\text{pred}}$ as Q^2 :

$$Q^2 = 1 - \frac{\|\underline{\mathbf{Y}}^{\text{pred}} - \underline{\mathbf{Y}}\|_F^2}{\|\underline{\mathbf{Y}}\|_F^2}. \quad (1.9)$$

In order to measure quality of proposed method we performed cross validation. Here we split accelerometer data $\underline{\mathbf{X}}$ by the first dimension on 5 folds. In our model $L_2 = K_2 = 6$, $L_3 = K_3 = 1$.

The goal of the computational experiment is to get dependence of validation quality on number of latent vectors R . We also compare proposed method with N-PLS. In fact, HOPLS can be simplified to N-PLS if we define all $L_n = 1$ and $K_m = 1$.

From Table 1 it can be seen that HOPLS method gives better solution in terms of Q^2 . In addition, variance of both methods is comparable.

From Figure 1 it can be seen that the optimal $R^* = 4$. Moreover, the higher R is, the more the variance of Q^2 is.

¹<https://github.com/arthurdehgan/HOPLS>

²<https://github.com/Konstantin-Iakovlev/MathMethodsOfforecasting>

Table 1: Dependence of Q^2 on R on validation dataset						
Method	$R = 1$	$R = 4$	$R = 10$	$R = 13$	$R = 16$	$R = 19$
HOPLS	0.41 ± 0.12	0.51 ± 0.19	0.41 ± 0.12	0.04 ± 0.47	-0.22 ± 0.68	-0.32 ± 0.68
N-PLS	0.42 ± 0.15	0.45 ± 0.11	0.35 ± 0.18	0.35 ± 0.19	0.33 ± 0.20	0.30 ± 0.20




Figure 1: Dependence of Q^2 on number of latent vectors R .

References

- [1] Jennifer R Kwapisz, Gary M Weiss, and Samuel A Moore. “Activity recognition using cell phone accelerometers”. In: *ACM SigKDD Explorations Newsletter* 12.2 (2011), pp. 74–82.
- [2] Qibin Zhao et al. “Higher order partial least squares (HOPLS): a generalized multilinear regression method”. In: *IEEE transactions on pattern analysis and machine intelligence* 35.7 (2012), pp. 1660–1673.