

# Computation of the mean of an inverse Gaussian distribution

Consider an inverse Gaussian distribution with location parameter  $\mu$  and scale parameter  $\lambda$ . The pdf is given as

$$f(x; \mu, \lambda) = \sqrt{\frac{\lambda}{2\pi x^3}} \exp\left(-\frac{\lambda(x - \mu)^2}{2\mu^2 x}\right).$$

We are interested in computing the mean, i.e.

$$\mathbb{E}[x] = \int_0^\infty dx x \sqrt{\frac{\lambda}{2\pi x^3}} \exp\left(-\frac{\lambda(x - \mu)^2}{2\mu^2 x}\right).$$

In order to do that consider  $t > 0$  and define a function  $g$

$$g(t) = \int_0^\infty dx \sqrt{\frac{\lambda}{2\pi x^3}} \exp\left(-\frac{\lambda(tx - \mu)^2}{2\mu^2 x}\right).$$

The integral can be solved by the substitution  $tx \rightarrow z$ :

$$\begin{aligned} g(t) &= \int_0^\infty dz \frac{1}{t} \sqrt{\frac{\lambda t^3}{2\pi x^3}} \exp\left(-\frac{\lambda t(z - \mu)^2}{2\mu^2 z}\right) \\ &= \int_0^\infty dz \sqrt{\frac{\lambda t}{2\pi x^3}} \exp\left(-\frac{\lambda t(z - \mu)^2}{2\mu^2 z}\right) \\ &= 1 \end{aligned}$$

In the last line we used that the integral is the normalization of an inverse Gaussian distribution with location parameter  $\mu$  and scale parameter  $\lambda t$ .

Now we compute

$$\begin{aligned} \frac{\partial g(t)}{\partial t} &= \int_0^\infty dx \sqrt{\frac{\lambda}{2\pi x^3}} \exp\left(-\frac{\lambda(tx - \mu)^2}{2\mu^2 x}\right) \left(-\lambda \frac{tx - \mu}{\mu^2 x} x\right) \\ &= -\frac{\lambda}{2\mu^2} (t\mathbb{E}[x] - \mu) \end{aligned}$$

If we compare that with above, where  $g(t) = 1$  and thus  $\frac{\partial g(t)}{\partial t} = 0$ , we find

$$t\mathbb{E}[x] - \mu = 0.$$

By setting  $t = 1$  we obtain

$$\mathbb{E}[x] = \mu.$$