



CONTENTS

1	Electromagnetic Waves	3
1.1	The greenhouse effect	4
2	Reflection and Photons	9
2.1	Photons	9
2.1.1	The Photoelectric Effect	10
2.1.2	Photons and Momentum	11
2.2	Reflection	12
2.2.1	Law of Reflection	13
2.3	Ellipses and Curved Mirrors	15
2.3.1	The Reflective Properties of Circles and Spheres	15
2.3.2	Ellipses and Ellipsoids	16
2.3.3	Elliptical Orbits	20
2.3.4	Ellipsoids	20
2.3.5	Parabolas and Parabolic Reflectors	22
3	Refraction	29
3.1	Refraction	29
3.2	Snell's Law	29
3.3	Diffraction	30
4	Lenses	31
4.1	Focal Length	32
4.2	Refractive Index	32

A	Answers to Exercises	33
Index		35

CHAPTER 1

Electromagnetic Waves

Sound is a compression wave — to travel, it needs a medium to compress: air, water, etc. (Regardless of what you have seen in movies, sound does not travel through a vacuum!)

Light is an electromagnetic wave — it causes fluctuations in the electric and magnetic fields that are everywhere. It can cross a vacuum, as it does to reach us from the sun.

Electromagnetic waves travel at about 300 million meters per second in a vacuum. The waves travel slower through different types of matter. For example, an electromagnetic wave travels at 225 million meters per second in water.

Electromagnetic waves come in different frequencies. For example, the light coming out of a red laser pointer is usually about 4.75×10^{14} Hz. The wifi data sent by your computer is carried on an electromagnetic wave too. It is usually close to 2.4×10^6 Hz or 5×10^6 Hz.

Because we know how fast the waves are moving, we sometimes talk about their wavelengths instead of their frequencies. Since we know its speed and its oscillation rate, we can calculate its wavelength. Since we want the distance per oscillation, we can divide the speed by the frequency. The light coming out of a laser pointer is $300 \times 10^6 / 4.75 \times 10^{14} = 630 \times 10^{-9}$ m, or 630 nm.

Exercise 1 Wavelengths

A green laser pointer emits light at 5.66×10^{14} Hz. What is its wavelength in a vacuum?

Working Space

Answer on Page 33

This can be simplified as:

$$\lambda = \frac{c}{f}$$

where $c = 3 \times 10^8$ m/s (the speed of light in a vacuum)

We have given names to different ranges of the electromagnetic spectrum in Figure 1.1

Name	Hertz	Meters
Gamma rays	$10^{20} - 10^{24}$	$10^{-14} - 10^{-11}$
X-rays	$10^{16} - 10^{20}$	$10^{-11} - 10^{-8}$
Ultraviolet	$10^{15} - 10^{16}$	$10^{-8} - 10^{-7}$
Blue	$\sim 6 \times 10^{14}$	$\sim 5 \times 10^{-7}$
Red	$\sim 4 \times 10^{14}$	$\sim 7 \times 10^{-7}$
Infrared	$10^{12} - 10^{14}$	$10^{-6} - 10^{-3}$
Microwaves	$10^9 - 10^{12}$	$10^{-3} - 10^{-1}$
Radio waves	$10^3 - 10^9$	$10^{-1} - 10^5$

Figure 1.1: A table of different ranges of the electromagnetic spectrum.

(You may have heard of “cosmic rays” and wonder why they are not listed in this table. Cosmic rays are actually the nuclei of atoms that have been stripped of their electron cloud. These particles come flying out of the sun at very high speeds. They were originally thought to be electromagnetic waves, and were mistakenly named “rays”.)

In general, the lower frequency the wave is, the better it passes through a mass. A radio wave, for example, can pass through the walls of your house, but visible light cannot. The people who designed the microwave oven chose the frequency of 2.45 GHz because the energy from those waves tended to get absorbed in the first few inches of food that it passed through.

1.1 The greenhouse effect

Humans have dug up a bunch of long carbon-based molecules (like oil and coal) and burned them, releasing large amounts of CO_2 into the atmosphere. It may not be obvious why that has made the planet warmer, but the answer is electromagnetic waves.

A warm object gives off infrared electromagnetic waves. That’s why, for example, motion detectors in security systems are actually infrared detectors: even in a dark room, your body gives off a lot of infrared radiation.

You may have heard of “heat-seeking missiles.” These are more accurately called “Infrared homing missiles” because they follow objects giving off infrared radiation – hot things like jet engines.

The sun beams a lot of energy to our planet in the form of electromagnetic radiation: visible light, infrared, ultraviolet. (How much? At the top of the atmosphere directly facing the sun, we get 1,360 watts of radiation per square meter. That is a lot of power!)

Some of that radiation just reflects back into space. 23% is reflected by the clouds and the atmosphere, while 7% makes it all the way to the surface of the planet and is reflected back into space.

The other 71% is absorbed. 48% is absorbed by the surface and 23% is absorbed by the atmosphere. All of that energy warms the planet and the atmosphere so that it gives off infrared radiation. The planet lives in equilibrium; the infrared radiation leaving our atmosphere is exactly the same amount of energy as that 71% of the radiation that it absorbs.

(If the planet absorbs more energy than it releases, the planet gets hotter. Hotter things release more infrared. When the planet is in equilibrium again, it stops getting hotter.)

So, what is the problem with CO₂ and other large molecules in the atmosphere? They absorb the infrared radiation instead of letting it escape into space. This means the planet must be hotter to maintain equilibrium.




Figure 1.2: Greenhouse gas effects from the solar rays.




Figure 1.3: A zoomed in version of greenhouse gases.

The planet is getting hotter, and it is creating a multitude of problems:

- Weather patterns are changing, which leads to extreme floods and droughts.
- Ice and snow in places like Greenland are melting and flowing into the oceans. This is raising sea levels.
- Biomes with biodiversity are resilient. Rapidly changing climate is destroying biodiversity everywhere, which is making these ecosystems very fragile.
- In many places, permafrost, which has trapped large amounts of methane in the ground for millennia, is melting.

That last item is particularly scary, because methane is a large gas molecule — it absorbs even more infrared radiation than CO_2 . As it escapes the permafrost, the problem will get worse.

Scientists are working on four kinds of solutions:

- **Stop increasing the amount of greenhouse gases in our atmosphere.** It is hoped that non-carbon based energy systems like solar, wind, hydroelectric, and nuclear could let us stop burning carbon. Given the methane already being released, it maybe too late for this solution to work on its own.
- **Take some of greenhouse gases out of our atmosphere and sequester them somewhere.** The trunk of a tree is largely made up of carbon molecules. When you grow a tree where there had not been one before, you are sequestering carbon inside the tree. There are also scientists that are trying to develop a process that pulls greenhouse gases out of the air and turn them into solids.
- **Decrease the amount of solar radiation that is absorbed by our planet and its atmosphere.** Clouds reflect a lot of radiation back into space. Could we increase the cloudiness of our atmosphere? Or maybe launch mirrors into orbit around our

planet?

- **Adapt to the changing climate.** These scientists are assuming that global warming will continue, and are working to minimize future human suffering. How will we relocate a billion people as the oceans claim their homes? When massive heat waves occur, how will we keep people from dying? As biodiversity decreases, how can we make sure that species that are important to human existence survive?

What are the greenhouse gases and how much does each contribute to keeping the heat from exiting to space? These numbers are still being debated, but this will give you a feel:

Water vapor	H ₂ O	36 - 72 %
Carbon dioxide	CO ₂	9 - 26 %
Methane	CH ₄	4 - 9 %
Ozone	O ₃	3 - 7 %

Notice that while we talk a lot about carbon dioxide, the most important greenhouse gas is actually water. Why don't we talk about it? Given the enormous surfaces of the oceans, it is difficult to imagine any way to permanently decrease the amount of water in the air. Additionally, a great deal of water in the air is in the form of clouds, which help reflect radiation before it is absorbed.

Reflection and Photons

What happens when light hits a mass?

2.1 Photons

In a previous chapter, we talked about light as a wave, and we mentioned that each color in the rainbow is a different wavelength. You can also think of light as particles of energy called *photons*. The term photons was coined by physicist Gilbert Newton Lewis, the same Lewis known for the Lewis dot structures we have talked about at the beginning of the sequence!

Every photon comes with an amount of energy that determines what color it is. When we are talking about light interacting with objects, your intuition will be right more often if you think of light as a beam of photons. These photons travel at the *vacuum speed of light*, more often called the *vacuum speed of light*. This speed, in empty space or vacuums, is given by 2.9979×10^8 m/s. Photons also

- behave like particles and a wave.
- have zero mass
- has zero charge
- can be created and destroyed when radiation is present
- release electrons upon collisions.

Formula for energy of a photon

For any given photon, its energy, E , is given by

$$E = hf = \frac{hc}{\lambda}$$

where c is the speed of light, λ is the wavelength of the electromagnetic wave, and h Planck's constant: 6.63×10^{-34} m²kg/s. Note that the angular frequency $f = \frac{c}{\lambda}$, and Planck's constant represents the quantum of action, establishing that energy is not continuous but exists in discrete packets called quanta. We will talk in a future

chapter about how charge and energy is quantized, or packeted in integerized values which we call discrete.

For example, a red laser light has a wavelength of about 630 nm. So, the energy in each photon is:

$$\frac{(300 \times 10^6)(6.63 \times 10^{-34})}{630 \times 10^{-9}} = 3.1 \times 10^{-19} \text{ joules}$$

For units sake, this is often abbreviated as $E = \frac{1240}{\lambda}$ where λ is in nanometers and E is in electronvolts (eV).

2.1.1 The Photoelectric Effect

In classical physics, scientists like Planck and Einstein thought that if you shined a light on a metal long enough or brightened it enough, it would eventually knock an electron loose, like a wave slowly pushing a boat to shore. Experiments showed that if the light was the wrong color, meaning too low of a frequency, nothing happened, no matter how bright or powerful the light was.

This showed that light is quantized, and in order to travel up or down the energy levels of an atom, it must absorb or emit a photon with the exact energy difference between those levels. This phenomenon is known as the *photoelectric effect*.

The effect is governed by the conservation of energy. The energy of an incoming photon (E_{photon}) must be greater than the bond holding the electron to the metal. This is described by

$$E_{\text{photon}} = \Phi + K_{\text{max}}$$

where

- $E_{\text{photon}} = hf$: The energy of the incoming photon, where h is Planck's constant and f is the frequency.
- Φ (*Work Function*): The minimum energy required to eject an electron from a specific metal. Each material has a different "cost" to let an electron go.
- f_0 (*Threshold Frequency*): The specific frequency where the photon has exactly enough energy to match the work function ($\Phi = hf_0$). If $f < f_0$, no electrons are ejected. If $f = f_0$, electrons are released but have zero kinetic energy ($K_{\text{max}} = 0$).
- K_{max} : The "leftover" kinetic energy of the electron once released from the metal.

This is how solar panels and digital camera sensors turn light into electricity. Solar panels work like a one-way road for light energy. First, sunlight hits the panel in the form of photons, which act like tiny kicks that knock electrons loose from their atoms. Because the panel has a built-in internal electric field, these loose electrons are forced to move in only one direction (back into the panel), creating a steady flow of Direct Current (DC) electricity that can power your home. We are going to look at camera sensors in depth in a future workbook.

2.1.2 Photons and Momentum

In classical mechanics, we define momentum as the product of mass and velocity ($p = mv$).

This leads to a common question: if a photon has zero rest mass, how can it possess momentum? Energy and momentum are linked, even for massless particles. For a photon, the momentum (p) is inversely proportional to its wavelength (λ):

$$p = \frac{h}{\lambda} \quad \text{or} \quad p = \frac{E}{c} \quad (2.1)$$

Application: For example, solar sails are spacecraft propulsion systems in space. These large, mirror-like sails in space can intercept millions of photons. As these photons reflect off the surface, they undergo a change in momentum (Δp), which, according to Newton's Second Law ($F = \frac{\Delta p}{\Delta t}$), exerts a constant force on the spacecraft, allowing for fuel-free propulsion.

Photonic Interactions

When a photon comes from the sun and hits an object, one of several things can happen:

- The energy of the photon is absorbed by the object. It makes the object a little warmer. If a large proportion of photons hitting the mass are absorbed like this, we say the object is “black”.
- The photon bounces off the object. If the surface is very smooth, the photons bounce in a predictable manner, and we call this *reflection* and we say the object is “shiny”.
- If the surface is rough and the photons are not absorbed, the photons are scattered in random directions. We call this *diffusion*. If most of the photons hitting an object are bounced in random directions, we say that the object is “white”.
- The photon passes through the mass. If the mass has smooth surfaces and a consistent composition, the photons will pass through the mass in a predictable manner. We say that the mass is “transparent”.

- If the photons pass through, but in an unpredictable, scattering manner, we say the mass is “translucent”.

No object absorbs every photon, but chemists are always coming up with “blackier” materials. Vantablack, for example, is a super-black paint that absorbs 99.965% of all photons in the visible spectrum.

No object reflects every photon, but a mirror is pretty close. Let’s talk about reflections in a mirror.

2.2 Reflection

When a beam of light hits a mirror, it bounces off the mirror at the same angle it approached from. That is, if it approaches nearly perpendicular to the mirror, it departs nearly perpendicular to the mirror. If it hits the mirror at a glancing angle, it departs at an angle close to the mirror’s surface. A good video to see this in action can be found here: https://www.youtube.com/shorts/qA_VQZfTiUA

2.2.1 Law of Reflection





Figure 2.1: The angle of *incidence* is equal to the angle of *reflection*.

Law of Reflection

The angle of incidence, denoted as θ_i , is equal to the angle of reflection, denoted as θ_r . This law can be mathematically expressed as:

$$\theta_i = \theta_r$$

where θ_i is the angle between the incident light ray and the normal to the surface, and θ_r is the angle between the reflected light ray and the normal.



Exercise 2 Law of Reflection

Working Space

You are standing 4 meters from a mirror hung on a wall. The bottom of the mirror is the same height as your chin, so you can't see your whole body. You stick a piece of masking tape to your body.

You walk forward until you are only 3 meters from the mirror, then put a piece of masking tape on your body at the new cut-off point. Is the new masking tape higher or lower on your body?

Answer on Page 33

Exercise 3 Photons and Color*Working Space*

There are red photons.

Are there black photons?

Are there white photons?

Are there yellow photons?

*Answer on Page 33***2.3 Ellipses and Curved Mirrors**

Flat mirrors are common and useful, but things get more interesting once you bend the mirrors. In this section, we are going to talk about a few different kinds of curved mirrors.

2.3.1 The Reflective Properties of Circles and Spheres

For example, if you were inside a circular room (a cylinder, actually), you could imagine standing in the center and pointing a flash light in any horizontal direction. The beam of light would bounce right back to you. See Figure 2.2




Figure 2.2: A circle would reflect back onto itself.

How do you know this? Because the tangent line is always perpendicular to the radius to the point of tangency:




Figure 2.3: The reflection on a circle uses a radius which hits a tangent line as a perpendicular bisector.

You could create a spherical room with mirror walls. You would create a platform in the center where you could stand, and if you pointed your flashlight in any direction, its beam of light would shine back at you.

2.3.2 Ellipses and Ellipsoids

Intuitively, you know what an ellipse is: an oval. However, the ellipse is actually an oval with some special properties. This is a good time to talk about those properties.


Mathematicians talk about a *standard* ellipse. A standard ellipse is centered on the origin $(0,0)$ and its long axis is parallel with the x -axis or the y -axis.

Equation for a Standard Ellipse

To be precise, a standard ellipse is the set of points (x,y) that are solutions to the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

Note that $(a,0)$, $(-a,0)$, $(0,b)$, $(0,-b)$ are all part of the set. The complete set looks like this:



We can now talk about two special points: the *foci*. Each focal point is on the long axis of the ellipse. Let's assume for a moment that $a > b$. (Everything works the same if $b > a$, but it gets confusing if we try to deal with both cases simultaneously.)

If p is a point on the ellipse, the distance from p to focal point 1 plus the distance from p to focal point 2 is always $2a$.




Figure 2.4: The both foci and point p forms a triangle with legs $2a - d$ or d .

How do we find the foci? We know they are on the long axis and that they are symmetrical across the short axis. All we need to know is how far are they from the short axis.

Distance from Center to the Foci

If you have an ellipse with a long axis that extends a from the center, and a short axis that extends b from the center, the foci lie on the long axis and are c from the center. Where


$$c = \sqrt{a^2 - b^2}$$

Exercise 4 Foci of an ellipse

Working Space

You need to draw an ellipse that is 12 cm long and 7 cm wide. You have a string, two pushpins, a ruler, and a pencil. Using the ruler, you draw two perpendicular axes.

You will stick one pin at each focal point. Each end of the string will be tied to a push pin. Using the pencil to keep the string taut, you will draw an ellipse.



How far from the short axis are the pushpins placed?

How long is the string between them?

Answer on Page 34

The Reflective Property of Ellipses

Here is something else that is wonderful about an ellipse: Pick any point p on the ellipse. Draw a line from p to each focal point. Draw the line tangent to the ellipse at p . You will see that the angle between the tangent and the line to focal point 1 is equal to the angle between the tangent and the line to focal point 2. See Figure 2.5




Figure 2.5: Reflective Property of Ellipses.

This is known as “The Reflective Property of Ellipses”. Imagine you and your friend Fred are at an ellipse-shaped skating rink, and the edge of the rink is mirrored. You sit at one focal point and your friend sits at the other. If you point a flashlight at the mirror (in any direction!), the beam will bounce off the wall and head directly for Fred.

If Fred ducks out of the way, the beam will bounce again and head back to you.




Figure 2.6: Your laser intersects Fred.

This will work for sound as well. If you whisper while on the focal point, Fred (at the other focal point) will hear you surprisingly well, because all the soundwaves that hit the wall will bounce (just like the light) straight at Fred.

2.3.3 Elliptical Orbits

One more fun fact about ellipses: We often imagine the planets traveling in circular orbits with the sun at the center — they actually travel in elliptical orbits, with the sun as one of the focal points.

The Earth is closest to the sun around January 3rd: 147 million km.

The Earth is farthest from the sun around July 3rd: 152 million km.

(Note that these dates are not the same as the solstices: The southern hemisphere is tilted the most toward the sun around December 21 and tilted most away around June 21.)

2.3.4 Ellipsoids

Just as we can pull the ideas of a circle into three dimensions to make a sphere, we can extend the ideas of the ellipse into three dimensions to talk about ellipsoids. Ellipsoids are like blimps.


The standard ellipsoids are centered at the origin and aligned with the three axes.

Equation for a Standard Ellipsoid

To be precise, a standard ellipse is the set of points (x, y, z) that satisfy the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

Note that $(a, 0, 0)$, $(0, b, 0)$, $(0, 0, c)$ are all part of the set. The complete set looks like



this:

The volume bounded by this ellipsoid is

$$V = \frac{4}{3}\pi abc$$

Of course, a , b , and c can be any positive number, but in the real world, we find ourselves working regularly with ellipsoids where two of the numbers are the same.

Oblate Spheroid

If two axes have the same length and one is shorter, you get something that looks like a sphere compressed in one direction — like a pumpkin. These are called *oblate spheroids*.

The Earth is actually an oblate spheroid; the axis that goes through the north and south pole is shorter than the axes that pass through the equator. How much shorter? Just a little, relatively speaking. The equator is 6,378 km from the center of the Earth; the north pole is 21 km closer.

Prolate Spheroid

If two axes have the same length and one is longer, you get something that looks like a sphere stretched in one direction — like a rugby ball. It is called a *prolate spheroid*.

Like an ellipse, prolate spheroids have two focal points.

Focal Points of a Prolate Spheroid

If the long axis has a radial length of a and the two shorter axes have radial length b , then the focal points are on the long axis. The distance from the center to the focal point is given by

$$c = \sqrt{a^2 - b^2}$$

For any point p on the prolate spheroid, the sum of the distances from p to the focus points will always be $2a$.

It has the reflective property: A photon shot in any direction from one focal point will bounce off the wall and head directly at the other.

Exercise 5 Volume of Ruby Ball

Working Space

Some jokesters once thought it would be fun to make something that looked like a rugby ball, but made out of lead.

A rugby ball is about 30 cm long and has a circumference of 60 cm at its midpoint. A cubic centimeter of lead has a mass of 11.34 grams.

How much would a solid (not hollow) lead ruby ball weigh?


Answer on Page 34

2.3.5 Parabolas and Parabolic Reflectors

You are familiar with quadratic functions:


$$y = ax^2 + bx + c$$

If a is not zero, the graph of a quadratic is a curved line called a *parabola*. The first parabola that most mathematicians think of is the graph of $y = x^2$:



Every parabola has a *focus* and a *directrix*. The focus is a point on the parabola's axis of symmetry. The directrix is a line perpendicular to the axis of symmetry. Every point on the parabola is equal distance from the focus and the directrix.

For the graph of $y = x^2$, the focus is the point $(0, \frac{1}{4})$. The directrix is the line $y = -\frac{1}{4}$:



For example, the point $(1, 1)$ is on this parabola. It is $5/4$ from the directorix. How far is it from the focus? 1 horizontally and $3/4$ vertically.

$$\sqrt{1^2 + \left(\frac{3}{4}\right)^2} = \frac{5}{4}$$


Thus, we have confirmed that $(1, 1)$ is equal distances from the focus and the directrix.

When we think about a parabola and its properties, we usually rotate and translate it to be symmetric around the y -axis, flip it so that it is low in the middle and rising on both sides, and push it up or down until the low point is on the x -axis.


Then, they can all be written:

$$y = \frac{a}{4}x^2$$

where $a > 0$. If a is small, the parabola opens wider.




The focus is at $(0, \frac{1}{a})$ and the directorix is the line $y = -\frac{1}{a}$.



Reflective Property of a Parabola

Assume you have a parabola-shaped mirror. A beam of light shot from the focus will



bounce off the mirror in the direction of the axis of symmetry:

This is why your flashlight has a parabolic mirror. The lightbulb is at the focus, so any photons that hit the mirror are redirected straight forward.

(Note that in the real world, we use parabolic dishes: a parabola rotated around its axis of symmetry.)

The reflection works exactly the same in reverse. There are solar cookers that are big parabolic mirrors. They let you put a pot on the focus point. You move the dish until its axis of symmetry is pointed at the sun.

You will also see a lot of antennas have parabolic dishes. Note that photons that come in parallel to the axis of symmetry are redirected to a single point: where the receiver is.



Figure 2.7: A real life parabolic dish as a satellite.

Sometimes in a science museum, you will see two parabolic dishes far apart and pointed at each other. One person speaks with their mouth at the focus of one. The other person listens with their ear at the focus of the other. Even though you are very far apart, it sounds like they are really, really close.



Figure 2.8: Two parabolic dishes send waves between each other.

This is because the speaker's parabolic wall focuses the sound energy in a nice beam the size of the wall pointed straight at the listener's parabolic wall. The listener's wall focuses

the energy of that beam at the listener's ear.

Refraction

3.1 Refraction

The refraction of light is a phenomenon where light changes its direction when it passes from one medium to another. The change in direction is due to a change in the speed of light as it moves from one medium to another. Do not confuse this with rarefaction, which sounds similar and is a similar wave concept for decompression!

3.2 Snell's Law

This phenomenon is explained by Snell's law, which states:

$$n_1 \cdot \sin(\theta_1) = n_2 \cdot \sin(\theta_2) \quad (3.1)$$

where:

- n_1 and n_2 are the indices of refraction for the first and second media, respectively. The index of refraction is the ratio of, c the speed of light in a vacuum to, v the speed of light in the medium, $n = \frac{c}{v}$. It is a dimensionless quantity. $n \geq 1$ for all materials, such that $n_{\text{vacuum}} = 1$ and $n_{\text{air}} \approx 1$. See a list of all mediums here: https://en.wikipedia.org/wiki/List_of_refractive_indices
- θ_1 and θ_2 are the angles of incidence and refraction, respectively. These angles are measured from the normal (perpendicular line) to the surface at the point where light hits the boundary.




Figure 3.1: Refraction occurs when light changes the medium it is in.

The angle of incidence (θ_1) is the angle between the incident ray and the normal to the interface at the point of incidence. Similarly, the angle of refraction (θ_2) is the angle between the refracted ray and the normal.


When light travels from a medium with a lower refractive index to a medium with a higher refractive index, it bends towards the normal. Conversely, when light travels from a medium with a higher refractive index to one with a lower refractive index, it bends away from the normal.

3.3 Diffraction


Lenses

Lenses are optical devices with perfect or approximate axial symmetry that transmit and refract light, converging or diverging the beam. There are two main types of lenses, distinguished by their shape and the way they refract light:

- **Converging (or Convex) Lenses:** These are thicker at the center than at the edges. When parallel light rays enter a convex lens, they converge to a point called the focal point. Examples of converging lenses include magnifying glasses and camera lenses.



- **Diverging (or Concave) Lenses:** These are thinner at the center than at the edges. When parallel light rays enter a concave lens, they diverge or spread out. These lenses are often used in glasses to correct nearsightedness.



4.1 Focal Length

The focal length of a lens is the distance between the center of the lens and the focal point. It is determined by the lens shape and the refractive index of the lens material. For a converging lens, the focal length is positive; for a diverging lens, the focal length is negative.

4.2 Refractive Index

The refractive index of a material is a measure of how much the speed of light is reduced inside the material. The refractive index n of a material is given by the ratio of the speed of light in a vacuum c to the speed of light v in the material:

$$n = \frac{c}{v}$$

The refractive index affects how much a light ray changes direction, or refracts, when entering the material at an angle. A higher refractive index indicates that light travels slower in that medium, and the light ray will bend more towards the normal.

Lenses work by refracting light at their two surfaces. By choosing the right lens shape and material, lenses can be designed to bring light to a focus, spread it out, or perform more complex transformations.

Answers to Exercises

Answer to Exercise 1 (on page 3)

$$\frac{300 \times 10^6}{5.66 \times 10^{14}} = 530 \times 10^{-9} = 530 \text{ nm}$$

Answer to Exercise 2 (on page 14)

Assuming the mirror is truly vertical and the floor is truly horizontal, the new cut off should be exactly the same as the old one: It should be below your chin the same amount that your eyes are above your chin.

Illustration Here

Answer to Exercise 3 (on page 15)

Are there white photons? No. What we call “white” is a blend of photons that are several different colors.

Some people like to say white light is the combination of all visible colors of photons in equal amounts. That seems oddly specific and unusual.

Maybe it is better to imagine it from the human experience of white light. In our eyes, we have three different types of color-sensing cones, which generally correspond to the red, green, and blue regions of the spectrum. When all three are excited about equal amounts, humans experience that as white. On your computer screen, for example, what you see as white is just a blend of three colors of photons: a red, a green, and a blue.

Are there black photons? No. What we call “black” is an absence of photons in the visible range.


Are there yellow photons? Yes! There is a region of the color spectrum that is yellow. It has a wavelength of about 527 nm. Photons at this energy level excite both our green-sensitive and red-sensitive cones (we will expand on this in the eye chapter!). Your computer

monitor does not actually create light with a 527 nm wavelength. Instead, it creates red light and green light, which our eyes interpret as yellow.

Answer to Exercise 4 (on page 18)

The length of the string is easy: $2 \times 12 = 24$ cm.

The distance from the center to the focal point is $\sqrt{12^2 - 7^2} \approx 6.78$ cm.



Answer to Exercise 5 (on page 22)

We need the distance from the center out to each of the three axes. We know that $a = \left(\frac{1}{2}\right) 30 = 15$ cm.

We can calculate the b and c (which are equal) using the circumference given: $2b\pi = 60$, so $c = b \approx 9.55$ cm.

The volume, then is

$$V = \frac{4}{3}\pi(15)(9.55)(9.55) \approx 5,730 \text{ cubic centimeters}$$

The mass would be $5,730 \times 11.34 = 64,973$ grams or about 65 kg.



INDEX

- absorption
 - photon, [11](#)
- circle
 - reflections in, [15](#)
- diffusion, [11](#)
- directix, [23](#)
- earth
 - shape of, [21](#)
- ellipse, [16](#)
 - focus points, [17](#)
 - reflective property of, [19](#)
- ellipsoid, [20](#)
- elliptical
 - orbit, [20](#)
- focal length, [32](#)
- foci, [17](#)
- focus, [23](#)
- lenses, [31](#)
- mirror, [12](#)
- oblate spheroid, [21](#)
- oblate spheroids, [21](#)
- parabola, [23](#)
- photoelectric effect, [10](#)
- photon, [9](#)
- photons, [9](#)
- prolate spheroid, [21](#)
- reflection, [11](#)
 - Law of, [13](#)
 - law of, [13](#)
- refractive index, [32](#)
- speed of light, [9](#)
- standard, [16](#)
- Threshold Frequency, [10](#)
- translucent, [12](#)
- transparent, [11](#)
- vacuum speed of light, [9](#)
- Vantablack, [12](#)
- Work Function, [10](#)