

Drag

The very first computers were created to do calculations of how artillery would fly when shot at different angles. The calculations were similar to the ones you did earlier for the flying hammer, with two important differences:

- They were interested in two dimensions: the height and the distance across the ground.
- However, artillery flies a lot faster than a hammer, so they also had to worry about drag from the air.




Figure 1.1: A cannonball shot has a parabolic trajectory.

1.1 Wind resistance

The first thing they did was put one of the shells in a wind tunnel. They measured how much force was created when they pushed 1 m/s of wind over the shell. Let's say it was 0.1 newtons.

One of the interesting things about the drag from the air (often called *wind resistance*) is that it increases with the *square* of the speed. Thus, if the wind pushing on the shell is 3

m/s, instead of 1 m/s, the resistance is $3^2 \times 0.1 = 0.9$ newtons.

(Why? Intuitively, three times as many air molecules are hitting the shell and each molecule is hitting it three times harder.)

So, if a shell is moving with the velocity vector v , the force vector of the drag points in the exact opposite direction. If μ is the force of wind resistance of the shell at 1 m/s, then the magnitude of the drag vector is $\mu|v|^2$ with μ being the wind resistance force.

1.2 Initial velocity and acceleration due to gravity

Let's say a shell is shot out of a tube at s m/s, and the tube is tilted θ radians above level. The initial velocity will be given by the vector $[s \cos(\theta), s \sin(\theta)]$




Figure 1.2: The initial velocity vector of the shell.

(The velocity of the shell is actually a 3-dimensional vector, but we are only going to worry about height and horizontal distance; we are assuming that the operator pointed it in the right direction.)

To figure out the path of the shell, we need to compute its acceleration. We remember that

$$F = ma$$

(Note that F and a are vectors.) Dividing both sides by m , we get:

$$a = \frac{F}{m}$$

Let's figure out the net force on the shell, so that we can calculate the acceleration vector.

If the shell has a mass of b , the force due to gravity will be in the downward direction, with a magnitude of $9.8b$ newtons.

To get the net force, we will need to add the force due to gravity with the force due to wind resistance.

1.3 Simulating artillery in Python

Create a file called `artillery.py`.

```

1  import numpy as np
2      import matplotlib.pyplot as plt
3
4      # Constants
5      mass = 45 # kg
6      start_speed = 300.0 # m/s
7      theta = np.pi/5 # radians (36 degrees above level)
8      time_step = 0.01 # s
9      wind_resistance = 0.05 # newtons in 1 m/s wind
10     force_of_gravity = np.array([0.0, -9.8 * mass]) # newtons
11
12     # Initial state
13     position = np.array([0.0, 0.0]) # [distance, height] in meters
14     velocity = np.array([start_speed * np.cos(theta), start_speed *
15         ↪ np.sin(theta)])
16     time = 0.0 # seconds
17
18     # Lists to gather data
19     distances = []
20     heights = []
21     times = []
22
23     # While shell is aloft
24     while position[1] >= 0:
25         # Record data
26         distances.append(position[0])
27         heights.append(position[1])
28         times.append(time)
29
30         # Calculate the next state
31         time += time_step
32         position += time_step * velocity
33
34         # Calculate the net force vector
35         force = force_of_gravity - wind_resistance * velocity**2


```

```
36     # Calculate the current acceleration vector
37     acceleration = force / mass
38
39     # Update the velocity vector
40     velocity += time_step * acceleration
41
42     print(f"Hit the ground {position[0]:.2f} meters away at {time:.2f} seconds.")
43
44     # Plot the data
45     fig, ax = plt.subplots()
46     ax.plot(distances, heights)
47     ax.set_title("Distance vs. Height")
48     ax.set_xlabel("Distance (m)")
49     ax.set_ylabel("Height (m)")
50     plt.show()
```

When you run it, you should get a message like:

```
1 Hit the ground 1696.70 meters away at 20.73 seconds.
```

You should also see a plot of the shell's path:



1.4 Terminal velocity

If you shot the shell very, very high in the sky, it would keep accelerating toward the ground until the force of gravity and the force of the wind resistance were equal. The speed at which this happens is called the *terminal velocity*. The terminal velocity of a falling human is about 53 m/s.

Note that kinematic equations do not apply to terminal velocity, because the acceleration is not constant. Instead, we can use the fact that at terminal velocity, the force of wind resistance equals the force of gravity.

Exercise 1 Terminal velocity

What is the terminal velocity of the shell described in our example?

Working Space

Answer on Page 7

This is a draft chapter from the Kontinua Project. Please see our website (<https://kontinua.org/>) for more details.

Answers to Exercises

Answer to Exercise 1 (on page 5)

The force of gravity is $9.8 \times 45 = 441$ newtons.

At any speed s , the force of wind resistance is $0.05 \times s^2 = 0.05s^2$ newtons.

At terminal velocity, $0.05s^2 = 441$.

Solving for s , we get $s = \sqrt{\frac{441}{0.05}}$

Thus, terminal velocity should be about 94 m/s.



INDEX

terminal velocity, [5](#)

wind resistance, [1](#)