

Functions and Their Graphs

Functions are a major part of science, engineering, and math. You can think of a function as a machine: you put something into the machine, it processes it, and out comes something else: a product. Just as we often use the variable x to stand in for a number, we often use the variable f to stand in for a function. $f(x)$ is said as “ f of x ”. However, this could be any letter or word that takes an input x , maybe even $zebra(x)$!

For example, we might ask, “Let the function f be defined like this:

$$f(x) = -5x^2 + 12x + 2$$

What is the value of $f(3)$, said as “ f of 3”?

You would run the number 3 through “the machine”: $-5(3^2) + 12(3) + 2 = -7$. The answer would be “ $f(3)$ is -7 ”.

However, some functions are not defined for every possible input. For example:

$$f(x) = \frac{1}{x}$$

This is defined for any x except 0, because you can’t divide 1 by 0. The set of values that a function can process is called its *domain*, and resulting output values are called the range. It is important to note that functions have only one *output* for each *input*. However, multiple inputs can have the same output. A relationship where one input can result in the more than one output is not a function, but a relation. This can be proven by the [vertical line test](#).

Exercise 1 Domain of a function

Working Space

Let the function f be given by $f(x) = \sqrt{x-3}$. What is its domain?

Answer on Page 17

1.1 Graphs of Functions

If you have a function, f , its graph is the set of pairs (x, y) such that $y = f(x)$. We usually draw a picture of this set, called a *graph*. The graph not only includes the picture, but also the values of x and y used to create it.

Here is the graph of the function $f(x) = -5x^2 + 12x + 2$:




Figure 1.1: The function $f(x) = -5x^2 + 12x + 2$ *graphed* on an xy -plane from $x = -1$ to $x = 3$. The inputs x are graphed along the x -axis, and the output of the function $f(x)$ is plotted along the y -axis.

(Note that this is just part of the graph; it goes infinitely in both directions. Remember your vectors!)

Here is the graph of the function $f(x) = \frac{1}{x}$:




Figure 1.2: The function $f(x) = \frac{1}{x}$ graphed on a coordinate plane.

Notice the *asymptotes* at $y = 0$ and $x = 0$. Here, the function is undefined at $x = 0$ and $y = 0$, but is continually approaching it. It never actually reaches these values though! $f(0)$ is not defined as you cannot divide by zero (recall division rules) and as x is increasing, $f(x)$ gets infinitesimally small but never equal to zero. We will talk more about these in the rational functions chapter.

To draw a graph, take each x value (usually in increments of 1) and determine its corresponding y value. Then, plot those points on a graph, using the origin as the starting point.

Exercise 2 Draw a graph

Working Space

Let the function f be given by
 $f(x) = -3x + 3$. Sketch its graph.

Answer on Page 17

1.2 Can this be expressed as a function?

Note that not all sets can be expressed as graphs of functions. For example, here is the set of points (x, y) such that $x^2 + y^2 = 9$:





Figure 1.4: The graph of $x^2 + y^2 = 9$.

This cannot be the graph of a function, because what would $f(0)$ be? 3 or -3? This set fails what we call “the vertical line test”: If any vertical line contains more than one point from the set, it isn’t the graph of a function. For example, the vertical line $x = 2$ would cross the graph twice:


Figure 1.5: Vertical line test on $x^2 + y^2 = 9$.

1.3 Inverses

Some functions have inverse functions. If a function f is a machine that turns number x into y , the inverse (usually denoted f^{-1}) is the machine that turns y back into x .

For example, let $f(x) = 5x + 1$. Its inverse is $f^{-1}(x) = (x - 1)/5$. (Spot check it: $f(3) = 16$ and $f^{-1}(16) = 3$)

Does the function $f(x) = x^3$ have an inverse? Yes, $f^{-1}(x) = \sqrt[3]{x}$. Let's plot the function (solid line) and its inverse (dashed):

Figure 1.6: The function $f(x) = x^3$ and its inverse.

The inverse is the same as the function, just with its axes swapped. The inverse function is *reflected* across the line $y = x$. This tells us how to solve for an inverse: We swap x and y and solve for y .

For example, if you are given the function $f(x) = 5x + 1$, its graph is all (x, y) such that $y = 5x + 1$. The graph of its inverse is all (x, y) , such that $x = 5y + 1$. Solving for y gives you $y = (x - 1)/5$. So we can say that $f^{-1}(x) = (x - 1)/5$, QED.

Not every function has an inverse. For example, $f(x) = x^2$. Note that $f(2) = f(-2) = 4$. What would $f^{-1}(4)$ be? 2 or -2? This implies the “horizontal line test”: If any horizontal line contains more than one point of a function’s graph, that function has no inverse. If a function passes the horizontal line test, it is called “one-to-one”, meaning there is exactly one x that gives each y .




Figure 1.7: The horizontal line test run on $f(x) = x^2$.

In some problems, you need an inverse, but you don’t need the whole domain, so you trim the domain to a set you can define an inverse on. This allow you to make claims such as “If we restrict the domain to the nonnegative numbers, the function $f(x) = x^2 - 5$ has an inverse: $f^{-1}(x) = \sqrt{x + 5}$.”

This raises the question: What is the domain of the inverse function f^{-1} ?

If we let X be the domain of f , we can run every member of X through “the machine” and gather them in a set on the other side. This set would be the *image* of the f “machine”. (This is the *range* of f .)

What is the image of $f(x) = x^2 - 5$? It is the set of all real numbers greater than or equal to -5. We write this:

$$\{x \in \mathbb{R} | x \geq -5\}$$

Now we can say: **The range (or image) of the function is the domain of the inverse function.**

In inverse functions, the domain and range get swapped: the domain of the function is the range of the inverse function, and visa versa. In our example, we can use any number greater than or equal to -5 as input into the inverse function.




Figure 1.8: Domain and range of functions and their inverses get swapped.

Exercise 3 Find the inverse

Working Space

Let $f(x) = (x-3)^2 + 2$. Sketch the graph.

Using all the real numbers as a domain, does this function have an inverse?

How would you restrict the domain to make the function invertible?

What is the inverse of that restricted function?

What is the domain of the inverse?

Answer on Page 17

Exercise 4

A function is given by a table of values, a graph, or a written description. Determine whether it is one-to-one.



Working Space

1.

x	1	2	3	4	5	6
$f(x)$	1.5	2.0	3.6	5.3	2.8	2.0

2.

x	1	2	3	4	5	6
$f(x)$	1.0	1.9	2.8	3.5	3.1	2.9



5. $f(t)$ is the height of a football t seconds after kickoff
6. $v(t)$ is the velocity of a dropped object


Answer on Page 18

1.4 Graphing Calculators

One really easy way to understand your function better is to use a graphing calculator. Desmos is a great, free online graphing calculator.

In a web browser, go to Desmos: <https://www.desmos.com/calculator>

In the field on the left, enter the function $y = x^2 - x - 6$. (For the exponent, just prefix it with a caret symbol: "x^2".)



1.5 Even and Odd Functions

1.5.1 Even Functions

An even function is symmetric about the y-axis. That means if you fold the graph along the y-axis, both sides will match perfectly. Note that the input value yields the same output regardless of whether the input value is positive or negative.

Even Functions

A function $f(x)$ is even if

$$f(-x) = f(x)$$

for all x in its domain.

Examples:

- $f(x) = x^2$
- $f(x) = \cos(x)$
- any $f(x) = x^n$ where n is an even number
- $f(x) = |x|$

On a graph, a function will be a mirror image across the y-axis.

1.5.2 Odd Functions

An odd function is symmetric about the origin. That means if you rotate the graph 180° about the origin, it lands on itself. Algebraically, when you input a value k , you get some output n ; if you negate that input as $-k$, the output is also negated resulting in $-n$.

Odd Functions

A function $f(x)$ is odd if

$$f(-x) = -f(x)$$

for all x in its domain.

Equivalently,

$$f(x) + f(-x) = 0$$

for all x in its domain.

- $f(x) = x^3$
- $f(x) = \sin(x)$
- $f(x) = \tan(x)$
- any $f(x) = x^n$ where n is an odd number

On a graph, rotating the graph 180° about the origin will yield the same graph.

1.5.3 Neither Even Nor Odd

Some functions are neither even nor odd.

For example, $f(x) = x^3 + 1$ does not satisfy either condition.

1.6 Piecewise Functions

One other thing to mention is the existence of *piecewise functions*. A piecewise function is one defined by two different equations. You chose which function to evaluate dependent on the domain. Let's look at a simple example:

$$f(x) = \begin{cases} \frac{1}{x} & x < 0 \\ x^2 & x \geq 0 \end{cases}$$

This piecewise function has two cases: when $x < 0$, we use the equation $\frac{1}{x}$; when $x \geq 0$, we use x^2 . Notice that we don't consider $x = 0$ for $\frac{1}{x}$ since it would be undefined, but it is defined under x^2 . The domain of $f(x)$ is defined for all values of x ! Let's make a table of values:

x	-2	-1	0	0.1	1	2
$f(x)$	$-\frac{1}{2}$	-1	0	0.01	1	4

We can graph this like the following




Figure 1.10: The piecewise function $f(x)$ graphed.

Notice the *closed circle*. This indicates that the value at the is included at a disjunction. An *open circle* would indicate that the value at that point is not included. Say we have the following function:

$$g(x) = \begin{cases} x^2 + 1 & x < 0 \\ x^2 & x \geq 0 \end{cases}$$

Graphing this, we obtain an open circle at $(0, 1)$ and a closed circle at $(0, 0)$.




Figure 1.11: The piecewise function $g(x)$ graphed.

Exercise 5 **Absolute Value**

Working Space

Rewrite the absolute value function $|x|$
as a piecewise function.

Answer on Page 19


Exercise 6 Piecewise 1

Working Space

$$f(x) = \begin{cases} x + 2 & \text{if } -5 \leq x < -1 \\ x^2 & \text{if } -1 \leq x \leq 2 \\ 3 - x & \text{if } 2 < x \leq 5 \end{cases}$$

For the equation above, create a table for every integer from -5 to 5 (inclusive), and graph.

Here is a blank coordinate plane for plotting the piecewise function:



Answer on Page 19

1.6.1 Floor and Ceiling Functions

Another common piecewise function you may see are the *floor function* and *ceiling function*. The floor function, which takes the any real number x , is defined as

$$\lfloor x \rfloor \text{ or } \text{floor}(x) = \begin{cases} n & \text{if } n \leq x < n + 1, \quad n \in \mathbb{Z} \end{cases}$$

This means that you take any number x and find the closest integer less than or equal to it. For example, $\lfloor 0.7 \rfloor$ would be floored to 0.




Figure 1.12: The flooring function, $f(x) = \lfloor x \rfloor$.

Similarly, the ceiling function is defined as follows:

$$\lceil x \rceil \text{ or } \text{ceil}(x) = \begin{cases} n & \text{if } n-1 < x \leq n, \quad n \in \mathbb{Z} \end{cases}$$

The ceiling function takes in any real numbers and returns the next greatest integer. For example, $\lceil 8.00001 \rceil$ returns 9, the next greatest integer. For any valid integer like $x = 7$, the $\lfloor 7 \rfloor$ and $\lceil 7 \rceil$ are both equal to 7.




Figure 1.13: The ceiling function, $f(x) = \lceil x \rceil$.

Exercise 7 More TVs is Better*Working Space*

A store sells televisions with pricing that depends on how many units a customer buys:

- For the first 5 TVs, the cost per TV is \$500
- For the next 10 TVs, the cost per TV is \$350
- Any additional TVs cost \$300

A hotel is buying TVs for each hotel room.

1. Construct a piecewise function $C(x)$ for the cost of TVs. Assume TVs must be a discrete integer (you cannot buy 8.5 TVs).
2. Find the cost of 12 TVs using your function.
3. When is average cost $< \$360$?

*Answer on Page 19***1.6.2 Invalid Piecewise Functions**

One thing to note when creating piecewise functions is to make sure your cases do not contradict each other, or in other words, create cases that define two different possibilities for x at a given value. For example,

$$f(x) = \begin{cases} 10.2 & x = 4 \\ x^2 & x \geq 0 \end{cases}$$

This creates an impossible and thus invalid piecewise function, as at $x = 4$, $f(x)$ is both defined as 10.2 and 16. This would be impossible ($\rightarrow\leftarrow$). In this case, either you define

the value at DNE or undefined, or revise the piecewise function at that point. Most often, it is done as follows:

$$f(x) = \begin{cases} 10.2 & x = 4 \\ x^2 & x \geq 0, x \neq 4 \end{cases}$$

Then our value at $x = 4$ is equal to 10.2.

1.7 Summary

This chapter talked about functions. Functions take in an input and spit out an output, like a magician putting a banana in a hat and releasing a white bunny. However, this is not magic, but math. Each input is given exactly one output. Not all functions have equations that are defined for all real numbers.

We can also graph functions according to their inputs (the x values on the x -axis) and outputs (the returns of the function $f(x)$ plotted on the y axis).

We established that relations must pass the vertical line test to be functions. Some functions have inverse functions, which reverse the action of the original function. A function has an inverse only if it is one-to-one, which can be checked with the horizontal line test. When a function and its inverse are graphed, they are reflections of each other across the line $y = x$ and the domain and range of a function switch roles in its inverse.

Functions can also be classified by symmetry (even or odd). Piecewise functions are by different formulas on different parts of the domain.

This is a draft chapter from the Kontinua Project. Please see our website (<https://kontinua.org/>) for more details.

Answers to Exercises

Answer to Exercise 1 (on page 2)

You can only take the square root of nonnegative numbers, so the function is only defined when $x - 3 \geq 0$. Thus, the domain is all real numbers greater than or equal to 3.

Answer to Exercise 2 (on page 4)

The graph of this function is a line, its slope is -3 , and it intersects the y axis at $(0, 3)$.





Figure 1.3: The resulting graph of $f(x) = -3x + 3$.

Answer to Exercise 3 (on page 7)

This graph is the graph of $y = x^2$ that has been moved to the right by three units and up two units:

Figure 1.9: The graph of $f(x) = (x - 3)^2 + 2$.

To prevent any horizontal line from containing more than one point of the graph, you would need to use the left or the right side — either $\{x \in \mathbb{R} \mid x \leq 3\}$ or $\{x \in \mathbb{R} \mid x \geq 3\}$. Most people will choose the right side; the rest of the solution will assume that you did too.

To find the inverse we swap x and y : $x = (y - 3)^2 + 2$

Next, we solve for y to get the inverse: $y = \sqrt{x - 2} + 3$

You can take the square root of nonnegative numbers. So the function $f^{-1}(x) = \sqrt{x - 2} + 3$ is defined whenever x is greater than or equal to 2.

Answer to Exercise 4 (on page 8)

1. This function is not one-to-one. From $x = 3$ to $x = 4$, the function increases from 3.6 to 5.3, which means it must pass through $f(x_1) = 4.0$. From $x = 4$ to $x = 5$, the function decreases from 5.3 to 2.8, which means it must pass through $f(x_2) = 4.0$ again.
2. This function is not one-to-one by a similar argument in the above solution
3. This function is not one-to-one, because it fails the horizontal line test
4. This function is one-to-one, because it passes the horizontal line test
5. $f(t)$ would not be one-to-one because the football must pass through each height (except the peak height) both on the way up and on the way back down
6. $v(t)$ would be one-to-one because a falling object only speeds up. Therefore, every time has a unique speed.

Answer to Exercise 5 (on page 12)


From $x \in [0, \infty)$, we say that $|x| = x$. For $x < 0$, we say $|x| = -x$, since we want to invert the magnitude. So,

$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0 \end{cases}$$

Answer to Exercise 6 (on page 13)

First, let's create a table of values for each integer from -5 to 5:

x	-5	-4	-3	-2	-1	0	1	2	3	4	5
$f(x)$	-3	-2	-1	0	1	0	1	4	0	-1	-2



Answer to Exercise 7 (on page 15)

1.

$$C(x) = \begin{cases} 400x & 0 < x \leq 5 \\ 2000 + 350(x - 5) & 5 < x \leq 15 \\ 5500 + 300(x - 15) & x > 15 \end{cases}$$

2. $C(12) = 2000 + 350(12 - 5) = \4450

3. The average cost is \$360 when $\frac{C(x)}{x} < 360$. This will only happen during the 3rd case, when $x \geq 15$.

$$C(x) = 5500 + 300(x - 15) = 5500 + 300x - 4500 = 1000 + 300x$$

Now,

$$\frac{1000 + 300x}{x} < 360$$

$$300 + \frac{1000}{x} < 360$$

$$\frac{1000}{x} < 60$$

$$1000 < 60x$$

$$x > \frac{1000}{60} \approx 16.67 \rightarrow 17$$

There must be 17 TVs sold for the average cost to fall to \$360



INDEX

asymptote, [3](#)

asymptotes, [3](#)

ceiling function, [13](#)

domain, [1](#)

floor function, [13](#)

function, [1](#)

horizontal line test, [6](#)

piecewise functions, [10](#)

range, [6](#)

reflections, [5](#)

vertical line test, [5](#)