

# Units and Conversions

Accurate measurements are at the heart of good data and good problem solving. Engineers need to be able to describe many different types of phenomena, such as distance, sound, light, force, and more.

At this point, you are working with a lot of units: grams for weight, joules for energy, newtons for force, meters for distance, seconds for time, and so on. For each type of measurement, there are several different units. For example, distance can be measured in feet, miles, and light-years.

## Some Equivalencies

| Distance                |                                       |
|-------------------------|---------------------------------------|
| 1 mile                  | 1.6093 kilometers                     |
| 1 foot                  | 0.3048 meters                         |
| 1 inch                  | 2.54 centimeters                      |
| 1 light-year            | $9.461 \times 10^{12}$ kilometers     |
| Volume                  |                                       |
| 1 milliliter            | 1 cubic centimeter                    |
| 1 quart                 | 0.9461 liters                         |
| 1 gallon                | 3.7854 liters                         |
| 1 fluid ounce           | 29.6 milliliters                      |
| Mass                    |                                       |
| 1 pound                 | 0.4535924 kilograms                   |
| 1 ounce                 | 0.4535924 grams                       |
| 1 metric ton            | 1000 kilograms                        |
| Force                   |                                       |
| 1 newton                | 1 kilogram meter per sec <sup>2</sup> |
| Pressure                |                                       |
| 1 pascal                | 1 newton per square meter             |
| 1 bar                   | 0.98692 atmosphere                    |
| 1 pound per square inch | 6897 pascals                          |
| Energy                  |                                       |
| 1 joule                 | 1 newton meter                        |
| 1 calorie               | 4.184 joules                          |
| 1 kilowatt-hour         | $3.6 \times 10^6$ joules              |

(You don't need to memorize these! Just remember that this page is here.)

In the metric system, prefixes are often used to express a multiple. Here are the common prefixes:

### Common Prefixes for Metric Units

|       |               |
|-------|---------------|
| giga  | $\times 10^9$ |
| mega  | $\times 10^6$ |
| kilo  | $\times 10^3$ |
| milli | $\div 10^3$   |
| micro | $\div 10^6$   |
| nano  | $\div 10^9$   |

(These are worth memorizing. Here's a mnemonic for the most common ones: "King Henry Doesn't Usually Drink Chocolate Milk" Or Kilo ( $\times 10^3$ ), Hecto ( $\times 10^2$ ), Deca ( $\times 10^1$ ), Unit (for example: gram) ( $\times 10^0$ ), Deci ( $\times 10^{-1}$ ), Centi ( $1 \times 10^{-2}$ ), Milli ( $\times 10^{-3}$ ).)

## 1.1 Conversion Factors

Here is a really handy trick to remembering how to do conversions between units.

Often, you will be given a table like the one above, and someone will ask you "How many miles are in 0.23 light-years?" You know that 1 mile = 1.6093 kilometers and that 1 light-year is  $9.461 \times 10^{12}$  kilometers. How do you do the conversion?

The trick is to treat the two parts of the equality as a fraction that equals 1. In other words, you think:

$$\frac{1 \text{ miles}}{1.6093 \text{ km}} = \frac{1.6093 \text{ km}}{1 \text{ miles}} = 1$$

and

$$\frac{1 \text{ light-years}}{9.461 \times 10^{12} \text{ km}} = \frac{9.461 \times 10^{12} \text{ km}}{1 \text{ light-years}} = 1$$

We call these fractions *conversion factors*.

Now, your problem is

$$0.23 \text{ light-years} \times \text{Some conversion factors} = ? \text{ miles}$$

Note that when you multiply fractions together, things in the numerators can cancel with things in the denominator:

$$\left(\frac{31\pi}{47}\right)\left(\frac{11}{37\pi}\right) = \left(\frac{31\cancel{\pi}}{47}\right)\left(\frac{11}{37\cancel{\pi}}\right) = \left(\frac{31}{47}\right)\left(\frac{11}{37}\right)$$

When working with conversion factors, you will do the same with the units:

$$\begin{aligned} 0.23 \text{ light-years} \left(\frac{9.461 \times 10^{12} \text{ km}}{1 \text{ light-years}}\right) \left(\frac{1 \text{ miles}}{1.6093 \text{ km}}\right) &= \\ 0.23 \cancel{\text{ light-years}} \left(\times \frac{9.461 \times 10^{12} \cancel{\text{ km}}}{1 \cancel{\text{ light-years}}}\right) \left(\frac{1 \text{ miles}}{1.6093 \cancel{\text{ km}}}\right) &= \frac{(0.23)(9.461 \times 10^{12})}{1.6093} \text{ miles} \end{aligned}$$

### Exercise 1 Simple Conversion Factors

*Working Space*

How many calories are in 4.5 kilowatt-hours?

*Answer on Page 7*

## 1.2 Conversion Factors and Ratios

Conversion factors also work on ratios. For example, if you are told that a bug is moving 0.5 feet every 120 milliseconds, what is that in meters per second?

The problem then is

$$\frac{0.5 \text{ feet}}{120 \text{ milliseconds}} = \frac{? \text{ m}}{\text{second}}$$

So you will need conversion factors to replace the “feet” with “meters” and to replace “milliseconds” with “seconds”:

$$\left(\frac{0.5 \text{ feet}}{120 \text{ milliseconds}}\right) \left(\frac{0.3048 \text{ meters}}{1 \text{ foot}}\right) \left(\frac{1000 \text{ milliseconds}}{1 \text{ second}}\right) = \frac{(0.5)(0.3048)(1000)}{120} \text{ m/second}$$

## Exercise 2 Conversion Factors

Working Space

The hole in the bottom of the boat lets in 0.1 gallons every 2 minutes. How many milliliters per second is that?

Answer on Page 7

## 1.3 When Conversion Factors Don't Work

Conversion factors only work when the units being converted are proportional to each other. Gallons and liters, for example, are proportional to each other: If you have  $n$  gallons, you have  $n \times 3.7854$  liters.

Degrees celsius and degrees fahrenheit are *not* proportional to each other. If your food is  $n$  degrees celsius, it is  $n \times \frac{9}{5} + 32$  degrees fahrenheit. You can't use conversion factors to convert celsius to fahrenheit.

## 1.4 SI units

In standardized sciences and studies, there has a set of units for all scientists to commonly use. This is done so that scientific research, other papers, and different international communities can easily communicate their research with one another. This system is referred to as **SI Units**, or the **international system of units**.

Here is the very basic table for SI units:

|                     |               |
|---------------------|---------------|
| Time                | second (s)    |
| Length              | meter (m)     |
| Mass                | kilogram (kg) |
| Electric Current    | ampere (A)    |
| Temperature         | Kelvin (K)    |
| Number of substance | mole (mol)    |
| Luminous intensity  | candela (cd)  |

All other units can be derived from these 7 base units from the SI system.

Often, you will see units represented in the form of exponents. Let's look at velocity as an example. Velocity is often described as **meters per second**. This can be written as a fraction, or with seconds to a negative power:

$$v = \frac{\text{m}}{\text{s}} = \text{m} \cdot \text{s}^{-1}$$

Similarly, we can represent acceleration (meters per second per second) with exponents

$$a = \frac{\text{m}}{\text{s}^2} = \text{m} \cdot \text{s}^{-2}$$

### 1.4.1 Converting from SI units

There are many common units that are often used in place of SI units. Here is another table.

| Physical Quantity | Name    | Symbol | Expression in Base Units                         |
|-------------------|---------|--------|--------------------------------------------------|
| Force             | Newton  | N      | $\text{kg} \cdot \text{m} \cdot \text{s}^{-2}$   |
| Energy/Work       | Joule   | J      | $\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$ |
| Power             | Watt    | W      | $\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-3}$ |
| Frequency         | Hertz   | Hz     | $\text{s}^{-1}$                                  |
| Electric Charge   | Coulomb | C      | $\text{A} \cdot \text{s}$                        |

These units can all be derived manually, and it is good to know how to! Once you learn acceleration and force, try deriving the units for Newtons yourself!

---

*This is a draft chapter from the Kontinua Project. Please see our website (<https://kontinua.org/>) for more details.*



# Answers to Exercises

## Answer to Exercise 1 (on page 3)

$$4.5 \text{ kWh} \left( \frac{3.6 \times 10^6 \text{ joules}}{1 \text{ kWh}} \right) \left( \frac{1 \text{ calories}}{4.184 \text{ joules}} \right) = \frac{(4.5)(3.6 \times 10^6)}{4.184} = 1.08 \times 10^6 \text{ calories}$$

## Answer to Exercise 2 (on page 4)

$$\frac{0.1 \text{ gallons}}{2 \text{ minutes}} \left( \frac{3.7854 \text{ liters}}{1 \text{ gallons}} \right) \left( \frac{1000 \text{ milliliters}}{1 \text{ liters}} \right) \left( \frac{1 \text{ minutes}}{60 \text{ seconds}} \right) =$$
$$\frac{(0.1)(3.7854)(1000)}{(2)(60)} \text{ ml/second} = 3.1545 \text{ ml/second}$$







---

# INDEX

conversion factors, [2](#)

metric system  
    prefixes, [2](#)

SI units, [4](#)

units table, [1](#)