



CONTENTS

1	Atmospheric Pressure	3
1.1	Altitude and Atmospheric Pressure	5
1.2	How a Drinking Straw Works	6
1.2.1	The Longest Usable Straw	7
1.2.2	Millimeters Mercury	9
1.3	How Siphon Works	10
1.4	How a Toilet Works	12
2	Solving Quadratics	15
2.1	The Traditional Quadratic Formula	17
2.2	Factoring Quadratics	18
3	Complex Numbers	21
3.1	Definition	21
3.2	Why Are Complex Numbers Necessary?	21
3.2.1	Roots of Negative Numbers	21
3.2.2	Polynomial Equations	22
3.2.3	Physics and Engineering	22
3.3	Adding Complex Numbers	22
3.4	Multiplying Complex Numbers	22
4	Introduction to Sequences	23
4.1	Arithmetic sequences	23
4.1.1	Formulas for arithmetic sequences	24

4.2	Geometric sequences	27
4.2.1	Formulas for geometric sequences	28
A	Answers to Exercises	31
Index		33

Atmospheric Pressure

The air you breathe is a blend of gases:

1. 78% nitrogen in the form of N_2
2. 21% oxygen in the form O_2
3. 1% other gases (mostly argon)

If you fill a balloon with helium (He), the helium will push against the interior of the balloon with a certain amount of pressure. The pressure is the same at every point in the interior of the balloon. Pressure, then, is force spread over some area. Force is commonly measured in newtons. Pressure is measured in *pascals*. A pascal is 1 newton per square meter.

We don't usually think about it, but the air outside the balloon is also pushing against the exterior of the balloon. We call this *barometric pressure* or *atmospheric pressure*, and it is caused by gravity pulling on the gas molecules above the balloon.




Figure 1.1: Air pushes both ways on a balloon.

Imagine a square meter on the ground at sea level. Now imagine the column of air above it — reaching all the way to the top of the atmosphere.




Figure 1.2: An air column surrounding a person experiences a large amount of pressure.

The air inside that column has a mass of about 10,340 kg. One kilogram on the earth experiences a gravitational force of 9.8 N. So the atmospheric pressure all around you is about 101,332 pascals. When dealing with such large numbers, we often use kilopascals. We'd say the barometric pressure at sea level is about 101.3 kPa.

That's a lot of pressure! Why doesn't your ribcage collapse crushing your lungs? The air *inside* your lungs is the same pressure as the air pushing on the outside of your rib cage.

This is why we go about our lives relatively oblivious to this huge force that is all around us — but you can see it sometimes. For example, if you suck the air out of a plastic bottle, the bottle will be crushed by the barometric pressure.

1.1 Altitude and Atmospheric Pressure

If you let go of the balloon, as it rises through this column there will be less and less air mass above it, and thus less and less atmospheric pressure on the outside of the balloon.




Figure 1.3: As a balloon rises, it reaches a point of less and less outer pressure.

What would be the atmospheric pressure at h meters above sea level? Here is a handy formula for that:

$$p = 101,332 \times \left(1 - \left(2.25577 \times 10^{-5} \times h\right)\right)^{5.25588}$$

where p is the atmospheric pressure in pascals.

Exercise 1 Atmospheric Pressure

Working Space

You are thinking about riding your bicycle to the top of Mount Everest. You are worried when the atmospheric pressure outside the tire drops, the tire will fail. (I have had a tire fail before; It is very, very loud.)

Calculate the atmospheric pressure at the top of Mount Everest (9,144 meters above sea level).

Answer on Page 31

1.2 How a Drinking Straw Works

When you suck on a drinking straw, why does the beverage rise? It is actually pushed by atmospheric pressure.

Before you put your mouth on the straw, the atmospheric pressure is pressing on the entire surface of the liquid (even inside the straw) evenly. Gravity pulls on the liquid making the surface level.

When you suck some air out of the straw, the pressure on the surface inside the straw drops. The atmospheric pressure on the surface outside the straw pushes into the straw and the beverage rises.





Figure 1.4: You use pressure to counteract gravity in a straw.

Of course, gravity is still trying to pull the liquid inside the straw back down. And for every inch that you lift the liquid in the straw, the force of gravity gets greater, demanding more suction.

1.2.1 The Longest Usable Straw

Assuming you are drinking water in a place with 100 kPa of atmospheric pressure, how high could you suck water with a perfect vacuum? That is, given a very, very long and very, very stiff drinking straw, if you created a pressure of 0 Pa inside, how far above the surface of the glass could you get the water?

Let's say a cross-section of the straw has an area of A square meters, and the very top of the column of water is h meters above the surface in the glass.



With how many newtons of force is the atmosphere pushing the water upward? 100 kPa = 100,000 newtons per square meter. So:

$$F_u = (100,000)a$$

With how many newtons of force is gravity pulling the water in the straw downward? The volume of the water is ah . A cubic meter of liquid water weighs 1000 kg. The force of gravity is 9.8 Newtons per kg.

$$F_d = (ah)(1000)(9.8)$$

The water will stop rising when $F_u = F_d$. So, to find h , we substitute in:

$$(100,000)a = (ah)(1000)(9.8)$$

Notice that we can divide both sides by a getting:

$$h = \frac{100,000}{9,800} = 10.2 \text{ meters}$$

A perfect vacuum would only be able to drag the water up 10.2 meters.


1.2.2 Millimeters Mercury

The density of mercury is 13,500 kg per cubic meter. How far up a straw would a perfect vacuum pull mercury?

$$h = \frac{100,000}{(9.8)(13,500)} \approx 0.756 \text{ meters}$$

When the atmospheric pressure is 100kPa, the mercury will rise 756 mm into a vacuum.

This is actually how scientists measure atmospheric pressure. They have a long glass tube filled with mercury. One end is closed off and pointed into the sky (exactly opposite the direction of gravity). The other end is placed into a dish of mercury. There are millimeter marks on the glass tube.



We use fluctuations in the atmospheric pressure to help us predict the weather. You might

hear a weather nerd with a barometer in their house say, "Wow, the barometer has gone from 752 to 761 millimeters mercury in the last hour. A high-pressure system is moving in."

1.3 How Siphon Works


Let's say you had two cups on a table; one is filled with water, one is empty. And you connected them with an empty U-shaped straw. Water will not crawl up the empty straw — the pressure on each end of the straw is the same, and crawling the straw (against gravity) would require energy.

However, what if the straw were filled with water? Then the force of gravity is pulling the water on each side of the hump in different directions. However, the side going into the empty glass pulls a little harder. That is sufficient to create enough suction to pull the water in the other side up and over the hump.

In turn, this will pull more water. Water will continue to flow from the full glass to the empty one until their surfaces are at the same level. At this point, the pull of gravity is the same on each side of the tube.

This is known as a *siphon*. Notice that atmospheric pressure makes the siphon possible. When the water on one side is pulled down by gravity, the atmospheric pressure pushes the other side up. If you were on a planet with plenty of gravity, but no atmosphere, a siphon wouldn't work.

A siphon is especially useful when you want to get liquid out of a container that is too big to pour. For example, if you wanted to take the gasoline out of a car, you could use any flexible tubing to make a siphon. You would put the hose in the gas tank, suck enough gas up into the hose to get the siphon going, then put the hose into your jug. (If you ever do this, be extra careful not to suck any of the gasoline into your mouth; ingesting even a little bit of gasoline can make you incredibly sick, or even kill you.)



There are two rules to siphons:


- The peak of the siphon needs to be low enough for atmospheric pressure to push the liquid that high. For water at sea-level, for example, the peak of the siphon can't be more than 10.2 meters above the surface of the source liquid.
- The tube has to carry the liquid to a lower level than the surface of the source liquid. If the destination end of the siphon is submerged, the surface of the liquid it is submerged in must be lower than the surface of the source liquid. If the destination

end of the siphon is not submerged, its opening must be lower than the surface of the source liquid.

As long as you follow these two rules, your siphons can be very creative. For example, every toilet has a siphon in it.

1.4 How a Toilet Works

Before we talk about toilets, you should know about P-traps. The drain from every sink, shower, and toilet in your house curves up and then down. This is known as a *P-trap*. The P-trap should always have some water in it. That keeps stinky (and flammable!) gases in the sewer from coming up and into your house.



If one of your fixtures (especially one that hasn't been used in a while) smells like raw sewage, run some water to ensure that the P-trap is full.

Now, back to the toilet! The drain in the bottom of the toilet is connected to a siphon into the sewer. The siphon is filled with air most of the time. However, when you flush the toilet, water rushes from the tank into the bowl, which also fills the siphon. Once the siphon is filled, it pulls water out of the toilet until air starts to enter the siphon. At that point, the water stops flowing.




Figure 1.5: The steps a toilet takes when siphoning out waste.

The toilet tank is pretty simple: it has a float and valve that opens with the float is too low. So, anytime the water-level is too low, the value is open and slowly filling the tank. This means the tank is nearly always filled with a precise amount of water.

When you flush, a small door in the bottom of the tank is opened and the water rushes into the bowl. When the water is out, the door closes again so the tank can refill.

CHAPTER 2

Solving Quadratics

A quadratic function has three terms: $ax^2 + bx + c$. a , b , and c are known as the *coefficients*. The coefficients can be any constant, except that a can never be zero. (If a is zero, it is a linear function, not a quadratic.)

When you have an equation with a quadratic function on one side and a zero on the other, you have a quadratic equation. For example:

$$72x^2 - 12x + 1.2 = 0$$

How can you find the values of x that will make this equation true?

You can always reduce a quadratic equation so that the first coefficient is 1, so that your equation looks like this:

$$x^2 + bx + c = 0$$

For example, if you are asked to solve $4x^2 + 8x - 19 = -2x^2 - 7$

$$4x^2 + 8x - 19 = -2x^2 - 7$$

$$6x^2 + 8x - 12 = 0$$

$$x^2 + \frac{4}{3}x - 2 = 0$$

Here, $b = \frac{4}{3}$ and $c = -2$.

$x^2 + bx + c = 0$ when


$$x = -\frac{b}{2} \pm \frac{\sqrt{b^2 - 4c}}{2}$$

Only when $a = 1$


What does this mean?

For any b and c , the graph of $x^2 + bx + c$ is a parabola that goes up on each end. Its low point is at $x = -\frac{b}{2a}$. This is referred to as the vertex formula.


If there are no real roots ($b^2 - 4c < 0$), which means the parabola never gets low enough to cross the x -axis:



If there is one real root ($b^2 - 4c = 0$), it means that the parabola only touches the x -axis.



If there are two real roots ($b^2 - 4c > 0$), it means that the parabola crosses the x -axis twice as it dips below and then returns:



Exercise 2 Roots of a Quadratic

Working Space

In the last chapter, you found that the function for the height of your flying hammer is:

$$p = -\frac{1}{2}9.8t^2 + 12t + 2$$

At what time will the hammer hit the ground?

Answer on Page 31

2.1 The Traditional Quadratic Formula

If the last explanation was a little tricky to understand, the quadratic formula is a nifty tool.

The Quadratic Formula

$ax^2 + bx + c = 0$ when

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

2.2 Factoring Quadratics

Sometimes, the quadratic gets a bit clunky and hard to solve, especially without a calculator. That's when factoring quadratics becomes a good tool to use as well.

Factoring quadratics is a way to separate a quadratic, usually in the form $x^2 + bx + c$ where ($a = 1$), into two multiplied equations in the form $(px + q)(rx + s) = 0$. Solving these for x also gives the roots of the equation.

Types of Factoring Techniques

1. Factoring out the GCF (Greatest Common Factor)

Always check for a common factor first:

$$2x^2 + 4x = 2x(x + 2)$$

2. Factoring Trinomials: $a = 1$

For expressions like $x^2 + bx + c$, find two numbers that:

- Multiply to c
- Add to b

$$x^2 + 5x + 6 = (x + 2)(x + 3)$$

3. Factoring Trinomials: $a \neq 1$

Use the "AC method":

- (a) Multiply $a \times c$
- (b) Find two numbers that multiply to ac and add to b
- (c) Split the middle term and factor by grouping

$$6x^2 + 11x + 4 = 6x^2 + 3x + 8x + 4 = (3x + 2)(2x + 2)$$

4. Special Cases:

- Perfect Square Trinomials:

$$a^2 + 2ab + b^2 = (a + b)^2$$

$$x^2 + 6x + 9 = (x + 3)^2$$

- Difference of Squares:

$$a^2 - b^2 = (a - b)(a + b)$$

$$x^2 - 16 = (x - 4)(x + 4)$$

Tips

- Always factor out the GCF first.

- Check your result by expanding (FOIL). The inverse of factoring is expanding using the FOIL method.
- Not all quadratics are factorable over real integers.

Complex Numbers

Complex numbers are an extension of real numbers, which in turn are an extension of rational numbers. In mathematics, the set of complex numbers is a number system that extends the real number line to a full two dimensions, using the imaginary unit, which is denoted by i , with the property that $i^2 = -1$.

An imaginary number is a number that is a multiple of the imaginary unit i . For example, $5i$ and $-3i$ are both imaginary numbers. While imaginary numbers are a subset of complex numbers (with real part equal to zero), not all complex numbers are imaginary. Complex numbers can have both a real and an imaginary part, while imaginary numbers specifically refer to those with no real part.

3.1 Definition

A complex number is a number of the form $a + bi$, where a and b are real numbers, and i is the imaginary unit, with the property that $i^2 = -1$. The real part of the complex number is a , and the imaginary part is bi .

3.2 Why Are Complex Numbers Necessary?

Complex numbers are essential to many fields of science and engineering. Here are a few reasons why:

3.2.1 Roots of Negative Numbers

In the real number system, the square root of a negative number does not exist, because there is no real number that you can square to get a negative number. The introduction of the imaginary unit i , which has the property that $i^2 = -1$, allows us to take square roots of negative numbers and gives rise to complex numbers.

3.2.2 Polynomial Equations

The fundamental theorem of algebra states that every non-constant polynomial equation with complex coefficients has a complex root. This theorem guarantees that polynomial equations of degree n always have n roots in the complex plane.

3.2.3 Physics and Engineering

In physics and engineering, complex numbers are used to represent waveforms in control systems, in quantum mechanics, and many other areas. Their properties make many mathematical manipulations more convenient.

3.3 Adding Complex Numbers

The addition of complex numbers is straightforward. If we have two complex numbers $z_1 = a + bi$ and $z_2 = c + di$, their sum is defined as:

$$z_1 + z_2 = (a + c) + (b + d)i \quad (3.1)$$

In other words, you add the real parts to get the real part of the sum, and add the imaginary parts to get the imaginary part of the sum.

3.4 Multiplying Complex Numbers

The multiplication of complex numbers is a bit more involved. If we have two complex numbers $z_1 = a + bi$ and $z_2 = c + di$, their product is defined as:

$$z_1 \cdot z_2 = (a + bi) \cdot (c + di) = ac + adi + bci - bd = (ac - bd) + (ad + bc)i \quad (3.2)$$

Note the last term comes from $i^2 = -1$. You multiply the real parts and the imaginary parts just as you would in a binomial multiplication, and remember to replace i^2 with -1 . See the Khan academy video in the digital resources for a more in-depth explanation.

Introduction to Sequences

A sequence is a list of numbers in a particular order. $\{1, 3, 5, 7, 9\}$ is a sequence. So is $\{\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\}$. There are many types of sequences. We will present two of the most common types in this chapter: arithmetic and geometric sequences.

Sequences are generally represented like this:

$$a_1, a_2, a_3, a_4, \dots, a_n, \dots$$

The first number, a_1 , is called the *first term*, a_2 is the *second term*, and a_n is the *nth term*. A sequence can be finite or infinite. If the sequence is infinite, we represent that with ellipses ($\dots$) at the end of the list, to indicate that there are more numbers.

We can also write formulas to represent a sequence. Take the first example, the finite sequence $\{1, 3, 5, 7, 9\}$. Notice that each term is two more than the previous term. We can define the sequence *recursively* by defining the n^{th} term as a function of the $(n - 1)^{\text{th}}$ term. In our example, we see that $a_n = a_{n-1} + 2$ with $a_1 = 1$ for $1 \leq n \leq 5$. This is called a recursive formula, because you have to already know the $(n - 1)^{\text{th}}$ term to find the n^{th} term.

Another way to write a formula for a sequence is to find a rule for the n^{th} term. In our example sequence, the first term is 1 plus 0 times 2, the second term is 1 plus 1 times 2, the third term is 1 plus 2 times 2, and so on. Did you notice the pattern? The n^{th} term is 1 plus $(n-1)$ times 2. We can write this mathematically:

$$a_n = 1 + 2(n - 1) \text{ for } 1 \leq n \leq 5$$

This is called the *explicit* formula because each term is explicitly defined. Notice that for the second way of writing a formula, we don't have to state what the first term is — the formula tells us.

4.1 Arithmetic sequences

Our first example sequence, $\{1, 3, 5, 7, 9\}$ is a *finite, arithmetic* sequence. We know it is finite because there is a limited number of terms in the sequence (in this case, 5). How do we know it is arithmetic?

An arithmetic sequence is one where you add the same number every time to get the next term. Our example is an arithmetic sequence because you add 2 to get the next term every time. That number that you add is called the *common difference*, so we can say the sequence $\{1, 3, 5, 7, 9\}$ has a common difference of 2. The common difference can be positive (in the case of an increasing arithmetic sequence) or negative (in the case of a decreasing arithmetic sequence). Formally, we can find the common difference of an arithmetic sequence by subtracting the $(n - 1)^{\text{th}}$ term from the n^{th} term:

$$d = a_n - a_{n-1}$$

Exercise 3

Which of the following are arithmetic sequences? For the arithmetic sequences, find the common difference.

1. $\{\frac{1}{2}, \frac{1}{4}, \frac{1}{6}, \frac{1}{8}, \dots\}$
2. $\{5, 8, 11, 14, 17, \dots\}$
3. $\{3, -1, -5, -9, \dots\}$
4. $\{-1, 2, -3, 4, -5, 6, \dots\}$

Working Space

Answer on Page 31

4.1.1 Formulas for arithmetic sequences

If you are given an arithmetic sequence, you can write an explicit or recursive formula. You can think of the formula as a function where the domain (input) is restricted to integers greater than or equal to one. Let's write explicit and recursive formulas for the sequence $\{3, -1, -5, -9, \dots\}$.

For either type of formula, we need to identify the common difference. Since each term is 4 less than the previous term, the common difference is -4 (see figure 4.1). This means the n^{th} term is the $(n - 1)^{\text{th}}$ term minus 4. The general form of a recursive formula is $a_n = a_{n-1} + d$, where d is the common difference. For our example, the common difference is -4, so we can write a recursive formula:

$$a_n = a_{n-1} - 4$$

However, this formula doesn't tell us what a_1 is! For recursive formulas, you have to specify the first term in the sequence. So, the *complete* recursive formula for the sequence is:

$$a_n = a_{n-1} - 4$$

$$a_1 = 3$$




Figure 4.1: The common difference in the sequence $\{3, -1, -5, -9, \dots\}$ is -4

Recursive formulas make it easy to see how each term is related to the next term. However, it is difficult to use recursive formulas to find a specific term. Say we wanted to know the 7th term in the sequence. Well, from the formula, we know that:

$$a_7 = a_6 - 4$$

What is a_6 ? Again, we see that

$$a_6 = a_5 - 4$$

Now we have to find a_5 ! If we keep going, we see that:

$$a_5 = a_4 - 4$$

$$a_4 = a_3 - 4$$

$$a_3 = a_2 - 4$$

$$a_2 = a_1 - 4$$

Since we were told a_1 , we can find a_2 and propagate our terms back up the chain to find a_7 :

$$a_2 = 3 - 4 = -1$$

$$a_3 = a_2 - 4 = -1 - 4 = -5$$

$$a_4 = a_3 - 4 = -5 - 4 = -9$$

$$a_5 = a_4 - 4 = -9 - 4 = -13$$

$$a_6 = a_5 - 4 = -13 - 4 = -17$$

$$a_7 = a_6 - 4 = -17 - 4 = -21$$

Ultimately, we see that $a_7 = -21$. That was a lot of work! You can imagine that for higher n terms, such as the 100^{th} or 1000^{th} term, this method becomes cumbersome. This is where the explicit formula is more useful.

The general form of an explicit formula for an arithmetic sequence is

$$a_n = a_1 + d \times (n - 1)$$

where d is the common difference. For our example sequence, $\{3, -1, -5, -9, \dots\}$, the common difference is -4 . So the explicit formula is

$$a_n = 3 + (-4)(n - 1) = 3 - 4(n - 1)$$

You may be tempted to distribute and simplify, which is fine and yields an equivalent formula:

$$a_n = 7 - 4n$$

Now, to find the 7^{th} term, all we have to do is substitute $n = 7$:

$$a_7 = 3 - 4(7 - 1) = 3 - 4(6) = 3 - 24 = -21$$

We get the same answer with much less effort!

Another example of a recursive sequence is the Fibonacci sequence which we will cover in-depth later

Exercise 4

An arithmetic sequence is defined by the recursive formula $a_n = a_{n-1} + 5$ with $a_1 = -4$. Write the first 5 terms of the sequence and determine an explicit formula for the same sequence.

Working Space

Answer on Page 32

Exercise 5

The first four terms of an arithmetic sequence are $\{\pi, \frac{3\pi}{2}, 2\pi, \frac{5\pi}{2}\}$. What is the common difference? Write explicit and recursive formulas for the infinite sequence.

Working Space

Answer on Page 32

4.2 Geometric sequences

Let's look at the other sequence given as an example at the beginning of the chapter: $\{\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\}$. How is each term related to the previous term? Well, $\frac{1}{4}$ is half of $\frac{1}{2}$, and $\frac{1}{8}$ is half of $\frac{1}{4}$, so each term is the previous term multiplied by $\frac{1}{2}$. When each term in a sequence is a multiple of the previous term, this is a *geometric* sequence. The number we multiply by each time (in our example, this is $\frac{1}{2}$), which is called the *common ratio*. The common ratio can be positive or negative, but not zero.

An easy way to determine the common ratio (r) is to divide the n^{th} term by the $(n-1)^{\text{th}}$ term. In our example sequence, the first term is $\frac{1}{2}$ and the second is $\frac{1}{4}$.

$$r = \frac{a_2}{a_1} = \frac{1/4}{1/2} = \frac{1}{2}$$

which returns the common ratio we already identified, $r = \frac{1}{2}$.

If the common ratio is negative, then the sequence will "flip" back and forth from positive to negative. For example, suppose there is a geometric sequence such that $a_1 = 1$ and $r = -2$. Then the first 5 terms are $\{1, -2, 4, -8, 16\}$. Whenever you see a sequence going back and forth from positive to negative, that means the common ratio is negative.

For positive ratios ($r > 1$), the sequence is increasing. And for negative ratios, ($r < 1$), the sequence is decreasing.

4.2.1 Formulas for geometric sequences

Like arithmetic sequences, we can write recursive and explicit formulas. For geometric sequences, the recursive formula has the general form:

$$a_n = r(a_{n-1})$$

where r is the common ratio and a_1 is specified. For our example sequence, $\{\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\}$, the recursive formula is:

$$a_n = \frac{1}{2}a_{n-1}$$
$$a_1 = \frac{1}{2}$$

In a geometric sequence, each term is the first term, a_1 , multiplied by the common ratio, r , $n - 1$ times. Therefore, the general form of an explicit formula for a geometric function is:

$$a_n = (a_1)r^{n-1}$$

Again, for our example sequence, $a_1 = \frac{1}{2}$ and $r = \frac{1}{2}$, so the explicit formula is:

$$a_n = (\frac{1}{2})(\frac{1}{2})^{(n-1)}$$

Exercise 6

Which of the following are geometric sequences? For each geometric sequence, determine the common ratio.

1. $\{2, -4, 6, -8, \dots\}$
2. $\{4, 2, 1, \frac{1}{2}, \dots\}$
3. $\{-5, 25, -125, 525, \dots\}$
4. $\{2, 0, -2, -4, \dots\}$

Working Space

Answer on Page 32

Exercise 7

A geometric sequence is defined by the recursive formula $a_n = a_{n-1} \times \frac{3}{2}$ with $a_1 = 1$. Write the first five terms of the sequence and determine an explicit formula for the same sequence.

*Working Space**Answer on Page 32***Exercise 8**

The first four terms of a geometric sequence are $\{-4, 2, -1, \frac{1}{2}\}$. What is the common ratio? Write recursive and explicit formulas for the infinite sequence.

*Working Space**Answer on Page 32*

Answers to Exercises

Answer to Exercise 1 (on page 6)

$$p = 101,332 \times \left(1 - 2.25577 \times 10^{-5} \times h\right)^{5.25588}$$

and $h = 9,144$. Thus,

$$p \approx 30.1 \text{ kPa}$$

Answer to Exercise 2 (on page 17)

For what t is $-4.9t^2 + 12t + 2 = 0$? Start by dividing both sides of the equation by -4.9 .

$$t^2 - 2.45t - 0.408 = 0$$

The roots of this are at

$$x = -\frac{b}{2} \pm \frac{\sqrt{b^2 - 4c}}{2} = -\frac{-2.45}{2} \pm \frac{\sqrt{(-2.45)^2 - 4(-0.408)}}{2} = 1.22 \pm 1.36$$

We only care about the root after we release the hammer ($t > 0$).

$1.22 + 1.36 = 2.58$ seconds after releasing the hammer, it will hit the ground.

Answer to Exercise 3 (on page 24)

1. not arithmetic
2. arithmetic, common difference is 3

3. arithmetic, common difference is -4
4. not arithmetic

Answer to Exercise 4 (on page 26)

The first five terms are $\{-4, 1, 6, 11, 16\}$ and an explicit formula is $a_n = -4 + 5(n - 1)$.

Answer to Exercise 5 (on page 27)

The common difference is $\frac{3\pi}{2} - \pi = \frac{\pi}{2}$. The recursive formula is $a_n = a_{n-1} + \frac{\pi}{2}$ with $a_1 = \pi$. The explicit formula is $a_n = \pi + \frac{\pi}{2}(n - 1)$.

Answer to Exercise 6 (on page 28)

1. not geometric
2. geometric sequence with common ratio $r = \frac{1}{2}$
3. geometric sequence with common ratio $r = -5$
4. not geometric

Answer to Exercise 7 (on page 29)

The first five terms are $\{1, \frac{3}{2}, \frac{9}{4}, \frac{27}{8}, \frac{81}{16}\}$. An explicit formula for this sequence is $a_n = 1(\frac{3}{2})^{(n-1)}$.

Answer to Exercise 8 (on page 29)

The common ratio is $\frac{a_n}{a_{n-1}} = \frac{2}{-4} = -\frac{1}{2}$. A recursive formula would be $a_n = a_{n-1} \times -\frac{1}{2}$ with $a_1 = -4$. An explicit formula would be $a_n = (-4)(-\frac{1}{2})^{(n-1)}$.



INDEX

atmospheric pressure, [3](#)

barometric pressure, [3](#)

Complex Numbers, [21](#)

millimeters mercury, [9](#)

P-trap, [12](#)

pascals, [3](#)

pressure, [3](#)

siphon, [10](#)

straw

 drinking, [6](#)

Vertex Formula, [15](#)