

Linear Algebra I HW1

B13902024 張沂魁

September 7, 2025

Problem 0.0.1. Let W be a set of all $(x_1, x_2, x_3, x_4, x_5)$ in \mathbb{R}^5 which satisfy

$$\begin{array}{rrrrr} 2x_1 - x_2 & +\frac{4}{3}x_3 - x_4 & & & = 0 \\ x_1 & +\frac{2}{3}x_3 & & -x_5 & = 0 \\ 9x_1 - 3x_2 & +6x_3 - 3x_4 & -3x_5 & & = 0 \end{array}$$

Find a finite set of vectors which spans W .

Proof. We can first write the system of equations into the matrix form and then use Gaussian elimination.

$$\begin{pmatrix} 2 & -1 & \frac{4}{3} & -1 & 0 & 0 \\ 1 & 0 & \frac{2}{3} & 0 & -1 & 0 \\ 9 & -3 & 6 & -3 & -3 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -\frac{1}{2} & \frac{2}{3} & -\frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 & \frac{1}{2} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Hence, we can need to solve

$$\begin{cases} x_1 - \frac{1}{2}x_2 + \frac{2}{3}x_3 - \frac{1}{2}x_4 = 0 \\ \frac{1}{2}x_2 + \frac{1}{2}x_4 - x_5 = 0. \end{cases}$$

So we know $(x_1, x_2, x_3, x_4, x_5) = (t - \frac{2}{3}a, b, a, 2t - b, t)$ for some $a, b, t \in \mathbb{R}^5$, and thus we know the set

$$S = \left\{ (1, 0, 0, 2, 1), \left(-\frac{2}{3}, 0, 1, 0, 0\right), (0, 1, 0, -1, 0) \right\}$$

spans W . ■

Problem 0.0.2. Prove that a subspace of \mathbb{R}^2 is \mathbb{R}^2 , or the zero subspace, or consists of all scalar multiples of some fixed vector in \mathbb{R}^2 . (The last type of subspace is, intuitively, a straight line through the origin.)

Proof. We first give a claim:

Claim. Suppose V is a vector space, and if W is a subspace of V , then $\dim W \leq \dim V$.

Proof. Since $W \subseteq V$, so suppose $k = \dim V$ and B_1 is a basis of V , then $W \subseteq V = \text{span } B_1$, which means if there is a basis of W , say B_2 , then $|B_2| \leq |B_1|$, which means $\dim W \leq \dim V$. ⊗

Now also we know $\dim \mathbb{R}^2 = 2$ since $\{(0, 1), (1, 0)\}$ is a linearly independent set and spans \mathbb{R}^2 . Thus, if there is a subspace of \mathbb{R}^2 , say W , then $\dim W = 0, 1, 2$. If $\dim W = 0$, then W is the zero subspace. If $\dim W = 1$, then W consists of all scalar multiples of some fixed vector in \mathbb{R}^2 . If $\dim W = 2$, then we can give a claim first:

Claim. Suppose $W \subseteq V$ and they are both vector spaces, then if $\dim V = \dim W$, then $V = W$.

Proof. Suppose by contradiction, there exists $v \in V \setminus W$, and suppose B is a basis of W , then we know $B \cup \{v\}$ is linearly independent in V . However, $|B \cup \{v\}| > \dim V$, which is a contradiction. ⊗

By this claim, we know $W = \mathbb{R}^2$ if $\dim W = 2$. ■

Problem 0.0.3. Let W_1 and W_2 be subspaces of a vector space V such that the set-theoretic union of W_1 and W_2 is also a subspace. Prove that one of the spaces W_i is contained in the other.

Proof. Suppose each W_i is not contained in the other, then there exists u, v s.t.

$$u \in W_2 \setminus W_1 \quad v \in W_1 \setminus W_2.$$

Thus, we know $u + v \in W_1 \cup W_2$ since $W_1 \cup W_2$ is a vector space. Also, we know $u + v \in W_1$ or $u + v \in W_2$. Now if $u + v \in W_1$, then $u = u + v + (-v) \in W_1$, which is a contradiction, and if $u + v \in W_2$, we have $v = u + v + (-u) \in W_2$, which is also a contradiction. Hence, we know either $W_1 \subseteq W_2$ or $W_2 \subseteq W_1$ should happen. ■

Problem 0.0.4. Let V be the vector space of all functions from \mathbb{R} into \mathbb{R} ; let V_e be the subset of even functions, $f(-x) = f(x)$; let V_o be the subset of odd functions, $f(-x) = -f(x)$.

- (a) Prove that V_e and V_o are subspaces of V .
- (b) Prove that $V_e + V_o = V$.
- (c) Prove that $V_e \cap V_o = \{0\}$.

Proof.

- (a) First note that $V_e \subseteq V$ and $V_o \subseteq V$, then

- For all $f, g \in V_e$ and $\alpha \in F$, we define $h(x) = \alpha f(x) + g(x)$, and we know $h \in V_e$ since

$$h(-x) = \alpha f(-x) + g(-x) = \alpha f(x) + g(x) = h(x).$$

Hence, V_e is a subspace of V .

- For all $f, g \in V_o$ and $\alpha \in F$, we define $h(x) = \alpha f(x) + g(x)$, and we know $h \in V_o$ since

$$h(-x) = \alpha f(-x) + g(-x) = -\alpha f(x) - g(x) = -h(x).$$

Hence, V_o is a subspace of V .

- (b) We first show that $V \subseteq V_e + V_o$. Since for all $f \in V$, we know

$$f(x) = \left(\frac{f(x) + f(-x)}{2} \right) + \left(\frac{f(x) - f(-x)}{2} \right),$$

where

$$f_e(x) = \left(\frac{f(x) + f(-x)}{2} \right) \in V_e \quad f_o(x) = \left(\frac{f(x) - f(-x)}{2} \right) \in V_o$$

since

$$\begin{aligned} f_e(-x) &= \left(\frac{f(-x) + f(x)}{2} \right) = f_e(x) \\ f_o(-x) &= \left(\frac{f(-x) - f(x)}{2} \right) = -f_o(x). \end{aligned}$$

Now we show that $V_e + V_o \subseteq V$. This is trivial since for all $f \in V_e + V_o$ we have $f = g + h$ for some $g \in V_e \subseteq V$ and $h \in V_o \subseteq V$, so $f = g + h \in V$ since V is a vector space. Hence, $V_e + V_o \subseteq V$.

Now we have $V \subseteq V_e + V_o$ and $V_e + V_o \subseteq V$, so we can conclude that $V = V_e + V_o$.

- (c)

■

Problem 0.0.5. Let W_1 and W_2 be subspaces of a vector space V such that $W_1 + W_2 = V$ and $W_1 \cap W_2 = \{0\}$. Prove that for each vector α in V there are unique vectors α_1 in W_1 and α_2 in W_2 such that $\alpha = \alpha_1 + \alpha_2$.

Proof. Since $W_1 + W_2 = V$, so we know for each vector $\alpha \in V$, it can be representend as $\alpha = \alpha_1 + \alpha_2$ for some $\alpha_1 \in W_1$ and $\alpha_2 \in W_2$. If there are two differet (α_1, α_2) -pairs to represent α , say

$\alpha = \alpha_1 + \alpha_2 = \alpha'_1 + \alpha'_2$ where $\alpha_1, \alpha'_1 \in W_1$ and $\alpha_2, \alpha'_2 \in W_2$, then we know $\alpha_1 - \alpha'_1 = \alpha'_2 - \alpha_2$. However, since $\alpha_1 - \alpha'_1 \in W_1$ and $\alpha'_2 - \alpha_2 \in W_2$, so $\alpha_1 - \alpha'_1 = \alpha'_2 - \alpha_2 \in W_1 \cap W_2 = \{0\}$, which means $\alpha_1 = \alpha'_1$ and $\alpha_2 = \alpha'_2$ and thus the (α_1, α_2) -pair is unique. ■