

Linear Algebra I HW1

B13902024 張沂魁

September 6, 2025

Problem 0.0.1. Let W be a set of all $(x_1, x_2, x_3, x_4, x_5)$ in \mathbb{R}^5 which satisfy

$$\begin{array}{cccccc} 2x_1 - x_2 & +\frac{4}{3}x_3 - x_4 & & & & = 0 \\ x_1 & +\frac{2}{3}x_3 & & -x_5 & & = 0 \\ 9x_1 - 3x_2 & +6x_3 - 3x_4 & -3x_5 & & & = 0 \end{array}$$

Find a finite set of vectors which spans W .

Proof. We can first write the system of equations into the matrix form and then use Gaussian elimination.

$$\begin{pmatrix} 2 & -1 & \frac{4}{3} & -1 & 0 & 0 \\ 1 & 0 & \frac{2}{3} & 0 & -1 & 0 \\ 9 & -3 & 6 & -3 & -3 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -\frac{1}{2} & \frac{2}{3} & -\frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 & \frac{1}{2} & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Hence, we can need to solve

$$\begin{cases} x_1 - \frac{1}{2}x_2 + \frac{2}{3}x_3 - \frac{1}{2}x_4 = 0 \\ \frac{1}{2}x_2 + \frac{1}{2}x_4 - x_5 = 0. \end{cases}$$

So we know $(x_1, x_2, x_3, x_4, x_5) = (t - \frac{2}{3}a, b, a, 2t - b, t)$ for some $a, b, t \in \mathbb{R}$, and thus we know the set

$$S = \left\{ (1, 0, 0, 2, 1), \left(-\frac{2}{3}, 0, 1, 0, 0\right), (0, 1, 0, -1, 0) \right\}$$

spans W . ■

Problem 0.0.2. Prove that a subspace of \mathbb{R}^2 is \mathbb{R}^2 , or the zero subspace, or consists of all scalar multiples of some fixed vector in \mathbb{R}^2 . (The last type of subspace is, intuitively, a straight line through the origin.)

Proof. We first give a claim:

Claim. Suppose V is a vector space, and if W is a subspace of V , then $\dim W \leq \dim V$.

Proof. Since $W \subseteq V$, so suppose $k = \dim V$ and B_1 is a basis of V , then $W \subseteq V = \text{span} B_1$, which means if there is a basis of W , say B_2 , then $|B_2| \leq |B_1|$, which means $\dim W \leq \dim V$. ⊗

Now also we know $\dim \mathbb{R}^2 = 2$ since $\{(0, 1), (1, 0)\}$ is a linearly independent set and spans \mathbb{R}^2 . Thus, if there is a subspace of \mathbb{R}^2 , say W , then $\dim W = 0, 1, 2$. If $\dim W = 0$, then W is the zero subspace. If $\dim W = 1$, then W consists of all scalar multiples of some fixed vector in \mathbb{R}^2 . If $\dim W = 2$, then we can give a claim first:

Claim. Suppose $W \subseteq V$ and they are both vector spaces, then if $\dim V = \dim W$, then $V = W$.

Proof. Suppose by contradiction, there exists $v \in V \setminus W$, and suppose B is a basis of W , then we know $B \cup \{v\}$ is linearly independent in V . However, $|B \cup \{v\}| > \dim V$, which is a contradiction. ⊗

By this claim, we know $W = \mathbb{R}^2$ if $\dim W = 2$. ■