

Introduction to Analysis II

Kon Yi

May 2, 2026

Abstract

Lecture note of Introduction to Analysis II.

Contents

1	Several Variable Differential Calculus	2
1.1	Linear Transformation	3
1.2	Derivatives in Several Variable Calculus	5
1.3	Partial and Directional Derivatives	8
1.4	Chain Rule	14
1.5	Double Derivatives and Clairaut's theorem	20
1.6	The Contraction Mapping Theorem	21
1.7	Inverse Function Theorem	24
1.8	The Implicit Function Theorem	28
2	Lebesgue measure	33
2.1	The goal: Lebsgue measure	33
2.2	First attempt: Outer Measure	34
2.3	Outer measure is not additive	39
2.4	Measurable sets	42
2.5	Measurable Functions	48
3	Lebesgue Integration	54
3.1	Simple Functions	56
3.2	Integration of non-negative measurable functions	61
3.3	Integration of Absolutely Integrable Functions	70
3.4	Comparison with the Riemann Integral	76
A	Midterm Practice	79
A.1	HW6	79
A.2	HW1	80
A.3	HW2	80
A.4	HW3	80
A.5	HW4	80
A.6	HW5	81
A.7	Random Stuff	81
B	Additional proof	82

Chapter 1

Several Variable Differential Calculus

Lecture 1

In this chapter, we want to approximate non-linear functions by linear maps. If we consider

24 Feb.

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad f(\underbrace{x_1, x_2, \dots, x_n}_x) = (f_1(x), f_2(x), \dots, f_m(x)),$$

where $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ for all i . Now given a real-valued function $F : \mathbb{R}^n \rightarrow \mathbb{R}$ which is differentiable at a point x_0 . We know that near $x_0 \in \mathbb{R}^n$ we can approximate $F(x)$ in the following way:

$$F(x) \approx F(x_0) + \nabla F(x_0) \cdot (x - x_0)$$

where

$$\nabla F(x_0) = \left(\frac{\partial F(x_0)}{\partial x_1}, \frac{\partial F(x_0)}{\partial x_2}, \dots, \frac{\partial F(x_0)}{\partial x_n} \right) \in \mathbb{R}^n \text{ with } x_0 = (x_1, x_2, \dots, x_n)$$

and thus

$$\begin{aligned} \nabla F(x_0) \cdot (x - x_0) &= \left\langle \frac{\partial F(x_0)}{\partial x_1}, \frac{\partial F(x_0)}{\partial x_2}, \dots, \frac{\partial F(x_0)}{\partial x_n} \right\rangle \cdot \langle x_1, x_2, \dots, x_n \rangle \\ &= \sum_{i=1}^n \frac{\partial F(x_0)}{\partial x_i} x_i. \end{aligned}$$

Hence,

$$f(x) = \begin{pmatrix} f_1(x_0) \\ f_2(x_0) \\ \vdots \\ f_n(x_0) \end{pmatrix} \approx \begin{pmatrix} f_1(x_0) + \nabla f_1(x)(x - x_0) \\ f_2(x_0) + \nabla f_2(x)(x - x_0) \\ \vdots \\ f_n(x_0) + \nabla f_n(x)(x - x_0) \end{pmatrix},$$

which gives

$$f(x) - f(x_0) \approx \begin{pmatrix} \nabla f_1(x_0)(x - x_0) \\ \nabla f_2(x_0)(x - x_0) \\ \vdots \\ \nabla f_n(x_0)(x - x_0) \end{pmatrix} = \begin{pmatrix} \nabla f_1(x_0) \\ \nabla f_2(x_0) \\ \vdots \\ \nabla f_n(x_0) \end{pmatrix} \cdot \underbrace{(x - x_0)}_{\text{column vector}}.$$

1.1 Linear Transformation

Definition 1.1.1 (Row vectors). Let $n \geq 1$ be an integer. We refer to elements of $\mathbb{R}^n$ as n -dimensional row vectors. A typical row vector is $x = (x_1, x_2, \dots, x_n)$ which we abbreviate as $(x_i)_{1 \leq i \leq n}$. The components $x_1, x_2, \dots, x_n$ are real numbers. If x and y are two row vectors in $\mathbb{R}^n$, we can define vector sum by

$$x + y = (x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) := (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n).$$

If $c \in \mathbb{R}$ is any real number, we define scalar multiplications by

$$cx = c(x_1, x_2, \dots, x_n) = (cx_1, cx_2, \dots, cx_n).$$

Remark 1.1.1.

- (1) $-x := (-1) \cdot x = (-x_1, -x_2, \dots, -x_n)$.
- (2) zero vector is denoted by 0 , i.e. $(0, 0, \dots, 0)$.

Lemma 1.1.1 ($\mathbb{R}^n$ is a vector space). Let x, y, z be vectors in $\mathbb{R}^n$, and let $c, d \in \mathbb{R}$. Then the following properties hold:

- (a) $x + y = y + x$.
- (b) $(x + y) + z = x + (y + z)$.
- (c) $x + 0 = 0 + x = x$.
- (d) $x + (-x) = (-x) + x = 0$.
- (e) $(c \cdot d)x = c \cdot (dx)$.
- (f) $c(x + y) = cx + cy$.
- (g) $(c + d)x = cx + dx$.
- (h) $1x = x$.

Definition 1.1.2. Let $x = (x_1, x_2, \dots, x_n)$ be row vector. Its transpose is the n -dimensional column vector

$$x^T = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}.$$

Definition 1.1.3. The standard basis of $\mathbb{R}^n$ consists of $e_1, e_2, \dots, e_n$, where e_j has 1 in the j -th position and 0 elsewhere:

$$e_j = (0, \dots, 0, \underbrace{1}_{j\text{-th}}, 0, \dots, 0).$$

Every row vector

$$x = (x_1, x_2, \dots, x_n) = \sum_{j=1}^n x_j e_j.$$

Similarly,

$$x^T = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \sum_{j=1}^n x_j e_j^T.$$

Definition 1.1.4 (Linear transformation). A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is any function from one Euclidean space to another that satisfies the following two properties:

- (a) Additivity: For $x, y \in \mathbb{R}^n$, $T(x + y) = T(x) + T(y)$.
- (b) Homogeneity: For $x \in \mathbb{R}^n$ and all scalars $c \in \mathbb{R}$, $T(cx) = cT(x)$.

Remark 1.1.2. This definition is equivalent to the following:

$$T(c_1v_1 + \cdots + c_kv_k) = c_1T(v_1) + \cdots + c_kT(v_k)$$

where $v_1, \dots, v_k \in \mathbb{R}^n$ and $c_1, c_2, \dots, c_n \in \mathbb{R}$.

Definition 1.1.5. Let $m, n \geq 1$ be integers. An $m \times n$ ordered matrix is an ordered rectangular array of real numbers

$$A = (a_{ij})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

consisting of m rows and n columns, where

- (a) The entry a_{ij} denote the number in the i -th row and j -th column.
- (b) We denote the set of all $m \times n$ matrices by $\mathbb{R}^{m \times n}$.
- (c) In particular, a row vector is a $1 \times n$ matrix, a column vector is a $n \times 1$ vector.

Definition 1.1.6 (Matrix multiplication). Given an $m \times n$ matrix $A = (a_{ij})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}}$ and an $n \times p$ matrix $B = (b_{jk})_{\substack{1 \leq j \leq n \\ 1 \leq k \leq p}}$, we define AB to be the $m \times p$ matrix $(c_{ik})_{\substack{1 \leq i \leq m \\ 1 \leq k \leq p}}$ where

$$c_{ik} = \sum_{j=1}^n a_{ij}b_{jk}.$$

Definition 1.1.7 (Matrix-vector multiplication). Let $A = (a_{ij}) \in \mathbb{R}^{m \times n}$ and $x = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n$ be a column vector. We define

$$Ax = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{pmatrix}$$

Remark 1.1.3. In our class, we just treat $\mathbb{R}^n, \mathbb{R}^m$ as column vector spaces, and $L_A(x) = Ax$ is a $m \times 1$ column vector.

Theorem 1.1.1. Let A be a $m \times n$ matrix, then $L_A(x) = Ax$ is a linear transformation from $\mathbb{R}^n$ to $\mathbb{R}^m$.

Proof. ■

DIY

Proposition 1.1.1. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. For each $j = 1, 2, \dots, n$, let e_j denote the j -th standard basis vector in $\mathbb{R}^n$ and write $T(e_j) = \begin{pmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{pmatrix}$. Define the matrix $A = (a_{ij})$, then $T(x) = Ax$.

Proof. Let $x = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}$. We can write $x = \sum_{j=1}^n x_j e_j$, then we know

$$T(x) = T\left(\sum_{j=1}^n x_j e_j\right) = \sum_{j=1}^n x_j T(e_j) = \sum_{j=1}^n x_j \begin{pmatrix} a_{1j} \\ \vdots \\ a_{mj} \end{pmatrix} = Ax.$$

■

Lemma 1.1.2. Let A be a $m \times n$ matrix and let B be a $n \times p$ matrix. Then $L_A \circ L_B = L_{(AB)}$.

Proof. It suffices to show that $(L_A \circ L_B)(x) = L_{AB}(x)$ for $x \in \mathbb{R}^p$, and the rest is easy. ■

As previously seen. $f : E \rightarrow \mathbb{R}$ where E is a subset of $\mathbb{R}$, then

$$f'(x_0) = \lim_{\substack{x \rightarrow x_0 \\ x \in E \setminus \{x_0\}}} \frac{f(x) - f(x_0)}{x - x_0}.$$

Suppose now $f : E \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$, we can't define

$$f'(x) = \lim_{\substack{x \rightarrow x_0 \\ x \in E \setminus \{x_0\}}} \frac{f(x) - f(x_0)}{x - x_0}$$

since the denominator and the numerator are vectors in $\mathbb{R}^n$ and $\mathbb{R}^m$.

1.2 Derivatives in Several Variable Calculus

Lemma 1.2.1. Let $E \subseteq \mathbb{R}$, let $f : E \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$ and x_0 be a limit point of E . Then the following two statements are equivalent:

(a) f is differentiable at x_0 and $f'(x_0) = L$.

(b) $\lim_{\substack{x \rightarrow x_0 \\ x \in E \setminus \{x_0\}}} \frac{|f(x) - f(x_0) - L(x - x_0)|}{|x - x_0|} = 0$.

Proof. Note that

$$\frac{f(x) - f(x_0)}{x - x_0} = L + \frac{f(x) - f(x_0) - L(x - x_0)}{x - x_0} \text{ if } x \neq x_0,$$

so we have

$$\frac{f(x) - f(x_0)}{x - x_0} - L = \frac{f(x) - f(x_0) - L(x - x_0)}{x - x_0} \text{ if } x \neq x_0,$$

and thus

$$0 = \lim_{\substack{x \rightarrow x_0 \\ x \in E \setminus \{x_0\}}} \left| \frac{f(x) - f(x_0)}{x - x_0} - L \right| = \lim_{\substack{x \rightarrow x_0 \\ x \in E \setminus \{x_0\}}} \left| \frac{f(x) - f(x_0) - L(x - x_0)}{x - x_0} \right|.$$

Lecture 2

26 Feb.

Definition 1.2.1 (Differentiability). Let E be a subset of $\mathbb{R}^n$, let $f : E \rightarrow \mathbb{R}^m$ be a function, and let x_0 be a limit point of E . Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. We say f is differentiable at x_0 with derivative L if

$$\lim_{\substack{x \rightarrow x_0 \\ x \in E \setminus \{x_0\}}} \frac{\|f(x) - f(x_0) - L(x - x_0)\|}{\|x - x_0\|} = 0,$$

or equivalently, given $\varepsilon > 0$, $\exists \delta > 0$ s.t. for all $x \in E$ satisfying $0 < \|x - x_0\| < \delta$, we have

$$\frac{\|f(x) - f(x_0) - L(x - x_0)\|}{\|x - x_0\|} < \varepsilon.$$

Remark 1.2.1. $\|f(x) - f(x_0) - L(x - x_0)\|$ is the length of a vector in $\mathbb{R}^m$ and $\|x - x_0\|$ is the length of a vector in $\mathbb{R}^n$.

Remark 1.2.2. x_0 is a limit point of $E \subseteq \mathbb{R}^n$ if for any $r > 0$, $B(x_0, r) \cap (E \setminus \{x_0\}) \neq \emptyset$. That is, for every $r > 0$, $\exists x \in E$ s.t. $0 < \|x - x_0\| < r$.

Example 1.2.1. Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ has

$$f(x) = \begin{cases} 0, & \text{if } x \in E = \{(x_1, x_2, \dots, x_n) \mid x_n \geq 0\}; \\ x, & \text{if } x \in \mathbb{R}^n \setminus E \end{cases}$$

then f is differentiable on E and $f'(x) = 0$ and f is not differentiable on $\mathbb{R}^n$ at 0. This is because on E , f is a constant function, so

$$\frac{\|f(x) - f(x_0) - L(x - x_0)\|}{\|x - x_0\|} = 0$$

for $x_0 \in E, x \in E \setminus \{x_0\}$. On $\mathbb{R}^n$ and $x_0 = 0$, then for $x \in \mathbb{R}^n \setminus \{x_0\} = \mathbb{R}^n \setminus \{0\}$, then if we approach to 0 along E , then it gives

$$\lim_{\substack{x \rightarrow x_0 \\ x \in E}} \frac{\|f(x) - f(x_0) - L(x - x_0)\|}{\|x - x_0\|} = 0,$$

but if we approach 0 along $x_n < 0$, then it becomes

$$\lim_{\substack{x \rightarrow x_0 \\ x \in \mathbb{R}^n \setminus E}} \frac{\|f(x) - f(x_0) - L(x - x_0)\|}{\|x - x_0\|} = \lim_{\substack{x \rightarrow x_0 \\ x \in \mathbb{R}^n \setminus E}} \frac{\|x - 0 - L(x - x_0)\|}{\|x\|} = 1$$

for any linear map L since any linear map is differentiable and thus continuous. Thus, f is not differentiable at 0 on $\mathbb{R}^n$.

Remark 1.2.3. Recall that x_0 is said to be an interior point of $E \subseteq \mathbb{R}^n$ if there exists $r > 0$ s.t.

$$B(x_0, r) = \{x \in \mathbb{R}^n : \|x - x_0\| < r\} \subseteq E.$$

If x_0 is an interior point of E , then f is differentiable at x_0 is equivalent to

$$\lim_{\substack{h \rightarrow 0 \\ x_0 + h \in E}} \frac{\|f(x_0 + h) - f(x_0) - L(h)\|}{\|h\|} = 0$$

since if $\|h\| < r$, then $x_0 + h \in B(x_0, r)$, which gives $x_0 + h \in E$, and thus $f(x_0 + h)$ is well-defined.

Remark 1.2.4. Here $\|\cdot\|$ denoted the standard Euclidean norm on $\mathbb{R}^n$ (and on $\mathbb{R}^m$):

$$\|(x_1, x_2, \dots, x_n)\| = (x_1^2 + x_2^2 + \dots + x_n^2)^{\frac{1}{2}}.$$

Example 1.2.2. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be given by

$$f(x, y) = (x^2, y^2),$$

let $x_0 = (1, 2)$, and let $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear map

$$L(x, y) = (2x, 4y).$$

We claim that f is differentiable at x_0 with derivative L .

Proof. By definition, f is differentiable at x_0 with derivative L if and only if

$$\lim_{h \rightarrow 0} \frac{\|f(x_0 + h) - f(x_0) - L(h)\|}{\|h\|} = 0,$$

where $h = (a, b)$. Thus, we know the above equation becomes

$$\frac{\|f(1 + a, 2 + b) - f(1, 2) - L(a, b)\|}{\|(a, b)\|} = \frac{\sqrt{a^4 + b^4}}{\sqrt{a^2 + b^2}},$$

and since

$$0 \leq \frac{\sqrt{a^4 + b^4}}{\sqrt{a^2 + b^2}} \leq \sqrt{a^2 + b^2},$$

so by the Squeeze Theorem, we have

$$\lim_{(a,b) \rightarrow (0,0)} \frac{\sqrt{a^4 + b^4}}{\sqrt{a^2 + b^2}} = 0,$$

and we're done. *

Lemma 1.2.2 (Uniqueness of derivative). Let E be a subset of $\mathbb{R}^n$, and let $f : E \rightarrow \mathbb{R}^m$ be a function, and let $x_0 \in E$ be an interior point of E . Suppose $L_1, L_2 : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are linear transformations s.t. f is differentiable at x_0 with derivative L_1 and also differentiable at x_0 with derivative L_2 . Then $L_1 = L_2$.

Proof. Since x_0 is an interior point. By Remark 1.2.3, we have

$$\lim_{h \rightarrow 0} \frac{\|f(x_0 + h) - f(x_0) - L_1(h)\|}{\|h\|} = 0 \text{ and } \lim_{h \rightarrow 0} \frac{\|f(x_0 + h) - f(x_0) - L_2(h)\|}{\|h\|} = 0.$$

Thus,

$$\begin{aligned} 0 &\leq \frac{\|L_1(h) - L_2(h)\|}{\|h\|} = \frac{\|L_1(h) - f(x_0 + h) + f(x_0) + f(x_0 + h) - f(x_0) - L_2(h)\|}{\|h\|} \\ &\leq \frac{\|L_1(h) - f(x_0 + h) + f(x_0)\|}{\|h\|} + \frac{\|f(x_0 + h) + f(x_0) - L_2(h)\|}{\|h\|}. \end{aligned}$$

Thus, by squeeze theorem, we know

$$\lim_{h \rightarrow 0} \frac{\|L_1(h) - L_2(h)\|}{\|h\|} = 0.$$

Now take $v \in \mathbb{R}^n \setminus \{0\}$, and let $h = tv$, so

$$\lim_{t \rightarrow 0} \frac{\|L_1(tv) - L_2(tv)\|}{\|tv\|} = 0.$$

That is,

$$\lim_{t \rightarrow 0} \frac{|t| \|L_1(v) - L_2(v)\|}{|t| \|v\|} = 0.$$

Hence, we have

$$\lim_{t \rightarrow 0} \frac{\|L_1(v) - L_2(v)\|}{\|v\|} = 0,$$

and since $v \neq 0$, so we know $L_1(v) = L_2(v)$, and since v can be arbitrary non-zero vector in $\mathbb{R}^n \setminus \{0\}$, so $L_1 = L_2$. ($L_1(0) = L_2(0) = 0$) ■

Remark 1.2.5. Because of [Lemma 1.2.2](#), the derivative of f at interior points x_0 is unique, and thus we may safely write it as $f'(x_0)$ or $Df(x_0)$. Thus $f'(x_0)$ is the unique linear transformation $f'(x_0) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that

$$\lim_{x \rightarrow x_0, x \neq x_0} \frac{\|f(x) - (f(x_0) + f'(x_0)(x - x_0))\|}{\|x - x_0\|} = 0.$$

Informally, this condition means that near x_0 , the function f can be approximated by its linearization:

$$f(x) \approx f(x_0) + f'(x_0)(x - x_0).$$

This approximation is sometimes referred to as Newton's approximation in higher dimension.

Remark 1.2.6. A useful consequence of [Lemma 1.2.2](#) is the following: If two functions f and g satisfy $f(x) = g(x)$ for all $x \in E$, and both are differentiable at an interior point x_0 , then their derivatives coincide at x_0 , i.e. $f'(x_0) = g'(x_0)$. This will be important in later arguments, where one extends functions to larger domains or modifies them on sets of measure zero.

Remark 1.2.7. As we have emphasized, [Lemma 1.2.2](#) guarantees the uniqueness of the derivative only at interior points of the domain E . If x_0 is instead a boundary point of E , the derivative may fail to be uniquely determined.

1.3 Partial and Directional Derivatives

We now begin to relate the concept of differentiability to the more classical notions of partial and directional derivatives. Directional derivatives describe how f changes when we move from a point x_0 in a fixed direction v .

Definition 1.3.1 (Directional derivative). Let $E \subseteq \mathbb{R}^n$, let $f : E \rightarrow \mathbb{R}^m$ be a function, let x_0 be an interior point of E and let $v \in \mathbb{R}^n$. If the limit

$$\lim_{\substack{t \rightarrow 0, t > 0 \\ x_0 + tv \in E}} \frac{f(x_0 + tv) - f(x_0)}{t} = \lim_{\substack{t \rightarrow 0^+ \\ x_0 + tv \in E}} \frac{f(x_0 + tv) - f(x_0)}{t}$$

exists, then we say f is differentiable in the direction v at x_0 . This limit is called the directional derivative of f at x_0 in the direction v , and we denote it by

$$D_v f(x_0) := \lim_{t \rightarrow 0^+} \frac{f(x_0 + tv) - f(x_0)}{t} \in \mathbb{R}^m.$$

Remark 1.3.1. Under this definition,

$$D_{-v}f(x_0) = \lim_{t \rightarrow 0^+} \frac{f(x_0 + t(-v)) - f(x_0)}{t} = \lim_{t \rightarrow 0^+} \frac{f(x_0 - tv) - f(x_0)}{t}.$$

In usual definition, one define

$$D_vf(x_0) = \lim_{t \rightarrow 0} \frac{f(x_0 + tv) - f(x_0)}{t}.$$

This usual definition implies that

$$D_vf(x_0) = -D_{-v}f(x_0)$$

since

$$\begin{aligned} D_vf(x_0) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + tv) - f(x_0)}{t} = \lim_{t \rightarrow 0^-} \frac{f(x_0 + tv) - f(x_0)}{t} \\ &= \lim_{s \rightarrow 0^+} \frac{f(x_0 + (-s)v) - f(x_0)}{(-s)} = - \lim_{s \rightarrow 0^+} \frac{f(x_0 - sv) - f(x_0)}{s} = -(D_{-v}f(x_0)). \end{aligned}$$

Remark 1.3.2. This definition should be compared with the definition of differentiability. Here we divide by the scalar t rather than by a vector, so the expression always makes sense algebraically. The directional derivative $D_vf(x_0)$ is a vector in $\mathbb{R}^m$.

Remark 1.3.3. It is sometimes possible to define directional derivatives at boundary points of E , provided that the vector v points inward toward the domain. However, we shall restrict attention to interior points in what follows.

Lemma 1.3.1. Let $E \subseteq \mathbb{R}^n$, let $f : E \rightarrow \mathbb{R}^m$, let $x_0 \in \text{Int}(E)$ and $v \in \mathbb{R}^n$. If f is differentiable at x_0 , then f is differentiable in the direction v at x_0 , and

$$D_vf(x_0) = f'(x_0)(v).$$

Proof. Since $x_0 \in \text{Int}(E)$ and f is differentiable at x_0 , so

$$\lim_{h \rightarrow 0} \frac{\|f(x_0 + h) - f(x_0) - f'(x_0)(h)\|}{\|h\|} = 0.$$

Note that for $t \neq 0$, we have

$$\frac{f(x_0 + tv) - f(x_0)}{t} = \frac{f(x_0 + tv) - f(x_0) - f'(x_0)(tv)}{t} + f'(x_0)(v),$$

so

$$\frac{f(x_0 + tv) - f(x_0)}{t} - f'(x_0)(v) = \frac{f(x_0 + tv) - f(x_0) - f'(x_0)(tv)}{t}.$$

Now we show that

$$\lim_{t \rightarrow 0^+} \frac{f(x_0 + tv) - f(x_0) - f'(x_0)(tv)}{t} = 0.$$

Recall that f is differentiable at x_0 and $x_0 \in \text{Int}(E)$, so

$$\lim_{h \rightarrow 0} \frac{\|f(x_0 + h) - f(x_0) - f'(x_0)(h)\|}{\|h\|} = 0.$$

Let $h = tv$ when $v \neq 0$, then

$$\lim_{t \rightarrow 0^+} \frac{\|f(x_0 + tv) - f(x_0) - f'(x_0)(tv)\|}{\|tv\|} = 0 \Rightarrow \lim_{t \rightarrow 0^+} \frac{\|f(x_0 + tv) - f(x_0) - f'(x_0)(tv)\|}{|t|} = 0,$$

which is equivalent to

$$\lim_{t \rightarrow 0^+} \frac{f(x_0 + tv) - f(x_0) - f'(x_0)(tv)}{t} = 0,$$

(since the numerator should tend to 0 so the limit will exist), so we know

$$\lim_{t \rightarrow 0^+} \frac{f(x_0 + tv) - f(x_0)}{t} - f'(x_0)(v) = 0,$$

i.e.

$$\lim_{t \rightarrow 0^+} \frac{f(x_0 + tv) - f(x_0)}{t} = f'(x_0)(v).$$

■

Remark 1.3.4. One consequence of this lemma is that total differentiability implies directional differentiability. However, the converse is not true: the existence of all directional derivatives does not guarantee differentiability.

Definition 1.3.2 (Partial Derivative). Let $E \subseteq \mathbb{R}^n$, let $f : E \rightarrow \mathbb{R}^m$, and let $x_0 \in \text{Int}(E)$, and let $1 \leq j \leq n$. The partial derivative of f with respect to the variable x_j at x_0 , denoted $\frac{\partial f}{\partial x_j}(x_0)$, is defined by

$$\frac{\partial f}{\partial x_j}(x_0) := \lim_{t \rightarrow 0, t \neq 0} \frac{f(x_0 + te_j) - f(x_0)}{t}$$

provided the limit exists.

Remark 1.3.5. If $f = \begin{pmatrix} f_1 \\ \vdots \\ f_m \end{pmatrix}$, then differentiation is componentwise:

$$\frac{\partial f}{\partial x_j}(x_0) = \begin{pmatrix} \frac{\partial f_1}{\partial x_j}(x_0) \\ \vdots \\ \frac{\partial f_m}{\partial x_j}(x_0) \end{pmatrix}.$$

We say that f is continuously differentiable if all partial derivatives exist and are continuous.

Remark 1.3.6. If f is differentiable at x_0 , then

$$\frac{\partial f(x_0)}{\partial x_j} = D_{e_j} f(x_0) = f'(x_0)(e_j).$$

Remark 1.3.7. If f is differentiable at x_0 , then

$$D_v f(x_0) = f'(x_0)(v) = f'(x_0) \left(\sum_{j=1}^n v_j e_j \right) = \sum_{j=1}^n v_j f'(x_0)(e_j) = \sum_{j=1}^n v_j \frac{\partial f(x_0)}{\partial x_j}.$$

Theorem 1.3.1. Let $E \subseteq \mathbb{R}^n$, let $f : E \rightarrow \mathbb{R}^m$, $F \subseteq E$, and let $x_0 \in \text{Int}(F)$ (the partial derivative exists on a neighborhood of x_0). If all partial derivative $\frac{\partial f}{\partial x_j}$ exists on F and are continuous at x_0 , then f is differentiable at x_0 , and

$$f'(x_0)(v) = \sum_{j=1}^n v_j \frac{\partial f}{\partial x_j}(x_0)$$

for all $v \in \mathbb{R}^n$.

Proof. Define a linear map $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by

$$L(v) := \sum_{j=1}^n v_j \frac{\partial f}{\partial x_j}(x_0).$$

We prove that f is differentiable at x_0 with derivative L . It suffices to show that for any $\varepsilon > 0$, $\exists \delta > 0$ s.t.

$$\|f(x) - f(x_0) - L(x - x_0)\| < \varepsilon \|x - x_0\| \text{ whenever } x \in F \text{ and } \|x - x_0\| < \delta$$

Since $x_0 \in \text{Int}(F)$, $\exists r > 0$ s.t. $B(x_0, r) \subseteq F$. Because each partial derivative $\frac{\partial f_i}{\partial x_j}$ is continuous at x_0 , for every pair (i, j) there exists $\delta_{ij} > 0$ s.t.

$$\left| \frac{\partial f_i}{\partial x_j}(x) - \frac{\partial f_i}{\partial x_j}(x_0) \right| < \frac{\varepsilon}{nm} \text{ whenever } \|x - x_0\| < \delta_{ij}.$$

Let $\delta := \min \{r, \delta_{ij} : 1 \leq i \leq m, 1 \leq j \leq n\}$. Now fix $x \in F$ with $\|x - x_0\| < \delta$ and write

$$x - x_0 = \sum_{j=1}^n v_j e_j, \text{ so } x = x_0 + \sum_{j=1}^n v_j e_j.$$

Then

$$\|x - x_0\| = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}, \quad |v_j| \leq \|x - x_0\|,$$

Write $f = (f_1, f_2, \dots, f_m)$. To estimate $f(x) - f(x_0)$, we vary the coordinates one at a time. Consider

$$\begin{aligned} x^{(0)} &= x_0, \\ x^{(j)} &= x_0 + \sum_{k=1}^j v_k e_k \text{ for } j = 1, 2, \dots, n. \end{aligned}$$

Thus, $x^{(n)} = x$ and

$$f(x) - f(x_0) = \sum_{j=1}^n \left(f(x^{(j)}) - f(x^{(j-1)}) \right) = \sum_{j=1}^n \sum_{i=1}^m \left(f_i(x^{(j)}) - f_i(x^{(j-1)}) \right) e_i.$$

Fix j and consider the j -th increment. For each component f_i , define

$$\varphi_i^j(t) = f_i(x^{(j-1)} + t v_j e_j), \quad 0 \leq t \leq 1.$$

Then

$$f_i(x^{(j)}) - f_i(x^{(j-1)}) = \varphi_i^j(1) - \varphi_i^j(0).$$

Since f_i has continuous partial derivatives, φ_i^j is differentiable. By MVT, there exists $t_j \in (0, 1)$ s.t.

$$\varphi_i^j(1) - \varphi_i^j(0) = \varphi_i^{j'}(t_j).$$

By the chain rule (See [Lemma 1.3.2](#)),

$$\varphi_i^{j'}(t) = \nabla f_i(x^{(j-1)} + t v_j e_j) \cdot (v_j e_j) = \frac{\partial f_i}{\partial x_j}(x^{(j-1)} + t v_j e_j) v_j.$$

Hence,

$$f_i(x^{(j)}) - f_i(x^{(j-1)}) = \frac{\partial f_i}{\partial x_j}(\xi_j) v_j,$$

where

$$\xi_j = x^{(j-1)} + t_j v_j e_j.$$

Subtracting $\frac{\partial f_i}{\partial x_j}(x_0)v_j$, we obtain

$$\left| f_i(x^{(j)}) - f_i(x^{(j-1)}) - \frac{\partial f_i}{\partial x_j}(x_0)v_j \right| = \left| \frac{\partial f_i}{\partial x_j}(\xi_j) - \frac{\partial f_i}{\partial x_j}(x_0) \right| |v_j|.$$

Because $\|\xi_j - x_0\| < \delta$, continuity of partial derivatives yields,

$$\left| f_i(x^{(j)}) - f_i(x^{(j-1)}) - \frac{\partial f_i}{\partial x_j}(x_0)v_j \right| < \frac{\varepsilon}{nm} |v_j|.$$

Summing over $i = 1, 2, \dots, m$ and using $\|u\| \leq \sum_i |u_i|$, we conclude

$$\begin{aligned} \left\| f(x^{(j)}) - f(x^{(j-1)}) - \frac{\partial f}{\partial x_j}(x_0)v_j \right\| &= \left\| \sum_{i=1}^m \left(f_i(x^{(j)}) - f_i(x^{(j-1)}) - \frac{\partial f_i}{\partial x_j}(x_0)v_j \right) e_i \right\| \\ &\leq \sum_{i=1}^m \left| f_i(x^{(j)}) - f_i(x^{(j-1)}) - \frac{\partial f_i}{\partial x_j}(x_0)v_j \right| \leq m \frac{\varepsilon}{nm} |v_j| = \frac{\varepsilon}{n} |v_j| \leq \frac{\varepsilon}{n} \|x - x_0\|. \end{aligned}$$

Finally, summing over $j = 1, 2, \dots, n$ and applying the triangle inequality,

$$\begin{aligned} \left\| f(x) - f(x_0) - \sum_{j=1}^n \frac{\partial f}{\partial x_j}(x_0)v_j \right\| &= \left\| \sum_{j=1}^n \left(f(x^{(j)}) - f(x^{(j-1)}) - \frac{\partial f}{\partial x_j}(x_0)v_j \right) \right\| \\ &\leq \sum_{j=1}^n \left\| \left(f(x^{(j)}) - f(x^{(j-1)}) - \frac{\partial f}{\partial x_j}(x_0)v_j \right) \right\| \leq \sum_{j=1}^n \frac{\varepsilon}{n} \|x - x_0\| = \varepsilon \|x - x_0\|. \end{aligned}$$

Since $L(x - x_0) = \sum_{j=1}^n v_j \frac{\partial f}{\partial x_j}(x_0)$, so we have

$$\|f(x) - f(x_0) - L(x - x_0)\| \leq \varepsilon \|x - x_0\|,$$

and we're done. ■

Remark 1.3.8. From [Theorem 1.3.1](#) and [Lemma 1.3.1](#) we conclude the following important fact:

If the partial derivatives of a function $f : E \rightarrow \mathbb{R}^m$ exist and are continuous on a set $F \subseteq E$, then at every interior point x_0 of F all directional derivatives exist, and they are given by

$$D_{(v_1, v_2, \dots, v_n)} f(x_0) = \sum_{j=1}^n v_j \frac{\partial f}{\partial x_j}(x_0).$$

- The scalar-valued case. If $f : E \rightarrow \mathbb{R}$ is real-valued, we define the gradient of f at x_0 to be the row vector

$$\nabla f(x_0) := \left(\frac{\partial f}{\partial x_1}(x_0), \dots, \frac{\partial f}{\partial x_n}(x_0) \right).$$

Whenever x_0 lies in the interior of a region where the partial derivatives exist and are continuous, the directional derivative takes the familiar form

$$D_v f(x_0) = \nabla f(x_0) \cdot v.$$

- The vector-valued case. Now let $f : E \rightarrow \mathbb{R}^m$ be vector-valued, say

$$f = \begin{pmatrix} f_1 \\ \vdots \\ f_m \end{pmatrix}.$$

If x_0 lies in the interior of a region where all partial derivatives exist and are continuous, then

$$f'(x_0)(v_1, v_2, \dots, v_n) = \sum_{j=1}^n v_j \frac{\partial f}{\partial x_j}(x_0).$$

Writing this componentwise,

$$f'(x_0)(v_1, \dots, v_n) = \sum_{j=1}^n v_j \begin{pmatrix} \frac{\partial f_1}{\partial x_j}(x_0) \\ \vdots \\ \frac{\partial f_m}{\partial x_j}(x_0) \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^n v_j \frac{\partial f_1}{\partial x_j}(x_0) \\ \vdots \\ \sum_{j=1}^n v_j \frac{\partial f_m}{\partial x_j}(x_0) \end{pmatrix} = \begin{pmatrix} \nabla f_1(x_0) \cdot v \\ \vdots \\ \nabla f_m(x_0) \cdot v \end{pmatrix} = \begin{pmatrix} \nabla f_1(x_0) \\ \vdots \\ \nabla f_m(x_0) \end{pmatrix} v$$

- The derivative matrix (Jacobian matrix). We therefore define the derivative matrix (or Jacobian matrix) of f at x_0 by

$$Df(x_0) := \left(\frac{\partial f_i}{\partial x_j}(x_0) \right)_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}}.$$

Explicitly,

$$Df(x_0) = \begin{pmatrix} \frac{\partial f_1}{\partial x_1}(x_0) & \cdots & \frac{\partial f_1}{\partial x_n}(x_0) \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1}(x_0) & \cdots & \frac{\partial f_m}{\partial x_n}(x_0) \end{pmatrix}.$$

With this notation, the derivative acts by matrix multiplication:

$$D_v f(x_0) = f'(x_0)v = Df(x_0)v.$$

Lecture 3

Lemma 1.3.2. Let $E \subseteq \mathbb{R}^n$ and let $f = (f_1, f_2, \dots, f_m) : E \rightarrow \mathbb{R}^m$. Fix $1 \leq i \leq m$ and $1 \leq j \leq n$. Let $z \in E$, $v \in \mathbb{R}$ and define $\phi_i^j(t) = f_i(z + tve_j)$ for $0 \leq t \leq 1$. Assume that for some $t_0 \in (0, 1)$, $\exists \rho > 0$ s.t.

$$z + (t_0 + s)ve_j \in E \quad \text{for all } |s| < \rho,$$

and that the partial derivative $\frac{\partial f_i}{\partial x_j}$ exists at $p = z + t_0ve_j$. Then, ϕ_i^j is differentiable at t_0 and

$$\left(\phi_i^j \right)'(t_0) = \frac{\partial f_i}{\partial x_j}(p)(v).$$

Proof. Note that

$$\begin{aligned} \left(\phi_i^j \right)'(t_0) &= \lim_{h \rightarrow 0} \frac{\phi_i^j(t_0 + h) - \phi_i^j(t_0)}{h} = \lim_{h \rightarrow 0} \frac{f_i(z + (t_0 + h)ve_j) - f_i(z + t_0ve_j)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f_i(p + hve_j) - f_i(p)}{h}. \end{aligned}$$

Now if $v = 0$, then this limit is equal to 0. Now if $v \neq 0$, then

$$\lim_{h \rightarrow 0} \frac{f_i(p + hve_j) - f_i(p)}{h} = \lim_{h \rightarrow 0} \frac{f_i(p + hve_j) - f_i(p)}{vh} \cdot v = \lim_{s \rightarrow 0} \frac{f_i(p + se_j) - f_i(p)}{s} \cdot v = \frac{\partial f_i}{\partial x_j}(p) \cdot v. \quad \blacksquare$$

Example 1.3.1. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by $f(x, y) = (x^2 + xy, y^2)$, then

$$Df(x, y) = \begin{pmatrix} \frac{\partial}{\partial x}(x^2 + xy) & \frac{\partial}{\partial y}(x^2 + xy) \\ \frac{\partial}{\partial x}(y^2) & \frac{\partial}{\partial y}(y^2) \end{pmatrix} = \begin{pmatrix} 2x + y & x \\ 0 & 2y \end{pmatrix}.$$

Thus,

$$D_{(v,w)}f(x,y) = \begin{pmatrix} 2x+y & x \\ 0 & 2y \end{pmatrix} \begin{pmatrix} v \\ w \end{pmatrix} = \begin{pmatrix} (2x+y)v + xw \\ 2yw \end{pmatrix}.$$

1.4 Chain Rule

If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$, then their composition $g \circ f : X \rightarrow Z$ is defined by $(g \circ f)(x) = g(f(x))$. We want to show that

$$D(g \circ f)(x) = (Dg(f(x))) \cdot (Df(x))$$

under suitable conditions.

Definition 1.4.1. Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ be square matrix. The trace of A , denoted $\text{Tr}(A)$, is defined by $\text{Tr}(A) = \sum_{i=1}^n a_{ii}$. Let $M = (m_{ij})$ be an $r \times s$ matrix. We may regard M as an element of $\mathbb{R}^{r \times s}$. We can define its Frobenius norm by

$$\|M\|_F = \left(\sum_{i=1}^r \sum_{j=1}^s m_{ij}^2 \right)^{\frac{1}{2}},$$

which can be written as $\|M\|_F = (\text{Tr}(MM^t))^{\frac{1}{2}}$.

Theorem 1.4.1. Let $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$. Define the Frobenius norm by

$$\|M\|_F = \sqrt{\text{Tr}(MM^t)} = \left(\sum_{i=1}^r \sum_{j=1}^s m_{ij}^2 \right)^{\frac{1}{2}}$$

with $M \in \mathbb{R}^{r \times s}$. Then,

$$\|AB\|_F \leq \|A\|_F \|B\|_F.$$

In particular, for any column vector $v \in \mathbb{R}^n$, we have

$$\|Av\| \leq \|A\|_F \|v\|,$$

and the linear map $x \mapsto Ax$ is continuous on $\mathbb{R}^n$.

Proof. Note that

$$(AB)_{ik} = \sum_{j=1}^n a_{ij} b_{jk},$$

so

$$|(AB)_{ik}| = \left| \sum_{j=1}^n a_{ij} b_{jk} \right| \leq \left(\sum_{j=1}^n a_{ij}^2 \right)^{\frac{1}{2}} \left(\sum_{j=1}^n b_{jk}^2 \right)^{\frac{1}{2}}.$$

Hence, we have

$$\|AB\|_F^2 = \sum_i \sum_k (AB)_{ik}^2 \leq \sum_i \sum_k \left(\sum_{j=1}^n a_{ij}^2 \right) \left(\sum_{j=1}^n b_{jk}^2 \right) = \|A\|_F^2 \|B\|_F^2.$$

For the second part, since we have

$$\|Av\|_F \leq \|A\|_F \|v\|_F,$$

and we know for column vector, Frobenius norm coincides with the Euclidean norm, so we're done.

Finally, for $x, x_0 \in \mathbb{R}^n$, we know $Ax - Ax_0 = A(x - x_0)$, and thus

$$0 \leq \|Ax - Ax_0\| = \|A(x - x_0)\| \leq \|A\|_F \|x - x_0\|,$$

and if $x \rightarrow x_0$, we know

$$\|A\|_F \|x - x_0\| \rightarrow 0$$

since $\|A\|_F$ is a constant, and thus $\|A(x - x_0)\|$ also converges to 0, which shows the map $x \mapsto Ax$ is linear. ■

Theorem 1.4.2. Let $E \subseteq \mathbb{R}^n$, let $f : E \rightarrow \mathbb{R}^m$, and let x_0 be an interior point of E . If f is differentiable at x_0 , then f is continuous at x_0 .

Proof. If f is differentiable at x_0 , then there exists $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ s.t.

$$\lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - L(x - x_0)\|}{\|x - x_0\|} = 0.$$

Thus,

$$\lim_{x \rightarrow x_0} \|f(x) - f(x_0) - L(x - x_0)\| = \lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - L(x - x_0)\|}{\|x - x_0\|} \lim_{x \rightarrow x_0} \|x - x_0\| = 0.$$

Hence, we know

$$\begin{aligned} \|f(x) - f(x_0)\| &= \|f(x) - f(x_0) - L(x - x_0) + L(x - x_0)\| \\ &\leq \|f(x) - f(x_0) - L(x - x_0)\| + \|L(x - x_0)\| \end{aligned}$$

and the former converges to 0 as $x \rightarrow x_0$, while the latter also converges to 0 as $x \rightarrow x_0$ since L is continuous. ■

Theorem 1.4.3 (Chain Rule). Let $E \subseteq \mathbb{R}^n$, let $F \subseteq \mathbb{R}^m$, and let $f : E \rightarrow F$ and $g : F \rightarrow \mathbb{R}^p$. Let $x_0 \in \text{Int}(E)$. Suppose f is differentiable at x_0 , that $f(x_0)$ lies in the interior of F , and that g is differentiable at $f(x_0)$. Then, $g \circ f : E \rightarrow \mathbb{R}^p$ is differentiable at x_0 , and

$$(g \circ f)'(x_0) = g'(f(x_0)) \circ f'(x_0)$$

Proof. Set $y_0 = f(x_0)$ and denote the derivatives by $A = f'(x_0) : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $B : g'(f(x_0)) = g'(y_0) : \mathbb{R}^m \rightarrow \mathbb{R}^p$. Since g is differentiable at y_0 , define $\eta : \mathbb{R}^m \rightarrow \mathbb{R}^p$ by

$$\eta(k) = \begin{cases} \frac{g(y_0 + k) - g(y_0) - Bk}{\|k\|}, & \text{if } k \neq 0; \\ 0, & \text{if } k = 0. \end{cases}$$

Note that $\lim_{k \rightarrow 0} \eta(k) = 0$ since g is differentiable at y_0 , so η is continuous at 0. Moreover, we have

$$g(y_0 + k) - g(y_0) - Bk = \eta(k)\|k\| \quad \text{true for all } k.$$

Since $x_0 \in \text{Int}(E)$, $x_0 + h \in E$ for small enough $\|h\|$. Note that

$$\begin{aligned} (g \circ f)(x_0 + h) - g(f(x_0)) &= g(f(x_0 + h)) - g(f(x_0)) = g(f(x_0) + (f(x_0 + h) - f(x_0))) - g(f(x_0)) \\ &= B(f(x_0 + h) - f(x_0)) + \eta(f(x_0 + h) - f(x_0)) \cdot \|f(x_0 + h) - f(x_0)\|, \end{aligned}$$

so we have

$$(g \circ f)(x_0 + h) - g(f(x_0)) - BAh = B(f(x_0 + h) - f(x_0) - Ah) + \eta(f(x_0 + h) - f(x_0)) \cdot \|f(x_0 + h) - f(x_0)\|.$$

Now since

$$\lim_{h \rightarrow 0} \frac{\|f(x_0 + h) - f(x_0) - Ah\|}{\|h\|} = 0,$$

so

$$\frac{\|B(f(x_0 + h) - f(x_0) - Ah)\|}{\|h\|} \leq \frac{\|B\|_F \|f(x_0 + h) - f(x_0) - Ah\|}{\|h\|}$$

implies

$$\lim_{h \rightarrow 0} \frac{\|B(f(x_0 + h) - f(x_0) - Ah)\|}{\|h\|} = 0.$$

Now we claim that

$$\lim_{h \rightarrow 0} \frac{\eta(f(x_0 + h) - f(x_0)) \cdot \|f(x_0 + h) - f(x_0)\|}{\|h\|} = 0.$$

Note that

$$\frac{\eta(f(x_0 + h) - f(x_0)) \cdot \|f(x_0 + h) - f(x_0)\|}{\|h\|} = \eta(f(x_0 + h) - f(x_0)) \frac{\|f(x_0 + h) - f(x_0)\|}{\|h\|}.$$

Now since η is continuous at 0, so

$$\lim_{h \rightarrow 0} \eta(f(x_0 + h) - f(x_0)) = \eta(0) = 0.$$

Also,

$$\begin{aligned} \frac{\|f(x_0 + h) - f(x_0)\|}{\|h\|} &= \frac{\|f(x_0 + h) - f(x_0) - f'(x_0)(h) + f'(x_0)(h)\|}{\|h\|} \\ &\leq \frac{\|f(x_0 + h) - f(x_0) - f'(x_0)(h)\| + \|f'(x_0)(h)\|}{\|h\|} \\ &\leq \frac{\|f(x_0 + h) - f(x_0) - f'(x_0)(h)\|}{\|h\|} + \frac{\|h\| \|f'(x_0)\|_F}{\|h\|} \\ &\rightarrow \|A\|_F \text{ as } h \rightarrow 0. \end{aligned}$$

Thus,

$$\lim_{h \rightarrow 0} \frac{\|(g \circ f)(x_0 + h) - g(f(x_0)) - BAh\|}{\|h\|} = 0.$$

■

Corollary 1.4.1. We may express the chain rule in matrix form.

$$D(g \circ f)(x_0) = Dg(f(x_0))Df(x_0).$$

Remark 1.4.1. We prove the chain rule of linear transformations in the last theorem, and earlier we have found that in fact the linear transformation $f'(x_0)$ is the Jacobian matrix $Df(x_0)$, so we can write the matrix form of the chain rule, i.e. the chain rule on Jacobian matrix.

Example 1.4.1. Let $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable functions. Recall that

$$\nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right), \quad \nabla g = \left(\frac{\partial g}{\partial x_1}, \frac{\partial g}{\partial x_2}, \dots, \frac{\partial g}{\partial x_n} \right),$$

and

$$\nabla f = \sum_{i=1}^n \frac{\partial f}{\partial x_i} e_i,$$

then we have

$$\nabla(fg) = \sum_{i=1}^n \frac{\partial}{\partial x_i}(fg)e_i = \sum_{i=1}^n \left(\frac{\partial f}{\partial x_i}g + f \frac{\partial g}{\partial x_i} \right) e_i = g\nabla f + f\nabla g$$

by ordinary product rule(前微後不微加上前不微後微).

Example 1.4.2. Consider $f : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$ defined by

$$f(A) = A^t A,$$

then $Df(A) = ?$

Proof. We use two methods to solve this problem.

Method 1: Direct Expansion (Using the Definition) To find the derivative of f at A , we apply a small matrix perturbation $H \in \mathbb{R}^{n \times n}$ and expand $f(A + H)$:

$$f(A + H) = (A + H)^t (A + H) = (A^t + H^t) (A + H) = A^t A + A^t H + H^t A + H^t H.$$

Hence, we have

$$f(A + H) - f(A) = (A^t H + H^t A) + H^t H.$$

Hence, we may choose $L : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$ by

$$L(H) = A^t H + H^t A.$$

Thus, we have

$$\lim_{H \rightarrow 0} \frac{\|f(A + H) - f(A) - L(H)\|}{\|H\|} = \lim_{H \rightarrow 0} \frac{\|H^t H\|}{\|H\|} \leq \lim_{H \rightarrow 0} \frac{\|H^t\| \|H\|}{\|H\|} = \lim_{H \rightarrow 0} \|H^t\| = 0.$$

Method 2: Decomposition and the Chain Rule We can deconstruct f into

$$A \xrightarrow{\Phi} (A^t, A) \xrightarrow{\Psi} A^t A$$

Note that

$$\Phi(A + h) - \Phi(A) - (H^t, H) = 0,$$

so

$$\lim_{\|H\| \rightarrow 0} \frac{\|\Phi(A + H) - \Phi(A) - (H^t, H)\|}{\|H\|} = 0.$$

Thus, $\Phi(A)'(H) = (H^t, H)$. Also, we can check $\Psi'(B, C)(K, L) = KC + BL$, i.e.

$$\lim_{(K, L) \rightarrow (0, 0)} \frac{\|\Psi(B + K, C + L) - \Psi(B, C) - (KC + BL)\|}{\|(K, L)\|} = 0.$$

Note that

$$\Psi(B + K, C + L) - \Psi(B, C) - (KC + BL) = KL.$$

and

$$\|(K, L)\| = \sqrt{\|K\|^2 + \|L\|^2} \geq \frac{\|K\| + \|L\|}{\sqrt{2}}.$$

Thus,

$$\begin{aligned} \frac{\|\Psi(B + K, C + L) - \Psi(B, C) - (KC + BL)\|}{\|(K, L)\|} &\leq \sqrt{2} \frac{\|KL\|}{\|K\| + \|L\|} \leq \sqrt{2} \frac{\|K\| \|L\|}{\|K\| + \|L\|} \\ &\leq \sqrt{2} \frac{\|K\| \|L\|}{\|L\|} = \sqrt{2} \|K\|. \end{aligned}$$

This shows

$$0 \leq \lim_{(K,L) \rightarrow (0,0)} \frac{\|\Psi(B+K, C+L) - \Psi(B, C) - (KC + BL)\|}{\|(K, L)\|} \leq \lim_{(K,L) \rightarrow (0,0)} \sqrt{2}\|K\| = 0.$$

By chain rule, we know

$$f'(A)(H) = \Psi'(\Phi(A))(\Phi'(A)(H)) = H^t A + A^t H.$$

⊛

Lecture 4

If $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear, then $T'(x) = T$ for every x . This is because for every x

5 March.

$$\lim_{h \rightarrow 0} \frac{\|T(x+h) - T(x) - T(h)\|}{\|h\|} = 0.$$

Hence, for any differentiable $f : E \rightarrow \mathbb{R}^n$, the composition Tf is differentiable and

$$(Tf)'(x_0) = T'(f(x_0)) \circ f'(x_0) = T \circ f'(x_0) = T(f'(x_0)).$$

This generalizes the familiar one-variable rule $(cf)' = cf'$.

Another useful special case of the chain rule is the following. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable and $x_j : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable functions for $j = 1, 2, \dots, n$, then

$$\frac{d}{dt} f(x_1(t), x_2(t), \dots, x_n(t)) = \sum_{j=1}^n x'_j(t) \frac{\partial f}{\partial x_j}(x_1(t), x_2(t), \dots, x_n(t))$$

since we can suppose $f = (f_1, f_2, \dots, f_m)$ and $g(t) = (x_1(t), x_2(t), \dots, x_n(t))^t$ and thus

$$\begin{aligned} \frac{d}{dt} f(x_1(t), x_2(t), \dots, x_n(t)) &= \frac{d}{dt} f \circ g(t) = f'(g(t)) \circ g'(t) \\ &= \begin{pmatrix} \frac{\partial f_1}{\partial x_1}(g(t)) & \frac{\partial f_1}{\partial x_2}(g(t)) & \cdots & \frac{\partial f_1}{\partial x_n}(g(t)) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1}(g(t)) & \frac{\partial f_m}{\partial x_2}(g(t)) & \cdots & \frac{\partial f_m}{\partial x_n}(g(t)) \end{pmatrix} \begin{pmatrix} x'_1(t) \\ \vdots \\ x'_n(t) \end{pmatrix} \\ &= \begin{pmatrix} \sum_{i=1}^n \frac{\partial f_1}{\partial x_i}(g(t)) x'_i(t) \\ \vdots \\ \sum_{i=1}^n \frac{\partial f_m}{\partial x_i}(g(t)) x'_i(t) \end{pmatrix} = \sum_{i=1}^n x'_i(t) \begin{pmatrix} \frac{\partial f_1}{\partial x_i}(g(t)) \\ \vdots \\ \frac{\partial f_m}{\partial x_i}(g(t)) \end{pmatrix} \\ &= \sum_{i=1}^n x'_i(t) \frac{\partial f}{\partial x_i}(g(t)) = \sum_{i=1}^n x'_i(t) \frac{\partial f}{\partial x_i}(x_1(t), x_2(t), \dots, x_n(t)). \end{aligned}$$

The Mean Value Theorem for functions $f : \mathbb{R} \rightarrow \mathbb{R}$ asserts that

$$f(y) - f(x) = f'(z)(y - x)$$

where $z \in (x, y)$. We begin by extending this statement to real-valued functions defined on open subsets of $\mathbb{R}^n$.

Theorem 1.4.4 (Mean Value Theorem for Real-Valued Functions). Let V be open in $\mathbb{R}^n$ and suppose that $f : V \rightarrow \mathbb{R}$ is differentiable on V . Given $a, x \in V$, write $L(x; a)$ for the (closed) line segment joining a to x , namely

$$L(x; a) := \{a + t(x - a) : 0 \leq t \leq 1\}.$$

Assume that $L(x; a) \subseteq V$. Then there exists $c \in L(x; a)$ such that

$$f(x) - f(a) = \nabla f(c) \cdot (x - a).$$

Proof. Suppose $g(t) := a + t(x - a)$ for $t \in \mathbb{R}$. Then, $g : \mathbb{R} \rightarrow \mathbb{R}^n$ is differentiable and

$$Dg(t) = x - a \quad \forall t \in \mathbb{R}.$$

Since V is open and the compact segment $L(x; a) = \{g(t) : t \in [0, 1]\}$ is contained in V , there exists $\delta > 0$ s.t.

$$g(t) \in V \quad \text{for all } t \in I_\delta = (-\delta, 1 + \delta).$$

Hence, $f \circ g : I_\delta \rightarrow \mathbb{R}$ is well-defined and differentiable. By the chain rule, for each $t \in I_\delta$ we have

$$D(f \circ g)(t) = Df(g(t))Dg(t) = Df(g(t))(x - a).$$

Because f is real-valued, the differential $Df(y)$ may be identified with the gradient by

$$Df(y)(h) = \nabla f(y) \cdot h, y \in V, h \in \mathbb{R}^n.$$

Therefore the above equation becomes

$$(f \circ g)'(t) = \nabla f(g(t)) \cdot (x - a), t \in I_\delta.$$

Now apply the one-dimensional MVT to $f \circ g$ on $[0, 1]$. There exists $t_0 \in (0, 1)$ such that

$$(f \circ g)(1) - (f \circ g)(0) = (f \circ g)'(t_0).$$

Since $(f \circ g)(1) = f(x)$ and $(f \circ g)(0) = f(a)$, we obtain

$$f(x) - f(a) = \nabla f(g(t_0)) \cdot (x - a).$$

We can set $c := g(t_0) \in L(x; a)$, and we're done. ■

The conclusion of [Theorem 1.4.4](#) relies essentially on the fact that f is real-valued. The following remark shows that no analogous statement can hold in general for vector-valued functions, even when the domain is one-dimensional.

Remark 1.4.2. The function $f(t) = (\cos t, \sin t)$ is differentiable on $\mathbb{R}$ and satisfies $f(w\pi) = f(0)$, but there is no $c \in \mathbb{R}$ s.t. $Df(c) = (0, 0)$.

Now we want to extend the MVT above to $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $m > 1$, one can recover a valid statement by testing against linear functionals on $\mathbb{R}^m$. Equivalently, we take the dot product with an arbitrary vector $a \in \mathbb{R}^m$ and apply the real-valued mean value theorem to the scalar function $t \mapsto a \cdot f(x + tu)$. This yields the following useful generalization.

In the following statement, we let

$$L(x; y) = \{tx + (1 - t)y : 0 \leq t \leq 1\}.$$

Theorem 1.4.5 (Mean-Value Theorem). Let S be an open subset of $\mathbb{R}^n$ and assume that $f : S \rightarrow \mathbb{R}^m$ is differentiable at each point of S . Let x and y be two points in S s.t. $L(x; y) \subseteq S$. Then for every vector $a \in \mathbb{R}^m$ there is a point $z \in L(x; y)$ s.t.

$$a \cdot \{f(y) - f(x)\} = a \cdot \{f'(z)(y - x)\}.$$

Proof. Let $u = y - x$. Since S is open and $L(x; y) \subseteq S$, there is a $\delta > 0$ s.t.

$$x + tu \in S \quad \text{for all real } t \in (-\delta, 1 + \delta).$$

Let a be a fixed vector in $\mathbb{R}^m$ and define the real-valued function F on $(-\delta, 1 + \delta)$ by

$$F(t) = a \cdot f(x + tu).$$

Then F is differentiable on $(-\delta, 1 + \delta)$ since $F = T \circ f$ where $T(x) = a \cdot x$ is a linear map, and note that

$$F'(t) = a \cdot \{f'(x + tu)(u)\}.$$

By the usual MVT for real-valued functions, we have

$$F(1) - F(0) = F'(\theta) \text{ for some } \theta \text{ with } 0 < \theta < 1.$$

Now

$$F'(\theta) = a \cdot \{f'(x + \theta u)(u)\} = a \cdot \{f'(z)(y - x)\},$$

where $z = x + \theta u \in L(x; y)$. But

$$F(1) - F(0) = a \cdot \{f(y) - f(x)\},$$

so we're done. ■

Remark 1.4.3. If S is convex, then $L(x; y) \subseteq S$ for all $x, y \in S$, so [Theorem 1.4.5](#) holds for all x and y in S .

1.5 Double Derivatives and Clairaut's theorem

Definition 1.5.1. Let $E \subseteq \mathbb{R}^n$ be open and let $f : E \rightarrow \mathbb{R}^m$. We say f is twice continuously differentiable on E if

- (1) f is continuously differentiable on E , i.e. $\frac{\partial f}{\partial x_i}$ exists and are continuous on E
- (2) for each $j = 1, 2, \dots, n$, the partial derivative $\frac{\partial f}{\partial x_j} : E \rightarrow \mathbb{R}^m$ is continuously differentiable on E .

Theorem 1.5.1 (Clairaut's theorem). Let E be an open set in $\mathbb{R}^n$, and let $f : E \rightarrow \mathbb{R}^m$ be twice continuously differentiable on E . Then, for all $1 \leq i, j \leq n$ and all $x_0 \in E$

$$\frac{\partial^2 f(x_0)}{\partial x_i \partial x_j} = \frac{\partial^2 f(x_0)}{\partial x_j \partial x_i}.$$

Proof. Since $f = (f_1, f_2, \dots, f_m)$ where $f_i : E \rightarrow \mathbb{R}$, we may just assume $f : E \rightarrow \mathbb{R}$. We may also assume $x_0 = 0$. The $i = j$ case is trivial. Given any $\varepsilon > 0$, $\exists \delta_0 > 0$ s.t.

$$\left| \frac{\partial^2 f(x)}{\partial x_i \partial x_j} - \frac{\partial^2 f(0)}{\partial x_i \partial x_j} \right| < \varepsilon \text{ and } \left| \frac{\partial^2 f(x)}{\partial x_j \partial x_i} - \frac{\partial^2 f(0)}{\partial x_j \partial x_i} \right| < \varepsilon$$

whenever $\|x\| \leq 2\delta_0$ since the second order derivative is continuous at 0. Fix $\delta \in (0, \delta_0)$. Consider the square in the (x_i, x_j) -plane with vertices $0, \delta e_i, \delta e_j, \delta e_i + \delta e_j$ and define the second finite difference

$$X := f(\delta e_i + \delta e_j) - f(\delta e_i) - f(\delta e_j) + f(0).$$

Define

$$\phi(t) := f(te_i + \delta e_j) - f(te_i), \quad t \in [0, \delta].$$

Then, $X = \phi(\delta) - \phi(0)$. By MVT, $\exists c_i \in (0, \delta)$ s.t.

$$X = \delta \phi'(c_i) = \delta \left[\frac{\partial f}{\partial x_i}(c_i e_i + \delta e_j) - \frac{\partial f}{\partial x_i}(c_i e_i) \right].$$

Now consider

$$g(s) := \frac{\partial f}{\partial x_i}(c_i e_i + s e_j), \quad s \in [0, \delta].$$

Then

$$g(\delta) - g(0) = \frac{\partial f}{\partial x_i}(c_i e_i + \delta e_j) - \frac{\partial f}{\partial x_i}(c_i e_i).$$

Applying MVT one more time, then we know there exists $c_j \in (0, \delta)$ s.t.

$$g(\delta) - g(0) = g'(c_j)\delta = \delta \frac{\partial^2 f}{\partial x_j \partial x_i}(c_i e_i + c_j e_j).$$

Substituting into the expression for X gives

$$X = \delta^2 \frac{\partial^2 f}{\partial x_j \partial x_i}(\xi), \quad \xi = c_i e_i + c_j e_j.$$

Since $\|\xi\| = \sqrt{c_i^2 + c_j^2} < \delta\sqrt{2} < 2\delta_0$, continuity implies

$$\left| \frac{X}{\delta^2} - \frac{\partial^2 f}{\partial x_j \partial x_i}(0) \right| \leq \varepsilon.$$

If we write X as

$$X = f(\delta e_i + \delta e_j) - f(\delta e_j) - f(\delta e_i) + f(0)$$

and define

$$\psi(s) := f(\delta e_i + s e_j) - f(s e_j), \quad s \in [0, \delta].$$

Then, we can use similar method to show

$$\left| \frac{X}{\delta^2} - \frac{\partial^2 f}{\partial x_i \partial x_j}(0) \right| \leq \varepsilon.$$

Thus, by triangle inequality, we know

$$\left| \frac{\partial^2 f}{\partial x_j \partial x_i}(0) - \frac{\partial^2 f}{\partial x_i \partial x_j}(0) \right| \leq 2\varepsilon$$

for any $\varepsilon > 0$, and we're done. ■

Lecture 5

1.6 The Contraction Mapping Theorem

10 March.

Definition 1.6.1. Let (X, d) be a metric, and let $f : X \rightarrow X$ be a map. We say f is a contraction if

$$d(f(x), f(y)) \leq d(x, y) \quad \text{for all } x, y \in X.$$

We say f is a strict contraction if there exists a constant $0 < c < 1$ s.t.

$$d(f(x), f(y)) \leq cd(x, y) \quad \text{for all } x, y \in X.$$

We call c a contraction constant of f .

Example 1.6.1. Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$,

(a) if $f(x) = x + 1$, then

$$f(x) - f(y) = (x + 1) - (y + 1) = x - y \Rightarrow |f(x) - f(y)| = |x - y|,$$

so f is a contraction but not a strict contraction.

(b) if $f(x) = \frac{x}{2}$, then

$$|f(x) - f(y)| = \frac{|x - y|}{2},$$

so f is a strict contraction.

Example 1.6.2. Suppose $f : [0, 1] \rightarrow [0, 1]$ and $f(x) = x - x^2$, then $f'(x) = 1 - 2x$, so

$$|f'(x)| \leq 1 \quad \text{on } [0, 1].$$

By MVT,

$$f(x) - f(y) = f'(c)(x - y) \text{ for all } x, y \in [0, 1] \text{ for some } c \in (x, y).$$

Now since $|f'(c)| \leq 1$, so we know f is a contraction. But f is not a strict contraction. Suppose it is a strict contraction, then

$$|f(x)| = |f(x) - f(0)| \leq c|x| \text{ for some } 0 < c < 1 \text{ for all } x \in [0, 1].$$

However,

$$\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = f'(0) = 1,$$

which is impossible.

Definition 1.6.2. Let $f : X \rightarrow X$ be a map and let $x \in X$. We say that x is a fixed point of f if $f(x) = x$.

Remark 1.6.1. Not every contraction has a fixed point.

Remark 1.6.2. Suppose $X \subseteq \mathbb{R}$. The existence of a fixed point x s.t. $f(x) = x$ means that the graph of $y = f(x)$ intersect with $y = x$.

Definition 1.6.3. A metric space (X, d) is said to be complete if every Cauchy sequence in X converges to a point of X .

Example 1.6.3. In $\mathbb{R}^n$, $\mathbb{R}^n$ is complete with the standard metric

$$d(x, y) = \|x - y\| = \sqrt{\sum_{i=1}^n (x_i - y_i)^2} \text{ with } x = (x_1, x_2, \dots, x_n), y = (y_1, y_2, \dots, y_n).$$

Remark 1.6.3. Recall that

- Any closed subset of a complete metric space is also a complete metric space.
- Any closed set in $\mathbb{R}^n$ is complete. In particular, a closed ball

$$\overline{B(x_0, r)} = \{y : \|y - x_0\| \leq r\}$$

is complete.

Theorem 1.6.1 (Contraction Mapping Theorem). Let (X, d) be a metric space and $f : X \rightarrow X$ be a strict contractor. Then f have at most one fixed point. Moreover, if we also assume X is non-empty and complete, then f has exactly one fixed point.

Proof. Suppose x and y are fixed points of a strict contractor f . Then, $f(x) = x$ and $f(y) = y$, and $\exists 0 < c < 1$ s.t.

$$d(x, y) = d(f(x), f(y)) \leq cd(x, y),$$

so $0 \leq (1 - c)d(x, y) \leq 0$, which shows $x = y$.

Now we assume X is non-empty and complete. Choose any point $x_0 \in X$. Consider the sequence $x_1 = f(x_0), x_2 = f(x_1), \dots$. For $n \geq 1$, we have

$$d(x_{n+1}, x_n) = d(f(x_n), f(x_{n-1})) \leq cd(x_n, x_{n-1}) \leq \dots \leq c^n d(x_1, x_0).$$

This shows $\{x_n\}_{n=0}^\infty$ is a Cauchy sequence. Now since X is complete, so $\lim_{n \rightarrow \infty} x_n = x^*$ for some $x^* \in X$. Now we claim $f(x^*) = x^*$. By triangle inequality and $f(x_{n-1}) = x_n$,

$$\begin{aligned} d(x^*, f(x^*)) &\leq d(x^*, f(x_n)) + d(f(x_n), f(x^*)) \leq d(x^*, x_{n+1}) + cd(x_n, x^*) \\ &\leq d(x^*, x_{n+1}) + d(x_n, x^*) \leq d(x_{n+1}, x_n) \rightarrow 0. \end{aligned}$$

■

Lemma 1.6.1. Let $B(0, r)$ be an open ball in $\mathbb{R}^n$, and let $g : B(0, r) \rightarrow \mathbb{R}^n$ be a map with $g(0) = 0$ and

$$\|g(x) - g(y)\| \leq \frac{1}{2}\|x - y\| \text{ for all } x, y \in B(0, r).$$

Then, the function $f : B(0, r) \rightarrow \mathbb{R}^n$ defined by $f(x) = x + g(x)$ is one to one and the image of $f(B(0, r))$ contains the ball $B(0, \frac{r}{2})$.

Proof. First, we show that f is one to one. Suppose $f(x) = f(y)$, then

$$x + g(x) = y + g(y) \Leftrightarrow x - y = g(y) - g(x) \Leftrightarrow \|x - y\| = \|g(x) - g(y)\| \leq \frac{1}{2}\|x - y\|,$$

so $x = y$.

Now we show $B(0, \frac{r}{2}) \subseteq f(B(0, r))$. It suffices to show that for any $y \in B(0, \frac{r}{2})$, $\exists x \in B(0, r)$ s.t. $f(x) = y$, or equivalently, $x = y - g(x)$. Define $F(x) = y - g(x)$, then we want to show that for any $y \in B(0, \frac{r}{2})$, $\exists x \in B(0, r)$ s.t. $F(x) = x$. Now $y \in B(0, \frac{r}{2})$, then $\exists \varepsilon > 0$ s.t. $2\|y\| \leq r - \varepsilon$.

Claim 1.6.1. $F : \overline{B(0, r - \varepsilon)} \rightarrow \overline{B(0, r - \varepsilon)}$ is a strict contraction.

Proof. We know

$$\|F(x)\| \leq \|y\| + \|g(x)\| \leq \frac{r - \varepsilon}{2} + \|g(x)\|.$$

Since $g(0) = 0$, so

$$\|g(x)\| = \|g(x) - g(0)\| \leq \frac{1}{2}\|x - 0\| \leq \frac{r - \varepsilon}{2}.$$

Thus,

$$\|F(x)\| \leq \frac{r - \varepsilon}{2} + \frac{r - \varepsilon}{2} = r - \varepsilon.$$

Thus, $F(x) \in \overline{B(0, r - \varepsilon)}$. Also,

$$\|F(x_1) - F(x_2)\| = \|g(x_1) - g(x_2)\| \leq \frac{1}{2}\|x_1 - x_2\|.$$

Thus, F is a strict contraction. ⊗

Now since closed ball is non-empty and complete, so by [Theorem 1.6.1](#), F has a fixed point. ■

Lecture 6

12 March.

1.7 Inverse Function Theorem

Recall that if $f : \mathbb{R} \rightarrow \mathbb{R}$ is invertible, i.e. $f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$ exists s.t. $f \circ f^{-1}(x) = f^{-1} \circ f(x) = x$ for all $x \in \mathbb{R}$, and $f'(x_0) \neq 0$ for some $x_0 \in \mathbb{R}$, then f^{-1} is differentiable at $f(x_0)$ and

$$(f^{-1})'(f(x_0)) = \frac{1}{f'(x_0)},$$

which can be proved by chain rule.

In fact, one can say something even when f' is not invertible, as long as we know that f is continuously differentiable. If $f'(x_0) \neq 0$, then $f'(x_0)$ must be either strictly positive or strictly negative (since we assume f' to be continuous by [Theorem 1.4.2](#)), which implies $f'(x)$ is strictly positive for x near x_0 , or strictly negative for x near x_0 , and thus one-to-one in such small interval. In either case, f will become invertible if we restrict the domain and codomain of f to be sufficiently close to x_0 and to $f(x_0)$, respectively.

Now if $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $f \in C^1$, i.e. f is continuously differentiable. Suppose

$$f'(x_0) = Df(x_0) = \left[\frac{\partial f_i(x_0)}{\partial x_j} \right]$$

is invertible. Then, $\exists$ a neighborhood U of x_0 and V of $f(x_0)$ s.t. $f : U \rightarrow V$ is bijective and its inverse $f^{-1} : V \rightarrow U$ is continuous and differentiable at $f(x_0)$ with

$$(f^{-1})'(f(x_0)) = [f'(x_0)]^{-1}.$$

This is the Inverse Function Theorem, and we will prove it later.

First, we prepare a lemma.


Lemma 1.7.1. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an invertible linear transformation. Then $T^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is also linear.

Proof.

DIY

Theorem 1.7.1 (The Inverse Function Theorem). Let E be an open subset of $\mathbb{R}^n$ and let $f : E \rightarrow \mathbb{R}^n$ be a function which is continuously differentiable on E . Suppose $x_0 \in E$ is the point s.t. the linear map $f'(x_0) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is invertible, then there exists an open set $U \subseteq E$ s.t. $x_0 \in U$ and an open set $V \subseteq \mathbb{R}^n$ s.t. $f(x_0) \in V$ so that $f : U \rightarrow V$ is a bijection. In particular, there is an inverse map $f^{-1} : V \rightarrow U$. Furthermore, this inverse map is differentiable at $f(x_0)$, and

$$(f^{-1})'(f(x_0)) = (f'(x_0))^{-1}.$$



Proof. Suppose such f^{-1} exists, i.e. $\exists U, V$ s.t. $f : U \rightarrow V$ and $f^{-1} : V \rightarrow U$ are bijective where $x_0 \in U$ and $f(x_0) \in V$. Since $f'(x_0)$ is invertible and $f^{-1}(f(x_0))$ exists, we know

$$f^{-1}(f(x)) = x \text{ for } x \in U \text{ and } (f^{-1})'(f(x_0)) \circ (f'(x_0)) = Id \text{ by chain rule.}$$

Now we have to show such U, V exists. Let $A = f'(x_0)$, then

$$\lim_{x \rightarrow 0} \frac{\|f(x_0 + x) - f(x_0) - A(x)\|}{\|x\|} = 0.$$

Note that A^{-1} exists and

$$\begin{aligned} A^{-1}(f(x_0 + x) - f(x_0) - Ax) &= A^{-1}(f(x_0 + x) - f(x_0)) - A^{-1}Ax \\ &= A^{-1}(f(x_0 + x) - f(x_0)) - x. \end{aligned}$$

Note that

$$\lim_{x \rightarrow 0} \frac{\|A^{-1}(f(x_0 + x) - f(x_0)) - x\|}{\|x\|} = 0$$

since

$$\begin{aligned} 0 &\leq \lim_{x \rightarrow 0} \frac{\|A^{-1}(f(x_0 + x) - f(x_0)) - x\|}{\|x\|} = \lim_{x \rightarrow 0} \frac{\|A^{-1}(f(x_0 + x) - f(x_0) - Ax) + A^{-1}Ax - x\|}{\|x\|} \\ &\leq \lim_{x \rightarrow 0} \frac{\|A^{-1}\| \|f(x_0 + x) - f(x_0) - Ax\|}{\|x\|} = 0. \end{aligned}$$

Let $g(x) = A^{-1}(f(x_0 + x) - f(x_0)) - x$. Thus,

$$\lim_{x \rightarrow 0} \frac{\|g(x)\|}{\|x\|} = 0.$$

Note that $g(0) = 0$. Thus,

$$\lim_{x \rightarrow 0} \frac{\|g(x) - g(0) - 0\|}{\|x\|} = 0,$$

which shows g is differentiable at 0 with derivative $g'(0) = 0$. Now let's analyze $g(x)$. Since f is continuously differentiable, g is also continuously differentiable. Since E is open and $x_0 \in E$, there exists $r_0 > 0$ s.t. $B(x_0, r_0) \subseteq E$. Hence, g is defined on $B(0, r_0)$. (For $x \in B(0, r_0)$, $\|x\| < r_0$, so $\|(x + x_0) - x_0\| < r_0$, i.e. $x + x_0 \in B(x_0, r_0) \subseteq E$, so $g(x)$ is well-defined.)

Thus, for all i, j , $\exists r > 0$ s.t.

$$\left| \frac{\partial g_j}{\partial x_i} \right| \leq \frac{1}{2\sqrt{n^5}} \text{ for } x \in B(0, r)$$

since $\frac{\partial g_j(0)}{\partial x_i} = 0$ (since $g'(0) = 0$ and thus g 's Jacobian matrix at 0 is 0 matrix) and $\frac{\partial g_j}{\partial x_i}$ is continuous, and hence

$$\left| \frac{\partial g_j(x)}{\partial x_i} - \frac{\partial g_j(0)}{\partial x_i} \right| \leq \frac{1}{2\sqrt{n^5}} \text{ if } \|x\| < r \text{ for some small enough } r.$$

By this, we can choose r small enough so that g is defined on $B(0, r)$ and

$$\left\| \frac{\partial g(x)}{\partial x_j} \right\| = \left(\sum_{i=1}^n \left(\frac{\partial g_i(x)}{\partial x_j} \right)^2 \right)^{\frac{1}{2}} \leq \frac{1}{2n^2} \text{ for all } x \in B(0, r), \quad j = 1, 2, \dots$$

Now if $x, y \in B(0, r)$, then $x + t(y - x) \in B(0, r)$ if $0 \leq t \leq 1$. Also,

$$\begin{aligned} g(y) - g(x) &= \int_0^1 \left(\frac{d}{dt} [g(x + t(y - x))] \right) dt = \int_0^1 [Dg(x + t(y - x)) \circ (y - x)] dt \\ &= \int_0^1 (D_{(y-x)}g(x + t(y - x))) dt. \end{aligned}$$

Now we estimate $D_v g(x)$ for $v = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$.

$$\|D_v g(x)\| = \left\| \sum_{j=1}^n v_j \frac{\partial g(x)}{\partial x_j} \right\| \leq \sum_{j=1}^n \|v_j\| \left\| \frac{\partial g(x)}{\partial x_j} \right\| \leq \sum_{j=1}^n \|v\| \frac{1}{2n^2} = \frac{\|v\|}{2n} \leq \frac{\|v\|}{2}.$$

Thus,

$$\|D_{(y-x)}g(x + t(y - x))\| \leq \frac{\|x - y\|}{2}.$$

Hence, we have

$$\|g(y) - g(x)\| = \left\| \int_0^1 (D_{(y-x)}g(x + t(y - x))) dt \right\| \leq \frac{\|x - y\|}{2}.$$

Recall $A^{-1}(f(x_0 + x) - f(x_0)) = g(x)$ and $g(0) = 0$, and we have shown

$$\|g(x) - g(y)\| \leq \frac{\|x - y\|}{2} \text{ for } x, y \in B(0, r).$$

Let $\Phi(x) = x + g(x)$. By [Lemma 1.6.1](#), $\Phi : B(0, r) \rightarrow \mathbb{R}^n$ is one to one and $B(0, \frac{r}{2}) \subseteq \Phi(B(0, r))$. Since Φ is one to one on $B(0, r)$. Let $W = \Phi^{-1}(B(0, \frac{r}{2}))$, then $\Phi : W \rightarrow B(0, \frac{r}{2})$ is one to one and onto. Now we know Φ is continuous (since g is composition of differentiable and thus continuous map, so g is continuous, and thus Φ is continuous), so W is open (preimage of open set is open for continuous map). Also, $0 \in W$ since $\Phi(0) = 0 \in B(0, \frac{r}{2})$. Now we claim $\Phi^{-1} : B(0, \frac{r}{2}) \rightarrow W$ is continuous. Note that for all $x, y \in W$

$$\|\Phi(x) - \Phi(y)\| = \|x - y + g(x) - g(y)\| \geq \|x - y\| - \|g(x) - g(y)\| \geq \|x - y\| - \frac{1}{2}\|x - y\| = \frac{\|x - y\|}{2}.$$

Let $x = \Phi^{-1}(u)$ and $y = \Phi^{-1}(v)$ for $u, v \in B(0, \frac{r}{2})$, then

$$\|\Phi(\Phi^{-1}(u)) - \Phi(\Phi^{-1}(v))\| \geq \frac{1}{2}\|\Phi^{-1}(u) - \Phi^{-1}(v)\|.$$

Thus,

$$\|u - v\| \geq \frac{1}{2}\|\Phi^{-1}(u) - \Phi^{-1}(v)\|.$$

This gives

$$\|\Phi^{-1}(u) - \Phi^{-1}(v)\| \leq 2\|u - v\|,$$

and implies Φ^{-1} is continuous. Thus, $\Phi : W \rightarrow B(0, \frac{r}{2})$ is a homeomorphism, i.e. Φ is continuous and Φ^{-1} is also continuous. Since $\Phi(0) = 0$ and Φ is one-to-one, $\Phi^{-1}(0) = 0$. Also, note that

$$\lim_{z \rightarrow 0} \frac{\|\Phi^{-1}(z) - \Phi^{-1}(0) - z\|}{\|z\|} = \lim_{z \rightarrow 0} \frac{\|\Phi^{-1}(z) - z\|}{\|z\|}.$$

Note that for $z \neq 0$, if we let $\Phi^{-1}(z) = x$, then $z = \Phi(x) = x + g(x)$, and $x \neq 0$, so

$$\frac{\|\Phi^{-1}(z) - z\|}{\|z\|} = \frac{\|x - (x + g(x))\|}{\|z\|} = \frac{\|g(x)\|}{\|z\|} = \frac{\|g(x)\|}{\|x\|} \cdot \frac{\|x\|}{\|z\|}.$$

Hence,

$$\lim_{z \rightarrow 0} \frac{\|\Phi^{-1}(z) - z\|}{\|z\|} = \lim_{z \rightarrow 0} \frac{\|g(x)\|}{\|x\|} \cdot \frac{\|x\|}{\|z\|} = \lim_{z \rightarrow 0} \frac{\|g(x)\|}{\|x\|} \cdot \frac{\|\Phi^{-1}(z)\|}{\|z\|}$$

Now if $z \rightarrow 0$, then since Φ^{-1} is continuous, so $x = \Phi^{-1}(z) \rightarrow \Phi^{-1}(0) = 0$. Also, when $z \rightarrow 0$, we know

$$\|\Phi^{-1}(z) - \Phi^{-1}(0)\| \leq 2\|z - 0\| \Rightarrow \frac{\|\Phi^{-1}(z)\|}{\|z\|} \leq 2,$$

and since

$$\lim_{z \rightarrow 0} \frac{\|g(x)\|}{\|x\|} = \lim_{x \rightarrow 0} \frac{\|g(x)\|}{\|x\|} = 0,$$

so by Squeeze theorem,

$$\lim_{z \rightarrow 0} \frac{\|\Phi^{-1}(z) - z\|}{\|z\|} = 0.$$

This shows $(\Phi^{-1})'(0) = I$. Recall that

$$\Phi(x) = x + g(x) = A^{-1}(f(x + x_0) - f(x_0))$$

and $\Phi : W \rightarrow B(0, \frac{r}{2})$ is a homomorphism. Now set

$$U = x_0 + W, \quad V = f(x_0) + A\left(B\left(0, \frac{r}{2}\right)\right).$$

Since W is open and A is a linear isomorphism, both U and V are open subsets of $\mathbb{R}^n$. Moreover, $x_0 \in U$ and $f(x_0) \in V$. We claim that f maps U bijectively to V . Let $u = x + x_0 \in U$, where $x \in W$, then since $\Phi(W) = B(0, \frac{r}{2})$, we have $\Phi(x) \in B(0, \frac{r}{2})$, and thus

$$f(u) = f(x + x_0) = f(x_0) + A\Phi(x) \in f(x_0) + A\left(B\left(0, \frac{r}{2}\right)\right) = V.$$

This shows $f(U) \subseteq V$. Conversely, let $y \in V$, then

$$A^{-1}(y - f(x_0)) \in B\left(0, \frac{r}{2}\right).$$

Because $\Phi : W \rightarrow B(0, \frac{r}{2})$ is surjective, there exists $x \in W$ s.t.

$$\Phi(x) = A^{-1}(y - f(x_0)).$$

For this x we have

$$f(x + x_0) = f(x_0) + A\Phi(x) = y.$$

This shows $V \subseteq f(U)$ and thus $f(U) = V$, i.e. $f : U \rightarrow V$ is surjective. Since Φ is one to one, so the x above is unique. It follows that f is one to one on U . Thus, $f : U \rightarrow V$ is bijective.

Now for $y \in V$ we know $y = f(x + x_0)$ for some $x \in W$, so $x + x_0 = f^{-1}(y)$. Recall

$$\Phi(x) = A^{-1}(f(x + x_0) - f(x_0)) = A^{-1}(y - f(x_0)).$$

Hence,

$$x = \Phi^{-1}(A^{-1}(y - f(x_0))) \Rightarrow f^{-1}(y) - x_0 = \Phi^{-1}(A^{-1}(y - f(x_0))).$$

Thus,

$$f^{-1}(y) = \Phi^{-1}(A^{-1}(y - f(x_0))) + x_0.$$

This shows

$$(f^{-1})'(f(x_0)) = (\Phi^{-1})'(A^{-1}(f(x_0) - f(x_0)))(A^{-1})'(0) = (\Phi^{-1})'(0) \circ A^{-1} = IA^{-1} = (f'(x_0))^{-1}$$

since $x_0 \in U$ so $f(x_0) \in V$ and every linear transformation is differentiable at every point where $(A^{-1})(x) = A^{-1}$ for all x . Thus, we're done. $\blacksquare$

Remark 1.7.1. Since $\det f'(x_0) \neq 0$ and $\det$ is continuous, there exists a neighborhood B of x_0 such that $f'(x)$ is invertible for all $x \in B$.

Applying the inverse function theorem at each point $x \in B$, we obtain neighborhoods U_x such that f^{-1} is differentiable on $f(U_x)$ and

$$(f^{-1})'(f(x)) = (f'(x))^{-1}.$$

Restrict to a sufficiently small neighborhood $U' \subset B$ of x_0 and set $V' = f(U')$. Then f^{-1} is differentiable on V' .

Moreover, for $y \in V'$,

$$(f^{-1})'(y) = (f'(f^{-1}(y)))^{-1}.$$

Since f^{-1} is continuous, f' is continuous, and matrix inversion is continuous on the set of invertible matrices, it follows that $(f^{-1})'$ is continuous on V' .

Hence, f^{-1} is continuously differentiable on V' .

Lecture 7

Remark 1.7.2. The requirement of $f'(x_0)$ is invertible is necessary to get the differentiability of f^{-1} at x_0 .

17 March.


1.8 The Implicit Function Theorem

In $\mathbb{R}^2$, $S = \{(x, y) \mid x^2 + y^2 = 1\}$ is a circle. However, S is not the graph of one function. S can be expressed as the union of the graph of four functions:

$$\begin{cases} y = \sqrt{1 - x^2} & \text{for } -1 < x < 1 \\ y = -\sqrt{1 - x^2} & \text{for } -1 < x < 1 \\ x = \sqrt{1 - y^2} & \text{for } -1 < y < 1 \\ x = -\sqrt{1 - y^2} & \text{for } -1 < y < 1. \end{cases}$$

For a $\mathbb{R}^3$ case, consider $S = \{(x, y, z) \mid x^2 + y^2 - z^2 + 1 = 0\}$, then S is the union of 2 graph:


$$\begin{cases} z = \sqrt{x^2 + y^2 + 1} & \text{for } z > 0 \\ z = -\sqrt{x^2 + y^2 + 1} & \text{for } z < 0. \end{cases}$$



For another example, consider $S = \{(x, y, z) \mid x^2 + y^2 = \cosh^2(z)\}$, where

$$\cosh(z) = \frac{e^z + e^{-z}}{2}.$$

Then S is symmetric with respect to z .



Theorem 1.8.1 (Implicit Function Theorem). Let E be an open subset of $\mathbb{R}^n$, let $f : E \rightarrow \mathbb{R}$ be continuously differentiable, and let $y = (y_1, y_2, \dots, y_n) \in E$ s.t. $f(y) = 0$ and $\frac{\partial f(y)}{\partial x_n} \neq 0$. Then, $\exists$ an open subset $U \subseteq \mathbb{R}^{n-1}$ containing $(y_1, y_2, \dots, y_{n-1})$, an open subset $V \subseteq E$ containing y , and a function $g : U \rightarrow \mathbb{R}$ s.t.

$$g(y_1, y_2, \dots, y_{n-1}) = y_n$$

and

$$\{x = (x_1, \dots, x_n) \in V : f(x) = 0\} = \{(x_1, \dots, x_{n-1}, g(x_1, \dots, x_{n-1})) : (x_1, \dots, x_{n-1}) \in U\}.$$

In other words, the set $\{x \in V : f(x) = 0\}$ is a graph of a function over U . Moreover, g is differentiable at $(y_1, \dots, y_{n-1})$ and we have

$$\frac{\partial g}{\partial x_j}(y_1, y_2, \dots, y_{n-1}) = -\frac{\frac{\partial f}{\partial x_j}(y)}{\frac{\partial f}{\partial x_n}(y)}$$

for all $1 \leq j \leq n-1$.

Proof. Define $F : E \rightarrow \mathbb{R}^n$ by

$$(x_1, x_2, \dots, x_n) \rightarrow (x_1, x_2, \dots, x_{n-1}, f(x_1, x_2, \dots, x_n)).$$

Since $f \in C^1(E)$, so $F \in C^1(E)$. Thus,

$$DF(x) = \begin{bmatrix} \nabla x_1 \\ \nabla x_2 \\ \vdots \\ \nabla x_{n-1} \\ \nabla f \end{bmatrix} = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ & & \ddots & \\ 0 & \cdots & 1 & 0 \\ \frac{\partial f}{\partial x_1}(x) & \cdots & \frac{\partial f}{\partial x_{n-1}}(x) & \frac{\partial f}{\partial x_n}(x) \end{bmatrix},$$

which gives

$$\det(DF(x)) = \frac{\partial f(x)}{\partial x_n}.$$

Hence,

$$\det(DF(y)) = \frac{\partial f(y)}{\partial x_n} \neq 0,$$

so $F'(y)$ is invertible. Now we can apply inverse function theorem, so $\exists$ an open set $V \subseteq E$ and $W \subseteq \mathbb{R}^m$ with $y \in V$ and $F(y) = (y_1, \dots, y_{n-1}, 0) \in W$ s.t. $F : V \rightarrow W$ is a C^1 diffeomorphism. Write the inverse in coordinate

$$F^{-1}(\underbrace{u_1, u_2, \dots, u_n}_u) = (h_1(u), h_2(u), \dots, h_n(u)).$$

Since $F(F^{-1}(u)) = u$ for $u \in W$, so

$$(h_1(u), \dots, h_{n-1}(u), f(h_1(u), \dots, h_n(u))) = F(h_1(u), h_2(u), \dots, h_n(u)) = (u_1, u_2, \dots, u_n).$$

Hence,

$$\begin{aligned} h_1(u) &= u_1 \\ h_2(u) &= u_2 \\ &\vdots \\ h_{n-1}(u) &= u_{n-1} \\ f(h_1(u), \dots, h_n(u)) &= f(u_1, u_2, \dots, u_{n-1}, h_n(u)) = u_n. \end{aligned}$$

Thus, for $u_1, \dots, u_n \in W$,

$$F^{-1}(u_1, \dots, u_n) = (u_1, \dots, u_{n-1}, h_n(u)).$$

By this, we can plug $u_n = 0$ into it and get

$$F^{-1}(u_1, \dots, u_{n-1}, 0) = (u_1, \dots, u_{n-1}, h_n(u_1, \dots, u_{n-1}, 0)),$$

where

$$0 = u_n = f(u_1, \dots, u_{n-1}, h_n(u_1, \dots, u_{n-1}, 0)).$$

Now we want to restrict $u_n = 0$ and define g . Set

$$U = \{x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}, (x', 0) \in W\}.$$

Then U is open and contains $(y_1, \dots, y_{n-1})$ since

$$F(y_1, \dots, y_n) = (y_1, \dots, y_{n-1}, f(y_1, \dots, y_n)) = (y_1, \dots, y_{n-1}, 0) \in W.$$

Define $g : U \rightarrow \mathbb{R}$ by

$$g(x') = g(x_1, \dots, x_{n-1}) = h_n(x_1, \dots, x_{n-1}, 0).$$

Since $f(y_1, \dots, y_n) = 0$, we know $F(y) = (y_1, \dots, y_n, 0)$. Hence,

$$(y_1, \dots, y_n) = y = F^{-1}(F(y)) = F^{-1}(y_1, \dots, y_n, 0) = (y_1, \dots, y_{n-1}, h_n(y_1, \dots, y_{n-1}, 0)).$$

Thus,

$$y_n = h_n(y_1, \dots, y_{n-1}, 0) = g(\underbrace{y_1, \dots, y_{n-1}}_{y'}) \Rightarrow g(y') = y_n.$$

Moreover, g is differentiable since h_n is differentiable.

Now we want to show the zero set is the graph of g .

Claim 1.8.1. $\{x \in V : f(x) = 0\} = \{(x', g(x')) : x' \in U\}.$

Proof. Let $A = \{x \in V : f(x) = 0\}$ and $B = \{(x', g(x')) : x' \in U\}$. We first show $A \subseteq B$. Let $x = (x', x_n) \in V$ and $f(x', x_n) = 0$. Then

$$F(x) = (x', f(x', x_n)) = (x', 0) \in W.$$

Since $x' \in U$,

$$x = F^{-1}(F(x)) = F^{-1}(x', 0) = (x', h_n(x', 0)) = (x', g(x')).$$

Hence, $A \subseteq B$.

Now we show $B \subseteq A$. Suppose $x' \in U$, then

$$x = (x', g(x')) = (x', h_n(x', 0)).$$

Thus,

$$F^{-1}(x', 0) = (x', h_n(x', 0)) = (x', g(x')).$$

Thus,

$$(x', 0) = F(F^{-1}(x', 0)) = F(x', g(x')) = (x', f(x', g(x'))).$$

Thus,

$$f(x', g(x')) = 0,$$

which shows $x \in A$. ⊗

Now for $(x_1, x_2, \dots, x_{n-1}) \in U$, we know

$$f(x_1, x_2, \dots, x_{n-1}, g(x_1, x_2, \dots, x_{n-1})) = 0.$$

Fix $j \in [n-1]$, and set $y' = (y_1, \dots, y_{n-1})$. Consider

$$\psi(t) := f(y' + te_j, g(y' + te_j)),$$

defined for t near 0. Note that $\psi(t) \equiv 0$ for all t sufficiently close to 0. Hence, $\psi'(0) = 0$. Since f is differentiable at $y = (y', g(y'))$ and g is differentiable at y' , we may apply the chain rule to compute $\psi'(0)$. Suppose $\psi(t) = f(\gamma(t))$, where $\gamma : \mathbb{R} \rightarrow \mathbb{R}^n$ is defined by

$$t \mapsto (y' + te_j, g(y' + te_j)).$$

Now since

$$\psi'(t) = f'(\gamma(t)) \circ \gamma'(t),$$

and f is a function from a subset of $\mathbb{R}^n$ to $\mathbb{R}$, so

$$f'(\gamma(t)) = \nabla f(\gamma(t)) = \left(\frac{\partial f(\gamma(t))}{\partial x_1}, \frac{\partial f(\gamma(t))}{\partial x_2}, \dots, \frac{\partial f(\gamma(t))}{\partial x_n} \right),$$

which gives

$$f'(\gamma(0)) = \left(\frac{\partial f(y)}{\partial x_1}, \frac{\partial f(y)}{\partial x_2}, \dots, \frac{\partial f(y)}{\partial x_n} \right).$$

Also,

$$\gamma'(t) = \left(0, \dots, 1, \dots, 0, \frac{dg(y' + te_j)}{dt} \right) \Rightarrow \gamma'(0) = \left(0, \dots, 1, \dots, 0, \frac{dg(y')}{dt} \right),$$

and note that

$$\frac{dg(y' + te_j)}{dt} = \nabla g(y' + te_j) \circ e_j \Rightarrow \frac{dg(y')}{dt} = \nabla g(y') \circ e_j = \frac{\partial g(y')}{\partial x_j}.$$

Hence,

$$\begin{aligned}\psi'(0) &= \left(\frac{\partial f(y)}{\partial x_1}, \frac{\partial f(y)}{\partial x_2}, \dots, \frac{\partial f(y)}{\partial x_n} \right) \circ \left(0, \dots, 1, \dots, 0, \frac{\partial g(y')}{\partial t} \right) \\ &= \frac{\partial f}{\partial x_j}(y) + \frac{\partial f}{\partial x_n}(y) \frac{\partial g}{\partial x_j}(y'),\end{aligned}$$

and thus

$$0 = \psi'(0) = \frac{\partial f}{\partial x_j}(y) + \frac{\partial f}{\partial x_n}(y) \frac{\partial g}{\partial x_j}(y').$$

Now since $\frac{\partial f}{\partial x_n}(y) \neq 0$, we have

$$\frac{\partial g}{\partial x_j}(y') = -\frac{\frac{\partial f}{\partial x_j}(y)}{\frac{\partial f}{\partial x_n}(y)}.$$

■

Example 1.8.1. Let $S = \{(x, y, z) \in \mathbb{R}^3 \mid xy + yz + xz = -1\}$, which we rewrite as

$$\{(x, y, z) \in \mathbb{R}^3 : f(x, y, z) = 0\},$$

where $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is the function

$$f(x, y, z) = xy + yz + xz + 1.$$

Clearly f is continuously differentiable, and $\frac{\partial f}{\partial z} = y + x$. Thus, for any $(x_0, y_0, z_0) \in S$ with $y_0 + x_0 \neq 0$, one can write this surface (near (x_0, y_0, z_0)) as a graph of the form

$$\{(x, y, g(x, y)) : (x, y) \in U\}$$

for some open set U containing (x_0, y_0) , and some function g which is differentiable at (x_0, y_0) . Indeed one can implicitly obtain

$$\frac{\partial g}{\partial x}(x_0, y_0) = -\frac{y_0 + z_0}{y_0 + x_0}, \quad \text{and} \quad \frac{\partial g}{\partial y}(x_0, y_0) = -\frac{x_0 + z_0}{y_0 + x_0}.$$

Chapter 2

Lebesgue measure

Lecture 8

To overcome the limitations of the Riemann integral, particularly its inability to handle pointwise limits of functions, this chapter constructs a more robust theory of integration starting with the concept of measure. 19 March.

2.1 The goal: Lebesgue measure

Our goal is to define a concept of measurable set, which will be a special kind of subset of $\mathbb{R}^n$. For every such measurable set $\Omega \subseteq \mathbb{R}^n$, we'll define the Lebesgue measure $m(\Omega)$ to be a number in $[0, \infty]$.

Definition 2.1.1 (Measurable set). The concept of measurable set will obey the following properties:

- (i) **(Borel property)** Every open set in $\mathbb{R}^n$ is measurable, as is every closed set.
- (ii) **(Complementarity)** If Ω is measurable, then $\mathbb{R}^n \setminus \Omega$ is also measurable.
- (iii) **(Boolean algebra property)** If $(\Omega_j)_{j \in J}$ is any finite collection of measurable sets, then $\bigcup_{j \in J} \Omega_j$ and $\bigcap_{j \in J} \Omega_j$ are also measurable.
- (iv) **(σ -algebra property)** If $(\Omega_j)_{j \in J}$ are any countable collection of measurable sets (so J is countable), then $\bigcup_{j \in J} \Omega_j$ and $\bigcap_{j \in J} \Omega_j$ are also measurable.

Note that some of these properties are redundant; for instance, (iv) will imply (iii), and once one knows all open sets are measurable, (ii) will imply that all closed sets are measurable also. The properties (i)-(iv) will ensure that virtually every set one cares about is measurable; though there do exist non-measurable sets.

Definition 2.1.2 (Lebesgue measure). To every measurable set Ω , we associate the Lebesgue measure $m(\Omega)$ of Ω , which will obey the following properties:

- (v) **(Empty set)** The empty set $\emptyset$ has measure $m(\emptyset) = 0$.
- (vi) **(Positivity)** We have $0 \leq m(\Omega) \leq +\infty$ for every measurable set Ω .
- (vii) **(Monotonicity)** If $A \subseteq B$, and A and B are both measurable, then $m(A) \leq m(B)$.
- (viii) **(Finite sub-additivity)** If $(A_j)_{j \in J}$ are a finite collection of measurable sets, then

$$m\left(\bigcup_{j \in J} A_j\right) \leq \sum_{j \in J} m(A_j).$$

(ix) **(Finite additivity)** If $(A_j)_{j \in J}$ are a finite collection of *disjoint* measurable sets, then

$$m\left(\bigcup_{j \in J} A_j\right) = \sum_{j \in J} m(A_j).$$

(x) **(Countable sub-additivity)** If $(A_j)_{j \in J}$ are a countable collection of measurable sets, then

$$m\left(\bigcup_{j \in J} A_j\right) \leq \sum_{j \in J} m(A_j).$$

(xi) **(Countable additivity)** If $(A_j)_{j \in J}$ are a countable collection of *disjoint* measurable sets, then

$$m\left(\bigcup_{j \in J} A_j\right) = \sum_{j \in J} m(A_j).$$

(xii) **(Normalization)** The unit cube

$$[0, 1]^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n : 0 \leq x_j \leq 1 \text{ for all } 1 \leq j \leq n\}$$

has measure $m([0, 1]^n) = 1$.

(xiii) **(Translation invariance)** If Ω is a measurable set, and $x \in \mathbb{R}^n$, then $x + \Omega := \{x + y : y \in \Omega\}$ is also measurable, and $m(x + \Omega) = m(\Omega)$.

Again, many of these properties are redundant; for instance, the countable additivity property can be used to deduce the finite additivity property, which in turn can be used to derive monotonicity (when combined with the positivity property). One can also obtain the sub-additivity properties from the additivity ones. Note that $m(\Omega)$ can be $+\infty$, and so in particular some of the sums in the above properties may also equal $+\infty$; in this chapter we adopt the convention that an infinite sum $\sum_{j \in J} a_j$ of non-negative quantities a_j is equal to $+\infty$ if the sum is not absolutely convergent. (Since everything is non-negative we will never have to deal with indeterminate forms such as $(-\infty) + (+\infty)$.)

Our goal for this chapter can then be stated:

Theorem 2.1.1 (Existence of Lebesgue measure). There exists a concept of a measurable set and a way to assign a number $m(\Omega)$ to every measurable set $\Omega \subseteq \mathbb{R}^n$, which obeys all of the properties (i) to (xiii).

Lecture 9

2.2 First attempt: Outer Measure

24 March.

Definition 2.2.1. An open box B in $\mathbb{R}^n$ is any set of the form

$$B = \prod_{i=1}^n (a_i, b_i) := \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_i \in (a_i, b_i) \text{ for all } 1 \leq i \leq n\},$$

where $b_i \geq a_i$ are real numbers. We define the volume $\text{vol}(B)$ of this box to be the number

$$\text{vol}(B) := \prod_{i=1}^n (b_i - a_i).$$

Definition 2.2.2. Let $\Omega \subseteq \mathbb{R}^n$ be a subset of $\mathbb{R}^n$. We say a collection $(B_j)_{j \in J}$ of boxes cover Ω iff $\Omega \subseteq \bigcup_{j \in J} B_j$.

Remark 2.2.1. An index set J is countable if it is finite or $\exists$ a one-to-one map $f : J \rightarrow \mathbb{N}$ where $\mathbb{N} = \{1, 2, \dots\}$ is the set of natural numbers.

Remark 2.2.2. The set of rational numbers is countable. We usually denote it by

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid q \neq 0, p, q \in \mathbb{Z} \right\}.$$

Definition 2.2.3 (Outer measure). If Ω is a set in $\mathbb{R}^n$, we define the outer measure $m^*(\Omega)$ of Ω to be the quantity

$$m^*(\Omega) := \inf \left\{ \sum_{j \in J} \text{vol}(B_j) : (B_j)_{j \in J} \text{ is a collection of boxes covers } \Omega \text{ and } J \text{ at most countable.} \right\}.$$

Note that $m^*(\Omega)$ may be ∞ . Also, since $\text{vol}(B_j) \geq 0$, so $m^*(\Omega) \geq 0$.

Lemma 2.2.1. The space $\mathbb{R}^n$ can be covered by countably many translates of the open cube $\left(-\frac{1}{2}, \frac{1}{2}\right)^n$.

Proof. For each $m \in \mathbb{Z}^n$, define

$$Q_m := \frac{m}{2} + \left(-\frac{1}{2}, \frac{1}{2}\right)^n.$$

We claim $\mathbb{R}^n = \bigcup_{m \in \mathbb{Z}^n} Q_m$. Let $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. For each i , set $m_i := \lfloor 2x_i \rfloor \in \mathbb{Z}$, then

$$\frac{m_i}{2} \leq x_i < \frac{1}{2} + \frac{m_i}{2} \Rightarrow x \in \frac{m}{2} + \left(-\frac{1}{2}, \frac{1}{2}\right)^n.$$

Note that $\mathbb{Z}^n$ is countable since $\mathbb{Z}$ is countable, and thus we're done. ■

Lemma 2.2.2 (Properties of outer measure). Outer measure has the following 6 properties:

- (v) **(Empty set)** The empty set $\emptyset$ has outer measure 0, i.e. $m^*(\emptyset) = 0$.
- (vi) **(Positivity)** We have $0 \leq m^*(\Omega) \leq \infty$ for all measurable set Ω .
- (vii) **(Monotocity)** If $A \subseteq B \subseteq \mathbb{R}^n$, then $m^*(A) \leq m^*(B)$.
- (viii) **(Finite sub-additivity)** If $(A_j)_{j \in J}$ are finite collection of subsets of $\mathbb{R}^n$, then

$$m^* \left(\bigcup_{j \in J} A_j \right) \leq \sum_{j \in J} m^*(A_j).$$

- (x) **(Countable sub-additivity)** If $(A_j)_{j \in J}$ are countable collection of subsets of $\mathbb{R}^n$, then

$$m^* \left(\bigcup_{j \in J} A_j \right) \leq \sum_{j \in J} m^*(A_j).$$

- (xiii) **(Translation invariance)** If Ω is a subset of $\mathbb{R}^n$, and $x \in \mathbb{R}^n$, then $m^*(x + \Omega) = m^*(\Omega)$.

proof of (v). From the definition we know $m^*(\Omega) \geq 0$ for all sets $\Omega \subseteq \mathbb{R}^n$. Hence, $m^*(\emptyset) \geq 0$. Now given $\varepsilon > 0$, we know $\left(0, \varepsilon^{\frac{1}{n}}\right)^n$ is an open box that covers $\emptyset$. Thus,

$$m^*(\emptyset) \leq \text{vol} \left(\left(0, \varepsilon^{\frac{1}{n}}\right)^n \right) = \varepsilon.$$

Thus, $m^*(\Omega) = 0$. ■

proof of (vi). It follows from the definition of m^* . ■

proof of (vii). If $m^*(B) = \infty$, then this is trivial. Now suppose $m^*(B) < \infty$. Recall

$$m^*(B) = \inf \left\{ \sum_{j \in J} \text{vol}(B_j) : (B_j)_{j \in J} \text{ are open boxes cover } B \right\}.$$

Also, for any countable collection of boxes covering B , say $(B_j)_{j \in J}$, it also covers A , so

$$\left\{ \sum_{j \in J} \text{vol}(B_j) : (B_j)_{j \in J} \text{ are open boxes cover } B \right\} \subseteq \left\{ \sum_{j \in J} \text{vol}(A_j) : (A_j)_{j \in J} \text{ are open boxes cover } A \right\},$$

which shows

$$m^*(A) \leq m^*(B).$$

proof of (x). If one of $m^*(A_j) = \infty$, then ■

$$\infty = m^*(A_j) \leq m^* \left(\bigcup_{j \in J} A_j \right),$$

so we're done. Now we assume $m^*(A_j) < \infty$ for all $j \in J$. Given $\varepsilon > 0$, $\exists$ open boxes $B_{j,k}$ s.t. $A_j \subseteq \bigcup_{k=1}^{\infty} B_{j,k}$ and

$$\sum_{k=1}^{\infty} \text{vol}(B_{j,k}) \leq m^*(A_j) + \frac{\varepsilon}{2^j}$$

since the definition of m^* is infimum. Now since J is countable, so we can re-label it to $\mathbb{N}$. Hence,

$$\bigcup_{j=1}^{\infty} A_j \subseteq \bigcup_{j=1}^{\infty} \bigcup_{k=1}^{\infty} B_{j,k},$$

and note that R.H.S. is the union of a collection of boxes, so $\bigcup_{j=1}^{\infty} A_j$ is covered by them. This shows

$$m^* \left(\bigcup_{j=1}^{\infty} A_j \right) \leq \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \text{vol}(B_{j,k}) \leq \sum_{j=1}^{\infty} m^*(A_j) + \frac{\varepsilon}{2^j} = \sum_{j=1}^{\infty} m^*(A_j) + \varepsilon.$$

Now since this is true for all $\varepsilon > 0$, so

$$m^* \left(\bigcup_{j=1}^{\infty} A_j \right) \leq \sum_{j=1}^{\infty} m^*(A_j).$$

Proposition 2.2.1 (Outer measure of closed box). For any closed box

$$B = \prod_{i=1}^n [a_i, b_i] = \{(x_1, x_2, \dots, x_n) \mid a_i \leq x_i \leq b_i \text{ for } 1 \leq i \leq n\},$$

we have

$$m^*(B) = \prod_{i=1}^n (b_i - a_i).$$

Proof. Write

$$\text{vol}(B) = \prod_{i=1}^n (b_i - a_i).$$

We first show that

$$m^*(B) \leq \text{vol}(B).$$

Let $\varepsilon > 0$, then the box

$$U_\varepsilon := \prod_{i=1}^n (a_i - \varepsilon, b_i + \varepsilon)$$

contains B . Thus,

$$m^*(B) \leq \text{vol}(U_\varepsilon) = \prod_{i=1}^n (b_i - a_i + 2\varepsilon).$$

Take $\varepsilon \rightarrow 0$, then we can show

$$m^*(B) \leq \text{vol}(B).$$

Now we show

$$m^*(B) \geq \prod_{i=1}^n (b_i - a_i).$$

Claim 2.2.1. For any countable family of open boxes $\{B_j\}_{j \in J}$ covering B . We have

$$\text{vol}(B) \leq \sum_{j \in J} \text{vol}(B_j).$$

Proof. Since B is compact, $\exists$ a finite subcover of $(B_j)_{j \in J}$

$$(U_j)_{j=1}^N \text{ s.t. } B \subseteq \bigcup_{j=1}^N U_j.$$

Thus,

$$\sum_{j=1}^N \text{vol}(U_j) \leq \sum_{j \in J} \text{vol}(B_j).$$

Choose a closed box Q s.t. $B \subseteq Q$ and $U_k \subseteq Q$ for $1 \leq k \leq N$. Define functions on Q by

$$f(x) = \begin{cases} 1, & \text{if } x \in B; \\ 0, & \text{if } x \in Q \setminus B. \end{cases} \quad f_k(x) = \begin{cases} 1, & \text{if } x \in U_k; \\ 0, & \text{if } x \in Q \setminus U_k. \end{cases}$$

Thus,

$$\int_Q f(x) \, dx = \text{vol}(B) \text{ and } \int_Q f_k(x) \, dx = \text{vol}(U_k).$$

Note that this is integrable since $B, (U_k)_{k=1}^N$ are all boxes and Q is also a box. Now since

$$f(x) \leq \sum_{j=1}^k f_j(x) \text{ on } Q.$$

(If $x \in B$, then $x \in U_j$ for some j .) Thus,

$$\text{vol}(B) = \int_Q f(x) \, dx \leq \int_Q \sum_{j=1}^k f_j(x) \, dx = \sum_{j=1}^k \int_Q f_j(x) \, dx = \sum_{j=1}^k \text{vol}(U_j) \leq \sum_{j \in J} \text{vol}(B_j).$$

⊛

This implies $\text{vol}(B)$ is a lower bound of

$$\left\{ \sum_{j \in J} \text{vol}(B_j) : (B_j)_{j \in J} \text{ are boxes covering } B \right\},$$

so by the definition of m^* ,

$$m^*(B) \geq \text{vol}(B).$$

■

Lecture 10

Corollary 2.2.1. For any open box

$$B = \prod_{i=1}^n (a_i, b_i) = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid a_i < x_i < b_i \text{ for } 1 \leq i \leq n\},$$

we have $m^*(B) = \prod_{i=1}^n (b_i - a_i)$.

Proof. First since B cover itself, we have $m^*(B) \leq \text{vol}(B) = \prod_{i=1}^n (b_i - a_i)$. Choose $\varepsilon_i < \frac{b_i - a_i}{2}$. Then,

$$[a_i + \varepsilon_i, b_i - \varepsilon_i] \subseteq (a_i, b_i).$$

Hence,

$$\prod_{i=1}^n [a_i + \varepsilon_i, b_i - \varepsilon_i] \subseteq \prod_{i=1}^n (a_i, b_i),$$

which shows

$$m^* \left(\prod_{i=1}^n [a_i + \varepsilon_i, b_i - \varepsilon_i] \right) \leq m^*(B),$$

and by previous proposition, we have

$$\prod_{i=1}^n (b_i - a_i - 2\varepsilon_i) = m^* \left(\prod_{i=1}^n [a_i + \varepsilon_i, b_i - \varepsilon_i] \right) \leq m^*(B).$$

Now pick all $\varepsilon_i \rightarrow 0$, then we have $m^*(B) \geq \text{vol}(B)$. ■

Example 2.2.1. Let $r > 0$, then $(-r, r) \subseteq \mathbb{R}$, and thus

$$2r = m^*((-r, r)) \leq m^*(\mathbb{R})$$

for all r , so $m^*(\mathbb{R}) = \infty$. Similarly, $m^*(\mathbb{R}^n) = \infty$.

Example 2.2.2.

$$\mathbb{Q} = \text{the set of rational numbers} = \bigcup_{q \in \mathbb{Q}} \{q\},$$

and $\mathbb{Q}$ is countable, so

$$m^*(\mathbb{Q}) \leq \sum_{q \in \mathbb{Q}} m^*(\{q\}) = 0.$$

In particular, $\exists$ an countable family of open boxes (B_j) s.t. $\mathbb{Q} \subseteq \bigcup_{j \in J} B_j$ and

$$\sum_{j \in J} \text{vol}(B_j) \leq m^*(\mathbb{Q}) + \varepsilon = \varepsilon.$$

Example 2.2.3. Since $\mathbb{R} = (\mathbb{R} \setminus \mathbb{Q}) \cup \mathbb{Q}$, we have

$$m^*(\mathbb{R}) \leq m^*(\mathbb{R} \setminus \mathbb{Q}) + m^*(\mathbb{Q}),$$

and since $m^*(\mathbb{R}) = \infty$ and $m^*(\mathbb{Q}) = 0$, we know $m^*(\mathbb{R} \setminus \mathbb{Q}) = \infty$.

Claim 2.2.2. $m^*([0, 1] \setminus \mathbb{Q}) = 1$.

Proof. Since

$$[0, 1] = ([0, 1] \setminus \mathbb{Q}) \cup ([0, 1] \cap \mathbb{Q}),$$

we have

$$1 = m^*([0, 1]) \leq m^*([0, 1] \setminus \mathbb{Q}) + m^*([0, 1] \cap \mathbb{Q}) = m^*([0, 1] \setminus \mathbb{Q})$$

since

$$[0, 1] \cap \mathbb{Q} \subseteq \mathbb{Q} \text{ and } 0 \leq m^*([0, 1] \cap \mathbb{Q}) \leq m^*(\mathbb{Q}) = 0.$$

Now since

$$[0, 1] \setminus \mathbb{Q} \subseteq [0, 1],$$

so

$$m^*([0, 1] \setminus \mathbb{Q}) \leq m^*([0, 1]) = 1.$$

Thus,

$$m^*([0, 1] \setminus \mathbb{Q}) = 1.$$

⊛

Claim 2.2.3. Let

$$L = \{(x, 0) \mid x \in \mathbb{R}\} \subseteq \mathbb{R}^2,$$

then $m^*(L) = 0$.

Proof. Let $L_n = \{(x, 0) \mid -n \leq x \leq n\} \subseteq \mathbb{R}^2$, then

$$L_n \subseteq [-n, n] \times [-\varepsilon, \varepsilon],$$

so

$$0 \leq m^*(L_n) \leq (2n) \cdot (2\varepsilon).$$

Let $\varepsilon \rightarrow 0$, then we have

$$m^*(L_n) = 0.$$

Now since

$$L = \bigcup_{n=1}^{\infty} L_n,$$

so

$$m^*(L) \leq \sum_{n=1}^{\infty} m^*(L_n) = 0,$$

which shows $m^*(L) = 0$.

⊛

2.3 Outer measure is not additive

We only know that

$$m^*\left(\bigcup_{j \in J} A_j\right) \leq \sum_{j \in J} m^*(A_j)$$

if J is countable. To construct a measure, we require that if $A_i \cap A_j = \emptyset$ for all $i, j \in J$ and $i \neq j$,

$$m^* \left(\bigcup_{j \in J} A_j \right) = \sum_{j \in J} m^*(A_j).$$

Proposition 2.3.1 (Failure of countable additivity). There exists a countable collection $(A_j)_{j \in J}$ of disjoint subsets of $\mathbb{R}$ s.t.

$$m^* \left(\bigcup_{j \in J} A_j \right) \neq \sum_{j \in J} m^*(A_j).$$

Proof. We shall construct such a family from the coset of $\mathbb{Q}$ in $\mathbb{R}$.

- Step 1. Coset of $\mathbb{Q}$. For $x \in \mathbb{R}$, let

$$x + \mathbb{Q} = \{x + q : q \in \mathbb{Q}\}.$$

This is called a coset of $\mathbb{Q}$. Two properties of cosets:

- (1) $(x + \mathbb{Q}) \cap (y + \mathbb{Q}) \neq \emptyset$ implies $x + \mathbb{Q} = y + \mathbb{Q}$.
- (2) Every coset meets the interval $[0, 1]$. (We can pick $-x \leq q \leq 1 - x$ so that $0 \leq q + x \leq 1$)
- Step 2. Choose one representative from each coset. Let $\mathbb{R}/\mathbb{Q}$ denote the collection of all cosets. For each coset $A \in \mathbb{R}/\mathbb{Q}$. By (2) in Step 1, there exists $x_A \in [0, 1]$ s.t. $x_A \in A \cap [0, 1]$ (By axiom of choice), we can choose one and define

$$E = \{x_A : A \in \mathbb{R}/\mathbb{Q}, x_A \in [0, 1]\} \subseteq [0, 1].$$

- Step 3. Rational translates of E . Consider the set

$$X := \bigcup_{q \in \mathbb{Q} \cap [-1, 1]} (q + E).$$

Since $E \subseteq [0, 1]$ and $-1 \leq q \leq 1$, we have $X \subseteq [-1, 2]$. Now we claim $[0, 1] \subseteq X$. Let y be any number in $[0, 1]$. Let $y \in [0, 1]$, then y is in some coset of $\mathbb{R}/\mathbb{Q}$. Thus, $\exists$ cosets A and $x_A \in E \subseteq [0, 1]$ s.t. $y \in A = x_A + \mathbb{Q} = y + \mathbb{Q}$. Let $y = x_A + q$ for some $q \in \mathbb{Q}$. Thus, $q = y - x_A \in [-1, 1]$. Thus, $q \in \mathbb{Q} \cap [-1, 1]$. Hence,

$$y = q + x_A \in q + E \subseteq X.$$

Thus,

$$[0, 1] \subseteq X \subseteq [-1, 2].$$

- Step 4. the sets $q + E$ are pairwise disjoint. We now show that the family

$$\{q + E\}_{q \in \mathbb{Q} \cap [-1, 1]}$$

is pairwise disjoint. Suppose, for contradiction, there are distinct $q, q' \in \mathbb{Q} \cap [-1, 1]$ we had

$$(q + E) \cap (q' + E) \neq \emptyset.$$

Then, there exists cosets $A, A' \in \mathbb{R}/\mathbb{Q}$ s.t.

$$q + x_A = q' + x_{A'}.$$

Hence,

$$x_{A'} - x_A = q - q' \in \mathbb{Q}.$$

Now since $x_A \in A = x_A + \mathbb{Q}$ and $x_{A'} \in A' = x_{A'} + \mathbb{Q}$, so $A = A'$. But E contains exactly one representative from each coset, which means $x_A = x_{A'}$. Thus, $q = q'$, which contradicts to the assumption, so the sets $q + E$ are pairwise disjoint.

- Step 5. Comparing the two sides. Now since $[0, 1] \subseteq X \subseteq [-1, 2]$,

$$1 = m^*([0, 1]) \leq m^*(X) \leq m^*([-1, 2]) = 3.$$

On the other hand,

$$m^*(q + E) = m^*(E) \text{ for every } q \in \mathbb{Q} \cap [-1, 1].$$

Thus,

$$\sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(q + E) = \sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(E).$$

Note that

$$\sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(E) = \begin{cases} 0, & \text{if } m^*(E) = 0; \\ \infty, & \text{if } m^*(E) > 0. \end{cases}$$

In particular, it cannot lie in the interval $[1, 3]$. But $m^*(X) \in [1, 3]$. Hence,

$$m^*(X) \neq \sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(q + E).$$

Since $\mathbb{Q} \cap [-1, 1]$ is countably infinite, so we're done. ■

Proposition 2.3.2. There exists a finite collection $(A_j)_{j \in J}$ of disjoint subsets of $\mathbb{R}$ s.t.

$$m^*\left(\bigcup_{j \in J} A_j\right) \neq \sum_{j \in J} m^*(A_j).$$

Proof. Suppose m^* was finitely additive, and let E and X to be same as previous proposition, then

$$m^*(X) \leq \sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(q + E) = \sum_{q \in \mathbb{Q} \cap [-1, 1]} m^*(E).$$

Since we know that $1 \leq m^*(X) \leq 3$, we thus have $m^*(E) \neq 0$, or otherwise we will have $m^*(X) \leq 0$, but we have $m^*(X) \in [1, 3]$.

Since $m^*(E) > 0$, so $m^*(E) > \frac{1}{n}$ for some $n \in \mathbb{N}$. Now let J be a finite subset of $\mathbb{Q} \cap [-1, 1]$ of cardinality $3n$. If m^* were finitely additive, then we would have

$$m^*\left(\bigcup_{j \in J} (q + E)\right) = \sum_{q \in J} m^*(q + E) = \sum_{q \in J} m^*(E) > 3n \cdot \frac{1}{n} = 3.$$

But we know $\bigcup_{j \in J} (q + E)$ is a subset of X , so it has outer measure at most 3, so we have a contradiction. ■

Lecture 11

31 March.

2.4 Measurable sets

In this chapter, our goal is to construct a measure. Note that we constructed outer measure in previous lecture, but countable additivity fails on it.

Definition 2.4.1 (Lebesgue measurability). Let $E \subseteq \mathbb{R}^n$. We say E is Lebesgue measurable, or measurable, iff we have the identity

$$m^*(A) = m^*(A \cap E) + m^*(A \setminus E) \text{ for all } A \subseteq \mathbb{R}^n.$$

Notation. If E is measurable, then we define $m(E) = m^*(E)$.

Remark 2.4.1. We know $A = (A \cap E) \cup (A \setminus E)$, so

$$m^*(A) \leq m^*(A \cap E) + m^*(A \setminus E).$$

Thus, E is measurable iff

$$m^*(A) \geq m^*(A \cap E) + m^*(A \setminus E).$$

Note that $\mathbb{R}^n$ is measurable since $A \cap \mathbb{R}^n = A$ and $A \setminus \mathbb{R}^n = \emptyset$. Also, $\emptyset$ is measurable since $A \cap \emptyset = \emptyset$ and $A \setminus \emptyset = A$.

Exercise 2.4.1. If A is an open box in $\mathbb{R}^n$ and $E = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : x_n > 0\}$, then

$$m^*(A) = m^*(A \cap E) + m^*(A \setminus E).$$

Proof. We first give a claim:

Claim 2.4.1. If A is an open interval in $\mathbb{R}$, then

$$m^*(A) = m^*(A \cap (0, \infty)) + m^*(A \setminus (0, \infty)).$$

Proof. Suppose $A = (a_i, b_i)$ s.t. $a_i \leq b_i$, then

- Case 1: $0 \leq a_i \leq b_i$, then $A \cap (0, \infty) = A$ and $A \setminus (0, \infty) = \emptyset$, so this is true.
- Case 2: $a_i \leq 0 \leq b_i$. Similar.
- Case 3: $a_i \leq b_i \leq 0$. Similar.

⊗

Suppose $A = \prod_{i=1}^n I_i$. Note that

$$m^*(A) = \text{vol}(A) = \prod_{i=1}^n \text{vol}(I_i) = \prod_{i=1}^n m_1^*(I_i).$$

Also,

$$A \cap E = \left(\prod_{i=1}^{n-1} I_i \right) \times (I_n \cap (0, \infty)) \quad A \setminus E = \left(\prod_{i=1}^{n-1} I_i \right) \times (I_n \setminus (0, \infty)).$$

Thus, $A \cap E$ and $A \setminus E$ are both open boxes. By the claim, we have

$$m_1^*(I_n) = m_1^*(I_n \cap (0, \infty)) + m_1^*(I_n \setminus (0, \infty)).$$

Thus,

$$\begin{aligned}
 m^*(A) &= \prod_{i=1}^n m_1^*(I_i) = \prod_{i=1}^{n-1} m_1^*(I_i) \times m_1^*(I_n) = \prod_{i=1}^{n-1} m_1^*(I_i) \times (m_1^*(I_n \cap (0, \infty)) + m_1^*(I_n \setminus (0, \infty))) \\
 &= \prod_{i=1}^{n-1} m_1^*(I_i) \times m_1^*(I_n \cap (0, \infty)) + \prod_{i=1}^{n-1} m_1^*(I_i) \times m_1^*(I_n \setminus (0, \infty)) \\
 &= m^*(A \cap E) + m^*(A \setminus E).
 \end{aligned}$$

■

Lemma 2.4.1 (Half-spaces are measurable). The half space $\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_n > 0\}$ is measurable.

Proof. Let $E = \{x \in \mathbb{R}^n \mid x_n > 0\}$. Then

$$m^*(A) \leq m^*(A \cap E) + m^*(A \setminus E)$$

is true for all $A \subseteq \mathbb{R}^n$, so it suffices to show that

$$m^*(A) \geq m^*(A \cap E) + m^*(A \setminus E) \text{ for all } A \subseteq \mathbb{R}^n.$$

If $m^*(A) = \infty$, then this is obviously true. Now we assume $m^*(A) < \infty$. Given $\varepsilon > 0$, $\exists$ a countable collection of open boxes (B_k) s.t. $A \subseteq \bigcup_{k=1}^{\infty} B_k$ and

$$\sum_{k=1}^{\infty} \text{vol}(B_k) \leq m^*(A) + \varepsilon.$$

We know that

$$m^*(B_k) = m^*(B_k \cap E) + m^*(B_k \setminus E) \text{ by Exercise 2.4.1.}$$

Thus,

$$\text{vol}(B_k) = m^*(B_k \cap E) + m^*(B_k \setminus E).$$

This gives

$$\begin{aligned}
 \sum_{k=1}^{\infty} \text{vol}(B_k) &= \sum_{k=1}^{\infty} m^*(B_k \cap E) + \sum_{k=1}^{\infty} m^*(B_k \setminus E) \geq m^*\left(\left(\bigcup_{k=1}^{\infty} B_k\right) \cap E\right) + m^*\left(\left(\bigcup_{k=1}^{\infty} B_k\right) \setminus E\right) \\
 &\geq m^*(A \cap E) + m^*(A \setminus E).
 \end{aligned}$$

Hence,

$$m^*(A) + \varepsilon \geq \sum_{k=1}^{\infty} \text{vol}(B_k) \geq m^*(A \cap E) + m^*(A \setminus E),$$

and since this is true for all $\varepsilon > 0$, so we're done. ■

Remark 2.4.2. A similar argument will also show that any half-space of the form

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n : x_j > 0\} \text{ or } \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_j < 0\}$$

for all $1 \leq j \leq n$ is measurable.

Lemma 2.4.2 (Properties of measurable sets).

- (a) If E is measurable, then $\mathbb{R}^n \setminus E$ is also measurable.
- (b) If E is measurable, and $x \in \mathbb{R}^n$, then $x + E$ is also measurable.

- (c) If E_1 and E_2 are measurable, then $E_1 \cap E_2$ and $E_1 \cup E_2$ are measurable.
- (d) If $E_1, \dots, E_N$ are measurable, then $\bigcup_{j=1}^N E_j$ and $\bigcap_{j=1}^N E_j$ are measurable.
- (e) Every open box and close box is measurable.
- (f) Any set E of outer measure zero is measurable.

proof of (a). Since E is measurable, $m^*(A) = m^*(A \cap E) + m^*(A \setminus E)$ for any $A \subseteq \mathbb{R}^n$. Also,

$$A \cap (\mathbb{R}^n \setminus E) = A \setminus E \text{ and } A \setminus (\mathbb{R}^n \setminus E) = A \cap E.$$

Thus,

$$m^*(A) = m^*(A \setminus (\mathbb{R}^n \setminus E)) + m^*(A \cap (\mathbb{R}^n \setminus E))$$

for all $A \subseteq \mathbb{R}^n$, which means $\mathbb{R}^n \setminus E$ is measurable. ■

proof of (b). Suppose E is measurable, then $m^*(B) = m^*(B \cap E) + m^*(B \setminus E)$ for any $B \subseteq \mathbb{R}^n$. To show $x + E$ is measurable, if we let $A \subseteq \mathbb{R}^n$ arbitrarily, we have

$$m^*(A - x) = m^*((A - x) \cap E) + m^*((A - x) \setminus E).$$

Now since $m^*(A) = m^*(A - x)$, and

$$(A - x) \cap E = (A \cap (E + x)) - x \text{ and } (A - x) \setminus E = (A \setminus (E + x)) - x,$$

we have

$$\begin{aligned} m^*(A) &= m^*(A - x) = m^*((A - x) \cap E) + m^*((A - x) \setminus E) \\ &= m^*((A \cap (E + x)) - x) + m^*((A \setminus (E + x)) - x) = m^*(A \cap (E + x)) + m^*(A \setminus (E + x)). \end{aligned}$$

Thus, $E + x$ is measurable. ■

proof of (c). It suffices to show that $E_1 \cup E_2$ is measurable. Then, by Demorgan's law,

$$E_1 \cap E_2 = \mathbb{R}^n \setminus ((\mathbb{R}^n \setminus E_1) \cup (\mathbb{R}^n \setminus E_2)),$$

so we can show $E_1 \cap E_2$ is measurable. Now we want to show

$$m^*(A) = m^*(A \cap (E_1 \cup E_2)) + m^*(A \setminus (E_1 \cup E_2)).$$

Since E_1 is measurable,

$$m^*(A) = m^*(A \cap E_1) + m^*(A \setminus E_1).$$

Next, since E_2 is measurable,

$$m^*(A \setminus E_1) = m^*((A \setminus E_1) \cap E_2) + m^*((A \setminus E_1) \setminus E_2).$$

Hence,

$$m^*(A) = m^*(A \cap E_1) + m^*(A \setminus E_1) = m^*(A \cap E_1) + m^*((A \setminus E_1) \cap E_2) + m^*((A \setminus E_1) \setminus E_2).$$

Notice that $(A \setminus E_1) \cap E_2 = A \cap (E_2 \setminus E_1)$ and $(A \setminus E_1) \setminus E_2 = A \setminus (E_1 \cup E_2)$. Thus,

$$m^*(A) = m^*(A \cap E_1) + m^*(A \cap (E_2 \setminus E_1)) + m^*(A \setminus (E_1 \cup E_2)).$$

Note that

$$(A \cap E_1) \cup (A \cap (E_2 \setminus E_1)) = A \cap (E_1 \cup E_2),$$

so

$$m^*(A \cap (E_1 \cup E_2)) \leq m^*(A \cap E_1) + m^*(A \cap (E_2 \setminus E_1)).$$

This gives

$$m^*(A) \geq m^*(A \cap (E_1 \cup E_2)) + m^*(A \setminus (E_1 \cup E_2)).$$

On the other hand, we have $A = (A \cap (E_1 \cup E_2)) \cup (A \setminus (E_1 \cup E_2))$, so we have

$$m^*(A) = m^*(A \cap (E_1 \cup E_2)) + m^*(A \setminus (E_1 \cup E_2)).$$

proof of (d). This can be proved by induction on N with (c). ■

proof of (e). Note that every open box is the intersection of many half-spaces of the form

$$\{x : x_j > a_j\} \text{ or } \{x : x_j < b_j\}.$$

Thus, by Lemma 2.4.1 and (d), we know every open box is measurable. Closed boxes are complement of union of open half spaces and thus measurable. ■

proof of (f). Suppose $m^*(E) = 0$. For any $A \subseteq \mathbb{R}^n$, we have $A \cap E \subseteq E$, so

$$m^*(A \cap E) \leq m^*(E) = 0.$$

Hence,

$$m^*(A \cap E) + m^*(A \setminus E) = m^*(A \setminus E) \leq m^*(A).$$

Since

$$m^*(A \cap E) + m^*(A \setminus E) \geq m^*(A)$$

is trivial, so we're done. ■

Lemma 2.4.3 (Finite additivity). If $(E_j)_{j \in J}$ are a finite collection of disjoint measurable sets, then for any set A (not necessarily measurable), we have

$$m^* \left(A \cap \left(\bigcup_{j \in J} E_j \right) \right) = \sum_{j \in J} m^*(A \cap E_j).$$

Furthermore, we have

$$m \left(\bigcup_{j \in J} E_j \right) = \sum_{j \in J} m(E_j).$$

Proof. It suffices to show the case of two sets: If E and F are disjoint measurable sets, then for any $A \subseteq \mathbb{R}^n$,

$$m^*(A \cap (E \cup F)) = m^*(A \cap E) + m^*(A \cap F).$$

Assume the statement is true for k pairwise disjoint measurable sets, and let $E_1, E_2, \dots, E_k, E_{k+1}$ be pairwise disjoint measurable sets. Applying the two sets case,

$$m^*(A \cap (\bigcup_{i=1}^k E_i \cup E_{k+1})) = m^*(A \cap \bigcup_{i=1}^k E_i) + m^*(A \cap E_{k+1}).$$

Now the induction hypothesis gives

$$m^*(A \cap \bigcup_{i=1}^k E_i) = \sum_{i=1}^k m^*(A \cap E_i),$$

so

$$m^*(A \cap \left(\bigcup_{i=1}^{k+1} E_i \right)) = \sum_{i=1}^{k+1} m^*(A \cap E_i).$$

Now we show the two set cases. Suppose E and F are measurable and disjoint, then $U := E \cup F$ is measurable, and since E is measurable,

$$m^*(A \cap U) = m^*((A \cap U) \cap E) + m^*((A \cap U) \setminus E).$$

Note that

$$A \cap U \cap E = A \cap (E \cup F) \cap E = A \cap E \text{ and } (A \cap U) \setminus E = (A \cap (E \cup F)) \setminus E = A \cap F$$

since $E \cap F = \emptyset$. Hence,

$$m^*(A \cap U) = m^*(A \cap E) + m^*(A \cap F),$$

and we're done.

Now if we have pairwise disjoint $E_1, E_2, \dots, E_k$, we can pick $A = \mathbb{R}^n$, and thus

$$m(\bigcup_{j \in J} E_j) = \sum_{j \in J} m(E_j).$$

■

Corollary 2.4.1. If $A \subseteq B$ and A, B are both measurable, then $B \setminus A$ is measurable and

$$m(B \setminus A) + m(A) = m(B).$$

Proof. Since

$$B \setminus A = B \cap (\mathbb{R}^n \setminus A),$$

and A, B are measurable, so we can deduce $B \setminus A$ is measurable. Also, since $A \cap (B \setminus A) = \emptyset$, so by [Lemma 2.4.3](#), we know

$$m(B) = m(A) + m(B \setminus A).$$

■

Lemma 2.4.4 (Countable additivity). Let $(E_j)_{j \in J}$ be a countable collection of pairwise disjoint measurable subsets of $\mathbb{R}^n$, then

$$E := \bigcup_{j \in J} E_j$$

is measurable, and

$$m(E) = \sum_{j \in J} m(E_j).$$

Proof. Since J is countable, we may enumerate it as $\{j_1, j_2, \dots\}$. For each $N \in \mathbb{N}$, define $F_N = \bigcup_{k=1}^N E_{j_k}$. Thus, F_N is measurable. By previous result,

$$m(F_N) = \sum_{k=1}^N m(E_{j_k}).$$

Let $E = \bigcup_{k=1}^{\infty} E_{j_k}$, we have $E = \bigcup_{N=1}^{\infty} F_N$ and $F_1 \subseteq F_2 \subseteq \dots$

We first show E is measurable. It suffices to show that

$$m^*(A \cap E) + m^*(A \setminus E) \leq m^*(A) \text{ for all } A \subseteq \mathbb{R}^n.$$

Since F_N is measurable and $F_N \subseteq E$, we have

$$m^*(A) = m^*(A \cap F_N) + m^*(A \setminus F_N)$$

and

$$A \setminus E \subseteq A \setminus F_N \Rightarrow m^*(A \setminus F_N) \geq m^*(A \setminus E).$$

Also,

$$m^*(A \cap F_N) = \sum_{k=1}^N m^*(A \cap E_{j_k}).$$

Hence,

$$m^*(A) \geq \sum_{k=1}^N m^*(A \cap E_{j_k}) + m^*(A \setminus E) \text{ true for all } N.$$

Thus,

$$m^*(A) \geq \sum_{k=1}^{\infty} m^*(A \cap E_{j_k}) + m^*(A \setminus E).$$

Since

$$\bigcup_{k=1}^{\infty} (A \cap E_{j_k}) = A \cap \left(\bigcup_{k=1}^{\infty} E_{j_k} \right) = A \cap E,$$

we have

$$m^*(A \cap E) \leq \sum_{k=1}^{\infty} m^*(A \cap E_{j_k}),$$

so

$$m^*(A) \geq m^*(A \cap E) + m^*(A \setminus E).$$

This shows E is measurable.

Now we show

$$m(E) = \sum_{k=1}^{\infty} m(E_{j_k}).$$

First, since $E = \bigcup_{i=1}^{\infty} E_{j_k}$, so

$$m(E) \leq \sum_{k=1}^{\infty} m(E_{j_k}).$$

Now since $F_N \subseteq E$ for all N , so

$$\sum_{k=1}^N m(E_{j_k}) = m(F_N) \leq m(E),$$

so

$$m(E) = \sum_{k=1}^{\infty} m(E_{j_k}).$$

■

Lemma 2.4.5 (σ -algebra property). If $(\Omega_j)_{j \in J}$ is a countable collection of measurable sets, then $\bigcup_{j \in J} \Omega_j$ and $\bigcap_{j \in J} \Omega_j$ are both measurable.

Proof. Write $J = \{j_1, j_2, \dots\}$ and define $E_1 = \Omega_{j_1}$ and

$$E_N = \Omega_N \setminus \left(\bigcup_{k=1}^{N-1} \Omega_{j_k} \right) \text{ for all } N \geq 2.$$

Hence, E_i is measurable for all $i \in \mathbb{N}$ and disjoint by construction. Note that

$$\bigcup_{k=1}^{\infty} E_k = \bigcup_{k=1}^{\infty} \Omega_{j_k}.$$

By this, $\bigcup_{k=1}^{\infty} \Omega_{j_k}$ is measurable since we have shown that the union of countably infinitely many pairwise disjoint measurable sets is measurable.

Now since

$$\bigcap_{k=1}^{\infty} \Omega_{j_k} = \mathbb{R}^n \setminus \bigcup_{k=1}^{\infty} (\mathbb{R}^n \setminus \Omega_{j_k}),$$

we know $\bigcap_{k=1}^{\infty} \Omega_{j_k}$ is measurable.

Lemma 2.4.6. Every open set can be written as a countable or finite union of open boxes. ■

Proof. We say an open box is rational if $B = \prod_{i=1}^n (a_i, b_i)$ where $a_i, b_i \in \mathbb{Q}$. The set of rational open boxes is countable (since this set is isomorphic to a subset of $\mathbb{Q}^{2n}$).

We make the following claim: given any open ball $B(x, r)$, $\exists$ a rational box B with $x \in B \subseteq B(x, r)$. To show this, let $x = (x_1, \dots, x_n)$ be any point in $\mathbb{R}^n$. Let a_i, b_i be rational numbers s.t.

$$x_i - \frac{r}{n} < a_i < x_i < b_i < x_i + \frac{r}{n},$$

then we show that $B \subseteq B(x, r)$. Let $y \in B$, then

$$x_i - \frac{r}{n} < a_i < y_i < b_i < x_i + \frac{r}{n},$$

so

$$|y_i - x_i| < \frac{r}{n}.$$

Hence,

$$\|y - x\| = \sqrt{\sum_{i=1}^n (y_i - x_i)^2} \leq \sum_{i=1}^n |y_i - x_i| < \frac{r}{n} \cdot n = r.$$

Hence, $y \in B(x, r)$. Thus, we're done.

Now let E be an open set in $\mathbb{R}^n$. For any $x \in E$, $\exists r_x > 0$ s.t. $B(x, r_x) \subseteq E$. Thus, $E = \bigcup_{x \in E} B(x, r_x)$. Also, for any $B(x, r_x)$, there exists a rational box B_x s.t. $x \in B_x \subseteq B(x, r_x)$. Thus,

$$\bigcup_{x \in E} B_x = E. \quad \blacksquare$$

Lemma 2.4.7. Every open set, and every closed set, is Lebesgue measurable.

Proof. By previous lemma, every open set can be written as the union of countable or finite rational open boxes, and every rational open box is measurable, so every open set is measurable. Since every closed box is $\mathbb{R}^n \setminus E$ for some open E , so it is also measurable. ■

Lecture 12

2.5 Measurable Functions

7 April.

Definition 2.5.1 (Measurable functions). Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f : \Omega \rightarrow \mathbb{R}^m$ be a function. Then, f is measurable iff $f^{-1}(V)$ is measurable for every open set $V \subseteq \mathbb{R}^m$, where

$$f^{-1}(V) = \{x \in \Omega \mid f(x) \in V\}$$

is the pre-image(or inverse image) of V under f .

Remark 2.5.1. In measure theory, this type of function is called Borel measurable.

Definition 2.5.2. Given a set X . A collection of subsets $\sigma(X)$ of X is called a σ -algebra if

- (1) $X \in \sigma(X)$.

- (2) $E \in \sigma(X)$ implies $X \setminus E \in \sigma(X)$.
- (3) $E_i \in \sigma(X)$ for all $i \in \mathbb{N}$ implies $\bigcup_{i=1}^{\infty} E_i \in \sigma(X)$.

Remark 2.5.2. Since

$$\bigcap_{i=1}^{\infty} E_i = X \setminus \left(\bigcup_{i=1}^{\infty} (X \setminus E_i) \right),$$

so $E_i \in \sigma(X)$ for all $i \in \mathbb{N}$ implies $\bigcap_{i=1}^{\infty} E_i \in \sigma(X)$.

Definition 2.5.3. In $\mathbb{R}^n$, a set is called Borel measurable if it is in the smallest σ -algebra that contains all open sets.

Lemma 2.5.1. Let Ω be a measurable subset of $\mathbb{R}^n$, and $f : \Omega \rightarrow \mathbb{R}^m$ be continuous, then f is also measurable.

Proof. Let V be any open subset of $\mathbb{R}^m$, then since f is continuous, so $f^{-1}(V)$ is open, and since all open sets are measurable, we know $f^{-1}(V)$ is measurable, and thus f is measurable. ■

Lemma 2.5.2. Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f : \Omega \rightarrow \mathbb{R}^m$ be a function. Then f is measurable if and only if $f^{-1}(B)$ is measurable for every open box B .

Proof.

($\Rightarrow$) If f is measurable, then since every open box B is open, $f^{-1}(B)$ is measurable.

($\Leftarrow$) Given any open set V , $\exists$ a countable union of open boxes s.t. $V = \bigcup_{\ell=1}^{\infty} B_{\ell}$ by [Lemma 2.4.6](#), and

$$f^{-1}(V) = f^{-1}\left(\bigcup_{\ell=1}^{\infty} B_{\ell}\right) = \bigcup_{\ell=1}^{\infty} f^{-1}(B_{\ell}),$$

and since $f^{-1}(B_{\ell})$ is measurable for all $\ell \in \mathbb{N}$, and thus $\bigcup_{\ell=1}^{\infty} f^{-1}(B_{\ell})$ is measurable by [Lemma 2.4.5](#). ■

Corollary 2.5.1. Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f : \Omega \rightarrow \mathbb{R}^m$ be a function. Suppose $f = (f_1, f_2, \dots, f_m)$, where $f_j : \Omega \rightarrow \mathbb{R}$ is the j -th coordinate of f . Then f is measurable if and only if f_j is measurable for all $1 \leq j \leq m$.

Proof.

($\Rightarrow$) Fix $j \in \{1, 2, \dots, m\}$. Let $I \subseteq \mathbb{R}$ be an open interval. Consider the open set $B_j = \mathbb{R} \times \dots \times I \times \dots \times \mathbb{R}$, where I is in the j -th coordinate, then for $x \in \Omega$, we know

$$x \in f^{-1}(B_j) \Leftrightarrow f(x) \in B_j \Leftrightarrow f_j(x) \in I \Leftrightarrow x \in f_j^{-1}(I).$$

Hence, $f^{-1}(B_j) = f_j^{-1}(I)$. Since f is measurable, $f^{-1}(B_j)$ is measurable since B_j is an open box. Hence, $f_j^{-1}(I)$ is measurable, and since this is true for all open box I of $\mathbb{R}$, so by [Lemma 2.5.2](#), f_j is measurable for all $1 \leq j \leq m$.

($\Leftarrow$) Let $B = I_1 \times I_2 \times \cdots \times I_m$ be an open box in $\mathbb{R}^m$. Then,

$$\begin{aligned} x \in f^{-1}(B) &\Leftrightarrow f(x) = (f_1(x), f_2(x), \dots, f_m(x)) \in B = I_1 \times I_2 \times \cdots \times I_m \\ &\Leftrightarrow f_j(x) \in I_j \text{ for all } 1 \leq j \leq m \Leftrightarrow x \in f_j^{-1}(I_j) \text{ for all } 1 \leq j \leq m \\ &\Leftrightarrow x \in \bigcap_{j=1}^m f_j^{-1}(I_j). \end{aligned}$$

Hence, $f^{-1}(B) = \bigcap_{j=1}^m f_j^{-1}(I_j)$. Since for all $1 \leq j \leq m$, f_j is measurable and I_j is an open box in $\mathbb{R}$, so $f_j^{-1}(I_j)$ is measurable and thus $\bigcap_{j=1}^m f_j^{-1}(I_j)$ is measurable, so $f^{-1}(B)$ is measurable, and thus f is measurable. ■

Lemma 2.5.3. Let Ω be a measurable subset of $\mathbb{R}^n$, and let W be an open subset of $\mathbb{R}^m$. If $f : \Omega \rightarrow W$ is measurable and $g : W \rightarrow \mathbb{R}^p$ is continuous, then $g \circ f : \Omega \rightarrow \mathbb{R}^p$ is measurable.

Proof. Let V be an open set in $\mathbb{R}^p$, then $(g \circ f)^{-1}(V) = f^{-1}(g^{-1}(V))$. Since g is continuous and V is open, $g^{-1}(V)$ is open in W , so $g^{-1}(V) = U \cap W$, where U is open in $\mathbb{R}^m$. Since W is also open in $\mathbb{R}^m$, $W \cap U$ is also open in $\mathbb{R}^m$, so $g^{-1}(V)$ is open in $\mathbb{R}^m$. Now since f is measurable, $f^{-1}(g^{-1}(V))$ is measurable, i.e. $(g \circ f)^{-1}(V)$ is measurable. ■

Corollary 2.5.2. Let Ω be a measurable subset of $\mathbb{R}^n$. If $f : \Omega \rightarrow \mathbb{R}$ is measurable, then so is $|f|$, $\max\{f, 0\}$, and $\min\{f, 0\}$.

Proof. Since $g_1(x) := |x|$, $g_2(x) := \max\{x, 0\}$, and $g_3(x) := \min\{x, 0\}$ are all continuous, so by Lemma 2.5.3, $g_1 \circ f$, $g_2 \circ f$, $g_3 \circ f$ are all measurable.

Remark 2.5.3.

$$\max\{x, 0\} = \frac{x + |x|}{2}, \quad \min\{x, 0\} = \frac{x - |x|}{2}.$$
■

Corollary 2.5.3. Let Ω be a measurable subset of $\mathbb{R}^n$. If $f : \Omega \rightarrow \mathbb{R}$ and $g : \Omega \rightarrow \mathbb{R}$ are measurable, then so are $f + g$, $f - g$, $f \cdot g$, $\max\{f, g\}$, and $\min\{f, g\}$. If $g(x) \neq 0$ for all $x \in \Omega$, then f/g is also measurable.

Proof. Consider the map $F : \Omega \rightarrow \mathbb{R}^2$ defined by $F(x) = (f(x), g(x))$, then by Corollary 2.5.1, F is also measurable. Define $h_+(u, v) = u + v$, where $h_+ : \mathbb{R}^2 \rightarrow \mathbb{R}$ is continuous. Then $h_+ \circ F$ is measurable, where $(h_+ \circ F)(x) = f(x) + g(x)$. Similarly, we can define

$$\begin{aligned} h_*(u, v) &= uv, \\ h_{\max}(u, v) &= \max(u, v) = \frac{u + v + |u - v|}{2}, \\ h_{\min}(u, v) &= \min(u, v) = \frac{u + v - |u - v|}{2}, \end{aligned}$$

and they are all continuous, so $h_* \circ F$, $h_{\max} \circ F$, $h_{\min} \circ F$ are all measurable. As for the division, it is similar. ■

Lemma 2.5.4. Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f : \Omega \rightarrow \mathbb{R}$ be a function, then f is measurable iff $f^{-1}((a, \infty))$ is measurable for any $a \in \mathbb{R}$.

Proof.

($\Rightarrow$) Since (a, ∞) is open, so $f^{-1}((a, \infty))$ is measurable.

($\Leftarrow$) Let I be an open interval in $\mathbb{R}$ with $I = (\alpha, \beta)$ where $\alpha < \beta$, then it suffices to show $f^{-1}(I)$ is measurable by [Lemma 2.5.2](#). Note that $(\alpha, \beta) = (\alpha, \infty) \cap (\mathbb{R} \setminus [\beta, \infty))$, and

$$[\beta, \infty) = \bigcap_{m=1}^{\infty} \left(\beta - \frac{1}{m}, \infty \right).$$

Therefore,

$$f^{-1}([\beta, \infty)) = f^{-1} \left(\bigcap_{m=1}^{\infty} \left(\beta - \frac{1}{m}, \infty \right) \right) = \bigcap_{m=1}^{\infty} f^{-1} \left(\left(\beta - \frac{1}{m}, \infty \right) \right),$$

which is measurable since $f^{-1}(a, \infty)$ is measurable for all $a \in \mathbb{R}$. Hence,

$$f^{-1}(\alpha, \beta) = f^{-1}(\alpha, \infty) \cap (\Omega \setminus f^{-1}(\beta, \infty))$$

is measurable. ■

Definition 2.5.4 (Measurable functions in the extended reals). Let Ω be a measurable subset of $\mathbb{R}^n$. Set

$$\mathbb{R}^* := \mathbb{R} \cup \{+\infty\} \cup \{-\infty\}.$$

A function $f : \Omega \rightarrow \mathbb{R}^*$ is said to be measurable if $f^{-1}((a, \infty])$ is measurable for all $a \in \mathbb{R}$, where

$$f^{-1}((a, \infty]) = f^{-1}((a, \infty)) \cup f^{-1}(\{\infty\}).$$

Lecture 13

Example 2.5.1. Suppose $f(x) = \frac{1}{x^2}$ is defined on $\mathbb{R}$, then we can extend it to $f : \mathbb{R} \rightarrow \mathbb{R}^*$ so that

$$f(x) = \begin{cases} \frac{1}{x^2}, & \text{if } x \neq 0; \\ \infty, & \text{if } x = 0. \end{cases}$$

Then f is measurable. Note that if we define f by $f : \Omega \rightarrow \mathbb{R}$, then $f^{-1}((a, \infty]) = f^{-1}((a, \infty))$ for all $a \in \mathbb{R}$.

Lemma 2.5.5. Let $\Omega \subseteq \mathbb{R}^n$ be measurable. For each positive integer n , let $f_n : \Omega \rightarrow \mathbb{R}^*$ be a measurable function. Then, $\sup_{n \in \mathbb{N}} f_n, \inf_{n \in \mathbb{N}} f_n, \limsup_{n \rightarrow \infty} f_n, \liminf_{n \rightarrow \infty} f_n$ are measurable. In particular, if $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ for $f : \Omega \rightarrow \mathbb{R}^*$, then f is also measurable.

Proof. Let $g(x) = \sup_{n \in \mathbb{N}} f_n(x)$, then g is measurable iff $g^{-1}((a, \infty])$ is measurable for any $a \in \mathbb{R}$. Also,

$$x \in g^{-1}((a, \infty]) \Leftrightarrow g(x) > a \Leftrightarrow \sup_{n \in \mathbb{N}} f_n(x) > a \Leftrightarrow \exists k \in \mathbb{N} \text{ s.t. } f_k(x) > a$$

$$\Leftrightarrow \exists k \in \mathbb{N} \text{ s.t. } x \in f_k^{-1}((a, \infty]) \Leftrightarrow x \in \bigcup_{n=1}^{\infty} f_n^{-1}((a, \infty]).$$

Hence, $g^{-1}((a, \infty]) = \bigcup_{n=1}^{\infty} f_n^{-1}((a, \infty])$, so it suffices to show that $\bigcup_{n=1}^{\infty} f_n^{-1}((a, \infty])$ is measurable. However, since for all $n \in \mathbb{N}$, f_n is measurable, so $f_n^{-1}((a, \infty])$ is measurable, and thus we're done.

Let $h(x) = \inf_{n \in \mathbb{N}} f_n(x)$, then h is measurable iff $h^{-1}((a, \infty])$ is measurable for all $a \in \mathbb{R}$. Now

since

$$\begin{aligned} x \in h^{-1}((a, \infty]) &\Leftrightarrow h(x) \in (a, \infty] \Leftrightarrow \inf_{n \in \mathbb{N}} f_n(x) \in (a, \infty] \Leftrightarrow \inf_{n \in \mathbb{N}} f_n(x) > a \Leftrightarrow \forall n \in \mathbb{N}, f_n(x) > a \\ &\Leftrightarrow \forall n \in \mathbb{N}, f_n(x) \in (a, \infty] \Leftrightarrow \forall n \in \mathbb{N}, x \in f_n^{-1}((a, \infty]) \Leftrightarrow x \in \bigcap_{n=1}^{\infty} f_n^{-1}((a, \infty]). \end{aligned}$$

Hence, $h^{-1}((a, \infty]) = \bigcap_{n=1}^{\infty} f_n^{-1}((a, \infty])$. However, since for all $n \in \mathbb{N}$, f_n is measurable, so $f_n^{-1}((a, \infty])$ is measurable, and thus we're done.

Now we define $g_N(x) = \sup_{n \geq N} f_n(x)$, then we know g_N is measurable for all $N \in \mathbb{N}$ by what we've proved. Now since

$$\limsup_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \sup_{k \geq n} f_k(x) = \inf_{N \geq 1} \sup_{n \geq N} f_n(x) = \inf_{N \geq 1} g_N(x),$$

so by what we've proved, we know $\inf_{N \geq 1} g_N(x)$ is measurable, and we're done.

Similar argument can be used to prove $\liminf_{n \rightarrow \infty} f_n(x)$ is measurable.

Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on a set Ω , and suppose that

$$f_n(x) \rightarrow f(x) \quad \text{for every } x \in \Omega.$$

We will show that for every $x \in \Omega$,

$$\limsup_{n \rightarrow \infty} f_n(x) = f(x) = \liminf_{n \rightarrow \infty} f_n(x).$$

Fix $x \in \Omega$. Then $\{f_n(x)\}$ is a sequence of real numbers converging to $f(x)$.

Recall the definitions:

$$\limsup_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \left(\sup_{k \geq n} f_k(x) \right), \quad \liminf_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \left(\inf_{k \geq n} f_k(x) \right).$$

We first show that

$$\limsup_{n \rightarrow \infty} f_n(x) \leq f(x).$$

Let $\varepsilon > 0$. Since $f_n(x) \rightarrow f(x)$, there exists $N \in \mathbb{N}$ such that for all $n \geq N$,

$$|f_n(x) - f(x)| < \varepsilon,$$

which implies

$$f_n(x) \leq f(x) + \varepsilon.$$

Hence, for all $n \geq N$,

$$\sup_{k \geq n} f_k(x) \leq f(x) + \varepsilon.$$

Taking the limit as $n \rightarrow \infty$, we obtain

$$\limsup_{n \rightarrow \infty} f_n(x) \leq f(x) + \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary,

$$\limsup_{n \rightarrow \infty} f_n(x) \leq f(x).$$

Next, we show that

$$\liminf_{n \rightarrow \infty} f_n(x) \geq f(x).$$

Again let $\varepsilon > 0$. Since $f_n(x) \rightarrow f(x)$, there exists $N \in \mathbb{N}$ such that for all $n \geq N$,

$$|f_n(x) - f(x)| < \varepsilon,$$

which implies

$$f_n(x) \geq f(x) - \varepsilon.$$

Hence, for all $n \geq N$,

$$\inf_{k \geq n} f_k(x) \geq f(x) - \varepsilon.$$

Taking the limit as $n \rightarrow \infty$, we obtain

$$\liminf_{n \rightarrow \infty} f_n(x) \geq f(x) - \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary,

$$\liminf_{n \rightarrow \infty} f_n(x) \geq f(x).$$

Combining the two inequalities, we have

$$\limsup_{n \rightarrow \infty} f_n(x) \leq f(x) \leq \liminf_{n \rightarrow \infty} f_n(x).$$

But in general,

$$\liminf_{n \rightarrow \infty} f_n(x) \leq \limsup_{n \rightarrow \infty} f_n(x),$$

so all three quantities are equal:

$$\limsup_{n \rightarrow \infty} f_n(x) = f(x) = \liminf_{n \rightarrow \infty} f_n(x).$$

Since $x \in \Omega$ was arbitrary, the result holds for all x , hence

$$f = \limsup_{n \rightarrow \infty} f_n = \liminf_{n \rightarrow \infty} f_n.$$

This shows that if $f_n \rightarrow f$ pointwise, then f is measurable. ■

Chapter 3

Lebesgue Integration

What is Lebesgue integration?

Lebesgue integration is a way of defining the integral of a function by grouping together points that have *similar output values*. Instead of cutting the x -axis into many thin intervals, as in Riemann integration, Lebesgue integration studies the sets of points where the function lies in certain value ranges and measures the size of those sets.

Informally, the two viewpoints are:

- **Riemann integration:** slice the domain into vertical strips and add “width $\times$ height.”
- **Lebesgue integration:** slice the range into horizontal bands and add “value $\times$ measure of the set having that value.”

This change of perspective makes Lebesgue integration much more flexible. It can integrate many functions that are badly behaved from the Riemann point of view, and it works especially well in probability, Fourier analysis, and modern analysis.

The main idea

Suppose a function f takes the value approximately y on a set of points whose length is approximately m . Then its contribution to the integral is approximately



$$y \cdot m.$$

Adding these contributions over all output levels leads to the Lebesgue integral.

For nonnegative functions, one can think of the construction as approximating f by step functions built from *horizontal* layers. In this way, the Lebesgue integral measures area by organizing the graph according to function values rather than according to input intervals.

How is it different from Riemann integration?

The essential difference is the direction of the partition.



- In **Riemann integration**, the interval on the x -axis is divided into many small subintervals. One samples the function in each interval and sums rectangle areas.
- In **Lebesgue integration**, the y -axis is divided into small bands. For each band, one looks at all x such that $f(x)$ falls inside that band, then measures how large that set is.

So Riemann organizes the area by *where the input lies*, while Lebesgue organizes it by *what values the function takes*.

Why Lebesgue integration is more powerful

Lebesgue integration handles limits of functions much better. For example, if a sequence of functions converges in a reasonable way, the Lebesgue integral often allows us to interchange limit and integral. This is one of the main reasons it became the standard notion of integration in advanced mathematics.

It also integrates some functions that are not Riemann integrable. The classical example is the Dirichlet function on $[0, 1]$:

$$f(x) = \begin{cases} 1, & x \in \mathbb{Q}, \\ 0, & x \notin \mathbb{Q}. \end{cases}$$


This function is discontinuous at every point, so it is not Riemann integrable. However, the rational numbers in $[0, 1]$ have Lebesgue measure 0, so from the Lebesgue point of view the function is equal to 0 almost everywhere. Therefore,

$$\int_0^1 f(x) dx = 0$$

in the Lebesgue sense.

A visual example

The next picture shows the Dirichlet function schematically. The dense blue points near $y = 1$ represent rational inputs, and the dense red points near $y = 0$ represent irrational inputs. Since the rational set has measure zero, the upper layer contributes no area in the Lebesgue sense.



Summary

- **Riemann integration** partitions the domain and adds areas of vertical rectangles.
- **Lebesgue integration** partitions the range and measures the size of the sets where the function takes certain values.
- Lebesgue integration is better suited to complicated functions and to taking limits.
- Every Riemann integrable function is Lebesgue integrable, but not every Lebesgue integrable function is Riemann integrable.

In short, Lebesgue integration is a more robust and more powerful notion of area, built to handle modern analysis.

3.1 Simple Functions

Definition 3.1.1. Let $\Omega \subseteq \mathbb{R}^n$ be a measurable set, and let $f : \Omega \rightarrow \mathbb{R}$ be a measurable function. We say f is a simple function if the image of $f(\Omega)$ is finite, i.e. $\exists c_1, \dots, c_N$ s.t. for any $x \in \Omega$, $f(x) = c_j$ for some $1 \leq j \leq N$.

Example 3.1.1. Let $\Omega \subseteq \mathbb{R}^n$ be a measurable set, and let E be a measurable subset of Ω . Define the characteristic function $\chi_E : \Omega \rightarrow \mathbb{R}$ by

$$\chi_E(x) = \begin{cases} 1, & \text{if } x \in E; \\ 0, & \text{if } x \in \Omega \setminus E. \end{cases}$$

Then, we claim that χ_E is measurable. Note that for any open interval I , we have

$$\chi_E^{-1}(I) = \begin{cases} \emptyset, & \text{if } 0 \notin I, 1 \notin I; \\ E, & \text{if } 0 \notin I, 1 \in I; \\ \Omega \setminus E, & \text{if } 0 \in I, 1 \notin I; \\ \Omega, & \text{if } 0 \in I, 1 \in I. \end{cases}$$

Thus, χ_E is measurable.

Lemma 3.1.1. Let $\Omega \subseteq \mathbb{R}^n$ be measurable, and let $f : \Omega \rightarrow \mathbb{R}$ and $g : \Omega \rightarrow \mathbb{R}$ be simple function. Then, $f + g$ is also simple and for any $c \in \mathbb{R}$, the function cf is also simple.

Proof. Suppose $f(\Omega) = \{a_1, \dots, a_N\}$ and $g(\Omega) = \{b_1, \dots, b_M\}$, then $(f + g)(\Omega) = \{a_i + b_j\}_{\substack{1 \leq i \leq N, \\ 1 \leq j \leq M}}$, which is finite, so $f + g$ is simple.

Also, $(cf)(\Omega) = \{ca_1, \dots, ca_N\}$, which is also finite, so cf is simple. ■

Lemma 3.1.2. Let $\Omega \subseteq \mathbb{R}^n$ be measurable, and let $f : \Omega \rightarrow \mathbb{R}$ be a simple function. Then, there exists finitely many distinct real numbers $c_1, \dots, c_N$ and finitely many pairwise disjoint measurable sets $E_1, \dots, E_N \subseteq \Omega$ s.t. $f = \sum_{j=1}^N c_j \chi_{E_j}$.

Proof. Suppose $f(\Omega) = \{c_1, \dots, c_N\}$, where $c_i \neq c_j$ for all $i \neq j$. Define $E_j = f^{-1}(\{c_j\})$ for each $1 \leq j \leq N$, then $(E_j)_{j=1}^N$ are pairwise disjoint. Since $f(x) = c_\ell$ for some $1 \leq \ell \leq N$ for all $x \in \Omega$, $\Omega = \bigcup_{j=1}^N E_j$. Note that

$$E_j = f^{-1}(\{c_j\}) = \bigcap_{k=1}^{\infty} f^{-1}\left(\left(c_j - \frac{1}{k}, c_j + \frac{1}{k}\right)\right),$$

then since f is measurable, we know E_j is measurable. Now for $x \in \Omega$, we know $x \in E_k$ for some $1 \leq k \leq N$. Hence, $f(x) = c_k$, and thus

$$\sum_{j=1}^N c_j \chi_{E_j}(x) = c_k \chi_{E_k}(x) = c_k = f(x).$$

Thus, $f = \sum_{j=1}^N c_j \chi_{E_j}$. ■

Remark 3.1.1. This shows f is simple implies f is measurable since we can show $f^{-1}((a, \infty))$ is measurable for all $a \in \mathbb{R}$ with knowing that in fact $f^{-1}((a, \infty))$ is some intersection and union of E_i 's.

Lemma 3.1.3. Let $\Omega \subseteq \mathbb{R}^n$ be measurable, and let $f : \Omega \rightarrow [0, \infty]$ be a measurable function. Then there exists a sequence of simple functions $\{f_n\}_{n=1}^\infty$ with $f_n : \Omega \rightarrow \mathbb{R}$ s.t. each $f_n \geq 0$, and the sequence is increasing:

$$0 \leq f_1(x) \leq f_2(x) \leq \dots \quad \text{for all } x \in \Omega,$$

and $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ for all $x \in \Omega$.

Proof. For each integer $n \geq 1$, define

$$E_{n,k} := \left\{ x \in \Omega : \frac{k}{2^n} \leq f(x) < \frac{k+1}{2^n} \right\}, \quad k = 0, 1, \dots, n2^n - 1,$$

and $F_n := \{x \in \Omega : f(x) \geq n\}$. Then define

$$f_n(x) := \sum_{k=0}^{n2^n-1} \frac{k}{2^n} \chi_{E_{n,k}}(x) + n \chi_{F_n}(x).$$

Thus, f_n is obtained from f by truncating at height n and then rounding down to the nearest multiple of 2^{-n} . Note that if $f(x) < n$, then $\exists 0 \leq k \leq n2^n - 1$ s.t.

$$\frac{k}{2^n} \leq f(x) < \frac{k+1}{2^n},$$

so $f_n(x) = \frac{k}{2^n}$, and if $f(x) \geq n$, then $f_n(x) = n$.

Now we try to write down the explicit formula for f_n . Fix $x \in \Omega$, then write $a := f(x) \in [0, \infty]$.

Claim 3.1.1.

$$f_n(x) = \min \left(n, \frac{\lfloor 2^n a \rfloor}{2^n} \right) = \min \left(n, \frac{\lfloor 2^n f(x) \rfloor}{2^n} \right).$$

Proof. If $a = f(x) \geq n$, then $f_n(x) = n$, and $n \leq \frac{\lfloor 2^n f(x) \rfloor}{2^n}$, so this is true. If $a = f(x) < n$, then $\exists 0 \leq k \leq n2^n - 1$ s.t.

$$\frac{k}{2^n} \leq a < \frac{k+1}{2^n} \Leftrightarrow k \leq 2^n a < k+1,$$

so $k = \lfloor 2^n a \rfloor$. Since $f(x) \in E_{n,k}$, we have

$$f_n(x) = \frac{k}{2^n} = \frac{\lfloor 2^n a \rfloor}{2^n}.$$

Thus, in this case, $f_n(x) = \min \left(n, \frac{\lfloor 2^n a \rfloor}{2^n} \right)$.

⊛

By this claim, we know $0 \leq f_n(x) \leq f(x)$ for all n .

Now we show $f_n(x) \leq f_{n+1}(x)$.

- Case 1: $n \leq f(x)$, then $f_n(x) = n$, and $f_{n+1}(x) = \min \left(n+1, \frac{\lfloor 2^{n+1} f(x) \rfloor}{2^{n+1}} \right) \geq n$, so we're done.
- Case 2: $f(x) < n$, then

$$f_n(x) = \min \left(n, \frac{\lfloor 2^n f(x) \rfloor}{2^n} \right) = \frac{\lfloor 2^n f(x) \rfloor}{2^n}.$$

Thus, $\exists$ unique $m = \lfloor 2^n f(x) \rfloor \leq 2^n f(x) < m+1$, and

$$f_{n+1}(x) = \min \left(n+1, \frac{\lfloor 2^{n+1} f(x) \rfloor}{2^{n+1}} \right),$$

and note that

$$2m \leq 2^{n+1}f(x) < 2(m+1),$$

so

$$f_n(x) = \frac{\lfloor 2^n f(x) \rfloor}{2^n} = \frac{m}{2^n} = \frac{2m}{2^{n+1}} \leq \frac{\lfloor 2^{n+1}f(x) \rfloor}{2^{n+1}} = f_{n+1}(x),$$

where the last equality holds since $f(x) < n < n+1$.

Now we prove $\lim_{n \rightarrow \infty} f_n(x) = f(x)$.

- Case 1: $f(x) = \infty$, then $f_n(x) = n$, so $\lim_{n \rightarrow \infty} f_n(x) = x$.
- Case 2: $f(x) < \infty$, then $\exists N \in \mathbb{N}$ s.t. $f(x) < N$. If $n \geq N$, then

$$f_n(x) = \frac{k}{2^n} \text{ where } \frac{k}{2^n} \leq f(x) < \frac{k+1}{2^n}.$$

Thus, $f_n(x) \leq f(x) \leq f_n(x) + \frac{1}{2^n}$, i.e.

$$|f(x) - f_n(x)| < \frac{1}{2^n}.$$

Thus, $\lim_{n \rightarrow \infty} f_n(x) = f(x)$. ■

Lecture 14

Definition 3.1.2. Let $\Omega \subseteq \mathbb{R}^n$ be measurable, and let $f : \Omega \rightarrow [0, \infty)$ be a simple function which is non-negative, thus f is measurable and the image $f(\Omega)$ is finite and contained in $[0, \infty)$. Thus, we can define the Lebesgue integral $\int_{\Omega} f$ of f on Ω by

$$\int_{\Omega} f = \sum_{\lambda \in f(\Omega), \lambda > 0} \lambda m(\{x \in \Omega : f(x) = \lambda\}) = \sum_{\lambda \in f(\Omega)} \lambda m(\{x \in \Omega : f(x) = \lambda\}).$$

We'll also sometimes write $\int_{\Omega} f$ as $\int_{\Omega} f \, dm$.

Remark 3.1.2. By definition, we know simple function can be written as

$$f = \sum_{j=1}^N c_j \chi_{E_j},$$

where $\{c_1, \dots, c_N\}$ are distinct and $\{E_j\}_{j=1}^N$ are disjoint, measurable subsets of Ω defined by $E_j = f^{-1}(\{c_j\})$, and thus

$$\int_{\Omega} f = \sum_{j=1}^N c_j m(E_j).$$

Example 3.1.2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be the function which is equal to 3 on $[1, 2]$ and equal to 4 on $(2, 4)$ and is zero elsewhere. Then,

$$\int_{\mathbb{R}} f = 3m([1, 2]) + 4m([2, 4]) = 11.$$

Now we show the simple integral of a simple function can equal $+\infty$. (The reason why we restrict this integral to non-negative functions is to avoid ever encountering the indefinite form $+\infty + (-\infty)$.)

21 April.

Example 3.1.3. Suppose $g : \mathbb{R} \rightarrow \mathbb{R}$ is the function which is 1 on $[0, \infty)$ and is zero on $(-\infty, 0)$, then $g = 1 \cdot \chi_{[0, \infty)}$ and

$$\int_{\mathbb{R}} g = 1 \cdot m([0, \infty)) = \infty.$$

Lemma 3.1.4. Let $\Omega \subseteq \mathbb{R}^n$ be measurable and let $E_1, E_2, \dots, E_N$ be pairwise disjoint measurable subset of Ω . Let $c_1, \dots, c_N$ be non-negative real numbers (not required to be distinct). Then,

$$\int_{\Omega} \sum_{j=1}^N c_j \chi_{E_j} = \sum_{j=1}^N c_j m(E_j).$$

Proof. Define $f := \sum_{j=1}^N c_j \chi_{E_j}$, then f is simple by definition. Let $d_1, \dots, d_L$ be the distinct positive numbers that occurs among $c_1, \dots, c_N$. Let

$$F_{\ell} := \bigcup_{\{j: c_j = d_{\ell}\}} E_j.$$

Then since each E_j is measurable, so F_{ℓ} is measurable. Note that $F_1, F_2, \dots, F_L$ are pairwise disjoint since $E_1, \dots, E_N$ are pairwise disjoint. Also, note that

$$f = \sum_{\ell=1}^L d_{\ell} \chi_{F_{\ell}}.$$

Now since $d_1, \dots, d_L$ are distinct and $F_1, F_2, \dots, F_L$ are pairwise disjoint, by definition we have

$$\int_{\Omega} f = \sum_{\ell=1}^L d_{\ell} m(F_{\ell}) = \sum_{\ell=1}^L d_{\ell} \sum_{j: c_j = d_{\ell}} m(E_j) = \sum_{\ell=1}^L \sum_{j: c_j = d_{\ell}} c_j m(E_j) = \sum_{j=1}^N c_j m(E_j).$$

■

Proposition 3.1.1. Let $\Omega \subseteq \mathbb{R}^n$ be measurable, and let $f : \Omega \rightarrow [0, \infty)$ and $g : \Omega \rightarrow [0, \infty)$ be non-negative simple functions.

- (a) We have $0 \leq \int_{\Omega} f \leq \infty$. Furthermore, we have $\int_{\Omega} f = 0$ iff $m(\{x \in \Omega : f(x) \neq 0\}) = 0$.
- (b) $\int_{\Omega} f + g = \int_{\Omega} f + \int_{\Omega} g$.
- (c) For any $c > 0$, we have $\int_{\Omega} cf = c \int_{\Omega} f$.
- (d) If $f(x) \leq g(x)$ for all $x \in \Omega$, then $\int_{\Omega} f \leq \int_{\Omega} g$.

Remark 3.1.3. We make a very convenient notational convention: if a property $P(x)$ holds for all points in Ω , except for a set of measure zero, then we say that P holds for almost every point in Ω . Thus (a) asserts that $\int_{\Omega} f = 0$ if and only if f is zero for almost every point in Ω .

Proof. Since

$$f = \sum_{\lambda \in f(\Omega) \setminus \{0\}} \lambda \cdot \chi_{\{x \in \Omega : f(x) = \lambda\}},$$

we can write $f = \sum_{j=1}^N c_j \chi_{E_j}$, where $E_1, \dots, E_N$ are pairwise disjoint, and each of c_j is positive. Similarly, we write $g = \sum_{j=1}^M d_j \chi_{F_j}$, where $F_1, \dots, F_M$ are pairwise disjoint and $d_1, \dots, d_M$ are positive.

- (a) Since $\int_{\Omega} f = \sum_{j=1}^N c_j m(E_j)$, so it is clear that $0 \leq \int_{\Omega} f \leq \infty$. If f is zero almost everywhere, then all of E_j must have measure 0 (because $E_j \subseteq \{x \in \Omega : f(x) \neq 0\}$), and thus $\int_{\Omega} f = 0$.

Conversely, if $\int_{\Omega} f = 0$, then $\sum_{j=1}^N c_j m(E_j) = 0$, which can only happen when all of $m(E_j)$ are 0, and hence f is zero almost everywhere.

(b) Set

$$E_0 = \Omega \setminus \bigcup_{j=1}^N E_j, \quad c_0 = 0,$$

then $E_0, E_1, \dots, E_N$ are pairwise disjoint, and their union is Ω , and $f = \sum_{j=0}^N c_j \chi_{E_j}$. Similarly, we can define

$$F_0 = \Omega \setminus \bigcup_{k=1}^M F_k, \quad d_0 = 0,$$

and $g = \sum_{k=0}^M d_k \chi_{F_k}$, where $F_0, F_1, \dots, F_M$ are pairwise disjoint and their union is Ω . Now we know

$$\{E_j \cap F_k : 0 \leq j \leq N, 0 \leq k \leq M\}$$

forms a partition of Ω . Now since

$$\chi_{E_j} = \sum_{k=0}^M \chi_{E_j \cap F_k}, \quad \chi_{F_k} = \sum_{j=0}^N \chi_{E_j \cap F_k},$$

we have

$$f = \sum_{j=0}^N \sum_{k=0}^M c_j \chi_{E_j \cap F_k}, \quad g = \sum_{j=0}^N \sum_{k=0}^M d_k \chi_{E_j \cap F_k}.$$

(Note that we need to define E_0, F_0 to write this thing.) Hence,

$$f + g = \sum_{j=0}^N \sum_{k=0}^M (c_j + d_k) \chi_{E_j \cap F_k},$$

which shows

$$\int_{\Omega} (f + g) = \sum_{j=0}^N \sum_{k=0}^M (c_j + d_k) m(E_j \cap F_k) = \int_{\Omega} f + \int_{\Omega} g$$

by previous lemma ($c_j + d_k$ may repeat, so we have to use the lemma) since

$$\int_{\Omega} f = \sum_{j=0}^N c_j m(E_j) = \sum_{j=0}^N \sum_{k=0}^M c_j m(E_j \cap F_k),$$

and similarly we have

$$\int_{\Omega} g = \sum_{k=0}^M d_k m(F_k) = \sum_{k=0}^M \sum_{j=0}^N d_k m(E_j \cap F_k).$$

(c) Since $cf = \sum_{j=1}^N (cc_j) \chi_{E_j}$, we have

$$\int_{\Omega} cf = \sum_{j=1}^N cc_j m(E_j) = c \int_{\Omega} f.$$

(d) Write $h := g - f$, then h is simple and non-negative and $g = h + f$. Hence, by (b), $\int_{\Omega} g = \int_{\Omega} h + \int_{\Omega} f$, but by (a) $\int_{\Omega} h \geq 0$, so we're done. ■

3.2 Integration of non-negative measurable functions

We now pass from integrating non-negative simple functions to integrating general non-negative measurable functions. In this section we allow our measurable functions to take the value $+\infty$.

Definition 3.2.1. Let $f : \Omega \rightarrow \mathbb{R}$ and $g : \Omega \rightarrow \mathbb{R}$ be functions, we say that f majorizes g , or equivalently that g minorizes f , if

$$f(x) \geq g(x) \quad \forall x \in \Omega.$$

We sometimes use the phrase “ f dominates g ” instead of “ f majorizes g ”

Definition 3.2.2. Let $\Omega \subseteq \mathbb{R}^n$ be measurable, and $f : \Omega \rightarrow [0, \infty]$ be measurable and non-negative. We define the Lebesgue integral $\int_{\Omega} f$ of f on Ω by

$$\begin{aligned} \int_{\Omega} f &:= \sup \left\{ \int_{\Omega} s : s \text{ is simple and non-negative, and } s \text{ minorizes } f \right\} \\ &= \sup \left\{ \int_{\Omega} s : s \text{ is simple and non-negative, and } s(x) \leq f(x) \quad \forall x \in \Omega \right\}. \end{aligned}$$

Remark 3.2.1. If $\Omega' \subseteq \Omega$ is a measurable subset of Ω , then

$$\int_{\Omega'} f = \int_{\Omega'} f|_{\Omega'}.$$

Proposition 3.2.1. Let Ω be a measurable set, and let $f : \Omega \rightarrow [0, \infty]$ and $g : \Omega \rightarrow [0, \infty]$ be non-negative measurable functions.

- (a) We have $0 \leq \int_{\Omega} f \leq \infty$. Furthermore, $\int_{\Omega} f = 0$ iff $f(x) = 0$ for almost every $x \in \Omega$.
- (b) For any positive c , $\int_{\Omega} cf = c \int_{\Omega} f$.
- (c) If $f(x) \leq g(x)$ for all $x \in \Omega$, then $\int_{\Omega} f \leq \int_{\Omega} g$.
- (d) If $f(x) = g(x)$ for almost every $x \in \Omega$, then $\int_{\Omega} f = \int_{\Omega} g$.
- (e) If $\Omega' \subseteq \Omega$ is measurable, then

$$\int_{\Omega'} f = \int_{\Omega} f \chi_{\Omega'} \leq \int_{\Omega} f.$$

Proof.

- (a) $0 \leq \int_{\Omega} f \leq \infty$ since zero function is simple and non-negative, and minorizes f for all f . Now we show

$$\int_{\Omega} f = 0 \Leftrightarrow f(x) = 0 \text{ almost everywhere in } \Omega.$$

If $f \equiv 0$ almost everywhere in Ω , then for any simple non-negative function s minorizes f , we have $s(x) = 0$ whenever $f(x) = 0$. This shows $s \equiv 0$ almost everywhere in Ω . Hence, $\int_{\Omega} s = 0$ by definition. This implies $\int_{\Omega} f = 0$.

Now if $\int_{\Omega} f = 0$. For each $k \in \mathbb{N}$, define

$$A_k := \left\{ x \in \Omega : f(x) \geq \frac{1}{k} \right\}.$$

Since f is measurable, so $A_k = f^{-1}((\frac{1}{k}, \infty))$ is measurable for all $k \in \mathbb{N}$. Now define

$s_k := \frac{1}{k} \chi_{A_k}$. Then, s_k is a non-negative simple function. Note that $s_k \leq f$ on Ω . Thus,

$$\int_{\Omega} s_k \leq \int_{\Omega} f = 0$$

by definition, which shows $\int_{\Omega} s_k = 0$. Note that

$$0 = \int_{\Omega} s_k = \frac{1}{k} m(A_k) \Rightarrow m(A_k) = 0.$$

This is true for all $k \in \mathbb{N}$, and since

$$\{x \in \Omega : f(x) > 0\} = \bigcup_{k=1}^{\infty} A_k,$$

we have

$$m(\{x \in \Omega : f(x) > 0\}) \leq \sum_{k=1}^{\infty} m(A_k) = 0,$$

and we're done.

- (b) We first show $\int_{\Omega} cf \geq c \int_{\Omega} f$. Let s be any simple non-negative function minorizing f , then cs minorizes cf . Note that

$$\int_{\Omega} cs = c \int_{\Omega} s,$$

Hence,

$$\int_{\Omega} cf \geq \int_{\Omega} cs = c \int_{\Omega} s.$$

Since this is true for all simple non-negative s minorizing f , so

$$\int_{\Omega} cf \geq c \sup \left\{ \int_{\Omega} s : s \text{ simple and non-negative and minorizes } f \right\} = c \int_{\Omega} f.$$

For the reverse inequality, let t be any simple non-negative function s.t. $t \leq cf$, then since $c > 0$, we have $\frac{1}{c}t$ simple, non-negative, and $\frac{1}{c}t \leq f$. Hence,

$$\int_{\Omega} t = c \int_{\Omega} \frac{1}{c}t \leq c \int_{\Omega} f.$$

Since this is true for all simple, non-negative t s.t. $t \leq cf$, so we can take supremum over all such t and gets

$$\int_{\Omega} cf \leq c \int_{\Omega} f.$$

- (c) Let s be any simple, non-negative function s.t. $s \leq f$, then $s \leq g$. Thus,

$$\sup \left\{ \int_{\Omega} s : s \text{ simple, non-negative, and } s \leq f \right\} \leq \sup \left\{ \int_{\Omega} s : s \text{ simple, non-negative, and } s \leq g \right\},$$

i.e. $\int_{\Omega} f \leq \int_{\Omega} g$.

- (d) Let $N = \{x \in \Omega : f(x) \neq g(x)\}$, then $m(N) = 0$ and N is measurable, and we have $f = g$ on $\Omega \setminus N$. We first prove that $\int_{\Omega} f \leq \int_{\Omega} g$. Let s be any non-negative simple function s.t. $s \leq f$ on Ω , then define $t := s \chi_{\Omega \setminus N}$. Note that t is simple and non-negative. Moreover, if $x \in \Omega \setminus N$, then

$$t(x) = s(x) \leq f(x) = g(x),$$

while if $x \in N$, then $t(x) = 0 \leq g(x)$. Thus, $t \leq g$ on Ω . Since s and t only differ on N , so $s = t$ almost everywhere in Ω . In other words, $s - t \equiv 0$ almost everywhere in Ω . This shows $\int_{\Omega} s - t = 0$, i.e. $\int_{\Omega} s = \int_{\Omega} t$. Note that we have

$$\int_{\Omega} s = \int_{\Omega} t \leq \int_{\Omega} g$$

by definition of $\int_{\Omega} g$, and this is true for any simple, non-negative $s \leq f$, so taking supremum we have $\int_{\Omega} f \leq \int_{\Omega} g$. Now we can interchanging the role of f and g , and we can get $\int_{\Omega} g \leq \int_{\Omega} f$.

(e) We first show that $\int_{\Omega'} f = \int_{\Omega} f \chi_{\Omega'}$. First note that $\int_{\Omega'} f = \int_{\Omega'} f|_{\Omega'}$. We then first show

$$\int_{\Omega'} f \leq \int_{\Omega} f \chi_{\Omega'}.$$

Let s be any simple non-negative function on Ω' s.t. $s \leq f|_{\Omega'}$. Extend s to a function $\tilde{s}$ on Ω by

$$\tilde{s}(x) := \begin{cases} s(x), & \text{if } x \in \Omega'; \\ 0, & \text{if } \Omega \setminus \Omega'. \end{cases}$$

Then, $\tilde{s}$ is simple and non-negative on Ω , and we have $\tilde{s} \leq f \chi_{\Omega'}$. Moreover, we have

$$\int_{\Omega} \tilde{s} = \int_{\Omega'} s$$

by definition. Hence, every simple, non-negative s with $s \leq f|_{\Omega'}$ produces a simple, non-negative $\tilde{s}$ with $\tilde{s} \leq f \chi_{\Omega'}$ and $s, \tilde{s}$ has same Lebesgue integral. Hence, by taking supremum,

$$\int_{\Omega'} f|_{\Omega'} \leq \int_{\Omega} f \chi_{\Omega'} \Rightarrow \int_{\Omega'} f \leq \int_{\Omega} f \chi_{\Omega'}.$$

Now we show $\int_{\Omega} f \chi_{\Omega'} \leq \int_{\Omega'} f$. Let t be any simple, non-negative function on Ω s.t. $t \leq f \chi_{\Omega'}$, then $t(x) = 0$ for every $x \notin \Omega'$. Also, we have $t|_{\Omega'} \leq f|_{\Omega'}$, and $t|_{\Omega'}$ is a simple, non-negative function on Ω' . Hence,

$$\int_{\Omega} t = \int_{\Omega'} t|_{\Omega'} \leq \int_{\Omega'} f|_{\Omega'} = \int_{\Omega'} f.$$

Since this is true for all simple, non-negative t s.t. $t \leq f \chi_{\Omega'}$, so by taking supremum, we have

$$\int_{\Omega} f \chi_{\Omega'} \leq \int_{\Omega'} f.$$

Now we have shown $\int_{\Omega'} f = \int_{\Omega} f \chi_{\Omega'}$, and since $f \chi_{\Omega'} \leq f$ on Ω , so

$$\int_{\Omega} f \chi_{\Omega'} \leq \int_{\Omega} f.$$

■

Lecture 15

Remark 3.2.2. (d) of [Proposition 3.2.1](#) shows that one can modify the values of a function on any measure zero set (e.g. every rational number), and not affect its integral at all.

23 April.

Remark 3.2.3. We haven't shown that $\int_{\Omega}(f + g) = \int_{\Omega} f + \int_{\Omega} g$. From definition, it is easy to show $\int_{\Omega}(f + g) \geq \int_{\Omega} f + \int_{\Omega} g$, but proving equality requires more work and will be done later.

As we have seen in previous chapters, we cannot always interchange an integral with a limit (or with limit-like concepts such as supremum). However, for the Lebesgue integral it is possible to do so when the functions are increasing:

Theorem 3.2.1 (Lebesgue monotone convergence theorem). Let Ω be a measurable subset of $\mathbb{R}^n$, and let $(f_n)_{n=1}^\infty$ be a sequence of non-negative measurable functions from Ω to $[0, \infty]$ which are increasing in the sense that

$$0 \leq f_1(x) \leq f_2(x) \leq \dots \quad \forall x \in \Omega.$$

(Note that we are assuming $f_n(x)$ is increasing with respect to n ; this is different notion from $f_n(x)$ increasing with respect to x .) Then we have

$$0 \leq \int_{\Omega} f_1 \leq \int_{\Omega} f_2 \leq \int_{\Omega} f_3 \leq \dots$$

and

$$\int_{\Omega} \sup_n f_n = \sup_n \int_{\Omega} f_n.$$

Proof. Since $f_n(x) \leq f_{n+1}(x)$ for all $x \in \Omega$, then by (c) of [Proposition 3.2.1](#), we have

$$\int_{\Omega} f_n \leq \int_{\Omega} f_{n+1} \quad \forall n \geq 1.$$

In particular,

$$0 \leq \int_{\Omega} f_1 \leq \int_{\Omega} f_2 \leq \int_{\Omega} f_3 \leq \dots$$

It remains to prove that

$$\int_{\Omega} \sup_n f_n = \sup_n \int_{\Omega} f_n.$$

Set $f := \sup_n f_n$, then by [Lemma 2.5.5](#), f is also measurable and takes values in $[0, \infty]$. We first show

$$\int_{\Omega} f \geq \sup_n \int_{\Omega} f_n.$$

Since $f_n \leq f$ pointwise for every n , (c) of [Proposition 3.2.1](#) yields

$$\int_{\Omega} f_n \leq \int_{\Omega} f \quad \forall n.$$

Taking supremum over n , then we're done.

Now we show

$$\int_{\Omega} f \leq \sup_n \int_{\Omega} f_n.$$

Since

$$\int_{\Omega} f = \sup \left\{ \int_{\Omega} s : s \text{ is simple, non-negative, and } s \leq f \right\},$$

it suffices to show that for every simple, non-negative function s satisfying $s \leq f$, one has

$$\int_{\Omega} s \leq \sup_n \int_{\Omega} f_n.$$

Thus, we fix a non-negative simple function s with

$$s(x) \leq f(x) = \sup_n f_n(x) \quad \forall x \in \Omega.$$

We will prove that $\int_{\Omega} s \leq \sup_n \int_{\Omega} f_n$.

Now let $0 < \varepsilon < 1$. For each $n \geq 1$, define

$$E_n := \{x \in \Omega : f_n(x) \geq (1 - \varepsilon)s(x)\}.$$

Now we have f_n and s are measurable, we know that each set E_n is measurable since

$$E_n = (f_n - (1 - \varepsilon)s)^{-1}([0, \infty]),$$

and since

$$A_{n,k} := (f_n - (1 - \varepsilon)s)^{-1}((-\frac{1}{k}, \infty])$$

is measurable by definition for all $k \in \mathbb{N}$, we have $E_n = \bigcup_{k=1}^{\infty} A_{n,k}$ measurable.

Claim 3.2.1. $E_1 \subseteq E_2 \subseteq E_3 \subseteq \dots$ and $\bigcup_{n=1}^{\infty} E_n = \Omega$.

Proof. We first show $E_n \subseteq E_{n+1}$. This follows from $f_n \leq f_{n+1}$, so if $x \in E_n$, i.e. $f_n(x) \geq (1 - \varepsilon)s(x)$, then

$$f_{n+1}(x) \geq f_n(x) \geq (1 - \varepsilon)s(x) \Rightarrow x \in E_{n+1}.$$

To prove that $\bigcup_{n=1}^{\infty} E_n = \Omega$, fix $x \in \Omega$. Since

$$s(x) \leq f(x) = \sup_n f_n(x),$$

there exists some index $N(x)$ s.t. $f_{N(x)}(x) \geq s(x) \geq (1 - \varepsilon)s(x)$. Thus, $x \in E_{N(x)}$, and we're done. $\otimes$

Since s is a non-negative simple function, we know

$$s = \sum_{j=1}^N c_j \chi_{F_j}$$

for some $c_1, \dots, c_N \geq 0$ and $F_1, \dots, F_N$ are pairwise disjoint measurable subsets of Ω . For each n ,

$$\chi_{F_j} \chi_{E_n} = \chi_{F_j \cap E_n},$$

and hence

$$s \chi_{E_n} = \sum_{j=1}^N c_j \chi_{F_j} \chi_{E_n} = \sum_{j=1}^N c_j \chi_{F_j \cap E_n}.$$

Therefore,

$$\int_{E_n} s = \int_{\Omega} s \chi_{E_n} = \int_{\Omega} \sum_{j=1}^N c_j \chi_{F_j \cap E_n} = \sum_{j=1}^N c_j m(F_j \cap E_n).$$

Now since $E_n \uparrow \Omega$ by the claim, so for each j ,

$$F_j \cap E_1 \subseteq F_j \cap E_2 \subseteq \dots \quad \text{and} \quad \bigcup_{n=1}^{\infty} (F_j \cap E_n) = F_j.$$

Since each $F_j \cap E_n$ is measurable, continuity from below of Lebesgue measure gives

$$m(F_j \cap E_n) \uparrow m(F_j) \quad \text{as } n \rightarrow \infty.$$

Thus,

$$\int_{E_n} s = \sum_{j=1}^N c_j m(F_j \cap E_n) \rightarrow \sum_{j=1}^N c_j m(F_j) = \int_{\Omega} s \quad \text{as } n \rightarrow \infty.$$

Note that since $E_1 \subseteq E_2 \subseteq \dots$, we have

$$\int_{E_1} s \leq \int_{E_2} s \leq \int_{E_3} s \leq \dots$$

and $\lim_{n \rightarrow \infty} \int_{E_n} s = \int_{\Omega} s$, so

$$\sup_n \int_{E_n} s = \int_{\Omega} s.$$

Now by definition of E_n , we have

$$f_n(x) \geq (1 - \varepsilon)s(x) \quad \text{for every } x \in E_n.$$

Hence, by (c), (e) of [Proposition 3.2.1](#),

$$\int_{\Omega} f_n \geq \int_{E_n} f_n \geq \int_{E_n} (1 - \varepsilon)s = (1 - \varepsilon) \int_{E_n} s.$$

Taking supremum over n ,

$$\sup_n \int_{\Omega} f_n \geq (1 - \varepsilon) \sup_n \int_{E_n} s = (1 - \varepsilon) \int_{\Omega} s.$$

Since this holds for all $0 < \varepsilon < 1$, letting $\varepsilon \rightarrow 0$ gives

$$\sup_n \int_{\Omega} f_n \geq \int_{\Omega} s.$$

Hence, we're done. ■

Lemma 3.2.1 (Interchange of addition and integration). Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f, g : \Omega \rightarrow [0, \infty]$ be measurable functions. Then,

$$\int_{\Omega} (f + g) = \int_{\Omega} f + \int_{\Omega} g.$$

Proof. By [Lemma 3.1.3](#), there exist increasing sequence of non-negative simple functions

$$s_1 \leq s_2 \leq \cdots \leq f, \quad t_1 \leq t_2 \leq \cdots \leq g$$

s.t. $s_n(x) \uparrow f(x)$ and $t_n(x) \uparrow g(x)$ for every $x \in \Omega$. For each n , $s_n + t_n$ is again a non-negative simple function. Moreover, since both (s_n) and (t_n) are increasing, so is $(s_n + t_n)$. For each $x \in \Omega$, we have

$$s_n(x) + t_n(x) \uparrow f(x) + g(x),$$

and hence $s_n + t_n \uparrow f + g$ pointwise on Ω . Applying the monotone convergence theorem to the three increasing sequences

$$s_n \uparrow f, \quad t_n \uparrow g, \quad s_n + t_n \uparrow f + g,$$

we obtain

$$\int_{\Omega} f = \sup_n \int_{\Omega} s_n, \quad \int_{\Omega} g = \sup_n \int_{\Omega} t_n, \quad \int_{\Omega} (f + g) = \sup_n \int_{\Omega} (s_n + t_n).$$

Note that

$$\int_{\Omega} (s_n + t_n) = \int_{\Omega} s_n + \int_{\Omega} t_n \quad \text{for every } n$$

since s_n, t_n are non-negative simple functions. Thus,

$$\int_{\Omega} (f + g) = \sup_n \int_{\Omega} (s_n + t_n) = \sup_n \left(\int_{\Omega} s_n + \int_{\Omega} t_n \right)$$

Now let $a_n := \int_{\Omega} s_n$ and $b_n := \int_{\Omega} t_n$, then (a_n) and (b_n) are increasing sequences in $[0, \infty]$. Consequently,

$$a_n + b_n \uparrow \left(\sup_n a_n \right) + \left(\sup_n b_n \right),$$

Thus, $\sup_n (a_n + b_n) = \sup_n a_n + \sup_n b_n$. It follows that

$$\int_{\Omega} (f + g) = \sup_n \left(\int_{\Omega} s_n + \int_{\Omega} t_n \right) = \sup_n \int_{\Omega} s_n + \sup_n \int_{\Omega} t_n = \int_{\Omega} f + \int_{\Omega} g.$$

■

Lecture 16

Once we can interchange an integral with a sum of two functions, we can handle finite sums by induction. 28 April.
More surprisingly, we can handle infinite sums of non-negative functions as well:

Corollary 3.2.1. If $\Omega \subseteq \mathbb{R}^n$ is measurable and $g_1, g_2, \dots$ are a sequence of non-negative measurable functions from Ω to $[0, \infty]$, then

$$\int_{\Omega} \sum_{n=1}^{\infty} g_n = \sum_{n=1}^{\infty} \int_{\Omega} g_n.$$

Proof. For each integer $N \geq 1$, define $G_N := \sum_{n=1}^N g_n$. Since g_n is measurable and non-negative, it follows that each G_N is also measurable and non-negative. Moreover, the sequence (G_N) is increasing pointwise, because

$$G_{N+1}(x) = G_N(x) + g_{N+1}(x) \geq G_N(x) \quad \forall x \in \Omega.$$

Hence, we may apply the Lebesgue monotone convergence theorem to the sequence (G_N) . We obtain

$$\int_{\Omega} \sup_N G_N = \sup_N \int_{\Omega} G_N.$$

Now since for each $x \in \Omega$, $G_1(x) \leq G_2(x) \leq \dots$, so $\sup_N G_N(x) = \lim_{N \rightarrow \infty} G_N(x) = \sum_{n=1}^{\infty} g_n(x)$. This gives

$$\int_{\Omega} \sup_N G_N = \int_{\Omega} \sum_{n=1}^{\infty} g_n.$$

Now since for each $N \in \mathbb{N}$,

$$\int_{\Omega} G_N = \int_{\Omega} \sum_{n=1}^N g_n = \sum_{n=1}^N \int_{\Omega} g_n.$$

Hence,

$$\sup_N \int_{\Omega} G_N = \sup_N \sum_{n=1}^N \int_{\Omega} g_n.$$

Note that the sequence $\left(\sum_{n=1}^N \int_{\Omega} g_n\right)_{N=1}^{\infty}$ is increasing, so

$$\sup_N \sum_{n=1}^N \int_{\Omega} g_n = \sum_{n=1}^{\infty} \int_{\Omega} g_n.$$

Thus,

$$\int_{\Omega} \sum_{n=1}^{\infty} g_n = \int_{\Omega} \sup_N G_N = \sup_N \int_{\Omega} G_N = \sum_{n=1}^{\infty} \int_{\Omega} g_n.$$

■

Remark 3.2.4. We do not need to assume anything about convergence of above sums; it may well happen that both sides equal $+\infty$. However, we do need to assume non-negativity.

One might similarly ask whether we can interchange limits and integrals; i.e. whether

$$\int_{\Omega} \lim_{n \rightarrow \infty} f_n = \lim_{n \rightarrow \infty} \int_{\Omega} f_n$$

holds in general. Unfortunately this is not true: For $n = 1, 2, 3, \dots$ let $f_n : \mathbb{R} \rightarrow \mathbb{R}$ be $f_n = \chi_{[n, n+1)}$. Then, $f_n(x) \rightarrow 0$ for every x , but $\int_{\mathbb{R}} f_n = 1$ for every n since f_n is a simple function, and thus

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}} f_n = 1 \neq 0.$$

However, the following lemma of Fatou shows that the reverse cannot happen: the limiting function cannot have larger integral than the (liminf of the) original integrals.

Lemma 3.2.2 (Fatou's lemma). Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f_1, f_2, \dots : \Omega \rightarrow [0, \infty]$ be a sequence of non-negative measurable functions. Then

$$\int_{\Omega} \liminf_{n \rightarrow \infty} f_n \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f_n.$$

Proof. For each $N \in \mathbb{N}$, define

$$g_N(x) := \inf_{n \geq N} f_n(x) \quad \forall x \in \Omega.$$

Then g_N is non-negative and measurable. Moreover, $g_1(x) \leq g_2(x) \leq \dots$ for all $x \in \Omega$. By the definition of $\liminf$,

$$\liminf_{n \rightarrow \infty} f_n(x) = \sup_{N \geq 1} g_N(x).$$

Since $(g_N(x))$ is increasing, this means that $g_N(x) \uparrow \liminf_{n \rightarrow \infty} f_n(x)$ for all $x \in \Omega$. Note that we can apply Lebesgue monotone convergence theorem since (g_N) is increasing:

$$\int_{\Omega} \liminf_{n \rightarrow \infty} f_n(x) = \int_{\Omega} \sup_{N \geq 1} g_N = \sup_{N \geq 1} \int_{\Omega} g_N.$$

On the other hand, for every N and every $n \geq N$, we have $g_N \leq f_n$. Hence,

$$\int_{\Omega} g_N \leq \int_{\Omega} f_n \quad \forall n \geq N.$$

Taking infimum over all $n \geq N$, we get $\int_{\Omega} g_N \leq \inf_{n \geq N} \int_{\Omega} f_n$. Now take the supremum over N to obtain

$$\sup_{N \geq 1} \int_{\Omega} g_N \leq \sup_{N \geq 1} \inf_{n \geq N} \int_{\Omega} f_n.$$

Combining previous result, we have

$$\int_{\Omega} \liminf_{n \rightarrow \infty} f_n(x) = \sup_{N \geq 1} \int_{\Omega} g_N \leq \sup_{N \geq 1} \inf_{n \geq N} \int_{\Omega} f_n = \liminf_{n \rightarrow \infty} \int_{\Omega} f_n.$$

■

Note that we are allowing our functions to take the value $+\infty$ at some points. It is even possible for a function to take the value $+\infty$ but still have a finite integral; for instance, if E is a set of measure zero, and $f : \Omega \rightarrow \mathbb{R}$ is equal to $+\infty$ on E but equal 0 everywhere else, then $\int_{\Omega} f = 0$ by (a) of [Proposition 3.2.1](#). However, if the integral is finite, the function must be finite almost everywhere:

Lemma 3.2.3. Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f : \Omega \rightarrow [0, \infty]$ be a non-negative measurable function s.t. $\int_{\Omega} f$ is finite. Then f is finite almost everywhere, i.e. the set $\{x \in \Omega : f(x) = +\infty\}$ has measure zero.

Proof. For each integer $k \geq 1$, define $A_k := \{x \in \Omega : f(x) \geq k\}$. Since f is measurable and $[k, \infty]$ is a Borel subset of $[0, \infty]$, each set A_k is measurable.

On the set A_k , we have $f(x) \geq k$, while outside A_k the function $k\chi_{A_k}$ vanishes. Thus, for every $x \in \Omega$, $k\chi_{A_k}(x) \leq f(x)$. Because both $k\chi_{A_k}$ and f are non-negative measurable functions, the monotonicity property gives $\int_{\Omega} k\chi_{A_k} \leq \int_{\Omega} f$. Now

$$\int_{\Omega} k\chi_{A_k} = k \int_{\Omega} \chi_{A_k} = km(A_k),$$

so we obtain $km(A_k) \leq \int_{\Omega} f$. Since $\int_{\Omega} f < \infty$, it follows that $m(A_k) \leq \frac{1}{k} \int_{\Omega} f$, and thus

$$m(A_k) \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Note that

$$\{x \in \Omega : f(x) = +\infty\} = \bigcap_{k=1}^{\infty} A_k.$$

Also, $A_1 \supseteq A_2 \supseteq A_3 \supseteq \dots$ by definition, and by continuity of measure (HW5 P2),

$$m\left(\bigcap_{k=1}^{\infty} A_k\right) = \lim_{k \rightarrow \infty} m(A_k).$$

Since we have shown $\lim_{k \rightarrow \infty} m(A_k) = 0$, so we have

$$m(\{x \in \Omega : f(x) = +\infty\}) = m\left(\bigcap_{k=1}^{\infty} A_k\right) = 0.$$

■

Lemma 3.2.4 (Borel-Cantelli lemma). Let $\Omega_1, \Omega_2, \dots$ be measurable subsets of $\mathbb{R}^n$ such that

$$\sum_{n=1}^{\infty} m(\Omega_n) < \infty.$$

Then the set

$$\{x \in \mathbb{R}^n : x \in \Omega_k \text{ for infinitely many } k\}$$

has measure zero. Equivalently, for almost every $x \in \mathbb{R}^n$, there exists $N = N(x)$ s.t. $x \notin \Omega_k$ for all $k \geq N$.

Proof. Define

$$G(x) := \sum_{n=1}^{\infty} \chi_{\Omega_n}(x), \quad \forall x \in \mathbb{R}^n,$$

where the value $+\infty$ is allowed.

For each n , χ_{Ω_n} is measurable and non-negative. Hence, the partial sums

$$G_N(x) := \sum_{n=1}^N \chi_{\Omega_n}(x)$$

are non-negative measurable functions. Since $G_N(x) \uparrow G(x)$ for every $x \in \mathbb{R}^n$. It follows that $G = \sup_N G_N$ is also measurable and non-negative. By [Corollary 3.2.1](#), we have

$$\int_{\mathbb{R}^n} G = \int_{\mathbb{R}^n} \sum_{n=1}^{\infty} \chi_{\Omega_n} = \sum_{n=1}^{\infty} \int_{\mathbb{R}^n} \chi_{\Omega_n}.$$

Since $\int_{\mathbb{R}^n} \chi_{\Omega_n} = m(\Omega_n)$, this becomes

$$\int_{\mathbb{R}^n} G = \sum_{n=1}^{\infty} m(\Omega_n) < \infty.$$

By [Lemma 3.2.3](#),

$$m(\{x \in \mathbb{R}^n : G(x) = +\infty\}) = 0.$$

Note that

$$G(x) = +\infty \Leftrightarrow \sum_{n=1}^{\infty} \chi_{\Omega_n}(x) = \infty \Leftrightarrow x \in \Omega_k \text{ for infinitely many } k,$$

so

$$\{x \in \mathbb{R}^n : G(x) = +\infty\} = \{x \in \mathbb{R}^n : x \in \Omega_k \text{ for infinitely many } k\},$$

and we're done. ■

3.3 Integration of Absolutely Integrable Functions

We have now completed the theory of the Lebesgue integral for non-negative functions. We now consider how to integrate functions which can be both positive and negative. However, we wish to avoid the indefinite expression $+\infty + (-\infty)$, so we restrict our attention to a subclass of measurable functions: the absolutely integrable functions.

Definition 3.3.1. Let Ω be a measurable subset of $\mathbb{R}^n$. A measurable function $f : \Omega \rightarrow \mathbb{R}^*$ is said to be absolutely integrable if the integral $\int_{\Omega} |f|$ is finite.

Of course, $|f|$ is always non-negative, so this definition makes sense even if f changes sign. Absolutely integrable functions are also known as $L^1(\Omega)$ functions.

If $f : \Omega \rightarrow \mathbb{R}^*$ is a function, we define the positive part $f^+ : \Omega \rightarrow [0, \infty]$ and the negative part $f^- : \Omega \rightarrow [0, \infty]$ by

$$f^+ := \max(f, 0), \quad f^- := -\min(f, 0).$$

Equivalently, for each $x \in \Omega$,

$$f^+(x) = \max\{f(x), 0\}, \quad f^-(x) = \max\{-f(x), 0\}.$$

From [Corollary 2.5.3](#), which can be extended to $\mathbb{R}^*$ -valued functions without difficulty, we know that f^+, f^- are measurable. Clearly f^+ and f^- are non-negative.

We also have the pointwise identities

$$f = f^+ - f^-, \quad |f| = f^+ + f^-.$$

Definition 3.3.2 (Lebesgue integral). Let $f : \Omega \rightarrow \mathbb{R}^*$ be an absolutely integrable function. We define the Lebesgue integral $\int_{\Omega} f$ of f to be the quantity

$$\int_{\Omega} f := \int_{\Omega} f^+ - \int_{\Omega} f^-.$$

Remark 3.3.1. Note that since f is absolutely integrable, $\int_{\Omega} f^+$ and $\int_{\Omega} f^-$ are less than or equal to $\int_{\Omega} |f|$ and hence are finite. Thus, $\int_{\Omega} f$ is always finite; we are never encountering the indeterminate form $+\infty - (+\infty)$.

Remark 3.3.2. We have the useful triangle inequality

$$\left| \int_{\Omega} f \right| \leq \int_{\Omega} |f|.$$

This is because

$$\left| \int_{\Omega} f \right| = \left| \int_{\Omega} f^+ - \int_{\Omega} f^- \right| \leq \left| \int_{\Omega} f^+ \right| + \left| \int_{\Omega} f^- \right| = \int_{\Omega} f^+ + \int_{\Omega} f^- = \int_{\Omega} f^+ + f^- = \int_{\Omega} |f|.$$

Proposition 3.3.1. Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f, g : \Omega \rightarrow \mathbb{R}^*$ be absolutely integrable functions. Then the following statements hold.

- (a) $\forall c \in \mathbb{R}$, the function cf is absolutely integrable, and $\int_{\Omega} cf = c \int_{\Omega} f$.
- (b) The function $f + g$ is absolutely integrable, and $\int_{\Omega} (f + g) = \int_{\Omega} f + \int_{\Omega} g$.
- (c) If $f(x) \leq g(x)$ for all $x \in \Omega$, then $\int_{\Omega} f \leq \int_{\Omega} g$.
- (d) If $f(x) = g(x)$ for almost every $x \in \Omega$, then $\int_{\Omega} f = \int_{\Omega} g$.

Proof.

- (a) We first show cf is absolutely integrable. Since $|cf| = |c||f|$, we have $\int_{\Omega} |cf| = |c| \int_{\Omega} |f|$ since f is absolutely integrable and thus $\int_{\Omega} |f| < \infty$.

Now it remains to compute its integral. We consider two cases.

Case 1: $c \geq 0$. In this case, $(cf)^+ = cf^+$ and $(cf)^- = cf^-$. Thus,

$$\begin{aligned} \int_{\Omega} cf &= \int_{\Omega} (cf)^+ - \int_{\Omega} (cf)^- = \int_{\Omega} cf^+ - \int_{\Omega} cf^- = c \int_{\Omega} f^+ - c \int_{\Omega} f^- \\ &= c \left(\int_{\Omega} f^+ - \int_{\Omega} f^- \right) = c \int_{\Omega} f. \end{aligned}$$

Case 2: $c < 0$. Now we have $(cf)^+ = -cf^-$ and $(cf)^- = -cf^+$. Hence,

$$\begin{aligned} \int_{\Omega} cf &= \int_{\Omega} (cf)^+ - \int_{\Omega} (cf)^- = \int_{\Omega} (-c)f^- - \int_{\Omega} (-c)f^+ = (-c) \int_{\Omega} f^- - (-c) \int_{\Omega} f^+ \\ &= c \int_{\Omega} f^+ - c \int_{\Omega} f^- = c \left(\int_{\Omega} f^+ - \int_{\Omega} f^- \right) = c \int_{\Omega} f. \end{aligned}$$

- (b) We first show that $f + g$ is absolutely integrable. Note that

$$\int_{\Omega} |f + g| \leq \int_{\Omega} |f| + |g| = \int_{\Omega} |f| + \int_{\Omega} |g| < \infty$$

by the properties of Lebesgue integral on non-negative measurable functions, and hence we're done.

Now since

$$f + g = (f^+ - f^-) + (g^+ - g^-) = (f^+ + g^+) - (f^- + g^-),$$

where $f^+ + g^+$ and $f^- + g^-$ are non-negative measurable functions. Hence,

$$\begin{aligned} \int_{\Omega} (f + g) &= \int_{\Omega} (f^+ + g^+) - \int_{\Omega} (f^- + g^-) \\ &= \int_{\Omega} (f^+ + g^+) - \int_{\Omega} (f^- + g^-) = \left(\int_{\Omega} f^+ + \int_{\Omega} g^+ \right) - \left(\int_{\Omega} f^- + \int_{\Omega} g^- \right) \\ &= \left(\int_{\Omega} f^+ - \int_{\Omega} f^- \right) + \left(\int_{\Omega} g^+ - \int_{\Omega} g^- \right) = \int_{\Omega} f + \int_{\Omega} g. \end{aligned}$$

- (c) Assume that $f(x) \leq g(x)$ for all $x \in \Omega$, then $g - f$ is non-negative everywhere on Ω . By (a),(b), we know $g - f = g + (-f)$ is absolutely integrable. Also,

$$\int_{\Omega} g = \int_{\Omega} (f + (g - f)) = \int_{\Omega} f + \int_{\Omega} (g - f).$$

Since $g - f$ is non-negative and measurable, $\int_{\Omega} (g - f) \geq 0$, and thus $\int_{\Omega} g \geq \int_{\Omega} f$, and we're done.

- (d) Assume that $f(x) = g(x)$ for almost every $x \in \Omega$. Define $h := f - g$. Then $h = 0$ almost everywhere. Also, since f, g are absolutely integrable, (a) and (b) shows that h is absolutely integrable. Because $h = 0$ almost everywhere, we also have $|h| = 0$ almost everywhere. This shows $\int_{\Omega} |h| = 0$ by properties of Lebesgue integral for non-negative measurable functions. Now apply the triangle inequality:

$$\left| \int_{\Omega} h \right| \leq \int_{\Omega} |h| = 0,$$

which shows $\int_{\Omega} h = 0$. This shows

$$0 = \int_{\Omega} h = \int_{\Omega} f - g = \int_{\Omega} f - \int_{\Omega} g,$$

and thus $\int_{\Omega} f = \int_{\Omega} g$. ■

Lecture 17

As discussed in the previous section, limits and integrals cannot in general be interchanged. That is, 30 April.
even if $f_n \rightarrow f$ pointwise on Ω , it may happen that

$$\lim_{n \rightarrow \infty} \int_{\Omega} f_n \neq \int_{\Omega} \lim_{n \rightarrow \infty} f_n.$$

Thus, in order to justify passing a limit through an integral, one needs an additional hypothesis. The Lebesgue dominated convergence theorem provides one of the most useful such criteria: if the sequence is dominated by a single absolutely integrable function, then the interchange of limit and integral is valid.

Theorem 3.3.1 (Lebesgue dominated convergence theorem). Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f_1, f_2, \dots$ be measurable functions from Ω to $\mathbb{R}^*$ such that $f_n(x)$ converges pointwise for every $x \in \Omega$. Suppose moreover that there exists an absolutely integrable function $F : \Omega \rightarrow [0, \infty]$ such that $|f_n(x)| \leq F(x)$ for all $x \in \Omega$ and all $n \geq 1$. Then, if we write $f(x) := \lim_{n \rightarrow \infty} f_n(x)$, we have

$$\int_{\Omega} f = \lim_{n \rightarrow \infty} \int_{\Omega} f_n.$$

Proof. Since F is non-negative, measurable, and absolutely integrable, we know $F < \infty$ almost everywhere by Lemma 3.2.3. In other words, $E := \{x \in \Omega : F(x) = +\infty\}$ has measure zero. If we replace Ω by $\Omega \setminus E$, then none of the relevant integrals changes, because deleting a measure zero set does not affect the integral of an absolutely integrable function. Hence, without loss of generality, we may assume from now on that $F(x) < \infty$ for every $x \in \Omega$. Since $|f_n(x)| \leq F(x)$ for all n , this also implies that each $f_n(x)$ is finite for every $x \in \Omega$.

Because each f_n is measurable and $f_n \rightarrow f$ pointwise, f is measurable by Lemma 2.5.5. Next, since $|f_n(x)| \leq F(x)$ for all n and all $x \in \Omega$, we may pass to the limit in n and obtain

$$|f(x)| \leq F(x) \quad \text{for all } x \in \Omega.$$

Therefore,

$$\int_{\Omega} |f| \leq \int_{\Omega} F < \infty,$$

so f is absolutely integrable. The same argument can show that f_n is absolutely integrable, because

$$\int_{\Omega} |f_n| \leq \int_{\Omega} F < \infty.$$

Thus, all of the integrals appearing below are finite and well-defined.

For each n and each $x \in \Omega$, we have $|f_n(x)| \leq F(x)$. Hence,

$$-F(x) \leq f_n(x) \leq F(x).$$

It follows that $F(x) + f_n(x) \geq 0$ for all $x \in \Omega$. So the function $F + f_n$ are non-negative and measurable. Moreover, since $f_n(x) \rightarrow f(x)$, we also have

$$F(x) + f_n(x) \rightarrow F(x) + f(x) \quad \text{pointwise on } \Omega.$$

Fatou's lemma therefore gives

$$\int_{\Omega} (F + f) = \int_{\Omega} \liminf_{n \rightarrow \infty} (F + f_n) \leq \liminf_{n \rightarrow \infty} \int_{\Omega} (F + f_n).$$

Using additivity of the integral for absolutely integrable functions, this becomes

$$\int_{\Omega} F + \int_{\Omega} f \leq \liminf_{n \rightarrow \infty} \left(\int_{\Omega} F + \int_{\Omega} f_n \right).$$

Since $\int_{\Omega} F$ is a fixed finite number, we may subtract it from both sides and obtain

$$\int_{\Omega} f \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f_n.$$

Similarly, since $|f_n(x)| \leq F(x)$, we have $F(x) - f_n(x) \geq 0$ for all $x \in \Omega$. Thus, $F - f_n$ is non-negative and measurable for all n . Since $f_n(x) \rightarrow f(x)$ pointwise, we also have

$$F(x) - f_n(x) \rightarrow F(x) - f(x) \quad \text{pointwise on } \Omega.$$

Applying **Fatou's lemma** once more, we obtain

$$\int_{\Omega} (F - f) = \int_{\Omega} \liminf_{n \rightarrow \infty} (F - f_n) \leq \liminf_{n \rightarrow \infty} \int_{\Omega} (F - f_n).$$

By additivity,

$$\int_{\Omega} F - \int_{\Omega} f \leq \liminf_{n \rightarrow \infty} \left(\int_{\Omega} F - \int_{\Omega} f_n \right) = \int_{\Omega} F - \limsup_{n \rightarrow \infty} \int_{\Omega} f_n.$$

This gives

$$\int_{\Omega} f \geq \limsup_{n \rightarrow \infty} \int_{\Omega} f_n.$$

Now we have

$$\limsup_{n \rightarrow \infty} \int_{\Omega} f_n \leq \int_{\Omega} f \leq \liminf_{n \rightarrow \infty} \int_{\Omega} f_n.$$

But for real sequence (a_n) we always has

$$\liminf_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} a_n.$$

Therefore, the only possibility is that

$$\liminf_{n \rightarrow \infty} \int_{\Omega} f_n = \limsup_{n \rightarrow \infty} \int_{\Omega} f_n = \int_{\Omega} f.$$

Hence, the sequence $(\int_{\Omega} f_n)$ converges, and its limit is $\int_{\Omega} f$ by **Theorem B.0.1**. That is,

$$\lim_{n \rightarrow \infty} \int_{\Omega} f_n = \int_{\Omega} f.$$

This proves the dominated convergence theorem. ■

Finally, we record a lemma which is not particularly interesting in itself, but we will have some useful consequences later in these notes.

Definition 3.3.3 (Upper and lower Lebesgue integral). Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f : \Omega \rightarrow \mathbb{R}$ (not necessarily measurable). We define the upper Lebesgue integral $\overline{\int_{\Omega}} f$ to be

$$\overline{\int_{\Omega}} f := \inf \left\{ \int_{\Omega} g : g \text{ is an absolutely integrable function from } \Omega \text{ to } \mathbb{R} \text{ that majorizes } f \right\},$$

and the lower Lebesgue integral $\underline{\int_{\Omega}} f$ to be

$$\underline{\int_{\Omega}} f := \sup \left\{ \int_{\Omega} g : g \text{ is an absolutely integrable function from } \Omega \text{ to } \mathbb{R} \text{ that minorizes } f \right\}.$$

We first record two elementary consequences of this definition.

It is easy to see that

$$\underline{\int_{\Omega}} f \leq \overline{\int_{\Omega}} f.$$

Next, suppose f is absolutely integrable. Then f is both an absolutely integrable minorant and an absolutely integrable majorant of itself. Therefore,

$$\underline{\int_{\Omega}} f = \sup \left\{ \int_{\Omega} u : u \text{ is absolutely integrable and } u \leq f \right\} \geq \int_{\Omega} f.$$

On the other hand, if $u \leq f$ for some absolutely integrable u , then

$$\int_{\Omega} u \leq \int_{\Omega} f.$$

Taking the supremum over all absolutely integrable $u \leq f$, we obtain

$$\underline{\int_{\Omega}} f \leq \int_{\Omega} f.$$

Hence, $\underline{\int_{\Omega}} f = \int_{\Omega} f$.

Similarly, since f is an absolutely integrable majorant of itself,

$$\overline{\int_{\Omega}} f = \inf \left\{ \int_{\Omega} v : v \text{ is absolutely integrable and } f \leq v \right\} \leq \int_{\Omega} f.$$

Conversely, if $f \leq v$ for some absolutely integrable v , then

$$\int_{\Omega} f \leq \int_{\Omega} v.$$

Taking the infimum over all absolutely integrable $v \geq f$, we get $\int_{\Omega} f \leq \overline{\int_{\Omega}} f$. Thus, $\overline{\int_{\Omega}} f = \int_{\Omega} f$. Consequently, whenever f is absolutely integrable,

$$\underline{\int_{\Omega}} f = \int_{\Omega} f = \overline{\int_{\Omega}} f.$$

The converse is also true:

Lemma 3.3.1. Let Ω be a measurable subset of $\mathbb{R}^n$, and let $f : \Omega \rightarrow \mathbb{R}$ be a function, not assumed a priori to be measurable. Suppose that there exists $A \in \mathbb{R}$ s.t.

$$\overline{\int_{\Omega}} f = \underline{\int_{\Omega}} f = A.$$

Then f is absolutely integrable, and

$$\int_{\Omega} f = \overline{\int_{\Omega}} f = \underline{\int_{\Omega}} f = A.$$

Proof. We will show that f agree almost everywhere with an absolutely integrable function, and hence f itself is absolutely integrable with integral A .

By definition,

$$\overline{\int_{\Omega}} f = \inf \left\{ \int_{\Omega} g : g \text{ is absolutely integrable and } g \geq f \right\}.$$

Since this infimum is equal to A , for every $n \geq 1$, there exists an absolutely integrable function

$f_n^+ : \Omega \rightarrow \mathbb{R}$ s.t. f_n^+ majorizes f and

$$\int_{\Omega} f_n^+ \leq A + \frac{1}{n}.$$

Similarly, since

$$A = \int_{\Omega} f = \sup \left\{ \int_{\Omega} g : g \text{ is absolutely integrable and } g \leq f \right\},$$

for every integer $n \geq 1$ there exists an absolutely integrable function $f_n^- : \Omega \rightarrow \mathbb{R}$ s.t. f_n^- minorizes f and

$$\int_{\Omega} f_n^- \geq A - \frac{1}{n}.$$

Thus, for every n we have $f_n^- \leq f \leq f_n^+$ on Ω .

Now set

$$F^+(x) := \inf_{n \geq 1} f_n^+(x), \quad F^-(x) := \sup_{n \geq 1} f_n^-(x), \quad x \in \Omega.$$

Since each f_n^+ and f_n^- is measurable, we know F^+ and F^- are measurable by [Lemma 2.5.5](#). Since $f(x) \leq f_n^+(x)$ for all n , we have

$$f(x) \leq \inf_{n \geq 1} f_n^+(x) = F^+(x).$$

Similarly, we have

$$F^-(x) = \sup_{n \geq 1} f_n^-(x) \leq f(x).$$

Hence,

$$F^-(x) \leq f(x) \leq F^+(x) \quad \forall x \in \Omega.$$

We first consider F^+ . Since $F^+ \leq f_1^+$ pointwise and f_1^+ is absolutely integrable, we would like to conclude that F^+ is also absolutely integrable. For this we also need a lower bound by an absolutely integrable function.

Because $f_n^- \leq f \leq F^+$ for every n , in particular we have

$$f_1^- \leq F^+ \leq f_1^+.$$

Hence,

$$|F^+| \leq \max\{|f_1^-|, |f_1^+|\} \leq |f_1^-| + |f_1^+|.$$

Since f_1^- and f_1^+ are absolutely integrable, so is $|f_1^-| + |f_1^+|$. By comparison, F^+ is absolutely integrable.

The same argument applies to F^- , by $f_1^- \leq F^- \leq f_1^+$, we have

$$|F^-| \leq |f_1^-| + |f_1^+|$$

and thus F^- is also absolutely integrable.

Now since $F^+ \leq f_n^+$ for all n and they are both absolutely integrable,

$$\int_{\Omega} F^+ \leq \int_{\Omega} f_n^+ \leq A + \frac{1}{n} \quad \forall n \geq 1.$$

Letting $n \rightarrow \infty$, we obtain

$$\int_{\Omega} F^+ \leq A.$$

Similarly, we have

$$A - \frac{1}{n} \leq \int_{\Omega} f_n^- \leq \int_{\Omega} F^-.$$

Letting $n \rightarrow \infty$, we have

$$A \leq \int_{\Omega} F^-.$$

On the other hand, since $F^- \leq F^+$ pointwise, we have

$$\int_{\Omega} F^- \leq \int_{\Omega} F^+.$$

Combining the three inequalities,

$$A \leq \int_{\Omega} F^- \leq \int_{\Omega} F^+ \leq A.$$

Thus,

$$\int_{\Omega} F^- = \int_{\Omega} F^+ = A.$$

Now since

$$\int_{\Omega} (F^+ - F^-) = \int_{\Omega} F^+ - \int_{\Omega} F^- = A - A = 0,$$

we know $F^+ - F^-$ is zero almost everywhere since $F^+ - F^-$ is a non-negative measurable function. Hence, $F^+ = F^-$ almost everywhere on Ω .

Recall that

$$F^-(x) \leq f(x) \leq F^+(x) \quad \forall x \in \Omega.$$

At every point where $F^+(x) = F^-(x)$, we have

$$f(x) = F^+(x) = F^-(x).$$

Since $F^+ = F^-$ almost everywhere, it follows that $f = F^+$ almost everywhere. Now F^+ is measurable and absolutely integrable. A function which agrees almost everywhere with a measurable function is measurable by [Theorem B.0.2](#), so f is measurable. Also, $|f| = |F^+|$ almost everywhere, and therefore

$$\int_{\Omega} |f| = \int_{\Omega} |F^+| < \infty.$$

Thus, f is absolutely integrable.

Finally, since $f = F^+$ almost everywhere, we have

$$\int_{\Omega} f = \int_{\Omega} F^+ = A.$$

Therefore,

$$\int_{\Omega} f = \overline{\int_{\Omega} f} = \underline{\int_{\Omega} f} = A.$$

■

3.4 Comparison with the Riemann Integral

We have spent a lot of effort constructing the Lebesgue integral, but have not yet addressed the question of how to actually compute any Lebesgue integrals, and whether Lebesgue integration is any different from the Riemann integral (say for integrals in one dimension). We now show that the Lebesgue integral is a generalization of the Riemann integral. To clarify the following discussion, we shall temporarily distinguish the Riemann integral from the Lebesgue integral by writing the Riemann integral as $R. \int_I f$. Recall that, in the definition of the Riemann integral on a bounded interval $I \subseteq \mathbb{R}$, one starts with a bounded function $f : I \rightarrow \mathbb{R}$. The boundedness of both the interval I and the function f will be important. Our objective in this section is to prove that every Riemann integrable function on a bounded interval is Lebesgue integrable, indeed absolutely integrable, and that the two integrals agree.

Proposition 3.4.1. Let $I \subseteq \mathbb{R}$ be a bounded interval, and let $f : I \rightarrow \mathbb{R}$ be a Riemann integrable

function. Then f is also absolutely integrable, and

$$\int_I f = R. \int_I f.$$

Proof. Write $A := R. \int_I f$. Since f is Riemann integrable, f 's upper integral and lower integral agrees, and thus for every $\varepsilon > 0$, there exists a partition $\mathcal{P}$ of I into finitely many subintervals J s.t.

$$A - \varepsilon \leq \sum_{J \in \mathcal{P}} |J| \inf_{x \in J} f(x) \leq A \leq \sum_{J \in \mathcal{P}} |J| \sup_{x \in J} f(x) \leq A + \varepsilon,$$

where $|J|$ represents the length of J . Note that $m(J) = |J|$ since J is a one-dimensional box.

Define simple functions

$$f_\varepsilon^-(x) = \sum_{J \in \mathcal{P}} \left(\inf_{t \in J} f(t) \right) \chi_J(x), \quad f_\varepsilon^+(x) = \sum_{J \in \mathcal{P}} \left(\sup_{t \in J} f(t) \right) \chi_J(x).$$

By construction, we have $f_\varepsilon^- \leq f \leq f_\varepsilon^+$ pointwise on I . Note that $f_\varepsilon^-, f_\varepsilon^+$ are simple, and thus measurable. Moreover, since I is bounded, so it has finite measure, so f_ε^- and f_ε^+ are absolutely integrable. Also,

$$\int_I f_\varepsilon^- = \sum_{J \in \mathcal{P}} \inf_{t \in J} f(t) \cdot |J|, \quad \int_I f_\varepsilon^+ = \sum_{J \in \mathcal{P}} \sup_{t \in J} f(t) \cdot |J|.$$

Therefore,

$$A - \varepsilon \leq \int_I f_\varepsilon^- \leq A \leq \int_I f_\varepsilon^+ \leq A + \varepsilon.$$

Since f_ε^+ majorizes f , and f_ε^+ is absolutely integrable, by definition we have $\overline{\int_I f} \leq \int_I f_\varepsilon^+$. Similarly, since f_ε^- minorizes f and is absolutely integrable, we have $\underline{\int_I f} \geq \int_I f_\varepsilon^-$. Hence,

$$A - \varepsilon \leq \underline{\int_I f} \leq \overline{\int_I f} \leq A + \varepsilon \quad \forall \varepsilon > 0.$$

This shows that $\underline{\int_I f} = \overline{\int_I f} = A$. By the previous lemma, f is absolutely integrable and $\int_I f = A$. Thus,

$$\int_I f = R. \int_I f.$$

■

Appendix

Appendix A

Midterm Practice

A.1 HW6

Problem A.1.1. Let A be an open interval in $\mathbb{R}$. Show that

$$m^*(A) = m^*(A \cap (0, \infty)) + m^*(A \setminus (0, \infty)).$$

Proof. Let $A = (\alpha, \beta)$, then

$$A \cap (0, \infty) = \begin{cases} \emptyset, & \text{if } \beta \leq 0; \\ (\alpha, \beta), & \text{if } \alpha, \beta > 0; \\ (0, \beta), & \text{if } \alpha \leq 0, \beta > 0, \end{cases}$$

and

$$A \setminus (0, \infty) = \begin{cases} (\alpha, \beta), & \text{if } \beta \leq 0; \\ \emptyset, & \text{if } \alpha, \beta > 0; \\ (\alpha, 0], & \text{if } \alpha \leq 0, \beta > 0. \end{cases}$$

Thus, in each case

$$m^*(A) = m^*(A \cap (0, \infty)) + m^*(A \setminus (0, \infty)).$$

■

Problem A.1.2. Let A be an open box in $\mathbb{R}^n$, and let $E = \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_n > 0\}$. Show that

$$m^*(A) = m^*(A \cap E) + m^*(A \setminus E).$$

Proof. Let $A = \prod_{i=1}^{\infty} (a_i, b_i)$, then

$$A \cap E = \left(\prod_{i=1}^{n-1} (a_i, b_i) \right) \times ((a_n \cap b_n) \cap (0, \infty)).$$

Now since $(a_n, b_n) \cap (0, \infty)$ is an interval,

$$m^*(A \cap E) = \text{vol}(A_{n-1}) \cdot m^*(A_n \cap (0, \infty)) \text{ where } A_{n-1} = \prod_{i=1}^{n-1} (a_i, b_i), A_n = (a_n, b_n).$$

Similarly,

$$A \setminus E = A_{n-1} \times (A_n \setminus (0, \infty)),$$

so

$$m^*(A \setminus E) = \text{vol}(A_{n-1}) \cdot m^*(A_n \setminus (0, \infty)).$$

Hence,

$$\begin{aligned} m^*(A) &= \text{vol}(A) = \text{vol}(A_{n-1})(m^*(A_n)) = \text{vol}(A_{n-1})(m^*(A_n \cap (0, \infty)) + m^*(A_n \setminus (0, \infty))) \\ &= \text{vol}(A_{n-1}) \cdot m^*(A_n \cap (0, \infty)) + \text{vol}(A_{n-1}) \cdot m^*(A_n \setminus (0, \infty)) \\ &= m^*(A \cap (0, \infty)) + m^*(A \setminus (0, \infty)). \end{aligned}$$

■

A.2 HW1

Problem A.2.1. Let

$$D = \{(x, y) \in \mathbb{R}^n, x \geq 0, 0 \leq y \leq x^3\},$$

and $x_0 = (0, 0)$. Define $f : D \rightarrow \mathbb{R}$ by $f(x, y) = 2x + \sqrt{y}$.

- (a) Prove that f is differentiable at $(0, 0)$ on D .
- (b) Describe the set of all derivative $L(x, y)$ for f on $(0, 0)$ on D .

A.3 HW2

Problem A.3.1. Define

$$\begin{aligned} \text{GL}(n, \mathbb{R}) &= \{A \text{ is a } n \times n \text{ matrix and } A \text{ is invertible}\} \\ &= \{A \mid \det(A) \neq 0\}, \end{aligned}$$

which is open in $\mathbb{R}^{n \times n}$. Consider the inverse map $g : \text{GL}(n, \mathbb{R}) \rightarrow \text{GL}(n, \mathbb{R})$, i.e. $g(A) = A^{-1}$. Show that $g'(A) : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$ is $g'(A)(H) = A^{-1}HA^{-1}$ for every $H \in \mathbb{R}^{n \times n}$.

Remark A.3.1. We can use the fact that if $M \in \mathbb{R}^{n \times n}$ has $\|M\| < 1$, then

$$(I + M)^{-1} = I - M + M^2 - M^3 + \dots$$

converges in $\mathbb{R}^{n \times n}$.

A.4 HW3

Problem A.4.1. Let $f : S \rightarrow S$ be a map between a complete metric space S . Assume there is a real sequence $\{\alpha_n\}$ which converges to 0 s.t.

$$d(f^n(x), f^n(y)) \leq \alpha_n d(x, y)$$

for all $n \geq 1$ and all $x, y \in S$. Prove that f has a unique fixed point. Hint. Apply the fixed-point theorem to f^m for a suitable m .

A.5 HW4

Problem A.5.1. Let $n \geq 2$, and let

$$f : \mathbb{R}^n \rightarrow \mathbb{R}$$

be a smooth function. Define

$$S := \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : f(x) = 0\}.$$

Assume that

$$\nabla f(x) \neq 0 \quad \text{for every } x \in S.$$

Then for every point $p \in S$, there exists an index $j \in \{1, \dots, n\}$, an open neighborhood $W \subset \mathbf{R}^n$ of p , an open set $U \subset \mathbf{R}^{n-1}$, and a differentiable function

$$g : U \rightarrow \mathbf{R}$$

such that, after possibly reordering the coordinates, the set $S \cap W$ can be written in the form

$$S \cap W = \left\{ (x_1, \dots, x_{j-1}, g(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_n), x_{j+1}, \dots, x_n) : (x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_n) \in U \right\}.$$

In other words, near every point of S , the hypersurface S is locally the graph of a differentiable function of $n - 1$ variables.

A.6 HW5

Problem A.6.1. Recall that a function $f : [a, b] \rightarrow \mathbf{R}$ is *Lipschitz* if there exists $L \geq 0$ such that $|f(x) - f(y)| \leq L|x - y|$ for all $x, y \in [a, b]$. Define the graph of f as:

$$\Gamma_f := \{(x, f(x)) : x \in [a, b]\} \subset \mathbf{R}^2.$$

Prove that $m_2^*(\Gamma_f) = 0$. *Hint: Partition $[a, b]$ into n sub-intervals and cover the corresponding pieces of the graph with rectangles. Consider the limit as $n \rightarrow \infty$.*

A.7 Random Stuff

Theorem A.7.1. If f is continuous and I is an open interval, then $f(I)$ is also an interval.

Theorem A.7.2. Any interval in $\mathbb{R}$ is measurable (open, closed, or half-open, half-closed)

Theorem A.7.3. A set E is Lebesgue measurable iff $m^*(A) = m^*(A \cap E) + m^*(A \setminus E)$ for all $A \subseteq \mathbb{R}^n$. To show E is measurable, it suffices to show

$$m^*(A) \geq m^*(A \cap E) + m^*(A \setminus E).$$

Appendix B

Additional proof

Theorem B.0.1. If $f = \liminf_{n \rightarrow \infty} f_n = \limsup_{n \rightarrow \infty} f_n$, then $f_n \rightarrow f$ pointwise.

Proof. Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on a set Ω , and suppose that for every $x \in \Omega$,

$$f(x) = \liminf_{n \rightarrow \infty} f_n(x) = \limsup_{n \rightarrow \infty} f_n(x).$$

We will show that $f_n(x) \rightarrow f(x)$ for every $x \in \Omega$.

Fix $x \in \Omega$. Then $\{f_n(x)\}$ is a sequence of real numbers satisfying

$$\liminf_{n \rightarrow \infty} f_n(x) = \limsup_{n \rightarrow \infty} f_n(x) = f(x).$$

We prove that $f_n(x) \rightarrow f(x)$.

Let $\varepsilon > 0$.

By the definition of $\limsup$, we have

$$\limsup_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \sup_{k \geq n} f_k(x) = f(x).$$

Hence there exists $N_1 \in \mathbb{N}$ such that for all $n \geq N_1$,

$$\sup_{k \geq n} f_k(x) < f(x) + \varepsilon.$$

In particular, for all $k \geq n \geq N_1$,

$$f_k(x) \leq \sup_{j \geq n} f_j(x) < f(x) + \varepsilon.$$

Thus for all $k \geq N_1$,

$$f_k(x) < f(x) + \varepsilon.$$

Similarly, by the definition of $\liminf$, we have

$$\liminf_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \inf_{k \geq n} f_k(x) = f(x).$$

Hence there exists $N_2 \in \mathbb{N}$ such that for all $n \geq N_2$,

$$\inf_{k \geq n} f_k(x) > f(x) - \varepsilon.$$

In particular, for all $k \geq n \geq N_2$,

$$f_k(x) \geq \inf_{j \geq n} f_j(x) > f(x) - \varepsilon.$$

Thus for all $k \geq N_2$,

$$f_k(x) > f(x) - \varepsilon.$$

Let $N = \max\{N_1, N_2\}$. Then for all $n \geq N$,

$$f(x) - \varepsilon < f_n(x) < f(x) + \varepsilon,$$

which implies

$$|f_n(x) - f(x)| < \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, we conclude that

$$f_n(x) \rightarrow f(x).$$

Since $x \in \Omega$ was arbitrary, the convergence is pointwise on Ω . ■

Theorem B.0.2. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a Lebesgue measurable function, and let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ be a function such that $f = g$ almost everywhere. Then g is also Lebesgue measurable.

Proof. Since $f = g$ almost everywhere, there exists a measurable set $N \subset \mathbb{R}^n$ such that $m(N) = 0$ and

$$f(x) = g(x) \quad \text{for all } x \in \mathbb{R}^n \setminus N.$$

To prove that g is measurable, it suffices to show that for every $a \in \mathbb{R}$, the set

$$E := \{x \in \mathbb{R}^n : g(x) > a\}$$

is measurable.

Fix $a \in \mathbb{R}$ and define

$$F := \{x \in \mathbb{R}^n : f(x) > a\}.$$

Since f is measurable, the set F is measurable.

We claim that

$$E \Delta F \subseteq N,$$

where $E \Delta F = (E \setminus F) \cup (F \setminus E)$ denotes the symmetric difference.

Indeed, let $x \notin N$. Then $f(x) = g(x)$, and hence

$$g(x) > a \Leftrightarrow f(x) > a.$$

Thus $x \in E$ if and only if $x \in F$, which implies $x \notin E \Delta F$. Therefore $E \Delta F \subseteq N$.

Since N has measure zero, it is measurable. It follows that $E \Delta F$ is measurable. As F is measurable and the collection of measurable sets is closed under symmetric difference, we conclude that

$$E = F \Delta (E \Delta F)$$

is measurable.

Since $a \in \mathbb{R}$ was arbitrary, this shows that $\{x : g(x) > a\}$ is measurable for all a , and hence g is measurable. ■