

PATIENT ID:



PATIENT NAME:



CHIANA MIKOVA

DATE OF BIRTH:



24/01/1977

SAMPLE CODE:



5245059654

QR-CODE:



02BJT0E9

ANALYZED ON:



03/05/2024

TESTED ALLERGENS:



296

TEST METHOD:



ALEX²

REFERRING PHYSICIAN:

ADDITIONAL INFORMATION:

The internal QC (Plausibility check for GD) was within acceptance range.

Lab report: Summary on detectable sensitisations

POLLEN

Grass Pollen



Tree Pollen



Weed Pollen



MITES

House Dust Mites & Storage Mites



PLANT-BASED FOOD

Legumes



Grains



Spices



Fruits



Vegetables



Nuts & Seeds



INSECTS & VENOMS

Ant, Bee, Wasp



Cockroach



MICROORGANISMS

Fungal Spores & Yeast



ANIMAL-DERIVED FOOD

Milk



Egg



Fish & Seafood



Meat



EPITHELIAL TISSUES OF ANIMALS

Pets



Farm Animals



OTHERS

Latex



Ficus




CCD



Parasite



Highest measured IgE concentration per allergen group



Name	E/M	Allergen	Function	kU _A /L
------	-----	----------	----------	--------------------

POLLEN

Grass Pollen

Bermuda grass	●●●●	Cyn d		0.10	
	●	Cyn d 1	Beta-Expansin	≤ 0.10	
Perennial Ryegrass	●	Lol p 1	Beta-Expansin	0.11	
Bahia grass	●●●●	Pas n		≤ 0.10	
Timothy grass	●	Phl p 1	Beta-Expansin	0.21	
	●	Phl p 2	Expansin	≤ 0.10	
	●	Phl p 5.0101	Grass Group 5/6	≤ 0.10	
	●	Phl p 6	Grass Group 5/6	≤ 0.10	
	●	Phl p 7	Polcalcin	≤ 0.10	
	●	Phl p 12	Profilin	≤ 0.10	
Common reed	●●●●	Phr c		≤ 0.10	
Cultivated rye, Pollen	●●●●	Sec c_pollen		≤ 0.10	

Tree Pollen

Acacia	●●●●	Aca m		0.13	
Tree of Heaven	●●●●	Ail a		≤ 0.10	
Alder	●	Aln g 1	PR-10	≤ 0.10	
	●	Aln g 4	Polcalcin	≤ 0.10	
Silver birch	●	Bet v 1	PR-10	≤ 0.10	
	●	Bet v 2	Profilin	≤ 0.10	
	●	Bet v 6	Isoflavon Reductase	≤ 0.10	
Paper mulberry	●●●●	Bro pa		≤ 0.10	
Hazel pollen	●●●●	Cor a_pollen		≤ 0.10	
	●	Cor a 1.0103	PR-10	0.13	
Sugi	●	Cry j 1	Pectate Lyase	≤ 0.10	
Cypress	●	Cup a 1	Pectate Lyase	≤ 0.10	
	●●●●	Cup s		≤ 0.10	
Beech	●	Fag s 1	PR-10	≤ 0.10	
Ash	●●●●	Fra e		≤ 0.10	
	●	Fra e 1	Ole e 1-Family	≤ 0.10	
Walnut pollen	●●●●	Jug r_pollen		0.13	
Mountain cedar	●●●●	Jun a		0.14	
Mulberry	●●●●	Mor r		0.11	
Olive	●	Ole e 1	Ole e 1-Family	≤ 0.10	

Name	E/M	Allergen	Function	kU _A /L
	⊙	Ole e 9	1,3 β Glucanase	≤ 0.10
Date palm	⊙	Pho d 2	Profilin	≤ 0.10
London plane tree	⊙	Pla a 1	Plant Invertase	≤ 0.10
	⊙	Pla a 2	Polygalacturonase	≤ 0.10
	⊙	Pla a 3	nsLTP	≤ 0.10
Cottonwood	⬢	Pop n		≤ 0.10
Elm	⬢	Ulm c		≤ 0.10

Weed Pollen

Common Pigweed	⬢	Ama r		≤ 0.10
Ragweed	⬢	Amb a		≤ 0.10
	⊙	Amb a 1	Pectate Lyase	≤ 0.10
	⊙	Amb a 4	Plant Defensin	≤ 0.10
Mugwort	⬢	Art v		≤ 0.10
	⊙	Art v 1	Plant Defensin	≤ 0.10
	⊙	Art v 3	nsLTP	≤ 0.10
Hemp	⬢	Can s		≤ 0.10
	⊙	Can s 3	nsLTP	≤ 0.10
Lamb's quarter	⬢	Che a		≤ 0.10
	⊙	Che a 1	Ole e 1-Family	≤ 0.10
Annual mercury	⊙	Mer a 1	Profilin	≤ 0.10
Wall pellitory	⬢	Par j		0.10
	⊙	Par j 2	nsLTP	≤ 0.10
Ribwort	⬢	Pla l		0.11
	⊙	Pla l 1	Ole e 1-Family	0.48
Russian thistle	⬢	Sal k		≤ 0.10
	⊙	Sal k 1	Pectin Methylesterase	≤ 0.10
Nettle	⬢	Urt d		≤ 0.10

MITES

House Dust Mite

American house dust mite	⊙	Der f 1	Cysteine protease	≤ 0.10
	⊙	Der f 2	NPC2 Family	≤ 0.10
European house dust mite	⊙	Der p 1	Cysteine protease	≤ 0.10
	⊙	Der p 2	NPC2 Family	≤ 0.10
	⊙	Der p 5	unknown	≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
	⊙	Der p 7	Mites, Group 7	0.15
	⊙	Der p 10	Tropomyosin	0.30
	⊙	Der p 11	Myosin, heavy chain	≤ 0.10
	⊙	Der p 20	Arginine kinase	≤ 0.10
	⊙	Der p 21	unknown	≤ 0.10
	⊙	Der p 23	Peritrophin-like protein domain	≤ 0.10

Storage Mite

Acarus siro	⦿	Aca s		≤ 0.10
Blomia tropicalis	⊙	Blo t 5	Mites, Group 5	≤ 0.10
	⊙	Blo t 10	Tropomyosin	≤ 0.10
	⊙	Blo t 21	unknown	≤ 0.10
Glycyphagus domesticus	⊙	Gly d 2	NPC2 Family	≤ 0.10
Lepidoglyphus destructor	⊙	Lep d 2	NPC2 Family	≤ 0.10
Tyrophagus putrescentiae	⦿	Tyr p		≤ 0.10
	⊙	Tyr p 2	NPC2 Family	≤ 0.10

MICROORGANISMS & SPORES

Yeast

Malassezia sympodialis	⊙	Mala s 5	unknown	≤ 0.10
	⊙	Mala s 6	Cyclophilin	≤ 0.10
	⊙	Mala s 11	Mn Superoxid-Dismutase	≤ 0.10
Yeast	⦿	Sac c		≤ 0.10

Moulds

Alternaria alternata	⊙	Alt a 1	Alt a 1-Family	0.12
	⊙	Alt a 6	Enolase	≤ 0.10
Aspergillus fumigatus	⊙	Asp f 1	Mitogillin Family	≤ 0.10
	⊙	Asp f 3	Peroxisomal Protein	0.12
	⊙	Asp f 4	unknown	≤ 0.10
	⊙	Asp f 6	Mn Superoxid-Dismutase	≤ 0.10
Cladosporium herbarum	⦿	Cla h		≤ 0.10
	⊙	Cla h 8	Short Chain Dehydrogenase	≤ 0.10
Penicillium chrysogenum	⦿	Pen ch		≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
------	-----	----------	----------	--------------------

PLANT FOOD

Legumes

Peanut	⊙	Ara h 1	7/8S Globulin	≤ 0.10
	⊙	Ara h 2	2S Albumin	0.24
	⊙	Ara h 3	11S Globulin	≤ 0.10
	⊙	Ara h 6	2S Albumin	≤ 0.10
	⊙	Ara h 8	PR-10	≤ 0.10
	⊙	Ara h 9	nsLTP	≤ 0.10
	⊙	Ara h 15	Oleosin	≤ 0.10
Chickpea	⬢	Cic a		≤ 0.10
Soy	⊙	Gly m 4	PR-10	≤ 0.10
	⊙	Gly m 5	7/8S Globulin	≤ 0.10
	⊙	Gly m 6	11S Globulin	≤ 0.10
	⊙	Gly m 8	2S Albumin	0.16
Lentil	⬢	Len c		≤ 0.10
White bean	⬢	Pha v		≤ 0.10
Pea	⬢	Pis s		≤ 0.10

Cereals

Oat	⬢	Ave s		≤ 0.10
Quinoa	⬢	Che q		≤ 0.10
Common buckwheat	⬢	Fag e		≤ 0.10
	⊙	Fag e 2	2S Albumin	≤ 0.10
Barley	⬢	Hor v		≤ 0.10
Lupine seed	⬢	Lup a		≤ 0.10
Rice	⬢	Ory s		0.24
Millet	⬢	Pan m		≤ 0.10
Cultivated rye	⬢	Sec c_flour		≤ 0.10
Wheat	⊙	Tri a aA_TI	Alpha-Amylase Trypsin-Inhibitor	≤ 0.10
	⊙	Tri a 14	nsLTP	≤ 0.10
	⊙	Tri a 19	Omega-5-Gliadin	≤ 0.10
Spelt	⬢	Tri s		≤ 0.10
Maize	⬢	Zea m		≤ 0.10
	⊙	Zea m 14	nsLTP	≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
------	-----	----------	----------	--------------------

Spices

Paprika	●●●●	Cap a		≤ 0.10
Caraway	●●●●	Car c		≤ 0.10
Oregano	●●●●	Ori v		≤ 0.10
Parsley	●●●●	Pet c		≤ 0.10
Anise	●●●●	Pim a		≤ 0.10
Mustard	●●●●	Sin		≤ 0.10
	●	Sin a 1	2S Albumin	≤ 0.10

Fruits

Kiwi	●	Act d 1	Cysteine protease	0.17
	●	Act d 2	TLP	0.22
	●	Act d 5	Kiwellin	≤ 0.10
	●	Act d 10	nsLTP	≤ 0.10
Papaya	●●●●	Car p		≤ 0.10
Orange	●●●●	Cit s		≤ 0.10
Melon	●	Cuc m 2	Profilin	≤ 0.10
Fig	●●●●	Fic c		≤ 0.10
Strawberry	●	Fra a 1+3	PR-10+LTP	≤ 0.10
Apple	●	Mal d 1	PR-10	0.10
	●	Mal d 2	TLP	≤ 0.10
	●	Mal d 3	nsLTP	≤ 0.10
Mango	●●●●	Man i		≤ 0.10
Banana	●●●●	Mus a		≤ 0.10
Avocado	●●●●	Pers a		≤ 0.10
Cherry	●●●●	Pru av		≤ 0.10
Peach	●	Pru p 3	nsLTP	≤ 0.10
Pear	●●●●	Pyr c		≤ 0.10
Blueberry	●●●●	Vac m		≤ 0.10
Grapes	●	Vit v 1	nsLTP	≤ 0.10

Vegetables

Onion	●●●●	All c		0.12
Garlic	●●●●	All s		≤ 0.10
Celery	●	Api g 1	PR-10	≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
	⊙	Api g 2	nsLTP	≤ 0.10
	⊙	Api g 6	nsLTP	≤ 0.10
Carrot	⬢	Dau c		≤ 0.10
	⊙	Dau c 1	PR-10	≤ 0.10
Potato	⬢	Sol t		≤ 0.10
Tomato	⬢	Sola l		≤ 0.10
	⊙	Sola l 6	nsLTP	≤ 0.10

Nuts

Cashew	⬢	Ana o		≤ 0.10
	⊙	Ana o 2	11S Globulin	≤ 0.10
	⊙	Ana o 3	2S Albumin	0.20
Brazil nut	⬢	Ber e		≤ 0.10
	⊙	Ber e 1	2S Albumin	≤ 0.10
Pecan	⬢	Car i		≤ 0.10
Hazelnut	⊙	Cor a 1.0401	PR-10	≤ 0.10
	⊙	Cor a 8	nsLTP	≤ 0.10
	⊙	Cor a 9	11S Globulin	≤ 0.10
	⊙	Cor a 11	7/8S Globulin	≤ 0.10
	⊙	Cor a 14	2S Albumin	≤ 0.10
Walnut	⊙	Jug r 1	2S Albumin	≤ 0.10
	⊙	Jug r 2	7/8S Globulin	0.12
	⊙	Jug r 3	nsLTP	≤ 0.10
	⊙	Jug r 4	11S Globulin	≤ 0.10
	⊙	Jug r 6	7/8S Globulin	≤ 0.10
Macadamia	⊙	Mac i 2S Albumin	2S Albumin	0.10
	⬢	Mac inte		≤ 0.10
Pistachio	⊙	Pis v 1	2S Albumin	0.25
	⊙	Pis v 2	11S Globulin subunit	≤ 0.10
	⊙	Pis v 3	7/8S Globulin	≤ 0.10
Almond	⬢	Pru du		≤ 0.10

Seed

Pumpkin seed	⬢	Cuc p		≤ 0.10
Sunflower seed	⬢	Hel a		≤ 0.10
Poppy seed	⬢	Pap s		≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
	⊙	Pap s 2S Albumin	2S Albumin	≤ 0.10
Sesame	⊙	Ses i		≤ 0.10
	⊙	Ses i 1	2S Albumin	≤ 0.10
Fenugreek seeds	⊙	Tri fo		≤ 0.10

ANIMAL FOOD

Milk

Cow, milk	⊙	Bos d_milk		≤ 0.10
	⊙	Bos d 4	α-Lactalbumin	≤ 0.10
	⊙	Bos d 5	β-Lactoglobulin	≤ 0.10
	⊙	Bos d 8	Casein	≤ 0.10
Camel	⊙	Cam d		≤ 0.10
Goat, milk	⊙	Cap h_milk		≤ 0.10
Mare's milk	⊙	Equ c_milk		0.15
Sheep, milk	⊙	Ovi a_milk		≤ 0.10

Egg

Egg white	⊙	Gal d_white		0.10
Egg yolk	⊙	Gal d_yolk		≤ 0.10
Egg white	⊙	Gal d 1	Ovomucoid	≤ 0.10
	⊙	Gal d 2	Ovalbumin	≤ 0.10
	⊙	Gal d 3	Ovotransferrin	≤ 0.10
	⊙	Gal d 4	Lysozym C	≤ 0.10
Egg yolk	⊙	Gal d 5	Serum Albumin	0.23

Seafood

Herring worm	⊙	Ani s 1	Kunitz Serin Protease Inhibitor	≤ 0.10
	⊙	Ani s 3	Tropomyosin	≤ 0.10
Crab	⊙	Chi spp.		≤ 0.10
Herring	⊙	Clu h		≤ 0.10
	⊙	Clu h 1	β-Parvalbumin	≤ 0.10
Brown shrimp	⊙	Cra c 6	Troponin C	≤ 0.10
Carp	⊙	Cyp c 1	β-Parvalbumin	≤ 0.10
Atlantic cod	⊙	Gad m		≤ 0.10
	⊙	Gad m 2+3	β-Enolase & Aldolase	≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
	⊙	Gad m 1	β-Parvalbumin	≤ 0.10
Lobster	⬢	Hom g		≤ 0.10
Shrimp	⬢	Lit s		≤ 0.10
Squid	⬢	Lol spp.		≤ 0.10
Common mussel	⬢	Myt e		≤ 0.10
Oyster	⬢	Ost e		≤ 0.10
Shrimp	⬢	Pan b		≤ 0.10
Scallop	⬢	Pec spp.		≤ 0.10
Black Tiger Shrimp	⊙	Pen m 1	Tropomyosin	≤ 0.10
	⊙	Pen m 2	Arginine kinase	≤ 0.10
	⊙	Pen m 3	Myosin, light chain	≤ 0.10
	⊙	Pen m 4	Sarcoplasmic Calcium Binding Protein	≤ 0.10
Thornback ray	⬢	Raj c		≤ 0.10
	⊙	Raj c Parvalbumin	α-Parvalbumin	≤ 0.10
Clam	⬢	Rud spp.		≤ 0.10
Salmon	⬢	Sal s		≤ 0.10
	⊙	Sal s 1	β-Parvalbumin	≤ 0.10
Atlantic mackerel	⬢	Sco s		≤ 0.10
	⊙	Sco s 1	β-Parvalbumin	≤ 0.10
Tuna	⬢	Thu a		≤ 0.10
	⊙	Thu a 1	β-Parvalbumin	≤ 0.10
Swordfish	⊙	Xip g 1	β-Parvalbumin	≤ 0.10

Meat

House cricket	⬢	Ach d		≤ 0.10
Cattle, meat	⬢	Bos d_meat		≤ 0.10
	⊙	Bos d 6	Serum Albumin	≤ 0.10
Horse, meat	⬢	Equ c_meat		≤ 0.10
Chicken meat	⬢	Gal d_meat		0.14
Migratory locust	⬢	Loc m		≤ 0.10
Turkey	⬢	Mel g		≤ 0.10
Rabbit, meat	⬢	Ory_meat		0.14
Sheep, meat	⬢	Ovi a_meat		≤ 0.10
Pork	⬢	Sus d_meat		≤ 0.10
	⊙	Sus d 1	Serum Albumin	≤ 0.10
Mealworm	⬢	Ten m		≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
------	-----	----------	----------	--------------------

INSECTS & VENOMS

Fire ant poison

Fire ant	•••••	Sol spp.		≤ 0.10
----------	-------	----------	--	--------

Honey Bee Venom

Honey bee	•••••	Api m		0.17
	•	Api m 1	Phospholipase A2	≤ 0.10
	•	Api m 10	Icarapin Variant 2	0.18

Wasp Venom

Hornet	•••••	Dol spp		≤ 0.10
Paper wasp venom	•••••	Pol d		≤ 0.10
	•	Pol d 5	Antigen 5	≤ 0.10
Wasp venom	•••••	Ves v		≤ 0.10
	•	Ves v 1	Phospholipase A1	≤ 0.10
	•	Ves v 5	Antigen 5	≤ 0.10

Cockroach

German Cockroach	•	Bla g 1	Cockroach Group 1	≤ 0.10
	•	Bla g 2	Aspartyl protease	≤ 0.10
	•	Bla g 4	Lipocalin	≤ 0.10
	•	Bla g 5	Glutathione S-transferase	≤ 0.10
	•	Bla g 9	Arginine kinase	≤ 0.10
American Cockroach	•••••	Per a		≤ 0.10
	•	Per a 7	Tropomyosin	≤ 0.10

ANIMAL ORIGIN

Pet

Dog	•	Can f_Fd1	Uteroglobulin	≤ 0.10
Male dog urine (incl. Can f 5)	•••••	Can f_male urine		≤ 0.10
Dog	•	Can f 1	Lipocalin	≤ 0.10
	•	Can f 2	Lipocalin	≤ 0.10
	•	Can f 3	Serum Albumin	≤ 0.10

Name	E/M	Allergen	Function	kU _A /L
	⊙	Can f 4	Lipocalin	≤ 0.10
	⊙	Can f 6	Lipocalin	≤ 0.10
Guinea pig	⊙	Cav p 1	Lipocalin	≤ 0.10
Cat	⊙	Fel d 1	Uteroglobin	≤ 0.10
	⊙	Fel d 2	Serum Albumin	≤ 0.10
	⊙	Fel d 4	Lipocalin	≤ 0.10
	⊙	Fel d 7	Lipocalin	≤ 0.10
House mouse	⊙	Mus m 1	Lipocalin	≤ 0.10
Rabbit, epithel	⊙	Ory c 1	Lipocalin	≤ 0.10
	⊙	Ory c 2	Lipophilin	≤ 0.10
	⊙	Ory c 3	Uteroglobin	≤ 0.10
Djungarian hamster	⊙	Phod s 1	Lipocalin	≤ 0.10
Rat	⊙	Rat n		≤ 0.10

Farm Animals

Cattle	⊙	Bos d 2	Lipocalin	≤ 0.10
Goat, epithel	⊙	Cap h_epithelia		≤ 0.10
Horse, epithel	⊙	Equ c 1	Lipocalin	≤ 0.10
	⊙	Equ c 3	Serum Albumin	≤ 0.10
	⊙	Equ c 4	Latherin	0.54
Sheep, epithel	⊙	Ovi a_epithelia		≤ 0.10
Pig	⊙	Sus d_epithelia		≤ 0.10

OTHERS

Latex

Latex	⊙	Hev b 1	Rubber elongation factor	≤ 0.10
	⊙	Hev b 3	Small rubber particle protein	≤ 0.10
	⊙	Hev b 5	unknown	0.20
	⊙	Hev b 6.02	Hevein	≤ 0.10
	⊙	Hev b 8	Profilin	≤ 0.10
	⊙	Hev b 11	Class 1 Chitinase	0.10

Ficus

Weeping fig	⊙	Fic b		≤ 0.10
-------------	---	-------	--	--------

Name	E/M	Allergen	Function	kU _A /L
------	-----	----------	----------	--------------------

CCD

Hom s Lactoferrin	●	Hom s LF	CCD	≤ 0.10
-------------------	---	----------	-----	--------

Parasite

Pigeon tick	●	Arg r 1	Lipocalin	≤ 0.10
-------------	---	---------	-----------	--------

Total IgE result: 81 kU/L

Reference range total-IgE
Adults: < 100 kU/L

SAMPLED ON
02/05/2024

PRINTED ON
03/05/2024

Information to cross-reactive allergens

Ole e 1 Family

Ole e 1 family allergens show a high degree of cross-reactivity within plant family borders.

The major olive pollen allergen, Ole e 1, represents the prototype of all Ole e 1 family allergens and is the primary sensitiser in regions with olive pollen exposure. Further Ole e 1 related allergens were identified in other trees from the Oleaceae family (ash, lilac, privet). There are further Ole e 1 family members present in grass- and weed pollen.

Tropomyosins

Tropomyosins show a very high degree of cross-reactivity.

Tropomyosins can induce diverse symptoms including anaphylaxis. Sensitisation to tropomyosins can occur by ingestion (seafood), inhalation (mites, cockroaches) or parasite infection (ascariasis, anisakiasis). Tropomyosins are stable to processing.

ALEX² – Number of tested allergen sources: 296



GRASS POLLEN 6
Bahia grass, Bermuda grass, Common reed, Perennial ryegrass, Rye, Timothy grass



COCKROACH 2
American cockroach, German cockroach



TREE POLLEN 19
Acacia, Alder, Arizona Cypress, European Ash, Beech, Cottonwood, Date palm, Elm, Hazel, London Plane Tree, Mediterranean Cypress, Mountain cedar, Mulberry, Olive, Paper mulberry, Silver birch, Sugi, Tree of Heaven, Walnut



INSECT VENOMS 5
Common wasp venom, Fire ant venom, Honeybee venom, Long-headed wasp venom, Paper wasp venom



WEED POLLEN 10
Annual mercury, Hemp, Lamb's quarter, Mugwort, Nettle, Pigweed, Ragweed, Ribwort, Russian thistle, Wall pellitory



FUNGAL SPORES & YEAST 6
Alternaria alternata, Aspergillus fumigatus, Baker's yeast, Cladosporium herbarum, Malassezia sympodialis, Penicillium chrysogenum



HOUSE DUST MITES & STORAGE MITES 7
Acarus siro, American house dust mite, Blomia tropicalis, European house dust mite, Glycyphagus domesticus, Lepidoglyphus destructor, Tyrophagus putrescentiae



EGG 2
Egg white, Egg yolk



LEGUMES 6
Chickpea, White bean, Lentil, Pea, Peanut, Soy



FISH & SEAFOOD 20
Anisakis simplex, Atlantic cod, Atlantic herring, Atlantic mackerel, Black-Tiger shrimp, Brown shrimp, Carp, Common mussel, Crab, Lobster, Northern prawn, Oyster, Salmon, Scallop, Shrimp mix, Squid, Swordfish, Thornback ray, Tuna, Venus clam



GRAINS 11
Barley, Buckwheat, Corn, Cultivated rye, Lupine, Millet, Oat, Quinoa, Rice, Spelt, Wheat



MEAT 10
Beef, Chicken, Horse, House cricket, Lamb, Mealworm, Migratory locust, Pig, Rabbit, Turkey



SPICES 6
Anise, Caraway, Mustard, Oregano, Paprika, Parsley



PETS 7
Cat, Djungarian hamster, Dog, Guinea pig, Mouse, Rabbit, Rat



FRUITS 15
Avocado, Apple, Banana, Blueberry, Cherry, Fig, Grape, Kiwi, Mango, Muskmelon, Orange, Papaya, Peach, Pear, Strawberry



FARM ANIMALS 5
Cattle, Goat, Horse, Pig, Sheep



VEGETABLES 6
Carrot, Celery, Garlic, Onion, Potato, Tomato



OTHERS 4
Latex, Hom s lactoferrin, Pigeon tick, Weeping fig



NUTS & SEEDS 13
Almond, Brazil nut, Cashew, Hazelnut, Macadamia, Pecan, Pistachio, Walnut, Fenugreek seeds, Poppy seed, Pumpkin seed, Sesame, Sunflower seed

Raven Interpretation Summary

Sample Information

The sample was tested on ALEX² Barcode 02BJT0E9, interpretation date 03/05/2024.

Of the tested 295 allergens, 3 were/was above the cut off of 0.3 kU_A/L. A sensitisation can be an indicator of an IgE dependent allergy. For all positive ALEX² allergens, comments for interpretation guidance are listed below.

Total IgE: 81 kU/L

The measured total IgE was 81 kU/L. With a total IgE titre of below 100 kU/L, allergy is possible but unlikely.

Cross-Reactive allergen sensitisation detected

Sensitisations against molecular allergens which are markers of (broad) cross-reactivity between different allergen sources were detected.

Detected cross-reactive allergen sensitisations:

- Tropomyosins: Der p 10

Tropomyosins

Members of the Tropomyosin allergen family can cause inhalative, as well as mild to severe reactions after consumption of seafood. Allergens of the TM allergen family have been identified in ,fish-parasites, insects (e.g. cockroach), mites and seafood. The degree of cross-reactivity between TM members is high.

Weed Pollen

Ribwort

Sensitisation to pollen from ribwort was detected. Allergic symptoms associated with this allergen source range from allergic rhinoconjunctivitis to allergic asthma.

Pla I 1 is a member of the Ole e 1 allergen family. The degree of cross-reactivity to Ole e 1 allergens from other plantain species is high, that to Ole e 1-like allergens from other plant families is low. Pla I 1 serves as a marker for AIT indication, if corresponding clinical symptoms are present.

Causal treatment is possible via AIT. Symptomatic treatment includes anti-histamines and local corticosteroids in various formulations (tablet, spray).

Furry Animals

Horse

Sensitisation to horse was detected. Allergic symptoms associated with this allergen source range from allergic rhino-conjunctivitis to allergic asthma.

Equ c 4 is a member of the Latherin allergen family. Significantly higher levels are found in stallions, compared to mares and geldings. So far only one other member of this allergen family has been described (from cat) - the degree of cross-reactivity can be considered low.

If avoidance of horses is not possible an AIT can be prescribed - Equ c 1 serves a marker for AIT indication, if corresponding clinical symptoms are present. Symptomatic treatment includes anti-histamines as well as local corticosteroids in various formulations (tablet, spray). Avoidance is strongly recommended.

Mites and Cockroaches

House dust mites

Sensitisation to house dust mite was detected. Allergic symptoms associated with this allergen source range from allergic rhinoconjunctivitis to asthma.

Der p 10 is a member of the Tropomyosin allergen family. The degree of cross-reactivity between Der p 10 and other Tropomyosins is high. Sensitisation to Der p 10 can be the cause for cross-reactions to shrimp and other seafood species (except fish).

Allergen avoidance is advised. Encasings for blankets, mattresses and pillows can reduce the allergen load. Der f 1/Der p 1 and Der f 2/Der p 2 are major allergens from house dust mite and serve as markers for AIT indication, if corresponding clinical symptoms are present. Symptomatic treatment includes anti-histamines as well as local corticosteroids in various formulations (tablet, spray).

DISCLAIMER: THE PRESENCE OF IgE-ANTIBODIES IMPLIES A RISK OF ALLERGIC REACTIONS AND HAS TO BE ANALYZED IN CONJUNCTION WITH THE CLINICAL HISTORY AND OTHER DIAGNOSTIC TEST RESULTS. THE RAVEN INTERPRETATION GUIDANCE SOFTWARE IS A TOOL TO SUPPORT PHYSICIANS IN THE INTERPRETATION OF ALEX² RESULTS. RAVEN COMMENTS DO NOT REPLACE THE DIAGNOSIS BY A PHYSICIAN. NO LIABILITY IS ACCEPTED FOR RAVEN COMMENTS AND RESULTING THERAPEUTIC INTERVENTIONS. THE STATED COMMENTS ARE DESIGNED EXCLUSIVELY FOR ALEX² RESULTS.

