

Finite-size effects in quantum thermodynamics

Kamil Korzekwa

*Faculty of Physics, Astronomy and Applied Computer Science,
Jagiellonian University, Poland*



TEAM-NET

Outline

1. Motivation
2. Thermodynamic setting
3. Asymptotic reversibility
4. Finite-size irreversibility
5. Resource resonance effect
6. Fluctuation-dissipation relations
7. Outlook

Quantum 2, 108 (2018)
Phys. Rev. A 99, 032332 (2019)
Phys. Rev. Lett. 122, 110403 (2019)
Phys. Rev. E 105, 054127 (2022)



Marco Tomamichel
National University of Singapore



Christopher Chubb
ETH Zurich



Tanmoy Biswas
University of Gdańsk



Alexssandre de Oliveira Junior
Jagiellonian University




Michał Horodecki
University of Gdańsk

Motivation


What can we say about the dynamics without solving equations of motion?

Closed systems



Energy conservation

Open systems




Entropy growth

Quantum thermodynamics:

Using minimal assumptions of the quantum theory, find constraints on the evolution of a quantum system interacting with thermal baths


Motivation

Open dynamics approach:



Exact time evolution
for a given model

Resource-theoretic approach:




Allowed final states
compatible with the laws
of thermodynamics

Thermodynamic setting

Thermodynamic transformations modelled by **thermal operations***:

$$\mathcal{E}^T(\cdot) = \text{Tr}_B \left(U (\cdot \otimes \gamma_B) U^\dagger \right) \quad \text{with} \quad [U, H + H_B] = 0$$



Gibbs state γ of the system at temperature T : $\gamma = e^{-\frac{H}{T}} / \mathcal{Z}, \quad \mathcal{Z} = \text{Tr} \left(e^{-\frac{H}{T}} \right)$

Note: all results with units such that $k_B = 1$.

*M. Horodecki, J. Oppenheim
Nature Commun. 4, 2059 (2013)

Thermodynamic setting

Setting: Initial state ρ , target state σ , background temperature T

Single-shot interconversion: Does there exist \mathcal{E}^T such that $\mathcal{E}^T(\rho) = \sigma$?

Many-copies interconversion: Does there exist \mathcal{E}^T such that $\mathcal{E}^T(\rho^{\otimes n}) \approx_{\epsilon} \sigma^{\otimes R_n n}$?

Optimal rate R_n for error ϵ ?

Note: $\sigma \approx_{\epsilon} \tilde{\sigma}$ means $1 - F(\sigma, \tilde{\sigma}) \leq \epsilon$ with fidelity F

Restrictions:


Focus on many copies (large but finite n) and *energy-incoherent* states:

$[\rho, H] = [\sigma, H] = 0 \implies$ states represented by: $\mathbf{p} = \text{eig}(\rho)$, $\mathbf{q} = \text{eig}(\sigma)$.

$[\gamma, H] = 0 \implies$ thermal state represented by: $\gamma = \text{eig}(\gamma)$

Asymptotic reversibility

Asymptotic rate for $n \rightarrow \infty^*$: $R_\infty(\mathbf{p} \rightarrow \mathbf{q}) = \frac{D(\mathbf{p} \parallel \boldsymbol{\gamma})}{D(\mathbf{q} \parallel \boldsymbol{\gamma})}$



Relative entropy: $D(\mathbf{p} \parallel \boldsymbol{\gamma}) := \sum_{i=1}^d p_i \log \frac{p_i}{\gamma_i}$

Physical interpretation:

$$\frac{1}{T} \left[\underbrace{\langle E \rangle_{\mathbf{p}} - TH(\mathbf{p})}_{\text{Free energy } F = U - TS} - \underbrace{(-T \log \mathcal{Z})}_{\text{Free energy of } \boldsymbol{\gamma}} \right]$$

No dissipation of free energy in the thermodynamic limit!

*F. Brandão *et al.*,
Phys. Rev. Lett. 111, 250404 (2013)

Finite-size irreversibility

Rate for large but finite n :

$$R_n = R_\infty - f(\mathbf{p}, \mathbf{q}, \boldsymbol{\gamma}, n, \epsilon)$$


Relevant quantity quantifying irreversibility:

Relative
entropy
variance:

$$V(\mathbf{p} \parallel \boldsymbol{\gamma}) := \sum_{i=1}^d p_i \left(\log \frac{p_i}{\gamma_i} - D(\mathbf{p} \parallel \boldsymbol{\gamma}) \right)^2$$

Physical
interpretation:

$$V(\boldsymbol{\gamma}' \parallel \boldsymbol{\gamma}) = \underbrace{\frac{\partial \langle E \rangle_{\boldsymbol{\gamma}'}}{\partial T'}}_{\text{Specific heat capacity}} \cdot \underbrace{\left(1 - \frac{T'}{T} \right)^2}_{\text{Carnot factor}}$$



Finite-size irreversibility


Optimal conversion rate R_n with constant error ϵ :

Irreversibility parameter:

$$R_n(\epsilon) \simeq R_\infty + \sqrt{\frac{V(\mathbf{p} \parallel \boldsymbol{\gamma})}{D(\mathbf{q} \parallel \boldsymbol{\gamma})^2}} \frac{Z_\nu^{-1}(\epsilon)}{\sqrt{n}}$$

$$\nu = \frac{V(\mathbf{q} \parallel \boldsymbol{\gamma})/D(\mathbf{q} \parallel \boldsymbol{\gamma})}{V(\mathbf{p} \parallel \boldsymbol{\gamma})/D(\mathbf{p} \parallel \boldsymbol{\gamma})}$$

Rayleigh-normal distribution Z_ν^* :




Z_0 - standard normal distribution Φ

Z_1 - Rayleigh distribution ($Z_1(x) = 0$ for $x \leq 0$)

*W. Kumagai *et al.*, IEEE Trans. Inf. Theory **63**, 1829–1857 (2017)

Finite-size irreversibility


Numerical verification of the obtained second-order asymptotic expression for optimal rate:




Finite-size irreversibility

Effects of finite-size irreversibility on work distillation and dilution processes:


Work distillation process:



Work dilution process:



Finite-size irreversibility




Dissipation of free energy beyond the thermodynamic limit!

Resource resonance

Optimal conversion rate R_n with vanishing error $\epsilon = e^{-n^\alpha}$ and $\alpha \in (0, 1)$:

$$R_n(\epsilon) \simeq R_\infty - \sqrt{\frac{V(\textcolor{blue}{p} \parallel \textcolor{red}{\gamma})}{D(\textcolor{green}{q} \parallel \textcolor{red}{\gamma})^2}} \frac{|\sqrt{1/\nu} - 1|}{\sqrt{n^{1-\alpha}}}$$

When $\nu = 1$ correction term disappears for every error ϵ




No free energy dissipation!
(at least up to second order asymptotics)

Recall that $\nu = 1$ means that the relative fluctuations of free energy are the same for the initial state ρ and target state σ


Resource resonance

Resonance example: Heat engine with a finite-size working body:



Resource resonance

Working body: $n = 200$ qubits, energy gap E
Background (hot) bath: $T_h = 10E$



Fluctuation-dissipation relations

Einstein-Smoluchowski relation for a Brownian particle:




Thermodynamic distillation process

Non-zero free energy:

$$F^N := \frac{1}{\beta} \sum_{n=1}^N D(\rho_n^N \| \gamma_n^N)$$

Non-zero free energy fluctuations:

$$\sigma^2(F^N) := \frac{1}{\beta^2} \sum_{n=1}^N V(\rho_n^N \| \gamma_n^N)$$



Non-zero free energy, but
vanishing free energy fluctuations

Free energy
fluctuations

?

Free energy
dissipated in
the process

Fluctuation-dissipation relations

Optimal error in thermodynamic distillation process:

$$\lim_{N \rightarrow \infty} \epsilon_N = \lim_{N \rightarrow \infty} \Phi \left(-\frac{\Delta F^N}{\sigma(F^N)} \right)$$

ΔF^N - Free energy difference between initial and **target** state

Minimal amount of free energy dissipated in the optimal distillation process:

$$F_{\text{diss}}^N \simeq a(\epsilon_N) \sigma(F^N)$$

F_{diss}^N - Free energy difference between initial and **final** state

$$a(\epsilon) = -\Phi^{-1}(\epsilon)(1 - \epsilon) + \exp(-[\Phi^{-1}(\epsilon)]^2/2)/\sqrt{2\pi}$$

Three regimes:

$$\lim_{N \rightarrow \infty} \frac{\Delta F^N}{\sqrt{N}} = \begin{cases} \infty, \\ -\infty, \\ \alpha \in \mathbb{R} \end{cases} \begin{matrix} \Rightarrow \\ \Rightarrow \\ \Rightarrow \end{matrix} \begin{cases} \epsilon = 0, \\ \epsilon = 1, \\ \end{cases} \begin{matrix} F_{\text{diss}}^N = \Delta F^N \\ F_{\text{diss}}^N = 0 \end{matrix}$$

Also holds for initial pure states with coherence!

Outlook

- Extend finite-size analysis to other resource-theories (asymmetry, contextuality).
- Design experimental protocols employing the resonance phenomenon.
- Generalise the formalism to include quantum states with coherence.
- Look for resonance phenomena in other quantum information processing tasks.
- Extend resource-theoretic fluctuation-dissipation theorem to continuous variable systems

Quantum 2, 108 (2018)
Phys. Rev. A 99, 032332 (2019)
Phys. Rev. Lett. 122, 110403 (2019)
Phys. Rev. E 105, 054127 (2022)

Thank you!