



Wrocław University of Technology

Decoherence-driven mechanism for initialization of hole spins in a p-doped semiconductor quantum well

K. Korzekwa, P. Machnikowski

Institute of Physics, Wrocław University of Technology

M. Kugler, C. Gradl, S. Furthmeier, M. Griesbeck, M. Hirmer, D. Schuh,
C. Schüller, T. Korn

Institut für Experimentelle und Angewandte Physik, Universität Regensburg

T. Kuhn

Institut für Festkörpertheorie Westfälische Wilhelms-Universität






Contents


1. Motivation
2. Modeled experiment
3. Model of hole-trion spin dynamics
4. Kerr signal for non-resonant excitation
5. Decoherence- and precession-driven mechanisms for initialization of resident hole spin polarization
6. Competition of mechanisms in resonant spin amplification experiment
7. Conclusions

Motivation

- Spin dynamics and coherence is crucial for many applications in emerging technologies (spintronics)
- Time-resolved Kerr rotation is a very efficient method for the investigation of spin dynamics; interesting experiments have been performed quite recently
- Resonant spin amplification signal shows interesting behaviour for non-resonant excitation, including the formation of a zero-field peak
- Extended life times are observed in semiconductor nanostructures, especially for hole spin states (for which reduced hyperfine interaction makes the spin decoherence slower)




Experiment



Sample:

Single-side p-modulation-doped GaAs/Al_{0.3}Ga_{0.7}As quantum well,
width: 4 nm, hole density: $1.1 \times 10^{11} \text{ cm}^{-2}$, mobility: $1.3 \times 10^4 \text{ cm}^2/\text{Vs}$


Model




Selection rules for σ^+ pump pulse

$$|\downarrow\rangle \longleftrightarrow |T\downarrow\rangle \quad |\uparrow\rangle \nleftrightarrow |T\downarrow\rangle$$

Hole states




$|\uparrow\rangle$




$|\downarrow\rangle$

Trion states




$|T\uparrow\rangle$



$|T\downarrow\rangle$

$$\rho(t) = \begin{pmatrix} \frac{1-N_t(t)+\Sigma_h(t)}{2} & \frac{1}{2}X_h(t) - \frac{i}{2}Y_h(t) & 0 & 0 \\ \frac{1}{2}X_h(t) + \frac{i}{2}Y_h(t) & \frac{1-N_t(t)-\Sigma_h(t)}{2} & 0 & 0 \\ 0 & 0 & \frac{N_t(t)+\Sigma_t(t)}{2} & \frac{1}{2}X_t(t) - \frac{i}{2}Y_t(t) \\ 0 & 0 & \frac{1}{2}X_t(t) + \frac{i}{2}Y_t(t) & \frac{N_t(t)-\Sigma_t(t)}{2} \end{pmatrix}$$



- 1.** The electric field couples to the interband transitions via a dipole moment:

$$H_{\text{p}} = \frac{1}{2}f(t)e^{i\Delta t - i\psi}|\uparrow\rangle\langle T\uparrow| + \text{H.c.}$$

$$f(t) = -\mathbf{d} \cdot \boldsymbol{\mathcal{E}}_0^* (1 + r) \eta(t/\tau_p)$$

- 2.** The system evolution (Larmor precession, recombination and spin decoherence) is modeled in terms of the Markovian Master equation:

$$\dot{\rho} = -\frac{i}{\hbar}[H_0, \rho] + \mathcal{L}_h[\rho] + \mathcal{L}_t[\rho] + \mathcal{L}_r[\rho]$$

$$H_0 = -\frac{1}{2}\mu_{\text{B}}\mathbf{B}\hat{g}_{\text{h}}\boldsymbol{\sigma}_{\text{h}} - \frac{1}{2}g_{\text{t}}\mu_{\text{B}}\mathbf{B} \cdot \boldsymbol{\sigma}_{\text{t}}$$


- 3.** The optical response, that is, the rotation of the polarization plane of the reflected or transmitted probe pulse, is proportional to*:

$$\text{TRKR} = \Sigma_t - \Sigma_h$$

*P. Machnikowski and T. Kuhn, Phys. Rev. B **81**, 115306 (2010)
I. A. Yugova *et al.* Phys. Rev. B **80**, 104436 (2009)


Kerr signal for non-resonant excitation

$T = 15 \text{ K}$ $B = 0 \text{ T}$



Fast initial hole spin decoherence

Additional decoherence mechanism due to the excess energy delivered to the system




Described by occupation relaxation parameter u and pure dephasing parameter w :

$$\begin{aligned}
 \langle \downarrow | \rho_2 | \uparrow \rangle &= \langle \downarrow | \rho_1 | \uparrow \rangle e^{-u/2-w}, \\
 \langle \uparrow | \rho_2 | \downarrow \rangle &= \langle \uparrow | \rho_1 | \downarrow \rangle e^{-u/2-w}, \\
 \langle \uparrow | \rho_2 | \uparrow \rangle &= \frac{1}{2} \langle \uparrow | \rho_1 | \uparrow \rangle (1 + e^{-u}) + \frac{1}{2} \langle \downarrow | \rho_1 | \downarrow \rangle (1 - e^{-u}), \\
 \langle \downarrow | \rho_2 | \downarrow \rangle &= \frac{1}{2} \langle \downarrow | \rho_1 | \downarrow \rangle (1 - e^{-u}) + \frac{1}{2} \langle \uparrow | \rho_1 | \uparrow \rangle (1 + e^{-u}).
 \end{aligned}$$



Kerr signal for non-resonant excitation


T = 15 K B = 0 T



$$\text{TRKR} = ae^{-\gamma_1 t} - be^{-\kappa t}$$

$$a = \left(\frac{\gamma_1}{\gamma_1 - \kappa} + 1 \right) \Sigma_t^{(0)}$$

$$b = \left(\frac{\gamma_1}{\gamma_1 - \kappa} - e^{-u} \right) \Sigma_t^{(0)}$$




Hole spin
decoherence rate

Recombination rate


Hole spin coherence time $T_2 = 10$ ns





Decoherence-driven mechanism




Precession-driven mechanism



Resonant spin amplification



RSA signal obtained by finding the stationary point of the spin polarization transformation corresponding to one repetition of the pulsed laser:




Response from an **inhomogeneous ensemble** of hole spins obtained by averaging the result according to a **Gaussian distribution** of hole g-factors

**Analytical expression
for RSA signal**

$$\text{RSA} \sim f \frac{P}{Q}$$

$$f = 1 - e^{-u} - \frac{\omega_t^2}{\gamma_R^2 + \omega_t^2}$$


RSA signal: competition of mechanisms




$$f = 1 - e^{-u} - \frac{\omega_t^2}{\gamma_R^2 + \omega_t^2}$$

RSA for non-resonant excitation

T=1.2 K



Intrinsic hole spin coherence time
 T_2 reaching **100 ns**





Conclusions

- The negative Kerr signal observed in the experiment for nonresonant excitation is linked to the ratio of the hole-to-trion polarization after the fast initial decay
- Detuning does not directly contribute to the RSA signal. Its role in the appearance of the zero-field peak may be to provide excess energy that increases the initial dephasing
- It is possible to control the sign and magnitude of long-living spin polarization through the interplay of decoherence- and precession-induced mechanisms
- Hole spin coherence time in zero magnetic field (intrinsic) was experimentally observed to reach 100 ns

[Phys. Rev. B **84**, 085327 (2011)]

This work was supported by the TEAM programme of the Foundation for Polish Science co-financed from the European Regional Development Fund

