

Magnetic propeties of a two dimentional electron gas strongly coupled to lights

Kosala Herath

May 14, 2021

v.2.0.0.beta

1 Current Operator in Landau Levels

Now consider about the current density operator for N th Landau level. Since we have already found the extact solution for our time depenedent Hamiltonian and we have identify them as Floquet states with quasi energies. From these solutions we can identify the *Floquet modes* as given in Eq. (??) and using quantum numbers we can represent those states as follows

$$|\phi_\alpha\rangle = |\phi_{n,m}\rangle \equiv |n, k_x\rangle \quad \text{where} \quad k_x = m \frac{2\pi}{L_x} \quad (1.1)$$

Using above complete set of eigenstates of Floquet Hamiltonian we can represent the single particle current operator's matrix element as

$$(\mathbf{j})_{nm,n'm'} = \langle n, k_x | \hat{\mathbf{j}} | n', k'_x \rangle \quad (1.2)$$

where particle current operator for this system will be

$$\hat{\mathbf{j}} = \frac{1}{m} \left(\hat{\mathbf{P}} - e[\mathbf{A}_s + \mathbf{A}_d(t)] \right). \quad (1.3)$$

However, we are only consider the transverse conductivity in x direction we can identify that x directional current operator as

$$\hat{j}_x = \frac{1}{m} \left(-i\hbar \frac{\partial}{\partial x} + eBy \right). \quad (1.4)$$

Now we can calculate the matrix elements of x directional current operator's matrix in Floquet mode basis as

$$(j_x)_{nm,n'm'} = \langle n, k_x | \hat{j}_x | n', k'_x \rangle = \langle n, k_x | \frac{1}{m} \left(-i\hbar \frac{\partial}{\partial x} + eBy \right) | n', k'_x \rangle \quad (1.5)$$

and we can evaluate these using Floquet modes derived in Eq.(??) as follows

$$\begin{aligned} (j_x)_{nm,n'm'} &= \int dx \int dy \frac{1}{\sqrt{L_x}} \chi_n(y - y_0 - \zeta(t)) \exp(-ik_x x) \\ &\times \frac{1}{m} \left(-i\hbar \frac{\partial}{\partial x} + eBy \right) \frac{1}{\sqrt{L_x}} \chi_{n'}(y - y_0 - \zeta(t)) \exp(ik'_x x) \end{aligned} \quad (1.6)$$

and this can be simplified as

$$\begin{aligned} (j_x)_{nm,n'm'} &= \frac{1}{mL_x} \int dx \exp(-i(k_x - k'_x)x) \int dy \chi_n(y - y_0 - \zeta(t)) \\ &\times (\hbar k'_x + eBy) \chi_{n'}(y - y_0 - \zeta(t)) \end{aligned} \quad (1.7)$$

and

$$(j_x)_{nm,n'm'} = \frac{1}{m} \delta_{k_x, k'_x} \int dy (\hbar k'_x + eBy) \chi_n(y - y_0 - \zeta(t)) \chi_{n'}(y - y_0 - \zeta(t)). \quad (1.8)$$

Now let $y - y_0 - \zeta(t) = \bar{y}$ and we will get

$$(j_x)_{nm,n'm'} = \frac{1}{m} \delta_{k_x, k'_x} \int d\bar{y} (\hbar k'_x + eB\bar{y} + eBy_0 + eB\zeta(t)) \chi_n(\bar{y}) \chi_{n'}(\bar{y}). \quad (1.9)$$

using definition of y_0 given in Eq. (??) this will be modified to

$$(j_x)_{nm,n'm'} = \frac{1}{m} \delta_{k_x, k'_x} \int d\bar{y} (\hbar k'_x + eB\bar{y} - \hbar k'_x + eB\zeta(t)) \chi_n(\bar{y}) \chi_{n'}(\bar{y}) \quad (1.10)$$

and using integral identities of Gauss-Hermite functions

$$\int dy \chi_n(y) \chi_{n'}(y) = \delta_{n',n} \quad (1.11)$$

$$\int dy y \chi_n(y) \chi_{n'}(y) = \left(\sqrt{\frac{n+1}{2}} \delta_{n',n+1} + \sqrt{\frac{n}{2}} \delta_{n',n-1} \right) \quad (1.12)$$

this becomes

$$(j_x)_{nm,n'm'} = \frac{1}{m} \delta_{k_x, k'_x} eB \left[\left(\sqrt{\frac{n+1}{2}} \delta_{n',n+1} + \sqrt{\frac{n}{2}} \delta_{n',n-1} \right) + \zeta(t) \delta_{n',n} \right] \quad (1.13)$$

Due to complexity we can only consider the constant contribution and we allows only the one-cycle averaged current flow and then we can derive the $s = 0$ components of the Fourier series as

$$(j_{s=0}^x)_{nm,n'm'} = \frac{1}{T} \int_0^T dt \frac{1}{m} \delta_{k_x, k'_x} eB \left[\left(\sqrt{\frac{n+1}{2}} \delta_{n',n+1} + \sqrt{\frac{n}{2}} \delta_{n',n-1} \right) + \frac{eE}{m(\omega_0^2 - \omega^2)} \sin(\omega t) \delta_{n',n} \right] \quad (1.14)$$

and this can be evaluate and get

$$(j_{s=0}^x)_{nm,n'm'} = \frac{eB}{m} \delta_{k_x, k'_x} \left(\sqrt{\frac{n+1}{2}} \delta_{n',n+1} + \sqrt{\frac{n}{2}} \delta_{n',n-1} \right) \quad (1.15)$$

For electric current operator we can introduce the electron's charge and effective mass

$$(j_{s=0}^x)_{nm,n'm'} = \frac{e^2 B}{m} \delta_{k_x, k'_x} \left(\sqrt{\frac{n+1}{2}} \delta_{n',n+1} + \sqrt{\frac{n}{2}} \delta_{n',n-1} \right) \quad (1.16)$$

■

2 Floquet-Drude Conductivity in Quantum Hall Systems

The general expression for the conductivity [*Ref: Martin Wackerl Thesis 1.250] with the disorder averaging can be represent as follows

$$\lim_{\omega \rightarrow 0} \text{Re}[\sigma^{xx}(0, \omega)] = \frac{-1}{4\pi\hbar A} \int_{\lambda-\hbar\Omega/2}^{\lambda+\hbar\Omega/2} d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) \times \text{tr} [j_0^x(\mathbf{G}^r(\varepsilon) - \mathbf{G}^a(\varepsilon))j_0^x(\mathbf{G}_0^r(\varepsilon) - \mathbf{G}_0^a(\varepsilon))]. \quad (2.1)$$

where j_0^x and $\mathbf{G}^{r,a}(\varepsilon)$ are x directional current operator matrix and white noise disorder averaged Green function matrix respectively defined against to the *Floquet modes* of the system. Here we have assumed that only $s = 0$ Fourier component of the current operator is contributing to the conductivity.

Now this can be expand in off resonant regime ($\omega\tau_0 \gg 1$) using only central entry Fourier components ($l = l' = 0$) of *Floquet modes* mentioned in Eq. (1.1) as

$$\lim_{\omega \rightarrow 0} \text{Re}[\sigma^{xx}(0, \omega)] = \frac{-1}{4\pi\hbar A} \int_{\lambda-\hbar\Omega/2}^{\lambda+\hbar\Omega/2} d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) \times \frac{1}{L_x} \sum_{k_x} \sum_n \langle n, k_x | j_0^x(\mathbf{G}^r(\varepsilon) - \mathbf{G}^a(\varepsilon)) j_0^x(\mathbf{G}_0^r(\varepsilon) - \mathbf{G}_0^a(\varepsilon)) | n, k_x \rangle \quad (2.2)$$

and one can evaluate these matrix elements as follows

$$\begin{aligned} \lim_{\omega \rightarrow 0} \text{Re}[\sigma^{xx}(0, \omega)] &= \frac{-1}{4\pi\hbar A} \int_{\lambda-\hbar\Omega/2}^{\lambda+\hbar\Omega/2} d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) \frac{1}{L_x} \sum_{k_x} \sum_n \frac{1}{L_x^3} \sum_{k_{x1}, k_{x2}, k_{x3}} \sum_{n_1, n_2, n_3} \\ &\times \langle n, k_x | j_0^x | n_1, k_{x1} \rangle \langle n_1, k_{x1} | (\mathbf{G}^r(\varepsilon) - \mathbf{G}^a(\varepsilon)) | n_2, k_{x2} \rangle \\ &\times \langle n_2, k_{x2} | j_0^x | n_3, k_{x3} \rangle \langle n_3, k_{x3} | (\mathbf{G}^r(\varepsilon) - \mathbf{G}^a(\varepsilon)) | n, k_x \rangle \end{aligned} \quad (2.3)$$

Since we can diagonalize the impurity averaged Green's function using unitary transformation ($\mathbf{T} = |n, k_x\rangle$) [*Ref: Martin Wackerl - Paper] and we can evaluate the matrix element of difference between retarded and advanced Green's function as follows [*Ref: My report 2.535]

$$\langle n_1, k_{x1} | \mathbf{T}^\dagger (\mathbf{G}^r(\varepsilon) - \mathbf{G}^a(\varepsilon)) \mathbf{T} | n_2, k_{x2} \rangle = \left[\frac{2i\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T}) \delta_{n_1, n_2} \delta_{k_{x1}, k_{x2}}}{\left(\frac{1}{\hbar}\varepsilon - \frac{1}{\hbar}\varepsilon_{n_1}\right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})]^2} \right] \quad (2.4)$$

and

$$\langle n_3, k_{x3} | \mathbf{T}^\dagger (\mathbf{G}^r(\varepsilon) - \mathbf{G}^a(\varepsilon)) \mathbf{T} | n, k_x \rangle = \left[\frac{2i\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T}) \delta_{n_3, n} \delta_{k_{x3}, k_x}}{\left(\frac{1}{\hbar}\varepsilon - \frac{1}{\hbar}\varepsilon_n\right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})]^2} \right] \quad (2.5)$$

Then applying the results we derived in previous section (??) we can calculate the conductivity

$$\begin{aligned} \lim_{\omega \rightarrow 0} \text{Re}[\sigma^{xx}(0, \omega)] &= \frac{-1}{4\pi\hbar A} \int_{\lambda-\hbar\Omega/2}^{\lambda+\hbar\Omega/2} d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) \frac{1}{L_x} \sum_{k_x} \sum_n \frac{1}{L_x^3} \sum_{k_{x1}, k_{x2}, k_{x3}} \sum_{n_1, n_2, n_3} \\ &\times \frac{e^2 B}{m} \delta_{k_x, k_{x1}} \left(\sqrt{\frac{n+1}{2}} \delta_{n_1, n+1} + \sqrt{\frac{n}{2}} \delta_{n_1, n-1} \right) \left[\frac{2i\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T}) \delta_{n_1, n_2} \delta_{k_{x1}, k_{x2}}}{\left(\frac{1}{\hbar}\varepsilon - \frac{1}{\hbar}\varepsilon_{n_1}\right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})]^2} \right] \\ &\times \frac{e^2 B}{m} \delta_{k_{x2}, k_{x3}} \left(\sqrt{\frac{n_2+1}{2}} \delta_{n_3, n_2+1} + \sqrt{\frac{n_2}{2}} \delta_{n_3, n_2-1} \right) \left[\frac{2i\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T}) \delta_{n_3, n} \delta_{k_{x3}, k_x}}{\left(\frac{1}{\hbar}\varepsilon - \frac{1}{\hbar}\varepsilon_n\right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})]^2} \right] \end{aligned} \quad (2.6)$$

and this will be modified to

$$\begin{aligned}
\lim_{\omega \rightarrow 0} \text{Re}[\sigma^{xx}(0, \omega)] &= \frac{-1}{4\pi\hbar A} \int_{\lambda-\hbar\Omega/2}^{\lambda+\hbar\Omega/2} d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) \frac{1}{L_x} \sum_{k_x} \sum_n \sum_{n_1, n_2} \\
&\times \frac{e^2 B}{m} \left(\sqrt{\frac{n+1}{2}} \delta_{n_1, n+1} + \sqrt{\frac{n}{2}} \delta_{n_1, n-1} \right) \left[\frac{2i \text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T}) \delta_{n_1, n_2}}{\left(\frac{1}{\hbar} \varepsilon - \frac{1}{\hbar} \varepsilon_{n_1} \right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})]^2} \right] \\
&\times \frac{e^2 B}{m} \left(\sqrt{\frac{n_2+1}{2}} \delta_{n, n_2+1} + \sqrt{\frac{n_2}{2}} \delta_{n, n_2-1} \right) \left[\frac{2i \text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})}{\left(\frac{1}{\hbar} \varepsilon - \frac{1}{\hbar} \varepsilon_n \right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})]^2} \right]
\end{aligned} \tag{2.7}$$

and the only non-zero term would be

$$\begin{aligned}
\lim_{\omega \rightarrow 0} \text{Re}[\sigma^{xx}(0, \omega)] &= \frac{-1}{4\pi\hbar A} \frac{e^4 B^2}{m^2} \int_{\lambda-\hbar\Omega/2}^{\lambda+\hbar\Omega/2} d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) \frac{1}{L_x} \sum_{k_x} \sum_n \left(\frac{n+1}{2} \right) \\
&\times \left[\frac{2i \text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})_{n+1}}{\left(\frac{1}{\hbar} \varepsilon - \frac{1}{\hbar} \varepsilon_{n+1} \right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})_{n+1}]^2} \right] \left[\frac{2i \text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})_n}{\left(\frac{1}{\hbar} \varepsilon - \frac{1}{\hbar} \varepsilon_n \right)^2 + [\text{Im}(\mathbf{T}^\dagger \sum^r \mathbf{T})_n]^2} \right]
\end{aligned} \tag{2.8}$$

■