


Factors Influencing Substance Use

Peer pressure, family substance use, and mental health risk
tested against substance use among survey respondents

196 VALID RESPONSES · CHI-SQUARE TESTS OF INDEPENDENCE



Each dot is one respondent: 124 reported use (coral),
72 reported no use (sage).

What This Analysis Covers

Three factors tested against substance use

1

Peer Pressure

Is experiencing peer pressure associated with a respondent's substance use?

2

Family Members' Use

Is substance use by family members associated with the respondent's own use?

3

Mental Health Risk

Is a higher mental health risk associated with substance use?

METHOD

196 valid responses · SPSS crosstabs · Pearson Chi-Square tests of independence · $\alpha = 0.05$

How the Analysis Was Conducted

Quantitative design using survey data from respondents

1

Quantitative Design

Survey data collected from respondents and analysed statistically.

2

Sample of 196

All 196 valid cases were included in the analysis.

3

Chi-Square Tests

SPSS crosstabs with the Pearson Chi-Square test of independence for each factor.


4

Decision Rule

Reject the null hypothesis if $p < 0.05$ (95% confidence level).

Substance Use Among Respondents

Each dot represents one of the 196 valid responses



124

reported substance use

63.3% of respondents

72

reported no substance use

36.7% of respondents

Nearly two in three respondents reported substance use, the baseline against which the three factors were tested.

All Three Relationships Are Significant

Pearson Chi-Square tests of independence at the 0.05 level

Peer Pressure

vs. substance use

100.675

chi-square statistic

df = 2

p = 0.000

SIGNIFICANT · REJECT H0

Family Members' Use

vs. substance use

74.258

chi-square statistic

df = 2

p = 0.000

SIGNIFICANT · REJECT H0

Mental Health Risk

vs. substance use

24.002

chi-square statistic

df = 1


p = 0.000

SIGNIFICANT · REJECT H0

Decision rule: the null hypothesis is rejected where $p < 0.05$. All three null hypotheses were rejected.

Peer Pressure and Substance Use

Share of each group reporting substance use



KEY FINDING


76.9%

of respondents who reported peer pressure also reported substance use (80 of 104).

Chi-square 100.675 · p = 0.000

Family Substance Use and Respondents' Use

Share of each group reporting substance use



KEY FINDING


100%

of respondents whose family members use substances also reported substance use (46 of 46).

Chi-square 74.258 · p = 0.000

Mental Health Risk and Substance Use

Share of each group reporting substance use



KEY FINDING

78.2%

of respondents with high mental health risk reported substance use (86 of 110), against 44.2% of the low-risk group.

Chi-square 24.002 · $p = 0.000$

Reading the Results Carefully

PEER PRESSURE: NO

100%

of respondents who reported no peer pressure (40 of 40) still reported substance use, so the result is not simply "more pressure, more use".

Observed in the peer pressure crosstab.

WHY IT MATTERS

- 1 Chi-square tests show that variables are associated, not that one causes the other.
- 2 The "Not Sure" groups show much lower use (7.7% for peer pressure, 22.6% for family use) and account for roughly 69% and 60% of the respective chi-square values.
- 3 Further analysis, such as regression on a larger sample, would help confirm the direction and strength of each factor.

What Should Be Done

01

Peer Education Programs

Schools and communities should run peer mentorship and awareness programs to reduce negative peer influence.

02

Family-Based Interventions

Educate families on the impact of substance use within the home environment.

03

Mental Health Support Services

Increase access to counseling and mental health services for individuals at risk.

04

Public Awareness Campaigns

Government agencies and NGOs should intensify campaigns on the dangers of substance use.

05

Early Detection and Prevention

Schools and healthcare institutions should adopt early screening strategies for risk factors.

THE COMMON THREAD

Target peer influence, family environments, and mental health support systems together.

Conclusion & Call to Action

1

All three factors are significantly associated with substance use: peer pressure, family members' use, and mental health risk.

2

All three null hypotheses were rejected, with p-values below the 0.05 threshold.

3

The highest rates appear with high mental health risk (78.2%) and family exposure (up to 100%).

IMMEDIATE PRIORITY

Peer & family-focused prevention

Social influences show strong associations

NEXT MOST URGENT

Mental health support & early screening

High-risk respondents report the highest use

Without targeted action on peers, families, and mental health support, substance use risk will remain high.