

Collatz-Thwaites-Ulam-Hasse-Syracuse-Kakutani (CTUHSK) Theorem  
: Convergence of Collatz  $(3n+1)$  Sequence to the Trivial Cycle Proved

© Dr.(Prof.) Keshava Prasad Halemane,  
Professor - retired (2017JAN31) from  
Department of Mathematical And Computational Sciences,  
National Institute of Technology Karnataka Surathkal,  
Srinivasnagar, Mangaluru - 575025, India.  
<https://orcid.org/0000-0003-3483-3521>

Independent Researcher  
(no affiliations; no funding; no data used; no conflict of interests)  
SASHESHA, 8-129/12 Sowjanya Road, Naigara Hills,  
Bikarnakatte, Kulshekar Post,  
Mangaluru-575005. Karnataka State, India  
[https://www.linkedin.com/in/keshavaprasadahalemane/  
k.prasad.h@gmail.com](https://www.linkedin.com/in/keshavaprasadahalemane/k.prasad.h@gmail.com)

Abstract

This paper presents the *Collatz-Thwaites-Ulam-Hasse-Syracuse-Kakutani* (CTUHSK) theorem, which asserts the convergence of the Collatz  $(3n+1)$  Sequence to the trivial cycle, thereby proving the Collatz Conjecture, a long-standing *unsolved problem*. The proof is in two parts. The necessary condition is provided by the *order-preserving isomorphism* (along with an invariant-base-element) established between the relevant component  $H^s$  of a structured system framework  $H$  and the set of positive integers. The structured system framework  $H$  itself has been designed by a two-stage bijective mapping: (1) from the Collatz-domain to

BELnet, that is the network of *binary exponential ladders* defined on the set of positive odd numbers; and (2) from BELnet to the structured system framework H. The sufficient condition is provided by a reductio-ad-absurdum argument (along with a uniquely special modular arithmetic characteristic property of the Collatz system) that is used to demonstrate *domain exhaustion*; having already captured all the modular (MOD3) residue classes in  $H^s$ ; logically excluding the existence of any extraneous elements or objects or sub-systems such as disjoint loops/cycles  $H^\infty$  and/or divergent chains  $H^\&$  or even any/all non-standard objects, in H. The proof uses the most fundamental Dedekind-Peano axioms and modular arithmetic properties of the Collatz  $(3n+1)$  System; just enough mathematics, without any unnecessary sophistications.

Some directions for possible future research work on algorithmic, computational and/or dynamic characteristics of the Collatz system have also been presented. A situation has been identified wherein the *emergence of global system properties through persistent local subsystem characteristics* can be clearly demonstrated; with  $\{(31 \leftarrow 41 \leftarrow 27)\}$  and  $\{(1 \leftarrow 5 \leftarrow 3)\}$  as exceptional and limiting cases.

Keywords: Binary-Exponential-Ladder; Modular-Periodicity; Arborescence;

CTUHSK-Theorem; CTUHSK-Generative-Parameters;

Collatz-Thwaites-Ulam-Hasse-Syracuse-Kakutani (CTUHSK) Sequence;

Collatz-Thwaites-Ulam-Hasse-Syracuse-Kakutani (CTUHSK) Conjecture;

Order-Preserving Isomorphism; Dedekind-Peano Axioms; Convergence.

AMS MSC Mathematics Subject Classification: 11B83; 11Y55; 93A05; 93A99.

## 1. Introduction

The Collatz-Thwaites-Ulam-Hasse-Syracuse-Kakutani (CTUHSK) Conjecture, simply referred to as the *Collatz Conjecture* [1] states that the Collatz Sequence, also referred to as the Collatz  $(3n+1)$  Sequence, converges to the trivial cycle, that is,  $\{(4 \rightarrow 2 \rightarrow 1)\}$ , starting from any positive integer.

Lagarias [3]&[4] gives an exhaustive annotated bibliography, whereas Lagarias [5]&[6] gives an elaborate overview of the Collatz Problem, also referred to as the  $3x+1$  problem, referring to it as “The Ultimate Challenge”. Guy [2] has been somewhat pessimistic in advising researchers “Don’t Try to Solve These Problems”. Lagarias [5]&[6] refers to the statement by the most revered Mathematician and Number Theory Expert Paul Erdos, who said that - "Mathematics is not yet ready for such problems" - about the Collatz Conjecture. The Collatz  $(3n+1)$  Sequence is indeed a highly complex chaotic system defined or generated with one of the simplest deterministic well-defined equations, that too, just two in number.

We will not get into any more of the reported literature except the problem definition; but will simply focus on a rather shockingly simple approach to solve this amazing long-standing unsolved problem. This paper presents the *Collatz-Thwaites-Ulam-Hasse-Syracuse-Kakutani* (CTUHSK) Theorem, asserting the convergence of the Collatz  $(3n+1)$  Sequence to the trivial cycle. The proof is two-fold. First is the application of the most fundamental *Dedekind-Peano Axioms* and *Modular Arithmetic* to a meticulously designed *Structured System Framework* of *Binary-Exponential-Ladders* defined on the set of positive odd numbers, and establishing an *order-preserving isomorphism* between the set of natural numbers

and the *structured system framework* of *Binary Exponential Ladders*. Secondly a reductio-ad-absurdum argument (along with a uniquely special modular arithmetic characteristic property of the Collatz system) ensures the non-existence of any extraneous elements.

## 2. Problem Description

We define the *Collatz Function*  $C(n)$  with a positive integer  $n$  as its input argument, in terms of a ‘pull-Down’ operator  $D(n)$  and a ‘push-Up’ operator  $U(n)$  as follows:

$$\text{if } (n \text{ is even}) \ C(n) := D(n) = (n/2); \text{ else } C(n) := U(n) = (3*n+1); \quad [\text{Eqn.1}]$$

where the ‘pull-Down’  $D(n)$  operator takes only an even number as its input argument whereas the ‘push-Up’ operator  $U(n)$  takes only an odd number as its input argument and gives an output that is an even number.

For convenience in our study of the Collatz Sequence, we define the *Compact Collatz Function*  $T(m)$  by the repeated application of the ‘pull-down’ operator  $D(m)$  wherever applicable, say,  $(p \geq 1)$  times, that is,  $D^*(m) := D^p(m)$  so as to get an output  $D^\#(m)$  that is an odd number:

$$\begin{aligned} \text{if } (m \text{ is even}) \ T(m) &:= D^*(m) := D^p(m) = (m / 2^p) := D^\#(m); \\ \text{else } (m \text{ is odd}) \ T(m) &:= U(m) = (3*m + 1) := U^\#(m); \end{aligned} \quad [\text{Eqn.2}]$$

where  $D^\#(m)$  is called the “D-floor number” associated with the input argument  $m$ ; and  $U^\#(m)$  is called the “U-ceiling number” associated with the input argument  $m$ .

The *Compact Collatz Function*  $T(m)$  may as well be considered to have been redefined with these newly introduced two operators, the “D-floor operator”  $D^\#(m)$  and the “U-ceiling operator”  $U^\#(m)$  as given in [Eqn.2] above.

This new definition for the *Compact Collatz Function*  $T(m)$  facilitates our study of the corresponding *Compact Collatz Sequence*; which is no different from its equivalent standard Collatz Sequence, once we understand that the repeated application, say,  $(p \geq 1)$  times, of the ‘pull-Down’ operator  $D(m)$  has now been collapsed into an equivalent single “D-floor operator”  $D^\#(m)$  giving the D-floor number  $D^\#(m)$  as its output. The push-Up operator  $U$  has been simply redefined as the “U-ceiling operator”  $U^\#$  for uniformity and elegant completeness.

The *Compact Collatz Sequence* is obtained by the repeated sequential application of the *Compact Collatz Function*  $T(m)$  starting with the given initial input number  $m$  - represented by an alternating series of  $D^\#$  number and  $U^\#$  number - except possibly the starting initial ‘seed’ number  $m$  and the final terminating number, which as per the Collatz Conjecture, is anyway a  $D^\#$  number that is unity.

### 3. Observations on the Pull-Down Operator

The pull-Down operator  $D$  *always* takes only an even number  $n$  as its input argument. Every application of this pull-down operator results in an alternating (toggling) effect on the  $n \bmod 3$  property of the input argument number; that is, a  $1 \bmod 3$  input gives a  $2 \bmod 3$  output and a  $2 \bmod 3$  input gives a  $1 \bmod 3$  output; whereas a  $0 \bmod 3$  input gives a  $0 \bmod 3$  output. Repeated application of  $D$ , in

case applicable, results in a final output that is an odd number and therefore becomes an input for the push-Up operator. In such a case, we call it a “D-floor operator”  $D^\#$  as defined in [Eqn.2] above, and its output a “D-floor number”  $D^\#(n)$  characterized by being a odd number;  $D^\#(n)$  may be in any one of the three possible types:

- (1) a 1MOD6 odd number, being a 1MOD3 odd number of the type  $(6m-5)$ ;
- (2) a 5MOD6 odd number, being a 2MOD3 odd number of the type  $(6m-1)$ ;
- (3) a 3MOD6 odd number, being a 0MOD3 odd number of the type  $(6m-3)$ .

#### 4. Observations on the Push-Up Operator

The push-Up operator  $U$  *always* takes only an odd number  $m$  as its input argument, and *always* gives an output that is a 4MOD6 number, being a 1MOD3 even number that is of the type  $(6m-2)$ ; irrespective of whether the input is a 1MOD6 odd number or a 3MOD6 odd number or a 5MOD6 odd number. Note that one single application of the ‘push-Up’ operator  $U$  transforms any input odd number  $m$  into a 4MOD6 even number that becomes an input to the “D-floor operator  $D^\#$ ”. That is why we may as well consider the push-Up operator  $U$  as the “U-ceiling operator”  $U^\#$  as defined in [Eqn.2] above.

#### 5. Observations on the Compact Collatz Function

Start with any positive integer. (1) If the starting initial number  $n$  is even, then we apply the D-floor operator  $D^\#$  giving an output that is the D-floor number  $D^\#(n)$  which is given as input to the U-ceiling operator. Of course, if the starting

number is a power of 2 we terminate at unity. Else, we have a D-floor number  $D^\#(n)$  that is an odd number greater than unity, in any non-trivial case; as the initial  $D^\#$  node in the Compact Collatz Sequence. (2) If on the other hand the starting initial seed number  $n$  is an odd number, we treat that itself as the initial  $D^\#$  node in the Compact Collatz Sequence.

Having thus obtained the initial  $D^\#$  node in the Compact Collatz Sequence, we apply the U-ceiling operator  $U^\#$  to get the U-ceiling number  $U^\#$  that is a  $4\text{MOD}6$  even number. That in turn is given as input to the D-floor operator  $D^\#$ . Now the process continues.

Note that the *Compact Collatz Sequence* can therefore be defined by a *trajectory* generated by an *alternating sequence* of a “D-floor number”  $D^\#$  and a “U-ceiling number”  $U^\#$ , *with its starting initial node being a  $D^\#$  number*. The Compact Collatz Function as presented in [Eqn.2] defines the unique link (directed arc) from any given D-floor number  $D^\#$  as the predecessor node to its corresponding unique U-ceiling number  $U^\#$  as the successor node and also the unique link (directed arc) from any given U-ceiling number  $U^\#$  as the predecessor node to its corresponding unique D-floor number  $D^\#$  as the successor node. The unique link (directed arc) from a starting initial even “seed” number leading to the first node (D-floor number  $D^\#$ ) in the *trajectory* is similarly defined.

As mentioned earlier, the application of the D-floor operator  $D^\#$  on a U-ceiling number  $U^\#$  that is a  $4\text{MOD}6$  even number of the form  $(6m-2)$  can lead to a D-floor number  $D^\#$  that is an odd number that can be either: (1) a  $1\text{MOD}6$  odd number, being a  $1\text{MOD}3$  odd number that is of the type  $(6m-5)$ ; (2) a  $5\text{MOD}6$  odd number, being a  $2\text{MOD}3$  odd number that is of the type  $(6m-1)$ ; but can never be (3) a

3MOD6 odd number, that is a 0MOD3 odd number of the type  $(6m-3)$ . The only exception, when the D-floor operator  $D^\#$  gives an output D-floor number  $D^\#$  that is a 3MOD6 odd number of the type  $(6m-3)$  is the situation when its input is a 0MOD6 even number, which is impossible for any U-ceiling number  $U^\#$ ; although such an input may come in those special cases wherein the starting initial ‘seed’ number itself is a 0MOD6 even number that is of the form  $(6m-3).2^p$  leading to an output  $D^\#$  that is a 3MOD6 odd number of the form  $(6m-3)$ .

## 6. Analysis of the Compact Collatz Sequence

From the above observations, it is clear that corresponding to every positive integer  $n$  as the starting initial ‘seed’ number, there is a starting initial node in the trajectory representing the *Compact Collatz Sequence*, that is a  $D^\#$  number in exactly one of the three possible forms as mentioned above - that can be an input argument to the U-ceiling operator  $U^\#$  giving exactly one unique output  $U^\#$  which itself can be an input to the D-floor operator  $D^\#$  so that the process continues. Successive application of each of these two operators ( $U^\#$  and  $D^\#$ ) wherever applicable, traces a unique *trajectory*, wherein each node is represented by the unique output number of the appropriate operation applied to the input number represented by the preceding node in the trajectory.

The anticipated terminating trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\}$  can be obtained only through a final application of the D-floor operator  $D^\#$  on a 4MOD6 even number that is of the form  $(6m-2).2^p$  with  $m=1$  and  $0 \leq p$ .



## 7. Binary-Exponential-Ladder with its Defining-Base-Rung $D^\#$

Here, we present a meticulously designed *Structured System Framework* that *partitions* the *set of positive integers* to facilitate a *general systems analysis* of the *Compact Collatz Sequence*.

Let every positive odd number be associated with a *Binary-Exponential-Ladder*, denoted by  $BEL(2m-1)$  and defined as a sequence  $\{(2m-1).2^u \mid (u \geq 0)\}$ ; with its *defining-base-rung* ( $u=0$ ) given by the odd number  $(2m-1)$ .

$$BEL(2m-1) := \{(2m-1).2^u \mid (u \geq 0); (m \geq 0)\}; \quad [Eqn.3]$$

Thus, we establish an exact one-to-one correspondence (bijection) between the *set of positive odd numbers* that form the  $D^\#$  value and the *defining-base-rung* of the corresponding *Binary Exponential Ladder*  $BEL(D^\#)$ .

Every *positive even number* in the form  $\{(2m-1).2^u \mid (u > 0)\}$ ; for which there exists its corresponding  $D^\#$  value,  $D^\#((2m-1).2^u) = (2m-1)$ ; there exists exactly one corresponding *Binary-Exponential-Ladder*  $BEL(2m-1)$  that contains the given even number  $(2m-1).2^u$  as  $B((2m-1), u)$  as one of the higher rungs in  $BEL(2m-1)$ .

$$B((2m-1), u) := \{(2m-1).2^u \mid (u > 0)\}; \quad [Eqn.4]$$

Therefore, *the set of all Binary-Exponential-Ladders form a partition of the set of all positive integers*; with an *exact one-to-one correspondence (bijection)* between each positive odd number  $D^\#$  and the corresponding *Binary-Exponential-Ladder*

for which it is the *defining-base-rung*  $D^\#$ ; whereas each of the positive even numbers correspond to exactly one of the higher rungs of a specific Binary-Exponential-Ladder identified by the D-floor number  $D^\#$  associated with that given positive even number.

This partitioned framework of positive integers goes another step deeper because of the fact that the *defining-base-rung*  $D^\#$  of a Binary-Exponential-Ladder  $BEL(D^\#)$  can itself be in one of the three possible forms 1MOD6 or 5MOD6 or 3MOD6; whereas all the upper rungs of the Binary-Exponential-Ladder are either (1) alternatingly 2MOD6 and 4MOD6; or (2) all being 0MOD6 numbers.

The Collatz Conjecture states that every Collatz Sequence, starting from any positive integer, converges to the trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\}$  which is in  $BEL(1)$  that is uniquely identified by its defining-base-rung  $D^\#$  value that is unity. Therefore, our focus will be the Binary-Exponential-Ladders  $BEL(1)$  and its relationship with every other Binary-Exponential-Ladder  $BEL(D^\#)$ .

Note that  $D^\#$  can be a positive odd number in any one of the three possible forms: (1) a 1MOD6 number of the form  $(6m-5)$ ; (2) a 5MOD6 number of the form  $(6m-1)$ ; (3) a 3MOD6 number of the form  $(6m-3)$ .  $BEL(6m-5)$  contains the output of  $U^\#$  at a higher rung  $(6m-5)2^W$  with  $w$  being a positive even exponent; the input of  $U^\#$  being given by  $\lfloor \{(6m-5).2^W - 1\} / 3 \rfloor$ .  $BEL(6m-1)$  contains the output of  $U^\#$  at a higher rung  $(6m-1)2^V$  with  $v$  being a positive odd exponent; the input of  $U^\#$  being given by  $\lfloor \{(6m-1).2^V - 1\} / 3 \rfloor$ . However,  $BEL(6m-3)$  cannot contain any such output of the U-ceiling operator  $U^\#$  irrespective of any input argument.

## 8. Immediate Neighborhood of a Binary-Exponential-Ladder

The relationship between a pair of Binary-Exponential-Ladders  $BEL(m)$  and  $BEL(n)$  can be considered to be defined and characterized by the relationship between the corresponding pair of the *defining-base-rung*  $D^\#$  values  $m$  and  $n$  along with the corresponding pair  $U^\#(m)$  and  $U^\#(n)$ .

The immediate-neighborhood of a given Binary-Exponential-Ladder  $BEL(D^\#)$  is defined by the *immediate-predecessors* and *immediate-successors*, considering the  $U$ -ceiling operator  $U^\#$ ; since the  $D$ -floor operator  $D^\#$  is applicable only within a given Binary-exponential-Ladder and not between a pair of them.

### 8.1 Single Unique Immediate Successor – CTUHSK function

It turns out that the only *one single unique immediate successor* of  $BEL(m)$  is  $BEL(D^\#(U^\#(m)))$  that contains  $U^\#(m)$  as one of its higher rungs; its identifying characteristic  $D$ -floor number associated with its defining-base-rung being given by  $n := D^\#(U^\#(m))$ . That is,

$$S(BEL(m)) = BEL(D^\#(U^\#(m))) := BEL(n); \text{ with } n := D^\#(U^\#(m)); \quad [\text{Eqn.5}]$$

Note that [Eqn.5] describes the forward tracing movement (transfer function) from one  $BEL$  to its immediate successor  $BEL$  defined by the very same Compact Collatz Function [Eqn.2] above; wherein the downward movement ( $D^\#$  operator) from a higher rung to the defining-base-rung within the same  $BEL$  is compressed and left as implicit; whereas the upward movement from the defining-base-rung of one  $BEL$

to a higher rung of its immediate-successor BEL is kept explicit; thus redefining the compact Collatz sequence by its equivalent, a *condensed compact Collatz* (CTUHSK) *sequence*; which is simply a hop from one BEL to its immediate successor BEL. Here, [Eqn.5] can therefore be considered as the condensed compact Collatz (CTUHSK) function; representing one hop from a positive odd number to its immediate successor odd number; thus defining the condensed compact Collatz (CTUHSK) sequence, that is equivalent to skipping the intermediate even numbers in the standard Collatz sequence. Our study will focus on the characteristics of this condensed compact Collatz (CTUHSK) sequence defined by the condensed compact Collatz (CTUHSK) function; or equivalently the characteristics of BELnet, that is, the network of binary exponential ladders.

## 8.2 Multiple Immediate Predecessors – inverse CTUHSK function

There exists a *set of immediate-predecessors* for each  $BEL(D^\#)$  of the form  $BEL(6m-5)$  and  $BEL(6m-1)$  although none for  $BEL(6m-3)$ . Note that if  $S(BEL(m))$  is  $BEL(n)$  than  $BEL(m)$  is one of the predecessors of  $BEL(n)$ .

The *set of immediate-predecessors* for a given  $BEL(n)$  is defined by considering the *inverse of the immediate-successor relationship*; as the set of all BELs each of which having its single unique immediate-successor as  $BEL(n)$ .

$$\{P(BEL(n))\} := \{BEL(m) \mid BEL(n) = S(BEL(m))\}; \quad [Eqn.6]$$

$BEL(1 \bmod 6)$  or equivalently  $BEL(6m-5)$  has, as its set of immediate-predecessors,  $\{BEL([(1 \bmod 6) \cdot 2^w - 1]/3)\}$  or equivalently  $\{BEL([(6m-5) \cdot 2^w - 1]/3)\}$  with  $w$  being a positive even exponent. Here, the input of  $U^\#$  is given by  $\{[(1 \bmod 6) \cdot 2^w - 1]/3\}$

or equivalently  $\{((6m-5).2^W - 1)/3\}$  and the output of  $U^\#$  is  $\{(1 \bmod 6).2^W\}$  or equivalently  $\{(6m-5).2^W\}$  that is contained in  $BEL(1 \bmod 6)$  or equivalently  $BEL(6m-5)$ . Each of the three possible classes of  $BEL$ , namely,  $BEL(1 \bmod 6)$  and  $BEL(5 \bmod 6)$  and  $BEL(3 \bmod 6)$  can be the immediate-predecessor of  $BEL(1 \bmod 6)$ .

$$\{P(BEL(6m-5))\} = \{BEL(|(6m-5).2^W - 1|/3)\}; \quad [Eqn.7]$$

$BEL(5 \bmod 6)$  or equivalently  $BEL(6m-1)$  has, its set of immediate-predecessors,  $\{BEL(|(5 \bmod 6).2^V - 1|/3)\}$  or equivalently  $\{BEL(|(6m-1).2^V - 1|/3)\}$  with  $v$  being a positive odd exponent. Here, the input of  $U^\#$  is given by  $\{(5 \bmod 6).2^V - 1\}/3$  or equivalently  $\{(6m-1).2^V - 1\}/3$  and the output of  $U^\#$  is  $\{(5 \bmod 6).2^V\}$  or equivalently  $\{(6m-1).2^V\}$  that is contained in  $BEL(5 \bmod 6)$  or equivalently  $BEL(6m-1)$ . Each of the three possible classes of  $BEL$ , namely,  $BEL(1 \bmod 6)$  and  $BEL(5 \bmod 6)$  and  $BEL(3 \bmod 6)$  can be the immediate-predecessor of  $BEL(5 \bmod 6)$ .

$$\{P(BEL(6m-1))\} = \{BEL(|(6m-1).2^V - 1|/3)\}; \quad [Eqn.8]$$

$BEL(3 \bmod 6)$  or equivalently  $BEL(6m-3)$  has no immediate-predecessors.

$$\{P(BEL(6m-3))\} = \phi; \quad [Eqn.9]$$

Note that [Eqn.6] or equivalently, [Eqn.7]&[Eqn.8]&[Eqn.9] together define the algebra for the inverse of the condensed compact Collatz (CTUHSK) function.

The CTUHSK function in [Eqn.5] along with its inverse in [Eqn.6] or equivalently in [Eqn.7]&[Eqn.8]&[Eqn.9] together provide a complete mathematical model for the network of binary exponential ladders.

### 8.3 Quaternary-Exponential-Ladder

The above mentioned property, that *only* the alternating rungs, defined by  $\{(1 \bmod 6).4^u \mid u > 0\}$  of  $BEL(1 \bmod 6)$  or  $\{(5 \bmod 6).2.4^u \mid u \geq 0\}$  of  $BEL(5 \bmod 6)$  are the ‘active’ higher rungs of BEL forming the nodes in the CTUHSK-Sequence; makes it convenient to define a system of *Quaternary-Exponential-Ladders* QEL wherein every rung of QEL becomes an active node in the CTUHSK-Sequence. This concept is not directly needed for proving the convergence of the Collatz Sequence, and therefore we will not take up this line of study in this paper.

### 8.4 BEL(1) as the Central Focus

Considering  $BEL(1)$  as our central focus of interest, which itself belongs to the type  $BEL(1 \bmod 6)$  or equivalently  $BEL(6m-5)$ ; it is interesting to note that it has its *single unique immediate-successor*; as  $S(BEL(1)) = BEL(D^\#(U^\#(1))) = BEL(1)$ ; that is,  $BEL(1)$  itself is its single unique immediate-successor, and that it has no other immediate-successor distinct from itself; because of the fact that the trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\}$  contained within  $BEL(1)$ .

### 8.5. BELnet : Network of Binary-Exponential-Ladders

The above discussion about the successor predecessor relationship among the binary-exponential-ladders and its neighborhood leads to the observation that the

*network of binary-exponential-ladders*, BELnet, has countably infinite number of each of the three classes/types of nodes: (1) BEL(1MOD6) or equivalently BEL(6m-5); (2) BEL(5MOD6) or equivalently BEL(6m-1); and (3) BEL(3MOD6) or equivalently BEL(6m-3). Each BEL being a node of the BELnet has a single unique outward directed arc that points towards its single unique immediate-successor, specifically linking onto some higher rung. Multiple (countably infinite number of) inward directed arcs, each linked onto some specific higher rung of a given BEL, emanate from each of its immediate-predecessors. BEL(1) is an invariant-base-element or equivalently a sink node in BELnet, the network of binary exponential ladders.

Thus, the set of binary-exponential-ladders is an exact representation of the Collatz-domain and the network of binary-exponential-ladders BELnet is an exact representation of the Collatz-map.

The *connectedness* of the network of binary-exponential-ladders BELnet will be analyzed from the *design of a structured system framework* consisting of the entire set of binary-exponential-ladders, merely as a re-organized *condensation* of the very same BELnet, as presented below.

## 9. Structured System Framework H

From the above discussion we find that it is convenient for our study to consider a *Structured System Framework* H as an infinite (well-ordered) sequence of terms each of which being a set of BELs; that is,  $H := \{ \dots H_{k-1} < H_k < H_{k+1} \dots \}$  wherein the *well-ordering relationship* between the adjacent terms of the sequence is derived

from the successor predecessor relationships among the BELs that form the member elements of these adjacent terms in the sequence.

Specifically,  $H_k$  is defined as the set formed by the unique immediate-successor of each BEL belonging to  $H_{k+1}$  and also the set of immediate-predecessors of each BEL belonging to  $H_{k-1}$ ; that is,

$$H := \{ \dots H_{k-1} < H_k < H_{k+1} \dots \}; \quad [\text{Eqn.10}]$$

and

$$H_k := \{S(\text{BEL}(m)) \mid \text{BEL}(m) \in H_{k+1}\} \cup \{\text{BEL}(m) \mid S(\text{BEL}(m)) \in H_{k-1}\}; \quad [\text{Eqn.11}]$$

Note that the second part of [Eqn.11] here is required to ensure that BELs of the class/type  $\text{BEL}(3\text{MOD}6)$  can be included in each term  $H_k$  since each of them have immediate-successor in  $H_{k-1}$  although none of them have any predecessors in  $H_{k+1}$ .

Now, we may as well define the predecessor relationship as the inverse of the above defined successor relationship, as –

$$H_{k-1} := S(H_k) \quad \text{and} \quad H_k := S(H_{k+1}); \quad [\text{Eqn.12}]$$

and

$$P(H_{k-1}) := H_k \quad \text{and} \quad P(H_k) := H_{k+1}; \quad [\text{Eqn.13}]$$

The multiplicity of the *immediate-predecessor* relationship among the BELs requires that the set of all immediate-predecessors of every element of  $H_{k-1}$  form the elements of the set  $H_k$  so as to guarantee the strict and complete linear ordering relation  $H_{k-1} < H_k < H_{k+1}$  among these sets, in spite of only a partial ordering



relationship among the BELs; and also to guarantee that the entire set of all the BELs are present in  $H$  thus making it as a *re-organized structure* for BELnet.

### 9.1 Closed Chains and Unbounded Chains and Sink Nodes in $H$

The design of the structured system framework  $H$  can *in general* allow for the existence of sink nodes (invariant-base-elements) and/or unbounded open chains and/or closed chains (loops). That is, the structured system framework  $H$  can in general be *partitioned into three mutually disjoint and independent components*,

$$H := H^s \cup H^{\&} \cup H^\infty \quad [\text{Eqn.14}]$$

where (1)  $H^s$  corresponds to the set of all possible terms in  $H$  connected with sink nodes; (2)  $H^{\&}$  corresponds to the set of all possible terms in  $H$  connected with unbounded open chains; and (3)  $H^\infty$  corresponds to the set of all possible terms in  $H$  connected with closed chains (loops). In such a situation, each of these components,  $H^s$  and  $H^{\&}$  and  $H^\infty$  needs to satisfy the well-ordering conditions expressed above in [Eqn.10], [Eqn.11], [Eqn.12] & [Eqn.13].

### 9.2 A Sink Node $H_0$ in $H^s$

We have observed earlier that  $BEL(1)$  itself is its single unique immediate-successor and does not have any immediate-successor distinct from itself, although it has multiple immediate-predecessors. That is,  $BEL(1)$  is an invariant-base-element or equivalently a sink node in BELnet. Therefore, the component  $H^s$  must necessarily

have a term  $H_0$  as its *invariant-base-element* or equivalently a *sink node*, that is,  $H_0 := \{\text{BEL}(1)\}$ ; with  $\text{BEL}(1)$  being its singleton member element. That is,

$$H^s := \{H_0, H_1, H_2, \dots\}; \quad [\text{Eqn.15}]$$

From the above discussion we observe that  $H^s := \{H_0, H_1, H_2, \dots\}$  is, by its very design, an infinite (well-ordered) sequence of terms, each term being a countably infinite set of BELs with an exception that the invariant base element or the ‘root’  $H_0 := \{\text{BEL}(1)\}$  is a singleton set. The set of  $k^{\text{th}}$  immediate predecessors of  $\text{BEL}(1)$  form the set  $H_k$  at tier- $k$  level in the hierarchy, if one wishes to consider it as a hierarchy.

Note that the entire Collatz Domain has been mapped to the structured system framework  $H$  through a two-stage bijective mapping.

## 10. Collatz-Thwaites-Ulam-Hasse-Syracuse-Kakutani Theorem

### STATEMENT OF THE CTUHSK THEOREM

The Collatz Sequence converges to the trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\}$ .

### PROOF

There are two parts to the proof – a *necessary condition* and a *sufficient condition*. The necessary condition is provided by establishing an order-isomorphism between

$H^s$  and the set of natural numbers, by the application of the Dedekind-Peano axioms; thus proving the convergence of the CTUHSK sequence, starting from any given positive integer contained in any BEL that is in  $H^s$  to the trivial cycle at the base of  $BEL(1)$  which itself is in  $H_0 \in H^s$ . The sufficient condition is provided by a reductio-ad-absurdum argument (along with a uniquely special modular arithmetic characteristic property of the Collatz system) that establishes *domain exhaustion*; having already captured all the modular (MOD3) residue classes in  $H^s$ ; and guarantees the non-existence of any extraneous elements.

### 10.1 Proof of CTUHSL Theorem - Necessary Condition

We show that  $H^s$  satisfies the Dedekind-Peano's axioms (replacing the 'successor' by the 'predecessor') and therefore  $H^s$  is order-isomorphic with the set of natural numbers; and satisfies the necessary condition for the above convergence statement.

DEDEKIND-PEANO AXIOM : Existence of 1 as the invariant-base-element.

$H_0 \in H^s$ .  $H_0$  is the invariant-base-element of  $H^s$ .

The trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\} \in BEL(1)$  is contained in  $H_0 \in H^s$ .

DEDEKIND-PEANO AXIOM : Existence of a *successor function*.

By the very design of  $H^s := \{H_0, H_1, H_2, \dots\}$ , for every positive integer  $k$ ,

$H_k \in H^s$  is the *predecessor* of  $H_{k+1} \in H^s$ .

Application of the CTUHSK function with the input from numbers contained in some BEL that is a member of  $H_k$  yields the single unique output number contained in some immediate-successor BEL that is a member of  $H_{k+1}$ ; because of the definition of the successor predecessor relationship between  $H_k$  and  $H_{k+1}$ .

DEDEKIND-PEANO AXIOM : 1 is not a successor; 1 has no predecessor; 1 is a source node in the sequence of natural numbers.

$H_0$  is not a predecessor to any other  $H_k$ . There *does not exist any*  $H_k \in H^s$ ,  $k \neq 0$ ; that is distinct from  $H_0$ ; with  $H_k \neq H_0$ ; such that  $H_0$  is the predecessor of  $H_k$ .

$H_0$  does not have any successor distinct from itself.

$H_0$  is a sink node in the sequence  $H^s := \{H_0, H_1, H_2, \dots\}$ .

Once the CTUHSK Sequence reaches the trivial cycle/sink there is no exit from it.

DEDEKIND-PEANO AXIOM : Successor function is a unique one-to-one mapping.

If  $H_u$  is the predecessor of  $H_v$  and also  $H_u$  is the predecessor of  $H_w$ ;

then it necessarily implies  $H_v = H_w$  by the very design of  $H^s$ ;

and,

If  $H_v$  is the predecessor of  $H_u$  and also  $H_w$  is the predecessor of  $H_u$ ;

then it necessarily implies  $H_v = H_w$  by the very design of  $H^s$ ;

This is because  $H^s$  is a complete linear order sequence.

Also, note that for each positive integer  $k$  there corresponds a unique set  $H_k \in H^s$ , and for each  $H_k \in H^s$  there corresponds a unique positive integer  $k$ ; thus, establishing a *bijection* between  $H^s$  and the set of positive integers.

DEDEKIND-PEANO AXIOM : Principle of induction.

Collatz Sequence starting with any number from  $BEL(1) \in H_0$  converges in the trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\} \in BEL(1) \in H_0$ .

Collatz Sequence starting with any positive number that passes through  $H_k$  must necessarily pass through  $H_{k-1}$  because by design  $H_{k-1} := S(H_k)$ .

Thus, we establish a direct *order-preserving isomorphism* between  $H^s$  and the set of natural numbers.

This guarantees the CTUHSK sequence starting from any positive integer that is contained in  $H^s$  passes through a linear directed path (chain) with no forking or merging in  $H^s$  (although merging is observed deeper at the level of the BELs); moving through  $\dots H_{k+1}$  to  $H_k$  to  $H_{k-1} \dots$  etc. to  $H_0$  in that order; and converging in the trivial cycle.

## 10.2 Proof of CTUHSL Theorem - Sufficient Condition

The proof for the sufficient condition is provided in two parts; that is, by establishing the non-existence of any extraneous elements/objects/sub-systems as (1) disjoint loops/cycles  $H^\infty$ ; and/or (2) divergent chains  $H^\&$  in the structured system framework  $H$ ; so that  $H^s = H$ .

### 10.2.1 Non-existence of Disjoint Loops/Cycles $H^\infty$ in $H$

The non-existence of any loop/cycle  $H^\infty$  (finite or infinite) other than the trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\}$  in BELnet can be proved here by a *reductio-ad-absurdum* argument. If indeed there exists such a loop/cycle, it must necessarily be a sink loop with no exit, because of the characteristic property that there exists only a

single unique immediate-successor for every BEL in BELnet. Also, the non-existence of any immediate-predecessor for BEL(6m-3) implies that BEL(6m-3) cannot be in any such loop. Being a sink loop consisting of only BEL(6m-5) and BEL(6m-1) - each with multiple immediate-predecessors; implies that such a loop can be collapsed into an element, say,  $X_0$  in  $H^\infty$  (that is similar to  $H_0$  as in  $H^s$ ); and a sequence of predecessors can be developed similar to [Eqn.15] as in [Eqn.16];

$$H^\infty := \{X_0, X_1, X_2, \dots\}; \quad [\text{Eqn.16}]$$

Applying the same arguments as in the necessary condition of the theorem presented above, we can establish the order-isomorphism between the set of natural numbers and  $H^\infty$ ; with  $H^\infty$  itself having the loop/cycle as a sink in the super-node  $X_0$ . This implies that there is a parallel system that is equivalent to  $H^s$  that is also order-isomorphic to the set of natural numbers. As explained above,  $X_0$  can contain only  $\{\text{BEL}(6m-5)\}$  and  $\{\text{BEL}(6m-1)\}$  but cannot contain  $\{\text{BEL}(6m-3)\}$ .

Suppose among the set of BELs of the form  $\text{BEL}(6m-1) \in X_0$ ; we assume that there exists a  $\text{BEL}(6k-1) \in X_0$ ; that is the BEL with the minimum numerical value associated with its defining base rung of lowest denomination. Applying [Eqn.8] with  $v=1$  leads to a situation wherein we get the set of its immediate predecessors of lowest denomination as -

$$\{P(\text{BEL}(6k-1))\} = \{\text{BEL}([((6m-1) \cdot 2^1 - 1)/3])\} = \{\text{BEL}(4k-1)\}; \quad [\text{Eqn.17}]$$

establishing that  $\text{BEL}(4k-1) \in X_0$ .

However,  $(4k-1) \bmod 3$  must necessarily be 1 because it can neither be 2 since we assumed that  $BEL(6k-1)$  is of the lowest denomination and  $(4k-1) < (6k-1)$ ; nor can it be 0 since it is in  $X_0$  which cannot contain  $BEL(6m-3)$  as already observed earlier. The lowest value of  $k$  with  $(4k-1) \bmod 3 = 1$  and  $(6k-1) \bmod 3$  being equal to 2 is  $k=2$  thus making  $BEL(4k-1) = BEL(7) \in X_0$  and  $BEL(6k-1) = BEL(11) \in X_0$ . This is a direct contradiction to the above assumption, since  $BEL(7) \in H^s$  and  $BEL(11) \in H^s$  and also that  $H^s$  and  $X_0 \in H^\infty$  are mutually disjoint.

Therefore, we conclude the non-existence of any such loops/cycles  $H^\infty$  in BELnet whether finite or infinite, and hence the non-existence of  $H^\infty$  in  $H$ .

### 10.2.2 Non-existence of Divergent Chains $H^\&$ in $H$

Suppose among the set of BELs of the form  $BEL(6m-1) \in H^\&$ ; we assume that there exists a  $BEL(6k-1) \in H^\&$ ; that is the BEL with the minimum numerical value associated with its defining base rung of lowest denomination. We can use exactly the similar reductio-ad-absurdum argument banking on [Eqn.17] in establishing the non-existence of  $H^\infty$  as used above; to show that  $BEL(6k-1) \in H^\&$  must necessarily implies that  $BEL(4k-1) \in H^\&$ . Because  $k$  is necessarily a natural number belonging to the Collatz domain; although  $H^\&$  is assumed to be a divergent chain; such a possibility leads to the limiting situation wherein  $k$  can take its least possible value,  $k=1$  thus making  $BEL(4k-1) = BEL(3) \in H^\&$  and  $BEL(6k-1) = BEL(5) \in H^\&$ . This is a direct contradiction to the above assumption, since  $BEL(3) \in H^s$  and  $BEL(5) \in H^s$  and also that  $H^s$  and  $H^\&$  are mutually disjoint. Therefore, there cannot exist any divergent chain (infinite threads); and  $H^\& = \phi$ .

### 10.2.3 Existence of any/every/all Non-standard Rogue Elements in H

Note that the Collatz-domain does not allow for the existence of any non-standard *rogue elements* etc. The study of non-standard extraneous objects (of whatever kind) is immaterial to the present proof, because the Collatz  $(3n+1)$  Conjecture is stated strictly within the domain of natural numbers.

Having established that  $H^\infty = \phi$  and  $H^\& = \phi$  in [Eqn.14] above; we get  $H = H^\&$ ; therefore, asserting the Collatz Conjecture, that starting from any positive integer the Collatz  $(3n+1)$  Sequence converges to the trivial cycle  $\{(4->2->1)\}$ .

END OF PROOF

## 11. Some Explicit Forms for the BEL Neighborhood

We can perform some simple algebraic manipulation to get the parametric relation [Eqn.18] that gives a generic form for the set  $H_k$  that is the set of  $k^{\text{th}}$  predecessors of  $H_0 = \{\text{BEL}(1)\}$ ; that is, the set  $H_k$  corresponds to the set of tier-k level of the hierarchy with the set of Binary-Exponential-Ladders  $\{\text{BEL}(m)\}$  each with its defining-base-rung  $m$  being a positive odd number  $m > 1$ .

$$m = [2^z - \{3^0.2^{z0} + 3^1.2^{z1} + 3^2.2^{z2} + \dots + 3^{k-1}.2^{zk-1}\}] / 3^k; \quad [\text{Eqn.18}]$$



wherein  $k > 0$  is the tier-level and  $z > z_0$  whereas the  $k$ -tuple  $(z_0, z_1, z_2, \dots, z_{k-1})$  form the set of strictly decreasing non-negative integer exponents values in [Eqn.18] each of which takes a unique value corresponding to each positive odd number  $m > 1$ . That is, each positive odd number  $m > 1$  can be considered to be defined by the corresponding unique set of these parameters. The set of values for the  $k$ -tuple  $(z_0, z_1, z_2, \dots, z_{k-1})$  are of strictly decreasing non-negative integer exponent values, all less than  $z$ , that is,  $z > z_0 > z_1 > z_2 > \dots > z_{k-1}$ ; ( $z_k := 0$ ;  $z_{k-1} = 0$  for positive odd number  $m > 1$ ).

Now, define  $p_0 := (z - z_0)$ ;  $p_j := (z_{j-1} - z_j)$ ; where  $p_j$  corresponds to the number of rungs in  $BEL\{H_j\}$  above the *defining-base-rung* of  $BEL\{H_j\}$  for the node located in  $BEL\{H_j\}$  that the Collatz sequence/trajectory passes through;  $BEL\{H_j\}$  being the *Binary-Exponential-Ladder* at tier- $j$  with  $j=0,1,2, \dots, k$ . Thus, we may as well redefine the set of  $(k+1)$  parameters as a tuple  $(GPT) := (p_0, p_1, p_2, \dots, p_k)$  the set of  $(k+1)$  CTUHSK generative parameters that generate each positive integer  $n$  as per the parametric relation [Eqn.18] given above ( $p_k = 0$  for positive odd number  $m$ ).

For any  $k > 0$ , the above set of exponents  $z, z_0, z_1, z_2, z_3, \dots, z_k$ , can be redefined in terms of the newly defined CTUHSK generative parameters, by rewriting the above definition as  $z := (z_0 + p_0)$ ;  $z_{j-1} := (z_j + p_j)$ ;  $z_k := 0$ ;  $p_k = 0$  for any positive odd number  $m$ .

Table-1 gives some of the possible set of valid values for the CTUHSK generative parameters and therefore the corresponding valid values of the exponents in [Eqn.18] above along with their resultant  $n(GPT) := n(p_0, p_1, p_2, \dots, p_k)$  values.

| Table-1 : Some typical CTUHSK generative parameter tuples GPT(n) |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|------------------------------------------------------------------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| p0                                                               | p1 | p2 | p3 | p4 | p5 | p6 | p7 | p8 | n  | z  | z0 | z1 | z2 | z3 | z4 | z5 | z6 | z6 | z7 |
| 0                                                                |    |    |    |    |    |    |    |    | 1  | 2  | 0  |    |    |    |    |    |    |    |    |
| 1                                                                |    |    |    |    |    |    |    |    | 2  |    |    |    |    |    |    |    |    |    |    |
| 2                                                                |    |    |    |    |    |    |    |    | 4  |    |    |    |    |    |    |    |    |    |    |
| 3                                                                |    |    |    |    |    |    |    |    | 8  |    |    |    |    |    |    |    |    |    |    |
| 4                                                                |    |    |    |    |    |    |    |    | 16 |    |    |    |    |    |    |    |    |    |    |
| 4                                                                | 0  |    |    |    |    |    |    |    | 5  | 4  | 0  | 0  |    |    |    |    |    |    |    |
| 4                                                                | 1  | 0  |    |    |    |    |    |    | 3  | 5  | 1  | 0  | 0  |    |    |    |    |    |    |
| 4                                                                | 3  | 0  |    |    |    |    |    |    | 13 | 7  | 3  | 0  | 0  |    |    |    |    |    |    |
| 4                                                                | 3  | 2  | 0  |    |    |    |    |    | 17 | 9  | 5  | 2  | 0  | 0  |    |    |    |    |    |
| 4                                                                | 3  | 2  | 1  | 0  |    |    |    |    | 11 | 10 | 6  | 3  | 1  | 0  | 0  |    |    |    |    |
| 4                                                                | 3  | 2  | 1  | 1  | 0  |    |    |    | 7  | 11 | 7  | 4  | 2  | 1  | 0  | 0  |    |    |    |
| 4                                                                | 3  | 2  | 1  | 1  | 2  | 0  |    |    | 9  | 13 | 9  | 6  | 4  | 3  | 2  | 0  | 0  |    |    |
| 6                                                                | 0  |    |    |    |    |    |    |    | 21 | 6  | 0  | 0  |    |    |    |    |    |    |    |
| 4                                                                | 3  | 2  | 1  | 3  | 0  |    |    |    | 29 | 13 | 9  | 6  | 4  | 3  | 0  | 0  |    |    |    |
| 4                                                                | 3  | 2  | 1  | 3  | 1  | 0  |    |    | 19 | 14 | 10 | 7  | 5  | 4  | 1  | 0  | 0  |    |    |
| 4                                                                | 3  | 2  | 1  | 3  | 1  | 2  | 0  |    | 25 | 16 | 12 | 9  | 7  | 6  | 3  | 2  | 0  | 0  |    |
| 4                                                                | 3  | 2  | 1  | 3  | 1  | 2  | 2  | 0  | 33 | 18 | 14 | 11 | 9  | 8  | 5  | 4  | 2  | 0  | 0  |
| p0                                                               | p1 | p2 | p3 | p4 | p5 | p6 | p7 | p8 | n  | z  | z0 | z1 | z2 | z3 | z4 | z5 | z6 | z6 | z7 |
| Table-1 : Some typical CTUHSK generative parameter tuples GPT(n) |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |

## 12. A Challenge to the Cool-Headed Brave-Hearts

If you can prove that corresponding to every positive odd number  $m > 1$  there exists a unique valid set of CTUHSK generative parameters  $GPT(m) = \{p_0, p_1, p_2, \dots, p_k\}$  and therefore the corresponding valid set of exponents  $\{z, z_0, z_1, z_2, z_3, \dots, z_{k-1}, z_k\}$  in the parametric equation [Eqn.18] given above that generates every positive odd number  $m > 1$ , then you can directly prove the Collatz Conjecture establishing the convergence of the Collatz Sequence to the trivial cycle  $\{(4 \rightarrow 2 \rightarrow 1)\}$ .

## 12.1 Restrictions on the CTUHSK Generative Parameters

Note that the set of valid values for the CTUHSK generative parameters and therefore for the exponents in [Eqn.18] above, are governed by certain rules as can be seen from the earlier observations, regarding the matching relationship between the  $((D^\#) \text{MOD} 3)$  of the predecessor and the  $((U^\#) \text{MOD} 3)$  of the successor in the Collatz Sequence.

Specifically, [Eqn.19] states the relationship satisfied among (i) the  $[x] \text{MOD} 3$  value of the exponent  $x$  for  $U^\# = \{(6m-5).4^x\}$  at some higher rung in QEL(6m-5); (ii) with its defining-base-rung at (6m-5); and (iii) its predecessor  $D^\# = \lfloor \{(6m-5).4^x - 1\} / 3 \rfloor$ . Similarly, [Eqn.20] states the relationship satisfied among (i) the  $[y] \text{MOD} 3$  value of the exponent  $x$  for  $U^\# = \{(6m-1).2.4^y\}$  at some higher rung in QEL(6m-1); (ii) with its defining-base-rung at (6m-1); and (iii) its predecessor  $D^\# = \lfloor \{(6m-1).2.4^y - 1\} / 3 \rfloor$ .

$$\lfloor \{(6m-5).4^x - 1\} / 3 \rfloor \text{MOD} 3 = \lfloor [x] \text{MOD} 3 - [m] \text{MOD} 3 + 1 \rfloor \text{MOD} 3; \quad [\text{Eqn.19}]$$

and

$$\lfloor \{(6m-1).2.4^y - 1\} / 3 \rfloor \text{MOD} 3 = \lfloor [y] \text{MOD} 3 + [m] \text{MOD} 3 - 1 \rfloor \text{MOD} 3; \quad [\text{Eqn.20}]$$

Rewriting [Eqn.19] & [Eqn.20] for the Binary-Exponential-Ladders, we get the equivalent set of equations as:

$$\lfloor \{(6m-5).2^w - 1\} / 3 \rfloor \text{MOD} 3 = \lfloor [w/2] \text{MOD} 3 - [m] \text{MOD} 3 + 1 \rfloor \text{MOD} 3; \quad [\text{Eqn.21}]$$

and

$$\lfloor \{(6m-1).2^v - 1\} / 3 \rfloor \text{MOD} 3 = \lfloor [(v-1)/2] \text{MOD} 3 + [m] \text{MOD} 3 - 1 \rfloor \text{MOD} 3; \quad [\text{Eqn.22}]$$

## 12.2 Most Intriguing and Reassuring Observation

Suppose we move forward from  $H_{k-1}$  to  $H_k$  and then onto  $H_{k+1}$ ; that is, trace backwards along the Collatz Sequence, in the reverse direction; moving from  $BEL(n_{k-1})$  to  $BEL(n_k)$  and then onto  $BEL(n_{k+1})$ ; with  $BEL(n_{k-1}) = S(BEL(n_k))$  and  $BEL(n_k) = S(BEL(n_{k+1}))$ ; every time selecting the immediate-predecessor  $BEL$  of the *lowest denomination* with the *lowest value of the defining-base-rung*, that is, selecting  $BEL(n_{k-1}) \in H_{k-1}$ ; as represented by the lowest value of 2 for  $w$  in [Eqn.7] or the lowest value of 1 for  $v$  in [Eqn.8] above.

We find that [Eqn.7] with  $w=2$  leads to a situation wherein  $n_{k-1} < n_k$ ; that is,

$$BEL(6m-5) = S(BEL([(6m-5).2^2 - 1]/3)) = S(BEL(8m-7)); \quad [Eqn.23]$$

whereas [Eqn.8] with  $v=1$  leads to a situation wherein  $n_{k-1} > n_k$ ; that is,

$$BEL(6m-1) = S(BEL([(6m-1).2^1 - 1]/3)) = S(BEL(4m-1)); \quad [Eqn.24]$$

which incidentally the same as [Eqn.17] as seen earlier.

Thus, we find that moving forward from  $H_{k-1}$  to  $H_k$  and then onto  $H_{k+1}$ ; every time selecting the immediate-predecessor  $BEL$  of the lowest denomination, we encounter the three distinct situations:– (1)  $BEL(6m-3) \in H_{k-1}$  being a leaf node of the BELnet, does not have any predecessors; (2)  $BEL(6m-5) \in H_{k-1}$  has its immediate-predecessor  $BEL$  of lowest denomination being  $BEL(8m-7) \in H_k$ ; (3)  $BEL(6m-1) \in H_{k-1}$  has its immediate-predecessor  $BEL$  of lowest denomination being  $BEL(4m-1) \in H_k$ .

This establishes the fact that apart from the one-third of the BELs that lead to the leaf-nodes, the remaining two-thirds of the BELs are equally distributed between the two cases - BEL(6m-5) leads to the situation wherein the arborescence grows forward with increasing values for the defining-base-rung; whereas BEL(6m-1) leads to the situation wherein the arborescence grows forward with decreasing values for the defining-base-rung, thus *reaching-down* to some positive odd number of the form (4m-1) that was *left out in the earlier stages/tier-levels* of the hierarchy.

Note that there can be only three types of positive odd numbers corresponding to the three modular (nMOD3) residue classes; odd multiples of three that are of the form (6m-3) or numbers of the form (6m-5) or numbers of the form (6m-1); of which the odd multiples of three correspond to the defining-base-rung of the leaf nodes in the BELnet arborescence. Of the remaining two cases, it is of utmost significance that every positive odd number of the form (6m-1) forming the defining-base-rung of  $BEL(6m-1) \in H_{k-1}$  having its immediate-predecessor BEL of lowest denomination that is of the form  $BEL(4m-1) \in H_k$ . This is how as we move from  $H_{k-1}$  to  $H_k$  and then onto  $H_{k+1}$ ; we encounter some positive odd numbers of the form (4m-1) that *could not be reached in the earlier stages/tier-levels*; as indicated by the presence of 1 in the CTUHSK generative parameter tuple.

The CTUHSK generative parameter tuple corresponding to every positive odd integer of the form (4m-1) must necessarily end with the sub-sequence (... , 1, 0); as for example,  $GPT(11) = (4, 3, 2, 1, 0)$ ; whereas the presence of a positive odd integer of the form (4m-1) anywhere in the Collatz Sequence is indicated by the presence of a 1 in the corresponding position in its CTUHSK generative parameter tuple; as for example,  $GPT(9) = (4, 3, 2, 1, 1, 2, 0)$ ; with the Collatz Sequence for 9 is  $9 \rightarrow 7 \rightarrow 11 \rightarrow 17 \rightarrow 13 \rightarrow 5 \rightarrow 1$ ; the two repeated occurrence of 1s corresponding to

the sub-sequence 7->11->17 in the trajectory. It is indeed intriguing to observe that this specific process of reaching-down to such positive odd numbers of the form  $(4m-1)$  can indeed be a repeated contiguous operation at times, as can be seen by the presence of repeated 1's in the CTUHSK generative parameter tuple.

This is a situation wherein the emergence of global system properties through persistent local component subsystem characteristics can be clearly demonstrated! Banking on this global property, the author claims that the length of the CTUHSK generative parameter tuple  $|GPT(m)|$  required to generate a given positive odd number  $m$  is limited to be *no more than*  $m$  itself; whereas the triad  $|GPT(27)| = 42$  and  $|GPT(31)| = 40$  with  $|GPT(41)| = 41$  is an exception. Note that the triad  $|GPT(3)| = 3$  and  $|GPT(1)| = 1$  with  $|GPT(5)| = 2$  is another limiting case.

We will delve deeper into such algorithmic & computational details in a sequel to this research work, wherein a systematic (algorithmic) procedure for building the BELnet using the CTUHSK generative parameter tuples, corresponding to a systematic (algorithmic) procedure to produce a dictionary (lexicographic) of the CTUHSK generative parameter tuples, is planned to be presented.

### 13. Self-Similar Structural Symmetry of BELnet

From the above discussion one can notice that BELnet forms an arborescence in  $H^s$  with a self-similar structural symmetry.  $\{BEL(1)\}$  stands at the center, with its trivial cycle at its defining-base-rung. At every tier-level  $k$  corresponding to the  $k^{th}$  term  $H_k$  in the sequence  $H^s$ , Binary-Exponential-Ladders of all the three modular (MOD3) residue classes are present in BELnet, each being countably

infinite and equal in number; each having its single unique immediate-successor in  $H_{k-1}$ . Each of the  $\{BEL(6m-5)\}$  and each of the  $\{BEL(6m-1)\}$  has its immediate-predecessors in  $H_{k+1}$ ; whereas of each the  $\{BEL(6m-3)\}$  remain as leaf-nodes since they can't have any immediate-predecessors. Thus, one-third of the BELs remain as leaf nodes; the other two-thirds become intermediate nodes that propagate the arborescence structure unboundedly to infinity. The self-similar structural symmetry in the network of binary-exponential-ladders BELnet is a self-similar derived from the modular-periodicity w.r.t both the defining-base-rung value among the neighboring BELs as well as the binary-exponent value within any given BEL, as defined by [Eqn.19] & [Eqn.20] or equivalently [Eqn.21] & [Eqn.22] above.

Note that each of the  $\{BEL(6m-5)\}$  with  $(6m-5)$  as its defining-base-rung, has its immediate-predecessor of lowest denomination being  $\{BEL(8m-7)\}$  as defined in [Eqn.23] where  $(8m-7) > (6m-5)$ ; facilitating the outward growth of the BELnet arborescence by pushing upward in the number-line. Also note that each of the  $\{BEL(6m-1)\}$  with  $(6m-1)$  as its defining-base-rung, has its immediate-predecessor of lowest denomination being  $\{BEL(4m-1)\}$  as defined in [Eqn.24] where  $(4m-1) < (6m-1)$ ; facilitating the inward growth of the BELnet arborescence by pulling downward in reaching-down the not-yet-reached in the number-line (positive odd numbers). Therefore, we observe that the specific instance of [Eqn.8] represented by [Eqn.24]; in the inverse of the CTUHSK function; acts as the precise mechanism that systematically fills the gaps in the number line (positive odd numbers) that otherwise gets left out by [Eqn.7] or even [Eqn.23]; while constructing the lexicographic dictionary of CTUHSK generative parameter tuples for growing the BELnet arborescence.

This self-similar structural symmetry with the modular-periodicity (triadic/hexadic) both with respect to the defining-base-rung value and the binary-exponent value, along with the above mentioned unique feature of reaching-down the not-yet-reached numbers in the number-line, and using all the three modular (nMOD3) residue classes among the set of positive odd numbers as the defining-base-rung for the BELs, together guarantee the connectedness of the network of binary-exponential-ladders BELnet; which has of course been established above by showing the non-existence of any disjoint loops and/or divergent chains.

#### 14. Conclusion

We established a two stage bijective mapping from the Collatz-domain to a meticulously designed *structured system framework*  $H$  with a careful reorganization & condensation :- in the first stage, a bijective mapping from the Collatz-domain to BELnet, that is the network of Binary-Exponential-Ladders defined on the set of positive odd numbers; and then in the second stage, a bijective mapping from BELnet to the structured system framework  $H$ . Then we established the *order-preserving isomorphism* between the relevant component  $H^s$  (that is shown to be an infinite well-ordered sequence or rather a hierarchy or indeed an arborescence of the *Binary-Exponential-Ladders* having its root with an invariant base element  $H_0$ ) of the *structured system framework*  $H$  and the set of positive integers; providing the necessary condition. A sufficient condition is provided through a reductio-ad-absurdum argument (along with a uniquely special modular arithmetic characteristic property of the Collatz system) that establishes *domain exhaustion*; having already captured all the modular (MOD3) residue classes in  $H^s$ ; and guarantees the non-existence of any extraneous objects like disjoint loops/cycles



and/or divergent chains. Since the Collatz  $(3n+1)$  Conjecture is restricted to the set of natural numbers, the existence of any non-standard ‘rogue’ and/or ‘ghost’ elements are of no concern here.

As a direction towards future research work, we have also presented a possible alternative approach in proving this convergence, using modular arithmetic for the conditions to be satisfied by the CTUHSK generative parameters or equivalently the exponents in a closed-form expression for generating the reverse Collatz sequence corresponding to any given positive odd number.

It is observed that the consistent & persistent application of the modular arithmetic characteristics associated with the local neighborhood of BELnet imposed by the CTUHSK function gives rise to the emergence of the global system property associated with the structured system framework that forms an arborescence (hierarchy) of BELnet – this leads to an upper limit on the depth in the arborescence or the rank/level in the hierarchy that one needs to analyse in order to reach for the existence of a binary-exponential-ladder with the desired positive odd number as its defining base rung in BELnet.

## 15. Recommended Reading

- [1]. Wikipedia Page – [https://en.wikipedia.org/wiki/Collatz\\_conjecture](https://en.wikipedia.org/wiki/Collatz_conjecture)
- [2]. Richard K Guy;  
“Don’t Try to Solve These Problems”;  
The American Mathematical Monthly: Vol. 90, No. 1, pp. 35-41; 1983.

<https://doi.org/10.1080/00029890.1983.11971148>

- [3]. Jeffrey C Lagarias;  
“The  $3x+1$  problem: An annotated bibliography (1963--1999)  
(sorted by author)”;  
<https://arxiv.org/abs/math/0309224>
- [4]. Jeffrey C Lagarias;  
“The  $3x+1$  Problem: An Annotated Bibliography, II (2000-2009)”;  
<https://arxiv.org/abs/math/0608208>
- [5]. Jeffrey C Lagarias;  
“The  $3x + 1$  Problem: An Overview”  
<https://arxiv.org/abs/2111.02635>
- [6]. Jeffrey C Lagarias;  
“The Ultimate Challenge: The  $3x+1$  Problem”;  
American Mathematical Society; 2010.

## 16. Declaration Regarding Affiliation and Funding

I, Dr(Prof) Keshava Prasad Halemane, hereby declare that I am a Professor retired as on 2017JAN31 from National Institute of Technology Karnataka Surathkal India, and I am not affiliated to any institution or organization or corporation or any other agency or whatever. This research work has been conducted entirely by me on my own as an Independent Researcher, and that I have not received any funding from any source other than my own savings, and I do not have any obligations or encumbrances of any kind, neither financial nor legal nor of any other kind, regarding the contents of the manuscript - of which I am the original author and creator. Also, I hereby declare there is no conflict of interests.

## 17. Author Rights Retention Notice

The author as the original creator of this work asserts exclusive ownership and retains full-&-complete Intellectual Property Rights (IPR) and CopyRight for this (as well as all its previous versions, including all the versions archived in various repositories, all of them with CC-BY-NC-ND 4.0 International Public Copyright License) and any/every Author Accepted Manuscript (AAM) version arising from this manuscript) in perpetuity, for the maximum duration permitted by law. For the purpose of publication, the author hereby grants a non-exclusive publishing rights to the publisher. The author's IPR including all commercial and derivative rights are retained in full as the exclusive intellectual property of the author and shall pass to his designated legal heirs (the three children mentioned herein below) in perpetuity, for the maximum duration permitted by law.

## 18. Dedication

To my ಅಜ್ಜ(ajja) Karinja Halemane Keshava Bhat & ಅಜ್ಜಿ(ajji) Thirumaleshwari, ಅಪ್ಪ(appa) Shama Bhat & ಅಮ್ಮ(amma) Thirumaleshwari, for their *teachings through love, that quality matters more than quantity*; to my wife Vijayalakshmi for her *ever consistent love & support*; to my daughter [Sriwidya.Bharati](#) and my twin sons [Sriwidya.Ramana](#) & [Sriwidya.Prawina](#) for their *love & affection*.