



# GATE WALLAH

→ TOPICWISE ←  
**GATE PREVIOUS  
YEAR QUESTIONS**

2008 – 2023

**ENGINEERING MATHEMATICS &  
GENERAL APTITUDE**

## **EDITION: First**

**Published By:**



**ISBN:** 978-81-19192-70-0

**Mobile App:** Physics Wallah (Available on Play Store)



**Website:** [www.pw.live](http://www.pw.live)

**Youtube Channel:** GATE Wallah (@GATEWallahbyPW)

GATE Wallah English (@GateWallahEnglish)

GATE Wallah - EE, EC & CS (@GATEWallah\_EE\_EC\_CS)

GATE Wallah - ME, CE & XE (@GATEWallah\_ME\_CE\_XE)

**Email:** support@pw.live

## **Rights**

All rights are reserved with the Publisher. No part of this book may be used or reproduced in any manner whatsoever without written permission from the author or publisher.

In the interest of the student community:

Circulation of soft copy of Book(s) in PDF or other equivalent format(s) through any social media channels, emails, etc. or any other channels through mobiles, laptops or desktop is a criminal offense. Anybody circulating, downloading, or storing, a soft copy of the book on his device(s) is in breach of the Copyright Act. Further photocopying of this book or any of its material is also illegal. Do not download or forward in case you come across any such soft copy material.

## **Disclaimer**

A team of PW GATE Wallah experts and faculties with a deep understanding of the subject has worked hard for the creation and curation of this book.

While the content creators, editors and publisher have used their best efforts in preparing these books. The content has been checked for accuracy. As the book is intended for educational purposes, the author shall not be responsible for any errors contained in the book.

The publication has designed the content to provide accurate and authoritative information with regard to the subject matter covered.

This book and the individual contribution contained in it are protected under Copyright Act by the publisher.

*(This book shall only be used for educational purposes.)*



# PREFACE

A highly skilled professional team of GATE Wallah works arduously to ensure that the students receive the best content for their **GATE** exam.

A plethora of GATE Study Material is available in the market but GATE Wallah professionals at PW are continuously working to provide supreme quality study material for the GATE students.

From the beginning, the content team comprising Subject Matter Experts, Content Creators, Reviewers, DTP operators, Proofreaders, and others is involved in shaping the material to their best knowledge and experience to produce powerful content for the students.

GATE Wallah Faculties have adopted a novel style of presenting the content in easy-to-understand language and have provided the content team with expert guidance and supervision throughout the creation and curation of this book.

PW's GATE Wallah strongly believes in conceptual and fun-based learning. GATE Wallah provides highly exam-oriented content to bring quality and clarity to the students.

This book adopts a multi-faceted approach to mastering and understanding the concepts by having a rich diversity of questions asked in the examination and equipping the students with the knowledge for this highly competitive exam.

The main objective of this book is to provide an edge to your preparation with high-quality content & video solutions.

## BOOK FEATURES

This book, especially designed for GATE aspirants, contains

- A topic-wise set of questions from the past 16 years (2008-2023)
- Embedded QR codes for Video Solutions (by expert faculties)
- Complete explanations to help you ace your exam
- Detailed Chapter Wise Analysis of PYQs
- Covers Engineering Mathematics and Aptitude based PYQs from the 5 branches:
  - Mechanical Engineering
  - Civil Engineering
  - Electrical Engineering
  - Electronics & Communication Engineering
  - Computer Science & Information Technology

## SPECIAL ABOUT THIS BOOK

We take pride in providing high-quality study material for the **GATE** exam. Our team at **GATE WALLAH** has meticulously prepared each solution to ensure they are error-free and easy to understand.

The **GATE** exam pattern has shifted in recent years, with an increased emphasis on topics that previously received fewer questions. At **GATE WALLAH**, we understand the importance of identifying these crucial topics, and that's why we've provided a chapter-wise analysis for the last 16 years. Our analysis will help you understand the significance of each chapter and focus your preparation accordingly. We believe that this will be a valuable resource in your journey to success in the **GATE** exam.

To ensure that you focus on the most relevant questions, we have arranged the questions in this book topic-wise and year-wise in descending order. This means that the questions from the most recent years are emphasized, as we believe they are the most relevant for your **GATE** exam preparation. By studying these questions, you'll gain a deeper understanding of the exam pattern and the type of questions asked in recent years, which will help you perform better in the upcoming **GATE** exam.

We've designed the text solutions in this book to be in sync with our video solutions recorded by expert faculties, making it easier for you to understand the concepts. We understand that every student has a unique learning style, and that's why we've included both text and video solutions to cater to your needs. By using our comprehensive guide and video solutions together, you'll gain a better understanding of the concepts and be better prepared to tackle the **GATE** exam.

### Steps to Open Video solutions through mobile:

- (1) Scan the given embedded QR Code for a particular solution.



- (2) Visit the link generated & you'll be redirect to the video solution.



# GATE

## ENGINEERING MATHEMATICS & GENERAL APTITUDE

### CONTENTS

|    |                               |             |
|----|-------------------------------|-------------|
| 1. | Engineering Mathematics ..... | 1.1 – 1.359 |
| 2. | General Aptitude .....        | 2.1 – 2.258 |





# Engineering Mathematics

|    |                                  |               |
|----|----------------------------------|---------------|
| 1. | Linear Algebra .....             | 1.1 – 1.66    |
| 2. | Basic Calculus .....             | 1.67 – 1.152  |
| 3. | Probability and Statistics ..... | 1.153 – 1.206 |
| 4. | Differential Equations.....      | 1.207 – 1.258 |
| 5. | Vector Calculus.....             | 1.259 – 1.290 |
| 6. | Complex Calculus .....           | 1.291 – 1.318 |
| 7. | Fourier Series .....             | 1.319 – 1.325 |
| 8. | Laplace Transforms .....         | 1.326 – 1.335 |
| 9. | Numerical Methods.....           | 1.336 – 1.359 |



# Engineering Mathematics

## Syllabus

(Mechanical Engineering)

- **Linear Algebra:** Matrix algebra, systems of linear equations, eigenvalues and eigenvectors.
- **Calculus:** Functions of single variable, limit, continuity and differentiability, mean value theorems, indeterminate forms; evaluation of definite and improper integrals; double and triple integrals; partial derivatives, total derivative, Taylor series (in one and two variables), maxima and minima, Fourier series; gradient, divergence and curl, vector identities, directional derivatives, line, surface and volume integrals, applications of Gauss, Stokes and Green's theorems.
- **Differential Equations:** First order equations (linear and nonlinear); higher order linear differential equations with constant coefficients; Euler-Cauchy equation; initial and boundary value problems; Laplace transforms; solutions of heat, wave and Laplace's equations.
- **Complex Variables:** Analytic functions; Cauchy-Riemann equations; Cauchy's integral theorem and integral formula; Taylor and Laurent series.
- **Probability and Statistics:** Definitions of probability, sampling theorems, conditional probability; mean, median, mode and standard deviation; random variables, binomial, Poisson and normal distributions.
- **Numerical Methods:** Numerical solutions of linear and non-linear algebraic equations; integration by trapezoidal and Simpson's rules; single and multi-step methods for differential equations.

## ME Chapter wise Weightage Analysis (Marks)

| <b>Chapter<br/>Paper<br/>Year</b> | <b>Ch.1</b> | <b>Ch.2</b> | <b>Ch.3</b> | <b>Ch.4</b> | <b>Ch.5</b> | <b>Ch.6</b> | <b>Ch.7</b> | <b>Ch.8</b> | <b>Ch.9</b> |
|-----------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>2008</b>                       | 5           | 10          | 1           | 3           | 3           | 2           | 0           | 0           | 0           |
| <b>2009</b>                       | 1           | 6           | 3           | 2           | 1           | 2           | 0           | 1           | 0           |
| <b>2010</b>                       | 2           | 3           | 2           | 1           | 2           | 1           | 0           | 2           | 2           |
| <b>2011</b>                       | 3           | 3           | 2           | 2           | 0           | 1           | 0           | 0           | 2           |
| <b>2012</b>                       | 4           | 4           | 2           | 2           | 1           | 0           | 0           | 2           | 0           |
| <b>2013</b>                       | 1           | 4           | 3           | 4           | 0           | 0           | 0           | 2           | 1           |
| <b>2014 (P1)</b>                  | 4           | 3           | 2           | 0           | 3           | 1           | 0           | 0           | 0           |
| <b>2014 (P2)</b>                  | 3           | 4           | 1           | 2           | 1           | 0           | 0           | 0           | 2           |
| <b>2014 (P3)</b>                  | 0           | 1           | 4           | 3           | 0           | 2           | 0           | 1           | 2           |
| <b>2014 (P4)</b>                  | 0           | 2           | 3           | 1           | 0           | 3           | 0           | 0           | 4           |
| <b>2015 (P1)</b>                  | 1           | 5           | 3           | 2           | 0           | 1           | 0           | 0           | 1           |
| <b>2015 (P2)</b>                  | 1           | 3           | 3           | 2           | 1           | 0           | 0           | 1           | 2           |
| <b>2015 (P3)</b>                  | 3           | 3           | 1           | 0           | 1           | 0           | 0           | 2           | 3           |
| <b>2016 (P1)</b>                  | 1           | 2           | 1           | 2           | 0           | 3           | 0           | 1           | 3           |
| <b>2016 (P2)</b>                  | 1           | 3           | 2           | 0           | 0           | 3           | 0           | 1           | 3           |
| <b>2016 (P3)</b>                  | 3           | 5           | 3           | 1           | 0           | 0           | 0           | 0           | 1           |
| <b>2017 (P1)</b>                  | 3           | 3           | 1           | 2           | 2           | 0           | 0           | 0           | 2           |
| <b>2017 (P2)</b>                  | 3           | 2           | 2           | 2           | 1           | 2           | 0           | 1           | 0           |
| <b>2018 (P1)</b>                  | 1           | 3           | 4           | 0           | 0           | 1           | 0           | 2           | 2           |
| <b>2018 (P2)</b>                  | 1           | 0           | 2           | 4           | 3           | 2           | 1           | 0           | 0           |
| <b>2019 (P1)</b>                  | 3           | 3           | 3           | 1           | 0           | 2           | 0           | 0           | 1           |
| <b>2019 (P2)</b>                  | 1           | 2           | 3           | 3           | 3           | 1           | 0           | 0           | 0           |
| <b>2020 (P1)</b>                  | 1           | 3           | 3           | 0           | 1           | 3           | 0           | 1           | 2           |
| <b>2020 (P2)</b>                  | 2           | 1           | 3           | 0           | 2           | 2           | 0           | 1           | 2           |
| <b>2021 (P1)</b>                  | 2           | 3           | 1           | 1           | 0           | 2           | 0           | 1           | 1           |
| <b>2021 (P2)</b>                  | 3           | 2           | 1           | 2           | 0           | 1           | 0           | 1           | 3           |
| <b>2022 (P1)</b>                  | 3           | 2           | 2           | 0           | 2           | 2           | 1           | 1           | <b>0</b>    |
| <b>2022 (P2)</b>                  | 4           | 2           | 0           | 2           | 1           | 2           | 1           | 0           | 1           |
| <b>2023</b>                       | 0           | 4           | 1           | 2           | 1           | 1           | 0           | 2           | 2           |



# Engineering Mathematics

## Syllabus

(Civil Engineering)

---

---

- **Linear Algebra:** Matrix algebra, systems of linear equations; Eigen values and Eigen vectors.
- **Calculus:** Functions of single variable; Limit, continuity and differentiability; Mean value theorems, local maxima and minima; Taylor series; Evaluation of definite and indefinite integrals, application of definite integral to obtain area and volume; Partial derivatives; Total derivative; Gradient, Divergence and Curl, Vector identities; Directional derivatives; Line, Surface and Volume integrals.
- **Ordinary Differential Equation (ODE):** First order (linear and non-linear) equations; higher order linear equations with constant coefficients; Euler-Cauchy equations; initial and boundary value problems.
- **Partial Differential Equation (PDE):** Fourier series; separation of variables; solutions of one-dimensional diffusion equation; first and second order one-dimensional wave equation and two-dimensional Laplace equation.
- **Probability and Statistics:** Sampling theorems; Conditional probability; Descriptive statistics - Mean, median, mode and standard deviation; Random Variables - Discrete and Continuous, Poisson and Normal Distribution; Linear regression.
- **Numerical Methods:** Error analysis. Numerical solutions of linear and non-linear algebraic equations; Newton's and Lagrange polynomials; numerical differentiation; Integration by trapezoidal and Simpson's rule; Single and multi-step methods for first order differential equations.

## CE Chapter wise Weightage Analysis (Marks)

| <b>Chapter<br/>Paper<br/>Year</b> | <b>Ch.1</b> | <b>Ch.2</b> | <b>Ch.3</b> | <b>Ch.4</b> | <b>Ch.5</b> | <b>Ch.6</b> | <b>Ch.7</b> | <b>Ch.8</b> | <b>Ch.9</b> |
|-----------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>2008</b>                       | 5           | 4           | 6           | 3           | 2           | 0           | 0           | 0           | 0           |
| <b>2009</b>                       | 1           | 0           | 2           | 2           | 3           | 3           | 0           | 2           | 2           |
| <b>2010</b>                       | 2           | 7           | 1           | 3           | 0           | 0           | 0           | 0           | 2           |
| <b>2011</b>                       | 1           | 4           | 1           | 2           | 2           | 2           | 0           | 0           | 1           |
| <b>2012</b>                       | 2           | 1           | 3           | 2           | 2           | 0           | 0           | 0           | 3           |
| <b>2013</b>                       | 1           | 2           | 2           | 2           | 0           | 0           | 0           | 0           | 2           |
| <b>2014 (P1)</b>                  | 2           | 4           | 3           | 2           | 0           | 0           | 0           | 0           | 0           |
| <b>2014 (P2)</b>                  | 3           | 2           | 2           | 1           | 0           | 1           | 0           | 0           | 0           |
| <b>2015 (P1)</b>                  | 3           | 0           | 1           | 2           | 2           | 2           | 0           | 0           | 3           |
| <b>2015 (P2)</b>                  | 3           | 2           | 2           | 2           | 0           | 1           | 0           | 0           | 3           |
| <b>2016 (P1)</b>                  | 1           | 4           | 3           | 4           | 0           | 0           | 0           | 0           | 1           |
| <b>2016 (P2)</b>                  | 2           | 8           | 4           | 0           | 0           | 0           | 1           | 0           | 0           |
| <b>2017 (P1)</b>                  | 3           | 2           | 3           | 3           | 0           | 0           | 0           | 0           | 2           |
| <b>2017 (P2)</b>                  | 3           | 5           | 1           | 2           | 1           | 0           | 0           | 0           | 0           |
| <b>2018 (P1)</b>                  | 2           | 3           | 0           | 4           | 0           | 0           | 0           | 0           | 2           |
| <b>2018 (P2)</b>                  | 4           | 0           | 3           | 1           | 2           | 0           | 0           | 2           | 1           |
| <b>2019 (P1)</b>                  | 0           | 6           | 0           | 2           | 0           | 0           | 0           | 0           | 2           |
| <b>2019 (P2)</b>                  | 3           | 1           | 2           | 2           | 2           | 0           | 0           | 1           | 1           |
| <b>2020 (P1)</b>                  | 2           | 2           | 0           | 3           | 2           | 0           | 0           | 0           | 3           |
| <b>2020 (P2)</b>                  | 2           | 1           | 1           | 4           | 0           | 0           | 2           | 0           | 1           |
| <b>2021 (P1)</b>                  | 2           | 2           | 1           | 2           | 0           | 0           | 0           | 0           | 4           |
| <b>2021 (P2)</b>                  | 4           | 2           | 0           | 2           | 3           | 0           | 0           | 0           | 2           |
| <b>2022 (P1)</b>                  | 1           | 5           | 2           | 1           | 0           | 0           | 1           | 0           | 3           |
| <b>2022 (P2)</b>                  | 4           | 3           | 2           | 2           | 1           | 0           | 0           | 0           | <b>0</b>    |
| <b>2023 (P1)</b>                  | 3           | 3           | 1           | 1           | 0           | 0           | 1           | 0           | 4           |
| <b>2023 (P2)</b>                  | 5           | 0           | 2           | 5           | 1           | 0           | 0           | 0           | <b>0</b>    |



# Engineering Mathematics

## Syllabus

(Electrical Engineering)

- **Linear Algebra:** Matrix Algebra, Systems of linear equations, Eigenvalues, Eigenvectors.
- **Calculus:** Mean value theorems, Theorems of integral calculus, Evaluation of definite and improper integrals, Partial Derivatives, Maxima and minima, Multiple integrals, Fourier series, Vector identities, Directional derivatives, Line integral, Surface integral, Volume integral, Stokes's theorem, Gauss's theorem, Divergence theorem, Green's theorem.
- **Differential Equations:** First order equations (linear and nonlinear), Higher order linear differential equations with constant coefficients, Method of variation of parameters, Cauchy's equation, Euler's equation, Initial and boundary value problems, Partial Differential Equations, Method of separation of variables.
- **Complex Variables:** Analytic functions, Cauchy's integral theorem, Cauchy's integral formula, Taylor series, Laurent series, Residue theorem, Solution integrals.
- **Probability and Statistics:** Sampling theorems, Conditional probability, Mean, Median, Mode, Standard Deviation, Random variables, Discrete and Continuous distributions, Poisson distribution, Normal distribution, Binomial distribution, Correlation analysis, Regression analysis.

## EE Chapter wise Weightage Analysis (Marks)

| <b>Chapter<br/>Paper Year</b> | <b>Ch.1</b> | <b>Ch.2</b> | <b>Ch.3</b> | <b>Ch.4</b> | <b>Ch.5</b> | <b>Ch.6</b> | <b>Ch.7</b> | <b>Ch.8</b> | <b>Ch.9</b> |
|-------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>2008</b>                   | 6           | 2           | 1           | 2           | 0           | 0           | 0           | 2           | 6           |
| <b>2009</b>                   | 1           | 4           | 2           | 0           | 2           | 0           | 0           | 2           | 1           |
| <b>2010</b>                   | 4           | 3           | 2           | 2           | 1           | 0           | 0           | 0           | 4           |
| <b>2011</b>                   | 2           | 3           | 0           | 1           | 2           | 1           | 0           | 2           | 2           |
| <b>2012</b>                   | 2           | 2           | 3           | 3           | 2           | 2           | 1           | 0           | 2           |
| <b>2013</b>                   | 3           | 2           | 1           | 0           | 2           | 3           | 0           | 2           | 3           |
| <b>2014 (P1)</b>              | 3           | 3           | 2           | 1           | 2           | 1           | 1           | 0           | 3           |
| <b>2014 (P2)</b>              | 1           | 5           | 3           | 1           | 0           | 1           | 0           | 0           | 1           |
| <b>2014 (P3)</b>              | 1           | 0           | 3           | 0           | 2           | 2           | 0           | 2           | 1           |
| <b>2015 (P1)</b>              | 3           | 1           | 3           | 2           | 1           | 0           | 0           | 0           | 3           |
| <b>2015 (P2)</b>              | 1           | 2           | 2           | 2           | 0           | 1           | 0           | 0           | 1           |
| <b>2016 (P1)</b>              | 5           | 3           | 2           | 1           | 0           | 1           | 0           | 0           | 5           |
| <b>2016 (P2)</b>              | 3           | 2           | 2           | 0           | 1           | 3           | 3           | 0           | 3           |
| <b>2017 (P1)</b>              | 1           | 3           | 0           | 2           | 0           | 3           | 0           | 2           | 1           |
| <b>2017 (P2)</b>              | 2           | 3           | 4           | 0           | 0           | 2           | 0           | 2           | 2           |
| <b>2018</b>                   | 3           | 4           | 0           | 0           | 3           | 3           | 0           | 0           | 3           |
| <b>2019</b>                   | 4           | 0           | 1           | 1           | 3           | 3           | 1           | 0           | 4           |
| <b>2020</b>                   | 2           | 4           | 0           | 1           | 4           | 1           | 0           | 0           | 2           |
| <b>2021</b>                   | 3           | 4           | 2           | 0           | 2           | 1           | 0           | 0           | 3           |
| <b>2022</b>                   | 5           | 2           | 2           | 0           | 6           | 0           | 0           | 0           | 5           |
| <b>2023</b>                   | 1           | 3           | 4           | 0           | 3           | 0           | 0           | 0           | 1           |



# Engineering Mathematics

(Electronic and Communication Engineering)

## Syllabus

---

---

- **Linear Algebra:** Vector space, basis, linear dependence and independence, matrix algebra, Eigen values and eigen vectors, rank, solution of linear equations- existence and uniqueness.
- **Calculus:** Mean value theorems, theorems of integral calculus, evaluation of definite and improper integrals, partial derivatives, maxima and minima, multiple integrals, line, surface and volume integrals, Taylor series.
- **Differential Equations:** First order equations (linear and nonlinear), higher order linear differential equations, Cauchy's and Euler's equations, methods of solution using variation of parameters, complementary function and particular integral, partial differential equations, variable separable method, initial and boundary value problems.
- **Vector Analysis:** Vectors in plane and space, vector operations, gradient, divergence and curl, Gauss's, Green's and Stokes' theorems.
- **Complex Analysis:** Analytic functions, Cauchy's integral theorem, Cauchy's integral formula, sequences, series, convergence tests, Taylor and Laurent series, residue theorem.
- **Probability and Statistics:** Mean, median, mode, standard deviation, combinatorial probability, probability distributions, binomial distribution, Poisson distribution, exponential distribution, normal distribution, joint and conditional probability.

## ECE Chapter wise Weightage Analysis (Marks)

| Chapter<br>Paper Year \ Ch. | Ch.1 | Ch.2 | Ch.3 | Ch.4 | Ch.5 | Ch.6 | Ch.7 | Ch.8 | Ch.9     |
|-----------------------------|------|------|------|------|------|------|------|------|----------|
| <b>2008</b>                 | 4    | 4    | 0    | 1    | 4    | 3    | 2    | 4    | 4        |
| <b>2009</b>                 | 2    | 2    | 1    | 3    | 2    | 1    | 0    | 2    | 2        |
| <b>2010</b>                 | 1    | 2    | 2    | 1    | 2    | 2    | 2    | 1    | 2        |
| <b>2011</b>                 | 2    | 0    | 2    | 1    | 0    | 1    | 2    | 2    | 0        |
| <b>2012</b>                 | 2    | 2    | 2    | 1    | 0    | 2    | 0    | 2    | 2        |
| <b>2013</b>                 | 3    | 1    | 0    | 0    | 0    | 0    | 0    | 3    | 1        |
| <b>2014 (P1)</b>            | 4    | 4    | 1    | 0    | 0    | 1    | 0    | 4    | 4        |
| <b>2014 (P2)</b>            | 5    | 2    | 0    | 1    | 2    | 2    | 0    | 5    | 2        |
| <b>2014 (P3)</b>            | 2    | 4    | 3    | 1    | 0    | 0    | 1    | 2    | 4        |
| <b>2014 (P4)</b>            | 0    | 3    | 4    | 3    | 2    | 0    | 0    | 0    | 3        |
| <b>2015 (P1)</b>            | 2    | 5    | 1    | 2    | 0    | 1    | 0    | 2    | 5        |
| <b>2015 (P2)</b>            | 1    | 1    | 2    | 3    | 0    | 2    | 0    | 1    | 1        |
| <b>2015 (P3)</b>            | 1    | 1    | 2    | 2    | 0    | 1    | 2    | 1    | 1        |
| <b>2016 (P1)</b>            | 3    | 3    | 1    | 1    | 2    | 2    | 0    | 3    | 3        |
| <b>2016 (P2)</b>            | 1    | 2    | 0    | 2    | 3    | 1    | 0    | 1    | 2        |
| <b>2016 (P3)</b>            | 3    | 3    | 1    | 3    | 0    | 3    | 0    | 3    | 3        |
| <b>2017 (P1)</b>            | 2    | 3    | 1    | 1    | 1    | 0    | 1    | 2    | 3        |
| <b>2017 (P2)</b>            | 1    | 2    | 1    | 1    | 2    | 2    | 0    | 1    | 2        |
| <b>2018</b>                 | 2    | 4    | 1    | 4    | 0    | 2    | 0    | 2    | 4        |
| <b>2019</b>                 | 1    | 3    | 1    | 3    | 2    | 2    | 0    | 1    | 3        |
| <b>2020</b>                 | 3    | 1    | 0    | 3    | 3    | 0    | 0    | 3    | 1        |
| <b>2021</b>                 | 3    | 0    | 2    | 1    | 1    | 2    | 0    | 3    | <b>0</b> |
| <b>2022</b>                 | 3    | 4    | 1    | 2    | 3    | 1    | 0    | 3    | 4        |
| <b>2023</b>                 | 4    | 0    | 0    | 0    | 5    | 2    | 0    | 4    | <b>0</b> |

# Engineering Mathematics

(Computer Science & IT)

## Syllabus

- Linear Algebra:** Matrices, determinants, system of linear equations, eigenvalues and eigenvectors, LU decomposition.
- Calculus:** Limits, continuity and differentiability. Maxima and minima. Mean value theorem. Integration.
- Probability and Statistics:** Random variables. Uniform, normal, exponential, Poisson and binomial distributions. Mean, median, mode and standard deviation. Conditional probability and Bayes theorem.

### CSE & IT Chapter wise Weightage Analysis

| Chapter<br>Paper Year | Ch.1 | Ch.2 | Ch.3 |
|-----------------------|------|------|------|
| 2008                  | 3    | 4    | 6    |
| 2009                  | 0    | 2    | 0    |
| 2010                  | 2    | 1    | 4    |
| 2011                  | 2    | 2    | 6    |
| 2012                  | 1    | 1    | 3    |
| 2013                  | 0    | 1    | 1    |
| 2014 (P1)             | 3    | 3    | 1    |
| 2014 (P2)             | 2    | 0    | 0    |
| 2014 (P3)             | 1    | 2    | 2    |
| 2015 (P1)             | 3    | 2    | 0    |
| 2015 (P2)             | 1    | 2    | 0    |
| 2015 (P3)             | 1    | 4    | 0    |
| 2016 (P1)             | 1    | 1    | 3    |
| 2016 (P2)             | 2    | 0    | 1    |
| 2017 (P1)             | 3    | 2    | 2    |
| 2017 (P2)             | 3    | 1    | 5    |
| 2018                  | 2    | 1    | 1    |
| 2019                  | 3    | 1    | 4    |
| 2020                  | 0    | 1    | 2    |
| 2021 (P1)             | 1    | 2    | 3    |
| 2021 (P2)             | 1    | 1    | 1    |
| 2022                  | 3    | 1    | 0    |
| 2023                  | 2    | 2    | 0    |

# CHAPTER

1

# **LINEAR ALGEBRA**

## Algebra of Matrices

- 1. [MCQ] [GATE-CE-2023:2M]**

Cholesky decomposition is carried out on the following square matrix [A].

$$[A] = \begin{bmatrix} 8 & -5 \\ -5 & a_{22} \end{bmatrix}$$

Let  $l_{ij}$  and  $a_{ij}$  be the  $(i,j)^{\text{th}}$  elements of matrices  $[L]$  and  $[A]$ , respectively. If the element  $l_{22}$  of the decomposed lower triangular matrix  $[L]$  is 1.968, what is the value (rounded off to the nearest integer) of the element  $a_{22}$ ?




2. [MCQ] [GATE-EE-2023:1M]

For a given vector  $w = [1 \ 2 \ 3]^T$ , the vector normal to the plane defined by  $w^T x = 1$  is

- (a)  $[-2 -2 2]^T$       (b)  $[3 0 -1]^T$   
 (c)  $[3 2 1]^T$       (d)  $[1 2 3]^T$

- 3. [MCQ] [GATE-EE-2023:1M]**

In the figure, the vectors  $u$  and  $v$  are related as:  $Au = v$  by a transformation matrix  $A$ . The correct choice of  $A$  is



- $$(a) \begin{bmatrix} \frac{4}{5} & \frac{3}{5} \\ -\frac{3}{5} & \frac{4}{5} \end{bmatrix} \quad (b) \begin{bmatrix} \frac{4}{5} & -\frac{3}{5} \\ \frac{3}{5} & \frac{4}{5} \end{bmatrix}$$

- $$(c) \begin{bmatrix} \frac{4}{5} & \frac{3}{5} \\ \frac{5}{5} & \frac{5}{5} \\ \frac{3}{5} & \frac{4}{5} \\ \hline \frac{5}{5} & \frac{5}{5} \end{bmatrix} \quad (d) \begin{bmatrix} \frac{4}{5} & \frac{-3}{5} \\ \frac{5}{5} & \frac{5}{5} \\ \frac{3}{5} & \frac{-4}{5} \\ \hline \frac{5}{5} & \frac{5}{5} \end{bmatrix}$$

4. [MCQ] [GATE-CS-2023:2M]

$$\text{Let } A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 4 & 1 & 2 & 3 \\ 3 & 4 & 1 & 2 \\ 2 & 3 & 4 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 3 & 4 & 1 & 2 \\ 4 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \end{bmatrix}$$

Let  $\det(A)$  and  $\det(B)$  denote the determinants of the matrices A and B, respectively. Which one of the options given below is TRUE?

- (a)  $\det(A) = \det(B)$
  - (b)  $\det(B) = -\det(A)$
  - (c)  $\det(A) = 0$
  - (d)  $\det(AB) = \det(A) + \det(B)$

5. [MCQ] [GATE-CS-2022:1M]

Consider the following two statements  
the matrices  $A_{m \times n}$ ,  $B_{n \times m}$ ,  $C_{n \times n}$  and  $D_{n \times n}$ .

Statement 1:  $\text{tr}(AB) \equiv \text{tr}(BA)$

Statement 2:  $\text{tr}(\mathbf{CD}) \equiv \text{tr}(\mathbf{DC})$

Where  $\text{tr}()$  represents the trace of a matrix. Which one of the following holds?

- (a) Statement 1 is correct and Statement 2 is wrong.
  - (b) Statement 1 is wrong and Statement 2 is correct.
  - (c) Both Statement 1 and Statement 2 are correct.
  - (d) Both Statement 1 and Statement 2 are wrong.

**6. [MCQ]****[GATE-CE-2022:2M]**

The Cartesian coordinates of a point P in a right handed coordinate system are (1, 1, 1). The transformed coordinates of P due to a  $45^\circ$  clockwise rotation of the coordinate system about the positive x-axis are

- (a) (1, 0,  $\sqrt{2}$ )      (b) (1, 0,  $-\sqrt{2}$ )  
 (c) (-1, 0,  $\sqrt{2}$ )      (d) (-1, 0,  $-\sqrt{2}$ )

**7. [NAT]****[GATE-CE-2022:1M]**

The components of pure shear strain in a sheared

material are given in the matrix form  $E = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$

Here, Trace (E) = 0, Given, P = Trace (E<sup>8</sup>) and Q = Trace (E<sup>11</sup>)

The numerical value of (P + Q) is \_\_\_\_\_ (in integer)

**8. [MCQ]****[GATE-ME-2022: 1M]**

If  $A = \begin{bmatrix} 10 & 2k+5 \\ 3k-3 & k+5 \end{bmatrix}$  is a symmetric matrix, the value of k is \_\_\_\_\_.

- (a) 8      (b) 5  
 (c) -0.4      (d)  $\frac{1+\sqrt{1561}}{12}$

**9. [MCQ]****[GATE-CE-2021:2M]**

If  $P = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$  and  $Q = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  then  $Q^T P^T$  is

- (a)  $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$       (b)  $\begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$   
 (c)  $\begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}$       (d)  $\begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}$

**10. [MCQ]****[GATE-CE-2021:1M]**

If A is a square matrix then orthogonality property mandates

- (a)  $AA^T = I$       (b)  $AA^T = 0$   
 (c)  $AA^T = A^{-1}$       (d)  $AA^T = A^2$

**11. [MCQ]****[GATE-ME-2020: 1M]**

Multiplication of real valued square matrices of same dimension is

- (a) associative  
 (b) commutative  
 (c) always positive definite  
 (d) not always possible to compute

**12. [MCQ]****[GATE-ME-2020: 1M]**

A matrix P is decomposed into its symmetric part S and skew symmetric part V. If

$$S = \begin{pmatrix} -4 & 4 & 2 \\ 4 & 3 & 7/2 \\ 2 & 7/2 & 2 \end{pmatrix}, V = \begin{pmatrix} 0 & -2 & 3 \\ 2 & 0 & 7/2 \\ -3 & -7/2 & 0 \end{pmatrix}$$

then matrix P is

- (a)  $\begin{pmatrix} -4 & 6 & -1 \\ 2 & 3 & 0 \\ 5 & 7 & 2 \end{pmatrix}$   
 (b)  $\begin{pmatrix} -4 & 2 & 5 \\ 6 & 3 & 7 \\ -1 & 0 & 2 \end{pmatrix}$   
 (c)  $\begin{pmatrix} -4 & -6 & 1 \\ -2 & -3 & 0 \\ -5 & -7 & -2 \end{pmatrix}$   
 (d)  $\begin{pmatrix} -2 & 9/2 & -1 \\ -1 & 81/4 & 11 \\ -2 & 45/2 & 73/4 \end{pmatrix}$

**13. [MCQ]****[GATE-EE-2020: 2M]**

The number of purely real elements in a lower triangular representation of the given  $3 \times 3$  matrix, obtained through the given decomposition is\_\_\_\_\_.

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} a_{11} & 0 & 0 \\ a_{12} & a_{22} & 0 \\ a_{13} & a_{23} & a_{33} \end{bmatrix} \begin{bmatrix} a_{11} & 0 & 0 \\ a_{12} & a_{22} & 0 \\ a_{13} & a_{23} & a_{33} \end{bmatrix}^T$$

- (a) 5      (b) 6  
 (c) 8      (d) 9

**14. [MCQ]****[GATE-CE-2019:2M]**

The inverse of the matrix  $\begin{bmatrix} 2 & 3 & 4 \\ 4 & 3 & 1 \\ 1 & 2 & 4 \end{bmatrix}$  is.

(a)  $\begin{bmatrix} 10 & -4 & -9 \\ -15 & 4 & 14 \\ 5 & -1 & -6 \end{bmatrix}$

(b)  $\begin{bmatrix} -10 & 4 & 9 \\ 15 & -4 & -14 \\ -5 & 1 & 6 \end{bmatrix}$

(c)  $\begin{bmatrix} 2 & -\frac{4}{5} & -\frac{9}{5} \\ -3 & \frac{4}{5} & \frac{14}{5} \\ 1 & -\frac{1}{5} & -\frac{6}{5} \end{bmatrix}$

(d)  $\begin{bmatrix} -2 & \frac{4}{5} & \frac{9}{5} \\ 3 & -\frac{4}{5} & -\frac{14}{5} \\ -1 & \frac{1}{5} & \frac{6}{5} \end{bmatrix}$

**15. [MCQ]****[GATE-CS-2019:1M]**

Let X be a square matrix. Consider the following two statements on X.

I: X is invertible.

II: Determinant of X is non-zero.

Which one of the following is TRUE?

- (a) I implies II; II does not imply I.
- (b) II implies I; I does not imply II.
- (c) I does not imply II; II does not imply I.
- (d) I and II are equivalent statements.

**16. [MCQ]****[GATE-EE-2019: 2M]**

Consider a  $2 \times 2$  matrix  $M = [v_1, v_2]$ , where,  $v_1$  and  $v_2$

are the column vectors. Suppose  $M^{-1} = \begin{bmatrix} u_1^T \\ u_2^T \end{bmatrix}$ , where

$u_1^T$  and  $u_2^T$  are the row vectors. Consider the following statements:

**Statement 1:**  $u_1^T v_1 = 1$  and  $u_2^T v_2 = 1$ **Statement 2:**  $u_1^T v_2 = 0$  and  $u_2^T v_1 = 0$ 

Which of the following option is correct?

- (a) Statement 1 is true and statement 2 is false
- (b) Statement 2 is true and statement 1 is false
- (c) Both the statements are true
- (d) Both the statements are false

**17. [MCQ]****[GATE-CE-2018:1M]**

For the given orthogonal matrix Q =

$$\begin{bmatrix} \frac{3}{7} & \frac{2}{7} & \frac{6}{7} \\ \frac{7}{7} & \frac{7}{7} & \frac{7}{7} \\ -\frac{6}{7} & \frac{3}{7} & \frac{2}{7} \end{bmatrix}$$

The inverse is

$$\begin{bmatrix} \frac{3}{7} & \frac{2}{7} & \frac{6}{7} \\ \frac{7}{7} & \frac{7}{7} & \frac{7}{7} \\ \frac{2}{7} & \frac{6}{7} & -\frac{3}{7} \end{bmatrix}$$

$$\begin{bmatrix} \frac{3}{7} & \frac{2}{7} & \frac{6}{7} \\ -\frac{6}{7} & \frac{3}{7} & \frac{2}{7} \\ \frac{2}{7} & \frac{6}{7} & -\frac{3}{7} \end{bmatrix}$$

$$\begin{bmatrix} -\frac{3}{7} & -\frac{2}{7} & -\frac{6}{7} \\ \frac{6}{7} & -\frac{3}{7} & -\frac{2}{7} \\ -\frac{2}{7} & -\frac{6}{7} & \frac{3}{7} \end{bmatrix}$$

$$\begin{bmatrix} \frac{3}{7} & -\frac{6}{7} & \frac{2}{7} \\ \frac{2}{7} & \frac{3}{7} & \frac{6}{7} \\ \frac{6}{7} & \frac{2}{7} & -\frac{3}{7} \end{bmatrix}$$

$$\begin{bmatrix} -\frac{3}{7} & \frac{6}{7} & -\frac{2}{7} \\ -\frac{2}{7} & -\frac{3}{7} & -\frac{6}{7} \\ -\frac{6}{7} & -\frac{2}{7} & \frac{3}{7} \end{bmatrix}$$

**18. [MCQ]****[GATE-CE-2018:1M]**

Which one of the following matrices is singular

(a)  $\begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$

(b)  $\begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix}$

(c)  $\begin{bmatrix} 2 & 4 \\ 3 & 6 \end{bmatrix}$

(d)  $\begin{bmatrix} 4 & 3 \\ 6 & 2 \end{bmatrix}$

**19. [NAT]****[GATE-ME-2018: 1M]**

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 1 \end{bmatrix}$$

If  $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 1 \end{bmatrix}$  then  $\det(A^{-1})$  is \_\_\_\_\_ (Correct to

two decimal places).

**20. [MCQ]****[GATE-CE-2017:2M]**

If  $A = \begin{bmatrix} 1 & 5 \\ 6 & 2 \end{bmatrix}$  and  $B = \begin{bmatrix} 3 & 7 \\ 8 & 4 \end{bmatrix}$ ,  $AB^T$  is equal to

(a)  $\begin{bmatrix} 38 & 28 \\ 32 & 56 \end{bmatrix}$

(b)  $\begin{bmatrix} 3 & 40 \\ 42 & 8 \end{bmatrix}$

(c)  $\begin{bmatrix} 43 & 27 \\ 34 & 50 \end{bmatrix}$

(d)  $\begin{bmatrix} 38 & 32 \\ 28 & 56 \end{bmatrix}$

**21. [MCQ]****[GATE-CE-2017:1M]**

The matrix P is the inverse of a matrix Q. If I denotes the identity matrix, which one of the following option is correct?

(a)  $PQ = I$  but  $QP \neq I$       (b)  $QP = I$  but  $PQ \neq I$ (c)  $PQ = I$  and  $QP = I$       (d)  $PQ - QP = I$

**22. [NAT]****[GATE-ME-2017: 2M]**

Consider the matrix  $A = \begin{bmatrix} 50 & 70 \\ 70 & 80 \end{bmatrix}$  whose eigenvectors corresponding to eigenvalues  $\lambda_1$  and  $\lambda_2$  are  $x_1 = \begin{bmatrix} 70 \\ \lambda_1 - 50 \end{bmatrix}$  and  $x_2 = \begin{bmatrix} \lambda_2 - 80 \\ 70 \end{bmatrix}$ , respectively.

The value of  $x_1^T x_2$  is \_\_\_\_\_.

**23. [MCQ]****[GATE-ME-2017: 2M]**

Consider the matrix

$$P = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{-1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Which one of the following statements about P is INCORRECT?

- (a) Determinant of P is equal to 1.
- (b) P is orthogonal.
- (c) Inverse of P is equal to its transpose.
- (d) All eigenvalues of P are real numbers.

**24. [NAT]****[GATE-EE-2017: 1M]**

Let X and y be integers satisfying the following equations

$$2x^2 + y^2 = 34$$

$$x + 2y = 11$$

The value of  $(x + y)$  is \_\_\_\_\_.

**25. [MCQ]****[GATE-ME-2016: 1M]**

A real square matrix A is called skew-symmetric if

- (a)  $A^T = A$
- (b)  $A^T = A^{-1}$
- (c)  $A^T = -A$
- (d)  $A^T = A + A^{-1}$

**26. [NAT]****[GATE-EC-2016: 2M]**

The matrix  $\begin{bmatrix} a & 0 & 3 & 7 \\ 2 & 5 & 1 & 3 \\ 0 & 0 & 2 & 4 \\ 0 & 0 & 0 & b \end{bmatrix}$  has  $\det(A) = 100$  and trace

$(A) = 14$ . The value of  $|a-b|$  is \_\_\_\_\_.

**27. [MCQ]****[GATE-EC-2016: 1M]**

Let  $M^4 = I$ , (where I denotes the identity matrix) and  $M \neq I$ ,  $M^2 \neq I$  and  $M^3 \neq I$ . Then, for any natural number k,  $M^{-1}$  equals:

- (a)  $M^{4k} + 1$
- (b)  $M^{4k+2}$
- (c)  $M^{4k+3}$
- (d)  $M^{4k}$

**28. [NAT]****[GATE-EC-2016: 2M]**

A sequence  $x[n]$  is specified as

$$\begin{bmatrix} x[n] \\ x[n-1] \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \text{ for } n \geq 2.$$

The initial conditions are  $x[0] = 1$ ,  $x[1] = 1$  and  $x[n] = 0$  for  $n < 0$ . The value of  $x[12]$  is \_\_\_\_\_.

**29. [MCQ]****[GATE-EE-2016: 2M]**

Let  $P = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Consider the set S of all vectors

$\begin{pmatrix} x \\ y \end{pmatrix}$  such that  $a^2 + b^2 = 1$  where  $\begin{pmatrix} a \\ b \end{pmatrix} = P \begin{pmatrix} x \\ y \end{pmatrix}$ . Then S

is

(a) a circle of radius  $\sqrt{10}$

(b) a circle of radius  $\frac{1}{\sqrt{10}}$

(c) An ellipse with major axis along  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$

(d) An ellipse with major axis along  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$

**30. [NAT]****[GATE-CS-2016: 1M]**

Suppose that the eigenvalues of matrix A are 1, 2, 4.

The determinant of  $(A^{-1})^T$  is \_\_\_\_\_.

**31. [NAT]****[GATE-CS-2015: 1M]**

In the LU decomposition of the matrix  $\begin{bmatrix} 2 & 2 \\ 4 & 9 \end{bmatrix}$  if the

diagonal elements of U are both 1, then the lower diagonal entry  $l_{22}$  of L is \_\_\_\_\_.

**32. [MCQ]****[GATE-ME-2015: 1M]**

If any two columns of a determinant  $P = \begin{vmatrix} 4 & 7 & 8 \\ 3 & 1 & 5 \\ 9 & 6 & 2 \end{vmatrix}$

are interchanged, which one of the following statements regarding the value of the determinant is **CORRECT?**

- (a) Absolute value remains unchanged but sign will change.
- (b) Both absolute value and sign will change.
- (c) Absolute value will change but sign will not change.
- (d) Both absolute value and sign will remain unchanged.

**33. [MSQ]****[GATE-CE-2015:2M]**

$P$  and  $Q$  are two square matrices of the same order. Which of the following statement(s) is/are correct?

- (a) If  $P$  and  $Q$  are invertible then  $[PQ]^{-1} = Q^{-1}P^{-1}$
- (b) If  $P$  and  $Q$  are invertible then  $[QP]^{-1} = P^{-1}Q^{-1}$
- (c) If  $P$  and  $Q$  are invertible then  $[PQ]^{-1} = P^{-1}Q^{-1}$
- (d) If  $P$  and  $Q$  are not invertible then  $[PQ]^{-1} = Q^{-1}P^{-1}$

**34. [MCQ]****[GATE-ME-2015: 2M]**

For a given matrix  $P = \begin{bmatrix} 4+3i & -i \\ i & 4-3i \end{bmatrix}$ , where

$i = \sqrt{-1}$ , the inverse of matrix  $P$  is

- (a)  $\frac{1}{24} \begin{bmatrix} 4-3i & i \\ -i & 4+3i \end{bmatrix}$
- (b)  $\frac{1}{25} \begin{bmatrix} i & 4-3i \\ 4+3i & -i \end{bmatrix}$
- (c)  $\frac{1}{24} \begin{bmatrix} 4+3i & -i \\ i & 4-3i \end{bmatrix}$
- (d)  $\frac{1}{25} \begin{bmatrix} 4+3i & -i \\ i & 4-3i \end{bmatrix}$

**35. [MCQ]****[GATE-ME-2014: 1M]**

Given that the determinant of the matrix  $\begin{bmatrix} 1 & 3 & 0 \\ 2 & 6 & 4 \\ -1 & 0 & 2 \end{bmatrix}$

is  $-12$ , the determinant of the matrix  $\begin{bmatrix} 2 & 6 & 0 \\ 4 & 12 & 8 \\ -2 & 0 & 4 \end{bmatrix}$  is

- (a)  $-96$
- (b)  $-24$
- (c)  $24$
- (d)  $96$

**36. [MCQ]****[GATE-ME-2014: 1M]**

Which one of the following equations is a correct identity for arbitrary  $3 \times 3$  real matrices  $P$ ,  $Q$  and  $R$ ?

- (a)  $P(Q + R) = PQ + RP$
- (b)  $(P - Q)^2 = P^2 - 2PQ + Q^2$
- (c)  $\det(P + Q) = \det P + \det Q$
- (d)  $(P + Q)^2 = P^2 + PQ + QP + Q^2$

**37. NAT****[GATE-EC-2014:2M]**

The maximum value of the determinant among all  $2 \times 2$  real symmetric matrices with trace  $14$  is \_\_\_\_\_

**38. NAT****[GATE-EC-2014:2M]**

Consider the matrix

$$J_6 = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Which is obtained by reversing the order of the columns of the identity matrix  $I_6$ .

Let  $P = I_6 + \alpha J_6$ , where  $\alpha$  is a non-negative real number. The value of  $\alpha$  for which  $\det(P) = 0$  is \_\_\_\_\_.

**39. [MCQ]****[GATE-EC-2014:1M]**

For matrices of same dimension  $M$ ,  $N$  and scalar  $c$ , which one of these properties DOES NOT ALWAYS hold?

- (a)  $(M^T)^T = M$
- (b)  $(cM)^T = c(M)^T$
- (c)  $(M + N)^T = M^T + N^T$
- (d)  $MN = NM$



**51. [MCQ]****[GATE-EE-2011: 2M]**

The matrix  $[A] = \begin{bmatrix} 2 & 1 \\ 4 & -1 \end{bmatrix}$  is decomposed into a product of a lower triangular matrix  $[L]$  and an upper triangular matrix  $[U]$ . The properly decomposed  $[L]$  and  $[U]$  matrices respectively are

- (a)  $\begin{bmatrix} 1 & 0 \\ 4 & -1 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 1 \\ 0 & -2 \end{bmatrix}$
- (b)  $\begin{bmatrix} 2 & 0 \\ 4 & -1 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$
- (c)  $\begin{bmatrix} 1 & 0 \\ 4 & 1 \end{bmatrix}$  and  $\begin{bmatrix} 2 & 1 \\ 0 & -1 \end{bmatrix}$
- (d)  $\begin{bmatrix} 2 & 0 \\ 4 & -3 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix}$

**52. [MCQ]****[GATE-CE-2011:2M]**

The Eigen values of the matrix  $\begin{bmatrix} 9 & 5 \\ 5 & 8 \end{bmatrix}$  are

- (a) -2.42 and 6.86
- (b) 3.48 and 13.53
- (c) 4.70 and 6.86
- (d) 6.86 and 9.50

**53. [MCQ]****[GATE-ME-2009: 1M]**

For a matrix  $[M] = \begin{bmatrix} \frac{3}{x} & \frac{4}{5} \\ \frac{5}{x} & \frac{5}{5} \\ x & \frac{3}{5} \end{bmatrix}$ , the transpose of the

matrix is equal to the inverse of the matrix,  $[M]^T = [M]^{-1}$ . The value of x is given by

- (a)  $-\frac{4}{5}$
- (b)  $-\frac{3}{5}$
- (c)  $\frac{3}{5}$
- (d)  $\frac{4}{5}$

**54. [MCQ]****[GATE-CE-2009:2M]**

The inverse of the matrix  $\begin{bmatrix} 3+2i & i \\ -i & 3-2i \end{bmatrix}$  is,

- (a)  $\frac{1}{12} \begin{bmatrix} 3+2i & -i \\ i & 3-2i \end{bmatrix}$
- (b)  $\frac{1}{12} \begin{bmatrix} 3-2i & -i \\ i & 3+2i \end{bmatrix}$
- (c)  $\frac{1}{14} \begin{bmatrix} 3+2i & -i \\ i & 3-2i \end{bmatrix}$
- (d)  $\frac{1}{14} \begin{bmatrix} 3+2i & -i \\ i & 3+2i \end{bmatrix}$

**55. [MCQ]****[GATE-CE-2008:1M]**

The product of matrices  $(PQ)^{-1} P$  is

- (a)  $P^{-1}$
- (b)  $Q^{-1}$
- (c)  $P^{-1}Q^{-1}P$
- (d)  $PQ P^{-1}$

**56. [MCQ]****[GATE-EE-2008: 2M]**

Let  $P$  be a  $2 \times 2$  real orthogonal matrix and  $\vec{x}$  is a real vector  $[x_1, x_2]^T$  with length  $\|\vec{x}\| = (x_1^2 + x_2^2)^{\frac{1}{2}}$ .

Then, which one of the following statements is correct?

- (a)  $\|P\vec{x}\| \leq \|\vec{x}\|$  where at least one vector satisfies  $\|P\vec{x}\| < \|\vec{x}\|$
- (b)  $\|P\vec{x}\| = \|\vec{x}\|$  for all vectors  $\vec{x}$
- (c)  $\|P\vec{x}\| \geq \|\vec{x}\|$  where at least one vector satisfies  $\|P\vec{x}\| > \|\vec{x}\|$
- (d) No relationship can be established between  $\|\vec{x}\|$  and  $\|P\vec{x}\|$

**57. [MCQ]****[GATE-EE-2008: 2M]**

$A$  is  $m \times n$  full rank matrix with  $m > n$  and  $I$  is an identity matrix. Let matrix  $A' = (A^T A)^{-1} A^T$ . Then which one of the following statements is FALSE?

- (a)  $AA'A = A$
- (b)  $(AA')^2 = AA'$
- (c)  $A'A = I$
- (d)  $AA'A = A'$

**Rank of Matrices****58. [NAT]****[GATE-CS-2021: 1M]**

Suppose that  $P$  is a  $4 \times 5$  matrix such that every solution of the equation  $Px = 0$  is a scalar multiple of  $[2 \ 5 \ 4 \ 3 \ 1]^T$ . The rank of  $P$  is \_\_\_\_\_.

**59. [MCQ]****[GATE-CE-2021:1M]**

The rank of matrix  $\begin{bmatrix} 1 & 2 & 2 & 3 \\ 3 & 4 & 2 & 5 \\ 5 & 6 & 2 & 7 \\ 7 & 8 & 2 & 9 \end{bmatrix}$  is

- (a) 1
- (b) 2
- (c) 3
- (d) 4



**72. [MCQ]****[GATE-EE-2008: 1M]**

If the rank of a  $(5 \times 6)$  matrix Q is 4, then which one of the following statements is correct?

- (a) Q will have four linearly independent rows and four linearly independent columns.
- (b) Q will have four linearly independent rows and five linearly independent columns
- (c)  $QQ^T$  will be invertible
- (d)  $Q^TQ$  will be invertible

**System of Equations****73. [MSQ]****[GATE-CE-2023:2M]**

For the matrix

$$[A] = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix}$$

Which of the following statements is/are TRUE?

- (a)  $[A]\{x\}=\{b\}$  has a unique solution
- (b)  $[A]\{x\}=\{b\}$  does not have a unique solution
- (c)  $[A]$  has three linearly independent eigenvectors
- (d)  $[A]$  is a positive definite matrix

**74. [MCQ]****[GATE-EC-2022:1M]**

Consider a system of linear equations  $Ax = b$ , where

$$\begin{bmatrix} 1 & -\sqrt{2} & 3 \\ -1 & \sqrt{2} & -3 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

This system of equations admits \_\_\_\_\_.

- (a) a unique solution for  $x$
- (b) infinitely many solutions for  $x$
- (c) no solutions for  $x$
- (d) exactly two solutions for  $x$

**75. [MCQ]****[GATE-CS-2022:2M]**

Consider solving the following system of simultaneous equations using LU decomposition.

$$x_1 + x_2 - 2x_3 = 4$$

$$x_1 + 3x_2 - x_3 = 7$$

$$2x_1 + x_2 - 5x_3 = 7$$

where L and U are denoted as  $\begin{pmatrix} L_{11} & 0 & 0 \\ L_{21} & L_{22} & 0 \\ L_{31} & L_{32} & L_{33} \end{pmatrix}$ ,

$$U = \begin{pmatrix} U_{11} & U_{12} & U_{13} \\ 0 & U_{22} & U_{23} \\ 0 & 0 & U_{33} \end{pmatrix}$$

Which one of the following is the correct combination of values for  $L_{32}$ ,  $U_{33}$ , and  $x_1$ ?

- (a)  $L_{32} = 2$ ,  $U_{33} = -1/2$ ,  $x_1 = -1$
- (b)  $L_{32} = -1/2$ ,  $U_{33} = 2$ ,  $x_1 = 0$
- (c)  $L_{32} = 2$ ,  $U_{33} = 2$ ,  $x_1 = -1$
- (d)  $L_{32} = -1/2$ ,  $U_{33} = -1/2$ ,  $x_1 = 0$

**76. [MSQ]****[GATE-ME-2022: 2M]**

The system of linear equations in real  $(x, y)$  given by

$$(xy) \begin{bmatrix} 2 & 5-2\alpha \\ \alpha & 1 \end{bmatrix} = (0 \ 0) \text{ involves a real parameter}$$

$\alpha$  and has infinitely many non-trivial solutions for special value(s) of  $\alpha$ . Which one or more among the following options is/are non-trivial solution(s) of  $(x, y)$  for such special value(s) of  $\alpha$ ?

- (a)  $x = 2, y = -2$
- (b)  $x = -1, y = 4$
- (c)  $x = 1, y = 1$
- (d)  $x = 4, y = -2$

**77. [MSQ]****[GATE-ME-2022: 1M]**

A is a  $3 \times 5$  real matrix of rank 2. For the set of homogeneous equations  $Ax = O$ , where O is a zero vector and x is a vector of unknown variables, which of the following is/are true?

- (a) The given set of equations will have a unique solution.
- (b) The given set of equations will be satisfied by a zero vector of appropriate size.
- (c) The given set of equations will have infinitely many solutions.
- (d) The given set of equations will have many but a finite number of solutions.

**78. [NAT]****[GATE-CE-2020:2M]**

$$\begin{bmatrix} 1 & 3 & 2 \\ 2 & 2 & -3 \\ 4 & 4 & -6 \\ 2 & 5 & 2 \end{bmatrix} \begin{matrix} x_1 \\ x_2 \\ x_3 \end{matrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}.$$

Consider the system of equations

The value of  $x_3$  (round off to the nearest integer), is \_\_\_\_\_.

**79. [MCQ]****[GATE-EC-2020:2M]**

Consider the following system of linear equations.

$$\begin{array}{ll} x_1 + 2x_2 = b_1; & 2x_1 + 4x_2 = b_2; \\ 3x_1 + 7x_2 = b_3; & 3x_1 + 9x_2 = b_4 \end{array}$$

Which one of the following conditions ensures that a solution exists for the above system?

- (a)  $b_2 = 2b_1$  and  $6b_1 - 3b_3 + b_4 = 0$
- (b)  $b_3 = 2b_1$  and  $6b_1 - 3b_3 + b_4 = 0$
- (c)  $b_2 = 2b_1$  and  $3b_1 - 6b_3 + b_4 = 0$
- (d)  $b_3 = 2b_1$  and  $3b_1 - 6b_3 + b_4 = 0$

**80. [MCQ]****[GATE-ME-2019: 2M]**

The set of equations

$$\begin{aligned} x + y + z &= 1 \\ ax - ay + 3z &= 5 \\ 5x - 3y + az &= 6 \end{aligned}$$

has infinite solutions, if  $a =$

- (a) -3
- (b) 3
- (c) 4
- (d) -4

**81. [NAT]****[GATE-EC-2018:1M]**

Consider matrix  $A = \begin{bmatrix} k & 2k \\ k^2 - k & k^2 \end{bmatrix}$  and vector  $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ . The number of distinct real values of  $k$  for

which the equation  $Ax = 0$  has infinitely many solutions is \_\_\_\_\_.

**82. [MCQ]****[GATE-CS-2017: 1M]**

Let  $c_1, \dots, c_n$  be scalars, not all zero, such that

$$\sum_{i=1}^n a_i x_i = 0 \text{ where } a_i \text{ are column vectors in } \mathbb{R}^n.$$

Consider the set of linear equations

$$Ax = b$$

where  $A = [a_1, \dots, a_n]$  and  $b = \sum_{i=1}^n a_i$ . The set of

equations has

- (a) a unique solution at  $x = J_n$  where  $J_n$  denotes a  $n$ -dimensional vector of all 1
- (b) no solution
- (c) infinitely many solutions
- (d) finitely many solutions

**83. [MCQ]****[GATE-CE-2017:2M]**

Consider the following simultaneous equations (with  $c_1$  and  $c_2$  being constants)

$$3x_1 + 2x_2 = c_1$$

$$4x_1 + x_2 = c_2$$

The characteristics equation for these simultaneous equation is

- (a)  $\lambda^2 - 4\lambda - 5 = 0$
- (b)  $\lambda^2 - 4\lambda + 5 = 0$
- (c)  $\lambda^2 + 4\lambda - 5 = 0$
- (d)  $\lambda^2 + 4\lambda + 5 = 0$

**84. [MCQ]****[GATE-ME-2016: 1M]**

The solution to the system of equations

$$\begin{bmatrix} 2 & 5 \\ -4 & 3 \end{bmatrix} \begin{Bmatrix} x \\ y \end{Bmatrix} = \begin{Bmatrix} 2 \\ -30 \end{Bmatrix}$$

is

- (a) 6, 2
- (b) -6, 2
- (c) -6, -2
- (d) 6, -2

**85. [MCQ]****[GATE-CS-2016:1M]**

Consider the systems, each consisting of  $m$  linear equations in  $n$  variables.

- I. If  $m < n$ , then all such systems have a solution
- II. If  $m > n$ , then none of these systems has a solution
- III. If  $m = n$ , then there exists a system which has a solution

Which one of the following is CORRECT?

- (a) I, II and III are true
- (b) Only II and III are true
- (c) Only III is true
- (d) None of them is true

**86. [MCQ]****[GATE-CE-2016:1M]**

Consider the following linear system

$$x + 2y - 3z = a$$

$$2x + 3y + 3z = b$$

$$5x + 9y - 6z = c$$

This system is consistent if  $a$ ,  $b$  and  $c$  satisfy the equation

- (a)  $7a - b - c = 0$
- (b)  $3a + b - c = 0$
- (c)  $3a - b + c = 0$
- (d)  $7a - b + c = 0$

**87. [NAT]****[GATE-EC-2015: 1M]**

Consider a system of linear equations:

$$x - 2y + 3z = -1$$

$$x - 3y + 4z = 1, \text{ and}$$

$$-2x + 4y - 6z = k$$

The value of  $k$  for which the system has infinitely many solutions is \_\_\_\_\_.

**88. [NAT]****[GATE-CE-2015:1M]**

For what value of  $P$  the following set of equations will have no solutions

$$2x + 3y = 5$$

$$3x + py = 10$$

**89. [MCQ]****[GATE-EC-2014:2M]**

The system of linear equations

$$\begin{pmatrix} 2 & 1 & 3 \\ 3 & 0 & 1 \\ 1 & 2 & 5 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 5 \\ -4 \\ 14 \end{pmatrix} \text{ has}$$

- (a) a unique solution
- (b) infinitely many solutions
- (c) no solution
- (d) exactly two solutions

**90. [MCQ]****[GATE-EE-2014: 1M]**

Given a system of equation:

$$x + 2y + 2z = b_1$$

$$5x + y + 3z = b_2$$

Which of the following is true regarding its solutions

- (a) The system has a unique solution for any given  $b_1$  and  $b_2$
- (b) The system will have infinitely many solutions for any given  $b_1$  and  $b_2$
- (c) Whether or not a solution exists depends on the given  $b_1$  and  $b_2$
- (d) The system would have no solution for any value of  $b_1$  and  $b_2$

**91. [MCQ]****[GATE-ME-2014: 1M]**

The matrix form of the linear system  $\frac{dx}{dt} = 3x - 5y$

and  $\frac{dy}{dt} = 4x + 8y$  is

$$(a) \frac{d}{dt} \begin{Bmatrix} x \\ y \end{Bmatrix} = \begin{bmatrix} 3 & -5 \\ 4 & 8 \end{bmatrix} \begin{Bmatrix} x \\ y \end{Bmatrix}$$

$$(b) \frac{d}{dt} \begin{Bmatrix} x \\ y \end{Bmatrix} = \begin{bmatrix} 3 & 8 \\ 4 & -5 \end{bmatrix} \begin{Bmatrix} x \\ y \end{Bmatrix}$$

$$(c) \frac{d}{dt} \begin{Bmatrix} x \\ y \end{Bmatrix} = \begin{bmatrix} 4 & -5 \\ 3 & 8 \end{bmatrix} \begin{Bmatrix} x \\ y \end{Bmatrix}$$

$$(d) \frac{d}{dt} \begin{Bmatrix} x \\ y \end{Bmatrix} = \begin{bmatrix} 4 & 8 \\ 3 & -5 \end{bmatrix} \begin{Bmatrix} x \\ y \end{Bmatrix}$$

**92. [MCQ]****[GATE-CE-2014:2M]**

The following simultaneous equations

$$x + y + z = 3$$

$$x + 2y + 3z = 4$$

$$x + 4y + kz = 6$$

will NOT have a unique solution for  $k$  equal to

- (a) 0
- (b) 5
- (c) 6
- (d) 7

**93. [NAT]****[GATE-CS-2014:1M]**

Consider the following system of equations:

$$3x + 2y = 1$$

$$4x + 7z = 1$$

$$x + y + z = 3$$

$$x - 2y + 7z = 0$$

The number of solutions for this system is \_\_\_\_\_.

**94. [MCQ]****[GATE-ME-2012: 2M]**

$$x + 2y + z = 4$$

$$2x + y + 2z = 5$$

$$x - y + z = 1$$

The system of algebraic equations given above has

- (a) A unique solution of  $x = 1, y = 1$  and  $z = 1$
- (b) Only the two solutions of  $(x = 1, y = 1, z = 1)$  and  $(x = 2, y = 1, z = 0)$
- (c) Infinite number of solutions
- (d) No feasible solution

**95. [MCQ]****[GATE-EC-2011:2M]**

The system of equations

$$x + y + z = 6$$

$$x + 4y + 6z = 20$$

$$x + 4y + \lambda z = \mu$$

has No solution for values of  $\lambda$  and  $\mu$  given by

- |                                |                                   |
|--------------------------------|-----------------------------------|
| (a) $\lambda = 6, \mu = 20$    | (b) $\lambda = 6, \mu \neq 20$    |
| (c) $\lambda \neq 6, \mu = 20$ | (d) $\lambda \neq 6, \mu \neq 20$ |

**96. [MCQ]****[GATE-ME-2011: 2M]**

Consider the following system of equations

$$2x_1 + x_2 + x_3 = 0$$

$$x_2 - x_3 = 0$$

$$x_1 + x_2 = 0$$

The system has

- (a) A unique solution
- (b) No solution
- (c) Infinite number of solutions
- (d) Five solutions

**97. [MCQ]****[GATE-EE-2010: 2M]**

For the set of equations

$$x_1 + 2x_2 + x_3 + 4x_4 = 2$$

$$3x_1 + 6x_2 + 3x_3 + 12x_4 = 6$$

The following statement is true:

- (a) Only the trivial solutions  $x_1 = x_2 = x_3 = x_4 = 0$
- (b) There are no solutions
- (c) A unique non-trivial solution exists
- (d) Multiple non-trivial solutions exist.

**98. [MCQ]****[GATE-CS-2008:1M]**

The following system of equations

$$x_1 + x_2 + 2x_3 = 1$$

$$x_1 + 2x_2 + 3x_3 = 2$$

$$x_1 + 4x_2 + \alpha x_3 = 4$$

Has a unique solution. The only possible value(s) for  $\alpha$  is/are

- (a) 0
- (b) either 0 or 1
- (c) One of 0, 1 or -1
- (d) any real number

**99. [MCQ]****[GATE-EC-2008:1M]**

The system of linear equations

$$4x + 2y = 7$$

$$2x + y = 6$$

has

- (a) a unique solution
- (b) no solution
- (c) an infinite number of solution
- (d) exactly two distinct solutions

**100.[MCQ]****[GATE-ME-2008: 2M]**

For what value of  $a$ , if any, will the following system of equations in  $x$ ,  $y$  and  $z$  have a solution?

$$2x + 3y = 4$$

$$x + y + z = 4$$

$$x + 2y - z = a$$

- (a) Any real number
- (b) 0
- (c) 1
- (d) There is no such value

**Eigen Values and Eigen Vectors****101.[MSQ]****[GATE-CE-2023:2M]**

If  $M$  is an arbitrary real  $n \times n$  matrix, then which of the following matrices will have non-negative eigenvalues?

- |             |               |
|-------------|---------------|
| (a) $M^2$   | (b) $MM^T$    |
| (c) $M^T M$ | (d) $(M^T)^2$ |

**102. [MCQ]****[GATE-EC-2023: 1M]**

Let the sets of eigenvalues and eigenvectors of a matrix  $B$  be  $\{\lambda_k \mid 1 \leq k \leq n\}$  and  $\{v_k \mid 1 \leq k \leq n\}$ , respectively. For any invertible matrix  $P$ , the sets of eigenvalues and eigenvectors of the matrix  $A$ , where  $B = P^{-1}AP$ , respectively, are

- (a)  $\{\lambda_k \det(A) \mid 1 \leq k \leq n\}$  and  $\{Pv_k \mid 1 \leq k \leq n\}$
- (b)  $\{\lambda_k \mid 1 \leq k \leq n\}$  and  $\{v_k \mid 1 \leq k \leq n\}$
- (c)  $\{\lambda_k \mid 1 \leq k \leq n\}$  and  $\{Pv_k \mid 1 \leq k \leq n\}$
- (d)  $\{\lambda_k \mid 1 \leq k \leq n\}$  and  $\{P^{-1}v_k \mid 1 \leq k \leq n\}$

**103.[MSQ]****[GATE-CE-2023:2M]**

For the matrix

$$[A] = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 2 \end{bmatrix}$$

which of the following statements is/are TRUE?

- (a) The eigenvalues of  $[A]^T$  are same as the eigenvalues of  $[A]$
- (b) The eigenvalues of  $[A]^{-1}$  are the reciprocals of the eigenvalues of  $[A]$
- (c) The eigenvectors of  $[A]^T$  are same as the eigenvectors of  $[A]$
- (d) The eigenvectors of  $[A]^{-1}$  are same as the eigenvectors of  $[A]$

**104.[MSQ]****[GATE-CE-2022:1M]**

Let  $y$  be a non-zero vector of the size  $2022 \times 1$ . Which of the following statement(s) is/are TRUE?

- (a)  $yy^T$  is symmetric matrix
- (b)  $yy^T$  is invertible
- (c)  $yy^T$  has a rank of 2022
- (d)  $y^T y$  is an eigen value of  $yy^T$

**105.[MSQ]****[GATE-CE-2022:1M]**

The matrix  $M$  is defined as  $M = \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix}$  and has eigenvalues 5 and -2. The matrix  $Q$  is formed as  $Q = M^3 - 4M^2 - 2M$

Which of the following is/are the eigenvalue(s) of matrix  $Q$ ?

- (a) 15
- (b) 28
- (c) -20
- (d) -30

**106.[NAT]****[GATE-ME-2022: 2M]**

If the sum and product of eigenvalues of a  $2 \times 2$  real matrix  $\begin{bmatrix} 3 & p \\ p & q \end{bmatrix}$  are 4 and -1 respectively, then  $|p|$  is \_\_\_\_\_ (in integer).

**107.[MCQ]****[GATE-EE-2022: 2M]**

Consider a matrix  $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & -2 \\ 0 & 1 & 1 \end{bmatrix}$

The matrix  $A$  satisfies the equation  $6A^{-1} = A^2 + cA + dI$ , where  $c$  and  $d$  are scalars and  $I$  is the identity matrix. Then  $(c + d)$  is equal to

- (a) 5
- (b) 17
- (c) -6
- (d) 11

**108.[MCQ]****[GATE-EE-2022: 2M]**

$e^A$  denotes the exponential of a square matrix  $A$ . Suppose  $\lambda$  is an eigenvalue and  $v$  is the corresponding eigen-vector of matrix  $A$ .

Consider the following two statements:

- Statement 1:  $e^\lambda$  is an eigenvalue of  $e^A$
- Statement 2:  $v$  is an eigen-vector of  $e^A$ .

Which one of the following options is correct?

- (a) Statement 1 is true and statement 2 is false
- (b) Statement 1 is false and statement 2 is true
- (c) Both the statements are correct
- (d) Both the statements are false.

**109.[MCQ]****[GATE-EE-2022: 1M]**

Consider a  $3 \times 3$  matrix  $A$  whose  $(i, j)^{\text{th}}$  element,  $a_{i,j} = (i-j)^3$ . Then the matrix  $A$  will be

- (a) symmetric
- (b) skew-symmetric
- (c) unitary
- (d) null

**110.[MCQ]****[GATE-ME-2021: 2M]**

Consider a vector  $p$  in 2-dimensional space. Let its direction (counter clockwise angle with the positive x-axis) be  $\theta$ . Let  $p$  be an eigenvector of a  $2 \times 2$  matrix  $A$  with corresponding eigenvalue  $\lambda$ ,  $\lambda > 0$ . If we denote the magnitude of a vector  $v$  by  $\|v\|$ , identify the VALID statement regarding  $p'$ , where  $p' = Ap$ .

- (a) Direction of  $P' = \lambda\theta$ ,  $\|p'\| = \|p\|$
- (b) Direction of  $P' = \theta$ ,  $\|p'\| = \lambda \|p\|$
- (c) Direction of  $P' = \lambda\theta$ ,  $\|p'\| = \lambda \|p\|$
- (d) Direction of  $P' = \theta$ ,  $\|p'\| = \|p\|/\lambda$



**123.[NAT]****[GATE-EE-2018: 2M]**

Let  $A = \begin{bmatrix} 1 & 0 & -1 \\ -1 & 2 & 0 \\ 0 & 0 & -2 \end{bmatrix}$  and  $B = A^3 - A^2 - 4A + 5I$ ,

where  $I$  is the  $3 \times 3$  identity matrix. The determinant of  $B$  is \_\_\_\_\_ (up to 1 decimal place).

**124.[MCQ]****[GATE-CE-2018:2M]**

The matrix  $\begin{bmatrix} 2 & -4 \\ 4 & -2 \end{bmatrix}$  has

- (a) Real eigenvalues and eigenvectors
- (b) Real eigenvalue but complex eigenvectors
- (c) Complex eigenvalues but real eigenvectors
- (d) Complex eigenvalues and eigenvectors

**125.[NAT]****[GATE-CS-2018: 2M]**

Consider a matrix  $P$  whose only eigenvectors are the multiples of  $\begin{bmatrix} 1 \\ 4 \end{bmatrix}$ .

Consider the following statements.

- I:  $P$  does not have an inverse
- II:  $P$  has a repeated eigenvalue
- III:  $P$  cannot be diagonalized

Which one of the following options is correct?

- (a) Only I and III are necessarily true
- (b) Only II is necessarily true
- (c) Only I and II are necessarily true
- (d) Only II and III are necessarily true

**126.[NAT]****[GATE-EE-2018: 1M]**

Consider a non-singular  $2 \times 2$  square matrix  $A$ . If  $\text{trace}(A) = 4$  and  $\text{trace}(A^2) = 5$ , the determinant of the matrix  $A$  is \_\_\_\_\_ (up to 1 decimal place).

**127. [MSQ]****[GATE-EC-2018:1M]**

Let  $M$  be a real  $4 \times 4$  matrix. Consider the following statements:

- S1:  $M$  has 4 linearly independent eigenvectors.
- S2:  $M$  has 4 distinct eigenvalues.
- S3:  $M$  is non-singular (invertible).

Which one among the following is TRUE?

- (a) S1 implies S2
- (b) S1 implies S3
- (c) S2 implies S1
- (d) S3 implies S2

**128. [MCQ]****[GATE-CS-2017: 2M]**

Let  $A$  be  $n \times n$  real valued square symmetric matrix of rank 2 with  $\sum_{i=1}^n \sum_{j=1}^n A_{ij}^2 = 50$  Consider the following statements.

- (I) One eigenvalue must be in  $[-5, 5]$
- (II) The eigenvalue with the largest magnitude must be strictly greater than 5

Which of the above statements about eigenvalues of  $A$  is/are necessarily CORRECT?

- (a) Both (I) and (II)
- (b) (I) only
- (c) (II) only
- (d) Neither (I) nor (II)

**129.[NAT]****[GATE-CS-2017:2M]**

If the characteristic polynomial of a  $3 \times 3$  matrix  $M$  over  $\mathbb{R}$  (the set of real numbers) is  $\lambda^3 - 4\lambda^2 + a\lambda + 30$ ,  $a \in \mathbb{R}$ , and one eigenvalue of  $M$  is 2, then the largest among the absolute values of the eigenvalues of  $M$  is \_\_\_\_\_.

**130.[MCQ]****[GATE-CE-2017:2M]**

Consider the matrix  $\begin{bmatrix} 5 & -1 \\ 4 & 1 \end{bmatrix}$  which one of the following statements is TRUE for the eigenvalues and eigen vectors of this matrix?

- (a) Eigenvalue 3 has multiplicity of 2, and only one independent eigenvector exists.
- (b) Eigenvalue 3 has a multiplicity of 2, and two independent eigenvector exists.
- (c) Eigenvalue 3 has a multiplicity of 2, and no independent eigenvector exists.
- (d) Eigenvalue are 3 and  $-3$  and two independent eigenvector exists.

**131.[MCQ]****[GATE-EC-2017: 1M]**

Consider the  $5 \times 5$  matrix

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 1 & 2 & 3 & 4 \\ 4 & 5 & 1 & 2 & 3 \\ 3 & 4 & 5 & 1 & 2 \\ 2 & 3 & 4 & 5 & 1 \end{bmatrix}$$

It is given that  $A$  has only one real eigenvalue. Then the real eigenvalue of  $A$  is

- (a)  $-2.5$
- (b)  $0$
- (c)  $15$
- (d)  $25$



**142.[MCQ]****[GATE-EE-2016: 2M]**

Let the eigenvalues of a  $2 \times 2$  matrix A be 1, -2 with eigenvectors  $x_1$  and  $x_2$  respectively. Then the eigenvalues and eigenvectors of the matrix  $A^2 - 3A + 4I$  would, respectively, be

- (a) 2, 14 ;  $x_1, x_2$
- (b) 2, 14 ;  $x_1 + x_2, x_1 - x_2$
- (c) 2, 0 ;  $x_1, x_2$
- (d) 2, 0 ;  $x_1 + x_2, x_1 - x_2$

**143.[MCQ]****[GATE-EE-2016: 1M]**

A  $3 \times 3$  matrix P is such that,  $P^3 = P$ . Then the eigenvalues of P are

- (a) 1, 1, -1
- (b)  $1, 05 + j0.866, 0.5 - j0.866$
- (c)  $1, -0.5 + j0.868, -0.5 - 0.866$
- (d) 0, 1, -1

**144.[NAT]****[GATE-EE-2016: 1M]**

Consider a  $3 \times 3$  matrix with every element being equal to 1. Its only non-zero eigenvalue is \_\_\_\_\_.

**145.[MCQ]****[GATE-EE-2015: 1M]**

The maximum value of 'a' such that the matrix

$$\begin{bmatrix} -3 & 0 & -2 \\ 1 & -1 & 0 \\ 0 & a & -2 \end{bmatrix}$$

has three linearly independent real

eigenvectors is

- (a)  $\frac{2}{3\sqrt{3}}$
- (b)  $\frac{1}{3\sqrt{3}}$
- (c)  $\frac{1+2\sqrt{3}}{3\sqrt{3}}$
- (d)  $\frac{1+\sqrt{3}}{3\sqrt{3}}$

**146.[MCQ]****[GATE-EE-2015: 1M]**

We have a set of 3 linear equations in 3 unknowns.

'X ≡ Y' means X and Y are equivalent statements.

'X ≠ Y' means X and Y are not equivalent statements.

P. There is a unique solution.

Q. The equations are linearly independent.

R. All eigenvalues of the coefficient matrix are nonzero.

S. The determinant of the coefficient matrix is nonzero.

Which one of the following is TRUE?

- (a) P ≡ Q ≡ R ≡ S
- (b) P ≡ Q ≠ R ≡ S
- (c) P = Q ≠ R = 0
- (d) P ≠ Q ≠ R ≠ S

**147.[NAT]****[GATE-EE-2015: 1M]**

If the sum of the diagonal elements of a  $2 \times 2$  matrix is -6, then the maximum possible value of determinant of the matrix is \_\_\_\_\_.

**148.[MCQ]****[GATE-CS-2015:2M]**

Consider the following  $2 \times 2$  matrix A where two elements are unknown and are marked by a and b. The eigenvalues of this matrix are -1 and 7. What are the values of a and b?

$$A = \begin{pmatrix} 1 & 4 \\ b & a \end{pmatrix}$$

- (a) a = 6, b = 4
- (b) a = 4, b = 6
- (c) a = 3, b = 5
- (d) a = 5, b = 3

**149.[MCQ]****[GATE-CE-2015:2M]**

The smallest and largest Eigen value of the following matrix are

$$\begin{bmatrix} 3 & -2 & 2 \\ 4 & -4 & 6 \\ 2 & -3 & 5 \end{bmatrix}$$

- (a) 1.5 and 2.5
- (b) 0.5 and 2.5
- (c) 1.0 and 3.0
- (d) 1.0 and 2.0

**150.[MCQ]****[GATE-CE-2015:2M]**

The two Eigen values of the matrix  $\begin{bmatrix} 2 & 1 \\ 1 & p \end{bmatrix}$  have a ratio of 3 : 1 for p = 2. What is the another value of P if eigen value have the same ratio of 3 : 1 ?

- (a) -2
- (b) 1
- (c) 7/3
- (d) 14/3

**151.[NAT]****[GATE-CS-2015: 1M]**

The largest of the two eigenvalues of the matrix  $\begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix}$  is \_\_\_\_.

**152.[MCQ]****[GATE-CS-2015:1M]**

In the given matrix  $\begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 1 & 2 & 1 \end{bmatrix}$  one of the eigenvalues is 1. The eigenvectors corresponding to the eigenvalues 1 are

- (a)  $\{\alpha(4, 2, 1) | \alpha \neq 0, \alpha \in \mathbb{R}\}$
- (b)  $\{\alpha(-4, 2, 1) | \alpha \neq 0, \alpha \in \mathbb{R}\}$
- (c)  $\{\alpha(\sqrt{2}, 0, 1) | \alpha \neq 0, \alpha \in \mathbb{R}\}$
- (d)  $\{\alpha(-\sqrt{2}, 0, 1) | \alpha \neq 0, \alpha \in \mathbb{R}\}$

**153.[MCQ]****[GATE-EC-2015:1M]**

The value of x for which all the eigen-values of the

matrix given below are real is  $\begin{bmatrix} 10 & 5+j & 4 \\ x & 20 & 2 \\ 4 & 2 & -10 \end{bmatrix}$

- (a)  $5+j$
- (b)  $5-j$
- (c)  $1-5j$
- (d)  $1+5j$

**154.[MCQ]****[GATE-ME-2015: 1M]**

At least one eigenvalue of a singular matrix is

- (a) Positive
- (b) Zero
- (c) Negative
- (d) Imaginary

**155.[NAT]****[GATE-ME-2015: 1M]**

The lowest eigenvalue of the  $2 \times 2$  matrix  $\begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}$  is \_\_\_\_.

**156.[MCQ]****[GATE-ME-2014: 1M]**

One of the eigenvectors of the matrix  $\begin{bmatrix} -5 & 2 \\ -9 & 6 \end{bmatrix}$  is

- (a)  $\begin{Bmatrix} -1 \\ 1 \end{Bmatrix}$
- (b)  $\begin{Bmatrix} -2 \\ 9 \end{Bmatrix}$
- (c)  $\begin{Bmatrix} 2 \\ -1 \end{Bmatrix}$
- (d)  $\begin{Bmatrix} 1 \\ 1 \end{Bmatrix}$

**157.[MCQ]****[GATE-ME-2014: 1M]**

Consider a  $3 \times 3$  real symmetric matrix S such that two of its eigenvalues are  $a \neq 0, b \neq 0$  with respective

eigenvectors  $\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}, \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ . If  $a \neq b$ , then

$x_1 y_1 + x_2 y_2 + x_3 y_3$  equals

- (a) a
- (b) b
- (c) ab
- (d) 0

**158.[MCQ]****[GATE-EE-2014: 1M]**

Which one of the following statements is true for all real symmetric matrices?

- (a) All the eigenvalues are real
- (b) All the eigenvalues are positive
- (c) All the eigenvalues are distinct
- (d) Sum of all the eigenvalues is zero

**159.[NAT]****[GATE-EE-2014: 2M]**

A system matrix is given as follows:

$$A = \begin{bmatrix} 0 & 1 & -1 \\ -6 & -11 & 6 \\ -6 & -11 & 5 \end{bmatrix}$$

The absolute value of the ratio of the maximum eigenvalue to the minimum eigenvalue is \_\_\_\_.

**160.[NAT]****[GATE-CS-2014:1M]**

The value of the dot product of the eigenvectors corresponding to any pair of different eigenvalues of a 4-by-4 symmetric positive definite matrix is \_\_\_\_.

**161.[NAT]****[GATE-CS-2014:2M]**

$$\text{If the matrix } A \text{ is such that } A = \begin{bmatrix} 2 \\ -4 \\ 7 \end{bmatrix} \begin{bmatrix} 1 & 9 & 5 \end{bmatrix}$$

then the determinant of A is equal to \_\_\_\_.

**162.[MCQ]****[GATE-CE-2014:1M]**

The sum of Eigen values of matrix, [M] is where

$$[M] = \begin{bmatrix} 215 & 650 & 795 \\ 655 & 150 & 835 \\ 485 & 355 & 550 \end{bmatrix}$$

- (a) 915
- (b) 1355
- (c) 1640
- (d) 2180

**163.[MCQ]****[GATE-EC-2014:2M]**

Which one of the following statements is NOT true for a square matrix A?

- (a) "If A is upper triangular, the eigen values of A are the diagonal elements of it"
- (b) "If A is real, symmetric, the eigen values of A are always real and positive"
- (c) "If A is real, the eigen values of A and "  $A^T$  " are always the same."
- (d) "If all the principal minors of A are positive, all the eigen values of A are also positive."

**164.[MCQ]****[GATE-ME-2013: 1M]**

The eigenvalues of a symmetric matrix are all

- (a) Complex with non-zero positive imaginary part.
- (b) Complex with non-zero negative imaginary part.
- (c) Real.
- (d) Pure imaginary.

**165.[MCQ]****[GATE-EE-2013: 2M]**

A matrix has eigenvalues  $-1$  and  $-2$ . The corresponding eigen vectors are  $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$  and  $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$  respectively. The matrix is

- |  |  |
|--|--|
| (a) $\begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix}$ | (b) $\begin{bmatrix} 1 & 2 \\ -2 & -4 \end{bmatrix}$ |
| (c) $\begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix}$ | (d) $\begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$ |

**166.[MCQ]****[GATE-EE-2013: 1M]**

The equation  $\begin{bmatrix} 2 & -2 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$  has

- (a) no solution
- (b) only one solution  $\begin{bmatrix} X_1 \\ X_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$
- (c) non-zero unique solution
- (d) multiple solutions

**167.[MCQ]****[GATE-EC-2013:1M]**

The minimum eigen value of the following matrix is

$$\begin{bmatrix} 3 & 5 & 2 \\ 5 & 12 & 7 \\ 2 & 7 & 5 \end{bmatrix}$$

- (a) 0
- (b) 1
- (c) 2
- (d) 3

**168.[MCQ]****[GATE-ME-2012: 2M]**

For the matrix  $A = \begin{bmatrix} 5 & 3 \\ 1 & 3 \end{bmatrix}$ , one of the normalized

eigen vector is given as

- |   |   |
|---|---|
| (a) $\begin{pmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{pmatrix}$           | (b) $\begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} \end{pmatrix}$ |
| (c) $\begin{pmatrix} \frac{3}{\sqrt{10}} \\ \frac{-1}{\sqrt{10}} \end{pmatrix}$ | (d) $\begin{pmatrix} \frac{1}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \end{pmatrix}$  |

**169.[MCQ]****[GATE-EE-2012: 2M]**

Given that  $A = \begin{bmatrix} -5 & -3 \\ 2 & 0 \end{bmatrix}$  and  $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  the value of  $A^3$  is

- (a)  $15A + 12I$
- (b)  $19A + 30I$
- (c)  $17A + 15I$
- (d)  $17A + 21I$

**170.[NAT]****[GATE-CS-2012:1M]**

Let A be the  $2 \times 2$  matrix with elements  $a_{11} = a_{12} = a_{21} = +1$  and  $a_{22} = -1$ . Then the eigenvalues of the matrix  $A^{19}$  are

- (a)  $1024$  and  $-1024$
- (b)  $1024\sqrt{2}$  and  $-1024\sqrt{2}$
- (c)  $4\sqrt{2}$  and  $-4\sqrt{2}$
- (d)  $512\sqrt{2}$  and  $-512\sqrt{2}$

**171.[MCQ]****[GATE-CS-2011:2M]**

Consider the matrix as given below.

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 7 \\ 0 & 0 & 3 \end{bmatrix}$$

Which one of the following options provides the CORRECT values of the Eigen values of the matrix?

- (a) 1, 4, 3
- (b) 3, 7, 3
- (c) 7, 3, 2
- (d) 1, 2, 3

**172.[MCQ]****[GATE-ME-2011: 1M]**

Eigenvalues of a real symmetric matrix are always

- (a) Positive
- (b) Negative
- (c) Real
- (d) Complex

**173.[MCQ]****[GATE-ME-2010: 2M]**

One of the eigen vectors of the matrix  $A = \begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$  is

- |   |   |
|---|---|
| (a) $\begin{Bmatrix} 2 \\ -1 \end{Bmatrix}$ | (b) $\begin{Bmatrix} 2 \\ 1 \end{Bmatrix}$  |
| (c) $\begin{Bmatrix} 4 \\ 1 \end{Bmatrix}$  | (d) $\begin{Bmatrix} 1 \\ -1 \end{Bmatrix}$ |

**174.[MCQ]****[GATE-EC-2010:1M]**

The eigenvalues of a skew symmetric matrix are

- (a) always zero
- (b) Always pure imaginary
- (c) Either zero or pure imaginary
- (d) Always real

**175.[MCQ]****[GATE-CS-2010:2M]**

Consider the following matrix  $A = \begin{bmatrix} 2 & 3 \\ x & y \end{bmatrix}$ . If the

Eigen values of A are 4 and 8, then

- (a)  $x = 4, y = 10$
- (b)  $x = 5, y = 8$
- (c)  $x = 3, y = 9$
- (d)  $x = -4, y = 10$

**176.[MCQ]****[GATE-EE-2010: 2M]**

An eigenvector of  $P = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix}$  is

- (a)  $[-1 \ 1 \ 1]^T$
- (b)  $[1 \ 2 \ 1]^T$
- (c)  $[1 \ -1 \ 2]^T$
- (d)  $[2 \ 1 \ -1]^T$

**177.[MCQ]****[GATE-EE-2009: 1M]**

The trace and determinant of a  $2 \times 2$  matrix are known to be -2 and -35 respectively its eigenvalues are

- (a) -30 and -5
- (b) -37 and -1
- (c) -7 and 5
- (d) 17.5 and -2

**178.[MCQ]****[GATE-EC-2009:2M]**

The eigen values of the following matrix are

$$\begin{bmatrix} -1 & 3 & 5 \\ -3 & -1 & 6 \\ 0 & 0 & 3 \end{bmatrix}$$

- (a)  $3, 3 + 5j, 6 - j$
- (b)  $-6 + 5j, 3 + j, 3 - j$
- (c)  $3 + j, 3 - j, 5 + j$
- (d)  $3, -1 + 3j, -1 - 3j$

**179.[MCQ]****[GATE-CE-2008:1M]**

The Eigen values of the matrix  $[P] = \begin{bmatrix} 4 & 5 \\ 2 & -5 \end{bmatrix}$  are

- (a) -7 and 8
- (b) -6 and 5
- (c) 3 and 4
- (d) 1 and 2

**180.[MCQ]****[GATE-CS-2008:2M]**

How many of the following matrices have an Eigen value 1?

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \text{ and } \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix}$$

- (a) One
- (b) Two
- (c) Three
- (d) Four

**181.[MCQ]****[GATE-ME-2008: 1M]**

The matrix  $\begin{bmatrix} 1 & 2 & 4 \\ 3 & 0 & 6 \\ 1 & 1 & p \end{bmatrix}$  has one eigenvalue equal to 3.

The sum of the other two eigenvalue is

- (a) p
- (b)  $p - 1$
- (c)  $p - 2$
- (d)  $p - 3$

**182.[MCQ]****[GATE-ME-2008: 2M]**

The eigenvectors of the matrix  $\begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}$  are written in

the form  $\begin{bmatrix} 1 \\ a \end{bmatrix}$  and  $\begin{bmatrix} 1 \\ b \end{bmatrix}$ . What is  $a + b$ ?

- (a) 0
- (b)  $\frac{1}{2}$
- (c) 1
- (d) 2

**183.[MCQ]****[GATE-EC-2008: 1M]**

All the four entries of the  $2 \times 2$  matrix  $P = \begin{bmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \end{bmatrix}$

are nonzero, and one of its eigenvalues is zero. Which of the following statements is true?

- (a)  $P_{11}P_{22} - P_{12}P_{21} = 1$
- (b)  $P_{11}P_{22} - P_{12}P_{21} = -1$
- (c)  $P_{11}P_{22} - P_{12}P_{21} = 0$
- (d)  $P_{11}P_{22} + P_{12}P_{21} = 0$

**Basis and Spaces****184.[MCQ]****[GATE-CE-2023:2M]**

Two vectors  $[2 \ 1 \ 0 \ 3]^T$  and  $[1 \ 0 \ 1 \ 2]^T$  belong to the null space of a  $4 \times 4$  matrix of rank 2. Which one of the following vectors also belongs to the null space?

- (a)  $[1 \ 1 \ -1 \ 1]^T$
- (b)  $[2 \ 0 \ 1 \ 2]^T$
- (c)  $[0 \ -2 \ 1 \ -1]^T$
- (d)  $[3 \ 1 \ 1 \ 2]^T$

**185.[MCQ]****[GATE-EC-2015:1M]**

$A = \begin{bmatrix} 1 & \tan x \\ -\tan x & 1 \end{bmatrix}$ , the determinant of  $A^T A^{-1}$  is

- (a)  $\sec^2 x$
- (b)  $\cos 4x$
- (c) 1
- (d) 0

**186.[NAT]****[GATE-CS-2014:1M]**

If  $V_1$  and  $V_2$  are 4-dimensional subspaces of a 6-dimensional vector space  $V$ , then the smallest possible dimension of  $V_1 \cap V_2$  is \_\_\_\_\_.


**ANSWER KEY**

- |                    |                   |                    |                    |
|--------------------|-------------------|--------------------|--------------------|
| 1. (b)             | 2. (d)            | 3. (a)             | 4. (b)             |
| 5. (c)             | 6. (a)            | 7. (32)            | 8. (a)             |
| 9. (c)             | 10. (a)           | 11. (a)            | 12. (b)            |
| 13. (marks to all) | 14. (d)           | 15. (d)            | 16. (c)            |
| 17. (c)            | 18. (c)           | 19. (0.25 to 0.25) | 20. (a)            |
| 21. (c)            | 22. (0 to 0)      | 23. (d)            | 24. (7)            |
| 25. (c)            | 26. (3)           | 27. (c)            | 28. (233 to 233)   |
| 29. (d)            | 30. (0.125)       | 31. (5)            | 32. (a)            |
| 33. (a, b)         | 34. (a)           | 35. (a)            | 36. (d)            |
| 37. (49)           | 38. (1)           | 39. (d)            | 40. (200)          |
| 41. (c)            | 42. (23 to 23)    | 43. (a)            | 44. (88 to 88)     |
| 45. (a)            | 46. (b)           | 47. (16)           | 48. (b)            |
| 49. (d)            | 50. (b)           | 51. (d)            | 52. (b)            |
| 53. (a)            | 54. (b)           | 55. (b)            | 56. (b)            |
| 57. (d)            | 58. (4)           | 69. (b)            | 60. (c)            |
| 61. (3)            | 62. (b)           | 63. (b)            | 64. (b)            |
| 65. (c)            | 66. (2)           | 67. (4)            | 68. (17)           |
| 69. (b)            | 70. (c)           | 71. (2 to 2)       | 72. (a)            |
| 73. (b, c)         | 74. (c)           | 75. (d)            | 76. (a, b)         |
| 77. (b, c)         | 78. (3 to 3)      | 79. (a)            | 80. (c)            |
| 81. (2 to 2)       | 82. (c)           | 83. (a)            | 84. (d)            |
| 85. (c)            | 86. (b)           | 87. (2)            | 88. (4.49 to 4.51) |
| 89. (b)            | 90. (b)           | 91. (a)            | 92. (d)            |
| 93. (1)            | 94. (c)           | 95. (b)            | 96. (a)            |
| 97. (d)            | 98. (c)           | 99. (b)            | 100. (b)           |
| 101. (b, c)        | 102. (c)          | 103. (a, b, d)     | 104. (a, d)        |
| 105. (a, c)        | 106. (2 to 2)     | 107. (a)           | 108. (c)           |
| 109. (b)           | 110. (b)          | 111. (d)           | 112. (1)           |
| 113. (b)           | 114. (3 to 3)     | 115. (b)           | 116. (d)           |
| 117. (1 to 1)      | 118. (3 to 3)     | 119. (b)           | 120. (d)           |
| 121. (d)           | 122. (12 to 12)   | 123. (1.0)         | 124. (d)           |
| 125. (d)           | 126. (5.5 to 5.5) | 127. (c)           | 128. (b)           |
| 129. (5)           | 130. (a)          | 131. (c)           | 132. (b)           |
| 133. (5 to 5)      | 134. (a)          | 135. (c)           | 136. (a)           |
| 137. (2 to 2)      | 138. (d)          | 139. (1)           | 140. (d)           |
| 141. (15 to 15)    | 142. (a)          | 143. (d)           | 144. (3)           |

- |                 |                 |                      |                      |
|-----------------|-----------------|----------------------|----------------------|
| <b>145.</b> (b) | <b>146.</b> (a) | <b>147.</b> (9)      | <b>148.</b> (d)      |
| <b>149.</b> (d) | <b>150.</b> (d) | <b>151.</b> (6 to 6) | <b>152.</b> (b)      |
| <b>153.</b> (b) | <b>154.</b> (b) | <b>155.</b> (2 to 2) | <b>156.</b> (d)      |
| <b>157.</b> (d) | <b>158.</b> (a) | <b>159.</b> (0.33)   | <b>160.</b> (0 to 0) |
| <b>161.</b> (0) | <b>162.</b> (a) | <b>163.</b> (b)      | <b>164.</b> (c)      |
| <b>165.</b> (d) | <b>166.</b> (d) | <b>167.</b> (a)      | <b>168.</b> (b)      |
| <b>169.</b> (b) | <b>170.</b> (d) | <b>171.</b> (a)      | <b>172.</b> (c)      |
| <b>173.</b> (b) | <b>174.</b> (c) | <b>175.</b> (d)      | <b>176.</b> (b)      |
| <b>177.</b> (c) | <b>178.</b> (d) | <b>179.</b> (b)      | <b>180.</b> (a)      |
| <b>181.</b> (c) | <b>182.</b> (b) | <b>183.</b> (c)      | <b>184.</b> (a)      |
| <b>185.</b> (c) | <b>186.</b> (2) |                      |                      |

## SOLUTIONS

**1. (b)**As per Cholesky decomposition;  $L L^T = A$ 

$$\begin{bmatrix} L_{11} & 0 \\ L_{21} & L_{22} \end{bmatrix} \begin{bmatrix} L_{11} & L_{21} \\ 0 & L_{22} \end{bmatrix} = \begin{bmatrix} 8 & -5 \\ -5 & 0_{22} \end{bmatrix}$$

Let  $L$  be the lower triangular matrix =  $\begin{bmatrix} L_{11} & 0 \\ L_{21} & L_{22} \end{bmatrix}$

$$\Rightarrow L_{11} = 2\sqrt{2} \quad L_{21}^2 + L_{22}^2 a_{22}$$

$$\Rightarrow L_{11} L_{21} = -5 \left( -\frac{5}{2\sqrt{2}} \right)^2 + (1.968)^2 = a_{22}$$

$$L_{21} = -\frac{5}{2\sqrt{2}}$$

$$a_{22} = 6.99 \approx 7$$



Scan for Video solution




$$\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = 1$$

$$x + 2y + 3z = 1$$

$$f \rightarrow x + 2y + 3z - 1$$

$$\text{grad } f \rightarrow \frac{\partial f}{\partial x} \hat{i} + \frac{\partial f}{\partial y} \hat{j} + \frac{\partial f}{\partial z} \hat{k} = \hat{i} + 2\hat{j} + 3\hat{k} \quad [1 \ 2 \ 3]^T$$

▶
Scan for Video solution

**3. (a)**

For anti-clockwise rotation by  $\theta$   $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$

But for clockwise rotation by  $\theta$   $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$

**2. (d)**

$$w^T x = 1$$


$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}^T \begin{bmatrix} x \\ y \\ z \end{bmatrix} = 1$$

$(\theta < 0)$

$$\therefore A = \begin{bmatrix} \cos 36.86^\circ & \sin 36.86^\circ \\ -\sin 36.86^\circ & \cos 36.86^\circ \end{bmatrix}$$

$$A = \begin{bmatrix} 4/5 & 3/5 \\ -3/5 & 4/5 \end{bmatrix}$$

$$\therefore AU = v$$




4. (b)

$$\text{Given } A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 4 & 1 & 2 & 3 \\ 3 & 4 & 1 & 2 \\ 2 & 3 & 4 & 1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 3 & 4 & 1 & 2 \\ 4 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \end{bmatrix}$$

'B' is obtained by interchanging 1<sup>st</sup> and 3<sup>rd</sup> rows of 'A'. If two rows (or) columns of a determinant are interchanged then determinant changes its sign.

$$\Rightarrow \det(B) = -\det(A)$$



5. (c)


Given,

order of 'A' is  $m \times n$ ; order of 'B' is ' $n \times m$ '

Order of 'C' is  $n \times m$ ; order of 'O' is ' $n \times m$ '

For any two matrices A and B, if both AB and BA exist, then  $\text{tr}(AB) = \text{tr}(BA)$

$\therefore$  Both statements 1 and statements 2 are correct



6. (a)

Let the coordinate of point P in transformed system is

$$(x', y', z') \xrightarrow[\text{about x-axis}]{45^\circ \text{ rotation}} (x, y, z)$$


The potential matrix for transformation about x-axis

$$\begin{bmatrix} x' \\ y' \\ z' \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

Since angle of rotation  $\theta = 45^\circ$

$$\begin{bmatrix} x' \\ y' \\ z' \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ \sqrt{2} \end{bmatrix}$$

The transformed coordinates of P is  $(1, 0, \sqrt{2})$



7. (32)

$$\text{Given } E = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} P_1 \text{ Let eigen values of } E \text{ be } \lambda_1 \text{ and } \lambda_2.$$

$$\lambda_1 + \lambda_2 = 0$$

$$\lambda_1 \lambda_2 = -2$$

$$\therefore \lambda_1 = \sqrt{2}; \lambda_2 = -\sqrt{2}$$

$$\therefore \text{Eigen values of } E^8 = (\sqrt{2})^8 \text{ and } (-\sqrt{2})^8 = 16 \text{ and } 16$$


$$\therefore \text{Eigen values of } E^{11} = (\sqrt{2})^{11} \text{ and } (-\sqrt{2})^{11} = 32\sqrt{2} \text{ and } -32\sqrt{2}$$

$$\therefore P = \text{Trance}(E^8) = 16 + 16 = 32$$

$$\therefore Q = \text{Trance}(E^{11}) = 32\sqrt{2} - 32\sqrt{2} = 0$$

$$\therefore P + Q = 32 + 0$$

$$P + Q = 32$$



8. (a)

**Given:**

Matrix  $A = \begin{bmatrix} 10 & 2k+5 \\ 3k-3 & k+5 \end{bmatrix}$  is symmetric.

$$\Rightarrow 2k+5 = 3k-3 \Rightarrow k=8$$

 Scan for Video solution 

9. (c)

$$\begin{aligned} Q^T P^T &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^T \\ &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \end{aligned}$$

 Scan for Video solution 

10. (a)

If  $A A^T = I$  or  $A^T = A^{-1}$  then the matrix is said to be orthogonal

 Scan for Video solution 

11. (a)

- Matrix multiplication is associative.
  - Matrix multiplication NEED not be commutative.
  - A matrix 'A' is said to be definite if for any non-zero column vector 'z',  $z^T A z$  is always definite.
- In actual cases,  $z^T A z$  need not be positive always.

 Scan for Video solution 

12. (b)

- Every square matrix 'P' can be decomposed to symmetric and skew-symmetric parts.

$$\Rightarrow P = S + V$$

$$\Rightarrow P = \begin{bmatrix} -4 & 4 & 2 \\ 4 & 3 & 7/2 \\ 2 & 7/2 & 2 \end{bmatrix} + \begin{bmatrix} 0 & -2 & 3 \\ 2 & 0 & 7/2 \\ -3 & -7/2 & 0 \end{bmatrix}$$

$$\Rightarrow P = \begin{pmatrix} -4 & 2 & 5 \\ 6 & 3 & 7 \\ -1 & 0 & 2 \end{pmatrix}$$

 Scan for Video solution 

13. (marks to all)

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} a_{11} & 0 & 0 \\ a_{12} & a_{22} & 0 \\ a_{13} & a_{23} & a_{33} \end{bmatrix} \begin{bmatrix} a_{11} & 0 & 0 \\ a_{12} & a_{22} & 0 \\ a_{13} & a_{23} & a_{33} \end{bmatrix}^T$$

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} a_{11} & 0 & 0 \\ a_{12} & a_{22} & 0 \\ a_{13} & a_{23} & a_{33} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix}^{-1}$$

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} a_{11}^2 & a_{11}a_{12} & a_{11}a_{13} \\ a_{11}a_{12} & a_{12}^2 + a_{22}^2 & a_{12}a_{13} + a_{22}a_{23} \\ a_{11}a_{13} & a_{12}a_{13} + a_{22}a_{23} & a_{13}^2 + a_{23}^2 + a_{33}^2 \end{bmatrix}$$

$$\Rightarrow a_{11}^2 = 2 \Rightarrow a_{11} = \sqrt{2}$$

$$\Rightarrow a_{11}a_{12} = 3 \Rightarrow \sqrt{2} a_{12} = 3 \Rightarrow a_{12} = 3/\sqrt{2}$$

$$\Rightarrow a_{11}a_{13} = 3 \Rightarrow \sqrt{2} a_{13} = 3 \Rightarrow a_{13} = 3/\sqrt{2}$$

$$\Rightarrow a_{12}^2 + a_{12}^2 = 2 \Rightarrow (3/\sqrt{2})^2 + a_{22}^2 = 2$$

$$\Rightarrow a_{13} = \sqrt{-\frac{\sqrt{5}}{2}} = 1.58i$$

$$\Rightarrow a_{11}a_{13} + a_{22}a_{23} = 1$$

$$\sqrt{2} \times \frac{3}{\sqrt{2}} + (1.58i)a_{23} = 1$$

$$a_{23} = 2.21i$$

$$\Rightarrow a_{13}^2 + a_{23}^2 + a_{33}^2 = 7$$

$$\left(\frac{3}{\sqrt{2}}\right)^2 + (2.21i)^2 + a_{33}^2 = 7$$

$$a_{33} = \sqrt{7.38}$$

Now,

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} \sqrt{2} & 0 & 0 \\ 3/\sqrt{2} & 1.58i & 0 \\ 3/\sqrt{2} & 2.21i & \sqrt{7.38} \end{bmatrix} \begin{bmatrix} \sqrt{2} & 0 & 0 \\ 3/\sqrt{2} & 1.58i & 0 \\ 3/\sqrt{2} & 2.21i & \sqrt{7.38} \end{bmatrix}^T$$

$$[A = LL^T]$$

∴ Number of purely real elements in matrix L = 7  
Final answer is not matching with any option hence this question was **marks to all**.



Scan for Video solution



14. (d)

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 4 & 3 & 1 \\ 1 & 2 & 4 \end{bmatrix}$$

$$\text{We know, } A^{-1} = \frac{\text{adj } A}{|A|}$$

Adj. (A) (Cofactor matrix)<sup>T</sup>

$$\begin{bmatrix} 10 & -15 & 5 \\ -4 & 4 & -1 \\ -9 & 14 & -6 \end{bmatrix}^T = \begin{bmatrix} 10 & -4 & -9 \\ -15 & 4 & 14 \\ 5 & -1 & -6 \end{bmatrix}$$

$$|A| = 2(12 - 2) - 2(16 - 1) + 4(8 - 3) = 2 \times 10 - 3 \times 15 + 4 \times 5 = -5$$

$$A^{-1} = \frac{\text{Adj}}{|A|} = -\frac{1}{5} \begin{bmatrix} 10 & -4 & -9 \\ -15 & 4 & 14 \\ 5 & -1 & -6 \end{bmatrix} = \begin{bmatrix} -2 & \frac{4}{5} & \frac{9}{5} \\ 3 & -\frac{4}{5} & -\frac{14}{5} \\ -1 & \frac{1}{5} & \frac{6}{5} \end{bmatrix}$$



Scan for Video solution



15. (d)

For the inverse of matrix to exist, the matrix should be non-singular

(i.e.) determinant is non-zero

∴ I and II are equivalent statements



Scan for Video solution



16. (c)

$$\text{Let } M_{2 \times 2} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} V_1 & V_2 \end{bmatrix}$$

$$V_1 = \begin{bmatrix} a \\ c \end{bmatrix} \quad V_2 = \begin{bmatrix} b \\ d \end{bmatrix}$$

$$\therefore M^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \begin{bmatrix} U_1^T \\ U_2^T \end{bmatrix}$$

$$U_1^T = \begin{bmatrix} \frac{d}{ad - bc} & \frac{-b}{ad - bc} \end{bmatrix}, U_2^T = \begin{bmatrix} \frac{-c}{ad - bc} & \frac{a}{ad - bc} \end{bmatrix}$$

Now,

$$U_1^T \cdot V_1 = \begin{bmatrix} \frac{d}{ad - bc} & \frac{-b}{ad - bc} \end{bmatrix} \begin{bmatrix} a \\ c \end{bmatrix} = \frac{ad - bc}{ad - bc} = 1$$

$$U_2^T \cdot V_2 = \begin{bmatrix} \frac{-c}{ad - bc} & \frac{a}{ad - bc} \end{bmatrix} \begin{bmatrix} b \\ d \end{bmatrix} = \frac{ad - bc}{ad - bc} = 1$$

$$\therefore U_1^T \cdot V_1 = 1 \text{ and } U_2^T \cdot V_2 = 1$$

Hence, statement 1 is true.

Now,

$$U_1^T \cdot V_2 = \begin{bmatrix} \frac{d}{ad - bc} & \frac{-b}{ad - bc} \end{bmatrix} \begin{bmatrix} b \\ d \end{bmatrix} = \frac{bd - bd}{ad - bc} = 0$$

$$U_2^T \cdot V_1 = \begin{bmatrix} \frac{-c}{ad - bc} & \frac{a}{ad - bc} \end{bmatrix} \begin{bmatrix} a \\ c \end{bmatrix} = \frac{-ca + ca}{ad - bc} = 0$$

$$\therefore U_1^T \cdot V_2 = 0 \text{ and } \therefore \rho(A^T A) = 2$$

Hence statement 2 is also true.



Scan for Video solution



17. (c)

Since matrix Q is orthogonal hence

$$Q Q^T = I$$

Pre multiply by  $Q^{-1}$ 

$$Q^{-1} Q Q^T = Q^{-1} I$$

$$Q^T = Q^{-1}$$

$$\text{Hence } Q^{-1} = Q^T = \begin{bmatrix} \frac{3}{7} & -\frac{6}{7} & \frac{2}{7} \\ \frac{2}{7} & \frac{3}{7} & \frac{6}{7} \\ \frac{6}{7} & \frac{2}{7} & -\frac{3}{7} \end{bmatrix}$$



Scan for Video solution



18. (c)

If  $|A| = 0$  (Singular matrix)

**Option a:-**  $\Delta = 6 - 5 = 1 \neq 0$

**Option b:-**  $\Delta = 9 - 4 = 5 \neq 0$

**Option c:-**  $\Delta = 12 - 12 = 0$  Hence it is singular

**Option d:-**  $\Delta = 8 - 18 = 10 \neq 0$



Scan for Video solution



19. (0.25 to 0.25)

**Given,**

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 1 \end{bmatrix}$$

 $\Rightarrow$  'A' is an upper triangular matrix.

For any upper triangular matrix, determinant is product of principal diagonal elements.

$$\Rightarrow |A| = (1)(4)(1) = 4$$

Now  $|A^{-1}| = \frac{1}{|A|} = \frac{1}{4} = 0.25$

$$\therefore |A^{-1}| = 0.25$$



Scan for Video solution



20. (a)

$$\begin{aligned} AB^T &= \begin{bmatrix} 1 & 5 \\ 6 & 2 \end{bmatrix} \begin{bmatrix} 3 & 7 \\ 8 & 4 \end{bmatrix}^T \\ &= \begin{bmatrix} 1 & 5 \\ 6 & 2 \end{bmatrix} \begin{bmatrix} 3 & 8 \\ 7 & 4 \end{bmatrix} = \begin{bmatrix} 38 & 28 \\ 32 & 56 \end{bmatrix} \end{aligned}$$



Scan for Video solution



21. (c)

Given:-  $P = Q^{-1}$ 

- Post multiply by Q

$$PQ = Q^{-1} Q \quad [\because A^{-1} A = I]$$

$$PQ = I$$

- Pre multiply by Q

$$QP = Q Q^{-1} \quad [\because AA^{-1} = I]$$

$$QP = I$$

Hence  $PQ = I$  and  $QP = I$ 

Scan for Video solution



22. (0 to 0)

**Given:** Matrix  $A = \begin{bmatrix} 50 & 70 \\ 70 & 80 \end{bmatrix}$

'A' is symmetric matrix

If  $x_1, x_2$  are eigen vectors of 'A' corresponding to  $\lambda_1$  and  $\lambda_2$ , then

$$x_1^T \cdot x_2 = 0 \quad (\text{In general, } x_i^T \cdot x_j = 0 \ \forall i \neq j)$$

**OR**

Given,  $x_1 = \begin{bmatrix} 70 \\ \lambda_1 - 50 \end{bmatrix}; \quad x_2 = \begin{bmatrix} \lambda_2 - 80 \\ 70 \end{bmatrix}$

$$\begin{aligned} \Rightarrow x_1^T \cdot x_2 &= [70 \ \lambda_1 - 50] \begin{bmatrix} \lambda_2 - 80 \\ 70 \end{bmatrix} \\ &= 70(\lambda_2 - 80) + 70(\lambda_1 - 50) \\ &= 70(\lambda_1 + \lambda_2 - 130) \\ &= 70(\text{Sum of diagonal elements} - 130) \\ &= 70(130 - 130) = 0 \end{aligned}$$

$$\Rightarrow x_1^T \cdot x_2 = 0$$



Scan for Video solution



23. (d)

Given

$$P = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{-1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\Rightarrow |P| = \frac{1}{\sqrt{2}} \left( \frac{1}{\sqrt{2}} - 0 \right) + \frac{1}{\sqrt{2}} \left( \frac{1}{\sqrt{2}} \right) = \frac{1}{2} + \frac{1}{2} = 1$$

 $P \cdot P^T$  is given by

$$= \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{-1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & -1 \\ 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Since  $P \cdot P^T = I \Rightarrow P$  is orthogonal  $\Rightarrow P^{-1} = P^T$ 

An orthogonal matrix need not have all the eigen values being real.

Scan for Video solution



24. (7)

$$2x^2 + y^2 = 34$$

.....(1)

$$x + 2y = 11$$

$$y = \frac{11-x}{2}$$

... (2)

From value of  $y$  from equation (2) to equation (1),

$$2x^2 + \left( \frac{11-x}{2} \right)^2 = 34$$

$$\Rightarrow 8x^2 + 121 + x^2 - 22x = 136$$

$$\Rightarrow 9x^2 - 22x - 15 = 0$$

$$\Rightarrow x = \frac{-10}{18}, 3$$

Discarding  $x = -10/18$  because it is not an integer.Hence  $x = 3$ .Put value of  $x$  in equation (2),

$$y = \frac{11-x}{2} = \frac{11-3}{2} = 4$$

Hence,

$$\begin{aligned} x + y &= 3 + 4 \\ \Rightarrow x + y &= 7 \end{aligned}$$

Scan for Video solution



25. (c)

A square matrix 'A' is said to be Skew-symmetric if  $A = -A^T$ 

$$\Rightarrow A^T = -A$$

Scan for Video solution



26. (3)

Let the given matrix is

$$A = \begin{bmatrix} a & 0 & 3 & 7 \\ 2 & 5 & 1 & 3 \\ 0 & 0 & 2 & 4 \\ 0 & 0 & 0 & b \end{bmatrix}; |A| = 100, \text{tr}(A) = 14$$

$$\text{tr}(A) = a + 5 + 2 + b = 14$$

$$\Rightarrow a + b = 7$$

$$|A| = \begin{bmatrix} a & 0 & 3 & 7 \\ 2 & 5 & 1 & 3 \\ 0 & 0 & 2 & 4 \\ 0 & 0 & 0 & b \end{bmatrix} =$$

$$|A| = a \begin{vmatrix} 5 & 1 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & b \end{vmatrix} + 3 \begin{vmatrix} 2 & 5 & 3 \\ 0 & 0 & 4 \\ 0 & 0 & b \end{vmatrix} - 7 \begin{vmatrix} 2 & 5 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{vmatrix}$$

$$= a(10b) = 10ab$$

$$\Rightarrow 10ab = 100 \Rightarrow ab = 10$$

$$\begin{aligned} \text{Since } (a-b)^2 &= (a+b)^2 - 4ab \\ &= (7)^2 - 40 = 9 \end{aligned}$$

$$\Rightarrow (a-b)^2 = 9$$

$$\Rightarrow |(a-b)| = 3$$

Scan for Video solution



27. (c)

$$\text{Given } M^4 = I \Rightarrow (M^4)^k = I^k = I \Rightarrow M^{4k} = I$$

Multiplying both sides with  $M^{-1}$

$$\Rightarrow M^3 = M^{-1}$$

$$\Rightarrow M^{-1} = M^3 = M^3 \cdot I$$

$$\Rightarrow M^{-1} = M^3 \cdot M^{4k}$$

$$\Rightarrow M^{-1} = M^{4k+3}$$



Scan for Video solution



28. (233 to 233)

Given that a sequence is defined by  $x[n]$  and it is given by

$$\begin{bmatrix} x[n] \\ x[n-1] \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\text{And } x[0] = 1; x[1] = 1$$

$$\text{For } n=2, \begin{bmatrix} x[2] \\ x[1] \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\therefore x[2] = 2$$

Given sequence is a Fibonacci sequence

$\therefore$  The given sequence is equal to

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, \dots$$

$$\therefore x[12] = 233$$

$$\therefore 13^{\text{th}} \text{ term of the sequence} = x[12] = 233$$



Scan for Video solution



29. (d)

$$\text{Given } \begin{pmatrix} a \\ b \end{pmatrix} = P \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$3x + y = a \dots \text{(i)}$$

$$x + 3y = b \dots \text{(ii)}$$

On squaring (i) and (ii) and then adding, we get

$$a^2 + b^2 = (3x + y)^2 + (x + 3y)^2 = 1$$

$$9x^2 + y^2 + 6xy + x^2 + 9y^2 + 6xy = 1$$

$$10x^2 + 10y^2 + 12xy = 1 \dots \text{(i)}$$

On comparing with std from  $ax^2 + by^2 + 2hxy + 2gx + 2fy + c = 0$


$$\therefore a = 10, b = 10, h = 6$$

$h^2 - ab = 36 - 10 \times 10 = -64 < 0 \Rightarrow$  Ellipse

$\therefore$  Equation (i) represents an ellipse

With major axis along  $y = -x \approx \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  and with

minor axis along  $y = x \approx \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .



Scan for Video solution



30. (0.125)

Given, eigen values of  $A = 1, 2, 4$


$$\Rightarrow |A| = \text{product of eigen values} = 1 \cdot 2 \cdot 4 = 8$$

Determinant of  $(A^{-1})^T$

$$= \text{Determinant of } (A^{-1}) = \frac{1}{|A|} = \frac{1}{8} = 0.125$$



Scan for Video solution



31. (5)

$$\text{Let, } A = \begin{bmatrix} 2 & 2 \\ 4 & 9 \end{bmatrix}$$

LU decomposition  $\Rightarrow A = LU$

$$\Rightarrow \begin{bmatrix} 2 & 2 \\ 4 & 9 \end{bmatrix} = \begin{bmatrix} L_{11} & 0 \\ L_{12} & L_{22} \end{bmatrix} \begin{bmatrix} 1 & U_{12} \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 2 \\ 4 & 9 \end{bmatrix} = \begin{bmatrix} L_{11} & L_{11}U_{12} \\ L_{12} & L_{12}U_{12} + L_{22} \end{bmatrix}$$

$$L_{11} = 2, L_{12} = 4, U_{12} = 1$$

$$\begin{aligned} L_{12}U_{12} + L_{22} &= 9 \\ \Rightarrow L_{22} &= 9 - (4 \times 1) = 5 \\ \therefore L_{22} &= 5 \end{aligned}$$



Scan for Video solution

**32. (a)**

When two rows (or) two columns of a determinants are interchanged, then the absolute value of the determinant does not change but the sign of the value changes.



Scan for Video solution

**33. (a, b)**

If P and Q are invertible, then  
 $(PQ)^{-1} = Q^{-1}P^{-1}$  and  $(QP)^{-1} = P^{-1}Q^{-1}$   
→ If P and Q are not invertible, then  
 $(PQ)^{-1} \neq Q^{-1}P^{-1}$   
 $(QP)^{-1} \neq P^{-1}Q^{-1}$



Scan for Video solution

**34. (a)**

For a matrix A =  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\Rightarrow P = \begin{bmatrix} 4+3i & -i \\ i & 4-3i \end{bmatrix},$$

$$P^{-1} = \frac{1}{(4+3i)(4-3i)+i^2} \begin{bmatrix} 4-3i & i \\ -i & 4+3i \end{bmatrix}$$

$$\Rightarrow P^{-1} = \frac{1}{24} \begin{bmatrix} 4-3i & i \\ -i & 4+3i \end{bmatrix}$$



Scan for Video solution

**35. (a)****Given**


$$\begin{vmatrix} 1 & 3 & 0 \\ 2 & 6 & 4 \\ -1 & 0 & 2 \end{vmatrix} = -12$$

$$\begin{vmatrix} 2 & 6 & 0 \\ 4 & 12 & 8 \\ -2 & 0 & 4 \end{vmatrix} = \begin{vmatrix} 2 \times 1 & 2 \times 3 & 2 \times 0 \\ 2 \times 2 & 2 \times 6 & 2 \times 4 \\ 2 \times -1 & 2 \times 0 & 2 \times 2 \end{vmatrix}$$

$$= 2 \times 2 \times 2 \begin{vmatrix} 1 & 3 & 0 \\ 2 & 6 & 4 \\ -1 & 0 & 2 \end{vmatrix} = 8 \times (-12) = -96$$

$$\left\{ \begin{vmatrix} k_1 a_1 & k_1 a_2 & k_1 a_3 \\ k_2 b_1 & k_2 b_2 & k_2 b_3 \\ k_3 c_1 & k_3 c_2 & k_3 c_3 \end{vmatrix} = k_1 \cdot k_2 \cdot k_3 \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \right\}$$

$$\therefore \begin{vmatrix} 2 & 6 & 0 \\ 4 & 12 & 8 \\ -2 & 0 & 4 \end{vmatrix} = -96$$

**36. (d)**

$$(i) P(Q + R) = PQ + PR$$


PR need not be equal to RP since matrix multiplication is not commutative.

$$(ii) (P - Q)^2 = (P - Q)(P - Q) = P^2 - QP - PQ + Q^2$$

(PQ need not be equal to QP)

$$(iii) |P + Q| \neq |P| + |Q| \text{ (Since } |P + Q| \leq |P| + |Q|)$$

$$(iv) (P + Q)^2 = (P + Q)(P + Q) = P^2 + QP + PQ + Q^2$$

**37. (49)**

Let the  $2 \times 2$  symmetric matrix = A =  $\begin{bmatrix} a & b \\ b & c \end{bmatrix}$

$$\text{Given } \text{tr}(A) = 14 \Rightarrow a + c = 14 \Rightarrow c = 14 - a$$

$$\text{Maximum value of } |A| = ac - b^2$$

$$\text{For } |A| \text{ to be maximum, } b^2 = 0 \text{ (}\because b^2 \geq 0\text{)}$$

$$\therefore |A| = ac = a(14 - a)$$

For  $|A|$  to be maximum,

$$\frac{d}{da}\{a(14-a)\} = 0$$

$$\Rightarrow 14 - 2a = 0$$

$$\Rightarrow a = 7$$

Considering  $\left. \frac{d^2 |A|}{da^2} \right|_{a=7} = -2 < 0$

$\therefore |A|$  is maximum at  $a = 7$

$$\Rightarrow c = 14 - a \Rightarrow c = 7$$

$$\therefore |A|_{\max} = ac = 7(7) = 49$$



Scan for Video solution



38. (1)

$$J_6 = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Given  $P = I_6 + \alpha J_6$

$$\Rightarrow P = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & \alpha \\ 0 & 1 & 0 & 0 & \alpha & 0 \\ 0 & 0 & 1 & \alpha & 0 & 0 \\ 0 & 0 & \alpha & 1 & 0 & 0 \\ 0 & \alpha & 0 & 0 & 1 & 0 \\ \alpha & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_6 \rightarrow R_6 - \alpha R_1$$

$$R_5 \rightarrow R_5 - \alpha R_2$$

$$R_4 \rightarrow R_4 - \alpha R_3$$

$$\Rightarrow P \sim \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & \alpha \\ 0 & 1 & 0 & 0 & \alpha & 0 \\ 0 & 0 & 1 & \alpha & 0 & 0 \\ 0 & 0 & 0 & 1 - \alpha^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 - \alpha^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 - \alpha^2 \end{bmatrix}$$

The matrix got converted to upper triangular matrix.

$\Rightarrow |P| = \text{Product of Principal diagonal elements}$

$$= (1 - \alpha^2)^3 = 0$$

$$\Rightarrow 1 - \alpha^2 = 0$$

$$\Rightarrow \boxed{\alpha = +1} \text{ (Since } \alpha \text{ is non-negative)}$$



Scan for Video solution



39. (d)

(i)  $(M^T)^T = M$  (Always hold)

(ii)  $(cM)^T = cM^T$  (Always hold)

(iii)  $(M + N)^T = M^T + N^T$  (Always hold)

(iv)  $MN$  need not be equal to  $NM$  since matrix multiplication need not be commutative

$\therefore$  option 4 does not Always hold



Scan for Video solution



40. (200)

Given  $|A| = 5$  and  $|B| = 40$

$$\therefore |AB| = |A||B| = 5(40) = 200$$

$$\therefore |AB| = 200$$



Scan for Video solution



41. (c)

Given function is

$$f(\theta) = \begin{vmatrix} \sin\theta & \cos\theta & \tan\theta \\ \frac{1}{2} & \frac{\sqrt{3}}{2} & \frac{1}{\sqrt{3}} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & \sqrt{3} \end{vmatrix}$$

$$\Rightarrow f(\theta) = \begin{vmatrix} \cos\theta & -\sin\theta & \sec^2\theta \\ \frac{1}{2} & \frac{\sqrt{3}}{2} & \frac{1}{\sqrt{3}} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} & \sqrt{3} \end{vmatrix}$$

$$= k_1 \cdot \cos\theta + k_2 \cdot \sin\theta + k_3 \cdot \sec^2\theta$$

Where

$k_1, k_2$  and  $k_3$  are constants

By LaGrange's' mean value theorem

$\exists$  at least one  $\theta \in \left[ \frac{\pi}{6}, \frac{\pi}{3} \right]$  such that

$$f'(\theta) = \frac{f\left(\frac{\pi}{3}\right) - f\left(\frac{\pi}{6}\right)}{\frac{\pi}{3} - \frac{\pi}{6}}$$

Since  $f\left(\frac{\pi}{3}\right) = f\left(\frac{\pi}{6}\right) = 0$  because two rows become

identical for  $\theta = \frac{\pi}{3}$  and  $\theta = \frac{\pi}{6}$

$\therefore \exists$  some ' $\theta$ ' where  $\theta \in \left[ \frac{\pi}{6}, \frac{\pi}{3} \right]$  such that  $f'(\theta) = 0$

Also, since  $f'(\theta)$  is not an identity,  $\exists$  some ' $\theta$ ' where  $\theta \in \left[ \frac{\pi}{6}, \frac{\pi}{3} \right]$  such that  $f'(\theta) \neq 0$

$\therefore$  Both statements are true.



Scan for Video solution



#### 42. (23 to 23)

$$J = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 6 \end{bmatrix} \quad k = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

$$K^T = [1 \ 2 \ -1]$$

$$K^T \cdot J \cdot K = [1 \ 2 \ -1] \begin{bmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 6 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} = [6 \ 8 \ -1]$$

$$[1 \ 2 \ -1] \begin{bmatrix} 6 \\ 8 \\ -1 \end{bmatrix}$$


$$= 6 + 16 + 1 = 23.$$



Scan for Video solution



#### 43. (a)



$$\text{Area of triangle} = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

$$= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

$$= \frac{1}{2} [-1 + 6 - 8] = \left| \frac{-3}{2} \right| = \frac{3}{2}$$

$$\boxed{\text{Area of } \Delta = \frac{3}{2}}$$

#### 44. (88 to 88)

$$\left| \begin{array}{cccc} 0 & 1 & 2 & 3 \\ 1 & 0 & 3 & 0 \\ 2 & 3 & 0 & 1 \\ 3 & 0 & 1 & 2 \end{array} \right| \xrightarrow{R_3 \rightarrow R_3 - 3R_1} \left| \begin{array}{cccc} 0 & 1 & 2 & 3 \\ 1 & 0 & 3 & 0 \\ 2 & 0 & -6 & -8 \\ 3 & 0 & 1 & 2 \end{array} \right|$$

Interchange column 1 and column 2 and expand about first column.

$$\begin{aligned} \left| \begin{array}{cccc} 1 & 0 & 2 & 3 \\ 0 & 1 & 3 & 0 \\ 0 & 2 & -6 & -8 \\ 0 & 3 & 1 & 2 \end{array} \right| &= -1 \begin{vmatrix} 1 & 3 & 0 \\ 2 & -6 & -8 \\ 3 & 1 & 2 \end{vmatrix} = \{1(-12 + 8) - 3(4 + 24) + 0(2 + 18)\} \\ &- \{-4 - 84 + 0\} = \boxed{88} \end{aligned}$$



Scan for Video solution



45. (a)

A square matrix is defined as skew – Symmetric matrix if and only if  $A^T = -A$

$$\text{Ex: } A = \begin{bmatrix} o & h & g \\ -h & o & f \\ -g & -f & o \end{bmatrix}$$

$$A^T = \begin{bmatrix} o & -h & -g \\ h & o & -f \\ g & f & o \end{bmatrix}$$

$$A^T = -\begin{bmatrix} o & h & g \\ -h & o & f \\ -g & -f & o \end{bmatrix}$$

$$A^T = -A$$



Scan for Video solution



46. (b)

$$\text{Given matrix is (Say C)} = \begin{bmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{bmatrix}$$

Converting 'C' to  $(I+AB)$  form

$$\Rightarrow \begin{bmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

Since both the matrices C and  $I+AB$  are same, the determinant is same.

Writing the matrix AB as Product of a  $4 \times 1$  and  $1 \times 4$  matrices,

$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}$$

$$\Rightarrow BA = \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = [4]_{1 \times 1}$$

$$\therefore I_{1 \times 1} + BA = [5] \Rightarrow |I_{1 \times 1} + BA| = 5$$

Since  $|c| = \text{determinant } (I_4 + AB) = \text{determinant } (I_1 + BA) = 5$

By above property, determinant of given Matrix is 5.



Scan for Video solution



47. (16)

$$P \rightarrow 4 \times 2 (m \times p)$$

$$Q \rightarrow 2 \times 2 (p \times n)$$

$$R \rightarrow 4 \times 1 (n \times p)$$

$$\text{Total number of multiplications} = m \cdot np$$

$$PQ \rightarrow 4 \times 2$$

$$QR \rightarrow 2 \times 4$$

$$2 \times 4$$

$$4 \times 1$$

$$= 4 \times 4 = 16$$

$$= 2 \times 1$$

$$m \times n$$

$$m \times n = 2$$

$$PQ = mnp = 16 \times 2 = 32 \quad mnp = 2 \times 4 = 8$$

$$PQR \rightarrow 4 \times 4 = 4 \times 1 \quad PQR = 4 \times 2 = 2 \times 1 \\ = (4 \times 1) \quad = 4 \times 1 = 4$$

$$mnp = 4 \times 4 = 16$$

$$mnp = 2 \times 4 = 8$$

$$pmn + mnp$$

$$mnp + mnp$$

$$= 16 + 32 = 48$$

$$= 8 + 8 = 16$$

Minimum number of multiplications = 16.



Scan for Video solution



48. (b)

$$\text{Given matrix is } A = \begin{bmatrix} -5 & -3 \\ 2 & 0 \end{bmatrix};$$

Characteristic equation of 'A' is given by

$$\begin{vmatrix} -5 - \lambda & -3 \\ 2 & -\lambda \end{vmatrix} = 0$$

$$\Rightarrow -(5 + \lambda) \cdot (+\lambda) + 6 = 0$$

$$\Rightarrow \lambda^2 + 5\lambda + 6 = 0$$

$\therefore$  By cayley-Hamilton theorem,

$$\begin{aligned} A^2 + 5A + 6I &= 0 \quad \dots(1) \\ \Rightarrow A^2 &= -5A - 6I \quad \dots(2) \\ (1) \Rightarrow A^2 + 5A + 6I &= 0 \\ \Rightarrow A(A^2 + 5A + 6I) &= 0 \\ \Rightarrow A^3 + 5A^2 + 6A &= 0 \\ \Rightarrow A^3 + 5(-5A - 6I) + 6A &= 0 \\ \Rightarrow A^3 &= 19A + 30I \end{aligned}$$



Scan for Video solution

**49. (d)**

Since  $(A + A^T)^T = A^T + (A^T)^T$   
 $= A^T + A \quad [\because (A^T)^T = A]$   
 $A + A^T = (A + A^T)^T$

$[S] = [S]^T$  hence  $[S]$  is symmetric.

- Since  $(A - A^T)^T = A^T - (A^T)^T$   
 $= A^T - A \quad [\because (A^T)^T = A]$   
 $-(A - A^T) = (A - A^T)^T$   
 $-[D] = [D]^T$  hence  $[D]$  is skew symmetric.



Scan for Video solution

**50. (b)**

We know cube roots of unity are  $1, w, w^2$ .

$$1, w = -\frac{1}{2} + \frac{\sqrt{3}}{2}j, w^2 = -\frac{1}{2} - \frac{\sqrt{3}}{2}j$$

$$\vec{x} = [1 \ 1 \ 1] \quad \vec{x}_2 = [1 \ a \ a^2]$$

$$\vec{x}_1 \cdot \vec{x}_2^T = [1 \ 1 \ 1] \begin{bmatrix} 1 \\ a \\ a^2 \end{bmatrix}$$

$$= [1 + a + a^2]$$

$$\vec{x}_1 \cdot \vec{x}_2^T = 0$$

$\therefore x_1$  and  $x_2$  are orthogonal vectors.



Scan for Video solution

**51. (d)**

$$A = LU$$

$$\begin{bmatrix} 2 & 1 \\ 4 & -1 \end{bmatrix} = \begin{bmatrix} L_{11} & 0 \\ L_{21} & L_{22} \end{bmatrix} \begin{bmatrix} 1 & U_{12} \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 4 & -1 \end{bmatrix} = \begin{bmatrix} L_{11} & L_{11}L_{12} \\ L_{21} & L_{21}L_{12} + L_{22} \end{bmatrix}$$

$$\therefore L_{11} = 2$$

$$\therefore L_{21} = 4$$

$$L_{11}U_{12} = 1$$

$$\therefore U_{12} = 0.5$$

$$L_{21} = U_{12} + L_{21} = -1$$

$$4 \times 0.5 + L_{22} = -1$$

$$\therefore L_{22} = -3$$

$$\therefore L = \begin{bmatrix} 2 & 0 \\ 4 & -3 \end{bmatrix}$$

$$U = \begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix}$$



Scan for Video solution

**52. (b)**

$$\text{Let } A = \begin{bmatrix} 9 & 5 \\ 5 & 8 \end{bmatrix}$$

The characteristic equation  $|A - \lambda I| = 0$

$$\begin{bmatrix} 9-\lambda & 5 \\ 5 & 8-\lambda \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 0$$

$$\begin{vmatrix} 9-\lambda & 5 \\ 5 & 8-\lambda \end{vmatrix} = 0$$

$$(9-\lambda)(8-\lambda) - 5 \times 5 = 0$$

$$\lambda^2 - 17\lambda + 47 = 0$$

$$\lambda_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-(-17) \pm \sqrt{(-17)^2 - 4 \times 1 \times 47}}{2 \times 1}$$

$$\lambda = 3.48, 13.53$$



Scan for Video solution



53. (a)

Given:

$$\text{Matrix } M = \begin{bmatrix} 3 & 4 \\ 5 & 5 \\ x & 3 \\ 5 & 5 \end{bmatrix} \text{ and } M^T = M^{-1}$$

$\Rightarrow$  'M' is orthogonal matrix ( $\because MM^T = MM^{-1} = I$ )

$$\Rightarrow |M| = \pm 1$$

$$\text{Let } |M| = 1$$

$$\Rightarrow \frac{9}{25} - \frac{4x}{5} = 1$$

$$\Rightarrow \frac{4x}{5} = \frac{-16}{25} \Rightarrow x = \frac{-4}{5}$$

$$(\text{Or}) \text{ Let } |M| = -1$$

$$\Rightarrow \frac{9}{25} - \frac{4x}{5} = -1$$

$$\Rightarrow \frac{4x}{5} = \frac{34}{25} \Rightarrow x = \frac{17}{10} \text{ (Not in given options)}$$

$$\therefore \boxed{x = -4/5}$$



Scan for Video solution



54. (b)

We know that;

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{(ad - bc)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\text{Alternatively; } A^{-1} = \frac{\text{adj}A}{|A|}$$

$$\begin{bmatrix} 3+2i & i \\ -i & 3-2i \end{bmatrix} = \frac{1}{(3+2i)(3-2i) + i^2} \begin{bmatrix} 3-2i & -i \\ +i & 3+2i \end{bmatrix}$$

$$= \frac{1}{12} \begin{bmatrix} 3-2i & -i \\ i & 3+2i \end{bmatrix}$$



Scan for Video solution



55. (b)

$$(PQ)^{-1} P$$

$$[Q^{-1} P^{-1}] P$$

$$Q^{-1} (P^{-1} P)$$

$$Q^{-1} (I)$$

$$Q^{-1}$$

$$\{\therefore (AB)^{-1} = B^{-1} A^{-1}\}$$

$$(ABC)^{-1} = C^{-1} B^{-1} A^{-1}$$

Scan for Video solution



56. (b)

We know  $PP^T = I$  ( $\because P$  is orthogonal)Let  $P = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  which is an orthogonal matrix

$$P \cdot x = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$Px = x$$

$$\therefore \|Px\| = \|x\| = (x_1^2 + x_2^2)^{1/2} \text{ for all vectors } \vec{x}$$

Scan for Video solution



57. (d)

Given

$$A^1 = (A^T \cdot A) \cdot AT$$

$$A^1 = A^T \cdot (AT)^{-1} \cdot AT \quad \because (AB)^{-1} = B^{-1} A^{-1}$$

$$A \cdot A^1 = (A^T)^{-1} \cdot A^T$$

From equation

$$\left. \begin{array}{l} A \cdot A^1 = I \\ A \cdot A^1 A = I \end{array} \right\} \dots(i) \quad \left. \begin{array}{l} (A \cdot A^1)(A \cdot A^1) = A \cdot A^1 \\ (A \cdot A^1)^2 = A \cdot A^1 \end{array} \right\} \dots(ii)$$

Scan for Video solution



**58. (4)**Given P is a  $4 \times 5$  matrix.Every solution of  $Px = 0$  is a scalar multiple of  $[2 \ 5 \ 4 \ 3 \ 1]^T$ . $\Rightarrow [0 \ 0 \ 0 \ 0 \ 0]^T$  is the only independent (trivial) solution of  $Px = 0$ 

\therefore Number of linearly independent solutions = 1

\therefore Number of unknowns -  $\rho(P) = 1$ 

$$\Rightarrow 5 - \rho(P) = 1$$

$$\Rightarrow \rho(P) = 4$$

\therefore Rank of matrix 'P' is 4.



Scan for Video solution

**59. (b)**

$$A = \begin{bmatrix} 1 & 2 & 2 & 3 \\ 3 & 4 & 2 & 5 \\ 5 & 6 & 2 & 7 \\ 7 & 8 & 2 & 9 \end{bmatrix} \xrightarrow{\substack{R_2 \rightarrow R_2 - 3R_1 \\ R_3 \rightarrow R_3 - 5R_1 \\ R_4 \rightarrow R_4 - 4R_1}} \begin{bmatrix} 1 & 2 & 2 & 3 \\ 0 & -2 & -4 & -4 \\ 0 & -4 & -8 & -8 \\ 0 & -6 & -12 & -12 \end{bmatrix}$$

$$\xrightarrow{\substack{R_3 \rightarrow R_3 - 2R_2 \\ R_4 \rightarrow R_4 - 3R_2}} \begin{bmatrix} 1 & 2 & 2 & 3 \\ 0 & -2 & -4 & -4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

No of non-zero rows in echelon form = 2

Hence Rank of A = 2



Scan for Video solution

**60. (c)**

$$\begin{bmatrix} 5 & 0 & -5 & 0 \\ 0 & 2 & 0 & 1 \\ -5 & 0 & 5 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix} \xrightarrow{R_3 \rightarrow R_3 + R_1} \begin{bmatrix} 5 & 0 & -5 & 0 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}$$

$$\xrightarrow{R_4 \rightarrow R_4 - \frac{R_2}{2}} \begin{bmatrix} 5 & 0 & -5 & 0 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3/2 \end{bmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_4} \begin{bmatrix} 5 & 0 & -5 & 0 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 3/2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Number of non-zero rows in echelon form = 3

\therefore Rank of A = 3

 Scan for Video solution



**61. (3)**

$$|M| = 0 - 1(0 - 1) + 1(1 - 0) = 2$$

Since  $|M| \neq 0 \therefore \rho(M) = 3$ 

\therefore Rank = order of largest non-zero minor.

 Scan for Video solution



**62. (b)**

$$\text{We know, } \rho(A) = \rho(A^T A)$$

$$\therefore \rho(A^T A) = 2$$

 Scan for Video solution



**63. (b)****Given**

$$\begin{pmatrix} -4 & 1 & -1 \\ -1 & -1 & -1 \\ 7 & -3 & 1 \end{pmatrix} = A \text{ (Say)}$$

Converting 'A' to Row-echelon form

$$R_2 \rightarrow 4R_2 - R_1 \text{ and } R_3 \rightarrow 4R_3 + 7R_1$$

$$\Rightarrow A \sim \begin{bmatrix} -4 & 1 & -1 \\ 0 & -5 & -3 \\ 0 & -5 & -3 \end{bmatrix}$$

$$R_3 \rightarrow R_3 - R_2$$

$$\Rightarrow A \sim \begin{bmatrix} -4 & 1 & -1 \\ 0 & -5 & -3 \\ 0 & 0 & 0 \end{bmatrix}$$

Since number of non-zero rows of 'A' in Row-echelon form is 2,

$$\text{Rank of } A = \rho(A) = 2$$



Scan for Video solution



64. (b)

$$\begin{array}{c} \left[ \begin{array}{cccc} 1 & 1 & 0 & -2 \\ 2 & 0 & 2 & 2 \\ 4 & 1 & 3 & 1 \end{array} \right] \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 4R_1}} \left[ \begin{array}{cccc} 1 & 1 & 0 & -2 \\ 0 & -2 & 2 & 6 \\ 0 & -3 & 3 & 9 \end{array} \right] \\ \xrightarrow{R_3 \rightarrow R_3 - \frac{3}{2}R_2} \left[ \begin{array}{cccc} 1 & 1 & 0 & -2 \\ 0 & -2 & 2 & 6 \\ 0 & 0 & 0 & 0 \end{array} \right] \end{array}$$

Number of non-zero rows = 2

Hence Rank = 2



Scan for Video solution



65. (c)

Given:

$$\text{Matrix } M = \begin{bmatrix} 5 & 10 & 10 \\ 1 & 0 & 2 \\ 3 & 6 & 6 \end{bmatrix}$$

Converting M to row echelon form,

$$R_2 \rightarrow 5R_2 - R_1$$

$$R_3 \rightarrow 5R_3 - 3R_1$$

$$\Rightarrow M \sim \begin{bmatrix} 5 & 10 & 10 \\ 0 & -10 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

The above matrix is in row-echelon form

Number of non-zero rows = 2,  $\therefore \rho(M) = 2$



Scan for Video solution



66. (2)

$$\text{Given } P = \begin{bmatrix} 1 & 1 & -1 \\ 2 & -3 & 4 \\ 3 & -2 & 3 \end{bmatrix}; Q = \begin{bmatrix} -1 & -2 & -1 \\ 6 & 12 & 6 \\ 5 & 10 & 5 \end{bmatrix}$$

$$\Rightarrow P + Q = \begin{bmatrix} 0 & -1 & -2 \\ 8 & 9 & 10 \\ 8 & 8 & 8 \end{bmatrix}$$

$$\therefore |P + Q| = 0 \begin{vmatrix} 9 & 10 \\ 8 & 8 \end{vmatrix} + 1 \begin{vmatrix} 8 & 10 \\ 8 & 8 \end{vmatrix} - 2 \begin{vmatrix} 8 & 9 \\ 8 & 8 \end{vmatrix}$$


$$= 0 + 1(-16) - 2(-8) = 0$$

$$\therefore |P + Q| = 0$$

There exist a  $2 \times 2$  non-zero minor of  $P + Q$

$$\begin{vmatrix} 8 & 9 \\ 8 & 8 \end{vmatrix} \neq 0$$

$\therefore$  Rank of  $P + Q = 2$



Scan for Video solution



67. (4)

Let the given matrix is

$$A = \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

Converting to row-echelon form

$$R_4 \leftrightarrow R_2$$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

$$R_5 \rightarrow R_1 + R_2 + R_3 + R_4 + R_5$$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R_2 \rightarrow R_2 + R_1; R_4 \rightarrow R_4 + R_3$$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R_3 \rightarrow R_3 + R_2$$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The above matrix is in Row- echelon form.

$$\therefore \text{Number of non-zero rows of 'A' in echelon form} = 4$$

$$\Rightarrow \rho(A) = 4$$

Scan for Video solution



68. (17)

Let the given matrix be

$$A = \begin{bmatrix} 4 & 1 & 2 \\ p & 2 & 1 \\ 14 & -4 & 10 \end{bmatrix} \text{ and eigen vector is } x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

For an eigen values  $\lambda$ ,  $AX = \lambda X$

$$\therefore \begin{bmatrix} 4 & 1 & 2 \\ p & 2 & 1 \\ 14 & -4 & 10 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \lambda \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 12 \\ p+7 \\ 36 \end{bmatrix} = \begin{bmatrix} \lambda \\ 2\lambda \\ 3\lambda \end{bmatrix}$$

$$\Rightarrow \lambda = 12 \text{ and } p+7 = 2\lambda \Rightarrow p+7 = 24 \Rightarrow p = 17$$

Scan for Video solution



Determinant of sub-matrix  $\begin{bmatrix} 6 & 0 \\ -2 & 14 \end{bmatrix} \neq 0$

$\therefore$  Rank is 2.



Scan for Video solution



72. (a)

Given  $\rho(Q) = 4$

- Rank = Row Rank = No. of linearly independent rows
  - Rank = Column rank = No. of Linearly independent columns.
- $\therefore$  option (a) is correct

$$Q_{5 \times 6} Q_{6 \times 5}^T = QQ^T_{5 \times 5}$$

$$Q_{6 \times 5}^T Q_{5 \times 6} = Q^T Q_{6 \times 6}$$

We know,  $\rho(Q) = \rho(QQ^T) = \rho(Q^T Q) = 4$

$$\therefore |QQ^T| = |Q^T Q| = 0$$

$\therefore QQ^T$  and  $Q^T Q$  are not invertible



Scan for Video solution



73. (b, c)

$$A = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix}$$

For eigen values;  $|A - \lambda I| = 0$

$$\begin{vmatrix} 1-\lambda & -1 & 0 \\ -1 & 2-\lambda & -1 \\ 0 & -1 & 1-\lambda \end{vmatrix} = 0$$

$$(1-\lambda)\{(2-\lambda)(1-\lambda)-1\} + 1\{-(1-\lambda)\} = 0$$

$$(1-\lambda)\{(2-\lambda)(1-\lambda)-1-1\} = 0$$

$$(1-\lambda)\lambda(\lambda-3) = 0$$

$$\lambda = 0, 1, 3$$

Since there are three eigen values so number of linearly independent eigen vector are 3.

Since; product or eigen values  $= |A| = 0$

Since  $|A| = 0$  hence its  $\text{Rank}(A) < 3$

$\therefore \text{Rank } (A) < \text{number of unknowns}$

$\Rightarrow [A]\{x\} = [B]$  does not have a unique solution. For the positive definite matrix, all the eigen values must be positive. But here one eigen value is  $\lambda = 0$  so A is not a possible definite matrix.



Scan for Video solution



74. (c)

Given, system is

$$\begin{bmatrix} 1 & -\sqrt{2} & 3 \\ -1 & \sqrt{2} & -3 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

Let,  $x_1, x_2, x_3$  be the unknowns.

$$\Rightarrow A \times = b \Rightarrow \begin{bmatrix} 1 & -\sqrt{2} & 3 \\ -1 & \sqrt{2} & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

$$\Rightarrow x_1 - \sqrt{2} x_2 + 3x_3 = 1 \quad \dots(1)$$

$$-x_1 + \sqrt{2} x_2 - 3x_3 = 3 \quad \dots(2)$$

From equation (1) and (2),

$$1 = -3 \text{ (False)}$$

$\because$  LHS of both (1) and (2) differ by '-' sign

$\therefore$  Given system has no solution.



Scan for Video solution



75. (d)

Given system of equation is

$$x_1 + x_2 - 2x_3 = 4; x_1 + 3x_2 - x_3 = 7; 2x_1 + x_2 - 5x_3 = 7$$

The given system can be written as

$$\left[ \begin{array}{ccc|c} 1 & 1 & -2 \\ 1 & 3 & -1 \\ 2 & 1 & -5 \end{array} \right] \left[ \begin{array}{c} x_1 \\ x_2 \\ x_3 \end{array} \right] = \left[ \begin{array}{c} 4 \\ 7 \\ 7 \end{array} \right] \Rightarrow A \times B \text{ and } A = LU$$

Considering (LU = A)

$$LU = \left[ \begin{array}{ccc|c} 1 & 0 & 0 \\ L_{21} & 1 & 0 \\ L_{31} & L_{32} & 1 \end{array} \right] \left[ \begin{array}{ccc} U_{11} & U_{12} & U_{13} \\ 0 & U_{22} & U_{23} \\ 0 & 0 & U_{33} \end{array} \right]$$

$$\begin{aligned} LU &= \left[ \begin{array}{ccc|c} U_{11} & U_{12} & U_{13} \\ L_{21}U_{11} & L_{21}U_{12} + U_{22} & L_{21}U_{13} + U_{23} \\ U_{11}L_{31} & L_{31}U_{12} + L_{32}U_{22} & L_{31}U_{13} + L_{32}U_{23} + U_{33} \end{array} \right] \\ &= \left[ \begin{array}{ccc} 1 & 1 & -2 \\ 1 & 3 & -1 \\ 2 & 1 & -5 \end{array} \right] \end{aligned}$$

By equating the corresponding elements

$$U_{11} = 1; U_{12} = 1, U_{13} = -2$$

$$L_{21}U_{11} = 1 \Rightarrow L_{21} = 1$$

$$L_{21}U_{12} + U_{22} = 3 \Rightarrow (1 \times 1) + U_{22} = 3 \Rightarrow U_{22} = 2$$

$$L_{21}U_{13} + U_{23} = -1 \Rightarrow (1 \times -2 + U_{23}) = -1 \Rightarrow U_{23} = 1$$

$$U_{11}L_{31} = 2 \Rightarrow L_{31} = 2$$

$$L_{31}U_{12} + L_{32}U_{22} = 1 \Rightarrow (2 \times 1) + L_{32} \times 2 = 1 \Rightarrow L_{32} = -1/2$$

$$L_{31}U_{13} + L_{32}U_{23} + U_{33} = -5$$

$$2 \times -2 + \left( -\frac{1}{2} \times 1 \right) + U_{33} = -5$$

$$\Rightarrow U_{33} = -1 + \frac{1}{2} = -\frac{1}{2}$$

$$\Rightarrow U_{33} = -\frac{1}{2}$$



Scan for Video solution



$$\begin{vmatrix} 2 & 5-2\alpha \\ \alpha & 1 \end{vmatrix} = 0$$

$$\Rightarrow 2 - \alpha(5 - 2\alpha) = 0 \Rightarrow 2 - 5\alpha + 2\alpha^2 = 0$$

$$\Rightarrow 2\alpha^2 - 5\alpha + 2 = 0 \Rightarrow \alpha = 2, \frac{1}{2}$$

$$\text{For } \alpha = 2: \text{ system is} \quad 2x + 2y = 0 \\ x + y = 0$$

$\therefore (2, -2)$  is a solution.

$$\text{For } \alpha = \frac{1}{2}, 2x + \frac{y}{2} = 0 \Rightarrow 4x + y = 0$$

$\therefore (-1, 4)$  is a solution.

Scan for Video solution

### 77. (b, c)

**Given**

System is  $Ax = 0$

$A$  is of order  $3 \times 5$  and  $\rho(A) = 2$

For the system

$Ax = 0$  if  $\rho(A) < n$  (Number of unknowns),

Along with the trivial solution (zero-vector), the system has infinitely many solutions.

Scan for Video solution

### 78. (3 to 3)

$$(A / B) = \left[ \begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 2 & 2 & -3 & 1 \\ 4 & 4 & -6 & 2 \\ 2 & 5 & 2 & 1 \end{array} \right] \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 4R_1 \\ R_4 \rightarrow R_4 - 2R_1}} \left[ \begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 0 & -4 & -7 & -1 \\ 0 & -8 & -14 & -2 \\ 0 & -1 & -2 & -1 \end{array} \right]$$

$$\xrightarrow{\substack{R_4 \rightarrow R_4 - 2R_3}} \left[ \begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 0 & -1 & -2 & -1 \\ 0 & -4 & -7 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

### 76. (a, b)

**Given**

$$[xy] \begin{bmatrix} 2 & 5-2\alpha \\ \alpha & 1 \end{bmatrix} = [0 \ 0]$$

For the system to have non-trivial solution,

$$\xrightarrow{R_3 \rightarrow R_3 - 4R_2} \left[ \begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 0 & -1 & -2 & -1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

After converting to echelon form, equations have reduced to

$$\Rightarrow \left[ \begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 0 & 1 & -2 & -1 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right] \left[ \begin{array}{c} x_1 \\ x_2 \\ x_3 \\ x_4 \end{array} \right] = \left[ \begin{array}{c} 1 \\ -1 \\ 3 \\ 0 \end{array} \right]$$

From 3<sup>rd</sup>  $x_3 = 3$  equation of system.



Scan for Video solution



79. (a)

**Given:** System is

$$\begin{aligned} x_1 + 2x_2 &= b_1; & 2x_1 + 4x_2 &= b_2 \\ 3x_1 + 7x_2 &= b_3; & 3x_1 + 9x_2 &= b_4 \end{aligned}$$

The above system can be written in matrix form as

$$\left[ \begin{array}{cccc} 1 & 2 & 0 & 0 \\ 2 & 4 & 0 & 0 \\ 3 & 7 & 0 & 0 \\ 3 & 9 & 0 & 0 \end{array} \right] \left[ \begin{array}{c} x_1 \\ x_2 \\ x_3 \\ x_4 \end{array} \right] = \left[ \begin{array}{c} b_1 \\ b_2 \\ b_3 \\ b_4 \end{array} \right]$$

$$AX = B \text{ (say)}$$

$$\therefore x_1 + 2x_2 = b_1$$

$$\Rightarrow 2x_1 + 4x_2 = 2b_1 = b_2 \quad \therefore b_2 = 2b_1$$

For the system to have a solution,

$$\rho(A|B) = 2 = \text{number of unknowns}$$

All the  $4 \times 4$  and  $3 \times 3$  minors of  $[A|B]$  should be 0.

The  $3 \times 3$  minor of  $[A|B]$  involving  $b_1, b_3, b_4$  is

$$\begin{vmatrix} 1 & 2 & b_1 \\ 3 & 7 & b_3 \\ 3 & 9 & b_4 \end{vmatrix} = 0$$

$$\Rightarrow 1(7b_4 - 9b_3) - 2(3b_4 - 3b_3) + b_1(27 - 21) = 0$$

$$\Rightarrow b_4 - 3b_3 + 6b_1 = 0 \Rightarrow 6b_1 - 3b_3 + b_4 = 4$$

$\therefore$  For the system to have a solution,

$$b_2 = 2b_1 \text{ and } 6b_1 - 3b_3 + b_4 = 0$$



Scan for Video solution



80. (c)

**Given**

System is

$$\begin{aligned} x + y + z &= 1 \\ ax - ay + 3z &= 5 \\ 5x - 3y + az &= 6 \end{aligned}$$

$$\Rightarrow \left[ \begin{array}{ccc|c} 1 & 1 & 1 & x \\ a & -a & 3 & y \\ 5 & -3 & a & z \end{array} \right] = \left[ \begin{array}{c} 1 \\ 5 \\ 6 \end{array} \right] \quad (\text{Say } Ax = B)$$

$$\Rightarrow [A|B] = \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & 1 \\ a & -a & 3 & 5 & 5 \\ 5 & -3 & a & 6 & 6 \end{array} \right]$$

Converting to row echelon form

$$R_2 \rightarrow R_2 - aR_1 \text{ and } R_3 \rightarrow R_3 - 5R_1$$

$$\Rightarrow [A|B] \sim \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & 1 \\ 0 & -2a & 3-a & 5-a & 5 \\ 0 & -8 & a-5 & 1 & 1 \end{array} \right]$$

$$R_3 \rightarrow R_3 + 4R_2$$

$$\Rightarrow [A|B] = \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & 1 \\ 0 & -2a & 3-a & 5-a & 5 \\ 0 & 0 & a^2 - 5a - 12 + 4a & a - 20 + 4a & 1 \end{array} \right]$$

For the system to have infinite solutions,

$$a^2 - a - 12 = 0, 5a - 20 = 0$$

$$\Rightarrow \therefore a = 4 \text{ for infinitely many solutions}$$



Scan for Video solution



81. (2 to 2)

**Given:**

$$A = \begin{bmatrix} k & 2k \\ k^2 - k & k^2 \end{bmatrix}; \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

For the system  $Ax = 0$  to have infinitely many solutions,

$$|A| = 0$$

$$\Rightarrow k(k^2 - 2k) = 0 \Rightarrow -k^3 + 2k^2 = 0$$

$$\Rightarrow -k^2(k - 2) = 0 \Rightarrow k = 0, 0, 2$$

$$\therefore \text{No of distinct real value of } k = 2.$$



Scan for Video solution



82. (c)

Given  $a_i$  are column vectors in  $\mathbb{R}^n$ .

$$\sum_{i=1}^n c_i a_i = 0 \text{ for } c_1, c_2, \dots, c_n \text{ not all zero.}$$

$\Rightarrow a_1, a_2, \dots, a_n$  are linearly dependent.

$$\therefore |A| = |a_1 \ a_2 \ a_3 \ \dots \ a_n| = 0$$

$$\Rightarrow \rho(A) < n$$

$\therefore$  The system has infinitely many solutions.



Scan for Video solution



83. (a)

$$A = \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix}$$

The characteristic equation is  $|A - \lambda I| = 0$

$$\begin{bmatrix} 3-\lambda & 2 \\ 4 & 1-\lambda \end{bmatrix} = 0$$

$$(3-\lambda)(1-\lambda) - 8 = 0$$

$$\lambda^2 - 4\lambda + 3 - 8 = 0$$

$$\lambda^2 - 4\lambda - 5 = 0$$



Scan for Video solution



84. (d)

Given

$$\begin{bmatrix} 2 & 5 \\ -4 & 3 \end{bmatrix} \begin{Bmatrix} x \\ y \end{Bmatrix} = \begin{Bmatrix} 2 \\ -30 \end{Bmatrix}$$

$$\Rightarrow 2x + 5y = 2 \Rightarrow 4x + 10y = 4 \quad \dots (1)$$

$$\Rightarrow -4x + 3y = -30 \Rightarrow -4x + 3y = -30 \quad \dots (2)$$

$$(1) + (2) \Rightarrow 13y = -26$$

$$\Rightarrow y = -2 \Rightarrow x = 6$$

$$\therefore \begin{Bmatrix} x \\ y \end{Bmatrix} = \begin{Bmatrix} 6 \\ -2 \end{Bmatrix}$$



Scan for Video solution



85. (c)

Given system has 'm' linear equations in n variables

(i) For  $m < n$  (or)  $m > n$  (or)  $m = n$ , we can have systems with no solution, infinitely many solutions (or) single solution depending on Homogeneous or Non-Homogeneous system.

$\therefore$  First two statements (I) and (II) are generalized statements which need not be true.

(ii)  $\exists$  a system which has a solution for  $m = n$  which is correct.



Scan for Video solution



86. (b)

For system to be consistent;

Rank (A) Rank (A/B)

Augmented matrix for the system of equations

$$(A / B) = \left[ \begin{array}{ccc|c} 1 & 2 & -3 & a \\ 2 & 3 & 3 & b \\ 5 & 9 & -6 & c \end{array} \right]$$

$$\xrightarrow{\begin{array}{l} R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 5R_1 \end{array}} \left[ \begin{array}{ccc|c} 1 & 2 & -3 & a \\ 0 & -1 & 9 & b - 2a \\ 0 & -1 & 9 & c - 5a \end{array} \right]$$

$$\left[ \begin{array}{ccc|c} 1 & 2 & -3 & a \\ 0 & -1 & 9 & b - 2a \\ 0 & 0 & 0 & c - 5a - (b - 2a) \end{array} \right] \xleftarrow{R_3 \rightarrow R_3 - R_2}$$

Rank(A) = 2 Hence Rank(A/B) should be 2 for this to happen  $C - 5a - b + 2a = 0$

$$3a + b - c = 0$$



Scan for Video solution



87. (2)

$$\text{Given system is } \left[ \begin{array}{ccc|c} 1 & -2 & 3 & x \\ 1 & -3 & 4 & y \\ -2 & 4 & -6 & z \end{array} \right] \left[ \begin{array}{c} x \\ y \\ z \end{array} \right] = \left[ \begin{array}{c} -1 \\ 1 \\ k \end{array} \right]$$

$$AX = B$$

$$\text{Augmented matrix} = \left[ \begin{array}{ccc|c} 1 & -2 & 3 & -1 \\ 1 & -3 & 4 & 1 \\ -2 & 4 & -6 & k \end{array} \right]$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 + 2R_1$$

$$\Rightarrow [A/B] \sim \left[ \begin{array}{cccc} 1 & -2 & 3 & -1 \\ 0 & -1 & 1 & 2 \\ 0 & 0 & 0 & k-2 \end{array} \right]$$

If  $\rho(A) = \rho(A: B) < n$  for  
Infinitely many solutions

$$\therefore k-2=0$$

$$\Rightarrow k=2$$



Scan for Video solution



### 88. (4.49 to 4.51)

For no solution the given lines must be parallel and hence their slope should be equal.

$$-\frac{2}{3} = -\frac{3}{P}$$

$$P = \frac{9}{2} = 4.5$$



Scan for Video solution



### 89. (b)

Given system is

$$\begin{pmatrix} 2 & 1 & 3 \\ 3 & 0 & 1 \\ 1 & 2 & 5 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 5 \\ -4 \\ 14 \end{pmatrix}$$

Let  $Ax = B$

$\Rightarrow$  Augmented matrix;  $[A|B]$

$$\begin{bmatrix} 2 & 1 & 3 & 5 \\ 3 & 0 & 1 & -4 \\ 1 & 2 & 5 & 14 \end{bmatrix}$$

$$R_2 \rightarrow 2R_2 - 3R_1$$

$$R_3 \rightarrow 2R_3 - R_1$$

$$\Rightarrow [A | B] \sim \left[ \begin{array}{cccc} 2 & 1 & 3 & 5 \\ 0 & -3 & -7 & -23 \\ 0 & 3 & 7 & 23 \end{array} \right]$$

$$R_3 \rightarrow R_3 + R_2$$

$$\Rightarrow [A | B] \sim \left[ \begin{array}{cccc} 2 & 1 & 3 & 5 \\ 0 & -3 & -7 & -23 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\Rightarrow \rho(A) = \rho(A | B) = 2 < 3 \text{ (No. of unknowns)}$$

$\therefore$  The system has infinitely many solutions



Scan for Video solution



### 90. (b)

$$\text{Augmented matrix } (A: B) = \left[ \begin{array}{ccc|c} 1 & 2 & 2 & b_1 \\ 5 & 1 & 3 & b_2 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow R_2 - 5R_1} \left[ \begin{array}{ccc|c} 1 & 2 & 3 & b_1 \\ 0 & -9 & -7 & b_2 - 5b_1 \end{array} \right]$$

$$\rho(A) = \rho(A / B) = 2 < \text{No. of unknowns (3)}$$

$\therefore$  Infinite solution exists.



Scan for Video solution



### 91. (a)

Given system of equation is

$$\frac{dx}{dt} = 3x - 5y \text{ and } \frac{dy}{dt} = 4x + 8y$$

$$\Rightarrow \begin{bmatrix} \frac{dx}{dt} \\ \frac{dy}{dt} \end{bmatrix} = \begin{bmatrix} 3 & -5 \\ 4 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} \frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 3 & -5 \\ 4 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$



Scan for Video solution



## 92. (d)

The augmented matrix for the given system is

$$(A / B) = \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 1 & 2 & 3 & 4 \\ 1 & 4 & k & 6 \end{array} \right]$$

Convert it to row echelon form which will reduce it into upper triangular matrix to investigate its rank.

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 1 & 2 & 3 & 4 \\ 1 & 4 & k & 6 \end{array} \right] \xrightarrow{\substack{R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - R_1}} \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 2 & 1 \\ 0 & 3 & k-1 & 3 \end{array} \right]$$

$$R_3 \rightarrow R_3 - 3R_2$$

$$(A / B) = \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & k-7 & 0 \end{array} \right]$$

- For unique solution; Rank (A) = Rank (A/B) = no. of unknowns and for that  $k \neq 7$   
 $\Rightarrow$  Rank (A) = 3  
 $\Rightarrow$  Rank (A/B) = 3
- If  $k = 7$ ; Rank (A) = Rank (A/B) = 2 < no of unknowns (3). This is the condition for infinite solution and hence unique solution is not possible



Scan for Video solution



## 93. (1)

Given system is

$$3x + 2y = 1; 4x + 7z = 1$$

$$x + y + z = 3; x - 2y + 7z = 0$$

The matrix form of given system,

$$Ax = B \Rightarrow \left[ \begin{array}{ccc|c} 3 & 2 & 0 & 0 \\ 4 & 0 & 7 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & -2 & 7 & 0 \end{array} \right] \left[ \begin{array}{c} x \\ y \\ z \\ w \end{array} \right] = \left[ \begin{array}{c} 1 \\ 1 \\ 3 \\ 0 \end{array} \right]$$

$\Rightarrow$  Augmented matrix is

$$[A / B] = \left[ \begin{array}{cccc|c} 3 & 2 & 0 & 0 & 1 \\ 4 & 0 & 7 & 0 & 1 \\ 1 & 1 & 1 & 0 & 3 \\ 1 & -2 & 7 & 0 & 0 \end{array} \right]$$

$$R_2 \rightarrow 3R_2 - 4R_1$$

$$R_3 \rightarrow 3R_3 - R_1$$

$$R_4 \rightarrow 3R_4 - R_1$$

$$\Rightarrow [A / B] \sim \left[ \begin{array}{cccc|c} 3 & 2 & 0 & 0 & 1 \\ 0 & -8 & 21 & 0 & -1 \\ 0 & -1 & 3 & 0 & 8 \\ 0 & -8 & 21 & 0 & -1 \end{array} \right]$$

$$R_2 \leftrightarrow R_3$$

$$\Rightarrow [A / B] \sim \left[ \begin{array}{cccc|c} 3 & 2 & 0 & 0 & 1 \\ 0 & -1 & 3 & 0 & 8 \\ 0 & -8 & 21 & 0 & -1 \\ 0 & -8 & 21 & 0 & -1 \end{array} \right]$$

$$R_3 \leftrightarrow R_3 - 8R_2$$

$$\therefore [A / B] \sim \left[ \begin{array}{cccc|c} 3 & 2 & 0 & 0 & 1 \\ 0 & -1 & 3 & 0 & 8 \\ 0 & 0 & -3 & 0 & -65 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$\rho(A/B) = 3 = \rho(A) = \text{Number of unknowns}$$

$\therefore$  The system has unique solution.

$\therefore$  Number of solutions = 1



Scan for Video solution



## 94. (c)

**Given:** The system is

$$x + 2y + z = 4$$

$$2x + y + 2z = 5$$

$$x - y + z = 1$$

$$\Rightarrow \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 4 \\ 2 & 1 & 2 & 5 \\ 1 & -1 & 1 & 1 \end{array} \right] \left[ \begin{array}{c} x \\ y \\ z \\ x \\ y \\ z \end{array} \right] = \left[ \begin{array}{c} 4 \\ 5 \\ 1 \end{array} \right]$$

(say A) X B

For the system  $Ax = B$  to have a solution

$$\rho(A) = \rho(A/B)$$

$A \sim A/B$

$$\Rightarrow \left[ \begin{array}{ccc|c} 1 & 2 & 1 \\ 2 & 1 & 2 \\ 1 & -1 & 1 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 4 \\ 2 & 1 & 2 & 5 \\ 1 & -1 & 1 & 1 \end{array} \right]$$

Converting to Row-echelon form

$$R_2 \rightarrow R_2 - 2R_1 \text{ and } R_3 \rightarrow R_3 - R_1$$

$$\Rightarrow \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 4 \\ 0 & -3 & 0 & -3 \\ 0 & -3 & 0 & -3 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 4 \\ 0 & -3 & 0 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$R_3 \rightarrow R_3 - R_2$$

$$\Rightarrow \rho(A) = \rho(A/B) = 2 < 3$$

$\therefore$  Given system has infinitely many solutions.



Scan for Video solution



### 95. (b)

**Given:** System of equations

$$x + y + z = 6$$

$$x + 4y + 6z = 20$$

$$x + 4y + \lambda z = \mu$$

$$\Rightarrow AX = B \rightarrow \left[ \begin{array}{ccc|c} 1 & 1 & 1 & x \\ 1 & 4 & 6 & y \\ 1 & 4 & \lambda & z \end{array} \right] \left[ \begin{array}{c} 6 \\ 20 \\ \mu \end{array} \right]$$

Augmented matrix  $[A|B]$  is

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 1 & 4 & 6 & 20 \\ 1 & 4 & \lambda & \mu \end{array} \right]$$

Converting to row echelon form

$$R_2 \rightarrow R_2 - R_1 \text{ and } R_3 \rightarrow R_3 - R_1$$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 1 & 4 & 6 & 20 \\ 1 & 4 & \lambda & \mu \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 3 & 5 & 14 \\ 0 & 3 & \lambda - 1 & \mu - 6 \end{array} \right]$$

$$\text{Now, } R_3 \rightarrow R_3 - R_2$$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 3 & 5 & 14 \\ 0 & 3 & \lambda - 1 & \mu - 6 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 3 & 5 & 14 \\ 0 & 0 & \lambda - 6 & \mu - 20 \end{array} \right]$$

For the system to have no solution,  $\rho[A] \neq \rho[A|B]$

$$\Rightarrow \lambda - 6 = 0 \text{ and } \mu - 20 \neq 0$$

$$\therefore \boxed{\lambda = 6 \text{ and } \mu \neq 20}$$



Scan for Video solution



### 96. (a)

**Given:** system is

$$2x_1 + x_2 + x_3 = 0$$

$$x_2 - x_3 = 0$$

$$x_1 + x_2 = 0$$

$$\Rightarrow \left[ \begin{array}{ccc|c} 2 & 1 & 1 & x_1 \\ 0 & 1 & -1 & x_2 \\ 1 & 1 & 0 & x_3 \end{array} \right] = \left[ \begin{array}{c} 0 \\ 0 \\ 0 \end{array} \right] \rightarrow \text{Homogeneous system}$$

For this system to have non-trivial (or) infinity many solutions,

$$\left| \begin{array}{ccc} 2 & 1 & 1 \\ 0 & 1 & -1 \\ 1 & 1 & 0 \end{array} \right| = 0$$

$$\Rightarrow 2(1) - 1(1) + 1(-1) = 0 \text{ (True)}$$

$$\therefore \boxed{\text{Given system has infinitely many solutions.}}$$



Scan for Video solution



### 97. (d)

Augmented matrix

$$(A : B) = \left[ \begin{array}{cccc|c} 1 & 2 & 1 & 4 & 2 \\ 3 & 6 & 3 & 12 & 6 \end{array} \right]$$

$$\underline{R_2 \rightarrow R_2 - 3R_1} \left[ \begin{array}{cccc|c} 1 & 2 & 1 & 4 & 2 \\ 1 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$\therefore \rho(A) = \rho(A|B) = 1 < \text{Number of unknowns (4)}$$

$\therefore$  Infinite solutions exist.



Scan for Video solution



**98. (c)**

Given System is

$$x_1 + x_2 + 2x_3 = 1$$

$$x_1 + 2x_2 + 3x_3 = 2$$

$$x_1 + 4x_2 + \alpha x_3 = 4$$

If the system has unique solution,  $|A| \neq 0$ 

$$\begin{vmatrix} 1 & 1 & 2 \\ 1 & 2 & 3 \\ 1 & 4 & \alpha \end{vmatrix} \neq 0$$

$$\Rightarrow (2\alpha - 12) - (\alpha - 3) + 2(4 - 2) \neq 0$$

$$\Rightarrow 2\alpha - 12 - \alpha + 3 + 4 \neq 0$$

$$\Rightarrow \alpha - 5 \neq 0$$

$$\Rightarrow \alpha \neq 5$$

∴ Best option is (c).



Scan for Video solution

**99. (b)**

Given: equations are

$$4x + 2y = 7, \quad 2x + y = 6$$

$$\Rightarrow 2(2x + y) = 7 \quad \Rightarrow 2x + y = 7/2$$

The given equations can be written as

$$2x + y = 7/2 \text{ and } 2x + y = 6$$

∴  $7/2 = 6$  (Not possible), the system has no solution.

Scan for Video solution

**100. (b)**

Given: The system of equations

$$2x + 3y = 4$$

$$x + y + z = 4$$

$$x + 2y - z = a$$

$$\Rightarrow \begin{bmatrix} 2 & 3 & 0 \\ 1 & 1 & 1 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ a \end{bmatrix}$$

(say A)      (X)      (B)

For the system  $AX = B$ , to have a solution,

$$\text{Rank } \rho(A) = \text{Rank } \rho(A / B)$$

$$\Rightarrow \begin{bmatrix} 2 & 3 & 0 \\ 1 & 1 & 1 \\ 0 & 2 & -1 \end{bmatrix} \sim \begin{bmatrix} 2 & 3 & 0 & 4 \\ 1 & 1 & 1 & 4 \\ 1 & 2 & -1 & a \end{bmatrix}$$

$$R_2 \rightarrow 2R_2 - R_1 \text{ and } R_3 \rightarrow 2R_3 - R_1$$

$$\Rightarrow \begin{bmatrix} 2 & 3 & 0 \\ 0 & -1 & 2 \\ 0 & 1 & -2 \end{bmatrix} \sim \begin{bmatrix} 2 & 3 & 0 & 4 \\ 0 & -1 & 2 & 4 \\ 0 & 1 & -2 & 2a - 4 \end{bmatrix}$$

$$R_3 \rightarrow R_3 + R_2$$

$$\Rightarrow \begin{bmatrix} 2 & 3 & 0 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \sim \begin{bmatrix} 2 & 3 & 0 & 4 \\ 0 & -1 & 2 & 4 \\ 0 & 0 & 0 & 2a \end{bmatrix}$$

$$\Rightarrow \rho(A) = 2$$

For  $\rho(A / B) = 2$ , All elements in 3<sup>rd</sup> row of matrix (A/B) should be zero.

$$\Rightarrow 2a = 0$$

$$\Rightarrow a = 0$$



Scan for Video solution

**101. (b, c)**

We know,

$$Mx = \lambda x$$

∴  $\lambda$  is eigen value of M.

$$\text{Multiply by } M; \quad M^2x = \lambda Mx$$

$$M^2x = \lambda(\lambda x)$$

$$M^2x = \lambda^2x$$

∴  $\lambda^2$  is eigen value of  $M^2$ Since M and  $M^T$  have same eigen values,

$$\therefore \text{We can write } M^T x = \lambda x$$

$$\text{Multiply by } M^T; \quad (M^T)^2 x = \lambda M^T x$$

$$(M^T)^2 x = \lambda (\lambda x)$$

$$(MT)^2 x = \lambda^2 x$$

∴  $\lambda^2$  is eigen value of  $(M^T)^2$ 

$$MM^T x = \lambda Mx$$

$$MM^T x = \lambda(\lambda x)$$

$$MM^T x = \lambda^2 x$$

∴  $\lambda^2$  is eigen value of  $MM^T$ .

$$M^T M x = \lambda M^T x$$

- $M^T M x = \lambda(\lambda x)$   
 $M^T M x = \lambda^2 x$   
 $\therefore \lambda^2$  is eigen value of  $M^T M$   
 Since  $MM^T$  and  $M^T M$  is always symmetric matrix.  
 Eigen values of symmetric matrices are always real.  
 Hence  $\lambda^2$  will non-negative iff  $\lambda$  is real  
 Hence option b and c are correct.



Scan for Video solution

**102. (c)**

Given that  $\{\lambda_k | 1 \leq k \leq n\}$  and  $\{v_k | 1 \leq k \leq n\}$  are eigen values and eigen vectors of 'B' respectively.

$$B = P^{-1} AP$$

$\Rightarrow A$  is the matrix obtained from diagonalization of 'B'.

$\therefore B$  and  $A$  have some eigen values.

Let 'v' be the eigen vector of 'B' for an eigen values  $\lambda$ .

$$\Rightarrow BV = \lambda v$$

Multiplying with 'P'

$$\Rightarrow (AP)V = \lambda(PV)$$

$$\Rightarrow A(PV) = \lambda(PV)$$

i.e. eigen vector of 'A' is  $PV$

$\therefore$  Eigen vectors of 'A' are  $\{PV_k | 1 \leq k \leq n\}$

$\therefore$  Eigen values and eigen vectors of 'A' are

$$\{\lambda | 1 \leq k \leq n\} \text{ and } \{PV_k | 1 \leq k \leq n\}$$



Scan for Video solution

**103. (a, b, d)**

Using standard properties of eigen values and eigen vectors and vectors:

- Eigen values of  $A$  and  $A^T$  are same
- Eigen values of  $A^{-1}$  is reciprocal of eigen values of  $A$ .
- Eigen vectors of  $A^{-1}$  are same as that eigen vectors of  $A$ .

Row space and column space for any matrix are not same.



Scan for Video solution

**104. (a, d)**

Let  $y = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}_{3 \times 1}$  Hence  $y_T = [1 \ 1 \ 1]_{1 \times 3}$

Then  $yy^T = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}_{3 \times 3}$

Then  $Y^T y = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = [3]_{1 \times 1}$

- $\text{Rank}(y) = \text{Rank}(y^T) = \text{Rank}(yy^T) = \text{Rank}(y^T y) = 1$
- $\therefore$  Since  $yy^T = (yy^T)$  hence  $yy^T$  is symmetric.
- $\therefore$  Since determinate of  $yy^T$  is zero hence its inverse does not exists.
- Eigen values for  $yy^T$  are 0,0,3  
 Hence  $y^T y$  is an eigen value for  $yy^T$  i.e.3
- Rank of  $y^T y$  is 1



Scan for Video solution

**105. (a, c)**

Since  $M$  has  $\lambda_1 = 5$  and  $\lambda_2 = -2$

Hence matrix  $Q = M^3 - 4M^2 - 2M$  will have its eigen values as

$\rightarrow$  1<sup>st</sup> eigen value =

$$\lambda_1^3 - 4\lambda_1^2 - 2\lambda_1 = 5^3 - 4(5)^2 - 2(5) = 15$$

$\rightarrow$  2<sup>nd</sup> eigen value =

$$\lambda_2^3 - 4\lambda_2^2 - 2\lambda_2 = (-2)^3 - 4(-2)^2 - 2(-2) = -20$$

using eigen value properties.



Scan for Video solution

**106. (2 to 2)**

Let  $\lambda_1, \lambda_2$  are Eigen values of  $\begin{bmatrix} 3 & p \\ p & q \end{bmatrix} = A$  (say)

Given  $\lambda_1 + \lambda_2 = 4$  and  $\lambda_1 \cdot \lambda_2 = -1$

$$\Rightarrow \lambda_1 + \lambda_2 = \text{tr}(A) = 3 + q = 4 \Rightarrow q = 1$$

$$\begin{aligned}\lambda_1, \lambda_2 &= |A| = -1 \\ \Rightarrow 3q - p^2 &= -1 \Rightarrow p^2 = 3q + 1 \\ \Rightarrow p^2 &= 3(1) + 1 \Rightarrow p^2 = 4 \\ \Rightarrow |p| &= 2\end{aligned}$$



Scan for Video solution

**107. (a)**

$$|A - \lambda I| = 0 \Rightarrow \begin{vmatrix} 1-\lambda & 0 & 0 \\ 0 & 4-\lambda & -2 \\ 0 & 1 & 1-\lambda \end{vmatrix} = 0$$

$$\Rightarrow (1-\lambda)[(4-\lambda)(1-\lambda) + 2] = 0$$

$$(1-\lambda)(\lambda^2 - 5\lambda + 6) = 0$$

$$-\lambda^3 + 5\lambda^2 - 6\lambda + \lambda^2 - 5\lambda + 6 = 0$$

$$\lambda^3 - 6\lambda^2 + 11\lambda - 6 = 0 \text{ (Characteristic equation)}$$

By Cayley Hamilton theorem,

$$A^3 - 6A^2 + 11A - 6I = 0$$

$$A^2 - 6A + 11I - 6A^{-1} = 0$$

$$6A^{-1} = A^2 - 6A + 11I \text{ [Multiply by } A^{-1}]$$

$$6A^{-1} = A^2 + cA + dI$$

$\therefore$  On comparing;  $c = -6$

$$d = 11$$

$$\therefore c + d = -6 + 11 = 5$$



Scan for Video solution

**108. (c)**

$$\text{We know; } e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$e^x V = V + V_x + \frac{Vx^2}{2!} + \frac{Vx^3}{3!} + \dots \text{ [Multiply by } V]$$

$$e^A V = V + vA + V \frac{A^2}{2!} + \frac{vA^3}{3!} + \dots \text{ [ } x \rightarrow A \text{ ]}$$

$$e^A V = V \left[ 1 + \lambda + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \dots \right] \quad [ A \rightarrow \lambda ]$$

$$e^A V = e^\lambda V$$

$\therefore e^\lambda$  is an eigen value of  $e^A$ .

$V$  is an eigen vector of  $e^A$ .



Scan for Video solution

**109. (b)**

$$a_{ij} = (i - j)^3$$

$$A = \begin{bmatrix} 0 & -1 & -8 \\ 1 & 0 & -1 \\ 8 & 1 & 0 \end{bmatrix} \quad a_{ij} = -a_{ji}$$

$$A^T = \begin{bmatrix} 0 & 1 & 8 \\ -1 & 0 & 1 \\ -8 & -1 & 0 \end{bmatrix} = -A$$

$\therefore A$  is skew-symmetric.




Scan for Video solution

**110. (b)****Given**

'P' is a two-dimensional vector and also Eigen vector of 'A' corresponding to  $\lambda$ .

$$\Rightarrow AP = \lambda P = P'$$



So, since Eigen value is just a scalar,  $P'$  and  $P$  will be in the same direction.

$$\Rightarrow \text{Direction of } P' = \theta \text{ and } \|P'\| = \lambda \cdot \|P\|$$



Scan for Video solution



**111. (d)****Given**

For a non-zero column vector 'P'

$$AP = \lambda P \text{ where } \lambda \text{ is an Eigen value.}$$

$$\text{Also, } AP = \alpha^2 P$$

 $\Rightarrow \lambda = \alpha^2$  is Eigen value of 'A'.

$$\therefore \text{Eigen value of } A^2 = \lambda^2 = (\alpha^2)^2$$

$$\therefore \text{Eigen value of } A^2 = \alpha^4$$



Scan for Video solution

**112. (1)**

$$A \rightarrow 10 \times 10 \text{ matrix}$$

$$A^5 = 0$$

$$\therefore A^{10} = A^5 \cdot A^5 = 0 \cdot 0 = 0$$

 $\therefore \text{Eigen values of } A^{10} \text{ are all } 0.$ 
Let eigen values of A are  $\lambda_1, \lambda_2, \dots, \lambda_{10}$ .Then eigen values of  $A^{10}$  are  $\lambda_1^{10}, \lambda_2^{10}, \lambda_3^{10}, \dots, \lambda_{10}^{10}$ 

$$\therefore \lambda_1^{10} = \lambda_2^{10} = \dots = 0$$

$$\therefore \lambda_1, \lambda_2, \dots, \lambda_{10} = 0$$

$$\begin{aligned} \therefore \text{Eigen values of } A + I &= \lambda_1 + 1, \lambda_2 + 1, \dots, \lambda_{10} + 1 \\ &= 1, 1, \dots, 10 \text{ times} \end{aligned}$$

 $\therefore \text{Determinant of } A + I = |A + I| = \text{Product of eigen values}$ 

$$= 1 \times 1 \times \dots \times 10 \text{ times}$$

$$|A+I| = 1$$



Scan for Video solution

**113. (b)**

$$\text{Let } A = \begin{bmatrix} p & q \\ q & -p \end{bmatrix}$$

$$|A - \lambda I| = 0 \Rightarrow \begin{bmatrix} p & q \\ q & -p \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} p - \lambda & q \\ q & -p - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (p - \lambda)(-p - \lambda) - q^2 = 0$$

$$-p^2 - \lambda p + \lambda p + \lambda^2 - q^2 = 0$$

$$\lambda^2 - (p^2 + q^2) = 0 \quad [\because p^2 + q^2 = 1]$$

$$\lambda^2 - 1 = 0$$

$$\lambda = \pm 1$$



Scan for Video solution

**114. (3 to 3)**

Given matrix is

$$\begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

If sum of all the elements of a row (or) column is constant (say k) for all the rows (or) columns, then 'k' is the largest eigen value of the matrix.

For given matrix, sum of all the elements of a row (or) column = 3 (constant for any row or column)

 $\therefore \text{Largest eigen value of given matrix} = 3$ 


Scan for Video solution

**115. (b)**

$$\text{Let } A = \begin{bmatrix} 2 & -2 \\ -1 & 6 \end{bmatrix}$$

The characteristic equation is  $|A - \lambda I| = 0$ 

$$= \begin{bmatrix} 2 - \lambda & -2 \\ -1 & 6 - \lambda \end{bmatrix} = 0$$

$$= (2 - \lambda)(6 - \lambda) - 2 = 0$$

$$= \lambda^2 - 8\lambda + 10 = 0$$

$$\Rightarrow \lambda = 4 \pm \sqrt{6}$$

Hence smallest eigen value is  $4 - \sqrt{6}$

For  $\lambda = 4 - \sqrt{6}$  find eigen vector;  $(A - \lambda I) X = 0$

$$\begin{bmatrix} 2-\lambda & -2 \\ -1 & 6-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2-(4-\sqrt{6}) & -2 \\ -1 & 6-(4-\sqrt{6}) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0$$

$$(2+\sqrt{6})x_1 - 2x_2 = 0$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -2+\sqrt{6} \end{bmatrix} = \begin{bmatrix} 2 \\ 0.45 \end{bmatrix}$$

Hence smallest eigen value is 1.55 its corresponding

eigen vector is  $\begin{bmatrix} 2 \\ 0.45 \end{bmatrix}$



Scan for Video solution



### 116. (d)

Sum of eigen values = Trace of matrix

$$\text{Trace}(P) = 0 + 3 + 6 + 6 = 15$$

Product of eigen values = Determinant of matrix

$$= -(-2) \begin{vmatrix} 1 & 3 & 0 \\ 0 & 6 & 1 \\ 0 & 1 & 6 \end{vmatrix} = 2 \times 35 = 70$$

Only **option (c)** is correct because sum of eigen values = 15 & product of eigen values = 70



Scan for Video solution



### 117. (1 to 1)

**Given**

'I' is an identity matrix of dimensions  $100 \times 100$

→ All the eigen values of an identity matrix are '1'.

∴ The number of distinct eigen values of 'A' is 1.

$$\therefore E = \{1\}$$

⇒ Number of elements in E = 1



Scan for Video solution



### 118. (3 to 3)

**Given:**

$$\text{Matrix } A = \begin{bmatrix} 2 & 2 & 3 & 3 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

'A' is an upper triangular matrix

∴ Eigen values of 'A' are principal diagonal elements of 'A' i.e., 2, 1, 3, and 2.

∴ Number of distinct eigen values = 3



Scan for Video solution



### 119. (b)

**Given**

$$P = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \text{ (Upper triangular matrix)}$$

Eigen values of upper triangular matrix are diagonal elements.

∴ Eigen values of P are 1, 1, 1

∴ Number of distinct Eigen values of P = 1



Scan for Video solution



### 120. (d)

**Given**

$$[A] = \begin{bmatrix} 4 & 8 & 4 \\ 8 & 16 & -4 \\ 4 & -4 & 15 \end{bmatrix}, \{X\} = \begin{Bmatrix} 2 \\ 1 \\ 4 \end{Bmatrix} \text{ and } \{R\} = \begin{Bmatrix} 32 \\ 16 \\ 64 \end{Bmatrix}$$

For 'X' to be an Eigen vector of 'A',  $AX = \lambda X$

$$\Rightarrow \begin{bmatrix} 4 & 8 & 4 \\ 8 & 16 & -4 \\ 4 & -4 & 15 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix} = \lambda \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix} = \begin{bmatrix} 32 \\ 16 \\ 64 \end{bmatrix}$$

$$\Rightarrow \lambda \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix} = 16 \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix} \Rightarrow \lambda = 16$$

$\therefore$  One of the Eigen values = 16



Scan for Video solution



121. (d)

$$M \rightarrow \lambda_1, \lambda_2$$

$$\Rightarrow M^2 \rightarrow \lambda_1^2, \lambda_2^2$$

Eigen values of M are 4 and 9.

$\therefore$  Eigen values of  $M^2$  are  $4^2$  and  $9^2$

$$\Rightarrow 16 \text{ and } 81.$$

[By property of eigen values]



Scan for Video solution



122. (12 to 12)

$$\text{Given } R = \begin{bmatrix} 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \\ 1 & 5 & 25 & 125 \end{bmatrix}$$

Absolute value of product of eigen values = modulus of  $|R|$

$$\therefore \det(R) = R = \begin{bmatrix} 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \\ 1 & 5 & 25 & 125 \end{bmatrix}$$

$$R_2 \Rightarrow R_2 - R_1$$

$$R_3 \Rightarrow R_3 - R_1$$

$$R_4 \Rightarrow R_4 - R_1$$

$$\Rightarrow |R| = \begin{vmatrix} 1 & 2 & 4 & 8 \\ 0 & 1 & 5 & 19 \\ 0 & 2 & 12 & 56 \\ 0 & 3 & 21 & 117 \end{vmatrix}$$

$$\text{Since } |R| = |R^T| = \begin{vmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 2 & 3 \\ 4 & 5 & 12 & 21 \\ 8 & 19 & 56 & 117 \end{vmatrix}$$

$$= \begin{vmatrix} 1 & 2 & 3 \\ 5 & 12 & 21 \\ 19 & 56 & 117 \end{vmatrix}$$

$$= 1 ((12 \times 117) - (21 \times 56)) - 2 ((5 \times 117) - (19 \times 21)) + 3 ((5 \times 56) - (19 \times 12)) = 12$$

$\therefore$  Absolute value of product of eigen values  
 $= \text{mod}(|R|) = 12$



Scan for Video solution



123. (1.0)

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 1 - \lambda & 0 & -1 \\ -1 & 2 - \lambda & 0 \\ 0 & 0 & -2 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (1 - \lambda)(-2 - \lambda)(2 - \lambda) = 0$$

$$\lambda = 1, 2, -2$$

$\therefore$  eigen value of  $B = A^3 - A^2 - 4A + 5I$

$$\text{For } \lambda = 1, \lambda_1 = 1^3 - 1^2 - 4 \cdot 1 + 5 \cdot 1 = 1$$

$$\text{For } \lambda = 2, \lambda_2 = 2^3 - 2^2 - 4 \cdot 2 + 5 \cdot 1 = 1$$

$$\text{For } \lambda = -2, \lambda_3 = (-2)^3 - (-2)^2 - 4(-2) + 5 \cdot 1 = 1$$

$$\therefore |B| = \lambda_1 \cdot \lambda_2 \cdot \lambda_3$$

$$= 1 \times 1 \times 1 = 1$$



Scan for Video solution



124. (d)

$$\text{Let } A = \begin{bmatrix} 2 & -4 \\ 4 & -2 \end{bmatrix}$$

The characteristic equation is  $|A - \lambda I| = 0$

$$\begin{vmatrix} 2-\lambda & -4 \\ 4 & -2-\lambda \end{vmatrix} = 0$$

$$(2-\lambda)(-2-\lambda) + 16 = 0$$

$$\lambda^2 - 2\lambda + \lambda - 4 + 16 = 0$$

$$\lambda^2 = 12 = 0$$

$\lambda = \pm 2\sqrt{3}$ ; (Hence complex eigen values) for 1<sup>st</sup> eigen vector, put  $\lambda = 2\sqrt{3}i$   $(A - \lambda I)x = 0$

$$\begin{bmatrix} 2-\lambda & -4 \\ 4 & -2-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2-2\sqrt{3}i & -4 \\ 4 & -2-2\sqrt{3}i \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$(2-\sqrt{3}i)x_1 - 4x_2 = 0$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4 \\ 2-2\sqrt{3}i \end{bmatrix}$$

For 2<sup>nd</sup> eigen vector, put  $\lambda = -2\sqrt{3}i$

$$\begin{bmatrix} 2+2\sqrt{3}i & -4 \\ 4 & 02+2\sqrt{3}i \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$(2+2\sqrt{3}i)x_1 - 4x_2 = 0$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4 \\ 2+2\sqrt{3}i \end{bmatrix}$$

∴ The eigen values and eigen vector are complex.

**NOTE:** If eigen value is real the eigen vector is also real, If eigen value is complex then eigen vector is also complex.

Scan for Video solution



### 125. (d)

The condition of diagonalization of matrix is that all its eigen vectors should be linearly Independent. (The eigen values may not be distinct).

Linearly dependent eigen vectors are due to the repeated eigen Values.

So, statements (II) and statement (III) are necessarily true.

Scan for Video solution



### 126. (5.5 to 5.5)

Let eigen values of  $A \rightarrow \lambda_1, \lambda_2$

Then eigen values of  $A^2 \rightarrow \lambda_1^2, \lambda_2^2$

Given  $T_r(A) = \lambda_1 + \lambda_2 = 4$  ----(1)

$T_r(A^2) = \lambda_1^2 + \lambda_2^2 = 5$  ----(2)

We know,  $(\lambda_1 + \lambda_2)^2 = \lambda_1^2 + \lambda_2^2 + 2\lambda_1\lambda_2$

$$4^2 = 5 + 2\lambda_1\lambda_2$$

$$\therefore \lambda_1\lambda_2 = 11/2$$

∴ Determinant of  $A = \lambda_1\lambda_2 = 5.5$

Scan for Video solution



### 127. (c)

Eigen vectors of distinct eigen values are always linearly independent

∴ If eigen values of a matrix are distinct then it implies that eigen vectors are linearly independent

∴ S2 implies S1

**Note:** Invertible (Non-singularity) depends on non-zero Eigen values.

Scan for Video solution



### 128. (b)

Given 'A' is a square matrix.

$$\Rightarrow A = A^T$$

$$\sum_{i=1}^n \sum_{j=1}^n A_{ij}^2 = 50$$

For any square matrix;

Sum of squares of all elements = trace of  $(A \cdot A^T)$

$$\Rightarrow \sum_{i=1}^n \sum_{j=1}^n A_{ij}^2 = \text{tr}(A \cdot A^T) = \text{tr}(A \cdot A^T) = \text{tr}(A^2)$$

Given  $\rho(A) = 2$

$$\Rightarrow n - \{\text{Algebraic multiplicity of '0' as Eigen value}\} = 2$$

$\Rightarrow$  Algebraic multiplying of '0' as Eigen value =  $n - 2$

$\therefore$  Let  $\lambda_1, \lambda_2, 0, 0, \dots$  are Eigen values of 'A'.

$\Rightarrow \lambda_1^2, \lambda_2^2, 0, 0, \dots, 0$  are

$$\Rightarrow 50 = \lambda_1^2 + \lambda_2^2 \quad \dots(1)$$

$$\text{Let } |\lambda_1| = 5 \Rightarrow |\lambda_2| = 5$$

$$|\lambda_1| = 4 \Rightarrow |\lambda_2| \leq 6$$

So, for any combination, one Eigen value lies in  $[-5, 5]$ .

(ii) The Eigen value with largest magnitude need not be greater than 5. Since  $\lambda_1 = 5$  and  $\lambda_2 = 5$  satisfy equation (i).

$\therefore$  Only statement I is true.



Scan for Video solution



### 129. (5)

Given characteristic polynomial of  $M_{3 \times 3}$  is

$$C(\lambda) = \lambda^3 - 4\lambda^2 + a\lambda + 30$$

Let  $\lambda_1, \lambda_2, \lambda_3$ , are eigen values of M

$$\Rightarrow \lambda_1 + \lambda_2 + \lambda_3 = 4 \text{ and } \lambda_1, \lambda_2, \lambda_3 = -30$$

Let  $\lambda_1 = 2$

$$\Rightarrow \lambda_2 + \lambda_3 = 2 \text{ and } \lambda_2, \lambda_3 = -15$$

On solving the above two equations,  $\lambda_2 = 5, \lambda_3 = -3$

$\therefore$  Eigen values of M are 2, 5, -3

$\therefore$  Absolute values of eigen values of M are 2, 5, 3

$\therefore$  Largest among the absolute Eigen values of M = 5



Scan for Video solution



### 130. (a)

$$\text{Let } A = \begin{bmatrix} 5 & -1 \\ 4 & 1 \end{bmatrix}$$

Hence characteristic equation is  $|A - \lambda I| = 0$

$$|A - \lambda I| = \begin{vmatrix} 5 - \lambda & -1 \\ 4 & 1 - \lambda \end{vmatrix} = 0$$

$$(5 - \lambda)(1 - \lambda) + 4 = 0$$

$$5 - 6\lambda + \lambda^2 + 4 = 0$$

$$\lambda^2 - 6\lambda + 9 = 0$$

$$(\lambda - 3)^2 = 0$$

$$\lambda = 3, 3$$

Hence eigen value 3 has algebraic multiplicity of 2 for eigen vector  $(A - \lambda I)x = 0$

$$\begin{bmatrix} 5 - \lambda & -1 \\ 4 & 1 - \lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\text{For } \lambda = 3; \begin{bmatrix} 5 - 3 & -1 \\ 4 & 1 - 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$


$$2x_1 - x_2 = 0$$

Let  $x_1 = k_1$  then  $x_2 = 2k_1$

$$\text{Hence eigen vector } x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = k \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$\text{Eigen vector is } \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Hence only one independent eigen vector.



Scan for Video solution



### 131. (c)

Given:

$$\text{Matrix } A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 1 & 2 & 3 & 4 \\ 4 & 5 & 1 & 2 & 3 \\ 3 & 4 & 5 & 1 & 2 \\ 2 & 3 & 4 & 5 & 1 \end{bmatrix}$$

Scan for Video solution

To calculate the eigen values of A, we need  $|A - \lambda I| = 0$

$$\Rightarrow \begin{vmatrix} 1-\lambda & 2 & 3 & 4 & 5 \\ 5 & 1-\lambda & 2 & 3 & 4 \\ 4 & 5 & 1-\lambda & 2 & 3 \\ 3 & 4 & 5 & 1-\lambda & 2 \\ 2 & 3 & 4 & 5 & 1-\lambda \end{vmatrix} = 0$$

Applying  $C_1 \rightarrow C_1 + C_2 + C_3 + C_4 + C_5$

$$\Rightarrow \begin{vmatrix} 15-\lambda & 2 & 3 & 4 & 5 \\ 15-\lambda & 1-\lambda & 2 & 3 & 4 \\ 15-\lambda & 5 & 1-\lambda & 2 & 3 \\ 15-\lambda & 4 & 5 & 1-\lambda & 2 \\ 15-\lambda & 3 & 4 & 5 & 1-\lambda \end{vmatrix} = 0$$

$\Rightarrow (15-\lambda)$  is a factor of above equation

$\Rightarrow \lambda = 15$  is an eigen value of 'A'

$\therefore$  The one and only real values of 'A' is 15.

**Note:** If sum of all the elements of a row (or) column is constant (k) for all the rows (or) columns of a matrix, then 'k' is an eigen values of the matrix.



Scan for Video solution



### 132. (b)

**Given**

$$\text{Matrix } P = \begin{bmatrix} 2 & 0 & 1 \\ 4 & -3 & 3 \\ 0 & 2 & -1 \end{bmatrix}$$

Product of Eigen values of a matrix is equal to its determinant.

$$\Rightarrow \lambda_1 \cdot \lambda_2 \cdot \lambda_3 = |P| = 2(3 - 6) - 0(-4) + 1(8) \\ = -6 + 8 = 2$$

$$\therefore \lambda_1 \cdot \lambda_2 \cdot \lambda_3 = 2$$



Scan for Video solution



$$\begin{bmatrix} 3/2 & 0 & 1/2 \\ 0 & -1 & 0 \\ 1/2 & 0 & 3/2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = \lambda X = 1 \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$\therefore$  Option c) is correct.



Scan for Video solution



### 136. (a)

Given

$$\text{Matrix } A = \begin{bmatrix} 2 & 1 \\ 1 & k \end{bmatrix}$$

Let  $\lambda_1, \lambda_2$  be the eigen values of 'A'.

Since the product of eigenvalues of a matrix is equal to its determinant.

$$\therefore \lambda_1 \cdot \lambda_2 = |A| = 2k - 1$$

If Eigen values are positive then  $\lambda_1, \lambda_2 > 0$

$$\Rightarrow 2k - 1 > 0 \Rightarrow k > \frac{1}{2}$$



Scan for Video solution



### 137. (2 to 2)

Given

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

$\Rightarrow$  'A' is an upper Triangular matrix.

- Number of linearly independent Eigen vectors of a matrix is equal to number of distinct Eigen values of the Matrix.

Since 'A' is an upper triangular matrix, the eigen values of 'A' are diagonal elements.

$\therefore$  The Eigen values of 'A' are 2, 2, 3.

$$\therefore \left\{ \begin{array}{l} \text{Number of distinct} \\ \text{Eigen values} \end{array} \right\} = \left\{ \begin{array}{l} \text{Number of linearly} \\ \text{independent Eigen vectors} = 2 \end{array} \right\}$$


Scan for Video solution



### 138. (d)

Given:  $A = \begin{bmatrix} \sigma & x \\ \omega & \sigma \end{bmatrix}$  and eigen values of A are

$$\sigma + j\omega \text{ and } \sigma - j\omega,$$

$$\text{Product of eigen values of } A = |A|$$

$$\Rightarrow (\sigma + j\omega)(\sigma - j\omega) = \sigma^2 - x\omega$$

$$\Rightarrow \sigma^2 + \omega^2 = \sigma^2 - x\omega$$

$$\therefore x = -\omega$$


Scan for Video solution



### 139. (1)

$$\text{Given } A = \begin{bmatrix} 3 & 2 & 4 \\ 9 & 7 & 13 \\ -6 & -4 & -9+x \end{bmatrix}$$

For zero to be an eigen values,  $|A| = 0$ :

$$\Rightarrow \begin{vmatrix} 3 & 2 & 4 \\ 9 & 7 & 13 \\ -6 & -4 & -9+x \end{vmatrix} = 0$$

$$R_3 \rightarrow R_3 + 2R_1$$

$$\Rightarrow \begin{vmatrix} 3 & 2 & 4 \\ 9 & 7 & 13 \\ 0 & 0 & -1+x \end{vmatrix} = 0$$

This determinant is zero if  $-1 + x = 0$

$$\Rightarrow x = 1$$


Scan for Video solution



### 140. (d)

Consider a  $2 \times 2$  matrix  $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

Hence characteristic equation will be  $|M - \lambda I| = 0$

$$\begin{aligned}
 &= \begin{vmatrix} a - \lambda & b \\ c & d - \lambda \end{vmatrix} = 0 \\
 &= (a - \lambda)(d - \lambda) - bc = 0 \\
 &= \lambda^2 - (a + d)\lambda + (ad - bc) = 0 \\
 &= \lambda^2 - (a + d)\lambda + ad - (1 - d)(1 - a) = 0
 \end{aligned}$$

Now put  $\lambda = 1$ ,

$$1 - a - d + ad - 1 + a + d - ad = 0$$

Hence  $\lambda = 1$  satisfies the characteristic equation  
similarly put  $\lambda = 2, 3, 4$  but this value will not satisfy  
characteristic equation for any value of  $a$  &  $d$



Scan for Video solution



#### 141. (15 to 15)

Let  $\lambda_1, \lambda_2, \lambda_3$ , are eigen values of 'P'

$$\text{And } \lambda_1 = 2 + \sqrt{-1} \text{ and } \lambda_2 = 3$$

Complex roots always occur as conjugate pairs.

$$\Rightarrow \lambda_3 = 2 - \sqrt{-1}$$

$$\therefore |P| = \lambda_1, \lambda_2, \lambda_3,$$

$$= (2 + \sqrt{-1})(2 - \sqrt{-1})3 = 5(3) = 15$$

$\therefore$  Determinant of 'P' is 15.



Scan for Video solution



#### 142. (a)

By property, if eigen values of A are  $\lambda_1, \lambda_2$  then then  
the eigen values of  $A^m$  is  $\lambda_1^m, \lambda_2^m$  but eigen vector of  
A &  $A^m$  are same respectively.

Eigen values of A are 1, -2

$\therefore$  Eigen values of  $A^2 - 3A + 4I$

$$\Rightarrow \lambda_1 = 1^2 - 3 \times 1 + 4 \times 1 = 2$$

$$\Rightarrow \lambda_2 = (-2)^2 - 3(-2) + 4 \times 1 = 14$$

&  $\therefore$  using property eigen vector of  $A^2 - 3A + 4I$  are  
 $x_1$  and  $x_2$ .



Scan for Video solution



#### 143. (d)

$$p^3 = p$$

According to cayley hamilton theorem every square  
matrix satisfies its own characteristic equation.

$\therefore$  Characteristic equation matrix P is  $\lambda^3 = \lambda$

If eigen values of P are  $\lambda_1, \lambda_2$  and  $\lambda_3$ ,

$$\lambda^3 = \lambda$$

$$\lambda^3 - \lambda = 0$$

$$\lambda(\lambda^2 - 1) = 0$$

$$\lambda(\lambda+1)(\lambda-1) = 0$$

$$\lambda = 0, 1, -1$$



Scan for Video solution



#### 144. (3)

$$\text{Let } A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$|A - \lambda I| = 0 \Rightarrow \begin{bmatrix} 1-\lambda & 1 & 1 \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{bmatrix} = 0$$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$\begin{bmatrix} 3-\lambda & 3-\lambda & 3-\lambda \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{bmatrix} = 0$$

$$(3-\lambda) \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{bmatrix} = 0$$

$$(3-\lambda)[(1-\lambda)^2 - 1] - 1(1-\lambda-1) + 1(1-(1-\lambda)) = 0$$

$$(3-\lambda)[1+\lambda^2 - 2\lambda - 1 + \lambda + \lambda] = 0$$

$$\lambda^2(3-\lambda) = 0$$

$$\lambda = 0, 0, 3$$

Hence only non-zero eigen value is 3.



Scan for Video solution



**145. (b)**

$$A = \begin{bmatrix} -3 & 0 & -2 \\ 1 & -1 & 0 \\ 0 & a & -2 \end{bmatrix}$$

$$|A - \lambda I| = \begin{bmatrix} -3 - \lambda & 0 & -2 \\ 1 & -1 - \lambda & 0 \\ 0 & a & -2 - \lambda \end{bmatrix}$$

$$= (-3-\lambda)[(-1-(-\lambda)(-2-\lambda)] - 2a$$

$$= -(\lambda+3)[2 + \lambda + 2\lambda + \lambda^2] - 2a$$

$$= -(\lambda+3)[\lambda^2 + 3\lambda + 2] - 2a$$

$$= -(\lambda^3 + 6\lambda^2 + 11\lambda + 6) - 2a = 0$$

$$a = -\frac{1}{2}(\lambda^3 + 6\lambda^2 + 11\lambda + 6)$$

$a = f(\lambda)$  Hence, to find maximum value of  $a$  we can use concept of maximum & maxima.

$$\frac{da}{d\lambda} = 0 \Rightarrow \frac{da}{d\lambda} \left( -\frac{1}{2}(\lambda^3 + 6\lambda^2 + 11\lambda + 6) \right) = 0$$

$$3\lambda^2 + 12\lambda + 11 = 0$$

$$\lambda = \frac{-12 \pm \sqrt{12^2 - 4 \times 3 \times 11}}{2 \times 3}$$

$$\lambda = -2 \pm \frac{1}{\sqrt{3}}$$

$$\lambda = -22.577, -1.422$$

We can write,

$$\frac{d^2a}{d\lambda^2} = \frac{d}{d\lambda} \left( -\frac{1}{2}(3\lambda^2 + 12\lambda + 11) \right)$$

$$\frac{d^2a}{d\lambda^2} = -\frac{1}{2}(6\lambda + 12)$$

For  $\lambda = -2.577$

$$\frac{d^2a}{d\lambda^2} = -\frac{1}{2}(6 \times -2.577 + 12) = 1.731 > 0$$

$\therefore$  At  $\lambda = -2.577$ ,  $a$  is minimum.

For  $\lambda = 1.422$

$$\frac{d^2a}{d\lambda^2} = -\frac{1}{2}(6 \times -1.422 + 12) = -1.734 < 0$$

$\therefore$  At  $\lambda = 1.422$ ,  $a$  is maximum

$$\therefore a = -\frac{1}{2}(\lambda^3 + 6\lambda^2 + 11\lambda + 6)$$

$$= -\frac{1}{2}[(-1.422)^2 + 6(-1.422)^2 + 11(-1.422) + 6]$$

$$= 0.1924$$

$$= \frac{1}{3\sqrt{3}}$$

Scan for Video solution



**146. (a)**

$$\text{Let } a_1 x + b_1 y + c_1 z = d_1$$

$$a_2 x + b_2 y + c_2 z = d_2$$

$$a_3 x + b_3 y + c_3 z = d_3$$

$$AX = B$$

For unique solution  $\delta(A) = \delta(A:B) = \text{No of unknowns}$   
(3)

$\therefore$  if solutions are unique the all 3 equations are independent and if equation are independent

$$\therefore |A| \neq 0$$

Let eigen values of coefficient matrix A are  $\lambda_1, \lambda_2, \lambda_3$

$\therefore$  Determinant = Product of eigen values of  $\lambda_1, \lambda_2, \lambda_3$ .

$\therefore$  If  $|A| \neq 0$  then none of the eigen values are zero.

$\therefore P \equiv Q \equiv R \equiv S$



Scan for Video solution



#### 147. (9)

Let A be  $2 \times 2$  matrix whose eigen values are  $\lambda_1$  and  $\lambda_2$ .

Sum of eigen values of  $\lambda_1 + \lambda_2 = -6$

We know, determinant = Product of eigen values

$$D = \lambda_1 \lambda_2$$

$$D = \lambda_1 (-6 - \lambda_1)$$

$$D = -6\lambda_1 - \lambda_1^2$$

$$\frac{dD}{d\lambda_1} = -6 - 2\lambda_1 = 0$$

$$\lambda_1 = -3$$

$$\frac{d^2D}{d\lambda_1^2} = -6 < 0$$

$\therefore$  At  $\lambda_1 = -3$ , D is maximum.

$\therefore$  Maximum value of determinant

$$\begin{aligned} D &= -6\lambda_1 - \lambda_1^2 \\ &= -6(-3) - (-3)^2 \\ &= 18 - 9 = 9 \end{aligned}$$

$$D_{\max} = 9$$



Scan for Video solution



**150. (d)**

$$A = \begin{bmatrix} 2 & 1 \\ 1 & p \end{bmatrix}$$

Let  $\lambda_1$  and  $\lambda_2$  be the eigen values of matrix A.

Sum of eigen values =  $\lambda_1 + \lambda_2 = 2 + p$  .....(i)

Products of eigen value  $\lambda_1\lambda_2 = 2p = 1$ .....(ii)

Given  $\lambda_1/\lambda_2 = 3 = \lambda_1 \Rightarrow 3\lambda_2$

From equation (i)  $\Rightarrow 3\lambda_2 + \lambda_2 = 2 + P$

$$\lambda_2 = \frac{2 + P}{4}$$

From equation (ii)

$$3\lambda_2^2 = 2p - 1$$

$$3\left(\frac{2+p}{4}\right)^2 = 2p - 1$$

$$3(4 + p^2 + 4p) = 32p - 16$$

$$12 + 3p^2 + 12p = 32p - 16$$

$$3p^2 - 20p + 28 = 0$$

$$3p^2 - 6p - 14p + 28 = 0$$

$$3p(p-2) - 14(p-2) = 0$$

$$(p-2)(3p-14) = 0$$

$$p = 2, \frac{14}{3}$$

Hence another value of  $p = \frac{14}{3}$

 Scan for Video solution

**151. (6 to 6)**

Given matrix is  $\begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix} = A$  (say)

Let  $\lambda_1, \lambda_2$ , are eigen value of 'A'

$$\Rightarrow \lambda_1 + \lambda_2 = \text{trace}(A) = 5 \quad \dots(1)$$

$$\lambda_1, \lambda_2 = |A| = -6$$

$$\text{since } (\lambda_1 - \lambda_2)^2 = (\lambda_1 + \lambda_2)^2 - 4\lambda_1\lambda_2$$

$$\Rightarrow (\lambda_1 - \lambda_2)^2 = 5^2 - 4 \times -6$$

$$\Rightarrow (\lambda_1 - \lambda_2)^2 = 25 + 24 = 49$$

$$\Rightarrow \lambda_1 - \lambda_2 = 7 \quad \dots(2)$$

adding equation (1) and (2),

$$\lambda_1 = 6$$

$$\lambda_2 = -1$$

Largest eigen value = 6

 Scan for Video solution

**152. (b)**

Let the given matrix be

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 1 & 2 & 1 \end{bmatrix}$$

And  $\lambda_1 = 1$  For  $\lambda_1 = 1$

$$AX = \lambda_1 X$$

$$\Rightarrow AX = X$$

$$\begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 1 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\Rightarrow x_1 - x_2 + 2x_3 = x_1$$

$$\Rightarrow x_2 - 2x_3 = 0 \Rightarrow x_3 = \frac{x_2}{2}$$

$$\Rightarrow x_1 + 2x_2 + x_3 = x_3$$

$$\Rightarrow x_1 + 2x_2 = 0 \Rightarrow x_2 = -\frac{x_1}{2}, x_3 = -\frac{x_1}{4}$$

$$\therefore \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ -x_1/2 \\ -x_1/4 \end{bmatrix} = \frac{-x_1}{4} \begin{bmatrix} -4 \\ 2 \\ 1 \end{bmatrix}$$

$$\therefore \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \alpha [-4, 2, 1]$$

 Scan for Video solution

**153. (b)**

Given matrix is

$$\text{Let } P = \begin{bmatrix} 10 & 5+j & 4 \\ x & 20 & 2 \\ 4 & 2 & -10 \end{bmatrix}$$

All the eigen values of Hermitian matrix are always real.

$\therefore$  For 'P' to be Hermitian matrix,  $x = 5-j$



Scan for Video solution

**154. (b)**

For a singular matrix, Determinant (i.e., product of all Eigen Values) is zero.

$\Rightarrow$  At least one eigen value should be zero.



Scan for Video solution

**155. (2 to 2)**

Given

$$\text{Matrix } \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix} = A \text{ (say)}$$

Let  $\lambda_1, \lambda_2$  are the eigenvalues of this matrix.

$$\Rightarrow \lambda_1 + \lambda_2 = \text{tr}(A) = 7$$

$$\text{and } \lambda_1 \cdot \lambda_2 = |A| = 10$$

$\Rightarrow$  Eigen values are 2, 5.

$\therefore$  Lowest Eigen value = 2



Scan for Video solution

**156. (d)**

$$\text{Let } A = \begin{bmatrix} -5 & 2 \\ -9 & 6 \end{bmatrix}$$

The Characteristic equation is  $|A - \lambda I| = 0$

$$\Rightarrow \begin{vmatrix} -5 - \lambda & 2 \\ -9 & 6 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow -(5 + \lambda)(6 - \lambda) + 18 = 0$$

$$\Rightarrow (\lambda + 5)(\lambda - 6) + 18 = 0$$

$$\Rightarrow \lambda^2 - \lambda - 12 = 0$$

$$\Rightarrow \lambda = 4, -3$$

$$\text{For } \lambda = 4, \quad Ax = \lambda x$$

$$\Rightarrow \begin{bmatrix} -5 & 2 \\ -9 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -5x_1 + 2x_2 \\ -9x_1 + 6x_2 \end{bmatrix} = \begin{bmatrix} 4x_1 \\ 4x_2 \end{bmatrix} \Rightarrow 9x_1 = 2x_2$$

$$\therefore \text{Eigen Vector} = \begin{bmatrix} x_1 \\ \frac{2}{9}x_1 \end{bmatrix} = \begin{bmatrix} 9 \\ 2 \end{bmatrix}$$

$$\text{For } \lambda = -3, \quad Ax = \lambda x$$

$$\Rightarrow \begin{bmatrix} -5 & 2 \\ -9 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -3x_1 \\ -3x_2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -5x_1 + 2x_2 \\ -9x_1 + 6x_2 \end{bmatrix} = \begin{bmatrix} -3x_1 \\ -3x_2 \end{bmatrix} \Rightarrow 2x_2 = 2x_1 \Rightarrow x_1 = x_2$$

$$\therefore \text{Eigen vector} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\therefore \text{One of the Eigen vectors} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Scan for Video solution

**157. (d)**

Eigen vectors of a symmetric matrix are orthogonal to each other.

$$\therefore x_1 y_1 + x_2 y_2 + x_3 y_3 = [x_1 x_2 x_3] [y_1 y_2 y_3]^T = 0$$

Scan for Video solution

**158. (a)**

Eigen value of real symmetric matrix, it may be distinct positive or negative.



Scan for Video solution

**159. (0.33)**

Let us find characteristic equation;

$$|A - \lambda I| = 0 \Rightarrow \begin{vmatrix} 0 - \lambda & 1 & -1 \\ -6 & -11 - \lambda & 6 \\ -6 & -11 & 5 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow \lambda^3 + 6\lambda^2 + 11\lambda + 6 = 0$$

$$(\lambda + 1)(\lambda + 2)(\lambda + 3) = 0$$

$$\lambda = -1, -2, -3$$

$$\lambda_{\min} = -3, \lambda_{\max} = -1$$

$$\left| \frac{\lambda_{\max}}{\lambda_{\min}} \right| = \left| \frac{-1}{-3} \right| = 0.33$$



Scan for Video solution

**160. (0 to 0)**

The Eigen vectors corresponding to different Eigen values of a symmetric matrix are orthogonal to each others. (i.e.,  $X_i^T X_j = 0 \forall i \neq j$ )

$\therefore$  Dot product of eigen vectors = 0



Scan for Video solution

**161. (0)**

$$\text{Given } A = \begin{bmatrix} 2 \\ -4 \\ 7 \end{bmatrix}_{3 \times 1} [1 \ 9 \ 5]_{1 \times 3} = \begin{bmatrix} 2 & 18 & 10 \\ -4 & -36 & -20 \\ 7 & 63 & 35 \end{bmatrix}_{3 \times 3}$$

Since  $R_1$  and  $R_2$  are proportional to each other,  $|A| = 0$

$$\therefore |A| = 0$$



Scan for Video solution

**162. (a)**

Sum of eigen values of a given matrix = Trace of that given matrix

= sum of principal diagonal elements

$$= 215 + 150 + 550$$

$$= 915$$



Scan for Video solution

**163. (b)**

For a Real symmetric matrix 'A'.

Eigen values are always Real but need not be positive.

All other given options are correct.



Scan for Video solution

**164. (c)**

The Eigen values of a symmetric matrix are always real numbers.



Scan for Video solution

**165. (d)**

We know  $AX - \lambda X$

$$\text{Let } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\rightarrow \text{for } \lambda = -1 \rightarrow X = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = -1 \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$a - b = -1 \dots \text{(i)}$$

$$c - d = 1 \dots \text{(ii)}$$

$$\rightarrow \text{for } \lambda = -2 \rightarrow X = \begin{bmatrix} 1 \\ -2 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = -2 \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

$$a - 2b = -2 \dots \text{(iii)}$$

$$c - 2d = 4 \dots \text{(iv)}$$

On solving (i) and (iii);  $a = 0$  and  $b = 1$

On solving (ii) and (iv);  $c = -2$  and  $d = -3$

$$\therefore A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$$



Scan for Video solution



#### 166. (d)

$$2x_1 - 2x_2 = 0$$

$$x_1 - x_2 = 0$$

$$A = \begin{bmatrix} 2 & -2 \\ 1 & -1 \end{bmatrix}$$

$$|A| = 0 \therefore \rho(A) = 1$$

Since  $\rho(A) <$  Number of unknowns  $\{x_1, x_2\}$

$$1 < 2$$

$\therefore$  Infinite solution exists.

Let,  $x_1 = k$ ,

Then,  $x_2 = k$



Scan for Video solution



#### 167. (a)

Let the given matrix is

$$A = \begin{bmatrix} 3 & 5 & 2 \\ 5 & 12 & 7 \\ 2 & 7 & 5 \end{bmatrix}$$

$$C_1 \rightarrow C_1 + C_3$$

$$\Rightarrow A \sim \begin{bmatrix} 5 & 5 & 2 \\ 12 & 12 & 7 \\ 7 & 7 & 5 \end{bmatrix}$$

Since two columns are identical,

$\Rightarrow |A| = \text{Product of eigen values of } A = 0$

$\Rightarrow '0'$  is an eigen value of 'A'

$\therefore$  From given options, '0' is the least Eigen value



Scan for Video solution



#### 168. (b)

$$\text{Given: } A = \begin{bmatrix} 5 & 3 \\ 1 & 3 \end{bmatrix}.$$

$\Rightarrow$  Characteristic equation is  $|A - \lambda I| = 0$ .

$$\Rightarrow \begin{bmatrix} 5 - \lambda & 3 \\ 1 & 3 - \lambda \end{bmatrix} = 0$$

$$\Rightarrow (\lambda - 5)(\lambda - 3) - 3 = 0$$

$$\Rightarrow \lambda^2 - 8\lambda + 12 = 0$$

$$\Rightarrow \lambda = 2, 6$$

**(i) Eigen vector of '2'**

$$\Rightarrow \begin{bmatrix} 5 & 3 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix}$$

$$\Rightarrow 5x_1 + 3x_2 = 2x_1$$

$$\Rightarrow x_1 = -x_2$$

$$x_1 + 3x_2 = 2x_1 \Rightarrow x_1 = -x_2$$

$$\therefore \text{Eigen vector} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$\therefore \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an eigen vector of 'A'.

$$\Rightarrow \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} \end{bmatrix} \text{ is a normalized Eigen vector.}$$



Scan for Video solution



**169. (b)**

We know characteristic equation is  $|A - \lambda I| = 0$

$$\begin{vmatrix} -5 - \lambda & -3 \\ 2 & 0 - \lambda \end{vmatrix} = 0$$

$$\lambda(\lambda + 5) + 6 = 0$$

$$\lambda^2 + 5\lambda + 6 = 0$$

According to Cayley-Hamilton Theorem,

$$A^2 + 5A + 6I = 0$$

$$A^2 = -5A - 6I$$

$$A^2 \cdot A = (-5A - 6I)A$$

$$A^3 = -5A^2 - 6IA$$

$$= -5(-5A - 6I) - 6A$$

$$A^3 = 19A + 30I$$



Scan for Video solution

**170. (d)**

$$\text{Let } A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

Let Eigen values of 'A' be  $\lambda_1$  and  $\lambda_2$ .

$$\Rightarrow \lambda_1 + \lambda_2 = \text{tr}(A) = 1 - 1 = 0$$

$$\Rightarrow \lambda_1 + \lambda_2 = 0$$

$$\Rightarrow \lambda_1 = -\lambda_2$$

$$\lambda_1, \lambda_2 = |A| = -2$$

$$\therefore \lambda_1 = \sqrt{2} \text{ and } \lambda_2 = -\sqrt{2}$$

Eigen value of  $A^{19}$  are

$$\lambda_1^{19}, \lambda_2^{19} = (\sqrt{2})^{19}, (-\sqrt{2})^{19} = 512\sqrt{2}, -512\sqrt{2}$$



Scan for Video solution

**171. (a)**

$$\text{Let the given matrix is } A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 7 \\ 0 & 0 & 3 \end{bmatrix}$$

Since the given matrix is upper triangular matrix, Eigen values are Principal diagonal elements.

$\therefore$  Eigen values of given matrix are 1, 4, 3.



Scan for Video solution

**172. (c)**

- Eigen values of real symmetric matrix are always real. (Ex: Principal stresses of a stress tensor are always real.)
- Eigen values of skew symmetric matrix are either zero (or) purely imaginary.



Scan for Video solution

**173. (b)**

**Given:**

$$\text{Matrix } A = \begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$$

The characteristic equation of 'A' is  $|A - \lambda I| = 0$

$$\Rightarrow (2 - \lambda)(3 - \lambda) - 2 = 0 \Rightarrow \lambda^2 - 5\lambda + 6 - 2 = 0$$

$$\Rightarrow \lambda^2 - 5\lambda + 4 = 0 \Rightarrow \lambda = 1, 4$$

**Eigen vector for  $\lambda = 1$ :**  $Ax = \lambda x$

$$\Rightarrow \begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 1 \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\Rightarrow 2x_1 + 2x_2 = x_1 \Rightarrow x_2 = -2x_1$$

$$\text{and } x_1 + 3x_2 = x_1 \Rightarrow x_1 = -2x_2$$

$$\therefore \begin{bmatrix} 2 \\ -1 \end{bmatrix} \text{ is an Eigen vector}$$



Scan for Video solution



**174. (c)**

- Eigen values of a symmetric matrix are always real.
- Eigen values of a skew-symmetric matrix are either zero (or) Purely Imaginary complex Number.



Scan for Video solution

**175. (d)**

Given  $A = \begin{bmatrix} 2 & 3 \\ x & y \end{bmatrix}$  and 4, 8 are Eigen values of A.

Since sum of eigen values = trace (A)

$$\Rightarrow 2 + y = 4 + 8$$

$$\Rightarrow y = 10$$

Product of Eigen Values =  $|A|$

$$\Rightarrow \lambda_1 \cdot \lambda_2 = 2y - 3x$$

$$\Rightarrow 4 \times 8 = (2 \times 10) - 3x$$

$$\Rightarrow 3x = -12$$

$$\Rightarrow x = -4$$



Scan for Video solution

**176. (b)**

We know  $PX = \lambda X$

$$\text{From option(a)} \quad \left[ \begin{array}{ccc|c} 1 & 1 & 0 & -1 \\ 0 & 2 & 2 & 1 \\ 0 & 0 & 3 & 1 \end{array} \right] = PX$$

$$PX = \left[ \begin{array}{c} 0 \\ 4 \\ 3 \end{array} \right] \neq \lambda \left[ \begin{array}{c} -1 \\ 1 \\ 1 \end{array} \right] (\therefore PX \neq \lambda X)$$

$$\text{From option (b)} \quad \left[ \begin{array}{ccc|c} 1 & 1 & 0 & 1 \\ 0 & 2 & 2 & 2 \\ 0 & 0 & 3 & 1 \end{array} \right] = PX$$

$$PX = \left[ \begin{array}{c} 3 \\ 6 \\ 3 \end{array} \right] = \lambda \left[ \begin{array}{c} 1 \\ 2 \\ 1 \end{array} \right]$$

Since for  $\lambda=3$   
 $PX = \lambda X$  Hence option (b)  
is correct



Scan for Video solution

**177. (c)**

Let the eigen of  $2 \times 2$  matrix are  $\lambda_1$  and  $\lambda_2$ .

Trace of matrix  $= \lambda_1 + \lambda_2 = -2 \dots \text{(i)}$

Determinant of matrix  $= \lambda_1 \cdot \lambda_2 = -35 \dots \text{(ii)}$

$$\lambda_1 + \frac{-35}{\lambda_1} = -2$$

$$\lambda_1^2 + 2\lambda_1 - 35 = 0$$

$$(\lambda_1 + 7)(\lambda_1 - 5) = 0$$

$$\lambda_1 = -7, \quad 5$$



$$\lambda_2 = 5, \quad -7$$

$\therefore$  Eigen values are -7 and 5.

▶
Scan for Video solution

**178. (d)**

**Given:**

$$\text{Matrix } A = \begin{bmatrix} -1 & 3 & 5 \\ -3 & -1 & 6 \\ 0 & 0 & 3 \end{bmatrix}$$

Eigen values of 'A' are obtained by the equation  $|A - \lambda I| = 0$ . In this case the characteristic equation will be of 3<sup>rd</sup> degree.

All options involve complex roots of  $|A - \lambda I| = 0$

We know complex roots occurs as conjugate pairs:

i.e Given matrix should have 1 real eigen value and two conjugates complex eigen values.

$\therefore$  Option 'D' is correct

▶
Scan for Video solution

**179. (b)**

$$[P] = \begin{bmatrix} 4 & 5 \\ 2 & -5 \end{bmatrix}$$

Hence the characteristic equation for matrix P,

$$|P - \lambda I| = 0$$

$$\left| \begin{bmatrix} 4 & 5 \\ 2 & -5 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right| = 0$$

$$\begin{vmatrix} 4-\lambda & 5 \\ 2 & -5-\lambda \end{vmatrix} = 0$$

$$(4-\lambda)(-5-\lambda) - 2 \times 5 = 0$$

$$-20 - 4\lambda + 5\lambda + \lambda^2 - 10 = 10$$

$$\lambda^2 + \lambda - 30 = 0$$

$$\lambda^2 - 5\lambda + 6\lambda - 30 = 0$$

$$1(\lambda - 5) + 6(\lambda - 5) = 0$$

$$(\lambda - 5)(\lambda + 6) = 0$$

$$\lambda = 5, -6$$



Scan for Video solution



### 180. (a)

Given matrices are

(i)  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  → Diagonal matrix.

∴ Eigen values are 1, 0.

(ii)  $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$  → Upper triangular matrix.

∴ Eigen values are 0, 0.

(iii)  $\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$  → Let  $\lambda_1, \lambda_2$  are Eigen values

$$\Rightarrow \lambda_1 + \lambda_2 = 2; \lambda_1 \lambda_2 = 2$$

$$\Rightarrow \lambda_1 + \frac{2}{\lambda_1} = 2$$

$$\Rightarrow \lambda^2 - 2\lambda + 2 = 0$$

$$\Rightarrow (\lambda - 1)^2 + 1 = 0$$

$$\Rightarrow \lambda = 1 + i, 1 - i$$

(iv)  $\begin{bmatrix} -1 & 0 \\ 1 & -1 \end{bmatrix}$  → Lowest triangular matrix

⇒ Eigen values are -1, -1.

∴ Number of matrices with eigen values '1' is 1.



Scan for Video solution



### 181. (c)

**Given:**

Matrix,  $A = \begin{bmatrix} 1 & 2 & 4 \\ 3 & 0 & 6 \\ 1 & 1 & p \end{bmatrix}$

Let  $\lambda_1, \lambda_2, \lambda_3$  are eigenvalues of A

⇒  $\lambda_1 + \lambda_2 + \lambda_3 = \text{trace}(A) = \text{Sum of diagonal element}$

$$\text{Let } \lambda_1 = 3,$$

$$\Rightarrow 3 + \lambda_2 + \lambda_3 = 1 + 0 + p \Rightarrow \lambda_2 + \lambda_3 = p - 2$$

∴ Sum of other two eigen values =  $p - 2$



Scan for Video solution



### 182. (b)

**Given:**

Matrix  $A = \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}$

∴ 'A' is an upper triangular matrix. Therefore, Eigen values of 'A' are principal diagonal elements.

$$\therefore \text{Let } \lambda_1 = 1 \& \lambda_2 = 2$$

For  $\lambda_1 = 1$ , let the eigen vector be  $\begin{bmatrix} 1 \\ a \end{bmatrix}$

$$\Rightarrow A \begin{bmatrix} 1 \\ a \end{bmatrix} = \lambda_1 \begin{bmatrix} 1 \\ a \end{bmatrix} \quad \Rightarrow \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ a \end{bmatrix} = \begin{bmatrix} 1 \\ a \end{bmatrix}$$


$$\Rightarrow \begin{bmatrix} 1+2a \\ 2a \end{bmatrix} \begin{bmatrix} 1 \\ a \end{bmatrix} \quad \Rightarrow a = 0$$

Let  $\lambda_2 = 2$ , let the Eigen vector be  $\begin{bmatrix} 1 \\ b \end{bmatrix}$

$$\Rightarrow A \begin{bmatrix} 1 \\ b \end{bmatrix} = \lambda_2 \cdot \begin{bmatrix} 1 \\ b \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ b \end{bmatrix} = 2 \begin{bmatrix} 1 \\ b \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1+2b \\ 2b \end{bmatrix} = \begin{bmatrix} 2 \\ 2b \end{bmatrix} \Rightarrow 1+2b=2 \Rightarrow b=1/2$$

$$\therefore a+b=0+\frac{1}{2}=\frac{1}{2}$$




183. (c)

**Given:**

$$P = \begin{bmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \end{bmatrix} \text{ and 'P' has '0' as an eigen value.}$$

 $\Rightarrow$  Product of eigen value of  $P = |P| = 0$ 

$$\Rightarrow |P| = 0 \Rightarrow P_{11}P_{22} - P_{12}P_{21} = 0$$



184. (a)

$$\text{Rank}(A_{4 \times 4}) = 2$$

$$\text{Nullity}(A) = \text{Number of columns} - \text{Rank} = 4 - 2 = 2$$


i.e., Null space of A will consist only two linear independent vectors which are given.

Eigen vectors of A are  $c_1 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 3 \end{bmatrix}$  and  $c_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 2 \end{bmatrix}$

As these are linearly independent eigen vectors so remaining eigen vectors of null space must be linearly dependent



Hence  $C_1 - C_2 \begin{bmatrix} 1 \\ 1 \\ -1 \\ 1 \end{bmatrix} = [1 \ 1 \ -1 \ 1]^T$




185. (c)

For a square matrix 'A'

$$\begin{aligned} |A^T \cdot A^{-1}| &= |A^T| \cdot |A^{-1}| \\ &= |A| \cdot \left| \frac{1}{A} \right| = 1 \\ (\because |A^T| &= |A| \text{ & } |A^{-1}| = |A|^{-1}) \end{aligned}$$

$$\therefore |A^T \cdot A^{-1}| = 1$$



186. (2)


Given  $\dim(v) = 6$  and $\dim(V_1) = \dim(V_2) = 4$ Since  $V_1, V_2$  are subspaces of 'V'

$$\dim(V) \geq \dim(V_1) + \dim(V_2) - \dim(V_1 \cap V_2)$$

$$\Rightarrow 6 \geq 4 + 4 - \dim(V_1 \cap V_2)$$

$$\Rightarrow -2 \geq -\dim(V_1 \cap V_2)$$

$$\Rightarrow \dim(V_1 \cap V_2) \geq 2$$



## CHAPTER


# 2

# BASIC CALCULUS

### Differential Calculus

- 1. [MCQ]** [GATE-ME-2023: 1M]

The figure shows the plot of a function over the interval  $[-4, 4]$ , which one of the options given CORRECTLY identifies the function?



- (a)  $|2 - x|$       (b)  $|2 - |x||$   
 (c)  $|2 + |x||$       (d)  $2 - |x|$

- 2. [MCQ]** [GATE-ME-2022: 1M]

The limit  $P = \lim_{x \rightarrow \pi} \left( \frac{x^2 + \alpha x + 2\pi^2}{x - \pi + 2 \sin x} \right)$  has a finite value

for a real  $\alpha$ . The value of  $\alpha$  and the corresponding limit  $p$  are

- (a)  $\alpha = -3\pi$ , and  $p = \pi$       (b)  $\alpha = -2\pi$ , and  $p = 2\pi$   
 (c)  $\alpha = \pi$ , and  $p = \pi$       (d)  $\alpha = 2\pi$ , and  $p = 3\pi$

- 3. [NAT]** [GATE-CS-2022: 1M]

The value of the following limit is \_\_\_\_\_.

$$\lim_{x \rightarrow 0^+} \frac{\sqrt{x}}{1 - e^{2\sqrt{x}}}.$$

- 4. [NAT]** [GATE-CS-2021: 2M]

Consider the following Expression.  $\lim_{x \rightarrow -3} \frac{\sqrt{2x + 22} - 4}{x + 3}$

The value of the above expression (rounded to 2 decimal place) is \_\_\_\_\_

- 5. [MCQ]**

[GATE-ME-2021: 1M]

The value of  $\lim_{x \rightarrow 0} \left( \frac{1 - \cos x}{x^2} \right)$  is

- (a)  $1/4$       (b)  $1/3$   
 (c)  $1/2$       (d)  $1$

- 6. [MCQ]**

[GATE-CE-2021: 1M]

The value of  $\lim_{x \rightarrow \infty} \frac{x \ln(x)}{1 + x^2}$  is

- (a)  $1.0$       (b)  $0.5$   
 (c)  $\infty$       (d)  $0$

- 7. [NAT]**

[GATE-CE-2021: 1M]

Consider the limit:  $\lim_{x \rightarrow 1} \left( \frac{1}{\ln x} - \frac{1}{x-1} \right)$ .

The limit (correct up to one decimal place) is \_\_\_\_\_.

- 8. [MCQ]**

[GATE-CE-2020: 1M]

The value of  $\lim_{x \rightarrow \infty} \frac{\sqrt{9x^2 + 2020}}{x + 7}$  is

- (a)  $1$       (b)  $3$   
 (c)  $\frac{7}{9}$       (d) Indeterminable

- 9. [MCQ]**

[GATE-CE-2020: 1M]

The true value of  $\ln(2)$  is 0.69. If the value of  $\ln(2)$  is obtained by linear interpolation between  $\ln(1)$  and  $\ln(6)$ , the percentage of absolute error (round off the nearest integer), is

- (a) 35      (b) 69  
 (c) 69      (d) 48



$$f(x) = \begin{cases} 1-x & x \leq 0 \\ x^2, & x > 0 \end{cases}$$

Consider the composition of f and g. i.e.,  $(f \circ g)(x) = f(g(x))$ . The number of discontinuities in  $(f \circ g)(x)$  present in the interval  $(-\infty, 0)$  is:

- |       |       |
|-------|-------|
| (a) 0 | (b) 1 |
| (c) 2 | (d) 4 |

**19. [MCQ] [GATE-EE-2017 : 2M]**

A function  $f(x)$  is defined as

$$f(x) = \begin{cases} e^x & ; \quad x < 1 \\ \ln x + ax^2 + bx & ; \quad x \geq 1 \end{cases},$$

where  $x \in \mathbb{R}$ . Which one of the following statement is TRUE?

- (a)  $f(x)$  is NOT differentiable at  $x = 1$  for any values of a and b.
- (b)  $f(x)$  is differentiable at  $x = 1$  for the unique value of a and b.
- (c)  $f(x)$  is differentiable at  $x = 1$  for all values of a and b such that  $a + b = e$ .
- (d)  $f(x)$  is differentiable at  $x = 1$  for all values of a and b

**20. [MCQ] [GATE-CS-2017:2M]**

$$\text{The value of } \lim_{x \rightarrow 1} \frac{x^7 - 2x^5 + 1}{x^3 - 3x^2 + 2}$$

- |          |                    |
|----------|--------------------|
| (a) Is 0 | (b) Is -1          |
| (c) is 1 | (d) Does not exist |

**21. [NAT] [GATE-CE-2017:1M]**

$$\lim_{x \rightarrow 0} \left( \frac{\tan x}{x^2 - x} \right) \text{ is equal to } \underline{\hspace{2cm}}$$

**22. [MCQ] [GATE-CE-2017:1M]**

Let  $W = f(x,y)$ , where x and y are function of t. Then, according to the chain rule,  $\frac{dw}{dt}$  is equal

- |   |   |
|---|---|
| (a) $\frac{dw}{dx} \frac{dx}{dt} + \frac{dw}{dy} \frac{dy}{dt}$                                 | (b) $\frac{\partial w}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial t}$ |
| (c) $\frac{\partial w}{\partial x} \frac{dx}{dt} + \frac{\partial w}{\partial y} \frac{dy}{dt}$ | (d) $\frac{dw}{dx} \frac{\partial x}{\partial t} + \frac{dw}{dy} \frac{\partial t}{\partial t}$                                 |

**23. [MCQ]**
**[GATE-ME-2017: 1M]**

The value of  $\lim_{x \rightarrow 0} \frac{x^3 - \sin(x)}{x}$  is

- |       |        |
|-------|--------|
| (a) 0 | (b) 3  |
| (c) 1 | (d) -1 |

**24. [NAT]**
**[GATE-EC-2017: 1M]**

The smaller angle (in degrees) between the planes  $x + y + z = 1$  and  $2x - y + 2z = 0$  is \_\_\_\_\_.

**25. [MCQ]**
**[GATE-EC-2016:1M]**

Given the following statements about a function  $f: \mathbb{R} \rightarrow \mathbb{R}$ , select the right option.

P. If  $f(x)$  is continuous at  $x = x_0$ , then it is also differentiable at  $x = x_0$ .

Q. If  $f(x)$  is continuous at  $x = x_0$ , then it may not be differentiable at  $x = x_0$

R. If  $f(x)$  is differentiable at  $x = x_0$ , then it is also continuous at  $x = x_0$

- (a) P is true, Q is false, R is false
- (b) P is false, Q is true, R is true
- (c) P is false, Q is true, R is false
- (d) P is true, Q is false, R is true

**26. [NAT]**
**[GATE-CS-2016: 1M]**

$$\lim_{x \rightarrow 4} \frac{\sin(x-4)}{x-4} = \underline{\hspace{2cm}}$$

**27. [MCQ]**
**[GATE-ME-2016: 1M]**

$\text{Lt}_{x \rightarrow 0} \frac{\log_e(1+4x)}{e^{3x}-1}$  is equal to

- |                   |                    |
|-------------------|--------------------|
| (a) 0             | (b) $\frac{1}{12}$ |
| (c) $\frac{4}{3}$ | (d) 1              |

- |  |   |   |                                   |
|--|---|---|-----------------------------------|
| <b>28. [MCQ]</b>   | <b>[GATE-ME-2016: 2M]</b>                     | <b>34. [NAT]</b>  | <b>[GATE-ME-2015: 1M]</b>         |
| $\lim_{x \rightarrow \infty} \sqrt{x^2 + x - 1} - x$ is  |   | The value of $\lim_{x \rightarrow 0} \left( \frac{-\sin x}{2 \sin x + x \cos x} \right)$ is ____.   |                                   |
| (a) 0<br>(c) $1/2$   | (b) $\infty$<br>(d) $-\infty$                 |   |                                   |
| <b>29. [MCQ]</b>   | <b>[GATE-ME-2016: 1M]</b>                     | <b>35. [NAT]</b>  | <b>[GATE-ME-2015: 2M]</b>         |
| The values of $x$ for which the function   |   | Consider an ant crawling along the curve $(x - 2)^2 + y^2 = 4$ , where $x$ and $y$ are in meters. The ant starts at the point $(4, 0)$ and moves counter-clockwise with a speed of 1.57 meters per second. The time taken by the ant to reach the point $(2, 2)$ is (in seconds) ____ |                                   |
| $f(x) = \frac{x^2 - 3x - 4}{x^2 + 3x - 4}$   |   |   |                                   |
| is NOT continuous are  |   |   |                                   |
| (a) 4 and $-1$<br>(c) $-4$ and 1   | (b) 4 and 1<br>(d) $-4$ and $-1$              |   |                                   |
| <b>30. [NAT]</b>   | <b>[GATE-EE- 2016: 2M]</b>                    | <b>36. [MCQ]</b>  | <b>[GATE-CS-2015: 1M]</b>         |
| Let $S = \sum_{n=0}^{\infty} n\alpha^n$ where $ \alpha  < 1$ . The value of $\alpha$ in the range $0 < \alpha < 1$ , such that $S = 2\alpha$ is _____. |   | $\lim_{x \rightarrow \infty} x^{1/x}$   |                                   |
|  |   | (a) $\infty$<br>(c) 1   | (b) 0<br>(d) Not defined          |
| <b>31. [MCQ]</b>   | <b>[GATE-EE-2015 : 1M]</b>                    | <b>37. [MCQ]</b>  | <b>[GATE- CS-2015:1M]</b>         |
| If a continuous function $f(x)$ does not have a root in the interval $[a, b]$ , then which one of the following statements is TRUE?                    |   | The value of $\lim_{x \rightarrow \infty} (1+x^2)e^{-x}$ is   |                                   |
| (a) $f(a).f(b) = 0$<br>(c) $f(a).f(b) > 0$   | (b) $f(a).f(b) < 0$<br>(d) $f(a)/f(b) \leq 0$ |   |                                   |
| <b>32. [MCQ]</b>   | <b>[GATE-CE-2015:1M]</b>                      |   |                                   |
| $\lim_{x \rightarrow \infty} (1+1/x)^{2x}$ is equal to   |   |   |                                   |
| (a) $e^{-2}$<br>(c) $e$  | (b) 1<br>(d) $e^2$                            | (a) 0<br>(c) 1  | (b) $\frac{1}{2}$<br>(d) $\infty$ |
| <b>33. [MCQ]</b>   | <b>[GATE-ME-2015: 1M]</b>                     | <b>38. [MCQ]</b>  | <b>[GATE-EC-2014: 1M]</b>         |
| The value of $\lim_{x \rightarrow 0} \frac{1 - \cos(x^2)}{2x^4}$ is  |   | The value of $\lim_{x \rightarrow \infty} \left( 1 + \frac{1}{x} \right)^x$ is  |                                   |
| (a) 0<br>(c) $\frac{1}{4}$   | (b) $\frac{1}{2}$<br>(d) Undefined            | (a) $\ln 2$<br>(c) $e$  | (b) 1.0<br>(d) $\infty$           |
| <b>39. [NAT]</b>   | <b>[GATE-CS-2014:2M]</b>                      | <b>40. [MCQ]</b>  | <b>[GATE-CE-2014:1M]</b>          |
| The function $f(x) = x \sin x$ satisfies the following equation: $f'(x) + f(x) + t \cos x = 0$ . The value of $t$ is _____.                            |   | $\lim_{x \rightarrow \infty} \left( \frac{x + \sin x}{x} \right)$ equal to  |                                   |
|  |   | (a) $-\infty$<br>(c) 0  | (b) 1<br>(d) $\infty$             |

**41. [MCQ]****[GATE-CE-2014:2M]**

The expression  $\lim_{\alpha \rightarrow 0} \frac{x^\alpha - 1}{\alpha}$  is equal to

- (a)  $\log x$       (b)  $x \log x$   
 (c) 0      (d)  $\infty$

**42. [MCQ]****[GATE-CE-2014:2M]**

What should be the value of  $\lambda$  such that the function defined below is continuous at  $x = \pi/2$ ?

$$f(x) = \begin{cases} \frac{\lambda \cos x}{2} & \text{if } x \neq \frac{\pi}{2} \\ \frac{\pi}{2} - x & \\ 1 & \text{if } x = \frac{\pi}{2} \end{cases}$$

- (a) 0      (b)  $2/\pi$   
 (c) 1      (d)  $\pi/2$

**43. [MCQ]****[GATE-ME-2014: 1M]**

$\lim_{x \rightarrow 0} \frac{x - \sin x}{1 - \cos x}$  is

- (a) 0      (b) 1  
 (c) 3      (d) Not defined.

**44. [MCQ]****[GATE-ME-2014: 1M]**

$\lim_{x \rightarrow 0} \left( \frac{e^{2x} - 1}{\sin(4x)} \right)$  is equal to

- (a) 0      (b) 0.5  
 (c) 1      (d) 2

**45. [MCQ]****[GATE-ME-2014: 1M]**

If a function is continuous at a point,

- (a) The limit of the function may not exist at the point.  
 (b) The function must be derivable at the point.  
 (c) The limit of the function at the point tends to infinity.  
 (d) The limit must exist at the point and the value of limit should be same as the value of the function at that point.

**46. [MCQ]****[GATE-CS-2013:1M]**

Which one of the following functions is continuous at  $x = 3$ ?

$$(a) f(x) = \begin{cases} 2, & \text{if } x = 3 \\ x - 1, & \text{if } x > 3 \\ \frac{x+3}{3}, & \text{if } x < 3 \end{cases}$$

$$(b) f(x) = \begin{cases} 4, & \text{if } x = 3 \\ 8 - x & \text{if } x \neq 3 \end{cases}$$

$$(c) f(x) = \begin{cases} x + 3, & \text{if } x \leq 3 \\ x - 4 & \text{if } x > 3 \end{cases}$$

$$(d) f(x) = \frac{1}{x^3 - 27}, \text{ if } x \neq 3$$

**47. [MCQ]****[GATE-ME-2013: 1M]**

Choose the CORRECT set of functions, which are linearly dependent.

- (a)  $\sin x, \sin^2 x$  and  $\cos^2 x$   
 (b)  $\cos x, \sin x$  and  $\tan x$   
 (c)  $\cos 2x, \sin^2 x$  and  $\cos^2 x$   
 (d)  $\cos 2x, \sin x$  and  $\cos x$

**48. [MCQ]****[GATE-ME-2012: 1M]**

Consider the function  $f(x) = |x|$  in the interval  $-1 \leq x \leq 1$ . At the point  $x = 0$ ,  $f(x)$  is

- (a) Continuous and differentiable.  
 (b) Non-continuous and differentiable.  
 (c) Continuous and non-differentiable.  
 (d) Neither continuous nor differentiable.

**49. [MCQ]****[GATE-ME-2012: 1M]**

$$\lim_{x \rightarrow 0} \left( \frac{1 - \cos x}{x^2} \right)$$

- (a) 1 / 4      (b) 1 / 2  
 (c) 1      (d) 2

**50. [MCQ]****[GATE-ME-2011: 1M]**

What is  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta}$  equal to?

- (a) 0      (b)  $\sin \theta$   
 (c) 0      (d) 1

**51. [MCQ]****[GATE- ME -2010: 1M]**The function  $y = |2 - 3x|$ 

- (a) is continuous  $\forall x \in \mathbb{R}$  and differentiable  $\forall x \in \mathbb{R}$
- (b) is continuous  $\forall x \in \mathbb{R}$  and differentiable  $\forall x \in \mathbb{R}$  except at  $x = 3/2$
- (c) is continuous  $\forall x \in \mathbb{R}$  and differentiable  $\forall x \in \mathbb{R}$  except at  $x = 2/3$
- (d) is continuous  $\forall x \in \mathbb{R}$  except at  $x = 3$  and differentiable  $\forall x \in \mathbb{R}$ .

**52. [MCQ]****[GATE-CS-2010:1M]**What is the value of  $\lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^{2n}$ 

- (a) 0
- (b)  $e^{-2}$
- (c)  $e^{-1/2}$
- (d) 1

**53. [MCQ]****[GATE-CE-2010:1M]**The  $\lim_{x \rightarrow 0} \frac{\sin\left[\frac{2}{3}x\right]}{x}$  is

- (a)  $2/3$
- (b) 1
- (c)  $3/2$
- (d)  $\infty$

**54. [MCQ]****[GATE-CS-2008:2M]** $\lim_{x \rightarrow \infty} \frac{x - \sin x}{x + \cos x}$  equals

- (a) 1
- (b)  $-1$
- (c)  $\infty$
- (d)  $-\infty$

**55. [MCQ]****[GATE-ME-2008: 1M]**The value of  $\lim_{x \rightarrow 8} \frac{x^{1/3} - 2}{(x - 8)}$  is

- (a)  $\frac{1}{16}$
- (b)  $\frac{1}{12}$
- (c)  $\frac{1}{8}$
- (d)  $\frac{1}{4}$

**Applications of Differentiation****56. [NAT]****[GATE 2023-EE :2M]**In the following differential equation, the numerically obtained value of  $y(t)$ , at  $t = 1$ , is \_\_\_\_\_. (Round off to 2 decimal places)

$$\frac{dy}{dt} = \frac{e^{-\alpha t}}{2 + \alpha t}, \alpha = 0.01 \text{ and } y(0) = 0$$

**57. [MCQ]****[GATE-2023-EC : 2M]**

Consider the following series:

$$\sum_{n=1}^{\infty} \frac{n^d}{c^n}$$

For which of the following combinations of  $c, d$  values does this series converge?

- |                        |                     |
|------------------------|---------------------|
| (a) $c = 1, d = -1$    | (b) $c = 2, d = 1$  |
| (c) $c = 0.5, d = -10$ | (d) $c = 1, d = -2$ |

**58. [MCQ]****[GATE-2023-EC : 2M]**The function  $f(x) = 8 \log_e x - x^2 + 3$  attains its minimum over the interval  $[1, e]$  at  $x = _____$ .(Here  $\log_e x$  is the natural logarithm of  $x$ .)

- |         |                     |
|---------|---------------------|
| (a) 2   | (b) 1               |
| (c) $e$ | (d) $\frac{1+e}{2}$ |

**59. [MCQ]****[GATE-CS-2023: 1M]**Let  $f(x) = x^3 + 15x^2 - 33x - 36$  be a real-valued function.

Which of the following statements is/are TRUE?

- (a)  $f(x)$  does not have a local maximum.
- (b)  $f(x)$  has a local maximum.
- (c)  $f(x)$  does not have a local minimum.
- (d)  $f(x)$  has a local minimum.

**60. [MCQ]****[GATE-CE-2023:2M]**For the function  $f(x) = e^x |\sin x|; x \in \mathbb{R}$ , which of the following statements is/are TRUE?

- (a) The function is continuous at all  $x$
- (b) The function is differentiable at all  $x$
- (c) The function is periodic
- (d) The function is bounded

**61. [NAT]****[GATE-ME-2023: 2M]**

The smallest perimeter that a rectangle with area of 4 square units can have is \_\_\_\_\_. units. (Answer in integer)

**62. [MCQ]****[GATE-CE-2022:1M]**

Consider the polynomial  $f(x) = x^3 - 6x^2 + 11x - 6$  on the domain S, given by  $1 \leq x \leq 3$ . The first and second derivatives are  $f'(x)$  and  $f''(x)$ . Consider the following statements:

- I) The given polynomial is zero at the boundary points  $x = 1$  and  $x = 3$
- II) There exists one local maxima of  $f(x)$  within the domain S.
- III) The second derivative  $f''(x) > 0$  throughout the domains S.
- IV) There exists one local minima  $f(x)$  within the domain S.
- (a) Only statements II and IV are correct
- (b) Only statements I and IV are correct.
- (c) Only statements I, II and III are correct.
- (d) Only statements I, II and IV are correct

**63. [MSQ]****[GATE-CE-2022:2M]**

Let  $\max(a, b)$  denote the maximum of two real numbers a and b. Which of the following statement(s) is/are TRUE about the function  $f(x) = \max\{3 - x, x - 1\}$ ?

- (a) It is continuous on its domain
- (b) It has a local minimum at  $x = 2$
- (c) It has a local maximum at  $x = 2$
- (d) It is differentiable on its domain

**64. [MCQ]****[GATE-ME-2021: 2M]**

Let the superscript T represent the transpose operation.

Consider the function  $f(x) = \frac{1}{2} x^T Q x - r^T x$ , where x and r are  $n \times 1$  vectors and Q is a symmetric  $n \times n$  matrix.

The stationary point of  $f(x)$  is

- (a)  $Q^T r$
- (b)  $Q^{-1} r$
- (c)  $\frac{r}{r^T r}$
- (d) r

**65. [NAT]****[GATE-CS-2021:1M]**

Suppose that  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function on the interval  $[-3, 3]$  and a differentiable function in the interval  $(-3, 3)$  such that for every  $x$  in the interval,  $|f'(x)| \leq 2$ . If  $f(-3) = 7$ , then  $f(3)$  is at most \_\_\_\_\_.

**66. [MCQ]****[GATE-EE-2021 : 1M]**

Let  $f(x)$  be a real-valued function such that  $f'(x_0) = 0$  for some  $x_0 \in (0, 1)$ , and  $f''(x) > 0$  for all  $x \in (0, 1)$ . Then  $f(x)$  has

- (a) no local minimum in  $(0, 1)$
- (b) one local maximum in  $(0, 1)$
- (c) exactly one local minimum in  $(0, 1)$
- (d) two distinct local minima in  $(0, 1)$

**67. [NAT]****[GATE-EE-2021: 1M]**

Suppose the circles  $x^2 + y^2 = 1$  and  $(x - 1)^2 + (y - 1)^2 = r^2$  intersect each other orthogonally at the point  $(u, v)$ . Then  $u + v = \underline{\hspace{2cm}}$ .

**68. [MCQ]****[GATE-EE-2021: 2M]**

In the open interval  $(0, 1)$ , the polynomial  $p(x) = x^4 - 4x^3 + 2$  has

- (a) two real roots
- (b) one real root
- (c) three real roots
- (d) no real roots

**69. [MSQ]****[GATE-EE-2020: 1M]**

$ax^3 + bx^2 + cx + d$  is a polynomial on real x over real coefficients a, b, c, d wherein  $a \neq 0$ . Which of the following statements is true?

- (a) d can be chosen to ensure that  $x=0$  is a root for any given set a, b, c. identical.
- (b) No choice of coefficients can make all roots identical.
- (c) a, b, c, d can be chosen to ensure that all roots are complex
- (d) c alone cannot ensure that all roots are real

**70. [MCQ]****[GATE-EE-2020: 2M]**

For real numbers, x and y, with  $y = 3x^2 + 3x + 1$ , the maximum and minimum value of y for  $x \in [-2, 0]$  are respectively, \_\_\_\_\_.

- (a) 7 and  $1/4$
- (b) 7 and 1
- (c) -2 and  $-1/2$
- (d) 1 and  $1/4$

**71. [MCQ]****[GATE-CS-2020:1M]**

Consider the functions

I:  $e^{-x}$

II:  $x^2 - \sin x$









**117.[MCQ]****[GATE-EC-2010:2M]**

- If  $e^y = x^{1/x}$ , then y has a
- maximum at  $x = e$
  - minimum at  $x = e$
  - maximum at  $x = e^{-1}$
  - minimum at  $x = e^{-1}$

**118.[MCQ]****[GATE-EC-2009: 2M]**

- The Taylor series expansion of  $\frac{\sin x}{x - \pi}$  at  $x = \pi$  is given by

- $1 + \frac{(x - \pi)^2}{3!} + \dots$
- $-1 - \frac{(x - \pi)^2}{3!} + \dots$
- $1 - \frac{(x - \pi)^2}{3!} + \dots$
- $-1 + \frac{(x - \pi)^2}{3!} + \dots$

**119.[MCQ]****[GATE-ME-2009: 2M]**

- The distance between the origin and the point nearest to it on the surface  $z^2 = 1 + xy$  is

- 1
- $\frac{\sqrt{3}}{2}$
- $\sqrt{3}$
- 2

**120.[MCQ]****[GATE-EE-2009 : 1M]**

- The Taylor series expansion of  $\frac{\sin x}{x - \pi}$  at  $x = \pi$  is given by

- $1 + \frac{(x - \pi)^2}{3!} + \dots$
- $-1 - \frac{(x - \pi)^2}{3!} + \dots$
- $1 - \frac{(x - \pi)^2}{3!} + \dots$
- $-1 + \frac{(x - \pi)^2}{3!} + \dots$

**121.[MCQ]****[GATE-EE -2008 : 2M]**

- Consider function  $f(x) = (x^2 - 4)^2$  where x is real number. Then the function has

- only one minima
- only two minima
- three minima
- three maxima

**122.[MCQ]****[GATE-EC-2008: 2M]**

- In the Taylor series expansion of  $\exp(x) + \sin(x)$  about the point  $x = \pi$ , the coefficient of  $(x - \pi)^2$  is

- $\exp(\pi)$
- $0.5 \exp(\pi)$
- $\exp(\pi) + 1$
- $\exp(\pi) - 1$

**123.[MCQ]****[GATE-EC-2008:1M]**

- For real values of x, the minimum value of the function  $f(x) = \exp(x) + \exp(-x)$  is

- 2
- 1
- 0.5
- 0

**124.[MCQ]****[GATE-EC-2008: 1M]**

- Which of the following functions would have only odd powers of x in its Taylor series expansion about the point  $x = 0$ ?

- $\sin(x^3)$
- $\sin(x^2)$
- $\cos(x^3)$
- $\cos(x^2)$

**125.[MCQ]****[GATE-CS-2008:2M]**

- A point on a curve is said to be an extremum if it is a local minimum or a local maximum. The number of distinct extrema for the curve  $3x^4 - 16x^3 + 24x^2 + 37$  is

- 0
- 1
- 2
- 3

**126.[MCQ]****[GATE-ME-2008: 1M]**

- In the Taylor series expansion of  $e^x$  about  $x = 2$ , the coefficient of  $(x - 2)^4$  is

- $\frac{1}{4!}$
- $\frac{e^4}{4!}$
- $\frac{e^2}{4!}$
- $\frac{e^4}{4!}$

**Integral Calculus****127.[MCQ]****[GATE-CE-2023:1M]**

- For the integral

$$I = \int_{-1}^1 \frac{1}{x^2} dx$$

- which of the following statements is TRUE?



**138.[MCQ]****[GATE-CS-2017:1M]**

If  $f(x) = R \sin\left(\frac{\pi x}{2}\right) + S$ ,  $f'\left(\frac{1}{2}\right) = \sqrt{2}$  and  $\int_0^1 f(x) dx = \frac{2R}{\pi}$  then the constants  $R$  and  $S$  are, respectively

- |  |  |
|--|--|
| (a) $\frac{2}{\pi}$ and $\frac{16}{\pi}$ | (b) $\frac{2}{\pi}$ and 0                |
| (c) $\frac{4}{\pi}$ and 0                | (d) $\frac{4}{\pi}$ and $\frac{16}{\pi}$ |

**139.[MCQ]****[GATE-CE-2017:2M]**

Consider the following definite integral:

$$I = \int_0^1 \frac{(\sin^{-1} x)^2}{\sqrt{1-x^2}} dx$$

The value of the integral is

- |                        |                        |
|------------------------|------------------------|
| (a) $\frac{\pi^3}{24}$ | (b) $\frac{\pi^3}{12}$ |
| (c) $\frac{\pi^3}{48}$ | (d) $\frac{\pi^3}{64}$ |

**140.[MCQ]****[GATE-CE-2017:1M]**

Let  $x$  be a continuous variable defined over the interval  $(-\infty, \infty)$  and  $f(x) = e^{-x-e^{-x}}$ . The integral  $g(x) = \int f(x) dx$  is equal to

- |                  |                   |
|------------------|-------------------|
| (a) $e^{e^{-x}}$ | (b) $e^{-e^{-x}}$ |
| (c) $e^{-e^x}$   | (d) $e^{-x}$      |

**141.[MCQ]****[GATE-ME-2017: 2M]**

A parametric curve defined by  $x = \cos\left(\frac{\pi u}{2}\right)$ ,  $y = \sin\left(\frac{\pi u}{2}\right)$  in the range  $0 \leq u \leq 1$  is rotated about the X-axis by 360 degrees. Area of the surface generated is

- |                     |            |
|---------------------|------------|
| (a) $\frac{\pi}{2}$ | (b) $\pi$  |
| (c) $2\pi$          | (d) $4\pi$ |

**142.[NAT]****[GATE-CS-2015: 2M]**

$$\int_{\pi}^{\frac{2}{\pi}} \frac{\cos\left(\frac{1}{x}\right)}{x^2} dx = \underline{\hspace{2cm}}$$

**143.[MCQ]****[GATE-CS-2015:2M]**

If for non-zero  $x$ ,  $a \cdot f(x) + b \cdot f\left(\frac{1}{x}\right) = \frac{1}{x} - 25$  where

$$a \neq b \text{ then } \int_1^2 f(x) dx$$

- |  |
|--|
| (a) $\frac{1}{a^2 - b^2} \left[ a(\ln 2 - 25) + \frac{47b}{2} \right]$   |
| (b) $\frac{1}{a^2 - b^2} \left[ a(2 \ln 2 - 25) - \frac{47b}{2} \right]$ |
| (c) $\frac{1}{a^2 - b^2} \left[ a(2 \ln 2 - 25) + \frac{47b}{2} \right]$ |
| (d) $\frac{1}{a^2 - b^2} \left[ a(\ln 2 - 25) - \frac{47b}{2} \right]$   |

**144.[NAT]****[GATE-CS-2014:1M]**

$$\text{If } \int_0^{2x} |x \cdot \sin x| dx = k\pi \text{ then the value of } k \text{ is equal to } \underline{\hspace{2cm}}$$

**145.[MCQ]****[GATE-CS-2014:2M]**

The value of the integral given below is  $\int_0^{\pi} x^2 \cos x dx$

- |             |            |
|-------------|------------|
| (a) $-2\pi$ | (b) $\pi$  |
| (c) $-\pi$  | (d) $2\pi$ |

**146.[NAT]****[GATE-CE-2014:2M]**

The value determined from

$\iiint 8xyz dV$  for  $V = [2,3] \times [1,2] \times [0,1]$  will be (in integer)



**158.[MCQ]****[GATE-ME-2008: 2M]**

Which of the following integrals is unbounded?

- |                                      |   |
|--------------------------------------|---|
| (a) $\int_0^{\pi/4} \tan x \, dx$    | (b) $\int_0^{\infty} \frac{1}{x^2 + 1} \, dx$ |
| (c) $\int_0^{\infty} x e^{-x} \, dx$ | (d) $\int_0^1 \frac{1}{1-x} \, dx$            |

**Applications of Integration****159.[MSQ]****[GATE-EE-2023: 2M]**

Consider the following equation in a 2-D real-space

$$|x_1|^p + |x_2|^p = 1 \text{ for } p > 0$$

Which of the following statement(s) is/are true.

- (a) When  $p = 2$ , the area enclosed by the curve is  $\pi$
- (b) When  $p$  tends to  $\infty$ , the area enclosed by the curve tends to 4.
- (c) When  $p$  tends to 0, the area enclosed by the curve is 1.
- (d) When  $p = 1$ , the area enclosed by the curve is 2.

**160.[NAT]****[GATE-ME-2023: 1M]**

A linear transformation maps a point  $(x, y)$  in the plane to the point  $(\hat{x}, \hat{y})$  according to the rule


$$\hat{x} = 3y, \hat{y} = 2x.$$

Then, the disc  $x^2 + y^2 < 1$  gets transformed to a region with an area equal to \_\_\_\_\_. (Rounded off to two decimals)

Use  $\pi = 3.14$

**161.[NAT]****[GATE-EC-2023 : 2M]**


The value of the integral  $\iint_R xy \, dx \, dy$  over the region  $R$ , given in the figure, is \_\_\_\_\_ (rounded off to the nearest integer).

**162.[NAT]****[GATE-EC-2022 : 2M]**

The value of the integral,

$$\iint_D 3(x^2 + y^2) \, dx \, dy,$$

where  $D$  is the shaded triangular region shown in the diagram, is \_\_\_\_\_ (rounded off to the nearest integer).

**163.[MCQ]****[GATE-CE-2020:1M]**


The area of an ellipse represented by an equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ is}$$

- |                         |                        |
|-------------------------|------------------------|
| (a) $\frac{4\pi ab}{3}$ | (b) $\pi ab$           |
| (c) $\frac{\pi ab}{2}$  | (d) $\frac{\pi ab}{4}$ |

**164.[MCQ]****[GATE-ME-2019: 1M]**

A parabola  $x = y^2$  with  $0 \leq x \leq 1$  is shown in the figure. The volume of the solid of rotation obtained by rotating the shaded area by  $360^\circ$  around the x-axis is



- |                     |                     |
|---------------------|---------------------|
| (a) $\frac{\pi}{4}$ | (b) $\frac{\pi}{2}$ |
| (c) $\pi$           | (d) $2\pi$          |

**165.[MCQ]****[GATE-CE-2018:1M]**

The value of the integral  $\int_0^\pi x \cos^2 x dx$

(a)  $\frac{\pi^2}{8}$

(b)  $\frac{\pi^2}{4}$

(c)  $\frac{\pi^2}{2}$

(d)  $\pi^2$

**166.[NAT]****[GATE-CE-2016:2M]**

The area between the parabola  $x^2 = 8y$  and the straight-line  $y = 8$  is \_\_\_\_\_.

**167.[MCQ]****[GATE-CE-2016:2M]**

The area of the region bounded by the parabola  $y = x^2 + 1$  and the straight-line  $x + y = 3$  is

(a)  $\frac{59}{6}$

(b)  $\frac{9}{2}$

(c)  $\frac{10}{3}$

(d)  $\frac{7}{6}$

**168.[NAT]****[GATE-ME-2015: 2M]**

Consider a spatial curve in three-dimensional space given in parametric form by

$$x(t) = \cos t, y(t) = \sin t, z(t) = \frac{2}{\pi} t; \quad 0 \leq t \leq \frac{\pi}{2}.$$

The length of the curve is \_\_\_\_\_.

**169.[MCQ]****[GATE-ME-2012: 1M]**

The area enclosed between the straight-line  $y = x$  and the parabola  $y = x^2$  in the  $x - y$  plane is

(a) 1/6

(b) 1/4

(c) 1/3

(d) 1/2

**170.[MCQ]****[GATE-2010-CE:2M]**

The parabolic cable is held between two supports at the same level. The horizontal span between the supports is  $L$ . The sag at the mid-span is  $h$ . The equation of the parabola is  $y = 4h \frac{x^2}{L^2}$ , where  $x$  is the horizontal coordinate and  $y$  is the vertical coordinate with the

origin at the centre of the cable. The expression for the total length of the cable is

(a)  $\int_0^L \sqrt{1 + 64 \frac{h^2 x^2}{L^4}} dx$

(b)  $\int_0^L \sqrt{1 + 64 \frac{h^2 x^2}{L^4}} dx$

(c)  $2 \int_0^{L/2} \sqrt{1 + 64 \frac{h^2 x^2}{L^4}} dx$

(d)  $2 \int_0^{L/2} \sqrt{1 + 64 \frac{h^2 x^2}{L^4}} dx$

**171.[MCQ]****[GATE-ME-2010: 1M]**

The parabolic arc  $y = \sqrt{x}$ ,  $1 \leq x \leq 2$  is revolved around the  $x$ -axis. The volume of the solid revolution is

- |              |              |
|--------------|--------------|
| (a) $\pi/4$  | (b) $\pi/2$  |
| (c) $3\pi/4$ | (d) $3\pi/2$ |

**172.[MCQ]****[GATE-ME-2009: 2M]**

The area enclosed between the curves  $y^2 = 4x$  and  $x^2 = 4y$  is

- |                    |        |
|--------------------|--------|
| (a) $\frac{16}{3}$ | (b) 8  |
| (c) $\frac{32}{3}$ | (d) 16 |

**173.[MCQ]****[GATE-ME-2008: 2M]**

The length of curve  $y = \frac{2}{3}x^{3/2}$  between  $x = 0$  and  $x = 1$

- is
- |          |          |
|----------|----------|
| (a) 0.27 | (b) 0.67 |
| (c) 1    | (d) 1.22 |

### Multi Variable Calculus

**174.[NAT]****[GATE-CS-2023:1M]**

The value of the definite integral

$$\int_{-3}^3 \int_{-2}^2 \int_{-1}^1 (4x^2 y - z^3) dz dy dx$$

is \_\_\_\_\_ . (Rounded off to the nearest integer)

**175.[MCQ]****[GATE-ME-2021: 2M]**

The value of  $\int_0^{\pi/2} \int_0^{\cos\theta} r \sin\theta dr d\theta$  is

- (a) 0
- (b)  $\frac{1}{6}$
- (c)  $\frac{4}{3}$
- (d)  $\pi$

**176.[MCQ]****[GATE-EC-2020: 1M]**

The partial derivative of the function

$$f(x, y, z) = e^{1-x} \cos y + x z e^{-1/(1+y^2)}$$

with respect to x at the point (1, 0, e) is

- (a) -1
- (b) 0
- (c) 1
- (d)  $\frac{1}{e}$

**177.[MCQ]****[GATE-ME-2020: 1M]**

Let  $I = \int_{x=0}^1 \int_{y=0}^{x^2} xy^2 dy dx$ . Then, I may also be expressed as

- (a)  $\int_{y=0}^1 \int_{x=0}^{\sqrt{y}} xy^2 dx dy$
- (b)  $\int_{y=0}^1 \int_{x=\sqrt{y}}^0 yx^2 dx dy$
- (c)  $\int_{y=0}^1 \int_{x=\sqrt{y}}^1 xy^2 dx dy$
- (d)  $\int_{y=0}^1 \int_{x=0}^{\sqrt{y}} yx^2 dx dy$

**178.[MCQ]****[GATE-EC-2018: 1M]**

Let,  $f(x, y) = \frac{ax^2 + by^2}{xy}$ , where a and b are constants.

If  $\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y}$  at  $x = 1$  and  $y = 2$ , then the relation between

a and b is

- (a)  $a = \frac{b}{4}$
- (b)  $a = \frac{b}{2}$
- (c)  $a = 2b$
- (d)  $a = 4b$

**179.[NAT]****[GATE-EC-2018:2M]**

Let  $r = x^2 + y - z$  and  $z^3 - xy + yz + y^3 = 1$ . Assume that x and y are independent variables. At  $(x, y, z) = (2, -1, 1)$ , the value (correct to two decimal places) of  $\frac{\partial r}{\partial x}$  is

**180.[NAT]****[GATE-EE-2017: 1M]**


Consider a function  $f(x, y, z)$  given by

$$f(x, y, z) = (x^2 + y^2 - 2z^2)(y^2 + z^2)$$

The partial derivative of this function with respect x at the point  $x = 2, y = 1, z = 3$  is \_\_\_\_\_.

**181.[NAT]****[GATE-EE-2017: 1M]**

Let  $I = c \int_R xy^2 dx dy$  where R is the region shown in the figure and  $c = 6 \times 10^{-4}$ . The value of I equals \_\_\_\_\_. (Give the answer up to two decimal places).

**182.[NAT]****[GATE-EC-2017:2M]**

A three-dimensional region R of finite volume is described by  $x^2 + y^2 \leq z^3; 0 \leq z \leq 1$ . Where x, y, z are real. The volume of R (up to two decimal place) is \_\_\_\_\_.

**183.[NAT]****[GATE-EC-2016: 2M]**

The region specified by  $\{(\rho, \phi, z) : 3 \leq \rho \leq 5, \frac{\pi}{8} \leq \phi \leq \frac{\pi}{4}, 3 \leq z \leq 4.5\}$  in cylindrical coordinates has

volume of \_\_\_\_\_.

**184.[NAT]****[GATE-EC-2016: 2M]**

A triangle in the xy-plane is bounded by the straight lines  $2x = 3y$ ,  $y = 0$  and  $x = 3$ . The volume above the triangle and under the plane  $x + y + z = 6$  is \_\_\_\_\_.

**185.[NAT]****[GATE-EE-2015: 2M]**

The volume enclosed by the surface  $f(x, y) = e^x$  over the triangle bounded by the lines  $x = y$ ;  $x = 0$ ;  $y = 1$  in the xy plane is \_\_\_\_\_.

**186.[MCQ]****[GATE-EC-2015: 1M]**

The contour on the x-y plane, where the partial derivative of  $x^2 + y^2$  with respect to y is equal to the partial derivative of  $6y + 4x$  with respect to x, is

- (a)  $y = 2$       (b)  $x = 2$   
 (c)  $x + y = 4$       (d)  $x - y = 0$

**187.[MCQ]****[GATE-EE-2014: 2M]**

To evaluate the double integral

$$\int_0^8 \left( \int_{y/2}^{(y/2)+1} \left( \frac{2x-y}{2} \right) dx \right) dy, \text{ we make the substitution}$$

$u = \left( \frac{2x-y}{2} \right)$  and  $v = \frac{y}{2}$ . The integral will reduce to

- (a)  $\int_0^4 \left( \int_0^1 2udu \right) dv$   
 (c)  $\int_0^4 \left( \int_0^1 udu \right) dv$       (d)  $\int_0^4 \left( \int_0^2 udu \right) dv$

**188.[NAT]****[GATE-EC-2014: 2M]**

The volume under the surface  $z(x,y) = x + y$  and above the triangle in the x - y plane defined by  $\{0 \leq y \leq x \text{ and } 0 \leq x \leq 12\}$  is \_\_\_\_\_

**189.[MCQ]****[GATE-ME-2014: 2M]**

The value of the integral  $\iint_0^2 e^{x+y} \cdot dy \cdot dx$  is

- (a)  $\frac{1}{2}(e-1)$       (b)  $\frac{1}{2}(e^2-1)^2$   
 (c)  $\frac{1}{2}(e^2-e)$       (d)  $\frac{1}{2}\left(e-\frac{1}{e}\right)^2$

**190.[MCQ]****[GATE-EE-2009: 2M]**

$F(x, y)$  is a continuous function defined over  $(x, y) \in [0, 1] \times [0, 1]$ . Given the two constraints,  $x > y^2$  and  $y > x^2$ , the volume under  $f(x, y)$  is

$$(a) \int_{y=0}^1 \int_{x=y^2}^{x=\sqrt{y}} f(x, y) dx dy$$


$$(b) \int_{y=x}^{y=1} \int_{x=y^2}^{x=1} f(x, y) dx dy$$

$$(c) \int_{y=0}^{y=1} \int_{x=0}^{x=1} f(x, y) dx dy$$

$$(d) \int_{y=0}^{y=\sqrt{x}} \int_{x=0}^{x=\sqrt{y}} f(x, y) dx dy$$

**191.[MCQ]****[GATE-ME-2009: 2M]**

A path AB in the form of one quarter of a circle of unit radius is shown in the figure. Integration of  $(x + y)^2$  on path of AB traversed in a counter-clockwise sense



$$(a) \frac{\pi}{2} - 1$$


$$(b) \frac{\pi}{2} + 1$$

$$(c) \frac{\pi}{2}$$

$$(d) 1$$

**192.[MCQ]****[GATE-ME-2008: 2M]**

Consider the shaded triangular region P shown in the figure. Where is  $\iint_P xy \, dx \, dy$  ?



$$(a) \frac{1}{6}$$

$$(b) \frac{2}{9}$$

$$(c) \frac{7}{16}$$

$$(d) 1$$




**ANSWER KEY**


- |             |                  |             |                  |             |              |             |                |
|-------------|------------------|-------------|------------------|-------------|--------------|-------------|----------------|
| <b>1.</b>   | (b)              | <b>2.</b>   | (a)              | <b>3.</b>   | (−0.5)       | <b>4.</b>   | (0.25)         |
| <b>5.</b>   | (c)              | <b>6.</b>   | (d)              | <b>7.</b>   | (0.5)        | <b>8.</b>   | (b)            |
| <b>9.</b>   | (d)              | <b>10.</b>  | (c)              | <b>11.</b>  | (c)          | <b>12.</b>  | (a)            |
| <b>13.</b>  | (c)              | <b>14.</b>  | (d)              | <b>15.</b>  | (d)          | <b>16.</b>  | (b)            |
| <b>17.</b>  | (d)              | <b>18.</b>  | (a)              | <b>19.</b>  | (b)          | <b>20.</b>  | (c)            |
| <b>21.</b>  | (−1)             | <b>22.</b>  | (c)              | <b>23.</b>  | (d)          | <b>24.</b>  | (54 to 55)     |
| <b>25.</b>  | (b)              | <b>26.</b>  | (1)              | <b>27.</b>  | (c)          | <b>28.</b>  | (c)            |
| <b>29.</b>  | (c)              | <b>30.</b>  | (0.28 to 0.31)   | <b>31.</b>  | (c)          | <b>32.</b>  | (d)            |
| <b>33.</b>  | (c)              | <b>34.</b>  | (−0.35 to −0.30) | <b>35.</b>  | (1.9 to 2.1) | <b>36.</b>  | (c)            |
| <b>37.</b>  | (a)              | <b>38.</b>  | (c)              | <b>39.</b>  | (−2)         | <b>40.</b>  | (b)            |
| <b>41.</b>  | (a)              | <b>42.</b>  | (c)              | <b>43.</b>  | (a)          | <b>44.</b>  | (b)            |
| <b>45.</b>  | (d)              | <b>46.</b>  | (a)              | <b>47.</b>  | (c)          | <b>48.</b>  | (c)            |
| <b>49.</b>  | (b)              | <b>50.</b>  | (d)              | <b>51.</b>  | (c)          | <b>52.</b>  | (b)            |
| <b>53.</b>  | (a)              | <b>54.</b>  | (a)              | <b>55.</b>  | (b)          | <b>56.</b>  | (0.50)         |
| <b>57.</b>  | (b)              | <b>58.</b>  | (b)              | <b>59.</b>  | (d)          | <b>60.</b>  | (a)            |
| <b>61.</b>  | (7.999 to 8.001) | <b>62.</b>  | (d)              | <b>63.</b>  | (a, b)       | <b>64.</b>  | (b)            |
| <b>65.</b>  | (19)             | <b>66.</b>  | (c)              | <b>67.</b>  | (1)          | <b>68.</b>  | (b)            |
| <b>69.</b>  | (a, d)           | <b>70.</b>  | (a)              | <b>71.</b>  | (a)          | <b>72.</b>  | (5.18 to 5.2)  |
| <b>73.</b>  | (c)              | <b>74.</b>  | (a)              | <b>75.</b>  | (c)          | <b>76.</b>  | (d)            |
| <b>77.</b>  | (11.5 to 12.5)   | <b>78.</b>  | (7)              | <b>79.</b>  | (a)          | <b>80.</b>  | (c)            |
| <b>81.</b>  | (−100)           | <b>82.</b>  | (c)              | <b>83.</b>  | (0 to 0)     | <b>84.</b>  | (b)            |
| <b>85.</b>  | (b)              | <b>86.</b>  | (d)              | <b>87.</b>  | (d)          | <b>88.</b>  | (−5.1 to −4.9) |
| <b>89.</b>  | (2)              | <b>90.</b>  | (1)              | <b>91.</b>  | (b)          | <b>92.</b>  | (b)            |
| <b>93.</b>  | (c)              | <b>94.</b>  | (d)              | <b>95.</b>  | (a)          | <b>96.</b>  | (a)            |
| <b>97.</b>  | (6)              | <b>98.</b>  | (d)              | <b>99.</b>  | (a)          | <b>100.</b> | (c)            |
| <b>101.</b> | (−1.00 to −0.94) | <b>102.</b> | (2)              | <b>103.</b> | (b)          | <b>104.</b> | (c)            |
| <b>105.</b> | (a)              | <b>106.</b> | (b)              | <b>107.</b> | (c)          | <b>108.</b> | (c)            |
| <b>109.</b> | (b)              | <b>110.</b> | (d)              | <b>111.</b> | (c)          | <b>112.</b> | (c)            |
| <b>113.</b> | (c)              | <b>114.</b> | (b)              | <b>115.</b> | (d)          | <b>116.</b> | (a)            |
| <b>117.</b> | (a)              | <b>118.</b> | (c)              | <b>119.</b> | (a)          | <b>120.</b> | (d)            |
| <b>121.</b> | (b)              | <b>122.</b> | (b)              | <b>123.</b> | (a)          | <b>124.</b> | (a)            |
| <b>125.</b> | (b)              | <b>126.</b> | (c)              | <b>127.</b> | (d)          | <b>128.</b> | (b)            |
| <b>129.</b> | (a)              | <b>130.</b> | (0)              | <b>131.</b> | (c)          | <b>132.</b> | (a, c, d)      |
| <b>133.</b> | (b)              | <b>134.</b> | (2.090 to 2.104) | <b>135.</b> | (0.50)       | <b>136.</b> | (a)            |
| <b>137.</b> | (0.27 to 0.3)    | <b>138.</b> | (c)              | <b>139.</b> | (a)          | <b>140.</b> | (b)            |

- |                           |                           |                          |                              |
|---------------------------|---------------------------|--------------------------|------------------------------|
| <b>141.</b> (c)           | <b>142.</b> (-1)          | <b>143.</b> (a)          | <b>144.</b> (4)              |
| <b>145.</b> (a)           | <b>145.</b> (15)          | <b>147.</b> (b)          | <b>148.</b> (c)              |
| <b>149.</b> (d)           | <b>150.</b> (c)           | <b>151.</b> (c)          | <b>152.</b> (*)              |
| <b>153.</b> (b)           | <b>154.</b> (b)           | <b>155.</b> (d)          | <b>156.</b> (d)              |
| <b>157.</b> (a)           | <b>158.</b> (d)           | <b>159.</b> (a, b, d)    | <b>160.</b> (18.80 to 18.90) |
| <b>161.</b> (0)           | <b>162.</b> (171)         | <b>163.</b> (b)          | <b>164.</b> (b)              |
| <b>165.</b> (b)           | <b>166.</b> (85 to 85.5)  | <b>167.</b> (b)          | <b>168.</b> (1.85 to 1.87)   |
| <b>169.</b> (a)           | <b>170.</b> (d)           | <b>171.</b> (d)          | <b>172.</b> (a)              |
| <b>173.</b> (d)           | <b>174.</b> (0)           | <b>175.</b> (b)          | <b>176.</b> (b)              |
| <b>177.</b> (c)           | <b>178.</b> (d)           | <b>179.</b> (4.4 to 2.6) | <b>180.</b> (40)             |
| <b>181.</b> (1.00)        | <b>182.</b> (0.7 to 0.85) | <b>183.</b> (4 to 5)     | <b>184.</b> (10)             |
| <b>185.</b> (0.7 to 0.76) | <b>186.</b> (a)           | <b>187.</b> (b)          | <b>188.</b> (864)            |
| <b>189.</b> (b)           | <b>190.</b> (a)           | <b>191.</b> (b)          | <b>192.</b> (a)              |

## SOLUTIONS

**1. (b)**

Let the given function be  $f(x)$ .



- (i)  $f(x)$  is symmetric about y-axis.  
 $\Rightarrow f(-x) = -f(x)$
- (ii)  $f(x) > 0 \forall x \in [-4, 4]$
- (iii) At  $x = 2$ ,  $f(x) = 0$
- (iv)  $0 \leq f(x) \leq 2 \forall x \in [-4, 4]$

The only function that satisfies the above conditions is option B.

$$\Rightarrow f(x) = |2 - |x||$$



Scan for Video solution



**2. (a)**

Given limit is

$$P = \lim_{x \rightarrow \pi} \left( \frac{x^2 + \alpha x + 2\pi^2}{x - \pi + 2 \sin x} \right)$$

$$P = \lim_{x \rightarrow \pi} \frac{x^2 + \alpha x + 2\pi^2}{x - \pi + 2 \sin x} = \frac{3\pi^2 + \alpha\pi}{0}$$

But for 'P' has to be finite,

$$3\pi^2 + \alpha\pi = 0 \Rightarrow \alpha = -3\pi$$

If  $\alpha = -3\pi$ ,

$$P = \lim_{x \rightarrow \pi} \frac{x^2 - 3\pi x + 2\pi^2}{x - \pi + 2 \sin x} \quad \left[ \left( \frac{0}{0} \right) \text{form} \right]$$

[Apply L - Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form}$ ]  
 $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

$$P = \lim_{x \rightarrow \pi} \frac{\frac{d}{dx}(x^2 - 3\pi x + 2\pi^2)}{\frac{d}{dx}(x - \pi + 2 \sin x)}$$

$$P = \lim_{x \rightarrow \pi} \frac{2x - 3\pi}{1 + 2 \cos x} = \frac{-\pi}{-1} = \pi$$

$$\therefore \alpha = -3\pi \text{ and } P = \pi$$

Scan for Video solution

## 3. (-0.5)

Given limit is

$$P = \lim_{x \rightarrow 0} + \frac{\sqrt{x}}{1 - e^{2\sqrt{x}}} = \frac{\sqrt{0}}{1 - e^{2(\sqrt{0})}} = \frac{0}{1 - 1} = \left( \begin{array}{c} 0 \\ 0 \end{array} \right)$$

$\left[ \text{Apply L-Hospital rule, For } \left( \begin{array}{c} 0 \\ 0 \end{array} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \right]$

$$\left[ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \right]$$

$$P = \lim_{x \rightarrow 0^+} \frac{\frac{d}{dx}(\sqrt{x})}{\frac{d}{dx}(1 - e^{2\sqrt{x}})} = \lim_{x \rightarrow 0} + \frac{\frac{1}{2\sqrt{x}}}{0 - e^{2\sqrt{x}} \cdot 2 \cdot \frac{1}{2\sqrt{x}}} = \frac{1}{0 - 1} = -1$$

$$P = \frac{1}{-2} = -0.5$$

$$\therefore P = \lim_{x \rightarrow 0} + \frac{\sqrt{x}}{1 - e^{2\sqrt{x}}} = -0.5$$



Scan for Video solution



## 4. (0.25)

$$\text{Given limit is } P = \lim_{x \rightarrow -3} \frac{\sqrt{2x + 22} - 4}{x + 3} = \left( \begin{array}{c} 0 \\ 0 \end{array} \right)$$

$\left[ \text{Apply L-Hospital rule, For } \left( \begin{array}{c} 0 \\ 0 \end{array} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \right]$

$$\left[ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \right]$$

$$P = \lim_{x \rightarrow -3} \frac{\frac{d}{dx}\sqrt{2x + 22} - 4}{\frac{d}{dx}(x + 3)} = \lim_{x \rightarrow -3} \frac{\frac{1}{\sqrt{2x + 22}} \cdot 2 - 4}{1} = \frac{1}{\sqrt{2(-3) + 22}} \cdot 2 - 4 = \frac{1}{\sqrt{16}} \cdot 2 - 4 = 2 - 4 = -2$$

$$P = \lim_{x \rightarrow -3} \frac{1}{\sqrt{2x + 22}}$$

$$P = \frac{1}{\sqrt{16}} = \frac{1}{4} = 0.25$$

$$\therefore \lim_{x \rightarrow -3} \frac{\sqrt{2x + 22} - 4}{x + 3} = 0.25$$

Scan for Video solution



## 5. (c)

$$\text{Given } P = \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$$

$$P = \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1 - 1}{0} = \left( \begin{array}{c} 0 \\ 0 \end{array} \right)$$

$$\left[ \left( \begin{array}{c} 0 \\ 0 \end{array} \right) \text{ form} \right]$$

$\left[ \text{Apply L-Hospital rule, For } \left( \begin{array}{c} 0 \\ 0 \end{array} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \right]$

$$\left[ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \right]$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(1 - \cos x)}{\frac{d}{dx}(x^2)} = \lim_{x \rightarrow 0} \frac{\sin x}{2x}$$

$$P = \frac{1}{2} \times \lim_{x \rightarrow 0} \frac{\sin x}{x}$$

$$\left[ \because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right]$$

$$P = \frac{1}{2} \times 1 = \frac{1}{2}$$

$$\therefore \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2}$$

Scan for Video solution



## 6. (d)

$$P = \lim_{x \rightarrow \infty} \frac{x \ln x}{1 + x^2} \quad \left( \frac{\infty}{\infty} \text{ form} \right)$$

Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form  
 $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

$$P = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(x \ln x)}{\frac{d}{dx}(1+x^2)} = \lim_{x \rightarrow \infty} \frac{x \cdot \frac{1}{x} + \ln x \cdot 1}{2x}$$

$$P = \lim_{x \rightarrow \infty} \frac{1 + \ln x}{2x}$$

$\left( \frac{\infty}{\infty} \text{ form, apply L Hospital's Rule} \right)$

$$P = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(1 + \ln x)}{\frac{d}{dx}(2x)} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{2} = \frac{0}{2} = 0$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \frac{x \ln x}{1+x^2} = 0}$$



Scan for Video solution



### 7. (0.5)

$$P = \lim_{x \rightarrow 1} \left( \frac{1}{\ln x} - \frac{1}{x-1} \right)$$

$$P = \lim_{x \rightarrow 1} \frac{(x-1) - \ln x}{(x-1)\ln x} \quad (0/0 \text{ form})$$

Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form  
 $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}((x-1) - \ln x)}{\frac{d}{dx}((x-1)\ln x)}$$

$$P = \lim_{x \rightarrow 1} \frac{1 - \frac{1}{x}}{(x-1)\frac{1}{x} + \ln x} = \left( \frac{0}{0} \right) \text{ form}$$

Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form  
 $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

$$P = \lim_{x \rightarrow 1} \frac{\frac{d}{dx}\left(1 - \frac{1}{x}\right)}{\frac{d}{dx}\left(\left(x-1\right)\frac{1}{x} + \ln x\right)} = \lim_{x \rightarrow 1} \frac{\frac{1}{x^2}}{\frac{1}{x^2} + \frac{1}{x}}$$

$$P = \lim_{x \rightarrow 1} \frac{-\left(-\frac{1}{x^2}\right)}{-\left(-\frac{1}{x^2}\right) + \frac{1}{x}} = \frac{1}{1+1} = \frac{1}{2} = 0.5$$

$$\therefore \boxed{\lim_{x \rightarrow 1} \left( \frac{1}{\ln x} - \frac{1}{x-1} \right) = 0.5}$$

Scan for Video solution



### 8. (b)

$$P = \lim_{x \rightarrow \infty} \frac{\sqrt{9x^2 + 2020}}{x+7} = \lim_{x \rightarrow \infty} \frac{3x \sqrt{1 + \frac{2020}{9x^2}}}{x \left(1 + \frac{7}{x}\right)}$$

$$P = \lim_{x \rightarrow \infty} \frac{3 \sqrt{1 + \frac{2020}{9x^2}}}{\left(1 + \frac{7}{x}\right)} = \lim_{x \rightarrow \infty} \frac{3\sqrt{1+0}}{(1+0)} = 3$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \frac{\sqrt{9x^2 + 2020}}{x+7} = 3}$$

Scan for Video solution



### 9. (d)

True value of  $\ln 2 = 0.69$

$$x_1 = 1$$

$$y_1 = \ln 1$$

$$x_2 = 6 \quad y_2 = \ln 6$$


$1, \ln 1$

$(2, y)$

$6, \ln 6$

$$\frac{y - \ln 1}{2 - 1} = \frac{\ln 6.0 - \ln 1}{6 - 1}$$

$$y = 0.358 \text{ (Approximate value of } \ln 2\text{)}$$



Hence % error =

$$\frac{\text{True value} - \text{Approximate value}}{\text{True value}} \times 100\%$$

$$= \frac{0.69 - 0.358}{0.69} \times 100 = 48.11\%$$

$$= 48\%$$



Scan for Video solution

**10. (c)****Method I:**

$$P = \lim_{x \rightarrow \infty} \frac{x^2 - 5x + 4}{4x^2 + 2x} = \left( \frac{\infty}{\infty} \text{ form} \right)$$

$$\left[ \begin{array}{l} \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \\ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \end{array} \right]$$

$$P = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(x^2 - 5x + 4)}{\frac{d}{dx}(4x^2 + 2x)} = \lim_{x \rightarrow \infty} \frac{2x - 5}{8x + 2} = \left( \frac{\infty}{\infty} \text{ form} \right)$$

$$\left[ \begin{array}{l} \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \\ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \end{array} \right] P$$

$$P = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(2x - 5)}{\frac{d}{dx}(8x + 2)} = \lim_{x \rightarrow \infty} \frac{2}{8} = \frac{2}{8} = \frac{1}{4}$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \frac{x^2 - 5x + 4}{4x^2 + 2x} = \frac{1}{4}}$$


**Method II:**

$$P = \lim_{x \rightarrow \infty} \frac{x^2 - 5x + 4}{4x^2 + 2x}$$

$$P = \lim_{x \rightarrow \infty} \frac{x^2 \left( 1 - \frac{5}{x} + \frac{4}{x^2} \right)}{x^2 \left( 4 + \frac{2}{x} \right)} = \lim_{x \rightarrow \infty} \frac{\left( 1 - \frac{5}{x} + \frac{4}{x^2} \right)}{\left( 4 + \frac{2}{x} \right)}$$

$$P = \frac{(1 - 0 + 0)}{(4 + 0)} = \frac{1}{4}$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \frac{x^2 - 5x + 4}{4x^2 + 2x} = \frac{1}{4}}$$

**11. (c)**

$$\text{Given } P = \lim_{x \rightarrow 1} \left( \frac{1 - e^{-c(1-x)}}{1 - xe^{-c(1-x)}} \right)$$

$$P = \lim_{x \rightarrow 1} \frac{1 - e^{-c(1-x)}}{1 - x \cdot e^{-c(1-x)}} = \frac{1 - e^{-0}}{1 - 1 \cdot e^{-0}} = \left( \frac{0}{0} \right)$$

$$\left[ \left( \frac{0}{0} \right) \text{ form} \right]$$

**Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form**

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow 1} \frac{\frac{d}{dx}(1 - e^{-c(1-x)})}{\frac{d}{dx}(1 - x \cdot e^{-c(1-x)})}$$

$$P = \lim_{x \rightarrow 1} \frac{0 - e^{-c(1-x)} \cdot c}{0 - \{x \cdot c \cdot e^{-c(1-x)} + e^{-c(1-x)} \cdot 1\}}$$

$$P = \lim_{x \rightarrow 1} \frac{-c \cdot e^{-c(1-x)}}{-e^{-c(1-x)} \{1 + cx\}}$$

$$P = \lim_{x \rightarrow 1} \frac{c}{1 + cx}$$

$$\lim_{x \rightarrow 1} \frac{1 - e^{-c(1-x)}}{1 - x \cdot e^{-c(1-x)}} = \frac{c}{1 + c}$$



Scan for Video solution

**12. (a)**

$$\text{Given; } x = \psi \ln \phi \quad \dots \text{(i)}$$

$$\text{and } y = \phi \ln \psi \quad \dots \text{(ii)}$$

Partially differentially equation (i) w.r.t. x

$$1 = \psi_x \ln \phi + \psi \cdot \frac{\phi_x}{\phi} \quad \dots \text{(iii)}$$

Partially differentially equation (ii) w.r.t. x

$$0 = \phi_x \ln \psi + \phi \frac{1}{\psi} \cdot \psi_x$$

$$\frac{\psi \phi_x}{\phi} = - \frac{\psi_x}{\ln \psi} \quad \dots \text{(iv)}$$

Put this term in (iii)

$$1 = \psi_x \ln \phi - \frac{\psi_x}{\ln \psi}$$

$$1 = \psi_x \left[ \ln \phi - \frac{1}{\ln \psi} \right]$$

$$1 = \psi_x \left[ \frac{\ln \phi \ln \psi - 1}{\ln \psi} \right]$$

$$\psi_x = \frac{\ln \psi}{\ln \phi \ln \psi - 1}$$

$$\frac{\partial \psi}{\partial x} = \frac{\ln \psi}{\ln \phi \ln \psi - 1}$$

Scan for Video solution

**13. (c)**

Given limits is

$$P = \lim_{x \rightarrow -3} \frac{x^4 - 81}{2x^2 - 5x - 3} = \frac{81 - 81}{18 - 15 - 3} = \left( \frac{0}{0} \right)$$

**Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form**

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow -3} \frac{\frac{d}{dx}(x^4 - 81)}{\frac{d}{dx}(2x^2 - 5x - 3)}$$

$$P = \lim_{x \rightarrow -3} \frac{4x^3}{4x - 5} = \frac{4(3)^3}{4(3) - 5} = \frac{4 \times 27}{7} = \frac{108}{7}$$

$$\therefore \lim_{x \rightarrow -3} \frac{x^4 - 81}{2x^2 - 5x - 3} = \frac{108}{7}$$

Scan for Video solution

**14. (d)**

**Method I:**

$$P = \lim_{x \rightarrow 0} \frac{x \sin x}{1 - \cos x}$$

$$P = \lim_{x \rightarrow 0} \frac{\sin x}{1 - \cos x}$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{x}{\sin x}}{\frac{1 - \cos x}{x^2}}$$

$$\left[ \because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \text{ and } \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2} \right]$$

$$P = \frac{1}{1/2} = 2$$

**Method II:**

$$P = \lim_{x \rightarrow 0} \frac{x \sin x}{1 - \cos x} = \left( \frac{0}{0} \right) \text{ form}$$

$$\left[ \begin{array}{l} \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \\ \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{f'(x)}{g'(x)} \end{array} \right]$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(x \sin x)}{\frac{d}{dx}(1 - \cos x)} = \lim_{x \rightarrow 0} \frac{x \cos x + \sin x}{\sin x} = \left( \frac{0}{0} \right) \text{ form}$$

$$\left[ \begin{array}{l} \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \\ \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{f'(x)}{g'(x)} \end{array} \right]$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(x \cos x + \sin x)}{\frac{d}{dx}(\sin x)}$$

$$P = \lim_{x \rightarrow 0} \frac{-x \sin x + \cos x + \cos x}{\cos x} = \frac{0+1+1}{1} = 2$$

$$\therefore \boxed{\lim_{x \rightarrow 0} \frac{x \sin x}{1 - \cos x} = 2}$$



Scan for Video solution



15. (d)

$$\text{Let, } P_1 = \lim_{x \rightarrow 0} \frac{\sin 4x}{\sin 2x}$$

**Method I:**

$$P_1 = \lim_{x \rightarrow 0} \frac{\sin 4x}{\sin 2x} = \left( \frac{0}{0} \right) \text{ Form}$$

$$\left[ \begin{array}{l} \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \\ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \end{array} \right]$$

$$P_1 = \lim_{x \rightarrow 0} \frac{\frac{d}{dx} \sin 4x}{\frac{d}{dx} \sin 2x} = \frac{4 \cos 4x}{2 \cos 2x}$$

$$P_1 = \frac{4}{2} = 2$$

**Method II:**

$$P_1 = \lim_{x \rightarrow 0} \frac{\sin 4x}{\sin 2x}$$

$$P_1 = \lim_{x \rightarrow 0} \frac{4x \cdot \left( \frac{\sin 4x}{4x} \right)}{2x \cdot \left( \frac{\sin 2x}{2x} \right)} \quad \left[ \because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right]$$

$$P_1 = 2 \cdot \frac{1}{1} = 2$$

$$\text{Let, } P_2 = \lim_{x \rightarrow 0} \frac{\tan x}{x}$$

$$P_2 = \lim_{x \rightarrow 0} \frac{\tan x}{x} = \left( \frac{0}{0} \right) \text{ form}$$

$$\left[ \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \right]$$


$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P_2 = \lim_{x \rightarrow 0} \frac{\frac{d}{dx} \tan x}{\frac{d}{dx} x} = \lim_{x \rightarrow 0} \frac{\sec^2 x}{1}$$

$$P_2 = \lim_{x \rightarrow 0} \frac{\sec^2 0}{1} = 1$$

$$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1 \leftarrow (\text{This expression we can remember also})$$

$$\therefore \boxed{\lim_{x \rightarrow 0} \frac{\sin 4x}{\sin 2x} = 2 \text{ & } \lim_{x \rightarrow 0} \frac{\tan x}{x} = 1}$$



## 16. (b)

**Given**

$$f(x) = \cos x \quad \dots(i)$$

$$\text{Approximate value of } f'(x) = \frac{f(x+h) - f(x-h)}{2h}$$

At  $x = \frac{\pi}{6}$  and  $h = 0.1$ :

$$\begin{bmatrix} \cos\left(\frac{\pi}{6} + 0.1\right) \\ -\cos\left(\frac{\pi}{6} - 0.1\right) \end{bmatrix}$$

$$\text{Approximate value of } f'(x) = \frac{\cos\left(\frac{\pi}{6} + 0.1\right) - \cos\left(\frac{\pi}{6} - 0.1\right)}{2(0.1)}$$

$$\text{Approximate value of } f'(x) = \frac{0.81178 - 0.9116}{0.2}$$

$$\text{Approximate value of } f'(x) = -0.49916$$

**Exact value of  $f'(x)$ :**Differentiating equation (i) with respect to  $x$ ,

$$\text{Exact value of } f'(x) = -\sin x$$

$$\text{At } x = \frac{\pi}{6} :$$

$$\text{Exact value of } f'(x) = -\sin \frac{\pi}{6}$$

$$\text{Exact value of } f'(x) = -0.5$$

$$\therefore \% \text{ error} = \left( \frac{-0.5 + 0.49916}{-0.5} \right) \times 100 = 0.166\%$$



Scan for Video solution



## 17. (d)

$$f(x) = \begin{cases} x^2 & ; x \geq 0 \\ -x^2 & ; x < 0 \end{cases}$$

Now,

**Left hand limit (LHL):**

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (x^2) = 0$$

**Right hand limit (RHL):**

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (-x^2) = 0$$

$$f(0) = 0^2 = 0$$

**Since**  $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$ , therefore

 **$f(x)$  is continuous at  $x = 0$ .**

Now,

$$f'(x) = \begin{cases} 2x & ; x \geq 0 \\ -2x & ; x < 0 \end{cases}$$

$$\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^-} (-2x) = 0$$

$$\lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0^+} (2x) = 0$$

$$f'(0) = 2 \times 0 = 0$$

**Since**  $\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^+} f'(x) = f'(0)$ , therefore

 **$f'(x)$  is continuous at  $x = 0$  i.e.  $f(x)$  is differentiable at  $x = 0$ .**


Now,

$$f''(x) = \begin{cases} 2 & ; x \geq 0 \\ -2 & ; x < 0 \end{cases}$$

$$\lim_{x \rightarrow 0^-} f''(x) = \lim_{x \rightarrow 0^-} (-2) = -2$$

$$\lim_{x \rightarrow 0^+} f''(x) = \lim_{x \rightarrow 0^+} (2) = 2$$

**Since**  $\lim_{x \rightarrow 0^-} f''(x) \neq \lim_{x \rightarrow 0^+} f''(x)$ , therefore  $f''(x)$  is not continuous at  $x = 0$  i.e.  $f'(x)$  is not differentiable at  $x = 0$ .



## 18. (a)

**Given Data:**

$$g(x) = \begin{cases} -x & ; x \leq 1 \\ x + 1 & ; x \geq 1 \end{cases} \quad \dots(i)$$

$$f(x) = \begin{cases} 1 - x & ; x \leq 0 \\ x^2 & ; x > 0 \end{cases} \quad \dots(ii)$$

$$(fog)(x) = f(g(x))$$

From equation (ii),

$$(fog)(x) = \begin{cases} 1 - g(x) & ; g(x) \leq 0 \\ [g(x)]^2 & ; g(x) > 0 \end{cases}$$

From equation (i).

$$(fog)(x) = \begin{cases} 1 - (-x) & ; -x \leq 0 \& x \leq 1 \\ 1 - (x+1) & ; x+1 \leq 0 \& x \geq 1 \\ (-x)^2 & ; -x > 0 \& x \leq 1 \\ (x+1)^2 & ; x+1 > 0 \& x \geq 1 \end{cases}$$

$$(fog)(x) = \begin{cases} 1 - (-x) & ; x \geq 0 \& x \leq 1 \Rightarrow 0 \leq x \leq 1 \\ 1 - (x+1) & ; x \leq -1 \& x \geq 1 \Rightarrow \text{No interval} \\ (-x)^2 & ; x < 0 \& x \leq 1 \Rightarrow x < 0 \\ (x+1)^2 & ; x > -1 \& x \geq 1 \Rightarrow x \geq 1 \end{cases}$$

$$(fog)(x) = \begin{cases} 1+x & ; 0 \leq x \leq 1 \\ x^2 & ; x < 0 \\ (x+1)^2 & ; x \geq 1 \end{cases}$$

In interval  $(-\infty, 0)$  i.e.  $x < 0$ , the  $(fog)(x)$  is  $x^2$  which is a continuous function. Therefore, number of discontinuities in interval  $(-\infty, 0)$  is 0.

**Note:**

If examiner is asking discontinuous point in interval  $(-\infty, \infty)$  for  $(fog)(x)$ , then we need to check continuity only at  $x = 0$  and  $x = 1$ .

**Checking continuity at  $x = 0$ :**

Left hand limit (LHL):

$$\lim_{x \rightarrow 0^-} (fog)(x) = \lim_{x \rightarrow 0^-} (x^2) = 0$$

Right hand limit (RHL):

$$\lim_{x \rightarrow 0^+} (fog)(x) = \lim_{x \rightarrow 0^+} (1+x) = 1$$

$\because LHL \neq RHL$ , therefore  $(fog)(x)$  is not continuous at  $x = 0$ .

**Checking continuity at  $x = 1$ :**

Left hand limit (LHL):

$$\lim_{x \rightarrow 1^-} (fog)(x) = \lim_{x \rightarrow 1^-} (1+x) = 2$$

Right hand limit (RHL):

$$\lim_{x \rightarrow 1^+} (fog)(x) = \lim_{x \rightarrow 1^+} ((x+1)^2) = 4$$

$\because LHL \neq RHL$ , therefore  $(fog)(x)$  is not continuous at  $x = 1$ .

Hence in the interval  $(-\infty, 0)$  there is no discontinuity but if we consider overall interval  $(-$

$\infty, \infty)$  then  $f[g(x)]$  is discontinuous at  $x = 0$  and  $x = 1$ .


Scan for Video solution


**19. (b)**

$$f(x) = \begin{cases} e^x & ; x < 1 \\ \ln x + ax^2 + bx & ; x \geq 1 \end{cases}$$

For  $f(x)$  to be differential, first  $f(x)$  should be continuous at  $x = 1$ . And for that,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1)$$

$$\lim_{x \rightarrow 1^-} (e^x) = \lim_{x \rightarrow 1^+} (\ln x + ax^2 + bx)$$

$$\Rightarrow e = (\ln 1 + a + b)$$

$$\Rightarrow e = a + b \quad \dots(i)$$

Now,

$$f'(x) = \begin{cases} e^x & ; x < 1 \\ \frac{1}{x} + 2ax + b & ; x \geq 1 \end{cases}$$

For  $f(x)$  to be differentiable

$$f'(1^-) = f'(1^+)$$

$$\Rightarrow e = 1 + 2a(1) + b(1) \quad \dots(ii)$$

From (i) and (ii)

$$a + b = 1 + 2a + b$$

$$\Rightarrow a = -1$$

Put value of  $a$  in equation (i),

$$b = 1 + e$$

**Hence,  $f(x)$  is differentiable at  $x = 1$  for the unique value of  $a$  and  $b$ .**


Scan for Video solution


**20. (c)**

Given limit is

$$P = \lim_{x \rightarrow 1} \frac{x^7 - 2x^5 + 1}{x^3 - 3x^2 + 2} = \frac{1 - 2 + 1}{1 - 3 + 2} = \left( \begin{array}{l} 0 \\ 0 \end{array} \right)$$

Apply L – Hospital rule, For  $\left( \begin{array}{l} 0 \\ 0 \end{array} \text{ (or) } \frac{\infty}{\infty} \right)$  form

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow 1} \frac{\frac{d}{dx}(x^7 - 2x^5 + 1)}{\frac{d}{dx}(x^3 - 3x^2 + 2)} = \lim_{x \rightarrow 1} \frac{7x^6 - 10x^4}{3x^2 - 6x}$$

$$P = \frac{7 - 10}{3 - 6} = \frac{-3}{-3} = 1$$

$$\therefore \lim_{x \rightarrow 1} \frac{x^7 - 2x^5 + 1}{x^3 - 3x^2 + 2} = 1$$



Scan for Video solution

**21. (-1 to -1)**

$$P = \lim_{x \rightarrow 0} \frac{\tan x}{x^2 - x} = \left( \begin{array}{l} 0 \\ 0 \end{array} \right) \quad \left[ \left( \begin{array}{l} 0 \\ 0 \end{array} \right) \text{ form} \right]$$

Apply L – Hospital rule, For  $\left( \begin{array}{l} 0 \\ 0 \end{array} \text{ (or) } \frac{\infty}{\infty} \right)$  form

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(\tan x)}{\frac{d}{dx}(x^2 - x)} = \lim_{x \rightarrow 0} \frac{\sec^2 x}{2x - 1}$$

$$P = \frac{\sec^2 0}{2(0) - 1} = \frac{1}{-1} = -1$$

$$\therefore \lim_{x \rightarrow 0} \frac{\tan x}{x^2 - x} = -1$$




Scan for Video solution

**22. (c)**

Given,  $w = f(x, y)$


By Chain rule of differentiation,

$$\frac{dw}{dt} = \frac{\partial w}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial w}{\partial y} \cdot \frac{dy}{dt}$$

**23. (d)**

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{x^3 - \sin x}{x} &= \lim_{x \rightarrow 0} x^2 - \frac{\sin x}{x} \\ &= \lim_{x \rightarrow 0} x^2 - \lim_{x \rightarrow 0} \frac{\sin x}{x} \\ &= 0 - 1 = -1 \end{aligned}$$

$$\therefore \lim_{x \rightarrow 0} \frac{x^3 - \sin x}{x} = -1$$

**24. (54 to 55)**

Given planes are  $x + y + z - 1 = 0$

and  $2x - y + 2z = 0$

The angle between the planes  $a_1x + b_1y + c_1z + d_1 = 0$  and  $a_2x + b_2y + c_2z + d_2 = 0$  is given by


$$\cos \theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \cdot \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

$$\Rightarrow \cos \theta = \frac{1(2) + 1(-1) + 1(2)}{\sqrt{1^2 + 1^2 + 1^2} \cdot \sqrt{2^2 + (-1)^2 + 2^2}}$$

$$\Rightarrow \cos \theta = \frac{3}{\sqrt{3} \times 3}$$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \theta = 54.73^\circ$$



**25. (b)**

(i) If  $f(x)$  is continuous at  $x = x_0$ , then it may not be differentiable at  $x_0$ , ex:  $|x|$  at  $x = 0$

(ii) If  $f(x)$  is differentiable at  $x = x_0$ , then it is definitely continuous at  $x = x_0$ .



Scan for Video solution

**26. (1)**

$$\lim_{x \rightarrow 4} \frac{\sin(x-4)}{(x-4)}$$

is the given limit

As  $x \rightarrow 4 \Rightarrow (x-4) \rightarrow 0$

Let  $x-4=t$

$$\Rightarrow \text{Given limits is } \lim_{t \rightarrow 0} \frac{\sin t}{t} = 1$$

$$\therefore \lim_{t \rightarrow 0} \frac{\sin(t)}{(t)} = 1$$



Scan for Video solution

**27. (c)**

$$\lim_{x \rightarrow 0} \frac{\log_e(1+4x)}{e^{3x}-1} = \frac{0}{0}$$

$$\left[ \begin{array}{l} \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \\ \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{f'(x)}{g'(x)} \end{array} \right]$$

$$\lim_{x \rightarrow 0} \frac{\frac{1}{1+4x} \times 4}{3 \cdot e^{3x}} = \frac{4 \times (1)}{3 \cdot e^0} = \frac{4}{3}$$

$$\therefore \lim_{x \rightarrow 0} \frac{\log_e(1+4x)}{e^{3x}-1} = \frac{4}{3}$$



Scan for Video solution

**28. (c)**

$$\lim_{x \rightarrow \infty} \sqrt{x^2 + x - 1} - x = \lim_{x \rightarrow \infty} \left( \sqrt{x^2 + x - 1} - x \right) \frac{\left( \sqrt{x^2 + x - 1} + x \right)}{\left( \sqrt{x^2 + x - 1} + x \right)}$$

$$= \lim_{x \rightarrow \infty} \frac{x^2 + x - 1 - x^2}{\sqrt{x^2 + x - 1} + x} = \lim_{x \rightarrow \infty} \frac{x \left( 1 - \frac{1}{x} \right)}{x \left( \sqrt{1 + \frac{1}{x} - \left( \frac{1}{x} \right)^2} + 1 \right)}$$

$$= \lim_{\frac{1}{x} \rightarrow 0} \frac{\left( 1 - \frac{1}{x} \right)}{\sqrt{1 + \left( \frac{1}{x} \right) - \left( \frac{1}{x} \right)^2} + 1}$$

$$\text{As } x \rightarrow \infty; \frac{1}{x} \rightarrow 0$$

$$= \frac{1}{1+1} = \frac{1}{2} = 0.5$$

$$\therefore \lim_{x \rightarrow \infty} \left( \sqrt{x^2 + x - 1} - x \right) = \frac{1}{2}$$



Scan for Video solution

**29. (c)**

The function  $f(x) = \frac{\phi(x)}{\psi(x)}$  where  $\phi(x)$  and  $\psi(x)$  are polynomial is discontinuous at the points where  $\psi(x) = 0$ .

Hence, given function is not continuous at the points Where,  $x^2 + 3x - 4 = 0$

$$\Rightarrow (x+4)(x-1) = 0$$

$$\Rightarrow x = -4, 1$$

**$\therefore f(x)$  is NOT continuous at  $x = -4, 1$**



Scan for Video solution

**30. (0.28 to 0.31)**

$$S = \sum_{n=0}^{\infty} n \alpha^n$$

$$\Rightarrow S = 0 + \alpha + 2\alpha^2 + 3\alpha^3 + \dots \quad \dots(1)$$

$$\Rightarrow \alpha S = 0 + \alpha^2 + 2\alpha^3 + 3\alpha^4 + \dots \quad \dots(2)$$

Equation (1) - (2)

$$\Rightarrow S(1 - \alpha) = \alpha + \alpha^2 + \alpha^3 + \dots \quad \dots(3)$$

Here,

$\alpha + \alpha^2 + \alpha^3 + \dots$  is infinite G.P. series with common ratio  $r = \alpha$

Now, sum of infinite G.P. series =  $\frac{a}{1-r}$  [If  $r < 1$ ]

Therefore,

$$\alpha + \alpha^2 + \alpha^3 + \dots = \frac{\alpha}{1-\alpha}$$

From equation (3),

$$S(1 - \alpha) = \frac{\alpha}{1 - \alpha}$$

$$\Rightarrow S = \frac{\alpha}{(1 - \alpha)^2} \quad [\because S = 2\alpha]$$

$$\Rightarrow 2\alpha = \frac{\alpha}{(1 - \alpha)^2}$$

$$\Rightarrow (1 - \alpha)^2 = \frac{1}{2}$$

$$\Rightarrow 1 - \alpha = \pm \frac{1}{\sqrt{2}}$$

$$\Rightarrow \alpha = 0.2928 \text{ and } \alpha = 1.7071$$

Discarding  $\alpha = 1.7071$  because  $0 < \alpha < 1$

$$\Rightarrow \therefore \boxed{\alpha = 0.2928}$$



Scan for Video solution



If  $f(x)$  does not have a root in the interval  $[a, b]$  then it implies that  $f(x)$  is not changing its sign in interval  $[a, b]$  i.e. both  $f(a)$  and  $f(b)$  are either positive or negative.

$$\therefore f(a) \cdot f(b) > 0$$

▶
Scan for Video solution

32. (d)

$$P = \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{2x}$$

$$P = \lim_{x \rightarrow \infty} \left[ \left(1 + \frac{1}{x}\right)^x \right]^2$$

$$\left[ \because \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e \right]$$

$$P = e^2$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{2x} = e^2}$$

▶
Scan for Video solution

33. (c)

Given:


$$P = \lim_{x \rightarrow 0} \frac{1 - \cos(x^2)}{2x^4}$$

$$P = \frac{1-1}{4} = \left(\frac{0}{0}\right)$$

$$\left[\left(\frac{0}{0}\right) \text{ form}\right]$$

Apply L - Hospital rule, For  $\left(\frac{0}{0} \text{ (or) } \frac{\infty}{\infty}\right) \text{ form}$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$



31. (c)

According to intermediate value theorem,

If  $f(x)$  is continuous in  $[a, b]$  and changing its sign between  $a$  and  $b$  (i.e.  $f(a)$  and  $f(b)$  have opposite signs) then there exists at least one root of  $f(x)$  in  $[a, b]$  as shown in figure.

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(1 - \cos(x^2))}{\frac{d}{dx}(2x^4)} = \lim_{x \rightarrow 0} \frac{\sin(x^2) \times 2x}{8x^3}$$

$$P = \frac{1}{4} \times \lim_{x \rightarrow 0} \frac{\sin(x^2)}{x^2}$$

As  $x \rightarrow 0$ ,  $x^2 \rightarrow 0$  and let  $x^2 = t$ , therefore,

$$P = \frac{1}{4} \times \lim_{t \rightarrow 0} \frac{\sin(t)}{t} \quad \left[ \because \lim_{t \rightarrow 0} \frac{\sin t}{t} = 1 \right]$$

$$P = \frac{1}{4} \times 1 = \frac{1}{4}$$

Hence,  $\boxed{\lim_{x \rightarrow 0} \frac{1 - \cos(x^2)}{2x^4} = \frac{1}{4}}$



Scan for Video solution



### 34. (-0.35 to -0.30)

Given:

$$P = \lim_{x \rightarrow 0} \left( \frac{-\sin x}{2 \sin x + x \cos x} \right)$$

$$P = \lim_{x \rightarrow 0} \left( \frac{-\left(\frac{\sin x}{x}\right)}{2\left(\frac{\sin x}{x}\right) + \left(\frac{x \cdot \cos x}{x}\right)} \right)$$

$$P = \frac{-\lim_{x \rightarrow 0} \frac{\sin x}{x}}{2 \cdot \lim_{x \rightarrow 0} \frac{\sin x}{x} + \lim_{x \rightarrow 0} x \cdot \cos x}$$

$$\left[ \because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right]$$

$$P = \frac{-1}{2+1} = \frac{-1}{3} = -0.33$$

$\therefore \boxed{\lim_{x \rightarrow 0} \frac{-\sin x}{\sin x + x \cdot \cos x} = -0.33}$



Scan for Video solution




### 35. (1.9 to 2.1)

Given

$$\text{curve } (x - 2)^2 + y^2 = 4$$

Starting point = (4, 0) and final point = (2, 2)

$$\text{Speed of the ant} = 1.57 \text{ m/s} = \frac{\pi}{2} \text{ m/s}$$



$$\text{Time required for travel} = \frac{\text{length of the arc AB}}{\text{speed of ant}}$$

$$= \frac{\frac{1}{4} \times 2\pi r}{\left(\frac{\pi}{2}\right)} = 2 \text{ sec}$$

$\therefore$  Time for travelling from A to B = 2 sec



Scan for Video solution



### 36. (c)

Given limit is  $\lim_{x \rightarrow \infty} x^{\frac{1}{x}} = \infty^\circ$  (undetermined form)

$$\text{Let } y = x^{\frac{1}{x}}$$

$$\Rightarrow \log_e y = \frac{1}{x} \log_e x$$

$$\Rightarrow \lim_{x \rightarrow \infty} (\log_e y) = \lim_{x \rightarrow \infty} \frac{\log_e x}{x} = \left( \frac{\infty}{\infty} \right)$$

Apply L-Hospital Rule

Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$\Rightarrow \log_e \left( \lim_{x \rightarrow \infty} y \right) = \lim_{x \rightarrow \infty} \frac{d}{dx} \log_e x = \lim_{x \rightarrow \infty} \frac{1}{x} = 0$$

$$\Rightarrow \log_e \left( \lim_{x \rightarrow \infty} y \right) = 0 \quad \Rightarrow \lim_{x \rightarrow \infty} y = e^0 = 1$$

$$\Rightarrow \boxed{\lim_{x \rightarrow \infty} x^x = 1}$$



Scan for Video solution

**37. (a)**

Given limit is

$$P = \lim_{x \rightarrow \infty} (1+x^2) \cdot e^{-x} = \lim_{x \rightarrow \infty} \frac{(1+x^2)}{e^x} = \left( \frac{\infty}{\infty} \right)$$

[ Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form ]  
 $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

$$\Rightarrow P = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(1+x^2)}{\frac{d}{dx}e^x} = \lim_{x \rightarrow \infty} \frac{2x}{e^x} = \left( \frac{\infty}{\infty} \right)$$

[ Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right)$  form ]  
 $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

$$\Rightarrow P = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx}(2x)}{\frac{d}{dx}e^x} = \lim_{x \rightarrow \infty} \frac{2}{e^x} = \frac{2}{e^0} = 0$$

$$\therefore \boxed{P = \lim_{x \rightarrow \infty} \frac{1+x^2}{e^x} = 0}$$



Scan for Video solution

**38. (c)**

$$\text{Let } P = \lim_{x \rightarrow \infty} \left( 1 + \frac{1}{x} \right)^x = 1^\infty$$

$$\text{For } \lim_{x \rightarrow a} (f(x))^{g(x)} = 1^\infty,$$

$$\lim_{x \rightarrow a} (f(x))^{g(x)} = e^{\lim_{x \rightarrow a} g(x) \cdot \{f(x)-1\}}$$

Therefore,

$$P = \lim_{x \rightarrow \infty} \left( 1 + \frac{1}{x} \right)^x = e^{\lim_{x \rightarrow \infty} x \left( 1 + \frac{1}{x} - 1 \right)}$$

$$P = e^{\lim_{x \rightarrow \infty} 1} = e^1 = e$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \left( 1 + \frac{1}{x} \right)^x = e}$$



Scan for Video solution

**39. (-2)**

$$\text{Given } f(x) = x \cdot \sin x$$

$$\Rightarrow f'(x) = x \cdot \cos x + \sin x$$

And,

$$\Rightarrow f''(x) = x \cdot (-\sin x) + \cos x + \cos x$$

$$\Rightarrow f''(x) = -x \cdot \sin x + 2\cos x$$

$$\text{Considering } f''(x) + f(x) + t \cdot \cos x = 0$$

$$\Rightarrow -x \cdot \sin x + 2\cos x + x \cdot \sin x + t \cdot \cos x = 0$$

$$\Rightarrow (2+t) \cos x = 0$$

$$\Rightarrow (2+t) = 0$$

$$\Rightarrow \boxed{t = -2}$$



Scan for Video solution



40. (b)

$$P = \lim_{x \rightarrow \infty} \left( \frac{x + \sin x}{x} \right)$$

$$P = \lim_{x \rightarrow \infty} \left( 1 + \frac{\sin x}{x} \right)$$

As  $x \rightarrow \infty$ ; the value of  $\sin x$  oscillated between +1 & -1 which is a finite value.

$$P = \lim_{x \rightarrow \infty} \left( 1 + \frac{\text{Finite value}}{\infty} \right)$$

$$P = 1 + 0 = 1$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \left( \frac{x + \sin x}{x} \right) = 1}$$



Scan for Video solution



41. (a)

$$P = \lim_{\alpha \rightarrow 0} \frac{x^\alpha - 1}{a} = \left( \frac{0}{0} \right)$$

$$\left[ \left( \frac{0}{0} \right) \text{form} \right]$$

Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form}$

$$\left[ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \right]$$

$$P = \lim_{\alpha \rightarrow 0} \frac{\frac{d}{d\alpha}(x^\alpha - 1)}{\frac{d}{d\alpha}(a)} = \lim_{\alpha \rightarrow 0} \frac{x^\alpha \log_e x - 0}{1}$$

$$P = \lim_{\alpha \rightarrow 0} \frac{x^0 \log_e x - 0}{1}$$

$$P = \log_e x$$

$$\therefore \boxed{\lim_{\alpha \rightarrow 0} \frac{x^\alpha - 1}{a} = \log_e x}$$



Scan for Video solution



42. (c)

$$\text{Given, } f(x) = \begin{cases} \frac{\lambda \cos x}{\frac{\pi}{2} - x} & \text{if } x \neq \frac{\pi}{2} \\ 1 & \text{if } x = \frac{\pi}{2} \end{cases}$$

For function to be continuous at  $x = \frac{\pi}{2}$ ,

$$\text{LHL} = \text{RHL} = f(\pi/2)$$

Now,

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{\lambda \cos x}{\frac{\pi}{2} - x} = \left( \frac{0}{0} \right) \text{ form}$$

$$\left[ \left( \frac{0}{0} \right) \text{ form} \right]$$

Apply L – Hospital rule, For  $\left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form}$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{\lambda \cos x}{\frac{\pi}{2} - x} = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\frac{d}{dx}(\lambda \cos x)}{\frac{d}{dx}\left(\frac{\pi}{2} - x\right)} = \lambda$$

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{\lambda \cos x}{\frac{\pi}{2} - x} = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\lambda(-\sin x)}{-1} = \lambda$$

Therefore,  $\text{LHL} = \text{RHL} = \lambda$

... (i)


For function to be continuous at  $x = \frac{\pi}{2}$ ,

$$\text{LHL} = \text{RHL} = f(\pi/2)$$

From equation 9i),

$$\lambda = f(\pi/2)$$

$$\boxed{\lambda = 1}$$



Scan for Video solution



43. (a)

**Given:**  $P = \lim_{x \rightarrow 0} \frac{x - \sin x}{1 - \cos x}$

$$P = \lim_{x \rightarrow 0} \frac{x - \sin x}{1 - \cos x} = \left( \begin{array}{c} 0 \\ 0 \end{array} \right)$$

$\left[ \left( \begin{array}{c} 0 \\ 0 \end{array} \right) \text{ form} \right]$

$\left[ \text{Apply L-Hospital rule, For } \left( \begin{array}{c} 0 \\ 0 \end{array} \right) \text{ (or) } \left( \begin{array}{c} \infty \\ \infty \end{array} \right) \text{ form} \right]$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(x - \sin x)}{\frac{d}{dx}(1 - \cos x)}$$

$$P = \lim_{x \rightarrow 0} \frac{1 - \cos x}{\sin x} = \left( \begin{array}{c} 0 \\ 0 \end{array} \right)$$

Again apply L-hospital rule.

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(1 - \cos x)}{\frac{d}{dx}(\sin x)}$$

$$P = \lim_{x \rightarrow 0} \frac{0 + \sin x}{\cos x} = \lim_{x \rightarrow 0} \tan x = 0$$

$$\therefore \boxed{\lim_{x \rightarrow 0} \frac{x - \sin x}{1 - \cos x} = 0}$$



Scan for Video solution



44. (b)

**Given:**  $P = \lim_{x \rightarrow 0} \frac{e^{2x} - 1}{\sin 4x}$

$$P = \lim_{x \rightarrow 0} \frac{e^{2x} - 1}{\sin 4x} = \frac{1 - 1}{0} = \left( \begin{array}{c} 0 \\ 0 \end{array} \right)$$

$\left[ \left( \begin{array}{c} 0 \\ 0 \end{array} \right) \text{ form} \right]$

$\left[ \text{Apply L-Hospital rule, For } \left( \begin{array}{c} 0 \\ 0 \end{array} \right) \text{ (or) } \left( \begin{array}{c} \infty \\ \infty \end{array} \right) \text{ form} \right]$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(e^{2x} - 1)}{\frac{d}{dx}(\sin 4x)} = \lim_{x \rightarrow 0} \frac{2 \cdot e^{2x} - 0}{4 \cdot \cos 4x}$$

$$P = \frac{2 \cdot e^0}{4 \cdot \cos 0} = \frac{2}{4} = \frac{1}{2} = 0.5$$

$$\therefore \boxed{\lim_{x \rightarrow 0} \frac{e^{2x} - 1}{\sin 4x} = 0.5}$$



Scan for Video solution



45. (d)

For  $f(x)$  to be continuous at a point  $x = a$

$$\lim_{x \rightarrow a} f(x) = f(a)$$

$\Rightarrow$  Limit should exist and must be equal to functional value at that point.



Scan for Video solution



46. (a)

For a function  $f(x)$  to be continuous at  $x = 3$

$$\lim_{x \rightarrow 3} f(x) = f(3)$$

$$2; \quad \text{if } x = 3$$

$$(i) f(x) = x - 1; \quad \text{if } x > 3$$

$$\frac{x+3}{3}; \quad \text{if } x < 3$$

$$\therefore \lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} \frac{x+3}{3} = 2$$

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} x - 1 = 2$$

$$\therefore \lim_{x \rightarrow 3} f(x) = 2$$

$$\text{and } f(3) = 2$$

Since  $\lim_{x \rightarrow 3} f(x) = f(3)$

$\therefore f(x)$  is continuous at  $x = 3$

It

In all other options,  $\lim_{x \rightarrow 3} f(x)$  doesn't exist. (Since left hand limits are not equal to right hand limits)



Scan for Video solution



Therefore, function is continuous at  $x = 0$ .

**Checking for differentiability:**

$$y' = f'(x) = \begin{cases} 1 & ; x > 0 \\ -1 & ; x < 0 \end{cases}$$

Left hand derivative (LHD):

$$\lim_{x \rightarrow 0^-} f'(x) = \lim_{x \rightarrow 0^-} (-1) = -1$$

Right hand derivative (RHD):

$$\lim_{x \rightarrow 0^+} f'(x) = \lim_{x \rightarrow 0^+} 1 = 1$$

$\because LHD \neq RHD$ , therefore, function is not differentiable at  $x = 0$ .

$|x|$  is continuous at  $x = 0$  and not differentiable at  $x = 0$ .

**Method II:**

$$\text{Given: } f(x) = |x|, -1 \leq x \leq 1$$

$|x|$  is continuous  $\forall x \in \mathbb{R}$  and differentiable  $\forall x \in \mathbb{R}$  except at  $x = 0$ .

$\therefore$  At  $x = 0$ ,  $|x|$  is continuous but not differentiable.

47. (c)

The functions  $f(x)$ ,  $g(x)$ ,  $h(x)$  are said to be linearly dependent if  $f(x) = C_1g(x) + C_2h(x)$

Where  $C_1, C_2$  are non-zero constants.

We have  $\cos 2x = \cos^2 x - \sin^2 x$

$$\Rightarrow \cos 2x = 1 \cdot \cos^2 x + (-1) \cdot \sin^2 x$$

$\therefore \cos 2x, \cos^2 x, \sin^2 x$  are linearly dependent.



Scan for Video solution



48. (c)

**Method I:**

$$y = f(x) = |x|$$

$$y = f(x) = \begin{cases} x & ; x \geq 0 \\ -x & ; x < 0 \end{cases}$$

**Checking for continuity:**

Left hand limit (LHL):

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

Right hand limit (RHL):

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$f(0)$ :

$$f(0) = |0| = 0$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

49. (b)

$$\text{Given: } P = \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}$$

**Method I:**

$$P = \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \left( \frac{0}{0} \right)$$

$\left[ \left( \frac{0}{0} \right) \text{ form} \right]$

$\left[ \text{Apply L - Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \right]$

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(1 - \cos x)}{\frac{d}{dx}(x^2)} = \lim_{x \rightarrow 0} \frac{\sin x}{2x}$$

$$P = \frac{1}{2} \times \lim_{x \rightarrow 0} \frac{\sin x}{x}$$

$$P = \frac{1}{2} \times 1 = \frac{1}{2}$$

$$\therefore \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2}$$

**Method II:**

$$\left( \lim_{x \rightarrow 0} \frac{1 - \cos ax}{x^2} = \frac{a^2}{2} \right)$$

$$P = \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1^2}{2} = \frac{1}{2}$$

$$\therefore \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2}$$



Scan for Video solution



50. (d)

$$\text{Given: } P = \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta}$$

$$P = \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = \left( \frac{0}{0} \right) \left[ \left( \frac{0}{0} \right) \text{form} \right]$$

$$\left[ \begin{array}{l} \text{Apply L-Hospital rule, For } \left( \frac{0}{0} \text{ (or) } \frac{\infty}{\infty} \right) \text{ form} \\ \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} \end{array} \right]$$

$$P = \lim_{\theta \rightarrow 0} \frac{\frac{d}{d\theta}(\sin \theta)}{\frac{d}{d\theta}(\theta)} = \lim_{\theta \rightarrow 0} \frac{\cos \theta}{1} = \cos 0 = 1$$

$$\therefore \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

Note: Remember above expression directly.



Scan for Video solution



51. (c)

**Method I:**

$$y = f(x) = |2 - 3x|$$

$$\Rightarrow y = f(x) = \begin{cases} 2 - 3x & ; 2 - 3x \geq 0 \\ -(2 - 3x) & ; 2 - 3x < 0 \end{cases}$$

$$\Rightarrow y = f(x) = \begin{cases} 2 - 3x & ; x \leq \frac{2}{3} \\ -(2 - 3x) & ; x > \frac{2}{3} \end{cases}$$

For  $f(x)$  we need to check its continuity and differentiability at  $x = \frac{2}{3}$  only.

**Checking for continuity:**

Left hand limit (LHL):

$$\lim_{x \rightarrow \frac{2}{3}^-} f(x) = \lim_{x \rightarrow \frac{2}{3}^-} (2 - 3x) = \left( 2 - 3 \times \frac{2}{3} \right)$$

$$\Rightarrow \lim_{x \rightarrow \frac{2}{3}^-} f(x) = 0$$

Right hand limit (RHL):

$$\lim_{x \rightarrow \frac{2}{3}^+} f(x) = \lim_{x \rightarrow \frac{2}{3}^+} -(2 - 3x) = -\left( 2 - 3 \times \frac{2}{3} \right)$$

$$\Rightarrow \lim_{x \rightarrow \frac{2}{3}^+} f(x) = 0$$

$$f\left(\frac{2}{3}\right):$$

$$f\left(\frac{2}{3}\right) = \left| 2 - 3 \times \frac{2}{3} \right| = 0$$

$$\therefore \lim_{x \rightarrow \frac{2}{3}^-} f(x) = \lim_{x \rightarrow \frac{2}{3}^+} f(x) = f\left(\frac{2}{3}\right)$$

Therefore, function is continuous at  $x = \frac{2}{3}$ .

**Checking for differentiability:**

$$y' = f'(x) = \begin{cases} -3 & ; x < \frac{2}{3} \\ 3 & ; x > \frac{2}{3} \end{cases}$$

Left hand derivative (LHD):

$$\lim_{x \rightarrow \frac{2}{3}^-} f'(x) = \lim_{x \rightarrow \frac{2}{3}^-} (-3) = -3$$

Right hand derivative (RHD):

$$\lim_{\substack{x \rightarrow \frac{2}{3}^+ \\ x \rightarrow \frac{2}{3}^-}} f'(x) = \lim_{x \rightarrow \frac{2}{3}} 3 = 3$$

$\therefore \text{LHD} \neq \text{RHD}$ , therefore, function is not differentiable at  $x = \frac{2}{3}$ .

$|2 - 3x|$  is continuous  $\forall x \in \mathbb{R}$  and not differentiable at  $x = 2/3$ .

### Method II:

Given: function  $y = f(x) = |2 - 3x|$

- In  $|f(x)|$  (Where  $f(x)$  is polynomial) is continuous  $\forall x \in \mathbb{R}$  and not differentiable at the points where  $f(x) = 0$ .  
 $|2 - 3x|$  is continuous  $\forall x \in \mathbb{R}$  and differentiable  $\forall x \in \mathbb{R}$  except at the value where  $2 - 3x = 0 \Rightarrow x = \frac{2}{3}$ .  
 $\therefore |2 - 3x|$  is continuous  $\forall x \in \mathbb{R}$  and not differentiable at  $x = 2/3$ .



Scan for Video solution



### 52. (b)

Given limit is

$$P = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^{2n} = 1^\infty$$

If  $\lim_{n \rightarrow \infty} (f(x))^{g(x)} = 1^\infty$ , then

$$\lim_{n \rightarrow \infty} (f(x))^{g(x)} = e^{\lim_{n \rightarrow \infty} g(x) \cdot \{f(x)-1\}}$$

Therefore,

$$\therefore P = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^{2n} = e^{\lim_{n \rightarrow \infty} 2n \left(1 - \frac{1}{n} - 1\right)}$$

$$\Rightarrow P = e^{\lim_{n \rightarrow \infty} -2} = e^{-2}$$

$$\therefore \boxed{\lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^{2n} = e^{-2}}$$



Scan for Video solution



### 53. (a)

$$P = \lim_{x \rightarrow 0} \frac{\sin \frac{2}{3}x}{x}$$

$$P = \lim_{x \rightarrow 0} \frac{\frac{2}{3} \sin \frac{2}{3}x}{\frac{2}{3}x}$$

If  $x \rightarrow 0$ , then  $\frac{2}{3}x \rightarrow 0$

$$P = \frac{2}{3} \times \lim_{x \rightarrow 0} \frac{\sin \frac{2}{3}x}{\frac{2}{3}x} \quad \left[ \because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right]$$

$$P = \frac{2}{3} \times 1 = \frac{2}{3}$$

$$\therefore \boxed{\lim_{x \rightarrow 0} \frac{\sin \frac{2}{3}x}{x} = \frac{2}{3}}$$



Scan for Video solution



### 54. (a)

Given limit is

$$P = \lim_{x \rightarrow \infty} \frac{x - \sin x}{x + \cos x} = \lim_{x \rightarrow \infty} \frac{x \left(1 - \frac{\sin x}{x}\right)}{x \left(1 + \frac{\cos x}{x}\right)}$$

For any  $x \in \mathbb{R}$ ,  $\sin x \in [-1, 1]$  and  $\cos x \in [-1, 1]$

$$P = \frac{1 - \frac{\sin x}{x}}{1 + \frac{\cos x}{x}}$$

$$P = \frac{1 - 0}{1 + 0} = 1$$

$$\therefore \boxed{\lim_{x \rightarrow \infty} \frac{x - \sin x}{x + \cos x} = 1}$$



Scan for Video solution



**55. (b)**

**Given:**  $P = \lim_{x \rightarrow 8} \frac{x^{1/3} - 2}{(x - 8)}$

$$P = \lim_{x \rightarrow 8} \frac{x^{1/3} - 2}{(x - 8)} = \frac{8^{1/3} - 2}{(8 - 8)} = \left( \begin{array}{c} 0 \\ 0 \end{array} \right)$$

$\left[ \left( \begin{array}{c} 0 \\ 0 \end{array} \right) \text{ form} \right]$

Apply L-Hospital rule, For  $\left( \begin{array}{c} 0 \\ 0 \end{array} \right)$  (or)  $\left( \begin{array}{c} \infty \\ \infty \end{array} \right)$  form

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$P = \lim_{x \rightarrow 8} \frac{\frac{d}{dx}(x^{1/3} - 2)}{\frac{d}{dx}(x - 8)} = \lim_{x \rightarrow 8} \frac{\frac{1}{3} \cdot x^{-2/3} - 0}{1 - 0} = \frac{1}{3} \cdot (8)^{-2/3}$$

$$P = \frac{1}{3} \cdot (8)^{-2/3} = \frac{1}{3} \cdot (2^3)^{-2/3} = \frac{1}{3} \cdot 2^{-2} = \frac{1}{12}$$

$$\therefore \lim_{x \rightarrow 8} \frac{x^{1/3} - 2}{(x - 8)} = \frac{1}{12}$$



Scan for Video solution

**56. (0.50)**

Given  $\frac{dy}{dt} = \frac{e^{-\alpha t}}{2 + \alpha t}, \alpha = 0.01$  and  $y(0) = 0$

$$\frac{dy}{dt} = \frac{e^{-\alpha t}}{2 + \alpha t} \quad \dots(i)$$

At  $t = 0$ :

$$\Rightarrow \frac{dy}{dt} = y'(0) = \frac{1}{2}$$

Differentiating equation (i) with respect to  $t$ ,

$$\frac{d^2y}{dt^2} = \frac{-\alpha(2 + \alpha t)e^{-\alpha t} - e^{-\alpha t}(\alpha)}{(2 + \alpha t)^2}$$

At  $t = 0$ :

$$\Rightarrow \frac{d^2y}{dt^2} = y''(0) = \frac{-\alpha(2) \times 1 - 1 \times (\alpha)}{(2)^2} = \frac{-3\alpha}{4}$$

By Taylor series,

$$y(t) = y(0) + ty'(0) + \frac{t^2}{2!}y''(0) \dots$$

$$\left[ \because y(0) = 0, y'(0) = \frac{1}{2}, y''(0) = -\frac{3\alpha}{4} \right]$$

$$\Rightarrow y(t) = 0 + t\left(\frac{1}{2}\right) + \frac{t^2}{2}\left(-\frac{3\alpha}{4}\right) + \dots$$

 $\therefore$  At  $t = 1$ 

$$\Rightarrow y(1) = 0 + \frac{1}{2} + \frac{1}{2} \times \left(-\frac{3\alpha}{4}\right) + \dots$$

 $[\because \alpha = 0.01]$ 

$$\Rightarrow y(1) = \frac{1}{2} + \frac{1}{2} \left(-\frac{3 \times 0.01}{4}\right) + \dots$$

$$\Rightarrow y(1) = 0.5 - \frac{3}{8} \times 0.01 = 0.496$$

$$\Rightarrow y(1) = 0.50$$

Scan for Video solution

**57. (b)**Given, series is  $\sum_{n=1}^{\infty} \frac{n^d}{c^n} = \sum u_n$ By ratio test,  $\sum u_n$  converges if  $\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| < 1$  $\therefore$  Series converges if

$$\lim_{n \rightarrow \infty} \left| \frac{\left( \frac{(n+1)^d}{c^{n+1}} \right)}{\left( \frac{n^d}{c^n} \right)} \right| < 1$$

$$\Rightarrow \lim_{n \rightarrow \infty} \left| \frac{(n+1)^d}{c^{n+1}} \times \frac{c^n}{n^d} \right| < 1$$


$$\Rightarrow \frac{1}{|c|} \times \lim_{n \rightarrow \infty} \left| \frac{n+1}{n} \right|^d < 1$$

$$\Rightarrow \frac{1}{|c|} \times \lim_{n \rightarrow \infty} \left| \frac{\left(1 + \frac{1}{n}\right)^d}{1} \right| < 1$$

$$\Rightarrow \frac{1}{|c|} \times 1 < 1 \quad (\text{For any finite number } d)$$

$$\Rightarrow |c| > 1 \text{ and } d \text{ is a finite number.}$$

**∴ out of the given options, c = 2 and d = 1 is possible.**



58. (b)

Given:

$$f(x) = 8\log_e x - x^2 + 3 \text{ in } [1, e] \quad \dots(i)$$

Calculating  $f'(x)$

$$f'(x) = \frac{8}{x} - 2x$$

Stationary point: [At stationary point  $f'(x) = 0$ ]

$$f'(x) = \frac{8}{x} - 2x = 0$$

$$\Rightarrow x = \pm 2$$

But,  $x = 2$  is the only value that lies in  $[1, e]$ .

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = 1$  and  $e$ ) and in stationary point (i.e. at  $x = 2$ ):

$$f(x) = 8\log_e x - x^2 + 3$$

At  $x = 1$ ,

$$f(1) = 8\log_e 1 - 1^2 + 3 = 8(0) - 1 + 3 = 2$$

At  $x = e$ ,

$$f(e) = 8\log_e e - e^2 + 3 = 8 - e^2 + 3 = 3.611$$

At  $x = 2$ ,

$$f(2) = 8\log_e 2 - 2^2 + 3 = 8\log_e 2 - 2^2 + 3 = 4.545$$

**∴ Minimum of  $f(x)$  is at  $x = 1$ .**



59. (d)

$$\text{Given } f(x) = x^3 + 15x^2 - 33x - 36$$

$$\Rightarrow f'(x) = 3x^2 + 30x - 33$$

$$\text{Equating } f'(x) = 0$$

$$\Rightarrow 3x^2 + 30x - 33 = 0$$

$$\Rightarrow x^2 + 10x - 11 = 0$$

$$\Rightarrow x = -11, x = 1$$


$$\text{Considering } f''(x) \Rightarrow f''(x) = 6x + 30$$

$$\text{At } x = -11, f''(-11) = -66 + 30 < 0$$

**∴  $f(x)$  has local maxima at  $x = -11$**

$$\text{At } x = 1; f''(1) = 6 + 30 = 36 > 0$$


**∴  $f(x)$  has local minima at  $x = 1$**



60. (a)

→ Since  $e^x$  and  $|\sin x|$  both are continuous.


∴  $e^x |\sin x|$  will also be continuous.



→  $f(x)$  is not differentiable for all  $x \in \mathbb{R}$  because there are sharp edges.


→  $f(x)$  is not periodic

→  $f(x)$  is unbounded



61. (7.999 to 8.001)

Let 'L' be length of Rectangle and 'b' be breadth of Rectangle.



Given : area of rectangle;  $Lb = 4$

Find: Minimum value of perimeter ( P )

$$P = 2(L + b)$$

$$\left[ \because Lb = 4 \Rightarrow b = \frac{4}{L} \right]$$

$$P = 2\left(L + \frac{4}{L}\right)$$

$\therefore$  For minimum of perimeter,

$$\frac{dP}{dL} = 0 \Rightarrow 2\left(L - \frac{4}{L^2}\right) = 0$$

$$L = 2$$

$\therefore$  For minimum perimeter,  $L = 2$

$$\text{and } b = \frac{4}{L} = \frac{4}{2} = 2$$

Hence minimum perimeter,

$$P = 2(L + b) = 2(4) = 8 \text{ units}$$



Scan for Video solution



## 62. (d)

$$f(x) = x^3 - 6x^2 + 11x - 6; x \in [1, 3]$$

for  $x = 1$

$$f(1) = 1^3 - 6(1)^2 + 11(1) - 6 = 0$$

for  $x = 3$

$$f(3) = 3^3 - 6(3^2) + 11(3) - 6 = 0$$

Therefore statement I is correct

Differentiating equation (i) with respect to x,

$$f'(x) = 3x^2 - 12x + 11 \quad \dots (\text{ii})$$

Differentiating equation (ii) with respect to x,

$$f''(x) = 6x - 12 \quad \dots (\text{iii})$$

Stationary point:

At stationary point  $f'(x) = 0$

$$f'(x) = 0$$


$$\Rightarrow 3x^2 - 12x + 11 = 0$$

$$\Rightarrow x = \frac{-(-12) \pm \sqrt{(-12)^2 - 4 \times 3 \times 11}}{2 \times 3}$$

$$\Rightarrow x = 2.577, 1.423 \in [1, 3]$$

$$\text{Now } f'(x) = 6(x - 2)$$

Drawing  $f'(x)$  on number line:



$$f''(x) = \begin{cases} < 0 & \text{for } x < 2 \\ > 0 & \text{for } x > 2 \end{cases}$$


At stationary point  $x = 1.423$  ( $x < 2$ )  $f''(x)$  is  $< 0$ , therefore  $x = 1.423$  is local maxima and at stationary point  $x = 2.577$  (ie.  $x > 2$ )  $f''(x)$  is  $> 0$ , therefore  $x = 2.577$  is local minima.

Hence in domain S. There exist a local maxima and a local minima.

$\therefore$  Both statement II and IV are correct.

Since  $f''(x)$  is not  $> 0$  for throughout domain s, therefore statement III is incorrect.

$\therefore$  Correct statements are I, II and III therefore option D is correct



## 63. (a, b)


For intersection of  $3 - x$  and  $x - 1$

$$3 - x = x - 1$$

$\therefore$  At  $x = 2$  these lines will intersect.

$$f(x) = \max \{3 - x, x - 1\}$$

$$f(x) = \begin{cases} 3 - x & x < 2 \\ x - 1 & x > 2 \\ 1 & x = 2 \end{cases}$$



Check for continuity:-

$$\text{LHL} = \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (3 - x) = 1$$

$$\text{LHL} = \lim_{x \rightarrow 2^=} f(x) = \lim_{x \rightarrow 2^+} (x - 1) = 1$$

$$f(2) = 1$$

Since at  $x=2$  LHL = RHL =  $f(2)$   $\therefore$  It is continuous in its domain.

#### Check for differentiability: –

From graph we can infer that  $f(x)$  has sharp edge at  $x=2$ , therefore, function is not differential at  $x=2$ .

From graph of this function it is evident that function has local minimum at  $x=2$



Scan for Video solution



**64. (b)**

Given

$$f(x) = \frac{1}{2} x^T Q x - r^T x; \text{ and}$$

$$Q \text{ is symmetric matrix } \Rightarrow Q = Q^T$$

$$f'(x) = \frac{1}{2} \times \frac{d}{dx} (x^T Q x) - \frac{d}{dx} (r^T x) \\ \left( \because \frac{d}{dx} (x^T Q x) = x^T (Q + Q^T) \right)$$

$$f'(x) = \frac{1}{2} \cdot \{x^T (Q + Q^T)\} - r^T \\ (\because Q^T = Q)$$

$$f'(x) = \frac{1}{2} \cdot \{x^T (2Q)\} - r^T$$

$$f'(x) = x^T Q - r^T$$

At stationary point,  $f'(x) = 0$ :

$$f'(x) = 0$$

$$x^T Q - r^T = 0$$

$$(x^T Q - r^T)^T = 0^T$$

$$(x^T Q - r^T)^T = 0$$

$$Q^T \cdot x - r = 0$$

$$x = Q^{-1} \cdot r$$



Scan for Video solution



**65. (19)**

Given  $f: \mathbb{R} \rightarrow \mathbb{R}$  is continuous in  $[-3, 3]$  and differentiable in  $(-3, 3)$

$$\text{By mean value theorem, } f'(c) = \frac{f(b) - f(a)}{b - a}$$

Where  $c \in (a, b)$

$$\Rightarrow f'(x) = \frac{f(3) - f(-3)}{3 - (-3)}$$

$$\Rightarrow f(3) = 6f'(x) + f(-3)$$

$$\text{Since } f'(x) \leq 2$$

$$\Rightarrow f(3) \leq 6(2) + f(-3)$$

$$\Rightarrow f(3) \leq 12 + 7 \leq 19$$

$$\therefore f(3) \leq 19$$

**$\Rightarrow f(3)$  is equal to 19 at most.**



Scan for Video solution



**66. (c)**

Given,  $f'(x_0) = 0$  at  $x_0 \in (0, 1)$ .

$\therefore x_0$  is a stationary or turning point. One stationary point is coming in interval  $(0, 1)$ . Now if  $f''(x_0) > 0$  at stationary point, then it will local minimum. Conversely, if  $f''(x_0) < 0$ , then it will be local maximum.

Since  $f''(x_0) > 0$  in all  $x$  in interval  $(0, 1)$ , it follows that at stationary point  $x_0$  also  $f''(x_0) > 0$ . Therefore, the stationary point  $x_0$  is local minimum. **Hence, there exists only one local minimum in interval  $(0, 1)$ .**



Scan for Video solution



**67. (1)**

Condition for orthogonality: -

$$m_1 \cdot m_2 = -1 \quad \dots (i)$$

$m_1 \rightarrow$  slope of curve  $-1$  at  $(u, v)$

$m_2 \rightarrow$  slope of curve  $-2$  at  $(u, v)$

Curve 1:

$$x^2 + y^2 = 1$$

Differentiate above equation with respect to x,

$$\Rightarrow 2x + 2y \frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = -\frac{x}{y}$$

Slope of curve -1 at (u,v):

$$m_1 = \left. \frac{dy}{dx} \right|_{(u,v)} = -\frac{u}{v} \quad \dots \text{(ii)}$$

Curve-2:-

$$(x-1)^2 + (y-1)^2 = r^2$$

Differentiate above equation with respect to x,

$$\Rightarrow 2(x-1) + 2(y-1) \frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = -\left( \frac{x-1}{1-y} \right)$$

Slope of curve -2 at (u,v):

$$m_2 = \left. \frac{dy}{dx} \right|_{(u,v)} = -\left( \frac{u-1}{1-v} \right) \quad \dots \text{(iii)}$$

Since both curves are intersecting at (u, v), therefore

$$m_1 \cdot m_2 = -1$$

$$\Rightarrow \left( -\frac{u}{v} \right) \cdot \left( \frac{u-1}{1-v} \right) = -1$$

$$\Rightarrow u^2 - u = v - v^2$$

$$\Rightarrow u + v = u^2 + v^2 \quad \dots \text{(v)}$$

Since (u, v) is intersection point of curve 1 and 2, therefore it will satisfy the equation of both curve 1 and 2. In equation of curve (i) put x=u and y=v,  $u^2 + v^2 = 1$

From equation (v),

$$\Rightarrow u + v = 1$$



Scan for Video solution



68. (b)

**Given:**

$$P(x) = x^4 - 4x^3 + 2 \dots \text{(i)}$$

To determine the number of real root in open interval (0, 1), we can analyze the behaviour of

polynomial and its derivative differentiating equation (i) with respect to x.

$$P'(x) = 4x^3 - 12x^2$$

$$\Rightarrow P'(x) = 4x^2(x-3)$$


Finding stationary point:

$$P'(x) = 0$$

$$\Rightarrow 4x^2(x-3) = 0$$

$$\Rightarrow x = 0 \text{ and } 3$$

Drawing  $P'(x)$  on number line;




Since in interval (0, 1)  $P'(x)$  is negative, therefore function in interval (0, 1) is decreasing function.

Now we will find value of function  $P(x)$  at  $x = 0$  and at  $x = 1$ .


$$P(0) = 0^4 - 4 \times (0)^3 + 2 = 2 \text{ (+ve)}$$

$$P(1) = 1^4 - 4 \times (1)^3 + 2 = 1 \text{ (-ve)}$$

Since sign of  $P(0)$  and  $P(1)$  is opposite and  $P(x)$  is decreasing function in interval [0, 1], therefore graph of  $p(x)$  in interval [0, 1] will be as given below.



(0, 1) in interval therefore  $f(x)$  has only one real root.



69. (a, d)

Given polynomial:  $-ax^3 + bx^2 + cx + d \quad a \neq 0$

**Option a)** If  $d = 0$ ;  $-ax^3 + bx^2 + cx = 0$

$$x(ax^2 + bx + c) = 0$$

$\therefore x = 0$  is a solution. Hence option (a) is correct.

**Option b)**

If all roots are identical then the given polynomial can be expressed as

$$ax^3 + bx^2 + cx + d = (x - r)^3$$
  

$$\Rightarrow ax^3 + bx^2 + cx + d = x^3 - 3x^2r + 3xr^2 - r^3$$
  
 On comparing,  $a = 1$ ,  $b = -3r$ ,  $c = 3r^2$ ,  $d = -r^3$   
 ∴ The above choice of coefficients can make all roots same. Hence option (b) is incorrect.

**Option c)**

All 3 roots cannot be complex because complex roots occur in conjugate pairs. Either 0 or 2 complex roots are possible. Hence option (c) is incorrect.

**Option d)** All coefficient along with c will decide the nature of roots. Hence option (d) is correct.



Scan for Video solution

**70. (a)**

$$y = f(x) = 3x^2 + 3x + 1 \quad \dots(i)$$

$f(x)$  is continuous and differential function in interval  $[-2, 0]$ .

Differentiate equation (i) with respect to 'x',

$$y' = f'(x) = 6x + 3$$

Stationary point: [At stationary point  $f'(x) = 0$ ]

$$y' = f'(x) = 6x + 3 = 0$$

$$\Rightarrow x = -1/2 \in [-2, 0]$$

Finding value of  $f(x)$  at initial and final point (i.e. at

$x = -2$  and  $0$ ) in stationary point (i.e. at  $x = -\frac{1}{2}$ ):

$$y = f(x) = 3x^2 + 3x + 1$$

At  $x = -2$ :

$$f(-2) = 3(-2)^2 + 3(-2) + 1 = 7$$

At  $x = 0$ :

$$f(0) = 3(0)^2 + 3(0) + 1 = 1$$

At  $x = \frac{1}{2}$ :

$$f\left(-\frac{1}{2}\right) = 3\left(-\frac{1}{2}\right)^2 + 3\left(-\frac{1}{2}\right) + 1 = \frac{1}{4}$$

∴ Maximum and minimum values of  $y$  in  $[-2, 0]$

are 7 and  $\frac{1}{4}$  respectively.


Scan for Video solution



**71. (a)**

Given function are

- (i)  $e^{-x}$
- (ii)  $x^2 - \sin x$
- (iii)  $\sqrt{x^3 + 1}$


For a function  $f(x)$  to be increasing in  $[0, 1]$ ,  $f'(x) > 0$  in  $[0, 1]$

(i) let,  $f(x) = e^{-x}$ :

$$\text{Then, } f'(x) = -e^{-x}$$

Since  $f'(x) = -e^{-x}$  is always less than zero therefore  $f(x) = e^{-x}$  is decreasing function  $\forall x \in \mathbb{R}$

(ii)  $f(x) = x^2 - \sin x \Rightarrow f'(x) = 2x - \cos x$



Since  $\cos x > 2x$  for some  $x \in [0, 1]$

$$\Rightarrow 2x - \cos x < 0 \text{ for some } x \in [0, 1]$$

∴  $x^2 - \sin x$  is not completely increasing in  $[0, 1]$

(iii)  $f(x) = \sqrt{x^3 + 1}$

$$\Rightarrow f'(x) = \frac{3x^2}{2\sqrt{x^3 + 1}} > 0 \quad \forall x \in [0, 1]$$

∴  $f(x) = \sqrt{x^3 + 1}$  is the only function that is increasing  $\forall x \in [0, 1]$


Scan for Video solution



**72. (5.18 to 5.2)**

**Given:** Diameter ( $D$ ) = 3 cm ; Height ( $H$ ) = 4 cm

Error ( $dD$  or  $dH$ ) =  $\pm 0.02$  cm

$$\text{Volume } V = \frac{\pi}{4} D^2 H$$

Absolute error in volume;

$$dV = \frac{\partial V}{\partial D} \cdot dD + \frac{\partial V}{\partial H} \cdot dH$$

$$dV = \left( \frac{\pi DH}{2} \right) dD + \left( \frac{\pi}{4} D^2 \right) dH$$

$$dV = \left( \frac{\pi}{2} \times 3 \times 4 \right) \times 0.02 + \left( \frac{\pi}{4} \times 3^2 \right) \times 0.02$$

$$dV = 1.65 \pi \text{ cm}^3 = 5.1836 \text{ cm}^3$$

**Absolute error in volume = 5.184 cm<sup>3</sup>**



Scan for Video solution



73. (c)

**Given:** Family of curves  $\frac{dy}{dx} = -\left(\frac{x}{y}\right)^n$

$$(i) \text{ for } n = -1, \frac{dy}{dx} = -\left(\frac{x}{y}\right)^{-1} = -\frac{y}{x}$$

$$\Rightarrow \frac{1}{y} dy = \frac{-1}{x} dx$$

$$\Rightarrow \int \frac{1}{y} dy = \int \frac{-1}{x} dx$$

$$\Rightarrow \ln y = -\ln x + c$$

$$\Rightarrow xy = c$$

$\therefore x y = c \rightarrow \text{hyperbolas}$

$$(ii) \text{ For } n = 1; \frac{dy}{dx} = -\frac{x}{y}$$

$$\Rightarrow ydx = -xdx$$

$$\Rightarrow \int ydx = \int -xdx$$

$$\Rightarrow \frac{y^2}{2} = \frac{-x^2}{2} + C \quad \Rightarrow x^2 + y^2 = C'$$

$\therefore x^2 + y^2 = C' \rightarrow \text{circles}$

**Hence, the family of curves for  $n = -1$  and  $n = +1$  are hyperbolas and circles respectively.**

▶
Scan for Video solution

74. (a)

Checking for function  $\sqrt[3]{x}$ :

$$\text{Let } y = \sqrt[3]{x} = x^{1/3}$$

$$\log y = \frac{\log x}{x}$$

$$y = e^{\frac{\log x}{x}}$$

Maxima and minima of  $y$  will depends on  $\frac{\log x}{x}$

$$\text{Let } f(x) = \frac{\log x}{x}$$

$$f'(x) = \frac{\frac{1}{x} - \log x \cdot 1}{x^2} = \frac{1 - \log x}{x^2}$$

At stationary point:

$$f'(x) = 0$$

$$\frac{1 - \log x}{x^2} = 0$$

$$\log x = 1$$

$$x = e$$

Now,

$$f''(x) = \frac{x^2 \cdot \left(-\frac{1}{x}\right) - (1 - \log x) \cdot 2x}{x^4} = \frac{-3x + 2x \log x}{x^4}$$

At  $x = e$ :

$$f''(e) = \frac{-3e + 2e \log e}{e^4} = -\frac{e}{e^4} < 0$$


Since at stationary point  $x = e$ ,  $f''(x)$  is less than 0,

hence, the function  $\sqrt[3]{x}, (x > 0)$  has the global

maxima at  $x = e$ .


Hence statement in options (a) is incorrect.

Checking for function  $x^3$ :



From graph of  $x^3$ , we infer it has neither global maxima nor global minima.

Checking for function  $|x|$ :



From graph of  $|x|$ , we conclude it has global minimum at  $x = 0$ .



Scan for Video solution



### 75. (c)

Taylor's series expansion about  $x$  for  $f(x + h)$  is  
 $f(x + h) = f(x) + (x + h - x)f'(x) + \dots$

$$f(x + h) = f(x) + (x + h - x)f'(x) + \frac{(x + h - x)^2}{2!}f''(x) + \dots$$

$$\therefore f(x + h) = f(x) + hf'(x) + \frac{h^2}{2!}f''(x) + \dots$$



Scan for Video solution



### 76. (d)

$$f(x) = x^3$$


$$f'(x) = 3x^2$$

For stationary points,

$$f'(x) = 3x^2 = 0$$

$\Rightarrow x = 0$  is a stationary point.

$f'(x)$  on number line:




$\therefore f'(x)$  is not changing its sign at  $x = 0$ ,  $\therefore f(x)$  at  $x = 0$  is neither local maxima nor minima now,

$$f''(x) = \frac{d}{dx}(3x^2) = 6x$$


At  $x = 0$ ,  $f''(x) = 0$

$f''(x) = 6x$  on number line



$\therefore f''(x)$  is changing its sign at

$x = 0$ , hence  $f(x) = x^3$  has point of inflection at  $x = 0$



From the graph of  $x^3$ , we can also conclude that at  $x = 0$ ; point of inflection occurs because nature of curve is changing at  $x = 0$ .



Scan for Video solution



### 77. (11.5 to 12.5)

$$f(x) = 3x^3 - 7x^2 + 5x + 6$$

... (i)

$f(x)$  is continuous and differential function in interval  $[0, 2]$ .

Differentiate equation (i) with respect to  $x$ ,

$$f'(x) = 9x^2 - 14x + 5$$

Stationary point: [At stationary point  $f'(x) = 0$ ]

$$f'(x) = 9x^2 - 14x + 5 = 0$$

$$\Rightarrow (9x - 5)(x - 1) = 0$$

$$\Rightarrow x = 1 \text{ and } \frac{5}{9} \in [0, 2]$$

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = 0$  and  $2$ ) in stationary points (i.e. at  $x = 1$  and  $\frac{5}{9}$ ):

$$f(x) = 3x^3 - 7x^2 + 5x + 6$$

At  $x = 0$ :

$$f(0) = 3(0)^3 - 7(0)^2 + 5(0) + 6 = 6$$

At  $x = 2$ :

$$f(2) = 3(2)^3 - 7(2)^2 + 5(2) + 6 = 12 \leftarrow \text{maximum}$$

At  $x = 1$ :

$$f(1) = 3(1)^3 - 7(1)^2 + 5(1) + 6 = 7$$

At  $x = \frac{5}{9}$ :

$$f\left(\frac{5}{9}\right) = 3\left(\frac{5}{9}\right)^3 - 7\left(\frac{5}{9}\right)^2 + 5\left(\frac{5}{9}\right) + 6 = 7 \cdot 13$$

Hence, Maximum value of  $f(x)$  in interval  $[0, 2]$  is 12.



Scan for Video solution



78. (7)

$$2x^2 + y^2 = 34 \quad \dots\dots(1)$$

$$x + 2y = 11$$

$$y = \frac{11-x}{2} \quad \dots (2)$$

From value of  $y$  from equation (2) to equation (1),

$$2x^2 + \left(\frac{11-x}{2}\right)^2 = 34$$

$$\Rightarrow 8x^2 + 121 + x^2 - 22x = 136$$

$$\Rightarrow 9x^2 - 22x - 15 = 0$$

$$\Rightarrow x = \frac{-10}{18}, 3$$

Discarding  $x = -10/18$  because it is not an integer.

Hence  $x = 3$ .

Put value of  $x$  in equation (2),

$$y = \frac{11-x}{2} = \frac{11-3}{2} = 4$$

Hence,

$$x + y = 3 + 4$$

$$\Rightarrow \boxed{x + y = 7}$$




Scan for Video solution



79. (a)

**Function is**  $y = x \ln x \quad \dots(i)$

Say point is A at which inclination of tangent with the x-axis is  $45^\circ$ .



Slope at point A =  $\tan 45^\circ$

$$\frac{dy}{dx} = \tan 45^\circ$$

$$\frac{d}{dx}(x \ln x) = 1$$

$$x \cdot \frac{1}{x} + \ln x \cdot 1 = 1$$

$$1 + \ln x = 1$$

$$\ln x = 0$$

$$x = 1$$

At  $x = 1$  from equation (i),

$$y = x \ln x = 1 \ln 1 = 1 \times 0 = 0$$

$\therefore \boxed{\text{Point is } (1, 0)}$



Scan for Video solution



80. (c)

Given  $f(x) = e^{x+x^2}$

The Taylor series expansion is written about  $x=0$

$$f(x) = e^{x+x^2}$$

$$f'(x) = e^{x+x^2} (1 + 2x)$$

$$\Rightarrow f'(x)|_{x=0} = f'(0) = 1$$

$$f''(x) = e^{x+x^2} (2) + (1+2x)^2 e^{x+x^2}$$

$$\Rightarrow f''(x) = e^{x+x^2} \{4x^2 + 4x + 3\}$$

$$\Rightarrow f''(x)|_{x=0} = f''(0) = 3$$

$$f'''(x) = \left[ e^{x+x^2} \{8x+4\} + (4x^2+4x+3)(1+2x)e^{x+x^2} \right]$$

$$\Rightarrow f'''(x)|_{x=0} = f'''(0) = 4 + 3 = 7$$

$$\therefore \text{coefficient of } x^2 = \frac{f''(0)}{2!} = \frac{3}{2!} = \frac{3}{2}$$

$$\text{Coefficient of } x^3 = \frac{f'''(0)}{3!} = \frac{7}{6}$$

$$\therefore f(x) = 1 + x + \frac{3}{2}x^2 + \frac{7}{6}x^3$$



Scan for Video solution



### 81. (-100)

$$\text{Given, } f(x) = \frac{1}{3}x(x^2 - 3); -100 \leq x \leq 100$$

$$f(x) = \frac{1}{3}(x^3 - 3x)$$

Calculating  $f'(x)$ :

$$f'(x) = \frac{1}{3}(3x^2 - 3) = x^2 - 1$$

At stationary point:

$$f'(x) = 0$$

$$\Rightarrow x^2 - 1 = 0$$

$$\Rightarrow x = 1, -1 \in [-100, 100]$$

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = -100$  and  $100$ ) in stationary point (i.e. at  $x = 1$  and  $-1$ ):

$$f(x) = \frac{1}{3}x(x^2 - 3)$$

At  $x = -100$ ,

$$f(-100) = \frac{1}{3}(-100)((-100)^2 - 3) = -333233.33$$

At  $x = 100$ ,

$$f(100) = \frac{1}{3}(100)((100)^2 - 3) = 333233.33$$

At  $x = 1$ ,


$$f(1) = \frac{1}{3}(1)((1)^2 - 3) = -0.667$$

At  $x = -1$ ,

$$f(-1) = \frac{1}{3}(-1)((-1)^2 - 3) = 0.667$$


$\therefore$  Minimum value of  $f(x)$  in  $[-100, 100]$   
 $= \min\{f(-100), f(1), f(-1), f(100)\}$   
 $= \min \{-333233.33, 0.667, -0.667, 333233.33\}$

$\therefore$  Min of  $f(x)$  in  $[-100, 100]$  is  $-333233.33$  at  $x = -100$




82. (c)

$$\text{Given, } \sin x = \frac{x}{2}$$



The curves  $y = \sin x$  and  $y = \frac{x}{2}$  intersect at 3 locations.

$\therefore$  Number of solutions of  $\sin x = \frac{x}{2}$  is 3.



**83. (0 to 0)**

$$f(x) = x(x-1)(x-2)$$

$$\Rightarrow f(x) = x^3 - 3x^2 + 2x \quad \dots(i)$$

$f(x)$  is continuous and differential function in interval  $[1, 2]$ .

Differentiate equation (i) with respect to  $x$ ,

$$f'(x) = 3x^2 - 6x + 2 \quad \dots(ii)$$

Stationary point: [At stationary point  $f'(x) = 0$ ]

$$f'(x) = 3x^2 - 6x + 2 = 0$$

$$\Rightarrow x = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(3)(2)}}{2(3)}$$

$$\Rightarrow x = \frac{6 \pm 2\sqrt{3}}{6}$$

$$\Rightarrow x = 1 + \frac{1}{\sqrt{3}}, 1 - \frac{1}{\sqrt{3}}$$

$$\Rightarrow x = 1 + \frac{1}{\sqrt{3}} \in [1, 2] \text{ & } x = 1 - \frac{1}{\sqrt{3}} \notin [1, 2]$$

Hence stationary point in interval  $[-3, 3]$  is

$$\Rightarrow x = 1 + \frac{1}{\sqrt{3}}.$$

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = 1$  and 2) in stationary points

$$(i.e. \text{ at } x = 1 + \frac{1}{\sqrt{3}}) :$$

$$f(x) = x(x-1)(x-2)$$

At  $x = 1$ :

$$f(1) = 1(1-1)(1-2) = 0$$

At  $x = 2$ :

$$f(2) = 2(2-1)(2-2) = 0$$

$$\text{At } x = 1 + \frac{1}{\sqrt{3}} :$$

$$f\left(1 + \frac{1}{\sqrt{3}}\right) = \left(1 + \frac{1}{\sqrt{3}}\right)\left(1 + \frac{1}{\sqrt{3}} - 1\right)\left(1 + \frac{1}{\sqrt{3}} - 2\right)$$

$$\Rightarrow f\left(1 + \frac{1}{\sqrt{3}}\right) = -0.3849$$

**Hence maximum value of  $f(x)$  is 0.**



Scan for Video solution

**84. (b)**

Given  $f(x) = x^3 - 3x^2 + 1$  in  $[-1, 3]$

$$\Rightarrow f'(x) = 3x^2 - 6x = 3x(x-2)$$

For  $f(x)$  to be increasing,

$$f'(x) > 0$$

$$\Rightarrow 3x(x-2) > 0$$

$$\Rightarrow x > 0 \text{ and } x > 2 \text{ or } x < 0 \text{ and } x < 2$$

$$\Rightarrow x > 2 \text{ or } x < 0$$

[In interval -1 to 3]

$$\Rightarrow (2, 3) \text{ or } (-1, 0)$$

For  $f(x)$  to be decreasing,

$$f'(x) < 0$$

$$\Rightarrow 3x(x-2) < 0$$

$$\Rightarrow x < 0 \text{ and } x > 2 \text{ or } x > 0 \text{ and } x < 2$$

$$\Rightarrow 0 < x < 2$$

[In interval -1 to 3]

$$\Rightarrow (0, 2)$$

Therefore, in interval -1 to 3,  $f(x)$  is initially increasing in  $(-1, 0)$  then decreasing in  $(0, 2)$  and again increasing in  $(2, 3)$ .


Scan for Video solution


**85. (b)**

By Taylor's expansion of  $f(x)$  about  $x = a$

$$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2!}f''(a) + \dots$$

About  $x = 0 \Rightarrow a = 0$

$$f(x) = f(0) + x f'(0) + \frac{x^2}{2!}f''(0) + \dots \quad \dots(i)$$

Now,

$$f(x) = x^3 - 3x^2 - 5 = 0^3 - 3(0)^2 - 5 = -5$$

$$f'(0) = 3x^2 - 6x = 3(0)^2 - 6(0) = 0$$


$$f''(0) = 6x - 6 = 6(0) - 6 = -6$$

Put the values in (i)

$$f(x) = -5 + (x \times 0) + \frac{x^2}{2}(-6) = -3x^2 - 5$$


Scan for Video solution


86. (d)

Slope of  $x^2 = 4y$ :

$$x^2 = 4y \Rightarrow y = \frac{x^2}{4}$$

$$\text{Slope, } \frac{dy}{dx} = \frac{2x}{4} = \frac{x}{2}$$

$$\text{Slope at } (0, 0) = \frac{x}{2} = \frac{0}{2} = 0 \text{ (hence tangent at origin)}$$

will be horizontal)

Slope of  $y^2 = 4x$ :

$$y^2 = 4x \Rightarrow y = 2\sqrt{x}$$

$$\text{Slope, } \frac{dy}{dx} = 2 \cdot \frac{1}{2\sqrt{x}} = \frac{1}{\sqrt{x}}$$

$$\text{Slope at } (0, 0) = \frac{1}{\sqrt{x}} = \frac{1}{\sqrt{0}} = \infty \text{ (hence tangent at origin will be vertical)}$$

Hence angle between the curves at  $(0, 0)$  is  $90^\circ$ 

Scan for Video solution



87. (d)

$$f(x) = x^2 - 4x + 2$$

$$f'(x) = 2x - 4 \text{ and}$$

$$f''(x) = 2$$

Find stationary points using  $f'(x) = 0$ 

$$f'(x) = 2x - 4 = 0$$

$$x = 2$$

At  $x = 2$ ,  $f''(x) = 2 > 0 \Rightarrow$  hence  $x = 2$  is minima.

Therefore,

$$f_{\min} = x^2 - 4x + 2 \text{ at } x = 2$$


$$f_{\min} = 2^2 - 4(2) + 2$$

$$f_{\min} = -2$$



Scan for Video solution



**90. (1)**Given ellipse is  $x^2 + 4y^2 = 1$ 

$$\Rightarrow \frac{x^2}{1} + \frac{y^2}{\left(\frac{1}{4}\right)} = 1$$

$$\Rightarrow \frac{x^2}{1^2} + \frac{y^2}{\left(\frac{1}{2}\right)^2} = 1$$

Let  $2a, 2b$  are lengths are breadths of the rectangle. $\therefore$  Area of the rectangle  $= A = 4ab$ And also  $a^2 + 4b^2 = 1$ 

$$\Rightarrow A = 4\sqrt{1 - 4b^2} \cdot b$$

For 'A' to be maximum

$$\frac{dA}{db} = 0 \Rightarrow \frac{d}{db} \left( b \cdot \sqrt{1 - 4b^2} \right) = 0$$

$$\Rightarrow b \frac{1}{2\sqrt{1 - 4b^2}} \times (-8b) + \sqrt{1 - 4b^2} \cdot (1) = 0$$

$$\Rightarrow -\frac{4b^2}{\sqrt{1 - 4b^2}} + \sqrt{1 - 4b^2} = 0$$

$$\Rightarrow \frac{-8b^2 + 1}{\sqrt{1 - 4b^2}} = 0$$

$$\Rightarrow 1 - 8b^2 = 0$$

$$\Rightarrow b = \frac{1}{2\sqrt{2}}$$

$$\therefore 2b = \frac{1}{\sqrt{2}}$$

Since  $a^2 + 4b^2 = 1$ 

$$\Rightarrow a^2 + \frac{1}{2} = 1 \Rightarrow a = \frac{1}{\sqrt{2}}$$

$$\Rightarrow 2a = \sqrt{2}$$

$$\therefore A_{\max} = 4ab = 4 \times \frac{1}{\sqrt{2}} \cdot \frac{1}{2\sqrt{2}} = 1 \text{ sq units}$$



Scan for Video solution

**91. (b)**Given  $f(x) = e^{-x} (x^2 + x + 1)$ 

$$= e^{-x} \left\{ \left( x + \frac{1}{2} \right)^2 + \frac{3}{4} \right\}$$

For  $x > 0, f(x) > 0$ 

$$\text{Consider } f'(x) = e^{-x} \{2x + 1\} + (x^2 + x + 1)(-e^{-x})$$

$$= e^{-x} \{2x + 1 - x^2 - x - 1\}$$

$$= e^{-x} \{x - x^2\} = e^{-x} x(1 - x)$$

For  $x > 0; e^{-x} > 0;$ And for  $0 < x < 1; x(1 - x) > 0$ and for  $x > 1; x(1 - x) < 0$ 

Therefore,

For  $0 < x < 1; f'(x) > 0$ For  $x > 1, f'(x) < 0$  $f(x)$  is increasing in  $0 < x < 1$  and decreasing in  $x > 1$ . $\therefore$  Option 'B' is correct

Scan for Video solution

**92. (b)**Given  $f(x) = 1 - x^2 + x^3; [-1, 1]$ By mean value theorem of  $f(x)$  in the interval  $[a, b]$ 

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

for  $f(x) = 1 - x^2 + x^3$ 

$$\Rightarrow f'(x) = 3x^2 - 2x$$

 $\Rightarrow$  By mean value theorem

$$f(c) = \frac{f(1) - f(-1)}{1 - (-1)}$$

$$\Rightarrow 3c^2 - 2c = \frac{1 - (-1)}{1 - (-1)}$$

$$\Rightarrow 3c^2 - 2c - 1 = 0$$

$$\Rightarrow 3c^2 - 3c + c - 1 = 0$$

$$\Rightarrow 3c(c-1) + 1(c-1) = 0 \Rightarrow c = \frac{-1}{3} \text{ and } c = 1$$

Since  $c \in (-1, 1)$ , the mean value 'c' is equal to  $\frac{-1}{3}$




Scan for Video solution



**93. (c)**

$$\text{Given } f(x) = x^{-\frac{1}{3}} = \frac{1}{x^{\frac{1}{3}}}$$



- (a) At  $x=0$ ;  $f(x)$  is not bounded  
As  $x \rightarrow 0$ ;  $f(x) \rightarrow \infty$
- (b)  $f(x)$  is not continuous at  $x=0$
- (c)  $A = \int_{-1}^1 x^{-1} dx = 3 \leftarrow \text{finite}$



Scan for Video solution



**94. (d)**

If  $f'(x_0) = 0$

If  $f''(x_0) > 0 \Rightarrow$  local minimum

If  $f''(x_0) < 0 \Rightarrow$  local maximum




Scan for Video solution



**95. (a)**

At  $x = 0$ , the function  $|x|$  has a minimum and minimum of  $|x| = 0$



Scan for Video solution



**96. (a)**

Given,

$$f(t) = e^{-t} - 2e^{-2t}$$

Calculating  $f'(t)$ ,

$$f'(t) = -e^{-t} + 4e^{-2t}$$

At stationary point:

$$\Rightarrow f'(t) = -e^{-t} + 4e^{-2t} = 0$$

$$\Rightarrow e^{-t}(-1 + 4e^{-t}) = 0$$

$$\Rightarrow e^{-t} = 0 \Rightarrow t \rightarrow \infty \notin [0, \infty)$$

$$\text{or } -1 + 4e^{-t} = 0$$

$$\Rightarrow e^{-t} = \frac{1}{4} \Rightarrow t = \log_e 4$$

Calculating  $f''(t)$

$$f''(t) = e^{-t} - 8e^{-2t} = e^{-t}(1 - 8e^{-t})$$

At  $t = \log_e 4$ ,

$$f''(t) = e^{-\log_e 4}(1 - 8e^{-\log_e 4}) = \frac{1}{4}\left(1 - \frac{8}{4}\right)$$

$$\Rightarrow f''(t) = -\frac{1}{4} < 0$$

Since at stationary point  $t = \log_e 4$ ,  $f''(t)$  is less than 0,

$$\therefore f(t) \text{ has maximum at } t = \log_e 4$$



Scan for Video solution



**97. (b)****Given:**

$$f(x) = 2x^3 - 9x^2 + 12x - 3 ; 0 \leq x \leq 3 \text{ i.e. } x \in [0, 3]$$

Calculating  $f'(x)$ ,

$$f'(x) = 6x^2 - 18x + 12$$

Stationary point: [At stationary point  $f'(x) = 0$ ]

$$f'(x) = 0$$

$$\Rightarrow 6x^2 - 18x + 12 = 0$$

$$\Rightarrow x^2 - 3x + 2 = 0$$

$$\Rightarrow (x-1)(x-2) = 0$$

$$\Rightarrow x = 1, 2 \in [0, 3]$$

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = 0$  and 3) and in stationary points (i.e. at  $x = 1$  and 2) :

$$f(x) = 2x^3 - 9x^2 + 12x - 3$$

At  $x = 0$ ,

$$f(0) = 2(0)^3 - 9(0)^2 + 12(0) - 3 = -3$$

At  $x = 3$ ,

$$f(3) = 2(3)^3 - 9(3)^2 + 12(3) - 3 = 6$$

At  $x = 1$ ,

$$f(1) = 2(1)^3 - 9(1)^2 + 12(1) - 3 = 2$$

At  $x = 2$ ,

$$f(2) = 2(2)^3 - 9(2)^2 + 12(2) - 3$$

∴ Maximum of  $f(x)$  in  $[0, 3]$ 

$$= \text{maximum } \{f(0), f(1), f(2), f(3)\}$$

$$= \text{maximum } \{-3, 2, 1, 6\} = 6$$

Maximum of  $f(x)$  in  $[0, 3] = 6$ 

Scan for Video solution

**98. (d)**

$$\sum_{n=0}^{\infty} \frac{1}{n!} = \frac{1}{0!} + \frac{1}{1!} + \frac{1}{2!} + \dots \infty$$

$$\Rightarrow \sum_{n=0}^{\infty} \frac{1}{n!} = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots \infty \quad \dots(i)$$

Now, expansion of  $e^x$ ,

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \infty$$

For  $x = 1$ ,

$$e = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots \infty \quad \dots(ii)$$

Comparing equation (i) and (ii),

$$\sum_{n=0}^{\infty} \frac{1}{n!} = e$$

Scan for Video solution

**99. (a)**Expansion of  $\sin x$  is:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

And expansion of  $\cos x$  is:

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

∴ expansion of  $3 \sin x + 2 \cos x$  is:


$$\begin{aligned} &= \left[ 3 \left( x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \right) \right] \\ &= \left[ +2 \left( 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \right) \right] \\ &= 2 + 3x - x^2 - \frac{x^3}{2} + \dots \end{aligned}$$

$$\therefore 3 \sin x + 2 \cos x = 2 + 3x - x^2 - \frac{x^3}{2} + \dots$$

Scan for Video solution

**100. (c)**

Given sum of length of a side and hypotenuse is constant.



$$\Rightarrow a + a \cos \theta = \text{constant} = c$$

$$\Rightarrow a(1 + \cos \theta) = c$$

$$\Rightarrow a = \frac{c}{(1 + \cos \theta)} \quad \dots(i)$$

$\therefore$  Area of triangle (A)

$$A = \frac{1}{2} \times a \cos \theta \times a \sin \theta$$

$$A = \frac{a^2}{2} \times \sin \theta \cos \theta$$

$$A = \frac{a^2}{4} \times \sin 2\theta$$

From equation (i),

$$A = \frac{c^2}{4(1 + \cos \theta)^2} \cdot \sin 2\theta$$

For area to be maximum,

$$\frac{d}{d\theta}(A) = 0$$

$$\frac{d}{d\theta}\left(\frac{c^2 \cdot \sin 2\theta}{4(1 + \cos \theta)^2}\right) = 0$$

$$\Rightarrow \frac{d}{d\theta}\left(\frac{\sin 2\theta}{(1 + \cos \theta)^2}\right) = 0$$

$$\Rightarrow (1 + \cos \theta)^2 \cdot 2 \cdot \cos 2\theta - \sin 2\theta \cdot 2(1 + \cos \theta) \cdot (-\sin \theta) = 0$$

$$\Rightarrow 2(1 + \cos \theta)\{\cos 2\theta(1 + \cos \theta) + \sin \theta \cdot \sin 2\theta\} = 0$$

$$\Rightarrow 2(1 + \cos \theta)\{\cos 2\theta + \cos 2\theta \cos \theta + \sin 2\theta \cdot \sin \theta\} = 0$$

$$\Rightarrow 2(1 + \cos \theta)\{\cos 2\theta + \cos \theta\} = 0$$

$$\Rightarrow \cos 2\theta + \cos \theta = 0$$

$$\Rightarrow 2\cos^2 \theta - 1 + \cos \theta = 0$$

$$\Rightarrow \cos \theta = 0.5, -1$$

$$\Rightarrow \theta = 60^\circ, 180^\circ$$

$\therefore$  For area to be maximum, angle between sides =  $60^\circ$ .



Scan for Video solution



### 101. (-1.00 to -0.94)


Let  $f(x) = e^{-x} \cdot \sin 100x$

We have for  $x > 0$ ,

$$-1 \leq \sin(100x) \leq 1$$

$$\Rightarrow -e^{-x} \leq e^{-x} \cdot \sin(100x) \leq e^{-x}$$

The graph of  $e^{-x} \cdot \sin(100x)$  is as shown



$\therefore$  Minimum value of  $e^{-x} \cdot \sin(100x)$  is at

$$100x = \frac{3\pi}{2} \Rightarrow x = \frac{3\pi}{200}$$

$$\begin{aligned} \Rightarrow \text{Min value} &= e^{-\left(\frac{3\pi}{200}\right)} \cdot \sin\left(100 \times \frac{3\pi}{200}\right) \\ &= e^{-\frac{3\pi}{200}} \cdot \sin\left(\frac{3\pi}{2}\right) = -0.9539 \end{aligned}$$

$\therefore$  Best approximate minimum value = -0.954



Scan for Video solution



### 102. (2)


$$\text{We know, } v = \frac{dx}{dt}$$

$\therefore$  The displacement of particle  $\int_0^x dx = \int_0^t v dt$

$$x = \int_0^t \frac{\pi}{2} \cos\left(\frac{\pi}{2} t\right) dt$$

$$\Rightarrow x = \left[ \sin \frac{\pi}{2} t \right]_0^t$$

$$\Rightarrow x = \sin \frac{\pi}{2} t$$


  
 $t = 0$

$\therefore$  At  $t = 0, x = 0$

$t = 1, x = 1$

$t = 2, x = 0$

$t = 3, x = -1$

In 3 sec, distance covered = 3 units

In 3 sec, displacement is = -1 units

$\therefore$  Required difference =  $3 - |-1| = 2$  units

Scan for Video solution



### 103. (b)

$$f(x) = x^3 - 3x^2 - 24x + 100$$

$f(x)$  is continuous and differential function in interval  $[-3, 3]$ .

Stationary points:

$$f'(x) = 0$$

$$\Rightarrow 3x^2 - 6x - 24 = 0$$

$$\Rightarrow x^2 - 2x - 8 = 0$$

$$\Rightarrow (x+2)(x-4) = 0$$

$$\Rightarrow x = -2, 4$$

$$\Rightarrow x = -2 \in [-3, 3] \text{ & } x = 4 \notin [-3, 3]$$

Hence stationary point in interval  $[-3, 3]$  is  $x = -2$ .

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = -3$  and 3) in stationary points (i.e. at  $x = -2$ ):


$$f(x) = x^3 - 3x^2 - 24x + 100$$

$$\Rightarrow f(-3) = (-3)^3 - 3(-3)^2 - 24(-3) + 100 = 118,$$

$$\Rightarrow f(3) = 3^3 - 3(3)^2 - 24(3) + 100 = 28, \leftarrow \text{minimum}$$

$$\Rightarrow f(-2) = (-2)^3 - 3(-2)^2 - 24(-2) + 100 = 128,$$

Hence minimum value of  $f(x)$  in interval  $[-3, 3] = 28$



### 104. (c)

$$f(x) = (x-1)^{2/3}$$

$f(x)$  is continuous function.

Differentiating above equation with respect to  $x$ ,

$$f'(x) = \frac{2}{3(x-1)^{1/3}} \quad \dots(i)$$

$f(x)$  is not differential at point where  $x-1=0$  i.e.  $x=1$ .

$f'(x)$  is never zero.

Here  $x=1$  is critical point.

Now, when  $x < 1$ ,  $f'(x) < 0$  and when  $x > 1$ ,  $f'(x) > 0$ . Therefore, in interval  $(-\infty, 1)$   $f(x)$  is decreasing function and in interval  $(1, \infty)$   $f(x)$  is increasing function.

$\therefore$  at  $x = 1$   $f(x)$  has minimum.

Scan for Video solution



### 105. (a)

$$f(x) = xe^{-x}$$

On Differentiating,

$$f'(x) = -x.e^{-x} + e^{-x}.1$$

$$\Rightarrow f'(x) = e^{-x}(1-x) \quad \dots(i)$$

Stationary point:

$$f'(x) = 0$$

$$\Rightarrow e^{-x}(1-x) = 0$$

$$\Rightarrow x = 1$$

Differentiating equation (i),

$$\begin{aligned} f''(x) &= \frac{d}{dx} \left\{ e^{-x} (1-x) \right\} \\ \Rightarrow f''(x) &= -e^{-x} (1-x) + e^{-x} (-1) \\ \Rightarrow f''(x) &= -e^{-x} (1-x+1) \\ \Rightarrow f''(x) &= -e^{-x} (2-x) \end{aligned}$$

At  $x = 1$

$$f''(1) = -e^{-1} (2-1) = -\frac{1}{e} < 0$$

Hence  $x = 1$  is maxima.

Hence maximum value of  $f(x)$

$$= f(1) = e^{-1}$$

Scan for Video solution



### 106. (b)

**Given:**  $y = f(x) = 5x^2 + 10x$  ... (i)

Since  $y$  is continuous in interval  $[1, 2]$  & is differentiable in interval  $(1, 2)$ .

Thus, both the condition of Lagrange's Mean Value Theorem is satisfying, therefore there exist at least one value  $c \in (1, 2)$  such that

$$f'(c) = \frac{f(b) - f(a)}{b - a} \quad \dots \text{(ii)}$$

Here,  $a = 1$  and  $b = 2$ , from equation (ii)

$$\Rightarrow f'(c) = \frac{f(2) - f(1)}{2 - 1} \quad \dots \text{(iii)}$$

Now, from equation (i);

$$f(1) = 5(1)^2 + 10(1) = 15$$

$$f(2) = 5(2)^2 + 10(2) = 40$$

Put value of  $f(1)$  and  $f(2)$  in equation (iii),

$$\therefore f'(c) = \frac{40 - 15}{1} = 25$$

Hence in interval  $(1, 2)$  atleast point will exist,

where  $\boxed{\frac{dy}{dx} = f'(c) = 25}.$

Scan for Video solution



### 107. (c)

Given error in  $\sin \theta \approx \theta$  holds is  $< 10\%$ .

Let ' $\theta$ ' be the value with in which error is less than 10%.

For small values of  $\theta$ ,  $\sin \theta \approx \theta - \frac{\theta^3}{3!}$

$$\therefore \frac{\sin \theta - \left( \theta - \frac{\theta^3}{3!} \right)}{\sin \theta} \times 100 < 10$$

$$\Rightarrow \sin \theta - \theta + \frac{\theta^3}{3!} < 0.1 \sin \theta$$

$$\Rightarrow \frac{\sin \theta}{\theta} - 1 + \frac{\theta^2}{3!} < 0.1 \frac{\sin \theta}{\theta}$$

$$\text{As } \theta \text{ is small, } \theta \rightarrow 0 \Rightarrow \frac{\sin \theta}{\theta} \rightarrow 1$$

$$\Rightarrow 1 - 1 + \frac{\theta^2}{6} < 0.1(1)$$

$$\Rightarrow \theta^2 < 0.6 \Rightarrow \theta < \sqrt{0.6}$$

$\therefore$  ' $\theta$ ' value should be less than  $\sqrt{0.6} \cong 0.7746$

$\therefore \theta \cong 44.38^\circ$  (Close to  $50^\circ$ ).

$\therefore$  For error to be less than 10%,  $\theta \leq 50^\circ$ .

Scan for Video solution



### 108. (c)

**Given:**

$$f(x) = x^3 - 9x^2 + 24x + 5 \quad \dots \text{(i)}$$

Differentiate equation (i) with respect to  $x$ ,

$$f'(x) = 3x^2 - 18x + 24$$

Stationary point: [At stationary point  $f'(x) = 0$ ]

$$f'(x) = 0$$

$$\Rightarrow 3x^2 - 18x + 24 = 0$$

$$\Rightarrow x^2 - 6x + 8 = 0$$

$$\Rightarrow (x - 2)(x - 4) = 0$$

$$\Rightarrow x = 2 \text{ (or) } x = 4 \in [1, 6]$$

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = 1$  and  $6$ ) and in stationary points (i.e. at  $x = 2$  and  $4$ ):

$$f(x) = x^3 - 9x^2 + 24x + 5$$

At  $x = 1$ ,

$$f(1) = (1)^3 - 9(1)^2 + 24(1) + 5 = 21$$

At  $x = 6$ ,

$$f(6) = (6)^3 - 9(6)^2 + 24(6) + 5 = 41$$

At  $x = 2$ ,

$$f(2) = (2)^3 - 9(2)^2 + 24(2) + 5 = 25$$

At  $x = 4$ ,

$$f(4) = (4)^3 - 9(4)^2 + 24(4) + 5 = 25$$

Maximum of  $f(x)$  in  $[1, 6] = \max\{f(2), f(4), f(1), f(6)\}$

$$= \max\{21, 25, 21, 41\} = 41$$

$\therefore$  Maximum of  $x^3 - 9x^2 + 24x + 5$  in  $[1, 6]$  is 41




Scan for Video solution



### 109. (b)

Given  $f(x) = \sin x$ ;  $x \in \left[\frac{\pi}{4}, \frac{7\pi}{4}\right]$



$\therefore f(x)$  has one minima at  $x = \frac{3\pi}{2}$



Scan for Video solution



### 110.(d)

Given  $f(x) = x^3 + 1$


$$\Rightarrow f'(x) = 3x^2$$

For stationary points,

$$f'(x) = 3x^2 = 0$$

$\Rightarrow x = 0$  is a stationary point.

$f'(x)$  on number line:



$\therefore f'(x)$  is not changing its sign at  $x = 0$ ,


$\therefore f(x)$  at  $x = 0$  is neither local maximum nor minimum.

Now,

$$f''(x) = \frac{d}{dx}(3x^2) = 6x$$

$$\text{at } x = 0, f''(x) = 0$$

$f''(x) = 6x$  on number line.



$\therefore f''(x)$  is changing its sign at  $x = 0$ , hence

$f(x) = x^3 + 1$  has point of inflection at  $x = 0$ .



Scan for Video solution



### 111. (c)

$$f(x) = x^3 - 9x^2 + 24x + 5 \quad \dots(i)$$

$f(x)$  is continuous and differentiable function in interval  $[1, 6]$ .

Differentiate equation (i) with respect to  $x$ ,

$$f'(x) = 3x^2 - 18x + 24$$

Stationary point: [At stationary point  $f'(x) = 0$ ]

$$f'(x) = 3x^2 - 18x + 24 = 0$$

$$\Rightarrow x^2 - 6x + 8 = 0$$

$$\Rightarrow (x - 2)(x - 4) = 0$$

$$\Rightarrow x = 2, 4 \in [1, 6]$$

Finding value of  $f(x)$  at initial and final point (i.e. at  $x = 1$  and 6) in stationary points (i.e. at  $x = 2$  and 4):

$$f(x) = x^3 - 9x^2 + 24x + 5$$

At  $x = 1$ :

$$f(1) = 1^3 - 9(1)^2 + 24(1) + 5 = 21$$

At  $x = 6$ :

$$f(6) = 6^3 - 9(6)^2 + 24(6) + 5 = 41 \leftarrow \text{maximum}$$

At  $x = 2$ :

$$f(2) = 2^3 - 9(2)^2 + 24(2) + 5 = 25$$

At  $x = 4$ :

$$f(4) = 4^3 - 9(4)^2 + 24(4) + 5 = 21$$

Values of  $f(x)$  at boundaries

**∴ Maximum value of  $f(x)$  occurs at  $x = 6$  and is equal to 41.**



Scan for Video solution



### Useful series

$$\cos \theta = 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \dots \infty$$

$$e^\theta = 1 + \frac{\theta}{1!} + \frac{\theta^2}{2!} + \frac{\theta^3}{3!} + \dots \infty$$



Scan for Video solution



### 112. (c)

$$f(x) = 2x - x^2 + 3 \quad \dots(i)$$

Differentiate equation (i) "ith respect to x,

$$f'(x) = 2 - 2x \quad \dots(ii)$$

Stationary point: '[At stationar' point  $f'(x) = 0]$

$$f'(x) = 2 - 2x = 0$$

$$\Rightarrow x = 1$$

Differentiate equation (ii) "ith respect to x,

$$f''(x) = -2$$

At  $x = 1$ :

$$f''(x) = -2 < 0$$

Since at st“tionary point  $x = 1$ ,  $f''(x) < 0$

**∴ At  $x = 1$ ,  $f(x)$  is a maximum**



Scan for Video solution



### 113. (c)

By McLaurin's series, expansion of  $e^x$  about  $x = 0$  is

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$



Scan for Video solution



### 114.(b)

**Given:**  $f(\theta) = \sin \theta$


The general Taylor series expansion of  $\sin \theta$  about  $\theta = 0$  is given by

$$\sin \theta = \theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \frac{\theta^7}{7!} + \dots \infty$$

### 115. (d)

$$f(t) = \frac{\sin t}{t}$$

From the graph of  $\frac{\sin t}{t}$



we can identify that maximum value of function lies at  $t$  tends to zero.

**Note: The maximum of the function can't be determined from the standard maxima and minima concept since the point  $x=0$  is a critical point.**



Scan for Video solution



### 116. (a)

Given  $f(x, y) = 4x^2 + 6y^2 - 8x - 4y + 8$

$$p = \frac{\partial f}{\partial x} = 8x - 8$$

$$q = \frac{\partial f}{\partial y} = 12y - 4$$

$$r = \frac{\partial^2 f}{\partial x^2} = 8$$

$$s = \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right) = \frac{\partial}{\partial x} (12y - 4) = 0$$

$$t = \frac{\partial^2 f}{\partial y^2} = 12$$

For stationary point:  $\frac{\partial f}{\partial x} = 0$  and  $\frac{\partial f}{\partial y} = 0$

$$\frac{\partial f}{\partial x} = 0 \Rightarrow 8x - 8 = 0 \Rightarrow x = 1$$

$$\frac{\partial f}{\partial y} = 0 \Rightarrow 12y - 4 = 0 \Rightarrow y = 1/3$$

Hence stationary point is  $(1, 1/3)$

Now,

$$rt - s^2 = 8 \times 12 - 0^2 = 96 > 0$$

and  $r = 8 > 0$

Since  $rt - s^2 > 0$  and  $r > 0$ , therefore stationary point  $(1, 1/3)$  is minima. And at stationary point  $(1, 1/3)$  function takes minimum value:

$$f_{\min} = 4x^2 + 6y^2 - 8x - 4y + 8$$

$$f_{\min} = 4(1)^2 + 6\left(\frac{1}{3}\right)^2 - 8(1) - 4\left(\frac{1}{3}\right) + 8$$

$$f_{\min} = 4 + \frac{6}{9} - 8 - \frac{4}{3} + 8 = 4 - \frac{2}{3}$$

$$f_{\min} = \frac{10}{3}$$



Scan for Video solution



Since at stationary point  $x = e$ ,  $y''$  is less than 0,

$\therefore y = \ln x^{1/x}$  has maximum at  $x = e$ .



Scan for Video solution



### 118.(c)

The Taylor series expansion of  $f(x) = \sin x$  at  $x = a$  is

$$\sin x = (x - a) - \frac{(x - a)^3}{3!} + \frac{(x - a)^5}{5!} - \frac{(x - a)^7}{7!} + \dots$$

Hence expansion of  $f(x) = \sin x$  at  $x = \pi$  is:

$$\sin x = (x - \pi) - \frac{(x - \pi)^3}{3!} + \frac{(x - \pi)^5}{5!} - \frac{(x - \pi)^7}{7!} + \dots$$

Divide both side by  $x - \pi$ ,

$$\frac{\sin x}{x - \pi} = 1 - \frac{(x - \pi)^2}{3!} + \frac{(x - \pi)^4}{5!} - \frac{(x - \pi)^6}{7!} + \dots$$



Scan for Video solution



Scan for Video solution



### 117.(a)

$$\text{Given: } e^y = x^{1/x} \Rightarrow y = \ln x^{1/x} \Rightarrow y = \frac{1}{x} \ln x$$

Calculating  $y'$ ,

$$y' = \frac{1}{x} \left( \frac{1}{x} \right) + (\ln x) \left( \frac{-1}{x^2} \right) \Rightarrow y' = \frac{1 - \ln x}{x^2}$$

At stationary point:

$$y' = 0 \Rightarrow \frac{1 - \ln x}{x^2} = 0 \Rightarrow 1 - \ln x = 0$$

$$\Rightarrow x = e$$

Calculating  $y''$

$$y'' = \frac{x^2 \left( \frac{-1}{x} \right) - (1 - \ln x) 2x}{x^4} = \frac{-x - 2x + 2x \ln x}{x^4}$$


$$\Rightarrow y'' = \frac{-3x + 2x \ln x}{x^4}$$

At  $x = e$ ,

$$\Rightarrow y''|_{x=e} = \frac{-3e + 2e \ln e}{e^4} = \frac{-e}{e^4} < 0$$

### 119.(a)

$$\text{Given: surface } z^2 = 1 + xy$$



Let 'P' be the nearest point on  $z^2 = 1 + xy$  to origin O.

For OP to be minimum,  $OP^2$  should be minimum.

$$OP^2 = (x - 0)^2 + (y - 0)^2 + (z - 0)^2$$

$$\Rightarrow OP^2 = x^2 + y^2 + z^2 = x^2 + y^2 + 1 + xy$$

For  $OP^2$  to be minimum,

$$x^2 + y^2 + 1 + xy \text{ should be minimum}$$

Let  $f = x^2 + y^2 + 1 + xy$

Stationary point;

$$p = \frac{\partial f}{\partial x} = 2x + y = 0 \quad \dots(i)$$

$$q = \frac{\partial f}{\partial y} = 2y + x = 0 \quad \dots(ii)$$

Solving equation (i) and (ii),

$$x=0, y=0$$

$\Rightarrow (0, 0)$  is stationary point

$$\text{Now } r = \frac{\partial^2 f}{\partial x^2} = 2; s = \frac{\partial^2 f}{\partial x \partial y} = 1; t = \frac{\partial^2 f}{\partial y^2} = 2$$

Since  $r > 0$  and  $rt - s^2 = 2(2) - 1^2 > 0$

$\therefore f$  has minimum at  $(0, 0)$

$$\therefore \text{Minimum of OP}^2 = 0^2 + 0^2 + 1 + (0, 0) = 1$$

$$\therefore \text{Minimum of OP} = \sqrt{1} = 1$$



Scan for Video solution



### 120. (d)

$$f(x) = \frac{\sin x}{x - \pi}$$

$$\Rightarrow f(x) = \frac{\sin(\pi + x - \pi)}{x - \pi} \quad (\because \sin(\pi + \theta) = -\sin \theta)$$

$$\Rightarrow f(x) = -\frac{\sin(x - \pi)}{(x - \pi)}$$

$$\left( \because \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \infty \right)$$

Therefore,

$$\Rightarrow f(x) = -\frac{1}{(x - \pi)} \left[ (x - \pi) - \frac{(x - \pi)^3}{3!} + \dots \right]$$

$$\Rightarrow f(x) = -1 + \frac{(x - \pi)^2}{3!} - \dots$$



Scan for Video solution



### 121. (b)

**123. (a)**

Given:  $f(x) = e^x + e^{-x}$  ... (i)

**Method I:**

For minimum value of  $f(x)$ ,  $f'(x) = 0$  and  $f''(x) > 0$

Differentiating equation (i) with respect to  $x$ ,

$$f'(x) = e^x - e^{-x} \quad \dots (\text{ii})$$

At stationary points:

$$f'(x) = 0$$

$$\Rightarrow e^x - e^{-x} = 0$$

$$\Rightarrow (e^x)^2 - 1 = 0$$

$$\Rightarrow e^x = 1$$

$$\Rightarrow x = 0$$

Now, Differentiating equation (ii) with respect to  $x$ ,

$$f''(x) = e^x + e^{-x} \quad \dots (\text{iii})$$

At stationary point  $x = 0$ ,

$$f''(0) = e^0 + e^{-0} = 2 > 0$$

Since at stationary point  $x = 0$ ,  $f''(x)$  is greater than 0,  $\therefore f(x)$  has minimum at  $x = 0$

The minimum value of  $f(x)$  is  $e^0 + e^{-0} = 2$

**Method II:**

For any two positive numbers  $a, b$ ,

$$A.M \geq G.M$$

$$\Rightarrow \frac{e^x + e^{-x}}{2} \geq \sqrt{e^x e^{-x}}$$

$$\Rightarrow e^x + e^{-x} \geq 2$$

**⇒ Minimum value of  $f(x) = 2$**



Scan for Video solution

**124. (a)**

The Taylor series expansion of  $f(x)$  about  $x = a$  is

$$f(x) = f(a) + \frac{f'(a)}{1!}(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \dots \infty$$

For the expansion to have only odd powers of  $x$  coefficients of even powers of ' $x$ ' at  $x = 0$  should be zero

$$\Rightarrow f(a) = f''(a) = f'''(a) = \dots = 0$$

**Option A:**

$$f(x) = \sin(x^3)$$

$$\Rightarrow f'(x) = 3x^2 \cos(x^3)$$

$$\Rightarrow f''(x) = 3\{x^2(-\sin(x^3))3x^2 + \cos(x^3)2x\} \\ = -9x^4 \sin(x^3) + 6x \cos(x^3)$$

At  $x = 0$ ,

$$f(0) = \sin(0) = 0,$$

$$\Rightarrow f''(0) = 0$$

$$\therefore \sin(x^3) = x^3 - \frac{x^9}{3!} + \frac{x^{15}}{5!} - \frac{x^{21}}{7!} + \dots \infty$$

$$\sin(x^2) = x^2 - \frac{x^6}{3!} + \frac{x^{10}}{5!} + \dots \infty$$

$$\cos(x^3) = 1 - \frac{x^6}{2!} + \frac{x^{12}}{4!} + \dots \infty$$

$$\cos(x^2) = 1 - \frac{x^4}{2!} + \frac{x^8}{4!} + \dots \infty$$

$\therefore$  Option A where  $f(x) = \sin(x^3)$ , is having odd powers of ' $x$ '.

▶
Scan for Video solution

**125. (b)**

$$\text{Given } f(x) = 3x^4 - 16x^3 + 24x^2 + 37$$

$$\Rightarrow f'(x) = 12x^3 - 48x^2 + 48x$$

$$\Rightarrow f'(x) = 12(x^3 - 4x^2 + 4x)$$

$$\Rightarrow f'(x) = 12x(x^2 - 4x + 4)$$

$$\Rightarrow f'(x) = 12x(x - 2)^2$$

For existance of maximum (or) minima,

$$f(x) = 0$$


$$\Rightarrow 12x(x - 2)^2 = 0$$

$$\Rightarrow x = 0, x = 2, 2$$

$\therefore$  The function has distinct stationary points  $x = 0, x = 2$ .

Now find the sign of  $f'(x)$  around stationary point.

$f'(x)$  on number line:



Since around  $x = 0$   $f'(x)$  is changing its sign from negative to positive. Therefore,  $x = 0$  is total minima.

Since around  $x = 2$ ,  $f'(x)$  is not changing its sign. Therefore,  $x = 2$  is neither maxima nor minima.

$\therefore$  The function has extreme only at  $x = 0$

$\therefore$  Number of points of extrema = 1



Scan for Video solution



### 126. (c)

$$\text{Given: } f(x) = e^x$$

The coefficient of  $(x-a)^n$  in Taylor series expansion

$$\text{of } f(x) \text{ about } x = a \text{ is } \frac{f^n(a)}{n!}$$

$$\therefore \text{The coefficient of } (x-2)^4 = \frac{f^4(2)}{4!}$$

$$\Rightarrow f^1(x) = f^2(x) = f^3(x) = f^4(x) = e^x \quad (\because f(x) = e^x)$$

$$\therefore \text{Coefficient of } (x-2)^4 = \frac{e^x|_{x=2}}{4!} = \frac{e^2}{4!}$$



Scan for Video solution



### 127. (d)

$$f(x) = \frac{1}{x^2}$$
 is not defined at  $x = 0$ .

$$f(x) = \int_{-1}^{+1} \frac{1}{x^2} dx \quad [\because f(x) = f(-x)]$$

$$= 2 \int_0^1 \frac{1}{x^2} dx$$

$$= 2 \left[ -\frac{1}{x} \right]_0^1 = 2(-1 + \infty) = \infty$$

Divergent


Scan for Video solution



### 128. (b)

$$\text{Given} \quad \int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$$

$$I = \int_{-\infty}^{\infty} e^{-a(x+b)^2} dx$$

$$\text{Let } x + b = t \Rightarrow dx = dt$$

$$\text{L.L: } x \rightarrow -\infty, x + b = t \rightarrow -\infty,$$

$$\text{U.L: } x \rightarrow \infty, x + b = t \rightarrow \infty$$

$$\therefore I = \int_{-\infty}^{\infty} e^{-at^2} \cdot dt = \int_{-\infty}^{\infty} e^{-(\sqrt{a}t)^2} \cdot dt$$

$$\text{Let } \sqrt{a}t = v \Rightarrow \sqrt{a} \cdot dt = dv$$

$$\text{L.L: } t \rightarrow -\infty; \sqrt{a}t \rightarrow -\infty$$

$$\text{U.L: } t \rightarrow \infty; \sqrt{a}t \rightarrow \infty$$

$$\therefore I = \frac{1}{\sqrt{a}} \cdot \int_{-\infty}^{\infty} e^{-v^2} \cdot dv$$

$$\left[ \because \int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi} \Rightarrow \int_{-\infty}^{\infty} e^{-v^2} \cdot dv = \sqrt{\pi} \right]$$

$$\Rightarrow I = \frac{1}{\sqrt{a}} \times \sqrt{\pi} = \sqrt{\frac{\pi}{a}}$$

$$\therefore \int_{-\infty}^{\infty} e^{-a(x+b)^2} dx = \sqrt{\frac{\pi}{a}}$$


Scan for Video solution



**129. (a)**

$$f(x) = \int_0^x e^t(t-1)(t-2)dt$$

To identify the nature of the function,  $f'(x)$  has to be calculated.

Differentiating the above function on both sides with respect to  $x$ , by Newton – Leibnitz rule,

$$f'(x) = \left[ e^x (x-1)(x-2) \frac{dx}{dx} \right] - \left[ e^0 (0-1)(0-2) \frac{d0}{dx} \right]$$

$$f'(x) = e^x(x-1)(x-2)$$

For  $f(x)$  to be decreasing,  $f'(x) < 0$

$$\Rightarrow f'(x) = e^x(x-1)(x-2) < 0$$

Since  $e^x$  is always positive, therefore

$$\Rightarrow (x-1)(x-2) < 0 \Rightarrow x \in (1, 2)$$

Therefore,  $f(x)$  is decreasing in  $x \in (1, 2)$ .



Scan for Video solution



**130. (0)**

$$\int_{-1}^{+1} x e^{|x|} dx$$

$$\text{Let, } f(x) = x e^{|x|}$$

$$\text{Then, } f(-x) = -x e^{-|x|} \Rightarrow f(-x) = -x e^{|x|} = -f(x)$$

Since  $f(x) = f(-x)$ , hence  $f(x)$  is an odd function. For odd function:

$$\int_{-a}^{+a} f(x) dx = 0$$

$$\int_{-1}^{+1} x e^{|x|} dx = 0$$



Scan for Video solution



**131. (c)**


Given

$$f(x) = x^2 - 2x + 2 \quad \text{in } x \in [1, 3]$$

$$f'(x) = 2x - 2$$

$$f(3) = 9 - 6 + 2 = 5$$

$$f(1) = 1 - 2 + 2 = 1$$



By Lagrange's mean value theorem,

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$\Rightarrow 2c - 2 = \frac{5 - 1}{3 - 1} = 2$$

$$\Rightarrow c = 2$$

$\therefore x = 2$  is the point where tangent is parallel to the straight line joining  $f(1)$  and  $f(3)$ .



Scan for Video solution



**132. (a, c, d)**

option (a)

$$I = \frac{1}{\pi} \int_0^\pi \sin m\theta \sin n\theta d\theta$$

$$\left[ \because \cos(A \pm B) = \cos A \cos B \mp \sin A \sin B \right]$$

$$\left[ \therefore \cos(A - B) - \cos(A + B) = \frac{1}{2} \sin A \sin B \right]$$

$$I = \frac{1}{2\pi} \int_0^\pi \cos(m-n)\theta - \cos(m+n)\theta \cdot d\theta$$

$$I = \frac{1}{2\pi} \left[ \frac{\sin(m-n)\theta}{m-n} - \frac{\sin(m+n)\theta}{m+n} \right]_0^\pi = 0$$

option (b)

$$I = \frac{1}{2\pi} \int_{-\pi/2}^{\pi/2} \sin p\theta \sin q\theta d\theta$$

$$\left[ \because \cos(A \pm B) = \cos A \cos B \mp \sin A \sin B \right]$$

$$\therefore \cos(A - B) - \cos(A + B) = \frac{1}{2} \sin A \sin B$$

$$I = \frac{1}{4\pi} \int_{-\pi/2}^{\pi/2} \cos(p-q)\theta - \cos(p+q)\theta d\theta$$

$$I = \frac{2}{4\pi} \left[ \frac{\sin(p-q)\theta}{p-q} - \frac{\sin(p+q)\theta}{p+q} \right]_0^{\pi/2}$$

$\rightarrow$  will depend on p and q it will not be necessarily be 0.

option (c)

$$I = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sin p\theta \cos q\theta d\theta$$

$$\left[ \because \sin(A \pm B) = \sin A \cos B \pm \cos A \sin B \right]$$

$$\therefore \sin A \cos B = \frac{1}{2} \{ \sin(A+B) + \sin(A-B) \}$$

$$I = \frac{1}{4\pi} \int_{-\pi}^{\pi} \{ \sin(p+q)\theta + \sin(p-q)\theta \} d\theta$$

Let,  $f(\theta) = \sin(p+q)\theta + \sin(p-q)\theta$  then,

$$f(-\theta) = \sin \{(p+q)(-\theta)\} + \sin \{(p-q)(-\theta)\}$$

$$f(-\theta) = -\sin(p+q)(\theta) - \sin(p-q)(\theta)$$

$$f(-\theta) = -\{\sin(p+q)\theta + \sin(p-q)\theta\} = -f(\theta)$$

Hence  $f(\theta) = \sin(p+q)\theta + \sin(p-q)\theta$  is odd function, therefore

$$I = 0$$

option (d)

$$L = \lim_{\alpha \rightarrow \infty} \frac{1}{2\alpha} \int_{-\alpha}^{\alpha} \sin p\theta \sin q\theta d\theta$$

$$L = \lim_{\alpha \rightarrow \infty} \frac{1}{2\alpha} \int_{-\alpha}^{\alpha} \cos(p-q)\theta - \cos(p+q)\theta d\theta$$

$$L = \lim_{\alpha \rightarrow \infty} \frac{2}{2\alpha} \left[ \frac{\sin(p-q)\theta}{p-q} - \frac{\sin(p+q)\theta}{p+q} \right]_0^{\alpha}$$

$$L = \lim_{\alpha \rightarrow \infty} \left[ \frac{\sin(p-q)\alpha}{(p-q)\alpha} - \frac{\sin(p+q)\alpha}{(p+q)\alpha} \right]$$

$$\left[ \because \lim_{x \rightarrow \infty} \frac{\sin x}{x} = 0 \right] \quad L = 0$$



Scan for Video solution



### 133. (b)

Given:  $f(-1) = 0$  and  $|f'(x)| \leq 2$

$$\Rightarrow -2 \leq f'(x) \leq 2$$

$$\Rightarrow -2 dx \leq f'(x) dx \leq 2 dx$$

On integrating

$$\int -2 dx \leq \int f'(x) dx \leq \int 2 dx$$

$$\Rightarrow -2x + c_1 \leq f(x) \leq 2x + c_2 \dots (i)$$

$\therefore f(-1) = 0$ ; hence from equation (i),

$$\Rightarrow 0 \leq -2 + c_1 \Rightarrow c_1 \geq 2$$

$$\text{Or, } 2 + c_2 \leq 0 \Rightarrow c_2 \leq -2$$

$$\therefore -2x - 2 \leq f(x) \leq 2x + 2 \dots (i)$$

$$\text{Or, } |f(x)| \leq 2x + 2$$

$$\Rightarrow f(x) \leq |2x + 2| \Rightarrow f(x) \leq 2|x + 1|$$

Scan for Video solution



### 134. (2.090 to 2.104)

$$I = \int_1^e (x \ln x) dx = ?$$

$$I = \int_1^e (x \ln x) dx = \int_1^e (\ln x) \cdot x \cdot dx$$

$$I = (\ln x) \cdot \frac{x^2}{2} - \int \frac{1}{x} \cdot \frac{x^2}{2} dx \Big|_1^e$$

$$I = \frac{x^2}{2} \ln x - \int \frac{x}{2} dx \Big|_1^e$$

$$I = \frac{x^2}{2} \ln x - \frac{x^2}{4} \Big|_1^e$$

$$I = \left( \frac{e^2}{2} \times \ln e - \frac{e^2}{4} \right) - \left( 0 - \frac{1}{4} \right)$$

$$I = \frac{e^2}{2}(1) - \frac{e^2}{4} + \frac{1}{4} = \frac{e^2 + 1}{4} = 2.097$$

$$\therefore \int_1^e (x \cdot \ln x) dx = 2.097$$

Scan for Video solution



**135. (0.50)**

$$I = \int_{0.25}^{1.25} f(x) dx = ?$$

$f(x) = x - [x]$

Since  $[x] = \begin{cases} 0 & ; 0.25 \leq x < 1 \\ 1 & ; 1 \leq x < 2 \end{cases}$

Hence,

$$\Rightarrow f(x) = \begin{cases} x - 0 = x & 0.25 \leq x < 1 \\ x - 1 & 1 \leq x < 2 \end{cases}$$

$$I = \int_{0.25}^{1.25} f(x) dx$$

$$\Rightarrow I = \int_{0.25}^1 x dx + \int_1^{1.25} (x - 1) dx$$

$$\Rightarrow I = \left[ \frac{x^2}{2} \right]_{0.25}^1 + \left[ \frac{x^2}{2} - x \right]_1^{1.25}$$

$$\Rightarrow I = \left[ \frac{1^2}{2} - \frac{(0.25)^2}{2} \right] + \left[ \left( \frac{1.25^2}{2} - 1.25 \right) - \left( \frac{1^2}{2} - 1 \right) \right]$$

$$\Rightarrow I = 0.5$$

$$\therefore \int_{0.25}^{1.25} f(x) dx = 0.5$$



Scan for Video solution

**136. (a)**

Mean value theorem of integration states that if a function  $f(x)$  is continuous in  $[a, b]$  and differentiable in  $(a, b)$  then in the interval  $[a, b]$  at least one value ' $\xi$ ' exists, such that,

$$f(\xi) = \frac{\int_a^b f(x) dx}{(b-a)}$$

$$\int_a^b f(x) dx = (b-a) \cdot f(\xi)$$



Scan for Video solution

**137. (0.27 to 0.3)**

Given Integral is

$$I = \int_0^{\frac{\pi}{4}} x \cdot \cos(x^2) dx$$

$$\text{Let } x^2 = t \Rightarrow 2x \cdot dx = dt = x \cdot dx = \frac{1}{2} \cdot dt$$

$$\text{L.L } x^2 = t \Rightarrow 0^2 = t \Rightarrow t = 0$$

$$\text{U.L } x^2 = t \Rightarrow \left(\frac{\pi}{4}\right)^2 = t \Rightarrow t = \frac{\pi^2}{16}$$

$$\therefore I = \int_0^{\frac{\pi^2}{16}} \cos t \frac{dt}{2} = \frac{1}{2} \times \sin \left| \frac{\pi^2}{16} \right|$$

$$= \frac{1}{2} \times \left\{ \sin \left( \frac{\pi}{16} \right)^2 - \sin 0 \right\}$$

$$= \frac{1}{2} \times 0.5784 = 0.2892$$

$$\therefore \int_0^{\frac{\pi}{4}} x \cdot \cos(x^2) dx = 0.2892$$

Scan for Video solution

**138. (c)**

Given

$$f(x) = R \sin\left(\frac{\pi x}{2}\right) + S, \quad f'\left(\frac{1}{2}\right) = \sqrt{2}; \quad \int_0^1 f(x) dx = \frac{2R}{\pi}$$

Now,

$$f(x) = R \sin\left(\frac{\pi x}{2}\right) + S$$

$$\Rightarrow f'(x) = R \left(\frac{\pi}{2}\right) \cos\left(\frac{\pi x}{2}\right)$$

$$\text{Given. } f'\left(\frac{1}{2}\right) = \sqrt{2}; \text{ therefore,}$$

$$\Rightarrow f'\left(\frac{1}{2}\right) = R \frac{\pi}{2} \cos\left(\frac{\pi}{4}\right) = \sqrt{2}$$

$$\Rightarrow \frac{R\pi}{2\sqrt{2}} = \sqrt{2}$$

$$\Rightarrow R = \frac{4}{\pi}$$

$$\text{Hence, } f(x) = R \sin\left(\frac{\pi x}{2}\right) + S = \frac{4}{\pi} \sin\left(\frac{\pi x}{2}\right) + S$$

Now,

$$\Rightarrow \int_0^1 f(x) dx = \frac{2R}{\pi} dx$$

$$\Rightarrow \int_0^1 \left( \frac{4}{\pi} \sin\left(\frac{\pi x}{2}\right) + S \right) dx = \frac{2}{\pi} \left( \frac{4}{\pi} \right)$$

$$\Rightarrow \frac{4}{\pi} \left[ -\cos\left(\frac{\pi x}{2}\right) \right]_0^1 + S \cdot x \Big|_0^1 = \frac{8}{\pi^2}$$

$$\Rightarrow \frac{8}{\pi^2} \{ -0 - (-1) \} + S = \frac{8}{\pi^2}$$

$$\Rightarrow \frac{8}{\pi^2} + S = \frac{8}{\pi^2}$$

$$\Rightarrow S = 0$$

$$\therefore R = \frac{4}{\pi} \text{ and } S = 0$$



Scan for Video solution



### 139. (a)

$$I = \int_0^1 \frac{(\sin^{-1} x)^2}{\sqrt{1-x^2}} dx$$

$$\text{Let } \sin^{-1} x = t \Rightarrow \frac{1}{\sqrt{1-x^2}} dx = dt$$

$$\text{At } x=0; t=\sin^{-1} 0=0$$

$$\text{At } x=1; t=\sin^{-1} 1 = \frac{\pi}{2}$$

Hence,

$$I = \int_0^{\pi/2} t^2 dt = \left[ \frac{t^3}{3} \right]_0^{\pi/2}$$

$$I = \frac{(\pi/2)^3}{3} = \frac{\pi^3}{24}$$

$$\therefore \int_0^1 \frac{(\sin^{-1} x)^2}{\sqrt{1-x^2}} dx = \frac{\pi^3}{24}$$



Scan for Video solution



### 140. (b)

$$g(x) = \int f(x) dx$$

$$g(x) = \int e^{-x} - e^{-x} dx$$

$$g(x) = \int e^{-x} \cdot e^{-e^{-x}} \cdot dx \quad \dots(1)$$

$$\text{Let } e^{-x} = t \Rightarrow -e^{-x} dx = dt$$

Then, equation (i) will become,

$$g(x) = - \int e^{-t} dt$$

$$g(x) = - \left[ \frac{e^{-t}}{-1} \right] = e^{-t} \quad [ \because e^{-x} = t ]$$

$$g(x) = e^{-e^{-x}}$$



Scan for Video solution



### 141. (c)

Given,


$$x = \cos\left(\frac{\pi u}{2}\right) \text{ and } y = \sin\left(\frac{\pi u}{2}\right)$$

$$\text{and } 0 \leq u \leq 1 \Rightarrow 0 \leq \frac{\pi u}{2} \leq \frac{\pi}{2}$$

$$\Rightarrow x^2 + y^2 = \cos^2\left(\frac{\pi u}{2}\right) + \sin^2\left(\frac{\pi u}{2}\right) = 1$$

The curve is  $x^2 + y^2 = 1$

$$\Rightarrow y = \sqrt{1-x^2} \text{ between } \theta = 0 \text{ to } \theta = \frac{\pi}{2}$$



Revolving of  $x^2 + y^2 = 1$  about x-axis by  $360^\circ$  forms a hemisphere of radius  $r=1$ .

$$\therefore \text{Surface area} = 2\pi r^2 = 2\pi(1)^2 = 2\pi$$



Scan for Video solution



**142. (-1)**

$$\text{Given Integral is } I = \int_{\frac{1}{\pi}}^{\frac{2}{\pi}} \cos\left(\frac{1}{x}\right) dx$$

$$\text{Let } \frac{1}{x} = t \Rightarrow \frac{-1}{x^2} dx = dt \Rightarrow \frac{1}{x^2} dx = -dt$$

$$\text{Lower limit : } \frac{1}{x} = t \Rightarrow \frac{1}{\left(\frac{1}{\pi}\right)} = t \Rightarrow t = \pi$$

$$\text{Upper limit : } \frac{1}{x} = t \Rightarrow \frac{1}{\left(\frac{2}{\pi}\right)} = t \Rightarrow t = \frac{\pi}{2}$$

$$\therefore I = \int_{\frac{\pi}{2}}^{\frac{2}{\pi}} \cos t (-dt) = \int_{\frac{\pi}{2}}^{\pi} \cos t (dt)$$

$$\Rightarrow I = \sin t \Big|_{\frac{\pi}{2}}^{\frac{2}{\pi}} = -1$$

$$\therefore I = \int_{\frac{1}{\pi}}^{\frac{2}{\pi}} \cos\left(\frac{1}{x}\right) dx = -1$$



Scan for Video solution



**143. (a)**

Given,

$$a \cdot f(x) + b \cdot f\left(\frac{1}{x}\right) = \frac{1}{x} - 25 \quad \dots\dots(1)$$

(Replacing x with  $\frac{1}{x}$ )

$$\Rightarrow a \cdot f\left(\frac{1}{x}\right) + b \cdot f(x) = x - 25 \quad \dots\dots(2)$$

In equation 1 multiply a both side,

$$a^2 \cdot f(x) + ab \cdot f\left(\frac{1}{x}\right) = \frac{a}{x} - 25a \quad \dots\dots(3)$$

In equation 2 multiply b both side,

$$ab \left(\frac{1}{x}\right) + b^2 \cdot f(x) = bx - 25b \quad \dots\dots(4)$$

Equation 3 – equation 4,

$$\Rightarrow (a^2 - b^2)f(x) = \frac{a}{x} - bx + 25(b - a)$$

$$\Rightarrow f(x) = \frac{1}{(a^2 - b^2)} \left\{ \frac{a}{x} - bx + 25(b - a) \right\}$$

On integrating,

$$\Rightarrow \int_1^2 f(x) dx = \frac{1}{(a^2 - b^2)} \left\{ a \cdot \ln x - \frac{bx^2}{2} + 25(b - a)x \right\} \Big|_1^2$$

$$\Rightarrow \int_1^2 f(x) dx = \frac{1}{(a^2 - b^2)} \left[ a \cdot \ln 2 - \frac{b}{2}(3) + 25(b - a) \right]$$

$$\Rightarrow \int_1^2 f(x) dx = \frac{1}{(a^2 - b^2)} \left\{ a(\ln 2 - 25) + \frac{47}{2}b \right\}$$

Scan for Video solution



**144. (4)**

$$\text{Given } \int_0^{2x} |x \cdot \sin x| dx = k\pi.$$

|x. sinx| is not differentiable at the points where,

$$x \cdot \sin x = 0$$

$$\Rightarrow x = 0, \pm\pi, \pm 2\pi, \dots$$

$\therefore \pi$  is a value in  $[0, 2\pi]$  where  $|x \cdot \sin x|$  is not differentiable.

$$\therefore \int_0^{2\pi} |x \cdot \sin x| dx = \int_0^{\pi} |x \cdot \sin x| dx + \int_{\pi}^{2\pi} |x \cdot \sin x| dx$$

When  $0 < x < \pi$ ,  $x \cdot \sin x > 0 \Rightarrow |x \cdot \sin x| = x \cdot \sin x$

$$\pi < x < 2\pi; x \cdot \sin x < 0 \Rightarrow |x \cdot \sin x| = -x \cdot \sin x$$

$$\therefore \int_0^{2\pi} |x \cdot \sin x| dx = \int_0^{\pi} x \cdot \sin x dx + \int_{\pi}^{2\pi} -x \cdot \sin x dx$$

$$= x \cdot (-\cos x) + 1 \cdot \sin x \Big|_0^\pi - (-x \cos x + \sin x) \Big|_0^{2\pi}$$

$$= \pi(1) - (-2\pi(1) + 0 - (+\pi)) = 4\pi$$

$$\therefore \int_0^{2\pi} |x \cdot \sin x| dx = 4\pi - K\pi$$

$$\Rightarrow K = 4$$



Scan for Video solution



#### 145. (a)

Given integral is

$$I = \int_0^\pi x^2 \cdot \cos x dx$$

$$\left[ \because \int u \cdot v dx = u \int v dx - \int \left( u' \int v dx \right) dx \right]$$

here, consider  $u = x^2$  &  $v = \cos x$

$$I = x^2 \cdot \sin x \Big|_0^\pi - \int_0^\pi 2x \sin x dx$$

$$I = (0 - 0) - 2 \int_0^\pi x \sin x dx$$

$$\left[ \because \int u \cdot v dx = u \int v dx - \int \left( u' \int v dx \right) dx \right]$$

In  $\int_0^\pi 2x \sin x dx$ , consider  $u = x$  &  $v = \sin x$

$$I = -2 \left[ -x \cos x \Big|_0^\pi - \int_0^\pi (-\cos x) dx \right]$$

$$I = -2 \left[ (\pi - 0) + \sin x \Big|_0^\pi \right]$$

$$I = -2 [\pi + (0 - 0)] = -2\pi$$

$$\therefore \int_0^\pi x^2 \cdot \cos x dx = -2\pi$$



Scan for Video solution



#### 146. (15)

$$2 \leq x \leq 3; 1 \leq y \leq 2; 0 \leq z \leq 1$$

Therefore,

$$I = \int_{z=0}^{z=1} \int_{y=1}^{y=2} \int_{x=2}^{x=3} 8xyz dx dy dz$$

$$I = 8 \left[ \frac{x^2}{2} \right]_2^3 \left[ \frac{y^2}{2} \right]_1^2 \left[ \frac{z^2}{2} \right]_0^1$$

$$I = 8 \left( \frac{9-4}{2} \right) \left( \frac{4-1}{2} \right) \left( \frac{1-0}{2} \right)$$

$$I = 8 \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2}$$

$$I = 15$$



Scan for Video solution



#### 147. (b)

$$\text{Find: } I = \int_0^2 \frac{(x-1)^2 \cdot \sin(x-1)}{(x-1)^2 + \cos(x-1)} dx$$

$$I = \int_0^2 \frac{(x-1)^2 \cdot \sin(x-1)}{(x-1)^2 + \cos(x-1)} dx \quad \dots(i)$$

$$\left. \begin{aligned} \because \int_a^b f(x) dx &= \int_a^b f(a+b-x) dx \\ \text{Hence replacing } x \text{ by } 0+2-x \text{ in equation (i)} \end{aligned} \right\}$$

$$I = \int_0^2 \frac{(0+2-x-1)^2 \cdot \sin(0+2-x-1)}{(0+2-x-1)^2 + \cos(0+2-x-1)} dx$$

$$I = \int_0^2 \frac{(1-x)^2 \cdot \sin(1-x)}{(1-x)^2 + \cos(1-x)} dx$$

$$I = \int_0^2 \frac{(x-1)^2 \cdot \sin(-(x-1))}{(x-1)^2 + \cos(-(x-1))} dx$$

$$I = \int_0^2 \frac{(x-1)^2 \cdot \{-\sin(x-1)\}}{(x-1)^2 + \cos(x-1)} dx$$

$$I = - \int_0^2 \frac{(x-1)^2 \cdot \sin(x-1)}{(x-1)^2 + \cos(x-1)} dx \quad \dots(ii)$$

Adding equation (i) and (ii),

$$2I = \int_0^2 \frac{(x-1)^2 \cdot \sin(x-1)}{(x-1)^2 + \cos(x-1)} dx - \int_0^2 \frac{(x-1)^2 \sin(x-1)}{(x-1)^2 + \cos(x-1)} dx$$

$$2I = 0$$

$$I = 0$$

$$\therefore I = \boxed{\int_0^2 \frac{(x-1)^2 \cdot \sin(x-1)}{(x-1)^2 + \cos(x-1)} dx = 0}$$



Scan for Video solution



#### 148. (c)

**Find:**  $I = \int_1^e \sqrt{x} \ln(x) dx$

$$I = \int_1^e (\sqrt{x} \cdot \ln x) dx = \int_1^e (\ln x) \cdot \sqrt{x} dx$$

$$I = \left\{ \ln x \cdot \frac{2}{3} \cdot x^{3/2} - \int \frac{1}{x} \cdot \frac{2}{3} \cdot x^{3/2} dx \right|_1^e \right\}$$

$$I = \left\{ \ln x \cdot \frac{2}{3} \cdot x^{3/2} - \frac{2}{3} \int x^{1/2} dx \right|_1^e \right\}$$

$$I = \left. \frac{2}{3} \cdot \ln x \cdot x^{3/2} - \frac{2}{3} \times \frac{2}{3} \times x^{3/2} \right|_1^e$$

$$I = \begin{bmatrix} \left( \frac{2}{3} \cdot \ln e \cdot e^{3/2} - \frac{2}{3} \times \frac{2}{3} \times e^{3/2} \right) \\ - \left( \frac{2}{3} \cdot \ln 1 \cdot 1^{3/2} - \frac{2}{3} \times \frac{2}{3} \times 1^{3/2} \right) \end{bmatrix}$$

$$I = \left[ \left( \frac{2}{3} e^{3/2} - \frac{4}{9} e^{3/2} \right) - \left( 0 - \frac{4}{9} \right) \right]$$

$$I = \frac{2}{3} e^{3/2} + \frac{4}{9} = \frac{2}{9} \sqrt{e^3} + \frac{4}{9}$$

$$\therefore \boxed{\int_1^e \sqrt{x} \ln(x) dx = \frac{2}{9} \sqrt{e^3} + \frac{4}{9}}$$



Scan for Video solution



#### 149. (d)

$$\text{Let } I = \int_0^{\pi/2} \frac{\cos x + i \cdot \sin x}{\cos x - i \sin x} dx = \int_0^{\pi/2} \frac{e^{ix}}{e^{-ix}} dx$$

$$= \int_0^{\pi/2} e^{i(2x)} dx$$

$$= \left. \frac{e^{i2x}}{2i} \right|_{x=0}^{x=\pi/2} = \frac{1}{2i} \{ e^{i2(\pi/2)} - e^{i2(0)} \}$$

$$= \frac{1}{2i} \{-1 - 1\} = \frac{-2}{2i} = \frac{-1}{i} = i$$

$$\therefore \boxed{\int_0^{\pi/2} \frac{\cos x + i \cdot \sin x}{\cos x - i \sin x} dx = i}$$



Scan for Video solution



#### 150. (c)

$$I = \int_0^{\pi/6} \cos^4 3\theta \sin^3 6\theta d\theta$$

$$\text{Let } 3\theta = t, \text{ then } 3 d\theta = dt \Rightarrow d\theta = \frac{dt}{3}$$

$$\text{If } \theta = 0; t = 3\theta = 0$$

$$\text{If } \theta = \pi/6; t = 3\theta = \frac{\pi}{2}$$

$$I = \int_0^{\pi/2} \cos^4 t \sin^3 2t \frac{dt}{3}$$

$$I = \int_0^{\pi/2} \cos^4 t (2 \sin t \cos t)^3 \frac{dt}{3}$$

$$I = \frac{8}{3} \int_0^{\pi/2} (\sin^3 t \times \cos^7 t) dt$$

$$\left[ \begin{array}{l} ((m-1)(m-3)\dots 2 \text{ (or) } 1) \\ \because \int_0^{\pi/2} \sin^m t \cos^n t dt = \frac{((n-1)(n-3)\dots 2 \text{ (or) } 1)}{((m+n)(m+n-2)\dots 2 \text{ (or) } 1)} \times k \\ k = \begin{cases} \pi/2, & \text{when both m and n are even} \\ 1, & \text{otherwise} \end{cases} \end{array} \right]$$

$$I = \frac{8}{3} \left[ \frac{2 \times 6 \times 4 \times 2}{10 \times 8 \times 6 \times 4 \times 2} \times 1 \right] = \frac{1}{15}$$

$$\therefore \int_0^{\pi/6} \cos^4 3\theta \sin^3 6\theta d\theta = \frac{1}{15}$$



Scan for Video solution


**151. (c)**

$$\left. \begin{array}{l} \text{Let } I = \int_0^a \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} dx \quad \dots \text{(i)} \\ \left\{ \begin{array}{l} \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \\ \text{Hence replacing } x \text{ by } a-x \text{ in equation (i)} \end{array} \right. \end{array} \right\}$$

$$I = \int_0^a \frac{\sqrt{a-x}}{\sqrt{(a-x)} + \sqrt{a-(a-x)}} dx$$

$$I = \int_0^a \frac{\sqrt{a-x}}{\sqrt{a-x} + \sqrt{x}} dx \quad \dots \text{(ii)}$$

On adding (i) and (ii),

$$2I = \int_0^a \frac{\sqrt{x} + \sqrt{a-x}}{\sqrt{x} + \sqrt{a-x}} dx$$

$$2I = \int_0^a 1 dx = a$$

$$I = \frac{a}{2}$$

$$\therefore I = \int_0^a \frac{\sqrt{x}}{\sqrt{x} + \sqrt{a-x}} dx = \frac{a}{2}$$


**152. (MTA)**

$$\begin{aligned} & \int \left( x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \right) dx \\ &= \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{12} - \frac{x^5}{20} + \dots \end{aligned}$$

Checking all the options follows

$$\begin{aligned} \frac{1}{1+x} &= (1+x)^{-1} = (1-x+x^2-x^3+\dots\infty) \\ -\frac{1}{1-x} &= (1-x)^{-1} = -(1+x+x^2+x^3+\dots\infty) \\ \frac{1}{1+x^2} &= (1+x^2)^{-1} = (1-x^2+x^4-x^6+\dots\infty) \\ -\frac{1}{1-x^2} &= (1-x^2)^{-1} = (1+x^2+x^4+x^6+\dots\infty) \end{aligned}$$

All options a, b, c and d are incorrect. (Hence marks to all)



Scan for Video solution


**153. (b)**
**Given:**  $f(x)$  is even function:  $a > 0$ 

$$\Rightarrow f(-x) = f(x)$$


$$\text{For any } a > 0 \text{ if } f(-x) = f(x)$$

Then,

$$\int_{-a}^a f(x) dx = 2 \times \int_0^a f(x) dx$$

**Important point:** if  $f(-x) = -f(x)$ , then

$$\int_{-a}^a f(x) dx = 0$$



Scan for Video solution



**154. (b)**

$$P = \int_0^1 xe^x dx$$

$$\left[ \because \int u.v dx = u \int v dx - \int \left( u' \int v dx \right) dx \right]$$

here, consider  $u = x$  &  $v = e^x$

$$\Rightarrow P = \left[ x e^x \right]_0^1 - \int_0^1 1.e^x dx \lim_{x \rightarrow \infty}$$

$$\Rightarrow P = \left( (1 \times e^1) - 0 \right) - \left[ e^x \right]_0^1$$

$$\Rightarrow P = e - \left( (e^1 - e^0) \right)$$

$$\Rightarrow P = e - (e - 1)$$


$$\Rightarrow P = 1$$



Scan for Video solution



$$\therefore \int_{-\infty}^{\infty} \frac{dx}{1+x^2} = \pi$$

**156. (d)**

$$\text{Given integral is } I = \int_0^{\pi/4} \frac{1 - \tan x}{1 + \tan x} dx$$

$$\Rightarrow I = \int_0^{\pi/4} \frac{1 - \tan x}{1 + \tan x} dx = \int_0^{\pi/4} \frac{\tan \frac{\pi}{4} - \tan x}{1 + \tan \frac{\pi}{4} \times \tan x} dx$$

$$\left( \because \tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \cdot \tan B} \right)$$

$$\therefore I = \int_0^{\pi/4} \tan \left( \frac{\pi}{4} - x \right) dx$$


$$\left( \because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right)$$

$$\Rightarrow I = \int_0^{\pi/4} \tan \left( \frac{\pi}{4} - \left( 0 + \frac{\pi}{4} - x \right) \right) dx$$

$$\Rightarrow I = \int_0^{\pi/4} \tan x dx$$

$$\Rightarrow I = \log_e |\sec x|_0^{\pi/4} = \log_e \frac{\sqrt{2}}{1} = \frac{1}{2} \cdot \log_e 2$$

$$\therefore \int_0^{\pi/4} \frac{1 - \tan x}{1 + \tan x} dx = \frac{1}{2} \ln 2$$

**157. (a)**

$$3y=x$$

$$\int_0^3 \int_{y=0}^{x=3y} (6-x-y) dy dx$$

**155. (d)**

$$I = \int_{-\infty}^{\infty} \frac{dx}{1+x^2} = ?$$

$$I = \int_{-\infty}^{\infty} \frac{dx}{1+x^2}$$

$$\left\{ \begin{array}{l} \because \int_a^{-a} f(x) dx = 2 \times \int_0^a f(x) dx \quad \text{if } f(-x) = f(x) \end{array} \right.$$

$$\text{Let, } f(x) = \frac{1}{1+x^2}, \text{ then}$$

$$f(-x) = \frac{1}{1+(-x)^2} = \frac{1}{1+x^2} = f(x)$$

Since,  $f(-x) = f(x)$ , hence

$$I = \int_{-\infty}^{\infty} \frac{dx}{1+x^2} = 2 \times \int_0^{\infty} \frac{dx}{1+x^2}$$

$$I = 2 \times \tan^{-1} x \Big|_0^{\infty} = 2 \times (\tan^{-1} \infty - \tan^{-1} 0)$$

$$I = 2 \times \left( \frac{\pi}{2} - 0 \right) = \pi$$

$$\begin{aligned}
 &= \int_0^3 \left[ 6y - xy - \frac{y^2}{2} \right]_{y=0}^{y=x} dx \\
 &= \int_0^3 \left( 6x - x^2 - \frac{x^2}{2} \right) dx \\
 &= \int_0^3 6x - \frac{3x^2}{2} dx = \left[ \frac{6x^2}{2} - \frac{3}{2} \times \frac{x^3}{3} \right]_0^3 \\
 &= \left[ 3x^2 - \frac{x^3}{2} \right]_0^3 \\
 &= \left( 3 \times 3^2 - \frac{3^3}{2} \right) = 27 - \frac{27}{2} = \frac{27}{2} = 13.5
 \end{aligned}$$

$$\therefore \int_0^3 \int_{y=0}^{y=x} (6 - x - y) dy dx = 13.5$$



Scan for Video solution

**158.(d)**

$$(i) I = \int_0^{\pi/4} \tan x dx = \ln |\sec x| \Big|_0^{\pi/4}$$

$$I = \ln \sqrt{2} - \ln(1) = \ln \sqrt{2} \rightarrow \text{finite}$$

$$(ii) I = \int_0^{\infty} \frac{1}{x^2 + 1} dx = \tan^{-1} x \Big|_0^{\infty} = \frac{\pi}{2} - 0 = \frac{\pi}{2}$$

 $\rightarrow \text{finite}$ 

$$\begin{aligned}
 (iii) I &= \int_0^{\infty} x \cdot e^{-x} dx = \int_0^{\infty} e^{-x} \cdot x dx = \int_0^{\infty} e^{-x} \cdot x^{2-1} dx \\
 &\quad \left( \because \int_0^{\infty} e^{-x} \cdot x^{n-1} dx = n! \right)
 \end{aligned}$$

$$I = \sqrt{2} = 1! = 1 \rightarrow \text{finite}$$

$$(iv) I = \int_0^1 \frac{1}{1-x} dx = \frac{\ln(1-x)}{-1} \Big|_0^1 = -1(\ln 0 - \ln 1)$$

 $I = -\ln 0 \rightarrow \infty \rightarrow \text{infinite hence unbounded}$ 

$\therefore \int_0^1 \frac{1}{1-x} dx$  is unbounded



Scan for Video solution

**159. (a, b, d)**


$$|x_1|^p + |x_2|^p = 1 \text{ for } p > 0$$

From option a):

$$p = 2$$

$$\therefore |x_1|^2 + |x_2|^2 = 1$$

Above equation is equation of circle of radius 1 unit.



$$\text{Area of circle} = \pi r^2 = \pi \times 1^2 = \pi$$


Hence option (a) is correct.

From option b):

$$p \rightarrow \infty \quad |x_1|^p + |x_2|^p = 1$$

Case I: If  $|x_1| < 1$  then  $\lim_{p \rightarrow \infty} |x_1|^p \rightarrow 0$ 

$$\therefore |x_2| = 1 \Rightarrow x_2 = 1 \text{ or } x_2 = -1$$

Case II: If  $|x_2| < 1$  then  $\lim_{p \rightarrow \infty} |x_2|^p \rightarrow 0$ 

$$\therefore |x_1| = 1 \\ \Rightarrow x_1 = 1 \text{ or } x_1 = -1$$

When  $p \rightarrow \infty$ , the given function represents asquare bounded by the lines  $|x_1| = 1$  and  $|x_2| = 1$ .

Side of square = 2, therefore,

Area enclosed in this case

$$= 2 \times 2 = 4$$

From option c):

When  $p$  tends to 0, the function does not enclose any area.


**From option (d):**

$$p = 1$$

$$|x_1| + |x_2| = 1$$

The above equation represents a square bounded by the lines  $x_1 + x_2 = 1$ ,  $x_1 - x_2 = 1$ ,  $-x_1 + x_2 = 1$  and  $-x_1 - x_2 = 1$ .

Side of square =  $\sqrt{1^2 + 1^2} = \sqrt{2}$ , therefore,



$$\text{Area} = (\sqrt{2})^2 = 2$$

Hence option a, b and d are correct



Scan for Video solution



### 160. (18.80 to 18.90)

Given


$$\hat{x} = 3y, \hat{y} = 2x \Rightarrow y = \frac{\hat{x}}{3}; x = \frac{\hat{y}}{2}$$

The disc  $x^2 + y^2 < 1$  is given.

$$\Rightarrow x^2 + y^2 < 1$$

$$\Rightarrow \left(\frac{\hat{y}}{2}\right)^2 + \left(\frac{\hat{x}}{3}\right)^2 < 1$$

$$\Rightarrow \frac{\hat{x}^2}{3^2} + \frac{\hat{y}^2}{2^2} < 1 \rightarrow \text{Elliptic Region}$$



The area of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is  $\pi ab$

Area of required region =  $\pi (3)(2) = (3.14)6 = 18.84$



Scan for Video solution




### 161. (0)

Given:

Region is a shown and the integral is (Say),

$$I = \iint_R xy \, dx \, dy$$



For any region 'R' that is symmetric about x-axis (or) y-axis,

$$\iint_R xy \, dx \, dy = 0$$

Since given region 'R' is symmetric about y-axis,

$$\iint_R xy \, dx \, dy = 0$$




Scan for Video solution



### 162. (171)

Given integral is,

$$I = \iint_D 3(x^2 + y^2) \, dx \, dy$$



From the given diagram,

Equation of OA  $\Rightarrow y = x$

Equation of OB  $\Rightarrow y = -x$

Taking vertical strip RS in required region D.

Now for entire region the limit of y at any x will be,

Lower limit  $\rightarrow y = -x$  [At R]

Upper limit  $\rightarrow y = x$  [At S]

Now by sliding strip RS in horizontal direction from extreme left to right in required region D we get the limit of x as given below,

Lower limit  $\rightarrow x = 0$

Upper limit  $\rightarrow x = 4$

Therefore,

$$I = \int_{x=0}^{x=4} \int_{y=-x}^{y=x} 3(x^2 + y^2) dy dx$$

$$I = \int_{x=0}^{x=4} 3 \left( x^2 y + \frac{y^3}{3} \right) \Big|_{-x}^x dx$$

$$I = \int_{x=0}^{x=4} 3 \left( x^2 (2x) + \frac{2x^3}{3} \right) dx$$

$$I = \int_{x=0}^{x=4} 8x^3 dx = 2x^4 \Big|_0^4$$

$$I = 2(4^4 - 0^4) = 512$$


$$\therefore I = \iint_D 3(x^2 + y^2) dx dy = 512$$



Scan for Video solution



### 163. (b)



$$\text{Given, curve is } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\Rightarrow \frac{y^2}{b^2} = \frac{a^2 - x^2}{a^2}$$

$$\Rightarrow y^2 = b^2 \left( \frac{a^2 - x^2}{a^2} \right)$$

$$\Rightarrow y = \pm \frac{b}{a} \sqrt{a^2 - x^2}$$

Area of ellipse (A)

$$A = 4 \int_{x=0}^{x=a} \int_{y=0}^{y=\frac{b}{a}\sqrt{a^2-x^2}} dy dx$$

$$A = 4 \int_{x=0}^{x=a} \frac{b}{a} \sqrt{a^2 - x^2} dx = 4 \frac{b}{a} \int_{x=0}^{x=a} \sqrt{a^2 - x^2} dx$$

$$\text{Since, } \int \sqrt{(a^2 - x^2)} dx = \frac{x\sqrt{(a^2 - x^2)}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a},$$


hence

$$A = \frac{4b}{a} \left[ \frac{x\sqrt{(a^2 - x^2)}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \Big|_0^a \right]$$

$$A = \frac{4b}{a} \left[ \left( 0 + \frac{a^2}{2} \sin^{-1} 1 \right) - \left( 0 + \frac{a^2}{2} \sin^{-1} 0 \right) \right]$$

$$A = \frac{4b}{a} \left[ \left( \frac{a^2}{2} \times \frac{\pi}{2} \right) - (0 + 0) \right] = 4 \left( \frac{\pi ab}{4} \right) = \pi ab$$

$$\therefore \text{Area of ellipse} = \pi ab$$



Scan for Video solution



**164. (b)****Given**

$$y^2 = x; \text{ and } 0 \leq x \leq 1$$

Volume of solid obtained by revolving a curve

$y = f(x); a \leq x \leq b$  by  $360^\circ$  about x-axis is  $\int_a^b \pi y^2 dx$ .

$$\therefore \text{Volume} = \int_0^1 \pi y^2 dx = \int_0^1 \pi(x) dx = \pi \frac{x^2}{2} \Big|_0^1 = \frac{\pi}{2}$$

$$\therefore \text{Volume of solid generated} = \frac{\pi}{2}$$



Scan for Video solution

**165. (b)**

$$\text{Let } I = \int_0^\pi x \cos^2 x dx \quad \dots(i)$$

$$\left\{ \begin{array}{l} \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \\ \text{Hence replacing } x \text{ by } \pi - x \text{ in equation (i)} \end{array} \right\}$$

$$I = \int_0^\pi (\pi - x) \cos^2(\pi - x) dx$$

$$I = \int_0^\pi (\pi - x) \cos^2 x$$

$$I = \int_0^\pi \pi \cos^2 x dx - \int_0^\pi x \cos^2 x dx$$

$$I = \int_0^\pi \pi \cos^2 x dx - I$$

$$2I = \pi \int_0^\pi \cos^2 x dx$$

$$\left[ \begin{array}{l} \text{Here } \cos^2(\pi - x) = \cos^2 x \\ \& \text{When } f(x) = f(2a - x) \\ \Rightarrow \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx \end{array} \right]$$

$$2I = 2\pi \int_0^{\pi/2} \cos^2 x dx$$

$$I = \pi \int_0^{\pi/2} \cos^2 x dx = \pi \times \frac{1}{2} \times \frac{\pi}{2}$$

$$I = \frac{\pi^2}{4}$$

Scan for Video solution


**166. (85 to 85.5)****Given:** curves are  $y = 8$  ... (i)and  $x^2 = 8y$  ... (ii)

Intersection points of curve (i) and (ii):

From (i) putting  $y = 8$  in equation (ii),

$$x^2 = 8 \times 8$$

$$x = \pm 8$$


For  $x = -8$ , from equation (i);  $y = 8$ And for  $x = 8$ , from equation (i);  $y = 8$ 

Area bounded by curves

$$= \int_{x_1}^{x_2} y_1 - y_2 dx$$

$$= \int_{-8}^8 8 - \frac{x^2}{8} dx = \left[ 8x - \frac{x^3}{24} \right]_{-8}^{+8}$$

$$= 2 \left( 64 - \frac{64}{3} \right) = 85.33$$



Scan for Video solution

**167. (b)**curves are  $x + y = 3$  ... (i)and  $y = x^2 + 1$  ... (ii)

Intersection point of curve (i) and (ii):

From (i) putting  $y = 3 - x$  in equation (ii),

$$3-x = x^2 + 1$$

$$x^2 + x - 2 = 0$$


$$(x+2)(x-1) = 0$$

$$x = -2, 1$$

For  $x = -2$ , from equation (i);  $y = 0$

And for  $x = 1$ , from equation (i);  $y = 1$

$$y = x^2 + 1 \text{ and } x + y = 3$$



Taking vertical strip AB in required region R. Now for entire region the limit of y at any x will be,

Lower limit  $\rightarrow y = y_2 = x^2 + 1$  [At A, from the equation of curve (ii)]

Upper limit  $\rightarrow y = y_1 = 3 - x$  [At B, from the equation of curve (i)]

Now by sliding strip AB in horizontal direction from extreme left to right in required region R we get the limit of x as given below,

Lower limit  $\rightarrow x = -2$

Upper limit  $\rightarrow x = 1$

Hence required area (A),

$$A = \int_{x=-2}^{x=1} \int_{y=x^2+1}^{y=3-x} dy dx$$

$$A = \int_{x=-2}^{x=1} \left( \int_{y=x^2+1}^{y=3-x} dy \right) dx$$

$$A = \int_{x=-2}^{x=1} (3-x) - (x^2 + 1) dx$$

$$A = \int_{x=-2}^{x=1} (-x^2 - x + 2) dx = \left[ -\frac{x^3}{3} - \frac{x^2}{2} + 2x \right]_{-2}^1$$

$$A = \left( -\frac{1}{3} - \frac{1}{2} + 2 \right) - \left( \frac{8}{3} - 2 - 4 \right)$$

$$A = \frac{9}{2}$$

Scan for Video solution



### 168. (1.85 to 1.87)

Given

$$x(t) = \cos t; y(t) = \sin t; z(t) = \frac{2}{\pi}t \text{ and}$$

$$0 \leq t \leq \frac{\pi}{2}$$

$$\Rightarrow \frac{dx}{dt} = -\sin t; \frac{dy}{dt} = \cos t; \frac{dz}{dt} = \frac{2}{\pi}$$

The length of the parametric curve (L) is

$$L = \int_{t=0}^{t=\frac{\pi}{2}} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt$$

$$L = \int_{t=0}^{t=\frac{\pi}{2}} \sqrt{(-\sin t)^2 + (\cos t)^2 + \left(\frac{2}{\pi}\right)^2} dt$$

$$L = \int_{t=0}^{t=\frac{\pi}{2}} \sqrt{1 + \frac{4}{\pi^2}} dt = \sqrt{1 + \frac{4}{\pi^2}} \times \frac{\pi}{2} = 1.862 \text{ units}$$

$$\Rightarrow \text{Length of the curve} = 1.862 \text{ units}$$

Scan for Video solution



### 169. (a)

Given: curves are  $y = x$  ... (i)

and  $y = x^2$  ... (ii)

Intersection point of curve (i) and (ii):

From (i) putting  $y = x$  in equation (ii),

$$x = x^2$$

$$x(x-1) = 0$$


$$x = 0 \text{ and } x = 1$$

For  $x = 0$ , from equation (i);  $y = 0$

And for  $x = 1$ , from equation (i);  $y = 1$

Hence intersection points of curve (i) and (ii) are O  $(0, 0)$  and P  $(1, 1)$

The region R bounded by the curve  $y = x^2$  and  $y = x$  as shown in figure by shaded region R.



Taking vertical strip AB in required region R. Now for entire region the limit of y at any x will be,  
Lower limit  $\rightarrow y = x^2$  [At A, from the equation of parabola  $y = x^2$ ]  
Upper limit  $\rightarrow y = x$  [At B, from the equation of line  $y = x$ ]

Now by sliding strip AB in horizontal direction from extreme left to right in required region R we get the limit of x as given below,

Lower limit  $\rightarrow x = 0$

Upper limit  $\rightarrow x = 1$

Hence required area (A),

$$A = \int_{x=0}^{x=1} \int_{y=x^2}^{y=x} dy dx$$


$$A = \int_{x=0}^{x=1} \left( \int_{y=x^2}^{y=x} dy \right) dx$$

$$A = \int_{x=0}^{x=1} (x - x^2) dx$$

$$A = \frac{x^2}{2} - \frac{x^3}{3} \Big|_0^1$$

$$A = \frac{1}{2} - \frac{1}{3} = \frac{1}{6}$$

$$\therefore \text{Area bounded} = \frac{1}{6} \text{ units}$$



Scan for Video solution




### 170. (d)

The length of the curve  $f(x)$  from  $x = a$  to  $x = b$  is given by

$$L = \int_a^b \sqrt{1 + \left( \frac{dx}{dy} \right)^2} dx \quad \dots(i)$$

$$\text{Now, } y = 4h \frac{x^2}{L^2} \Rightarrow \frac{dy}{dx} = \frac{8hx}{L^2}$$



Here, A & B are supports. Put  $y = h$  in the equation of parabola to get co-ordinates of the supports

$$h = 4h \frac{x^2}{L^2} \Rightarrow x^2 = \frac{L^2}{4} \Rightarrow x = \pm \frac{L}{2}$$

$$\therefore A \rightarrow \left( -\frac{L}{2}, 0 \right) \text{ and } B \rightarrow \left( \frac{L}{2}, 0 \right)$$

Therefore, from equation (i),

$$\text{Length of cable} = \int_{-L/2}^{L/2} \sqrt{1 + \left( \frac{8hx}{L^2} \right)^2} dx$$

Since  $\sqrt{1 + \left( \frac{8hx}{L^2} \right)^2}$  is even function, therefore

$$\text{Length of cable} = 2 \int_0^{L/2} \sqrt{1 + \frac{64h^2 x^2}{L^4}} dx$$



Scan for Video solution




171. (d)

**Given:**  $y = \sqrt{x}$ ;  $1 \leq x \leq 2$

The volume of the solid obtained by revolving  $y = f(x)$  about x-axis between  $x = a$  &  $x = b$  is given by

$$V = \int_{x=a}^{x=b} \pi y^2 dx$$

$$V = \int_{x=1}^{x=2} \pi y^2 dx$$



$$V = \int_{x=1}^{x=2} \pi (\sqrt{x})^2 dx = \pi \cdot \frac{x^2}{2} \Big|_1^2$$

$$V = \frac{\pi}{2}(4-1) = \frac{3\pi}{2}$$

$$\therefore \text{Volume of solid generated} = \frac{3\pi}{2}$$



Scan for Video solution



172. (a)

**Given:** curves are

$$y^2 = 4x \quad \dots(i)$$

$$\text{and } x^2 = 4y \quad \dots(ii)$$

Intersection point of curve (i) and (ii):

From (i) putting  $x = \frac{y^2}{4}$  in equation (ii),

$$\left(\frac{y^2}{4}\right)^2 = 4y$$

$$y^4 - 64y = 0$$

$$y(y^3 - 64) = 0$$


$$y = 0, y = 4$$

For  $y = 0$ , from equation (i);  $x = 0$

And for  $y = 4$ , from equation (i);  $x = 4$

Hence intersection points of curve (i) and (ii) are O (0, 0) and P (4, 4)

The region R bounded by the curve  $y = x^2$  and  $y = x$  is shown in figure.



Taking vertical strip AB in required region R. Now for entire region the limit of y at any x will be,

Lower limit  $\rightarrow y = 2\sqrt{x}$  [At A, from the equation of curve  $y^2 = 4x$ ]

Upper limit  $\rightarrow y = \frac{x^2}{4}$  [At B, from the equation of  $x^2 = 4y$ ]

Now by sliding strip AB in horizontal direction from extreme left to right in required region R we get the limit of x as given below,

Lower limit  $\rightarrow x = 0$

Upper limit  $\rightarrow x = 4$

Hence required area (A),

$$A = \int_{x=0}^{x=4} \int_{y=2\sqrt{x}}^{y=\frac{x^2}{4}} dy dx$$

$$A = \int_{x=0}^{x=1} \left( \int_{y=\frac{x^2}{4}}^{y=2\sqrt{x}} dy \right) dx$$

$$\text{Area between the curves } A = \int_{x=0}^{x=4} \left( 2\sqrt{x} - \frac{x^2}{4} \right) dx$$

$$A = \left( 2 \cdot \frac{2}{3} \cdot x^{3/2} - \frac{x^3}{12} \right) \Big|_0^4$$

$$A = \frac{4}{3}(4^{3/2}) - \frac{4^3}{12} = \frac{32}{3} - \frac{16}{3} = \frac{16}{3}$$

$$\therefore \text{Area bounded} = \frac{16}{3} \text{ units}$$



Scan for Video solution



**173. (d)****Given:**

$$y = \frac{2}{3}x^{3/2} \Rightarrow \frac{dy}{dx} = \frac{2}{3} \cdot \frac{3}{2} \cdot x^{1/2} = \sqrt{x}$$

The length of the curve

 $y = f(x)$  between  $x = a$  and  $x = b$  is given by

$$L = \int_{x=a}^{x=b} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \cdot dx$$

$$L = \int_{x=0}^{x=1} \sqrt{1 + (\sqrt{x})^2} dx$$

$$L = \int_{x=0}^{x=1} (\sqrt{1+x}) dx$$

$$L = \frac{2}{3}(1+x)^{3/2} \Big|_{x=0}^{x=1} = \frac{2}{3}(2^{3/2} - 1^{3/2})$$

$$L = \frac{2}{3}(2\sqrt{2} - 1) = 1.2189$$

$$\therefore \boxed{\text{Length of the curve} = 1.22}$$



Scan for Video solution

**174. (0)**

$$\begin{aligned} \text{Let } I &= \int_{-3}^3 \int_{-2}^2 \int_{-1}^1 (4x^2y - z^3) dz dy dx \\ &= \int_{-3}^3 \int_{-2}^2 \left( 4x^2yz - \frac{z^4}{4} \Big|_{-1}^1 \right) dy dx \\ &= \int_{-3}^3 \int_{-2}^2 \left\{ \left( 4x^2y - \frac{1}{4} \right) - \left( -4x^2y - \frac{1}{4} \right) \right\} dy dx \\ &= \int_{-3}^3 \int_{-2}^2 8x^2y dy dx \\ &= \int_{-3}^3 \frac{8x^2y^2}{2} \Big|_{-2}^2 dx = \int_{-3}^3 \left( (16x^2) - (16x^2) \right) dx = 0 \end{aligned}$$

$$\therefore \boxed{\int_{-3}^3 \int_{-2}^2 \int_{-1}^1 (4x^2y - z^3) dz dy dx = 0}$$



Scan for Video solution

**175. (b)**

$$I = \int_{\theta=0}^{\theta=\pi/2} \int_{r=0}^{r=\cos\theta} r \sin \theta dr d\theta$$

$$I = \int_{\theta=0}^{\theta=\pi/2} \int_{r=0}^{r=\cos\theta} r \sin \theta dr d\theta$$

$$I = \int_{\theta=0}^{\theta=\pi/2} \int_{r=0}^{r=\cos\theta} (r \sin \theta dr) d\theta$$

$$I = \int_{\theta=0}^{\theta=\frac{\pi}{2}} \left( \sin \theta \cdot \frac{r^2}{2} \Big|_0^{\cos \theta} \right) d\theta$$

$$I = \frac{1}{2} \times \int_{\theta=0}^{\theta=\frac{\pi}{2}} \sin \theta \cdot \cos^2 \theta \cdot d\theta$$

Let  $\cos \theta = t \Rightarrow -\sin \theta d\theta = dt$ At  $\theta = 0, t = \cos 0 = 1$  and  $\theta = \pi/2, t = \cos \pi/2 = 0$ 

$$I = \frac{1}{2} \times \int_1^0 t^2 (-dt) = -\frac{1}{2} \times \frac{t^3}{3} \Big|_1^0$$

$$I = -\frac{1}{2} \times \left( 0 - \frac{1}{3} \right) = \frac{1}{6}$$

$$\therefore \boxed{\int_{\theta=0}^{\theta=\frac{\pi}{2}} \int_{r=0}^{r=\cos\theta} r \cdot \sin \theta \cdot dr d\theta = \frac{1}{6}}$$



Scan for Video solution

**176. (b)****Given:**

$$f(x, y, z) = e^{1-x \cos y} + x z e^{-1/(1+y^2)}$$

$$\Rightarrow f(x, y, z) = e \cdot e^{(-\cos y)x} + x z e^{-1/(1+y^2)}$$

Differentiating above equation with respect to x,

$$\Rightarrow \frac{\partial f}{\partial x} = e \cdot e^{(-\cos y)x} \cdot (-\cos y) + z \cdot e^{-1/(1+y^2)}$$

At the point  $(1, 0, e)$ 

$$\frac{\partial f}{\partial x} \Big|_{(1,0,e)} = e \cdot e^{(-\cos 0)1} \cdot (-\cos 0) + e \cdot e^{-1/(1+0^2)}$$

$$\Rightarrow \frac{\partial f}{\partial x} \Big|_{(1,0,e)} = e \cdot e^{-1} \cdot (-1) + e \cdot e^{-1} = 0$$

∴ Partial deviation of  $f(x, y, z)$  at  $(1, 0, e)$  is 0.



Scan for Video solution




177. (c)

Given

$$I = \int_{x=0}^1 \int_{y=0}^{x^2} xy^2 dy dx$$

The curves involved are  $x = 0, x = 1$   
 $y = 0, y = x^2$

The region R bounded by curve  $y = 0, y = x^2, x = 0$  and  $x = 1$  is shown in figure.



Now to change the order of integration taking a horizontal strip AB in region R.

Now for region R the limit of x at any y will be,

Lower limit  $\rightarrow x = \sqrt{y}$  [At A]

Upper limit  $\rightarrow x = 1$  [At B]

Now by sliding strip AB in vertical direction from bottom to top in required region R we get the limit of y as given below,

Lower limit  $\rightarrow y = 0$

Upper limit  $\rightarrow y = 4$

On changing the order of integration, integral I will become,

$$I = \int_{y=0}^1 \int_{x=\sqrt{y}}^1 xy^2 dx dy$$



Scan for Video solution



178. (d)

Given:  $f(x, y) = \frac{ax^2 + by^2}{xy}$  &  $\left. \frac{\partial f}{\partial x} \right|_{\substack{x=1 \\ y=2}} = \left. \frac{\partial f}{\partial y} \right|_{\substack{x=1 \\ y=2}}$

$$f(x, y) = \frac{ax^2 + by^2}{xy}$$

$$\Rightarrow f(x, y) = a \cdot \frac{x}{y} + b \cdot \frac{y}{x} \quad \dots(i)$$

Differentiating equation (i) partially with respect to x,

$$\Rightarrow \frac{\partial f}{\partial x} = \frac{a}{y} - \frac{b \cdot y}{x^2}$$

At  $x = 1$  and  $y = 2$ :

$$\Rightarrow \left. \frac{\partial f}{\partial x} \right|_{\substack{x=1 \\ y=2}} = \frac{a}{2} - 2b$$

Differentiating equation (i) partially with respect to y,

$$\Rightarrow \frac{\partial f}{\partial y} = \frac{-ax}{y^2} + \frac{b}{x}$$

At  $x = 1$  and  $y = 2$ :

$$\Rightarrow \left. \frac{\partial f}{\partial y} \right|_{\substack{x=1 \\ y=2}} = \frac{-a}{4} + b$$

$$\therefore \left. \frac{\partial f}{\partial x} \right|_{\substack{x=1 \\ y=2}} = \left. \frac{\partial f}{\partial y} \right|_{\substack{x=1 \\ y=2}}$$

$$\Rightarrow \frac{a}{2} - 2b = \frac{-a}{4} + b \Rightarrow \frac{3a}{4} = 3b$$

$$\Rightarrow a = 4b$$

Scan for Video solution



179. (4.4 to 4.6)

Given  $r = x^2 + y - z$  and  $z^3 - xy + y^2 + y^3 = 1$

$$r = x^2 + y - z$$

Differentiating both sides partially with respect to 'x'

$$\Rightarrow \frac{\partial r}{\partial x} = 2x - \frac{\partial z}{\partial x} \dots\dots(1)$$

$$z^3 - xy + yz + y^3 = 1$$

Differentiating both sides partially with respect to 'x'

$$\Rightarrow 3z^2 \cdot \frac{\partial z}{\partial x} - y + y \frac{\partial z}{\partial x} + 0 = 0$$

$$\Rightarrow \frac{\partial z}{\partial x} = \frac{y}{(3z^2 + y)} \quad \dots\dots(2)$$

Substituting  $\frac{\partial z}{\partial x}$  in (1)

$$\Rightarrow \frac{\partial r}{\partial x} = 2x - \frac{y}{3z^2 + y}$$

At the point  $(2, -1, 1)$ ,

$$\left. \frac{\partial r}{\partial x} \right|_{(2,-1,1)} = 2 \times 2 - \frac{(-1)}{3 \times 1^2 + (-1)}$$

$$\left. \frac{\partial r}{\partial x} \right|_{(2,-1,1)} = 4.5$$



Scan for Video solution



### 180. (40)

$$f(x, y, z) = (x^2 + y^2 - 2z^2)(y^2 + z^2)$$

$$f(x, y, z) = x^2y^2 + x^2z^2 + y^4 + y^2z^2 - 2y^2z^2 - 2z^4$$

Differentiating partially with respect to  $x$ ,

$$\Rightarrow \frac{\partial f}{\partial x} = 2xy^2 + 2xz^2 + 0 + 0 - 0 - 0$$

$$\Rightarrow \frac{\partial f}{\partial x} = 2xy^2 + 2xz^2$$

At  $(2, 1, 3)$

$$\Rightarrow \frac{\partial f}{\partial x} = 2(2)(1)^2 + 2(2)(3)^2 = 4 + 36$$

$$\Rightarrow \left. \frac{\partial f}{\partial x} \right|_{(2,1,3)} = 40$$



Scan for Video solution




### 181. (1.00)

Equation of the line joining points  $(1, 2)$  and  $(5, 10)$  is:

$$\frac{x-1}{5-1} = \frac{y-2}{10-2}$$

$$\Rightarrow 2 \times (x-1) = y-2$$

$$\Rightarrow y = 2x$$



Taking vertical strip AB in required region R.

Now for entire region the limit of  $y$  at any  $x$  will be,

Lower limit  $\rightarrow y = 0$  [At A]

Upper limit  $\rightarrow y = 2x$  [At B]

Now by sliding strip AB in horizontal direction from extreme left to right in required region R we get the limit of  $x$  as given below,

Lower limit  $\rightarrow x = 1$

Upper limit  $\rightarrow x = 5$

Therefore,

$$I = c \int \int_R xy^2 dx dy$$

$$\Rightarrow I = c \int_{x=1}^{x=5} \int_{y=0}^{y=2x} xy^2 dx dy$$

$$\Rightarrow I = c \int_{x=1}^5 x \left[ \frac{y^3}{3} \right]_0^{2x} dx = c \int_{x=1}^5 x \left( \frac{8x^3}{3} \right) dx$$

$$\Rightarrow I = c \times \frac{8}{3} \int_{x=1}^5 x^4 dx$$


$$\Rightarrow I = 6 \times 10^{-4} \times \frac{8}{3} \times \left[ \frac{x^5}{5} \right]_1^5$$

$$\Rightarrow I = 6 \times 10^{-4} \times \frac{8}{3} \times \frac{5^5 - 1}{5} = 0.99968$$

$$\Rightarrow I = 1.00$$



Scan for Video solution

**182. (0.7 to 0.85)**Given  $x^2 + y^2 \leq z^3$ ;  $0 \leq z \leq 1$ .Let  $z^3 = r^2$ ,  $dv = \pi r^2 \cdot dz$ 

$$\therefore \text{Volume} = \int dv = \int_0^1 \pi z^3 dz = \frac{\pi z^4}{4} \Big|_0^1 = \frac{\pi}{4} = 0.7854$$


 $\therefore$  Volume of region R = 0.7854

Scan for Video solution

**183. (4 to 5)**

The region is given by

$$\left\{ (\rho, \phi, z) : 3 \leq \rho \leq 5, \frac{\pi}{8} \leq \phi \leq \frac{\pi}{4}, 3 \leq z \leq 4.5 \right\}$$



Volume of the region

= Volume of the portion of cylinder

= base area  $\times$  height

$$= \frac{\left(\frac{\pi}{4} - \frac{\pi}{8}\right)}{2\pi} \times \pi \times \left\{5^2 - 3^2\right\} \times (4.5 - 3)$$

$$= \frac{1}{16} \times \pi \times 16 \times 1.5$$


$$= 1.5\pi = 4.712$$

 $\therefore$  Volume of Region = 4.712 units

Scan for Video solution

**184. (10)**Given region in xy-plane is bounded by  $2x = 3y$ 

$$\Rightarrow \frac{2}{3}x, y = 0 \text{ and } x = 3$$



$$\text{Volume of the region} = \int_{x=0}^{x=3} \int_{y=0}^{y=\frac{2}{3}x} \int_{z=6-x-y}^{z=6} dz dy dx$$

$$= \int_{x=0}^{x=3} \int_{y=0}^{y=\frac{2}{3}x} (6 - x - y) dy dx$$

$$= \int_{x=0}^{x=3} (6 - x)y - \frac{y^2}{2} \Big|_0^{2/3} dx$$

$$= \int_{x=0}^{x=3} \left\{ (6 - x) \cdot \frac{2x}{3} - \frac{1}{2} \left( \frac{4}{9} x^2 \right) \right\} dx$$

$$= \int_{x=0}^{x=3} \left\{ 4x - \frac{2}{3}x^2 - \frac{2}{9}x^2 \right\} dx$$

$$= \int_{x=0}^{x=3} \left( 4x - \frac{8x^2}{9} \right) dx = 2x^2 - \frac{8x^3}{27} \Big|_0^3$$

$$= 18 - 8 = 10$$


 $\therefore$  Volume of the region = 10 units



Scan for Video solution

**185. (0.7 to 0.76)**

Triangle bounded by the lines  $x = y$ ;  $x = 0$ ;  $y = 1$  is shown in figure:



Required volume

$$\begin{aligned}
 &= \int \int f(x, y) dx dy \\
 &\quad \text{R} \\
 &= \int_{x=0}^{x=1} \int_{y=x}^{y=1} e^x dy dx \\
 &= \int_0^1 e^x [y]_x^1 dx = \int_0^1 e^x [1-x] dx \\
 &= \left[ e^x (1-x) \right]_0^1 - \int_0^1 (-1) \times e^x dx \\
 &= \left[ (0) - (1) \right] + \int_0^1 e^x dx \\
 &= -1 + \left[ e^x \right]_0^1 \\
 &= -1 + (e - 1) \\
 &= 0.718
 \end{aligned}$$



Scan for Video solution

**186. (a)**

$$\text{Given } \frac{\partial}{\partial y} (x^2 + y^2) = \frac{\partial}{\partial x} (6y + 4x)$$

$$\Rightarrow 2y = 4$$

$$\Rightarrow y = 2$$

$\therefore$  the required contour is  $y = 2$



Scan for Video solution

**187. (b)**

$$u = \frac{2x - y}{2} \quad \dots(i)$$

$$\text{And } v = \frac{y}{2} \Rightarrow y = 2v \quad \dots(ii)$$

$$\text{Put } y = 2v \text{ in equation (i), } u = \frac{2x - 2v}{2}$$

$$\Rightarrow u = x - v$$

$$\Rightarrow x = u + v \quad \dots(iii)$$

$$\text{Let, } I = \int_0^8 \left( \int_{y/2}^{(y/2)+1} \left( \frac{2x - y}{2} \right) dx \right) dy$$

By changing the variable above integration will become,

$$I = \int_{v_1}^{v_2} \left( \int_{u_1}^{u_2} u |J| du \right) dv \quad \dots(iv)$$

$$\left[ \because \frac{2x - y}{2} = u \right]$$

Where  $J$  = Jacobian of transformation.

$$x \rightarrow \frac{y}{2} \text{ to } \frac{y}{2} + 1 \quad \text{and} \quad y \rightarrow 0 \text{ to } 8$$

Lower limit of  $u$  ( $u_1$ ): Put  $x = \frac{y}{2}$  &  $y = 0$  in equation (i),

$$u_1 = \frac{2\left(\frac{y}{2}\right) - y}{2} = \frac{2\left(\frac{0}{2}\right) - 0}{2} = 0$$

Upper limit of  $u$  ( $u_2$ ): Put  $x = \frac{y}{2} + 1$  &  $y = 8$  in equation (i),

$$u_2 = \frac{2\left(\frac{y}{2} + 1\right) - y}{2} = \frac{2\left(\frac{8}{2} + 1\right) - 8}{2} = 1$$

Lower limit of  $v$  ( $v_1$ ): Put  $x = \frac{y}{2}$  &  $y = 0$  in equation (ii),

$$v_1 = \frac{0}{2} = 0$$

Upper limit of  $v$  ( $v_2$ ) : Put  $x = \frac{y}{2} + 1$  &  $y = 8$  in equation (ii),  $v_2 = \frac{8}{2} = 4$

Jacobian matrix

$$J = J\left(\frac{x, y}{u, v}\right) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$

$$\Rightarrow J = \begin{vmatrix} \frac{\partial(u+v)}{\partial u} & \frac{\partial(u+v)}{\partial v} \\ \frac{\partial(2v)}{\partial u} & \frac{\partial(2v)}{\partial v} \end{vmatrix}$$

$$\Rightarrow J = \begin{vmatrix} 1 & 1 \\ 0 & 2 \end{vmatrix} = 2$$

Hence from equation (iv),

$$I = \int_0^4 \left( \int_0^1 2udu \right) dv$$

Therefore,


$$I = \int_0^4 \left( \int_{y/2}^{(y/2)+1} \left( \frac{2x-y}{2} \right) dx \right) dy = \int_0^4 \left( \int_0^1 2udu \right) dv$$



Scan for Video solution



### 188. (864)



Given region is bounded by

$xy$  - plane ( $z = 0$ ) and  $z = x + y$   
 $\{0 \leq y \leq x \text{ and } 0 \leq x \leq 12\}$

$\therefore$  Volume

$$\begin{aligned}
 &= \int_{x=0}^{x=12} \int_{y=0}^{y=x} \int_{z=0}^{z=x+y} dz dy dx \\
 &= \int_{x=0}^{x=12} \int_{y=0}^{y=x} (x+y) dy dx = \int_{x=0}^{x=12} \left[ xy + \frac{y^2}{2} \right]_0^x dx \\
 &= \int_{x=0}^{x=12} \left( x^2 + \frac{x^2}{2} \right) dx \\
 &= \int_{x=0}^{x=12} \frac{3}{2} x^2 dx = \frac{x^3}{2} \Big|_0^{12} \\
 &= \frac{(12)^3}{2} = 864
 \end{aligned}$$

$\therefore$  Volume bounded by the given curves is 864 units.

▶
Scan for Video solution

### 189. (b)

$$I = \int_0^{2x} \int_0^x e^{x+y} \cdot dy \cdot dx$$

$$I = \int_0^{2x} \int_0^x e^x \cdot e^y \cdot dy \cdot dx$$

$$I = \int_0^2 e^x \cdot \left( e^y \Big|_0^x \right) dx$$

$$I = \int_0^2 e^x \cdot (e^x - 1) dx$$

$$I = \int_0^2 (e^{2x} - e^x) dx = \frac{e^{2x}}{2} - e^x \Big|_0^2$$

$$I = \left( \frac{e^4}{2} - e^2 \right) - \left( \frac{1}{2} - 1 \right)$$

$$I = \frac{e^4}{2} - e^2 + \frac{1}{2} = \frac{1}{2}(e^4 - 2e^2 + 1)$$

$$\therefore \int_0^{2x} \int_0^x e^{x+2y} dy dx = \frac{1}{2}(e^2 - 1)^2$$




Scan for Video solution

**190. (a)**

$$x > y^2 \quad x \in [0, 1]$$

$$y > x^2 \quad y \in [0, 1]$$



In case of horizontal strip limits will be  $x = y^2$  and  $x = \sqrt{y}$ , then this strip will move from  $y = 0$  to  $y = 1$  to give the required volume of  $f(x, y)$ . Hence volume under  $f(x, y)$  is


$$\int_{y=0}^{y=1} \int_{x=y^2}^{x=\sqrt{y}} f(x, y) dx dy$$



Scan for Video solution

**191. (b)****Given:** function is  $(x + y)^2$ Radius of quarter of a circle,  $r = 1$ 

$$\text{Let } I = \int_C (x + y)^2 ds \quad \dots(i)$$



$$ds = r \cdot d\theta = d\theta$$

$$[\because r = 1]$$

$$x = \cos\theta; y = \sin\theta$$

From equation (i),

$$I = \int_{\theta=0}^{\frac{\pi}{2}} (\cos\theta + \sin\theta)^2 \cdot d\theta$$

$$I = \int_{\theta=0}^{\frac{\pi}{2}} (\cos^2\theta + \sin^2\theta + 2\sin\theta\cos\theta) \cdot d\theta$$

$$I = \int_{\theta=0}^{\frac{\pi}{2}} (1 + \sin 2\theta) \cdot d\theta$$

$$I = \theta - \frac{\cos 2\theta}{2} \Big|_0^{\frac{\pi}{2}}$$

$$I = \left\{ \frac{\pi}{2} - \left( -\frac{1}{2} \right) \right\} - \left\{ 0 - \frac{1}{2} \right\} = \frac{\pi}{2} + 1$$


$$\therefore I = \int_{\theta=0}^{\frac{\pi}{2}} (x + y)^2 ds = \frac{\pi}{2} + 1$$



Scan for Video solution

**192. (a)****Given:**

$$\text{Integral } I = \iint_P xy dx dy$$



Equation of line RS:

$$\frac{x}{2} + \frac{y}{1} = 1 \Rightarrow x = 2(1 - y)$$

Taking horizontal strip AB in the region P.

Now for region P the limit of x at any y will be,  
Lower limit  $\rightarrow x = 0$  [At A]Upper limit  $\rightarrow x = 2(1 - y)$  [At B]

Now by sliding strip AB in vertical direction from bottom to top in required region P we get the limit of y as given below,

Lower limit  $\rightarrow y = 0$ Upper limit  $\rightarrow y = 1$

Therefore,

$$I = \iint_P xy \, dx \, dy = \int_{y=0}^{y=1} \int_{x=0}^{x=2(1-y)} xy \, dx \, dy$$

$$I = \int_{y=0}^{y=1} y \left( \frac{x^2}{2} \Big|_0^{2(1-y)} \right) dy$$

$$I = \int_{y=0}^{y=1} \frac{y}{2} \cdot 4(1-y)^2 dy$$

$$I = \int_{y=0}^{y=1} y(1-2y+y^2) dy = \int_{y=0}^{y=1} (y-2y^2+y^3) dy$$

$$I = 2 \times \left( \frac{y^2}{2} - \frac{2}{3}y^3 + \frac{y^4}{4} \Big|_0^1 \right)$$

$$I = 2 \left( \frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) = \frac{1}{6}$$

$$\therefore \iint_P xy \, dx \, dy = \frac{1}{6}$$

 Scan for Video solution



### Basic Probability

- 1. [NAT] [GATE-CE-2023:1M]**

The probabilities of occurrences of two independent events A and B are 0.5 and 0.8, respectively. What is the probability of occurrence of at least A or B (*Rounded off to one decimal place*)?

- 2. [NAT] [GATE-EE-2023:2M]**

The expected number of trials for first occurrence of a "head" in a biased coin is known to be 4. The probability of first occurrence of a "head" in the second trial is \_\_\_\_\_.  
(*Round off to 3 decimal places*)

- 3. [NAT] [GATE-CE-2022:2M]**

A pair of six-faced dice is rolled twice, the probability that the sum of the outcomes in each roll equals 4 in exactly two of the three attempts is \_\_\_\_\_ (*round off to three decimal places*).

- 4. [MCQ] [GATE-EE-2021:2M]**

Suppose the probability that a coin toss shows "head" is  $p$ , where  $0 < p < 1$ . The coin is tossed repeatedly until the first "head" appears. The expected number of tosses required is

- (a)  $p/(1 - p)$       (b)  $(1 - p)/p$   
(c)  $1/p$       (d)  $1/p^2$

- 5. [NAT] [GATE-2020-CE:1M]**

A fair (unbiased) coin is tossed 15 times. The probability of getting exactly 8 heads (*round off to three decimal places*), is \_\_\_\_\_.

- 6. [NAT] [GATE-ME-2020:1M]**

A company is hiring to fill four managerial vacancies. The candidates are five men and three women. If every candidate is equally likely to be chosen then the probability that at least one woman will be selected is \_\_\_\_\_  
(*round off to 2 decimal places*).

- 7. [NAT] [GATE-ME-2020:2M]**

A fair coin is tossed 20 times. The probability that 'head' will appear exactly 4 times in the first ten tosses, and 'tail' will appear exactly 4 times in the next ten tosses is \_\_\_\_\_.  
(*Round off to 3 decimal places*)

- 8. [NAT] [GATE-ME-2019:2M]**

The probability that a part manufactured by a company will be defective is 0.05. If 15 such parts are selected randomly and inspected, then the probability that at least two parts will be defective is \_\_\_\_\_. (*Round off to two decimal places*)

- 9. [NAT] [GATE-EE-2019:2M]**

The probability of a resistor being defective is 0.02. There are 50 such resistors in a circuit. The probability of two or more defective resistors in the circuit (*round off to two decimal places*) is \_\_\_\_\_.  
(*Round off to two decimal places*)

- 10. [NAT] [GATE-CE-2018:1M]**

Probability (up to one decimal place) of consecutively picking 3 red balls without replacement from a box containing 5 red balls and 1 white ball is \_\_\_\_\_.

**11. [MCQ]****[GATE-ME-2018:1M]**

Four red balls, four green balls and four blue balls are put in a box. Three balls are pulled out of the box at random one after another without replacement. The probability that all the three balls are red is

(a)  $\frac{1}{72}$

(b)  $\frac{1}{55}$

(c)  $\frac{1}{36}$

(d)  $\frac{1}{27}$

**12. [MCQ]****[GATE-ME-2018:1M]**

A six-faced fair dice is rolled five times. The probability (in %) of obtaining “ONE” at least four times is

(a) 33.3

(b) 3.33

(c) 0.33

(d) 0.0033

**13. [NAT]****[GATE-CS-2018:1M]**

Two people, P and Q, decide to independently roll two identical dice, each with 6 faces, numbered 1 to 6. The person with the lower number wins. In case of a tie, they roll the dice repeatedly until there is no tie. Define a trial as a throw of the dice by P and Q. Assume that all 6 numbers on each dice are equiprobable and that all trials are independent. The probability (rounded to 3 decimal places) that one of them wins on the third trial is \_\_\_\_\_.

**14. [MCQ]****[GATE-CS-2017:2M]**

P and Q are considering to apply for a job. The probability that P applies for the job is  $1/4$ , the probability that P applies for the job given that Q applies for the job is  $1/2$ , and the probability that Q applies for the job given that P applies for the job is  $1/3$ . Then the probability that P does not apply for the job given that Q does not apply for the job is

(a)  $\frac{4}{5}$

(b)  $\frac{5}{6}$

(c)  $\frac{7}{8}$

(d)  $\frac{11}{12}$

**15. [NAT]****[GATE-ME-2017:1M]**

A six-face fair dice is rolled a large number of times. The mean value of other outcomes is \_\_\_\_\_.

**16. [NAT]****[GATE-ME-2017:1M]**

Two coins are tossed simultaneously. The probability (upto two decimal points accuracy) of getting at least one head is \_\_\_\_\_.

**17. [NAT]****[GATE-CE-2017:1M]**

A two-faced coin has its faces designated as head (H) and tail (T). This coin is tossed three times in succession to record the following outcomes; H, H, H. If the coin is tossed one more time, the probability (up to one decimal place) of obtaining H again, given the previous realization of H, H and H, would be \_\_\_\_\_.

**18. [MCQ]****[GATE-EE-2017:1M]**

An urn contains 5 red balls and 5 black balls. In the first draw, one ball is picked at random and discarded without noticing its color. The probability to get a red ball in the second draw is

(a)  $\frac{1}{2}$

(b)  $\frac{4}{9}$

(c)  $\frac{5}{9}$

(d)  $\frac{6}{9}$

**19. [NAT]****[GATE-EC-2017:1M]**

Three fair cubical dice are thrown simultaneously. The probability that all three dice have the same number of dots on the faces showing up is (up to third decimal place) \_\_\_\_\_.

**20. [NAT]****[GATE-EC-2017:2M]**

Passengers try repeatedly to get a seat reservation in any train running between two stations until they are successful. If there is 40 % chance of getting reservation in any attempt by a passenger then the average number of attempts that passengers need to make to get a seat reserved is \_\_\_\_\_.

**21. [NAT]****[GATE-EC-2017:1M]**

Three fair cubical dice are thrown simultaneously. The probability that all three dice have the same number of dots on the faces showing up is (up to third decimal place) \_\_\_\_\_.

**22. [NAT]****[GATE-EE-2016:2M]**

Candidates were asked to come to an interview with 3 pens each. Black, Blue, green and red were the permitted pen colors that the candidate could bring. The probability that a candidate comes with all 3 pens having the same colour is\_\_\_\_\_.

**23. [NAT]****[GATE-ME-2016:2M]**

The probability that a screw manufactured by a company is defective is 0.1. The company sells screws in packets containing 5 screws and gives a guarantee of replacement if one or more screws in the packet are found to be defective. The probability that a packet would have to be replaced is\_\_\_\_\_.

**24. [MCQ]****[GATE-ME-2016:2M]**

Three cards were drawn from a pack of 52 cards. The probability that they are a king, a queen, and a jack is

(a)  $\frac{16}{5525}$

(b)  $\frac{64}{2197}$

(c)  $\frac{3}{13}$

(d)  $\frac{8}{16575}$

**25. [NAT]****[GATE-EC-2016:1M]**

The probability of getting a "head" in a single toss of a biased coin is 0.3. The coin is tossed repeatedly till a "head" is obtained. If the tosses are independent, then the probability of getting "head" for the first time in the fifth toss is\_\_\_\_\_.

**26. [NAT]****[GATE-CS-2016:2M]**

Consider the following experiment.

**Step 1.** Flip a fair coin twice.

**Step 2.** If the outcomes are (TAILS, HEADS) then output Y and stop.

**Step 3.** If the outcomes are either (HEADS, HEADS) or (HEADS, TAILS), then output N and stop.

**Step 4.** If the outcomes are (TAILS, TAILS), then go to Step1.

The probability that the output of the experiment is Y is (up to two decimal places)\_\_\_\_\_.

**27. [MCQ]****[GATE-ME-2015:2M]**

The probability of obtaining at least two "SIX" in throwing a fair dice 4 times is

- (a) 425/432      (b) 19/144  
(c) 13/144      (d) 125/432

**28. [NAT]****[GATE-ME-2015:1M]**

Three vendors were asked to supply a very high precision component. The respective probabilities of their meeting the strict design specifications are 0.8, 0.7 and 0.5. Each vendor supplies one component. The probability that out of total three components supplied by the vendors, at least one will meet the design specifications is\_\_\_\_\_.

**29. [MCQ]****[GATE-ME-2015:2M]**

The chance of a student passing an exam is 20%. The chance of a student passing the exam and getting above 90% marks in it is 5%. GIVEN that a student passes the examination, the probability that the student gets above 90% marks is

- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{1}{18}$ | (b) $\frac{1}{4}$  |
| (c) $\frac{2}{9}$  | (d) $\frac{5}{18}$ |

**30. [MCQ]****[GATE-ME-2015:1M]**

If  $P(X)=1/4$ ,  $P(Y)=1/3$ , and  $P(X \cap Y)=1/12$ , the value of  $P(Y/X)$  is

- (a) 1/4      (b) 4/25  
(c) 1/3      (d) 29/50

**31. [NAT]****[GATE-EC-2015:2M]**

Let the random variable X represent the number of times a fair coin needs to be tossed till two consecutive heads appear for the first time. The expectation of X is\_\_\_\_\_.

**32. [MCQ]****[GATE-EC-2015:1M]**

Suppose A and B are two independent events with probabilities  $P(A) \neq 0$  and  $P(B) \neq 0$ . Let  $\bar{A}$  and  $\bar{B}$  be their complements. Which one of the following statements is FALSE?

- (a)  $P(A \cap B) = P(A)P(B)$   
(b)  $P(A|B) = P(A)$   
(c)  $P(A \cup B) = P(A) + P(B)$   
(d)  $P(\bar{A} \cap \bar{B}) = P(\bar{A})P(\bar{B})$

**33. [NAT] [GATE-EC-2015:2M]**

A fair die with faces {1,2,3,4,5,6} is thrown repeatedly till '3' is observed for the first time. Let X denote the number of times the die is thrown. The expected value of X is \_\_\_\_\_.

**34. [MCQ] [GATE-EE-2015:2M]**

Two players, A and B, alternately keep rolling a fair dice. The person to get a six first wins the game. Given that player A starts the game, the probability that A wins the game is

- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{5}{11}$ | (b) $\frac{1}{2}$  |
| (c) $\frac{7}{13}$ | (d) $\frac{6}{11}$ |

**35. [MCQ] [GATE-EE-2015: 2M]**

Two coins R and S are tossed. The 4 joint events  $H_R H_s$ ,  $T_R T_s$ ,  $H_R T_s$ ,  $T_R H_s$  have probabilities 0.28, 0.18, 0.30, 0.24 respectively, where H represents head and T represents tail. Which one of the following is TRUE

- (a) The coin tosses are independent
- (b) R is fair, S is not.
- (c) S is fair, R is not
- (d) The coin tosses are dependent

**36. [MCQ] [GATE-ME-2014:1M]**

A box contains 25 parts of which 10 are defective. Two parts are being drawn simultaneously in a random manner from the box. The probability of both the parts being good is

- |                     |                      |
|---------------------|----------------------|
| (a) $\frac{7}{20}$  | (b) $\frac{42}{125}$ |
| (c) $\frac{25}{29}$ | (d) $\frac{5}{9}$    |

**37. [NAT] [GATE-ME-2014:1M]**

A group consists of equal number of men and women. Of this group 20% of the men and 50% of the women are unemployed. If a person is selected at random from this group, the probability of the selected person being employed is \_\_\_\_\_.

**38. [NAT] [GATE-ME-2014:2M]**

Consider an unbiased cubic dice with opposite faces coloured identically and each face coloured red, blue or green such that each colour appears only two times on the dice. If the dice is thrown thrice, the probability of obtaining red colour on top face of the dice at least twice is \_\_\_\_\_.

**39. [MCQ] [GATE-2014-CE:1M]**

A fair (unbiased) coin was tossed four times in succession and resulted in the following outcomes:  
(i) Head, (ii) Head, (iii) Head, (iv) Head. The probability of obtaining a 'Tail' when the coin is tossed again is

- |                   |                   |
|-------------------|-------------------|
| (a) 0             | (b) $\frac{1}{2}$ |
| (c) $\frac{4}{5}$ | (d) $\frac{1}{5}$ |

**40. [NAT] [GATE-EC-2014:1M]**

In a housing society, half of the families have a single child per family, while the remaining half have two children per family. The probability that a child picked at random, has a sibling is \_\_\_\_\_.

**41. [NAT] [GATE-EC-2014:1M]**

A fair coin is tossed repeatedly till both head and tail appear at least once. The average number of tosses required is \_\_\_\_\_.

**42. [MCQ] [GATE-EC-2014:1M]**

An unbiased coin is tossed an infinite number of times. The probability that the fourth head appears at the tenth toss is

- |           |           |
|-----------|-----------|
| (a) 0.067 | (b) 0.073 |
| (c) 0.082 | (d) 0.091 |

**43. [NAT] [GATE-EE-2014:1M]**

Consider a dice with the property that the probability of a face with n dots showing up is proportional to n. The probability of the face with three dots showing up is \_\_\_\_\_.

**44. [NAT] [GATE-CS-2014:2M]**

Let S be a sample space and two mutually exclusive events A and B be such that  $A \cup B = S$ . If  $P(\cdot)$  denotes the probability of the event, the maximum value of  $P(A)P(B)$  is \_\_\_\_\_.

**45. [MCQ]****[GATE-ME-2013:2M]**

The probability that a student knows the correct answer to a multiple choice question is  $\frac{2}{3}$ . If the student does not know the answer, then the student guesses the answer. The probability of the guessed answer being correct is  $\frac{1}{4}$ . Given that the student has answered the question correctly, the conditional probability that the student knows the correct answer is

- |                   |                   |
|-------------------|-------------------|
| (a) $\frac{2}{3}$ | (b) $\frac{3}{4}$ |
| (c) $\frac{5}{6}$ | (d) $\frac{8}{9}$ |

**46. [MCQ]****[GATE-ME-2012:2M]**

A box contains 4 red balls nad 6 black balls. Three balls are selected randomly from the box one after another, without replacement. The probability that the selected set contains one red ball and two black ball is

- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{1}{20}$ | (b) $\frac{1}{12}$ |
| (c) $\frac{3}{10}$ | (d) $\frac{1}{2}$  |

**47. [MCQ]****[GATE-CE-2012:2M]**

In an experiment, positive and negative values are equally likely to occur. The probability of obtaining at most one negative value in five trials is

- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{1}{32}$ | (b) $\frac{2}{32}$ |
| (c) $\frac{3}{32}$ | (d) $\frac{6}{32}$ |

**48. [MCQ]****[GATE-EE-2012:2M]**

A fair coin is tossed till a head appears for the first time. The probability that the number of required tosses is odd, is

- |                   |                   |
|-------------------|-------------------|
| (a) $\frac{1}{3}$ | (b) $\frac{1}{2}$ |
| (c) $\frac{2}{3}$ | (d) $\frac{3}{4}$ |

**49. [MCQ]****[GATE-EC-2012:2M]**

A fair coin is tossed till a head appears for the first time. The probability that the number of required tosses is odd, is

- |                   |                   |
|-------------------|-------------------|
| (a) $\frac{1}{3}$ | (b) $\frac{1}{2}$ |
| (c) $\frac{2}{3}$ | (d) $\frac{3}{4}$ |

**50. [MCQ]****[GATE-EC-2012:2M]**

A fair coin is tossed independently four times. The probability of the event "the number of times heads show up is more than the number of times tails show up" is

- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{1}{16}$ | (b) $\frac{1}{8}$  |
| (c) $\frac{1}{4}$  | (d) $\frac{5}{16}$ |

**51. [MCQ]****[GATE-CE-2011:1M]**

There are two containers, with one containing 4 Red and 3 Green balls and the other containing 3 Blue and

4 Green balls. One ball is drawn at random from each container. The probability that one of the balls is Red and the other is Blue will be

- |                     |                    |
|---------------------|--------------------|
| (a) $\frac{1}{7}$   | (b) $\frac{9}{49}$ |
| (c) $\frac{12}{49}$ | (d) $\frac{3}{7}$  |

**52. [MCQ]****[GATE-ME-2011:2M]**

A box contains 2 washers, 3 nuts and 4 bolts. Items are drawn from the box at random one at a time without replacement. The probability of drawing 2 washers first followed by 3 nuts and subsequently the 4 bolts is

- |                      |                      |
|----------------------|----------------------|
| (a) $\frac{2}{315}$  | (b) $\frac{1}{630}$  |
| (c) $\frac{1}{1260}$ | (d) $\frac{1}{2520}$ |

**53. [MCQ]****[GATE-ME-2011:2M]**

An unbiased coin is tossed five times. The outcome of each toss is either a head or a tail. The probability of getting at least one head is

- |                     |                     |
|---------------------|---------------------|
| (a) $\frac{1}{32}$  | (b) $\frac{13}{32}$ |
| (c) $\frac{16}{32}$ | (d) $\frac{31}{32}$ |

**54. [MCQ]****[GATE-EC-2011:2M]**

A fair dice is tossed two times. The probability that the second toss results in a value that is higher than the first toss is

- |                    |                   |
|--------------------|-------------------|
| (a) $\frac{2}{36}$ | (b) $\frac{2}{6}$ |
| (c) $\frac{5}{12}$ | (d) $\frac{1}{2}$ |

**55. [MCQ]****[GATE-CS-2011:1M]**

If two fair coins are flipped and at least one of the outcomes is known to be a head, what is the probability that both outcomes are heads?

- (a)  $1/3$       (b)  $1/4$   
 (c)  $1/2$       (d)  $2/3$

**56. [MCQ]****[GATE-CS-2010:2M]**

What is the probability that a divisor of  $10^{99}$  is a multiple of  $10^{96}$ ?

- (a)  $1/625$       (b)  $4/625$   
 (c)  $12/625$       (d)  $16/625$

**57. [MCQ]****[GATE-CE-2010:1M]**

Two coins are simultaneously tossed. The probability of two heads simultaneously appearing is

- (a)  $1/8$       (b)  $1/6$   
 (c)  $1/4$       (d)  $1/2$

**58. [MCQ]****[GATE-EE-2010:2M]**

A box contains 4 white balls and 3 red balls. In succession, two balls are randomly selected and removed from the box. Given that the first removed ball is white, the probability that the second removed ball is red is

- (a)  $1/3$       (b)  $3/7$   
 (c)  $1/2$       (d)  $4/7$

**59. [MCQ]****[GATE-EE-2009:1M]**

Assume for simplicity that N people, all born in April (a month of 30 days), are collected in a room. Consider the event of at least two people in the room being born on the same date of the month, even if in different years, e.g. 1980 and 1985. What is the smallest N so that the probability of this event exceeds 0.5?

- (a) 20      (b) 7  
 (c) 15      (d) 16

**60. [MCQ]****[GATE-EC-2009:1M]**

A fair coin is tossed 10 times. What is the probability that ONLY the first two tosses will yield heads?

- (a)  $\left(\frac{1}{2}\right)^2$       (b)  ${}^{10}C_2 \left(\frac{1}{2}\right)^2$   
 (c)  $\left(\frac{1}{2}\right)^{10}$       (d)  ${}^{10}C_2 \left(\frac{1}{2}\right)^{10}$

**61. [MCQ]****[GATE-ME-2009:1M]**

If three coins are tossed simultaneously, the probability of getting at least one head is

- (a)  $\frac{1}{8}$       (b)  $\frac{3}{8}$   
 (c)  $\frac{1}{2}$       (d)  $\frac{7}{8}$

**62. [MCQ]****[GATE-ME-2008:1M]**

A coin is tossed 4 times. What is the probability of getting heads exactly 3 times?

- (a)  $\frac{1}{4}$       (b)  $\frac{3}{8}$   
 (c)  $\frac{1}{2}$       (d)  $\frac{3}{4}$

**63. [MCQ]****[GATE-CE-2008:2M]**

A person on a trip has a choice between private car and public transport. The probability of using a private car is 0.45. While using the public transport, further choices available are bus and metro, out of which the probability commuting by a bus is 0.55. In such a situation, the probability (rounded up to two decimals) of using a car, bus and metro, respectively would be

- (a) 0.45, 0.30 and 0.25  
 (b) 0.45, 0.25 and 0.30  
 (c) 0.45, 0.55 and 0.00  
 (d) 0.45, 0.35 and 0.20



**70. [NAT] [GATE-ME-2022:2M]**

Let a random variable X follow Poisson distribution such that  $\text{Prob}(X = 1) = \text{Prob}(X = 2)$ .

The value of  $\text{Prob}(X = 3)$  is \_\_\_\_\_.  
*(Round off to 2 decimal places)*

**71. [MCQ] [GATE-ME-2021:1M]**

Consider a binomial random variable X. If  $X_1, X_2, \dots, X_n$  are independent and identically distributed samples from the distribution of X with sum

$$Y = \sum_{i=1}^n X_i,$$

can be approximated as

- (a) Exponential
- (b) Bernoulli
- (c) Binomial
- (d) Normal

**72. [MCQ] [GATE-ME-2021:1M]**

The mean and variance, respectively, of a binomial distribution for  $n$  independent trials with the probability of success as  $p$ , are

- (a)  $\sqrt{np}, np(1-2p)$
- (b)  $\sqrt{np}, \sqrt{np(1-p)}$
- (c)  $np, np$
- (d)  $np, np(1-p)$

**73. [NAT] [GATE-CS-2021:1M]**

The lifetime of a component of a certain type is a random variable whose probability density function is exponentially distributed with parameter 2. For a randomly picked component of this type, the probability that its lifetime exceeds the expected lifetime (rounded to 2 decimal places) is \_\_\_\_\_.  
**[GATE-CS-2021:1M]**

**74. [NAT] [GATE-CS-2021:1M]**

For a given biased coin, the probability that the outcome of a toss is a head is 0.4. This coin is tossed 1,000 times. Let X denote the random variable whose value is the number of times that head appeared in these 1,000 tosses. The standard deviation of X (rounded to 2 decimal places) is \_\_\_\_\_.  
**[GATE-CS-2021:1M]**

**75. [MCQ] [GATE-2021-CE:1M]**

The shape of the cumulative distribution function of Gaussian distribution is

- (a) Bell-shaped
- (b) S-shaped
- (c) Horizontal line
- (d) Straight line at 45-degree angle

**76. [NAT] [GATE-CS-2020:2M]**

For  $n > 2$ , let  $a \in \{0, 1\}^n$  be a non-zero vector. Suppose that x is chosen uniformly at random from  $\{0, 1\}^n$ . Then, the probability that  $\sum_{i=1}^n a_i x_i$  is an odd number is \_\_\_\_\_.  
**[GATE-CS-2020:2M]**

**77. [NAT] [GATE-ME-2020:2M]**

Consider two exponentially distributed random variables X and Y, both having a mean of 0.50. Let  $Z = X + Y$  and  $r$  be the correlation coefficient between X and Y. If the variance of Z equals 0, then the value of  $r$  is \_\_\_\_\_.  
*(Round off to 2 decimal places)*

**78. [MCQ] [GATE-ME-2020:1M]**

The sum of two normally distributed random variables X and Y is

- (a) always normally distributed.
- (b) normally distributed, only if X and Y are independent.
- (c) normally distributed, only if X and Y have the same standard deviation.
- (d) normally distributed, only if X and Y have the same mean.

**79. [NAT] [GATE-EC-2020:2M]**

x is a random variable with uniform probability density function in the interval  $[-2, 10]$ . For  $y = 2x - 6$ , the conditional probability  $P(y \leq 7|x \geq 5)$  (rounded off to three decimal places) is \_\_\_\_\_.  
**[GATE-EC-2020:2M]**

**80. [NAT] [GATE-EC-2019:1M]**

If X and Y are random variables such that  $E[2X+Y]=0$  And  $E[X+2Y]=33$ , then  $E[X] + E[Y] = _____$ .  
**[GATE-EC-2019:1M]**

**81. [NAT] [GATE-EC-2019:1M]**

Let z be an exponential random variable with mean 1. That is, the cumulative distribution function of z is given by

$$F_z(x) = \begin{cases} 1 - e^{-x} & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$$

Then  $\Pr(z > 2|z > 1)$ , rounded off to two decimal places, is equal to \_\_\_\_\_.

**82. [MCQ] [GATE-ME-2019:1M]**

The lengths of a large stock of titanium rods follow a normal distribution with a mean ( $\mu$ ) of 440 mm and a standard deviation ( $\sigma$ ) of 1 mm. What is the percentage of rods whose lengths lie between 438 mm and 441 mm?

- (a) 81.85%
- (b) 68.4%
- (c) 99.75%
- (d) 86.64%

**83. [MCQ] [GATE-ME-2019:2M]**

The variable  $x$  takes a value between 0 and 10 with uniform probability distribution. The variable  $y$  takes a value between 0 and 20 with uniform probability distribution. The probability of the sum of variables ( $x + y$ ) being greater than 20 is

- (a) 0
- (b) 0.25
- (c) 0.33
- (d) 0.50

**84. [NAT] [GATE-ME-2019:1M]**

If  $x$  is the mean of data 3,  $x$ , 2 and 4, then the mode is \_\_\_\_\_.

**85. [MCQ] [GATE-CE-2019:2M]**

The probability density function of a continuous random variable distributed uniformly between  $x$  and  $y$  (for  $y > x$ ) is


- (a)  $y - x$
- (b)  $1/(y - x)$
- (c)  $x - y$
- (d)  $1/(x - y)$

**86. [NAT] [GATE-CS-2019:2M]**

Suppose  $Y$  is distributed uniformly in the open interval (1,6). The probability that the polynomial  $3x^2 + 6xY + 3Y + 6$  has only real roots is (rounded off to 1 decimal place) \_\_\_\_\_.

**87. [MCQ] [GATE-CE-2018:1M]**

The graph of a function  $f(x)$  is shown in the figure.




For  $f(x)$  to be valid probability density function, the value of  $h$  is

- (a)  $1/3$
- (b)  $2/3$
- (c)  $1$
- (d)  $3$

**88. [MCQ] [GATE-CE-2018:1M]**

A probability distribution with right skew is shown in the figure.



The correct statement for the probability distribution is

- (a) Mean is equal to mode
- (b) Mean is greater than median but less than mode.
- (c) Mean is greater than median and mode
- (d) Mode is greater than median

**89. [NAT] [GATE-EC-2018:1M]**

Let  $X_1, X_2, X_3$  and  $X_4$  be independent normal random variables with zero mean and unit variance. The probability that  $X_4$  is the smallest among the four is \_\_\_\_\_.

**90. [MCQ]****[GATE-ME-2018:2M]**

Let  $X_1, X_2$  be two independent normal random variables with means  $\mu_1, \mu_2$  and standard deviations  $\sigma_1, \sigma_2$ , respectively. Consider  $Y = X_1 - X_2$ ;  $\mu_1 = \mu_2 = 1, \sigma_1 = 1, \sigma_2 = 2$ . Then,

- (a)  $Y$  is normally distributed with mean 0 and variance 1
- (b)  $Y$  is normally distributed with mean 0 and variance 5
- (c)  $Y$  has mean 0 and variance 5, but is NOT normally distributed
- (d)  $Y$  has mean 0 and variance 1, but is NOT normally distributed

**91. [MCQ]****[GATE-ME-2018:2M]**

Let  $X_1$  and  $X_2$  be two independent exponentially distributed random variables with means 0.5 and 0.25, respectively. Then  $Y = \min(X_1, X_2)$  is

- (a) exponentially distributed with mean  $1/6$
- (b) exponentially distributed with mean 2
- (c) normally distributed with mean  $3/4$
- (d) normally distributed with mean  $1/6$

**92. [MCQ]****[GATE-ME-2017:1M]**

A sample of 15 data is as follows: 17, 18, 17, 17, 13, 18, 5, 5, 6, 7, 8, 9, 20, 17, 3. The mode of the data is

- (a) 4
- (b) 13
- (c) 17
- (d) 20

**93. [NAT]****[GATE-EE-2017:1M]**

Assume that in a traffic junction, the cycle of the traffic signal lights is 2 minutes of green (vehicle does not stop) and 3 minutes of red (vehicle stops). Consider that the arrival time of vehicles at the junction is uniformly distributed over 5 minute cycle. The expected waiting time (in minutes) for the vehicle at the junction is \_\_\_\_\_.

**94. [MCQ]****[GATE-EE-2017:2M]**

A person decides to toss a fair coin repeatedly until he gets a head. He will make at most 3 tosses. Let the random variable  $Y$  denote the number of heads. The value of  $\text{var}(Y)$  where  $\text{var}(\cdot)$  denotes the variance, equals:

- |                    |                      |
|--------------------|----------------------|
| (a) $\frac{7}{8}$  | (b) $\frac{49}{64}$  |
| (c) $\frac{7}{64}$ | (d) $\frac{105}{64}$ |

**95. [NAT]****[GATE-CS-2017:1M]**

Let  $x$  be a Gaussian random variable with mean 0 and variance  $\sigma^2$ . Let  $y = \max(x, 0)$  where  $\max(a, b)$  is the maximum of  $a$  and  $b$ . The median of  $y$  is \_\_\_\_\_.

**96. [MCQ]****[GATE-CS-2017:2M]**

For any discrete random variable  $X$ , with probability mass function  $P(x = j) = p_j, p_j \geq 0, j \in \{0, \dots, N\}$ , and  $\sum_{j=0}^N p_j = 1$ , define the polynomial

function  $g_x(z) = \sum_{j=0}^N P_j z^j$ . For a certain discrete

random variable  $Y$ , there exists a scalar  $\beta \in [0, 1]$  such that

$$g_y(z) = (1 - \beta + \beta z)^N.$$

The expectation of  $y$  is

- (a)  $N\beta(1 - \beta)$
- (b)  $N\beta$
- (c)  $N(1 - \beta)$
- (d) Not expressible in terms of  $N$  and  $\beta$  alone

**97. [MCQ]****[GATE-CE-2017:1M]**

The number of parameters in the univariate exponential and Gaussian distributions, respectively, are

- (a) 2 and 2
- (b) 1 and 2
- (c) 2 and 1
- (d) 1 and 1

**98. [MCQ]****[GATE-CE-2017:2M]**

For the function  $f(x) = a + bx, 0 \leq x \leq 1$ , to be a valid probability density function, which one of the following statements is correct?

- (a)  $a = 1, b = 4$
- (b)  $a = 0.5, b = 1$
- (c)  $a = 0, b = 1$
- (d)  $a = 1, b = -1$

**99. [NAT]****[GATE-CS-2017:2M]**

If a random variable  $X$  has a Poisson distribution with mean 5, then the expectation  $E[(X + 2)^2]$  equals \_\_\_\_\_.

**100. [NAT]****[GATE-CS-2016:1M]**

Suppose that a shop has an equal number of LED bulbs of two different types. The probability of an LED bulb lasting more than 100 hours given that it is of Type 1 is 0.7, and given that it is of Type 2 is 0.4. The probability that an LED bulb chosen uniformly at random lasts more than 100 hours is \_\_\_\_\_.

**101. [MCQ]****[GATE-CE-2016:1M]**

Type II error in hypothesis testing is

- (a) Acceptance of the null hypothesis when it is false and should be rejected
- (b) Rejection of the null hypothesis when it is true and should be accepted
- (c) Rejection of the null hypothesis when it is false and should be rejected
- (d) Acceptance of the null hypothesis when it is true and should be accepted

**102. [MCQ]****[GATE-CE-2016:2M]**

Probability density function of a random variable  $X$  is given below

$$f(x) = \begin{cases} 0.25 & \text{if } 1 \leq x \leq 5 \\ 0 & \text{otherwise} \end{cases}$$

$P(X \leq 4)$  is

- |                   |                   |
|-------------------|-------------------|
| (a) $\frac{3}{4}$ | (b) $\frac{1}{2}$ |
| (c) $\frac{1}{4}$ | (d) $\frac{1}{8}$ |

**103. [NAT]****[GATE-CE-2016:1M]**

The spot speeds (expressed in km/hr) observed at a road section are 66, 62, 45, 79, 32, 51, 56, 60, 53 and 49. The median speed (expressed in km/hr) is \_\_\_\_\_.

(Note: Answer with one decimal accuracy)

**104. [MCQ]****[GATE-CE-2016:1M]**

$X$  and  $Y$  are two random independent events. It is known that  $P(X) = 0.40$  and  $P(X \cup Y^C) = 0.7$ .

Which one of the following is the value of  $P(X \cup Y)$ ?

- |         |         |
|---------|---------|
| (a) 0.7 | (b) 0.5 |
| (c) 0.4 | (d) 0.3 |

**105. [MCQ]****[GATE-CE-2016:2M]**

If  $f(x)$  and  $g(x)$  are two probability density functions,

$$f(x) = \begin{cases} \frac{x}{a} + 1 & : -a \leq x < 0 \\ -\frac{x}{a} + 1 & : 0 \leq x \leq a \\ 0 & : \text{otherwise} \end{cases}$$

$$g(x) = \begin{cases} -\frac{x}{a} & : -a \leq x < 0 \\ \frac{x}{a} & : 0 \leq x \leq a \\ 0 & : \text{otherwise} \end{cases}$$

Which one of the following statements is true?

- (a) Mean of  $f(x)$  and  $g(x)$  are same; Variance of  $f(x)$  and  $g(x)$  are same
- (b) Mean of  $f(x)$  and  $g(x)$  are same; Variance of  $f(x)$  and  $g(x)$  are different
- (c) Mean of  $f(x)$  and  $g(x)$  are different; Variance of  $f(x)$  and  $g(x)$  are same
- (d) Mean of  $f(x)$  and  $g(x)$  are different; Variance of  $f(x)$  and  $g(x)$  are different

**106. [MCQ]****[GATE-ME-2016:1M]**

Consider a Poisson distribution for the tossing of a biased coin. The mean for this distribution is  $\mu$ . The standard deviation for this distribution is given by

- |                  |             |
|------------------|-------------|
| (a) $\sqrt{\mu}$ | (b) $\mu^2$ |
| (c) $\mu$        | (d) $1/\mu$ |

**107. [NAT]****[GATE-ME-2016:1M]**

The area (in percentage) under standard normal distribution curve of random variable Z within limits from  $-3$  to  $+3$  is \_\_\_\_\_.

**108. [MSQ]****[GATE-EE-2016:2M]**

Let the probability density function of a random variable, X, be given as:

$$f_x(x) = \frac{3}{2}e^{-3x}u(x) + ae^{4x}u(-x)$$

where  $u(x)$  is the unit step function.

Then the value of 'a' and Probability  $\{X \leq 0\}$ , respectively, are

- |                      |                      |
|----------------------|----------------------|
| (a) $2, \frac{1}{2}$ | (b) $4, \frac{1}{2}$ |
| (c) $2, \frac{1}{4}$ | (d) $4, \frac{1}{4}$ |

**109. [NAT]****[GATE-EC-2016:2M]**

Two random variables x and y are distributed according to

$$f_{xy}(x, y) = \begin{cases} (x+y), & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$


The probability  $P(x+y \leq 1)$  is \_\_\_\_\_.

**110. [NAT]****[GATE-CS-2016:1M]**

A probability density function on the interval  $[a, 1]$  is given by  $1/x^2$  and outside this interval the value of the function is zero. The value of a is \_\_\_\_\_.

**111. [MCQ]****[GATE-ME-2015:1M]**

Among the four normal distributions with probability density functions as shown below, which one has the lowest variance?

**112. [NAT]****[GATE-CE-2015:1M]**

Consider the following probability mass function (p.m.f) of a random variable X.

$$p(X, q) = \begin{cases} q & \text{If } X = 0 \\ 1-q & \text{If } X = 1 \\ 0 & \text{otherwise} \end{cases}$$

If  $q = 0.4$ , the variance of X is \_\_\_\_\_.

**113. [NAT]****[GATE-CE-2015:2M]**

The probability density function of a random variable, x is

$$f(x) = \begin{cases} \frac{x}{4}(4-x^2) & 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

The mean,  $\mu_x$  of the random variable is \_\_\_\_\_.

**114. [NAT]****[GATE-CE-2014:1M]**

The probability density function of evaporation E on any day during a year in a watershed is given by

$$f(E) = \begin{cases} \frac{1}{5} & 0 < E < 5 \text{ mm/day} \\ 0 & \text{otherwise} \end{cases}$$

The probability that E lies in between 2 and 4 mm/day in a day in the watershed is (in decimal) \_\_\_\_\_.

**115. [MCQ]****[GATE-CE-2014:1M]**

If  $\{x\}$  is a continuous, real valued random variable defined over the interval  $(-\infty, +\infty)$  and its occurrence is defined by the density function given as:

$$f(x) = \frac{1}{\sqrt{2\pi} b} e^{-\frac{1}{2}\left(\frac{x-a}{b}\right)^2}$$

where 'a' and 'b' are the statistical attributes of the random variable  $\{x\}$ . The value of the integral

$$\int_{-\infty}^a \frac{1}{\sqrt{2\pi} b} e^{-\frac{1}{2}\left(\frac{x-a}{b}\right)^2} dx \text{ is}$$

- |           |             |
|-----------|-------------|
| (a) 1     | (b) 0.5     |
| (c) $\pi$ | (d) $\pi/2$ |

**116. [NAT]****[GATE-CE-2014:2M]**

A traffic office imposes on an average 5 number of penalties daily on traffic violators. Assume that the number of penalties on different days is independent and follows a Poisson distribution. The probability that there will be less than 4 penalties in a day is \_\_\_\_\_.

**117. [NAT]****[GATE-CE-2014:2M]**

In any given year, the probability of an earthquake greater than magnitude 6 occurring in the Garhwal Himalayas is 0.04. The average time between successive occurrence of such earthquake is \_\_\_\_\_ (in year).

**118. [NAT]****[GATE-EE-2014:1M]**

Lifetime of an electric bulb is a random variable with density  $f(x) = kx^2$ , where  $x$  is measured in years. If the minimum and maximum lifetimes of bulb are 1 and 2 years respectively, then the value of  $k$  is \_\_\_\_\_.

**119. [NAT]****[GATE-ME-2014:1M]**

A nationalized bank has found that the daily balance available in its savings accounts follows a normal distribution with a mean of Rs. 500 and a standard deviation of Rs. 50. The percentage of savings account holders, who maintain an average daily balance more than Rs. 500 is \_\_\_\_\_.

**120. [NAT]****[GATE-ME-2014:2M]**

The number of accidents occurring in a plant in a month follows Poisson distribution with mean as 5.2. The probability of occurrence of less than 2 accidents in the plant during a randomly selected month is

- |           |           |
|-----------|-----------|
| (a) 0.029 | (b) 0.034 |
| (c) 0.039 | (d) 0.044 |

**121. [MCQ]****[GATE-ME-2014:2M]**

In the following table,  $x$  is a discrete random variable and  $p(x)$  is the probability density. The standard deviation of  $x$  is

|        |     |     |     |
|--------|-----|-----|-----|
| $x$    | 1   | 2   | 3   |
| $p(x)$ | 0.3 | 0.6 | 0.1 |

- |          |          |
|----------|----------|
| (a) 0.18 | (b) 0.36 |
| (c) 0.54 | (d) 0.6  |

**122. [MCQ]****[GATE-ME-2014:2M]**

A machine produces 0, 1 or 2 defective pieces in a day with associated probability of  $1/6$ ,  $2/3$  and  $1/6$ , respectively. The mean value and the variance of the number of defective pieces produced by the machine in a day, respectively, are

- |                 |                     |
|-----------------|---------------------|
| (a) 1 and $1/3$ | (b) $1/3$ and 1     |
| (c) 1 and $4/3$ | (d) $1/3$ and $4/3$ |

**123. [NAT]****[GATE-EC-2014:1M]**

Let  $X_1$ ,  $X_2$  and  $X_3$  be independent and identically distributed random variables with the uniform distribution on  $[0,1]$ . The probability  $p\{X_1 \text{ is the largest}\}$  is \_\_\_\_\_.

**124. [NAT]****[GATE-EC-2014:1M]**

Let  $X$  be a zero mean unit variance Gaussian random variable.  $E[|X|]$  is equal to \_\_\_\_\_.

**125. [MCQ]****[GATE-EC-2014:2M]**

Let  $X$  be a real-valued random variable with  $E[X]$  and  $E[X^2]$  denoting the mean values of  $X$  and  $X^2$ , respectively. The relation which always holds true is

- |                         |                            |
|-------------------------|----------------------------|
| (a) $(E[x])^2 > E[x^2]$ | (b) $E[x^2] \geq (E[x])^2$ |
| (c) $E[x^2] = (E[x])^2$ | (d) $E[x^2] > (E[x])^2$    |

**126. [NAT]****[GATE-EE-2014:1M]**

Let  $X$  be a random variable with probability density function

$$f(x) = \begin{cases} 0.2, & \text{for } |x| \leq 1 \\ 0.1, & \text{for } 1 < |x| \leq 4 \\ 0, & \text{otherwise} \end{cases}$$

The probability  $P(0.5 < X < 5)$  is

**127. [NAT]****[GATE-CS-2014:1M]**

Suppose you break a stick of unit length at a point chosen uniformly at random. Then the expected length of the shorter stick is \_\_\_\_\_.

**128. [MCQ]****[GATE-EE-2013:1M]**

A Continuous Random Variable  $X$  has a probability density function  $f(x) = e^{-x}$ ,  $0 < x < \infty$ . Then  $P\{X > 1\}$  is

- |           |         |
|-----------|---------|
| (a) 0.368 | (b) 0.5 |
| (c) 0.632 | (d) 1.0 |

**129. [MCQ]****[GATE-ME-2013:1M]**

Let  $X$  be a normal random variable with mean 1 and variance 4. The probability  $P\{X < 0\}$  is

- (a) 0.5
- (b) Greater than zero and less than 0.5
- (c) Greater than 0.5 and less than 1.0
- (d) 1.0

**130. [MCQ]****[GATE-EC-2013:2M]**

Consider two identically distributed zero-mean random variables  $U$  and  $V$ . Let the cumulative distribution functions of  $U$  and  $2V$  be  $F(x)$  and  $G(x)$  respectively. Then, for all values of  $x$

- (a)  $F(x) - G(x) \leq 0$
- (b)  $F(x) - G(x) \geq 0$
- (c)  $(F(x) - G(x))x \leq 0$
- (d)  $(F(x) - G(x)).x \geq 0$

**131. [MCQ]****[GATE-EC-2013:2M]**

Let  $U$  and  $V$  be two independent zero mean Gaussian random variables of variances  $\frac{1}{4}$  and  $\frac{1}{9}$  respectively. The probability  $P(3V \geq 2U)$  is

- |                   |                   |
|-------------------|-------------------|
| (a) $\frac{4}{9}$ | (b) $\frac{1}{2}$ |
| (c) $\frac{2}{3}$ | (d) $\frac{5}{9}$ |

**132. [NAT]****[GATE-CE-2013:2M]**

Find the value of  $\lambda$  such that function  $f(x)$  is valid probability density function \_\_\_\_\_

$$f(x) = \begin{cases} \lambda(x-1)(2-x) & \text{for } 1 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

**133. [MCQ]****[GATE-CS-2013:1M]**

Suppose  $P$  is the number of cars per minute passing through a certain road junction between 5 PM and 6 PM, and  $P$  has a Poisson distribution with mean 3. What is the probability of observing fewer than 3 cars during any given minute in this interval?

- |                         |                         |
|-------------------------|-------------------------|
| (a) $\frac{8}{(3e^3)}$  | (b) $\frac{9}{(2e^3)}$  |
| (c) $\frac{17}{(2e^3)}$ | (d) $\frac{26}{(2e^3)}$ |

**134. [MCQ]****[GATE-CE-2012:1M]**

The annual precipitation data of a city is normally distributed with mean and standard deviation as 1000 mm and 200 mm, respectively. The probability that the annual precipitation will be more than 1200 mm is \_\_\_\_\_

- (a) < 50%
- (b) 50%
- (c) 75%
- (d) 100%

**135. [MCQ]****[GATE-EE-2012:1M]**

Two independent random variables  $X$  and  $Y$  are uniformly distributed in the interval  $[-1, 1]$ . The probability that  $\max[X, Y]$  is less than  $\frac{1}{2}$  is

- (a)  $\frac{3}{4}$
- (b)  $\frac{9}{16}$
- (c)  $\frac{1}{4}$
- (d)  $\frac{2}{3}$

**136. [MCQ]****[GATE-EC-2012:1M]**

Two independent random variables  $x$  and  $y$  are uniformly distributed in the interval  $[-1, 1]$ . The probability that  $\max[x, y]$  is less than  $\frac{1}{2}$  is

- (a)  $\frac{3}{4}$
- (b)  $\frac{9}{16}$
- (c)  $\frac{1}{4}$
- (d)  $\frac{2}{3}$

**137. [MCQ]****[GATE-CS-2012:1M]**

Consider a random variable  $X$  that takes values +1 and -1 with probability 0.5 each. The values of the cumulative distribution function  $F(x)$  at  $x = -1$  and +1 are

- (a) 0 and 0.5
- (b) 0 and 1
- (c) 0.5 and 1
- (d) 0.25 and 0.75

**138. [MCQ]****[GATE-CS-2012:2M]**

Suppose a fair six-sided die is rolled once. If the value on the die is 1, 2, or 3, the die is rolled a second time. What is the probability that the sum total of values that turn up is at least 6?

- |                     |                    |
|---------------------|--------------------|
| (a) $\frac{10}{21}$ | (b) $\frac{5}{12}$ |
| (c) $\frac{2}{3}$   | (d) $\frac{1}{6}$  |

**139. [MCQ]****[GATE-CS-2011:2M]**

Consider a finite sequence of random values  $X = [x_1, x_2, \dots, x_n]$ . Let  $\mu_x$  be the mean and  $\sigma_x$  be the standard deviation of  $X$ . Let another finite sequence  $Y$  of equal length be derived from this as  $y_i = a * x_i + b$ , where  $a$  and  $b$  are positive constants. Let  $\mu_y$  be the mean and  $\sigma_y$  be the standard deviation of this sequence. Which one of the following statements is INCORRECT?

- (a) Index position of mode of  $X$  in  $X$  is the same as the index position of mode of  $Y$  in  $Y$
- (b) Index position of median of  $X$  in  $X$  is the same as the index position of median of  $Y$  in  $Y$ .
- (c)  $\mu_y = a\mu_x + b$
- (d)  $\sigma_y = a\sigma_x + b$

**140. [MCQ]****[GATE-CS-2011:1M]**

If the difference between the expectation of the square of a random variable ( $E[X^2]$ ) and the square of the expectation of the random variable ( $E[X]$ )<sup>2</sup> is denoted by  $R$ , then?

- (a)  $R = 0$
- (b)  $R < 0$
- (c)  $R \geq 0$
- (d)  $R > 0$

**141. [MCQ]****[GATE-CS-2011:2M]**

A deck of 5 cards (each carrying a distinct number from 1 to 5) is shuffled thoroughly. Two cards are then removed one at a time from the deck. What is the probability that the two cards are selected with the number on the first card being one higher than the number on the second card?

- (a)  $1/5$
- (b)  $4/25$
- (c)  $1/4$
- (d)  $2/5$

**142. [MCQ]****[GATE-CS-2010:2M]**

Consider a company that assembles computers. The probability of a faulty assembly of any computer is  $p$ . The company therefore subjects each computer to a testing process. This testing process gives the correct result for any computer with a probability of  $q$ . What is the probability of a computer being declared faulty?

- (a)  $pq + (1-p)(1-q)$
- (b)  $(1-q)p$
- (c)  $(1-p)q$
- (d)  $pq$

**143. [MCQ]****[GATE-EC-2009:2M]**

Consider two independent random variables  $x$  and  $y$  with identical distributions. The variables  $x$  and  $y$  take values 0, 1 and 2 with probability  $1/2$ ,  $1/4$  and  $1/4$  respectively. What is the conditional probability  $P(x + y = 2 | x - y = 0)$ ?

- (a) 0
- (b)  $1/16$
- (c)  $1/6$
- (d) 1

**144. [MCQ]****[GATE-EC-2009:2M]**

A discrete random variable  $x$  takes values from 1 to 5 with probabilities as shown in the table. A student calculates the mean of  $x$  as 3.5 and her teacher calculates the variance to  $x$  as 1.5. Which of the following statement is true?

| $k$      | 1   | 2   | 3   | 4   | 5   |
|----------|-----|-----|-----|-----|-----|
| $P(x=k)$ | 0.1 | 0.2 | 0.4 | 0.2 | 0.1 |

- (a) Both the student and the teacher are right.
- (b) Both the student and the teacher are wrong.
- (c) The student is wrong but the teacher is right.
- (d) The student is right but the teacher is wrong.

**145. [MCQ]****[GATE-ME-2009:2M]**

The standard deviation of a uniformly distributed random variable between 0 and 1 is

- (a)  $\frac{1}{\sqrt{12}}$
- (b)  $\frac{1}{\sqrt{3}}$
- (c)  $\frac{5}{\sqrt{12}}$
- (d)  $\frac{7}{\sqrt{12}}$

**146. [MCQ]****[GATE- CE-2009:2M]**

The standard normal probability function can be approximate as

$$F(X_N) = \frac{1}{1 + \exp(-1.7255X_N|X_N|^{0.12})}$$

where  $X_N$  = standard normal deviate. If mean and standard deviation of annual precipitation are 102 cm and 27 cm respectively, the probability that the annual precipitation will be between 90 cm and 102 cm is

- (a) 66.7%
- (b) 33.3%
- (c) 50.0%
- (d) 16.7%

**147. [MCQ]****[GATE-CS-2009:2M]**

An unbalanced dice (with 6 faces, numbered from 1 to 6) is thrown. The probability that the face value is odd is 90 % of the probability that the face value is even. The probability of getting any even numbered face is the same.

If the probability that the face is even given that it is greater than 3 is 0.75, which one of the following options is closest to the probability that the face value exceeds 3?

- |           |           |
|-----------|-----------|
| (a) 0.453 | (b) 0.468 |
| (c) 0.485 | (d) 0.492 |

**148. [MCQ]****[GATE- CE-2008:2M]**

If probability density function of a random variable X is  $f(x) = x^2$  for  $-1 \leq x \leq 1$  and  
 $= 0$  for any other values of x

then, the percentage probability  $P(-1/3 \leq x \leq 1/3)$  is

- |           |          |
|-----------|----------|
| (a) 0.247 | (b) 2.47 |
| (c) 24.7  | (d) 247  |

**149. [MCQ]****[GATE-CS-2008:2M]**

$$\text{Let } P = \sum_{\substack{1 \leq i \leq 2k \\ i \text{ odd}}} i \text{ and } Q = \sum_{\substack{1 \leq i \leq 2k \\ i \text{ even}}} i \text{ where } k \text{ is a positive integer. Then}$$

- |                 |                  |
|-----------------|------------------|
| (a) $P = Q - k$ | (b) $P = Q + k$  |
| (c) $P = Q$     | (d) $P = Q + 2k$ |

**150. [MCQ]****[GATE-CS-2008:2M]**

Let X be a random variable following normal distribution with mean +1 and variance 4. Let Y be another normal variable with mean -1 and variance unknown. If  $P(X \leq -1) = P(Y \geq 2)$ , the standard deviation of Y is

- |                |       |
|----------------|-------|
| (a) 3          | (b) 2 |
| (c) $\sqrt{2}$ | (d) 1 |

**151. [MCQ]****[GATE-EE-2008:1M]**

X is a uniformly distributed random variable that takes values between 0 and 1. The value of  $E(X^3)$  will be

- |           |           |
|-----------|-----------|
| (a) 0     | (b) $1/8$ |
| (c) $1/4$ | (d) $1/2$ |

**152. [MCQ]****[GATE-EC-2008:2M]**

$P_x(x) = M \exp(-2|x|) + N \exp(-3|x|)$  is the probability density function for the real random variable X, over the entire x-axis. M and N are both positive real numbers. The equation relating M and N is

- |                            |                             |
|----------------------------|-----------------------------|
| (a) $M + \frac{2}{3}N = 1$ | (b) $2M + \frac{1}{3}N = 1$ |
| (c) $M + N = 1$            | (d) $M + N = 3$             |

**Correlation & Regression****153. [NAT]****[GATE-CE-2022:2M]**

A set of observations of independent variable (x) and the corresponding dependent variable (y) is given below.

|   |    |    |    |    |
|---|----|----|----|----|
| x | 5  | 2  | 4  | 3  |
| y | 16 | 10 | 13 | 12 |

Based on the data, the coefficient a of the linear regression model  $y = a + bx$  is estimated as 6.1. The coefficient b is \_\_\_\_\_.  
*(Round off to one decimal place)*

**154. [MCQ]****[GATE-CS-2021:2M]**

Consider the two statements.

S<sub>1</sub>: There exist random variables X and Y such that  $(\mathbb{E}[(X - \mathbb{E}(X))(Y - \mathbb{E}(Y))])^2 > \text{Var}[X] \text{Var}[Y]$

S<sub>2</sub>: For all random variables X and Y,

$$\text{Cov}[X, Y] = \mathbb{E} [|X - \mathbb{E}[X]| |Y - \mathbb{E}[Y]|]$$

Which one of the following choices is correct?

- |   |
|---|
| (a) Both S <sub>1</sub> and S <sub>2</sub> are true     |
| (b) S <sub>1</sub> is true, but S <sub>2</sub> is false |
| (c) S <sub>1</sub> is false, but S <sub>2</sub> is true |
| (d) Both S <sub>1</sub> and S <sub>2</sub> are false.   |

**155. [MCQ]****[GATE-2008-CE:2M]**

Three values of x and y are to be fitted in a straight line in the form  $y = a + bx$  by the method of least squares. Given:  $\Sigma x = 6$ ,  $\Sigma y = 21$ ,  $\Sigma x^2 = 14$  and  $\Sigma xy = 46$ , the value of a and b are respectively.

- |             |             |
|-------------|-------------|
| (a) 2 and 3 | (b) 1 and 2 |
| (c) 2 and 1 | (d) 3 and 2 |


**ANSWER KEY**

- |      |                  |      |                  |      |                  |      |                |
|------|------------------|------|------------------|------|------------------|------|----------------|
| 1.   | (0.9 to 0.9)     | 2.   | (0.187 to 0.188) | 3.   | (0.018 to 0.020) | 4.   | (c)            |
| 5.   | (0.190 to 0.200) | 6.   | (0.90 to 0.95)   | 7.   | (0.041 to 0.043) | 8.   | (0.16 to 0.18) |
| 9.   | (0.25 to 0.27)   | 10.  | (0.5 to 0.5)     | 11.  | (b)              | 12.  | (c)            |
| 13.  | (0.021 to 0.024) | 14.  | (a)              | 15.  | (3.5 to 3.5)     | 16.  | (0.75 to 0.75) |
| 17.  | (0.5 to 0.5)     | 18.  | (a)              | 19.  | (0.027 to 0.028) | 20.  | (2.4 to 2.6)   |
| 21.  | (0.027 to 0.028) | 22.  | (0.2 to 0.2)     | 23.  | (0.39 to 0.43)   | 24.  | (a)            |
| 25.  | (0.07 to 0.08)   | 26.  | (0.33 to 0.34)   | 27.  | (b)              | 28.  | (0.96 to 0.98) |
| 29.  | (b)              | 30.  | (c)              | 31.  | (1.5)            | 32.  | (c)            |
| 33.  | (6)              | 34.  | (d)              | 35.  | (d)              | 36.  | (a)            |
| 37.  | (0.64 to 0.66)   | 38.  | (0.25 to 0.27)   | 39.  | (b)              | 40.  | (0.65 to 0.68) |
| 41.  | (2.9 to 3.1)     | 42.  | (c)              | 43.  | (0.13 to 0.15)   | 44.  | (0.25 to 0.25) |
| 45.  | (d)              | 46.  | (d)              | 47.  | (d)              | 48.  | (c)            |
| 49.  | (c)              | 50.  | (d)              | 51.  | (c)              | 52.  | (c)            |
| 53.  | (d)              | 54.  | (c)              | 55.  | (a)              | 56.  | (a)            |
| 57.  | (c)              | 58.  | (c)              | 59.  | (b)              | 60.  | (c)            |
| 61.  | (d)              | 62.  | (a)              | 63.  | (a)              | 64.  | (c)            |
| 65.  | (0.99 to 1.01)   | 66.  | (a)              | 67.  | (4)              | 68.  | (a)            |
| 69.  | (b)              | 70.  | (0.17 to 0.19)   | 71.  | (d)              | 72.  | (d)            |
| 73.  | (0.35 to 0.39)   | 74.  | 15.00 to 16.00   | 75.  | (b)              | 76.  | (0.5 to 0.5)   |
| 77.  | (-1.00 to -0.98) | 78.  | (Marks to all)   | 79.  | (0.299 to 0.301) | 80.  | (11 to 11)     |
| 81.  | (0.36 to 0.38)   | 82.  | (a)              | 83.  | (b)              | 84.  | (3 to 3)       |
| 85.  | (b)              | 86.  | (0.8 to 0.8)     | 87.  | (a)              | 88.  | (c)            |
| 89.  | (0.25)           | 90.  | (b)              | 91.  | (a)              | 92.  | (c)            |
| 93.  | (0.9)            | 94.  | (c)              | 95.  | (0.0 to 0.0)     | 96.  | (b)            |
| 97.  | (b)              | 98.  | (b)              | 99.  | (54 to 54)       | 100. | (0.55 to 0.55) |
| 101. | (a)              | 102. | (a)              | 103. | (54.49 to 54.51) | 104. | (a)            |
| 105. | (b)              | 106. | (a)              | 107. | (99.6 to 99.8)   | 108. | (a)            |
| 109. | (0.33)           | 110. | (0.5 to 0.5)     | 111. | (d)              | 112. | (0.23 to 0.25) |
| 113. | (1.06 to 1.07)   | 114. | (0.4 to 0.4)     | 115. | (b)              | 116. | (0.26 to 0.27) |
| 117. | (25)             | 118. | (0.4 to 0.5)     | 119. | (49 to 51)       | 120. | (b)            |
| 121. | (d)              | 122. | (a)              | 123. | (0.32 to 0.34)   | 124. | (0.79 to 0.81) |
| 125. | (b)              | 126. | (0.35 to 0.45)   | 127. | (0.24 to 0.27)   | 128. | (a)            |
| 129. | (b)              | 130. | (d)              | 131. | (b)              | 132. | (6 to 6)       |
| 133. | (c)              | 134. | (a)              | 135. | (b)              | 136. | (b)            |
| 137. | (c)              | 138. | (b)              | 139. | (d)              | 140. | (c)            |
| 141. | (a)              | 142. | (a)              | 143. | (c)              | 144. | (b)            |
| 145. | (a)              | 146. | (d)              | 147. | (b)              | 148. | (b)            |
| 149. | (a)              | 150. | (a)              | 151. | (c)              | 152. | (a)            |
| 153. | (1.9 to 1.9)     | 154. | (d)              | 155. | (d)              |      |                |

## SOLUTIONS

### 1. (0.9 to 0.9)

**Given:**  $P(A) = 0.5$  and  $P(B) = 0.8$

Probability of occurrence of at least A or B

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= P(A) + P(B) - P(A) \cdot P(B) \end{aligned}$$

[ $\because$  A and B are independent events]

$$= 0.5 + 0.8 - (0.5 \times 0.8) = 0.9$$



Scan for Video solution



### 2. (0.187 to 0.188)

Let  $x \rightarrow$  Number of trials for first occurrence of a 'Head'

|      |      |       |        |      |
|------|------|-------|--------|------|
| x    | 1(H) | 2(TH) | 3(TTH) | .... |
| P(x) | P    | qp    | qqp    | .... |

p  $\rightarrow$  Head

q  $\rightarrow$  Tail

$$p + q = 1$$

$$\text{Given } E(x) = 4$$

$$\begin{aligned} E(x) &= \sum_{i=1}^n x P(x) = 1p + 2qp + 3q^2p + \dots \\ &= p(1 + 2q + 3q^2 + \dots) \\ &= p(1 - q)^{-2} \\ &= p(p)^{-2} = \frac{1}{p} = 4 \end{aligned}$$

$$\therefore p = \frac{1}{4} \text{ and } q = 1 - p = \frac{3}{4}$$

P (Getting Head in second trial)

$$= qp = \frac{3}{4} \cdot \frac{1}{4} = \frac{3}{16} = 0.1875 = 0.188$$



Scan for Video solution



### 3. (0.018 to 0.020)

Favorable outcomes for (sum = 4)

$$S = (1, 3) (2, 2) (3, 1)$$

Let  $x \rightarrow$  Event of getting (sum = 4)

$$P(x) = \frac{n(x)}{n(S)} = \frac{3}{36} = \frac{1}{12} \quad (\text{sum} = 4)$$

$$\therefore P(\bar{x}) = 1 - \frac{1}{12} = \frac{11}{12} \quad (\text{Sum} \neq 4)$$

Required probability is

$$\begin{aligned} P(x \bar{x} x) + P(x \bar{x} \bar{x}) + P(\bar{x} x \bar{x}) \\ = \frac{1}{12} \cdot \frac{1}{12} \cdot \frac{11}{12} + \frac{1}{12} \cdot \frac{11}{12} \cdot \frac{1}{12} + \frac{11}{12} \cdot \frac{1}{12} \cdot \frac{1}{12} \\ = \frac{3 \times 11}{12 \times 12 \times 12} = 0.019 \end{aligned}$$



Scan for Video solution



### 4. (c)

Let the probability of head is 'p'

The probability of tail is 'q'

Let head comes in  $r^{\text{th}}$  toss, then the favourable cases will be

H, TH, TTH, TTTH....( $r = 1, 2, 3, 4 \dots$ )

Probability distribution

|      |   |    |        |        |            |
|------|---|----|--------|--------|------------|
| x    | 1 | 2  | 3      | 4      | ...r       |
| P(x) | p | qp | $q^2p$ | $q^3p$ | $q^{r-1}p$ |

$$E(x) = \sum_{i=1}^r i p(x)$$

$$= p + 2qp + 3q^2p + 4q^3p + \dots + rq^{r-1}p$$

$$= p(1 + 2q + 3q^2 + 4q^3 + \dots + r q^{r-1})$$

$$= p(1 - q)^{-2} = \frac{p}{p^2} = \frac{1}{p} \quad [\text{By binomial expansion}]$$



Scan for Video solution



**5. (0.190 to 0.200)****Given:**  $n = 15$ 

$$p = P(H) = \frac{1}{2}, \quad q = P(T) = \frac{1}{2}$$

P (Probability of getting 8 Heads in 15 times)

$$P(x = 8) = {}^{15}C_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^{15-8} = 0.196$$



Scan for Video solution

**6. (0.90 to 0.95)****Given**

5 → Men

3 → Women

P (at least one women)

$$= 1 - P(\text{no woman})$$

$$= 1 - \frac{5c_4}{8c_4} = 1 - \frac{5}{\binom{8 \times 7 \times 6 \times 5}{4 \times 3 \times 2 \times 1}}$$

$$= 1 - \frac{5}{70} = \frac{65}{70} = 0.9285 \approx 0.93$$

$$\therefore P(\text{at least one woman}) = 0.93$$



Scan for Video solution

**7. (0.041 to 0.043)****Given**

A fair coin is tossed 20 times.

$$\Rightarrow \text{Number of trials } n = 20$$

$$\text{Let } p \rightarrow \text{Probability of getting a head in a toss} = \frac{1}{2}$$

$$q \rightarrow \text{Probability of getting a tail in a toss} = \frac{1}{2}$$

P (4 heads in 1<sup>st</sup> 10 tosses ∩ 4 tails in next 10 tosses)  
 $= P(\text{4 heads in 1<sup>st</sup> 10 tosses}) \times P(\text{4 tails in next 10 tosses})$

$$= \left({10}_C_4 \cdot p^4 \cdot q^6\right) \cdot \left({10}_C_4 \cdot q^4 \cdot p^6\right)$$

$$= \left({10}_C_4\right)^2 \cdot (pq)^{10} = \left(\frac{10 \times 9 \times 8 \times 7}{4 \times 3 \times 2 \times 1}\right)^2 \times \left(\frac{1}{4}\right)^{10}$$

$$= (210)^2 \times \frac{1}{4^{10}} = 0.042$$

$\therefore \text{Required Probability} = 0.042$



Scan for Video solution

**8. (0.16 to 0.18)****Given**

Probability of defective = 0.05 = p(say)

Let 'x' be the number of defective parts and 'q' be the probability of non-defective.

$$\Rightarrow q = 1 - p = 0.95$$

$$\begin{aligned} P(x \geq 2) &= 1 - P(x = 0) - P(x = 1) \\ &= 1 - {}^{15}C_0 \cdot p^0 \cdot q^{15} - {}^{15}C_1 \cdot p^1 \cdot q^{14} \\ &= 1 - q^{14} ({}^{15}C_0 \cdot q + {}^{15}C_1 \cdot p) \\ &= 1 - (0.95)^{14} (0.95 + 15(0.05)) \\ &= 1 - 0.829 = 0.171 \end{aligned}$$

$$\therefore P(\text{at least 2 defective}) = 0.17$$



Scan for Video solution

**9. (0.25 to 0.27)**

- probability of resistor being defective is  $p = 0.02$
- Number of resistors ( $n$ ) = 50

Approximate poisson distribution mean

$$(\mu) = np = 50 \times 0.02 = 1$$

$$P(x = r) = \frac{e^{-\mu} \mu^r}{r!}$$

$$p(\text{2 or more defective}) = 1 - p(\text{no resistor defective}) - p(\text{1 resistor defective})$$

$$= 1 - p(r = 0) - p(r = 1)$$

$$= 1 - \frac{e^{-\mu} \cdot \mu^0}{0!} - \frac{e^{-\mu} \mu^1}{1!}$$

$$= 1 - e^{-1} - e^{-1}$$

$$= 1 - 2e^{-1}$$

$$= 0.26$$




Scan for Video solution



**10. (0.5 to 0.5)**

**Given:** Picking 3 red balls consecutively without replacement



$$P(R_1 \cap R_2 \cap R_3)$$

$$= P(R_1) \cdot P(R_2 / R_1) \cdot P(R_3 / R_1 \cap R_2)$$

$$= \frac{5}{6} \cdot \frac{4}{5} \cdot \frac{3}{4} = 0.5$$



Scan for Video solution

**11. (b)**

**Given,**

4 → Red

4 → Green

4 → Blue

Three balls are pulled one after the other, without replacing

$$P(\text{Red} \cap \text{Red} \cap \text{Red}) = P(\text{Red in 1st draw}) \cdot P(\text{Red in 2nd draw}) \cdot P(\text{Red in 3rd draw})$$

$$= \frac{4}{12} \times \frac{3}{11} \times \frac{2}{10} = \frac{1}{55}$$

$$\therefore P(\text{Red} \cap \text{Red} \cap \text{Red}) = \frac{1}{55}$$



Scan for Video solution

**12. (c)**

Given that a six-faced fair dice is rolled

$$P(\text{getting one}) = \frac{1}{6}$$

Let P be the probability of getting one

q be the probability of not getting one

$$\Rightarrow P = \frac{1}{6} \text{ and } q = \frac{5}{6}$$

No. of Rolls = 5

Let 'x' be the number of times that ONE appears

$$\therefore P(x \geq 4) = P(x = 4) + P(x = 5)$$

$$= 5_{C_4} \cdot P^4 \cdot q + 5_{C_5} \cdot P^5 \cdot q^0$$

$$= 5 \cdot \left(\frac{1}{6}\right)^4 \left(\frac{5}{6}\right) + 1 \cdot \left(\frac{1}{6}\right)^5 = \left(\frac{1}{6}\right)^5 \{25 + 1\} = \frac{26}{6^5}$$

$$\therefore P(x \geq 4) = \frac{26}{6^5} = 0.00334 = 0.334\%$$

$$\therefore P(x \geq 4) = 0.33\%$$

Scan for Video solution

**13. (0.021 to 0.024)**

**Given:** A pair of dice is rolled.

Probability of getting same number (1,1), (2,2) .... (6,6)

$$P(\text{Tie}) = \frac{6}{36} = \frac{1}{6} \Rightarrow P(\text{Not a Tie}) = \frac{5}{6}$$

$$P(\text{Win in 3rd trials}) = P(\text{Tie} \cap \text{Tie} \cap \text{Not a Tie}) \\ = (P(\text{Tie}))^2 \cdot P(\text{Not a Tie})$$

$$= \left(\frac{1}{6}\right)^2 \cdot \left(\frac{5}{6}\right) = \frac{5}{216} = 0.023$$

$$\therefore P(\text{win in 3rd Trial}) = 0.023$$

Scan for Video solution

**14. (a)**

$$\text{Given: } P(P \text{ applies}) = \frac{1}{4}, P\left(\frac{P \text{ applies}}{Q \text{ applies}}\right) = \frac{1}{2}$$

$$\text{And } P\left(\frac{Q \text{ applies}}{P \text{ applies}}\right) = \frac{1}{3}$$

By multiplication theorem

$$P(P) \cdot P\left(\frac{Q}{P}\right) = P(Q) \cdot P\left(\frac{P}{Q}\right)$$

$$\Rightarrow \frac{1}{4} \times \frac{1}{3} = P(Q) \cdot \frac{1}{2}$$

$$\Rightarrow P(Q) = \frac{1}{6} \quad \Rightarrow P(P \cap Q) = \frac{1}{12}$$

$$\begin{aligned}
 P\left(\frac{P \text{ doesn't apply}}{Q \text{ doesn't apply}}\right) &= \frac{P(P' \cap Q')}{P(Q')} = \frac{1 - P(P \cup Q)}{1 - P(Q)} \\
 &= \frac{1 - \{P(Q) + P(P) - P(P \cap Q)\}}{1 - P(Q)} \\
 &= \frac{1 - \{1/6 + 1/4 - 1/12\}}{1 - 1/6} = \frac{1 - \{1/3\}}{5/6} = \frac{2/3}{5/6} = \frac{4}{5} \\
 \therefore P\left(\frac{P'}{Q'}\right) &= \frac{4}{5}
 \end{aligned}$$



Scan for Video solution


**15. (3.5 to 3.5)**

Let  $x$  be the number appeared when a dice is rolled

|        |       |       |       |       |       |       |
|--------|-------|-------|-------|-------|-------|-------|
| $x$    | 1     | 2     | 3     | 4     | 5     | 6     |
| $P(x)$ | $1/6$ | $1/6$ | $1/6$ | $1/6$ | $1/6$ | $1/6$ |

$$\begin{aligned}
 \text{Mean of the outcome} &= \sum_{i=1}^6 x_i P(x = x_i) \\
 &= \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) = \frac{7}{2} = 3.5
 \end{aligned}$$

$\therefore$  Mean value of outcomes = 3.5



Scan for Video solution


**16. (0.75 to 0.75)**

**Given**

Two coins are tossed simultaneously.

$$P(\text{at least one head}) = 1 - P(\text{No head})$$

$$= 1 - P(1^{\text{st}} \text{ coin tail} \cap 2^{\text{nd}} \text{ coin tail})$$

$$= 1 - (P(\text{tail}))^2 = 1 - \frac{1}{4} = 0.75$$

$\therefore P(\text{at least one head}) = 0.75$



Scan for Video solution


**17. (0.5 to 0.5)**

Probability of getting 3 heads (H, H, H)

$$= \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{8}$$

Probability of getting 4 Heads (H, H, H, H)


$$= \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{16}$$

Getting 4<sup>th</sup> Head when 3 heads are already realized.  
Conditional probability of getting next read after 3 Head is

$$P(4H/3H) = \frac{P(4H \cap 3H)}{P(3H)} = \frac{1/16}{1/8} = 0.5$$



Scan for Video solution


**18. (a)**


$P(\text{getting a red ball in 2}^{\text{nd}} \text{ draw})$

$$= P(R \cap R) + P(B \cap R)$$

$$= \frac{1}{2} \times \frac{4}{9} + \frac{1}{2} \times \frac{5}{9} = \frac{1}{2}$$



Scan for Video solution


**19. (0.027 to 0.028)**

**Given:** 3 fair dice are thrown simultaneously.

Total number of ways of outcomes =  $6 \times 6 \times 6 = 216$

For all the three dice to show the same number of dots i.e. the favorable outcomes are  $\{(1, 1, 1), (2, 2, 2), (3, 3, 3), (4, 4, 4), (5, 5, 5), (6, 6, 6)\}$

$$\begin{aligned}
 \therefore \text{Required probability} &= \frac{6}{216} = \frac{1}{36} \\
 &= 0.0277 \approx 0.028
 \end{aligned}$$



Scan for Video solution



Probability that all three dice show same number of dots is **0.028**.



Scan for Video solution



### 20. (2.4 to 2.6)

**Given:**

$$\text{Probability of getting a reservation} = 40\% \\ = 0.4 = p(\text{say})$$

$$\text{Probability of not getting a reservation} \\ = 1 - 0.4 = 0.6 = q \text{ (say)}$$

Let 'x' be the number of attempts made by the passenger

The probability distribution table is as shown below

|      |   |    |        |        |       |
|------|---|----|--------|--------|-------|
| x    | 1 | 2  | 3      | 4      | ..... |
| P(x) | p | qp | $q^2p$ | $q^3p$ | ..... |

∴ Average number of attempts

$$E(x) = \sum x \cdot P(x) = 1 \cdot p + 2 \cdot qp + 3 \cdot q^2p + 4 \cdot q^3p + \dots \\ = p(1 + 2q + 3q^2 + 4q^3 + \dots) \\ = p \cdot (1 - q)^{-2} = \frac{p}{(1 - q)^2} = \frac{p}{p^2} = \frac{1}{p} = \frac{1}{0.4} = 2.5$$

∴ Average number of attempts to get a seat is **2.5**.



Scan for Video solution



### 21. (0.027 to 0.028)

**Given:** 3 fair dice are thrown simultaneously.

Total number of ways of outcomes =  $6 \times 6 \times 6 = 216$

For all the three dice to show the same number of dots i.e. the favourable outcomes are  $\{(1, 1, 1), (2, 2, 2), (3, 3, 3), (4, 4, 4), (5, 5, 5), (6, 6, 6)\}$

∴ Required probability =  $= 0.0277 \approx 0.028$

Probability that all three dice show same number of dots is 0.028



Scan for Video solution



### 22. (0.2 to 0.2)

The number of ways in which all 3 pens are of same colour

$$\begin{matrix} \nearrow \text{Out of 4 colours} \\ = {}^4C_1 \\ \searrow \end{matrix} = 4$$

Choose 1 at a time

The number of ways in which all 3 pens are of different colour =  ${}^4C_3 = 4$

The number of ways in which 2 pens are of same colour and 3<sup>rd</sup> pen is of different colour.

$$= {}^4C_2 \cdot {}^2C_1 = 6 \times 2 = 12$$

Total number of ways 3 pens can be chosen  
 $= 4 + 4 + 12 = 20$  ways

P(All 3 pens are of same colour)

$$= \frac{\text{Favourable no. of ways}}{\text{Total no. of ways}} = \frac{4}{20} = \frac{1}{5} = 0.2$$

Scan for Video solution

### 23. (0.39 to 0.43)

Probability of a screw being defective =  $p(\text{say}) = 0.1$

Number of screws in a packet = 5

Let 'q' be probability of non-defective,  $q = 1 - p = 0.9$

If 'x' is the number of defective screws in a packet, then packet gets replaced if  $x \geq 1$

P(replacement) =  $P(x \geq 1)$

$$\begin{aligned} &= 1 - P(x = 0) \\ &= 1 - ({}^5C_0 \cdot P^0 \cdot q^5) \\ &= 1 - (1 \times 1 \times (0.9)^5) \\ &= 1 - (0.9)^5 = 0.40951 \end{aligned}$$

∴  $P(\text{packet being replaced}) = 0.41$

Scan for Video solution

### 24. (a)

**Given**

That '3' cards are drawn at random.

In a pack of cards, we have 4- queens, 4- kings, 4-jacks

P(picking a king, a queen, a jack)

$$\begin{aligned}
 &= \frac{\left( {}^4C_1 \times {}^4C_1 \times {}^4C_1 \right)}{{}^{52}C_3} \\
 &= \frac{4 \times 4 \times 4}{\binom{52 \times 51 \times 50}{3 \times 2 \times 1}} = \frac{16}{13 \times 17 \times 25} = \frac{16}{5525} \\
 \therefore \quad &\boxed{\text{Required probability} = \frac{16}{5525}}
 \end{aligned}$$



Scan for Video solution



### 25. (0.07 to 0.08)

Given  $P(\text{Head}) = 0.3$

$\Rightarrow P(\text{tail}) = 0.7$

$P(\text{1st head in 5th toss}) = P(\text{4 tails in 1st 4 tosses} \cap \text{head in 5th toss}) = (P(\text{Tail}))^4 P(\text{Head})$

$$= (0.7)^4 \cdot (0.3)$$

$$= 0.072$$

$$\therefore P(\text{1st head in 5th toss}) = 0.072$$



Scan for Video solution



### 26. (0.33 to 0.34)

$P(Y) = P(\text{Tail} \cap \text{Head}) + P(\text{Tail} \cap \text{Tail} \cap \text{Tail} \cap \text{Head}) + P(\text{Tail} \cap \text{Tail} \cap \text{Tail} \cap \text{Tail} \cap \text{Tail} \cap \text{Head}) + \dots$

Since all the tosses are independent

$$\begin{aligned}
 P(Y) &= P(T) \cdot P(H) + (P(T))^3 \cdot P(H) + (P(T))^5 \cdot P(H) \\
 &\quad + \dots
 \end{aligned}$$

$$= P(T) \cdot P(H) \{1 + P(T)^2 + P(T)^4 + \dots\}$$

$$= P(T) \cdot P(H) \left\{ \frac{1}{1 - (P(T))^2} \right\} = \frac{1}{2} \cdot \frac{1}{2} \left\{ \frac{1}{1 - 1/4} \right\}$$

$$= \frac{1}{4} \times \frac{4}{3} = 0.33$$

$$\Rightarrow \boxed{P(\text{output Y}) = 0.33}$$



Scan for Video solution



29. (b)

Given

$$P(\text{pass}) = 20\% = 0.2$$

$$P(\text{pass} \cap (> 90\%)) = 5\% = 0.05$$

$$\begin{aligned} P((> 90\%) | \text{pass}) &= \frac{P((> 90\%) \cap \text{pass})}{P(\text{pass})} \\ &= \frac{0.05}{0.2} = 0.25 \end{aligned}$$

$$\therefore P(\text{getting more than } 90\% | \text{pass}) = 0.25 = \frac{1}{4}$$



Scan for Video solution



30. (c)

Given

$$P(X) = \frac{1}{4}, \quad P(Y) = \frac{1}{3} \text{ and } P(X \cap Y) = \frac{1}{12}$$

$$\therefore P(Y/X) = \frac{P(Y \cap X)}{P(X)} = \frac{P(X \cap Y)}{P(X)} = \frac{\frac{1}{12}}{\frac{1}{4}} = \frac{1}{3}$$

$$\Rightarrow P(Y/X) = \frac{1}{3}$$



Scan for Video solution



31. (1.5)

Given

$X \rightarrow$  Number of times a coin is tossed until two consecutive heads

|      |     |      |                     |                    |       |
|------|-----|------|---------------------|--------------------|-------|
| X    | 2   | 3    | 4                   | 5                  | ..... |
| P(x) | p.p | qp.p | q <sup>2</sup> .p.p | q <sup>3</sup> p.p | ..... |

$\therefore$  Expectation of  $X = \sum x \cdot P(x)$

$$= 2p^2 + 3qp^2 + 4q^2p^2 + 5q^3p^2 + \dots$$

$$= p^2(2 + 3q + 4q^2 + 5q^3 + \dots)$$

Since

$$\therefore \frac{p^2}{q} (2q + 3q^2 + 4q^3 \dots) = \frac{p^2}{q} \left( \frac{1}{(1-q)^2} - 1 \right)$$

Since  $p = q = 0.5$

$\therefore$  Expectation of  $x = 0.5 \times 3 = 1.5$

$\therefore$  Average number of trials for getting two consecutive head = 1.5



Scan for Video solution



32. (c)

Given A, B are independent events.

$$1. \quad P(A \cap B) = P(A).P(B)$$

$$2. \quad P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A) \cdot P(B)}{P(B)} = P(A)$$

$$3. \quad P(A \cup B) = P(A) + P(B) - P(A \cap B) \\ = P(A) + P(B) - P(A) \cdot P(B)$$

$$4. \quad P(\bar{A} \cap \bar{B}) = P(\bar{A}) \cdot P(\bar{B})$$

$\therefore$  option (c) is correct.



Scan for Video solution



33. (6)

Given  $X \rightarrow$  Number of times the dice is rolled

$$P(3) = \frac{1}{6} \text{ and } P(3') = \frac{5}{6}$$

$$\therefore X = 1, 2, 3$$

$$P(3 \text{ on 1st roll}) = \frac{1}{6}$$

$$P(3 \text{ on 2nd roll}) = P(3' \cap 3) = \frac{5}{6} \times \frac{1}{6}$$

$$P(3 \text{ on 3rd roll}) = P(3' \cap 3' \cap 3) = \left(\frac{5}{6}\right)^2 \times \frac{1}{6} \text{ and so on}$$

$\therefore$  The probability distribution table

|      |               |                                  |   |   |       |
|------|---------------|----------------------------------|---|---|-------|
| X    | 1             | 2                                | 3   | 4   | ..... |
| P(3) | $\frac{1}{6}$ | $\frac{5}{6} \times \frac{1}{6}$ | $\left(\frac{5}{6}\right)^2 \times \frac{1}{6}$ | $\left(\frac{5}{6}\right)^3 \times \frac{1}{6}$ | ..... |

$\therefore$  Expectation of  $X = E[X]$

$$\begin{aligned}
 E[X] &= \left( \frac{1}{6} \times 1 \right) + \left( \frac{5}{6} \times \frac{1}{6} \times 2 \right) + \left( \left( \frac{5}{6} \right)^2 \times \frac{1}{6} \times 3 \right) + \\
 &\quad \left( \left( \frac{5}{6} \right)^3 \times \frac{1}{6} \times 4 \right) + \dots \\
 E[X] &= \frac{1}{6} \left[ 1 + \left( 2 \times \frac{5}{6} \right) + \left( 3 \times \left( \frac{5}{6} \right)^2 \right) + \left( 4 \times \left( \frac{5}{6} \right)^3 \right) + \dots \right] \\
 E[X] &= \frac{1}{6} \left( 1 - \frac{5}{6} \right)^{-2} \\
 \because (1-q)^{-2} &= 1 + 2q + 3q^2 + 4q^3 \dots
 \end{aligned}$$

$$E[X] = \frac{1}{6} \times \left( \frac{1}{6} \right)^{-2} = 6$$



Scan for Video solution

**34. (d)**

A → A wins if he gets a six

$$P(A) = 1/6$$

$$P(\bar{A}) = 1 - \frac{1}{6} = \frac{5}{6} \text{ (Probability of A not getting a six)}$$

$$\begin{aligned}
 P(\text{Probability of A winning}) &= P(A) + P(\bar{A}\bar{B}A) + P(\bar{A}\bar{B}\bar{A}\bar{B}A) + \dots \\
 &= \frac{1}{6} + \frac{5}{6} \cdot \frac{5}{6} \cdot \frac{1}{6} + \frac{5}{6} \cdot \frac{5}{6} \cdot \frac{5}{6} \cdot \frac{5}{6} \cdot \frac{1}{6} \\
 &= \frac{1}{6} \left[ 1 + \frac{25}{36} + \left( \frac{25}{36} \right)^2 + \dots \right] \\
 &= \frac{1}{6} \cdot \frac{1}{1 - \frac{25}{36}} = \frac{6}{11}
 \end{aligned}$$



Scan for Video solution

**35. (d)**

$$P(H_R H_S) = 0.28$$

$$P(T_R T_S) = 0.18$$

$$P(H_R T_S) = 0.30$$

$$P(T_R H_S) = 0.24$$

$$\begin{aligned}
 \therefore P(H_R) &= P(H_R H_S) + P(H_R T_S) \\
 &= 0.28 + 0.30 = 0.58
 \end{aligned}$$

$$\begin{aligned}
 \therefore P(T_R) &= P(T_R T_S) + P(T_R H_S) \\
 &= 0.18 + 0.24 = 0.42
 \end{aligned}$$

$$\begin{aligned}
 \therefore P(H_S) &= P(H_R H_S) + P(T_R H_S) \\
 &= 0.28 + 0.24 = 0.52
 \end{aligned}$$

$$\begin{aligned}
 \therefore P(T_S) &= P(T_R T_S) + P(H_R T_S) \\
 &= 0.18 + 0.30 = 0.18
 \end{aligned}$$

Since  $P(H_R) \neq P(T_R)$  &  $P(H_S) \neq P(T_S)$  $\therefore R$  and  $S$  are not fair coin.

Let us suppose coins tosses are independent

$$P(H_R H_S) = P(H_R) \cdot P(H_S)$$

$$\Rightarrow 0.58 \times 0.52 \neq 0.28$$

 $\therefore$  Coin tosses are dependent.

Scan for Video solution

**36. (a)**

Total number of parts = 25

Number of defective = 10

 $\Rightarrow$  Number of non-defective parts = 15

$$\begin{aligned}
 \therefore P(\text{both good}) &= \frac{^{15}C_2}{^{25}C_2} \\
 &= \frac{15 \times 14}{25 \times 24} = \frac{7}{20}
 \end{aligned}$$

$$\therefore P(\text{both good}) = \frac{7}{20}$$



Scan for Video solution



**37. (0.64 to 0.66)****Given**

A group consists of equal number of men and women

$$\Rightarrow P(\text{men}) = \frac{1}{2} \text{ and } P(\text{women}) = \frac{1}{2}$$

$$\text{Also, } P(\text{men} \cap \text{unemployed}) = 20\% = \frac{2}{10}$$

$$\text{And } P(\text{women} \cap \text{unemployed}) = 50\% = \frac{5}{10}$$

**P (Selecting Employed)**

$$\begin{aligned} &= P(\text{men} \cap \text{Employed}) + P(\text{Women} \cap \text{Employed}) \\ &= P(\text{men}) \cdot P(\text{Employed}) + P(\text{women}) \cdot P(\text{employed}) \\ &= \frac{1}{2} \cdot \left(1 - \frac{2}{10}\right) + \frac{1}{2} \cdot \left(1 - \frac{5}{10}\right) \\ &= \frac{1}{2} \left(\frac{8}{10}\right) + \frac{1}{2} \left(\frac{5}{10}\right) = \frac{13}{20} = 0.65 \\ \therefore & P(\text{Selecting Employed}) = 0.65 \end{aligned}$$



Scan for Video solution

**38. (0.25 to 0.27)**

Opposite faces of a dice are coloured same

$$P(\text{red}) = \frac{1}{3}; P(\text{green}) = \frac{1}{3}; P(\text{blue}) = \frac{1}{3}$$

 $P(\text{getting red at least twice}) = P(\text{exactly twice}) + P(\text{thrice})$ 

$$\begin{aligned} P(\text{exactly twice}) &= 3 \times P(\text{red} \cap \text{red} \cap (\text{blue} \cup \text{green})) \\ &= 3 \times P(\text{red}) \times P(\text{red}) \times P(\text{blue} \cup \text{green}) \\ &= 3 \times \frac{1}{3} \times \frac{1}{3} \times \frac{2}{3} = \frac{2}{9} \end{aligned}$$

$$P(\text{thrice}) = P(\text{red} \cap \text{red} \cap \text{red}) = (P(\text{red}))^3 = \frac{1}{27}$$

$$\therefore P(\text{red at least twice}) = \frac{2}{9} + \frac{1}{27} = \frac{7}{27} = 0.2592$$



Scan for Video solution

**39. (b)****Given:** A fair coin was tossed four times.

$$P(\text{obtaining 'Tail' when coin is tossed}) = \frac{n(E)}{n(S)}$$

$$n(E) = \{T\} = 1, \quad n(S) = \{H, T\} = 2$$

$$\Rightarrow P(E) = \frac{1}{2}$$

Scan for Video solution

**40. (0.65 to 0.68)**

Let the total families in society be 'x'.

$$\Rightarrow \text{No. of families with single child} = \frac{x}{2}$$

$$\text{No of families with two children} = \frac{x}{2}$$

 $\therefore \text{Total no of children in society} =$ 

$$= \left( \frac{x}{2} \times 1 + \frac{x}{2} \times 2 \right) = \frac{3x}{2}$$

$$\begin{aligned} \therefore p(\text{child having sibling}) &= \frac{\binom{\frac{x}{2} \times 2}{2}}{\binom{3x}{2}} = \frac{\frac{x}{2}}{\binom{3x}{2}} = \frac{2}{3} \\ &= 0.67 \end{aligned}$$

$$\therefore p(\text{child selected at random having a sibling}) = 0.67$$



Scan for Video solution

**41. (2.9 to 3.1)**

Given a coin is tossed repeatedly till both head and tail appear at least once.

Let  $x$  = Number of time,

T → Event of tail,

H → Event of tail

$$\therefore p(x = 2) = p(T \cap H) + P(H \cap T)$$

$$= P(T) \cdot P(H) + P(H) \cdot P(T) = 2 \cdot \left(\frac{1}{2}\right)^2$$

$$P(x = 3) = P(T \cap T \cap H) + P(H \cap H \cap T)$$

$$= P(T)^2 \cdot P(H) + (P(H))^2 \cdot P(T) = 2 \left(\frac{1}{2}\right)^3$$

$$P(x=4) = P(T \cap T \cap T \cap H) + P(H \cap H \cap H \cap T)$$

$$= (P(T))^3 \cdot P(H) + (P(H))^3 \cdot P(T) = 2 \cdot \left(\frac{1}{2}\right)^4$$

And so on

$\therefore$  Probability distribution table is

|      |                                      |                                      |                                      |                                      |       |
|------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|-------|
| x    | 2                                    | 3                                    | 4                                    | 5                                    | ..... |
| P(x) | $2 \cdot \left(\frac{1}{2}\right)^2$ | $2 \cdot \left(\frac{1}{2}\right)^3$ | $2 \cdot \left(\frac{1}{2}\right)^4$ | $2 \cdot \left(\frac{1}{2}\right)^5$ | ..... |

$$\therefore E(x) = \sum x \cdot P(x)$$

$$= 2 \times 2 \left(\frac{1}{2}\right)^2 + 3 \times 2 \left(\frac{1}{2}\right)^3 + 4 \times 2 \left(\frac{1}{2}\right)^4 + \dots$$

$$= 2 \times \left(\frac{1}{2}\right) + 3 \times \left(\frac{1}{2}\right)^2 + 4 \times \left(\frac{1}{2}\right)^3 + \dots + 1 - 1$$

$$= (1 + 2 \times \left(\frac{1}{2}\right) + 3 \times \left(\frac{1}{2}\right)^2 + 4 \times \left(\frac{1}{2}\right)^3 + \dots) - 1$$

$$\left( \because 1 + 2x + 3x^2 + \dots = (1-x)^{-2} \because |x| < 1 \right)$$

$$= \left(1 - \frac{1}{2}\right)^{-2} - 1 = 4 - 1 = 3$$

$\therefore$  Average number of tosses required = 3



Scan for Video solution



#### 42. (c)

Given coin is unbiased

$$\Rightarrow P(H) = P(T) = \frac{1}{2}$$

$P(4^{\text{th}} \text{ head in } 10^{\text{th}} \text{ toss}) = P(3 \text{ Heads in first 9 tosses} \cap \text{Head in } 10^{\text{th}} \text{ toss})$

$$= P(3 \text{ Heads in first 9 tosses}) \times P(\text{Head in } 10^{\text{th}} \text{ toss})$$

$$= {}^9C_3 \left(\frac{1}{2}\right)^3 \cdot \left(\frac{1}{2}\right)^6 \times \left(\frac{1}{2}\right)$$

$$= \frac{9 \times 8 \times 7}{3 \times 2 \times 1} \times \frac{1}{2^{10}} = \frac{84}{2^{10}} = 0.082$$

$$\therefore P(4^{\text{th}} \text{ Head in } 10^{\text{th}} \text{ toss}) = 0.082$$



Scan for Video solution



#### 43. (0.13 to 0.15)

Sample space = {1, 2, 3, 4, 5, 6 dots}

$$P(n \text{ dots}) \propto n$$

$$P(n \text{ dots}) = kn$$

Sum of all probabilities is 1.

$$k + 2k + 3k + 4k + 5k + 6k = 1$$

$$k = \frac{1}{21}$$

$\therefore$  Probability of dice showing 3 dots = 3k

$$= 3 \times \frac{1}{21} = \frac{1}{7} = 0.142$$



Scan for Video solution



#### 44. (0.25 to 0.25)

**Given:** A and B are mutually exclusive

$$\Rightarrow A \cap B = \emptyset \text{ and } A \cup B = S$$

By addition theorem,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\Rightarrow P(S) = P(A) + P(B) - 0$$

$$\Rightarrow P(A) + P(B) = 1$$

Since  $0 < P(A) < 1$  and  $0 < P(B) < 1$ ,

$$A.M \geq G.M$$

$$\Rightarrow \frac{P(A) + P(B)}{2} \geq \sqrt{P(A) \cdot P(B)}$$

$$\Rightarrow P(A) \cdot P(B) \leq \frac{1}{4} \cdot (P(A) + P(B))^2$$

$$\Rightarrow P(A) \cdot P(B) \leq \frac{1}{4} \cdot (1)^2$$

$\therefore$  Maximum value of  $P(A) \cdot P(B) = 0.25$



Scan for Video solution



45. (d)

**Given:**  $P(\text{know}) = \frac{2}{3} \Rightarrow P(\text{Guess}) = \frac{1}{3}$

And  $P(\text{correct/guess}) = \frac{1}{4}$

$P(\text{know/correct})$

$$= \frac{P(\text{know} \cap \text{correct})}{P(\text{know} \cap \text{correct}) + P(\text{guess} \cap \text{correct})}$$

$$= \frac{P(\text{know}) \cdot P(\text{correct / know})}{P(\text{know}) \cdot P(\text{correct / know}) + P(\text{guess}) \cdot P(\text{correct / guess})}$$

$$= \frac{\frac{2}{3} \cdot (1)}{\frac{2}{3}(1) + \frac{1}{3} \cdot \frac{1}{4}} = \frac{\frac{2}{3}}{\frac{2}{3} + \frac{1}{12}} = \frac{8}{9}$$

$$\therefore P(\text{know / correct}) = \frac{8}{9}$$



Scan for Video solution



46. (d)

**Given:** A box contains 4 red balls and 6 black balls.

**4 → red→ R**

**6 → black→ B**

$$P(1R \cap 2B) = P(R \cap B \cap B) + P(B \cap R \cap B) + P(B \cap B \cap R)$$

$$= P(R).P(B).P(B) + P(B).P(R).P(B) + P(B).P(B)$$

$$P(B).P(R)$$

$$= \left( \frac{4}{10} \times \frac{6}{9} \times \frac{5}{8} \right) + \left( \frac{6}{10} \times \frac{4}{9} \times \frac{5}{8} \right) + \left( \frac{6}{10} \times \frac{5}{9} \times \frac{4}{8} \right) = \frac{1}{2}$$

$$\therefore P(\text{red} \cap 2 \text{ black}) = \frac{1}{2}$$



Scan for Video solution



47. (d)

**Given:**

Since negative and positive value are equally likely, the distribution of number of negative values is binomial with  $n = 5$  and  $p = 1/2$ .

Let X represent number of negative values in 5 trials

$$X \rightarrow 0 \text{ to } 5, P(-ve) = p = 1/2, P(+ve) = q = 1/2$$

$$P(\text{at most 1 negative value}) = P(X = 0) + P(X = 1) = {}^5C_0 p^0 q^5 + {}^5C_1 p^1 q^4$$

$$= {}^5C_0 \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^5 + {}^5C_1 \left(\frac{1}{2}\right)^1 \left(\frac{1}{2}\right)^4$$

$$= \frac{1}{32} + \frac{5}{32} = \boxed{\frac{6}{32}}$$

Scan for Video solution



48. (c)

Let X be the random variable denoting number of tosses.

| X    | 1<br>(H)      | 3<br>(TTH)                   | 5<br>(TTTTH)                 | 7<br>(TTTTTTH)               |
|------|---------------|------------------------------|------------------------------|------------------------------|
| P(X) | $\frac{1}{2}$ | $\left(\frac{1}{2}\right)^3$ | $\left(\frac{1}{2}\right)^5$ | $\left(\frac{1}{2}\right)^7$ |

∴ The required probability =  $P(H) + P(TTH) + P(TTTTH) + \dots$

$$= \frac{1}{2} + \left(\frac{1}{2}\right)^3 + \left(\frac{1}{2}\right)^5 + \dots$$

$$= \frac{\frac{1}{2}}{1 - \frac{1}{4}} = \frac{2}{3}$$

Scan for Video solution



49. (c)

**Given:**

A fair coin is tossed till head appears for the first time  
Possibilities to get head in odd tosses are

H(1<sup>st</sup>), TTH(3<sup>rd</sup>), TTTTH(5<sup>th</sup>),.....

P(1<sup>st</sup> head in odd toss)

$$= P(H) + P(T \cap T \cap H) + P(T \cap T \cap T \cap T \cap H) + \dots$$



considering followable cases

For  $x = 1 \rightarrow \{(1, 2), (1, 3), (1, 4), (1, 5), (1, 6)\}$

$x = 2 \rightarrow \{(2, 3), (2, 4), (2, 5), (2, 6)\}$

$x = 3 \rightarrow \{(3, 4), (3, 5), (3, 6)\}$

$x = 4 \rightarrow \{(4, 5), (4, 6)\}$

$x = 5 \rightarrow \{(5, 6)\}$

No of favorable cases = 15

$$\text{Required probability} = \frac{15}{36} = \frac{5}{12}$$

$$P(2^{\text{nd}} \text{ roll value} > 1^{\text{st}} \text{ roll value}) = 5/12$$



Scan for Video solution



55. (a)

$P(\text{Both heads} | \text{at least 1 head})$

$$= \frac{P(\text{Both heads})}{P(\text{atleast one head})}$$

$$= \frac{P(H \cap H)}{P(T \cap H) + P(H \cap T) + P(H \cap H)}$$

$$= \frac{(P(H))^2}{2 \cdot P(T) \cdot P(H) + (P(H))^2} = \frac{\frac{1}{4}}{2\left(\frac{1}{4}\right) + \frac{1}{4}} = \frac{1}{3}$$

$$P(\text{Both Heads/at least one head}) = 1/3$$



Scan for Video solution



56. (a)

In general,

If  $N = p \times q \times r \times s \times \dots \times u$  (k terms) where p, q, r, s, ...u are prime factors.

The number of divisors of  $N^n = (n + 1)^k$

Number of divisors of  $10^{99} = (2 \times 5)^{99}$  is  
 $(99 + 1)^2 = 100^2 = 10^4$

Number of multiples of  $10^{96}$  that are factors of  $10^{99}$

$$= \text{number of divisors of } \left( \frac{10^{99}}{10^{96}} = 10^3 = (2 \times 5)^3 \right)$$

$$= (3+1)^2 = 16$$

$$\therefore \text{ Required probability} = \frac{16}{10^4} = \frac{16}{100 \times 100} = \frac{1}{625}$$



Scan for Video solution



57. (c)

**Given:** Two coins are tossed

Possible Outcomes = {HH, HT, TH, TT}

$$P(\text{2 Heads}) = \frac{\text{No of favourable outcomes } \{HH\}}{\text{No of total outcomes}}$$

$$\therefore \text{ The required probability} = \frac{1}{4}$$



Scan for Video solution



58. (c)

It is given that white ball is removed in first draw, therefore sample space reduces

$$\begin{matrix} 3W \\ 3R \end{matrix}$$

P(Probability of removing red ball in second draw)

$$= \frac{n(E)}{n(S)} = \frac{3}{6} = \frac{1}{2}$$



Scan for Video solution



59. (b)

P(at least two of them have same birthday)

= 1 - P(None of them have same birthday)

Assuming N = 2,

$$P = 1 - \frac{30}{30} \times \frac{29}{30} \not> \frac{1}{2}$$

Assuming  $N = 3$ ,

$$P = 1 - \frac{30}{30} \times \frac{29}{30} \times \frac{28}{30} > \frac{1}{2}$$

Assuming  $N = 7$ ,

$$\begin{aligned} P &= 1 - \frac{30}{30} \times \frac{29}{30} \times \frac{28}{30} \times \frac{27}{30} \times \frac{26}{30} \times \frac{25}{30} \times \frac{24}{30} \\ &= 0.5308 > \frac{1}{2} \quad \therefore N = 7 \end{aligned}$$



Scan for Video solution



60. (c)

**Given:** A fair coin is tossed 10 times.

$$P(1^{\text{st}} \text{ two heads} \cap \text{Next 8 tails})$$

$$\begin{aligned} &= P(H \cap H \cap T \cap T \cap \dots \cap T) \\ &= P(H) \cdot P(H) \cdot P(T) \cdot P(T) \dots (P(T) \text{ occurs 8 times}) \end{aligned}$$

$$= \frac{1}{2} \times \frac{1}{2} \times \left( \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \dots \times \frac{1}{2} \right)^8 = \left( \frac{1}{2} \right)^{10}$$

$$\therefore P(1^{\text{st}} \text{ two as heads} \cap 8 \text{ tails}) = \left( \frac{1}{2} \right)^{10}$$

$$P(\text{Number of heads} > \text{Number of tails}) = 5/16$$



Scan for Video solution



61. (d)

**Given:** Three coins are simultaneously tossed

$$P(\text{at least one head}) = 1 - P(\text{No head})$$

$$\begin{aligned} &= 1 - P(\text{tail} \cap \text{tail} \cap \text{tail}) \\ &= 1 - \{P(\text{tail}), P(\text{tail}), P(\text{tail})\} \\ &= 1 - (P(\text{tail}))^3 = 1 - \left( \frac{1}{2} \right)^3 = \frac{7}{8} \end{aligned}$$

$$\therefore P(\text{at least one head}) = 7/8$$



Scan for Video solution



62. (a)

**Given:** A coin is tossed four times

Let  $H \rightarrow$  denotes Head and  $T \rightarrow$  denotes Tail

Assuming all the tosses are independent

$$\begin{aligned} P(3 \text{ Heads}) &= P(H \cap H \cap H \cap T) + P(H \cap H \cap T \cap H) \\ &\quad + P(H \cap T \cap H \cap H) + P(T \cap H \cap H \cap H) \\ &= 4 \times \left( \frac{1}{2} \right)^4 = \frac{4}{16} = \frac{1}{4} \end{aligned}$$


$$\therefore \text{Probability of getting exactly 3 heads} = \frac{1}{4}$$



Scan for Video solution



63. (a)



Obviously, the probability of using a car = 0.45

$$\begin{aligned} \left( \text{The probability of using a bus} \right) &= \left( \text{Probability of using public transport} \right) \times \left( \text{Probability of choosing bus} \right) \\ &= (1 - 0.45) \times 0.55 \\ &= 0.3025 = \mathbf{0.30} \end{aligned}$$

$$\begin{aligned} \left( \text{The probability of using metro} \right) &= \left( \text{Probability of using public transport} \right) \times \left( \text{Probability of choosing a metro} \right) \\ &= (1 - 0.45)(1 - 0.55) \\ &= 0.2475 = \mathbf{0.25} \end{aligned}$$



Scan for Video solution



64. (c)

**Given:** The students studies either computer science (S) or mathematics (M) every day.

$$P(S \text{ and then } M) = 0.6$$

$$P(M \text{ and then } C) = 0.4$$

$$P(C \text{ on Wednesday} | C \text{ on Monday})$$

$$\begin{aligned} &= P(C \cap M \cap C) + P(C \cap C \cap C) \\ &= P(C \text{ and then } M) \cdot P(M \text{ and then } C) \\ &\quad + P(C \text{ and then } C) \cdot P(C \text{ and then } C) \\ &= (0.6 \times 0.4) + (0.4 + 0.4) = 0.24 + 0.16 = 0.40 \end{aligned}$$

$\therefore P(\text{computers on Wednesday}/\text{computers on Monday}) = 0.4$



Scan for Video solution



### 65. (0.99 to 1.01)

$$f(x) = ae^{-2|x|}$$

$$\therefore \int_{-\infty}^{+\infty} f(x) dx = 1 \quad [\text{for this function to be a valid probability density function.}]$$

$$\int_{-\infty}^{+\infty} ae^{-2|x|} dx = \int_{-\infty}^0 ae^{2x} dx + \int_0^{+\infty} ae^{-2x} dx = 1$$

$$\left[ \because |x| = \begin{cases} x & ; x \geq 0 \\ -x & ; x < 0 \end{cases} \right]$$

$$= \frac{a}{2} \left[ e^{2x} \right]_{-\infty}^0 + \frac{a}{-2} \left[ e^{-2x} \right]_0^{+\infty} = 1$$

$$= \frac{a}{2}(1-0) - \frac{a}{2}[0-1] = 1$$

$$\Rightarrow a = 1$$



Scan for Video solution



### 66. (a)

We know,

$$E[(x - \bar{x})^2] = \text{Var}$$

$$\text{Given } \bar{x} = 0, E(x^2) = \text{Var}$$

$$E(x_n^2) = \sigma_0^2$$

We have,

Standard deviation,

$$\sigma^2 = \frac{1}{N-1} \sum_{n=1}^N x_n^2 \Rightarrow \text{for sample}$$

$$= \frac{1}{N} \sum_{n=1}^N x_n^2 \Rightarrow \text{for population}$$

Here we took 10000 samples out of  $10^6$  numbers.

$$\therefore E(\hat{\sigma}_2^2) = \sigma_0^2 \text{ will be correct.}$$



Scan for Video solution



### 67. (4)

From the given graph:

|                      |   |   |   |    |    |    |   |   |   |   |    |
|----------------------|---|---|---|----|----|----|---|---|---|---|----|
| No. of wickets taken | 0 | 1 | 2 | 3  | 4  | 5  | 6 | 7 | 8 | 9 | 10 |
| No. of matches       | 5 | 7 | 8 | 25 | 20 | 17 | 8 | 4 | 3 | 2 | 1  |

Total number of matches

$$= 5 + 7 + 8 + 25 + 20 + 17 + 8 + 4 + 3 + 2 + 1 = 100$$

$\therefore$  Median is the average number of wickets taken in 50<sup>th</sup> and 51<sup>st</sup> match.

In the table above, data is arranged in ascending order.

$\therefore$  Number of wickets taken in 50<sup>th</sup> and 51<sup>st</sup> match are 4.

$$\therefore \text{Median} = \frac{4+4}{2} = 4$$

$\therefore$  Median number of wickets taken in a match = 4.



Scan for Video solution



### 68. (a)

Probability of a component being defective = 0.015

Number of components in a box = 200

$\Rightarrow$  Average number of (Mean of) defective components in a box =  $200 \times 0.015 = 3$

$$\Rightarrow \text{Mean} = \mu = 3$$





For Poisson distribution,

$$\text{Mean} = \text{Variance} = \mu = \sigma^2 = 3$$

$$\therefore \text{Mean} = \text{Variance} = 3$$



### 69. (b)





Scan for Video solution

**70. (0.17 to 0.19)**

Given 'X' follows Poisson distribution and

$$P(X=1) = P(X=2)$$

$$\Rightarrow \frac{e^{-\lambda} \cdot \lambda^1}{1!} = \frac{e^{-\lambda} \cdot \lambda^2}{2!} \Rightarrow \lambda = 2$$

$$\therefore P(X=3) = \frac{e^{-\lambda} \cdot \lambda^3}{3!} = \frac{e^{-2} \cdot (2)^3}{3!} = \frac{8}{6e^2}$$

$$\therefore P(x=3) = 0.180$$



Scan for Video solution

**71. (d)**

Binomial distribution is for finite samples.

When the number samples gets increased, the distribution can be approximated as continuous distribution.

So we can write the distribution as

$$Y = \sum_{i=1}^{\infty} X_i = \int X_i dx \text{ and } Y \text{ is normally distributed.}$$



Scan for Video solution

**72. (d)**

For a Binomial distribution with 'n' trials, if 'P' is the probability of success and  $q = 1 - P$  is the probability of failure, then

$$\text{Mean} = np, \quad \text{Variance} = npq = np(1-p)$$

$$\Rightarrow \text{Standard deviation} = \sqrt{npq} = \sqrt{np(1-p)}$$



Scan for Video solution

**73. (0.35 to 0.39)**

**Given:**  $P(x)$  is exponentially distribution with parameter '2'

$$\Rightarrow P(x) = \begin{cases} 2e^{-2x}; & x \geq 0 \\ 0; & x < 0 \end{cases}$$

Let 'x' be the life time of component

$$\text{Expected life time} = \text{Mean life time} = \frac{1}{\lambda} = \frac{1}{2}$$

$$P\left(x > \frac{1}{2}\right) = \int_{1/2}^{\infty} P(x) dx = \int_{1/2}^{\infty} 2e^{-2x} dx$$

$$= 2 \frac{e^{-2x}}{-2} \Big|_{1/2}^{\infty} = -1 \left( e^{-\infty} - e^{-2(1/2)} \right)$$

$$= e^{-1} = \frac{1}{e} = 0.3678 = 0.37 \because e^{-\infty} = 0$$

$$P(\text{life time exceeds expected life time}) = 0.37$$



Scan for Video solution

**74. (15.00 to 16.00)**

**Given:**

$$P(\text{Head}) = p = 0.4; \quad P(\text{Tail}) = q = 1 - 0.4 = 0.6$$

Let  $X \rightarrow$  Number of heads in 1000 tosses.

Given random variable follows Binomial distribution Since number of possible outcomes = 2  
For a random variable  $X$

Where  $n \rightarrow$  Total number of trials

$p \rightarrow$  Probability of success

$q \rightarrow$  Probability of failure.

$$\text{Standard deviation} = \sigma = \sqrt{npq}$$

$$= \sqrt{1000(0.4)(0.6)} = \sqrt{240} = 15.49$$




Scan for Video solution



**75. (b)**

The cumulative distribution function for the Gaussian distribution will be “S-shaped”.



Scan for Video solution

**76. (0.5 to 0.5)**

**Given:**

For  $n > 2$ ,  $a \in \{0, 1\}^n$

- ⇒ ‘a’ is a  $n \times 1$  vector with every element ‘0’ or ‘1’
- ⇒  $a = [a_{ij}] \forall 1 \leq i \leq n, j = 1$  where  $a_{ij} = 0$  (or) 1
- Also, ‘x’ is a  $n \times 1$  vector with element ‘0’ or ‘1’.
- ⇒  $x = [x_{ij}] \forall 1 \leq i \leq n; j = 1$  where  $x_{ij} = 0$  (or) 1.
- ∴  $a_{ij}, x_{ij}$  lies between 0 to ‘1’

$$\therefore \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_{ij} \text{ lies between 0 to } n$$

So, the probability distribution of  $\sum_{i=1}^n \sum_{j=1}^n a_{ij} x_{ij}$  is

$$\sum_{i=1}^n \sum_{j=1}^n a_{ij} x_{ij} \quad [0 \quad 1 \quad 2 \dots \quad n]$$

$$P\left(\sum_{i=1}^n \sum_{j=1}^n a_{ij} x_{ij}\right) \quad [n_{c_0} \quad n_{c_1} \quad n_{c_2} \quad n_{c_n}]$$

∴ Probability that  $a_{ij}, x_{ij}$  is odd number is

$$= \frac{n_{C_1} + n_{C_3} + \dots}{n_{C_0} + n_{C_1} + n_{C_2} + \dots} = \frac{2^{n-1}}{2^n} = \frac{1}{2} = 0.5$$



Scan for Video solution

**77. (-1.00 to -0.98)**

**Given**

X and Y are exponentially distributed with mean of 0.5.

$$\Rightarrow \mu_x = \mu_y = 0.5$$

$$\text{Also, } Z = X + Y$$

$$\therefore \text{Var}(Z) = \text{Var}(X+Y)$$

$$\Rightarrow 0 = \text{Var}(X) + \text{Var}(Y) + 2 \cdot \text{Cov}(X, Y)$$

$$\Rightarrow \text{Cov}(X, Y) = -\frac{1}{2}\{\text{Var}(X) + \text{Var}(Y)\}$$


Correlation coefficient is given by

$$r = \frac{\text{Cov}(X, Y)}{\sigma_X \cdot \sigma_Y} = \frac{-\frac{1}{2}\{\text{Var}(X) + \text{Var}(Y)\}}{\sigma_X \cdot \sigma_Y}$$

For exponential distribution;  $\mu_x = \sigma_x$  and  $\mu_y = \sigma_y$

$$\Rightarrow r = \frac{-\frac{1}{2}\{\mu_x^2 + \mu_y^2\}}{\mu_x \cdot \mu_y} = -\frac{1}{2} \frac{(2 \times 0.5^2)}{(0.5)^2} = -1$$

$$\Rightarrow r = -1$$

**78. (Marks to all (MTA))**

$$\text{If } x \sim N(\mu_x, \sigma_x^2)$$

$$y \sim N(\mu_y, \sigma_y^2)$$


Then

$$z = x + y \sim N(\mu_x + \mu_y, \sigma_x^2 + \sigma_y^2 + 2ab \text{ cov}(x, y))$$

If x and y are independent, then  $\text{cov}(x, y) = 0$

$$\Rightarrow z \sim N(\mu_x + \mu_y, \sigma_x^2 + \sigma_y^2)$$

∴ Z is normally distributed if x and y are independent.




**79. (0.299 to 0.301)**
**Given:**

'x' is random variable with uniform probability density function in  $[-2, 10]$

Let  $P(x) = k$

$$\Rightarrow \int_{-2}^{10} kdx = 1 \Rightarrow k = \frac{1}{12}$$




$$\text{Now } y \leq 7 \quad \Rightarrow 2x - 6 \leq 7$$

$$\Rightarrow x \leq \frac{13}{2} \quad \Rightarrow x \leq 6.5$$

$$\therefore P(y \leq 7 | x \geq 5) = \frac{P(y \leq 7 \cap x \geq 5)}{P(x \geq 5)}$$

$$= \frac{P(x \leq 6.5 \cap x \geq 5)}{P(x \geq 5)} = \frac{P(5 \leq x \leq 6.5)}{P(x \geq 5)}$$



$$\Rightarrow \frac{P(5 \leq x \leq 6.5)}{P(x \geq 5)} = \frac{1.5k}{5(k)} = \frac{1.5}{5} = 0.3$$

$$\therefore P(y \leq 7 | x \geq 5) = 0.3$$



Scan for Video solution


**80. (11 to 11)**
**Given:** X and Y are random variable and

$$E(2X + Y) = 0 \Rightarrow 2E(X) + E(Y) = 0 \quad \dots (1)$$

$$E(X + 2Y) = 33 \Rightarrow E(X) + 2E(Y) = 33 \quad \dots (2)$$


Adding equation (1) and (2)

$$\Rightarrow 3E(X) + 3E(Y) = 33 \Rightarrow 3(E(X) + E(Y)) = 33$$

$$\Rightarrow E(X) + E(Y) = 11$$



Scan for Video solution


**81. (0.36 to 0.38)**
**Given:** z is an exponential random variable with mean = 1


$$\Rightarrow P(z) = 1 \cdot e^{-1 \cdot z} = e^{-z}$$

$$\Pr(z > 2 | z > 1) = \frac{\Pr(z > 2 \cap z > 1)}{\Pr(z > 1)} = \frac{P(z > 2)}{P(z > 1)}$$

$$= \frac{\int_{\infty}^{\infty} e^{-z} dz}{\int_{\infty}^{\infty} e^{-z} dz} = \frac{\left. \frac{e^{-z}}{-1} \right|_2^{\infty}}{\left. \frac{e^{-z}}{-1} \right|_1^{\infty}} = \frac{e^{-\infty} - e^{-2}}{e^{-\infty} - e^{-1}}$$

$$= \frac{e^{-2}}{e^{-1}} = \frac{1}{e} = 0.3678 \approx 0.37$$

$$\therefore \boxed{\Pr(z > 2 | z > 1) = 0.37}$$



Scan for Video solution


**82. (a)**
**Given**


Length of rods follow normal distribution.

Let 'x' be the length of a randomly selected rod.

$$\Rightarrow x \sim N(440, 1^2)$$

$$x = 438 = \mu - 2\sigma$$

$$x = 441 = \mu + \sigma$$




$$\begin{aligned}
 P(438 < x < 441) &= P(\mu - 2\sigma < x < \mu + \sigma) \\
 &= P(\mu - 2\sigma < x < 0) + P(0 < x < \mu + \sigma) \\
 &= \frac{1}{2} \cdot P(\mu - 2\sigma < x < \mu + 2\sigma) + \frac{1}{2} \cdot (P(\mu - \sigma < x < \mu + \sigma)) \\
 &= \frac{1}{2} \times 0.9577 + \frac{1}{2} (0.6828) = 0.82025 = 82.025\% \\
 \therefore P(438 < x < 441) &= 82.025\%
 \end{aligned}$$



Scan for Video solution


**83. (b)**
**Given**

$0 < x < 10$  and  $0 < y < 20$  and  $x$  and  $y$  are uniformly distributed.



$$P(x + y > 20) = \frac{\text{area}(\Delta OBC)}{\text{area}(OABC)} = \frac{\frac{1}{2} \times BC \times BD}{BC \times AB}$$

$$\Rightarrow P(x + y > 20) = \frac{\frac{1}{2} \times 10 \times 10}{10 \times 20} = \frac{1}{4} = 0.25$$

$$\Rightarrow P(x + y > 20) = 0.25$$



Scan for Video solution


**84. (3 to 3)**
**Given**

$x$  is the mean of 3,  $x$ , 2, 4

$$\Rightarrow x = \frac{3+x+2+4}{4} \Rightarrow 3x = 9 \Rightarrow x = 3$$


$\therefore$  Given data is 3, 3, 2 and 4.

$\therefore$  Most repeating data point is  $x = 3$

Scan for Video solution

**85. (b)**

Uniform distribution is given by,



$$f(x) = \begin{cases} \frac{1}{y-x} & x \in (x, y) \\ 0 & \text{otherwise} \end{cases}$$

Scan for Video solution

**86. (0.8 to 0.8)**

**Given:**  $Y$  is uniformly distributed in (1, 6) and the polynomial is  $3x^2 + 6xY + (3Y + 6)$

For the polynomial,

$$Ax^2 + Bx + C \text{ to have real roots } B^2 - 4AC \geq 0$$

$$\Rightarrow (6Y)^2 - 4(3)(3Y + 6) \geq 0$$

$$\Rightarrow 36Y^2 - 12(3Y + 6) \geq 0$$


$$\Rightarrow 3Y^2 - 3Y - 6 \geq 0 \Rightarrow Y^2 - Y - 2 \geq 0$$

$$\Rightarrow Y^2 - 2Y + Y - 2 \geq 0 \Rightarrow (Y + 1)(Y - 2) \geq 0$$

$$\Rightarrow Y \leq -1 \text{ and } Y \geq 2$$

Since ' $y$ ' is distribution only in (1, 6)

For the polynomial to have real roots,  $y \geq 2$



$$P(Y \geq 2) = \frac{4.h}{5.h} = 0.8$$

Probability of polynomial having real roots = 0.8.

Scan for Video solution

**87. (a)**

For function to be a valid probability density function, the following condition must be satisfied,

$$\int_{-\infty}^{+\infty} f(x) dx = 1$$


$$\Rightarrow \int_0^1 f(x) dx + \int_1^2 f(x) dx + \int_2^3 f(x) dx = 1$$

$$\Rightarrow \left(\frac{1}{2} \times 1 \times h\right) + \left(\frac{1}{2} \times 1 \times 2h\right) + \left(\frac{1}{2} \times 1 \times 3h\right) = 1$$

$$\Rightarrow \frac{h}{2} + h + \frac{3h}{2} = 1 \Rightarrow h = \frac{1}{3}$$




Scan for Video solution

**88. (c)**


Right skewed curve

[Mode &lt; Median &lt; Mean]



Left skewed curve

[Mean &lt; Median &lt; Mode]

No skewness  
[Symmetrical]

[Mean = Median = Mode]



Scan for Video solution

**89. (0.25)**

Given  $X_1, X_2, X_3$  and  $X_4$  be independent normal random variables with same mean ( $= 0$ ) and variance ( $\sigma^2 = 1$ )

The probability that  $X_4$  is smallest =  $\frac{1}{4} = 0.25$ .

(Probability that a particular number from 4 randomly picked numbers to be smallest).



Scan for Video solution

**90. (b)**

Any normal random variable with mean ' $\mu$ ' and variance ' $\sigma^2$ ' is denoted as  $x \sim N(\mu, \sigma^2)$

Given,  $X_1 \sim N(1, 1)$  and  $X_2 \sim N(1, 4)$

For  $Y = aX_1 - bX_2$ ,

$Y \sim N(a\mu_1 - b\mu_2, a^2\sigma_1^2 + b^2\sigma_2^2 - 2ab \text{ Cov}(X_1, X_2))$

Since  $X_1, X_2$  are independent variables

$\Rightarrow \text{Cov}(X_1, X_2) = 0$

For  $a = 1$  and  $b = 1$

$\Rightarrow Y = X_1 - X_2$

$\Rightarrow Y \sim N(\mu_1 - \mu_2, \sigma_1^2 + \sigma_2^2)$

$\Rightarrow Y \sim N(0, 5)$



Scan for Video solution

**91. (a)****Given,**

$X_1 \rightarrow$  Exponentially distributed with Mean = 0.5

$$\Rightarrow X_1 = \left(\frac{1}{0.5}\right) \cdot e^{-x/(0.5)} = 2 \cdot e^{-2x} ; x > 0$$

$X_2 \rightarrow$  Exponentially distributed with Mean = 0.25

$$\Rightarrow X_2 = \left(\frac{1}{0.25}\right) \cdot e^{-x/0.25} = 4 \cdot e^{-4x} ; x > 0$$

$$\text{Since } Y = \min(X_1, X_2) = \begin{cases} 2 \cdot e^{-2x} ; 0 < x < \alpha \\ 4 \cdot e^{-4x} ; \alpha < x < \infty \end{cases}$$

Where ' $\alpha$ ' is value of 'x' where  $2e^{-2x}$  and  $4e^{-4x}$  intersect

$\therefore$  'Y' is exponentially distributed

$\Rightarrow$  Mean of Y < Min {Mean of  $X_1$ , Mean of  $X_2$ }

$\Rightarrow$  Mean of Y < Min {0.5, 0.25}

$\therefore$  Mean of Y < 0.25 = 1/6



Scan for Video solution

**92. (c)**


Mode is the most frequently occurring data point in a given data.

In the given data, 17 is the most repeating data point.

$$\therefore \text{Mode} = 17.$$



Scan for Video solution

**93. (0.9)**

Let the time of arrival be a random variable  $x$ , then

$$\text{its p.d.f. is } f(x) = \begin{cases} \frac{1}{5} & ; \quad 0 < x < 5 \\ 0 & ; \quad \text{otherwise} \end{cases}$$

Assume waiting time be  $h(x)$ , a function of arrival time

$$h(x) = \begin{cases} 0 & ; \quad 0 \leq x \leq 2 \text{ (GreenLight)} \\ 5 - x & ; \quad 2 < x \leq 5 \text{ (Red Light)} \end{cases}$$

Average waiting time =  $E[h(x)]$

$$\begin{aligned} &= \int_0^5 h(x) \cdot f(x) dx \\ &= \int_0^2 0 \cdot \frac{1}{5} dx + \int_2^5 (5-x) \cdot \frac{1}{5} dx \\ &= \int_2^5 \left(1 - \frac{x}{5}\right) dx \\ &= \left[x - \frac{x^2}{10}\right]_2^5 \\ &= (5 - 2.5) - (2 - 0.4) \end{aligned}$$


$$\text{Average waiting time} = 2.5 - 1.6 = 0.9$$



Scan for Video solution

**94. (c)**

The maximum number of tosses = 3



Hence possible outcomes = {H, TH, TTH, TTT}  
Here random variable  $Y$  denotes number of heads in the outcomes, hence possible values of  $Y$  will be 0 and 1.

$$p[y = 0] = P(\text{TTT}) = \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{8}$$

$$p[y = 1] = p(H) + p(TH) + p(TTH) \\ = \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{7}{8}$$

$\therefore$  Probability distribution

|      |     |     |
|------|-----|-----|
| y    | 0   | 1   |
| p(y) | 1/8 | 7/8 |

$$E(y) = \sum y(p(y)) = 0 \times \frac{1}{8} + 1 \times \frac{7}{8} = \frac{7}{8}$$

$$E(y^2) = \sum y^2 p(y) = 0^2 \times \frac{1}{8} + 1^2 \times \frac{7}{8} = \frac{7}{8}$$

$$\therefore \text{Var}(y) = E(y^2) - [E(y)]^2$$

$$= \frac{7}{8} - \left(\frac{7}{8}\right)^2 = \frac{7}{64}$$

$$\text{Var}(Y) = \frac{7}{64}$$




Scan for Video solution

**95. (0.0 to 0.0)****Given:**

$x$ : Gaussian random variable;  $\mu_x = 0$ ; & variance =  $\sigma^2$

$$\therefore y = \max(x, 0)$$

$$\Rightarrow y = \begin{cases} 0; & -\infty < x < 0 \\ x; & 0 < x < \infty \end{cases}$$



$\Rightarrow y = 0$  is the line that is in middle of the complete data of  $y$ .

$\therefore$  Median of  $y = 0$



Scan for Video solution



### 96. (b)

**Given:** 'X' is a discrete random variable and

$$P(X = j) = p_j, p_j \geq 0, j \in \{0, \dots, N\},$$

$$\Rightarrow \sum_{j=0}^N p_j = 1 \text{ (Sum of probabilities = 1)}$$

$$\therefore g_x(z) = \sum_{j=0}^N p_j z^j = P_0 + P_1 z + P_2 z^2 + \dots + P_N z^N$$

$\therefore g_x(z)$  is polynomial function.

$$\therefore g_y(z) = (1 - \beta + \beta z)^N = \sum_{j=0}^N N c_j (1 - \beta)^{N-j} \cdot (\beta z)^j$$

$\therefore$  Mean of  $Y = \text{coefficient of } z \text{ in above expansion}$

$$= N c_1 (1 - \beta)^{N-1} \beta = N \beta (1 - \beta)^{N-1}$$

For  $\beta \in [0, 1]$ ,  $(1 - \beta)^{N-1} = 1 - (N - 1) \beta$

$$\therefore \text{Expectation of } Y = N \beta (1 - (N - 1) \beta) \\ = N \beta - N(N - 1) \beta^2$$

Since ' $\beta$ ' is small, neglecting higher terms

$$\Rightarrow \text{Expectation of } Y = N \beta$$



Scan for Video solution



### 97. (b)

In exponential distribution,

$$f(x) = \lambda e^{-\lambda x}, \quad x \geq 0$$

Parameter is  $\lambda$ . (One parameter)

In Gaussian distribution,

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

Parameters are  $\mu$  and  $\sigma$ . (Two parameters)



Scan for Video solution



### 98. (b)

For function  $f(x)$  to be a valid probability density function, following condition must be satisfied;

$$\int_{-\infty}^{+\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^1 (a + bx) dx = \left[ ax + \frac{bx^2}{2} \right]_0^1$$

$$\Rightarrow a + \frac{b}{2} = 1$$

Hence option (b)  $a = 0.5, b = 1$  will satisfy the relation.



Scan for Video solution



### 99. (54 to 54)

**Given:** 'x' is a Poisson distributed variable

$\therefore$  Mean  $E(x) = \text{Variance} = 5$

$$E(x+2)^2 = E(x^2 + 4x + 4)$$

$$= E(x^2) + 4 E(x) + E(4)$$

$$= \text{Variance} + (E(x))^2 + 4 E(x) + 4$$

$$(\because \text{variance} = E(x^2) - E(x)^2)$$

$$\therefore E[(x+2)^2] = \text{Variance} + (E(x))^2 + 4 E(x) + 4 \\ = 5 + (5)^2 + 4(5) + 4 = 54$$

$$\therefore E[(x+2)^2] = 54$$



Scan for Video solution



**100. (0.55 to 0.55)****Given:**

Number of bulbs of type 1 = Number of bulbs of type 2

$$\Rightarrow P(\text{picking type 1}) = P(\text{picking type 2})$$

$$= 0.5 P(\text{type 1} > 100 \text{ hours})$$

$$= 0.7; P(\text{type 2} > 100 \text{ hours})$$

$$= 0.4 P(\text{bulb lasting} > 100 \text{ hours})$$

$$= P(\text{Type 1} \cap > 100 \text{ hrs} + P(\text{type 2}) \cap > 100 \text{ hrs})$$

$$= P(\text{Type 1}) \times P(\text{type 1} > 100 \text{ hrs}) + P(\text{type 2}) \times P(\text{type 2} > 100 \text{ hrs})$$

$$= (0.5 \times 0.7) + (0.5 \times 0.4) = 0.35 + 0.2 = 0.55$$

$$\therefore P(\text{bulb lasting} > 100 \text{ hours}) = 0.55$$



Scan for Video solution

**101. (a)**

The decisions to accept or reject a null hypothesis after inspecting only from it may be expressed exhaustively in the following table:-


|            | Reject Null             | Accept Null              |
|------------|-------------------------|--------------------------|
| Null True  | Wrong<br>(Type I error) | Correct                  |
| Null False | Correct                 | Wrong<br>(Type II Error) |



Scan for Video solution

**102. (a)****Given:**

$$f(x) = \begin{cases} 0.25 & \text{if } 1 \leq x \leq 5 \\ 0 & \text{otherwise} \end{cases}$$



$$P(x \leq 4) = \int_{-\infty}^4 f(x) dx$$

$$= \int_{-\infty}^1 f(x) dx + \int_1^4 f(x) dx$$

$$= \int_{-\infty}^1 0 dx + \int_1^4 0.25 dx = 0 + 0.25 [x]_1^4 = \frac{1}{4}(4 - 1)$$

$$P(x \leq 4) = \frac{3}{4}$$



Scan for Video solution

**103. (54.49 to 54.51)**

**Median speed** is the speed at the middle of series of spot speeds that are arranged in ascending order. 50% of speed values will be greater than the median and 50% of speed values will be less than median.

Ascending order of spot speed studies are

32, 45, 49, 51, 53, 56, 60, 62, 66, 79

$$\text{Median speed} = \frac{53 + 56}{2} = 54.5 \text{ km/hr}$$



Scan for Video solution

**104. (a)**

$$\text{Given: } P(X) = 0.4, \quad P(X \cup Y^c) = 0.7$$

$$\begin{aligned} P(X \cup Y^c) &= P(X) + P(Y^c) - P(X \cap Y^c) \\ &= P(X) + [1 - P(Y)] - P(X) \cdot P(Y^c) \\ &= P(X) + [1 - P(Y) - P(X) \{1 - P(Y)\}] \end{aligned}$$

$$\Rightarrow 0.7 = 0.4 + 1 - P(Y) - 0.4 \{1 - P(Y)\}$$

$$\Rightarrow P(Y) - 0.4 P(Y) = 0.4 + 1 - 0.7 - 0.4$$

$$\Rightarrow 0.6 P(Y) = 0.3 \Rightarrow P(Y) = 0.3/0.6 = 0.5$$

$$\begin{aligned} \therefore P(X \cup Y) &= P(X) + P(Y) - P(X \cap Y) \\ &= P(X) + P(Y) - P(X) \cdot P(Y) \end{aligned}$$

[Since X &amp; Y are independent events]

$$= 0.4 + 0.5 - 0.4 \times 0.5$$

$$\Rightarrow P(X \cup Y) = 0.7$$



Scan for Video solution



**105. (b)****Given:**

$$f(x) = \begin{cases} \frac{x}{a} + 1 & : -a \leq x \leq 0 \\ -\frac{x}{a} + 1 & : 0 \leq x \leq a \\ 0 & : \text{otherwise} \end{cases}$$

$$g(x) = \begin{cases} -\frac{x}{a} & : -a \leq x \leq 0 \\ \frac{x}{a} & : 0 \leq x \leq a \\ 0 & : \text{otherwise} \end{cases}$$

Mean of  $f(x)$ 

$$\begin{aligned} E[f(x)] &= \int_{-a}^0 x \left( \frac{x}{a} + 1 \right) dx + \int_0^a x \left( -\frac{x}{a} + 1 \right) dx \\ &= \left[ \frac{x^3}{3a} + \frac{x^2}{2} \right]_{-a}^0 + \left[ -\frac{x^3}{3a} + \frac{x^2}{2} \right]_0^a \\ &= -\left( -\frac{a^2}{3} + \frac{a^2}{2} \right) + \left( -\frac{a^2}{3} + \frac{a^2}{2} \right) = 0 \end{aligned}$$

Variance of  $f(x) = E[f(x)^2] - [E[f(x)]]^2$ 

$$\begin{aligned} E[f(x)^2] &= \int_{-a}^0 x^2 \left( \frac{x}{a} + 1 \right) dx + \int_0^a x^2 \left( -\frac{x}{a} + 1 \right) dx \\ &= \left[ \frac{x^4}{4a} + \frac{x^3}{3} \right]_{-a}^0 + \left[ -\frac{x^4}{4a} + \frac{x^3}{3} \right]_0^a = \frac{a^3}{6} \end{aligned}$$


Mean of  $g(x)$ 

$$\begin{aligned} E[g(x)] &= \int_{-a}^0 x \left( -\frac{x}{a} \right) dx + \int_0^a x \left( \frac{x}{a} \right) dx \\ &= \left[ -\frac{x^3}{3a} \right]_{-a}^0 + \left[ \frac{x^3}{3a} \right]_0^a = 0 \end{aligned}$$

Variance of  $g(x) = E[g(x)^2] - [E[g(x)]]^2$ 

$$\begin{aligned} E[g(x)^2] &= \int_{-a}^0 x^2 \left( -\frac{x}{a} \right) dx + \int_0^a x^2 \left( \frac{x}{a} \right) dx \\ &= \left[ -\frac{x^4}{4a} \right]_{-a}^0 + \left[ \frac{x^4}{4a} \right]_0^a = \frac{a^3}{2} \end{aligned}$$

- $\therefore$  Mean of  $f(x)$  and  $g(x)$  are same (i.e., equal to 0) but variance of  $f(x)$  and  $g(x)$  are different.



Scan for Video solution

**106. (a)**


For a Poisson distribution over a discrete random variable 'x', Mean = Variance

$$\Rightarrow \mu = \sigma^2 \Rightarrow \sigma = \sqrt{\mu}$$

$$\therefore \text{Standard deviation} = \sqrt{\mu}$$



Scan for Video solution

**107. (99.6 to 99.8)**If 'z' is standard normal variable, then the probability density function  $p(z)$  is as shown below**And**

$$P(-1 \leq z \leq 1) = 0.6828 = 68.28\%$$

$$P(-2 \leq z \leq 2) = 0.9577 = 95.77\%$$

$$P(-3 \leq z \leq 3) = 0.9973 = 99.73\%$$



Scan for Video solution

**108. (a)**

$$f(x) = \frac{3}{2} e^{-3x} u(x) + a e^{4x} u(-x)$$

$$\text{and, } u(x) = \begin{cases} 1 & ; \quad x \geq 0 \\ 0 & ; \quad x < 0 \end{cases}$$

$$u(-x) = \begin{cases} 0 & ; \quad x > 0 \\ 1 & ; \quad x \leq 0 \end{cases}$$

Since  $\int_{-\infty}^{+\infty} f(x)dx = 1$  for  $f(x)$  to a valid probability density function

$$\int_{-\infty}^{+\infty} \left[ \frac{3}{2} e^{-3x} u(x) + ae^{4x} u(-x) \right] dx$$

$$= \int_{-\infty}^0 ae^{4x} dx + \int_0^{+\infty} \frac{3}{2} e^{-3x} dx$$

$$= a \left[ \frac{e^{4x}}{4} \right]_{-\infty}^0 + \frac{3}{2} \left[ \frac{e^{-3x}}{-3} \right]_0^{+\infty} = 1$$

$$\frac{a}{4} + \frac{1}{2} = 1$$

$$a = 2$$

$$\text{Prob } \{x \leq 0\} = \int_{-\infty}^0 ae^{4x} dx = 2 \left[ \frac{e^{4x}}{4} \right]_{-\infty}^0$$

$$2 \left[ \frac{1}{4} - 0 \right]_{-\infty}^0 = \frac{1}{2}$$



Scan for Video solution




### 109. (0.33)

$$\text{Given } f(x, y) = \begin{cases} (x+y), & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0; & \text{otherwise} \end{cases}$$

Considering the region 'R' as the region, where  $x + y \leq 1$

$$\therefore P(x + y \leq 1) = \iint_R f(x, y) dy dx$$



$$\therefore P(x + y \leq 1) = \iint_R f(x, y) dy dx$$

$$= \int_{x=0}^{x=1} \int_{y=0}^{y=1-x} (x+y) dy dx$$

$$\begin{aligned} &= \int_{x=0}^{x=1} xy + \frac{y^2}{2} \Big|_0^{1-x} dx \\ &= \int_{x=0}^{x=1} (x(1-x) + \frac{(1-x)^2}{2}) dx \\ &= \frac{x^2}{2} - \frac{x^3}{3} + \frac{(1-x)^3}{-6} \Big|_0^1 = \frac{1}{2} - \frac{1}{3} + 0 - \left( \frac{-1}{6} \right) \\ &= \frac{1}{3} = 0.33 \end{aligned}$$

$$\therefore P(x + y \leq 1) = 0.33$$

Scan for Video solution



### 110. (0.5 to 0.5)

**Given:** Probability density function,

$$P(x) = \begin{cases} 1/x^2; & a \leq x \leq 1 \\ 0; & \text{otherwise} \end{cases}$$

For  $P(x)$  to be a valid p.d. function, following condition must be satisfied

$$\begin{aligned} \int_0^1 P(x) dx &= 1 \Rightarrow \int_a^1 \frac{1}{x^2} dx = 1 \\ \Rightarrow \frac{-1}{x} \Big|_a^1 &= 1 \Rightarrow \left\{ -1 + \frac{1}{a} \right\} = 1 \Rightarrow a = 0.5 \end{aligned}$$



Scan for Video solution



### 111. (d) For a given normal distribution,

$$\text{Mode} = \frac{1}{\sigma \cdot \sqrt{2\pi}} = \text{Max of } P(x)$$

For lowest variance,  $\sigma$  should be minimum.

If ' $\sigma$ ' is minimum, then  $\frac{1}{\sigma \cdot \sqrt{2\pi}}$  is maximum.

$\therefore$  The graph with highest peak is having least variance.

$\therefore$  Graph IV has least variance.



Scan for Video solution

**112. (0.23 to 0.25)**Given:  $q = 0.4$ 

|      |     |     |
|------|-----|-----|
| x    | 0   | 1   |
| P(x) | 0.4 | 0.6 |

$$E(x) = \sum_i x_i P(x_i) = 0 \times 0.4 + 1 \times 0.6 = 0.6$$

$$E(x^2) = \sum_i x_i^2 P(x_i) = 0^2 \times 0.4 + 1^2 \times 0.6 = 0.6$$

$$\begin{aligned} \therefore \text{Var}(x) &= E(x^2) - [E(x)]^2 \\ &= 0.6 - (0.6)^2 = 0.6 - 0.36 \end{aligned}$$

$$\text{Var}(x) = 0.24$$



Scan for Video solution

**113. (1.06 to 1.07)**

Given:

$$f(x) = \begin{cases} \frac{x}{4}(4-x^2) & 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

Since it is continuous random variable,

$$\therefore \mu_x = \int_0^2 x f(x) dx = \int_0^2 x \cdot \frac{x}{4}(4-x^2) dx$$

$$= \int_0^2 \left( x^2 - \frac{x^4}{4} \right) dx = \left[ \frac{x^3}{3} - \frac{x^5}{20} \right]_0^2$$

$$= \frac{8}{3} - \frac{32}{20} = \frac{16}{15}$$

$$\Rightarrow \mu_x = 1.066$$



Scan for Video solution


**114. (0.4 to 0.4)**

Given:

$$f(E) = \begin{cases} 1/5 & 0 < E < 5 \text{ mm/day} \\ 0 & \text{otherwise} \end{cases}$$

Required probability


$$P(2 < E < 4) = \int_2^4 f(E) dE = \int_2^4 \frac{1}{5} dE$$




$$= \frac{1}{5} [E]_2^4 = \frac{1}{5} (4 - 2) = \frac{2}{5} = 0.4$$



Scan for Video solution

**115. (b)**Given: The function  $f(x)$  is normal distributed curve with mean as 'a' and standard deviation as 'b'

$$\therefore \int_{-\infty}^a \frac{1}{\sqrt{2\pi} b} e^{-\frac{1}{2} \left( \frac{x-a}{b} \right)^2} dx = 0.5$$

**116. (0.26 to 0.27)**Given: Mean ( $\lambda$ ) = 5

For Poisson distribution

$$P(x=r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

Required probability

$$P(x < 4) = P(x=0) + P(x=1) + P(x=2) + P(x=3)$$

$$\begin{aligned}
 &= \frac{e^{-5} 5^0}{0!} + \frac{e^{-5} 5^1}{1!} + \frac{e^{-5} 5^2}{2!} + \frac{e^{-5} 5^3}{3!} \\
 &= e^{-5} \left[ 1 + 5 + \frac{25}{2} + \frac{125}{6} \right] = e^{-5} \left( \frac{118}{3} \right) \\
 &= 0.265
 \end{aligned}$$

$$\therefore P(x < 4) = 0.265$$



Scan for Video solution

**117. (25)**

**Given:** Probability of earth quake in an year is  
 $P(E) = 0.04$

$$\text{Average time interval } T = \frac{1}{P} = \frac{1}{0.04}$$

$$\Rightarrow T = 25 \text{ years}$$



Scan for Video solution

**118.(0.4 to 0.5)**

X → Random variable denoting lifetime of a bulb.  
 $f(x) = kx^2 ; 1 < x < 2$

For function to be a valid probability density function,

$$\therefore \int_1^2 f(x) dx = 1$$

$$\Rightarrow \int_1^2 kx^2 dx = 1$$

$$\Rightarrow k \left[ \frac{x^3}{3} \right]_1^2 = 1$$

$$\Rightarrow k \left( \frac{8 - 1}{3} \right) = 1$$

$$\Rightarrow k = \frac{3}{7}$$

$$\Rightarrow k = 0.4285$$



Scan for Video solution


**119. (49 to 51)**

Given that average daily balance follows normal distribution

Let 'x' be the normal random variable.

$$\mu = \text{Rs. } 500; \sigma = \text{Rs. } 50$$

$$P(x > 500) = P(x > \mu) = 0.5 = 50\%$$



$$\therefore P \left\{ \begin{array}{l} \text{Account having an} \\ \text{average daily balance} > 500 \end{array} \right\} = 50\%$$



Scan for Video solution



Scan for Video solution

**120. (b)**

Mean of Poisson distribution = 5.2

$$\text{Let mean} = \lambda \Rightarrow \lambda = 5.2$$

$$P(x < 2) = P(x = 0) + P(x = 1) \quad (x \rightarrow \text{number of accidents})$$

From Poisson distribution

$$P(x = r) = \frac{e^{-\lambda} \cdot \lambda^r}{r!}$$

$$P(x < 2) = P(x = 0) + P(x = 1) = \frac{e^{-\lambda} \cdot \lambda^0}{0!} + \frac{e^{-\lambda} \cdot \lambda^1}{1!}$$

$$= \frac{(1 + \lambda)}{e^\lambda} = \frac{1 + 5.2}{e^{5.2}} = 0.0342$$

$$\therefore P(x < 2) = 0.034$$



Scan for Video solution



**121. (d)****Given**

|      |     |     |     |
|------|-----|-----|-----|
| x    | 1   | 2   | 3   |
| p(x) | 0.3 | 0.6 | 0.1 |

$$\text{Standard deviation} = \sigma^2 = E(x^2) - (E(x))^2$$

$$\Rightarrow \sigma^2 = \left( \sum_{i=1}^n x_i^2 p(x=x_i) \right) - \left( \sum_{i=1}^n x_i \cdot p(x=x_i) \right)^2$$

$$\Rightarrow \sigma^2 = (1^2(0.3) + 2^2(0.6) + 3^2(0.1)) - (1(0.3) + 2(0.6) + 3(0.1))^2$$

$$\Rightarrow = 3.6 - (1.8)^2 = 0.36$$

$$\Rightarrow \boxed{\sigma = \text{Standard deviation} = 0.6}$$



Scan for Video solution

**122. (a)**

Let 'x' be the Number of defective pieces in a day

|      |     |     |     |
|------|-----|-----|-----|
| x    | 0   | 1   | 2   |
| P(x) | 1/6 | 2/3 | 1/6 |

$$\begin{aligned} \text{Mean} &= \sum_{i=1}^n x_i \cdot P(x=x_i) \\ &= 0\left(\frac{1}{6}\right) + 1\left(\frac{2}{3}\right) + 2\left(\frac{1}{6}\right) = \frac{2}{3} + \frac{2}{6} = 1 \end{aligned}$$

$$\therefore \text{Mean} = \mu = 1$$


$$\text{Variance} = \sum_{i=1}^n x_i^2 \cdot P(x=x_i) - \mu^2$$

$$\Rightarrow \sigma^2 = \left( 0^2 \left(\frac{1}{6}\right) + 1^2 \cdot \left(\frac{2}{3}\right) + 2^2 \cdot \left(\frac{1}{6}\right) \right) - 1^2$$

$$\Rightarrow = \left(\frac{2}{3} + \frac{4}{6}\right) - 1 = \frac{1}{3}$$

$$\Rightarrow \sigma^2 = \text{Variance} = \frac{1}{3}$$

$$\therefore \boxed{\text{Mean} = \mu = 1 \text{ and Variance} = \sigma^2 = \frac{1}{3}}$$



Scan for Video solution

**123. (0.32 to 0.34)**Given  $X_1, X_2$  and  $X_3$  be independent and identically distributed random valued variables

$$P\{X_1 \text{ is largest}\} = \frac{1}{3} (\text{X}_1 \text{ being largest of } X_1, X_2, X_3) = 0.33$$

$$\therefore P\{X_1 \text{ is largest}\} = 0.33$$



Scan for Video solution

**124. (0.79 to 0.81)**Given  $x$  be a zero mean unit variance gaussian random variable

$$\Rightarrow x \sim N(0, 1)$$

$\therefore$  Probability density function of  $x$  is

$$P(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

$\therefore$  Expectation of  $|x|$  is

$$E(|x|) = \int_{-\infty}^{\infty} |x| \cdot P(X) dx = \int_{-\infty}^{\infty} |x| \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$$

$$= \frac{1}{\sqrt{2\pi}} \times \left\{ 2 \times \int_0^{\infty} x \cdot e^{-\frac{x^2}{2}} dx \right\}$$

$$\text{Let } \frac{x^2}{2} = t \Rightarrow 2x dx = dt \Rightarrow x dx = dt$$

$$\text{Lower limit: } 0^2 = t \Rightarrow t = 0$$

$$\text{Upper limit: } t \rightarrow \infty$$

$$\therefore E|x| = \frac{2}{\sqrt{2\pi}} \times \int_0^{\infty} e^{-t} dt = \sqrt{\frac{2}{\pi}} \left( \left[ \frac{e^{-t}}{-1} \right]_0^{\infty} \right)$$

$$= \sqrt{\frac{2}{\pi}} \cdot \{0 + 1\} = 0.79788 \quad \therefore E|x| \approx 0.8$$



Scan for Video solution

**125. (b)**For a real valued random variable  $x$ ,

$$\text{Variance} = \sigma^2 = E(x^2) - (E(x))^2 \geq 0$$

$$E(x^2) \geq (E(x))^2$$


**Note:** If  $\sigma = 0 \Rightarrow$  All the data points in a sample should be identical.



Scan for Video solution

**126. (0.35 to 0.45)**

$$f(x) = \begin{cases} 0.2 & ; -1 \leq x \leq 1 \\ 0.1 & ; (-4 \leq x \leq -1) \cup (1 < x \leq 4) \\ 0 & ; (x < -4) \cup (x > 4) \end{cases}$$



$$P(0.5 < x < 5) = \int_{0.5}^5 f(x) dx$$

$$= \int_{0.5}^1 (0.2)dx + \int_1^4 (0.1)dx + \int_4^5 (0)dx$$

$$= 0.2(1 - 0.5) + 0.1(4 - 1) + 0$$

$$= 0.2 \times 0.5 + 0.1 \times 3 = 0.1 + 0.3 = 0.4$$



Scan for Video solution

**127. (0.24 to 0.27)****Given:**


Length of stick = 1 unit

For the broken stick to be shorter,


 $0 < \text{length of broken part} < 0.5$ .

Let 'x' be the length of shorter part,

$$\Rightarrow 0 < x < 0.5$$




$$E(x) = \text{average of } (0, 0.5) = \frac{0 + 0.5}{2} = 0.25$$

 $\therefore$  Expected length of shorter part is 0.25.**128.(a)**

$$f(x) = \begin{cases} e^{-x} & ; x > 0 \\ 0 & ; x \leq 0 \end{cases}$$

$$\therefore P(x > 1) = \int_{x=1}^{x=\infty} f(x) \cdot dx$$



$$= \int_1^\infty e^{-x} dx - \left[ e^{-x} \right]_1^\infty = -\left[ e^{-\infty} - e^{-1} \right]$$

$$= -\left[ 0 - \frac{1}{e} \right]$$

$$= \frac{1}{e} = 0.368$$




Scan for Video solution

**129.(b)****Given:**

'X' is a normal random variable with mean 1 and variance 4  $\Rightarrow X \sim N(1, 4)$

$\Rightarrow X = 0$  lies to the left of the mean  $X=1$ ,

$\therefore P(X < 0)$  for the given variable is as shown below

 $P(x)$ 

$$\Rightarrow P(X < 0) = 0.5 - P(0 < X < 1)$$

$$\Rightarrow 0 < P(X < 0) < 0.5$$



Scan for Video solution

**130. (d)**

Given U and V are identically distributed.

Let  $2V = W$

$$\Rightarrow W = 2U.$$

Cumulative function of 'U' is  $p(U \leq x)$

Cumulative function of 'W' is  $p(W \leq x)$

$$\Rightarrow (2U \leq x)$$

$$= p\left(U \leq \frac{x}{2}\right)$$

$$\therefore F(x) = p(U \leq x); G(x) = p\left(U \leq \frac{x}{2}\right)$$

So, when  $x > 0$ ;  $F(x) \geq G(x) \Rightarrow (F(x) - G(x)) \geq 0$

When  $x < 0$ ;  $F(x) < G(x) \Rightarrow (F(x) - G(x)) < 0$

$\therefore$  In both cases for all  $x$ ,

$$(F(x) - G(x))x \geq 0$$

$$\Rightarrow f(x) = \begin{cases} \lambda(-x^2 + 3x - 2) & \text{for } 1 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

$$\Rightarrow \int_1^2 f(x) dx = 1$$

$$\Rightarrow \int_1^2 \lambda(-x^2 + 3x - 2) dx = 1$$

$$\Rightarrow \lambda \left[ -\frac{x^3}{3} + \frac{3x^2}{2} - 2x \right]_1^2 = 1$$

$$\Rightarrow \lambda \left[ -\frac{(2^3 - 1^3)}{3} + 3 \frac{(2^2 - 1^2)}{2} - 2(2 - 1) \right] = 1$$

$$\Rightarrow \lambda \left[ -\frac{7}{3} + \frac{9}{2} - 2 \right] = 1 \Rightarrow [\lambda = 6]$$



Scan for Video solution



133. (c)

**Given:**Average number of cars per minute passing is  $\lambda$ 

'P' is following Poisson distribution with mean =  $\lambda$   
 $= 3$


$$P(x < 3) = P(x = 0) + P(x = 1) + P(x = 2)$$

$$= \frac{e^{-\lambda} \cdot \lambda^0}{0!} + \frac{e^{-\lambda} \cdot \lambda^1}{1!} + \frac{e^{-\lambda} \cdot \lambda^2}{2!}$$

$$\therefore P(x = r) = \frac{e^{-\lambda} \cdot \lambda^r}{r!}$$

$$= e^{-3} \left( \frac{1}{1} + \frac{3}{1} + \frac{9}{2} \right) = \frac{17}{2 \cdot e^3}$$

$$\therefore P(\text{less than 3 cars}) = \frac{17}{2 \cdot e^3}$$




134. (a)

**Given:** The annual precipitation is normally distributed with  $\mu = 1000 \text{ mm}$ ,  $\sigma = 200 \text{ mm}$

$$\begin{aligned} P(x > 1200 \text{ mm}) &= P\left(z > \frac{x - \mu}{\sigma}\right) \\ &= P\left(z > \frac{1200 - 1000}{200}\right) = P(z > 1) \end{aligned}$$

Where z is the standard normal variate.




$$\therefore P(-1 < z < 1) = 0.68$$

$$\Rightarrow P(0 < z < 1) = \frac{0.68}{2} = 0.34$$

$$\text{So, } P(z > 1) = 0.5 - 0.34 = 0.16$$

$$\approx 16\% < 50\%$$


Hence option (a) is correct



135. (b)

$$1 \leq x, y \leq +1$$

$$\text{Probability of } \max(x, y) < \frac{1}{2}$$



$$\begin{aligned}
 &= P\left(-1 < x, y < \frac{1}{2}\right) \\
 &= \frac{\text{Area of shaded region}}{\text{Total area}} = \frac{\frac{3}{2} \times \frac{3}{2}}{2 \times 2} = \frac{9}{16}
 \end{aligned}$$




Scan for Video solution

**136. (b)**


**Given:**  $x, y$  are independent random variables that are uniformly distributed in  $[-1, 1]$

The probability distributions of both  $x$  and  $y$  is as shown



$$\begin{aligned}
 P(\text{Max}(x, y) < 1/2) &= P\left(x < \frac{1}{2} \cap y < \frac{1}{2}\right) \\
 &= P\left(x < \frac{1}{2}\right) \cdot P\left(y < \frac{1}{2}\right) \\
 &= \frac{3}{4} \cdot \frac{3}{4} = \frac{9}{16}
 \end{aligned}$$

$$\therefore P\left(\max(x, y) < \frac{1}{2}\right) = 9/16$$

**137. (c)**

**Given:** ‘ $X$ ’ is a Random variable that can take values only  $+1$  and  $-1$  with equal probabilities. So the probability distribution of  $x$  is as below.

|      |     |     |
|------|-----|-----|
| x    | -1  | +1  |
| P(x) | 0.5 | 0.5 |


Cumulative distribution function is defined as

$$F(x) = P(X \leq x)$$

$$\therefore F(-1) = P(x \leq -1) = P(x = -1) = 0.5$$

$$\begin{aligned} \therefore F(+1) &= P(x \leq 1) = P(x = -1) + P(x = 1) \\ &= 0.5 + 0.5 = 1 \end{aligned}$$

$$\therefore F(-1) = 0.5 \text{ and } F(+1) = 1$$

**138. (b)**

**Given:** If the value on the die is 1, 2, or 3, the die is rolled a second time i.e., if (4 or 5 or 6) appear than dice is not rolled out second time.

$$P(\text{sum} \geq 6) = P(\text{sum} \geq 6)_{\text{1st roll}} + P(\text{sum} \geq 6)_{\text{2nd roll}}$$

$$\therefore P(\text{sum} \geq 6)_{\text{1st roll}} = P(6) = \frac{1}{6}$$

$$P(\text{sum} \geq 6)_{\text{2nd}}$$

$$= P(1 \cap \geq 5) + P(2 \cap \geq 4) + P(3 \cap \geq 3)$$

$$= P(1) \cdot P(\geq 5) + P(2) \cdot P(\geq 4) + P(3) \cdot P(\geq 3)$$

$$= \frac{1}{6} \cdot \left(\frac{2}{6}\right) + \left(\frac{1}{6}\right) \cdot \left(\frac{3}{6}\right) + \left(\frac{1}{6}\right) \cdot \left(\frac{4}{6}\right)$$

$$= \frac{1}{36} (2 + 3 + 4) = \frac{1}{4}$$

$$\therefore P(\text{sum} \geq 6) = \frac{1}{6} + \frac{1}{4} = \frac{5}{12}$$



Scan for Video solution

**139. (d)**

- If  $x_1, x_2, x_3, \dots, x_n$  are 'n' data points and  $y_i = a x_i + b$ ,

The index position of median and mode will be same as initial positions since all the points are modified uniformly.

$$\begin{aligned} E(y) &= E(a x_i + b) = a E(x_i) + b \\ \Rightarrow E(y) &= a E(x) + b \quad \Rightarrow \mu_y = a \mu_x + b \\ \text{But } \text{Var}(ax + b) &= a^2 \text{Var}(x) \quad \Rightarrow \sigma_y = a \cdot \sigma_x \\ \therefore \text{The statements } \sigma_y &= a \cdot \sigma_x + b \text{ is incorrect.} \end{aligned}$$



Scan for Video solution

**140. (c)**

For a random variable 'x'

$$\text{Variance } \sigma^2 = E(x^2) - (E(x))^2 = R \geq 0$$

$$\therefore R \geq 0$$

**Note:**  $R = 0$  if the data is ideal (i.e. all the data points are same).



Scan for Video solution

**141. (a)**

**Given:**

A deck contains 5 cards.

For 1<sup>st</sup> number to be more than the 2<sup>nd</sup> number by one, the possibilities are (5, 4), (4, 3), (3, 2), (2, 1)

Required probability

$$\begin{aligned} &= P(5).P(4) + P(4).P(3) + P(3).P(2) + P(2).P(1) \\ &= \left(\frac{1}{5} \times \frac{1}{4}\right) \times 4 = \frac{4}{20} = \frac{1}{5} \end{aligned}$$



Scan for Video solution

**142. (a)**

**Given:**

Probability of a computer being faulty =  $P(E) = p$

Probability that test gives correct result =  $P(C) = q$

A computer is declared faulty in two cases

- Faulty and test gave correct result
- Not faulty but test gave incorrect result.

$$P(\text{Declaring faulty}) = P(F \cap C) + P(F^c \cap C^c)$$

$$\begin{aligned} &= P(E) \cdot P(C) + P(F^c) \cdot P(C^c) \\ &= p \cdot q + (1-p)(1-q) \end{aligned}$$

$$\therefore P(\text{Declaring faulty}) = pq + (1-p)(1-q)$$



Scan for Video solution

**143. (c)**

**Given:** x, y are independent variables

|        |     |     |     |
|--------|-----|-----|-----|
| x or y | 0   | 1   | 2   |
| P      | 1/2 | 1/4 | 1/4 |

$$P(x+y=2 | x-y=0) = \frac{P(x+y=2 \cap x-y=0)}{P(x-y=0)}$$

$$P(x+y=2 \cap x-y=0) = P(x=1 \cap y=1)$$

$$= P(x=1) \cdot P(y=1) = \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{16}$$

$$\begin{aligned} P(x - y = 0) &= P(x = y) = P(x = y = 0) + P(x = y = 1) + P(x = y = 2) \\ &= P(x = 0)P(y = 0) + P(x = 1)P(y = 1) \end{aligned}$$

$$+ P(x = 2)P(y = 2)$$

$$= \left(\frac{1}{2}\right)^2 + \left(\frac{1}{4}\right)^2 + \left(\frac{1}{4}\right)^2 = \frac{3}{8}$$

$$\therefore P(x + y = 2 | x - y = 0) = \frac{1/16}{3/8} = \frac{1}{6}$$



Scan for Video solution



144. (b)

**Given:** 'x' is a random variable

|        |     |     |     |     |     |
|--------|-----|-----|-----|-----|-----|
| k      | 1   | 2   | 3   | 4   | 5   |
| P(x=k) | 0.1 | 0.2 | 0.4 | 0.2 | 0.1 |

$$\begin{aligned} \text{Mean of } x = E(x) &= \sum_{k=1}^5 k \cdot P(x = k) \\ &= 1(0.1) + 2(0.2) + 3(0.4) + 4(0.2) + 5(0.1) \\ &= 0.1 + 0.4 + 1.2 + 0.8 + 0.5 = 3 \end{aligned}$$

$$\begin{aligned} \text{Variance of } x = E(x^2) - (E(x))^2 &= \{1^2(0.1) + 2^2(0.2) + 3^2(0.4) + 4^2(0.2) + 5^2(0.1)\} - 3^2 \\ &= (0.1 + 0.8 + 3.6 + 3.2 + 2.5) - 9 = 1.2 \end{aligned}$$

So, both teacher and student are wrong.




Scan for Video solution



145.(a)

**Given:** A random variable is uniformly distributed.  
Let probability distribution function  $P(x) = k$ .



Since total probability = 1

$$\Rightarrow \int_0^1 P(x) dx = \int_0^1 k dx = 1 \Rightarrow P(x) = k = 1$$

$$\begin{aligned} \text{Variance } \sigma^2 &= \left( \int_0^1 x^2 \cdot P(x) dx \right) - \left( \int_0^1 x \cdot P(x) dx \right)^2 \\ &= \int_0^1 x^2 dx - \left( \int_0^1 x dx \right)^2 = \frac{x^3}{3} \Big|_0^1 - \left( \frac{x^2}{2} \Big|_0^1 \right)^2 \\ &= \frac{1}{3} - \frac{1}{4} = \frac{1}{12} \end{aligned}$$

$$\therefore \text{Standard deviation } = \sigma = \frac{1}{\sqrt{12}}$$

Scan for Video solution



146. (d)

$$\text{Given: } F(X_N) = \frac{1}{1 + \exp(-1.7255X_N |X_N|^{0.12})}$$

$\mu$  (Mean) = 102 cm, Standard deviation ( $\sigma$ ) = 27 cm


$P(90 \text{ cm} \leq X \leq 102 \text{ cm}) = ?$

$P(90 \text{ cm} \leq X \leq 102 \text{ cm})$

$$= P\left(\frac{90 - \mu}{\sigma} \leq X_N \leq \frac{102 - \mu}{\sigma}\right)$$

$$= P\left(\frac{90 - 102}{27} \leq X_N \leq \frac{102 - 102}{27}\right)$$

$$= P(-0.44 \leq X_N \leq 0) = F(0) - F(-0.44)$$



$$= \frac{1}{1 + \exp(0)} - \frac{1}{1 + \exp(-1.7255 \times (-0.44) | -0.44 |^{0.12})}$$

$$= \frac{1}{2} - \frac{1}{1 + \exp(0.688)} = \frac{1}{2} - \frac{1}{2.9897} = 0.1655$$

= **16.55%** (Approximate value)



Scan for Video solution



#### 147. (b)

**Given:**

$$P(\text{odd}) = 0.9 P(\text{even})$$

$$\therefore P(\text{odd}) + P(\text{even}) = 1$$

$$0.9 P(\text{even}) + P(\text{even}) = 1$$

$$\Rightarrow P(\text{even}) = \frac{1}{1.9}$$

Since probability of getting any even number is same.

$$P(2) = \frac{1}{3} \times \frac{1}{1.9}; P(4) = \frac{1}{3} \times \frac{1}{1.9}; P(6) = \frac{1}{3} \times \frac{1}{1.9}$$

$$P(\text{Even} | > 3) = 0.75 \Rightarrow \frac{P(\text{even} \cap > 3)}{P(> 3)} = 0.75$$

$$\Rightarrow \frac{P(4) + P(6)}{P(> 3)} = 0.75 \Rightarrow P(> 3) = \frac{P(4) + P(6)}{0.75}$$

$$= \frac{2 \left( \frac{1}{3 \times 1.9} \right)}{0.75} = 0.46783 \approx 0.468$$

$\therefore$  Required probability = 0.468


Scan for Video solution



#### 148. (b)

**Given:**  $f(x) = \begin{cases} x^2 & -1 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$

$$\therefore P\left(-\frac{1}{3} \leq x \leq \frac{1}{3}\right) = \int_{-1/3}^{+1/3} f(x) dx = \int_{-1/3}^{+1/3} x^2 dx$$

$$= 2 \int_0^{1/3} x^2 dx \quad [\because f(x) = f(-x)]$$

$$= 2 \left[ \frac{x^3}{3} \right]_0^{1/3} = 2 \left( \frac{1}{81} \right) = \frac{2}{81}$$

$$\therefore P\left(-\frac{1}{3} \leq x \leq \frac{1}{3}\right) = \frac{2}{81} \times 100 = 2.47 \%$$


Scan for Video solution



#### 149. (a)

**Given:**

$$P = \sum_{\substack{1 \leq i \leq 2k \\ i \text{ odd}}} i = 1 + 3 + 5 + \dots + (2k-1)$$

$$Q = \sum_{\substack{1 \leq i \leq 2k \\ i \text{ even}}} i = 2 + 4 + 6 + \dots + 2k$$

$$\therefore P = \text{sum of } 1^{\text{st}} 'k' \text{ odd numbers} = k^2$$

$$Q = \text{sum of } 1^{\text{st}} 'k' \text{ even numbers}$$

$$= 2 + 4 + 6 + \dots + 2k = 2(1 + 2 + 3 + \dots + k)$$

$$= 2 \cdot \left( \frac{k(k+1)}{2} \right) = k^2 + k$$

$$\therefore P = k^2; \text{ and } Q = k^2 + k$$

$$\Rightarrow Q = P + k \Rightarrow P = Q - k$$



Scan for Video solution



**150. (a)**

**Given:**

X is normal variable with Mean = +1 and variance = 4

Considering 'z' the standard normal variable as

$$z = \frac{X - \mu_X}{\sigma_X} \Rightarrow z = \frac{X - 1}{2} \quad \text{i.e. At } X = -1; z = -1$$

Y is normal variable with Mean = -1 & variance =

$\sigma_Y$


$$\Rightarrow z = \frac{Y - \mu_Y}{\sigma_Y} = \frac{Y + 1}{\sigma_Y} \quad \text{i.e. At } Y = 2; z = 3/\sigma_Y$$

$$\therefore P(X \leq -1) = P(Y \geq 2) \Rightarrow P(z < -1) = P$$

$$\left( z \geq \frac{3}{\sigma_Y} \right)$$

$$\Rightarrow \left| \frac{3}{\sigma_Y} \right| = 1 \quad \Rightarrow \sigma_Y = 3$$

$\therefore$  standard deviation of Y = 3




Scan for Video solution



**151. (c)**

Since X is uniformly distributed between [0, 1]



$$f(X) = \begin{cases} \frac{1}{1-0} & ; \quad 0 < X < 1 \\ 0 & ; \quad \text{otherwise} \end{cases}$$

$$\therefore f(X) = 1; 0 < X < 1 \text{ Now } E(X^3) = \int_0^1 X^3 f(x) dx$$

$$= \left[ \frac{X^4}{4} \right]_0^1 \therefore E(X^3) = \frac{1}{4}$$

Scan for Video solution



**152. (a)**

**Given:** Probability density function,

$$P_x(x) = M e^{-2|x|} + N e^{-3|x|}$$

Since Total probability = 1

$$\Rightarrow \int_{-\infty}^{\infty} P_x(x) dx = 1$$

$$\Rightarrow \int_{-\infty}^{\infty} (M e^{-2|x|} + N e^{-3|x|}) dx = 1$$

$$\Rightarrow 2 \times \int_0^{\infty} (M e^{-2|x|} + N e^{-3|x|}) dx = 1 \quad (\text{Since } |x| \text{ is even function})$$

$$\Rightarrow 2 \times \int_0^{\infty} (M e^{-2x} + N e^{-3x}) dx = 1$$

$$\Rightarrow 2 \times \left( \left. \frac{M e^{-2x}}{-2} + \frac{N e^{-3x}}{-3} \right|_0^{\infty} \right) = 1$$

$$\Rightarrow 2 \left( -\frac{M}{-2} - \frac{N}{-3} \right) = 1 \Rightarrow \boxed{M + \frac{2N}{3} = 1}$$



Scan for Video solution

**153. (1.9 to 1.9)****Given:**

|   |    |    |    |    |
|---|----|----|----|----|
| x | 5  | 2  | 4  | 3  |
| y | 16 | 10 | 13 | 12 |

| x | y  | $x^2$ | XY |
|---|----|-------|----|
| 5 | 16 | 25    | 80 |
| 2 | 10 | 4     | 20 |
| 4 | 13 | 16    | 52 |
| 3 | 12 | 9     | 36 |

The normal equation for fitting of straight lines is given by,

$$\Sigma y = an + b\Sigma x \quad \text{and} \quad \Sigma xy = a\Sigma x + b\Sigma x^2$$

Have,  $n = 4$ ,  $\Sigma x = 14$ ,  $\Sigma y = 51$ ,  $\Sigma x^2 = 54$ ,  $\Sigma xy = 188$

$$\Rightarrow 51 = 4a + 14b \quad \dots\dots(i)$$

$$\Rightarrow 188 = 14a + 54b \quad \dots\dots(ii)$$

After solving (i) and (ii)

$$\text{We get } a = 6.1, \quad b = 1.9$$



Scan for Video solution

**154. (d)****Given:** X, Y are random variables.

$$\Rightarrow \frac{[E(X - E(X)) \cdot (Y - E(Y))]^2}{\sigma_X^2 \cdot \sigma_Y^2} \leq 1$$



$$\Rightarrow [E(X - E(X)) \cdot (Y - E(Y))]^2 \leq \sigma_X^2 \cdot \sigma_Y^2$$

$$\therefore [E(X - E(X)) \cdot (Y - E(Y))]^2 \leq \text{Var}(X) \cdot \text{Var}(Y)$$

$\therefore$  Statement S<sub>1</sub> is false.

$$\text{Cov}[X, Y] = E(X - E(X)) \cdot (Y - E(Y))$$

Only if  $X > E(X)$  and  $Y > E(Y)$

Or  $X < E(X)$  and  $Y < E(Y)$

So it is not valid for any other relation between X and E(X).

$\therefore$  Statement-2 (for all X and Y) is also False.

$\Rightarrow$  Both S<sub>1</sub> and S<sub>2</sub> are False.

## CHAPTER

# 4

# DIFFERENTIAL EQUATIONS

### Order and Degree of Differential Equations

**1. [MCQ] [GATE-CE-2022: 1M]**

For the equation  $\frac{d^3y}{dx^3} + x\left(\frac{dy}{dx}\right)^{3/2} + x^2y = 0$  the correct describing is

- (a) An ordinary differential equation of order 3 and degree 2.
- (b) An ordinary differential equation of order 1 and degree 3.
- (c) An ordinary differential equation of order 2 and degree 3.
- (d) An ordinary differential equation of order 1 and degree 3/2

**2. [MCQ] [GATE-CE-2020:1M]**

The ordinary differential equation

$$\frac{d^2u}{dx^2} - 2x^2u + \sin x = 0 \text{ is}$$

- (a) Linear and homogeneous
- (b) Nonlinear and homogeneous
- (c) Nonlinear and nonhomogeneous
- (d) Linear and nonhomogeneous

**3. [MCQ] [GATE-EC-2014:1M]**

Which ONE of the following is a linear non-homogeneous differential equation, where x and y are the independent and dependent variables respectively?

- |                                   |                                  |
|-----------------------------------|----------------------------------|
| (a) $\frac{dy}{dx} + xy = e^{-x}$ | (b) $\frac{dy}{dx} + xy = 0$     |
| (c) $\frac{dy}{dx} + xy = e^{-y}$ | (d) $\frac{dy}{dx} + e^{-y} = 0$ |

**4. [MCQ] [GATE-ME-2013: 1M]**

The partial differential equation  $\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = \frac{\partial^2 u}{\partial x^2}$  is a

- (a) Linear equation of order 2
- (b) Non-linear equation of order 1
- (c) Linear equation of order 1
- (d) Non-linear equation of order 2

**5. [MCQ] [GATE-CE-2010:1M]**

The order and degree of the differential equation

$$\frac{d^3y}{dx^3} + 4\sqrt{\left(\frac{dy}{dx}\right)^3 + y^2} = 0 \text{ are respectively}$$

- |             |             |
|-------------|-------------|
| (a) 3 and 2 | (b) 2 and 3 |
| (c) 3 and 3 | (d) 3 and 1 |

**6. [MCQ] [GATE-ME-2010: 1M]**

The Blasius equation,  $\frac{d^3f}{d\eta^3} + \frac{f}{2} \frac{d^2f}{d\eta^2} = 0$ , is a

- (a) Second order nonlinear ordinary differential equation.
- (b) Third order nonlinear ordinary differential equation.
- (c) Third order linear ordinary differential equation.
- (d) Mixed order nonlinear ordinary differential equation.

**7. [MCQ] [GATE-EC-2009:1M]**

The order of the differential equation

$$\frac{d^2y}{dt^2} + \left(\frac{dy}{dt}\right)^3 + y^4 = e^{-t} \text{ is}$$

- |       |       |
|-------|-------|
| (a) 1 | (b) 2 |
| (c) 3 | (d) 4 |

## First Order Differential Equations

8. [NAT] [GATE-CE-2023: 1M]

In the differential equation  $\frac{dy}{dx} + \alpha xy = 0$ ,  $\alpha$  is a positive constant. If  $y = 1.0$  at  $x = 0.0$ , and  $y = 0.8$  at  $x = 1.0$ , the value of  $\alpha$  is \_\_\_\_\_ (rounded off to three decimal places).

10. [MCQ] [GATE-ME-2022: 2M]

For the exact differential equation

$$\frac{du}{dx} = \frac{-xu^2}{2+x^2u},$$

which one of the following is the solution?

- (a)  $u^2 + 2x^2 = \text{constant}$
  - (b)  $xu^2 + u = \text{constant}$
  - (c)  $\frac{1}{2}x^2u^2 + 2u = \text{constant}$
  - (d)  $\frac{1}{2}ux^2 + 2x = \text{constant}$
11. [MCQ] [GATE-ME-2021: 1M]

If  $y(x)$  satisfies the differential equation

$$(\sin x)\frac{dy}{dx} + y \cos x = 1,$$

subject to be condition  $y\left(\frac{\pi}{2}\right) = \frac{\pi}{2}$ , then  $y\left(\frac{\pi}{6}\right)$  is

- (a) 0
  - (b)  $\frac{\pi}{6}$
  - (c)  $\frac{\pi}{3}$
  - (d)  $\frac{\pi}{2}$
12. [MCQ] [GATE-ME-2021: 2M]

Consider the following differential equation

$$(1+y)\frac{dy}{dx} = y$$

The solution of the equation that satisfies the condition  $y(1) = 1$  is

- (a)  $2ye^y = e^x + e$
  - (b)  $y^2e^y = e^x$
  - (c)  $ye^y = e^x$
  - (d)  $(1+y)e^y = 2e^x$
13. [MCQ] [GATE-CE-2021:2M]

If  $k$  is a constant, the general solution of  $\frac{dy}{dx} - \frac{y}{x} = 1$

will be in the form of

- (a)  $y = x \ln(x)$
- (b)  $y = x k \ln(k)$
- (c)  $y = k \ln(kx)$
- (d)  $y = x \ln(kx)$

14. [MCQ] [GATE-EC-2021:1M]

Consider the differential equation given below.

$$\frac{dy}{dx} + \frac{x}{1-x^2}y = x\sqrt{y}$$

The integrating factor of the differential equation is

- (a)  $(1-x^2)^{-\frac{3}{4}}$
- (b)  $(1-x^2)^{-\frac{1}{4}}$
- (c)  $(1-x^2)^{-\frac{3}{2}}$
- (d)  $(1-x^2)^{-\frac{1}{2}}$

15. [NAT] [GATE-EE-2020 : 1M]

Consider the initial value problem below.

$$\frac{dy}{dx} = 2x - y, \quad y(0) = 1$$

The value of  $y$  at  $x = \ln 2$ , (rounded off to 3 decimal places) is \_\_\_\_\_.

16. [MCQ] [GATE-ME-2019: 1M]

For the equation  $\frac{dy}{dx} + 7x^2y = 0$ , if  $y(0) = \frac{3}{7}$ , then the value of  $y(1)$  is

- (a)  $\frac{7}{3}e^{-7/3}$
- (b)  $\frac{7}{3}e^{-3/7}$
- (c)  $\frac{3}{7}e^{-7/3}$
- (d)  $\frac{3}{7}e^{-3/7}$

17. [MCQ] [GATE-ME-2019: 1M]

The differential equation  $\frac{dy}{dx} + 4y = 5$  is valid in the domain  $0 \leq x \leq 1$  with  $y(0) = 2.25$ . The solution of the differential equation is

- (a)  $y = e^{-4x} + 5$
- (b)  $y = e^{-4x} + 1.25$
- (c)  $y = e^{4x} + 5$
- (d)  $y = e^{4x} + 1.25$

18. [MCQ] [GATE-CE-2019:2M]

An ordinary differential equation is given below.

$$\left(\frac{dy}{dx}\right)(x \ln x) = y$$

The solution for the above equation is

**Note:** K denotes a constant in the options

- (a)  $y = Kx e^x$
- (b)  $y = Kx e^{-x}$
- (c)  $y = K \ln x$
- (d)  $Y = Kx \ln x$



**28. [MCQ]****[GATE-EC-2015: 2M]**

Consider the differential equation  $\frac{dx}{dt} = 10 - 0.2x$  with initial condition  $x(0) = 1$ . The response  $x(t)$  for  $t > 0$  is

- (a)  $2 - e^{-0.2t}$       (b)  $2 - e^{0.2t}$   
 (c)  $50 - 49e^{-0.2t}$       (d)  $50 - 49e^{0.2t}$

**29. [NAT]****[GATE-EE-2015 : 2M]**

A differential equation  $\frac{di}{dt} - 0.2i = 0$  is applicable over  $-10 < t < 10$ . If  $i(4) = 10$ , then  $i(-5)$  is \_\_\_\_\_.

**30. [MCQ]****[GATE-CE-2014:2M]**

The integrating factor for differential equation  $\frac{dP}{dt} + k_2 P = k_1 L_0 e^{-k_1 t}$  is

- (a)  $e^{-k_1 t}$       (b)  $e^{-k_2 t}$   
 (c)  $e^{k_1 t}$       (d)  $e^{k_2 t}$

**31. [MCQ]****[GATE-ME-2014: 1M]**

The solution of the initial value problem  $\frac{dy}{dx} = -2xy$  ;  $y(0) = 2$  is

- (a)  $1 + e^{-x^2}$       (b)  $2 e^{-x^2}$   
 (c)  $1 + e^{x^2}$       (d)  $2 e^{x^2}$

**32. [MCQ]****[GATE-ME-2014: 2M]**

The general solution of the differential equation  $\frac{dy}{dx} = \cos(x + y)$ , with  $c$  as a constant, is

- (a)  $y + \sin(x + y) = x + c$   
 (b)  $\tan\left(\frac{x+y}{2}\right) = y + c$   
 (c)  $\cos\left(\frac{x+y}{2}\right) = x + c$   
 (d)  $\tan\left(\frac{x+y}{2}\right) = x + c$

**33. [MCQ]****[GATE-EE-2012 : 1M]**

With initial condition  $x(1) = 0.5$ , the solution of the differential equation,  $t \frac{dx}{dt} + x = t$  is

- (a)  $x = t - \frac{1}{2}$       (b)  $x = t^2 - \frac{1}{2}$   
 (c)  $x = \frac{t^2}{2}$       (d)  $x = \frac{t}{2}$

**34. [MCQ]****[GATE-EE-2011: 1M]**

With  $K$  as a constant the possible solution for the first order differential equation  $\frac{dy}{dx} = e^{-3x}$  is

- (a)  $-\frac{1}{3}e^{-3x} + K$       (b)  $-\frac{1}{3}e^{3x} + K$   
 (c)  $-3e^{-3x} + K$       (d)  $-3e^{-x} + K$

**35. [MCQ]****[GATE-ME-2011: 2M]**

Consider the differential equation  $\frac{dy}{dx} = (1 + y^2)x$ .

The general solution with constant  $c$  is

- (a)  $y = \tan\left(\frac{x^2}{2}\right) + \tan c$       (b)  $y = \tan^2\left(\frac{x}{2} + c\right)$   
 (c)  $y = \tan^2\left(\frac{x}{2}\right) + c$       (d)  $y = \tan\left(\frac{x^2}{2} + c\right)$

**36. [MCQ]****[GATE-CE-2011:2M]**

The solution of the differential equation  $\frac{dy}{dx} + \frac{y}{x} = x$ , with the condition that  $y = 1$  at  $x = 1$ , is

- (a)  $y = \frac{2}{3x^2} + \frac{x}{3}$       (b)  $y = \frac{x}{2} + \frac{1}{2x}$   
 (c)  $y = \frac{2}{3} + \frac{x}{3}$       (d)  $y = \frac{2}{3x} + \frac{x^2}{3}$

**37. [MCQ]****[GATE-CE-2011:2M]**

The solution of the ordinary differential equation  $\frac{dy}{dx} + 2y = 0$  for the boundary condition,  $y = 5$  at  $x = 1$  is

- (a)  $y = e^{-2x}$       (b)  $y = 2e^{-2x}$   
 (c)  $y = 10.95 e^{-2x}$       (d)  $y = 36.95 e^{-2x}$

**38. [MCQ]**

[GATE-EC-2011:1M]

The solution of the differential equation  $\frac{dy}{dx} = ky$ ,  
 $y(0) = c$  is

- (a)  $x = c e^{-ky}$       (b)  $x = k e^{cy}$   
 (c)  $y = c e^{kx}$       (d)  $y = c e^{-kx}$

**39. [MCQ]**

[GATE-ME-2009: 2M]

The solution of  $x \frac{dy}{dx} + y = x^4$  with the condition

$y(1) = \frac{6}{5}$  is

- (a)  $y = \frac{x^4}{5} + \frac{1}{x}$       (b)  $y = \frac{4x^4}{5} + \frac{4}{5x}$   
 (c)  $y = \frac{x^4}{5} + 1$       (d)  $y = \frac{x^5}{5} + 1$

**40. [MCQ]**

[GATE-CE-2009:2M]

Solution of the differential equation  $3y \frac{dy}{dx} + 2x = 0$

represents a family of

- (a) ellipses      (b) circles  
 (c) parabolas      (d) hyperbolas

**41. [MCQ]**

[GATE-EC- 2009:2M]

Match each differential equation in Group I to its family of solution curves from Group II

|    | <b>Group I</b>                 |    | <b>Group II</b> |
|----|--------------------------------|----|-----------------|
| P. | $\frac{dy}{dx} = \frac{y}{x}$  | 1. | Circles         |
| Q. | $\frac{dy}{dx} = -\frac{y}{x}$ | 2. | Straight lines  |
| R. | $\frac{dy}{dx} = \frac{x}{y}$  | 3. | Hyperbolas      |
| S. | $\frac{dy}{dx} = -\frac{x}{y}$ |    |                 |

Codes:

- |   |   |   |   |
|---|---|---|---|
| P | Q | R | S |
|---|---|---|---|
- (a) 2 3 3 1  
 (b) 1 3 2 1  
 (c) 2 1 3 3  
 (d) 3 2 1 2

**42. [MCQ]**

[GATE-CE-2008:2M]

Solution of  $\frac{dy}{dx} = -\frac{x}{y}$  at  $x = 1$   $y = \sqrt{3}$  is

- (a)  $x - y^2 = -2$   
 (b)  $x + y^2 = 4$   
 (c)  $x^2 - y^2 = -2$   
 (d)  $x^2 + y^2 = 4$

**43. [MCQ]**

[GATE-EC-2008:1M]

Which of the following is a solution to the differential equation  $\frac{dx(t)}{dt} + 3x(t) = 0$ ?

- (a)  $x(t) = 3e^{-t}$   
 (b)  $x(t) = 2e^{-3t}$   
 (c)  $x(t) = -\frac{3}{2}t^2$   
 (d)  $x(t) = 3t^2$

**Higher Order Equations**

**44. [MCQ]**

[GATE-CE-2023:2M]

The solution of the differential equation

$$\frac{d^3y}{dx^3} - 5.5 \frac{d^2y}{dx^2} + 9.5 \frac{dy}{dx} - 5y = 0$$

is expressed as  $y = C_1 e^{2.5x} + C_2 e^{\alpha x} + C_3 e^{\beta x}$ , where  $C_1$ ,  $C_2$ ,  $C_3$ ,  $\alpha$ , and  $\beta$  are constants, with  $\alpha$  and  $\beta$  being distinct and not equal to 2.5. Which of the following options is correct for the values of  $\alpha$  and  $\beta$ ?

- (a) 1 and 2      (b) -1 and -2  
 (c) 2 and 3      (d) -2 and -3

**45. [NAT]**

[GATE- EC-2023: 2M]

The general solution of the differential equation,

$$\frac{d^2y}{dx^2} + 2 \frac{dy}{dx} - 5y = 0$$

In terms of arbitrary constants  $K_1$  and  $K_2$  is

- (a)  $K_1 e^{(-1+\sqrt{6})x} + K_2 e^{(-1-\sqrt{6})x}$   
 (b)  $K_1 e^{(-1+\sqrt{8})x} + K_2 e^{(-1-\sqrt{8})x}$   
 (c)  $K_1 e^{(-2+\sqrt{6})x} + K_2 e^{(-2-\sqrt{6})x}$   
 (d)  $K_1 e^{(-2+\sqrt{8})x} + K_2 e^{(-2-\sqrt{8})x}$

**46. [MCQ]****[GATE-CE-2021:2M]**

The solution of the second-order differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0 \text{ with boundary condition } y(0) = 1$$

and  $y(1) = 3$  is

- (a)  $e^{-x} + (3e - 1)x e^{-x}$
- (b)  $e^{-x} + \left[ 3\sin\left(\frac{\pi x}{2}\right) - 1 \right] x e^{-x}$
- (c)  $e^{-x} - (3e - 1)x e^{-x}$
- (d)  $e^{-x} - \left[ 3\sin\left(\frac{\pi x}{2}\right) - 1 \right] x e^{-x}$

**47. [MCQ]****[GATE-CE-2020:2M]**

An ordinary differential equation is given below

$$6\frac{d^2y}{dx^2} + \frac{dy}{dx} - y = 0$$

The general solution of the above equation (with constants  $C_1$  and  $C_2$ ) is

- (a)  $y(x) = C_1 e^{-\frac{x}{3}} + C_2 x e^{\frac{x}{2}}$
- (b)  $y(x) = C_1 e^{-\frac{x}{3}} + C_2 e^{\frac{x}{2}}$
- (c)  $y(x) = C_1 x e^{-\frac{x}{3}} + C_2 e^{\frac{x}{2}}$
- (d)  $y(x) = C_1 e^{\frac{x}{3}} + C_2 e^{-\frac{x}{2}}$

**48. [MCQ]****[GATE-CE-2020:2M]**

For the Ordinary Differential Equation

$$\frac{d^2x}{dt^2} - 5\frac{dx}{dt} + 6x = 10, \text{ with initial condition } x(0)=0$$

and  $\frac{dx}{dt}(0) = 10$

- (a)  $-10e^{2t} + 10e^{3t}$
- (b)  $5e^{2t} + 6e^{3t}$
- (c)  $10e^{2t} + 10e^{3t}$
- (d)  $-5e^{2t} + 6e^{3t}$

**49. [NAT]****[GATE-ME-2018: 2M]**

Given the ordinary differential equation

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 0$$

with  $y(0) = 0$  and  $\frac{dy}{dx}(0) = 1$ , the value of  $y(1)$  is \_\_\_\_\_ (Correct to two decimal places).

**50. [NAT]****[GATE-CE-2018: 2M]**

The solution (up to three decimal places) at  $x = 1$  of the differential equation  $\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0$  subject to boundary conditions  $y(0) = 1$  and  $\frac{dy}{dx}(x=0) = -1$  is \_\_\_\_\_.

**51. [NAT]****[GATE-EC-2018: 2M]**

The position of a particle  $y(t)$  is described by the differential equation:

$$\frac{d^2y}{dt^2} = -\frac{dy}{dt} - \frac{5y}{4}.$$

The initial conditions are  $y(0) = 1$  and  $\frac{dy}{dt}\Big|_{t=0} = 0$ .

The position (accurate to two decimal places) of the particle at  $t = \pi$  is \_\_\_\_\_

**52. [MCQ]****[GATE-CE-2017:2M]**

Consider the following second-order differential equation:  $y'' - 4y' + 3y = 2t - 3t^2$

The particular solution of the differential equation is

- (a)  $-2 - 2t - t^2$
- (b)  $-2t - t^2$
- (c)  $2t - t^2$
- (d)  $-2 - 2t - 3t^2$

**53. [MCQ]****[GATE-ME-2017: 1M]**

The differential equation  $\frac{d^2y}{dx^2} + 16y = 0$  for  $y(x)$  with

the two boundary conditions  $\frac{dy}{dx}\Big|_{x=0} = 1$  and  $\frac{dy}{dx}\Big|_{x=\frac{\pi}{2}} = -1$  has

- (a) No solution
- (b) Exactly two solutions
- (c) Exactly one solution
- (d) Infinitely many solutions

- 54. [NAT]** [GATE-ME-2017: 2M]  
Consider the differential equation  $3y''(x) + 27 y(x) = 0$  with initial conditions  $y(0) = 0$  and  $y'(0) = 2000$ . The value of  $y$  at  $x = 1$  is \_\_\_\_\_.

- 55. [NAT]** [GATE-ME-2016: 2M]  
If  $y = f(x)$  satisfies the boundary value problem  $y'' + 9y = 0$ ,  $y(0) = 0$ ,  $y(\pi/2) = \sqrt{2}$  then  $y(\pi/4)$  is \_\_\_\_\_.

- 56. [MCQ]** [GATE-CE-2016: 2M]  
The respective expressions for complimentary function and particular integral part of the solution of the differential equation  $\frac{d^4y}{dx^4} + 3\frac{d^2y}{dx^2} = 108x^2$  are

(a)  $[C_1 + C_2x + C_3\sin\sqrt{3x} + C_4\cos\sqrt{3x}]$

and  $[3x^4 - 12x^2 + c]$

(b)  $[C_2 + C_3\sin\sqrt{3x} + C_4\cos\sqrt{3x}]$

and  $[5x^4 - 12x^2 + c]$

(c)  $[C_1 + C_3\sin\sqrt{3x} + C_4\cos\sqrt{3x}]$

and  $[3x^4 - 12x^2 + c]$

(d)  $[C_1 + C_2x + C_3\sin\sqrt{3x} + C_4\cos\sqrt{3x}]$

And  $[5x^4 - 12x^2 + c]$

- 57. [MCQ]** [GATE-EC-2016: 2M]  
The particular solution of the initial value problem given below is

$$\frac{d^2y}{dx^2} + 12\frac{dy}{dx} + 36y = 0$$

with  $y(0) = 3$   $\left.\frac{dy}{dx}\right|_{x=0} = -36$

(a)  $(3 - 18x)e^{-6x}$       (b)  $(3 + 25x)e^{-6x}$

(c)  $(3 + 20x)e^{-6x}$       (d)  $(3 - 12x)e^{-6x}$

- 58. [MCQ]** [GATE-EE-2016: 1M]  
A function  $y(t)$ , such that  $y(0) = 1$  and  $y(1) = 3e^{-1}$ , is a solution of the differential equation  $\frac{d^2y}{dt^2} + 2\frac{dy}{dt} + y = 0$ . Then  $y(2)$  is

- (a)  $5e^{-1}$       (b)  $5e^{-2}$   
(c)  $7e^{-1}$       (d)  $7e^{-2}$

- 59. [NAT]** [GATE-EE-2016: 2M]  
Let  $y(x)$  be the solution of the differential equation

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 0 \text{ with initial conditions } y(0) = 0 \text{ and } \left.\frac{dy}{dx}\right|_{x=0} = 1.$$

Then the value of  $y(1)$  is \_\_\_\_\_

- 60. [MCQ]** [GATE 2016-EE: 1M]  
The solution of the differential equation for  $t > 0$ ,  $y''(t) + 2y'(t) + y(t) = 0$  with initial condition  $y(0) = 0$  and  $y'(0) = 1$ , is ( $u(t)$  denotes the unit step function),

- (a)  $te^{-t}u(t)$   
(b)  $(e^{-t} - te^{-t})u(t)$   
(c)  $(-e^{-t} + te^{-t})u(t)$   
(d)  $e^{-t}u(t)$

- 61. [NAT]** [GATE-EC-2015: 2M]  
Consider the differential equation

$$\frac{d^2x(t)}{dt^2} + 3\frac{dx(t)}{dt} + 2x(t) = 0$$

Given  $x(0) = 20$  and  $x(1) = 10/e$ ,

where  $e = 2.718$ , the value of  $x(2)$  is \_\_\_\_\_.

- 62. [MCQ]** [GATE-EC-2015: 2M]  
The solution of the differential equation

$$\frac{d^2y}{dt^2} + 2\frac{dy}{dt} + y = 0 \text{ with } y(0) = y'(0) = 1 \text{ is}$$

- (a)  $(2-t)e^t$       (b)  $(1+2t)e^{-t}$   
(c)  $(2+t)e^{-t}$       (d)  $(1-2t)e^t$

- 63. [NAT]** [GATE-CE-2015: 2M]  
Consider the following second order linear

$$\text{differential equation } \frac{d^2y}{dx^2} = -12x^2 + 24x - 20.$$

The boundary conditions are: at  $x = 0$ ,  $y = 5$  and  $x = 2$ ,  $y = 21$

The value of  $y$  at  $x = 1$  is \_\_\_\_\_

**64. [MCQ]**

[GATE-ME-2015: 2M]

Find the solution of  $\frac{d^2y}{dx^2} = y$  which passes through the origin and the point  $(\ln 2, \frac{3}{4})$ .

- (a)  $y = \frac{1}{2}e^x - e^{-x}$       (b)  $y = \frac{1}{2}(e^x + e^{-x})$   
 (c)  $y = \frac{1}{2}(e^x - e^{-x})$       (d)  $y = \frac{1}{2}e^x + e^{-x}$

**65. [NAT]**

[GATE-ME-2014: 2M]

If  $y = f(x)$  is the solution of  $\frac{d^2y}{dx^2} = 0$  with the boundary conditions  $y = 5$  at  $x = 0$  and  $\frac{dy}{dx} = 2$  at  $x = 10, f(15) = \underline{\hspace{2cm}}$ .

**66. [MCQ]**

[GATE-ME-2014: 2M]

Consider two solutions  $x(t) = x_1(t)$  and  $x(t) = x_2(t)$  of the differential equation

$$\frac{d^2x(t)}{dt^2} + x(t) = 0, t > 0, \text{ such that } x_1(0) = 1,$$

$$\left. \frac{dx_1(t)}{dt} \right|_{t=0} = 0, x_2(0) = 0, \left. \frac{dx_2(t)}{dt} \right|_{t=0} = 1.$$

The Wronskian  $W(t) = \begin{vmatrix} x_1(t) & x_2(t) \\ \frac{dx_1(t)}{dt} & \frac{dx_2(t)}{dt} \end{vmatrix}$  at  $t = \frac{\pi}{2}$  is

- (a) 1      (b) -1  
 (c) 0      (d)  $\pi/2$

**67. [MCQ]**

[GATE-EC-2014: 1M]

If the characteristic equation of the differential

equation  $\frac{d^2y}{dx^2} + 2\alpha \frac{dy}{dx} + y = 0$  has two equal roots,

then the value of  $\alpha$  are

- (a)  $\pm 1$       (b) 0, 0  
 (c)  $\pm j$       (d)  $\pm \frac{1}{2}$

**68. [MCQ]**

[GATE-EC-2014: 1M]

If  $a$  and  $b$  are constants, the most general solution of

the differential equation  $\frac{d^2x}{dt^2} + 2 \frac{dx}{dt} + x = 0$  is

- (a)  $ae^{-t}$       (b)  $ae^{-t} + bte^{-t}$   
 (c)  $ae^t + bte^{-t}$       (d)  $ae^{-2t}$

**69. [NAT]**

[GATE-EC-2014: 2M]

With initial values  $y(0) = y'(0) = 1$ , the solution of

the differential equation  $\frac{d^2y}{dx^2} + 4 \frac{dy}{dx} + 4y = 0$  at  $x = 1$  is \_\_\_\_\_.

**70. [MCQ]**

[GATE-EE-2014: 1M]

Consider the differential equation  $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} - y = 0$ .

Which of the following is a solution to this differential equation for  $x > 0$ ?

- (a)  $e^x$       (b)  $X^2$   
 (c)  $1/x$       (d)  $\ln x$

**71. [MCQ]**

[GATE-EE-2014: 1M]

The solution for the differential equation

$$\frac{d^2x}{dt^2} = -9x,$$

With initial conditions  $x(0) = 1$  and  $\left. \frac{dx}{dt} \right|_{t=0} = 1$ , is

- (a)  $t^2 + t = 1$   
 (b)  $\sin 3t + \frac{1}{3}\cos 3t + \frac{2}{3}$   
 (c)  $\frac{1}{3}\sin 3t + \cos 3t$   
 (d)  $\cos 3t + t$

**72. [MCQ]**

[GATE-ME-2013: 2M]

The solution to the differential equation

$\frac{d^2u}{dx^2} - k \frac{du}{dx} = 0$  where  $k$  is a constant, subjected to the

boundary conditions  $u(0) = 0$  and  $u(L) = U$ , is

- (a)  $u = U \frac{x}{L}$       (b)  $u = U \left( \frac{1 - e^{kx}}{1 - e^{kL}} \right)$   
 (c)  $u = U \left( \frac{1 - e^{-kx}}{1 - e^{-kL}} \right)$       (d)  $u = U \left( \frac{1 + e^{kx}}{1 + e^{kL}} \right)$

**73. [MCQ]**

[GATE-CE-2010: 2M]

The solution to the ordinary differential equation

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 0$$

- (a)  $y = c_1 e^{3x} + c_2 e^{-2x}$       (b)  $y = c_1 e^{3x} + c_2 e^{2x}$   
 (c)  $y = c_1 e^{-3x} + c_2 e^{2x}$       (d)  $y = c_1 e^{-3x} + c_2 e^{-2x}$

**74. [MCQ]****[GATE-EC-2010: 1M]**

A function  $n(x)$  satisfies the differential equation  $\frac{d^2 n(x)}{dx^2} - \frac{n(x)}{L^2} = 0$  where  $L$  is a constant. The boundary conditions are  $n(0) = K$  and  $n(\infty) = 0$ . The solution to this equation is

- (a)  $n(x) = K \exp(x/L)$
- (b)  $n(x) = K \exp(-x/\sqrt{L})$
- (c)  $n(x) = K^2 \exp(-x/L)$
- (d)  $n(x) = K \exp(-x/L)$

**75 [MCQ]****[GATE-EE-2010: 2M]**

For the difference equation  $\frac{d^2 x}{dt^2} + 6 \frac{dx}{dt} + 8x = 0$

with initial condition  $x(0) = 1$  and  $\left. \frac{dx}{dt} \right|_{t=0} = 0$ .

- (a)  $x(t) = 2e^{-6t} - e^{-2t}$
- (b)  $x(t) = 2e^{-2t} - e^{-4t}$
- (c)  $x(t) = -e^{-6t} + e^{-4t}$
- (d)  $x(t) = e^{-2t} + 2e^{-4t}$

**76. [MCQ]****[GATE-ME-2008: 1M]**

Given that  $\ddot{x} + 3x = 0$ , and  $x(0) = 1$ ,  $\dot{x}(0) = 0$ , what is  $x(1)$ ?

- (a) -0.99
- (b) -0.16
- (c) 0.16
- (d) 0.99

**77. [MCQ]****[GATE-ME-2008: 2M]**

It is given  $y'' + 2y' + y = 0$ ,  $y(0) = 0$ ,  $y(1) = 0$ . What is  $y(0.5)$ ?

- (a) 0
- (b) 0.37
- (c) 0.62
- (d) 1.13

**78. [MCQ]****[GATE-CE-2008:1M]**

The general solution of  $\frac{d^2 y}{dx^2} + y = 0$  is

- (a)  $Y = P \cos x + Q \sin x$
- (b)  $y = p \cos x$
- (c)  $y = P \sin x$
- (d)  $y = P \sin 2x$

**Euler -Cauchy Equations****79. [NAT]****[GATE-ME-2023: 2M]**

Consider the second-order linear ordinary differential equation  $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - y = 0$ ,  $x \geq 1$

with the initial conditions

$$y(x=1) = 6, \quad \left. \frac{dy}{dx} \right|_{x=1} = 2$$

The value of  $y$  at  $x = 2$  equals \_\_\_\_\_. (Answer in integer)

**80. [MCQ]****[GATE-ME-2019: 2M]**

A differential equation is given as

$$x^2 \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + 2y = 4$$

The solution of the differential equation in terms of arbitrary constants  $C_1$  and  $C_2$  is

- (a)  $y = C_1 x^2 + C_2 x + 2$
- (b)  $y = \frac{C_1}{x^2} + C_2 x + 2$
- (c)  $y = C_1 x^2 + C_2 x + 4$
- (d)  $y = \frac{C_1}{x^2} + C_2 x + 4$

**81. [NAT]****[GATE-CE-2019:2M]**

Consider the ordinary differential equation  $x^2 \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + 2y = 0$ . Given the values of  $y(1) = 0$  and  $y(2) = 2$ , the value of  $y(3)$  (round off to 1 decimal place) is \_\_\_\_\_.

**82. [NAT]****[GATE-EC-2019: 2M]**

Consider the homogeneous ordinary differential equation,

$$x^2 \frac{d^2 y}{dx^2} - 3x \frac{dy}{dx} + 3y = 0, \quad x > 0$$

with  $y(x)$  as a general solution.

Given that  $y(1) = 1$  and  $y(2) = 14$  the value of  $y(1.5)$  (rounded off to two decimal places), is \_\_\_\_\_.

**83. [MCQ]****[GATE-ME-2012: 2M]**

Consider the differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - 4y = 0$$
 with the boundary conditions

of  $y(0) = 0$  and  $y(1) = 1$ . The complete solution of the differential equation is

- (a)  $x^2$
- (b)  $\sin\left(\frac{\pi x}{2}\right)$
- (c)  $e^x \sin\left(\frac{\pi x}{2}\right)$
- (d)  $e^{-x} \sin\left(\frac{\pi x}{2}\right)$

## Partial Differential Equations

- 84. [NAT] [GATE-CE-2023-1M]**  
A 5 cm long metal rod AB was initially at a uniform temperature of  $T_0$  °C. Thereafter, temperature at both the ends are maintained at 0 °C. Neglecting the heat transfer from the lateral surface of the rod, the heat transfer in the rod is governed by the one-dimensional diffusion equation  $\frac{\partial T}{\partial t} = D \frac{\partial^2 T}{\partial x^2}$ , where  $D$  is the thermal diffusivity of the metal, given as 1.0 cm<sup>2</sup>/s. The temperature distribution in the rod is obtained as  $T(x, t) = \sum_{n=1,3,5,\dots}^{\infty} C_n \sin \frac{n\pi x}{5} e^{-\beta n^2 t}$ , where  $x$  is in cm measured from A to B with  $x=0$  at A,  $t$  is in s,  $C_n$  are constants in °C,  $T$  is in °C, and  $\beta$  is in s<sup>-1</sup>. The value of  $\beta$  (in s<sup>-1</sup>, rounded off to three decimal places) is \_\_\_\_\_.

- 85. [NAT] [GATE-CE-2023-2M]**  
The steady-state temperature distribution in a square plate ABCD is governed by the 2-dimensional Laplace equation. The side AB is kept at a temperature of 100 °C and the other three sides are kept at a temperature of 0 °C. Ignoring the effect of discontinuities in the boundary conditions at the corners, the steady-state temperature at the center of the plate is obtained as  $T_0$  °C. Due to symmetry, the steady-state temperature at the center will be same ( $T_0$  °C), when any one side of the square is kept at a temperature of 100 °C and the remaining three sides are kept at a temperature of 0 °C. Using the principle of superposition, the value of  $T_0$  is \_\_\_\_\_ (rounded off to two decimal places).

- 86. [MCQ] [GATE-CE-2022-1M]**

Consider the following expression:

$$Z = \sin(y + it) + \cos(y - it)$$

Where  $z$ ,  $y$  and  $t$  are variables and  $i = \sqrt{-1}$  is a complex number. The partial differential equation derived from the above expression is

- (a)  $\frac{\partial^2 Z}{\partial t^2} + \frac{\partial^2 Z}{\partial y^2} = 0$       (b)  $\frac{\partial^2 Z}{\partial t^2} - \frac{\partial^2 Z}{\partial y^2} = 0$   
(c)  $\frac{\partial Z}{\partial t} - i \frac{\partial Z}{\partial y} = 0$       (d)  $\frac{\partial Z}{\partial t} + i \frac{\partial Z}{\partial y} = 0$

- 87. [MCQ] [GATE- CE-2022:1M]**

The function  $f(x, y)$  satisfies the Laplace equation  $\Delta^2 f(x, y) = 0$  on a circular domain of radius  $r = 1$  with its centre at point P with coordinates  $x = 0, y = 0$ . The value of this function on the circular boundary of this domain is equal to 3. The numerical value of  $f(0, 0)$  is

- (a) 1      (b) 0  
(c) 2      (d) 3

- 88. [MSQ] [GATE-EC-2022: 1M]**  
Consider the following partial differential equation (PDE),

$$a \frac{\partial^2 f(x, y)}{\partial x^2} + b \frac{\partial^2 f(x, y)}{\partial y^2} = f(x, y),$$

where  $a$  and  $b$  are distinct positive real numbers. Select the combination(s) of values of the real parameters  $\xi$  and  $\eta$  such that  $f(x, y) = e^{(\xi x + \eta y)}$  is a solution of the given PDE.

- (a)  $\xi = \frac{1}{\sqrt{2a}}, \eta = \frac{1}{\sqrt{2b}}$   
(b)  $\xi = \frac{1}{\sqrt{a}}, \eta = 0$   
(c)  $\xi = 0, \eta = 0$   
(d)  $\xi = \frac{1}{\sqrt{a}}, \eta = \frac{1}{\sqrt{b}}$

- 89. [MCQ] [GATE-CE-2020-1M]**

In the following partial differential equation,  $\theta$  is a function of  $t$  and  $z$ , and  $D$  and  $K$  are function of  $\theta$

$$D(\theta) \frac{\partial^2 \theta}{\partial z^2} + \frac{\partial K(\theta)}{\partial z} - \frac{\partial \theta}{\partial t} = 0$$

The above equation is

- (a) A second order linear equation  
(b) A second order non – linear equation  
(c) A second degree non-linear equation  
(d) A second degree linear equation

**90. [MCQ]**

**[GATE-CE-2020-1M]**

The following partial differential equation is defined for

$$u : u(x, y) \frac{\partial u}{\partial y} = \frac{\partial^2 y}{\partial x^2} : y \geq 0, x_1 \leq x \leq x_2$$

The set of auxiliary conditions necessary to solve the equation uniquely, is

- (a) One initial condition and two boundary conditions
- (b) Three initial conditions
- (c) Two initial conditions and one boundary condition
- (d) Three boundary conditions

**91. [MCQ]**

**[GATE-EE-2019 : 1M]**

The partial differential equation

$$\frac{\partial^2 u}{\partial t^2} - c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) = 0;$$

where  $c \neq 0$  is known as

- (a) heat equation
- (b) wave equation
- (c) Poisson's equation
- (d) Laplace equation

**92. [MCQ]**

**[GATE-ME-2018: 1M]**

Consider a function  $u$  which depends on position  $x$  and time  $t$ . The partial differential equation

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$$

is known as the

- (a) Wave equation
- (b) Heat equation
- (c) Laplace's equation
- (d) Elasticity equation

**93. [MCQ]**

**[GATE-ME-2017: 1M]**

Consider the following partial differential equation for  $u(x, y)$  with the constant  $c > 1$ :

$$\frac{\partial u}{\partial y} + c \frac{\partial u}{\partial x} = 0$$

Solution of this equation is

- (a)  $u(x, y) = f(x + cy)$
- (b)  $u(x, y) = f(x - cy)$
- (c)  $u(x, y) = f(cx + y)$
- (d)  $u(x, y) = f(cx - y)$

**94. [NAT]**

**[GATE-CE-2017-1M]**

Consider the following partial differential equation:

$$3 \frac{\partial^2 \phi}{\partial x^2} + B \frac{\partial^2 \phi}{\partial x \partial y} + 3 \frac{\partial^2 \phi}{\partial y^2} + 4\phi = 0$$

For this equation to be classified as parabolic, the value of  $B^2$  must be \_\_\_\_\_.

**95. [MCQ]**

**[GATE-CE-2017-1M]**

The solution at  $x = 1, t = 1$  of the partial differential

$$\text{equation } \frac{\partial^2 u}{\partial x^2} = 25 \frac{\partial^2 u}{\partial t^2} \text{ subject to initial conditions}$$

of  $u(x, 0) = 3x$  and  $\frac{\partial u}{\partial t}(t=0) = 3$  is \_\_\_\_\_.

- (a) 1
- (b) 2
- (c) 3
- (d) 6

**96. [MCQ]**

**[GATE-CE-2016-1M]**

The type of partial differential equation

$$\frac{\partial^2 P}{\partial x^2} + \frac{\partial^2 P}{\partial y^2} + 3 \frac{\partial^2 P}{\partial x \partial y} + 2 \frac{\partial P}{\partial x} - \frac{\partial P}{\partial y} = 0 \text{ is}$$

- (a) Elliptic
- (b) Parabolic
- (c) Hyperbolic
- (d) None of these

**97. [MCQ]**

**[GATE-CE-2016-1M]**

The solution of the partial differential equation

$$\frac{\partial u}{\partial t} = \alpha \frac{\partial^2 u}{\partial x^2} \text{ is of the form}$$

- (a)  $C \cos(kt) \left| C_1 e^{(\sqrt{k/\alpha})x} + C_2 e^{-(\sqrt{k/\alpha})x} \right|$
- (b)  $C e^{kt} \left| C_1 e^{(\sqrt{k/\alpha})x} + C_2 e^{-(\sqrt{k/\alpha})x} \right|$
- (c)  $C e^{kt} \left| C_1 \cos(\sqrt{k/\alpha})x + C_2 \sin(-\sqrt{k/\alpha})x \right|$
- (d)  $C \sin(kt) \left| C_1 \cos(\sqrt{k/\alpha})x + C_2 \sin(-\sqrt{k/\alpha})x \right|$

**98. [MCQ]**

**[GATE-ME-2015: 2M]**

Solutions of Laplace's equation having continuous second-order partial derivatives are called

- (a) Biharmonic functions
- (b) Harmonic functions
- (c) Conjugate harmonic functions
- (d) Error functions

**99. [MCQ]**

**[GATE-EC-2014: 1M]**

If  $z = xy \ln(xy)$ , then

- (a)  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 0$
- (b)  $y \frac{\partial z}{\partial x} = x \frac{\partial z}{\partial y}$
- (c)  $x \frac{\partial z}{\partial x} = y \frac{\partial z}{\partial y}$
- (d)  $y \frac{\partial z}{\partial x} + x \frac{\partial z}{\partial y} = 0$

**100. [MCQ]****[GATE-CE-2010-2M]**

The partial differential equation that can be formed from  $z = ax + by + ab$  has the form (with  $p = \frac{\partial z}{\partial x}$  and  $q = \frac{\partial z}{\partial y}$ ).

- (a)  $z = px + qy$       (b)  $z = px + pq$   
 (c)  $z = px + qy + pq$       (d)  $z = qy + pq$

**101. [MCQ]****[GATE-ME-2008: 2M]**

Let  $f = y^x$ . What is  $\frac{\partial^2 f}{\partial x \partial y}$  at  $x = 2, y = 1$ ?

- (a) 0      (b)  $\ln 2$   
 (c) 1      (d)  $\frac{1}{\ln 2}$

**102. [MCQ]****[GATE-CE-2008-2M]**

The equation  $k_x \frac{\partial^2 h}{\partial x^2} + k_z \frac{\partial^2 h}{\partial z^2} = 0$  . can be transformed to  $\frac{\partial^2 h}{\partial x_t^2} + \frac{\partial^2 h}{\partial z^2} = 0$  . by substituting.

- (a)  $x_t = x \frac{k_z}{k_x}$       (b)  $x_t = x \frac{k_x}{k_z}$   
 (c)  $x_t = x \sqrt{\frac{k_x}{k_z}}$       (d)  $x_t = x \sqrt{\frac{k_z}{k_x}}$




**ANSWER KEY**

- |                    |                    |                        |                     |
|--------------------|--------------------|------------------------|---------------------|
| 1. (a)             | 2. (d)             | 3. (a)                 | 4. (d)              |
| 5. (a)             | 6. (b)             | 7. (b)                 | 8. (0.446 to 0.446) |
| 9. (d)             | 10. (c)            | 11. (c)                | 12. (c)             |
| 13. (d)            | 14. (b)            | 15. (0.8774 to 0.8952) | 16. (c)             |
| 17. (b)            | 18. (c)            | 19. (c)                | 20. (c)             |
| 21. (a)            | 22. (d)            | 23. (d)                | 24. (a)             |
| 25. (c)            | 26. (c)            | 27. (c)                | 28. (c)             |
| 29. (1.6 to 1.7)   | 30. (d)            | 31. (b)                | 32. (d)             |
| 33. (d)            | 34. (a)            | 35. (d)                | 36. (d)             |
| 37. (d)            | 38. (c)            | 39. (a)                | 40. (a)             |
| 41. (a)            | 42. (d)            | 43. (b)                | 44. (a)             |
| 45. (a)            | 46. (a)            | 47. (d)                | 48. (a)             |
| 49. (1.45 to 1.48) | 50. (0.36 to 0.37) | 51. (- 0.23 to - 0.19) | 52. (a)             |
| 53. (a)            | 54. (93 to 95)     | 55. (-1.05 to -0.95)   | 56. (a)             |
| 57. (a)            | 58. (b)            | 59. (7.0 to 7.5)       | 60. (a)             |
| 61. (0.83 to 0.88) | 62. (b)            | 63. (18)               | 64. (c)             |
| 65. (34 to 36)     | 66. (a)            | 67. (a)                | 68. (b)             |
| 69. (0.53 to 0.55) | 70. (c)            | 71. (c)                | 72. (b)             |
| 73. (c)            | 74. (d)            | 75. (b)                | 76. (b)             |
| 77. (a)            | 78. (a)            | 79. (8.999 to 9.001)   | 80. (a)             |
| 81. (5.9 to 6.1)   | 82. (5.24 to 5.26) | 83. (a)                | 84. (0.395)         |
| 85. (25)           | 86. (a)            | 87. (d)                | 88. (a, b)          |
| 89. (b)            | 90. (a)            | 91. (b)                | 92. (b)             |
| 93. (b)            | 94. (36)           | 95. (d)                | 96. (c)             |
| 97. (b)            | 98. (b)            | 99. (c)                | 100. (c)            |
| 101. (c)           | 102. (d)           |                        |                     |


**SOLUTIONS**

1. (a)

$$\frac{d^3y}{dx^3} + x \left( \frac{dy}{dx} \right)^{3/2} + x^2 y = 0$$

$$\Rightarrow \frac{d^3y}{dx^3} + x^2 y = -x \left( \frac{dy}{dx} \right)^{3/2}$$

Square both side;

$$\Rightarrow \left[ \frac{d^3y}{dx^3} + x^2 y \right]^2 = x^2 \left( \frac{dy}{dx} \right)^3$$

$$\Rightarrow \left( \frac{d^3y}{dx^3} \right)^2 + x^4 y^2 + 2x^2 y \left( \frac{d^3y}{dx^3} \right) = x^2 \left( \frac{dy}{dx} \right)^3$$

We have successfully eliminated radicals or square roots from this differential equation.

Order = Order of highest derivative = 3

Degree = Exponent of highest derivative = 2



Scan for Video solution



2. (d)

**Given:**

$$\frac{d^2u}{dx^2} - 2x^2 u + \sin x = 0$$

A differential equation is considered linear if it satisfies the following conditions:

- (i) The dependent variable and differential coefficient appear in the differential equation only in a linear manner. This means that the power or exponent of the dependent variable and its derivatives is always 1.
- (ii) The coefficients of the dependent variable and differential coefficient are constants or functions of the independent variables. There should not be product of dependent variable and differential coefficient or product of two differential coefficient.

**Given that the provided differential equation satisfies the above conditions, it can be concluded that it is a linear differential equation.**

$$U = f(x)$$

Here,  $u$  is dependent variable &  $x$  is independent variable. Since this equation can't be expressed as

$$x^n f\left(\frac{y}{x}\right) \therefore \text{It is non-homogeneous.}$$


Scan for Video solution



3. (a)

$$(i) \frac{dy}{dx} + xy = e^{-x} \rightarrow \text{Linear Non-Homogeneous DE}$$

$$(ii) \frac{dy}{dx} + xy = 0 \rightarrow \text{Linear but Homogeneous DE}$$

$$(iii) \frac{dy}{dx} + xy = e^{-y} \Rightarrow \frac{dy}{dx} + xy = \left\{ 1 - \frac{y}{1!} + \frac{y^2}{2!} - \frac{y^3}{3!} + \dots \infty \right\}$$

→ Non-linear DE

$$(iv) \frac{dy}{dx} + e^{-y} = 0 \Rightarrow \frac{dy}{dx} = -e^{-y} \rightarrow \text{Non-linear DE}$$


Scan for Video solution



4. (d)

$$\text{Given: } \frac{\partial u}{\partial t} + u \cdot \frac{\partial u}{\partial x} = \frac{\partial^2 u}{\partial x^2}$$

$$u \cdot \frac{\partial u}{\partial x} \rightarrow \text{Non-linear term and } \frac{\partial^2 u}{\partial x^2} \rightarrow \text{order '2'}$$

∴ Given DE is a non-linear DE of 2<sup>nd</sup> order.


Scan for Video solution



5. (a)

$$\frac{d^3y}{dx^3} + 4\sqrt{\left(\frac{dy}{dx}\right)^3 + y^2} = 0$$

$$\Rightarrow \frac{d^3y}{dx^3} = -4\sqrt{\left(\frac{dy}{dx}\right)^3 + y^2}$$

Making differential equation free from radicals, we get

$$\Rightarrow \left(\frac{d^3y}{dx^3}\right)^2 = 16\left[\left(\frac{dy}{dx}\right)^3 + y^2\right]$$

Order = Order of highest differential coefficient = 3

Degree = Exponent of highest order derivative = 2



Scan for Video solution



6. (b)

$$\text{Given: } \frac{d^3f}{d\eta^3} + \frac{f}{2} \cdot \frac{d^2f}{d\eta^2} = 0$$

Clearly given equation is non-linear (**Due to presence of  $f \cdot \frac{d^2f}{d\eta^2}$  term**) and the highest order derivative that is involved is of 3<sup>rd</sup> order.

$\therefore$  Given DE is [3<sup>rd</sup> order, non-linear ordinary DE].



Scan for Video solution



7. (b)

**Given:**

$$\text{DE is } \frac{d^2y}{dt^2} + \left(\frac{dy}{dt}\right)^3 + y^4 = e^{-t}$$

For the given DE, highest order derivative that is involved is '2'

$\therefore$  **order of the given DE is 2**



Scan for Video solution



8. (0.446 to 0.446)

$y = 1.0$  at  $x = 0.0$ , and  $y = 0.8$  at  $x = 1.0$

$$\frac{dy}{dx} + \alpha xy = 0$$

$$\Rightarrow \frac{dy}{y} = -\alpha x dx \quad [\text{Variable separable form}]$$

Integrating both sides,

$$\Rightarrow \ln y = -\alpha \frac{x^2}{2} + c \quad \dots(i)$$

Given,  $y = 1$  at  $x = 0$ ; from equation (i),

$$\ln 1 = 0 + c \Rightarrow c = 0$$

Put value of c in equation (i),

$$\ln y = -\alpha \frac{x^2}{2} \quad \dots(ii)$$

Given,  $y = 0.8$  at  $x = 1$ ; from equation (ii),

$$\ln 0.8 = -\alpha \frac{1^2}{2}$$

$$\Rightarrow \boxed{\alpha = 0.446}$$



Scan for Video solution



9. (d)

$$\text{Given: } t \cdot \frac{dx}{dt} + x = t; \quad x(1) = 0.5$$

$$t \cdot \frac{dx}{dt} + x = t;$$

$$\Rightarrow \frac{dx}{dt} + \frac{1}{t}x = 1$$

This is **Leibnitz linear equation** of form

$$\frac{dx}{dt} + P'x = Q' \text{ where } P' = \frac{1}{t} \text{ and } Q' = 1.$$

Solution of above differentiable equation is given by,

$$x \cdot (\text{I.F}) = \int Q'(\text{IF}) dt + c$$

$$\Rightarrow x \cdot (\text{I.F}) = \int (\text{IF}) dt + c \quad \dots(i)$$

Integrating factor

$$\text{I.F} = e^{\int \frac{1}{t} dt} = e^{\ln t} = t$$

From equation (i),

$$x \times t = \int t dt + c$$

$$\Rightarrow x \cdot t = \frac{t^2}{2} + c$$

$$\Rightarrow x = \frac{t}{2} + \frac{c}{t} \quad \dots(ii)$$

At  $t = 1$ ;  $x = 0.5$ , from equation (ii),

$$\Rightarrow 0.5 = \frac{1}{2} + c \Rightarrow c = 0$$

Put value of  $c$  in equation (ii),

$$x = \frac{t}{2}$$



Scan for Video solution



10. (c)

**Method I:**

$$\frac{du}{dx} = \frac{-xu^2}{2+x^2u}$$

$$\Rightarrow 2du + x^2 \cdot u du = -xu^2 dx$$

$$\Rightarrow 2du + x^2 u du + xu^2 dx = 0$$

$$\Rightarrow 2du + d\left(\frac{x^2 u^2}{2}\right) = 0$$

On integrating

$$\Rightarrow 2u + \frac{x^2 u^2}{2} = \text{constant}$$

**Method II:**

$$\frac{du}{dx} = \frac{-xu^2}{2+x^2u}$$

$$\Rightarrow (2+x^2u)du + xu^2 dx = 0 \quad \dots(1)$$

Comparing with  $M du + N dx = 0$

$$M = 2+x^2u \text{ and } N = xu^2$$

The above differential equation

$$\text{Will be exact if, } \frac{\delta M}{\delta x} = \frac{\delta N}{\delta u}$$

$$\text{Now, } \frac{\delta M}{\delta x} = 2xu \text{ and } \frac{\delta N}{\delta u} = 2xu$$

$$\therefore \frac{\delta M}{\delta x} = \frac{\delta N}{\delta u}$$

$\therefore$  differential equation (i) is exact differential equation and solution of differential equation is:

$$\int_{x=c} M du + \int (\text{Terms of } N \text{ not containing } u) dx = c$$

$$\Rightarrow \int_{x=c} (2+x^2u) du + \int (0) dx = c$$

$$\Rightarrow 2u + \frac{x^2 u^2}{2} = c$$



Scan for Video solution



11. (c)

$$\text{Given } y\left(\frac{\pi}{2}\right) = \frac{\pi}{2} \text{ and } \sin x \frac{dy}{dx} + y \cos x = 1$$

$$\text{Find: } y\left(\frac{\pi}{6}\right) = ?$$

$$\sin x \frac{dy}{dx} + y \cos x = 1$$

$$\Rightarrow \frac{dy}{dx} + \cot x \cdot y = \operatorname{cosec} x$$

This is **Leibnitz linear equation** of form

$$\frac{dy}{dx} + P' y = Q' \text{ where } P' = \cot x \text{ and } Q' = \operatorname{cosec} x .$$

Solution of above differentiable equation is given by,

$$y \cdot (I.F) = \int Q' (I.F) dx + c$$

$$\Rightarrow y \cdot (I.F) = \int \operatorname{cosec} x (I.F) dx + c \quad \dots(i)$$

Here, Integrating factor:

$$I.F = e^{\int \cot x dx} = e^{\int \frac{\cos x}{\sin x} dx} = e^{\ln|\sin x|} = \sin x$$

From equation (i),

$$y (\sin x) = \int \sin x \cdot \operatorname{cosec} x dx + c$$

$$\Rightarrow y (\sin x) = \int dx + c$$

$$\Rightarrow y (\sin x) = x + c \quad \dots(ii)$$

Given, At  $x = \frac{\pi}{2}$ ,  $y = \frac{\pi}{2}$ ; from equation (ii)

$$\Rightarrow \frac{\pi}{2}(1) = \frac{\pi}{2} + c \Rightarrow c = 0$$

Put value of c in equation (ii),

$$y(\sin x) = x$$

$$\Rightarrow y = \frac{x}{\sin x}$$

$$\text{At } x = \frac{\pi}{6};$$

$$y\left(\frac{\pi}{6}\right) = \frac{\frac{\pi}{6}}{\left(\frac{1}{2}\right)} = \frac{\pi}{3}$$

$$\Rightarrow \boxed{y\left(\frac{\pi}{6}\right) = \frac{\pi}{3}}$$



Scan for Video solution



12. (c)

$$\text{Given, } (1+y) \frac{dy}{dx} = y$$

$$\Rightarrow \frac{1+y}{y} \cdot dy = dx \quad [\text{Variable separable form}]$$

$$\Rightarrow \left(\frac{1}{y} + 1\right) dy = dx$$

Since  $y(1) = 1$ , hence integrating above differential equation from  $x = 1$  to  $x$  and  $y = 1$  to  $y$

$$\Rightarrow \int_{y=1}^y \left(\frac{1}{y} + 1\right) dy = \int_{x=1}^x dx$$

$$\Rightarrow \ln y + y \Big|_1^y = x \Big|_1^x$$

$$\Rightarrow (\ln y + y) - (1) = x - 1$$

$$\Rightarrow x = y + \ln y$$

$$\Rightarrow x - y = \ln y$$

$$\Rightarrow e^{x-y} = y$$

$$\Rightarrow e^x = y \cdot e^y$$

$$\Rightarrow \boxed{ye^y = e^x}$$



Scan for Video solution



13. (d)

$$\frac{dy}{dx} - \frac{y}{x} = 1$$

This is Leibnitz linear equation of form

$$\frac{dy}{dx} + P'y = Q' \text{ where } P' = -\frac{1}{x} \text{ and } Q' = 1.$$

Solution of above differentiable equation is given by,

$$y \cdot (\text{I.F.}) = \int Q'(\text{I.F.}) dx + c$$

$$\Rightarrow y \cdot (\text{I.F.}) = \int 1 \times (\text{I.F.}) dx + c \quad \dots(1)$$

Here, Integrating factor (I.F.);

$$\text{I.F.} = e^{\int P' dx} = e^{\int -\frac{1}{x} dx} = e^{-\ln x}$$

$$\Rightarrow \text{I.F.} = e^{\ln \frac{1}{x}} = \frac{1}{x}$$

From equation (i),

$$y\left(\frac{1}{x}\right) = \int \frac{1}{x} dx + c \quad [c = \ln k]$$

$$\Rightarrow \frac{y}{x} = \ln x + \ln k$$

$$\Rightarrow \boxed{y = x \ln(kx)}$$



Scan for Video solution



14. (b)

Given:

$$\frac{dy}{dx} + \frac{x}{1-x^2} y = x \cdot \sqrt{y}$$

$$\Rightarrow \frac{1}{\sqrt{y}} \cdot \frac{dy}{dx} + \frac{x}{1-x^2} \cdot \sqrt{y} = x \quad \dots(1)$$

Let  $\sqrt{y} = z$

$$\Rightarrow \frac{1}{2\sqrt{y}} \cdot \frac{dy}{dx} = \frac{dz}{dx} \Rightarrow \frac{1}{\sqrt{y}} \cdot \frac{dy}{dx} = 2 \cdot \frac{dz}{dx}$$

Equation (i) will become,

$$\Rightarrow 2 \cdot \frac{dz}{dx} + \frac{x}{1-x^2} z = x$$

$$\Rightarrow \frac{dz}{dx} + \frac{x}{2(1-x^2)}z = \frac{x}{2}$$

This is **Leibnitz linear equation** of form

$$\frac{dz}{dx} + P'z = Q' \text{ where } P' = \frac{x}{2(1-x^2)} \text{ and } Q' = \frac{x}{2}.$$

$$\therefore \text{Integrating factor, I.F.} = e^{\int P' dx} = e^{\int \frac{x}{2(1-x^2)} dx}$$

$$= e^{\frac{-1}{4} \int \frac{-2x}{1-x^2} dx} = e^{\frac{-1}{4} \cdot \ln(1-x^2)}$$

$$= e^{\ln(1-x^2)^{-1/4}} = (1-x^2)^{-1/4}$$

$$\therefore \text{Integrating factor} = (1-x^2)^{-1/4}$$



Scan for Video solution



### 15. (0.8774 to 0.8952)

$$\text{Given: } \frac{dy}{dx} = 2x - y, \quad y(0) = 1$$

$$\frac{dy}{dx} = 2x - y$$

$$\Rightarrow \frac{dy}{dx} + y = 2x$$

This is **Leibnitz linear equation** of form

$$\frac{dy}{dx} + P'y = Q' \text{ where } P' = 1 \text{ and } Q' = 2x.$$

Solution of above differentiable equation is given by,

$$y \cdot (\text{I.F.}) = \int Q'(\text{I.F.}) dx + c$$

$$\Rightarrow y \cdot (\text{I.F.}) = \int 2x(\text{I.F.}) dx + c \dots (i)$$

Here, Integrating factor:

$$\text{I.F.} = e^{\int P dx} = e^{\int 1 dx} = e^x$$

From equation (i),

$$y \cdot (e^x) = \int 2x(e^x) dx + c$$

$$\Rightarrow ye^x = 2 \int xe^x dx + c$$

$$\left[ \because \int uv = u \int v dx - \int u' \left( \int v dx \right) dx \right]$$

In  $\int xe^x dx$ , consider  $u = x$  &  $v = e^x$

$$\Rightarrow y \cdot e^x = 2 \left[ xe^x - \int 1 \cdot e^x \right] + c$$

$$\Rightarrow y \cdot e^x = 2xe^x - 2e^x + c$$

$$\Rightarrow y = 2x - 2 + ce^{-x} \dots (ii)$$

Given  $y(0) = 1$ ; from equation (ii)

$$1 = 2(0) - 2 + ce^0$$

$$\Rightarrow c = 3$$

Put value of c in equation (ii)

$$y = 2x - 2 + 3e^{-x}$$

At  $x = \ln 2$ ;

$$y = 2 \ln 2 - 2 + 3e^{-\ln 2}$$

$$\Rightarrow \boxed{y = 0.886}$$



Scan for Video solution



### 16. (c)

**Given:**

$$\frac{dy}{dx} + 7x^2y = 0; \text{ and } y(0) = \frac{3}{7}$$

**Find:**  $y(1) = ?$

$$\frac{dy}{dx} + 7x^2y = 0$$

$$\Rightarrow \frac{dy}{dx} = -7x^2y$$

$$\Rightarrow \frac{1}{y} dy = -7x^2 dx \text{ [Variable separable form]}$$

$$y(0) = \frac{3}{7} \text{ i.e. at } x = 0 \Rightarrow y = \frac{3}{7} \text{ and let at } x = 1; y = \alpha$$

$\alpha$

Integrating above differential equation from  $x = 0$  to

$$x = 1 \text{ and } y = \frac{3}{7} \text{ to } y = \alpha$$

$$\Rightarrow \int_{y=3/7}^{y=\alpha} \frac{1}{y} dy = \int_{x=0}^{x=1} -7x^2 dx$$

$$\Rightarrow \ln y \Big|_{\frac{3}{7}}^{\alpha} = \frac{-7x^3}{3} \Big|_0^1$$

$$\Rightarrow \ln \left( \frac{\alpha}{\left( \frac{3}{7} \right)} \right) = \frac{-7}{3}(1 - 0) \Rightarrow \alpha = \frac{3}{7} \cdot e^{-7/3}$$

$$\Rightarrow y(1) = \frac{3}{7} \cdot e^{-7/3}$$



Scan for Video solution



17. (b)

Given  $\frac{dy}{dx} + 4y = 5$ ;  $y(0) = 2.25$

$$\frac{dy}{dx} + 4y = 5$$

$$\Rightarrow \frac{dy}{dx} = 5 - 4y$$

$$\Rightarrow \frac{1}{5 - 4y} dy = dx \quad [\text{Variable separable form}]$$

On integrating,

$$\Rightarrow \int \frac{1}{5 - 4y} dy = \int dx$$

$$\Rightarrow \frac{\ln(5 - 4y)}{-4} = x + \ln C$$

$$\Rightarrow \ln(5 - 4y) = -4x + \ln C$$

$$\Rightarrow \ln \frac{(5 - 4y)}{C} = -4x$$

$$\Rightarrow 5 - 4y = C e^{-4x}$$

$$\Rightarrow 4y = 5 - C e^{-4x} \quad \dots(i)$$

At  $x = 0$ ;  $y = 2.25$ ; from equation (i),

$$4 \times 2.25 = 5 - C e^{-4 \times 0} \Rightarrow 9 = 5 - C \Rightarrow C = -4$$

Put value of C in equation (i),

$$4y = 5 - (-4) \times e^{-4x}$$

$$\Rightarrow y = \frac{5}{4} + e^{-4x}$$

$$\Rightarrow y = e^{-4x} + 1.25$$



Scan for Video solution



18. (c)

$$\frac{dy}{dx} (x \ln x) = y$$

$$\Rightarrow \frac{dy}{y} = \frac{dx}{x \ln x}$$

[Variable separable form]

$$\Rightarrow \ln y = \int \frac{dx}{x \ln x} + \ln K \quad [\text{Integrating both sides}]$$

$$\text{Let, } \ln x = t \Rightarrow \frac{1}{x} dx = dt$$

$$\Rightarrow \ln y = \int \frac{dt}{t} + \ln K$$

$$\Rightarrow \ln y = \ln t + \ln K$$

$$\Rightarrow y = Kt \quad [\text{Put } t = \ln x]$$

$$\Rightarrow y = K \ln x$$



Scan for Video solution



19. (c)

$$x \frac{dy}{dx} + y = 0$$

$$\Rightarrow \frac{dy}{y} = -\frac{dx}{x}$$

[Variable separable

form]

Integrating both sides, we get

$$\ln y = -\ln x + c$$

At  $x = 1$ ,  $y = 1$ 

$$\ln 1 = -\ln 1 + c \Rightarrow c = 0$$

∴ The solution is,

$$\ln y = -\ln x \Rightarrow \ln(yx) = 0$$

$$\Rightarrow yx = 1 \Rightarrow y = x^{-1}$$



Scan for Video solution



20. (c)

$$\text{Given, } y^3 \cdot \frac{dy}{dx} + x^3 = 0; \quad y(0) = 1$$

Find:  $y(-1) = ?$ 

$$y^3 \cdot \frac{dy}{dx} + x^3 = 0$$

$$\Rightarrow y^3 \cdot dy = -x^3 \cdot dx \quad [\text{Variable separable form}]$$

$y(0) = 1$  i.e. at  $x = 0; y = 1$  and let at  $x = -1; y = \alpha$ , then integrating above differential equation from  $x = 0$  to  $x = -1$  and  $y = 1$  to  $y = \alpha$ ,

$$\begin{aligned} \Rightarrow \int_{y=1}^{y=\alpha} y^3 dy &= \int_{x=0}^{x=-1} -x^3 dx \\ \Rightarrow \frac{y^4}{4} \Big|_1^\alpha &= \frac{-x^4}{4} \Big|_0^{-1} \\ \Rightarrow \alpha^4 - 1 &= -(1 - 0) \Rightarrow \alpha^4 = 0 \\ \Rightarrow \alpha &= 0 \\ \therefore y(-1) &= 0 \end{aligned}$$



Scan for Video solution



21. (a)

$$\text{Given: } \frac{dy}{dx} = \frac{x^2 + y^2}{2y} + \frac{y}{x}$$

And the curve passes through  $(1, 0)$ .

$$\frac{dy}{dx} = \frac{x^2 + y^2}{2y} + \frac{y}{x} \quad \dots(i)$$

$$\text{Let } y = vx \Rightarrow \frac{dy}{dx} = v + x \cdot \frac{dv}{dx}$$

From equation (i),

$$\Rightarrow v + x \cdot \frac{dv}{dx} = \frac{x^2 + v^2 x^2}{2vx} + v$$

$$\Rightarrow \frac{dv}{dx} = \frac{1 + v^2}{2v}$$

$$\Rightarrow \int \frac{2v}{1 + v^2} dv = \int dx$$

Let,  $1 + v^2 = t$ , then  $2vdv = dt$

$$\Rightarrow \int \frac{dt}{t} = \int dx$$

$$\Rightarrow \ln t = x + c \quad \left[ \because 1 + v^2 = t \right]$$

$$\Rightarrow \ln(1 + v^2) = x + c \quad \left[ \because y = vx \Rightarrow v = \frac{y}{x} \right]$$

$$\Rightarrow \ln \left( 1 + \frac{y^2}{x^2} \right) = x + c \quad \dots(ii)$$

At  $x = 1, y = 0$ ; from equation (ii),

$$\Rightarrow \ln \left( 1 + \frac{0}{1} \right) = 1 + c \Rightarrow c = -1$$

Put value of c in equation (ii),

$$\Rightarrow \boxed{\ln \left( 1 + \frac{y^2}{x^2} \right) = x - 1}$$



Scan for Video solution



22. (d)

$$\text{Given: DE is } \frac{dy}{dx} = (x + y - 1)^2 \quad \dots(1)$$

$$\text{Let } x + y - 1 = t \Rightarrow 1 + \frac{dy}{dx} = \frac{dt}{dx} \Rightarrow \frac{dy}{dx} = \frac{dt}{dx} - 1$$

Put in equation (1),

$$\Rightarrow \frac{dt}{dx} - 1 = t^2$$

$$\Rightarrow \frac{dt}{dx} = 1 + t^2$$

$$\Rightarrow \frac{1}{1 + t^2} dt = dx$$

On integrating both side

$$\int \frac{1}{1 + t^2} dt = \int dx$$


$$\Rightarrow \tan^{-1} t = x + c$$

$$\Rightarrow t = \tan(x + c) \quad [\text{Since } x + y - 1 = t]$$

$$\Rightarrow x + y - 1 = \tan(x + c)$$

$$\Rightarrow y = 1 - x + \tan(x + c)$$

$\therefore y = 1 - x + \tan(x + c)$ , where c is a constant.



Scan for Video solution



23. (d)

$$\frac{dQ}{dt} + Q = 1 \text{ and } Q = 0 \text{ at } t = 0$$

This is **Leibnitz linear equation** of form

$$\frac{dy}{dx} + P'y = Q' \text{ or } \frac{dQ}{dt} + P'Q = Q' \text{ where } P' = 1 \\ \text{and } Q' = 1$$

Solution of above differentiable equation is given by,

$$Q(\text{I.F.}) = \int Q'(\text{I.F.}) dt + c \\ \Rightarrow Q \cdot (\text{I.F.}) = \int 1 \times (\text{I.F.}) dt + c \quad \dots(i)$$

Here, Integrating factor;

$$\text{I.F.} = e^{\int P'dt} = e^{\int dt} = e^t$$

From equation (i),

$$Q e^t = \int 1 \cdot e^t dt + c \Rightarrow Q e^t = e^t + c$$

$$\Rightarrow Q = 1 + ce^{-t} \quad \dots(ii)$$

when  $t = 0, Q = 0$

$$0 = 1 + c e^{-0} \Rightarrow c = -1$$

Therefore, from equation (ii),

$$Q(t) = 1 - e^{-t}$$



Scan for Video solution



24. (a)

$$\text{Given: } (t^2 - 81) \frac{dy}{dt} + 5ty = \sin t ; \quad y(1) = 2\pi$$

$$(t^2 - 81) \frac{dy}{dt} + 5ty = \sin t$$

$$\Rightarrow \frac{dy}{dt} + \frac{5t}{t^2 - 81}y = \frac{\sin t}{t^2 - 81}$$

This is **Leibnitz linear equation** of form

$$\frac{dy}{dt} + P'y = Q' \text{ where } P' = \frac{5t}{t^2 - 81} \text{ and } Q' = \frac{\sin t}{t^2 - 81}.$$

Solution of above differentiable equation is given by,

$$y \cdot (\text{I.F.}) = \int Q'(\text{I.F.}) dt + c$$

$$\Rightarrow y \cdot (\text{I.F.}) = \int \frac{\sin t}{t^2 - 81} (\text{I.F.}) dt + c \quad \dots(i)$$

Here, Integrating factor:

$$\text{I.F.} = e^{\int P'dt} = e^{\int \frac{5t}{t^2 - 81} dt}$$

$$\Rightarrow \text{I.F.} = e^{\frac{5}{2} \int \frac{2t}{t^2 - 81} dt}$$

Let,  $t^2 - 81 = 0 \Rightarrow 2tdt = du$ , then

$$\Rightarrow \text{I.F.} = e^{\frac{5}{2} \int \frac{du}{u}} = e^{\frac{5}{2} \ln u} = u^{\frac{5}{2}} \quad \left[ \because u = t^2 - 81 \right]$$

$$\Rightarrow \text{I.F.} = (t^2 - 81)^{5/2}$$

From equation (i),

$$y(t^2 - 81)^{5/2} = \int \frac{\sin t}{t^2 - 81} \cdot (t^2 - 81)^{5/2} dt + c$$

$$\Rightarrow y = \frac{\int \sin t \cdot (t^2 - 81)^{3/2} dt}{(t^2 - 81)^{5/2}} + \frac{c}{(t^2 - 81)^{5/2}}$$

For  $t^2 - 81 = 0 \Rightarrow t = \pm 9$  the solution will not be unique

$\therefore$  option b, c and d are incorrect as in their interval t becomes 9 or -9. Hence option b i.e. (-10, 10), c i.e. (-10, 2) and d i.e. (0, 10) are eliminated and correct option is option a i.e. (-2, 2).

Scan for Video solution

25. (c)

$$\text{Given } \frac{dy}{dx} = \frac{1 + \cos 2y}{1 - \cos 2x} = \frac{1 + 2\cos^2 y - 1}{1 - (1 - 2\sin^2 x)}$$

$$\Rightarrow \frac{dy}{dx} = \frac{2\cos^2 y}{2\sin^2 x} \Rightarrow \sec^2 y dy = \csc^2 x dx$$

On Integrating,

$$\Rightarrow \int \sec^2 y dy = \int \csc^2 x dx$$

$$\Rightarrow \tan y = -\cot x + c$$

$$\Rightarrow \tan y + \cot x = c$$

Scan for Video solution

26. (c)

Given:  $\frac{dy}{dt} = -5y$ ; At  $t = 0 \Rightarrow y = 2$

And at  $t = 3 \Rightarrow y = ?$

$$\frac{dy}{dt} = -5y$$

$$\Rightarrow \frac{1}{y} dy = -5 dt \quad [\text{Variable separable form}]$$

at  $t = 0 \Rightarrow y = 2$  and let at  $t = 3 \Rightarrow y = \alpha$ , then integrating above differential equation from  $t = 0$  to  $t = 3$  and  $y = 2$  to  $y = \alpha$ ,

$$\Rightarrow \int_{y=2}^{y=\alpha} \frac{1}{y} dy = \int_{t=0}^{t=3} -5 dt$$

$$\Rightarrow \ln y \Big|_2^{\alpha} = -5 \times t \Big|_0^3$$

$$\Rightarrow \ln\left(\frac{\alpha}{2}\right) = -5 \times (3 - 0) \Rightarrow \ln\left(\frac{\alpha}{2}\right) = -15$$

$$\Rightarrow \alpha = 2 \cdot e^{-15}$$

$$\Rightarrow y(3) = 2 \cdot e^{-15}$$



Scan for Video solution



27. (c)

$$x(y dx + x dy) \cos \frac{y}{x} = y(x dy - y dx) \sin \frac{y}{x}$$

$$\Rightarrow \frac{y dx + x dy}{x dy - y dx} = \frac{y}{x} \tan \frac{y}{x} \dots\dots\dots (i)$$

Let  $y = vx$

then,  $dy = v dx + x dv$ ,

$\therefore$  from equation (i),

$$\frac{v x dx + x(v dx + x dv)}{x(v dx + x dv) - vx dx} = v \tan v$$

$$\Rightarrow \frac{2vx dx + x^2 dv}{x^2 dv} = v \tan v$$

$$\Rightarrow \frac{2v}{x} \frac{dx}{dv} + 1 = v \tan v \quad [\text{divide both side by } v]$$

$$\Rightarrow 2 \frac{dx}{x dv} + \frac{1}{v} = \tan v$$

$$2 \frac{dx}{x} = \left( \tan v - \frac{1}{v} \right) dv$$

Integrating both sides,

$$\Rightarrow 2 \log x = \log |\sec v| - \log v + \log c$$

$$\Rightarrow \log x^2 = \log \frac{c \sec v}{v} \Rightarrow x^2 = \frac{c \sec v}{v}$$

$$\Rightarrow x^2 = \frac{c \sec \frac{y}{x}}{y/x} \Rightarrow xy \frac{1}{\sec \frac{y}{x}} = c$$

$$\Rightarrow xy \cos \frac{y}{x} = c$$



Scan for Video solution



28. (c)

Given  $\frac{dx}{dt} = 10 - 0.2x$ ;  $x(0) = 1$

$$\frac{dx}{dt} = 10 - 0.2x \Rightarrow \frac{1}{10 - 0.2x} dx = dt$$

Integrating above equation from  $t=0$  to  $t$  and  $x=1$  to  $x$ ,

$$\Rightarrow \int_{x=1}^{x=x} \frac{1}{10 - 0.2x} dx = \int_{t=0}^{t=t} dt$$

$$\Rightarrow \left. \frac{\ln(10 - 0.2x)}{-0.2} \right|_1^x = t$$

$$\Rightarrow (\ln(10 - 0.2x))|_1^x = -0.2t$$

$$\Rightarrow \ln\left(\frac{10 - 0.2x}{9.8}\right) = -0.2t$$

$$\Rightarrow 10 - 0.2x = 9.8(e^{-0.2t})$$

$$\Rightarrow x = \frac{10 - 9.8(e^{-0.2t})}{0.2}$$

$$\Rightarrow x = 50 - 49e^{-0.2t}$$

$$\therefore x(t) = 50 - 49e^{-0.2t}$$



Scan for Video solution



## 29. (1.6 to 1.7)

**Given:**  $\frac{di}{dt} - 0.2i = 0$  and  $i(4) = 10$

$$\frac{di}{dt} - 0.2i = 0$$

$$\Rightarrow \frac{di}{i} = 0.2dt \quad (\text{Variable separable form})$$

Integrating both side,

$$\ln i = 0.2t + \ln C$$

$$\Rightarrow i = Ce^{0.2t} \quad \dots(i)$$

**Given**  $i(4) = 10$ ; from equation (i)

$$10 = Ce^{0.2 \times 4}$$

$$\Rightarrow C = 10e^{-0.8}$$

Put value of c in equation (i),

$$i = 10e^{-0.8} \cdot e^{0.2t}$$

At  $t = -5$ ;

$$i(-5) = 10e^{-0.8} \cdot e^{0.2(-5)} = 10e^{-1.8}$$

$$\Rightarrow \boxed{i(-5) = 1.652}$$



Scan for Video solution



## 30. (d)

$$\frac{dP}{dt} + k_2 P = k_2 L_0 e^{-k_1 t}$$

This is **Leibnitz linear equation** of form

$$\frac{dP}{dt} + P' y = Q' \text{ where } P' = k_2 \text{ and } Q' = k_2 L_0 e^{-k_1 t}$$

Integrating factor (I.F.),

$$\text{I.F.} = e^{\int P' dt} \Rightarrow \boxed{\text{I.F.} = e^{\int k_2 dt}}$$



Scan for Video solution



## 31. (b)

**Given**  $\frac{dy}{dx} = -2xy$  and  $y(0) = 2$  i.e. At  $x = 0 \Rightarrow y = 2$

$$\frac{dy}{dx} = -2xy$$

$$\Rightarrow \frac{1}{y} dy = -2x dx \quad [\text{Variable separable form}]$$

Since  $y(0) = 2$ , hence integrating above differential equation from  $x = 0$  to  $x$  and  $y = 2$  to  $y$ ,

$$\Rightarrow \int_{y=2}^y \frac{1}{y} dy = \int_{x=0}^x -2x dx$$

$$\Rightarrow \ln y \Big|_2^y = -2 \frac{x^2}{2} \Big|_0^x \Rightarrow \ln \left( \frac{y}{2} \right) = -x^2$$

$$\Rightarrow \boxed{y = 2 \cdot e^{-x^2}}$$

Scan for Video solution



## 32. (d)

$$\text{Given: } \frac{dy}{dx} = \cos(x + y) \quad \dots(i)$$

Let  $x + y = t$

$$\text{Then, } 1 + \frac{dy}{dx} = \frac{dt}{dx} \Rightarrow \frac{dy}{dx} = \frac{dt}{dx} - 1$$

$\therefore$  Given equation (i) will become,

$$\frac{dt}{dx} - 1 = \cos t$$

$$\Rightarrow \frac{dt}{dx} = 1 + \cos t$$

$$\Rightarrow \frac{dt}{dx} = 2 \cos^2 t / 2$$

$$\Rightarrow \frac{1}{\cos^2 t / 2} dt = 2 \cdot dx$$

$$\Rightarrow \sec^2 \frac{t}{2} dt = 2 \cdot dx$$

On integrating,

$$\Rightarrow \int \sec^2 \frac{t}{2} dt = \int 2 \cdot dx$$

$$\text{Let } \frac{t}{2} = V \Rightarrow \frac{dt}{2} = dV$$

$$\Rightarrow \int \sec^2 V \cdot (2) dV = 2 \cdot \int dx$$

$$\Rightarrow \tan V = x + c \quad \left[ \because \frac{t}{2} = V \right]$$

$$\Rightarrow \tan \frac{t}{2} = x + c \quad [\because t = x + y]$$

$$\Rightarrow \boxed{\tan\left(\frac{x+y}{2}\right) = x + c}$$



Scan for Video solution

**33. (d)**

**Given:**  $t \frac{dx}{dt} + x = t$  ;  $x(1) = 0.5$ ,

$$t \frac{dx}{dt} + x = t$$

$$\Rightarrow \frac{dx}{dt} + \frac{x}{t} = 1$$

This is Leibnitz linear equation of form

$$\frac{dx}{dt} + P'x = Q' \text{ where } P' = \frac{1}{t} \text{ and } Q' = 1.$$

Solution of above differentiable equation is given by,

$$x \cdot (I.F) = \int Q'(I.F) dt + c$$

$$\Rightarrow x \cdot (I.F) = \int 1 \times (I.F) dt + c \quad \dots(i)$$

Here, Integrating factor:

$$I.F = e^{\int P' dt} = e^{\int \frac{1}{t} dt} = e^{\log t} = t$$

From equation (i)

$$xt = \int t dt + c$$

$$\Rightarrow xt = \frac{t^2}{2} + c \quad \dots(ii)$$

Given, at  $t = 1$   $x = 0.5$ ; from equation (ii),

$$(0.5)(1) = \frac{1^2}{2} + c \Rightarrow c = 0$$

Put value of  $c$  in equation (ii),

$$\boxed{x = \frac{t}{2}}$$


Scan for Video solution



**34. (a)**

$$\frac{dy}{dx} = e^{-3x}$$

$$\Rightarrow dy = e^{-3x} dx$$

(Variable separable form)

Integrating both sides.

$$y = -\frac{1}{3}e^{-3x} + K$$

$$\Rightarrow \boxed{y = -\frac{1}{3}e^{-3x} + K}$$


Scan for Video solution



**35. (d)**

**Given:**  $\frac{dy}{dx} = (1 + y^2)x$

$$\Rightarrow \frac{1}{1 + y^2} dy = x \cdot dx \quad [\text{Variable Separation form}]$$

On integrating both sides

$$\Rightarrow \int \frac{1}{1 + y^2} dy = \int x dx + c$$

$$\Rightarrow \tan^{-1} y = \frac{x^2}{2} + c$$

$$\Rightarrow \boxed{y = \tan\left(\frac{x^2}{2} + c\right)}$$


Scan for Video solution



**36. (d)**

**Given,**  $\frac{dy}{dx} + \frac{y}{x} = x$  and  $y = 1$  at  $x = 1$

This is **Leibnitz linear equation** of form

$$\frac{dy}{dx} + P'y = Q' \text{ where } P' = \frac{1}{x} \text{ and } Q' = x.$$

Solution of above differentiable equation is given by,

$$y \cdot (I.F.) = \int Q' (I.F.) dx + c$$

$$y \cdot (I.F.) = \int x (I.F.) dx + c \quad \dots(i)$$

Here, Integrating factor (I.F.):

$$I.F. = e^{\int P' dx} = e^{\int \frac{1}{x} dx} = e^{\ln x}$$

$$\Rightarrow I.F. = x$$

From equation (i),

$$yx = \int x \cdot x dx$$

$$\Rightarrow yx = \frac{x^3}{3} + c \quad \dots(ii)$$

At  $x = 1, y = 1;$

$\therefore$  from equation (ii)

$$1 = \frac{1}{3} + c \Rightarrow c = \frac{2}{3}, \text{ then}$$

$$yx = \frac{x^3}{3} + \frac{2}{3} \Rightarrow y = \frac{x^2}{3} + \frac{2}{3}$$



Scan for Video solution



### 37. (d)

$$\frac{dy}{dx} + 2y = 0$$

$$\Rightarrow \frac{dy}{y} = -2 dx \quad [\text{Variable separable form}]$$

On integrating, we get

$$\ln y = -2x + c$$

$$\Rightarrow y = e^{-2x+c}$$

$$\Rightarrow y = e^{-2x} \cdot e^c \quad [\text{replace } e^c \text{ by } c_1]$$

$$\Rightarrow y = c_1 e^{-2x} \quad \dots(i)$$

At  $x = 1, y = 5$  so

$$5 = c_1 e^{-2}$$

$$\Rightarrow c_1 = 5e^2 = 36.95$$

Put value of  $c_1$  in equation (i)

$$\boxed{y = 36.95 e^{-2x}}$$

Scan for Video solution

### 38. (c)

**Given:**

$$\frac{dy}{dx} = ky \text{ and } y(0) = c$$

$$\frac{dy}{dx} = ky$$

$$\Rightarrow \frac{dy}{y} = kdx$$

On integrating,

$$\Rightarrow \int \frac{1}{y} dy = \int k dx$$

$$\Rightarrow \ln y = kx + C_1 \quad \dots(i)$$

At  $x = 0; y = c$ ; from equation (i)

$$\Rightarrow \ln c = C_1$$

Put value of  $C_1$  in equation (i),

$$\ln y = kx + \ln c$$

$$\Rightarrow \ln y - \ln c = kx$$

$$\Rightarrow \ln y/c = kx$$

$$\Rightarrow \boxed{y = c \cdot e^{kx}}$$

Scan for Video solution

### 39. (a)

$$\text{Given: } x \cdot \frac{dy}{dx} + y = x^4; \quad y(1) = \frac{6}{5}$$

$$x \cdot \frac{dy}{dx} + y = x^4$$

$$\Rightarrow \frac{dy}{dx} + \frac{1}{x} \cdot y = x^3$$

This is **Leibnitz linear equation** of form

$$\frac{dy}{dx} + P'y = Q' \text{ where } P' = \frac{1}{x} \text{ and } Q' = x^3.$$

Solution of above differentiable equation is given by,

$$y \cdot (I.F.) = \int Q' (I.F.) dx + c$$

$$y \cdot (\text{I.F.}) = \int x^3 (\text{I.F.}) dx + c \quad \dots(\text{i})$$

Here, Integrating factor:

$$\text{I.F.} = e^{\int P' dx} = e^{\int \frac{1}{x} dx} = e^{\ln x} = x$$

From equation (i),

$$\begin{aligned} yx &= \int x^3 \cdot x dx + c \\ \Rightarrow yx &= \frac{x^5}{5} + c \\ \Rightarrow y &= \frac{x^4}{5} + \frac{c}{x} \end{aligned} \quad \dots(\text{ii})$$

Given,  $y(1) = \frac{6}{5}$ ; from equation (ii),

$$\frac{6}{5} = \frac{1}{5} + c \Rightarrow c = 1$$

Put value of c in equation (ii),

$$y = \frac{x^4}{5} + \frac{1}{x}$$

Scan for Video solution



40. (a)

$$3y \frac{dy}{dx} + 2x = 0$$

$$\Rightarrow 3y dy + 2x dx = 0$$

On integrating, we get

$$\frac{3y^2}{2} + \frac{2x^2}{2} = c$$

$$\Rightarrow 2x^2 + 3y^2 = 2c \Rightarrow [2x^2 + 3y^2 = c]$$

This equation represents family of ellipses.



Scan for Video solution



41. (a)

$$(P) \frac{dy}{dx} = \frac{y}{x} \Rightarrow \int \frac{1}{y} dy = \int \frac{1}{x} dx \Rightarrow \ln y = \ln x + c$$

$$\Rightarrow y = cx \quad \rightarrow \text{Straight line (2)}$$

$$(Q) \frac{dy}{dx} = \frac{-y}{x}$$

$$\Rightarrow \int \frac{1}{y} dy = -\int \frac{1}{x} dx \Rightarrow \ln y = -\ln x + c$$

$\Rightarrow xy = c \rightarrow \text{Rectangular Hyperbola (3)}$

$$(R) \frac{dy}{dx} = \frac{x}{y}$$

$$\Rightarrow \int y dy = \int x dx \Rightarrow y^2 - x^2 = c \rightarrow \text{Hyperbolas (3)}$$

$$(S) \frac{dy}{dx} = \frac{-x}{y}$$

$$\Rightarrow \int y dy = -\int x dx$$

$$\Rightarrow y^2 - x^2 = c \rightarrow \text{Circles (1)}$$

Scan for Video solution



42. (d)

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$\Rightarrow x dx + y dy = 0 \quad [\text{Variable Separation form}]$$

On integrating,

$$\frac{x^2}{2} + \frac{y^2}{2} = c \quad \dots(\text{i})$$

$$\text{At } x = 1, y = \sqrt{3}$$

$$\frac{1^2}{2} + \frac{\sqrt{3}^2}{2} = c \Rightarrow c = 2$$

Put value of c in equation (i),

$$\frac{x^2}{2} + \frac{y^2}{2} = c \Rightarrow [x^2 + y^2 = 4]$$

Scan for Video solution



43. (b)

Given:

$$\frac{d}{dt} x(t) + 3x(t) = 0$$

$$\Rightarrow \frac{d}{dt}x(t) = -3x(t) \Rightarrow \frac{1}{x(t)}dx(t) = -3dt$$

$$\Rightarrow \int \frac{1}{x(t)}dx(t) = \int -3dt$$

$$\Rightarrow \ln x(t) = -3t + c$$

$$\Rightarrow x(t) = e^{-3t+c} \quad [\text{Let } e^c = C]$$

$$\Rightarrow x(t) = C.e^{-3t}$$

∴ The solution should be of the form

$$x(t) = C.e^{-3t}$$

For  $C = 2$ ,

$$x(t) = 2.e^{-3t}$$



Scan for Video solution



#### 44. (a)

Given,

$$\frac{d^3y}{dx^3} - 5.5 \frac{d^2y}{dx^2} + 9.5 \frac{dy}{dx} - 5y = 0$$

Complete solution is,

$$y = C_1e^{2.5x} + C_2e^{\alpha x} + C_3e^{\beta x}$$

Replace,  $\frac{d}{dx}$  by  $D$  and  $\frac{d^2}{dx^2}$  by  $D^2$  in differential

equation,

$$(D^3 - 5.5D^2 + 9.5D - 5)y = 0$$

This is homogenous linear differential equation with constant coefficients. Therefore,

$$y = C.F. + P.I.$$

[Since homogenous equation ∴ P.I. = 0]

$$\Rightarrow C_1e^{2.5x} + C_2e^{\alpha x} + C_3e^{\beta x} = C.F.$$

$$\Rightarrow C.F. = C_1e^{2.5x} + C_2e^{\alpha x} + C_3e^{\beta x}$$

We will get above expression of C.F. when roots of the Auxiliary equation are 2.5,  $\alpha$  and  $\beta$ .

Now, Auxiliary equation is:

$$m^3 - 5.5m^2 + 9.5m - 5 = 0$$

Now we can solve this problem by two methods.

#### Method I:

Auxiliary equation:

$$m^3 - 5.5m^2 + 9.5m - 5 = 0$$

Since one root of the A.E. should be 2.5, therefore we can express A.E. in the form of

$$(m - 2.5)(am^2 + bm + c) :$$

$$m^3 - 2.5m^2 - 3m^2 + 7.5m + 2m - 5 = 0$$

$$\Rightarrow m^2(m - 2.5) - 3m(m - 2.5) + 2(m - 2.5) = 0$$

$$\Rightarrow (m - 2.5)(m^2 - 3m + 2) = 0$$

$$\Rightarrow (m - 2.5)(m^2 - m - 2m + 2) = 0$$

$$\Rightarrow (m - 2.5)[m(m - 1) - 2(m - 1)] = 0$$

$$\Rightarrow (m - 2.5)(m - 1)(m - 2) = 0$$

$$\Rightarrow m = 2.5, 1, 2$$

Since roots of A.E. are 2.5,  $\alpha$  and  $\beta$  therefore,

$$\alpha = 1 \text{ and } \beta = 2$$

#### Method II: (Check by options)

Auxiliary equation:

$$m^3 - 5.5m^2 + 9.5m - 5 = 0 \quad \dots(i)$$

Since  $\alpha$  and  $\beta$  are roots of A.E. (i), therefore correct option will satisfy equation (i)

#### Checking for Option A:

For  $m=1$

$$L.H.S = 1^3 - 5.5(1)^2 + 9.5(1) - 5 = 0 \Rightarrow L.H.S =$$

R.H.S

Hence  $m=1$  is root of A.E.

For  $m=2$

$$L.H.S = 2^3 - 5.5(2)^2 + 9.5(2) - 5 = 0 \Rightarrow L.H.S =$$

R.H.S

Hence  $m=2$  is also root of A.E.

Hence roots of A.E. other than 2.5 are 1 and 2, hence

$$\alpha = 1 \text{ and } \beta = 2$$



Scan for Video solution



#### 45. (a)

Given: DE is,  $\frac{d^2y}{dx^2} + 2 \cdot \frac{dy}{dx} - 5y = 0$

This is homogenous linear differential equation with constant coefficients.

Replace  $\frac{d}{dx}$  by D and  $\frac{d^2}{dx^2}$  by  $D^2$

$$(D^2 + 2D - 5)y = 0$$

Auxiliary equation is,

$$m^2 + 2m - 5 = 0$$

$$\Rightarrow (m+1)^2 - 6 = 0$$

$$\Rightarrow m = -1 \pm \sqrt{6}$$

$$\therefore m_1 = -1 + \sqrt{6}, m_2 = -1 - \sqrt{6}$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$C.F. = C_1 e^{(-1+\sqrt{6})x} + C_2 e^{(-1-\sqrt{6})x}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:

$$y = C.F. + P.I.$$

$$\Rightarrow y = C_1 e^{(-1+\sqrt{6})x} + C_2 e^{(-1-\sqrt{6})x}$$

$$\Rightarrow y = K_1 e^{(-1+\sqrt{6})x} + K_2 e^{(-1-\sqrt{6})x}$$



Scan for Video solution



46. (a)

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0; y(0) = 1 \text{ and } y(1)$$

This is homogenous linear differential equation with constant coefficients. Replace,

$\frac{d}{dx}$  by D and  $\frac{d^2}{dx^2}$  by  $D^2$

$$(D^2 + 2D + 1)y = 0$$

Auxiliary equation is:

$$m^2 + 2m + 1 = 0$$

$$\Rightarrow (m+1)^2 = 0$$

$$\Rightarrow m = -1, -1$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$C.F. = (C_1 + C_2 x)e^{-x}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$\Rightarrow y = (C_1 + C_2 x)e^{-x} \quad \dots(i)$$

Given,  $y(0) = 1$ ; from equation (i)

$$(C_1 + C_2 \times 0)e^{-0} \Rightarrow C_1 = 1$$

Given,  $y(1) = 3$ ; from equation (i)

$$3 = (C_1 + C_2 \times 1)e^{-1}$$

$$\Rightarrow C_1 + C_2 = 3e \quad [\text{Put } C_1 = 1]$$

$$\Rightarrow 1 + C_2 = 3e$$

$$\Rightarrow C_2 = 3e - 1$$

Put value of  $C_1$  and  $C_2$  in equation (i),

$$y = [1 + (3e - 1)x]e^{-x}$$

$$\Rightarrow y = e^{-x} + (3e - 1)xe^{-x}$$



Scan for Video solution



47. (d)

$$6\frac{d^2y}{dx^2} + \frac{dy}{dx} - y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace,

$\frac{d}{dx}$  by D and  $\frac{d^2}{dx^2}$  by  $D^2$

$$(6D^2 + D - 1)y = 0$$

Its auxiliary equation is:

$$6m^2 + m - 1 = 0$$

$$\Rightarrow 6m^2 + 3m - 2m - 1 = 0$$

$$\Rightarrow 3m(2m + 1) - (2m + 1) = 0$$

$$\Rightarrow (3m - 1)(2m + 1) = 0$$

$$\Rightarrow m = \frac{1}{3}, -\frac{1}{2}$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$C.F. = C_1 e^{\frac{x}{3}} + C_2 e^{-\frac{x}{2}}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$\Rightarrow y = C_1 e^{\frac{x}{3}} + C_2 e^{-\frac{x}{2}}$$



Scan for Video solution



$$x = -10e^{2t} + 10e^{3t}$$



Scan for Video solution



48. (a)

$$x(0)=0 \text{ and } \frac{dx}{dt}(0)=10 \text{ and}$$

$$\frac{d^2x}{dt^2} - 5 \frac{dx}{dt} + 6x = 10$$

This is homogenous linear differential equation with constant coefficients. Replace,

$$\frac{d}{dt} \text{ by } D \text{ and } \frac{d^2}{dt^2} \text{ by } D^2$$

$$(D^2 - 5D + 6)x = 0$$

Its auxiliary equation is:

$$m^2 - 5m + 6 = 0$$

$$\Rightarrow (m-3)(m-2) = 0$$

$$\Rightarrow m = 2, 3$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$C.F. = C_1 e^{2t} + C_2 e^{3t}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $x = C.F. + P.I.$

$$\Rightarrow x = C_1 e^{2t} + C_2 e^{3t} \quad \dots(i)$$

Differentiating equation (i) with respect to t,

$$\frac{dx}{dt} = 2C_1 e^{2t} + 3C_2 e^{3t} \quad \dots(ii)$$

Given,  $x(0) = 0$ ; from equation (i)

$$0 = C_1 e^{2 \times 0} + C_2 e^{3 \times 0}$$

$$\Rightarrow C_1 + C_2 = 0 \Rightarrow C_1 = -C_2 \quad \dots(iii)$$

Given,  $\frac{dx}{dt}(0) = 10$ ; from equation (ii)

$$10 = 2C_1 e^{2 \times 0} + 3C_2 e^{3 \times 0}$$

$$\Rightarrow 2C_1 + 3C_2 = 10 \quad \dots(iv)$$

Put  $C_1 = -C_2$ ;

$$2(-C_2) + 3C_2 = 10 \Rightarrow C_2 = 10$$

$$\therefore C_1 = -C_2 \Rightarrow C_1 = 10$$

Put value of  $C_1$  and  $C_2$  in equation (i),

49. (1.45 to 1.48)

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 0 ; y(0) = 0 \text{ and } \frac{dy}{dx}(0) = 1$$

Find:  $y(1) = ?$

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$(D^2 + D - 6)y = 0$$

The Auxiliary equation is

$$m^2 + m - 6 = 0 \Rightarrow m^2 + 3m - 2m - 6 = 0$$

$$\Rightarrow m(m+3) - 2(m+3) = 0$$

$$\Rightarrow (m+3)(m-2) = 0$$

$$\Rightarrow m = -3, 2$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$C.F. = C_1 e^{-3x} + C_2 e^{2x}$$

Since, this is homogenous equation, therefore

$$P.I. = 0$$

Now, Complete solution:

$$y = C.F. + P.I.$$

$$\Rightarrow y = C_1 e^{-3x} + C_2 e^{2x} \quad \dots(i)$$

Given,  $y(0) = 0$  i.e. At  $x = 0$ ,  $y = 0$ ; from equation (i),

$$0 = C_1 \cdot (1) + C_2$$

$$\Rightarrow C_2 = -C_1 \quad \dots(ii)$$

Put  $C_2 = -C_1$  in equation (i)

$$y = C_1(e^{-3x} - e^{2x}) \quad \dots(iii)$$

Differentiating above equation with respect to x,

$$\frac{dy}{dx} = C_1(-3 \cdot e^{-3x} - 2 \cdot e^{2x}) \quad \dots(iv)$$

Given  $\frac{dy}{dx}(0) = 1$  i.e. at  $x = 0 \Rightarrow \frac{dy}{dx} = 1$ ; from equation (iv),

$$1 = C_1 [-3(1) - 2(1)] \Rightarrow C_1 = -1/5$$

Put value of  $C_1$  in equation (iii)

$$y = \frac{-1}{5} (e^{-3x} - e^{2x})$$

At  $x = 1$ ,

$$y(1) = y = \frac{-1}{5} (e^{-3 \times 1} - e^{2 \times 1})$$

$$\Rightarrow y(1) = 1.47$$



Scan for Video solution



### 50. (0.36 to 0.37)

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0 \text{ and } y(0) = 1, \frac{dy}{dx}(x=0) = -1$$

**Find:**  $y(1) = ?$

This is homogenous linear differential equation with constant coefficients.

Replace,  $\frac{d}{dx}$  by  $D$  and  $\frac{d^2}{dx^2}$  by  $D^2$

$$(D^2 + 2D + 1)y = 0$$

Its auxiliary equation is:

$$m^2 + 2m + 1 = 0 \Rightarrow (m + 1)^2 = 0 \Rightarrow m = -1, -1$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$C.F. = (C_1 + C_2 x)e^{-x}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$\Rightarrow y = (C_1 + C_2 x)e^{-x} \quad \dots(i)$$

Differentiating with respect to  $x$ ,

$$\frac{dy}{dx} = (C_1 + C_2 x)(-e^{-x}) + C_2 e^{-x} \quad \dots(ii)$$

Now,  $y(0) = 1$  i.e. at  $x = 0$ ,  $y = 1$  then from equation (i),

$$1 = (C_1 + C_2 \times 0)e^{-0} \Rightarrow C_1 = 1$$

& at  $x = 0$ ,  $\frac{dy}{dx} = -1$  then from equation (ii),

$$-1 = (C_1 + C_2 \times 0)(-e^0) + C_2 e^0$$

$$\Rightarrow -1 = -C_1 + C_2 \Rightarrow -1 = -1 + C_2$$

$$\Rightarrow C_2 = 0$$

Put value of  $C_1$  and  $C_2$  in equation (i),

$$y = e^{-x}$$

$\therefore$  At  $x = 1$ ;

$$y(1) = e^{-1} \Rightarrow y(1) = 0.368$$

Scan for Video solution



### 51. (-0.23 to -0.19)

$$\text{Given: DE is } \frac{d^2y}{dt^2} = -\frac{dy}{dt} - \frac{5}{4}y$$

$$\text{And } y(0) = 1 \text{ and } \left. \frac{dy}{dt} \right|_{t=0} = 0.$$

$$\frac{d^2y}{dt^2} = -\frac{dy}{dt} - \frac{5}{4}y$$

$$\Rightarrow \frac{d^2y}{dt^2} + \frac{dy}{dt} + \frac{5}{4}y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$\frac{d}{dt}$  by  $D$  and  $\frac{d^2}{dt^2}$  by  $D^2$ ;

$$\Rightarrow (D^2 + D + \frac{5}{4})y = 0$$

The auxiliary equation:

$$m^2 + m + \frac{5}{4} = 0$$

$$\Rightarrow 4m^2 + 4m + 5 = 0$$

$$\Rightarrow m = \frac{-4 \pm \sqrt{16 - 4(4)(5)}}{8}$$

$$\Rightarrow m = \frac{-1}{2} \pm i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = -\frac{1}{2}$  and  $\beta = 1$ , therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = e^{\alpha t} (c_1 \cos \beta t + c_2 \sin \beta t)$$

$$\Rightarrow \text{C.F.} = e^{-t/2} \{c_1 \cos t + c_2 \sin t\}$$

Since, this is homogenous equation, therefore

$$\text{Particular integral (P.I.)} = 0$$

Now, Complete solution:

$$y(t) = \text{C.F.} + \text{P.I.}$$

$$\therefore y(t) = e^{-t/2} \{c_1 \cos t + c_2 \sin t\} \quad \dots(\text{i})$$

Now,

$$\frac{dy}{dt} = e^{-t/2} \left[ \{-c_1 \sin t + c_2 \cos t\} - \frac{1}{2} \{c_1 \cos t + c_2 \sin t\} \right] \quad \dots(\text{ii})$$

At  $t = 0$ ;  $y = 1$ ; from equation (i),

$$\Rightarrow 1 = 1 \{c_1 \cos 0 + c_2 \sin 0\} \Rightarrow c_1 = 1$$

$$\text{At } t = 0; \frac{dy}{dt} = 0; \text{ from equation (ii),}$$

$$\Rightarrow 0 = 1 [c_2 - \frac{1}{2} c_1]$$

$$\Rightarrow c_2 = \frac{c_1}{2} \quad [\text{Since } c_1 = 1]$$

$$\Rightarrow c_2 = \frac{1}{2}$$

Put value of  $c_1$  and  $c_2$  in equation (i),

$$y(t) = e^{-t/2} \left( \cos t + \frac{1}{2} \sin t \right)$$

At  $t = \pi$

$$y(t) = e^{-\pi/2} \left( \cos \pi + \frac{1}{2} \sin \pi \right)$$

$$\Rightarrow y(\pi) = e^{-\pi/2} (-1 + 0)$$

$$\Rightarrow \boxed{y(\pi) = -0.2079}$$



Scan for Video solution



52. (a)

$$y'' - 4y' + 3y = 2t - 3t^2 \text{ and find: P.I. = ?}$$

This is non homogenous linear differential equation with constant coefficients and can be written as:

$$(D^2 - 4D + 3)y = 2t - 3t^2$$

$$\text{P.I.} = \frac{1}{f(D)} (2t - 3t^2) = \frac{1}{D^2 - 4D + 3} (2t - 3t^2)$$

$$= \frac{1}{D^2 - 3D - D + 3} (2t - 3t^2)$$

$$= \frac{1}{-D(3-D) + 1(3-D)} (2t - 3t^2)$$

$$= \frac{1}{(1-D)(3-D)} (2t - 3t^2)$$

$$= \left[ \frac{1/2}{1-D} - \frac{1/2}{3-D} \right] (2t - 3t^2)$$

$$= \frac{1}{2}(1-D)^{-1} (2t - 3t^2) - \frac{1}{6} \left( 1 - \frac{D}{3} \right)^{-1} (2t + 3t^2)$$

$$\left[ \because (1-D)^{-1} = 1 + D + D^2 + D^3 + \dots \right]$$

$$= \left[ \frac{1}{2} (1 + D + D^2 + \dots) (2t - 3t^2) \right]$$

$$= \left[ -\frac{1}{6} \left( 1 + \frac{D}{3} + \frac{D^2}{9} + \dots \right) (2t - 3t^2) \right]$$


$$= \frac{1}{2} [(2t - 3t^2) + D(2t - 3t^2) + D^2(2t - 3t^2)]$$

$$= -\frac{1}{6} \left[ (2t - 3t^2) + \frac{D}{3} (2t - 3t^2) + \frac{D^2}{9} (2t - 3t^2) \right]$$

$$= \frac{1}{2} [2t - 3t^2 + 2 - 6t - 6] - \frac{1}{6} \left[ 2t - 3t^2 + \frac{2}{3} - 2t - \frac{2}{3} \right]$$

$$= \frac{1}{2} [-3t^2 - 4t - 4] - \frac{1}{6} [-3t^2] = -\frac{3t^2}{2} - 2t - 2 + \frac{t^2}{2}$$

$$\boxed{\text{P.I.} = -2 - 2t - t^2}$$



Scan for Video solution



53. (a)

$$\text{Given } \frac{d^2y}{dx^2} + 16y = 0; \quad \left. \frac{dy}{dx} \right|_{x=0} = 1 \text{ & } \left. \frac{dy}{dx} \right|_{x=\frac{\pi}{2}} = -1$$

$$\frac{d^2y}{dx^2} + 16y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$(D^2 + 16)y = 0$$

The Auxiliary equation is

$$m^2 + 16 = 0 \Rightarrow m = \pm 4i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = 4$ , therefore complimentary function (C.F.) is given by,

$$C.F. = e^{\alpha x} (c_1 \cos \beta x + c_2 \sin \beta x)$$

$$\Rightarrow C.F. = e^{0 \times x} (c_1 \cos(4 \times x) + c_2 \sin(4 \times x))$$

$$\Rightarrow C.F. = c_1 \cos 4x + c_2 \sin 4x$$

Since, this is homogenous equation, therefore

Particular integral (P.I.) = 0

Now, Complete solution:

$$y = C.F. + P.I.$$

$$\Rightarrow y = c_1 \cos 4x + c_2 \sin 4x \quad \dots(i)$$

Differentiating above equation with respect to x,

$$\frac{dy}{dx} = -4c_1 \sin 4x + 4c_2 \cos 4x \quad \dots(ii)$$

At  $x = 0$ ;  $\frac{dy}{dx} = 1$ , hence from equation (ii),

$$1 = 0 + 4c_2 \Rightarrow c_2 = \frac{1}{4}$$

At  $x = \frac{\pi}{2}$ ;  $\frac{dy}{dx} = -1$ , hence from equation (ii),

$$-1 + 0 + 4c_2 \Rightarrow c_2 = -\frac{1}{4}$$

Since  $c_2$  is different for both the boundary conditions.  $\therefore$  The differential equation has no solution that satisfy the given boundary conditions.



Scan for Video solution



#### 54. (93 to 95)

$$3 \cdot y''(x) + 27 \cdot y(x) = 0; y(0) = 0 \text{ and } y'(0) = 2000$$

**Find:**  $y(1) = ?$

$$3 \cdot y''(x) + 27 \cdot y(x) = 0$$

$$\Rightarrow y''(x) + 9 \cdot y(x) = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$(D^2 + 9)y(x) = 0$$

The Auxiliary equation is:

$$m^2 + 9 = 0 \Rightarrow m = \pm 3i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = 3$ , therefore complimentary function (C.F.) is given by,

$$C.F. = e^{\alpha x} (c_1 \cos \beta x + c_2 \sin \beta x)$$

$$\Rightarrow C.F. = e^{0 \times x} (c_1 \cos(3x) + c_2 \sin(3x))$$

$$\Rightarrow C.F. = c_1 \cos 3x + c_2 \sin 3x$$

Since, this is homogenous equation, therefore

Particular integral (P.I.) = 0

Now, Complete solution:

$$y(x) = C.F. + P.I.$$

$$\Rightarrow y(x) = c_1 \cos 3x + c_2 \sin 3x \quad \dots(i)$$

Given  $y(0) = 0$  i.e. at  $x = 0$ ,  $y = 0$ ; from equation (i),

$$0 = c_1 \cos 0 + c_2 \sin 0 \Rightarrow c_1 = 0$$

Put value of  $c_1$  in equation (i),

$$y(x) = c_2 \sin 3x \quad \dots(ii)$$

Differentiating above equation with respect to x,

$$y'(x) = 3c_2 \cos 3x \quad \dots(iii)$$

At  $x = 0$ ;  $y'(x) = 2000$ , from equation (iii),

$$2000 = 3c_2 \Rightarrow c_2 = \frac{2000}{3}$$

Put value of  $c_2$  in equation (ii),

$$y(x) = \frac{2000}{3} \cdot \sin 3x$$

At  $x = 1$ ,

$$y(1) = \frac{2000}{3} \cdot \sin(3 \times 1)$$

$$\Rightarrow \boxed{y(1) = 94.08}$$



Scan for Video solution



55. (-1.05 to -0.95)

$$y'' + 9y = 0, y(0) = 0, y(\pi/2) = \sqrt{2} \text{ and } y(\pi/4) = ?$$

$$y'' + 9y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2;$$

$$(D^2 + 9)y = 0$$

The Auxiliary equation is:

$$m^2 + 9 = 0 \Rightarrow m = \pm 3i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = 3$ , therefore complimentary function (C.F.) is given by,

$$C.F. = e^{\alpha x} (c_1 \cos \beta x + c_2 \sin \beta x)$$

$$\Rightarrow C.F. = e^{0 \times x} (c_1 \cos(3x) + c_2 \sin(3x))$$

$$\Rightarrow C.F. = c_1 \cos 3x + c_2 \sin 3x$$

Since, this is homogenous equation, therefore

$$\text{Particular integral (P.I.)} = 0$$

Now, Complete solution:

$$y = C.F. + P.I.$$

$$\Rightarrow y = c_1 \cos 3x + c_2 \sin 3x \quad \dots(i)$$

At  $x = 0; y = 0$ ; from equation (i),

$$0 = c_1(1) + c_2(0) \Rightarrow c_1 = 0$$

At  $x = \frac{\pi}{2}, y = \sqrt{2}$ ; and  $c_1 = 0$ ,  $\therefore$  from equation (i),

$$\sqrt{2} = 0 \times \cos\left(\frac{3\pi}{2}\right) + c_2 \sin\left(\frac{3\pi}{2}\right)$$

$$\Rightarrow \sqrt{2} = 0 + c_2 \times (-1)$$

$$\Rightarrow c_2 = -\sqrt{2}$$

Put value of  $c_1$  and  $c_2$  in equation (i),

$$y(x) = -\sqrt{2} \cdot \sin 3x$$

$$\text{At } x = \frac{\pi}{4};$$

$$y\left(\frac{\pi}{4}\right) = -\sqrt{2} \cdot \sin\left(\frac{3\pi}{4}\right) = -\sqrt{2} \cdot \left(\frac{1}{\sqrt{2}}\right)$$

$$\Rightarrow y\left(\frac{\pi}{4}\right) = -1$$



Scan for Video solution



56. (a)

Complimentary function (C.F.) = ?,

Particular Integral (P.I.) = ?

$$\frac{d^4 y}{dx^4} + 3 \frac{d^2 y}{dx^2} = 108x^2$$

This is non homogenous linear differential equation with constant coefficients.

$$\text{Replace } \frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$(D^4 + 3D^2)y = 108x^2$$

Its auxiliary equation is:

$$m^4 + 3m^2 = 0 \Rightarrow m^2(m^2 + 3) = 0$$

$$\Rightarrow m = 0, 0, \pm \sqrt{3}i$$

Here two roots 0 and 0 are real and repeated and two roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = \sqrt{3}$ , C.F. is given by,

$$C.F. = (C_1 + C_2x)e^{0x} + e^{0x}$$

$$(C_3 \sin \sqrt{3}x + C_4 \cos \sqrt{3}x)$$

$$\Rightarrow C.F. = C_1 + C_2x + C_3 \sin \sqrt{3}x + C_4 \cos \sqrt{3}x$$

Since, this is non-homogenous equation, therefore

$$P.I. = \frac{1}{f(D)} \cdot 108x^2 = 108 \left( \frac{1}{D^4 + 3D^2} x^2 \right)$$

$$P.I. = 108 \left( \frac{1}{3D^2 \left( 1 + \frac{D^2}{3} \right)} \right) x^2 = \frac{36}{D^2} \left( 1 + \frac{D^2}{3} \right)^{-1} (x^2)$$

$$P.I. = \frac{36}{D^2} \left[ 1 - \frac{D^2}{3} + \left( \frac{D^2}{3} \right)^2 - \dots \right] (x^2)$$

$$P.I. = \frac{36}{D^2} \left[ x^2 - \frac{2}{3} + 0 \right] = \frac{36}{D^2} \left[ x^2 - \frac{2}{3} \right]$$

$$\text{P.I.} = 36 \int \left( x^2 - \frac{2}{3} \right) dx = 36 \int \left( \frac{x^3}{3} - \frac{2}{3}x \right) dx$$

$$\text{P.I.} = 36 \left[ \frac{x^4}{12} - \frac{2}{3} \times \frac{x^2}{2} \right]$$

$$\boxed{\text{P.I.} = 3x^4 - 12x^2}$$



Scan for Video solution

**57. (a)**

$$\frac{d^2y}{dx^2} + 12 \cdot \frac{dy}{dx} + 36y = 0 ; y(0) = 3 \quad \left. \frac{dy}{dx} \right|_{x=0} = -36$$

$$\frac{d^2y}{dx^2} + 12 \cdot \frac{dy}{dx} + 36y = 0$$

This is homogenous linear differential equation with constant coefficients.

Replace  $\frac{d}{dx}$  by D and  $\frac{d^2}{dx^2}$  by  $D^2$

$$(D^2 + 12D + 36)y = 0$$

The Auxiliary equation is,

$$m^2 + 12m + 36 = 0$$

$$\Rightarrow (m+6)^2 = 0$$

$$\Rightarrow m = -6, -6$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = (c_1 x + c_2) \cdot e^{-6x}$$

Since, this is homogenous equation, therefore

$$\text{P.I.} = 0$$

Now, Complete solution:

$$y = \text{C.F.} + \text{P.I.}$$

$$\Rightarrow y = (c_1 x + c_2) \cdot e^{-6x} \quad \dots(i)$$

Differentiating above equation with respect to x,

$$\frac{dy}{dx} = -6(c_1 x + c_2) \cdot e^{-6x} + e^{-6x} c_1 \dots(ii)$$

Given,  $y(0) = 3$ ;  $\therefore$  from equation (i),

$$\Rightarrow 3 = c_2 \cdot e^{-6(0)} \Rightarrow c_2 = 3$$

$$\left. \frac{dy}{dx} \right|_{x=0} = -36 \text{ and } c_2 = 3, \text{ from equation (ii)},$$

$$\Rightarrow -36 = -6(0 + 3) \cdot e^0 + e^0 c_1$$

$$\Rightarrow -36 = -18 + c_1$$

$$\Rightarrow c_1 = -18$$

Put value of  $c_1$  and  $c_2$  in equation (i),

$$y = (-18x + 3) \cdot e^{-6x}$$

$$\therefore \text{The solution is, } \boxed{y = (3 - 18x)e^{-6x}}.$$



Scan for Video solution

**58. (b)**

$$\frac{d^2y}{dt^2} + 2 \frac{dy}{dt} + y = 0 ; y(0) = 1 \text{ and } y(1) = 3e^{-1}$$

$$\text{Find: } y(2) = ?$$

$$\frac{d^2y}{dt^2} + 2 \frac{dy}{dt} + y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$\frac{d}{dt}$  by D and  $\frac{d^2}{dt^2}$  by  $D^2$ ;

$$(D^2 + 2D + 1)y = 0$$

The Auxiliary equation is:

$$(m^2 + 2m + 1) = 0$$

$$\Rightarrow (m+1)^2 = 0$$

$$\Rightarrow m = -1, -1$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = (C_1 + C_2 t) e^{-t}$$

Since, this is homogenous equation, therefore

$$\text{Particular integral (P.I.)} = 0$$

Now, Complete solution:

$$y = \text{C.F.} + \text{P.I.}$$

$$\Rightarrow y = (C_1 + C_2 t) e^{-t} \quad \dots(i)$$

At  $t = 0$ ;  $y = 1$ ; hence from equation (i),

$$1 = (C_1 + 0)e^0 \Rightarrow C_1 = 1$$

At  $t = 1$ ;  $y = 3e^{-1}$  and  $C_1 = 1$ ; hence from equation (i),

$$3e^{-1} = (1 + C_2 \times 1)e^{-1}$$

$$\Rightarrow 1 + C_2 = 3$$

$$\Rightarrow C_2 = 2$$

Put value of  $C_1$  and  $C_2$  in equation (i),  
 $\therefore y = (1 + 2t)e^{-t}$

At  $t = 2$ ;

$$y(2) = (1 + 2 \times 2)e^{-2}$$

$$\boxed{y(2) = 5e^{-2}}$$



Scan for Video solution



### 59. (7.0 to 7.5)

$$\frac{d^2y}{dx^2} - 4 \frac{dy}{dx} + 4y = 0; y(0) = 0 \text{ and } \left. \frac{dy}{dx} \right|_{x=0} = 1$$

Find:  $y(1) = ?$

$$\frac{d^2y}{dx^2} - 4 \frac{dy}{dx} + 4y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$(D^2 - 4D - 4)y = 0$$

The Auxiliary equation is

$$m^2 - 4m + 4 = 0$$

$$\Rightarrow (m - 2)^2 = 0$$

$$\Rightarrow m = 2, 2$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$C.F. = (C_1 + C_2x)e^{2x}$$

Since, this is homogenous equation, therefore

$$P.I. = 0$$

Now, Complete solution:

$$y = C.F. + P.I.$$

$$\Rightarrow y = (C_1 + C_2x)e^{2x} \dots (i)$$

At  $x = 0$ ;  $y = 0$ ; hence from equation (i),

$$0 = (C_1 + C_2 \times 0)e^0$$

$$\Rightarrow C_1 = 0$$

Put value of  $C_1$  in equation (i),

$$y = (C_2x)e^{2x} \dots (ii)$$

Differentiating above equation with respect to x,

$$\frac{dy}{dx} = 2C_2xe^{2x} + C_2e^{2x} \dots (iii)$$

At  $x = 0$ ;  $\frac{dy}{dx} = 1$ ; hence from equation (ii),

$$1 = 2C_2(0)e^0 + C_2e^0$$

$$\Rightarrow C_2 = 1$$

Put value of  $C_2$  in equation (ii),

$$y = xe^{2x}$$

At  $x = 1$ ,

$$y = 1 \times e^{2 \times 1} = e^2$$

$$\Rightarrow \boxed{y = 7.389}$$



Scan for Video solution



### 60. (a)

$$y''(t) + 2y'(t) + y(t) = 0$$

Auxiliary equation is  $m^2 + 2m + 1 = 0$

$$(m + 1)^2 = 0$$

$$m = -1, -1$$

$$\therefore y(t) = \text{complementary function} = (C_1 + C_2t)e^{-t}$$

At  $t = 0$ ;  $y(t) = 0$

$$0 = (C_1 + C_2 \times 0)e^0$$

$$C_1 = 0$$

$$y(t) = C_2te^{-t}$$

$$y'(t) = C_2e^{-t} - C_2te^{-t}$$

At  $t = 0$ ;  $y'(0) = 1$

$$1 = C_2e^0 - C_2 \times 0 \times e^0$$

$$C_2 = 1$$

$$\therefore y(t) = te^{-t}$$

$$y(t) = te^{-t}u(t)$$

$$\{ \because t > 0; u(t) = 1 \}$$



Scan for Video solution



### 61. (0.83 to 0.88)

Given:

$$\frac{d^2x(t)}{dt^2} + \frac{dx(t)}{dt} + 2x(t) = 0;$$

$$x(0) = 20 \text{ and } x(1) = 10/e$$

$$\frac{d^2x(t)}{dt^2} + \frac{dx(t)}{dt} + 2x(t) = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dt} \text{ by } D \text{ and } \frac{d^2}{dt^2} \text{ by } D^2;$$

$$\Rightarrow (D^2 + 3D + 2)x(t) = 0$$

Auxiliary equation is

$$m^2 + 3m + 2 = 0$$

$$\Rightarrow (m+1)(m+2) = 0$$

$$\Rightarrow m = -1, m = -2$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = C_1 e^{-t} + C_2 e^{-2t}$$

Since, this is homogenous equation, therefore

$$\text{P.I.} = 0$$

Now, Complete solution:

$$x(t) = \text{C.F.} + \text{P.I.}$$

$$x(t) = C_1 e^{-t} + C_2 e^{-2t} \dots\dots\dots (i)$$

Given  $x(0) = 20$ ;  $\therefore$  from equation (i),

$$\Rightarrow 20 = C_1 + C_2 \dots\dots\dots (ii)$$

$$x(1) = \frac{10}{e}; \therefore \text{from equation (i),}$$

$$\Rightarrow \frac{10}{e} = \frac{C_1}{e} + \frac{C_2}{e^2} \Rightarrow 10 = C_1 + \frac{C_2}{e} \dots\dots\dots (iii)$$

Solving (ii) and (iii)

$$\Rightarrow C_2 = \frac{10e}{e-1}, C_1 = \frac{10(e-2)}{e-1}$$

Put value of  $C_1$  and  $C_2$  in equation (i),

$$x(t) = \frac{10(e-2)}{(e-1)} \cdot e^{-t} + \frac{10e}{(e-1)} e^{-2t}$$

At  $t=2$ ,

$$x(2) = \frac{10(e-2)}{(e-1)} \cdot e^{-2} + \frac{10e}{(e-1)} e^{-4}$$

$$\Rightarrow x(2) = \frac{10}{(e-1)} (e^{-1} - 2e^{-2} + e^{-3})$$

$$\Rightarrow x(2) = 0.855$$



Scan for Video solution



**62. (b)**

$$\frac{d^2y}{dt^2} + 2 \cdot \frac{dy}{dt} + y = 0; \quad y(0) = y'(0) = 1$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dt} \text{ by } D \text{ and } \frac{d^2}{dt^2} \text{ by } D^2;$$

$$(D^2 + 2D + 1)y = 0$$

$\therefore$  Auxiliary equation is:

$$m^2 + 2m + 1 = 0$$

$$\Rightarrow (m+1)^2 = 0$$

$$\Rightarrow m = -1, 1$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = (C_1 t + C_2) e^{-t}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = \text{C.F.} + \text{P.I.}$

$$y = (C_1 t + C_2) \cdot e^{-t} \dots\dots\dots (i)$$

Differentiating above equation with respect to t,

$$y'(t) = -(C_1 t + C_2) \cdot e^{-t} + e^{-t} \cdot C_1 \dots\dots\dots (ii)$$

$$y(0) = 0; \text{ from equation (i)}$$

$$\Rightarrow 1 = C_2 \cdot e^{-0} \Rightarrow C_2 = 1$$

$$y'(0) = 1 \text{ and } C_2 = 1; \therefore \text{from equation (i)}$$

$$\Rightarrow 1 = -1 + C_1 \Rightarrow C_1 = 2$$

Put value of  $C_1$  and  $C_2$  in equation (i),

$$y(t) = (2t + 1) \cdot e^{-t}$$



Scan for Video solution



**63. (18)**

**Find:**  $y(1) = ?$

$$\frac{d^2y}{dx^2} = -12x^2 + 24x - 20$$

Integrating both sides w.r.t. x

$$\Rightarrow \frac{dy}{dx} = -4x^3 + 12x^2 - 20x + c_1 \quad \dots(i)$$

Integrating both sides, w.r.t. x

$$\Rightarrow y = -x^4 + 4x^3 - 10x^2 + c_1x + c_2 \quad \dots(ii)$$

At  $x = 0, y = 5$ , hence from equation (ii)

$$c_2 = 5$$

At  $x = 2, y = 21$ , hence from equation (ii)

$$21 = -16 + 32 - 40 + 2c_1 + c_2$$

$$\Rightarrow 21 = -24 + 2c_1 + 5$$

$$\Rightarrow c_1 = 20$$

∴ From equation (ii),

$$y = -x^4 + 4x^3 - 10x^2 + 20x + 5$$

At  $x = 1$

$$y(1) = -1 + 4 - 10 + 20 + 5$$

$$y(1) = 18$$



Scan for Video solution



64. (c)

Given  $\frac{d^2y}{dx^2} = y$  and curve passes through  $(0, 0)$  i.e.

at  $x = 0; y = 1$  and  $\left(\ln 2, \frac{3}{4}\right)$  i.e. at  $x = \ln 2; y = \frac{3}{4}$ .

$$\frac{d^2y}{dx^2} = y \Rightarrow \frac{d^2y}{dx^2} - y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2;$$

$$(D^2 - 1)y = 0$$

The Auxiliary equation is

$$m^2 - 1 = 0 \Rightarrow m = \pm 1$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = c_1 e^x + c_2 e^{-x}$$

Since, this is homogenous equation, therefore

$$\text{P.I.} = 0$$

Now, Complete solution:

$$y = \text{C.F.} + \text{P.I.}$$

$$\Rightarrow y = c_1 e^x + c_2 e^{-x} \quad \dots(i)$$

Since at  $x = 0; y = 1$ ; ∴ from equation (i),

$$0 = c_1 + c_2 \Rightarrow c_2 = -c_1$$

Since at  $x = \ln 2; y = \frac{3}{4}$ , ∴ from equation (i),

$$\frac{3}{4} = c_1 e^{\ln 2} + c_2 e^{-\ln 2} \quad [\text{Since } c_2 = -c_1]$$

$$\Rightarrow \frac{3}{4} = 2c_1 - \frac{c_1}{2} \Rightarrow \frac{3}{4} = \frac{3}{2}c_1$$

$$\Rightarrow c_1 = \frac{1}{2}$$

$$\text{and } c_2 = -c_1 \Rightarrow c_2 = -\frac{1}{2}$$

Put value of  $c_1$  and  $c_2$  in equation (i),

$$y = \frac{1}{2} \times e^x + \left(-\frac{1}{2}\right) \times e^{-x}$$

$$\Rightarrow y = \frac{1}{2} (e^x - e^{-x})$$



Scan for Video solution



65. (34 to 36)

$$\frac{d^2y}{dx^2} = 0 ; \text{ at } x = 0; y = 5 \text{ and at } x = 10; \frac{dy}{dx} = 2$$

Find:  $f(15) = ?$

$$\frac{d^2y}{dx^2} = 0$$

$$\Rightarrow \frac{d}{dx} \left( \frac{dy}{dx} \right) = 0$$

On Integrating,

$$\frac{dy}{dx} = c_1 \quad \dots(i)$$

at  $x = 10; \frac{dy}{dx} = 2$ , hence from equation (i),

$$c_1 = 2$$

Put value of  $c_1$  in equation (i),

$$\frac{dy}{dx} = 2$$

On Integrating,

$$y = 2x + c_2 \quad \dots(\text{ii})$$

At  $x = 0$ ;  $y = 5$ ; hence from equation (ii),

$$5 = 0 + c_2 \Rightarrow c_2 = 5$$

Put value of  $c_2$  in equation (ii),

$$y = 2x + 5$$

$$\Rightarrow f(x) = 2x + 5$$

At  $x = 15$

$$f(15) = 2(15) + 5 = 35$$

$$\Rightarrow f(15) = 35$$



Scan for Video solution



66. (a)

$$\text{Given } \frac{d^2x(t)}{dt^2} + x(t) = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dt} \text{ by } D \text{ and } \frac{d^2}{dt^2} \text{ by } D^2;$$

$$(D^2 + 1)x(t) = 0$$

The Auxiliary equation is

$$m^2 + 1 = 0 \Rightarrow m = \pm i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = 1$ , therefore complimentary function (C.F.) is given by,

$$C.F. = e^{at} (c_1 \cos \beta t + c_2 \sin \beta t)$$

$$\Rightarrow C.F. = e^{0 \times t} (c_1 \cos(t) + c_2 \sin(t))$$

$$\Rightarrow C.F. = c_1 \cos(t) + c_2 \sin(t)$$

Since, this is homogenous equation, therefore

$$\text{Particular integral (P.I.)} = 0$$

Now, Complete solution:

$$x(t) = C.F. + P.I.$$

$$\Rightarrow x(t) = c_1 \cos t + c_2 \sin t$$

At  $t = 0$ ,  $\cos t$  is 1 and  $\sin t$  is 0. Since  $x_1(0) = 1$ , therefore  $\cos t$  is  $x_1(t)$  and hence  $\sin t$  will be  $x_2(t)$ .

$$\Rightarrow x_1(t) = \cos t \text{ and } x_2(t) = \sin t$$

$$W(t) = \begin{vmatrix} x_1(t) & x_2(t) \\ \frac{dx_1(t)}{dt} & \frac{dx_2(t)}{dt} \end{vmatrix}$$

$$\Rightarrow W(t) = \begin{vmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{vmatrix} = \cos^2 t + \sin^2 t$$

$$\Rightarrow W(t) = 1$$

$$\text{At } t = \frac{\pi}{2};$$

$$W(t) = 1$$



Scan for Video solution



67. (a)

$$\text{Given DE is } \frac{d^2y}{dx^2} + 2\alpha \cdot \frac{dy}{dx} + y = 0 \text{ and having two equal roots.}$$

$$\frac{d^2y}{dx^2} + 2\alpha \cdot \frac{dy}{dx} + y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace,

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$\Rightarrow (D^2 + 2\alpha D + 1)y = 0$$

$\therefore$  Auxiliary Equation is:

$$m^2 + 2am + 1 = 0.$$

Comparing with  $ax^2 + bx + c = 0$ :  $a = 1$ ,  $b = 2\alpha$ ,  $c = 1$ ,

For the equation to have equal root,

$$b^2 - 4ac = 0$$

$$\Rightarrow (2\alpha)^2 - 4(1)(1) = 0$$

$$\Rightarrow 4\alpha^2 - 4 = 0$$

$$\Rightarrow \alpha^2 - 1 = 0$$

$$\Rightarrow \alpha = 1, -1$$

$\therefore$  For the DE to have equal roots,  $\alpha = \pm 1$



Scan for Video solution

**68. (b)**

Given DE is  $\frac{d^2x}{dt^2} + 2\frac{dx}{dt} + x = 0$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dt} \text{ by } D \text{ and } \frac{d^2}{dt^2} \text{ by } D^2;$$

$$\Rightarrow (D^2 + 2D + 1)x = 0$$

$\therefore$  Auxiliary equation is:

$$m^2 + 2m + 1 = 0$$

$$\Rightarrow (m+1)^2 = 0$$

$$\Rightarrow m = -1, 1$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$C.F. = (a + bt)e^{-t} = ae^{-t} + bte^{-t}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:

$$x = C.F. + P.I.$$

$$\Rightarrow x = ae^{-t} + bte^{-t}$$



Scan for Video solution

**69. (0.53 to 0.55)**

$$\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4y = 0; y(0) = y'(0) = 1$$

$$\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$\Rightarrow (D^2 + 4D + 4)y = 0$$

$\therefore$  Auxiliary equation is:

$$m^2 + 4m + 4 = 0$$

$$\Rightarrow (m+2)^2 = 0$$

$$\Rightarrow m = -2, 2$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$C.F. = (C_1 + C_2x)e^{-2x}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$y(x) = (C_1 + C_2x)e^{-2x} \quad \dots(i)$$

Differentiating above equation with respect to x,

$$y'(x) = (C_1 + C_2x)(-2e^{-2x}) + C_2e^{-2x} \quad \dots(ii)$$

Since,  $y(0) = 1$ ; from equation (i)

$$\Rightarrow 1 = C_1 \cdot e^{-2(0)} \Rightarrow C_1 = 1$$

Since,  $y'(0) = 1$  and  $C_1 = 1$ ;  $\therefore$  from equation (ii)

$$1 = (1 + 0)(-2e^0) + C_2e^0$$

$$\Rightarrow -2 + C_2 = 1$$

$$\Rightarrow C_2 = 3$$

Put value of  $C_1$  and  $C_2$  in equation (i),

$$y(x) = (1 + 3x)e^{-2x}$$

$\therefore$  at  $x = 1$ ,

$$y(1) = (1 + 3 \times 1)e^{-2 \times 1}$$

$$\Rightarrow y(1) = \frac{4}{e^2}$$

$$\Rightarrow y(1) = 0.541$$



Scan for Video solution



**70. (c)** Scan for Video solution

**70. (c)**

$$x^2 \cdot \frac{d^2y}{dx^2} + x \cdot \frac{dy}{dx} - y = 0 \quad \dots(i)$$

**This differential equation is Euler Cauchy's form of linear differential equation.**

Let,  $x = e^z$  and  $\frac{d}{dz} = D$

$$\Rightarrow x \frac{d}{dx} = D \text{ & } x^2 \frac{d^2}{dx^2} = D(D-1)$$

Then equation (i) can be written as:

$$[D(D-1) + D-1] y = 0$$

$$\Rightarrow (D^2 - 1) y = 0 \quad \dots(\text{ii})$$

Equation (ii) is homogenous linear differential equation with constant coefficients.

Its auxiliary equation is:

$$m^2 + 1 = 0 \Rightarrow m = \pm 1$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$C.F. = C_1 e^z + C_2 e^{-z}$$

Since, Equation (ii) is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$y = C_1 e^z + C_2 e^{-z} \quad [ \because x = e^z ]$$

$$\Rightarrow y = C_1 x + \frac{C_2}{x} \quad \dots(\text{iii})$$

In given option  $y = \frac{1}{x}$  is one of the solution of

given differential equation when  $C_1$  is 0 and  $C_2$  is 1.



Scan for Video solution



71. (c)

$$\text{Given: } \frac{d^2x}{dt^2} + 9x = 0 ; x(0) = 1 \text{ and } \left. \frac{dx}{dt} \right|_{t=0} = 1,$$

$$\frac{d^2x}{dt^2} + 9x = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$\frac{d}{dt}$  by  $D$  and  $\frac{d^2}{dt^2}$  by  $D^2$ ;

$$(D^2 + 9)x = 0$$

The Auxiliary equation is

$$m^2 + 9 = 0 \Rightarrow m = \pm 3i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = 3$ , therefore complimentary function (C.F.) is given by,

$$C.F. = e^{at} (c_1 \cos \beta t + c_2 \sin \beta t)$$

$$\Rightarrow C.F. = e^{0 \times t} (c_1 \cos(3t) + c_2 \sin(3t))$$

$$\Rightarrow C.F. = c_1 \cos 3t + c_2 \sin 3t$$

Since, this is homogenous equation, therefore

$$\text{Particular integral (P.I.)} = 0$$

Now, Complete solution:

$$x = C.F. + P.I.$$

$$x = (c_1 \cos 3t + c_2 \sin 3t) \quad \dots(\text{i})$$

Differentiating above equation with respect to t,

$$\frac{dx}{dt} = -3c_1 \sin 3t + 3c_2 \cos 3t \quad \dots(\text{ii})$$

At  $t = 0$ ;  $x = 1$ ; hence from equation (i),

$$1 = c_1 \times 1 + c_2 \times 0$$

$$\Rightarrow c_1 = 1$$

At  $t = 0$ ;  $\frac{dx}{dt} = 1$ , and  $c_1 = 1$ ; hence from equation (ii),

$$1 = -3 \times 1 \times 0 + 3c_2 \times 1$$

$$\Rightarrow c_2 = 1/3$$

Put value of  $c_1$  and  $c_2$  in equation (ii),

$$x = \cos 3t + \frac{1}{3} \sin 3t$$



Scan for Video solution



72. (b)

$$\text{Given: } \frac{d^2u}{dx^2} - k \cdot \frac{du}{dx} = 0 \quad ; u(0) = 0 \text{ & } u(L) = U$$

$$\frac{d^2u}{dx^2} - k \cdot \frac{du}{dx} = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$\frac{d}{dx}$  by  $D$  and  $\frac{d^2}{dx^2}$  by  $D^2$ ;

$$(D^2 - kD) u = 0$$

The auxiliary equation is

$$m^2 - km = 0 \Rightarrow m(m-k) = 0$$

$$\Rightarrow m = 0 \text{ or } m = k$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = C_1 e^{0x} + C_2 e^{kx}$$

$$\Rightarrow \text{C.F.} = C_1 + C_2 e^{kx}$$

Since, this is homogenous equation, therefore

$$\text{P.I.} = 0$$

Now, Complete solution:

$$u = \text{C.F.} + \text{P.I.}$$

$$\Rightarrow u = C_1 + C_2 e^{kx} \quad \dots(i)$$

Given  $u(0) = 0$ ,  $\therefore$  from equation (i);

$$0 = C_1 + C_2 e^0$$

$$\Rightarrow C_1 + C_2 = 0$$

$$\Rightarrow C_2 = -C_1$$

Given  $u(L) = U$ ,  $\therefore$  from equation (i);

$$\Rightarrow U = C_1 + C_2 e^{kL} \quad [\text{Since } C_2 = -C_1]$$

$$\Rightarrow U = C_1 [1 - e^{kL}]$$

$$\Rightarrow C_1 = \frac{U}{1 - e^{kL}}$$

$$\text{And } C_2 = -C_1 \Rightarrow C_2 = \frac{-U}{1 - e^{kL}}$$

Put value of  $C_1$  and  $C_2$  in equation (i),

$$u = \frac{U}{1 - e^{kL}} + \frac{-U}{1 - e^{kL}} \cdot e^{kx}$$

$$\Rightarrow u = U \left( \frac{1 - e^{kx}}{1 - e^{kL}} \right)$$



Scan for Video solution



### 73. (c)

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 0$$

This is homogenous linear differential equation with constant coefficients.

$$\text{Replace } \frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$(D^2 + D - 6)y = 0$$

Its auxiliary equation is:

$$m^2 + m - 6 = 0$$

$$\Rightarrow m^2 + 3m - 2m - 6 = 0$$

$$\Rightarrow (m+3)(m-2) = 0$$

$$\Rightarrow m = -3, 2$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = C_1 e^{-3x} + C_2 e^{2x}$$

Since, this is homogenous equation, therefore

$$\text{P.I.} = 0$$

Now, Complete solution:

$$y = \text{C.F.} + \text{P.I.}$$

$$\Rightarrow y = C_1 e^{-3x} + C_2 e^{2x}$$



Scan for Video solution



### 74. (d)

$$\frac{d^2}{dx^2} n(x) - \frac{n(x)}{L^2} = 0; n(0) = K \text{ and } n(\infty) = 0$$

$$\frac{d^2}{dx^2} n(x) - \frac{n(x)}{L^2} = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2;$$

$$\left( D^2 - \frac{1}{L^2} \right) n(x) = 0$$

The Auxiliary equation is:

$$m^2 - \frac{1}{L^2} = 0 \Rightarrow m = \pm \frac{1}{L}$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = C_1 \cdot e^{\frac{x}{L}} + C_2 \cdot e^{-\frac{x}{L}}$$

Since, this is homogenous equation, therefore  
P.I. = 0

Now, Complete solution:

$$n(x) = \text{C.F.} + \text{P.I.}$$

$$\Rightarrow n(x) = C_1 \cdot e^{\frac{x}{L}} + C_2 \cdot e^{-\frac{x}{L}} \quad \dots(i)$$

$$n(0) = K; \text{ from equation (i),}$$

$$\Rightarrow K = c_1 + c_2 \quad \dots(ii)$$

$n(\infty) = 0$ ; from equation (i),

$$\Rightarrow 0 = c_1 \cdot e^{\frac{x}{L}} + c_2(0)$$

To avoid discontinuity, ' $c_1$ ' should be zero.

From equation (ii),

$$\Rightarrow K = 0 + c_2 \Rightarrow c_2 = K$$

Put value of  $c_1$  and  $c_2$  in equation (i),

$$\therefore n(x) = K \cdot e^{-\frac{x}{L}}$$

$$\Rightarrow n(x) = K \cdot \exp\left(-\frac{x}{L}\right)$$



Scan for Video solution



75. (b)

$$\frac{d^2x}{dt^2} + 6 \frac{dx}{dt} + 8x = 0; \quad x(0) = 1 \text{ and } \left. \frac{dx}{dt} \right|_{t=0} = 0$$

$$\frac{d^2x}{dt^2} + 6 \frac{dx}{dt} + 8x = 0$$

This is homogenous linear differential equation with constant coefficients.

Replace  $\frac{d}{dt}$  by  $D$  and  $\frac{d^2}{dt^2}$  by  $D^2$ ;

$$(D^2 + 6D + 8)x = 0$$

The Auxiliary equation is

$$m^2 + 6m + 8 = 0$$

$$\Rightarrow m^2 + 4m + 2m + 8 = 0$$

$$\Rightarrow (m+4)(m+2) = 0$$

$$\Rightarrow m = -2, -4$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$CF = C_1 e^{-2t} + C_2 e^{-4t}$$

Since, this is homogenous equation, therefore

$$P.I. = 0$$

Now, Complete solution:

$$x = C.F. + P.I.$$

$$\Rightarrow x = C_1 e^{-2t} + C_2 e^{-4t} \quad \dots(i)$$

Differentiating above equation with respect to t,

$$\frac{dx}{dt} = -2 C_1 e^{-2t} - 4 C_2 e^{-4t} \quad \dots(ii)$$

At  $t = 0$ ;  $x = 1$ ; hence from equation (i),

$$1 = C_1 + C_2 \quad \dots(iii)$$

$$\text{At } t = 0; \frac{dx}{dt} = 0; \text{ hence from equation (ii),}$$

$$0 = -2 C_1 - 4 C_2 \quad \dots(iv)$$

On solving (iii) and (iv)

$$C_1 = 2 \text{ and } C_2 = -1$$

Put value of  $C_1$  and  $C_2$  in equation (ii),

$$x = 2e^{-2t} - 1e^{-4t}$$

$$\Rightarrow x = 2e^{-2t} - e^{-4t}$$



Scan for Video solution



76. (b)

$$\text{Given: } \ddot{x} + 3x = 0; \quad x(0) = 1; \quad \dot{x}(0) = 0$$

Find:  $x(1) = ?$

$$\ddot{x} + 3x = 0 \Rightarrow \frac{d^2x}{dt^2} + 3x = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$\frac{d}{dt}$  by  $D$  and  $\frac{d^2}{dt^2}$  by  $D^2$ ;

$$(D^2 + 3)x = 0$$

The auxiliary equation is given by

$$m^2 + 3 = 0 \Rightarrow m = \pm \sqrt{3} i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = \sqrt{3}$ , therefore complimentary function (C.F.) is given by,

$$C.F. = e^{\alpha x} (c_1 \cos \beta t + c_2 \sin \beta t)$$

$$\Rightarrow C.F. = e^{0 \cdot t} (c_1 \cos(\sqrt{3}t) + c_2 \sin(\sqrt{3}t))$$

$$\Rightarrow C.F. = c_1 \cos \sqrt{3}t + c_2 \sin \sqrt{3}t$$

Since, this is homogenous equation, therefore

$$\text{Particular integral (P.I.)} = 0$$

Now, Complete solution:

$$x(t) = C.F. + P.I.$$

$$\Rightarrow x(t) = c_1 \cos \sqrt{3}t + c_2 \sin \sqrt{3}t \quad \dots(i)$$

Differentiating with respect to t,

$$\dot{x}(t) = -c_1 \sqrt{3} \cdot \sin \sqrt{3}t + c_2 \sqrt{3} \cdot \cos \sqrt{3}t \quad \dots(ii)$$

$\therefore x(0) = 1$ , hence from equation (i),

$$1 = c_1 + c_2(0) \Rightarrow c_1 = 1$$

$\therefore \dot{x}(0) = 0$  and  $c_1 = 1$ , hence from equation (ii),

$$0 = 0 + c_2 \cdot \sqrt{3} \cdot (1) \Rightarrow c_2 = 0$$

Put value of  $c_1$  and  $c_2$  in equation (i),

$$x(t) = \cos \sqrt{3}t$$

At  $t=1$ ,

$$x(1) = \cos \sqrt{3}(1) = -0.16$$

$$\Rightarrow x(1) = -0.16$$



Scan for Video solution



77. (a)

**Given:**  $y'' + 2y' + y = 0$ ;  $y(0) = 0$ ;  $y(1) = 0$

**Find:**  $y(1) = ?$

$$y'' + 2y' + y = 0$$

This is homogenous linear differential equation with constant coefficients. Replace

$$\frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2;$$

The auxiliary equation is:

$$m^2 + 2m + 1 = 0$$

$$\Rightarrow (m+1)^2 = 0$$

$$\Rightarrow m = -1, -1$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$C.F. = (c_1 + c_2 x) e^{-x}$$

Since, this is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$y = (c_1 + c_2 x) e^{-x} \quad \dots(i)$$

$\therefore y(0) = 0$ , hence from equation (i),

$$0 = c_1 \cdot (1) \Rightarrow c_1 = 0$$

$\therefore y(1) = 0$  and  $c_1 = 0$ , hence from equation (i),

$$0 = c_2 \cdot e^{-1} \Rightarrow c_2 = 0$$

Put value of  $c_1$  and  $c_2$  in equation (i),

$$y = 0$$

at  $x = 0.5$ ,

$$y(0.5) = 0$$



Scan for Video solution



78. (a)

$$\frac{d^2y}{dx^2} + y = 0$$

This is homogenous linear differential equation with constant coefficients.

$$\text{Replace } \frac{d}{dx} \text{ by } D \text{ and } \frac{d^2}{dx^2} \text{ by } D^2$$

$$(D^2 + 1)y = 0$$

Its auxiliary equation is:

$$m^2 + 1 = 0 \Rightarrow m = \pm i$$

Since roots are in the form of pair of complex number  $\alpha \pm i\beta$ , where  $\alpha = 0$  and  $\beta = 1$ , therefore complimentary function (C.F.) is given by,

$$C.F. = e^{\alpha x} (c_1 \cos \beta x + c_2 \sin \beta x)$$

$$\Rightarrow C.F. = e^{0 \times x} (c_1 \cos(1 \times x) + c_2 \sin(1 \times x))$$

$$\Rightarrow C.F. = c_1 \cos x + c_2 \sin x$$

Since, this is homogenous equation, therefore Particular integral (P.I.) = 0

Now, Complete solution:

$$y = C.F. + P.I.$$

$$\Rightarrow y = c_1 \cos x + c_2 \sin x$$

Replace arbitrary constants  $c_1$  by P and  $c_2$  by Q,

$$\Rightarrow y = P \cos x + Q \sin x$$



Scan for Video solution



**79. (8.999 to 9.001)**

$$x^2 \cdot \frac{d^2y}{dx^2} + x \cdot \frac{dy}{dx} - y = 0 ;$$

$$y(x=1) = 6, \quad \left. \frac{dy}{dx} \right|_{x=1} = 2$$

**Find:**  $y(2) = ?$

$$x^2 \cdot \frac{d^2y}{dx^2} + x \cdot \frac{dy}{dx} - y = 0 \quad \dots(i)$$

**This differential equation is Euler Cauchy's form of linear differential equation.**

$$\text{Let, } x = e^z \text{ and } \frac{d}{dz} = D$$

$$\Rightarrow x \frac{d}{dx} = D \text{ & } x^2 \frac{d^2}{dx^2} = D(D-1)$$

Then equation (i) can be written as:

$$[D(D-1) + D - 1] y = 0$$

$$\Rightarrow (D^2 - 1) y = 0 \quad \dots(ii)$$

Equation (ii) is homogenous linear differential equation with constant coefficients.

Its auxiliary equation is:

$$m^2 + 1 = 0 \Rightarrow m = \pm 1$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = C_1 e^z + C_2 e^{-z}$$

Since, Equation (ii) is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = \text{C.F.} + \text{P.I.}$

$$\Rightarrow y = C_1 e^z + C_2 e^{-z} \quad [ \because x = e^z ]$$

$$\Rightarrow y = C_1 x + \frac{C_2}{x} \quad \dots(iii)$$

$$\Rightarrow \frac{dy}{dx} = C_1 - \frac{C_2}{x^2} \quad \dots(iv)$$

At  $x = 1, y = 6$ , hence from equation (iii),

$$C_1 + C_2 = 6 \quad \dots(v)$$

At  $x = 1, \frac{dy}{dx} = 2$ , hence from equation (iv),

$$C_1 - C_2 = 2 \quad \dots(vi)$$

Solving equation (v) and (vi),

$$C_1 = 4 \text{ and } C_2 = 2$$

Put value of  $C_1$  and  $C_2$  in equation (iii),

$$y = 4x + \frac{2}{x}$$

At  $x = 2$ ,

$$y(2) = 8 + 1 = 9$$

$$\Rightarrow y(2) = 9$$

Scan for Video solution



**80. (a)**

**Given**

$$x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + 2y = 4$$

$$\left( x^2 \frac{d^2}{dx^2} - 2x \frac{d}{dx} + 2 \right) y = 4 \quad \dots(i)$$

**This differential equation is Euler Cauchy's form of linear differential equation.**

$$\text{Let, } x = e^z \text{ and } \frac{d}{dz} = D$$

$$\text{Then, } x \cdot \frac{d}{dx} = D \text{ and } x^2 \frac{d^2}{dx^2} = D(D-1)$$

Then equation (i) can be written as:

$$[D(D-1)-2D+2] y = 4$$

$$\Rightarrow (D^2 - 3D + 2) y = 4 \quad \dots(ii)$$

Equation (ii) is non-homogenous linear differential equation with constant coefficients.

Its auxiliary equation is:

$$m^2 - 3m + 2 = 0 \Rightarrow (m-2)(m-1) = 0$$

$$\Rightarrow m = 2, 1$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$\text{C.F.} = C_1 e^{2z} + C_2 e^z$$

Since, equation (ii) is non-homogenous equation, therefore Particular integral (P.I.)

$$\text{P.I.} = \frac{1}{(D^2 - 3D + 2)} \times 4$$

$$\Rightarrow P.I. = 4 \cdot \frac{1}{D^2 - 3D + 2} \cdot e^{0.z}$$

$$\Rightarrow P.I. = 4 \cdot \frac{1}{0^2 - 3 \times 0 + 2} \cdot e^{0.z} = \frac{4}{2}$$

$$\Rightarrow P.I. = 2$$

Now, Complete solution:  $y = C.F. + P.I.$

$$\Rightarrow y = C_1 e^{2z} + C_2 e^z + 2 \quad \dots(iii)$$

Since  $x = e^z \therefore$  from equation (iii),

$$y = C_1 x^2 + C_2 x + 2$$



Scan for Video solution



### 81. (5.9 to 6.1)

$$x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + 2y = 0 \therefore ; y(1) = 0 \text{ and } y(2) = 2$$

Given differential equation can be written as:

$$\left( x^2 \frac{d^2}{dx^2} - 2x \frac{d}{dx} + 2 \right) y = 0 \quad \dots(i)$$

This differential equation is Euler Cauchy's form of linear differential equation.

$$\text{Let, } x = e^z \text{ and } \frac{d}{dz} = D$$

$$\Rightarrow x \frac{d}{dx} = D \& x^2 \frac{d^2}{dx^2} = D(D-1)$$

Then equation (i) can be written as:

$$[D(D-1)-2D+2]y = 0$$

$$[D^2-3D+2]y = 0 \quad \dots(ii)$$

Equation (ii) is homogenous linear differential equation with constant coefficients.

Its auxiliary equation is:

$$m^2 - 3m + 2 = 0 \Rightarrow (m-1)(m-2) = 0$$

$$\Rightarrow m = 1, 2$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$C.F. = C_1 e^z + C_2 e^{2z}$$

Since, Equation (ii) is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$y = C_1 e^z + C_2 e^{2z} \quad \dots(iii)$$

Since  $x = e^z \therefore$  from equation (iii),

$$y = C_1 x + C_2 x^2 \quad \dots(iv)$$

Given,  $y(1) = 0$ ; from equation (iv)

$$0 = C_1 \times 1 + C_2 \times 1^2$$

$$\Rightarrow C_1 + C_2 = 0 \quad \dots(v)$$

Given,  $y(2) = 2$ ; from equation (iv)

$$2 = C_1 \times 2 + C_2 \times 2^2$$

$$\Rightarrow C_1 + 2C_2 = 1 \quad \dots(vi)$$

Equation (vi) – (v), we get

$$C_2 = 1 \quad [\text{Put in equation (v)}]$$

$$C_1 + 1 = 0 \Rightarrow C_1 = -1$$

Put value of  $C_1$  and  $C_2$  in equation (iv),

$$y = -x + x^2$$

At  $x = 3$ ;

$$y(3) = -3 + 3^2$$

$$y(3) = 6$$



Scan for Video solution



### 82. (5.24 to 5.26)

$$x^2 \frac{d^2y}{dx^2} - 3x \frac{dy}{dx} + 3y = 0 ; y(1) = 1 \text{ and } y(2) = 14$$

Find:  $y(1.5) = ?$

$$x^2 \frac{d^2y}{dx^2} - 3x \frac{dy}{dx} + 3y = 0 \quad \dots(i)$$

This differential equation is Euler Cauchy's form of linear differential equation.

$$\text{Let, } x = e^z \text{ and } \frac{d}{dz} = D$$

$$\Rightarrow x \frac{d}{dx} = D \& x^2 \frac{d^2}{dx^2} = D(D-1)$$

Then equation (i) can be written as:

$$(D(D-1) - 3D + 3)y = 0$$

$$\Rightarrow (D^2 - 4D + 3)y = 0 \quad \dots(ii)$$

Equation (ii) is homogenous linear differential equation with constant coefficients.

Its auxiliary equation is:

$$m^2 - 4m + 3 = 0$$

$$\Rightarrow (m-1)(m-3) = 0$$

$$\Rightarrow m = 1, 3$$

Since roots are real and same, therefore complimentary function (C.F.) is given by,

$$C.F. = c_1 e^z + c_2 e^{3z}$$

Since, Equation (ii) is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$\Rightarrow y = c_1 e^z + c_2 e^{3z} \quad [ \because x = e^z ]$$

$$\Rightarrow y = c_1 x + c_2 x^3 \quad \dots (iii)$$

Given,  $y(1) = 1$ , hence from equation (iii),

$$\Rightarrow 1 = c_1 + c_2 \quad \dots (iv)$$

Given,  $y(2) = 14$ , hence from equation (iii),

$$14 = 2c_1 + 8c_2 \quad \dots (v)$$

Solving equation (iv) and (v)

$$\Rightarrow c_1 = -1 \text{ and } c_2 = 2$$

Put value of  $c_1$  and  $c_2$  in equation (iii),

$$y = -x + 2x^3$$

At,  $x = 1.5$ ,

$$y(1.5) = -1.5 + 2(1.5)^3 = 5.25$$

$$\therefore y(1.5) = 5.25$$



Scan for Video solution



83. (a)

$$x^2 \cdot \frac{d^2 y}{dx^2} + x \cdot \frac{dy}{dx} - 4y = 0 ; y(0) = 0 \text{ & } y(1) = 1$$

$$x^2 \cdot \frac{d^2 y}{dx^2} + x \cdot \frac{dy}{dx} - 4y = 0 \quad \dots (i)$$

**This differential equation is Euler Cauchy's form of linear differential equation.**

$$\text{Let, } x = e^z \text{ and } \frac{d}{dz} = D$$

$$\Rightarrow x \frac{d}{dx} = D \text{ & } x^2 \frac{d^2}{dx^2} = D(D-1)$$

Then equation (i) can be written as:

$$[D(D-1) + D - 4] y = 0$$

$$\Rightarrow (D^2 - 4)y = 0 \quad \dots (ii)$$

Equation (ii) is homogenous linear differential equation with constant coefficients.

Its auxiliary equation is:

$$m^2 - 4 = 0 \Rightarrow (m-2)(m+2) = 0$$

$$\Rightarrow m = 2, -2$$

Since roots are real and different, therefore complimentary function (C.F.) is given by,

$$C.F. = c_1 e^{2z} + c_2 e^{-2z}$$

Since, Equation (ii) is homogenous equation, therefore P.I. = 0.

Now, Complete solution:  $y = C.F. + P.I.$

$$y = c_1 e^{2z} + c_2 e^{-2z} \quad [ \because x = e^z ]$$

$$\Rightarrow y = c_1 x^2 + \frac{c_2}{x^2}$$

$$\Rightarrow yx^2 = c_1 x^4 + c_2 \quad \dots (iii)$$

$\because y(0) = 0$ , hence from equation (iii)

$$0 = 0 + c_2 \Rightarrow c_2 = 0$$

$\because y(1) = 1$  and  $c_2 = 0$ , hence from equation (iii)

$$1 = c_1 \Rightarrow c_1 = 1$$

Put value of  $c_1$  and  $c_2$  in equation (iii),

$$yx^2 = x^4$$

$$\Rightarrow y = x^2$$




Scan for Video solution



84. (0.395)

Given 1-D diffusion equation :-



$$\frac{\partial T}{\partial t} = D \frac{\partial^2 T}{\partial x^2}$$

Also  $D = 1$ ,

$$T(x, 0) = T$$

$$T(0, t) = T(5, t) = 0$$

We know, solution of P.D.E is given by:-

$$T(x, t) = (C_1 \cos \alpha x + C_2 \sin \alpha x) C_3 e^{-\alpha^2 t} \quad \text{---(i)}$$

For  $T(0, t) = 0$

$$0 = (C_1 + 0) C_3 e^{-\alpha^2 t}$$

$$\Rightarrow C_1 = 0$$

For  $T(5, t) = 0$

$$0 = C_2 \sin 5\alpha C_3 e^{-\alpha^2 t}$$

$$\Rightarrow \sin 5\alpha = 0$$

$$\Rightarrow \alpha = \frac{n\pi}{5} \quad [n \rightarrow 1, 2, 3, \dots]$$

Now from equation (i)

$$T(x, t) = C_2 C_3 \sin\left(\frac{n\pi x}{5}\right) e^{-\frac{n^2 \pi^2 t}{25}}$$

∴ The generalized solution will be

$$T(x, t) = \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{5}\right) e^{-\frac{n^2 \pi^2 t}{25}}$$

From Fourier series,


$b_n = 0$  for even value of  $n$

$$\therefore T(x, t) = \sum_{n=1,3,5}^{\infty} b_n \sin\left(\frac{n\pi x}{5}\right) e^{-\frac{n^2 \pi^2 t}{25}}$$

$$\text{Given } T(x, t) = \sum_{n=1,3,5}^{\infty} C_n \sin\left(\frac{n\pi x}{5}\right) e^{-\beta n^2 t}$$

On comparison:


$$\beta = \frac{\pi^2}{25} s^{-1} = 0.395 \text{ sec}^{-1}$$



85. (25)

We know, Laplace equation is 2-dimension on a unit square with Dirichlet boundary condition:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad 0 < x, y < 1$$



This Laplace equation is also called harmonic function whose solution is given by

$$u(x, y) = (C_1 \cos \beta x + C_2 \sin \beta x) (C_3 e^{\beta y} + C_4 e^{-\beta y})$$

Let square ABCD is unit square.

Now given  $u(x, 0) = u(0, y) = u(1, y) = 0$

$$u(x, 1) = 100$$

For  $u(x, 0) = 0$

$$\Rightarrow 0 = (C_1 \cos \beta x + C_2 \sin \beta x)(C_3 + C_4)$$

$$[C_3 = -C_4]$$

For  $u(0, y) = 0$

$$\Rightarrow 0 = C_1 (C_3 e^{\beta y} + C_4 e^{-\beta y})$$

$$[C_1 = 0]$$

$$\text{Now } u(x, y) = C_2 \sin \beta x (C_3 e^{\beta y} - C_3 e^{-\beta y})$$

$$[c_2 c_3 = a_x]$$

$$u(x, y) = a_x \sin \beta x (e^{\beta y} - e^{-\beta y}) \quad \text{---(i)}$$

For  $u(1, y) = 0$

$$0 = a_x \sin \beta (e^{\beta y} - e^{-\beta y})$$

$$\sin \beta = 0$$

$$[\beta = n\pi]$$

For  $u(x, 1) = 100$ ,

$$100 = a_x \sin n\pi x (e^{n\pi} - e^{-n\pi})$$

$$\Rightarrow a_x = \frac{100}{\sin n\pi x (e^{n\pi} - e^{-n\pi})}$$

From equation (i)

$$u(x, y) = \sum_{n=1}^{\infty} \frac{100}{\sin n\pi x (e^{n\pi} - e^{-n\pi})} \sin n\pi x (e^{n\pi y} - e^{-n\pi y})$$

$$u(x, y) = \sum_{n=1}^{\infty} 100 \left( \frac{e^{n\pi y} - e^{-n\pi y}}{e^{n\pi} - e^{-n\pi}} \right)$$

$$\text{Now, at mid-point } (x, y) = \left( \frac{1}{2}, \frac{1}{2} \right)$$

$$\left( \frac{1}{2}, \frac{1}{2} \right) \text{ [Given]}$$

$$\begin{aligned} \therefore T_0 &= \sum_{n=1}^{\infty} 100 \frac{e^{n\pi/2} - e^{-n\pi/2}}{e^{n\pi} - e^{-n\pi}} \\ &= 100 \left[ \frac{e^{\pi/2} - e^{-\pi/2}}{e^\pi - e^{-\pi}} + \frac{e^{\pi} - e^{-\pi}}{e^{2\pi} - e^{-2\pi}} + \frac{e^{3\pi/2} - e^{-3\pi/2}}{e^{3\pi} - e^{-3\pi}} + \dots \right] \\ &= 100 [0.199 + 0.043 + 0.008 + \dots] \end{aligned}$$

$$T_0 \approx 25^\circ\text{C}$$



Scan for Video solution



### 86. (a)

$$Z = \sin(y + it) + \cos(y - it) \quad \dots(\text{i})$$

Differentiating equation (i) partially with respect to x,

$$\frac{\partial Z}{\partial y} = \cos(y + it) - \sin(y - it)$$

Again, differentiating equation (i) partially with respect to x,

$$\frac{\partial^2 Z}{\partial y^2} = -\sin(y + it) - \cos(y - it) \quad \dots(\text{ii})$$

Differentiating equation (i) partially with respect to t,

$$\frac{\partial Z}{\partial t} = i \cos(y + it) + i \sin(y - it)$$

Again, differentiating equation (i) partially with respect to t,

$$\frac{\partial^2 Z}{\partial t^2} = -t^2 \sin(y + it) - i^2 \cos(y - it)$$

[Since,  $i^2 = -1$ ]

$$\frac{\partial^2 Z}{\partial t^2} = \sin(y + it) + \cos(y - it) \quad \dots(\text{iii})$$

Adding equation (ii) and (iii)

$$\frac{\partial^2 Z}{\partial t^2} + \frac{\partial^2 Z}{\partial y^2} = 0$$



Scan for Video solution



### 87. (d)

According to given condition function  $f(x, y)$  is nothing but constant function i.e.  $f(x, y) = 3$  because this is the only function whose value is 3 at any point on the boundary of unit circle and it also satisfies Laplace's equation,

$$f(x, y) = 3$$

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0$$

$$\therefore f(0, 0) = 3$$



Scan for Video solution



### 88. (a, b)

Given, PDE is

$$a \cdot \frac{\partial^2 f}{\partial x^2} + b \cdot \frac{\partial^2 f}{\partial y^2} = f(x, y) \quad \dots(1)$$

Given,

$$f(x, y) = e^{(\xi x + \eta y)} = e^{\xi x} \cdot e^{\eta y} \quad \dots(2)$$

Differentiating partially with respect to x,

$$\Rightarrow \frac{\partial f}{\partial x} = \xi \cdot e^{\xi x} \cdot e^{\eta y}$$

Differentiating again partially with respect to x,

$$\Rightarrow \frac{\partial^2 f}{\partial x^2} = \xi^2 \cdot e^{\xi x} \cdot e^{\eta y} \quad \dots(3)$$

Differentiating equation (2) partially with respect to y,

$$\frac{\partial f}{\partial y} = \eta \cdot e^{\xi x} \cdot e^{\eta y}$$

Differentiating again partially with respect to y,

$$\Rightarrow \frac{\partial^2 f}{\partial y^2} = \eta^2 \cdot e^{\xi x} \cdot e^{\eta y} \quad \dots(4)$$

From equation (1), (2), (3) and (4),

$$\Rightarrow a(\xi^2 \cdot e^{\xi x} \cdot e^{\eta y}) + b(\eta^2 \cdot e^{\xi x} \cdot e^{\eta y}) = e^{\xi x} \cdot e^{\eta y}$$

$$\Rightarrow (a\xi^2 + b\eta^2) \cdot e^{\xi x} \cdot e^{\eta y} = e^{\xi x} \cdot e^{\eta y}$$

$$\Rightarrow a\xi^2 + b\eta^2 = 1$$

**Out of the given options, A and B satisfy the above equation. Hence option A and B are correct.**



Scan for Video solution



**89. (b)**

1<sup>st</sup> term of given equation contains product of dependent variable with its derivative, so it is non-linear and also we have 2<sup>nd</sup> order derivative so its order is 2

∴ It is second order non - linear differential equation.



Scan for Video solution



**90. (a)**

The given P.D.E is parabolic which requires one initial condition and two boundary conditions to solve it uniquely.



Scan for Video solution



**91. (b)**

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$$

This is 2-D wave equation.



Scan for Video solution



**92. (b)**

The equation  $\frac{\partial u}{\partial t} = C^2 \cdot \frac{\partial^2 u}{\partial x^2}$  is standard form of 1-dimensional heat equation.

$$\therefore \text{At } C = 1, \frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$$

∴ Given equation is heat equation.

**NOTE:**

$$(1) \frac{\partial u}{\partial t} = C^2 \cdot \frac{\partial^2 u}{\partial x^2} \rightarrow \text{Heat equation}$$

$$(2) \frac{\partial^2 u}{\partial t^2} = C^2 \cdot \frac{\partial^2 u}{\partial x^2} \rightarrow \text{Wave equation}$$

$$(3) \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \rightarrow \text{Laplace equation}$$



Scan for Video solution



**93. (b)**

**Given**

$$\frac{\partial u}{\partial y} + c \cdot \frac{\partial u}{\partial x} = 0 \quad \dots(i)$$

Let  $u = f(ax + by)$  is a solution, then

$$\Rightarrow \frac{\partial u}{\partial x} = a.f'(ax + by)$$

$$\text{And} \quad \frac{\partial u}{\partial y} = b.f'(ax + by)$$

Hence from equation (i0,

$$b \cdot f'(ax + by) + c \cdot a \cdot f'(ax + by) = 0$$

$$\Rightarrow (b + ac)f'(ax + by) = 0$$

$$\Rightarrow ac + b = 0$$

$$\Rightarrow b = -ac$$

The solution is:

$$u = f(ax + by) \quad [\text{Since } b = -ac]$$

$$\Rightarrow u = f(ax - acy)$$

$$\Rightarrow u = f(a(x - cy))$$

For  $a = 1$ ,

$$\Rightarrow \boxed{u = f(x - cy)}.$$



Scan for Video solution

**94. (36)****Given,**

$$3 \frac{\partial^2 \phi}{\partial x^2} + B \frac{\partial^2 \phi}{\partial x \partial y} + 3 \frac{\partial^2 \phi}{\partial y^2} + 4\phi = 0 \dots(i)$$

And equation is parabolic.

General form of second order partial differential equation,

$$A \frac{\partial^2 \phi}{\partial x^2} + B \frac{\partial^2 \phi}{\partial x \partial y} + C \frac{\partial^2 \phi}{\partial y^2} + f(\phi_x, \phi_y, x, y, c) = 0$$

| If $B^2 - 4AC$ | Type of curve |
|----------------|---------------|
| 0              | Parabolic     |
| Less than 0    | Elliptic      |
| Greater than 0 | Hyperbolic    |

Comparing General form of second order partial differential equation with given equation (i),

We have,  $A = 3$ ,  $B = B$ ,  $C = 3$ 

For P.D.E to be parabolic,

$$B^2 - 4AC = 0$$

$$B^2 - 4(3)(3) = 0$$

$$B^2 = 36$$



Scan for Video solution

**95. (d)**

Standard form of wave equation:

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$$

$$\text{Given, } \frac{\partial^2 u}{\partial x^2} = 25 \frac{\partial^2 u}{\partial t^2}$$

$$\frac{\partial^2 u}{\partial t^2} = \frac{1}{25} \frac{\partial^2 u}{\partial x^2}$$



Scan for Video solution



**97. (b)**

The solution of  $\frac{\partial u}{\partial t} = \alpha \frac{\partial^2 u}{\partial x^2}$  is

$$u(x, t) = (A \cos px + B \sin px) C e^{-p^2 \alpha t} \quad \text{.....(i)}$$

Let  $-p^2 \alpha = k$

$$\Rightarrow p = \sqrt{-\frac{k}{\alpha}} = \sqrt{\frac{k}{\alpha}} i$$

Putting value of p in equation (i)

$$u(x, t) = \left[ A \cos i \sqrt{\frac{k}{\alpha}} x + B \sin i \sqrt{\frac{k}{\alpha}} x \right] C e^{kt}$$

$$u(x, t) = \left[ A \cosh \sqrt{\frac{k}{\alpha}} x + i B \sinh \sqrt{\frac{k}{\alpha}} x \right] C e^{kt}$$

$$u(x, t) = \left[ A \left\{ \frac{e^{\sqrt{\frac{k}{\alpha}} x} + e^{-\sqrt{\frac{k}{\alpha}} x}}{2} \right\} + i B \left\{ \frac{e^{\sqrt{\frac{k}{\alpha}} x} - e^{-\sqrt{\frac{k}{\alpha}} x}}{2} \right\} \right] C e^{kt}$$

$$u(x, t) = C e^{kt} \left[ e^{\sqrt{\frac{k}{\alpha}} x} \left\{ \frac{A + iB}{2} \right\} + e^{-\sqrt{\frac{k}{\alpha}} x} \left\{ \frac{A - iB}{2} \right\} \right]$$

$$u(x, t) = C_1 e^{\sqrt{\frac{k}{\alpha}} x} + C_2 e^{-\sqrt{\frac{k}{\alpha}} x}$$



Scan for Video solution



**98. (b)**

Laplace equation is  $\nabla^2 f = 0$

Any solution of Laplace equation with 2<sup>nd</sup> order derivatives being continuous is called **Harmonic functions.**



Scan for Video solution



**99. (c)**

Given  $z = (xy) \ln(xy) \dots \text{(i)}$

Differentiating equation (i) partially with respect to x,

$$\frac{\partial z}{\partial x} = (xy) \frac{1}{xy} y + \ln(xy) y$$

$$\Rightarrow \frac{\partial z}{\partial x} = y(1 + \ln(xy))$$

Multiply above equation by x

$$\Rightarrow x \frac{\partial z}{\partial x} = xy(1 + \ln(xy)) \dots \text{(ii)}$$

Differentiating equation (i) partially with respect to y,

$$\frac{\partial z}{\partial y} = (xy) \frac{1}{xy} x + \ln(xy) x$$

$$\Rightarrow \frac{\partial z}{\partial y} = x(1 + \ln(xy))$$

Multiply above equation by y,

$$\Rightarrow y \frac{\partial z}{\partial y} = xy(1 + \ln(xy)) \dots \text{(iii)}$$

Equation (ii) ÷ (iii)

$$\frac{x \frac{\partial z}{\partial x}}{y \frac{\partial z}{\partial y}} = \frac{xy(1 + \ln(xy))}{xy(1 + \ln(xy))}$$

$$\Rightarrow x \frac{\partial z}{\partial x} = y \frac{\partial z}{\partial y}$$



Scan for Video solution



**100. (c)**

$z = ax + by + ab \dots \text{(i)}$

Differentiating partially equation (i) with respect to x,

$$p = \frac{\partial z}{\partial x} = a \Rightarrow a = p$$

Differentiating partially equation (i) with respect to y,

$$q = \frac{\partial z}{\partial y} = b \Rightarrow b = q$$

By substituting a and b in (i) in terms of p and q, we get

$$z = px + qy + pq$$



Scan for Video solution



101. (c)

**Given:**

$$f = y^x$$

$$\Rightarrow \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial x} \right)$$

$$\Rightarrow \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial y} (y^x \cdot \ln y)$$

$$\Rightarrow \frac{\partial^2 f}{\partial x \partial y} = y^x \cdot \frac{1}{y} + \ln y \times x \cdot y^{x-1}$$

$$\Rightarrow \frac{\partial^2 f}{\partial x \partial y} = y^x \left\{ \frac{1}{y} + \frac{x}{y} \cdot \log_e y \right\}$$

∴ At x = 2 & y = 1

$$\Rightarrow \frac{\partial^2 f}{\partial x \partial y} \Big|_{(2,1)} = 1^2 \left\{ \frac{1}{1} + \frac{2}{1} \cdot \log_e 1 \right\} = 1$$

$$\Rightarrow \boxed{\frac{\partial^2 f}{\partial x \partial y} \Big|_{(2,1)} = 1}$$



Scan for Video solution



102. (d)

**1 Transform**  $k_x \frac{\partial^2 h}{\partial x^2} + k_z \frac{\partial^2 h}{\partial z^2} = 0$  to

$$\frac{\partial^2 h}{\partial x_t^2} + \frac{\partial^2 h}{\partial z^2} = 0 .$$

Now,

$$\frac{\partial^2 h}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial h}{\partial x} \right) = \frac{\partial}{\partial x_t} \cdot \frac{\partial x_t}{\partial x} \left( \frac{\partial h}{\partial x_t} \cdot \frac{\partial x_t}{\partial x} \right)$$

$$\frac{\partial^2 h}{\partial x^2} = \frac{\partial}{\partial x_t} \left( \frac{\partial h}{\partial x_t} \right) \cdot \left( \frac{\partial x_t}{\partial x} \right)^2$$

$$\frac{\partial^2 h}{\partial x^2} = \left( \frac{\partial^2 h}{\partial x_t^2} \right) \cdot \left( \frac{\partial x_t}{\partial x} \right)^2 \quad \dots(i)$$

Now,

$$k_x \frac{\partial^2 h}{\partial x^2} + k_z \frac{\partial^2 h}{\partial z^2} = 0$$

From equation (i),

$$k_x \left( \frac{\partial^2 h}{\partial x_t^2} \right) \cdot \left( \frac{\partial x_t}{\partial x} \right)^2 + k_z \frac{\partial^2 h}{\partial z^2} = 0$$

$$\frac{k_x}{k_z} \left( \frac{\partial x_t}{\partial x} \right)^2 \left( \frac{\partial^2 h}{\partial x_t^2} \right) + \frac{\partial^2 h}{\partial z^2} = 0 \quad \dots(ii)$$

We have to transform above equation as:

$$\frac{\partial^2 h}{\partial x_t^2} + \frac{\partial^2 h}{\partial z^2} = 0 \quad \dots(iii)$$

Comparing equation (ii) and (iii),

$$\frac{k_x}{k_z} \left( \frac{\partial x_t}{\partial x} \right)^2 = 1$$

$$\frac{\partial x_t}{\partial x} = \sqrt{\frac{k_z}{k_x}}$$

$$\partial x_t = \sqrt{\frac{k_z}{k_x}} \partial x$$

$$x_t = \sqrt{\frac{k_z}{k_x}} x + \text{Constant}$$

Scan for Video solution

## CHAPTER

# 5

# VECTOR CALCULUS

### Vector Algebra

- 1. [NAT] [GATE-ME-2022: 2M]**

Consider two vectors:

$$\vec{a} = 5\hat{i} + 7\hat{j} + 2\hat{k}$$

$$\vec{b} = 3\hat{i} - \hat{j} + 6\hat{k}$$

Magnitude of the component of  $\vec{a}$  orthogonal to  $\vec{b}$  in the plane containing the vectors  $\vec{a}$  and  $\vec{b}$  is \_\_\_\_\_.(Round off to 2 decimal places)

- 2. [MCQ] [GATE-2021-CE:2M]**

The unit normal vector to the surface  $X^2 + Y^2 + Z^2 - 48 = 0$  at the point (4,4,4) is

(a)  $\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}$

(b)  $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$

(c)  $\frac{2}{\sqrt{2}}, \frac{2}{\sqrt{2}}, \frac{2}{\sqrt{2}}$

(d)  $\frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}, \frac{1}{\sqrt{5}}$

- 3. [MCQ] [GATE-2021-CE:1M]**

Let  $\phi$  be a scalar field, and  $\mathbf{u}$  be a vector field. Which of the following identities is true for  $\operatorname{div}(\phi\mathbf{u})$ ?

(a)  $\operatorname{div}(\phi\mathbf{u})=\phi\operatorname{div}(\mathbf{u})+\mathbf{u}\cdot\operatorname{grad}(\phi)$

(b)  $\operatorname{div}(\phi\mathbf{u})=\phi\operatorname{div}(\mathbf{u})+\mathbf{u}\times\operatorname{grad}(\phi)$

(c)  $\operatorname{div}(\phi\mathbf{u})=\phi\operatorname{grad}(\mathbf{u})+\mathbf{u}\cdot\operatorname{grad}(\phi)$

(d)  $\operatorname{div}(\phi\mathbf{u})=\phi\operatorname{grad}(\mathbf{u})+\mathbf{u}\times\operatorname{grad}(\phi)$

- 4. [NAT] [GATE-ME-2020: 1M]**

For three vectors

$$\vec{A} = 2\hat{j} - 3\hat{k}, \vec{B} = -2\hat{i} + \hat{k} \text{ and } \vec{C} = 3\hat{i} - \hat{j},$$

$\hat{i}, \hat{j}$  and  $\hat{k}$  are unit vectors along the axes of a right-handed rectangular /Cartesian coordinate system, the value of  $(\vec{A} \cdot (\vec{B} \times \vec{C})) + 6$  is \_\_\_\_.

- 5. [MCQ] [GATE-EE-2020: 2M]**

The vector function expressed by  $\mathbf{F} = a_x(5y - k_1 z) + a_y(3z + k_2 x) + a_z(k_3 y - 4x)$  represents a conservative field, where  $a_x, a_y, a_z$  are unit vectors along x, y and z directions, respectively. The values of constants  $k_1, k_2, k_3$  are given by:

(a)  $k_1=3, k_2=3, k_3=7$

(b)  $k_1=3, k_2=8, k_3=5$

(c)  $k_1=4, k_2=5, k_3=3$

(d)  $k_1=0, k_2=0, k_3=0$

- 6. [MCQ] [GATE-EC-2020: 1M]**

For a vector field  $\vec{A}$ , which one of the following is FALSE?

(a)  $\vec{A}$  is solenoidal if  $\nabla \cdot \vec{A} = 0$ .

(b)  $\nabla \times \vec{A}$  is another vector field.

(c)  $\vec{A}$  is irrotational if  $\nabla^2 \vec{A} = 0$ .

(d)  $\nabla \times (\nabla \times \vec{A}) = \nabla (\nabla \cdot \vec{A}) - \nabla^2 \vec{A}$ .

- 7. [NAT] [GATE-EE-2019 : 2M]**

If  $A = 2x\hat{i} + 3y\hat{j} + 4z\hat{k}$  and  $u = x^2 + y^2 + z^2$ , then  $\operatorname{div}(u\mathbf{A})$  at(1, 1, 1) is \_\_\_\_\_

- 8. [MCQ] [GATE-2019-CE:1M]**

Euclidean norm (length) of the vector  $[4 -2 -6]^T$  is

(a)  $\sqrt{48}$

(b)  $\sqrt{56}$

(c)  $\sqrt{24}$

(d)  $\sqrt{12}$

- 9. [MCQ] [GATE-2019-CE:1M]**

What is curl of the vector field  $2x^2y\hat{i} + 5z^2\hat{j} - 4yz\hat{k}$ ?

(a)  $6zi + 4xj - 2x^2k$

(b)  $6zi - 8xyj + 2x_2k$

(c)  $-14zi + 6yj + 2x^2k$

(d)  $-14zi - 2x^2k$

**10. [MCQ]****[GATE-CS-2017:2M]**

Let  $u$  and  $v$  be two vectors in  $\mathbb{R}^2$  whose Euclidean norms satisfy  $\|u\| = 2\|v\|$ . What is the value of  $a$  such that  $w = u + av$  bisects the angle between  $u$  and  $v$ ?


- (a) 2
- (b)  $\frac{1}{2}$
- (c) 1
- (d)  $-\frac{1}{2}$

**11. [NAT]****[GATE-2017-CE:1M]**

The divergence of the vector field  $\mathbf{V} = x^2\mathbf{i} + 2y^3\mathbf{j} + z^4\mathbf{k}$  at  $x=1$ ,  $y=2$ ,  $z=3$  is \_\_\_\_\_.

**12. [MCQ]****[GATE-EE-2017: 1M]**

The figures show diagrammatic representations of vector fields,  $\vec{X}$ ,  $\vec{Y}$  and  $\vec{Z}$ , respectively. Which one of the following choices is true?



- (a)  $\nabla \cdot \vec{X} = 0$ ,  $\nabla \times \vec{Y} \neq 0$ ,  $\nabla \times \vec{Z} = 0$
- (b)  $\nabla \cdot \vec{X} \neq 0$ ,  $\nabla \times \vec{Y} = 0$ ,  $\nabla \times \vec{Z} \neq 0$
- (c)  $\nabla \cdot \vec{X} \neq 0$ ,  $\nabla \times \vec{Y} \neq 0$ ,  $\nabla \times \vec{Z} \neq 0$
- (d)  $\nabla \cdot \vec{X} = 0$ ,  $\nabla \times \vec{Y} = 0$ ,  $\nabla \times \vec{Z} = 0$

**13. [NAT]****[GATE-EC-2016: 2M]**

Consider the time-varying vector  $I = \hat{x}15 \cos(\omega t) + \hat{y}5 \sin(\omega t)$  in Cartesian coordinates, where  $\omega > 0$  is a constant. When the vector magnitude  $|I|$  is at its minimum value, the angle  $\theta$  that  $I$  makes with the  $x$ -axis (in degree, such that  $0 \leq \theta \leq 180$ ) is \_\_\_\_\_.

**14. [MCQ]****[GATE-ME-2014: 1M]**

Which one of the following describes the relationship among the three vectors,  $\hat{i} + \hat{j} + \hat{k}$ ,  $2\hat{i} + 3\hat{j} + \hat{k}$  and  $5\hat{i} + 6\hat{j} + 4\hat{k}$ ?

- (a) The vectors are mutually perpendicular
- (b) The vectors are linearly dependent
- (c) The vectors are linearly independent
- (d) The vectors are unit vectors

**15. [NAT]****[GATE-EC-2014: 1M]**

The magnitude of the gradient for the function  $f(x,y,z) = x^2 + 3y^2 + z^3$  at the point  $(1, 1, 1)$  is \_\_\_\_\_.

**16. [MCQ]****[GATE-EE-2013 :1M]**

The curl of the gradient of the scalar field defined by  $\mathbf{V} = 2x^2\mathbf{y} + 3y^2\mathbf{z} + 4z^2\mathbf{x}$  is

- (a)  $4xy\mathbf{a}_x + 6yz\mathbf{a}_y + 8zx\mathbf{a}_z$
- (b)  $4\mathbf{a}_x + 6\mathbf{a}_y + 8\mathbf{a}_z$
- (c)  $(4xy + 4z^2)\mathbf{a}_x + (2x^2 + 6yz)\mathbf{a}_y + (3y^2 + 8zx)\mathbf{a}_z$
- (d) 0

**17. [MCQ]****[GATE-EC-2013: 1M]**

Consider a vector field  $\vec{A}(\vec{r})$ . The closed loop line integral  $\oint \vec{A} \cdot d\vec{l}$  can be expressed as

- (a)  $\iint (\nabla \times \vec{A}) \cdot \vec{ds}$  over the closed surface bounded by the loop
- (b)  $\iiint (\nabla \cdot \vec{A}) dv$  over the closed volume bounded by the loop
- (c)  $\iiint (\nabla \cdot \vec{A}) dv$  over the open volume bounded by the loop
- (d)  $\iint (\nabla \times \vec{A}) \cdot \vec{ds}$  over the open surface bounded by the loop


**18. [MCQ]****[GATE-EC-2013: 1M]**

The divergence of the vector field  $\vec{A} = x\hat{a}_x + y\hat{a}_y + z\hat{a}_z$  is

- (a) 0
- (b) 1/3
- (c) 1
- (d) 3

**19. [MCQ]****[GATE-2012-CE:2M]**

For the parallelogram OPQR shown in the sketch  $\overrightarrow{OP} = a\hat{i} + b\hat{j}$  and  $\overrightarrow{OR} = c\hat{i} + d\hat{j}$ . The area of the parallelogram is



- (a)  $ad - bc$
- (b)  $ac + bd$
- (c)  $ad + bc$
- (d)  $ab - cd$

**20. [MCQ]****[GATE-2011-CE:2M]**

If  $\vec{a}$  and  $\vec{b}$  are two arbitrary vectors with magnitudes  $a$  and  $b$ , respectively,  $|\vec{a} \times \vec{b}|^2$  will be equal to

- (a)  $a^2 b^2 - (\vec{a} \cdot \vec{b})^2$       (b)  $ab - \vec{a} \cdot \vec{b}$   
 (c)  $a^2 b^2 + (\vec{a} \cdot \vec{b})^2$       (d)  $ab + \vec{a} \cdot \vec{b}$

**21. [MCQ]****[GATE-EE-2010 :1M]**

Divergence of the three-dimensional radial vector field  $\vec{r}$  is

- (a) 3      (b)  $1/r$   
 (c)  $\hat{i} + \hat{j} + \hat{k}$       (d)  $3(\hat{i} + \hat{j} + \hat{k})$

**22. [MCQ]****[GATE-2009-CE:1M]**

For a scalar function  $f(x, y, z) = x^2 + 3y^2 + 2z^2$ , the gradient at the point P (1, 2, -1) is

- (a)  $2\vec{i} + 6\vec{j} + 4\vec{k}$       (b)  $2\vec{i} + 12\vec{j} - 4\vec{k}$   
 (c)  $2\vec{i} + 12\vec{j} + 4\vec{k}$       (d)  $\sqrt{56}$

**23. [MCQ]****[GATE-EC-2009: 1M]**

If a vector field  $\vec{V}$  is related to another field  $\vec{A}$  through  $\vec{V} = \nabla \times \vec{A}$ , which of the following is true?

**Note:** C and  $S_C$  refer to any closed contour and any surface whose boundary is C.

- (a)  $\oint_C \vec{V} \cdot d\vec{l} = \int_{S_C} \int \vec{A} \cdot \vec{ds}$   
 (b)  $\oint_C \vec{A} \cdot d\vec{l} = \int_{S_C} \int \vec{V} \cdot \vec{ds}$   
 (c)  $\oint_C \nabla \times \vec{V} \cdot d\vec{l} = \int_{S_C} \int \nabla \times \vec{A} \cdot \vec{ds}$   
 (d)  $\oint_C \nabla \times \vec{A} \cdot d\vec{l} = \int_{S_C} \int \vec{V} \cdot \vec{ds}$

**24. [MCQ]****[GATE-2008-CE:2M]**

The inner (dot) product of two vectors  $\vec{P}$  and  $\vec{Q}$  is zero.

The angle (degrees) between the two vectors is

- (a) 0      (b) 30  
 (c) 90      (d) 120

**Vector Differentiation****25. [MCQ]****[GATE-EC-2023: 1M]**

The rate of increase, of a scalar field  $f(x, y, z) = xyz$ , in the direction  $v = (2, 1, 2)$  at a point (0,2,1) is

- (a)  $\frac{2}{3}$       (b)  $\frac{4}{3}$   
 (c) 2      (d) 4

**26. [MCQ]****[GATE-EE-2022 :2M]**

Let,  $f(x, y, z) = 4x^2 + 7xy + 3xz^2$ . The direction in which the function  $f(x, y, z)$  increases most rapidly at point P = (1, 0, 2) is

- (a)  $20\hat{i} + 7\hat{j}$       (b)  $20\hat{i} + 7\hat{j} + 12\hat{k}$   
 (c)  $20\hat{i} + 12\hat{k}$       (d)  $20\hat{i}$

**27. [NAT]****[GATE-2021-CE:2M]**

A function is defined in Cartesian coordinate system as  $f(x, y) = xe^y$ . The value of the directional derivative of the function (in integer) at the point (2, 0) along the direction of the straight-line segment from point (2,0) to point  $(\frac{1}{2}, 2)$  is \_\_\_\_\_.

**28. [MCQ]****[GATE-ME-2020: 2M]**

The directional derivative of  $f(x, y, z) = xyz$  at point (-1, 1, 3) in the direction of vector  $\hat{i} - 2\hat{j} + 2\hat{k}$  is

- (a)  $3\hat{i} - 3\hat{j} - \hat{k}$       (b)  $-\frac{7}{3}$   
 (c)  $\frac{7}{3}$       (d) 7

**29. [MCQ]****[GATE-ME-2019: 1M]**

The directional derivative of the function  $f(x, y) = x^2 + y^2$  along a line directed from (0, 0) to (1, 1), evaluated at the point x = 1, y = 1 is

- (a)  $\sqrt{2}$       (b) 2  
 (c)  $2\sqrt{2}$       (d)  $4\sqrt{2}$

**30. [MCQ]****[GATE-ME-2018: 1M]**

The divergence of the vector field  $\vec{u} = e^x (\cos y \hat{i} + \sin y \hat{j})$

- is  
 (a) 0      (b)  $e^x \cos y + e^x \sin y$   
 (c)  $2e^x \cos y$       (d)  $2e^x \sin y$

**31. [MCQ]****[GATE-ME-2018: 2M]**

For a position vector  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  the norm of the vector can be defined as  $|\vec{r}| = \sqrt{x^2 + y^2 + z^2}$ . Given a function  $\phi = \ln |\vec{r}|$ , its gradient  $\nabla \phi$  is

- (a)  $\vec{r}$       (b)  $\frac{\vec{r}}{|\vec{r}|}$   
 (c)  $\frac{\vec{r}}{\vec{r} \cdot \vec{r}}$       (d)  $\frac{\vec{r}}{|\vec{r}|^3}$



**45. [MCQ]****[GATE-ME-2012: 1M]**

For the spherical surface  $x^2 + y^2 + z^2 = 1$ , the unit outward normal vector at the point  $\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0\right)$  is given by

(a)  $\frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{j}$

(b)  $\frac{1}{\sqrt{2}}\hat{i} - \frac{1}{\sqrt{2}}\hat{j}$

(c)  $\hat{k}$

(d)  $\frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} + \frac{1}{\sqrt{3}}\hat{k}$

**46. [MCQ]****[GATE-EE-2012 : 2M]**

The direction of vector A is radially outward from the origin, with  $|A| = Kr^n$  where  $r^2 = x^2 + y^2 + z^2$  and K is constant. The value of n for which  $\nabla \cdot A = 0$  is

(a) -2

(b) 2

(c) 1

(d) 0

**47. [MCQ]****[GATE-2009-CE:2M]**

For a scalar function  $f(x, y, z) = x^2 + 3y^2 + 2z^2$ , the directional derivative at the point P (1, 2, -1) in the direction of a vector  $\vec{i} - \vec{j} + 2\vec{k}$  is

(a) -18

(b)  $-3\sqrt{6}$

(c)  $3\sqrt{6}$

(d) 18

**48. [MCQ]****[GATE-ME-2009: 1M]**

The divergence of the vector field  $3xz\hat{i} + 2xy\hat{j} - yz^2\hat{k}$  at a point (1, 1, 1) is equal to

(a) 7

(b) 4

(c) 3

(d) 0

**49. [MCQ]****[GATE-ME-2008: 1M]**

The divergence of the vector field

$(x-y)\hat{i} + (y-x)\hat{j} + (x+y+z)\hat{k}$  is

(a) 0

(b) 1

(c) 2

(d) 3

**50. [MCQ]****[GATE-ME-2008: 2M]**

The directional derivative of the scalar function  $f(x, y, z) = x^2 + 2y^2 + z$  at the point P = (1, 1, 2) in the direction of the vector  $\vec{a} = 3\hat{i} - 4\hat{j}$  is

(a) -4

(b) -2


(c) -1

(d) 1

**Vector Integration****51. [NAT]****[GATE-EE-2023 : 2M]**

The closed curve shown in the figure is described by  $r = 1 + \cos \theta$ , where  $r = \sqrt{x^2 + y^2}$ ;  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

The magnitude of the line integral of the vector field  $\mathbf{F} = -y\hat{i} + x\hat{j}$  around the closed is \_\_\_\_\_ (round off to 2 decimal places).

**52. [NAT]****[GATE-EE-2023 : 2M]**

Three points in the x-y plane are (-1, 0.8), (0, 2.2) and (1, 2.8). The value of the slope of the best fit straight in the least square sense is \_\_\_\_\_ (Round off to 2 decimal places)

**53. [MCQ]****[GATE-EC-2023: 2M]**

The value of the line integral  $\int_P^Q (z^2 dx + 3y^2 dy + 2xz dz)$  along the straight line joining the points P (1, 1, 2) and Q (2, 3, 1) is

(a) 20

(b) 24

(c) 29

(d) -5

**54. [NAT]****[GATE-ME-2023: 1M]**


A vector field

$\vec{B}(x, y, z) = x\hat{i} + y\hat{j} - 2z\hat{k}$

is defined over a conical region having  $h = 2$ , base radius  $r = 3$  and axis along z, as shown in the figure. The base of the cone lies in the x-y plane and is centred at the origin. If  $\hat{n}$  denotes the unit outward normal to the curved surface S of the cone, the value of the integral

$$\int_S \vec{B} \cdot \hat{n} ds$$

equals \_\_\_\_\_. (Answer in integer)



**55. [MCQ]****[GATE-ME-2022: 1M]**

Given a function  $\varphi = \frac{1}{2}(x^2 + y^2 + z^2)$  in three-dimensional Cartesian space, the value of the surface integral  $\iint_s \hat{n} \cdot \nabla \varphi d\sigma$ , where  $s$  is the surface of a sphere

of unit radius and  $\hat{n}$  is the outward unit normal vector on  $s$ , is

- (a)  $4\pi$       (b)  $3\pi$   
 (c)  $4\pi/3$       (d)  $0$

**56. [MCQ]****[GATE-ME-2022: 1M]**

Consider a cube of unit edge length and sides parallel to co-ordinate axes, with its centroid at the point


(1, 2, 3). The surface integral  $\iint_A \vec{F} \cdot d\vec{A}$  of a vector field

$\vec{F} = 3x\hat{i} + 5y\hat{j} + 6z\hat{k}$  over the entire surface  $A$  of the cube is

- (a) 14      (b) 27  
 (c) 28      (d) 31

**57. [MCQ]****[GATE-EC-2022: 1M]**

Consider the two-dimensional vector field  $\vec{F}(x, y) = x\vec{i} + y\vec{j}$ , where  $\vec{i}$  and  $\vec{j}$  denote the unit vectors along the x-axis and the y-axis, respectively. A contour  $C$  in the x-y plane, as shown in the figure, is composed of two horizontal lines connected at the two ends by two semi-circular arcs of unit radius. The contour is traversed in the counter-clockwise sense. The value of the closed path integral  $\oint_C \vec{F}(x, y) \cdot (dx\vec{i} + dy\vec{j})$  is \_\_\_\_\_.




- (a) 0      (b) 1  
 (c)  $8 + 2\pi$       (d)  $-1$

**58. [MCQ]****[GATE-EE-2022: 2M]**


Let  $R$  be a region in the first quadrant of the  $xy$  plane enclosed by a closed curve  $C$  considered in counter-clockwise direction. Which of the following expressions does not represent the area of the region  $R$ ?

- (a)  $\iint_R dxdy$       (b)  $\int_C xdy$   
 (c)  $\int_C ydx$       (d)  $\frac{1}{2} \int_C (xdy - ydx)$

**59. [NAT]****[GATE-EE-2020 : 2M]**

Let  $\mathbf{a}_x$  and  $\mathbf{a}_y$  be unit vectors along  $x$  and  $y$  directions, respectively. A vector function is given by  $\mathbf{F} = \mathbf{a}_x y - \mathbf{a}_y x$ . The line integral of the above function  $\int_C \mathbf{F} \cdot d\mathbf{l}$  along the curve  $C$ , which follows the parabola

$y = x^2$  as shown below is \_\_\_\_\_ (rounded off to 2 decimal places).

**60. [NAT]****[GATE-2020-CE:2M]**

If  $C$  represents a line segment between  $(0,0,0)$  and  $(1,1,1)$  in Cartesian coordinate system, the value (expressed as integer) of the line integral  $\int_C [(y+z)dx + (x+z)dy + (x+y)dz]$  is \_\_\_\_\_.

**61. [MCQ]****[GATE-ME-2020: 2M]**

A vector field is defined as

$$\vec{f}(x, y, z) = \frac{x}{[x^2 + y^2 + z^2]^{3/2}} \hat{i} + \frac{y}{[x^2 + y^2 + z^2]^{3/2}} \hat{j} + \frac{z}{[x^2 + y^2 + z^2]^{3/2}} \hat{k}$$

Where,  $\hat{i}, \hat{j}, \hat{k}$  are unit vector along the axes of a right-handed rectangular/cartesian coordinate system. The surface integral  $\iint \vec{f} \cdot d\vec{s}$  (where  $d\vec{s}$  is an elemental surface area vector) evaluated over the inner and outer surfaces of a spherical shell formed by two concentric spheres with origin as the centre, and internal and external radii of 1 and 2, respectively is




**62. [NAT]**

[GATE-EC-2020: 2M]

For a solid S shown below, the value of  $\iiint_S x \cdot dx dy dz$

(rounded off to two decimal places) is \_\_\_\_\_.



**63. [NAT]**

[GATE-EE-2019 : 1M]

If  $f = 2x^3 + 3y^2 + 4z$ , the value of line integral  $\int_C \text{grad}(f) \cdot d\mathbf{r}$

.  $\mathbf{dr}$  evaluated over contour C formed by the segments  $(-3, -3, 2) \rightarrow (2, -3, 2) \rightarrow (2, 6, 2) \rightarrow (2, 6, -1)$  is \_\_\_\_\_.

**64. [MCQ]**

[GATE-ME-2019: 2M]


Given a vector  $\vec{u} = \frac{1}{3}(-y^3\hat{i} + x^3\hat{j} + z^3\hat{k})$  and  $\hat{n}$  is the unit normal vector to the surface of the hemisphere  $(x^2 + y^2 + z^2 = 1; z \geq 0)$ , the value of integral  $\int (\nabla \times \vec{u}) \cdot \hat{n} dS$  evaluated on the curved surface of the hemisphere S is

- (a)  $-\frac{\pi}{2}$       (b)  $\frac{\pi}{3}$   
 (c)  $\frac{\pi}{2}$       (d)  $\pi$

**65. [MCQ]**

[GATE-EC-2019: 2M]

Consider the line integral  $\int_C (xdy - ydx)$  the integral being taken in a counter clockwise direction over the closed curve C that forms the boundary of the region R shown in the figure below. The region R is the area enclosed by the union of a  $2 \times 3$  rectangle and a semi-circle of radius 1. The line integral evaluates to



- (a)  $6 + \frac{\pi}{2}$       (b)  $16 + 2\pi$   
 (c)  $8 + \pi$       (d)  $12 + \pi$

**66. [NAT]**


[GATE-2018-CE:2M]

The value (up to two decimal places) of a line integral  $\int_C \vec{F}(\vec{r}) \cdot d\vec{r}$  for  $\vec{F}(\vec{r}) = x^2\hat{i} + y^2\hat{j}$  along C which is a straight line joining (0, 0) to (1, 1) is \_\_\_\_\_.

67. [NAT]

[GATE-EE-2018 : 2M]

As shown in the figure, C is the arc from the point (3,0) to the point (0,3) on the circle  $x^2 + y^2 = 9$ . The value of integral  $\int_C (y^2 + 2yx)dx + (2xy + x^2)dy$  is \_\_\_\_\_ (up to 2 decimal places).



**68. [MCQ]****[GATE-ME-2018: 2M]**

The value of the integral

$$\iint_S \vec{r} \cdot \vec{n} \, dS$$

over the closed surface S bounding a volume V, where  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  is the position vector and  $\vec{n}$  is the normal to the surface S, is

- |        |        |
|--------|--------|
| (a) V  | (b) 2V |
| (c) 3V | (d) 4V |

**69. [NAT]****[GATE-ME-2017: 2M]**

The surface integral  $\iint_S \vec{F} \cdot \hat{n} \, dS$  over the surface S of the sphere  $x^2 + y^2 + z^2 = 9$ , where  $\vec{F} = (x+y)\hat{i} + (x+z)\hat{j} + (y+z)\hat{k}$  and  $\hat{n}$  is the unit outward surface normal, yields \_\_\_\_.

**70. [NAT]****[GATE-EC-2017: 2M]**

Let  $I = \int_C (2zdx + 2ydy + 2xdz)$  where x, y, z are real, and let C be the straight-line segment from point A:(0, 2, 1) to point B:(4, 1, -1). The value of I is \_\_\_\_.

**71. [MCQ]****[GATE-EE-2016 : 1M]**

The value of the line integral  $\int (2xy^2 dx + 2x^2 y dy + dz)$  along a path joining the origin (0, 0, 0) and the point (1, 1, 1) is

- |       |       |
|-------|-------|
| (a) 0 | (b) 2 |
| (c) 4 | (d) 6 |

**72. [NAT]****[GATE-EE-2016 :2M]**

The line integral of the vector field  $\vec{F} = 5xz\hat{i} + (3x^2 + 2y)\hat{j} + x^2 z\hat{k}$  along a path from (0, 0, 0) to (1, 1, 1) parameterized by  $(t, t^2, t)$  is \_\_\_\_.

**73. [NAT]****[GATE-EC-2016: 2M]**

Suppose C is the closed curve defined as the circle  $x^2 + y^2 = 1$  with C oriented anti-clockwise. The value of  $\oint (xy^2 dx + x^2 y dy)$  over the curve C equals \_\_\_\_.

**74. [NAT]****[GATE-ME-2016: 2M]**A scalar potential  $\phi$  has the following gradient:

$\nabla \phi = yz\hat{i} + xz\hat{j} + xy\hat{k}$ . Consider the integral  $\int_C \nabla \phi \cdot d\vec{r}$  on

the curve  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ .

The curve C is parameterized as follows:

$$\begin{cases} x = t \\ y = t^2 \\ z = 3t^2 \end{cases} \quad \text{and } 1 \leq t \leq 3$$

The value of the integral is \_\_\_\_.

**75. [NAT]****[GATE-ME-2016: 2M]**The value of the line integral  $\oint_C \vec{F} \cdot \vec{r}' ds$ , where C is acircle of radius  $\frac{4}{\sqrt{\pi}}$  is \_\_\_\_.

Here  $\vec{F}(x, y) = y\hat{i} + 2x\hat{j}$  and  $\vec{r}'$  is the UNIT tangent vector on the curve C at an arc length s from a reference point on the curve.  $\hat{i}$  and  $\hat{j}$  are the basis vectors in the x-y Cartesian reference. In evaluating the line integral, the curve has to be traversed in the counter-clockwise direction.

**76. [NAT]****[GATE-ME-2015: 2M]**

The surface integral  $\iint_S \frac{1}{\pi} (9x\hat{i} - 3y\hat{j}) \cdot \hat{n} \, dS$  over the sphere given by  $x^2 + y^2 + z^2 = 9$  is \_\_\_\_.

**77. [NAT]****[GATE-ME-2015: 2M]**

The value of

$$\int_C [(3x - 8y^2)dx + (4y - 6xy)dy],$$

(Where C is the boundary of the region bounded by  $x = 0$ ,  $y = 0$  and  $x + y = 1$ ) is \_\_\_\_.

**78. [MCQ]****[GATE-ME-2014: 2M]**

The integral  $\oint_C (ydx - xdy)$  is evaluated along the

circle  $x^2 + y^2 = \frac{1}{4}$  traversed in counter clockwise direction. The integral is equal to

- |                      |                      |
|----------------------|----------------------|
| (a) 0                | (b) $-\frac{\pi}{4}$ |
| (c) $-\frac{\pi}{2}$ | (d) $\frac{\pi}{4}$  |

**79. [MCQ]****[GATE-EE-2013: 1M]**

Given a vector field  $\vec{F} = y^2 x \hat{a}_x - y z \hat{a}_y - x^2 \hat{a}_z$ , the line integral  $\int_C \vec{F} \cdot d\vec{r}$  evaluated along a segment on the x-axis from  $x = 1$  to  $x = 2$  is

- |           |       |
|-----------|-------|
| (a) -2.33 | (b) 0 |
| (c) 2.33  | (d) 7 |

**80. [MCQ]****[GATE-ME-2013: 2M]**

The following surface integral is to be evaluated over a sphere for the given steady velocity vector field  $\vec{F} = x\hat{i} + y\hat{j} + z\hat{k}$  defined with respect to a Cartesian coordinate system having  $\hat{i}, \hat{j}$  and  $\hat{k}$  as unit base vectors.

$$\iint_S \frac{1}{4} (\vec{F} \cdot \hat{n}) dA$$

where  $S$  is the sphere,  $x^2 + y^2 + z^2 = 1$  and  $\hat{n}$  is the outward unit normal vector to the sphere. The value of the surface integral is

- |              |            |
|--------------|------------|
| (a) $\pi$    | (b) $2\pi$ |
| (c) $3\pi/4$ | (d) $4\pi$ |

**81. [MCQ]****[GATE-EC-2011:1M]**


Consider a closed surface  $s$  surrounding a volume  $v$ . If  $\vec{r}$  is the position vector of a point inside  $s$  with  $\hat{n}$  the unit normal on  $s$ , the value of the integral  $\iint_s 5\vec{r} \cdot \hat{n} ds$  is

- |           |           |
|-----------|-----------|
| (a) $3v$  | (b) $5v$  |
| (c) $10v$ | (d) $15v$ |

**82. [MCQ]****[GATE-EC-2010: 2M]**

If  $\vec{A} = xy\hat{x} + x^2\hat{y}$  then  $\oint_C \vec{A} \cdot d\vec{l}$  over the path shown

in the figure is



- |       |                          |
|-------|--------------------------|
| (a) 0 | (b) $\frac{2}{\sqrt{3}}$ |
| (c) 1 | (d) $2\sqrt{3}$          |

**83. [MCQ]****[GATE-EE-2009: 2M]**

$F(x, y) = (x^2 + xy)\hat{x} + (y^2 + xy)\hat{y}$ . It's line integral over the straight line from  $(x, y) = (0, 2)$  to  $(2, 0)$  evaluates to

- |        |       |
|--------|-------|
| (a) -8 | (b) 4 |
| (c) 8  | (d) 0 |

**84. [MCQ]****[GATE-EC-2008: 2M]**

Consider points P and Q in the x-y plane, with  $P = (1, 0)$  and  $Q = (0, 1)$ . The line integral

$$2 \int_P^Q (xdx + ydy)$$

along the semicircle with the line segment PQ as its diameter

- |  |
|--|
| (a) is -1  |
| (b) is 0   |
| (c) is 1   |
| (d) depends on the direction (clockwise or anti clockwise) of the semicircle |

**85. [MCQ]****[GATE-EC-2008: 2M]**

The value of the integral of the function

$g(x, y) = 4x^3 + 10y^4$  along the straight-line segment from the point  $(0,0)$  to the point  $(1, 2)$  in the x-y plane is

- |        |        |
|--------|--------|
| (a) 33 | (b) 35 |
| (c) 40 | (d) 56 |



 ANSWER KEY


- |                            |                              |                             |                           |
|----------------------------|------------------------------|-----------------------------|---------------------------|
| <b>1.</b> (7.90 to 8.70)   | <b>2.</b> (a)                | <b>3.</b> (a)               | <b>4.</b> (6 to 6)        |
| <b>5.</b> (c)              | <b>6.</b> (c)                | <b>7.</b> (45 to 45)        | <b>8.</b> (b)             |
| <b>9.</b> (d)              | <b>10.</b> (a)               | <b>11.</b> (133.9 to 134.1) | <b>12.</b> (c)            |
| <b>13.</b> (90 to 90)      | <b>14.</b> (b)               | <b>15.</b> (6.8 to 7.2)     | <b>16.</b> (d)            |
| <b>17.</b> (d)             | <b>18.</b> (d)               | <b>19.</b> (a)              | <b>20.</b> (a)            |
| <b>21.</b> (a)             | <b>22.</b> (b)               | <b>23.</b> (b)              | <b>24.</b> (c)            |
| <b>25.</b> (b)             | <b>26.</b> (b)               | <b>27.</b> (1 to 1)         | <b>28.</b> (c)            |
| <b>29.</b> (c)             | <b>30.</b> (c)               | <b>31.</b> (c)              | <b>32.</b> (a)            |
| <b>33.</b> (0 to 0)        | <b>34.</b> (b)               | <b>35.</b> (0 to 0)         | <b>36.</b> (a)            |
| <b>37.</b> (c)             | <b>38.</b> (-5.72 to -5.70)  | <b>39.</b> (a)              | <b>40.</b> (2.99 to 3.01) |
| <b>41.</b> (2.9 to 3.1)    | <b>42.</b> (a)               | <b>43.</b> (c)              | <b>44.</b> (a)            |
| <b>45.</b> (a)             | <b>46.</b> (a)               | <b>47.</b> (b)              | <b>48.</b> (c)            |
| <b>49.</b> (d)             | <b>50.</b> (b)               | <b>51.</b> (9 to 10)        | <b>52.</b> (0.9 to 1.1)   |
| <b>53.</b> (b)             | <b>54.</b> (-0.001 to 0.001) | <b>55.</b> (a)              | <b>56.</b> (a)            |
| <b>57.</b> (a)             | <b>58.</b> (a)               | <b>59.</b> (-3.05 to -2.95) | <b>60.</b> (3 to 3)       |
| <b>61.</b> (a)             | <b>62.</b> (2.25 to 2.25)    | <b>63.</b> (139 to 139)     | <b>64.</b> (c)            |
| <b>65.</b> (d)             | <b>66.</b> (0.60 to 0.70)    | <b>67.</b> (0 to 0)         | <b>68.</b> (c)            |
| <b>69.</b> (225 to 227)    | <b>70.</b> (-11.1 to -10.9)  | <b>71.</b> (b)              | <b>72.</b> (4.40 to 4.45) |
| <b>73.</b> (-0.03 to 0.03) | <b>74.</b> (725.9 to 726.1)  | <b>75.</b> (15.9 to 16.1)   | <b>76.</b> (214 to 218)   |
| <b>77.</b> (1.60 to 1.70)  | <b>78.</b> (c)               | <b>79.</b> (b)              | <b>80.</b> (a)            |
| <b>81.</b> (d)             | <b>82.</b> (c)               | <b>83.</b> (d)              | <b>84.</b> (b)            |
| <b>85.</b> (a)             |                              |                             |                           |


**SOLUTIONS**
**1. (7.90 to 8.70)**

**Given**  $\vec{a} = 5\hat{i} + 7\hat{j} + 2\hat{k}$   
 $\vec{b} = 3\hat{i} - \hat{j} + 6\hat{k}$

Component of  $\vec{a}$  orthogonal to  $\vec{b}$

$$|\vec{c}| = |\vec{a}| \cdot \sin \theta = |\vec{a}| \cdot \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| \cdot |\vec{b}|} = \frac{|\vec{a} \times \vec{b}|}{|\vec{b}|}$$



$$\therefore |\vec{a} \times \vec{b}| = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & 7 & 2 \\ 3 & -1 & 6 \end{vmatrix} = \hat{i}(44) - \hat{j}(24) + \hat{k}(-26)$$

$$|\vec{b}| = \sqrt{3^2 + (-1)^2 + 6^2} = \sqrt{46}$$

$$\therefore |\vec{a} \times \vec{b}| = \sqrt{44^2 + 24^2 + 26^2} = 56.46$$

$$\therefore \text{component of } \vec{a} \text{ orthogonal to } \vec{b} = \frac{56.46}{\sqrt{46}} = 8.325$$



Scan for Video solution

**2. (a)**

Let the surface be  $\phi = x^2 + y^2 + z^2 - 48$

$$\text{Grad } \phi = \frac{\partial \phi}{\partial x} \hat{i} + \frac{\partial \phi}{\partial y} \hat{j} + \frac{\partial \phi}{\partial z} \hat{k}$$

$$\text{Grad } \phi = (2x)\hat{i} + (2y)\hat{j} + (2z)\hat{k}$$


$$\text{At P (4, 4, 4) grad } \phi = 8\hat{i} + 8\hat{j} + 8\hat{k}$$

$$\therefore \text{Normal vector } \vec{n} = 8\hat{i} + 8\hat{j} + 8\hat{k}$$

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{8\hat{i} + 8\hat{j} + 8\hat{k}}{\sqrt{8^2 + 8^2 + 8^2}}$$

$$\therefore \text{Unit normal vector; } \hat{n} = \frac{\hat{i}}{\sqrt{3}} + \frac{\hat{j}}{\sqrt{3}} + \frac{\hat{k}}{\sqrt{3}}$$


$$\left( \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right)$$

**3. (a)**

For any Scalar  $\phi$  and vector  $\mathbf{u}$ ,

$$\text{div}(\phi \cdot \mathbf{u}) = \phi \text{ div } \mathbf{u} + \mathbf{u} \cdot \text{grad } \phi$$

The above result is obtained by applying the product rule of differentiation.

**4. (6 to 6)**

**Given**


$$\vec{A} = 2\hat{j} - 3\hat{k}, \vec{B} = -2\hat{i} - \hat{k}, \vec{C} = 3\hat{i} - \hat{j}$$

$$\vec{A} \cdot (\vec{B} \times \vec{C}) = (2\hat{j} - 3\hat{k}) \cdot \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -2 & 0 & 1 \\ 3 & -1 & 0 \end{vmatrix}$$

$$= (2\hat{j} - 3\hat{k}) \cdot (\hat{i}(1) - \hat{j}(-3) + \hat{k}(2)) = 6 - 6 = 0$$

$$\therefore \vec{A} \cdot (\vec{B} \times \vec{C}) = 0$$

$$\therefore \vec{A} \cdot (\vec{B} \times \vec{C}) + 6 = 0 + 6 = 6$$

**5. (c)**

$$\vec{F} = \hat{a}_x(5y - k_1z) + \hat{a}_y(3z + k_2x) + \hat{a}_z(k_3y - 4x)$$

Since  $\vec{F}$  is conservative field  $\therefore \text{curl } \vec{F} = 0$

$$\text{curl } \vec{F} = \vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 5y - k_1z & 3z + k_2x & k_3y - 4x \end{vmatrix} = 0$$

$$\Rightarrow \hat{i} \left( \frac{\partial(k_3 y - 4x)}{\partial y} - \frac{\partial(3z + k_2 x)}{\partial z} \right) \\ - \left( \frac{\partial(k_3 y - 4x)}{\partial x} - \frac{\partial(5y - k_1 z)}{\partial z} \right) \\ + \hat{k} \left( \frac{\partial(3z + k_2 x)}{\partial x} - \frac{\partial(5y - k_1 z)}{\partial y} \right)$$

$$\Rightarrow \hat{i}(k_3 - 3) - \hat{j}(-4 + k_1) + \hat{k}(k_2 - 5) = 0$$

$$\therefore k_3 - 3 = -4 + k_1 = k_2 - 5 = 0$$

$$\therefore k_1 = 4, k_2 = 5 \text{ and } k_3 = 3$$



Scan for Video solution


**6. (c)**

For any vector field  $\vec{A}$ ,

- If  $\nabla \cdot \vec{A} = 0$ , then  $\vec{A}$  is solenoidal
- If  $\nabla \times \vec{A} = \vec{0}$ , then  $\vec{A}$  is irrotational
- $\nabla \times (\nabla \times \vec{A}) = \nabla(\nabla \cdot \vec{A}) - \nabla^2 \cdot \vec{A}$

For  $\vec{A}$  to be irrotational,  $\nabla \times \vec{A} = \vec{0}$

$\therefore$  options 'c' is false statement



Scan for Video solution


**7. (45 to 45)**

$$\text{grad } u = \frac{\partial u}{\partial x} \hat{i} + \frac{\partial u}{\partial y} \hat{j} + \frac{\partial u}{\partial z} \hat{k}$$

$$= 2x\hat{i} + 2y\hat{j} + 2z\hat{k}$$

$$\therefore \text{grad } u \text{ at } (1,1,1) = 2\hat{i} + 2\hat{j} + 2\hat{k}$$

$$\text{and div } A = \frac{\partial A_1}{\partial x} + \frac{\partial A_2}{\partial y} + \frac{\partial A_3}{\partial z}$$

$$\therefore A = A_1 \hat{i} + A_2 \hat{j} + A_3 \hat{k}$$

$$\therefore \text{div } A = 2 + 3 + 4 = 9$$

Using vector identity,

$$\text{div}(uA) = u \text{div } A + A \text{ grad } u$$

$$\text{div}(uA) = (x^2 + y^2 + z^2)9 + (2x\hat{i} + 3y\hat{j} + 4z\hat{k}) \\ (2\hat{i} + 2\hat{j} + 2\hat{k})$$

$$\begin{aligned} \text{At } x = 1, y = 1, z = 1 \\ &= (1^2 + 1^2 + 1^2)9 + (4x + 6y + 8z) \\ &= 27 + 4 \times 1 + 6 \times 1 + 8 \times 1 \\ &= 27 + 4 + 6 + 8 \\ \text{div}(uA) &= 45 \end{aligned}$$

Scan for Video solution

**8. (b)**

$$\vec{x} = [4 \quad -2 \quad -6]^T$$

$$\vec{x} = \begin{bmatrix} 4 \\ -2 \\ -6 \end{bmatrix}$$

Euclidean norm length =  $|\vec{x}|$

$$= \sqrt{4^2 + (-2)^2 + (-6)^2} = \sqrt{56}$$

Scan for Video solution

**9. (d)**

$$\text{Curl } \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2x^2y & 5z^2 & -4yz \end{vmatrix}$$

$$= \hat{i} \left[ \frac{\partial(-4yz)}{\partial y} - \frac{\partial(5z^2)}{\partial z} \right] - \hat{j} \left[ \frac{\partial(-4yz)}{\partial x} - \frac{\partial(2x^2y)}{\partial z} \right]$$

$$+ \hat{k} \left[ \frac{\partial(5z^2)}{\partial x} - \frac{\partial(2x^2y)}{\partial y} \right]$$

$$= \hat{i}(-4z - 10z) - \hat{i}(0 - 0) + \hat{k}(0 - 2x^2)$$

$$= -14z\hat{i} - 2x^2\hat{k}$$


Scan for Video solution

**10. (a)**

Given

$$\|\vec{u}\| = 2\|\vec{v}\|$$

$\vec{w} = \vec{u} + a\vec{v}$  bisects angle between  
 $\vec{u}$  and  $\vec{v}$



$\Rightarrow$  let 'a' be angle between  
 $\vec{u}, \vec{w}$  and  $\vec{v}, \vec{w}$

By dot product

$$\Rightarrow \frac{\vec{u} \cdot \vec{w}}{\|\vec{u}\| \cdot \|\vec{w}\|} = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \cdot \|\vec{w}\|}$$

$$\Rightarrow \vec{u} \cdot \vec{w} = 2(\vec{v} \cdot \vec{w}) \quad (\text{since } \|\vec{u}\| = 2\|\vec{v}\|)$$

$$\Rightarrow \vec{u}(\vec{u} + a\vec{v}) = 2\vec{v}(\vec{u} + 2a\vec{v})$$

$$\Rightarrow \|\vec{u}\|^2 + a\vec{u} \cdot \vec{v} = 2\vec{v}\vec{u} + 2a\|\vec{v}\|^2$$

$$\Rightarrow \|\vec{u}\|^2 + a \cdot \vec{u} \cdot \vec{v} = 2\vec{v}\vec{u} + 2a \frac{\|\vec{u}\|^2}{4}$$

$$\Rightarrow \frac{2a}{4} = 1 \quad \text{and } a = 2$$

$$\Rightarrow a = 2$$



Scan for Video solution

**11. (133.9 to 134.1)**

$$\vec{V} = x^2\hat{i} + 2y^3\hat{j} + z^4\hat{k}$$

$$\vec{V} = V_1\hat{i} + V_2\hat{j} + V_3\hat{k}$$

$$\operatorname{div} \vec{V} = \frac{\partial V_1}{\partial x} + \frac{\partial V_2}{\partial y} + \frac{\partial V_3}{\partial z}$$

$$\operatorname{div} \vec{V} = 2x + 6y^2 + 4z^3$$

$$\text{At point } (1,2,3) \text{, } \operatorname{div} \vec{V} = 2 \times 1 + 6 \times 2^2 + 4 \times 3^3 = 2 + 24 + 108 = 134$$



Scan for Video solution

**12. (c)**From fig.(i)  $\vec{X}$  is a diverging field

$$\therefore \operatorname{div} \vec{X} \neq 0$$

$$\nabla \cdot \vec{X} \neq 0$$

from fig.(ii)  $\vec{Y}$  is a circularly rotating field

$$\therefore \operatorname{curl} \vec{Y} \neq 0$$

$$\nabla \times \vec{Y} \neq 0$$

from fig.(ii) is a circularly rotating field

$$\therefore \operatorname{curl} \vec{Z} \neq 0$$

$$\nabla \times \vec{Z} \neq 0$$

 Scan for Video solution



**13. (90 to 90)**

Given:

$$\vec{I} = \hat{x} 15 \cos(\omega t) + \hat{y} 5 \sin(\omega t)$$

$$\Rightarrow |\vec{I}| = ((15 \cos \omega t)^2 + (5 \sin \omega t)^2)^{1/2}$$

$$\Rightarrow |\vec{I}| = (225 + 25 \cos^2 \omega t)^{1/2}$$

For  $|\vec{I}|$  to be a minimum value,  $\cos \omega t = 0$ 

$$\Rightarrow \omega t = \frac{\pi}{2}$$

 $\therefore$  At minimum value of  $|\vec{I}|$ ,

$$\vec{I} = \hat{x} \cdot 15(0) + \hat{y} \cdot 5(1) = \hat{y} \cdot (5)$$

 $\therefore$  Angle made by  $\vec{I}$  with  $x$ -axis is 90 degrees.

 Scan for Video solution



**14. (b)**

Given Vectors are

$$\hat{i} + \hat{j} + \hat{k}, \quad 2\hat{i} + 3\hat{j} + \hat{k}, \quad 5\hat{i} + 6\hat{j} + 4\hat{k}$$

Let  $\vec{a} = \vec{i} + \vec{j} + \vec{k}$   
 $\vec{b} = 2\vec{i} + 3\vec{j} + \vec{k}$   
 $\vec{c} = 5\vec{i} + 6\vec{j} + 4\vec{k}$

For  $\vec{a}, \vec{b}, \vec{c}$  to linearly dependent,

Determinant of matrix  $\begin{bmatrix} \vec{a} & \vec{b} & \vec{c} \end{bmatrix} = 0$

$$\Rightarrow \begin{vmatrix} 1 & 1 & 1 \\ 2 & 3 & 1 \\ 5 & 6 & 4 \end{vmatrix} = 0$$

$$\Rightarrow 1 \times (6) - 1 \times (3) + 1 \times (-3) = 0 \text{ (True)}$$

$\therefore$  Given vectors are linearly Dependent.



Scan for Video solution



### 15. (6.8 to 7.2)

Given  $f(x, y, z) = x^2 + 3y^2 + z^3$

$$\Rightarrow \nabla f = \frac{\partial f}{\partial x} \vec{i} + \frac{\partial f}{\partial y} \vec{j} + \frac{\partial f}{\partial z} \vec{k}$$

$$= 2x\vec{i} + 6y\vec{j} + 3z^2\vec{k}$$

$$\Rightarrow \nabla f \Big|_{(1,1,1)} = 2\hat{i} + 6\hat{j} + 3\hat{k}$$

$$\Rightarrow |\nabla f|_{(1,1,1)} = \sqrt{2^2 + 6^2 + 3^2} = 7$$



Scan for Video solution



### 16. (d)

$$\nabla V = 2x^2y + 3y^2z + 4z^2x$$

$$\nabla \nabla V = (4xy + 4z^2)\hat{i} + (2x^2 + 6yz)\hat{j} + (3y^2 + 8zx)\hat{k}$$

$$\nabla \times \nabla V = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 4xy + 4z^2 & 2x^2 + 6yz & 3y^2 + 8zx \end{vmatrix}$$

$$= \hat{i}(6y - 6y) - \hat{j}(8z - 8z) + \hat{k}(4x - 4x)$$

$$\therefore \nabla \times \nabla V = \mathbf{0}$$



Scan for Video solution




### 17. (d)

Given vector field is  $\vec{A}(\vec{r})$

For any closed loop,

By stoke's theorem



$$\oint \vec{A} \cdot d\vec{l} = \iint_s \text{curl} \vec{A} \cdot d\vec{s}$$

$$= \iint_s (\nabla \times \vec{A}) \cdot d\vec{s}$$

$$\therefore \oint_c \vec{A} \cdot d\vec{l} = \iint_s (\nabla \times \vec{A}) \cdot d\vec{s}$$


### 18. (d)

$$\text{Given } \vec{A} = x\hat{a}_x + y\hat{a}_y + z\hat{a}_z$$

$$\Rightarrow \nabla \cdot \vec{A} = \frac{\partial}{\partial x}(x) + \frac{\partial}{\partial y}(y) + \frac{\partial}{\partial z}(z)$$

$$= 1 + 1 + 1 = 3$$

$$\therefore \nabla \cdot \vec{A} = 3$$



Scan for Video solution



### 19. (a)

$$\text{Area of parallelogram} = |\vec{OP} \times \vec{OR}|$$


$$\text{Given } \vec{OP} = a\hat{i} + b\hat{j}$$

$$\vec{OR} = c\hat{i} + d\hat{j}$$

$$\text{Area of parallelogram} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a & b & 0 \\ c & d & 0 \end{vmatrix}$$

$$= 0\hat{i} + 0\hat{j} + (ad - bc)\hat{k} = (ad - bc)\hat{k}$$

$$\text{Area} = ad - bc$$



Scan for Video solution



20. (a)

$$\vec{a} \cdot \vec{b} = a \cdot b \cos \theta$$

$$\vec{a} \times \vec{b} = a \cdot b \sin \theta \hat{n}$$

$$\therefore |\vec{a} \times \vec{b}|^2 = a^2 b^2 \sin^2 \theta |\hat{n}|^2$$

$$= a^2 b^2 (1 - \cos^2 \theta) \quad \left[ \because |\hat{n}|^2 = 1 \right]$$

$$= a^2 b^2 - a^2 b^2 \cos^2 \theta$$

$$= a^2 b^2 - (\vec{a} \cdot \vec{b})^2$$



Scan for Video solution



21. (a)

$$\text{We know, } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\nabla \vec{r} = \frac{\partial}{\partial x}(x)\hat{i} + \frac{\partial}{\partial y}(y)\hat{j} + \frac{\partial}{\partial z}(z)\hat{k}$$

$$= 1 + 1 + 1$$

$$\text{div } \vec{r} = 3$$



Scan for Video solution



22. (b)

$$f = x^2 + 3y^2 + 2z^2$$

$$\text{Grad } f = \frac{\partial f}{\partial x}\hat{i} + \frac{\partial f}{\partial y}\hat{j} + \frac{\partial f}{\partial z}\hat{k}$$

$$\text{Grad } f = (2x)\hat{i} + (6y)\hat{j} + (4z)\hat{k}$$

$$\text{At } P(1, 2, -1)$$

$$\text{Grad } f = 2\hat{i} + 12\hat{j} - 4\hat{k}$$



Scan for Video solution




23. (b)

$$\text{Given } \vec{V} = \nabla \times \vec{A} = \text{curl } \vec{A}$$

$S_c$  is any surface whose boundary is closed contour 'C',

By stokes theorem,



$$\oint_C \vec{A} \cdot d\vec{l} = \int \oint_{S_c} \text{curl } \vec{A} \cdot d\vec{s}$$

$$= \int \oint_{S_c} \vec{V} \cdot d\vec{s}$$

$$\therefore \oint_C \vec{A} \cdot d\vec{l} = \int_{S_c} \vec{V} \cdot d\vec{s} \rightarrow \text{From stokes theorem}$$




Scan for Video solution



24. (c)

$$\text{Given } \vec{P} \cdot \vec{Q} = 0$$



$$\Rightarrow |\vec{P}||\vec{Q}|\cos \theta = 0$$

$$\Rightarrow \cos \theta = 0$$


$$\Rightarrow \theta = 90^\circ$$



Scan for Video solution



25. (b)



$$f = xyz$$

Given scalar field,  $f(x, y, z) = xyz$

Point is  $P(0, 2, 1)$  and in the direction of  $(2, 1, 2)$

Rate of increase of  $f(x, y, z)$  at point 'P' in the direction of  $\vec{a}$  is given by

$$\nabla f|_P \cdot \hat{a}$$

$$= \nabla f|_P \cdot \frac{\vec{a}}{|\vec{a}|}$$

$$\therefore \nabla f = yzi + x^2j + xyk$$

$$\nabla f|_P = 2\hat{i} + 0\hat{j} + 0\hat{k}$$

$$\text{Direction is } \vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$$

$$\therefore \text{Rate of increase} = (2\hat{i}) \cdot \frac{(2\hat{i} + \hat{j} + 2\hat{k})}{\sqrt{2^2 + 1^2 + 2^2}}$$

$$= \frac{4}{\sqrt{9}} = \frac{4}{3}$$

$\therefore$  Rate of increase of  $f$  at  $(0, 2, 1)$  in the direction of  $(2, 1, 2)$  is  $\frac{4}{3}$



Scan for Video solution

**26. (b)**

$$f = 4x^2 + 7xy + 3xz^2$$

$$\nabla f = \frac{\partial f}{\partial x}\hat{i} + \frac{\partial f}{\partial y}\hat{j} + \frac{\partial f}{\partial z}\hat{k}$$

$$\nabla f = (8x + 7y + 3z^2)\hat{i} + (7x)\hat{j} + (6xz)\hat{k}$$

At point  $(1, 0, 2)$ 

$$\nabla f = (8 + 0 + 12)\hat{i} + 7\hat{j} + 12\hat{k}$$

$$= 20\hat{i} + 7\hat{j} + 12\hat{k}$$

In this direction function will increase most rapidly.



Scan for Video solution

**27. (1 to 1)**

$$f(x, y) = xe^y$$

$$P(2, 0) \text{ and } Q\left(\frac{1}{2}, 2\right)$$

$\therefore$  Required directional Derivative = grad  $f$ .  $PQ$  at point  $(2, 0)$

$$\text{Grad } f = \frac{\partial f}{\partial x}\hat{i} + \frac{\partial f}{\partial y}\hat{j}$$

$$= e^y\hat{i} + xe^y\hat{j}$$

$$\text{So, grad } f_{(2,0)} = e^0\hat{i} + 2e^0\hat{j}$$

$$= \hat{i} + 2\hat{j}$$

$$\text{Now } \overrightarrow{PQ} = \overrightarrow{OQ} - \overrightarrow{OP} = \left(\frac{1}{2}\hat{i} + 2\hat{j}\right) - (2\hat{i} + 0\hat{j})$$

$$\overrightarrow{PQ} = -\frac{3}{2}\hat{i} + 2\hat{j}$$

$$PQ = \frac{\overrightarrow{PQ}}{|\overrightarrow{PQ}|} = \frac{-\frac{3}{2}\hat{i} + 2\hat{j}}{\sqrt{\left(\frac{-3}{2}\right)^2 + 2^2}}$$

$\therefore$  Directional derivative = grad  $f$ .  $PQ$

$\therefore$  Directional derivative at  $(2, 0)$  =

$$\left(\hat{i} + 2\hat{j}\right) \cdot \frac{\left(-\frac{3}{2}\hat{i} + 2\hat{j}\right)}{\sqrt{\frac{9}{4} + 4}} = \frac{-\frac{3}{2} + 4}{\sqrt{\frac{9+16}{4}}} = \frac{\frac{5}{2}}{\frac{5}{2}} = 1$$



Scan for Video solution

**28. (c)****Given**

$$f(x, y, z) = xyz \text{ and } P(-1, 1, 3)$$

The directional derivative of  $f(x, y, z)$  at 'P' in the direction of  $\vec{a}$  is given by

$$\begin{aligned} \nabla f|_P \cdot \hat{a} &= \nabla(xyz)|_{(-1,1,3)} \cdot \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{|\sqrt{1^2 + (-2)^2 + 2^2}|} \\ &= y\hat{i} + xz\hat{j} + xy\hat{k}|_{(-1,1,3)} \cdot \frac{(\hat{i} - 2\hat{j} + 2\hat{k})}{3} \\ &= (3\hat{i} - 3\hat{j} - \hat{k}) \cdot \frac{(\hat{i} - 2\hat{j} + 2\hat{k})}{3} = \frac{7}{3} \end{aligned}$$

$\therefore$  Directional derivative =  $\frac{7}{3}$




Scan for Video solution

**29. (c)****Given**

$$f = x^2 + y^2; x = 1 \text{ & } y = 1$$

Directional derivative of a function 'f' at point 'P' in direction of  $\vec{a}$  is  $\nabla f|_P \cdot \hat{a}$



$$\text{Directional derivative} = \nabla f|_{(1,1)} \cdot \hat{a}$$

$$\begin{aligned} &= \nabla(x^2 + y^2)|_{(1,1)} \cdot \left( \frac{\vec{i} + \vec{j}}{\sqrt{2}} \right) \\ &= (2x\vec{i} + 2y\vec{j})|_{(1,1)} \cdot \left( \frac{\vec{i} + \vec{j}}{\sqrt{2}} \right) \\ &= (2\vec{i} + 2\vec{j}) \cdot \frac{(\vec{i} + \vec{j})}{\sqrt{2}} = 2\sqrt{2} \end{aligned}$$

$$\therefore \text{Directional derivative} = 2\sqrt{2}$$



Scan for Video solution



**30. (c)**

**Given,**

$$\vec{u} = e^x \cos y \hat{i} + e^x \cdot \sin y \hat{j}$$

$$\begin{aligned} \Rightarrow \nabla \cdot \vec{u} &= \frac{\partial}{\partial x}(e^x \cos y) + \frac{\partial}{\partial y}(e^x \cdot \sin y) \\ &= \cos y \cdot (e^x) + e^x \cdot (\cos y) \\ &= 2e^x \cdot \cos y \end{aligned}$$

$$\Rightarrow \boxed{\nabla \cdot \vec{u} = 2e^x \cdot \cos y}$$



Scan for Video solution



**31. (c)**

**Given,**

$$\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$$

$$\Rightarrow |\vec{r}| = \sqrt{x^2 + y^2 + z^2} = (x^2 + y^2 + z^2)^{1/2}$$

$$\phi = \ln |\vec{r}| = \frac{1}{2} \cdot \ln(x^2 + y^2 + z^2)$$

$$\begin{aligned} \Rightarrow \text{grad } \phi &= \nabla \phi = \frac{\partial \phi}{\partial x} \hat{i} + \frac{\partial \phi}{\partial y} \hat{j} + \frac{\partial \phi}{\partial z} \cdot \hat{k} \\ &= \frac{1}{2} \cdot \frac{1}{(x^2 + y^2 + z^2)} 2x\vec{i} + \frac{1}{2} \cdot \frac{1}{(x^2 + y^2 + z^2)} \cdot 2y\vec{j} \\ &\quad + \frac{1}{2} \cdot \frac{1}{(x^2 + y^2 + z^2)} \cdot 2z\vec{k} \\ &= \frac{x\vec{i} + y\vec{j} + z\vec{k}}{(x^2 + y^2 + z^2)} = \frac{\vec{r}}{|\vec{r}|^2} = \frac{\vec{r}}{\vec{r} \cdot \vec{r}} \end{aligned}$$

$$\therefore \boxed{\text{grad } \phi = \nabla \phi = \frac{\vec{r}}{\vec{r} \cdot \vec{r}}}$$

Scan for Video solution



**32. (a)**

$$\phi(x, y, z) = xy^2 + yz^2 + zx^2$$

$$\text{grad } \phi = \frac{\partial \phi}{\partial x} \hat{i} + \frac{\partial \phi}{\partial y} \hat{j} + \frac{\partial \phi}{\partial z} \hat{k}$$

$$\text{grad } \phi = (y^2 + 2xz)\hat{i} + (2xy + z^2)\hat{j} + (2yz + x^2)\hat{k}$$

$\therefore$  At point  $(2, -1, 1)$

$$\begin{aligned} \text{grad } \phi &= ((-1)^2 + 2 \cdot 2 \cdot 1)\hat{i} + (2 \cdot 2 \cdot (-1) + 1^2)\hat{j} \\ &\quad + (2 \cdot (-1) \cdot 1 + 2^2)\hat{k} \end{aligned}$$

$$\text{grad } \phi = 5\hat{i} - 3\hat{j} + 2\hat{k}$$

$\therefore$  The required directional derivative is given by,

$$\text{Directional Derivative} = \text{grad } \phi \cdot \frac{\vec{P}}{|\vec{P}|}$$

$$= (5\hat{i} - 3\hat{j} + 2\hat{k}) \cdot \frac{(\hat{i} + 2\hat{j} + 2\hat{k})}{\sqrt{1^2 + 2^2 + 2^2}}$$

$$= \frac{5 - 6 + 4}{\sqrt{9}} = \frac{3}{3} = 1$$



Scan for Video solution



**33. (0 to 0)**

Let  $\vec{F} = F_x \vec{i} + F_y \vec{j} + F_z \vec{k}$

$$\begin{aligned}\nabla \cdot (\nabla \times \vec{F}) &= \nabla \cdot \left| \begin{array}{ccc} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{array} \right| \\ &= \nabla \cdot \left( \vec{i} \left( \frac{\partial F_z}{\partial y} - \frac{\partial F_y}{\partial z} \right) - \vec{j} \left( \frac{\partial F_z}{\partial x} - \frac{\partial F_x}{\partial z} \right) + \vec{k} \left( \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) \right) \\ &= \frac{\partial}{\partial x} \left( \frac{\partial F_z}{\partial y} - \frac{\partial F_y}{\partial z} \right) - \frac{\partial}{\partial y} \left( \frac{\partial F_z}{\partial x} - \frac{\partial F_x}{\partial z} \right) + \frac{\partial}{\partial z} \left( \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right) \\ &= \frac{\partial^2 F_z}{\partial x \partial y} - \frac{\partial^2 F_y}{\partial x \partial z} - \frac{\partial^2 F_z}{\partial y \partial x} + \frac{\partial^2 F_x}{\partial y \partial z} + \frac{\partial^2 F_y}{\partial z \partial x} - \frac{\partial^2 F_x}{\partial z \partial y} = 0\end{aligned}$$

$\therefore$  For any vector point function  $\vec{F}$ ,  $\nabla \cdot (\nabla \times \vec{F}) = 0$

Scan for Video solution

**34. (b)**

Given

$$\vec{F} = \hat{a}_x (3y - k_1 z) + \hat{a}_y (k_2 x - 2z) - \hat{a}_z (k_3 y + z)$$

For  $\vec{F}$  to be irrotational,  $\nabla \times \vec{F} = 0$

$$\Rightarrow \left| \begin{array}{ccc} \hat{a}_x & \hat{a}_y & \hat{a}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3y - k_1 z & k_2 x - 2z & -k_3 y - z \end{array} \right| = 0$$

$$\Rightarrow \hat{a}_x (-k_3 + 2) - \hat{a}_y (k_1) + \hat{a}_z (k_2 - 3) = 0$$

$$\Rightarrow -k_3 + 2 = 0, k_1 = 0; k_2 - 3 = 0$$

$$\therefore k_1 = 0; k_2 = 3; k_3 = 2$$

Scan for Video solution



$$\begin{aligned}\therefore \nabla \cdot (\nabla \times \vec{V}) &= \frac{\partial^2 V_z}{\partial x \cdot \partial y} - \frac{\partial^2 V_y}{\partial x \cdot \partial z} - \frac{\partial^2 V_z}{\partial x \cdot \partial y} + \frac{\partial^2 V_x}{\partial y \cdot \partial z} \\ &\quad + \frac{\partial^2 V_y}{\partial z \cdot \partial x} - \frac{\partial^2 V_x}{\partial y \cdot \partial z} \\ &= 0\end{aligned}$$

For any Vector  $\vec{V}$ ,  $\text{Div Curl } \vec{V} = 0$



Scan for Video solution



### 38. (-5.72 to -5.70)

$$u = x^2 - 3yz$$

Directional derivative = grad  $u \cdot \hat{a}$

$$\Rightarrow \nabla u = \frac{\partial u}{\partial x} \hat{i} + \frac{\partial u}{\partial y} \hat{j} + \frac{\partial u}{\partial z} \hat{k}$$

$$\nabla u_{(2,-1,4)} = (2x) \hat{i} + (-3z) \hat{j} + (-3y) \hat{k} = 4\hat{i} - 12\hat{j} + 3\hat{k}$$

$$\hat{a} = \hat{i} + \hat{j} - 2\hat{k}$$

The required directional derivative is

$$\begin{aligned}&= \text{grad } u \cdot \frac{\hat{a}}{|\hat{a}|} \\&= (4\hat{i} - 12\hat{j} + 3\hat{k}) \cdot \frac{(\hat{i} + \hat{j} - 2\hat{k})}{\sqrt{1^2 + 1^2 + (-2)^2}} \\&= \frac{4 - 12 - 6}{\sqrt{6}} = -\frac{14}{\sqrt{6}} = -5.715\end{aligned}$$



Scan for Video solution



### 39. (a)

$$\vec{P} = x^3 y \hat{a}_x - x^2 y^2 \hat{a}_y - x^2 y z \hat{a}_z$$

Considering

$$\text{div } \vec{P} = \nabla \cdot \vec{P}$$

$$\begin{aligned}&= \frac{\partial}{\partial x}(x^3 y) + \frac{\partial}{\partial y}(-x^2 y^2) + \frac{\partial}{\partial z}(-x^2 y z) \\&= 3x^2 y - 2x^2 y - x^2 y = 0\end{aligned}$$

$$\therefore \nabla \cdot \vec{P} = 0$$

$$\text{Considering curl } \vec{P} = \nabla \times \vec{P} = \begin{vmatrix} \hat{a}_x & \hat{a}_y & \hat{a}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^3 y & -x^2 y^2 & -x^2 y z \end{vmatrix}$$

$$= \hat{a}_x (-x^2 z + 0) - \hat{a}_y (-2xyz - 0) + \hat{a}_z (-2xy^2 - x^3)$$

$$\neq \vec{0}$$

Since

$$\nabla \cdot \vec{P} = 0 \text{ & } \nabla \times \vec{P} \neq 0$$


$\vec{P}$  is solenoidal but not irrotational.



Scan for Video solution



### 40. (2.99 to 3.01)



$$\text{Given } f(x, y) = \frac{xy}{\sqrt{2}}(x + y)$$

Directional derivative of  $f$  at  $(1, 1)$ : In the direction of  $45^\circ$  to  $y$ -axis is

$$= \nabla f|_{(1,1)} \cdot \hat{a}$$

$$\nabla f|_{(1,1)} = \frac{1}{\sqrt{2}} \nabla(x^2 y + xy^2)|_{(1,1)}$$

$$= \frac{1}{\sqrt{2}} \left\{ (2xy + y^2) \hat{i} + (x^2 + 2xy) \hat{j} \right\}|_{(1,1)}$$

$$= \frac{1}{\sqrt{2}} \{3\hat{i} + 3\hat{j}\}$$

$$\hat{a} = \frac{\hat{i} + \hat{j}}{\sqrt{2}}$$

∴ Directional derivative

$$= \nabla f|_{(1,1)} \cdot \hat{a} = \frac{1}{\sqrt{2}} \cdot (3\hat{i} + 3\hat{j}) \cdot \frac{1}{\sqrt{2}} (\hat{i} + \hat{j})$$

$$= \frac{1}{2} (3 + 3) = 3$$

∴ Directional Derivative = 3



Scan for Video solution



Scan for Video solution



**41. (2.9 to 3.1)**

$$\text{Given } \vec{r} = x \cdot \hat{a}_x + y \cdot \hat{a}_y + z \cdot \hat{a}_z$$

$$|\vec{r}| = r$$

$$\nabla \cdot (r^2 \nabla \ln r) = \nabla \cdot (r^2 \nabla \left( \frac{1}{2} \cdot \ln(x^2 + y^2 + z^2) \right))$$

$$= \nabla \cdot \left( r^2 \cdot \frac{1}{2} \left( \frac{1}{r^2} (2x \hat{a}_x + 2y \hat{a}_y + 2z \hat{a}_z) \right) \right)$$

$$= \nabla \cdot (x \cdot \hat{a}_x + y \cdot \hat{a}_y + z \cdot \hat{a}_z) = 1 + 1 + 1 = 3$$

$$\therefore \nabla \cdot (r^2 \nabla \ln r) = 3$$



Scan for Video solution


**42. (a)**

Given

$$\vec{F} = x^2 z^2 \hat{i} - 2xy^2 z \hat{j} + 2y^2 z^3 \hat{k}$$

$$\Rightarrow \text{Curl } \vec{F} = \nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2 z^2 & -2xy^2 z & 2y^2 z^3 \end{vmatrix}$$

$$= \hat{i} \left( \frac{\partial}{\partial y} (2y^2 z^3) - \frac{\partial}{\partial z} (-2xy^2 z) \right)$$

$$- \hat{j} \left( \frac{\partial}{\partial x} (2y^2 z^3) - \frac{\partial}{\partial z} (x^2 z^2) \right)$$

$$+ \hat{k} \left( \frac{\partial}{\partial x} (-2xy^2 z) - \frac{\partial}{\partial y} (x^2 z^2) \right)$$

$$\Rightarrow \text{Curl } \vec{F} = \hat{i}(4yz^3 + 2xy^2) + \hat{j}(2x^2 z) - \hat{k}(2y^2 z)$$



Scan for Video solution


**43. (c)**

Given

$$\text{Vector field } x^2 z \hat{i} + xy \hat{j} - yz^2 \hat{k} = \vec{F} \text{ (say)}$$

$$\Rightarrow \nabla \cdot \vec{F} = \text{div } \vec{F} = \frac{\partial}{\partial x} (x^2 z) + \frac{\partial}{\partial y} (xy) + \frac{\partial}{\partial z} (-yz^2)$$

$$= 2xz + x - 2yz$$

$$\therefore \nabla \cdot \vec{F}|_{(1,-1,1)} = 2(1)(1) + 1 - 2(-1)(1) = 5$$

$$\therefore \boxed{\nabla \cdot \vec{F}|_{(1,-1,1)} = 5}$$



Scan for Video solution


**44. (a)**

$$\nabla \cdot (\vec{f} \vec{V}) = x^2 y + y^2 z + z^2 x$$

$$\vec{V} = y \hat{i} + z \hat{j} + x \hat{k}$$

$$\therefore \text{div } \vec{V} = \frac{\partial V_1}{\partial x} + \frac{\partial V_2}{\partial y} + \frac{\partial V_3}{\partial z}$$

 Using vector identity;  $\nabla \cdot \vec{V} = 0$ 

$$\vec{V} \cdot (\vec{f} \vec{V}) = \vec{f} \cdot (\nabla \cdot \vec{V}) + \vec{V} \cdot (\vec{f} \cdot \nabla \vec{V})$$


$$x^2 y + y^2 z + z^2 x = 0 + \vec{V} \cdot \nabla \vec{f}$$

$$\vec{V} \cdot \nabla \vec{f} = x^2 y + y^2 z + z^2 x$$



Scan for Video solution


**45. (a)**

 Given: surface is  $x^2 + y^2 + z^2 = 1$ 

 For any surface  $\phi(x, y, z) = c$ , the outward normal

 vector at point 'P' is given by  $\frac{\Delta \phi|_P}{|\Delta \phi|_P} = \hat{n}$ 

$$\Delta \phi|_P = \left( \frac{\partial \phi}{\partial x} \hat{i} + \frac{\partial \phi}{\partial y} \hat{j} + \frac{\partial \phi}{\partial z} \hat{k} \right)|_P$$

$$= 2x \hat{i} + 2y \hat{j} + 2z \hat{k} \Big|_{\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0\right)} = \sqrt{2} \hat{i} + \sqrt{2} \hat{j} + 0 \hat{k}$$

$$\therefore \hat{n} = \frac{\nabla \phi|_P}{|\nabla \phi|_P} = \frac{\sqrt{2}\hat{i} + \sqrt{2}\hat{j}}{2}$$

$$\Rightarrow \hat{n} = \frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{j}$$



Scan for Video solution

**46. (a)**

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

If  $\mathbf{f} \rightarrow f(r)$  then using standard result

$$\operatorname{div} \mathbf{f} = \vec{\nabla} \cdot \vec{\mathbf{f}} = \frac{1}{r^2} \left( \frac{\partial}{\partial r} r^2 |\mathbf{f}| \right)$$

 $\therefore \operatorname{div} \mathbf{A} = \vec{\nabla} \cdot \mathbf{A} = 0$  (Given)

$$\frac{1}{r^2} \left( \frac{\partial}{\partial r} \cdot r^2 |\mathbf{A}| \right) = 0$$

$$\frac{1}{r^2} \left( \frac{\partial}{\partial r} \cdot r^2 \cdot K r^n \right) = 0$$

$$\frac{1}{r^2} \left( \frac{\partial}{\partial r} \cdot K \cdot r^{n+2} \right)$$

$$\frac{1}{r^2} (n+2) K r^{n+1} = 0$$

$$(n+2)K r^{n-1} = 0$$

$$\therefore n+2=0$$

$$n=-2$$



Scan for Video solution

**47. (b)**

$$\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$$

Directional derivative is given by,

$$(\text{Grad. } f) \cdot \hat{a} = \operatorname{grad} f \cdot \frac{\vec{a}}{|\vec{a}|}$$

$$= \left( \frac{\partial f}{\partial x} \hat{i} + \frac{\partial f}{\partial y} \hat{j} + \frac{\partial f}{\partial z} \hat{k} \right) \cdot \frac{\hat{i} - \hat{j} + 2\hat{k}}{\sqrt{1^2 + (-1)^2 + 2^2}}$$

$$= (2x\hat{i} + 6y\hat{j} + 4z\hat{k}) \cdot \frac{\hat{i} - \hat{j} + 2\hat{k}}{\sqrt{6}}$$

The required directional derivative at P (1,2,-1) is

$$= (2(1)\hat{i} + 6(2)\hat{j} + 4(-1)\hat{k}) \cdot \frac{(\hat{i} - \hat{j} + 2\hat{k})}{\sqrt{6}}$$

$$= \frac{2-12-8}{\sqrt{6}} = -\frac{18}{\sqrt{6}} = -3\sqrt{6}$$



Scan for Video solution

**48. (c)****Given:**vector function  $\vec{F} = 3xz\hat{i} + 2xy\hat{j} - yz^2\hat{k}$ Divergence of  $\vec{F} = F_x\hat{i} + F_y\hat{j} + F_z\hat{k}$  is given by

$$\vec{\nabla} \cdot \vec{F} = \frac{\partial}{\partial x}(F_x) + \frac{\partial}{\partial y}(F_y) + \frac{\partial}{\partial z}(F_z)$$

$$= \frac{\partial}{\partial x}(3xz) + \frac{\partial}{\partial y}(2xy) + \frac{\partial}{\partial z}(-yz^2)$$

$$= 3z + 2x - 2yz$$

$$\vec{\nabla} \cdot \vec{F}|_{(1,1,1)} = 3z + 2x - 2yz|_{(1,1,1)}$$

$$= 3(1) + 2(1) - 2(1)(1) = 3$$

$$\therefore \vec{\nabla} \cdot \vec{F}|_{(1,1,1)} = 3$$



Scan for Video solution

**49. (d)****Given:**Vector  $\vec{F} = (x-y)\hat{i} + (y-x)\hat{j} + (x+y+z)\hat{k}$ If  $\vec{F} = F_x\hat{i} + F_y\hat{j} + F_z\hat{k}$ , then divergence of  $\vec{F}$  is

$$\vec{\nabla} \cdot \vec{F} = \frac{\partial}{\partial x}(F_x) + \frac{\partial}{\partial y}(F_y) + \frac{\partial}{\partial z}(F_z)$$

$$\Rightarrow \vec{\nabla} \cdot \vec{F} = \frac{\partial}{\partial x}(x-y) + \frac{\partial}{\partial y}(y-x) + \frac{\partial}{\partial z}(x+y+z)$$

$$= 1 + 1 + 1 = 3$$

$$\therefore \vec{\nabla} \cdot \vec{F} = 3$$



Scan for Video solution



50. (b)

Given:  $f(x, y, z) = x^2 + 2y^2 + z;$

$P(1, 1, 2)$ ; and  $\vec{a} = 3\vec{i} - 4\vec{j}$

The directional derivation of  $f(x, y, z)$  at  $P$  in direction of  $\vec{a}$  is given by  $\nabla f|_P \cdot \frac{\vec{a}}{|\vec{a}|}$

$$\begin{aligned}\nabla f|_P \cdot \frac{\vec{a}}{|\vec{a}|} &= \left( \frac{\partial f}{\partial x} \vec{i} + \frac{\partial f}{\partial y} \vec{j} + \frac{\partial f}{\partial z} \vec{k} \right)|_P \cdot \frac{\vec{a}}{|\vec{a}|} \\ &= ((2x)\vec{i} + (4y)\vec{j} + 1\vec{k})|_{(1,1,2)} \cdot \frac{(3\vec{i} - 4\vec{j})}{|3\vec{i} - 4\vec{j}|} \\ &= (2\vec{i} + 4\vec{j} + \vec{k}) \cdot \frac{(3\vec{i} - 4\vec{j})}{5} = \frac{6 - 16}{5} = -2\end{aligned}$$

$\therefore$  Directional derivative = -2



Scan for Video solution



51. (9 to 10)

C:  $r = 1 + \cos\theta$

$$\int_C \vec{F} \cdot d\vec{l} = \oint_C (-y\hat{i} + x\hat{j})(dx\hat{i} + dy\hat{j})$$

$$= \int_C -ydx + xdy$$

According to Green theorem;

$$\int_C Mdx + Ndy = \iint \left( \frac{\partial N}{\partial X} - \frac{\partial M}{\partial y} \right) dx dy$$

$$\therefore \int_C -ydx + xdy = \iint 1 - (-1) dx dy$$

$$= 2 \iint dx dy = 2 \text{ (Area of closed curve C)}$$

$$= 2 \int_0^{2\pi} \int_0^r r dr d\theta = 2 \int_0^{2\pi} \int_0^r \frac{r^2}{2} d\theta = \int_0^{2\pi} (1 + \cos\theta)^2 d\theta$$

$$= \int_0^{2\pi} [1 + \cos^2\theta + 2\cos\theta] d\theta$$

$$= [\theta]_0^{2\pi} + 4 \int_0^{\pi/2} \cos^2\theta + 4 \int_0^{\pi} \cos\theta$$

$$= 2\pi + 4 \frac{1}{2} \cdot \frac{\pi}{2} + 4 \times 0$$

$$= 2\pi + \pi + 0$$

$$= 3\pi = 9.42$$

Scan for Video solution



52. (0.9 to 1.1)

Given that

|   |     |     |     |
|---|-----|-----|-----|
| x | -1  | 0   | 1   |
| y | 0.8 | 2.2 | 2.8 |

Let line of best fit is  $y = ax + b$

By normal equations

$$\sum y = a \sum x + bn \quad \dots\dots(1)$$

$$\sum xy = a \sum x^2 + b \sum x \quad \dots\dots(2)$$

| x  | y   | $x^2$ | xy   |
|----|-----|-------|------|
| -1 | 0.8 | 1     | -0.8 |
| 0  | 2.2 | 0     | 0    |
| 1  | 2.8 | 1     | 2.8  |

$$\sum x = 0, \sum y = 5.8$$

$$\sum x^2 = 2, \sum xy = 2, n = 3$$

putting in (1) and (2)

$$5.8 = a(0) + b(3)$$

$$2 = a(2) + b(0)$$

$\therefore$  on solving  $a = 1, b = 1.93$

$\therefore$  line of best fit  $y = ax + b$

$$y = x + 1.93$$

$\therefore$  Slope of line of best fit = 1

Scan for Video solution



53. (b)

Given integral is  $\int_P^Q (z^2 dx + 3y^2 dy + 2xz dz) = I$  (say)

$$\therefore I = \int_P^Q (z^2 \vec{i} + 3y^2 \vec{j} + 2xz \vec{k}) \cdot (dx \vec{i} + dy \vec{j} + dz \vec{k})$$

For  $I = \int_A^B \vec{F} \cdot d\vec{r}$ , if  $\vec{F} = \nabla \phi$  for some scalar function  $\phi$ ,

$$\text{Then } \int_A^B \vec{F} \cdot d\vec{r} = \phi|_B - \phi|_A$$

$$\vec{F} = z^2 \vec{i} + 3y^2 \vec{j} + 2xz \vec{k} = \nabla(xz^2 + y^3)$$

$$\therefore \int_P^Q \vec{F} \cdot d\vec{r} = xz^2 + y^3 \Big|_P^Q = (xz^2 + y^3) \Big|_{(1,1,2)}^{(2,3,1)}$$

$$= (2(1)^2 + 3^3) - (1(2)^2 + 1^3) = 29 - 5 = 24$$

$$\therefore \int_P^Q z^2 dx + 3y^2 dy + 2xz dz = 24$$




Scan for Video solution



54. (-0.001 to 0.001)

Given

$\vec{B} = x \vec{i} + y \vec{j} - 2z \vec{k}$  and  $\hat{n}$  is unit normal to 'S'




$$\int_S \vec{B} \cdot \hat{n} ds = \iiint_V (\nabla \cdot \vec{B}) dv$$

(From Gauss-divergence theorem)

$$\nabla \cdot \vec{B} = \frac{\partial}{\partial x}(x) + \frac{\partial}{\partial y}(y) + \frac{\partial}{\partial z}(-2z) = 1 + 1 - 2 = 0$$

$$\Rightarrow \iiint_V (\nabla \cdot \vec{B}) dv = 0 \text{ Since } \nabla \cdot \vec{B} = 0$$

$$\Rightarrow \boxed{\int_S \vec{B} \cdot \hat{n} ds = 0}$$



55. (a)

$$\text{Given } \phi = \frac{1}{2}(x^2 + y^2 + z^2)$$

$$\Rightarrow \nabla \phi = x \vec{i} + y \vec{j} + z \vec{k} \quad \left\{ \because \nabla = \frac{\partial}{\partial x} \vec{i} + \frac{\partial}{\partial y} \vec{j} + \frac{\partial}{\partial z} \vec{k} \right\}$$

'S' is surface of sphere of unit radius.

$$\iint_S \hat{n} \cdot \nabla \phi ds = \iint_S \nabla \phi \cdot \hat{n} ds$$


From Gauss-divergence theorem,

$$\iint_S \nabla \phi \cdot \hat{n} ds = \iiint_V \nabla \cdot (\nabla \phi) dv$$

$$= \iiint_V \nabla \cdot (x \vec{i} + y \vec{j} + z \vec{k}) dv = 3 \cdot \iiint_V dv$$

$$= 3 \times \text{Volume of sphere} = 3 \times \frac{4}{3}\pi(1)^3 = 4\pi$$

$$\therefore \boxed{\iint_S \nabla \phi \cdot \hat{n} ds = 4\pi}$$



56. (a)

$$\text{Given } \vec{F} = 3x \hat{i} + 5y \hat{j} + 6z \hat{k}$$

and A: Area of surface of a cube


From Gauss-divergence Theorem

$$\iint_A \vec{F} \cdot d\vec{A} = \iiint_V (\nabla \cdot \vec{F}) dv = \iiint_V 14 dv$$

$$= 14 \times \text{Volume of the cube}$$

$$= 14 \times (a)^3 = 14 \times (1)^3 = 14$$

$$\therefore \boxed{\iint_A \vec{F} \cdot d\vec{A} = 14}$$



57. (a)


Given:  $\vec{F}(x, y) = x\vec{i} + y\vec{j}$

Integral is (Say)

$$I = \oint_c \vec{F}(x, y) \cdot (dx\vec{i} + dy\vec{j})$$

Curl,  $\vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & y & 0 \end{vmatrix}$

$$\vec{i}(0-0) - \vec{j}(0-0) + \vec{k}(0-0) = \vec{0}$$



Since, curl  $\vec{F}$  is zero,  $\vec{F}$  is irrotational (or) conservative field.

∴ For any closed path,

$$\oint_c \vec{F} \cdot d\vec{r} = \oint_c \vec{F}(x, y) \cdot (dx\vec{i} + dy\vec{j}) = 0$$



Scan for Video solution



58. (a)

$$\text{Area} = \iint dxdy$$

∴ option (a) is correct

From Green's theorem;

$$\int M dx + N dy = \iint \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dxdy$$

From option b

$$\int_c x dy = \iint (1-0) dxdy = \text{Area}$$

∴ option b is correct

From option c)

$$\int_c y dx = \iint (0-1) dxdy \neq \text{Area}$$

∴ Option c) is incorrect

From option d)

$$\frac{1}{2} \int_c x dy - y dx = \frac{1}{2} \iint 1 - (-1) dxdy =$$

$$\iint dxdy = \text{Area}$$

∴ option d) is also correct.



Scan for Video solution



59. (-3.05 to -2.95)

$$\vec{F} = a_x y \vec{i} - a_y x \vec{j}$$

$$\int_c \vec{F} \cdot d\ell = (a_x y - a_y x)(a_x dx + a_y dy)$$

$$\because d\ell = dx\vec{a}_x + dy\vec{a}_y$$

$$\int_c \vec{F} \cdot d\ell = \int y dx - x dy \quad x \rightarrow -1 \text{ to } 2$$

$$\therefore y = x^2, dy = 2xdx$$

$$= - \int_{x=-1}^{x=2} x^2 dx - x(2xdx)$$

$$= \int_{-1}^2 x^2 dx - \left[ \frac{x^3}{3} \right]_{-1}^2$$

$$= -\frac{1}{3}[8+1]$$

$$\int_c F \cdot d\ell = -3.00$$



Scan for Video solution



60. (3 to 3)

$$\int_c (y+z) dx + (x+z) dy + (x+y) dz = I \text{ (say)}$$

$$\Rightarrow I = \int_c y dx + z dx + x dy + z dy + x dz + y dz$$

$$\Rightarrow I = \int_c (x dy + y dx) + (y dz + z dy) + (x dz + z dx)$$

$$[\because d(xy) = xdy + ydx]$$

$$= \int_c d(xy) + d(yz) + d(xz)$$

$$= [xy + yz + xz]_{(0,0,0)}^{(1,1,1)} = (1+1+1) - (0+0+0) = 3$$



Scan for Video solution


**61. (a)****Given:** The function can be written as

$$\vec{f} = \frac{\vec{r}}{|\vec{r}|^3}$$

where  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  and  $|\vec{r}| = \sqrt{x^2 + y^2 + z^2}$ 

From Gauss-divergence theorem

$$\oint_S \vec{F} \cdot d\vec{s} = \iiint_V (\operatorname{div} \vec{f}) dv$$



$$\operatorname{div} \vec{f} = \nabla \cdot \vec{f} = \frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} + \frac{\partial f_3}{\partial z}$$

$$= \sum \frac{\partial}{\partial x} \left( \frac{x}{(x^2 + y^2 + z^2)^{3/2}} \right)$$

$$= \sum \frac{(x^2 + y^2 + z^2)^{\frac{3}{2}} \cdot (1) - x \cdot \frac{3}{2} \cdot (x^2 + y^2 + z^2)^{\frac{1}{2}} \cdot 2x}{(x^2 + y^2 + z^2)^3}$$

$$= \sum \frac{y^2 + z^2 - 2x^2}{(x^2 + y^2 + z^2)^{5/2}} = 0 \quad (\because \text{The term is cyclic})$$


$$\Rightarrow \iint_S \vec{F} \cdot d\vec{s} = 0$$



Scan for Video solution

**62. (2.25 to 2.25)****Given:**

Solid 's' is as shown below



$$\text{Let } I = \iiint_S x \cdot dx dy dz$$

$$\Rightarrow I = \int_{x=0}^{x=3} \int_{y=0}^{y=1} \int_{z=0}^{z=1-y} x \cdot dx dy dz$$

$$= \int_{x=0}^{x=3} \int_{y=0}^{y=1} \left( x \cdot z \Big|_0^{1-y} \right) dy dx$$

$$\Rightarrow = \int_{x=0}^{x=3} \int_{y=0}^{y=1} x(1-y) dy dx = \int_{x=0}^{x=3} \left( xy - \frac{x \cdot y^2}{2} \right)_0^1 dx$$

$$= \int_{x=0}^{x=3} \left( x - \frac{x}{2} \right) dx = \int_{x=0}^{x=3} \frac{x}{2} dx = \frac{x^2}{4} \Big|_0^3 = \frac{9}{4} = 2.25$$

$$\therefore \iint_S x \cdot dx dy dz = 2.25$$



Scan for Video solution

**63. (139 to 139)**We know  $f = 2x^3 + 3y^2 + 4z$ 

$$\operatorname{grad} f = \frac{\partial f}{\partial x} \hat{i} + \frac{\partial f}{\partial y} \hat{j} + \frac{\partial f}{\partial z} \hat{k}$$


$$\operatorname{grad} f = 6x^2 \hat{i} + 6y \hat{j} + 4 \hat{k}$$

$$\text{also } \vec{dr} = dx \hat{i} + dy \hat{j} + dz \hat{k}$$

$$\int_c \operatorname{grad} f \cdot d\vec{r} = \int_c (6x^2 \hat{i} + 6y \hat{j} + 4 \hat{k})$$

$$(dx \hat{i} + dy \hat{j} + dz \hat{k})$$


$$\begin{aligned}
 &= \int (6x^2 dx + 6y dy + 4dz) = \int d(2x^3 + 3y^2 + 2z^2) \\
 &= 2x^3 + 3y^2 + 2z^2 \Big|_{(-3,-3,-2)}^{(2,6,-1)} \\
 &= 139
 \end{aligned}$$



64. (c)

Given

$$\vec{u} = \frac{1}{3}(-y^3\hat{i} + x^3\hat{j} + z^3\hat{k}) \text{ and } S: x^2 + y^2 + z^2 = 1; z \geq 0$$



Let 'c' be the path enclosing the surface 'S'.  
From stoke's theorem,


$$\begin{aligned}
 \iint_S (\nabla \times \vec{u}) \cdot \hat{n} ds &= \oint_c \vec{u} \cdot d\vec{r} \\
 &= \oint_c \frac{-y^3}{3} dx + \frac{x^3}{3} dy
 \end{aligned}$$

From Green's theorem,

$$\begin{aligned}
 \oint_c \frac{-y^3}{3} dx + \frac{x^3}{3} dy &= \iint_R \frac{\partial}{\partial x} \left( \frac{x^3}{3} \right) - \frac{\partial}{\partial y} \left( \frac{-y^3}{3} \right) dx dy \\
 &=
 \end{aligned}$$


$$\iint_R (x^2 + y^2) dx dy = \frac{\pi}{2}(1)^4 = \frac{\pi}{2}$$

$$\therefore \iint_S (\nabla \times \vec{u}) \cdot \hat{n} ds = \frac{\pi}{2}$$



65. (d)


Given: Integral  $I = \int_C (xdy - ydx)$  and C is as shown



Since 'C' is a closed path, from Green's theorem

$$\begin{aligned}
 \int_C -ydx + xdy &= \iint_R \left( \frac{\partial}{\partial x}(x) - \frac{\partial}{\partial y}(-y) \right) dx dy \\
 &= 2 \cdot \iint_R dx \cdot dy = 2 \times \{\text{Area of region 'R'}\} \\
 &= 2 \times \{\text{Area of rectangle } R_1 + \text{Area of semicircle } R_2\} \\
 &= 2 \times \left\{ (3 \times 2) + \left( \frac{\pi}{2} \times (1)^2 \right) \right\} = 2 \times \left\{ 6 + \frac{\pi}{2} \right\} = 12 + \pi
 \end{aligned}$$

$$\oint_c (xdy - ydx) = 12 + \pi$$




66. (0.60 to 0.70)

$$\vec{F}(\vec{r}) = x^2\hat{i} + y^2\hat{j}$$

Let r be the position vector of point (x, y) then,

$$\vec{r} = x\hat{i} + y\hat{j} \quad d\vec{r} = dx\hat{i} + dy\hat{j}$$

$$\therefore \int_c \vec{F}(\vec{r}) \cdot d\vec{r} = \int_c (x^2\hat{i} + y^2\hat{j})(dx\hat{i} + dy\hat{j})$$



Equation of line from (0, 0) to (1, 1) is  $y = x$

$$\begin{aligned} \therefore \int_C \vec{F} \cdot d\vec{r} &= \int_C x^2 dx + y^2 dy \\ &= \int_{x=0}^{x=1} x^2 dx + x^2 dx \quad \because y = x \Rightarrow dy = dx \\ &= \int_0^1 2x^2 dx = \left[ 2 \frac{x^3}{3} \right]_0^1 = \frac{2}{3} \\ \therefore \int_C \vec{F}(\vec{r}) \cdot d\vec{r} &= 0.67 \end{aligned}$$



Scan for Video solution



### 67. (0 to 0)

$$\int_C (y^2 + 2xy)dx + (x^2 + 2xy)dy$$

$\therefore \vec{F} = (y^2 + 2xy)\hat{i} + (2xy + x^2)\hat{j}$  is conservative

field because  $\text{curl } \vec{F} = 0$ , therefore line integral is path independent and hence we will choose an easiest path i.e. a straight line. The equation of straight line will be,

$$\frac{x}{3} + \frac{y}{3} = 1 \Rightarrow x + y = 3 \Rightarrow dx = -dy$$

$$I = \int (y^2 + 2xy)dx + (2xy + x^2)dy$$

$$\begin{aligned} &= \int_{y=0}^{y=3} y^2 + 2(3-y)y(-dy) + \int_{x=0}^{x=3} (2x(3-x) + x^2)(-dx) \end{aligned}$$

$$= \int_0^3 -y^2 - 6y + 2y^2 + \int_3^0 -6x + 2x^2 - x^2$$

$$= \int_0^3 (y^2 - 6y)dy + \int_3^0 (x^2 - 6x)dx$$

$$= \left[ \frac{y^3}{3} - \frac{6y^2}{2} \right]_0^3 + \left[ \frac{x^3}{3} - \frac{6x^2}{2} \right]_3^0$$

$$= [9 - 27] + -(9 - 27) = 0$$



Scan for Video solution



### 68. (c)

$$\text{Given: } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

and S: closed surface with a volume 'V'

From Gauss-divergence theorem

$$\begin{aligned} \iint_S (\vec{r} \cdot \hat{n}) ds &= \iiint_V (\text{div } \vec{r}) dV = \iiint_V 3 dV \\ &= \iiint_V \left( \frac{\partial}{\partial x}(x) + \frac{\partial}{\partial y}(y) + \frac{\partial}{\partial z}(z) \right) dV = \iiint_V 3 dV \\ &= 3 \times \iiint_V dV = 3V \end{aligned}$$

$$\therefore \iint_S (\vec{r} \cdot \hat{n}) ds = 3V$$



Scan for Video solution



### 69. (225 to 227)

Given,

$$\vec{F} = (x+y)\hat{i} + (x+z)\hat{j} + (y+z)\hat{k}$$

and S:  $x^2 + y^2 + z^2 = 9 \rightarrow \text{Sphere}$

From Gauss-Divergence theorem

$$\iint_S \vec{F} \cdot \hat{n} ds = \iiint_V (\text{div } \vec{F}) dV$$

$$\begin{aligned} \text{div } \vec{F} &= \nabla \cdot \vec{F} = \frac{\partial}{\partial x}(x+y) + \frac{\partial}{\partial y}(x+z) + \frac{\partial}{\partial z}(y+z) \\ &= 1 + 0 + 1 = 2 \end{aligned}$$

$$\begin{aligned} \therefore \iint_S \vec{F} \cdot \hat{n} ds &= \iiint_V 2 \cdot dV = 2 \times \text{Volume of sphere} \\ &= 2 \times \frac{4}{3} \times \pi \times (3)^3 = 72\pi \\ &= 226.19 \text{ Cubic units} \end{aligned}$$

$$\Rightarrow \iint_S \vec{F} \cdot \hat{n} ds = 226.19$$



Scan for Video solution



### 70. (-11.1 to -10.9)

Given line integral is  $I = \int_C (2zdx + 2ydy + 2xdz)$

and A: (0, 2, 1) to point B:(4, 1, -1)

$$I = \int_C (2zdx + 2ydy + 2xdz)$$

$$I = \int_C (2z\hat{i} + 2y\hat{j} + 2x\hat{k}) \cdot (dx\hat{i} + dy\hat{j} + dz\hat{k})$$

$$I = \int_C \vec{F} \cdot d\vec{r} \text{ where } \vec{F} = 2(\vec{z}\hat{i} + \vec{y}\hat{j} + \vec{x}\hat{k})$$

Curl  $\vec{F} = 0$

$$\therefore \vec{F} = \nabla\phi \text{ where } \phi = 2xz + y^2$$

$$\therefore \int_A^B \vec{F} \cdot d\vec{r} = \phi \Big|_A^B = (2xz + y^2) \Big|_A^B$$

$$\Rightarrow \int_C 2zdx + 2ydy + 2xdz = (2xz + y^2) \Big|_{(0,2,1)}^{(4,1,-1)}$$

$$= 2(-4) + 1 - (4) = -11$$

$$\therefore I = \int_C (2zdx + 2ydy + 2xdz) = -11$$




Scan for Video solution



### 71. (b)

The equation of the path will be given as



$$\frac{x-0}{1-0} = \frac{y-0}{1-0} = \frac{z-0}{1-0}$$

Let  $x = y = z = t$

$$\therefore dx = dy = dz = dt$$

When  $x = y = z = 0 \Rightarrow t = 0$  &  $x = y = z = 1 \Rightarrow t = 1$

$$I = \int_C 2xy^2 dx + 2x^2 y dy + dz$$

$$= \int_{t=0}^{t=1} 2t(t)^2 dt + 2(t)^2 t dt + dt$$

$$= \int_0^1 (4t^3 + 1) dt = [t^4 + t]_0^1 = 2$$



Scan for Video solution



### 72. (4.40 to 4.45)

We know, Line integral =  $\int \vec{F} \cdot d\vec{r}$

$$\int_{(0,0,0)}^{(1,1,1)} [5xz\hat{i} + (3x^2 + 2y)\hat{j} + x^2z\hat{k}] \cdot [dx\hat{i} + dy\hat{j} + dz\hat{k}]$$

$$\int 5x^2 dx + (3x^2 + 2y) dy + x^2 z dz$$

$$x = t \quad dx = dt$$

$$y = t^2 \quad dy = 2t dt$$

$$z = t \quad dz = dt$$

$$\vec{F} \cdot d\vec{r} = 5(t)(t) \cdot dt + (3t^2 + 2t^2) 2t dt + t^3 dt$$

$$\int_0^1 5t^2 dt + 11t^3 dt = 0 + (3t^2 + 2t^2) 2t dt$$

$$\left[ \frac{5t^3}{3} + 11 \frac{t^4}{4} \right]_0^1$$

$$\frac{5}{3} + \frac{11}{4} = \frac{53}{12} = 4.4167$$



Scan for Video solution



### 73. (-0.03 to 0.03)

Given, Integral is  $I = \oint_c (xy^2 dx + x^2 y dy)$

$$\Rightarrow I = \oint_c d\left(\frac{x^2 y^2}{2}\right) = 0$$

Since, when the path is closed, initial and final points of integration are same.

$\therefore$  For any closed curve 'c',

$$\oint_c (xy^2 dx + x^2 y dy) = 0$$



Scan for Video solution



### 74. (725.9 to 726.1)

Given

$$\nabla\phi = y\hat{i} + x\hat{j} + xy\hat{k} \text{ and } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\int_C \nabla\phi \cdot d\vec{r} = \int_C yz \cdot dx + xz \cdot dy + xy \cdot dz$$

The curve 'C' is

$$x = t; y = t^2; z = 3t^2; \quad 1 \leq t \leq 3$$

$$\Rightarrow dx = dt; dy = 2t dt; dz = 6t dt$$

$$\begin{aligned}\therefore \int_C \nabla \phi \cdot d\vec{r} &= \int_1^3 3t^4 \cdot dt + 3t^3 \cdot 2t \cdot dt + t^3 \cdot 6t \cdot dt \\ &= \int_1^3 15t^4 \cdot dt = 3t^5 \Big|_1^3 = 3(3^5 - 1) = 726\end{aligned}$$

$\therefore$  Required value = 726



Scan for Video solution




### 75. (15.9 to 16.1)

Given

$$\vec{F} = y\hat{i} + 2x\hat{j}$$

$\vec{r}'$  is unit Tangent vector on 'c'

$$\Rightarrow \vec{r}' \cdot ds = d\vec{s}$$



$$\therefore \oint_C \vec{F} \cdot \vec{r}' ds = \oint_C \vec{F} \cdot d\vec{s} = \oint_C y dx + 2x dy$$

c is closed path

From Green's Theorem,

$$\begin{aligned}\oint_C y dx + 2x dy &= \iint_R \left( \frac{\partial}{\partial x}(2x) - \frac{\partial}{\partial y}(y) \right) dx dy \\ &= \iint_R dx dy = \text{Area of circle}\end{aligned}$$

$$\therefore \oint_C \vec{F} \cdot \vec{r} ds = \text{Area of circle} = \pi \left( \frac{4}{\sqrt{\pi}} \right)^2 = 16$$



Scan for Video solution



### 76. (214 to 218)

Given :

$$\text{Integral } \iint_S \frac{1}{\pi} (9x\hat{i} - 3y\hat{j}) \cdot \hat{n} ds \text{ and } S : x^2 + y^2 + z^2 = 9$$

S is surface of a sphere of radius '3'

By Gauss divergence theorem,

$$\iint_S \frac{1}{\pi} (9x\hat{i} - 3y\hat{j}) \cdot \hat{n} ds = \frac{1}{\pi} \iiint_V \nabla \cdot (9x\hat{i} - 3y\hat{j}) dV$$

$$= \frac{1}{\pi} \cdot \iiint_V 6 \cdot dV$$

$$= \frac{6}{\pi} \times \text{volume of the sphere}$$

$$= \frac{6}{\pi} \times \frac{4}{3} \times \pi \times (3)^3 = 216 \text{ units}$$

$$\therefore \iint_S \frac{1}{\pi} (9x\hat{i} - 3y\hat{j}) \cdot \hat{n} ds = 216$$



Scan for Video solution




### 77. (1.60 to 1.70)

Given

$$\text{Integral } \int_C (3x - 8y^2) dx + (4y - 6xy) dy$$

C: is region bounded by  $x = 0$ ,  $y = 0$ ,  $x + y = 1$



$\Rightarrow \Delta OAB$  is closed path.

From Green's Theorem

$$\oint_C M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx \cdot dy$$

$$\Rightarrow \oint_C (3x - 8y^2) dx + (4y - 6xy) dy$$

$$= \iint_R \left\{ \frac{\partial}{\partial x}(4y - 6xy) - \frac{\partial}{\partial y}(3x - 8y^2) \right\} dx \cdot dy$$

$$= \iint_R (-6y - (-16y)) dx \cdot dy$$

$$= \iint_R 10y dx \cdot dy$$

$$= 10 \times \iint_R y \, dx \cdot dy$$

$= 10 \times (\text{Area of } \Delta OAB) \times (y - \text{coordinate of centroid})$

$$\left( \because \iint_R y \, dA = A \times \bar{y} \right)$$

$$= 10 \times \left( \frac{1}{2} \times 1 \times 1 \right) \times \frac{1}{3} = \frac{10}{6}$$

$$\therefore \iint_C (3x - 8y^2) \, dx + (4y - 6xy) \, dy = \frac{10}{6} = 1.667$$




Scan for Video solution



78. (c)

$$\text{Integral, } \oint_c (y \, dx - x \, dy) \text{ and } c: x^2 + y^2 = \frac{1}{4}$$

Since path 'c' is closed,



By Green's theorem,

$$\oint_c M \, dx + N \, dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \, dx \, dy$$

$$\Rightarrow \oint_c (y \, dx - x \, dy) = \iint_R \left( \frac{\partial}{\partial x}(-x) - \frac{\partial}{\partial y}(y) \right) \, dx \, dy \\ = -2 \cdot \iint_R \, dx \, dy$$

$$= -2 \times \text{Area of Region 'R'} = -2 \times \pi \left( \frac{1}{4} \right) = \frac{-\pi}{2}$$


$$\therefore \oint_c (y \, dx - x \, dy) = \frac{-\pi}{2}$$



Scan for Video solution



79. (b)



Along x-axis

$$y = z = 0$$

$$dy = dz = 0$$

$$I = \int \vec{F} \cdot d\vec{l}$$

$$= \int (y^2 x a_x - y^2 a_y - x^2 a_z) (dx \, a_x + dy \, a_y + dz \, a_z)$$

$$= \int y^2 x \, dx - yz \, dy - x^2 \, dz$$

$$= \int 0^2 x \, dx - 0.0.0 - x^2 \cdot 0$$

$$\therefore I = 0$$



Scan for Video solution



80. (a)

Given:

$$\iint_S \frac{1}{4} (\vec{F} \cdot \hat{n}) \, dA$$

Where S  $\rightarrow$  surface of sphere  $x^2 + y^2 + z^2 = 1$

By Gauss-divergence theorem,

$$\begin{aligned} \iint_S \frac{1}{4} (\vec{F} \cdot \hat{n}) \, dA &= \iiint_V \frac{1}{4} (\nabla \cdot \vec{F}) \, dV \\ &= \frac{1}{4} \times \iiint_V \left( \frac{\partial(x)}{\partial x} + \frac{\partial(y)}{\partial y} + \frac{\partial(z)}{\partial z} \right) \, dV \\ &= \frac{3}{4} \times \iiint_V \, dV = \frac{3}{4} \times \text{volume of sphere} \\ &= \frac{3}{4} \times \frac{4}{3} \times \pi \times (1)^3 = \pi \end{aligned}$$

$$\therefore \iint_S \frac{1}{4} (\vec{F} \cdot \hat{n}) \, dA = \pi$$




Scan for Video solution



81. (d)

**Given:**

's' is a closed surface enclosing volume 'v'

Position vector  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ 

From Gauss-divergence theorem

$$\begin{aligned} \iint_s 5\vec{r} \cdot \vec{n} ds &= \iiint_v 5(\nabla \cdot \vec{r}) dv \\ &= \iiint_v 5 \left( \frac{\partial x}{\partial x} + \frac{\partial y}{\partial y} + \frac{\partial z}{\partial z} \right) dv = 15. \iiint_v dv = 15v \\ \therefore \iint_s 5\vec{r} \cdot \vec{n} ds &= 15v \end{aligned}$$



Scan for Video solution



82. (c)

$$\text{Given } \vec{A} = xy\hat{x} + x^2 \cdot \hat{y}$$

By Green's theorem,

$$\begin{aligned} \oint_c \vec{A} \cdot d\vec{l} &= \oint_c xy dx + x^2 dy \\ &= \iint_R \left( \frac{\partial}{\partial x}(x^2) - \frac{\partial}{\partial y}(xy) \right) dx dy \\ &= \iint_R (2x - x) dx dy = \iint_R x dx dy \end{aligned}$$

$\therefore \iint_R x dx dy = \text{Area of 'R' } \times \text{'x' co-ordinate of centroid.}$

$$= \left( \frac{1}{\sqrt{3}} \times 2 \right) \times \frac{1.5}{\sqrt{3}} = 1$$

$$\therefore \oint_c \vec{A} \cdot d\vec{l} = 1$$




Scan for Video solution



83. (d)

Equation of straight line;



$$\frac{x}{2} + \frac{y}{2} = 1$$

$$x + y = 2 \quad [\because d\vec{l} = dx \hat{x} + dy \hat{y}]$$

$$\int \vec{F}(x, y) \cdot d\vec{l} = \int \{(x^2 + xy)\hat{x} + (y^2 + xy)\hat{y}\} \cdot \{dx \hat{x} + dy \hat{y}\}$$

$$= \int (x^2 + xy) dx + (y^2 + xy) dy$$

$$= \int_{x=0}^{x=2} x^2 + x(2-x) dx + \int_{y=2}^{y=0} y^2 + (2-y)y dy$$

$$[x^2]_0^2 + [y^2]_2^0 = (4-0) + (0-4) = [0]$$



Scan for Video solution




84. (b)

**Given:** Points are P (1, 0) and Q (0, 1)

$$\text{Line integral } I = 2 \times \int_P^Q (xdx + ydy)$$

$$= 2 \times \int_P^Q (x\hat{i} + y\hat{j}) \cdot (dx\hat{i} + dy\hat{j})$$



$$\begin{aligned} \text{Along straight line} \\ x + y &= 1 \\ \Rightarrow y &= 1 - x \\ \Rightarrow dy &= -dx \end{aligned}$$

For the integral  $\int_c \vec{F} \cdot d\vec{r}$ , if  $\text{curl } \vec{F} = \vec{0}$ , then  $\int_c \vec{F} \cdot d\vec{r}$  is independent of path.

Consider  $\text{curl } \vec{F} = \nabla \times (x\hat{i} + y\hat{j}) = \vec{0}$ 

$$\therefore I = 2 \times \int_{y=0}^{y=1} (1-y)(-dy) + ydy$$

$$= 2 \times \int_0^1 -dy + 2y dy = 2 \times \{-y + y^2\Big|_0^1\} = 2 \times 0 = 0$$

$$\therefore 2 \times \int_P^Q x dx + y dy = 0$$



Scan for Video solution




85. (a)

$$\text{Given } g(x, y) = 4x^3 + 10y^4$$

The path is the line segment joining  $(0, 0)$  and  $(1, 2)$

$$\therefore g(x, y) = 4x^3 + 10y^4$$



Along OA,  $y = 2x$

$$\begin{aligned}\therefore \int_O^A g(x, y) dx &= \int_{x=0}^{x=1} 4x^3 + 10(2x)^4 \cdot dx \\ &= \int_{x=0}^{x=1} (4x^3 + 160x^4) \cdot dx \\ &= x^4 + 32 \cdot x^5 \Big|_0^1 = 1 + 32 = 33\end{aligned}$$

$\therefore$  The integration of  $g(x, y)$  along OA is 33.



Scan for Video solution



## CHAPTER

# 6

# COMPLEX CALCULUS

### Complex Numbers and their Properties

- 1. [MCQ]** **[GATE-EC-2023: 1M]**

Let  $w^4 = 16j$ . Which of the following cannot be a value of  $w$ ?

- |                            |                            |
|----------------------------|----------------------------|
| (a) $2e^{\frac{j2\pi}{8}}$ | (b) $2e^{\frac{j\pi}{8}}$  |
| (c) $2e^{\frac{j5\pi}{8}}$ | (d) $2e^{\frac{j9\pi}{8}}$ |

- 2. [MCQ]** **[GATE-ME-2021: 1M]**

Value of  $(1+i)^8$ , where  $i = \sqrt{-1}$ , is equal to

- |          |           |
|----------|-----------|
| (a) 4    | (b) 16    |
| (c) $4i$ | (d) $16i$ |

- 3. [MCQ]** **[GATE-EE-2017 : 1M]**

For a complex number  $z$ ,  $\lim_{z \rightarrow i} \frac{z^2 + 1}{z^3 + 2z - i(z^2 + 2)}$  is

- |           |          |
|-----------|----------|
| (a) $-2i$ | (b) $-i$ |
| (c) $i$   | (d) $2i$ |

- 4. [NAT]** **[GATE-EC-2015:1M]**

Let  $f(z) = \frac{az + b}{cz + d}$ , If  $f(z_1) = f(z_2)$  for all  $z_1 \neq z_2$ ,

$a = 2$ ,  $b = 4$  and  $c = 5$ , then  $d$  should be equal to \_\_\_\_\_.

- 5. [MCQ]** **[GATE-EE-2015 : 1M]**

Given  $f(z) = g(z) + h(z)$  where  $f, g, h$  are complex valued functions of a complex variable  $z$ . Which one of the following statements is TRUE?

- (a) If  $f(z)$  is differentiable at  $z_0$ , then  $g(z)$  and  $h(z)$  are also differentiable at  $z_0$

(b) If  $g(z)$  and  $h(z)$  are differentiable at  $z_0$ , then  $f(z)$  is also differentiable at  $z_0$ .

- (c) If  $f(z)$  is continuous at  $z_0$ , then it is differentiable at  $z_0$ .
- (d) If  $f(z)$  is differentiable at  $z_0$ , then so are its real and imaginary parts.

- 6. [MCQ]** **[GATE-ME-2015: 1M]**

Given two complex numbers  $z_1 = 5 + (5\sqrt{3})i$  and  $z_2 = \frac{2}{\sqrt{3}} + 2i$ , the argument of  $\frac{z_1}{z_2}$  in degrees is

- |        |        |
|--------|--------|
| (a) 0  | (b) 30 |
| (c) 60 | (d) 90 |

- 7. [MCQ]** **[GATE-ME-2014: 1M]**

The argument of the complex number  $\frac{1+i}{1-i}$ , where  $i = \sqrt{-1}$ , is

- |                     |                      |
|---------------------|----------------------|
| (a) $-\pi$          | (b) $\frac{-\pi}{2}$ |
| (c) $\frac{\pi}{2}$ | (d) $\pi$            |

- 8. [MCQ]** **[GATE-ME-2014: 2M]**

If  $z$  is a complex variable, the value of  $\int_5^{3i} \frac{dz}{z}$  is

- |                      |                      |
|----------------------|----------------------|
| (a) $-0.511 - 1.57i$ | (b) $-0.511 + 1.57i$ |
| (c) $0.511 - 1.57i$  | (d) $0.511 + 1.57i$  |

- 9. [MCQ]** **[GATE-EE-2014 : 1M]**

All the values of the multi-valued complex function  $1^i$ , where  $i = \sqrt{-1}$ , are

- |                           |
|---------------------------|
| (a) Purely imaginary      |
| (b) Real and non-negative |



**19. [MCQ]****[GATE-ME-2020: 1M]**

Which of the following function  $f(z)$ , of the complex variable  $z$ , is NOT analytic at all the points of the complex plane?

- (a)  $f(z) = z^2$       (b)  $f(z) = e^z$   
 (c)  $f(z) = \sin z$       (d)  $f(z) = \log z$

**20. [NAT]****[GATE-ME-2020: 2M]**

An analytic function of a complex variable  $z = x + iy$  ( $i = \sqrt{-1}$ ) is defined as

$$f(z) = x^2 - y^2 + i\psi(x, y),$$

where  $\psi(x, y)$ , is a real function. The value of the imaginary part of  $f(z)$  at  $z = (1 + i)$  is \_\_\_\_\_. (Round off to 2 decimal places).

**21. [MCQ]****[GATE-ME-2020: 2M]**

The function  $f(z)$  of complex variable  $z = x + iy$ , where  $i = \sqrt{-1}$ , is given as  $f(z) = (x^3 - 3xy^2) + i v(x, y)$ . For this function to be analytic,  $v(x, y)$  should be

- (a)  $(3xy^2 - y^3) + \text{constant}$   
 (b)  $(3x^2y^2 - y^3) + \text{constant}$   
 (c)  $(x^3 - 3x^2y) + \text{constant}$   
 (d)  $(3x^2y - y^3) + \text{constant}$

**22. [MCQ]****[GATE-EC-2019:1M]**

Which one of the following functions is analytic over the entire complex plane?

- (a)  $\ln(z)$       (b)  $e^{1/z}$   
 (c)  $\frac{1}{1-z}$       (d)  $\cos(z)$

**23. [MCQ]****[GATE-EE-2019 : 1M]**

Which one of the following functions is analytic in the region  $|z| \leq 1$ ?

- (a)  $\frac{z^2 - 1}{z}$       (b)  $\frac{z^2 - 1}{z + 2}$   
 (c)  $\frac{z^2 - 1}{z - 0.5}$       (d)  $\frac{z^2 - 1}{z + j0.5}$

**24. [MCQ]****[GATE-ME-2019: 2M]**

A harmonic function is analytic if it satisfies the Laplace equation. If  $u(x, y) = 2x^2 - 2y^2 + 4xy$  is a


harmonic function, then its conjugate harmonic function  $v(x, y)$  is

- (a)  $4xy - 2x^2 + 2y^2 + \text{constant}$   
 (b)  $4y^2 - 4xy + \text{constant}$   
 (c)  $2x^2 - 2y^2 + xy + \text{constant}$   
 (d)  $-4xy + 2y^2 - 2x^2 + \text{constant}$

**25. [MCQ]****[GATE-ME-2019: 2M]**

An analytic function  $f(z)$  of complex variable  $z = x + iy$  may be written as  $f(z) = u(x, y) + i v(x, y)$ . Then  $u(x, y)$  and  $v(x, y)$  must satisfy

- (a)  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$  and  $\frac{\partial u}{\partial y} = \frac{\partial v}{\partial x}$   
 (b)  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$  and  $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$   
 (c)  $\frac{\partial u}{\partial x} = -\frac{\partial v}{\partial y}$  and  $\frac{\partial u}{\partial y} = \frac{\partial v}{\partial x}$   
 (d)  $\frac{\partial u}{\partial x} = -\frac{\partial v}{\partial y}$  and  $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$

**26. [MCQ]****[GATE-ME-2018: 1M]**

$F(z)$  is a function of the complex variable  $z = x + iy$  given by  $F(z) = iz + k \operatorname{Re}(z) + i \operatorname{Im}(z)$ .

For what value of  $k$  will  $F(z)$  satisfy the Cauchy-Riemann equations?

- (a) 0      (b) 1  
 (c) -1      (d)  $y$

**27. [MCQ]****[GATE-ME-2017: 2M]**

If  $f(z) = (x^2 + ay^2) + i bxy$  is a complex analytic function of  $z = x + iy$ , where  $i = \sqrt{-1}$ , then

- (a)  $a = -1, b = -1$       (b)  $a = -1, b = 2$   
 (c)  $a = 1, b = 2$       (d)  $a = 2, b = 2$

**28. [MCQ]****[GATE-ME-2016: 1M]**

$f(z) = u(x, y) + iv(x, y)$  is an analytic function of complex variable  $z = x + iy$  where  $i = \sqrt{-1}$ .

If  $u(x, y) = 2xy$ , then  $v(x, y)$  may be expressed as

- (a)  $-x^2 + y^2 + \text{constant}$       (b)  $x^2 - y^2 + \text{constant}$   
 (c)  $x^2 + y^2 + \text{constant}$       (d)  $-(x^2 + y^2) + \text{constant}$

**29. [NAT]****[GATE-ME-2016: 1M]**

A function  $f$  of the complex variable  $z = x + iy$ , is given as  $f(x, y) = u(x, y) + iv(x, y)$ , where  $u(x, y) = 2kxy$  and  $v(x, y) = x^2 - y^2$ . The value of  $k$ , for which the function is analytic, is \_\_\_\_\_

**30. [NAT]****[GATE-EC-2016:1M]**

Consider the complex valued function  $f(z) = 2z^3 + b|z|^3$  where  $z$  is a complex variable. The value of  $b$  for which the function  $f(z)$  is analytic is \_\_\_\_\_.

**31. [MCQ]****[GATE-EE-2016 : 1M]**

Consider the function  $f(z) = z + z^*$  where  $z$  is a complex variable and  $z^*$  denotes its complex conjugate. Which one of the following is TRUE?

- (a)  $f(z)$  is both continuous and analytic
- (b)  $f(z)$  is a continuous but not analytic
- (c)  $f(z)$  is not continuous but is analytic
- (d)  $f(z)$  is neither continuous nor analytic

**32. [NAT]****[GATE-EC-2014:2M]**

The real part of an analytic function  $f(z)$  where  $z = x + iy$  is given by  $e^{-y} \cos(x)$ . The imaginary part of  $f(z)$  is

- (a)  $e^y \cos(x)$
- (b)  $e^{-y} \sin(x)$
- (c)  $-e^y \sin(x)$
- (d)  $-e^{-y} \sin(x)$

**33. [MCQ]****[GATE-EE-2014 : 1M]**

Let  $S$  be the set of points in the complex plane corresponding to the unit circle. (That is  $S = \{z : |z| = 1\}$ ). Consider the function  $f(z) = zz^*$  where  $z^*$  denotes the complex conjugate of  $z$ . The  $f(z)$  maps  $S$  to which one of the following in the complex plane

- (a) Unit circle
- (b) Horizontal axis line segment from origin to  $(1, 0)$
- (c) The point  $(1, 0)$
- (d) The entire horizontal axis

**34. [MCQ]****[GATE-ME-2014: 2M]**

An analytic function of a complex variable  $z = x + iy$  is expressed as  $f(z) = u(x, y) + iv(x, y)$ , where  $i = \sqrt{-1}$ . If  $u(x, y) = 2xy$ , then  $v(x, y)$  must be

- (a)  $x^2 + y^2 + \text{constant}$
- (b)  $x^2 - y^2 + \text{constant}$

- (c)  $-x^2 + y^2 + \text{constant}$
- (d)  $-x^2 - y^2 + \text{constant}$

**35. [MCQ]****[GATE-ME-2014: 2M]**

An analytic function of a complex variable  $z = x + iy$  is expressed as  $f(z) = u(x, y) + iv(x, y)$ , where  $i = \sqrt{-1}$ . If  $u(x, y) = x^2 - y^2$ , then expression for  $v(x, y)$  in terms of  $x, y$  and a general constant  $c$  would be

- (a)  $xy + c$
- (b)  $\frac{x^2 + y^2}{2} + c$
- (c)  $2xy + c$
- (d)  $\frac{(x-y)^2}{2} + c$

**36. [MCQ]****[GATE-ME-2009: 2M]**

An analytic function of a complex variable  $z = x + iy$  is expressed as  $f(z) = u(x, y) + iv(x, y)$  wheere  $i = \sqrt{-1}$ . If  $u = xy$ , then the expression for  $v$  should be

- (a)  $\frac{(x+y)^2}{2} + k$
- (b)  $\frac{x^2 - y^2}{2} + k$
- (c)  $\frac{y^2 - x^2}{2} + k$
- (d)  $\frac{(x-y)^2}{2} + k$

**37. [MCQ]****[GATE-EC-2008: 1M]**

The equation  $\sin(z) = 10$  has

- (a) no real (or) complex solution
- (b) exactly two distinct complex solutions
- (c) a unique solution
- (d) an infinite number of complex solutions

**Complex Integration****38. [MCQ]****[GATE-EC-2023:1M]**

The value of the contour integral,  $\oint \left( \frac{z+2}{z^2+2z+2} \right) dz$

, where the contour  $C$  is  $\left\{ z : \left| z + 1 - \frac{3}{2}j \right| = 1 \right\}$  taken in

the counter clockwise direction, is


- (a)  $-\pi(1+j)$
- (b)  $\pi(1+j)$
- (c)  $\pi(1-j)$
- (d)  $-\pi(1-j)$

**39. [NAT]****[GATE-EC-2022:2M]**

A simple closed path  $C$  in the complex plane is shown in the figure. If

$$\oint_C \frac{2^z}{z^2 - 1} dz = -i\pi A,$$

where  $i = \sqrt{-1}$ , then the value of  $A$  is \_\_\_\_\_ (rounded off to two decimal places).

**40. [MCQ]****[GATE-ME-2022: 2M]**

The value of the integral

$$\oint \left( \frac{6z}{2z^4 - 3z^3 + 7z^2 - 3z + 5} \right) dz$$

evaluated over a counter-clockwise circular contour in the complex plane enclosing only the pole  $z = i$ , where  $i$  is the imaginary unit, is

- |                   |                  |
|-------------------|------------------|
| (a) $(-1 + i)\pi$ | (b) $(1+i)\pi$   |
| (c) $2(1 - i)\pi$ | (d) $(2 + i)\pi$ |

**41. [NAT]****[GATE-ME-2022: 2M]**

Given  $z = x + iy$ ,  $i = \sqrt{-1}$ .  $C$  is a circle of radius 2 with the center at the origin.

If the contour  $C$  is traversed anticlockwise, then the value of the integral  $\frac{1}{2\pi j} \int_C \frac{1}{(z-i)(z+4i)} dz$  is \_\_\_\_\_.  
(Round off to one decimal place).

**42. [MCQ]****[GATE-ME-2021: 2M]**

Let  $C$  represent the unit circle centered at origin in the complex plane and complex variable,  $z = x + iy$ . The value of the contour integral  $\oint_C \frac{\cosh 3z}{2z} dz$  (where integration is taken counter clockwise) is

- |             |              |
|-------------|--------------|
| (a) 0       | (b) 2        |
| (c) $\pi i$ | (d) $2\pi i$ |

**43. [MCQ]****[GATE-EE-2021: 1 M]**

Let  $(-1 - j)$ ,  $(3 - j)$ ,  $(3 + j)$  and  $(-1 + j)$  be the vertices of a rectangle  $C$  in the complex plane. Assuming that  $C$  is traversed in counter-clockwise direction, the value of the contour integral  $\oint_C \frac{dz}{z^2(z-4)}$  is

- |               |               |
|---------------|---------------|
| (a) $j\pi/2$  | (b) 0         |
| (c) $-j\pi/8$ | (d) $j\pi/16$ |

**44. [MCQ]****[GATE-EE-2020 : 1M]**

The value of the following complex integral, with  $C$  representing the unit circle centered at origin in the counter clock wise sense, is:

$$\oint_C \frac{z^2 + 1}{z^2 - 2z} dz$$

- |              |               |
|--------------|---------------|
| (a) $8\pi i$ | (b) $-8\pi i$ |
| (c) $-\pi i$ | (d) $\pi i$   |

**45. [MCQ]****[GATE-EE-2019 : 2M]**

The closed loop line integral

$$\oint_{|z|=5} \frac{z^3 + z^2 + 8}{z + 2} dz$$

evaluated counter-clockwise, is

- |               |               |
|---------------|---------------|
| (a) $+8\pi j$ | (b) $-8\pi j$ |
| (c) $-4j\pi$  | (d) $+4j\pi$  |

**46. [NAT]****[GATE-EC-2019: 1M]**


The value of the contour integral

$$\frac{1}{2\pi j} \oint_C \left( z + \frac{1}{z} \right)^2 dz$$

Evaluated over the unit circle  $|z| = 1$  is \_\_\_\_\_.

**47. [NAT]****[GATE-EC-2018:2M]**

The contour  $C$  given below is on the complex plane  $z = x + iy$ , where,  $j = \sqrt{-1}$ .



The value of the integral  $\frac{1}{\pi j} \oint_C \frac{dz}{z^2 - 1}$  is \_\_\_\_\_

**48. [MCQ]****[GATE-EE-2018 : 1M]**

The value of the integral  $\oint_C \frac{z+1}{z^2-4} dz$  in counter clockwise direction around a circle C of radius 1 with center at the point  $z = -2$  is

- (a)  $\frac{\pi i}{2}$       (b)  $2\pi i$   
 (c)  $-\frac{\pi i}{2}$       (d)  $-2\pi i$

**49. [MCQ]****[GATE-EE-2018 : 2M]**

If C is a circle  $|Z|=4$  and  $f(z) = \frac{z^2}{(z^2 - 3z + 2)^2}$ ,

then  $\oint_C f(z) dz$  is

- (a) 1      (b) 0  
 (c)  $-1$       (d)  $-2$

**50. [MCQ]****[GATE-ME-2018: 2M]**

Let z be a complex variable. For a counter-clockwise integration around a unit circle C, centred at origin,

$\oint_C \frac{1}{5z-4} dz = A \pi i$  the value of A is

- (a)  $2/5$       (b)  $1/2$   
 (c) 2      (d)  $4/5$

**51. [MCQ]****[GATE-EC-2017:1M]**

The residues of a function  $f(z) = \frac{1}{(z-4)(z+1)^3}$  are,

- (a)  $\frac{-1}{27}$  and  $\frac{-1}{125}$       (b)  $\frac{1}{125}$  and  $\frac{-1}{125}$   
 (c)  $\frac{-1}{27}$  and  $\frac{1}{5}$       (d)  $\frac{1}{125}$  and  $\frac{-1}{5}$

**52. [MCQ]****[GATE-EC-2017:2M]**


An integral I over a counter clockwise circle C is given by  $I = \oint_C \frac{z^2-1}{z^2+1} e^z dz$ . If C is defined as  $|z| = 3$ ,

then the value of I is

- (a)  $-\pi i \sin(1)$       (b)  $-2\pi i \sin(1)$   
 (c)  $-3\pi i \sin(1)$       (d)  $-4\pi i \sin(1)$

**53. [MCQ]****[GATE-EE-2017 : 2M]**

Consider the line integral  $I = \int_C (x^2 + iy^2) dz$ , where  $z = x + iy$ . The line c is shown in the figure below.



The value of I is

- (a)  $\frac{1}{2} i$       (b)  $\frac{2}{3} i$   
 (c)  $\frac{3}{4} i$       (d)  $\frac{4}{5} i$

**54. [MCQ]****[GATE-EE-2017 : 2M]**

The value of the contour integral in the complex-plane  $\oint \frac{z^3 - 2z + 3}{z-2} dz$  along the contour  $|z| = 3$ , taken counter clockwise is:

- (a)  $-18\pi i$       (b) 0  
 (c)  $14\pi i$       (d)  $48\pi i$

**55. [MCQ]****[GATE-EE-2016 : 1M]**

The value of the integral  $\oint_C \frac{2z+5}{\left(z-\frac{1}{2}\right)(z^2-4z+5)} dz$

over the contour  $|z| = 1$ , taken in the anti-clockwise direction, would be

- (a)  $\frac{24\pi i}{13}$       (b)  $\frac{48\pi i}{13}$   
 (c)  $\frac{24}{13}$       (d)  $\frac{12}{13}$

**56. [MCQ]****[GATE-EC-2016:2M]**

The values of the integral  $\frac{1}{2\pi j} \oint_C \frac{e^z}{z-2} dz$  along a closed contour c in anti-clockwise direction for

- (i) The point  $z_0 = 2$  inside the contour c, and  
 (ii) The point  $z_0 = 2$  outside the contour c, respectively, are

- (a) (i) 2.72, (ii) 0  
 (c) (i) 0, (ii) 2.72

- (b) (i) 7.39, (ii) 0  
 (d) (i) 0, (ii) 7.39

**57. [NAT]****[GATE-EC-2016:2M]**


In the following integral, the contour C encloses the points  $2\pi j$  and  $-2\pi j$ .

$-\frac{1}{2\pi} \oint_C \frac{\sin z}{(z - 2\pi j)^3} dz$ . The value of the integral is \_\_\_\_\_.

**58. [MCQ]****[GATE-ME-2016: 2M]**

The value of  $\oint_{\Gamma} \frac{3z - 5}{(z - 1)(z - 2)} dz$  along a closed path

$\Gamma$  is equal to  $(4\pi i)$ , where  $z = x + iy$  and  $i = \sqrt{-1}$ . The correct path  $\Gamma$  is

**59. [MCQ]****[GATE-ME- 2016: 2M]**

The value of the integral

$$\int_{-\infty}^{\infty} \frac{\sin x}{x^2 + 2x + 2} dx$$

evaluated using contour integration and the residue theorem is

- (a)  $-\pi \sin(1)/e$   
 (c)  $\sin(1)/e$

- (b)  $-\pi \cos(1)/e$   
 (d)  $\cos(1)/e$

**60. [MCQ]****[GATE-EC-2015:1M]**

Let  $z = x + iy$  be a complex variable. Consider that contour integration performed along the unit circle in anticlockwise direction. Which one of the following statements is NOT TRUE?

- (a) The residue of  $\frac{z}{z^2 - 1}$  at  $z = 1$  is  $\frac{1}{2}$   
 (b)  $\oint_c z^2 dz = 0$   
 (c)  $\frac{1}{2\pi i} \oint_c \frac{1}{z} dz = 1$   
 (d)  $\bar{z}$  (complex conjugate of  $z$ ) is an analytical function

**61. [MCQ]****[GATE-EC-2015:1M]**

If  $C$  is a circle of radius  $r$  with centre  $z_0$ , in the complex  $z$ -plane and if  $n$  is a non-zero integer, then

$$\oint_c \frac{dz}{(z - z_0)^{n+1}}$$

- equals  
 (a)  $2\pi n j$   
 (b) 0  
 (c)  $\frac{n j}{2\pi}$   
 (d)  $2\pi n$

**62. [NAT]****[GATE-EC-2015:2M]**

If  $C$  denotes the counter clockwise unit circle, the value of the contour integral  $\frac{1}{2\pi i} \oint_C \operatorname{Re}\{z\} dz$  is \_\_\_\_\_.

**63. [MCQ]****[GATE-EC-2014:1M]**

$C$  is a closed path in the  $z$ -plane given by  $|z| = 3$ . The

$$\text{value of the integral } \oint_c \left( \frac{z^2 - z + 4j}{z + 2j} \right) dz$$

- (a)  $-4\pi(1+j2)$   
 (c)  $-4\pi(3+j2)$   
 (b)  $4\pi(3-j2)$   
 (d)  $4\pi(1-j2)$

**64. [MCQ]****[GATE-EE-2014 : 2M]**

Integration of the complex function  $f(z) = \frac{z}{z^2 - 1}$ , in the counter clockwise direction, around  $|z - 1| = 1$ , is

- (a)  $-\pi i$   
 (c)  $\pi i$

- (b) 0  
 (d)  $2\pi i$

**65. [MCQ]****[GATE-EE-2013 : 2M]**

$\oint \frac{z^2 - 4}{z^2 + 4} dz$  evaluated anticlockwise around the circle

- $|z - i| = 2$ , where  $i = \sqrt{-1}$ , is  
 (a)  $-4\pi$   
 (b) 0  
 (c)  $2 + \pi$   
 (d)  $2 + 2i$

**66. [MCQ]****[GATE-EE-2012:1M]**

Given  $f(z) = \frac{1}{z+1} - \frac{2}{z+3}$ . If C is a counterclockwise

path in the z-plane such that  $|z + 1| = 1$ , the value of

$$\frac{1}{2\pi j} \oint_C f(z) dz$$

- (a) -2  
 (b) -1  
 (c) 1  
 (d) 2

**67. [MCQ]****[GATE-EC-2012:1M]**

Given  $f(z) = \frac{1}{z+1} - \frac{2}{z+3}$ . If C is counterclockwise

path in the z-plane such that  $|z + 1| = 1$ , the value of

$$\frac{1}{2\pi j} \oint_C f(z) dz$$

- (a) -2  
 (b) -1  
 (c) 1  
 (d) 2

**68. [MCQ]****[GATE-EC-2011:1M]**

The value of the integral  $\oint_C \frac{-3z + 4}{(z^2 + 4z + 5)} dz$  where C is

the circle  $|z| = 1$  is given by

- (a) 0  
 (c)  $4/5$   
 (b)  $1/10$   
 (d) 1

**69. [MCQ]****[GATE-EC-2010:2M]**

The residues of a complex function

$$X(z) = \frac{1-2z}{z(z-1)(z-2)}$$

- at its poles are  
 (a)  $\frac{1}{2}, -\frac{1}{2}$  and 1  
 (b)  $\frac{1}{2}, \frac{1}{2}$  and -1

- (c)  $\frac{1}{2}, 1$  and  $-\frac{3}{2}$   
 (d)  $\frac{1}{2}, -1$  and  $-\frac{3}{2}$

**70. [MCQ]****[GATE-EC-2009:1M]**

If  $f(z) = c_0 + c_1 z^{-1}$ , then  $\oint_C \frac{1+f(z)}{z} dz$  is given by unit circle

- (a)  $2\pi c_1$   
 (c)  $2\pi j c_1$   
 (b)  $2\pi (1+c_0)$   
 (d)  $2\pi j (1+c_0)$

**71. [MCQ]****[GATE-EC-2008:2M]**

The residue of the function  $f(z) = \frac{1}{(z+2)^2(z-2)^2}$

at  $z = 2$  is

- (a)  $-\frac{1}{32}$   
 (b)  $-\frac{1}{16}$   
 (c)  $\frac{1}{16}$   
 (d)  $\frac{1}{32}$

**72. [MCQ]****[GATE-EE-2008 : 2M]**

Given  $X(z) = \frac{z}{(z-a)^2}$  with  $|z| > a$ , the residue of

$X(z) z^{n-1}$  at  $z = a$  for  $n \geq 0$  will be

- (a)  $a^{n-1}$   
 (b)  $a^n$   
 (c)  $n a^n$   
 (d)  $n a^{n-1}$

**73. [MCQ]****[GATE-ME-2008: 2M]**

The integral  $\oint_C f(z) dz$  evaluated around the unit

circle on the complex plane for  $f(z) = \frac{\cos z}{z}$  is

- (a)  $2\pi i$   
 (c)  $-2\pi i$   
 (b)  $4\pi i$   
 (d) 0

**Series of Complex Functions****74. [NAT]****[GATE-EC-2016:1M]**

For  $f(z) = \frac{\sin(z)}{z^2}$ , the residue of the pole at  $z = 0$  is

\_\_\_\_\_.


**ANSWER KEY**

- |                      |                         |                    |                    |
|----------------------|-------------------------|--------------------|--------------------|
| 1. (a)               | 2. (b)                  | 3. (d)             | 4. (10 to 10)      |
| 5. (b)               | 6. (a)                  | 7. (c)             | 8. (b)             |
| 9. (b)               | 10. (b)                 | 11. (a)            | 12. (a)            |
| 13. (d)              | 14. (d)                 | 15. (a)            | 16. (b)            |
| 17. (1.999 to 2.001) | 18. (d)                 | 19. (d)            | 20. (1.99 to 2.01) |
| 21. (d)              | 22. (d)                 | 23. (b)            | 24. (a)            |
| 25. (b)              | 26. (b)                 | 27. (b)            | 28. (a)            |
| 29. (-1.1 to -0.9)   | 30. (0 to 0)            | 31. (b)            | 32. (b)            |
| 33. (c)              | 34. (c)                 | 35. (c)            | 36. (c)            |
| 37. (d)              | 38. (b)                 | 39. (0.50 to 0.50) | 40. (a)            |
| 41. (0.2 to 0.2)     | 42. (c)                 | 43. (c)            | 44. (c)            |
| 45. (b)              | 46. (-0.0001 to 0.0001) | 47. (2 to 2)       | 48. (a)            |
| 49. (b)              | 50. (a)                 | 51. (b)            | 52. (d)            |
| 53. (b)              | 54. (c)                 | 55. (b)            | 56. (b)            |
| 57. (-136 to -132)   | 58. (b)                 | 59. (a)            | 60. (d)            |
| 61. (b)              | 62. (0.5 to 0.5)        | 63. (c)            | 64. (c)            |
| 65. (a)              | 66. (c)                 | 67. (c)            | 68. (a)            |
| 69. (c)              | 70. (d)                 | 71. (a)            | 72. (d)            |
| 73. (a)              | 74. (1 to 1)            |                    |                    |


**SOLUTIONS**
**1. (a)**Given  $W^4 = 16j$ 

$$\Rightarrow W = (16)^{1/4} \cdot (e^{j\pi/2})^{1/4}$$

By De-movier's theorem

$$\Rightarrow W = 2 \text{ cis} \left( \frac{2x\pi + \frac{\pi}{2}}{4} \right) \text{ where } n = 0, 1, 2, 3$$

$$\Rightarrow W = 2 \text{ cis} \left( \frac{\pi}{8} \right), 2 \cdot \text{Cis} \left( \frac{5\pi}{8} \right), 2 \cdot \text{Cis} \left( \frac{9\pi}{8} \right),$$

$$2 \cdot \text{Cis} \left( \frac{13\pi}{8} \right)$$

$$\Rightarrow W = 2 \cdot e^{j\pi/8}, 2 \cdot e^{j5\pi/8}, 2 \cdot e^{j9\pi/8}, 2 \cdot e^{j15\pi/8}$$

Out of the given options,  $2 \cdot e^{j2\pi/8}$  is not in the above values of z.

$\therefore 2 \cdot e^{j2\pi/8}$  is not value of 'w'.



Scan for Video solution



2. (b)

$$\begin{aligned}
 (1+i)^8 &= \left(\sqrt{2}\left(\frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}\right)\right)^8 \\
 &= \left(\sqrt{2}\left(\cos\frac{\pi}{4} + i \cdot \sin\frac{\pi}{4}\right)\right)^8 (\because e^{i\theta} = \cos\theta + i \sin\theta) \\
 &= \left(\sqrt{2} \cdot e^{i\frac{\pi}{4}}\right)^8 = 2^4 \cdot e^{i\left(\frac{8\pi}{4}\right)} \\
 &= 16 \cdot (e^{i(2\pi)}) \\
 &= 16\{\cos 2\pi + i \cdot \sin 2\pi\} \\
 &= 16(1) \\
 \therefore (1+i)^8 &= 16
 \end{aligned}$$



Scan for Video solution



3. (d)

$$\lim_{z \rightarrow i} \frac{z^2 + 1}{z^3 + 2z - i(z^2 + 2)}$$

Since it is  $\frac{0}{0}$  form use L'Hospital's rule and differentiate both numerator and denominator.

$$\lim_{z \rightarrow i} \frac{2z}{3z^2 + 2 - 2zi} = \frac{2i}{3i^2 + 2 - 2i^2} = \frac{2i}{-3 + 2 + 2} = 2i$$



Scan for Video solution



4. (10 to 10)

For all  $z_1 \neq z_2$  it is given that

$$F(z_1) = f(z_2)$$

$\Rightarrow f(z)$  is a constant function

$$\Rightarrow f'(z) = 0 \Rightarrow \frac{d}{dz} \left( \frac{az+b}{cz+d} \right) = 0$$

$$\Rightarrow \frac{ad-bc}{(cz+d)^2} = 0$$

$$\Rightarrow ad - bc = 0 \Rightarrow 2d - (20) = 0$$

$$\Rightarrow d = 10$$

Rationalizing the denominator

$$\Rightarrow z = \frac{1+i}{1-i} \left( \frac{1+i}{1+i} \right) = \frac{(1+i)^2}{1-i^2} = \frac{2i}{2} = i$$

$$\therefore z = i = 0 + 1i$$

$$\Rightarrow \arg(z) = \tan^{-1}(1/0) = \tan^{-1}\infty = \frac{\pi}{2}$$



Scan for Video solution



8. (b)

$$\text{Given: Integral } I = \int_5^{3i} \frac{dz}{z}$$

$$\Rightarrow I = \ln z \Big|_5^{3i} = \ln(3i) - \ln(5) = \ln\left(\frac{3}{5} \cdot i\right)$$

$$\Rightarrow I = \ln\left(\frac{3}{5}\right) + \ln i = \ln\left(\frac{3}{5}\right) + \ln e^{i\pi/2} \quad (\text{since } e^{i\pi/2} = i)$$

$$\Rightarrow I = \ln 0.6 + i\left(\frac{\pi}{2}\right)$$

$$\Rightarrow I = -0.5108 + i(1.5707)$$

$$\therefore \int_5^{3i} \frac{dz}{z} = -0.511 + i(1.571)$$



Scan for Video solution



9. (b)

$$\begin{aligned} f(z) &= 1^i \\ &= (e^{2\pi i})^i = e^{-2n\pi} \end{aligned}$$

Where n is any integer

Hence, f is real and non-negative.



Scan for Video solution



10. (b)

$$(-i)^{\frac{1}{2}} = \left[ e^{-i\pi/2} \right]^{\frac{1}{2}}$$

$$= \left[ \cos\left(-\frac{\pi}{2}\right) + i \sin\left(-\frac{\pi}{2}\right) \right]^{\frac{1}{2}}$$

$$= \left[ \cos\left(2n\pi - \frac{\pi}{2}\right) + i \sin\left(2n\pi - \frac{\pi}{2}\right) \right]^{\frac{1}{2}}$$

$$= \left[ \cos(4n-1)\frac{\pi}{4} + i \sin(4n-1)\frac{\pi}{4} \right]$$

(By De-Moivre's Theorem)

Now put n = 0 and 1

$$\sqrt{-i} = \cos\left(-\frac{\pi}{4}\right) + i \sin\left(-\frac{\pi}{4}\right), \cos\left(\frac{3\pi}{4}\right) + i \sin\left(\frac{3\pi}{4}\right)$$



Scan for Video solution



11. (a)

If  $x = \sqrt{-1} = i$

$$\therefore x^x = (i)^i = \left( e^{\frac{i\pi}{2}} \right)^i = e^{\frac{-\pi}{2}}$$

$$\left[ \therefore e^{\frac{i\pi}{2}} = \cos \frac{\pi}{2} + i \sin \frac{\pi}{2} = i \right]$$



Scan for Video solution



12. (a)

$$\text{Given: } x = \sqrt{-1} = j, (\because e^{i\pi/2} = \cos \pi/2 + j \sin \pi/2 = j)$$

$$\Rightarrow x^x = j^j = \left( e^{i\pi/2} \right)^j = e^{j^2\pi/2} = e^{-\pi/2}$$

For  $x = j$ ;  $x^x = e^{-\pi/2}$



Scan for Video solution



13. (d)

$$\Rightarrow x^3 + x^2 + x + 1 = 0$$

$$\Rightarrow x^2(x+1) + (x+1) = 0$$

$$\Rightarrow (x+1)(x^2+1) = 0$$

$$\Rightarrow (x+1)(x+j)(x-j) = 0$$

$$\Rightarrow x = -1, +j, -j$$



Scan for Video solution

**14. (d)**

We know,  $z = x + iy$  is in I<sup>st</sup> quadrant & within unit circle.

$$\therefore 0 < x < 1, 0 < y < 1 \text{ & } 0 < \sqrt{x^2 + y^2} < 1$$

$$\text{Now } y = \frac{1}{z} = \frac{1}{x + iy}$$

$$= \frac{1}{x + iy} \times \frac{x - iy}{x - iy} = \frac{x}{x^2 + y^2} - \frac{iy}{x^2 + y^2}$$

Since real part is positive and imaginary part is negative

$$\therefore y = \frac{1}{z} \text{ lies in IV<sup>th</sup> quadrant.}$$

$$\left| \frac{1}{z} \right| = \left| \frac{1}{x + iy} \right| = \frac{1}{\sqrt{x^2 + y^2}} > 1 \quad (\because 0 < \sqrt{x^2 + y^2} < 1)$$

$$\therefore y = \frac{1}{z} \text{ lies outside the unit circle.}$$



Scan for Video solution

**15. (a)****Given:**

$$(i+1). (2-5i) = 2 - 5i + i \quad (2-5i) = 2 - 5i + 2i + 5 \\ = 7 - 3i$$

$$\therefore (1+i) (2-5i) = 7 - 3i$$



Scan for Video solution

**16. (b)****Given**

$$\text{complex number is } \frac{3+4i}{1-2i} = z \text{ (say)}$$

$$\Rightarrow |z| = \left| \frac{3+4i}{1-2i} \right| = \frac{|3+4i|}{|1-2i|} \quad \left\{ \because \left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \right\}$$

$$\Rightarrow |z| = \frac{\sqrt{3^2 + (4)^2}}{\sqrt{1^2 + (-2)^2}} = \frac{\sqrt{25}}{\sqrt{5}} = \sqrt{5}$$

$$\Rightarrow \left| \frac{3+4i}{1-2i} \right| = \sqrt{5}$$



Scan for Video solution

**17. (1.999 to 2.001)****Given**

$$f(z) = e^{-kx} (\cos 2y - i \sin 2y) \\ = e^{-kx} \cdot \cos 2y + i (-e^{-kx} \cdot \sin 2y)$$

For  $f(z) = u(x, y) + i.v(x, y)$  to be analytic,

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\Rightarrow \frac{\partial}{\partial x} (e^{-kx} \cdot \cos 2y) = \frac{\partial}{\partial y} (-e^{-kx} \cdot \sin 2y)$$

$$\Rightarrow \cos 2y \cdot (-k \cdot e^{-kx}) = -e^{-kx} \cdot 2 \cos 2y$$

$$\Rightarrow k = 2$$



Scan for Video solution

**18. (d)**

$$ax^3 + bx^2 + cx + d = 0$$


Let the roots be p, q and r

We know, sum of the roots  $(p + q + r) = -b/a$ Product of the roots  $(pqr) = -d/a$ 

$$p(z) = z^3 + (1+j)z^2 + (2+j)z + 3$$

$$\therefore \text{Sum of roots} = -(1+j)$$

$$\text{Product of roots} = -3$$



$$P(\bar{z}) = (\bar{z})^3 + (1+j)(\bar{z})^2 + (z+j)\bar{z} + 3$$

$\therefore p(z) \neq p(\bar{z})$ . So only option - d is correct.



Scan for Video solution



$\therefore$

$$\psi(1, 1) = 2$$



Scan for Video solution



### 19. (d)

Given

$$(i) f(z) = z^2 \Rightarrow f'(z) = 2z \quad \exists \forall z \in C$$

$$(ii) f(z) = e^z \Rightarrow f'(z) = e^z \quad \exists \forall z \in C$$

$$(iii) f(z) = \sin z \Rightarrow f'(z) = \cos z \quad \exists \forall z \in C$$

$$(iv) f(z) = \log_e z \Rightarrow f'(z) = \frac{1}{z} \quad \nexists \text{ when } z = 0$$

$\therefore f(z) = \log_e z$  is not analytic  $z = 0$



Scan for Video solution



### 20. (1.99 to 2.01)

Given

$$f(z) = (x^2 - y^2) + i\psi(x, y) \\ = u(x, y) + i\psi(x, y)$$

$$\Rightarrow \frac{\partial u}{\partial x} = 2x; \text{ and } \frac{\partial u}{\partial y} = -2y$$

When  $f(z) = u + iv$  is analytic,

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial u}{\partial x} + i \left( \frac{-\partial u}{\partial y} \right)$$

$$\Rightarrow 2x - i(-2y) = 2(x + iy) = 2z$$

Integrating both sides

$$\Rightarrow \int f'(z) \cdot dz = \int 2z \cdot dz$$

$$\Rightarrow f(z) = z^2 + c$$

$$\Rightarrow f(z) = (x + iy)^2 + c = (x^2 - y^2) + i(2xy) + c$$

By taking  $c = 0$

$$\psi(x, y) = 2xy$$

At  $z = 1 + i$ ; i.e.,  $x = 1$  &  $y = 1$

$$\psi(1, 1) = 2(1)(1) = 2$$

### 21. (d)

Given  $f(z) = x^3 - 3xy^2 + i\psi(x, y)$  is analytic.

$$\Rightarrow f'(z) = 3x^2 - 3y^2 + i \frac{\partial \psi}{\partial x}$$

For analytic function  $f(z) = u + iv$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$$\Rightarrow f'(z) = 3x^2 - 3y^2 + i \left( \frac{-\partial u}{\partial y} \right)$$

$$\Rightarrow f'(z) = 3x^2 - 3y^2 - i(-6xy)$$

Apply Milne- Thomson method

Replace 'x' with 'z' and 'y' with '0'.

$$\Rightarrow \int f'(z) dz = \int 3z^2 dz$$

$$\Rightarrow f(z) = z^3 + c$$

$$\Rightarrow f(z) = (x + iy)^3 + c$$

$$= x^3 - 3xy^2 + i(3x^2y - y^3) + c$$

$$\therefore v(x, y) = 3x^2y - y^3 + \text{constant}$$



Scan for Video solution



### 22. (d)

$f(z) = \cos z \Rightarrow f'(z) = -\sin z$ ;  $-\sin z$  exists for all 'z'

$$f(z) = \ln(z) \Rightarrow f'(z) = \frac{1}{z}; f'(z) \text{ doesn't exist at } z = 0$$

$$f(z) = \frac{1}{1-z} \Rightarrow f'(z) = \frac{1}{(1-z)^2}; f'(z) \text{ doesn't exist at } z = 1$$

$$f(z) = e^{1/z} \Rightarrow f'(z) = \frac{-e^{1/z}}{z^2}; f'(z) \text{ doesn't exist at } z = 0$$

$\therefore f(z) = \cos z$  is the function that is analytic over the complete complex plane.



Scan for Video solution



### 23. (b)

For options a, c and d the singular points are lying within the given region  $|z| < 1$

$$\Rightarrow |z| \leq 1$$

$$\Rightarrow |x + iy| \leq 1$$

$$\Rightarrow x^2 + y^2 \leq 1$$

$|z| \leq 1$  represents a Circle with unit radius

Option a)  $z = 0$

Option b)  $z = -2$

Option c)  $z = 0.5$

Option d)  $z = -0.5j$

Only option b) will be analytic because it's singular point  $z = -2$  is not lying in the region  $|z| \leq 1$ .



Scan for Video solution



### 24. (a)

If  $f(z) = u + iv$  is analytic function, then  $u$  and 'v' are Harmonic conjugate pair

Given  $u = 2x^2 - 2y^2 + 4xy$

$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

$$\Rightarrow f'(z) = 4x + 4y - i(-4y + 4x)$$

Applying Milne Thomson method, Replace 'x' with 'z' & 'y' with '0'.

$$\Rightarrow \int f'(z) dz = \int (4z - i4z) dz$$

$$\Rightarrow f(z) = (1 - i)2z^2 + c$$

$$= (2 - 2i)(x^2 - y^2 + i2xy) + c$$

$$= 2(x^2 - y^2) + 4xy + i(4xy - 2x^2 + 2y^2) + c$$

$$\Rightarrow v(x, y) = 4xy - 2x^2 + 2y^2 + c$$

Scan for Video solution

### 25. (b)

Given

$$f(z) = u(x, y) + iv(x, y)$$

For  $f(z)$  to be analytic function,  $u(x, y)$  and  $v(x, y)$  should satisfy Cauchy-Riemann equations.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Scan for Video solution

### 26. (b)

Given,

$$\begin{aligned} F(z) &= iz + k \cdot \operatorname{Re}(z) + i \operatorname{Im}(z) \\ &= i(x + iy) + kx + iy \\ &= (kx - y) + i(x + y) \end{aligned}$$

For satisfying Cauchy-Riemann equations

$$\frac{\partial}{\partial x}(kx - y) = \frac{\partial}{\partial y}(x + y)$$

$$\Rightarrow k = 1$$

Scan for Video solution

### 27. (b)

Given,

$$f(z) = (x^2 + ay^2) + ibxy \text{ is analytic function.}$$

$$\therefore \frac{\partial}{\partial x}(x^2 + ay^2) = \frac{\partial}{\partial y}(bxy)$$

$$\Rightarrow 2x = bx$$


$$\Rightarrow b = 2$$

$$\text{and } \frac{\partial}{\partial y}(x^2 + ay^2) = -\frac{\partial}{\partial x}(bxy)$$

$$\Rightarrow 2ay = -by$$

$$\Rightarrow a = \frac{-b}{2} = -1$$

$$\Rightarrow [a = -1]$$



28. (a)

**Given**

$$f(z) = u(x, y) + i v(x, y) \text{ and } u(x, y) = 2xy$$

$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} + i \cdot \frac{\partial v}{\partial x}$$

When  $f(z)$  is analytic,

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} - i \cdot \frac{\partial u}{\partial y}$$

$$\Rightarrow f(z) = 2y - i(2x)$$


Applying Milne-Theorem method, Replace 'x' with z and 'y' with '0'.

$$\Rightarrow f(z) = -2iz$$

$$\Rightarrow \int f'(z) dz = -2i \int z \cdot dz$$

$$\begin{aligned} \Rightarrow f(z) &= -iz^2 + ic \\ &= -i(x+iy)^2 + ic \\ &= i(y^2 - x^2) + 2xy + ic \\ &= 2xy + i(y^2 - x^2 + \text{constant}) \end{aligned}$$

$$\therefore v(x, y) = y^2 - x^2 + \text{constant}$$



29. (-1.1 to -0.9)

**Given**


$$f(z) = u(x, y) + i v(x, y)$$

and  $u(x, y) = 2kxy$  and  $v(x, y) = x^2 - y^2$ 

$$\text{If } f(z) \text{ is analytic, } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\Rightarrow \frac{\partial}{\partial x}(2kxy) = \frac{\partial}{\partial y}(x^2 - y^2)$$

$$\Rightarrow 2ky = -2y \Rightarrow [k = -1]$$



30. (0 to 0)

$$\text{Given } f(z) = 2z^3 + b|z|^3$$

$$\Rightarrow f(z) = 2(x+iy)^3 + b(x^2+y^2)^{3/2}$$

$$= 2\{x^3 - iy^2 + i3x^2y - 3xy^2\} + b(x^2+y^2)^{3/2}$$

$$= 2x^3 - 6xy^2 + b(x^2+y^2)^{3/2} + i(6x^2y - 2y^3)$$

$\therefore$  For a  $f(z) = u(x, y) + iv(x, y)$  to be analytic

$$\frac{\partial u}{\partial y} = \frac{\partial v}{\partial x}$$


$$\Rightarrow \frac{\partial u}{\partial x} = \frac{\partial}{\partial x} \left\{ 2x^3 - 6xy^2 + b(x^2+y^2)^{3/2} \right\}$$

$$\Rightarrow \frac{\partial v}{\partial x} = \frac{\partial}{\partial y} \left\{ 6xy^2 - 2y^3 \right\}$$

$$\Rightarrow 6x^2 - 6y^2 + \frac{3b}{2} \cdot (2x)(x^2+y^2)^{1/2} = 6x^2 - 6y^2$$

$$\Rightarrow 3bx(x^2+y^2)^{1/2} = 0$$

$$\Rightarrow b = 0$$



31. (b)

$$f(z) = z + z^*$$

$$z = x + iy \quad z^* = x - iy$$

$$\therefore f(z) = z + z^* = x + iy + x - iy$$

$$f(z) = 2x$$


Since  $f(z)$  is polynomial and hence continuous.

$$f(z) = u + iv = 2x$$

$$\therefore u = 2x \text{ and } v = 0$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ & } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Since  $f(z)$  does not satisfy C-R equations therefore  $f(z)$  is not analytic.



**32. (b)**

Given  $f(z) = u(x, y) + iv(x, y)$  is analytic

$$u(x, y) \rightarrow \text{Real part} = e^{-y} \cos x$$

since  $f(z)$  is analytic,

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\Rightarrow e^{-y}(-\sin x) = \frac{\partial v}{\partial y}$$

$$\Rightarrow \int \partial v = \int -e^{-y} \cdot \sin x \cdot \partial y$$

$$\Rightarrow V = -\sin x \cdot \frac{e^{-y}}{-1} + \phi(x)$$

$$\Rightarrow v(x, y) = e^{-y} \cdot \sin x + \phi(x)$$

Where  $\phi(x)$  is arbitrary function

$$\text{For } \phi(x) = 0$$

$$V(x, y) = e^{-y} \cdot \sin x$$



Scan for Video solution

**33. (c)**

$$f(z) = zz^*$$

$$= |z|^2$$

$$(\because zz^* = |z|^2)$$

$$= 1^2$$

$$(\because |z| = 1)$$

$$f(z) = 1 + 0i = (1, 0)$$

$\therefore f(z)$  maps S to the point (1, 0) in the complex plane.



Scan for Video solution

**34. (c)**

**Given**

$$f(z) = u + iv$$
 is analytic and  $u = 2x y$

$$\Rightarrow \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

We have  $f(z) = u + iv$

$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\Rightarrow = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

$$\Rightarrow f'(z) = 2y - i(2x)$$

Applying Milne Thomson method replace 'x' with z and 'y' with '0'

$$\Rightarrow f'(z) = 2(0) - i(2z) = -i2z$$

$$\Rightarrow \int f'(z) dz = -i(2) \cdot \int z dz$$

$$\Rightarrow f(z) = -iz^2 + c = -i(x + iy)^2 + c = 2xy - i(x^2 - y^2) + ic$$

$$\therefore v(x, y) = -x^2 + y^2 + c$$

Scan for Video solution

**35. (c)**

**Given**

$$f(z) = u + iv$$
 is analytic, and  $u = x^2 - y^2$

$$\Rightarrow \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \Rightarrow 2x = \frac{\partial v}{\partial y}$$

$$\Rightarrow \int \partial v = \int 2x \partial y \Rightarrow v = 2xy + h(x)$$

Where  $h(x)$  is arbitrary function

From options,  $h(x) = \text{constant} = c$

$$\therefore v(x, y) = 2xy + c$$

Scan for Video solution

**36. (c)**

**Given:**

$$f(z) = u + iv$$
 is analytic function;  $u(x, y) = xy$

$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} + i \left( \frac{\partial v}{\partial x} \right)$$

$\because f(z)$  is analytic,  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$  and  $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$

$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

$$\Rightarrow f'(z) = \frac{\partial}{\partial x}(xy) - i \frac{\partial}{\partial y}(xy) = y - ix$$

Applying Milne-Thomson method (Replacing 'x' with z & 'y' with 0)

$$\Rightarrow f'(z) = 0 - iz \quad \Rightarrow \quad f'(z) = -iz$$

$$\Rightarrow f'(z) \cdot dz = -iz dz$$

On integrating both sides

$$f'(z) = \frac{-i}{2} z^2 + C = \frac{-i}{2} \{(x+iy)^2\} + C$$

$$\begin{aligned} f'(z) &= \frac{-i}{2} \{x^2 - y^2 + 2ixy\} + C \\ &= xy + i \left( \frac{y^2 - x^2}{2} + C \right) \end{aligned}$$

$$\therefore u = xy, \text{ and } v = \frac{y^2 - x^2}{2} + k$$

where  $k = C = \text{constant}$



Scan for Video solution



### 37. (d)

**Given:**  $\sin z = 10$

$$\Rightarrow \frac{e^{iz} - e^{-iz}}{2i} = 10 \quad \Rightarrow$$

$$e^{iz} - e^{-iz} - 20i = 0$$

$$\Rightarrow (e^{iz})^2 - 20i(e^{iz}) - 1 = 0$$

(Multiplied by  $e^{iz}$ )

$$\therefore e^{iz} = \frac{20i \pm \sqrt{(-20i)^2 - 4(1)(-1)}}{2} = \frac{20i \pm \sqrt{-400 + 4}}{2}$$

$$\Rightarrow e^{iz} = 10i \pm i3\sqrt{11} = i(10 \pm 3\sqrt{11})$$

Since  $e^{iz}$  is periodic function, for infinitely many values of 'z' that satisfy given equation.

$\therefore$  Number of complex solutions is infinite.



Scan for Video solution




### 38. (b)

Given integral is (say)  $I = \oint \left( \frac{z+2}{z^2+2z+2} \right) dz$

$$C: \left| z + 1 - \frac{3}{2}j \right| = 1$$

$$\Rightarrow \left| z - \left( -1 + \frac{3}{2}j \right) \right| = 1$$



$\therefore C$  is a circle with center at  $\left(-1, \frac{3}{2}\right)$  and radius 1.

Singular points of given integral

$$\text{Are at } z^2 + 2z + 2 = 0$$

$$\Rightarrow (z+1)^2 + 1 = 0$$

$$\Rightarrow z = -1 \pm j$$

i.e. at  $z = -1 + j$  and  $z = -1 - j$

out of the two singular points, only  $z = -1 + j$  lies inside 'c'

$\therefore$  By Cauchy's Residue theorem

$$\oint \left( \frac{z+2}{z^2+2z+2} \right) dz =$$

$$2\pi j \cdot \text{Res} \left( \frac{(z+2)}{(z - (-1+j))(z - (-1-j))} \Big|_{z=-1+j} \right)$$

$$= 2\pi j \left\{ \frac{-1+j+2}{(-1+j) - (-1-j)} \right\}$$

$$= 2\pi j \left\{ \frac{j+1}{2j} \right\} = \pi(1+j)$$

$$\therefore \oint \left( \frac{z+2}{z^2 + 2z + 2} \right) dz = \pi(1+j)$$




Scan for Video solution

**39. (0.50 to 0.50)**

Given, integral is

$$\oint_c \frac{2^z}{z^2 - 1} dz = -i\pi A$$

and the path 'C' is as shown.



Singularities of the given integral are at

$$z^2 - 1 = 0 \Rightarrow z = 1, -1$$

In the given curve, only  $z = -1$  lies with in 'C'.

$$\begin{aligned} \oint_c \frac{2^z}{(z-1)(z+1)} dz &= 2\pi i \left\{ \text{Res} \left( \frac{2^z}{(z-1)(z+1)} \right) \Big|_{z=-1} \right\} \\ &= 2\pi i \left\{ \frac{2^{-1}}{-2} \right\} = -\frac{\pi i}{2} \end{aligned}$$

$$\therefore A = \frac{1}{2} = 0.50$$



Scan for Video solution

**40. (a)**

$$\text{Given: } I = \oint_C \left( \frac{6z}{2z^4 - 3z^3 + 7z^2 - 3z + 5} \right) dz$$

where 'c' is enclosing the pole  $z = i$ 

From Cauchy's residue theorem

$$I = 2\pi i \times R \text{ es} \left( \frac{6z}{2z^4 - 3z^3 + 7z^2 - 3z + 5} \right)_{z=i}$$

$$\begin{aligned} I &= \left\{ \lim_{z \rightarrow i} \frac{6z}{2z^4 - 3z^3 + 7z^2 - 3z + 5} (z-i) \right\} \\ &= 2\pi i \times \left( \frac{0}{2+3i-7-3i+5} \right) = 2\pi i \times \left( \frac{0}{0} \right) \end{aligned}$$

Apply L-Hospital rule for the limit

$$\begin{aligned} \Rightarrow I &= 2\pi i \left\{ \lim_{z \rightarrow i} \frac{12z - 6i}{8z^3 - 9z^2 + 14z - 3} \right\} \\ &= 2\pi i \left\{ \frac{6i}{-8i + 9 + 14i - 3} \right\} = 2\pi i \left\{ \frac{6i}{6(1+i)} \right\} \end{aligned}$$

$$\therefore I = \frac{-2\pi}{(1+i)} \times \frac{(1-i)}{(1-i)} = \frac{-2\pi}{2}(1-i)$$


$$\Rightarrow I = -\pi(1-i) = \pi(-1+i)$$

$$\therefore \oint_c \frac{6z}{2z^4 - 3z^3 + 7z^2 - 3z + 5} dz = \pi(-1+i)$$

**41. (0.2 to 0.2)**

Given

$$\text{Integral is } I = \frac{1}{2\pi} \int_C \frac{1}{(z-i)(z+4i)} dz$$

Poles are  $z = i$ ,  $z = -4i$ The pole which is inside 'C' is only  $z = i$ 

$$I = \frac{1}{2\pi} \times 2\pi i \left\{ \text{Res} \left( \frac{1}{(z-i)(z+4i)} \right) \Big|_{z=i} \right\}$$

$$= i \left\{ \lim_{z \rightarrow i} \frac{1}{(z-i)(z+4i)} \times (z-i) \right\} = i \left( \frac{1}{5i} \right) = \frac{1}{5} = 0.2$$

$$\therefore \frac{1}{2\pi} \int_C \frac{1}{(z-i)(z+4i)} dz = 0.2$$



Scan for Video solution




42. (c)

**Given**

$$\oint_C \frac{\cosh 3z}{2z} dz = I \text{ (say)}$$

$z = 0$  is a singular point of  $\frac{\cosh 3z}{2z}$  within 'C'.



From Cauchy's residue theorem,

$$\oint_C \frac{\cosh 3z}{2z} dz = 2\pi i \{ \text{Res } f(z) \mid_{z=0} \}$$

$$= 2\pi i \left\{ \lim_{z \rightarrow 0} \frac{\cosh 3z}{2z} \times z \right\} = 2\pi i \left\{ \frac{e^{3z} + e^{-3z}}{2 \times 2} \Big|_{z=0} \right\}$$

$$= 2\pi i \left( \frac{1}{2} \right) = \pi i$$

$$\left\{ \because \cosh z = \left( \frac{e^z + e^{-z}}{2} \right) \right\}$$

$$\therefore \oint_C \frac{\cosh 3z}{2z} dz = \pi i$$




Scan for Video solution



43. (c)

$$I = \int_C \frac{dz}{z^2(z-4)}$$



Only pole at  $z = 0$  is there as it lies within the contour C.

Pole of order 2 at  $z = 0$  exists.

$$\text{Res}_{z \rightarrow a} [f(a)] = \frac{1}{(n-1)!} \left[ \frac{d^{n-1}}{dz^{n-1}} f(z) \times (z-a)^n \right]$$


$$= \frac{1}{(2-1)!} \cdot \left[ \frac{d}{dz} \cdot \frac{1}{z^2(z-4)} \cdot z^2 \right]_{z=0} = -\frac{1}{(z-4)^2} \Big|_{z=0}$$

$$\text{Res}[f(0)] = -\frac{1}{16}$$

By Cauchy Residue Theorem

$$\int f(z) dz = 2\pi j \text{Res}[f(a)]$$

$$\int_C \frac{dz}{z^2(z-4)} = 2\pi j \left( -\frac{1}{16} \right) = \frac{-\pi j}{8}$$



44. (c)

Using Cauchy's Integral theorem,

$$\int_C \frac{f(z)}{z-a} dz = 2\pi i (\text{Res}_{z=a})$$

$$I = \int_C \frac{z^2+1}{z^2-2z} dz = \int_C \frac{z^2+1}{z(z-2)} dz = \int_C \frac{\left( \frac{z^2+1}{z-2} \right)}{z} dz = \int_C \frac{f(z)}{z} dz$$

Poles are at  $z = 0$  and  $z = 2$  but only  $z = 0$  is lying inside the unit circle centered at origin.

$$\text{Residue } (z=0) = \lim_{z \rightarrow 0} f(z) = \frac{z^2+1}{z-2} = -\frac{1}{2}$$

$$\therefore \int_C \frac{z^2+1}{z^2-2z} dz = 2\pi i (\text{Res}_{z=0}) = 2\pi i \left( -\frac{1}{2} \right) = -\pi i$$



Scan for Video solution

**45. (b)**

Using Cauchy's Integral formula:-

$$f(a) = \frac{1}{2\pi j} \oint_C \frac{f(z)}{z-a} dz$$

(if  $f(z)$  is analytic within and on the closed curve  $C$ )

$$f(z) = z^3 + z^2 + 8$$

$$f(-2) = (-2)^3 + (-2)^2 + 8$$

$$f(-2) = -8 + 4 + 8$$

$$\therefore f(-2) = 4$$

$$I = \oint_C \frac{z^3 + z^2 + 8}{z+2} dz \quad \therefore a = -2$$

$$= 2\pi j f(a)$$

$$= 2\pi j f(-2)$$

$$= 2\pi j (-4)$$

$$= -8\pi j$$

$$\therefore \oint_{|z|=5} \frac{z^3 + z^2 + 8}{z+2} dz = -8\pi j$$



Scan for Video solution

**46. (-0.0001 to 0.0001)**

**Given:** Integral  $\frac{1}{2\pi j} \oint_C \left( z + \frac{1}{z} \right)^2 dz = I$  (say),  $c: |z| = 1$

$$\Rightarrow I = \frac{1}{2\pi j} \oint_C \left( z + \frac{1}{z} \right)^2 dz = \frac{1}{2\pi j} \oint_C \left( z^2 + 2 + \frac{1}{z^2} \right) dz$$

$$= \frac{1}{2\pi j} \oint_C \left( z^2 + 2 \right) dz + \frac{1}{2\pi j} \oint_C \left( \frac{1}{z^2} \right) dz$$

Since  $z^2 + 2$  is analytic over complex plane,

$$\oint_C \left( z^2 + 2 \right) dz = 0$$

$$\therefore I = \frac{1}{2\pi j} \oint_C \left( \frac{1}{z^2} \right) dz$$


 $z = 0$  is a pole of order 2 and  $z = 0$  lies in  $|z| = 1$ 

$$\begin{aligned} \therefore I &= \frac{1}{2\pi j} \oint_C \frac{1}{z^2} dz = \frac{1}{2\pi j} \times 2\pi j \times \left\{ \text{Res} \left( \frac{1}{z^2} \right) \Big|_{z=0} \right\} \\ &= \lim_{z \rightarrow 0} \frac{d}{dz} \left( z^2 \cdot \frac{1}{z^2} \right) = 0 \end{aligned}$$

$$\therefore \boxed{I = \frac{1}{2\pi j} \oint_C \left( z + \frac{1}{z^2} \right) dz = 0}$$



Scan for Video solution

**47. (2 to 2)****Given:**

$$\text{Integral } I = \frac{1}{\pi j} \oint_C \frac{dz}{z^2 - 1}$$

$$\Rightarrow I = \frac{1}{\pi j} \left\{ \oint_{C_1} \frac{dz}{z^2 - 1} - \oint_{C_2} \frac{dz}{z^2 - 1} \right\}$$

 $C_2$  is negative since it is in clockwise direction.

$$\Rightarrow \frac{1}{\pi j} \left\{ \text{Res} \left. \frac{1}{z^2 - 1} \right|_{z=1} \times 2\pi j - \text{Res} \left. \frac{1}{z^2 - 1} \right|_{z=-1} \times 2\pi j \right\}$$

$$\Rightarrow \left\{ \lim_{z \rightarrow 1} (z-1) \frac{1}{z^2 - 1} \times 2 - \lim_{z \rightarrow -1} (z+1) \frac{1}{(z+1)(z-1)} \times 2 \right\}$$

$$\left\{ \lim_{z \rightarrow 1} (z-1) \frac{1}{(z+1)(z-1)} \times 2 - \lim_{z \rightarrow -1} (z+1) \frac{1}{(z+1)(z-1)} \times 2 \right\}$$


$$\Rightarrow = 1 - (-1) = 2$$

$$\therefore \frac{1}{\pi j} \oint_C \frac{dz}{z^2 - 1} = 2$$



Scan for Video solution



**48. (a)**Equation of circle C:  $|z + 2| = 1$ 

$$I = \oint_c \frac{z+1}{z^2 - 4} dz$$

Only pole  $z = -2$  is lying within the contour therefore

$$\text{Res}(z = -2) = \lim_{z \rightarrow -2} [f(z)(z + 2)] = \lim_{z \rightarrow -2} \frac{z+1}{z-2} = \frac{1}{4}$$

$\therefore$  By Cauchy's Residue theorem,

$$\oint f(z) dz = 2\pi i [\text{Res } f(a)]$$

$$I = \oint_c \frac{z+1}{z^2 + 4} dz = 2\pi i \left( \frac{1}{4} \right) = \frac{\pi i}{2}$$



Scan for Video solution

**49. (b)**

$$f(z) = \frac{z^2}{(z^2 - 3z + 2)^2}$$

$$= \frac{z^2}{(z-1)^2(z-2)^2}$$

$\therefore$  At  $z = 1$  and  $z = 2$  poles of order 2 exists and they are lying within the contour  $|z| = 4$

$$\text{Res}[f(a)] = \frac{1}{(n-1)!} \lim_{z \rightarrow a} \left[ \lim_{n \rightarrow \infty} \frac{d^{n-1}}{dz^{n-1}} f(z)(z-a)^n \right]$$

$$\therefore \text{Res}[f(1)] = \frac{1}{(2-1)!} \lim_{z \rightarrow 1} \left[ \frac{d}{dz} \frac{z^2}{(z-2)^2} \right]$$

$$= \frac{(z-2)^2 \cdot 2z - z^2 \cdot 2(z-2)}{(z-2)^4} = 2 + 2 = 4$$

$$\therefore \text{Res}[f(2)] = \frac{1}{(2-1)!} \lim_{z \rightarrow 2} \left[ \frac{d}{dz} \frac{z^2}{(z-1)^2} \right]$$


$$= \frac{(z-1)^2 2z - z^2 2(z-1)}{(z-1)^4} = 4 - 8 = -4$$

By Cauchy Residue Theorem,

$$\oint f(z) dz = 2\pi i [\text{Res } f(a_1) + \text{Res } f(a_2)]$$

$$I = \oint_c \frac{z^2}{(z^2 - 3z + 2)^2} dz = 2\pi i [\text{Res } f(1) + \text{Res } f(2)]$$


$$= 2\pi i [4 - 4] = 0$$

**50. (a)**

Given,

$$\oint_c \frac{1}{5z-4} dz = A \pi i ; 'c' \text{ is unit circle centered at origin.}$$


For  $f(z) = \frac{1}{\left(z - \frac{4}{5}\right)}$ ,  $z = \frac{4}{5}$  is a singular point and it lies within 'C'



$\therefore$  From Cauchy's Residue theorem,

$$\begin{aligned} \frac{1}{5} \cdot \oint_c \frac{dz}{z - \frac{4}{5}} &= \frac{1}{5} \times 2\pi i \left( \text{Res } f(z) \Big|_{z=\frac{4}{5}} \right) \\ &= \frac{1}{5} \times 2\pi i \times (1) = \frac{2}{5} \pi i \end{aligned}$$

$$\Rightarrow A = \frac{2}{5}$$



51. (b)

$$\text{Given, } f(z) = \frac{1}{(z-4)(z+1)^3}$$

$\Rightarrow z=4$  is a simple pole.

$z=-1$  is a pole of order 3.

(i) Residue at  $z=4$

$$\begin{aligned} \text{Res } f(z)|_{z=4} &= (z-4) \cdot f(z)|_{z=4} = \frac{1}{(z+1)^3} \Big|_{z=4} \\ &= \frac{1}{5^3} = \frac{1}{125} \end{aligned}$$

(ii) Residue at  $z=-1$

$$\begin{aligned} \text{Res } f(z)|_{z=-1} &= \frac{1}{(3-1)!} \\ &\lim_{z \rightarrow -1} \frac{d^2}{dz^2} \left\{ (z+1)^3 \cdot \frac{1}{(z-4)(z+1)^3} \right\} \\ &= \frac{1}{2!} \cdot \lim_{z \rightarrow -1} \frac{2}{(z-4)^3} = \frac{1}{2} \times \left( \frac{2}{-125} \right) = \frac{-1}{125} \end{aligned}$$

$\therefore$  Residues of  $f(z)$  are  $\frac{1}{125}$  and  $\frac{-1}{125}$



Scan for Video solution



52. (d)

Given:

$$\oint_C \frac{z^2 - 1}{z^2 + 1} e^z dz = I ; \quad C: |z| = 3$$

Singular points of given function are

At  $z^2 + 1 = 0 \Rightarrow z^2 = -1 \Rightarrow z = \pm i$


Both the singular points  $z = i$ ;  $z = -i$  lies in 'C'

$$\oint_C \frac{z^2 - 1}{z^2 + 1} e^z dz = 2\pi i \left\{ \text{res } f(z)|_{z=i} + \text{res } f(z)|_{z=-i} \right\}$$

$$\begin{aligned} I &= 2\pi i \left\{ \lim_{z \rightarrow i} \frac{z^2 - 1}{(z+i)^2} e^z + \lim_{z \rightarrow -i} \frac{z^2 - 1}{(z-i)^2} e^z \right\} \\ &= 2\pi i \left\{ \frac{-1-1}{2i} e^i + \frac{-1-1}{-2i} e^{-i} \right\} = \pi \{-2e^i + 2e^{-i}\} \end{aligned}$$

$$= -2\pi \{e^i - e^{-i}\} = -2\pi \{2i \sin 1\} = -4\pi i \sin 1$$

$$\therefore I = -4\pi i \sin 1$$



Scan for Video solution



53. (b)

The line integral is being evaluated while moving from point  $(0, 0)$  to  $(1, i)$  therefore the equation of straight line will be  $y = x$  and  $z = x + iy$ .

$$\therefore dy = dx \quad \therefore dz = dx + idy$$

$$I = \int_c (x^2 + iy^2) dz$$

$$= \int_{x=0}^{x=1} (x^2 + ix^2)(dx + idx)$$

$$= \int_0^1 x^2 dx (1+i)^2 = \int_0^1 x^2 (1-1+2i) dx$$

$$= 2i \left[ \frac{x^3}{3} \right]_0^1 = \frac{2}{3} i$$



Scan for Video solution



54. (c)

$f(z) = \frac{z^3 - 2z + 3}{z-2}$  is having pole at  $z = 2$  which is

lying within the contour  $|z| = 3$ , therefore we have to find residue at  $z = 2$ .

$$\begin{aligned} \text{Res}(z=2) &= \lim_{z \rightarrow 2} f(z)(z-2) = \lim_{z \rightarrow 2} \frac{z^3 - 2z + 3}{z-2} (z-2) \\ &= 2^3 - 2(2) + 3 = 7 \end{aligned}$$

So, by Cauchy's Residue Theorem,

$$\oint f(z) dz = 2\pi i (\text{Res } f(a))$$

$$= 2\pi i [\text{Res } f(2)] = 2\pi i (7) = 14\pi i$$



Scan for Video solution



**55. (b)**

$$I = \oint_c \frac{2z+5}{\left(z - \frac{1}{2}\right)(z^2 - 4z + 5)} dz$$

Poles of the given function are  $z = \frac{1}{2}$ ,  $z = 2 + i$ ,  $z = 2 - i$ .

Only pole  $z = \frac{1}{2}$  is lying within the contour  $|z| = 1$

$\therefore$  By Cauchy's Residue theorem,

$$\begin{aligned} \oint_c \frac{f(z)}{z-a} dz &= 2\pi i \cdot [\text{Res } f(a)] \\ &= 2\pi i \left[ \frac{2 \times \frac{1}{2} + 5}{\left(\frac{1}{2}\right)^2 - 4\left(\frac{1}{2}\right) + 5} \right] \quad \left\{ \because a = \frac{1}{2} \right\} \\ &= 2\pi i \left( \frac{24}{13} \right) = \frac{48}{13}\pi i \end{aligned}$$



Scan for Video solution

**56. (b)**

Given integral is


$$I = \frac{1}{2\pi j} \oint_c \frac{e^z}{z-2} dz$$

(i) When  $z = 2$  is inside the contour

$$\begin{aligned} I &= \frac{1}{2\pi j} \oint_c \frac{e^z}{z-2} dz \\ &= I = \frac{1}{2\pi j} \times 2\pi j \times \text{Res} \left( \frac{e^z}{z-2} \right)_{z=2} \\ &= e^2 = 7.389 \end{aligned}$$

(ii) When  $z = 2$  is outside the contour

Then  $I = \frac{1}{2\pi j} \oint_c \frac{e^z}{z-2} dz = 0$  (From Cauchy's integral theorem).

**57. (-136 to -132)**

Given that contour 'C' encloses  $2\pi j$  and  $-2\pi j$


$$\text{Let } I = \frac{-1}{2\pi} \cdot \oint_C \frac{\sin z}{(z - 2\pi j)^3} dz$$

For the given integrand,  $z = 2\pi j$  is a pole of order '3'

$$\begin{aligned} \therefore \oint_C \frac{\sin z dz}{(z - 2\pi j)^3} &= 2\pi j \cdot \text{Res} \left\{ \frac{\sin z}{(z - 2\pi j)^3} \right\}_{z=2\pi j} \\ &= 2\pi j \left\{ \frac{1}{(3-1)!} \lim_{z \rightarrow 2\pi j} \frac{d^2}{dz^2} (\sin z) \right\} \\ &= \frac{2\pi j}{2!} \cdot -\sin(2\pi j) \\ &= -\pi j (\sin 2\pi j) = -\pi j \left( \frac{e^{j(2\pi j)} - e^{-j(2\pi j)}}{2j} \right) \\ &= \frac{-\pi}{2} (e^{-2\pi} - e^{2\pi}) \end{aligned}$$

$$\therefore I = \frac{-1}{2\pi} \cdot \oint_C \frac{\sin z}{(z - 2\pi j)^3} dz = \frac{-1}{2\pi} \cdot \left( -\frac{\pi}{2} (e^{-2\pi} - e^{2\pi}) \right) = \frac{+e^{-2\pi} - e^{2\pi}}{4} = -133.87$$

$$\therefore \frac{-1}{2\pi} \cdot \oint_C \frac{\sin z}{(z - 2\pi j)^3} dz = -133.87$$

**58. (b)**

Given

$$\text{Integral } \oint_\Gamma \frac{3z-5}{(z-1)(z-2)} dz = \oint_\Gamma f(z) dz$$

$$\text{Consider } \text{Res } f(z)|_{z=1} = \frac{3(1)-5}{(1-2)} = 2$$


$$\text{Res } f(z) \Big|_{z=2} = \frac{3(2)-5}{(2-1)} = 1$$

For  $\oint_{\Gamma} \frac{3z-5}{(z-1)(z-5)} dz = 4\pi i = 2\pi i(2)$ ,

$\therefore \text{Res } f(z) \Big|_{z=1} = 2$  is only to be considered.

$\Rightarrow z = 1$  has to be inside  $\Gamma$  and  $z = 2$  has to be outside  $\Gamma$ .

$\therefore$  option 'B' is correct.



59. (a)

Given integral is  $\int_{-\infty}^{\infty} \frac{\sin x}{x^2 + 2x + 2} dx$

As per the Contour Integration,

$-\infty < x < \infty \Rightarrow$  Implies the upper half of the circle with center at origin and radius  $\rightarrow \infty$ .


$$\therefore I = \int_{-\infty}^{\infty} \frac{\sin x}{x^2 + 2x + 2} dx = \oint_c \frac{\sin z}{z^2 + 2z + 2} dz$$

For the given functions, the poles are at

$$z^2 + 2z + 2 = 0$$

$$\Rightarrow (z+1)^2 + 1 = 0$$

$$\Rightarrow z = -1 \pm i$$




$\therefore$  Poles are at  $z = -1 + i$  and  $z = -1 - i$ .

Out of the above poles  $z = -1 + i$  lies in 'C'.

$$\begin{aligned} \therefore \oint_C \frac{\sin z}{z^2 + 2z + 2} dz &= 2\pi i \cdot \text{Res}\left(\frac{\sin z}{z^2 + 2z + 2}, z = -1 + i\right) \\ &= 2\pi i \cdot \frac{\sin(-1+i)}{2i} \\ &= \pi \cdot \sin(-1+i) = \pi \cdot \text{Im}(e^{i(-1+i)}) \end{aligned}$$

$$= \pi \cdot \text{Im}(e^{-i} \cdot e^{-1})$$


$$\Rightarrow \frac{\pi}{e} \cdot \text{Im}(\cos 1 - i \sin 1) = \frac{-\pi}{e} \cdot \sin 1$$



60. (d)

(i) Residue of  $\frac{z}{z^2 - 1}$  at  $z = 1$


$$\begin{aligned} &= \frac{1}{(1-1)!} \lim_{z \rightarrow 1} \frac{d^0}{dz^0} \left\{ (z-1) \frac{z}{(z+1)(z-1)} \right\} \\ &= \frac{1}{0!} \times \frac{1}{1+1} = \frac{1}{2} \end{aligned}$$



(ii)  $\oint_C z^2 dz = 0$ , since  $z^2$  is analytic at all points in complex plane.

$$(iii) \frac{1}{2\pi i} \oint_C \frac{1}{z} dz = \frac{1}{2\pi i} \times 2\pi i(1) = 1$$

(iv)  $\bar{z}$  is not an analytic function since C-R equations are not satisfied.



61. (b)

If the path 'C' is given by  
 $|z - z_0| = r$

Then  $\oint_C (z-a)^n dz = \begin{cases} 0; & \text{if } n \neq -1 \\ 2\pi i; & \text{if } n = -1 \end{cases} \rightarrow$  Cauchy's

integral theorem

For  $n = -(n+1) \neq -1$

$$\Rightarrow \oint_C \frac{dz}{(z - z_0)^{n+1}} = 0$$



Scan for Video solution



### 62. (0.5 to 0.5)


Given 'C' denotes a counter clockwise unit circle

Integral is

$$I = \frac{1}{2\pi j} \oint_C \operatorname{Re}(z) dz$$

$$= \frac{1}{2\pi j} \oint_C x \cdot (dx + jdy)$$

$$= \frac{1}{2\pi j} \left\{ \oint_C x dx + (jx) dy \right\}$$



Applying Green's theorem

$$\oint_C M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dy \cdot dx$$

$$\therefore \oint_C x dx + (jx) dy = \iint_R \left( \frac{\partial}{\partial x} (jx) - \frac{\partial}{\partial y} (x) \right) dy dx$$

$$= j \iint_R dx dy$$

$$\therefore \text{Integral } I = \frac{1}{2\pi j} \cdot j \cdot \iint_R dx dy$$

$$= \frac{1}{2\pi} \text{Area of 'R'} = \frac{1}{2\pi} \times \pi(1)^2 = \frac{1}{2} = 0.5$$

$$\therefore \oint_C \frac{1}{2\pi j} \cdot \operatorname{Re}(z) dz = 0.5$$



Scan for Video solution



### 63. (c)

Given


$$C: |z| = 3$$

$\Rightarrow$  C is a circle with center at origin and radius =

$$\text{Given Integral is } \oint_C \frac{z^2 - z + 4j}{z + 2j} dz$$

Singular point is  $z + 2j = 0 \Rightarrow z = -2j$

$\operatorname{Im}(z)$



$$\Rightarrow \oint_C \frac{z^2 - z + 4j}{z + 2j} dz = 2\pi j \cdot \operatorname{Res} \left\{ \frac{z^2 - z + 4j}{z + 2j} \Big|_{z=-2j} \right\}$$

$$= 2\pi j \{(-2j)^2 - (-2j) + 4j\}$$

$$= 2\pi j \{-4 + 6j\}$$

$$= 4\pi j \{-2 + 3j\} = 4\pi \{-3 - 2j\}$$

$$\therefore \oint_C \frac{z^2 - z + 4j}{z + 2j} dz = -4\pi(3 + 2j)$$



Scan for Video solution



### 64. (c)

$$f(z) = \frac{z^2}{z^2 - 1}$$


$$C : |z - 1| = 1$$

Poles of  $f(z)$  are at  $z^2 - 1 = 0$

$$\Rightarrow (z - 1)(z + 1) = 0$$

$$\Rightarrow z = \pm 1$$

Only pole  $z = 1$  lies within the contour C.



$$\text{Res}[f(a)] = \lim_{z \rightarrow a} f(z)(z-a)$$

$$= \lim_{z \rightarrow 1} \frac{z^2}{z^2 - 1} \cdot (z-1) = \lim_{z \rightarrow 1} \left( \frac{z^2}{z+1} \right) = \frac{1^2}{1+1} = \frac{1}{2}$$

By Cauchy Residue Theorem,

$$\int_c \frac{z^2}{z^2 - 1} dz = 2\pi i [\text{Res } f(1)] = 2\pi i \left( \frac{1}{2} \right) = \pi i$$



Scan for Video solution



65. (a)


$$I = \oint_c \frac{z^2 - 4}{z^2 + 4} dz \quad C : |z - i| = z$$

Poles of the given function are at  $z^2 + 4 = 0$

$$\Rightarrow (z + 2i)(z - 2i) = 0$$

$$\Rightarrow z = +2i, -2i$$

$\therefore$  Only pole  $z = 2i$  is lying within the contour C.



$$\text{Res}[f(a)] = \lim_{(z \rightarrow a)} f(z)(z-a)$$

$$\text{Res}[f(2i)] = \lim_{z \rightarrow 2i} \frac{z^2 - 4}{z^2 + 4} \cdot (z - 2i)$$

$$= \frac{4i^2 - 4}{2i + 2i} = \frac{-8}{4i} = -\frac{2}{i}$$

$\therefore$  By Cauchy Residue theorem,

$$I = \oint_c \frac{z^2 - 4}{z^2 + 4} dz = 2\pi i [\text{Res } f(2i)] = 2\pi i \left( -\frac{2}{i} \right) = -4\pi$$




Scan for Video solution



66. (c)

$$f(z) = \frac{1}{z+1} - \frac{2}{z+3} \quad C: |z+1|$$

Only  $z = -1$  is lying within the contour C.



$$\text{Res}[f(a)] = \lim_{(z \rightarrow a)} f(z)(z-a)$$

$$= \lim_{z \rightarrow -1} \left[ \frac{1}{z+1} - \frac{2}{z+3} \right] (z+1)$$

$$\text{Res}[f(-1)] = \left[ 1 - \frac{2(z+1)}{(z+3)} \right]_{z=-1} = 1$$

By Cauchy Residue Theorem,

$$= \frac{1}{2\pi j} \oint_c \left[ \frac{1}{z+1} - \frac{1}{z+3} \right] dz = \frac{1}{2\pi j} 2\pi j \text{Res}[f(-1)] = 1$$

$$\therefore \frac{1}{2\pi j} \oint_c f(z) dz = 1$$



Scan for Video solution




67. (c)

$$\text{Given: } f(z) = \frac{1}{z+1} - \frac{2}{z+3} \text{ and } c: |z+1| = 1$$

$\Rightarrow$  c is circle with centre  $(-1, 0)$  and radius '1'

The poles of  $f(z)$  are at  $z = -1, z = -3$

$\because z = -1$  is the only pole within 'c'



$$\frac{1}{2\pi j} \oint_c f(z) dz = \frac{1}{2\pi j} \oint_c \frac{-z+1}{(z+1)(z+3)} dz$$

$$= \frac{1}{2\pi j} \times 2\pi j \left\{ \text{Res} \left( \frac{1-z}{(z+1)(z+3)} \right)_{z=-1} \right\} = \frac{2}{2} = 1$$

$$\therefore \frac{1}{2\pi j} \oint_c \left( \frac{1}{z+1} - \frac{1}{z+3} \right) dz = 1$$




Scan for Video solution



68. (a)

**Given:** Integral  $\oint_c \frac{-3z + 4}{z^2 + 4z + 5} dz$ ; c:  $|z| = 1$ -unit circle



For  $\frac{-3z + 4}{z^2 + 4z + 5}$ , poles are at the points

$$\text{Where } z^2 + 4z + 5 = 0$$

$$\Rightarrow (z+2)^2 + 1 = 0 \Rightarrow z = -2 \pm i$$

Since both the poles are outside the path, given function is analytic within 'c'.

$$\therefore \oint_{|z|=1} \frac{-3z + 4}{z^2 + 4z + 5} dz = 0$$

Scan for Video solution



69. (c)

$$\text{Given function is } X(z) = \frac{1-2z}{z(z-1)(z-2)}$$

Poles of X(z) are at the points where,

$$z(z-1)(z-2) = 0$$

$\Rightarrow z = 0, z = 1, z = 2$  are simple poles.

$\therefore$  Residue of

$$(z)|_{z=0} = \lim_{z \rightarrow \infty} z \cdot X(z) = \lim_{z \rightarrow 0} \frac{1-2z}{(z-1)(z-2)} = \frac{1}{2}$$


Residue of

$$X(z)|_{z=1} = \lim_{z \rightarrow 1} (z-1) \cdot X(z) = \lim_{z \rightarrow 1} \frac{1-2z}{z(z-2)} = \frac{-1}{-1} = 1$$

Residue of

$$X(z)|_{z=2} = \lim_{z \rightarrow 2} (z-2) \cdot X(z) = \lim_{z \rightarrow 2} \frac{1-2z}{z(z-1)} = \frac{-3}{2}$$


$\therefore$  Residue at poles are  $\frac{1}{2}, 1$  and  $\frac{-3}{2}$ .



70. (d)

**Given:**

$$f(z) = c_0 + \frac{c_1}{z}$$




$$\oint_{\text{unit circle}} \frac{1+f(z)}{z} dz = \oint_{|z|=1} \frac{1+f(z)}{z} dz$$

$$= \oint_c \left[ \left( \frac{1+c_0}{z} \right) + \left( \frac{c_1}{z^2} \right) \right] dz$$

$$= 2\pi j \left\{ \text{Res} \left( \frac{1+c_0}{z} \right) \Big|_{z=0} + \text{Res} \left( \frac{c_1}{z^2} \right) \Big|_{z=0} \right\}$$

$$= 2\pi j \left\{ (1+c_0) + \frac{1}{1!} \lim_{z \rightarrow 0} \frac{d}{dz} \left( z^2 \cdot \frac{c_1}{z^2} \right) \right\} = 2\pi j(1+c_0)$$

$$\oint_{\text{unit circle}} \frac{1+f(z)}{z} dz = 2\pi j(1+c_0)$$



71. (a)

**Given:**  $f(z) = \frac{1}{(z+2)^2(z-2)^2}$  and  $z = 2$  is a pole

of order '2'.

For any  $f(z)$ , where  $z = a$  is a pole of order 'n'

$$\text{Res } f(z)|_{z=a} = \frac{1}{(n-1)!} \left\{ \lim_{z \rightarrow a} \frac{d^{n-1}}{dz^{n-1}} (z-a)^n f(z) \right\}$$

$$\text{Res } f(z)|_{z=2} = \frac{1}{(2-1)!} \lim_{z \rightarrow 2} \frac{d}{dz} \left\{ (z-2)^2 \frac{1}{(z-2)^2 (z+2)^2} \right\}$$

$$= \frac{1}{1!} \left\{ \lim_{z \rightarrow 2} -2(z+2)^{-3} \right\} = \frac{-2}{64} = \frac{-1}{32}$$

$\therefore \text{Res } f(z)|_{z=2} = \frac{-1}{32}$



Scan for Video solution



72. (d)

$$X(z) = \frac{z}{(z-a)^2}; |z| > a$$

$$f(z) = X(z) \cdot z^{n-1}$$

$$f(z) = \frac{z}{(z-a)^2} \cdot z^{n-1} = \frac{z^n}{(z-a)^2}$$

$\therefore$  We have a pole at  $z = a$  of order 2.

$$\text{Res}[f(a)] = \frac{1}{(n-1)!} \lim_{z \rightarrow a} \left[ \frac{d^{n-1}}{dz^{n-1}} f(z) \cdot (z-a)^n \right]$$


$$\begin{aligned} \text{Res}[f(a)] &= \frac{1}{(2-1)!} \lim_{z \rightarrow a} \left[ \frac{d}{dz} \frac{z^n}{(z-a)^2} \cdot (z-a)^n \right] \\ &= \lim_{z \rightarrow a} nz^{n-1} = na^{n-1} \end{aligned}$$



Scan for Video solution



73. (a)

**Given:**

$$f(z) = \frac{\cos z}{z}$$

 $\Rightarrow z = 0$  is a singular point

From Cauchy Residue Theorem

$$\oint f(z) dz = 2\pi i \times \{\text{sum of residue of } f(z)\}$$

$$\Rightarrow \oint_{|z|=1} \frac{\cos z}{z} dz = 2\pi i \{\text{Res of } f(z)|_{z=0}\}$$

$$= 2\pi i \{\cos z|_{z=0}\} = 2\pi i (1) = 2\pi i$$

$$\therefore \oint_{|z|=1} \frac{\cos z}{z} dz = 2\pi i$$

Scan for Video solution



74. (1 to 1)

Given

$$F(z) = \frac{\sin z}{z^2}$$

$$\frac{1}{z^2} \left\{ z - \frac{z^3}{3!} + \frac{z^5}{5!} \dots \right\}$$

$$\frac{1}{z} - \frac{z}{3!} + \frac{z^3}{5!} + \dots$$

$\text{Res}(z)|_{z=0} = \text{coefficient of } \frac{1}{z}$  in series expansion of  $f(z)$

$$\Rightarrow \text{res}(z)|_{z=0} = 1$$



Scan for Video solution



Scan for Video solution



## CHAPTER

# 7

# FOURIER SERIES

### General Fourier Expansions

- 1. [MSQ]** [GATE-CE-2023-2M]

The following function is defined over the interval  $[-L, L]$ :

$$f(x) = px^4 + qx^5$$

If it is expressed as a Fourier series,

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left\{ a_n \sin\left(\frac{\pi x}{L}\right) + b_n \cos\left(\frac{\pi x}{L}\right) \right\},$$

Which options amongst the following are true?

- (a)  $a_n, n = 1, 2, \dots, \infty$  depend on  $p$
- (b)  $a_n, n = 1, 2, \dots, \infty$  depend on  $q$
- (c)  $b_n, n = 1, 2, \dots, \infty$  depend on  $p$
- (d)  $b_n, n = 1, 2, \dots, \infty$  depend on  $q$

- 2. [NAT]** [GATE-CE-2022-1M]

The Fourier cosine series of a function is given by:

$$f(x) = \sum_{n=0}^{\infty} f_n \cos nx$$

For  $f(x) = \cos^4 x$ , the numerical value of  $(f_4 + f_5)$  is \_\_\_\_\_. (round off to three decimal places).

- 3. [MCQ]** [GATE-ME-2022:1M]


The Fourier series expansion of  $x^3$  in the interval  $-1 \leq x < 1$  with periodic continuation has

- (a) Only sine terms
- (b) Only cosine terms
- (c) Both sine and cosine terms
- (d) Only sine terms and a non-zero constant

- 4. [MCQ]**

[GATE-ME-2022:1M]

$F(t)$  is a periodic square wave function as shown. It takes only two values, 4 and 0, and stay at each of these values for 1 second before changing. What is the constant term in the Fourier series expansion of  $F(t)$ ?



- (a) 1
- (b) 2
- (c) 3
- (d) 4

- 5. [MCQ]**

[GATE-EE-2021:2M]

Let  $f(t)$  be an even function, i.e.  $f(-t) = f(t)$  for all  $t$ . Let the Fourier transform of  $f(t)$  be defined as

$$F(\omega) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt.$$

Suppose  $\frac{dF(\omega)}{d\omega} = -\omega F(\omega)$  for all  $\omega$ , and  $F(0) = 1$ .

Then

- (a)  $f(0) < 1$
- (b)  $f(0) > 1$
- (c)  $f(0) = 1$
- (d)  $f(0) = 0$

- 6. [NAT]**

[GATE-CE-2020-2M]

The Fourier series to represent  $x - x^2$  for  $-\pi \leq x \leq \pi$  is given by

$$x - x^2 = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$$

The value of  $a_0$  (round off to two decimal places), is

\_\_\_\_\_.

**7. [MCQ]****[GATE-EE-2019:2M]**

A periodic function  $f(t)$ , with a period of  $2\pi$ , is represented as its Fourier series,

$$f(t) = a_0 + \sum_{n=1}^{\infty} a_n \cos nt + \sum_{n=1}^{\infty} b_n \sin nt$$

If  $f(t) = \begin{cases} A \sin t, & 0 \leq t \leq \pi \\ 0, & \pi < t < 2\pi \end{cases}$ , the Fourier series coefficients  $a_1$  and  $b_1$  of  $f(t)$  are

- (a)  $a_1 = \frac{A}{\pi}; b_1 = 0$       (b)  $a_1 = \frac{A}{2}; b_1 = 0$   
 (c)  $a_1 = 0; b_1 = \frac{A}{\pi}$       (d)  $a_1 = 0; b_1 = \frac{A}{2}$

**8. [MCQ]****[GATE-ME-2018:1M]**

The fourier cosine series for an even function  $f(x)$  is

given by  $f(t) = a_0 + \sum_{n=1}^{\infty} a_n \cos(nx)$ . The value of the

coefficient  $a_2$  for the function  $f(x) = \cos^2(x)$  in  $[0, \pi]$  is

- (a) -0.5      (b) 0.00  
 (c) 0.5      (d) 1.0

**9. [MCQ]****[GATE-EC-2017:1M]**

For the function  $f(x) = \begin{cases} -2 & -\pi < x < 0 \\ 2 & 0 < x < \pi \end{cases}$ . The value

of  $a_n$  in the Fourier Series expansion of  $f(x)$  is

- (a) 2      (b) 4  
 (c) 0      (d) -2

**10. [MCQ]****[GATE-CE-2016-1M]**

The Fourier series of the function

$$f(x) = \begin{cases} 0 & -\pi < x \leq 0 \\ \pi - x & 0 < x < \pi \end{cases}$$

In the interval  $[-\pi, \pi]$  is

$$f(x) = \frac{\pi}{4} + \frac{2}{\pi} \left[ \frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \dots \right] + \left[ \frac{\sin x}{1} + \frac{\sin 2x}{2} + \frac{\sin 3x}{3} + \dots \right]$$

The convergence of the above Fourier series at  $x = 0$  gives

- (a)  $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$       (b)  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} = \frac{\pi^2}{12}$   
 (c)  $\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$       (d)  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n-1} = \frac{\pi}{4}$

**11. [NAT]****[GATE-EE-2014:2M]**

Let  $g: [0, \infty) \rightarrow [0, \infty)$  be a function defined by  $g(x) = x - [x]$ , where  $[x]$  represents the integer part of  $x$ . (That is, the largest integer which is less than or equal to  $x$ ). The value of the constant term in the Fourier Series expansion of  $g(x)$  is \_\_\_\_.

**12. [MCQ]****[GATE-EE-2010:1M]**

The period of the signal  $x(t) = 8 \sin \left( 0.8\pi t + \frac{\pi}{4} \right)$

- (a)  $0.4\pi$  s      (b)  $0.8\pi$  s  
 (c) 1.25 s      (d) 2.5 s




**ANSWER KEY**

- |                  |                            |                         |                |
|------------------|----------------------------|-------------------------|----------------|
| <b>1.</b> (b, c) | <b>2.</b> (0.120 to 0.130) | <b>3.</b> (a)           | <b>4.</b> (b)  |
| <b>5.</b> (a)    | <b>6.</b> (-6.28 to -6.68) | <b>7.</b> (d)           | <b>8.</b> (c)  |
| <b>9.</b> (c)    | <b>10.</b> (c)             | <b>11.</b> (0.5 to 0.5) | <b>12.</b> (d) |


**SOLUTIONS**
**1. (b, c)**

$$f(x) = px^4 + qx^5$$

Using Fourier coefficient formula,

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cdot \sin\left(\frac{n\pi x}{L}\right) dx$$

$$= \frac{1}{L} \int_{-L}^L [px^4 + qx^5] \sin\left(\frac{n\pi x}{L}\right) dx$$

$$= \frac{1}{L} \int_{-L}^L px^4 \sin\left(\frac{n\pi x}{L}\right) dx + \frac{1}{L} \int_{-L}^L qx^5 \sin\left(\frac{n\pi x}{L}\right) dx$$

$$a_n = 0 + \frac{1}{L} \int_{-L}^L qx^5 \sin\left(\frac{n\pi x}{L}\right) dx$$

$\left[ \because px^4 \sin\left(\frac{n\pi x}{L}\right)$  is an odd function  $\right]$

So,  $a_n$  depends on q.

$$f(x) = px^4 + qx^5$$

Using Fourier coefficient formula,


$$b_n = \frac{1}{L} \int_{-L}^L f(x) \cdot \cos\left(\frac{n\pi x}{L}\right) dx$$

$$= \frac{1}{L} \int_{-L}^L [px^4 + qx^5] \cos\left(\frac{n\pi x}{L}\right) dx$$

$$= \frac{1}{L} \int_{-L}^L px^4 \cos\left(\frac{n\pi x}{L}\right) dx + \frac{1}{L} \int_{-L}^L qx^5 \cos\left(\frac{n\pi x}{L}\right) dx$$

$$b_n = \frac{1}{L} \int_{-L}^L px^4 \cos\left(\frac{n\pi x}{L}\right) dx + 0$$

$\left[ \because qx^5 \cos\left(\frac{n\pi x}{L}\right)$  is an odd function  $\right]$

So,  $b_n$  depends on p.**2. (0.120 to 0.130)**

$$f(x) = \cos^4 x = (\cos^2 x)^2 = \left[ \frac{1 + \cos 2x}{2} \right]^2$$

$$\Rightarrow f(x) = \frac{1}{4} [1 + \cos^2 2x + 2 \cos 2x]$$

$$= \frac{1}{4} \left[ 1 + \frac{1 + \cos 4x}{2} + 2 \cos 2x \right]$$

$$= \frac{1}{4} + \frac{1}{8} + \frac{\cos 4x}{8} + \frac{\cos 2x}{2}$$

$$= \frac{3}{8} + 0 \cos x + \frac{1}{2} \cos 2x + 0 \cos 3x + \frac{1}{8} \cos 4x + 0 \cos 5x + \dots$$

∴ From cosine Fourier series

$$f_4 = \frac{1}{8} \text{ and } f_5 = 0$$

$$\therefore f_4 + f_5 = \frac{1}{8} = 0.125$$



Scan for Video solution



### 3. (a)

**Given data:**

$$f(x) = x^3 \quad [-1, 1]$$

The General Fourier Series Expansion is

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{c}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{c}\right)$$

Comparing ( $\alpha, \alpha + 2c$ ) with  $[-1, 1]$

$$\Rightarrow \alpha = -1; c = 1$$

$$a_0 = \frac{1}{c} \int_{-1}^1 f(x) dx = \frac{1}{c} \int_{-1}^1 x^3 dx = 0$$

$\{\because f(x) \text{ is an odd function}\}$

$$a_n = \frac{1}{c} \int_{-1}^1 f(x) \cos nx dx = \frac{1}{c} \int_{-1}^1 x^3 \cos nx dx = 0$$

$$b_n = \frac{1}{c} \int_{-1}^1 f(x) \sin nx dx = \frac{1}{c} \int_{-1}^1 x^3 \sin nx dx \neq 0$$

∴ The Fourier Series expansion of  $x^3$  in  $[-1, 1]$  has only sine terms in it.


Scan for Video solution



### 4. (b)

**Given data:**  $f(t)$  takes two values: 4 and 0.

Let us consider  $[-1, 1]$

( $\because$  since the period of function is 2 seconds)

The general Fourier Series expansion is given by

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{c}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{c}\right)$$

Where

$$a_0 = \frac{1}{c} \cdot \int_{\alpha}^{\alpha+2c} f(x) dx = \frac{1}{1} \int_{-1}^1 f(x) dx \\ = \int_{-1}^0 0 dx + \int_0^1 4 dx = 4$$

Since  $a_0 = 4$

$$\Rightarrow \frac{a_0}{2} = 2$$


Scan for Video solution



### 5. (a)

Given function is even

$$\frac{d}{d\omega} F(\omega) = -\omega F(\omega) \quad \dots (1)$$

From differentiation property,

$$t f(t) = j \frac{d}{d\omega} F(\omega)$$

Applying inverse Fourier transfer to the above equation,

$$-j t f(t) = j \frac{d}{d\omega} f(t)$$

$$\frac{d}{dt} f(t) = -t f(t) \quad \dots (2)$$

From equation (1) and (2) it is clear that  $f(t)$  is Gaussian function, it can be written as,

$$f(t) = \frac{1}{\sqrt{2\pi}} e^{-t^2/2}$$

$$\therefore f(0) = \frac{1}{\sqrt{2\pi}} = 0.3989$$

$$\therefore f(0) < 1$$



Scan for Video solution



#### 6. (-6.28 to -6.68)

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cdot dx$$

$$= \frac{1}{\pi} \int_{-\pi}^{\pi} (x - x^2) dx$$

$$= \frac{1}{\pi} \cdot 2 \int_0^{\pi} -x^2 dx \quad [\because x \text{ is an odd function}]$$

$$= -\frac{2}{\pi} \left[ \frac{x^3}{3} \right]_0^{\pi} = -\frac{2}{\pi} \cdot \frac{\pi^3}{3}$$

$$= -\frac{2\pi^2}{3} = -6.58$$



Scan for Video solution



#### 7. (d)

**Given data:**  $f(t) = \begin{cases} A \sin t; & 0 \leq t \leq \pi \\ 0; & \pi < t < 2\pi \end{cases}$

$$a_1 = \frac{1}{\pi} \cdot \int_0^{2\pi} f(t) \cdot \cos t \cdot dt$$

$$\Rightarrow a_1 = \frac{1}{\pi} \cdot \left\{ \int_0^{\pi} A \sin t \cdot \cos t \cdot dt \right\}$$

$$\Rightarrow a_1 = \frac{1}{\pi} \times \frac{A}{2} \times \int_0^{\pi} \sin 2t \cdot dt$$

$$\Rightarrow a_1 = \frac{A}{2\pi} \times \left\{ \frac{-\cos 2t}{2} \Big|_0^{\pi} \right\} = \frac{-A}{4\pi} \{1 - 1\} = 0$$

$$b_1 = \frac{1}{\pi} \cdot \int_0^{2\pi} f(t) \cdot \sin t \cdot dt$$

$$\Rightarrow b_1 = \frac{1}{\pi} \cdot \left\{ \int_0^{\pi} A \sin t \cdot \sin t \cdot dt \right\}$$

$$\Rightarrow b_1 = \frac{A}{\pi} \times \int_0^{\pi} \sin^2 t \cdot dt$$

$$\Rightarrow b_1 = \frac{2A}{\pi} \times \int_0^{\pi/2} \sin^2 t \cdot dt = \frac{2A}{\pi} \times \frac{1}{2} \times \frac{\pi}{2}$$

$$b_1 = \frac{A}{2} \text{ & } a_1 = 0$$



Scan for Video solution



#### 8. (c)

**Given data:**  $f(x) = \cos^2 x, [0, \pi]$

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cdot \cos nx$$

We know that

$$\cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\Rightarrow \cos^2 x = \frac{1}{2} + \frac{1}{2} \cdot \cos 2x = a_0 + a_2 \cdot \cos 2x$$

On comparing  $\Rightarrow a_2 = \frac{1}{2} = 0.5$



Scan for Video solution



9. (c)

**Given data:**

$$f(x) = \begin{cases} -2; & -\pi < x < 0 \\ 2; & 0 < x < \pi \end{cases}$$

The general Fourier Series expansion of  $f(x)$  is given by

$$\text{by } f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$$

$$\text{Where } a_n = \frac{1}{\pi} \cdot \int_{-\pi}^{\pi} f(x) \cdot \cos nx dx$$

$$\Rightarrow a_n = \frac{1}{\pi} \times \left\{ \int_{-\pi}^0 -2 \cdot \cos nx dx + \int_0^{\pi} 2 \cdot \cos nx dx \right\}$$

$$\Rightarrow \frac{1}{\pi} \cdot \left\{ -2 \cdot \frac{\sin nx}{n} \Big|_{-\pi}^0 + 2 \cdot \frac{\sin nx}{n} \Big|_0^{\pi} \right\}$$

$$\Rightarrow \frac{1}{\pi} \cdot \left\{ \frac{-2}{n}(0) + 2(0) \right\}$$

$$= 0$$

$$\therefore a_n = 0$$



Scan for Video solution



10. (c)

$$\text{The function } f(x) = \begin{cases} 0 & -\pi < x < 0 \\ \pi - x & 0 \leq x < \pi \end{cases}$$

And Fourier series is

$$f(x) = \frac{\pi}{4} + \frac{2}{\pi} \left[ \frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \dots \right]$$

$$+ \left[ \frac{\sin x}{1} + \frac{\sin 2x}{2} + \frac{\sin 3x}{3} + \dots \right]$$

At  $x = 0$ , it is a point of discontinuity, the Fourier series converges to  $\frac{1}{2} [f(0^-) + f(0^+)]$

$$\text{Where } f(0^-) = \lim_{x \rightarrow 0^-} (\pi - x) = \pi$$

$$f(0^+) = 0$$

Put  $x = 0$  in fouries series,

$$f(0) = \frac{\pi}{4} + \frac{2}{\pi} \left[ \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

$$\frac{\pi}{2} = \frac{\pi}{4} + \frac{2}{\pi} \left[ \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

$$\frac{\pi}{2} \left( \frac{\pi}{2} - \frac{\pi}{4} \right) = \left[ \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right]$$

$$\sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}$$




Scan for Video solution



11. (0.5 to 0.5)

**Given data:**  $f(x) = x - [x] = \{x\}$



From the graph, period of  $f(x) = 1$

The general Fourier Series expansion of  $f(x)$  in the interval  $(\alpha, \alpha + 2c)$  is given by

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{c}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{c}\right)$$

$$\text{Where } a_0 = \frac{1}{c} \int_{\alpha}^{\alpha+2c} f(x) dx$$

Considering the interval [0, 1]

$$\Rightarrow \alpha = 0 \text{ and } c = \frac{1}{2}$$

$$\begin{aligned} \frac{a_0}{2} &= \frac{1}{2} \left\{ \frac{1}{c} \int_{\alpha}^{\alpha+2c} f(x) dx \right\} \\ &= \frac{1}{2} \times \frac{1}{\left(\frac{1}{2}\right)} \times \int_0^1 (x - [x]) dx \\ \Rightarrow \frac{a_0}{2} &= \int_0^1 (x - 0) dx = \frac{1}{2} \\ \Rightarrow \frac{a_0}{2} &= \frac{1}{2} = 0.5 \end{aligned}$$



Scan for Video solution



12. (d)

The given signal is

$$x(t) = 8 \cdot \sin\left(0.8\pi t + \frac{\pi}{4}\right)$$

Comparing with the standard form of the signal

$$x(t) = A \cdot \sin(\omega t + \phi)$$

$$\Rightarrow \omega = 0.8\pi$$

$$\Rightarrow \frac{2\pi}{T} = 0.8\pi$$

$$\Rightarrow T = \frac{2}{0.8} = 2.5 \text{ s}$$

$$\therefore T = 2.5 \text{ s}$$



Scan for Video solution



## CHAPTER

# 8

# LAPLACE TRANSFORMS

### Laplace Transforms of Standard Functions

**1. [MCQ] [GATE-ME-2023:2M]**

Which of the following options given is the inverse Laplace transform of  $\frac{1}{s^3 - s}$ ?

$u(t)$  denotes the unit-step function.

- (a)  $\left(-1 + \frac{1}{2}e^{-t} + \frac{1}{2}e^t\right)u(t)$
- (b)  $\left(\frac{1}{3}e^{-t} - e^t\right)u(t)$
- (c)  $\left(-1 + \frac{1}{2}e^{-(t-1)} + \frac{1}{2}e^{(t-1)}\right)u(t-1)$
- (d)  $\left(-1 - \frac{1}{2}e^{-(t-1)} - \frac{1}{2}e^{(t-1)}\right)u(t-1)$

**2. [MCQ] [GATE-ME-2022:1M]**

Solution of  $\nabla^2 T = 0$  in a square domain ( $0 < x < 1$  and  $0 < y < 1$ ) with boundary conditions:

$T(x, 0) = x$ ;  $T(0, y) = y$ ;  $T(x, 1) = 1 + x$ ;  $T(1, y) = 1 + y$  is

- (a)  $T(x, y) = x - xy + y$
- (b)  $T(x, y) = x + y$
- (c)  $T(x, y) = -x + y$
- (d)  $T(x, y) = x + xy + y$

**3. [MCQ] [GATE-ME-2021:1M]**

The Dirac-delta function ( $\delta(t - t_0)$ ) for  $t, t_0 \in \mathbb{R}$ , has the following property.

$$\int_a^b \varphi(t)\delta(t - t_0)dt = \begin{cases} \varphi(t_0) & a < t_0 < b \\ 0 & \text{otherwise} \end{cases}$$

The Laplace transform of the Dirac-delta function

$\delta(t - a)$  for  $a > 0$ ;  $L(\delta(t - a)) = F(s)$  is

- (a) 0
- (b)  $\infty$
- (c)  $e^{sa}$
- (d)  $e^{-sa}$

**4. [MCQ] [GATE-ME-2021:1M]**

If the Laplace transform of a function  $f(t)$  is given by  $\frac{s+3}{(s+1)(s+2)}$  then  $f(0)$  is.

- (a) 0
- (b)  $\frac{1}{2}$
- (c) 1
- (d)  $\frac{3}{2}$

**5. [MCQ] [GATE-ME-2020:1M]**

The Laplace transform of a function  $f(t)$  is

$$L(f) = \frac{1}{(s^2 + \omega^2)}. \text{ Then, } f(t) \text{ is}$$

- (a)  $f(t) = \frac{1}{\omega^2}(1 - \cos \omega t)$
- (b)  $f(t) = \frac{1}{\omega} \cos \omega t$
- (c)  $f(t) = \frac{1}{\omega} \sin \omega t$
- (d)  $f(t) = \frac{1}{\omega^2}(1 - \sin \omega t)$

**6. [MCQ] [GATE-CE-2019:1M]**

The Laplace transform of  $\sin h(at)$  is

- (a)  $\frac{s}{s^2 + a^2}$
- (b)  $\frac{s}{s^2 - a^2}$
- (c)  $\frac{a}{s^2 - a^2}$
- (d)  $\frac{a}{s^2 + a^2}$

**7. [MCQ]**

[GATE-CE-2018:2M]

The Laplace transform  $F(s)$  of the exponential function  $f(t) = e^{at}$  when  $t \geq 0$ , where  $a$  is a constant and  $(s-a) > 0$ , is

- (a)  $\frac{1}{s+a}$       (b)  $\frac{1}{s-a}$   
 (c)  $\frac{1}{a-s}$       (d)  $\infty$

**8. [NAT]**

[GATE-ME-2018:2M]

$F(s)$  is the Laplace transform of the function

$$f(t) = 2t^2 e^{-t}$$

$F(1)$  is \_\_\_\_\_ (Correct to two decimal places).

**9. [MCQ]**

[GATE-ME-2017:1M]

The Laplace transform of  $t e^t$  is

- (a)  $\frac{s}{(s+1)^2}$       (b)  $\frac{1}{(s-1)^2}$   
 (c)  $\frac{1}{(s+1)^2}$       (d)  $\frac{s}{s-1}$

**10. [MCQ]**

[GATE-ME-2016:1M]

If  $f(t)$  is a function defined for all  $t \geq 0$ , its Laplace transform  $F(s)$  is defined as

- (a)  $\int_0^\infty e^{st} f(t) dt$       (b)  $\int_0^\infty e^{-st} f(t) dt$   
 (c)  $\int_0^\infty e^{ist} f(t) dt$       (d)  $\int_0^\infty e^{-ist} f(t) dt$

**11. [MCQ]**

[GATE-ME-2016:1M]

Laplace transform of  $\cos(\omega t)$  is

- (a)  $\frac{s}{s^2 + \omega^2}$       (b)  $\frac{\omega}{s^2 + \omega^2}$   
 (c)  $\frac{s}{s^2 - \omega^2}$       (d)  $\frac{\omega}{s^2 - \omega^2}$

**12. [MCQ]**

[GATE-CE-2016:2M]

The value of  $\int_0^\infty \frac{1}{1+x^2} dx + \int_0^\infty \frac{\sin x}{x} dx$  is

- (a)  $\frac{\pi}{2}$       (b)  $\pi$   
 (c)  $\frac{3\pi}{2}$       (d) 1

**13. [NAT]**

[GATE-2015-EC: 2M]

The value of the integral

$$\int_{-\infty}^{\infty} 12 \cos(2\pi t) \frac{\sin(4\pi t)}{4\pi t} dt \text{ is } \underline{\hspace{2cm}}$$

**14. [MCQ]**

[GATE-ME-2015:1M]

The Laplace transform of  $e^{it}$  where  $i = \sqrt{-1}$ , is


- (a)  $\frac{s-5i}{s^2-25}$       (b)  $\frac{s+5i}{s^2+25}$   
 (c)  $\frac{s+5i}{s^2-25}$       (d)  $\frac{s-5i}{s^2+25}$

**15. [MCQ]**

[GATE-ME-2015:2M]

Laplace transform of the function  $f(t)$  is given by

$F(s) = L\{f(t)\} = \int_0^\infty f(t) e^{-st} dt$ . Laplace transform of the function shown below is given by



- (a)  $\frac{1-e^{-2s}}{s}$       (b)  $\frac{1-e^{-s}}{2s}$   
 (c)  $\frac{2-2e^{-s}}{s}$       (d)  $\frac{1-2e^{-s}}{s}$

**16. [MCQ]**

[GATE-ME-2014:1M]

Laplace transform of  $\cos(\omega t)$  is  $\frac{s}{s^2 + \omega^2}$ . The

Laplace transformation of  $e^{-2t} \cos(4t)$  is

- (a)  $\frac{s-2}{(s-2)^2+16}$       (b)  $\frac{s+2}{(s-2)^2+16}$   
 (c)  $\frac{s-2}{(s+2)^2+16}$       (d)  $\frac{s+2}{(s+2)^2+16}$

**17. [MCQ] [GATE-ME-2013: 2M]**

The function  $f(t)$  satisfies the differential equation  $\frac{d^2 f}{dt^2} + f = 0$  and the auxiliary conditions,

$f(0) = 0, \frac{df}{dt}(0) = 4$ . The Laplace transform of  $f(t)$  is given by

- |                       |                       |
|-----------------------|-----------------------|
| (a) $\frac{2}{s+1}$   | (b) $\frac{4}{s+1}$   |
| (c) $\frac{4}{s^2+1}$ | (d) $\frac{2}{s^4+1}$ |

**18. [MCQ] [GATE-ME-2012:2M]**

The inverse Laplace transform of the function  $F(s) = \frac{1}{s(s+1)}$  is given by

- |                     |                            |
|---------------------|----------------------------|
| (a) $f(t) = \sin t$ | (b) $f(t) = e^{-t} \sin t$ |
| (c) $f(t) = e^{-t}$ | (d) $f(t) = 1 - e^{-t}$    |

**19. [MCQ] [GATE-ME-2010:2M]**

The Laplace transform of a function  $f(t)$  is  $\frac{1}{s^2(s+1)}$ .

The function  $f(t)$  is

- |                      |                      |
|----------------------|----------------------|
| (a) $t - 1 + e^{-t}$ | (b) $t + 1 + e^{-t}$ |
| (c) $-1 + e^{-t}$    | (d) $2t + e^t$       |

**20. [MCQ] [GATE-ME-2009:1M]**

The inverse Laplace transform of  $\frac{1}{(s^2+s)}$  is

- |                  |                  |
|------------------|------------------|
| (a) $1 + e^t$    | (b) $1 - e^t$    |
| (c) $1 - e^{-t}$ | (d) $1 + e^{-t}$ |

**21. [MCQ] [GATE-CE-2009:2M]**

Laplace transform for the function  $f(x) = \cos h(ax)$  is

- |                         |                         |
|-------------------------|-------------------------|
| (a) $\frac{a}{s^2-a^2}$ | (b) $\frac{s}{s^2-a^2}$ |
| (c) $\frac{a}{s^2+a^2}$ | (d) $\frac{s}{s^2+a^2}$ |

**Properties of Laplace Transforms****22. [MCQ] [GATE-ME-2020:1M]**

The solution of

$$\frac{d^2 y}{dt^2} - y = 1,$$

which additionally satisfies  $y|_{t=0} = \frac{dy}{dt}|_{t=0} = 0$  in the

Laplace s-domain is

- |                             |                        |
|-----------------------------|------------------------|
| (a) $\frac{1}{s(s+1)(s-1)}$ | (b) $\frac{1}{s(s+1)}$ |
| (c) $\frac{1}{s(s-1)}$      | (d) $\frac{1}{(s-1)}$  |

**23. [MCQ] [GATE-EE 2016:2M]**

The value of the integral  $2 \int_{-\infty}^{\infty} \left( \frac{\sin 2\pi t}{\pi t} \right) dt$  is equal to

- |       |         |
|-------|---------|
| (a) 0 | (b) 0.5 |
| (c) 1 | (d) 2   |

**24. [MCQ] [GATE-EE-2014:1M]**

Let  $X(s) = \frac{3s+5}{s^2+10s+21}$  be the Laplace Transform

of a signal  $x(t)$ . Then,  $x(0^+)$  is

- |       |        |
|-------|--------|
| (a) 0 | (b) 3  |
| (c) 5 | (d) 21 |

**25. [MCQ] [GATE-2012 : 2M]**

Consider the differential equation

$$\frac{d^2 y(t)}{dt^2} + 2 \frac{dy(t)}{dt} + y(t) = \delta(t) \text{ with}$$

$y(t)|_{t=0} = -2$  and  $\frac{dy}{dt}|_{t=0^+} = 0$  The numerical value of

$$\frac{dy}{dt}|_{t=0^-}$$

- |        |        |
|--------|--------|
| (a) -2 | (b) -1 |
| (c) 0  | (d) 1  |


**ANSWER KEY**

- |              |         |         |                   |
|--------------|---------|---------|-------------------|
| 1. (a)       | 2. (b)  | 3. (d)  | 4. (c)            |
| 5. (c)       | 6. (c)  | 7. (b)  | 8. (0.48 to 0.52) |
| 9. (b)       | 10. (b) | 11. (a) | 12. (b)           |
| 13. (3 to 3) | 14. (b) | 15. (c) | 16. (d)           |
| 17. (c)      | 18. (d) | 19. (a) | 20. (c)           |
| 21. (b)      | 22. (a) | 23. (d) | 24. (b)           |
| 25. (d)      |         |         |                   |


**SOLUTIONS**

1. (a)  
Given

$$L\{f(t)\} = \frac{1}{s^3 - s}$$

$u(t)$  is unit step function

$$\Rightarrow u(t) = 1 \quad \forall t > 0$$

$$\Rightarrow L\{f(t)\} = \frac{1}{s(s^2 - 1)}$$

$$\Rightarrow f(t) = L^{-1}\left\{\frac{1}{s(s-1)(s+1)}\right\}$$

$$\Rightarrow = L^{-1}\left\{-\frac{1}{s} + \frac{(1/2)}{s-1} + \frac{(1/2)}{s+1}\right\}$$

$$\Rightarrow f(t) = L^{-1}\left\{-\frac{1}{s}\right\} + \frac{1}{2} \cdot L^{-1}\left\{\frac{1}{s-1}\right\} + \frac{1}{2} \cdot L^{-1}\left\{\frac{1}{s+1}\right\}$$

$$\Rightarrow f(t) = -e^{0t} + \frac{1}{2} \cdot e^t + \frac{1}{2} \cdot e^{-t}$$

$$\left(\because L\{e^{at}\} = \frac{1}{s-a}\right)$$

$$\Rightarrow f(t) = -1 + \frac{1}{2}e^t + \frac{1}{2}e^{-t}$$

Since  $u(t) = 1$

$$\Rightarrow f(t) = \left(-1 + \frac{1}{2} \cdot e^t + \frac{1}{2} \cdot e^{-t}\right)u(t)$$




Scan for Video solution



2. (b)  
Given

$$\begin{aligned} \nabla^2 T &= 0 \\ T(x, 0) &= x; \\ T(x, 1) &= 1 + x; \\ T(0, y) &= y, \\ T(1, y) &= 1 + y \end{aligned}$$



From given options,  $T(x, y) = x + y$  is the only option that satisfies the boundary condition

$$T(x, 1) = 1 + x$$

$$\therefore T(x, y) = x + y$$

Scan for Video solution



3. (d)  
Given

For Dirac-delta function,  $\delta(t - t_0)$

$$\int_a^b \phi(t) \cdot \delta(t - t_0) dt = \begin{cases} \phi(t_0); & a < t < b \\ 0; & \text{otherwise} \end{cases}$$

$$\text{For } 0 < t < \infty \quad \int_0^\infty \phi(t) \cdot \delta(t - t_0) dt = \phi(t_0)$$

For  $\phi(t) = e^{-st}$  and  $t_0 = a$

$$\Rightarrow \int_0^\infty e^{-st} \cdot \delta(t-a) dt = \phi(a) = e^{-sa}$$

$$\Rightarrow L\{\delta(t-a)\} = e^{-sa} \quad \left\{ \because \int_0^\infty e^{-st} \cdot f(t) dt = L\{f(t)\} \right\}$$



Scan for Video solution



#### 4. (c)

**Given**

$$L\{f(t)\} = \frac{s+3}{(s+1)(s+2)}$$

$$\Rightarrow L\{f(t)\} = \frac{2}{s+1} - \frac{1}{s+2}$$

Applying  $L^{-1}$  on both sides

$$\Rightarrow L^{-1}\{L\{f(t)\}\} = L^{-1}\left\{\frac{2}{s+1} - \frac{1}{s+2}\right\}$$

$$\Rightarrow f(t) = 2 \cdot e^{-t} - e^{-2t}$$

$$\left( \because L\{e^{at}\} = \frac{1}{s-a} \right)$$

$$\therefore f(0) = 2 \cdot e^0 - e^{-2(0)} = 2 - 1 = 1$$

$$\Rightarrow f(0) = 1$$



Scan for Video solution



#### 5. (c)

**Given**

$$L\{f(t)\} = \frac{1}{s^2 + \omega^2}$$

$$\text{We have } L\{\sin \omega t\} = \frac{\omega}{s^2 + \omega^2}$$

$$\Rightarrow \frac{1}{s^2 + \omega^2} = \frac{1}{\omega} \cdot L\{\sin \omega t\} = L\left\{\frac{1}{\omega} \cdot \sin \omega t\right\}$$

$$\therefore f(t) = \frac{1}{\omega} \cdot \sin \omega t$$



Scan for Video solution



#### 6. (c)

$$\sinh at = \frac{e^{at} - e^{-at}}{2}$$

Taking Laplace Transform, we get

$$= \frac{1}{2} \left[ L\{e^{at}\} - L\{e^{-at}\} \right]$$

$$= \frac{1}{2} \left[ \frac{1}{s-a} - \frac{1}{s+a} \right]$$

$$= \frac{1}{2} \left[ \frac{1}{s-a} - \frac{1}{s+a} \right]$$

$$= \frac{1}{2} \left[ \frac{2a}{s^2 - a^2} \right]$$

$$L[\sinh at] = \frac{a}{s^2 - a^2}$$



Scan for Video solution



#### 7. (b)

$$L[f(t)] = \int_0^\infty e^{-st} \cdot f(t) dt$$

$$L[e^{at}] = \int_0^\infty e^{-st} \cdot e^{at} dt = \int_0^\infty e^{-(s-a)t} dt$$

$$\left[ \frac{e^{-(s-a)t}}{-(s-a)} \right]_0^\infty = -\frac{1}{s-a}(0-1)$$

$$L(e^{at}) = \frac{1}{s-a}$$



Scan for Video solution



#### 8. (0.48 to 0.52)

**Given,**

$$f(t) = 2 \cdot t^2 \cdot e^{-t}$$

$$L\{e^{-t}\} = \frac{1}{s+1} \quad \left\{ \because L\{e^{at}\} = \frac{1}{s-a} \right\}$$

$$L\{t^n \cdot e^{at}\} = (-1)^n \cdot \frac{d^n}{ds^n} \left( \frac{1}{s+a} \right)$$

$$\begin{aligned} \Rightarrow L\{t^2 \cdot e^{-t}\} &= (-1)^2 \cdot \frac{d^2}{ds^2} \left( \frac{1}{s+1} \right) \\ &= \frac{d}{ds} \left( \frac{-1}{(s+1)^2} \right) = \frac{2}{(s+1)^3} \end{aligned}$$

$$\therefore L\{2 \cdot t^2 \cdot e^{-t}\} = \frac{4}{(s+1)^3} = F(s)$$

$$\therefore F(s) = \frac{4}{(s+1)^3}$$

$$\Rightarrow F(1) = \frac{4}{(1+1)^3} = \frac{4}{8} = 0.5$$

$$\Rightarrow \boxed{F(1) = 0.5}$$



Scan for Video solution


**9. (b)**

$$L\{t \cdot e^t\} = ?$$

$$L\{e^{at}\} = \frac{1}{s-a}$$

$$\Rightarrow L\{t^n \cdot e^{at}\} = (-1)^n \cdot \frac{d^n}{ds^n} \left\{ \frac{1}{s-a} \right\}$$

$$\Rightarrow L\{t \cdot e^{at}\} = (-1) \cdot \frac{d}{ds} \left( \frac{1}{s-a} \right) = \frac{1}{(s-a)^2}$$

$$\Rightarrow \boxed{L\{t \cdot e^t\} = \frac{1}{(s-1)^2}} \quad \text{For } a=1$$



Scan for Video solution


**10. (b)**

For a function  $f(t)$  defined  $\forall t \geq 0$ ; the Laplace transform of  $f(t)$  is given by

$$\boxed{L\{f(t)\} = \int_0^\infty e^{-st} \cdot f(t) dt}$$



Scan for Video solution


**11. (a)**

$$L\{\cos \omega t\} = \int_0^\infty e^{-st} \cdot \cos \omega t dt = \frac{s}{s^2 + \omega^2}$$



Scan for Video solution


**12. (b)**

$$I = \int_0^\infty \frac{1}{1+x^2} dx + \int_0^\infty \frac{\sin x}{x} dx = ?$$

$$I = I_1 + I_2$$

Where,

$$I_1 = \int_0^\infty \frac{1}{1+x^2} dx = \left[ \tan^{-1} x \right]_0^\infty$$

$$I_2 = \int_0^\infty \frac{\sin x}{x} dx$$

 Considering Laplace Transform of  $\sin at$ 

$$\Rightarrow L\{\sin at\} = \frac{a}{s^2 + a^2}$$

By the Property of division with 't'

$$\Rightarrow L\left\{ \frac{\sin at}{t} \right\} = \int_s^\infty \frac{a}{s^2 + a^2} \cdot ds$$

$$\Rightarrow \int_s^\infty e^{-st} \cdot \frac{\sin at}{t} \cdot dt = \frac{1}{a} \cdot \left[ \tan^{-1} \left( \frac{s}{a} \right) \right]_s^\infty$$

$$\Rightarrow \int_s^\infty e^{-st} \cdot \frac{\sin at}{t} dt = \frac{1}{a} \cdot \left\{ \frac{\pi}{2} - \left[ \tan^{-1} \left( \frac{s}{a} \right) \right]_s^\infty \right\}$$

 Substituting  $s=0$  and  $a=1$  in both sides of above equation

$$\Rightarrow \int_s^{\infty} \frac{\sin at}{t} dt = 1 \left\{ \frac{\pi}{2} - 0 \right\} = \frac{\pi}{2}$$

$$\therefore \int_s^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}$$

**Therefore,**

$$I = \left[ \tan^{-1} x \right]_0^{\infty} + \int_0^{\infty} \frac{\sin x}{x} dx$$

$$\left[ \because \int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2} \right]$$

$$I = \tan^{-1} \infty - \tan^{-1} 0 + \frac{\pi}{2}$$

$$I = \frac{\pi}{2} - 0 + \frac{\pi}{2} = \pi$$

$$\therefore \int_0^{\infty} \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{\sin x}{x} dx = \pi$$



Scan for Video solution



### 13. (3 to 3)

Let the given integral,

$$I = \int_{-\infty}^{\infty} 12 \cdot \cos(2\pi t) \frac{\sin(4\pi t)}{4\pi t} dt$$

$$= 2 \times \int_0^{\infty} 12 \cdot \cos(2\pi t) \frac{\sin(4\pi t)}{4\pi t} dt$$

$$(\because \int_{-\infty}^{\infty} f(t) dt = 2 \times \int_0^{\infty} f(t) dt, \text{ If } f(-t) = f(t))$$

$$\Rightarrow I = 3 \times \int_0^{\infty} \frac{2 \sin(4\pi t) \cdot \cos(2\pi t)}{\pi t} dt$$

$$\Rightarrow I = \frac{3}{\pi} \times \int_0^{\infty} \frac{\sin(6\pi t) + \sin(2\pi t)}{t} dt$$

$$\left( \because \sin A + \sin B = 2 \sin \left( \frac{A+B}{2} \right) \cdot \cos \left( \frac{A-B}{2} \right) \right)$$

$$\Rightarrow I = 3 \times \left\{ \int_0^{\infty} \frac{\sin(6\pi t)}{\pi t} dt + \int_0^{\infty} \frac{\sin(2\pi t)}{\pi t} dt \right\}$$

$$I = I_1 + I_2$$

$$I_1 = 3 \times \int_0^{\infty} \frac{\sin 6\pi t}{\pi t} dt$$

Let when

$$\begin{aligned} 6\pi t &= u \\ 6\pi dt &= du \\ dt &= \frac{du}{6\pi} \end{aligned} \quad \left| \begin{array}{l} t=0, u=0 \\ t=\infty, u=\infty \end{array} \right.$$

$$\therefore I_1 = 3 \int_0^{\infty} \frac{\sin u}{u/6} \cdot \frac{du}{6\pi} = \frac{3}{\pi} \int_0^{\infty} \frac{\sin u}{u} du$$

$$\text{We know that } \int_0^{\infty} \frac{\sin t}{t} dt = \frac{\pi}{2}$$

$$\therefore I_1 = \frac{3}{\pi} \int_0^{\infty} \frac{\sin u}{u} dv = \frac{3}{\pi} \times \frac{\pi}{2} = \frac{3}{2}$$

$$\text{Now, } I_2 = 3 \int_0^{\infty} \frac{\sin 2\pi t}{\pi t} dt$$

Let when

$$\begin{aligned} 6\pi t &= v \\ 6\pi dt &= dv \\ dt &= \frac{dv}{2\pi} \end{aligned} \quad \left| \begin{array}{l} t=0, v=0 \\ t=\infty, v=\infty \end{array} \right.$$

$$\therefore I_2 = 3 \int_0^{\infty} \frac{\sin v}{v/2} \cdot \frac{dv}{2\pi} = \frac{3}{\pi} \int_0^{\infty} \frac{\sin v}{v} dv$$

Similarly,

$$I_2 = \frac{3}{\pi} \int_0^{\infty} \frac{\sin v}{v} dv = \frac{3}{\pi} \times \frac{\pi}{2} = \frac{3}{2}$$

$$\therefore I = I_1 + I_2 = \frac{3}{2} + \frac{3}{2} = 3$$



Scan for Video solution



14. (b)

$$\begin{aligned} L\{e^{i5t}\} &= L\{\cos 5t + i \sin 5t\} \\ &= L\{\cos 5t\} + i L\{\sin 5t\} \\ &= \frac{s}{s^2 + 5^2} + i \frac{5}{s^2 + 5^2} \\ \left( \because L\{\cos at\} = \frac{s}{s^2 + a^2}, L\{\sin at\} = \frac{a}{s^2 + a^2} \right) \\ &= \frac{s + i5}{s^2 + 25} \end{aligned}$$

$$\therefore L\{e^{i5t}\} = \frac{s + 5i}{s^2 + 25}$$



Scan for Video solution



15. (c)

From the given graph,

$$f(t) = \begin{cases} 2; & 0 < t < 1 \\ 0; & t > 1 \end{cases}$$

$$\begin{aligned} \therefore L\{f(t)\} &= \int_0^\infty e^{-st} \cdot f(t) dt \\ &= \int_0^1 e^{-st} \cdot 2 \cdot dt + \int_1^\infty e^{-st} \cdot (0) \cdot dt \\ \Rightarrow L\{f(t)\} &= 2 \cdot \left[ \frac{e^{-st}}{-s} \right]_0^1 = \frac{-2}{s} \{e^{-3} - 1\} = \frac{2}{s} (1 - e^{-s}) \\ \Rightarrow L\{f(t)\} &= \frac{2 - 2e^{-s}}{s} \end{aligned}$$



Scan for Video solution



16. (d)

$$L\{\cos \omega t\} = \frac{s}{s^2 + \omega^2}$$

By first shift property,

$$L\{e^{at} \cdot \cos \omega t\} = \frac{(s-a)}{(s-a)^2 + \omega^2}$$

Substitute  $a = -2$  &  $\omega = 4$

$$\Rightarrow L\{e^{-2t} \cdot \cos 4t\} = \frac{(s-(-2))}{(s-(-2))^2 + 4^2} = \frac{(s+2)}{(s+2)^2 + 16}$$

$$\Rightarrow L\{e^{-2t} \cdot \cos 4t\} = \frac{s+2}{(s+2)^2 + 16}$$



Scan for Video solution



17. (c)

$$\text{Given: } \frac{d^2 f}{dt^2} + f = 0; \quad f(0) = 0, \frac{df}{dt}(0) = 4$$

The auxiliary equation is  $m^2 + 1 = 0$

$$\Rightarrow m = \pm i$$

The solution is  $f(t) = c_1 \cos t + c_2 \sin t$

$$\text{Given } f(0) = 0 \Rightarrow 0 = c_1 \cos 0 + c_2 \sin 0 \Rightarrow c_1 = 0$$

$$\therefore f(t) = c_2 \sin t$$

$$\Rightarrow \frac{df(t)}{dt} = c_2 \cos t$$

$$\text{Given } \frac{df}{dt}(0) = 4 \Rightarrow 4 = c_2 \cdot (1) \Rightarrow c_2 = 4$$

$$\therefore f(t) = 4 \sin t$$

$$\Rightarrow L\{f(t)\} = 4 \cdot L\{\sin t\} = 4 \cdot \left( \frac{1}{s^2 + 1^2} \right)$$

$$= \frac{4}{s^2 + 1} \quad \left( \because L(\sin at) = \frac{a}{s^2 + a^2} \right)$$

$$\therefore L\{f(t)\} = \frac{4}{s^2 + 1}$$



Scan for Video solution



18. (d)

$$\text{Given: } F(s) = \frac{1}{s(s+1)}$$


$$\Rightarrow F(s) = \frac{1}{s} - \frac{1}{s+1}$$

Applying inverse Laplace on both sides

$$\Rightarrow L^{-1}\{F(s)\} = L^{-1}\left\{\frac{1}{s}\right\} - L^{-1}\left\{\frac{1}{s+1}\right\} = 1 - e^{-t}$$

$$\left( \because L\{t^n\} = \frac{n!}{s^{n+1}}; L\{e^{at}\} = \frac{1}{s-a} \right)$$

$$\Rightarrow f(t) = 1 - e^{-t}$$



19. (a)

Given:

$$\bar{F}(s) = \frac{1}{s^2(s+1)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s+1}$$

$$\Rightarrow \bar{F}(s) = \frac{As(s+1) + Bs + Cs^2}{s^2(s+1)} = \frac{1}{s^2(s+1)}$$

$$\Rightarrow A + C = 0; A + B = 0; B = 1$$

$$\Rightarrow A = -1; B = 1; C = 1$$

$$\Rightarrow \frac{1}{s^2(s+1)} = \frac{-1}{s} + \frac{1}{s^2} + \frac{1}{(s+1)}$$

Applying inverse Laplace on both sides


$$L^{-1}\left\{\frac{1}{s^2(s+1)}\right\} = L^{-1}\left\{\frac{-1}{s}\right\} + L^{-1}\left\{\frac{1}{s^2}\right\} + L^{-1}\left\{\frac{1}{(s+1)}\right\}$$

$$L^{-1}\left\{\frac{1}{s^2(s+1)}\right\} = -1 + t + e^{-t}$$

$$\left\{ \because L\{t^n\} = \frac{n!}{s^{n+1}} \right\}$$

$$\Rightarrow f(t) = -1 + t + e^{-t}$$

$$\left\{ \because L\{e^{at}\} = \frac{1}{s-a} \right\}$$



20. (c)

$$\text{Given: } \bar{F}(s) = \frac{1}{(s^2+s)} = \frac{1}{s(s+1)} = \frac{1}{s} - \frac{1}{s+1}$$

$$\Rightarrow \bar{F}(s) = \frac{1}{s} - \frac{1}{s+1}$$


$$\therefore L^{-1}\{\bar{F}(s)\} = L^{-1}\left\{\frac{1}{s}\right\} - L^{-1}\left\{\frac{1}{s+1}\right\}$$

$$= L^{-1}\left\{\frac{1}{s-0}\right\} - L^{-1}\left\{\frac{1}{s-(-1)}\right\}$$

$$= e^{0t} - e^{-1t} = 1 - e^{-t}$$

$$\left\{ \because L\{e^{at}\} = \frac{1}{s-a} \right\}$$

$$\Rightarrow L^{-1}\left\{\frac{1}{s^2+s}\right\} = 1 - e^{-t}$$



21. (b)

$$\cosh ax = \frac{e^{ax} + e^{-ax}}{2}$$


Taking Laplace transform, we get

$$= \frac{1}{2} L\{e^{ax}\} + L\{e^{-ax}\}$$

$$= \frac{1}{2} \left[ \frac{1}{s-a} + \frac{1}{s+a} \right]$$

$$\frac{1}{2} \left[ \frac{2s}{s^2 - a^2} \right]$$

$$L(\cosh ax) = \frac{s}{s^2 - a^2}$$



22. (a)

Given

$$\frac{d^2y}{dt^2} - y = 1; \quad y(0) = 0 \quad \& \quad \left. \frac{dy}{dt} \right|_{t=0} = 0$$

Applying Laplace transform on both sides of DE.

$$\Rightarrow (s^2 \cdot F(s) - s \cdot y(0) - y'(0)) - F(s) = \frac{1}{s}$$

$$\Rightarrow (s^2 - 1)F(s) = \frac{1}{s}$$

$$\Rightarrow F(s) = \frac{1}{s(s^2 - 1)}$$

$$\Rightarrow F(s) = \frac{1}{s(s-1)(s+1)}$$



Scan for Video solution



23. (d)

$$\text{Let } f(t) = \frac{\sin 2\pi t}{\pi t}$$

$$f(t) = f(-t) = \frac{\sin 2\pi t}{t}$$

$\therefore f(t)$  is even

$$I = 2 \int_{-\infty}^{\infty} \frac{\sin 2\pi t}{\pi t} dt = 4 \int_0^{\infty} \frac{\sin 2\pi t}{\pi t} dt = \frac{4}{\pi} \int_0^{\infty} \frac{\sin 2\pi t}{t} dt$$

$$\text{From Laplace transform } L\left[\frac{f(t)}{t}\right] = \int_s^{\infty} f(s) ds$$

$$\text{We know, } \int_0^{\infty} \frac{\sin at}{t} dt = \frac{\pi}{2}$$

$$\therefore \int_0^{\infty} \frac{\sin 2\pi t}{t} dt = \frac{\pi}{2}$$

$$\text{Hence, } I = \frac{4}{\pi} \cdot \frac{\pi}{2} = 2$$



Scan for Video solution



24. (b)

Using initial value theorem,

$$x(0^+) = \lim_{s \rightarrow \infty} s X(s)$$

$$= \lim_{s \rightarrow \infty} s \left( \frac{3s+5}{s^2 + 10s + 21} \right)$$

$$\lim_{s \rightarrow \infty} \frac{s}{1 + \frac{10}{s} + \frac{21}{s^2}} = 3$$



Scan for Video solution



25. (d)

$$\text{Given data, } \frac{d^2y(t)}{dt^2} + 2 \frac{dy(t)}{dt} + y(t) = \delta(t)$$

$$y(t)|_{t=0} = -2 \Rightarrow y(0) = -2$$

$$\frac{dy}{dt}|_{t=0} = 0 \Rightarrow y'(0) = 0$$

Differential equation can be written as:-

$$y'' + 2y' + y = \delta(t)$$

Taking Laplace transform on both sides

$$\left[ s^2 Y(s) - sy(0) - y'(0) \right] + 2[sY(s) - y(0)] + Y(s) = 1$$

$$[s^2 + 2s + 1]Y(s) + 2s + 4 = 1$$

$$Y(s) = \frac{-3 - 2s}{(s+1)^2}$$

$$Y(s) = -\frac{3}{(s+1)^2} - 2 \left[ \frac{(s+1)-1}{(s+1)^2} \right]$$

$$Y(s) = -\frac{3}{(s+1)^2} - \frac{2}{(s+1)} + \frac{2}{(s+1)^2} = -\frac{1}{(s+1)^2} - \frac{2}{s+1}$$

$$y(t) = L^{-1}Y(s) = L^{-1} \left[ -\frac{1}{(s+1)^2} - \frac{2}{(s+1)} \right]$$

$$y = -te^{-t} - 2e^{-t}$$

$$\frac{dy}{dt} = te^{-t} - e^{-t} + 2e^{-t}$$

$$= e^{-t}(t-1+2)$$

$$\frac{dy}{dt} = e^{-t}(t+1)$$

$$\text{At } t=0; \frac{dy}{dt} = 1$$



Scan for Video solution



# CHAPTER

9

# **NUMERICAL METHODS**

# Numerical Solutions of Linear Equations

- ## 1. [NAT] [GATE-ME-2016:2M]

Gauss-Seidel method is used to solve the following equations (as per the given order):

$$x_1 + 2x_2 + 3x_3 = 5$$

$$2x_1 + 3x_2 + x_3 = 1$$

$$3x_1 + 2x_2 + x_3 = 3$$

Assuming initial guess as  $x_1 = x_2 = x_3 = 0$ , the value of  $x_3$  after the first iteration is \_\_\_\_\_

- 2. [MCQ] [GATE- CE-2009:2M]**

In the solution of the following set of linear equations by Gauss elimination using partial pivoting  $5x + y + 2z = 34$ ;  $4y - 3z = 12$ ; and  $10x - 2y + z = -4$ ; the pivots for elimination of x and y are



# Numerical Solution of Algebraic & Transcendental Equations

3. [NAT] [GATE- CE-2022:1M]

Consider the following recursive iteration scheme for different values of variable P with the initial guess

$$x_1 = 1; \quad x_{n+1} = \frac{1}{2} \left( x_n + \frac{P}{x_n} \right), \quad n = 1, 2, 3, 4, 5$$

For  $P = 2$ ,  $x_5$  is obtained to be 1.414, rounded off to three decimal places. For  $P = 3$ ,  $x_5$  is obtained to be 1.732, rounded-off to three decimal places.

If  $P = 10$ , the numerically value of  $x_5$  is \_\_\_\_\_ (round off to three decimal place)

4. [NAT] [GATE-ME-2021:2M]

Find the positive real root of  $x^3 - x - 3 = 0$  using Newton-Raphson method. If the starting guess ( $x_0$ ) is 2, the numerical value of the root after two iterations ( $x_2$ ) is \_\_\_\_\_ (round off to two decimal places).

5. [NAT] [GATE- CE-2018:1M]

The quadratic equation  $2x^2 - 3x + 3 = 0$  is to be solved numerically starting with an initial guess as  $x_0 = 2$ . The new estimate of x after the first iteration using Newton-Raphson method is \_\_\_\_\_.

6. [NAT] [GATE-EE-2017: 2M]

Only one of the real roots  $f(x) = x^6 - x - 1$  lies in the interval  $1 \leq x \leq 2$  and bisection method is used to find its value. For achieving an accuracy of 0.001, the required minimum number of iterations is .

7. [NAT] [GATE-EE-2017 : 1M]

Let  $y^2 - 2y + 1 = x$  and  $\sqrt{x} + y = 5$ . The value of  $x + \sqrt{y}$  equals \_\_\_\_\_. (Given the answer up to three decimal)

8. [NAT] [GATE-EC-2017:2M]

Starting with  $x = 1$ , the solution of the equation  $x^3 + x = 1$ , after two iterations of Newton-Raphson's method (*up to two decimal places*) is \_\_\_\_.

9. [NAT] [GATE- CE-2016-:1M]

Newton-Raphson method is to be used to find root of equation  $3x - e^x + \sin x = 0$ . If the initial trial value for the roots is taken as 0.333, the next approximation for the root would be\_\_\_\_\_ (*answer up to three decimal places*).

- 10. [NAT] [GATE-ME-2016:1M]**  
Solve the equation  $x = 10 \cos(x)$  using the Newton-Raphson method. The initial guess is  $x = \pi/4$ . The value of the predicted root after the first iteration, up to second decimal, is \_\_\_\_\_.

**11. [MCQ] [GATE-ME-2016:1M]**  
The root of the function  $f(x) = x^3 + x - 1$  obtained after first iteration on application of Newton-Raphson scheme using an initial guess of  $x_0 = 1$  is  
 (a) 0.682      (b) 0.686  
 (c) 0.750      (d) 1.000

**12. [NAT] [GATE-ME-2015:2M]**  
Newton-Raphson method is used to find the roots of the equation,  $x^3 + 2x^2 + 3x - 1 = 0$ . If the initial guess is  $x_0 = 1$ , then the value of  $x$  after 2<sup>nd</sup> iteration is \_\_\_\_\_.

**13. [NAT] [GATE-EC-2015:2M]**  
The Newton-Raphson method is used to solve the equation  $f(x) = x^3 - 5x^2 + 6x - 8 = 0$ . Taking the initial guess as  $x = 5$ , the solution obtained at the end of the first iteration is \_\_\_\_\_.

**14. [NAT] [GATE-CE-2015:2M]**  
The quadratic equation  $x^2 - 4x + 4 = 0$  is to be solved numerically, starting with the initial guess  $x_0 = 3$ . The Newton-Raphson method is applied once to get a new estimated and then the Secant method is applied once using the initial guess and this new estimate. The estimated value of the root after the application of the Secant method is \_\_\_\_\_.

**15. [NAT] [GATE-CE-2015:1M]**  
In Newton-Raphson iterative method, the initial guess value ( $x_{ini}$ ) is considered as zero while finding the roots of the equation:  $f(x) = -2 + 6x - 4x^2 + 0.5x^3$ .  
The correction,  $\Delta x$ , to be added to  $x_{ini}$  in the first iteration is \_\_\_\_\_.

**16. [NAT] [GATE-ME-2014:2M]**  
The real root of the equation  $5x - 2\cos x - 1 = 0$  (up to two decimal accuracy) is \_\_\_\_\_.

**17. [NAT] [GATE-EE-2014:2M]**  
The function  $f(x) = e^x - 1$  is to be solved using Newton-Raphson method. If the initial value of  $x_0$  is taken as 1.0, then the absolute error observed at 2<sup>nd</sup> iteration is \_\_\_\_\_.

**18. [MCQ] [GATE-EE-2013:2M]**  
When the Newton-Raphson method is applied to solve the equation  $f(x) = x^3 + 2x - 1 = 0$ , the solution at the end of the first iteration with the initial guess value as  $x_0 = 1.2$  is  
 (a) -0.82      (b) 0.49  
 (c) 0.705      (d) 1.69

**19. [MCQ] [GATE-EE-2011:1M]**  
Solution of the variables  $x_1$  and  $x_2$  for the following equations is to be obtained by employing the Newton-Raphson iterative method.  
 Equation(i)  $10x_2 \sin x_1 - 0.8 = 0$   
 Equation(ii)  $10x_2^2 - 10x_2 \cos x_1 - 0.6 = 0$   
 Assuming the initial values  $x_1 = 0.0$  and  $x_2 = 1.0$ , the Jacobian matrix is  
 (a)  $\begin{bmatrix} 10 & -0.8 \\ 0 & -0.6 \end{bmatrix}$       (b)  $\begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$   
 (c)  $\begin{bmatrix} 0 & -0.8 \\ 10 & -0.6 \end{bmatrix}$       (d)  $\begin{bmatrix} 10 & 0 \\ 10 & -10 \end{bmatrix}$

**20. [MCQ] [GATE-EC-2011:2M]**  
A numerical solution of the equation  $f(x) = x + \sqrt{x} - 3 = 0$  can be obtained using Newton Raphson method. If the starting value is  $x = 2$  for the iteration, the value of  $x$  that is to be used in the next step is  
 (a) 0.306      (b) 0.739  
 (c) 1.694      (d) 2.306

**21. [MCQ] [GATE- CE-2011:1M]**  
The square root of a number  $N$  is to be obtained by applying the Newton Raphson iterations to the equation  $x^2 - N = 0$ . If  $i$  denotes the iteration index, the correct iterative scheme will be  
 (a)  $x_{i+1} = \frac{1}{2} \left( x_i + \frac{N}{x_i} \right)$       (b)  $x_{i+1} = \frac{1}{2} \left( x_i^2 + \frac{N}{x_i^2} \right)$   
 (c)  $x_{i+1} = \frac{1}{2} \left( x_i + \frac{N^2}{x_i} \right)$       (d)  $x_{i+1} = \frac{1}{2} \left( x_i - \frac{N}{x_i} \right)$

**22. [MCQ]****[GATE-EE-2009:2M]**

Let  $x^2 - 117 = 0$ . The iterative steps for the solution using Newton-Raphson's Method is given by

- (a)  $x_{k+1} = \frac{1}{2} \left( x_k + \frac{117}{x_k} \right)$
- (b)  $x_{k+1} = x_k - \frac{117}{x_k}$
- (c)  $x_{k+1} = x_k - \frac{x_k}{117}$
- (d)  $x_{k+1} = x_k - \frac{1}{2} \left( x_k + \frac{117}{x_k} \right)$

**23. [MCQ]****[GATE-EE-2008:2M]**

Equation  $e^x - 1 = 0$  is required to be solved using Newton's method with an initial guess  $x_0 = -1$ . Then, after one step of Newton's method, estimate  $x_1$  the solution will be given by

- (a) 0.71828
- (b) 0.36784
- (c) 0.20587
- (d) 0.00000


**24. [MCQ]****[GATE-EC-2008:2M]**

The recursion relation to solve  $x = e^{-x}$  using Newton-Raphson method is

- (a)  $x_{n+1} = e^{-x_n}$
- (b)  $x_{n+1} = x_n - e^{-x_n}$
- (c)  $x_{n+1} = (1 + x_n) \frac{e^{-x_n}}{1 + e^{-x_n}}$
- (d)  $x_{n+1} = \frac{x_n^2 - e^{-x_n}(1 + x_n) - 1}{x_n - e^{-x_n}}$

**Numerical Integration****25. [MCQ]****[GATE- CE-2023:1M]**

A function  $f(x)$ , that is smooth and convex-shaped between interval  $(x_l, x_u)$  is shown in the figure. This function is observed at odd number of regularly spaced points. If the area under the function is computed numerically, then.



- (a) the numerical value of the area obtained using the trapezoidal rule will be less than the actual
- (b) the numerical value of the area obtained using the trapezoidal rule will be more than the actual
- (c) the numerical value of the area obtained using the trapezoidal rule will be exactly equal to the actual
- (d) with the given details, the numerical value of area cannot be obtained using trapezoidal rule

**26. [MCQ]****[GATE-ME-2022:1M]**

Consider the definite integral  $\int_1^2 (4x^2 + 2x + 6) dx$ .

Let  $I_e$  be the exact value of the integral. If the same integral is estimated using Simpson's rule with 10 equal subintervals, the value is  $I_s$ . The percentage error is defined as  $e = 100 \times (I_e - I_s)/I_e$ . The value of  $e$  is

- (a) 2.5
- (b) 3.5
- (c) 1.2
- (d) 0

**27. [NAT]****[GATE-CE-2021:2M]**

The values of abscissa (x) and ordinate (y) of a curve are as follows:

| X   | Y     |
|-----|-------|
| 2.0 | 5.00  |
| 2.5 | 7.25  |
| 3.0 | 10.00 |
| 3.5 | 13.25 |
| 4.0 | 17.00 |

By Simpson's 1/3<sup>rd</sup> rule, the area under the curve (round off to two decimal places) is \_\_\_\_\_.

**28. [MCQ]****[GATE-CE-2021:2M]**

The value of  $\int_0^1 e^x dx$  using the trapezoidal rule with four equal subintervals is

- (a) 1.718
- (b) 2.192
- (c) 1.727
- (d) 2.718

**29. [NAT]****[GATE- CE-2021:2M]**

Numerically integrate,  $f(x) = 10x - 20x^2$  from lower limit  $a = 0$  to upper limit  $b = 0.5$ . Using Trapezoidal rule with five equal subdivisions. The value (in units, round off to two decimal places) obtained is \_\_\_\_\_.



**42. [NAT]****[GATE-ME-2014:1M]**

Using a unit step size, the value of integral  $\int_1^2 x \ln x \, dx$  by trapezoidal rule is \_\_\_\_\_

**43. [NAT]****[GATE-ME-2014:1M]**

The definite integral  $\int_1^3 \frac{1}{x} \, dx$  is evaluated using Trapezoidal rule with a step size of 1. The correct answer is \_\_\_\_\_

**44. [NAT]****[GATE-ME-2014:2M]**

The value of  $\int_{2.5}^4 \ln(x) \, dx$  calculated using the Trapezoidal rule with five subintervals is \_\_\_\_\_

**45. [NAT]****[GATE-ME-2014:2M]**

Using the trapezoidal rule, and dividing the interval of integration into three equal subintervals, the definite integral  $\int_{-1}^{+1} |x| \, dx$  is \_\_\_\_\_

**46. [NAT]****[GATE-CE-2013:2M]**

Find the magnitude of error (Correct to two decimal places) in the estimation of following integral using Simpson 1/3 rule. Take the step length as 1

$$\int_0^4 (x^4 + 10) \, dx$$

**47. [MCQ]****[GATE-ME-2013:1M]**

Match the CORRECT pairs

| Numerical Integration Scheme | Order of Fitting Polynomial |
|------------------------------|-----------------------------|
| P. Simpson's 3/8 rule        | 1. First                    |
| Q. Trapezoidal rule          | 2. Second                   |
| R. Simpson's 1/3 rule        | 3. Third                    |

- (a) P-2, Q-1, R-3  
 (c) P-1, Q-2, R-3

- (b) P-3, Q-2, R-1  
 (d) P-3, Q-1, R-2

**48. [MCQ]****[GATE-CE-2011:2M]**

The estimated of  $\int_{0.5}^{1.5} \frac{dx}{x}$  Obtained using Simpson's rule with three-point function evaluation exceeds the exact value by

- (a) 0.235  
 (c) 0.024

- (b) 0.068  
 (d) 0.012

**49. [MCQ]****[GATE-ME-2011:2M]**

The integral  $\int_1^3 \frac{1}{x} \, dx$ , when evaluated by using Simpson's 1/3 rule on two equal subintervals each of length 1, equals

- (a) 1.000  
 (c) 1.111

- (b) 1.098  
 (d) 1.120

**50. [MCQ]****[GATE-ME-2010:2M]**

Torque exerted on a flywheel over a cycle is listed in the table. Flywheel energy (in J per unit cycle) using Simpson's rule is

|                |   |      |      |     |     |      |     |
|----------------|---|------|------|-----|-----|------|-----|
| Angle (degree) | 0 | 60   | 12   | 180 | 240 | 300  | 360 |
| Torque (N m)   | 0 | 1066 | -323 | 0   | 323 | -355 | 0   |

- (a) 542  
 (c) 1444

- (b) 993  
 (d) 1986

**51. [MCQ]****[GATE-CE-2010:2M]**

The table below gives values of a function F(x) obtained for values of x at intervals of 0.25.

The value of the integral of the function between the limits 0 to 1 using Simpson's rule is

|      |   |        |     |      |      |
|------|---|--------|-----|------|------|
| x    | 0 | 0.25   | 0.5 | 0.75 | 1.0  |
| F(x) | 1 | 0.9412 | 0.8 | 0.64 | 0.50 |

- (a) 0.7854  
 (c) 3.1416

- (b) 2.3562  
 (d) 7.5000

### Numerical Solution of Differential Equations

**52. [MCQ]****[GATE-CE-2023:2M]**

The differential equation,  $\frac{du}{dt} + 2tu^2 = 1$ , is solved by employing a backward difference scheme within the finite difference framework. The value of  $u$  at the  $(n-1)^{th}$  time-step, for some  $n$ , is 1.75. The corresponding time ( $t$ ) is 3.14 s. Each time step is 0.01 s long. Then, the value of  $(u_n - u_{n-1})$  is \_\_\_\_\_ (round off to three decimal places).

**53. [NAT]****[GATE-ME-2023:2M]**

The initial value problem

$$\frac{dy}{dt} + 2y = 0, \quad y(0) = 1$$

is solved numerically using the forward Euler's method with a constant and positive time step of  $\Delta t$ . Let  $y_n$  represent the numerical solution obtained after  $n$  steps. The condition  $|y_{n+1}| - |y_n|$  is satisfied if and only if  $\Delta t$  does not exceed \_\_\_\_\_. (Answer in integer)

**54. [NAT]****[GATE-CE-2022:2M]**

Consider the differential equation  $\frac{dy}{dx} = 4(x+2) - y$

For the initial condition  $y = 3$  at  $x = 1$ , the value of  $y$  at  $x = 1.4$  obtained using Euler's method with a step-size of 0.2 is \_\_\_\_\_. (round off to one decimal place)

**55. [MCQ]****[GATE-ME-2022:1M]**

A polynomial  $\phi(s) = a_n s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0$  of degree  $n > 3$  with constant real coefficients  $a_n, a_{n-1}, \dots, a_0$  has triple roots at  $s = -\sigma$ . Which one of the following conditions must be satisfied?

(a)  $\phi(s) = 0$  at all the three values of  $s$  satisfying

$$s^3 + \sigma^3 = 0$$

(b)  $\phi(s) = 0, \frac{d\phi(s)}{ds} = 0$  and  $\frac{d^2\phi(s)}{ds^2} = 0$  at  $s = -\sigma$

(c)  $\phi(s) = 0, \frac{d^2\phi(s)}{ds^2} = 0$ , and  $\frac{d^4\phi(s)}{ds^4} = 0$ , at  $s = -\sigma$

(d)  $\phi(s) = 0$  and  $\frac{d^3\phi(s)}{ds^3} = 0$ , at  $s = -\sigma$

**56. [MCQ]****[GATE-ME-2021:1M]**

The ordinary differential equation  $\frac{dy}{dt} = -\pi y$  subjected to an initial condition  $y(0) = 1$  is solved numerically using the following scheme:

$$\frac{y(t_{n+1}) - y(t_n)}{h} = -\pi y(t_n)$$

where  $h$  is the time step,  $t_n = nh$ , and  $\pi = 0, 1, 2, \dots$ . This numerical scheme is stable for all values of  $h$  in the interval \_\_\_\_\_.

(a)  $0 < h < \frac{2}{\pi}$

(b)  $0 < h < 1$

(c)  $0 < h < \frac{\pi}{2}$

(d) for all  $h > 0$

**57. [MCQ]****[GATE- CE-2020:2M]**

A continuous function  $f(x)$  is defined. If the third derivative at  $x_i$  is to be computed by using the fourth order central finite-divided-difference scheme (with step length =  $h$ ), the correct formula is

(a)  $f'''(x_i) = \frac{-f(x_{i+3}) + 8f(x_{i+2}) - 13f(x_{i+1}) + 13f(x_{i-1}) - 8f(x_{i-2}) + f(x_{i-3})}{8h^3}$

(b)  $f'''(x_i) = \frac{f(x_{i+3}) - 8f(x_{i+2}) - 13f(x_{i+1}) + 13f(x_{i-1}) + 8f(x_{i-2}) + f(x_{i-3})}{8h^3}$

(c)  $f'''(x_i) = \frac{-f(x_{i+3}) - 8f(x_{i+2}) - 13f(x_{i+1}) + 13f(x_{i-1}) + 8f(x_{i-2}) - f(x_{i-3})}{8h^3}$

(d)  $f'''(x_i) = \frac{f(x_{i+3}) - 8f(x_{i+2}) + 13f(x_{i+1}) + 13f(x_{i-1}) - 8f(x_{i-2}) - f(x_{i-3})}{8h^3}$

**58. [NAT]****[GATE-ME-2018:2M]**

An explicit forward Euler method is used to numerically integrate the differential equation

$$\frac{dy}{dt} = y$$

using a time step of 0.1. With the initial condition  $y(0) = 1$ , the value of  $y(1)$  computed by this method is \_\_\_\_\_. (Correct to two decimal places).

**59. [NAT]****[GATE-CE-2017:2M]**

Consider the equation  $\frac{du}{dt} = 3t^2 + 1$  with  $u = 0$  at  $t = 0$ .

This is numerically solved by using the forward Euler method with a step size.  $\Delta t = 2$ . The absolute error in the solution in the end of the first-time step is

**60. [NAT]****[GATE-EC-2016:1M]**

Consider the first order initial value problem  $y' = y + 2x - x^2$ ,  $y(0) = 1$ ,  $(0, \leq x < \infty)$  with exact solutions  $y(x) = x^2 + e^x$ . For  $x = 0.1$ , the percentage difference between the exact solution and the solution obtained using a single iteration of the second-order Runge Kutta method with step-size  $h = 0.1$  is \_\_\_\_\_.

**61. [NAT]****[GATE-EC-2016:2M]**

The ordinary differential equation

$$\frac{dx}{dt} = -3x + 2; \text{ with } x(0) = 1$$

is to be solved using the forward Euler method. The largest time step that can be used to solve the equation without making the numerical solution unstable is \_\_\_\_\_.

**62. [MCQ]****[GATE-ME-2014:2M]**

Consider an ordinary differential equation  $\frac{dx}{dt} = 4t + 4$ .

If  $x = x_0$  at  $t = 0$ , the increment in  $x$  calculated using Runge-Kutta fourth order multi-step method with a step size of  $\Delta t = 0.2$  is

(a) 0.22 (b) 0.44

(c) 0.66 (d) 0.88

**63. MCQ**

[GATE-EC-2014:1M]

Match the application to appropriate numerical method.

**Application**

- P1: Numerical integration
- P2: Solution to a transcendental equation
- P3: Solution to a system of linear equation
- P4: Solution to a differential equation

**Numerical Method**

M1: Newton-Raphson Method

M2: Runge-Kutta Method

M3: Simpson's 1/3-rule

M4: Gauss Elimination Method

- (a) P1-M3, P2-M2, P3-M4, P4-M1
- (b) P1-M3, P2-M1, P3-M4, P4-M2
- (c) P1-M4, P2-M1, P3-M3, P4-M2
- (d) P1-M2, P2-M1, P3-M3, P4-M4

**64. [MCQ]**

[GATE-CE-2012:2M]

The error in  $\frac{d}{dx} f(x) \Big|_{x=x_0}$  for a continuous function estimated with  $h = 0.03$  using the central difference formula  $\frac{d}{dx} f(x) \Big|_{x=x_0} \approx \frac{f(x_0 + h) - f(x_0 - h)}{2h}$ , is  $2 \times 10^{-3}$ . The value of  $x_0$  and  $f(x_0)$  are 19.78 and 500.01, respectively. The corresponding error in the central difference estimate for  $h = 0.02$  is approximately

- (a)  $1.3 \times 10^{-4}$
- (b)  $4.5 \times 10^{-4}$
- (c)  $3.0 \times 10^{-4}$
- (d)  $9.0 \times 10^{-4}$

**65. [MCQ]**

[GATE-EC-2010: 2M]

Consider a differential equation  $\frac{dy(x)}{dx} - y(x) = x$  with the initial condition  $y(0) = 0$ . Using Euler's first order method with a step size of 0.1 the value of  $y(0.3)$  is

- (a) 0.01
- (b) 0.031
- (c) 0.0631
- (d) 0.1

**66. [MCQ]**

[GATE-EE-2008:2M]

A differential equation  $\frac{dx}{dt} = e^{-2t} u(t)$  has to be solved using trapezoidal rule of integration with a step size  $h = 0.01$ s. Function  $u(t)$  indicates a unit step function. If  $x(0^-) = 0$ , then value of  $x$  at  $t = 0.01$ s will be given by

- (a) 0.00099
- (b) 0.00495
- (c) 0.0099
- (d) 0.0198

**Interpolation****67. [MCQ]**

[GATE-CE-2019:1M]

The value of the function  $f(x)$  is given at  $n$  distinct values of  $x$  and its value is to be interpolated at the point  $x^*$ , using all the  $n$  points. The estimate is obtained first by the Lagrange polynomial denoted by  $I_L$  and then by the Newton polynomial, denoted by  $I_N$ . Which one of the following statements is correct?

- (a)  $I_L$  is always greater than  $I_N$
- (b) No definite relation exists between  $I_L$  and  $I_N$
- (c)  $I_L$  and  $I_N$  are always equal
- (d)  $I_L$  is always less than  $I_N$



 ANSWER KEY

- |                             |                           |                             |                               |
|-----------------------------|---------------------------|-----------------------------|-------------------------------|
| <b>1.</b> (-6 to -6)        | <b>2.</b> (a)             | <b>3.</b> (3.100 to 3.200)  | <b>4.</b> (1.66 to 1.68)      |
| <b>5.</b> (1 to 1)          | <b>6.</b> (10 to 10)      | <b>7.</b> (5.7 to 5.8)      | <b>8.</b> (0.65 to 0.72)      |
| <b>9.</b> (0.355 to 0.365)  | <b>10.</b> (1.53 to 1.59) | <b>11.</b> (c)              | <b>12.</b> (0.29 to 0.31)     |
| <b>13.</b> (4.25 to 4.35)   | <b>14.</b> (2.32 to 2.34) | <b>15.</b> (0.3 to 0.4)     | <b>16.</b> (0.53 to 0.56)     |
| <b>17.</b> (0.05 to 0.07)   | <b>18.</b> (c)            | <b>19.</b> (b)              | <b>20.</b> (c)                |
| <b>21.</b> (a)              | <b>22.</b> (a)            | <b>23.</b> (a)              | <b>24.</b> (c)                |
| <b>25.</b> (a)              | <b>26.</b> (d)            | <b>27.</b> (20.00 to 21.00) | <b>28.</b> (c)                |
| <b>29.</b> (0.38 to 0.42)   | <b>30.</b> (a)            | <b>31.</b> (d)              | <b>32.</b> (b)                |
| <b>33.</b> (5.1 to 5.5)     | <b>34.</b> (63 to 63)     | <b>35.</b> (a)              | <b>36.</b> (a)                |
| <b>37.</b> (0.175 to 0.195) | <b>38.</b> (2 to 2)       | <b>39.</b> (a)              | <b>40.</b> (1.36 to 1.37)     |
| <b>41.</b> (21.8 to 22.2)   | <b>42.</b> (0.68 to 0.70) | <b>43.</b> (1.1 to 1.2)     | <b>44.</b> (1.74 to 1.76)     |
| <b>45.</b> (1.10 to 1.12)   | <b>46.</b> (0.50 to 0.53) | <b>47.</b> (d)              | <b>48.</b> (d)                |
| <b>49.</b> (c)              | <b>50.</b> (b)            | <b>51.</b> (a)              | <b>52.</b> (-0.152 to -0.149) |
| <b>53.</b> (0.999 to 1.001) | <b>54.</b> (6.3 to 6.5)   | <b>55.</b> (b)              | <b>56.</b> (a)                |
| <b>57.</b> (a)              | <b>58.</b> (2.55 to 2.65) | <b>59.</b> (7.95 to 8.05)   | <b>60.</b> (0.060 to 0.063)   |
| <b>61.</b> (0.6 to 0.7)     | <b>62.</b> (d)            | <b>63.</b> (b)              | <b>64.</b> (d)                |
| <b>65.</b> (b)              | <b>66.</b> (c)            | <b>67.</b> (c)              |                               |


**SOLUTIONS**
**1. (-6 to -6)**

Given System is

$$\begin{aligned}x_1 + 2x_2 + 3x_3 &= 5 \\2x_1 + 3x_2 + x_3 &= 1 \\3x_1 + 2x_2 + x_3 &= 3\end{aligned}$$

Taking equation in same order

$$x_1^{(k+1)} = 5 - 2x_2^{(k)} - 3x_3^{(k)}$$

$$x_2^{(k+1)} = \frac{1}{3}\{1 - 2x_1^{(k+1)} - x_3^{(k)}\}$$

$$x_3^{(k+1)} = 3 - 3x_1^{(k+1)} - 2x_2^{(k+1)}$$

since  $x_1^{(0)} = x_2^{(0)} = x_3^{(0)} = 0$

$$\Rightarrow x_1^{(1)} = 5 - 2(0) - 3(0) = 5$$

$$x_2^{(1)} = \frac{1}{3}\{1 - 2(5) - 0\} = -3$$

$$x_3^{(1)} = 3 - 3(5) - 2(-3)$$

$$\Rightarrow x_3^{(1)} = 3 - 15 + 6 = -6$$

$\therefore x_3$  after 1<sup>st</sup> iteration = -6



Scan for Video solution

**2. (a)**The equation are,  $5x + y + 2z = 34$  $4y - 3z = 12$  and  $10x - 2y + z = -4$ 

The augmented matrix for gauss elimination is,

$$(A / B) = \left[ \begin{array}{ccc|c} 5 & 1 & 2 & 34 \\ 0 & 4 & -3 & 12 \\ 10 & -2 & 1 & -4 \end{array} \right]$$

Since in the first column maximum element in absolute value is 10 we need to exchange row 1 with row 3.

$$\left[ \begin{array}{ccc|c} 5 & 1 & 2 & 34 \\ 0 & 4 & -3 & 12 \\ 10 & -2 & 1 & -4 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_3} \left[ \begin{array}{ccc|c} 10 & -2 & 1 & -4 \\ 0 & 4 & -3 & 12 \\ 5 & 1 & 2 & 34 \end{array} \right]$$

So, the pivot for eliminating x is  $a_{11} = 10$ .

- Now to eliminate y, we need to eliminate in second column below the diagonal.

Since  $a_{22} = 4$  is already larger in absolute value when compared to  $a_{32} = 1$

- The pivots element for eliminating y is  $a_{22} = 4$  itself
- The pivots for eliminating x and y are 10 and 4 respectively.



Scan for Video solution

**3. (3.100 to 3.200)**We know,  $x_{n+1} = x_n = x$ ; when  $n \rightarrow \infty$ 

Recursive iteration formula

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{P}{x_n} \right)$$

$$x = \frac{1}{2} \left( x + \frac{P}{x} \right) \Rightarrow \frac{x}{2} = \frac{P}{2x}$$

$$x^2 = P \Rightarrow x = \sqrt{P}$$

$$\therefore \text{If } P = 10, \text{ then } x_5 = \sqrt{P} = \sqrt{10} = 3.162$$



Scan for Video solution

**4. (1.66 to 1.68)****Given**

$$x^3 - x - 3 = 0; \text{ and } x_0 = 2$$

For  $f(x) = 0$ , Newton-Raphson method is given by

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$x_{k+1} = x_k - \frac{(x_k^3 - x_k - 3)}{(3x_k^2 - 1)} = \frac{3x_k^3 - x_k - x_k^3 + x_k + 3}{3x_k^2 - 1}$$

$$\Rightarrow x_{k+1} = \frac{2x_k^3 + 3}{3x_k^2 - 1}$$

$$\text{For } k=0, x_1 = \frac{2x_0^3 + 3}{3x_0^2 - 1} = \frac{2(2)^3 + 3}{3(2)^2 - 1} = \frac{19}{11} = 1.7272$$

$$\text{For } k=1, x_2 = \frac{2x_1^3 + 3}{3x_1^2 - 1} = \frac{2(1.7272)^3 + 3}{3(1.7272)^2 - 1} = 1.6736$$

$$\therefore x_2 = 1.6736$$



Scan for Video solution



### 5. (1 to 1)

$$\text{Given: } f(x) = 2x^2 - 3x + 3, x_0 = 2$$

$$f'(x) = 4x - 3$$

By Newton-Raphson method,

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$x_1 = 2 - \frac{f(2)}{f'(2)} = 2 - \frac{(2 \times 2^2 - 3 \times 2 + 3)}{4 \times 2 - 3} = 2 - \frac{5}{5} = 1$$



Scan for Video solution



### 6. (10 to 10)

The minimum number of iterations in bisection method for error  $\epsilon$  is given by  $\frac{|b-a|}{2^n} < \epsilon$

a : Lower limit of interval

b: Upper limit of interval

n : No of iterations

$\epsilon$  : Error in approximation

$$\frac{|2-1|}{2^n} < 0.001 \Rightarrow 2^n > 1000$$

$$\therefore n = 10$$



Scan for Video solution



### 7. (5.7 to 5.8)

$$\text{Given: } y^2 - 2y + 1 = x \text{ and } \sqrt{x} + y = 5$$

$$x + \sqrt{y} = ?$$

$$\sqrt{x} + y = 5$$

$$\Rightarrow \sqrt{x} = 5 - y$$

$$\Rightarrow x = (5 - y)^2 \quad \dots(i)$$

And,

$$y^2 - 2y + 1 = x \quad \dots(ii)$$

From equation (i), put expression of x in equation (ii),

$$y^2 - 2y + 1 = (5 - y)^2$$

$$\Rightarrow y^2 - 2y + 1 = 25 + y^2 - 10y$$

$$\Rightarrow 8y = 24$$

$$\Rightarrow y = 3$$

From equation (i)

$$x = (5 - y)^2$$

$$\Rightarrow x = (5 - 3)^2 = 4$$

Therefore,

$$x + \sqrt{y} = 4 + \sqrt{3}$$

$$\Rightarrow x + \sqrt{y} = 5.732$$



Scan for Video solution



### 8. (0.65 to 0.72)

Given: Equation is,

$$x^3 + x = 1$$

$$\Rightarrow x^3 + x - 1 = 0$$

and initial guess,

$$x_0 = 1$$

The iterative scheme for  $f(x) = 0$  using Newton-Raphson iterative scheme is,

$$\Rightarrow x_{k+1} = x_k - \frac{(x_k^3 + x_k - 1)}{(3x_k^2 + 1)}$$

$$\Rightarrow x_{k+1} = \frac{2x_k^3 + 1}{3x_k^2 + 1}$$

For  $k = 1$ :

$$\Rightarrow x_1 = \frac{2x_0^3 + 1}{3x_0^2 + 1} = \frac{2(1) + 1}{3(1) + 1} = \frac{3}{4} = 0.75$$

For  $k = 2$ :

$$\Rightarrow x_2 = \frac{2x_1^3 + 1}{3x_1^2 + 1} = \frac{2(0.75)^3 + 1}{3(0.75)^2 + 1}$$

$$= \frac{1.84375}{2.6875} = 0.686$$

∴ The value after 2<sup>nd</sup> iteration is 0.69.



Scan for Video solution



### 9. (0.355 to 0.365)

According to newton Raphson method,

$$x_{N+1} = x_N - \frac{f(x_N)}{f'(x_N)}$$

$$f(x) = 3x - e^x + \sin x$$

$$f'(x) = 3 - e^x + \cos x$$

$$x_0 = 0.333$$

$$\Rightarrow x_1 = x_0 - \frac{f(0.333)}{f'(0.333)}$$

$$= 0.333 - \frac{(3 \times 0.33 - e^{0.333} + \sin 0.333)}{(3 - e^{0.333} + \cos 0.333)}$$

$$\Rightarrow x_1 = 0.36$$



Scan for Video solution



### 10. (1.53 to 1.59)

$$\text{Given: } x = 10 \cos x \quad \text{and} \quad x_0 = \frac{\pi}{4}$$

$$\Rightarrow x - 10 \cos x = 0$$

For  $f(x) = 0$ , Newton – Raphson scheme is given by

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$\Rightarrow x_{k+1} = x_k - \frac{(x_k - 10 \cos x_k)}{(1 + 10 \cdot \sin x_k)}$$

$$\Rightarrow x_{k+1} = \frac{10 x_k \cdot \sin x_k + 10 \cos x_k}{(1 + 10 \cdot \sin x_k)}$$

For  $k = 0$ ,

$$x_1 = \frac{10 x_0 \cdot \sin x_0 + 10 \cos x_0}{(1 + 10 \cdot \sin x_0)}$$

$$= \frac{10 \left( \frac{\pi}{4} \cdot \frac{1}{\sqrt{2}} \right) + 10 \left( \frac{1}{\sqrt{2}} \right)}{\left( 1 + 10 \left( \frac{1}{\sqrt{2}} \right) \right)} = \frac{10 \left( \frac{\pi}{4} + 1 \right)}{(10 + \sqrt{2})} = 1.564$$

∴ First iteration value =  $x_1 = 1.564$



Scan for Video solution



### 11. (c)

**Given:**  $f(x) = x^3 + x - 1$  and  $x_0 = 1$

By Newton-Raphson scheme,

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$\Rightarrow x_{k+1} = x_k - \frac{(x_k^3 + x_k - 1)}{(3x_k^2 + 1)}$$

$$\Rightarrow x_{k+1} = \frac{2x_k^3 + 1}{3x_k^2 + 1}$$

For  $k = 0$

$$\Rightarrow x_1 = \frac{2x_0^3 + 1}{3x_0^2 + 1} = \frac{2(1)^3 + 1}{3(1)^2 + 1} = \frac{3}{4} = 0.75$$

∴ First Iteration value = 0.750



Scan for Video solution



### 12. (0.29 to 0.31)

**Given:**  $x^3 + 2x^2 + 3x - 1 = 0$  and  $x_0 = 1$

For  $f(x) = 0$ , Newton-Raphson method is given by

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$\Rightarrow x_{k+1} = x_k - \frac{(x_k^3 + 2x_k^2 + 3x_k - 1)}{3x_k^2 + 4 \cdot x_k + 3}$$

$$\Rightarrow x_{k+1} = \frac{2x_k^3 + 2x_k^2 + 1}{3x_k^2 + 4 \cdot x_k + 3}$$

$$\text{For } k = 0, x_1 = \frac{2x_0^3 + 2x_0^2 + 1}{3x_0^2 + 4 \cdot x_0 + 3} = \frac{5}{10} = 0.5$$

For  $k = 1$ ;

$$x_2 = \frac{2x_1^3 + 2x_1^2 + 1}{3x_1^2 + 4x_1 + 3} = \frac{2(0.5)^3 + 2(0.5)^2 + 1}{3(0.5)^2 + 4(0.5) + 3} = 0.3043$$

$\therefore$  Value after 2<sup>nd</sup> iteration =  $x_2 = 0.304$



Scan for Video solution



### 13. (4.25 to 4.35)

Given equation is  $f(x) = x^3 - 5x^2 + 6x - 8 = 0$

Initial guess is  $x_0 = 5$

For any function  $f(x) = 0$ , newton-Raphson iterative scheme is given by

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$\Rightarrow x_{k+1} = x_k - \frac{(x_k^3 - 5x_k^2 + 6x_k - 8)}{(3x_k^2 - 10x_k + 6)}$$

$$\Rightarrow x_{k+1} = \frac{2x_k^3 - 5x_k^2 + 8}{3x_k^2 - 10x_k + 6}$$

$$\text{For } k = 0; x_1 = \frac{2x_0^3 - 5x_0^2 + 8}{3x_0^2 - 10x_0 + 6}$$

$$= \frac{2(125) - 5(25) + 8}{3(25) - 10(5) + 6} = \frac{133}{31} = 4.29$$

$\therefore$  The value at the end of 1<sup>st</sup> iteration is 4.29.



Scan for Video solution



### 14. (2.32 to 2.34)

$$f(x) = x^2 - 4x + 4$$

$$f'(x) = 2x - 4$$

$$x_0 = 3$$

$$f(3) = 1, f'(3) = 2$$

By Newton Raphson method

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = 3 - \frac{1}{2} = \frac{5}{2} = 2.5$$

$$(\because x_0 = 3, f(x_0) = 1, f'(x_0) = 2)$$

$$x_1 = \frac{5}{2} \quad f(x_1) = \left(\frac{5}{2}\right)^2 - 4\left(\frac{5}{2}\right) + 4 = \frac{1}{4}$$

By secant method.

$$x_2 = \frac{x_0 f(x_1) - x_1 f(x_0)}{f(x_1) - f(x_0)}$$

$$x_2 = \frac{3 \times \frac{1}{4} - \frac{5}{2} \times 1}{\frac{1}{4} - 1} = \frac{7}{3} = 2.33$$



Scan for Video solution



$\therefore$  Scan for Video solution



### 15. (0.3 to 0.4)

Given:  $f(x) = -2 + 6x - 4x^2 + 0.5x^3$ ,  $x_{\text{ini}} = 0$

$$f'(x) = 6 - 8x + 1.5x^2$$

By Newton Raphson method,

$$x_1 = x_{\text{ini}} - \frac{f(x_{\text{ini}})}{f'(x_{\text{ini}})}$$

$$x_1 = 0 - \frac{(-2)}{6} \quad [\because f(0) = -2, f'(0) = 6]$$

$$x_1 = \frac{1}{3}$$

$$\Delta x = x_1 - x_{\text{ini}} = \frac{1}{3} - 0$$

$$\Delta x = \frac{1}{3} = 0.33$$



Scan for Video solution



$\therefore$  Scan for Video solution



### 16. (0.53 to 0.56)

Given:  $5x - 2 \cos x - 1 = 0$

By Newton's Raphson Iterative Scheme

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$\Rightarrow x_{k+1} = x_k - \frac{(5x_k - 2 \cos x_k - 1)}{(5 + 2 \sin x_k)}$$

$$= \frac{2x_k \cdot \sin x_k + 2 \cdot \cos x_k + 1}{5 + 2 \sin x_k}$$

Let initial guess is  $x_0 = 0$

$$x_1 = \frac{2+1}{5} = 0.6$$

$$x_2 = \frac{2(0.6)\sin(0.6) + 2\cos(0.6) + 1}{5 + 2\sin(0.6)} = 0.543$$

$$x_3 = \frac{2(0.543)\cdot\sin(0.543) + 2\cos(0.543) + 1}{5 + 2\sin(0.543)} = 0.542$$

$\therefore$  The real root is 0.54.



Scan for Video solution



### 17. (0.05 to 0.07)

**Given:**  $f(x) = e^x - 1$ ,  $x_0 = 1$

$$\Rightarrow f'(x) = e^x$$

By Newton Raphson method;

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$x_1 = 1 - \frac{e^1 - 1}{e^1} = 1 - 1 + \frac{1}{e} = \frac{1}{e}$$

$$\text{now, } x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$= \frac{1}{e} - \left( \frac{e^{1/e} - 1}{e^{1/e}} \right) = \frac{1}{e} - 1 + \frac{1}{e^{1/e}}$$

$$= 0.37 - 1 + 0.69$$

$$x_2 = 0.06$$

We know actual root of  $e^x - 1 = 0$  is  $x = 0$ .

$$\therefore \text{Absolute error} = |0 - 0.06| = 0.06$$



Scan for Video solution



### 18. (c)

**Given:**  $f(x) = x^3 + 2x - 1$ ,  $x_0 = 1.2$

$$f'(x) = 3x^2 + 2$$

By Newton Raphson method;

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$x_1 = 1.2 - \frac{[1.2^3 + 2 \times 1.2 - 1]}{3 \times 1.2^2 + 2} = 0.705$$



Scan for Video solution



### 19. (b)

Let  $f(x_1, x_2) = 10x_2 \sin x_1 - 0.8$

$$g(x_1, x_2) = 10x_2^2 - 10x_2 \cos x_1 - 0.6$$

$$x_1 = 0.0 \text{ and } x_2 = 1.0$$

Then Jacobian matrix for  $(f, g) \rightarrow (x_1, x_2)$

$$J = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \frac{\partial f}{\partial x_2} \\ \frac{\partial g}{\partial x_1} & \frac{\partial g}{\partial x_2} \end{bmatrix} = \begin{bmatrix} 10x_2 \cos x_1 & 10 \sin x_1 \\ 10x_2 \sin x_1 & 20x_2 - 10 \cos x_1 \end{bmatrix}$$

$\therefore J$  at  $x_1 = 0.0$  and  $x_2 = 1.0$  is,

$$J = \begin{bmatrix} 10 \times 1 \times 1 & 10 \times 0 \\ 10 \times 1 \times 0 & 20 \times 1 - 10 \times 1 \end{bmatrix} = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$$



Scan for Video solution



### 20. (c)

**Given:**  $f(x) = x + \sqrt{x} - 3 = 0$  and initial guess  $x_0 = 2$ .

As per Newton Raphson iteration scheme,

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)} = x_k - \frac{(x_k + \sqrt{x_k} - 3)}{\left(1 + \frac{1}{2\sqrt{x_k}}\right)}$$

$$\Rightarrow x_{k+1} = \frac{x_k + \frac{\sqrt{x_k}}{2} - x_k - \sqrt{x_k} + 3}{\left(1 + \frac{1}{2\sqrt{x_k}}\right)}$$

$$\Rightarrow x_{k+1} = \frac{3 - \frac{\sqrt{x_k}}{2}}{\left(1 + \frac{1}{2\sqrt{x_k}}\right)}$$

For  $k = 0$

$$x_1 = \frac{3 - \frac{\sqrt{x_0}}{2}}{\left(1 + \frac{1}{2\sqrt{x_0}}\right)} = \frac{3 - \frac{\sqrt{2}}{2}}{\left(1 + \frac{1}{2\sqrt{2}}\right)} = 1.6939 \approx 1.694$$

$\therefore$  Value after 1<sup>st</sup> iteration = 1.694



Scan for Video solution



21. (a)

$$\text{Let } f(x) = x^2 - N$$

$$f'(x) = 2x$$

According to Newton – Raphson formula,

$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$

$$x_{i+1} = x_i - \frac{x_i^2 - N}{2x_i} = x_i - \frac{x_i}{2} + \frac{N}{2x_i}$$

$$x_{i+1} = \frac{x_i}{2} + \frac{N}{2x_i}$$

$$x_{i+1} = \frac{1}{2} \left( x_i + \frac{N}{x_i} \right)$$



Scan for Video solution



22. (a)

$$\text{Let } f(x) = x^2 - 117$$

$$\text{By Newton Raphson method, } x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

$$x_{k+1} = x_k - \frac{(x_k^2 - 117)}{2x_k}$$

$$= x_k - \frac{x_k}{2} + \frac{117}{2x_k} = \frac{x_k}{2} + \frac{117}{2x_k}$$

$$x_{k+1} = \frac{1}{2} \left( x_k + \frac{117}{x_k} \right)$$



Scan for Video solution



23. (a)

$$\text{Given: } f(x) = e^x - 1, x_0 = -1$$


By Newton's method;

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$x_1 = x_0 - \frac{e^{x_0} - 1}{e^{x_0}} \quad [\because f'(x) = e^x]$$

$$x_1 = -1 - \left( \frac{e^{-1} - 1}{e^{-1}} \right) = \frac{-e^{-1} - e^{-1} + 1}{e^{-1}}$$

$$= \frac{1 - 2e^{-1}}{e^{-1}} = \frac{1 - 2/e}{1/e} = e - 2 = 0.71828$$



24. (c)

$$\text{Given: } x = e^{-x}$$


$$\Rightarrow f(x) = x - e^{-x} = 0$$

For  $f(x) = 0$ , Newton – Raphson scheme is given by


$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)} = x_k - \frac{(x_k - e^{-x_k})}{(1 + e^{-x_k})}$$

$$\Rightarrow x_{k+1} = \frac{x_k e^{-x_k} + e^{-x_k}}{(1 + e^{-x_k})} = \frac{e^{-x_k} (1 + x_k)}{(1 + e^{-x_k})}$$

$$\therefore x_{n+1} = (1 + x_n) \frac{e^{-x_n}}{(1 + e^{-x_n})}$$



25. (a)



Area under approximated curve will be less as compared to actual smooth function.  
Numerically computed area < Actual area  
(Trapezoidal Rule)



Scan for Video solution



26. (d)

$$\text{Given: } I = \int_1^2 (4x^2 + 2x + 6) dx$$

For  $\int_a^b f(x) dx$ , if  $f(x)$  is polynomial of degree  $\leq 2$ ,

the integration by Simpson's  $\frac{1}{3}$ <sup>rd</sup> rule will be same as the exact value.

$\because f(x) = 4x^2 + 2x + 6$  is a polynomial of degree '2'

$$\Rightarrow I_e = I_s$$

$$\therefore \left| \frac{I_e - I_s}{I_e} \right| \times 100 = 0$$



Scan for Video solution



27. (20.00 to 21.00)

Given: Step size,  $h = 0.5$

By Simpson's  $1/3$ <sup>rd</sup> Rule, Area (A)

$$= \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots) + 2(y_2 + y_4 + \dots)]$$

$$= \frac{h}{3} [y_0 + 4y_1 + 2y_2 + 4y_3 + y_n]$$

$$= \frac{0.5}{3} [5 + 4 \times 7.25 + 2 \times 10 + 4 \times 13.25 + 17.00]$$

$$= 20.67 \text{ unit}^2$$



Scan for Video solution



28. (c)

Given:  $f(x) = e^x$ ,  $n = 4$ ,  $a = 0$ ,  $b = 1$

$$\text{Then } h = \frac{b-a}{n} = \frac{1-0}{4} = \frac{1}{4}$$

So,

|        |       |               |               |               |       |
|--------|-------|---------------|---------------|---------------|-------|
| x      | 0     | $\frac{1}{4}$ | $\frac{1}{2}$ | $\frac{3}{4}$ | 1     |
| $f(x)$ | 1     | $e^{1/4}$     | $e^{1/2}$     | $e^{3/4}$     | $e^1$ |
|        | $y_0$ | $y_1$         | $y_2$         | $y_3$         | $y_n$ |


Using Trapezoidal rule,

$$I = \int_0^1 e^x dx = \frac{h}{2} [y_0 + 2(y_1 + y_2 + y_3) + y_n]$$

$$= \frac{1/4}{2} [1 + 2(e^{1/4} + e^{1/2} + e^{3/4}) + e] = 1.727$$



Scan for Video solution



29. (0.38 to 0.42)

Given:  $f(x) = 10x - 20x^2$ ,  $n = 5$ ,  $a = 0$ ,  $b = 0.5$

$$\text{Then } h = \frac{b-a}{n} = \frac{0.5-0}{5} = 0.1$$

So,

|        |       |       |       |       |       |       |
|--------|-------|-------|-------|-------|-------|-------|
| x      | 0     | 0.1   | 0.2   | 0.3   | 0.4   | 0.5   |
| $f(x)$ | 0     | 0.8   | 1.2   | 1.2   | 0.8   | 0     |
|        | $y_0$ | $y_1$ | $y_2$ | $y_3$ | $y_4$ | $y_n$ |

$$\text{Using Trapezoidal rule, } I = \int_0^{0.5} (10x - 20x^2) dx$$


$$= \frac{h}{2} [y_0 + y_n + 2(y_1 + y_2 + y_3 + y_4)]$$

$$= \frac{0.1}{2} [0 + 2(0.8 + 1.2 + 1.2 + 0.8) + 0]$$

$$\therefore \int_0^{0.5} (10x - 20x^2) dx = 0.40$$



Scan for Video solution



30. (a)

Given: Integral  $\int_4^{5.2} \ln x dx$ ; step-size  $h = 0.3$

|         |         |           |           |           |           |
|---------|---------|-----------|-----------|-----------|-----------|
| x       | 4       | 4.3       | 4.6       | 4.9       | 5.2       |
| $\ln x$ | $\ln 4$ | $\ln 4.3$ | $\ln 4.6$ | $\ln 4.9$ | $\ln 5.2$ |

By Simpson's  $\frac{1}{3}$ <sup>rd</sup> rule,  $\int_4^{5.2} \ln x dx$

$$\begin{aligned} &= \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots) + 2(y_2 + y_4 + \dots)] \\ &= \frac{0.3}{3} [(\ln 4 + \ln 5.2) + 4(\ln 4.3 + \ln 4.9) + 2\ln(4.6)] \\ &= 1.8278 \\ \therefore \int_4^{5.2} \ln x dx &= 1.83 \end{aligned}$$



Scan for Video solution



31. (d)

Given: Integral  $I = \int_{-1}^{1.4} x|x| dx$  and step size  $h = 0.6$

|        |           |           |           |           |           |
|--------|-----------|-----------|-----------|-----------|-----------|
| x      | -1        | -0.4      | 0.2       | 0.8       | 1.4       |
| $x x $ | ( $y_0$ ) | ( $y_1$ ) | ( $y_2$ ) | ( $y_3$ ) | ( $y_n$ ) |

From Simpson's  $\frac{1}{3}$ <sup>rd</sup> rule,  $\int_a^b f(x) dx$  is given by

$$\begin{aligned} &= \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots) + 2(y_2 + y_4 + \dots)] \\ \therefore I &= \frac{0.6}{3} [(-1 + 1.96) + 4(-0.16 + 0.64) + 2(0.04)] \\ &= 0.592 \end{aligned}$$

$$\therefore \int_{-1}^{1.4} x|x| dx = 0.592$$



Scan for Video solution



32. (b)

The following rules will give zero error for polynomials up to the following degree:

Given, Polynomial is of 3 degree

|                                |                    |
|--------------------------------|--------------------|
| Rectangular Rule               | 0 degree ( $x^0$ ) |
| Trapezoidal Rule               | 1 degree ( $x^1$ ) |
| Simpson 1/3 <sup>rd</sup> Rule | 2 degree ( $x^2$ ) |
| Simpson 3/8 <sup>th</sup> Rule | 3 degree ( $x^3$ ) |

Since function is a polynomial of 3 degree hence Only Simpson's rule will give zero error.



Scan for Video solution



33. (5.1 to 5.5)

Given: Integral  $\int_0^{\frac{\pi}{2}} (8 + 4 \cos x) dx = I$  (say)

$$(i) \text{ Analytic Value} = \int_0^{\frac{\pi}{2}} (8 + 4 \cos x) dx$$

$$= 8x + 4 \sin x \Big|_0^{\frac{\pi}{2}} = 8\left(\frac{\pi}{2}\right) + 4 \sin\left(\frac{\pi}{2}\right)$$

$$= 4(1 + \pi) = A \text{ (say)}$$

(ii) Using Trapezoidal rule by considering end points,

$$\Rightarrow T = \frac{h}{2} [(y_0 + y_n)]$$

$$= \frac{\left(\frac{\pi}{2}\right)}{2} \left[ (8 + 4 \cos x) \Big|_{x=0} + (8 + 4 \cos x) \Big|_{x=\frac{\pi}{2}} \right]$$

$$= \frac{\pi}{4} [16 + 4] = 5\pi$$

$$\therefore T = 5\pi$$

$$\% \text{ error} = \left| \frac{A - T}{A} \right| \times 100$$

$$= \left| \frac{4 + 4\pi - 5\pi}{4 + 4\pi} \right| \times 100 = 5.182 \%$$

$$\text{Absolute \% error} = 5.2 \%$$



Scan for Video solution



34. (63 to 63)

Given: Integral  $I = \int_2^4 x^3 dx$

Number of sub-intervals = 2

$$\Rightarrow \text{step size} = \frac{4 - 2}{2} = 1 = h$$

|        |           |            |            |
|--------|-----------|------------|------------|
| x      | 2         | 3          | 4          |
| $f(x)$ | $y_0 = 8$ | $y_1 = 27$ | $y_n = 64$ |

By Trapezoidal rule,

$$\begin{aligned} I &= \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots)] \\ &= \frac{1}{2} [(8 + 64) + 2(27)] = 4 + 32 + 27 = 63 \\ \therefore \int_2^4 x^3 dx &= 63 \end{aligned}$$



Scan for Video solution



### 35. (a)

**Given:**

The Points are P (0, 3), Q (0.5, 4), R (1, 5) are collinear.

The difference in the values of  $\int_0^1 f(x) dx$  will be zero

in between Trapezoidal rule, Simpson's rule and analytical integration.

% Error between Trapezoidal and Simpson's rule = 0



Scan for Video solution



### 36. (a)

For a linear function ( $f(x) = ax + b$ ), the integration of  $f(x)$  between any two points using trapezoidal rule gives the same value as the Analytical Integration.



Scan for Video solution



### 37. (0.175 to 0.195)

**Given**

$$\text{Integral} = \int_0^\pi (\sin x + \cos x) dx$$

(i) **Analytic Value:** Let analytic value be 'A'

$$\begin{aligned} A &= \int_0^\pi (\sin x + \cos x) dx = -\cos x + \sin x \Big|_0^\pi \\ &= +1 + 0 - (-1 + 0) = 2 \end{aligned}$$

$$\Rightarrow A = 2$$

(ii) **Trapezoidal Rule:**

Number of sub-intervals = 3

$$\Rightarrow \text{step size} = h = \frac{\pi - 0}{3} = \frac{\pi}{3}$$

|        |           |                                |                                 |            |
|--------|-----------|--------------------------------|---------------------------------|------------|
| x      | 0         | $\pi/3$                        | $2\pi/3$                        | $\pi$      |
| $f(x)$ | $y_0 = 1$ | $y_1 = \frac{1 + \sqrt{3}}{2}$ | $y_2 = \frac{-1 + \sqrt{3}}{2}$ | $y_n = -1$ |

$$\begin{aligned} \therefore \int_0^\pi (\sin x + \cos x) dx &= \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2)] \\ &= \frac{\pi}{6} \left[ 1 - 1 + 2 \left( \frac{1 + \sqrt{3}}{2} + \frac{-1 + \sqrt{3}}{2} \right) \right] \\ &= \frac{\pi}{6} (2\sqrt{3}) = \frac{\pi}{\sqrt{3}} = 1.8138 \end{aligned}$$

$$\therefore \text{Error} = A - 1.8138 = 2 - 1.8138 = 0.186$$

Scan for Video solution

### 38. (2 to 2)

$$\text{Given: } f(x) = \frac{3}{5}x^2 + \frac{9}{5}$$

Least number of sub-intervals for Simpson's  $\frac{1}{3}$ <sup>rd</sup> rule is '2'.

$$\therefore \text{Step size} = \frac{b-a}{2} = \frac{1-0}{2} = 0.5$$

|        |                     |                                    |                      |
|--------|---------------------|------------------------------------|----------------------|
| x      | 0                   | 0.5                                | 1                    |
| $f(x)$ | $y_0 = \frac{9}{5}$ | $y_1 = \frac{3}{20} + \frac{9}{5}$ | $y_n = \frac{12}{5}$ |

$\therefore$  From Simpson's  $\frac{1}{3}$ <sup>rd</sup> rule,

$$\int_a^b f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4y_1]$$

$$= \frac{0.5}{3} \left[ \frac{9}{5} + \frac{12}{5} + 4 \left( \frac{3}{20} + \frac{9}{5} \right) \right]$$

= 2 (Value will be as Analytical value)

$$\therefore \int_0^1 \left( \frac{3}{5}x^2 + \frac{9}{5} \right) dx = 2$$




Scan for Video solution



39. (a)

For the function  $y = x^2$  between  $x_1$  and  $x_2$



$I =$  exact value of the integral is given by area under the curve  $y = x^2$  between  $x_1$  and  $x_2$  bounded by X-axis.

$J =$  Approximate value is obtained by the area under the straight-line AB between  $x_1$  and  $x_2$  bounded by X-axis.

So,  $J > I$



Scan for Video solution



40. (1.36 to 1.37)

Given:

$$f(x) = 0.2 + 2.5x - 200x^2 + 675x^3 - 900x^4 + 400x^5$$

Step size,  $h = 0.4$

|      |     |       |       |
|------|-----|-------|-------|
| x    | 0   | 0.4   | 0.8   |
| f(x) | 0.2 | 2.456 | 0.232 |

By Simpson's  $1/3$ <sup>rd</sup> Rule,

$$I = \int_0^{0.8} f(x) dx = \frac{h}{3} [y_0 + 4y_1 + y_n]$$

$$= \frac{0.4}{3} [0.2 + 4 \times 2.456 + 0.232]$$

$$I = 1.367$$


Scan for Video solution



41. (21.8 to 22.2)

Given: Step size  $h = 0.1$

|        |           |            |            |            |             |
|--------|-----------|------------|------------|------------|-------------|
| x      | 0         | 0.1        | 0.2        | 0.3        | 0.4         |
| $f(x)$ | $y_0 = 0$ | $y_1 = 10$ | $y_2 = 40$ | $y_3 = 90$ | $y_n = 160$ |

From trapezoidal rule,

$$\int_a^b f(x) dx = \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots)]$$

$$\Rightarrow \int_0^{0.4} f(x) dx = \frac{0.1}{2} [(0 + 160) + 2(10 + 40 + 90)] \\ = \frac{0.1}{2} [160 + 280] = 22$$

$$\therefore \int_0^{0.4} f(x) dx = 22$$


Scan for Video solution



42. (0.68 to 0.70)

Given:  $h = 1 \Rightarrow$  Number of sub intervals = 1

$\Rightarrow$  Integration is based on end point

|        |                        |                       |
|--------|------------------------|-----------------------|
| x      | 1                      | 2                     |
| $f(x)$ | $y_0 = 1 \cdot \ln(0)$ | $y_n = 2 \cdot \ln 2$ |

$$\therefore \int_1^2 x \ln x dx = \frac{h}{2} [(y_0 + y_n)] \\ = \frac{1}{2} [0 + 2 \cdot \ln 2] = \ln 2 = 0.693$$

$$\text{For } h = 1, \int_1^2 x \cdot \ln x \cdot dx = 0.693$$


Scan for Video solution



**43. (1.1 to 1.2)**

**Given:** Integral  $\int_1^3 \frac{1}{x} dx$  and step size  $h = 1$

|       |         |           |           |
|-------|---------|-----------|-----------|
| x     | 1       | 2         | 3         |
| $1/x$ | $y_0=1$ | $y_1=1/2$ | $y_n=1/3$ |

From Trapezoidal rule

$$\begin{aligned} \int_1^3 \frac{1}{x} dx &= \frac{h}{2} [(y_0 + y_n) + 2(y_1)] \\ \Rightarrow \int_1^3 \frac{1}{x} dx &= \frac{1}{2} \left[ \left( 1 + \frac{1}{3} \right) + 2 \left( \frac{1}{2} \right) \right] \\ &= \frac{1}{2} \left[ \frac{7}{3} \right] = \frac{7}{6} = 1.167 \end{aligned}$$

$$\Rightarrow \int_1^3 \frac{1}{x} dx = 1.167$$



Scan for Video solution

**44. (1.74 to 1.76)**

**Given:** Integral  $I = \int_{2.5}^4 \ln x dx$

Number of sub intervals = 5

$$\Rightarrow \text{step size, } h = \frac{4 - 2.5}{5} = 0.3$$

|         |           |           |           |           |           |           |
|---------|-----------|-----------|-----------|-----------|-----------|-----------|
| x       | 2.5       | 2.8       | 3.1       | 3.4       | 3.7       | 4.0       |
| $\ln x$ | $\ln 2.5$ | $\ln 2.8$ | $\ln 3.1$ | $\ln 3.4$ | $\ln 3.7$ | $\ln 4.0$ |

From Trapezoidal rule;

$$\begin{aligned} \int_a^b f(x) dx &= \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})] \\ \int_{2.5}^4 \ln x dx &= \frac{0.3}{2} \left[ (\ln 2.5 + \ln 4.0) + 2(\ln 2.8 + \ln 3.1 + \ln 3.4 + \ln 3.7) \right] \\ &= 0.15 (2.3025 + 9.3852) = 1.7533 \end{aligned}$$

$$\therefore \int_{2.5}^4 \ln x dx = 1.7533$$



Scan for Video solution

**45. (1.10 to 1.12)**

**Given:** Integral  $= \int_{-1}^1 |x| dx$ , no. of sub-intervals = 3

$$\therefore \text{step size, } h = \frac{1 - (-1)}{3} = \frac{2}{3}$$

|   |    |      |     |   |
|---|----|------|-----|---|
| x | -1 | -1/3 | 1/3 | 1 |
| x | 1  | 1/3  | 1/3 | 1 |

By Trapezoidal rule,

$$\begin{aligned} \int_b^a f(x) dx &= \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots)] \\ \Rightarrow \int_{-1}^1 |x| dx &= \frac{\left(\frac{2}{3}\right)}{2} \left[ (1+1) + 2\left(\frac{1}{3} + \frac{1}{3}\right) \right] \\ &= \frac{1}{3} \left[ 2 + \frac{4}{3} \right] = \frac{10}{9} = 1.111 \end{aligned}$$

$$\therefore \int_{-1}^1 |x| dx = 1.111$$



Scan for Video solution

**46. (0.50 to 0.53)**

**Given:**  $h = 1$

Exact value of integral  $\int_0^4 (x^4 + 10) dx$

$$= \left[ \frac{x^5}{5} + 10x \right]_0^4 = \frac{4^5}{5} + 10 \times 4 = 244.8$$

|      |    |    |    |    |     |
|------|----|----|----|----|-----|
| x    | 0  | 1  | 2  | 3  | 4   |
| f(x) | 10 | 11 | 26 | 91 | 266 |

Using Simpson's 1/3<sup>rd</sup> rule

$$\begin{aligned} \int_0^4 f(x) dx &= \frac{h}{3} [y_0 + 4y_1 + 2y_2 + 4y_3 + y_n] \\ &= \frac{1}{3} [10 + 4 \times 11 + 2 \times 26 + 4 \times 91 + 266] = 245.33 \end{aligned}$$

Approximate value = 245.33

$\therefore$  Magnitude of error =  $245.33 - 244.8 = 0.53$



Scan for Video solution




**47. (d)**

Iteration scheme → Fitting curve → order of fitting polynomial

1. Trapezoidal rule → straight line → first order
2. Simpson's  $1/3^{\text{rd}}$  rule → quadratic function → second order
3. Simpson's  $3/8^{\text{th}}$  rule → cubic function → third order.

$$\therefore P \rightarrow 3, Q \rightarrow 1, R \rightarrow 2$$

**48. (d)**

**Given:**  $f(x) = \frac{1}{x}$ ,  $h = \frac{1.5 - 0.5}{2} = 0.5$

Exact value of  $\int_{0.5}^{1.5} \frac{dx}{x} = [\log x]_{0.5}^{1.5} = \log 3 = 1.0986$

Approximate value by Simpson's rule with 3 point is,


|        |       |     |       |
|--------|-------|-----|-------|
| x      | 0.5   | 1.0 | 1.5   |
| $f(x)$ | 1/0.5 | 1/1 | 1/1.5 |

$$I = \frac{h}{3} [f(0.5) + 4f(1.0) + f(1.5)]$$

$$\therefore I = \frac{0.5}{3} \left[ \frac{1}{0.5} + 4 \times 1 + \frac{1}{1.5} \right] = 1.1111$$

So, approximate value exceeds the exact value by Approximate value – exact value

$$= 1.1111 - 1.0986 = 0.012$$

**49. (c)**

**Given:** integral is  $\int_1^3 \frac{1}{x} dx = I(\text{say})$

Number of sub intervals = 2


$$\Rightarrow \text{step size } h = \frac{b-a}{n} = \frac{3-1}{2} = 1$$

|              |           |             |             |
|--------------|-----------|-------------|-------------|
| x            | 1         | 2           | 3           |
| $f(x) = 1/x$ | $y_0 = 1$ | $y_1 = 1/2$ | $y_n = 1/3$ |

Applying Simpson's  $\frac{1}{3}^{\text{rd}}$  rule

$$\begin{aligned} \int_1^3 \frac{1}{x} dx &= \frac{h}{3} [(y_0 + y_n) + 4y_1] = \frac{1}{3} \left[ \left( 1 + \frac{1}{3} \right) + 4 \left( \frac{1}{2} \right) \right] \\ &= \frac{1}{3} \left[ 3 + \frac{1}{3} \right] = 1 + \frac{1}{9} = 1.111 \end{aligned}$$

$$\therefore \int_1^3 \frac{1}{x} dx = 1.111$$

**50. (b)**

**Given:** Step size  $h = \frac{360 - 0}{6} = 60^\circ = \frac{\pi}{3}$

|                |   |      |      |     |     |      |     |
|----------------|---|------|------|-----|-----|------|-----|
| Angle (degree) | 0 | 60   | 12   | 180 | 240 | 300  | 360 |
| Torque (N m)   | 0 | 1066 | -323 | 0   | 323 | -355 | 0   |

$$\text{Flywheel energy F.E.} = \int_{\theta=0^\circ}^{\theta=360^\circ} T \cdot d\theta$$


$$\text{Using Simpson's rule, } \int_a^b f(x) dx$$

$$= \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots) + 2(y_2 + y_4 + \dots)]$$

$$\text{F.E.} =$$

$$\frac{\pi}{3(3)} [(0+0) + 4(1066+0-355)] + 2(-323+323)$$

$$= 992.74 \text{ J} \Rightarrow \int T \cdot d\theta \approx 993 \text{ J}$$



**51. (a)****Given:**  $h = 0.25$ 

|      |   |        |     |      |      |
|------|---|--------|-----|------|------|
| x    | 0 | 0.25   | 0.5 | 0.75 | 1.0  |
| f(x) | 1 | 0.9412 | 0.8 | 0.64 | 0.50 |

According to Simpson's 1/3<sup>rd</sup> Rule,

$$\int_0^1 f(x) dx = \frac{h}{3} [(y_0 + y_n) + 2y_2 + 4(y_1 + y_3)] \\ = \frac{0.25}{3} [(1 + 0.5) + 2 \times 0.8 + 4 \times (0.9412 + 0.64)]$$

$$I = 0.7854$$



Scan for Video solution

**52. (-0.152 to -0.149)**

**Given:** D.E:  $\frac{du}{dt} = 1 - 2tu^2$

$u_{n-1} = 1.75, h = 0.01, t_{n-1} = 3.14$

By Euler's Backward method:

$u_n = u_{n-1} + h f(u_n, t_n) \quad [\because f(u, t) = 1 - 2tu^2] \\ u_n = 1.75 + 0.01 [1 - 2t_n u_n^2] \\ u_n = 1.75 + 0.01 [1 - 2 \times 3.15 u_n^2] \\ u_n = 1.76 - 0.063 u_n^2 \\ 0.063 u_n^2 + u_n - 1.76 = 0 \\ u_n = 1.599$

$\therefore u_n - u_{n-1} = 1.599 - 1.75 = -0.151$



Scan for Video solution

**53. (0.999 to 1.001)**

**Given:**  $\frac{dy}{dt} + 2y = 0; y(0) = 1$

 $\Delta t \rightarrow$  step size for forward Euler method.For stability of solution of  $\frac{dy}{dt} = \lambda y$ , solved using

forward Euler method,

$|1 + h\lambda| < 1 \quad \text{where 'h' is step size} \\ -1 < 1 + h\lambda < 1 \Rightarrow -2 < h\lambda < 0$

$\Rightarrow h \leq \frac{2}{|\lambda|}$

$\therefore \text{In given problem, } \Delta t \leq \frac{2}{|-2|} = 1$

For stability,

$\Delta t \leq 1$



Scan for Video solution

**54. (6.3 to 6.5)**

**Given:**  $\frac{dy}{dx} = 4(x+2) - y, f(x, y) = 4(x+2) - y$

$h = 0.2$

For finding  $y_2$  two iterations has to be followed

|   | $x_0$ | $x_1$ | $x_2$ |
|---|-------|-------|-------|
| x | 1     | 1.2   | 1.4   |
| y | 3     | ?     | ?     |

By Euler's method

$y_1 = y_0 + hf(x_0, y_0)$

$= 3 + 0.2 f(1, 3) \quad \therefore x_0 = 1, y_0 = 3$

$= 3 + 0.2 [4(1+2) - 3]$

$y_1 = 4.8$

$\therefore x_1 = 1.2, y_1 = 4.8$

$y_2 = y_1 + hf(x_1, y_1)$

$= 4.8 + 0.2 f(1.2, 4.8)$

$= 4.8 + 0.2 [4(1.2+2) - 4.8] = 4.8 + 1.6$

$y_2 = 6.4$



Scan for Video solution

**55. (b)**If  $s = \alpha$  is a root of  $\varphi(s) = a_n s^n + a_{n-1} s^{n-1} + \dots + a_0$  for 'k' times,then  $(f(\alpha) = f'(\alpha) = f''(\alpha) = \dots = f^{(k-1)}(\alpha) = 0)$ Given  $s = -\sigma$  is a triple root of  $\phi(s) = 0$  $\Rightarrow \phi(s) = 0; \phi'(s) = 0; \phi''(s) = 0$  at  $s = -\sigma$ 

$\Rightarrow \phi(s) = 0; \frac{d\phi}{ds} = 0; \frac{d^2\phi}{ds^2} = 0 \text{ at } s = -\sigma$



Scan for Video solution



**56. (a)****Given**

$$\frac{dy}{dt} = -\pi y$$

For Forward Euler scheme,

$$\frac{y(t_{n+1}) - y(t_n)}{h} = -\pi y(t_n)$$

to have a stable solution for  $\frac{dy}{dt} = \lambda y$ 

$$\text{Step size } h, \quad 0 < h < \frac{2}{|\lambda|}$$

$$\Rightarrow 0 < h < \frac{2}{|-\pi|} \Rightarrow 0 < h < \frac{2}{\pi}$$

$$\therefore \text{For stability, } 0 < h < \frac{2}{\pi}$$



Scan for Video solution

**57. (a)**

Using fourth order central divided difference formula for third derivative

$$f'''(x_i) = \frac{-f(x_{i+3}) + 8f(x_{i+2}) - 13f(x_{i+1}) + 13f(x_{i-1}) - 8f(x_{i-2}) + f(x_{i-3})}{8h^3}$$



Scan for Video solution

**58. (2.55 to 2.65)****Given**

$$\frac{dy}{dt} = y \text{ and } y(0) = 1$$

Explicit forward-Euler scheme for a DE  $\frac{dy}{dt} = f(t, y)$ where  $y(t_0) = y_0$ , is given by

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$\Rightarrow y_{i+1} = y_i + h \cdot y_i$$

$$\Rightarrow y_{i+1} = (1 + h) \cdot y_i$$

$$\therefore y(0.1) = (1 + h) \cdot y(0)$$

$$y(0.2) = (1 + h) \cdot y(0.1) = (1 + h)^2 \cdot y(0)$$

$$y(0.3) = (1 + h) \cdot y(0.2) = (1 + h)^3 \cdot y(0)$$

and so on.....

$$\text{So, } y(1) = (1 + h)^{10} \cdot y(0)$$

$$\Rightarrow y(1) = (1 + 0.1)^{10} \cdot 1 = (1.1)^{10} = 2.59$$

$$\Rightarrow y(1) = 2.59$$



Scan for Video solution

**59. (7.95 to 8.05)**

$$\text{Given: } \frac{du}{dt} = 3t^2 + 1, \quad f(u, t) = 3t^2 + 1$$

$$u_0 = 0, t_0 = 0, \Delta t = 2 = h$$

By Euler's method

$$u_1 = u_0 + h f(u_0, t_0)$$

$$u_1 = u_0 + h (3t_0^2 + 1)$$

$$= 0 + 2(3 \times 0^2 + 1)$$

$$u_1 = 2$$

After first iteration when  $t = 2$  then  $y = 2$ .

$$\frac{du}{dt} = 3t^2 + 1$$

Integrating both sides, we get

$$u = \frac{t^3}{3} + t + c$$

$$\text{When } t = 0, u = 0$$

$$0 = 0 + 0 + c$$

$$c = 0$$

$$u = t^3 + t$$

$$\text{When } t = 2; u = 2^3 + 2 = 10$$

$$\text{Absolute error} = \text{Exact value} - \text{Approximate value} \\ = 10 - 2 = 8$$



Scan for Video solution

**60. (0.060 to 0.063)**

$$\text{Given: } y' = y + 2x - x^2; \quad y(0) = 1$$

$$\text{Exact solution is } y = x^2 + e^x$$

$$\therefore \text{At } x = 0.1, y(0.1) = (0.1)^2 + e^{0.1} = 1.11517$$

Runge kutta 2<sup>nd</sup> order Method:

$$\text{Step size } h = 0.1$$

$$y_{i+1}^{(P)} = y_i + h \cdot f(x_i, y_i)$$

$$y_{i+1}^{(C)} = y_i + \frac{h}{2}(f(x_i, y_i) + f(x_i + h, y_{i+1}^{(P)}))$$

Here  $f(x, y) = y + 2x - x^2$ .

At  $x = 0; y = 1$

$$\Rightarrow y_1^{(P)} = y(0.1)^{(P)} = 1 + 0.1(1 + 2(0) - 0^2)$$

$$= 1 + 0.1(1) = 1.1$$

$$\therefore y_1^{(c)} = 1 + \frac{0.1}{2}[1 + f(0.1, 1.1)]$$

$$= 1 + \frac{0.1}{2}\left[1 + (1.1 + 2(0.1) - (0.1)^2)\right]$$

$$= 1 + \frac{0.1}{2}[2.29] = 1.1145$$

$$\therefore y(0.1) = 1.1145$$

$$\therefore \% \text{ Error} = \left| \frac{y(0.1)_{\text{Exact}} - y(0.1)_{\text{RK}}}{y(0.1)_{\text{Exact}}} \right| \times 100$$

$$= \left| \frac{1.11517 - 1.1145}{1.11517} \right| \times 100 = 0.061$$

$$\therefore \% \text{ Error} = 0.061$$



Scan for Video solution



### 61. (0.6 to 0.7)

Given equation is  $\frac{dx}{dt} = -3x + 2; x(0) = 1$

For the DE;  $\frac{dx}{dt} = \lambda x + k$  with  $x(t_0) = x_0$ , to have

a stable solution, step size  $h \leq \frac{2}{|\lambda|}$

$$\Rightarrow h \leq \frac{2}{|-3|} \Rightarrow h \leq \frac{2}{3}$$

$$\Rightarrow h = 0.67$$

$\therefore$  Largest time step that can be taken without instability is 0.67.



Scan for Video solution



### 62. (d)

$$\text{Given: } \frac{dx}{dt} = 4t + 4 = f(t, x)$$

At  $t = 0, x = x_0; h = \Delta t = 0.2$

From Rk - 4<sup>th</sup> order method

$$x_{i+1} = x_i + \frac{h}{6}(k_1 + k_2 + k_3 + k_4)$$

where  $k_1 = f(t_0, x_0)$

$$k_2 = f\left(t_0 + \frac{h}{2}, x_0 + \frac{k_1}{2}\right)$$

$$k_3 = f\left(t_0 + \frac{h}{2}, x_0 + \frac{k_2}{2}\right)$$

$$k_4 = f(t_0 + h, x + k)$$

$$\therefore k_1 = f(0, x_0) = 4(0) + 4 = 4$$

$$k_2 = f\left(0 + \frac{0.2}{2}, x_0 + \frac{k_1}{2}\right) = 4(0.1) + 4 = 4.4$$

$$k_3 = f\left(0 + \frac{0.2}{2}, x_0 + \frac{k_2}{2}\right) = 4(0.1) + 4 = 4.4$$

$$k_4 = f(0 + 0.2, x_0 + k_3) = 4(0.2) + 4 = 4.8$$

The increment is given by

$$x_1 - x_0 = \frac{0.2}{6}(4 + 8.8 + 8.8 + 4.8) = 0.88$$



Scan for Video solution



### 63. (b)

#### Numerical Integral

Trapezoidal, Simpson's  $\frac{1}{3}$ <sup>rd</sup>, Simpson's  $\frac{3}{8}$ <sup>th</sup> rule,

#### Solution to transcendental equation

Newton-Raphson Method.

#### Solution to system of linear equations

Gauss Elimination method

#### Solution of differential equation

Runge-kutta methods



Scan for Video solution



### 64. (d)

Error in central difference is  $O(h^2)$ .

This mean Error  $\propto h^2$

If error for  $h = 0.03$  is  $2 \times 10^{-3}$

Then, error for  $h = 0.02$  is approximately,

$$= 2 \times 10^{-3} \times \left( \frac{0.02}{0.03} \right)^2 = \frac{8}{9} \times 10^{-3} = \frac{80}{9} \times 10^{-4}$$

$$E \approx 9.0 \times 10^{-4}$$



Scan for Video solution



65. (b)

**Given:**  $\frac{dy(x)}{dx} - y(x) = x$ , Step size  $h = 0.1$

$$\Rightarrow \frac{dy}{dx} = x + y = f(x, y)$$

Initial condition is  $y(0) = 0$

At  $x_0 = 0, y_0 = 0$ .

Using Euler's first order method,

$$y_{i+1} = y_i + h \cdot f(x_i, y_i)$$

$$\therefore y_1 = y_0 + h \cdot f(x_0, y_0) = y_0 + h(x_0 + y_0)$$

$$y_1 = 0 + 0.1(0 + 0) = 0$$

$$\Rightarrow y(0.1) = 0$$

$$y(0.2) = y(0.1) + h(0.1 + 0)$$

$$= 0 + (0.1)(0.1 + 0) = 0.01$$

$$y(0.3) = y(0.2) + h(0.2 + 0.01)$$

$$= 0.01 + 0.1(0.2 + 0.01)$$

$$= 0.01 + 0.1(0.21) = 0.01 + 0.021 = 0.031$$

$$\therefore \text{At } x = 0.3, y(0.3) = 0.031$$



Scan for Video solution



66. (c)

**Given:** Step size  $h = 0.01$

At  $t = 0, x = 0$ , we have to find  $x$  at  $t = 0.01$

Using trapezoidal rule,  $x(a + nh)$

$$= x(a) + \frac{h}{2} [f(a) + f(a + h) + \dots + f(a + (n-1)h)]$$

$$\frac{dx}{dt} = e^{-2t} u(t)$$

$$\frac{dx}{dt} = e^{-2t}; \quad (\because u(t) = 1; t > 0)$$

$$f(t) = e^{-2t}$$

$$x(0.01) = x(0) + \frac{h}{2} [f(0) + f(0.01)]$$

$$= 0 + \frac{0.01}{2} [1 + e^{-0.02}]$$

$$x(0.01) = 0.005[1 + e^{-0.02}]$$

$$x(0.01) = 0.0099$$



Scan for Video solution



67. (c)

When interpolated with all the 'n' points,  $\exists$  a unique interpolating polynomial of degree  $\leq n$ .

$\therefore$  Whether the polynomial is generated using Lagrange's interpolation (or) Newton interpolation method, the polynomial remains same.

$$\therefore I_L = I_N.$$



Scan for Video solution



**NOTE:**



# General Aptitude

1. Quantitative Aptitude..... 2.1 – 2.126
2. Analytical Aptitude ..... 2.127 – 2.172
3. Spatial Aptitude ..... 2.173 – 2.182
4. Verbal Aptitude ..... 2.183 – 2.258



# General Aptitude

## Syllabus

---

---

**Verbal Aptitude:** Basic English grammar: tenses, articles, adjectives, prepositions, conjunctions, verb-noun agreement, and other parts of speech Basic vocabulary: words, idioms, and phrases in context Reading and comprehension Narrative sequencing.

**Quantitative Aptitude:** Data interpretation: data graphs (bar graphs, pie charts, and other graphs representing data), 2- and 3-dimensional plots, maps, and tables Numerical computation and estimation: ratios, percentages, powers, exponents and logarithms, permutations and combinations, and series Mensuration and geometry Elementary statistics and probability.

**Analytical Aptitude:** Logic: deduction and induction, Analogy, Numerical relations and reasoning.

**Spatial Aptitude:** Transformation of shapes: translation, rotation, scaling, mirroring, assembling, and grouping Paper folding, cutting, and patterns in 2 and 3 dimensions.

## ME Chapter wise Weightage Analysis

| Chapter<br>Paper<br>Year \ | Ch.1 | Ch.2 | Ch.3 | Ch.4 |
|----------------------------|------|------|------|------|
| <b>2010</b>                | 6    | 3    | 0    | 6    |
| <b>2011</b>                | 9    | 0    | 0    | 6    |
| <b>2012</b>                | 7    | 2    | 0    | 6    |
| <b>2013</b>                | 9    | 0    | 0    | 6    |
| <b>2014 (P1)</b>           | 7    | 4    | 0    | 4    |
| <b>2014 (P2)</b>           | 7    | 5    | 0    | 3    |
| <b>2014 (P3)</b>           | 9    | 3    | 0    | 3    |
| <b>2014 (P4)</b>           | 8    | 4    | 0    | 3    |
| <b>2015 (P1)</b>           | 6    | 2    | 0    | 7    |
| <b>2015 (P2)</b>           | 5    | 1    | 0    | 9    |
| <b>2015 (P3)</b>           | 7    | 3    | 0    | 5    |
| <b>2016 (P1)</b>           | 7    | 3    | 0    | 5    |
| <b>2016 (P2)</b>           | 7    | 3    | 0    | 5    |
| <b>2016 (P3)</b>           | 4    | 6    | 0    | 5    |
| <b>2017 (P1)</b>           | 8    | 1    | 0    | 6    |
| <b>2017 (P2)</b>           | 9    | 2    | 0    | 4    |
| <b>2018 (P1)</b>           | 11   | 2    | 0    | 2    |
| <b>2018 (P2)</b>           | 8    | 5    | 0    | 2    |
| <b>2019 (P1)</b>           | 6    | 4    | 0    | 5    |
| <b>2019 (P2)</b>           | 7    | 2    | 0    | 6    |
| <b>2020 (P1)</b>           | 7    | 2    | 0    | 6    |
| <b>2020 (P2)</b>           | 7    | 2    | 0    | 6    |
| <b>2021 (P1)</b>           | 7    | 2    | 2    | 4    |
| <b>2021 (P2)</b>           | 6    | 6    | 0    | 3    |
| <b>2022 (P1)</b>           | 8    | 3    | 1    | 3    |
| <b>2022 (P2)</b>           | 9    | 2    | 1    | 3    |
| <b>2023</b>                | 6    | 1    | 2    | 6    |

## CE Chapter wise Weightage Analysis

| <b>Chapter<br/>Paper<br/>Year</b> | <b>Ch.1</b> | <b>Ch.2</b> | <b>Ch.3</b> | <b>Ch.4</b> |
|-----------------------------------|-------------|-------------|-------------|-------------|
| <b>2010</b>                       | 6           | 3           | 0           | 6           |
| <b>2011</b>                       | 9           | 0           | 0           | 6           |
| <b>2012</b>                       | 7           | 2           | 0           | 6           |
| <b>2013</b>                       | 9           | 2           | 0           | 4           |
| <b>2014 (P1)</b>                  | 8           | 4           | 0           | 3           |
| <b>2014 (P2)</b>                  | 4           | 6           | 0           | 5           |
| <b>2015 (P1)</b>                  | 5           | 3           | 0           | 7           |
| <b>2015 (P2)</b>                  | 7           | 3           | 0           | 5           |
| <b>2016 (P1)</b>                  | 6           | 4           | 0           | 5           |
| <b>2016 (P2)</b>                  | 8           | 2           | 0           | 5           |
| <b>2017 (P1)</b>                  | 10          | 1           | 0           | 4           |
| <b>2017 (P2)</b>                  | 6           | 3           | 0           | 6           |
| <b>2018 (P1)</b>                  | 11          | 2           | 0           | 2           |
| <b>2018 (P2)</b>                  | 11          | 2           | 0           | 2           |
| <b>2019 (P1)</b>                  | 3           | 5           | 0           | 7           |
| <b>2019 (P2)</b>                  | 8           | 0           | 0           | 7           |
| <b>2020 (P1)</b>                  | 7           | 3           | 0           | 5           |
| <b>2020 (P2)</b>                  | 7           | 3           | 0           | 5           |
| <b>2021 (P1)</b>                  | 4           | 3           | 3           | 5           |
| <b>2021 (P2)</b>                  | 6           | 5           | 1           | 3           |
| <b>2022 (P1)</b>                  | 7           | 3           | 2           | 3           |
| <b>2022 (P2)</b>                  | 6           | 5           | 1           | 3           |
| <b>2023 (P1)</b>                  | 5           | 3           | 1           | 6           |
| <b>2023 (P2)</b>                  | 7           | 1           | 1           | 6           |

## EE Chapter wise Weightage Analysis

| <b>Chapter<br/>Paper<br/>Year</b> | <b>Ch.1</b> | <b>Ch.2</b> | <b>Ch.3</b> | <b>Ch.4</b> |
|-----------------------------------|-------------|-------------|-------------|-------------|
| <b>2010</b>                       | 6           | 3           | 0           | 6           |
| <b>2011</b>                       | 9           | 0           | 0           | 6           |
| <b>2012</b>                       | 7           | 2           | 0           | 6           |
| <b>2013</b>                       | 9           | 0           | 0           | 6           |
| <b>2014 (P1)</b>                  | 6           | 2           | 0           | 7           |
| <b>2014 (P2)</b>                  | 8           | 0           | 0           | 7           |
| <b>2014 (P3)</b>                  | 7           | 3           | 0           | 5           |
| <b>2015 (P1)</b>                  | 7           | 1           | 0           | 7           |
| <b>2015 (P2)</b>                  | 7           | 3           | 0           | 5           |
| <b>2016 (P1)</b>                  | 3           | 7           | 0           | 5           |
| <b>2016 (P2)</b>                  | 7           | 1           | 0           | 7           |
| <b>2017 (P1)</b>                  | 8           | 3           | 0           | 4           |
| <b>2017 (P2)</b>                  | 8           | 3           | 0           | 4           |
| <b>2018</b>                       | 11          | 2           | 0           | 2           |
| <b>2019</b>                       | 7           | 3           | 0           | 5           |
| <b>2020</b>                       | 9           | 2           | 0           | 4           |
| <b>2021</b>                       | 8           | 2           | 1           | 4           |
| <b>2022</b>                       | 7           | 5           | 0           | 3           |
| <b>2023</b>                       | 8           | 0           | 0           | 7           |

## ECE Chapter wise Weightage Analysis

| <b>Chapter<br/>Paper<br/>Year</b> | <b>Ch.1</b> | <b>Ch.2</b> | <b>Ch.3</b> | <b>Ch.4</b> |
|-----------------------------------|-------------|-------------|-------------|-------------|
| <b>2010</b>                       | 6           | 3           | 0           | 6           |
| <b>2011</b>                       | 9           | 0           | 0           | 6           |
| <b>2012</b>                       | 7           | 2           | 0           | 6           |
| <b>2013</b>                       | 9           | 0           | 0           | 6           |
| <b>2014 (P1)</b>                  | 7           | 4           | 0           | 4           |
| <b>2014 (P2)</b>                  | 7           | 5           | 0           | 3           |
| <b>2014 (P3)</b>                  | 9           | 3           | 0           | 3           |
| <b>2014 (P4)</b>                  | 8           | 4           | 0           | 3           |
| <b>2015 (P1)</b>                  | 1           | 5           | 0           | 9           |
| <b>2015 (P2)</b>                  | 6           | 2           | 0           | 7           |
| <b>2015 (P3)</b>                  | 5           | 1           | 0           | 9           |
| <b>2016 (P1)</b>                  | 7           | 3           | 0           | 5           |
| <b>2016 (P2)</b>                  | 4           | 6           | 0           | 5           |
| <b>2016 (P3)</b>                  | 7           | 4           | 0           | 4           |
| <b>2017 (P1)</b>                  | 10          | 4           | 0           | 6           |
| <b>2017 (P2)</b>                  | 6           | 8           | 0           | 6           |
| <b>2018</b>                       | 9           | 0           | 0           | 6           |
| <b>2019</b>                       | 6           | 4           | 0           | 5           |
| <b>2020</b>                       | 9           | 0           | 0           | 6           |
| <b>2021</b>                       | 8           | 2           | 1           | 4           |
| <b>2022</b>                       | 6           | 4           | 2           | 3           |
| <b>2023</b>                       | 5           | 4           | 0           | 6           |

## CSE & IT Chapter wise Weightage Analysis

| <b>Chapter<br/>Paper<br/>Year</b> | <b>Ch.1</b> | <b>Ch.2</b> | <b>Ch.3</b> | <b>Ch.4</b> |
|-----------------------------------|-------------|-------------|-------------|-------------|
| <b>2010</b>                       | 6           | 3           | 0           | 6           |
| <b>2011</b>                       | 9           | 0           | 0           | 6           |
| <b>2012</b>                       | 7           | 2           | 0           | 6           |
| <b>2013</b>                       | 9           | 0           | 0           | 6           |
| <b>2014 (P1)</b>                  | 6           | 2           | 0           | 7           |
| <b>2014 (P2)</b>                  | 8           | 0           | 0           | 7           |
| <b>2014 (P3)</b>                  | 7           | 3           | 0           | 5           |
| <b>2015 (P1)</b>                  | 7           | 1           | 0           | 7           |
| <b>2015 (P2)</b>                  | 7           | 3           | 0           | 5           |
| <b>2015 (P3)</b>                  | 5           | 3           | 0           | 7           |
| <b>2016 (P1)</b>                  | 6           | 4           | 0           | 5           |
| <b>2016 (P2)</b>                  | 3           | 7           | 0           | 5           |
| <b>2017 (P1)</b>                  | 8           | 3           | 0           | 4           |
| <b>2017 (P2)</b>                  | 8           | 3           | 0           | 4           |
| <b>2018</b>                       | 12          | 1           | 0           | 2           |
| <b>2019</b>                       | 2           | 8           | 0           | 5           |
| <b>2020</b>                       | 7           | 2           | 0           | 6           |
| <b>2021 (P1)</b>                  | 7           | 2           | 2           | 4           |
| <b>2021 (P2)</b>                  | 6           | 2           | 3           | 4           |
| <b>2022</b>                       | 6           | 6           | 0           | 3           |
| <b>2023</b>                       | 6           | 0           | 3           | 6           |

**NOTE:**

# CHAPTER

1

# **QUANTITATIVE APTITUDE**

## Data Interpretation


---

**1. [MCQ]**

[GATE-ME-2023: 1M]

A certain country has 504 universities and 25951 colleges. These are categorised into Grades I, II, and III as shown in the given pie charts.

What is the percentage, correct to one decimal place, of higher education institutions (colleges and universities) that fall into Grade III?







## 2. [MCQ]

[GATE-CE-2023: 2M]


Which one of the options can be inferred about the mean, median and mode for the given probability distribution (i.e., probability mass function),  $P(x)$ , of a variable  $x$ ?



- (a) mean = median  $\neq$  mode
  - (b) mean = median = mode
  - (c) mean  $\neq$  median = mode
  - (d) mean  $\neq$  mode = median

### 3. [MCQ]

[GATE-CE-2022:2M]



The above frequency chart shows the frequency distribution of marks obtained by a set of students in an exam.

From the data presented above, which one of the following is CORRECT?

- (a) mean > mode > median
  - (b) mode > median > mean
  - (c) mode > mean > median
  - (d) median > mode > mean

#### **4. [MCQ]**

[GATE-CE-2022:1M]


$$x:y:z = \frac{1}{2} : \frac{1}{3} : \frac{1}{4}.$$

What is the value of  $\frac{x+z-y}{y}$ ?

## 5. [MCQ]

[GATE-CS-2021: 2M]

The number of units of a product sold in three different years and the respective net profits are presented in the figure below. The cost/unit in Year 3 was Rs. 1, which was half the cost/unit in Year 2. The cost/ unit in Year 3 was one-third of the cost/unit in Year 1. Taxes were paid on the selling price at 10% ,13% and 15% respectively for the three years. Net profit is calculated as the difference between the selling price and the sum of cost and taxes paid in that year.




The ratio of the selling price in Year 2 to the selling price in Year 3 is \_\_\_\_\_.



## 6. [MCQ]

[GATE-EC-2021:2M]




The number of minutes spent by two students, X and Y, exercising every day in a given week are shown in the bar chart above.

The number of days in the given week in which one of the students spent a minimum of 10% more than the other student, on a given day, is



## 7. [MCQ]

[GATE-ME -2021: 2M]



| Company | Ratio |
|---------|-------|
| C1      | 3:2   |
| C2      | 1:4   |
| C3      | 5:3   |
| C4      | 2:3   |
| C5      | 9:1   |
| C6      | 3:4   |

The distribution of employees at the rank of executives, across different companies C1, C2, ..., C6 is presented in the chart given above. The ratio of executives with a management degree to those without a management degree in each of these companies is provided in the table above. The total number of executives across all companies is 10,000.

The total number of management degree holders among the executives in companies C2 and CS together is .




## 8. [MCQ]

[GATE-EE-2020 : 2M]

The revenue and expenditure of four different companies P, Q, R and S in 2015 are shown in the figure. If the revenue of company Q in 2015 was 20% more than that in 2014, and company Q had earned a profit of 10% on expenditure in 2014. then its expenditure (in million rupees) in 2014 was.


**Revenue and Expenditure (in million rupees) of four companies P, Q, R and S in 2015**



### 9. [MCQ]

[GATE-CS-2020: 2M]

The total revenue of a company during 2014-2018 is shown in the bar graph. If the total expenditure of the company in each year is 500 million rupees, then the aggregate profit or loss (in percentage) on the total expenditure of the company during 2014-2018 is \_\_\_\_.




- (a) 16.67% profit      (b) 16.67% loss  
 (c) 20% profit      (d) 20% loss

**10. [MCQ]**

[GATE-EC-2020:2M]

The following figure shows the data of students enrolled in 5 years (2014 to 2018) for two schools P and Q. During this period, the ratio of the average number of the students enrolled in school P to the average of the difference of the number of students enrolled in schools P and Q is .







### 11. [MCQ]


[GATE-ME -2020: 2M]

The two pie-charts given below show the data of total students and only girls registered in different streams in a university. If the total number of students registered in the university is 5000, and the total number of the registered girls is 1500; then, the ratio of boys enrolled in Arts to the girls enrolled in Management is

Percentage of students enrolled in different streams in a University



### Percentage of girls enrolled in difference streams







**12. [MCQ]**

[GATE-CE-2020-2M]

The monthly distribution of 9 Watt LED bulbs sold by two firms X and Y from January to June 2018 is shown in the pie-chart and the corresponding table. If the total number of LED bulbs sold by two firms during April-June 2018 is 50000, then the number of LED bulbs sold by the firm Y during April-June 2018 is \_\_\_\_\_.

**Percentage of 9 Watt LED bulbs sold by the firms X and Y from January 2018 to June, 2018**




| Month    | Ratio of LED bulbs sold by two firms (X : Y) |
|----------|--|
| January  | 7 : 8  |
| February | 2 : 3  |
| March    | 2 : 1  |
| April    | 3 : 2  |
| May      | 1 : 4  |
| June     | 9 : 11                                       |

- (a) 11250      (b) 9750  
 (c) 8750      (d) 8250

**13. [MCQ] [GATE-CE-2020: 2M]**

The total expenditure of a family, on different activities in a month, is shown in the pie-chart. The extra money spent on education as compared to transport (in percent) is




- (a) 5      (b) 33.3  
 (c) 50      (d) 100

**14. [MCQ] [GATE-ME -2020: 2M]**

The bar graph shows the data of the students who appeared and passed in an examination for four schools

P, Q, R and S. The average of success rates (in percentage) of these four schools is \_\_\_\_.

Performance of Schools P, Q, R and S







- (a) 58.5 %      (b) 58.8 %  
 (c) 59.0 %      (d) 59.3 %

**15. [MCQ]**

**[GATE-ME-2020: 2M]**


Select the graph that schematically represents BOTH  $y = x^m$  and  $y = x^{1/m}$  properly in the interval  $0 \leq x \leq 1$ , for integer values of m, where  $m > 1$ .




**16. [MCQ]****[GATE-EC-2019:2M]**

The bar graph in Panel (a) shows the proportion of male and female illiterates in 2001 and 2011. The proportions of males and females in 2001 and 2011 are given in Panel (b) and (c), respectively. The total population did not change during this period.


The percentage increase in the total number of literates from 2001 to 2011 is \_\_\_\_.



Panel (a)



Panel (b)



Panel (c)

- (a) 30.43      (b) 33.43  
 (c) 34.43      (d) 35.43

**17. [MCQ]****[GATE-ME-2019: 2M]**

Mola is a digital platform for taxis in a city. It offers three types of rides - Pool, Mini and Prime. The Table

below presents the number of rides for the past four months. The platform earns one US dollar per ride. What is the percentage share of revenue contributed by Prime to the total revenues of Mola, for the entire duration?


| Type  | Month   |          |       |       |
|-------|---------|----------|-------|-------|
|       | January | February | March | April |
| Pool  | 170     | 320      | 215   | 190   |
| Mini  | 110     | 220      | 180   | 70    |
| Prime | 75      | 180      | 120   | 90    |

- (a) 16.24      (b) 23.97  
 (c) 25.86      (d) 38.74

**18. [MCQ]****[GATE-ME-2019: 2M]**

A firm hires employee at five different skill levels P, Q, R, S, T. The shares of employment at these skill levels of total employment in 2010 is given in the pie chart as shown. There were a total of 600 employees in 2010 and the total employment increased by 15% from 2010 to 2016. The total employment at skill levels P, Q and R remained unchanged during this period. If the employment at skill level S increased by 40% from 2010 to 2016, how many employees were there at skill level T in 2016?

Percentage share of skills in 2010



- (a) 30      (b) 35  
 (c) 60      (d) 72

**19. [MCQ]****[GATE-EE-2018 : 2M]**

P, Q, R and S crossed a lake in a boat that can hold a maximum of two persons, with only one set of oars. The following additional facts are available.

- (i) The boat held two persons on each of the three forward trips across the lake and one person on each of the two return trips.

- (ii) P is unable to row when someone else is in the boat.
  - (iii) Q is unable to row with anyone else except R.
  - (iv) Each person rowed for at least one trip.
  - (v) Only one person can row during a trip.  
Who rowed twice?

| <b>T<sub>0</sub></b> | <b>p</b> | <b>T</b> |
|----------------------|----------|----------|
| 25                   | 2        | 32.4     |
| 30                   | 5        | 42.0     |



**20. [MCQ]**

[GATE-EE-2018 : 2M]

A designer uses marbles of four different colours for his designs. The cost of each marble is the same, irrespective of the colour. The table below shows the percentage of marbles of each colour used in the current design. The cost of each marble increased by 25%. Therefore, the designer decided to reduce equal numbers of marbles of each colour to keep the total cost unchanged. What is the percentage of blue marbles in the new design?

| Blue | Black | Red | Yellow |
|------|-------|-----|--------|
| 40%  | 25%   | 20% | 15%    |



**21. [MCQ]**

[GATE-CS-2018: 2M]

In appreciation of the social improvements completed in a town, a wealthy philanthropist decided to gift Rs 750 to each male senior citizen in the town and Rs 1000 to each female senior citizen. Altogether, there were 300 senior citizens eligible for this gift. However, only  $\frac{8}{9}$ th of the eligible men and  $\frac{2}{3}$ rd of the eligible women claimed the gift. How much money (in Rupees) did the philanthropist give away in total?



**22. [MCQ]**

[GATE-CE-2018:1M]


The temperature  $T$  in a room varies as a function of the outside temperature  $T_o$  and the number of persons in the room  $p$ , according to the relation  $T = K (\Theta p + T_o)$ , where  $\Theta$  and  $K$  are constants. What would be the value of  $\Theta$  given the following data?



**23. [MCQ]**

[GATE-CE-2018: 2M]

Which of the following function(s) is an accurate description of the graph for the range(s) indicated?




- (i)  $y = 2x + 4$  for  $-3 \leq x \leq -1$   
 (ii)  $y = |x - 1|$  for  $-1 \leq x \leq 2$   
 (iii)  $y = ||x| - 1|$  for  $-1 \leq x \leq 2$   
 (iv)  $y = 1$  for  $2 \leq x \leq 3$

(a) (i), (ii) and (iii) only.  
 (b) (i), (ii) and (iv) only.  
 (c) (i) and (iv) only.  
 (d) (ii) and (iv) only.

**24. [MCQ]**

[GATE-CE-2018: 2M]

The annual average rainfall in a tropical city is 1000 mm. On a particular rainy day (24-hour period), the cumulative rainfall experienced by the city is shown in the graph. Over the 24-hour period, 50% of the rainfall falling on a rooftop, which had an obstruction free area of  $50 \text{ m}^2$ , was harvested into a tank. What is the total volume of water collected in the tank in liters?







**25. [MCQ]**

[GATE-CS,EE-2017 : 2M]

An air pressure contour line joins locations in a region having the same atmospheric pressure. The following is an air pressure contour plot of a geographical region. Contour lines are shown at 0.05 bar intervals in this plot.

If the possibility of a thunderstorm is given by how fast air pressure rises or drops over a region, which of the following regions is most likely to have a thunderstorm?







**26. [MCQ]**

[GATE-EC-2017:2M]

A contour line joins locations having the same height above the mean sea level. The following is a contour plot of a geographical region. Contour lines are shown at 25 m intervals in this plot.




Which of the following is the steepest path leaving from P?



27

[GATE-EC-2017:2M]

A contour line joins locations having the same height above the mean sea level. The following is a contour plot of a geographical region. Contour lines are shown at 25 m intervals in this plot.



The path from P to Q is best described by

- (a) Up-Down-Up-Down
  - (b) Down-Up-Down-Up
  - (c) Down-Up-Down
  - (d) Up-Down-Up


**28. [MCQ]**

[GATE-EC-2017:1M]

40 % of deaths on city roads may be attributed to drunken driving. The number of degrees needed to represent this as a slice of a pie chart is

**29. [MCQ]****[GATE-CE-2017: 2M]**

The points in the graph below represent the halts of a lift for durations of 1 minute, over a period of 1 hour.



Which of the following statements are correct?

- The elevator never moves directly from any non-ground floor to another non-ground floor over the one-hour period
  - The elevator stays on the fourth floor for the longest duration over the one hour-period
- (a) Only i                    (b) Only ii  
 (c) Both i and ii            (d) Neither i nor ii

**30. [MCQ]****[GATE-CE-2017: 2M]**

Students applying for hostel rooms are allotted rooms in order of seniority. Students already staying in a room will move if they get a room in their preferred list. Preferences of lower ranked applicants are ignored during allocation.


Given the data below, which room will Ajit stay in?

| Names   | Student seniority | Current room | Room preference list |
|---------|-------------------|--------------|----------------------|
| Amar    | 1                 | P            | R, S, Q              |
| Akbar   | 2                 | None         | R, S                 |
| Anthony | 3                 | Q            | P                    |
| Ajit    | 4                 | S            | Q, P, R              |

- (a) P                            (b) Q  
 (c) R                            (d) S

**31. [MCQ]****[GATE-CE-2017: 2M]**

The bar graph below shows the output of five carpenters over one month, each of whom made different items of furniture: chairs, tables, and beds.



Consider the following statements.


- The number of beds made by carpenter C2 is exactly the same as the number of tables made by carpenter C3.
- The total number of chairs made by all carpenters is less than the total number of tables.

Which one of the following is true?

- (a) Only i                            (b) Only ii  
 (c) Both i and ii                    (d) Neither i nor ii

**32. [MCQ]****[GATE-CS,EE-2017 : 2M]**


A contour line joins locations having the same height above the mean sea level. The following is a contour plot of a geographical region. Contour lines are shown at 25 m intervals in this plot. If in a flood, the water level rises to 525 m. which of the villages P, Q, R, S, T get submerged?



- (a) P, Q                            (b) P, Q, T  
 (c) R, S, T                            (d) Q, R, S

**33. [MCQ]****[GATE-ME-2017: 2M]**

The growth of bacteria (*lactobacillus*) in milk leads to curd formation. A minimum bacterial population density of 0.8 (in suitable units) is needed to form curd. In the graph below, the population density of *lactobacillus* in 1 litre of milk is plotted as a function of time, at two different temperatures, 25°C and 37°C.



Consider the following statements based on the data shown above:


- i. The growth in bacterial population stops earlier at 37°C as compared to 25°C
- ii. The time taken for curd formation at 25°C is twice the time taken at 37°C

Which one of the following options is correct?


- (a) Only i
- (b) Only ii
- (c) Both i and ii
- (d) Neither i nor ii

**34. [NAT]****[GATE-EE-2016 : 2M]**

The following graph represents the installed capacity for cement production (in tonnes) and the actual production (in tonnes) of nine cement plants of a cement company. Capacity utilization of a plant is defined as ratio of actual production of cement to installed capacity. A plant with installed capacity of at least 200 tonnes is called a large plant and a plant with lesser capacity is called a small plant. The difference between total production of large plants and small plants, in tonnes is \_\_\_\_.

**35. [MCQ]****[GATE-CS,EE-2016 : 2M]**

Choose the correct expression for  $f(x)$  given in the graph



- (a)  $f(x) = 1 - |x - 1|$
- (b)  $f(x) = 1 + |x - 1|$
- (c)  $f(x) = 2 - |x - 1|$
- (d)  $f(x) = 2 + |x - 1|$

**36. [MCQ]****[GATE-CS,CE-2016: 2M]**

A shaving set company sells 4 different types of razors, Elegance, Smooth, Soft and Executive. Elegance sells at Rs. 48, Smooth at Rs. 63, Soft at Rs. 78 and Executive at Rs. 173 per piece. The table below shows the numbers of each razor sold in each quarter of a year.

| Quarter \ Product | Elegance | Smooth | Soft  | Executive |
|-------------------|----------|--------|-------|-----------|
| Q1                | 27300    | 20009  | 17602 | 9999      |
| Q2                | 25222    | 19392  | 18445 | 8942      |
| Q3                | 28976    | 22429  | 19544 | 10234     |
| Q4                | 21012    | 18229  | 16595 | 10109     |


Which product contributes the greatest fraction to the revenue of the company in that year?

- (a) Elegance
- (b) Executive
- (c) Smooth
- (d) Soft

## 37. [MCQ]

[GATE-EC-2016: 2M]

The velocity  $V$  of a vehicle along a straight line is measured in m/s and plotted as shown with respect to time in seconds. At the end of the 7 seconds, how much will the odometer reading increase by (in m)?





- (a) 0
- (b) 3
- (c) 4
- (d) 5

## 38. [MCQ]

[GATE-ME-2016: 2M]

Which of the following curves represents the function  $y = \ln(|e^{[\sin(|x|)]}|)$  for  $|x| < 2\pi$ ? Here,  $x$  represents the abscissa and  $y$  represents the ordinate.




## 39. [MCQ]

[GATE-CE-2016: 2M]

Two finance companies, P and Q, declared fixed annual rates of interest on the amounts invested with them. The rates of interest offered by these companies may differ from year to year. Year-wise annual rates of interest offered by these companies are shown by the line graph provided below.

If the amounts invested in the companies, P and Q, in 2006 are in the ratio 8:9, then the amounts received after one year as interests from companies P and Q would be in the ratio:






**40. [MCQ] [GATE-EE,CS-2015 : 2M]**

The number of students in a class who have answered correctly, wrongly, or not attempted each question in an exam, are listed in the table below. The marks for each question are also listed. There is no negative or partial marking.


What is the average of the marks obtained by the class in the examination?

| Q.No. | Marks | Answered<br>Correctly | Answered<br>Wrongly | Not<br>Attempted |
|-------|-------|-----------------------|---------------------|------------------|
| 1     | 2     | 21                    | 17                  | 6                |
| 2     | 3     | 15                    | 27                  | 2                |
| 3     | 1     | 11                    | 29                  | 4                |
| 4     | 2     | 23                    | 18                  | 3                |
| 5     | 5     | 31                    | 12                  | 1                |




41. [NAT] [GATE-EE,CS-2015 : 2M]

The pie chart below has the breakup of the number of students from different departments in an engineering college for the year 2012. The proportion of male to female students in each department is 5:4. There are 40 males in Electrical Engineering. What is the difference between the numbers of female students in the Civil department and the female students in the Mechanical department?



42. [NAT] [GATE-CS,CE-2015: 2M]

The exports and imports (in crores of Rs.) of a country from the year 2000 to 2007 are given in the following bar chart. In which year is the combined percentage increase in imports and exports the highest?



**43. [MCQ]** [GATE-CE-2015: 2M]

Read the following table giving sales data of five types of batteries for years 2006 to 2012:

Out of the following, which type of battery achieved highest growth between the years 2006 and 2012?

| Year | Type I | Type II | Type III | Type IV | Type V |
|------|--------|---------|----------|---------|--------|
| 2006 | 75     | 144     | 114      | 102     | 108    |
| 2007 | 90     | 126     | 102      | 84      | 126    |
| 2008 | 96     | 114     | 75       | 105     | 135    |
| 2009 | 105    | 90      | 150      | 90      | 75     |
| 2010 | 90     | 75      | 135      | 75      | 90     |
| 2011 | 105    | 60      | 1065     | 45      | 120    |
| 2012 | 115    | 85      | 160      | 100     | 145    |

- (a) Type V
- (b) Type III
- (c) Type II
- (d) Type I

**44. [MCQ] [GATE-ME,EC-2015: 1M]**

An electric bus has onboard instruments that report the total electricity consumed since the start of the trip as well as the total distance covered. During a single day of operation, the bus travels on stretches M, N, O and P, in that order. The cumulative distances travelled and the corresponding electricity consumption are shown in the Table below:

| Stretch | Cumulative distance (km) | Electricity used (kWh) |
|---------|--------------------------|------------------------|
| M       | 20                       | 12                     |
| N       | 45                       | 25                     |
| O       | 75                       | 45                     |
| P       | 100                      | 57                     |

The stretch where the electricity consumption per km is minimum is

- (a) M
- (b) N
- (c) O
- (d) P


**45. [NAT] [GATE-EE,CS-2014 : 2 M]**

In a survey, 300 respondents were asked whether they own a vehicle or not. If yes, they were further asked to mention whether they own a car or scooter or both. Their responses are tabulated below. What percent of respondents do not own a scooter?

|                    |         | Men | Women |
|--------------------|---------|-----|-------|
| Own Vehicle        | Car     | 40  | 34    |
|                    | Scooter | 30  | 20    |
|                    | Both    | 60  | 46    |
| Do not own vehicle |         | 20  | 50    |

**46. [NAT] [GATE-EE,CS-2014 : 2M]**

The ratio of male to female students in a college for five years is plotted in the following line graph. If the number of female students doubled in 2009, by what percent did the number of male students increase in 2009?


**47. [MCQ] [GATE-EE,CS-2014 : 1M]**

The table below has question-wise data on the performance of students in an examination. The marks for each question are also listed. There is no negative or partial marking in the examination.


What is the average of the marks obtained by the class in the examination?

| Q.N<br>o | Mark<br>s | Answered<br>Correctly | Answered<br>Wrongly | Not<br>Attempted |
|----------|-----------|-----------------------|---------------------|------------------|
| 1        | 2         | 21                    | 17                  | 6                |
| 2        | 3         | 15                    | 27                  | 2                |
| 3        | 2         | 23                    | 18                  | 3                |

- (a) 1.34
- (b) 1.74
- (c) 3.02
- (d) 3.91

**48. [MCQ] [GATE-EE,CS-2014 : 2M]**

The ratio of male to female students in a college for five years is plotted in the following line graph. If the number of female students in 2011 and 2012 is equal, what is the ratio of male students in 2012 to male students in 2011?







49. [NAT]

[GATE-ME,EC-2014: 2M]

A firm producing air purifiers sold 200 units in 2012. The following pie chart presents the share of raw material, labour, energy, plant & machinery, and transportation costs in the total manufacturing cost of the firm in 2012. The expenditure on labour in 2012 is Rs. 4,50,000. In 2013, the raw material expenses increased by 30% and all other expenses increased by 20%. What is the percentage increase in total cost for the company in 2013?




**50. [NAT]**

[GATE-ME,EC-2014: 2M]

A firm producing air purifiers sold 200 units in 2012. The following pie chart presents the share of raw material, labour, energy, plant & machinery, and transportation costs in the total manufacturing cost of the firm in 2012. The expenditure on labour in 2012 is Rs. 4,50,000. In 2013, the raw material expenses increased by 30% and all other expenses increased by 20%. If the company registered a profit of Rs. 10 lakhs


in 2012, at what price (in Rs.) was each air purifier sold?



**51. [MCQ]**

[GATE-ME,EC-2014: 2M]


The multi-level hierarchical pie chart shows the population of animals in a reserve forest. The correct conclusions from this information are:




- (i) Butterflies are birds
  - (ii) There are more tigers in this forest than red ants
  - (iii) All reptiles in this forest are either snakes or crocodiles
  - (iv) Elephants are the largest mammals in this forest
  - (a) (i) and (ii) only
  - (b) (i), (ii), (iii) and (iv)
  - (c) (i), (iii) and (iv) only
  - (d) (i), (ii) and (iii) only

**52. [MCQ]****[GATE-ME,EC-2014: 2M]**


The total exports and revenues from the exports of a country are given in the two charts shown below. The pie chart for exports shows the quantity of each item exported as a percentage of the total quantity of exports. The pie chart for the revenues shows the percentage of the total revenue generated through export of each item. The total quantity of exports of all the items is 500 thousand tonnes and the total revenues are 250 crore rupees. Which item among the following has generated the maximum revenue per kg?



- (a) Item 2      (b) Item 3  
(c) Item 6      (d) Item 5

**53. [MCQ]****[GATE-ME,EC-2014: 2M]**

The exports and imports (in crores of Rs.) of a country from 2000 to 2007 are given in the following bar chart. If the trade deficit is defined as excess of imports over exports, in which year is the trade deficit 1/5th of the exports?



- (a) 2005      (b) 2004  
(c) 2007      (d) 2006

**54. [MCQ]****[GATE-ME,EC-2014: 1M]**


The statistics of runs scored in a series by four batsmen are provided in the following table. Who is the most consistent batsman of these four?

| Batsman | Average | Standard deviation |
|---------|---------|--------------------|
| K       | 31.2    | 5.21               |
| L       | 46.0    | 6.35               |
| M       | 54.4    | 6.22               |
| N       | 17.9    | 5.90               |

- (a) K      (b) L  
(c) M      (d) N

**55. [MCQ]****[GATE-CE-2014: 2M]**

The monthly rainfall chart based on 50 years of rainfall in Agra is shown in the following figure. Which of the following are true? (k percentile is the value such that k percent of the data fall below that value)




- (i) On average, it rains more in July than in December  
(ii) Every year, the amount of rainfall in August is more than that in January  
(iii) July rainfall can be estimated with better confidence than February rainfall  
(iv) In August, there is at least 500 mm of rainfall  
(a) (i) and (ii)      (b) (i) and (iii)  
(c) (ii) and (iii)      (d) (iii) and (iv)

**56. [MCQ]****[GATE-CE-2014: 2M]**

The total exports and revenues from the exports of a country are given in the two pie charts below. The pie chart for exports shows the quantity of each item as a

percentage of the total quantity of exports. The pie chart for the revenues shows the percentage of the total revenue generated through export of each item. The total quantity of exports of all the items is 5 lakh tonnes and the total revenues are 250 crore rupees. What is the ratio of the revenue generated through export of Item 1 per kilogram to the revenue generated through export of Item 4 per kilogram?



- 57. [MCQ] [GATE-CE-2013: 2M]**

Following table provides figures (in rupees) on annual expenditure of a firm for two years – 2010 and 2011.

In 2011, which of the following two categories have registered increase by same percentage?

| Category               | 2010  | 2011  |
|------------------------|-------|-------|
| Raw material           | 5200  | 6240  |
| Power & fuel           | 7000  | 9450  |
| Salary & wages         | 9000  | 12600 |
| Plant & machinery      | 20000 | 25000 |
| Advertising            | 15000 | 19500 |
| Research & Development | 22000 | 26400 |

- (a) Raw material and Salary & wages
  - (b) Salary & wages and Advertising
  - (c) Power & fuel and Advertising
  - (d) Raw material and Research & Development

- 58. [MCQ] [GATE-EE,EC-2012 : 2M]**

The data given in the following table summarizes the monthly budget of an average household.


| <b>Category</b> | <b>Amount (Rs)</b> |
|-----------------|--------------------|
| Food            | 4000               |
| Clothing        | 1200               |
| Rent            | 2000               |
| Savings         | 1500               |
| Other expenses  | 1800               |

The approximate percentage of the monthly budget  
**NOT** spent on savings is



- 59. [MCQ] [GATE-EC,EE-2011 : 2M]**

The fuel consumed by a motorcycle during a journey while traveling at various speeds is indicated in the graph below.



The distances covered during four laps of the journey are listed in the table below

| Lap | Distance<br>(Kilometres) | Average Speed<br>(Kilometres per hour) |
|-----|--------------------------|--|
| P   | 15                       | 15                                     |
| Q   | 75                       | 45                                     |
| R   | 40                       | 75                                     |
| S   | 10                       | 10                                     |


From the given data, we can conclude that the fuel consumed per kilometre was least during the lap



- 60. [MCQ] [GATE-ME,CE,CS-2011: 2M]**

P, Q, R and S are four types of dangerous microbes recently found in a human habitat. The area of each circle with its diameter printed in brackets represents

the growth of a single microbe surviving human immunity system within 24 hours of entering the body. The danger to human beings varies proportionately with the toxicity, potency and growth attributed to a microbe shown in the figure below:




A pharmaceutical company is contemplating the development of a vaccine against the most dangerous microbe. Which microbe should the company target in its first attempt?



## Numerical Computation

- 61. [MCQ] [GATE-EE-2023 : 2M]**


A square with sides of length 6 cm is given. The boundary of the shaded region is defined by two semi-circles whose diameters are the sides of the square, as shown. The area of the shaded region is  $\text{cm}^2$ .



- (a)  $6\pi$
  - (b) 18
  - (c) 20
  - (d)  $9\pi$

- 62. [MCQ]**

Which one of the following options represents the given graph?



- (a)  $f(x) = x^2 2^{-|x|}$       (b)  $f(x) = x 2^{-|x|}$   
 (c)  $f(x) = |x| 2^{-x}$       (d)  $f(x) = x 2^{-x}$

63. [MCQ] [GATE-EE-2023 : 1M]

Given a fair six-faced dice where the faces are labelled ‘1’, ‘2’, ‘3’, ‘4’, ‘5’, and ‘6’, what is the probability of getting a ‘1’ on the first roll of the dice and a ‘4’ on the second roll?

- |                    |                   |
|--------------------|-------------------|
| (a) $\frac{1}{36}$ | (b) $\frac{1}{6}$ |
| (c) $\frac{5}{6}$  | (d) $\frac{1}{3}$ |

64. [MCQ] [GATE-CS-2023:2M]

Consider two function of time ( $t$ )

$$f(t) = 0.01 t^2$$

$$g(t) = 4t$$

Where  $0 < t < \infty$

Now consider the following two statements:

- (i) For some  $t > 0$ ,  $g(t) > f(t)$
  - (ii) There exists a  $T$ , such that  $f(t) > g(t)$  for all  $t > T$

Which one of the following options is TRUE?

- (a) Only (i) is correct
  - (b) Only (ii) is correct
  - (c) Both (i) and (ii) are correct
  - (d) Neither (i) nor (ii) is correct

65. [MCQ] [GATE-CS-2023:2M]

$f(x)$  and  $g(y)$  are functions of  $x$  and  $y$ , respectively, and  $f(x) = g(y)$  for all real values of  $x$  and  $y$ . Which one of the following options is necessarily TRUE for all  $x$  and  $y$ ?

- (a)  $f(x) = 0$  and  $g(y) = 0$
- (b)  $f(x) = g(y) = \text{constant}$
- (c)  $f(x) \neq \text{constant}$  and  $g(y) \neq \text{constant}$
- (d)  $f(x) + g(y) = f(x) - g(y)$

**66. [MCQ] [GATE-ME-2023: 2M]**

How many pairs of sets  $(S, T)$  are possible among the subsets of  $\{1, 2, 3, 4, 5, 6\}$  that satisfy the condition that  $S$  is a subset of  $T$ ?

- (a) 729
- (b) 728
- (c) 665
- (d) 664

**67. [MCQ] [GATE-ME-2023: 2M]**

Consider the following inequalities

$$p^2 - 4q < 4$$

$$3p + 2q < 6$$

Where  $p$  and  $q$  are positive integers.

The value of  $(p + q)$  is \_\_\_\_\_.


- (a) 2
- (b) 1
- (c) 3
- (d) 4

**68. [MCQ] [GATE-CE-2023: 2M]**

A square of side length 4 cm is given. The boundary of the shaded region is defined by one semi-circle on the top and two circular arcs at the bottom each of radius 2 cm, as shown.


The area of the shaded region is \_\_\_\_\_ cm<sup>2</sup>.

- (a) 8
- (b) 4
- (c) 12
- (d) 10


**69. [MCQ] [GATE-CE-2023: 1M]**

In the given figure,  $PQRSTV$  is a regular hexagon with each side of length 5 cm. A circle is drawn with its

centre at  $V$  such that it passes through  $P$ . What is the area (in cm<sup>2</sup>) of the shaded region? (The diagram is representative)



- (a)  $\frac{25\pi}{3}$
- (b)  $\frac{20\pi}{3}$
- (c)  $6\pi$
- (d)  $7\pi$

**70. [MCQ] [GATE-CE-2023: 2M]**

Let  $a = 30!$ ,  $b = 50!$  and  $c = 100!$  Consider the following numbers:  $\log_{ac}a$ ,  $\log_{ca}a$ ,  $\log_ba$ ,  $\log_ba$

Which one of the following inequalities is CORRECT?

- (a)  $\log_{ca}a < \log_{ba}a < \log_{ab}a < \log_{ac}a$
- (b)  $\log_{ca}a < \log_{ab}a < \log_{ba}a < \log_{bc}a$
- (c)  $\log_{ca}a < \log_{ba}a < \log_{ac}a < \log_{ab}a$
- (d)  $\log_{ba}a < \log_{ca}a < \log_{ab}a < \log_{ac}a$

**71. [MCQ] [GATE-EE-2022 : 2M]**

There are two identical dice with a single letter on each of the faces. The following six letters: Q, R, S, T, U, and V, one on each of the faces. Any of the six outcomes are equally likely.


The two dice are thrown once independently at random. What is the probability that the outcomes on the dice were composed only of any combination of the following possible outcomes: Q, U and V?

- (a) 1/4
- (b) 3/4
- (c) 1/6
- (d) 5/36

**72. [MCQ] [GATE-EE-2022 : 1M]**

The figure below shows the front and rear view of a disc, which is shaded with identical patterns. The disc is flipped once with respect to any one of the fixed axes 1-1, 2-2 or 3-3 chosen uniformly at random.

What is the probability that the disc DOES NOT retain the same front and rear views after the flipping operation?



- (a) 0  
(c)  $\frac{2}{3}$

- (b)  $\frac{1}{3}$   
(d) 1

**73. [MCQ]**
**[GATE-CS-2022:1M]**

A function  $y(x)$  is defined in the interval  $[0, 1]$  on the  $x$ -axis as

$$y(x) = \begin{cases} 2 & \text{if } 0 \leq x < \frac{1}{3} \\ 3 & \text{if } \frac{1}{3} \leq x < \frac{3}{4} \\ 1 & \text{if } \frac{3}{4} \leq x \leq 1 \end{cases}$$

Which one of the following is the area under the curve for the interval  $[0, 1]$  on the  $x$ -axis?

- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{5}{6}$  | (b) $\frac{6}{5}$  |
| (c) $\frac{13}{6}$ | (d) $\frac{6}{13}$ |

**74. [MCQ]**
**[GATE-EC-2022:2M]**

Four points P (0, 1), Q (0, -3), R (-2, -1), and S (2, -1) represent the vertices of a quadrilateral.

What is the area enclosed by the quadrilateral?

- |       |                 |
|-------|-----------------|
| (a) 4 | (b) $4\sqrt{2}$ |
| (c) 8 | (d) $8\sqrt{2}$ |

**75. [MCQ]**
**[GATE-EC-2022:2M]**

Consider the following inequalities.

- (i)  $2x - 1 > 7$   
(ii)  $2x - 9 < 1$

Which one of the following expressions below satisfies the above two inequalities?

- |                 |                     |
|-----------------|---------------------|
| (a) $x \leq -4$ | (b) $-4 < x \leq 4$ |
| (c) $4 < x < 5$ | (d) $x \geq 5$      |

**76. [MCQ]**
**[GATE-EC-2022:1M]**

A trapezium has vertices marked as P, Q, R and S (in that order anticlockwise).

The side PQ is parallel to side SR.

Further, it is given that, PQ = 11 cm, QR = 4 cm, RS = 6 cm and SP = 3 cm.

What is the shortest distance between PQ and SR (in cm)?

- |          |          |
|----------|----------|
| (a) 1.80 | (b) 2.40 |
| (c) 4.20 | (d) 5.76 |

**77. [MCQ]**
**[GATE-CE-2022:2M]**

Consider the following equations of straight lines:

Line L1:  $2x - 3y = 5$

Line L2:  $3x + 2y = 8$

Line L3:  $4x - 6y = 5$

Line L4:  $6x - 9y = 6$

Which one among the following is the correct statement?

- |  |
|--|
| (a) L1 is parallel to L2 and L1 is perpendicular to L3 |
| (b) L2 is parallel to L4 and L2 is perpendicular to L1 |
| (c) L3 is perpendicular to L4 and L3 is parallel to L2 |
| (d) L4 is perpendicular to L2 and L4 is parallel to L3 |

**78. [MCQ]**
**[GATE-ME-2022: 2M]**

Consider the following functions for non-zero positive integers,  $p$  and  $q$ .

$$f(p, q) = \underbrace{p \times p \times p \times \dots \times p}_{q \text{ terms}} = p^q; \quad f(p, 1) = p$$

:(up to  $q$  terms)

$$g(p, q) = p^p \times p^p \times \dots \times p^p = p^{pq}; \quad g(p, 1) = p$$

Which one of the following options is correct based on the above?

- |                                     |
|-------------------------------------|
| (a) $f(2, 2) = g(2, 2)$             |
| (b) $f(g(2, 2), 2) < f(2, g(2, 2))$ |
| (c) $h(2, 1) \neq f(2, 1)$          |
| (d) $F(3, 2) > g(3, 2)$             |

**79. [MCQ]**
**[GATE-ME-2022: 1M]**

If  $f(x) = 2\ln(\sqrt{e^x})$  what is the area bounded by  $f(x)$  for the interval  $[0, 2]$  on the  $x$ -axis?


- (a)  $\frac{1}{2}$       (b) 1  
 (c) 2      (d) 4

**80. [MCQ]****[GATE-ME-2022: 1M]**

Which one of the following is a representation (not to scale and in bold) of all values of  $x$  satisfying the

inequality  $2 - 5x \leq -\frac{6x - 5}{3}$  on the real number line?

- (a)
- (b)
- (c)
- (d)

**81. [MCQ]****[GATE-EE-2021 : 1M]**

The number of students passing or failing in an exam for a particular subject are presented in the bar chart above. Students who pass the exam cannot appear for the exam again. Students who fail the exam in the first attempt must appear for the exam in the following year. Students always pass the exam in their second attempt. The number of students who took the exam for the first time in the year 2 and the year 3 respectively, are.

- (a) 65 and 53      (b) 60 and 50  
 (c) 55 and 53      (d) 55 and 48

**82. [MCQ]****[GATE-EE-2021 : 1M]**

Let  $X$  be a continuous random variable denoting the temperature measured. The range of temperature is  $[0, 100]$  degree Celsius and let the probability density function of  $X$  be  $f(x) = 0.01$  for  $0 \leq X \leq 100$ . The mean of  $X$  is \_\_\_\_\_.

- (a) 2.5      (b) 5.0  
 (c) 25.0      (d) 50.0

**83. [MCQ]****[GATE-CS-2021: 1M]**

If  $\theta$  is the angle, in degrees, between the longest diagonal of the cube and any one of the edges of the cube, then,  $\cos \theta =$

- (a)  $\frac{1}{2}$       (b)  $\frac{1}{\sqrt{3}}$   
 (c)  $\frac{1}{\sqrt{2}}$       (d)  $\frac{\sqrt{3}}{2}$

**84. [MCQ]****[GATE-CS-2021: 1M]**

If  $\left(x - \frac{1}{2}\right)^2 - \left(x - \frac{3}{2}\right)^2 = x + 2$ , then the value of  $x$  is

- (a) 2      (b) 4  
 (c) 6      (d) 8


**85. [MCQ]****[GATE-CE-2021: 2M]**

A function  $\lambda$ , is defined by

$$\lambda(p, q) = \begin{cases} (p-q)^2, & \text{if } p \geq q \\ p+q, & \text{if } p < q \end{cases}$$

The value of the expression  $\lambda(-(3+2), (-2+3))/(((-2+1)))$  is:

- (a) -1      (b) 0  
 (c)  $\frac{16}{3}$       (d) 16

**86. [MCQ]****[GATE-EC-2021:2M]**

Corners are cut from an equilateral triangle to produce a regular convex hexagon as shown in the figure above.

The ratio of the area of the regular convex hexagon to the area of the original equilateral triangle is

- (a) 2 : 3      (b) 3 : 4  
 (c) 4 : 5      (d) 5 : 6

## 87. [MCQ]

[GATE-EC-2021:2M]

Consider a square sheet of side 1 unit. In the first step, it is cut along the main diagonal to get two triangles. In the next step, one of the cut triangles is revolved about its short edge to form a solid cone. The volume of the resulting cone, in cubic units, is \_\_\_\_\_

- (a)  $\pi/3$       (b)  $2\pi/3$   
 (c)  $3\pi/2$       (d)  $3\pi$

## 88. [MCQ]

[GATE-EC-2021:1M]

$p$  and  $q$  are positive integers and  $\frac{p}{q} + \frac{q}{p} = 3$ ,

$$\text{then, } \frac{p^2}{q^2} + \frac{q^2}{p^2} =$$

- (a) 3      (b) 7  
 (c) 9      (d) 11

## 89. [MCQ]

[GATE-CE-2021: 1M]

Two identical cube shaped dice each with faces numbered 1 to 6 are rolled simultaneously. The probability that an even number is rolled out on each dice is:


- (a)  $\frac{1}{36}$       (b)  $\frac{1}{8}$   
 (c)  $\frac{1}{12}$       (d)  $\frac{1}{4}$

## 90. [MCQ]

[GATE-CE-2021: 2M]

In the figure shown above, PQRS is a square. The shaded portion is formed by the intersection of sectors of circles with radius equal to the side of the square and centers at S and Q.

The probability that any point picked randomly within the square falls in the shaded area is.




- (a)  $4 - \frac{\pi}{2}$       (b)  $\frac{1}{2}$   
 (c)  $\frac{\pi}{2} - 1$       (d)  $\frac{\pi}{4}$

## 91. [MCQ]

[GATE-EE-2020 : 2M]

Given a semicircle with  $O$  as the centre, as shown in the figure, the ratio  $\frac{\overline{AC} + \overline{CB}}{\overline{AB}}$  is

Where,  $\overline{AC}$ ,  $\overline{AC}$  and  $\overline{AB}$  are chords.



- (a)  $\sqrt{2}$       (b)  $\sqrt{3}$   
 (c) 2      (d) 3

## 92. [MCQ]

[GATE-EE-2020 : 2M]

In four-digit integer numbers from 1001 to 9999, the digit group “37” (in the same sequence) appears \_\_\_\_\_ times.

- (a) 270      (b) 279  
 (c) 280      (d) 299

## 93. [MCQ]

[GATE-EE-2020 : 1M]


If P, Q, R, S are four individuals, how many teams of size exceeding one can be formed, with Q as a member?

- (a) 5      (b) 6  
 (c) 7      (d) 8

## 94. [MCQ]

[GATE-CS-2020: 2M]

The figure below shows an annular ring with outer and inner radii as  $b$  and  $a$ , respectively. The annular space has been painted in the form of blue color circles touching the outer and inner periphery of annular space. If maximum  $n$  number of circles can be painted, then the unpainted area available in annular space is \_\_\_\_.



- (a)  $\pi \left[ (b^2 - a^2) - \frac{n}{4}(b-a)^2 \right]$
- (b)  $\pi \left[ (b^2 - a^2) - n(b-a)^2 \right]$
- (c)  $\pi \left[ (b^2 - a^2) + \frac{n}{4}(b-a)^2 \right]$
- (d)  $\pi \left[ (b^2 - a^2) + n(b-a)^2 \right]$

## 95. [MCQ]

[GATE-CS-2020: 2M]

Two straight lines are drawn perpendicular to each other in X-Yplane. If  $\alpha$  and  $\beta$  are the acute angles the straight lines make with the X-axis, then  $\alpha + \beta$  is \_\_\_\_.

- (a)  $60^\circ$
- (b)  $90^\circ$
- (c)  $120^\circ$
- (d)  $180^\circ$

## 96. [MCQ]

[GATE-EC-2020:2M]


$a, b, c$  are real numbers. The quadratic equation  $ax^2 - bx + c = 0$  has equal roots, which is  $\beta$ , then

- (a)  $\beta = b/a$
- (b)  $\beta^2 = ac$
- (c)  $\beta^3 = bc/(2a^2)$
- (d)  $b^2 \neq 4ac$

## 97. [MCQ]

[GATE-EC-2020:2M]

A circle with centre  $O$  is shown in the figure. A rectangle PQRS of a maximum possible area is inscribed in the circle. If the radius of the circle is  $a$ , then the area of the shaded portion is \_\_\_\_.



- (a)  $\pi a^2 - a^2$
- (b)  $\pi a^2 - \sqrt{2}a^2$
- (c)  $\pi a^2 - 2a^2$
- (d)  $\pi a^2 - 3a^2$

## 98. [MCQ]

[GATE-EC-2020:1M]

A superadditive function  $f(\cdot)$  satisfies the following property

$$f(x_1 + x_2) \geq f(x_1) + f(x_2)$$

Which of the following functions is a superadditive function for  $x > 1$ ?

- (a)  $e^x$
- (b)  $\sqrt{x}$
- (c)  $1/x$
- (d)  $e^{-x}$

## 99. [MCQ]

[GATE-CE-2020: 1M]

If  $f(x) = x^2$  for each  $x \in (\infty, \infty)$  then  $\frac{f(f(f(x)))}{f(x)}$  is equal


to \_\_\_\_.

- (a)  $f(x)$
- (b)  $(f(x))^2$
- (c)  $(f(x))^3$
- (d)  $(f(x))^4$

## 100.[MCQ]

[GATE-ME-2020: 2M]

An engineer measures THREE quantities X, Y and Z in an experiment. She finds that they follow a relationship that is represented in the figure below: (the product of X and Y linearly varies with Z)



Then, which of the following statements is FALSE?

- (a) For fixed Z; X is proportional to Y
- (b) For fixed Y; X is proportional to Z
- (c) For fixed X; Z is proportional to Y
- (d) XY/Z is constant

### 101.[MCQ]

[GATE-ME-2020: 2M]

The sum of the first n terms in the sequence 8, 88, 888, 8888, ..... is

- (a)  $\frac{81}{80}(10^n - 1) + \frac{9}{8}n$
- (b)  $\frac{81}{80}(10^n - 1) - \frac{9}{8}n$
- (c)  $\frac{80}{81}(10^n - 1) + \frac{8}{9}n$
- (d)  $\frac{80}{81}(10^n - 1) - \frac{8}{9}n$

### 102.[MCQ]

[GATE-ME-2020: 1M]

Define  $[x]$  as the greatest integer less than or equal to x, for each  $x \in (-\infty, \infty)$ . If  $y = [x]$ , then area under y for  $x \in [1,4]$  is \_\_\_\_.

- (a) 1
- (b) 3
- (c) 4
- (d) 6

### 103.[MCQ]

[GATE-EE-2019 : 2M]

Given two sets  $X = \{1, 2, 3\}$  and  $Y = \{2, 3, 4\}$ , we construct a set  $Z$  of all possible fractions where the numerators belong to set  $X$  and the denominators belong to set  $Y$ . The product of elements having minimum and maximum values in the set  $Z$  is \_\_\_\_\_.

- (a) 1/12
- (b) 1/8
- (c) 1/6
- (d) 3/8

### 104.[MCQ]

[GATE-CE-2019: 1M]

On a horizontal ground, the base of a straight ladder is 6 m away from the base of a vertical pole. The ladder makes an angle of  $45^\circ$  to the horizontal. If the ladder is resting at a point located at one-fifth of the height of the pole from the bottom, the height of the pole is \_\_\_\_\_ meters.

- (a) 15
- (b) 25
- (c) 30
- (d) 35

### 105.[MCQ]

[GATE-EE-2018 : 2M]

A class of twelve children has two more boys than girls. A group of three children are randomly picked from this class to accompany the teacher on a field trip. What is the probability that the group accompanying the teacher contains more girls than boys?

- (a) 0
- (b)  $\frac{325}{864}$
- (c)  $\frac{525}{864}$
- (d)  $\frac{5}{12}$

### 106.[MCQ]

[GATE-EE-2018 : 2M]

An e-mail password must contain three characters. The password has to contain one numeral from 0 to 9, one upper case and one lower case character from the English alphabet. How many distinct passwords are possible?

- (a) 6,760
- (b) 13,520
- (c) 40,560
- (d) 1,05,456

### 107.[MCQ]

[GATE-EE-2018 : 1M]

Functions  $F(a, b)$  and  $G(a, b)$  are defined as follows:  
 $F(a, b) = (a - b)^2$  and  $G(a, b) = |a - b|$ , where  $|x|$  represents the absolute value of  $x$ . What would be the value of  $G(F(1, 3), G(1, 3))$ ?

- (a) 2
- (b) 4
- (c) 6
- (d) 36

### 108.[MCQ]

[GATE-EE-2018 : 1M]

The three roots of the equation  $f(x) = 0$  are  $x = \{-2, 0, 3\}$ . What are the three values of  $x$  for which  $f(x - 3) = 0$ ?

- (a) -5, -3, 0
- (b) -2, 0, 3
- (c) 0, 6, 8
- (d) 1, 3, 6

**109.[MCQ]****[GATE-CS-2018: 1M]**

The area of a square is  $d$ . What is the area of the circle which has the diagonal of the square as its diameter?

- (a)  $\pi d$       (b)  $\pi d^2$   
 (c)  $\frac{1}{4}\pi d^2$       (d)  $\frac{1}{2}\pi d$


**110.[MCQ]****[GATE-CS-2018: 2M]**

If  $pqr \neq 0$  and  $p^{-x} = \frac{1}{q}$ ,  $q^{-y} = \frac{1}{r}$ ,  $r^{-z} = \frac{1}{p}$ , what is the value of the product  $xyz$ ?

- (a) -1      (b)  $\frac{1}{pqr}$   
 (c) 1      (d)  $pqr$

**111.[MCQ]****[GATE-CS-2018: 2M]**

In the figure below,  $\angle DEC + \angle BFC$  is equal to \_\_\_\_\_.



- (a)  $\angle BCD - \angle BAD$       (b)  $\angle BAD + \angle BCF$   
 (c)  $\angle BAD + \angle BCD$       (d)  $\angle CBA + \angle ADC$

**112.[MCQ]****[GATE-EC-2018:1M]**

A 1.5 m tall person is standing at a distance of 3 m from a lamp post. The light from the lamp at the top of the post casts her shadow. The length of the shadow is twice her height. What is the height of the lamp post in meters?

- (a) 1.5      (b) 3  
 (c) 4.5      (d) 6

**113.[MCQ]****[GATE-CE-2018: 2M]**

$$\underbrace{a+a+a+\dots+a}_{n \text{ time}} = a^2b \text{ and } \underbrace{b+b+b+\dots+b}_{m \text{ time}} = ab^2,$$

where  $a$ ,  $b$ ,  $n$  and  $m$  are natural numbers. What is the value of

$$\left( \underbrace{m+m+m+\dots+m}_{n \text{ time}} \right) \left( \underbrace{n+n+n+\dots+n}_{m \text{ time}} \right)$$

(a)  $2a^2b^2$       (b)  $a^4b^4$   
 (c)  $ab(a+b)$       (d)  $a^2 + b^2$

**114.[MCQ]****[GATE-CE-2018: 1M]**

Tower A is 90 m tall and tower B is 140 m tall. They are 100 m apart. A horizontal skywalk connects the floors at 70 m in both the towers. If a taut rope connects the top of tower A to the bottom of tower B, at what distance (in meters) from tower A will the rope intersect the skywalk?

- (a) 22.22      (b) 50  
 (c) 57.87      (d) 77.78

**115.[MCQ]****[GATE-CE-2018: 2M]**

For non-negative integers,  $a$ ,  $b$ ,  $c$ , what would be the value of  $a + b + c$  if  $\log a + \log b + \log c = 0$ ?

- (a) 3      (b) 1  
 (c) 0      (d) -1

**116.[MCQ]****[GATE-CE-2018: 2M]**

Given that  $\frac{\log P}{y-z} = \frac{\log Q}{z-x} = \frac{\log R}{x-y} = 10$  for  $x \neq y \neq z$

what is the value of the product  $PQR$ ?

- (a) 0      (b) 1  
 (c)  $xyz$       (d)  $10^{xyz}$

**117.[MCQ]****[GATE-ME-2018: 1M]**

The value of the expression


$$\frac{1}{1+\log_u vw} + \frac{1}{1+\log_v wu} + \frac{1}{1+\log_w uw} \text{ is } \underline{\hspace{2cm}}.$$

- (a) -1      (b) 0  
 (c) 1      (d) 3

**118.[MCQ]**

[GATE-ME-2018: 2M]

Which of the following functions describe the graph shown in the below figure?



- (a)  $y = ||x| + 1| - 2$       (b)  $y = ||x| - 1| - 1$   
 (c)  $y = ||x| + 1| - 1$       (d)  $y = ||x - 1| - 1|$

**119.[MCQ]**

[GATE-ME-2018: 2M]

For integers  $a$ ,  $b$  and  $c$ , what would be the minimum and maximum values respectively of

- $a + b + c$  if  $\log |a| + \log |b| + \log |c| = 0$ ?  
 (a) -3 and 3      (b) -1 and 1  
 (c) -1 and 3      (d) 1 and 3

**120.[MCQ]**

[GATE-EE,CE-2017 : 2M]

The expression  $\frac{(x+y)-|x-y|}{2}$  is equal to

- (a) The maximum of  $x$  and  $y$   
 (b) The minimum of  $x$  and  $y$   
 (c) 1  
 (d) None of the above

**121.[MCQ]**

[GATE-EE,CS-2017 : 2M]

The number of roots of  $e^x + 0.5x^2 - 2 = 0$  in the range  $[-5, 5]$  is

- (a) 0      (b) 1  
 (c) 2      (d) 3

**122.[MCQ]**

[GATE-EC-2017:2M]

The number of 3-digit numbers such that the digit 1 is never to the immediate right of 2 is

- (a) 781      (b) 791  
 (c) 881      (d) 891

**123.[MCQ]**

[GATE-EC-2017:2M]

There are 3 Indians and 3 Chinese in a group of 6 people. How many subgroups of this group can we choose so that every subgroup has at least one Indian?

- (a) 56      (b) 52  
 (c) 48      (d) 44

**124.[MCQ]**

[GATE-CE-2017: 1M]

Two dice are thrown simultaneously. The probability that the product of the numbers appearing on the top faces of the dice is a perfect square is

- (a)  $\frac{1}{9}$       (b)  $\frac{2}{9}$   
 (c)  $\frac{1}{3}$       (d)  $\frac{4}{9}$

**125.[MCQ]**

[GATE-ME-2017: 2M]


Let  $S_1$  be the plane figure consisting of the points  $(x, y)$  given by the inequalities  $|x - 1| \leq 2$  and  $|y + 2| \leq 3$ . Let  $S_2$  be the plane figure given by the inequalities  $x - y \geq -2$ ,  $y \geq 1$ , and  $x \leq 3$ . Let  $S$  be the union of  $S_1$  and  $S_2$ . The area of  $S$  is

- (a) 26      (b) 28  
 (c) 32      (d) 34

**126.[MCQ]**

[GATE-ME-2017: 1M]

A right-angled cone (with base radius 5 cm and height 12 cm), as shown in the figure below, is rolled on the ground keeping the point P fixed until the point Q (at the base of the cone, as shown) touches the ground again.



By what angle (in radians) about P does the cone travel?

- (a)  $\frac{5\pi}{12}$       (b)  $\frac{5\pi}{24}$   
 (c)  $\frac{24\pi}{5}$       (d)  $\frac{10\pi}{13}$

**127.[NAT]****[GATE-EE-2016 : 2M]**

The numeral in the units position of  $211^{870} + 146^{127} \times 3^{424}$  is \_\_\_\_\_.

**128.[MCQ]****[GATE-EE-2016 : 2M]**

Shaquille O' Neal is a 60% career free throw shooter, meaning that he successfully makes 60 free throws out of 100 attempts on average. What is the probability that he will successfully make exactly 6 free throws in 10 attempts?

- |            |            |
|------------|------------|
| (a) 0.2508 | (b) 0.2816 |
| (c) 0.2934 | (d) 0.6000 |

**129.[MCQ]****[GATE-EE-2016 : 1M]**

If  $|9y - 6| = 3$ , then  $y^2 - 4y/3$  is \_\_\_\_\_.

- |            |               |
|------------|---------------|
| (a) 0      | (b) $+1/3$    |
| (c) $-1/3$ | (d) undefined |

**130.[MCQ]****[GATE-EE,CS-2016 : 1M]**

In a quadratic function, the value of the product of the roots ( $\alpha, \beta$ ) is 4. Find the value of

$$\frac{\alpha^n + \beta^n}{\alpha^{-n} + \beta^{-n}}$$

- |                |               |
|----------------|---------------|
| (a) $n^4$      | (b) $4^n$     |
| (c) $2^{2n-1}$ | (d) $4^{n-1}$ |

**131.[MCQ]****[GATE-CS,CE-2016: 2M]**

If  $f(x) = 2x^7 + 3x - 5$ , which of the following is a factor of  $f(x)$ ?

- |                 |               |
|-----------------|---------------|
| (a) $(x^3 + 8)$ | (b) $(x - 1)$ |
| (c) $(2x - 5)$  | (d) $(x + 1)$ |

**132.[MCQ]****[GATE-CS,CE-2016: 2M]**

In a process, the number of cycles to failure decreases exponentially with an increase in load. At a load of 80 units, it takes 100 cycles for failure. When the load is halved, it takes 10000 cycles for failure. The load for which the failure will happen in 5000 cycles is \_\_\_\_\_.

- |           |           |
|-----------|-----------|
| (a) 40.00 | (b) 46.02 |
| (c) 60.01 | (d) 92.02 |

**133.[MCQ]****[GATE-EC-2016:2M]**

A straight line is fit to a data set  $(\ln x, y)$ . This line intercepts the abscissa at  $\ln x = 0.1$  and has a slope of  $-0.02$ . What is the value of  $y$  at  $x = 5$  from the fit?

- |            |            |
|------------|------------|
| (a) -0.030 | (b) -0.014 |
| (c) 0.014  | (d) 0.030  |

**134.[MCQ]****[GATE-EC-2016:2M]**

Find the area bounded by the lines  $3x + 2y = 14$ ,  $2x - 3y = 5$  in the first quadrant.

- |           |           |
|-----------|-----------|
| (a) 14.95 | (b) 15.25 |
| (c) 15.70 | (d) 20.23 |

**135. [MCQ]****[GATE-ME,EC-2016: 2M]**

If  $q^{-a} = \frac{1}{r}$  and  $r^{-b} = \frac{1}{s}$  and  $s^{-c} = \frac{1}{q}$ , the value of abc is \_\_\_\_\_.

- |                  |                 |
|------------------|-----------------|
| (a) $(rqs)^{-1}$ | (b) 0           |
| (c) 1            | (d) $r + q + s$ |

**136.[MCQ]****[GATE-CE-2016: 2M]**

A square pyramid has a base perimeter x, and the slant height is half of the perimeter. What is the lateral surface area of the pyramid?

- |                |                |
|----------------|----------------|
| (a) $x^2$      | (b) $0.75 x^2$ |
| (c) $0.50 x^2$ | (d) $0.25 x^2$ |

**137.[MCQ]****[GATE-EE,CS-2015 : 1M]**

Given Set A = {2, 3, 4, 5} and Set B = {11, 12, 13, 14, 15}, two numbers are randomly selected, one from each set. What is the probability that the sum of the two numbers equals 16?

- |          |          |
|----------|----------|
| (a) 0.20 | (b) 0.25 |
| (c) 0.30 | (d) 0.33 |

**138.[MCQ]****[GATE-EE,CS-2015 : 1M]**

Consider a function  $f(x) = 1 - |x|$  on  $-1 \leq x \leq 1$ . The value of  $x$  at which the function attains a maximum, and the maximum value of the function are:

- |           |           |
|-----------|-----------|
| (a) 0, -1 | (b) -1, 0 |
| (c) 0, 1  | (d) -1, 2 |



**148.[MCQ]****[GATE-EE,CS-2014 : 1M]**

The roots of  $ax^2 + bx + c = 0$  are real and positive. a, b and c are real, Then  $ax^2 + b|x| + c = 0$  has

- |                  |                  |
|------------------|------------------|
| (a) No roots     | (b) 2 real roots |
| (c) 3 real roots | (d) 4 real roots |

**149.[MCQ]****[GATE-EE,CS-2014 : 2M]**

If x is real and  $|x^2 - 2x + 3| = 11$ , then possible values of  $-x^3 + x^2 - x$  include

- |           |            |
|-----------|------------|
| (a) 2, 4  | (b) 2, 14  |
| (c) 4, 52 | (d) 14, 52 |

**150.[MCQ]****[GATE-EE,CS-2014 : 2M]**

Consider the equation:  $(7526)_8 - (Y)_8 = (4364)_8$ , where  $(X)_N$  stands for X to the base N. Find Y.

- |          |          |
|----------|----------|
| (a) 1634 | (b) 1737 |
| (c) 3142 | (d) 3162 |

**151.[MCQ]****[GATE-CE-2014: 1M]**

If  $y = 5x^2 + 3$ , then the tangent at  $x = 0$ ,  $y = 3$

- |                                      |
|--------------------------------------|
| (a) passes through $x = 0$ , $y = 0$ |
| (b) has a slope of +1                |
| (c) is parallel to the x-axis        |
| (d) has a slope of -1                |

**152.[NAT]****[GATE-CE-2014: 2M]**

The smallest angle of a triangle is equal to two thirds of the smallest angle of a quadrilateral. The ratio between the angles of the quadrilateral is 3:4:5:6. The largest angle of the triangle is twice its smallest angle. What is the sum, in degrees, of the second largest angle of the triangle and the largest angle of the quadrilateral?

**153.[NAT]****[GATE-CE-2014: 1M]**

In any given year, the probability of an earthquake greater than Magnitude 6 occurring in the Garhwal Himalayas is 0.04. The average time between successive occurrences of such earthquakes is \_\_\_\_ years.

**154.[MCQ]****[GATE-ME,EC-2014: 1M]**

Let  $f(x, y) = x^n y^m = P$ . If x is doubled and y is halved, the new value of f is

- |                 |                 |
|-----------------|-----------------|
| (a) $2^{n-m} P$ | (b) $2^{m-n} P$ |
| (c) $2(n-m)P$   | (d) $2(m-n)P$   |

**155.[MCQ]****[GATE-EE,EC-2013 : 2M]**

What is the chance that a leap year, selected at random, will contain 53 Saturdays?

- |           |           |
|-----------|-----------|
| (a) $2/7$ | (b) $3/7$ |
| (c) $1/7$ | (d) $5/7$ |

**156.[MCQ]****[GATE-EE,EC-2013 : 2M]**

The set of values of p for which the roots of the equation  $3x^2 + 2x + p(p-1) = 0$  are of opposite sign is

- |                    |                   |
|--------------------|-------------------|
| (a) $(-\infty, 0)$ | (b) $(0, 1)$      |
| (c) $(1, \infty)$  | (d) $(0, \infty)$ |

**157.[MCQ]****[GATE-EE,EC-2013 : 2M]**

Find the sum to n terms of the series

$$10 + 84 + 734 + \dots$$

- |                                 |                                  |
|---------------------------------|----------------------------------|
| (a) $\frac{9(9^n + 1)}{10} + 1$ | (b) $\frac{9(9^n - 1)}{8} + 1$   |
| (c) $\frac{9(9^n - 1)}{8} + n$  | (d) $\frac{9(9^n - 1)}{8} + n^2$ |

**158.[MCQ]****[GATE-CE-2013: 2M]**

X and Y are two positive real numbers such that  $2X + Y \leq 6$  and  $X + 2Y \leq 8$ . For which of the following values of (X, Y) the function  $f(X, Y) = 3X + 6Y$  will give maximum value?

- |                   |                   |
|-------------------|-------------------|
| (a) $(4/3, 10/3)$ | (b) $(8/3, 20/3)$ |
| (c) $(8/3, 10/3)$ | (d) $(4/3, 20/3)$ |

**159.[MCQ]****[GATE-CE-2013: 2M]**

If  $|4X - 7| = 5$  then the values of  $2|X| - |X|$  is:

- |                      |                      |
|----------------------|----------------------|
| (a) $2, \frac{1}{3}$ | (b) $\frac{1}{2}, 3$ |
| (c) $\frac{3}{2}, 9$ | (d) $\frac{2}{3}, 9$ |



**171.[MCQ]****[GATE-CS-2023:1M]**

A series of natural numbers  $F_1, F_2, F_3, F_4, F_5, F_6, F_7, \dots$  obeys

$F_{n+1} = F_n + F_{n-1}$  for all integers  $n \geq 2$ .

If  $F_6 = 37$ , and  $F_7 = 60$ , then what is  $F_1$ ?

- |       |       |
|-------|-------|
| (a) 4 | (b) 5 |
| (c) 8 | (d) 9 |

**172.[MCQ]****[GATE-CS-2023:1M]**

A survey for a certain year found that 90% of pregnant women received medical care at least once before giving birth. Of these women, 60% received medical care from doctors, while 40% received medical care from other healthcare providers. Given this information, which one of the following statements can be inferred with certainty?

- (a) More than half of the pregnant women received medical care at least once from a doctor.
- (b) Less than half of the pregnant women received medical care at least once from a doctor.
- (c) More than half of the pregnant women received medical care at most once from a doctor.
- (d) Less than half of the pregnant women received medical care at most once from a doctor.

**173.[MCQ]****[GATE-EC-2023:2M]**

Out of 1000 individuals in a town, 100 unidentified individuals are covid positive. Due to lack of adequate covid-testing kits, the health authorities of the town devised a strategy to identify these covid-positive individuals. The strategy is to:

- (i) Collect saliva samples from all 1000 individuals and randomly group them into sets of 5.
- (ii) Mix the samples within each set and test the mixed sample for covid.
- (iii) If the test done in (ii) gives a negative result, then declare all the 5 individuals to be covid negative.
- (iv) If the test done in (ii) gives a positive result, then all the 5 individuals are separately tested for covid.

Given this strategy, no more than \_\_\_\_\_ testing kits will be required to identify all the 100 covid positive individuals irrespective of how they are grouped.

- (a) 700

- (b) 600

- (c) 800

- (d) 1000


**174.[MCQ]****[GATE-EC-2023:1M]**

What is the smallest number with distinct digits whose digits add up to 45?

- |               |               |
|---------------|---------------|
| (a) 123555789 | (b) 123457869 |
| (c) 123456789 | (d) 99999     |

**175.[MCQ]****[GATE-CE-2023: 1M]**

In how many ways can cells in a  $3 \times 3$  grid be shaded, such that each row and column have exactly one shaded cell? An example of one valid shading is shown.



- (a) 2

- (b) 9

- (c) 3

- (d) 6

**176.[MCQ]****[GATE-CE-2023: 2M]**

Three husband-wife pairs are to be seated at a circular table that has six identical chairs. Seating arrangements are defined only by the relative position of the people. How many seating arrangements are possible such that every husband sits next to his wife?

- |         |         |
|---------|---------|
| (a) 16  | (b) 4   |
| (c) 120 | (d) 720 |

**177.[MCQ]****[GATE-CE-2023: 2M]**

If  $x$  satisfies the equation  $4^8 = 256$ , then  $x$  is equal to

$$\frac{1}{2}$$

- (b)  $\log_{16}8$

$$\frac{2}{3}$$

- (d)  $\log_48$


**178.[MCQ]****[GATE-ME-2023: 1M]**

The minute-hand and second-hand of a clock cross each other \_\_\_\_\_ times between 09:15:00 AM and 09:45:00 AM on a day.



**187.[MCQ]****[GATE-CE-2022:1M]**

In the following diagram, the point R is the center of the circle. The lines PQ and ZV are tangential to the circle. The relation among the areas of the squares, PXWR, RUVZ and SPQT is.



- (a) Area of SPQT = Area of RUVZ = Area of PXWR
- (b) Area of SPQT = Area of PXWR – Area of RUVZ
- (c) Area of PXWR = Area of SPQT – Area of RUVZ
- (d) Area of PXWR = Area of RUVZ – Area of SPQT

**188.[MCQ]****[GATE-CE-2022:2M]**

P invested ₹ 5000 per month for 6 months of a year and Q invested ₹  $x$  per month for 8 months of the year in a partnership business. The profit is shared in proportion to the total investment made in that year.

If at the end of that investment year, Q receives  $\frac{4}{9}$  of the total profit, what is the value of  $x$  (in ₹)?

- (a) 2500
- (b) 3000
- (c) 4687
- (d) 8437

**189.[MCQ]****[GATE-CE-2022:2M]**

In a partnership business the monthly investment by three friends for the first six months is in the ratio 3 : 4 : 5. After six months, they had to increase their monthly investments by 10%, 15% and 20%, respectively, of their initial monthly investment. The new investment ratio was kept constant for the next six months.

What is the ratio of their shares in the total profit (in the same order) at the end of the year such that the share is proportional to their individual total investment over the year?

- (a) 22 : 23 : 24
- (b) 22 : 33 : 50
- (c) 33 : 46 : 60
- (d) 63 : 86 : 110

**190.[MCQ]****[GATE-CE-2022:1M]**

Both the numerator and the denominator of  $\frac{3}{4}$  are increased by a positive integer,  $x$ , and those of  $\frac{15}{17}$  are decreased by the same integer. This operation results in the same value for both the fractions.


What is the value of  $x$ ?

- (a) 1
- (b) 2
- (c) 3
- (d) 4

**191.[MCQ]****[GATE-ME-2022: 2M]**

Equal sized circular regions are shaded in a square sheet of paper of 1 cm side length. Two cases, case M and case N, are considered as shown in the figures below. In the case M, four circles are shaded in the square sheet and in the case N, nine circles are shaded in the square sheet as shown.

What is the ratio of the areas of unshaded regions of case M to that of case N?



Case M      Case N

- (a) 2 : 3
- (b) 1 : 1
- (c) 3 : 2
- (d) 2 : 1

**192.[MCQ]****[GATE-ME-2022: 2M]**

For the past  $m$  days, the average daily production at a company was 100 units per day.

If today's production of 180 units changes the average to 110 units per day, what is the value of  $m$ ?

- (a) 18
- (b) 10
- (c) 7
- (d) 5

**193.[MCQ]****[GATE-ME-2022: 1M]**

A person was born on the fifth Monday of February in a particular year.

Which one of the following statements is correct based on the above information?



**202.[MCQ]****[GATE-CS-2021: 1M]**

The ratio of boys to girls in a class is 7 to 3.

Among the options below, an acceptable value for the total number of students in the class is:

- |        |        |
|--------|--------|
| (a) 21 | (b) 37 |
| (c) 50 | (d) 73 |

**203.[MCQ]****[GATE-CS-2021: 2M]**

We have 2 rectangular sheets of paper. M and N. of dimensions 6 cm x 1 cm each. Sheet M is rolled to form an open cylinder by bringing the short edges of the sheet together. Sheet N is cut into equal square patches and assembled to form the largest possible closed cube. Assuming the ends of the cylinder are closed, the ratio of the volume of the cylinder to that of the cube is\_\_\_\_\_.

- |                     |                     |
|---------------------|---------------------|
| (a) $\frac{\pi}{2}$ | (b) $\frac{3}{\pi}$ |
| (c) $\frac{9}{\pi}$ | (d) $3\pi$          |

**204.[MCQ]****[GATE-CS-2021: 2M]**

Details of prices of two items P and Q are presented in the above table. The ratio of cost of item P to cost of item Q is 3:4. Discount is calculated as the difference between the marked price and the selling price. The profit percentage is calculated as the ratio of the difference between selling price and cost, to the cost

| Items | Cost Rs. | Profit% | Marked Price Rs. |
|-------|----------|---------|------------------|
| P     | 5,400    | ....    | 5,860            |
| Q     | 7200     | 25      | 10,000           |

$$\text{profit \%} = (\text{Selling price} - \text{cost})/\text{cost} \times 100$$

The discount on item Q, as a percentage of its marked price is\_\_\_\_\_.

- |        |          |
|--------|----------|
| (a) 25 | (b) 12.5 |
| (c) 10 | (d) 5    |

**205.[MCQ]****[GATE-CS-2021: 2M]**

There are five bags each containing identical sets of ten distinct chocolates. One chocolate is picked from each bag.

The probability that at least two chocolates are identical is \_\_\_\_\_.

- |            |            |
|------------|------------|
| (a) 0.3024 | (b) 0.4235 |
| (c) 0.6976 | (d) 0.8125 |

**206.[MCQ]****[GATE-CS-2021: 2M]**

The number of students in three classes is in the ratio 3:13:6. If 18 students are added to each class, the ratio changes to 15:35:21.


The total number of students in all the three classes in the beginning was:

- |        |         |
|--------|---------|
| (a) 22 | (b) 66  |
| (c) 88 | (d) 110 |

**207.[MCQ]****[GATE-EC-2021:1M]**

The current population of a city is 11,02,500. If it has been increasing at the rate of 5% per annum, what was its population 2 years ago?

- |               |               |
|---------------|---------------|
| (a) 9,92,500  | (b) 9,95,006  |
| (c) 10,00,000 | (d) 12,51,506 |

**208.[MCQ]****[GATE-CE-2021: 2M]**

Five-line segments of equal lengths, PR, PS, QS, QT and RT are used to form a star as shown in the figure above.

The value of  $\theta$ , in degrees is\_\_\_\_\_.

- |        |         |
|--------|---------|
| (a) 36 | (b) 45  |
| (c) 72 | (d) 108 |

**209.[MCQ]****[GATE-CE-2021: 1M]**

Four persons P, Q, R and S are to be seated in a row. R should not be seated at the second position from the left end of the row. The number of distinct seating arrangements possible is:

- |        |        |
|--------|--------|
| (a) 6  | (b) 9  |
| (c) 18 | (d) 24 |

**210.[MCQ]****[GATE-CE-2021: 2M]**


In an equilateral triangle PQR, side PQ is divided into four equal parts, side QR is divided into six equal parts and side PR is divided into eight equal parts. The length of each subdivided part in cm is an integer.

The minimum area of the triangle PQR possible, in  $\text{cm}^2$ , is

- |                  |                   |
|------------------|-------------------|
| (a) 18           | (b) 24            |
| (c) $48\sqrt{3}$ | (d) $144\sqrt{3}$ |

**211.[MCQ]****[GATE-ME-2021: 2M]**

The ratio of the area of the inscribed circle to the area of the circumscribed circle of an equilateral triangle is



- |                   |                   |
|-------------------|-------------------|
| (a) $\frac{1}{8}$ | (b) $\frac{1}{6}$ |
| (c) $\frac{1}{4}$ | (d) $\frac{1}{2}$ |

**212.[MCQ]****[GATE-ME-2021: 2M]**

A box contains 15 blue balls and 45 black balls. If 2 balls are selected randomly, without replacement, the probability of an outcome in which the first selected is a blue ball and the second selected is a black ball, is

- .
- |                    |                      |
|--------------------|----------------------|
| (a) $\frac{3}{16}$ | (b) $\frac{45}{236}$ |
| (c) $\frac{1}{4}$  | (d) $\frac{3}{4}$    |

**213.[MCQ]****[GATE-ME-2021: 1M]**

A digital watch X beeps every 30 seconds while watch Y beeps every 32 seconds. They beeped together at 10 AM. The immediate next time that they will beep together is

- |              |              |
|--------------|--------------|
| (a) 10.08 AM | (b) 10.42 AM |
| (c) 11.00 AM | (d) 10.00 PM |

**214.[MCQ]****[GATE-ME-2021: 1M]**

Five persons P, Q, R, S and T are to be seated in a row, all facing the same direction, but not necessarily in the same order. P and T cannot be seated at either end of the row. P should not be seated adjacent to S. R is to be seated at the second position from the left end of the row. The number of distinct seating arrangements possible is:

- |       |       |
|-------|-------|
| (a) 2 | (b) 3 |
| (c) 4 | (d) 5 |

**215.[MCQ]****[GATE-ME-2021: 2M]**

Five persons P, Q, R, S and T are sitting in a row not necessarily in the same order. Q and R are separated by one person, and S should not be seated adjacent to Q.

The number of distinct seating arrangements possible is:

- |        |        |
|--------|--------|
| (a) 4  | (b) 8  |
| (c) 10 | (d) 16 |

**216.[MCQ]****[GATE-ME-2021: 2M]**


The number of hens, ducks and goats in farm P are 65, 91 and 169, respectively. The total number of hens, ducks and goats in a nearby farm Q is 416. The ratio of hens: ducks: goats in farm Q is 5:14:13. All the hens, ducks and goats are sent from farm Q to farm P.

The new ratio of hens: ducks: goats in farm P is

- .
- |              |              |
|--------------|--------------|
| (a) 5:7:13   | (b) 5:14:13  |
| (c) 10:21:26 | (d) 21:10:26 |

**217.[MCQ]****[GATE-ME-2021: 1M]**

In the above figure, O is the center of the circle and, M and N lie on the circle.



The area of the right triangle MON is  $50 \text{ cm}^2$ . What is the area of the circle in  $\text{cm}^2$ ?

- (a)  $2\pi$       (b)  $50\pi$   
 (c)  $75\pi$       (d)  $100\pi$

**218.[MCQ]****[GATE-EE-2020 : 2M]**

Non-performing Assets (NPAs) of a bank in India is defined as an asset, which remains unpaid by a borrower for a certain period of time in terms of interest, principal, or both. Reserve Bank of India (RBI) has changed the definition of NPA thrice during 1993-2004, in terms of the holding period of loans. The holding period was reduced by one quarter each time. In 1993, the holding period was four quarters (360 days).


Based on the above paragraph, the holding period of loans in 2004 after the third revision was \_\_\_\_\_ days.

- (a) 45      (b) 90  
 (c) 135      (d) 180

**219.[MCQ]****[GATE-CS-2020: 1M]**

There are multiple routes to reach from node 1 to node 2, as shown in the network.

The cost of travel on an edge between two nodes is given in rupees. Nodes 'a', 'b', 'c', 'd', 'e' and 'f' are toll booths. The toll price at toll booths marked 'a' and 'e' is Rs. 200, and is Rs. 100 for the other toll booths. Which is the cheapest route from node 1 to node 2?



- (a) 1-a-c-2  
 (c) 1-b-2

- (b) 1-f-b-2  
 (d) 1-f-e-2

**220.[MCQ]****[GATE-EC-2020:2M]**

It is quarter past three in your watch. The angle between the hour hand and the minute hand is

- (a)  $0^\circ$       (b)  $7.5^\circ$   
 (c)  $15^\circ$       (d)  $22.5^\circ$

**221.[MCQ]****[GATE-CE-2020: 1M]**

The sum of two positive numbers is 100. After subtracting 5 from each number, the product of the resulting numbers is 0. One of the original numbers is.

- (a) 80      (b) 85  
 (c) 90      (d) 95

**222.[MCQ]****[GATE-CE-2020: 1M]**

The ratio of 'the sum of the odd positive integers from 1 to 100' to 'the sum of the even positive integers from 150 to 200' is \_\_\_\_\_.

- (a) 45 : 95      (b) 1 : 2  
 (c) 50 : 91      (d) 1 : 1

**223.[MCQ]****[GATE-CE-2020: 2M]**

For the year 2019, which of the previous year's calendar can be used?

- (a) 2011      (b) 2012  
 (c) 2013      (d) 2014

**224.[MCQ]****[GATE-CE-2020: 2M]**

The unit's place in  $26591749^{110016}$  is \_\_\_\_\_.

- (a) 1      (b) 3  
 (c) 6      (d) 9



complete the same order. If both machines work simultaneously at their respective constant rates, the time taken to complete the same order is \_\_\_\_\_ hours.

- |                   |                   |
|-------------------|-------------------|
| (a) $\frac{2}{3}$ | (b) $\frac{3}{4}$ |
| (c) $\frac{4}{3}$ | (d) $\frac{7}{3}$ |

**235.[MCQ]****[GATE-CE-2019: 2M]**

An oil tank can be filled by pipe X in 5 hours and pipe Y in 4 hours, each pump working on its own. When the oil tank is full and the drainage hole is open, the oil is drained in 20 hours. If initially the tank was empty and someone started the two pumps together but left the drainage hole open, how many hours will it take for the tank to be filled? (Assume that the rate of drainage is independent of the Head)

- |          |          |
|----------|----------|
| (a) 1.50 | (b) 2.00 |
| (c) 2.50 | (d) 4.00 |

**236.[MCQ]****[GATE-CE-2019: 2M]**

A square has sides 5 cm smaller than the sides of a second square. The area of the larger square is four times the area of the smaller square. The side of the larger square is \_\_\_\_\_ cm.

- |           |           |
|-----------|-----------|
| (a) 18.50 | (b) 15.10 |
| (c) 10.00 | (d) 8.50  |

**237.[MCQ]****[GATE-CE-2019: 1M]**

Suresh wanted to lay a new carpet in his new mansion with an area of  $70 \times 55$  sq. mts. However, an area of 550 sq. mts. had to be left out for flower pots. If the cost of carpet is Rs. 50 per sq. mts. how much money (in Rs.) will be spent by Suresh for the carpet now?

- |                 |                 |
|-----------------|-----------------|
| (a) Rs.1,65,000 | (b) Rs.1,92,500 |
| (c) Rs.2,75,000 | (d) Rs.1,27,500 |

**238.[MCQ]****[GATE-CE-2019: 1M]**

A retaining wall with measurements  $30\text{ m} \times 12\text{ m} \times 6\text{ m}$  was constructed with bricks of dimensions  $8\text{ cm} \times 6\text{ cm} \times 6\text{ cm}$ . If 60% of the wall consists of bricks, the number of bricks used for the construction is \_\_\_\_\_ lakhs.

- |        |        |
|--------|--------|
| (a) 30 | (b) 40 |
| (c) 45 | (d) 75 |

**239.[MCQ]****[GATE-CE-2019: 2M]**

Mohan, the manager, wants his four workers to work in pairs. No pair should work for more than 5 hours. Ram and John have worked together for 5 hours. Krishna and Amir have worked as a team for 2 hours. Krishna does not want to work with Ram. Whom should Mohan allot to work with John, if he wants all the workers to continue working?

- |          |                       |
|----------|-----------------------|
| (a) Amir | (b) Krishna           |
| (c) Ram  | (d) None of the three |

**240.[MCQ]****[GATE-CE-2019: 2M]**

Population of state X increased by  $x\%$  and the population of state Y increased by  $y\%$  from 2001 to 2011. Assume that  $x$  is greater than  $y$ . Let  $P$  be the ratio of the population of state X to state Y in a given year. The percentage increase in  $P$  from 2001 to 2011 is.

- |                                  |                                  |
|----------------------------------|----------------------------------|
| (a) $\frac{x}{y}$                | (b) $y - x$                      |
| (c) $\frac{100(x - y)}{100 + x}$ | (d) $\frac{100(x - y)}{100 + y}$ |

**241.[MCQ]****[GATE-ME-2019: 2M]**

Two pipes P and Q can fill a tank in 6 hours and 9 hours respectively, while a third pipe R can empty the tank in 12 hours. Initially, P and R are open for 4 hours. Then P is closed and Q is opened. After 6 more hours R is closed. The total time taken to fill the tank (in hours) is \_\_\_\_\_.

- |           |           |
|-----------|-----------|
| (a) 13.50 | (b) 14.50 |
| (c) 15.50 | (d) 16.50 |

**242.[MCQ]****[GATE-ME-2019: 2M]**

Fiscal deficit was 4% of the GDP in 2015 and that increased to 5% in 2016. If the GDP increased by 10% from 2015 to 2016, the percentage increase in the actual fiscal deficit is \_\_\_\_\_.

- |           |           |
|-----------|-----------|
| (a) 37.50 | (b) 35.70 |
| (c) 25.00 | (d) 10.00 |

**243.[MCQ]****[GATE-ME-2019: 1M]**

The product of three integers X, Y and Z is 192. Z is equal to 4 and P is equal to the average of X and Y. What is the minimum possible value of P?

- |       |         |
|-------|---------|
| (a) 6 | (b) 7   |
| (c) 8 | (d) 9.5 |

**244.[MCQ]****[GATE-ME-2019: 2M]**

A person divided an amount of Rs. 100,000 into two parts and invested in two different schemes. In one he got 10% profit and in the other he got 12%. If the profit percentages are interchanged with these investments he would have got Rs.120 less. Find the ratio between his investments in the two schemes.

- |             |             |
|-------------|-------------|
| (a) 9 : 16  | (b) 11 : 14 |
| (c) 37 : 63 | (d) 47 : 53 |

**245.[MCQ]****[GATE-ME-2019: 1M]**

The sum and product of two integers are 26 and 165 respectively. The difference between these two integers is .

- |       |       |
|-------|-------|
| (a) 2 | (b) 3 |
| (c) 4 | (d) 6 |

**246.[MCQ]****[GATE-ME-2019: 1M]**

A worker noticed that the hour hand on the factory clock had moved by 225 degrees during her stay at the factory. For how long did she stay in the factory?

- |                |                         |
|----------------|-------------------------|
| (a) 3.75 hours | (b) 4 hours and 15 mins |
| (c) 8.5 hours  | (d) 7.5 hours           |

**247.[MCQ]****[GATE-EE-2018 : 1M]**

For what values of  $k$  given below is  $\frac{(k+2)^2}{k-3}$  an integer?

- |              |               |
|--------------|---------------|
| (a) 4, 8, 18 | (b) 4, 10, 16 |
| (c) 4, 8, 28 | (d) 8, 26, 28 |

**248.[MCQ]****[GATE-CS-2018: 1M]**

What would be the smallest natural number which when divided either by 20 or by 42 or by 76 leaves a remainder of 7 in each case?

- (a) 3047

- (c) 7987

- (b) 6047

- (d) 63847

**249.[MCQ]****[GATE-CS-2018: 2M]**

In a party, 60% of the invited guests are male and 40% are female. If 80% of the invited guests attended the party and if all the invited female guests attended, what would be the ratio of males to females among the attendees in the party?

- |         |         |
|---------|---------|
| (a) 2:3 | (b) 1:1 |
| (c) 3:2 | (d) 2:1 |

**250.[MCQ]****[GATE-CS-2018: 2M]**

A six-sided unbiased die with four green faces and two red faces is rolled seven times. Which of the following combinations is the most likely outcome of the experiment?

- (a) Three green faces and four red faces.
- (b) Four green faces and three red faces.
- (c) Five green faces and two red faces.
- (d) Six green faces and one red face.

**251.[MCQ]****[GATE-EC-2018:2M]**

A cab was involved in a hit and run accident at night. You are given the following data about the cabs in the city and the accident.

- (i) 85% of cabs in the city are green and the remaining cabs are blue.
- (ii) A witness identified the cab involved in the accident as blue.
- (iii) It is known that a witness can correctly identify the cab colour only 80% of the time.

Which of the following options is closest to the probability that the accident was caused by a blue cab?

- |         |         |
|---------|---------|
| (a) 12% | (b) 15% |
| (c) 41% | (d) 80% |



would be the loss in Rupees for a deviation of 4 units from the target?

- (a) 400
- (b) 1200
- (c) 1600
- (d) 2800

**262.[MCQ]****[GATE-CE-2018: 2M]**

A faulty wall clock is known to gain 15 minutes every 24 hours. It is synchronized to the correct time at 9 AM on 11th July. What will be the correct time to the nearest minute when the clock shows 2 PM on 15th July of the same year?

- (a) 12:45 PM
- (b) 12:58 PM
- (c) 1:00 PM
- (d) 2:00 PM

**263.[MCQ]****[GATE-ME-2018: 2M]**

An unbiased coin is tossed six times in a row and four different such trials are conducted. One trial implies six tosses of the coin. If H stands for head and T stands for tail, the following are the observations from the four trials:

- |            |              |
|------------|--------------|
| (1) HTHTHT | (2) TTHHHT   |
| (3) HTTHHT | (4) HHHT___. |

Which statement describing the last two-coin tosses of the fourth trial has the highest probability of being correct?

- (a) Two T will occur.
- (b) One H and one T will occur.
- (c) Two H will occur.
- (d) One H will be followed by one T.

**264.[MCQ]****[GATE-ME-2018: 2M]**

A contract is to be completed in 52 days and 125 identical robots were employed, each operational for 7 hours a day. After 39 days, five-seventh of the work was completed. How many additional robots would be required to complete the work on time, if each robot is now operational for 8 hours a day?

- (a) 50
- (b) 89
- (c) 146
- (d) 175

**265.[MCQ]****[GATE-ME-2018: 2M]**

A wire would enclose an area of  $1936 \text{ m}^2$ , if it is bent into a square. The wire is cut into two pieces. The longer piece is thrice as long as the shorter piece. The long and the short pieces are bent into a square and a circle, respectively. Which of the following choices is closest to the sum of the areas enclosed by the two pieces in square meters?

- (a) 1096
- (b) 1111
- (c) 1243
- (d) 2486

**266.[MCQ]****[GATE-ME-2018: 1M]**

The perimeters of a circle, a square and an equilateral triangle are equal. Which one of the following statements is true?

- (a) The circle has the largest area.
- (b) The square has the largest area.
- (c) The equilateral triangle has the largest area.
- (d) All the three shapes have the same area.

**267.[MCQ]****[GATE-ME-2018: 2M]**

From the time the front of a train enters a platform, it takes 25 seconds for the back of the train to leave the platform, while travelling at a constant speed of 54 km/h. At the same speed, it takes 14 seconds to pass a man running at 9 km/h in the same direction as the train. What is the length of the train and that of the platform in meters, respectively?

- (a) 210 and 140
- (b) 162.5 and 187.5
- (c) 245 and 130
- (d) 175 and 200

**268.[MCQ]****[GATE-ME-2018: 2M]**

Given that  $a$  and  $b$  are integers and  $a + a^2 b^3$  is odd, which one of the following statements is correct?

- (a)  $a$  and  $b$  are both odd
- (b)  $a$  and  $b$  are both even
- (c)  $a$  is even and  $b$  is odd
- (d)  $a$  is odd and  $b$  is even

**269.[MCQ]****[GATE-ME-2018: 1M]**

A number consists of two digits. The sum of the digits is 9. If 45 is subtracted from the number, its digits are interchanged. What is the number?

- |  |   |  |  |
|--|---|--|--|
| (a) 63   | (b) 72  | <b>275.[MCQ]</b>   | <b>[GATE-EE,CS-2017 : 1M]</b>  |
| (c) 81   | (d) 90  |  | There are 3 red socks. 4 green socks and 3 blue socks. You choose 2 socks. The probability that they are of the same colour is   |
| <b>270.[MCQ]</b> <b>[GATE-ME-2018: 1M]</b>   |   |  |  |
| A rectangle becomes a square when its length and breadth are reduced by 10 m and 5 m, respectively. During this process, the rectangle loses $650 \text{ m}^2$ of area. What is the area of the original rectangle in square meters?   | (a) 1125  | (b) 2250   | (a) 1/5  |
| (c) 2924   | (d) 4500  | (c) 1/4  | (b) 7/30   |
| <b>271.[MCQ]</b> <b>[GATE-EE,CS-2017 : 1M]</b>   |   |  | (d) 4/15   |
| Find the smallest number $y$ such that $y \times 162$ is a perfect cube.   | (a) 24  | (b) 27   | <b>276.[MCQ]</b> <b>[GATE-EE,CS-2017 : 2M]</b>   |
| (c) 32   | (d) 36  | X is a 30 digit number starting with the digit 4 followed by the digit 7. Then the number $X^3$ will have  |  |
| <b>272.[MCQ]</b> <b>[GATE-EE,CS-2017 : 1M]</b>   |   |  | (a) 90 digits  |
| The probability that a k-digit number does NOT contain the digits 0, 5, or 9 is  | (a) $0.3^k$   | (b) $0.6^k$  | (c) 92 digits  |
| (c) $0.7^k$  | (d) $0.9^k$   | (d) 93 digits  | <b>277.[MCQ]</b> <b>[GATE-EC-2017:2M]</b>  |
| <b>273.[MCQ]</b> <b>[GATE-EE,CS-2017 : 2M]</b>   |   |  | 1200 men and 500 women can build a bridge in 2 weeks. 900 men and 250 women will take 3 weeks to build the same bridge. How many men will be needed to build the bridge in one week? |
| Arun, Gulab, Neel and Shweta must choose one shirt each from a pile of four shirts coloured red, pink, blue and white respectively. Arun dislikes the colour red and Shweta dislikes the colour white. Gulab and Neel like all the colours. In how many different ways can they choose the shirts so that no one has a shirt with a colour he or she dislikes? | (a) 21  | (b) 18   | (a) 3000   |
| (c) 16   | (d) 14  | (c) 3600   | (b) 3300   |
| <b>274.[MCQ]</b> <b>[GATE-EE,CS-2017 : 1M]</b>   |   |  | (d) 3900   |
| A test has twenty questions worth 100 marks in total. There are two types of questions. Multiple choice questions are worth 3 marks each and essay questions are worth 11 marks each. How many multiple choice questions does the exam have?   | (a) 12  | (b) 15   | <b>278.[MCQ]</b> <b>[GATE-EC-2017:2M]</b>  |
| (c) 18   | (d) 19  | Trucks (10 m long) and cars (5 m long) go on a single lane bridge. There must be a gap of at least 20 m after each truck and a gap of at least 15 m after each car. Trucks and cars travel at a speed of 36 km/h. If cars and trucks go alternately, what is the maximum number of vehicles that can use the bridge in one hour? |  |
|  |   | (a) 1440   | (b) 1200   |
|  |   | (c) 720  | (d) 600  |
| <b>279.[MCQ]</b> <b>[GATE-EC-2017:1M]</b>  |   |  | <b>280.[MCQ]</b> <b>[GATE-EC-2017:1M]</b>  |
| In the summer, water consumption is known to decrease overall by 25%. A Water Board official states that in the summer household consumption decreases by 20%, while other consumption increases by 70%. Which of the following statements is correct?   | (a) The ratio of household to other consumption is $8/17$ |  |  |
|  | (b) The ratio of household to other consumption is $1/17$ |  |  |
|  | (c) The ratio of household to other consumption is $17/8$ |  |  |
|  | (d) There are errors in the official's statement.         |  |  |

**280.[MCQ]****[GATE-CE-2017: 1M]**

If the radius of a right circular cone is increased by 50%, its volume increases by

- |          |            |
|----------|------------|
| (a) 75%  | (b) 100%   |
| (c) 125% | (d) 237.5% |

**281.[MCQ]****[GATE-CE-2017: 1M]**

The following sequence of numbers is arranged in increasing order: 1, x, x, x, y, y, 9, 16, 18. Given that the mean and median are equal, and are also equal to twice the mode, the value of y is

- |       |       |
|-------|-------|
| (a) 5 | (b) 6 |
| (c) 7 | (d) 8 |

**282.[MCQ]****[GATE-CE-2017: 2M]**

Budhan covers a distance of 19 km in 2 hours by cycling one fourth of the time and walking the rest. The next day he cycles (at the same speed as before) for half the time and walks the rest (at the same speed as before) and covers 26 km in 2 hours. The speed in km/h at which Budhan walks is

- |       |       |
|-------|-------|
| (a) 1 | (b) 4 |
| (c) 5 | (d) 6 |

**283.[MCQ]****[GATE-CE-2017: 2M]**

The last digit of  $(2171)^7 + (2172)^9 + (2173)^{11} + (2174)^{13}$  is

- |       |       |
|-------|-------|
| (a) 2 | (b) 4 |
| (c) 6 | (d) 8 |

**284.[MCQ]****[GATE-CE-2017: 2M]**

Two machines M1 and M2 are able to execute any of four jobs P, Q, R and S. The machines can perform one job on one object at a time. Jobs P, Q, R and S take 30 minutes, 20 minutes, 60 minutes and 15 minutes each respectively. There are 10 objects each requiring exactly 1 job. Job P is to be performed on 2 objects. Job Q on 3 objects. Job R on 1 object and Job S on 4 objects. What is the minimum time needed to complete all the jobs?

- |             |               |
|-------------|---------------|
| (a) 2 hours | (b) 2.5 hours |
| (c) 3 hours | (d) 3.5 hours |

**285.[MCQ]****[GATE-CE-2017: 1M]**

What is the value of x when

$$81 \times \left(\frac{16}{25}\right)^{x+2} \div \left(\frac{3}{5}\right)^{2x+4} = 144$$

- |        |                          |
|--------|--------------------------|
| (a) 1  | (b) -1                   |
| (c) -2 | (d) Cannot be determined |

**286.[MCQ]****[GATE-ME-2017: 2M]**

There are 4 women P, Q, R, S, and 5 men V, W, X, Y, Z in a group. We are required to form pairs each consisting of one woman and one man. P is not to be paired with Z, and Y must necessarily be paired with someone. In how many ways can 4 such pairs be formed?

- |        |        |
|--------|--------|
| (a) 74 | (b) 76 |
| (c) 78 | (d) 80 |

**287.[MCQ]****[GATE-ME-2017: 2M]**

X bullocks and Y tractors take 8 days to plough a field. If we halve the number of bullocks and double the number of tractors, it takes 5 days to plough the same field. How many days will it take X bullocks alone to plough the field?

- |        |        |
|--------|--------|
| (a) 30 | (b) 35 |
| (c) 40 | (d) 45 |

**288.[MCQ]****[GATE-ME-2017: 1M]**

P looks at Q while Q looks at R. P is married, R is not. The number of pairs of people in which a married person is looking at an unmarried person is

- |       |                          |
|-------|--------------------------|
| (a) 0 | (b) 1                    |
| (c) 2 | (d) Cannot be determined |

**289.[MCQ]****[GATE-ME-2017: 1M]**

A couple has 2 children. The probability that both children are boys if the older one is a boy is

- |         |         |
|---------|---------|
| (a) 1/4 | (b) 1/3 |
| (c) 1/2 | (d) 1   |



- (a) 3 inches = 0.5 yards
- (b) 9 inches = 1.5 yards
- (c) 9 inches = 0.25 yards
- (d) 81 inches = 0.0625 yards

**300. [MCQ] [GATE-ME-2016: 2M]**

The binary operation  $\square$  is defined as  $a \square b = ab + (a + b)$ , where  $a$  and  $b$  are any two real numbers. The value of the identity element of this operation, defined as the number  $x$  such that  $a \square x = a$ , for any  $a$ , is \_\_\_\_\_.

- (a) 0
- (b) 1
- (c) 2
- (d) 10

**301.[MCQ] [GATE-ME-2016: 2M]**

Students taking an exam are divided into two groups, P and Q such that each group has the same number of students. The performance of each of the students in a test was evaluated out of 200 marks. It was observed that the mean of group P was 105, while that of group Q was 85. The standard deviation of group P was 25, while that of group Q was 5. Assuming that the marks were distributed on a normal distribution, which of the following statements will have the highest probability of being TRUE?

- (a) No student in group Q scored less marks than any student in group P.
- (b) No student in group P scored less marks than any student in group Q.
- (c) Most students of group Q scored marks in a narrower range than students in group P.
- (d) The median of the marks of group P is 100.

**302.[MCQ] [GATE-ME-2016: 1M]**

A window is made up of a square portion and an equilateral triangle portion above it. The base of the triangular portion coincides with the upper side of the square. If the perimeter of the window is 6 m, the area of the window in  $m^2$  is \_\_\_\_\_.

- (a) 1.43
- (b) 2.06
- (c) 2.68
- (d) 2.88

**303. [MCQ]**
**[GATE-ME,EC-2016: 2M]**

P, Q, R and S are working on a project. Q can finish the task in 25 days, working alone for 12 hours a day. R can finish the task in 50 days, working alone for 12 hours per day. Q worked 12 hours a day but took sick leave in the beginning for two days. R worked 18 hours a day on all days. What is the ratio of work done by Q and R after 7 days from the start of the project?

- (a) 10:11
- (b) 11:10
- (c) 20:21
- (d) 21:20

**304. [MCQ]**
**[GATE-ME,EC-2016: 2M]**

A person moving through a tuberculosis prone zone has a 50% probability of becoming infected. However, only 30% of infected people develop the disease. What percentage of people moving through a tuberculosis prone zone remains infected but does not show symptoms of disease?

- (a) 15
- (b) 33
- (c) 35
- (d) 37

**305. [MCQ]**
**[GATE-ME,EC-2016: 1M]**

In a huge pile of apples and oranges, both ripe and unripe mixed together, 15% are unripe fruits. Of the unripe fruits, 45% are apples. Of the ripe ones, 66% are oranges. If the pile contains a total of 5692000 fruits, how many of them are apples?

- (a) 2029198
- (b) 2467482
- (c) 2789080
- (d) 3577422

**306.[MCQ]**
**[GATE-EE,CS-2015 : 2M]**

The probabilities that a student passes in Mathematics, Physics and Chemistry are  $m$ ,  $p$  and  $c$  respectively. Of these subjects, the student has 75% chance of passing in at least one, a 50% chance of passing in at least two and a 40% chance of passing in exactly two. Following relations are drawn in  $m$ ,  $p$ ,  $c$ :

- (I)  $p + m + c = 27/20$
- (II)  $p + m + c = 13/20$
- (III)  $(p) \times (m) \times (c) = 1/10$
- (a) Only relation I is true
- (b) Only relation II is true
- (c) Relations II and III are true
- (d) Relations I and III are true

**307.[MCQ]****[GATE-EE,CS-2015 : 2M]**

If p, q, r, s are distinct integers such that:

$$f(p, q, r, s) = \max(p, q, r, s)$$

$$g(p, q, r, s) = \min(p, q, r, s)$$

$h(p, q, r, s) = \text{remainder of } (p \times q) / (r \times s) \text{ if } (p \times q) > (r \times s) \text{ or remainder of } (r \times s) / (p \times q) \text{ if } (r \times s) > (p \times q)$

Also a function  $fgh(p, q, r, s) = f(p, q, r, s) \times g(p, q, r, s) \times h(p, q, r, s)$

Also the same operations are valid with two variable functions of the form  $f(p, q)$ .

What is the value of  $fg(h(2,5,7,3), 4,6,8)$ ?

- |       |       |
|-------|-------|
| (a) 6 | (b) 7 |
| (c) 8 | (d) 9 |

**308.[MCQ]****[GATE-CS,CE-2015: 1M]**

A function  $f(x)$  is linear and has a value of 29 at  $x = -2$  and 39 at  $x = 3$ . Find its value at  $x = 5$ .

- |        |        |
|--------|--------|
| (a) 59 | (b) 45 |
| (c) 43 | (d) 35 |

**309.[MCQ]****[GATE-CE-2015: 2M]**

The given question is followed by two statements: select the most appropriate option that solves the question.

Capacity of a solution tank A is 70% of the capacity of tank B. How many gallons of solution are in tank A and tank B?

Statements:

- (I) Tank A is 80% full and tank B is 40% full.
  - (II) Tank A if full contains 14,000 gallons of solution.
- |   |   |
|---|---|
| (a) Statement I alone is sufficient.              | (b) Statement II alone is sufficient.                     |
| (c) Either statement I or II alone is sufficient. | (d) Both the statements I and II together are sufficient. |

**310.[NAT]****[GATE-CE-2015: 2M]**

How many four digit numbers can be formed with the 10 digits 0, 1, 2, ..., 9 if no number can start with 0 and if repetitions are not allowed?


**311.[MCQ]****[GATE-ME-2015: 2M]**

A coin is tossed thrice. Let X be the event that head occurs in each of the first two tosses. Let Y be the event that a tail occurs on the third toss. Let Z be the event that two tails occur in three tosses. Based on the above information, which one of the following statements is TRUE?

- (a) X and Y are not independent
- (b) Y and Z are dependent
- (c) Y and Z are independent
- (d) X and Z are independent

**312.[NAT]****[GATE-ME-2015: 2M]**

In the given figure angle Q is a right angle,  $PS : QS = 3 : 1$ ,  $RT : QT = 5 : 2$  and  $PU : UR = 1 : 1$ . If area of triangle QTS is  $20 \text{ cm}^2$ , then the area of triangle PQR in  $\text{cm}^2$  is \_\_\_\_\_.

**313.[MCQ]****[GATE-ME-2015: 1M]**

Five teams have to compete in a league, with every team playing every other team exactly once, before going to the next round. How many matches will have to be held to complete the league round of matches?

- |        |        |
|--------|--------|
| (a) 20 | (b) 10 |
| (c) 8  | (d) 5  |

**314.[NAT]****[GATE-ME,EC-2015: 2M]**

From a circular sheet of paper of radius 30 cm, a sector of 10% area is removed. If the remaining part is used to make a conical surface, then the ratio of the radius and height of the cone is \_\_\_\_\_.

**315.[NAT]****[GATE-ME,EC-2015: 2M]**

A tiger is 50 leaps of its own behind a deer. The tiger takes 5 leaps per minute to the deer's 4. If the tiger and the deer cover 8 meter and 5 metre per leap respectively what distance in meters will the tiger have to run before it catches the deer?

**316.[MCQ]****[GATE-ME,EC-2015: 1M]**

Ram and Ramesh appeared in an interview for two vacancies in the same department. The probability of Ram's selection is  $1/6$  and that of Ramesh is  $1/8$ . What is the probability that only one of them will be selected?

- |             |             |
|-------------|-------------|
| (a) $47/48$ | (b) $1/4$   |
| (c) $13/48$ | (d) $35/48$ |

**317.[NAT]****[GATE-EE,CS-2014 : 1M]**

If  $(z + 1/z)^2 = 98$ , compute  $(z^2 + 1/z^2)$ .

**318.[NAT]****[GATE-EE,CS-2014 : 2M]**

Round-trip tickets to a tourist destination are eligible for a discount of 10% on the total fare. In addition, groups of 4 or more get a discount of 5% on the total fare. If the one way single person fare is Rs 100, a group of 5 tourists purchasing round-trip tickets will be charged Rs \_\_\_\_\_.

**319.[MCQ]****[GATE-EE,CS-2014 : 1M]**

What is the average of all multiples of 10 from 2 to 198?

- |         |         |
|---------|---------|
| (a) 90  | (b) 100 |
| (c) 110 | (d) 120 |

**320.[MCQ]****[GATE-EE,CS-2014 : 1M]**

The value of  $\sqrt{12 + \sqrt{12 + \sqrt{12 + \dots}}}$  is

- |           |           |
|-----------|-----------|
| (a) 3.464 | (b) 3.932 |
| (c) 4.000 | (d) 4.444 |

**321.[MCQ]****[GATE-EE,CS-2014 : 2M]**

At what time between 6 a.m. and 7 a.m. will the minute hand and hour hand of a clock make an angle closest to  $60^\circ$ ?

- |               |               |
|---------------|---------------|
| (a) 6:22 a.m. | (b) 6:27 a.m. |
| (c) 6:38 a.m. | (d) 6:45 a.m. |

**322.[MCQ]****[GATE-EE,CS-2014 : 2M]**

The Gross Domestic Product (GDP) in Rupees grew at 7% during 2012-2013. For international comparison, the GDP is compared in US Dollars (USD) after conversion based on the market exchange rate. During the period 2012-2013 the exchange rate for the USD increased from Rs. 50/USD to Rs. 60/USD. India's GDP in USD during the period 2012-2013

- (a) increased by 5 %
- (b) decreased by 13%
- (c) decreased by 20%
- (d) decreased by 11%

**323.[NAT]****[GATE-CE-2014: 1M]**

A foundry has a fixed daily cost of Rs 50,000 whenever it operates and a variable cost of Rs  $800Q$ , where  $Q$  is the daily production in tonnes. What is the cost of production in Rs per tonne for a daily production of 100 tonnes?

**324.[MCQ]****[GATE-CE-2014: 2M]**

One percent of the people of country X are taller than 6 ft. Two percent of the people of country Y are taller than 6 ft. There are thrice as many people in country X as in country Y. Taking both countries together, what is the percentage of people taller than 6 ft?

- |         |          |
|---------|----------|
| (a) 3.0 | (b) 2.5  |
| (c) 1.5 | (d) 1.25 |

**325.[MCQ]****[GATE-CE-2014: 1M]**

The population of a new city is 5 million and is growing at 20% annually. How many years would it take to double at this growth rate?

- |               |               |
|---------------|---------------|
| (a) 3-4 years | (b) 4-5 years |
| (c) 5-6 years | (d) 6-7 years |

**326.[MCQ]****[GATE-ME,EC-2014: 2M]**

A five-digit number is formed using the digits 1,3,5,7 and 9 without repeating any of them. What is the sum of all such possible five-digit numbers?

- |             |             |
|-------------|-------------|
| (a) 6666660 | (b) 6666600 |
| (c) 6666666 | (d) 6666606 |

**327. [MCQ]****[GATE-ME,EC-2014: 2M]**

Industrial consumption of power doubled from 2000-2001 to 2010-2011. Find the annual rate of increase in percent assuming it to be uniform over the years.

- |          |          |
|----------|----------|
| (a) 5.6  | (b) 7.2  |
| (c) 10.0 | (d) 12.2 |

**328. [NAT]****[GATE-ME,EC-2014: 1M]**

In a sequence of 12 consecutive odd numbers, the sum of the first 5 numbers is 425. What is the sum of the last 5 numbers in the sequence?

**329. [NAT]****[GATE-ME-2014: 2M]**

A batch of one hundred bulbs is inspected by testing four randomly chosen bulbs. The batch is rejected if even one of the bulbs is defective. A batch typically has five defective bulbs. The probability that the current batch is accepted is \_\_\_\_\_.

**330. [NAT]****[GATE-ME,EC-2014: 2M]**

A man can row at 8 km per hour in still water. If it takes him thrice as long to row upstream, as to row downstream, then find the stream velocity in km per hour.

**331. [MCQ]****[GATE-ME,EC-2014: 1M]**

In which of the following options will the expression  $P < M$  be definitely true?

- (a)  $M < R > P > S$
- (b)  $M > S < P < F$
- (c)  $Q < M < F = P$
- (d)  $P = A < R < M$

**332. [MCQ]****[GATE-ME,EC-2014: 2M]**

It takes 30 minutes to empty a half-full tank by draining it at a constant rate. It is decided to simultaneously pump water into the half-full tank while draining it. What is the rate at which water has to be pumped in so that it gets fully filled in 10 minutes?

- (a) 4 times the draining rate
- (b) 3 times the draining rate
- (c) 2.5 times the draining rate
- (d) 2 times the draining rate

**333. [NAT]****[GATE-ME,EC-2014: 2M]**

The sum of eight consecutive odd numbers is 656. The average of four consecutive even numbers is 87. What is the sum of the smallest odd number and second largest even number?

**334. [MCQ]****[GATE-ME,EC-2014: 1M]**

A regular die has six sides with numbers 1 to 6 marked on its sides. If a very large number of throws show the following frequencies of occurrence: 1 → 0.167; 2 → 0.167; 3 → 0.152; 4 → 0.166; 5 → 0.168; 6 → 0.180.

We call this die

- (a) Irregular
- (b) Biased
- (c) Gaussian
- (d) Insufficient

**335. [MCQ]****[GATE-ME,EC-2014: 2M]**

You are given three coins: one has heads on both faces, the second has tails on both faces, and the third has a head on one face and a tail on the other. You choose a coin at random and toss it, and it comes up heads. The probability that the other face is tails is

- (a)  $1/4$
- (b)  $1/3$
- (c)  $1/2$
- (d)  $2/3$

**336. [NAT]****[GATE-ME,EC-2014: 2M]**

A train that is 280 metres long, travelling at a uniform speed, crosses a platform in 60 seconds and passes a man standing on the platform in 20 seconds. What is the length of the platform in metres?

**337. [MCQ]****[GATE-EE,EC-2013 : 2M]**

A car travels 8 km in the first quarter of an hour, 6 km in the second quarter and 16 km in the third quarter. The average speed of the car in km per hour over the entire journey is

- (a) 30
- (b) 36
- (c) 40
- (d) 24

**338. [MCQ]****[GATE-EE,EC-2013 : 1M]**

In the summer of 2012, in New Delhi, the mean temperature of Monday to Wednesday was  $41^{\circ}\text{C}$  and of Tuesday to Thursday was  $43^{\circ}\text{C}$ . If the temperature on Thursday was 15% higher than that of Monday, then the temperature in  $^{\circ}\text{C}$  on Thursday was

- (a) 40
- (b) 43
- (c) 46
- (d) 49

**339.[MCQ]****[GATE-CE-2013: 1M]**

A number is as much greater than 75 as it is smaller than 117. The number is:

- |        |        |
|--------|--------|
| (a) 91 | (b) 93 |
| (c) 89 | (d) 96 |

**340.[MCQ]****[GATE-CE-2013: 2M]**

A firm is selling its product at Rs. 60 per unit. The total cost of production is Rs. 100 and firm is earning total profit of Rs. 500. Later, the total cost increased by 30%. By what percentage the price should be increased to maintained the same profit level.

- |        |        |
|--------|--------|
| (a) 5  | (b) 10 |
| (c) 15 | (d) 30 |

**341.[MCQ]****[GATE-ME,CS-2013: 2M]**

The current erection cost of a structure is Rs. 13,200. If the labour wages per day increase by  $\frac{1}{5}$  of the current wages and the working hours decrease by  $\frac{1}{24}$  of the current period, then the new cost of erection in Rs. is

- |            |            |
|------------|------------|
| (a) 16,500 | (b) 15,180 |
| (c) 11,000 | (d) 10,120 |

**342.[MCQ]****[GATE-ME,CS-2013: 2M]**

A tourist covers half of his journey by train at 60 km/h, half of the remainder by bus at 30 km/h and the rest by cycle at 10 km/h. The average speed of the tourist in km/h during his entire journey is

- |        |        |
|--------|--------|
| (a) 36 | (b) 30 |
| (c) 24 | (d) 18 |

**343.[MCQ]****[GATE-ME,CS-2013: 2M]**

Out of all the 2-digit integers between 1 and 100, a 2-digit number has to be selected at random. What is the probability that the selected number is not divisible by 7?

- |                     |                     |
|---------------------|---------------------|
| (a) $\frac{13}{90}$ | (b) $\frac{12}{90}$ |
| (c) $\frac{78}{90}$ | (d) $\frac{77}{90}$ |

**344.[MCQ]****[GATE-ME,CS-2013: 1M]**

What will be the maximum sum of 44, 42, 40, ....?

- |         |         |
|---------|---------|
| (a) 502 | (b) 504 |
| (c) 506 | (d) 500 |

**345.[MCQ]****[GATE-EE,EC-2012 : 2M]**

Raju has 14 currency notes in his pocket consisting of only Rs. 20 notes and Rs. 10 notes. The total money value of the notes is Rs. 230. The number of Rs. 10 notes that Raju has is

- |       |        |
|-------|--------|
| (a) 5 | (b) 6  |
| (c) 9 | (d) 10 |

**346.[MCQ]****[GATE-EE,EC-2012 : 1M]**

If  $(1.001)^{1259} = 3.52$  and  $(1.001)^{2062} = 7.85$ , then  $(1.001)^{3321} =$

- |           |           |
|-----------|-----------|
| (a) 2.23  | (b) 4.33  |
| (c) 11.37 | (d) 27.64 |

**347.[MCQ]****[GATE-ME,CE,CS-2012: 2M]**

Which of the following assertions are CORRECT?

- P: Adding 7 to each entry in a list adds 7 to the mean of the list
  - Q: Adding 7 to each entry in a list adds 7 to the standard deviation of the list
  - R: Doubling each entry in a list doubles the mean of the list
  - S: Doubling each entry in a list leaves the standard deviation of the list unchanged
- |          |          |
|----------|----------|
| (a) P, Q | (b) Q, R |
| (c) P, R | (d) R, S |

**348.[MCQ]****[GATE-ME,CE,CS-2012: 2M]**

An automobile plant contracted to buy shock absorbers from two suppliers X and Y. X supplies 60% and Y supplies 40% of the shock absorbers. All shock absorbers are subjected to a quality test. The ones that pass the quality test are considered reliable. Of X's shock absorbers, 96% are reliable. Of Y's shock absorbers, 72% are reliable. The probability that a randomly chosen shock absorber, which is found to be reliable, is made by Y is.

- |           |           |
|-----------|-----------|
| (a) 0.288 | (b) 0.334 |
| (c) 0.667 | (d) 0.720 |

**349.[MCQ]****[GATE-EE,EC-2011 : 2M]**

Three friends, R, S and T shared toffee from a bowl. R took  $\frac{1}{3}$ rd of the toffees, but returned four to the bowl. S took  $\frac{1}{4}$ th of what was left but returned three toffees to the bowl. T took half of the remainder but returned two back into the bowl. If the bowl had 17 toffees left, how many toffees were originally there in the bowl?

- |        |        |
|--------|--------|
| (a) 38 | (b) 31 |
| (c) 48 | (d) 41 |

**350.[MCQ]****[GATE-EE, ,EC-2011 : 1M]**

There are two candidates P and Q in an election. During the campaign, 40% of the voters promised to vote for P, and rest for Q. However, on the day of election 15% of the voters went back on their promise to vote for P and instead voted for Q. 25% of the voters went back on their promise to vote for Q and instead voted for P. Suppose, P lost by 2 votes, then what was the total number of voters?

- |         |         |
|---------|---------|
| (a) 100 | (b) 110 |
| (c) 90  | (d) 95  |

**351.[MCQ]****[GATE-ME,CE,CS-2011: 2M]**

A transporter receives the same number of orders each day. Currently, he has some pending orders (backlog) to be shipped. If he uses 7 trucks, then at the end of the 4<sup>th</sup> day he can clear all the orders. Alternatively, if he uses only 3 trucks, then all the orders are cleared at the end of the 10<sup>th</sup> day. What is the minimum number of

trucks required so that there will be no pending roder at the end of the 5<sup>th</sup> day?

- |       |       |
|-------|-------|
| (a) 4 | (b) 5 |
| (c) 6 | (d) 7 |

**352.[MCQ]****[GATE-ME,CE,CS-2011: 2M]**

A container orginally contains 10 litres of pure spirit. From this container 1 litre of spirit is repalced with 1 litre of water. Subsequently, 1 litre of the mixture is again replaced with 1 litre of water and this process is repeated one more time. How much spirit is now left in the container?

- |                 |                 |
|-----------------|-----------------|
| (a) 7.58 litres | (b) 7.84 litres |
| (c) 7 litres    | (d) 7.29 litres |

**353.[MCQ]****[GATE-ME,CE,EE,EC,CS-2010: 2M]**

Given digits 2, 2, 3, 3, 3, 4, 4, 4, 4 how many distinct 4 digit numbers greater than 3000 can be formed?

- |        |        |
|--------|--------|
| (a) 50 | (b) 51 |
| (c) 52 | (d) 54 |

**354.[MCQ]****[GATE-ME,CE,EE,EC,CS-2010: 2M]**

5 skilled workers can build a wall in 20 days, 8 semi-skilled workers can build a wall in 25 days, 10 unskilled workers can build a wall in 30 days. If a team has 2 skilled, 6 semi-skilled and 5 unskilled workers, how long will it take to build the wall?

- |             |             |
|-------------|-------------|
| (a) 20 days | (b) 18 days |
| (c) 16 days | (d) 15 days |




**ANSWER KEY**

- |                     |                      |               |                    |
|---------------------|----------------------|---------------|--------------------|
| 1. (a)              | 2. (a)               | 3. (b)        | 4. (b)             |
| 5. (a)              | 6. (c)               | 7. (c)        | 8. (c)             |
| 9. (c)              | 10. (b)              | 11. (d)       | 12. (Marks to All) |
| 13. (d)             | 14. (c)              | 15. (a)       | 16. (a)            |
| 17. (b)             | 18. (c)              | 19. (c)       | 20. (c)            |
| 21. (b)             | 22. (b)              | 23. (b)       | 24. (c)            |
| 25. (c)             | 26. (b)              | 27. (c)       | 28. (b)            |
| 29. (d)             | 30. (b)              | 31. (c)       | 32. (c)            |
| 33. (a)             | 34. (120 to 120)     | 35. (c)       | 36. (b)            |
| 37. (d)             | 38. (c)              | 39. (d)       | 40. (c)            |
| 41. (32 to 32)      | 42. (2006 to 2006)   | 43. (d)       | 44. (d)            |
| 45. (48 to 48)      | 46. (140 to 140)     | 47. (c)       | 48. (c)            |
| 49. (22 to 22)      | 50. (20000 to 20000) | 51. (d)       | 52. (d)            |
| 53. (d)             | 54. (a)              | 55. (b)       | 56. (d)            |
| 57. (d)             | 58. (d)              | 59. (b)       | 60. (d)            |
| 61. (b)             | 62. (b)              | 63. (a)       | 64. (c)            |
| 65. (b)             | 66. (a)              | 67. (a)       | 68. (a)            |
| 69. (a)             | 70. (a)              | 71. (a)       | 72. (c)            |
| 73. (c)             | 74. (c)              | 75. (c)       | 76. (b)            |
| 77. (d)             | 78. (a)              | 79. (c)       | 80. (c)            |
| 81. (d)             | 82. (d)              | 83. (b)       | 84. (b)            |
| 85. (b)             | 86. (a)              | 87. (a)       | 88. (b)            |
| 89. (d)             | 90. (c)              | 91. (a)       | 92. (c)            |
| 93. (c)             | 94. (a)              | 95. (b)       | 96. (c)            |
| 97. (c)             | 98. (a)              | 99. (c)       | 100. (a)           |
| 101. (d)            | 102. (d)             | 103. (d)      | 104. (c)           |
| 105. (Marks to All) | 106. (c)             | 107. (a)      | 108. (d)           |
| 109. (d)            | 110. (c)             | 111. (a)      | 112. (b)           |
| 113. (b)            | 114. (a)             | 115. (a)      | 116. (b)           |
| 117. (c)            | 118. (b)             | 119. (a)      | 120. (b)           |
| 121. (c)            | 122. (c)             | 123. (a)      | 124. (b)           |
| 125. (c)            | 126. (d)             | 127. (7 to 7) | 128. (a)           |
| 129. (c)            | 130. (b)             | 131. (b)      | 132. (b)           |
| 133. (a)            | 134. (b)             | 135. (c)      | 136. (d)           |
| 137. (a)            | 138. (c)             | 139. (b)      | 140. (b)           |
| 141. (b)            | 142. (a)             | 143. (c)      | 144. (c)           |

- |                 |          |          |                     |
|-----------------|----------|----------|---------------------|
| 145. (a)        | 146. (b) | 147. (b) | 148. (d)            |
| 149. (d)        | 150. (c) | 151. (c) | 152. (180 to 180)   |
| 153. (25 to 25) | 154. (a) | 155. (a) | 156. (b)            |
| 157. (d)        | 158. (a) | 159. (b) | 160. (b)            |
| 161. (b)        | 162. (c) | 163. (a) | 164. (c)            |
| 165. (d)        | 166. (a) | 167. (b) | 168. (c)            |
| 169. (a)        | 170. (c) | 171. (a) | 172. (a)            |
| 173. (a)        | 174. (c) | 175. (d) | 176. (a)            |
| 177. (c)        | 178. (a) | 179. (b) | 180. (a)            |
| 181. (a)        | 182. (d) | 183. (d) | 184. (d)            |
| 185. (d)        | 186. (d) | 187. (b) | 188. (b)            |
| 189. (d)        | 190. (c) | 191. (b) | 192. (c)            |
| 193. (a)        | 194. (a) | 195. (b) | 196. (a)            |
| 197. (c)        | 198. (a) | 199. (c) | 200. (d)            |
| 201. (d)        | 202. (c) | 203. (c) | 204. (c)            |
| 205. (c)        | 206. (c) | 207. (c) | 208. (a)            |
| 209. (c)        | 210. (d) | 211. (c) | 212. (b)            |
| 213. (a)        | 214. (b) | 215. (d) | 216. (c)            |
| 217. (d)        | 218. (b) | 219. (b) | 220. (b)            |
| 221. (d)        | 222. (c) | 223. (c) | 224. (a)            |
| 225. (c)        | 226. (a) | 227. (d) | 228. (b)            |
| 229. (c)        | 230. (b) | 231. (b) | 232. (c)            |
| 233. (b)        | 234. (c) | 235. (c) | 236. (c)            |
| 237. (a)        | 238. (c) | 239. (b) | 240. (d)            |
| 241. (b)        | 242. (a) | 243. (b) | 244. (d)            |
| 245. (c)        | 246. (d) | 247. (c) | 248. (c)            |
| 249. (b)        | 250. (c) | 251. (c) | 252. (b)            |
| 253. (b)        | 254. (d) | 255. (b) | 256. (d)            |
| 257. (c)        | 258. (b) | 259. (c) | 260. (d)            |
| 261. (c)        | 262. (b) | 263. (b) | 264. (Marks to All) |
| 265. (c)        | 266. (a) | 267. (d) | 268. (d)            |
| 269. (b)        | 270. (b) | 271. (d) | 272. (c)            |
| 273. (d)        | 274. (b) | 275. (d) | 276. (a)            |
| 277. (c)        | 278. (a) | 279. (d) | 280. (c)            |
| 281. (d)        | 282. (d) | 283. (b) | 284. (a)            |
| 285. (b)        | 286. (c) | 287. (a) | 288. (b)            |
| 289. (c)        | 290. (d) | 291. (d) | 292. (b)            |
| 293. (a)        | 294. (a) | 295. (a) | 296. (c)            |

- |                            |                            |                            |                          |
|----------------------------|----------------------------|----------------------------|--------------------------|
| <b>297.</b> (b)            | <b>298.</b> (b)            | <b>299.</b> (c)            | <b>300.</b> (a)          |
| <b>301.</b> (c)            | <b>302.</b> (b)            | <b>303.</b> (c)            | <b>304.</b> (c)          |
| <b>305.</b> (a)            | <b>306.</b> (d)            | <b>307.</b> (c)            | <b>308.</b> (c)          |
| <b>309.</b> (d)            | <b>310.</b> (4536 to 4536) | <b>311.</b> (b)            | <b>312.</b> (280 to 280) |
| <b>313.</b> (b)            | <b>314.</b> (1.9 to 2.2)   | <b>315.</b> (800 to 800)   | <b>316.</b> (b)          |
| <b>317.</b> (96 to 96)     | <b>318.</b> (850 to 850)   | <b>319.</b> (b)            | <b>320.</b> (c)          |
| <b>321.</b> (a)            | <b>322.</b> (d)            | <b>323.</b> (1300 to 1300) | <b>324.</b> (d)          |
| <b>325.</b> (a)            | <b>326.</b> (b)            | <b>327.</b> (b)            | <b>328.</b> (495 to 495) |
| <b>329.</b> (0.80 to 0.82) | <b>330.</b> (4 to 4)       | <b>331.</b> (d)            | <b>332.</b> (a)          |
| <b>333.</b> (163 to 163)   | <b>334.</b> (b)            | <b>335.</b> (b)            | <b>336.</b> (560 to 560) |
| <b>337.</b> (c)            | <b>338.</b> (c)            | <b>339.</b> (d)            | <b>340.</b> (a)          |
| <b>341.</b> (b)            | <b>342.</b> (c)            | <b>343.</b> (d)            | <b>344.</b> (c)          |
| <b>345.</b> (a)            | <b>346.</b> (d)            | <b>347.</b> (c)            | <b>348.</b> (b)          |
| <b>349.</b> (c)            | <b>350.</b> (a)            | <b>351.</b> (c)            | <b>352.</b> (d)          |
| <b>353.</b> (b)            | <b>354.</b> (d)            |                            |                          |

## SOLUTIONS

**1. (a)**

Universities = 504

Colleges = 25951

Percentage of higher education institutions that fall into grade III

$$= \frac{7 \times 504 + 23 \times 25951}{504 + 25951}$$

= 22.7%



Scan for Video solution



**2. (a)**

Mode = x with highest probability  
= -13, 13

Median =  $\frac{-1+1}{2} = 0$

Mean = 0

**3. (b)**

Mode = 7 (data with maximum frequency)  
Median

$$= \left( \frac{\frac{n}{2}^{\text{th}} + \left( \frac{n}{2} + 1 \right)^{\text{th}}}{2} \right) = \frac{25^{\text{th}} + 26^{\text{th}}}{2} = \frac{6+6}{2} = 6$$

Mean

$$\begin{aligned} &= \frac{3 \times 3 + 9 \times 4 + 11 \times 5 + 7 \times 6 + 14 \times 7 + 2 \times 8 + 4 \times 9}{3 + 9 + 11 + 7 + 14 + 2 + 4} \\ &= \frac{9 + 36 + 55 + 42 + 98 + 16 + 36}{50} \\ &= \frac{292}{50} = < 6 \\ \therefore \quad &\text{mode} > \text{median} > \text{mean} \end{aligned}$$



Scan for Video solution



## 4. (b)

Given:

$$x:y:z = \frac{1}{2}:\frac{1}{3}:\frac{1}{4} = 6:4:3 = k \text{ (say)}$$

$$\Rightarrow x = 6k, y = 4k, z = 3k$$

$$\therefore \frac{x+z-y}{y} = \frac{6k+3k-4k}{4k} = \frac{5}{4} = 1.25$$



Scan for Video solution



## 5. (a)

Given cost/unit for year 3 = 1

Cost/unit for year 2 = 2

Cost/unit for year 1 = 3

For year 2

Net profit = selling price - (cost + taxes)

$$296 = SP_2 - \left( 2 \times 200 + \frac{13}{100} \times SP_2 \right)$$

$$SP_2 = 800$$

For year 3

$$210 = SP_3 - \left( 1 \times 300 + \frac{15}{100} \times SP_3 \right)$$

$$SP_3 = 600$$

$$\text{Then } \frac{SP_2}{SP_3} = \frac{800}{600} = \frac{4}{3}$$



Scan for Video solution



## 6. (c)

% of more minutes spent than the other on Sunday

$$= \frac{65-55}{55} \times 100 = 18.18\%$$

% of more minutes spent than the other on Saturday

$$= \frac{60-50}{50} \times 100 = 20\%$$

% of more minutes spent than the other on Friday

$$= \frac{35-20}{20} \times 100 = 75\%$$

% of more minutes spent than the other on Thursday

$$= \frac{60-55}{55} \times 100 = 9.09\%$$

% of more minutes spent than the other on Wednesday

$$= \frac{60-50}{50} \times 100 = 20\%$$


% of more minutes spent than the other on Tuesday

$$= \frac{65-55}{55} \times 100 = 18.18\%$$

% of more minutes spent than the other on Monday

$$= \frac{70-45}{45} \times 100 = 55.56\%$$

So only on Thursday percentage is less than 10 and hence for remaining 6-days, percentage is more than 10.




## 7. (c)

$$\text{Total executives in } C_2 = \frac{5}{100} \times 10000 = 500$$

$$\text{Total executives in } C_5 = \frac{20}{100} \times 10000 = 2000$$

Total number of executives with management degree in  $C_2$  &  $C_5$  (from the chart)

$$= \frac{1}{5} \times 500 + \frac{9}{10} \times 2000 = 1900$$



**8. (c)**

Revenue of company Q in 2015 = 45 million Rs.  
Let the revenue of company Q in 2014 = x million Rs

Then,  
The revenue of company Q in 2015 = 1.2x million Rs.

$$\therefore 1.2x = 45$$

We get, x = 37.5 million Rs.

As per given in question, company Q had earned a profit of 10%.,

$$\therefore \text{Expenditure of Q in 2014} = \frac{37.5}{1.1} = 34.1 \text{ million Rs.}$$

So, option 'C' is correct.

 Scan for Video solution
**9. (c)**

Total expenditure during 2014 – 2018

$$5 \times 500 = 2500 \text{ million rupees}$$

Total revenue during 2014 – 2018

$$\begin{aligned} &= 500 + 700 + 800 + 600 + 400 \\ &= 3000 \text{ million rupees.} \end{aligned}$$

$$\begin{aligned} \text{Total profit} &= \frac{\text{revenue} - \text{expenditure}}{\text{expenditure}} \\ &= \frac{3000 - 2500}{2500} = 20\% \end{aligned}$$

 Scan for Video solution
**10. (b)**

Average number of students in school P

$$P_{\text{avg}} = \frac{3+5+5+6+4}{5} = \frac{23}{5}$$

Average number of students in school Q

$$Q_{\text{avg}} = \frac{4+7+8+7+5}{5} = \frac{31}{5}$$

$$\therefore Q_{\text{avg}} - P_{\text{avg}} = \frac{31}{5} - \frac{23}{5} = \frac{8}{5}$$

$$\therefore P_{\text{avg}} : (Q_{\text{avg}} - P_{\text{avg}}) = \frac{23}{5} : \frac{8}{5} = 23 : 8$$

 Scan for Video solution
**11. (d)**

$$\text{Total number of students in art} = \frac{20}{100} \times 5000 = 1000$$

$$\text{Number of girls in art} = \frac{30}{100} \times 1500 = 450$$

$$\text{So, the number of boys in art is} = 1000 - 450 = 550$$

$$\text{Number of girls in management} = \frac{15}{100} \times 1500 = 225$$

$$\text{So, the required ratio} = \frac{550}{225} = \frac{22}{9}$$

 Scan for Video solution
**12. (Marks to All)**

As per given information

Total bulb sold day two firms X and Y during April-June in year (2018) = 50,000

Assume total numbering of bulb sold by both firm in all months = P

$$\therefore 0.15 P + 0.1 P + 0.1 P = 50,000$$

$$0.35 P = 50,000$$

$$P = \frac{1000000}{7} = 142857.1429$$

LED bulbs sold by firm Y from April to June 2018

$$y_{\text{april}} + y_{\text{may}} + y_{\text{june}}$$

$$= \left( \frac{2}{5} \times 15 + \frac{4}{5} \times 10 + \frac{11}{20} \times 10 \right) \% \text{ of } P$$

$$\begin{aligned}
 &= (6 + 8 + 5.5)\% \text{ of P.} \\
 &= 19.5\% \text{ P} \\
 &= \frac{19.5 \times 142857.1429}{100} = 27857.42
 \end{aligned}$$

Total sold bulb by firm Y from April to Yance  
= 27858

Answer does not match with any of the given options



Scan for Video solution



**13. (d)**

Money spent on transport = 0.1 x

Money spent on education = 0.15 x

$$\text{Extra expenditure} = \frac{0.15x - 0.1x}{0.1x} \times 100 = 50\%$$



Scan for Video solution



**14. (c)**

Success rate of each schools:

$$P = \frac{280}{500} \times 100 = 56\%$$

$$Q = \frac{330}{600} \times 100 = 55\%$$

$$R = \frac{455}{700} \times 100 = 65\%$$

$$S = \frac{240}{400} \times 100 = 60\%$$

$$\begin{aligned}
 \text{Average of success rate} &= \frac{56 + 55 + 65 + 60}{4} \\
 &= 59\%
 \end{aligned}$$



Scan for Video solution



**15. (a)**

$\Rightarrow$  For  $y = x^m$ , the slope at  $x = 0$  &  $x = 1$

$$y' = mx^{m-1}$$

$$y'(0) = 0$$

$$y'(1) = +m$$

$\Rightarrow$  For  $y = x^{1/m}$ , the slope at  $x = 0$  &  $x = 1$

$$y' = \frac{1}{m} x^{\frac{1}{m}-1} = \frac{1}{m} \frac{1}{x^{\frac{1}{m}-1}}$$

$$y'(0) = \infty$$

$$y'(1) = \frac{1}{m}$$

These conditions are satisfied only by option A



Scan for Video solution



**16. (a)**

Let the total population be x.

Number of males in 2001 = 0.60 x

Number of females in 2001 = 0.40 x

Number of illiterates males in 2001

$$= 0.50 \times 0.60 x$$

$$= 0.30 x$$

Number of illiterates females in 2001

$$= 0.60 \times 0.40 x$$

$$= 0.24 x$$

Total number of illiterates in 2001

$$= 0.30 x + 0.24 x$$

$$= 0.54 x$$

So, total number of literates in 2001 =  $x - 0.54 x$

$$= 0.46 x$$

Similarly,

Number of males in 2011 = 0.50 x

Number of females in 2011 = 0.50 x

Number of illiterates males in 2011 =  $0.40 \times 0.50 x$   
= 0.20 x

Number of illiterates females in 2011

$$= 0.40 \times 0.50 x$$

$$= 0.20 x$$

Total number of illiterates in 2011

$$= 0.20 x + 0.20 x$$

$$= 0.40 x$$

So, number of literates in 2011 =  $x - 0.40 x$

$$\begin{aligned} \left\{ \begin{array}{l} \% \text{ increase} \\ \text{of literates} \end{array} \right\} &= \frac{\left\{ \begin{array}{l} \text{number of} \\ \text{literate in 2011} \end{array} \right\} - \left\{ \begin{array}{l} \text{number of} \\ \text{literate in 2001} \end{array} \right\}}{\text{number of literates in 2001}} = 0.60x \\ &= \frac{0.60x - 0.46x}{0.46x} \times 100 = \frac{14}{46} \times 100 = 30.43\% \end{aligned}$$



Scan for Video solution

**17. (b)**

The contribution of Pool in total revenues is

$$170 + 320 + 215 + 190 = 895$$

The contribution of Mini

$$110 + 220 + 180 + 70 = 580$$

The contribution of Prime

$$75 + 180 + 120 + 90 = 465$$

The percentage of revenue contributed by Prime

$$= \frac{465}{895 + 580 + 465} \times 100 = 23.97\%$$



Scan for Video solution

**18. (c)**

Given, total employee in 2010 was 600, according to the pie chart S had 25% and T had 5% of them.

The increase in total employee is 15% so, the number of employees increased from 2010 to 2016 is:

$$= \frac{15}{100} \times 600 = 90$$

Employee increased in S

$$= \left( \frac{25}{100} \times 600 \right) \times \frac{40}{100} = 60$$

Since employment at levels P, Q & R remained unchanged so, employees increased in T is.

$$90 - 60 = 30$$

Hence, total number of employees at level T in 2016 is

$$\frac{5}{100} \times 600 + 30 = 60$$

Scan for Video solution

**19. (c)**

Q can row only with R, so that let Q and R will go in first forward trip. In this trip Q will row. Now in return trip R will row. Q will not return, it means Q has crossed the lake.

P and R will go in second forward trip. In this trip R will row because P cannot row when anyone else is in the boat. Now in return trip P will row because P is alone in the boat. R will not return, it means R has crossed the lake too.

P and S will go in last forward trip. In this trip S will row, it means P and S have crossed the lake too.

Hence, only R rowed two times.

So, option 'C' is correct.

Scan for Video solution

**20. (c)**

Let, the number of marbles = 100

Cost of each marbles = 1 Rs.

As cost increased by 25%,

New cost = 1 + 25% of 1 = 1.25 Rs.

$$\text{New number of marbles} = \frac{100 \times 1}{1.25} = 80$$

Now, consider  $p$  number of marbles are reduced of each colour, then

$$(40 - p) + (25 - p) + (20 - p) + (15 - p) = 80$$

$$100 - 4p = 80 \Rightarrow p = 5$$

The percentage of blue marbles in new design,

$$= \frac{40 - 5}{80} \times 100 = 43.75\%$$

So, option 'C' is correct.



Scan for Video solution



21. (b)

$$\text{Given, } M + F = 300 \rightarrow (i)$$

$$= \frac{8}{9}M \times 750 + \frac{2}{3}F \times 1000$$

$$= \frac{6000}{9}M + \frac{2000}{3}F$$

$$= \frac{6000}{9}M + \frac{6000}{9}F$$

$$= \frac{6000}{9}(M + F)$$

$$TA = \frac{6000}{9} \times 300 = 200000$$



Scan for Video solution



22. (b)

$$T = K(\Theta p + T_o)$$

$$\text{For } p = 2 \text{ & } T_o = 25, T = 32.4$$

$$32.4 = K(\Theta \times 2 + 25) \quad (1)$$

$$\text{For } p = 5 \text{ & } T_o = 30, T = 42.0$$

$$42 = K(\Theta \times 5 + 30) \quad (2)$$

Equation (1)  $\div$  equation (2)

$$\frac{32.4}{42} = \frac{\Theta \times 2 + 25}{\Theta \times 5 + 30}$$

$$162 \Theta + 972 = 84 \Theta + 1050$$

$$78 \Theta = 78$$

$$\Theta = 1$$


Scan for Video solution



23. (b)

(i)  $y = 2x + 4$  for  $-3 \leq x \leq -1$   
 $\left. \begin{array}{l} x=-3, y=-2 \\ x=-1, y=2 \end{array} \right\}$  Matches with the graph

(ii)  $y = |x - 1|$  for  $-1 \leq x \leq 2$   
 $\left. \begin{array}{l} x=-1, y=2 \\ x=2, y=1 \\ x=1, y=0 \end{array} \right\}$  Matches with the graph

(iii)  $y = ||x| - 1|$  for  $-1 \leq x \leq 2$   
 $\left. \begin{array}{l} x=-1, y=0 \\ x=2, y=1 \end{array} \right\}$  Does not match with the graph

(iv)  $y = 1$  for  $2 \leq x \leq 3$  } Matches with the graph

So, (i), (ii) & (iv) are accurate descriptions of graph for the range(s) indicated.


Scan for Video solution



24. (c)

Total rainfall in 24-hour period is 300 mm.

$$\text{Area of roof top} = 50 \text{ m}^2$$

Then volume collected in the tank

$$= 50\% \text{ of } (50 \text{ m}^2 \times 0.3 \text{ m})$$

$$= 7.5 \text{ m}^3 \text{ or } 7500 \text{ liters.}$$


Scan for Video solution



25. (c)

The air pressure changes by 0.05 in the regions P, Q and S but it changes by 0.20 in the region R. So that region R is most likely to have a thunderstorm.

So, option 'C' is correct.


Scan for Video solution



**26. (b)**

Difference in heights in moving P to Q

$$= 575 - 475 = 100 \text{ m}$$

Difference in heights in moving P to R

$$= 575 - 400 = 175 \text{ m}$$



Difference in heights in moving P to S

$$= 575 - 500 = 75 \text{ m}$$

Difference in heights in moving P to T

$$= 575 - 500 = 75 \text{ m}$$

$\therefore$  The steepest path is P to R

**27. (c)**

During the path P to A

$$575 \text{m} \rightarrow 550 \text{m} \rightarrow 525 \text{m} \rightarrow 500 \text{m} \rightarrow 475 \text{m}$$

So, the movement is downwards

During the path A to B

$$475 \text{m} \rightarrow 500 \text{m} \rightarrow 525 \text{m} \rightarrow 550 \text{m} \rightarrow 575 \text{m}$$

The movement is upwards

During the path B to Q,  $575 \text{m} \rightarrow 550 \text{m}$ 

The movement is downwards


So, the total movement from P to Q is

Down  $\rightarrow$  up  $\rightarrow$  Down**28. (b)**


Pie chart is a circle representing total deaths in the city.

$$\Rightarrow 100\% \text{ of area} \rightarrow 360^\circ$$

$$\therefore 40\% \text{ of area} \rightarrow \frac{360^\circ \times 40}{100} = 144^\circ$$

**29. (d)**

The elevator does move directly from any non-ground floor to another non-ground floor over the one-hour period. Thus (i) is wrong. The elevator stays on the fourth floor for 19 min and stays on the ground floor for 21 min so, statement ii is wrong.

**30. (b)**


The allocation of rooms according to their preferences are as below.

Amar – R

Akbar – S

Anthony – P

Ajit – Q

**31. (c)**(i) no. beds by C<sub>2</sub> = no. tables by C<sub>3</sub>  
 $8 = 8$ 

(ii) no. of chairs by all &lt; no. of tables by all

$$2 + 10 + 5 + 2 + 4 < 7 + 2 + 8 + 3 + 10$$

$$23 < 30$$



Scan for Video solution



32. (c)

As mentioned in the question that water level rises to 525 m. It clearly indicates that village P and Q will not be submerged because height of these villages is above 525 m. But village R, S, T will get submerged because height of these villages is below 525 m. So, option 'C' is correct.



Scan for Video solution



33. (a)

From the graph it is clear that growth of bacteria at 37°C stops earlier than that at 25°C. So, statements I is correct.

Curd formation needs minimum density of 0.8 so at 37°C, Curd form after 90 min and Curd formation at 25° is after 130 min. So statement II is wrong.



Scan for Video solution



34. (120 to 120)

The plants 1, 4, 8, 9 have installed capacity more than 200 tones, which means these are large plants.

∴ Total production of large plant is

$$160 + 190 + 230 + 190 = 770$$

The plants 2, 3, 5, 6, 7 have installed capacity less than 200, which means these are small plants.

∴ Total production of small plant is

$$= 150 + 160 + 120 + 100 + 120 = 650$$

∴ The difference between large and small plants production =  $770 - 650 = 120$



Scan for Video solution



35. (c)

Function  $f(x)$  is zero at  $x = -1$  and  $x = 3$ , also  $f(x)$  is 2 at  $x = 1$ . These three conditions are satisfied by only option (c).

$$f(x) = 2 - |x - 1|$$

$$f(-1) = 2 - |-1 - 1| = 2 - 2 = 0$$

$$f(3) = 2 - |3 - 1| = 2 - 2 = 0$$

$$f(1) = 2 - |1 - 1| = 2 - 0 = 2$$

So, option 'C' is correct.



Scan for Video solution



36. (b)

| Quarter \ Product | Elegance | Smooth | Soft  | Executive |
|-------------------|----------|--------|-------|-----------|
| Q1                | 27300    | 20009  | 17602 | 9999      |
| Q2                | 25222    | 19392  | 18445 | 8942      |
| Q3                | 28976    | 22429  | 19544 | 10234     |
| Q4                | 21012    | 18229  | 16595 | 10109     |
| Total             | 1,02510  | 80,059 | 72186 | 39,284    |

$$\text{Revenue of Elegance} \rightarrow 1,02,510 \times 48 = 49,20,480$$

$$\text{Revenue of Smooth} \rightarrow 80,059 \times 63 = 50,43,717$$

$$\text{Revenue of Soft} \rightarrow 72,186 \times 78 = 56,30,508$$

Revenue of Executive  $\rightarrow 39,284 \times 173 = 67,96,132$

Executive contributes the highest to the revenue.



Scan for Video solution



37. (d)

Increase in odometer reading

= Distance travelled by vehicle in 7 seconds

= Area under the V-t graph

$$\begin{aligned} &= \frac{1}{2} \times 1 \times 1 + 1 \times 1 + 1 \times 1 + \frac{1}{2} \times 1 \times 1 + \frac{1}{2} \times 2 \times 1 \\ &\quad + \frac{1}{2} \times 1 \times 1 + \frac{1}{2} \times 1 \times 1 \end{aligned}$$

$$= \frac{1}{2} + 1 + 1 + \frac{1}{2} + 1 + \frac{1}{2} + \frac{1}{2} = 5 \text{ m}$$



Scan for Video solution




38. (c)

$e^x$  = positive for  $-\infty < x < +\infty$

$$y = \ln \left( \left| e^{\lceil \sin(|x)| \rceil} \right| \right) = \ln \left( e^{\lceil \sin(|x)| \rceil} \right)$$

$$= |\sin|x|| \ln e$$

$$= |\sin|x||$$



Scan for Video solution



39. (d)

Given,  $\frac{A_P}{A_Q} = \frac{8}{9}$  and rate of interest by P & Q in 2006 are 6% and 4% respectively.

Interest  $\propto A \times r$

$$\frac{\text{Interest}_P}{\text{Interest}_Q} = \frac{A_P \times r_P}{A_Q \times r_Q} = \frac{8}{9} \times \frac{6}{4} = \frac{4}{3}$$



Scan for Video solution



40. (c)

Total number of students in the class = 44

Total marks scored by the class,

$$= 21 \times 2 + 15 \times 3 + 11 \times 1 + 23 \times 2 + 31 \times 5 = 299$$

$$\therefore \text{average marks} = \frac{\text{Total marks scored by the class}}{\text{Total number of students in the class}}$$

$$= \frac{299}{44} = 6.795$$

So, option 'C' is correct.



Scan for Video solution



41. (32 to 32)

Number of male students in electrical = 40

Let the number of female students in electrical =  $F_e$

$$\therefore \frac{40}{F_e} = \frac{5}{4} \Rightarrow F_e = 32$$

Therefore, total student in electrical =  $40 + 32 = 72$

Now, let total student in college =  $S$

$$\text{Then, } \therefore S \times \frac{20}{100} = 72 \Rightarrow S = 360$$

Let the number of female students in civil =  $F_c$

$$\therefore F_c = \frac{30}{100} \times 360 \times \frac{4}{9} = 48$$

Let the number of female students in mechanical =  $F_m$

$$\therefore F_m = \frac{10}{100} \times 360 \times \frac{4}{9} = 16$$

Then the difference =  $48 - 16 = 32$



Scan for Video solution



#### **42. (2006 to 2006)**

⇒ For 2004

$$\text{Export} \rightarrow \frac{10}{60} = \frac{1}{6} \left[ \frac{1}{6} + \frac{1}{7} = \frac{13}{42} \right] \times 100 = 30.95\%$$

$$\text{Import} \rightarrow \frac{10}{70} = \frac{1}{7}$$

⇒ For 2005

Export = 0

$$\text{Import} \rightarrow \frac{10}{80} = \frac{1}{8} \times 100 = 12.5\%$$

⇒ For 2006

$$\text{Export} \rightarrow \frac{30}{70} = \frac{3}{7} \left[ \frac{3}{7} + \frac{3}{9} = \frac{48}{63} \right] = 76.19\%$$

$$\text{Import} \rightarrow \frac{30}{90} = \frac{3}{9}$$

⇒ For 2007

$$\text{Export} = \frac{10}{100} = 10\% \quad \left. \right\} = 10\%$$

$$\text{Import} = 0$$



Scan for Video solution



#### **43. (d)**

$$\text{Growth of type I battery} = \frac{115 - 75}{75} \times 100 = 53.33\%$$

$$\text{Growth of type II battery} = \frac{85 - 144}{144} \times 100 = -40.97\%$$

$$\text{Growth of type III battery} = \frac{160 - 114}{114} \times 100 = 40.35\%$$

$$\text{Growth of type IV battery} = \frac{100 - 102}{102} \times 100 = -1.96\%$$

$$\text{Growth of type V battery} = \frac{145 - 108}{108} \times 100 = 34.26\%$$

So, type I battery has the highest growth between 2006 to 2012.



Scan for Video solution



#### **44. (d)**

| Stretch | Cumulative distance | Electricity used | Electricity/Distance                |
|---------|---------------------|------------------|-------------------------------------|
| M       | 20                  | 12               | $= \frac{12}{20} = 0.6$             |
| N       | 45                  | 25               | $= \frac{25 - 12}{45 - 20} = 0.52$  |
| O       | 75                  | 45               | $= \frac{45 - 25}{75 - 45} = 0.66$  |
| P       | 100                 | 57               | $= \frac{57 - 45}{100 - 75} = 0.48$ |

Electricity consumption per km is minimum in P.



Scan for Video solution



**45. (48 to 48)**

Total people have only scooter =  $30 + 20 = 50$   
 Total people have both (scooter + car) =  $60 + 46 = 106$   
 Total people do not have scooter,  
 $= 40 + 34 + 20 + 50 = 144$   
 Then the percentage of people do not have a scooter,  
 $= \frac{144}{300} \times 100 = 48\%$



Scan for Video solution

**46. (140 to 140)**

Let the female students in the 2008 = F  
 And the male students in the 2008 = M  
 $\therefore$  Ratio of male to female in 2008

$$\Rightarrow \frac{M}{F} = 2.5 \Leftrightarrow M = 2.5F$$

Now the female students in the 2009 = 2F  
 Now let the male students in the 2009 =  $M_1$   
 And given that Ratio of male to female in 2009

$$\Rightarrow \frac{M_1}{2F} = 3 \Leftrightarrow M_1 = 6F$$

The increment in percentage of male from 2008 to 2009

$$= \frac{6F - 2.5F}{2.5F} \times 100 = 140\%$$



Scan for Video solution

**47. (c)**

Total number of students in the class = 44  
 Total marks scored by the class,  
 $= 21 \times 2 + 15 \times 3 + 23 \times 2 = 133$   
 $\therefore$  average marks =  $\frac{\text{Total marks scored by the class}}{\text{Total number of students in the class}}$

$$= \frac{133}{44} = 3.0227 \approx 3.02$$

So, option 'C' is correct.



Scan for Video solution

**48. (c)**

Let the female students in the 2011 = F  
 And the male students in the 2011 = M  
 $\therefore$  Ratio of male to female in 2011

$$\Rightarrow \frac{M}{F} = 1 \Leftrightarrow M = F$$

Now the female students in the 2012 = F

Now let the male students in the 2012 =  $M_1$

And given that Ratio of male to female in 2012

$$\Rightarrow \frac{M_1}{F} = 1.5 \Leftrightarrow M_1 = 1.5F$$

Then the ratio of male student from 2012 to 2011

$$= \frac{1.5F}{F} = 1.5 : 1$$

So, option 'C' is correct.



Scan for Video solution

**49. (22 to 22)**

2013  $\rightarrow$  Raw material expenses =  $1.3 \times 20 = 26\%$   
 Other expenses =  $1.2 \times 80$   
 $= 96\%$

$$26\% + 96\% = 122\%$$

$$122 - 100 = 22\%$$



Scan for Video solution

**50. (20000 to 20000)**

$0.15x = 4,50,000$   
 $x = 4,50,000 / 0.15$   
 $x = 30,00,000$  (total manufacturing cost)  
 Profit = 10,00,000  
 $S.P. = x + \text{Profit} = 40,00,000$

$$\text{Each unit S.P.} = \frac{40,00,000}{200} \\ = 20,000$$



Scan for Video solution

**51. (d)**

The pie chart shows us population of animals not the size of animals so, statement (iv) is false. All other statements are true.





Scan for Video solution

**52. (d)**

$$\text{Revenue generated per kg} \propto \frac{\% \text{ of revenue of the item}}{\% \text{ of export of the item}}$$

$$\text{For item } [2], \frac{0.2}{0.2} = 1$$

$$\text{For item } [3], \frac{0.23}{0.19} = 1.210$$



$$\text{For item } [6], \frac{0.19}{0.18} = 1.1875$$

$$\text{For item } [5], \frac{0.2}{0.12} = 1.6$$



Scan for Video solution

**53. (d)**

Given,

$$\text{Trade Deficit (TD)} = \frac{\text{Import (I)} - \text{Export (E)}}{\text{Export (E)}}$$

$$\text{For 2005, TD} = (90-70)/70 = 2/7$$

$$\text{Similarly for 2004, TD} = 1/7$$

$$\text{for 2007, TD} = 1/11$$

$$\text{for 2006, TD} = 1/5$$



Scan for Video solution

**54. (a)**

The standard deviation gives an idea about how the data is spread. For a low standard deviation, most of the scores will be near mean value (average).

$$\text{consistency} \propto \frac{1}{\text{standard deviation}}$$

Since standard deviation is lowest for K, hence K is the most consistent batsman.



Scan for Video solution

**55. (b)**

In July, 5% of rainfall is below 5 percentile line and in January, 5% of rainfall is above 95 percentile line. So, there might be a year for which the rainfall in January is more than that in July. Similarly, there might be a year in which the rainfall in August is below 500 mm. So, statements (ii) and (iv) are not true. The variation in July is very less hence July rainfall can be estimated with more confidence.



Scan for Video solution

**56. (d)**

$$\text{Total export} = 5 \times 10^8 \text{ kgs}$$

$$\text{Total revenue} = 250 \times 10^7 \text{ Rs}$$

$$\text{Item 1 export} = 0.11 \times 5 \times 10^8 \text{ kg} = 55 \times 10^6 \text{ kg}$$

$$\text{Item 2 export} = 0.22 \times 5 \times 10^8 \text{ kg} = 1.1 \times 10^8 \text{ kg}$$

Item 1 revenue =  $0.12 \times 250 \times 10^7$  Rs =  $30 \times 10^7$  Rs  
 Item 4 revenue =  $0.06 \times 250 \times 10^7$  Rs =  $15 \times 10^7$  Rs

$$\text{Revenue per kg for item 1: } - \frac{30 \times 10^7}{0.55 \times 10^8}$$

$$\text{Revenue per kg for item 4: } - \frac{15 \times 10^7}{1.1 \times 10^8}$$

$$\Rightarrow \frac{\frac{30 \times 10^7}{0.55 \times 10^8}}{\frac{1.5 \times 10^7}{1.1 \times 10^8}} = \frac{220}{55} = 4 : 1$$



Scan for Video solution



**57. (d)**

$$(1) \frac{6240 - 5200}{5200} \times 100 = \uparrow 20\%$$

$$(2) \frac{9450 - 7000}{7000} \times 100 = \uparrow 35\%$$

$$(3) \frac{12600 - 9000}{9000} \times 100 = \uparrow 40\%$$

$$(4) \frac{25000 - 20000}{20000} \times 100 = \uparrow 25\%$$

$$(5) \frac{19500 - 15000}{15000} \times 100 = \uparrow 30\%$$

$$(6) \frac{26400 - 22000}{22000} \times 100 = \uparrow 20\%$$



Scan for Video solution



**58. (d)**

| Category | Amount |
|----------|--------|
| Food     | 4000   |
| Clothing | 1200   |
| Rent     | 2000   |
| Savings  | 1500   |
| Others   | 1800   |
| Total    | 10500  |

So, total monthly budget of the household = 10500

Total monthly saving = 1500

Then, the percentage of the monthly budget not spent on saving =  $\frac{9,000}{10,500} \times 100 = 85.71\% \approx 86\%$

So, option 'D' is correct.



Scan for Video solution



**59. (b)**

For lap 'P'

Average speed = 15 km/hr

Distance travelled = 15 km

Distance travelled per liter at 15 km/hr = 60 km

Fuel consumption per km at 15 km/hr =  $\frac{1}{60}$  liter

For lap 'Q'

Average speed = 45 km/hr

Distance travelled = 75 km

Distance travelled per liter at 45 km/hr = 90 km

Fuel consumption per km at 45 km/hr =  $\frac{1}{90}$  liter

For lap 'R'

Average speed = 75 km/hr

Distance travelled = 40 km

Distance travelled per liter at 75 km/hr = 75 km

Fuel consumption per km at 75 km/hr =  $\frac{1}{75}$  liter

For lap 'S'

Average speed = 10 km/hr

Distance travelled = 10 km

Distance travelled per liter at 10 km/hr = 30 km

Fuel consumption per km at 10 km/hr =  $\frac{1}{30}$  liter

Hence, from the above four laps, lap 'Q' has least consumption of fuel per km.

So, option 'B' is correct.



Scan for Video solution



**60. (d)**


$$\text{Level of danger} \propto \frac{\text{Growth (G)} \times \text{Potency (P)}}{\text{Toxicity (T)}}$$

Where G = area of the circle and P and T can be taken from the graph

From the above formula, level of danger is maximum for S hence will be targeted first.



Scan for Video solution

**61. (b)**

$$\text{Area of triangle } ABC = \frac{1}{2} \times 3 \times 3 = 4.5 \text{ cm}^2$$

$$\text{Area of quarter circle} = \frac{1}{4} \pi r^2$$

$$= \frac{1}{4} \times \pi \times (3)^2 = \frac{9}{4} \pi \text{ cm}^2$$

$$\text{Now area of arc (blank portion)} = 2 \times \left( \frac{9}{4} \pi - 4.5 \right)$$

$\therefore$  Area of dark portion =

$$2 \left( \frac{\pi}{2} \times (3)^2 \right) - 2 \times \left( \frac{9}{4} \pi - 4.5 \right)$$

$$= 9\pi - 9\pi + 18 = 18 \text{ cm}^2$$

$\therefore$  Option 'B' is correct.



Scan for Video solution

**62. (b)**

Given graph is an odd function.

From the given all 4 options only option 'B' represents an odd function.

$\therefore$  Option (b) is correct.



Scan for Video solution

**63. (a)****Given Data:**

Total number = 6 (i.e. 1, 2, 3, 4, 5, 6)

$$\text{Probability of getting '1'} = P(1) = \frac{1}{6}$$

$$\text{Probability of getting '4'} = P(4) = \frac{1}{6}$$

$\therefore$  As we can see here, both events are independent to each other.

$$\therefore P(A \cap B) = P(A) \cdot P(B)$$

$$P(1 \cap 4) = P(1) \cdot P(4) = \frac{1}{6} \times \frac{1}{6} = \frac{1}{36}$$

$\therefore$  Option (a) is correct.



Scan for Video solution

**64. (c)**


$$\text{Given } f(t) = 0.01t^2$$

$$g(t) = 4t$$

$$\text{For point A, } g(t) = f(t)$$

$$0.01t^2 = 4t$$

$$t = 20$$



(1) Since for some  $t > 0$  ( $0 < t < 20$ )  $g(t) > f(t)$

(2) After  $t = 20$  (say  $T$ )  $f(t) > g(t)$  for all  $t > T$ .



Scan for Video solution




### 65. (b)

Given  $f(x) = g(y)$  for all real values of  $x$  &  $y$ . So,

$$f(4) = g(5)$$

or

$$f(11) = g(6)$$



This is only possible when

$$f(4) = f(11) = k \text{ (constant)}$$

$$g(5) = g(6) = k \text{ (constant)}$$



Scan for Video solution



### 66. (a)

No. of pairs of 6-digit subsets & its subsets

= no. of 6-digit subsets  $\times$  no. of subsets of each 6-digit subset

$$= {}^6C_6 \times 2^6 = 64$$

No. of pairs of 5-digit subsets & its subsets

= no. of 5-digit subsets  $\times$  no. of subsets of each 5-digit subset

$$= {}^6C_5 \times 2^5 = 6 \times 32 = 192$$

No. of pairs of 4-digit subsets & its subset

$$= {}^6C_4 \times 2^4 = 15 \times 16 = 240$$

No. of pairs of 3-digit subset & its subset

$$= {}^6C_3 \times 2^3 = 20 \times 8 = 160$$

No. of 2-digit subset & its subset

$$= {}^6C_2 \times 2^2 = 15 \times 4 = 60$$

No. of 1-digit subset & its subset

$$= {}^6C_1 \times 2^1 = 6 \times 2 = 12$$

No. of pairs of empty subsets & its subset ( $\phi, \phi$ )

$$= 1$$

Total pairs =  $64 + 192 + 240 + 160 + 60 + 12 + 1 =$

$$729$$



Scan for Video solution



### 67. (a)

Positive integers  $\Rightarrow 1, 2, 3, \dots$

$\Rightarrow$  For  $p + q = 2$  (option A)

The only possibility is  $p = q = 1$

This condition satisfies both the inequalities

$$p^2 = 4q = -3 < 4$$

$$3p + 2q = 5 < 6$$

$\Rightarrow p + q = 1$  is not possible because zero is not a positive integer.

$\Rightarrow p + q = 3$  can have two possibilities  $p = 1, q = 2$  and  $p = 2, q = 1$ . Both the possibilities do not satisfy both the inequalities.



Scan for Video solution



### 68. (a)


Total shaded area =  $2x + 2y$

$$\text{Area of quadrant (y)} = \frac{1}{4}\pi r^2 = \frac{1}{4}\pi(2)^2 = \pi\text{cm}^2$$

Area of sector (x)

$$= \frac{\text{Area of square}}{4} - \text{Area of quadrant}$$

$$= \frac{4^2}{4} - \pi = \{4 - \pi\}\text{cm}^2$$



Total shaded area =  $2x + 2y$

$$= 2(4 - \pi) + 2\pi = 8 \text{ cm}^2$$



Scan for Video solution



69. (a)



Each angle of a regular hexagon: -

$$\Rightarrow \frac{(n-2) \times 180}{n} = \frac{(6-2) \times 180}{6}$$

$$\Rightarrow \frac{720}{6} = 120^\circ$$

$$\text{Area of sector} = \frac{\theta}{360^\circ} \pi r^2$$

$$= v \propto r = \frac{25}{3} \times \pi \text{ cm}^2$$



Scan for Video solution



70. (a)

$$\log_n m = \frac{\log m}{\log n}$$

$$\log_a c = \frac{\log 100!}{\log 30!}, \log_c a = \frac{\log 30!}{\log 100!},$$

$$\log_b a = \frac{\log 30!}{\log 50!}, \log_a b = \frac{\log 50!}{\log 30!}$$

$\log_c a < \log_b a < \log_a b < \log_a c$



Scan for Video solution



71. (a)

Q, R, S, T, U, and V are total 6 faces of each dice,  
Total possibilities, when two dice are thrown once independently at random =  $6 \times 6 = 36$

Possible outcome combinations due to Q, U, V = 9  
i.e. (Q, Q), (Q, U), (U, Q), (Q, V), (V, Q), (U, U), (V, V), (U, V), (V, U).

$$\therefore \text{Required probability} = \frac{9}{36} = \frac{1}{4}$$

$\therefore$  option (a) is correct.



Scan for Video solution



72. (c)

As we can observe that in front and rear view of the disc, the view is only symmetrical about 1-1 axis.  
Now if we do flip operation, it will retain the same front and rear views in only one case.

It means it will not retain the same front and rear views with respect to 2-2 and 3-3.

Hence out of three cases, two case does not retain the same front and rear views.

$$\therefore \text{Required probability} = \frac{2}{3}$$

$\therefore$  option (c) is correct



Scan for Video solution




73. (c)

The graph for  $y(x)$  is drawn below.

Area under the curve is

$$= \frac{1}{3} \times 2 + \left( \frac{3}{4} - \frac{1}{3} \right) \times 3 + \left( 1 - \frac{3}{4} \right) \times 1$$




$$= \frac{2}{3} + \frac{5}{4} + \frac{1}{4} = \frac{8+15+3}{12} = \frac{26}{12} = \frac{13}{6}$$



Scan for Video solution


**74. (c)**

The given quadrilateral is as shown in the figure



Length of side PS = Distance between the points (0, 1) and (2, -1)

$$\Rightarrow PS = \sqrt{(0-2)^2 + (1+1)^2} = \sqrt{4+4} = \sqrt{8}$$

Similarly

$$PR = \sqrt{(0+2)^2 + (1+1)^2} = \sqrt{8}$$

$$SQ = \sqrt{(2-0)^2 + (-1+3)^2} = \sqrt{8}$$

$$QR = \sqrt{(0+2)^2 + (-3+1)^2} = \sqrt{8}$$

That implies all the sides are equal to  $\sqrt{8}$  unit

$$\text{Length of diagonal } PQ = \sqrt{(0-0)^2 + (1+3)^2} = 4$$

$$\text{Length of Diagonal RS} = \sqrt{(-2-2)^2 + (1+1)^2} = 4$$

That implies diagonals are equal to 4 unit.

Hence the given quadrilateral PSQR is a square

$$\therefore \text{Area} = (\text{length of the side})^2$$

$$= (\sqrt{8})^2 = 8 \text{ square units}$$

Scan for Video solution


**75. (c)**

$$\text{Given: } 2x - 1 > 7$$

$$\Rightarrow 2x > 8 \Rightarrow x > 4$$

$$\text{Also, } 2x - 9 < 1 \Rightarrow 2x < 10 \Rightarrow x < 5$$


So, we can write,  $4 < x < 5$

$\Rightarrow$  Hence option (c) is correct

Scan for Video solution


**76. (b)**

**Given:** Trapezium PQRS



The shortest distance between PQ and SR will be the perpendicular distance  $h$  between them.

$$\therefore PQ = 11 \Rightarrow x + 6 + y = 11$$

$$\Rightarrow x + y = 5 \quad \dots(a)$$

For the triangles we can write

$$h^2 + x^2 = 3^2 = 9 \quad \dots(1)$$

$$\text{and } h^2 + y^2 = 4^2 = 16 \quad \dots(2)$$

Equation (2) – equation (1)

$$y^2 - x^2 = 7 \Rightarrow (y-x)(y+x) = 7$$

$$\Rightarrow y - x = \frac{7}{5} \quad \dots (3)$$

Equation (a) + Equation (3)

$$2y = \frac{7}{5} + 5 = \frac{32}{5} \Rightarrow y = \frac{16}{5}$$

From equation (2)

$$h^2 = 16 - \left(\frac{16}{5}\right)^2 = 16 - \frac{256}{25} = \frac{144}{25}$$

$$\Rightarrow h = \sqrt{\frac{144}{25}} = \frac{12}{5} = 2.4$$



Scan for Video solution



**77. (d)**

- Lines are parallel  $\Rightarrow$  their slopes are equal.
- Lines are perpendicular  $\Rightarrow$  the product of their slopes is  $-1$ .

$$\text{Line L1: } 2x - 3y = 5 \Rightarrow y = \frac{2x}{3} - \frac{5}{3}$$

$$\text{slope} = \frac{2}{3}$$

$$\text{Line L2: } 3x + 2y = 8 \Rightarrow y = -\frac{3x}{2} + \frac{8}{2}$$

$$\text{slope} = -\frac{3}{2}$$

$$\text{Line L3: } 4x - 6y = 5 \Rightarrow y = \frac{4x}{6} - \frac{5}{6}$$

$$\text{slope} = \frac{2}{3}$$

$$\text{Line L4: } 6x - 9y = 6 \Rightarrow y = \frac{6x}{9} - \frac{6}{9}$$

$$\text{slope} = \frac{2}{3}$$

$\therefore$  Slope (L4) = Slope (L3) and

Slope (L4)  $\times$  Slope (L2) =  $-1$ , **L4 is perpendicular to L2 and L4 is parallel to L3**



Scan for Video solution



**78. (a)**

$$\Rightarrow \begin{cases} f(2,2) = 2^2 = 4 \\ g(2,2) = 2^2 = 4 \end{cases} \quad f(2,2) = g(2,2)$$

$$\Rightarrow f(g(2,2),2) = f(4,2) = 4^2 = 16$$

$$f(2,g(2,2)) = f(2,4) = 2^4 = 16$$

$$\Rightarrow \begin{cases} g(2,1) = 2 \\ f(2,1) = 2 \end{cases} \quad f(2,1) = g(2,1)$$

$$\Rightarrow \begin{cases} f(3,2) = 3^2 = 9 \\ g(3,2) = 3^3 = 27 \end{cases} \quad f(3,2) < g(3,2)$$



Scan for Video solution




**79. (c)**

Given,

$$f(x) = 2 \ln \sqrt{e^x}$$

$$= 2 \ln e^{\frac{x}{2}} = 2 \frac{x}{2} = x$$

$$\text{Area} = \frac{1}{2} \cdot 2 \times 2 = 2$$



Scan for Video solution



**80. (c)**

$$2 - 5x \leq -\frac{6x - 5}{3}$$


$$6 - 15x \leq -6x + 5 \quad \leftarrow \begin{array}{c} 0 \quad \frac{1}{9} \end{array}$$

$$15x - 6x + 5 - 6 \geq 0$$

$$9x - 1 \geq 0$$

$$x \geq \frac{1}{9}$$

This result is similar to option C.



Scan for Video solution



**81. (d)**

Number of students appear in year 2 =  $60 + 5 = 65$   
 But 10 students who have failed in year 1 are also appear in year 2,  
 So that, students appear in year 2 for the very first time =  $65 - 10 = 55$ .  
 Similarly,  
 Number of students appear in year 3 =  $50 + 3 = 53$   
 But 5 students who have failed in year 2 are also appear in year 3,  
 So that, students appear in year 3 for the very first time =  $53 - 5 = 48$   
 So, option 'D' is correct.



Scan for Video solution

**82. (d)**

Given  $f(x) = 0.01$  for  $0 \leq x \leq 100$   
 The mean value of the X will be,

$$\mu = \int_{-\infty}^{\infty} xf(x)dx$$


But the function lies in the interval of  $0 \leq x \leq 100$ .

$$\text{Therefore, } \mu = \int_0^{100} 0.01x dx = 0.01 \times \left[ \frac{x^2}{2} \right]_0^{100} = 50$$

So, option 'D' is correct.




Scan for Video solution

**83. (b)**

$$\sin \theta = \frac{\sqrt{2}a}{\sqrt{3}a} = \frac{\sqrt{2}}{\sqrt{3}}$$

$$\cos \theta = \frac{a}{\sqrt{3}a} = \frac{1}{\sqrt{3}}$$



▶
Scan for Video solution

**84. (b)**

$$\left( x - \frac{1}{2} \right)^2 - \left( x - \frac{3}{2} \right)^2 = x + 2$$

$$\left( x - \frac{1}{2} + x - \frac{3}{2} \right) \left( x - \frac{1}{2} - x + \frac{3}{2} \right) = x + 2$$

$$(2x - 2)(1) = x + 2 \Rightarrow x = 4$$

▶
Scan for Video solution

**85. (b)**

Given,

$$\lambda(p, q) = \begin{cases} (p - q)^2, & \text{if } p \geq q \\ (p + q), & \text{if } p < q \end{cases}$$

$$f(1) = \lambda(-(-3+2), (-2+3)) / ((-(-2+1)))$$

$$f(1) = \lambda(1, 1)$$

$$\text{as } p = 1 \text{ } \$ \text{ } q = 1 ; p = q$$


We can write

$$f(1) = \lambda(1, 1) = (1 - 1)^2 = 0$$

▶
Scan for Video solution

86. (a)

Regular hexagon has equal sides which is equal to the base of corner triangles which are equilateral also as shown in figure.



Area of  $\Delta ABC$

$$A = \frac{\sqrt{3}}{4} (\text{side})^2 = \frac{\sqrt{3}}{4} (3a)^2 = \frac{9\sqrt{3}}{4} a^2$$

Area of hexagon

$A_H = \text{Area of } \Delta ABC - 3 \times \text{Area of corner triangle}$

$$= \frac{9\sqrt{3}}{4} a^2 - 3 \times \frac{\sqrt{3}}{4} \cdot a^2$$

$$= \frac{\sqrt{3}}{4} a^2 [9 - 3] = 6 \frac{\sqrt{3}}{4} a^2$$

$$\therefore A_H : A = 6 \frac{\sqrt{3}}{4} a^2 : 9 \frac{\sqrt{3}}{4} a^2 = 6 : 9 = 2 : 3$$





Scan for Video solution



87. (a)

Let the triangle ABC is revolved about side BC, resulting in a cone as shown



So,  $r = 1$  unit and  $H = 1$  unit

$$\text{Volume of cone, } V = \frac{1}{3} \pi r^2 H = \frac{\pi}{3} \text{ cubic unit}$$



Scan for Video solution



88. (b)

$$\text{Given: } \frac{p}{q} + \frac{q}{p} = 3 \Rightarrow \left( \frac{p}{q} + \frac{q}{p} \right)^2 = 3^2$$

$$\Rightarrow \frac{p^2}{q^2} + \frac{q^2}{p^2} + 2 = 9$$

$$\Rightarrow \frac{p^2}{q^2} + \frac{q^2}{p^2} = 9 - 2 = 7$$

Scan for Video solution



89. (d)

$$P(\text{even on both dice}) = P(\text{even on I}^{\text{st}} \& \text{ even on II}^{\text{nd}})$$

$$= P(\text{even - I}) \times P(\text{even - II})$$


$$= \frac{1}{2} \times \frac{1}{2}$$

$$= \frac{1}{4}$$

Scan for Video solution



90. (c)



$$\text{Area of quadrant of a circle} = \frac{\pi r^2}{4}$$

$$\text{Shaded area} = 2 \times \frac{\pi r^2}{4} - r^2$$

$$P(\text{Any point in shaded area}) = \frac{\text{Shaded Area}}{\text{Area of square}}$$

$$= \frac{\frac{\pi r^2}{4} \times 2 - r^2}{r^2} = \frac{\pi}{2} - 1$$



Scan for Video solution

**91. (a)**

In the given figure,

OA = OB = OC = radius of circle

And two right angle triangles are formed with these radii.

$$\therefore AC = BC = \sqrt{OA^2 + OC^2} = \sqrt{2}OA = \sqrt{2}OB$$

$$\therefore \frac{\overline{AC} + \overline{CB}}{\overline{AB}} = \frac{\sqrt{2}OA + \sqrt{2}OB}{OA + OB} = \sqrt{2}$$

So, option 'A' is correct.



Scan for Video solution

**92. (c)**

Four-digit integer numbers from 1001 to 9999, the digit "37" in the same sequence appears in following cases,

**Case-1:** When 37 is at first two digits,

i.e. 3 7 \_\_

In this case, last two digits can be filled by 0 to 9, i.e. total 10 digit. So that total numbers =  $10 \times 10 = 100$ .**Case-2:** When 37 is at middle two digits,

i.e. \_\_ 3 7 \_\_

In this case, last digit can be filled by 0 to 9, i.e. total 10 digit, but first digit can be filled by 1 to 9, i.e. total 9 digit. So that total numbers =  $9 \times 10 = 90$ .**Case-3:** When 37 is at last two digits,

i.e. \_\_ \_\_ 3 7

In this case, second digit can be filled by 0 to 9, i.e. total 10 digit, but first digit can be filled by 1 to 9, i.e. total 9 digit. So that total numbers =  $9 \times 10 = 90$ .Total number =  $100 + 90 + 90 = 280$ .

So, option 'C' is correct.



Scan for Video solution

**93. (c)**

If Q is compulsory a member of team,

For 2-member team, we need to select 1 member out of 3, therefore number of 2-member team =  ${}^3C_1$ For 3-member team, we need to select 2 members out of 3, therefore number of 3-member team =  ${}^3C_2$ For 4-member team, we need to select 3 members out of 3, therefore number of 4-member team =  ${}^3C_3$ Total number of teams =  ${}^3C_1 + {}^3C_2 + {}^3C_3 = 7$ 

PQ, RQ, SQ, PRQ, PSQ, RSQ, PQRS

So, option 'C' is correct.



Scan for Video solution

**94. (a)**The area annular space =  $\pi(b^2 - a^2)$ The radius of blue circles =  $\frac{(b-a)}{2}$ The area of n blue circles =  $\frac{n\pi}{4}(b-a)^2$ 

The unpainted area in annular space

$$= \pi(b^2 - a^2) - \frac{n\pi}{4}(b-a)^2$$


$$= \pi \left[ (b^2 - a^2) - \frac{n}{4}(b-a)^2 \right]$$



Scan for Video solution



95. (b)



From the graph of  $\Delta ABC$

$$\alpha + \beta + 90^\circ = 180^\circ$$

$$\alpha + \beta = 90^\circ$$

Scan for Video solution



96. (c)

Given:  $ax^2 - bx + c = 0$

$$\Rightarrow x = \frac{-(-b) \pm \sqrt{(-b)^2 - 4 \times a \times c}}{2a} = \frac{b \pm \sqrt{b^2 - 4ac}}{2a}$$

For equal roots (one repeated root)

$$b^2 - 4ac = 0 \Rightarrow b^2 = 4ac$$

And the root is given by

$$\beta = \frac{b}{2a}$$

$$\therefore \beta^3 = \frac{b \cdot b^2}{(2a)^3} = \frac{b \cdot 4ac}{8a^3} = \frac{bc}{2a^2}$$

Scan for Video solution



97. (c)

Given: radius of circle  $r = a$

$\therefore O$  is the centre of circle

$OP = OQ = OR = OS = \text{radius of circle} = a$

The length of side 'SR' of square PQRS

$$= \sqrt{a^2 + a^2} = \sqrt{2}a$$

Area of shaded region

$$A = \text{Area of circle} - \text{Area of PQRS}$$

$$= \pi a^2 - (\sqrt{2}a)^2 = \pi a^2 - 2a^2$$



Scan for Video solution



98. (a)

$$(i) f(x) = e^x$$

$$\Rightarrow f(x_1 + x_2) = e^{x_1 + x_2}$$

$$\text{Let } x_1 = 2, x_2 = 3 \quad \therefore x > 1$$

For super additive function

$$f(x_1 + x_2) \geq f(x_1) + f(x_2)$$

$$\Rightarrow e^{x_1 + x_2} \geq e^{x_1} + e^{x_2} \Rightarrow e^{2+3} \geq e^2 + e^3$$

$$\Rightarrow e^5 \geq e^2 + e^3 \Rightarrow 148.4 \geq 27.5$$

Hence  $e^x$  is a super additive function

Similarly, we can check for  $\sqrt{x}$ ,  $\frac{1}{x}$ ,  $e^{-x}$

$$\sqrt{2+3} \geq \sqrt{2} + \sqrt{3} \Rightarrow \sqrt{5} \geq \sqrt{2} + \sqrt{3} \text{ which is wrong}$$

$$\frac{1}{2+3} \geq \frac{1}{2} + \frac{1}{3} \Rightarrow \frac{1}{5} \geq \frac{5}{6} \text{ Which is wrong}$$

$$\text{And } \frac{1}{e^{2+3}} \geq \frac{1}{e^2} + \frac{1}{e^3} \text{ which is also wrong}$$



Scan for Video solution



99. (c)

As per given information

$$f(x) = x^2$$

$$f(x) = x^2 \text{ for } x \in (-\infty, +\infty)$$

$$f(f(f(x))) = f(f(x^2)) = f(f(x^4))$$

$$= (x^4)^2 = x^8$$

$$\frac{f(f(f(x)))}{f(x)} = \frac{x^8}{x^2} = x^6 = (x^2)^3 = (f(x))^3$$



Scan for Video solution



**100. (a)****Given**

$$X \cdot Y \propto Z$$

$$X \cdot Y = C Z$$

where C is constant.

$$\text{So, for fixed } Z ; X \propto \frac{1}{Y}$$

 $\Rightarrow$  For fixed Y ;  $X \propto Z$  $\Rightarrow$  For fixed X ;  $Z \propto Y$ 

$$\Rightarrow \frac{X \cdot Y}{Z} = C \text{ (constant)}$$

So, statement in option A is wrong.

Scan for Video solution

**101. (d)**

$$S = 8 + 88 + 888 + \dots$$

$$= 8(1 + 11 + 111 + \dots)$$

$$= \frac{8}{9}(9 + 99 + 999 + \dots)$$

$$= \frac{8}{9}((10 - 1) + (100 - 1) + (1000 - 1) + \dots)$$


$$= \frac{8}{9}\{(10 + 100 + 1000 + \dots) - n\}$$

$$= \frac{8}{9} \frac{10(10^n - 1)}{10 - 1} - \frac{8}{9}n$$

$$= \frac{80}{81}(10^n - 1) - \frac{8}{9}n$$



Scan for Video solution

**102. (d)**The graph for  $y = [x]$  is drawn below.Area under graph is  $= 1 \times 1 + 2 \times 1 + 3 \times 1$  $= 6$ 

Scan for Video solution

**103. (d)**

Given two functions,

$$X = \{1, 2, 3\}$$

$$Y = \{2, 3, 4\}$$

The combinations are in which numerator belongs to X and denominator belongs to Y are given below,

$$Z = \left\{ \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{2}{2}, \frac{2}{3}, \frac{2}{4}, \frac{3}{2}, \frac{3}{3}, \frac{3}{4} \right\}$$

$$\text{Minimum value in } \{Z\} = \frac{1}{4}$$

$$\text{Maximum value in } \{Z\} = \frac{3}{2}$$

$$\therefore \text{Product} = \frac{1}{4} \times \frac{3}{2} = \frac{3}{8}$$


So, option 'D' is correct.



Scan for Video solution

**104. (c)**


Let the height of pole is L then from the condition given



$$\tan 45^\circ = \frac{L/5}{6} \Rightarrow L = 30 \text{ m}$$



Scan for Video solution



Scan for Video solution

**107. (a)**

$$\text{Given, } G(F(1,3), G(1,3))$$

$$G(F(1,3), G(1,3)) = G((1-3)^2, |1-3|)$$

$$\Rightarrow G(4, 2) = |4 - 2| = 2$$

So, option 'A' is correct.



Scan for Video solution

**108. (d)**

$$\text{Given, } f(x) = 0 \text{ have roots at } x = \{-2, 0, 3\}$$

$$\text{Now, } f(x-3) = 0$$


New function  $f(x-3) = 0$  is right shifted by 3, which indicates that value of the function, where its value is zero will be right shifted by three.

$$\therefore f(x-3) = 0 \text{ have roots at } x = \{1, 3, 6\}$$

So, option 'D' is correct.



Scan for Video solution

**109. (d)**

$$\text{Area of square} = s^2$$

$$(s)^2 = d$$

**105. (Marks to All)**

Let the number of boys =  $x$

And the number of girls =  $y$

Then,  $x = y + 2$  and  $x + y = 12$

$$\therefore x = 7 \text{ and } y = 5$$

Now, in the group of 3 children, in which number of girls will be more than number of boys,

There are only 2 possibilities for this type of group,

(i) 2 girls and 1 boy

(ii) All 3 girls

Now, the number of ways of selection of 2 girls out of 5 girls and 1 boy out of 7 boys

$$= {}^5C_2 \times {}^7C_1 = 10 \times 7 = 70$$

And, the number of ways of selection of 3 girls out of 5 girls =  ${}^5C_3 = 10$

Total ways of selection of 3 students out of 12

$$= {}^{12}C_3 = 220$$

$$\text{Required Probability} = \frac{70+10}{220} = \frac{80}{220} = \frac{4}{11}$$

So, no option is correct.

This question was 'Marks to All' in official IIT answer key.



Scan for Video solution

**106. (c)**

Total numeral number = 10

Total lower-case English alphabet = 26

Total upper-case English alphabet = 26

$$\therefore \text{Total password} = 10 \times 26 \times 26 = 6760$$

Now, these three characters can be interchanging their place in  $3!$  Ways, i.e.  $3! = 6$  ways.

$$\text{Hence, Total distinct password} = 6760 \times 6 = 40560$$

So, option 'C' is correct.

$$s = \sqrt{d}$$

$$(\sqrt{d})^2 + (\sqrt{d})^2 = (\text{diameter})^2$$

$$d + d = (\text{diameter})^2$$

$$\text{diameter} = \sqrt{2d}$$

$$\text{radius} = \frac{\sqrt{2d}}{2}$$

$$\text{Area of circle} = \pi r^2$$

$$= \pi \left( \frac{\sqrt{2d}}{2} \right)^2 = \pi \frac{2d}{4} = \pi \frac{d}{2}$$

$$= \frac{1}{2} \pi d$$



Scan for Video solution


**110. (c)**

$$P^{-x} = \frac{1}{q}$$

$$\Rightarrow \frac{1}{P^x} = \frac{1}{q}$$

$$\Rightarrow P^x = q$$

$$q^{-y} = \frac{1}{r}$$

$$\Rightarrow \frac{1}{q^y} = \frac{1}{r} \Rightarrow [q^y = r]$$

$$r^{-z} = \frac{1}{p}$$

$$\frac{1}{r^z} = \frac{1}{p}$$

$$r^z = p$$

$$P^x = q$$

$$(P^x)^y = q^y$$

$$(P^x)^y = r \quad [\because q^y = r]$$


$$((P^x)^y)^z = r^z$$

$$P^{xyz} = p \quad [\because r^z = p]$$

So,  $xyz = 1$


Scan for Video solution



**111. (a)**


In quadrilateral DCBA

$$p + q + r + s = 360^\circ \dots\dots\dots (1)$$

In  $\triangle AFD$

$$p + q + \beta = 180^\circ \dots\dots\dots (2)$$

In  $\triangle AEB$

$$p + r + \alpha = 180^\circ \dots\dots\dots (3)$$

From equations 2 & 3

$$p + q + \beta + p + r + \alpha = 360^\circ$$

From equation 1

$$p + q + \beta + p + r + \alpha = p + q + r + s$$


$$\boxed{\alpha + \beta = s - p}$$

Or,  $\alpha + \beta = \angle BCD - \angle BAD$ , or

$\angle DEC + \angle BFC = \angle BCD - \angle BAD$


Scan for Video solution



**112. (b)**


OAB and ODC are similar triangles

$$\frac{AB}{OB} = \frac{CD}{OD} \Rightarrow \frac{1.5}{3} = \frac{h}{6} \Rightarrow h = 3\text{m}$$



Scan for Video solution



113. (b)

$$na = a^2b \Rightarrow n = ab$$

$$mb = ab^2 \Rightarrow m = ab$$

$$(mn)(nm) \Rightarrow (a^2b^2)(a^2b^2) = a^4b^4$$




Scan for Video solution



114. (a)

Let the intersection point P is x distance away from tower A. Then from similar triangles.



$$\frac{x}{20} = \frac{(100-x)}{70}$$

$$70x = 2000 - 20x$$

$$90x = 2000$$

$$x = 22.22$$



Scan for Video solution



115. (a)

$$\log a + \log b + \log c = 0$$

$$\log(a \times b \times c) = 0$$

$$\text{or } a \times b \times c = 1$$

∴ integers are non-negative

∴ above condition is only possible if

$$a = b = c = 1$$

$$a + b + c = 3$$



Scan for Video solution



116. (b)

$$\frac{\log P}{y-z} = \frac{\log Q}{z-x} = \frac{\log R}{x-y} = 10$$

$$\log P = 10(y - z)$$

$$\log Q = 10(z - x)$$

$$\log R = 10(x - y)$$

$$\log P + \log Q + \log R = 10(y - z + z - x + x - y)$$

$$\log(PQR) = 0 \text{ or } PQR = 1$$



Scan for Video solution



Scan for Video solution



117. (c)

$$\frac{1}{1 + \log_u vw} + \frac{1}{1 + \log_v wu} + \frac{1}{1 + \log_w uv}$$

$$\frac{1}{\log_u u + \log_u vw} + \frac{1}{\log_v v + \log_v wu} + \frac{1}{\log_w w + \log_w uv}$$

$$\frac{1}{\log_u uvw} + \frac{1}{\log_v uvw} + \frac{1}{\log_w uvw}$$

$$\log_{uvw} u + \log_{uvw} v + \log_{uvw} w$$

$$\log_{uvw} uvw = 1$$



Scan for Video solution



Scan for Video solution



118. (b)

Let the graph is for  $y = f(x)$ .

From the graph,

$$f(0) = 0, f(1) = f(-1) = -1$$

This condition is only satisfied by  $y = |x| - 1$



Scan for Video solution



### 119. (a)

$$\log|a| + \log|b| + \log|c| = 0$$

$$|a| \cdot |b| \cdot |c| = 0$$

$$|a| \cdot |b| \cdot |c| = 1$$

Since a, b & c are integers hence the values of a, b & c can either be 1 or -1.

For  $(a+b+c)_{\min}$  we will have to put  $a=b=c=-1$

$$(a+b+c)_{\min} = -1 - 1 - 1 = -3$$

Maximum value of  $(a+b+c)$  can be obtained when  $a=b=c=1$

$$(a+b+c)_{\max} = +1 + 1 + 1 = +3.$$



Scan for Video solution



### 120. (b)

**Case - 1,** if  $x=1, y=2$

$$\frac{(x+y)-|x-y|}{2} = \frac{(1+2)-|1-2|}{2} = 1 = x$$

**Case - 2,** if  $x=2, y=1$

$$\frac{(x+y)-|x-y|}{2} = \frac{(2+1)-|2-1|}{2} = 1 = y$$

From the above two cases, it is clear that value of this expression will be minimum value of x and y.

So, option 'B' is correct.



Scan for Video solution



### 121. (c)


Given,

$$\Rightarrow e^x + 0.5x^2 - 2 = 0$$

$$\Rightarrow e^x = -0.5x^2 + 2$$

Above equation can be divided into two parts i.e. L.H.S. is an exponential function and R.H.S. is a parabola.

These two curves can be drawn on same plane. The number of intersections between them will be the roots of the given equation in the range of  $[-5, 5]$ .



The above graph shows that both curves will intersect two times each other. It means roots of the given equation will be two.

So, option 'C' is correct.

Scan for Video solution

### 122. (c)

$$\text{Total number of 3-digit number} = 999 - 99 = 900$$

Total number of 3-digits numbers with **21** together

$$\begin{array}{ccc} 21 & & 21 \\ \hline 10 \text{ possibility} & & 9 \text{ possibility} \\ (0 \text{ to } 9) \text{ unit digit} & & (1 \text{ to } 9) \text{ hundredth place} \\ \downarrow & & \downarrow \\ & & = 10 + 9 = 19 \end{array}$$

$$\text{So, the required number} = 900 - 19 = 881$$

Scan for Video solution

**123. (a)**

Sub groups containing only Indians

$$= 3C_1 + 3C_2 + 3C_3 = 3 + 3 + 1 = 7$$

Subgroups containing one Indian and rest Chinese

$$= 3C_1 [3C_1 + 3C_2 + 3C_3] = 3[3 + 3 + 1] = 21$$

Sub groups containing two Indian and remaining Chinese

$$= 3C_2 [3C_1 + 3C_2 + 3C_3] = 21$$

Sub groups containing three Indian and remaining Chinese

$$= 3C_3 [3C_1 + 3C_2 + 3C_3] = 7$$

Total number of sub groups =  $7 + 21 + 21 + 7 = 56$



Scan for Video solution



**124. (b)**

Perfect square

1, 4, 9, 16, 25, 36

So, the outcomes whose product is perfect square are  
 (1,1), (2,2), (3,3), (4,4), (5,5), (6,6),  
 (1,4) & (4,1)

$$P(E) = \frac{8}{36} \Rightarrow \boxed{\frac{2}{9}}$$



Scan for Video solution




**125. (c)**

From  $|x - 1| \leq 2$ , x can lie between

$$-1 \leq x \leq 3$$

From  $|y + 2| \leq 3$ , x can lie between


$$-5 \leq y \leq 1$$



From  $x - y \geq -2$  or  $y \leq x + 2$ ,  $y \geq 1$  and  $x \leq 3$ , we can find the area  $S_2$ .

Area of  $S_1 \cup S_2$

$$= 6 \times 4 + \frac{1}{2} \times 4 \times 4 = 24 + 8 = 32$$



**126. (d)**

Given base radius  $r = 5$  cm


Height  $h = 12$  cm

Slanted height  $l = \sqrt{5^2 + 12^2} = 13$  cm

Distance covered by base circle in  $2\pi$  = Distance covered by slanted height in  $\theta$  radians.

$$2\pi r = \theta \times l$$

$$\theta = \frac{2\pi \times 5}{13} = \frac{10\pi}{13}$$



**127. (7 to 7)**

Given,  $211^{870} + 146^{127} \times 3^{424}$

Unit place of  $211^{870}$  will be 1 always. It's because unit digit of 211 is 1. And we know that if we multiply a number which have unit digit as 1 with itself as many as times then the result will have unit digit as 1.

Unit place of  $146^{127}$  will be 6 always. It's because unit digit of 146 is 6. And we know that if we multiply a number which have unit digit as 6 with same number as many as times then the result will have unit digit as 6.

If unit place of a number is 3 and it multiplied 4 times with itself, it repeats as  $\rightarrow 3 \rightarrow 9 \rightarrow 7 \rightarrow 1 \rightarrow 3$  at unit places, which means unit place of  $3^{424}$  is 1.  
 $\therefore$  The unit digit of  $211^{870} + 146^{127} \times 3^{424}$  is  
 $= 1 + (6 \times 1) = 7$



Scan for Video solution



128. (a)

$$\text{Probability of success (p)} = \frac{60}{100} = 0.6$$

$$\text{Probability of success (q)} = 1 - 0.6 = 0.4$$

$$\therefore \text{Required probability} = {}^n C_r (p)^r (q)^{n-r}$$

$$= {}^{10} C_6 (0.6)^6 (0.4)^4 = 0.2508$$

So, option 'A' is correct.



Scan for Video solution



129. (c)

$$\text{Given, } |9y - 6| = 3$$

$$\text{Let, } f(y) = y^2 - \frac{4y}{3}$$

**Case – 1,**

$$9y - 6 = 3 \Rightarrow y = 1$$

**Case – 2,**

$$\text{At, } y = 1 \rightarrow f(y) = (1)^2 - \frac{4 \times 1}{3} = -\frac{1}{3}$$

$$\text{At, } y = \frac{1}{3} \rightarrow f(y) = \left(\frac{1}{3}\right)^2 - \frac{4}{3} \times \frac{1}{3} = -\frac{1}{3}$$

So, option 'C' is correct.



Scan for Video solution



130. (b)

Given,  $\alpha\beta = 4$

$$\therefore \frac{\alpha^n + \beta^n}{\alpha^{-n} + \beta^{-n}} = \frac{\alpha^n + \beta^n}{\frac{1}{\alpha^n} + \frac{1}{\beta^n}}$$

$$= \frac{\alpha^n + \beta^n}{\left(\frac{\beta^n + \alpha^n}{\alpha^n \beta^n}\right)} = \left(\frac{\alpha^n + \beta^n}{(\alpha\beta)^n}\right) = (\alpha\beta)^n$$

$$\therefore (\alpha\beta)^n = 4^n$$

So, option 'B' is correct.



Scan for Video solution



131. (b)

$$x^3 + 8 = 0 \Rightarrow x^3 = -8 \Rightarrow x = -2$$

$$f(-2) = 2 \times (-2)^7 + 3(-2) - 5 = 2 \times (-128) - 6 - 5$$

$$= -256 - 6 - 5$$

= -267 (not satisfies the equation)

$$x - 1 = 0 \Rightarrow x = 1$$

$$f(1) = 2(1)^7 + 3(1) - 5$$

= 2 + 3 - 5 = 0 (satisfies the equation)

$\therefore x - 1$  is factor of  $f(x)$



Scan for Video solution



132. (b)

|      |         |
|------|---------|
| Load | Failure |
|------|---------|

$$80 \quad 10^2$$

$$x \quad 5 \times 10^3$$

$$40 \quad 10^4$$

5000 is between 100 and 10000 corresponding to the loads of 80 and 40 respectively. Since the variation is exponential hence, the load should be little more than 40.



Scan for Video solution



**133. (a)**

Equation of line for abscissa  $\ln x$  and ordinate  $y$

$$y = m(\ln x) + C$$

where,  $m \rightarrow$  slope,  $C \rightarrow$  intercept at  $y$  axis


Given  $m = -0.02$  and for  $\ln x = 0.1$ ,  $y = 0$

$$\therefore 0 = -0.02 \times 0.1 + C \Rightarrow C = 0.002$$

$$\therefore y = -0.02 \ln x + 0.002$$

At  $x = 5$ ,  $\ln x = \ln 5$

$$\therefore y(5) = -0.02 \times \ln 5 + 0.002 = -0.030$$




Scan for Video solution



**134. (b)**

$$3x + 2y = 14 \Rightarrow \frac{x}{\left(\frac{14}{3}\right)} + \frac{y}{7} = 1 \quad \dots (1)$$

$$\text{And } 2x - 3y = 5 \Rightarrow \frac{x}{\left(\frac{5}{2}\right)} + \frac{y}{-\frac{5}{3}} = 1 \quad \dots (2)$$



$$\text{Equation (1)} \times 2 - \text{Equation (2)} \times 3$$

$$13y = 13 \Rightarrow y = 1$$

$$\Rightarrow x = \frac{5 + 3 \times 1}{2} = 4 \Rightarrow \text{Point B}(4, 1)$$

So, the required area is

$$A = \text{area of } \Delta ABC + \text{Area of trapezium CBDO}$$

$$= \frac{1}{2} \times 6 \times 4 + \frac{5}{2} \times 1 + \frac{1}{2} \times 1 \left( 4 - \frac{5}{2} \right)$$

$$= 12 + 2.5 + \frac{3}{4} = 15.25 \text{ square unit}$$

Scan for Video solution



**135. (c)**

Taking log

$$-a \ln q = -1 \ln r$$

$$a = \frac{\ln r}{\ln q} \quad \dots (\text{i})$$

$$-b \ln r = -1 \ln s$$

$$b = \frac{\ln s}{\ln r} \quad \dots (\text{ii})$$

$$-c \ln s = -1 \ln q$$

$$c = \frac{\ln q}{\ln s} \quad \dots (\text{iii})$$


From equations i, ii & iii

$$a \times b \times c = 1$$

Scan for Video solution



**136. (d)**



Area of one face

$$= \frac{1}{2} \text{ base} \times \text{height}$$

$$= \frac{1}{2} \times \frac{x}{4} \times \frac{x}{2} = \frac{x^2}{16}$$

So, total lateral surface area

$$= 4 \times \left( \frac{x^2}{16} \right) = \frac{x^2}{4} = 0.25 x^2$$



Scan for Video solution



### 137. (a)

Given two sets,  $A = \{2, 3, 4, 5\}$ ,  $B = \{11, 12, 13, 14, 15\}$

Total cases for selection of two numbers, one from each set  $= {}^4C_1 \times {}^5C_1 = 20$

Total cases for selection of two numbers, when sum is 16 = 4, i.e. (2, 14), (3, 13), (4, 12), (5, 11)

$$\text{Required probability} = \frac{\text{favourite}}{\text{Total}} = \frac{4}{20} = 0.20$$



So, option 'A' is correct.




Scan for Video solution



### 138. (c)



$$\therefore f(x) = f_1(x) - f_2(x) = 1 - |x|$$



$\therefore$  The above graph represents the function  $f(x)$ , which attains its maximum value at  $x = 0$  and its maximum value is 1.

So, option '(c)' is correct.

Scan for Video solution

### 139. (b)

If we subtract or multiply by a same number in an arithmetic sequence then sequence cannot be change. It will remain an arithmetic sequence, because common difference between the terms will remain same.


Therefore statement (I) and statement (II) are correct. By squaring each term of an arithmetic sequence, common difference cannot be remained same so that it cannot be an arithmetic sequence.

Therefore statement (III) is incorrect.

So, option 'B' is correct.

Scan for Video solution

### 140. (b)



Area of a triangle,

$$= \frac{1}{2} \times (\text{product of any two sides}) \times \sin(\text{angle formed by same two sides})$$

$$\therefore \text{Area}(\triangle PQR) \Rightarrow \text{Area}(\triangle PQS) + \text{Area}(\triangle PRS)$$

$$\Rightarrow \frac{1}{2}q \times r \times \sin 120^\circ = \frac{1}{2}r \times PS \times \sin 60^\circ + \frac{1}{2}q \times PS \sin 60^\circ$$


$$\Rightarrow q \times r \times \frac{\sqrt{3}}{2} = r \times PS \times \frac{\sqrt{3}}{2} + q \times PS \times \frac{\sqrt{3}}{2}$$

$$\Rightarrow q \times r = r \times PS + q \times PS$$


$$\Rightarrow q \times r = PS(q + r)$$

$$PS = \left[ \frac{qr}{r+q} \right]$$

So, option 'B' is correct.



**141. (b)**



If 'x = 0, then 'y' = 0

If 'x' = 2, then 'y' = -1

Option A  $\Rightarrow x = y - |y|$

$$2 = -1 - |-1| \Rightarrow 2 = -1 - 1$$

2 = -2 not possible

Option B  $\Rightarrow x = -(y - |y|)$

$$2 = -(-1 - |-1|)$$

2 = 2 correct

Option C  $\Rightarrow x = y + |y|$


$$2 = -1 + |-1|$$

$2 = -1 + 1 = 0$  not possible

Option D  $\Rightarrow x = -(y + |y|)$

$$2 = -(-1 + |-1|)$$

$2 = 0$  not possible.




**142. (a)**

We know  $\log_b a = x \Rightarrow b^x = a$

$$\therefore \log_x \left( \frac{5}{7} \right) = -\frac{1}{3} \Rightarrow (x)^{-\frac{1}{3}} = \frac{5}{7}$$

$$\Rightarrow (x)^3 = \frac{7}{5} \Rightarrow (x) = \left( \frac{7}{5} \right)^3 = \frac{343}{125}$$



**143. (c)**

$-4 \leq x \leq 5$ ; x can have 10 values

$6 \leq y \leq 16$ ; y can have 11 values.

Coordinates of P can be chosen from these values & the number of different coordinates of P will be

$${}^{10}C_1 \times {}^{11}C_1 = 110$$

Since PR line is parallel to x-axis so y-coordinate of R is same as P so the number of different coordinates of R can be


$${}^9C_1 = 9 \quad \begin{cases} \because \text{Coordinate of } x \text{ can not be} \\ \text{Same as } x \text{ coordinate of } P. \end{cases}$$

Similarly, for Q, x-coordinate will be same as x-coordinate of P so number of different coordinates of Q is

$${}^{10}C_1 = 10 \quad \begin{cases} \because \text{Coordinate of } y \text{ can not be} \\ \text{Same as } y \text{ coordinate of } P. \end{cases}$$

total number of triangles =  $110 \times 9 \times 10$

$$= 9900$$



**144. (c)**


$$\tan 89^\circ = \cot 1^\circ$$

$$\tan 88^\circ = \cot 2^\circ$$

so

$$\log \tan 1^\circ + \log \tan 2^\circ + \dots + \log \tan 89^\circ$$

$$\begin{aligned}
 &= \log (\tan 1^\circ, \tan 2^\circ, \tan 3^\circ, \dots, \tan 45^\circ, \dots, \cot 2^\circ, \cot 1^\circ) \\
 &= \log (\tan 45^\circ) = \log 1 = 0
 \end{aligned}$$

**145. (a)**

For  $x > y > 1$


$\ln x > \ln y$  and  $e^x > e^y$

but  $y^x > x^y$  is not always true

If  $x = 3$  &  $y = 2 \Rightarrow 2^3 < 3^2$

but when  $x = 5$  &  $y = 2$

$2^5 > 5^2$  and  $\cos x < \cos y$

**146. (b)**

$$(a + b + c)^2 \geq 0$$

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) \geq 0$$

$$1 + 2(ab + bc + ca) \geq 0$$

$$ab + bc + ca \geq -1/2 \quad \text{(1)}$$


$$(a - b)^2 + (b - c)^2 + (c - a)^2 \geq 0$$

$$2(a^2 + b^2 + c^2) - 2(ab + bc + ca) \geq 0$$

$$(a^2 + b^2 + c^2) - (ab + bc + ca) \geq 0$$


$$1 - (ab + bc + ca) \geq 0$$

$$(ab + bc + ca) \leq 1 \quad \text{(2)}$$

**147. (b)**

Given that first two drawn cards are kings then there will be 2 kings remaining in the pack of 50. So, the probability of 3<sup>rd</sup> drawn card is king, is

$$P = \frac{2}{50}$$

**148. (d)**

Given a quadratic equation,  $ax^2 + bx + c = 0$  has real and positive roots,

The roots of a quadratic equation are real and positive if  $b^2 - 4ac > 0$ ,  $b^2 - 4ac < -b$ ,  $b < 0$


$$\text{Now, we know that } |x| = \begin{cases} x; & x > 0 \\ -x; & x < 0 \end{cases}$$

Then, we will get two different equations,

$$ax^2 + bx + c = 0 \text{ and } ax^2 - bx + c = 0$$

As above we find that,  $b^2 - 4ac > 0$ , will remain same for both equations, which indicates that both roots of the 2<sup>nd</sup> equation will be real. So that we will get total 4 real roots.

So, option 'D' is correct.

**149. (d)**

Given that  $x$  is a real number, then

$$|x^2 - 2x + 3| = 11 \Rightarrow \begin{cases} x^2 - 2x + 3 = 11 \\ -(x^2 - 2x + 3) = 11; \end{cases}$$

**Case – 1,**

$$x^2 - 2x + 3 = 11$$

$$x^2 - 2x - 8 = 0$$

$$x = 4, -2$$

**Case – 2,**

$$-(x^2 - 2x + 3) = 11$$

$$x^2 - 2x + 14 = 0$$

Above equation will have only complex roots, i.e. no real roots.

Then,

$$|-x^3 + x^2 - x| \text{ at } x = 4;$$

$$\Rightarrow |-64 + 16 - 4| = 52$$

$$|-x^3 + x^2 - x| \text{ at } x = -2;$$

$$\Rightarrow |8 + 4 + 2| = 14$$

So, option 'D' is correct.



Scan for Video solution



### 150. (c)

$$\text{Given, } (7526)_8 - (Y)_8 = (4364)_8$$

$$\therefore (Y)_8 = (7526)_8 - (4364)_8$$

$$= (7526 - 4364)_8 = (3142)_8$$

So, option 'C' is correct.



Scan for Video solution



### 151. (c)

The derivative of a function gives us the slope of the line tangent to the function at any point on the graph.

$$y = 5x^2 + 3$$

$$\frac{dy}{dx} = 10x$$

$$\frac{dy}{dx} = 0 \text{ (at } x = 0, y = 3\text{)}$$

Since, slope is 0 hence, the tangent is parallel to the x-axis.



Scan for Video solution



### 152. (180 to 180)

Smallest angle of quadrilateral  $\rightarrow q$

Smallest angle of triangle  $\rightarrow t$

$$t = \frac{2}{3}q$$

Quadrilateral: -

$$3 : 4 : 5 : 6$$

$$3x + 4x + 5x + 6x = 360^\circ$$

$$x = 20^\circ$$

$$q = 3 \times 20 = 60^\circ$$

$$t = \frac{2}{3}(60^\circ) = 40^\circ$$

$$\begin{aligned} \text{Largest angle of } \Delta &= 2 \times \text{smallest angle of } \Delta \\ &= 2 \times t = 2 \times 40^\circ = 80^\circ \end{aligned}$$

$$\text{Second largest angle of } \Delta + \text{largest angle of quadrilateral} = 60 + 6x \quad (x = 20^\circ) = 180$$



Scan for Video solution



### 153. (25 to 25)

$$P(\text{magnitude} > 6) = 0.04$$

Average time interval: -

$$T = \frac{1}{P} = \frac{1}{0.04} = \frac{100}{4} = 25 \text{ years}$$



Scan for Video solution



### 154. (a)

$$\text{Given: } f(x, y) = x^n y^m = P$$

$$f(x', y')$$

$$= (2x)^n \times \left(\frac{y}{2}\right)^m$$

$$= 2^n \times x^n \times y^m \times 2^{-m}$$

$$= 2^{n-m} \times x^n \times y^m = 2^{n-m} P$$



Scan for Video solution



### 155. (a)

Number of days in a leap year = 366

Number of weeks in a leap year = 52 weeks + 2 days

It means that there are two odd days in a leap year, Then there are two possibilities that 53 Saturdays in a leap year.

**Case – 1:** when a leap year starts with Saturday then 365<sup>th</sup> day will be 53<sup>rd</sup> Saturday.

**Case – 2:** when a leap year starts with Friday then 366<sup>th</sup> day will be 53<sup>rd</sup> Saturday.

A leap year can start with any day out of 7, so that total cases of start a leap year is 7. Out of 7, only two cases are those which have 53 Saturday.

$$\therefore \text{Required Probability} = \frac{2}{7}$$

So, option ‘A’ is correct.



Scan for Video solution



### 156. (b)

Given a quadratic equation,

$$3x^2 + 2x + p(p-1) = 0$$

The roots of a quadratic equation are of opposite sign, if product of both roots will be negative or less than zero.

Let roots of a quadratic equation  $ax^2 + bx + c = 0$ ,

are  $\alpha$  and  $\beta$ , then product of root will be  $\alpha\beta = \frac{c}{a}$ , then

above information can be written as,

$$\frac{p(p-1)}{3} < 0$$

$$p(p-1) < 0$$

Using wavy curve,



Then, product of roots will be negative, if value of  $p$  will lie in the interval of  $(0, 1)$ .

So, option ‘B’ is correct.



Scan for Video solution



### 157. (d)

Let the sum equal to ‘S’; then

$$S = 10 + 84 + 734 + \dots$$

$$S = (9+1) + (9^2+3) + (9^2+5) \dots$$

$$S = (9 + 9^2 + 9^3 \dots) + (1 + 3 + 5 \dots)$$

$$S = \frac{9(9^n - 1)}{(9-1)} + \frac{n}{2} [2 \times 1 + (n-1) \times 2]$$

$$S = \frac{9(9^n - 1)}{8} + n^2$$

So, option ‘D’ is correct.



Scan for Video solution



### 158. (a)

$$2x + y \leq 6 \dots\dots(1)$$

$$x + 2y \leq 8 \dots\dots(2)$$


Equation (1) - 2×equation 2

$$3y \leq 10 \Rightarrow y \leq \frac{10}{3}$$

Putting  $y = \frac{10}{3}$  in equation (1)

$$2x + \frac{10}{3} \leq 6 \Rightarrow x \leq \frac{4}{3}$$

(4/3, 10/3)



### 159. (b)

$$\text{From } |4X-7|=5, X = 3 \text{ or } 1/2$$

$$2|X| - |-X| = 2|X| - |X|$$

$$2|X| - |X|$$

$$\begin{array}{ccc} & \text{for } x = 3 & \text{for } x = 1/2 \\ & \swarrow & \searrow \\ 6-3 & = 3 & 1-1/2 \\ & & = 1/2 \end{array}$$



Scan for Video solution




Scan for Video solution



**160. (b)**

$$\begin{aligned} \text{Sum} &= \frac{1}{\sqrt{2} + \sqrt{1}} + \frac{1}{\sqrt{3} + \sqrt{2}} + \dots + \frac{1}{\sqrt{81} + \sqrt{80}} \\ &= \frac{\sqrt{2} - \sqrt{1}}{(\sqrt{2})^2 - (\sqrt{1})^2} + \frac{\sqrt{3} - \sqrt{2}}{(\sqrt{3})^2 - (\sqrt{2})^2} + \dots + \frac{\sqrt{81} - \sqrt{80}}{(\sqrt{81})^2 - (\sqrt{80})^2} \\ &= \sqrt{2} - \sqrt{1} + \sqrt{3} - \sqrt{2} + \sqrt{4} - \sqrt{3} + \sqrt{5} - \sqrt{4} \dots \\ &\quad \dots + \sqrt{81} - \sqrt{80} \\ \text{Sum} &= \sqrt{81} - \sqrt{1} = 9 - 1 = 8 \end{aligned}$$



**161. (b)**

$$y = 2x - 0.1x^2$$

→ For  $x = 8$  m

$$y = 16 - 0.1 \times 64 = 9.6 \text{ m}$$

→ For  $x = 10$  m

$$y = 20 - 0.1 \times 100 = 10 \text{ m}$$


→ For  $x = 12$  m

$$y = 24 - 0.1 \times 144 = 9.6 \text{ m}$$

→ For  $x = 14$  m


$$y = 28 - 0.1 \times 196 = 8.4 \text{ m}$$

So, the maximum height is 10 m when  $x = 10$




**162. (c)**

The wait duration cannot be more than 15 minutes, which is applicable for both 'A' and 'B'; the above information can be drawn into a graph given below, unfavourite



Then the probability that 'A' and 'B' will meet on that day =  $\frac{\text{Favourite area}}{\text{Total area}} = \frac{7}{16}$

So, option 'C' is correct.



**163. (a)**

$$CP = 5q^2, SP = 50q, P = SP - CP$$

$$P = 50q - 5q^2$$

→ For  $q = 5$

$$P = 50 \times 5 - 5 \times 25 = 125$$

→ For  $q = 10$


$$P = 500 - 500 = 0$$

→ For  $q = 15$

$$P = 50 \times 15 - 5(15)^2 = -375$$

→ For  $q = 25$

$$P = 50 \times 25 - 5 \times 625 = -1875$$



**164. (c)**

$$= 4 + 44 + 444 \dots$$

$$= \frac{4}{9}[9 + 99 + 999 \dots]$$


$$= \frac{4}{9}[(10-1) + (10^2 - 1) + (10^3 - 1) \dots]$$

$$= \frac{4}{9}[10 + 10^2 + 10^3 \dots - 1 - 1 - 1 \dots]$$

$$= \frac{4}{9} \left[ \frac{10(10^n - 1)}{(10-1)} - n \right]$$

$$= \frac{4}{81}[10^{n+1} - 9n - 10]$$

So, option 'C' is correct.



**165. (d)**

$$\text{Given, } f(y) = \frac{|y|}{y}$$

$$\text{Then, } |f(q) - f(-q)| = \left| \frac{|q|}{q} - \frac{|-q|}{-q} \right| \\ = \left| \frac{|q|}{q} + \frac{|-q|}{q} \right| = \left| \frac{|q|}{q} + \frac{|q|}{q} \right|$$

$\therefore |-q| = |q| = q$ ;  $q$  is non-zero real number.

$$|f(q) - f(-q)| = \left| \frac{q}{q} + \frac{q}{q} \right| = 2$$

So, option 'D' is correct.



Scan for Video solution

**166. (a)**

$$\text{Given data: } V = 4q; \quad F = 100/q$$

$$\rightarrow V + F = 4q + \frac{100}{q}$$

$$\text{For } q = 5 \rightarrow V + F = 20 + 20 = 40$$

$$\text{For } q = 4 \rightarrow V + F = 16 + 25 = 41$$

$$\text{For } q = 7 \rightarrow V + F = 28 + 14.28 = 42.28$$

$$\text{For } q = 6 \rightarrow V + F = 24 + 16.\bar{6} = 40.\bar{6}$$

So minimum  $(V + F)$  is for  $q = 5$



Scan for Video solution

**167. (b)**

$$\log P = \frac{1}{2} \log Q = \frac{1}{3} \log R = K$$

$$\frac{1}{2} \log Q = K \Rightarrow \log Q = 2K \Rightarrow Q = 10^{2K}$$

$$Q^2 = (10^{2K})^2 = 10^{4K}$$

$$\text{Similarly, } P \times R = 10^K \times 10^{3K} = 10^{4K}$$

$$\text{Hence, } Q^2 = PR$$



Scan for Video solution

**168. (c)**

$$\begin{array}{r} 137 \\ +276 \\ \hline 435 \end{array}$$

In base 10,  $137 + 276 = 413 < 435$ , so the base is other than 10. To find the base we add unit digits (7 + 6) of the numbers and find out which base can give us 5 in the unit digit of summed value.

$7 + 6 = 13 = 8 + 5$ , so the base is 8, now

$$\begin{array}{r} 731 \\ +672 \\ \hline 1623 \end{array}$$



Scan for Video solution



Scan for Video solution

**169. (a)**

Here, if 3 multiplied 4 times with itself, it repeats  $\rightarrow 3 \rightarrow 9 \rightarrow 7 \rightarrow 1$  at unit places, and again 3 which means unit place of  $3^{999}$  is 7

Now, if 7 multiplied 4 times with itself it repeats  $\rightarrow 7 \rightarrow 9 \rightarrow 3 \rightarrow 1$  at unit place and again 7 on next move, which means unit place of  $7^{1000}$  is 1.

$$\therefore 3^{999} \times 7^{1000}$$

$$= 7 \times 1 = 7$$


$\therefore$  Option (a) is correct.



Scan for Video solution

**170. (c)**

Triangle in the half portion of the curve = 8



$$\text{Total} = 8 \times 2 = 16$$

But when we add both half portions then 4 triangles are also formed,

$$\therefore \text{Total} = 16 + 4 = 20$$

$\therefore$  Option (c) is correct.



Scan for Video solution



### 171. (a)

$$\text{Given } F_{n+1} = F_n + F_{n-1}$$

If we put  $n = 6$

$$F_7 = F_6 + F_5$$

$$F_5 = F_7 - F_6 = 60 - 37 = 23$$

$$F_4 = F_6 - F_5 = 37 - 23 = 14$$

$$F_3 = F_5 - F_4 = 23 - 14 = 9$$

$$F_2 = F_4 - F_3 = 14 - 9 = 5$$

$$F_1 = F_3 - F_2 = 9 - 5 = 4$$



Scan for Video solution



### 172. (a)

The percentage of pregnant women received medical care from doctor at least once a year.

$$= \left( \frac{60}{100} \times \frac{90}{100} \right) \times 100$$

$$= 54\%$$

So, more than half of the pregnant women received medical care at least once from a doctor.



Scan for Video solution




### 173. (a)

To find the maximum number of testing kit needed, we have to take worst case scenario i.e, there is only one covid positive among each set of 5 people.

$$\text{Total number of sets} = \frac{1000}{5} = 200$$

So, 200 test kit is needed to first identify the positive 100 sets of 5. Also 5 test kits are needed to identify positive among each set.


$$\text{So, Total maximum kits required} = 200 + 100 \times 5 = 700$$



### 174. (c)


Only the numbers 123457869 and 123456789 have distinct digits whose sum add to 45.

Out of these two 123456789 is smallest and hence option (c) is correct.



### 175. (d)

In the top row any one of the three cells can be shaded so, number of ways is 3. Now that a cell in the top row is shaded, we have only 2 number of ways (cells) in which a cell in second row can be shaded. And similarly, there is only one cell which we can shade in the third row. So, total number of ways =  $3 \times 2 \times 1 = 6$



**176. (a)**

Assuming husband-wife combination as one then total number of ways in which these 3 pairs can be arranged is  $2!$  and the positions of husband wife in each pair can be interchanged in 2 ways and for all three pairs  $2^3$  so, total number of possible arrangements are

$$\Rightarrow 2! \times (2)^3 = 16$$



Scan for Video solution

**177. (c)**

$$4^{8^x} = 4^4$$

$$8^x = 4$$

$$(2^3)^x = 2^2$$

$$2^{3x} = 2^2$$

$$\Rightarrow 3x = 2 \quad \Rightarrow \quad x = \frac{2}{3}$$



Scan for Video solution

**178. (a)**

From 9:15 to 9:16 minute hand & second hand will cross each other once, from 9:15 to 9:17 they will cross twice, from 9:15 to 9:18 they will cross thrice, similarly from 9:15 to 9:45 they will cross 30 times.



Scan for Video solution

**179. (b)**

Consider the price of an item in store S = x  
And the price of same item in store M = y  
Therefore,  $x = 0.9y$

Store S charges ₹ 150 for delivery,  
So, as per given condition,

$$y - (x + 150) = 100$$

$$y - 0.9y - 150 = 100$$

$$0.1y = 250$$

$$y = 2500 \text{ ₹}$$

∴ The price of that item in store S = x

And, we know that  $x = 0.9y = 0.9 \times 2500 = 2250 \text{ ₹}$

∴ option (b) is correct.



Scan for Video solution

**180. (a)****Given Data:**

We have to find the LCM of individual times of every bell to find the same time of the bells to ring together,

$$20 = 2 \times 2 \times 5$$

$$30 = 2 \times 3 \times 5$$

$$50 = 2 \times 5 \times 5$$

Therefore, LCM of 20, 30, 50

$$= 2 \times 2 \times 3 \times 5 \times 5 = 300 \text{ minutes} = 5 \text{ hours}$$

It means they will ring together after 5 hours of 12 pm i.e. at 5 pm.

∴ option (a) is correct.



Scan for Video solution

**181. (a)**

Let the speeds of P and Q are  $3x \text{ m/sec}$  and  $4x \text{ m/sec}$  respectively.

Total distance of the race = 500 m

Leading distance of P = 140 m

Distance to be covered by P =  $500 - 140 = 360 \text{ m}$

Time taken by P to finish the race after Q starts to

$$\text{run} = \frac{360}{3x} = \frac{120}{x} \text{ sec}$$

$$\text{Distance covered by Q in } \frac{120}{x} \text{ sec}$$

$$= \frac{120}{x} \times 4x = 480 \text{ m}$$

$$\therefore \text{P wins the race by } 500 - 480 = 20 \text{ m.}$$

∴ option (a) is correct.



Scan for Video solution



**182. (d)**

If  $r$  is the root of the equation  $x^2 + 2x + 6 = 0$ , then it satisfies the equation  $r^2 + 2r + 6 = 0$

$$r^2 = -2r - 6$$

$$\begin{aligned} \Rightarrow & (r+2)(r+3)(r+4)(r+5) \\ = & (r^2 + 5r + 6)(r^2 + 9r + 20) \\ = & (-2r - 6 + 5r + 6)(-2r - 6 + 9r + 20) \\ = & 3r(7r + 14) \\ = & 21(r^2 + 2r) = 21 \times (-6) = -126 \end{aligned}$$



Scan for Video solution

**183. (d)**

Given, the boys constituted 65% of those who appeared and girls accounted for 60% of the qualified candidates. Let total candidates are  $x$  and qualified candidates are  $y$ , then number of girls and number of boys who appeared is given by:

$$\left. \begin{array}{l} B_A = 0.65x \\ G_A = 0.35x \end{array} \right\} B_A > G_A$$


The number of boys and girls who qualified is given by:

$$\left. \begin{array}{l} B_Q = 0.4y \\ G_Q = 0.6y \end{array} \right\} B_Q < G_Q$$

From above analysis it is clear that the number of boys who qualified the test is less than the number of girls who Qualified.



Scan for Video solution

**184. (d)**

So total probability of 2<sup>nd</sup> drawn ball is orange.

$$\begin{aligned} &= \frac{3}{5} \times \frac{2}{4} + \frac{2}{5} \times \frac{2}{5} \\ &= \frac{6}{20} + \frac{4}{25} = \frac{30+16}{100} = \frac{46}{100} = \frac{23}{50} \end{aligned}$$



Scan for Video solution

**185. (d)**

**Given:**  $P : Q : R : S = 5 : 2 : 4 : 3$

Let  $P'$  share =  $5x$ ,  $Q'$  share =  $2x$ ,  $R'$  share =  $4x$   
 $S'$  share =  $3x$

$\therefore R'$  share =  $1000 + S'$  Share

$$4x = 1000 + 3x \Rightarrow x = 1000$$

$$\therefore Q' \text{ share} = 2x = 2 \times 1000 = ₹ 2000$$



Scan for Video solution

**186. (d)**

**Given:**  $u$  is 50% more than  $s$

$$\Rightarrow u = 1.5s \Rightarrow \frac{s}{u} = \frac{1}{1.5} \Rightarrow s : u = 1 : 1.5$$

$$\therefore p : q = 1 : 2 = 6 : 12$$

$$q : r = 4 : 3 = 12 : 9$$

$$\Rightarrow p : q : r = 6 : 12 : 9 = 24 : 48 : 36$$

$$r : s = 4 : 5 = 36 : 45$$

$$\Rightarrow p : q : r : s = 24 : 48 : 36 : 45$$

$$s : u = 45 : 67.5$$

$$\Rightarrow p : q : r : s : u = 24 : 48 : 36 : 45 : 67.5$$

$$\therefore p : u = 24 : 67.5 = 240 : 675 = 16 : 45$$



Scan for Video solution



### 187. (b)

PQ is tangent to circle so,  $\angle PQR = 90^\circ$

So, in right angled  $\triangle PQR$

$$(PR)^2 = (PQ)^2 + (QR)^2 \quad [\text{Pythagoras theorem}]$$

$$\text{Area of } \triangle PXW = \text{Area of } \triangle SPQT + \text{Area of } \triangle RZVU \\ (\text{Since } QR = RZ \text{ (radius)})$$

$$\text{Area of } \triangle SPQT = \text{Area of } \triangle PXW - \text{Area of } \triangle RZVU$$



Scan for Video solution



### 188. (b)

Let profit be 'T'

$$\text{Total investment by P} = 5000 \times 6 = 30000$$

$$\text{Total investment by Q} = x \times 8 = 8x$$

$$Q \text{ receives profit} = \frac{4}{9}T \Rightarrow P \text{ receives profit} = \frac{5}{9}T$$

$$\frac{\text{Total investment by P}}{\text{Total investment by Q}} = \frac{P \text{ 's profit}}{Q \text{ 's profit}}$$

$$\frac{30000}{8x} = \frac{5/9T}{4/9T} \Rightarrow x = 3000$$



Scan for Video solution



192. (c)

Given,

For Past  $m$  days, average daily production = 100 units

Today's production = 180

New average daily production

$$110 = \frac{100 \times m + 180}{m + 1}$$

$$110m + 110 = 100m + 180$$

$$10m = 70$$

$$m = 7$$



Scan for Video solution



193. (a)

5 Mondays in a month will require at least 29 days, and for that, the month should start with Monday.

So, 2<sup>nd</sup> Feb will be Tuesday and 1, 8, 15, 22, 29 will be the dates having Monday.



Scan for Video solution



194. (a)

(Marks to all in Answer key)

3pm to 11pm, there will be 8 coincidences.

From 11pm to 1am there will be only 1 coincidence.




From 1am to 3am there will 2 coincidences. Hence total of 11 coincidences.



Scan for Video solution



195. (b)



Let

$$A_{\Delta} = A_{O} = A_{\square} = K$$

$$\frac{\sqrt{3}}{4} a^2 = \pi r^2 = b^2 = K$$

$$a = \sqrt{\frac{4K}{\sqrt{3}}}, r = \sqrt{\frac{K}{\pi}}, b = \sqrt{K}$$

$$P_{\Delta} : P_{\square} : P_O$$

$$3a : 4b : 2\pi r$$

$$3\sqrt{\frac{4K}{\sqrt{3}}} : 4\sqrt{K} : 2\pi\sqrt{\frac{K}{\pi}}$$

$$2\sqrt{3}\sqrt{3} : 4 : 2\sqrt{\pi}$$

$$\sqrt{3}\sqrt{3} : 2 : \sqrt{\pi}$$




Scan for Video solution



196. (a)

Since the sides of rhombus are of equal length & are perpendicular to each other hence rhombus is a square here.



The side of rhombus is  $\frac{1}{\sqrt{2}}$

So, the diameter of largest circle drawn inside rhombus is  $\frac{1}{\sqrt{2}}$



Scan for Video solution



**197. (c)**

Let distance of first part of the 80 km journey is  $x$  km for which the speed is 10 kmph. The speed for second part is 18 kmph.

$$\text{Total time} = \frac{x}{10} + \frac{(80-x)}{18}$$

$$6 = \frac{x}{10} - \frac{x}{18} + \frac{80}{18}$$

$$x = \left(6 - \frac{80}{18}\right) \times \frac{10 \times 18}{8}$$

$$x = 35$$

$$\text{Percentage} = \frac{35}{80} = 43.75\%$$



Scan for Video solution

**198. (a)**

Given,

$$\frac{M+N+S}{3} = 4,000$$

Or

$$M+N+S = 12,000 \quad \dots(a)$$

$$\frac{N+S+P}{3} = 5,000$$

$$N+S+P = 15,000 \quad \dots(b)$$

From equation (b)

$$N+S = 15,000 - 6,000 = 9,000$$

From equation a

$$M = 12,000 - 9,000 = 3000$$

$$\frac{M}{P} = \frac{3000}{6000} = 50\%$$



Scan for Video solution

**199. (c)**

As each inner square is formed by joining the mid-point of the side of the outer square then the ratio of the inner square side to outer square side =  $\frac{1}{\sqrt{2}}$

This process repeats itself 5 times,

$$\text{Therefore, the side of smallest square} = \frac{10}{(\sqrt{2})^5} \text{ cm}$$

Then the area of smallest square,

$$= \left( \frac{10}{(\sqrt{2})^5} \right)^2 = 3.125 \text{ cm}^2$$

So, option 'C' is correct.



Scan for Video solution

**200. (d)**

$$\text{Using } a^2 - b^2 = (a+b)(a-b),$$

Only option (d) can be factorize using this property,

$$\begin{aligned} (11^{52} - 1) &= (11^{26})^2 - (1)^2 = (11^{26} + 1)(11^{26} - 1) \\ &= (11^{26} + 1)(11^{13} - 1)(11^{13} + 1) \end{aligned}$$

As we can see that,  $(11^{13} + 1)$  is the factor of  $(11^{52} - 1)$ ,

So that,  $(11^{52} - 1)$  is exactly divisible by  $(11^{13} + 1)$ .

So, option 'D' is correct.



Scan for Video solution

**201. (d)**

$$\text{Sum of all angles of 'n' side polygon} = (n-2) \times 180^\circ$$

Given regular polygon of 10 sides,

$\therefore$  Interior angle between two sides,

$$= \left( \frac{n-2}{n} \right) \times 180^\circ = \left( \frac{10-2}{10} \right) \times 180^\circ = 144^\circ$$

So, option 'D' is correct.



Scan for Video solution



### 202. (c)

The ratio of boys and girls in the class is 7 : 3. Let us say the number of boys and girls are  $7x$  and  $3x$  respectively. Then total number of students  $= 7x + 3x = 10x$

So, the number of students should be multiple of 10 hence option (c) 50 is correct



Scan for Video solution



### 203. (c)

The radius of cylinder

$$r = \frac{6}{2\pi} \text{ cm}$$

Length of the cylinder  $= l = 1$  cm

$$\text{Volume of cylinder} = \pi r^2 \times l = \pi \left( \frac{36}{4\pi^2} \right) \times 1$$

$$V_{\text{cylinder}} = \frac{9}{\pi} \text{ cm}^3$$

The sheet  $6 \text{ cm} \times 1 \text{ cm}$  can be cut into 6 small patches of  $1 \text{ cm} \times 1 \text{ cm}$  each to form a cube. The volume of cube

$$V_{\text{cube}} = 1 \times 1 \times 1 = 1 \text{ cm}^3$$

$$\text{So, } \frac{V_{\text{cylinder}}}{V_{\text{cube}}} = \frac{9}{\pi}$$



Scan for Video solution



### 204. (c)

$$\frac{\text{cost of P}}{\text{cost of Q}} = \frac{3}{4} = \frac{5400}{\text{cost of Q}}$$

$$\text{Cost of Q} = 7200$$

Since profit on Q is 25% so, selling price of Q ( $SP_Q$ )

$$SP_Q = 7200 \times 1.25$$

$$SP_Q = 9000$$

Discount on Q

$$= \frac{MP_Q - SP_Q}{MP_Q} = \frac{10000 - 9000}{10000} = 10\%$$


Scan for Video solution



### 205. (c)

The probability of no chocolate identical

$$P = \frac{10 \times 9 \times 8 \times 7 \times 6}{10^5} = 0.3024$$

The probability of at least two chocolates identical

$$Q = 1 - P$$

$$= 1 - 0.3024$$

$$= 0.6976$$


Scan for Video solution



### 206. (c)

Given the ratio of students in classes A, B and C are 3 : 13 : 6. Let the students in these classes are

$$A \rightarrow 3x, B \rightarrow 13x, C \rightarrow 6x$$

$$(3x + 18) : (13x + 18) : (6x + 18) = 15 : 35 : 21$$

$$\frac{3x + 18}{13x + 18} = \frac{15}{35} = \frac{3}{7}$$

$$21x + 126 = 39x + 54$$

$$18x = 72$$

$$x = 4$$

Then total number of students at the beginning  
 $= 3x + 13x + 6x = 22x$   
 $= 22 \times 4 = 88$



Scan for Video solution



So left ways =  $3 \times 3! = 18$




Scan for Video solution



210. (d)

| PQ       | QR       | RP       |
|----------|----------|----------|
| 4        | 6        | 8        |
| a        | a        | a        |
| <u>a</u> | <u>a</u> | <u>a</u> |
| 4        | 6        | 8        |

Let side of equilateral  $\Delta = a$



$$\frac{a}{4} \quad \frac{q}{6} \quad \frac{a}{8}$$

LCM of (4, 6, 8)  $\rightarrow 24$

$$\begin{aligned} \text{Area of equilateral } \Delta &= \frac{\sqrt{3}}{4} (\text{LCM})^2 \\ &= \frac{\sqrt{3}}{4} (24)^2 \\ &= 144\sqrt{3} \end{aligned}$$



Scan for Video solution



211. (c)

From the diagram below,

$$\sin 30^\circ = \frac{r}{R} = \frac{1}{2}$$

$$\frac{\pi r^2}{\pi R^2} = \left( \frac{r}{R} \right)^2 = \frac{1}{4}$$

207. (c)


Let the population 2 years ago was x.  
 $\therefore x \times 1.05 \times 1.05 = 1102500$   
 $\Rightarrow x = 10,00,000$



Scan for Video solution



208. (a)



Given,

$$PR = PS = QS = QT = RT$$

So inside it will make regular pentagon each angle of

$$\text{reg pentagon} = \frac{540}{5} = 108^\circ$$

$$\angle PAB = 180^\circ - 108^\circ = 72^\circ$$

$$\angle PBA = 180^\circ - 108^\circ = 72^\circ$$

$$\angle APB = \theta = 180^\circ - 72^\circ - 72^\circ = 36^\circ$$




Scan for Video solution



209. (c)

1<sup>st</sup> 2<sup>nd</sup> 3<sup>rd</sup> 4<sup>th</sup>  
In 2<sup>nd</sup> position R cannot sit



Scan for Video solution

**212. (b)**

The probability of first selected ball is blue & second selected ball is black, is.

$$= \frac{^{15}C_1 \times ^{45}C_1}{^{60}C_1 \times ^{59}C_1} = \frac{15 \times 45}{60 \times 59} = \frac{45}{236}$$



Scan for Video solution

**213. (a)**

$$30 = 5 \times 2 \times 3$$

$$32 = 2^5$$

The LCM of 30 & 32 is  $= 2^5 \times 3 \times 5 = 480$  seconds or 8 minutes

Hence next beep together will be at 10:08 AM



Scan for Video solution

**214. (b)**

For the given conditions, the possible arrangements are:

|   |   |   |   |   |
|---|---|---|---|---|
| Q | R | P | T | S |
| S | R | P | T | Q |
| S | R | T | P | Q |

So, total number of possibilities is 3.



Scan for Video solution

**215. (d)**

Fixing the positions of Q & R

$$Q \_ R \_ \_ \Rightarrow \text{possibilities: } 2 \times 2 \times 1 = 4$$

$\because$  S cannot sit adjacent to Q so, at position 2, there can be 2 possibilities ( $P/T$ ) , in 4 there are 2 possibilities ( $P$  or  $T/S$ )

$$\_ Q \_ R \_ S \Rightarrow \text{possibilities: } 2 \times 1 \times 1 = 2$$

$$S \_ Q \_ R \Rightarrow \text{possibilities: } 2 \times 1 \times 1 = 2$$

On flipping the positions of Q and R, similarly there will also be 8 possibilities hence total 16 possibilities.



Scan for Video solution

**216. (c)**

In farm Q

$$5x + 14x + 13x = 416$$

$$32x = 416$$

$$x = 13$$

After sending to farm P

$$\text{Hens} = 65 + 5 \times 13 = 130$$

$$\text{Ducks} = 91 + 14 \times 13 = 273$$

$$\text{Goats} = 169 + 13 \times 13 = 338$$


$$H : D : G = 10 : 21 : 26$$



Scan for Video solution

**217. (d)**

Given area of triangle =  $50 \text{ cm}^2$



$$\frac{1}{2}r^2 = 50$$

$$r = 10 \text{ cm}$$

$$\text{Area of circle} = \pi r^2 = 100\pi$$



Scan for Video solution



### 218. (b)

$$\text{Holding period in 1993} = 360 \text{ days}$$

As per the given data, holding period was revised 3 times, each time it was reduced by one quarter. It means it was reduced by 3 quarter in all 3 revision.

$$\text{Number of days in one quarter} = 90$$

$$\therefore \text{Days in three quarter} = 90 \times 3 = 270$$

$$\text{So that, after third revision, the holding period} = 360 - 270 = 90 \text{ days}$$

So, option 'B' is correct.



Scan for Video solution



### 219. (b)

$$\text{Cost of route } 1 - a - c - 2 = 700$$

$$\text{Cost of route } 1 - f - b - 2 = 500$$

$$\text{Cost of route } 1 - b - 2 = 600$$

$$\text{Cost of route } 1 - f - e - 2 = 700$$

So, the cheapest route is 1-f-b-2



Scan for Video solution



### 220. (b)

$$\text{Angular speed of hour hand} = 0.5^\circ/\text{minute}$$

$$\text{Angular speed of minute hand} = 6^\circ/\text{minute}$$

$$\begin{aligned} \text{Relative velocity of minute hand w.r.t. hour hand} \\ = 6 - 0.5 = 5.5^\circ/\text{minute} \end{aligned}$$

At quarter part three implies minute hand has move 15 min.

So, the angle between the minute hand and hour hand.

= Angle between hands at 3'O clock – Relative angle turned by minute hand in 15 min.

$$= 90^\circ - 5.5 \times 15 = 7.5^\circ$$



Scan for Video solution



### 221. (d)

Let's assume those two positive numbers are A and B

$$A + B = 100$$

$$(A - 5) \times (B - 5) = 0$$

$$(A - 5) = 0$$

$$A = 5, B = 95$$

Or

$$(B - 5) = 0 \Rightarrow B = 5$$

$$A = 95$$



Scan for Video solution



### 222. (c)

As per given information

Sum of odd positive between 1,100

$$= 1 + 3 + 5 + \dots + 99$$

$$= \frac{n}{2} [2a + (n - 1)d] \quad \because (n = 50)$$

$$= \frac{50}{2} [2 \times 1 + (49) \times 2]$$

$$= 25[2 \times 1 \times 88]$$

$$= 25 \times 100$$

Sum of even positive number between

150 to 200 is

$$= 150 + 152 + \dots + 200$$

$$\frac{n}{2} [2 \times a + (n - 1)d] \quad \because (n = 26)$$

$$\frac{26}{2} [2 \times 150 + (26 - 1) \times 2]$$

$$26[150 + 25]$$

Ratio =

$$\frac{25 \times 100}{26 \times 175} = \frac{100}{26 \times 7} = \frac{50}{91}$$



Scan for Video solution

**223. (c)**

A per given information

In order to use same calender as for the year 2019

First check, is it leap year or not?

$$\frac{2019}{400} \neq 0 \text{ therefore it is not leap year}$$

For normal year number of odd days = 1

$$\frac{365}{7} \text{ days} = 1 \text{ (Remainder)}$$

To find the same calender as for the year 2019, the number of odd days from that year to 2019 must be zero

Total number of odd days

2011 to 2018 =

Odd days (2011) + odd days (2012) + ..... odd days (2018)

$$= 1 + 2 + 1 + 1 + 1 + 2 + 1 + 1 = \frac{10}{7} = 3 \text{ days}$$

Hence 2011 not possible

Similary

$$\text{Odd dyas (2012 - 2018)} = \frac{10-1}{7} = \frac{9}{7} = 2 \text{ days}$$

Similarly

$$\text{Odd days (2013-2018)} = \frac{10-1-2}{7} = \frac{7}{7} = 0 \text{ days}$$

Hence the 2013 year calendar is the same as that of the 2019.



Scan for Video solution

**224. (a)**

Concept of cyclicity

Number =  $(26591749)^{110016}$ 

The cycle of 9<sup>th</sup> is (9, 18, 729) Hence cycle of 2 term It means 2 term 9 again appear and

$$\text{Remainder} = \frac{110016}{2} = 0$$

Therefore, the last term of the cycle is consider as unit digit

Unitl place of  $9^{\text{even}} = 1$ 

Scan for Video solution

**225. (c)**

As per given informatin

1<sup>st</sup> term → 2, +7 more term+34 → 9<sup>th</sup> term

$$T_n = (a + (n - 1)d)$$

$$34 = 2 + (9 - 1) \times d$$

$$d = 4$$

therefore obtained sereis is:

$$\rightarrow 2, 6, 10, 14, 18, 22, \dots 34$$

$$S_n = \frac{n}{2} (2a + (n - 1)d)$$

$$= \frac{9}{2} (2 \times 2 + (9 - 1) \times 4) = 126$$



Scan for Video solution

**226. (a)**

From Man-Day-Hour-Work formula

$$\frac{m_1 d_1 h_1}{w_1} = \frac{m_2 d_2 h_2}{w_2} \Rightarrow \frac{52 \times 10 \times h}{w} = \frac{40 \times d_2 \times h}{w}$$

Where m = number of men

d = number of days

h = working hours in a day

w = part of work done


d<sub>2</sub> = 13 days

So, 3 more days than the estimated, will be required to complete the strip.



Scan for Video solution



**227. (d)**

Before entering into next level, the price is increased at that level by 25% hence the T received the product at Rs. 292.968 and after increasing 25% it arrives to customers at Rs. 366.21.



Scan for Video solution

**228. (b)**

Let, the number of boys =  $4x$

And the number of girls =  $3x$

$$\text{Then the total passed candidates} = 7x \times \frac{80}{100} = 5.6x$$

$$\text{The number of girls passed} = 3x \times \frac{90}{100} = 2.7x$$

$$\therefore \text{The number of boys passed} = 5.6x - 2.7x = 2.9x$$

$$\begin{aligned} \text{The percentage of boys passed} \\ = \frac{2.9x}{4x} \times 100 = 72.5\% \end{aligned}$$

So, option 'B' is correct.



Scan for Video solution

**229. (c)**

The even digits are = 0, 2, 4, 6, 8

The numbers between 100 to 1000 can be a 3-digit number, then we need to fill 3 places (i.e. \_\_\_) with 5 digits and repetition of the digits are allowed, but a 3-digit number cannot start with 0, so that at hundredth place we have only 4 entries,

$$\therefore \text{Then the total numbers are} = 4 \times 5 \times 5 = 100$$

So, option 'C' is correct.



Scan for Video solution

**230. (b)**

$$X \text{ can mow in 1 hour} = \frac{1}{2} \text{ portion of lawn}$$

$$Y \text{ can mow in 1 hour} = \frac{1}{4} \text{ portion of lawn}$$

X and Y both can mow together in 1 hour

$$= \frac{1}{2} + \frac{1}{4} = \frac{3}{4} \text{ portion of lawn}$$

$\therefore$  Time required to X and Y both to mow the

$$\text{complete lawn} = \frac{4}{3} \text{ hour} = \frac{4}{3} \times 60 = 80 \text{ min}$$

So, option 'B' is correct.



Scan for Video solution

**231. (b)**

$$V_A = 50 \text{ km/h}$$

$$V_B = 60 \text{ km/h}$$

The relative velocity  $V = 10 \text{ km/h}$

For a gap of 20km, the time required is

$$T = \frac{20}{10} = 2 \text{ hours}$$



Scan for Video solution

**232. (c)**

Let the cost of gift is  $X$

For 10 friends, the share on each one will be  $\frac{X}{10}$

Now after 2 friends are not contributing then

$$\frac{X}{8} - \frac{X}{10} = 150 \Rightarrow X = 6000$$



Scan for Video solution



**233. (b)**

Angular velocity of hour hand =  $0.5^\circ/\text{minute}$

So, time taken by hour hand to move  $210^\circ$

$$= \frac{210}{0.5} = 420 \text{ minutes}$$

Similarly, time taken by hour hand to move  $240^\circ$

$$= \frac{240}{0.5} = 480 \text{ minutes}$$

So, working time for P = 420 minutes

Working time for Q = 480 minutes

Effective working time for P =  $420 - 30 = 390$  minutes

Effective working time for Q =  $480 - 20 = 460$  minutes

Total effective working time for P and Q

$$= \frac{390 + 460}{60} = \frac{85}{6} \text{ hours}$$

$$\begin{aligned} \text{USD paid to consultants} &= 200 \times \frac{85}{6} \\ &= 2833.33 \text{ USD} \end{aligned}$$

So, the remaining USD with the client

$$= 3000 - 2833.33 = 166.67$$



Scan for Video solution

**234. (c)**

Machine A's 1 hour work =  $1/4$  part

Another machine B's 1 hour work =  $1/2$  part

1 hour's work of both the machines working together

$$= \frac{1}{4} + \frac{1}{2} = \frac{1}{4} + \frac{2}{4} = \frac{3}{4} \text{ part}$$

So, the time taken to complete the task by both machines is  $4/3$  hours.



Scan for Video solution

**235. (c)**

Let 'V' is the volume of tank.

$$\text{Capacity of pipe X} = +\frac{V}{5} \text{ litre / hour}$$

$$\text{Capacity of pipe Y} = +\frac{V}{4} \text{ litre / hour}$$



$$\text{Capacity of drainage pipe (Z)} = -\frac{V}{20} \text{ litre / hour}$$

If both the pumps are started with the drainage hole open then let after T hours, the tank is full, then

$$\left( \frac{V}{5} + \frac{V}{4} - \frac{V}{20} \right) T = V$$

$$\left( \frac{4+5-1}{20} \right) T = 1$$

$$T = \frac{20}{8} = 2.5 \text{ hours}$$

**236 (c)**

$$s_1 = s_2 - 5 \text{ cm} \quad \text{_____ (1)}$$

$$A_2 = 4A_1$$

$$s_2^2 = 4 s_1^2$$

$$s_2 = 2 s_1 \quad \text{_____ (2)}$$

From equation (1) & (2)

$$s_1 = 5 \text{ cm}$$

$$s_2 = 10 \text{ cm}$$

**237. (a)**

$$\begin{aligned} \text{Total surface area of mansion} &= 70 \times 55 \\ &= 3850 \text{ m}^2 \end{aligned}$$

$$\text{Area for flower pots} = 550 \text{ m}^2$$

$$\text{Carpet area required} = 3850 - 550 = 3300 \text{ m}^2$$

Cost of carpet =  $50 \times 3300 = \text{Rs. } 1,65,000$



Scan for Video solution



238. (c)

Volume of wall =  $30 \times 12 \times 6 = 2160 \text{ m}^3$

$$\begin{aligned}\text{Volume of a brick} &= \frac{8}{100} \times \frac{6}{100} \times \frac{6}{100} \\ &= 288 \times 10^{-6} \text{ m}^3\end{aligned}$$

Since, 60% of wall consists of bricks, then number of bricks required

$$= \frac{0.6 \times 2160}{288 \times 10^{-6}} = 45 \times 10^5 \text{ or } 45 \text{ lakhs}$$



Scan for Video solution



239. (b)

John has already worked with Ram for 5 hours so, Ram can not work with John any more & Ram can not work with Krishna. So, now Ram can only work with Amir. Hence, Krishna should be allotted to work with John.



Scan for Video solution



240. (d)

$\Rightarrow 2001$

$$P_1 = \frac{\text{Population of state X in } 2001 (N_X)}{\text{Population of state Y in } 2001 (N_Y)} = \frac{N_X}{N_Y}$$

$\Rightarrow 2011$

$$P_2 = \frac{\left(1 + \frac{x}{100}\right) N_X}{\left(1 + \frac{y}{100}\right) N_Y}$$

Percentage increase in P

$$= \left( \frac{P_2}{P_1} - 1 \right) \times 100$$

$$= \left( \frac{1 + \frac{x}{100}}{\left(1 + \frac{y}{100}\right)} - 1 \right) \times 100 \Rightarrow \frac{100(x-y)}{100+y}$$



Scan for Video solution



241. (b)

Let the total volume of tank is V then capacities of P,

$$Q \text{ & } R \text{ are } \frac{V}{6}, \frac{V}{9} \text{ and } -\frac{V}{12} \text{ per hour.}$$

When P & R are opened for 4 hours, then the volume in tank after 4 hour is.

$$\left( \frac{V}{6} - \frac{V}{12} \right) \times 4 = \frac{V}{3} \text{ is filled.}$$

Then P is closed and Q is opened for 6 more hours.

Volume in tank after 6 more hours is.

$$\frac{V}{3} + \left( \frac{V}{9} - \frac{V}{12} \right) 6 = \frac{V}{2} \text{ is filled.}$$

Then R is closed and volume to be filled is  $\frac{V}{2}$

$$\frac{V}{9} \times x = \frac{V}{2}$$

$$x = \frac{9}{2} \text{ or } 4.5 \text{ hour took by Q to fill } \frac{V}{2} \text{ volume.}$$

Total time

$$= 4 + 6 + 4.5 = 14.5 \text{ hours}$$



Scan for Video solution



**242. (a)**

Let in 2015 then GDP was  $G_1$  & that in 2016 is  $G_2$ . Given  $G_2 = 1.1 G_1$ , then percentage increase in the actual fiscal deficit is.

$$\frac{5\% \text{ of } G_2 - 4\% G_1}{4\% G_1} = \frac{\frac{5}{100} \times 1.1 G_1 - \frac{4}{100} \times G_1}{\frac{4}{100} \times G_1}$$

= 0.375 or 37.5%



Scan for Video solution



**243. (b)**

$$X \cdot Y \cdot Z = 192$$

$$X \cdot Y = \frac{192}{4} = 48$$

$$AM > GM$$

$$\frac{x+y}{2} \geq \sqrt{xy}$$

$$P \geq \sqrt{48}$$

$$P \geq 6.928$$

$P = 7$  (Minimum possible value of  $P$  for integers  $X$ ,  $Y$ , &  $Z$ )



Scan for Video solution



**244. (d)**

Let the amounts of two parts are  $x$  and  $(100,000 - x)$

$$\{x \times 1.1 + (100,000 - x) \times 1.12\} - \{x \times 1.12 + (100,000 - x) \times 1.1\} = 120$$

$$x(1.1 - 1.12 - 1.12 + 1.1) + 100,000 (1.12 - 1.1) = 120$$

$$x = 47,000$$

So, the required ratio is  $\frac{47,000}{100,000 - 47,000} = \frac{47}{53}$



Scan for Video solution



**245. (c)**

$$A + B = 26$$

$$A \times B = 165$$

$$(A - B)^2 = (A + B)^2 - 4A \times B$$

$$(A - B)^2 = 26^2 - 4 \times 165$$

$$A - B = 4$$



Scan for Video solution



**246. (d)**

For 1 hour, the hour hand moves by  $\frac{360^\circ}{12} = 30^\circ$ .

So, for  $225^\circ \Rightarrow \frac{225}{30} = 7.5$  hour



Scan for Video solution



**247. (c)**

Given,  $\frac{(k+2)^2}{k-3}$  will be an integer, if denominator is

factor or the numerator,

By option checking, let option (c).

$$\text{At, } k = 4 \Rightarrow \frac{(4+2)^2}{4-3} = 36$$

$$\text{At, } k = 8 \Rightarrow \frac{(8+2)^2}{8-3} = \frac{100}{5} = 20$$

$$\text{At, } k = 28 \Rightarrow \frac{(28+2)^2}{28-3} = \frac{900}{25} = 36$$

So, option 'C' is correct.



Scan for Video solution

**248. (c)**

|   |            |
|---|------------|
| 2 | 20, 42, 76 |
| 2 | 10, 21, 38 |
|   | 5, 21, 19  |

L.C.M of 20, 42, 76

$$= 20 \times 21 \times 19 = 7980$$

So, the number which leaves a remainder of 7 when divided by 20, 42 and 76 is

$$= 7980 + 7 = 7987$$



Scan for Video solution

**249. (b)**Let, the number of invited guests =  $x$ 

Invited male (I.M) = 0.6x

Invited female (I.F.) = 0.4x

The sum of female attendees (F.A.) and male attendees (M.A.) is 80% of all invited.

$$\text{Attendees} = \text{F.A.} + \text{M.A.} = 0.8x$$

$$0.4x + \text{M.A.} = 0.8x$$

$$\Rightarrow \text{M.A.} = 0.4x$$

$$\frac{\text{M.A.}}{\text{F.A.}} = \frac{0.4x}{0.4x} = \frac{1}{1}$$



Scan for Video solution

**250. (c)**

$$\text{Probability of getting green P(G)} = \frac{4}{6} = \frac{2}{3}$$

$$\text{Probability of getting red P(R)} = \frac{2}{6} = \frac{1}{3}$$

Option (A)

$$P(G = 3) = {}^7C_3 \cdot \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{1}{3} \times \frac{1}{3} \times \frac{1}{3} \times \frac{1}{3}$$

$$= {}^7C_3 \cdot \frac{8}{3^7} = \boxed{\frac{280}{3^7}}$$

Option (B)

$$P(G = 4) = {}^7C_4 \cdot \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{1}{3} \times \frac{1}{3} \times \frac{1}{3}$$

$${}^7C_4 \cdot \frac{16}{3^7} = \boxed{\frac{560}{3^7}}$$

Option (C)


$$P(G = 5) = {}^7C_5 \cdot \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{1}{3} \times \frac{1}{3}$$

$${}^7C_5 \cdot \frac{32}{3^7} = \boxed{\frac{672}{3^7}} \text{ (Maximum probability)}$$

Option (D)

$$P(G = 6) = {}^7C_6 \cdot \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{2}{3} \times \frac{1}{3}$$

$${}^7C_6 \cdot \frac{64}{3^7} = \boxed{\frac{448}{3^7}}$$


**251. (c)**

The sample space consists of blue cab which he identified as blue and the green cabs which he misidentifies as blue.

$$P\left(\frac{\text{Blue}}{\text{Correct}}\right) = \frac{P\left(\frac{\text{Correct}}{\text{Blue}}\right)}{P\left(\frac{\text{Correct}}{\text{blue}}\right) + P\left(\frac{\text{incorrect}}{\text{green}}\right)}$$

$$= \frac{0.80 \times 0.15}{0.80 \times 0.15 + 0.20 \times 0.85}$$

$$= \frac{1200}{2900} \times 100 = 41.37\%$$



**252. (b)**

$$(\text{Gold})_A : (\text{Copper})_A = 2 : 3$$

$$(\text{Gold})_B : (\text{Copper})_B = 3 : 7$$

Let the alloy C is made by taking  $x$  kg of both A and B alloys.

Gold in alloy C,  $(\text{Gold})_C = \text{Gold in A} + \text{Gold in B}$

$$= \frac{2}{5}x + \frac{3}{10}x = \frac{7x}{10}$$

Copper in alloy C

$(\text{Copper})_C = \text{Copper in A} + \text{Copper in B}$

$$= \frac{3}{5}x + \frac{7}{10}x = \frac{13x}{10}$$

$$\therefore (\text{Gold})_C : (\text{Copper})_C = \frac{7x}{10} : \frac{13x}{10} = 7 : 13$$



Scan for Video solution

**253. (b)**

Amount after 5 years  $A = 10,00,000$

Principal amount  $P = ?$

$$\text{We know } A = P \left(1 + \frac{r}{100}\right)^n$$

$$\Rightarrow 10,00,000 = P \left(1 + \frac{10}{100}\right)^5$$

$$\Rightarrow P = 620921.3231 \approx 6,21,000$$



Scan for Video solution

**254. (d)**

**Given:** Series is  $1 + \frac{1}{4} + \frac{1}{16} + \frac{1}{64} + \frac{1}{256} + \dots$

First term  $a = 1$

$$\text{Common ratio} = \frac{1}{4} \div 1 = \frac{1}{16} \div \frac{1}{4} = \dots = \frac{1}{4}$$

Sum of infinite geometric progression is given by

$$S_{\infty} = \frac{a}{1 - r} = \frac{1}{1 - \frac{1}{4}} = \frac{4}{3}$$

Scan for Video solution

**255. (b)**

A whole number is said to be divisible by 3 if the sum of all its digits is exactly divisible by 3.

Let the number is  $x$ .

$$\therefore \frac{7+1+5+x+4+2+3}{3} = n \quad (n \rightarrow \text{whole number})$$

$$\Rightarrow x = 3n - 22$$

for  $x = 0, n = \frac{22}{3}$  which is not a whole number

$x = 2, n = 8$  which is a whole number

$x = 5, n = 9$  which is a whole number

$x = 6, n = \frac{28}{3}$  which is not whole number

So, the minimum whole number in place of ■ is 2.

Scan for Video solution

**256. (d)**

Let the ages of Hema, Hari & Suresh are represented by A, B & C respectively.

$$A = 2B + 5 \quad \text{_____ (1)}$$

$$C = 10B - 13 \quad \text{_____ (2)}$$

$$C = 3A \quad \text{_____ (3)}$$

From equation 2 & 3

$$3A = 10B - 13 \quad \text{_____ (4)}$$

Equation (4) – equation (1)  $\times$  (3)

$$0 = 4B - 13 - 15$$

$$B = 7$$

From equation (1)

$$A = 19$$



Scan for Video solution

**257. (c)**

$$\frac{\text{cost price} - (\text{Selling price})_1}{\text{cost price}} = 12.5\% = \frac{1}{8}$$

$$\frac{S.P_1}{C.P.} = \frac{7}{8} \quad \text{_____ (1)}$$

For  $S.P_2 = S.P_1 + 108$

$$\frac{S.P_2 - CP}{CP} = 10\% = \frac{1}{10}$$

$$\frac{S.P_2}{CP} = \frac{11}{10} \quad \text{_____ (2)}$$

Equation (2) – equation (1)

$$\frac{S.P_2}{CP} - \frac{S.P_1}{CP} = \frac{11}{10} - \frac{7}{8}$$

$$\frac{S.P_2 - S.P_1}{CP} = \frac{44 - 35}{40}$$

$$\frac{108}{CP} = \frac{9}{40}$$

$$CP = 480$$

From equation (1)

$$S.P_1 = 480 \times \frac{7}{8} = 420$$

So, the loss in rupees is  $480 - 420 = 60$



Scan for Video solution

**258. (b)**

Given

$$\text{Price. } P \propto (\text{length, } L)^2$$

$$P = KL^2 \text{ where } K \text{ is constant}$$

$$1600 = K \cdot 100 \Rightarrow K = 16$$

$$P = 16L^2$$

Price for two wires of lengths 4 m & 6 m

$$P = 16 \times 4^2 + 16 \times 6^2 = 16(4^2 + 6^2) = 832 \text{ Rs.}$$



Scan for Video solution

**259. (c)**

$$a_1 + a_2 + a_3 + \dots + a_{50}$$

$$= \left( \frac{1}{1} - \frac{1}{1+2} \right) + \left( \frac{1}{2} - \frac{1}{2+2} \right) + \left( \frac{1}{3} - \frac{1}{3+2} \right) \dots + \left( \frac{1}{50} - \frac{1}{50+2} \right)$$

$$= \left( \frac{1}{1} - \frac{1}{3} \right) + \left( \frac{1}{2} - \frac{1}{4} \right) + \left( \frac{1}{3} - \frac{1}{5} \right) \dots \dots \left( \frac{1}{50} - \frac{1}{52} \right)$$

$$= \frac{1}{1} + \frac{1}{2} - \frac{1}{51} - \frac{1}{52} = 1 + \frac{1}{2} - \left( \frac{1}{51} + \frac{1}{52} \right)$$



Scan for Video solution

**260. (d)**

The number of distinct committees of 3-members out of 9 people possible is –

$${}^9C_3 = \frac{9 \times 2 \times 7}{3 \times 2 \times 1} = 84$$



Scan for Video solution



**261. (c)**

Loss,  $L \propto (\text{Deviation}, D)^2$

$L = KD^2$  where K is a constant

For  $D = 7$  units,  $L = \text{Rs. } 4900$

$$4900 = K \times 7^2$$

$$K = 100$$

$$L = 100 D^2$$

for  $D = 4$  units

$$L = 100 \times 4^2 = \text{Rs. } 1600$$



Scan for Video solution




Probability of one H & one T

$$= P(\text{TH}) + P(\text{HT}) = \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ (highest probability)}$$

Probability of one H followed by one T

$$P(\text{HT}) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$

**262. (b)**

For every 24 hours, 15 minutes in faulty clock, there is 24 hours in accurate clock

$$24.25 \equiv 24$$

∴ From 9 AM on 11 July to 2 PM on 15 July, there are  $(4 \times 24 + 5)$  hours.

$$\therefore \text{for } (4 \times 24 + 5) \equiv \frac{24}{24.25} \times (4 \times 24 + 5)$$

101 hours in faulty clock  $\equiv 99.96$  hours in accurate clock

∴ The difference in hour in faulty clock & accurate clock is approximately 1.04 hours or 1 hour, 2 minutes.

∴ The correct time when faulty clock shows 2 PM, on 15 July is 12:58 PM in accurate clock on 15 July.



Scan for Video solution

**263. (b)**

Probability of two T

$$P(2\text{T}) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$

Similarly, the probability of two H =  $\frac{1}{2}$


**264. (Marks to All)**

From men-day-hour-work formula

$$\frac{m_1 d_1 h_1}{w_1} = \frac{m_2 d_2 h_2}{w_2}$$

$$\frac{125 \times 39 \times 7}{\left(\frac{5}{7}\right)} = \frac{(125+x) \times (52-39) \times 8}{\left(\frac{2}{7}\right)}$$

$x = 6.25$  or 7 more robots.

**265. (c)**

Area of square = 1936 m<sup>2</sup>

$$l^2 = 1936 \text{ m}^2$$

$$l = 44 \text{ m}$$

Total length of wire =  $4l = 176$  m.

Let the length of shorter piece after cutting the wire a is

$$a + 3a = 176 \Rightarrow a = 44 \text{ m}$$

So, the lengths of longer & shorter wires are 132m & 44m.

$$\text{Total area} = \left(\frac{3a}{4}\right)^2 + \pi \left(\frac{a}{2\pi}\right)^2 = 1243.06 \text{ m}^2$$



Scan for Video solution

**266. (a)**

A circle encloses maximum area for the given parameter among all other shapes.



Scan for Video solution

**267. (d)**

Let length of train & platform are a and b in meters respectively.

$\Rightarrow$  When train passes platform

$$\frac{a+b}{25} = 54 \times \frac{5}{18} = 15 \text{ m/s} \quad \dots(1)$$

$\Rightarrow$  When train passes the man

$$\frac{a}{14} = (54 - 9) \times \frac{5}{18} = 12.5 \text{ m/s} \quad \dots(2)$$

$$a = 175 \text{ m}$$

From equation – 1

$$a + b = 15 \times 25$$

$$b = 200 \text{ m}$$



Scan for Video solution

**268. (d)**


$a + a^2 b^3$  is odd

$a(1 + ab^3)$  is odd only if a &  $(a + ab^3)$  both are odd.

$1 + ab^3$  is odd

or  $ab^3$  is even

a is odd hence  $b^3$  should be even and  $b^3$  is even if b is even. So, a is odd & b is even.



Scan for Video solution

**269. (b)**

Let in the two-digit number, y is the units digit and x is the tens digit, then the digit can be written as

$$10x + y$$

$$\text{Given, } x + y = 9$$

$$10x + y - 45 = 10y + x$$

$$9x = 9y + 45$$


$$9x = 9(9 - x) + 45$$

$$18x = 81 + 45$$

$$x = 7$$

$$\text{then } y = 2$$

$\therefore$  The number is 72



Scan for Video solution

**270. (b)**

Let the length & breadth are l & b.

$$l - 10 = b - 5$$

$$b = l - 5$$

$$lb - (l - 10)(b - 5) = 650$$

$$l(l - 5) - (l - 10)(l - 10) = 650$$


$$l^2 - 5l - (l^2 - 20l + 100) = 650$$

$$15l = 750$$

$$l = 50$$

$$b = 45$$

$$\text{Area of rectangle} = 50 \times 45 = 2250 \text{ m}^2$$



Scan for Video solution



**271. (d)**

Given,  $y \times 162 = y \times 6 \times 27$

$$\Rightarrow y \times 6 \times (3)^3$$

Above number will be a perfect cube if  $y$  will be  $6^2$ .

$$\therefore y = 6^2 = 36$$

So, option 'D' is correct.



Scan for Video solution

**272. (c)**

If we consider the one-digit number, then only seven numbers are possible, i.e. 1, 2, 3, 4, 6, 7, 8. Then probability of one-digit number is

$$\left(\frac{7}{10}\right) = 0.7 = (0.7)^1 = (0.7)^k$$

Similarly, if we consider the two-digit number, then only 49 numbers are possible which excludes the digit 0, 5, 9. Then probability of two-digit number is

$$\left(\frac{49}{90}\right) = 0.54 \approx (0.7)^2 = (0.7)^k$$

Similarly, if we consider the three-digit number, then only 343 numbers are possible which excludes the digit 0, 5, 9. Then probability of three-digit number is

$$\left(\frac{343}{900}\right) = 0.38 \approx (0.7)^3 = (0.7)^k$$

If we see this cycle, no option is correct in this question, but if we approximate our answer then only valid option is (c). It may be happened because examiner had taken total two-digit number as 100 but in actual, these are 90. Similarly, total three-digit number as 1000 but in actual, these are 900.

So, option 'C' is correct.



Scan for Video solution

**273. (d)**

There are four different people and four different colours, if all four people select any colour out of four, then total 16 cases are possible, but Arun does not like the colour red and Shweta does not like the colour white, so that two cases will reduce. Hence the number of ways they can choose the shirt =  $16 - 2 = 14$  ways.

So, option 'D' is correct.



Scan for Video solution

**274. (b)**

Let,

The number of MCQ questions =  $p$

The number of essay questions =  $q$

$$\therefore p + q = 20 \quad \dots \text{(i)}$$

$$\text{And, } 3p + 11q = 100 \quad \dots \text{(ii)}$$

On solving (i) and (ii), we get

$$p = 15, q = 5$$

$\therefore$  The number of MCQ questions =  $p = 15$

So, option 'b' is correct.



Scan for Video solution



Scan for Video solution

**275. (d)**

Number of ways, if both socks are Red =  ${}^3C_2$

Number of ways, if both socks are Green =  ${}^4C_2$

Number of ways, if both socks are Blue =  ${}^3C_2$

Number of ways, if both are of any colour =  ${}^{10}C_2$

Probability of both socks are of the same colour,

$$= \frac{{}^3C_2 + {}^4C_2 + {}^3C_2}{{}^{10}C_2} = \frac{3 + 6 + 3}{45} = \frac{12}{45} = \frac{4}{15}$$

So, option 'D' is correct.



Scan for Video solution



Scan for Video solution



**276. (a)**

Smallest 30-digit number, starts with  $47 = 47 \times 10^{28}$   
 Greatest 30-digit number, starts with  $47 = 48 \times 10^{28} - 1$

$$\text{Now, } (47 \times 10^{28})^3 = 103823 \times 10^{84}$$

And,

$$(48 \times 10^{28} - 1)^3 < (110592 \times 10^{84} = (48 \times 10^{28})^3)$$

Hence this value will lie in the interval of  $103823 \times 10^{84}$  to  $110592 \times 10^{84}$ . Which clearly indicate that it will be a 90-digit number.

So, option 'A' is correct.



Scan for Video solution

**277. (c)**

1200 men and 500 women can build in 2 weeks.

$$(1200 \text{ Men} + 500 \text{ Women}) \text{ one-week work} = \frac{1}{2}$$

part... (i)

Similarly,

$$(900 \text{ men} + 250 \text{ women}) \text{ one-week work} = \frac{1}{3}$$

part... (ii)

Equation (ii)  $\times 2$  – equation (i)

$$(1800 \text{ men} - 1200 \text{ men}) \text{ one-week work} = \frac{2}{3} - \frac{1}{2} = \frac{1}{6}$$

600 men will complete the work in 6 weeks


Men required to complete the work in one week

$$\Rightarrow 600 \times 6 = x \times 1 \Rightarrow x = 3600$$

So, 3600 men will complete the work in 1 week.



Scan for Video solution

**278. (a)**

So, distance to be covered by 1 batch of vehicles  
 $= 20 + 10 + 15 + 5 = 50 \text{ m}$

$\therefore$  Speed is 36km/1hr


In 1 hr distance travelled is 36 km = 36000 m

$\therefore$  50 m distance is covered by 1 batch

$$36000 \text{ m distance will be covered by} = \frac{1}{50} \times 36000 \\ = 720 \text{ batches}$$

So, the maximum number of vehicles

$$= 720 \times 2 = 1440$$

**279. (d)**

water consumption (W) = household consumption (H) + other consumption (O)


In summer,  $0.75 W = 0.80 H + 1.70 O$

$$\Rightarrow 0.75 (H + O) = 0.80 H + 1.70 O$$

$$\Rightarrow (0.75 - 0.80) H = (1.70 - 0.75) O$$

$$\Rightarrow -0.05 H = 0.95 O \Rightarrow \frac{H}{O} = -\frac{0.95}{0.05} = -19$$

As the ratio of household to other consumption is negative there must be some errors in the official's statement.




280. (c)

$$V = \frac{1}{3}\pi r^2 h$$

$$= r^2 = 1.5 \times 1.5 = 2.25$$

more than 1 by 1.25 thus 125%



Volume of right circular cone

$$= \frac{1}{3}\pi r^2 \times h$$

$$V \propto r^2$$

Percentage increase in volume

$$\begin{aligned} &= \frac{r_2^2 - r_1^2}{r_1^2} = \frac{(1.5 r_1)^2 - r_1^2}{r_1^2} \\ &= \frac{2.25 r_1^2 - r_1^2}{r_1^2} \\ &= 1.25 \text{ or } 125\% \end{aligned}$$

Scan for Video solution



281. (d)

The sequence of numbers in increasing order

1, x, x, x, y, y, 9, 16, 18

There is total nine numbers so,

Median  $\Rightarrow$  5<sup>th</sup> term

Median = y

Mode  $\Rightarrow$  highest repeating number = x

Given, median = 2 mode

$$y = 2x$$

$$\text{Mean} = \frac{1+x+x+x+y+y+9+16+18}{9} = y$$

$$\frac{3x + 2y + 44}{9} = y$$

$$3x + 2y + 44 = 9y$$

$$3x + 44 = 7y$$

$$3x + 44 = 7(2x) \quad \{ \because y = 2x \}$$

$$11x = 44$$

$$x = 4, y = 8$$

Scan for Video solution



282. (d)

Let the Budhan speed by cycle is 'x' and that by walking is 'y' in km/h

$$\Rightarrow 19 = x \times \frac{t}{4} + y \times \frac{3t}{4}$$

$$19 = x \frac{2}{4} + y \frac{3 \times 2}{4}$$

$$38 = x + 3y \quad \text{_____ (1)}$$

$$\Rightarrow 26 = x \times \frac{t}{2} + y \times \frac{t}{2}$$

$$26 = x \times \frac{2}{2} + y \times \frac{2}{2}$$

$$26 = x + y \quad \text{_____ (2)}$$

From equations 1 & 2

$$2y = 12$$

$$y = 6 \text{ km/h}$$

Scan for Video solution



283. (b)

$$1 + 2^1 + 3^3 + 4^1$$

$$= 1 + 2 + 7 + 4 = 14$$

unit digit is 4

Scan for Video solution



284. (a)

|            |         |                        |
|------------|---------|------------------------|
| P → 30 min | overall | $30 \times 2 = 60$ min |
| Q → 20 min | overall | $20 \times 3 = 60$ min |
| R → 60 min | overall | $60 \times 1 = 60$ min |
| S → 15 min | overall | $15 \times 4 = 60$ min |

| P                            | Q | R | S |
|------------------------------|---|---|---|
| 2                            | 3 | 1 | 4 |
| M <sub>1</sub> → P P R       |   |   |   |
| M <sub>2</sub> → Q Q S S S S |   |   |   |



Scan for Video solution



285. (b)

$$9^2 \times \frac{\left(\frac{4}{5}\right)^{2x+4}}{\left(\frac{3}{5}\right)^{2x+4}} = 3^2 \times 4^2$$

$$\Rightarrow \left(\frac{4}{3}\right)^{2x+4} = \left(\frac{4}{3}\right)^2 \Rightarrow 2x + 4 = 2 \Rightarrow x = -1$$



Scan for Video solution



286. (c)

Total number of ways =  $5 \times 4 \times 3 \times 2 = 120$   
Number of ways, in which (P is paired with Z) =  $4 \times 3 \times 2 = 24$   
Number of ways in which (Y is not paired) =  $4 \times 3 \times 2 \times 1 = 24$   
Number of ways in which (Y is not paired and also P and Z are paired together) =  $3 \times 2 \times 1 = 6$   
So, the number of ways in which the required pairs can be formed  
=  $120 - 24 - 24 + 6 = 78$



Scan for Video solution



287. (a)

Let Y tractor is equivalent to AX bullocks. The work done by X bullocks and Y tractors in 8 days will be same as work done by X/2 bullocks and 2Y tractors in 5 days. So,

$$(X + Y)8 = \left(\frac{X}{2} + 2Y\right) \times 5$$

$$(X + AX)8 = \left(\frac{X}{2} + 2AX\right) \times 5$$

$$(1 + A) \times 8 = \left(\frac{1}{2} + 2A\right)5$$

$$8 + 8A = 2.5 + 10A$$

$$A = 2.75$$

Let 'n' be the number of days taken to plough the field when only X bullocks plough the field. Then similarly,

$$(X + Y).8 = X. n$$

$$X(1 + A) . 8 = X. n$$

$$(1 + 2.75) \times 8 = n$$

$$n = 30 \text{ days}$$



Scan for Video solution



288. (b)

Given P looks at Q while Q looks at R. P is married, R is unmarried.

Assuming Q is married.

The pair we get is Q – R, so one pair.

Or Assuming Q is unmarried

Then the pair will be P – Q, here also one pair so in either of the cases, the pair is only 1.



Scan for Video solution



**289. (c)**

Let,  $P(A)$  : probability of 1<sup>st</sup> child being a boy =  $\frac{1}{2}$

$P(B)$  : probability of 2<sup>nd</sup> child being boy =  $\frac{1}{2}$

∴ Both the events are independent.

$$\therefore P(A \cap B) = P(A) \times P(B)$$

$$P\left(\frac{B}{A}\right) = \frac{P(A \cap B)}{P(A)} = \frac{P(B) \cdot P(A)}{P(A)} = P(B) = \frac{1}{2}$$



Scan for Video solution

**290. (d)**

If  $a$  &  $b$  are integers and  $a - b$  is even then  $a$  &  $b$  both can either be even or both can be odd integers.

$$\Rightarrow ab - b$$

=  $b(a - 1)$  will also be even because if  $a$  &  $b$  are even numbers then  $a - 1$  will be odd & multiplication of odd & even numbers is even.

Similarly if  $a$  &  $b$  are odd numbers then  $a - 1$  will be even and  $b(a - 1)$  will be even.



Scan for Video solution

**291. (d)**

$$\begin{array}{r} 5 \underline{0} 1 \underline{0} 0 \\ - 4 8 \underline{9} 8 9 \\ \hline 1 1 1 1 \end{array} \quad \& \quad \begin{array}{r} 5 \underline{0} \underline{0} 0 0 \\ - 4 8 \underline{8} 8 9 \\ \hline 1 1 1 1 \end{array}$$

There are both possibilities so the sum of missing digits cannot be determined.



Scan for Video solution

**292. (b)**

To find median, the 100 employees can be numbered from 1 to 100 in increasing order of their salaries.

1 to 45 earn 20,000, 46 to 70 earn 30,000, 71 to 90 earn 40,000, 91 to 98 earn 60,000 and 99 to 100 earn 150,000. Since 100 is an even number then median

$$\begin{aligned} &= \frac{\left(\frac{100}{2}\right)^{\text{th}} \text{ employee} + \left(\frac{100}{2} + 1\right)^{\text{th}} \text{ employee}}{2} \\ &= \frac{50^{\text{th}} \text{ employee salary} + 51^{\text{th}} \text{ employee salary}}{2} \\ &= \frac{30,000 + 30,000}{2} = 30,000 \end{aligned}$$



Scan for Video solution

**293. (a)**

$$\text{Time to pass a post} = \frac{\text{Length of train}}{\text{Speed of train}}$$

$$\Rightarrow 10 = \frac{120}{S_1} \quad \Rightarrow S_1 = 12 \text{ m/s}$$

$$\text{Also, } 15 = \frac{150}{S_2} \quad \Rightarrow S_2 = 10 \text{ m/s}$$

So, difference in the speed

$$\Delta S = S_1 - S_2 = 12 - 10 = 2 \text{ m/s}$$



Scan for Video solution

**294. (a)**

$$\Rightarrow x\% \text{ of } y + y\% \text{ of } x$$

$$= \frac{x}{100}y + \frac{y}{100}x$$

$$= \frac{2}{100}xy \text{ or } 2\% \text{ of } xy$$



Scan for Video solution



**295. (a)**

The two-digit number can be represented by “ $x \times 10 + y$ ”, where  $x$  &  $y$  are digits at tens place & units place respectively.

Given,  $x + y = 12$  \_\_\_\_\_ (1)

After reversing, the digits new number is “ $y \times 10 + x$ ”.

$$(y \times 10 + x) - (x \times 10 + y) = 54$$

$$9y - 9x = 54$$

$$y - x = 6 \quad \text{_____ (2)}$$

From equation (1) & (2)

$$x = 3, y = 9$$

then the original number is

$$= x \times 10 + y$$

$$= 30 + 9 = 39$$



Scan for Video solution

**296. (c)**

The reading capacity of Ananth =  $\frac{1}{6}$  books/hour

The reading capacity of Bharath =  $\frac{1}{4}$  books/hour

Number of pages to be read means number of pages left.

Let after  $x$  hours Ananth is left with the number of pages twice that by Bharath

$$\left(1 - \frac{x}{6}\right) = 2\left(1 - \frac{x}{4}\right)$$

$$\frac{6-x}{6} = \frac{4-x}{2}$$

$$6-x = 12-3x$$

$$2x = 6$$

$$x = 3$$



Scan for Video solution

**297. (b)**

Square  $\frac{340-x}{4}$

$$\frac{340-x}{4}$$

Rectangle  $b = \frac{x}{6}$

$$a = \frac{x}{3} \quad a = 2b$$

Let the length of two parts are  $(340 - x)$  and  $x$ .

For rectangle,

$$x = 2a + 2b$$

Since, b: a = 1:2, hence

$$x = 6b$$

$$b = \frac{x}{6} \quad \text{and} \quad a = \frac{x}{3}$$

$$\text{Combined Area} = \left(\frac{340-x}{4}\right)^2 + \frac{x}{6} \times \frac{x}{3}$$

$$A = \frac{(340-x)^2}{16} + \frac{x^2}{18}$$

For minimum area, differentiating A with respect to  $x$  and equating it to 0, we get the value of  $x$  for which area will be minimum.

$$\frac{dA}{dx} = -2 \frac{(340-x)}{16} + \frac{2x}{18} = 0$$

$$\frac{340-x}{8} = \frac{x}{9} \Rightarrow x = 180$$

$$\frac{d^2A}{dx^2} \Big|_{x=180} = +ve, \text{ hence } A \text{ is minimum at } x = 180.$$

So, the length of the square corresponding to, minimum combined area is

$$\frac{340-180}{4} = 40 \text{ mm.}$$



Scan for Video solution




**298. (b)**

Let  $x$  be the number of days E worked. Since the efficiency of M is twice that of others, hence

$$\text{M's work} = 10 \sin 45^\circ$$

$$\text{E's work} = \left( 12 \frac{\text{Hours}}{\text{Day}} \times x \right) = 12x$$


The ratio is  $6 : 12 , 1 : 2$

**299. (c)**

$$(9 \text{ inches})^{\frac{1}{2}} = (0.25 \text{ yards})^{\frac{1}{2}}$$

Squaring both sides.

$$9 \text{ inches} = 0.25 \text{ yards}$$

**300. (a)**

$$a \square x = a$$

$$ax + (a + x) = a$$

$$\Rightarrow \text{For } x = 0$$

$$a \times 0 + a + 0 = a$$

$$a = a$$

$$\Rightarrow \text{For } x = 1$$


$$a + a + a \neq a$$

$$\Rightarrow \text{For } x = 2$$

$$2a + a + 2 \neq a$$

$$\Rightarrow \text{For } x = 10$$

$$10x + a + 10 \neq a$$

**301. (c)**

For normal distributions


Mean = Median = mode.

Median of P = 105

Median of Q = 85

This shows that most of the students in Q got less marks than P, but not necessarily all.

Standard deviation shows the variation or dispersion of a set of data values. Since standard deviation of P is more than that of Q so marks of students in Q is in a narrower range.

**302. (b)**

Perimeter of window = 6m


$$5a = 6$$

$$a = \frac{6}{5}$$

$$\text{Area of window} = a^2 + \frac{1}{2} a \times \left( \frac{\sqrt{3}}{2} a \right)$$

$$= a^2 \left( 1 + \frac{\sqrt{3}}{4} \right) = \left( \frac{6}{5} \right)^2 \left( 1 + \frac{\sqrt{3}}{4} \right)$$

$$= 2.06 \text{ m}^2$$



**303. (c)**

$$\text{Capacity of Q to do work per hour per day} = \frac{1}{12 \times 25}$$

$$\text{Capacity of R to do work per hour per day} = \frac{1}{12 \times 50}$$

Work done by Q in 5 days working for 12 hours a day

$$W_Q = \frac{1}{12 \times 25} \times 5 \times 12 = \frac{1}{5}$$


Work done by R in 7 days working for 18 hours a day

$$W_R = \frac{1}{12 \times 50} \times 7 \times 18 = \frac{21}{100}$$

So  $W_Q : W_R = 20 : 21$



Scan for Video solution

**304. (c)**

The percentage of people who does not show symptom of disease after being infected.

$$= \frac{50}{100} \times \frac{70}{100} = 0.35 \text{ or } 35\%$$



Scan for Video solution

**305. (a)**

Out of the total 5692000 fruits, 15% are unripe.

$$\text{The apples in unripe fruits} = 0.45 \times 0.15 \times 5692000$$

$$= 384210$$


Out of total fruits, 85% are ripe.

$$\text{The apple in ripe fruit} = 0.34 \times 0.85 \times 5692000$$

$$= 1644988$$

$$\text{Total apples} = 384210 + 1644988$$

$$= 2029198$$


**306. (d)**

Let;

$$\text{Probability of passes in Math } P(m) = M$$

$$\text{Probability of passes in Physics } P(p) = P$$

$$\text{Probability of passes in Chemistry } P(c) = C$$



Therefore;

The probability of passing in at least one subject,

$$P(m \cup p \cup c) = P(m) + P(p) + P(c) - P(m \cap p) - P(p \cap c) - P(c \cap m) + P(m \cap p \cap c)$$

$$0.75 = M + P + C - MP - PC - CM + MPC \quad \dots(i)$$

The probability of passing in at least two subjects,

$$0.5 = P(m \cap p) + P(p \cap c) + P(c \cap m) - 2P(m \cap p \cap c)$$

$$0.5 = MP + PC + CM - 2MPC \quad \dots(ii)$$

The probability of passing in exactly two subjects,

$$0.4 = P(m \cap p) + P(p \cap c) + P(c \cap m) - 3P(m \cap p \cap c)$$

$$0.4 = MP + PC + CM - 3MPC \quad \dots(\text{iii})$$

From the equation (ii) and (iii), we get

$$MPC = 0.1 = \frac{1}{10} \quad \dots(\text{iv})$$

From the equation (iii) and (iv), we get

$$MP + PC + CM = 0.7 = \frac{14}{20} \quad \dots(\text{v})$$

From the equation (i), (iv) and (v), we get

$$M + P + C = 1.3 = \frac{27}{20} \quad \dots(\text{vi})$$

So, option 'D' is correct.



Scan for Video solution



### 307. (c)

$$h(2, 5, 7, 3) = \frac{7 \times 3}{2 \times 5} \rightarrow \text{Remainder} = 1$$

$$fg(h(2, 5, 7, 3), 4, 6, 8) = fg(1, 4, 6, 8)$$

$$\Rightarrow f(1, 4, 6, 8) \times g(1, 4, 6, 8) = 8 \times 1 = 8$$



Scan for Video solution



### 308. (c)

$$f(x) = mx + c$$

$$f(-2) = -2m + c$$

$$29 = -2m + c \quad \dots(\text{i})$$

$$f(3) = 3m + c$$

$$39 = 3m + c \quad \dots(\text{ii})$$

$$(\text{ii}) - (\text{i})$$

$$5m = 10$$

$$m = \frac{10}{5} = 2$$

From equation (i)

$$29 = -4 + c$$

$$c = 33$$

$$f(5) = 2 \times 5 + 33$$

$$= 10 + 33 = 43$$


Scan for Video solution


### 309. (d)

Volume of A = 70% of volume of B

$$V_A = 0.7 V_B$$

Volume of A from statement II

$$V_A = 14,000 \text{ gallons}$$

$$\text{Then, } V_B = \frac{14000}{0.7} = 20,000 \text{ gallons}$$

Total gallons of solution in A & B

from statement I

$$= 80\% \text{ of } V_A + 40\% \text{ of } V_B$$

$$= 0.8 \times 14,000 + 0.4 \times 20,000$$


$$= 19,200 \text{ gallons}$$

So, to answer the question both the statements together are sufficient.


Scan for Video solution


### 310. (4536 to 4536)

Let the four boxes represent four digits. The number of possible digits at each place is shown below the box



So, total number that can be formed, is

$$9 \times 9 \times 8 \times 7 = 4536$$



Scan for Video solution



### 311. (b)

Event Z is that two tails occur at three tosses and event Y is that a tail occurs on the third toss. Event [TTH] is possible in Z but not possible in Y so, Z and Y are dependent.

Events Z and X are opposite to each other so these are dependent.

Events X and Y are not opposing each other hence these events are independent.



Scan for Video solution



### 312. (280 to 280)

$$\frac{PS}{QS} = \frac{3}{1}; \quad PS = 3x, QS = 1x$$

$$\frac{RT}{QT} = \frac{5}{2}; \quad RT = 5y, QT = 2y$$

Area of  $\Delta QTS$

$$\frac{1}{2} QS \times QT = 20 \text{ cm}^2$$

$$\frac{1}{2}(x \times 2y) = 20 \text{ cm}^2$$

$$xy = 20 \text{ cm}^2$$

From  $\Delta PQR$

Area =  $\frac{1}{2} PQ \times QR$

$$= \frac{1}{2} (3x + 1x) \times (5y + 2y) = \frac{1}{2} 28xy$$

$$= 14 \times 20 = 280 \text{ cm}^2$$



Scan for Video solution



### 313. (b)

Number of matches to complete the league round of matches is

$${}^5 C_2 = \frac{5 \times 4}{2} = 10$$





Scan for Video solution



### 314. (1.9 to 2.2)

Surface area of cone = 0.9 area of a circular sheet



$$\pi r R = 0.9 \times \pi R^2$$

$$r = 27 \text{ cm}$$

$$H = \sqrt{R^2 + r^2}$$

$$H = \sqrt{30^2 + 27^2} = 39.00 \text{ cm}$$

$$\frac{r}{H} = \frac{27}{39.00} = 0.69$$



Scan for Video solution



### 315. (800 to 800)

Tiger's speed = 5 leaps/min =  $5 \times 8$  meter/min = 40 meter/min

Dear's speed = 4 leaps/min =  $4 \times 5$  meter/min = 20 meter/min

Distance between tiger & dear =  $50 \times 8 = 400$  meters.

$$\text{Time to catch the dear} = \frac{400}{40 - 20} = 20 \text{ min}$$

In 20 minutes, tiger will cover  $40 \times 20 = 800$  meters




Scan for Video solution



### 316. (b)

$$\text{Given } P(A) = \frac{1}{6} \text{ and } P(B) = \frac{1}{8}$$

Probability of only one of Ram and Ramesh to be selected



$$= P(A) + P(B) - 2P(A \cap B)$$

$$= \frac{1}{6} + \frac{1}{8} - 2 \times \frac{1}{8} \times \frac{1}{6}$$

$$= \frac{14}{48} - \frac{2}{48} = \frac{12}{48} = \frac{1}{4}$$



Scan for Video solution



### 317. (96 to 96)


$$\text{Given } \left( z + \frac{1}{z} \right)^2 = 98$$

$$\text{Using } (a+b)^2 = a^2 + b^2 + 2ab$$

$$\left( z + \frac{1}{z} \right)^2 = z^2 + \frac{1}{z^2} + 2 \times z \times \frac{1}{z} = 98$$

$$z^2 + \frac{1}{z^2} + 2 = 98$$

$$z^2 + \frac{1}{z^2} = 98 - 2 = 96$$



### 318. (850 to 850)

Charge for single person of two-way fare =  $2 \times 100$


There are total 5 person in the group, then,

Total fare charge without discount =  $2 \times 100 \times 5$

Total discount =  $10\% + 5\% = 15\%$

$\therefore$  Total charge to a group of 5 people

$$= 1000 \times \frac{100 - 15}{100} = 850 \text{ Rs.}$$



### 319. (b)

From 2 to 198, first multiple of 10 will be 10 and last will be 190.


Let total multiple of 10 in between 2 and 198 is n.

$$\text{Then, } 190 = 10 + (n - 1)10 \Rightarrow n = 19$$

Sum of these 19 numbers =  $S_n$

$$S_n = \frac{19}{2} [2 \times 10 + (19 - 1) \times 10] = 1900$$

$$\text{Now, the avg of these numbers} = \frac{1900}{19} = 100$$



### 320. (c)


$$\text{Let, } p = \sqrt{12 + \sqrt{12 + \sqrt{12 + \dots}}}$$

$$\text{Then, } p = \sqrt{12 + p}$$

Squaring both sides, we get


$$p^2 = 12 + p$$

$$\begin{aligned} p^2 - p - 12 &= 0 \\ \Rightarrow (p - 4)(p + 3) &= 0 \\ p &= 4, -3 \\ \text{Valid value of } p \text{ will be positive, then } p &= 4 \\ \text{So, option 'C' is correct.} \end{aligned}$$

**321. (a)**

The required time =  $x : (30x \pm \theta) \times \frac{2}{11}$

Where,  $x$  = starting time of span  
 $\theta$  = angle between minute and hour hand  
 Take '+' for 0 to 6 span and '-' for 6 to 12 span  
 $\therefore$  Time =  $6 : (30 \times 6 - 60) \times \frac{2}{11} = 6 : 21 \frac{9}{11} \approx 6 : 22$   
 So, option 'A' is correct.

**322. (d)**

Consider the GDP before year 2012-2013 =  $x$   
 GDP increased during 2012-2013 by 7%.  
 $\therefore$  GDP during 2012-2013 =  $1.07x$   
 Similarly, USD price increased from 50 to 60 during this period, which reduced the value of rupee by a ratio  $5/6$   
 $\therefore$  New GDP =  $1.07x \times \frac{5}{6} = 0.8916x$


The percentage change is GDP

$$= \frac{0.8916x - x}{x} \times 100$$

$$= -10.84\% \approx -11\%$$


Negative sign indicates that GDP will decrease by 11% approximately.

So, option 'D' is correct.

**323. (1300 to 1300)**

$$\begin{aligned} \text{Total cost} &= \text{Fixed cost} + \text{Variable cost} \\ &= 50000 + 800 Q \\ &= 50000 + 800 \times 100 \end{aligned}$$

$$\begin{aligned} \text{Total cost for } 100 \text{ tonnes} &= 1,30,000 \\ \text{For } 1 \text{ tonne} &= 1,30,000/100 = 1300 \end{aligned}$$

**324. (d)**


Let the populations in country X and Y are  $x$  and  $y$  respectively.

$$x = 3y$$


$$\begin{array}{ccc} x & & y \\ \downarrow & & \downarrow \\ 3y & & y \end{array}$$

The percentage of people taller than 6 ft

$$\frac{0.01 \times 3y + 0.02y}{4y} \times 100 = 1.25$$

**325. (a)**

$$\begin{aligned} 5 &\rightarrow 6 \rightarrow 7.2 \rightarrow 8.64 \rightarrow 10 \Rightarrow 3 \text{ to 4 years} \\ 5 \times (1.2)^n &= 10 \\ (1.2)^n &= 2 \\ n &= \frac{\log 2}{\log 1.2} \Rightarrow n = 3.8 \end{aligned}$$



**326. (b)**

$$\begin{array}{cccccc}
 \text{T.Th} & \text{Th} & \text{H} & \text{T} & \text{U} \\
 \boxed{\phantom{0}} & \boxed{\phantom{0}} & \boxed{\phantom{0}} & \boxed{\phantom{0}} & \boxed{\phantom{0}} \\
 5 & \times & 4 & \times & 3 & \times & 2 & \times & 1 \\
 = & 120
 \end{array}$$

Out of these 120, our 5 possible values will each appear equal number of times in the unit place, the tens place, the hundreds place etc. So,  $120/5 = 24$  or there will be 24 numbers starting with 1, 24 starting with 3 and 24 starting with 5 etc.

Sum of the numbers =

$$\begin{aligned}
 & 24 \times [10000(1+3+5+7+9) + 1000] \\
 & (1+3+5+7+9) + 100(1+3+5+7+9) + 10 \\
 & (1+3+5+7+9) + (1+3+5+7+9)] = 6666600
 \end{aligned}$$



Scan for Video solution



**327. (b)**

$$A = P \left(1 + \frac{r}{100}\right)^n$$

$$2P = P \left(1 + \frac{r}{100}\right)^{10}$$

$$r = 7.2\%$$



Scan for Video solution



**328. (495 to 495)**

1<sup>st</sup> five numbers =  $x, x+2, x+4, x+6, x+8$

$$\text{Sum} \Rightarrow 5x + 20 = 425$$

$$x = \frac{405}{5} = 81$$

Last five =  $x+14, x+16, x+18, x+20, x+22$

$$\text{Sum} \Rightarrow 81 \times 5 + 90 = \boxed{495}$$



Scan for Video solution



**329. (0.80 to 0.82)**

$$\text{Probability to be defective } \frac{5}{100}$$

$$\text{Probability to be non-defective} = \frac{95}{100} = 0.95$$

$$\text{Probability of batch is accepted} = (0.95)^4 = 0.8145$$



Scan for Video solution



**330. (4 to 4)**

$$T = \frac{\text{Distance}}{\text{Speed}}$$

Speed of Man = 8 km/h

Speed of Stream = 'x' km/h

Since, the time to row upstream is thrice the time to row downstream. Hence,

$$3 \left( \frac{D}{8+x} \right) = \frac{D}{8-x}$$

$$\Rightarrow 24 - 3x = 8 + x$$

$$4x = 16$$

$$'x' = 4 \text{ km/h}$$



Scan for Video solution



**331. (d)**

In options A and B, the direct relation between P and M is not clear. In option C,  $P > M$ . In option D,  $P < M$ . So option D is correct.



Scan for Video solution

**332. (a)**

Let the volume of half tank is 1 litre, then drainage (E) and pumping (F) rates are respectively

$$E = \frac{-1}{30}, F = \frac{1}{x}$$

Given that the tank should be filled in 10 min, so

$$\frac{1}{x} - \frac{1}{30} = \frac{1}{10}$$

$$\frac{1}{x} = \frac{1}{10} + \frac{1}{30} = \frac{4}{30}$$

$$\Rightarrow x = \frac{30}{4} = 7.5 \text{ min}$$

$$7.5 \times 4 = \boxed{30}$$

Hence water has to be pumped with a rate 4 times that of drainage.



Scan for Video solution

**333. (163 to 163)**

ODD No.

$$x, x+2, x+4, x+6, x+8, x+10, x+12, x+14$$

$$\text{sum} = 8x + 56 = 656$$

$$8x = 600$$

$$x = \boxed{75}$$

EVEN No.

$$y, y+2, \underline{y+4}, y+6$$

$$\text{sum} = 4y + 12 = 87 \times 4$$

$$y = 84$$

sum of smallest odd number and second largest even number  $= x + (y+4) = 75 + (84+4) = 163$



Scan for Video solution

**334. (b)**

$$0.167 + 0.167 + 0.152 + 0.166 + 0.168 + 0.180 = 1$$


Since the probabilities of all the numbers are not same hence the dice is biased.



Scan for Video solution

**335. (b)**

Given that the tossed side is heads



then the probability of other side being tails is  $1/3$



Scan for Video solution



**336. (560 to 560)**

Train crossing Man

$$D = 280 \text{ m} \text{ (length of train)}$$

$$T = 20 \text{ sec}$$

$$\text{Speed (S)} = \frac{280}{20} = 14 \text{ m/sec}$$

Train crossing platform of length P

$$D = 280 + P$$

$$T = 60 \text{ sec}$$

$$S = 14 \text{ m/sec}$$

$$D = S \times T$$

$$280 + P = 14 \times 60$$

$$\Rightarrow P = 840 - 280 = 560 \text{ meter}$$

Scan for Video solution

**337. (c)**

$$\text{Average speed} = \frac{\text{Total distance (km)}}{\text{Total time (hr)}}$$

$$\text{Average speed} = \frac{8 + 6 + 16}{(15 + 15 + 15) \times \frac{1}{60}}$$

$$\text{Average speed} = \frac{30 \times 60}{45} = 40 \text{ km/hr}$$

So, option 'C' is correct.

Scan for Video solution

**338. (c)**

Let the temperature of Monday, Tuesday, Wednesday and Thursday are w, x, y, z respectively;

$$\text{Then, } \frac{w + x + y}{3} = 41$$

$$\therefore w + x + y = 123 \quad \dots \text{(i)}$$

$$\text{And, } \frac{x + y + z}{3} = 43$$

$$\therefore x + y + z = 129 \quad \dots \text{(ii)}$$

From equation (i) and (ii),

$$z - w = 6$$

Also, given that,  $z = w + 0.15w = 1.15w$ 

$$\therefore 1.15w - w = 6$$

$$0.15w = 6$$

$$w = 40^\circ\text{C}$$

$$z = 1.15w = 1.15 \times 40 = 46^\circ\text{C}$$

So, option 'c' is correct.

Scan for Video solution

**339. (d)**

$$x - 75 = 117 - x$$

$$x = 96$$

Scan for Video solution

**340. (a)**

Let n units are sold

S.P. - C.P. = Profit

$$60 \times n - 100 = 500$$

$$n = 10$$

$\Rightarrow$  If total cost is increased by 30% then let x% is increased in selling price to maintain the same level of profit.

$$\left(1 + \frac{x}{100}\right)60 \times 10 - 1.30 \times 100 = 500$$

$$x = 5\%$$

Scan for Video solution




**341. (b)**

Given that labour wages per day increase by  $1/5$  and the working hours per day decrease by  $1/24$ , so the new erection cost is

$$= \left(1 + \frac{1}{5}\right) \times \left(1 - \frac{1}{24}\right) \times 13,200 = 15,180$$



Scan for Video solution

**342. (c)**

$$\frac{2x}{60} + \frac{x}{30} + \frac{x}{10} = \frac{4x}{V_{avg}}$$

$$\frac{10x}{60} = \frac{4x}{V_{avg}}, \Rightarrow V_{avg} = 6 \times 4 = 24 \text{ km/h}$$



Scan for Video solution

**343. (d)**

Two-digit numbers between 1 to 100 = 90

$$\begin{array}{cc} 14 & 98 \\ (7 \times 2) & (7 \times 14) \end{array}$$

Numbers divisible by 7 =  $7 \times 2, 7 \times 3, 7 \times 4, \dots, 7 \times 14$ .

So, total numbers divisible by 7 = 13

Not divisible by 7 =  $90 - 13 = 77$

The probability of the selected number is not divisible by 7 =  $\frac{77}{90}$



Scan for Video solution

**344. (c)**

$$44 + 42 + 40 + \dots \dots \dots 2$$

$$2(22 + 21 + 20 + 19 \dots \dots \dots 1)$$

Sum of n terms of an Arithmetic series

$$= \frac{n(a_1 + a_n)}{2}$$

$$= 2\left(\frac{22 + 23}{2}\right) = 506$$



Scan for Video solution

**345. (a)**

Let,

$$\text{Total notes of Rs. } 20 = x$$

$$\text{Total notes of Rs. } 10 = y$$

$$\text{So, } x + y = 14 \dots \text{(i)}$$

And;

$$20x + 10y = 230 \dots \text{(ii)}$$

On solving (i) & (ii);

$$\text{We get, } x = 9, \text{ and } y = 5$$

Hence the number of notes of Rs. 10 are 5.

So, option 'a' is correct.



Scan for Video solution

**346. (d)**

Given;

$$(1.001)^{1259} = 3.52$$

$$(1.001)^{2062} = 7.85$$

Then,

$$(1.001)^{3321} = (1.001)^{1259} \times (1.001)^{2062}$$

$$= 3.52 \times 7.85 = 27.632 \approx 27.64$$

So, option 'D' is correct.



Scan for Video solution



**347. (c)**

$$\text{Mean } (M) = \frac{\sum_{i=1}^n x_i}{n}$$

$$M' = \frac{\sum_{i=1}^n (x_i + 7)}{n} = \frac{\sum_{i=1}^n x_i}{n} + \frac{7n}{n} = M + 7$$

$$\text{Standard deviation } \sigma = \sqrt{\frac{1}{N} \sum_{i=1}^n (M - x_i)^2}$$

On doubling each entry, the mean and  $x_i$  both will be doubled.

$$\text{Standard deviation} = \sqrt{\frac{1}{N} \sum_{i=1}^n 2^2 (M - x_i)^2} = 2\sigma$$



Scan for Video solution

**348. (b)**

The percentage of total shock absorber which is reliable and made by  $X = 0.6 \times 0.96 = 0.576$  or 57.6%

The percentage of total shock absorber which is reliable and made by  $Y = 0.4 \times 0.72 = 0.288$  or

28.8%.

The probability that a chosen shock absorber which is reliable, is made by Y

$$= \frac{0.288}{0.576 + 0.288} = \frac{0.288}{0.864}$$

$$\frac{288}{864} = 0.33 = 0.\bar{3}$$



Scan for Video solution

**349. (c)**

Only option (c) can be exactly divide by 3 so that it can be the answer. Its because no other option can be divided into three equal parts.

**Calculations:**

Let the number of toffees = x

R took  $1/3^{\text{rd}}$  but returned four, then the remaining toffees in the bowl =  $x - \frac{x}{3} + 4 = \frac{2x}{3} + 4$

S took  $1/4^{\text{th}}$  but returned three, then the remaining toffees in the bowl

$$= \frac{2x}{3} + 4 - \frac{1}{4} \left( \frac{2x}{3} + 4 \right) + 3 = \frac{x}{2} + 6$$

T took  $1/2$  of remaining but returned two, then the remaining toffees in the bowl

$$= \frac{1}{2} \left( \frac{x}{2} + 6 \right) + 2 = \frac{x}{4} + 5$$

$$\therefore \frac{x}{4} + 5 = 17 \Rightarrow x = 48$$



Scan for Video solution

**350. (a)**

Let total number of candidates = x

Initial voters of P = 0.4x

Initial voters of Q = 0.6x

As per new change in voters,

New voters of P will be,

$$= 0.4x - 0.15 \times 0.4x + 0.25 \times 0.6x = 0.49x$$

New voters of Q will be,

$$= 0.6x - 0.25 \times 0.6x + 0.15 \times 0.4x = 0.51x$$

$$\therefore 0.51x - 0.49x = 2$$

$$x = 100$$

So, option 'A' is correct.



Scan for Video solution

**351. (c)**

Let A and B be the number of orders each day and backlogs respectively.

$$4A + B = 7 \times 4 = 28$$

$$10A + B = 3 \times 10 = 30$$

Solving above equations we get  $A=1/3$  and  $B=80/3$ .

For 5 days to ship the orders

$$5A + B = T \times 5 \Rightarrow 5/3 + 80/3 = T \times 5$$

$$T = 5.66$$

So, minimum trucks required will be 6.



Scan for Video solution



### 352. (d)

$$\text{Quantity left after } n \text{ operations} = x \left(1 - \frac{y}{x}\right)^n$$

Where  $x$  and  $y$  are initial amount and amount of mixture replaced each time respectively.

$$= 10 \times 0.9^3$$

$$= 10 \times 0.729$$

$$= 7.29 \text{ lit}$$



Scan for Video solution



### 353. (b)

Four digits can be named as unit place, ten's place, hundred's place and thousand's place.





If the thousand's place is 3 (fixed) then total number of 4-digit numbers is



$3 \times 3 \times 3 = 27 - 2 = 25$  ( $\because$  3 unique digits are there and two numbers 3222 & 3333 are not possible)

If the thousand's place is 4 (fixed) then total number of 4-digit numbers is

$3 \times 3 \times 3 = 27 - 1 = 26$  ( $\because$  4222 is not possible)

Total numbers possible =  $25 + 26 = 51$



Scan for Video solution



### 354. (d)

Capacities (per day per worker) of skilled (s), semiskilled (ss) and unskilled (us) workers are calculated below:

$$5 \text{ sk} = \frac{1}{20}$$

$$8 \text{ ssk} = \frac{1}{25}$$

$$1 \text{ sk} = \frac{1}{100}$$

$$1 \text{ ssk} = \frac{1}{200}$$

$$10 \text{ usk} = \frac{1}{30}$$

$$1 \text{ usk} = \frac{1}{300}$$

So, 2 skilled, 6 semiskilled and 5 unskilled workers will have combined capacity (per day) of

$$\frac{2}{100} + \frac{6}{200} + \frac{5}{300} = \frac{12+18+10}{600}$$

$$= \frac{40}{600} = \frac{1}{15}$$

So, the time taken to build the wall by this team will be 15 days.



Scan for Video solution



# CHAPTER

2

# **ANALYTICAL APTITUDE**

## Logical Reasoning


- 1. [MCQ] [GATE-ME-2023: 1M]**

The symbols  $\bigcirc$ ,  $*$ ,  $\Delta$  and  $\square$  are to be filled, one in each box, as shown below.

The rules for filling in the four symbols are as follows.

- (1) Every row and every column must contain each of the four symbols.
  - (2) Every  $2 \times 2$  square delineated by bold lines must contain each of the four symbols.

Which symbol will occupy the box marked with ‘?’ in the partially filled figure?



- |                |               |
|----------------|---------------|
| (a) $\bigcirc$ | (b) $*$       |
| (c) $\Delta$   | (d) $\square$ |

2. [MCO] [GATE-EC-2023:2M]

A  $100 \text{ cm} \times 32 \text{ cm}$  rectangular sheet is folded 5 times. Each time the sheet is folded, the long edge aligns with its opposite side. Eventually, the folded sheet is a rectangle of dimensions  $100 \text{ cm} \times 1 \text{ cm}$ .

The total number of creases visible when the sheet is unfolded is



- 3. [MCQ] [GATE-EC-2023:1M]**

In a class of 100 students,

- (i) there are 30 students who neither like romantic movies nor comedy movies,
  - (ii) the number of students who like romantic movies is twice the number of students who like comedy movies, and
  - (iii) the number of students who like both romantic movies and comedy movies is 20.

How many students in the class like romantic movies?



4. [MCQ] [GATE-CE-2023: 1M]

There are 4 red, 5 green, and 6 blue balls inside a box. If  $N$  number of balls are picked simultaneously, what is the smallest value of  $N$  that guarantees there will be at least two balls of the same colour?

One cannot see the colour of the balls until they are picked.



5. [MCO] [GATE-CE-2023: 2M]

Consider a spherical globe rotating about an axis passing through its poles

There are three points P, Q and R situated respectively on the equator, north pole and midway between the equator and the north pole in the northern hemisphere. Let P, Q and R move with speeds  $v_p$ ,  $v_Q$ , and  $v_R$ , respectively.

Which one of the following options are CORRECT?

- (a)  $v_p < v_r < v_Q$       (b)  $v_p < v_Q < v_r$   
 (c)  $v_p > v_r > v_Q$       (d)  $v_p = v_r \neq v_Q$

## 6. [MCQ] [GATE-CE-2023: 1M]

A duck named Donald Duck says “All ducks always lie.”

Based only on the information above, which one of the following statements can be logically inferred with certainty?

- (a) Donald Duck always lies.
- (b) Donald Duck always tells the truth.
- (c) Donald Duck’s statement is true.
- (d) Donald Duck’s statement is false.

## 7. [MCQ] [GATE-CS-2023:1M]

A palindrome is a word that reads the same forwards and backwards. In a game of words, a player has the following two plates painted with letters.



From the additional plates given in the options, which one of the combinations of additional plates would allow the player to construct a five-letter palindrome. The player should use all the five plates exactly once. The plates can be rotated in their plane.

- |     |  |  |  |
|-----|--|--|--|
| (a) |  |  |  |
| (b) |  |  |  |
| (c) |  |  |  |
| (d) |  |  |  |

## 8. [MCQ] [GATE-ME-2022: 1M]

Four girls P, Q, R and S are studying languages in a University. P is learning French and Dutch. Q is learning Chinese and Japanese. R is learning Spanish and French. S is learning Dutch and Japanese.

Given that: French is easier than Dutch; Chinese is harder than Japanese; Dutch is easier than Japanese, and Spanish is easier than French.

Based on the above information, which girl is learning the most difficult pair of languages?

- (a) P
- (b) Q
- (c) R
- (d) S

## 9. [MCQ] [GATE-EC-2022:2M]

In a class of five students P, Q, R, S and T, only one student is known to have copied in the exam. The disciplinary committee has investigated the situation and recorded the statements from the students as given below.

**Statement of P:** R has copied in the exam.

**Statement of Q:** S has copied in the exam.

**Statement of R:** P did not copy in the exam.

**Statement of S:** Only one of us is telling the truth.

**Statement of T:** R is telling the truth.

The investigating team had authentic information that S never lies.


Based on the information given above, the person who has copied in the exam is

- (a) R
- (b) P
- (c) Q
- (d) T

## 10. [MCQ] [GATE-EC-2022:1M]

An art gallery engages a security guard to ensure that the items displayed are protected. The diagram below represents the plan of the gallery where the boundary walls are opaque. The location the security guard posted is identified such that all the inner space (shaded region in the plan) of the gallery is within the line of sight of the security guard.


If the security guard does not move around the posted location and has a 360° view, which one of the following correctly represents the set of ALL possible locations among the locations P, Q, R and S, where the security guard can be posted to watch over the entire inner space of the gallery.



- (a) P and Q
- (b) Q
- (c) Q and S
- (d) R and S

**11. [MCQ]****[GATE-EC-2022:1M]**

The figure shows a grid formed by a collection of unit squares. The unshaded unit square in the grid represents a hole.








What is the maximum number of squares without a “hole in the interior” that can be formed within the  $4 \times 4$  grid using the unit squares as building blocks

- (a) 15
  - (b) 20
  - (c) 21
  - (d) 26
- 12. [MCQ]** **[GATE-EE-2022: 2M]**

An ant is at the bottom-left corner of a grid (point P) as shown above. It aims to move to the top-right corner of the grid. The ant moves only along the lines marked in the grid such that the current distance to the top-right corner strictly decreases.

Which one of the following is a part of a possible trajectory of the ant during the movement?



- (a) 
- (b) 
- (c) 
- (d) 

**13. [MCQ]****[GATE-EE-2022: 2M]**

The letters P, Q, R, S, T and U are to be placed one per vertex on a regular convex hexagon, but not necessarily in the same order.

**Consider the following statements:**

- ★ The line segment joining R and S is longer than the line segment joining P and Q.
- ★ The line segment joining R and S is perpendicular to the line segment joining P and Q.
- ★ The line segment joining R and U is parallel to the line segment joining T and Q.

Based on the above statements, which one of the following options is CORRECT?

- (a) The line segment joining R and T is parallel to the line segment joining Q and S
- (b) The line segment joining T and Q is parallel to the line joining P and U
- (c) The line segment joining R and P is perpendicular to the line segment joining U and Q
- (d) The line segment joining Q and S is perpendicular to the line segment joining R and P

**14. [MCQ]****[GATE-EE-2022 : 1M]**

Given below are two statements and four conclusions drawn based on the statements.

**Statement 1:** Some bottles are cups.

**Statement 2:** All cups are knives.

**Conclusion I:** Some bottles are knives.

**Conclusion II:** Some knives are cups.

**Conclusion III:** All cups are bottles.

**Conclusion IV:** All knives are cups.

Which one of the following options can be logically inferred?

- (a) Only conclusion I and conclusion II are correct
- (b) Only conclusion II and conclusion III are correct
- (c) Only conclusion II and conclusion IV are correct
- (d) Only conclusion III and conclusion IV are correct

## 15. [MCQ]

[GATE-CE-2022:1M]

Given the statements:

- ★ P is the sister of Q.
- ★ Q is the husband of R.
- ★ R is the mother of S.
- ★ T is the husband of P.

Based on the above information, T is \_\_\_\_\_ of S.

- (a) the grandfather      (b) the father  
 (c) an uncle              (d) a brother

## 16. [MCQ]


[GATE-CE-2022:2M]

In the square grid shown on the left, a person standing at P2 position is required to move to P5 position.


The only movement allowed for a step involves, “two moves along one direction followed by one move in a perpendicular direction”. The permissible directions for movement are shown as dotted arrows in the right.

For example, a person at a given position Y can move only to the positions marked X on the right.

Without occupying any of the shaded squares at the end of each step, the minimum number of steps required to go from P2 to P5 is



- (a) 4  
 (c) 5




- (b) 6  
 (d) 7

## 17. [MCQ]

[GATE-CE-2022:2M]

An ant walks in a straight line on a plane leaving behind a trace of its movement. The initial position of the ant is at point P facing east.


The ant first turns  $72^\circ$  anticlockwise at P, and then does the following two steps in sequence exactly FIVE times before halting.




1. moves forward for 10 cm.

2. turns  $144^\circ$  clockwise.


The pattern made by the trace left behind by the ant is




$$PQ = QR = RS = ST = TP = 10 \text{ cm}$$



$$PQ = QR = RS = ST = TU = UP = 10 \text{ cm}$$



$$SQ = QT = TR = RP = PS = 10 \text{ cm}$$



$$SW = WR = RP = PT = TQ = QU = US = 10 \text{ cm}$$

## 18. [MCQ]

[GATE-CS-2022:2M]

The corners and mid-points of the sides of a triangle are named using the distinct letters P, Q, R, S, T and U, but not necessarily in the same order. Consider the following statements:

- The line joining P and R is parallel to the line joining Q and S.

- P is placed on the side opposite to the corner T.
- S and U cannot be placed on the same side.

Which one of the following statements is correct based on the above information?

- P cannot be placed at a corner
- S cannot be placed at a corner
- U cannot be placed at a mid-point
- R cannot be placed at a corner

**19. [MCQ] [GATE-ME-2021: 1M]**

Ms. X came out of a building through its front door to find her shadow due to the morning sun falling to her right side with the building to her back. From this, it can be inferred that building is facing

- |           |           |
|-----------|-----------|
| (a) North | (b) East  |
| (c) West  | (d) South |

**20. [MCQ] [GATE-EE-2021 : 1M]**

Seven cars P, Q, R, S, T, U and V are parked in a row not necessarily in that order. The cars T and U should be parked next to each other. The cars S and V also should be parked next to each other, whereas P and Q cannot be parked next to each other. Q and S must be parked next to each other. R is parked to the immediate right of V. T is parked to the left of U.

Based on the above statements, the only INCORRECT option given below is:

- There are two cars parked in between Q and V.
- Q and R are not parked together.
- V is the only car parked in between S and R.
- Car P is parked at the extreme end.

**21. [MCQ] [GATE-CE-2021: 1M]**


Four persons P, Q, R and S are to be seated in a row, all facing the same direction, but not necessarily in the same order. P and R cannot sit adjacent to each other. S should be seated to the right of Q. The number of distinct seating arrangements possible is:

- |       |       |
|-------|-------|
| (a) 2 | (b) 4 |
| (c) 6 | (d) 8 |

**22. [MCQ] [GATE-CE-2020: 1M]**

After the inauguration of the new building, the Head of the Department (HoD) collated faculty preferences for office space. P wanted a room adjacent to the lab. Q wanted to be close to the lift. R wanted a view of the playground and S wanted a corner office.

Assuming that everyone was satisfied, which among the following shows a possible allocation?



**23. [MCQ] [GATE-CE-2020: 2M]**

Five friends P, Q, R, S and T went camping. At night, they had to sleep in a row inside the tent. P, Q and T refused to sleep next to R since he snored loudly. P and S wanted to avoid Q as he usually hugged people in sleep.

Assuming everyone was satisfied with the sleeping arrangements, what is the order in which they slept?

- |           |           |
|-----------|-----------|
| (a) RSPTQ | (b) SPRTQ |
| (c) QRSPQ | (d) QTSPR |

**24. [MCQ]****[GATE-ME-2019: 2M]**

M and N had four children P, Q, R and S. Of them, only P and R were married. They had children X and Y respectively. If Y is a legitimate child of W, which one of the following statements is necessarily FALSE?

- (a) M is the grandmother of Y
- (b) R is the father of Y
- (c) W is the wife of R
- (d) W is the wife of P

**25. [MCQ]****[GATE-ME-2019: 2M]**

Under a certain legal system, prisoners are allowed to make one statement. If their statement turns out to be true then they are hanged. If the statement turns out to be false then they are shot. One prisoner made a statement and the judge had no option but to set him free. Which one of the following could be that statement?

- (a) I did not commit the crime
- (b) I committed the crime
- (c) I will be shot
- (d) You committed the crime

**26. [MCQ]****[GATE-EC-2019:2M]**

Five people P, Q, R, S and T work in a bank. P and Q don't like each other but have to share an office till T gets a promotion and moves to the big office next to the garden. R, who is currently sharing an office with T wants to move to the adjacent office with S, the handsome new intern. Given the floor plan, what is the current location of Q, R and T?

(O = Office, WR = Washroom)

|    |           |    |         |         |
|----|-----------|----|---------|---------|
| WR | O1<br>P,Q | O2 | O3<br>R | O4<br>S |
|    |           |    |         |         |
|    |           |    |         |         |
|    |           |    |         |         |

(a) Manager T      Teller 1      Teller 2

Entry      Garden

|    |           |    |         |            |
|----|-----------|----|---------|------------|
| WR | O1<br>P,Q | O2 | O3<br>T | O4<br>R, S |
|    |           |    |         |            |
|    |           |    |         |            |
|    |           |    |         |            |

(b) Manager      Entry      Teller 1      Teller 2

Garden

|    |           |    |            |         |
|----|-----------|----|------------|---------|
| WR | O1<br>P,Q | O2 | O3<br>R, T | O4<br>S |
|    |           |    |            |         |
|    |           |    |            |         |
|    |           |    |            |         |

(c) Manager      Entry      Teller 1      Teller 2

Garden

|    |         |    |         |         |
|----|---------|----|---------|---------|
| WR | O1<br>P | O2 | O3<br>R | O4<br>S |
|    |         |    |         |         |
|    |         |    |         |         |
|    |         |    |         |         |

(d) Manager      Entry      Teller 1      Teller 2

Garden

**27. [MCQ]****[GATE-EC-2019:1M]**

Five different books (P, Q, R, S, T) are to be arranged on a shelf. The books R and S are to be arranged first and second, respectively from the right side of the shelf. The number of different orders in which P, Q and T may be arranged is\_\_\_\_\_.

- (a) 2
- (b) 6
- (c) 12
- (d) 120

**28. [MCQ]****[GATE-EC-2019:1M]**

When he did not come home, she \_\_\_\_\_ him lying dead on the roadside somewhere.

- (a) concluded      (b) looked  
(c) notice      (d) pictured

**29. [MCQ] [GATE-EC-2019:2M]**

Four people are standing in a line facing you. They are Rahul, Mathew, Seema and Lohit. One is an engineer, one is a doctor, one a teacher and another a dancer. You are told that:

You are told that:

1. Mathew is not standing next to Seema
  2. There are two people standing between Lohit and the engineer
  3. Rahul is not a doctor
  4. The teacher and the dancer are standing next to each other
  5. Seema is turning to her right to speak to the doctor standing next to her.

Who among them is an engineer?



**30. [MCQ] [GATE-CE-2019: 2M]**

P, Q, R, S and T are related and belong to the same family. P is the brother of S. Q is the wife of P. R and T are the children of the siblings P and S respectively. Which one of the following statements is necessarily FALSE

- (a) S is the aunt of R
  - (b) S is the aunt of T
  - (c) S is the sister-in-law of Q
  - (d) S is the brother of P

31. [MCO] [GATE-CS-2019:2M]

The police arrested four criminals – P, Q, R and S. The criminals knew each other. They made the following statements:

P says “Q committed the crime.”

Q says “S committed the crime.”

R says “I did not do it.”

S says “What Q said about me is false.”

Assume only one of the arrested four committed the crime and only one of the statements made above is true. Who committed the crime?



32. [MCQ] [GATE-CS-2019:2M]



Three of the five students are allocated to a hostel put in special requests to the warden. Given the floor plan of the vacant rooms, select the allocation plan that will accommodate all their requests.



Request by X: Due to pollen allergy, I want to avoid a wing next to the garden.

Request by Y: I want to live as far from the washrooms as possible, since I am very sensitive to smell.

Request by Z: I believe in Vaastu and so I want to stay in the South-west wing.

The shaded rooms are already occupied. WR is washroom.



**33. [MCQ]****[GATE-ME-2017: 2M]**

All people in a certain island are either 'Knights' or 'Knaves' and each person knows every other person's identity. Knights NEVER lie, and knaves ALWAYS lie.

P says "Both of us are knights". Q says "None of us are knaves".

Which one of the following can be logically inferred from the above?

- (a) Both P and Q are knights
- (b) P is a knight; Q is a knave
- (c) Both P and Q are knaves
- (d) The identities of P, Q cannot be determined

**34. [MCQ]****[GATE-ME-2017: 1M]**

P, Q, and R talk about S's car collection. P states that S has at least 3 cars. Q believes that S has less than 3 cars. R indicates that to his knowledge, S has at least one car. Only one of P, Q and R is right. The number of cars owned by S is

- (a) 0
- (b) 1
- (c) 3
- (d) Cannot be determined

**35. [MCQ]****[GATE-EC-2017:2M]**

Each of P, Q, R, S, W, X, Y and Z has been married at most once. X and Y are married and have two children P and Q. Z is the grandfather of the daughter S of P. Further, Z and W are married and are parents of R. Which one of the following must necessarily be FALSE?

- (a) X is the mother-in-law of R
- (b) P and R are not married to each other
- (c) P is a son of X and Y
- (d) Q cannot be married to R

**36. [MCQ]****[GATE-EC-2017:1M]**

A rule states that in order to drink beer, one must be over 18 years old. In a bar, there are 4 people. P is 16 years old. Q is 25 years old. R is drinking milkshake and S is drinking a beer. What must be checked to ensure that the rule is being followed?

- (a) Only P's drink
- (b) Only P's drink and S's age
- (c) Only S's age
- (d) Only P's drink, Q's drink and S's age

**37. [MCQ]****[GATE-EC-2017:2M]**

S, T, U, V, W, X, Y and Z are seated around a circular table. T's neighbours are Y and V. Z is seated third to the left of T and second to the right of S. U's neighbours are S and Y; and T and W are not seated opposite each other. Who is third to the left of V?

- (a) X
- (b) W
- (c) U
- (d) T

**38. [MCQ]****[GATE-CS, EE-2017: 1M]**

Rahul, Murali, Srinivas and Arul are seated around a square table. Rahul is sitting to the left of Murali. Srinivas is sitting to the right of Arul. Which of the following pairs are seated opposite each other?

- (a) Rahul and Murali
- (b) Srinivas and Arul
- (c) Srinivas and Murali
- (d) Srinivas and Rahul



Which one of the following statements must be TRUE based on the above?

- (a) When Shiva plays chess with Leela and Pavithra, he often loses.
- (b) Leela is the oldest of the three.
- (c) Shiva is a better chess player than Pavithra.
- (d) Pavithra is the youngest of the three.


**46. [MCQ] [GATE-EC-2016:1M]**

M has a son Q and a daughter R. He has no other children. E is the mother of P and daughter-in law of M. How is P related to M?

- (a) P is the son-in-law of M
- (b) P is the grandchild of M
- (c) P is the daughter in law of M
- (d) P is the grandfather of M

**47. [MCQ] [GATE-CS, EE-2016 : 2M]**

In a  $2 \times 4$  rectangle grid shown below, each cell is a rectangle. How many rectangles can be observed in the grid?



- |        |        |
|--------|--------|
| (a) 21 | (b) 27 |
| (c) 30 | (d) 36 |

**48. [MCQ] [GATE-CS, EE-2016 : 2M]**

All hill-stations have a lake. Ooty has two lakes.

Which of the statement(s) below is/are logically valid and can be inferred from the above sentences?

- (i) Ooty is not a hill-station.
  - (ii) No hill-station can have more than one lake.
- |                       |                          |
|-----------------------|--------------------------|
| (a) (i) only          | (b) (ii) only            |
| (c) Both (i) and (ii) | (d) Neither (i) nor (ii) |

**49. [MCQ] [GATE-CS, CE-2016: 1M]**

If ‘relftaga’ means carefree, ‘otaga’ means careful and ‘fertaga’ means careless, which of the following could mean ‘aftercare’?

- |             |             |
|-------------|-------------|
| (a) zentaga | (b) tagafer |
| (c) tagazen | (d) refffer |

**50. [MCQ]**

[GATE-ME-2015: 1M]

Tanya is older than Eric.

Cliff is older than Tanya

Eric is older than cliff.

If the first two statements are true, than the third statement is:

- |               |                       |
|---------------|-----------------------|
| (a) True      | (b) False             |
| (c) Uncertain | (d) Data insufficient |

**51. [MCQ]**

[GATE-EC-2015:2M]

A cube of side 3 units is formed using a set of smaller cubes of side 1 unit. Find the proportion of the number of faces of the smaller cubes visible to those which are NOT visible.

- |           |           |
|-----------|-----------|
| (a) 1 : 4 | (b) 1 : 3 |
| (c) 1 : 2 | (d) 2 : 3 |

**52. [MCQ]**

[GATE-EE, CS-2015: 2M]

Four branches of a company are located at M, N, O, and P. M is north of N at a distance of 4 km. P is south of O at a distance of 2 km N is southeast of O by 1 km What is the distance between M and P in km?

- |          |           |
|----------|-----------|
| (a) 5.34 | (b) 6.74  |
| (c) 28.5 | (d) 45.49 |

**53. [MCQ]**

[GATE-CS, CE-2015: 2M]

The head of a newly formed government desires to appoint five of the six selected members P, Q, R, S, T and U to portfolios of Home, Power, Defense, Telecom, and Finance. U does not want any portfolio if S gets one of the five. R wants either Home or Finance or no portfolio. Q says that if S gets either Power or Telecom, then she must get the other one T insists on a portfolio if P gets one

Which is the valid distribution of portfolios?

- |  |
|--|
| (a) P-Home Q-Power. R-Defense. S-Telecom. T-Finance  |
| (b) R-Home, S-Power. P-Defense, Q-Telecom, T-Finance |
| (c) P-Home. Q-Power. T-Defense. S-Telecom. U-Finance |
| (d) Q Home. U Power. T-Defense. R-Telecom P-Finance  |

**54. [MCQ] [GATE-ME, EC-2014: 2M]**

If ‘KCLFTSB’ stands for ‘best of luck’ and ‘SHSWDG’ stands for ‘good wishes’, which of the following indicates ‘ace the exam’?

- (a) MCHTX      (b) MXHTC
- (c) XMHCT      (d) XMHTC

**55. [MCQ] [GATE-ME, EC-2014: 2M]**

Lights of four colors (red, blue, green, yellow) are hung on a ladder. On every step of the ladder there are two lights. If one of the lights is red, the other light on that step will always be blue. If one of the lights on a step is green, the other light on that step will always be yellow. Which of the following statements is not necessarily correct?

- (a) The number of red lights is equal to the number of blue lights
- (b) The number of green lights is equal to the number of yellow lights
- (c) The sum of the red and green lights is equal to the sum of the yellow and blue lights
- (d) The sum of the red and blue lights is equal to the sum of the green and yellow lights

**56. [NAT] [GATE-EE, CS-2014 : 2M]**

When a point inside of a tetrahedron (a solid with four triangular surfaces) is connected by straight lines to its corners, how many (new) internal planes are created with these lines? \_\_\_\_\_.

**57. [MCQ] [GATE-EE, CS-2014 : 2M]**

A dance programme is scheduled for 10.00 a.m. Some students are participating in the programme and they need to come an hour earlier than the start of the event. These students should be accompanied by a parent. Other students and parents should come in time for the programme. The instruction you think that is appropriate for this is

- (a) Students should come at 9.00 a.m. and parents should come at 10.00 a.m
- (b) Participating students should come at 9.00 a.m. accompanied by a parent, and other

parents and students should come by 10.00 a.m.

- (c) Students who are not participating should come by 10.00 a.m. and they should not bring their parents. Participating students should come at 9.00 a.m.
- (d) Participating students should come before 9.00 a.m. Parents who accompany them should come at 9.00 a.m. All others should come at 10.00 a.m.

**58. [MCQ] [GATE-CE-2014: 1M]**

Anuj, Bhola, Chandan, Dilip, Eswar and Faisal live on different floors in a six-storeyed building (the ground floor is numbered 1, the floor above it 2, and so on). Anuj lives on an even-numbered floor. Bhola does not live on an odd numbered floor. Chandan does not live on any of the floors below Faisal’s floor. Dilip does not live on floor number 2. Eswar does not live on a floor immediately above or immediately below Bhola. Faisal lives three floors above Dilip. Which of the following floor-person combinations is correct?

|     | Anuj | Bhola | Chandan | Dilip | Eswar | Faisal |
|-----|------|-------|---------|-------|-------|--------|
| (A) | 6    | 2     | 5       | 1     | 3     | 4      |
| (B) | 2    | 6     | 5       | 1     | 3     | 4      |
| (C) | 4    | 2     | 6       | 3     | 1     | 5      |
| (D) | 2    | 4     | 6       | 1     | 3     | 5      |

**59. [MCQ] [GATE-CE-2014: 2M]**

In a group of four children, Som is younger to Riaz. Shiv is elder to Ansu. Ansu is youngest in the group. Which of the following statements is/are required to find the eldest child in the group?

**Statements**

1. Shiv is younger to Riaz.
  2. Shiv is elder to Som.
- (a) Statement 1 by itself determines the eldest child.
  - (b) Statement 2 by itself determines the eldest child.
  - (c) Statements 1 and 2 are both required to determine the eldest child.
  - (d) Statements 1 and 2 are not sufficient to determine the eldest child.

**60. [MCQ]****[GATE-CE-2013: 2M]**

Abhishek is elder to Savar.

Savar is younger to Anshul.

Which of the given conclusions is logically valid and is inferred from the above statements?

- (a) Abhishek is elder to Anshul
- (b) Anshul is elder to Abhishek
- (c) Abhishek and Anshul are of the same age
- (d) No conclusion follows


**61. [MCQ]****[GATE-EC, EE-2012 : 2M]**

There are eight bags of rice looking alike, seven of which have equal weight and one is slightly heavier. The weighing balance is of unlimited capacity. Using this balance, the minimum number of weighings required to identify the heavier bag is

- |       |       |
|-------|-------|
| (a) 2 | (b) 3 |
| (c) 4 | (d) 8 |

**Logical Venn Diagrams****62. [MCQ]****[GATE-EC-2023:2M]**

Which one of the following options represents the given graph?



- |                           |                         |
|---------------------------|-------------------------|
| (a) $f(x) = x^2 2^{- x }$ | (b) $f(x) = x 2^{- x }$ |
| (c) $f(x) =  x  2^{-x}$   | (d) $f(x) = x 2^{-x}$   |

**63. [MCQ]****[GATE-ME-2022: 2M]**

Given below are three conclusions drawn based on the following three statements.

Statement 1: All teachers are professors.

Statement 2: No professor is a male.

Statement 3: Some males are engineers.

Conclusion I: No engineer is a professor.

Conclusion II: Some engineers are professors.

Conclusion III: No male is a teacher.

Which one of the following options can be logically inferred?

- (a) Only conclusion III is correct
- (b) Only conclusion I and conclusion II are correct
- (c) Only conclusion II and conclusion III are correct
- (d) Only conclusion I and conclusion III are correct

**64. [MCQ]****[GATE-CE-2022:1M]**

A survey of 450 students about their subjects of interest resulted in the following outcome.

- 150 students are interested in Mathematics.
  - 200 students are interested in Physics.
  - 175 students are interested in Chemistry.
  - 50 students are interested in Mathematics and Physics.
  - 60 students are interested in Physics and Chemistry.
  - 40 students are interested in Mathematics and Chemistry.
  - 30 students are interested in Mathematics, Physics and Chemistry.
  - Remaining students are interested in Humanities.
- Based on the above information, the number of student's interest in Humanities is.
- |        |        |
|--------|--------|
| (a) 10 | (b) 30 |
| (c) 40 | (d) 45 |

**65. [MCQ]****[GATE-CE-2022:2M]**

Given below are two statements and four conclusions drawn based on the statements.

Statement 1: Some soaps are clean.

Statement 2: All clean objects are wet.

Conclusion I: Some clean objects are soaps.

Conclusion II: No clean object is a soap.

Conclusion III: Some wet objects are soaps.

Conclusion IV: All wet objects are soaps.

Which one of the following options can be logically inferred?

- (a) Only conclusion I is correct
- (b) Either conclusion I or conclusion II is correct
- (c) Either conclusion III or conclusion IV is correct
- (d) Only conclusion I and conclusion III are correct

**66. [MCQ] [GATE-CS-2022:1M]**

Given below are four statements.

**Statement 1:** All students are inquisitive.

**Statement 2:** Some students are inquisitive.

**Statement 3:** No student is inquisitive.

**Statement 4:** Some students are not inquisitive.

From the given four statements, find the two statements that CANNOT BE TRUE simultaneously, assuming that there is at least one student in the class.

- (a) Statement 1 and Statement 3
- (b) Statement 1 and Statement 2
- (c) Statement 2 and Statement 4
- (d) Statement 3 and Statement 4

**67. [MCQ] [GATE-CE-2021: 1M]**

$\oplus$  and  $\odot$  are two operators on numbers p and q such that  $p \oplus q = \frac{p^2+q^2}{pq}$  and  $p \odot q = \frac{p^2}{q}$ . If  $x \oplus y = 2 \odot 2$ , then  $x =$

- |                    |          |
|--------------------|----------|
| (a) $\frac{y}{2}$  | (b) $y$  |
| (c) $\frac{3y}{2}$ | (d) $2y$ |

**68. [MCQ] [GATE-CE-2021: 1M]**

$\oplus$  and  $\odot$  are two operators on numbers p and q such that  $p \odot q = p - q$ , and  $p \oplus q = p \times q$ .

Then,  $(9 \odot (6 \oplus 7)) \odot (7 \oplus (6 \odot 5)) =$

- (a) 40
- (b) -26
- (c) -33
- (d) -40

**69. [MCQ] [GATE-CE-2021: 2M]**

On a planar field, you travelled 3 units East from a point O. Next you travelled 4 units South to arrive at point P. Then you travelled from P in the North-East direction such that you arrive at a point that is 6 units East of point O. Next, you travelled in the North-West direction, so that you arrive at point Q that is 8 units North of point P.

The distance of point Q to point O, in the same units, should be\_\_\_\_\_

- (a) 3
- (b) 5
- (c) 4
- (d) 6

**70. [MCQ] [GATE-ME-2021: 2M]**

Given below are two statements 1 and 2, and two conclusions I and II.

**Statement 1:** All entrepreneurs are wealthy.

**Statement 2:** All wealthy are risk seekers.

**Conclusion I:** All risk seekers are wealthy.

**Conclusion II:** Only some entrepreneurs are risk seekers.

Based on the above statements and conclusions, which one of the following options is CORRECT?

- (a) Only conclusion I is correct
- (b) Only conclusion II is correct
- (c) Neither conclusion I nor II is correct
- (d) Both conclusions I and II are correct

**71. [MCQ] [GATE-EC-2021:2M]**

Given below are two statements and two conclusions.

**Statement 1:** All purple are green.

**Statement 2:** All black are green.

**Conclusion I:** Some black are purple.

**Conclusion II:** No black is purple.

Based on the above statements and conclusions, which one of the following options is logically CORRECT?

- (a) Only conclusion I is correct.
- (b) Only conclusion II is correct
- (c) Either conclusion I or II is correct.
- (d) Both conclusion I and II are correct

**72. [MCQ] [GATE-CE-2021: 1M]**

In a company, 35% of the employees drink coffee, 40% of the employees drink tea and 10% of the employees drink both tea and coffee. What % of employees drink neither tea nor coffee?

- (a) 15
- (b) 25
- (c) 35
- (d) 40

**73. [MCQ] [GATE-CE-2021: 2M]**

Among the option given below, the statements that logically follows from the two statements 1 and 2 above, is:

1. Some football players play cricket.
  2. All cricket players play hockey.
- (a) No football player plays hockey.
  - (b) Some football players play hockey.
  - (c) All football players play hockey.
  - (d) All hockey players play football.

**74. [MCQ] [GATE-CS-2021: 2M]**

Given below are two statements 1 and 2, and two conclusions I and II.

Statement 1: All bacteria are microorganisms.

Statement 2: All pathogens are microorganisms.

Conclusion I: Some pathogens are bacteria.

Conclusion II: All pathogens are not bacteria.

Based on the above statements and conclusions, which one of the following options is logically CORRECT?

- (a) Only conclusion I is correct
- (b) Only conclusion II is correct
- (c) Either conclusion I or II is correct.
- (d) Neither conclusion I nor II is correct.

**75. [MCQ] [GATE-CE-2020: 1M]**

If 0, 1, 2, ..., 7, 8, 9 are coded as O, P, Q, ..., V, W, X, then 45 will be coded as \_\_\_\_.

- (a) TS
- (b) ST
- (c) SS
- (d) SU

**76. [MCQ] [GATE-CE-2020: 2M]**

In a school of 1000 students, 300 students play chess and 600 students play football. If 50 students play both chess and football, the number of students who play neither is \_\_\_\_.

- (a) 200
- (b) 150
- (c) 100
- (d) 5

**77. [MCQ] [GATE-CE-2019: 1M]**

If  $E = 10$ ;  $J = 20$ ;  $O = 30$ ; and  $T = 40$ , what will be  $P + E + S + T$ ?

- (a) 51
- (b) 82
- (c) 120
- (d) 164

**78. [MCQ] [GATE-CE-2019: 2M]**

In a sports academy of 300 people, 105 play only cricket, 70 play only hockey, 50 play only football, 25 play both cricket and hockey, 15 play both hockey and football and 30 play both cricket and football. The rest


of them play all three sports. What is the percentage of people who play at least two sports?

- (a) 23.30
- (b) 25.00
- (c) 28.00
- (d) 50.00

**79. [MCQ] [GATE-CS-2019:2M]**

In the given diagram, teachers are represented in the triangle, researchers in the circle and administrators in the rectangle. Out of the total number of the people, the percentage of administrators shall be in the range of \_\_\_\_\_.

Teachers



- (a) 0 to 15
- (b) 16 to 30
- (c) 31 to 45
- (d) 46 to 60

**80. [MCQ] [GATE-CS-2019:2M]**

In a college, there are three student clubs. Sixty students are only in the Drama club. 80 students are only in the Dance club. 30 students are only in the Maths club. 40 students are in both Drama and Dance clubs. 12 students are in both Dance and Maths clubs. 7 students are in both Drama and Maths clubs, and 2 students are in all the clubs. If 75% of the students in the college are not in any of these clubs, then the total number of students in the college is \_\_\_\_.

- (a) 1000
- (b) 975
- (c) 900
- (d) 225

**81. [MCQ] [GATE-CE-2018: 2M]**

Each of the letters arranged as below represents a unique integer from 1 to 9. The letters are positioned in the figure such that  $(A \times B \times C)$ ,  $(B \times G \times E)$  and  $(D \times E \times F)$  are equal. Which integer among the following choices cannot be represented by the letters A, B, C, D, E, F or G?

|   |   |   |
|---|---|---|
| A |   | D |
| B | G | E |
| C |   | F |



**82. [MCQ] [GATE-CE-2018: 2M]**

Each of the letters in the figure below represents a unique integer from 1 to 9. The letters are positioned in the figure such that each of  $(A + B + C)$ ,  $(C + D + E)$ ,  $(E + F + G)$  and  $(G + H + K)$  is equal to 13. Which integer does  $E$  represent?

|   |   |   |
|---|---|---|
| A | B | C |
|   | D |   |
| E | F | G |
|   | H | K |

- (a) 1
  - (b) 4
  - (c) 6
  - (d) 7

83. [MCQ] [GATE-ME-2018: 2M]

Forty students watched films A, B and C over a week. Each student watched either only one film or all three. Thirteen students watched film A, sixteen students watched film B and nineteen students watched film C. How many students watched all three films?



**84. [MCQ] [GATE-ME-2018: 2M]**

Consider the following three statements:

- (i) Some roses are red.
  - (ii) All red flowers fade quickly.
  - (iii) Some roses fade quickly

Which of the following statements can be logically inferred from the above statements?

- (a) If (i) is true and (ii) is false, then (iii) is false.
  - (b) If (i) is true and (ii) is false, then (iii) is true.
  - (c) If (i) and (ii) are true, then (iii) is true.
  - (d) If (i) and (ii) are false, then (iii) is false.

**85. [MCQ] [GATE-EC-2017:1M]**

500 students are taking one or more courses out of Chemistry, Physics, and Mathematics. Registration records indicate course enrolment as follows: Chemistry (329), Physics (186), Mathematics (295), Chemistry and Physics (83), Chemistry and Mathematics (217), and Physics and Mathematics (63). How many students are taking all 3 subjects?



**86. [MCQ] [GATE-EC-2017:1M]**  
Some tables are shelves. Some shelves are chairs. All chairs are benches. Which of the following conclusions can be deduced from the preceding sentences?

- i. At least one bench is a table
  - ii. At least one shelf is a bench
  - iii. At least one chair is a table
  - iv. All benches are chairs

(a) Only i

(b) Only ii

(c) Only ii and iii

(d) Only iv

87. [MCQ] [GATE-CE-2017: 1M]

Consider the following sentences:  
All benches are beds. No bed is a bulb. Some bulbs  
are lamps.

Which of the following can be inferred?



88. [MCQ] [GATE-CE-2017: 2M]  
P, Q, R, S, T and U are seated around a circular-table. R is seated two places to the right of Q. P is seated three places to the left of R. S is seated opposite U. If P and U now switch seats, which of the following must necessarily be true?


- (a) P is immediately to the right of R
  - (b) T is immediately to the left of P
  - (c) T is immediately to the left of P or P is immediately to the right of Q
  - (d) U is immediately to the right of R or P is immediately to the left of T



## Numerical Relations

- 102. [MCQ] [GATE-EC-2023:1M]**

How many rectangles are present in the given figure?






- 103. [MCQ] [GATE-ME-2022: 2M]**

Four cities P, Q, R and S are connected through one-way routes as shown in the figure. The travel time between any two connected cities is one hour. The boxes beside each city name describe the starting time of first train of the day and their frequency of operation. For example, from city P, the first trains of the day start at 8 AM with a frequency of 90 minutes to each of R and S. A person does not spend additional time at any city other than the waiting time for the next connecting train.

If the person starts from R at 7 AM and is required to visit S and return to R, what is the minimum time required?



- (a) 6 hours, 30 minutes  
 (b) 3 hours, 45 minutes  
 (c) 4 hours, 30 minutes  
 (d) 5 hours, 15 minutes

**104. [MCQ]****[GATE-ME-2021: 2M]**

Consider a square sheet of side 1 unit. The sheet is first folded along the main diagonal. This is followed by a fold along its line of symmetry. The resulting folded shape is again folded along its line of symmetry. The area of each face of the final folded shape, in square units, equal to

- |                    |                    |
|--------------------|--------------------|
| (a) $\frac{1}{4}$  | (b) $\frac{1}{8}$  |
| (c) $\frac{1}{16}$ | (d) $\frac{1}{32}$ |

**105. [MCQ]****[GATE-ME-2021: 1M]**

The front door of Mr. X's house faces East. Mr. X leaves the house, walking 50 m straight from the back door that is situated directly opposite to the front door. He then turns to his right, walks for another 50 m and stops. The direction of the point Mr. X is now located at with respect to the starting point is

- |                |                |
|----------------|----------------|
| (a) South-East | (b) North-East |
| (c) West       | (d) North-West |

**106. [MCQ]****[GATE-ME-2021: 1M]**

If  $\oplus \div \odot = 2$ ;  $\oplus \div \Delta = 3$ ;  $\odot + \Delta = 5$ ;  $\Delta \times \otimes = 10$ ,  
 then the value of  $(\otimes - \oplus)^2$ , is :

- |       |        |
|-------|--------|
| (a) 0 | (b) 1  |
| (c) 4 | (d) 16 |

**107. [MCQ]****[GATE-ME-2021: 1M]**

If  $\begin{cases} \oplus \text{"means"} -, \\ \otimes \text{"means"} \div, \\ \Delta \text{"means"} +, \\ \nabla \text{"means"} \times, \end{cases}$

then, the value of the expression

$$\Delta \oplus 3\Delta((4 \otimes 2) \nabla 4) =$$

- |        |          |
|--------|----------|
| (a) -1 | (b) -0.5 |
| (c) 6  | (d) 7    |


**108. [MCQ]****[GATE-CS-2020: 2M]**

If values of  $P = 3$ ,  $R = 27$ ,  $T = 243$ , then  $Q + S =$   
 \_\_\_\_\_.

- |        |         |
|--------|---------|
| (a) 40 | (b) 80  |
| (c) 90 | (d) 110 |

**109. [MCQ]****[GATE-ME-2020: 2M]**

Find the missing element in the following figure



- |       |       |
|-------|-------|
| (a) d | (b) e |
| (c) w | (d) y |

**110. [MCQ]****[GATE-ME-2020: 2M]**

P, Q, R and S are to be uniquely coded using  $\alpha$  and  $\beta$ . If P is coded as  $\alpha\alpha$  and Q as  $\alpha\beta$ , then R and S, respectively, can be coded as

- |                                     |                                     |
|-------------------------------------|-------------------------------------|
| (a) $\beta\alpha$ and $\alpha\beta$ | (b) $\beta\beta$ and $\alpha\alpha$ |
| (c) $\alpha\beta$ and $\beta\beta$  | (d) $\beta\alpha$ and $\beta\beta$  |

**111. [MCQ]****[GATE-EE-2020 : 2M]**

Select the next element of the series: Z, WV, RQP,

- 
- |          |          |
|----------|----------|
| (a) LKJI | (b) JIHG |
| (c) KJIH | (d) NMLK |

**112. [MCQ]****[GATE-ME-2019: 1M]**

If IMHO = JNIP; IDK = JEL; and SO = TP, then IDC = \_\_\_\_\_.

- |         |         |
|---------|---------|
| (a) JDE | (b) JED |
| (c) JDC | (d) JCD |

**113. [MCQ]****[GATE-EE-2019 : 2M]**

Consider five people – Mita, Ganga, Rekha, Lakshmi and Sana. Ganga is taller than both Rekha and Lakshmi. Lakshmi is taller than Sana. Mita is taller than Ganga.

Which of the following conclusions are true?

1. Lakshmi is taller than Rekha
  2. Rekha is shorter than Mita
  3. Rekha is taller than Sana
  4. Sana is shorter than Ganga
- |             |            |
|-------------|------------|
| (a) 1 and 3 | (b) 3 only |
| (c) 2 and 4 | (d) 1 only |

**114. [MCQ]****[GATE-EE-2019 : 1M]**

The missing number in the given sequence 343, 1331, \_\_\_\_\_, 4913 is









**ANSWER KEY**

- |                          |          |                 |                   |
|--------------------------|----------|-----------------|-------------------|
| 1. (b)                   | 2. (c)   | 3. (c)          | 4. (a)            |
| 5. (c)                   | 6. (d)   | 7. (b)          | 8. (b)            |
| 9. (b)                   | 10. (c)  | 11. (b)         | 12. (c)           |
| 13. (a)                  | 14. (a)  | 15. (c)         | 16. (c)           |
| 17. (c)                  | 18. (b)  | 19. (d)         | 20. (a)           |
| 21. (c)                  | 22. (c)  | 23. (a)         | 24. (d)           |
| 25. (c)                  | 26. (c)  | 27. (b)         | 28. (d)           |
| 29. (d)                  | 30. (b)  | 31. (c)         | 32. (d)           |
| 33. (d)                  | 34. (a)  | 35. (d)         | 36. (b)           |
| 37. (a)                  | 38. (c)  | 39. (a)         | 40. (a)           |
| 41. (b)                  | 42. (d)  | 43. (c)         | 44. (c)           |
| 45. (d)                  | 46. (b)  | 47. (c)         | 48. (d)           |
| 49. (c)                  | 50. (b)  | 51. (c)         | 52. (a)           |
| 53. (b)                  | 54. (b)  | 55. (d)         | 56. (6 to 6)      |
| 57. (b)                  | 58. (b)  | 59. (a)         | 60. (d)           |
| 61. (a)                  | 62. (a)  | 63. (a)         | 64. (d)           |
| 65. (d)                  | 66. (a)  | 67. (b)         | 68. (d)           |
| 69. (b)                  | 70. (c)  | 71. (c)         | 72. (c)           |
| 73. (b)                  | 74. (d)  | 75. (b)         | 76. (b)           |
| 77. (c)                  | 78. (b)  | 79. (c)         | 80. (c)           |
| 81. (b)                  | 82. (b)  | 83. (c)         | 84. (c)           |
| 85. (d)                  | 86. (b)  | 87. (d)         | 88. (c)           |
| 89. (d)                  | 90. (c)  | 91. (a)         | 92. (b)           |
| 93. (a)                  | 94. (c)  | 95. (d)         | 96. (a)           |
| 97. (d)                  | 98. (c)  | 99. (d)         | 100. (d)          |
| 101. (0.4896 to 0.04896) | 102. (c) | 103. (a)        | 104. (b)          |
| 105. (d)                 | 106. (b) | 107. (d)        | 108. (c)          |
| 109. (a)                 | 110. (d) | 111. (c)        | 112. (b)          |
| 113. (c)                 | 114. (c) | 115. (c)        | 116. (d)          |
| 117. (b)                 | 118. (b) | 119. (d)        | 120. (d)          |
| 121. (b)                 | 122. (d) | 123. (c)        | 124. (c, d)       |
| 125. (d)                 | 126. (c) | 127. (c)        | 128. (b)          |
| 129. (a)                 | 130. (c) | 131. (3 to 3)   | 132. (c)          |
| 133. (c)                 | 134. (c) | 135. (c)        | 136. (b)          |
| 137. (16 to 16)          | 138. (d) | 139. (45 to 45) | 140. (725 to 725) |
| 141. (a)                 | 142. (b) |                 |                   |

## SOLUTIONS

**1. (b)**

The box obtained after filling the symbols is.



Hence the symbol in place of ? is \*. Option (B) is correct.



Scan for Video solution

**2. (c)**

Number of creases after 1<sup>st</sup> folding = 1


Number of creases after 2<sup>nd</sup> folding =  $2^1$

Number of creases after 3<sup>rd</sup> folding =  $2^2$


Number of creases after 4<sup>th</sup> folding =  $2^3$

Number of creases after 5<sup>th</sup> folding =  $2^4$

Total number of creases after unfolding



Scan for Video solution

**3. (c)**

n (neither romantic nor comedy)

= total students – n (Romantic  $\cup$  comedy)

$$30 = 100 - (2x + x - 20) \Rightarrow 3x - 20 = 70$$

$$x = \frac{90}{3} = 30$$


$$n(\text{romantic}) = 2x = 2 \times 30 = 60$$

Scan for Video solution

**4. (a)**

There are balls of three different colours. In worst case the three picked ball are of different colours. But as the fourth ball is picked that has to be one of these three colours so, out of four balls there are definitely two balls of same colour. Hence, the minimum number of N is 4.

Scan for Video solution

**5. (c)**

For rotational motion, the speed is given by  $v = r\omega$   
Where,  $r$  is the distance of the point from the axis of rotation and  $\omega$  is the angular speed which is same for all the points here hence,

$$v \propto r$$

$$\therefore r_P > r_R > r_Q$$

$$\therefore v_P > v_R > v_Q$$

Scan for Video solution

6. (d)

If Donald Duck statements is true then he is also lying which contradicts the statement. So, Donald duck's statement is wrong.



Scan for Video solution



7. (b)

To construct 5 letter pelindrom out of 5 alphabets, first alphabet should be same as 5<sup>th</sup> one and 2<sup>nd</sup> alphabet should be same as 4<sup>th</sup> one while 3<sup>rd</sup> alphabet should be unique.

In options (A), (C) & (D) this condition can not be fulfilled. For option (B), the possible pelindromes are:

R A D A R

A R D R A



Scan for Video solution



8. (b)

From the given data, the languages in descending difficulty order is given below.

From the given data, the language in descending order of difficulty is given below.

Chinese &gt; Japanese &gt; Dutch &gt; French &gt; Spanish

Since girl Q is studying Chinese and Japanese hence Q is learning the most difficult pair of languages.




Scan for Video solution



9. (b)


It is given that S never lies and S is saying "**only one of us is telling the truth**" implies all the other's statements of P, Q, R & T are not true. So, we can say

- (i) R has not copied in the exam
- (ii) S has not copied in the exam
- (iii) P did copy in the exam
- (iv) R is not telling the truth




10. (c)

As the walls are opaque, the security guard from position P and R, cannot see the region CDB and ABE respectively as shown in the figure



From position Q and S, the whole gallery can be watched.




11. (b)


Number of squares without hole of size  $(1 \times 1)$   
 $= 4 \times 4 - 1 = 15$

Number of squares without hole of size  $(2 \times 2)$

$= ABCD, DCFE, PQHG, JADI, KNML = 5$




$\therefore$  Total number of squares without hole  
 $= 15 + 5 = 20$



12. (c)

The ant starts at the position P, and move up to top-right corner of the grid, the only possible way in the option is given in option (c). It is because in option (a) and (b), starting position of the ant is not P so that they both eliminated and in option (d) final destination is not given,

$\therefore$  option (c) is correct.




Scan for Video solution




13. (a)

As per the given statement in the question, a regular convex hexagon can be drawn as given below,



From the above figure, we can see that the line segment joining R and T is parallel to the line segment joining Q and S. And other given options is not satisfied.

$\therefore$  option (a) is correct.




Scan for Video solution




14. (a)

Given two statements can be represented in the given Venn diagram,



From the Venn diagram, only conclusion (I) and conclusion (II) follows.

$\therefore$  option (a) is correct



Scan for Video solution




15. (c)

$$T^+ \sim P^- \cdot Q^+ \sim R^- \\ \downarrow \\ S$$

$\cdot \rightarrow$  Sibling       $+ \rightarrow$  male  
 $- \rightarrow$  female       $\sim \rightarrow$  couple

$\therefore T$  is an uncle of s.



Scan for Video solution



16. (c)



P2 to Q4 – step (1)

Q4 to R2 – step (2)

R2 to P3 – step (3)

P3 to R4 – step (4)


R4 to P5 – step (5)



Scan for Video solution



17. (c)




After turning  $72^\circ$  anticlockwise at P the ant will walk 10 cm along PS. At S it takes a  $144^\circ$  clockwise turn and then move along SQ and it repeat these two operations 5 times before halting.




Scan for Video solution

**18. (b)**

From the given three conditions, the possible triangle is drawn below.




If we have to put S at a corner then according to the condition III, the possible triangle will be.



Since P is at the side opposite to corner T so, P can possibly be at 1 or 2 but this will not satisfy condition (I), hence S can not be at the corner.



Scan for Video solution

**19. (d)**

So, the building is facing south.



Scan for Video solution

**20. (a)**

From the given information, the following pattern can be written,

- (i) PTUQSVR
- (ii) TUQSVRP
- (iii) QSVRTUP
- (iv) QSVRPTU

From the above pattern, it is clear that, only one car is parked between Q and V.

So, option 'a' is correct.



Scan for Video solution

**21. (c)**

Given, P & R cannot sit adjacent to each other and S should be seated to the right side of Q

Hence, PQRS, PQSR, RQPS, RQSP, QRSP, QPRS

To separate R & S,

Q first and S at third place = 2 ways

Q second and S at third place = 2 ways

Q second and S at fourth place = 2 ways.

Total = 6 ways.



Scan for Video solution

**22. (c)**

From given information the best allocation of faculties is in option c.

P → near lab

Q → near lift

R → near play ground

S → at the corner



Scan for Video solution





**31. (c)**

If P's statement is correct then all other statements are wrong and so R's statements is also wrong which contradicts the P's statement.

Similaly, if we consider Q's and R's statements correct once at a time then there will be contradictions, but if we consider S's statements correct then there will not be any contradiction and R will be the criminal.



Scan for Video solution

**32. (d)**

Options B is wrong as X's condition is not satisfied.  
 Options A is wrong as Z's conditions is not satisfied.  
 Options C is wrong as Z's condition is not satisfied.  
 Options D satisfies all the conditions of X, Y, Z.



Scan for Video solution

**33. (d)**

P says "Both of us are knights"

Q says "None of us are knave"

If P and Q are knights then this condition supports both the statements.

If P & Q are knave then this condition also supports both the statements.

So, no clear indication for P & Q.



Scan for Video solution

**34. (a)**

If S has 0 cars, then only Q is right.

If S has 1 car, then both Q & R are right.

If S has 3 cars, then both P & R are right.


Since only one of P, Q and R can be right so S has 0 cars.




Scan for Video solution

**35. (d)**

X and Y are married and have two children P and Q



Z is grandfather of the daughters S of P and Z and W are married and are parents of R



So, P and R are a couple and since everyone is married at most once, Q cannot be married to R.



Scan for Video solution

**36. (b)**

**Given:** Age/drink for all the persons are

P → 16 /?, Q → 25 /?

R → ? / milkshake S → ? / beer

Rule for beer is age should be over 18 years.

So, we need to check P's drink and S's age to follow the rule.




Scan for Video solution


**37. (a)**

**Given:**


T's neighbours are Y and V



Z is seated third to the left of T and second to the right of S



U's neighbours are S and Y; T and W are not seated opposite each other.



∴ The person third to the left of V is X.




Scan for Video solution



38. (c)

The given information can be drawn in a figure given below,



So, option (c) is correct.




Scan for Video solution



39. (a)

As clearly mentioned that there are at least two men and two women, and given that none of the woman is right handed. It means two women are left handed. Also given that there are 3 right handed person. It means, there are at least 3 men because no woman can be right handed.

Now every woman has a left-handed person to her immediate right. It means both women are sitting next to each other and third person sitting right to 2<sup>nd</sup> woman is also a left-handed person, which cannot be a woman because if it is a woman then next person will be a left-handed person. But there are at least 3 right handed person. So that there are 4 men and 2 women.



So, option 'A' is correct.




Scan for Video solution




40. (a)

The given information can be drawn in a figure given below,



We can see in the above figure that the building 'V' is in the middle

So, option 'A' is correct.



Scan for Video solution




41. (b)

**Given Data:**


In this case, you will choose the box which was labeled as apples and oranges. It is because, all are labeled as wrong so that this box must be a single fruit box, i.e. apples box or oranges box. By this way one box can be identified, now other two can be identified easily because other two also wrongly labeled in which one has single fruit and other has two fruits.

So, option 'b' is correct.



42. (d)

We can use minimum of 3 additional ropes to divide the plot into four similar shapes.



43. (c)

Card with number 2 should be turned to check if the colour is red or blue. If the colour is red then proposition is true. Now a blue-coloured card should be turned to check if the number is even or odd. If the number is odd then the proposition is true.

$p \Rightarrow q$  Direct

$p \Rightarrow q$  Converse


$rp \Rightarrow rq$  Inverse

for ' $\Rightarrow$ '


Direct & contrapositive

are true

$rq \Rightarrow rp$  Contrapositive    Boy  $\Rightarrow$  Human being




44. (c)




The position of M is  $10 + 10\cos 45^\circ$  in x-direction and  $10\sin 45^\circ$  in y-direction i.e. (17.07, 7.07). Similarly, the position of N is (2.83, -7.83). The shortest distance

$$\begin{aligned} &= \sqrt{(17.07 - 2.83)^2 + (7.07 - (-7.83))^2} \\ &= \sqrt{202.78 + 222.01} \\ &= 20.61 \text{ km} \end{aligned}$$




45. (d)

From the passage it is clear that Shiva is oldest and Pavithra is youngest. Hence option D is correct.




46. (b)

M has a son Q and a daughter R and he has no other children



E is the mother of P and daughter-in law of M



As gender of  $P$  is not known, we can say  $P$  is the grandchild of  $M$ .



Scan for Video solution



**47. (c)**

To form a rectangle, we require 2 horizontal and two vertical lines. Here we need to select, 2 out of 5 vertical and 2 out of 3 horizontal lines.

Therefore, Number of rectangles =  ${}^5C_2 \times {}^3C_2 = 30$   
So, option 'C' is correct.



Scan for Video solution



**48. (d)**

Ooty is a hill-station or not which is not mentioned here so that statement (i) is wrong.

Also, it is not mentioned that no hill-station can have more than one lake, so that statement (ii) is wrong.  
So, option 'd' is correct.



Scan for Video solution



**49. (c)**

From the observation of the given words, it is clear that 'taga' refers to 'care'.

Fertaga → careless

Tagafer → lesscare

So, the correct word for 'aftercare' is 'tagazen'.




Scan for Video solution



**50. (b)**


From the two statements it is clear that cliff is oldest and Eric is youngest. So, statement three is wrong as Eric is younger than cliff.



**51. (c)**

$$\left\{ \begin{array}{l} \text{Volume of} \\ \text{unit 3 cube} \end{array} \right\} = \left\{ \begin{array}{l} \text{number of} \\ \text{unit 1 cube} \end{array} \right\} \times \left\{ \begin{array}{l} \text{volume of} \\ \text{unit 1 cube} \end{array} \right\}$$

$$(3)^3 = n \times 1^3 \Rightarrow n = 27$$



Total number of faces of unit 1 cube =  $6 \times 27 = 162$

Number of faces of larger cube which are visible = 6

Each face of unit 3 cube = 9 faces of unit 1 cube


So, number of faces of unit 1 which are visible

$$= 9 \times 6 = 54$$

Hence number of faces of unit 1 which are not visible


$$= 162 - 54 = 108$$

Required ratio =  $54 : 108 = 1 : 2$



**52. (a)**

The given information can be drawn into a figure as shown below,



In the  $\Delta$ NOS,

$$\sin 45^\circ = \frac{OS}{1} \Rightarrow OS = \frac{1}{\sqrt{2}} \text{ km}$$

$$PS = OP - OS = \left( 2 - \frac{1}{\sqrt{2}} \right) \text{ km}$$

$$PR = MN + PS = 4 + \left( 2 - \frac{1}{\sqrt{2}} \right) = \left( 6 - \frac{1}{\sqrt{2}} \right) \text{ km}$$

$$\text{And, } MR = NS = OS = \left( \frac{1}{\sqrt{2}} \right) \text{ km}$$

$$MP = \sqrt{PR^2 + MR^2}$$

$$MP = \sqrt{\left( 6 - \frac{1}{\sqrt{2}} \right)^2 + \left( \frac{1}{\sqrt{2}} \right)^2} = 5.34 \text{ km}$$

So, option 'A' is correct.



Scan for Video solution



### 53. (b)

According to the given conditions, if S gets any portfolio, then U does not want any portfolio so, option C is wrong. R wants either Home or Finance or no portfolio so, options A and D is also wrong. If S gets either Power or Telecom then Q wants other one. All these conditions are satisfied by option B.



Scan for Video solution



### 54. (b)

BEST OF LUCK

$\rightarrow$  BSTFLCK

$\rightarrow$  KCLFTSB

GOOD WISHES  $\rightarrow$  GDWSHS

$\leftarrow$  SHSWDG

ACE THE EXAM  $\rightarrow$  CTHXM

MXHTC



Scan for Video solution



### 55. (d)

Since red light is hung with blue and green light is hung with yellow hence, the number of red lights = number of blue lights the number of green lights = number of yellow lights and the sum of red and green lights = sum of blue and yellow lights.



Scan for Video solution



### 56. (6 to 6)

As given 4 corner points of a tetrahedron forms four planes, Now, if we add a point which increase the count to 5. We know that more than 2 points are required to form a plane. To make maximum number of planes, we require minimum 3 points to form a plane. So that total number of planes will be  ${}^5C_3 = 10$  planes.

$\therefore$  New internal planes =  $10 - 4 = 6$



Scan for Video solution



### 57. (b)

All the students, who are participating have to come at 9:00 am with their parents, but the other students and their parents should come at 10:00 am, therefore option (b) is correct.



Scan for Video solution



### 58. (B)

Since, Eswar does not live on a floor immediately above or immediately below Bhola so, options A, C & D are eliminated. All other conditions are fulfilled in option B is correct.



Scan for Video solution



**59. (a)**

If only statement I is provided then we can confidently say that Riaz is eldest but we cannot determine between Shiv & Som that who is elder. So, statement I is sufficient to determine the eldest of four.

Riaz > Shiv > Som > Ansu.



Scan for Video solution

**60. (d)**

Abhishek > Savar

Anshul > Savar

From these two information, we can decide between Abhishek and Anshul that who is elder.



Scan for Video solution

**61. (a)****Case – 1**

Take any 6 bags and put 3, 3 bags on each side of weighting balance. If both are equal then put rest two bags each on both side, this will indicate heavier bag. So, only two weighing's are required.

**Case – 2**

Take any 6 bags and put 3, 3 bags on each side of weighting balance. If both are not equal then rest two are equal of weight then no need to compare them, now out of three bags which was heavier in 1<sup>st</sup> weighting, put 2 bags on each side, if they are equal then remaining 3<sup>rd</sup> will be heavier or if they are not equal then one will be heavier. So, only two weighing's are required.

So, option 'A' is correct.



Scan for Video solution

**62. (a)**

(a)  $\rightarrow f(x) = x^2 2^{-|x|} \Rightarrow f(-x) = (-x)^2 2^{-|-x|} = x^2 2^{-|x|} = f(x)$   
**→ even**

(b)  $\rightarrow f(x) = x 2^{-|x|} \Rightarrow f(-x) = -x 2^{-|-x|} = -x 2^{-|x|} = -f(x)$   
**→ Not even**

(c)  $\rightarrow f(x) = |x| 2^{-x} \Rightarrow f(-x) = |-x| 2^{-(-x)} = |x| 2^x \neq f(x)$

**→ Not even**

(d)  $\rightarrow f(x) = x 2^{-x} \Rightarrow f(-x) = (-x) 2^{-(-x)} = -x 2^x \neq f(x)$

**→ Not even**

so, as the given graph is an even function, hence option (a) is correct.



Scan for Video solution

**63. (a)**

Conclusion I is wrong as some engineers can be professors.

Conclusion II is also wrong because it is not certain that can some engineers be professors are not.

Conclusion III is correct because No male is professors so also no male can be teacher.


So, option A is correct.



Scan for Video solution

**64. (d)**

Given,



$$n(H) = \text{Total} - n(M \cup P \cup C)$$

$$= 450 - (150 + 120 + 30 + 105) = 450 - 405$$



$$n(H) = 45$$



Scan for Video solution





65. (d)



66. (a)

Statements ‘All students are inquisitive’ and ‘No student is inquisitive’ cannot be true simultaneously.  
Statements 1 and 2 can be shown by single diagram.



67. (b)

Given data:

$$p \cap q = \frac{p^2 + q^2}{pq}, p \cup q = \frac{p^2}{q}$$

$$\$ x \oplus y = 2 \odot 2$$


By observing given data we can write

$$x \cap y = \frac{x^2 + y^2}{xy} \quad \$ 2 \odot 2 = \frac{2^2}{2} = 2$$

Now putting these values in given x, y equation:

$$\frac{x^2 + y^2}{xy} = 2 \Rightarrow x^2 + y^2 - 2xy$$

$$(x - y)^2 = 0 \Rightarrow x = y$$



68. (d)

Given,

$$\Rightarrow p \odot q = p - q$$

$$\Rightarrow p \oplus q = p * q$$

From these equations we can observe


$$\odot = - \quad \$ \oplus = \times$$

Put these values in given equation



$$(9 \odot (6 \oplus 7)) \odot (7 \oplus (6 \odot 5))$$

$$(9 - (6 * 7)) - (7 * (6 - 5))$$

$$9 - 42 - 7 = -40$$




69. (b)



Using Pythagoras theorem:


$$(OQ)^2 = 3^2 + 4^2$$

$$OQ = \sqrt{9 + 16} = 5$$



**70. (c)**


Conclusion I is wrong because only some risk seekers are wealthy conclusion II is also wrong because all the entrepreneurs are risk seekers.



Scan for Video solution


**71. (c)**

Possible cases for all purple (P) are green (G) and All black (B) are green (G) is as shown in figure.



Case I

or



Case II

So, if we take case I, conclusion (ii) is true but conclusion (i) is false.

If we take case II, conclusion (i) is true but conclusion (ii) is false.

So, either conclusion I or II is correct not both.

Scan for Video solution


**72. (c)**

Given data:

% of employees drink coffee = 35%


% of employees drink coffee = 40%

% of employees drink both = 10%




$$100\% - (30\% + 10\% + 25\%)$$

$$= 35\%$$

**73. (b)**

Some football player plays hockey.




Scan for Video solution

**74. (d)**

Statement I is not correct as there is also a possibility of no pathogens are bacteria.

Statement II is not correct as this indicates at least few pathogens are not bacterial

So, option (d) is correct.

**75. (b)**

O, P, Q, R, S, T, U, V, W, X  
 0 1 2 3 4 5 6 7 8 9


TS → 54

ST → 45

SS → 44

SU → 46

So, correct option is (b).

**76. (b)**

As per given information

Sample size =  $n(s) = 100$  students

Students play chess  $n(A) = 300$

Students play foot ball  $n(B) = 600$

Students play both game  $n(A \cap B) = 50$

Students who atleast play one of the game:

$$n(A \cup B) = n(A) + n(B) - n(A \cap B)$$

$$= 300 + 600 - 50$$

$$= 900 - 50 = 850$$

Number of students who play neither of the game

$$n(s) - n(A \cup B) = 1000 - 850 = 150$$



Scan for Video solution



77. (c)

$$E = 10 \Rightarrow 5 \times 2$$

$$J = 20 \Rightarrow 10 \times 2$$

$$O = 30 \Rightarrow 15 \times 2$$

$$T = 40 \Rightarrow 20 \times 2$$

Then,

$$P = 16 \times 2 = 32$$

$$S = 19 \times 2 = 38$$


$$P + E + S + T = 32 + 10 + 38 + 40 = 120$$



Scan for Video solution



78. (b)



From given scenario, the Venn diagram is drawn below. Percentage of people who play at least two games

$$= \frac{25 + 15 + 30 + 5}{300} \times 100 = 25\%$$



Scan for Video solution




79. (c)


Total number of people =  $70 + 10 + 20 + 20 + 40 = 160$

Total administrator = 50

$$\% \text{ of administrators} = \frac{50}{160} \times 100 = 31.25\%$$




80. (c)



From the venn diagram, total students in the clubs =  $60 + 80 + 30 + 38 + 10 + 5 + 2 = 225$

Since 25% of all the students are in clubs, then


$$225 = \frac{25}{100} \times x \Rightarrow x = 900$$



81. (b)

Let us assume  $A = 5$ , then in  $A \times B \times C$ , the possible unit digits are 0 or 5. But according to the condition  $(A \times B \times C) = (B \times G \times E)$  which mean  $B \times G \times E$  should also have 0 or 5 at its unit digit which is only possible when one of the digits (B or G or E) is 5. Which is not possible as each letter should represent unique integer 1 to 9.

⇒ If we consider  $B = 5$  then same case will happen with  $D \times E \times F$  i.e., one of these 3 letters should represent 5 to get unit digit 0 or 5 which is not possible.



82. (b)

$$(A + B + C) + (C + D + E) + (E + F + G) + (G + H + K) = 4 \times 13$$

$$\therefore A + B + C + D + E + F + G + H + K = 1 + 2 + \dots + 9 = 45$$

$$\therefore 45 + C + E + G = 4 \times 13$$

$$C + E + G = 7$$

Only possible digits we can assign to C, E & G are {1, 2, 4}.

If E = 1, then either C = 2 or G = 2 which means D should be 10 or F should be 10 respectively to make the summation 13 which is not possible as number should be among 1 to 9. So, out of (1, 2, 4) only 4 can be the correct option.




Scan for Video solution



83. (c)

Let x students watches all three movies, then from Venn diagram-



Total number of students = 40

$$13 + 16 + 19 - 2x = 40$$

{ ∵ x is counted three times so 2x is subtracted }

$$2x = 48 - 40$$

$$x = 4$$





Scan for Video solution



84. (c)

Using statements (i) and (ii) we can draw following Venn diagram and shaded portion representing that same roses fade quickly (statement III).

So, if statement (i) & (ii) are true, then statement (iii) is also true.



85. (d)

**Given:** Chemistry(C), Physics(P), and Mathematics(M)


$$n(C) = 329, n(P) = 186, n(M) = 295$$

$$n(C \cap P) = 83, n(C \cap M) = 217$$

$$n(P \cap M) = 63, n(P \cup C \cup M) = 500$$


$$n(P \cup C \cup M) = n(P) + n(C) + n(M) - n(P \cap C) - n(C \cap M) - n(M \cap C) + n(P \cap C \cap M)$$

$$500 = 186 + 329 + 295 - 83 - 217 - 63 + n(P \cap C \cap M) \Rightarrow n(P \cap C \cap M) = 53$$



86. (b)

Possible Venn diagram is as shown



Conclusion (i) is false as "No bench is a table" is a possibility.

Conclusion (ii) is true as “Some chair will always be a table” in all possible cases.

Conclusion (iii) is false as “No chair is a table” is a possibility.

Conclusion (iv) is false as “Some benches are not chair” is a possibility.



Scan for Video solution



87. (d)

From both the statements it is not certain that some beds are lamps or some lamps are beds or no bed is a lamp.


So, neither (i) nor (ii) is true.




Scan for Video solution



88. (c)



Case-1



Case-2

As per the given conditions, two cases as shown above, are possible. If P & U now switch seats, then it is clear that T is immediate left of P or P is immediately to the right of Q.




Scan for Video solution



89. (d)

The shaded area represents the students who like to read books or play sports. Hence the number of students



$$= 13 + 12 + 44 + 7 + 15 + 17 = 108$$



Scan for Video solution



90. (c)

If it rains then the field is wet & if the field is not wet then it did not rain, so option B & D are possible. If the field is wet it does not necessarily due to rain so option A is possible but option C is not possible.



Scan for Video solution



91. (a)

$$\text{Total faculty members} = 150$$

$$\text{Faculty members do not have any Facebook or WhatsApp} = 30$$


$$\text{Faculty members have Facebook or WhatsApp} = 150 - 30 = 120$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$120 = 55 + 85 - P(A \cap B)$$

$$P(A \cap B) = 20$$

The above information can be drawn as,



Which means that only Facebook users is 35.

So, option ‘A’ is correct.




Scan for Video solution



**92. (b)**


From the given facts, we can draw Venn diagram as shown below.



So, we can certainly say that some humans are not engineers.



Scan for Video solution

**93. (a)**

$$VF = \sqrt{2^2 + 2^2}$$

$$VF = 2\sqrt{2}$$




Scan for Video solution

**94. (c)****statements:**



No manager is leader

All leaders are executives



but manager can be executive and executive can be manager. So, both the conclusions do not follow.

**95. (d)**

From the Venn diagram it can be analyzed that all film directors are playback singers and some film stars are film directors.


**96. (a)**

From the given conditions we can draw the Venn diagram as shown below. From Venn diagram, we can say that the number of teachers who can teach ES & EM but not TD is 1.





97. (d)

|   |   |   |   |   |
|---|---|---|---|---|
| A | L | R | V | X |
| 3 | 3 | 3 | 3 | 3 |
| E | P | V | Z | B |
| 3 | 3 | 3 | 3 | 3 |
| I | T | Z | D | F |
| 5 | 4 | 4 | 4 | 4 |
| O | Y | E | I | K |




98. (c)

The positions of X, Y, Z, W, P and Q are shown below.




99. (d)

The possible venn diagrams from the statements




It is clear that if all women are entrepreneurs and some women are doctors then some entrepreneurs will definitely be doctors



100. (d)

$$H \rightarrow 15 \quad F \rightarrow 17$$




$$\text{Total number of persons} = 25$$


$$\text{The number of persons playing either Hockey or Football or both} = 5 + 10 + 7 = 22$$

The number of persons playing neither hockey nor football is:

$$25 - 22 = 3$$




101. (0.4896 to 0.04896)




$$P(A/B)$$

$$P(\text{actually HIV}^+ / \text{HIV}^+)$$

$$= \frac{0.1 \times 0.95}{0.1 \times 0.95 + 0.9 \times 0.11} = \frac{0.0950}{0.194} = 0.4896$$



**102.(c)**

Note: A square is a special type of rectangle because it has all the characteristic of a rectangles.

Number of rectangles of size unit  $1 \times 2$

$$= '56ca', '8ac7', '8db5', '7db6' = 4$$

Number of rectangles (square) of size  $1 \times 1$

$$= '5aeb', '6ceb', '7dec', '8acd' = 4$$

Number of rectangles (square) of size  $2 \times 2$

$$= '1\ 2\ 3\ 4', '5\ 6\ 7\ 8' = 2$$


$$\therefore \text{Total number of rectangles} = 4 + 4 + 2 = 10$$



Scan for Video solution

**103.(a)**

The only way for the person to move from R to S & returning to R is.




Person will board the train at times 7am (R), 9am (Q), 11am (P), 12:30pm (S) and reaches the station R after 6:30 hours.



Scan for Video solution



**104.(b)**

When the sheet is folded as mentioned in the question, we act the final shape as below.



$$\text{The area of final shape} = \frac{1}{2} \left(\frac{1}{2}\right)^2 = \frac{1}{8}$$

Scan for Video solution

**105.(d)**

So, the end location of Mr. X from is House is North-West

Scan for Video solution

**106.(b)**

$$\frac{\oplus}{\odot} = 2, \frac{\oplus}{\Delta} = 3$$

$$\frac{\odot}{\Delta} = \frac{3}{2} \quad \& \quad \odot + \Delta = 5$$

$$\therefore \odot = 3, \Delta = 2$$

$$\Delta \times \odot = 10$$

$$\otimes = \frac{10}{\Delta} = \frac{10}{2} = 5$$

$$\therefore (\otimes - \oplus)^2 = (5 - 2 \times 3)^2 = 1$$

Scan for Video solution

**107.(d)**

$$\Delta 2 \oplus 3\Delta ((4 \otimes 2) \nabla 4)$$

$$+ 2 - 3 + ((4 \div 2) \times 4)$$

$$- 1 + (2 \times 4) = 7$$



Scan for Video solution

**108. (c)**

$$Q = 3P = 3 \times 3 = 9$$

$$R = 3Q = 3 \times 9 = 27$$


$$S = 3R = 3 \times 27 = 81$$

$$T = 3S = 3 \times 81 = 243$$

$$\text{So, } Q + S = 9 + 81 = 90.$$



Scan for Video solution

**109. (a)**

The position of h, n & t alphabets are 8, 14 & 20. So, the difference of the numbers of n-h & t-n are 6 & 6. Similarly, the difference x-n is 10 so, the missing element should be 'd' so that n-d = 10.



Scan for Video solution

**110. (d)**

There are 4 codes that can be made by  $\alpha$  and  $\beta$ .

$\alpha\alpha, \alpha\beta, \beta\alpha$  and  $\beta\beta$ .

P is coded as  $\alpha\alpha$  and Q as  $\alpha\beta$  so, R and S can be coded as  $\beta\alpha$  and  $\beta\beta$ .

So, option d is correct.



Scan for Video solution

**111. (c)**

Given sequence following a special pattern.

$$Z - 3 = WV, V - 4 = RQP, P - 5 = KJIH$$

$$\text{Or, } (26), (23, 22), (18, 17, 16), (11, 10, 9, 8)$$

So, option 'C' is correct.



Scan for Video solution

**112. (b)**

$$\begin{array}{ccccccc} I & M & H & O \\ \downarrow +1 & \downarrow +1 & \downarrow +1 & \downarrow +1 \\ J & N & I & P \end{array}$$

$$\begin{array}{ccc} I & D & K \\ \downarrow +1 & \downarrow +1 & \downarrow +1 \\ J & E & L \end{array}$$

$$\begin{array}{ccc} I & D & C \\ \downarrow +1 & \downarrow +1 & \downarrow +1 \\ J & E & D \end{array}$$

**(b) Makes to all in answer key**

Scan for Video solution

**113. (c)**

From the given information, it is concluded that Mita is tallest, as Ganga is taller than Laxmi and Rekha then Rekha must be shorter than Mita. As Laxmi is taller than Sana, then Sana must be shorter than Ganga. Statement (2) and (4) are true. Statement (1) and (3) are invalid so these are not true.

So, option 'C' is correct.



Scan for Video solution

**114. (c)**

Given series is the cube of consecutive prime numbers,

$$7^3, 11^3, 13^3, 17^3 = 343, 1331, 2197, 4913$$

Missing term = 2197

So, option 'C' is correct.



Scan for Video solution

**115. (c)**

The given two sequence AMCF = EQGJ and NKUF = ROYJ, following the same pattern. In this pattern, all letters have difference of 3 letters at their respective places. i.e. A = E, M = Q, C = G and F = J Therefore, DHLP = HLPT

So, option 'C' is correct.



Scan for Video solution

**116. (d)**

Multiple of 3 between 50 to 59 are {51, 54, 57}, all three are non-multiple of 4 & 6.

Non-multiple of 4 between 60 to 69 are {61, 62, 63, 65, 66, 67, 69}, no number is a multiple of 6 and non-multiple of 3.

Non-multiple of 6 between 70 to 79 are {70, 71, 73, 74, 75, 76, 77, 79}, only 76 is the multiple of 6 and non-multiple of 3.



Scan for Video solution

**117. (b)**

In the series, before BC, one alphabet (A) is missing, before FGH, two alphabets (D & E) are missing, before LMNO three alphabets are missing so after LMNO, four alphabets will be missing so next term of the series starts with T and will have five alphabets in that term, so it will be TUVWX.



Scan for Video solution

**118. (b)**

$$\begin{array}{ccccccc} \times 6 & \times 5 & \times 4 & \times 3 & \times 2 & \times 1 & \times 0 \\ 2, & 12, & 60, & 240, & 720, & 1440, & \underline{1440}, & 0 \end{array}$$

So, the missing number is 1440.




Scan for Video solution

**119. (d)**

Surface area of 1 face

$$4 \times 4 = 16$$

$$\text{Surface area of 6 faces} = 16 \times 6 = 96$$



Removing one cubic block from each corner will not change the surface area of the cube. So, the surface area is 96 unit<sup>2</sup>.



Scan for Video solution

**120. (d)**

P always beats Q, Q always beats R, R always beats S but S loses to P only sometimes. From above observation, we can say that P is not the best player among all the players and S loses to P only sometimes so, S is not the worst player.



Scan for Video solution

**121. (b)**

Given series contains prime numbers 13, 23, 43, 53, but 33 is not a prime number. So that odd one is 33. So, option 'B' is correct.



Scan for Video solution

**122. (d)**

Option (a), (b), (c) following the same pattern, but only option (d) following a different pattern.

Hence, odd one is option (d).

Pattern: -2, +3, -2, +3 ...


$$(a) C - 2 = A + 3 = D - 2 = B + 3 = E$$

$$(b) J - 2 = H + 3 = K - 2 = I + 3 = L$$

$$(c) X - 2 = V + 3 = Y - 2 = W + 3 = Z$$


$$(d) O - 2 = M + 3 = P - 2 = N + 3 = Q$$

So, option 'd' is correct.




### 123. (c)


Manu is 2 months older than Sravan, who (Sravan) is 3 months younger than Trideep



Pavan is one-month older than Sravan



So, we can see that Trideep is the Oldest



### 124. (c, d)

$$324 = 18^2, 441 = 21^2, 64 = 8^2$$

So, 97 is not perfect square and only prime number hence least fits the given set.

Also,


$$324 \rightarrow 3+2+4 = 9 = 3^2 = \text{perfect square}$$

$$441 \rightarrow 4+4+1 = 9 = 3^2 = \text{perfect square}$$


$$97 \rightarrow 9 + 7 = 16 = 4^2 = \text{perfect square}$$

$$64 \rightarrow 6 + 4 = 10 = \text{not perfect square}$$


So, option (c) and (d) both are the least fitting in the set.





### 125. (d)



Two and quarter hours back, if in mirror view, the time is 1:30 then in actual clock it is 10:30. So after 2 hour, 15 minutes, the current time will be 12:45.




### 126. (c)



### 127. (c)


Arun lives farther away than Ahmed (5 km) but closer than Susan (7 km). So, the possible distance of Susan is 6.02 km.



### 128. (b)

$\overset{+3}{R} \overset{+3}{O} \overset{+3}{A} \overset{+3}{D} \rightarrow \overset{+3}{U} \overset{+3}{R} \overset{+3}{D} \overset{+3}{G}$

$\overset{+3}{S} \overset{+3}{W} \overset{+3}{A} \overset{+3}{N} \rightarrow \overset{+3}{V} \overset{+3}{Z} \overset{+3}{D} \overset{+3}{Q}$



### 129. (a)

As per given data, we have to cover only height to go from first floor to second floor. So that we have taken care only height of the step. As there is nothing mentioned about the width to be taken, then we can ignore the width of the step.

Hence only statement (I) is alone sufficient and statement (II) is not sufficient in any condition.

$$\text{Total steps} = \frac{9}{\cancel{3}/4} = \frac{9 \times 4}{3} = 12$$

So, option 'A' is correct.



Scan for Video solution



### 130. (c)

Let the weight of a pole =  $x$  kg

As per data given in statement (I):

$$\frac{x}{4} = 5 \Rightarrow x = 20 \text{ kg}$$

As per data given in statement (II):

$$10x = 2x + 160 \Rightarrow x = 20 \text{ kg}$$

From either statement (I) or statement (II), we can calculate the weight of single pole. It means we can identify the weight of 10 poles.

Hence, either statement (I) or statement (II) is sufficient.

So, option 'C' is correct.



Scan for Video solution



### 131. (3 to 3)

The pattern followed

$$5 = \frac{6+4}{2}$$

$$7 = \frac{7+4+2+1}{2}$$

$$8 = \frac{1+9+2+1+2+1}{2}$$

⋮

$$\text{So, } x = \frac{3+3}{2} = 3$$




Scan for Video solution





### 132. (c)

$$(66 \square 6) \rightarrow (66 \blacklozenge 6) = \left( \frac{66-6}{66+6} \right) \rightarrow \left( \frac{66+6}{66-6} \right)$$

$$= \frac{60}{72} \times \frac{72}{60} = 1$$



### 133. (c)



### 134. (c)

$$2, 5, 10, 17, 26, 37, 50, 64$$


$$2, 2+3, 5+5, 10+7, 17+9, 26+11, 37+13, 50+15,$$

$$2, 5, 10, 17, 26, 37, 50, 65$$

Here a next odd integer is added to previous term.

Hence, 64 is wrong term.

So, option 'C' is correct.




### 135. (c)

$$13, 17, 19 \underline{(23)} \text{ (Prime No.)}$$

$$M+6 \rightarrow S \text{ similarly } Q+6 \rightarrow W$$

Hence the next term in the series is 23W



### 136. (b)

Next prime no. is 17 and  $G+6 \rightarrow 13$  so  $K+6 \rightarrow Q$ .  
So, the term in the sequence is 17Q



Scan for Video solution

**137. (16 to 16)**

$$\frac{81}{1.5} = 54, \frac{54}{1.5} = 36, \frac{36}{1.5} = 24, \frac{24}{1.5} = 16$$



Scan for Video solution

**138. (d)**

W, E, K, O  
 $23, 5, 11, 15$   
 $\begin{matrix} \curvearrowleft & \curvearrowright \\ +8, & +6 & +4 \end{matrix}$

I, Q, W, A  
 $9, 17, 23, 1$   
 $\begin{matrix} \curvearrowleft & \curvearrowright \\ +8, & +6 & +4 \end{matrix}$

F, N, T, X  
 $6, 14, 20, 24$   
 $\begin{matrix} \curvearrowleft & \curvearrowright \\ +8, & +6 & +4 \end{matrix}$

N, V, B, D  
 $14, 22, 2, 4$   
 $\begin{matrix} \curvearrowleft & \curvearrowright \\ +8, & +6 & +2 \end{matrix}$



Scan for Video solution

**139. (45 to 45)**

2  $\times 1.5$  3  $\times 2$  6  $\times 2.5$  15  $\times 3$  45  $\boxed{45}$  157.5  $\times 3.5$  630  $\times 4$



Scan for Video solution

**140. (725 to 725)**

$\begin{matrix} \curvearrowleft & \curvearrowright \\ \times 2+11 & \times 2+11 & \times 2+11 & \times 2+11 & \times 2+11 \end{matrix}$   
 $12, \quad 35, \quad 81, \quad 173, \quad 357, \quad \underline{\hspace{2cm}}$

$$357 \times 2 + 11 = 725$$



## CHAPTER

# 3

# SPATIAL APTITUDE

### Transformations of Shapes


#### 1. [MCQ] [GATE-CS-2023:1M]

Looking at the surface of a smooth 3-dimensional object from the outside, which one of the following options is TRUE?

- (a) The surface of the object must be concave everywhere.
- (b) The surface of the object must be convex everywhere.
- (c) The surface of the object may be concave in some places and convex in other places.
- (d) The object can have edges, but no corners.

#### 2. [MCQ] [GATE-ME-2023: 2M]

An opaque pyramid (shown below), with a square base and isosceles faces, is suspended in the path of a parallel beam of light, such that its shadow is cast on a screen oriented perpendicular to the direction of the light beam. The pyramid can be reoriented in any direction within the light beam. Under these conditions, which one of the shadows P, Q, R, and S is NOT possible?



- (a) P
- (b) Q
- (c) R
- (d) S


#### 3. [MCQ] [GATE-CE-2023:1M]

Consider a circle with its centre at the origin (O), as shown. Two operations are allowed on the circle.

Operation 1: Scale independently along the x and y axes.

Operation 2: Rotation in any direction about the origin.

Which figure among the options can be achieved through a combination of these two operations on the given circle?




## 4. [MCQ]

[GATE-ME-2022: 1M]

A block with a trapezoidal cross-section is placed over a block with rectangular cross section as shown above.


Which one of the following is the correct drawing of the view of the 3D object as viewed in the direction indicated by an arrow in the below figure?



## 5. [MCQ]

[GATE-EC-2022:2M]

Consider the following square with the four corners and the center marked as P, Q, R, S and T respectively.



Let X, Y and Z represent the following operations:

X: rotation of the square by 180 degree with respect to the S-Q axis.

Y: rotation of the square by 180 degree with respect to the P-R axis.

Z: rotation of the square by 90 degree clockwise with respect to the axis perpendicular, going into the screen and passing through the point T.

Consider the following three distinct sequences of operation (which are applied in the left to right order).

- (1) XYZZ
- (2) XY
- (3) ZZZZ


Which one of the following statements is correct as per the information provided above?

- (a) The sequence of operations (1) and (2) are equivalent
- (b) The sequence of operations (1) and (3) are equivalent
- (c) The sequence of operations (2) and (3) are equivalent
- (d) The sequence of operations (1), (2) and (3) are equivalent

## 6. [MCQ]

[GATE-CE-2022:1M]


For the picture shown below, which one of the following is the correct picture representing reflection with respect to the mirror shown as the dotted line?






## 7. [MCQ]

[GATE-CS-2021: 1M]

A transparent square sheet shown below is folded along the dotted line. The folded sheet will look like\_\_\_\_\_.




- (a) 
- (b) 
- (c) 
- (d) 

## 8. [MCQ]

[GATE-CE-2021: 1M]

The mirror image of the above text about the X-axis is.



- (a) **LBIANGLER**
- (b) **LBIANGLER**
- (c) **LBIANGLER**
- (d) **EGRNLAIBT**

## 9. [MCQ]

[GATE-CE-2021: 1M]




The mirror image of the below text about the X-axis is.

- (a) **PHΛΓVXΗ**
- (b) **拜ΗΓVXΗ**
- (c) **SIΧΙΑΓΛΥΗΡ**
- (d) **拜SIΧΙΑΓΛΥΗ**


## 10. [MCQ]

[GATE-CS-2017:1M]





Which one of the options best describes the transformation of the 2-dimensional figure P to Q, and then to R, as shown?



- (a) Operation 1: A clockwise rotation by 90° about an axis perpendicular to the plane of the figure  
Operation 2: A reflection along a horizontal line.
- (b) Operation 1: A counterclockwise rotation by 90° about an axis perpendicular to the plane of the figure  
Operation 2: A reflection along a horizontal line.
- (c) Operation 1: A clockwise rotation by 90° about an axis perpendicular to the plane of the figure  
Operation 2: A reflection along a vertical line.
- (d) Operation 1: A counterclockwise rotation by 180° about an axis perpendicular to the plane of the figure  
Operation 2: A reflection along a vertical line.

**Classification and Grouping****11. [MCQ]****[GATE-ME-2022: 1M]**


Which one of the groups given below can be assembled to get the shape that is shown above using each piece only once without overlapping with each other? (rotation and translation operations may be used).

- (a) 
- (b) 
- (c) 
- (d) 

**Patterns in 2D & 3D****12. [MCQ]****[GATE-CE-2023: 1M]**

A line of symmetry is defined as a line that divides a figure into two parts in a way such that each part is a mirror image of the other part about that line.

The figure below consists of 20-unit squares arranged as shown. In addition to the given black squares, upto 5 more may be coloured black. Which one among the following options depicts the minimum number of boxes that must be coloured black to achieve two lines of symmetry? (The figure is representative)




- (a) d
- (b) c, d, i
- (c) c, i
- (d) c, d, i, f, g

**13. [MCQ]****[GATE-CE-2022:2M]**

Consider a cube made by folding a single sheet of paper of appropriate shape. The interior faces of the cube are all blank. However, the exterior faces that are not visible in the above view may not be blank.

Which one of the following represents a possible unfolding of the cube?



**14. [MCQ]**

[GATE-CE-2021: 2M]

Consider two rectangular sheets, Sheet M and Sheet N of dimensions 6 cm x 4 cm each.

Folding operation 1: The sheet is folded into half by joining the short edges of the current shape.

Folding operation 2: The sheet is folded into half by joining the long edges of the current shape.


Folding operation 1 is carried out on Sheet M three times.

Folding operation 2 is carried out on Sheet N three times.

- (a)  $13 : 7$       (b)  $3 : 2$   
 (c)  $7 : 5$       (d)  $5 : 13$


**15. [MCQ]**

[GATE-ME-2021: 2M]



A jigsaw puzzle has 2 pieces. One of the pieces is shown above. Which one of the given options for the missing piece when assembled will form a rectangle? The piece can be moved, rotated or flipped to assemble with the above piece.


- Figure 1 consists of two parts, (a) and (b), each showing a stepped pattern. Part (a) shows a single stepped pattern with a central vertical line. The steps are arranged in a roughly triangular shape. Part (b) shows two stepped patterns side-by-side. The left pattern has a central vertical line and a small horizontal extension on the right. The right pattern has a central vertical line and a small horizontal extension on the left.



### 16. [MCQ]

[GATE-EC-2021:1M]


The least number of squares that must be added so that the line P-Q becomes the line of symmetry is






**17. [MCQ]**

[GATE-EE-2021: 1M]






A transparent square sheet shown above is folded along the dotted line. The folded sheet will look like

- Figure 1 consists of four panels labeled (a), (b), (c), and (d). Each panel contains a solid black rectangle with a dashed border. Inside each rectangle, there is a smaller solid black triangle pointing downwards. The orientation of the triangle varies by panel: (a) is rotated 45 degrees clockwise, (b) is vertical, (c) is rotated 45 degrees counter-clockwise, and (d) is rotated 90 degrees clockwise.






**18. [MCQ]****[GATE-CS-2021: 1M]**

A polygon is convex if, for every pair of points, P and Q belonging to the polygon, the line segment PQ lies completely inside or on the polygon.






Which one of the following is NOT a convex polygon?

**19. [MCQ]****[GATE-CS-2021: 1M]**

A circular sheet of paper is folded along the lines in the directions shown. The paper, after being punched in the final folded state as shown and unfolded in the reverse order of folding, will look like \_\_\_\_\_.

**20. [MCQ]****[GATE-CS-2021: 2M]**

A jigsaw puzzle has 2 pieces. One of the pieces is shown below. Which one of the given options for the missing piece when assembled will form a rectangle? The piece can be moved, rotated or flipped to assemble with the below piece.




**ANSWER KEY**

- |           |         |            |         |
|-----------|---------|------------|---------|
| 1. (c)    | 2. (b)  | 3. (a)     | 4. (a)  |
| 5. (b)    | 6. (a)  | 7. (b)     | 8. (b)  |
| 9. (b)    | 10. (a) | 11. (b, c) | 12. (b) |
| 13. (MTA) | 14. (a) | 15. (a)    | 16. (c) |
| 17. (c)   | 18. (a) | 19. (a)    | 20. (a) |


**SOLUTIONS**
**1. (c)**

If the object has edges, then it surely shall have corners:  
option D wrong

Just seeing the surface of SMOOTH 3D object from outside, we cannot say it has concave or convex faces so, options A & B are also wrong and option C is correct



Scan for Video solution

**2. (b)**

Shadow Q is not possible

Depending upon the angle of incidence of the light, the shadows P, R and S can be obtained.




Scan for Video solution

**3. (a)**


Even after both the operations, the centre of the new shape will also lie on the origin. When operation 1 is performed then we can get an ellipse and when operation 2 is performed then a tilted ellipse with centre at origin can be achieved, hence option A is correct.


**4. (a)**

When this 3D object is viewed in the direction of arrow, then the drawing obtained is.




Scan for Video solution



**5. (b)**





So, output (1) and (3) are equivalent.




6. (a)



7. (b)




8. (b)  
**TRIANGLE**  
**LATERAL** Mirror



9. (b)  
**PHYLAXIS**






10. (a)



11. (b, c)

(b) and (c) both are possible



**12. (b)**

The boxes to be coloured black to achieve two lines of symmetry are c, d and i.




Scan for Video solution

**13. (MTA)**

The black shade should be in the direction of line and the grey surface should be in the direction perpendicular to the line.



One of the possible unfolding of cube is shown below.



No option matches with the unfolding.



Scan for Video solution


**14. (a)**

$$\text{Perimeter} = 2(a + b)$$


$$\text{Sheet M's perimeter} = 2(1.5 + 2) = 7$$

$$\text{Sheet N's perimeter} = 2(6 + 0.5) = 13$$

$$\frac{\text{ratio of sheet N perimeter}}{\text{ratio of sheet M perimeter}} = \frac{13}{7}$$

**15. (a)**

The piece in option A when flipped and rotated, will form a rectangle on assembling with first part.



**16. (c)**

For lower part of the figure to be symmetric about PQ, we have to add 1 square on the left side and 3 squares on the right side as shown in figure.

For upper part, we have to add 2 squares on the left side only as shown in figure.

So, total number of minimum squares needed is

$$= 1 + 3 + 2 = 6$$



**17. (c)**

As the paper is folded with respect to dotted line, it will form two triangles vertically one above other. Which is available in option (c).


So, option 'C' is correct.

**18. (a)**


In polygon as shown in figure below, if line from 1 to 5 is drawn then it lies outside the polygon hence it is not a convex polygon

**19. (a)**


The unfolding of the punched paper is shown below.



From the above diagram it is clear that the unfolded paper will be like in option A.

**20. (a)**

The piece in option A if flipped and rotated by 90° can fit in the piece in question.





- |  |                                    |   |
|--|------------------------------------|---|
| 9. [MCQ]   | [GATE-EE-2021:1M]                  | (c) (i) is grammatically correct and (ii) is ambiguous" |
| The people _____ were at the demonstration             | were from all sections of society. |   |
| (a) whose  | (b) which                          | (d) whom  |
| (c) who  | (d) whom                           |   |
| 10. [MCQ]  | [GATE-CE-2021:1M]                  |   |
| Getting to the top is _____ than staying on top        |                                    |   |
| (a) more easy  | (b) much say                       | (d) easier  |
| (c) easiest  | (d) easier                         |   |
| 11. [MCQ]  | [GATE-CE-2021:2M]                  |   |
| Statement: Either P marries Q or X marries Y           |                                    |   |
| Among the options below, the logical NEGATION of       |                                    |   |
| the above statements.                                  |                                    |   |
| (a) P does not marry Q and X marries Y.                |                                    |   |
| (b) Neither P marries Q nor X marries Y.               |                                    |   |
| (c) X does not marry Y and P marries Q.                |                                    |   |
| (d) P marries Q and X marries Y.                       |                                    |   |
| 12. [MCQ]  | [GATE-CE-2021:1M]                  |   |
| (i) Arun and Aparna are here.                          |                                    |   |
| (ii) Arun and Aparna is here.                          |                                    |   |
| (iii) Arun's families is here.                         |                                    |   |
| (iv) Arun's family is here.                            |                                    |   |
| Which of the above sentence are grammatically          |                                    |   |
| CORRECT?   |                                    |   |
| (a) (i) And (ii)                                       | (b) (ii) And (iv)                  |   |
| (c) (i) And (iv)                                       | (d) (iii) And (iv)                 |   |
| 13. [MCQ]  | [GATE-CS-2021:1M]                  |   |
| Consider the following sentences:                      |                                    |   |
| (i) Everybody in the class is prepared for the exam.   |                                    |   |
| (ii) Babu invited Danish to his home because he        |                                    |   |
| enjoys playing chess.                                  |                                    |   |
| Which of the following is the CORRECT observation      |                                    |   |
| about the above two sentences?                         |                                    |   |
| (a) (i) is grammatically correct and (ii) is           |                                    |   |
| unambiguous  |                                    |   |
| (b) (i) is grammatically incorrect and (ii) is         |                                    |   |
| unambiguous"   |                                    |   |
| 14. [MCQ]  | [GATE-CS-2021:1M]                  |   |
| Gauri said that she can play the keyboard_____         |                                    |   |
| her sister.  |                                    |   |
| (a) as well as   |                                    | (b) as better as  |
| (c) as nicest as                                       |                                    | (d) as worse as   |
| 15. [MCQ]  | [GATE-ME-2020:1M]                  |   |
| While I agree ____his proposal this time, I do not     |                                    |   |
| often agree ____him                                    |                                    |   |
| (a) to, with   |                                    | (b) with, to  |
| (c) with, with   |                                    | (d) to, to  |
| 16. [MCQ]  | [GATE-ME-2020:1M]                  |   |
| Jofra Archer, the England fast bowler, is _____ than   |                                    |   |
| accurate.  |                                    |   |
| (a) more fast  |                                    | (b) faster  |
| (c) less fast  |                                    | (d) more faster   |
| 17. [MCQ]  | [GATE-EC-2020:1M]                  |   |
| He was not only accused of theft _____of               |                                    |   |
| conspiracy.  |                                    |   |
| (a) rather   |                                    | (b) but also  |
| (c) but even   |                                    | (d) rather than   |
| 18. [MCQ]  | [GATE-EC-2020:1M]                  |   |
| The untimely loss of life is a cause of serious global |                                    |   |
| concern as thousands of people get killed _____        |                                    |   |
| accidents every year while many other die _____        |                                    |   |
| diseases like cardio vascular disease, cancer, etc.    |                                    |   |
| (a) in, of   |                                    | (b) from, of  |
| (c) during, from                                       |                                    | (d) from, from  |
| 19. [MCQ]  | [GATE-EE-2020:1M]                  |   |
| People were prohibited _____ their vehicles near       |                                    |   |
| the entrance of the main administrative building.      |                                    |   |
| (a) to park  |                                    |   |
| (b) from parking                                       |                                    |   |
| (c) parking  |                                    |   |
| (d) to have parked                                     |                                    |   |

**20. [MCQ]****[GATE-EE-2020:1M]**

This book, including all its chapters, \_\_\_\_\_ interesting. The students as well as the instructor \_\_\_\_\_ in agreement about it.

- |             |               |
|-------------|---------------|
| (a) is, was | (b) are, are  |
| (c) is, are | (d) were, was |

**21. [MCQ]****[GATE-CE-2020:1M]**

Rescue teams deployed \_\_\_\_\_ disaster hit areas combat \_\_\_\_\_ a lot of difficulties to save the people.

- |                |              |
|----------------|--------------|
| (a) with, at   | (b) in, with |
| (c) with, with | (d) to, to   |

**22. [MCQ]****[GATE-CS-2020:1M]**

Raman is confident of speaking English \_\_\_\_\_ six months as he has been practicing regularly \_\_\_\_\_ the last three weeks.

- |                 |                 |
|-----------------|-----------------|
| (a) during, for | (b) for, since  |
| (c) for, in     | (d) within, for |

**23. [MCQ]****[GATE-ME-2019:1M]**

Once the team of analysts identify the problem, we \_\_\_\_\_ in a better position to comment on the issue.

Which one of the following choices CANNOT fill the given blank?

- |                     |                |
|---------------------|----------------|
| (a) will be         | (b) were to be |
| (c) are going to be | (d) might be   |

**24. [MCQ]****[GATE-ME-2019:1M]**

I permitted him to leave, I wouldn't have had any problem with him being absent, \_\_\_\_\_ I?

- |                   |                    |
|-------------------|--------------------|
| (a) had, wouldn't | (b) have, would    |
| (c) had, would    | (d) have, wouldn't |

**25. [MCQ]****[GATE-EC-2019:1M]**

The boat arrived \_\_\_\_\_ dawn.

- |        |           |
|--------|-----------|
| (a) in | (b) at    |
| (c) on | (d) under |

**26. [MCQ]****[GATE-EC-2019:1M]**

The strategies that the company \_\_\_\_\_ to sell its products \_\_\_\_\_ house-to-house marketing.

- |                    |                     |
|--------------------|---------------------|
| (a) use, includes  | (b) uses, include   |
| (c) used, includes | (d) uses, including |

**27. [MCQ]****[GATE-EE-2019:1M]**

The passengers were angry \_\_\_\_\_ the airline staff about the delay.

- |          |             |
|----------|-------------|
| (a) on   | (b) about   |
| (c) with | (d) towards |

**28. [MCQ]****[GATE-CE-2019:1M]**

They have come a long way in \_\_\_\_\_ trust among the users.

- |              |             |
|--------------|-------------|
| (a) creating | (b) created |
| (c) creation | (d) create  |

**29. [MCQ]****[GATE-CE-2019:1M]**

The CEO's decision to quit was as shocking to the Board as it was to \_\_\_\_\_.

- |        |            |
|--------|------------|
| (a) i  | (b) me     |
| (c) my | (d) myself |

**30. [MCQ]****[GATE-CE-2019:1M]**

Hima Das was \_\_\_\_\_ only Indian athlete to win \_\_\_\_\_ gold for India.

- |               |             |
|---------------|-------------|
| (a) the, many | (b) the, a  |
| (c) an, a     | (d) an, the |

**31. [MCQ]****[GATE-CS-2019:1M]**

The expenditure on the project \_\_\_\_\_ as follows: equipment Rs.20 lakhs, salaries Rs. 12 lakhs, and contingency Rs.3 lakhs.

- |                 |            |
|-----------------|------------|
| (a) break down  | (b) break  |
| (c) breaks down | (d) breaks |

**32. [MCQ]****[GATE-CS-2018:1M]**

"From where are they bringing their books? \_\_\_\_\_ bringing \_\_\_\_\_ books from \_\_\_\_\_. "

The words that best fill the blanks in the above sentence are

- |                           |                           |
|---------------------------|---------------------------|
| (a) their, they're, there | (b) they're, their, there |
| (c) there, their, they're | (d) they're, there, there |

**33. [MCQ]****[GATE-ME-2017:1M]**

The ways in which this game can be played \_\_\_\_\_ potentially infinite.

- |         |               |
|---------|---------------|
| (a) is  | (b) is being  |
| (c) are | (d) are being |

**34. [MCQ]****[GATE-ME-2017:1M]**

He was one of my best \_\_\_\_\_ and I felt his loss \_\_\_\_\_.

- (a) friend, keenly
- (b) friends, keen
- (c) friend, keener
- (d) friends, keenly

**35. [MCQ]****[GATE-CS, EE-2017:1M]**

After Rajendra Chola returned from his voyage to Indonesia, he \_\_\_\_\_ to visit the temple in Thanjavur.

- (a) was wishing
- (b) is wishing
- (c) wished
- (d) had wished

**36. [MCQ]****[GATE-CS, EE-2017:1M]**

Saturn is \_\_\_\_\_ to be seen on a clear night with the naked eye.

- (a) enough bright
- (b) bright enough
- (c) as enough bright
- (d) bright as enough

**37. [MCQ]****[GATE-EC-2017:1M]**

It is \_\_\_\_\_ to read this year's textbook \_\_\_\_\_ the last year's.

- (a) easier, than
- (b) most easy, than
- (c) easier, from
- (d) easiest, from

**38. [MCQ]****[GATE-EC-2017:1M]**

I \_\_\_\_\_ made arrangements had I \_\_\_\_\_ informed earlier.

- (a) could have, been
- (b) would have, being
- (c) had, have
- (d) had been, been

**39. [MCQ]****[GATE-CE-2017:1M]**

The bacteria in milk are destroyed when it \_\_\_\_\_ heated to 80 degree Celsius.

- (a) would be
- (b) will be
- (c) is
- (d) was

**40. [MCQ]****[GATE-CE-2017:1M]**

The event would have been successful if you \_\_\_\_\_ able to come.

- (a) are
- (b) had been
- (c) have been
- (d) would have been

**41. [MCQ]****[GATE-ME, EC-2016:1M]**

Based on the given statements, select the appropriate option with respect to grammar and usage.

Statements

- (i) The height of Mr. X is 6 feet.
- (ii) The height of Mr. Y is 5 feet.
- (a) Mr. X is longer than Mr. Y.
- (b) Mr. X is more elongated than Mr. Y.
- (c) Mr. X is taller than Mr. Y.
- (d) Mr. X is lengthier than Mr. Y.

**42. [MCQ]****[GATE-ME-2016:1M]**

The unruly crowd demanded that the accused be \_\_\_\_\_ without trial.

- (a) hanged
- (b) hanging
- (c) hankering
- (d) hung

**43. [MCQ]****[GATE-ME-2016:1M]**

The volume of a sphere of diameter 1 unit is \_\_\_\_\_ than the volume of a cube of side 1 unit.

- (a) least
- (b) less
- (c) lesser
- (d) low

**44. [MCQ]****[GATE-ME, EC-2016:1M]**

Which of the following is CORRECT with respect to grammar and usage?

Mount Everest is \_\_\_\_\_.

- (a) the highest peak in the world
- (b) highest peak in the world
- (c) one of highest peak in the world
- (d) one of the highest peaks in the world

**45. [MCQ]****[GATE-CS, EE-2016:1M]**

The man who is now Municipal Commissioner worked as \_\_\_\_\_.

- (a) the security guard at a university
- (b) a security guard at the university
- (c) a security guard at university
- (d) the security guard at the university

**46. [MCQ]****[GATE-EC-2016:1M]**

An apple costs Rs. 10. An onion costs Rs. 8.

Select the most suitable sentence with respect to grammar and usage.

- (a) The price of an apple is greater than an onion.
- (b) The price of an apple is more than onion.
- (c) The price of an apple is greater than that of an onion.
- (d) Apples are more costlier than onions.

**47. [MCQ] [GATE-CE-2016:1M]**

If I were you, I \_\_\_\_\_ that laptop. It's much too expensive.

- |                  |                |
|------------------|----------------|
| (a) won't buy    | (b) shan't buy |
| (c) wouldn't buy | (d) would buy  |

**48. [MCQ] [GATE-CE-2016:1M]**

Choose the most appropriate set of words from the options given below to complete the following sentence \_\_\_\_\_ is a will \_\_\_\_\_ is a way.

- |                         |                         |
|-------------------------|-------------------------|
| (a) wear, there, their  | (b) were, their, there  |
| (c) where, there, there | (d) where, their, their |

**49. [MCQ] [GATE-CS, CE-2016:1M]**

Out of the following four sentences, select the most suitable sentence with respect to grammar and usage.

- (a) I will not leave the place until the minister does not meet me.
- (b) I will not leave the place until the minister doesn't meet me.
- (c) I will not leave the place until the minister meet me.
- (d) I will not leave the place until the minister meets me.

**50. [MCQ] [GATE-ME-2015:2M]**

Select the appropriate option in place of underlined part of the sentence.

Increased productivity necessary reflects greater efforts made by the employees.

- (a) Increase in productivity necessary
- (b) Increase productivity is necessary
- (c) Increase in productivity necessarily
- (d) No improvement required

**51. [MCQ] [GATE-ME, EC-2015:2M]**

Ram and Shyam shared a secret and promised to each other that it would remain between them. Ram expressed himself in one of the following ways as

given in the choices below. Identify the correct way as per standard English.

- (a) It would remain between you and me.
- (b) It would remain between I and you.
- (c) It would remain between you and I
- (d) It would remain with me.

**52. [MCQ] [GATE-ME, EC-2015:1M]**

Choose the most appropriate word from the options given below to complete the following sentence.

If the athlete had wanted to come first in the race, he \_\_\_\_\_ several hours every day.

- (a) should practice
- (b) should have practiced
- (c) practiced
- (d) should be practicing

**53. [MCQ] [GATE-ME, EC-2015:2M]**

In the following sentence certain parts are underlined and marked P, Q and R. One of the parts may contain certain error or may not be acceptable in standard written communication. Select the part containing an error. Choose D as your answer if there is no error.

The student corrected all the errors (P) that the instructor marked (Q) on the answer book (R).

- |       |              |
|-------|--------------|
| (a) P | (b) Q        |
| (c) R | (d) No Error |

**54. [MCQ] [GATE-ME, EC-2015:1M]**

What is the adverb for the given word below?

Misogynous

- |                    |                |
|--------------------|----------------|
| (a) Misogynousness | (b) Misogyny   |
| (c) Misogynously   | (d) Misogynous |

**55. [MCQ] [GATE-ME, EC-2015:1M]**

Choose the appropriate word/phrase, out of the four options given below, to complete the following sentence.

Dhoni, as well as the other team members of Indian team, \_\_\_\_\_ present on the occasion

- (a) were
- (b) was
- (c) has
- (d) have

**56. [MCQ] [GATE-EE, CS-2015:1M]**

We \_\_\_\_\_ our friend's birthday and we \_\_\_\_\_ how to make it up to him

- (a) completely forgot – don't just know
- (b) forgot completely - don't just know
- (c) completely forgot - just don't know
- (d) forgot completely - just don't know

**57. [MCQ] [GATE-EE, CS-2015:1M]**

Didn't you buy \_\_\_\_\_ when you went shopping?

- (a) any paper
- (b) much paper
- (c) no paper
- (d) a few paper

**58. [MCQ] [GATE-EE, CS-2015:2M]**

Out of the following four sentences, select the most suitable sentence with respect to grammar and usage

- (a) Since the report lacked needed information, it was of no use to them
- (b) The report was useless to them because there were no needed information in it.
- (c) Since the report did not contain the needed information, it was not real useful to them
- (d) Since the report lacked needed information, it would not had been useful to them

**59. [MCQ] [GATE-EC-2015:2M]**

The following question presents a sentence, part of which is underlined. Beneath the sentence you find four ways of phrasing the underlined part. Following the requirements of the standard written English, select the answer that produces the most effective sentence.

Tuberculosis, together with its effects, ranks one of the leading causes of death in India

- (a) ranks as one of the leading causes of death
- (b) rank as one of the leading causes of death
- (c) has the rank of one of the leading causes of death
- (d) are one of the leading causes of death

**60. [MCQ] [GATE-CS, CE-2015:1M]**

The Tamil version of \_\_\_\_\_ John Abraham-starrer *Madras Cafe* \_\_\_\_\_ cleared by the Censor Board with no cuts last week but the film's

distributors \_\_\_\_\_ no takers among the exhibitors for a release in Tamil Nadu \_\_\_\_\_ this Friday.

- (a) Mr., was, found, on
- (b) a, was, found, at
- (c) the, was, found, on
- (d) a, being, find, at

**61. [MCQ] [GATE-ME, EC-2014:1M]**

'Advice' is \_\_\_\_\_.

- (a) a verb
- (b) a noun
- (c) an adjective
- (d) both a verb and a noun

**62. [MCQ] [GATE-ME, EC-2014:1M]**

Which of the options given below best completes the following sentence?

She will feel much better if she \_\_\_\_\_.

- (a) will get some rest
- (b) gets some rest
- (c) will be getting some rest
- (d) is getting some rest

**63. [MCQ] [GATE-ME, EC-2014:1M]**

Choose the most appropriate word from the options given below to complete the following sentence.

Communication and interpersonal skills are \_\_\_\_\_ important in their own ways.

- (a) each
- (b) both
- (c) all
- (d) either

**64. [MCQ] [GATE-ME, EC-2014:1M]**

Choose the most appropriate phrase from the options given below to complete the following sentence.

The aircraft \_\_\_\_\_ take off as soon as its flight plan was filed.

- (a) is allowed to
- (b) will be allowed to
- (c) was allowed to
- (d) has been allowed to

**65. [MCQ] [GATE-EE, CS-2014:1M]**

Who \_\_\_\_\_ was coming to see us this evening?

- (a) you said
- (b) did you say
- (c) did you say that
- (d) had you said





**88. [MCQ]****[GATE-ME-2022:1M]**

Writing too many things on the \_\_\_\_\_ while teaching could make the students get\_\_\_\_\_.

- (a) bored / board
- (b) board / bored
- (c) board / board
- (d) bored / bored

**89. [MCQ]****[GATE-ME-2022:1M]**

After playing \_\_\_\_\_ hours of tennis, I am feeling \_\_\_\_\_ tired to walk back.

- (a) too / too
- (b) too / two
- (c) two / two
- (d) two / too

**90. [MCQ]****[GATE-EE-2022:1M]**

As you grow older, an injury to your \_\_\_\_\_ may take longer to \_\_\_\_\_.

- (a) heel/heel
- (b) heal/heel
- (c) heal/heal
- (d) heel/heal

**91. [MCQ]****[GATE-EC-2022:1M]**

Mr. X speaks \_\_\_\_\_ Japanese \_\_\_\_\_ Chinese.

- (a) neither / or
- (b) either / nor
- (c) neither / nor
- (d) also / but

**92. [MCQ]****[GATE-CE-2022:1M]**

You should \_\_\_\_\_ when to say \_\_\_\_\_

- (a) no/no
- (b) no/know
- (c) know/know
- (d) know/no

**93. [MCQ]****[GATE-CS-2022:1M]**

The \_\_\_\_\_ is too high for it to be considered \_\_\_\_\_.

- (a) fair/ fare
- (b) fair/ fair
- (c) fare/ fare
- (d) fare/ fair

**94. [MCQ]****[GATE-EC-2021:1M]**

*Nostalgia* is to *anticipation* as \_\_\_\_\_ is to \_\_\_\_\_.

Which one of the following options maintains a similar logical relation in the above sentence?

- (a) present, past
- (b) future, past
- (c) past, future
- (d) future, present

**95. [MCQ]****[GATE-EE-2021:1M]**

Oasis is to sand as island is to \_\_\_\_\_.

Which one of the following options maintains a similar logical relation in the above sentence?

- (a) stone
- (b) land
- (c) water
- (d) mountain

**96. [MCQ]****[GATE-CS-2021:1M]**

\_\_\_\_\_ is to surgery as writer is to \_\_\_\_\_

Which one of the following options maintains a similar logical relation in the above sentence?

- (a) Plan, outline
- (b) Hospital, library
- (c) Doctor, book
- (d) Medicine, grammar

**97. [MCQ]****[GATE-CS-2021:1M]**

Pen : Write :: Knife : \_\_\_\_\_

Which one of the following options maintains a similar logical relation in the above?

- (a) Vegetables
- (b) Sharp
- (c) Cut
- (d) Blunt

**98. [MCQ]****[GATE-ME-2020:1M]**

In one of the greatest innings ever seen in 142 years of Test history, Ben Stokes upped the tempo in a five-and-a-half hour long stay of 219 balls including 11 fours and 8 sixes that saw him finish on a 135 not out as England squared the five-match series.

Based on their connotations in the given passage, which one of the following meanings DOES NOT match?

- (a) upped = increased
- (b) squared = lost
- (c) tempo = enthusiasm
- (d) saw = resulted in

**99. [MCQ]****[GATE-ME-2020:1M]**

Select the word that fits the analogy:

White : Whitening :: Light

- (a) Lightning
- (b) Lightening
- (c) Lighting
- (d) Enlightening

**100. [MCQ]****[GATE-ME-2020:1M]**

The recent measure to improve the output would \_\_\_\_\_ the level of production to our satisfaction.

- (a) increase
- (b) decrease
- (c) speed
- (d) equalize

**101.[MCQ]****[GATE-ME-2020:1M]**

I do not think you know the case well enough to have opinions. Having said that, I agree with your other point. What does the phrase "having said that" mean in the given text?

- (a) as opposed to what I have said
- (b) despite what I have said
- (c) in addition to what I have said
- (d) contrary to what I have said

**102.[MCQ]****[GATE-ME-2020:1M]**

Select the word that fits the analogy:

Build : Building :: Grow : \_\_\_\_\_

- |            |            |
|------------|------------|
| (a) Grown  | (b) Grew   |
| (c) Growth | (d) Growed |

**103.[MCQ]****[GATE-ME-2020:1M]**

He is known for his unscrupulous ways. He always sheds \_\_\_\_\_ tears to deceive people.

- |               |                 |
|---------------|-----------------|
| (a) fox's     | (b) crocodile's |
| (c) crocodile | (d) fox         |

**104.[MCQ]****[GATE-EE-2020:1M]**

Stock markets \_\_\_\_\_ at the news of the coup.

- |             |             |
|-------------|-------------|
| (a) poised  | (b) plunged |
| (c) plugged | (d) probed  |

**105.[MCQ]****[GATE-EE-2020:1M]**

Select the word that fits the analogy:

Do : Undo :: Trust : \_\_\_\_\_

- |              |             |
|--------------|-------------|
| (a) Entrust  | (b) Intrust |
| (c) Distrust | (d) Untrust |

**106.[MCQ]****[GATE-EC-2020:1M]**

Select the word that fits the analogy:

Explicit : Implicit :: Express: \_\_\_\_\_.

- |              |              |
|--------------|--------------|
| (a) Impress  | (b) Repress  |
| (c) Compress | (d) Suppress |

**107.[MCQ]****[GATE-CE-2020:1M]**

Select the word that fits the analogy:

Fuse : Fusion :: Use : \_\_\_\_\_.

- |           |           |
|-----------|-----------|
| (a) Usage | (b) User  |
| (c) Uses  | (d) Usion |

**108.[MCQ]****[GATE-CE-2020:2M]**

It is a common criticism that most of the academicians live in their \_\_\_\_\_ so, they are not aware of the real-life challenges.

- |                   |                  |
|-------------------|------------------|
| (a) homes         | (b) ivory towers |
| (c) glass palaces | (d) big flats    |

**109.[MCQ]****[GATE-CE-2020:1M]**

His hunger for reading is insatiable. He reads indiscriminately. He is most certainly a/an \_\_\_\_\_ reader.

- |               |                |
|---------------|----------------|
| (a) all-round | (b) precocious |
| (c) voracious | (d) wise       |

**110.[MCQ]****[GATE-CE-2020:1M]**

Select the most appropriate word that can replace the underlined word without changing the meaning of the sentence:

Now a days, most children have a tendency to belittle the legitimate concerns of their parents.

- |               |              |
|---------------|--------------|
| (a) disparage | (b) applaud  |
| (c) reduce    | (d) begrudge |

**111.[MCQ]****[GATE-CS-2020:1M]**

His knowledge of the subject was excellent, but his classroom performance was \_\_\_\_\_.

- |                    |                  |
|--------------------|------------------|
| (a) extremely poor | (b) good         |
| (c) desirable      | (d) praiseworthy |

**112.[MCQ]****[GATE-CS-2020:1M]**

Select the word that fits the analogy:

Cook : Cook :: Fly :

- |           |              |
|-----------|--------------|
| (a) Flyer | (b) Flying   |
| (c) Flew  | (d) Flighter |

**113.[MCQ]****[GATE-ME-2019:1M]**

Are there enough seats here? There are \_\_\_\_\_ People here than I expected.

- |           |          |
|-----------|----------|
| (a) many  | (b) most |
| (c) least | (d) more |

**114.[MCQ]****[GATE-ME-2019:1M]**

A final examination is the \_\_\_\_\_ of a series of evaluations that a student has to go through.

- |                 |                  |
|-----------------|------------------|
| (a) culmination | (b) consultation |
| (c) desperation | (d) insinuation  |

**115.[MCQ]****[GATE-ME-2019:1M]**

The minister avoided any mention of the issue of women's reservation in the private sector. He was accused of \_\_\_\_\_ the issue.

- (a) collaring
- (b) skirting
- (c) tying
- (d) belting

**116.[MCQ]****[GATE-ME-2019:1M]**

John Thomas, an \_\_\_\_\_ writer, passed away in 2018.

- (a) Imminent
- (b) Prominent
- (c) Eminent
- (d) Dominant

**117.[MCQ]****[GATE-EE-2019:1M]**

Newspapers are a constant source of delight and recreation for me. The \_\_\_\_\_ trouble is that I read \_\_\_\_\_ many of them.

- (a) even, quite
- (b) even, too
- (c) only, quite
- (d) only, too

**118.[MCQ]****[GATE-EE-2019:1M]**

I am not sure if the bus that has been booked will be able to \_\_\_\_\_ all the students.

- (a) sit
- (b) deteriorate
- (c) fill
- (d) accommodate

**119.[MCQ]****[GATE-CE-2019:1M]**

The lecture was attended by quite \_\_\_\_\_ students, so the hall was not very \_\_\_\_\_.

- (a) a few, quite
- (b) few, quiet
- (c) a few, quiet
- (d) few, quite

**120.[MCQ]****[GATE-CE-2019:1M]**

Daytime temperature in Delhi can \_\_\_\_\_ 40°C.

- (a) get
- (b) stand
- (c) reach
- (d) peak

**121.[MCQ]****[GATE-CE-2019:1M]**

The growth rate of ABC motors in 2017 was the same \_\_\_\_\_ XYZ motors in 2016

- (a) as off
- (b) as those of
- (c) as that off
- (d) as that of

**122.[MCQ]****[GATE-CE-2019:2M]**

"The increasing interest in tribal characters might be a mere coincidence, but the timing is of interest. None of this, though, is to say that the tribal hero has arrived

in Hindi cinema, or that the new crop of characters represents the acceptance of the tribal character in the industry. The films and characters are too few to be described as a pattern."

What does the word 'arrived' mean in the paragraph above?

- (a) reached a terminus
- (b) came to a conclusion
- (c) attained a status
- (d) went to a place

**123.[MCQ]****[GATE-CS-2019:1M]**

The search engine's business model \_\_\_\_\_ around the fulcrum of trust.

- (a) revolves
- (b) plays
- (c) sinks
- (d) bursts

**124.[MCQ]****[GATE-CS-2019:1M]**

A court is to a judge as \_\_\_\_\_ is to a teacher.

- (a) a student
- (b) a punishment
- (c) a syllabus
- (d) a school

**125.[MCQ]****[GATE-ME-2018:1M]**

"The judge's standing in the legal community, though shaken by false allegations of wrongdoing, remained \_\_\_\_\_."

The word that best fills the blank in the above sentence is

- (a) undiminished
- (b) damaged
- (c) illegal
- (d) uncertain

**126.[MCQ]****[GATE-ME-2018:1M]**

"The dress \_\_\_\_\_ her so well that they all immediately \_\_\_\_\_ her on her appearance."

The words that best fill the blanks in the above sentence are

- (a) complemented, complemented
- (b) complimented, complemented
- (c) complimented, complimented
- (d) complemented, complimented

**127.[MCQ]****[GATE-ME-2018:1M]**

"Her \_\_\_\_\_ should not be confused with miserliness; she is ever willing to assist those in need."



The word that best fills the blank in the above sentence is

- (a) meandering
- (b) timely
- (c) consistent
- (d) systematic

**138.[MCQ] [GATE-ME-2017:2M]**

"If you are looking for a history of India, or for an account of the rise and fall of the British Raj, or for the reason of the cleaving of the subcontinent into two mutually antagonistic parts and the effects this mutilation will have in the respective sections, and ultimately on Asia, you will not find it in these pages; for though I have spent a lifetime in the country. I lived too near the seat of events, and was too intimately associated with the actors, to get the perspective needed for the impartial recording of these matters."

Which of the following is closest in meaning to 'cleaving'?

- (a) Deteriorating
- (b) Arguing
- (c) Departing
- (d) Splitting

**139.[MCQ] [GATE-ME-2017:1M]**

If you choose plan P, you will have to \_\_\_\_\_ plan Q, as these two are mutually \_\_\_\_\_.

- (a) forgo, exclusive
- (b) forget, inclusive
- (c) accept, exhaustive
- (d) adopt, intrusive

**140.[MCQ] [GATE-ME-2017:1M]**

As the two speakers became increasingly agitated, the debate became \_\_\_\_\_.

- (a) lukewarm
- (b) poetic
- (c) forgiving
- (d) heated

**141.[MCQ] [GATE-CS, EE-2017:1M]**

Research in the workplace reveals that people work for many reasons\_\_\_\_\_.

- (a) money beside
- (b) beside money
- (c) money besides
- (d) besides money

**142.[MCQ] [GATE-CS, EE-2017:1M]**

Choose the option with words that are not synonyms.

- (a) aversion, dislike
- (b) luminous, radiant
- (c) plunder, loot
- (d) yielding, resistant

**143.[MCQ] [GATE-EC-2017:1M]**

The ninth and the tenth of this month are Monday and Tuesday \_\_\_\_\_.

- (a) figuratively
- (b) retrospectively
- (c) respectively
- (d) rightfully

**144.[MCQ] [GATE-EC-2017:2M]**

"If you are looking for a history of India, or for an account of the rise and fall of the British Raj, or for the reason of the cleaving of the subcontinent into two mutually antagonistic parts and the effects this mutilation will have in the respective sections, and ultimately on Asia, you will not find it in these pages: for though I have spent a lifetime in the country. I lived too near the seat of events, and was too intimately associated with the actors, to get the perspective needed for the impartial recording of these matters".

Here, the word 'antagonistic' is closest in meaning to

- (a) Impartial
- (b) argumentative
- (c) separated
- (d) hostile

**145.[MCQ] [GATE-EC-2017:1M]**

She has a sharp tongue and it can occasionally turn \_\_\_\_\_.

- (a) hurtful
- (b) left
- (c) methodical
- (d) vital

**146.[MCQ] [GATE-CE-2017:1M]**

\_\_\_\_\_ with someone else's email account is now a very serious offence.

- (a) involving
- (b) Assisting
- (c) Tampering
- (d) Incubating

**147.[MCQ] [GATE-CE-2017:1M]**

There was no doubt that their work was thorough

Which of the words below is closest in meaning to the underline word above?

- (a) Pretty
- (b) Complete
- (c) Sloppy
- (d) Haphazard

**148.[MCQ] [GATE-ME, EC-2016:1M]**

After India's cricket world cup victory in 1983, Shrotria who was playing both tennis and cricket till then, decided to concentrate only on cricket. And the rest is history.

What does the underlined phrase mean in this context?

- (a) History will rest in peace
- (b) Rest is recorded in history books
- (c) Rest is well known
- (d) Rest is archaic

**149.[MCQ] [GATE-ME, EC-2016:1M]**

The students \_\_\_\_\_ the teacher on teachers' day for twenty years of dedicated teaching.

- (a) facilitated
- (b) felicitated
- (c) fantasized
- (d) facillitated

**150.[MCQ] [GATE-ME-2016:1M]**

Choose the statement(s) where the underlined word is used correctly:

- (i) A prone is a dried plum.
- (ii) He was lying prone on the floor.
- (iii) People who eat a lot of fat are prone to heart disease.
- (a) (i) and (iii) only
- (b) (iii) only
- (c) (i) and (ii) only
- (d) (ii) and (iii) only

**151.[MCQ] [GATE-ME, EC-2016:1M]**

The policeman asked the victim of a theft, "What did you \_\_\_\_\_?"

- (a) loose
- (b) lose
- (c) loss
- (d) louse

**152.[MCQ] [GATE-ME, EC-2016:1M]**

Despite the new medicine's \_\_\_\_\_ in treating diabetes, it is not \_\_\_\_\_ widely.

- (a) effectiveness --- prescribed
- (b) availability --- used
- (c) prescription --- available
- (d) acceptance --- proscribed

**153.[MCQ] [GATE-EE-2016:1M]**

Identify the correct spelling out of the given options:

- (a) Managable
- (b) Manageable
- (c) Mangaeble
- (d) Managible

**154.[MCQ] [GATE-EE-2016:1M]**

The chairman requested the aggrieved shareholders to \_\_\_\_\_ him.

- (a) bare with
- (b) bore with
- (c) bear with
- (d) bare

**155.[MCQ]**

**[GATE-CS, EE-2016:1M]**

Nobody knows how the Indian cricket team is going to cope with the difficult and seamer-friendly wickets in Australia.

Choose the option which is closest in meaning to the underlined phrase in the above sentence.

- (a) put up with
- (b) put in with
- (c) put down to
- (d) put up against

**156.[MCQ]**

**[GATE-CS, EE-2016:1M]**

Find the odd one in the following group of words.  
mock, deride, praise, jeer

- (a) mock
- (b) deride
- (c) praise
- (d) jeer

**157.[MCQ]**

**[GATE-EC-2016:1M]**

The Buddha said, "Holding on to anger is like grasping a hot coal with the intent of throwing it at someone else; you are the one who gets burnt."

Select the word below which is closest in meaning to the word underlined above.

- (a) Burning
- (b) Igniting
- (c) Clutching
- (d) Flinging

**158.[MCQ]**

**[GATE-CE-2016:1M]**

He turned a deaf ear to my request

What does the underlined phrasal verb mean?

- (a) ignored
- (b) appreciated
- (c) twisted
- (d) returned

**159.[MCQ]**

**[GATE-CS, CE-2016:1M]**

A rewording of something written or spoken is a \_\_\_\_\_.

- (a) paraphrase
- (b) paradox
- (c) paradigm
- (d) paraffin

**160.[MCQ]**

**[GATE-CS, CE-2016:1M]**

Archimedes said, "Give me a lever long enough and a fulcrum on which to place it, and I will move the world."

The sentence above is an example of a \_\_\_\_\_ statement.

- (a) figurative
- (b) collateral
- (c) literal
- (d) figurine

**161.[MCQ]****[GATE-ME-2015:1M]**

Fill in the blank with the correct idiom/phrase.

That boy from the town was a \_\_\_\_\_ in the sleepy village.

- (a) Dog out of herd
- (b) Sheep form the heap
- (c) fish out of water
- (d) bird from the flock

**162.[MCQ]****[GATE-ME-2015:1M]**

Choose the statement where underlined word is used correctly.

- (a) When the teacher eludes to different authors, he is being elusive.
- (b) When the thief keeps eluding the police, he is being elusive.
- (c) Matters that are difficult to understand, identify or remember are allusive.
- (d) Mirages can be allusive, but a better way to express them is illusory

**163.[MCQ]****[GATE-ME-2015:1M]**

Choose the appropriate word/phrase, out of the four option given below, to complete the following sentence:

Apparent lifelessness \_\_\_\_\_ dormant life.

- (a) harbours
- (b) leads to
- (c) supports
- (d) affects

**164.[MCQ]****[GATE-ME, EC-2015:1M]**

Choose the correct verb to fill in the blank below:

Let us \_\_\_\_\_.

- (a) introvert
- (b) alternate
- (c) atheist
- (d) altruist

**165.[MCQ]****[GATE-ME, EC-2015:1M]**

Choose the most suitable one word substitute for the following expression:

Connotation of a road or way

- (a) Pertinacious
- (b) Viaticum
- (c) Clandestine
- (d) Ravenous

**166.[MCQ]****[GATE-ME, EC-2015:1M]**

Choose the word most similar in meaning to the given word:

Awkward

- (a) Inept
- (b) Graceful
- (c) Suitable
- (d) Dreadful

**167.[MCQ]****[GATE-EE, CS-2015:1M]**

Which of the following options is the closest in meaning to the sentence below?

She enjoyed herself immensely at the party.

- (a) She had a terrible time at the party
- (b) Site had a horrible time at the party
- (c) She had a terrific time at the party
- (d) She had a terrifying time at the party

**168.[MCQ]****[GATE-EE, CS-2015:1M]**

Which one of the following combinations is incorrect?

- (a) Acquiescence – Submission
- (b) Wheedle - Roundabout
- (c) Flippancy – Lightness
- (d) Profligate – Extravagant

**169.[MCQ]****[GATE-EE, CS-2015:2M]**

Select the alternative meaning of the underlined part of the sentence.

The chain snatchers took to their heels when the police party arrived.

- (a) took shelter in a thick jungle
- (b) open indiscriminate fire
- (c) took to flight
- (d) unconditionally surrendered

**170.[MCQ]****[GATE-EE, CS-2015:1M]**

Choose the statement where underlined word is used correctly.

- (a) The industrialist had a personnel jet
- (b) I write my experience in my personnel diary
- (c) All personnel are being given the day off
- (d) Being religious is a personnel aspect

**171.[MCQ]****[GATE-EE, CS-2015:1M]**

A generic term that includes various items of clothing such as a skirt, a pair of trousers and a shirt is

- (a) Fabric
- (b) Textile
- (c) Fibre
- (d) Apparel

**172.[MCQ]****[GATE-EC-2015:1M]**

Choose the appropriate word/phrase, out of the four options given below, to complete the following sentence:

Frogs \_\_\_\_\_.

- (a) croak
- (b) roar
- (c) hiss
- (d) patter

**173.[MCQ] [GATE-EC-2015:1M]**

Choose the word most similar in meaning to the given word:

Educe

- (a) Exert
- (b) Educate
- (c) Extract
- (d) Extend

**174.[MCQ] [GATE-EC-2015:1M]**

Choose the most appropriate word from the options given below to complete the following sentence.

The principal presented the chief guest with a \_\_\_\_\_ as token of appreciation.

- (a) momento
- (b) memento
- (c) momentum
- (d) moment

**175.[MCQ] [GATE-CE-2015:1M]**

Choose the most appropriate word from the option given below to complete the following sentence.

The official answered \_\_\_\_\_ that the complaints of the citizen would be looked into.

- (a) respectfully
- (b) respectfully
- (c) reputably
- (d) respectively

**176.[MCQ] [GATE-CE-2015:1M]**

Choose the statement where underlined word is used correctly.

- (a) The minister insured the victims that everything would be all right.
- (b) He ensured that the company will not have to bear any loss
- (c) The actor got himself ensured against any accident.
- (d) The teacher insured students of good results

**177.[MCQ] [GATE-CE-2015:1M]**

Which word is not a synonym for the word vernacular?

- (a) Regional
- (b) Indigenous
- (c) Indigent
- (d) colloquial

**178.[MCQ]**

**[GATE-CE-2015:2M]**

The word similar in meaning to 'dreary' is

- (a) cheerful
- (b) dreamy
- (c) hard
- (d) dismal

**179.[MCQ]**

**[GATE-CS, CE-2015:1M]**

Extreme focus on syllabus and studying for tests has become such a dominant concern of Indian students that they close their minds to anything \_\_\_\_\_ to the requirements of the exam.

- (a) related
- (b) extraneous
- (c) outside
- (d) useful

**180.[MCQ]**

**[GATE-CS, CE-2015:1M]**

Select the pair that best expresses a relationship similar to that expressed in the pair:

Children : Pediatrician

- (a) Adult : Orthopaedist
- (b) Females : Gynaecologist
- (c) Kidney : Nephrologist
- (d) Skin : Dermatologist

**181.[MCQ]**

**[GATE-ME, EC-2014:1M]**

While receiving the award, the scientist said, "I feel vindicated". Which of the following is closest in meaning to the word 'vindicated'?

- (a) Punished
- (b) Substantiated
- (c) Appreciated
- (d) Chastened

**182.[MCQ]**

**[GATE-ME, EC-2014:1M]**

After the discussion, Tom said to me, 'Please revert!'. He expects me to \_\_\_\_\_.

- (a) retract
- (b) get back to him
- (c) move in reverse
- (d) retreat

**183.[MCQ]**

**[GATE-ME, EC-2014:1M]**

Which of the following options is the closest in meaning to the word underlined in the sentence below?

In a democracy, everybody has the freedom to disagree with the government.

- (a) Dissent
- (b) Descent
- (c) Decent
- (d) Decadent



**193.[MCQ]****[GATE-CE-2014:1M]**

A student is required to demonstrate a high level of comprehension of the subject especially in the social sciences.

The word closest in meaning to comprehension is

- (a) Understanding
- (b) Meaning
- (c) Concentration
- (d) Stability

**194.[MCQ]****[GATE-CE-2014:1M]**

Choose the most appropriate word from the option given below to complete the following sentence.

One of his biggest \_\_\_\_\_ was his ability to forgive

- |             |              |
|-------------|--------------|
| (a) Vice    | (b) Virtues  |
| (c) Choices | (d) Strength |

**195.[MCQ]****[GATE-CE-2014:1M]**

Choose the most appropriate word from the options given below to complete the following sentence.

\_\_\_\_\_ is the key to their happiness; they are satisfied with what they have.

- |                  |              |
|------------------|--------------|
| (a) Contentment  | (b) Ambition |
| (c) Perseverance | (d) Hunger   |

**196.[MCQ]****[GATE-CE-2014:1M]**

Which of the following option is the closest in meaning to the sentence below?

As a women, I have no country

- (a) Women have no country.
- (b) Women are not citizens of any country
- (c) Women's solidarity knows no national boundaries
- (d) Women of all countries have equal legal rights

**197.[MCQ]****[GATE-ME, CS-2013:1M]**

Which one of the following options is the closest in meaning to the word given below?

**Nadir**

- (a) Highest
- (b) Lowest
- (c) Medium
- (d) Integration

**198.[MCQ]****[GATE-ME, CS-2013:1M]**

Complete the sentence:

Universalism is to particularism as diffuseness is to \_\_\_\_\_.

- (a) specificity
- (b) neutrality
- (c) generality
- (d) adaptation

**199.[MCQ]****[GATE-EC, EE-2013:1M]**

They were requested not to **quarrel** with others.

Which one of the following options is the closest in meaning to the word **quarrel**?

- (a) Make out
- (b) Call out
- (c) Dig out
- (d) Fall out

**200.[MCQ]****[GATE-EC, EE-2013:1M]**

Complete the sentence:

Dare \_\_\_\_\_ mistakes.

- (a) commit
- (b) to commit
- (c) committed
- (d) committing

**201.[MCQ]****[GATE-EC, EE-2013:1M]**

**Statement:** You can always give me a ring whenever you need.

Which one of the following is the best inference from the above statement?

- (a) Because I have a nice caller tune.
- (b) Because I have a better telephone facility.
- (c) Because a friend in need is a friend indeed.
- (d) Because you need not pay towards the telephone bills when you give me a ring.

**202.[MSQ]****[GATE-CE-2013:1M]**

Which of the following option is the closest in meaning to the word given below:

**Primeval**

- (a) Modern
- (b) Historic
- (c) Primitive
- (d) Antique

**203.[MCQ]****[GATE-CE-2013:1M]**

Friendship no matter how \_\_\_\_\_ it is, has its limitations

- (a) cordial
- (b) intimate
- (c) secret
- (d) pleasant

**204.[MCQ]****[GATE-CE-2013:1M]**

Select the pair that best expresses a relationship similar to that expressed in the pair:

Medicine: Health

- (a) Science: Experiment
- (b) Wealth: Peace
- (c) Education: Knowledge
- (d) Money: Happiness

**205.[MCQ]****[GATE-ME, CE, CS-2012:1M]**

Which one of the following options is the closest in meaning to the word given below?

Mitigate

- |              |             |
|--------------|-------------|
| (a) Diminish | (b) Divulge |
| (c) Dedicate | (d) Denote  |

**206.[MCQ]****[GATE-ME, CE, CS-2012:1M]**

Choose the most appropriate alternative from the options given below to complete the following sentence:

Despite several \_\_\_\_\_ the mission succeeded in its attempt to resolve the conflict

- |              |                 |
|--------------|-----------------|
| (a) attempts | (b) setbacks    |
| (c) meetings | (d) delegations |

**207.[MCQ]****[GATE-EC, EE-2012:1M]**

Which one of the following options is the closest in meaning to the word given below?

**Latitude**

- |                 |                    |
|-----------------|--------------------|
| (a) Eligibility | (b) Freedom        |
| (c) Coercion    | (d) Meticulousness |

**208.[MCQ]****[GATE-EC, EE-2012:1M]**

Choose the most appropriate word from the options given below to complete the following sentence:

**Given the seriousness of the situation that he had to face, his\_\_\_\_\_ was impressive.**

- |              |                  |
|--------------|------------------|
| (a) beggary  | (b) nomenclature |
| (c) jealousy | (d) nonchalance  |

**209.[MCQ]****[GATE-ME, CE, CS-2011:2M]**

Few school curricula include a unit on how to deal with bereavement and grief, and yet all students at some point in their lives suffer from losses through death and parting.

Based on the above passage which topic would not be included in a unit on bereavement?

- (a) How to write a letter of condolence
- (b) What emotional stages are passed through in the healing process
- (c) What the leading causes of death are
- (d) How to give support to a grieving friend

**210.[MCQ]****[GATE-ME, CE, CS-2011:1M]**

Choose the most appropriate word from the options given below to complete the following sentence.

If you are trying to make a strong impression on your audience, you cannot do so by being understated, tentative or \_\_\_\_\_.

- |                   |                 |
|-------------------|-----------------|
| (a) hyperbolic    | (b) restrained  |
| (c) argumentative | (d) indifferent |

**211.[MCQ]****[GATE-ME, CE, CS-2011:1M]**

Which of the following options is the closest in the meaning to the word below:

**Inexplicable**

- |                      |                |
|----------------------|----------------|
| (a) Incomprehensible | (b) Indelible  |
| (c) Inextricable     | (d) Infallible |

**212.[MCQ]****[GATE-ME, CE, CS-2011:1M]**

Choose the word from the options given below that is most nearly opposite in meaning to the given word:

**Amalgamate**

- |             |              |
|-------------|--------------|
| (a) merge   | (b) split    |
| (c) collect | (d) separate |

**213.[MCQ]****[GATE-EC, EE-2011:1M]**

Choose the most appropriate word from the options given below to complete the following sentence:

**Under ethical guidelines recently adopted by the Indian Medical Association, human genes are to be manipulated only to correct diseases for which \_\_\_\_\_ treatments are unsatisfactory.**

- |              |               |
|--------------|---------------|
| (a) similar  | (b) most      |
| (c) uncommon | (d) available |

**214.[MCQ]****[GATE-EC, EE-2011:1M]**

Choose the word from the options given below that is most nearly opposite in meaning to the given word:



**223.[MCQ]****[GATE-EE-2023:2M]**

Which one of the options does NOT describe the passage below or follow from it?

We tend to think of cancer as a ‘modern’ illness because its metaphors are so modern. It is a disease of overproduction, of sudden growth, a growth that is unstoppable, tipped into the abyss of no control. Modern cell biology encourages us to imagine the cell as a molecular machine. Cancer is that machine unable to quench its initial command (to grow) and thus transform into an indestructible, self-propelled automaton. [Adapted from *The Emperor of All Maladies* by Siddhartha Mukherjee]

- (a) It is a reflection of why cancer seems so modern to most of us.
- (b) It tells us that modern cell biology uses and promotes metaphors of machinery.
- (c) Modern cell biology encourages metaphors of machinery, and cancer is often imagined as a machine.
- (d) Modern cell biology never uses figurative language, such as metaphors, to describe or explain anything.

**224.[MCQ]****[GATE-EE-2023:2M]**

Students of all the departments of a college who have successfully completed the registration process are eligible to vote in the upcoming college elections. However, by the time the due date for registration was over, it was found that surprisingly none of the students from the Department of Human Sciences had completed the registration process. Based only on the information provided above, which one of the following sets of statement(s) can be logically inferred with certainty?

- (i) All those students who would not be eligible to vote in the college elections would certainly belong to the Department of Human Sciences.
  - (ii) None of the students from departments other than Human Sciences failed to complete the registration process within the due time.
  - (iii) All the eligible voters would certainly be students who are not from the Department of Human Sciences.
- |                  |                   |
|------------------|-------------------|
| (a) (i) and (ii) | (b) (i) and (iii) |
| (c) only (i)     | (d) only (iii)    |

**225.[MCQ]****[GATE-EE-2023:1M]**

A recent survey shows that 65% of tobacco users were advised to stop consuming tobacco. The survey also shows that 3 out of 10 tobacco users attempted to stop using tobacco. Based only on the information in the above passage, which one of the following options can be logically inferred with certainty?

- (a) A majority of tobacco users who were advised to stop consuming tobacco made an attempt to do so.
- (b) A majority of tobacco users who were advised to stop consuming tobacco did not attempt to do so.
- (c) Approximately 30% of tobacco users successfully stopped consuming tobacco.
- (d) Approximately 65% of tobacco users successfully stopped consuming tobacco.

**226.[MCQ]****[GATE-EC-2023:2M]**

Which one of the following options can be inferred from the given passage alone?

When I was a kid, I was partial to stories about other worlds and interplanetary travel. I used to imagine that I could just gaze off into space and be whisked to another planet.

[Excerpt from *The Truth about Stories* by T. King]

- (a) It is a child’s description of what he or she likes.
- (b) It is an adult’s memory of what he or she liked as a child.
- (c) The child in the passage read stories about interplanetary travel only in parts.
- (d) It teaches us that stories are good for children.

**227.[MCQ]****[GATE-EC-2023:2M]**

Forestland is a planet inhabited by different kinds of creatures. Among other creatures, it is populated by animals all of whom are ferocious. There are also creatures that have claws, and some that do not. All creatures that have claws are ferocious.

Based only on the information provided above, which one of the following options can be logically inferred with *certainty*?

- (a) All creatures with claws are animals.
- (b) Some creatures with claws are non-ferocious.
- (c) Some non-ferocious creatures have claws.
- (d) Some ferocious creatures are creatures with claws.

**228.[MCQ]****[GATE-CS-2023:2M]**

The country of Zombieland is in distress since more than 75% of its working population is suffering from serious health issues. Studies conducted by competent health experts concluded that a complete lack of physical exercise among its working population was one of the leading causes of their health issues. As one of the measures to address the problem, the Government of Zombieland has decided to provide monetary incentives to those who ride bicycles to work.

Based only on the information provided above, which one of the following statements can be logically inferred with certainty?

- All the working population of Zombieland will henceforth ride bicycles to work.
- Riding bicycles will ensure that all of the working population of Zombieland is free of health issues.
- The health experts suggested to the Government of Zombieland to declare riding bicycles as mandatory.
- The Government of Zombieland believes that riding bicycles is a form of physical exercise.

**229.[MCQ]****[GATE-CE-2023:2M]**

Elvesland is a country that has peculiar beliefs and practices. They express almost all their emotions by gifting flowers. For instance, if anyone gifts a white flower to someone, then it is always taken to be declaration of one's love for that person. In a similar manner, the gifting of a yellow flower to someone often means that one is angry with that person.

Based only on the information provided above, which one of the following sets of statements(s) can be logically inferred with certainty?

- In Elvesland, one always declares one's love by gifting a white flower.
- In Elvesland, all emotions are declared by gifting flowers.
- In Elvesland, sometimes one expresses one's anger by gifting a flower that is not yellow.
- In Elvesland, sometimes one expresses one's love by gifting a white flower.

- |                         |                         |
|-------------------------|-------------------------|
| (a) Only (ii)           | (b) (i), (ii) and (iii) |
| (c) (i), (iii) and (iv) | (d) Only (iv)           |

**230.[MCQ]****[GATE-CE-2023:2M]**

Based only on the following passage, which one of the options can be inferred with certainty?

When the congregation sang together, Apenyo would also join, though her little screams were not quite audible because of the group singing. But whenever there was a special number, trouble would begin. Apenyo would try singing along, much to the embarrassment of her mother. After two or three such mortifying Sunday evenings, the mother stopped going to church altogether until Apenyo became older and learnt to behave.

At home too, Apenyo never kept quiet; she hummed or made up silly songs to sing by herself, which annoyed her mother at times but most often made her become pensive. She was by now convinced that her daughter had inherited her love of singing from her father who had died unexpectedly away from home.

[Excerpt from These Hills Called Home by Temsula Ao]

- The mother was embarrassed about her daughter's singing at home.
- The mother's feelings about her daughter's singing at home were only of annoyance.
- The mother was not sure if Apenyo had inherited her love of singing from her father.
- When Apenyo hummed at home, her mother tended to become thoughtful.

**231.[MCQ]****[GATE-CE-2023:2M]**

The James web telescope, recently launched in space, is giving humankind unprecedented access to the depths of time by imaging very old stars formed almost 13 billion years ago. Astrophysicists and cosmologists believe that this odyssey in space may even shed light on the existence of dark matter. Dark matter is supposed to interact only via the gravitational interaction and not through the electromagnetic-, the weak- or the strong-interaction. This may justify the epithet "dark" in dark matter.

Based on the above paragraph, which one of the following statements is FALSE?

- (a) No other telescope has captured images of stars older than those captured by the James web telescope
- (b) People other than astrophysical and cosmologists may also believe in the existence of dark matter.
- (c) The James Web telescope could be of use in the research on dark matter.
- (d) If dark matter was known to interact via the strong interaction, then the epithet “dark” would be justified.

**232.[MCQ]****[GATE-CE-2022:2M]**

Healthy eating is a critical component of healthy aging. When should one start eating healthy? It turns out that it is never too early. For example, babies who start eating healthy in the first year are more likely to have better overall health as they get older.

Which one of the following is the CORRECT logical inference based on the information in the above passage?

- (a) Healthy eating is important for those with good health condition, but not for others
- (b) Eating healthy can be started at any age, earlier the better
- (c) Eating healthy and better overall health are more correlated at a young age, but not elder age
- (d) Healthy eating is more important for adults than kids

**233.[MCQ]****[GATE-CE-2022:2M]**

In the last few years, several new shopping malls were opened in the city. The total number of visitors in the malls is impressive. However, the total revenue generated through sales in the shops in these malls is generally low.

Which one of the following is the CORRECT logical inference based on the information in the above passage?

- (a) Fewer people are visiting the malls but spending more
- (b) More people are visiting the malls but not spending enough
- (c) More people are visiting the malls and spending more
- (d) Fewer people are visiting the malls and not spending enough

**234.[MCQ]****[GATE-ME-2022:2M]**

Fish belonging to species S in the deep sea have skins that are extremely black (ultra-black skin). This helps them not only to avoid predators but also sneakily attack their prey. However, having this extra layer of black pigment results in lower collagen on their skin, making their skin more fragile.

Which one of the following is the CORRECT logical inference based on the information in the above passage?

- (a) Having ultra-black skin is only advantageous to species S
- (b) Species S with lower collagen in their skin are at an advantage because it helps them avoid predators
- (c) Having ultra-black skin has both advantages and disadvantages to species S
- (d) Having ultra-black skin is only disadvantageous to species S but advantageous only to their predators

**235.[MCQ]****[GATE-ME-2022:2M]**

Humans are naturally compassionate and honest. In a study using strategically placed wallets that appear “lost”, it was found that wallets with money are more likely to be returned than wallets without money. Similarly, wallets that had a key and money are more likely to be returned than wallets with the same amount of money alone. This suggests that the primary reason for this behavior is compassion and empathy.

Which one of the following is the CORRECT logical inference based on the information in the above passage?

- (a) Wallets with a key are more likely to be returned because people do not care about money
- (b) Wallets with a key are more likely to be returned because people relate to suffering of others
- (c) Wallets used in experiments are more likely to be returned than wallets that are really lost
- (d) Money is always more important than keys

**236.[MCQ]****[GATE-EE-2022:2M]**

Altruism is the human concern for the wellbeing of others. Altruism has been shown to be motivated more by social bonding, familiarity and identification of belongingness to a group. The notion that altruism may be attributed to empathy or guilt has now been rejected.

Which one of the following is the CORRECT logical inference based on the information in the above passage?

- (a) Humans engage in altruism due to guilt but not empathy
- (b) Humans engage in altruism due to empathy but not guilt
- (c) Humans engage in altruism due to group identification but not empathy
- (d) Humans engage in altruism due to empathy but not familiarity

**237.[MCQ]****[GATE-EC-2022:2M]**

Mosquitoes pose a threat to human health. Controlling mosquitoes using chemicals may have undesired consequences. In Florida, authorities have used genetically modified mosquitoes to control the overall mosquito population. It remains to be seen if this novel approach has unforeseen consequences.

Which one of the following is the correct logical inference based on the information in the above passage?

- (a) Using chemicals to kill mosquitoes is better than using genetically modified mosquitoes because genetic engineering is dangerous
- (b) Using genetically modified mosquitoes is better than using chemicals to kill mosquitoes because they do not have any side effects
- (c) Both using genetically modified mosquitoes and chemicals have undesired consequences and can be dangerous
- (d) Using chemicals to kill mosquitoes may have undesired consequences but it is not clear if using genetically modified mosquitoes has any negative consequence

**238.[MCQ]****[GATE-CS-2022:2M]**

Some people believe that “what gets measured, improves”. Some others believe that “what gets measured, gets gamed”. One possible reason for the difference in the beliefs is the work culture in organizations. In organizations with good work culture, metrics help improve outcomes. However, the same metrics are counterproductive in organizations with poor work culture.

Which one of the following is the CORRECT logical inference based on the information in the above passage?

- (a) Metrics are useful in organizations with poor work culture
- (b) Metrics are useful in organizations with good work culture
- (c) Metrics are always counterproductive in organizations with good work culture
- (d) Metrics are never useful in organizations with good work culture

**239.[MCQ]****[GATE-ME-2021:2M]**

The world is going through the worst pandemic in the past hundred years. The air travel industry is facing a crisis, as the resulting quarantine requirement for travelers led to weak demand.

In relation to the first sentence above, what does the second sentence do?

- (a) Restates an idea from the first sentence.
- (b) Second sentence entirely contradicts the first sentence.
- (c) The two statements are unrelated.
- (d) States an effect of the first sentence.

**240.[MCQ]****[GATE-ME-2021:2M]**

Oxpeckers and rhinos manifest a symbiotic relationship in the wild. The oxpeckers warn the rhinos about approaching poachers, thus possibly saving the lives of the rhinos. Oxpeckers also feed on the parasitic ticks found on rhinos.

In the symbiotic relationship described above, the primary benefits for oxpeckers and rhinos respectively are,

- (a) Oxpeckers get a food source, rhinos have no benefit.
  - (b) Oxpeckers save their habitat from poachers while the rhinos have no benefit.
  - (c) Oxpeckers get a food source, rhinos may be saved from the poachers.
  - (d) Oxpeckers save the lives of poachers, rhinos save their own lives.

241.[MCQ]

[GATE-ME-2021:1M]

"The increased consumption of leafy vegetables in the recent months is a clear indication that the people in the state have begun to lead a healthy lifestyle"

Which of the following can be logically inferred from the information presented in the above statement?

- (a) The people in the state did not consume leafy vegetables earlier.
  - (b) Consumption of leafy vegetables may not be the only indicator of healthy lifestyle.
  - (c) Leading a healthy lifestyle is related to a diet with leafy vegetables.
  - (d) The people in the state have increased awareness of health hazards causing by consumption of junk foods.

**242. [MCQ]**

[GATE-EE-2021:1M]

The importance of sleep is often overlooked by students when they are preparing for exams. Research has consistently shown that sleep deprivation greatly reduces the ability to recall the material learnt. Hence, cutting down on sleep to study longer hours can be counterproductive.

Which one of the following statements is the CORRECT inference from the above passage?

- (a) Sleeping well alone is enough to prepare for an exam. Studying has lesser benefit.
  - (b) Students are efficient and are not wrong in thinking that sleep is a waste of time.
  - (c) If a student is extremely well prepared for an exam, he needs little or no sleep.
  - (d) To do well in an exam, adequate sleep must be part of the preparation.

243. [MCQ]

[GATE-EC-2021:2M]

Computers are ubiquitous. They are used to improve efficiency in almost all fields from agriculture to space exploration. Artificial intelligence (AI) is currently a hot topic. AI enables computers to learn, given enough training data. For humans, sitting in front of a computer for long hours can lead to health issues.

Which of the following can be deduced from the above passage?

- (i) Now a days, computers are present in almost all places.
  - (ii) Computers cannot be used for solving problems in engineering.
  - (iii) For humans, there are both positive and negative effects of using computers.
  - (iv) Artificial intelligence can be done without data.
  - (a) (ii) and (iii)
  - (b) (ii) and (iv)
  - (c) (i), (iii) and (iv)
  - (d) (i) and (iii)

**244. [MCQ]**

[GATE-CE-2021:2M]

Humans have the ability to construct worlds entirely in their minds, which don't exist in the physical world. So far as we know, no other species possesses this ability. This skill is so important that we have different words to refer to its different flavors, such as imagination, invention and innovation.

Based on the above passage, which one of the following is TRUE?

- (a) No species possess the ability to construct worlds in their minds.
  - (b) The terms imagination, invention and innovation refer to unrelated skills.
  - (c) We do not know of any species other than humans who possess the ability to construct mental worlds.
  - (d) Imagination, invention and innovation are unrelated to the ability to construct mental worlds.

245. [MCQ]

[GATE-CE-2021:2M]

The author said, "Musicians rehearse before their concerts. Actors rehearse their roles before the opening of a new play. On the other hand, I find it

strange that many public speakers think they can just walk on to the stage and start speaking. In my opinion, it is no less important for public speakers to rehearse their talks."

Based on the above passage, which one of the following is TRUE?

- (a) The author is of the opinion that rehearsing is important for musicians, actors and public speakers.
- (b) The author is of the opinion that rehearsing is less important for public speakers than for musicians and actors.
- (c) The author is of the opinion that rehearsing is more important only for musicians than public speakers.
- (d) The author is of the opinion that rehearsal is more important for actors than musicians.

**246.[MCQ]****[GATE-CS-2021:2M]**

Some people suggest anti-obesity measures (AOM) such as displaying calorie information in restaurant menus. Such measures sidestep addressing the core problems that cause obesity: poverty and income inequality.

Which one of the following statements summarizes the passage?

- (a) The proposed AOM addresses the core problems that cause obesity.
- (b) If obesity reduces, poverty will naturally reduce, since obesity causes poverty.
- (c) AOM are addressing the core problems and are likely to succeed.
- (d) AOM are addressing the problem superficially.

**247.[MCQ]****[GATE-CS-2021:2M]**

Listening to music during exercise improves exercise performance and reduces discomfort. Scientists researched whether listening to music while studying can help students learn better and the results were inconclusive. Students who needed external stimulation for studying fared worse while students who did not need any external stimulation benefited from music.

Which one of the following statements is the CORRECT inference of the above passage?

- (a) Listening to music has no effect on learning and a positive effect on physical exercise.
- (b) Listening to music has a clear positive effect both on physical exercise and on learning.
- (c) Listening to music has a clear positive effect on physical exercise. Music has a positive effect on learning only in some students.
- (d) Listening to music has a clear positive effect on learning in all students. Music has a positive effect only in some students who exercise.

**248.[MCQ]****[GATE-ME-2020:2M]**

Climate change and resilience deal with two aspects - reduction of sources of non-renewable energy resources and reducing vulnerability of climate change aspects. The terms 'mitigation' and 'adaptation' are used to refer to these aspects, respectively.

Which of the following assertions is best supported by the above information?

- (a) Mitigation deals with consequences of climate change.
- (b) Adaptation deals with causes of climate change.
- (c) Mitigation deals with actions taken to reduce the use of fossil fuels.
- (d) Adaptation deals with actions taken to combat green-house gas emissions.

**249.[MCQ]****[GATE-ME-2020:2M]**

Crowd funding deals with mobilisation of funds for a project from a large number of people, who would be willing to invest smaller amounts through web-based platforms in the project.

Based on the above paragraph, which of the following is correct about crowd funding?

- (a) Funds raised through unwilling contributions on web-based platforms.
- (b) Funds raised through large contributions on web-based platforms.
- (c) Fund raised through coerced contributions on web-based platform.
- (d) Fund raised through voluntary contribution on web-based platform.

**250.[MCQ]****[GATE-EC-2020:2M]**

The global financial crisis in 2008 is considered to be the most serious world-wide financial crisis, which started with the sub-prime lending crisis in USA in 2007. The subprime lending crisis led to the banking crisis in 2008 with the collapse of Lehman Brothers in 2008. The sub-prime lending refers to the provision of loans to those borrowers who may have difficulties in repaying loans, and it arises because of excess liquidity following the East Asian crisis.

Which one of the following sequences shows the correct precedence as per the given passage?

- East Asian crisis → subprime lending crisis → banking crisis → global financial crisis.
- Subprime lending crisis → global financial crisis → banking Crisis → East Asian crisis.
- Banking crisis → subprime lending crisis → global financial Crisis → East Asian crisis.
- Global financial crisis → East Asian crisis → Banking crisis → subprime lending crisis.

**251.[MCQ]****[GATE-EC-2020:1M]**

The Canadian constitution requires that equal importance be given to English and French. Last year, Air Canada lost a lawsuit, and had to pay a six-figure fine to a French-speaking couple after they filed complaints about formal in-flight announcements in English lasting 15 seconds, as opposed to informal 5 second message in French.

The French-speaking couple were upset at \_\_\_\_\_.

- The in-flight announcements being made in English.
- The English announcements being clearer than the French ones.
- The English announcements being longer than the French ones.
- Equal importance being given to English and French.

**252.[MCQ]****[GATE-CE-2020:2M]**

The American psychologist Howard Gardner expounds that human intelligence can be

subcategorised into multiple kinds, in such a way that individuals differ with respect to their relative competence in each kind. Based on this theory, modern educationists insist on prescribing multi-dimensional curriculum and evaluation parameters that enable development and assessment of multiple intelligences.

Which of the following statements can be inferred from the given text?

- Howard Gardner insists that the teaching curriculum and evaluation needs to be multidimensional.
- Howard Gardner wants to develop and assess the theory of multiple intelligences.
- Modern educationists want to develop and assess the theory of multiple intelligences.
- Modern educationists insist that the teaching curriculum and evaluation needs to be multi-dimensional.

**253.[MCQ]****[GATE-CE-2020:2M]**

Nominal interest rate is defined as the amount paid by the borrower to the lender for using the borrowed amount for a specific period of time. Real interest rate calculated on the basis of actual value (inflation-adjusted), is approximately equal to the difference between nominal rate and expected rate of inflation in the economy.

Which of the following assertions is best supported by the above information?

- Under high inflation, real interest rate is low and borrowers get benefited.
- Under low inflation, real interest rate is high and borrowers get benefited.
- Under high inflation, real interest rate is low and lenders get benefited.
- Under low inflation, real interest rate is low and borrowers get benefited.

**254.[MCQ]****[GATE-CE-2020:1M]**

Select the word that fits the analogy:

Partial : Impartial :: Popular

- |                |                |
|----------------|----------------|
| (a) Impopular  | (b) Dispopular |
| (c) Mispopular | (d) Unpopular  |

**255.[MCQ]****[GATE-CS-2020:1M]**

The dawn of the 21st century witnessed the melting glaciers oscillating between giving too much and too little to billions of people who depend on them for fresh water. The UN climate report estimates that without deep cuts to man-made emissions, at least 30% of the northern hemisphere's surface permafrost could melt by the end of the century. Given this situation of imminent global exodus of billions of people displaced by rising seas, nation-states need to rethink their carbon footprint for political concerns, if not for environmental ones.

Which one of the following statements can be inferred from the given passage?

- (a) Nation-states do not have environmental concerns.
- (b) Nation-states are responsible for providing fresh water to billions of people.
- (c) Billions of people are responsible for man-made emissions.
- (d) Billions of people are affected by melting glaciers.

**256.[MCQ]****[GATE-CS-2020:2M]**

Goods and Services Tax (GST) is an indirect tax introduced in India in 2017 that is imposed on the supply of goods and services, and it subsumes all indirect taxes except few. It is a destination-based tax imposed on goods and services used, and it is not imposed at the point of origin from where goods come. GST also has a few components specific to state governments, central government and Union Territories (UTs).

Which one of the following statements can be inferred from the given passage?

- (a) GST is imposed on the production of goods and services.
- (b) GST includes all indirect taxes.
- (c) GST does not have a components specific to UT.
- (d) GST is imposed at the point of usage of goods and services.

**257.[MCQ]****[GATE-ME-2019:2M]**

X is an online media provider. By offering unlimited and exclusive online content at attractive prices for a loyalty membership, X is almost forcing its customers towards its loyalty membership. If its loyalty membership continues to grow at its current rate, within the next eight years more households will be watching X than cable television.

Which one of the following statements can be inferred from the above paragraph?

- (a) Most households that subscribe to X's loyalty membership discontinues watching cable television
- (b) Non-members prefer to watch cable television
- (c) Cable television operators don't subscribe to X's loyalty membership
- (d) The X is canceling accounts of non-members

**258.[MCQ]****[GATE-ME-2019:2M]**

While teaching a creative writing class in India, I was surprised at receiving stories from the students that were all set in distant places: in the American West with cowboys and in Manhattan penthouses with clinking ice cubes. This was, till an eminent Caribbean writer gave the writers in the once-colonised countries the confidence to see the shabby lives around them as worthy of being "told".

The writer of the passage is surprised by the creative writing assignments of his students. Because \_\_\_\_\_.

- (a) Some of the students had written stories set in foreign places
- (b) None of the students had written stories set in India
- (c) None of the students had written about ice cubes and cowboys
- (d) Some of the students had written about ice cubes and cowboys

**259.[MCQ]****[GATE-ME-2019:2M]**

Congo was named by Europeans. Congo's dictator Mobuto later changed the name of the country and the river to Zaire with the objective of Africanising names of persons and spaces. However, the name Zaire was a Portuguese alteration of *Nzadi o Nzere*, a local

African term meaning 'River that swallows Rivers'. Zaire was the Portuguese name for the Congo river in the 16th and 17th centuries.

Which one of the following statements can be inferred from the paragraph above?

- (a) Mobuto was not entirely successful in Africanising the name of his country
- (b) The term *Nzadi o Nzere* was of Portuguese origin
- (c) Mobuto's desire to Africanise names was prevented by the Portuguese
- (d) As a dictator Mobuto ordered the Portuguese to alter the name of the river to Zaire

**260.[MCQ]****[GATE-EE-2019:2M]**

An award-winning study by a group of researchers suggests that men are as prone to buying on impulse as women but women feel more guilty about shopping.

Which one of the following statements can be inferred from the given text?

- (a) Some men and women indulge in buying on impulse
- (b) All men and women indulge in buying on impulse
- (c) Few men and women indulge in buying on impulse
- (d) Many men and women indulge in buying on impulse

**261.[MCQ]****[GATE-EC-2019:2M]**

"Indian history was written by British historians - extremely well documented and researched, but not always impartial. History had to serve its purpose: Everything was made subservient to the glory of the Union Jack. Latter-day Indian scholars presented a contrary picture."

From the text above, we can infer that:

- Indian history written by British historians \_\_\_\_\_
- (a) was well documented and not researched but was always biased
  - (b) was not well documented and researched and was always biased
  - (c) was well documented and researched but was sometimes biased
  - (d) was not well documented and researched and was sometimes biased

**262.[MCQ]****[GATE-CE-2019:2M]**

*The Newspaper* reports that over 500 hectares of tribal land spread across 28 tribal settlements in Mohinitampuram forest division have already been "alienated". A top forest official said. "First the tribals are duped out of their land holdings. Second, the families thus rendered landless are often forced to encroach further into the forests".

On the basis of the information available in the paragraph, \_\_\_\_\_ is/are responsible for duping the tribals.

- (a) Forest officials
- (b) Landless families
- (c) The Newspaper
- (d) It cannot be inferred who

**263.[MCQ]****[GATE-CE-2019:2M]**

"Popular Hindi fiction, despite or perhaps because of its wide reach, often does not appear in our cinema. As ideals that viewers are meant to look up to rather than identify with. Hindi film protagonists usually read books of aspirational value. Textbooks. English books or high value literature."

Which one of the following CANNOT be inferred from the paragraph above?

- (a) Though popular Hindi fiction has wide reach it often does not appear in movies.
- (b) Protagonists in Hindi movies, being ideals for viewers read only books of aspirational value
- (c) Textbooks, English books or high literate have aspirational value but not popular Hindi fiction
- (d) People do not look up to writers of textbooks, English books or high value literature

**264.[MCQ]****[GATE-CE-2019:2M]**

'The new cotton technology', Bollgard-II, with herbicide tolerant traits has developed into a thriving business in India. However, the commercial use of this technology is not legal in India. Notwithstanding that, reports indicate that the herbicide tolerant cotton had been purchased by farmers at an average of Rs 200 more than the control price of ordinary cotton, and planted in 15% of the cotton growing area in the 2017 Kharif season.

Which one of the following statements can be inferred from the given passage?

- (a) Farmers want to access the new technology if India benefits from it
- (b) Farmers want to access the new technology even if it is not legal
- (c) Farmers want to access the new technology for experimental purposes
- (d) Farmers want to access the new technology by paying high price

**265.[MCQ]****[GATE-CS-2019:2M]**

“A recent High Court judgement has sought to dispel the idea of begging as a disease — which leads to its stigmatization and criminalization — and to regard it as a symptom. The underlying disease is the failure of the state to protect citizens who fall through the social security net.”

Which one of the following statements can be inferred from the given passage?

- (a) Beggars are lazy people who beg because they are unwilling to work
- (b) Beggars are created because of the lack of social welfare schemes
- (c) Begging is an offence that has to be dealt with firmly
- (d) Begging has to be banned because it adversely affects the welfare of the state

**266.[MCQ]****[GATE-EC-2018:2M]**

A coastal region with unparalleled beauty is home to many species of animals. It is dotted with coral reefs and unspoilt white sandy beaches. It has remained inaccessible to tourists due to poor connectivity and lack of accommodation. A company has spotted the opportunity and is planning to develop a luxury resort with helicopter service to the nearest major city airport. Environmentalists are upset that this would lead to the region becoming crowded and polluted like any other major beach resorts.

Which one of the following statements can be logically inferred from the information given in the above paragraph?

- (a) The culture and tradition of the local people will be influenced by the tourists.
- (b) The region will become crowded and polluted due to tourism.
- (c) The coral reefs are on the decline and could soon vanish.
- (d) Helicopter connectivity would lead to an increase in tourists coming to the region.

**267.[MCQ]****[GATE-EC-2018:2M]**

The Cricket Board has long recognized John's potential as a leader of the team. However, his on-field temper has always been a matter of concern for them since his junior days. While this aggression has filled stadium with die-hard fans, it has taken a toll on his own batting. Until recently, it appeared that he found it difficult to convert his aggression into big scores. Over the past three seasons though, that picture of John has been replaced by a cerebral, calculative and successful batsman-captain. After many years, it appears that the team has finally found a complete captain.

Which of the following statements can be logically inferred from the above paragraph?

- (i) Even as a junior cricketer, John was considered a good captain.
  - (ii) Finding a complete captain is a challenge.
  - (iii) Fans and the Cricket Board have differing views on what they want in a captain.
  - (iv) Over the past three seasons John has accumulated big scores.
- (a) (i), (ii) and (iii) only    (b) (iii) and (iv) only  
 (c) (ii) and (iv) only    (d) (i), (ii), (iii) and (iv)

**268.[MCQ]****[GATE-ME-2017:2M]**

Two very famous sportsmen Mark and Steve happened to be brothers, and played for country K. Mark teased James, an opponent from country E. “There is no way you are good enough to play for your country.” James replied, “Maybe not, but at least I am the best player in my own family.”

Which one of the following can be inferred from this conversation?

- (a) Mark was known to play better than James
- (b) Steve was known to play better than Mark
- (c) James and Steve were good friends
- (d) James played better than Steve

**269.[MCQ]****[GATE-ME-2017:2M]**

"Here, throughout the early 1820s, Stuart continued to fight his losing battle to allow his sepoys to wear their caste-marks and their own choice of facial hair on parade, being again reprimanded by the commander-in-chief. His retort that 'A stronger instance than this of European prejudice with relation to this country has never come under my observations' had no effect on his superiors."

According to this paragraph, which of the statements below is most accurate?

- (a) Stuart's commander-in-chief was moved by this demonstration of his prejudice.
- (b) The Europeans were accommodating of the sepoys' desire to wear their caste-marks.
- (c) Stuart's 'losing battle' refers to his inability to succeed in enabling sepoys to wear caste-marks.
- (d) The commander-in-chief was exempt from the European prejudice that dictated how the sepoys were to dress.

**270.[MCQ]****[GATE-CS, EE-2017:2M]**

"The hold of the nationalist imagination on our colonial past is such that anything inadequately or improperly nationalist is just not history."

Which of the following statements best reflects the author's opinion?

- (a) Nationalists are highly imaginative.
- (b) History is viewed through the filter of nationalism.
- (c) Our colonial past never happened.
- (d) Nationalism has to be both adequately and properly imagined.

**271.[MCQ]****[GATE-CS, EE-2017:2M]**

"We lived in a culture that denied any merit to literary works, considering them important only when they were handmaidens to something seemingly more urgent - namely ideology. This was a country where all gestures, even the most private, were interpreted in political terms."

The author's belief that ideology is not as important as literature is revealed by the word:

- |               |                 |
|---------------|-----------------|
| (a) 'culture' | (b) 'seemingly' |
| (c) 'urgent'  | (d) 'political' |

**272.[MCQ]****[GATE-EC-2017:2M]**

"If you are looking for a history of India, or for an account of the rise and fall of the British Raj, or for the reason of the cleaving of the subcontinent into two mutually antagonistic parts and the effects this mutilation will have in the respective sections, and ultimately on Asia, you will not find it in these pages: for though I have spent a lifetime in the country. I lived too near the seat of events, and was too intimately associated with the actors, to get the perspective needed for the impartial recording of these matters."

Which of the following statements best reflects the author's opinion?

- (a) An intimate association does not allow for the necessary perspective.
- (b) Matters are recorded with an impartial perspective.
- (c) An intimate association offers an impartial perspective.
- (d) Actors are typically associated with the impartial recording of matters.

**273.[MCQ]****[GATE-CE-2017:2M]**

The old concert hall was demolished because of fears that the foundation would be affected by the construction of the new metro line in the area. Modern technology for underground metro construction tried to mitigate the impact of pressurized air pockets created by the excavation of large amounts of soil. But even with these safeguards, it was feared that the soil below the concert hall would not be stable.

From this, one can infer that

- (a) The foundations of old buildings create pressurized air pockets underground, which are difficult to handle during metro construction.
- (b) Metro construction has to be done carefully considering its impact on the foundations of existing buildings.

- (c) Old buildings in an area form an impossible hurdle to metro construction in that area.
  - (d) Pressurized air can be used to excavate large amounts of soil from underground areas.

274.[MCQ]

[GATE-CE-2017:1M]

Bhaichung was observing the pattern of people entering and leaving a car service centre. There was a single window where customers were being served. He saw that people inevitably came out of the centre in the order that they went in. However, the time they spent inside seemed to vary a lot: some people came out in a matter of minutes while for others it took much longer.

From this, what can one conclude?

- (a) The centre operates on a first-come-first-served basis, but with variable service times, depending on specific customer needs.
  - (b) Customers were served in an arbitrary order, since they took varying amounts of time for service completion in the centre.
  - (c) Since some people came out within a few minutes of entering the centre, the system is likely to operate on a last-come-first-served basis.
  - (d) Entering the centre early ensured that one would have shorter service times and most people attempted to do this.

275.[MCQ]

[GATE-CE-2017:2M]

A map shows the elevations of Darjeeling, Gangtok, Kalimpong, Pelling and Siliguri. Kalimpong is at a lower elevation than Gangtok. Pelling is at a lower elevation than Gangtok. Pelling is at a higher elevation than Siliguri. Darjeeling is at a higher elevation than Gangtok.

Which of the following statements can be inferred from the paragraph above?

- (i) Pelling is at a higher elevation than Kalimpong
  - (ii) Kalimpong is at a lower elevation than Darjeeling
  - (iii) Kalimpong is at a higher elevation than Siliguri
  - (iv) Siliguri is at a lower elevation than Gangtok
  - (a) Only (ii)
  - (b) Only (ii) and (iii)
  - (c) Only (ii) and (iv)
  - (d) Only (iii) and (iv)

**276. [MCQ]**

[GATE-ME, EC-2016:2M]

Social science disciplines were in existence in an amorphous form until the colonial period when they were institutionalized. In varying degrees, they were intended to further the colonial interest. In the time of globalization and the economic rise of postcolonial countries like India, conventional ways of knowledge production have become obsolete.

Which of the following can be logically inferred from the above statements?

- (i) Social science disciplines have become obsolete.
  - (ii) Social science disciplines had a pre-colonial origin.
  - (iii) Social science disciplines always promotes colonialism.
  - (iv) Social science must maintain disciplinary boundaries.
    - (a) (ii) only
    - (b) (i) and (iii) only
    - (c) (ii) and (iv) only
    - (d) (iii) and (iv) only

277.[MCQ]

[GATE-ME-2016:2M]

A smart city integrates all modes of transport, uses clean energy and promotes sustainable use of resources. It also uses technology to ensure safety and security of the city, something which critics argue, will lead to a surveillance state.

Which of the following can be logically inferred from the above paragraph?

- (i) All smart cities encourage the formation of surveillance states.
  - (ii) Surveillance is an integral part of a smart city.
  - (iii) Sustainability and surveillance go hand in hand in a smart city.
  - (iv) There is a perception that smart cities promote surveillance.  
  - (a) (i) and (iv) only
  - (b) (ii) and (iii) only
  - (c) (iv) only
  - (d) (i) only

278.[MCQ]

[GATE-ME, EC-2016:2M]

In a world filled with uncertainty, he was glad to have many good friends. He had always assisted them in times of need and was confident that they would reciprocate. However, the events of the last week proved him wrong.

Which of the following inference(s) is/are logically valid and can be inferred from the above passage?

- (i) His friends were always asking him to help them.
  - (ii) He felt that when in need of help, his friends would let him down.
  - (iii) He was sure that his friends would help him when in need.
  - (iv) His friends did not help him last week.
- |                  |                    |
|------------------|--------------------|
| (a) (i) and (ii) | (b) (iii) and (iv) |
| (c) (iii) only   | (d) (iv) only      |

**279.[MCQ]****[GATE-EE-2016:2M]**

Sourya committee had proposed the establishment of Sourya Institutes of Technology (SITs) in line with Indian Institutes of Technology (IITs) to cater to the technological and industrial needs of a developing country.

Which of the following can be logically inferred from the above sentence?

Based on the proposal,

- (i) In the initial years, SIT students will get degrees from IIT.
  - (ii) SITs will have a distinct national objective.
  - (iii) SIT like institutions can only be established in consultation with IIT.
  - (iv) SITs will serve technological needs of a developing country.
- |                         |
|-------------------------|
| (a) (iii) and (iv) only |
| (b) (i) and (iv) only   |
| (c) (ii) and (iv) only  |
| (d) (ii) and (iii) only |

**280.[MCQ]****[GATE-EE-2016:2M]**

A poll of students appearing for masters in engineering indicated that 60 % of the students believed that mechanical engineering is a profession unsuitable for women. A research study on women with masters or higher degrees in mechanical engineering found that 99 % of such women were successful in their professions.

Which of the following can be logically inferred from the above paragraph?

- (a) Many students have misconceptions regarding various engineering disciplines.
- (b) Men with advanced degrees in mechanical engineering believe women are well suited to be mechanical engineers.
- (c) Mechanical engineering is a profession well suited for women with masters or higher degrees in mechanical engineering.
- (d) The number of women pursuing higher degrees in mechanical engineering is small.

**281.[MCQ]****[GATE-EE-2016:1M]**

R2D2 is a robot. R2D2 can repair aeroplanes. No other robot can repair aeroplanes.

Which of the following can be logically inferred from the above statements?

- (a) R2D2 is a robot which can only repair aeroplanes.
- (b) R2D2 is the only robot which can repair aeroplanes.
- (c) R2D2 is a robot which can repair only aeroplanes.
- (d) Only R2D2 is a robot.

**282.[MCQ]****[GATE-CS, EE-2016:2M]**

Computers were invented for performing only high-end useful computations. However, it is no understatement that they have taken over our world today. The internet, for example, is ubiquitous.

Many believe that the internet itself is an unintended consequence of the original invention. With the advent of mobile computing on our phones, a whole new dimension is now enabled. One is left wondering if all these developments are good or, more importantly, required.

Which of the statement(s) below is/are logically valid and can be inferred from the above paragraph?

- (i) The author believes that computers are not good for us.
  - (ii) Mobile computers and the internet are both intended inventions
- |                       |                          |
|-----------------------|--------------------------|
| (a) (i) only          | (b) (ii) only            |
| (c) both (i) and (ii) | (d) neither (i) nor (ii) |

**283.[MCQ]****[GATE-EC-2016:2M]**

The overwhelming number of people infected with rabies in India has been flagged by the World Health Organization as a source of concern. It is estimated that inoculating 70% of pets and stray dogs against rabies can lead to a significant reduction in the number of people infected with rabies.

Which of the following can be logically inferred from the above sentences?

- The number of people in India infected with rabies is high.
- The number of people in other parts of the world who are infected with rabies is low.
- Rabies can be eradicated in India by vaccinating 70 % of stray dogs.
- Stray dogs are the main source of rabies worldwide

**284.[MCQ]****[GATE-CS, CE-2016:2M]**

Indian currency notes show the denomination indicated in at least seventeen languages. If this is not an indication of the nation's diversity, nothing else is.

Which of the following can be logically inferred from the above sentences?

- India is a country of exactly seventeen languages.
- Linguistic pluralism is the only indicator of a nation's diversity.
- Indian currency notes have sufficient space for all the Indian languages.
- Linguistic pluralism is strong evidence of India's diversity.

**285.[MCQ]****[GATE-CE-2016:2M]**

Today, we consider Ashoka as a great ruler because of the copious evidence he left behind in the form of stone carved edicts. Historians tend to correlate greatness of a king at his time with the availability of evidence today.

Which of the following can be logically inferred from the above sentences?

- Emperors who do not leave significant sculpted evidence are completely forgotten.
- Ashoka produced stone carved edicts to ensure that later historians will respect him.

- Statues of kings are a reminder of their greatness.
- A king's greatness, as we know him today, is interpreted by historians..

**286.[MCQ]****[GATE-ME, EC-2015:2M]**

Ms. X will be in Bagdogra from 01/05/2014 to 20/05/2014 and from 22/05/2014 to 31/05/2014. On the morning of 21/05/2014, she will reach Kochi via Mumbai.

Which one of the statements below is logically valid and can be inferred from the above sentences?

- Ms. X will be in Kochi for one day, only in May.
- Ms. X will be in Kochi for only one day in May
- Ms. X will be only in Kochi for one day in May
- Only Ms. X will be in Kochi for one day in May

**287.[MCQ]****[GATE-ME, EC-2015:2M]**

In the following question, the first and the last sentence of the passage are in order and numbered 1 and 6. The rest of the passage is split into 4 parts and numbered as 2, 3, 4 and 5. These 4 parts are not arranged in proper order. Read the sentences and arrange them in a logical sequence to make a passage and choose the correct sequence from the given options.

- On Diwali, the family rises early in the morning
  - The whole family, including the young and the old enjoy doing this.
  - Children let off fireworks later in the night with their friends.
  - At sunset, the lamps are lit and the family performs various rituals.
  - Father, mother, and children visit relatives and exchange gifts and sweets.
  - Houses look so pretty with lighted lamps all around.
- |             |             |
|-------------|-------------|
| (a) 2,5,3,4 | (b) 5,2,4,3 |
| (c) 3,5,4,2 | (d) 4,5,2,3 |

**288.[MCQ]****[GATE-ME, EC-2015:2M]**

Lamenting the gradual sidelining of the arts in school curricula, a group of prominent artists wrote to the Chief Minister last year, asking him to allocate more funds to support arts education in schools. However, no such increase has been announced in this year's Budget. The artists expressed their deep anguish at their request not being approved, but many of them remain optimistic about funding in the future.

Which of the statement(s) below is/are logically valid and can be inferred from the above statements?

- (i) The artists expected funding for the arts to increase this year.
  - (ii) The Chief Minister was receptive to the idea of increasing funding for the arts.
  - (iii) The Chief Minister is a prominent artist.
  - (iv) Schools are giving less importance to arts education nowadays
- (a) (iii) and (iv)  
 (b) (i) and (iv)  
 (c) (i), (ii) and (iv)  
 (d) (i) and (iii)

**289.[MCQ] [GATE-EE, CS-2015:2M]**

The given statement is followed by some courses of action. Assuming the statement to be true, decide the correct option.

**Statement:**

There has been a significant drop in the water level in the lakes supplying water to the city.

**Course of action:**

- (I) The water supply authority should impose a partial cut in supply to tackle the situation.
  - (II) The government should appeal to all the residents through mass media for minimal use of water.
  - (III) The government should ban the water supply in lower areas.
- (a) Statements I and II follow  
 (b) Statements I and III follow  
 (c) Statements II and III follow  
 (d) All statements follow

**290.[MCQ] [GATE-EC-2015:2M]**

Humpty Dumpty sits on a wall every day while having lunch. The wall sometimes breaks. A person sitting on the wall falls if the wall breaks.

Which one of the statements below is logically valid and can be inferred from the above sentences?

- (a) Humpty Dumpty always falls while having lunch  
 (b) Humpty Dumpty does not fall sometimes while having lunch  
 (c) Humpty Dumpty never falls during dinner  
 (d) When Humpty Dumpty does not sit on the wall, the wall does not break

**291.[MCQ]**
**[GATE-EC-2015:2M]**

Read the following paragraph and choose the correct statement.

Climate change has reduced human security and threatened human well-being. An ignored reality of human progress is that human security largely depends upon environmental security. But on the contrary, human progress seems contradictory to environmental security. To keep up both at the required level is a challenge to be addressed by one and all. One of the ways to curb the climate change may be suitable scientific innovations, while the other may be the Gandhian perspective on small scale progress with focus on sustainability.

- (a) Human progress and security are positively associated with environmental security.  
 (b) Human progress is contradictory to environmental security.  
 (c) Human security is contradictory to environmental security.  
 (d) Human progress depends upon environmental security.

**292.[MCQ]**
**[GATE-CS, CE-2015:2M]**

Alexander turned his attention towards India, since he had conquered Persia

Which one of the statements below is logically valid and can be inferred from the above sentence?

- (a) Alexander would not have turned his attention towards India had he not conquered Persia.  
 (b) Alexander was not ready to rest on his laurels and wanted to march to India.  
 (c) Alexander was completely in control of his army and could command it to move towards India  
 (d) Since Alexander's kingdom extended to Indian borders after the conquest of Persia, he was keen to move further.

**293.[MCQ]**
**[GATE-CS, CE-2015:2M]**

Most experts feel that in spite of possessing all the technical skills required to be a batsman of the highest order, he is unlikely to be so due to lack of requisite temperament. He was guilty of throwing away his wicket several times after working hard to lay a strong

foundation. His critics pointed out that until he addressed this problem, success at the highest level will continue to elude him.

Which of the statement(s) below is/are logically valid and can be inferred from the above passage?



**294.[MCQ] [GATE-ME, EC-2014:1M]**

“India is a country of rich heritage and cultural diversity.”

Which one of the following facts best supports the claim made in the above sentence?

- (a) India is a union of 28 states and 7 union territories.
  - (b) India has a population of over 1.1 billion.
  - (c) India is home to 22 official languages and thousands of dialects.
  - (d) The Indian cricket team draws players from over ten states.

**295.[MCQ] [GATE-ME, EC-2014:2M]**

For submitting tax returns, all resident males with annual income below Rs 10 lakh should fill up Form P and all resident females with income below Rs 8 lakh should fill up Form Q. All people with incomes above Rs 10 lakh should fill up Form R, except nonresidents with income above Rs 15 lakhs, who should fill up Form S. All others should fill Form T. An example of a person who should fill Form T is

(c) A solid angle  $\Omega$  with area  $A = \Omega R^2$  and radius  $R = 0.1\text{ m}$

- (a) A resident male with annual income Rs 9 lakh
  - (b) A resident female with annual income Rs 9 lakh
  - (c) A non-resident male with annual income Rs 16 lakh
  - (d) A non-resident female with annual income Rs 16 lakh

296. [MCQ]

[GATE-EE, CS-2014:2M]

The Palghat Gap (or Palakkad Gap), a region about 30 km wide in the southern part of the Western Ghats in India, is lower than the hilly terrain to its north and south. The exact reasons for the formation of this gap are not clear. It results in the neighbouring regions of Tamil Nadu getting more rainfall from the South West monsoon and the neighbouring regions of Kerala having higher summer temperatures.

What can be inferred from this passage?

- (a) The Palghat gap is caused by high rainfall and high temperatures in southern Tamil Nadu and Kerala
  - (b) The regions in Tamil Nadu and Kerala that are near the Palghat Gap are low-lying
  - (c) The low terrain of the Palghat Gap has a significant impact on weather patterns in neighbouring parts of Tamil Nadu and Kerala
  - (d) Higher summer temperatures result in higher rainfall near the Palghat Gap area

297.[MCQ]

[GATE-EE, CS-2014:2M]

Geneticists say that they are very close to confirming the genetic roots of psychiatric illnesses such as depression and schizophrenia, and consequently, that doctors will be able to eradicate these diseases through early identification and gene therapy.

On which of the following assumptions does the statement above rely?

- (a) Strategies are now available for eliminating psychiatric illnesses
  - (b) Certain psychiatric illnesses have a genetic basis
  - (c) All human diseases can be traced back to genes and how they are expressed
  - (d) In the future, genetics will become the only relevant field for identifying psychiatric illnesses

**298.[MCQ]**

[GATE-EE, CS-2014:2M]

The old city of Koenigsberg, which had a German majority population before World War 2, is now called Kaliningrad. After the events of the war, Kaliningrad is now a Russian territory and has a predominantly Russian population. It is bordered by the Baltic Sea on the north and the countries of Poland

to the south and west and Lithuania to the east respectively. Which of the statements below can be inferred from this passage?

- (a) Kaliningrad was historically Russian in its ethnic make up
- (b) Kaliningrad is a part of Russia despite it not being contiguous with the rest of Russia
- (c) Koenigsberg was renamed Kaliningrad, as that was its original Russian name
- (d) Poland and Lithuania are on the route from Kaliningrad to the rest of Russia

**299.[MCQ] [GATE-EE, CS-2014:2M]**

The number of people diagnosed with dengue fever (contracted from the bite of a mosquito) in north India is twice the number diagnosed last year. Municipal authorities have concluded that measures to control the mosquito population have failed in this region. Which one of the following statements, if true, does not contradict this conclusion?

- (a) A high proportion of the affected population has returned from neighbouring countries where dengue is prevalent
- (b) More cases of dengue are now reported because of an increase in the Municipal Office's administrative efficiency
- (c) Many more cases of dengue are being diagnosed this year since the introduction of a new and effective diagnostic test
- (d) The number of people with malarial fever (also contracted from mosquito bites) has increased this year

**300.[MCQ] [GATE-EE, CS-2014:2M]**

By the beginning of the 20th century, several hypotheses were being proposed, suggesting a paradigm shift in our understanding of the universe. However, the clinching evidence was provided by experimental measurements of the position of a star which was directly behind our sun.

Which of the following inference(s) may be drawn from the above passage?

- (i) Our understanding of the universe changes based on the positions of stars
- (ii) Paradigm shifts usually occur at the beginning of centuries

- (iii) Stars are important objects in the universe
- (iv) Experimental evidence was important in confirming this paradigm shift
- (a) (i), (ii) and (iv)      (b) (iii) only
- (c) (i) and (iv)      (d) (iv) only

**301.[MCQ]**
**[GATE-CE-2014:2M]**

Moving into a world of big data will require us to change our thinking about the merits of exactitude. To apply the conventional mindset of measurement to the digital, connected world of the twenty-first century is to miss a crucial point. As mentioned earlier, the obsession with exactness is an artefact of the information-deprived analog era. When data was sparse, every data point was critical, and thus great care was taken to avoid letting any point bias the analysis. *From "BIG DATA" Viktor Mayer-Schonberger and Kenneth Cukier*

The main point of the paragraph is:

- (a) The twenty-first century is a digital world
- (b) Big data is obsessed with exactness
- (c) Exactitude is not critical in dealing with big data
- (d) Sparse data leads to a bias in the analysis

**302.[MCQ]**
**[GATE-CE-2014:1M]**

Rajan was not happy that Sajan decided to do the project on his own. On observing his unhappiness, Sajan explained to Rajan that he preferred to work independently.

Which one of the statements below is logically valid and can be inferred from the above sentences?

- (a) Rajan has decided to work only in a group.
- (b) Rajan and Sajan were formed into a group against their wishes.
- (c) Sajan had decided to give in to Rajan's request to work with him.
- (d) Rajan had believed that Sajan and he would be working together.

**303.[MCQ]**
**[GATE-ME, CS-2013:2M]**

After several defeats in wars, Robert Bruce went in exile and wanted to commit suicide. Just before committing suicide, he came across a spider attempting tirelessly to have its net. Time and again, the spider failed but that did not deter it to refrain from

making attempts. Such attempts by the spider made Bruce curious. Thus, Bruce started observing the near-impossible goal of the spider to have the net. Ultimately, the spider succeeded in having its net despite several failures. Such act of the spider encouraged Bruce not to commit suicide. And then, Bruce went back again and won many a battle, and the rest is history. Which of the following assertions is best supported by the above information?

- (a) Failure is the pillar of success
- (b) Honesty is the best policy
- (c) Life begins and ends with adventures
- (d) No adversity justifies giving up hope

**304.[MCQ] [GATE-EC, EE-2013:2M]**

**Statement:** There were different streams of freedom movements in colonial India carried out by the moderates, liberals, radicals, socialists, and so on.

Which one of the following is the best inference from the above statement?

- (a) The emergence of nationalism in colonial India led to our Independence.
- (b) Nationalism in India emerged in the context of colonialism.
- (c) Nationalism in India is homogeneous.
- (d) Nationalism in India is heterogeneous.

**305.[MCQ] [GATE-EC, EE-2012:2M]**

One of the legacies of the Roman legions was discipline. In the legions, military law prevailed and discipline was brutal. Discipline on the battlefield kept units obedient, intact and fighting, even when the odds and conditions were against them.

Which one of the following statements best sums up the meaning of the above passage?

- (a) Thorough regimentation was the main reason for the efficiency of the Roman legions even in adverse circumstances
- (b) The legions were treated inhumanly as if the men were animals
- (c) Discipline was the armies' inheritance from their seniors
- (d) The harsh discipline to which the legions were subjected led to the odds and conditions being against them.

**306.[MCQ]**
**[GATE-EC, EE-2011:2M]**

The horse has played a little known but very important role in the field of medicine. Horses were injected with toxins of diseases until their blood built up immunities. Then a serum was made from their blood. Serums to fight with diphtheria and tetanus were developed this way.

It can be inferred from the passage, that horses were

- (a) given immunity to diseases
- (b) generally quite immune to diseases
- (c) given medicines to fight toxins
- (d) given diphtheria and tetanus serums

**307.[MCQ] [GATE-ME, CE, CS-2011:2M]**

Wanted Temporary, Part-time persons for the post of Field Interviewer to conduct personal interviews to collect and collate economic data. Requirements: High School-pass, must be available for Day, Evening and Saturday work. Transportation paid, expenses reimbursed.

Which one of the following is the best inference from the above advertisement?

- (a) Gender-discriminatory
- (b) Xenophobic
- (c) Not designed to make the post attractive
- (d) Not gender-discriminatory

**308.[MCQ] [GATE-ME,CE,EE,EC,CS-2010:2M]**

Modern warfare has changed from large scale clashes of armies to suppression of civilian populations. Chemical agents that do their work silently appear to be suited to such warfare; and regrettably, there exist people in military establishments who think that chemical agents are useful tools for their causes. Which of the following statements best sums up the meaning of the above passage:

- (a) Modern warfare has resulted civil strife
- (b) Chemical agents are useful in modern warfare
- (c) Use of chemical agents in warfare would be undesirable
- (d) People in military establishments like to use chemical agents in war




**ANSWER KEY**

- |          |          |          |          |
|----------|----------|----------|----------|
| 1. (a)   | 2. (d)   | 3. (a)   | 4. (c)   |
| 5. (c)   | 6. (a)   | 7. (d)   | 8. (b)   |
| 9. (c)   | 10. (d)  | 11. (b)  | 12. (c)  |
| 13. (c)  | 14. (a)  | 15. (a)  | 16. (a)  |
| 17. (b)  | 18. (a)  | 19. (b)  | 20. (c)  |
| 21. (b)  | 22. (d)  | 23. (b)  | 24. (c)  |
| 25. (b)  | 26. (b)  | 27. (c)  | 28. (a)  |
| 29. (b)  | 30. (b)  | 31. (c)  | 32. (b)  |
| 33. (c)  | 34. (d)  | 35. (c)  | 36. (b)  |
| 37. (a)  | 38. (a)  | 39. (c)  | 40. (b)  |
| 41. (c)  | 42. (a)  | 43. (b)  | 44. (a)  |
| 45. (b)  | 46. (c)  | 47. (c)  | 48. (c)  |
| 49. (d)  | 50. (c)  | 51. (a)  | 52. (b)  |
| 53. (b)  | 54. (c)  | 55. (b)  | 56. (c)  |
| 57. (a)  | 58. (a)  | 59. (a)  | 60. (c)  |
| 61. (b)  | 62. (b)  | 63. (b)  | 64. (c)  |
| 65. (b)  | 66. (d)  | 67. (c)  | 68. (d)  |
| 69. (c)  | 70. (a)  | 71. (d)  | 72. (b)  |
| 73. (d)  | 74. (a)  | 75. (b)  | 76. (a)  |
| 77. (c)  | 78. (a)  | 79. (b)  | 80. (a)  |
| 81. (c)  | 82. (a)  | 83. (c)  | 84. (d)  |
| 85. (b)  | 86. (b)  | 87. (d)  | 88. (b)  |
| 89. (d)  | 90. (d)  | 91. (c)  | 92. (d)  |
| 93. (d)  | 94. (c)  | 95. (c)  | 96. (c)  |
| 97. (c)  | 98. (b)  | 99. (b)  | 100. (a) |
| 101. (b) | 102. (c) | 103. (c) | 104. (b) |
| 105. (c) | 106. (b) | 107. (a) | 108. (b) |
| 109. (c) | 110. (a) | 111. (a) | 112. (a) |
| 113. (d) | 114. (a) | 115. (b) | 116. (c) |
| 117. (d) | 118. (d) | 119. (c) | 120. (c) |
| 121. (d) | 122. (c) | 123. (a) | 124. (d) |
| 125. (a) | 126. (d) | 127. (c) | 128. (a) |

- |          |          |          |          |
|----------|----------|----------|----------|
| 129. (d) | 130. (a) | 131. (a) | 132. (b) |
| 133. (a) | 134. (a) | 135. (a) | 136. (a) |
| 137. (a) | 138. (d) | 139. (a) | 140. (d) |
| 141. (d) | 142. (d) | 143. (c) | 144. (d) |
| 145. (a) | 146. (c) | 147. (b) | 148. (c) |
| 149. (b) | 150. (d) | 151. (b) | 152. (a) |
| 153. (b) | 154. (c) | 155. (a) | 156. (c) |
| 157. (c) | 158. (a) | 159. (a) | 160. (a) |
| 161. (c) | 162. (b) | 163. (a) | 164. (b) |
| 165. (b) | 166. (a) | 167. (c) | 168. (b) |
| 169. (c) | 170. (c) | 171. (d) | 172. (a) |
| 173. (c) | 174. (b) | 175. (b) | 176. (b) |
| 177. (c) | 178. (d) | 179. (b) | 180. (b) |
| 181. (b) | 182. (b) | 183. (a) | 184. (b) |
| 185. (c) | 186. (b) | 187. (b) | 188. (c) |
| 189. (c) | 190. (a) | 191. (a) | 192. (c) |
| 193. (a) | 194. (b) | 195. (a) | 196. (c) |
| 197. (b) | 198. (a) | 199. (d) | 200. (b) |
| 201. (c) | 202. (c) | 203. (b) | 204. (c) |
| 205. (a) | 206. (b) | 207. (b) | 208. (d) |
| 209. (c) | 210. (b) | 211. (a) | 212. (b) |
| 213. (d) | 214. (b) | 215. (c) | 216. (a) |
| 217. (b) | 218. (b) | 219. (a) | 220. (d) |
| 221. (c) | 222. (d) | 223. (d) | 224. (d) |
| 225. (b) | 226. (b) | 227. (d) | 228. (d) |
| 229. (d) | 230. (d) | 231. (d) | 232. (b) |
| 233. (b) | 234. (c) | 235. (b) | 236. (c) |
| 237. (d) | 238. (b) | 239. (d) | 240. (c) |
| 241. (c) | 242. (d) | 243. (d) | 244. (c) |
| 245. (a) | 246. (d) | 247. (c) | 248. (c) |
| 249. (d) | 250. (a) | 251. (c) | 252. (d) |
| 253. (a) | 254. (d) | 255. (d) | 256. (d) |
| 257. (a) | 258. (b) | 259. (a) | 260. (a) |
| 261. (c) | 262. (d) | 263. (b) | 264. (b) |
| 265. (b) | 266. (d) | 267. (c) | 268. (b) |
| 269. (c) | 270. (b) | 271. (b) | 272. (a) |
| 273. (b) | 274. (a) | 275. (c) | 276. (a) |
| 277. (c) | 278. (b) | 279. (c) | 280. (c) |

- |                 |                 |                 |                 |
|-----------------|-----------------|-----------------|-----------------|
| <b>281.</b> (b) | <b>282.</b> (d) | <b>283.</b> (a) | <b>284.</b> (d) |
| <b>285.</b> (d) | <b>286.</b> (b) | <b>287.</b> (b) | <b>288.</b> (b) |
| <b>289.</b> (a) | <b>290.</b> (b) | <b>291.</b> (b) | <b>292.</b> (a) |
| <b>293.</b> (b) | <b>294.</b> (c) | <b>295.</b> (b) | <b>296.</b> (c) |
| <b>297.</b> (b) | <b>298.</b> (b) | <b>299.</b> (d) | <b>300.</b> (d) |
| <b>301.</b> (c) | <b>302.</b> (d) | <b>303.</b> (d) | <b>304.</b> (d) |
| <b>305.</b> (a) | <b>306.</b> (b) | <b>307.</b> (d) | <b>308.</b> (c) |

## SOLUTIONS

**1. (a)**

He did not manage to fix the car himself, so he got it fixed in the garage.



Scan for Video solution



**2. (d)**

Direct Speech - Indirect Speech

Present Continuous Tense - Past Continuous Tense

∴ Indirect Speech of the given sentence will be,

Rafi told Mary that he **was thinking** of watching a film that weekend.

∴ Option (d) is correct.



Scan for Video solution



**3. (a)**

When it goes from one side to other, we use the word 'across'.



Scan for Video solution



**4. (c)**

As it is tentatively, thus may past 'might' will be used.



Scan for Video solution



**5. (c)**

'couldn't help laughing' is used when a person finds situation very funny.



Scan for Video solution



**6. (a)**

"The number" is followed by a singular verb while "A number" is followed by plural verb hence sentences I & II are correct.



Scan for Video solution



**7. (d)**

Hardly or barely comes before the main verb of a sentence. But when there is a modal or auxiliary verb, hardly/barely usually comes after it.



Scan for Video solution



8. (b)

**Wake:** emerge or cause to emerge from sleep. Its second and third form is woke and woken respectively.

So, statements (i) and (iii) are correct.



Scan for Video solution



9. (c)

Whose is used for belongs to someone,  
 Which is a relative cause  
 Who is used for a person/object  
 Whom is a preposition  
 ∴ Option (c) is correct.



Scan for Video solution



10. (d)

The use of ‘than’ suggest that it should be followed by comparative adjective degree. The comparative adjective degree of easy is ‘easier’



Scan for Video solution



11. (b)

Negation means opposite,  
 Opposite of either is neither.  
 Opposite of or is nor.  
 Hence, “Neither P marries Q nor X marries Y.”



Scan for Video solution



12. (c)

In statement (ii) “is” is incorrect.  
 In statement (iii) “families” is incorrect.  
 So (i) and (iv) are correct.



Scan for Video solution



13. (c)

1<sup>st</sup> sentence is grammatically correct as everybody or everyone here is considered a single group so singular form of verb ‘is’ is used.

Second sentence is ambiguous because it is not clear that who among the two, enjoy to play chess.



Scan for Video solution



14. (a)

The correct option is ‘as well as’ because it is in first degree. Option B and D are in comparative degree so should be followed by ‘than’.



Scan for Video solution



15. (a)

While I agree to his proposal this time, I do not often agree with him.

‘Agree with’ is used for a person, and the ‘agree to’ is used for a non-person.



Scan for Video solution



16. (a)

Jofra Archer, the England fast bowler, is more fast than accurate.

In the sentence, we are comparing the two different qualities of the bowler: speed & accuracy. For such comparison, we use ‘more + positive degree adjective’.



Scan for Video solution



17. (b)

The construction **not only ... but (also)** is called a co relative conjunction. It is used to present two related pieces of information. Both pieces of information are being presented as surprising or unexpected, with the second one being even more surprising than the first.



Scan for Video solution



18. (a)

**'In'** is used to indicate opinion, belief, feeling, event etc and **'of'** is used if someone dies of a specific disease or some reasons which is well-known. **'From'** is used if something leads to death.



Scan for Video solution



19. (b)

If something is prohibited then it will be from someone,

So that from parking is the suitable word in this sentence.

∴ Option (b) is correct



Scan for Video solution



20. (c)

This book is a singular noun and the students as well as the instructor, it means two person so that it is a plural noun, then in 1<sup>st</sup> blank space 'is' and in 2<sup>nd</sup> 'are' will be used.

∴ Option (c) is correct.



Scan for Video solution



21. (b)

'in' is used to indicate location or position. 'with' is used in opposition to or against. Thus option (b) is the correct answer.



Scan for Video solution



22. (d)

'Within' is used where something will occur in a specified period of time. 'For' is used for duration of time (Hour, Days, Week, Month) whereas 'Since' is used for specific point in time.



Scan for Video solution



23. (b)

The first part of the sentence is in future tense  
So, "were to be" cannot be used because it is in past tense. All other options can be used.



Scan for Video solution



24. (c)

Had I permitted him to leave, I wouldn't have had any problem with him being absent, would I?

The sentence is in past tense and having third form of verb (V<sub>3</sub>) hence 'had' will be there. Since the second part of the sentence contains 'wouldn't' so, the question tab 'would I' will be used.



Scan for Video solution



25. (b)

**At** is used to refer to time or place.



Scan for Video solution



26. (b)

The subject **company** is singular so the verb will be singular also i.e. **uses**. And as the subject **products** is plural the associated verb will be plural i.e., **include**.



Scan for Video solution



27. (c)

In this sentence ‘with’ is the suitable word for anger/fight with someone.

So, option (c) is correct.



Scan for Video solution



28. (a)

The sentence should be in present perfect continuous tense, therefore verb+ing is to be used, i.e. ‘Creating’.



Scan for Video solution



29. (b)

The given sentence has a blank in object. The objective pronoun is ‘me’.



Scan for Video solution



30. (b)

When a specific or definite noun is mentioned, ‘the’ article is used. For indefinite ‘a’ and ‘an’ is used based on consonant and vowel respectively. Here the first blank should consist definite article ‘the’ and second blank indefinite article ‘a’.



Scan for Video solution



31. (c)

For singular noun, singular verb is used while for plural noun, plural verb is used.

‘The expenditure’ is singular noun so singular verb ‘breaks down’ should be used.



Scan for Video solution



32. (b)

They’re → They are

Their → Pronounce for they

There → Denotes the place



Scan for Video solution



33. (c)

The ways in which this game can be played are infinite.

Since the subject of the sentence is plural hence ‘are’ will be used.



Scan for Video solution



34. (d)

He was one of my best friends and I felt his loss keenly.

‘One of the’ is used to describe something/someone from the same group.

Keenly means intensely, in a lager manner.



Scan for Video solution



35. (c)

Here first sentence in past perfect the second must be in past indefinite, so wished is the second form of wish which is used in past indefinite tense.

So, option ‘c’ is correct.



Scan for Video solution



36. (b)

Enough is used after the adjective or adverb and before noun, so that bright enough is suitable for this sentence.

So, option 'b' is correct.



Scan for Video solution



37. (a)

The comparative forms of the adjective "easy" are "easier" and "easiest" not "Most easier".

Than is used in comparisons as a conjunction.



Scan for Video solution



38. (a)

We use 'could have' to talk about something of a past possibility that didn't happen. And we use third form of verb with had.



Scan for Video solution



39. (c)

For facts or universal truth, simple present tense is used. So 'is'.



Scan for Video solution



40. (b)

Here 'If clause' should have past perfect tense, therefore 'Had been' is suitable in the given blank.



Scan for Video solution



41. (c)

'Tall' is used to measure height of a person or object. 'Long' is used to measure length of a thing.

Mr. X is taller than Mr. Y.



Scan for Video solution



42. (a)

The unruly crowd demanded that the correct be hanged without trial.



Scan for Video solution



43. (b)

The volume of a sphere of diameter 1 unit is less than, the volume of a cube of side 1 unit.

Lesser than would be incorrect because lesser & than both imply a comparison making them redundant. Less is comparative form of little.



Scan for Video solution



44. (a)

Mountain Everest is the highest peak in the world.

'Highest peak' is specific so we need to use the definite article 'The' before it.



Scan for Video solution



45. (b)

Security guard is a common man so that 'a' will be used as an article, and 'the' will be used with university so that option (b) is correct.



Scan for Video solution



46. (c)

We use "**than**" when we compare two objects or things directly with each other, as in "Your laptop is more powerful **than** my laptop" or, more succinctly, "Your laptop is more powerful **than** mine."

On the other hand, we use "**than that of**" when we compare not the two objects or things directly but an attribute, possession, or part of theirs, as in "The

processor of your laptop is more powerful than that of mine."

More costlier is wrong because we cannot use "two comparative degrees together in a single sentence".



Scan for Video solution



**47. (c)**

When 'If clause' is in past tense 'main clause' uses the word like 'could / would / should'. Here the blank should have negative, thus suitable word in blank is 'wouldn't buy'.



Scan for Video solution



**48. (c)**

This is famous quote 'Where there is a will, there is a way.' It means if someone has strong wish to do something, they would definitely find a way to do it.



Scan for Video solution



**49. (d)**

It is useless to use negative sentence after until. Until is not followed by a negative sentence.

Also, minister is singular thus singular verb 'meets' is to be used.



Scan for Video solution



**50. (c)**

The underline part can be replaced with "Increase in productivity necessarily" so the complete sentence is 'Increase in productivity necessarily reflects greater efforts made by the employees.'



Scan for Video solution



**51. (a)**

"It would remain between you & me" is the correct statement.



Scan for Video solution



**52. (b)**

The most appropriate word from the options given below is "should have practiced"

If the athlete wanted to come first in the race, he should have practiced several hours every day.



Scan for Video solution



**53. (b)**

Part Q is wrong. It should be "the instructor had marked"



Scan for Video solution



**54. (c)**

Misogynous: Showing feelings of hating women. The adverb of misogynous is "Misogynously". Adverb is the word or phrase that modifies an adjective, verb or a word group.



Scan for Video solution



**55. (b)**

Dhoni as well as other members of Indian team, was present on the occasion. The noun preceding "as well as" carries stronger emphasis than the noun following it.



Scan for Video solution



56. (c)

Completely must be used before verb so ‘completely forgot’ must be used in the 1<sup>st</sup> blank space. Just cannot be in between the don’t and know, therefore, ‘just don’t know will be used in 2<sup>nd</sup> blank space.

So, option (c) is correct.



Scan for Video solution



57. (a)

Much is used for uncountable things, no/few are not inferred with the statement, so that only option (a) is left, which is correct.



Scan for Video solution



58. (a)

- (a) Since the report lacked needed information, it was of no use to them
- (b) The report was useless to them because there was no needed information in it.
- (c) Since the report did not contain the needed information, it was not **really** useful to them
- (d) Since the report lacked needed information, it would not **have** been useful to them

So, option (a) is correct.



Scan for Video solution



59. (a)

If connective/ appositives like **along with, together with, as well as, accompanied by** etc are used to combine two subjects, the verb agrees with the subject mentioned first.

So, verb must agree with Tuberculosis which is a singular and hence **ranks** should be used.



Scan for Video solution



60. (c)

The suitable words to complete the sentence are:

The Tamil version of the John Abraham-starrer *Madras Cafe* was cleared by the Censor Board with no cuts last week but the film's distributors found no takers among the exhibitors for a release in Tamil Nadu on this Friday.



Scan for Video solution



61. (b)

Advice is a noun that means guidance whereas advise is a verb which means to guide someone.



Scan for Video solution



62. (b)

In the given sentence “If clause” is given. In this, the principal clause is in future tense (will/shall) so the conditional clause (if) should be in present indefinite tense.

Hence “gets some rest” is the correct option.



Scan for Video solution



63. (b)

Since communication and interpersonal skills are considered together here hence ‘both’ is the correct word.

‘Each’ is used for separate things, ‘all’ is not used for 2 things and ‘either’ is used to present alternative hence these three words are not appropriate here.



Scan for Video solution



64. (c)

The second part of the sentence is in past tense so first sentence should also be in past tense.



Scan for Video solution



**65. (b)**

This is an interrogative sentence, so that who is a question word, therefore 'did' is helping verb. Now that cannot be used here, therefore, option (b) is correct.



Scan for Video solution

**66. (d)**

Part (iv) is not appropriate, because the sentence is in the past that it must be 'lost consciousness'



Scan for Video solution

**67. (c)**

If first clause in the past perfect then main clause must be in the past indefinite.

Therefore, had known and could have best suited words in this sentence.



Scan for Video solution

**68. (d)**

For facts or universal truth, simple present tense is used. Thus 'experiences' is the suitable word.



Scan for Video solution

**69. (c)**

'An European' is wrong. It should be 'A European'.



Scan for Video solution

**70. (a)**

In a conditional sentence, if the part of the sentence containing "If I" is in past then resultant part of the sentence will contain 'would'.



Scan for Video solution

**71. (d)**

The sentence 'Two and two make four' is correct as the plural verb make agree with the plural compound subject 'two and two'.

So, option (d) is correct.



Scan for Video solution

**72. (b)**

'to' is used before the verb, here 'ordered to' is incorrect. It should be 'ordered'.



Scan for Video solution

**73. (d)**

In statement (D), 'very less' should be 'much less' or very little. Very less implies that it is less than something else or less than it was before.



Scan for Video solution

**74. (a)**

That → Essential clause

Which → Non – essential clause

Who and whom are for people not dogs.



Scan for Video solution

**75. (b)**

The sentence is in active voice as the speaker requested it, "should be given" is in passive voice and is an incorrect usage. It should be 'should give' for should be given.

Hence option (b) correct.



Scan for Video solution



76. (a)

For excreting outcome that are completely unrealistic or will not likely happen in future, are the simple past tense in the if-clause and an auxiliary modal verb (e.g., could, should, would, might) in the main clause (the one that express the unrealistic or unlikely outcome). Hence option (a) correct.



Scan for Video solution



77. (c)

Contemplate is a transitive verb so it is followed by gerund (a verb form which functions as noun) hence the correct sentence is “I contemplated **visiting** Singapore for my vacation but decided against it.”



Scan for Video solution



78. (a)

Planting : Seed :: Raising : Child

Planting a Seed is similar to raising a child



Scan for Video solution



79. (b)

‘Enforce’ and ‘Relax’ are opposite words to each other

∴ ‘Permit’ and ‘Forbid’ also have the same relation (opposite) to each other.

∴ Option (b) is correct.



Scan for Video solution



80. (a)

The legislature works in the parliament, similarly the judiciary performs their duty or work in the courts.



Scan for Video solution



81. (c)

**Conscious:** aware of and responding to one's surroundings

**Consensus:** a general agreement

**Conscience:** a person's moral sense of right and wrong, viewed as acting as a guide to one's behaviour.

**Consent:** permission for something to happen or agreement to do something.



Scan for Video solution



82. (a)

Seldom is antonym of often similarly cruel is antonym of kind.



Scan for Video solution



83. (c)

Insert is antonym of eject similarly retreat is antonym of advance.



Scan for Video solution



84. (d)

Since, some humans are intelligent means at least one human is intelligent so, option A is wrong. Option B is wrong because may be some humans are not intelligent. Option C is wrong because may be all the intelligent are humans.



Scan for Video solution



85. (b)

We reached the station late, and nearly missed the train.

Utterly → Completely

Mostly → Almost all the time.



Scan for Video solution



86. (b)

Frequently is the synonym of often similarly type is the synonym of kind referring to classification.



Scan for Video solution



87. (d)

The correct sequence that creates a coherent narrative is – (iii), (ii), (i), (iv).



Scan for Video solution



88. (b)

Writing too many things on the board while teaching could make the students get bored.



Scan for Video solution



89. (d)

After playing two hours of tennis, I am feeling too tired to walk back.



Scan for Video solution



90. (d)

Heel = back part of the foot  
 Heal = to become healthy again  
 ∴ Option (d) is correct.



Scan for Video solution



91. (c)

Neither/nor is used together to state “two or more things that are untrue or won’t happen”.



Scan for Video solution



92. (d)

No → rejection

Know → to be aware



Scan for Video solution



93. (d)

Fare → The amount paid for a journey on a public transport.

Fair → Impartial and just.

So, correct sentence is

The fare is too high for it to be considered fair.



Scan for Video solution



94. (c)

**Nostalgia:** Excessively sentimental yearning for return to or of some past period or irrecoverable condition.

**Anticipation:** A feeling of excitement about something that is going to happen in the near future.



Scan for Video solution



95. (c)

Oasis is the place in the sand which is wet.

Same island is the place in the water which is dry.

So, option (c) is correct.



Scan for Video solution



**96. (c)**

Doctor performs the surgery similarly writer writes the book.



Scan for Video solution

**97. (c)**

Pen is used to write similarly knife is used to cut.



Scan for Video solution

**98. (b)**

Squared the series means to have same number of wins/ points as the opponent in the series.



Scan for Video solution

**99. (b)**

Whitening makes the objects white similarly lightening makes the objects light (in weight).



Scan for Video solution

**100. (a)**

The recent measures to improve the output would increase the level of production to our satisfaction



Scan for Video solution

**101. (b)**

In the sentence “Having said that” means despite what I have said.



Scan for Video solution

**102. (c)**

Build is a verb and building is a noun similarly grow is a verb and growth is a noun



Scan for Video solution

**103. (c)**

Shed crocodile tear means to pretend to be sad. Hence the correct sentence is “He is known for his unscrupulous ways. He always sheds crocodile tears to deceive people.”



Scan for Video solution

**104. (b)**

Coup = seizure of power,

Poised = poison

Plunged = rise/fall

Plugged is use to switch on some electrical equipment.

Rise or fall is the part of stock market,

∴ Option (b) is correct.



Scan for Video solution

**105. (c)**

Do and undo are opposite to each other,

So, opposite word of trust is distrust.

∴ Option (c) is correct



Scan for Video solution

**106. (b)**

**Explicit** describes something that is very clear and without vagueness or ambiguity. **Implicit** often functions as the opposite, referring to something that is understood, but not described clearly or directly.

**Express:** To make known (as an idea, emotion or opinion)

**Repress:** To repress is to hold something back or to prevent an act of volition, especially by force.

**Suppress:** To suppress something is to put an end to it forcibly, among other things.



Scan for Video solution



### 107. (a)

'Fuse' is verb and 'Fusion' is its noun form. Similarly, 'Use' is verb and 'Usage' is its noun form.



Scan for Video solution



### 108. (b)

Ivory towers → unaware of reality. Thus option (b) is the answer.



Scan for Video solution



### 109. (c)

Voracious → doing something with enthusiasm.

Precocious → Advance, Ahead.

Thus suitable word for the given blank is voracious.



Scan for Video solution



### 110. (a)

Disparage → To lower reputation

Applaud → praise

Reduce → decrease

Begrudge → to be envy or jealous



Scan for Video solution



### 111. (a)

'But' in the sentence shows that last part of sentence contradicts to the first part of sentence. So correct option is 'extremely poor'.



Scan for Video solution



### 112. (a)

Here 'cook' is verb then 'cook' is noun.

Cook (subject) cooks (verb) the food similarly, the flyer flies the aircraft.



Scan for Video solution



### 113. (d)

In the sentence, 'more' is correct because 'than' comes for comparative degree.

"Are there enough seats here? There are more people than I expected."



Scan for Video solution



### 114. (a)

Culmination means highest or end point of something especially attained after a long time.

So, correct sentence is "A final examination is the culmination" of a series of evaluations that a student has to go through.



Scan for Video solution



**115. (b)**

The verb ‘skirt’ means avoiding the discussion of a subject hence the correct sentence is:

The minister avoided any mention of the issue of women’s reservation in the private sector. He was accused of skirting the issue.



Scan for Video solution

**116. (c)**

John Thomas an eminent writer, passes away in 2018.

Eminent means famous and respected.

After ‘an’ the word should start with vowel sound.



Scan for Video solution

**117. (d)**

In first sentence, discussion about delight or joy but in second sentence, it is for trouble, so even cannot be used here. Because both sentences have some different meaning, trouble is due to news paper so it should be too.

So, option (d) is correct.



Scan for Video solution

**118. (d)**

Deteriorate means decline or worse. So, it cannot be the proper word in this sentence.

Accommodate means occupied something, or providing sufficient space. Therefore, accommodate is the suitable word to this sentence.



Scan for Video solution

**119. (c)**

‘quite a few’ means a fairly large number. ‘Quiet’ means silence. Thus option (c) is appropriate in the given blanks.



Scan for Video solution

**120. (c)**

‘Reach’ is appropriate word for the given blank. It means extend or stretch out. Peak is highest point or top.



Scan for Video solution

**121. (d)**

In comparison ‘off’ word is inappropriate. Also in a singular noun ‘those’ can’t be used. Thus, the blank should consist ‘as that of’.



Scan for Video solution

**122. (c)**

From the passage, the meaning of word “arrived” can be taken as ‘attained a status’.



Scan for Video solution

**123. (a)**

The word ‘revolve’ in the sentence infer to ‘an element which is treated as the most important’ so correct sentence is.

The search engine’s business model revolves around the fulcrum of trust.



Scan for Video solution



**124. (d)**

A court is to a judge as school is to a teacher.  
 The workplace for a judge is a court  
 The workplace for a teacher is a school.



Scan for Video solution

**125. (a)**

Undiminished: Not reduced or lessened.  
 “The judge’s standing in the legal community, through shaken by false allegation of wrongdoing, remained undiminished.



Scan for Video solution

**126. (d)**

Complement: Refers to something that completes something else.  
Compliment: A polite expression of praise or admiration.  
 So, complete sentence is “The dress Complemented her so well that they all immediately Complimented her on her appearance.



Scan for Video solution

**127. (c)**

Miserliness: Extreme meanness or excessive desire to save money (Negative meaning)  
Frugality: Quality of being economical with money (Positive meaning)  
 Her Frugality should not be confused with miserliness, she is ever willing to assist those in need.



Scan for Video solution

**128. (a)**

Going by the Principle that many hands make light work, then school Principal involved all the students in the task.

Principle → A fundamental truth.

Principal → The most important person in an organization.



Scan for Video solution

**129. (d)**

As per given in the question, that ideas must be complicated means confused, so that detailed/simple/clear words are not suitable. Convoluted means difficult which is inferred to complicated.

So, option (d) is correct.



Scan for Video solution

**130. (a)**

Course = way or direction or path  
Coarse = rough area or harsh land

“Since you have gone off the way, the rough area sand is likely to damage the car.”

So, option (a) is correct.



Scan for Video solution

**131. (a)**

**Improvement**: A thing that makes something better.

**Neglected**: Not receiving proper attention.

**Glum**: Looking or feeling dejected

**Disinterested**: Not influenced by considerations of personal advantage.

“Even though there is a vast scope for its **Improvement**, tourism has remained a **Neglected** area.”



Scan for Video solution




**132. (b)**

**Piece:** a portion of an object produced by cutting or breaking the whole.

**Peace:** A lack of conflict and freedom from fear of violence between individuals or groups.


**Peas:** A type of vegetable.

**133. (a)**


Brake → Attempt to stop

Break → relax / rest.

Thus the sentence should be ‘The driver applied the brake as soon as she approached the hotel where she wanted to take a break.’

**134. (a)**


As we cannot be aware of the codes or rules of different society, thus the proper word here in blank should be ‘Unpredictable’.

**135. (a)**


Shone → shine → glitter

Shown → show → exhibit.


Hence the correct option is (a).

**136. (a)**


‘Pioneering ideas’ means new ideas. Thus, the suitable word is ‘Innovative’.

**137. (a)**

Meandering → Indirect, not organized

**138. (d)**

Cleaving mean splitting.


**139. (a)**

Forgo: go without (something desirable)

Exclusive: can not occur at the same time (events).


So, the correct sentence is.

If you choose plan P, you will have to forgo plan Q, as these two are mutually exclusive.

**140. (d)**


As the two speakers become increasingly agitated, the debate become heated.

Here heated means angry, furious, fierce, intense.

**141. (d)**

As per research, people doing work for many reasons besides money.

So, option (d) is correct.



## 142. (d)

Aversion means strong dislike,  
 Luminous or radiant means lighting  
 Plunder or loot means robbing  
 Yielding means giving and resistance means stopping or to oppose.  
 So, option (d) is correct.



Scan for Video solution



## 143. (c)

**Figuratively:** used to indicate a departure from a literal use of words; metaphorically  
**Retrospectively:** with consideration of past events or situations  
**Respectively:** Separately or individually and in the order already mentioned  
**Rightfully:** In accordance with a legitimate right.



Scan for Video solution



## 144. (d)

**Antagonistic:** Showing or feeling active opposition or hostility towards someone or something.  
**Impartial:** Treating all rivals or disputants equally.  
**Argumentative:** using or characterized by systematic reasoning  
**Hostile:** Showing or feeling opposition or dislike; unfriendly



Scan for Video solution



## 145. (a)

Sharp tongued means to be someone who often criticizes and speaks in a severe way.



Scan for Video solution



## 146. (c)

Involving → participation  
 Assisting → Guiding  
 Tampering → Modification/change  
 Incubating → keeping warm the egg to make it hatch. Thus ‘tampering’ is suitable word in the given blank.



Scan for Video solution



## 147. (b)

Pretty → Lovely  
 Sloppy → careless  
 Haphazard → Not systematic  
 Thorough means systematic and complete. Thus option (b) is correct.



Scan for Video solution



## 148. (c)

The phrase ‘rest is history’ means rest is well known.



Scan for Video solution



## 149. (b)

Felicitated: Congratulated.

The students felicitated the teacher on teacher’s day for twenty years of dedicated teaching.



Scan for Video solution



**150. (d)**

Prone: Likely to suffer from, lying flat especially face downwards.

Prune is a fruit which is a dried plum.

So, statement (ii) & (iii) are correct.



Scan for Video solution

**151. (b)**

The verb that follows ‘did’ should be used in its 1<sup>st</sup> form. So, lose is the correct word.



Scan for Video solution

**152. (a)**

The correct option to create a meaningful sentence is A

Despite the new medicine’s effectiveness in treating diabetes, it is not prescribed widely.



Scan for Video solution

**153. (b)**

Manage + able = manageable, is the correct word here, so that So, option (b) is correct.



Scan for Video solution

**154. (c)**

Bare means without cover

Bear means tolerate with

So that bear with is best suited in this statement.

So, option (c) is correct.



Scan for Video solution

**155. (a)**

Cope with means deal with or put up with  
So, option (a) is correct.



Scan for Video solution

**156. (c)**

Mock means to copy someone or deride or fun as someone or jeer.

Praise means to compliment someone.

Praise is the odd one here

So, option (c) is correct.



Scan for Video solution

**157. (c)**

**Grasp:** To take and hold (something) with your fingers, hand etc.

**Clutching:** Clutch suggests anxiety in seizing or grasping and may imply less success in holding.



Scan for Video solution

**158. (a)**

Turning deaf ear is not listening or ignoring. Thus option (a) is correct.



Scan for Video solution

**159. (a)**

Paraphrase → Restatement of something already written or spoken

Paradox → A Self-contradictory statement

Paradigm → A specific pattern or model

Paraffin → A wax like solid obtained from petroleum.

So, the correct word to complete the sentence is ‘paraphrase’.



Scan for Video solution



### 160. (a)

**Figurative** → Representing examples from real life.

**Collateral** → A security for loan repayment.

**Literal** → Simple meaning of given word or statement.

**Figurine** → Human stature.

So, the correct word to complete the sentence is ‘figurative’.



Scan for Video solution



### 161. (c)

The meaning of idiom "fish out of water" is 'a person who feels awkward or unhappy because they are in a situation that is not familiar or because they are different from the people around them. Other options have no meaning.



Scan for Video solution



### 162. (b)

**Elusive**: Difficult to find, achieve.

**Allusive**: To describe something that makes a reference, hint.

**Elude**: Scape from or avoid.

So, the correct statement is ‘When the thief keeps eluding the police, he is being elusive’.



Scan for Video solution



### 163. (a)

**Harbour**: Keep a thought or felling (typically a negative one) in one’s mind, so, the sentence will be “apparent lifelessness harbours dormant life.



Scan for Video solution



### 164. (b)

Introvert, atheist and altruist all are nouns hence ‘alternate’ is the correct answer as it is a verb.



Scan for Video solution



### 165. (b)

**Connotation**: connotations of a particular word are the ideas or qualities which the word make you think of.

**Pertinacious**: holding firmly to an opinion

**Viaticum**: provision for a journey

**Clandestine**: kept secret or done secretly

**Ravenous**: Extremely hungry



Scan for Video solution



### 166. (a)

**Awkward**: Causing difficulty, hard to deal with

**Inept**: Generally incompetent (showing no skills)

**Graceful**: having elegance

**Suitable**: Right for the purpose or situation.

**Dreadful**: Causing great suffering

So, most similar word to awkward is ‘inept’.



Scan for Video solution



**167. (c)**

Immensely means great/terrific/good time.  
 Terrible means bad/horrible/terrifying  
 So, option (c) is correct.



Scan for Video solution

**168. (b)**

- (a) Acquiescence/acceptance/submission - same words.
  - (b) Wheedle/flat/direct and Roundabout/not direct – opposite words
  - (c) Flippancy/Lightness/lack of respect – same words
  - (d) Profligate/Extravagant/reckless – same words
- So, option (b) is correct.



Scan for Video solution

**169. (c)**

'took to their heels' means run away or took to flight  
 So, option (c) is correct.



Scan for Video solution

**170. (c)**

Personnel means employees or staff  
 Personal means private or individual

- (a) The industrialist had a **personel** jet
  - (b) I write my experience in my **personel** diary
  - (c) All **personnel** are being given the day off
  - (d) Being religious is a **personel** aspect
- So, option (c) is correct.



Scan for Video solution

**171. (d)**

Fabric/textile = cloth made by fibre  
 Fibre = cotton or thread by which fabric is made  
 Apparel = attire/garments/shirt/trousers/shirt  
 So, option (d) is correct.



Scan for Video solution

**172. (a)**

**Croak:** A characteristic deep hoarse sound made by a frog or a crow.



Scan for Video solution

**173. (c)**

**Educe:** Bring out / extract or develop something latent or potential.



Scan for Video solution

**174. (b)**

**Memento:** An object kept as a reminder of a person or event.



Scan for Video solution

**175. (b)**

Respectably → Acceptable standard.  
 Respectfully → with respect  
 Respectively → As mentioned in order or sequence  
 Reputably → Image or standard  
 Hence the suitable word in the blank is 'Respectfully'.



Scan for Video solution



**176. (b)**

Insured → to give protection

Ensured → to approve/make sure.

Thus in statement (b) the word is used correctly



Scan for Video solution

**177. (c)**

Vernacular → Native Speech, Common language

Regional → particular area

Indigenous → originally natural, native

Indigent → poor

Colloquial → used is ordinary language.

Except ‘Indigent’ all other words have same meaning.



Scan for Video solution

**178. (d)**

Dreary → Dull or boring

Dismal → Less attractive or without hope

Thus option (d) is correct.



Scan for Video solution

**179. (b)**

Extraneous → Irreverent/ which is not related

So, the correct word that fits in the sentence is ‘extraneous’.



Scan for Video solution

**180. (b)**

Pediatrician → Doctor for kids

Orthopaedist → Bone doctor

Gynaecologist → Doctor for women

Nephrologist → Kidney doctor

Dermatologist → Skin doctor



Scan for Video solution

**181. (b)**

**Vindicated** → justified

Substantiated → justified

Chastened → to bring down

So, the correct word will be ‘substantiated’



Scan for Video solution

**182. (b)**

**Revert** → reply

Retract → to take the statement back

Get back to him → reply

Move in reverse → to move backward

Retreat → moving away

So, the correct word will be ‘Get back to him’.



Scan for Video solution

**183. (a)**

Dissent → having opinions that are different from majority

Descent → to go downward

Decent → proper/civilized

Decadent → to decline

So, the correct word will be ‘dissent’.



Scan for Video solution



**184. (b)**

Depressed → Sad

Depreciated → Loosing value

Appreciated → Praise

Stabilized → To be stable

Hence most appropriate word is ‘depreciated’.



Scan for Video solution

**185. (c)**

Bear → to carry/stand a thought

Bare → uncovered part of the body

Losing → to set free

Losing → suffering a loss.

So, the appropriate pair of word is ‘Bear, Losing’



Scan for Video solution

**186. (b)**

Impel → To force someone

Dispel → To dissappear

Propel → To push

Repel → To get back

Hence most appropriate word is ‘dispel’.



Scan for Video solution

**187. (b)**

‘cope with’ means to face someone or to deal with or adjust with

Adopt to means accepting legally

Adapt to means to adjust with

Adept in means skillful

Accept with means agree with

∴ Option (b) is correct



Scan for Video solution

**188. (c)**

The person was not sure about her performance that she should be awarded by first prize because her performance was mediocre.

∴ Option (c) is correct.



Scan for Video solution

**189. (c)**

‘The buck stops here’ means that I will take the responsibility.

∴ Option (c) is correct.



Scan for Video solution

**190. (a)**

Post means after any duration,

In early days, due to the exisance of the British colony, now India is a post-colonial country.

∴ Option (a) is correct.



Scan for Video solution

**191. (a)**

Eradicate means to remove completely which is equivalent to destroy utterly. 1 → S

Distort means to mislead or Misrepresent. 2 → P

Saturate means Thorley soaked. 3 → Q

Utilize means thoroughly use. 4 → R

∴ Option (a) is correct.



Scan for Video solution




**192. (c)**

Coherent means reasonable or logical or clear meaning.

Sticky means attachable


Rambling means confusing

Therefore, option (c) is correct.

**193. (a)**


Comprehension → Ability to understand or grasp.

The correct answer is ‘Understanding’.

**194. (b)**

Vice → bad behavior

Virtues → good behavior. Thus the correct option is ‘Virtues’


**195. (a)**

Contentment → satisfaction


Ambition → goal, aim

Perseverance → to be firm


Hunger → desire to eat. Thus the appropriate word is ‘Contentment’.

**196. (c)**

‘As a women, I have no country’ means there cannot be boundaries or barriers to divide women. All women are united. Thus option (c) is the closest in the meaning to the given sentence.

**197. (b)**

Nadir → The lowest or the most unsuccessful point in a situation.

**198. (a)**

Universalism → All

Particularism → Specific


Diffuseness → Spread in direction

Specificity → The quality of belonging uniquely to a particular subject

Neutrality → Neutral in quarrel

Generality → General

Adaptation → Oneself Suitable to environment

**199. (d)**


**Quarrel** means to have an angry argument or disagreement.

**Make out** means manage with some difficulty/to see or hear someone or something.

**Call out** means for gathering or to deal with an emergency or provide a service.

**Dig out** means to remove earth or debris from by digging, to hollow out by digging, to find by searching.


**Fall out** means quarrel or to cut off relations over a quarrel. Former friends who have fallen out.  
∴ Option (d) is correct.



**200. (b)**

A preposition must be connected between these two words so that sentence will be dare to commit mistakes.


∴ Option (b) is correct.



**201. (c)**


“A friend in need is a friend indeed” is a proverb which means a person who helps at a difficult time is a person who you can rely on.

So, option (c) is correct.



**202. (c)**


Primeval → Ancient / past  
Modern → Recent/ present  
Historic → famous/significant  
Primitive → Earliest/ Ancient  
Antique → old valuable thing/ Treasure.  
Thus Primeval means ‘Primitive’.



**203. (b)**


Cordial → friendly

Intimate → familiar/close. Thus, the option (b).



**204. (c)**

Medicine is taken to improve health similarly education is taken to improve knowledge.



**205. (a)**


Mitigate → to reduce

Diminish → reduce the size

Divulge → make something

Dedicate → to offer


Denote → to represent



**206. (b)**

Setbacks → an event that make someone to lose progress.

The correct sentence is “Despite several **setbacks** the mission succeeded in its attempt to resolve the conflict.”



**207. (b)**

**Latitude:** scope for freedom of action or thought

**Eligibility:** the state of having the right to do or obtain something through satisfaction of the appropriate conditions.

**Freedom:** the power or right to act, speak, or think as one wants.

**Coercion:** the practice of persuading someone to do something by using force or threats.

**Meticulousness:** marked by extreme or excessive care in the consideration or treatment of details  
Hence option (b) is correct.



Scan for Video solution



208. (d)

**Beggary:** a state of extreme poverty

**Nomenclature:** the devising or choosing of names for things, especially in a science or other discipline.

**Jealousy:** the state or feeling of being jealous.

**Nonchalance:** (of a person or manner) feeling or appearing casually calm and relaxed; not displaying anxiety, interest, or enthusiasm

Hence option (d) is correct.



Scan for Video solution



209. (c)

The topics in options a, b and d should be included to deal with bereavement and grief to help students in times of losses though death and parting but topic in option c cannot help.



Scan for Video solution



210. (b)

Understated → Not trying to attract people

Tentative → Uncertain or with hesitation

Restrained → Doing something in limit



Scan for Video solution



211. (a)

Inexplicable → Puzzled, Unclear, Unsolved

Incomprehensible → Not able to understand

Indelible → Cannot be erased

Inextricable → Which cannot be separated

Infallible → In capable of doing something



Scan for Video solution



212. (b)

**Amalgamate** → To unite, to merge

So opposite word will be split.



Scan for Video solution



213. (d)

The sentence implies human genes are to be manipulated only to correct diseases for which the satisfactory treatments are not available.

Hence option (d) is correct.



Scan for Video solution



214. (b)

**Frequency:** the number of times something is repeated, or the fact of something happening often.

**Periodicity:** the quality of character of being periodic; the tendency to recur at intervals.

**Rarity:** (of an event, situation or condition) not occurring very often.

**Gradualness:** occurring or developing slowly or by small increments.

**Persistency:** the fact of continuing in an opinion or course of action in spite of difficulty or opposition.

So, ‘Rarity’ is opposite to ‘Frequency’ and hence option (b) is correct.



Scan for Video solution



**215. (c)**

The sentence implies that foreign technocrats were harmful to the country. So, only suitable opinion is **Exacerbate** – meaning aggravate or make things worse or increase the severity

∴ Option (c) is correct.



Scan for Video solution

**216. (a)**

Arena is the combat place of the gladiator or some other animals. It means gladiator performs in the arena. The same way a dancer performs on the stage.

So, option (a) is correct.



Scan for Video solution

**217. (b)**

Belied → Being in contradiction or unable to give a true impression.



Scan for Video solution

**218. (b)**

Circuitous → Route longer than direct way.



Scan for Video solution

**219. (a)**

**Fallow** → refers to the land which is ploughed but left for a period without being sown in order to restore fertility.

Worker who is not working is unemployed similarly land which is not producing anything is fallow.



Scan for Video solution

**220. (d)**

- (a) Uphold → Cause to remain
- (b) Restrain → Keep under control
- (c) Cherish → Be fond of
- (d) Conserve → Keep in safety



Scan for Video solution

**221. (c)**

The correct sequence of sentence that creates a coherent narrative is: II → I → IV → III.



Scan for Video solution

**222. (d)**

In the paragraph it is given that almost all of Ravi's Friends at school were hard working and kind too. So, it is not necessary that Ravi's Friends outside the school are hardworking and kind too, and also almost 'all' infer 'some' so, from the passage it is certain that "Some of Ravi's Friends are hardworking and kind."



Scan for Video solution

**223. (d)**

From the conclusion of the given passage, we find that option a, b, c, describe the passage so option a, b, c is incorrect.

Option (d) is correct because modern cell biology uses figurative language, such as metaphors, to describe or explain anything.

∴ Option (d) is correct.



Scan for Video solution



**224. (d)**

It may be possible that student, who would not eligible to vote will belong to any other department not only to the Human Sciences, hence statement (i) is not correct.

It may be possible that student, who failed to complete the registration process within time will belong to any other department not only to the Human Sciences, hence statement (ii) is not correct.

Statement (iii) is correct because no student from the department of Human Sciences was eligible to vote as they failed to complete the registration within time.

∴ Hence only statement (iii) is correct.

∴ Option (d) is correct.



Scan for Video solution

**225. (b)**

As given in the statement that 65% of the tobacco users were advised to stop consuming tobacco but only 3 out of 10 users attempted to stop using tobacco, it means only 30% of that 65% tobacco users attempted to stop using tobacco. Which indicates that majority of the tobacco users who were advised to stop consuming tobacco did not attempted to do so. Only some of the majority attempted to stop using tobacco.

∴ Option (b) is correct.



Scan for Video solution

**226. (b)**


The sentence “when I was a kid” implies it is an adult’s memory of what he or she liked as a child.



Scan for Video solution


**227. (d)**

Option (a) is wrong because




Option (b) is wrong because “All creatures that have clause are ferocious”.

Option (c) is wrong because



**Option (d) is right** because “same ferocious creatures may not have claws”



Scan for Video solution

**228. (d)**

From reading the paragraph we certainly can infer that the government of zombieland believes that riding bicycles is a form of physical exercise.




Scan for Video solution




## 229. (d)

In Elvesland, gifting a white flower to someone is taken as declaration of love but one not always declare one's love by gifting white flower. Yellow flower is gifted to show anger. So, statement iv is correct.




## 230. (d)

The mother's feelings about her daughter's singing at home were not only of annoyance but most often this made the mother pensive (thoughtful). So, option (d) is correct.




## 231. (d)

"Dark matter is supposed to interact only via the gravitational interaction and not through the electromagnetic-, the weak- or the strong-interaction. This may justify the epithet "dark" in dark matter." So, option (d) is false and all other are true.




## 232. (b)

"For example, babies who start eating healthy in the first year are more likely to have better overall health as they get older." **implies Eating healthy can be started at any age, earlier the better.**




## 233. (b)

"The total number of visitors in the malls is impressive. However, the total revenue generated through sales in the shops in these malls is generally low." Implies **More people are visiting the malls but not spending enough.**




## 234. (c)

Having ultra black skin is advantageous because it helps them to avoid predators & also attack their prey. It also has disadvantage because it makes their skin fragile.



## 235. (b)

In the passage it is clear that the primary reason for people to return the wallet is empathy which means the ability to understand the feeling of another. So, wallets with a key are more likely to be returned because people relate suffering of others.




## 236. (c)

As given in the passage that altruism may be attributed to empathy or guilt has now been rejected. So that option (a), (b) and (d) is not correct.

Humans engage in altruism due to group identification and familiarity

∴ Option (c) is correct.



**237. (d)**

Controlling mosquitoes using chemicals may have undesired consequences implies “**Using chemicals to kill mosquitoes may have undesired consequences**”

And it remains to be seen if this novel approach has unforeseen consequences implies “**it is not clear if using genetically modified mosquitoes has any negative consequence**”.



Scan for Video solution

**238. (b)**

The passage clearly indicates that the matrices are useful in the organizations with good work culture. So, option (c) is wrong and options (a) and (d) also wrong as discussed above.



Scan for Video solution

**239. (d)**

In the two given sentences, it is observed that facing a crisis is the effect of pandemic, hence option (d) is correct.



Scan for Video solution

**240. (c)**

In the symbiotic relationship described in passage, the primary benefit for Oxpeckers is that they get food source & that for rhinos is that they may be saved from the poachers.



Scan for Video solution

**241. (c)**

From the passage it is clear that the indication of healthy lifestyle is related to consumption of leafy vegetable.



Scan for Video solution

**242. (d)**

The conclusion of the passage is that to do well in an exam, adequate sleep must be part of the preparation.

∴ Options (d) is correct.



Scan for Video solution

**243. (d)**

“Computers are ubiquitous” implies **Nowadays, computers are present in almost all places**. “They are used to improve efficiency in almost all fields from agriculture to space exploration. For humans, sitting in front of a computer for long hours can lead to health issues” implies **for humans, there are both positive and negative effects of using computers**.

“AI enables computers to learn, given enough training data” implies

**Artificial intelligence cannot be done without data.**

So, option (d): (i) & (iii) is correct.



Scan for Video solution

**244. (c)**

We do not know of any species other than humans who posses the ability to construct mental worlds.



Scan for Video solution



**245. (a)**

The author is of the opinion that rehearsing is important for musicians, actors and public speakers.



Scan for Video solution

**246. (d)**

The passage indicates that the core problems that causes obesity is poverty and income inequality. Since AOM is not addressing the core problems, so AOM is superficially addressing the problem.



Scan for Video solution

**247. (c)**

Since it is clear from the passage that only some students who did not need any external stimulation, benefited from music so option c is correct.



Scan for Video solution

**248. (c)**

Mitigation deals with actions taken to reduce the use of fossils fuels.

Adaptation deals with the process of adjustment to actual climate, and its effects in order to moderate harm due to climate change.



Scan for Video solution

**249. (d)**

In a crowd funding, fund is raised from a large number of people willing to (voluntarily) contribute on web-based platforms.



Scan for Video solution

**250. (a)**

It is clear from the passage that **world-wide financial crisis** started with the sub-prime lending crisis in USA in 2007. And the subprime lending crisis led to the **banking crisis** in 2008 with the collapse of Lehman Brothers in 2008. The **sub-prime lending crisis** arised because of excess liquidity following the **East Asian crisis**.



Scan for Video solution

**251. (c)**

The French-speaking couple were upset at the English announcements being longer than the French ones.



Scan for Video solution

**252. (d)**

As per given information

Theory given by the American psychologist Howard Gardner. Based on this theory, Modern educationist insist that the teaching curriculum and evaluation needs to be multidimension. Therefore option D is correct.



Scan for Video solution

**253. (a)**

As the inflation is high real interest rate will be low borrow get benefited.

As the under low inflation, the interest rate will be high lenders get benefited. Therefore option A is correct.



Scan for Video solution



**254. (d)**

Partial : Impartial :: Popular : unpopular



Scan for Video solution

**255. (d)**

From the passage, melting glaciers are affecting billions of people and nation-states of UN need to rethink their carbon footprint, infers that some nation-states are not very serious for environment change.



Scan for Video solution

**256. (d)**

It is clearly mentioned in passage that GST is not imposed on the point of origin rather it is imposed on point of destination/usage.



Scan for Video solution

**257. (a)**

From the paragraph it can be inferred that most households that subscribe X's loyalty membership discontinue watching cable television.



Scan for Video solution

**258. (b)**

The teacher was surprised because he/she received the stories from the students that were **all** set in distant places and none of the students had written stories set in India.



Scan for Video solution

**259. (a)**

Since the name Zaira was a Portuguese alteration of *Nzadi o Nzere* which is an African term. So, Mobuto was not entirely successful in Africanising the name of his country.



Scan for Video solution

**260. (a)**

In the question statement about men are prone to buying on impulse as women but not about many/few/all men/women but it can be equivalent to some men and women, therefore option (a) is correct.



Scan for Video solution

**261. (c)**

"Indian history was written by British historians-extremely well documented and researched, but not always impartial" implies **Indian history written by British historians was well documented and researched but was sometimes biased.**



Scan for Video solution

**262. (d)**

In the passage it is not mentioned who duped the tribals out of their land holdings. Hence, option D is correct.



Scan for Video solution

**263. (b)**

In the given paragraph it is said 'Protagonists usually read books of aspirational values'. Here the word usually does not mean only. They may read other books too. Therefore option (b) cannot be inferred.



Scan for Video solution



**264. (b)**

From the passage it is clear that the new cotton technology, Bollgard-II is not legal in India, but farmers are using this technology, this indicates that farmers want to access the new technology even if it is not legal.



Scan for Video solution

**265. (b)**

The passage infers that beggars are created because of the lack of social welfare schemes.



Scan for Video solution

**266. (d)**

"It has remained inaccessible to tourists due to poor connectivity and lack of accommodation. A company has spotted the opportunity and is planning to develop a luxury resort with helicopter service to the nearest major city airport" implies **Helicopter connectivity would lead to an increase in tourists coming to the region.**



Scan for Video solution

**267. (c)**

"However, his on-field temper has always been a matter of concern for them since his junior days" implies **john's temper as a player not captain.** "After many years, it appears that the team has finally found a complete captain" implies **finding a complete captain is a challenge.** "Over the past three seasons though, that picture of John has been replaced by a cerebral, calculative and successful

batsman-captain" implies **Over the past three seasons John has accumulated big scores.**



Scan for Video solution

**268. (b)**

James was the best player in his own family this means Mark was not best in his family so, Steve was known to play better than Mark.



Scan for Video solution

**269. (c)**

Stuart's demonstration had no effect on his superior so option A is not correct. The Europeans were prejudiced so option B is not correct Stuart was reprimanded by commander-in-chief, so option D is also not correct. So, option (c) is correct.



Scan for Video solution

**270. (b)**

As mentioned in the statement inadequately or improperly nationalist is just not history, but it is viewed through the filter of nationalism. Which is not based on the imagination or past.

So, option (b) is correct.



Scan for Video solution

**271. (b)**

The author's belief that ideology is not as important as literature is revealed by the word seemingly. So, option (b) is correct.



Scan for Video solution



**272. (a)**

"for though I have spent a lifetime in the country. I lived too near the seat of events, and was too intimately associated with the actors, to get the perspective needed for the impartial recording of these matters." Implies **An intimate association offers partial perspective and hence matters are not recorded with an impartial perspective.**

Actors are typically associated with the partial recording of matters.

So, we can say "An intimate association does not allow for the necessary perspective."



Scan for Video solution

**273. (b)**

From the passage, air pockets are created by excavation of large amount of soil. So, option A & D is wrong. Old buildings do not form impossible hurdle to metro construction in that area.



Scan for Video solution

**274. (a)**

Since, the customers came out of the centre in the order that they went in infers that the service pattern in first-come-first-served basis.



Scan for Video solution

**275. (c)**

Kalimpong < Gangtok

Pelling < Gangtok

Pelling > Siliguri

Darjeeling > Gangtok

From the above conditions, statements (ii) & (iv) are correct.



Scan for Video solution

**276. (a)**

From the paragraph it can be inferred that social science disciplines had a pre-colonial origin because the disciplines were in existence in an amorphous form until the colonial period.



Scan for Video solution

**277. (c)**

From the paragraph it can be inferred that surveillance state is an perception of use of technology to ensure safety & security of smart city.



Scan for Video solution

**278. (b)**

From the passage it can be inferred that he was sure that his friends would help him when in need but his friends did not help him last week.



Scan for Video solution

**279. (c)**

It is not mentioned that SIT student will get an IIT degree so that statement (i) is incorrect.

Any institute but not only SIT can only be established in consultation with IIT so that statement (iii) is also incorrect.

SIT is doing this as a national objective so that they can provide good student to the country by which any country can develop.

So, option (c) is correct.



Scan for Video solution



**280. (c)**

As per the passage the woman who had done their masters in mechanical engineering are successful. Which means that only option (c) is best suited this statement. As most of the people have believed that masters in mechanical engineering are not suitable for woman so that option (b) is wrong.

Option (a) and (d) do not have any sense with the passage.

So, option (c) is correct.



Scan for Video solution

**281. (b)**

R2D2 is a robot. R2D2 can repair aeroplanes. No other robot can repair aeroplanes.

The conclusion of the above statement is R2D2 is the only robot which can repair aeroplanes.

So, option (b) is correct.



Scan for Video solution

**282. (d)**

Author believes that computers have advantages as well as disadvantages, so that statement (i) is not correct.

Mobile computers and the internet are both unintended inventions, so that statement (ii) is not correct.

So, option (d) is correct.



Scan for Video solution

**283. (a)**

“The overwhelming number of people infected with rabies in India has been flagged by the World Health Organization as a source of concern” implies **the number of people in India infected with rabies is high.**

Also inoculating 70% of pets and stray dogs against rabies can lead to a significant reduction not eradication.



Scan for Video solution

**284. (d)**

From the passage it can be inferred that the linguistic pluralism is one of the indicators of nation's diversity. In India there are 22 officially registered spoken languages. Linguistic pluralism is strong evidence of India's diversity.



Scan for Video solution

**285. (d)**

From the passage “Historians tend to correlate greatness of a king at his time with the availability of evidence today” infers that a king's greatness as we know him today, is interpreted by historians.



Scan for Video solution

**286. (b)**

From the paragraph it can be inferred that Ms. x will reach Kochi on 21 may & will be Bagdogra on 22 may but she may visit Kochi again in June or any other month.



Scan for Video solution

**287. (b)**

The correct order of the sentences for a meaningful paragraph is

1, 5, 2, 4, 3, 6.



Scan for Video solution



**288. (b)**

From the paragraph its can be inferred that the artists expected funding for the arts to increase this year & schools are giving less importance to arts education nowadays.



Scan for Video solution

**289. (a)**

The government shouldnot ban the water supply in lower areas. Therefore statement (iii) is not correct. So, option ‘a’ is correct.



Scan for Video solution

**290. (b)**

Because the wall sometimes breaks, we can say **Humpty Dumpty does not fall sometimes while having lunch.**



Scan for Video solution

**291. (b)**

“But on the contrary, human progress seems contradictory to environmental security” implies **Human progress is contradictory to environmental security.**



Scan for Video solution

**292. (a)**

From the statement, it can be inferred that since Alexander conquered Persia, he headed for India. Now if he had not conquered Persia then he would not move towards India.



Scan for Video solution

**293. (b)**

From the passage it is clear that he already had all the technical skills and to become a great batsman he has to work on his temperament. He was guilty of throwing away his wickets several times after working hard to lay a strong foundation.



Scan for Video solution

**294. (c)**

In option C, 22 official languages and thousands of dialects represents the rich heritage and cultural diversity of India.



Scan for Video solution

**295. (b)**

Male above 10 lakhs → FORM-R, Male below 10 lakhs → FORM-P

Female above 10 lakhs → FORM-R, Female below 8 lakhs → FORM-Q

Non-residence with income above 15 lakhs → FORM-S

All others should fill the FORM-T

From the options, a residence female with annual income Rs. 9 lakhs will be in others and so will the FORM-T.



Scan for Video solution

**296. (c)**

Palghat is surrounded by the hills which is lower than the hilly terrain to its north and south. Which impact on weather patterns in neighbouring parts of Tamil Nadu and Kerala.

∴ Option (c) is correct.



Scan for Video solution



**297. (b)**

Certain psychiatric illnesses have a genetic basis i.e. depression and schizophrenia, and consequently.  
∴ Option (b) is correct.



Scan for Video solution

**298. (b)**

Before world war 2, Kaliningrad was a German majority population area, but currently it is a Russian territory even it is not close to the Russia. Therefore,  
∴ Option (b) is correct.



Scan for Video solution

**299. (d)**

The dengue cases were increases because municipal authorities have concluded that measures to control the mosquito population have failed in this region. Therefore, the number of people with malarial fever (also contracted from mosquito bites) has increased this year.

∴ Option (d) is correct.



Scan for Video solution

**300. (d)**

In the above statement only, statement (iv) inferred with the passage, therefore, option (d) is correct.



Scan for Video solution

**301. (c)**

From the passage, the main point we can take is that exactitude is not critical in dealing with big data.



Scan for Video solution

**302. (d)**

From the sentences it can be inferred that Rajan had believed that Sajan and he would be working together but Sajan decided to do the project on his own.



Scan for Video solution

**303. (d)**

From the paragraph it can be inferred that ‘No adversity justifies giving up hope’. The spider failed many times but that did not deter it to refrain from making attempts. This event encouraged Bruce not to lose hope and he won many battles thereafter.



Scan for Video solution

**304. (d)**

There were different streams of freedom movement in colonial India, so that we can say the nationalism in India is heterogeneous or they are of different thinking.

∴ Option (d) is correct.



Scan for Video solution

**305. (a)**

“In the legions, military law prevailed and discipline was brutal. Discipline on the battlefield kept units obedient, intact and fighting, even when the odds and conditions were against them” implies “Thorough regimentation was the main reason for the efficiency of the Roman legions even in adverse circumstances”.

∴ Option (a) is correct.



Scan for Video solution



**306. (b)**

In the passage doctors are injecting toxin to horses because they know that horses will develop immunity automatically. They are not giving any immunity to horses.

∴ Option (b) is correct.



Scan for Video solution

**307. (d)**

In the advertisement there is nothing mention about gender so the advertisement is not gender-discriminatory.

**308. (c)**

From the paragraph it can be inferred that use of chemical agents would be undesirable in warfare.



Scan for Video solution



**NOTE:**

**NOTE:**