# ISTA 116: Statistical Foundations for the Information Age

Discrete Random Variables

31 October 2011

#### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

#### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

A single random sample may have more than one characteristic that we can observe (i.e., it may be bi-/multivariate data).

- A single random sample may have more than one characteristic that we can observe (i.e., it may be bi-/multivariate data).
- We can represent each characteristic (e.g., sex, weight, cancer status, etc.) using a **random variable**

- A single random sample may have more than one characteristic that we can observe (i.e., it may be bi-/multivariate data).
- We can represent each characteristic (e.g., sex, weight, cancer status, etc.) using a **random variable**

#### (Definition: Random Variable)

A **random variable** connects each element of the sample space to a value or quantity of interest.

- A single random sample may have more than one characteristic that we can observe (i.e., it may be bi-/multivariate data).
- We can represent each characteristic (e.g., sex, weight, cancer status, etc.) using a **random variable**

#### (Definition: Random Variable)

A **random variable** connects each element of the sample space to a value or quantity of interest.

 $\blacksquare X$ : People  $\mapsto$  Their Sex


- A single random sample may have more than one characteristic that we can observe (i.e., it may be bi-/multivariate data).
- We can represent each characteristic (e.g., sex, weight, cancer status, etc.) using a **random variable**

#### (Definition: Random Variable)


A **random variable** connects each element of the sample space to a value or quantity of interest.

- X: People  $\mapsto$  Their Sex
- $lue{Y}$ : Sequences of coin flips  $\mapsto$  Number of Heads


# Examples of Random Variables



# Examples of Random Variables



# Examples of Random Variables



#### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

# Types of Random Variables

We can classify random variables based on the types of values they take on.

# Types of Random Variables

- We can classify random variables based on the types of values they take on.
- **Discrete** random variables take on discrete values (e.g., categories or integers)

# Types of Random Variables

- We can classify random variables based on the types of values they take on.
- **Discrete** random variables take on discrete values (e.g., categories or integers)
- Continuous random variables take on continuous values (e.g., real numbers).


#### Outline


- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution


When a random variable is discrete, its distribution is characterized by the probabilities assigned to each distinct value.


- When a random variable is discrete, its distribution is characterized by the probabilities assigned to each distinct value.
- These probabilities are determined by the probabilities on the sample space itself

- When a random variable is discrete, its distribution is characterized by the probabilities assigned to each distinct value.
- These probabilities are determined by the probabilities on the sample space itself
- If the sample space is a finite population and we make a simple random draw, then the probability of a value is the proportion of individual outcomes assigned to it.










x	Male	Female
P(X=x)	1/2	1/2



# Properties of Discrete Distributions

Note that each value of a discrete random variable corresponds to an event in the original sample space.

- Note that each value of a discrete random variable corresponds to an event in the original sample space.
- The probability associated with the value is the probability associated with the event.

- Note that each value of a discrete random variable corresponds to an event in the original sample space.
- The probability associated with the value is the probability associated with the event.
- Moreover, every outcome in the sample space is associated with exactly one value of the random variable.

- Note that each value of a discrete random variable corresponds to an event in the original sample space.
- The probability associated with the value is the probability associated with the event.
- Moreover, every outcome in the sample space is associated with exactly one value of the random variable.
- Therefore, the values of a discrete random variable give us a set of **disjoint** events whose **union** is the entire sample space.

■ What must be true of the set of probabilities then?

```
(Properties of Discrete Distributions)
```

■ What must be true of the set of probabilities then?

#### (Properties of Discrete Distributions)

1 For every x in the range of X,  $P(X = x) \ge 0$ .

■ What must be true of the set of probabilities then?

#### (Properties of Discrete Distributions)

- **1** For every x in the range of X,  $P(X = x) \ge 0$ .
- 2

$$\sum_{x \in \mathsf{Range}(X)} P(X = x) = 1$$

#### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

# Visualizing Discrete Distributions




- Very simple distributions can be visualized with a pie chart.
- Can imagine a spinner mechanism that lands on a slice according to its probability.

# Visualizing Discrete Distributions




- Very simple distributions can be visualized with a pie chart.
- Can imagine a spinner mechanism that lands on a slice according to its probability.
- But, like pie charts, this is limited in its ability to convey information.

# The Spike Plot



An alternative is the spike plot

# The Spike Plot



- An alternative is the spike plot
- Like a bar plot, but with probabilities, instead of frequencies or proportions, on the y-axis.

### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

└ The PMF and the CDF

### Probability Mass Function

■ The **distribution** of a random variable is characterized by the set of values and their probabilities.

- The **distribution** of a random variable is characterized by the set of values and their probabilities.
- For finite sets of values, can think of a table.

- The **distribution** of a random variable is characterized by the set of values and their probabilities.
- For finite sets of values, can think of a table.
- One way to use such a table is to start with a value and look up its probability.

- The **distribution** of a random variable is characterized by the set of values and their probabilities.
- For finite sets of values, can think of a table.
- One way to use such a table is to start with a value and look up its probability.
- This process is characterized by the probability mass function, which takes a value and returns its probability.

#### (Definition: The Probability Mass Function)

A discrete random variable, X, can be characterized by its **probability mass function**,  $f_X$ , which takes values and returns probabilities:

$$f_X(x) = P(X = x)$$

#### (Definition: The Probability Mass Function)

A discrete random variable, X, can be characterized by its **probability mass function**,  $f_X$ , which takes values and returns probabilities:

$$f_X(x) = P(X = x)$$

■ In the most general case, we just have to consult a table.

#### (Definition: The Probability Mass Function)

A discrete random variable, X, can be characterized by its **probability mass function**,  $f_X$ , which takes values and returns probabilities:

$$f_X(x) = P(X = x)$$

- In the most general case, we just have to consult a table.
- However, we will see examples later of PMFs that have algebraic expressions.

Discrete Distributions

☐ The PMF and the CDF

#### The Cumulative Distribution Function

Often times we are interested in the probability of falling in some *range* of values.

- Often times we are interested in the probability of falling in some *range* of values.
- For this purpose, we can use the cumulative distribution function (or CDF), which gives the "accumulated probability" up to a particular value.

- Often times we are interested in the probability of falling in some range of values.
- For this purpose, we can use the cumulative distribution function (or CDF), which gives the "accumulated probability" up to a particular value.

#### (Definition: The Cumulative Distribution Function)

A random variable, X, can be characterized by its **cumulative distribution function**,  $F_X$ , which takes values and returns *cumulative* probabilities:

$$F_X(x) = P(X \le x)$$

■ How can we calculate  $F_X(x)$  from the distribution table?

- How can we calculate  $F_X(x)$  from the distribution table?
- How would we calculate P(X > x)?

- How can we calculate  $F_X(x)$  from the distribution table?
- How would we calculate P(X > x)?
- How about  $P(X \ge x)$ ?

- How can we calculate  $F_X(x)$  from the distribution table?
- How would we calculate P(X > x)?
- How about  $P(X \ge x)$ ?
- How would we calculate  $P(a < X \le b)$ ?

- How can we calculate  $F_X(x)$  from the distribution table?
- How would we calculate P(X > x)?
- How about  $P(X \ge x)$ ?
- How would we calculate  $P(a < X \le b)$ ?
- How about  $P(a \le X \le b)$ ?

- How can we calculate  $F_X(x)$  from the distribution table?
- How would we calculate P(X > x)?
- How about  $P(X \ge x)$ ?
- How would we calculate  $P(a < X \le b)$ ?
- How about  $P(a \le X \le b)$ ?
- $P(a \le X < b)$ ?

### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

## Summarizing Random Variables

As with data, it is useful to characterize the center and spread of a probability distribution.

## Summarizing Random Variables

- As with data, it is useful to characterize the center and spread of a probability distribution.
- Most of the measures we've seen can be computed; but the most common are the mean and variance.

### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

☐ The Mean, or "Expected Value"

### Expected Value

The mean of a random variable is also called its expected value. The Mean, or "Expected Value"

## Expected Value

- The mean of a random variable is also called its expected value.
- As with a sample mean, it represents an average over the possible values; but it is weighted by the probabilities.

# Example: Mean Number of Heads

$$\mu_X = E(X) =$$

# Example: Mean Number of Heads

$$\mu_X = E(X) = 0 \times P(X = 0) + 1 \times P(X = 1) + 2 \times P(X = 2)$$

☐ The Mean, or "Expected Value"

# Example: Mean Number of Heads

$$\mu_X = E(X) = 0 \times P(X = 0) + 1 \times P(X = 1)$$
  
  $+2 \times P(X = 2)$   
  $= 0 \times \frac{1}{4} + 1 \times \frac{1}{2} + 2 \times \frac{1}{4}$ 

The Mean, or "Expected Value"

# Example: Mean Number of Heads

$$\mu_X = E(X) = 0 \times P(X = 0) + 1 \times P(X = 1)$$

$$+2 \times P(X = 2)$$

$$= 0 \times \frac{1}{4} + 1 \times \frac{1}{2} + 2 \times \frac{1}{4}$$

$$= \frac{1}{2} + \frac{2}{4} = 1$$

## Expected Value of a Discrete Random Variable

■ In general, we have:

(Definition: Expected Value of a Discrete Random Variable)

$$\mu_X = E(X) = \sum_{x \in \mathsf{Range}(X)} x P(X = x)$$

### Expected Value of a Discrete Random Variable

■ In general, we have:

(Definition: Expected Value of a Discrete Random Variable)

$$\mu_X = E(X) = \sum_{x \in \mathsf{Range}(X)} x P(X = x)$$

If X is a numeric variable with values from 0 to some number n, we have

(Expected Value of Finite, Integer-Valued Random Variable)

$$\mu_X = E(X) = \sum_{x=0}^{n} x P(X = x)$$

### Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

### Variance of a Discrete Random Variable

■ The variance is the expected squared deviation:

(Definition: Variance of a Discrete Random Variable)

$$\sigma_X^2 = E((X-\mu_X)^2) = \sum_{x \in \mathsf{Range}(X)} (x-\mu_X)^2 P(X=x)$$

### Variance of a Discrete Random Variable

■ The variance is the expected squared deviation:

(Definition: Variance of a Discrete Random Variable)

$$\sigma_X^2 = E((X-\mu_X)^2) = \sum_{x \in \mathsf{Range}(X)} (x-\mu_X)^2 P(X=x)$$

If X is a numeric variable with values from 0 to some number n, we have

(Variance of Finite, Integer-Valued Random Variable)

$$\sigma_X^2 = E((X - \mu_X)^2) = \sum_{x=0}^n (x - \mu_x)^2 P(X = x)$$

# Example: Variance of Number of Heads

■ To compute the variance in the number of heads in two coin tosses:

$$\sigma_X^2 \ =$$

# Example: Variance of Number of Heads

■ To compute the variance in the number of heads in two coin tosses:

$$\sigma_X^2 = (0 - \mu_X)^2 \times P(X = 0) + (1 - \mu_X)^2 \times P(X = 1) + (2 - \mu_X)^2 \times P(X = 2)$$

└Variance of a Discrete Random Variable

# Example: Variance of Number of Heads

To compute the variance in the number of heads in two coin tosses:

$$\sigma_X^2 = (0 - \mu_X)^2 \times P(X = 0) + (1 - \mu_X)^2 \times P(X = 1)$$

$$+ (2 - \mu_X)^2 \times P(X = 2)$$

$$= (0 - 1)^2 \times \frac{1}{4} + (1 - 1)^2 \times \frac{1}{2}$$

$$+ (2 - 1)^2 \times \frac{1}{4}$$

# Example: Variance of Number of Heads

■ To compute the variance in the number of heads in two coin tosses:

$$\sigma_X^2 = (0 - \mu_X)^2 \times P(X = 0) + (1 - \mu_X)^2 \times P(X = 1)$$

$$+ (2 - \mu_X)^2 \times P(X = 2)$$

$$= (0 - 1)^2 \times \frac{1}{4} + (1 - 1)^2 \times \frac{1}{2}$$

$$+ (2 - 1)^2 \times \frac{1}{4}$$

$$= \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

└─Variance of a Discrete Random Variable

# Standard Deviation

■ The standard deviation is then \_

## Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

## Named Distributions

- Any discrete set of values, along with associated probabilities, forms a discrete probability distribution
  - as long as \_\_\_\_\_\_
  - and \_\_\_\_\_

## Named Distributions

- Any discrete set of values, along with associated probabilities, forms a discrete probability distribution
  - as long as \_\_\_\_\_\_
  - and \_\_\_\_\_
- However, some sets show up so often, they are given names.

# Named Distributions

- Any discrete set of values, along with associated probabilities, forms a discrete probability distribution
  - as long as \_\_\_\_\_
  - and \_\_\_\_\_
- However, some sets show up so often, they are given names.
- We will look at three of these:
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

# Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

Some Common Discrete Distributions

☐ The Bernoulli Distribution

### The Bernoulli Distribution


■ The simplest possible probability distribution is the **Bernoulli Distribution**, named after Jakob Bernoulli, who is also credited with discovering the constant *e*.

- The simplest possible probability distribution is the **Bernoulli Distribution**, named after Jakob Bernoulli, who is also credited with discovering the constant *e*.
- Any random variable with only two possible outcomes has a Bernoulli distribution with "success" probability p (where we choose one of the outcomes to call a "success").

- The simplest possible probability distribution is the Bernoulli Distribution, named after Jakob Bernoulli, who is also credited with discovering the constant e.
- Any random variable with only two possible outcomes has a Bernoulli distribution with "success" probability p (where we choose one of the outcomes to call a "success").
- Examples:
  - Flipping a single coin
  - Playing the lottery
  - Getting or not getting cancer
  - Winning or losing a baseball game
  - Graduating or not graduating from college
  - etc. etc.



# Example: Playing the Lottery



x	Lose	Win
P(X=x)		

### The Bernoulli Distribution

If a "success" has probability p, then a failure has probability \_\_\_\_\_

- If a "success" has probability p, then a failure has probability \_\_\_\_\_
- Typically, we label the success outcome as "1", and the failure outcome as "0".
- We can then compute a mean and variance.

- If a "success" has probability p, then a failure has probability \_\_\_\_\_
- Typically, we label the success outcome as "1", and the failure outcome as "0".
- We can then compute a mean and variance.
- In terms of the **parameter** *p*, the mean of a Bernoulli distribution is \_\_\_\_\_

- If a "success" has probability *p*, then a failure has probability \_\_\_\_\_
- Typically, we label the success outcome as "1", and the failure outcome as "0".
- We can then compute a mean and variance.
- In terms of the **parameter** *p*, the mean of a Bernoulli distribution is \_\_\_\_\_
- The variance is \_\_\_\_\_

- If a "success" has probability *p*, then a failure has probability \_\_\_\_\_
- Typically, we label the success outcome as "1", and the failure outcome as "0".
- We can then compute a mean and variance.
- In terms of the **parameter** *p*, the mean of a Bernoulli distribution is \_\_\_\_\_
- The variance is \_\_\_\_\_

$$\sigma^2 = (0-p)^2(1-p) + (1-p)^2p$$

- If a "success" has probability p, then a failure has probability \_\_\_\_\_
- Typically, we label the success outcome as "1", and the failure outcome as "0".
- We can then compute a mean and variance.
- In terms of the **parameter** *p*, the mean of a Bernoulli distribution is \_\_\_\_\_\_
- The variance is \_\_\_\_\_\_

$$\sigma^{2} = (0-p)^{2}(1-p) + (1-p)^{2}p$$
$$= p^{2}(1-p) + (1-2p+p^{2})p$$

- If a "success" has probability p, then a failure has probability \_\_\_\_\_
- Typically, we label the success outcome as "1", and the failure outcome as "0".
- We can then compute a mean and variance.
- In terms of the **parameter** *p*, the mean of a Bernoulli distribution is \_\_\_\_\_\_
- The variance is \_\_\_\_\_

$$\sigma^{2} = (0-p)^{2}(1-p) + (1-p)^{2}p$$

$$= p^{2}(1-p) + (1-2p+p^{2})p$$

$$= p^{2} - p^{3} + p - 2p^{2} + p^{3}$$

- If a "success" has probability p, then a failure has probability \_\_\_\_\_
- Typically, we label the success outcome as "1", and the failure outcome as "0".
- We can then compute a mean and variance.
- In terms of the **parameter** *p*, the mean of a Bernoulli distribution is \_\_\_\_\_
- The variance is \_\_\_\_\_

$$\sigma^{2} = (0-p)^{2}(1-p) + (1-p)^{2}p$$

$$= p^{2}(1-p) + (1-2p+p^{2})p$$

$$= p^{2} - p^{3} + p - 2p^{2} + p^{3}$$

$$= p - p^{2} = p(1-p)$$

# Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

Some Common Discrete Distributions

The Discrete Uniform Distribution

## The Discrete Uniform Distribution

Another common distribution is the discrete uniform distribution.

## The Discrete Uniform Distribution

- Another common distribution is the discrete uniform distribution.
- Here, we have n possible outcomes (labeled 1 through n) which are all equally likely.

## The Discrete Uniform Distribution

- Another common distribution is the discrete uniform distribution.
- Here, we have n possible outcomes (labeled 1 through n) which are all equally likely.
- In terms of the **parameter** n, f(k) = ? for k = 1, ..., n
- What are some examples?


## The Discrete Uniform Distribution

- Another common distribution is the discrete uniform distribution.
- Here, we have n possible outcomes (labeled 1 through n) which are all equally likely.
- In terms of the **parameter** n, f(k) = ? for k = 1, ..., n
- What are some examples?
- What is F(k) equal to?
- How would we find the mean and variance?

Some Common Discrete Distributions

☐ The Discrete Uniform Distribution

# Example: Card Suit



# Outline

- 1 Random Variables
  - Types of Random Variables
- 2 Discrete Distributions
  - Visualizing Discrete Distributions
  - The PMF and the CDF
- 3 Summarizing (Numeric) Random Variables
  - The Mean, or "Expected Value"
  - Variance of a Discrete Random Variable
- 4 Some Common Discrete Distributions
  - The Bernoulli Distribution
  - The Discrete Uniform Distribution
  - The Binomial Distribution

### The Binomial Distribution

Many times we have some basic process with two outcomes (i.e., a Bernoulli process), which is repeated some number of times.

- Many times we have some basic process with two outcomes (i.e., a Bernoulli process), which is repeated some number of times.
- We may be interested in the *number* of "successes".

- Many times we have some basic process with two outcomes (i.e., a Bernoulli process), which is repeated some number of times.
- We may be interested in the *number* of "successes".
- If both
- (a) the success probability stays constant
- (b) success events are mutually independent
- then the number of successes has a Binomial Distribution with parameters n (number of trials) and p (individual success probability)

Some Common Discrete Distributions

The Binomial Distribution

#### The Binomial Distribution

■ We've seen an example already. What was it?

- We've seen an example already. What was it?
- The number of heads seen in 2 tosses is Binomial with n=? and p=?

- We've seen an example already. What was it?
- The number of heads seen in 2 tosses is Binomial with n=? and p=?
- Q: How do we find the probability of seeing k heads in 2 tosses?

- We've seen an example already. What was it?
- The number of heads seen in 2 tosses is Binomial with n=? and p=?
- Q: How do we find the probability of seeing k heads in 2 tosses?
- A: Find all the individual sequences with k heads, and add their probabilities (since individual sequences are disjoint events).

- We've seen an example already. What was it?
- The number of heads seen in 2 tosses is Binomial with n=? and p=?
- Q: How do we find the probability of seeing k heads in 2 tosses?
- A: Find all the individual sequences with k heads, and add their probabilities (since individual sequences are disjoint events).
- Q: What is the probability of any particular sequence?

- We've seen an example already. What was it?
- The number of heads seen in 2 tosses is Binomial with n=? and p=?
- Q: How do we find the probability of seeing k heads in 2 tosses?
- A: Find all the individual sequences with k heads, and add their probabilities (since individual sequences are disjoint events).
- Q: What is the probability of any particular sequence?
- A: In two independent tosses, it's
   P(First Outcome) × P(Second Outcome)

### The Binomial Distribution

• Q: In general, for a sequence of n **independent** trials, each with success probability p, how would we find the probability of a specific sequence of successes and failures?

Q: In general, for a sequence of n independent trials, each with success probability p, how would we find the probability of a specific sequence of successes and failures?

$$\begin{array}{rcl} P(\mathsf{Sequence}) &=& P(\mathsf{First\ Outcome} \times P(\mathsf{Second\ Outcome}) \\ && \times \cdots \times P(\mathsf{Last\ Outcome}) \\ &=& \end{array}$$

Q: In general, for a sequence of n independent trials, each with success probability p, how would we find the probability of a specific sequence of successes and failures?

$$\begin{array}{rcl} P(\mathsf{Sequence}) & = & P(\mathsf{First\ Outcome} \times P(\mathsf{Second\ Outcome}) \\ & & \times \cdots \times P(\mathsf{Last\ Outcome}) \\ & = & p^{\#\ \mathsf{successes}} \times (1-p)^{\#\ \mathsf{failures}} \end{array}$$

Q: In general, for a sequence of n independent trials, each with success probability p, how would we find the probability of a specific sequence of successes and failures?

$$\begin{array}{rcl} P(\mathsf{Sequence}) & = & P(\mathsf{First\ Outcome} \times P(\mathsf{Second\ Outcome}) \\ & & \times \cdots \times P(\mathsf{Last\ Outcome}) \\ & = & p^{\#\ \mathsf{successes}} \times (1-p)^{\#\ \mathsf{failures}} \end{array}$$

■ If there were *k* successes, how many failures were there?

Some Common Discrete Distributions

The Binomial Distribution

#### The Binomial Distribution

```
P(\mathsf{Each} \; \mathsf{sequence} \; \mathsf{with} \; k \; \mathsf{successes}) \; = \;
```

$$P(\mathsf{Each} \; \mathsf{sequence} \; \mathsf{with} \; k \; \mathsf{successes}) = p^k \times (1-p)^{n-k}$$

■ Therefore, every sequence of n trials with k successes has probability :

$$P(\mathsf{Each} \; \mathsf{sequence} \; \mathsf{with} \; k \; \mathsf{successes}) = p^k \times (1-p)^{n-k}$$

So, is this the probability of k successes in n (identical, independent) tries?

$$P(\mathsf{Each} \; \mathsf{sequence} \; \mathsf{with} \; k \; \mathsf{successes}) \; = \; p^k \times (1-p)^{n-k}$$

- So, is this the probability of k successes in n (identical, independent) tries?
- **No!** This is the probability of *each sequence!* What else do we need to know?