



Tirzepatide

Information sheet written for the 20 mg vial TrueHold holds in stock. Not a generic internet protocol.


THIS VIAL — STARTING DOSE (read this first)

Tirzepatide · 20 mg lyophilized · mix with 2 mL bacteriostatic water · 10 mg/mL · 1 unit = 100 mcg

Start: draw **25 units** on a U-100 syringe (**0.25 mL = 2.5 mg**). On a 0.5 mL (50-unit) U-100 syringe, draw to the 25 mark.

Route: Subcutaneous (belly or back of upper arm)

Rhythm: Once weekly. Stay on each level at least 4 weeks before going up.



1. What it is — the molecule, how large it is

Tirzepatide is a 39-amino-acid synthetic peptide with a C20 fatty-diacid chain that lets it bind albumin and last about five days. Molecular formula C₂₂₅H₃₄₈N₄₈O₆₈. Molecular weight 4,813.53 daltons — a large peptide hormone analog, not a small-molecule pill. TrueHold stocks a 20 mg lyophilized research vial.

How incretin shots talk to the body

GLP-1 receptor

Appetite down
Stomach slower
Insulin (if glucose high)

GIP receptor

Insulin support
Fat-cell signaling

Glucagon receptor

Energy burn
(retatrutide only)

How large is it? (daltons, log-ish bars — bigger bar = bigger molecule)

GHK-Cu	404 Da
NAD+	663 Da
SS-31	749 Da
Semax	814 Da
MOTS-c	2175 Da
Retatrutide	4731 Da
Tirzepatide	4814 Da

2. How it works in the body

It is a dual incretin agonist: it turns on both the GIP receptor and the GLP-1 receptor. In the body that means stronger glucose-dependent insulin release, less glucagon when glucose is high, slower stomach emptying, and quieter appetite signaling in the brain. That is why people notice less 'food noise' and smaller portions. It is injected under the skin once weekly because the



fatty-acid tail keeps it in circulation.

3. Testing data — animal

Rodent and primate incretin studies showed appetite reduction, improved insulin secretion, and weight loss versus GLP-1 alone. The rodent thyroid C-cell finding that produced the boxed warning on the approved pens was seen with this drug class in rats and mice; it has not been confirmed as a human cancer signal, but it is why people with a personal or family history of medullary thyroid carcinoma or MEN2 are told to stay away from GLP-1/GIP drugs.

4. Testing data — human, what is expected, hopes

Large Phase 3 programs (SURPASS for type 2 diabetes, SURMOUNT for obesity) enrolled thousands of adults. On the 15 mg approved pen, SURMOUNT-1 reported on the order of 15–22% mean body-weight reduction versus placebo depending on dose and diabetes status. The molecule is FDA-approved as Mounjaro (type 2 diabetes, 2022) and Zepbound (chronic weight management 2023; later sleep apnea).

- Less hunger and fewer cravings
- Lower weight and waist when food and protein are tracked
- Better glucose and insulin-response numbers in people who have that goal
- A once-weekly rhythm that is easier to keep than daily shots

5. Have people hurt themselves? Risks

Yes — people have been harmed, mostly from the gastrointestinal effect and from rare but serious class events. Very common: nausea, vomiting, diarrhea, constipation, reduced appetite, stomach pain. Serious and documented: acute pancreatitis (including rare fatal reports in post-marketing systems), gallbladder disease, kidney injury from dehydration after vomiting, delayed stomach emptying that can raise aspiration risk under anesthesia, and low blood sugar when stacked with insulin or sulfonylureas. Stop and get emergency care for severe persistent abdominal pain radiating to the back.

6. FDA — approved or not

FDA-APPROVED as the branded pens Mounjaro and Zepbound — not as this vial. TrueHold's 20 mg lyophilized research vial is not an FDA-approved drug product, not a pen, and not interchangeable with pharmacy Mounjaro/Zepbound. The milligram start on this sheet (2.5 mg weekly) matches the approved starting milligrams so the math is honest; it does not make this vial a licensed medicine.

7. Reconstitution for this exact vial

Check the label: it must say Tirzepatide and 20 mg powder in a standard ~3 mL glass vial. You need bacteriostatic water (BAC), alcohol swabs, and U-100 insulin syringes (0.5 mL / 50-unit for every starting draw on this sheet; 1 mL / 100-unit if a later step exceeds 50 units).

- Wash hands. Wipe the rubber top with an alcohol swab.
- Draw **2 mL** of BAC into a mixing syringe.
- Insert through the stopper. Inject slowly down the inside glass wall — not straight onto the powder.
- Gently roll or swirl until clear. Do not shake.
- Wipe the top again. Write today's date on the vial. Refrigerate 36–46°F. Use within 28 days.
- Mix and store in this vial only. Do not transfer.

8. Loading the 0.5 mL syringe — schedule for this vial

U-100 math is the same on a 0.5 mL or 1 mL syringe: 100 units = 1.00 mL, so 50 units = 0.50 mL (a completely full 0.5 mL syringe). Starting draws on this sheet all fit 0.5 mL. The green row is the start.

When	Units (U-100)	mg	mL	0.5 mL syringe	How often
Weeks 1–4 (start here)	25	2.5	0.25	Draw to 25	Once per week
Weeks 5–8 (if tolerated)	50	5	0.50	Draw to 50	Once per week
Weeks 9–12 (optional)	75	7.5	0.75	Too big — 1 mL syringe or 50+ remainder	Once per week



Week 13+ (optional)	100	10	1.00	Too big — 1 mL syringe or 50+ remainder	Once per week
---------------------	-----	----	------	---	---------------

- Use 2 mL bacteriostatic water only — this vial cannot hold more. Do not pour into another container.
- 25 units = 2.5 mg, which is the same milligram start used on FDA-approved tirzepatide pens. This sheet is for TrueHold's 20 mg research vial, not Mounjaro or Zepbound.
- At 25 units/week one mixed vial lasts about 8 weekly doses.

Fulfillment

- Prep and local delivery: Las Vegas residents only.
- Shipping: dry (lyophilized) vials only — we do not ship mixed liquid.
- Interest orders are confirmed by phone before payment.

Working with the team

- Share a phone number so the team can call.
- Confirm Las Vegas residency for prep and local delivery.
- Complete required documentation with the team.
- If anything on this sheet disagrees with your vial label or COA, stop and call before mixing.

Sources

- FDA Zepbound / Mounjaro prescribing information (Lilly): <https://pi.lilly.com/us/zepbound-uspi.pdf>
- PubChem tirzepatide: MW 4813.53 Da, 39 amino acids, C225H348N48O68
- TrueHold locked 20 mg vial protocol: 2 mL BAC, 25/50/75/100 units weekly
- TrueHold Standard Starting Protocol locked to the 20 mg vial (2 mL BAC, 25/50/75/100 units weekly).

Protocol note

TrueHold writes these sheets for the milligram vials we hold. Mix volume and unit marks were double-checked so milligrams, milliliters, and U-100 units agree. This is educational research information, not a prescription and not medical advice. The Telegram bot does not dose in chat — read this PDF. For an individualized plan, use /schedule. Research use only.

