



KLOW

Information sheet written for the 80 mg vial TrueHold holds in stock. Not a generic internet protocol.


THIS VIAL — STARTING DOSE (read this first)

KLOW · 80 mg lyophilized · mix with 2 mL bacteriostatic water · 40 mg/mL · 1 unit = 400 mcg

Start: draw **10 units** on a U-100 syringe (**0.10 mL = 4 mg**). On a 0.5 mL (50-unit) U-100 syringe, draw to the 10 mark.


Route: Subcutaneous (belly or love-handle). Faint blue-green tint from GHK-Cu is expected.

Rhythm: Once daily, 5 days on / 2 days off, for 8–12 weeks, then a 4-week break.



1. What it is — the molecule, how large it is

KLOW is a four-peptide regenerative blend totaling 80 mg in TrueHold's vial. US research catalogs use this split: GHK-Cu 50 mg (copper tripeptide, ~404 Da) + BPC-157 10 mg (15 amino acids, gastric-juice fragment) + TB-500 10 mg (43 amino-acid thymosin-β4 fragment) + KPV 10 mg (Lys-Pro-Val, the C-terminal of α-MSH). Confirm that split on your COA before mixing.



2. How it works in the body

The four pieces hit different repair stories at once: GHK-Cu on extracellular-matrix genes and copper-dependent enzymes; BPC-157 on nitric-oxide / growth-factor signaling in tissue models; TB-500 on actin and cell migration; KPV on NF-κB inflammatory signaling. That is why it is sold as one 80 mg vial instead of four bottles. It is not a GLP-1 drug and not a steroid.

3. Testing data — animal

Each component has animal data (wound, tendon, gut, inflammation models). The four-peptide combination as KLOW does not have a published multi-center animal program of its own — you are stacking four literatures.

4. Testing data — human, what is expected, hopes



There is no Phase 3 trial of KLOW. GHK-Cu has topical cosmetic human use. BPC-157, TB-500, and KPV do not have FDA-approved human drugs. The 10-unit draw on this sheet is the handling math that puts each component into its usual research milligram/microgram window in one shot.

- Tissue comfort and recovery from training
- Skin and connective-tissue notes
- A quieter inflammatory background (KPV / BPC themes)
- One daily draw instead of four separate peptides

5. Have people hurt themselves? Risks

This blend has not been safety-tested as a licensed combination. Injection-site redness is common. GHK-Cu can tinge the liquid blue-green. Copper load matters if you have Wilson disease. BPC-157 has been flagged in US compounding policy debates — quality and contamination risk on gray-market vials is real. People have reported dizziness or water-retention anecdotes on TB-500-type fragments; that is not a clean trial database. Stop for hives, breathing trouble, severe swelling, or yellowing skin.

6. FDA — approved or not

NOT FDA-APPROVED. None of the four peptides is an approved injectable drug in this blend. This is a research 80 mg vial, not a wound-healing medicine.

7. Reconstitution for this exact vial

Check the label: it must say KLOW and 80 mg KLOW blend powder in a standard ~3 mL glass vial. You need bacteriostatic water (BAC), alcohol swabs, and U-100 insulin syringes (0.5 mL / 50-unit for every starting draw on this sheet; 1 mL / 100-unit if a later step exceeds 50 units).

- Wash hands. Wipe the rubber top with an alcohol swab.
- Draw **2 mL** of BAC into a mixing syringe.
- Insert through the stopper. Inject slowly down the inside glass wall — not straight onto the powder.
- Gently roll or swirl until clear. Do not shake.
- Wipe the top again. Write today's date on the vial. Refrigerate 36–46°F. Use within 28 days.
- Mix and store in this vial only. Do not transfer.

8. Loading the 0.5 mL syringe — schedule for this vial

U-100 math is the same on a 0.5 mL or 1 mL syringe: 100 units = 1.00 mL, so 50 units = 0.50 mL (a completely full 0.5 mL syringe). Starting draws on this sheet all fit 0.5 mL. The green row is the start.

When	Units (U-100)	mg	mL	0.5 mL syringe	How often
Weeks 1–4 (start here)	10	4	0.10	Draw to 10	Once daily, 5 days on / 2 days off
Weeks 5–12 (if tolerated)	10	4	0.10	Draw to 10	Once daily, 5 days on / 2 days off (same draw; do not double)

- 80 mg in 2 mL → 40 mg/mL total blend. 10 units = 0.10 mL = 4.0 mg total.
- Standard 80 mg KLOW split used in US research catalogs: GHK-Cu 50 mg + BPC-157 10 mg + TB-500 10 mg + KPV 10 mg. A 10-unit draw then delivers about 2.5 mg GHK-Cu and 0.50 mg of each of the other three — each piece sits in its usual research window at once.
- If your vial's COA lists a different milligram split, stop and call the team before mixing.
- At 10 units five days per week one mixed vial lasts 4 weeks.

Fulfillment

- Prep and local delivery: Las Vegas residents only.
- Shipping: dry (lyophilized) vials only — we do not ship mixed liquid.
- Interest orders are confirmed by phone before payment.

Working with the team



- Share a phone number so the team can call.
- Confirm Las Vegas residency for prep and local delivery.
- Complete required documentation with the team.
- If anything on this sheet disagrees with your vial label or COA, stop and call before mixing.

Sources

- Standard 80 mg KLOW catalog split: GHK-Cu 50 + BPC-157 10 + TB-500 10 + KPV 10
- 2 mL BAC → 40 mg/mL; 10 units = 4 mg total (2.5 mg GHK-Cu + 0.5 mg × 3)
- 80 mg TrueHold vial / 2 mL BAC. 10 units = 4 mg total blend, the documented 80 mg KLOW handling draw.

Protocol note

TrueHold writes these sheets for the milligram vials we hold. Mix volume and unit marks were double-checked so milligrams, milliliters, and U-100 units agree. This is educational research information, not a prescription and not medical advice. The Telegram bot does not dose in chat — read this PDF. For an individualized plan, use /schedule. Research use only.

