



Tirzepatide

Locked peptide information sheet

Short Description

Tirzepatide is a dual incretin agonist that activates GIP and GLP-1 receptor pathways. It is widely discussed in metabolic-health, appetite-control, and body-composition contexts.

How It Works

Tirzepatide works through two incretin pathways: GIP and GLP-1. These pathways are associated with appetite signaling, delayed gastric emptying, insulin-response biology, and metabolic regulation.

Key Research Themes

- Appetite control
- Hunger and cravings
- Body-composition support
- Blood-sugar and insulin-response research
- Long-term lifestyle consistency

Who Commonly Asks About It

- People focused on weight loss
- People struggling with appetite or food noise
- People tracking metabolic health
- People building a lifestyle plan around nutrition and movement



What To Track

- Weight
- Waist
- Hunger and cravings
- Energy
- Digestion and comfort
- Exercise consistency

TrueHold Standard Starting Protocol

Written for the vial in your TrueHold order. Follow the unit numbers on your syringe — no calculations needed.

Check your label

- Product: Tirzepatide
- Should say: 20 mg (powder in a standard ~3 mL glass vial)

What you need

- Bacteriostatic water (BAC)
- Alcohol swabs
- Insulin syringes — U-100, 1 mL (100-unit) size

Step-by-step: mix your vial

- Wash hands. Wipe the rubber top with an alcohol swab.
- Draw 2 mL of BAC into a syringe.
- Insert needle through the vial top. Inject slowly down the inside glass wall — not onto the powder.
- Gently roll or swirl until clear. Do not shake.
- Wipe the top again. Write today's date on the vial.
- Refrigerate (36–46°F). Use within 28 days.

Important: Use 2 mL only — do not add more liquid; this vial cannot hold it. Do not pour into another container.

Your weekly schedule (syringe units)

Use a 1 mL / 100-unit syringe. Draw to the number below, then inject under the skin once that week (belly or back of upper arm).

When	Draw to this number	How often
Weeks 1–4 (start here)	25	Once per week
Weeks 5–8 (if tolerated)	50	Once per week



Weeks 9–12 (optional)	75	Once per week
Week 13+ (optional)	100	Once per week

Stay on each level at least 4 weeks before going up. If side effects are rough, stay at your current number longer.

How long one mixed vial lasts

- At 25 units/week → about 8 weekly doses from one mixed vial
- Higher numbers use more per week — plan refills accordingly

Protocol Note

TrueHold shares standard starting protocols for general education only. For individualized plans, use /schedule to book with the team.

