



Semax

Information sheet written for the 10 mg vial TrueHold holds in stock. Not a generic internet protocol.


THIS VIAL — STARTING DOSE (read this first)

Semax · 10 mg lyophilized · mix with 2 mL bacteriostatic water · 5 mg/mL · 1 unit = 50 mcg

Start: draw **6 units** on a U-100 syringe (**0.06 mL = 0.3 mg**). On a 0.5 mL (50-unit) U-100 syringe, draw to the 6 mark.

Route: Intranasal drops (clinical Semax is a nasal medicine in Russia). Draw, then drip — do not inject this starting protocol.

Rhythm: Once or twice daily for 10–14 days, then take a break of at least 7 days.



1. What it is — the molecule, how large it is

Semax is a 7-amino-acid neuropeptide: Met-Glu-His-Phe-Pro-Gly-Pro (MEHFPGP). Formula C₃₇H₅₁N₉O₁₀S. Molecular weight 813.93 daltons — small next to tirzepatide. It is a stabilized ACTH(4-10) analog with a Pro-Gly-Pro tail so enzymes do not chew it as fast. It does not tell the adrenal glands to make cortisol the way full ACTH does. TrueHold stocks a 10 mg lyophilized vial.

How large is it? (daltons, log-ish bars — bigger bar = bigger molecule)

GHK-Cu	404 Da
NAD+	663 Da
SS-31	749 Da
Semax	814 Da
MOTS-c	2175 Da
Retatrutide	4731 Da
Tirzepatide	4814 Da

2. How it works in the body

In the brain it is discussed for BDNF-related signaling, melanocortin receptors, and enkephalinase effects. People notice focus and stress tolerance, not appetite collapse. Russian medicine uses it as nasal drops so it can reach the brain along olfactory pathways. That is why this TrueHold sheet reconstitutes the 10 mg vial and then drips — it does not start you on an injection.

3. Testing data — animal

Rodent work reports neuroprotection after ischemia, improved learning tasks, and BDNF changes. Those models are why it was developed for stroke and cognition in Russia.

4. Testing data — human, what is expected, hopes



Semax has been a registered medicine in Russia since the 1990s (0.1% cognitive drops; 1% hospital stroke drops) and is on the Russian vital-drugs list. Published English-language trials are smaller than Western Phase 3 programs. There are no FDA trials. Cognitive protocols commonly sit around 200–900 mcg/day nasally for 10–30 days.

- Sharper focus and mental endurance
- Stress tolerance during heavy cognitive load
- Neuroprotective research interest after brain insult (that is a medical use in Russia, not a TrueHold claim)

5. Have people hurt themselves? Risks

Published Russian use mostly reports mild nasal irritation, occasional headache, and sleep disruption if taken late. Serious events are not a feature of the English-language literature, but Western long-term RCT safety does not exist. Do not treat this as a stroke medicine. Do not drive the 1% hospital dose at home. If you get a severe headache, nosebleed that will not stop, or mood crash, stop and call the team.

6. FDA — approved or not

NOT FDA-APPROVED in the United States. Approved as a nasal medicine in Russia/Ukraine. TrueHold's 10 mg lyophilized vial is a research formula, not a US drug.

7. Reconstitution for this exact vial

Check the label: it must say Semax and 10 mg powder in a standard ~3 mL glass vial. You need bacteriostatic water (BAC), alcohol swabs, and U-100 insulin syringes (0.5 mL / 50-unit for every starting draw on this sheet; 1 mL / 100-unit if a later step exceeds 50 units).

- Wash hands. Wipe the rubber top with an alcohol swab.
- Draw **2 mL** of BAC into a mixing syringe.
- Insert through the stopper. Inject slowly down the inside glass wall — not straight onto the powder.
- Gently roll or swirl until clear. Do not shake.
- Wipe the top again. Write today's date on the vial. Refrigerate 36–46°F. Use within 28 days.
- Mix and store in this vial only. Do not transfer.

8. Loading the 0.5 mL syringe — schedule for this vial

U-100 math is the same on a 0.5 mL or 1 mL syringe: 100 units = 1.00 mL, so 50 units = 0.50 mL (a completely full 0.5 mL syringe). Starting draws on this sheet all fit 0.5 mL. The green row is the start.

When	Units (U-100)	mg	mL	0.5 mL syringe	How often
Days 1–10 (start here)	6	0.3	0.06	Draw to 6	Once in the morning
If tolerated, still in the 10-day block	6	0.3	0.06	Draw to 6	Morning and early afternoon (not late evening)
Optional upper end of the cognitive range	12	0.6	0.12	Draw to 12	Once in the morning

- 10 mg in 2 mL → 5 mg/mL. 1 unit = 50 mcg. 6 units = 300 mcg, inside the Russian 0.1% cognitive range (about 200–900 mcg/day).
- Drip half the draw into each nostril while seated. Do not inject Semax on this sheet — published human use is nasal.
- At 6 units once daily, one mixed vial lasts about 33 days; TrueHold still cycles 10–14 days on, then a break.

Fulfillment

- Prep and local delivery: Las Vegas residents only.
- Shipping: dry (lyophilized) vials only — we do not ship mixed liquid.
- Interest orders are confirmed by phone before payment.

Working with the team

- Share a phone number so the team can call.
- Confirm Las Vegas residency for prep and local delivery.



- Complete required documentation with the team.
- If anything on this sheet disagrees with your vial label or COA, stop and call before mixing.

Sources

- PubChem / Wikipedia Semax: C37H51N9O10S, 813.93 g/mol, sequence MEHFPGP
- Russian 0.1% cognitive range ~200–900 mcg/day mapped to 6 units (300 mcg) on this 10 mg / 2 mL vial
- Russian 0.1% Semax cognitive range mapped onto TrueHold's 10 mg vial with 2 mL BAC so unit marks are round.

Protocol note

TrueHold writes these sheets for the milligram vials we hold. Mix volume and unit marks were double-checked so milligrams, milliliters, and U-100 units agree. This is educational research information, not a prescription and not medical advice. The Telegram bot does not dose in chat — read this PDF. For an individualized plan, use /schedule. Research use only.

