



**PART 1**

# **The Tate Pairing**

**KU LEUVEN**

## The Weil Pairing

The iconic pairing, that always helps us when we are dealing with  $m$ -torsion  $E[m]$ ,

$$e_m : E[m] \times E[m] \rightarrow \mu_m,$$

where  $\mu_m = \{ \zeta \in \overline{\mathbb{F}_q}^* \mid \zeta^m = 1 \}$ . The  $m$ -Weil pairing is *bilinear*, *non-degenerate*, and *alternating*.

**Destructive** use: the MOV attack on elliptic curves, reducing elliptic-curve DLOG to  $\mathbb{F}_q^*$ -DLOG.

**Constructive** use\* in isogeny-based cryptography: decomposing  $K \in E[m]$  into  $K = [a]P + [b]Q$  for a given basis  $P, Q$  for  $E[m]$ .



Many other applications are out there in identity-



**Definition 3.** A *pairing* on an elliptic curve  $E$  is a bilinear map  $e : A \times B \rightarrow \mathbb{F}_q^*$ , where  $A$  and  $B$  are subgroups of  $E$ . We say  $e$  is *non-degenerate* when for every  $a \in A$ , there is at least one  $b \in B$  such that  $e(a, b) \neq 1$ , and vice versa. We say  $e$  is *alternating* when  $A = B$  and for every  $a \in A$  we have  $e(a, a) = 1$ .



**Lefty and His Gang (1987)**





**Lefty and His Gang (1987)**