

PART 1  
The Tate Pairing

**Corollary.** Let  $2^n$  divide  $p + 1$  for some prime  $p$ , and take a supersingular Montgomery curve over  $\mathbb{F}_{p^2}$  given by

$$E_A : y^2 = x^3 + Ax^2 + x, \quad \text{with } A \in \mathbb{F}_{p^2}.$$

If a point  $P = (x_P, y_P) \in E(\mathbb{F}_{p^2})$  has  $x_P$  non-square, then  $P$  has order divisible by  $2^n$

**Theorem 3.** For an elliptic curve  $E : y^2 = (x - r_1)(x - r_2)(x - r_3)$  with  $r_i \in \mathbb{F}_{p^2}$ , we have

$$P \in [2]E(\mathbb{F}_{p^2}) \quad \text{if and only if} \quad (x_P - r_1), (x_P - r_2), \text{ and } (x_P - r_3) \text{ are squares.}$$

Consequently,  $P \in E(\mathbb{F}_{p^2}) \setminus [2]E(\mathbb{F}_{p^2})$  if any  $(x_P - r_i)$  is a non-square.

Tate  
Pairing?

PART 1  
The Tate Pairing

**Corollary.** Let  $2^n$  divide  $p + 1$  for some prime  $p$ , and take a supersingular Montgomery curve over  $\mathbb{F}_{p^2}$  given by

$$E_A : y^2 = x^3 + Ax^2 + x, \quad \text{with } A \in \mathbb{F}_{p^2}.$$

If a point  $P = (x_P, y_P) \in E(\mathbb{F}_{p^2})$  has  $x_P$  non-square, then  $P$  has order divisible by  $2^n$

---

**Theorem 3.** For an elliptic curve  $E : y^2 = (x - r_1)(x - r_2)(x - r_3)$  with  $r_i \in \mathbb{F}_{p^2}$ , we have

$$P \in [2]E(\mathbb{F}_{p^2}) \quad \text{if and only if} \quad (x_P - r_1), (x_P - r_2), \text{ and } (x_P - r_3) \text{ are squares.}$$

Consequently,  $P \in E(\mathbb{F}_{p^2}) \setminus [2]E(\mathbb{F}_{p^2})$  if any  $(x_P - r_i)$  is a non-square.

Tate  
Pairing???