

Tutorial 3

Presented by Jiawei Xu

jiaweixu1@link.cuhk.edu.cn

The Chinese University of Hong Kong, Shenzhen

September 25, 2024

Methods of Proofs

There are endless possibilities for how to construct mathematical proof.

Most common ones used in practice:

- Proof by direct construction $P(a) \implies \exists x P(x)$
- Proof by contraposition $(\neg B \rightarrow \neg A) \equiv (A \rightarrow B)$
- Proof by contradiction $\neg\neg A \equiv A$
- Proof by cases $(A \rightarrow B) \wedge (\neg A \rightarrow B) \equiv B$

Direct Proof

Direct Proof

In a direct proof, we assume the antecedent is true, then use rules of inference, laws, definitions and/or previously proven theorems to show the consequent is true.

Prove “The sum of two even integers is even”.

Direct Proof

Direct Proof

In a direct proof, we assume the antecedent is true, then use rules of inference, laws, definitions and/or previously proven theorems to show the consequent is true.

Prove “The sum of two even integers is even”.

Let $n = 2r$, where $r \in \mathbb{Z}$, $m = 2s$, where $s \in \mathbb{Z}$, then we can get
 $n + m = 2r + 2s = 2(r + s)$, where $(r + s) \in \mathbb{Z}$.

Exercise

Prove “if n is an odd integer, then n^2 is odd”.

Exercise

Prove “if n is an odd integer, then n^2 is odd”.

Let $n = 2r + 1$, where $r \in \mathbb{Z}$, then we can get

$$n^2 = (2r + 1)^2 = 4 + 4r + 1 = 2(2 + 2r) + 1.$$

Proof by Contraposition

Contraposition

A type of indirect proof that makes use of the fact that $p \rightarrow q$ is logically equivalent to its contrapositive $\neg q \rightarrow \neg p$. So we assume $\neg q$ is true, then work to prove $\neg p$ is true.

Prove “if n is an integer and $3n + 2$ is odd, then n is odd”.

Proof by Contraposition

Contraposition

A type of indirect proof that makes use of the fact that $p \rightarrow q$ is logically equivalent to its contrapositive $\neg q \rightarrow \neg p$. So we assume $\neg q$ is true, then work to prove $\neg p$ is true.

Prove “if n is an integer and $3n + 2$ is odd, then n is odd”.

Let $p : 3n + 2$ is odd, $q : n$ is odd. Then we can prove $p \rightarrow q$ by showing $\neg q \rightarrow \neg p$.

Let $n = 2k$, where $k \in \mathbb{Z}$, then $3n + 2 = 2 \cdot (3k + 1)$, where $(3k + 1) \in \mathbb{Z}$.

Exercise

Prove “if n is an integer and $3n + 2$ is even, then n is even”.

Exercise

Prove “if n is an integer and $3n + 2$ is even, then n is even”.

Let $p : 3n + 2$ is even, $q : n$ is even. Then we can prove $p \rightarrow q$ by showing $\neg q \rightarrow \neg p$.

Let $n = 2k + 1$, where $k \in \mathbb{Z}$, then

$$3n + 2 = 6k + 5 = 2(3k + 2) + 1, \text{ where } (3k + 2) \in \mathbb{Z}.$$

Proof by Contradiction

Contradiction

In this method of proof, we assume a proposition is not true, then through that premise and logic find a contradiction that shows our original premise must be incorrect, and therefore, the proposition was true.

Prove “if n is an integer and $3n + 2$ is even, then n is even”.

Proof by Contradiction

Contradiction

In this method of proof, we assume a proposition is not true, then through that premise and logic find a contradiction that shows our original premise must be incorrect, and therefore, the proposition was true.

Prove “if n is an integer and $3n + 2$ is even, then n is even”.

Let $p : 3n + 2$ is even, $q : n$ is even, then we can prove $p \rightarrow q$ by showing $\neg(p \rightarrow q) \equiv \neg(\neg p \vee q) \equiv p \wedge \neg q$ is incorrect.

Assume $3n + 2$ is even, and n is odd, then we can get $3n$ is even, and $3n - n$ is odd. However, $3n - n = 2n$ is even, which is a contradiction.

Exercise

Prove by contradiction that “ $\sqrt{3} + \sqrt{2}$ is irrational”.

Exercise

Prove by contradiction that “ $\sqrt{3} + \sqrt{2}$ is irrational”.

Let p be the statement: $\sqrt{3} + \sqrt{2}$ is irrational. We will prove p by showing that $\neg p$ leads to a contradiction.

Assume that $\sqrt{3} + \sqrt{2}$ is rational. Then, $(\sqrt{3} + \sqrt{2})^2$ must also be rational.

Consider $(\sqrt{3} + \sqrt{2})^2 = 3 + 2\sqrt{3}\sqrt{2} + 2 = 5 + 2\sqrt{6}$.

For $\neg p$ to be incorrect, we need to show that $\sqrt{6}$ is irrational, which would contradict the initial assumption that $\sqrt{3} + \sqrt{2}$ is rational.

Proof by Cases

Proof by Cases

In a proof by cases, we must cover all possible cases that arise and cover all of the domain.

Prove “if n is an integer, then $n \leq n^2$ ”.

Proof by Cases

Proof by Cases

In a proof by cases, we must cover all possible cases that arise and cover all of the domain.

Prove “if n is an integer, then $n \leq n^2$ ”.

Case 1: $n \leq -1$. We can get $n^2 > 0 > n$.

Case 2: $n = 0$. We can get $n^2 = 0 = n$.

Case 3: $n \geq 1$. We can get $n \cdot n \geq 1 \cdot n \implies n^2 \geq n$.

Exercise

Prove “if n is an integers, $n^2 + 3n + 2$ is even”.

Exercise

Prove “if n is an integers, $n^2 + 3n + 2$ is even”.

Case 1: n is even. We can write $n = 2k$, where $k \in \mathbb{Z}$. Then we have

$$n^2 + 3n + 2 = 4k^2 + 6k + 2 = 2(2k^2 + 3k + 1).$$

Case 2: n is odd. We can write $n = 2k + 1$, where $k \in \mathbb{Z}$. Then we have

$$n^2 + 3n + 2 = 4k^2 + 10k + 6 = 2(2k^2 + 5k + 3).$$

Existence Proofs

Definition

Many theorems are assertions that objects of a particular type exist. A theorem of this type is a proposition of the form $\exists xP(x)$, where P is a predicate. A proof of a proposition of the form $\exists xP(x)$ is called an **existence proof**.

Methods

- 1 **Constructive Proof**: sometimes an existence proof of $\exists xP(x)$ can be given by finding an element a , called a witness, such that $P(a)$ is true.
- 2 **Nonconstructive Proof**; we do not find an element a such that $P(a)$ is true, but rather directly prove $\exists xP(x)$ is true in some other way.

Constructive Proof

Show that there is a positive integer that can be written as the sum of cubes of positive integers in two different ways.

Constructive Proof

Show that there is a positive integer that can be written as the sum of cubes of positive integers in two different ways.

$$1729 = 10^3 + 9^3 = 12^3 + 1^3$$

Nonconstructive Proof

Proof that there exists a solution to the equation $x^2 - 2x + 1 = 0$.

Nonconstructive Proof

Proof that there exists a solution to the equation $x^2 - 2x + 1 = 0$.

$$x^2 - 2x + 1 = (x - 1)^2 = 0 \implies x = \pm 1$$

Uniqueness Proof

Definition

Some theorems assert there is **exactly one** element with a particular property. To prove a statement of this type we must show that an element with this property exists and that no other element has this property.

Method

Existence: We show that an element x with the desired property exists.

Uniqueness: We show that if x and y both have the desired property, then $x = y$.

Example

Prove “If a and b are real numbers, and $a \neq 0$, then there is a unique real number r such that $a \cdot r + b = 0$ ”.

Example

Prove “If a and b are real numbers, and $a \neq 0$, then there is a unique real number r such that $a \cdot r + b = 0$ ”.

Existence: We show an element x with the property.

Let $r = \frac{-b}{a}$, then r is a solution to $a \cdot r + b = 0$.

Uniqueness: We show $x \neq y$, and y have the property.

Suppose s is a real number, such that $a \cdot s + b = 0$, then

$a \cdot r + b = a \cdot s + b \implies r = s$. Therefore, $x \neq y$ is not true.

Exercise

Prove that there is no positive integers n such that $n^2 + n^3 = 100$

Exercise

Prove that there is no positive integers n such that $n^2 + n^3 = 100$

$\because n^3 > 100$ for all $n > 4$.

\therefore we only need to show that $n \in \{1, 2, 3, 4\}$ do not satisfy $n^2 + n^3 = 100$.

Exercise

Prove that between every two rational numbers there is an irrational number.

Exercise

Prove that between every two rational numbers there is an irrational number.

We can assume that the given rational numbers are a and b , where $a < b$.

Then we know $a + \frac{b-a}{2}$ is rational and in between a and b .

Thus, $x = a + \frac{b-a}{2} \cdot \frac{\sqrt{2}}{2}$ is also in between a and b because $0 < \frac{\sqrt{2}}{2} < 1$.

Since $\sqrt{2}$ is irrational, x is also irrational.

Exercise

Prove that $k(k+1)(k+2)$ is always divisible by 6.

Exercise

Prove that $k(k+1)(k+2)$ is always divisible by 6.

We notice that it is enough to prove that it is always divisible by 3 and 2.

Case 1: k is even. If k is even, then $k+1$ is odd and $k+2$ is even.

Therefore, either k or $k+2$ is divisible by 2, and k , $k+1$, and $k+2$ are consecutive integers, so one of them is divisible by 3.

Case 2: k is odd. If k is odd, then $k+1$ is even and $k+2$ is odd.

Therefore, $k+1$ is divisible by 2, and k , $k+1$, and $k+2$ are consecutive integers, so one of them is divisible by 3.

Exercise

Shorter proof without cases:

Numbers $k, k + 1, k + 2$ are 3 sequential integers, among them there must be at least one even and at least one divisible by 3.

Conclusion

- Four methods for mathematical proof.
- Two methods for existence proof.
- Two steps in the uniqueness proof.

Thank You

Thank you for your attention!