

Tutorial 7: Max-Flow and Min-Cut Problems

Kangran ZHAO


School of Data Science (SDS)
The Chinese University of Hong Kong, Shenzhen

June 26, 2025


Maximum Flow Problem

The maximum flow problem can be described as follows:

- ▶ Given a directed, weighted graph $G = (V, E)$ and a pair of nodes s and t (V is the set of nodes, E is the set of edges)
- ▶ One can think this as a traffic network
- ▶ There is an edge capacity w_{ij} on each edge
- ▶ Question: What is the largest amount of flow one can send from s to t , subject to the capacity constraints?




A Concrete Example



One Transformation

Assume there is an imaginary node o , with edges (o, s) and (t, o) . There is no capacity constraint on those two edges

- The problem becomes a closed system. One wants to maximize the flow from o to s , which we denote by Δ .



LP Formulation

Using this transformation, we can write down the LP formulation.
Let x_{ij} denote the amount of flow across edge (i, j) .

$$\begin{aligned} & \text{maximize}_{\mathbf{x}, \Delta} && \Delta \\ & \text{subject to} && \sum_{j:(j,i) \in E} x_{ji} - \sum_{j:(i,j) \in E} x_{ij} = 0, && \forall i \neq s, t \\ & && \sum_{j:(j,s) \in E} x_{js} - \sum_{j:(s,j) \in E} x_{sj} + \Delta = 0 \\ & && \sum_{j:(j,t) \in E} x_{jt} - \sum_{j:(t,j) \in E} x_{tj} - \Delta = 0 \\ & && x_{ij} \leq w_{ij}, && \forall (i, j) \in E \\ & && x_{ij} \geq 0, && \forall (i, j) \in E \end{aligned}$$

- ▶ The first constraint is the flow balancing constraints for all nodes other than s and t
- ▶ The second (third, resp.) constraint is the flow balancing constraints for node s (t , resp.)

Dual of the Maximum Flow Problem

We construct the dual problem:

$$\begin{array}{ll}\text{minimize} & \sum_{(i,j) \in E} w_{ij} z_{ij} \\ \text{subject to} & z_{ij} \geq y_i - y_j, \quad \forall (i,j) \in E \\ & y_s - y_t = 1 \\ & z_{ij} \geq 0\end{array}$$

What does the dual problem mean?

First assume all y 's are 0 or 1. Then

- ▶ We assign a label (0 or 1) to each node, 1 to s and 0 to t .
- ▶ If i has a larger label than j for $(i,j) \in E$, there is a cost w_{ij} .

Interpretation of the Dual

The dual problem is equivalent to finding a subset S of vertices containing s but not t , that minimizes the weight of the cut, i.e.

$$\sum_{i \in S, j \notin S} w_{ij}$$

- This is called the min-cut problem

