



PHY 1001: Mechanics

Tutorial Session 10

T-05: Apr - 17 - 2024, Wednesday, 19:00~19:50

T-12: Apr - 17 - 2024, Wednesday, 20:00~20:50

Problem 1



香港中文大學(深圳)理工学院
School of Science and Engineering

* Buoyancy

A rock is suspended by a light string. When the rock is in air, the tension in the string is **39.2 N**. When the rock is totally immersed in water, the tension is **28.4 N**. When the rock is totally immersed in an unknown liquid, the tension is **18.6 N**. What is the density of the unknown liquid?

Solution:

Identify:

The vertical forces on the rock sum to zero. The buoyant force equals the weight of liquid displaced by the rock. $V = \frac{4}{3}\pi R^3$

Set Up:

The density of water is **$1.00 \times 10^3 \text{ kg/m}^3$** .

Problem 1



香港中文大學(深圳)理工学院
School of Science and Engineering

Solution:

Execute:

- (1) Rock displaces a volume of water whose weight is $39.2 \text{ N} - 28.4 \text{ N} = 10.8 \text{ N}$.
- (2) The mass of this much water is thus $10.8 \text{ N} / (9.80 \text{ m/s}^2) = 1.102 \text{ kg}$ and its volume, equal to the rock's volume, is $1.102 \text{ kg} / 1.00 \times 10^3 \text{ kg/m}^3 = 1.102 \times 10^{-3} \text{ m}^3$.
- (3) The weight of unknown liquid displaced is $39.2 \text{ N} - 18.6 \text{ N} = 20.6 \text{ N}$, and its mass is $20.6 \text{ N} / (9.80 \text{ m/s}^2) = 2.102 \text{ kg}$.
- (4) The liquid's density is thus $2.102 \text{ kg} / 1.102 \times 10^{-3} \text{ m}^3 = 1.91 \times 10^3 \text{ kg/m}^3$.

Evaluate:

The density of the unknown liquid is roughly twice the density of water.

QED.

Problem 2




香港中文大學(深圳)理工学院
School of Science and Engineering

* (Halliday C14-P2)

In the figure below, the fresh water behind a reservoir dam has depth $D=12$ m. A horizontal pipe 4.0 cm in diameter passes through the dam at depth $d=6.0$ m. A plug secures the pipe opening.

Questions:

- (a) Find the magnitude of the frictional force between plug and pipe wall.
- (b) The plug is removed. What water volume exits the pipe in 3.0 h?



Problem 2



Solution (a):

(a) The magnitude of the friction force must balance the gauge pressure force, thus it is

$$f = \Delta p A = \rho g d (\pi r^2) = 74 \text{ N}.$$


Solution (b):

(1) The speed of water flowing out of the hole is $v = \sqrt{2gd}$, which can be derived from the Bernoulli's equation .

(2) Thus, the volume of water flowing out of the pipe in $t = 3.0\text{h}$ rounded off to 2 significant figures is

$$V = Avt = 1.5 \times 10^2 \text{ m}^3.$$

QED.



Problem 3



香港中文大學(深圳)理工学院
School of Science and Engineering

* (Halliday C14-P3)


In the figure below, water stands at depth $D = 30.0 \text{ m}$ behind the vertical upstream face of a dam of width $W = 250 \text{ m}$.

Question (a):

Find the net horizontal force on the dam from the gauge pressure of the water (gauge pressure is the pressure difference w.r.t. the external air).

Solution (a):

- (1) At depth y the gauge pressure of the water is $p = \rho g y$, where ρ is the density of the water.
- (2) We consider a horizontal strip of width W at depth y , with (vertical) thickness dy , across the dam.



Problem 3



香港中文大學(深圳)理工学院
School of Science and Engineering

Solution (a):

(3) Its area is $dA = Wdy$ and the force it exerts on the dam is $dF = pdA = \rho gyWdy$.

The total force of the water on the dam is

$$F = \int_0^D \rho gyW dy = \frac{1}{2} \rho g W D^2 = \frac{1}{2} (1.00 \times 10^3 \text{ kg/m}^3) (9.80 \text{ m/s}^2) (250 \text{ m}) (30.0 \text{ m})^2 \\ = 1.10 \times 10^9 \text{ N}.$$


Question (b):

Find the net torque due to that force about a horizontal line through O parallel to the width of the dam. This torque tends to rotate the dam around that line, which would cause the dam to fail.

Solution (b):

(1) Again we consider the strip of water at depth y . Its moment arm for the torque it exerts about O is $D - y$ so the torque it exerts is

$$d\tau = dF(D - y) = \rho gyW(D - y)dy$$



Gao Qian

Problem 3



香港中文大學(深圳)理工学院
School of Science and Engineering

Solution (b):

(2) and the total torque of the water is

$$\begin{aligned}\tau &= \int_0^D \rho g y W (D - y) dy = \rho g W \left(\frac{1}{2} D^3 - \frac{1}{3} D^3 \right) = \frac{1}{6} \rho g W D^3 \\ &= \frac{1}{6} (1.00 \times 10^3 \text{ kg/m}^3) (9.80 \text{ m/s}^2) (250 \text{ m}) (30.0 \text{ m})^3 = 1.10 \times 10^{10} \text{ N} \cdot \text{m}.\end{aligned}$$

Question (c):


Find the effective moment arm of the torque $\mathbf{r}_{eff} \equiv \boldsymbol{\tau}/F$.

Solution (c):

We write $\boldsymbol{\tau} = \mathbf{r}F$, where \mathbf{r} is the effective moment arm. Then,

$$r = \frac{\tau}{F} = \frac{\frac{1}{6} \rho g W D^3}{\frac{1}{2} \rho g W D^2} = \frac{D}{3} = \frac{30.0 \text{ m}}{3} = 10.0 \text{ m}.$$

QED.



Gao Qian

Problem 4



香港中文大學(深圳)理工学院
School of Science and Engineering

* Fluid Flow.

Water runs into a fountain, filling all the pipes, at a steady rate of **$0.750 \text{ m}^3/\text{s}$** .

Questions:

- (a) How fast will it shoot out of a hole **4.50 cm** in diameter?
- (b) At what speed will it shoot out if the diameter of the hole is three times as large?

Solutions:

Identify: The volume flow rate is Av . **Set Up:** $Av = 0.750 \text{ m}^3/\text{s}$. $A = \pi D^2/4$.

Execute (a): $v\pi D^2/4 = 0.750 \text{ m}^3/\text{s}$. $v = \frac{4(0.750 \text{ m}^3/\text{s})}{\pi(4.50 \times 10^{-2} \text{ m})^2} = 472 \text{ m/s}$.

Execute (b): vD^2 must be constant, so $v_1 D_1^2 = v_2 D_2^2$.

$$v_2 = v_1 \left(\frac{D_1}{D_2} \right)^2 = (472 \text{ m/s}) \left(\frac{D_1}{3D_1} \right)^2 = 52.4 \text{ m/s}$$

QED.

Gao Qian

Problem 5



香港中文大學(深圳) 理工学院
School of Science and Engineering

*** * (Halliday C14-P57).**

In Fig. below, water flows through a horizontal pipe and then out into the atmosphere at a speed $v_1 = 23.0 \text{ m/s}$. The diameters of the left and right sections of the pipe are **5.00 cm** and **3.00 cm**.

Question (a): What volume of water flows into the atmosphere during a **20.0 min** period?


Solution (a): The volume of water (during **20 minutes**) is

$$V = (v_1 t) A_1 = (23 \text{ m/s})(20 \text{ min})(60 \text{ s/min}) \left(\frac{\pi}{4} \right) (0.03 \text{ m})^2 = 19.5 \text{ m}^3.$$

Question (b): In the left section of the pipe, what is the speed v_2 ?

Solution (b): The speed in the left section pipe is

$$v_2 = v_1 \left(\frac{A_1}{A_2} \right) = v_1 \left(\frac{d_1}{d_2} \right)^2 = (23 \text{ m/s}) \left(\frac{3.0 \text{ cm}}{5.0 \text{ cm}} \right)^2 = 8.28 \text{ m/s}.$$



Problem 5



香港中文大學(深圳)理工学院
School of Science and Engineering

Question (c): Find the gauge pressure in the left section of the pipe.


Solution (c):

Since

$$p_1 + \frac{1}{2}\rho v_1^2 + \rho g h_1 = p_2 + \frac{1}{2}\rho v_2^2 + \rho g h_2$$

and $h_1 = h_2$, $p_1 = p_0$, where p_0 is the atmospheric pressure,

$$\begin{aligned} p_2 &= p_0 + \frac{1}{2}\rho(v_1^2 - v_2^2) \\ &= 1.01 \times 10^5 \text{ Pa} + \frac{1}{2}(1.0 \times 10^3 \text{ kg/m}^3)[(23 \text{ m/s})^2 - (8.28 \text{ m/s})^2] \\ &= 3.31 \times 10^5 \text{ Pa} = 3.28 \text{ atm.} \end{aligned}$$



Thus, the gauge pressure is

$$(3.28 \text{ atm} - 1.00 \text{ atm}) = 2.28 \text{ atm} = 2.30 \times 10^5 \text{ Pa.}$$

QED.



Gao Qian

Problem 6



* * (Halliday C14-P48).

A pitot tube (Figures below) is used to determine the air-speed of an airplane (relative to the air). It consists of an outer tube with a number of small holes **B** (four are shown) that allow air into the tube; that tube is connected to one arm of a *U*-tube. The other arm of the *U*-tube is connected to hole **A** at the front end of the device, which points in the direction the plane is headed. At **A** the air becomes stagnant so that $\mathbf{v}_A = \mathbf{0}$. At **B**, however, the speed of the air presumably equals the airspeed \mathbf{v} of the plane.



Problem 6



Question (a): Use Bernoulli's equation to show that

$$v = \sqrt{\frac{2\rho gh}{\rho_{air}}},$$


where ρ is the density of the liquid in the U -tube and h is the difference in the liquid levels in that tube.

Solution (a):

(1) Bernoulli's equation gives $p_A + \frac{1}{2}\rho_{air}0^2 = p_B + \frac{1}{2}\rho_{air}v^2$.

(2) However, $\Delta p = p_A - p_B = \rho gh$ in order to balance the pressure in the two arms of the U -tube.

(3) Thus $\rho gh = \frac{1}{2}\rho_{air}v^2$, or $v = \sqrt{\frac{2\rho gh}{\rho_{air}}}$.



Problem 6





Question (b): Suppose that the level difference h is **20.0 cm** in the tube. What is the plane's speed relative to the air? The density of the air is **1.03 kg/m³** and that of the liquid is **810 kg/m³**.

Solution (b):

The plane's speed relative to the air is

$$v = \sqrt{\frac{2\rho g h}{\rho_{air}}} = \sqrt{\frac{2(810 \text{ kg/m}^3)(9.8 \text{ m/s}^2)(0.200 \text{ m})}{1.03 \text{ kg/m}^3}} = 55.5 \text{ m/s.}$$




QED.

Problem 7



** As shown below, the Venturi meter can be used to measure flow speed in a pipe. The narrow pipe part of the pipe is called the throat. Derive an expression for the flow speed v_1 in terms of the cross-sectional areas A_1 and A_2 and the difference in height h of the liquid levels in the two vertical tubes.



Problem 7



Solution:

- (1) We assume the fluid is incompressible and steady and has negligible internal friction.
- (2) Hence we can use Bernoulli's equation and apply this equation to the wide part (**point 1**) and narrow part (**point 2**, the throat) of the pipe.


- (3) **Points 1** and **2** have the same vertical coordinate $y_1 = y_2$, so

$$p_1 + \rho g y_1 + \frac{1}{2} \rho v_1^2 = p_2 + \rho g y_2 + \frac{1}{2} \rho v_2^2.$$

- (4) From the continuity equation, $v_2 = (A_1/A_2)v_1$. Substituting this into the Bernoulli's equation, we get

$$p_1 - p_2 = \frac{1}{2} \rho v_1^2 \left[\frac{A_1^2}{A_2^2} - 1 \right].$$

Difference in height results from reduced pressure in throat (point 2).



Problem 7



Solution:

(5) Now let us look at the vertical pressure difference.

(6) It is important to note that the fluid line is indicating the fluid motion along the horizontal direction while the fluid is stagnant along the vertical direction.

(7) For static fluid, one finds $p_1 - p_2 = \rho gh$.


(8) Therefore, combining this with the above result and solving for v_1 , we get

$$v_1 = \sqrt{\frac{2gh}{(A_1/A_2)^2 - 1}}.$$

Comments:

Since $v_2 > v_1$, pressure differences produce a net force to the right that makes the fluid speed up as it enters the throat, and a net force to the left that slows it as it leaves.

QED.



Problem 8



* * * A large bucket of height H and cross-sectional area A_1 is filled with water. The top is open to the atmosphere. There is an opening faucet of area A_2 , which is much smaller than A_1 , at the bottom of the bucket.

Question (a):

Show that when the height of the water is h , the speed of the water leaving the faucet is approximately $\sqrt{2gh}$.

Solution (a):

From Bernoulli's equation between the surface and the bottom of the bucket, one gets

$$p_0 + \rho gh + \frac{1}{2}\rho \cdot 0^2 = p_0 + \rho g \cdot 0 + \frac{1}{2}\rho \cdot v^2$$
$$\Rightarrow v = \sqrt{2gh}$$

with p_0 the atmosphere pressure.

Problem 8



* * * A large bucket of height H and cross-sectional area A_1 is filled with water. The top is open to the atmosphere. There is an opening faucet of area A_2 , which is much smaller than A_1 , at the bottom of the bucket.

Question (b):

Show that if $A_2 \ll A_1$, the rate of change of the height h of the water is given by

$$\frac{dh}{dt} = -\frac{A_2}{A_1} \sqrt{2gh}.$$

Solution (b):

During short time dt , according to the conservation of fluid matter volume, one gets the volume that goes out of the bucket from the bottom equals the volume decrease from the top surface. Thus

$$v dt A_2 = -dh A_1 \quad \Rightarrow \quad \frac{dh}{dt} = -\frac{A_2}{A_1} v = -\frac{A_2}{A_1} \sqrt{2gh},$$

where $dh < 0$ means the height of the water surface is decreasing.

Problem 8



香港中文大學(深圳)理工学院
School of Science and Engineering

Question (c):

Find h as a function of time if $h = H$ at $t = 0$.

Solution (c):

Solve ordinary differential equation by separating the variables h from t , then write

$$\frac{dh}{\sqrt{h}} = -\frac{A_2}{A_1} \sqrt{2g} dt,$$

$$\Rightarrow \int \frac{dh}{\sqrt{h}} = -\frac{A_2}{A_1} \sqrt{2g} \int dt + C$$

$$\Rightarrow 2\sqrt{h} = -\frac{A_2}{A_1} \sqrt{2g} t + C$$

From initial condition $h(0) = H \rightarrow C = 2\sqrt{H}$, one finds $\sqrt{h} = -\frac{A_2}{A_1} \sqrt{\frac{g}{2}} t + \sqrt{H}$,

and eventually writes $h = \left(-\frac{A_2}{A_1} \sqrt{\frac{g}{2}} t + \sqrt{H} \right)^2$.

Problem 8



香港中文大學(深圳)理工学院
School of Science and Engineering

* * * A large bucket of height H and cross-sectional area A_1 is filled with water. The top is open to the atmosphere. There is an opening faucet of area A_2 , which is much smaller than A_1 , at the bottom of the bucket.

Question (d):

Find the total time needed to drain the bucket if $H = 2.00 \text{ m}$, $A_1 = 0.800 \text{ m}^2$, and $A_2 = 1.00 \times 10^{-4} A_1$.

Solution (d):

From the above results, one can solve for time (t) as a function of h and obtain

$$t = (\sqrt{H} - \sqrt{h}) \frac{A_1}{A_2} \sqrt{\frac{2}{g}} = (\sqrt{2.00} - 0) \frac{A_1}{1.00 \times 10^{-4} A_1} \sqrt{\frac{2}{9.81}} = 6.39 \times 10^3 \text{ s.}$$

QED.

Problem 9



香港中文大學(深圳)理工学院
School of Science and Engineering

* Flight plan to the planet Mars

* The position of a particle is given by $x = 2.5 \cos \pi t$, where x is in meters and t is in seconds.

Questions:

- (a) Find the maximum speed and maximum acceleration of the particle.
- (b) Find the velocity and acceleration of the particle when $x = 1.5 \text{ m}$.

Solution: For the harmonic oscillation

$$x = 2.5 \cos \pi t,$$

$$v = \frac{dx}{dt} = -2.5 \pi \sin \pi t,$$

$$a = \frac{dv}{dt} = -2.5 \pi^2 \cos \pi t.$$

Solution (a): $v_{\max} = 2.5\pi = 7.9 \text{ m/s}$, $a_{\max} = 2.5 \pi^2 = 25 \text{ m/s}^2$.

Gao Qian

Problem 9



Solution (b):

(1) When $x = 1.5 = 2.5 \cos \pi t$, we have $\cos \pi t = 0.6$ and $\sin \pi t = \pm \sqrt{1 - 0.6^2} = \pm 0.8$.

(2) As shown in the left figure, we can find that there are two sets of solution (given by the two points where two curves coincide) within a cycle.

(3) If we compute t explicitly, we can find that $t = 0.30$ and $t = 1.7$. Therefore,


$$v = \mp 2.5\pi \times 0.8 = \mp 6.28 \text{ m/s}$$

$$a = -2.5\pi^2 \times 0.6 = -14.8 \text{ m/s}^2$$

(4) For v , the minus sign comes from $t = 0.30$ while the solution with the plus sign comes from $t = 1.7$.

(5) However, for the acceleration $a = -\omega^2 x$, it is always opposite to x . So a only has one solution.

QED.



Problem 10



* Simple Harmonic Oscillation

Questions:

- (a) Show that the term $A_0 \cos(\omega t + \delta)$ can be written as $A_s \sin(\omega t) + A_c \cos(\omega t)$, and determine A_s and A_c in terms of A_0 and δ .
- (b) Relate A_c and A_s to the initial position and velocity of a particle undergoing simple harmonic motion.

Solution: Given $x(t) = A_0 \cos(\omega t + \delta)$, then by differentiation one gets

$$v = \frac{dx}{dt} = -A_0 \omega \sin(\omega t + \delta).$$

Solution (a):

(1) We can begin with

$$\begin{aligned} x(t) &= A_0 \cos(\omega t + \delta) = A_0 \cos \omega t \cos \delta - A_0 \sin \omega t \sin \delta \\ &= A_0 \cos \delta \cos \omega t - A_0 \sin \delta \sin \omega t \\ &= A_c \cos \omega t + A_s \sin \omega t \end{aligned}$$

Problem 10



香港中文大學(深圳)理工学院
School of Science and Engineering

Solution (a):

(2) By comparing the coefficients of $\cos \omega t$ and $\sin \omega t$, we can find

$$A_s = -A_0 \sin \delta, \quad \text{and} \quad A_c = A_0 \cos \delta.$$

Solution (b):

(1) At $t = 0$, $x(0) = A_0 \cos \delta$ and $v(0) = -A_0 \omega \sin \delta$.

(2) Therefore, $A_c = x(0)$ and $A_s = v(0)/\omega$.

QED.

Comments: It is also useful to write

$$x(t) = x(0) \cos \omega t + \frac{v(0)}{\omega} A_0 \sin \omega t,$$

and

$$v(t) = -x(0)\omega \sin \omega t + v(0) \cos \omega t.$$