

Provider Risk Stratification Policy

Synthetic Training Policy - Non-PHI

Defines provider risk tiers based on claim volume and historical denial patterns. Covers provider_risk_score, provider_claim_count, and low_volume_provider_risk features.

Policy ID	CLAIMOPS-POL-011
Effective Date	2026-04-25
Applies To	Synthetic claims, provider, diagnosis, and cost datasets in this repository
Source Type	Locally generated demo policy; not a real payer coverage policy

Policy Summary

- provider_risk_score aggregates denial frequency, charge outliers, and practice-pattern metrics into a single risk score.
- Provider risk is stratified into tiers based on claim volume and denial history.
- Micro-volume and low-volume providers have statistically unreliable risk scores and are assessed primarily on claim-level signals.
- High-volume providers with elevated risk scores signal systemic documentation or coding issues that may affect all their claims.

Rules

Rule ID	Requirement
RISK-01	provider_risk_score baseline is 0.15 (synthetic prior for unknown providers). Scores between 0.15 and 0.30 are within normal range. Scores above 0.30 are elevated and trigger enhanced review for all claims from that provider.
RISK-02	provider_risk_score above 0.50 is high risk. Claims from high-risk providers receive the most intensive review tier: mandatory clinical documentation check, cost benchmark comparison, and dx-px compatibility verification before payer submission.
RISK-03	Micro-volume tier: provider_claim_count < 5. Individual risk scores are unstable due to insufficient data. Claims are assessed using dx-px compatibility, cost benchmarks, and completeness signals rather than provider history. low_volume_provider_risk = 1.
RISK-04	Low-volume tier: provider_claim_count between 5 and 20. Risk scores should be used with caution. Supplement provider-history signals with claim-level coding and cost signals. low_volume_provider_risk = 1.
RISK-05	Adequate-volume tier: provider_claim_count between 20 and 100. Risk score estimates are moderately reliable. Use provider history as one signal among several.
RISK-06	High-volume tier: provider_claim_count > 100. Risk score estimates are statistically reliable. Provider history carries full weight in the denial-risk model alongside claim-level signals.

Remediation Guidance

- For micro-volume providers, focus remediation on claim-level completeness and coding accuracy.

- For high-volume, high-risk providers, initiate a provider education or documentation improvement program.
- Recompute provider risk scores weekly to capture changing practice patterns.
- Do not include provider identifiers, claim counts, or risk scores in LLM prompts.

RAG Safety Note

This document contains only synthetic policy text. It intentionally avoids patient names, patient identifiers, service dates, and real billed amounts. Use policy_id, rule_id, denial reason, diagnosis category, and procedure status as retrieval metadata.