

stationary_combustion_r4.R

April 4, 2025

Contents

Stationary_combustion	1
---------------------------------	---

Index	6
--------------	----------

Stationary_combustion	<i>Create gridded stationary combustion methane emissions maps</i>
-----------------------	--

Description

‘Stationary_combustion’ writes up to 56 netcdf files of gridded methane emissions from stationary combustion sources, as well as optional visuals

Usage

```
Stationary_combustion(  
  input_directory,  
  domain,  
  domain_template,  
  state_name_list,  
  output_directory,  
  inventory_year,  
  verbose,  
  County_Tigerlines,  
  Use_ACES,  
  Use_Vulcan,  
  ACES_directory,  
  vulcan_directory,  
  ACES_year,  
  vulcan_band,  
  stationary_combustion_by_state,  
  stationary_combustion_by_domain,  
  stationary_combustion_GHGI_data,  
  stationary_combustion_emission_factors,  
  EIA_API_key,  
  plot_directory,  
  State_Tigerlines  
)
```

Arguments

input_directory	Character providing the full filepath to save/load raw input data
domain	SpatVector polygon outlining the desired output area
domain_template	SpatRaster providing the desired output grid, including the desired resolution and coordinate reference system
state_name_list	Character vector listing all states within the desired domain
output_directory	Character providing the full filepath to save processed data
inventory_year	Character indicating the desired year of data to use.
verbose	Logical indicating whether to save additional output. This includes plots of the gridded methane emissions for each fuel-sector-inventory-variation combination as well as 2 summed plots for each inventory-variation combination - one for wood and one for all other sectors.
County_Tigerlines	SpatVector. United States Census Bureau county shapefile downloaded in Main.
Use_ACES	Logical. Pulled from config file. indicating whether or not to use ACES to disaggregate from county-level to pixel level emissions. Either ACES or Vulcan must be used, though both can be.
Use_Vulcan	Logical. Pulled from config file. indicating whether or not to use Vulcan to disaggregate from county-level to pixel level emissions. Either ACES or Vulcan must be used, though both can be.
ACES_directory	Character providing the full path to a folder containing the ACES sectoral CO2 inventories. Must include the residential, commercial, electric (elec), and industrial sectors. ACES v2.0 is available at https://doi.org/10.3334/ORNLDAAAC/1943 , though the hourly file should be averaged across hours to create an annually averaged inventory. There is a function to do this as well, though it is quite time consuming given the amount of data that needs to be downloaded. The year closest to "inventory_year" is used, but those further from that year are considered if the closest is unavailable.
vulcan_directory	Character providing the full path to a folder containing the Vulcan sectoral CO2 inventories. Must include the residential, commercial, electric (elec_prod), and industrial sectors. Vulcan v3.0 is available at https://doi.org/10.3334/ORNLDAAAC/1741 , and the annual mean files should be used. The year closest to "inventory_year" is used. As all years are contained in the same file, it does not search for other years.
ACES_year	Numeric providing the year of ACES data to use, calculated in Main.
vulcan_band	Numeric providing the band of Vulcan data to use (1-6 = 2010 - 2015), calculated in Main.
stationary_combustion_by_state	Logical. Pulled from config file. indicating whether state-toal emissions should be distributed to the county scale as is. Either bystate or bydomain must be used, though both can be.
stationary_combustion_by_domain	Logical. Pulled from config file. indicating whether state-toal emissions should be aggregated to the domain and then distributed to the county scale. Either bystate or bydomain must be used, though both can be.

stationary_combustion_GHGI_data

Data frame. Pulled from config file. 1 by 15 data frame with consumption for each sector-fuel combination from the GHGI. Although the data is national, there is a state entry set to US_EPA. Consumption is in thousands of British Thermal Units (BTU). The GHGI is available at <https://www.epa.gov/ghgemissions/inventory-us-greenhouse-gas-emissions-and-sinks> and the values can be found in the Annexes in a table titled "Fuel Consumption by Stationary Combustion for Calculating CH₄ and N₂O Emissions (TBtu)". Includes every combination of residential (res), commercial (com), electric (elec), and residential (res) sectors with coal, petroleum (petr), natural gas (gas), and wood fuels with names using the abbreviations in parentheses (e.g., com_coal, elec_petr). Excludes res_coal and res_gas as residential coal does not exist in the U.S. anymore and residential gas is accounted for elsewhere.

stationary_combustion_emission_factors

Data frame. Pulled from config file. 1 by 14 data frame with emission factors for each sector-fuel combination from the IPCC. Built equivalently to the GHGI_data, but without an entry for the state. Emission factors are in g/GJ, equivalent to kg/TJ. Default emission factors are available in IPCC 2006 volume 2: Energy, tables 2.2 through 2.5 <https://www.ipcc-nggip.iges.or.jp/public/2006gl/vol2.html>. The natural gas electric sector emission factor is instead pulled from Hajny et al., 2019 <https://doi.org/10.1021/acs.est.9b01875>. This is 5.7 kg/TJ, within uncertainties of the GHGI value of 4.1 kg/TJ, both of which are larger than the IPCC default of 1 kg/TJ.

EIA_API_key

Character. Pulled from config file. API key to access SEDS data API. The API is described at <https://www.eia.gov/opendata/> and one can register for a key with a link on the right hand side of this page.

plot_directory

Character providing the full filepath to save figures. Only relevant if verbose = TRUE.

State_Tigerlines

SpatVector. United States Census Bureau county shapefile. Available at <https://www.census.gov/geographies/mapping-files/time-series/geo/tiger-line-file.html>. Only relevant if verbose=TRUE.

Details

This function calculates and grids methane emissions from stationary combustion. It uses the Energy Information Administration's (EIA) State Energy Data System (SEDS), Environmental Protection Agency's (EPA) Greenhouse Gas Inventory (GHGI), EPA National Emissions Inventory (NEI) and the Vulcan and/or Anthropogenic Carbon Emission System (ACES) CO₂ inventories. First, the ratio of SEDS national consumption by fuel-sector combination to the equivalent GHGI data is applied to SEDS state data so that they are approximately consistent with the GHGI. The state total consumption is then converted to emissions using emission factors. There are two variations at this step.

bystate: The state level emissions are then distributed to the county level using NEI CO concentrations to calculate weighting values for each county (i.e., county CO / state total CO).

bydomain: The state level emissions are summed to get a total for all states in the domain. This is distributed to the county level using NEI CO concentrations to calculate weighting values for each county (i.e., county CO / domain total CO).

Lastly, county total emissions are distributed using the ACES and/or Vulcan CO₂ inventories.

This entire process is done separately for each fuel and sector using the GHGI, SEDS, and NEI data that are broken down by sector and fuel and CO₂ inventories that are broken down by sector.


```

inventory_year=2018,
verbose=TRUE,
Use_ACES=TRUE,
Use_Vulcan=TRUE,
ACES_directory=~/.//Desktop/in/Inventories/ACES_v2.0",
vulcan_directory=~/.//Desktop/in/Inventories/Vulcan_v3.0",
ACES_year=2017,
vulcan_band=6,
stationary_combustion_by_state=TRUE,
stationary_combustion_by_domain=TRUE,
stationary_combustion_GHGI_data=data.frame(
  "State"="US_EPA",
  "com_coal"=17,
  "ind_coal"=517,
  "elec_coal"=10554,
  "res_petr"=975,
  "com_petr"=801,
  "ind_petr"=2062,
  "elec_petr"=42,
  "com_gas"=3647,
  "ind_gas"=9484,
  "elec_gas"=11553,
  "res_wood"=544,
  "com_wood"=84,
  "ind_wood"=1407,
  "elec_wood"=68),
stationary_combustion_emission_factors=data.frame(
  "com_coal"=10,
  "ind_coal"=10,
  "elec_coal"=1,
  "res_petr"=10,
  "com_petr"=10,
  "ind_petr"=3,
  "elec_petr"=3,
  "com_gas"=5,
  "ind_gas"=1,
  "elec_gas"=5.4/(1.0550559*0.9), #g/mmbtu to g/GJ and low to high heating value (0.9)
  "res_wood"=300,
  "com_wood"=300,
  "ind_wood"=30,
  "elec_wood"=30),
EIA_API_key=user_key,
State_Tigerlines=vect("~/.//Desktop/in/State_Tigerlines/tl_2018_us_state.shp"),
County_Tigerlines=vect("~/.//Desktop/in/County_Tigerlines/tl_2018_us_county.shp"),
plot_directory=~/.//Desktop/plots/")

```

Index

Stationary_combustion, [1](#)