

EXERCISE 3-44 Show that a nonsymmetric doubly contravariant or covariant second-rank tensor cannot be diagonalized by a linear transformation.

3-15 Magnitude of Second-Rank Tensor

If \mathbf{T} is a second-rank tensor, and \mathbf{u} is a unit vector, the operation $\mathbf{T}\mathbf{u}$ produces a new vector \mathbf{v} ; that is,

$$\mathbf{v} = \mathbf{T}\mathbf{u} \quad (3-134)$$

The magnitude $T(\mathbf{u})$ of the tensor \mathbf{T} in the direction of the unit vector \mathbf{u} is equal to the length of the projection of \mathbf{v} onto \mathbf{u} . That is,

$$T(\mathbf{u}) = \mathbf{u} \cdot \mathbf{v} = \mathbf{u}\mathbf{T}\mathbf{u} \quad (3-135)$$

In terms of components

$$T(\mathbf{u}) = T^{ij}u_i u_j = T_{ij} u^i u^j \quad (3-136)$$

If \mathbf{u} is an eigenvector of \mathbf{T} , $T(\mathbf{u})$ is of course just the corresponding eigenvalue. A physical interpretation of the magnitude of a second-rank tensor will be described in Section 5-2.

EXERCISE 3-45 Derive a formula for the magnitude in the direction of reciprocal axis \mathbf{a}^k of an anisotropic temperature factor tensor with components β^{ij} .

EXERCISE 3-46 Calculate the magnitude of the anisotropic temperature factor tensor of Exercise 3-38 in the crystal directions (a) $[1, 0, 0]$; (b) $[0, 1, 0]$; (c) $[0, 0, 1]$; (d) $[-0.1097, 0.0210, 0.0492]$.

3-16 Rigid-Body Motion

The thermal motion of atoms in a molecular crystal often may be resolved into the intermolecular movements of the molecules as units and the intramolecular movements of the atoms within the molecules relative to each other. Separation of the rigid-body molecular motion from the total movement is particularly clearcut if the amplitudes of molecular motion are large compared with the vibrational amplitudes within the molecules. Determination of the rigid-body motion from the observed anisotropic temperature factors was first described by Cruickshank (1956). A rigorous theory, which permitted noninter-

secting
blood (1
tion. Th
dyadic r
lucid ma
Johnson
great em
body mo
problems

The ge
tion and
rotation.
sequently
quate trea
the quadri
tions, whi
torial rep
cartesian
tor \mathbf{e}_3 mo
small, the
and this is
 $\lambda \Delta \mathbf{r}$, whe
general dis
The tota
equilibrium

or, in suffi

The averag
the doubly

TABLE 5-1. CONDITIONS IMPOSED BY SYMMETRY UPON THE COMPONENTS OF A SYMMETRIC SECOND-RANK TENSOR REFERRED TO CRYSTAL AXES

Crystal system	σ^{11}	σ^{12}	σ^{13}	σ^{22}	σ^{23}	σ^{33}
Triclinic	σ^{11}	σ^{12}	σ^{13}	σ^{22}	σ^{23}	σ^{33}
Monoclinic	σ^{11}	0	σ^{13}	σ^{22}	0	σ^{33}
Orthorhombic	σ^{11}	0	0	σ^{22}	0	σ^{33}
Tetragonal	σ^{11}	0	0	σ^{11}	0	σ^{33}
Hexagonal, trigonal	σ^{11}	$\frac{1}{2}\sigma^{11}$	0	σ^{11}	0	σ^{33}
Cubic	σ^{11}	0	0	σ^{11}	0	σ^{11}

TABLE 5-2. CONDITIONS IMPOSED BY SYMMETRY UPON THE COMPONENTS OF A SYMMETRIC SECOND-RANK TENSOR REFERRED TO CARTESIAN AXES

Crystal system	σ^{11}	σ^{12}	σ^{13}	σ^{22}	σ^{23}	σ^{33}
Triclinic	σ^{11}	σ^{12}	σ^{13}	σ^{22}	σ^{23}	σ^{33}
Monoclinic	σ^{11}	0	σ^{13}	σ^{22}	0	σ^{33}
Orthorhombic	σ^{11}	0	0	σ^{22}	0	σ^{33}
Tetragonal	σ^{11}	0	0	σ^{11}	0	σ^{33}
Hexagonal, trigonal	σ^{11}	0	0	σ^{11}	0	σ^{33}
Cubic	σ^{11}	0	0	σ^{11}	0	σ^{11}

EXERCISE 5-1 The specific resistance of bismuth metal at 20°C is 109×10^{-6} ohm-cm perpendicular to \mathbf{a}_3 and 138×10^{-6} ohm-cm parallel to \mathbf{a}_3 . Bismuth is rhombohedral, space group $R\bar{3}m$, with $a_1 = 4.537$, $a_3 = 11.838$ Å for the hexagonal cell.

(a) Calculate the components of the conductance tensor referred to cartesian coordinates with \mathbf{e}_3 along \mathbf{a}_3 .

(b) Calculate the components of the conductance tensor referred to the hexagonal axes.

(c) Calculate the resistance of a single crystal bismuth rod 3.00 cm long and 1.00 cm in diameter cut with the cylinder axis along [001].

(d) Suppose that a single crystal of bismuth is cut into a cylinder 3.00 cm long and 1.00 cm in diameter with the cylinder axis along the [211] direction (a rhombohedral axis). Calculate the resistance to the flow of current along [211] when an electric field is applied to the ends of the rod.

EXERCISE 5-2 Calculate the specific conductance of bismuth (see Exercise 5-1) along the hexagonal [211], $[\bar{1}11]$, and $[\bar{1}\bar{2}1]$ directions.

EXERCISE 5-3 Evaluate $\int \bar{\mathbf{u}} \chi \mathbf{u} d\tau / \int c d\theta d\phi$, θ varies from

EXERCISE 5-4 Measurements on a zirconium crystal. The values are

(a) Calculate all components of the tensor.

(b) Find the principal values.

(c) Zircon is tetragonal. What are the tetragonal axes?

(d) In what direction are the principal values?

(e) Positive values of the tensor are diamagnetic? What are the negative values?

EXERCISE 5-5 The principal values of the tensor of the

may be determined if points are placed along the A^{22} axis. A^{22} is the figure circle centered at C . Derive equations for the angle θ in the cartesian axes through

EXERCISE 5-6 The tensor of the watt/deg cm parallel to the \mathbf{a}_3 axis (Ho, Powell, and Lil

University of Illinois at Urbana-Champaign
Online Library Catalog

Library Gateway	Hours & Locations
E-Journals	Ask A Librarian

Try [VuFind](#) (an alternate catalog)

Your session will expire in 07:17

[NewSearch](#) [I-ShareCatalog](#) [OtherLibraries](#) [Headings](#) [Titles](#) [MyAccount / Renew Books](#) [Request](#) [History](#) [Quit](#) [Help](#)

Now Searching: University of Illinois at Urbana-Champaign Library

You Searched: Boolean Search =

TKEY vectors AND TKEY tensors AND TKEY crystallography and GKEY vectors AND GKEY tensors AND GKEY crystallography

Search Results: Displaying 1 of 1 entries.

[Full View](#)[Brief View](#)[More Like This](#)[Staff \(MARC\) View](#)

Cover Image:

[Click for more information on this title.](#)**Author:** [Sands, Donald, 1929-](#)**Title:** Vectors and tensors in crystallography / Donald E. Sands.**Published:** Reading, Mass. : Addison-Wesley, 1982.**Physical Description:** xvi, 228 p. : ill. ; 24 cm.**Includes:** Bibliography: p. 199-201.**Subject (LCSH):** [Crystallography, Mathematical](#)[Vector analysis](#)[Tensor algebra](#)**Notes:** Includes index.**ISBN:** 0201071479**Persistent link to this page:** <https://i-share.carli.illinois.edu/uiu/cgi-bin/Pwebrecon.cgi?DB=local&v1=1&BBRecID=570096>**Institution:** University of Illinois at Urbana-Champaign Library**Location:** Engineering**Call Number:** [548.7 SA57V](#) → [Text me this call number](#)**Copy:** 1**Status:** c.1 Overdue - Due on 04-10-12 [Find copies in other I-Share libraries.](#)

Print / Download / Email / Store	
Select Download Format	Full Record <input type="button" value="Print/Save"/> <input type="button" value="Save Search Strategy"/>
Enter your email address: <input type="button" value="Email"/>	
<input type="button" value="Put Selected Titles in MyBookbag"/>	

[NewSearch](#) [I-ShareCatalog](#) [OtherLibraries](#) [Headings](#) [Titles](#) [MyAccount / Renew Books](#) [Request](#) [History](#) [Quit](#) [Help](#)Comments on the Library Catalog may be sent to: opac@library.uiuc.edu