

# Carter (1968)

K. R.

2025 3/31

## 0.1 Tetrads

Carter researched metrics of spacetime that the Hamilton–Jacobi equation of a particle is separable in terms of two abelian groups. The metrics have six functions  $(P_\lambda, P_\mu, Q_\lambda, Q_\mu, \Delta_\lambda, \Delta_\mu)$  of coordinates  $\lambda, \mu$ . These are subscripts and are called non-ignorable because parameters of two abelian groups are not  $\lambda, \mu$ . These are similar to the radius coordinate  $r$  of the spherical coordinates. The remaining coordinates are denoted by  $\psi$  and  $\chi$ , which is called ignorable. Using the coordinate basis  $(d\lambda, d\mu, d\psi, d\chi)$  of the dual tangent space, the metric is restricted to certain symmetric forms

$$\begin{aligned} ds^2 &= \frac{Z}{\Delta_\lambda} d\lambda^2 + \frac{Z}{\Delta_\mu} d\mu^2 + \frac{\Delta_\mu}{Z} [P_\lambda d\psi - Q_\lambda d\chi]^2 - \frac{\Delta_\lambda}{Z} [P_\mu d\psi - Q_\mu d\chi]^2, \\ Z &= P_\lambda Q_\mu - P_\mu Q_\lambda. \end{aligned} \quad (0.1.1)$$

Carter discussed the separability of variables of Hamilton–Jacobi equations with electromagnetic fields. Electromagnetic fields without source fields are described by four-potential in spacetime. A classical four-potential is now one-form

$$A = \frac{P_\lambda X_\mu + P_\mu X_\lambda}{Z} d\psi - \frac{Q_\lambda X_\mu + Q_\mu X_\lambda}{Z} d\chi, \quad (0.1.2)$$

with functions  $X_\lambda$  and  $X_\mu$  and the corresponding electromagnetic field (strength) tensor is  $F = dA$  ( $\times 2$  on the Carter’s paper).

Carter classifies metrics of

$$\text{type } A \begin{cases} P_\lambda = \lambda^2, P_\mu = -\mu^2, Q_\lambda = 1, Q_\mu = 1, \\ \Delta_\lambda = \frac{1}{3}\Lambda\lambda^4 + h\lambda^2 - 2m_+\lambda + n + e^2, \Delta_\mu = \frac{1}{3}\Lambda\mu^4 - (h\mu^2 - 2m_-\mu - n), \end{cases} \quad (0.1.3)$$

$$\text{type } B(+) \begin{cases} P_\lambda = \lambda^2 + 1, P_\mu = 2\mu, Q_\lambda = 0, Q_\mu = 1, \\ \Delta_\lambda = \Lambda \left( \frac{1}{3}\lambda^4 + 2\lambda^2 - 1 \right) + (h(\lambda^2 - 1) - 2m_+\lambda + e^2), \Delta_\mu = -(h\mu^2 - 2m_-\mu + n), \end{cases} \quad (0.1.4)$$

$$\text{type } C(+) \begin{cases} P_\lambda = \lambda^2, P_\mu = 0, Q_\lambda = 0, Q_\mu = 1, \\ \Delta_\lambda = \Lambda\lambda^4 + (h\lambda^2 - 2m_+\lambda + e^2), \Delta_\mu = -(h\mu^2 - 2m_-\mu + n), \end{cases} \quad (0.1.5)$$

$$\text{type } D \begin{cases} P_\lambda = \frac{1}{2}, P_\mu = \frac{1}{2}, Q_\lambda = 1, Q_\mu = 1, \\ \Delta_\lambda = \Lambda\lambda^2 + (e^2\lambda^2 - 2m_+\lambda + n_+), \Delta_\mu = \Lambda\mu^2 - (e^2\mu^2 - 2m_-\mu + n_-). \end{cases} \quad (0.1.6)$$

We can deduce ortho-normal tetrads whose superscripts are numbered as non-ignorable and ignorable using a condition  $U = Z$ ,

$$\omega^i = \left( \frac{U}{\Delta_i} \right)^{1/2} d\lambda^i, \quad \omega^{2i} = \frac{(U\Delta_{-i})^{1/2}}{Z} ((r) Z_i^r d\varphi^{2r}) \quad (0.1.7)$$

for an integer index  $i \in \{+1, -1\}$  and summed indices  $r = +1, -1$ . The non-ignorable coordinates correspond to two ones that the metric component depends on, in other words,  $(\lambda^{+1}, \lambda^{-1}) \equiv (\lambda, \mu)$ . On the other hand, the ignorable ones  $(\varphi^{+2}, \varphi^{-2})$  are equivalent to  $(\psi, \chi)$ . A round bracket  $(r)$  means summed signs, namely  $(r) Z_i^r d\varphi^{2r} = +Z_i^{+1} d\varphi^{+2} - Z_i^{-1} d\varphi^{-2} = P_i d\psi - Q_i d\chi$ . Then non-ignorable-numbered tetrads are represented by those greek letters as

$$\omega^{+1} = \left( \frac{Z}{\Delta_\lambda} \right)^{1/2} d\lambda, \quad \omega^{-1} = \left( \frac{Z}{\Delta_\mu} \right)^{1/2} d\mu \quad (0.1.8)$$

using  $U = Z$  and the remainder are

$$\omega^{+2} = \frac{(Z\Delta_\mu)^{1/2}}{Z} (+P_\lambda d\psi - Q_\lambda d\chi), \quad \omega^{-2} = \frac{(Z\Delta_\lambda)^{1/2}}{Z} (+P_\mu d\psi - Q_\mu d\chi). \quad (0.1.9)$$

These relations are summarized by a matrix

$$\hat{E} = \begin{bmatrix} \left( \frac{Z}{\Delta_\lambda} \right)^{1/2} & 0 & 0 & 0 \\ 0 & \left( \frac{Z}{\Delta_\mu} \right)^{1/2} & 0 & 0 \\ 0 & 0 & \frac{(Z\Delta_\mu)^{1/2}}{Z} P_\lambda & -\frac{(Z\Delta_\mu)^{1/2}}{Z} Q_\lambda \\ 0 & 0 & \frac{(Z\Delta_\lambda)^{1/2}}{Z} P_\mu & -\frac{(Z\Delta_\lambda)^{1/2}}{Z} Q_\mu \end{bmatrix}, \quad (0.1.10)$$

which combines the ortho-normal tetrads  $\omega^j$ s ( $j \in \{\pm 1, \pm 2\}$ )

$$\begin{bmatrix} \omega^{+1} \\ \omega^{-1} \\ \omega^{+2} \\ \omega^{-2} \end{bmatrix} = \hat{E} \begin{bmatrix} d\lambda \\ d\mu \\ d\psi \\ d\chi \end{bmatrix}. \quad (0.1.11)$$

The determinant of the matrix  $\hat{E}$  equals  $-Z$ :

$$\begin{aligned} \det \hat{E} &= \left( \frac{Z}{\Delta_\lambda} \right)^{1/2} \left( \frac{Z}{\Delta_\mu} \right)^{1/2} \frac{(Z\Delta_\lambda)^{1/2}}{Z} \frac{(Z\Delta_\mu)^{1/2}}{Z} (-P_\lambda Q_\mu + P_\mu Q_\lambda) \\ &= \left( \frac{Z^2}{Z\Delta_\lambda} \right)^{1/2} \left( \frac{Z^2}{Z\Delta_\mu} \right)^{1/2} \frac{(Z\Delta_\lambda)^{1/2} (Z\Delta_\mu)^{1/2}}{Z^2} (-Z) \\ &= |Z|^2 / Z \times (-1) = -Z \end{aligned} \quad (0.1.12)$$

Moreover, we can deduce the ortho-normal basis. Let  $\{\partial_\lambda, \partial_\mu, \partial_\psi, \partial_\chi\}$  be coordinate basis vectors and  $\{e_{+1}, e_{-1}, e_{+2}, e_{-2}\}$  be ortho-normal basis vectors corresponding to the tetrads  $\omega^j$ s. These ortho-normal vectors satisfy

$$\begin{bmatrix} e_{+1} & e_{-1} & e_{+2} & e_{-2} \end{bmatrix} = \begin{bmatrix} \partial_\lambda & \partial_\mu & \partial_\psi & \partial_\chi \end{bmatrix} \hat{E}^{-1}, \quad (0.1.13)$$

where the inverse matrix  $\hat{E}^{-1}$  of the transformation matrix Eq. (0.1.10) is

$$\hat{E}^{-1} = \begin{bmatrix} \frac{1}{\sqrt{Z/\Delta_\lambda}} & 0 & 0 & 0 \\ 0 & \frac{1}{\sqrt{Z/\Delta_\mu}} & 0 & 0 \\ 0 & 0 & \frac{Q_\mu}{\sqrt{Z\Delta_\mu}} & -\frac{Q_\lambda}{\sqrt{Z\Delta_\lambda}} \\ 0 & 0 & \frac{P_\mu}{\sqrt{Z\Delta_\mu}} & -\frac{P_\lambda}{\sqrt{Z\Delta_\lambda}} \end{bmatrix} \quad (0.1.14)$$

If  $\mathbf{E}^{-1}$  is expand concretely, each ortho-normal vector in  $\{e_{+1}, e_{-1}, e_{+2}, e_{-2}\}$  is represented by

$$\begin{aligned} e_{+1} &= \left( \frac{\Delta_\lambda}{Z} \right)^{1/2} \partial_\lambda, \quad e_{-1} = \left( \frac{\Delta_\mu}{Z} \right)^{1/2} \partial_\mu, \\ e_{+2} &= \frac{1}{(Z\Delta_\mu)^{1/2}} (Q_\mu \partial_\psi + P_\mu \partial_\chi), \quad e_{-2} = \frac{1}{(Z\Delta_\lambda)^{1/2}} (-Q_\lambda \partial_\psi - P_\lambda \partial_\chi). \end{aligned} \quad (0.1.15)$$

using the coordinate basis  $\{\partial_\lambda, \partial_\mu, \partial_\psi, \partial_\chi\}$ . The tetrads construct the metric tensor

$$g = \omega^{+1} \otimes \omega^{+1} + \omega^{-1} \otimes \omega^{-1} + \omega^{+2} \otimes \omega^{+2} - \omega^{-2} \otimes \omega^{-2}. \quad (0.1.16)$$

We can also check the tetrads are the dual basis of the ortho-normal basis:

$$\begin{bmatrix} \omega^{+1} \\ \omega^{-1} \\ \omega^{+2} \\ \omega^{-2} \end{bmatrix}^T [e_{+1} \ e_{-1} \ e_{+2} \ e_{-2}] = \hat{E} \begin{bmatrix} d\lambda \\ d\mu \\ d\psi \\ d\chi \end{bmatrix} [\partial_\lambda \ \partial_\mu \ \partial_\psi \ \partial_\chi] \hat{E}^{-1} = 1. \quad (0.1.17)$$

That is why we can confirm the ortho-normality

$$g(e_j, e_j) = +1 \text{ for } j \in \{+1, -1, +2\}, \ g(e_{-2}, e_{-2}) = -1, \text{ otherwise vanish.} \quad (0.1.18)$$

- Lorentzian Manifold  $M$
- An open set  $U$
- coordinates:  $(\lambda, \mu, \psi, \chi)$
- Independent functions:  $\Delta_\lambda, \Delta_\mu, P_\lambda, P_\mu, Q_\lambda, Q_\mu$
- $Z = P_\lambda Q_\mu - P_\mu Q_\lambda$

$$A = \sum_i \frac{X_{-i}}{Z} ((r) Z_i^r d\varphi^{2r}), \ A = \sum_i \frac{X_{-i}}{(U\Delta_{-i})^{1/2}} \omega^{2i} \quad (0.1.19)$$

$$A = \frac{P_\lambda X_\mu + P_\mu X_\lambda}{Z} d\psi - \frac{Q_\lambda X_\mu + Q_\mu X_\lambda}{Z} d\chi$$

Cartan's first structure formula  $d\theta^\mu + \omega_\nu^\mu \wedge \theta^\nu = \tau^\mu$  with torsion free, i.e.,  $\tau^\mu = 0$  gives conditions of the connection form

$$\omega_\nu^\mu \wedge \theta^\nu = -d\theta^\mu. \quad (0.1.20)$$

We use the interior product  $\iota$  to solve  $\omega_\nu^\mu$ s. The interior product