




# The Next Generation: Refining the Standard Population Synthesis Model to Better Predict Stellar Population Properties

Sean Richards


A thesis submitted in fulfilment of the requirements of the degree of Doctor of  
Philosophy in Physics, the University of Auckland, 2024. This thesis is for  
examination purposes only and is confidential to the examination process.

# Abstract

A short description of the work.


**Cover Image:** The Whirlpool Galaxy (M51). Image is by NASA and released into the public domain:  
[https://images.nasa.gov/details/GSFC\\_20171208\\_Archive\\_e001925](https://images.nasa.gov/details/GSFC_20171208_Archive_e001925)





For my grandparents: Alan and  
Nanette, and Walter and Eileen.

Swing low, sweet chariot.



That's the trouble with science. It's never done.  
Always upending itself. Ruining perfect systems  
for the little inconvenience of them being wrong.

*Zahel*

# Acknowledgements

*Ho there, wanderer! Stay thy course a moment and indulge an old man.*

---

ELMINSTER AUMAR

Waipapa Taumata Rau (The University of Auckland) sits within the rohe of several Māori iwi/hapū, including Ngāti Whātua, Ngāti Whātua o Ōrākei, Te Kawerau a Maki, Ngāti Tamaoho, Te Ākitai Waiohū, Ngāti Maru, Te Patukirikiri, Ngāti Pāoa, Ngāti Tamatera, Ngāi Tai ki Tāmaki, Ngāti Te Ata, and Ngāti Whanaunga, and Waikato-Tainui. Ipipiri (the Bay of Islands) sits within the rohe of Ngāpuhi and Ngāti Rēhia. I acknowledge these iwi and hapū as the tāngata whenua of Tāmaki Makaurau, Ipipiri, and Aotearoa, I acknowledge that sovereignty was never ceded under Te Tiriti O Waitangi, and I pay my respects to kuia and kaumātua of these iwi and hapū, past, present, and future. Tūturu whakamaui, kia tina! Tihei mauri ora!

Throughout my travels for this PhD, I was privileged to visit several lands. In Australia, I worked on lands traditionally occupied by peoples of the Kulin Nations (predominantly the Wurundjeri people), and the Jagera, Yuggera and Ugarapul peoples. I acknowledge their sovereignty and traditional ownership and custodianship of the lands and waterways on which Swinburne University of Technology and the University of Southern Queensland (Springfield) are located. I acknowledge their Elders, past, present, future and emerging, and I acknowledge that sovereignty was never ceded. I thank the kind hospitality of the Polish, Belgian, Singaporean, German, Dutch, and Australian peoples, all of whose lands I visited throughout this thesis. I have learned so much, sitting on the shores of the Eternal Sea that is the Milky Way.

The author wishes to acknowledge the use of New Zealand eScience Infrastructure (NeSI) high performance computing facilities, consulting support and/or training services as part of this research. New Zealand's national facilities are provided by NeSI and funded jointly by NeSI's collaborator institutions and through the Ministry of Business, Innovation & Employment's Research Infrastructure programme. URL <https://www.nesi.org.nz>.

This work made use of v2.2 of the Binary Population and Spectral Synthesis (BPASS) models as described in Eldridge et al. (2017) and Stanway & Eldridge (2018).

*Raise a glass to all of us  
Tomorrow there'll be more of us  
Telling the story of tonight.*

---

FOUND/TONIGHT (BEN PLATT AND LIN-MANUEL MIRANDA)

I am the proud grandchild of immigrants, and by my count, the first person who identifies as Sāmoan to graduate with a PhD from the Department of Physics at the University of Auckland in (at least) 23 years. I'm also one of the few gender non-conforming students in my Department. To all my fellow BIPOC students, all my queer whānau, and all my other underrepresented students in the sciences, I see you, I recognise you, I feel you. Kia kaha, kia māia, kia manawanui.

*Who am I?  
I am a girl who loves my island  
And the girl who loves the sea, it calls me  
I am the daughter of the village chief  
We are descended from voyagers  
Who found their way across the world  
They call me*

---

It would be impossible for me to list my entire whakapapa. It is known as far back as nine generations in parts, tracing back to around the mid-1700s – including to the first Lord Mayor of the City of Sheffield, a Prime Minister of Samoa, a former Minister of Finance and Chair of Samoa’s National Bank, and some of the major clans in Samoa. We suspect there may be an American President in there too. At last count we know of about 450 relatives – my Great-Great-Aunt’s father-in-law had thirteen children in the mid-1800s. But I want to especially mihi to my mother Sue, my late aunt Leslie, my late grandfather Alan, my grandmother Nanette and her late mother Mary “Sugalu” Scott, my aunt Alison and uncle Pete, my great-uncle Ben and great-aunt Lulu, my cousins Jeremy, Jaime, Charlotte, Candice, Ngareta, Wairaka, Marihi, and Tama. To the rest of my ‘aiga: *e lautele le vateatea, ae le o le avanoa iinei: Ou te alofa ia te outou uma.*

*Life before death. Strength before weakness. Journey before destination.*

---

FIRST IDEAL OF THE KNIGHTS RADIANT

For Jaffe, Bailey, Riegel, Ray, Mercer, Johnson, Willingham, and O’Brien: Vox Machina, the Mighty Nein, Bells Hells, Gilmore, Allura, Delilah, Pumat, Xandis, and Ryn. For Iyengar, Carazo, Ray, Riegel, Willingham, Wilson, and Mulligan: the Ring of Brass, Vespene, Maya, and Kir. For Lewis: Peter, Edmund, Susan, Lucy, Aslan and Jadis. For Sanderson: Kaladin, Syl, Shallan, Jasnah, Vin, Elend, Kelsier, Rysn, and Hoid. For Vincke, Wilde, English, Newbon, Downie, Solomon, Béart, Jones, Wiles, Mercer, Gregory, Simmons, Isaacs, and Robertson: Lae’zel, Shadowheart, Astarion, Gale, Wyll, Karlach, Halsin, Jaheira, Minsc, Minthara, Ketheric, Gortash, and Orin the Red. For those who imagine the strange and beautiful and complex and weird and mystical worlds of fantasy, whether betwixt the pages of a book, on the silver screen, or upon the roll of a d20, I pay my enormous regards to you.

Thank you to Nick from Nick’s Kitchen, I am a convert to Adobo from your kai. Thanks also to the crew at Standing Room Espresso, Toasted Cafe, and whoever supplies the beans to the Level 6 Coffee Machine. As Jake Peralta once said, my brain needs its fast juice, and you all supplied that in *spades*.

*Can you imagine a piece of the universe  
More fit for princes and kings?  
I’ll trade you ten of your cities  
For Marion Bridge and the pleasure it brings*

---

*Song for the Mira* BY JOHN McDERMOTT

It occurs to me that there are precious few moments in one’s life to acknowledge and thank the people who contributed to where one is now. I hold the belief that a person is defined by the sum of their actions and interactions, so I would like to acknowledge a few of the people that have made me the person I am today.

I’d like at this point to acknowledge friends: Connor, Frazer, Girish, Hunter, Pierce, Ryan, Scott, and Sean, with whom I shared some of my first physics and mathematics courses. Emma, the self-appointed creative director of this thesis. Josh, Tessa, Hamish, Jasmine, Ryan: my first experience with D&D was with Eldrin. Andrew, George, Hamish, Connor, Callum, and Mark: Katerina was my longest player experience; playing her taught me so much about myself (and Andrew). Josh, Liam, Arthur, Connor, Andrew, Oliver: Ayo and the rest of the Blue Originators (you never learned their party’s name) owe you a debt of gratitude. Even good old Necromancer Bob says hi.

Thank you to my office mates, Max, Bushra, Jonathan, Petra, Hamish, Erin, and Laura. On the topic of Big Dog and Big L: I can’t say what I got told they’d do if I didn’t acknowledge them. They know what they said. But thank you for long walks on beaches, fish and chips, drinks at My Bar (we need to get a better bar), and generally being a welcome distraction whilst writing this thesis. Also, get therapy, the both of you.

I've lost touch with some of the people I did high school with but the formative experiences there still sit inscribed in indelible ink in my mind. Caitlin, who has been one of my best friends for the better part of a quarter-century (and who would *not* let me live it down were I to forget her). Callum, Susannah, Kyle, Grayson, Tomek "if you're going to try to pronounce my last name, put some **EMPHASIS** on it" Pietkiewicz and the entire business and commerce crew. Sam, Louise, Crooks, the IT Ninjas. Isabella, Chyna, Evan, Parth, Jay, Matt, all the people at NZSDC 2015, even if I had to argue in favour of alternative medicine and exclusive schools for LGBTQIA youth, those moments will be with me forever. Finn and Susannah: I know they said they never ranked us in maths classes, but they ranked us in maths classes, and I know we always vied for the top three places. Finn is an architect now and Susannah is admitted to the High Court – does that mean I finally win the ranking?

I'd also like to thank Di Eastwood, Maddy Haydock, Melissa Johansson, and Lauren Mulcahy (our debating coach, and my Year 13, 12, and 11 English teachers). Not all of you are teaching anymore, but I still remember vivid parts of your lessons. Miss Mulcahy commending my essay structure (for not opening an essay with "The question I am answering is..." but actually motivating why I enjoyed *To Kill a Mockingbird* and *Romeo+Juliet*). Miss Johansson, for showing faith in the process of my creative writing portfolio ("Sean, I want to send you to counselling for this story. Change nothing, this is an E."). Mrs Haydock, firstly for Mufasa but also for believing in my weird and experimentalist poetry (I was going through a bit of a fourth-wall breaking phase with poetry), for letting me rant for 30 minutes on why *The Matrix* is actually a Marxist allegory and the (concerningly frequent) themes from Freudian psychoanalysis we can pull from it. Moreover, I remember in English class, you congratulating me for using "moreover" correctly in a sentence. Mrs Eastwood for always challenging me to do better in debating. I've said it before and I will say it again: I credit a mix of you four for my erudition and eloquence.

On the topic of teachers, I would like to mihi to Robbie Lumkong, Andy Dyson, Chris Whitaker. Physics lessons in Year 12 and 13 with Mr Lumkong (and in Year 13 with Mr Dyson) were absolutely batshit crazy – can I say batshit in a thesis? – but it is the undeniable infective enthusiasm for physics that someone who has played international rugby<sup>1</sup> brings that made me even think about doing PHYSICS 120 when I came to university and had a free paper in first year. Mr Whitaker, I still remember sitting on a bus behind you in Year 11 Physical Science and asking you a tonne of questions about the big bang and the standard model.

Right, I'm almost done here, I swear.

Physics is a collaborative endeavor, and I would be remiss to not acknowledge my research compatriots. Thank you to the BPASS team, both in Auckland and abroad: John, Heloise, Max, Sohan, Wouter, Petra, Erin, Adib, Conor, Gareth, Ashley, and Ankur. We've also been joined by a few students who have since left the team (because they graduated!): Gleb, Emma, Louis, Ethan, Siris, Celina, and Amanda. Thank you to our PIs, Dr Elizabeth Stanway and Prof. Jan Eldridge.

Speaking of Jan! Thank you to Prof. Jan Eldridge and Assoc. Prof. Nick Rattenbury. I genuinely cannot image a more perfect supervision team. Both carry a large amount of mana, and I have learned an immense amount from each of them. You'll notice that all of my contractions of people's titles either omit (in the case of "Dr") or contain periods (in the case of basically every other title). Nick was very pedantic about this in my Masters thesis and now I subconsciously check for this when reading other people's prose. Thank you by extension to Prof. Richard Easter. Jan, Nick, and Richard were my lecturers for PHYSICS 107/G in Semester Two of 2016. I needed extra papers to fill out my first year, as there were only three required papers for my (at that time) Computer Science degree. I took PHYSICS 120 and 150 because I was good at it in High School (see Mr

---

<sup>1</sup>This is not a joke or a quip, my Year 13 physics teacher played as a number eight in the 1999 Rugby World Cup for the USA.

Lumkong, Mr Dyson, and Mr Whitaker above), and PHYSICS 107/G because it sounded fun. I left that course a physics double major, and no joke, restructured my entire degree around taking PHYSICS 356. Jan, Nick, and Richard are a large part of why I am a scientist today. I cannot thank them enough for that. From the bottom of my heart, thank you, ngā mihi, vinaka vakalevu, fa'afetai lava.

I'd also like to mihi to a few of my inspirational heroes in science as a whole. Prof. Stephen Hawking, Prof. Kip Thorne, Prof. Brian Greene, Prof. Brian Cox, Dr Derek Miller (Veritasium), Mrs Dianna Covern (PhysicsGirl), Mr Destin Sandlin (SmarterEveryDay), Mr John D. Boswell (melodysheep), and Mr Tim Blais (acapellascience).

*You'll be in my heart.  
No matter what they say.  
You'll be here in my heart.  
Always.*

---

*You'll be in my heart* BY PHIL COLLINS

When I started this PhD, I had two alive grandparents. I now have only one, my grandmother Nanette. For Alan, Walter, and Eileen, I carry your memories with me always. In the words of Liam O'Brien: *I was richer for having known you, friend.*

*I could not be more keen, for some plain old boring precedented times.*

---

CR. RICHARD HILLS

I've noticed that the more theses I write, the more people I thank. My PGDipSci dissertation thanked five people. My Masters thesis thanked some fifty-five, a Government, and several coffee companies. Did I beat it this time?

*Run, you clever [girl], and be a Doctor.*

---

CLARA OSWIN OSWALD, *Doctor Who*

*No, thank YOU. One thing about trains... it doesn't  
matter where they're going. What matters is  
deciding to get on.*

---

THE CONDUCTOR, *The Polar Express*



# Contents

|                                                                    |              |
|--------------------------------------------------------------------|--------------|
| <b>Contents</b>                                                    | <b>xi</b>    |
| <b>List of Figures</b>                                             | <b>xv</b>    |
| <b>List of Tables</b>                                              | <b>xvii</b>  |
| <b>Glossary</b>                                                    | <b>xix</b>   |
| <b>Thesis To-do List</b>                                           | <b>xxiii</b> |
| <b>1 Introduction</b>                                              | <b>1</b>     |
| 1.1 Observational astronomy . . . . .                              | 2            |
| 1.2 Stellar Modelling . . . . .                                    | 4            |
| 1.2.1 Population Synthesis Codes . . . . .                         | 5            |
| Detailed Population Synthesis . . . . .                            | 5            |
| Rapid Population Synthesis . . . . .                               | 5            |
| 3D MHD Simulations . . . . .                                       | 5            |
| 1.3 This work . . . . .                                            | 6            |
| 1.3.1 Research aims . . . . .                                      | 6            |
| 1.3.2 Thesis outline . . . . .                                     | 6            |
| <b>2 Theoretical Background</b>                                    | <b>9</b>     |
| 2.1 Stellar Formation . . . . .                                    | 9            |
| 2.2 Stellar Evolution . . . . .                                    | 10           |
| 2.2.1 The chemical composition of stars . . . . .                  | 11           |
| 2.2.2 Evolutionary Pathways . . . . .                              | 11           |
| Low-Mass Stars: $0.08 M_{\odot} < M < 2 M_{\odot}$ . . . . .       | 11           |
| Intermediate-Mass Stars: $2 M_{\odot} < M < 8 M_{\odot}$ . . . . . | 12           |
| Massive Stars: $8 M_{\odot} < M < 80 M_{\odot}$ . . . . .          | 13           |
| 2.2.3 Relativistic corrections . . . . .                           | 13           |
| 2.2.4 Stellar Atmospheres . . . . .                                | 16           |
| 2.3 Supernovae . . . . .                                           | 16           |
| 2.4 Two to tango: Binary Evolution . . . . .                       | 18           |
| 2.4.1 Binary star kinematics . . . . .                             | 18           |
| 2.4.2 Roche Lobes . . . . .                                        | 19           |
| The $\mathcal{L}_1$ point . . . . .                                | 22           |
| The $\mathcal{L}_2$ point . . . . .                                | 23           |
| 2.4.3 Mass Transfer . . . . .                                      | 24           |
| Wind mass-loss . . . . .                                           | 24           |
| Roche Lobe overflow (RLOF) . . . . .                               | 26           |
| Accretion . . . . .                                                | 28           |
| Bondi-Hoyle accretion . . . . .                                    | 28           |

|          |                                                                                    |           |
|----------|------------------------------------------------------------------------------------|-----------|
| 2.5      | Post-Supernova . . . . .                                                           | 28        |
| 2.5.1    | Merger Rates of compact binaries . . . . .                                         | 29        |
| 2.6      | Numerical implementation . . . . .                                                 | 30        |
| 2.6.1    | The Cambridge STARS code . . . . .                                                 | 30        |
|          | Brief History . . . . .                                                            | 30        |
|          | Matrix Inversion . . . . .                                                         | 31        |
|          | Initial Condition Selection . . . . .                                              | 32        |
|          | Mesh spacing . . . . .                                                             | 32        |
|          | Timestep control . . . . .                                                         | 33        |
|          | Numerical instabilities . . . . .                                                  | 33        |
|          | Critical modifications . . . . .                                                   | 34        |
| 2.6.2    | The TUI code . . . . .                                                             | 36        |
| 2.6.3    | The KAITIAKI grid-watcher code . . . . .                                           | 37        |
| 2.7      | Chapter Summary . . . . .                                                          | 38        |
| <b>3</b> | <b>Detailed Models of Overcontact Binary Stars</b>                                 | <b>39</b> |
| 3.1      | Observations of contact binaries . . . . .                                         | 39        |
| 3.2      | Data matching with BPASS v2.2 . . . . .                                            | 39        |
| 3.2.1    | Progenitor genealogy . . . . .                                                     | 39        |
|          | LSS 3074 . . . . .                                                                 | 39        |
|          | MY Camelopardalis . . . . .                                                        | 41        |
|          | SMC 108086 . . . . .                                                               | 41        |
|          | TU Muscae . . . . .                                                                | 41        |
|          | V382 Cygni . . . . .                                                               | 41        |
|          | VFTS 352 . . . . .                                                                 | 43        |
| 3.3      | Forward evolution . . . . .                                                        | 43        |
| 3.4      | Exotic binaries arising from contact systems . . . . .                             | 43        |
| 3.4.1    | Double contact systems . . . . .                                                   | 43        |
| 3.4.2    | Reverse mass transfer . . . . .                                                    | 43        |
| 3.4.3    | Overflow through $\mathcal{L}_2$ . . . . .                                         | 43        |
| <b>4</b> | <b>Helium Rich Mass Transfer</b>                                                   | <b>45</b> |
| 4.1      | Evolution overview . . . . .                                                       | 45        |
| 4.1.1    | Pre-interaction . . . . .                                                          | 45        |
| 4.1.2    | Pre-contact . . . . .                                                              | 45        |
| 4.1.3    | Contact phase . . . . .                                                            | 48        |
| 4.1.4    | Post-contact . . . . .                                                             | 48        |
| 4.1.5    | Post-interaction . . . . .                                                         | 48        |
| 4.2      | Atypical abundance profiles . . . . .                                              | 48        |
| 4.3      | Comparison between variable and non-variable accretion . . . . .                   | 48        |
| 4.4      | Chapter Summary . . . . .                                                          | 50        |
| <b>5</b> | <b>Population properties of overcontact binaries and helium rich mass transfer</b> | <b>53</b> |
| <b>6</b> | <b>Post-Supernova Kinematics: The Natal Kick</b>                                   | <b>55</b> |
| 6.1      | Overview . . . . .                                                                 | 55        |
| 6.1.1    | Parameter space . . . . .                                                          | 56        |

|          |                                                                                 |           |
|----------|---------------------------------------------------------------------------------|-----------|
| 6.1.2    | Natal Kick Prescriptions . . . . .                                              | 58        |
| 6.1.3    | The TAKAHE orbital evolution code . . . . .                                     | 58        |
| 6.2      | Merger Rates of BNS systems . . . . .                                           | 59        |
| 6.3      | Period-Eccentricity Distributions . . . . .                                     | 60        |
| 6.4      | Single-Star Kick Velocities . . . . .                                           | 65        |
| 6.4.1    | Pulsar velocity dataset . . . . .                                               | 65        |
| 6.4.2    | Comparison . . . . .                                                            | 65        |
| 6.5      | Production of Ultra-Stripped Supernovae . . . . .                               | 67        |
| 6.6      | Refinement of the parameter space . . . . .                                     | 69        |
| 6.7      | Discussion . . . . .                                                            | 73        |
| 6.8      | Chapter Summary . . . . .                                                       | 74        |
| <b>7</b> | <b>Conclusion</b>                                                               | <b>75</b> |
| 7.1      | Summary . . . . .                                                               | 75        |
| 7.2      | Limitations . . . . .                                                           | 75        |
| 7.3      | Suggested Future Work . . . . .                                                 | 75        |
|          | <b>Appendices</b>                                                               | <b>77</b> |
| <b>A</b> | <b>Code Samples</b>                                                             | <b>79</b> |
| <b>B</b> | <b>Code Versions</b>                                                            | <b>83</b> |
| <b>C</b> | <b>Period-Eccentricity distributions along the LVK Collaboration O3 contour</b> | <b>85</b> |
| <b>D</b> | <b>Parameter Configuration</b>                                                  | <b>87</b> |
|          | <b>Bibliography</b>                                                             | <b>91</b> |



# List of Figures

|      |                                                                                                                                                                         |    |
|------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| 1.1  | The locations of the $\sim 9000$ stars in the night sky visible to the naked eye. . . . .                                                                               | 2  |
| 2.1  | The evolution of an $4 M_{\odot}$ star from ZAMS until CO core development. . . . .                                                                                     | 12 |
| 2.2  | The evolution of stars in the range $10 - 80 M_{\odot}$ . . . . .                                                                                                       | 13 |
| 2.3  | The evolution of an $8 M_{\odot}$ star. . . . .                                                                                                                         | 14 |
| 2.4  | A rough outline of supernova classification. . . . .                                                                                                                    | 17 |
| 2.5  | The binding energy per nucleon for elements up to Uranium-238. . . . .                                                                                                  | 18 |
| 2.6  | A schematic of a simple binary star system. . . . .                                                                                                                     | 19 |
| 2.7  | The dimensionless Roche potential as a function of mass ratio $q$ . . . . .                                                                                             | 20 |
| 2.8  | A schematic of the Roche lobe formula Equation 2.30. . . . .                                                                                                            | 22 |
| 2.9  | The volume of the Roche Lobe computed numerically through Monte-Carlo integration vs the<br>volume assuming Equation 2.30. . . . .                                      | 23 |
| 2.10 | As for Figure 2.9, but for the $\mathcal{L}_2$ point. . . . .                                                                                                           | 23 |
| 2.11 | An illustration of the bi-stability jump at $\log T \approx 4.4$ . . . . .                                                                                              | 27 |
| 2.12 | Two stars of a similar mass that undergo a helium flash and two thermal pulses. . . . .                                                                                 | 35 |
| 2.13 | The helium flash shown on a Kippenhahn diagram. . . . .                                                                                                                 | 36 |
| 3.1  | BPASS v2.2.1 primary models for the matches obtained by HOKI for the parameters in Table 3.1. .                                                                         | 40 |
| 4.1  | The main evolution of a binary with variable accretion enabled. . . . .                                                                                                 | 46 |
| 4.2  | As for Figure 4.1, but absent variable accretion. . . . .                                                                                                               | 47 |
| 4.3  | The evolution of the radius, rotation rate, and luminosity in the presence of variable accretion,<br>shown around the contact phase. . . . .                            | 48 |
| 4.4  | The chemical composition of each star as a function of the stellar radius for the donor and accretor<br>at the five key stages of the binary's lifetime. . . . .        | 50 |
| 4.5  | An HR diagram showing the evolution of our model both with and without the variable accretion<br>prescription. . . . .                                                  | 51 |
| 6.1  | A schematic of how FINEGRID and WIDEGRID are related. . . . .                                                                                                           | 57 |
| 6.2  | The effect of varying $d_e$ in Equation 6.5 on the coalescence time. . . . .                                                                                            | 60 |
| 6.3  | The effect of varying $d_e$ in Equation 6.5 on the runtime and performance. . . . .                                                                                     | 61 |
| 6.4  | The merger rate as a function of kick parameters $\alpha$ and $\beta$ . . . . .                                                                                         | 62 |
| 6.5  | The period-eccentricity distributions for six selected Bray & Eldridge (2016) kicks. . . . .                                                                            | 63 |
| 6.6  | The $\ln \mathcal{K}$ values across $\alpha - \beta$ space. . . . .                                                                                                     | 64 |
| 6.7  | The PDF and CDF of the distributions using kicks from Hobbs et al. (2005); Verbunt et al. (2017);<br>Bray & Eldridge (2018); Willcox et al. (2021). . . . .             | 66 |
| 6.8  | The K-S statistic for FINEGRID and WIDEGRID. . . . .                                                                                                                    | 67 |
| 6.9  | The PDFs and CDFs of four systems, alongside the Hobbs CDF and Willcox et al. (2021) CDF. . .                                                                           | 68 |
| 6.10 | The four constraints we apply overlaid on our FINEGRID parameter space. . . . .                                                                                         | 70 |
| 6.11 | The CDF for a population of isolated pulsars and the period-eccentricity distribution for $\alpha =$<br>$115 \text{ km s}^{-1}, \beta = 15 \text{ km s}^{-1}$ . . . . . | 71 |

|      |                                                                                                                 |    |
|------|-----------------------------------------------------------------------------------------------------------------|----|
| 6.12 | The peak of the event rate $\mathcal{R}(t)$ as a function of $\alpha$ and $\beta$ . . . . .                     | 71 |
| 6.13 | The event rate $\mathcal{R}(z)$ for a region surrounding $\alpha = 330, \beta = -70$ on FINEGRID. . . . .       | 72 |
| C.1  | The period-eccentricity distributions of populations governed by Bray kicks across the LIGO O3 contour. . . . . | 86 |

# List of Tables

|     |                                                                                                                                                                                                                                                                                                                                                                                                          |    |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| 2.1 | The mass loss rate constants for interpolation around the bistability jump. . . . .                                                                                                                                                                                                                                                                                                                      | 25 |
| 3.1 | Search parameters for the six close-contact binaries compiled in Abdul-Masih et al. (2022), as used in our fitting to the BPASS v2.2.1 grid. . . . .                                                                                                                                                                                                                                                     | 41 |
| 3.2 | Matches from the HOKI (Stevance et al., 2020a) software. . . . .                                                                                                                                                                                                                                                                                                                                         | 42 |
| 4.1 | The key evolutionary parameters of both stars and the binary configuration at five critical points: the initial parameters, and the beginning and ends of both epochs of Roche Lobe overflow (RLOF). All values are rounded to 2 decimal places. The letter in parentheses represents which segment of Figure 4.4a (for Table 4.1a) or Figure 4.4b (for Table 4.1b) the abundances are given in. . . . . | 49 |
| 6.1 | The populations under study. . . . .                                                                                                                                                                                                                                                                                                                                                                     | 56 |
| 6.2 | The systems we use in our study. . . . .                                                                                                                                                                                                                                                                                                                                                                 | 64 |
| B.1 | The code versions we use in this thesis. *TUI was described in earlier papers, but Ghodla et al. (2023) was the first to name it. . . . .                                                                                                                                                                                                                                                                | 83 |
| D.1 | Our general initial conditions. . . . .                                                                                                                                                                                                                                                                                                                                                                  | 87 |



# Glossary

---

| Notation | Description                               |
|----------|-------------------------------------------|
| 3D MHD   | 3D Magnetohydrodynamics.                  |
| AGB      | Asymptotic Giant Branch.                  |
| BH       | Black Hole.                               |
| BNS      | Binary Neutron Star.                      |
| BPASS    | Binary Population and Spectral Synthesis. |
| CCSN     | core-collapse supernovae.                 |
| CDF      | Cumulative Distribution Function.         |
| CEE      | Common Envelope Evolution.                |
| CME      | coronal mass ejection.                    |
| CO core  | Carbon-Oxygen core.                       |
| CPTA     | Chinese Pulsar Timing Array.              |
| DTD      | Delay-Time Distribution.                  |
| EPTA     | European Pulsar Timing Array.             |
| GR       | General Relativity.                       |
| IMF      | Initial Mass Function.                    |
| ISCO     | innermost stable circular orbit.          |
| ISM      | Interstellar Medium.                      |
| JWST     | James Webb Space Telescope.               |

---

| Notation          | Description                                                   |
|-------------------|---------------------------------------------------------------|
| K-S               | Kolmogorov-Smirnov.                                           |
| LIGO              | Laser Interferometer Gravitational Wave Observatory.          |
| LISA              | Laser Interferometer Space Antenna.                           |
| LMC               | Large Magellanic Cloud.                                       |
| LVK Collaboration | LIGO-Virgo-KAGRA Collaboration.                               |
| MLT               | Mixing Length Theory.                                         |
| MWEG              | Milky-Way Equivalent Galaxy.                                  |
| NANOGrav          | North American Nanohertz Observatory for Gravitational Waves. |
| NS                | Neutron Star.                                                 |
| NS-BH             | Neutron Star - Black Hole Binary.                             |
| O3                | Operating Run 3.                                              |
| O4                | Operating Run 4.                                              |
| ONe core          | Oxygen-Neon core.                                             |
| PDF               | Probability Density Function.                                 |
| PPTA              | Parkes Pulsar Timing Array.                                   |
| RLOF              | Roche Lobe overflow.                                          |
| SFH               | Stellar Formation History.                                    |
| SFRD              | Star Formation Rate Density.                                  |
| SMC               | Small Magellanic Cloud.                                       |
| SPH               | Smoothed-particle hydrodynamics.                              |
| TAMS              | Terminal-Age Main Sequence.                                   |

| Notation | Description               |
|----------|---------------------------|
| USSN     | ultra-stripped supernova. |
| WD       | White Dwarf.              |
| WR       | Wolf Rayet.               |
| ZAMS     | Zero-Age Main Sequence.   |



# Thesis To-do List

|                                                                |    |
|----------------------------------------------------------------|----|
| 2.1 but why? “its what the code does” is a bad reason. . . . . | 32 |
| D.2 Richards et al 2024 . . . . .                              | 87 |
| D.3 Richards et al 2024 . . . . .                              | 87 |
| D.4 Richards et al 2024 . . . . .                              | 89 |
| D.5 cite >:) . . . . .                                         | 89 |

## Co-Authorship Form

Email:

This form is to accompany the submission of any PhD that contains published or unpublished co-authored work. **Please include one copy of this form for each co-authored work.** Completed forms should be included in all copies of your thesis submitted for examination and library deposit (including digital deposit), following your thesis Acknowledgements. Co-authored works may be included in a thesis if the candidate has written all or the majority of the text and had their contribution confirmed by all co-authors as not less than 65%.

Please indicate the chapter/section/pages of this thesis that are extracted from a co-authored work and give the title and publication details or details of submission of the co-authored work.

*Parts of Chapter 4 are based on "The variable evolution of accretor stars in binary systems due to accretion of increasingly helium-rich material", submitted to MNRAS and available on the NASA ADS here:*  
<https://ui.adsabs.harvard.edu/abs/2024arXiv241103000R/abstract>

|                                             |                                                 |  |
|---------------------------------------------|-------------------------------------------------|--|
| Nature of contribution by PhD candidate     | Performed data analysis, wrote majority of text |  |
| Extent of contribution by PhD candidate (%) | 70                                              |  |


### CO-AUTHORS

| Name         | Nature of Contribution                                |
|--------------|-------------------------------------------------------|
| Jan Eldridge | Wrote part of text, criticism, commentary, and review |
| Max Briel    | Criticism, commentary, and review                     |
| Sohan Ghodla | Criticism, commentary, and review                     |
|              |                                                       |

### Certification by Co-Authors

The undersigned hereby certify that:

- ❖ the above statement correctly reflects the nature and extent of the PhD candidate's contribution to this work, and the nature of the contribution of each of the co-authors; and
- ❖ that the candidate wrote all or the majority of the text.

| Name         | Signature                                                                            | Date       |
|--------------|--------------------------------------------------------------------------------------|------------|
| Jan Eldridge |  | 11/11/2024 |
| Max Briel    |   | 19/11/2024 |
| Sohan Ghodla |   | 11/11/2024 |
|              |                                                                                      |            |

## Co-Authorship Form

Email:

This form is to accompany the submission of any PhD that contains published or unpublished co-authored work. **Please include one copy of this form for each co-authored work.** Completed forms should be included in all copies of your thesis submitted for examination and library deposit (including digital deposit), following your thesis Acknowledgements. Co-authored works may be included in a thesis if the candidate has written all or the majority of the text and had their contribution confirmed by all co-authors as not less than 65%.

Please indicate the chapter/section/pages of this thesis that are extracted from a co-authored work and give the title and publication details or details of submission of the co-authored work.

*Chapter 6 is based on work from "New constraints on the Bray conservation-of-momentum natal kick model from multiple distinct observations", published in the Monthly Notices of the Royal Astronomical Society (MNRAS), here: [dx.doi.org/10.1093/mnras/stad977](https://doi.org/10.1093/mnras/stad977)*

|                                             |                                                                                                                                    |
|---------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|
| Nature of contribution by PhD candidate     | Wrote majority of the paper, developed most parts of the testing pipeline, later data generation, performed the statistical tests. |
| Extent of contribution by PhD candidate (%) | 90                                                                                                                                 |


### CO-AUTHORS

| Name             | Nature of Contribution                                                                                                                                   |
|------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Jan Eldridge     | Original data generation, guidance on project, wrote clarification on the difference between BPASS primary and secondary models, criticism on manuscript |
| Max Briel        | Developing ideas, code commentary, criticism on manuscript                                                                                               |
| Héloïse Stevance | Developing the test for Ultra-Stripped Supernovae, guidance on project, criticism on manuscript                                                          |
| Reinhold Willcox | Providing single-star pulsar dataset from Willcox+ 2021, criticism on manuscript                                                                         |

### Certification by Co-Authors

The undersigned hereby certify that:

- ❖ the above statement correctly reflects the nature and extent of the PhD candidate's contribution to this work, and the nature of the contribution of each of the co-authors; and
- ❖ that the candidate wrote all or the majority of the text.

| Name             | Signature                                                                            | Date       |
|------------------|--------------------------------------------------------------------------------------|------------|
| Jan Eldridge     |  | 11/11/2024 |
| Max Briel        |   | 19/11/2024 |
| Héloïse Stevance |  | 21/11/2024 |
| Reinhold Willcox |  | 21/11/2024 |



CHAPTER 1

---

# Introduction

*Okay then. That's what I'll do. I'll tell you a story.*

---

THE ELEVENTH DOCTOR

**A**RGUABLY THE GREATEST ADVANCEMENT in modern astronomy was the direct detection of gravitational waves – the rippling of space-time as a result of the motion of massive celestial bodies – on September 14, 2015. The Laser Interferometer Gravitational Wave Observatory (LIGO) collaboration detected the chirp of merging binary black holes, with masses  $m_1 = 36^{+5}_{-4} M_\odot$  and  $m_2 = 29^{+4}_{-4} M_\odot$  (Abbott et al., 2016). This detection sparked the so-called “gravitational wave era”, with detections in Operating Run 4 (O4) reading a cadence of hundreds per year. These detections afford researchers probes of various physical phenomena. For instance, these merger events are some of the most energetic events in the Universe (GW150914 released over 50 times the energy of all stars in the observable Universe, per Abbott et al. (2016)), and the merger of binary neutron stars are the predominant formation channels for most of the elements heavier than iron (see, among others, Eichler et al., 1989; Arcavi et al., 2017; Pian et al., 2017; Smartt et al., 2017).

In addition, the stochastic background of gravitational waves is a direct probe on early-Universe cosmology (Christensen, 2018), though the frequency band that these gravitational waves span is between  $10^{-9}$  and  $10^{14}$  Hz (Christensen, 1992, 2018), making detection of the stochastic background difficult. Indeed, it took eight years after the detection of gravitational waves for the stochastic background to be detected by the collaborative effort of North American Nanohertz Observatory for Gravitational Waves (NANOGrav) (Agazie et al., 2023), the European Pulsar Timing Array (EPTA) (Antoniadis et al., 2023), the Parkes Pulsar Timing Array (PPTA) (Reardon et al., 2023), and the Chinese Pulsar Timing Array (CPTA) (Xu et al., 2023).

Any attempt at understanding gravitational waves must be founded in a strong theoretical understanding of stellar structure and evolution. Without theory, any observation is no more than a curiosity; it is with theoretical underpinning and models that insight can be gained. As most astrophysical sources of gravitational waves (i.e., not the stochastic background) are formed by the merger of two “compact remnants” – black holes, white dwarfs, and neutron stars – one must understand their formation and dynamics, and thus the evolution of progenitor stars into these remnants, as well.

Furthermore, a sound understanding of the evolution of stars has impacts beyond gravitational wave astronomy. As the interiors of stars are complex nuclear furnaces, they are responsible for producing most of the elements in the Universe heavier than Hydrogen (Burbidge et al., 1957). In fact, of the elements lighter than Uranium, the only elements not produced in stellar processes (such as through kilonovae or in supernovae) are technetium, promethium (which both lack stable isotopes), boron, beryllium (which both are produced via cosmic ray fission), and some formation channels of lithium.


From the perspective of Universal building blocks, stars are also the focal point of planetary systems, and provide the theoretical right conditions for the formation of life. Without a solid theory of stellar evolution, one is unable to describe the likelihood of extraterrestrial life within the Universe.

## 1.1 | Observational astronomy

*If the stars should appear one night in a thousand years, how would  
men believe and adore; and preserve for many generations the  
remembrance of the city of God which had been shown!*

*Nature*, BY RALPH WALDO EMERSON

There are approximately 9000 stars visible in the night sky to the naked eye (with magnitude lower than 6.5), across both hemispheres (Hoffleit, 1964), which are displayed in Figure 1.1. The dark-adapted eye can see approximately as far as  $\eta$  Carinae,<sup>1</sup> which is only approximately 2300 pc away.



**Figure 1.1:** The locations of the  $\sim 9000$  stars in the night sky visible to the naked eye, as reported in Hoffleit (1964). The colors are computed by converting the B-V color indices to temperatures using the fits in Ballesteros (2012), and then converted to RGB via the algorithm in this StackOverflow post: <https://stackoverflow.com/a/45497817>. The Southern Cross (Crucis) is shown fully opaque in blue towards the bottom left of the plot as a point of reference, and  $\eta$  Carinae is shown fully opaque in black in the bottom right.

2300 pc pales in comparison to the size of the Galaxy, which is approximately 26.8 kpc in diameter (Goodwin et al., 1998), which in turn pales in comparison to the Local Group at  $\sim 3$  Mpc (Redd, 2018), and the observable Universe at 28.5 Gpc (Bars & Terning, 2010). There are, in addition,  $\sim 1.7 \times 10^{22}$  stars in the observable Universe, assuming  $1.7 \times 10^{11}$  galaxies, with approximately  $10^{11}$  stars in each (Marov, 2015). Thus, the birth, evolution, and death of stars is of fundamental importance to understanding the local structure and evolution of astrophysical objects on any scale.

Given the number of stars visible in the vast expanse of night, and their relatively stable positions in the sky, it is no surprise that they are a fixture in many religions and cultures around the world. Ancient Greece saw a difference in movement of some celestial objects, and called these  $\pi\lambda\alpha\nu\eta\tau\eta\varsigma$ , which morphed into *planētes* and eventually the English word “planets”. Babylonia knew of Halley’s comet and tracked observations of it as late as

<sup>1</sup>This is a debated topic; we use the definition that a “star” includes binaries and that the star is in the catalogue of Hoffleit (1964), which rules out V762 Cas, the other commonly held “farthest star”.

87 BCE (Stephenson et al., 1985). Records of eclipses in Chinese culture exist as far back as 1600 BCE (Fang, 2015). Knowledge of the movements of stars guided Polynesian wayfarers as they crossed the South Pacific (Buck, 1938; Hamacher, 2022), were embedded in the cultural mythos of First Nations peoples in Australia (Noon & De Napoli, 2022), allowed them to forecast weather patterns (Hamacher et al., 2019), and were plausibly recorded into oral histories (Fuller & Hamacher, 2017). Even in the arts, the knowledge of the lunar perigee was written into *Othello* by Shakespeare:

It is the very error of the moon;  
She comes more near the earth than she was wont,  
And makes men mad.

– Shakespeare (ca. 1603)

Knowledge of oceanic tides existed in the time of Ancient Greece, and was described by Ptolemy, who wrote in *Tetrabiblos* that

ἀμπώτεις καὶ αἱ παλίρροιαί, καὶ αἱ τῶν πνευμάτων  
δὲ τροπαὶ μάλιστα περὶ τὰς τοιαύτας τῶν φωτῶν  
κεντρώσεις ἀποτελοῦνται πρὸς οὓς ἂν τῶν ἀνέμων  
ἐπὶ τὰ αὐτὰ τὸ πλάτος τῆς σελήνης τὰς  
προσνεύσεις ποιούμενον καταλαμβάνηται.

Ptolemy & Proclus (ca. 168)

*For the hour by hour intensifications and  
relaxations of the weather vary in response to  
such positions of the stars as these, in the same  
way that the ebb and flow of the tide respond to  
the phases of the moon, and the changes in the  
air-currents are brought about especially at such  
appearances of the luminaries at the angles, in  
the direction of those winds towards which the  
latitude of the moon is found to be inclining.*

Robbins (1940)

This knowledge flowed through to Dante, who wrote in the third part of the Divine Comedy, *Paradiso*:

*E come 'l volger del ciel de la luna  
cuopre e discuopre i liti sanza posa,  
così fa di Fiorenza la Fortuna:*

Alighieri (ca. 1317)

*And, as the turning of the lunar sphere covers  
and endlessly uncovers the edges of the shore,  
thus does fortune deal with Florence.*

Corbett & Webb (2016)

In religious scripture, celestial navigation has oft been a fixture, such as in Chapter 2 of the Gospel according to Matthew:<sup>2</sup>

2:7 – Then Herod, when he had priuily called the Wise men, enquired of them diligently what time the Starre appeared:

2:8 – And he sent them to Bethlehem, and said, Goe, and search diligently for the yong child, and when ye haue found him, bring me word againe, that I may come and worship him also.

<sup>2</sup>This text is as reproduced from the King James Version of the Bible, as written in Early Modern English as faithful as possible to the script.

2:9 – When they had heard the King, they departed, and loe, the Starre which they saw in the East, went before them, till it came and stood ouer where the young childe was.

– Various Authors (1611)

Despite this deep cultural connection to the stars, what we would now call Astronomy is a somewhat recent invention. Arguably it arose in the 16th century CE with Copernicus proposing the heliocentric model of the solar system, which was confirmed by Galileo in the 17th century CE (Carroll & Ostlie, 2017) – although Galileo’s claims did result in his being convicted of heresy for 300 years.

The early 20th century saw a veritable renaissance in both the observational and theoretical spheres. On the one hand, developments such as the theories of Special Relativity in 1905 (Einstein, 1905), General Relativity (GR) in 1917 (Einstein, 1917), and the predictions of gravitational waves (Poincaré, 1906; Einstein, 1916) led to theoretical revolutions. On the other hand, the innovations of Payne (1925) in showing that stars are composed of mostly Hydrogen and Helium, Baade & Zwicky (1934) in distinguishing “super-novae” from “common novae”, Hoyle (1946) in developing the foundation of the nuclear reaction of elements heavier than Hydrogen in stars, Salpeter (1955) in developing the Initial Mass Function (IMF), Burbidge et al. (1957) in truly developing the theory of stellar nucleosynthesis, Henyey et al. (1955) and Hayashi (1961) in exploring the pre-main sequence evolution of stars were key to kickstarting entire branches of astronomy such as spectroscopy and computational astrophysics.

The pace of discovery of new and exotic astronomical phenomena has not slowed into the 21st century. Thorne-Żytkow objects – giant stars with Neutron star cores (Thorne & Żytkow, 1977) – once only existed as theoretical curiosities, yet now have credible observational candidates in Levesque et al. (2014). The aforementioned gravitational waves were detected in 2015 (LIGO Scientific Collaboration and Virgo Collaboration, 2016), a century after they were demonstrated by Einstein (1916) to be a central prediction of GR. More recently, Lau et al. (2022) directly imaged 17 carbon-rich dust rings around WR140 with James Webb Space Telescope (JWST), enriching the Interstellar Medium (ISM) with carbon-rich compounds. In the coming years, the Vera C. Rubin Observatory (Ivezić et al., 2019) and Laser Interferometer Space Antenna (LISA) are expected to vastly increase the number of short-period eclipsing double white dwarfs (Burdge et al., 2020), with the latter able to detect gravitational waves with frequencies between 0.1 mHz and 1 Hz (Amaro-Seoane et al., 2012), compared to the  $\sim 10$  Hz – 1000 Hz that LIGO-Virgo-KAGRA Collaboration (LVK Collaboration) can detect (Schmitz, 2021). The stochastic gravitational-wave background was detected by Agazie et al. (2023); Antoniadis et al. (2023); Reardon et al. (2023); Xu et al. (2023), via a search of pulsar-timing arrays.

Astronomy as a field has blossomed since the naked-eye era – entering the era of the telescope, the era of the observatory, and now the gravitational wave era. The next era of astronomy is the big data era. With observations reaching sub-day cadences, a large array of theoretical models, and perhaps the development of “on-demand” detailed models is required.

## 1.2 | Stellar Modelling

Modelling the evolution of a star is a nontrivial endeavour. Stars as objects obey General Relativity (GR) owing to their masses and sizes, and reach internal temperatures, pressures, and densities unsustainable on Earth. For instance, the core of a star is capable of nuclear fusion, which has not been sustainably recreated on Earth. Moreover, the density of Sirius B – a White Dwarf (WD) in the constellation of Canis Major – is  $2.14 \times 10^{10} \text{ kg m}^{-3}$  (Barstow et al., 2005) and the mean density of the Sun is  $\sim 1410 \text{ kg m}^{-3}$  (Eddy & Ise, 1979). On Earth, the highest density measured in a laboratory is  $\sim 27\,000 \text{ kg m}^{-3}$ , requiring pressures on the order of 2.3 TPa (Fratanduono et al., 2020), and observed for only nanoseconds.

Because of these factors, one often turns to computer simulations to model the interiors of stars, as it is nigh impossible to achieve and sustain these conditions to experimentally test any physical phenomena. That said, missions such as the Parker Solar Probe have made flyby missions into the atmosphere of the Sun, though broadening the sample size to two stars – using Proxima Centauri, the nearest star to Earth aside from the Sun – would require travelling at approximately  $0.1c$  for 40 years to reach. If instead we were to send the Parker Solar Probe travelling at a constant velocity of its top velocity of  $\sim 176\,471\text{ m s}^{-1}$ , it would take approximately 7200 years to reach Proxima Centauri. Needless to say, physical experimentation with a star is difficult.

### 1.2.1 | Population Synthesis Codes

The diversity and complexity of stellar profiles, interiors, and characteristics discussed earlier demands a careful and varied approach. One cannot abstract away much of the important physics. Although it is common to use approximations such as spherical symmetry to reduce the dimensionality of the problem, and to only evaluate the equations of stellar structure at a fixed number of locations in this one remaining dimension, chosen sufficiently intelligently, a complex computational effort must be undertaken.

Broadly speaking, there are three approaches to modelling stars, all of which have both merits and drawbacks. These three methods are:

1. Rapid Population Synthesis,
2. Detailed Population Synthesis, and
3. 3D Magnetohydrodynamics (3D MHD) Simulations.

Each of these approaches informs the other two, and all play important roles in stellar modelling. Although we focus on detailed methods, we will briefly outline all three approaches here.

#### Detailed Population Synthesis

Detailed stellar models explicitly solve the equations of hydrostatic equilibrium, usually via some form of matrix-inversion method. For instance, as will be discussed in subsection 2.6.1, the Cambridge STARS code inverts the Henyey matrix  $\mathbf{H}_{ij} = \frac{\partial(\text{equation } i)}{\partial(\text{variable } j)}$ . MESA (Paxton et al., 2011, 2013, 2015, 2018, 2019; Jermyn et al., 2023) uses a Newton-Raphson solver in order to converge to a solution. In particular, we use a variant of the Cambridge STARS code, which solves the equations of *both* stars simultaneously.

#### Rapid Population Synthesis

Rapid population synthesis methods exploit the statistical properties of stars in order to rapidly infer properties of populations of stars. However, they still require as input detailed population synthesis models – for example, COMPAS (Team COMPAS: Riley et al., 2022) uses MESA (Paxton et al., 2011, 2013, 2015, 2018, 2019; Jermyn et al., 2023) models as its input models.

#### 3D MHD Simulations

3D Magnetohydrodynamics (3D MHD) and Smoothed-particle hydrodynamics (SPH) simulations are the most detailed method of stellar modelling. The base principle is that the star is evaluated as a network of particles in a continuous media, and the evolution of the star is followed in theoretically atomic detail. For example, PHANTOM (Price et al., 2018) is a SPH code which essentially models the star as a fluid, where the Lagrangian hydrodynamical equations are solved for in two-dimensional cells (termed “particles”). Although these present the most detailed models, they are also the most time-intensive to solve. For a SPH code, the famous Navier-Stokes

equation is invoked, which has no analytical solution but is computed in excruciating detail. As a result, most SPH models take weeks to run and simulate a star for kiloseconds of its lifetime, rather than for the millions (if not billions) of years detailed models are able to compute.

Though it may seem that they serve the same niche and are therefore competing to be the “better” method of stellar modelling, detailed 1D models and 3D MHD/SPH models exist harmoniously with each other. The advancements in 3D MHD/SPH modelling inform the evolution of 1D models in a process known as “3-to-1”, and the computational flexibility of 1D models offers insights into where 3D models wish to target to evolve, in a lock-step advancement method.

## 1.3 | This work

*Come with me, and you'll be, in a world of pure imagination.*

*Pure Imagination*, BY GENE WILDER

### 1.3.1 | Research aims

There are essentially three major themes guiding this research.

- (I) Can we methodologically overcome the typical cases in which stellar models fail:
  - (a) RLOF
  - (b) Common Envelope Evolution (CEE)
  - (c) The helium flash
- (II) Can we develop a testing regime for the natal supernova kick, which relies on more observations than just isolated pulsar velocities?
- (III) What modifications are required to be made to our stellar modelling code in preparation for the next major BPASS release?

### 1.3.2 | Thesis outline

This thesis is, broadly, a tour of stellar evolution from Zero-Age Main Sequence (ZAMS) to the eventual merger of two compact objects, with a few omissions along the way (such as the moment of supernova). The thesis is structured as follows:

1. In chapter 2, we provide the necessary theoretical background for future chapters, critically examine the pieces of software and numerical methods employed in this work, and discuss the logistics of running this single-thread grid-based code as a multiprocessed code on a distributed cluster.
2. In chapter 3, we present the results of our stellar evolution codes in the interesting case when both stars almost (or in some instances, do) touch. (Theme I)
3. In chapter 4, we explore the regime in which mass transfer is assumed to become helium-rich, and the effects on the future evolution of these binaries. (Themes I, III)
4. In chapter 5, we explore the population properties of binaries which underwent helium-rich mass transfer (Themes I, III).

5. In chapter 6, we examine the kinematics of the post-supernova inspiral and the mechanics of the natal supernova kick. (Theme II)
6. In chapter 7, we review the thesis and conclude. (Themes I, II, III)
7. In Appendix A, we provide the KAITIAKI code necessary to generate the models for certain stellar evolution plots throughout the thesis. (Themes I, III)
8. In Appendix B, we list the codes employed and their versions.
9. In Appendix C, we present period-eccentricity distributions for the LVK Collaboration O3 contour (Theme II)
10. In Appendix D, we list the default parameter choices we make for our stellar evolution models (Themes I, III).



CHAPTER 2

---

# Theoretical Background

*Se'i lua'i lou le 'ulu taumamao.**Pick the breadfruits on the far-off branches first*

---

SĀMOAN PROVERB

**I**N THIS CHAPTER, WE PROVIDE A brief account of the evolution of a star from a theoretical perspective, from its formation to post-supernova. We will discuss both the case of a single isolated star, and an isolated binary. It is pertinent to note that our binaries will be assumed to have formed at birth, rather than by, for example, the dynamical capture of a runaway star in a globular cluster, which introduces extra effects and considerations. This process can be modelled by evolving a single star to a certain point, and then enabling binary evolution.

## 2.1 | Stellar Formation

*Nature and nature's laws lay hid in night: God said, "Let Newton be!" and all was light.*

---


ALEXANDER POPE

From the point of view of stellar evolution, which starts in earnest at the ZAMS, the formation of stars is interesting but ultimately broadly unimportant. We provide a short recap of stellar formation here as scene-setting, but our stellar evolution models assume that a gas cloud has already collapsed and we have a star at its ZAMS.


Stars are born in clouds of dust in rich H(II) regions of space. Due to inhomogeneities in the dust formation, some regions become more dense than others. When a region of this gas with adiabatic constant  $\gamma$  and mean molecular weight  $\mu$  reaches the so-called Jeans Mass,

$$M_J = \left( \frac{10k_B T}{3(\gamma - 1)\mu m_H G} \right)^{\frac{3}{2}} \left( \frac{3}{4\pi\rho} \right)^{\frac{1}{2}} \sim \left( \frac{10k_B T}{G} R \right)^{\frac{3}{2}} \left( \frac{1}{M} \right)^{\frac{1}{2}} \quad (2.1)$$

where the similarity is taking  $\gamma = \frac{5}{3}, \mu = \frac{1}{2}$  for an adiabatic gas composed of H(II), then gravity becomes the dominant force in the clumps of gas. This causes a protostar to form. This protostar slowly gains more mass from its surroundings and, as it does so, increases in temperature. When the core temperature reaches  $10^7$  K, the hydrogen in the core begins to fuse into helium via the reaction chain



This is aided by two further hydrogen to helium reactions:



or, via a master equation,



with intermediate products which remain stable and are thus omitted for simplicity. This energy release will sustain the star throughout its lifetime, providing outwards pressure resisting gravity pulling the star inwards, in a balance termed *hydrostatic equilibrium*. Once hydrogen fusion has begun, we say that the star has reached ZAMS, and is now on the *main sequence*, where it will spend approximately 90% of its lifetime.

## 2.2 | Stellar Evolution

*It did not last: the Devil howling “Ho! Let Einstein be!” restored the status quo.*

*Answer to Pope’s Epitaph for Sir Isaac Newton, BY J. C. SQUIRE*

Now that our protostar has evolved onto the main sequence, it has become a star in its own right. Over the next few million years, it will continue to burn hydrogen into helium, forming a Helium core. It is at this point that the evolution encounters a number of bifurcation points, depending on the mass and chemical composition of the star. In general, while the star lives, the four equations of stellar structure hold:

$$\left. \begin{aligned} \frac{dm}{dr} &= 4\pi r^2 \rho && \text{(mass continuity)} \\ \frac{dp}{dr} &= -\frac{GM\rho}{r^2} && \text{(hydrostatic equilibrium)} \\ \frac{dT}{dr} &= -\frac{3\kappa\rho L}{64\pi r^2\sigma T^3} && \text{(energy transport)} \\ \frac{dL}{dr} &= 4\pi r^2 \rho \epsilon && \text{(thermal equilibrium)} \end{aligned} \right\} \quad (2.3)$$

where  $\epsilon$  is the rate of energy eneration per unit mass,  $\kappa$  is the opacity of the material,  $\sigma$  is the Stefann-Boltzmann constant, and all other symbols have their usual meanings.

These equations represent conservation of mass, hydrostatic equilibrium, energy transport, and thermal equilibrium, respectively. Knowing constants  $\epsilon, \kappa$ , we have four equations for four unknowns. It is usually convenient to take the *Rosseland mean* opacity,

$$\frac{1}{\kappa} = \frac{\pi}{acT^3} \int_0^\infty \frac{1}{\kappa_\nu} \frac{\partial B}{\partial T} d\nu \quad (2.4)$$

where  $a$  is the radiation density constant ( $a = 7.57 \times 10^{-15} \text{ erg cm}^{-3} \text{ K}^{-4}$ ) and  $B$  is the black-body intensity at frequency  $\nu$  and temperature  $T$ . The function  $\kappa_\nu$  is obtainable from nuclear physics, and is related to the scattering cross-section  $\sigma_\nu = \int_0^{2\pi} \int_0^\pi \frac{d\sigma}{d\Omega} \sin\theta d\theta d\phi$ . The Rosseland mean becomes

$$\kappa = \frac{T^5 ac^3 k_B}{\pi h^2 \int_0^\infty \frac{\nu^4}{\left(\cosh\left(\frac{h\nu}{Tk_B}\right) - 1\right) \kappa_\nu} d\nu}, \quad (2.5)$$

though the integral cannot be analytically solved further. On the other hand,  $\epsilon$  is obtainable from knowing the nuclear reactions occurring in the star’s core:

$$\epsilon = \sum_i^N \text{energy output of reaction } i \quad (2.6)$$

All told, in the simplest sense, the internal *structure* of the star can be modelled by hand by solving Equation 2.3 from  $m = 0$  to  $m = M$ .

### 2.2.1 | The chemical composition of stars

A star contains at its core a complex nuclear furnace. Since stars are born in regions of space rich in H(II) – molecular hydrogen – and obtain high enough pressures in the core to ignite fusion, the star will be sustained by the burning of predominantly Hydrogen and Helium. We thus group the three main stages into Hydrogen, Helium, and Metals (which we define as anything with atomic number  $\geq 3$ , i.e., Lithium and heavier elements). It is convenient to speak of the mass fraction of the element, i.e., the fraction by mass of a star that is a certain element. For our purposes, this is expressible as

$$X + Y + \sum_{i=0}^{115} Z_i = 1, \quad (2.7)$$

where  $X$  is the mass fraction of Hydrogen,  $Y$  of Helium, and  $Z_i$  as the metal<sup>1</sup> with atomic number  $i+3$ . When speaking about the chemical composition of a star it is also convenient to speak of the *metallicity*, the fraction at ZAMS of the star that is metal. For this, we define the metallicity as

$$Z \equiv \sum_{i=0}^{115} Z_i, \quad (2.8)$$

such that Equation 2.7 becomes

$$X + Y + Z = 1. \quad (2.9)$$

### 2.2.2 | Evolutionary Pathways

In this section, we examine the four major regimes of stellar evolution for single-stars. The precise boundaries are debatable, but are generally accepted to be logarithmically spaced.

#### Low-Mass Stars: $0.08 M_{\odot} < M < 2 M_{\odot}$

Low-mass stars ignite helium degenerately in their cores, leading to thermal runaway. The star can peak at a luminosity close to  $10^{11} L_{\odot}$  (approximately 100 times the peak luminosity of SN 1987A, per Figure 5 of Arnett et al. (1989)). However, this peak in luminosity is attributable to nuclear fusion in the core, and the energy does not reach the surface, let alone escape into the ISM – rather, it is absorbed into the outer layers of the star. During this extremely brief thermal runaway, the luminosity begins to settle and the star is able to continue evolving. Models of low mass stars fail at this stage and care is required to evolve them. This phenomena is called the *Helium flash* and is discussed further in subsubsection 1.

After the ignition of helium, these stars cross the so-called Hertzsprung Gap – the region where very few stars are observed empirically. This region is bounded (on a colour-magnitude diagram, which is a transform of the HR diagram)<sup>2</sup> by the points  $(M_v, B - V) = \{(2, 0.5), (-7, 0.1), (-6, 1.6)\}$  (Hoyle, 1960), where  $M_v$  is the V-band magnitude (magnitude in the visual band), and  $B - V$  is the colour index of the star. The phenomenon of a gap


<sup>1</sup>It is custom in astronomy to refer to anything with an atomic number greater than 2 as a metal, rather than the conventional definition of a material conductive at absolute zero, including most of the d-block, s-block, f-block, and about half of the p-block on the periodic table.

<sup>2</sup>The relationship between a colour-magnitude diagram and the HR diagram is that  $\log L = 0.4(M_{\text{bol},\odot} - M_{\text{bol},*}) \simeq 0.4(4.74 - M_{v,*} - \text{BC})$ ,  $\log T = \log \left( 4600 \left( \frac{1}{0.92(B-V)+1.7} + \frac{1}{0.92(B-V)+0.62} \right) \right)$ , where BC is a bolometric correction which depends on the stellar type. This expression for the temperature is a fit provided by Ballesteros (2012).

existing is attributable to the fact that stars cross this gap exceedingly quickly: whereas it spends  $\sim 90\%$  of its life on the main sequence, it spends  $\sim 2\%$  of its life crossing the Hertzsprung gap (Hoyle, 1960).

### Intermediate-Mass Stars: $2 M_{\odot} < M < 8 M_{\odot}$


Intermediate mass stars exist above the threshold for the Helium flash to occur. Helium ignition is not degenerate and the development of a helium core is not impeded. As the star ascends the Asymptotic Giant Branch (AGB), the core becomes helium-exhausted and it becomes a red giant, with a radius many hundreds or even thousands of times greater than the sun. Its mass, sustained prior to the AGB via nuclear fusion, begins to decrease. An example of intermediate-mass evolution is shown in Figure 2.1.



**Figure 2.1:** The evolution of an  $4 M_{\odot}$  star from ZAMS until CO core development. The final core abundances are  $Z_C \approx 0.44$  and  $Z_O \approx 0.53$ .

### Massive Stars: $8 M_{\odot} < M < 80 M_{\odot}$

Massive stars are the key focal objects of this study, and in many ways offer some of the most exciting phenomena we can probe. In Figure 2.2 we show the variety of evolution that this regime has, alongside the ZAMS track from  $8 M_{\odot}$  to  $90 M_{\odot}$ . The key features are as follows. When a massive star exhausts its supply of hydrogen in its core and evolves off the main sequence to begin burning helium, it generally bypasses the Hertzsprung gap and evolves up the AGB. From here it generally engages in a “blue loop”, where it remains at a roughly constant radius whilst increasing in temperature, before cooling at a slightly higher luminosity. The question of the physical process powering a blue loop is unanswered (Zhao et al., 2023), though this thesis is unconcerned with their generation. We note that our models include blue loops, as shown in Figure 2.2 and Figure 2.3.



**Figure 2.2:** The evolution of stars in the range  $10 - 80 M_{\odot}$ . The dashed gray line represents the Zero-Age Main Sequence (ZAMS), which is the origin point of all of our models, and (in this plot) extends from  $1 M_{\odot}$  to  $300 M_{\odot}$ . The blue triangle is the (highly approximate) bounds of the Hertzsprung gap.


### 2.2.3 | Relativistic corrections

*There’s a trick you can use in mathematics called “not worrying about it”*

MATT PARKER

This analysis has presumed Newtonian mechanics. However, the mass of stars puts them formally into the general relativistic regime, and thus strictly speaking we should treat this as a problem in full GR. In this approach, we would solve the Einstein Field Equations for the interior of the star:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu} \quad (2.10)$$



**Figure 2.3:** The evolution of an  $8 M_{\odot}$  star. Highlighted is the star's Zero-Age Main Sequence (ZAMS), Terminal-Age Main Sequence (TAMS), the blue loop, and the Asymptotic Giant Branch (AGB). The Hertzsprung gap is also highlighted. In this particular model, numerical instabilities in the leftmost (hottest) region of the plot have been highlighted in pink. This is normally removed in postprocessing though has been left in as an illustration.

with cosmological constant  $\Lambda$ , stress-energy  $T_{\mu\nu}$ , and the Einstein tensor  $G_{\mu\nu}$  given by

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R \quad (2.11)$$

$$= R^\lambda{}_{\mu\lambda\nu} - \frac{1}{2}g_{\mu\nu}R^\sigma{}_\sigma \quad (2.12)$$

$$= R^\lambda{}_{\mu\lambda\nu} - \frac{1}{2}g_{\mu\nu}g^{\sigma\gamma}R_{\sigma\gamma} \quad (2.13)$$

$$= \sum_\lambda R^\lambda{}_{\mu\lambda\nu} - \frac{1}{2}g_{\mu\nu} \sum_\sigma \sum_\gamma \sum_\alpha g^{\sigma\gamma} R^\alpha{}_{\sigma\alpha\gamma} \quad (2.14)$$

$$= \sum_\lambda \left( \Gamma^\lambda{}_{\mu\nu,\lambda} - \Gamma^\lambda{}_{\mu\lambda,\nu} + \sum_\xi \Gamma^\lambda{}_{\xi\lambda} \Gamma^\xi{}_{\mu\nu} - \sum_\xi \Gamma^\lambda{}_{\xi\nu} \Gamma^\xi{}_{\mu\lambda} \right) - \frac{1}{2}g_{\mu\nu} \sum_\sigma \sum_\gamma \sum_\alpha g^{\sigma\gamma} \left( \Gamma^\alpha{}_{\sigma\gamma,\alpha} - \Gamma^\alpha{}_{\sigma\alpha,\gamma} + \sum_\xi \Gamma^\alpha{}_{\xi\alpha} \Gamma^\xi{}_{\sigma\gamma} - \sum_\xi \Gamma^\alpha{}_{\xi\gamma} \Gamma^\xi{}_{\sigma\alpha} \right), \quad (2.15)$$

where there is still tensorial shorthand:

$$\Gamma^\gamma{}_{\beta\mu} \equiv \frac{1}{2} \sum_\alpha g^{\alpha\gamma} (g_{\alpha\beta,\mu} + g_{\alpha\mu,\beta} - g_{\beta\mu,\alpha}) \quad (2.16)$$

and the comma-derivative notation is

$$\phi_{,\mu} \equiv \frac{\partial \phi}{\partial x^\mu}, \quad (2.17)$$

such that, for example,

$$g_{\alpha\beta,\mu} = \frac{\partial g_{\alpha\beta}}{\partial x^\mu}.$$

In Equation 2.14, we explicitly reinsert the summation symbols to illustrate the number of simultaneous summations that are taking place implicitly. As one can see, this reasonably simple expression balloons quickly into something intractable by hand and time-consuming to evaluate computationally. Although the metric tensor is generally four-dimensional,<sup>3</sup> such as the Friedmann-Lemaître-Robertson-Walker metric (Friedmann, 1922; Lemaître, 1931; Robertson, 1935; Walker, 1937) describing the expansion of the Universe,

$$g_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & a(t)^2 & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix},$$

it often lacks a closed form, especially on large-scale chaotic behaviour that cannot be locally “washed out” (such as how on Universal scales, individual stars can be ignored to describe the homogeneity and isotropy of the Universe), and is obtained via numerical methods.

This is also not considering complications of writing the stress-energy tensor  $T_{\mu\nu}$ , which may have its own complex form (again, assuming a closed-form solution *exists*) determined via fluid-mechanical processes. All told, there are 64 equations to be solved instead of 4. In addition, the 64 solutions are “worse” to solve: harder to

<sup>3</sup>And thus the summation indices run from 1 to 4

integrate by hand, and requiring greater time complexity. Numerical relativity codes typically simulate one star for the order of kiloseconds of its lifetime (see, for an example, ud-Doula et al., 2013), and even the most efficient codes take longer than our codes to run. For instance, the PHANTOM 3D MHD code (Price et al., 2018) can take several days to run high-resolution models. Consequently, for the number of stars we wish to simulate, we choose to adopt a Newtonian framework.

### 2.2.4 | Stellar Atmospheres

As a final note in the evolution of a single star, we must briefly address what occurs about the photosphere of the star. A star is not a planet – there is no firm boundary demarcating the inner of a star from the outer. The common definition of the “surface” of a star is the surface of last scattering (photosphere), which is the point radially in which the mean free path  $\ell$  of a photon emitted from the core of the star becomes meaningfully long (internally,  $\ell_\gamma \sim 2$  cm, externally,  $\ell_\gamma \sim 4.4$  pc).<sup>4</sup> However, outflows from the star as well as the potential interaction of a partner in a common-envelope event mean that this surface is not necessarily firm: there is no sharp turn from stellar material to empty space. Thus, the simulation of stellar atmospheres is of interest to the stellar astrophysicist. However, this introduces another rather large layer of complexity. The synthesis of a stellar spectra, for example, is not a trivial task, as well as simulating the matter content. This thesis is concerned with detailed modelling of the stellar *interior*, and thus the atmosphere is an interesting sidenote, but one we shall not pursue.

## 2.3 | Supernovae

*And on the pedestal these words appear:  
“My name is Ozymandias, king of kings:  
Look on my works, ye Mighty, and despair!”  
Nothing beside remains.*

---

PERCY BYSSHE SHELLEY, *Ozymandias*

*Pallida mors æquo pulsat pede pauperum  
tabernas regumque turres.  
Pale death knocks with the same tempo upon the  
huts of the poor and the towers of kings.*

---

HORACE


A star’s lifetime is, unfortunately, not infinite. The key assumption underpinning stellar evolution is that of hydrostatic equilibrium: the counterplay of gravity against outwards pressure. Whilst this equilibrium holds, the star is stable and resists explosion. It is when this equilibrium is broken – with gravity the dominant force in this scenario – that the star implodes in a core-collapse supernovae (CCSN). From a thermodynamic perspective, as the star evolves, it shrinks as energy is bled off into the interstellar medium. This shrinkage compresses and heats the core, meaning the star’s specific heat capacity is negative. This system is thermodynamically unstable, and thus every star either collapses into a remnant (cf. black holes, neutron stars) or is sustained in some form by quantum mechanical phenomena (cf. White Dwarfs, Thorne-Żytkow objects, and other exotica).

In any case, the breaking of hydrostatic equilibrium results generally results in<sup>5</sup> a supernova, which can be classified into subtypes, as shown in Figure 2.4. In this event, gravity pulls the stellar core inwards, before “bouncing” and ejecting the envelope into the interstellar medium. The physics of this approximately second-long event are highly uncertain, and are in general unable to be modelled by detailed stellar evolution codes. All models we generate – assuming they have not been terminated earlier – will terminate at the moment of supernova.

---


<sup>4</sup>The mean free path is a function of the particle under consideration: for a neutrino, internally  $\ell_\nu \sim 3.68$  pc and externally  $\ell_\nu \sim 81$  Rpc  $\sim 3 \times 10^{18}$  times the diameter of the observable Universe.

<sup>5</sup>The extent of the veracity of this statement is an area of active research: Reynolds et al. (2015) presents a short recap.



**Figure 2.4:** A rough outline of supernova classification. One can see that the presence of hydrogen lines distinguishes a Type I from Type II supernova. Type II supernova subtypes are determined by the behaviour of their light curves – how the apparent magnitude changes against time – and thus are determined by the luminosity of the supernova, something detailed models are not able to model. On the other hand, the presence of hydrogen/silicon/helium in the supernova is something that detailed codes *are* able to model.

Importantly for the nucleosynthesis of heavier-than-Iron elements (which cannot be formed via fusion,<sup>6</sup> as shown in Figure 2.5), the supernova shockwave propagating through the core causes a new phase of nucleosynthesis, albeit briefly.



**Figure 2.5:** The binding energy per nucleon for elements up to Uranium-238. The most stable element of fusion here is Iron-56. Elements lighter than Iron-56 will fuse towards it, and elements heavier will undergo fission to reach it. The Bethe-Weizsäcker formula is the semi-empirical mass formula devised by Bethe and Weizsäcker to describe the liquid drop model of Gamow (1930). The marked atoms on the plot represent the six elements we generally consider in our stellar modelling code –  $^1\text{H}$ ,  $^4\text{He}$ ,  $^{12}\text{C}$ ,  $^{14}\text{N}$ ,  $^{16}\text{O}$ ,  $^{20}\text{Ne}$  – with the balance assumed to be insufficiently energetic to alter the structure or evolution of a star (Stancliffe et al., 2004). Graph adapted from Fastfission on Wikimedia Commons (who released it into the public domain).

## 2.4 | Two to tango: Binary Evolution

Introducing a second star into the mix complicates the evolution. Whereas before a star losing mass (through, for example, a coronal mass ejection (CME)) was assumed to eject it into the ISM, it may now accrete that material directly onto its companion. In addition, either star may in principle increase in radius enough to overlap with its companion, or even touch surfaces – introducing new dynamics into the mix.

### 2.4.1 | Binary star kinematics

There are several formation channels for a binary star system. Binaries may be formed from birth (as discussed in section 2.1, with the caveat that in this scenario two clumps of gas form near each other), via fragmentation of a protostar’s disc, dynamical capture of one star by another (comparable to one of the suggested origins of

<sup>6</sup>Strictly speaking, fusion can occur for elements heavier than Iron-56, but the reaction is endothermic; it requires more energy than is released, meaning it is unsustainable.

Phobos and Deimos, per Burns (1978)), or via a higher multiplicity system ejecting one or more of its constituent stars, perhaps via a natal supernova kick.


In any case, once formed and the orbit is stabilised, binary systems obey Kepler’s three laws. In the generalised binary regime, however, the assumption that  $M_1 \gg M_2$  (which underpins most statements of Kepler’s laws) is not necessarily true – for instance, the system LSS 3074 has two stellar components with ZAMS masses of approximately  $30 M_\odot$  and  $10 M_\odot$  (Raucq et al., 2017). Moreover, the fact that there are two stars in orbit rather than one object orbiting another means that Kepler’s laws must be stated more precisely. Formally, they can be expressed as (see the appendix of Richards (2021) for derivations)

1. All orbits are ellipses, with both stars orbiting a common barycentre at the focus of the ellipse,
2. A line segment joining the two stars sweeps out equal areas in equal times,
3. The square of the period of the system is proportional to the cube of the semimajor axis of the ellipse (the binary separation), with the proportionality term dependent on the sum of the masses.

The third law can be stated more precisely as,

$$T^2 = \frac{4\pi^2}{G(M_1 + M_2)} a^3, \quad (2.18)$$

and a schematic of the orbit of a binary system is given in Figure 2.6.



**Figure 2.6:** A schematic of a simple binary star system. The centre of mass (barycentre) is shown as a blue dot. Here,  $M_1$  is the heavier star, which orbits at a smaller distance  $x_1$  to the barycentre by Kepler’s first and Newton’s second laws. Figure adapted from Richards (2021).

## 2.4.2 | Roche Lobes


*Algebra’s like sheet music. The important thing isn’t can you read music, it’s can you hear it. Can you hear the music, Robert?*

*Niels Bohr, IN OPPENHEIMER*

A binary system is complicated by the presence of its companions in several different ways. As alluded to in the previous section, the presence of a companion warps space-time in such a way that material can be quite distorted from a sphere or oblate spheroid but still remain gravitationally bound to the system. In the extremal case, a star can grow to fill or even exceed its Roche Lobe, which is the tear-drop equipotential shaped surface within which material remains gravitationally bound to its star. In Figure 2.7 we display these equipotential surfaces for a range of different mass ratios. It is pertinent to note that the shape of all of the Roche Lobes change rapidly at low mass ratios, however at higher mass ratios the shapes change less rapidly. In an equal-mass ratio binary, the binary configuration has rotational symmetry about the  $\mathcal{L}_1$  point, and reflectional symmetry about the  $x$ -axis and the line  $y = \mathcal{L}_1$ .

There are two primary Roche Lobe points we consider: the  $\mathcal{L}_1$  point and the  $\mathcal{L}_2$  point. The  $\mathcal{L}_1$  point (also known as the inner Lagrangian point) is the nexus between the two stars, the  $\mathcal{L}_2$  point (the outer Lagrangian point) is located on the other side of the secondary to the inner point.

We follow the derivation given in Rose (1998) to find an expression for the Roche potential. First we note that in a rotating reference frame in Cartesian coordinates, we have



**Figure 2.7:** The dimensionless Roche potential as a function of mass ratio  $q$ . Note that the shape of the Roche potential changes markedly at low mass ratio, and changes more slowly as the mass ratio approaches unity. Distances have been normalised such that the primary star is at  $(x, y, z) = (0, 0, 0)$  and the secondary star is at  $(x, y, z) = (1, 0, 0)$ , i.e., we measure distances in units of separation and that the coordinate system is centered on the primary star.

$$\overbrace{\frac{GM_1M_2}{a^2}\hat{\mathbf{x}}}^{\text{gravitational force}} = \underbrace{-M_1\vec{\omega} \times (\vec{\omega} \times \vec{r}_1)}_{\text{centrifugal force}}, \quad (2.19)$$

where  $\hat{\mathbf{x}}$  is the line between the center of mass of both stars,  $a$  the separation,  $\vec{\omega}$  the angular velocity, and  $\vec{r}_1$  the position vector of the primary star. The x-ordinates of the centres of mass are given by

$$\left. \begin{aligned} x_1 &= \frac{M_2}{M_1+M_2}a \\ x_2 &= \frac{-M_1}{M_1+M_2}a \end{aligned} \right\} \quad (2.20)$$

Since we are considering a rotating coordinate system we have that the centrifugal force is given by

$$-M_1\vec{\omega} \times (\vec{\omega} \times \vec{r}_1) = M_1\omega^2 x_1 \quad (2.21)$$

and hence

$$\begin{aligned} \frac{GM_1M_2}{a^2}\hat{\mathbf{x}} &= M_1\omega^2 x_1 \\ \omega^2 &= \frac{GM_1M_2}{M_1x_1a^2}\hat{\mathbf{x}} \\ &= \frac{GM_2}{x_1a^2}\hat{\mathbf{x}} \\ &= \frac{GM_2(M_1+M_2)}{M_2 \cdot a \cdot a^2}\hat{\mathbf{x}} \\ &= \frac{G(M_1+M_2)}{a^3}\hat{\mathbf{x}}. \end{aligned} \quad (2.22)$$

The centrifugal potential of a test mass  $m$  at  $(x, y)$  is hence

$$V = -\frac{1}{2}m\omega^2 (x^2 + y^2) = -\frac{Gm(M_1+M_2)(x^2 + y^2)}{2a^3}, \quad (2.23)$$

and thus its energy is

$$E = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) + V(x, y), \quad (2.24)$$

where

$$V(x, y) = -\overbrace{\frac{GmM_1}{\sqrt{(x-x_1)^2 + y^2}}}^{\text{G.P.E. due to primary}} - \overbrace{\frac{GmM_2}{\sqrt{(x-x_2)^2 + y^2}}}^{\text{G.P.E. due to secondary}} - \overbrace{\frac{Gm(M_1+M_2)(x^2 + y^2)}{2a^3}}^{\text{centrifugal potential}}. \quad (2.25)$$

We introduce the dimensionless coordinates measured in units of the separation of the binary (or reduced masses)

$$\xi = \frac{x}{a}, \quad \eta = \frac{y}{a}, \quad \xi_1 = \frac{M_2}{M_1+M_2}, \quad \xi_2 = \frac{M_1}{M_1+M_2} \quad (2.26)$$

Our potential now becomes

$$V(\xi, \eta) = -\frac{Gm(M_1 + M_2)}{a} \times \left[ \frac{\xi_2}{\sqrt{(\xi - \xi_1)^2 + \eta^2}} + \frac{\xi_1}{\sqrt{(\xi - \xi_2)^2 + \eta^2}} + \frac{1}{2}(\xi^2 + \eta^2) \right]. \quad (2.27)$$

Taking the partial derivative along the  $\xi$ -axis (the rescaled  $x$ -axis), we get

$$\frac{\partial V}{\partial \xi} = -\frac{Gm(M_1 + M_2)}{a} \times \left[ -\frac{\xi_2(\xi - \xi_1)}{\left((\xi - \xi_1)^2 + \eta^2\right)^{\frac{3}{2}}} - \frac{\xi_1(\xi - \xi_2)}{\left((\xi - \xi_2)^2 + \eta^2\right)^{\frac{3}{2}}} + \xi \right] \quad (2.28)$$

Hence the extrema of the potential  $V$  are at the points  $\xi$  where (taking  $\eta = 0$  on the  $\xi$ -axis)

$$0 = \xi - \frac{\xi_2}{(\xi - \xi_1)^2} - \frac{\xi_1}{(\xi - \xi_2)^2}. \quad (2.29)$$


No analytical function for  $\xi$  exists unless either or both of  $\xi_1, \xi_2$  are zero (which would only be the case for a single star, and thus the Roche lobe would not exist in the first instance), thus we must make approximations.

### The $\mathcal{L}_1$ point


The inner Lagrangian point has the approximation from Eggleton (1983), which is

$$\frac{R_L}{a} \approx \frac{0.49q^{\frac{2}{3}}}{0.6q^{\frac{2}{3}} + \ln\left(1 + q^{\frac{1}{3}}\right)}. \quad (2.30)$$

We show the circle with this radius  $R_L$  in Figure 2.8 plotted against the exact solution for  $\eta = \mathcal{L}_1$ . Eggleton's formula is a volume-equivalent one; although his radius  $R_L < \xi$ , the volume of an Eggleton-Roche sphere  $\frac{4}{3}\pi R_L^3$  is approximately equal (to within 1 per cent) of the numerically integrated volume, as is shown in Figure 2.9.



**Figure 2.8:** A schematic of the Roche lobe formula Equation 2.30. Here, the exact solution to the Roche lobe formula  $\xi$  in Equation 2.29 (orange) vs the fit provided by Eggleton (1983) (blue). Although precise details of the Roche lobe, such as the apex of the teardrop at the point  $\xi$ , are lost, the volume is preserved to within 1 per cent accuracy over all mass ratios.




**Figure 2.9:** The volume of the Roche Lobe computed numerically through Monte-Carlo integration (orange) vs the volume assuming Equation 2.30 (blue). The Eggleton approximation fits well to the volume, with around 1 per cent error across the entire mass ratio range (truncated here to  $q \leq 1$ ).

### The $\mathcal{L}_2$ point

The outer Lagrangian point,  $\mathcal{L}_2$ , is given by a different zero of the potential function. The approximation we use is that of Marchant et al. (2016), where

$$R_{L2} = R_L(q) \left( 1 + \frac{0.299 \arctan(1.84q^{-0.397})}{q^{0.520}} \right), \quad (2.31)$$

and  $R_L(q)$  is given by Equation 2.30.



**Figure 2.10:** As for Figure 2.9, but for the  $\mathcal{L}_2$  point with formula given by Equation 2.31 (Marchant et al., 2016).

We note that Marchant et al. (2016) uses an inverse mass ratio definition compared to Eggleton (1983), that is, Marchant et al. (2016) takes  $q_{\text{Marchant}} = \frac{M_2}{M_1}$  whereas Eggleton (1983) takes  $q_{\text{Eggleton}} = \frac{M_2}{M_1} = \frac{1}{q_{\text{Marchant}}}$ . This

is accounted for in Equation 2.31 such that the mass ratios are consistent (and we take  $q = q_{\text{Eggleton}}$ ), but other texts do not necessarily differentiate or describe the difference.

### 2.4.3 | Mass Transfer

When two stars are sufficiently close in proximity, events such as CMEs can cause two stars to exchange mass (whereas in the single-star case, this would be matter lost to the ISM). The nature of mass transfer is a complicated affair; most binary star population synthesis codes typically inherit their mass loss prescriptions from papers such as Bondi & Hoyle (1944); Hurley et al. (2002), and this approach is what we detail in this section.

Mass transfer can broadly be sectioned into four main categories:

1.  $\dot{M}_{\text{wind}}$ , where matter is transferred through stellar winds,
2.  $\dot{M}_{\text{RLOF}}$ , where matter is transferred through one (or more) of a binary's Lagrange points (cf. subsection 2.4.2),
3.  $\dot{M}_{\text{acc}}$ , where matter is transferred through accretion (or interaction with a disk), and
4.  $\dot{M}_{\text{Bondi}}$ , where matter proceeds through a Bondi & Hoyle (1944) type mechanism.

Our mass loss rate for each star is constructed from these four terms as

$$\dot{M} = \dot{M}_{\text{wind}} - \dot{M}_{\text{Bondi}} + \dot{M}_{\text{RLOF}} - \dot{M}_{\text{acc}}. \quad (2.32)$$

#### Wind mass-loss

Wind mass loss computations are (generally) performed using the prescription of de Jager et al. (1988) and Vink et al. (2001). The precise scheme is detailed in Eldridge et al. (2017).<sup>7</sup> There are two critical modifications we make to their scheme.

The first modification surrounds the transition into the Wolf Rayet (WR) regime. Traditionally this has been a sharp transition when the surface hydrogen mass fraction becomes less than 0.4 and the surface temperature is above  $10^4$  K. This is a reasonable approximation, though if a star reaches this phase and is *not* yet a WR star, the code will fail. To account for this we gradually turn on these enhanced WR rates when this condition approaches, smoothing the transition.

The second modification we have made is that if a star has a hydrogen exhausted core and begins approaching the WR temperature range, we assume that WR mass loss must begin and thus turn on the slowly-turning on WR regime.

Mathematically, the computation for  $\dot{M}_{\text{wind}}$  is more involved, though follows the same essential structure as Vink et al. (2001). We construct it as follows. The total pre-WR mass loss is given by de Jager et al. (1988), namely

$$\log \dot{M}_{\text{de Jager}} = - \sum_{n=0}^5 \sum_{\substack{i=0 \\ j=n-i}}^n a_{ij} \cos(i \arccos x_1) \cos(j \arccos x_2), \quad (2.33)$$

$$\dot{M}_{\text{pre-WR}} = \sqrt{\frac{Z}{Z_{\odot}}} 10^{\log \dot{M}_{\text{de Jager}}}, \quad (2.34)$$

<sup>7</sup>In other codes such as MESA, this is referred to as the ‘‘Dutch’’ mass loss prescription owing to the nationality of the authors of de Jager et al. (1988) and Vink et al. (2001).

where  $x_1 = \frac{\log T - 4.05}{0.75}$ ,  $x_2 = \frac{\log L - 4.6}{2.1}$ , and the coefficients  $a_{ij}$  are given in Table VI of de Jager et al. (1988). We interpolate the bi-stability jump around 25 kK (see, for example, Pauldrach & Puls, 1990; Vink et al., 1999) with the following regime. Define the constants

$$R_1 = \left( \frac{Z}{Z_\odot} \right)^{0.13}, \quad (2.35)$$

$$R_2 = -c_1 + c_2 \log \left( \frac{L}{c_{16}} \right) - c_3 \log \left( \frac{M}{c_{15}} \right) - c_4 \log (1.3R_1) + c_5 \log \left( \frac{T}{c_{14}} \right) - c_6 \left( \log \left[ \frac{T}{c_{14}} \right] \right)^2 + c_7 \log \left( \frac{Z}{Z_\odot} \right), \quad (2.36)$$

and

$$R_3 = -c_8 + c_9 \log \left( \frac{L}{c_{16}} \right) - c_{10} \log \left( \frac{M}{c_{15}} \right) - c_{11} \log (0.65R_1) + c_{12} \log \left( \frac{T}{2 \times 10^4} \right) + c_{13} \log \left( \frac{Z}{Z_\odot} \right), \quad (2.37)$$

where the constants  $c_i$  are given in Table 2.1.

| Constant | Value | Constant | Value             |
|----------|-------|----------|-------------------|
| $c_1$    | 6.697 | $c_9$    | 2.210             |
| $c_2$    | 2.194 | $c_{10}$ | 1.339             |
| $c_3$    | 1.313 | $c_{11}$ | 1.601             |
| $c_4$    | 1.226 | $c_{12}$ | 1.07              |
| $c_5$    | 0.933 | $c_{13}$ | 0.85              |
| $c_6$    | 10.92 | $c_{14}$ | $4 \times 10^4$ K |
| $c_7$    | 0.85  | $c_{15}$ | 30 $M_\odot$      |
| $c_8$    | 6.688 | $c_{16}$ | $10^5$ $L_\odot$  |

**Table 2.1:** The mass loss rate constants for interpolation around the bistability jump (Equation 2.35, Equation 2.36, Equation 2.37) (shown pictorially in Figure 2.11).

Our mass loss rate becomes:

$$\dot{M}_{\text{BS}} = \begin{cases} \frac{(T-T_1)10^{R_3} + (T_2-T)\dot{M}_{\text{pre-WR}}}{2.5 \times 10^3} & T_1 < T < T_2 \\ 10^{R_2} & T_2 \leq T \leq T_3 \\ \frac{(T-T_3)10^{R_2} + (T_4-T)10^{R_3}}{5 \times 10^3} & T_3 < T < T_4 \\ \frac{(6T_1-T)10^{R_2} + (T-5T_1)\dot{M}_{\text{pre-WR}}}{10^4} & 5T_1 < T < 6T_1 \\ \dot{M}_{\text{pre-WR}} & \text{otherwise} \end{cases} \quad (2.38)$$

where  $T_1 = 10$  kK,  $T_2 = 12.5$  kK,  $T_3 = 22.5$  kK,  $T_4 = 27.5$  kK. This interpolation is shown in Figure 2.11. We define

$$\mathcal{M} = \sqrt{\frac{Z}{Z_\odot}} \times 10^{-13.6 + 1.63 \log \left( \frac{L}{L_\odot} \right) + 2.22 \log X}, \quad (2.39)$$

where  $X$  is the surface hydrogen abundance of the primary, and then define

$$\dot{M}_{\text{WNL}} = \begin{cases} 5 \left( \mathcal{M} - \dot{M}_{\text{BS}} \right) (0.42 - X) + \dot{M}_{\text{BS}} & 0.42 > X > 0.4 \\ \mathcal{M} & \text{otherwise} \end{cases}. \quad (2.40)$$

Our total mass loss (provided the star has not depleted its surface helium, which will be discussed below) is now

$$\dot{M}_{\text{A}} = \begin{cases} 10\dot{\mu} (\log T - 3.9) + \dot{M}_{\text{BS}} & X < 0.42 \text{ and } \log T > 3.9 \\ \dot{M}_{\text{WNL}} & X > 0.42 \text{ and } \log T > 4.0, \\ \dot{M}_{\text{BS}} & \text{otherwise} \end{cases}, \quad (2.41)$$

where  $\dot{\mu} = \dot{M}_{\text{WNL}} - \dot{M}_{\text{BS}}$ . Finally, if the surface helium depletes we modify the mass loss rate further, defining

$$\dot{M}_{\text{WC}} = \sqrt{\frac{Z}{Z_{\odot}}} \times 10^{-8.3+0.84 \log\left(\frac{Z}{Z_{\odot}}\right)+2.04 \log X_{\text{He}}+1.04 \log(1-X_{\text{He}})} \quad (2.42)$$

such that our total mass loss rate becomes (provided  $X_{\text{H}} < 10^{-3}$  and  $\log T > 4$ )

$$\dot{M}_{\text{B}} = \begin{cases} 100 \left( \dot{M}_{\text{WC}} - \dot{M}_{\text{WNL}} \right) (\mathcal{X} - 0.02) + \dot{M}_{\text{WNL}} & \mathcal{X} > 0.02 \\ \dot{M}_{\text{WNL}} & \mathcal{X} < 0.02 \end{cases}, \quad (2.43)$$

where  $\mathcal{X} = \frac{\frac{1}{3}X_{\text{C}} + \frac{1}{4}X_{\text{O}}}{X_{\text{He}}}$ . The total Vink et al. (2001) mass-loss rate is then

$$\dot{M}_{\text{wind}} = \begin{cases} \dot{M}_{\text{B}} & \text{surface helium depleted} \\ \dot{M}_{\text{A}} & \text{otherwise} \end{cases}. \quad (2.44)$$


### Roche Lobe overflow (RLOF)

Matter extruded from the Lagrangian points is assumed to transfer with a mass loss rate similar to Hurley et al. (2002). We take

$$\dot{M}_{\text{RLOF}} = \min \left( k \mathcal{R} \left( \frac{R}{R_{\text{RL}}} \right)^3, 10^{-2} \right) M_{\odot} \text{ yr}^{-1} \quad (2.45)$$

Here,  $k$  (referred to in our code as RMT) parameterises effectively how much of the star we permit to exist outside of its Roche lobe before it is accreted – i.e., the voracity of RLOF. The choice of this can be somewhat arbitrary. Hurley et al. (2002) for example took  $k = 3 \times 10^{-6} \min \{M_1, 5\}^2$ , where the coefficient was “chosen by experiment to ensure that the mass transfer is steady”, and warned that for degenerate stars a more appropriate choice would be to increase this by a factor of  $10^3$ . Instead, we choose to adopt  $k = 10^3$  outright with no dependence on the mass, finding instead that this replicated other work – allowing efficient RLOF – best.

Matter exceeding  $\mathcal{L}_2$  (the outer Lagrange point) does not currently receive special treatment. The implications of  $\mathcal{L}_2$  overflow are significant. This ejection will lead to unstable mass transfer, shrinking the orbit and likely inducing a CEE event (Fragos et al., 2023). There is debate about the appropriate treatment here. Linial & Sari (2017) indexes the  $\mathcal{L}_2$  rate to the  $\mathcal{L}_1$  rate via an efficiency parameter. Li & Li (2013) considers the case analogous to  $\mathcal{L}_1$  outflow, where material outflowing from  $\mathcal{L}_2$  acts to retard the angular momentum. In their notation  $\frac{dJ}{dt} = -\epsilon \dot{M}_1 a_{L2}^2 \omega$  (compare  $\frac{dJ}{dt} = -\beta \dot{M}_1 a_{L1}^2 \omega$  for RLOF through  $\mathcal{L}_1$ ) for controllable parameters  $\beta$  and  $\epsilon$ . The ambiguity of treatment leads to some codes to terminate at the onset of  $\mathcal{L}_2$  outflow, such as POSYDON (Fragos et al., 2023).  $\mathcal{L}_2$  outflow requires stars to be in reasonably tight orbits ( $\sim 2$  d), which we will approach,



**Figure 2.11:** An illustration of the bi-stability jump at  $\log T \approx 4.4$ . This illustration was computed for a  $21 M_{\odot}$  star. Shown in blue is the cool side of the bistability jump, where  $R_3$  (Equation 2.37) is the interpolation, and shown in red is the hot side, where  $R_2$  (Equation 2.36) is the interpolation.

but none of our stars will enter this regime, and thus we do not afford it special treatment.

### Accretion

Our accretion rates are taken from Bondi & Hoyle (1944); Hurley et al. (2002), and limits the accretion rate by the thermal timescale of the star. The prescription is that

$$\dot{M}_{\text{acc}} = \min \left( \dot{M}_{\text{RLOF}}, \frac{M}{T_{\text{KH}}} \mathcal{R} \left( 1 - \frac{\omega}{0.8\omega_{\text{crit}}} \right) \right) \quad (2.46)$$

where  $\omega$  and  $\omega_{\text{crit}}$  are the spin rate and critical spin rate of the star. We generally assume that the star is composed of individually rotating shells of variable thickness. If  $N$  is the number of meshpoints that the star is composed of (for our cases, normally  $N = 199$ ), then the moment of inertia is given by

$$I = \frac{2}{5} M_c R_c^2 + \sum_{k=1}^N \frac{2}{3} (m_k - m_{k+1}) r_k^2, \quad (2.47)$$

where  $m_k$  is the mass internal to meshpoint  $k$  (sim. for  $r_k$ ), and a subscript of  $c$  refers to the central meshpoint. It is assumed that the central meshpoint sits atop a solid sphere, hence  $\frac{2}{5}$  as its prefactor.

### Bondi-Hoyle accretion

Bondi-Hoyle accretion for a circular orbit is given by

$$\dot{M}_{\text{Bondi}} = \min \left[ - \left( \frac{GM_2}{v_W^2} \right)^2 \frac{\alpha_W}{2a^2} \frac{1}{(1+v^2)^{3/2}}, 0.8 \right] \dot{M}_{\text{wind}}, \quad (2.48)$$

where we have assumed that the orbital velocity is given by

$$v^2 = \frac{G(M_1 + M_2)}{av_W^2}, \quad (2.49)$$

and that the wind velocity is given by

$$v_W = 2\beta_W \frac{GM_1}{R_1}. \quad (2.50)$$

Generalising this to an eccentric orbit causes the first argument in Equation 2.48 to pick up a coefficient of  $\frac{1}{\sqrt{1-e^2}}$ . There are two free parameters to select:  $\alpha_W$  and  $\beta_W$ . These ostensibly depend on stellar types, though we select for simplicity  $\alpha_W = \frac{3}{2}$  (Boffin & Jorissen, 1988) and  $\beta_W = 7$  (Lamers et al., 1995) for O-type stars. This parameterisation is likely incomplete; Hurley et al. (2002) argues that as per Lamers et al. (1995),  $\beta_W$  should vary between 7 for O-type stars to 0.5 for A and F-type stars, perhaps descending as low as  $\beta_W \approx \frac{1}{8}$  for cool supergiants. We leave this implementation for future work under the assumption that our mass-loss rate is correct to within an order of magnitude.

## 2.5 | Post-Supernova

*Wit nodded.*

*“Oh, is this the place where I talk?” Kaladin said.*

*“If you wish.”*

*“I don’t wish. Get on with the story.”*

*Rhythm of War, BY BRANDON SANDERSON*

After a supernova, the star leaves behind its core as a compact remnant. Generally, these are either white dwarves, black holes, or neutron stars. The interior structure of each is either complicated or, in the case of black holes, impossible to observationally probe without violating causality. Their orbital motions are now deterministic from four initial parameters:  $m_1, m_2, a$ , and  $e$ . Motion is now wholly determined by the emission of gravitational radiation through (detectable) gravitational waves. The pertinent equations of motion (all derived more formally in Richards 2021) are given by Peters (1964), namely,

$$\frac{da}{dt} = \frac{-\beta}{a^3 (1 - e^2)^{\frac{7}{2}}} \left( 1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right) \quad (2.51)$$

$$\frac{de}{dt} = -\frac{19}{12} \frac{\beta}{a^4 (1 - e^2)^{\frac{5}{2}}} \left( e + \frac{121}{304} e^3 \right) \quad (2.52)$$

with the constant  $\beta$  defined as

$$\beta \equiv \frac{64}{5} \frac{G^3 m M (m + M)}{c^5}. \quad (2.53)$$

Now if one defines the constant  $c_0$  through

$$c_0 = a_0 (1 - e_0^2) e_0^{-\frac{12}{19}} \left( 1 + \frac{121}{304} e_0^2 \right)^{-\frac{870}{2299}}, \quad (2.54)$$

then the time from the second supernova to the conclusion of the inspiral, termed the *coalescence time*, will be given by

$$T_c(a_0, e_0) = \frac{12}{19} \frac{c_0^4}{\beta} \int_0^{e_0} \frac{e^{\frac{22}{19}} \left( 1 + \frac{121}{304} e^2 \right)^{\frac{1181}{2299}}}{(1 - e^2)^{\frac{3}{2}}} de. \quad (2.55)$$

The integral has no clean closed-form solution, and a numerical integrator is typically used to compute the coalescence time. However, some approximations such as Mandel (2021) do exist and approximate the coalescence time rather well over the entire range of mass ratios.

### 2.5.1 | Merger Rates of compact binaries

The merger rate of a population of compact binaries at time  $t$  after formation can be formalised via the heuristic

$$m(t) = \text{number of binaries for which } T_c + T_{\text{rej}} + T_{\text{evo}} < t, \quad (2.56)$$

where  $T_c$  is the coalescence time,  $T_{\text{rej}}$  is the rejuvenation age (the age at which the primary star last accreted mass), and  $T_{\text{evo}}$  is the evolution time-scale (the length of time the primary spends on the main sequence).  $m(t)$  is called the Delay-Time Distribution (DTD) of a population.

However, the underlying assumption in this is that the population was formed after an instantaneous burst of star-forming activity at time  $t = 0$ , which is clearly unphysical. We must account for the so-called Star Formation Rate Density (SFRD)  $\psi(z)$ , assumed to be taken from Madau & Dickinson (2014):

$$\psi(z) = 0.015 \frac{(1 + z)^{2.7}}{1 + \left( \frac{1+z}{2.9} \right)^{5.6}} \text{ M}_\odot \text{ yr}^{-1} \text{ Mpc}^{-3}, \quad (2.57)$$

where  $z$  is the redshift. For the purposes of converting between lookback time  $t$  and redshift  $z$ , we assume a standard  $\Lambda$ CDM cosmology with  $\Omega_M = 0.3, \Omega_\Lambda = 0.7, \Omega_k = 0, H_0 = 69.6 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

Of course, this ignores the effect of mixed-metallicity models, which we account for using the prescription of Langer & Norman (2006), whereupon the SFRD is multiplied by a factor  $\chi$ :

$$\chi(z, Z) = \frac{\hat{\Gamma}\left[0.84, \left(\frac{Z}{Z_{\odot}}\right)^2 \cdot 10^{0.3z}\right]}{\Gamma(0.84)}, \quad (2.58)$$

where  $\hat{\Gamma}$  and  $\Gamma$  are the upper incomplete and complete Gamma functions respectively. This means our Stellar Formation History (SFH) – which takes into account both redshift and metallicity – denoted  $\Psi(z, Z)$ , is given by

$$\Psi(z, Z) = 0.015 \int_{z_1}^{z_2} \frac{(1+z)^{2.7}}{1 + \left(\frac{1+z}{2.9}\right)^{5.6}} \frac{\hat{\Gamma}\left[0.84, \left(\frac{Z}{Z_{\odot}}\right)^2 \cdot 10^{0.3z}\right]}{\Gamma(0.84)} dz, \quad (2.59)$$

where  $z_1$  and  $z_2$  are the redshifts of our bin edges. The event rate as a function of time,  $\mathcal{R}(t)$ , is therefore given by

$$\mathcal{R}(t) = \Psi(z_t) * m(t), \quad (2.60)$$

where  $z_t$  is the redshift corresponding to lookback time  $t$ . Detail on how this convolution is computed may be found in Richards (2021).

## 2.6 | Numerical implementation

*I think there is a world market for maybe five computers.*

THOMAS WATSON, PRESIDENT OF IBM, 1943

Having now discussed the evolution of stars from a theoretical perspective, it becomes imperative to discuss the methods of modelling them. In this section, we provide an overview of the four codes we either use extensively or have developed to carry out the work of this thesis. These four codes are:

1. The Cambridge STARS code (subsection 2.6.1), originally developed by Peter Eggleton, which we use to solve the equations of stellar structure and evolution,
2. The TUI code (subsection 2.6.2), developed by Jan Eldridge, which performs our binary population synthesis,
3. The KAITIAKI code (subsection 2.6.3), developed for this thesis, which is our grid-manager code and allows modification of the input parameters to the Cambridge STARS code programmatically, and
4. The TAKAHE code (subsection 6.1.3), developed by Sean Richards for Richards (2021), which is our binary compact remnant evolution code.

### 2.6.1 | The Cambridge STARS code

#### Brief History

*And through the night, behind the wheel, the mileage clicking west  
I think upon Mackenzie, David Thompson and the rest  
Who cracked the mountain ramparts and did show a path for me  
To race the roaring Fraser to the sea.*

Northwest Passage, BY STAN ROGERS

Stellar modelling became a field of active research in the 1950s. Prior to the invention of the computer, a model was treated as a mathematical process and thus was computed – often labouriously – by hand. For instance, in Eggleton (1967), Eggleton writes that

Since we have seen in Section 3 that the structure of the flame zone can be obtained independently (almost) of the rest of the star, it is reasonable to expect that instabilities of the type considered by Schwarzschild & Härm can also be examined in this way, so that the considerable labour of computing the structure of an entire star can be avoided.

– Eggleton (1967)

The Cambridge STARS code was one of the first detailed stellar evolution codes to be written, though initially it only modelled one star in detail. To model a binary system, tweaks were made whereupon the primary would be modelled in detail whilst approximating the secondary with the equations of Hurley et al. (2002) until the primary supernova, at which point the secondary would be evolved in detail. This had its merits; notably computation time was saved, and the results were generally accurate anyway.

However, Stancliffe & Eldridge (2009) introduced detailed modelling of *both* stars in a binary system into the STARS code – that is, the interiors of both stars were computed in detail, alongside the simultaneous computation of the orbital parameters. Since then, a wide array of improvements have been made to the numerical stability of the code, as well as introducing or refining new physics.

There are four variants of this code:

1. The Cambridge STARS code, which is the version originally developed by Peter Eggleton and originally described in Eggleton (1971). This version was originally only capable of solving the equations for a single star.
2. The ELDRIDGE STARS code described in Eldridge & Tout (2004b,a); Eldridge et al. (2017), which is the version used to generate the fiducial BPASS v2 grids. This version models the primary in detail while approximating the secondary using the equations of Hurley et al. (2002).
3. The BSTARS code, which is the unmodified version lately available from Cambridge, described in Stancliffe & Eldridge (2009). This version models both stars in detail throughout their lives.
4. The AOTEAROA STARS code, which is the version of the code incorporating the changes we have made.

Unless context otherwise requires, when we refer to “the STARS code” with no further qualification, we are referring specifically to the AOTEAROA STARS code.

## Matrix Inversion

The heart of STARS is a matrix inversion algorithm. Although computing the inverse of a matrix is a polynomial time process, it can still take considerable time to invert. STARS implements a Henyey matrix (Henyey et al., 1964), which is of size  $n = (4 + n_{\text{comp}})N_{\text{mesh}}$ , with  $n_{\text{comp}}$  the number of equations to be solved and  $N_{\text{mesh}}$  the number of meshpoints in the model. An individual element of the Henyey matrix is given by

$$\mathbf{H}_{ij} = \frac{\partial (\text{equation } i)}{\partial (\text{variable } j)}. \quad (2.61)$$

For a “typical” binary computation, with  $n_{\text{comp}} = 30$  equations and  $N_{\text{mesh}} = 199$  meshpoints, the matrix is already  $6766 \times 6766$ , for a binary which evolves up the AGB (the theoretically most complex calculation we

perform) with  $N_{\text{mesh}} = 999$ , the matrix is  $33966 \times 33966$ . Introducing  $k$  new equations into the code causes the runtime of the matrix inversion calculation to scale like  $\mathcal{O}(n^3 + k^3 N_{\text{mesh}}^3)$ .

The matrix inversion is simple Gauss-Jordan inversion, which is of complexity class (for an  $n \times n$  matrix)  $\mathcal{O}(n^3)$ . Coppersmith & Winograd (1987) showed that this could be optimised to  $\mathcal{O}(n^{2.376})$  and conjectured reducibility to  $\mathcal{O}(n^2)$ .<sup>8</sup> Since our code is embarassingly parallelisable, and can be executed in parallel simply by increasing the number of cores available, the payoff for decreasing the runtime by one polynomial order is inconsequential, and thus we use the straightforward implementation.

## Initial Condition Selection

In this section we provide an account for the selection of our general initial conditions. Unless otherwise noted in this thesis, it is assumed that the values will be using the parameters described here and in Appendix D.

In our implementation of Mixing Length Theory (MLT) (Vitense, 1953; Böhm-Vitense, 1958; Cox & Giuli, 1968), we use an MLT parameter of  $\alpha_{\text{MLT}} = 1.985$ , selected via calibration against a model of the Sun. This is effectively a dissipativity parameter, and represents the depth at which a parcel of matter penetrates into a lower layer before becoming fully dissipated. The value was selected by modelling a  $1 M_{\odot}$  star and varying only the mixing length parameter until it had a luminosity of  $1 L_{\odot}$  and a radius of  $1 R_{\odot}$ , at the present age of the Sun.

## Mesh spacing

A natural question that arises when modelling a star is where internally to model it – there is a counterbalance between the level of detail and the time cost of the matrix inversion. The obvious answer would be to distribute a mesh equally spaced within radius of the star, however the nonhomogeneity of a star, plus the potential for very thin shells of physically important material, necessitate a more intelligent approach.

Instead of being equally spaced *radially*, these mesh points are taken to be equally spaced within a function  $\mathcal{Q}$ , which is given for a binary that does not have AGB-specific treatment enabled by

$$\begin{aligned} \mathcal{Q}(P, T, M, R) = & c_4 \ln P + c_5 \ln(P + c_{\text{H}}) + c_2 \ln(P + 3P_{\text{He}}) + c_2 \ln(P + 0.3P_{\text{He}}) + c_7 \ln\left(\frac{T}{T + 2 \times 10^4}\right) \\ & + \ln\left(\frac{c_6 M^{\frac{2}{3}}}{M + M^{\frac{2}{3}}}\right) - c_3 \ln\left(\frac{R^2}{c_8} + 1\right), \end{aligned} \quad (2.62)$$

where  $P, T, M, R$  are the total pressures, temperatures, masses, and radii of each star, the constants  $c_i$  are user controllable and given in Appendix D,  $P_{\text{He}}$  is the pressure in the Helium-burning shell, and

$$c_{\text{H}} = \begin{cases} 0.1P_{\text{H}} & X_{\text{H}} \geq 0.15 \\ 0.1\mathcal{P} & \text{otherwise} \end{cases} \quad (2.63)$$

where  $P_{\text{H}}$  is the pressures in the Hydrogen-burning shell. Here,  $\mathcal{P}$  represents the pressure in the Hydrogen-burning shell when  $X_{\text{H}} = 0.15$ . The choice of mesh spacing coefficients  $c_1 \dots c_{10}$  leads to different areas of the star being prioritised. Coordinates  $c_1, c_9$ , and  $c_{10}$  are not used except in AGB-specific evolution, which is not required for most cases (and requires a change of parameterisation in the pressure-shell terms).  $c_5$  is hardcoded to be 0.15, and  $c_2$  to be 0.0. **[TODO: but why? “its what the code does” is a bad reason.]**

Placement of the mesh is controlled by the derivative  $\frac{\partial \mathcal{Q}}{\partial k}$ , where  $k$  is the meshpoint number (Stancliffe, 2016). Specifically, meshpoints are placed at locations where

<sup>8</sup>The current “best” optimisation is  $\mathcal{O}(n^{2.3727})$  (Williams, 2012). DeepMind’s AlphaTensor has claimed a faster result in Fawzi et al. (2022), though they do not state a complexity class. This is because “...this does not yield practical algorithms.”

$$\left. \frac{\partial \mathcal{Q}}{\partial k} \right|_k - \left. \frac{\partial \mathcal{Q}}{\partial k} \right|_{k+1} = 0. \quad (2.64)$$

The mesh is thus adaptive (insofar as it changes to place meshpoints where they are needed for evolution) and non-Lagrangian (insofar as the derivative is not at constant mass coordinate). If one desires to make the mesh viscous (i.e., partially Lagrangian), one can follow Stancliffe et al. (2004) and write, for  $0 \leq \gamma \leq 1$ ,

$$\left( \left. \frac{\partial \mathcal{Q}}{\partial k} \right|_k - \left. \frac{\partial \mathcal{Q}}{\partial k} \right|_{k+1} \right) (1 - \gamma) + \gamma \left( \left. \frac{\partial m}{\partial t} \right|_k \right) = 0, \quad (2.65)$$

where now the parameter  $\gamma$  is a weighting coefficient such that if  $\gamma = 1$ , then the mesh spacing function is irrelevant and the parameters are spaced equally in mass, and if  $\gamma = 0$ , the original mesh spacing function is used. In between, if  $\gamma = \gamma(k, dt)$ , then parts of the mesh are allowed to be Lagrangian and parts non-Lagrangian; the mesh is said to be *viscous*. We typically keep the mesh non-viscous and non-Lagrangian, as a viscous mesh is most useful for AGB evolution and evolving through thermal pulses.

More detail is provided in Eggleton (1971) for the original mesh spacing function, and Stancliffe et al. (2004) for the binary mesh spacing function.

## Timestep control

The STARS code utilises an adaptive timestep control mechanism, whereby the timestep is allowed to shrink to model rapid changes or grow if the star is broadly inert and unchanging. The core computation is a parameter  $\Delta$ , given by

$$\Delta = \frac{d_{\text{opt}}}{\sum_i \sum_j |\delta x_{ij}|} \quad (2.66)$$

where  $d_{\text{opt}}$  is a pre-defined target timestep (typically taken to be  $\sim 1$ , unless otherwise specified; see Appendix D for our selections), and  $\delta x_{ij}$  is the change in variable  $x_i$  at meshpoint  $j$ . In this calculation, luminosity is explicitly excluded as this would cause the timestep to explode during certain points of the star's life-time. A new timestep,  $d'$ , is selected from the previous timestep  $d$  by


$$d' = \begin{cases} \min[d_1, \Delta] d & \Delta < 1 \\ d & \Delta = 1 \\ \max[d_2, \Delta] d & \Delta > 1 \end{cases} \quad (2.67)$$

$d_1$  and  $d_2$  are pre-determined control parameters, most commonly we take  $d_1 = 0.1$  and  $d_2 = 1.2$  unless otherwise specified.

## Numerical instabilities

Stellar modelling is a nontrivial endeavour and there are many cases where the code can fail seemingly arbitrarily. The largest cause of numerical instability is the helium flash, which occurs in low-mass stars between about  $0.8 M_{\odot} \lesssim M_{\text{ZAMS}} \lesssim 2.5 M_{\odot}$ . In this band, the triple- $\alpha$  process, given by the reactions (Carroll & Ostlie, 2017)<sup>9</sup>

<sup>9</sup>The reversible reaction is due to Beryllium-8 being unstable; its half-life is on the order of  $1 \times 10^{-16}$  s, and it decays back to  $2^4_2\text{He}$  if it does not react further.



occurs within a highly degenerate helium core. As a result, the temperature of the star increases sharply during this process, leading to thermal runaway. This temperature increase in turn results in an increase in the luminosity (which peaks at  $\sim 1 \times 10^{10} L_{\odot}$ ). The flash itself – being the period of thermal runaway – occurs on the order of a few minutes, though the spike in luminosity lasts approximately 7 Myr (for the  $2.51 M_{\odot}$  star shown in Figure 2.12).

A method termed “pseudoevolution” can be used to overcome this source of error. Briefly, a complete model (termed the *reference model*) of a  $3 M_{\odot}$  star is evolved with a full model description saved at every timestep. When the helium flash occurs in a model (heuristically,  $0.1 \leq M_{\text{He}}/M_{\odot} \leq 0.55$  and  $\log T_c/\text{K} > 7.9$ ), we query the reference model for a model with the same helium core mass as the failed model. Using a custom subroutine in STARS, we turn off all nuclear and thermal burning and remove mass from the reference model to match the mass of the failed model. This model is then allowed to evolve as per normal, having overcome the helium flash. We note that this is not a *perfect* method, as this does induce discontinuities in the plots, as shown in Figure 2.13, though these are essentially insignificant.

The other stage where low-mass stars are prone to fail is during a thermal pulse (Schwarzschild & Härm, 1965, 1967). The mass range for thermal pulses is broader, from  $\sim 0.8 M_{\odot}$  to  $\sim 8 M_{\odot}$ . A thermal pulse occurs as a star ascends the AGB; as a Carbon-Oxygen core (CO core) or Oxygen-Neon core (ONe core) has developed, shells of burning helium and hydrogen (separated by a region of almost pure helium) sit atop the core (Pols et al., 2001). As the intershell region dissipates due to dredge up or helium burning, hydrogen burning reignites in the hydrogen shell. As both shells burn (at the same rate), the helium-burning shell becomes unstable and eventually a process of thermal runaway begins (similar to the helium flash, except off-core). This causes a spike in the luminosity, termed a thermal pulse. This can persist for many pulses – Stancliffe (2006) observes pulses before model termination, while Stancliffe (2005) observes 17 pulses for a model similar to those in Figure 2.12.

The method to overcome this numerical instability is involved and not fully implemented in STARS. Briefly, the number of meshpoints must be drastically increases near the onset of thermal pulses; up to 999 instead of the typical 199/499. The timestep must also be forcibly decreased; thermal pulses typically last for around a year (Pols et al., 2001) so the timestep must be reduced to not miss them (and to not have the modulus of the change in parameters –  $\sum_i \sum_j |\delta x_{ij}|$  in Equation 2.66 – and thus the error, increase too much). This method was implemented to generate Figure 2.12, though only two pulses were achievable.


## Critical modifications

The major modifications we have made to the STARS code are as follows.


Firstly, the floating-point precision has been increased to double precision for all variables. This has minor effects on the evolved models, insofar as they are generally able to evolve to slightly further stages. This step has previously been undertaken, however not all variables were correctly converted. This did not invalidate any models previous versions of the STARS code has generated; rather it means they terminated slightly earlier than they may have been able to evolve to.

As part of this, we also undertook a reasonably extensive refactor of the codebase. Whilst some of the code remains as originally written, it is now more compliant with modern programming practices and is in some sense easier to alter with new routines.

We have also improved mass loss interpolation in the MASSLOSS.F routine. Whilst not changing the underlying



**Figure 2.12:** Two stars of a similar mass that undergo a helium flash and two thermal pulses. Shown in the inset is a zoomed-in helium flash that lasts  $\sim 7$  Myr. Both stars undergo two thermal pulses before the code crashes due to the relative error in the calculation increasing too rapidly. The two stars could, in principle, undergo many more thermal pulses before their deaths.



**Figure 2.13:** The helium flash (gray dashed line) shown on a Kippenhahn diagram. There is a slight discontinuity in the helium core mass (blue) due to a perfect match not existing in the  $3 M_{\odot}$  reference model. The total mass is shown as the solid gray line, and the CO core mass is shown in red. The green lines show the boundaries of the convective regions of the star.

assumption that mass transfer is governed by de Jager et al. (1988); Vink et al. (2001) (subsection 2.4.3), we have improved the transition between typical O-type mass loss and WR mass loss rates by slowing turning on the WR mass loss rate as the surface hydrogen abundance drops below 0.4. The transition was previously abrupt and would cause the code to fail due to numerical instability if a non-WR star entered this regime.

Similarly to this, the code now manually reduces the timestep in the run-up to, during, and after the contact binary phase. Evolution during this phase breaks thermal equilibrium (i.e., the star evolves on a longer timescale than its Kelvin-Helmholtz time-scale) and hence the evolution can change rapidly as the star attempts to adjust back to thermal equilibrium. The contact phase, for example, typically last for between 100-200 kyr in our models.

We have also begun implementing an  $\alpha - \lambda$  treatment of CEE, though we stress that this has not been rigorously tested yet, and is thus disabled for the majority of our models.

## 2.6.2 | The TUI code

TUI is the core population synthesis code within the BPASS project. TUI takes as input the output from the STARS code. TUI was first described in Ghodla et al. (2023), and was more fulsomely described in Stevance et al. (2023), though we note that their version is technically a refactor of the version we use. The predominant function we use TUI for is the matching of pre- and post- supernova models for chapter 6, and in this section we provide a brief account of that functionality of TUI (though the control of the population synthesis models is beyond the scope of this thesis).

TUI has precompiled “input\_” files which report a STARS plot file, an IMF weight, and a model type (merger/normal primary/normal secondary/single model/quasi-homogeneous evolution binary/single star). After loading in a model, TUI applies a given kick and computes the new orbital parameters from Tauris & Takens (1998). Crucially, a core assumption of BPASS is that the orbit remains circular after the first supernova and is allowed to be eccentric after the second. Of course, nature cares little for our assumptions, and the supernova may cause the binary to become eccentric after the first supernova; TUI accounts for this by setting the separation after the first supernova to be the semi-latus rectum of the eccentric binary. TUI repeats this process after the second supernova.

For the purposes of chapter 6, if the binary remains bound (has  $0 \leq e < 1$ ) after the second supernova, we

record the following parameters:

1. The mass of the primary,
2. The mass of the secondary,
3. The orbital separation,
4. The eccentricity,
5. A weight term proportional to the IMF weighting,
6. The rejuvenation age, which is the time since the companion last accreted material, if it accreted more than five per cent of its initial mass (see Richards et al., 2023), and
7. The evolution age, which is the length of time from the secondary’s ZAMS until its death.

This provides us with the post-secondary supernova orbital parameters which determine the properties of the merger, such as the event rate  $\mathcal{R}(z)$  and its path through period-eccentricity space.

### 2.6.3 | The KAITIAKI grid-watcher code

KAITIAKI is a grid-watcher code we have developed that sits atop the STARS code and manages its execution. It was developed primarily to manipulate the fixed-width text files that STARS uses, in contrast to the “typical” manipulation done through the command line. However development proceeded in such a way that it now provides an interface to all of the (relevant to this thesis) output and input files. Further, it also now allows the user to control jobs through a PYTHON interface, greatly simplifying grid calculations.

The core of KAITIAKI is the `STARSController()` interface. The most naïve run would be to initialise a model of, say,  $8 M_{\odot}$  and evolve it until it evolves up the AGB. A sample of code for this is shown in Code Sample 2.1 which evolves a  $10 M_{\odot}$  star.

Code Sample 2.1: A KAITIAKI script to evolve a  $10 M_{\odot}$  star

```

1 import kaitiaki
2
3 # First, instantiate the STARS code. run_bs is the location of the run_bs
4 # bash script, which sets up the symlinks and actually runs the code.
5 STARS = kaitiaki.STARS.STARSController(run_bs='..')
6
7 # Loads the default data file and solar metallicity opacity table
8 STARS.blit('z020')
9
10 # Loads a 1 Msun model
11 STARS.load_default_modin()
12
13 # Sets the code up to target a ZAMS mass of 10 Msun.
14 STARS.setup_zams_inflation(10)
15
16 # Executes the code, such that the final modout will be a 10 Msun model.
17 STARS.run()
18
19 # Equivalent to "tail -399 modout > modin" -- transfers the last 2*199+1
20 # lines from modout (the last converged model) to modin
21 STARS.modout_to_modin()
22
23 # Again, to reload data

```

```

24 STARS.blit()
25
26 # Set up the key parameters for our run
27 STARS.configure_parameters({
28     'DT1': 0.1, # Fraction of timestep to allow for lower variation
29                 # (if the timestep is cut, it can be cut to 10% of its
30                 # present value)
31     'DDD': 2,   # Modulus of total desired timestep
32     'IML1': 5,  # WR mass-loss rates
33     'RML': 0,   # This was set to 10 Msun during inflation -- should be
34                 # zero so no mass is added except via the usual ML channels
35                 # (this parameter, formally, sets a constant ML rate so
36                 # that in the absence of other mechanisms, the mass would
37                 # always decrease)
38     'IMODE': 1, # Single star mode
39     'IX': 1,   # Hydrogen burning
40     'IY': 1,   # Helium burning
41     'IZ': 1,   # Metal burning
42     'ITH': 1,  # Thermal energy generation
43     'ISTART': 1 # Reset age, NMOD, and the timestep (to 10% of T_KH)
44 })
45
46 # Execute the model, run until the AGB.
47 STARS.run()

```

## 2.7 | Chapter Summary

*Looking up, he saw a thousand stars glittering in the deep velvet of a night with no moon. He knew them all, their stories and their names. He knew them in a familiar way, the way he knew his own hands.*

---

*The Name of the Wind*, BY PATRICK ROTHFUSS

In this Chapter, we have explored the evolution of stars from pre-main sequence evolution to supernova, accounting for the effects of binary interaction. This represents the state of the field as at the commencement of this work. In future chapters we will detail the relevant changes we have made to these prescriptions, motivating the work we have performed.

# Detailed Models of Overcontact Binary Stars

*There are more things in Heaven and Earth, Horatio, than are  
dreamt of in your philosophy.*

HAMLET

**A**S PREVIOUSLY DISCUSSED, STARS undergoing mass transfer will generally experience RLOF during their lifetimes. In a binary where both stars are undergoing mass transfer, both stars may undergo RLOF simultaneously, resulting in a contact binary system in which both stars physically touch. Contact binaries are not uncommon – they form one in every approximately 100 main-sequence stars (Rucinski, 1998) and 39 in every 100 binary systems (Green et al., 2023).

The evolution is also complex. Sufficiently close binaries may overflow their  $\mathcal{L}_2$  points, or create mergers on exceptionally short timescales – typically  $P_{\text{orb}} \lesssim 2$  d, but periods as low as  $P_{\text{orb}} = 85$  min (El-Badry et al., 2022) are possible in theory or  $\sim 167$  min in practice (Koen, 2022).

## 3.1 | Observations of contact binaries

Observations of contact binaries date back to the 1940s (Kuiper, 1941), however only a handful of contact binaries are known. The binaries we study in this chapter are summarised in Table 3.1, and were originally compiled in Abdul-Masih et al. (2022). Their sample focused on measuring the period stability of contact binaries and is a subset of the known sample, of which somewhere between  $\sim 42\,000$  (Petrosky et al., 2021) and  $\sim 86\,000$  (Qian et al., 2020) binaries exist.

## 3.2 | Data matching with BPASS v2.2

We perform our data matching using the HOKI code (Stevance et al., 2020a). The search parameters are shown in Table 3.1. We query the BPASS v2.2.1 dataset and record whether or not they enter the (widened) parameter values. Parameters are widened as some of the periods are tighter than the BPASS values and thus can be missed in the search.

The matching values HOKI finds are shown in Table 3.2. The primary models are also shown on Figure 3.1.


### 3.2.1 | Progenitor genealogy

We now examine each of the progenitor histories in detail, focusing in particular on the ZAMS parameters of each model, as well as providing observational context to each system.

#### LSS 3074


The binarity of LSS 3074 (Raucq et al., 2017) was first observed in 1985 by Morrell & Niemela (1990) with an observed period of 2.19 d, with its HeI and HeII spectra causing the authors to conclude that it was an O4f+ and O6-7 binary, with inclination-weighted masses of  $M_1 \sin^3 i = 8 M_\odot$  and  $M_2 \sin^3 i = 9 M_\odot$ . With later measurements of the inclinations to be  $i \approx 24^\circ$ , these masses were refined to  $M_1 = 17.2 M_\odot$  and  $M_2 = 14.8 M_\odot$ . It is located behind the Coalsack Nebula, at a distance of 2256 pc, with a B-band magnitude of 12.81 mag (Stephenson & Sanduleak, 1971).

LSS 3074 matches according to HOKI




(a) LSS 3074

MY Cam matches according to HOKI




(b) MY Camelopardalis

SMC 108086 matches according to HOKI




(c) SMC 108086

TU Mus matches according to HOKI




(d) TU Muscae

V382 Cyg matches according to HOKI



(e) V382 Cygni

VFTS 352 matches according to HOKI



(f) VFTS 352

**Figure 3.1:** BPASS v2.2.1 primary models for the matches obtained by HOKI for the parameters in Table 3.1. The legend key is the BPASS stellar model file name, and is broken down into its parts in Table 3.2.

| Binary name           | $M_1$          | $M_2$          | $P$                   | $Z$       | $\log T_1$      | $\log L_1$      | $\log T_2$     | Ref. |
|-----------------------|----------------|----------------|-----------------------|-----------|-----------------|-----------------|----------------|------|
| LSS 3074              | $14.8 \pm 2.2$ | $17.2 \pm 4.2$ | $2.183 \pm 1$         | $0.020^a$ | $4.6 \pm 0.2$   | $5.15 \pm 0.2$  | $4.53 \pm 0.5$ | A    |
| MY Cam                | $37.7 \pm 3.2$ | $31.6 \pm 4.2$ | $1.175^{+6}_{-1.175}$ | $0.020^a$ | $4.62 \pm 0.2$  | $5.2 \pm 1.0$   | $4.59 \pm 0.5$ | B    |
| SMC 108086            | $16.9 \pm 1.2$ | $14.3 \pm 5.1$ | $0.883 \pm 0.88$      | $0.004^b$ | $4.53 \pm 0.1$  | $4.608 \pm 0.4$ | $4.48 \pm 0.5$ | C    |
| TU Mus                | $16.7 \pm 1.6$ | $10.4 \pm 1.4$ | $1.387 \pm 1$         | $0.020^c$ | $4.57 \pm 0.2$  | $5.02 \pm 0.4$  | $4.54 \pm 0.5$ | D    |
| V382 Cyg <sup>d</sup> | $26.1 \pm 3.2$ | $19.0 \pm 4.2$ | $1.886 \pm 1$         | $0.020^e$ | $4.57 \pm 0.2$  | $5.47 \pm 0.4$  | $4.58 \pm 0.5$ | E    |
| VFTS 352              | $28.6 \pm 3.6$ | $28.9 \pm 1.8$ | $1.124^{+2}_{-1.124}$ | $0.010^f$ | $4.63 \pm 0.45$ | $5.19 \pm 1.0$  | $4.61 \pm 0.5$ | F    |

**Table 3.1:** Search parameters for the six close-contact binaries compiled in Abdul-Masih et al. (2022), as used in our fitting to the BPASS v2.2.1 grid. The uncertainties presented are not the physical uncertainties on their parameters, but instead represent the size of the range that we search within, e.g.,  $P = 1.175^{+6}_{-1.175}$  indicates that we match on all periods between 0 and 7.165 d inclusive. Masses and Luminosities are measured relative to solar, periods are measured in days, and temperatures measured in Kelvin. **Notes:** <sup>a</sup>Source paper assumes solar metallicity with no further qualification <sup>b</sup>Using the mean metallicity of the SMC <sup>c</sup>Based on Andersen & Grønbech (1975) claiming 0.021 and Terrell et al. (2003) confirming a comparison with synthetic spectra “does not support a carbon content, and therefore metallicity, far from the solar value” <sup>d</sup> A mass ratio inversion occurred for this binary (see Martins et al., 2017) and hence we use the definition of “primary” from within the paper when reporting in this table. <sup>e</sup>Using the mean metallicity of the Milky Way <sup>f</sup>As Crowther et al. (2018) reports “half-solar” for the Tarantula Nebula, the host nebula of VFTS 352. **References:** A: Raucq et al. (2017), B: Lorenzo et al. (2014), C: Hilditch et al. (2005), D: Penny et al. (2008), E: Martins et al. (2017), F: Almeida et al. (2015)

### MY Camelopardalis

MY Camelopardalis (MY Cam; Lorenzo et al., 2014) is a binary in the cluster Alicante 1, and its binarity was established in 2003 by Negueruela & Marco (2003). It is likely to be a very massive merger progenitor, though is barely in contact with a fillout factor  $f = 1.01$  (Abdul-Masih et al., 2022).<sup>1</sup>

### SMC 108086

OGLE SMC-SC10 108086 (SMC 108086; Hilditch et al., 2005) is a O9 + O9.5/B0 binary. It is a system deeply in contact, with  $f = 1.70$  (Hilditch et al., 2005; Abdul-Masih et al., 2022) – the deepest known – and has a short period of  $P = 0.88$  d. It is located at a distance of 56.5 kpc, in the Small Magellanic Cloud (SMC).

### TU Muscae

TU Muscae (TU Mus; Penny et al., 2008) is an O8V + B0V binary discovered by Oosterhoff (1928, 1930). Its binarity was confirmed as late as 1971, in Knipe (1971). The system was profiled in Andersen & Grønbech (1975), who concluded that, based on comparison to theoretical models by Stothers (1972, 1973), the system was of solar composition ( $Z = 0.021 \approx Z_\odot$ ) and on the main sequence with  $X_H = 0.739$ .

### V382 Cygni

V382 Cygni (V382 Cyg; Martins et al., 2017) is an O6.5 + O6 binary whose components are oblate spheroids (deduced from the  $\beta$ -Lyrae light curves; Pearce 1952). It was originally discovered by Morgenroth (1935) and A. A. Petrov in 1946 (Landolt, 1964; Yaşarsoy & Yakut, 2012).<sup>2</sup>

<sup>1</sup>A fillout factor is a measurement of the degree of which a system overfills its Roche lobes, with  $1 < f < 2$  denoting an overcontact binary, and a high fillout factor representing a more swelled and in-contact system.

<sup>2</sup>Petrov’s paper in *Peremennye Zvezdy* is not available digitally, and given the ongoing Russia-Ukraine conflict, the author is unable to access a physical copy to verify the citation and cite this statement.

| Binary     | BPASS v2.2.1 matches    | $M_{1,ZAMS}$ | $M_{2,ZAMS}$ | $q$ | $P_{ZAMS}$ | log Age |
|------------|-------------------------|--------------|--------------|-----|------------|---------|
| LSS 3074   | sneplot-z020-30-0.3-0.6 | 30.0         | 9.0          | 0.3 | 3.98       | 6       |
|            | sneplot-z020-25-0.4-0.6 | 25.0         | 10.0         | 0.4 | 3.98       | 6       |
|            | sneplot-z020-25-0.5-0.6 | 25.0         | 12.5         | 0.5 | 3.98       | 6       |
|            | sneplot-z020-24-0.4-0.6 | 24.0         | 9.6          | 0.4 | 3.98       | 6       |
|            | sneplot-z020-24-0.5-0.6 | 24.0         | 12.0         | 0.5 | 3.98       | 6       |
| MY Cam     | sneplot-z020-70-0.3-1   | 70.0         | 21.0         | 0.3 | 10.00      | 6.4     |
|            | sneplot-z020-60-0.3-0.8 | 60.0         | 18.0         | 0.3 | 6.31       | 6.4     |
|            | sneplot-z020-60-0.4-0.8 | 60.0         | 24.0         | 0.4 | 6.31       | 6.4     |
|            | sneplot-z020-60-0.3-1   | 60.0         | 18.0         | 0.3 | 10.00      | 6.4     |
|            | sneplot-z020-50-0.5-0.6 | 50.0         | 25.0         | 0.5 | 3.98       | 6.4     |
|            | sneplot-z020-50-0.4-0.8 | 50.0         | 20.0         | 0.4 | 6.31       | 6.4     |
|            | sneplot-z020-50-0.5-0.8 | 50.0         | 25.0         | 0.5 | 6.31       | 6.4     |
|            | sneplot-z020-50-0.6-0.8 | 50.0         | 30.0         | 0.6 | 6.31       | 6.4     |
|            | sneplot-z020-40-0.7-0.8 | 40.0         | 28.0         | 0.7 | 6.31       | 6.4     |
| SMC 108086 | sneplot-z004-24-0.5-0.4 | 24.0         | 12.0         | 0.5 | 2.51       | 7       |
|            | sneplot-z004-23-0.4-0.4 | 23.0         | 9.2          | 0.4 | 2.51       | 7       |
|            | sneplot-z004-23-0.5-0.4 | 23.0         | 11.5         | 0.5 | 2.51       | 7       |
|            | sneplot-z004-22-0.4-0.4 | 22.0         | 8.8          | 0.4 | 2.51       | 7       |
|            | sneplot-z004-22-0.5-0.4 | 22.0         | 11.0         | 0.5 | 2.51       | 7       |
|            | sneplot-z004-21-0.4-0.4 | 21.0         | 8.4          | 0.4 | 2.51       | 7       |
|            | sneplot-z004-21-0.5-0.4 | 21.0         | 10.5         | 0.5 | 2.51       | 7       |
|            | sneplot-z004-20-0.4-0.4 | 20.0         | 8.0          | 0.4 | 2.51       | 7       |
|            | sneplot-z004-20-0.5-0.4 | 20.0         | 10.0         | 0.5 | 2.51       | 7       |
|            | sneplot-z004-19-0.4-0.4 | 19.0         | 7.6          | 0.4 | 2.51       | 7       |
| TU Mus     | sneplot-z020-25-0.3-0.4 | 25.0         | 7.5          | 0.3 | 2.51       | 6.2     |
|            | sneplot-z020-24-0.3-0.4 | 24.0         | 7.2          | 0.3 | 2.51       | 6.2     |
|            | sneplot-z020-23-0.3-0.4 | 23.0         | 6.9          | 0.3 | 2.51       | 6.2     |
|            | sneplot-z020-16-0.6-0.4 | 16.0         | 9.6          | 0.6 | 2.51       | 6.2     |
| V382 Cyg   | sneplot-z020-50-0.2-0.6 | 50.0         | 10.0         | 0.2 | 3.98       | 6.4     |
|            | sneplot-z020-40-0.3-0.6 | 40.0         | 12.0         | 0.3 | 3.98       | 6.4     |
|            | sneplot-z020-40-0.4-0.6 | 40.0         | 16.0         | 0.4 | 3.98       | 6.4     |
|            | sneplot-z020-35-0.4-0.6 | 35.0         | 14.0         | 0.4 | 3.98       | 6.4     |
|            | sneplot-z020-30-0.4-0.6 | 30.0         | 12.0         | 0.4 | 3.98       | 6.4     |
|            | sneplot-z020-30-0.5-0.6 | 30.0         | 15.0         | 0.5 | 3.98       | 6.4     |
| VFTS 352   | sneplot-z010-40-0.5-0.6 | 40.0         | 20.0         | 0.5 | 3.98       | 6.5     |

**Table 3.2:** Matches from the HOKI (Stevance et al., 2020a) software. Ages were computed via HOKI’s AgeWizard utility, described in Stevance et al. (2020b), and are the most likely ages computed via a bootstrapping method.

## VFTS 352

VFTS 352 was detected in the VLT-FLAMES survey of the Tarantula Nebula (Evans et al., 2011) and is the most recent detection in our sample. The Tarantula Nebula / 30 Doradus is a large H(II) region in the Large Magellanic Cloud (LMC). The progenitor of VFTS 352 is likely an equal mass, short period binary with no strong helium or nitrogen surface abundances (Abdul-Masih et al., 2019), which may be due to biases in the CNO abundances, or similar, in that atmospheric study.

## 3.3 | Forward evolution

Having now established the ZAMS parameters for our binaries, we can take these as input to our STARS code to determine their future history.

## 3.4 | Exotic binaries arising from contact systems

### 3.4.1 | Double contact systems

### 3.4.2 | Reverse mass transfer

### 3.4.3 | Overflow through $\mathcal{L}_2$



# Helium Rich Mass Transfer

*We, all of us, are what happens when a primordial mixture of hydrogen and helium evolves for so long that it begins to ask where it came from.*

JILL TARTER

SINCE THE FORCE BETWEEN TWO objects due to gravity scales like  $F \sim \frac{1}{r_{\text{sep}}^2}$ , another question naturally arises during the consideration of contact binaries. For sufficiently close binaries, one can imagine that mass transfer through RLOF could become reasonably strong, stripping the hydrogen envelope of a main sequence (or shortly post-main sequence) star and exposing the helium-enriched core underneath. If this stripping does not cause the star to retreat within its Roche lobe, then this core material would continue to be accreted, creating helium-enriched matter streams accreting onto its companion.

As discussed in section 2.4, the presence of a companion measurably alters the evolution of a star. Typically this manifests as allowing matter to be exchanged between the components or the gravity of one star warping the photosphere of the other. In this chapter we focus on the first instance, where mass transfer occurs between two stars. We consider a binary system with ZAMS masses of  $30 + 10 M_{\odot}$ , and an initial period of  $10^{0.6} \approx 3.98$  d. We specifically explore this case as a case study on the effect of Helium enriched mass transfer, and in chapter 5 we will explore the effect that this has on a grid of close binaries.

The general background for this section remains as elaborated in chapter 2, however our parameterisation requires us to delineate between two cases: that when variable composition accretion is enabled (referred to hereafter as “variable accretion”) and the case where it is disabled (hereafter “non-variable accretion”). These correspond to setting the parameter IVARACC in Appendix D to 1 or 0 respectively.

## 4.1 | Evolution overview


The major stages of the binary’s evolution can be broken up into five main stages, explored in the following section. Unless otherwise indicated, at every stage we shall be discussing the non-variable accretion case. The full evolution for the variable (non-variable) accretion case is shown in Figure 4.1 (Figure 4.2), with subplots showing the behaviour around the contact phase.

### 4.1.1 | Pre-interaction


The pre-interaction phase of both stars is identical in both cases. They evolve as a fully detached binary for 4.51 Myr, with the donor completing its main sequence evolution. Hydrogen is exhausted in the core, and a helium core develops. In both cases, as the donor begins to cross the Hertzsprung gap, it inflates to fill its Roche lobe, causing matter to begin to be exchanged via RLOF and kick-starting the pre-contact phase.

### 4.1.2 | Pre-contact


When the donor swells to fill its Roche lobe, it begins to efficiently transfer material to its companion. The increased drag causes the orbit to become hardened; shrinking by approximately 20 per cent. The accretor begins to accrete material, bleeding angular momentum into the system and causing the accretor to spin up. As the contact phase approaches, it approaches 15 per cent of the critical rotation rate, as shown in Figure 4.3.



**Figure 4.1:** The main evolution of a  $30 + 10 M_{\odot}$  binary with variable accretion enabled. Clockwise from top: the HR diagram with stages in Figure 4.4 and Table 4.1, the mass transfer efficiency, the separation / radius, the mass, and the surface abundances. In the lower four plots, the gray region shows the contact phase, and the plots are cropped to show the  $\sim 15$  kyr on each side of the contact phase. The dashed line on each HR diagram track shows when the relevant star is overfilling  $\mathcal{L}_1$ , and contact is when both stars are dashed.



**Figure 4.2:** As for Figure 4.1, but absent variable accretion.



**Figure 4.3:** The evolution of the radius, rotation rate, and luminosity in the presence of variable accretion, shown around the contact phase.

#### 4.1.3 | Contact phase

#### 4.1.4 | Post-contact

#### 4.1.5 | Post-interaction

### 4.2 | Atypical abundance profiles

Because the stars are quite close together and have a mass ratio of  $1/3$ , when RLOF starts it is quite voracious, and becomes even more so during the contact phase. The donor star sheds its hydrogen-rich surface and material from the helium-rich core is exposed, which too becomes part of the accreted material. The material stream becomes increasingly helium-rich, depositing this material on top of a hydrogen-rich surface, causing the accretor to have an atypical abundance profile, as shown pictorially in Figure 4.4. The net effect is that there is a helium-rich surface sitting atop a hydrogen-rich layer sitting atop a helium-rich core. At this stage thermohaline mixing is projected to occur, as there is an unstable composition gradient (owing to helium having a higher mean molecular weight than hydrogen).

### 4.3 | Comparison between variable and non-variable accretion


In Figure 4.5 we display a comparative HR diagram between the cases of variable and non-variable accretion. In each case an  $30 + 10 M_{\odot}$  binary was evolved with the only difference between the two being whether material was allowed to have variable composition accretion or the standard accretion prescription. In the bottom panel only the companion is shown, as the donor has identical evolution in both cases.

The first and most striking conclusion that we can draw from this result is that the companion is, in the variable accretion case, hotter and more luminous than the non-variable case.

|          |                             | $t = 0$<br>(a) | Star 1  |         | Star 2  |         | Star 1  |         | Star 2  |         |
|----------|-----------------------------|----------------|---------|---------|---------|---------|---------|---------|---------|---------|
|          |                             |                | Begin   | End     | Begin   | End     | Begin   | End     | Begin   | End     |
|          |                             |                | (b)     | (e)     | (c)     | (d)     | (b)     | (e)     | (c)     | (d)     |
| Donor    | $M / M_{\odot}$             | 30.00          | 27.93   | 7.56    | 26.91   | 17.43   | 27.93   | 7.57    | 26.91   | 17.27   |
|          | $R / R_{\odot}$             | 8.82           | 16.32   | 10.20   | 13.88   | 9.76    | 16.32   | 10.15   | 13.88   | 9.77    |
|          | $\log L / L_{\odot}$        | 5.07           | 5.30    | 5.25    | 5.16    | 4.94    | 5.30    | 5.25    | 5.16    | 4.98    |
|          | $X_{\text{surf}}$           | 0.70           | 0.70    | 0.09    | 0.70    | 0.57    | 0.70    | 0.09    | 0.70    | 0.57    |
|          | $Y_{\text{surf}}$           | 0.28           | 0.28    | 0.89    | 0.28    | 0.41    | 0.28    | 0.89    | 0.28    | 0.42    |
| Accretor | $M / M_{\odot}$             | 10.00          | 9.99    | 20.97   | 11.00   | 14.57   | 9.99    | 20.77   | 11.00   | 14.52   |
|          | $R / R_{\odot}$             | 4.87           | 4.36    | 4.02    | 9.23    | 8.98    | 4.36    | 6.53    | 9.23    | 9.02    |
|          | $\log L / L_{\odot}$        | 3.73           | 3.81    | 4.89    | 4.90    | 5.06    | 3.81    | 4.73    | 4.90    | 4.95    |
|          | $X_{\text{surf}}$           | 0.70           | 0.70    | 0.09    | 0.70    | 0.57    | 0.70    | 0.70    | 0.70    | 0.70    |
|          | $Y_{\text{surf}}$           | 0.28           | 0.28    | 0.89    | 0.28    | 0.41    | 0.28    | 0.28    | 0.28    | 0.28    |
| Binary   | $P_{\text{bin}} / \text{d}$ | 3.98           | 3.85    | 4.41    | 3.14    | 2.52    | 3.85    | 4.38    | 3.14    | 2.54    |
|          | $a / R_{\odot}$             | 36.14          | 34.74   | 34.56   | 30.28   | 24.71   | 34.74   | 34.33   | 30.28   | 24.78   |
|          | Age / kyr                   | 0.00           | 4707.16 | 6450.23 | 4712.75 | 4716.42 | 4707.16 | 6448.11 | 4712.75 | 4716.70 |


(a) Variable composition accretion
(b) Non-variable accretion

**Table 4.1:** The key evolutionary parameters of both stars and the binary configuration at five critical points: the initial parameters, and the beginning and ends of both epochs of RLOF. All values are rounded to 2 decimal places. The letter in parentheses represents which segment of Figure 4.4a (for Table 4.1a) or Figure 4.4b (for Table 4.1b) the abundances are given in.



**Figure 4.4:** The chemical composition of each star as a function of the stellar radius for the donor and accretor at the five key stages of the binary's lifetime as detailed in Table 4.1. Letters are as given in the same table. Each grid marking in the radial direction represents  $1 R_{\odot}$ .

## 4.4 | Chapter Summary



**Figure 4.5:** An HR diagram showing the evolution of our model both with (top left) and without (top right) the variable accretion prescription. The bottom panel shows only the evolution of the secondary. The point of divergence between the two models is shortly into the contact phase. The two dashed lines on the bottom top HR diagram represent models evolved as the final model of the binary secondary, as a single star. On the top HR diagrams, the pink and brown models are single stars evolved with ZAMS masses of 21 and  $26 M_\odot$  respectively.



CHAPTER 5

---

# Population properties of overcontact binaries and helium rich mass transfer



# Post-Supernova Kinematics: The Natal Kick

*And the memories were lost long ago  
But at least you have beautiful ghosts.*

*Beautiful Ghosts*, BY TAYLOR SWIFT

**T**HE TERMINUS OF A MASSIVE STAR’S life is a core-collapse supernova, when the hydrostatic equilibrium between the pressure of the star and its self-gravity is broken, and the stellar core collapses in on itself. Once the star reaches nuclear densities, neutron degeneracy is able to stabilise the remnant, causing a shockwave to propagate outwards from the core. The energy released is roughly an order of magnitude greater than the binding energy of the star (Eldridge & Tout, 2018), resulting in the ejection of an amount of mass we denote  $M_{\text{ej}}$  and leaving a remnant of mass  $M_{\text{rem}}$  behind. If the ejection is asymmetric, the collapse endows the nascent remnant with a velocity  $v_k$  whose overall direction is antialigned to the more massive ejecta. This motion is termed a natal supernova kick.

Given an observation of the proper velocity of a compact remnant, it is in principle possible to trace its motion back and ascertain its kick velocity. However, this is not necessarily a straightforward endeavour. If the proper velocity is high and the remnant old, it has travelled quite a distance from its explosion site to be located at the point the observation has been taken. In addition, the star has potentially had its proper velocity modified by gravitational attraction from either other runaway stars or its host galaxy’s potential. One would require the exact location in 3D space as well as the 3D velocity and remnant age in order to ascertain the motion, but from there the inverse problem becomes straightforward.

## 6.1 | Overview

As a consequence of conservation of momentum, it can be shown that if the ejecta is assumed to eject along a preferred axis, the velocity of the kick can be written as

$$v_k = \alpha \left( \frac{M_{\text{ej}}}{M_{\text{rem}}} \right) + \beta \left( \frac{M_{\text{NS}}}{M_{\text{rem}}} \right), \quad (6.1)$$

for free parameters  $\alpha$  and  $\beta$  (both measured in  $\text{km s}^{-1}$ ). Here,  $M_{\text{NS}}$  is the mass of a Neutron Star (NS); if the kick is applied to a NS, then  $M_{\text{NS}} = M_{\text{rem}}$  and the coefficient reduces to unity. If the kick is applied to a Black Hole (BH), then we can take  $M_{\text{NS}} = 1.4 M_{\odot}$ , which serves to convert the distribution from one in velocity-space to one in momentum-space; see Bray & Eldridge (2016, 2018); Richards et al. (2023) for further detail.

This parameterization introduces, effectively, four free parameters in  $\alpha, \beta, \theta, \phi$  (the latter two being azimuthal and equatorial angles of the ejecta mass). We assume that the preferential axis is sampled uniformly across the  $4\pi$  steradians of the star, and that our only two free parameters are  $\alpha$  and  $\beta$ . It is pertinent to note two things due to this assumption: firstly, we are not assuming that the ejection occurs along one axis and that the kick is aligned along a different, random, axis. Rather, we are assuming that the asymmetric ejection occurs along a randomly sampled axis and that the kick is antialigned to that. The second note, explored further in subsection 6.1.2, is that there is considerable uncertainty around the nature of the kick and the precise mechanism that causes it to eject along a given axis. This is the reason we sample the kick direction uniformly, though we will simulate sufficiently many systems (on the order of  $\mathcal{O}(10^{12})$ ) that this will be marginalised over.

The existence of two free parameters,  $\alpha$  and  $\beta$ , mean that a choice on the values must be made. The “traditional” way to choose these parameters, as outlined in Bray & Eldridge (2016, 2018), is to refine them

based on isolated pulsar velocities. This methodology, however, has significant drawbacks. Notably, the reliance on one observation can engender biases in the result at the expense of no longer predicting accurate results for other observables. For instance, in the limit of ultra-stripped supernovae (USSNe) where the ejecta-remnant mass ratio becomes small, the fiducial kick in Bray & Eldridge (2018) would produce kick velocities that are strongly negative (indicating a strong kick aligned with the direction of motion of the ejecta), contrasted with the weak kicks we expect from these supernovae (e.g., Willcox et al., 2021). It becomes evident that a new series of empirical tests are needed, whereupon more constraints than simply pulsar velocities are tested. In this Chapter, we outline four such tests that form the basis of the work of Richards et al. (2023). These tests – using observed LVK Collaboration merger rates, period-eccentricity distributions of Binary Neutron Stars (BNSs), isolated pulsar velocities, and production of USSNe – allow us to narrow the parameter space significantly, with the promise of future refinement with more observations (such as through LVK Collaboration O4).

### 6.1.1 | Parameter space

We perform our computation on three populations, termed FINEGRID, WIDEGRID, and HOBBS, whose properties are outlined in Table 6.1. These populations simulate 1,000 kicks per system sampled. The aim of the three populations is twofold. Firstly, by generating FINEGRID and WIDEGRID, we are able to look both at fine detail in the region at which the most likely value is expected to lie and also the large-scale behaviour over wider parameter ranges.

| Population Name | $\alpha_{\text{low}}$ [km s <sup>-1</sup> ] | $\alpha_{\text{high}}$ [km s <sup>-1</sup> ] | $\beta_{\text{low}}$ [km s <sup>-1</sup> ] | $\beta_{\text{high}}$ [km s <sup>-1</sup> ] | Step Size [km s <sup>-1</sup> ] |
|-----------------|---------------------------------------------|----------------------------------------------|--------------------------------------------|---------------------------------------------|---------------------------------|
| FINEGRID        | 0                                           | 400                                          | -200                                       | 200                                         | 10                              |
| WIDEGRID        | 0                                           | 700                                          | -350                                       | 350                                         | 50                              |
| HOBBS           | -                                           | -                                            | -                                          | -                                           | -                               |

**Table 6.1:** The populations under study. HOBBS is a fiducial kick prescription as given by Hobbs et al. (2005), which is a Maxwellian distribution with distribution parameter  $\sigma = 265 \text{ km s}^{-1}$ , based on an empirical fit to the velocities of 233 pulsars. All three prescriptions simulated 1,000 kicks for each system, with the resultant kick being the mean velocity sampled, and the direction of the kick being sampled uniformly across the  $4\pi$  steradians of the orbit.


Secondly, by using Hobbs et al. (2005), we are able to create a population governed by a kick that has been well-tested in literature (see, for example, Bouffanais et al., 2021; Broekgaarden et al., 2021; Fragione, 2021; Igoshev et al., 2021; Mandel & Broekgaarden, 2022) which serves as a fiducial comparison. From a Bayesian perspective, we are essentially choosing the HOBBS distribution as our prior for the express purpose of a Bayesian test in section 6.3.

The choice of extremal values for  $\alpha$  and  $\beta$  are somewhat arbitrary. Figure 6.1 shows a wider parameter space with  $\alpha \in [0, 1000]$ ,  $\beta \in [-500, 500]$ . Regions in the upper right (bottom left) corner of the plot have quite high (low) mean kick velocities. Although both grids encroach on these regions somewhat, there is a tradeoff between exploring the grid space fully and using computational resources efficiently. Our mean kick velocities should not be too extreme; the mean for a Hobbs et al. (2005) kick is  $\sigma = 265 \text{ km s}^{-1}$ , for a Verbunt et al. (2017) is  $\sigma_1 = 75 \text{ km s}^{-1}$  and  $\sigma_2 = 316 \text{ km s}^{-1}$ , and for the observed Willcox et al. (2021) kick  $\sigma_1 \approx 30 \text{ km s}^{-1}$ ,  $\sigma_2 \approx 230 \text{ km s}^{-1}$ . These<sup>1</sup> are shown on the grid as black lines.

Whilst the exact parameters can be shifted somewhat and maintain the statistical integrity given these constraints, there are a few additional heuristic constraints we place on the regions to select. First, we wish to capture

<sup>1</sup>Due to the grid size and the random sampling, a true contour is difficult to observe due to sampling noise. The contours shown are thus linear fits to the actual contours.

low-velocity USSN kicks (as will be discussed in section 6.5), necessitating a region near  $\alpha = \beta = 0$  (where the kick would be identically zero for all ejecta-remnant mass ratios). Secondly, the fiducial work of Bray & Eldridge (2018) found  $\alpha = 100, \beta = -170$ . Although a goal was to (ideally) remove  $\beta < 0$  as a possible value (for USSN again), to impose that as a prior when the fiducial value sat within that range would have gone contrary to typical Bayesian practices. Thirdly, we expected but could not guarantee that the magnitude of each parameter would shrink with further tests, given the large uncertainty bands that Bray & Eldridge (2018) found (hence broadening our search area beyond  $|\alpha| = 100, |\beta| = 170$ ). Finally, we elected to increase the grid search region by 75 per cent for WIDEGRID as this would allow us to just reach  $\bar{v}_k = 700 \text{ km s}^{-1}$ , which is where the Cumulative Distribution Function (CDF) of the Hobbs et al. (2005); Verbunt et al. (2017); Willcox et al. (2021) distributions all (approximately) begin to level out (signifying few velocities above those), as shown in Figure 6.7.



**Figure 6.1:** A schematic of how FINEGRID and WIDEGRID are related. The mean kick velocity  $v_k$  was computed by averaging the kick velocities of 1,000 simulated kicks for each combination of  $\alpha$  and  $\beta$ , with  $m_{\text{rem}} \in [0.1, 2.5]$  and  $m_{\text{ej}} \in [0.1, m_{\text{rem}}]$  sampled uniformly. The pink cross represents the point  $\alpha = 100, \beta = -170$ , which are the fiducial kick parameters from Bray & Eldridge (2018).

### 6.1.2 | Natal Kick Prescriptions

There are conflicting accounts as to the cause of the kick (see, for example, Fryer et al., 1998; Arzoumanian et al., 2002; Bombaci & Popov, 2004; Hobbs et al., 2005; Bray & Eldridge, 2016; Verbunt et al., 2017). There are conflicting accounts as to the form and distribution of the kick too, as discussed in, for example, Blaauw (1961); Gunn & Ostriker (1970); de Loore et al. (1975); Sutantyo (1978); Hobbs et al. (2005); Baker et al. (2008); Wongwathanarat et al. (2013); Bray & Eldridge (2018); Giacobbo & Mapelli (2020); Mandel & Müller (2020).

We stress that we are not attempting to prescribe *which* prescription is somehow “best”, but seek an easy-to-configure kick which allows us to configure kicks without changing any of the physics of how our population was formed. This can be done by modifying the kick of Bray & Eldridge (2016), which takes two free parameters which can be changed without imposing constraints on physics. This prescription is hereinafter referred to as the “Bray kick”. Similarly, we require for the purposes of computing log-Likelihoods in section 6.3 a comparison kick, for which we select the kick of Hobbs et al. (2005), a well-known empirically derived kick. This prescription is hereinafter referred to as the “Hobbs kick”.

The Bray kick velocity is prescribed by Equation 6.1. The coefficient to the  $\beta$  term,  $\frac{M_{\text{NS}}}{M_{\text{rem}}}$ , serves to generalise the kick from purely NSs to BHs as well. When applied to a BH,  $M_{\text{NS}}$  is taken to be  $1.4 M_{\odot}$ , and conversely when applied to a NS,  $M_{\text{NS}}$  is by definition the remnant mass, so the kick velocity becomes

$$v_k = \alpha \left( \frac{M_{\text{ej}}}{M_{\text{rem}}} \right) + \beta. \quad (6.2)$$

On the other hand, the Hobbs kick velocity is given by the Probability Density Function (PDF)

$$P(v_k) dv_k = \sqrt{\frac{2}{\pi}} \frac{v_k^2}{\sigma^3} \exp \left( -\frac{v_k^2}{2\sigma^2} \right) dv_k, \quad (6.3)$$

where  $\sigma = 265 \text{ km s}^{-1}$  is the distribution parameter found by fit in Hobbs et al. (2005) to the velocities of 233 pulsars.

### 6.1.3 | The TAKAHE orbital evolution code

As part of this work, we have developed the TAKAHE binary evolution code, which solves the coupled equations of motion Equation 2.51 and Equation 2.52. Originally, to facilitate the equations being solved rapidly, we wrote the code as a PYTHON wrapper to a JULIA (Bezanson et al., 2017) integrator. However, the PYJULIA library (Arakaki et al., 2023) proved to be fragile<sup>2</sup> and is not in fact supported officially on recent Python versions. To facilitate the potential interoperability between KAITIAKI and TAKAHE (and to remove a prolonged loading time when the JULIA runtime was loaded, given that PYJULIA precompiles our modules), we elected to rewrite the core integrator into PYTHON and to optimise it using traditional PYTHON methods instead.

There are two relevant functions in TAKAHE. The remaining functions are utility functions that allow us to (for instance) convert between lookback time / redshift, and period / separation. These functions are `integrate_eoms()` (Code Sample A.1), which solves the coupled differential equations for  $\frac{da}{dt}$  and  $\frac{de}{dt}$  (Equation 2.51 and Equation 2.52), and `integrate_timescale()` (Code Sample A.2), which solves Equation 2.55 for the coalescence time of the binary. With sufficient performance optimisations,<sup>3</sup> `integrate_timescale()` is able to achieve speeds of approximately 18,000 numerical integrations per second on an office PC<sup>4</sup> on a sample of Neutron Star - Black Hole Binary (NS-BH) binaries in the LISA dataset (Tang, priv. comm.).

<sup>2</sup>It requires custom installation procedures (so cannot be included as a dependency in e.g. a PYTHON package), would hang with no discernible output or ability to ascertain why, and could not handle KeyboardInterrupt exceptions in PYTHON

<sup>3</sup>Primarily, just-in-time compilation with the NUMBA (Lam et al., 2015) library.

<sup>4</sup>A Dell OptiPlex 7480 AIO with a 12-core 3.1 GHz Intel Core i5-10500 CPU, running no other programs except for during the final few iterations

Algorithmically, both codes employ a simple Euler integration routine (where  $f_{n+1} = f_n + \frac{df_n}{dt} dt$ ) and the choices work as follows:

1. When `integrate_eoms()` takes a step in time, it evaluates  $da$  and  $de$ . It then computes the relative changes in both  $e$  and  $a$  that arise by taking this timestep (through  $\delta f = \left| 0.01 \frac{f}{f} \right|$ ), and takes a timestep corresponding to the smallest change between  $a$  and  $e$ , or half of the width of the smallest BPASS time bin.
2. `integrate_timescale()` breaks up Equation 2.55 into four cases dependent on the initial eccentricity. Let  $\tau$  be the coalescence time of a circular binary ( $\tau = \frac{a_0^4}{4\beta}$ ), then the cases are:

$$t_c = \begin{cases} \tau & e_0 = 0 \\ \frac{c_0^4}{4\beta} e_0^{\frac{48}{19}} & e_0 < 0.01 \\ \frac{12}{19} \frac{c_0^4}{\beta} \int_0^{e_0} \frac{e^{\frac{22}{19}} \left(1 + \frac{121}{304} e^2\right)^{\frac{1181}{2299}}}{(1-e^2)^{\frac{3}{2}}} de & 0.01 \leq e_0 \leq 0.99 \\ \tau \frac{768}{425} (1 - e_0^2)^{\frac{7}{2}} & 0.99 < e_0 \end{cases}, \quad (6.4)$$

where all symbols have the same meaning as in section 2.5. The integral in the intermediate case is evaluated numerically, taking a fixed timestep of  $de = \frac{e_0}{1000}$  such that the discretised sum becomes nothing more than


$$t_c (0.01 \leq e_0 \leq 0.99) = \frac{e_0}{1000} \frac{12}{19} \frac{c_0^4}{\beta} \sum_{e=0}^{e_0} \frac{e^{\frac{22}{19}} \left(1 + \frac{121}{304} e^2\right)^{\frac{1181}{2299}}}{(1 - e^2)^{\frac{3}{2}}}, \quad (6.5)$$

and dependence of the coalescence time  $t_c$  on the initial separation  $a_0$  is handled through the  $c_0$  term. Since the step size is an arbitrary choice, a tradeoff can be made between accuracy and runtime. Figure 6.2 shows the effect of changing the step size ( $\frac{e_0}{1000}$  in Equation 6.5) on convergence to a final result, and Figure 6.3 shows the runtime as a function of step size on an office computer.

## 6.2 | Merger Rates of BNS systems

The first test is to match the observed merger rates of BNS systems. To do this, we use the LVK Collaboration catalogue of results from Operating Run 3 (O3). The simplest test is to match the merger rate today, which we denote as  $\mathcal{R}_0$ , to that observed by LVK Collaboration (LIGO-Virgo, 2021; LIGO-Virgo-KAGRA, 2021). We display this in Figure 6.4 alongside the previously reported fiducial values of  $\alpha = 100 \text{ km s}^{-1}$  and  $\beta = -170 \text{ km s}^{-1}$ , the theoretical merger rate calculation performed by Eldridge et al. (2019), and the observed merger rates from LIGO-Virgo (2021) and LIGO-Virgo-KAGRA (2021). We term the contour defined by LIGO-Virgo (2021) (in black) the “LIGO contour”.

Selecting the value to calibrate to is not a straightforward task. Although the LIGO-Virgo (2021) contour is superseded by the LIGO-Virgo-KAGRA (2021) contour, the separation into different inference models means that the merger rates for BNS systems have intractably large error bars covering the entire plane; for instance, their MULTISOURCE (MS) model alone spans  $2.11 \leq \log \mathcal{R}_0 \leq 3.23$  which covers FINEGRID entirely and almost entirely covers WIDEGRID. In addition to the lack of constraint, we would be imposing additional parameters into the model insofar as the selection of model – whether, for example, PDB was taken as truth over BGP, MS, or one of the other population models. Whilst an interesting and assuredly insightful endeavour, the aim of this work is instead to develop a simple test that can be used in future on extra populations such as these. A simple criterion which can be extended is valuable, and future work may extend these tests in multiple directions.



**Figure 6.2:** The effect of varying  $de$  in Equation 6.5 on the coalescence time. Although hard to see, the pink line does oscillate above and below the brown line. This binary was assumed to have  $m_1 = m_2 = 1.44 M_\odot$  and  $a_0 = 1970.5 R_\odot$  (the mean separation of a sample of binaries with  $\alpha = 100 \text{ km s}^{-1}$ ,  $\beta = -170 \text{ km s}^{-1}$ ).

### 6.3 | Period-Eccentricity Distributions


Since in the absence of any mass transfer the motion of two compact binaries is wholly dependent on their initial separation and orbital eccentricity, they trace out paths in a two-dimensional phase space. In general, since binaries circularise over time, and by the emission of gravitational waves their orbits shrink, a binary will start in the high-period, high-eccentricity region of period-eccentricity space, and evolve towards the low-period, low-eccentricity region. As a result, given a population of binary stars, we can design a density plot, where the density is effectively the likelihood of a binary selected at random having the parameters we sample.

Computationally, for each binary in our sample, we trace its path through period-eccentricity space, and at every timestep we add  $w \times dt$  to the density plot, where  $dt$  is the timestep, and  $w$  is the weight term output from the TUI code proportional to the IMF weight (where the constant of proportionality is related to the uniformly sampled kick direction). The result of this computation, for the parameter values that lie along the LIGO contour, are shown in Appendix C. We also show the same result in Figure 6.5. Given a catalogue of  $N$  (observed) binary stars, we can then compute the likelihood of the model as


$$\ln \mathcal{L} = \sum_{n=0}^N \ln p(\ln P_n, e_n), \quad (6.6)$$

where  $p(\ln P_n, e_n)$  is the content of the density plot at the location of the  $n^{\text{th}}$  observed BNS system. Then, the Bayes factor (defined as the likelihood ratio) can be computed as


$$\mathcal{K} \equiv \frac{\mathcal{L}}{\mathcal{L}_{\text{Hobbs}}} \implies \ln \mathcal{K} = \ln \mathcal{L} - \ln \mathcal{L}_{\text{Hobbs}}, \quad (6.7)$$



**Figure 6.3:** The effect of varying  $d_e$  in Equation 6.5 on the runtime, which is demonstrably  $\mathcal{O}(n)$ , and the performance of the algorithm. The input data to this algorithm were the LISA binary sample from Tang (priv. comm.). The performance is shown in units of thousands of iterations per second.



**Figure 6.4:** The merger rate as a function of kick parameters  $\alpha$  and  $\beta$ . The cyan contour is the theoretical merger rate as reported by Eldridge et al. (2019). The black contour is the merger rate observed in LIGO-Virgo (2021). The remaining contours are the merger rates reported in LIGO-Virgo-KAGRA (2021) (pink: BGP, yellow: MS, orange: PDB [ind], see LIGO-Virgo-KAGRA (2021) for an explanation of each model). Shaded regions represent uncertainties, and uncertainties are not reported for the LIGO-Virgo-KAGRA (2021) results as they would cover the entire plane. The blue cross is the fiducial value of  $\alpha$  and  $\beta$  from Bray & Eldridge (2018). This value lies off the previous work’s contours as the rejuvenation age calculation has been improved in TUI in preparation for the next major BPASS release.




**Figure 6.5:** The period-eccentricity distributions for six selected Bray & Eldridge (2016) kicks. Shown are the kicks for  $\alpha = 180 \text{ km s}^{-1}, \beta = 10 \text{ km s}^{-1}$  as the system with the highest  $\ln \mathcal{K}$  across the LIGO contour,  $\alpha = 170 \text{ km s}^{-1}, \beta = -10 \text{ km s}^{-1}$  as the lowest, and the balance to illustrate structural differences. White regions show regions where no systems evolve to. The white line represents a constant-time coalescence time of the Hubble time ( $\sim 13.8$  Gyr), computed using Mandel (2021). The cyan diamonds are the observed systems enumerated in Table 6.2. In white text are the values of  $(\alpha, \beta)$ , and the  $\ln \mathcal{K}$  values for each plot.

| Name       | $\log(P/\text{days})$ | $e$   | Original Reference     | Name       | $\log(P/\text{days})$ | $e$   | Original Reference     |
|------------|-----------------------|-------|------------------------|------------|-----------------------|-------|------------------------|
| J0453+1559 | 0.609808              | 0.113 | Martinez et al. (2015) | J0737-3039 | -0.991400             | 0.088 | Kramer et al. (2006)   |
| B1534+12   | -0.375718             | 0.274 | Fonseca et al. (2014)  | J1756-2251 | -0.494850             | 0.181 | Faulkner et al. (2005) |
| B1913+16   | -0.490797             | 0.617 | Hulse & Taylor (1975)  | J1913+1102 | -0.686133             | 0.090 | Lazarus et al. (2016)  |
| J1757-1854 | -0.735182             | 0.606 | Cameron et al. (2018)  | J1518+4904 | 0.936212              | 0.249 | Janssen et al. (2008)  |
| J1811-1736 | 1.273672              | 0.828 | Corongiu et al. (2007) | J1829+2456 | 0.070407              | 0.139 | Champion et al. (2004) |
| J1930-1852 | 1.653791              | 0.399 | Swiggum et al. (2015)  | J1753-2240 | 1.134751              | 0.304 | Keith et al. (2009)    |
| J1411+2551 | 0.417638              | 0.169 | Martinez et al. (2017) | J1946+2052 | -1.107905             | 0.064 | Stovall et al. (2018)  |

**Table 6.2:** The systems we use in our study. These data were originally compiled in Vigna-Gómez et al. (2018).

where  $\mathcal{L}_{\text{Hobbs}}$  is the log-Likelihood of the Hobbs kick. We use a catalogue of fourteen BNS systems originally compiled in Vigna-Gómez et al. (2018), shown in Table 6.2. From this catalogue, we sample the period-eccentricity distribution at that space and take that as the probability of observing that binary, thus allowing us to use the content of the period-eccentricity distribution as the  $\ln p$  term in our summand of Equation 6.6.

By doing this we effectively construct a Bayes-factor map, where  $\ln \mathcal{K}$  can be shown evaluated at every combination of  $\alpha$  and  $\beta$ . We term this value  $\ln \mathcal{K}_{\alpha\beta}$ . This is shown in Figure 6.6 for both of our considered grids. In general we see that approximately 60% of FINEGRID has a positive log-Bayes factor (indicating any level of support of the model compared to Hobbs), whereas approximately 40% of WIDEGRID has a positive log-Bayes factor. In a Bayesian analysis, the constraint that  $\ln \mathcal{K} \geq 3.4$  indicates strong support for the model contrasted against the reference (i.e., Hobbs et al., 2005), which contains approximately 9.1% of FINEGRID and 4.4% of WIDEGRID.



**Figure 6.6:** The  $\ln \mathcal{K}$  values across  $\alpha - \beta$  space. The orange line demarcates  $\ln \mathcal{K} = 3.4$ , which in a Bayesian test represents conclusive evidence in favour of the distribution compared to the Hobbs et al. (2005) kick.

## 6.4 | Single-Star Kick Velocities

The traditional methodology for refining a kick prescription is to consider the velocity distribution of isolated pulsars. Thus, given a suitable observational dataset, we would be able to compute the expected kick distribution and compare it, via a Kolmogorov-Smirnov (K-S) test, to the observed data set. The K-S statistic between distributions  $\mathcal{F}$  (the observed distribution) and  $\mathcal{G}$  (the compared distribution) is given by

$$\text{KS} = \max |\text{cdf}(\mathcal{F}) - \text{cdf}(\mathcal{G})|, \quad (6.8)$$

where cdf means the CDF function, defined as the integral of the PDF. Given the CDF is bounded between 0 and 1 inclusive, the K-S statistic is also bounded between this region. A K-S statistic of 1 means that the compared distribution diverges perfectly from the observed distribution, whereas a K-S statistic of 0 means it matches perfectly. This test of divergence was the refinement performed in Bray & Eldridge (2016, 2018), hence we select this (with an independently generated dataset to those works) to be consistent with prior work.

### 6.4.1 | Pulsar velocity dataset

We use the same dataset of 81 pulsars from Willcox et al. (2021), and filter the dataset in the same way that they do. We remove millisecond pulsars, pulsars in globular clusters, and pulsars known to be in binaries, in order to homogenise the dataset. The PDF and CDF of Willcox et al. (2021), as well as those generated from Hobbs et al. (2005) and Bray & Eldridge (2018), is displayed in Figure 6.7.


This homogenised dataset gives the transverse velocities of each pulsar alongside the RA/Dec, the proper motion in the RA/Dec directions, and parallax. We assume that the transverse velocity is approximately the original kick velocity. As mentioned earlier in this chapter, old remnants with high velocities may have moved substantially – a 20 Myr remnant with a transverse velocity of  $500 \text{ km s}^{-1}$  will have moved  $\sim 10 \text{ kpc}$ , absent external forces (Willcox et al., 2021). However, this is likely only to be an issue for pulsars whose transverse velocity substantially exceeds the transverse Galactic rotational velocity of the Sun  $v_{\text{TGRV}} \approx 230 \text{ km s}^{-1}$  (Bhattacharjee et al., 2014), shown in yellow on Figure 6.7. Although the dataset of Willcox et al. (2021) has pulsars with velocities higher than this value – peaking at  $\sim 850 \text{ km s}^{-1}$  – this is not a grave concern for this work, for two reasons. Firstly, the kick velocities output from TUI are taken immediately after the supernova. Secondly, the analysis Willcox et al. (2021) performs is focused on low-velocity pulsars, and this test limits the kick velocity of an USSNe to  $30 \text{ km s}^{-1}$ , well below  $v_{\text{TGRV}}$ .

The Hobbs et al. (2005) kick has a closed-form expression for the PDF  $P(v_k)$ , given by Equation 6.3. Consequently, it has a smooth curve in Figure 6.7, whereas the Bray & Eldridge (2018) and Willcox et al. (2021) PDFs are presented as histograms as they are computed.


### 6.4.2 | Comparison

Figure 6.8 shows the K-S statistic as a function of  $\alpha$  and  $\beta$  across our two parameter spaces. We also show the K-S statistic of the Hobbs kick, which has a value of 0.34. We find in general a valley of models with good agreement to Willcox et al. (2021), and find that the global minimum K-S statistic is at  $\alpha = 100, \beta = -180$  – agreeing with the prior work of Bray & Eldridge (2018) (who find  $\alpha = 100, \beta = -170$ ).

We also show in Figure 6.9 a selection of four PDFs, alongside their associated CDF and the CDFs for Willcox et al. (2021) and the Hobbs kick. We note that the numerical values shown in Figure 6.9 differs somewhat from those shown in Richards et al. (2023). This was due to a minor bug in the code incorrectly labelling the KS statistics for the figure. The actual result of the paper does not change with the correction of the bug, however the values reported in Figure 5 of Richards et al. (2023) do change.



**Figure 6.7:** The PDF and CDF of the distributions using kicks from Hobbs et al. (2005); Verbunt et al. (2017); Bray & Eldridge (2018); Willcox et al. (2021). Hobbs et al. (2005) has a closed-form expression, whereas the others are computed empirically, hence the Bray and Willcox CDFs are less smooth than that of Hobbs. The two dashed lines indicate where and how the K-S statistic is measured for the associated distribution. The yellow region indicates the region for which  $v_{\text{kick}} \geq v_{\text{TGRV}}$ , the transverse Galactic rotation velocity of the Sun,  $\sim 230 \text{ km s}^{-1}$ .



**Figure 6.8:** The K-S statistic for FINEGRID and WIDEGRID. The orange line denotes the K-S statistic for the Hobbs kick, which has a value of 0.34.

The four values we present on the left side of Figure 6.9 correspond to  $\alpha = 100 \text{ km s}^{-1}, \beta = -170 \text{ km s}^{-1}$  for the fiducial Bray kick,  $\alpha = 180 \text{ km s}^{-1}, \beta = 10 \text{ km s}^{-1}$  to represent the system with the highest  $\ln \mathcal{K}$  value from section 6.3,  $\alpha = 170 \text{ km s}^{-1}, \beta = -10 \text{ km s}^{-1}$  the lowest, and  $\alpha = 40 \text{ km s}^{-1}, \beta = 0 \text{ km s}^{-1}$  the system with low  $\alpha$  and zero  $\beta$ , corresponding to a kick that would *a priori* generate logical velocities in the USSNe limit.


## 6.5 | Production of Ultra-Stripped Supernovae

As discussed earlier, the fiducial values of the Bray kick from Bray & Eldridge (2018) admit some problematic extremal cases. If  $m_{\text{ej}} \rightarrow 0$ , as is the case in a USSN, the kick velocity would become negative. For instance, an “average” USSN with  $m_{\text{ej}} = 0.15 M_{\odot}$  and  $m_{\text{rem}} = 1.45 M_{\odot}$  would have a kick velocity of  $\sim -159 \text{ km s}^{-1}$ . The concern here is twofold. Firstly, a negative kick velocity would mean that the velocity vector of the kick would be antialigned with the direction of the more massive remnant; this is a violation of conservation of momentum. Secondly, a kick with high magnitude means that, irrespective of the alignment of the kick, the kick itself is strong. As discussed in Willcox et al. (2021), USSNe are associated with weak kicks, peaking at  $v_k \sim 30 \text{ km s}^{-1}$ . This means that the fiducial Bray kick is unable to correctly account for USSNe.

To account for these supernova kicks, we introduce a new constraint onto our parameters. Following Tauris et al. (2015), we define a USSN as follows:

1. The ejecta mass is in the range  $0 \leq m_{\text{ej}} \leq 0.3 M_{\odot}$ , admitting the USSN SN2019dge from Yao et al. (2020),
2. The remnant mass is in the range  $1.1 \leq m_{\text{rem}} \leq 1.8 M_{\odot}$ ,
3. The envelope mass  $m_{\text{env}} \leq 0.2 M_{\odot}$ .

For every combination  $\alpha, \beta$ , we record whether it is capable of producing one USSN. The constraint is that this heuristic is met. This constraint is extremely conservative, insofar as we do not constrain on the prevalence



**Figure 6.9:** The PDFs and CDFs of four systems, alongside the Hobbs CDF and Willcox et al. (2021) CDF. The dashed line on the PDF plots is the PDF of the observed distribution from Willcox et al. (2021). On the right hand plot, the number in parentheses represents the K-S statistic for the distribution, measured relative to Willcox et al. (2021), which by definition has no K-S statistic.

of USSNe in our sample, but rather by whether or not they are possible at all. The reason for the conservatism here is that the rates of Type Ic USSNe are expected to be very low. In the BPASS dataset, they comprise 1.8 per cent of all CCSNe (Stevance, 2022). The paucity of data alongside the paucity of observational evidence suggests a low rate, but since the prevalence is unknown, we place the conservative constraint of at least one USSN to be agnostic. This effectively widens our error bars such that when the USSNe rate is refined further, it will shrink the parameter space.

## 6.6 | Refinement of the parameter space

We are now in a position to impose constraints on our parameter space based on the analysis provided above. We apply the following four constraints:


1. **DNS Merger Rate** (section 6.2): Does the merger rate produced by these parameters sit along the LIGO contour?
2. **DNS Period-Eccentricity Distributions** (section 6.3): Is  $\ln \mathcal{K}_{\alpha\beta} \geq 3.4$  compared to the Hobbs et al. (2005) kick (representing conclusive evidence in favour of the distribution)?
3. **Isolated Pulsar Velocities** (section 6.4): Is the K-S statistic produced by this choice of parameters, in comparison to the Willcox et al. (2021) kick, less than 0.5 (indicating it agrees with observational data more than it disagrees)?
4. **Logical USSN velocities** (section 6.5): Is this choice of parameters able to produce at least one USSN?

These can be overlaid onto our parameter space, which is shown in Figure 6.10. The center of mass of our fit region is located at  $\alpha = 117_{-51}^{+36} \text{ km s}^{-1}$ ,  $\beta = 13_{-14}^{+8} \text{ km s}^{-1}$ , however this value is computed on a finer grid than the parameter space and is thus likely to be an overfit to the data. We thus report the final recommended model value to the nearest half-bin-width of the parameter space, broadening error bars where necessary, and find  $\alpha = 115_{-55}^{+40} \text{ km s}^{-1}$ ,  $\beta = 15_{-15}^{+10} \text{ km s}^{-1}$  as the recommended model value. Error bars are always broadened here; traditional rounding would result in the upper bound on  $\alpha$  (being  $153 \text{ km s}^{-1}$ ) becoming  $150 \text{ km s}^{-1}$ , which artificially removes potentially permissible regions of the parameter space. Broadening the bars removes this at the cost of potentially admitting values that do not belong within the “acceptable” zone. Further constraints, however, will shrink the region, so this is an acceptable tradeoff.


In Figure 6.11, we present the two major inferred properties of our recommended model value. First, we present the CDFs of a population governed by a Bray kick with the recommended parameters, contrasted with the associated Hobbs kick and observed Willcox et al. (2021) data. We also show the period-eccentricity distribution of the population, with the observed systems from Table 6.2 overlaid atop the distribution.

We are also able to extrapolate this to comment on the expected peak of the event-rate distribution. Given that the event rate is a convolution between the merger rate of our population and the SFRD we choose (cf. Equation 2.60), by affixing the SFRD across our populations, by Equation 2.56 we can vary  $\mathcal{R}(t)$  by varying  $\alpha$  and  $\beta$ . Since the SFRD is of a Gaussian form (with a peak at  $z \sim 1.863$ ), we can also extract the peak of the distribution assuming it is governed by a particular  $\alpha$ - $\beta$  combination. The general results across the parameter space are shown in Figure 6.12. The best model choice parameters we find fall within the valley of  $z = 0$ , indicating that the peak of the event rate distribution is the present day.


We note that there is a single peak at  $\alpha = 330 \text{ km s}^{-1}$ ,  $\beta = -70 \text{ km s}^{-1}$  which appears to be an outlier when compared to its neighbours. We display in Figure 6.13 the local behaviour of the 9 cells around this region. One can see that this beak is bimodal with a second peak at  $z = 0$ , which is only 0.098 per cent lower than the reported peak, clearly resulting in a simple outlier.




**Figure 6.10:** The four constraints we apply overlaid on our FINEGRID parameter space. The cyan region represents the region which fits all four parameters, and the blue cross represents the “center of mass” of this region.



**Figure 6.11:** The CDF for a population of isolated pulsars (left) and the period-eccentricity distribution (right) for  $\alpha = 115 \text{ km s}^{-1}$ ,  $\beta = 15 \text{ km s}^{-1}$ . The blue region on the CDF plot shows the uncertainty region for the reported value. The red region shows the uncertainty band on the Willcox et al. (2021) CDF, which was computed by sampling the PDF 10,000 times, taking 2,000 samples each time, and plotting the extent of the simulated distribution. The black band is the uncertainty on the Hobbs distribution, taking  $\sigma_{\text{max}} = 291 \text{ km s}^{-1}$  and  $\sigma_{\text{min}} = 239 \text{ km s}^{-1}$  from Verbut et al. (2017).



**Figure 6.12:** The peak of the event rate  $\mathcal{R}(t)$  as a function of  $\alpha$  and  $\beta$ . Varying these two parameters generally brings the peak closer to the present day ( $z = 0$ ). Absent a merger rate (or, more formally, assuming the merger rate is instantaneous and given by a Dirac-delta function), the SFRD peaks at  $z \sim 1.863$ . The region surrounding the peak at  $\alpha = 330$ ,  $\beta = -70$  is shown in Figure 6.13.



**Figure 6.13:** The event rate  $\mathcal{R}(z)$  for a region surrounding  $\alpha = 330, \beta = -70$  on FINEGRID. This region corresponds to the “outlier” in Figure 6.12. Shown on the left are the outlier bin in the center and its neighbours in both  $\alpha$ -space and  $\beta$ -space. The black line on each plot denotes the location in redshift of the event rate distribution peak. The right plot shows the outlier in black and its neighbours in faded red behind.

Unfortunately, there is a paucity of observational evidence to confirm whether or not the peak is truly the present day. Given that a redshift of  $z = 2$  corresponds to a lookback time of  $t_{\text{LB}} = 10$  Gyr, we lack evidence to corroborate this prediction. The next generation of telescopes, including JWST, may yet be able to constrain this result further. In the interim, indirect evidence may be able to be applied via the r-process enrichment history of the Milky Way. Van de Voort et al. (2022), for example, finds that greater velocity natal kicks would be correlated with fewer r-process enhanced stars in the Milky Way, as higher kicks would unbind more binaries, meaning that r-process elements are no longer able to accrete onto star-forming regions. Kobayashi et al. (2023) indicates that current population synthesis models<sup>5</sup> are unable to reproduce the abundance distributions of certain high-mass isotopes, including the abundances of Oxygen, Iron, Europium, Gold, and Uranium.

## 6.7 | Discussion

There are limitations to this analysis. The first is that we did not preferentially weight our binaries. Tauris et al. (2017) for example places particular weight on PSR J1930-1852, which is a particularly wide binary. Per Swiggum et al. (2015) (who reported it), “its unique spin and orbital parameters challenge models that describe BNS formation”, which implies that any credible model of BNS formation should replicate its orbital parameters. We did find in Figure C.1 that some parameters had non-zero probability of replicating this pulsar, which indicates that the Bray kick is able to reproduce it.

Secondly, we have assumed Newtonian mechanics. Formally, the derivations presented earlier (cf. subsection 2.4.1) assumed general relativity for the existence of the gravitational wave equation,<sup>6</sup>

$$\square \bar{h}_{\mu\nu} = -\frac{1}{2}\sqrt{32\pi G}T_{\mu\nu}, \quad (6.9)$$

but the injected physics was Newtonian (see Richards (2021) for a derivation). Strictly speaking, relativistic corrections apply to these kinematic equations too. The logic outlined in subsection 2.2.3 ceases to apply, as we are not modelling the interior of a star. Instead, one would need to apply corrections to the two-body equation of motion, termed the post-Newtonian approximation. The general equation becomes

$$\frac{d\vec{v}}{dt} = \frac{G(m_1 + m_2)}{r^2} \left( -\hat{n} + \sum_{k=2}^N \frac{1}{c^k} \vec{A}_{\frac{k}{2}\text{PN}} \right), \quad (6.10)$$

where  $N$  is the highest order term considered, and  $\vec{A}_{\frac{k}{2}\text{PN}}$  is termed the “ $\frac{k}{2}$ -th Post-Newtonian” coefficient. Testing with the RIRORIRO code from van Zeist et al. (2021) indicated that including terms up to  $N = 6$  (3PN) increases the merger time by fractions of a year, and thus does not significantly alter the merger rate density we observe. Typically 3.5PN ( $N = 7$ ) is used as an upper bound (Itoh & Futamase, 2003), though up to 10PN ( $N = 20$ ) has been used before (Blanchet, 2014, and references therein).

Relativistic corrections also typically manifest in using more sophisticated spacetime metrics. Equation 6.9 assumes a metric which is locally Minkowski, that is

$$g_{\mu\nu} = \eta_{\mu\nu} + \sqrt{32\pi G}h_{\mu\nu}, \quad (6.11)$$

where  $\eta_{\mu\nu}$  is the positive-definite Minkowski metric. This “weak-field” approximation breaks down when a compact object enters within its companion’s innermost stable circular orbit (ISCO), given by  $r_{\text{isco}} = \frac{6GM}{c^2}$ , where matter can be exchanged. Given gravitational waves result in strains on the order of  $10^{-19}$  (van Zeist et al., 2023),

<sup>5</sup>Explicitly considered in the paper are COMPAS (Team COMPAS: Riley et al., 2022), StarTrack (Belczynski et al., 2020), Brussels (Mennekens & Vanbeveren, 2014, 2016), ComBinE (Kruckow et al., 2018), and BPASS (Eldridge et al., 2017).

<sup>6</sup>Here,  $\bar{h}_{\mu\nu}$  is the trace inverse of a weak perturbation to locally Minkowski spacetime with positive-definite trace, and  $\square$  is the d’Alembertian operator.

representing observed fractional shifts on Earth smaller than the diameter of a proton, (Moore et al., 2014), the weak field regime suits, and again modelling the relativistic corrections to the post-ISCO regime only increases the merger rate by fractions of years.

To contextualise this case study, other authors such as Mapelli & Giacobbo (2018); Wiktorowicz et al. (2019); Giacobbo & Mapelli (2020); Santoliquido et al. (2021); Kapil et al. (2023) have explored the parameter space of BNS kicks, though none have performed the analysis in the way we have. These works typically vary the progenitor population properties, in such a way that the formed binaries can in principle vary strongly. However, in our population synthesis procedure, we are able to effectively affix the physics between different models such that *only* the kick velocities are expected to vary between individual populations. Since the Bray kick has two parameters which can be altered, we are able to modify the kick velocity without changing the physics, thus allowing us to isolate and study the kick mechanism alone.

## 6.8 | Chapter Summary

We have presented a formalism for evaluating and quantifying the goodness-of-fit of the natal kick mechanism against four independent tests. The tests we have presented – constraining the merger rate of BNS systems, the period-eccentricity distribution of Galactic BNS systems, the kick velocities of single-star pulsars, and the production of ultra-stripped supernovae – greatly constrain the parameter space to be less than 1 per cent of the possible parameters. Furthermore, the data suggest that the peak of the merger rate distribution is the present day, with  $z \simeq 1.5$  as the upper bound of the entire parameter space, and  $z = 0$  for our recommended value.

## CHAPTER 7

---

# Conclusion

*Oh! I have slipped the surly bonds of Earth  
And danced the skies on laughter-silvered wings;*

...

*And, while with silent lifting mind I've trod  
The high untrespassed sanctity of space,  
Put out my hand, and touched the face of God.*

---

*High Flight*, BY JOHN MAGEE JR.

*Take-off is optional. Landing is mandatory.*

---

FIRST RULE OF AVIATION

## 7.1 | Summary

## 7.2 | Limitations

*Though in reviewing the incidents of my administration I am  
unconscious of intentional error, I am nevertheless too sensible of  
my defects not to think it probable that I may have committed many  
errors.*

---

GEORGE WASHINGTON

1. circular orbits
2. o-type stars
3. parameter selection

## 7.3 | Suggested Future Work



---

# Appendices



# Code Samples

Code Sample A.1: The TAKAHE function `integrate_eoms()`

```

1 def integrate_eoms(a0, e0, p):
2     """
3     General purpose integrator for Nyadzani & Razzaque eqns for a & e.
4
5     Params:
6         a0 - The initial semimajor axis, measured in solar radii
7         e0 - The initial eccentricity, dimensionless
8         p - A vector of parameters:
9             p[1] = m1 (units: Solar Mass)
10            p[2] = m2 (units: Solar Mass)
11            p[3] = unused (will be removed in next version)
12            p[4] = unused (will be removed in next version)
13            p[5] = Lifetime (evolution + rejuvenation)
14
15     Returns:
16         A - An array of the semimajor axes of the binary system over time. (Solar
17         ↪ Mass)
18         E - An array of the eccentricities of the binary system over time. (no dim.)
19     """
20
21     Solar_Mass = 1.989e30      # kg
22     Solar_Radius = 696340000.0 # m
23     G = 6.67e-11              # m^3 kg^-1 s^-2
24     c = 299792458.0           # m/s
25
26     A = np.array([])
27     E = np.array([])
28     H = np.array([])
29
30     m1, m2, _, _, evotime = p[0], p[1], p[2], p[3], p[4]
31
32     # number of seconds in a year
33     seconds_per_year = 60 * 60 * 24 * 365.25
34
35     #####
36     #      Unit Check      #
37     #####
38     a = a0 * Solar_Radius  # Meters
39     e = e0                  # Dimensionless
40     m1 = m1 * Solar_Mass    # Kilogram
41     m2 = m2 * Solar_Mass    # Kilogram
42     #####
43     #      End Unit Check      #
44     #####
45
46     # Beta has units m^4 / s
47     beta = ((64/5) * G**3 * m1 * m2 * (m1 + m2) / (c**5))

```

```

48     A = np.append(A, a)
49     E = np.append(E, e)
50     H = np.append(H, 0.0)
51
52     total_time = 0
53
54     # Integrate until past the end of the universe, or a 10km orbit
55     while total_time/seconds_per_year + evotime < 1e11 and a > 1e4:
56         # an euler integrator: work out da/dt then times it by dt
57         # to get da, which then we can work out as a = a + da/dt * dt.
58         initial_da = (- beta / ((a**3) * (1 - e**2)**(7/2)))
59         da = initial_da * (1 + (73/24) * e**2 + (37/96) * e**4)
60
61         intial_de = (((-19/12) * beta) / (a**4*(1-e**2)**(5/2)))
62         de = intial_de * (e + (121/304) * e**3)
63         # Units: s^-1
64
65         timeA = abs(1e-2 * a/da)
66
67         if e > 1e-10:
68             timeE = abs(1e-2 * e/de)
69         else:
70             de = 0
71             e = 1e-10
72             timeE = timeA * 10
73
74         # maximum timestep is half of the width of the smallest BPASS time bin
75         conv_frac = 0.23076752*0.5*seconds_per_year
76
77         dt2 = (evotime + total_time/seconds_per_year)*conv_frac
78
79         # Take a timestep that results in the smallest change: either a
80         # change in E, a change in A, 1/2 the smallest BPASS bin.
81         dt = min(timeE, timeA, dt2)
82
83         a = a + dt * da
84         e = e + dt * de
85
86         A = np.append(A, a)
87         E = np.append(E, e)
88         H = np.append(H, dt)
89
90         total_time = total_time + dt
91
92         # Determine why we stopped, so the parent function can tell what went
93         # "wrong".
94         stop_reason = "flag_not_set"
95
96         if total_time/seconds_per_year + evotime >= 1e11:
97             stop_reason = "out_of_time"
98
99         if a <= 1e4:
100             stop_reason = "merged"
101
102         # Solar Radii, Dimensionless
103         return A / Solar_Radius, E, H, stop_reason

```

Code Sample A.2: The TAKAHE function `integrate_timescale()`

```

1 def integrate_timescale(m1, m2, p0, e0):
2     """
3     General purpose integrator for eqn(5.14) of Peters, 1964 [1].
4
5     [1] https://ui.adsabs.harvard.edu/abs/1964PhRv..136.1224P/abstract
6
7     Params:
8         m1 -- The mass of the first star (Solar Mass)
9         m2 -- The mass of the second star (Solar Mass)
10        p0 -- The initial period (days)
11        e0 -- The initial eccentricity
12
13     Returns:
14         tC -- The coalescence time (seconds)
15     """
16     Solar_Mass = 1.989e30 # kg
17     Solar_Radius = 696340000.0 # m
18     G = 6.67e-11 # m^3 kg^-1 s^-2
19     c = 299792458.0 # m/s
20
21     m1 = float(m1)
22     m2 = float(m2)
23     p0 = float(p0)
24     e0 = float(e0)
25
26     #####
27     # BEGIN UNIT CHECK #
28     #####
29     m1 = m1 * Solar_Mass # Kilograms ##
30     m2 = m2 * Solar_Mass # Kilograms ##
31     p0 = p0 * 24 * 60 * 60 # Seconds ##
32     e0 = e0 # Dimensionless ##
33     #####
34     # END UNIT CHECK #
35     #####
36
37     # Use Kepler's Third Law to compute the semimajor axis, in meters
38     a0 = (p0**2.0 * (G * (m1+m2))) / (4.0 * np.pi**2.0))**(1.0/3.0) # meters
39
40     # compute the constant c0
41     c0 = a0 * (1.0-e0**2.0) * e0**(-12.0/19.0) * (1.0+(121.0 / 304.0 *
42     ↪ e0**2.0))**(-870.0/2299.0) # meters
43
44     # Compute the constant beta
45     beta = 64.0 / 5.0 * G**3.0 * m1 * m2 * (m1 + m2) / c**5.0
46     # meters^4 / s
47
48     # Circular binary coalescence time
49     tC = a0**4.0 / (4.0*beta)
50
51     if e0 != 0:
52         if e0 < 0.01:
53             # Low ecc - see eqn after eqn(5.14) of Peters, 1964
54             tC = c0**4.0 * e0**(48.0/19.0) / (4.0*beta)

```

```
54     elif e0 > 0.99:
55         # High ecc - see eqn after eqn after eqn(5.14) of Peters, 1964.
56         tC = tC * ((768.0 / 425.0) * (1.0-e0**2.0)**3.5)
57     else:
58         # Medium ecc - see eqn(5.14) of Peters, 1964
59         e = 0.0
60         de = e0 / 10000.0
61         summand = 0.0
62
63         while e < e0:
64             this_integral = de * e**(29.0/19.0)
65             this_integral *= (1.0 + (121.0/304.0) * e**2.0)**(1181.0/2299.0)
66             this_integral /= (1.0 - e**2.0)**(1.5)
67
68             summand += this_integral
69             e += de
70
71             e = float(e)
72
73         tC = (12.0/19.0) * (c0**4.0 / beta) * summand
74
75     return tC
```

APPENDIX B

---

# Code Versions

| Code Name  | Version | Reference               |
|------------|---------|-------------------------|
| HOKI       | 1.7.1   | Stevance et al. (2020a) |
| KAITIAKI   | 1.0.0   | This work               |
| MATPLOTLIB | 3.8.2   | Hunter (2007)           |
| NUMBA      | 0.58.1  | Lam et al. (2015)       |
| NUMPY      | 1.26.3  | Harris et al. (2020)    |
| PANDAS     | 2.1.4   | McKinney (2010)         |
| PYTHON     | 3.11.5  |                         |
| SCIPY      | 1.11.4  | Virtanen et al. (2020)  |
| TAKAHE     | 1.0.0   | Richards et al. (2023)  |
| TUI        | 8       | Ghodla et al. (2023)*   |

**Table B.1:** The code versions we use in this thesis. \*TUI was described in earlier papers, but Ghodla et al. (2023) was the first to name it.




## APPENDIX C

---

# Period-Eccentricity distributions along the LVK Collaboration O3 contour

In Figure C.1 we display the period-eccentricity distributions along the LVK Collaboration O3 contour. We also display a white dashed line corresponding to a merger time contour of the Hubble time. Most plots have a white region in the lower right corner of the plot, which corresponds to no systems passing through those bins. We use the definition of a Milky-Way Equivalent Galaxy (MWEG) from Kasen et al. (2017), namely that there is one MWEG per  $(4.4 \text{ Mpc})^3$ .



**Figure C.1:** The period-eccentricity distributions of populations governed by Bray kicks across the LIGO O3 contour. The combination of  $\alpha$  and  $\beta$  used in the kick prescription is given in white along the left side of each plot. White regions denote no systems passing through those bins.

## APPENDIX D

## Parameter Configuration

In this appendix we list the default parameters chosen for our binary evolution runs, as detailed in subsection 2.6.1.

| Parameter | Value     | Comment        | Parameter | Value     | Comment | Parameter | Value     | Comment        |
|-----------|-----------|----------------|-----------|-----------|---------|-----------|-----------|----------------|
| NM2       | 199       | Meshpoints     | ICL       | 1         |         | IT1       | 40        |                |
| ION       | 2         |                | IT2       | 20        |         | IAM       | 1         |                |
| JIN       | 15        |                | IOP       | 5         |         | OUT       | 15        |                |
| INUC      | 0         |                | NCH       | 3         |         | IBC       | 0         |                |
| JP        | 1         |                | ICN       | 0         |         | ITH       | 1         |                |
| IML1      | 5         | Mass loss (*1) | IX        | 1         |         | IML2      | 5         | Mass loss (*2) |
| IY        | 1         |                | ISGTH     | 0         |         | IZ        | 1         |                |
| IMO       | 0         |                | IMODE     | 2         |         | IDIFF     | 0         |                |
| NWRT1     | 100       |                | EPS       | $10^{-6}$ |         | NWRT2     | 1         |                |
| DEL       | 0.01      |                | NWRT3     | 2         |         | DHO       | $10^{-8}$ |                |
| NWRT4     | 1         |                | DT3       | 0         |         | NWRT5     | 0         |                |
| DDD       | 1.0       |                | NSAVE     | 10        |         | NMONT     | 0         |                |
| DT1       | 0.1       |                | DT2       | 1.2       |         | CT1       | 3.0       |                |
| CT2       | 3.0       |                | CT3       | 0.05      |         | CT4       | 0.01      |                |
| CT5       | 0.6       |                | CT6       | 0.35      |         | CT7       | 0.45      |                |
| CT8       | $10^{-4}$ |                | CT9       | 0.1       |         | CT10      | 0.3       |                |

**Table D.1:** Our general initial conditions. Unless otherwise described, these are assumed to be enabled for all runs.

The possible mass loss prescriptions (IML1/IML2) are:

0. No mass loss

1. Reimers (1975)

2. Blöcker (1995)

3. Vassiliadis & Wood (1993)

4. de Jager et al. (1988); Nugis & Lamers (2000); Vink et al. (2001) with interpolation errors described in **[TODO: Richards et al 2024]**

5. de Jager et al. (1988); Nugis & Lamers (2000); Vink et al. (2001) (described in Eldridge et al., 2017)

6. de Jager et al. (1988); Nugis & Lamers (2000); Vink et al. (2001) with slowly turning on WR rates described in **[TODO: Richards et al 2024]**

| Parameter | Value            | Comment | Parameter | Value           | Comment | Parameter | Value  | Comment          |
|-----------|------------------|---------|-----------|-----------------|---------|-----------|--------|------------------|
| ZS        | 0.02             |         | ALPHA     | 1.925           |         | CH        | 0.7    |                  |
| CC        | 0.173            |         | CN        | 0.053           |         | CO        | 0.482  |                  |
| CNE       | 0.099            |         | CMG       | 0.038           |         | CSI       | 0.08   |                  |
| CFE       | 0.072            |         | RCD       | 10 <sup>6</sup> |         | OS        | 0.12   |                  |
| RML       | 0                |         | RMG       | 0               |         | ECA       | 0      |                  |
| XF        | 0.1              |         | DR        | 0.001           |         | RMT       | 1000.0 |                  |
| RHL       | 0                |         | AC        | 0               |         | AK1       | 1.0    |                  |
| AK2       | 10 <sup>-4</sup> |         | ECT       | 0               |         | TRB       | 0      |                  |
| IRAM      | 1                |         | IRS1      | 1               |         | VROT1     | 0      |                  |
| IRS2      | 1                |         | VROT2     | 10.0            |         | FMAC      | 1.0    |                  |
| FAM       | 0                |         | IVMC      | 0               |         | TRC1      | 700.0  |                  |
| IVMS      | 0                |         | TRC2      | 0               |         | MWTS      | 0      |                  |
| IAGB      | 0                |         | ISGFAC    | 0               |         | FACSGMIN  | 0.01   |                  |
| SGTHFAC   | 0                |         | ISTART    | 0               |         | HKH       | 1.0    |                  |
| GFF       | 10.0             | Removed | NNMOD     | 0               |         | CEPR      | 0      | CEE prescription |
| ALPHACE   | 1.0              |         | -         | -               | -       | -         | -      | -                |

Table D.1: Continued

7. de Jager et al. (1988); Nugis & Lamers (2000); Vink et al. (2001) with no WR rates, described in **[TODO: Richards et al 2024]**
8. Unimplemented
9. ZAMS evolution to a specific ZAMS mass, custom routine by Jan Eldridge  
**[TODO: cite >:)]**



# Bibliography

- Abbott B. P., et al., 2016, Phys. Rev. Lett., 116, 061102
- Abdul-Masih M., et al., 2019, The Astrophysical Journal, 880, 115
- Abdul-Masih M., Escorza A., Menon A., Mahy L., Marchant P., 2022, A&A, 666, A18
- Agazie G., et al., 2023, The Astrophysical Journal Letters, 951, L8
- Alighieri D., ca. 1317, Paradiso
- Almeida L. A., et al., 2015, ApJ, 812, 102
- Amaro-Seoane P., et al., 2012, Classical and Quantum Gravity, 29, 124016
- Andersen J., Grønbech B., 1975, A&A, 45, 107
- Antoniadis J., et al., 2023, Astronomy & Astrophysics, 678, A50
- Arakaki T., et al., 2023, JuliaPy/pyjulia: v0.6.2, doi:10.5281/zenodo.10443741, <https://doi.org/10.5281/zenodo.10443741>
- Arcavi I., et al., 2017, Nature, 551, 64
- Arnett W. D., Bahcall J. N., Kirshner R. P., Woosley S. E., 1989, ARA&A, 27, 629
- Arzoumanian Z., Chernoff D. F., Cordes J. M., 2002, ApJ, 568, 289
- Baade W., Zwicky F., 1934, Contributions from the Mount Wilson Observatory, 3, 73
- Baker J. G., Boggs W. D., Centrella J., Kelly B. J., McWilliams S. T., Miller M. C., van Meter J. R., 2008, ApJ, 682, L29
- Ballesteros F. J., 2012, EPL (Europhysics Letters), 97, 34008
- Bars I., Terning J., 2010, Extra Dimensions in Space and Time, doi:10.1007/978-0-387-77638-5, <https://ui.adsabs.harvard.edu/abs/2010edst.book.....B>
- Barstow M. A., Bond H. E., Holberg J. B., Burleigh M. R., Hubeny I., Koester D., 2005, MNRAS, 362, 1134
- Belczynski K., et al., 2020, Astronomy & Astrophysics, 636, A104
- Bezanson J., Edelman A., Karpinski S., Shah V. B., 2017, SIAM Review, 59, 65
- Bhattacharjee P., Chaudhury S., Kundu S., 2014, The Astrophysical Journal, 785, 63
- Blaauw A., 1961, Bull. Astron. Inst. Netherlands, 15, 265
- Blanchet L., 2014, Living Reviews in Relativity, 17, 2
- Blöcker T., 1995, Astronomy and Astrophysics, 297, 727
- Boffin H. M. J., Jorissen A., 1988, Astronomy and Astrophysics, 205, 155

- Bombaci I., Popov S. B., 2004, *A&A*, 424, 627
- Bondi H., Hoyle F., 1944, *Monthly Notices of the Royal Astronomical Society*, 104, 273
- Bouffanais Y., Mapelli M., Santoliquido F., Giacobbo N., Di Carlo U. N., Rastello S., Artale M. C., Iorio G., 2021, *Monthly Notices of the Royal Astronomical Society*, 507, 5224
- Bray J. C., Eldridge J. J., 2016, *MNRAS*, 461, 3747
- Bray J. C., Eldridge J. J., 2018, *MNRAS*, 480, 5657
- Broekgaarden F. S., et al., 2021, *Monthly Notices of the Royal Astronomical Society*, 508, 5028
- Buck P. H., 1938, *Vikings of the sunrise*. New Zealand Electronic Text Centre, Wellington, N.Z
- Burbidge E. M., Burbidge G. R., Fowler W. A., Hoyle F., 1957, *Reviews of Modern Physics*, 29, 547
- Burdge K. B., et al., 2020, *ApJ*, 905, L7
- Burns J. A., 1978, *Vistas in Astronomy*, 22, 193
- Böhm-Vitense E., 1958, *Z. Astrophys.*, 46, 108
- Cameron A. D., et al., 2018, *MNRAS*, 475, L57
- Carroll B. W., Ostlie D. A., 2017, *An Introduction to Modern Astrophysics*, 2 edn. Cambridge University Press, doi:10.1017/9781108380980
- Champion D. J., Lorimer D. R., McLaughlin M. A., Cordes J. M., Arzoumanian Z., Weisberg J. M., Taylor J. H., 2004, *MNRAS*, 350, L61
- Christensen N., 1992, *Phys. Rev. D*, 46, 5250
- Christensen N., 2018, *Reports on Progress in Physics*, 82, 016903
- Coppersmith D., Winograd S., 1987, in *Proceedings of the Nineteenth Annual ACM Symposium on Theory of Computing*. STOC '87. Association for Computing Machinery, New York, NY, USA, pp 1–6, doi:10.1145/28395.28396, <https://doi.org/10.1145/28395.28396>
- Corbett G., Webb H., 2016, *Vertical Readings in Dante's Comedy: Volume 2*. JSTOR Open Access monographs, Open Book Publishers, <https://books.google.co.nz/books?id=5hmuDQAAQBAJ>
- Corongiu A., Kramer M., Stappers B. W., Lyne A. G., Jessner A., Possenti A., D'Amico N., Löhmer O., 2007, *A&A*, 462, 703
- Cox J. P., Giuli R. T., 1968, *Principles of stellar structure*. <https://ui.adsabs.harvard.edu/abs/1968pss.book.....C>
- Crowther P. A., Castro N., Evans C., Vink J., Melnick J., Selman F., 2018, Technical report, *Dissecting the Core of the Tarantula Nebula with MUSE*, <https://ui.adsabs.harvard.edu/abs/2018arXiv180100855C>, <https://ui.adsabs.harvard.edu/abs/2018arXiv180100855C>
- de Jager C., Nieuwenhuijzen H., van der Hucht K. A., 1988, *A&AS*, 72, 259
- de Loore C., de Grève J. P., de Cuyper J. P., 1975, *Ap&SS*, 36, 219
- Eddy J. A., Ise R., 1979, *A new sun : the solar results from SKYLAB*. National Aeronautics and Space Administration, <https://ui.adsabs.harvard.edu/abs/1979nssr.book.....E>

- Eggleton P. P., 1967, MNRAS, 135, 243
- Eggleton P. P., 1971, MNRAS, 151, 351
- Eggleton P. P., 1983, ApJ, 268, 368
- Eichler D., Livio M., Piran T., Schramm D. N., 1989, Nature, 340, 126
- Einstein A., 1905, Annalen der Physik, 322, 891
- Einstein A., 1916, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften, pp 688–696
- Einstein A., 1917, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften, pp 142–152
- El-Badry K., Conroy C., Fuller J., Kiman R., van Roestel J., Rodriguez A. C., Burdge K. B., 2022, MNRAS, 517, 4916
- Eldridge J. J., Tout C. A., 2004a, MNRAS, 348, 201
- Eldridge J. J., Tout C. A., 2004b, MNRAS, 353, 87
- Eldridge J. J., Tout C. A., 2018, The Structure And Evolution Of Stars. World Scientific
- Eldridge J. J., Stanway E. R., Xiao L., McClelland L. a. S., Taylor G., Ng M., Greis S. M. L., Bray J. C., 2017, Publ. Astron. Soc. Australia, 34, e058
- Eldridge J. J., Stanway E. R., Tang P. N., 2019, MNRAS, 482, 870
- Evans C. J., et al., 2011, Astronomy & Astrophysics, 530, A108
- Fang C., 2015, Highlights of Astronomy, 16, 19
- Faulkner A. J., et al., 2005, ApJ, 618, L119
- Fawzi A., et al., 2022, Nature, 610, 47
- Fonseca E., Stairs I. H., Thorsett S. E., 2014, ApJ, 787, 82
- Fragione G., 2021, The Astrophysical Journal, 923, L2
- Fragos T., et al., 2023, The Astrophysical Journal Supplement Series, 264, 45
- Fratanduono D. E., et al., 2020, Phys. Rev. Lett., 124, 015701
- Friedmann A., 1922, Z. Phys., 10, 377
- Fryer C., Burrows A., Benz W., 1998, ApJ, 496, 333
- Fuller R. S., Hamacher D. W., 2017, Journal of Astronomical History and Heritage, 20, 349
- Gamow G., 1930, Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character, 126, 632
- Ghodla S., Eldridge J. J., Stanway E. R., Stevance H. F., 2023, Monthly Notices of the Royal Astronomical Society, 518, 860
- Giacobbo N., Mapelli M., 2020, ApJ, 891, 141
- Goodwin S. P., Gribbin J., Hendry M. A., 1998, The Observatory, 118, 201

- Green M. J., Maoz D., Mazeh T., Faigler S., Shahaf S., Gorn R., El-Badry K., Rix H.-W., 2023, *MNRAS*, 522, 29
- Gunn J. E., Ostriker J. P., 1970, *ApJ*, 160, 979
- Hamacher D., 2022, *The First Astronomers: How Indigenous Elders read the stars*. Allen & Unwin, Crows Nest, New South Wales
- Hamacher D. W., Barsa J., Passi S., Tapim A., 2019, Indigenous use of stellar scintillation to predict weather and seasonal change, doi:10.48550/arXiv.1903.01060, <https://ui.adsabs.harvard.edu/abs/2019arXiv190301060H>
- Harris C. R., et al., 2020, *Nature*, 585, 357
- Hayashi C., 1961, *PASJ*, 13, 450
- Heney L. G., Lelevier R., Levée R. D., 1955, *PASP*, 67, 154
- Heney L. G., Forbes J. E., Gould N. L., 1964, *ApJ*, 139, 306
- Hilditch R. W., Howarth I. D., Harries T. J., 2005, *MNRAS*, 357, 304
- Hobbs G., Lorimer D. R., Lyne A. G., Kramer M., 2005, *MNRAS*, 360, 974
- Hoffleit D., 1964, *Catalogue of Bright Stars*
- Hoyle F., 1946, *MNRAS*, 106, 343
- Hoyle F., 1960, *MNRAS*, 120, 22
- Hulse R. A., Taylor J. H., 1975, *ApJ*, 195, L51
- Hunter J. D., 2007, *Computing in Science & Engineering*, 9, 90
- Hurley J. R., Tout C. A., Pols O. R., 2002, *MNRAS*, 329, 897
- Igoshev A. P., Chruslinska M., Dorozsmai A., Toonen S., 2021, *Monthly Notices of the Royal Astronomical Society*, 508, 3345
- Itoh Y., Futamase T., 2003, *Physical Review D*, 68, 121501
- Ivezić u., et al., 2019, *ApJ*, 873, 111
- Janssen G. H., Stappers B. W., Kramer M., Nice D. J., Jessner A., Cognard I., Purver M. B., 2008, *A&A*, 490, 753
- Jermyn A. S., et al., 2023, *ApJS*, 265, 15
- Kapil V., Mandel I., Berti E., Müller B., 2023, *Monthly Notices of the Royal Astronomical Society*, 519, 5893
- Kasen D., Metzger B., Barnes J., Quataert E., Ramirez-Ruiz E., 2017, *Nature*, 551, 80
- Keith M. J., Kramer M., Lyne A. G., Eatough R. P., Stairs I. H., Possenti A., Camilo F., Manchester R. N., 2009, *MNRAS*, 393, 623
- Knipe G. F. G., 1971, *Republic Observatory Johannesburg Circular*, 8, 8
- Kobayashi C., et al., 2023, *The Astrophysical Journal*, 943, L12


- Koen C., 2022, *Monthly Notices of the Royal Astronomical Society*, 510, 1857
- Kramer M., et al., 2006, *Science*, 314, 97
- Kruckow M. U., Tauris T. M., Langer N., Kramer M., Izzard R. G., 2018, *Monthly Notices of the Royal Astronomical Society*, 481, 1908
- Kuiper G. P., 1941, *The Astrophysical Journal*, 93, 133
- LIGO Scientific Collaboration and Virgo Collaboration 2016, *Phys. Rev. Lett.*, 116, 061102
- LIGO-Virgo 2021, *ApJ*, 913, L7
- LIGO-Virgo-KAGRA 2021, arXiv e-prints, p. arXiv:2111.03634
- Lam S. K., Pitrou A., Seibert S., 2015, in *Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in HPC. LLVM '15*. Association for Computing Machinery, New York, NY, USA, pp 1–6, doi:10.1145/2833157.2833162, <https://dl.acm.org/doi/10.1145/2833157.2833162>
- Lamers H. J. G. L. M., Snow T. P., Lindholm D. M., 1995, *The Astrophysical Journal*, 455, 269
- Landolt A. U., 1964, *The Astrophysical Journal*, 140, 1494
- Langer N., Norman C. A., 2006, *ApJ*, 638, L63
- Lau R. M., et al., 2022, *Nature Astronomy*, 6, 1308
- Lazarus P., et al., 2016, *ApJ*, 831, 150
- Lemaître G., 1931, *MNRAS*, 91, 483
- Levesque E. M., Massey P., Żytkow A. N., Morrell N., 2014, *MNRAS: Letters*, 443, L94
- Li T., Li X.-D., 2013, *The Astrophysical Journal*, 780, 47
- Linial I., Sari R., 2017, *Monthly Notices of the Royal Astronomical Society*, 469, 2441
- Lorenzo J., Negueruela I., Val Baker A. K. F., García M., Simón-Díaz S., Pastor P., Méndez Majuelos M., 2014, *A&A*, 572, A110
- Madau P., Dickinson M., 2014, *ARA&A*, 52, 415
- Mandel I., 2021, *RNAAS*, 5, 223
- Mandel I., Broekgaarden F. S., 2022, *Living Reviews in Relativity*, 25, 1
- Mandel I., Müller B., 2020, *MNRAS*, 499, 3214
- Mapelli M., Giacobbo N., 2018, *Monthly Notices of the Royal Astronomical Society*, 479, 4391
- Marchant P., Langer N., Podsiadlowski P., Tauris T. M., Moriya T. J., 2016, *Astronomy and Astrophysics*, 588, A50
- Marov M. Y., 2015, *The Fundamentals of Modern Astrophysics*, doi:10.1007/978-1-4614-8730-2, <https://ui.adsabs.harvard.edu/abs/2015fma...book.....M>
- Martinez J. G., et al., 2015, *ApJ*, 812, 143
- Martinez J. G., et al., 2017, *ApJ*, 851, L29

- Martins F., Mahy L., Hervé A., 2017, *A&A*, 607, A82
- McKinney W., 2010, in Walt S. v. d., Millman J., eds, *Proceedings of the 9th Python in Science Conference*. pp 56 – 61, doi:10.25080/Majora-92bf1922-00a
- Mennekens N., Vanbeveren D., 2014, *Astronomy & Astrophysics*, 564, A134
- Mennekens N., Vanbeveren D., 2016, *Astronomy & Astrophysics*, 589, A64
- Moore C. J., Cole R. H., Berry C. P. L., 2014, *Classical and Quantum Gravity*, 32, 015014
- Morgenroth O., 1935, *Astronomische Nachrichten*, 255, 425
- Morrell N. I., Niemela V. S., 1990, in Garmany C. D., ed., *Astronomical Society of the Pacific Conference Series Vol. 7, Properties of Hot Luminous Stars*. pp 57–59
- Negueruela I., Marco A., 2003, *Astronomy & Astrophysics*, 406, 119
- Noon K., de Napoli K., 2022, *First Knowledges Astronomy: Sky Country*. Thames & Hudson
- Nugis T., Lamers H. J. G. L. M., 2000, *A&A*, 360, 227
- Oosterhoff P. T., 1928, *Bulletin of the Astronomical Institutes of the Netherlands*, 4, 183
- Oosterhoff P. T., 1930, *Bulletin of the Astronomical Institutes of the Netherlands*, 5, 156
- Pauldrach A. W. A., Puls J., 1990, *Astronomy and Astrophysics*, 237, 409
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, *ApJS*, 192, 3
- Paxton B., et al., 2013, *ApJS*, 208, 4
- Paxton B., et al., 2015, *ApJS*, 220, 15
- Paxton B., et al., 2018, *ApJS*, 234, 34
- Paxton B., et al., 2019, *ApJS*, 243, 10
- Payne C. H., 1925, PhD thesis, Harvard, <https://ui.adsabs.harvard.edu/abs/1925PhDT.....1P>
- Pearce J. A., 1952, *Publications of the Astronomical Society of the Pacific*, 64, 219
- Penny L. R., Ouzts C., Gies D. R., 2008, *ApJ*, 681, 554
- Peters P. C., 1964, *Phys. Rev. Lett.*, 136, 1224
- Petrosky E., Hwang H.-C., Zakamska N. L., Chandra V., Hill M. J., 2021, *Monthly Notices of the Royal Astronomical Society*, 503, 3975
- Pian E., et al., 2017, *Nature*, 551, 67
- Poincaré M. H., 1906, *Rendiconti del Circolo Matematico di Palermo (1884-1940)*, 21, 129
- Pols O. R., Tout C. A., Lattanzio J. C., Karakas A. I., 2001, in Podsiadlowski P., Rappaport S., King A. R., D’Antona F., Burderi L., eds, *Astronomical Society of the Pacific Conference Series Vol. 229, Evolution of Binary and Multiple Star Systems*. p. 31
- Price D. J., et al., 2018, *Publ. Astron. Soc. Australia*, 35, e031

- Ptolemy C., Proclus ca. 168, *Tetrabiblos*
- Qian S.-B., Zhu L.-Y., Liu L., Zhang X.-D., Shi X.-D., He J.-J., Zhang J., 2020, *Research in Astronomy and Astrophysics*, 20, 163
- Raucq F., Gosset E., Rauw G., Manfroid J., Mahy L., Mennekens N., Vanbeveren D., 2017, *A&A*, 601, A133
- Reardon D. J., et al., 2023, *The Astrophysical Journal Letters*, 951, L6
- Redd N. T., 2018, *Proceedings of the National Academy of Science*, 115, 12836
- Reimers D., 1975, *Memoires of the Societe Royale des Sciences de Liege*, 8, 369
- Reynolds T. M., Fraser M., Gilmore G., 2015, *MNRAS*, 453, 2885
- Richards S., 2021, Master's thesis, University of Auckland, doi:10.5281/zenodo.5523933, <https://zenodo.org/record/5523933>
- Richards S. M., Eldridge J. J., Briel M. M., Stevance H. F., Willcox R., 2023, *MNRAS*, 522, 3972
- Robbins F. E., 1940, *Tetrabiblos*. Harvard University Press Cambridge, MA
- Robertson H. P., 1935, *ApJ*, 82, 284
- Rose W. K., 1998, *Advanced Stellar Astrophysics*. Cambridge University Press
- Rucinski S. M., 1998, *ApJ*, 116, 2998
- Salpeter E. E., 1955, *ApJ*, 121, 161
- Santoliquido F., Mapelli M., Giacobbo N., Bouffanais Y., Artale M. C., 2021, *Monthly Notices of the Royal Astronomical Society*, 502, 4877
- Schmitz K., 2021, *JHEP*, 2021, 97
- Schwarzschild M., Härm R., 1965, *ApJ*, 142, 855
- Schwarzschild M., Härm R., 1967, *The Astrophysical Journal*, 150, 961
- Shakespeare W., ca. 1603, *The Tragedy of Othello, the Moor of Venice*
- Smartt S. J., et al., 2017, *Nature*, 551, 75
- Stancliffe R. J., 2005, PhD thesis, <https://ui.adsabs.harvard.edu/abs/2005PhDT.....352S>
- Stancliffe R. J., 2006, *Memorie della Societa Astronomica Italiana*, 77, 792
- Stancliffe R. J., 2016, *IAU Focus Meeting*, 29B, 600
- Stancliffe R. J., Eldridge J. J., 2009, *MNRAS*, 396, 1699
- Stancliffe R. J., Tout C. A., Pols O. R., 2004, *MNRAS*, 352, 984
- Stanway E. R., Eldridge J. J., 2018, *MNRAS*, 479, 75
- Stephenson C. B., Sanduleak N., 1971, *Publications of the Warner & Swasey Observatory*, 1, 1
- Stephenson F. R., Yau K. K. C., Hunger H., 1985, *Nature*, 314, 587
- Stevance H., 2022, Private Communication

- Stevance H., Eldridge J., Stanway E., 2020a, *The Journal of Open Source Software*, 5, 1987
- Stevance H. F., Eldridge J. J., McLeod A., Stanway E. R., Chrimes A. A., 2020b, *MNRAS*, 498, 1347
- Stevance H. F., Eldridge J. J., Stanway E. R., Lyman J., McLeod A. F., Levan A. J., 2023, *Nature Astronomy*, 7, 444
- Stothers R., 1972, *The Astrophysical Journal*, 175, 431
- Stothers R., 1973, *Publications of the Astronomical Society of the Pacific*, 85, 363
- Stovall K., et al., 2018, *ApJ*, 854, L22
- Sutantyo W., 1978, *Ap&SS*, 54, 479
- Swiggum J. K., et al., 2015, *ApJ*, 805, 156
- Tauris T. M., Takens R. J., 1998, *Astronomy and Astrophysics*, 330, 1047
- Tauris T. M., Langer N., Podsiadlowski P., 2015, *MNRAS*, 451, 2123
- Tauris T. M., et al., 2017, *The Astrophysical Journal*, 846, 170
- Team COMPAS: Riley J., et al., 2022, *ApJS*, 258, 34
- Terrell D., Munari U., Zwitter T., Nelson R., 2003, *ApJ*, 126, 2988
- Thorne K. S., Żytkow A. N., 1977, *ApJ*, 212, 832
- ud-Doula A., Sundqvist J. O., Owocki S. P., Petit V., Townsend R. H. D., 2013, *MNRAS*, 428, 2723
- van de Voort F., Pakmor R., Bieri R., Grand R. J. J., 2022, *MNRAS*, 512, 5258
- Various Authors 1611, *The Holy Bible: Conteyning the Old Testament, and the New: Newly Translated out of the Originall Tongues: & with the Former Translations Diligently Compared and Reuised*, by His Maiesties Speciall Commandement. Robert Barker
- Vassiliadis E., Wood P. R., 1993, *The Astrophysical Journal*, 413, 641
- Verbunt F., Igoshev A., Cator E., 2017, *A&A*, 608, A57
- Vigna-Gómez A., et al., 2018, *MNRAS*, 481, 4009
- Vink J. S., de Koter A., Lamers H. J. G. L. M., 1999, *Astronomy and Astrophysics*, 350, 181
- Vink J. S., de Koter A., Lamers H. J. G. L. M., 2001, *A&A*, 369, 574
- Virtanen P., et al., 2020, *Nature Methods*, 17, 261
- Vitense E., 1953, *Z. Astrophys.*, 32, 135
- Walker A. G., 1937, *Proc. London Math. Soc.*, 42, 90
- Wiktorowicz G., Wyrzykowski Ł., Chruslinska M., Klencki J., Rybicki K. A., Belczynski K., 2019, *The Astrophysical Journal*, 885, 1
- Willcox R., Mandel I., Thrane E., Deller A., Stevenson S., Vigna-Gómez A., 2021, *ApJ*, 920, L37

- Williams V. V., 2012, in Proceedings of the forty-fourth annual ACM symposium on Theory of computing. STOC '12. Association for Computing Machinery, New York, NY, USA, pp 887–898, doi:10.1145/2213977.2214056, <https://dl.acm.org/doi/10.1145/2213977.2214056>
- Wongwathanarat A., Janka H.-T., Müller E., 2013, A&A, 552, A126
- Xu H., et al., 2023, Research in Astronomy and Astrophysics, 23, 075024
- Yao Y., et al., 2020, ApJ, 900, 46
- Yaşarsoy B., Yakut K., 2012, The Astronomical Journal, 145, 9
- Zhao L., et al., 2023, Astronomy & Astrophysics, 674, A92
- van Zeist W., Stevance H., Eldridge J., 2021, The Journal of Open Source Software, 6, 2968
- van Zeist W. G. J., Eldridge J. J., Tang P. N., 2023, Monthly Notices of the Royal Astronomical Society, 524, 2836



Isa lei – Nanoqu rarawa,  
Miko sana vodo e na mataka,  
Bau nonuma, na nodatau lasa,  
Mai Suva, nonuma tikoga.

---

Isa Lei. The purple shadows fall,  
Sad the morrow will dawn upon my sorrow.  
Ah, forget not when you are far away  
Precious moments beside dear Suva Bay.

---

*Isa Lei, "The Fijian Farewell Song"*