

List of Figures



1.1	The locations of the ~ 9000 stars in the night sky visible to the naked eye.	4
2.1	A schematic of the reactions leading up to the CNO cycle.	17
2.2	The evolution of an $4 M_{\odot}$ star from ZAMS until CO core development.	20
2.3	The evolution of stars in the range $0.5 - 80 M_{\odot}$	21
2.4	The evolution of an $8 M_{\odot}$ star.	22
2.5	A rough outline of supernova classification.	24
2.6	The binding energy per nucleon and half-lives for elements up to Oganesson.	25
2.7	The HRD for an $21 M_{\odot}$ star with an (unshown) $10 M_{\odot}$ companion with varying period.	26
2.8	A schematic of a simple binary star system.	27
2.9	The dimensionless Roche potential as a function of mass ratio q	28
2.10	A schematic of the Roche lobe formula Eqn(2.15).	29
2.11	The volume of the Roche Lobe computed numerically through Monte-Carlo integration vs the volume assuming Eqn(2.15).	30
2.12	As for Figure 2.11, but for the $\mathcal{L}_2$ point.	30
2.13	Two stars of a similar mass that undergo a helium flash and two thermal pulses.	42
2.14	The helium flash shown on a Kippenhahn diagram.	43
3.1	The value of the meshpoint sum $\mathcal{C}$ in Eqn(3.2) throughout VCA.	51
3.2	The main evolution of a binary with VCA enabled.	53
3.3	As for Figure 3.2, but for the NVCA model.	54
3.4	The evolution of the radius, rotation rate, and luminosity for the VCA and NVCA models, shown around the contact phase.	55
3.5	The Kelvin-Helmholtz time-scales of both stars in the VCA and NVCA regime.	56
3.6	A Hertzsprung-Russell Diagram showing the evolution of our model both with and without the VCA prescription.	57
3.7	The abundance of helium at the five critical stages of the binary evolution.	59
3.8	The helium abundance in the accretor near when the model detaches.	61
3.9	An Hertzsprung-Russell Diagram of the post-mass transfer secondary model evolved as a single star with different thermohaline mixing coefficients.	62

3.10	The helium abundance profile for each model of thermohaline mixing, as a function of the internal mass of the star.	63
3.11	The (evolutionary) mass-luminosity relationship in the presence and absence of VCA.	66
4.1	The PDF and CDF of the distributions using kicks from Hobbs et al. (2005) ; Verbunt et al. (2017) ; Bray & Eldridge (2018) ; Willcox et al. (2021)	72
4.2	A schematic of how FINEGRID and WIDEGRID are related.	73
4.3	The effect of varying d_e in Eqn(4.5) on the coalescence time.	76
4.4	The effect of varying d_e in Eqn(4.5) on the runtime and performance.	76
4.5	The merger rate as a function of kick parameters α and β	77
4.6	The period-eccentricity distributions for six selected Bray & Eldridge (2016) kicks.	78
4.7	The $\ln \mathcal{K}$ values across $\alpha - \beta$ space.	80
4.8	The KS statistic for FINEGRID and WIDEGRID.	81
4.9	The PDFs and CDFs of four systems, alongside the Hobbs CDF and Willcox et al. (2021) CDF.	82
4.10	The four constraints we apply overlaid on our FINEGRID parameter space.	84
4.11	The CDF for a population of isolated pulsars and the period-eccentricity distribution for $\alpha = 115 \text{ km s}^{-1}, \beta = 15 \text{ km s}^{-1}$	85
4.12	The peak of the event rate $\mathcal{R}(t)$ as a function of α and β	86
4.13	The event rate $\mathcal{R}(z)$ for a region surrounding $\alpha = 330, \beta = -70$ on FINEGRID.	87
5.1	BPASS v2.2.1 primary models for the matches obtained by HOKI for the parameters in Table 5.1.	99
5.2	BPASS v2 model tracks for our six close-contact binaries.	100
5.3	Detailed model tracks for our six close-contact binaries.	103
5.4	The mass-radius plots for our models.	104
5.5	HRDs for the matching overcontact binary models.	106
5.6	The three fits for TU Muscae we recover.	109
5.7	The dependence of the mean contact length on the initial parameters $\log P$, M_1 , and q	111
6.1	The geometry of an ellipse and the circle with a radius equal to the semi-latus rectum.	116
A.1	A schematic of a simple binary star system.	124
A.2	An infinitesimal area segment of an orbital trajectory.	127
A.3	The Chebyshev polynomial expansion of Eqn(A.92).	135
A.4	An illustration of the bi-stability jumps in the mass loss prescription.	138
C.1	The period-eccentricity distributions of populations governed by Bray kicks across the LIGO O3 contour.	145
D.1	The results of our bespoke binary evolution for normal periods.	153
D.2	The results of our bespoke binary evolution for wider periods.	157

D.3 The results of our bespoke binary evolution for shorter periods.	161
--	-----

