



CT scanning: patterns of use and dose

Fred A Mettler Jr, Philip W Wiest, Julie A Locken and Charles A Kelsey

Published under licence by IOP Publishing Ltd

Journal of Radiological Protection, Volume 20, Number 4

Citation Fred A Mettler Jr *et al* 2000 *J. Radiol. Prot.* **20** 353

DOI 10.1088/0952-4746/20/4/301

1. Received 31 July 2000
2. Accepted 22 August 2000

[Buy this article in print](#)

 Journal RSS

 Sign up for new issue notifications

Abstract

CT scanning is a relatively high-dose procedure. In spite of the use of magnetic resonance imaging, with faster CT scanners and helical techniques CT is becoming more common. There are few data from practice in the United States regarding the age and sex distribution of patients receiving CT scans, what type of scan and how many scans they receive, or how much radiation dose CT scans contribute.

We reviewed over 33 700 consecutive CT examinations done at our institution in 1998 and 1999. Information on the types of scans as well as the age and sex distribution of the patients was determined.

Between 1990 and 1999, CT examinations in our institution increased from 6.1% to 11.1% of all radiology procedures. Nineteen per cent of all patients seen in our department in the last year had at least one CT scan and more than half had multiple scans on the same day. Thirty-six per cent of all patients had a prior CT examination done on an earlier date. The male/female ratio of patients was 56/44. Studies of children age 0-15 years comprised 11.2% of scans. The highest percentage of scans was done in the 36-50-year-old age group. CT scanning accounted for 67% of the effective dose from diagnostic radiology.

In most large hospitals in the United States CT scanning probably accounts for more than 10% of diagnostic radiology examinations and about two-thirds of the radiation dose. Most patients have multiple scan sequences. Studies done on children are probably more common than previously thought.

Export citation and abstract

BibTeX

RIS

← **Previous** article in issue

Next article in issue →

Access this article

The computer you are using is not registered by an institution with a subscription to this article. Please choose one of the options below.

Login

Access through your institution

[IOPscience login](#)

[SRP member access](#)

Find out more about journal [subscriptions at your site](#).

Purchase from

[Article Galaxy](#).

[CCC RightFind](#)

Purchase this article from our trusted document delivery partners.

Make a recommendation

To gain access to this content, please complete the [Recommendation Form](#) and we will follow up with your librarian or Institution on your behalf.

For corporate researchers we can also follow up directly with your R&D manager, or the information management contact at your company. Institutional subscribers have access to the current volume, plus a 10-year back file (where available).

You may also like

JOURNAL ARTICLES

Feasibility of iodine contrast enhanced CT-scan during a 18F-fluorodeoxyglucose Positron Emission Tomography

Multi-scale CT Scanning and Digital Rock Modeling of Tight Sandstone Reservoirs

Risks from CT scans—what do recent studies tell us?

Absolute and relative dose–surface and dose–volume histograms of the bladder: which one is the most representative for the actual treatment?

Spiral computed tomography phase-space source model in the BEAMnrc/EGSnrc Monte Carlo system: implementation and validation

Evaluation of Effective Doses of Radiation and Risk Factors in Thorax-Testing CT Scans of Patients with Lung Cancer Cases

IOPSCIENCE[Journals](#)[Books](#)[IOP Conference Series](#)[About IOPscience](#)[Contact Us](#)[Developing countries access](#)[IOP Publishing open access policy](#)[Accessibility](#)**IOP PUBLISHING**[Copyright 2024 IOP Publishing](#)[Terms and Conditions](#)[Disclaimer](#)[Privacy and Cookie Policy](#)[Text and Data mining policy](#)**PUBLISHING
SUPPORT**[Authors](#)[Reviewers](#)[Conference Organisers](#)